

BISPECIFIC PROTEINS AND METHODS FOR PREPARING SAME

Technical Field

The present invention relates to a bispecific protein with a high heterodimerization rate
5 and a method of preparing a bispecific protein.

Background Art

A bispecific antibody (BsAb), which is an antibody having two paratopes capable of recognizing two different types of target antigens, is a generic term for an antibody capable of
10 binding simultaneously to two different target antigens or for an antigen binding fragment thereof. This feature allows for the establishment of a therapeutic strategy which is impossible with conventional monoclonal antibodies. Bispecific antibodies rarely occur naturally and are in most part artificially constructed to simultaneously bind to two different types of biological targets. The double-targeting ability can provide such BsAbs with new applicable fields which
15 have not been managed by monospecific antibodies (MsAbs). When it comes to therapeutic purposes, considerable interest is arising in (1) reliable recruitment of immune cells into the proximity of target cells, (2) inhibition or activation of two distantly separated signaling pathways in target cells to create a synergistic effect, and (3) specific and regulatory delivery of therapeutic substance, medications, toxins, etc. to target cells.

20 A variety of BsAb-related techniques (45 different formats) has been developed. These techniques are classified into four categories on a structural basis: first, heterologous bispecification of heavy chains by various methods comprising structural complementarity, also known as Knob-into-Hole or simply KiH, electrostatic steering effect, and CH3 domain shuffling (called SEEDbodyTM); second, various antibody fragment formats such as DiabodyTM
25 (Diabody: dimeric antibody fragments) BiTETM (BiTE: Bi-specific T-cell engagers), and DARTTM (DART: dual affinity retargeting bispecific antibody); third, technology using one or more functional domains combined with intact antibodies, such as Modular AntibodyTM, ZybobyTM, dAbsTM (dAbs : Single domain antibodies), and DVD-IgTM (DVD-Ig: dual-variable-domain immunoglobulin); and fourth, techniques adopting a full-length IgG-like scheme such as
30 DuobodyTM (Fab-Arm Exchange), CrossMabTM, AzymetricTM, and kI bodyTM have been developed.

For example, U. S. Patent No. 2013-892198 A to Zymeworks discloses a heteropolymer structure of immunoglobulin chains having mutations in the Fc domain, specifically stating that antibodies of the heteropolymer structure can be constructed by modifying cysteine residues involved in disulfide bonds into charged amino acids to exert an electrical interaction.

5 There are problems with conventional techniques of constructing bispecific antibodies as follows.

First, there are undesirable combinations of heavy chains. When heavy chains (A and B) originating respectively from two antibodies targeting different epitopes randomly combine with each other, two combinations of the same origins (AA and BB) and one combination of different
10 origins (AB or BA) are formed. Combinations between heavy chains from the same origins act as undesired impurities in constructing bispecific antibodies, decreasing the purity of bispecific antibodies and requiring a process of removing the combinations. Therefore, they may have disadvantageous effects, such as provoking difficulty in the isolation and purification of antibodies, inciting undesired immune responses or signal transduction to cause side effects, etc.
15 Hence, ongoing needs exist for forming only combinations between heavy chains of heterogeneous origins allowing none or up to a considerably restricted level of combinations between heavy chains of homogeneous origin.

Second, although light and heavy chains of homogeneous origins are specifically combined with each other, cases occur in which the light and heavy chains that combine with
20 each other are of heterogeneous origins. A bispecific antibody targets two different kinds of epitopes. Each epitope can be recognized by an antibody in which light and heavy chains of the same origin are combined. Antibodies that are created through combinations of light and heavy chains from heterogeneous origins may recognize new epitopes other than the desired two epitopes, act as an impurity making it difficult to isolate and purify antibodies of interest, and
25 cause a side effect. Therefore, the production of bispecific antibodies requires combinations of correct pairs of light and heavy chains while a non-specific combination of light and heavy chains does not occur or occurs at up to an insignificant level.

In consideration of the above-mentioned problems with the provision of bispecific antibodies, requirements for construction of bispecific antibodies are summarized as follows. A
30 total of ten combinations is possible when bispecific antibodies are constructed with light and heavy chains from two different kinds of antibodies respectively recognizing two different

epitopes, as shown in FIG. 1. Of them, the combination marked by a dotted line circle meets the requirements that (1) heavy chains derived from different antibodies are combined with each other and (2) light and heavy chains derived from the same antibodies are coupled to each other. The other nine combinations should be not formed or should be formed at a minimum level.

5 WO2014142591, which is a previous patent application of the present inventor, introduces “protein in which electrical interaction is introduced within a hydrophobic interaction site and preparation method therefor”. Disclosed in the patent application is a protein or an antibody having an electrical interaction introduced in a hydrophobic interaction site thereof wherein the electrical interaction is made by a positive charge and a negative charge on a
10 positively charged substance and a negatively charged substance which are changed from a pair of hydrophobic amino acids selected in the hydrophobic interaction site. Leading to the present invention, intensive and thorough research made by the present inventors resulted in the finding that bispecific antibodies can be formed using electrostatic interaction in a non-hydrophobic interaction site which was not taken into consideration in the previous patent application and can
15 be constructed at high yield even by size-dependent coupling and/or amino acid change (swapping) between coupled pairs.

[Prior Art Document]

[Patent Document]

20 (Patent Document 1) Korean Patent Number 10-2014-0019385 A (February 14, 2014)

Disclosure

Technical Problem

In order to solve the above problems, the present invention provides a bispecific
25 antibody with high purity and a method of preparing the same.

An embodiment provides a dimer comprising a first CH3 domain and a second CH3 domain of an antibody, or Fc regions comprising the CH3 domains, wherein

the first CH3 domain and the second domain are mutated such that at least one selected from among amino acid pairs forming amino acid-amino acid bonds between the first CH3
30 domain and the second CH3 domain is modified by at least one of the following mutations:

(1) a mutation in which, of at least one amino acid pair between the CH3 domains, one

amino acid is substituted with an amino acid having a positive charge and the other is substituted with an amino acid having a negative charge;

(2) a mutation in which amino acids in at least one amino acid pair between the CH3 domains are swapped with each other; and

5 (3) a mutation in which, of at least one amino acid pair between the CH3 domains, one amino acid is substituted with a large hydrophobic amino acid while the other is substituted with a small hydrophobic amino acid.

The first CH3 domain and the second CH3 domain may be derived from the same or different kinds of antibodies (immunoglobulins).

10 Another embodiment provides a nucleic acid molecule encoding the modified CH3 domain or a modified Fc region comprising the modified CH3 domain, a recombinant vector carrying the nucleic acid molecule, and a recombinant cell containing the recombinant vector therein.

As used herein the terms “one amino acid” and “the other amino acid” in the expression
15 “of at least one amino acid pair, one amino acid... and the other amino acid...” mean one amino acid and the other amino acid of the two amino acids in each of one or more amino acid pairs, respectively (hereinafter the same definition will be applied).

Another embodiment provides a bispecific protein for targeting two different kinds of targets, the bispecific protein comprising a first CH3 domain or a first Fc region comprising the
20 first CH3 domain and a second CH3 domain or a second Fc region comprising the second CH3 domain, wherein the first CH3 domain and the second CH3 domain are mutated such that at least one selected from amino acid pairs forming amino acid-amino acid bonds between the first CH3 domain and the second CH3 domain is modified by at least one of the following mutations:

(1) a mutation in which, of at least one amino acid pair between the CH3 domains, one
25 amino acid is substituted with an amino acid having a positive charge while the other is substituted with an amino acid having a negative charge;

(2) a mutation in which amino acids in at least one amino acid pair between the CH3 domains are swapped with each other; and

(3) a mutation in which, of at least one amino acid pair between the CH3 domains, one
30 amino acid is substituted with a large hydrophobic amino acid while the other is substituted with a small hydrophobic amino acid.

The amino acid pair that is substituted with amino acids having opposite charges in step (1) may be an amino acid under hydrophobic interaction and/or an amino acid under non-hydrophobic interaction. In one embodiment, at least one of the two amino acids constituting an amino acid pair may be not a hydrophobic amino acid (i.e., a hydrophilic amino acid). The electrostatic interaction that is introduced by step (1) of substitution with amino acids having opposite charges may contribute to improving the formation of a heterodimer between Fc regions.

As used herein, the term “heterodimer” means a fusion body in which two different proteins are coupled to each other and is intended to encompass any bispecific protein in which two proteins targeting respective different targets are coupled to each other (e.g., bispecific antibody), etc.

The amino acid switching (swapping) in step (2) may be conducted in any amino acid pair forming an amino acid-amino acid linkage in the Fc regions or CH3 domains and, for example, in at least one amino acid pair selected from all the amino acid pairs which are each bonded through interaction other than electrostatic interaction, hydrophobic interaction, and amino acid size difference-based interaction. As such, two amino acid residues that interact with each other (e.g., bond themselves to each other) in each amino acid pair (i.e., an amino acid residue of the first CH3 domain and the other amino acid of the second CH3 domain in the interacting amino acid pair) may be exchanged with each other to decrease the possibility of forming a homodimer because the presence of the same amino acid at the counterpart positions for amino acid interaction between the CH3 domains of homogeneous origins makes the bonding therebetween difficult, thereby contributing to improving a heterodimerization rate (for example, when an amino acid pair of S364 and K370 undergoes S364K mutation in the first domain and K370S mutation in the second CH3 domain, there is no interaction between the homogeneous CH3 domains because both amino acids at positions 364 and 370 become lysine (K) in the first CH3 domain and serine (S) in the second CH3 domain, but interaction occurs between the heterogeneous CH3 domain to form a heterodimer only).

The step (3) of substitution with amino acids different in size improves structural engagement suitability between a large amino acid and a small amino acid (that is, a large amino acid is inserted into a spare space established by a small amino acid, thereby increasing bonding efficiency), with the consequent increase of heterodimerization rates. Particularly, an interacting

amino acid pair is mutated such that one amino acid is substituted with a large hydrophobic amino acid while the other amino acid is substituted with a small hydrophobic amino acid whereby advantage is taken of the difficulty in making a bond between large amino acids or between small amino acids to minimize a homodimerization rate (large amino acids, if existing
5 respectively in two opposite chains, render the two chains distant from the each other to obstruct dimerization whereas two small amino acids, if existing in two opposite chains, interact with each other at low possibility because of a long distance therebetween and have difficulty in interaction therebetween). On the contrary, a large hydrophobic amino acid in one CH3 domain or Fc and a small hydrophobic amino acid in the other CH3 domain or Fc undergo hydrophobic
10 interaction with each other at a closer distance compared to the pre-mutation amino acids, thus making a condition good for heterodimerization. Therefore, a large and a small amino acid to be substituted in the step of substitution with amino acids different in size may be both selected from among hydrophobic amino acids. For example, the large amino acid may be at least one selected from the group consisting of tryptophan and phenylalanine, which are both
15 hydrophobic. In addition, the small amino acid may be at least one selected from the group consisting of alanine, glycine, and valine, which are all hydrophobic.

The bispecific protein comprising the mutant Fc regions or CH3 domains may be selected from among any type of proteins targeting (e.g., specifically recognizing and/or binding to) two different kinds of targets. For targeting, the bispecific protein comprising the mutant Fc
20 regions or CH3 domains may comprise two targeting domains capable of targeting (specifically recognizing and/or binding to) two different kinds of targets, respectively (for example, a first targeting domain for targeting a first target and a second targeting domain for targeting a second target). The targeting domains may form a covalent or non-covalent bond (linkage) to the mutant Fc regions or CH3 domains, respectively, in a direct or indirect (e.g., via a linker)
25 manner. For example, the bispecific proteins comprising the mutant Fc regions or CH3 domains may be at least one selected from the group consisting of a bispecific antibody, an antigen-binding fragment of a bispecific antibody (e.g., (scFv-Fc)₂, etc.), a bispecific antibody analog (e.g., nanobody, peptibody, peptide, aptide, etc.), and a fusion protein of a target-specific binding polypeptide and the mutant Fc region or CH3 domain.

30 The target-specific binding polypeptide may be any polypeptide that binds specifically to a biological target substance (any compounds present in the body comprising proteins, nucleic

acids, and the like) and may be at least one polypeptide selected from, for example, the group consisting of a paratope (e.g., e.g., a CDR or variable region of a heavy chain and/or a light chain), single-chain Fv (scFv), a membrane protein (e.g., various receptors, etc.), a membrane protein ectodomain, and a ligand (e.g., various growth factors, cytokines, etc.). In one
5 embodiment, the fusion protein of a target-specific binding polypeptide and the mutant Fc region or CH3 domain may be at least one selected from the group consisting of a fusion protein of a membrane protein and the mutant Fc region or CH3 domain, a fusion protein of a membrane protein ectodomain and the mutant Fc region or CH3 domain, a fusion protein of a ligand and the mutant Fc region or CH3 domain, and a fusion protein of scFv and the mutant Fc region or
10 CH3 domain.

When the bispecific protein comprising the mutant Fc region or CH3 domain is an antibody, an antigen-binding fragment of an antibody, or an antibody analog, the targeting domain may be a paratope (e.g., e.g., a CDR or variable region of a heavy chain and/or a light chain). For the above-mentioned fusion protein of a target-specific binding polypeptide and the
15 mutant Fc region or CH3 domain, the target-specific binding polypeptide may be at least one selected from the group consisting of a membrane protein (e.g., various receptors), a membrane protein ectodomain, a ligand (e.g., various growth factors, cytokines, etc.), and a paratope (e.g., a CDR or variable region of a heavy chain and/or a light chain).

The two different kinds of targets may refer to two different kinds of biological
20 substances (e.g., proteins) or different regions within one biological substance (e.g., one protein). The bispecific protein comprising the mutant Fc region or CH3 domain is characterized by an increase in heterodimerization rate, a decrease in homodimerization, and/or stability, compared to a bispecific protein comprising a non-mutant (wild-type) Fc region or CH3 domain.

Another embodiment provides a bispecific antibody or an antigen-binding fragment
25 thereof, the bispecific antibody comprising a first CH1 domain and a first CL (light chain constant region) domain derived respectively from the heavy chain and light chain of an antibody recognizing a first epitope and a second CH1 domain and a second CL domain derived respectively from the heavy chain and light chain of an antibody recognizing a second epitope, wherein the CH1 domains and the CL domains are mutated to contain at least one of the
30 following mutations:

a mutation in which, of the two amino acids constituting each pair of one or more first

amino acid pairs selected from among amino acid pairs forming respective amino acid-amino acid bonds between the first CH1 domain and the first CL domain, one is substituted with an amino acid having a positive charge and the other is substituted with an amino acid having a negative charge; and

5 a mutation in which, of the two amino acids constituting each pair of one or more second amino acid pairs selected from among amino acid pairs forming respective amino acid-amino acid bonds between the second CH1 domain and the second CL domain, one is substituted with an amino acid having a positive charge and the other is substituted with an amino acid having a negative charge.

10 The first epitope and the second epitope may exist in respective different proteins (antibodies) or in different (discriminative) regions of one protein (antigen).

In the bispecific antibody or the antigen-binding fragment thereof according to one embodiment, the amino acids substituted respectively in the first CH1 domain and the second CH1 domain have opposite charges, the amino acids substituted respectively in the first CL domain and the first CH1 domain have opposite charges, and the amino acids substituted
15 respectively in the second CL domain and the second CH1 domain have opposite charges.

In the bispecific antibody or the antigen-binding fragment thereof, for example,
the amino acid, positioned in the first CH1 domain, as a member of at least one first amino acid pair selected from among amino acid pairs forming respective amino acid-amino
20 acid bonds between the first CH1 domain and the first CL domain may be substituted with an amino acid having a positive charge while the other member positioned in the first CL domain may be substituted with an amino acid having a charge different from that of the amino acid substituted in the first CH1 domain, that is, a negative charge, and

the amino acid, positioned in the second CH1 domain, as a member of at least one
25 second amino acid pair selected from among amino acid pairs forming respective amino acid-amino acid bonds between the second CH1 domain and the second CL domain may be substituted with an amino acid having a charge different from that of the amino acid substituted in the first CH1 domain, that is, a negative charge while the other member positioned in the second CL domain may be substituted with an amino acid having a charge different from that of
30 the amino acid substituted in the second CH1 domain, that is, a positive charge.

In the bispecific antibody or the antigen-binding fragment thereof comprising mutant

CH1 CL domains, the first and the second amino acid pair to be substituted may be the same or different, the positively charged amino acids substituted in the first and the second amino acid pair may be the same or different, and the negatively charged amino acids substituted in the first and the second amino acid pair may be the same or different. The antigen-binding fragment of the bispecific antibody comprising a mutant CH1 domain and a mutant CL domain may be, for example, a F(ab')₂ fragment. The bispecific antibody comprising mutant CH1 and CL domains or the antigen-binding fragment thereof targets the same epitopes as those for a bispecific antibody comprising non-mutant (wild-type) CH1 and CL domains and exhibits higher heavy chain (or heavy chain variable region-CH1)-light chain dimerization rates and/or stability, compared to a bispecific antibody comprising non-mutant (wild-type) CH1 and CL domains.

The bispecific antibody comprising mutant CH1 and CL domains may comprise modified CH3 domains inclusive of a first CH3 domain derived from an antibody recognizing a first epitope and a second CH3 domain derived from an antibody recognizing a second epitope, or an Fc region comprising the modified CH3 domain, wherein the first CH3 domain and the second CH3 domain have at least one of the following mutations:

- (1) a mutation in which one member of at least one amino acid pair between the CH3 domains is substituted with an amino acid having a positive charge and the other member is substituted with an amino acid having a negative charge;
- (2) a mutation in which the members of at least one amino acid pair between the CH3 domains are switched with each other; and
- (3) a mutation in which one member of at least one amino acid pair between the CH3 domains is substituted with a large amino acid (e.g., large hydrophobic amino acids such as tryptophan, phenylalanine, etc.) and the other member is substituted with a small amino acid (e.g., small hydrophobic amino acids such as alanine, glycine, valine, etc.).

A bispecific protein or bispecific antibody comprising the mutant CH1 domain, the mutant CL domain, and the mutant Fc region or modified CH3 domain may exhibit an improvement in heterodimerization rate, dimerization rate between a heavy chain (or heavy chain variable region-CH1) and a light chain, both targeting the same epitope, and/or stability, compared to a bispecific protein or antibody comprising a CH1 domain, a CL domain, and an Fc region or CH3 domain none of which are mutant.

The bispecific antibody comprising mutant CH1 and CL domains or the antigen-binding

fragment thereof targets the same epitopes as those for a bispecific antibody comprising non-mutant (wild-type) CH1 and CL domains and exhibits higher heavy chain (or heavy chain variable region-CH1)-light chain dimerization rates and/or stability, compared to a bispecific antibody comprising non-mutant (wild-type) CH1 and CL domains.

5 Another embodiment provides a method for enhancing heterodimerization of a bispecific protein for targeting different targets, the bispecific protein comprising modified CH3 domains or an Fc region comprising the modified CH3 domains, said method comprising one of the following mutation introducing steps:

(1) substituting one of the two amino acids in at least one amino acid pair selected from
10 amino acid pairs forming amino-amino acid bonds between the CH3 domains with an amino acid having a positive charge and the other with an amino acid having a negative charge;

(2) switching the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between a first CH3 domain and a second CH3 domain with each other; and

15 (3) substituting one of the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain and the second CH3 domain with a large hydrophobic amino acid (for example, tryptophan, phenylalanine, etc.) and the other with a small hydrophobic amino acid (for example, alanine, glycine, valine, etc.).

20 Another embodiment provides a method for constructing a bispecific antibody or an antigen-binding fragment thereof or for enhancing a dimerization rate between a heavy chain (or heavy chain variable region-CH1) and a light chain, both targeting the same epitope, the method comprising the following CH1 and CL domain mutating steps of:

substituting one of the two amino acid residues constituting at least one selected from
25 amino acid pairs forming amino acid-amino acid bonds between a first CH3 domain derived from the heavy chain and a first CL domain of an antibody recognizing a first epitope with an amino acid having a positive charge and the other with an amino acid having a negative charge; and

substituting one of the two amino acid residues constituting at least one selected from
30 amino acid pairs forming amino acid-amino acid bonds between a second CH3 domain derived from the heavy chain and a second CL domain of an antibody recognizing a second epitope with

an amino acid having a positive charge and the other with an amino acid having a negative charge.

The method for constructing a bispecific antibody or an antigen-binding fragment thereof or for enhancing a dimerization rate between a heavy chain (or heavy chain variable region-CH1) and a light chain may comprise, in addition to the CH1 and CL domain mutating steps, at least one of the following CH3 domain mutation steps:

(1) substituting one of the two amino acids in at least one selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain derived from the heavy chain of an antibody recognizing a first epitope and the second CH3 domain derived from the heavy chain of an antibody recognizing a second epitope with an amino acid having a positive charge and the other with an amino acid having a negative charge;

(2) switching the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain and the second CH3 domain with each other; and

(3) substituting one of the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain and the second CH3 domain with a large hydrophobic amino acid (for example, tryptophan, phenylalanine, etc.) and the other with a small hydrophobic amino acid (for example, alanine, glycine, valine, etc.).

Another embodiment provides a method for constructing a bispecific antibody or an antigen-binding fragment thereof and for enhancing heterodimerization of a bispecific antibody or an antigen-binding fragment thereof for targeting different targets, the method comprising one of the following mutation introducing steps to introduce at least one mutation into at least one selected from amino acid pairs forming amino acid-amino acid bonds between a first CH3 domain and a second CH3 domain:

(1) substituting one of the two amino acids in at least one selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain derived from the heavy chain of an antibody recognizing a first epitope and the second CH3 domain derived from the heavy chain of an antibody recognizing a second epitope with an amino acid having a positive charge and the other with an amino acid having a negative charge;

(2) switching the two amino acids in at least one amino acid pair selected from amino

acid pairs forming amino-amino acid bonds between the first CH3 domain and the second CH3 domain with each other; and

(3) substituting one of the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain and the second CH3 domain with a large hydrophobic amino acid (for example, tryptophan, phenylalanine, etc.) and the other with a small hydrophobic amino acid (for example, alanine, glycine, valine, etc.).

The method for constructing a bispecific antibody or an antigen-binding fragment thereof may further comprise the following CH1 and CL domain mutating steps of:

substituting one of the two amino acid residues constituting at least one selected from amino acid pairs forming amino acid-amino acid bonds between a first CH3 domain derived from the heavy chain and a first CL domain of an antibody recognizing a first epitope with an amino acid having a positive charge and the other with an amino acid having a negative charge; and

substituting one of the two amino acid residues constituting at least one selected from amino acid pairs forming amino acid-amino acid bonds between a second CH3 domain derived from the heavy chain and a second CL domain of an antibody recognizing a second epitope with an amino acid having a positive charge and the other with an amino acid having a negative charge.

The method for constructing a bispecific antibody or an antigen-binding fragment thereof can enhance heterodimerization between CH3 domains or Fc regions derived from antibodies recognizing different epitope as well as between CH1 domains (heavy chains) and CL domains (light chains) derived from antibodies recognizing the same epitope.

Technical Solution

Below, a detailed description will be given of the present invention.

The present invention provides a bispecific protein comprising an Fc constant region and/or an Fab constant region and targeting different targets and a construction method therefor, wherein an amino acid mutation is introduced to the Fc constant regions (CH3 domains) linked (fused) respectively to targeting domains different from each other to increase coupling between the Fc constant regions linked to different targeting domains, thereby increasing a

heterodimerization rate between the Fc constant regions linked to different targeting domains and decreasing a homodimerization rate between the Fc constant regions linked to the same targeting domain; and/or

an amino acid mutation is introduced to Fab constant regions linked to targeting domains
5 different from each other to increase coupling between the Fab constant regions linked to different targeting domains, thereby a dimerization rate between the same targeting domains and between the Fab constant regions linked thereto,

whereby a bispecific protein having different targeting domains can be produced at high yield.

10 Amino acid positions in the antibodies (heavy and light chains), CH1 domains, CL domains, Fc regions, and CH3 domains disclosed in the description are all given as numbered according to the EU numbering system [the EU-index set forth in “Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)”] and differ from numbers accounting for positions of sequences in the
15 sequencing list.

The antibody may be at least one selected from among all kinds of immunoglobulins originating from mammals or birds. For example, the antibody used in the description may be at least one selected from the group consisting of IgG (e.g., IgG type 1 (IgG1), IgG type 2 (IgG2), IgG type 3 (IgG3), and IgG type 4 (IgG4)), IgA (e.g., IgA type 1 (IgA1) and IgA type 2 (IgA2)),
20 IgD, IgE, and IgM. The antibody may be an immunoglobulin derived from mammals such as primates comprising humans, monkeys, etc. and rodents comprising mice, rats, etc. and may be, for example, a human-derived immunoglobulin. In one embodiment, the antibody may be at least one selected from among human IgG1 (constant region; protein: GenBank Accession No. AAC82527.1, gene: GenBank Accession No. J00228.1), human IgG2 (constant region; protein:
25 GenBank Accession No. AAB59393.1, gene: GenBank Accession No. J00230.1), human IgG3 (constant region; protein: GenBank Accession No. P01860, gene: GenBank Accession No. X03604.1), human IgG4 (constant region; protein: GenBank Accession No. AAB59394.1, gene: GenBank Accession No. K01316.1), human IgA1 (constant region; protein: GenBank Accession No. AAT74070.1, gene: GenBank Accession No. AY647978.1), human IgA2
30 (constant region; protein: GenBank Accession No. AAB59396.1, gene: GenBank Accession No. J00221.1), human IgD (constant region; protein: GenBank Accession No. AAA52771.1,

AAA52770.1), human IgE (constant region; protein: GenBank Accession No. AAB59395.1, gene: GenBank Accession No. J00222.1), and human IgM (constant region; protein: GenBank Accession No. CAB37838.1, gene: GenBank Accession No. X57086.1). In one embodiment, the antibody may be at least one selected from the group consisting of human-derived IgG1, IgG2, IgG3, and IgG4, but is not limited thereto. The 1st CH3 domain and the 2nd CH3 domain, the 1st CH1 domain and the 1st CL domain, and the 2nd CH1 domain and the 2nd CL domain into all of which a mutation is introduced may each be independently selected from among identical or different immunoglobulin types.

As can be seen in FIG. 33a depicting sequence alignment results of the human IgG1 heavy chain constant region (SEQ ID NO: 33) and the human IgA1 heavy chain constant region (SEQ ID NO: 34) and in FIG. 33b depicting sequence alignment results of the kappa constant region (SEQ ID NO: 35) and lambda constant region (SEQ ID NO: 36) of the human immunoglobulin light chain, the heavy chain constant region and light chain constant region exhibit highly conserved amino acid sequences between subtypes.

In addition, immunoglobulin sequences are highly conserved among species and subtypes from which the sequences are derived. For instance, as shown by the sequence alignment results of heavy chain constant regions among human, mouse, and rat in FIG. 33c (CH1 domain sequence alignment) and FIG. 33d (CH3 domain sequence alignment), the amino acid sequences of heavy chain constant region of immunoglobulins are of high interspecies conservation.

In the description, hence, reference is given to the human IgG1 to designate amino acid positions in the CH1 domains and CH3 domains and to the human kappa constant region to designate amino acid positions in the CL domains. The amino acid positions designated with the human IgG1 and the human kappa constant region serving as references allows corresponding amino acid positions in immunoglobulins of other subtypes and immunoglobulins of species other than humans to be explicitly designated through a typical sequence alignment means (e.g., <https://blast.ncbi.nlm.nih.gov/Blast.cgi>) (see Table 1).

In addition, amino acid positions in the CH1 domain, CL domain, and CH3 domain, provided in the description, are represented according to the EU numbering system, and with respect to the details thereof, reference may be made to “http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGHGnber.html” (heavy chain

constant region)”, “http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGLCnber.html (light chain lambda region)” and “http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGKCnber.html (light chain kappa region)”.

5 Using the human IgG1 as a reference, the EU numbering system numbers:

(1) the CH1 domain (SEQ ID NO: 1) consecutively, with the first amino acid residue (Ala) given position 118 (i.e., 108 amino acid residues of the CH1 domain of SEQ ID NO: 1 correspond respectively to positions 118 to 215 in IgG1); and

(2) the CH3 domain (SEQ ID NO: 15) consecutively, with the first amino acid residue
10 (Lys) given position 340 (i.e., 108 amino acid residues of the CH3 domain of SEQ ID NO: 15 corresponds respectively to positions 340 to 447 in IgG1) (see, http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGHGnber.html).

In the description, amino acid positions in the CH1 and CH3 domains and amino acid kinds corresponding thereto are depicted, with the human IgG1 serving as a reference.

15 Further, according to the EU numbering,

the CL domain (SEQ ID NO: 10) of the human kappa constant region (protein: GenBank Accession No. AAA58989.1 gene: GenBank Accession No. J00241.1) is numbered consecutively, with the first amino acid residue (Val) given position 110 (i.e., 105 amino acid residues of the CL domain of SEQ ID NO: 10 correspond respectively to positions 110 to 214;
20 see http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGKCnber.html); and

the CL domains (SEQ ID NO: 11 (Lambda1), SEQ ID NO: 12 (Lambda2), SEQ ID NO: 13 (Lambda3), and SEQ ID NO: 14 (Lambda7)) of the human lambda constant region are numbered consecutively, with the first amino acid residue (Lys) given position 110 (for the lambda constant region, positions 169, 201, and 203 are omitted from the serial number
25 established; that is, the 103 amino acid residues of the CL domain of SEQ ID NO: 11 or 12 are numbered from position 110 to position 168, from position 170 to position 200, and from position 203 to position 215; see http://www.imgt.org/IMGTScientificChart/Numbering/Hu_IGLCnber.html). In the description, amino acid positions in the CL domains and amino acid kinds corresponding thereto are
30 depicted, with the human kappa constant region serving as a reference.

The Fab constant region may comprise one heavy chain constant region (i.e., CH1

domain) selected from the group consisting of heavy chain constant regions of Fab fragments of IgG (IgG1, IgG2, IgG3, and IgG 4), IgA (IgA1 and IgA2), IgD, IgE, and IgM and one light chain constant region (i.e. CL domain) selected from the group consisting of the kappa type and lambda types (e.g., lambda type 1, lambda type 2, lambda type 3, and lambda type 7) of immunoglobulin light chains.

By way of example, the CH1 domains of IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM, which are each available as a heavy chain constant region (CH1 domain) of the Fab fragment may comprise the amino acid sequences of SEQ ID NO: 1 (corresponding to positions 118 to 215 according to EU numbering), SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, and SEQ ID NO: 9, respectively. In addition, the CL domains of the kappa type, lambda type 1, lambda type 2, lambda type 3, and lambda type 7 are each available as the light chain constant region (CL domain) and may comprise the amino acid sequences of SEQ ID NO: 10 (corresponding to positions 110 to 214 according to EU numbering), SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, and SEQ ID NO: 14, respectively. In an embodiment, the Fab constant region the CH1 domain (SEQ ID NO: 1) of IgG type 1 and the light chain constant region (CL domain) (SEQ ID NO: 10) of the kappa type. In order to enhance dimerization between molecules targeting the same subject, the amino acid substituted with a negatively charged amino acid or a positively charged amino acid in the CH1 domain may be at least one residue, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) residues selected from the group consisting of leucine at position 145, lysine at position 147, phenylalanine at position 170, serine at position 183, and valine at position 185 in IgG type 1 (SEQ ID NO: 1) as numbered according to the EU numbering system. In another embodiment, the amino acid substituted with a negatively charged amino acid or a positively charged amino acid in the CH1 domain may be at least one residue selected from the group consisting of amino acids of other IgG subtypes (IgG2, IgG3, and IgG4), IgA1, IgA2, IgD, IgE, and IgM (respectively SEQ ID NOS: 2 to 9) at positions corresponding to leucine at position 145, lysine at position 147, phenylalanine at position 170, serine at position 183, and valine at position 185 on the amino acid sequence of SEQ ID NO: 1.

As used herein, “amino acids at positions corresponding to” can be determined by typical sequence alignment of the amino acid sequence of SEQ ID NO: 1 with target amino acid

sequences (i.e., SEQ ID NOS: 2 to 9) without difficulty (hereinafter, the same will be applied).

The amino acid having a positive charge (positively charged amino acid) may be selected from basic amino acids and may be, for example, lysine or arginine. When an amino acid having a positive charge is introduced into the CH1 domain, at least one residue, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) residues selected from the group consisting of leucine at position 145, lysine at position 147, phenylalanine at position 170, proline at position 171, serine at position 183, and valine at position 185 on the amino acid sequence of SEQ ID NO: 1, and amino acids at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 2 to 9 may each be independently substituted by a basic amino acid, for example, lysine or arginine.

For example, in order to introduce an amino acid having a positive charge thereinto, the CH1 domain may comprise at least one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) of the following mutations therein (on the amino acid sequence of SEQ ID NO: 1; and also applied to amino acid residues at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 2 to 9):

- substitution of leucine at position 145 with lysine or arginine (e.g., lysine);
- substitution of serine at position 183 with lysine or arginine (e.g., lysine);
- substitution of lysine at position 147 with arginine;
- substitution of phenylalanine at position 170 with lysine or arginine (e.g., lysine);
- substitution of proline at position 171 with lysine or arginine (e.g., lysine); and
- substitution of valine at position 185 with lysine or arginine (e.g., arginine).

The amino acid having a negative charge may be selected from among acidic amino acid residues and may be, for example, aspartic acid or glutamic acid. Hence, when an amino acid having a negative charge is introduced into the CH1 domain, at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) residues selected from the group consisting of leucine at position 145, lysine at position 147, phenylalanine at position 170, proline at position 171, serine at position 183, and valine at position 185 on the amino acid sequence of SEQ ID NO: 1, and amino acids at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 2 to 9 may each be independently substituted by an acidic amino acid, for example, aspartic acid or glutamic acid. For example, in order to introduce an amino acid having a positive charge thereinto, the CH1 domain may comprise at

least one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) of the following mutations (on the amino acid sequence of SEQ ID NO: 1; and also applied to amino acid residues at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 2 to 9):

- 5 substitution of leucine at position 145 with aspartic acid or glutamic acid (e.g., glutamic acid);
- substitution of lysine at position 147 with aspartic acid or glutamic acid (e.g., aspartic acid);
- substitution of serine at position 183 with aspartic acid or glutamic acid (e.g., glutamic
- 10 acid);
- substitution of valine at position 185 with aspartic acid or glutamic acid (e.g., aspartic acid);
- substitution of phenylalanine at position 170 with aspartic acid or glutamic acid (e.g., aspartic acid); and
- 15 substitution of proline at position 171 with aspartic acid or glutamic acid (e.g., aspartic acid).

In order to enhance dimerization between molecules targeting the same subject, the amino acid substituted with a negatively charged amino acid or a positively charged amino acid in the light chain constant region (CL domain) may be at least one residue, for example, one, two

20 or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) residues selected from the group consisting of serine at position 131, valine at position 133, leucine at position 135, serine at position 162, and threonine at position 180 in the kappa type (SEQ ID NO: 10). In another embodiment, the amino acid substituted with a negatively charged amino acid or a positively charged amino acid in the CL domain may be at least one residue selected from the

25 group consisting of amino acids of the CL domains of lambda types (lambda type 1, lambda type 2, lambda type 3, and lambda type 7) (respectively SEQ ID NOS: 11 to 14) at positions corresponding to serine at position 131, valine at position 133, leucine at position 135, serine at position 162, and threonine at position 180 on the amino acid sequence of SEQ ID NO: 10.

The amino acid having a positive charge (positively charged amino acid) may be

30 selected from basic amino acids and may be, for example, lysine or arginine. Hence, when an amino acid having a positive charge is introduced into the CL domain, at least one residue, for

example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) residues selected from the group consisting of serine at position 131, valine at position 133, leucine at position 135, serine at position 162, and threonine at position 180 on the amino acid sequence of SEQ ID NO: 10, and amino acids at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 11 to 14 may each be independently substituted by a basic amino acid, for example, lysine or arginine.

By way of example, in order to introduce an amino acid having a positive charge thereinto, the CL domain may comprise at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) of the following mutations (for the amino acid sequence of SEQ ID NO: 10; and also applied to the amino acids at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 11 to 14):

- substitution of serine at position 131 with lysine or arginine (e.g., lysine);
- substitution of valine at position 133 with lysine or arginine (e.g., lysine);
- substitution of leucine at position 135 with lysine or arginine (e.g., arginine);
- substitution of serine at position 162 with lysine or arginine (e.g., lysine); and
- substitution of threonine at position 180 with lysine or arginine (e.g., arginine).

In order to further enhance homodimerization, the mutant CH1 domain and/or the mutant CL domain may comprise two or more mutations simultaneously.

For example, lysine at position 147 and valine at position 185 in the CH1 domain may be substituted by an amino acid having a positive or negative charge. By way of example, lysine at position 147 and valine at position 185 in one of the first and the second CH1 domain may be substituted by an amino acid having a positive charge (e.g., lysine or arginine) while lysine at position 147 and valine at position 185 in the other CH1 domain may be substituted with an amino acid having a negative charge (glutamic acid or aspartic acid). In addition, leucine at position 135 and threonine at position 180 in the CL domain may be substituted with an amino acid having a positive or negative charge. By way of example, leucine at position 135 and threonine at position 180 in one of the first and the second CH1 domain may be substituted with an amino acid having a positive charge (e.g., lysine or arginine) while leucine at position 135 and threonine at position 180 in the other CL domain may be substituted with an amino acid having a negative charge (glutamic acid or aspartic acid).

In another embodiment, phenylalanine at position 170 and proline at position 171 in the

CH1 domain may be substituted with an amino acid having a positive or negative charge. By way of example, phenylalanine at position 170 and proline at position 171 in one of the first CH1 domain and the second CH1 domain may be substituted with an amino acid having a positive charge (e.g., lysine or arginine) while phenylalanine at position 170 and proline at position 171 in the other CH1 domain may be substituted with an amino acid having a negative charge (glutamic acid or aspartic acid). In addition, substitution with an amino acid having a positive or negative charge may be carried out for leucine at position 135 and serine at position 162 in the CL domain. By way of example, leucine at position 135 and serine at position 162 in one of the first CL domain and the second CL domain may be substituted with an amino acid having a positive charge (e.g., lysine or arginine) while leucine at position 135 and serine at position 162 in the other CL domain may be substituted with an amino acid having a negative charge (e.g., glutamic acid or aspartic acid).

The amino acid having a negative charge may be selected from among acidic amino acids and may be, for example, aspartic acid or glutamic acid. Hence, when an amino acid having a negative charge is introduced into the CL domain, at least one residue, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) residues selected from the group consisting of serine at position 131, valine at position 133, leucine at position 135, serine at position 162, and threonine at position 180 on the amino acid sequence of SEQ ID NO: 10, and amino acids at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 11 to 14 may each be independently substituted by a basic amino acid, for example, aspartic acid or glutamic acid. By way of example, in order to introduce an amino acid having a positive charge thereinto, the CL domain may comprise at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) of the following mutations (on the amino acid sequence of SEQ ID NO: 10; and also applied to amino acid residues at positions corresponding thereto on the amino acid sequences of SEQ ID NOS: 11 to 14):

substitution of serine at position 131 with aspartic acid or glutamic acid (e.g., glutamic acid);

substitution of valine at position 133 with aspartic acid or glutamic acid (e.g., glutamic acid);

substitution of leucine at position 135 with aspartic acid or glutamic acid (e.g., aspartic

acid);

substitution of serine at position 162 with aspartic acid or glutamic acid (e.g., aspartic acid); and

substitution of threonine at position 180 with aspartic acid or glutamic acid (e.g., aspartic acid).

In one embodiment, a set of two amino acids forming an amino acid pair between the CH1 domain and the CL domain that are to be substituted with a pair of amino acids having opposite charges may be at least one, for example, one, two or more (e.g. two), three or more (e.g., three), or four or more (e.g., four) pairs selected from the group consisting of a pair of leucine at position 145 in the CH1 domain and serine at position 131 in the CL domain, a pair of leucine at position 145 in the CH1 domain and valine at position 133 in the CL domain, a pair of lysine at position 147 in the CH1 domain and threonine at position 180 in the CL domain, a pair of serine at position 183 in the CH1 domain and valine at position 133 in the CL domain, a pair of valine at position 185 in the CH1 domain and leucine at position 135 in the CL domain, a pair of phenylalanine at position 170 in the CH1 domain and leucine at position 135 in the CL domain, and a pair of proline at position 171 in the CH1 domain and serine at position 162 in the CL domain, as numbered on the basis of the amino acid sequences of SEQ ID NO: 1 (CH1 domain) and SEQ ID NO: 10 (CL domain). For example, the amino acid pair between the CH1 and the CL domain to which the mutation is introduced may be one or two or more (e.g., two) pairs selected from the group consisting of a pair of leucine at position 145 in the CH1 domain and serine at position 131 in the CL domain, a pair of leucine at position 145 in the CH1 domain and valine at position 133 in the CL domain, a pair of lysine at position 147 in the CH1 domain and threonine at position 180 in the CL domain, a pair of serine at position 183 in the CH1 domain and valine at position 133 in the CL domain, and a pair of valine at position 185 in the CH1 domain and leucine at position 135 in the CL domain, as numbered on the basis of the amino acid sequences of SEQ ID NO: 1 (CH1 domain) and SEQ ID NO: 10 (CL domain).

The amino acid pairs to which the mutations are introduced between the first CH1 domain and the first CL domain may be the same as or different from those between the second CH1 domain and the second CL domain.

In one embodiment, an amino acid having a positive charge is introduced to the first CH1 domain (with the introduction of an amino acid having a negative charge to the first CL

domain) while an amino acid having a negative charge is introduced to the second CH1 domain (with the introduction of an amino acid having a positive charge to the second CL domain).

In order to enhance heterodimerization, at least one, for example, one, two, or three of the following mutations may be introduced to the Fc region of a heavy chain, in detail, the CH3 domain in the Fc region:

(1) mutation in which one amino acid residue of at least one amino acid pair (at least one of the two amino acid residues is not hydrophobic) between CH3 domains is substituted with an amino acid having a positive charge while the other residue is substituted with an amino acid having a negative charge (hereinafter referred to as “electrostatic interaction-induced mutation”);

(2) mutation in which amino acid residues in at least one amino acid pair between the CH3 domains are exchanged with each other (hereinafter referred to as “swapping mutation”); and

(3) mutation in which one amino acid residue in at least one amino acid pair between the CH3 domains is substituted with a large amino acid (e.g., a large hydrophobic amino acid such as tryptophan, phenylalanine, etc.) while the other amino acid residue is substituted with a small amino acid (e.g., a small hydrophobic amino acid such as alanine, glycine, valine, etc.) (hereinafter referred to as “size mutation”).

As such, the CH3 domain to which the mutations are introduced may be selected from the group consisting of the CH3 domain of human IgG1 (SEQ ID NO: 15; corresponding to positions 340 to 447 according to the EU numbering), the CH3 domain of human IgG2 (SEQ ID NO: 16), the CH3 domain of human IgG3 (SEQ ID NO: 17), the CH3 domain of human IgG4 (SEQ ID NO: 18), the CH3 domain of human IgA1 (SEQ ID NO: 19), the CH3 domain of human IgA2 (SEQ ID NO: 20), the CH3 domain of human IgD (SEQ ID NO: 21), the CH3 domain of human IgE (SEQ ID NO: 22), and the CH3 domain of human IgM (SEQ ID NO: 23).

The first CH3 domain and the second CH3 domain to which the mutations are introduced may be derived from the same or different immunoglobulin types which may each be independently selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM. In one embodiment, the first CH3 domain and the second CH3 domain may both be the CH3 domain of human IgG1 having the amino acid sequence of SEQ ID NO: 15, but is not limited thereto.

The following amino acid pairs between the first CH3 domain and the second CH3 domain are based on the CH3 domain of human IgG1 having the amino acid of SEQ ID NO: 15, and the basis is true of the amino acid pairs corresponding thereto between CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM (respectively, SEQ ID NOS: 16 to 23).

5 The amino acid pair between the first and second CH3 domains to which one of the mutations is introduced may be at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) amino acid pairs selected from the group consisting of the amino acid pairs between CH3 domains and amino acid pairs at positions corresponding thereto between CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM (respectively,
10 SEQ ID NOS: 16 to 23), listed in Table 1, below.

TABLE 1

Amino acid pair No.	IgG (1-4)		IgA1		IgA2		IgE		IgM	
	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B
1	Q347	K360	E347	L360	E347	L360	E444	R457	Q347	K360
2	Y349	S354	H349	P354	H349	P354	Y446	P451	Y349	S354
3	Y349	E357	H349	E357	H349	E357	Y446	P454	Y349	E357
4	Y349	K360	H349	L360	H349	L360	Y446	R457	Y349	K360
5	L351	L351	L351	L351	L351	L351	F448	F448	L351	L351
6	P352	P352	P352	P352	P352	P352	A449	A449	P352	P352
7	S354	Y349	P354	H349	P354	H349	P451	Y446	S354	Y349
8	D356	K439	E356	K439	E356	K439	W453	R539	D356	K439
9	E357	Y349	E357	H349	E357	H349	P454	Y446	E357	Y349
10	E357	K370	E357	R370	E357	R370	P454	Q467	E357	K370
11	K360	Q347	L360	E347	L360	E347	R457	E444	K360	Q347
12	K360	Y349	L360	H349	L360	H349	R457	Y446	K360	Y349
13	S364	L368	T364	L368	T364	L368	T461	L465	S364	L368
14	S364	K370	T364	R370	T364	R370	T461	Q467	S364	K370
15	T366	T366	T366	T366	T366	T366	A463	A463	T366	T366
16	T366	Y407	T366	T407	T366	T407	A463	F506	T366	Y407
17	L368	S364	L368	T364	L368	T364	L465	T461	L368	S364
18	L368	K409	L368	I409	L368	I409	L465	R508	L368	K409
19	K370	E357	R370	E357	R370	E357	Q467	P454	K370	E357
20	K370	S364	R370	T364	R370	T364	Q467	T461	K370	S364
21	K370	T411	R370	R411	R370	R411	Q467	E510	K370	T411
22	N390	S400	L390	S400	L390	S400	R489	K499	N390	S400
23	K392	L398	L392	Q398	L392	Q398	S491	K497	K392	L398
24	T394	T394	W394	W394	W394	W394	T493	T493	T394	T394
25	T394	V397	W394	R397	W394	R397	T493	R496	T394	V397
26	P395	P395	A395	A395	A395	A395	Q494	Q494	P395	P395
27	P395	V397	A395	R397	A395	R397	Q494	R496	P395	V397
28	V397	T394	R397	W394	R397	W394	R496	T493	V397	T394

29	V397	P395	R397	A395	R397	A395	R496	Q494	V397	P395
30	L398	K392	Q398	L392	Q398	L392	K497	S491	L398	K392
31	S400	N390	S400	K390	S400	K390	K499	R489	S400	N390
32	F405	K409	A405	I409	A405	I409	F504	R508	F405	K409
33	Y407	T366	T407	T366	T407	T366	F506	A463	Y407	T366
34	Y407	Y407	T407	T407	T407	T407	F506	F506	Y407	Y407
35	Y407	K409	T407	I409	T407	I409	F506	R508	Y407	K409
36	K409	L368	I409	L368	I409	L368	R508	L465	K409	L368
37	K409	F405	I409	A405	I409	A405	R508	F504	K409	F405
38	K409	Y407	I409	T407	I409	T407	R508	F506	K409	Y407
39	T411	K370	R411	R370	R411	R370	E510	Q467	T411	K370
40	K439	D356	K439	K439	K439	K439	R539	W453	K439	D356

(Chain A: first CH3 domain; Chain B: second CH3 domain)

In greater detail, the amino acid pair between the first CH3 domain and the second CH3 domain to which at least one of the following mutations: (1) electrostatic interaction-induced mutation (represented as charge (J) in Table 2 and FIG. 2); (2) swapping mutation (represented as swap (O) in Table 2 and FIG. 2); and (3) size mutation (represented as size (B) in Table 2 and FIG. 2) may be at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) amino acid pairs selected from the amino acid pairs between CH3 domains of IgG1 (SEQ ID NO: 15) and amino acid pairs at positions corresponding thereto between CH3 domains of IgG2, IgG3 IgG4 IgA1 IgA2 IgD, IgE, and IgM (respectively, SEQ ID NOS: 16 to 23), suggested in Table 2 and FIG. 2:

TABLE 2

Amino acid pair No.	Charge (J)		Swap (O)		Size (B)	
	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B
1	Q347	K360	Q347	K360	Q347	K360
2	Y349	S354			Y349	S354
3	Y349	E357			Y349	E357
4	Y349	K360			Y349	K360
5	L351	L351			L351	L351
6	P352	P352			P352	P352
7	S354	Y349	S354	Y349	S354	Y349
8	D356	K439			D356	K439
9	E357	Y349	E357	Y349	E357	Y349
10	E357	K370	E357	K370	E357	K370
11	K360	Q347			K360	Q347
12	K360	Y349	K360	Y349	K360	Y349
13	S364	L368	S364	L368	S364	L368
14	S364	K370	S364	K370	S364	K370
15	T366	T366			T366	T366
16	T366	Y407			T366	Y407

17	L368	S364			L368	S364
18	L368	K409	L368	K409	L368	K409
19	K370	E357			K370	E357
20	K370	S364			K370	S364
21	K370	T411			K370	T411
22	N390	S400	N390	S400	N390	S400
23	K392	L398			K392	L398
24	T394	T394			T394	T394
25	T394	V397	T394	V397	T394	V397
26	P395	P395			P395	P395
27	P395	V397			P395	V397
28	V397	T394			V397	T394
29	V397	P395			V397	P395
30	L398	K392	L398	K392	L398	K392
31	S400	N390			S400	N390
32	F405	K409	F405	K409	F405	K409
33	Y407	T366	Y407	T366	Y407	T366
34	Y407	Y407			Y407	Y407
35	Y407	K409			Y407	K409
36	K409	L368			K409	L368
37	K409	F405			K409	F405
38	K409	Y407			K409	Y407
39	T411	K370	T411	K370	T411	K370
40					K439	D356

(Chain A: first CH3 domain; Chain B: second CH3 domain; the mutations listed in Table 2 are given to positions based on IgG1, but are applied to corresponding positions in the CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM).

As used herein, for example, “Q347” means glutamine at position 347 on the amino acid sequence of SEQ ID NO: 15 in the CH3 domain of human IgG1, and such notation is true of amino acid residues at positions corresponding thereto in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM (respectively, SEQ ID NOS: 16 to 23) (hereinafter, the same definition is applied).

Hereinafter, amino acid pairs between CH3 domains to which the mutations are introduced are numbered on the basis of the amino acid sequence of SEQ ID NO: 15 and unless otherwise described, the numbering is construed to be applied to amino acids at denoted positions in IgG1 as well as at positions corresponding thereto in CH3 domains of other type immunoglobulins (IgG2, IgG3, IgG4, IgA1, IgA2, IgD, IgE, and IgM).

The electrostatic interaction-induced mutation is intended to substitute a positively charged amino acid for one amino acid residue of at least one amino acid pair between Fc

regions or CH3 domains (at least one residue of the paired two amino acids is not hydrophobic) and a negatively charged amino acid for the other amino acid residue to introduce electrostatic interaction to a hydrophobic interaction-lacking site, thereby contributing to an increase of electrostatic interaction-induced binding force.

5 The hydrophobic amino acid may be selected from the group consisting of glycine, alanine, valine, leucine, isoleucine, methionine, proline, phenylalanine, and tryptophan.

 The amino acid having a negative charge may be selected from among acidic amino acids and may be, for example, aspartic acid or glutamic acid. The amino acid having a positive charge may be selected from among basic amino acid and may be, for example, lysine or
10 arginine.

 The amino acid pair between the first CH3 domain and the second CH3 domain to which an electrostatic interaction-introduced mutation is applicable may be at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) selected from among amino acid pair numbers 1 to 39 in Table 2 and may be at least one, for
15 example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) selected from the group consisting of, for example, a pair of serine at position 364 and leucine at position 368, a pair of threonine at position 394 and threonine at position 394, a pair of glutamic acid at position 357 and lysine at position 370, a pair of glutamic acid at position 357 and tyrosine at position 349, a pair of threonine at position 366 and tyrosine at position 407, and a
20 pair of threonine at position 394 and valine at position 397.

 That is, the electrostatic interaction-introduced mutation in the CH3 domain may comprise substitution of an amino acid having a positive charge for one amino acid residue of each of the amino acid pairs, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) selected from among the amino acid pair number 1 to 39 in
25 Table 2; and an amino acid having a negative charge for the other amino acid residue. By way of example, in each of the amino acid pairs, for example, one or more (e.g., one), two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) amino acid pairs selected from the group consisting of a pair of serine at position 364 and leucine at position 368, a pair of threonine at position 394 and threonine at position 394, a pair of glutamic acid at position 357
30 and lysine at position 370, a pair of glutamic acid at position 357 and tyrosine at position 349, a pair of threonine at position 366 and tyrosine at position 407, and a pair of threonine at position

394 and valine at position 397, one amino acid residue is substituted with an amino acid having a positive charge while the other amino acid residue is substituted with an amino acid having a negative charge.

For instance, the electrostatic interaction-introduced mutation in CH3 domains may
5 comprise at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) of the following mutations:

substitution of serine at position 364 with an amino acid having a positive charge and leucine at position 368 with an amino acid having a negative charge;

substitution of threonine at position 394 with an amino acid having a positive charge and
10 threonine at position 394 with an amino acid having a negative charge;

substitution of glutamic acid at position 357 with an amino acid having a positive charge and lysine at position 370 with an amino acid having a negative charge;

substitution of glutamic acid at position 357 with an amino acid having a positive charge and tyrosine at position 349 with an amino acid having a negative charge;

15 substitution of threonine at position 366 with an amino acid having a positive charge and tyrosine at position 407 with an amino acid having a negative charge;

substitution of threonine at position 394 with an amino acid having a positive charge and valine at position 397 with an amino acid having a negative charge; and

substitution of tyrosine at position 349 with an amino acid having a positive charge and
20 glutamic acid at position 357 with an amino acid having a negative charge.

Such an electrostatic interaction introduction in CH3 domains may achieve a heterodimerization rate of 60% or higher, 65% or higher, 70% or higher, 73% or higher, 75% or higher, 78% or higher, 80% or higher, 85% or higher, 90% or higher, 95% or higher, or 100%.

The swapping mutation means a mutation in which two amino acid residues constituting
25 an amino acid pair are exchanged (swapped) with each other.

As illustrated in Table 2 and FIG. 2, the amino acid pair between the first CH3 domain and the second CH3 domain to which the swapping mutation is applicable may be at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) selected from the group consisting of a pair of glutamine at position 354 and lysine at position
30 360, a pair of serine at position 357 and tyrosine at position 349, a pair of glutamic acid at position 357 and lysine at position 370, a pair of lysine at position 360 and tyrosine at position

349, a pair of serine at position 364 and leucine at position 368, a pair of serine at position 364 and lysine at position 370, a pair of leucine at position 368 and lysine at position 409, a pair of asparagine at position 390 and serine at position 400, a pair of threonine at position 394 and valine at position 397, a pair of leucine at position 398 and lysine at position 392, a pair of phenylalanine at position 405 and lysine at position 409, a pair of tyrosine at position 407 and threonine at position 366, and a pair of threonine at position 411 and lysine at position 370 and particularly from the group consisting of a pair of serine at position 364 and lysine at position 370, a pair of tyrosine at position 407 and threonine at position 366, a pair of glutamic acid at position 357 and lysine at position 370, a pair of phenylalanine at position 405 and lysine at position 409, and a pair of serine at position 357 and tyrosine at position 349.

That is, the swapping mutation in CH3 domains may comprise substitution in which exchange (swapping) is made between two paired amino acid residues in each of at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) amino acid pairs selected from the group consisting of a pair of glutamine at position 354 and lysine at position 360, a pair of serine at position 357 and tyrosine at position 349, a pair of glutamic acid at position 357 and lysine at position 370, a pair of lysine at position 360 and tyrosine at position 349, a pair of serine at position 364 and leucine at position 368, a pair of serine at position 364 and lysine at position 370, a pair of leucine at position 368 and lysine at position 409, a pair of asparagine at position 390 and serine at position 400, a pair of threonine at position 394 and valine at position 397, a pair of leucine at position 398 and lysine at position 392, a pair of phenylalanine at position 405 and lysine at position 409, a pair of tyrosine at position 407 and threonine at position 366, and a pair of threonine at position 411 and lysine at position 370; and for example, from the group consisting of a pair of serine at position 364 and lysine at position 370, a pair of tyrosine at position 407 and threonine at position 366, a pair of glutamic acid at position 357 and lysine at position 370, a pair of phenylalanine at position 405 and lysine at position 409, and a pair of serine at position 357 and tyrosine at position 349.

For example, the swapping mutation in CH3 domains may comprise at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) of the following mutations:

substitution of serine at position 364 with lysine and lysine at position 370 with serine;
substitution of phenylalanine at position 405 with lysine and lysine at position 409 with

phenylalanine;

substitution of tyrosine at position 407 with threonine and threonine at position 366 with tyrosine;

substitution of glutamic acid at position 357 with lysine and lysine at position 370 with
5 glutamic acid; and

substitution of serine at position 357 with tyrosine and tyrosine at position 349 with serine.

Such a swapping mutation in CH3 domains may achieve a heterodimerization rate of 60% or higher, 65% or higher, 70% or higher, 73% or higher, 75% or higher, 78% or higher,
10 80% or higher, 85% or higher, 90% or higher, 95% or higher, or 100%.

The size mutation means a mutation in which, of the two paired amino acids in at least one amino acid pair between CH3 domains, one residue is substituted with a large hydrophobic amino acid (e.g., tryptophan, phenylalanine, etc.) and the other is substituted with a small hydrophobic amino acid (e.g., alanine, glycine, valine, etc.) so that the large amino acid is fitted
15 to the space secured by the small amino acid, thereby contributing to heterodimerization.

The large amino acid may comprise a cyclic residue and may be selected from the group consisting of tryptophan and phenylalanine, and particularly tryptophan. The small amino acid may be selected from the group consisting of alanine, glycine, and valine and may be, for example, alanine.

20 The amino acid pair between the first CH3 domain and the second CH3 domain to which the size mutation is applicable may be at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) selected from among amino acid pair numbers 1 to 40 in Table 2 and may be, for example, a pair of lysine at position 409 and tyrosine at position 407, a pair of lysine at position 409 and phenylalanine at position 405, or a
25 combination thereof.

That is, the size mutation may comprise substitution in which, of two paired amino acid residues in each of at least one, for example, one, two or more (e.g., two), three or more (e.g., three), or four or more (e.g., four) amino acid pairs selected from among amino acid pair numbers 1 to 40 in Table 2, one is substituted with a large hydrophobic amino acid (e.g.,
30 tryptophan, phenylalanine, etc.), for example, tryptophan while the other is substituted with a small hydrophobic amino acid (e.g., alanine, glycine, valine, etc.), for example, alanine. By way

of example, in a pair of lysine at position 409 and tyrosine at position 407 or a pair of lysine at position 409 and phenylalanine at position 405, or in each of them, one residue may be substituted with a large hydrophobic amino acid, for example, phenylalanine or tryptophan while the other residue may be substituted with a small hydrophobic amino acid, for example, alanine, glycine, or valine.

For example, the size mutation in CH3 domains may comprise at least one of the following mutations:

substitution of lysine at position 409 with tryptophan and tyrosine at position 407 with alanine; and

substitution of lysine at position 409 with tryptophan and phenylalanine at position 405 with alanine.

The modified CH3 domains may comprise at least one, for example, one or two of the three mutations described above, that is, electrostatic interaction-introduced mutation, swapping mutation, and size mutation.

In order to exert the most advantageous effect on dimerization, the modified CH3 domains may comprise at least one, for example, one, two, or three mutations selected from the group consisting of substitution of one residue in a pair of serine at position 364 and leucine at position 368 with an amino acid having a positive charge and the other with an amino acid having a negative charge (electrostatic interaction-introduced mutation), exchange of the two residues of a pair of serine at position 364 and lysine at position 370 with each other (swapping mutation), and exchange of the two residues of a pair of phenylalanine at position 405 and lysine at position 409 with each other (swapping mutation).

For example, the modified CH3 domains may comprise one, two, or three of the following mutations:

(a) substitution of serine at position 364 with an amino acid having a positive charge and leucine at position 368 with an amino acid having a negative charge;

(b) substitution of serine at position 364 with lysine and lysine at position 370 with serine; and

(c) substitution of phenylalanine at position 405 with lysine and lysine at position 409 with phenylalanine.

In order to decrease monomerization rates, but increase dimerization rates, an additional

amino acid modification may be introduced after the three selected mutations ((a)-(c)). As such, the amino acid to which the additional amino acid mutation can be introduced may be serine at position 364, phenylalanine at position 405, and/or lysine at position 409. For example, in a pair of serine at position 364 and leucine at position 368 which is to undergo an electrostatic
5 interaction-induced mutation, when leucine at position 368 may be substituted with an amino acid having a negative charge (aspartic acid or glutamic acid, e.g., aspartic acid), serine at position 364 may be substituted with an amino acid having a positive charge (lysine or arginine) (S364K or S364R; electrostatic interaction-introduced mutation) or with asparagine (S364N). In addition, in a pair of lysine at position 370 and serine at position 364, which is to undergo a
10 swapping mutation, lysine at position 370 may be substituted with serine and serine at position 364 may be substituted with lysine (S364K; swapping mutation) or with arginine or asparagine (S364R or S364N).

Further, in a pair of lysine at position 409 and phenylalanine at position 405, which is to undergo swapping mutation, lysine at position 409 may be substituted with phenylalanine (for
15 swapping mutation) or with tryptophan and phenylalanine at position 405 may be substituted with lysine (F405K; for swapping mutation) or with arginine, glutamine, or asparagine (F405R, F405Q, or F405N).

In one embodiment, the modified CH3 domains may comprise at least one of the following mutations:

- 20 substitution of serine at position 364 with lysine and leucine at position 368 with aspartic acid;
- substitution of serine at position 364 with lysine and lysine at position 370 with serine;
- substitution of phenylalanine at position 405 with lysine and lysine at position 409 with phenylalanine;
- 25 substitution of phenylalanine at position 405 with arginine and lysine at position 409 with phenylalanine;
- substitution of phenylalanine at position 405 with lysine and lysine at position 409 with tryptophan; and
- substitution of phenylalanine at position 405 with arginine and lysine at position 409
30 with tryptophan.

In one embodiment, the modified CH3 domains may comprise at least one of the

following double mutations:

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second
5 CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second
CH3 domain with arginine;

10 substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at
15 position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position
20 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second
25 CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with lysine; or

30 substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in

the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with arginine.

Such a double mutation in the CH3 domain may result in a dimerization rate of 70% or higher, 75% or higher, 80% or higher, 85% or higher, 90% or higher, 92% or higher, 93% or higher, 94% or higher, 95% or higher, or 96% or higher.

Another aspect of the present invention provides an anti-influenza B antibody comprising a heavy chain variable region composed of the amino acid sequence of SEQ ID NO: 27 and a light chain variable region composed of the amino acid sequence of SEQ ID NO: 31, or an antigen-binding fragment thereof. Another aspect of the present invention provides an anti-influenza A antibody comprising a heavy chain variable region composed of the amino acid sequence of SEQ ID NO: 29 and a light chain variable region composed of the amino acid sequence of SEQ ID NO: 31, or an antigen-binding fragment thereof.

Another aspect of the present invention provides an anti-influenza A/anti-influenza B bispecific antibody comprising an anti-influenza B antibody comprising a heavy chain variable region composed of the amino acid sequence of SEQ ID NO: 27 and a light chain variable region composed of the amino acid sequence of SEQ ID NO: 31, and an anti-influenza A antibody comprising a heavy chain variable region composed of the amino acid sequence of SEQ ID NO: 29 and a light chain variable region composed of the amino acid sequence of SEQ ID NO: 31, or an antigen-binding fragment thereof. The anti-influenza A/anti-influenza B bispecific antibody may comprise (1) the modified CH3 domain (as mentioned above, a mutation pair of CH3-CH3 introduced); (2) the mutant CH1 domain and the mutant CL domain (as mentioned above, a mutation pair of CH1-CL introduced); or (3) both the modified CH3 domain, and the mutant CH1 domain and the mutant CL domain.

The bispecific protein or bispecific antibody of the present invention is a bispecific matter constructed according to the **Correlated and Harmonious Interfacial Mutation** between **Protein Subunits** (hereinafter referred to as “Chimps”).

As used herein, the term “antibody” refers to a class of structurally related glycoproteins consisting of two pairs of polypeptide chains, one pair of light (L) low molecular weight chains and one pair of heavy (H) chains, all four inter-connected by disulfide bonds. The structure of antibodies has been well characterized (see, for instance, [Fundamental Immunology Ch. 7 (Paul, W., 2nd ed. Raven Press, N. Y. (1989))]. In brief, each heavy chain typically is comprised

of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region typically is comprised of three domains, CH1, CH2, and CH3. The heavy chains are inter-connected via disulfide bonds in the so-called “hinge region”. Each light chain typically is comprised of a light chain variable region (abbreviated herein as VL) and
5 a light chain constant region. The light chain constant region typically is comprised of one domain, CL. Typically, the numbering of amino acid residues in the constant region is performed according to the EU-index as described in the document [Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. (1991)]. The VH and VL regions may be further subdivided into regions of
10 hypervariability (or hypervariable regions which may be hypervariable in sequence and/or form of structurally defined loops), also termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FRs). Each VH and VL is typically composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4 (see also
15 document [Chothia and Lesk J. Mol. Biol. 196, 901 917 (1987)]).

As used herein, the term “Fab-arm” refers to one heavy chain-light chain pair.

The term “Fc region”, as used herein, refers to an antibody region comprising a CH2 domain and a CH3 domain and may further comprise the hinge region, optionally.

As used herein, the term “bispecific antibody” refers to an antibody having specificities
20 for at least two different epitope, typically non-overlapping epitopes.

The term “full-length antibody”, as used herein, refers to an antibody which contains all heavy and light chain constant and variable domains corresponding to those that are normally found in an antibody of that isotype. In an embodiment, a full-length antibody comprises two full-length heavy chains and two full-length light chains. As used herein, “isotype” refers to the
25 antibody class (for instance IgG1, IgG2, IgG3, IgG4, IgD, IgA, IgE, or IgM) that is encoded by heavy chain constant region genes.

The term “antigen-binding fragment” is intended to mean a portion of an antibody containing variable domains of the antibody and may be selected from among Fab, F(ab')₂, scFv, (scFv)₂, scFv-Fc, (scFv-Fc), etc., but is not limited thereto.

30 The term “epitope” means a protein determinant capable of specifically binding to an antibody. Epitopes usually consist of surface groupings of molecules such as amino acids or

sugar side chains and usually have specific three dimensional structural characteristics as well as specific charge characteristics.

The term “bispecific antibody” is generic to the antibody that recognizes and/or binds to two different antigens or two different (non-overlapping) epitopes on one antigen. In an embodiment, the bispecific antibody may comprise one antigen-binding site directed against a tumor cell antigen and another antigen-binding site directed against a cytotoxic trigger molecule. By way of example, the bispecific antibody may be selected from the group consisting of an anti-FcγRI/anti-CD15 antibody, an anti-p185HER2/anti-FcγRIII (CD16) antibody, an anti-CD3/anti-malignant B-cell (1D10) antibody, an anti-CD 3/anti-p185HER2 antibody, an anti-CD3/anti-p97 antibody, an anti-CD3/anti-renal cell carcinoma antibody, an anti-CD3/anti-OVCAR-3 antibody, an anti-CD3/anti-L-D1 (anti-colon cancer) antibody, an anti-CD3/anti-melanocyte stimulating hormone analog antibody, an anti-EGF receptor/anti-CD3 antibody, an anti-CD3/anti-CAMA1 antibody, an anti-CD3/anti-CD19 antibody, an anti-CD3/MoV18 antibody, an anti-neural cell adhesion molecule (NCAM)/anti-CD3 antibody, an anti-folate binding protein (FBP)/anti-CD3 antibody, an anti-pan carcinoma associated antigen (AMOC-31)/anti-CD3 antibody, but are not limited thereto. In another embodiment, bispecific antibodies with one antigen-binding site which binds specifically to a tumor antigen and another antigen-binding site which binds to a toxin may comprise, but is not limited to, an anti-saporin/anti-Id-1 antibody, an anti-CD22/anti-saporin antibody, an anti-CD7/anti-saporin antibody, an anti-CD38/anti-saporin antibody, an anti-CEA/anti-ricin A chain antibody, an anti-interferon-α (IFN-α)/anti-hybridoma idiotype antibody, and an anti-CEA/anti-vinca alkaloid antibody. In another embodiment, the bispecific antibody may be selected from among antibodies for converting enzyme activated prodrugs such as anti-CD30/anti-alkaline phosphatase (which catalyzes conversion of mitomycin phosphate prodrug to mitomycin alcohol), but is not limited thereto. In another embodiment, the bispecific antibody may be selected from among those that can be used as fibrinolytic agents such as anti-fibrin/anti-tissue plasminogen activator (tPA), anti-fibrin/anti-uropkinase-type plasminogen activator (uPA), etc., but is not limited thereto. In another embodiment, the bispecific antibody may be selected from among those for targeting immune complexes to cell surface receptors such as anti-low density lipoprotein (LDL)/anti-Fc receptor (e.g. FcγRI, FcγRII or FcγRIII), but is not limited thereto. In another embodiment, the bispecific antibody may be selected from those for use in therapy of infectious diseases (e.g., viral infection

diseases) such as anti-influenza A/anti-influenza B, anti-CD3/anti-herpes simplex virus (HSV), anti-T-cell receptor:CD3 complex/anti-influenza, anti-Fc γ R/anti-HIV, etc., but is not limited thereto. In another embodiment, the bispecific antibody may be selected from those for tumor detection in vitro or in vivo such as anti-CEA/anti-EOTUBE, anti-CEA/anti-DPTA, anti-p185

5 HER2/anti-hapten, etc., but is not limited thereto. In another embodiment, the bispecific antibody may be selected from those for use as diagnostic tools such as anti-rabbit IgG/anti-ferritin, anti-horse radish peroxidase (HRP)/anti-hormone, anti-somatostatin/anti-substance P, anti-HRP/anti-FITC, anti-CEA/anti- β -galactosidase, etc. In another embodiment, examples of the bispecific antibody comprise, but are not limited to, an antibody inclusive of a first antigen-binding site directed against CD30 and an antigen-binding site directed against erbB2; an

10 antibody inclusive of a first antigen-binding site directed against CD30 and a second antigen-binding site directed against Pseudomonas exotoxin (PE); an antibody inclusive of a first antigen-binding site directed against CD30 and a second antigen-binding site directed against streptavidin.

15 According to another embodiment, the targeting domain of the bispecific protein may comprise at least one target specific binding polypeptide selected from the group consisting of various membrane proteins, for example, various receptors (e.g., receptor tyrosine kinase (RTKs), etc.), ectodomains (extracellular domains) of the receptors, and various ligands (e.g., various growth factors, cytokines, etc.). Examples of the receptor comprise, but are not limited

20 to, tumor necrosis factor receptor (TNFR) (e.g., TNFR1, TNFR2, etc.), epidermal growth factor receptor (EGFR) (e.g., Her1(epidermal growth factor receptor 1), Her2 (human epidermal growth factor receptor 2), Her3 (human epidermal growth factor receptor 3), etc.), angiopoietin receptor (e.g., Tie1, Tie2, etc.), transforming growth factor receptor (e.g. TGFbR1, TGFbR2, TGFbR3, TGFaR1, etc.), bone morphogenetic protein receptor (e.g. BMPR1b), interleukin

25 receptor (e.g., interleukin 12 receptor subunit beta 1 (IL-12R-b1)), IL-4Ra, IL-12A, IL-4, IL-1R1L, IL-17RA, IL-17A, IL-12R-b2, IL-13Ra1, IL-12B, IL-13, IL-1RAP, IL-17RC, IL-17F, etc.), integrin (e.g., integrin alpha 4 (ITGA4), integrin subunit alpha 2b (ITGA2B), ITGB1, ITGB3, etc.), interferon receptor (e.g., interferon-alpha/beta receptor 1 (IFNAR1), IFNAR2, IFNGR, etc.), Fas (tumor necrosis factor receptor superfamily member 6; TNFRSF6), VEGF

30 receptor (e.g., Flt1 (fms related tyrosine kinase 1), etc.), hepatocyte growth factor receptor (e.g., Met, etc.), and Interferon gamma receptor (IFNGR). The ligand may be at least one selected

from the group consisting of tumor necrosis factor (TNF), epidermal growth factor (EGF), vascular endothelial cell growth factor (VEGF-A, VEGF-B, VEGF-C, VEGF-D, etc.), angiopoietin (e.g., Ang1, Ang2, etc.), transforming growth factor (TGF), hepatocyte growth factor (HGF), bone morphogenetic protein (e.g., BMP2, BMP7, etc.), interleukin, and interferon,
5 but is not limited thereto.

The term "host cell", as used herein, is intended to refer to a cell into which an expression vector has been introduced, e.g. an expression vector encoding an antibody of the invention. Recombinant host cells comprise, for example, transfectomas, such as CHO cells, HEK293 cells, NS/0 cells, and lymphocytic cells.

10 CL domains, CH1 domains, and Fc regions (e.g., CH3 domains) in the antibodies of the present invention may be obtained from any antibody such as IgG1, IgG2, IgG3, or IgG4 subtype, IgA, IgE, IgD, or IgM. The antibodies may be derived from mammals comprising primates such as humans, monkeys, etc. and rodents such as mice, rats, etc. Because antibodies derived from mammals exhibit high sequence homology and structural homology among
15 species, an explanation given of the CL domain, CH1 domain, and CH3 domain in the description is generally applicable to antibodies derived from mammals. In one embodiment, the CL domain, CH1 domain, and CH3 domain may be derived from IgG (e.g., IgG1), but is not limited thereto. As mentioned above, the Fc region in the antibodies described in the present invention comprise two different heavy chains (e.g., different in the sequence of variable
20 domain). Of the two different heavy chains, at least one undergoes an amino acid mutation to increase a possibility of stably forming a heterodimer between the two different heavy chains, but decrease a possibility of stably forming a homodimer between two identical heavy chains.

Bispecific proteins, bispecific antibodies, and antigen-binding fragments thereof provided in the description can be constructed using any means and, for example, by a chemical
25 synthesis or recombinant method. The proteins, the antibodies, and the fragments may be non-naturally occurring.

Mutation sites necessary for dimerization of Fc in the present invention are depicted in FIG. 2. Targets amount to 39 sites for electrostatic interaction-introduced mutation and 14 sites for swapping mutation, and 40 sites for size mutation. In FIG. 2, amino acid residues are
30 expressed as capital letters according to a typical method. The numbering of amino acid residues in the constant region is performed according to the EU-index as described in the document

[Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD. (1991)].

Another aspect of the present invention provides a pharmaceutical composition comprising the aforementioned bispecific protein or bispecific antibody and optionally a
5 pharmaceutically acceptable carrier. Another aspect provides a use of the aforementioned bispecific protein or bispecific antibody in preparing a pharmaceutical composition. Another aspect provides a method for preparing a pharmaceutical composition comprising the aforementioned bispecific protein or bispecific antibody.

An antibody and a composition comprising the same (e.g., pharmaceutical composition)
10 can be applied to diagnosis and treatment, and as such, can be contained in a therapeutic or diagnostic kit.

As used herein, “pharmaceutically acceptable carrier” comprises any and all physiologically compatible solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonicity agents, absorption delaying agents, typical vehicles used for preparation of
15 other drugs, excipients, and additives. Preferably, the carrier may be suitable for intravenous, intramuscular, subcutaneous, parenteral, intraspinal, or epidermal administration (e.g., by injection or infusion).

A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by a person skilled in the art, the route and/or mode of
20 administration will vary depending upon the desired results. To administer a compound of the invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents comprise saline and aqueous buffer solutions.
25 Pharmaceutical carriers comprise sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art.

The phrases “parenteral administration” and “administered parenterally”, as used herein, means modes of administration other than enteral and topical administration, usually by
30 injection, and comprises intravenous, intramuscular, intra-arterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular,

intra-articular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, *supra*, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to comprise isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin. Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors comprising the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

The administration subject may be selected from among mammals comprising primates such as humans, monkeys, etc., rodents such as mice, rats, etc., and the like, isolates therefrom comprising cells, tissues, and body fluid (e.g., blood, etc.), and cultured products thereof.

Advantageous Effects

The present invention provide a highly pure heterodimeric protein as a Chimps protein (e.g., antibody), which is significantly free of contaminants such as homodimers or monomers, and a construction technique therefor. Another advantage of the present invention is to increase

the purity of the bispecific antibody and to introduce a minimal number of mutations to a natural antibody, thereby causing no significant structural changes of natural antibodies and reducing the risk of inducing the antibody to undergo functional loss or abnormality and/or to elicit immune rejection.

5

Description of Drawings

FIG. 1 is a schematic diagram showing only one perfect heterodimeric bispecific antibody (dotted line circle) among a variety of antibody forms possible for construction of bispecific antibodies (10 in total). A and B represent respective heavy chains different from each other and a and b represent respective light chain different from each other.

10

FIG. 2 shows residue positions of antibodies used in electrostatic interaction, swapping, and size methods for inducing heterodimerization in Fc constant regions according to one embodiment, and residue positions related thereto.

FIG. 3 is an SDS-PAGE profile showing heterodimerization results from electrostatic interaction-induced mutations listed in Table 4.

15

FIG. 4 is an SDS-PAGE profile showing heterodimerization results from swapping-associated mutations listed in Table 5.

FIG. 5 is an SDS-PAGE profile showing heterodimerization results from size-associated mutations listed in Table 6.

FIG. 6a shows comparison of 12 mutations that allow outstanding heterodimerization among the single mutations listed in Table 7.

20

FIG. 6b shows heterodimerization results after S364K of two key mutations was substituted with other amino acids according to one embodiment.

FIG. 6c shows heterodimerization results after F405K of two key mutations was substituted with other amino acids according to one embodiment.

25

FIG. 6d shows heterodimerization results after additional mutations were introduced to the three selected key-lock mutation pairs S364K-L368D, S364K-K370S, and F405K-K409F according to one embodiment, wherein x-axis accounts for molecular weights (kD).

FIG. 7 shows heterodimerization efficiency of the single mutations S364K-L368D, S364K-K370S, and F405K-K409F in comparison with conventional techniques (KiH, CPC, and AzS controls). When amino acid residues corresponding to each key are changed to other

30

different amino acids, K is best for S364 and K and R (arginine) show almost the same effect for F405. Positions at which single mutations are made on A and B chains are indicated and heterodimerization quantity is numerically expressed.

FIG. 8 shows heterodimerization efficiencies of a total of four double mutation pairs in comparison with conventional techniques (KiH, CPC, AzS controls), wherein the four double mutations are obtained by selecting two double mutation pairs from combinations of the three mutation pairs S364K-L368D, S364K-K370S, and F405K-K409F and mutating F405 into two types K and R for each pair.

FIGS. 9a to 9c are SDS-PAGE profiles after L368, K370, and K409 corresponding to lock mutations in double mutation pairs are substituted with other amino acids as indicated in Table 6 in order to identify better effects when amino acid residues corresponding to lock mutations are changed to other mutations.

FIG. 10 shows positions of electrostatic interaction-associated mutations, size-associated mutations, and swap-associated mutations in Fab of antibodies.

FIG. 11 is a schematic diagram of a competitive pairing (CPP) assay procedure.

FIG. 12 shows pairing modes established through the process of FIG. 11 between heavy and light chains and visualized on SDS-PAGE after 4D9 and 2B9 having conventional mutation pairs C1, DuetMab, and V23 between heavy and light chains were cloned and co-expressed.

FIG. 13 is an SDS-PAGE profile showing pairing modes when an electrostatic interaction-associated mutation between heavy and light chains is established by substituting K(lysine) for target amino acids on A chain (2B9 heavy chain) and D (aspartic acid) for target amino acids on B chain (4D9 heavy chain).

FIG. 14 shows pairing modes of the mutation at position 30 resulting from substituting heavy chain L145 and light chain V133 with K and D, respectively, which exhibit relatively high pairing accuracy among the list of mutation pairs as the 4D9 and 2B9 antibodies are compared on SDS-PAGE.

FIGS. 15 and 16 show pairing modes after mutation at position 29 S131D and/or S131K is introduced to the light chain of the antibody in which heavy chain L145 is substituted with E or D and light chain V133 is substituted with R, as analyzed by SDS-PAGE.

FIG. 17 shows pairing modes of mutation pairs at position 48 (heavy chain S183 and light chain V133 were substituted with K(R) and D(E), respectively), as analyzed by SDS-

PAGE.

FIG. 18 shows pairing modes analyzed by SDS-PAGE after the mutation pair c29c30c48F Φ 29f30f48 shown in Table 19 and variations thereof are introduced.

FIG. 19 shows pairing modes analyzed by SDS-PAGE after the mutation pairs of Table
5 20 are introduced.

FIG. 20 shows pairing modes analyzed by SDS-PAGE after the mutation pairs of Table 21 are introduced.

FIG. 21 shows pairing ratios of chains after combinations of the mutation pairs at position 34 and 51 selected from among the heavy and light chain mutation pairs are introduced
10 according to one embodiment.

FIG. 22 shows pairing ratios of chains after c34 Φ f51 mutation pair selected from among the heavy and light chain mutation pairs is introduced according to one embodiment.

FIG. 23 shows pairing ratios of chains after c40 Φ f44 mutation pair selected from among the heavy and light chain mutation pairs is introduced according to one embodiment.

FIG. 24 is a schematic diagram of a bispecific antibody in which the heavy and light chains are mutated according to one embodiment.
15

FIG. 25 is a graph showing thermal stability of A chain and B chain for the heavy chain and a chain and b chain for the light chain in antibodies constructed according to one embodiment.

FIG. 26 is a cleavage map of pcDNA3.
20

FIG. 27 is a graph showing hydrophobic interaction chromatography (HIC) results of the bispecific antibodies Trabev and Adabev constructed according to one embodiment, each having c'29c'30c48 Φ f29f30f48 mutation pair and AWBB mutation pair introduced thereto.

FIG. 28 is a graph showing size exclusion chromatography (SEC) analysis results of the bispecific antibody Trabev, constructed according to one embodiment, having
25 c'29c'30c48 Φ f29f30f48 mutation pair and AWBB mutation pair introduced thereto.

FIG. 29 is a graph showing size exclusion chromatography (SEC) analysis results of the bispecific antibodies Trabev and Adabev constructed according to one embodiment, each having c'29c'30c48 Φ f29f30f48 mutation pair and AWBB mutation pair introduced thereto.

FIG. 30 shows dimerization modes of the bispecific antibodies Trabev and Adabev constructed according to one embodiment, each having c'29c'30c48 Φ f29f30f48 mutation
30

pair and AWBB mutation pair introduced thereto.

FIG. 31 is a graph showing affinity for antigens (Her2 and VEGF) of the bispecific antibody Trabev, constructed according to one embodiment, having c`29c`30c48Φf`29f`30f`48 mutation pair and AWBB mutation pair introduced thereto.

5 FIG. 32 is a graph showing affinity for antigens (TNF-alpha and VEGF) of the bispecific antibody Adabev, constructed according to one embodiment, having c`29c`30c48Φf`29f`30f`48 mutation pair and AWBB mutation pair introduced thereto.

FIG. 33a shows sequence alignment between the human IgG1 heavy chain constant region and the human IgA1 heavy chain constant region (human IgG1 heavy chain constant
10 region: SEQ ID NO: 33; human IgA1 heavy chain constant region: SEQ ID NO: 34).

FIG. 33b shows sequence alignment between the kappa constant region and lambda constant region of human immunoglobulin light chain (kappa constant region of human immunoglobulin light chain: SEQ ID NO: 35; and lambda constant region of human immunoglobulin light chain: SEQ ID NO: 36).

15 FIGS. 33c and 33d show sequence alignment of heavy chain constant regions among human, mouse, and rat IgG subtypes (CH1 domain sequences in FIG. 33c and CH3 domain sequences in FIG. 33d) (Human IgG1: SEQ ID NO: 1, Human IgG2: SEQ ID NO: 2, Human IgG3: SEQ ID NO: 3, Human IgG4: SEQ ID NO: 4, Mouse IgG1: SEQ ID NO: 37, Mouse IgG2a^b: SEQ ID NO: 38, Mouse IgG2a^a: SEQ ID NO: 39, Mouse IgG2b: SEQ ID NO: 40,
20 Mouse IgG3: SEQ ID NO: 41, Rat IgG1: SEQ ID NO: 42, Rat IgG2a: SEQ ID NO: 43, Rat IgG2b: SEQ ID NO: 44, Rat IgG2c: SEQ ID NO: 45 in FIG. 33c (CH1 domain); and Human IgG1: SEQ ID NO: 15, Human IgG2: SEQ ID NO: 16, Human IgG3: SEQ ID NO: 17, Human IgG4: SEQ ID NO: 18, Mouse IgG1: SEQ ID NO: 46, Mouse IgG2a^b: SEQ ID NO: 47, Mouse IgG2a^a: SEQ ID NO: 48, Mouse IgG2b: SEQ ID NO: 49, Mouse IgG3: SEQ ID NO: 50, Rat
25 IgG1: SEQ ID NO: 51, Rat IgG2a: SEQ ID NO: 52, Rat IgG2b: SEQ ID NO: 53, and Rat IgG2c: SEQ ID NO: 54 in FIG. 33d (CH3 domain)).

FIG. 34 shows paring levels between light and heavy chains in antibodies, constructed according to one embodiment, having c34Φf51 mutation pair introduced thereto.

FIG. 35 is a graph showing a HIC result of the bispecific antibody Trabev, constructed
30 according to one embodiment, having c34Φf51 mutation pair and AWBB mutation pair.

FIG. 36 shows graphs of HIC results of the bispecific antibody Adabev, constructed

according to one embodiment, having c34Φf51 mutation pair and AWBB mutation pair.

FIG. 37 shows dimerization modes of the bispecific antibodies Trabev and Adabev, constructed according to one embodiment, each having c34Φf51 mutation pair and AWBB mutation pair introduced thereto.

5 FIG. 38 shows pairing levels between light and heavy chains in an antibody, constructed according to one embodiment, having c40Φf44 mutation pair introduced thereto.

FIG. 39 shows graphs of HIC results of the bispecific antibody Trabev, constructed according to one embodiment, having c40Φf44 mutation pair and AWBB mutation pair.

FIG. 40 shows graphs of HIC results of the bispecific antibody Adabev, constructed
10 according to one embodiment, having c40Φf44 mutation pair and AWBB mutation pair.

FIG. 41 shows dimerization modes of the bispecific antibodies Trabev and Adabev, constructed according to one embodiment, each having c40Φf44 mutation pair and AWBB mutation pair introduced thereto.

FIG. 42 is an SDS-PAGE profile showing homodimerization levels upon single
15 transfection of Enb-Fc and Fas-Fc, separately, according to a Comparative Example.

Mode for Invention

Hereinafter, the present invention will be described in further detail with reference to examples. It is to be understood, however, that these examples are for illustrative purposes only
20 and are not to be construed to limit the scope of the present invention.

For the practice of the present invention, all samples were prepared after the following procedure.

Protein Expression

- 25 1. A target gene was cloned to an expression vector (pcDNA3 (Invitrogen)).
2. HEK293E cells (ATCC) were cultured in high-glucose DMEM (Dulbecco's modified Eagle's medium) supplemented with 5 % FBS (fetal bovine serum) in a humidified, CO₂ incubator.
3. A prepared plasmid DNA was introduced by transient transfection into HEK393E
30 cells grown to full confluency. Before transfection, the cells were washed with PBS (phosphate-buffered saline) and, followed by exchanging the culture medium with serum-free high-glucose

DMEM.

4. After incubation for one week, a conditioned medium was used to harvest the proteins which were then filtered. Fc-fusion proteins and antibodies were isolated by protein A chromatography.

5 5. Isolated proteins were quantitatively determined as analyzed at 280nm.

EXAMPLE 1: Selection of Mutation for Heterodimerization of Two Fc Regions

Mutation positions for heterodimerization of two Fc regions (CH3 domain; SEQ ID NO: 15) (on the basis of IgG1) are depicted in FIG. 2. Section was made of 39 positions for electrostatic interaction-associated mutation (electrostatic interaction-introduced mutation) 10 (represented by “Charge J”), 14 positions for swapping-associated mutation (swapping mutation) (represented by “Swap O”), and 40 positions for size-associated mutation (size mutation) (represented by “Size B”). Finally selected positions in CH3 domains and mutated amino acids thereat are summarized in Table 3. Each of the mutation pairs was applied two Fc 15 regions (respectively represented by A chain and B chain) derived from different antibodies, followed by cloning for co-expression. For electrostatic interaction-associated mutation, mutations were carried out to substitute the corresponding amino acids on A chain with K (lysine) representative of amino acids having a positive charge and on B chain with D (aspartic acid) representative of amino acids having a negative charge. Swapping-associated mutation 20 was carried out to exchange the corresponding amino acids on A chain and B chain with each other. For size-associated mutation, the corresponding amino acids were substituted with W (tryptophan) on A chain and with the small size amino acid A (alanine) on B chain.

TABLE 3

Amino acid pair No.	Charge (J)		Swap (O)		Size (B)	
	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B
1	Q347K	K360D	Q347K	K360Q	Q347W	K360A
2	Y349K	S354D			Y349W	S354A
3	Y349K	E357			Y349W	E357A
4	Y349K	K360D			Y349W	K360D
5	L351K	L351D			L351W	L351A
6	P352K	P352D			P352W	P352A
7	S354K	Y349D	S354Y	Y349S	S354W	Y349A
8	D356K	K439D			D356W	K439A
9	E357K	Y349D	E357Y	Y349E	E357W	Y349A
10	E357K	K370D	E357K	K370E	E357W	K370A

11	K360K	Q347D			K360W	Q347A
12	K360K	Y349D	K360Y	Y349K	K360W	Y349A
13	S364K	L368D	S364L	L368S	S364W	L368A
14	S364K	K370D	S364K	K370S	S364W	K370A
15	T366K	T366D			T366W	T366A
16	T366K	Y407D			T366W	Y407A
17	L368K	S364D			L368W	S364A
18	L368K	K409D	L368K	K409L	L368W	K409A
19	K370	E357D			K370W	E357A
20	K370	S364D			K370W	S364A
21	K370	T411D			K370W	T411A
22	N390K	S400D	N390S	S400N	N390W	S400A
23	K392	L398D			K392W	L398A
24	T394K	T394D			T394W	T394A
25	T394K	V397D	T394V	V397T	T394W	V397A
26	P395K	P395D			P395W	P395A
27	P395K	V397D			P395W	V397A
28	V397K	T394D			V397W	T394A
29	V397K	P395D			V397W	P395A
30	L398K	K392D	L398K	K392L	L398W	K392A
31	S400K	N390D			S400W	N390A
32	F405K	K409D	F405K	K409F	F405W	K409A
33	Y407K	T366D	Y407T	T366Y	Y407W	T366A
34	Y407K	Y407D			Y407W	Y407A
35	Y407K	K409D			Y407W	K409A
36	K409	L368D			K409W	L368A
37	K409	F405D			K409W	F405A
38	K409	Y407D			K409W	Y407A
39	T411K	K370D	T411K	K370T	T411W	K370A
40					K439W	D356A

In order to easily identify homodimers or heterodimers of Fc-fusion proteins through SDS-PAGE, fusion was made of the ectodomain (coding sequence of region 1-771 of SEQ ID NO: 24) of TNF-alpha receptor (TNFRSF1B: NP_001057.1 (coding gene: CDS of NM_001066.2; SEQ ID NO: 24)) to Fc (coding gene: SEQ ID NO: 26) on one chain (chain A: 5 Enbrel) and the ectodomain (coding sequence of region 1-519 of SEQ ID NO: 25) of Fas receptor (NP_000034.1 (coding gene: CDS of NM_000043.5; SEQ ID NO: 25)) to Fc (coding gene: SEQ ID NO: 26) on the other chain (chain B: Fas), using pcDNA3 vector (see FIG. 26; Invitrogen) as a backbone. The monomer of chain A has a size of 53 kD while the monomer of chain B has a size of about 44 kD. Because chains A and B, each having an Fc-ectodomain 10 fusion, were different in size from each other, two homodimers (AA and BB) and one heterodimer (AB) could be easily discriminated on SDS-PAGE.

Percentages (%) of dimerization between chains A and between chains B

(homodimerization) are expressed as S_{AA} and S_{BB} , respectively whereas S_{AB} accounts for percentages (%) of dimerization between chains A and B (heterodimerization).

When modes of dimerization were observed by SDS-PAGE, ratios (%) of homodimerization (AA and BB) and heterodimerization (AB) between chains A and B having the mutations of Table 3 introduced therein were compared with those between chains A and B that had not been mutated (represented by WT). Results are given Table 4 and FIG. 3 for the electrostatic interaction-induced mutation, Table 5 and FIG. 4 for the size mutation, and Table 6 and FIG. 5 for the swapping-associated mutation.

TABLE 4

A	WT	Y349K	Y349K	S354K	E356K	E357K	E357K	S364K	T366K	T394K	T394K	T411K
B	WT	E357D	K360D	Y349D	K439D	Y349D	K370D	L368D	Y407D	T394D	V397D	K370D
S_{AA}	28	11	5	27	10	15	12	0	0	11	10	15
S_{AB}	46	70	60	63	64	75	78	100	75	79	73	67
S_{BB}	26	19	35	10	26	10	10	0	25	10	17	18

TABLE 5

A	E357Y	E357K	S364L	S364K	F405K	Y407T	T411K
B	Y349E	K370E	L368S	K370S	K409F	T366Y	K370T
S_{AA}	8	4	8	0	0	10	13
S_{AB}	71	80	61	90	70	83	60
S_{BB}	21	16	31	10	30	7	27

TABLE 6

A	WT	K409W	K409W
B	WT	F405A	Y407A
S_{AA}	22	3	2
S_{AB}	56	60	85
S_{BB}	23	37	13

In Tables 4 to 6, mutations accounting for a heterodimerization rate (%) of 70 % or higher are expressed in bold. As is understood from data of Tables 4 to 6, the tested mutation pairs exhibited a heterodimerization rate of 60 % or higher.

Through the results, selection was made of 12 mutation pairs that gave higher rates to heterodimerization than homodimerization, with the heterodimerization rate being 70 % or higher (electrostatic interaction-introduced mutation: Y349K-E357D, E357K-Y349D, E357K-K370D, S364K-L368D, T366K-Y407D, T394K-T394D, T394K-V397D; and swapping mutation: E357Y-Y349E, E357K-K370E, S364K-K370S, F405K-K409F, Y407T-T366Y)

(expressed in bold in Tables 4 and 5).

Each of the 12 amino acid residue mutations contained in the selected 12 amino acid pairs was introduced into the Fas-Fc fusion protein to express mutant Fas-Fc fusion proteins that contained single mutations at the amino acid positions, respectively. In the same condition, homodimer and monomer survival rates (S_{sm} ; 1=100%) were compared, and the results are depicted in Table 7 and FIG. 6a.

TABLE 7

SM	S_{SM}
E357Y	0.03
Y349D	0.03
K370D	0.06
L368D	0.06
S364K	0.77
K370S	0.05
S400N	0.05
T394D	0.11
F405K	0.67
K409F	0.05
T366A	0.09
T366Y	0.46

In Table 7 and FIG. 6a, two mutations S364K and F405K, which were low in homodimerization rate but high in monomer survival rate (50 % or higher), were selected.

It was postulated that the two selected mutations acted as “key” mutations while the CH3 domain sites interacting therewith on the other chain were “lock” mutations. As combinations therebetween, three key-lock pairs S364K-L368D, S364K-K370S, and F405K-K409F were selected. These mutation pairs were introduced into chain A (key mutation) and chain B (lock mutation) to make three different single mutation pairs (see TABLE 8).

TABLE 8

Mutation type	Sample	Key Single Mutation	Lock Single Mutation
Introduction of electrostatic interaction	J13	S364K	L368D
Swapping	O14	S364K	K370S
Swapping	O32	F405K	K409F

The amino acids at respective key mutation positions were substituted with different

amino acid residues and tested for the single mutation-induced heterodimerization effect, as described above, so as to identify mutation types effective for heterodimerization at the positions. For S364, substitution with K (lysine) was observed to bring about the highest heterodimerization effect. An outstanding effect was detected upon substitution with N (asparagine) and R (arginine) (see FIG. 6b). F405 exhibited excellent similar heterodimerization effects upon substitution with K and R and outstanding heterodimerization effects upon substitution with N and Q (glutamine) (see FIG. 6c).

The additional mutations S364N, S364R, F405R, F405N, and F405Q, which were identified in FIGS. 6b and 6c, were applied to the three selected key-lock mutation pairs S364K-L368D, S364K-K370S, and F405K-K409F, to introduce the lock mutations to chain A (TNFR2-Fc) and the key mutations to chain B (Fas-Fc), followed by analyzing heterodimerization effects on SDS-PAGE.

The results thus obtained are depicted in Table 9 and FIG. 6d:

TABLE 9

Sample	A (TNFR2)	B (Fas)	S _{AA}	S _{AB}	S _{BB}
	Key Single Mutation	Lock Single Mutation			
J13(K:D)	S364K	L368D	8	92	0
J13(N:D)	S364N	L368D	8	82	9
J13(R:D)	S364R	L368D	20	79	1
O14(K:S)	S364K	K370S	25	65	9
O14(N:S)	S364N	K370S	13	84	3
O14(R:S)	S364R	K370S	23	76	1
O32(K:F)	F405K	K409F	0	75	25
O32(N:F)	F405N	K409F	21	62	17
O32(Q:F)	F405Q	K409F	5	68	27
O32(R:F)	F405R	K409F	5	89	6

(S_{AA}: AA homodimerization rate (%);

S_{AB}: AB heterodimerization rate (%);

S_{BB}: BB homodimerization rate (%))

EXAMPLE 2: Test for Heterodimerization of Fc Region by Single Mutation

The three key-lock mutation pairs S364K-L368D, S364K-K370S, and F405K-K409F, which were selected in Example 1, and the mutation pair F405R-K409F, which resulted from

substituting F405 with R instead of K, were tested for heterodimerization on SDS-PAGE. The heterodimerization effects were compared with those obtained with the conventional heterodimer Fc mutation pairs KiH, CPC, and AzS used as controls.

SDS-PAGE data in this Example and all the following Examples were obtained by quantitating the band intensities with the aid of GelQuant.NET Software.

The results are depicted in Table 10 and FIG. 7:

TABLE 10

Sample	Chain A (eTNFR2)	Chain B (eFas)	S_{AB}	S_A	S_B	S_M	T_m
J13	S364K	L368D	0.93	0.86	0.10	0.48	66.8
O14	S364K	K370S	0.87	0.86	0.07	0.47	67.6
O32	F405K	K409F	0.72	0.91	0.08	0.49	65.2
O32'	F405R	K409F	0.72	0.91	0.08	0.49	65.2
KiH	T366S/L368A/Y407V	T366W	0.90	0.68	0.64	0.66	67.4
CPC	K392D/K409D	E356K/D399K	0.74	0.85	0.30	0.58	66.4
AzS	T350V/T366L/K392L/T394W	T350V/L351Y/F405A/Y407V	0.84	0.84	0.64	0.74	69.2

(S_{AB} : AB heterodimerization rate (%); S_A : AA homodimerization rate (%);

S_B : BB homodimerization rate (%); S_M : monomer survival rate (%))

T_m was measured as follows:

Reagent: Invitrogen 4461146 "Protein Thermal ShiftTM" Dye Kit

Instrument: Chromo4-PTC200 (MJ Research)

Reaction mixture: 20 $\mu\ell$ in total

Protein 10 $\mu\ell$

DW 3 $\mu\ell$

Protein Thermal ShiftTM Buffer 5 $\mu\ell$

1/100 diluted Protein Thermal ShiftTM Dye 2 $\mu\ell$

Protocol:

1. Incubate at 50.0 $^{\circ}\text{C}$ for 30sec;

2. Melting Curve from 50.0 $^{\circ}\text{C}$ to 90.0 $^{\circ}\text{C}$, read every 0.2 $^{\circ}\text{C}$, hold for 2 sec;

3. Incubate at 90.0 $^{\circ}\text{C}$ for 2 min

4. Incubate at 10.0□ forever

5. End

As shown in Table 10 and FIG. 7, when chain A and chain B were each separately expressed, chain A retaining the key did not form a homodimer, but is expressed as a monomer.

5 A sample resulting from co-expression was observed to contain no homodimers, but mostly heterodimers (see FIG. 7).

EXAMPLE 3: Heterodimerization of Fc Region by Double Mutation

Less plausibility of homodimerization might result from existence of the key mutations
10 on both of chains A and B than on either of the chains. Two double mutation pairs were made from combinations of the three mutation pairs selected in Example 1. For each double mutation pair, F405 was mutated into two types K and R. Thus, a total of four double mutation pairs were obtained. These found double mutations pairs were analyzed for heterodimerization on SDS-PAGE and the dimerization effects were compared with those of the controls KiH, CPC, and
15 AzS.

The results are given in Table 11 and FIG. 8:

TABLE 11

DMP	Chain A (eTNFR2)	Chain B (eFas)	S_{AB}	S_A	S_B	S_M	T_m
J13/O32	S364K/K409F	L368D/F405K	0.96	1.00	0.74	0.87	58.2
J13/O32'	S364K/K409F	L368D/F405R	0.95	1.00	0.73	0.87	61.5
O14PO32	S364K/K409F	K370S/F405K	0.95	1.00	0.72	0.86	64.1
O14/O32'	S364K/K409F	K370S/F405R	0.93	1.00	0.71	0.86	64.4
KiH	T366S/L368A/Y407V	T366W	0.91	0.64	0.61	0.63	67.4
CPC	K392D/K409D	E356K/D399K	0.73	0.81	0.30	0.58	66.4
AzS	T350V/T366L/K392L/T394W	T350V/L351Y/F405A/Y407V	0.84	0.82	0.62	0.72	69.2

As shown in Table 11 and FIG. 8, the key mutations present in both chains allowed the
20 homodimerization of the chains for none of the four mutation pairs upon separate expression. All of the proteins in a sample resulting from co-expression were observed to be heterodimers

(See FIG. 8). In addition, higher thermal stability was measured in a double-mutation pair having F405R introduced thereto than F405K introduced thereto (Table 11).

In order to examine whether a better effect was obtained when amino acids corresponding to lock mutation were substituted with other residues, L368, K370, and K409, which were the lock in the double mutation pairs, were substituted with other amino acid residues. Mutation combinations obtained by variously mutating L368, K370, and K409 are summarized in Table 12. The mutation combinations were tested for heterodimerization on SDS-PAGE (NR: 8% SDS-PAGE gel; Sample: 24ul Loading), and the results are depicted in FIGS. 9a to 9c.

TABLE 12

Lock variants

X	Chain A	Chain B	Y	Chain A	Chain B	Z	Chain A	Chain B
UA	S364K/ K409F	L368A/ F405R	XA	S364K/ K409F	K370A/ F405R	ZA	S364K/ K409A	K370S/ F405R
UC	S364K/ K409F	L368C/ F405R	XC	S364K/ K409F	K370C/ F405R	ZC	S364K/ K409C	K370S/ F405R
UD	S364K/ K409F	L368D/ F405R	XD	S364K/ K409F	K370D/ F405R	ZD	S364K/ K409D	K370S/ F405R
UE	S364K/ K409F	L368E/ F405R	XE	S364K/ K409F	K370E/ F405R	ZE	S364K/ K409E	K370S/F405 R
UF	S364K/ K409F	L368F/ F405R	XF	S364K/ K409F	K370F/ F405R	ZF	S364K/ K409F	K370S/ F405R
UG	S364K/ K409F	L368G/ F405R	XG	S364K/ K409F	K370G/ F405R	ZG	S364K/ K409G	K370S/ F405R
UH	S364K/ K409F	L368H/ F405R	XH	S364K/ K409F	K370H/ F405R	ZH	S364K/ K409H	K370S/ F405R
UI	S364K/ K409F	L368I/ F405R	XI	S364K/ K409F	K370I/ F405R	ZI	S364K/ K409I	K370S/ F405R
UK	S364K/ K409F	L368K/ F405R	XK	S364K/ K409F	K370/ F405R	ZK	S364K/ K409K	K370S/ F405R
UL	S364K/ K409F	L368/ F405R	XL	S364K/ K409F	K370L/ F405R	ZL	S364K/ K409L	K370S/ F405R
UM	S364K/ K409F	L368M/ F405R	XM	S364K/ K409F	K370M/ F405R	ZM	S364K/ K409M	K370S/ F405R
UN	S364K/ K409F	L368N/ F405R	XN	S364K/ K409F	K370N/ F405R	ZN	S364K/ K409N	K370S/ F405R

UQ	S364K/ K409F	L368Q/ F405R	XQ	S364K/ K409F	K370Q/ F405R	ZQ	S364K/ K409Q	K370S/ F405R
UR	S364K/ K409F	L368R/ F405R	XR	S364K/ K409F	K370R/ F405R	ZR	S364K/ K409R	K370S/ F405R
US	S364K/ K409F	L368S/F 405R	XS	S364K/ K409F	K370S/ F405R	ZS	S364K/ K409S	K370S/ F405R
UT	S364K/ K409F	L368T/ F405R	XT	S364K/ K409F	K370T/ F405R	ZT	S364K/ K409T	K370S/ F405R
UV	S364K/ K409F	L368V/ F405R	XV	S364K/ K409F	K370V/ F405R	ZV	S364K/ K409V	K370S/ F405R
UW	S364K/ K409F	L368W/ F405R	XW	S364K/ K409F	K370W/ F405R	ZW	S364K/ K409W	K370S/ F405R
UY	S364K/ K409F	L368Y/ F405R	XY	S364K/ K409F	K370Y/ F405R	ZY	S364K/ K409Y	K370S/ F405R

The combinations were measured for thermal stability (see Example 2) and the results are given in Table 13.

TABLE 13

	Chain A	Chain B	T_m
UC	S364K/K409F	L368C/F405R	63.8
UD	S364K/K409F	L368D/F405R	60.8
UL	S364K/K409F	L368/F405R	63.0
UW	S364K/K409F	L368W/F405R	61.2
UY	S364K/K409F	L368Y/F405R	61.2
ZF	S364K/K409F	K370S/F405R	65.6
ZH	S364K/K409H	K370S/F405R	64.4
ZI	S364K/K409I	K370S/F405R	62.4
ZN	S364K/K409N	K370S/F405R	61.0
ZR	S364K/K409R	K370S/F405R	62.8
ZT	S364K/K409T	K370S/F405R	65.4
ZV	S364K/K409V	K370S/F405R	66.0
ZW	S364K/K409W	K370S/F405R	67.0

ZY	S364K/K409Y	K370S/F405R	63.8
----	-------------	-------------	------

Analysis showed higher thermal stability of K409W (tryptophan) than K409F.

Based on the data obtained above, mutations were made of S364K and K409W into chain A and K370S and F405R into chain B to produce a double mutation pair, termed AWBB mutation pair, for use in the following Fc heterodimerization test of CH3 domain.

EXAMPLE 4: Selection of Mutation for Heavy Chain and Light Chain in Antibody Fab

To select mutations in heavy and light chains of antibodies, electrostatic interaction-associated mutations, size-associated mutations, and swapping-associated mutations were carried out. Positions of interaction at heavy and light chains are depicted in FIG. 10.

In order to easily identify mutations in heavy and light chains, antibodies having the same light chain were cloned. 4D9 antibody (anti-Influenza A antibody) and 2B9 antibody (anti-Influenza B antibody), which are both anti-influenza antibodies, have the same light chain (common light chain: CLC). Because they have the same light chain, SDS-PAGE analysis allows interaction between heavy and light chains to be easily understood. Although identical in the amino acid sequence of light chain, the two antibodies 4D9 and 2B9 are different from each other with respect to the sequence and size of the heavy chain (the heavy chain of 2B9 has more amino acid residues by six than that of 4D9 and a size (50130.62 Daltons) greater than that of 4D9 (49499.98 Daltons): they can be clearly discriminated on SDS-PAGE). Thus, size analysis on SDS-PAGE makes it possible to understand which of the two chains interacts with the light chain.

Amino acid sequences and coding nucleic acid sequences thereof in the heavy chain variable regions and light chain variable regions of the two antibodies 4D9 and 2B9 are listed in Table 14, below.


TABLE 14

	Amino Acid Sequence	Nucleic Acid Sequence
2B9 Heavy Chain Variable Region	EVQLVESGGGLVQPGKSLRLSC AATGFTFDDYAMHWVRQAPG KGLEWVSSLNWKGNSVDYAD SVRGRFTMSRDNAKKLVYLM NGLRGDDTAVYFCAKDNKAD	GAAGTGCAGCTGGTGGAGTCTGGGGGAG GCTTGGTACAGCCTGGCAAGTCCCTGAG ACTCTCCTGTGCAGCCACTGGATTACAT TTGACGATTACGCCATGCACTGGGTCCGC CAAGCTCCAGGGAAGGGCCTGGAGTGGG

	ASMDYYYHHGMDVWGQGT VTVSS (SEQ ID NO: 27)	TCTCAAGTCTTAATTGGAAGGGAAATAGT GTAGACTACGCGGACTCTGTGAGGGGCC GATTCACCATGTCCAGAGACAACGCCAA GAAACTAGTGTATCTGCAAATGAACGGT CTGAGAGGTGACGACACGGCCGTCTATTT TTGTGCAAAAGATAATAAAGCGGATGCA TCTATGGACTACTACTACCACCACGGTAT GGACGTCTGGGGCCAAGGGACCACGGTC ACCGTCTCCTCG (SEQ ID NO: 28)
4D9 Heavy Chain Variable Region	QVTILRESGPGLVKPSETLSLTCT ISGASINTDYWSWIRQPPGKGL WIGYIYYRGRITNYPNPSLRSRVTI SVDTSKNQFSL QMTSMTAADTAVYYCARDVT GISRENAFDIWGQGTILTVSS (SEQ ID NO: 29)	CAGGTCACCTTGAGGGAGTCGGGGCCAG GACTGGTGAAGCCTTCGGAGACCCTGTCC CTCACCTGCACTATCTCCGGTGCCTCCAT CAATACTGACTACTGGAGCTGGATCCGG CAGCCCCAGGGAAGGGACTGGAGTGGA TTGGCTATATCTATTACAGAGGGCGCACC AACTACAACCCCTCCCTCAGGAGCCGAG TCACCATATCAGTAGACACGTCCAAGAA TCAATTCTCCCTG CAGATGACGTCTATGACCGCTGCTGACAC GGCCGTATATTACTGTGCGAGAGATGTG ACTGGCATCAGTCGAGAAAATGCTTTTGA TATCTGGGGCCAAGGCACCCTGGTCACC GTCTCCTCG (SEQ ID NO: 30)
Light Chain (CLC) Variable Region	AIRMTQSPSSLSASVGDRTTTC RASQISGYLNWYQKPKGAP KLLIYAASSLQSGVPSRFSGSGS GTDFTLTISLQPEDFATYYCQQ SYSIPTTFGQGTIRLEIK (SEQ ID NO: 31)	GCCATCCGGATGACCCAGTCTCCATCCTC CCTGTCTGCATCTGTAGGAGACAGAGTCA CCATCACTTGCCGGGCAAGTCAGAGCATT AGCGGCTATTTAAATTGGTATCAGCAGA AACCAGGGAAGCCCTAAGCTCTTGAT CTATGCTGCATCCAGTTTGCAGAGTGGGG TCCCATCAAGGTTCAGTGGCAGTGGATCT GGGACAGATTTCACTCTACCATCAGCAG TCTGCAACCTGAAGATTTTGCAACTTACT ACTGTCAACAGAGCTACAGTATCCCCACC ACCTTCGGCCAAGGGACACGACTGGAGA TTAAA (SEQ ID NO: 32)

The two antibodies 4D9 and 2B9 employ the constant region of IgG1 as a heavy chain constant region and the kappa constant region as a light chain constant region.

Examination was made to see whether a light chain having a mutation introduced
5 thereinto pairs with only a heavy chain having a mutation interacting with the mutation of the
light chain. In this regard, a light chain, a heavy chain pairing with the light chain, and a heavy
chain forming a mispair with the light chain were co-expressed to afford antibodies. SDS-
PAGE analysis of the antibodies in a reducing condition can identify the extent to which the
pairings are accurately formed. For convenience, 4D9 heavy chain is referred to as A chain and
10 a light chain pairing therewith as a chain, and 2B9 heavy chain is referred to as B chain and a
light chain pairing therewith as b chain (see the following illustrations).



First, 4D9 and 2B9 having conventional mutation pairs between heavy and light chains, already known to be effective, were cloned and co-expressed. Pairing modes established through competition between the resulting mutant chains, that is, two heavy chains and one light chain were examined on SDS-PAGE. (Competitive Pairing (CPP) Assay). The process of performing a competitive pairing (CPP) assay is schematically illustrated in FIG. 11, and the result thus obtained is depicted in FIG. 12. As shown in FIG. 12, all the already known mutation pairs undergo normal pairing and abnormal mispairing, concurrently.

For electrostatic interaction-associated mutations between heavy and light chains, antibodies in which various mutation pairs were introduced with the substitution of corresponding amino acids in B chain (2B9 heavy chain) with K (lysine) and in A chain (4D9 heavy chain) with D (aspartic acid) were subjected to CPP assay on SDS-PAGE (see FIG. 11).

As a result, seven candidate mutation pairs that were identified to undergo relatively accurate pairing were screened in the electrostatic interaction-associated mutation group. CPP assay results on SDS-PAGE of the antibodies into which the seven screened mutation pairs were introduced are given in Table 15 and depicted in FIG. 13.

TABLE 15

No	SOP (Symmetric Orthogonal Pairs)	A (4D9)	B (2B9)	a	b	a^{CPP}	b^{CPP}	S^{CPP}
c	cΦf	L145	L145	S131	S131			
1	c29Φf29	L145D	L145K	S131K	S131D	58	50	54
2	c30Φf30	L145D	L145K	V133K	V133D	50	79	65
3	c34Φf34	K147D	K147K	T180K	T180D	49	87	68
4	c40Φf40	F170D	F170K	L135K	L135D	59	61	60
5	c44Φf44	P171D	P171K	S162K	S162D	59	51	55
6	c48Φf48	S183D	S183K	V133K	V133D	62	80	72
7	c51Φf51	V185D	V185K	L135K	L135D	67	73	70

$$(CPP \text{ Score } (S^{CPP}) = \frac{1}{2}(a^{CPP} + b^{CPP}))$$

$$a^{CPP} = 100(C_{aA})/(C_{aA} + C_{aB}) = 100(C_A)/(C_A + C_B)$$

$$b^{CPP} = 100(C_{bB})/(C_{bB} + C_{bA}) = 100(C_B)/(C_A + C_B)$$

- 5 Comparison on SDS-PAGE found some mutation pairs that underwent relatively accurate pairing in the list of mutation pairs using the 4D9 (A chain) and 2B9 (B chain) antibodies (expressed in bold in Table 15). For further study, mutation at position 30 (expressed as c30Φf30) was modified as in Table 16.

TABLE 16

Mutation c30Φf30			
	Heavy Chain	Light Chain	Pairing Accuracy
Aa (4D9)	L145D	V133R	75%
Bb (2B9)	L145R	V133D	60%

10

- The mutation at position 30 resulted from substitution heavy chain L145 and light chain V133 with K and D, respectively. As is understood from the data of Table 16, the effect (pairing accuracy: Aa pairing or Bb pairing ratio) was observed to be good (Aa pairing accuracy 75%, Bb pairing accuracy: 60%) for the variations in which heavy chain L145 and light chain V133 were substituted with R and D, respectively (see FIG. 14).

- Addition of mutation S131D on the light chain (termed mutation at position 29) to the mutation at position 30 or variations thereof (heavy chain L145 and light chain V133 were substituted with K or R, and D or E, respectively) improved the accuracy of pairing (see Table 17 and FIG. 15 (c29c^R30Φf^R30 result; Aa pairing accuracy: 80% and Bb pairing accuracy: 70%) and FIG. 16 (c^R29c^{RE}30Φf^R29f^{RE}30 result; Aa pairing accuracy: 90% and Bb pairing accuracy: 80%)).

20

TABLE 17

	Heavy Chain	Light Chain	Paring Accuracy
c^{29c}R³⁰Φ^R30			
Aa (4D9)	L145D	V133R	80%
Bb (2B9)	L145R	S131D/V133D	70%
c^R29c^{RE}30Φ^R29f^{RE}30			
Aa (4D9)	L145E	S131K/V133R	90%
Bb (2B9)	L145R	S131D/V133E	80%

As can be seen in FIGS. 14 and 15, outstanding pairing accuracy was detected in the cases where heavy chain L145 and light chain V133 were substituted with R and D, respectively and where substitution of light chains S131 and V133 were respectively made by K and R, or D and E as well as in mutation at position 30 wherein heavy chain L145 and light chain V133 were substituted with K and D, respectively.

In addition, a mutation pair in which heavy chain S183 and light chain V133 were respectively substituted with K (or R) and D (or E) (termed mutation at position 48) was also observed to be effective (Table 18 and FIG. 17 (Aa pairing accuracy (low bands): 45%, Bb pairing accuracy (upper bands): 95%).


TABLE 18

LC	V133K (a)	V133D (b)
2B9 Heavy Chain (B)	S183K	
4D9 Heavy Chain (A)	S183D	
mg/L	2.6	2.4

Antibodies containing a combination of the aforementioned mutation at position 29, mutation at position 30, and mutation at position 48 (c^{29c}30c^{48F}Φ^{29f}30f⁴⁸) or variations thereof were constructed (see Table 19) and then tested for pairing accuracy. The results are depicted in FIG. 18 (lower bands: Aa pairing; and upper bands: Bb paring):

TABLE 19

	Heavy Chain	Light Chain	Paring Accuracy
c^R29c^{RE}30c^{48F}Φ^R29f^{RE}30f⁴⁸			
Aa (4D9)	L145E/S183D	S131K/V133R	100%
Bb (2B9)	L145R/S183K	S131D/V133E	90%
Mutation 1			
Aa (4D9)	L145D/S183D	S131K/V133K	85%
Bb (2B9)	L145K/S183K	S131D/V133D	90%
Mutation 2			
Aa (4D9)	L145D/S183D	S131D/V133K	95%



In addition, antibodies containing a variant mutation pair of at least one of mutation at position 29, mutation at position 30, and mutation at position 48 (see Table 20) were examined for pairing accuracy and the results are depicted in FIG. 19 (lower bands: Aa pairing; and upper bands: Bb pairing):

5

TABLE 20

No.	1		2		3		4		5	
C	c30Φf30		c ^R 30Φf ^R 30		c ^R 29c ^R 30Φf ^R 30		c ^R 29c ^{RE} 30Φf ^R 29f ^{RE} 30		c ^R 29c ^{RE} 30f48Φf ^R 29f ^{RE} 30f48	
LC	V133K (a)	V133D (b)	V133R	V133D	V133R	S131D/ V133D	S131K/ V133R	S131D/ V133E	S131K/ V133R	S131D/ V133E
4D9 (A)	L145D		L145D		L145D		L145E		L145E/S183D	
2B9 (B)	L145K		L145R		L145R		L145R		L145R/S183K	
Aa	79%		67%		80%		90%		100%	
Bb	50%		62%		70%		80%		90%	

(Aa: Aa pairing accuracy; Bb: Bb pairing accuracy)

Further, combinations of mutation at position 29, mutation at position 30, and mutation at position 48 were subjected to swapping between D and E to search for combinations that allowed for relative accurate pairing (Table 21 and FIG. 20).

10

TABLE 21

Code	A	B	a	b	Aa	Bb
0	L145K/S183K	L145D/S183D	S131D/V133D	S131K/V133K	65%	100%

ab	L145K/S183K	L145E/S183E	S131D/V133D	S131K/V133K	95%	95%
gh	L145K/S183K	L145D/S183D	S131E/V133E	S131K/V133K	100%	100%
abgh	L145K/S183K	L145E/S183E	S131E/V133E	S131K/V133K	100%	100%

(Aa: Aa pairing accuracy; Bb: Bb pairing accuracy)

As can be seen in Table 21 and FIG. 20, all the tested mutation pairs ab, gh, and abgh exhibited a pairing accuracy of 65% or higher, or 95% or higher. Of them, abgh was selected and termed c'29c'30c48Φf 29f 30f 48 mutation pair.

- 5 Chains into which combinations of the mutation pairs at position 34 and 51 selected from among the mutation pairs identified in Table 15 were introduced were tested for pairing and the results are given in Table 22 and FIG. 21.

TABLE 22

LC	L135K/T180K (a)	L135D/T180D (b)
4D9 Heavy Chain (A)	K147D/V185D	
2B9 Heavy Chain (B)	K147/V185K	
Aa (%)	95%	
Bb (%)	62%	

- 10 In addition, mutation at position 34 and mutation at position 51 of Table 15 were subjected to change from K to R and from D to E, followed by a pairing test to detect combinations having improved pairing accuracy. As a result, a combination in which L135 and T180 on the light chain were each substituted with E was improved in pairing accuracy and termed c34Φf51 mutation pair (see Table 23 and FIG. 22).

TABLE 23

LC	L135K/T180K (a)	L135E/T180E (b)
4D9 Heavy Chain (A)	K147D/V185D	
2B9 Heavy Chain (B)	K147/V185K	
Aa(%)	95%	
Bb(%)	86%	

- 15 Of various mutation pairs found in Table 15, a combination of mutation at position 40 and mutation at position 44 was also obverted to improve in pairing accuracy.

- In addition, all the mutation pairs were subjected to swapping between K and R and between D and E to search for a combination that allowed for the most accurate pairing. The combination was a mutation pair in which light chain L135 was substituted with R and E, and
20 was termed c40Φf44 mutation pair (see Table 24 and FIG. 23):

TABLE 24

LC	L135R/S162K (a)	L135E/S162D (b)
----	-----------------	-----------------

4D9 Heavy Chain (A)	F170D/P171D
2B9 Heavy Chain (B)	F170K/P171K
Aa(%)	95%
Bb(%)	84%

EXAMPLE 5: Bispecific Antibody Formation by Coupling between Heavy Chains and between Heavy Chain and Light Chain

5.1. Bispecific antibody having c'29c'30c48Φf'29f'30f'48 mutation pair introduced thereto

5.1.1. Test of coupling between heavy and light chain using 4D9/2B9 antibody

Antibodies were constructed by coupling heavy and light chains containing the c'29c'30c48Φf'29f'30f'48 mutation pair (see Table 25):

TABLE 25

Antibody	Heavy Chain Constant Region (HC: CH1)	Light Chain Constant Region (LC)
4D9 (Aa)	L145E/S183E	S131K/V133K
2B9 (Bb)	L145K/S183K	S131E/V133E

In order to examine the accuracy of pairing among A chain, B chain, a chain, and b chain in the bispecific antibodies, individual heavy chains and light chains were co-expressed in all possible combinations of normal pairs and abnormal mispairs thereamong. Expression levels measured by SDS-PAGE are given in Table 26, below. Thermal stability (T_m) of the combinations was measured and the results are given in Table 27 and FIG. 25:

TABLE 26

HC	A		B	
LC	a	b	a	b
Expression Amount	89.8mg/l	18.8 mg/l	24.5 mg/l	68.5 mg/l

TABLE 27

Pairs	T _m
4D9 WT	69.9 ±2.00
2B9 WT	69.1 ±1.50
Aa	70.3 ±1.70
Ab	58.5 ±11.96
Bb	68.7 ±0.92

Ba	58.6 ±0.53
----	------------

Thermal stability (Tm) was measured with reference to the method explained in Example 2.

As shown in Tables 26 and 27 and FIG. 25, the bispecific antibodies having c'29c'30c48Φf29f30f48 mutation pair introduced thereto were found to be higher in expression level and thermal stability for normal pairing (Aa/Bb) than for abnormal pairing (Ab/Ba).

5.1.2. Trastuzumab/Bevacizumab bispecific antibody or Adalimumab/Bevacizumab bispecific antibody

Trastuzumab (Herceptin®; Roche), Bevacizumab (Avastin™; Roche), and Adalimumab (Humira®; AbbVie) were purchased and subjected to amino acid sequencing (the Korea Basic Science Institute, Korea). cDNAs corresponding to the amino acid sequences were synthesized and used to construct bispecific antibodies to which the c'29c'30c48Φf29f30f48 mutation pair and the AWBB mutation pair of CH3 domain selected in Example 3 were introduced in the combinations shown in Table 28, below (pcDNA3 vector (see FIG. 26) used).

TABLE 28

Antibody	Heavy Chain Constant Region (HC)		Light Chain Constant Region (LC)
	CH1	CH3	
Trabev			
Trastuzumab (Aa)	L145E/S183E	S364K/K409W	S131K/V133K
Bevacizumab (Bb)	L145K/S183K	K370S/F405R	S131E/V133E
Adabev			
Adalimumab (Aa)	L145E/S183E	S364K/K409W	S131K/V133K
Bevacizumab (Bb)	L145K/S183K	K370S/F405R	S131E/V133E

The bispecific antibodies Trabev and Adabev were subjected to hydrophobic interaction chromatography (HIC) in the following condition and the results are depicted in FIG. 27 (y-axis: Value (mAU); and x-axis: time (min)):

Instrument: HPLC-U3000
Column: MAbPac Hic-20
Flow rate: 0.2mL/min
Detection: UV, 280nm
Mobile Phase: 0.10 M Ammonium acetate, pH 7.0

Passage of the obtained protein through the HIC column formed peaks at different time points depending on the hydrophobicity of the protein, whereby the construction of accurate bispecific antibodies could be accounted for. As can be seen in FIG. 27, peaks for the heterodimerized bispecific antibodies (Trabev and Adabev) are distinctively observed between peaks for two different homodimer antibodies (Trastuzumab and Bevacizumab, or Adalimumab and Bevacizumab), indicating the fine formation of bispecific antibodies, each composed of halves from two different antibodies.

In addition, the bispecific antibodies Trabev and Adabev thus obtained were subjected to size exclusion chromatography (SEC) analysis, and the results are given in Table 29 and FIGS.

28 and 29 (x-axis: time (min)):

Instrument: HPLC-U3000

Column: size-exclusion chromatography T SKgel G3000SWXL Tosoh Bioscience

Flow rate: 1.0mL/min

Detection: UV, 280nm

Mobile Phase: 25mM Tris-HCl (pH 8.5), 150 mM NaCl

TABLE 29

WT (Monospecific Antibody)		BsAb (Bispecific Antibody)	
1A.Trastuzumab	8.007min.	1" Trabev	7.847 min.
1B.Bevacizumab	7.743 min.	2" Adabev	7.833 min.
2A.Adalimumab	7.987 min.		

In the SEC analysis, peaks are detected according to protein size and can elucidate protein aggregation. As shown in Table 29 and FIGS. 28 and 29, peaks for the bispecific antibodies are present between peaks of the two corresponding antibodies on the time axis, indicating the fine formation of the bispecific antibodies.

Further, heterodimerization modes of the bispecific antibodies Trabev and Adabev on SDS-PAGE are depicted in FIG. 30. As can be seen in FIG. 30, the bispecific antibodies Trabev and Adabev, which are heterodimers, were detected as single bands at intermediate sizes between Trastuzumab and Bevacizumab and between Adalimumab and Bevacizumab, respectively. These results imply that the bispecific antibodies are not homodimers, but are constructed only as a result of normal pairing.

Through ELISA, tests were conducted to examine whether the BsAbs bind effectively to

respective antigens (Trastuzumab: Her2), Bevacizumab (VEGF), and Adalimumab (TNF-alpha).

Affinity for antigen was measured as follows:

Reagent

5 Detection antibody: goat anti-human kappa-HRP (southern biotech, 2060-05)
TMB single solution (LIFE TECHNOLOGY, 002023)

Instrument

Emax precision microplate reader (Molecular devices)

Protocol

10 Coating buffer: Carbonate buffer pH 9.6
Blocking buffer: protein-free(TBS) blocking buffer (Thermo scientific)
Wash buffer: 0.05%(w/v) Tween20 in TBS, pH7.4 (TBST)
Diluent: 0.05%(w/v) Tween20 in TBS, pH7.4
Stop buffer: 1N Hydrochloric acid solution (HCl)

15 Protocol

Coating: dilute antigen in coating buffer, load 100 ul of dilution to each well, and incubate at 4°C overnight (Her2, VEGF: 50ng/well, TNF-alpha: 100ng/well);

Washing 3 times with washing buffer;

Blocking: load 300 ul of blocking buffer, incubate at room temperature (RT) for 1 hour;

20 Washing 3 times with washing buffer;

Binding: load antibodies at an aliquot of 100ng/well, and incubate at RT for 1 hr;

Washing 3 times with washing buffer;

Detection Antibody: dilute goat anti-human kappa-HRP in TBST at a ratio of 1:4000 and incubate at RT for 1 hr;

25 Washing 3 times with washing buffer

Detection: load 100 ul of TMB solution per well, incubate at RT for 3 min in the dark;

Stop solution: load 100 ul of 1N HCl per well;

Reading: read at optical density 450nm

The results thus obtained are given in Table 30 and FIG. 31 (for Trabev) and in Table 31
30 and FIG. 32 (for Adabev):

TABLE 30

Antigen	Her2			VEGF		
Anti-body	Trastuzumab	Bevacizumab	Tra-bev	Trastuzumab	Bevacizumab	Tra-bev
O.D	2.50	0.05	2.54	0.65	2.89	2.04
	2.96	0.05	2.60	0.70	2.88	2.25
	2.58	0.04	2.49	0.60	3.12	2.36
Mean	2.68	0.05	2.54	0.65	2.96	2.22
blank	0.05			0.05		

TABLE 31

Antigen	TNF-alpha			VEGF		
Anti body	Adalimumab	Bevacizumab	Ada-bev	Adalimumab	Bevacizumab	Ada-bev
O.D	1.24	0.23	0.93	0.07	2.24	2.37
	1.26	0.22	1.16	0.06	2.27	2.28
	1.19	0.27	1.09	0.07	2.60	2.47
Mean	1.23	0.24	1.06	0.07	2.37	2.37
blank	0.06			0.06		

As shown in Table 29 and FIG. 31, the bispecific antibody Trabev was found to bind to both Her2 and VEGF, which are antigens of Trastuzumab and Bevacizumab, respectively. Also, data in Table 30 and FIG. 32 proved that the bispecific antibody Adabev binds to TNF-alpha and VEGF, which are antigens of Adalimumab and Bevacizumab, respectively. These results confirmed that the two bispecific antibodies normally exerting desired functions were successfully constructed.

5.2. Bispecific antibody having c34Φf51 mutation pair introduced thereto

With reference to Example 5.1.1, antibodies having c34Φf51 mutation pair (see Table 23) introduced to 4D9 (Aa) and 2B9 (Bb) thereof were constructed. For each heavy chain, a and b light chains were co-expressed, and inter-light/heavy chain pairing ratios were measured on SDS-PAGE (conducted for heavy chains A and B, each). T_m was measured as in Example 2, and the results are given in Table 32, below and FIG. 34.

TABLE 32

	Test 1		Test 2	
HC	A (K147D/V185D)		B (K147/V185K)	
LC	a (L135K/T180K)	b (L135E/T180E)	b (L135E/T180E)	a (L135K/T180K)

<i>q</i>	90%	10%	89%	11%
<i>T_m</i>	65.8	N/A	65.8	N/A

(*q*: light chain/heavy chain pairing ratio (%); N/A: not available)

As shown in Table 32 and FIG. 34, normal heavy chain/light chain pairs (Aa and Bb) were formed at a ratio of as high as 90% and 89%, respectively, and exhibited high thermal stability.

5 In addition, bispecific antibodies Trabev (Aa: Trastuzumab; Bb: Bevacizumab) and Adabev (Aa: Adalimumab; Bb: Bevacizumab) to each of which the c34Φf51 mutation pair and the CH3 domain AWBB mutation pair (Aa: S364K/K409W; Bb: K370S/F405R) selected in Example 3 were introduced were constructed using Trastuzumab (Herceptin®; Roche), Bevacizumab (Avastin™; Roche), and Adalimumab (Humira®; AbbVie) in a manner similar to
10 the construction procedure for bispecific antibodies of Example 5.1.2.

The constructed bispecific antibodies Trabev and Adabev, each having the c34Φf51 mutation pair and AWBB mutation pair introduced therein were analyzed using hydrophobic interaction chromatography (HIC) with reference to the method of Example 5.1.2.

The resulting analysis data are given in Table 33 and FIG. 35 (for Trabev) and in Table
15 34 and FIG. 36 (for Adabev).

TABLE 33

# Set1	Antibody	Time (min.)
Aa	Trastuzumab	23.41
Bb	Bevacizumab	38.05
1AB	Trabev	29.13

TABLE 34

# Set2	Antibody	Time (min.)
Aa	Adalimumab	22.07
Bb	Bevacizumab	38.05
2AB	Adabev	30.21

As can be seen in Table 33 and FIG. 35 and in Table 34 and FIG. 36, the peak of each of the heterodimeric bispecific antibodies (Trabev and Adabev) is distinctively observed between
20 the peaks of two corresponding monospecific antibodies (Trastuzumab and Bevacizumab, or Adalimumab and Bevacizumab). The result implies the fine formation of bispecific antibodies

having the c34Φf51 mutation pair and AWBB mutation pair introduced thereto, each composed of halves from two different antibodies.

Further, heterodimization modes of the bispecific antibodies having the c34Φf51 mutation pair and AWBB mutation pair introduced thereto are depicted in FIG. 37 (SDS-PAGE gel 6%, Non-Reducing condition). As can be seen in FIG. 37, concurrent introduction of the c34Φf51 mutation pair and AWBB mutation pair into already known antibodies was found to construct pure heterodimers as analyzed by SDS-PAGE.

5.3. Bispecific antibody having c40Φf44 mutation pair introduced thereto

With reference to Example 5.1.1, antibodies having c40Φf44 mutation pair (see Table 24) introduced to 4D9 (Aa) and 2B9 (Bb) thereof were constructed. For each heavy chain, a and b light chains were co-expressed, and inter-light/heavy chain pairing ratios were measured on SDS-PAGE (conducted for heavy chains A and B, each). T_m was measured as in Example 2, and the results are given in Table 35, below and FIG. 38.

TABLE 35

HC	A (F170D/P171D)		B (F170K/P171K)	
LC	a (L135R/S162K)	b (L135E/S162D)	b (L135E/S162D)	a (L135R/S162K)
q	99%	1%	99%	1%
T _m	65.0	N/A	63.4	N/A

(q: light chain/heavy chain pairing ratio (%); N/A: not available)

As shown in Table 35 and FIG. 38, normal heavy chain/light chain pairs (Aa and Bb) were both formed at a ratio of as high as 99%, and exhibited high thermal stability.

In addition, bispecific antibodies Trabev (Aa: Trastuzumab; Bb: Bevacizumab) and Adabev (Aa: Adalimumab; Bb: Bevacizumab) to each of which the c40Φf44 mutation pair and the AWBB mutation pair (A: S364K/K409W; B: K370S/F405R) were introduced were constructed using Trastuzumab (Herceptin®; Roche), Bevacizumab (Avastin™; Roche), and Adalimumab (Humira®; AbbVie) in a manner similar to the construction procedure for bispecific antibodies of Example 5.1.2.

The constructed bispecific antibodies Trabev and Adabev, each having the c40Φf44 mutation pair and AWBB mutation pair introduced thereto, were analyzed using hydrophobic

interaction chromatography (HIC) with reference to the method of Example 5.1.2.

The resulting analysis data are given in Table 36 and FIG. 39 (for Trabev) and in Table 37 and FIG. 40 (for Adabev).

TABLE 36

# Set1	Antibody	Time (min.)
Aa	Trastuzumab	23.41
Bb	Bevacizumab	38.05
1AB	Trabev	26.11

TABLE 37

# Set2	Antibody	Time (min.)
Aa	Adalimumab	22.07
Bb	Bevacizumab	38.05
2AB	Adabev	31.97

As can be seen in Table 36 and FIG. 39 and in Table 37 and FIG. 40, the peak of each of the heterodimeric bispecific antibodies (Trabev and Adabev) is distinctively observed between the peaks of two corresponding monospecific antibodies (Trastuzumab and Bevacizumab, or Adalimumab and Bevacizumab). The result implies the fine formation of bispecific antibodies having the c40Φf44 mutation pair and AWBB mutation pair introduced thereto, each composed of halves from two different antibodies. In addition, only one peak distinctively appearing between the two monospecific antibodies indicates that only one kind of normal pairing was made without mispairs between two heavy chains and between heavy and light chains.

Further, heterodimerization modes of the bispecific antibodies having the c40Φf44 mutation pair and AWBB mutation pair introduced thereto are depicted in FIG. 41 (SDS-PAGE gel 6%, Non-Reducing condition). As can be seen in FIG. 41, concurrent introduction of the c40Φf44 mutation pair and AWBB mutation pair into already known antibodies was found to construct pure heterodimers as analyzed by SDS-PAGE.

COMPARATIVE EXAMPLE 1

Heterodimerization rates were examined for cases where the mutations of CH3 domain proposed herein, which were different from the mutations suggested in Examples 1 to 3 although

being identical amino acid pairs, were introduced.

For comparison, TNFRSF1B-Fc fusion protein (Enbrel; Enb) and Fas-Fc fusion protein (Fas) to which mutations of Table 38 were introduced were constructed with reference to Example 1 (expressed as BEAT-A and BEAT-B).

TABLE 38

	Enb	Fas
AW/BB	AW (S364K/K409W)	BB (K370S/F405R)
BEAT-A	S364K/K409W	K370T/F405A
BEAT-B	S364T/K409R	K370T/F405S

TNFRSF1B-Fc fusion protein (A chain) and Fas-Fc fusion protein having CH3 domain mutations selected in Example 3, which are representative of CH3 domain mutations proposed in the description (e.g., A chain having S364K and K409W introduced thereto and B chain having K370S and F405R introduced thereto) (expressed as AW/BB in Table 38) were prepared.

When Enb-Fc and Fas-Fc which had the mutation pairs listed in Table 38 were subjected to single transfection, homodimerization modes were observed on SDS-PAGE. The results are given in FIG. 42.

As can be seen in FIG. 42, AW/BB exhibited high percentages of monomers whereas higher percentages of homodimers than monomers were detected in BEAT-A and BEAT-B. On the basis of the result, BEAT-A and BEAT-B are more likely to form homodimers than AW/BB when there is a difference in expression level between A chain and B chain. If a large quantity of homodimers is produced when heavy and light chains are combined so as to construct bispecific antibodies, an accurate heterodimer is difficult to separate from the homodimers because there are almost no differences in physical properties between the heterodimer and the homodimers. Accordingly, minimalizing homodimerization is very important for constructing a bispecific antibody of high purity. A combination of mutations that permits heterodimerization as little as possible is advantageous for the construction of bispecific antibodies.

Taken together, the data show that BEAT-A and BEAT-B have low heterodimerization potentials due to high homodimerization rates compared to AW/BB and make it difficult to isolate accurate heterodimers.

CLAIMS

1. A bispecific protein for targeting two different kinds of targets, the bispecific protein comprising a first CH3 domain or a first Fc region comprising the first CH3 domain and a second CH3 domain or a second Fc region comprising the second CH3 domain, wherein the first CH3 domain and the second CH3 domain are mutated such that at least one selected from among amino acid pairs forming respective amino acid-amino acid bonds between the first CH3 domain and the second CH3 domain is modified by at least one of the following mutations:

(1) a mutation in which the two amino acids in at least one amino acid pair between the CH3 domains are swapped with each other (swapping mutation);

(2) a mutation in which, of at least one amino acid pair between the CH3 domains, one amino acid is substituted with an amino acid having a positive charge while the other is substituted with an amino acid having a negative charge, at least one of the two amino acid residues in the amino acid pair not being hydrophobic (electrostatic interaction-introduced mutation); and

(3) a mutation in which, of at least one amino acid pair between the CH3 domains, one amino acid is substituted with a large hydrophobic amino acid while the other is substituted with a small hydrophobic amino acid (size mutation), wherein the large amino acid is selected from the group consisting of tryptophan and phenylalanine and the small amino acid is selected from the group consisting of alanine, glycine, and valine.

2. The bispecific protein of claim 1, wherein the first CH3 domain and the second CH3 domain are each independently derived from an immunoglobulin selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM.

3. The bispecific protein of claim 1, wherein the amino acid pair forming an amino acid-amino acid bond is at least one selected from amino acid numbers 1 to 40 listed in the following table according to Eu numbering in IgG1-derived CH3 domains or from the group consisting of amino acid pairs at positions corresponding thereto in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

Amino	IgG (1-4)	IgA1	IgA2	IgE	IgM
-------	-----------	------	------	-----	-----

acid pair No.	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B
1	Q347	K360	E347	L360	E347	L360	E444	R457	Q347	K360
2	Y349	S354	H349	P354	H349	P354	Y446	P451	Y349	S354
3	Y349	E357	H349	E357	H349	E357	Y446	P454	Y349	E357
4	Y349	K360	H349	L360	H349	L360	Y446	R457	Y349	K360
5	L351	L351	L351	L351	L351	L351	F448	F448	L351	L351
6	P352	P352	P352	P352	P352	P352	A449	A449	P352	P352
7	S354	Y349	P354	H349	P354	H349	P451	Y446	S354	Y349
8	D356	K439	E356	K439	E356	K439	W453	R539	D356	K439
9	E357	Y349	E357	H349	E357	H349	P454	Y446	E357	Y349
10	E357	K370	E357	R370	E357	R370	P454	Q467	E357	K370
11	K360	Q347	L360	E347	L360	E347	R457	E444	K360	Q347
12	K360	Y349	L360	H349	L360	H349	R457	Y446	K360	Y349
13	S364	L368	T364	L368	T364	L368	T461	L465	S364	L368
14	S364	K370	T364	R370	T364	R370	T461	Q467	S364	K370
15	T366	T366	T366	T366	T366	T366	A463	A463	T366	T366
16	T366	Y407	T366	T407	T366	T407	A463	F506	T366	Y407
17	L368	S364	L368	T364	L368	T364	L465	T461	L368	S364
18	L368	K409	L368	I409	L368	I409	L465	R508	L368	K409
19	K370	E357	R370	E357	R370	E357	Q467	P454	K370	E357
20	K370	S364	R370	T364	R370	T364	Q467	T461	K370	S364
21	K370	T411	R370	R411	R370	R411	Q467	E510	K370	T411
22	N390	S400	L390	S400	L390	S400	R489	K499	N390	S400
23	K392	L398	L392	Q398	L392	Q398	S491	K497	K392	L398
24	T394	T394	W394	W394	W394	W394	T493	T493	T394	T394
25	T394	V397	W394	R397	W394	R397	T493	R496	T394	V397
26	P395	P395	A395	A395	A395	A395	Q494	Q494	P395	P395
27	P395	V397	A395	R397	A395	R397	Q494	R496	P395	V397
28	V397	T394	R397	W394	R397	W394	R496	T493	V397	T394
29	V397	P395	R397	A395	R397	A395	R496	Q494	V397	P395
30	L398	K392	Q398	L392	Q398	L392	K497	S491	L398	K392
31	S400	N390	S400	K390	S400	K390	K499	R489	S400	N390
32	F405	K409	A405	I409	A405	I409	F504	R508	F405	K409
33	Y407	T366	T407	T366	T407	T366	F506	A463	Y407	T366
34	Y407	Y407	T407	T407	T407	T407	F506	F506	Y407	Y407
35	Y407	K409	T407	I409	T407	I409	F506	R508	Y407	K409
36	K409	L368	I409	L368	I409	L368	R508	L465	K409	L368
37	K409	F405	I409	A405	I409	A405	R508	F504	K409	F405
38	K409	Y407	I409	T407	I409	T407	R508	F506	K409	Y407
39	T411	K370	R411	R370	R411	R370	E510	Q467	T411	K370
40	K439	D356	K439	K439	K439	K439	R539	W453	K439	D356

4. The bispecific protein of claim 1, wherein the electrostatic interaction-introduced mutation is a mutation in which, of the two amino acids constituting: at least one amino acid pair selected from the group consisting of serine at position 364 and leucine at position 368, threonine

at position 394 and threonine at position 394, glutamic acid at position 357 and lysine at position 370, glutamic acid at position 357 and tyrosine at position 349, threonine at position 366 and tyrosine at position 407, and threonine at position 394 and valine at position 397 in an IgG1 CH3 domain (EU numbering); or an amino acid pair at a position corresponding to the at least one
5 amino acid pair in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM,

one is substituted with an amino acid having a positive charge and the other is substituted with an amino acid having a negative charge.

5. The bispecific protein of claim 4, wherein the amino acid having a negative charge is
10 aspartic acid or glutamic acid and the amino acid having a positive charge is lysine or arginine.

6. The bispecific protein of claim 4, wherein the electrostatic interaction-introduced mutation comprises at least one of the following mutations on the basis of IgG1 (EU numbering), or at least one of corresponding mutations in the CH3 domains of IgG2, IgG3,
15 IgG4, IgA1, IgA2, IgE, and IgM:

substitution of serine at position 364 with an amino acid having a positive charge and leucine at position 368 with an amino acid having a negative charge;

substitution of threonine at position 394 with an amino acid having a positive charge and threonine at position 394 with an amino acid having a negative charge;

20 substitution of glutamic acid at position 357 with an amino acid having a positive charge and lysine at position 370 with an amino acid having a negative charge;

substitution of glutamic acid at position 357 with an amino acid having a positive charge and tyrosine at position 349 with an amino acid having a negative charge;

25 substitution of threonine at position 366 with an amino acid having a positive charge and tyrosine at position 407 with an amino acid having a negative charge;

substitution of threonine at position 394 with an amino acid having a positive charge and valine at position 397 with an amino acid having a negative charge; and

substitution of tyrosine at position 349 with an amino acid having a positive charge and glutamic acid at position 357 with an amino acid having a negative charge.

30

7. The bispecific protein of claim 6, wherein the amino acid having a negative charge is

aspartic acid or glutamic acid and the amino acid having a positive charge is lysine or arginine.

8. The bispecific protein of claim 1, wherein the swapping mutation is substitution in which exchange is made between two paired amino acid residues in:

- 5 at least one amino acid pair selected from the group consisting of a pair of serine at position 364 and lysine at position 370, a pair of phenylalanine at position 405 and lysine at position 409, a pair of glutamine at position 354 and lysine at position 360, a pair of serine at position 357 and tyrosine at position 349, a pair of glutamic acid at position 357 and lysine at position 370, a pair of lysine at position 360 and tyrosine at position 349, a pair of serine at position 364 and leucine at position 368, a pair of leucine at position 368 and lysine at position 409, a pair of asparagine at position 390 and serine at position 400, a pair of threonine at position 394 and valine at position 397, a pair of leucine at position 398 and lysine at position 392, a pair of tyrosine at position 407 and threonine at position 366, and a pair of threonine at position 411 and lysine at position 370 on the basis of CH3 domain IgG1 (EU numbering); or
- 15 an amino acid pair at a position corresponding to the at least one amino acid pair in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM.

9. The bispecific protein of claim 8, wherein the swapping mutation comprises at least one of the following mutations on the basis of the CH3 domain of IgG1 (EU numbering) or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

- substitution of serine at position 364 with lysine and lysine at position 370 with serine;
substitution of phenylalanine at position 405 with lysine and lysine at position 409 with phenylalanine;
- 25 substitution of tyrosine at position 407 with threonine and threonine at position 366 with tyrosine;
- substitution of glutamic acid at position 357 with lysine and lysine at position 370 with glutamic acid; and
- substitution of serine at position 357 with tyrosine and tyrosine at position 349 with
- 30 serine.

10. The bispecific protein of claim 1, wherein the size mutation comprises substitution in which, of two paired amino acid residues in: at least one amino acid pair selected from the group consisting of a pair of lysine at position 409 and tyrosine at position 407, a pair of lysine at position 409 and phenylalanine at position 405, and a combination thereof on the basis of CH3 domain of IgG1 (EU numbering); or an amino acid pair at a position corresponding to the at least one amino acid pair in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM,

one is substituted with a large hydrophobic amino acid while the other is substituted with a small hydrophobic amino acid wherein the large hydrophobic amino acid is selected from the group consisting of tryptophan and phenylalanine and the small hydrophobic amino acid is selected from the group consisting of alanine, glycine, and valine.

11. The bispecific protein of claim 10, wherein the large amino acid is tryptophan and the small amino acid is alanine.

12. The bispecific protein of claim 10, wherein the size mutation comprises: at least one mutation selected from among the following mutations on the basis of the CH3 domain of IgG1 (EU numbering); and a mutation corresponding the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of lysine at position 409 with tryptophan and tyrosine at position 407 with alanine; and

substitution of lysine at position 409 with tryptophan and phenylalanine at position 405 with alanine.

13. The bispecific protein of claim 1, comprising at least two of the electrostatic interaction-introduced mutation, the swapping mutation, and the size mutation.

14. The bispecific protein of claim 1, comprising:

at least one mutation selected from the group consisting of substitution of one of serine at position 364 and leucine at position 368 with an amino acid having a positive charge and the other with an amino acid having a negative charge, exchange of serine at position 364 with lysine at position 370, and exchange phenylalanine at position 405 with lysine at position 409, on

the basis of the CH3 domain of IgG1 (EU numbering); or

a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM.

5 15. The bispecific protein of claim 14, comprising at least one of the following mutations on the basis of the CH3 of IgG1, or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3 IgG4 IgA1 IgA2 IgD, IgE, and IgM:

(a) substitution serine at position 364 with lysine or arginine, and leucine at position 368 with aspartic acid or glutamic acid;

10 (b) substitution of serine at position 364 with lysine, and lysine at position 370 with serine; and

(c) substitution of phenylalanine at position 405 with lysine, and lysine at position 409 with phenylalanine.

15 16. A bispecific protein for targeting two different kinds of targets, the bispecific protein comprising at least one of the following mutations on the basis of CH3 domain of IgG1 (EU numbering), or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of lysine at position 409 with phenylalanine or tryptophan, and
20 phenylalanine at position 405 with lysine, arginine, glutamine, or asparagine;

substitution of leucine at position 368 with aspartic acid, glutamic acid, or for example, aspartic acid), and serine at position 364 with lysine, arginine, or asparagine; and

substitution of lysine at position 370 with serine, and serine at position 364 with lysine, arginine, or asparagine.

25

17. The bispecific protein of claim 16, comprising at least one of the following mutations on the basis of the CH3 domain of IgG1 (EU numbering), or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of phenylalanine at position 405 with arginine, and lysine at position 409
30 with tryptophan;

substitution of serine at position 364 with lysine, and leucine at position 368 with

aspartic acid;

substitution of serine at position 364 with lysine, and lysine at position 370 with serine;

substitution of phenylalanine at position 405 with lysine, and lysine at position 409 with phenylalanine;

5 substitution of phenylalanine at position 405 with arginine, and lysine at position 409 with phenylalanine; and

substitution of phenylalanine at position 405 with lysine, and lysine at position 409 with tryptophan.

10 18. The bispecific protein of claim 17, comprising at least one of the following mutations on the basis of the CH3 domain of IgG1 (EU numbering), or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 15 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second 20 CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with lysine;

25 substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at 30 position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second

CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second

5 CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with lysine; or

10 substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with arginine.

15 19. The bispecific protein of one of claims 1 to 18, wherein bispecific protein is at least one selected from the group consisting of a bispecific antibody; an antigen-binding fragment of a bispecific antibody, a bispecific antibody analog, and a fusion protein of a target-specific binding polypeptide and an Fc region or CH3 domain.

20 20. The bispecific protein of claim 19, wherein the target-specific binding polypeptide is selected from the group consisting of a receptor, an ectodomain, and a ligand.

21. A bispecific antibody or an antigen-binding fragment thereof, the bispecific antibody comprising a first CH1 domain and a first CL (light chain constant region) domain derived
25 respectively from the heavy chain and light chain of an antibody recognizing a first epitope and a second CH1 domain and a second CL domain derived respectively from the heavy chain and light chain of an antibody recognizing a second epitope, wherein the CH1 domains and the CL domains are mutated to contain at least one of the following mutations:

a mutation in which, of the two amino acids constituting each pair of one or more first
30 amino acid pairs selected from among amino acid pairs forming respective amino acid-amino acid bonds between the first CH1 domain and the first CL domain, one is substituted with an

amino acid having a positive charge and the other is substituted with an amino acid having a negative charge; and

a mutation in which, of the two amino acids constituting each pair of one or more second amino acid pairs selected from among amino acid pairs forming respective amino acid-amino acid bonds between the second CH1 domain and the second CL domain, one is substituted with an amino acid having a positive charge and the other is substituted with an amino acid having a negative charge.

22. The bispecific antibody or the antigen-binding fragment of claim 21, wherein:
the amino acids substituted respectively in the first CH1 domain and the second CH1 domain have opposite charges,
the amino acids substituted respectively in the first CL domain and the first CH1 domain have opposite charges, and
the amino acids substituted respectively in the second CL domain and the second CH1 domain have opposite charges.

23. The bispecific antibody or antigen-binding fragment of claim 21, wherein the amino acid having a positive charge is lysine or arginine, the amino acid having a negative charge is aspartic acid or glutamic acid, and the amino acid to be substituted with an amino acid having a positive or negative charge in the CH1 domain is at least one of the following amino acids on the basis of the CH1 domain of IgG1 (EU numbering), or an amino acid at a position corresponding to the at least one amino acid in CH1 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

leucine at position 145,
serine at position 183,
lysine at position 147,
phenylalanine at position 170,
proline at position 171, and
valine at position 185, and

the amino acid to be substituted with an amino acid having a positive or negative charge in the CL domain is at least one of the following amino acids on the basis of the CL domain of kappa type (EU numbering), or an amino acid at a position corresponding to the at least one

amino acid in the CL domain of lambda type:

serine at position 131,
valine at position 133,
leucine at position 135,
5 serine at position 162, and
threonine at position 180.

24. The bispecific antibody or antigen-binding fragment of claim 21, wherein
the amino acid having a positive charge is lysine or arginine,
10 the amino acid having a negative charge is aspartic acid or glutamic acid, and
a set of two amino acids forming an amino acid pair between the CH1 domain and the
CL domain is at least one of the following amino acid pairs on the basis of the CH1 domain of
IgG1 and the CL domain of kappa type (EU numbering), or an amino acid pair at a position
corresponding to the at least one amino acid pair in CH1 domains and CL domain of lambda
15 type of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

a pair of leucine at position 145 in the CH1 domain and serine at position 131 in the CL
domain,
a pair of leucine at position 145 in the CH1 domain and valine at position 133 in the CL
domain,
20 a pair of serine at position 183 in the CH1 domain and valine at position 133 in the CL
domain,
a pair of lysine at position 147 in the CH1 domain and threonine at position 180 in the
CL domain,
a pair of valine at position 185 in the CH1 domain and leucine at position 135 in the CL
25 domain,
a pair of phenylalanine at position 170 in the CH1 domain and leucine at position 135 in
the CL domain, and
a pair of proline at position 171 in the CH1 domain and serine at position 162 in the CL
domain.

30
25. The bispecific antibody or antigen-binding fragment of one of claims 21 to 24,

wherein the bispecific antibody comprises modified CH3 domains or a mutant Fc region inclusive of the modified CH3 domains, the modified CH3 domains comprising a first CH3 domain and a second CH3 domain and being mutated such that at least one selected from amino acid pairs forming amino acid-amino acid bonds between the first CH3 domain derived from the heavy chain of an antibody recognizing a first epitope and the second CH3 domain derived from the heavy chain of an antibody recognizing a second epitope has at least one of the following mutations:

a mutation in which amino acids in at least one amino acid pair between the CH3 domains are switched with each other (swapping mutation);

a mutation in which, of at least one amino acid pair between the CH3 domains, one amino acid is substituted with an amino acid having a positive charge while the other is substituted with an amino acid having a negative charge, at least one of the two amino acid residues in the amino acid pair not being hydrophobic (electrostatic interaction-introduced mutation); and

a mutation in which, of at least one amino acid pair between the CH3 domains, one amino acid is substituted with a large hydrophobic amino acid while the other is substituted with a small hydrophobic amino acid (size mutation), wherein the large amino acid is selected from the group consisting of tryptophan and phenylalanine and the small amino acid is selected from the group consisting of alanine, glycine, and valine.

26. The bispecific antibody or antigen-binding fragment of claim 25, wherein the first CH3 domain and the second CH3 domain are each independently derived from an immunoglobulin selected from the group consisting of IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM.

27. The bispecific antibody or antigen-binding fragment of claim 25, wherein the amino acid pair forming an amino acid-amino acid bond is at least one selected from amino acid numbers 1 to 40 listed in the following table according to Eu numbering in IgG1-derived CH3 domains or from the group consisting of amino acid pairs at positions corresponding thereto in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

Amino	IgG (1-4)	IgA1	IgA2	IgE	IgM
-------	-----------	------	------	-----	-----

acid pair No.	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B	Chain A	Chain B
1	Q347	K360	E347	L360	E347	L360	E444	R457	Q347	K360
2	Y349	S354	H349	P354	H349	P354	Y446	P451	Y349	S354
3	Y349	E357	H349	E357	H349	E357	Y446	P454	Y349	E357
4	Y349	K360	H349	L360	H349	L360	Y446	R457	Y349	K360
5	L351	L351	L351	L351	L351	L351	F448	F448	L351	L351
6	P352	P352	P352	P352	P352	P352	A449	A449	P352	P352
7	S354	Y349	P354	H349	P354	H349	P451	Y446	S354	Y349
8	D356	K439	E356	K439	E356	K439	W453	R539	D356	K439
9	E357	Y349	E357	H349	E357	H349	P454	Y446	E357	Y349
10	E357	K370	E357	R370	E357	R370	P454	Q467	E357	K370
11	K360	Q347	L360	E347	L360	E347	R457	E444	K360	Q347
12	K360	Y349	L360	H349	L360	H349	R457	Y446	K360	Y349
13	S364	L368	T364	L368	T364	L368	T461	L465	S364	L368
14	S364	K370	T364	R370	T364	R370	T461	Q467	S364	K370
15	T366	T366	T366	T366	T366	T366	A463	A463	T366	T366
16	T366	Y407	T366	T407	T366	T407	A463	F506	T366	Y407
17	L368	S364	L368	T364	L368	T364	L465	T461	L368	S364
18	L368	K409	L368	I409	L368	I409	L465	R508	L368	K409
19	K370	E357	R370	E357	R370	E357	Q467	P454	K370	E357
20	K370	S364	R370	T364	R370	T364	Q467	T461	K370	S364
21	K370	T411	R370	R411	R370	R411	Q467	E510	K370	T411
22	N390	S400	L390	S400	L390	S400	R489	K499	N390	S400
23	K392	L398	L392	Q398	L392	Q398	S491	K497	K392	L398
24	T394	T394	W394	W394	W394	W394	T493	T493	T394	T394
25	T394	V397	W394	R397	W394	R397	T493	R496	T394	V397
26	P395	P395	A395	A395	A395	A395	Q494	Q494	P395	P395
27	P395	V397	A395	R397	A395	R397	Q494	R496	P395	V397
28	V397	T394	R397	W394	R397	W394	R496	T493	V397	T394
29	V397	P395	R397	A395	R397	A395	R496	Q494	V397	P395
30	L398	K392	Q398	L392	Q398	L392	K497	S491	L398	K392
31	S400	N390	S400	K390	S400	K390	K499	R489	S400	N390
32	F405	K409	A405	I409	A405	I409	F504	R508	F405	K409
33	Y407	T366	T407	T366	T407	T366	F506	A463	Y407	T366
34	Y407	Y407	T407	T407	T407	T407	F506	F506	Y407	Y407
35	Y407	K409	T407	I409	T407	I409	F506	R508	Y407	K409
36	K409	L368	I409	L368	I409	L368	R508	L465	K409	L368
37	K409	F405	I409	A405	I409	A405	R508	F504	K409	F405
38	K409	Y407	I409	T407	I409	T407	R508	F506	K409	Y407
39	T411	K370	R411	R370	R411	R370	E510	Q467	T411	K370
40	K439	D356	K439	K439	K439	K439	R539	W453	K439	D356

28. The bispecific antibody or antigen-binding fragment of claim 25, wherein the electrostatic interaction-introduced mutation is a mutation in which, of the two amino acids constituting: at least one amino acid pair selected from the group consisting of serine at position

364 and leucine at position 368, threonine at position 394 and threonine at position 394, glutamic acid at position 357 and lysine at position 370, glutamic acid at position 357 and tyrosine at position 349, threonine at position 366 and tyrosine at position 407, and threonine at position 394 and valine at position 397 in an IgG1 CH3 domain (EU numbering); or an amino acid pair at a position corresponding to the at least one amino acid pair in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM,

one is substituted with an amino acid having a positive charge and the other is substituted with an amino acid having a negative charge.

29. The bispecific antibody or antigen-binding fragment of claim 28, wherein the amino acid having a negative charge is aspartic acid or glutamic acid and the amino acid having a positive charge is lysine or arginine.

30 The bispecific antibody or antigen-binding fragment of claim 28, wherein the electrostatic interaction-introduced mutation comprises at least one of the following mutations on the basis of IgG1 (EU numbering), or at least one of corresponding mutations in the CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of serine at position 364 with an amino acid having a positive charge and leucine at position 368 with an amino acid having a negative charge;

substitution of threonine at position 394 with an amino acid having a positive charge and threonine at position 394 with an amino acid having a negative charge;

substitution of glutamic acid at position 357 with an amino acid having a positive charge and lysine at position 370 with an amino acid having a negative charge;

substitution of glutamic acid at position 357 with an amino acid having a positive charge and tyrosine at position 349 with an amino acid having a negative charge;

substitution of threonine at position 366 with an amino acid having a positive charge and tyrosine at position 407 with an amino acid having a negative charge;

substitution of threonine at position 394 with an amino acid having a positive charge and valine at position 397 with an amino acid having a negative charge; and

substitution of tyrosine at position 349 with an amino acid having a positive charge and glutamic acid at position 357 with an amino acid having a negative charge.

31. The bispecific antibody or antigen-binding fragment of claim 30, wherein the amino acid having a negative charge is aspartic acid or glutamic acid and the amino acid having a positive charge is lysine or arginine.

5

32. The bispecific antibody or antigen-binding fragment of claim 25, wherein the swapping mutation is substitution in which exchange is made between two paired amino acid residues in:

at least one amino acid pair selected from the group consisting of a pair of glutamine at
10 position 354 and lysine at position 360, a pair of serine at position 357 and tyrosine at position 349, a pair of glutamic acid at position 357 and lysine at position 370, a pair of lysine at position 360 and tyrosine at position 349, a pair of serine at position 364 and leucine at position 368, a pair of serine at position 364 and lysine at position 370, a pair of leucine at position 368 and lysine at position 409, a pair of asparagine at position 390 and serine at position 400, a pair of
15 threonine at position 394 and valine at position 397, a pair of leucine at position 398 and lysine at position 392, a pair of phenylalanine at position 405 and lysine at position 409, a pair of tyrosine at position 407 and threonine at position 366, and a pair of threonine at position 411 and lysine at position 370 on the basis of CH3 domain IgG1 (EU numbering); or

an amino acid pair at a position corresponding to the at least one amino acid pair in CH3
20 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM.

33. The bispecific antibody or antigen-binding fragment of claim 32, wherein the swapping mutation comprises at least one of the following mutations on the basis of the CH3 domain of IgG1 (EU numbering) or a mutation corresponding to the at least one mutation in
25 CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of serine at position 364 with lysine and lysine at position 370 with serine;
substitution of phenylalanine at position 405 with lysine and lysine at position 409 with phenylalanine;
substitution of tyrosine at position 407 with threonine and threonine at position 366 with
30 tyrosine;
substitution of glutamic acid at position 357 with lysine and lysine at position 370 with

glutamic acid; and

substitution of serine at position 357 with tyrosine and tyrosine at position 349 with serine.

5 34. The bispecific antibody or antigen-binding fragment of claim 25, wherein the size mutation comprises substitution in which, of two paired amino acid residues in: at least one amino acid pair selected from the group consisting of a pair of lysine at position 409 and tyrosine at position 407, a pair of lysine at position 409 and phenylalanine at position 405, and a combination thereof on the basis of CH3 domain of IgG1 (EU numbering); or an amino acid
10 pair at a position corresponding to the at least one amino acid pair in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM,

one is substituted with a large hydrophobic amino acid while the other is substituted with a small hydrophobic amino acid wherein the large hydrophobic amino acid is selected from the group consisting of tryptophan and phenylalanine and the small hydrophobic amino acid is
15 selected from the group consisting of alanine, glycine, and valine.

35. The bispecific antibody or antigen-binding fragment of claim 34, wherein the large amino acid is tryptophan and the small amino acid is alanine.

20 36. The bispecific antibody or antigen-binding fragment of claim 34, wherein the size mutation comprises: at least one mutation selected from among the following mutations on the basis of the CH3 domain of IgG1 (EU numbering); and a mutation corresponding the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of lysine at position 409 with tryptophan and tyrosine at position 407 with
25 alanine; and

substitution of lysine at position 409 with tryptophan and phenylalanine at position 405 with alanine.

37. The bispecific antibody or antigen-binding fragment of claim 25, wherein the
30 bispecific antibody comprises at least two of the electrostatic interaction-introduced mutation, the swapping mutation, and the size mutation.

38. The bispecific antibody or antigen-binding fragment of claim 25, wherein the bispecific antibody comprises:

at least one mutation selected from the group consisting of substitution of one of serine at
5 position 364 and leucine at position 368 with an amino acid having a positive charge and the
other with an amino acid having a negative charge, exchange of serine at position 364 with
lysine at position 370, and exchange phenylalanine at position 405 with lysine at position 409, on
the basis of the CH3 domain of IgG1 (EU numbering); or

a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3,
10 IgG4, IgA1, IgA2, IgE, and IgM.

39. The bispecific antibody or antigen-binding fragment of claim 38, wherein the
bispecific antibody comprises at least one of the following mutations on the basis of the CH3
domain of IgG1, or a mutation corresponding to the at least one mutation in CH3 domains of
15 IgG2, IgG3 IgG4 IgA1 IgA2 IgD, IgE, and IgM:

(a) substitution of serine at position 364 with lysine or arginine, and leucine at position
368 with aspartic acid or glutamic acid;

(b) substitution of serine at position 364 with lysine, and lysine at position 370 with
serine; and

20 (c) substitution of phenylalanine at position 405 with lysine, and lysine at position 409
with phenylalanine.

40. The bispecific antibody or antigen-binding fragment of claim 25, wherein the
bispecific protein comprises at least one of the following mutations on the basis of CH3 domain
25 of IgG1 (EU numbering), or a mutation corresponding to the at least one mutation in CH3
domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of lysine at position 409 with phenylalanine or tryptophan, and
phenylalanine at position 405 with lysine, arginine, glutamine, or asparagine;

substitution of leucine at position 368 with aspartic acid, glutamic acid, or aspartic acid,
30 and serine at position 364 with lysine, arginine, or asparagine; and

substitution of lysine at position 370 with serine, and serine at position 364 with lysine,

arginine, or asparagine.

41. The bispecific antibody or antigen-binding fragment of claim 40, wherein the bispecific antibody comprises at least one of the following mutations on the basis of the CH3 domain of IgG1 (EU numbering), or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of phenylalanine at position 405 with arginine, and lysine at position 409 with tryptophan;

substitution of serine at position 364 with lysine, and leucine at position 368 with aspartic acid;

substitution of serine at position 364 with lysine, and lysine at position 370 with serine;

substitution of phenylalanine at position 405 with lysine, and lysine at position 409 with phenylalanine;

substitution of phenylalanine at position 405 with arginine, and lysine at position 409 with phenylalanine; and

substitution of phenylalanine at position 405 with lysine, and lysine at position 409 with tryptophan.

42. The bispecific antibody or antigen-binding fragment of claim 41, wherein the bispecific antibody comprises at least one of the following mutations on the basis of the CH3 domain of IgG1 (EU numbering), or a mutation corresponding to the at least one mutation in CH3 domains of IgG2, IgG3, IgG4, IgA1, IgA2, IgE, and IgM:

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at

position 368 in the second CH3 domain with aspartic acid, and substitution of lysine in the first CH3 domain at position 409 with phenylalanine and phenylalanine at position 405 in the second CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at
5 position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at
position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in
10 the first CH3 domain with phenylalanine and phenylalanine at position 405 in the second CH3 domain with arginine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at
position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position
409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second
15 CH3 domain with lysine;

substitution of serine at position 364 in the first CH3 domain with lysine and leucine at
position 368 in the second CH3 domain with aspartic acid, and substitution of lysine at position
409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second
CH3 domain with arginine; or

substitution of serine at position 364 in the first CH3 domain with lysine and lysine at
20 position 370 in the second CH3 domain with serine, and substitution of lysine at position 409 in the first CH3 domain with tryptophan and phenylalanine at position 405 in the second CH3 domain with lysine.

25 43. A method for constructing a bispecific protein for targeting different targets, the method comprising one of the following mutation introducing steps to introduce at least one mutation into at least one selected from amino acid pairs forming amino acid-amino acid bonds between a first CH3 domain and a second CH3 domain:

switching the two amino acids in at least one amino acid pair selected from amino acid
30 pairs forming amino-amino acid bonds between a first CH3 domain and a second CH3 domain with each other;

substituting one of the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between the CH3 domains with an amino acid having a positive charge and the other with an amino acid having a negative charge; and

substituting one of the two amino acids in at least one amino acid pair selected from
5 amino acid pairs forming amino-amino acid bonds between the CH3 domains with a large hydrophobic amino acid and the other with a small hydrophobic amino acid, wherein the large amino acid is selected from the group consisting of tryptophan and phenylalanine and the small amino acid is selected from the group consisting of alanine, glycine, and valine.

10 44. A method for constructing a bispecific antibody or an antigen-binding fragment thereof, comprising the following CH1 and CL domain mutating steps of:

substituting one of the two amino acid residues constituting at least one selected from amino acid pairs forming amino acid-amino acid bonds between a first CH1 domain derived from the heavy chain and a first CL domain of an antibody recognizing a first epitope with an
15 amino acid having a positive charge and the other with an amino acid having a negative charge; and

substituting one of the two amino acid residues constituting at least one selected from amino acid pairs forming amino acid-amino acid bonds between a second CH1 domain derived from the heavy chain and a second CL domain of an antibody recognizing a second epitope with
20 an amino acid having a positive charge and the other with an amino acid having a negative charge.

45. The method of claim 44, further comprising at least one of the following CH3 domain mutating steps:

25 switching the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between a first CH3 domain and a second CH3 domain with each other;

substituting one of the two amino acids in at least one selected from amino acid pairs forming amino-amino acid bonds between the CH3 domains with an amino acid having a
30 positive charge and the other with an amino acid having a negative charge; and

substituting one of the two amino acid residues constituting at least one selected from

amino acid pairs forming amino acid-amino acid bonds between a first CH3 domain derived from the heavy chain of an antibody recognizing a first epitope and a second CH3 domain derived from the heavy chain of an antibody recognizing a second epitope with an amino acid having a positive charge and the other with an amino acid having a negative charge; and

5 substituting one of the two amino acids in at least one amino acid pair selected from amino acid pairs forming amino-amino acid bonds between the first CH3 domain and the second CH3 domain with a large hydrophobic amino acid and the other with a small hydrophobic amino acid, wherein the large amino acid is selected from the group consisting of tryptophan and phenylalanine and the small amino acid is selected from the group consisting of
10 alanine, glycine, and valine.

46. An anti-influenza B antibody or an antigen-binding fragment thereof, the antibody comprising the heavy chain variable region of SEQ ID NO: 27 and the light chain variable region of SEQ ID NO: 31.

15

47. An anti-influenza A antibody or an antigen-binding fragment thereof, the antibody comprising the heavy chain variable region of SEQ ID NO: 29 and the light chain variable region of SEQ ID NO: 31.

20 48. An anti-influenza A/anti-influenza B bispecific antibody or an antigen-binding fragment thereof, the bispecific antibody comprising an anti-influenza B antibody comprising the heavy chain variable region of SEQ ID NO: 27 and the light chain variable region of SEQ ID NO: 31; and an anti-influenza A antibody comprising the heavy chain variable region of SEQ ID NO: 29 and the light chain variable region of SEQ ID NO: 31.

25

49. The bispecific antibody or antigen-binding fragment of one of claims 21 to 24, wherein

the antibody recognizing a first epitope is an anti-influenza B antibody comprising the heavy chain variable region of SEQ ID NO: 27 and the light chain variable region of SEQ ID
30 NO: 31; and

the antibody recognizing a second epitope is an anti-influenza A antibody comprising

the heavy chain variable region of SEQ ID NO: 29 and the light chain variable region of SEQ ID NO: 31.

50. The bispecific antibody or antigen-binding fragment of claim 25, wherein
- 5 the antibody recognizing a first epitope is an anti-influenza B antibody comprising the heavy chain variable region of SEQ ID NO: 27 and the light chain variable region of SEQ ID NO: 31; and
- the antibody recognizing a second epitope is an anti-influenza A antibody comprising the heavy chain variable region of SEQ ID NO: 29 and the light chain variable region of SEQ ID
- 10 NO: 31.