



(51) International Patent Classification:

C07D 513/18 (2006.01) A61K 31/55 (2006.01)
C07D 515/18 (2006.01) A61P 31/14 (2006.01)

(21) International Application Number:

PCT/EP2009/004942

(22) International Filing Date:

8 July 2009 (08.07.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

08159965.6	8 July 2008 (08.07.2008)	EP
08160254.2	11 July 2008 (11.07.2008)	EP
08161743.3	4 August 2008 (04.08.2008)	EP

(71) Applicant (for all designated States except US): **TIBOTEC PHARMACEUTICALS** [IE/IE]; Eastgate Village, Eastgate Little Island, County Cork (IE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VENDEVILLE, Sandrine, Marie, Hélène** [FR/BE]; c/o Tibotec-Virco Virology BVBA, Generaal De Wittelaan L11B3, B-2800 Mechelen (BE). **RABOISSON, Pierre, Jean-Marie, Bernard** [FR/BE]; c/o Tibotec-Virco Virology BVBA, Generaal De Wittelaan L11B3, B-2800 Mechelen (BE). **LIN, Tse-I** [DE/BE]; c/o Tibotec-Virco Virology BVBA, Generaal De Wittelaan L11B3, B-2800 Mechelen (BE).

(74) Agent: **DAELEMANS, Frank**; Tibotec-Virco Comm. VA, J&J Patent Law Department, Generaal De Wittelaan L 11B 3, B-2800 Mechelen (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(54) Title: MACROCYCLIC INDOLE DERIVATIVES USEFUL AS HEPATITIS C VIRUS INHIBITORS



(57) Abstract: Inhibitors of HCV replication of formula (I) including stereochemically isomeric forms, and salts, hydrates, solvates thereof, wherein R¹, R², R⁴, R⁵, R⁶ and R⁷ have the meaning defined in the claims. The present invention also relates to processes for preparing said compounds, pharmaceutical compositions containing them and their use in HCV therapy.

MACROCYCLIC INDOLE DERIVATIVES USEFUL AS HEPATITIS C VIRUS INHIBITORS

Field of the invention

The present invention is concerned with macrocyclic indole derivatives having
5 inhibitory activity on the replication of the hepatitis C virus (HCV). It further concerns compositions comprising these compounds as active ingredients as well as processes for preparing these compounds and compositions.

Background of the invention

Hepatitis C virus is the leading cause of chronic liver disease worldwide and has
10 become a focus of considerable medical research. HCV is a member of the *Flaviviridae* family of viruses in the *hepacivirus* genus, and is closely related to the *flavivirus* genus, which includes a number of viruses implicated in human disease, such as dengue virus and yellow fever virus, and to the animal *pestivirus* family, which includes bovine viral diarrhoea virus (BVDV). HCV is a positive-sense, single-stranded RNA virus, with a
15 genome of around 9,600 bases. The genome comprises both 5' and 3' untranslated regions that adopt RNA secondary structures, and a central open reading frame that encodes a single polyprotein of around 3,010-3,030 amino acids. The polyprotein encodes ten gene products, which are generated from the precursor polyprotein by an orchestrated series of co- and posttranslational endoproteolytic cleavages mediated by
20 both host and viral proteases. The viral structural proteins include the core nucleocapsid protein, and two envelope glycoproteins E1 and E2. The non-structural (NS) proteins encode some essential viral enzymatic functions (helicase, polymerase, protease), as well as proteins of unknown function. Replication of the viral genome is mediated by an RNA-dependent RNA polymerase, encoded by non-structural protein 5b (NS5B). In
25 addition to the polymerase, the viral helicase and protease functions, both encoded in the bifunctional NS3 protein, have been shown to be essential for replication of HCV RNA. In addition to the NS3 serine protease, HCV also encodes a metalloproteinase in the NS2 region.

HCV replicates preferentially in hepatocytes but is not directly cytopathic, leading to
30 persistent infection. In particular, the lack of a vigorous T-lymphocyte response and the high propensity of the virus to mutate appear to promote a high rate of chronic infection. There are 6 major HCV genotypes and more than 50 subtypes, which are differently distributed geographically. HCV type 1 is the predominant genotype in the US and Europe. For instance, HCV type 1 accounts for 70 to 75 percent of all HCV
35 infections in the United States. The extensive genetic heterogeneity of HCV has

- important diagnostic and clinical implications, perhaps explaining difficulties in vaccine development and the lack of response to therapy. An estimated 170 million persons worldwide are infected with hepatitis C virus (HCV). Following the initial acute infection, a majority of infected individuals develops chronic hepatitis, which can
- 5 progress to liver fibrosis leading to cirrhosis, end-stage liver disease, and HCC (hepatocellular carcinoma) (National Institutes of Health Consensus Development Conference Statement: Management of Hepatitis C. *Hepatology*, 36, 5 Suppl. S3-S20, 2002). Liver cirrhosis due to HCV infection is responsible for about 10,000 deaths per year in the U.S.A. alone, and is the leading cause for liver transplantations.
- 10 Transmission of HCV can occur through contact with contaminated blood or blood products, for example following blood transfusion or intravenous drug use. The introduction of diagnostic tests used in blood screening has led to a downward trend in post-transfusion HCV incidence. However, given the slow progression to the end-stage liver disease, the existing infections will continue to present a serious medical and
- 15 economic burden for decades (Kim, W.R. *Hepatology*, 36, 5 Suppl. S30-S34, 2002).
- Current HCV therapies are based on (pegylated) interferon-alpha (IFN- α) in combination with ribavirin. This combination therapy yields a sustained virologic response in more than 40% of patients infected by genotype 1 viruses and about 80% of those infected by genotypes 2 and 3. Beside the limited efficacy on HCV type 1,
- 20 combination therapy has significant side effects and is poorly tolerated in many patients. For instance, in registration trials of pegylated interferon and ribavirin, significant side effects resulted in discontinuation of treatment in approximately 10 to 14 percent of patients. Major side effects of combination therapy include influenza-like symptoms, hematologic abnormalities, and neuropsychiatric symptoms. The
- 25 development of more effective, convenient and tolerated treatments is a major public health objective. Thus, the treatment of this chronic disease is an unmet clinical need, since current therapy is only partially effective and limited by undesirable side effects.
- One area of particular focus has been the search for inhibitors of the NS5b RNA-dependent RNA polymerase (RdRp). Close structural homologs of this
- 30 polymerase do not exist within the uninfected host cell and the finding of inhibitors of said polymerase would provide a more specific mode of action. Inhibitors that are currently under investigation can be classified as either nucleoside inhibitors (NIs) or non-nucleoside inhibitors (NNIs). NIs directly compete with nucleotide substrates for binding to highly conserved active sites. Greater specificity may be achieved by NNIs,
- 35 which may interact outside of the highly conserved active site at a unique allosteric site common only to structurally related polymerases.


Indole derivatives have been described for HCV inhibitory activity. WO 2007/092000 discloses tetracyclic indole derivatives as HCV NS5B inhibitors for the treatment and/or prevention of HCV virus infection. US 2008/0146537 discloses cyclopropyl fused indolobenzazepine HCV NS5B inhibitors. WO 2008/075103 discloses
 5 macrocyclic indole derivatives useful for the treatment or prevention of infection by hepatitis C virus.

To date, preliminary clinical trials have resulted in a high failure rate, thereby highlighting the need to pursue the search for novel NS5b inhibitors. There is a high medical need for safe and effective anti-HCV treatment. Such HCV inhibitors may
 10 overcome the disadvantages of current HCV therapy such as side effects, limited efficacy, the emergence of resistance, and compliance failures, as well as improve the sustained viral response. In particular wherein the therapeutic compounds have good bioavailability and a favorable pharmacokinetic and metabolic profile.

Summary


15 It has been found that certain macrocyclic indole derivatives exhibit antiviral activity in subjects infected with HCV with useful properties regarding one or more of the following parameters: antiviral efficacy, favorable mutant prophile, lack of toxicity, favorable pharmacokinetic and metabolic profile, and ease of formulation and administration. These compounds are therefore useful in treating or combating HCV
 20 infections.


The present disclosure concerns inhibitors of HCV replication. In a first aspect, there is provided a compound of formula (I),



25 including stereochemically isomeric forms, and N-oxides, salts, hydrates, and solvates thereof, wherein:

– R^1 is a bivalent chain selected from





- each R^3 is independently selected from the group comprising hydrogen, C_{1-4} alkyl and C_{3-5} cycloalkyl;
 - a is 3, 4, 5 or 6;
 - each b is independently 1 or 2;
 - c is 1 or 2;
 - macrocycle A has 14 to 18 member atoms;
 - each R^2 is independently hydrogen, halo or C_{1-4} alkoxy;
 - R^4 and R^5 are hydrogen or R^4 and R^5 together form a double bond or a methylene group to form a fused cyclopropyl;
 - R^6 is hydrogen or methyl; and
 - R^7 is a C_{3-7} cycloalkyl optionally substituted with halo.
- 10
- 15

The disclosure further relates to methods for the preparation of the compounds of formula (I), including stereochemically isomeric forms, and *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof, their intermediates, and the use of the intermediates in the preparation of the compounds of formula (I).

20

In a second aspect, there is provided compounds of formula (I) *per se*, including stereochemically isomeric forms, and *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof, for use as a medicament. In a third aspect, there is provided a compound as defined above or a pharmaceutical composition as defined above, for inhibiting HCV replication. The disclosure relates to the compounds of formula (I) *per se*, including stereochemically isomeric forms, and *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof, for treating hepatitis C. In a fourth aspect, there is provided pharmaceutical compositions comprising a carrier and as active ingredient an anti-virally effective amount of a compound of formula (I) as specified herein. The pharmaceutical compositions may comprise combinations of the aforementioned compounds with other anti-HCV agents. The pharmaceutical compositions may comprise combinations of the aforementioned compounds with anti-HIV agents. The disclosure further relates to the aforementioned pharmaceutical compositions for administration to a subject suffering from HCV infection.

In a fifth aspect, there is provided use of a compound of formula (I), including stereochemically isomeric forms, or *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof, for the manufacture of a medicament for inhibiting HCV replication. The disclosure also relates to the use of a compound of formula (I), including stereochemically isomeric forms, or *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof, for the manufacture of a medicament for preventing or treating conditions associated with HCV. The disclosure also relates to a method of inhibiting HCV replication in a warm-blooded animal said method comprising the administration of an effective amount of a compound of formula (I), including stereochemically isomeric forms, or *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof. The disclosure also relates to a method for preventing or treating conditions associated with HCV in a warm-blooded animal said method comprising the administration of an effective amount of a compound of formula (I), including stereochemically isomeric forms, or *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof.

Detailed description

The present invention will now be further described. In the following passages, different aspects or embodiments of the invention are defined in more detail. Each aspect or embodiment so defined may be combined with any other aspect(s) or embodiment(s) unless clearly indicated to the contrary. In particular, any feature indicated as being preferred or advantageous may be combined with any other feature or features indicated as being preferred or advantageous to formulate a particular embodiment.

As used in the foregoing and hereinafter, the following definitions apply unless otherwise noted.

For the purpose of the present invention, the terms "subject" or "infected subject" or "patient" refers to an individual infected with HCV, in need of treatment.

- 5 The term "halo" or "halogen" is generic to fluoro, chloro, bromo and iodo.

- As used herein "C₁₋₄alkyl" as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as for example methyl, ethyl, prop-1-yl, prop-2-yl, but-1-yl, but-2-yl, isobutyl, 2-methylprop-1-yl; "C₁₋₃alkyl" as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 3 carbon atoms such as for example methyl, ethyl, prop-1-yl, prop-2-yl.
- 10

- The term "C₁₋₆alkylene" as a group or part of a group refers to C₁₋₆alkyl groups that are divalent, i.e., with two single bonds for attachment to two other groups. Non-limiting examples of alkylene groups includes methylene, ethylene, methylenemethylene, propylene, ethylethylene, 1-methylethylene and 1,2-dimethylethylene.
- 15

"C₃₋₇cycloalkyl" is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The term "C₃₋₅cycloalkyl" is meant to comprise cyclopropyl, cyclobutyl and cyclopentyl.

- The term "C₁₋₄alkoxy" or "C₁₋₄alkyloxy" as a group or part of a group refers to a radical having the Formula -OR^a wherein R^a is C₁₋₄alkyl as defined above. Non-limiting examples of suitable C₁₋₄alkoxy include methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, *sec*-butoxy and *tert*-butoxy.
- 20

It should be noted that the radical positions on any molecular moiety used in the definitions may be anywhere on such moiety as long as it is chemically stable.

- 25 Radicals used in the definitions of the variables include all possible isomers unless otherwise indicated. For instance, piperidinyl includes piperidin-1-yl, piperidin-2-yl, piperidin-3-yl, and piperidin-4-yl; pentyl includes pent-1-yl, pent-2-yl and pent-3-yl.

When any variable occurs more than one time in any constituent, each definition is independent.

- 30 Whenever used hereinafter, the term "compounds of formula (I)", or "the present compounds" or similar terms, it is meant to include the compounds of formula (I), including stereochemically isomeric forms, and their *N*-oxides, quaternary amines, metal complexes, salts, hydrates or solvates thereof. One embodiment comprises the compounds of formula (I) or any subgroup thereof specified herein, including the


possible stereochemically isomeric forms, as well as the *N*-oxides, salts, hydrates, and solvates thereof. Another embodiment comprises the compounds of formula (I) or any subgroups thereof specified herein, including the possible stereochemically isomeric forms, as well as the *N*-oxides, salts, hydrates, and solvates thereof.

- 5 Whenever used hereinafter, the term "optionally substituted" is meant to include unsubstituted as well as substituted with at least one of the specified substituting radicals. For the purpose of example, "C₁₋₄alkyl optionally substituted with chloro" is meant to include unsubstituted C₁₋₄alkyl as well as C₁₋₄alkyl substituted with chloro.




- 10 The compounds of formula (I) may have one or more centers of chirality and may exist as stereochemically isomeric forms. The term "stereochemically isomeric forms" as used herein defines all the possible compounds made up of the same atoms bonded by the same sequence of bonds but having different three-dimensional structures, which the compounds of formula (I) may possess.

- With reference to the instances where (*R*) or (*S*) is used to designate the absolute
15 configuration of a chiral atom within a substituent, the designation is done taking into consideration the whole compound and not the substituent in isolation.


In one aspect, the present invention provides compounds of formula (I)




- 20 including stereochemically isomeric forms, and *N*-oxides, salts, hydrates, and solvates thereof, wherein R¹, R², R⁴, R⁵, R⁶, R⁷ and A have the same meaning as defined herein. Embodiments of the present inventions concerns compounds of formula (I), or any subgroup thereof as defined herein, wherein one or more of the definitions for R¹, R², R⁴, R⁵, R⁶, and R⁷ as specified in the embodiments herein-under apply:
- 25 Particular subgroups of the compounds of formula (I) are compounds of formula (II), (III) or (IV)




In one embodiment, R^1 is a bivalent chain selected from




In a particular embodiment, R^1 is selected from










10 wherein a is 4 or 5 and c is 1 or 2. In another particular embodiment R^1 is selected from





amine is connected to the carbonyl group, or, the piperazinyl moiety is connected to the carbonyl group and the aliphatic amine is connected to the sulfonamide group.


Preferably, when R^1 is , the piperazinyl moiety is connected to the carbonyl group and the aliphatic amine is connected to the sulfonamide group.


- 5 In a more particular embodiment, R^1 is selected from , , and .

Alternatively, R^1 is selected from , , , and .

In a preferred embodiment, R^1 is .

- 10 In another embodiment, R^1 is .

In another embodiment R^1 is .



In another preferred embodiment, R^1 is .


- Each R^3 is independently selected from the group comprising hydrogen, C_{1-4} alkyl and C_{3-5} cycloalkyl. In a particular embodiment, R^3 is independently selected from the group consisting of hydrogen, methyl, ethyl, isopropyl and cyclopropyl. In a more particular embodiment, each R^3 is independently selected from the group consisting of hydrogen and methyl; or, R^3 is methyl.
- 15

Macrocycle A has 14 to 18 member atoms. In a particular embodiment, macrocycle A has 16, 17 or 18 member atoms. In a more particular embodiment, A has 17 member atoms.

- R^2 is selected from the group comprising hydrogen, halo or C_{1-4} alkoxy. In a particular embodiment, R^2 is selected from the group comprising hydrogen, chloro, fluoro or methoxy. In a more particular embodiment, R^2 is hydrogen or methoxy or chloro; or, alternatively, R^2 is fluoro or methoxy; or, in a preferred embodiment, R^2 is methoxy. In another embodiment, R^2 is positioned on the benzene ring in meta or para with respect to the bond linking the benzene to the indole group. In a preferred embodiment, R^2 is positioned on the benzene ring in para with respect to the bond linking this benzene to the indole group.
- R^4 and R^5 are hydrogen or R^4 and R^5 together form a double bond or a methylene group to form a fused cyclopropyl. In a particular embodiment, R^4 and R^5 are hydrogen or R^4 and R^5 together form a methylene group to form a fused cyclopropyl.
- In another particular embodiment, R^4 and R^5 together form a double bond.
- In another embodiment, R^6 is selected from hydrogen and methyl. In a particular embodiment, R^6 is hydrogen when the compound of formula (I) is a compound of formula (III) or (IV). In another particular embodiment, R^6 is methyl when the compound of formula (I) is a compound of formula (II).
- R^7 is a C_{3-7} cycloalkyl optionally substituted with halo. In a particular embodiment, R^7 is selected from cyclopentyl, cyclohexyl, and fluorocyclohexyl (in particular, 2-fluorocyclohexyl). In a preferred embodiment, R^7 is cyclohexyl.

- A particular subgroup of compounds of formula (I) are compounds of formula (I) wherein R^4 and R^5 together form a double bond, and wherein one or more of the definitions for R^1 , R^2 , R^6 , and R^7 as specified in the embodiments herein apply. A more particular subgroup of compounds of formula (I) are compounds of formula (II), wherein R^1 , R^2 , R^4 , R^5 , R^6 , R^7 and A have the same meaning as defined herein. More particular are those compound represented by the following structural formulae (II-1), (II-2), and (II-3) wherein R^2 , R^6 and R^7 have the same meaning as defined herein for compounds of formula (I) or subgroups thereof.





In a particular embodiment, the invention provides compounds of, independently, formula (II), (II-1), (II-2) and (II-3) wherein R^6 is hydrogen or methyl, more in particular, wherein R^6 is a methyl.





- 5 In another embodiment, the invention provides compounds of formula (II) or subgroups thereof wherein R^7 is cyclohexyl or 2-fluorocyclohexyl.

In another embodiment, the invention provides compounds of formula (II) or subgroups thereof wherein R^2 is hydrogen, methoxy or chloro. Alternatively, the invention provides compounds of formula (II) or subgroups thereof wherein R^2 is fluoro or methoxy.

10

A particular subgroup of compounds of formula (I) are compounds of formula (III), wherein R^1 , R^2 , R^4 , R^5 , R^6 , R^7 and A have the same meaning as defined herein. More particular are those compounds represented by the following structural formulae (III-1), (III-2), (III-3) and (III-4) wherein R^2 , R^6 and R^7 have the same meaning as defined

- 15 herein for compounds of formula (I).




In particular, the invention provides compounds of, independently, formula (III), (III-1), (III-2), (III-3) and (III-4) wherein R^6 is hydrogen.

In another embodiment, the invention provides compounds of formula (III) or subgroups thereof wherein R^7 is cyclohexyl or 2-fluorocyclohexyl.

In another embodiment, the invention provides compounds of formula (III) or subgroups thereof wherein R^2 is hydrogen, methoxy or chloro.

- 5 A particular subgroup of compounds of formula (I) are compounds of formula (IV), wherein R^1 , R^2 , R^4 , R^5 , R^6 , R^7 and A have the same meaning as defined herein. More particular are those compound represented by the following structural formulae (IV-1), (IV-2), and (IV-3) wherein R^2 , R^6 and R^7 have the same meaning as defined herein for compounds of formula (I).




In another embodiment, the invention provides compounds of formula (IV) or subgroups thereof wherein R^7 is cyclohexyl or 2-fluorocyclohexyl.


- 15 In another embodiment, the invention provides compounds of formula (IV) or subgroups thereof wherein R^2 is hydrogen, methoxy or chloro.

In a particular embodiment, the present invention concerns compounds of formula (II-1), (III-1), and (IV-1). Another embodiment of the present invention concerns compounds of formula (II-2), (III-2) and (IV-2). Another embodiment of the present invention concerns compounds of formula (II-3), (III-3) and (IV-3).


- 20 In a particular embodiment, the invention provides compounds of formula (I) selected from the group comprising




More in particular, the present invention provides compounds of formula (I) selected




5 Alternatively, the present invention provides compounds of formula (I) selected from





More in particular, the present invention provides compounds of formula (I) selected from



Unless otherwise mentioned or indicated, the chemical designation of a compound encompasses the mixture of some or all possible stereochemically isomeric forms, which said compound might possess. Said mixture may contain all diastereomers and/or enantiomers of the basic molecular structure of said compound. All stereochemically isomeric forms of the compounds of the present invention both in

pure form or mixed with each other are intended to be embraced within the scope of the present invention.





Pure stereoisomeric forms of the compounds and intermediates as mentioned herein are defined as isomers substantially free of other enantiomeric or diastereomeric forms of the same basic molecular structure of said compounds or intermediates. In particular, the term "stereoisomerically pure" concerns compounds or intermediates having a stereoisomeric excess of at least 80% (i.e. minimum 90% of one isomer and maximum 10% of the other possible isomers) up to a stereoisomeric excess of 100% (i.e. 100% of one isomer and none of the other), more in particular, compounds or intermediates having a stereoisomeric excess of 90% up to 100%, even more in particular having a stereoisomeric excess of 94% up to 100% and most in particular having a stereoisomeric excess of 97% up to 100%. The terms "enantiomerically pure" and "diastereomerically pure" should be understood in a similar way, but then having regard to the enantiomeric excess, and the diastereomeric excess, respectively, of the mixture in question.

Pure stereoisomeric forms of the compounds and intermediates of this invention may be obtained by the application of art-known procedures. For instance, enantiomers may be separated from each other by the selective crystallization of their diastereomeric salts with optically active acids or bases. Examples thereof are tartaric acid, dibenzoyltartaric acid, ditoluoyltartaric acid and camphosulfonic acid. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided the reaction occurs stereospecifically. Preferably, if a specific stereoisomer is desired, said compound is synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The diastereomeric racemates of the compounds of formula (I) or any subgroup thereof, can be obtained separately by conventional methods. Appropriate physical separation methods that may advantageously be employed are, for example, selective crystallization and chromatography, e.g. column chromatography.





For some of the compounds of formula (I), their *N*-oxides, salts, hydrates, solvates, quaternary amines, or metal complexes, and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined. A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction.











In an embodiment, the present invention concerns compounds of formula (IIIA), (IIIB), (IVA) and (IVB),



5 wherein R^1 , R^2 , R^6 , R^7 and A have the same meaning as that defined herein.


In a more particular embodiment, the present invention concerns compounds of Formula (IIIA-1), (IIIA-2), (IIIA-3), (IIIA-4), (IIIB-1), (IIIB-2), (IIIB-3), (IIIB-4), (IVA-1), (IVA-2), (IVA-3), (IVB-1), (IVB-2) and (IVB-3)





wherein R^2 , R^6 and R^7 have the same meaning as that defined herein.

In another embodiment, where applicable, compounds of formula (I) or subgroups thereof have the stereochemical configuration as illustrated by formula (IA).



The present invention is also intended to include all isotopes of atoms occurring on the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

For therapeutic use, salts of the compounds of formula (I) are those wherein the counter-ion is pharmaceutically acceptable. However, salts of acids and bases, which are non-pharmaceutically acceptable, may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

The pharmaceutically acceptable acid and base salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid and base addition salt forms that the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (i.e. ethanedioic), malonic, succinic (i.e. butanedioic acid), maleic, fumaric, malic (i.e. hydroxybutanedioic acid), tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-amino-salicylic, pamoic and the like acids.

Conversely, said salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) or any subgroup thereof containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium,

sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

5 The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) or any subgroup thereof are able to form by reaction between a basic nitrogen of a compound of formula (I) or any subgroup thereof and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl
10 trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.

15 The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) or any subgroup thereof wherein one or several nitrogen atoms are oxidized to the so-called *N*-oxide.

It will be appreciated that the compounds of formula (I) or any subgroup thereof may have metal binding, chelating, complex forming properties and therefore may exist as metal complexes or metal chelates. Such metalated derivatives of the compounds of
20 formula (I) or any subgroup thereof are intended to be included within the scope of the present invention.


Some of the compounds of formula (I) or any subgroup thereof and intermediates may also exist in one or more tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present
25 invention. Accordingly, the compounds and intermediates may be present as a mixture of tautomers or as an individual tautomer.

In the invention, particular preference is given to compounds of Formula I or any subgroup thereof, that in the inhibition assays described below have an inhibition value of less than 100 μM , preferably less than 50 μM , more preferably less than 10 μM ,
30 preferably less than 5 μM , even more preferably less than 1 μM preferably less than 100 nM, and in particular less than 10 nM, as determined by a suitable assay, such as the assays used in the Examples below.

It is to be understood that the above defined subgroups of compounds of formula (I) as well as any other subgroup defined herein, are meant to include stereochemically
35 isomeric forms, and any *N*-oxides, salts, quaternary amines, hydrates, solvates and metal complexes of such compounds.

Preparation of the compounds of formula (I)General synthetic schemes

Compounds of formula (I) may be prepared following the different methods A, B, C, D, E, F and G described below, from indole derivatives A-1



wherein R^2 , R^4 , R^5 , R^6 and R^7 are as defined for compounds of formula (I) or subgroups thereof, and R_a is selected from methyl and *tert*-butyl and R_b is selected from methyl. The compounds of formula (A-1) are either known in the art or may be obtained as described in US20070270406A1, WO2007/054741 and WO2007/092000.

10

Method A

A schematic overview for the synthesis of the compounds of formula (I) is given in scheme 1. The method starts from a compound of formula A-1.


15 Compounds of formula A-2 may be prepared by the regioselective hydrolysis of the ester bearing the R_b group, under basic conditions, using a hydroxide such as LiOH or NaOH, in polar solvents such as water, an alcohol such as methanol or ethanol, tetrahydrofuran (THF), or a mixture thereof. This method may be used when R_b is a methyl group and R_a is a *tert*-butyl group, or R_a is a methyl group.

20 A monoprotected bifunctional R^1 derived reagent of formula $PG-R^1-H$ wherein R^1 is as defined for formula (I) or subgroups thereof; may then be coupled to the carboxylic acid of compounds A-2 to form an amide bond, leading to compounds A-3. "PG", as used herein, is a suitable amine protecting group, chosen from the ones known in the art. Preferably PG is a *tert*-butoxycarbonyl (Boc) protecting group or a 4-nitrobenzenesulfonyl (nosyl) group.

25 The formation of amide bonds can be carried out using standard procedures such as those used for coupling amino acids in peptide synthesis. The latter involves the dehydrative coupling of a carboxyl group of one reactant with an amino group of the other reactant to form a linking amide bond. The amide bond formation may be performed by reacting the starting materials in the presence of a coupling agent or by
30 converting the carboxyl functionality into an active form such as an active ester, mixed anhydride or a carboxyl acid chloride or bromide. General descriptions of such

coupling reactions and the reagents used therein can be found in general textbooks on peptide chemistry, for example, M. Bodanszky, "Peptide Chemistry", 2nd rev. ed., Springer-Verlag, Berlin, Germany, (1993).

5 Scheme 1



Examples of coupling reactions with amide bond formation include the azide method, mixed carbonic-carboxylic acid anhydride (isobutyl chloroformate) method, the carbodiimide (dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIC), or water-soluble carbodiimide such as *N*-ethyl-*N'*-[3-(dimethylamino)propyl]carbodiimide (EDC)) method, the active ester method (e.g. *p*-nitrophenyl, *p*-chlorophenyl, trichlorophenyl, pentachlorophenyl, pentafluorophenyl, *N*-hydroxysuccinic imide and the like esters), the Woodward reagent K-method, the 1,1-carbonyldiimidazole (CDI or *N,N'*-carbonyldiimidazole) method, the phosphorus reagents or oxidation-reduction methods. Some of these methods can be enhanced by adding suitable catalysts, e.g. in the carbodiimide method by adding 1-hydroxybenzotriazole, or 4-dimethylaminopyridine (4-DMAP). Further coupling agents are (benzotriazol-1-yloxy)-*tris*-(dimethylamino) phosphonium hexafluorophosphate, either by itself or in the presence of 1-hydroxy-benzotriazole or 4-DMAP; or 2-(1*H*-benzotriazol-1-yl)-*N,N,N',N'*-tetra-methyluronium tetrafluoroborate, or O-(7-azabenzotriazol-1-yl)-

N,N,N',N'-tetramethyluronium hexafluorophosphate. These coupling reactions can be performed in either solution (liquid phase) or solid phase.

The coupling reactions preferably are conducted in an inert solvent, such as halogenated hydrocarbons, e.g. dichloromethane (DCM), chloroform, dipolar aprotic solvents such as acetonitrile, dimethylformamide (DMF), dimethylacetamide, DMSO, HMPT, ethers such as tetrahydrofuran (THF).

In many instances the coupling reactions are done in the presence of a suitable base such as a tertiary amine, e.g. triethylamine, diisopropylethylamine (DIPEA), *N*-methyl-morpholine, *N*-methylpyrrolidine, 4-DMAP or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). The reaction temperature may range between 0 °C and 50 °C and the reaction time may range between 15 min and 24 h.

Removal of the protecting group following methods known in the art may lead to compounds A-4. These methods include the reaction of compounds A-3 with trifluoroacetic acid (TFA) in a suitable solvent such as DCM, when PG is a Boc-protecting group, or the reaction of compounds A-3 with a thiol like mercapto acetic acid or thiophenol, in solution or in solid phase, in the presence of a base, such as cesium carbonate or LiOH, in a suitable solvent, such as DMF, THF when PG is nosyl. When Ra is a *tert*-butyl group and PG is a Boc-protecting group, removal of PG as described above, may lead to a compound A-4, with Ra being OH.

Compounds A-4 are then reacted with sulfamide, in a suitable solvent, for example dioxane, under heating conditions, ie 100°C. This reaction may take place under microwave irradiation and lead to compounds A-5. Another method to introduce the sulfamide moiety may consist of the reaction of compound A-4 with aminosulfonyl-chloride, in the presence of a suitable base, such as triethylamine, DIPEA, or pyridine, in a suitable solvent, such as a chlorinated solvent like DCM, or DMF, THF.


The ester function of compounds A-5, i.e. -CO-O-Ra, may then be hydrolyzed, using conditions known in the art, and including the saponification in basic media as described above, leading to compounds A-6. Heating may be required to complete this reaction. Acidic conditions may also be used to hydrolyze the ester function of compounds A-5, for example TFA in a suitable solvent like DCM, when Ra is a *tert*-butyl group.

Compounds (I) may be obtained by macrocyclisation by forming the intramolecular acylsulfamide bond, in the presence of coupling agents, such as CDI which converts the carboxylic acid group to a reactive species acylimidazole, under heating. This acylimidazole may then be purified before adding a suitable base such as DBU, in order to perform the ring closure, which may take place under heating conditions. Solvents

used for these reactions may include acetonitrile or THF. Other coupling agents, such as those known in the art, may also be used to achieve the ring closure.

Method B

5 Scheme 2




An alternate method leading to compounds A-4 as illustrated in scheme 2, may be the formation of an amide bond between compounds A-2 and a symmetrical bivalent chain R^1 , used in excess compared to compounds A-2. This amide bond may be synthesized as described above, in particular using a coupling agent such as [dimethylamino-
 10 ([1,2,3]triazolo[4,5-b]pyridin-3-yloxy)-methylene]-dimethyl-ammonium hexafluorophosphate (HATU), in the presence of a base such as DIPEA and in a suitable solvent like DCM, DMF, or more preferably THF. Compounds A-4 may then be reacted as described above in method A in order to prepare compounds (I).

15

Method C

Scheme 3



Compounds may be prepared directly from compounds A-2, in a similar way as described above for the synthesis of compounds A-3, but using a bivalent chain R^1
 20 bearing one sulfamide moiety instead of a protecting group. Such a sulfamide chain R^1 may be introduced on $H-R^1-H$ by heating a reagent of formula $H-R^1-H$, which can either be mono-protected by a suitable protecting group (i.e. $PG-R^1-H$), or not if it is symmetrical, with sulfamide in a suitable solvent, such as dioxane, under microwave
 25 irradiation. The protecting group may then be removed by methods known in the art,


for example by reaction with TFA in dichloromethane when the protecting group is a Boc-protecting group, leading to the monosulfamide derivatized R¹ chain.

Method D

- 5 Compounds of formula A-3 or A-4 may undergo functional group manipulation, such as alkylation or reductive amination, before PG removal of compounds A-3 and/or reaction leading to the sulfamide A-4.

Method E

Scheme 4



10

The ester bearing the Ra group of compounds A-1 (Ra being for example a *tert*-butyl group and Rb a methyl group) may be hydrolyzed as described above, in acidic conditions, using for instance TFA in a suitable solvent like DCM, to yield the carboxylic acid derivative E-2.

- 15 Reaction of compounds E-2 with the sulfamide moiety introduced on a mono-protected bivalent chain R¹, may lead to the acyl sulfamide compounds E-3, using the conditions described for the last step of method A. Preferably the coupling agent used to activate the carboxylic acid group may be CDI, in a suitable solvent like acetonitrile or THF, under heating conditions. Addition of the sulfamide chain in the presence of a base
- 20 such as DBU may subsequently lead to compounds E-3. PG is a suitable amine protecting group, chosen from the ones known in the art. Preferably, within method E, PG is a Boc-protecting group.

Removal of the protecting group PG of compounds E-3 following methods known in the art may lead to compounds E-4. These methods include the reaction of compounds

- 25 E-3 with TFA in a suitable solvent such as DCM, when PG is a Boc-protecting group.


The ester function of compounds **E-4** (Rb is a methyl group) may then be hydrolyzed, using conditions known in the art, and including the saponification in basic media as described above, leading to compounds **E-5**.

Alternatively, compounds **E-3** may undergo the saponification reaction in basic media to hydrolyze the ester bearing Rb, prior to the removal of the amine protecting group using the conditions described above, and leading to compounds **E-5**.

Compounds (**I**) may be obtained by macrocyclisation of compounds **E-5** by forming the intramolecular amide bond, in the presence of coupling agents, as described in method A. Preferably this amide formation step may be performed under high dilution conditions.

Method F


Scheme 5



Compounds **F-3** may be obtained by an amide forming reaction, starting from compounds **A-2** and an alkenylamine, as described for the second step of method A. Subsequent ester hydrolysis under basic or acidic conditions as described previously may lead to compounds **F-4**. The acylsulfamide bond may then be formed using the method described for the last step of method A, using an alkenyl sulfamide compound and leading to compounds **F-5**.

Alternatively, the acylsulfamide group may be introduced on a compound of formula **E-2**, prior to the hydrolysis of the ester bearing the Rb group and coupling of the obtained carboxylic acid with an alkenamine as described above, leading to compound **F-5**.

Formation of the macrocycle, i.e. compound of formula **F-6**, which is a compound of



formula **(I)** bearing the following bivalent chain as R^1 :

- can be carried out via an olefin metathesis reaction in the presence of a suitable metal catalyst such as e.g. the Ru-based catalyst reported by Miller, S.J., Blackwell, H.E., Grubbs, R.H. J. Am. Chem. Soc. 118, (1996), 9606-9614; Kingsbury, J. S., Harrity, J. P. A., Bonitatebus, P. J., Hoveyda, A. H., J. Am. Chem. Soc. 121, (1999), 791-799; and Huang et al., J. Am. Chem. Soc. 121, (1999), 2674-2678; for example a Hoveyda-Grubbs catalyst.

- Air-stable ruthenium catalysts such as bis(tricyclohexylphosphine)-3-phenyl-1H-inden-1-ylidene ruthenium chloride (Neolyst M1[®]) or bis(tricyclohexylphosphine)-[(phenylthio)methylene]ruthenium (IV) dichloride can be used. Other catalysts that can be used are Grubbs first and second generation catalysts, i.e. benzylidene-bis(tricyclohexylphosphine)dichlororuthenium and (1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene)dichloro(phenylmethylene)(tricyclohexylphosphine)ruthenium, respectively. Of particular interest are the Hoveyda-Grubbs first and second generation catalysts, which are dichloro(*o*-isopropoxyphenylmethylene)(tricyclohexylphosphine)-ruthenium(II) and 1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene)dichloro(*o*-isopropoxyphenylmethylene)ruthenium respectively. Also, other catalysts containing other transition metals such as Mo can be used for this reaction.


- The metathesis reactions may be conducted in a suitable solvent such as for example ethers, e.g. THF, dioxane; halogenated hydrocarbons, e.g. dichloromethane, CHCl_3 , 1,2-dichloroethane and the like, hydrocarbons, e.g. toluene. These reactions are conducted at increased temperatures under nitrogen atmosphere.

- Compounds of formula **(I)** or any subgroup thereof or any subgroups thereof may be converted into each other following art-known functional group transformation reactions. For example, amino groups may be *N*-alkylated, nitro groups reduced to amino groups, a halo atom may be exchanged for another halo.

- Compounds of formula **F-6** may be submitted to catalytic hydrogenation, using for example Pd/C as a catalyst, in a suitable solvent such as methanol, ethanol, THF, acetic acid or a mixture thereof, to yield compounds of formula **F-7**, where the alkene of the bivalent chain R^1 is reduced to the corresponding alkane. Compounds of formula **F-6** belonging to the group of compounds of formula **(II)** may lead to compounds **F-7** having the structure of compounds of formula **(IV)** after this hydrogenation step.

More generally, a compound of formula (II) may be transformed to a compound of formula (IV) by catalytic hydrogenation as shown below.

Scheme 6



- 5 The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth
- 10 alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxoperoxoic acid or halo substituted benzenecarboxoperoxoic acid, e.g. 3-chlorobenzenecarboxoperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *tert*-butyl hydroperoxide. Suitable solvents are, for example, water, lower
- 15 alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques,

20 e.g., counter-current distribution, liquid chromatography and the like.

The compounds of formula (I) may be obtained as racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I), which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable

25 chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure


30 stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the

reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound may be synthesized by stereospecific methods of preparation. These methods may advantageously employ enantiomerically pure starting materials.

Method G describes the synthesis of enantiomerically pure starting materials A-2, belonging to the groups of compounds (III) and (IV).

Method G

Scheme 7



A racemic mixture A-2 may be reacted with a chiral auxiliary, such as (S)-4-benzyl-2-oxazolidinone, after having being transformed to its acylchloride using methods known in the art, such as reaction of A-2 with oxalyl chloride in a suitable solvent like THF, in the presence of a catalytic amount of DMF. The acid chloride may then be reacted with the anion of (S)-4-benzyl-2-oxazolidinone formed by the reaction with a strong base, such as butyl lithium, in a suitable solvent such as THF, at low temperatures, typically -78°C, and under an inert atmosphere, leading to the diastereoisomers G1 and G2, which can be isolated by methods known in the art, such as chromatography on silica gel.

Removal of the chiral auxiliary from each of the diastereoisomers G1 and G2 may then be performed with a base such as NaOH in a suitable solvent, such as methanol, water, THF, leading to the enantiomerically pure compounds A-2' and A-2''. Using these

enantiomerically pure starting materials may lead to enantiomerically pure compounds of formula (I) bearing one stereocenter, such as compounds of formula (IIIA), (IIIB).

Pure stereochemically isomeric forms of the compounds of formula (I) or any subgroups thereof may be obtained by the application of art-known procedures.

- 5 Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., counter-current distribution, liquid chromatography and the like.

- The compounds of formula (I) or any subgroups thereof may be obtained as racemic mixtures of enantiomers, which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) or any subgroups thereof, which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) or any subgroups thereof involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound may be synthesized by stereospecific methods of preparation. These methods may advantageously employ enantiomerically pure starting materials.
- 10
15
20

- In a further aspect, the present invention concerns a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I) or any subgroups thereof, as specified herein, and a pharmaceutically acceptable carrier. A therapeutically effective amount in this context is an amount sufficient to prophylactically act against, to stabilize or to reduce viral infection, and in particular HCV viral infection, in infected subjects or subjects being at risk of being infected. In still a further aspect, this invention relates to a process of preparing a pharmaceutical composition as specified herein, which comprises intimately mixing a pharmaceutically acceptable carrier with a therapeutically effective amount of a compound of formula (I) or any subgroups thereof, as specified herein.
- 25
30

- Therefore, according to an embodiment of the present invention, the compounds of formula (I) or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. It is understood that all compositions usually employed for systemically administering drugs are included as appropriate
- 35

compositions. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in salt form or a metal complex, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations that are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

The compounds of the present invention may also be administered via oral inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder, a solution being preferred. Any system developed for the delivery of solutions, suspensions or dry powders via oral inhalation or insufflation are suitable for the administration of the present compounds.

Thus, the present invention also provides a pharmaceutical composition adapted for administration by inhalation or insufflation through the mouth comprising a compound of formula (I) or any subgroups thereof and a pharmaceutically acceptable carrier.

Preferably, the compounds of the present invention are administered via inhalation of a solution in nebulized or aerosolized doses.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage.

- 5 Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, suppositories, powder packets, wafers, 10 injectable solutions or suspensions and the like, and segregated multiples thereof.

- The compounds of formula (I) and any subgroup thereof show antiviral properties. Viral infections and their associated diseases treatable using the compounds and methods of the present invention include those infections brought on by HCV and other pathogenic flaviviruses such as Yellow fever, Dengue fever (types 1-4), St. Louis 15 encephalitis, Japanese encephalitis, Murray valley encephalitis, West Nile virus and Kunjin virus. The diseases associated with HCV include progressive liver fibrosis, inflammation and necrosis leading to cirrhosis, end-stage liver disease, and HCC; and for the other pathogenic flaviviruses the diseases include yellow fever, dengue fever, hemorrhagic fever and encephalitis.

- 20 However, compounds of the invention may also be attractive due to the fact that they lack activity against other viruses, in particular against HIV. HIV infected patients often suffer from co-infections such as HCV. Treatment of such patients with an HCV inhibitor that also inhibits HIV may lead to the emergence of resistant HIV strains.

- Due to their antiviral properties, particularly their anti-HCV properties, the compounds 25 of formula (I) or any subgroup thereof, including stereochemically isomeric forms, and their *N*-oxides, quaternary amines, metal complexes, salts, hydrates and solvates, are useful in the treatment of individuals experiencing a viral infection, particularly a HCV infection, and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected 30 with viruses, in particular flaviviruses such as HCV.

- The compounds of the present invention or any subgroup thereof may therefore be used as medicines. Said use as a medicine or method of treatment comprises the systemic administration to virally infected subjects or to subjects susceptible to viral infections of an amount effective to combat the conditions associated with the viral infection, in 35 particular the HCV infection.

The present invention also relates to the use of the present compounds or any subgroup thereof in the manufacture of a medicament for the treatment or the prevention of viral infections, particularly HCV infection.

5 The present invention furthermore relates to a method of treating a warm-blooded animal infected by a virus, or being at risk of infection by a virus, in particular by HCV, said method comprising the administration of an anti-virally effective amount of a compound of formula (I), or any subgroups thereof, as specified herein.

The present invention also concerns combinations of a compound of formula (I) or any subgroup thereof, as specified herein with other anti-HCV agents. In an embodiment, 10 the invention concerns combination of a compound of Formula (I) or any subgroup thereof with at least one anti-HCV agent. In a particular embodiment, the invention concerns combination of a compound of Formula (I) or any subgroup thereof with at least two anti-HCV agents. In a particular embodiment, the invention concerns combination of a compound of Formula (I) or any subgroup thereof with at least three 15 anti-HCV agents. In a particular embodiment, the invention concerns combination of a compound of Formula (I) or any subgroup thereof with at least four anti-HCV agents.

The combination of previously known anti-HCV compound, such as interferon- α (IFN- α), pegylated interferon- α , ribavirin or a combination thereof, and, a compound of formula (I) or any subgroup thereof can be used as a medicine in a combination 20 therapy. In an embodiment, the term "combination therapy" relates to a product containing mandatory (a) a compound of formula (I), and (b) at least one other anti-HCV compound, as a combined preparation for simultaneous, separate or sequential use in treatment of HCV infections, in particular, in the treatment of infections with HCV.

25 Anti-HCV compounds encompass agents selected from HCV polymerase inhibitors, R-7128, MK-0608, VCH759, PF-868554, GS9190, NM283, valopicitabine, PSI-6130, XTL-2125, NM-107, R7128 (R4048), GSK625433, R803, R-1626, BILB-1941, HCV-796, JTK-109 and JTK-003, ANA-598, IDX-184, MK-3281, MK-1220, benzimidazole derivatives, benzo-1,2,4-thiadiazine derivatives, phenylalanine 30 derivatives, A-831 and A-689; HCV proteases (NS2-NS3 and NS3-NS4A) inhibitors, the compounds of WO02/18369 (see, e.g., page 273, lines 9-22 and page 274, line 4 to page 276, line 11), BI-1335, TMC435350, MK7009, ITMN-191, BILN-2061, VX-950, BILN-2065, BMS-605339, VX-500, SCH 503034; inhibitors of other targets in the HCV life cycle, including helicase, and metalloprotease inhibitors, ISIS-14803; 35 immunomodulatory agents such as, α -, β -, and γ - interferons such as rIFN- α 2b, rIFN- α 2ba, consensus IFN- α (infergen), feron, reaferon, intermax α , rIFN- β , infergen +

actimmune, IFN-omega with DUROS, albuferon, locteron, Rebif, Oral IFN- α , IFN- α 2b XL, AVI-005, pegylated-infergen, pegylated derivatized interferon- α compounds such as pegylated rIFN- α 2b, pegylated rIFN- α 2a, pegylated IFN- β , compounds that stimulate the synthesis of interferon in cells, interleukins, Toll like receptor (TLR) agonists, compounds that enhance the development of type 1 helper T cell response, and thymosin; other antiviral agents such as ribavirin, ribavirin analogs such as rebetol, copegus and viramidine (taribavirin), amantadine, and telbivudine, inhibitors of internal ribosome entry, alpha-glucosidase 1 inhibitors such as MX-3253 (celgosivir) and UT-231B, hepatoprotectants such as IDN-6556, ME-3738, LB-84451 and MitoQ, broad-spectrum viral inhibitors, such as IMPDH inhibitors (e.g., compounds of US5,807,876, US6,498,178, US6,344,465, US6,054,472, WO97/40028, WO98/40381, WO00/56331, mycophenolic acid and derivatives thereof, and including, but not limited to VX-497, VX-148, and/or VX-944); and other drugs for treating HCV such as zadaxin, nitazoxanide, BIVN-401 (virostat), PYN-17 (altirex), KPE02003002, actilon (CPG-10101), KRN-7000, civacir, GI-5005, ANA-975, XTL-6865, ANA-971, NOV-205, tarvacin, EHC-18, NIM811, DEBIO-025, VGX-410C, EMZ-702, AVI 4065, Bavituximab, and Oglufanide; or combinations of any of the above.

Thus, to combat or treat HCV infections, the compounds of formula (I) or any subgroups thereof may be co-administered in combination with for instance, interferon- α (IFN- α), pegylated interferon- α , ribavirin or a combination thereof, as well as therapeutics based on antibodies targeted against HCV epitopes, small interfering RNA (si RNA), ribozymes, DNazymes, antisense RNA, small molecule antagonists of for instance NS3 protease, NS3 helicase and NS5B polymerase.

The combinations of the present invention may be used as medicaments. Accordingly, the present invention relates to the use of a compound of formula (I) or any subgroup thereof as defined above for the manufacture of a medicament useful for inhibiting HCV activity in a mammal infected with HCV viruses, wherein said medicament is used in a combination therapy, said combination therapy preferably comprising a compound of formula (I) and at least one other HCV inhibitory compound, e.g. IFN- α , pegylated IFN- α , ribavirin or a combination thereof.

Furthermore, it is known that a large percentage of patients infected with human immunodeficiency virus 1 (HIV) are also infected with HCV, i.e. they are HCV/HIV co-infected. HIV infection appears to adversely affect all stages of HCV infection, leading to increased viral persistence and accelerated progression of HCV-related liver disease. In turn, HCV infection may affect the management of HIV infection, increasing the incidence of liver toxicity caused by antiviral medications.

The present invention therefore also concerns combinations of a compound of Formula (I) or any subgroup thereof with anti-HIV agents. Also, the combination of one or more additional anti-HIV compounds and a compound of Formula (I) or any subgroups thereof can be used as a medicine. In particular, said combination can be used for inhibition HCV and HIV replication.

The term "combination therapy" also encompasses a product comprising (a) a compound of Formula (I) or any subgroup thereof, and (b) at least one anti-HIV compound, and (c) optionally at least one other anti-HCV compound, as a combined preparation for simultaneous, separate or sequential use in treatment of HCV and HIV infections, in particular, in the treatment of infections with HCV and HIV, or for preventing or treating conditions associated with HCV and HIV.

Thus, the present invention also relates to a product containing (a) at least one compound of Formula (I) or any subgroup thereof, and (b) one or more additional anti-HIV compounds, as a combined preparation for simultaneous, separate or sequential use in anti-HCV and anti-HIV treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. Said anti-HIV compounds may be any known antiretroviral compounds such as suramine, pentamidine, thymopentin, castanospermine, dextran (dextran sulfate), foscarnet-sodium (trisodium phosphono formate); nucleoside reverse transcriptase inhibitors (NRTIs), e.g. zidovudine (AZT), didanosine (ddI), zalcitabine (ddC), lamivudine (3TC), stavudine (d4T), emtricitabine (FTC), abacavir (ABC), amdoxovir (DAPD), elvucitabine (ACH-126,443), AVX 754 ((-)-DOTC), fozivudine tidoxil (FZT), phosphazide, HDP-990003, KP-1461, MIV-210, racivir (PSI-5004), UC-781 and the like; non-nucleoside reverse transcriptase inhibitors (NNRTIs) such as delavirdine (DLV), efavirenz (EFV), nevirapine (NVP), dapivirine (TMC120), etravirine (TMC125), rilpivirine (TMC278), DPC-082, (+)-Calanolide A, BILR-355, and the like; nucleotide reverse transcriptase inhibitors (NtRTIs), e.g. tenofovir ((R)-PMPA) and tenofovir disoproxil fumarate (TDF), and the like; nucleotide-competing reverse transcriptase inhibitors (NcRTIs), e.g. NcRTI-1 and the like; inhibitors of trans-activating proteins, such as TAT-inhibitors, e.g. RO-5-3335, BI-201, and the like; REV inhibitors; protease inhibitors e.g. ritonavir (RTV), saquinavir (SQV), lopinavir (ABT-378 or LPV), indinavir (IDV), amprenavir (VX-478), TMC126, nelfinavir (AG-1343), atazanavir (BMS 232,632), darunavir (TMC114), fosamprenavir (GW433908 or VX-175), brecanavir (GW-640385, VX-385), P-1946, PL-337, PL-100, tipranavir (PNU-140690), AG-1859, AG-1776, Ro-0334649 and the like; entry inhibitors, which comprise fusion inhibitors (e.g. enfuvirtide (T-20)), attachment inhibitors and co-receptor inhibitors, the latter comprise the CCR5 antagonists (e.g.

ancriviroc, CCR5mAb004, maraviroc (UK-427,857), PRO-140, TAK-220, TAK-652, vicriviroc (SCH-D, SCH-417,690)) and CXR4 antagonists (e.g. AMD-070, KRH-27315), examples of entry inhibitors are PRO-542, TNX-355, BMS-488043, BlockAide/CR™, FP 21399, hNM01, nonakine, VGV-1; a maturation inhibitor for example is PA-457; inhibitors of the viral integrase e.g. raltegravir (MK-0518), elvitegravir (JTK-303, GS-9137), BMS-538158; ribozymes; immunomodulators; monoclonal antibodies; gene therapy; vaccines; siRNAs; antisense RNAs; microbicides; Zinc-finger inhibitors.

Therefore, HCV infected patients also suffering from conditions associated with HIV or even other pathogenic retroviruses, such as AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic CNS diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis, can conveniently be treated with the present composition.

The compositions may be formulated into suitable pharmaceutical dosage forms such as the dosage forms described above. Each of the active ingredients may be formulated separately and the formulations may be co-administered or one formulation containing both and if desired further active ingredients may be provided.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients, as well as any product that results, directly or indirectly, from the combination of the specified ingredients.

The term "therapeutically effective amount" as used herein means that amount of active compound or component or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought, in the light of the present invention, by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease being treated. Since the instant invention refers as well to combinations comprising two or more agents, the "therapeutically effective amount" in the context of combinations is also that amount of the agents taken together so that the combined effect elicits the desired biological or medicinal response. For example, the therapeutically effective amount of a composition comprising (a) the compound of formula (I) and (b) another anti-HCV agent, would be the amount of the compound of formula (I) and the amount of the other anti-HCV agent that when taken together have a combined effect that is therapeutically effective.

In general, it is contemplated that an antiviral effective daily amount would be from 0.01 mg/kg to 500 mg/kg body weight, more preferably from 0.1 mg/kg to 50 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four

or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

5 The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject
10 and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines.



In one embodiment of the present invention there is provided an article of manufacture comprising a composition effective to treat an HCV infection or to inhibit the NS5B
15 polymerase of HCV; and packaging material comprising a label which indicates that the composition can be used to treat infection by the hepatitis C virus; wherein the composition comprises a compound of the formula (I) or any subgroup thereof, or the combination as described herein.

Another embodiment of the present invention concerns a kit or container comprising a
20 compound of the formula (I) or any subgroup thereof, in an amount effective for use as a standard or reagent in a test or assay for determining the ability of potential pharmaceuticals to inhibit HCV NS5B polymerase, HCV growth, or both. This aspect of the invention may find its use in pharmaceutical research programs.


The compounds and combinations of the present invention can be used in
25 high-throughput target-analyte assays such as those for measuring the efficacy of said combination in HCV treatment.

Examples

The following examples are intended to illustrate the present invention and not to limit it thereto. Unless otherwise indicated, purification of the synthesized compounds by
30 column chromatography or flash chromatography is performed on a silica gel column.

Example 1- synthesis of compound 1**1**Step 1**1a****1b**


A solution of NaOH (6.38 g) in 25 mL of water was added to a stirred solution of **1a** (10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate, synthesized as described in US 2007270406 A1) in THF (100 mL) and MeOH (150 mL). After 1 hour the reaction was concentrated under reduced pressure, then diluted with ice-cold water (150 mL). The pH of the resulting solution was adjusted to 6 with acetic acid (AcOH). The precipitate was collected by filtration, washed with water and dried under vacuum to give 1.90g (98%) of **1b** as a yellowish powder: $m/z = 488$ ($M+H$)⁺

Step 2**1b****1c**

HATU (1.17 g, 3.08 mmol) was added under nitrogen to a stirred solution of **1b** (1.00 g, 2.05 mmol), DIPEA (1.07 mL, 6.15 mmol) and 2,2'-oxybis(*N*-methylethanamine) (1.08 g, 8.20 mmol) in 30 mL of dry THF. After 1 h, the reaction mixture was quenched with water (100 mL) and extracted with ethyl acetate (EtOAc). The organic layer was successively dried (Na_2SO_4), filtered and evaporated. The


residue was triturated in water, filtered and dried to give 1.15 g (93%) the target compound **1c** as a yellowish powder: $m/z = 602$ ($M+H$)⁺

Step 3




A solution of **1c** (1.15 g, 1.91 mmol) and sulfamide (1.84 g, 19.1 mmol) in dioxane (10 mL) was heated at 100°C in a microwave oven for 20 minutes. The reaction mixture was cooled down to room temperature, then evaporated under vacuum. The residue was triturated in water, filtered and washed with water. The powder was reconstituted in EtOAc, dried (Na_2SO_4) and evaporated to give 1.15 g (88%) of the desired product **1d** as a yellowish powder: $m/z = 681$ ($M+H$)⁺


Step 4



TFA (3.0 g, 26.3 mmol) was added to a solution of **1d** (1.15 g, 1.70 mmol) in dichloromethane (3 mL). After 1h, the reaction mixture was concentrated under vacuum. The residue was triturated in ether, filtered and washed with ether, then purified by chromatography (gradient EtOAc to EtOAc/EtOH, 9:1) to give 802 mg (76%) of the desired product **1e**: $m/z = 625$ ($M + H$)⁺

Step 5


Carbonyldiimidazole (389 mg, 2.40 mmol) was added to a stirred solution of **1e** (500 mg, 0.80 mmol) in dry THF (3 mL). The reaction mixture was stirred at room temperature for 1h: complete conversion to intermediate **1f** was observed. The resulting solution was evaporated, then the residue was purified by flash chromatography (gradient EtOAc to CH₃CN 1:0 to 0:1) to give 550 mg of the target product **1f** which was used as such in the next step: $m/z = 675$ ($M + H$)⁺

10 Step 6

DBU (244 mg, 0.32 mmol) was added to a solution of **1f** (550 mg) in acetonitrile (25 mL). The reaction mixture was stirred overnight at room temperature, then concentrated under reduced pressure. The residue was dissolved in water (30 mL) and the pH of the resulting solution was adjusted to 5. The precipitate was collected by filtration, washed with water and dried. Recrystallization from ethanol followed by a purification by column chromatography (gradient EtOAc to EtOAc/EtOH 9:1) provided 380 mg (78%) of the title product **1** as a white powder: $m/z = 607$ ($M + H$)⁺, ¹H NMR (DMSO-*d*₆) δ 1.15 (m, 1H), 1.40 (m, 3H), 1.71 (m, 2H), 1.88 (m, 1H), 2.01 (m, 3H), 2.56 (m, 3H), 2.77 (m, 1H), 2.99 (s, 3H), 3.26 (m, 2H), 3.50-3.71 (m, 6H), 3.87 (s, 3H), 4.44 (d, $J = 14.1$ Hz, 1H), 5.09 (d, $J = 15.0$ Hz, 1H), 6.95 (s, 1H), 7.13 (s, 1H), 7.19 (d,

$J = 8.6$ Hz, 1H), 7.47 (d, $J = 8.0$ Hz, 1H), 7.54 (d, $J = 8.3$ Hz, 1H), 7.88 (d, $J = 7.8$ Hz, 1H), 8.33 (s, 1H), 11.40 (s, 1H).


Example 2 – synthesis of compound 2



A solution of **1** (56 mg, 0.092 mmol) in MeOH (15 mL) and THF (5 mL) was hydrogenated in an H-cube apparatus using a 10% Pd on Carbon cartridge. Then, solvent was evaporated and the residue was purified by column chromatography (CH₂Cl₂/CH₃CN, 9:1) to give 23 mg (41%) of the desired product **2** as a white powder: $m/z = 609$ (M+H)⁺.



10

Examples 3 and 4 – synthesis of compounds 3 and 4




The racemic mixture **2** was purified by SFC, using a 6.5 minutes run on a chiral CHIRALCEL OD-H column (250 x 10 mm, coated on 5 μ m silica gel) and 55% methanol / 45% CO₂ as mobile phase, at a flow rate of 10 mL/min and lead to the two pure enantiomers **3** and **4**. Retention times under these conditions were observed at 4.25 min and 5.54 min.

20

Example 5 – synthesis of compound 5Step 1

5


The compound **5a** was synthesized in 96% yield from intermediate **1b** and 2-[4-(*tert*-butyloxycarbonyl)piperazin-1-yl]ethylamine following the procedure reported for the synthesis of intermediate **1c**: $m/z = 699$ ($M+H$)⁺.

Step 2


10

Trifluoroacetic acid (5.00 g, 43.9 mmol) was added to 740 mg of intermediate **5a**. After 1 hour at room temperature, the solvent was evaporated. The residue was triturated in EtOH/Et₂O, filtered and dried under high vacuum to give 380 mg (64%) of the desired product **5b** as a yellowish powder: $m/z = 543$ ($M+H$)⁺.


15


Step 3

5 A solution of **5b** (380 mg, 0.700 mmol) and sulfamide (673 mg, 7.00 mmol) in dioxane (10 mL) was heated at 100 °C in a microwave oven for 15 minutes. Then, the reaction mixture was successively cooled down at room temperature, concentrated under vacuum, triturated in water and filtered. Purification by column chromatography (gradient EtOAc/CH₂Cl₂ 1:1 to 1:0) gave 210 mg (46%) of the desired product **5c**: $m/z = 622$ (M+H)⁺.


10 Step 4

The title product **5d** was synthesized in 11 % yield following the procedure (steps 5 and 6) reported for the synthesis of compound **1**, followed by a recrystallization from ethanol, affording the desired product as white powder, $m/z = 604$ (M+H)⁺.


Example 6 - synthesis of compound 6

Step 1


- To a solution of *N*₁-(2-aminoethyl)-*N*₁-methylethane-1,2-diamine (10.58 g, 90 mmol) in DCM (350 mL) was added slowly a solution of 2-nitrobenzenesulfonyl chloride dissolved in DCM (50 mL). After 2 h at RT, the reaction mixture (RM) was washed with water, dried over MgSO₄, filtered and concentrated. The residue was purified by flash chromatography on silica gel with a gradient of methanol in DCM (0 to 10%), yielding to 6.9 g of *N*-(2-((2-aminoethyl)(methyl)amino)ethyl)-2-nitrobenzenesulfonamide **6a** and 3.9 g of *N,N'*-(2,2'-(methylazanediyl)-bis(ethane-2,1-diyl))bis(2-nitrobenzenesulfonamide) **6b**; *m/z* (**6a**) = 303 (M+H)⁺, *m/z* (**6b**) = 488 (M+H)⁺.

Step 2

- To a solution of carboxylic acid **1b** (500 mg, 1.025 mmol), HATU (585 mg, 1.5 eq) and diisopropylethylamine (212 mg, 1.6 eq) in dry DMF (10 mL) was added *N*-(2-((2-aminoethyl)(methyl)amino)ethyl)-2-nitrobenzenesulfonamide **6a** (341 mg, 1.1 eq). After 30 minutes at RT, the RM was diluted with water. The yellow precipitate was filtered off and washed with water. It was then reconstituted in EtOAc, dried over MgSO₄, filtered, concentrated and dried under vacuum to give 800 mg of the desired product 13-Cyclohexyl-3-methoxy-6-(2-{methyl[2-(2-nitrobenzenesulfonylamino)ethyl]-amino}-ethylcarbamoyl)-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **6c** as a yellow powder; *m/z* = 772 (M+H)⁺.


Step 3

To a solution of 13-cyclohexyl-3-methoxy-6-(2-{methyl-[2-(2-nitro-benzenesulfonyl)-amino]-ethyl}-amino)-ethylcarbamoyl]-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **6c** (650 mg, 0.842 mmole) and cesium carbonate (1.646-g, 6 eq) in dry DMF (10 mL) was added slowly a solution of methyl iodide (122 mg, 1.02 mmole) in dry DMF (2 mL). After stirring for 1h at RT, the RM was diluted with water and extracted with EtOAc. The organic layer was then washed with water, dried over MgSO₄, filtered and concentrated. The residue was purified by flash chromatography on silica gel, using a gradient of EtOAc in DCM (0 to 100%), yielding to 550 mg (83% yield) of the desired product 13-Cyclohexyl-3-methoxy-6-[2-(methyl-{2-[methyl-(2-nitro-benzenesulfonyl)-amino]-ethyl}-amino)-ethylcarbamoyl]-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **6d** as a yellow solid; $m/z = 786 (M+H)^+$.

Step 4


A mixture of 13-cyclohexyl-3-methoxy-6-[2-(methyl-{2-[methyl-(2-nitro-benzene-sulfonyl)amino]-ethyl}-amino)-ethylcarbamoyl]-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **6d** (380 mg, 0.483 mmole), cesium carbonate (315 mg, 2 eq) and thiophenol (107 mg, 2 eq) in DMF (5 mL) was stirred at RT overnight. Cesium carbonate (315 mg, 2 eq) and thiophenol (107 mg, 2 eq) were then added to the RM and the RM was stirred for 1h. Upon completion of the reaction, the RM was filtered and loaded on a cartridge containing a SCX-resin, pre-washed with

- DCM. After rinsing the cartridge with DCM (several times, until a colorless fraction was obtained) the product was eluted with NH_3 in MeOH, yielding to 240 mg of the desired product 13-cyclohexyl-3-methoxy-6-{2-[methyl-(2-methylamino-ethyl)-amino]-ethylcarbamoyl}-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **6e**, which was further purified by preparative HPLC; $m/z = 601$ ($\text{M}+\text{H}$)⁺.
- Step 5




- The synthesis of product **6f** was performed using the procedure described for the synthesis of compound **1d**, using intermediate **6e** instead of intermediate **1c**, yielding 200 mg (50%) of the target product; $m/z = 680$ ($\text{M}+\text{H}$)⁺.

Step 6





- The synthesis of product **6g** was performed using the procedure described for the synthesis of compound **1e**, using intermediate **6f** instead of intermediate **1d**, yielding 187 mg (quantitative yield) of the target product; $m/z = 624$ ($\text{M}+\text{H}$)⁺.


Step 7



- The synthesis of product **6** was performed using the procedure described for the synthesis of compound **1**, using intermediate **6g** instead of intermediate **1e**, yielding 43 mg (22% yield) of the target product; $m/z = 606$ ($\text{M}+\text{H}$)⁺.

Example 7 – synthesis of compound 7Step 1


To a solution of *N,N'*-(2,2'-(methylazanediyl)bis(ethane-2,1-diyl))bis(2-nitrobenzenesulfonamide) **6b** (3 g, 6.15 mmol) in dry DMF (50 mL) was added portion wise sodium hydride (738 mg, 3 eq, 60% in mineral oil) at 0°C. After 20 minutes, a solution of methyl iodide dissolved in dry DMF (5 mL) was added slowly to the RM. After stirring for 1 h at RT, the RM was quenched with water and extracted with EtOAc. The organic layer was washed with water, dried over MgSO₄, filtered and concentrated. Purification by flash chromatography with a gradient of EtOAc in DCM (20 to 80%) afforded 1.94 g (61%) of the desired product *N,N'*-(2,2'-(methylazanediyl)bis(ethane-2,1-diyl))bis(*N*-methyl-2-nitrobenzenesulfonamide) **7a**; *m/z* = 516 (M+H)⁺.

Step 2

A mixture of *N,N'*-(2,2'-(methylazanediyl)bis(ethane-2,1-diyl))bis(*N*-methyl-2-nitrobenzenesulfonamide) **7a** (1.24 g, 2.405 mmol), cesium carbonate (2.35 g, 3 eq) and thiophenol (795 mg, 3 eq) in DMF (25 mL) was stirred at RT during 1 h. Upon completion of the reaction, the RM was filtered and loaded on a MP-TsOH cartridge, prewashed with DCM. After rinsing the cartridge with DCM (several times, until a colorless fraction was obtained) the product was eluted with NH₃ in MeOH, yielding to


220 mg (63%) of *N*₁,*N*₂-dimethyl-*N*₁-(2-(methylamino)ethyl)ethane-1,2-diamine **7b**, which was used directly in the next step; *m/z* = 146 (*M*+*H*)⁺.

Step 3




The synthesis of 13-cyclohexyl-3-methoxy-6-(methyl-{2-[methyl-(2-methylamino-ethyl)-amino]-ethyl}-carbamoyl)-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **7c** was performed following the procedure reported for the synthesis of compound **1c**, using *N*₁,*N*₂-dimethyl-*N*₁-(2-(methylamino)ethyl)ethane-1,2-diamine **7b** instead of Methyl-[2-(2-methylamino-ethoxy)-ethyl]-amine. After purification by flash chromatography with a gradient of ammonia in methanol 7M in EtOAc (5 to 15%), 100 mg of the desired product **7c** were obtained as a yellow oil; *m/z* = 615 (*M*+*H*)⁺.


Step 4





The synthesis of **7d** was performed following the procedure reported for the synthesis of compound **1d**, using 13-cyclohexyl-3-methoxy-6-(methyl-{2-[methyl-(2-methylamino-ethyl)-amino]-ethyl}-carbamoyl)-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **7c** instead of 13-cyclohexyl-3-methoxy-6-{methyl-[2-(2-methylamino-ethoxy)-ethyl]-carbamoyl}-7H-benzo[3,4]azepino[1,2-a]indole-10-carboxylic acid *tert*-butyl ester **1c**. After purification by flash chromatography with a gradient of methanol in EtOAc (0 to 10%), 50 mg of the desired product **7d** were obtained; *m/z* = 694 (*M*+*H*)⁺.

Step 5

The synthesis of **7e** was performed following the procedure reported for the synthesis of compound **1e**, using intermediate **7d** instead of intermediate **1d**; $m/z = 638$ ($M+H$)⁺.


Step 6

The synthesis of product **7** is being performed using the procedure described for the synthesis of compound **1**, using intermediate **7e** instead of intermediate **1e**.

Example 8 – synthesis of compound 8Step 1

The synthesis of **8b** was performed following the procedure reported for the synthesis of compound **1b**, using methyl ester **8a** instead of **1a**. The desired product **8b** was obtained in 95% yield as a light yellow solid; $m/z = 502$ ($M+H$)⁺.

Step 2




At 0 °C and under protective atmosphere, oxalyl chloride (4.07 ml, 47.4 mmol) was added to a solution of carboxylic acid **8b** (19.83 g, 39.5 mmol) and DMF (5 drops) in tetrahydrofuran (dry) (100 mL). Upon addition of oxalyl chloride immediate formation of gas was observed. The reaction was stirred at 0 °C for 1.5 hour. Then, an extra amount of 0.5 eq of oxalyl chloride was added and the reaction was stirred for 1 more hour (repeated once until full conversion was obtained). The reaction was then evaporated to dryness in vacuo to afford 20.5 g (97%) of the acid chloride **8c** as a white solid; m/z (methyl ester formed by addition of methanol prior to analysis) = 516 ($M+H$)⁺.

10

15

Step 3




To a solution of (*S*)-4-benzyl-2-oxazolidinone (7.50 g, 42.3 mmol) in tetrahydrofuran (dry) (60 ml) under nitrogen atmosphere *n*-butyllithium (26.4 ml, 42.3 mmol) was added slowly at -78°C. The reaction mixture was stirred for 40 minutes at -78°C. After 40 minutes, the anion solution was added via a canula to a solution of the acid chloride **8c** (20 g, 38.5 mmol) in 60 mL THF at -78°C. The reaction mixture was stirred for 1.5 hours at -78°C. When the reaction was finished, it was quenched with an ammonia chloride solution at -70°C. The reaction mixture was then warmed up to room temperature and extracted with EtOAc, washed with brine and dried over Na₂SO₄. The organic layer was filtered and concentrated to afford 26.34 g of a yellow solid. The two

20

25

enantiomers **8d** and **8e** were separated by flash column chromatography using 5:1 Heptane/EtOAc and were obtained as light yellow solids; $m/z = 661$ ($M+H$)⁺.

Step 4



5


8d**8f**

Diastereoisomer **8d** (11.17 g, 16.90 mmol) was first dissolved in THF (130 ml) then methanol (130 ml) was added. 1N NaOH solution (101 mL, 101 mmol) was added slowly so that the temperature was kept below 30°C. The reaction mixture was stirred at room temperature for 2 hours. When the reaction was finished 1 N HCl solution was added until the pH reached 2. 500 mL H₂O was then added and the reaction mixture was extracted with EtOAc, washed with brine and concentrated. Purification by flash column chromatography using 1:1 Heptane/EtOAc afforded 5.24 g (60%) of the desired enantiomer (4b*R*,5a*S*)-9-(*tert*-butoxycarbonyl)-12-cyclohexyl-3-methoxy-4b,5,5a,6-tetrahydrobenzo[3,4]cyclopro[5,6]azepino[1,2-*a*]indole-5a-carboxylic acid **8f** with an ee of 97%; $m/z = 502$ ($M+H$)⁺.

10

15

Step 5

**8f****8g**


To a stirred solution of intermediate **8f** (2 g, 3.99 mmol) in dry DMF (50 mL) at 0°C, were added di-isopropyl ethylamine (DIEA, 1.54 g, 11.9 mmol), HATU (2.27 g, 5.98 mmol) and 2,2'-oxybis(*N*-methylethanamine) (2.1 g, 15.95 mmol). The resulting mixture was stirred at 0°C for 1 hour then kept at room temperature for 12 hours. The reaction mixture was then successively poured into an iced water solution, extracted with dichloromethane, dried over MgSO₄ then concentrated. The residue was purified by column chromatography using a gradient of methanol in dichloromethane (0 to 10%) to yield 0.94 g (38% yield) of the desired product *tert*-butyl(1a*R*,12b*S*)-

20

25

8-cyclohexyl-11-methoxy-1a-(methyl{2-[2-(methylamino)ethoxy]ethyl} carbamoyl)-1,1a,2,12b-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylate **8g** as a white solid; m/z 616 ($M+H$)⁺.


Step 6



The synthesis of (1*aR*,12*bS*)-8-cyclohexyl-11-methoxy-1a-(methyl{2-[2-(methylamino)ethoxy]ethyl} carbamoyl)-1,1a,2,12b-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]-benzazepine-5-carboxylic acid **8h** was performed following the procedure reported for the preparation of compound **1e**, using intermediate **8g** instead of **1d**. The obtained residue was further dissolved in DCM, washed with water, dried over magnesium sulfate, filtered and concentrated to dryness, leading to 0.474 g (56% yield) of title compound **8h**; m/z = 560 ($M+H$)⁺.


10

Step 7





To a solution of intermediate **8h** (0.474 g, 0.847 mmol) in dioxane (10 mL) was added sulfamide (0.814 g, 8.47 mmol). The resulting mixture was stirred at 100°C in a microwave oven for 4 hours. The reaction mixture was then cooled down to room temperature and concentrated. The residue was purified by column chromatography using a gradient of methanol in dichloromethane (0 to 10%) to give 143 mg (26%) of the title product (1*aR*,12*bS*)-1a-[(2-{2-[(aminosulfonyl)(methyl)amino]ethoxy}ethyl) (methyl)carbamoyl]-8-cyclohexyl-11-methoxy-1,1a,2,12b-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylic acid **8i**; m/z = 639 ($M+H$)⁺.

20

Step 8


The synthesis of (1a*R*,12b*S*)-8-cyclohexyl-11-methoxy-16,22-dimethyl-1,12b-dihydro-
 5 5,1a-(methanoiminothioiminoethanoxyethanoiminomethano)cyclopropa[*d*]-
 indolo[2,1-*a*][2]benzazepine-13,23(2*H*)-dione 15,15-dioxide **8** was performed
 following the 2-step procedure reported for the synthesis of compound **1**, using
 intermediate **8i** instead of **1e**, yielding to 90 mg (52% yield) of a white solid; $m/z = 621$
 (M+H)⁺. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.3 - 1.5 (m, 3 H) 1.75 - 1.8
 10 (m, 5 H) 1.85 - 2.05 (m, 6 H) 2.5-3 (m, 3 H) 3.2 (s, 3 H) 3.22 (s, 3H) 3.4-3.7 (m, 6 H)
 3.87 (s, 3 H) 3.75-3.9 (m, 1 H) 4.9-5.1 (m, 1 H) 6.95-7.16 (d, $J=8.39$ Hz, 1 H) 7.28 (s, 1
 H) 7.44-7.55 (m, 2 H) 7.80 (d, $J=8.39$ Hz, 1 H) 9.4 (br. s., 1 H).

Example 9 – synthesis of compound 9Step 1

Enantiomer (4b*S*,5a*R*)-9-(*tert*-butoxycarbonyl)-12-cyclohexyl-3-methoxy-4b,5,5a,6-
 20 tetrahydrobenzo[3,4]cyclopropa[5,6]azepino[1,2-*a*]indole-5a-carboxylic acid **9a** was

obtained in 27% yield, and 96% ee, following the procedure reported for the synthesis of compound **8f**, starting from the diastereoisomer **8e** instead of **8d**; $m/z = 502 (M+H)^+$.

Step 2




5

9a**9b**

tert-butyl(1*aS*,12*bR*)-8-cyclohexyl-11-methoxy-1*a*-(methyl{2-[2-(methylamino)ethoxy]ethyl}carbamoyl)-1,1*a*,2,12*b*-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylate **9b** was prepared in 60% yield from **9a** and 2,2'-oxybis(*N*-methylethanamine) following the procedure used for the preparation of compound **1c**; $m/z = 616 (M+H)^+$.

10

Step 3


**9b****9c**

To a solution of intermediate **9b** (0.73 g, 1.185 mmol) in dioxane (10 mL) was added sulfamide (1.14 g, 11.85 mmol). The resulting mixture was stirred at 100°C in a microwave oven for 3 hours. The reaction mixture was cooled down to room temperature then concentrated. The residue was purified by column chromatography using a gradient of methanol in dichloromethane (0 to 10%) to yield 743 mg (80%) of the title product *tert*-butyl(1*aS*,12*bR*)-1*a*-[(2-{2-[(aminosulfonyl)(methyl)amino]ethoxy}ethyl)(methyl)carbamoyl]-8-cyclohexyl-11-methoxy-1,1*a*,2,12*b*-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylate **9c**.

15


20

Step 4





The synthesis of (1a*S*,12b*R*)-1a-[(2-{2-[(aminosulfonyl)(methyl)amino]ethoxy} ethyl)(methyl)carbamoyl]-8-cyclohexyl-11-methoxy-1,1a,2,12b-tetrahydrocyclopropa [d]indolo[2,1-*a*][2]benzazepine-5-carboxylic acid **9d** was performed following the procedure reported for the preparation of compound **1e**, using intermediate **9c** instead of **1d**, yielding 517 mg (79% yield) of a brownish foam; $m/z = 639$ ($M+H$)⁺.


Step 5



The synthesis of (1a*S*,12b*R*)-8-cyclohexyl-11-methoxy-16,22-dimethyl-1,12b-dihydro-5,1a-(methanoiminothioiminoethanoxyethanoiminomethano)cyclopropa[d]indolo-[2,1-*a*][2]benzazepine-13,23(2*H*)-dione 15,15-dioxide **9** was performed following the 2-step procedure reported for the synthesis of compound **1**, using intermediate **9d** instead of **1e**, yielding to 80 mg (16% yield) of a white solid; $m/z = 621$ ($M+H$)⁺. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.3 - 1.5 (m, 3 H) 1.75 - 1.8 (m, 5 H) 1.85 - 2.05 (m, 6 H) 2.5-3 (m, 3 H) 3.2 (s, 3 H) 3.22 (s, 3H) 3.4-3.7 (m, 6 H) 3.87 (s, 3 H) 3.75-3.9 (m, 1 H) 4.9-5.1 (m, 1 H) 6.95-7.16 (d, $J=8.39$ Hz, 1 H) 7.28 (s, 1 H) 7.44-7.55 (m, 2 H) 7.80 (d, $J=8.39$ Hz, 1 H) 9.4 (br. s., 1 H).

Example 10 – synthesis of compound 10Step 1


To a solution of 5-*tert*-butyl 1a-methyl 8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a,5(2*H*)-dicarboxylate **8a** (2 g, 3.88 mmol) in dichloromethane (25 mL) was added TFA (22.34 g, 194 mmol). The resulting mixture was stirred at room temperature for 6 hours then concentrated to dryness. The residue was successively dissolved in dichloromethane, washed with water, dried under MgSO_4 , filtered and concentrated. The residue was then purified by column chromatography using dichloromethane and ethyl acetate as eluent to yield 1.7 g (95%) of the title product 8-cyclohexyl-11-methoxy-1a-(methoxycarbonyl)-1,1a,2,12b-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylic acid **10a** as a white powder; m/z 460 ($\text{M}+\text{H}$)⁺.

Step2

1,1'-Carbonyldiimidazole (0.847 g, 5.22 mmol) was added to a stirred solution of 8-cyclohexyl-11-methoxy-1a-(methoxycarbonyl)-1,1a,2,12b-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylic acid **10a** (0.8 g, 1.74 mmol) in THF (15 mL) at 25°C. The evolution of CO_2 was instantaneous and when it slowed the solution was heated at 50°C for 2 hours and then cooled to room temperature. *Tert*-butyl{4-[(amino-sulfonyl)(methyl)amino]butyl}carbamate **10b** (0.735 g, 2.61 mmol) was added


followed by the addition of DBU (0.53 g, 3.48 mmol). Stirring was continued for 12 hours at 50°C. The mixture was cooled to room temperature then partitioned between dichloromethane and water. The water was extracted with dichloromethane. The organic layers were dried over MgSO₄ then concentrated to dryness. The residue was purified by column chromatography using dichloromethane and ethyl acetate to yield 0.66 g (53%) of the title product methyl 5-({[4-[(*tert*-butoxycarbonyl)amino]butyl} (methyl)amino]sulfonyl}carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylate **10c** as a white foam; m/z 723 (M+H)⁺.

10 Step3



To a solution of intermediate **10c** (0.65 g, 0.899 mmol) in THF (20 mL) was added lithium hydroxide (0.75 g, 1.8 mmol) dissolved in water (5 mL). The resulting mixture was stirred at room temperature overnight then diluted with water and neutralized with a 2M HCl aqueous solution. The resulting mixture was extracted with dichloromethane, dried over MgSO₄ then concentrated. The residue was purified by column chromatography using a gradient of methanol in CH₂Cl₂ to yield 0.55 g (86%) of the title product 5-({[4-[(*tert*-butoxycarbonyl)amino]butyl} (methyl)amino]sulfonyl}-carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid **10d** as a white solid; m/z 709 (M+H)⁺.


Step 4



25 TFA (2.5 g, 22 mmol) was added to a solution of intermediate **10d** (0.52 g, 0.734 mmol) in DCM (10 mL). The resulting mixture was stirred at RT for approximately 10 hours. The reaction was then evaporated to dryness and the residue was purified by column chromatography using a gradient of methanol in DCM to afford 0.3 g (68%) of the title compound 5-({[4-aminobutyl} (methyl)amino]sulfonyl}carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid.


(methylamino)sulfonyl carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid **10e** as a TFA salt; *m/z* 609 (M+H)⁺


Step 5




To a stirred solution of intermediate **10e** (0.22 g, 0.36 mmol) in dry DMF (100 mL), at 0°C, were added DIPEA (0.14 g, 1.08 mmol) and HATU (0.206 g, 0.542 mmol). The resulting mixture was stirred at 0°C for 1 hour then kept at room temperature for 12 hours. The reaction mixture was then successively poured into an iced watered solution, extracted with dichloromethane, dried over MgSO₄ and then concentrated. The residue was purified by column chromatography to yield 0.188 g (88%) of the title product 8-cyclohexyl-11-methoxy-16-methyl-1,12b-dihydro-5,1a-(methanoiminothioiminobutanoimino)cyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-13,22(2*H*)-dione 15,15-dioxide **10** as a white solid. ¹H NMR (DMSO-*d*₆) : 11.5 (s, 1H), 8.4 (s, 1H), 8.3 (s, 1H), 7.8 (d, *J*=8.2 Hz, 1H), 7.3 (d, *J*=8.2 Hz, 1H), 7.25 (d, *J*=8.4 Hz, 1H), 7.15 (s, 1H), 7 (d, *J*=8.4 Hz, 1H), 5.6 (d, *J*=16 Hz, 1H), 3.85 (s, 3H), 3.55 (d, *J*=16 Hz, 1H), 3-3.2 (m, 2 H), 3 (s, 3H), 2.7-2.9 (m, 4H), 1.8-2.1 (m, 5H), 1.6-1.7 (m, 2H), 1.3-1.6 (m, 6H), 1-0.7 (m, 3H); *m/z* 609 (M+H)⁺

Example 11 – synthesis of compound 11



Step 1


1,1'-carbonyldiimidazole (0.522 g, 3.22 mmol) was added to a stirred solution of 8-cyclohexyl-11-methoxy-1a-(methoxycarbonyl)-1,1a,2,12b-tetrahydrocyclopropa-
 5 [d]indolo[2,1-a][2]benzazepine-5-carboxylic acid **10a** (0.74 g, 1.61 mmol) in THF (15 mL) at 25°C. The evolution of CO₂ was instantaneous and when it slowed the solution was heated at 50°C for 2 hours and then cooled to room temperature. *Tert*-butyl 4-{{2-[(aminosulfonyl)(methyl)amino]ethyl}piperazine-1-carboxylate} **11a** (1.038
 10 g, 3.22 mmol) was added followed by the addition of DBU (0.49 g, 3.22 mmol). Stirring was continued for 12 hours at 50°C. The mixture was cooled to room temperature then partitioned between dichloromethane and water. The water was extracted with dichloromethane and the organic layers were dried over MgSO₄ then concentrated to dryness. The residue was purified by column chromatography using
 15 dichloromethane and ethyl acetate to yield 0.83 g (68%) of the title compound methyl 5-({[2-[4-(*tert*-butoxycarbonyl)piperazin-1-yl]ethyl}(methyl)amino]sulfonyl}carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[d]indolo[2,1-a][2]benzazepine-1a(2*H*)-carboxylate **11b** as a white foam; *m/z* 764 (M+H)⁺

20 Step 2

To a solution of intermediate **11b** (0.6 g, 0.785 mmol) in THF (20 mL) was added LiOH (0.82 g, 1.96 mmol) in water (5 mL). The resulting mixture was stirred at room
 25 temperature overnight then diluted with water and neutralized with a 2M HCl aqueous solution. The resulting mixture was extracted with dichloromethane, dried over MgSO₄ then concentrated. The resulting residue was purified by column chromatography using CH₂Cl₂ and methanol to yield 0.4 g (68%) of the title compound 5-({[2-[4-(*tert*-butoxycarbonyl)piperazin-1-yl]ethyl}(methyl)amino]sulfonyl}carbamoyl)-

8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid **11c** as a white solid; m/z 750 ($M+H$)⁺


Step 3



TFA (1.44 g, 12.7 mmol) was added to a solution of intermediate **11c** (0.38 g, 0.507 mmol) in dichloromethane (10 mL). The resulting mixture was stirred at RT for approximately 10 hours. The reaction was then evaporated to dryness and the residue was purified by column chromatography using dichloromethane and methanol to afford the title compound 8-cyclohexyl-11-methoxy-5-({[methyl(2-piperazin-1-yl-ethyl)amino]sulfonyl}carbamoyl)-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]-benzazepine-1a(2*H*)-carboxylic acid **11d** (0.24 g, 73 %); m/z 650 ($M+H$)⁺

10



Step 4




To a stirred solution of intermediate **11d** (0.24 g, 0.37 mmol) in dry DMF (100 mL), at 0°C, were added di-isopropyl ethylamine (0.143 g, 1.1 mmol) and HATU (0.211 g, 0.554 mmol). The resulting mixture was stirred at 0°C for 1 hour then kept at room temperature for 12 hours. The reaction mixture was then successively poured into an iced watered solution, extracted with dichloromethane, dried over MgSO₄ and concentrated. The residue was purified by column chromatography using dichloromethane / methanol to yield 0.018 g (18%) of the title compound 31-cyclohexyl-8-methoxy-22-methyl-21-thia-1,13,20,22,25-pentaazaheptacyclo-[23.2.2.1^{3,13}.1^{12,15}.1^{14,18}.0^{3,5}.0^{6,11}]dotriaconta-6,8,10,12(31),14(30),15,17-heptaene-2,19-dione 21,21-dioxide **11** as a white solid; m/z 632 ($M+H$)⁺

20

25

Example 12 – synthesis of compound 12Step 1


To a solution of 5-*tert*-butyl-1a-methyl-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a,5(2*H*)-dicarboxylate **8a** (2 g, 3.88 mmol) in dichloromethane (25 mL) was added trifluoroacetic acid (22.34 g, 194 mmol). The resulting mixture was stirred at room temperature for 6 hours then concentrated to dryness. The residue was successively dissolved in dichloromethane, washed with water, dried under MgSO₄, filtered and concentrated. The residue was purified by column chromatography using dichloromethane and ethyl acetate as eluent to yield 1.7 g (95%) of the title product 8-cyclohexyl-11-methoxy-1a-(methoxycarbonyl)-1,1a,2,12b-tetrahydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-5-carboxylic acid **12a** as a white powder; *m/z* 460 (M+H)⁺

Step 2

To a solution of intermediate **12a** (1.73 g, 3.76 mmol) in THF (25 mL) at 0°C were added successively 4-dimethylaminopyridine (DMAP) (1.38 g, 3.76 mmol), *N*1-((ethylimino)methylene)-*N*3,*N*3-dimethylpropane-1,3-diamine hydrochloride (EDC) (2.16 g, 11.29 mmol) and allyl(methyl)aminosulfonamide **12b** (1.3 g, 8.66 mmol). The resulting mixture was stirred at 0°C for 2 h then at room temperature for 8 h. Water was then added and the reaction mixture was filtered. The resulting solid


was purified by column chromatography using dichloromethane and ethyl acetate to yield 500 mg (23%) of the title product methyl 5-({[allyl(methyl)amino]sulfonyl}-carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]-benzazepine-1a(2*H*)-carboxylate **12c**; m/z 592 ($M+H$)⁺

5 Step 3



To a solution of intermediate **12c** (0.5 g, 0.845 mmol) in THF (20 mL) was added lithium hydroxide (0.73 g, 1.69 mmol) in water (5 mL). The resulting mixture was stirred at room temperature overnight then diluted with water and neutralized with a 2M HCl aqueous solution. The resulting mixture was extracted with dichloromethane, dried over MgSO₄ then concentrated. The resulting residue was purified by column chromatography using CH₂Cl₂ and methanol to yield 0.4 g (75%) of the title product 5-({[allyl(methyl)amino]sulfonyl}carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid **12d** as a white solid; m/z 578 ($M+H$)⁺


20 Step 4



To a solution of intermediate **12d** (0.2 g, 0.346 mmol) in THF (15 mL), at 0°C, were added successively 4-dimethylaminopyridine (DMAP) (0.127 g, 1.04 mmol), *N*1-((ethylimino)methylene)-*N*3,*N*3-dimethylpropane-1,3-diamine hydrochloride (EDC) (0.199 g, 1.04 mmol) and but-3-en-1-amine (0.062 g, 0.866 mmol). The resulting mixture was stirred at 0°C for 2 h then at room temperature for 8 h. Water was then added and the resulting mixture was filtered. The solid was washed with dichloromethane then the filtrate was successively extracted with dichloromethane, dried over MgSO₄, filtered and concentrated. The resulting residue was purified by column chromatography with dichloromethane and ethyl acetate to yield 70 mg (32%)

of the title product N^5 -{[allyl(methyl)amino]sulfonyl}- N^{1a} -but-3-en-1-yl-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a,5(2*H*)-dicarboxamide **12e**; m/z 631 ($M+H$)⁺

Step 5




A solution of intermediate **12e** (0.1 g, 0.16 mmol) in dichloroethane (50 mL) was degassed with argon for 10 minutes then Hoveyda-Grubbs 1st generation catalyst (0.03 mg, 0.032 mmol) was added. The resulting mixture was warmed to 70°C and kept under argon overnight. The mixture was then cooled down to room temperature and the solvent was removed under vacuum. The resulting dark residue was purified by column chromatography using DCM and ethyl acetate to give 15 mg (16%) of the title product 8-cyclohexyl-11-methoxy-16-methyl-1,12b-dihydro-5,1a-(methaniminothioiminopent[2]enoiminomethano)cyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-13,23(2*H*)-dione 15,15-dioxide **12** as a white solid; m/z 603 ($M+H$)⁺


10

15

Example 13 – synthesis of compound 13



Step 1




13-Cyclohexyl-3-methoxy-7*H*-benzo[3,4]azepino[1,2-*a*]indole-6,10-dicarboxylic acid 10-*tert*-butyl ester 6-methyl ester **1a** (1 g, 1 eq) was dissolved in dry dichloromethane under N₂, followed by the addition of trifluoroacetic acid (TFA) (8.88 mL, 60 eq). The

solution was stirred at RT for 24 h. The solvent was then removed under reduced pressure. The crude product was triturated with diethyl ether. The crystals were filtered off and dried under vacuum overnight to afford the title product 13-Cyclohexyl-3-methoxy-7H-benzo[3,4]azepino[1,2-a]indole-6,10-dicarboxylic acid 6-methyl ester


5 **13a** (89%, 0.86 g); LC-MS: Rt. 3.19 min., m/z 446 $[M+H]^+$.

Step 2




- 13-Cyclohexyl-3-hydroxy-7H-benzo[3,4]azepino[1,2-a]indole-6,10-dicarboxylic acid 6-methyl ester **13a** (0.86 g, 1 eq), *N*-methyl-*N*-allyl-sulfuric diamide **12b** (0.67 g, 2.03 eq), *N*¹-((ethylimino)methylene)-*N*³,*N*³-dimethylpropane-1,3-diamine hydrochloride (EDCI) (1.14 g, 3.06 eq) and dimethyl-pyridin-4-yl-amine (DMAP) (0.67 g, 3.04 eq) were dissolved in dry dimethylformamide (20 ml) under N₂. The solution was stirred at RT for 3 days. This solution was slowly added into ice water.
- 15 The water layer was extracted with ethyl acetate (3 x 50 ml) and washed with tetrahydrofurane (3 x 50 ml). The combined organic layers were dried over magnesium sulfate, filtered and evaporated under reduced pressure. The crude product was purified by preparative HPLC to give 0.63 g (55%) of the title product **13b**; LC-MS: Rt. 6.16 min., m/z 578 $[M+H]^+$. ¹H-NMR (DMSO) δ (ppm) 1.13-1.20 (m, 1H, CH₂), 1.30-1.47 (m, 3H, CH₂ (2x)), 1.62-1.78 (m, 2H, CH₂), 1.81-1.93 (m, 1H, CH₂), 1.93-2.12 (m, 3H, CH₂ (2x)), 2.70-2.82 (m, 1H, CH), 2.86 (s, 3H, CH₃N), 3.79 (s, 3H, CH₃O), 3.88 (s, 3H, CH₃O), 3.90-3.98 (m, 2H, CH₂), 4.21 (d, 1H, J = 12.97 Hz, CH₂), 5.21 (d, 1H, J = 10.15 Hz, CH₂), 5.31 (d, 1H, J = 17.13 Hz, CH₂), 5.61 (d, 1H, J = 13.12 Hz, CH₂), 5.78-5.90 (m, 1H, CH_{arom}), 7.25 (dd, 1H, J = 2.50 and J = 8.60 Hz, CH_{arom}), 7.32-7.35 (m, 1H, CH_{arom}), 7.54 (d, 1H, J = 8.60 Hz, CH_{arom}), 7.61 (d, 1H, J = 8.45 Hz, CH_{arom}), 7.88 (d, 1H, J = 9.01 Hz, CH_{arom}), 7.91 (s, 1H, CH), 8.31-8.34 (brs, 1H, NHSO₂).
- 20
25

Step 3




Compound **13b** (0.60 g, 1 eq) was dissolved in a mixture of tetrahydrofurane:methanol (1:1) (20 ml), followed by the addition of a LiOH solution in water (0.09 g, 2 eq). The solution was stirred overnight at RT for several days. The solvents were then evaporated under reduced pressure and the water layer was acidified with a 3 N HCl solution until pH 2. The resulting crystals were filtered off, washed with water and isopropyl ether and dried under vacuum overnight to afford 0.44 g (74%) of the title product **13c**; LC-MS: Rt. 5.84 min., m/z 562 $[M-H]^-$.

Step 4




Compound **13c** (0.44 g, 1 eq) and HATU (0.47 g, 1.6 eq) were dissolved in dimethylformamide under N_2 , followed by the addition of DIPEA (0.15 g, 0.20 ml, 1.5 eq) and allylamine (0.07 ml, 1.2 eq). The solution was stirred at RT for 3 days. The dimethylformamide solution was then slowly poured into ice water. The resulting crystals were filtered off, washed with water and dried under vacuum overnight to afford 0.47 g (100%) of the title product **13d**; LC-MS: Rt. 3.01 min., m/z 603 $[M+H]^+$.

Step 5




1.62-1.78 (m, 2H, CH₂), 1.82-1.93 (m, 1H, CH₂), 1.93-2.12 (m, 3H, CH₂ (2x)), 2.65-2.90 (m, 4H, CH and CH₃N), 3.56 (d, 2H, *J* = 18.10 Hz, CH₂), 3.80-3.97 (brs, 5H, CH₂ and CH₃O), 4.21 (d, 1H, *J* = 15.12 Hz, CH₂), 4.28-4.46 (m, 1H, CH), 5.72 (d, 1H, *J* = 14.15 Hz, CH₂), 5.78-5.88 (m, 1H, CH), 6.53 (s, 1H, CH_{arom}), 7.18-7.28 (m, 2H, CH_{arom} (2x)), 7.39-7.49 (m, 1H, CH_{arom}), 7.55 (d, 1H, *J* = 8.38 Hz, CH_{arom}), 7.62 (s, 1H, CH_{arom}), 7.72-7.84 (m, 1H, NH), 8.29 (s, 1H, CH), 8.51-8.62 (brs, 1H, NH).

Example 14 – synthesis of compound 14




Step 1



13-Cyclohexyl-3-methoxy-7H-benzo[3,4]azepino[1,2-a]indole-6,10-dicarboxylic acid 6-methyl ester **13a** (0.60 g, 1 eq) was dissolved in dry acetonitrile (50 ml) under N₂, followed by the addition of diimidazol-1-yl-methanone (CDI) (0.66 g, 3 eq). The solution was stirred overnight at 50°C. The solvent was then evaporated under reduced pressure and the crude product was purified by flash chromatography on elution with heptane:acetonitrile and finally ethyl acetate. The product was recrystallized from ethyl acetate to give 0.50 g (75%) of the title product **14a**.


Step 2




Compound **14a** (0.50 g, 1 eq) was dissolved in dry acetonitrile (50 ml), followed by the addition of *tert*-butyl 4-(2-(sulfamoylamino)ethyl)piperazine-1-carboxylate **14b**

- (0.47 g, 1.50 eq) and 2,3,4,6,7,8,9,10-octahydro-pyrimido[1,2-a]azepine (DBU) (0.31 g, 2 eq). The solution was heated at 50°C overnight, then evaporated under reduced pressure. The resulting residue was stirred in a 0.1 N citric acid water solution. The crystals were filtered off and dried under vacuum overnight. The product was
- 5 purified by column chromatography on elution with dichloromethane to remove the first impurity. The other obtained fractions were added together. This product was further purified by flash chromatography on elution with dichloromethane:methanol (100 to 99:1) to afford 0.41 g (55%) of the title product **14c**; LC-MS: Rt. 5.59 min., m/z 736 $[M+H]^+$. 1H -NMR ($CDCl_3$) δ (ppm) 1.18-1.34 (m, 1H, CH_2), 1.35-1.50 (brs, 10H, CH_2 and $C(CH_3)_3$), 1.70-1.85 (m, 3H, CH_2 (2x)), 1.90-2.12 (m, 5H, CH_2 (3x)), 2.30-2.41 (m, 4H, CH_2 (2x)), 2.52-2.62 (m, 2H, CH_2), 2.77-2.90 (m, 1H, CH), 3.13-3.22 (m, 2H, CH_2), 3.43-3.57 (m, 4H, CH_2 (2x)), 3.83 (s, 3H, CH_3O), 3.92 (s, 3H, CH_3O), 4.16-4.23 (m, 1H, CH_2), 5.58-5.69 (m, 1H, CH_2), 7.00 (d, 1H, J = 2.54 Hz, CH_{arom}), 7.11 (dd, 1H, J = 2.67 and J = 8.59 Hz, CH_{arom}), 7.48 (d, 1H, J = 8.44 Hz, CH_{arom}), 7.53 (d, 1H, J = 8.61 Hz, CH_{arom}), 7.83 (s, 1H, CH_{arom}), 7.90 (d, 1H, J = 8.48 Hz, CH_{arom}), 8.09 (s, 1H, CH).
- 15


Step 3




- 20 Compound **14c** (0.41 g, 1 eq) was dissolved in dry dichloromethane (10 ml) under N_2 followed by the addition of trifluoroacetic acid (1.30 ml, 30 eq). The solution was stirred at RT overnight. The solvent was then removed under reduced pressure and the crude product was stirred in diethyl ether. The resulting crystals were filtered off and
- 25 dried under reduced pressure to afford 0.31 g (87%) of the title product **14d**; LC-MS: RT. 3.81 min., m/z 634 $[M-H]^+$.


Step 4

Compound **14d** (0.31 g, 1 eq) was dissolved in a mixture of tetrahydrofuran:methanol (1:1), followed by the addition of 50% NaOH-water solution (1 ml). The solution was stirred at RT overnight then evaporated under reduced pressure. The water layer was acidified with acetic acid to pH 4, and extracted with ethyl acetate (7 x 50 ml). The combined ethyl acetate layers were dried over sodium sulfate, filtered off and evaporated under reduced pressure to obtain the desired compound **14e** as a yellow powder (0.30 g, 100%); LC-MS: Rt. 3.64 min., m/z 622 $[M+H]^+$.


Step 5

The synthesis of the title compound **14** is being performed following the procedure reported for the synthesis of compound **11**, using intermediate **14e** instead of **11d**.


Example 15 – synthesis of compound 15

Step 1


Compound **13a** (0.20 g, 1 eq) was dissolved in dry acetonitrile under N₂, followed by the addition of CDI (0.1 g, 1.3 eq). The solution was stirred at 60°C for 1 h. According to TLC, the reaction went to completion. DBU (0.10 ml, 1.52 eq) and diaminosulfuric diamide **15a** (0.29 g, 2 eq) were then added. The solution was stirred at 60°C for 3 h, then was evaporated under reduced pressure. A citric acid water solution (0.1 N) cooled in ice, was added to the crude product. The residual solution was extracted with ethyl acetate (3 x 50 ml). The combined organic layers were washed with brine (50 ml), dried over sodium sulfate, filtered off and evaporated under reduced pressure to afford 0.21 g (62%) of the title product **15b**; LC-MS: Rt: 5.63 min., *m/z* 750 [M+H]⁺.

Step 2



The synthesis of compound **15c** was performed following the procedure reported for the synthesis of compound 5-({[2-[4-(*tert*-butoxycarbonyl)piperazin-1-yl]ethyl}(methyl)amino)sulfonyl}carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid (**11c**), using intermediate **15b** instead of methyl 5-({[2-[4-(*tert*-butoxycarbonyl)piperazin-1-yl]ethyl}(methyl)amino)sulfonyl}carbamoyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylate (**11b**); *m/z* 736 [M+H]⁺.

Step 3


The synthesis of the title compound **15d** was performed following the procedure reported for the synthesis of compound 8-cyclohexyl-11-methoxy-5-({[methyl-(2-piperazin-1-ylethyl)amino]sulfonyl}carbonyl)-1,12b-dihydrocyclopropa[*d*]-indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid (**11d**), using intermediate **15c** instead of 5-({[2-[4-(*tert*-butoxycarbonyl)piperazin-1-yl]ethyl}(methyl)amino]-sulfonyl}carbonyl)-8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid (**11c**), yielding to 448 mg (quantitative yield) of the desired product; m/z 636 $[M+H]^+$.

Step 4


The synthesis of the title compound **15** was performed following the procedure reported for the synthesis of compound 31-cyclohexyl-8-methoxy-22-methyl-21-thia-1,13,20,22,25-pentaazaheptacyclo[23.2.2.1^{3,13}.1^{12,15}.1^{14,18}.0^{3,5}.0^{6,11}]dotriaconta-6,8,10,12(31),14(30),15,17-heptaene-2,19-dione 21,21-dioxide **11**, using intermediate **15d** instead of 8-cyclohexyl-11-methoxy-5-({[methyl(2-piperazin-1-ylethyl)amino]-sulfonyl}carbonyl)-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a(2*H*)-carboxylic acid **11d**, yielding 150 mg (34% yield) of a cream solid; m/z 618 $[M+H]^+$. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.06 - 1.18 (m, 1 H) 1.19 - 1.31 (m, 2 H) 1.31 - 1.50 (m, 2 H) 1.62 - 1.78 (m, 2 H) 1.81 - 1.93 (m, 1 H) 1.93 - 2.10 (m, 2 H) 2.53 - 3.21 (m, 12 H) 3.31 - 3.67 (m, 4H) 3.86 (s, 3 H) 4.33 - 4.51 (m, 1 H) 4.99 - 5.16 (m, 1 H) 7.06 - 7.14 (m, 2 H) 7.17 (d, $J=8.02$ Hz, 1 H) 7.52 (d, $J=8.22$ Hz, 1 H) 7.55 - 7.68 (m, 1 H) 7.77 (m, 1 H) 8.39 (m, 1 H).

Example 16 – synthesis of compound 16Step 1


A solution of 50% NaOH w/w in water (9.31 g) was added to a stirred solution of **16a** (3.0 g, 5.82 mmol) in THF (100 mL) and MeOH (150 mL). After 1 hour the reaction mixture was concentrated under reduced pressure, and subsequently diluted with ice-cold water (150 mL). The pH of the resulting solution was adjusted to 6 with diluted HCl. A precipitate was formed, which was collected by filtration, washed with water and dried under vacuum to give 3.17g (89%) of **16b** as a yellowish powder. The product was used without any further purification in the next step; $m/z = 502$ (M+H)⁺.

Step 2


HATU (3.6 g, 9.48 mmol) was added under nitrogen to a stirred solution of **16b** (3.17 g, 6.32 mmol), DIPEA (3.3 mL, 3 eq) and 2,2'-oxybis(*N*-methylethanamine) (3.34 g, 4 eq) in 60 mL of dry THF. After 1 h, the reaction mixture was quenched with water (100 mL) and extracted with ethyl acetate (EtOAc). The organic layer was successively dried (Na₂SO₄), filtered and evaporated. The residue was triturated in water, filtered and dried to give 4.05 g (quantitative yield) of the target compound **16c**, used directly in the next step: $m/z = 616$ (M+H)⁺

Step 3


- 5 A solution of **16c** (3.90g, 6.33 mmol) and sulfamide (3.04g, 6 eq) in dioxane (100 mL) was refluxed at 100°C overnight. The reaction mixture was cooled down to room temperature, then evaporated under vacuum. The residue was redissolved in DCM, washed with water, dried over magnesium sulfate, filtered and concentrated to give 4.48 g (quantitative yield) of the desired product **16d**, used directly in the next step: $m/z = 695 (M+H)^+$


10 Step 4

- 15 TFA (14.7 g, 129 mmol) was added to a solution of **16d** (4.48 g, 6.45 mmol) in dichloromethane (50 mL). After 1h, the reaction mixture was concentrated under vacuum. The residue was triturated in ether, filtered and washed with ether, then purified by chromatography (gradient EtOAc to EtOAc/EtOH, 9:1) to give 3.05 g (68%) of the desired product **16e**: $m/z = 639 (M + H)^+$


Step 5

- Carbonyldiimidazole (1.07 g, 6.59 mmol) was added to a stirred solution of **16e** (3.05 mg, 4.39 mmol) in dry ACN (40 mL). The reaction mixture was stirred at 60°C for 1h: complete conversion to the acyl imidazole intermediate was observed. The resulting solution was cooled down to RT, diluted with dry ACN (300 mL) and DBU (1.34 g, 2 eq) was added. The reaction mixture was stirred overnight at room temperature, then concentrated under reduced pressure. The residue was redissolved in DCM, washed with water, dried, filtered and concentrated. Purification by column chromatography (gradient DCM to DCM/MeOH 9:1) provided 930 mg (33%) of the title product **16** as a white powder: $m/z = 621$ ($M + H^+$), 1H NMR (400 MHz, CHLOROFORM- d) δ ppm 1.15 - 1.31 (m, 1 H) 1.31 - 1.52 (m, 3 H) 1.69 - 1.81 (m, 2 H) 1.84 (s, 3 H) 1.88 - 2.13 (m, 7 H) 2.45 (d, $J=14.87$ Hz, 1 H) 2.76 - 2.92 (m, 1 H) 3.14 (s, 3 H) 3.40 (d, $J=15.65$ Hz, 1 H) 3.54 - 3.70 (m, 3 H) 3.81 - 3.90 (m, 1 H) 3.93 (s, 3 H) 4.03 - 4.18 (m, 1 H) 4.37 (d, $J=14.67$ Hz, 1 H) 4.64 - 4.80 (m, 2 H) 7.06 (d, $J=8.80$ Hz, 1 H) 7.09 (s, 1 H) 7.48 (d, $J=8.22$ Hz, 1 H) 7.57 (s, 1 H) 7.70 (d, $J=8.22$ Hz, 1 H) 7.89 (d, $J=8.41$ Hz, 1 H) 10.01 (br. s., 1 H)


Example 17 – synthesis of compound 17

Step 1


5 The synthesis of the title compound **17a** was performed following the procedure reported for the synthesis of compound **16c**, using N,N' -dimethylbutane-1,4-diamine instead of 2,2'-oxybis(N -methylethanamine), yielding 1.25 g (quant. yield) of a white solid; m/z 600 $[M+H]^+$.

Step 2


The synthesis of the title compound **17b** was performed following the procedure reported for the synthesis of compound **16d**, using compound **17a** instead of compound **16c**, yielding 1 g (54% yield) of a slightly yellow solid; m/z 679 $[M+H]^+$.

Step 3


15 The synthesis of the title compound **17c** was performed following the procedure reported for the synthesis of compound **16e**, using compound **17b** instead of compound **16d**, yielding 538 mg (62% yield) of a slightly brown solid; m/z 623 $[M+H]^+$.

Step 4


- The synthesis of the title compound **17** was performed following the procedure reported for the synthesis of compound **16**, using compound **17c** instead of compound **16e**, yielding 70 mg (15% yield) of a white solid; m/z 605 $[M+H]^+$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ ppm 1.08 - 1.20 (m, 1 H) 1.22 - 1.79 (m, 13 H) 1.88 (s, 6 H) 2.40 - 2.47 (m, 1 H) 2.69 - 2.83 (m, 1 H) 2.92 - 3.14 (m, 4 H) 3.56 - 3.72 (m, 1 H) 3.89 (s, 3 H) 3.92 - 4.04 (m, 1 H) 4.26 (d, $J=14.67$ Hz, 1 H) 4.86 (d, $J=14.09$ Hz, 1 H) 7.18 (dd, $J=8.61, 2.15$ Hz, 1 H) 7.22 (d, $J=2.15$ Hz, 1 H) 7.46 - 7.57 (m, 2 H) 7.80 - 7.92 (m, 1 H) 8.48 (s, 1 H) 11.39 (br. s., 1 H)

Example 18 – synthesis of compound 18

- The synthesis of the title compound **18** was performed following the 4-step procedure reported for the synthesis of compound **17**, starting from intermediate **1b** instead of **16b**, and yielding 0.5 g of a white solid; m/z 591 $[M+H]^+$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ ppm 1.01 - 1.19 (m, 1 H) 1.18 - 1.52 (m, 5 H) 1.54 - 1.79 (m, 4 H) 1.80 - 2.08 (m, 4 H) 2.42 - 2.48 (m, 1 H) 2.63 - 2.80 (m, 1 H) 2.93 (s, 3 H) 2.98 - 3.14 (m, 1 H) 3.43 - 3.75 (m, 5 H) 3.85 (s, 3 H) 4.43 (d, $J=14.87$ Hz, 1 H) 5.04 (d, $J=14.48$ Hz, 1 H) 6.84 (br. s., 1 H) 7.09 (s, 1 H) 7.18 (d, $J=8.22$ Hz, 1 H) 7.45 (d, $J=8.22$ Hz, 1 H) 7.55 (d, $J=8.41$ Hz, 1 H) 7.87 (d, $J=8.41$ Hz, 1 H) 8.35 (br. s., 1 H) 11.33 (br. s., 1 H)

Example 19 – synthesis of compound 19**19**


The synthesis of the title compound **19** was performed following the 5-step procedure reported for the synthesis of compound **1**, starting from intermediate 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-fluoro-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **19a** instead of 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **1a**, and yielded 180 mg of a white solid; *m/z* 595 [M+H]⁺. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.11 - 1.29 (m, 1 H) 1.29 - 1.53 (m, 3 H) 1.67 - 1.83 (m, 3 H) 1.87 - 2.11 (m, 4 H) 2.30 (br. s., 3 H) 2.69 - 2.82 (m, 1 H) 2.81 - 2.98 (m, 1 H) 3.11 (s, 3 H) 3.46 - 3.58 (m, 1H) 3.59 - 3.79 (m, 3 H) 3.90 - 4.08 (m, 1 H) 4.24 - 4.38 (m, 1 H) 4.43 (dd, *J*=14.73, 1.27 Hz, 1 H) 4.97 (d, *J*=14.63 Hz, 1 H) 6.73 (s, 1 H) 7.11 (dd, *J*=9.27, 2.63 Hz, 1 H) 7.17 - 7.30 (m, 1 H) 7.57 (dd, *J*=8.68, 5.76 Hz, 1 H) 7.69 (s, 1 H) 7.67 (dd, *J*=8.78, 1.56 Hz, 1 H) 7.90 (d, *J*=8.78 Hz, 1 H) 9.84 (br. s., 1 H)

Example 20 – synthesis of compound 20**20**

The synthesis of the title compound **20** was performed following the 5-step procedure reported for the synthesis of compound **1**, starting from intermediate 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-fluoro-5-methyl-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **20a** instead of 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **1a**, and yielded 130 mg of a white solid; *m/z* 609 [M+H]⁺. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.14 - 1.31 (m, 1 H) 1.32 - 1.46 (m, 3 H) 1.64 - 1.81 (m, 3 H) 1.84 (s, 3 H) 1.87 - 1.99 (m, 3 H) 2.01 (s, 3 H) 2.47 (d, *J*=14.63 Hz, 1 H) 2.73 - 2.87 (m, 1 H) 3.14 (s, 3 H) 3.43 (d, *J*=15.02 Hz, 1 H) 3.56 - 3.64 (m, 2 H) 3.65 (d, *J*=3.12 Hz, 1 H) 3.74 - 3.88 (m, 1 H)

4.00 - 4.12 (m, 1H) 4.35 (d, $J=14.83$ Hz, 1 H) 4.64 - 4.75 (m, 1 H) 4.81 (d, $J=14.63$ Hz, 1 H) 7.16 - 7.32 (m, 2 H) 7.53 (dd, $J=8.39, 6.05$ Hz, 1 H) 7.64 (s, 1 H) 7.70 (d, $J=8.39$ Hz, 1 H) 7.91 (d, $J=8.39$ Hz, 1 H) 10.09 (br. s., 1 H)

Example 21 – synthesis of compound 21




5

21

The synthesis of the title compound **21** was performed following the 5-step procedure reported for the synthesis of compound **1**, starting from intermediate 10-*tert*-butyl 6-methyl 3-chloro-13-cyclohexyl-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **21a** instead of 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-


10 *a*][2]benzazepine-6,10-dicarboxylate **1a**, and yielding 270 mg of a white solid; m/z 611 $[M+H]^+$. 1H NMR (400 MHz, DMSO- d_6) δ ppm 1.08 - 1.22 (m, 1 H) 1.31 - 1.52 (m, 3 H) 1.63 - 1.78 (m, 2 H) 1.81 - 2.09 (m, 4 H) 2.50 (s, 3H) 2.69 - 2.80 (m, 1 H) 3.00 (s, 3 H) 3.08 - 3.19 (m, 1 H) 3.19 - 3.28 (m, 1 H) 3.46 - 3.88 (m, 6 H) 4.52 (d, $J=14.87$ Hz, 1 H) 5.12 (d, $J=13.11$ Hz, 1 H) 6.97 (s, 1 H) 7.49 (d, $J=7.83$ Hz, 1 H) 7.58 - 7.70 (m, 15 3 H) 7.94 (d, $J=8.61$ Hz, 1 H) 8.36 (s, 1 H) 11.39 (br. s., 1 H)

Example 22 – synthesis of compound 22




22

The synthesis of the title compound **22** was performed following the 5-step procedure reported for the synthesis of compound **1**, using *N*¹,*N*⁶-dimethylhexane-1,6-diamine instead of 2,2'-oxybis(*N*-methylethanamine) in step 2, and yielded 50 mg of a white solid; m/z 619 $[M+H]^+$. 1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.00 - 1.64 (m, 11 H) 1.66 - 1.87 (m, 3 H) 1.87 - 2.15 (m, 4H) 2.47 (s, 3 H) 2.66 - 2.91 (m, 2 H) 3.23 (s, 3 H) 3.25 - 3.33 (m, 1 H) 3.33 - 3.45 (m, 1 H) 3.90 (s, 3 H) 4.09 - 4.25 (m, 1 H) 4.39 (d, $J=14.28$ Hz, 1 H) 5.14 (d, $J=14.48$ Hz, 1 H) 6.81 (s, 1 H) 6.90 (s, 1 H) 20 7.06 (dd, $J=8.61, 2.15$ Hz, 1 H) 7.45 (d, $J=8.22$ Hz, 1 H) 7.50 (d, $J=8.61$ Hz, 1 H) 25 7.81 - 7.96 (m, 2 H) 8.94 (br. s., 1 H)


Example 23 – synthesis of compound 23**23**

The synthesis of the title compound **23** was performed following the 5-step procedure reported for the synthesis of compound **1**, using *N*¹,*N*²-dimethyl-*N*¹-(2-



- 5 (methylamino)ethyl)ethane-1,2-diamine instead of 2,2'-oxybis(*N*-methylethanamine) in step 2, and yielded 20 mg of a white solid; *m/z* 592 [*M*+*H*]⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.04 (d, *J*=5.87 Hz, 1 H) 1.06 - 1.22 (m, 1 H) 1.27 - 1.51 (m, 3 H) 1.60 - 1.78 (m, 2 H) 1.80 - 1.92 (m, 1 H) 1.92 - 2.07 (m, 3 H) 2.12 (s, 3 H) 2.27 - 2.41 (m, 1 H) 2.69 - 2.83 (m, 2 H) 2.83 - 2.97 (m, 2 H) 3.01 - 3.15 (m, 2 H) 3.17 - 3.28 (m, 2 H) 3.86 (s, 3 H) 4.21 (d, *J*=15.65 Hz, 1 H) 5.54 (d, *J*=15.65 Hz, 1 H) 7.11 - 7.25 (m, 2H) 7.35 (s, 1 H) 7.47 (d, *J*=8.22 Hz, 1 H) 7.53 (d, *J*=9.00 Hz, 1 H) 7.70 - 7.83 (m, 1 H) 8.32 (br. s., 1 H) 8.37 - 8.50 (m, 1 H)
- 10

Example 24 – synthesis of compound 24**24**

- 15 The synthesis of the title compound **24** was performed following the 4-step procedure reported for the synthesis of compound **17**, starting from intermediate 10-(*tert*-butoxycarbonyl)-2-chloro-13-cyclohexyl-7*H*-indolo[2,1-*a*][2]benzazepine-6-carboxylic acid **24b** instead of 10-(*tert*-butoxycarbonyl)-13-cyclohexyl-3-methoxy-5-methyl-7*H*-indolo[2,1-*a*][2]benzazepine-6-carboxylic acid **16b**, and yielded 0.25 g of a white
- 20 solid; *m/z* 595 [*M*+*H*]⁺. ¹H NMR (400 MHz, Chloroform-*d*) δ ppm 1.25 - 1.5 (m, 4 H) 1.5 - 1.8 (m, 4 H) 1.9 - 2.1 (m, 4 H) 1.8 (s, 3 H) 2.8-2.13 (m, 3 H) 2.5 - 2.6 (m, 2 H) 3.2 (s, 3 H) 3.6 (br. s., 1 H) 4.1 (br. s., 1H) 4.45 (d, *J*=15 Hz, 1 H) 5 (d, *J*=15 Hz, 1 H) 6.6 (s, 1 H) 7.25 (d, *J*=8.4 Hz, 1 H) 7.4 (dd, *J*=8.5, *J*=2.5 Hz, 1 H) 7.5-7.6 (m, 2H) 7.69 (s, 1 H) 7.9 (d, *J*=8.4 Hz, 1 H) 9.1 (br. s., 1 H)


Example 25 – synthesis of compound 25**25**

The synthesis of the title compound **25** was performed following the 5-step procedure reported for the synthesis of compound **10**, starting from intermediate 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **1a** instead of 5-*tert*-butyl 1a-methyl 8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropano[*d*]indolo[2,1-*a*][2]benzazepine-1a,5(2*H*)-dicarboxylate **8a**, and yielded 45 mg of a white solid; m/z 577[M+H]⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.03 - 1.19 (m, 1 H) 1.25 - 1.49 (m, 4 H) 1.49 - 2.29 (m, 10 H) 2.67 - 2.82 (m, 1 H) 2.84 - 3.04 (m, 1 H) 3.05 - 3.24 (m, 1 H) 3.48 - 3.72 (m, 5 H) 3.86 (s, 3 H) 4.42 (d, $J=14.67$ Hz, 1 H) 5.00 (d, $J=14.28$ Hz, 1 H) 6.84 (br. s., 1 H) 7.09 (s, 1 H) 7.18 (d, $J=8.41$ Hz, 1 H) 7.47 (d, $J=7.83$ Hz, 1 H) 7.55 (d, $J=8.41$ Hz, 1 H) 7.75 - 7.92 (m, 1 H) 8.19 - 8.41 (m, 1 H) 11.27 (br. s., 1 H)

Example 26 – synthesis of compound 26**26**Step 1

A solution of 606 mg (1.24 mmole) of **1b**, 410 mg (1.1 eq) of **26a**, 710 mg (1.5 eq) of HATU and 0.65 mL (3 eq) of diisopropylethyl amine in dry DMF (10 mL) was stirred at RT during 1h. The RM was then diluted with water and the resulting yellow precipitate was filtered off, washed with water, and purified by flash chromatography (eluant DCM to DCM/MeOH 0.5%) to give a quantitative yield of the desired product **26b** as a yellow powder; m/z 771 $[M+H]^+$.

Step 2




10 **26b**

26c

To a solution of 1.1g (1.44 mmole) of **26b** and thiophenol (0.32 g, 2 eq) in dry DMF (15 mL) was added cesium carbonate (0.94 g, 2 eq) at RT. After 2h, the RM was diluted with water and extracted with EtOAc. The organic layer was washed with brine, dried over magnesium sulfate, filtered and concentrated. The resulting residue was further purified by flash chromatography (eluant: DCM to DCM/ NH₃ in MeOH 85/15) to give 0.77 g (90% yield) of **26c** as a yellow powder; m/z 586 $[M+H]^+$.


Step 3




26c

26d


A mixture of **26c** (0.72g, 1.23 mmole) and sulfamide (0.35 g, 3 eq) in dioxane (15 mL) was refluxed until completion (~7h). The RM was then concentrated under vacuo and the residue was triturated in DCM. The resulting precipitate of sulfamide in excess was filtered off. The organic layer was concentrated and purified by flash chromatography (eluant: DCM to DCM/MeOH 1%) to give 776 mg (95 % yield) of the desired product **26d** as a light yellow powder; m/z 665 $[M+H]^+$.

Step 4

A solution of **26d** (0.72 g, 1.086 mmole) in 10 mL of HCl in isopropanol and 5 mL of DCM was stirred at RT for 3h. The RM was then concentrated under vacuo, and the residue was triturated in diethyl ether. The resulting precipitate was filtered off, washed with ether and dried in a vacuum oven overnight to give 661 mg (97% yield) of the desired product **26e** as a light yellow powder; m/z 609 $[M+H]^+$.


Step 5

A solution of **26e** (0.6g, 0.971 mmole) and CDI (0.205 g, 1.3 eq) in acetonitrile (10 mL) was heated at 60°C until complete formation of the acyl imidazole intermediate (~1h). The RM was then diluted with 20 mL of acetonitrile and DBU (0.296 g, 2 eq) was added at RT. The RM was stirred at RT until completion, then was concentrated. The residue was redissolved in water and acetic acid was then added until pH 2. The resulting precipitate was filtered off, washed with water, and purified by flash chromatography (eluant: DCM to DCM/MeOH 5%) to give 0.315 g (55% yield) of the desired product **26** as a slightly yellow powder; m/z 591 $[M+H]^+$. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.07 - 2.09 (m, 16 H) 2.71 - 2.84 (m, 1 H) 2.94 (s, 3 H) 3.01 - 3.18 (m, 2 H) 3.19 - 3.31 (m, 2 H) 3.87 (s, 3 H) 4.25 (d, $J=15.06$ Hz, 1 H) 5.52 (d, $J=15.26$ Hz, 1 H) 7.16 - 7.26 (m, 2H) 7.32 - 7.44 (m, 2 H) 7.54 (d, $J=9.19$ Hz, 1 H) 7.86 (d, $J=8.61$ Hz, 1 H) 8.26 (s, 1 H) 8.40 - 8.51 (m, 1 H) 11.61 (br. s., 1 H)


Example 27 – synthesis of compound 27

The synthesis of the title compound **27** was performed following the 5-step procedure reported for the synthesis of compound **26**, starting from intermediate 10-(*tert*-



- 5 butoxycarbonyl)-2-chloro-13-cyclohexyl-7*H*-indolo[2,1-*a*][2]benzazepine-6-carboxylic acid **24b** instead of 10-(*tert*-butoxycarbonyl)-13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-*a*][2]benzazepine-6-carboxylic acid **1b**, and yielded 85 mg of a yellow solid; m/z 596 $[M+H]^+$. ¹H NMR (400 MHz, Chloroform-*d*) δ ppm 1.21 - 1.5 (m, 10 H) 1.75 - 1.8 (m, 2 H) 1.9 - 2.1 (m, 4 H) 2.75 (br. s., 1 H) 3.01 (s, 3 H) 3.1 - 3.2 (m, 2 H) 3.5 - 3.6 (m, 2 H) 4.23 (dd, $J=15.28, 1.27$ Hz, 1 H) 5.6 (d, $J=15.28$ Hz, 1 H) 7.4 (s, 1 H) 7.5-7.6 (m, 3 H) 7.65 (d, $J=8.5$ Hz, 1 H) 7.8 (s, 1 H) 7.9 (d, $J=8.5$ Hz, 1 H) 7.69 (s, 1 H) 8.64 (br. s., 1 H)
- 10

Example 28 – synthesis of compound 28


- 15 The synthesis of the title compound **28** was performed following the 5-step procedure reported for the synthesis of compound **10**, starting from intermediate 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **1a** instead of 5-*tert*-butyl 1a-methyl 8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a,5(2*H*)-dicarboxylate **8a**, and
- 20 using *N*-[4-(methylamino)butyl]-*N*-(1-methylethyl)sulfamide **28a** instead of *N*-(4-aminobutyl)-*N*-methylsulfamide **10b**, and yielded 50 mg of the desired product **28**; m/z 619 $[M+H]^+$. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.05 - 1.15 (m, 1 H) 1.18 (d, $J=6.65$ Hz, 3 H) 1.25 (d, $J=6.46$ Hz, 3 H) 1.28 - 1.51 (m, 4 H) 1.53 - 2.31 (m, 13 H) 2.67 - 2.85 (m, 1 H) 3.01 - 3.19 (m, 1 H) 3.51 - 3.73 (m, 1H) 3.89 (s, 3 H) 3.95 -
- 25 4.15 (m, 1 H) 4.42 (d, $J=14.48$ Hz, 1 H) 4.52 - 4.72 (m, 1 H) 5.01 (d, $J=14.48$ Hz, 1 H) 6.68 (s, 1 H) 6.87 (s, 1 H) 7.05 (d, $J=8.41$ Hz, 1 H) 7.52 (d, $J=8.41$ Hz, 1 H) 7.63 (d, $J=8.22$ Hz, 1 H) 7.77 -7.99 (m, 2 H) 9.42 (br. s., 1 H)

Example 29 – synthesis of compound 10b


A mixture of *tert*-butyl 4-(methylamino)butylcarbamate (4 g, 19.77 mmol) and sulfuric diamide (7.6 g, 4 eq) in dioxane (10 mL) was heated at 100°C in a microwave oven during 30 minutes. The RM was then concentrated in vacuo, and DCM was added. The resulting white precipitate of sulfuric diamide in excess was filtered off, and the filtrate was successively washed with diluted HCl, then brine, dried over magnesium sulfate, filtered and concentrated. Trituration in diisopropyl ether afforded 3.55 g (64% yield) of *tert*-butyl 4-(methyl(sulfamoyl)amino)butylcarbamate **10b** as a white solid; *m/z* 282 [M+H]⁺.

Example 30 – synthesis of compound 26aStep 1


To a solution of *tert*-butyl 5-aminopentylcarbamate **30a** (20 g, 99 mmol) and 2-nitrobenzene-1-sulfonyl chloride **30b** (23 g, 1.05 eq) in DCM (200 mL) was added dropwise diisopropyl ethyl amine (19.2 g, 1.5 eq) at 0°C. After stirring at RT overnight, the RM was successively washed with an aqueous solution of citric acid, then brine, dried over magnesium sulfate, filtered and concentrated. Trituration in diisopropyl ether afforded 32.61 g (85% yield) of *tert*-butyl 5-(2-nitrophenylsulfonamido)pentylcarbamate **30c** as a white solid; *m/z* 388 [M+H]⁺.


Step 2

To a mixture of *tert*-butyl 5-(2-nitrophenylsulfonamido)pentylcarbamate **30c** (32.61 g, 84 mmol) and potassium carbonate (13.96 g, 1.2 eq) in acetone (300 mL) was added methyl iodide (5.5 mL, 1.05 eq). After stirring at RT overnight, more methyl iodide (1 eq) and potassium carbonate (0.6 eq) was added and the RM was stirred at RT until completion. The RM was then diluted with water and extracted with DCM. The organic layer was separated, washed with brine, dried over magnesium sulfate, filtered and concentrated. Trituration in diisopropyl ether afforded 31.59 g (93% yield) of *tert*-butyl 5-(*N*-methyl-2-nitrophenylsulfonamido)pentylcarbamate **30d** as a white solid; m/z 402 $[M+H]^+$.


Step 3

A solution of *tert*-butyl 5-(*N*-methyl-2-nitrophenylsulfonamido)pentylcarbamate **30d** (31.5 g, 79 mmol) and trifluoroacetic acid (29.2 mL, 5 eq) in DCM (300 mL) was stirred at RT until completion (~16 h). The RM was then concentrated under vacuo, redissolved in DCM, washed with a saturated sodium bicarbonate aqueous solution (2 times), then brine, dried over magnesium sulfate, filtered and concentrated. Trituration in diisopropyl ether afforded 23.7 g (quantitative yield) of *N*-(5-aminopentyl)-*N*-methyl-2-nitrobenzenesulfonamide **26a** as a slightly yellow solid; m/z 302 $[M+H]^+$.


Example 31 – synthesis of compound 28a


Step 1

- A mixture of *tert*-butyl 4-aminobutyl(methyl)carbamate **31a** (287 mg, 1.42 mmole), acetone (75 mg, 1.29 mmole) and sodium triacetoxymethylborohydride (383 mg, 1.8 mmole) was stirred under nitrogen at RT until completion. The RM was then concentrated, diluted with a saturated sodium bicarbonate aqueous solution, and extracted with ether (2 times). The organic layers were combined, dried over magnesium sulfate, filtered and concentrated to give 200 mg (63% yield) of the desired product *tert*-butyl 4-(isopropylamino)butyl(methyl)carbamate **31b**, used without further purification in the next step; m/z 245 $[M+H]^+$.


Step 2

- A mixture of *tert*-butyl 4-(isopropylamino)butyl(methyl)carbamate **31b** (3.38 g, 13.8 mmole) and sulfamic acid (3.99 g, 3 eq) in dioxane (10 mL) was heated at 110°C in a microwave oven during 60 minutes. The RM was then concentrated in vacuo, and DCM was added. The resulting white precipitate of sulfamic acid in excess was filtered off, and the filtrate was concentrated in vacuo. The residue was purified by flash chromatography (eluant: DCM to DCM/MeOH 20%) to give 1.7 g (38% yield) of the desired product *tert*-butyl 4-(isopropyl(sulfamoyl)amino)butyl(methyl)carbamate **28a**; m/z 324 $[M+H]^+$.

Example 32 – synthesis of compound 19a


Step 1**32a****32b**

A mixture of *tert*-butyl 2-bromo-3-cyclohexyl-1H-indole-6-carboxylate **32a** (5 g, 13.22 mmoles, synthesized as described in US 2007270406 A1), pinacolborane (5.75 mL, 3 eq) and triethylamine (7.35 mL, 4 eq) in THF (50 mL) was stirred at RT during 3 h. Palladium acetate (90 mg, 0.03 eq) and biphenyl-2-ylidicyclohexylphosphine (556 mg, 0.12 eq) were then added and the RM was heated at 80°C during 2h. The reaction mixture was then allowed to cool down to RT and poured in a solution of watered NH₄Cl then extracted with ethyl acetate. The organic layers were dried over magnesium sulfate, filtered and concentrated. The residue was purified by column chromatography using a gradient of ethyl acetate in heptane to give 3.5 g (70% yield) of the desired product *tert*-butyl 3-cyclohexyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-6-carboxylate **32b**; *m/z* 426 [M+H]⁺.

15 Step 2**32b****32c****32d**


To a mixture of *tert*-butyl 3-cyclohexyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-6-carboxylate **32b** (2.77 g, 6.5 mmoles) and 2-bromo-5-fluorobenzaldehyde **32c** (1.58 g, 1.2 eq) in DME (40 mL) was added a solution of sodium carbonate (2.07 g, 3 eq) in water (15 mL). The resulting mixture was then flushed with nitrogen at RT during 10 minutes. After the addition of palladium tetrakis triphenylphosphine (376 mg, 0.05 eq), the RM was heated at 70°C during 1 h. The mixture was then allowed to cool down to RT and poured in water, then extracted with ethyl acetate (3 times). The organic layers were combined, dried over MgSO₄, filtered and concentrated. The residue was recrystallized from di-isopropyl ether / heptane to give 2 g (73% yield) of the desired product *tert*-butyl 3-cyclohexyl-2-(4-fluoro-2-formylphenyl)-1H-indole-6-carboxylate **32d** as a white solid; *m/z* 422 [M+H]⁺.

Step 3


**32d****19a**

- A mixture of *tert*-butyl 3-cyclohexyl-2-(4-fluoro-2-formylphenyl)-1H-indole-6-carboxylate **32d** (2 g, 4.75 mmol), cesium carbonate (1.85 g, 1.2 eq) and methyl 2-(dimethoxyphosphoryl)acrylate (16.475 mL, 0.36 M solution in toluene, 1.25 eq) in DMF (80 mL) was stirred at 60°C during 2 h. The reaction mixture was then allowed to cool down to room temperature, poured in water and extracted with ethyl acetate. The organic layer was then dried over MgSO₄, filtered and concentrated. The residue was purified by column chromatography using heptanes / dichloromethane to give 2 g (86% yield) of the desired product 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-fluoro-7H-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **19a**; *m/z* 490 [M+H]⁺..

Example 33 – synthesis of compound 20a

**20a**


Step 1

**33a****33b**

- To a solution of 2-bromo-5-fluorobenzonitrile **33a** (10 g, 50 mmol) in dry tetrahydrofuran (100 mL) under nitrogen was added methylmagnesium bromide (3.2 M in ether, 19 mL, 60.0 mmol), and the resulting mixture was heated to reflux for 4 hours.


The RM was then cooled down to RT, poured into a 2 N HCl solution (100 mL) and then diluted with methanol (100 mL). The resulting green solution was concentrated on a steam bath for 1 h at which point the organic solvents had been removed and the crude product had precipitated. The reaction mixture was then extracted with ethyl acetate, dried over MgSO₄ and concentrated. The residue was purified by column chromatography using heptane and dichloromethane to give 4.88 g (45% yield) of the desired product 1-(2-bromo-5-fluorophenyl)ethanone **33b** as a pink oil; *m/z* 218 [M+H]⁺.

Step 2




The title product *tert*-butyl 2-(2-(2-acetyl-4-fluorophenyl)-3-cyclohexyl-1H-indole-6-carboxylate **33c** was synthesized following the procedure reported for the synthesis of *tert*-butyl 3-cyclohexyl-2-(4-fluoro-2-formylphenyl)-1H-indole-6-carboxylate **32d**, using 1-(2-bromo-5-fluorophenyl)ethanone **33b** instead of 2-bromo-5-fluorobenzaldehyde **32c**, and was obtained in 65% yield as a white solid; *m/z* 436 [M+H]⁺.

Step 3




The title product 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-fluoro-5-methyl-7H-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **20a** was synthesized following the procedure reported for the synthesis of 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-fluoro-7H-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **19a**, using *tert*-butyl 2-(2-(2-acetyl-4-fluorophenyl)-3-cyclohexyl-1H-indole-6-carboxylate **33c** instead of *tert*-butyl 3-cyclohexyl-2-(4-fluoro-2-formylphenyl)-1H-indole-6-carboxylate **32d**, and was obtained in 11% yield as a white solid; *m/z* 504 [M+H]⁺.

Example 34 – synthesis of compound 21a



Step 1




5 32a

34a

34b

The bromo indole derivative **32a** (5 g, 13.22 mmol), 4-chloro-2-formylphenylboronic acid **34a** (3.17 g, 17.18 mmol) and potassium carbonate (4.20 g, 30.4 mmol) were dissolved in 100mL of 1,2-dimethoxyethane (80 ml)/water (20 ml) 4/1 and the obtained solution was flushed thoroughly with argon. Then bis(triphenylphosphine)palladium(II) chloride (0.464 g, 0.661 mmol) was added and the reaction was heated to 63°C under argon during 3h. The reaction was then diluted with EtOAc, washed with water and with sat. aq. NaHCO₃, dried (brine, sulfate) and evaporated. The residue was stripped with DIPE and stirred and sonicated in heptane with a few mL DIPE added. The solid was filtered off and dried to afford 4.97 g (86% yield) of the desired product *tert*-butyl 2-(4-chloro-2-formylphenyl)-3-cyclohexyl-1H-indole-6-carboxylate **34b**; *m/z* 437 [M+H]⁺.


Step 2



20 The indole derivative **34b** (4.95 g, 11.30 mmol) and cesium carbonate (4.42 g, 13.56 mmol) were dissolved in *N,N*-dimethylformamide (dry) (50 ml) and methyl 2-(dimethoxyphosphoryl)acrylate (3.23 g, 14.13 mmol) was added. The RM was stirred at 65°C during 2h. It was then cooled to rt and dropped onto 300 ml of water


vigorously stirred. The resulting yellowish solid was filtered off, washed with water and dried to afford 5.40 g (94% yield) of the desired product 10-*tert*-butyl 6-methyl 3-chloro-13-cyclohexyl-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **21a**, used without further purification in the next step; m/z 507 $[M+H]^+$.

5 Example 35 – synthesis of compound 24a





The title compound **24a** was synthesized following the 2-step procedure reported for the synthesis of 10-*tert*-butyl 6-methyl 3-chloro-13-cyclohexyl-7*H*-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **21a**, using 5-chloro-2-formylphenylboronic acid in the first step, instead of 4-chloro-2-formylphenylboronic acid **34a**, and was obtained
10 in the first step, instead of 4-chloro-2-formylphenylboronic acid **34a**, and was obtained with an overall yield of 70% as a yellowish solid; m/z 507 $[M+H]^+$.


Example 36 – synthesis of compound 24b



15 A solution of NaOH (6.38 g) in 25 mL of water was added to a stirred solution of the indole derivative **24a** in THF (100 mL) and MeOH (150 mL). After 1 hour the reaction was concentrated under reduced pressure, then diluted with ice-cold water (150 mL). The pH of the resulting solution was adjusted to 6 with HCl then extracted with dichloromethane and dried over MgSO_4 . The solvent was removed then the residue was
20 purified by column chromatography using DCM/MeOH as eluant to give 1.7 g (87% yield) of a yellowish solid; m/z 492 $[M+H]^+$.

Example 37 – synthesis of compound 16aStep 1


Ethane-1,2-diol (4.06 g) and Tos-OH (0.41 g) were added to a solution of 1-(2-bromo-5-methoxyphenyl)ethanone **37a** (5 g) in toluene (950 ml). The solution was heated under reflux with stirring in a 3-neck round-bottomed flask fitted with a Dean-Stark receiver for 3 hours. The reaction mixture was then cooled to room temperature. The mixture was transferred to a separating funnel and a sodium carbonate solution (1 M, 50 ml) was added. The mixture was agitated and two phases formed. The organic layer was separated, washed with water (2 x 50 ml), dried over MgSO_4 and concentrated under vacuum to afford 6.5 g (quantitative yield) of the desired product 2-(2-bromo-5-methoxyphenyl)-2-methyl-1,3-dioxolane **37b** as a white solid.

15 Step 2

The bromo derivative **37b** (6 g) was dissolved in dry THF (60 ml) and the solution was cooled down to -78°C . Then n-BuLi (16.5 ml) was added carefully at such a rate that the temperature did not exceed -60°C . After 1 h, B(O-i-Pr)_3 (6.2 g) was added neatly dropwise at -78°C . After all was added, the cooling bath was removed. The mixture was stirred at 0°C for 2.5 h, then 2 N HCl (60 ml) was added, and the RM stirred at r.t. for 2 h. The organic solvent was then removed under vacuum and the

aqueous layer was saturated with NaCl and extracted with EtOAc. The organic layer was dried over Na₂SO₄ and evaporated under vacuum to give 3 g of the desired product 2-acetyl-4-methoxyphenylboronic acid **37c**.


Step 3



The title product *tert*-butyl 2-(2-acetyl-4-methoxyphenyl)-3-cyclohexyl-1H-indole-6-carboxylate **37d** was synthesized by following a similar procedure to that used for the synthesis of *tert*-butyl 2-(4-chloro-2-formylphenyl)-3-cyclohexyl-1H-indole-6-carboxylate **34b**, using 2-acetyl-4-methoxyphenylboronic acid **37c** instead of 4-chloro-2-formylphenylboronic acid **34a**.

10


Step 4



The title product 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-methoxy-5-methyl-7H-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **16a** was synthesized by following a similar procedure to that used for the synthesis of 10-*tert*-butyl 6-methyl 13-cyclohexyl-3-fluoro-5-methyl-7H-indolo[2,1-*a*][2]benzazepine-6,10-dicarboxylate **20a**, using *tert*-butyl 2-(2-acetyl-4-methoxyphenyl)-3-cyclohexyl-1H-indole-6-carboxylate **37d** instead of *tert*-butyl 2-(2-acetyl-4-fluorophenyl)-3-cyclohexyl-1H-indole-6-carboxylate **33c**.

15

20

Example 38 – synthesis of compound 38**38**

The synthesis of the title compound **38** was performed following the 5-step procedure reported for the synthesis of compound **10**, starting from intermediate 13-cyclohexyl-3-methoxy-7*H*-benzo[3,4]azepino[1,2-*a*]indole-6,10-dicarboxylic acid 10-*tert*-butyl ester 6-methyl ester **1a** instead of 5-*tert*-butyl 1a-methyl 8-cyclohexyl-11-methoxy-1,12b-dihydrocyclopropa[*d*]indolo[2,1-*a*][2]benzazepine-1a,5(2*H*)-dicarboxylate **8a**, and yielded 60 mg of a beige solid; *m/z* 577 [M+H]⁺.

Example 39 – Activity of compounds of formula (I)**10 Replicon assay**

The compounds of formula (I) were examined for activity in the inhibition of HCV RNA replication in a cellular assay. The assay demonstrated that the compounds of formula (I) inhibited a HCV functional cellular replicating cell line, also known as HCV replicons. The cellular assay was based on a bicistronic expression construct, as described by Lohmann et al. (1999) Science vol. 285 pp. 110-113 with modifications described by Krieger et al. (2001) Journal of Virology 75: 4614-4624, in a multi-target screening strategy. In essence, the method was as follows.

The assay utilized the stably transfected cell line Huh-7 luc/neo (hereafter referred to as Huh-Luc). This cell line harbors an RNA encoding a bicistronic expression construct comprising the wild type NS3-NS5B regions of HCV type 1b translated from an Internal Ribosome Entry Site (IRES) from encephalomyocarditis virus (EMCV), preceded by a reporter portion (FfL-luciferase), and a selectable marker portion (neo^R, neomycine phosphotransferase). The construct is bordered by 5' and 3' NTRs (non-translated regions) from HCV type 1b. Continued culture of the replicon cells in the presence of G418 (neo^R) is dependent on the replication of the HCV RNA. The stably transfected replicon cells that express HCV RNA, which replicates autonomously and to high levels, encoding *inter alia* luciferase, are used for screening the antiviral compounds.

The replicon cells were plated in 384 well plates in the presence of the test and control compounds which were added in various concentrations. Following an incubation of three days, HCV replication was measured by assaying luciferase activity (using standard luciferase assay substrates and reagents and a Perkin Elmer ViewLux™ ultraHTS microplate imager). Replicon cells in the control cultures have high luciferase expression in the absence of any inhibitor. The inhibitory activity of the compound was monitored on the Huh-Luc cells, enabling a dose-response curve to be generated for each test compound. EC50 values were then calculated, which value represents the amount of the compound required to decrease by 50% the level of detected luciferase activity, or more specifically, the ability of the genetically linked HCV replicon RNA to replicate.

Enzymatic assay

1. HCV NS5B 1bJ4

1.a) Protein purification

The cDNA encoding NS5B amino acid 1-570 (HC-J4, genotype 1b, pCV-J4L6S, genebank accession number AF054247) was subcloned into the Nhe I and Xho I restriction sites of pET-21b. Expression of the subsequent His-tagged C-terminal 21 amino acid deleted NS5B was performed as follows:

The NS5B expression construct was transformed into *E. coli* BL21(DE3) (Novagen, Madison, WI). Five milliliter of LB-medium supplemented with ampicillin (50 µg/mL) was inoculated with one colony. When the pre-culture reached an optical density of 0.6 measured at 600 nm, it was transferred to fresh LB-medium supplemented with ampicillin, at a ratio of 1:200. Cells were grown to an optical density at 600 nm of 0.6, after which the expression cultures were shifted to a growth temperature of 20°C following induction with isopropyl-1-thio-β-D-galactopyranoside and MgCl₂ at a final concentration of 0.4 mM and 10 µM, respectively. After 10 h of induction, cells were harvested by centrifugation and resuspended in 20 mM Tris-HCl, pH 7.5, 300 mM NaCl, 10% glycerol, 0.1% NP40, 4 mM MgCl₂, 5 mM DTT supplemented with EDTA-free Complete Protease Inhibitor (Roche, Basel, Switzerland). Cell suspensions were disrupted by sonication and incubated with 10-15 mg/L of DNase I (Roche, Basel, Switzerland) for 30 min. Cell debris was removed through ultracentrifugation at 30,000x g for 1 hour and clarified cell lysate was flash frozen and stored at -80°C prior to purification.

Clarified cell lysate was thawed and subsequently loaded onto a 5 mL pre-packed HisTrap FF column equilibrated with 25 mM HEPES, pH 7.5, 500 mM NaCl, 10% glycerol and 5 mM DTT. Proteins were eluted with 500 mM imidazole at a flow rate of

1 mL/min. Fractions containing the protein of interest were applied onto a pre-packed 26/10 HiPrep Desalting Column equilibrated with 25 mM HEPES, pH 7.5, 150 mM NaCl, 10% glycerol and 5 mM DTT. The buffer-exchanged NS5B peak was then applied onto a 20 mL Poly-U Sepharose column. Protein was eluted with an increasing salt gradient and fractions collected. Protein purity was assessed on Nu-PAGE pre-cast gels (Invitrogen, Carlsbad, CA). Purified NS5B samples were concentrated using Centri-Prep concentrators (Millipore, Billerica, MA, USA) and protein concentrations were determined by Bradford assay (Pierce, Rockford, IL, USA).

1.b) Protein Sequence

10 PDB: 1nb4, Apo form

The protein sequence is as described in WO 2007/026024. Calc. Mol. Properties: 64941.4 g/mol.

1.c) Inhibition assay with NS5b 1bJ4

Measurement of HCV NS5B polymerization activity was performed by evaluating the amount of radiolabeled GTP incorporated by the enzyme in a newly synthesized RNA using heteropolymeric RNA template/primer. The RdRp assay was carried out in 384-well plates using 50 nM of purified NS5B enzyme, which was incubated with 300 nM 5'-biotinylated oligo(rG₁₃)/poly(rC) or oligo (rU₁₅)/poly(rA) primer-template, 600 nM of GTP, and 0.1 µCi of [³H]GTP or [³H]UTP in 25 mM Tris-HCl, pH 7.5, 5 mM MgCl₂, 25 mM KCl, 17 mM NaCl and 3 mM of DTT. The 30 µL reaction mixture was incubated at room temperature for 2 h before stopping the reaction by adding 30 µL of streptavidin coated SPA-beads (GE Healthcare, Uppsala, Sweden) in 0.5 M EDTA. The 30 µL reaction was terminated after 2 hours at 25°C upon addition of 30 µL streptavidin-coated SPA beads (GE Healthcare, Uppsala, Sweden 5 mg/ml in 0.5 M EDTA). After incubation at 25°C for 30 min, the plate was counted using a Packard TopCount microplate reader (30 sec/well, 1 min count delay) and IC₅₀ values were calculated (Table 1: IC₅₀ 1bJ4). IC₅₀ values represent the concentration of compound required to decrease by 50% the amount of RNA produced which is measured by the detection of incorporated radiolabeled GTP.

30 2. HCV NS5B con1b

2.a) Cloning, expression and purification of NS5B con1b.

The coding sequence for NS5B (genotype 1b consensus strain Con1) lacking 21 C-terminal residues was amplified from plasmid pFKI₃₈₉/ns3-3' _N (Genbank accession no. AJ242654) and subcloned into the pET21b plasmid as described previously (Pauwels et al, 2007, J Virol 81:6909-19). The NS5BAC21 expression construct was


transformed into *E. coli* Rosetta 2 (DE3) (Novagen, Madison, WI). One hundred milliliters of LB-medium supplemented with carbenicillin (50 µg/mL) and chloramphenicol (34 µg/mL) was inoculated with one colony, grown overnight, and transferred to fresh LB-medium supplemented with 3% ethanol, carbenicillin and chloramphenicol, at a ratio of 1:200. The remaining of the procedure was as described previously (Pauwels et al, 2007, J Virol **81**:6909-19), except that the column used for ion-exchange chromatography was a 6 mL Resource S column (GE Healthcare), and that protein concentrations were determined with the Nanodrop (Nanodrop Technologies, Wilmington, DE, USA).

10 2.b) RNA-dependent RNA polymerase assay.






Fifty-percent inhibitory concentrations (Table 1: IC₅₀ con1b) were determined according to the method as described previously (Pauwels et al, 2007, J Virol **81**:6909-19) using a primer-dependent transcription assay. Following a 10 minute preincubation with the inhibitor, 20 nM of purified Con1b NS5B enzyme was incubated for 10 min. with 150 nM 5'-biotinylated oligo (rG₁₃) primer, 15 nM poly (rC) template, 19 mM Tris-HCl, 5 mM MgCl₂, 17 mM NaCl, 21 mM KCl, and 2.5 mM DTT. 600 nM GTP and 0.13 µCi of [³H]GTP was then added to initiate the 40-µl reaction mixture, which was then incubated at room temperature for 2h before the reaction was stopped by addition of 40-µl streptavidin-coated SPA beads.






20 The following Table 1 lists compounds according to any one of the above examples. The activities of the compounds tested are also depicted in Table 1.






Table 1






Nr.	Structure	EC ₅₀ (µM)	IC ₅₀ 1bJ4 (µM)	IC ₅₀ Con1b (µM)
2		= 0.09	= 0.41	


Nr.	Structure	EC_{50} (μM)	IC_{50} 1bJ4 (μM)	IC_{50} Con1b (μM)
1		= 0.07	= 0.24	= 0.038
11		= 0.055	= 0.22	
5		= 0.92	= 0.29	
12		= 0.29	= 0.93	
10		= 0.06	= 0.28	

Nr.	Structure		EC ₅₀ (μ M)		IC ₅₀ 1bJ4 (μ M)		IC ₅₀ Con1b (μ M)
13		=	0.39	=	0.19		
3		=	0.079			=	0.53
4		=	3.83			=	2.42
6		=	0.550				
8		=	0.040				

Nr.	Structure	EC_{50} (μM)	IC_{50} 1bJ4 (μM)	IC_{50} Con1b (μM)
9		= 0.800		
15		= 0.180		
16		= 0.078		= 0.040
17		= 0.170		
18		= 0.079		= 0.026

Nr.	Structure	EC ₅₀ (μ M)	IC ₅₀ 1bJ4 (μ M)	IC ₅₀ Con1b (μ M)
19		= 0.072		= 0.031
20		= 0.081		= 0.038
21		= 0.280		
22		= 0.160		
23		= 0.470		

Nr.	Structure		EC ₅₀ (μ M)	IC ₅₀ 1bJ4 (μ M)	IC ₅₀ Con1b (μ M)
24		=	14.48		
25		=	0.550		
26		=	0.130		
27		=	10.20		
28		=	0.330		

Nr.	Structure		EC ₅₀ (μ M)	IC ₅₀ 1bJ4 (μ M)	IC ₅₀ Con1b (μ M)
38		=	0.912		

Enzyme binding affinity

The compounds of formula (I) were examined for their enzymatic binding kinetics using a Surface Plasmon Resonance (SPR)-based method, i.e. Biacore. A slow dissociation of the inhibiting compound from its viral target (low k_{off} , low K_d) is believed to potentially reduce the development of drug resistance against anti-viral drugs (Dierynck et al. 2007. Journal of Virology, vol.81, No. 24, 13845-13851). All measurements were performed on a Biacore T100 instrument (GE Healthcare). The purified HIS₆-tagged NS5BAC21 polymerases were immobilized using non-covalent capturing to an NTA sensor chip (GE Healthcare) in immobilization buffer (20 mM MOPS pH 7.4, 500 mM NaCl, 0.005 % Tween-P20, 1 mM DTT, 50 μ M EDTA). Interaction studies were all performed at 25°C. Inhibitors were serially diluted in running buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 50 μ M EDTA, 1 mM DTT, 0.005 % Tween-P20) containing 5 % dimethyl sulfoxide (DMSO). Single-cycle kinetics were used, in which 5 increasing concentrations of compound were injected for a period of 300 s each in 1 single cycle, and dissociation was monitored for a period of 1200 s. The sensor surface was completely regenerated in between the cycles. Data were analyzed using simultaneous nonlinear regression analysis (global fitting) adapted for single-cycle kinetics with Biacore T100 BiaEval evaluation software 2.0 (GE Healthcare). The individual rate constants k_{on} and k_{off} and a derived affinity constant, $K_d = k_{\text{off}}/k_{\text{on}}$, were determined by a kinetic evaluation of the sensorgrams. The kinetic models accounted for bulk and limited mass transport effects. Every analysis was performed at least in two independent experiments. The dissociation rate of a kinetic interaction can be translated into a compound residence time (dissociative half-life $t_{1/2} = \ln(2)/k_{\text{off}}$) representative for the interaction time between the polymerase and its inhibitor.

The observed association rate constants (k_{on}), dissociation rate constants (k_{off}), derived affinity constant (K_d) and dissociative half-life ($t_{1/2}$) measured for compounds of formula (I) or subgroups thereof on NS5B wild-type enzyme (genotype 1b, Con1b) are given in Table 2.

- 5 Table 3 lists binding affinity data for compound nr. 1 on different forms of HCV NS5B polymerase. The different forms studied (NS5B Target) comprise different clinical isolates of different genotypes of the wild type enzyme, and, different mutant NS5B polymerases. The mutant enzymes were obtained by site directed mutagenesis of the 1bJ4 or Con1b NS5B enzyme. Mutations P495L, V494A and L392I are located in the
10 binding pocket of the compounds of the invention to NS5B polymerase.

- It was observed that the strong binding of compounds of formula (I) or subgroups thereof is consistent within one genotype, that the compounds of formula (I) or subgroups thereof show affinity for NS5B polymerase of the different genotypes, as well as for NS5B polymerases with mutation in the indole binding pocket, and that
15 binding of the compounds of formula (I) or subgroups thereof is not affected by mutations to other sites in the enzyme.

Table 2

Nr.	k_{on} (1/Ms)	k_{off} (1/s)	K_d (M)	$t_{1/2}$ (min)
16	2,2E+04	3,6E-05	1,6E-09	321,5
18	2,0E+04	4,8E-05	2,4E-09	241,0
1	2,0E+04	9,0E-05	4,4E-09	128,4
28	7,3E+03	6,6E-05	9,0E-09	175,5
25	2,9E+04	3,1E-04	1,1E-08	37,8
17	8,7E+03	1,6E-04	1,8E-08	72,0
27	9,5E+03	3,8E-03	4,0E-07	3,1
24	4,8E+03	3,7E-03	7,6E-07	3,1
38	5,3E+03	4,1E-05	7,7E-09	283,8


Table 3

NS5B target	k_{on} (1/Ms)	k_{off} (1/s)	K_d (M)	$t_{1/2}$ (min)
1a isolate 1	7,4E+04	4,8E-04	6,5E-09	24,2
1a isolate 2	3,9E+04	3,7E-04	9,4E-09	31,3
1a isolate 3	8,1E+04	6,0E-04	7,4E-09	19,2
1a isolate 4	7,4E+04	7,7E-04	1,1E-08	14,9
1a isolate 5	1,1E+05	3,1E-04	2,8E-09	37,2
1b isolate 1	2,6E+04	1,0E-04	4,0E-09	110,2

1b isolate 2	2,9E+04	6,7E-05	2,3E-09	172,1
1b isolate 3	3,7E+04	1,2E-04	3,3E-09	96,4
1b isolate 4	3,5E+04	1,7E-04	4,9E-09	67,5
2b isolate 1	1,8E+04	1,4E-02	8,2E-07	0,8
2b isolate 2	4,3E+04	1,2E-02	2,7E-07	1,0
2b isolate 3	4,4E+03	1,7E-02	3,8E-06	0,7
3a isolate 1	9,5E+04	3,7E-04	3,9E-09	31,1
3a isolate 2	2,5E+04	4,7E-04	1,9E-08	24,6
3a isolate 3	6,0E+04	3,6E-04	6,1E-09	31,7
4a isolate 1	2,0E+05	4,3E-04	2,1E-09	26,7
4a isolate 2	2,8E+05	3,8E-04	1,4E-09	30,1
4a isolate 3	1,8E+05	6,0E-04	3,4E-09	19,1
5a isolate 4	4,3E+04	9,7E-04	2,2E-08	12,0
6a isolate 5	5,0E+04	1,6E-03	3,2E-08	7,3
1bJ4	2,0E+04	1,0E-04	5,2E-09	110,2
Con1b	2,0E+04	9,0E-05	4,4E-09	128,4
P495L (1bJ4)	5,9E+03	2,2E-02	3,8E-06	0,5
L392I (Con1b)	1,8E+04	9,7E-04	5,5E-08	11,9
P495L (Con1b)	3,4E+03	2,2E-02	6,4E-06	0,5
V494A (Con1b)	5,5E+04	8,4E-04	1,5E-08	13,8
M414T (1bJ4)	2,6E+04	1,5E-04	5,9E-09	75,4
M423T (1bJ4)	2,5E+04	1,5E-04	6,3E-09	75,0
S282T (1bJ4)	3,4E+04	1,3E-04	3,9E-09	88,9
C316Y (Con1b)	3,5E+04	7,6E-05	2,2E-09	151,7



Claims

1. A compound of Formula (I)









5 including stereochemically isomeric forms, and N-oxides, salts, hydrates, and solvates thereof, wherein:

— R^1 is a bivalent chain selected from



10







- each R^3 is independently selected from the group comprising hydrogen, C_{1-4} alkyl and C_{3-5} cycloalkyl;
- a is 3, 4, 5 or 6;
- 5 - each b is independently 1 or 2;
- c is 1 or 2;
- macrocycle **A** has 14 to 18 member atoms, in particular macrocycle **A** has 17 or 18 member atoms;
- each R^2 is independently hydrogen, halo or C_{1-4} alkoxy;
- 10 - R^4 and R^5 are hydrogen or R^4 and R^5 together form a double bond or a methylene group to form a fused cyclopropyl;
- R^6 is hydrogen or methyl; and
- R^7 is a C_{3-7} cycloalkyl optionally substituted with halo.

2. A compound according to claim 1 wherein,





- 15 - R^1 is selected from $-N(R^3)-(CH_2)_4-N(R^3)-$,





- each R^3 is independently selected from hydrogen and methyl.
3. A compound according to any of claims 1 or 2 wherein R^2 is positioned the benzene group in para with respect to the bond linking this benzene to the indole group.
- 20 4. A compound according to any of claims 1 to 3 wherein R^2 is selected from fluoro and methoxy.
5. A compound according to any of claims 1 to 4 wherein R^7 is selected from cyclohexyl and 2-fluorocyclohexyl.
- 25 6. A compound according to any of claims 1 to 5 wherein R^4 and R^5 together form a double bond.
7. A compound according to any of claims 1 to 5 wherein the compounds of formula (I) have the stereochemical configuration as illustrated by formula (IA).




8. A compound according to any of claims 1 to 3, having one of the structural formula II-1, II-2, II-3, III-1, III-2, III-3, III-4, IV-1, IV-2 or IV-3




5







(III-4),



(IV-1),




(IV-2),



(IV-3).

9. A compound having the following structure:



- 5 including stereochemically isomeric forms, and N-oxides, salts, hydrates, and solvates thereof.
10. A pharmaceutical composition comprising a carrier, and as active ingredient an anti-virally effective amount of a compound as claimed in any of claims 1 to 9.
11. A pharmaceutical composition according to claim 10, further comprising at least one other anti-HCV compound.
12. A pharmaceutical composition according to claim 10 or 11, further comprising at least one anti HIV compound.
13. A compound according to any of claims 1 to 9 or a pharmaceutical composition according to any of claims 10 to 12, for use as a medicament.
14. A compound according to any of claims 1 to 9 or a pharmaceutical composition according to any of claims 10 to 12, for inhibiting HCV replication.
15. Use of a compound according to any of claims 1 to 9, for the manufacture of a medicament for inhibiting HCV replication.