

which is further substituted by one, two or three substituents **Z**,

whereby each **Z** is a substituent independently selected from the group consisting of

H, -OH, oxo, -CN, halogen, -C₁₋₆-alkyl, -O-C₁₋₆-alkyl, -C₁₋₆-haloalkyl, three- to seven-membered cycloalkyl, **Het**, **Hetaryl**, -CO-N(CH₃)₂, -CO-NHCH₃, -CO-NH₂, -(C₁₋₃-alkylene)-O-(C₁₋₃-alkyl), -O-**Het**,

which is further substituted by one, two or three substituents **X**,

whereby each **X** is independently selected from the group consisting of halogen, oxo, -C₁₋₄-alkyl, -O-C₁₋₄-alkyl, -C₁₋₄-haloalkyl, -O-(C₁₋₄-alkylene)-**Het**, **Het**, -NH₂,

whereby substituent **X** is further substituted by one, two or three substituents of a group selected from H, oxo, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and tert-butyl,

and the pharmaceutically acceptable salts of the aforementioned compounds.

18. The compound of formula **1c** or of formula **1c'** of claim 17, wherein

R¹ is

a) either selected from the group consisting of **Het** and **Hetaryl**;

which is further substituted by one, two or three substituents **Z**,

whereby each **Z** is a substituent independently selected from the group consisting of -OH, oxo, -CN, halogen, -C₁₋₆-alkyl, -O-C₁₋₆-alkyl, -C₁₋₆-haloalkyl, three- to seven-membered cycloalkyl, **Het**, **Hetaryl**, -CO-N(CH₃)₂, -CO-NHCH₃, -CO-NH₂, -(C₁₋₃-alkylene)-O-(C₁₋₃-alkyl), -O-**Het**,

which is further substituted by one, two or three substituents **X**,

whereby each **X** is independently selected from the group consisting of H, halogen, oxo, -C₁₋₄-alkyl,

-O-C₁₋₄-alkyl, -C₁₋₄-haloalkyl, -O-(C₁₋₄-alkylene)-**Het**, **Het**, **Hetaryl**, -NH₂,

whereby substituent **X** is further substituted by one, two or three substituents of a group selected from H, oxo, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and tert-butyl, and the pharmaceutically acceptable salts of the aforementioned compounds.

21. The compound of formula **1c** or of formula **1c'** according to claim 17, wherein

R¹ is phenyl,

which is further substituted by one, two or three substituents **Z**,

whereby each **Z** is a substituent independently selected from the group consisting of

H, -OH, oxo, -CN, halogen, -C₁₋₆-alkyl, -O-C₁₋₆-alkyl, -C₁₋₆-haloalkyl, three- to seven-membered cycloalkyl, **Het**, **Hetaryl**, -CO-N(CH₃)₂, -CO-NHCH₃, -CO-NH₂, -(C₁₋₃-alkylene)-O-(C₁₋₃-alkyl), -O-**Het**,

which is further substituted by one, two or three substituents **X**,

whereby each **X** is independently selected from the group consisting of halogen, oxo, -C₁₋₄-alkyl, -O-C₁₋₄-alkyl, -C₁₋₄-haloalkyl, -O-(C₁₋₄-alkylene)-**Het**, **Het**, -NH₂,

whereby substituent **X** is further substituted by one, two or three substituents of a group selected from H, oxo, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and tert-butyl,

and the pharmaceutically acceptable salts of the aforementioned compounds.

22. The compound of formula **1c** or of formula **1c'** according to claim 17, which is selected from the group consisting of