Production of Nucleic Acid

Field of invention

The present invention relates to a process for the production of nucleic acid encoding a target protein and target protein obtainable thereby, including enzymes such as nucleic acid processing enzymes, and in particular reverse transcriptase.

Background of the invention

Protein evolution is a known technology for selection and directed evolution of proteins from large libraries. The basic principle of selection is to ensure that there is a linkage between a specific phenotype (protein) and its encoding genotype. This phenotype-genotype linkage can be realized in three different ways:

- covalent linkage such as mRNA display, and to some extent phage display, bacterial display, yeast display etc.,
- non-covalent linkage which use affinity interaction. Examples are ribosome display, CIS display, plasmid display etc.,
- compartmentalization such as *in vitro* compartmentalisation (IVC), compartmentalized self-replication (CSR), simple bacterial screening, high throughput screening etc.

As indicated in the above paragraph, one example of covalent phenotype-genotype linkage is achieved using mRNA display. As described by Roberts and Szostak (1997) covalent fusions between an mRNA and the peptide or protein that it encodes can be generated by in vitro translation of synthetic mRNAs that carry puromycin, a peptidyl acceptor antibiotic, at their 3' end.

Non-covalent linkage between phenotype and genotype can be achieved with ribosome display. In ribosome display, an array of RNAs including one or more encoding a target protein is subjected to an in vitro translation system so as to form a non-covalent ternary complex of ribosome, mRNA and protein coupled to tRNA. This array of ternary complexes must be stabilized. Accordingly, each ternary complex formed during in vitro translation uses mRNA lacking a STOP codon at the end. The ternary complexes are further stabilized at low temperature (4°C) and high concentration of magnesium (50mM). In the stable complex the linkage between phenotype and genotype is preserved. A selection step follows whereby target

protein is selected on the basis of a property of the protein whilst still attached to the ternary complex. Selected ternary complexes may then be disassembled and the mRNA associated with the target protein is amplified by RT-PCR.

In general, ribosome display is successfully applied for the selection of peptides (Mattheakis et al., 1994; Matsuura and Pluckthun, 2003) and proteins (Hanes and Pluckthun, 1997; He and Taussig, 1997; Irving et al., 2001), which bind to different targets. In some cases it is possible to use ribosome display to select for enzymatic activities, performing affinity selection of proteins with suicide inhibitor (Amstutz et al., 2002) or active site ligand (Takahashi et al., 2002).

In in vitro compartmentalization (IVC) phenotype and genotype linkage is realized by in vitro compartmentalization of genes in a water-in-oil emulsion. The number of genes used to prepare the emulsion is calculated such that most of water compartments contain no more than a single gene. The compartmentalized genes are transcribed and translated. The activity of the synthesized proteins is then assessed. Subsequent selection on the basis of protein activity results in amplification of DNA encoding active proteins with desired properties. Water droplets used in most IVC applications are 2-3 μm in size, giving ~5 femtolitres reaction volume and ~10^{10} water-in-oil compartments (50 μl water phase) per 1 ml of emulsion. The first successful example of IVC selection system was based on target-specific DNA methylation activity (Tawfik and Griffiths, 1998). Genes of HaeIII methyltransferase were compartmentalized, transcribed and translated. In vitro synthesized methyltransferase in the presence of a cofactor was able to methylate its own DNA. Based on methylated DNA (reaction product) resistance to digestion with HaeIII restriction endonuclease, genes encoding methyltransferase were selected from a 10^7 fold excess of other DNA molecules.

To date many more modifications of IVC have been designed and realized. The easiest way to perform IVC selection is to use DNA-modifying enzymes, in particular DNA-methyltransferases (Lee et al., 2002; Cohen et al., 2004). A similar experimental strategy was applied in order to select for active variants of FokI restriction endonuclease (Doi et al., 2004). Compartmentalized DNA, encoding active restriction endonucleases, was

digested and selected by subsequent incorporation of biotin-dUTP and binding to streptavidin beads.

A different IVC selection strategy was applied for evolution of DNA polymerases (Ghadessy et al., 2001; Ghadessy et al., 2004; Ong et al., 2006). This new selection method is based on 'compartmentalized self-replication' (CSR) of genes encoding active DNA polymerase. Contrary to usual IVC, where proteins of interest are expressed in situ, CSR is performed by compartmentalization of bacterial cells expressing thermophilic DNA polymerase. Cells resuspended in PCR buffer, supplemented with primers and dNTP's, are emulsified yielding compartments ~15 μm in size. Each water droplet serves as a separate PCR compartment. During the initial PCR denaturation step the bacterial cells are broken releasing the expressed thermophilic DNA polymerase and its encoding gene into the reaction mixture, allowing self-replication to proceed meanwhile other bacterial proteins are denatured by high temperature.

A modification of IVC is the use of double water-in-oil-in-water emulsions. Water droplets surrounded by an oil layer can be analyzed and sorted in a FACS at a rate of $>10^4$ variants per second (Bernath et al., 2004; Mastrobattista et al., 2005).

Selection of proteins for binding may also be performed by IVC. A protein of interest expressed in water-in-oil compartments is coupled covalently (Bertschinger and Neri, 2004) or non-covalently (Doi and Yanagawa, 1999; Yonezawa et al., 2003; Sepp and Choo, 2005) to the gene that encodes it.

Other applications of IVC are known where microbeads entrapped into compartments are used as intermediators for protein and gene coupling. Single genes attached to microbeads are transcribed and translated in water droplets. Newly synthesized protein is captured onto the same bead within reaction compartment. After emulsion is broken isolated beads can be used further for affinity selection (Sepp et al., 2002). Emulsions usually are broken using organic solvents (hexane, ether, chloroform), which can also decrease activity of certain enzymes displayed on the beads, limiting the application of this technology. For selection of catalytic activity microbeads can be easily washed, resuspended in different reaction buffer and

compartmentalized again by second emulsification step (Griffiths and Tawfik, 2003). However sometimes rigid enzyme-bead-gene complexes (because of sterical hindrance and enzyme mobility limitations) can not fulfil essential reaction requirements and the activity of the attached enzymes may be lower than that of the free enzyme. In addition, such methods are technically complicated since many additional components have to be used (i.e. affinity tags, antibodies, and beads).

The composition of emulsions used in IVC is designed to ensure stability of water compartments and efficient in vitro transcription of mRNA and subsequent translation of target proteins. In vitro evolution has a broad range of targets to be improved. Some proteins and enzymes of interest are robust hard-workers and can be active enough in different buffers, in particular in reaction mixtures used in IVC. However there are many complicated enzymes, which can work only under optimized or specific conditions. In addition to that the first law of in vitro evolution says - "you will evolve what you are selecting for". That means, an enzyme evolved and optimized in transcription/translation reaction mixture will work well in that particular mixture and most likely will perform much worse in its own buffer. In some cases enzyme working conditions are incompatible with in vitro transcription and translation mixture used for protein expression in compartments. A partial solution is a nano-droplets delivery system (Bernath et al., 2005) used to transport different solutes into emulsion compartments. Even more sophisticated manipulations with water-in-oil compartments can be done employing microfluidics devices. Highly monodisperse single or double emulsions (Thorsen et al., 2001; Okushima et al., 2004) can be prepared at a rate of up to 10000 aqueous droplets per second. Generated water compartments can be transported in microfluidic channels, fused, subdivided and sorted (Song et al., 2003; Link et al., 2006). Nevertheless full buffer exchange in the compartments is still a problem.

Reverse transcriptases are very important commercial enzymes used to synthesize cDNA from an mRNA target. A lot of research has been done in order to improve properties of reverse transcriptases. However no properly working selection system suitable for in vitro evolution of reverse transcriptase was known to date. Almost all improvements are made and mutants of reverse transcriptase selected using high throughput screening and rational design.

Neither ribosome display (RD), nor in vitro compartmentalization (IVC) may be used for selection of fully active reverse transcriptase. Ternary complexes used in ribosome display usually are not stable at higher temperatures required to perform reverse transcriptase selection and as a consequence linkage between phenotype and genotype will be lost. Whilst relatively stable ternary complexes used in ribosome display can be produced using synthetic in vitro translation extract WakoPURE (Matsuura et al., 2007), in vitro translated reverse transcriptase immobilized to ribosome and mRNA will encounter significant steric hindrance during synthesis of full-length cDNA. There is also the possibility that immobilized enzyme will act in trans as well as in cis, which again is incompatible with protein evolution strategy, because the phenotype-genotype linkage will not be preserved.

IVC generally employs DNA as genetic material. In vitro transcription of mRNA and target protein translation is performed in spatially separated emulsified water compartments in the presence of the DNA. In the case of reverse transcriptase in vitro evolution, the presence of coding DNA sequence abolishes the main prerequisite of selection for reverse transcriptase activity - cDNA has to be synthesized de novo. In other words, selection for better reverse transcriptase variants is based on enzyme ability to synthesize their own coding cDNA from mRNA. Newly synthesized cDNA has to be amplified by PCR, therefore cDNA should be the only source of DNA in the reaction. DNA used in IVC selection will amplify together with cDNA canceling basic selection scheme.

More sophisticated variants of in vitro compartmentalization, such as usage of microbeads entrapped in water compartments (Sepp et al., 2002; Griffiths and Tawfik, 2003), allow complete exchange of reaction buffer. In this approach the phenotype-genotype linkage is realized via microbeads yielding a rigid selection unit mRNA-microbead-protein, which, in case of reverse transcriptase selection, again can cause steric hindrance and as a result inefficient cDNA synthesis.

Taq DNA polymerase able to synthesize ~300 nucleotides long cDNA was selected using modification of phage display technology (Vichier-Guerre et.al, 2006). Although this approach works, it has some shortcomings: 1) not all proteins can be displayed on phages; 2) absolute requirement to use biotin labeled nucleotides for selection; 3) displayed enzyme can work in

trans as well as in cis; 4) because of steric hindrance and enzyme mobility limitations phage-enzyme-DNA/RNA complex can interfere with efficient synthesis of cDNA.

The possibility of selection for reverse transcriptase is also mentioned in WO0222869, which is related to the compartmentalized self replication (CSR) method. CSR technology is used to select thermophilic DNA polymerases, in particular Taq DNA polymerase (Ghadessy et al., 2001; Ghadessy et al., 2004; Ong et al., 2006). Bacterial cells, expressing mutants library of thermophilic DNA polymerase are suspended in PCR mixture and emulsified yielding separate PCR compartments for in vitro selection of more active polymerases.

Real selection for reverse transcriptase activity will be prevented by the presence of bacterial RNases, which remain active at moderate temperatures and will degrade target mRNA. There will also be DNA contamination from non-selected plasmid DNA (released from bacterial cell) as well as the presence of all E.coli enzymes, structural proteins, ribosomes, NTP, RNases, DNases and small molecular weight molecules.

Summary of the Invention

The present invention aims to provide an improved method of protein evolution which does not suffer from drawbacks of the prior art methods.

Accordingly, in a first aspect, the present invention provides a process for the production of nucleic acid encoding a target protein, which comprises:

(a) providing an array of RNA or DNA molecules including one or more encoding the target protein;

(b) generating a target protein from the array to form RNA-protein or DNA-protein complexes in which the RNA or DNA molecule is non-covalently or covalently bound to the complex;

(c) separating the complexes into compartments wherein most or all of the compartments contain no more than one complex;

(d) subjecting the complexes to reaction conditions which allow target protein activity; and

(e) selecting nucleic acid encoding the target protein on the basis of the activity associated therewith,

wherein when the complex is a DNA-protein complex in which the DNA is non-covalently bound, step b) is performed in the absence of separate compartments for each complex.

The present inventors have devised a method which utilizes two distinct types of phenotype-genotype linkage and achieves a new selection system for use in methods of protein evolution within the laboratory. The new features of the method, described further below, allow it to be applied to an extended range of target proteins in comparision to the methods of the prior art, with increased ease of use and flexibility. In particular, the present invention provides for the first time the possibility of evolving and improving the properties of reverse transcriptase enzymes – one of the most important enzyme groups in the toolbox of molecular biologists.

In one embodiment of this aspect the invention provides a process for the production of nucleic acid encoding a target protein, which comprises:

(a) providing an array of RNA or DNA molecules including one or more encoding the target protein;

(b) generating a target protein from the array to form RNA-protein or DNA-protein complexes;

(c) separating the complexes into compartments wherein most or all of the compartments contain no more than one complex;

(d) subjecting the complexes to reaction conditions which allow target protein activity; and

(e) selecting nucleic acid encoding the target protein on the basis of the activity associated therewith,

wherein in the RNA-protein complex the RNA is non-covalently or covalently bound thereto and in the DNA-protein complex the DNA is covalently bound thereto.

In a second embodiment of this aspect the invention provides process for the production of nucleic acid encoding a target protein, which comprises:

(a) providing an array of RNA or DNA molecules including one or more encoding the target protein;

(b) generating a target protein from the array to form RNA-protein or DNA-protein complexes in which the RNA or DNA molecule is non-covalently or covalently bound to the complex;

(c) separating the complexes into compartments wherein most or all of the compartments contain no more than one complex;

(d) subjecting the complexes to reaction conditions which allow target protein activity; and

(e) selecting nucleic acid encoding the target protein on the basis of the activity associated therewith,

wherein step b) is performed in the absence of separate compartments for each complex.

Covalent linkages between DNA or RNA and the target protein can be generated by any technique known in the art, for example, mRNA display, phage display, bacterial display or yeast display. In particular, a covalent RNA-protein linkage can be generated using the technique of mRNA display, while a covalent DNA-protein linkage can be generated using the technique of covalent antibody display (CAD) (Reiersen et al., 2005), performing translation in compartments by covalent DNA display (Bertschinger and Neri, 2004), or using similar covalent display techniques such as those described by Stein et al., (2005).

Non-covalent linkages between DNA or RNA and the target protein can also be generated by any technique known in the art, for example, ribosome display, CIS display, or plasmid display. In particular, a non-covalent DNA-protein linkage can be generated in the absence of a compartment using CIS display (Odergrip et al., 2004), while a non-covalent RNA-protein linkage can be generated using the technique of ribosome display.

As indicated above, when the complex is a DNA-protein complex in which the DNA is non-covalently bound, step (b) of the process of the invention is performed in the absence of separate compartments for each complex. In other words, step (b) is un-compartmentalised. Specifically, when the complex generated is a DNA-protein complex in which the DNA is non-covalently bound, the generation step is performed without separating each member of the array from one another. In particular, the generation step is performed without separating each member of the array by *in vitro* compartmentalization (IVC). In a particularly preferred

embodiment the generation step is performed without separating each member of the array in a water-in-oil emulsion.

Compartmentalisation can also be performed by any method known in the art which enables the complexes to be generated or separated such that all or substantially all of the compartments contain no more than one complex. In particular, it is preferred that at least 70%, at least 80% or at least 90% of the compartments contain no more than one complex. For example, compartmentalization can be performed by separating members of the array or each complex into different wells on a microtitre or nanotitre plate, or by *in vitro* compartmentalization (IVC). In particularly, separation by IVC can involve separation into aqueous droplets in a water-in-oil emulsion or a water-in-oil-in-water emulsion.

The method of the present invention combines at least two different types of genotype-phenotype linkage selected from the list: covalent linkage, non-covalent linkage and compartmentalization. In a preferred aspect of the invention the method utilized no more than two of these linkages. Thus in a particularly preferred embodiment the method utilizes the covalent or non-covalent linkage as the only genotype-phenotype linkage in step b). In other words, in this embodiment there is no compartmentalization in step b).

Covalent/non-covalent linkages between DNA or RNA and the protein in the absence of a compartment can be established in many different ways, for example, by ribosome display, mRNA display (Roberts and Szostak, 1997), CIS display (Odergrip et al., 2004) or covalent antibody display (CAD) (Reiersen et al., 2005). Specifically, covalent DNA-protein linkage can be realized in the absence of compartments by using CAD technique, while covalent RNA-protein linkage and non-covalent RNA-protein linkage can be established by mRNA display and ribosome display, respectively.

The present invention can be realized through a combination of many different linkages. For example, the present invention can be realized through a compbination of ribosome display and in vitro compartmentalization, or through a combination of mRNA display, CIS display, or CAD display and IVC.

In a preferred aspect the process for the production of nucleic acid encoding a target protein is realized through a combination of ribosome display and in vitro compartmentalization. In particular such an process comprises:

(a) providing an array of mRNAs including one or more encoding the target protein;

(b) incubating the array of mRNAs under conditions for ribosome translation to generate an array of ternary complexes each comprising an mRNA, a ribosome and protein translated from the mRNA;

(c) incorporating the array of ternary complexes into aqueous phase droplets of a water-in-oil or a water-in-oil-in-water emulsion, wherein most or all of the aqueous phase droplets contain no more than one ternary complex;

(d) subjecting the aqueous phase droplets to reaction conditions which allow protein activity; and

(e) selecting nucleic acid encoding the target protein on the basis of the enzyme activity associated therewith.

Such a process according to the present invention may be termed "compartmentalized ribosome display" (CRD). CRD is applicable to a wide range of target proteins, including enzymes. CRD has the advantage that the linkage between the enzyme and the mRNA is non-covalent. Thus if the reaction conditions used in step (d) involve a raised temperature the ternary complexes generated in step (b) will fall apart and the enzyme will be released. This avoids the problems associated with enzyme mobility described above for prior art methods in which the enzyme is immobilized on a bead.

Emulsion droplets with ribosome display complexes inside can be sorted or selected in step (e) in many ways. Preferably they are sorted by fluorescence activated cell sorting (FACS) or using microfluidic techniques. Both techniques mainly exploit fluorescence based droplet sorting. However, droplets can also be searpated by size, light diffraction or light absorption, depending on the reaction conditions used in step (d) and the protein activity that is being selected for.

Fluorescent based sorting methods are preferably used when the target protein is an enzyme. In this embodiment the reaction conditions employed in step (d) include a non-fluorescent substrate capable of being converted to a fluorescent product. Activity by the enzyme generates the fluorescent product, allowing FACs to be used to distinguish between fluorescent droplets containing an active enzyme and non- or less fluorescent droplets which contain no active enzyme or a less active enzyme.

In particular, CRD is applicable to nucleic acid processing enzymes such as reverse transcriptases, and allows for fast and efficient in vitro evolution.

In a further aspect the present invention provides a process for the production of nucleic acid encoding a target protein, which comprises:

(a) providing an array of mRNAs including one or more encoding the target protein, wherein the mRNAs comprise a substrate for an enzyme comprising the target protein or a co-enzyme thereof;

(b) incubating the array of mRNAs under conditions for ribosome translation to generate an array of ternary complexes each comprising an mRNA, a ribosome and protein translated from the mRNA;

(c) incorporating the array of ternary complexes, and optionally the co-enzyme, into aqueous phase droplets of a water-in-oil or a water-in-oil-in-water emulsion, wherein most or all of the aqueous phase droplets contain no more than one ternary complex;

(d) subjecting the aqueous phase droplets to reaction conditions which allow enzyme activity; and

(e) selecting nucleic acid encoding the target protein on the basis of the enzyme activity associated therewith.

In one embodiment of this aspect of the invention, where the nucleic acid processing enzyme is a DNA dependent DNA polymerase, the mRNA of step (a) can be ligated to a double stranded DNA adaptor molecule to provide the substrate.

CRD diversity is ~10^9-10^{10} variants and is limited by IVC step. The new method is much more efficient, less time consuming and cheaper compared to high throughput screening (HTS),

which can be used to screen ~10^5-10^6 mutant variants of reverse transcriptase. CRD diversity is about four orders of magnitude higher compared to HTS; thus many more beneficial mutants missed by HTS can be easily fished-out by compartmentalized ribosome display selection.

According to step (a), an array of mRNAs which are typically synthesized mRNAs is provided which includes one or more members of the array encoding the target protein. Where the target protein is reverse transcriptase the mRNAs comprise a substrate for an enzyme comprising the target protein or a co-enzyme thereof. In the subsequent selection step (e), the nucleic acid encoding the target protein is selected on the basis of the enzyme activity associated therewith. In this way, two embodiments of the invention are contemplated: one in which the enzyme activity is provided by the target protein and one in which the enzyme activity is provided by a co-enzyme of the target protein in the presence of the target protein. In the embodiment requiring the co-enzyme, this is incorporated into aqueous phase droplets of the water-in-oil emulsion of step (c), as is the array of ternary complexes. Where the enzyme comprises the target protein, no additional co-enzyme need to be incorporated into the aqueous phase droplets.

In step (b) of the process, the array of mRNAs is treated with ribosomes to generate an array of ternary complexes, each comprising an mRNA, a ribosome and protein translated from the mRNA. This step may be performed under any conditions suitable for typical in vitro translation of mRNA, as used for example in the technique of ribosome display. At this point, the ternary complexes may be purified, although this is not essential. The ternary complexes may be supplemented at this point with any co-substrates necessary for the subsequent enzyme activity whereupon the reaction mixture is typically emulsified to give approximately 10^{10} water-in-oil compartments each typically having a mean diameter of approximately 2 to 3 μm. Even a small volume (25 μl) of in vitro translation reaction generates approximately 10^{11} to 10^{12} molecules of stored ribosomal complexes. A typical ribosome display method uses mRNAs lacking STOP codons, although STOP codon may be present (Matsuura et al., 2007). In order to achieve aqueous phase droplets in which most or all contain no more than one ternary complex the concentration of ternary complexes would have to be reduced by about two orders of magnitude as compared with corresponding concentration used in a typical

ribosome display technique. Only a very small concentration of ternary complexes is used in this step of the process.

The enzyme may comprise a nucleic acid processing enzyme, which may be an RNA processing enzyme. The nucleic acid processing enzyme may comprise the target protein and may be selected from a nucleic acid polymerase, a nucleic acid ligase and a terminal deoxynucleotidyl transferase. As decribed in further detail herein, the nucleic acid polymerase may comprise a reverse transcriptase. In this embodiment, the mRNA encoding the reverse transcriptase is itself the substrate for the reverse transcriptase. Step (e) of selecting nucleic acid encoding the target protein, comprises selecting cDNA produced by the action of the reverse transcriptase, which cDNA encodes reverse transcriptase.

Where the target protein is a nucleic acid ligase, selection for RNA (DNA) ligases able to ligate RNA to RNA or DNA to RNA can be performed. Preferably, the reaction conditions which allow enzyme activity include a co-substrate comprising a nucleic acid linker or adaptor, which co-substrate further comprises an affinity ligand for attachment to a ligand binding partner or sequence tag for specific amplification of processed mRNA in RT-PCR. In the first case, the mRNA encoding the target protein is ligated to the co-substrate in those aqueous phase droplets which incorporate mRNA encoding a nucleic acid ligase. Preferably, the affinity ligand comprises biotin and the ligand binding partner comprises streptavidin. The step of selecting nucleic acid encoding the target protein comprises selecting mRNA incorporating the co-substrate by attachment to a solid phase comprising the ligand binding partner. In a typical process, a mutant library of ligase is translated in vitro and purified ternary complexes are diluted and emulsified in reaction buffer with biotin labeled DNA/RNA linker and/or adaptor. After bringing the emulsion to a temperature of 37°C ribosome ternary complexes disassemble. Ligase will be released and the 3' end of the mRNA will become accessible for the biotin labeled adaptor and subsequent ligation reaction. Biotin labeled mRNA encoding only active (or more active) variants of ligase will be purified on the streptavidin beads and may be amplified by RT-PCR.

In the second case the step of selecting nucleic acid encoding the target protein comprises selecting mRNA with attached sequence specific tag, which can be used for selective annealing site of primer for reverse transcription and subsequent PCR.

In a typical process, a mutant library of ligase is translated in vitro and purified ternary complexes are diluted and emulsified in reaction buffer with DNA/RNA linker and/or adaptor. After bringing the emulsion to a temperature of 37°C ribosome ternary complexes disassemble. Ligase will be released and the 3' end of the mRNA will become accessible for the adaptor and subsequent ligation reaction. RNA encoding only active (or more active) variants of ligase will have specific linker sequence required for specific annealing of primer used in reverse transcription and may be efficiently amplified by RT-PCR.

It is also possible to select for terminal deoxynucleotidyl transferase (TdT). This enzyme works on RNA and incorporates deoxyribonucleotides, ribonucleotides, nucleotide analogues and similar. In this embodiment, the reaction conditions which allow enzyme activity include a co-substrate comprising dNTP which further comprises an affinity ligand for attachment to a ligand binding partner. As with the nucleic acid ligase, the affinity ligand may be biotin and the ligand binding partner streptavidin. Selecting nucleic acid encoding the target protein may comprise selecting mRNA incorporating the co-substrate by attachment to a solid phase comprising the ligand binding partner. A mutant library of TdTs may be translated in vitro and purified ribosome ternary complexes may have to be diluted and emulsified in reaction buffer with biotin labeled nucleotides, such as biotin-dUTP. The optimal working temperature for wild type enzyme is 37°C. At this temperature the ribosome ternary complexes will disassemble and the 3' end of mRNA will become accessible for template independent polymerization reaction. TdT-encoding mRNA incorporating biotin labeled nucleotide is selected on the streptavidin beads and may subsequently be reversed transcribed and amplified by RT-PCR.

In a further embodiment, the target protein comprises a reverse transcriptase helper enzyme such as a helicase, pyrophosphatase, processivity factor, RNA binding protein or other protein able to improve a reverse transcription reaction in the presence of reverse transcriptase. In this embodiment, the nucleic acid processing enzyme comprises a reverse transcriptase, which is

the co-enzyme incorporated into the aqueous phase droplets of the water-in-oil emulsion. In step (e) of selecting nucleic acid target protein, cDNA is selected, which cDNA is produced by the action of the reverse transcriptase and which encodes the reverse transcriptase helper enzyme. The presence of the reverse transcriptase helper enzyme in the aqueous phase facilitates reverse transcription of the mRNA which encodes the helper. Thus, mRNA which is reverse transcribed forms cDNA which encodes the helper and this may be PCR amplified.

In a further embodiment, the target protein comprises an RNase inhibitor. In this embodiment, the nucleic acid processing enzyme comprises an RNase. The step (e) of selecting nucleic acid encoding the target protein comprises selecting mRNA undegraded by RNase. In this embodiment, the RNase is incorporated as the co-enzyme into the aqueous phase droplets of the water-in-oil emulsion. Once reaction conditions allow enzyme activity, any droplets not containing effective RNase inhibitor would exhibit RNase activity whereby the mRNA would be degraded. Thus, mRNA encoding RNase inhibitor effective at the reaction conditions used would survive. Typically, a mutant library of RNase inhibitors is translated in vitro and purified ribosome ternary complexes are diluted and emulsified in reaction buffer with appropriate RNase. In an alternative arrangement RNase can be delivered later by emulsion micro droplets. mRNA encoding only active (or more stable) RNase inhibitor will be purified and amplified by RT-PCR.

Compartmentalized ribosome display can also be used for reaction buffer exchange in *in vitro* compartmentalization where selection buffer is incompatible with in vitro translation mixture and substrate conversion to product has to be performed under strictly controlled reaction conditions.

The nucleic acid encoding the target protein which has been selected on the basis of the enzyme activity associated therewith may be DNA or RNA, as discussed herein. The array may be converted or amplified to form DNA or RNA. In a preferred arrangement, the array is converted or amplified to form the array of mRNAs of step (a) of the process and is subject to one or more further cycles of steps (b) to (e) so as to enrich further the array with increasing amounts of mRNAs encoding the target protein.

The step (d) of subjecting the aqueous phase droplets to reaction conditions which allow enzyme activity provides the basis for selection step (e) where those nucleic acids encoding the target protein are selected. A wide range of reaction conditions may be used in step (d) to provide selection pressure. In one example, the reaction conditions include a temperature above the optimum temperature for a wild type enzyme. These reaction conditions may be used to select a mutant enzyme which is more thermostable than wild type enzyme or which has a greater reaction velocity at that temperature or an altered temperature-activity profile. Mutant enzymes may have to operate at higher sensitivities than wild type enzymes because concentrations of mRNA in the aqueous phase droplets are approximately 400pM. Mutant enzymes may also have to perform more accurately. All of these selection pressures are particularly important in relation to reverse transcriptases. As well as physical conditions, the reaction conditions may include alterations in buffer, concentrations of other factors such as metal ions and pH.

Many more different selection pressures can be applied in CRD selecting for better reverse transcriptases: 1) selection for more soluble enzymes which are less prone to aggregation - ternary complexes (before emulsification) have to be preincubated with hydrophobic material in order to eliminate proteins with surface exposed hydrophobic residues; 2) selection for very fast enzymes - reverse transcription reaction times have to be gradually reduced during selection cycles; 3) selection for enzymes synthesizing long cDNA - gradual prolongation of mRNA library used in CRD and as a consequence synthesis of longer cDNA; 4) selection for enzymes able to transcribe through secondary structures - secondary structure forming sequences have to be introduced into mRNA library used in CRD; 5) selection for enzymes working in buffers which are different from RT buffer (for example in PCR, one step RT-PCR buffer or with denaturing agents) - CRD selection has to be performed in buffer of our choice; 6) selection for enzymes able to incorporate nucleotide analogues - selection has to be performed in RT buffer with biotin labeled nucleotide analogues with subsequent cDNA purification on streptavidin beads.

Compartmentalized ribosome display (CRD) is also suitable for many fluorescence activated cell sorting (FACS) applications. Protein of interest has to be displayed in ribosome display format. Optionally purified (or just diluted many times) ternary complex comprising mRNA-

ribosome-protein (tRNA) mixed with non fluorescent substrate (S) in reaction buffer should be emulsified producing double water-in-oil-in-water emulsions (Bernath et al., 2004; Mastrobattista et al., 2005). Active variants of compartmentalized enzymes will convert substrate (S) to fluorescent product (P) allowing FACS to distinguish between fluorescent (active enzyme inside) and "dark" (inactive enzyme inside) droplets. Contrary to previously published examples, where enzymatic reaction has to be performed in transcription/translation mixture, CRD allows for complete buffer exchange and selection for active enzymes in more native (required) conditions (FIG 10.)

Selection and evolution of thermostable DNA polymerases using CRD is also possible (FIG 11.). Polymerase of interest has to be displayed in ribosome display format. Optionally purified (or just diluted many times) ternary complex comprising mRNA-ribosome-polymerase can be used to prepare reaction mixture with reverse transcriptase (helper enzyme), dNTP's and primer set in PCR buffer. Reaction solution should be emulsified producing water-in-oil emulsion. In first RT step - reverse transcriptase has to synthesize cDNA, which subsequently will serve as a target for second PCR step - cDNA amplification by ribosome displayed DNA polymerase. One of the primers used in PCR can have biotin for optional subsequent purification with streptavidin beads and non-complementary 5' end. After RT-PCR emulsions should be broken, newly synthesized DNA fragment can be purified via biotin and reamplified using new primer set, which will contain one primer with sequence non-complementary to cDNA, but identical to 5' part of primer used in first amplification reaction (selective amplification of DNA over cDNA background). More active variants of DNA polymerase will be enriched over less active variants and can be used for further analysis or next selection round (FIG 11.).

Important features of the compartmentalized ribosome display technology are:

1) the genotype is maintained by an mRNA library, which is especially useful in selecting RNA processing enzymes;
2) the selection unit is a ternary complex of mRNA-ribosome-protein (tRNA) and this can readily purified by ultracentrifugation, gel-filtration, affinity tag purification and other simple means;

3) the reaction buffer can be exchanged thereby enabling the selection unit to be transferred (with or without purification) to a new reaction mixture and at the same time be diluted by a factor of 100 to 200 (in order to adjust the number of ribosome complexes after emulsification to less than one molecule per reaction compartment);
4) the mRNA library diversity of CRD is limited only by the diversity of the in vitro compartmentalization and is less than 10^{10} different variants;
5) once emulsified, the selection reaction can be performed at a broad range of temperatures from 4° to 94°C because emulsions are stable over these temperatures;
6) if selections performed at elevated temperatures (30°C and above), the ternary complexes will dissociate but remain compartmentalized so as not to lose the genotype-phenotype link, releasing mRNA and in vitro translated protein.

The compartmentalized ribosome display process of the present invention has been found to work particularly well in promoting evolution of new reverse transcriptase enzymes. As an example, M-MuLV reverse transcriptase (Gerard et al., 1986, pRT601) may be used for selection (the enzyme including a N terminal His tag for purification). This reverse transcriptase has a temperature optimum for activity at 42°C and is active at temperatures up to 50°C. An array of mRNAs encoding the M-MuLV reverse transcriptase may be subjected in step (d) of the process of the present invention to reaction conditions which include primer and dNTPs required for cDNA synthesis at an incubation temperature in the range of from 50°C to 60°C. At these elevated temperatures stored ribosomal complexes stable at 4°C quickly dissociate releasing into solution a reverse transcriptase substrate (mRNA) and enzyme.

In one embodiment, in vitro translated reverse transcriptase M-MuLV released from ribosomal complex has C terminal fusion with phage lambda outer surface protein D used in ribosome display construct as a spacer to remain in ribosomal tunnel (Matsuura and Pluckthun, 2003) and covalently bound tRNA, because translation was not terminated properly. Protein D is very well expressed, soluble, stable protein with unfolding transition temperature ~57°C (Forrer and Jaussi, 1998) and therefore is good fusion partner for selection of thermostable reverse transcriptase.

In compartmentalized ribosome display selection method only fully active variants of reverse transcriptase can perform synthesis of cDNA, which encodes the same active enzyme. In 1 hr, after the reverse transcription reaction is completed, emulsion is broken, cDNA is purified and amplified by nested PCR. Because of the selection nature of CRD only cDNA, which encodes active variants of reverse transcriptase, able to perform full length cDNA synthesis, will be amplified and if necessary transferred to the next selection round. In order to eliminate undesirable mutations in T7 polymerase promoter region, ribosome binding site (RBS) and protein D sequence, amplified DNA, encoding only M-MuLV sequence, may be ligated to native 5' and 3' terminal fragments in such way that original ribosome display construct is restored and next selection round can be performed.

In five selection rounds enzyme variants with specific activities at 50° have been identified, which are 2-4 times better as compared to the activity of the primary enzyme used for library preparation. Some of proteins are faster, some - more thermostable. Many selected M-MuLV variants have mutations D524G or D583N, which turn off RNase H activity of reverse transcriptase and improve cDNA synthesis as well as thermostability (Gerard et al., 2002). Many more variants of selected M-MuLV reverse transcriptases have other different beneficial mutations (H204R; H638R; T197A; M289V; E302K; T306A; N454K; Y64C; E69G; Q190R; V223M; F309S; L435P; E562K) mentioned and described before (US7056716; US20060094050A1; US7078208; US20050232934A1; WO07022045A2). In addition to that we have found many new hot spots in reverse transcriptase amino acids sequence. Some mutations repeat very frequently and are of very high importance, what was shown analyzing purified mutants (Example 2). Thus we can state, that our CRD technology is very fast and robust selection method, efficiency of which was confirmed by direct evolution and improvement of M-MuLV reverse transcriptase. As proof of principle we have selected variants of M-MuLV reverse transcriptase working better at higher temperatures.

In a further aspect, the present invention provides a reverse transcriptase enzyme obtainable by the process as described herein.

In a further aspect, the present invention provides a reverse transcriptase enzyme having an optimum activity at a temperature above 42°C, preferably at least 50°C and more preferably in

the range of from 50°C to 60°C. Reverse transcriptase enzymes may be selected according to the process described herein by applying reaction conditions having an elevated temperature, preferably of at least 50°C. In this aspect of the invention, a reverse transcriptase enzyme may be selected which has an activity-temperature profile which is shifted in comparison with wild type enzyme to increase the temperature at which optimum activity is observed.

In a further aspect, the invention provides a reverse transcriptase enzyme which comprises a MMLV reverse transcriptase amino acid sequence with a mutation at one or more of the following amino acid positions:

D200,	D653,	L603,	T330,	L139,	Q221,
T287,	I49,	N479,	H594,	F625,	H126,
A502,	E607,	K658,	P130,	Q237,	N249,
A307,	Y344,	Q430,	D449,	A644,	N649,
L671,	E673,	M39,	Q91,	M66,	W388,
I179	E302	L333	R390	Q374 and	E5

Where the mutation is at D653 it is preferred that the mutation is not D653N. Where the mutation is at L603 it is preferred that the mutation is not L603A. Further, where the mutation is at H594 it is preferred that the mutation is not H594A.

It is preferred that the mutations at the above positions are point mutations.

Preferably, the reverse transcriptase has one or more of the following mutations:

D200N,A or, G,	D653N,G,A,H or, V,	L603W or M,	T330P,	L139P,	Q221R,
T287A,	I49V or T,	N479D,	H594R or Q,	F625S or L,	H126S or R,
A502V,	E607K, G or A,	K658R or Q	P130S,	Q237R,	N249D,
A307V,	Y344H,	Q430R,	D449G or A,	A644V or T,	N649S,
L671P,	E673G or K,	M39V or L,	Q91R or L,	M66L,	W388R,
I179T or V	E302K	L333Q	R390W	Q374R and	E5K

Each of these mutations is found, for example, in mutant enzymes having a higher activity at 50°C as compared with the corresponding wild type enzyme. Further details of these mutations are described in the specific examples.

In particularly preferred aspect of the invention the mutant enzyme has at least two mutations. In one embodiment the two mutations are at D200 and at L603. For example the mutations are D200N and L603W. In an alternative embodiment the mutations are at N479 and H 594. For example the mutations are N479D and H594R.

In a further aspect, the invention provides a reverse transcriptase enzyme having an optimum activity at a temperature above 37°C, wherein the activity at 50°C is at least 120% of the activity at 37°C. Preferably, the activity at 50°C is at least 130%, more preferably at least 160% of the activity at 37°C.

In a further aspect, the present invention provides a mutant reverse transcriptase enzyme having an activity at 50°C which is at least twice that of the corresponding wild type enzyme.

In a further aspect, the present invention provides a mutant reverse transcriptase enzyme having a specific activity at 37°C which is at least 130% of the corresponding wild type enzyme. Preferably, the specific activity of the mutant reverse transcriptase enzyme is at least 140%, more preferably at least 150% and particularly preferably at least 160% of the specific activity of the corresponding wild type enzyme. It has been found as described herein that partially purified wild type enzyme specific activity at 37°C is approximately 200000U/mg. Specific mutant reverse transcriptase enzymes obtainable in accordance with the present invention are discussed in further detail in the specific examples.

In a further aspect, the present invention provides a mutant reverse transcriptase enzyme having a thermostability of at least 1.5 times that of the corresponding wild type enzyme. Thermostability is measured in the present application as residual activity at 37°C following treatment at 50°C for 5 minutes. Preferably, the thermostability of the mutant reverse transcriptase enzyme is at least 1.5 times, more preferably at least 2 times, still more preferably at least 2.5 times that of the corresponding wild type enzyme. Typically, residual activity at

37°C of the wild type reverse transcriptase enzyme is approximately 11% as compared with untreated enzyme.

It is preferred that the reverse transcriptase enzyme according to the invention comprises an MMLV reverse transcriptase.

In a further aspect, the present invention provides a polynucleotide, such as an mRNA or DNA, encoding a reverse transcriptase as described herein.

The reverse transcriptases according to the present invention may be used in a variety of molecular biology techniques such as RT-PCR (qRT-PCR, etc). A kit for RT-PCR may be provided in which the reverse transcriptase of the kit is a reverse transcriptase according to the present invention.

Detailed Description of Invention

The invention will now be described in further detail, by way of example only, with reference to the accompanying figures and appendices:

Figure 1. The experimental scheme of Example 1. Two plasmids pET_his_MLV_pD (encoding Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase fused to protein D spacer) and pET_his_del_pD (encoding inactivated (57 amino acids deletion in pol domain) Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase fused to protein D spacer) were used to synthesize PCR fragments. PCR fragments furtheron are used in transcription reaction and synthesis of mRNA lacking STOP codon at the 3' end. Purified mRNA is mixed with ratio 1 : 50 = MLV (active RT) : del (inactive RT) and used for *in vitro* translation reaction. During translation reaction ribosomal complex synthesizes protein and stops at the end of mRNA lacking STOP codon. Mixture of ternary complexes (TC) is purified by ultracentrifugation on sucrose cushions. Purified ternary complexes ($<3*10^9$ molecules taken) already containing mRNA linked to *in vitro* translated MLV reverse transcriptase are used to prepare reverse transcription reaction mix supplemented with external dNTP set and primer for RT reaction. Ice-cold RT reaction mixture is emulsified giving $\sim1*10^{10}$ water in oil compartments ~2μm in size. Emulsified RT reaction mixture (less than one TC (mRNA +

MLV RT) per compartment is incubated for 1 hr at 42°C in order to perform RT reaction. After the temperature of compartmentalized RT reaction mixture is raised most of TC dissociate releasing mRNA and reverse transcriptase. Successful RT reaction is performed only in compartments containing active MLV reverse transcriptase (MLV_pD) and no cDNA is synthesized in compartments with inactive reverse transcriptase (del_pD). Subsequent PCR amplifies cDNA and enrichment of active reverse transcriptase (MLV_pD) genes over inactive reverse transcriptase (del_pD) is observed.

Figure 2. The structural scheme of pET_his_MLV_pD plasmid.

Figure 3. Example 1 - the agarose gel electrophoresis of first PCR performed on cDNA synthesized during CRD selection. Primers used: RD_Nde (SEQ ID No: 9) and pD_55 (SEQ ID No: 10). Expected length of PCR fragments was 2185 bp for MLV_pD and 2014 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 μl of PCR mix per well.

Figure 4. Example 1 - the agarose gel electrophoresis of nested PCR for partial gene amplification performed on first PCR product. Primers used: M_F (SEQ ID No: 11) and M_2R (SEQ ID No: 12). Expected length of PCR fragments was 907 bp for MLV_pDa and 736 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 μl of PCR mix per well.

Figure 5. Example 1 - the agarose gel electrophoresis of nested PCR for full gene amplification performed on first PCR product. Primers used: M_Esp (SEQ ID No: 13) and M_Eri (SEQ ID No: 14). Expected length of PCR fragments was 2077 bp for MLV_pDa and 1906 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 μl of PCR mix per well.

Figure 6. The experimental scheme of CRD selection in Example 2. PCR fragments encoding mutants library of reverse transcriptase (in fusion with protein D) MLV_pD was used to synthesize mRNA. Purified mRNA was used for *in vitro* translation reaction. Ternary complexes (TC) of mRNA-ribosome-MLV_pD(tRNA) were formed in translation mixture and stabilized by low temperature and high concentration of Mg^{2+} ions. Mixture of TC was purified by ultracentrifugation on sucrose cushions. Precipitated TC was dissolved in ice-cold buffer

(50mM Mg^{2+}) and used to prepare reverse transcription reaction mix supplemented with external dNTP set and primer for RT reaction. Ice-cold RT reaction mixture was emulsified giving ~$1*10^{10}$ water in oil compartments ~2μm in size. Optimal reaction temperature of MLV RT is ~42°C. In order to select for reverse transcriptase variants, which are working better at higher temperatures emulsified RT reaction mixture (less than one TC (mRNA + MLV RT) per compartment) was incubated for 1 hr at 50°C. At this temperature successful synthesis of full length cDNA was performed better in compartments containing more active or thermostable MLV reverse transcriptase variants. Subsequent PCR was used to amplify full length cDNA and enrichment of more active and thermostable reverse transcriptase genes was performed. By PCR amplified genes were moved back to CRD format restoring intact 5' (START fragment - T7 polymerase promoter, SD and his-tag coding sequences) and 3' (END fragment - gs linker, protein D and second gs linker) sequences by ligation PCR. Reconstructed PCR fragment, containing enriched library of reverse transcriptase genes, was used for subsequent mRNA transcription and next CRD selection round. Each selection round was performed at higher and higher temperatures of RT reaction: 50°C (1^{st} round); 52.5°C (2^{nd} round); 55°C (3^{rd} round); 57.5°C (4^{th} round) and 60°C (5^{th} round).

Figure 7. The scheme of PCR fragment reconstruction before new round of CRD selection. Mutated MLV RT library was digested with Esp3I (NcoI compatible end) and EcoRI and ligated with START (244 bp) and END (398 bp) fragments in order to get PCR fragment suitable for CRD selection. START fragment (containing T7 polymerase promoter, SD and his-tag coding sequences) was constructed by PCR amplification of initial 983 bp START fragment (target - plasmid pET_his_del_pD (SEQ ID No: 2), primers - pro-pIVEX (SEQ ID No: 3) and M_1R (SEQ ID No: 15)) and subsequent digestion with NcoI (recognition sequence C↓CATGG) giving 244 bp DNA fragment. END fragment (containing gs linker, protein D and second gs linker sequences) was constructed by PCR amplification of initial 1039 bp END fragment (target - plasmid pET_his_del_pD (SEQ ID No: 2), primers - M_3F (SEQ ID No: 16) and pD-ter (SEQ ID No: 4)) and subsequent digestion with EcoRI (recognition sequence G↓AATTC) giving 398 bp DNA fragment.

Figure 8. Reverse transcriptase activities of mutant RT variants measured at 37°C, 50°C and residual activity at 37°C after 5 min incubation at 50°C. Reverse transcriptase activity at 37°C

is normalized to be always 100% and is omitted. Thus only two types of columns (percents of RT activity at 50°C and residual RT activity at 37°C after 5 min incubation at 50°C) are shown. As a control is given wt M-MuLV reverse transcriptase used for mutants library construction. This primary enzyme is expressed in the same vector and purified in the same way as mutant variants of RT. An average value of mutant RT activity at 50°C for all tested mutants is about ~92% and is more than 2 times higher comparing to wt enzyme (45%). An average residual activity of mutant RT variants at 37°C after 5 min preincubation at 50°C is 12% (wt enzyme - 11%).

Figure 9. Specific activity (u/mg of protein) of partially purified wt and mutant RT variants measured 10 min at 37°C.

Figure 10. The proposed experimental scheme of CRD selection using FACS. Protein of interest is displayed in ribosome display format. Purified (or just diluted many times) ternary complex comprising mRNA-ribosome-protein (tRNA) is mixed with non fluorescent substrate (S) in reaction buffer and subsequently emulsified producing double water-in-oil-in-water emulsions. Active variants of compartmentalized enzymes will convert substrate (S) to fluorescent product (P) allowing FACS to distinguish between fluorescent (active enzyme inside) and "dark" (inactive enzyme inside) droplets.

Figure 11. The proposed experimental scheme for selection and evolution of thermostable DNA polymerases using CRD. Polymerase of interest has to be displayed in ribosome display format. Optionally purified (or just diluted many times) ternary complex comprising mRNA-ribosome-polymerase can be used to prepare reaction mixture with reverse transcriptase (helper enzyme), dNTP's and primer set in PCR buffer. Reaction solution should be emulsified producing water-in-oil emulsion. In first RT step - reverse transcriptase has to synthesize cDNA, which subsequently will serve as a target for second PCR step - cDNA amplification by ribosome displayed DNA polymerase. One of the primers used in PCR can have biotin for optional subsequent purification with streptavidin beads and non-complementary 5' end. After RT-PCR emulsions should be broken, newly synthesized DNA fragment can be purified via biotin and reamplified using new primer set, which will contain one primer with sequence non-complementary to cDNA, but identical to 5' part of primer used in first amplification reaction

(selective amplification of DNA over cDNA background). More active variants of DNA polymerase will be enriched over less active variants and can be used for further analysis or next selection round.

Figure 12. The experimental scheme of Example 4. In this experimental setup M-MuLV reverse transcriptase is used as DNA dependent DNA polymerase. Two plasmids pET_his_MLV_D583N_pD (encoding RNase H minus Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase fused to protein D spacer) and pET_his_del_pD (encoding inactivated reverse transcriptase fused to protein D spacer - 57 amino acids deletion in pol domain and point mutation D583N in RNase H domain) were used to synthesize PCR fragments. PCR fragments furtheron are used in transcription reaction. Purified mRNA is mixed with ratio 1 : 20 = MLV_D583N_pD (active RT) : del_pD (inactive RT) and used to prepare mRNA/dsDNA complex by dsDNA ligation to mRNA mix using T4 DNA ligase. mRNA/dsDNA complex was used for *in vitro* translation reaction. During translation reaction ribosomal complex synthesizes protein and stops at the end of mRNA (at the beginning of mRNA/DNA hybrid). Mixture of ternary complexes (TC) is purified by ultracentrifugation on sucrose cushions. Purified ternary complexes ($<3*10^9$ molecules taken) already containing mRNA/dsDNA linked to *in vitro* translated polymerase (M-MuLV reverse transcriptase) are used to prepare elongation reaction mix supplemented with external biotin-dUTP. Ice-cold reaction mixture is emulsified giving $\sim1*10^{10}$ water in oil compartments ~2μm in size. Emulsified elongation reaction mixture (less than one TC (mRNA/dsDNA + polymerase) per compartment is incubated for 30 min at 37°C in order to incorporate biotinylated nucleotide. After the temperature of compartmentalized reaction mixture is raised most of TC dissociate releasing mRNA/dsDNA complex and polymerase. Successful incorporation reaction in to dsDNA substrate is performed only in compartments containing active polymerase (reverse transcriptase - MLV_D583N_pD) and no cDNA is synthesized in compartments with inactive polymerase (del_pD). After the emulsions are broken excess of biotin-dUTP is removed using gel-filtration mini-column. Biotinylated mRNA/dsDNA complex is purified on streptavidin beads and used to synthesize cDNA. Subsequent PCR amplifies cDNA and enrichment of active polymerse (reverse transcriptase - MLV_D583N_pD) genes over inactive polymerase (del_pD) is observed.

Figure 13. Determination of biotin-dUTP incorporation efficiencies into mRNA/dsDNA complex and into self primed mRNA. Picture of RT-PCR performed on samples of mRNA/dsDNA (MLV_D583N_pD) and mRNA (del_pD) after the incorporation of dTTP or biotin-dUTP. Predicted amplicons size are 907 bp for MLV_D583N_pD and 736 bp for del_pD cDNA. PCR products were analyzed on 1% agarose gel loading 10 μl of PCR mix per well.

Figure 14. General control of mRNA/dsDNA complex existence by incorporation of dTTP (biotin-dUTP) and [α-P^{33}]dATP. Radioactive dATP should be introduced into dsDNA substrate subsequentialy after the incorporation of initial dTTP or biotin-dUTP. A - etidium bromide visualized agarose gel (mRNA or mRNA/dsDNA bands ~2.5 kb). B - the same gel as in A dried on filter paper and visualized using Cyclone Phosphor Imager (Perkin-Elmer, Wellesley, MA). Labelled mRNA/dsDNA complex and/or only dsDNA bands are observed. C - structure and sequence of dsDNA counterpart in mRNA/dsDNA complex.

Figure 15. The resultant final RT-PCR fragments analysis of Example 4. Predicted amplicons size are 907 bp for MLV_D583N_pD and 736 bp for del_pD cDNA. PCR products were analyzed on 1% agarose gel loading 10 μl of PCR mix per well. RT-PCR samples before streptavidin beads corresponds to the ratio of active and inactive polymerase genes 1:20 (almost only del_pD fragment ~736 bp is visibile). RT-PCR samples after the purification on streptavidin beads corresponds to the ratio of active and inactive polymerase genes after the single selection round - 1:1. An enrichment factor of ~20 is observed in this selection.

Figure 16. Some examples of alkaline agarose gels used to determine highest temperature of 1 kb and 4.5 kb cDNA synthesis reaction. PCR machine - Eppendor Mastercycle Gradient. A-D - 1 kb cDNA synthesis (M-MuLV (wt), D200N, L603W and Q221R); temperature gradient 41.9°C, 43.6°C, 45.5°C, 47.8°C, 50.4°C, 53.1°C, 55.8°C, 58.1°C, 60.1°C, 62.1°C; size standart - DNA Fast Ruler Middle Range (Fermentas). E-G - 4.5 kb cDNA synthesis (M2, M3 and M4); temperature gradient 49.8°C, 51.5°C, 53.4°C, 55.7°C, 58.3°C, 61.0°C, 63.7°C, 66.1°C, 68.0°C, 70.0°C; size standart - Zip Ruler Express DNA ladder 2 (Fermentas).

Appendix 1. The general scheme of mutations found in initial MLV RT library (sequence between NcoI and EcoRI restriction sites – SEQ ID No. 24). 23 nucleotide mutations were found among 10 sequenced genes (1 transversion, 20 transitions - mutated positions are underlined, mutations indicated above the sequence, 2 deletions – underlined and indicated as dashed line above the sequence) giving 15 amino acids exchanges, 6 silent mutations, 1 stop codon and 2 frame shifts of coding frame - on average 1-2 amino acids substitutions per gene.

Appendix 2. The CLUSTALW alignment of all 104 protein sequences without N terminal His tag in order to have the same numeration of amino acids as is usually used in literature. Wild type sequence denoted as MLV (SEQ ID No: 25) is always given as first sequence (mutated sequences represent SEQ ID Nos: 26 to 128 based on the order in which they are shown). Mutations are marked using white font color in black background. Amino acids positions, mutations of which somehow improve M-MuLV reverse transcriptase properties and are described in different patent applications are marked in the alignment as columns of amino acids (white font) highlighted in grey. Mutations originating from our selection and located in grey columns indicate that our selection procedure precisely targeted the beneficial hot spot or even exact amino acid mutations described elsewhere. Sequences of analyzed proteins, activity of which at 50°C was substantially better as compared to the primary wt M-MuLV (70% and more as compared to 45% of wt activity) are highlighted in grey.

Appendix 3. List of mutations found in all selected RT variants. Proteins are sorted by decreasing number of mutations.

Appendix 4. Mutations frequency (in decreasing order) of selected RT variants. Names of analyzed proteins, which activity at 50°C was substantially better as compared to primary wt M-MuLV (70% and more comparing to 45% of wt activity) are highlighted in grey.

Appendix 5. Summarized table of data on M-MuLV (wt) reverse transcriptase and single mutants, which contains: name of protein; selection frequency (number of sequenced mutants, which had exact mutation and the number in the parentheses indicates total number of particular amino acid mutations found in selection); protein concentration (mg/ml); reverse transcriptase specific activity at 37°C (u/mg); relative activity at 50°C (%); relative residual

activity at 37°C after 5 min enzyme incubation at 50°C (%); specific RNase H activity of protein (u/mol); relative RNase H activity (%) and highest temperature of 1 kb cDNA synthesis reaction.

Appendix 6. Summarized table of data on M-MuLV (wt) reverse transcriptase and single mutants, which contains: name of protein; protein concentration (mg/ml); reverse transcriptase specific activity at 37°C (u/mg); relative activity at 50°C (%) and highest temperature of 1 kb and 4.5 kb cDNA synthesis reaction.

Appendix 7. Sequence and information relating to SEQ ID Nos: 1 to 23.

Example 1 - *CRD - proof of principle*

To provide proof of principle for Compartmentalized Ribosome Display selection system test selection was performed. Typical proof of principle experiment should give positive signal for active enzyme (in our case RT-PCR fragment for original MLV reverse transcriptase) and no signal for inactive enzyme (no RT-PCR fragment for inactivated MLV reverse transcriptase). More sophisticated experiment is to use mixture of genes with defined ratio encoding active and inactive enzymes. As a result of successful experiment genes encoding active enzyme should be enriched over genes encoding inactive enzyme.

General scheme of experiment is shown in FIG 1. Two plasmids pET_his_MLV_pD (encoding Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase fused to protein D spacer) and pET_his_del_pD (encoding inactivated (57 amino acids deletion in pol domain) Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase fused to protein D spacer) were used to synthesize PCR fragments. Further PCR fragments were used in transcription reaction for synthesis of mRNA, which lacks STOP codon at the 3' end. Purified mRNAs resulting from the two abovementioned PCR fragments were mixed with a ratio 1 : 50 = MLV (active RT) : del (inactive RT) and used for *in vitro* translation reaction. During translation reaction ribosomal complex synthesizes protein and stops at the end of mRNA lacking STOP codon. Translation reaction was stopped by dilution with ice-cold buffer containing 50 mM Mg^{2+}. Low temperature, high concentration of Mg^{2+} ions and absence of STOP codon at the end of mRNA stabilize ternary complexes (TC) of mRNA-ribosome-protein(tRNA). Mixture of

ternary complexes (TC) was purified by ultracentrifugation on sucrose cushions. Ultracentrifugation was optimized in such a way that TC (~3.5 MDa) were precipitated at the bottom of ultracentrifugation tube, meanwhile small molecular weight molecules, proteins and most of free mRNA (~0.9 MDa) remained in the supernatant. Precipitated TCs were dissolved in the ice-cold buffer (50mM Mg^{2+}). Purified ternary complexes (<$3*10^9$ molecules taken) already containing mRNA linked to *in vitro* translated MLV reverse transcriptase were used to prepare reverse transcription reaction mix supplemented with external dNTP set and primer for RT reaction. Ice-cold RT reaction mixture was emulsified yielding ~$1*10^{10}$ water in oil compartments ~2μm in size. Emulsified RT reaction mixture (less than one TC (mRNA + MLV RT) per compartment was incubated for 1 hr at 42°C in order to perform RT reaction. After the temperature of compartmentalized RT reaction mixture is raised most of TCs dissociate releasing mRNA and reverse transcriptase. Successful RT reaction is performed only in compartments containing active MLV reverse transcriptase (MLV_pD) and no cDNA is synthesized in compartments with inactive reverse transcriptase (del_pD). Subsequent PCR ensures the amplification of synthesized cDNA and enrichment of active reverse transcriptase (MLV_pD) genes over inactive reverse transcriptase (del_pD) is observed.

Methods and materials

Initial plasmid pET_his_MLV_pD (SEQ ID No: 1 and FIG 2) was constructed by modification of pET type plasmid in T7 polymerase promoter and Shine-Dalgarno sequences region and insertion of MLV H+ reverse transcriptase coding sequence (306-2363 on SEQ ID No: 1) with N-terminal His-tag (258-305 on SEQ ID No: 1) and C-terminal fusion with glycine-serine (gs) linker (2364-2393 on SEQ ID No: 1), part of protein D (pD) from phage lambda (2394-2669 on SEQ ID No: 1) and second glycine-serine (gs) linker (2670-2759 on SEQ ID No: 1). N-terminal His-tag is used for protein express purification. C-terminal fusion has to remain in ribosome tunnel during protein in vitro translation and formation of mRNA-ribosome-MLV(tRNA) ternary complex.

M-MuLV reverse transcriptase has two main enzymatic activities: RNA dependent DNA polymerase and RNase H. Reverse transcriptase RNase H activity was turned off introducing point mutation D583N (single nucleotide G to A exchange at position 2055 in plasmid pET_his_MLV_pD, SEQ ID No: 1). Aspartate 583 is located in RNase H active site, is

involved in Mg ion binding and is crucial for RNase H activity. New plasmid is identified as pET_his_MLV_D583N_pD and was used for further construction of next plasmid pET_his_del_pD (SEQ ID No: 2), which encodes inactivated reverse transcriptase. Plasmid pET_his_MLV_D583N_pD was digested with restriction endonuclease XmaJI (recognition sequence C↓CTAGG - positions 1047 and 1218 on SEQ ID No: 1). Gene fragment 171bp in length was removed and digested plasmid was self - ligated, yielding plasmid pET_his_del_pD (SEQ ID No: 2), which encodes reverse transcriptase gene shorter by 171 nucleotides or 57 amino acids, without shift in protein translation frame.

It was important to have the same reverse transcriptase gene: 1) shorter in length (for easy PCR detection); 2) inactive (it was confirmed experimentally that deletion of 57 amino acids in polymerase domain completely inactivated polymerase activity and mutation D583N inactivated RNase H activity); and 3) without frameshift (any frameshift will result in appearance of STOP codons, which are not compatible with ribosome display format).

Preparation of PCR fragments for in vitro transcription. PCR mixture was prepared on ice: 20 µl - 10X Taq buffer with KCl (Fermentas); 20 µl - 2 mM of each dNTP (Fermentas); 12 µl - 25 mM $MgCl_2$ (Fermentas); 16 µl - DMSO (D8418 - Sigma); 4 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 1 µl - 100 µM pro-pIVEX primer (SEQ ID No: 3); 1 µl - 100 µM pD-ter primer (SEQ ID No: 4); 122 µl water - mixture divided into two tubes 2x 98 µl. To 2x 98 µl of PCR master mix were added either 2 µl of pET_his_MLV_pD (diluted to ~1 ng/µl) or 2 µl of pET_his_del_pD (diluted to ~1 ng/µl). The cycling protocol was: initial denaturation step 3 min at 94°C, 30 cycles (45 sec at 94°C, 45 sec at 53°C, and 2 min at 72°C) and final elongation 5 min at 72°C. Amplification was ~7000 fold from 2 ng of plasmid (7873 bp) target to ~5 µg (50 ng/µl) of amplified product (2702 bp PCR fragment for pET_his_MLV_pD; 2531 bp PCR fragment for pET_his_del _pD).

Transcription mixture was prepared: 40 µl - 5x T7 transcription buffer (1 M HEPES-KOH pH 7.6; 150 mM Mg acetate; 10 mM spermidin; 0.2 M DTT); 56 µl - 25 mM of each NTP (Fermentas); 8 µl - 20 u/µl T7 RNA polymerase (Fermentas); 4 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 52 µl nuclease-free water - mixture divided into two tubes 2x 80 µl and

add 20 µl - 50 ng/µl of MLV_pD(pro-pIVEX//pD-ter) or 20 µl - 50 ng/µl of del_pD(pro-pIVEX//pD-ter) PCR mixture. Transcription was performed 3 hr at 37°C.

Both transcription mixtures were diluted to 200 µl with ice-cold nuclease-free water and 200 µl of 6 M LiCl solution were added. Mixtures incubated 30 min at +4°C and centrifuged for 30 min at +4°C in cooling centrifuge at max speed (25'000 g). Supernatant was discarded and RNA pellet washed with 500 µl of ice-cold 75% ethanol. Tubes again were centrifuged for 5 min at +4°C in cooling centrifuge at max speed and supernatant was discarded. RNA pellet was dried for 12 min at room temperature and subsequently resuspended in 200 µl of nuclease-free ice-cold water by shaking for 15 min at +4°C and 1400 rpm. Tubes again were centrifuged for 5 min at +4°C in cooling centrifuge at max speed in order to separate not dissolved RNA. About 180 µl of supernatant were moved to new tube with 20 µl of 10x DNase I buffer (Mg^{2+}) (Fermentas); 1 µl - 1 u/µl DNaseI (RNase-free) (Fermentas) and incubated for 20 min at +37°C in order to degrade DNA. To each tube were added 20 µl of 3 M Sodium acetate pH 5.0 solution and 500 µl of ice-cold 96% ethanol. Finally RNA was precipitated by incubation for 30 min at -20°C and centrifugation for 30 min at +4°C in cooling centrifuge at max speed (25'000 g). Supernatant was discarded and RNA pellet washed with 500 µl of ice-cold 75% ethanol. Tubes again were centrifuged for 5 min at +4°C in cooling centrifuge at max speed and supernatant was discarded. RNA pellet was dried for 12 min at room temperature and subsequently resuspended in 43 µl of nuclease-free ice-cold water by shaking for 15 min at +4°C and 1400 rpm. RNA solution was aliqouted 4x 10 µl and liquid nitrogen frozen. Concentration of mRNA was measured spectrophotometrically and double checked on agarose gel using RiboRuler™ RNA Ladder, High Range (Fermentas) - MLV_pD mRNA ~1.2 µg/µl; del_pD mRNA ~1.2 µg/µl.

Purified mRNA is mixed with ratio 1:50 = MLV (active RT) : del (inactive RT). MLV_pD mRNA was diluted 25 times to ~48 ng/µl and 1 µl (~48 ng) was mixed with 2 µl ~1.2 µg/µl del_pD mRNA (2.4 µg) giving mRNA mixture ~0.8 µg/µl with ratio 1:50. In vitro translation was performed using two translation systems RTS 100 E.coli HY Kit (03 186 148 001 - Roche) and synthetic WakoPURE (295-59503 - Wako). Proteins translation sequences are given in SEQ ID No: 6 for MLV_pD and SEQ ID No: 7 for del_pD.

Translation mixture for RTS HY system (25 µl): 6 µl - E.coli lysate (Roche); 5 µl - Reaction Mix (Roche); 6 µl - amino acids (Roche); 0.5 µl - 100 mM Met (Roche); 0.5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 0.4 µl - 200 µM assrA oligonucleotide (SEQ ID No: 5); 0.25 µl - 1 M DTT; 2.5 µl reconstitution buffer (Roche); 2.5 µl nuclease-free water and 1.5 µl - 0.8 µg/µl mRNA mixture 1:50=MLV_pD:del_pD (~1200 ng). In vitro translation was performed for 20 min at 30°C.

Translation mixture for WakoPURE system (25 µl): 12.5 µl - A solution (Wako); 5 µl - B solution (Wako); 0.5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 0.4 µl - 200 µM αssrA oligonucleotide (SEQ ID No: 6); 0.25 µl - 1 M DTT; 5 µl nuclease-free water and 1.5 µl - 0.8 µg/µl mRNA mixture 1:50=MLV_pD:del_pD (~1200 ng). In vitro translation was performed for 30 min at 37°C.

Both translations (~25 µl) were stopped by adding 155 µl of ice-cold stopping buffer WBK_{500}+DTT+triton (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - triton x-100 (T8787 - Sigma)) and centrifuged for 5 min at +4°C and 25'000 g. Very carefully 160 µl of centrifuged translation mixture was pippeted on the top of 840 µl 35% (w/v) sucrose solution in WBK_{500}+DTT+triton (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - triton x-100 (T8787 - Sigma); 35% (w/v) - sucrose (84097 - Fluka)). In order to purify ternary complexes (TC) of mRNA-ribosome-protein(tRNA) ultracentrifugation was performed using TL-100 Beckman ultracentrifuge; TLA100.2 fixed angle rotor (Beckman); transparent 1 ml ultracentrifugation tubes (343778 - Beckman) for 9 min at +4°C and 100'000 rpm. In order to keep small transparent pellet of TC at the bottom of ultracentrifugation tube intact tubes were handled with care. Initially 750 µl of solution was removed from the very top of the centrifugation tube. Then (very carefully) tube walls were washed with 750 µl of WBK_{500} (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl). Finally all solution was removed starting from the very top of the centrifugation tube and pellet was dissolved in 30 µl of ice-cold stopping buffer WBK_{500}+DTT+triton (50 mM tris-acetate pH 7.5

at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - triton x-100 (T8787 - Sigma)).

As it was determined using radioactively labeled mRNA after ultracentrifugation 5% - 30 % of input mRNA is located in ternary complex pellet. Therefore it was expected to have less than 360 ng (30% from 1200 ng mRNA used in translation reaction) of mRNA in 30 µl of buffer (~12 ng/µl or $9*10^9$ molecules/µl of ternary complex).

Reverse transcription reaction mixture for selection was prepared on ice: 60 µl - 5X reaction buffer for Reverse Transcriptase (Fermentas); 7.5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 15 µl - 20 µM pD_42 oligonucleotide (SEQ ID No: 8); 188 µl nuclease-free water - mixture divided into two tubes 2x 135 µl and 0.9 µl of purified ($<8*10^9$ molecules) TC (translation in Roche - RTS HY kit) or 0.9 µl of purified ($<8*10^9$ molecules) TC (translation in Wako - WakoPURE) were added. Each reaction mixture (~135 µl) again was divided into two tubes 45 µl and 90 µl. To the first part - 45 µl of RT mixture 5 µl of nuclease-free water were added. This sample is considered to be the *negative selection control* (without dNTP) and has to prove that there is no DNA contamination in the reaction mixture and cDNA synthesis is strictly linked to reverse transcriptase functional activity coming from MLV RT in ternary complex. To the second part - 90 µl of RT mixture 10 µl - 10 mM each dNTP Mix (Fermentas) were added and reaction mixture again was divided into two tubes - 50 µl for *selection control* and 50 µl supplemented with 1 µl - 200 u/µl of RevertAid H- M-MuLV Reverse Transcriptase (Fermentas) for *positive selection control.* According to the protocol each reverse transcription reaction mix contains $<2.7*10^9$ molecules of ternary complex in 50 µl volume.

Oil-surfactant mixture for emulsification was prepared by mixing ABIL EM 90 (Goldschmidt) into mineral oil (M5904 - Sigma) to final concentration of 4% (v/v) (Ghadessy and Holliger, 2004; US2005064460). Emulsions were prepared at +4°C in 5 ml cryogenic vials (430492 - Corning) by mixing 950 µl of oil-surfactant mixture with 50 µl of RT mixture. Mixing was performed using MS-3000 magnetic stirrer with speed control (Biosan) at ~2100 rpm; Rotilabo®- (3x8 mm) magnetic followers with centre ring (1489.2 - Roth); water phase was added by 10 µl aliquots every 30 sec, continuing mixing for 2 more minutes (total mixing time

- 4 min). According to optical microscopy data compartments in our emulsions vary from 0.5 μm to 10 μm in size with average diameter of ~2 μm. Therefore it was expected to have ~$1*10^{10}$ water in oil compartments after the emulsification of 50 μl reverse transcription reaction mixture, which contains less than $2.7*10^{9}$ molecules of ternary complex (about 1 mRNA and reverse transcriptase molecules per 3-4 compartments).

All six emulsions representing RT reactions with TC, translation in Roche - RTS HY kit (*negative selection control*, *selection control* and *positive selection control*) and with TC, translation in Wako - WakoPURE (*negative selection control*, *selection control* and *positive selection control*) were incubated 1 hr at +42°C.

To recover the reaction mixtures emulsions were moved to 1.5 ml tube, centrifuged for 1 min at room temperature and 25'000 g. Oil phase was removed leaving concentrated (but still intact) emulsion at the bottom of the tube and 250 μl of PB buffer (Qiagen PCR purification kit) were added. Finally emulsions were broken by extraction with 0.9 ml water-saturated ether; 0.9 ml water-saturated ethyl-acetate (in order to remove ABIL EM 90 detergent) and again 0.9 ml water-saturated ether. Water phase was dried for 5 min under vacuum at room temperature. Synthesized cDNA was purified with Qiagen PCR purification kit and eluted in 30 μl of EB buffer (Qiagen PCR purification kit).

Amplification of cDNA was performed by nested PCR. Initial PCR mixture was prepared on ice: 16 μl - 10X Taq buffer with KCl (Fermentas); 16 μl - 2 mM of each dNTP (Fermentas); 9.6 μl - 25 mM $MgCl_2$ (Fermentas); 3.2 μl - 1 u/μl LC (recombinant) Taq DNA Polymerase (Fermentas); 0.8 μl - 100 μM RD_Nde primer (SEQ ID No: 9); 0.8 μl - 100 μM pD_55 primer (SEQ ID No: 10); 74 μl water - mixture was divided into 6 samples x 15 μl (6x15 μl) and 30 μl. To 6x 15 μl of PCR master mix were added 5 μl of cDNA (1-6 RT samples); to 30 μl of PCR master mix were added 9 μl water and mixture again divided into two tubes 2x 19.5 μl for negative PCR control (plus 0.5 μl water) and positive PCR control (plus 0.5 μl - 1:1 mixture (~1ng) of pET_his_MLV_pD and pET_his_del_pD plasmids). The cycling protocol was: initial denaturation step 3 min at 94°C, 25 cycles (45 sec at 94°C, 45 sec at 58°C, and 2 min at 72°C) and final elongation 5 min at 72°C. Expected length of PCR fragments was 2185 bp for

MLV_pD and 2014 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 µl of PCR mix per well (FIG 3).

Nested PCR was performed using two different sets of primers, giving either partial gene amplification (for better resolution of MLV:del cDNA ratio in RT samples) or full gene amplification (to prove possibility of full gene recovery).

Nested PCR mixture for partial gene amplification was prepared on ice: 28 µl - 10X Taq buffer with KCl (Fermentas); 28 µl - 2 mM of each dNTP (Fermentas); 16.8 µl - 25 mM $MgCl_2$ (Fermentas); 5.6 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 1.4 µl - 100 µM M_F primer (SEQ ID No: 11); 1.4 µl - 100 µM M_2R primer (SEQ ID No: 12); 185 µl water - mixture divided 2x 19 µl and 6x 38 µl. To 2x 19 µl of PCR master mix was added 1 µl of positive or negative controls of first PCR (primers set RD_Nde//pD_55) - 30 PCR cycles amplification; to 6x 38 µl of PCR master mix were added 2 µl of first PCR (primers set RD_Nde//pD_55) (1-6 samples) - each sample again was divided into two 2x 20 µl for 23 or 30 PCR cycles amplification. The cycling protocol was: initial denaturation step 3 min at 94°C, 23 or 30 cycles (45 sec at 94°C, 45 sec at 57°C, and 1 min at 72°C) and final elongation 5 min at 72°C. Expected length of PCR fragments was 907 bp for MLV_pD and 736 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 µl of PCR mix per well (FIG 4).

Nested PCR mixture for full gene amplification was prepared on ice: 28 µl - 10X Taq buffer with KCl (Fermentas); 28 µl - 2 mM of each dNTP (Fermentas); 16.8 µl - 25 mM $MgCl_2$ (Fermentas); 5.6 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 1.4 µl - 100 µM M_Esp primer (SEQ ID No: 13); 1.4 µl - 100 µM M_Eri primer (SEQ ID No: 14); 185 µl water - mixture divided 2x 19 µl and 6x 38 µl. To 2x 19 µl of PCR master mix was added 1 µl of positive or negative controls of first PCR (primers set RD_Nde//pD_55) - 30 PCR cycles amplification; to 6x 38 µl of PCR master mix were added 2 µl of first PCR (primers set RD_Nde//pD_55) (1-6 samples) - each sample again was divided into 2x 20 µl for 23 or 30 PCR cycles amplification. The cycling protocol was: initial denaturation step 3 min at 94°C, 23 or 30 cycles (45 sec at 94°C, 45 sec at 55°C, and 2 min at 72°C) and final elongation 5 min at

72°C. Expected length of PCR fragments was 2077 bp for MLV_pD and 1906 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 μl of PCR mix per well (FIG 5).

Results

To demonstrate proof of principle for Compartmentalized Ribosome Display (CRD) method selection was performed using starting 1:50=MLV:del mixture of two mRNA encoding active (MLV) and inactive (del) reverse transcriptases fused to protein D spacer (FIG 1). In vitro translation was performed using two different translation systems Roche - RTS 100 E.coli HY or Wako - WakoPURE in order to understand which translation system is better in our experimental setup. For each translation system three compartmentalized RT reactions were performed: *negative selection control* without dNTP, which has to prove that there is no DNA contamination in the reaction mixture; *selection control*, which has to demonstrate the enrichment of genes encoding active (MLV) reverse transcriptase over genes encoding inactivated enzyme (del), because only active enzyme can synthesize cDNA; and *positive selection control* supplemented with external RevertAid H- commercial reverse transcriptase, which has to serve as positive RT control synthesizing cDNA from both MLV_pD and del_pD mRNA in all compartments, showing real ratio of genes in reaction mixture without selection pressure applied.

Synthesyzed cDNA was amplified by nested PCR. The picture of agarose gel electrophoresis of initial PCR (25 cycles) (FIG 3) shows weak PCR fragments bands only in case of both positive selection controls (translation systems - Roche and Wako). This is normal, because these samples contain external RT enzyme, which synthesizes cDNA much more efficiently comparing to reactions containing only one in vitro synthesized reverse transcriptase molecule per compartment.

The picture of agarose gel electrophoresis of nested PCR (partial gene amplification) is shown in FIG 4. Amplification results after 23 and 30 PCR cycles are consistent:

1) there is no amplification (no DNA contamination) in *negative selection controls* (w/o dNTP);

2) very efficient amplification of del_pD cDNA (736 bp DNA fragment) is observed in *positive selection controls* (external RT enzyme) and no MLV_pD cDNA amplification is visible, because initial ratio of MLV_pD to del_pD mRNA is 1:50;

3) amplification of both cDNA MLV_pD (907 bp DNA fragment) and del_pD (736 bp DNA fragment) are observed in case of *selection controls*;

4) ratio of MLV_pD:del_pD ~ 1:1 is observed in case of reverse transcriptase synthesized by Roche in vitro translation system, what means ~50 times enrichment of MLV_pD genes over del_pD genes starting from initial 1:50 ratio;

5) ratio of MLV_pD:del_pD ~ 1:3 is observed in case of reverse transcriptase synthesized by Wako in vitro translation system, what means ~16 times enrichment of MLV_pD genes over del_pD genes starting from initial 1:50 ratio.

The picture of agarose gel electrophoresis of nested PCR (full gene amplification) is shown in FIG 5. Amplification results after 23 and 30 PCR cycles are consistent in between and comparing to results of nested PCR used for partial gene amplification (FIG 5):

1) there is no amplification (no DNA contamination) in *negative selection controls* (w/o dNTP);

2) very efficient amplification of del_pD cDNA (1906 bp DNA fragment) is observed in *positive selection controls* (external RT enzyme) and no MLV_pD cDNA amplification is visible, because initial ratio of MLV_pD to del_pD mRNA is 1:50;

3) amplification of both cDNA MLV_pD (2077 bp DNA fragment) and del_pD 1906 bp DNA fragment) are observed in case of *selection controls*;

4) it is difficult to determine ratio of MLV_pD:del_pD in case of full gene amplification, because relative difference between 2077 bp (MLV_pD) and 1906 bp (del_pD) DNA fragments is not big enough, but in general ratios are similar to results of nested PCR used for partial gene amplification.

As a result of this example we can conclude that during reverse transcription reaction performed in CRD format we have enriched genes encoding active MLV reverse transcriptase over the genes encoding inactive enzyme by a factor of 50 in case of Roche translation system or by a factor of 16 in case of Wako translation system used to synthesize enzymes in vitro.

Example 2 - *CRD - selection for reverse transcriptase, which shows improved performance at higher temperatures*

In order to understand, how efficiently Compartmentalized Ribosome Display (CRD) selection works, evolution experiment of Moloney Murine Leukemia Virus reverse transcriptase (M-MuLV RT) was performed. General scheme of experiment is shown in FIG 6. Initial mutants library of reverse transcriptase was constructed by error-prone PCR using nucleotide analogues dPTP and 8-oxo-dGTP. Whole gene (~2 kb) mutagenesis was performed introducing 2-3 nucleotides or 1-2 amino acids mutations per gene. PCR fragment encoding mutants library of reverse transcriptase (in fusion with protein D) MLV_pD was used to synthesize mRNA. Purified mRNA was used for *in vitro* translation reaction. Ternary complexes (TC) of mRNA-ribosome-MLV_pD(tRNA) were formed in translation mixture and stabilized by low temperature and high concentration of Mg^{2+} ions. Mixture of TC was purified by ultracentrifugation on sucrose cushions. Precipitated TC was dissolved in ice-cold buffer (50mM Mg^{2+}) and used to prepare reverse transcription reaction mix supplemented with external dNTP set and primer for RT reaction. Ice-cold RT reaction mixture was emulsified giving ~$1*10^{10}$ water in oil compartments ~2μm in size. Optimal reaction temperature of MLV RT is ~42°C. In order to select for reverse transcriptase variants, which are working better at higher temperatures emulsified RT reaction mixture (less than one TC (mRNA + MLV RT) per compartment) was incubated for 1 hr at 50°C. At this temperature successful synthesis of full length cDNA was performed better in compartments containing more active or thermostable MLV reverse transcriptase variants. Subsequent PCR was used to amplify full length cDNA and enrichment for more active and thermostable reverse transcriptase genes was performed. By PCR amplified genes were moved back to CRD format restoring intact 5' (START fragment - T7 polymerase promoter, SD and his-tag coding sequences) and 3' (END fragment - gs linker, protein D and second gs linker) sequences by ligation-PCR.

Reconstructed PCR fragment, containing enriched library of reverse transcriptase genes, was used for subsequent mRNA transcription and next CRD selection round. Each selection round was performed at higher and higher temperatures of RT reaction: 50°C (1^{st} round); 52.5°C (2^{nd} round); 55°C (3^{rd} round); 57.5°C (4^{th} round) and 60°C (5^{th} round).

Amplified library of reverse transcriptase genes (without C terminal pD linker) after 5th selection round was cloned into plasmid vector. Individual clones were sequenced and analyzed. Pool of evolved proteins as well as individual mutants were purified via his-tag using affinity chromatography. MLV reverse transcriptase specific activities at 37°C, 50°C and residual activity at 37°C after 5 min enzyme incubation at 50°C were determined.

Methods and materials

Initial plasmid pET_his_MLV_pD (SEQ ID No: 1 and FIG 2) was used as a starting material for error-prone PCR. Mutations were introduced using nucleotide analogues dPTP and 8-oxo-dGTP. PCR mixture for error prone PCR was prepared on ice: 10 µl - 10X Taq buffer with KCl (Fermentas); 10 µl - 2 mM of each dNTP (Fermentas); 6 µl - 25 mM $MgCl_2$ (Fermentas); 2 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 0.5 µl - 100 µM M_Esp primer (SEQ ID No: 13); 0.5 µl - 100 µM M_Eri primer (SEQ ID No: 14); 1 µl - 10 µM dPTP (TriLink BioTechnolgies); 5 µl - 100 µM 8-oxo-dGTP (TriLink BioTechnolgies); 3.75 µl - 40 ng/µl (totally 150 ng) of pET_his_MLV_pD plasmid; 61.25 µl water. The cycling protocol was: initial denaturation step 3 min at 94°C, 30 cycles (30 sec at 94°C, 30 sec at 55°C, and 2 min at 72°C) and final elongation 5 min at 72°C. Amplification was 150-300 fold from 150 ng of plasmid (7873 bp) target to ~6-12 µg of amplified product (2077 bp PCR fragment for pET_his_MLV_pD). PCR fragment was purified using Qiagen PCR purification kit, digested with Esp3I (recognition sequence CGTCTC (1/5)) and EcoRI (recognition sequence G↓AATTC) and finaly purified from agarose gel using Qiagen Gel extraction kit giving DNA concentration ~50 ng/µl.

Mutagenesis efficiency and library quality was checked by sequencing of individual clones subcloned back into original pET_his_MLV_pD plasmid digested with NcoI and EcoRI. As it was expected mutations were distributed randomly all over the amplified sequence of MLV RT gene (Appendix 1). Among 10 sequenced genes 23 nucleotide mutations (1 transversion, 20 transitions, 2 deletions – labelled red in Appendix 1) were found giving 15 amino acids exchanges, 6 silent mutations, 1 stop codon and 2 frame shifts of coding frame - on average 1-2 amino acids substitutions per gene.

Mutated library was ligated with START (244 bp) and END (398 bp) fragments in order to get PCR fragment suitable for CRD selection (FIG 7). START fragment (containing T7 polymerase promoter, SD and his-tag coding sequences) was constructed by PCR amplification of initial 983 bp START fragment (target - plasmid pET_his_del_pD (SEQ ID No: 2), primers - pro-pIVEX (SEQ ID No: 3) and M_1R (SEQ ID No: 15)) and subsequent digestion with NcoI (recognition sequence C↓CATGG) giving 244 bp DNA fragment. END fragment (containing gs linker, protein D and second gs linker sequences) was constructed by PCR amplification of initial 1039 bp END fragment (target - plasmid pET_his_del_pD (SEQ ID No: 2), primers - M_3F (SEQ ID No: 16) and pD-ter (SEQ ID No: 4)) and subsequent digestion with EcoRI (recognition sequence G↓AATTC) giving 398 bp DNA fragment.

Ligation reaction (150µl) was prepared at room temperature: 15 µl - 10X ligation buffer for T4 DNA Ligase (Fermentas); 15 µl - 1 u/µl T4 DNA ligase (Fermentas); 26 µl - 50 ng/µl mutated MLV RT library digested with Esp3I (NcoI compatible end) and EcoRI (~1300ng or ~$5.9*10^{11}$ molecules); 9.4 µl - 35 ng/µl START fragment digested with NcoI (~329 ng or ~$1.2*10^{12}$ molecules); 15.7 µl - 35 ng/µl END fragment digested with EcoRI (~548 ng or ~$1.2*10^{12}$ molecules); 68.9 µl - water. Ligation was performed overnight at +4°C. Reaction mixture was treated once with phenol and twice with chloroform, precipitated and dissolved in 53 µl of water. Approximate ligation yield ~20% was determined comparing amplification efficiency of ligation mixture and known amount of plasmid pET_his_MLV_pD. Taking into account 20% ligation yield diversity of MLV RT mutants library was defined as ~$1.2*10^{11}$ molecules (50 µl).

Ligated MLV RT library was amplified by PCR (1 ml - prepared on ice): 100 µl - 10X Taq buffer with KCl (Fermentas); 100 µl - 2 mM of each dNTP (Fermentas); 60 µl - 25 mM $MgCl_2$ (Fermentas); 80 µl - DMSO (D8418 - Sigma); 20 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 5 µl - 100 µM pro-pIVEX primer (SEQ ID No: 3); 5 µl - 100 µM pD-ter- primer (SEQ ID No: 17); 20 µl - ligated MLV RT library (~$5*10^{10}$ molecules); 610 µl - water. The cycling protocol was: initial denaturation step 3 min at 94°C, 15 cycles (30 sec at 94°C, 30 sec at 53°C, and 3 min at 72°C) and final elongation 5 min at 72°C. Amplification

was ~200 fold from ~$5*10^{10}$ molecules (corresponds to ~150 ng) of final ligation fragment 2702 bp in size target to ~30 µg (30 ng/µl) of amplified product (2702 bp PCR fragment).

1st selection round

Transcription mixture (100 µl) was prepared: 20 µl - 5x T7 transcription buffer (1 M HEPES-KOH pH 7.6; 150 mM Mg acetate; 10 mM spermidin; 0.2 M DTT); 28 µl - 25 mM of each NTP (Fermentas); 4 µl - 20 u/µl T7 RNA polymerase (Fermentas); 2 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 30 µl - 30 ng/µl of mutants library (pro-pIVEX//pD-ter-) PCR mixture (~900 ng or ~$3*10^{11}$ molecules); 16 µl nuclease-free water. Transcription was performed 3 hr at 37°C (library diversity ~$5*10^{10}$ molecules).

Transcription mixture was diluted to 200 µl with ice-cold nuclease-free water and 200 µl of 6 M LiCl solution were added. Mixture incubated 30 min at +4°C and centrifuged for 30 min at +4°C in cooling centrifuge at max speed (25'000 g). Supernatant was discarded and RNA pellet washed with 500 µl of ice-cold 75% ethanol. Tube again was centrifuged for 5 min at +4°C in cooling centrifuge at max speed and supernatant was discarded. RNA pellet was dried for 12 min at room temperature and subsequently resuspended in 200 µl of nuclease-free ice-cold water by shaking for 15 min at +4°C and 1400 rpm. Tube again was centrifuged for 5 min at +4°C in cooling centrifuge at max speed in order to separate not dissolved RNA. About 180 µl of supernatant were moved to new tube with 20 µl of 10x DNase I buffer (Mg^{2+}) (Fermentas); 1 µl - 1 u/µl DNaseI (RNase-free) (Fermentas) and incubated for 30 min at +37°C in order to degrade DNA. To reaction mixture were added 20 µl of 3 M Sodium acetate pH 5.0 solution and 500 µl of ice-cold 96% ethanol. Finally RNA was precipitated by incubation for 30 min at -20°C and centrifugation for 30 min at +4°C in cooling centrifuge at max speed (25'000 g). Supernatant was discarded and RNA pellet washed with 500 µl of ice-cold 75% ethanol. Tube again was centrifuged for 5 min at +4°C in cooling centrifuge at max speed and supernatant was discarded. RNA pellet was dried for 12 min at room temperature and subsequently resuspended in 33 µl of nuclease-free ice-cold water by shaking for 10 min at +4°C and 1400 rpm. RNA solution was aliqouted 3x 10 µl and liquid nitrogen frozen. Concentration of mRNA was measured spectrophotometrically and double checked on agarose

gel using RiboRuler™ RNA Ladder, High Range (Fermentas) - MLV RT library mRNA ~2.1 μg/μl.

In vitro translation was performed using RTS 100 E.coli HY (03 186 148 001 - Roche) translation system (25 μl): 6 μl - E.coli lysate (Roche); 5 μl - Reaction Mix (Roche); 6 μl - amino acids (Roche); 0.5 μl - 100 mM Met (Roche); 0.5 μl - 40 u/μl RiboLock RNase inhibitor (Fermentas); 0.4 μl - 200 μM assrA oligonucleotide (SEQ ID No: 5); 0.25 μl - 1 M DTT; 3 μl reconstitution buffer (Roche); 2.5 μl nuclease-free water and 0.6 μl - 2.1 μg/μl mRNA (~1200 ng). Reaction mixture was incubated for 20 min at 30°C. Translation was stopped adding 155 μl of ice-cold stopping buffer WBK_{500}+DTT+triton (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - triton x-100 (T8787 - Sigma)) and centrifuged for 5 min at +4°C and 25'000 g. Very carefully 160 μl of centrifuged translation mixture was pippeted on the top of 840 μl 35% (w/v) sucrose solution in WBK_{500}+DTT+triton (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - triton x-100 (T8787 - Sigma); 35% (w/v) - sucrose (84097 - Fluka)). In order to purify ternary complexes (TC) of mRNA-ribosome-MLV(tRNA) ultracentrifugation was performed using TL-100 Beckman ultracentrifuge; TLA100.2 fixed angle rotor (Beckman); transparent 1 ml ultracentrifugation tubes (343778 - Beckman) for 9 min at +4°C and 100'000 rpm. In order to keep small transparent pellet of TC at the bottom of ultracentrifugation tube intact tubes were handled with care. Initially 750 μl of solution was removed from the very top of the centrifugation tube. Than (very carefully) tube walls were washed with 750 μl of WBK_{500} (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl). Finally all solution was removed starting from the very top of the centrifugation tube and pellet was dissolved in 30 μl of ice-cold stopping buffer WBK_{500}+DTT+triton (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - triton x-100 (T8787 - Sigma)).

As it was determined experimentally, using radioactively labeled mRNA, ultracentrifugation yields 5% - 30 % of input mRNA in ternary complex pellet. Therefore it was expected to have less than 360 ng (30% from 1200 ng mRNA used in translation reaction) of mRNA in 30 μl of buffer (~12 ng/μl or $9*10^9$ molecules/μl of ternary complex).

Reverse transcription reaction mixture for selection was prepared on ice: 60 µl - 5X reaction buffer for Reverse Transcriptase (Fermentas); 7.5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 15 µl - 20 µM pD_42 oligonucleotide (SEQ ID No: 8); 186 µl nuclease-free water and 1.8 µl of purified ($<1.8*10^{10}$ molecules) TC (translation in Roche - RTS HY kit). Reaction mixture was divided into two tubes 45 µl and 225 µl. To first part - 45 µl of RT mixture were added 5 µl of nuclease-free water. This sample was considered to be *negative selection control* (without dNTP) and has to prove that there is no DNA contamination in the reaction mixture and cDNA synthesis is strictly linked to reverse transcriptase functional activity coming from MLV RT in ternary complex. To second part - 225 µl of RT mixture were added 25 µl - 10 mM each dNTP Mix (Fermentas) and reaction mixture again was divided into two tubes - 200 µl (4x 50 µl) for *selection control* ($<1.2*10^{10}$ molecules of TC totally) and 50 µl supplemented with 1 µl - 200 u/µl of RevertAid H- M-MuLV Reverse Transcriptase (Fermentas) for *positive selection control.* According to protocol each reverse transcription reaction mix contains $<3*10^{9}$ molecules of ternary complex in 50 µl volume.

Oil-surfactant mixture for emulsification was prepared by mixing ABIL EM 90 (Goldschmidt) into mineral oil (M5904 - Sigma) to final concentration of 4% (v/v) (Ghadessy and Holliger, 2004; US2005064460). Emulsions were prepared at +4°C in 5 ml cryogenic vials (430492 - Corning) by mixing 950 µl of oil-surfactant mixture with 50 µl of RT mixture. Mixing was performed using MS-3000 magnetic stirrer with speed control (Biosan) at ~2100 rpm; Rotilabo®- (3x8 mm) magnetic followers with centre ring (1489.2 - Roth); water phase was added by 10 µl aliquots every 30 sec, continuing mixing for 2 more minutes (total mixing time - 4 min). According to optical microscopy data compartments in our emulsions vary from 0.5 µm to 10 µm in size with average diameter of ~2 µm. Therefore it was expected to have $\sim1*10^{10}$ water in oil compartments after the emulsification of 50 µl reverse transcription reaction mixture, which contains less than $3*10^{9}$ molecules of ternary complex (about 1 mRNA and reverse transcriptase molecules per 3-4 compartments).

All emulsions were incubated 1 hr at +50°C in order to select for reverse transcriptase variants, which work better at higher temperatures. To recover the reaction mixtures emulsions were

moved to 1.5 ml tube, centrifuged for 10 min at room temperature and 25'000 g. Oil phase was removed leaving concentrated (but still intact) emulsion at the bottom of the tube. Emulsions were broken by extraction with 0.9 ml water-saturated ether; 0.9 ml water-saturated ethyl-acetate (in order to remove ABIL EM 90 detergent) and again 0.9 ml water-saturated ether. Water phase was dried for 5 min under vacuum at room temperature and 250 µl of PB buffer (Qiagen PCR purification kit) were added. Four selection samples were merged into two tubes. Synthesized cDNA was further purified with Qiagen PCR purification kit and eluted in 30 µl of EB buffer (Qiagen PCR purification kit) in case of negative and positive selection controls and 2x 30 µl in case of selection control.

Amplification of cDNA was performed by nested PCR. First of all small PCR amplification was performed in order to check negative and positive selection controls and determine minimal number of PCR cycles required for efficient amplification of cDNA in selection samples. PCR mixture (200 µl) was prepared on ice: 20 µl - 10X Taq buffer with KCl (Fermentas); 20 µl - 2 mM of each dNTP (Fermentas); 12 µl - 25 mM $MgCl_2$ (Fermentas); 2.5 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 1 µl - 2.5 u/µl Pfu DNA Polymerase (Fermentas); 1 µl - 100 µM RD_Nde primer (SEQ ID No: 9); 1 µl - 100 µM pD_55 primer (SEQ ID No: 10); 50 µl - purified cDNA of selection controls; 92 µl water. The cycling protocol for 2185 bp PCR fragment was: initial denaturation step 3 min at 94°C, 25 cycles (45 sec at 94°C, 45 sec at 58°C, and 2 min at 72°C) and final elongation 5 min at 72°C.

Nested PCR mixture (500 µl) for full gene amplification was prepared on ice: 50 µl - 10X Taq buffer with KCl (Fermentas); 50 µl - 2 mM of each dNTP (Fermentas); 30 µl - 25 mM $MgCl_2$ (Fermentas); 6.25 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 2.5 µl - 2.5 u/µl Pfu DNA Polymerase (Fermentas); 2.5 µl - 100 µM M_Esp primer (SEQ ID No: 13); 2.5 µl - 100 µM M_Eri primer (SEQ ID No: 14); 50 µl of first PCR (primers set RD_Nde//pD_55); 306 µl water. The cycling protocol for 2077 bp PCR fragment was: initial denaturation step 3 min at 94°C, 22 cycles (45 sec at 94°C, 45 sec at 55°C, and 2 min at 72°C) and final elongation 5 min at 72°C.

Final PCR fragment of selection sample was agarose-gel purified using Qiagen gel extraction kit (elution in 60 µl ~50 ng/µl). Purified PCR fragment was digested with EcoRI and Esp3I for 1 hr at 37°C and again agarose-gel purified (elution in 30 µl ~50 ng/µl).

Recovered MLV reverse transcriptase library after 1st selection round was ligated with START and END fragments (construction described earlier in this example) in order to get PCR fragment (FIG 7) suitable for 2nd round of CRD selection (FIG 6). Ligation reaction (40µl) was prepared at room temperature: 4 µl - 10X ligation buffer for T4 DNA Ligase (Fermentas); 2 µl - 1 u/µl T4 DNA ligase (Fermentas); 4 µl - 50 ng/µl selected library digested with Esp3I and EcoRI (~200ng or ~$0.9*10^{11}$ molecules); 1.1 µl - 35 ng/µl START fragment digested with NcoI (~35 ng or ~$1.5*10^{11}$ molecules); 1.76 µl - 35 ng/µl END fragment digested with EcoRI (~61 ng or ~$1.5*10^{11}$ molecules); 27.2 µl - water. Ligation was performed 1 hr at room temperature.

Ligated MLV RT library was amplified by PCR (300 µl - prepared on ice): 30 µl - 10X Taq buffer with KCl (Fermentas); 30 µl - 2 mM of each dNTP (Fermentas); 18 µl - 25 mM $MgCl_2$ (Fermentas); 24 µl - DMSO (D8418 - Sigma); 3.7 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 1.5 µl - 2.5 u/µl Pfu DNA Polymerase (Fermentas); 1.5 µl - 100 µM pro-pIVEX primer (SEQ ID No: 3); 1.5 µl - 100 µM pD-ter- primer (SEQ ID No: 17); 25.5 µl - ligated MLV RT library (<$0.6*10^{10}$ molecules); 164.3 µl - water. The cycling protocol for 2702 bp PCR fragment was: initial denaturation step 3 min at 94°C, 15 cycles (45 sec at 94°C, 45 sec at 53°C, and 3 min at 72°C) and final elongation 5 min at 72°C. PCR fragment was agarose-gel purified using Qiagen gel exraction kit (elution in 30 µl ~100 ng/µl).

2nd selection round

Second selection round was performed following general setup of 1st selection round of experimental scheme with minor modifications in PCR cycles, emulsified RT reaction temperature and few more details.

All modifications are given below:

- transcription - 10 µl - 100 ng/µl (~1000ng) of agarose-gel purified PCR fragment were taken; final concentration of mRNA used in 2nd slection round was 1.5 µg/µl;
- translation - 0.8 µl - 1.5 µg/µl (~1.2 µg) of mRNA were taken;
- emulsified RT reaction was performed 1 hr at 52.5°C;
- 1st PCR (RD_Nde//pD_55) - 24 cycles were performed;
- 2nd (nested) PCR (M_Esp//M_Eri) - 23 cycles were performed;
- final concentration of digested PCR fragment - 80 ng/µl;
- ligation - 200 ng (~$0.9*10^{11}$ molecules) of MLV RT library were taken;
- PCR (on ligation mix) - <$0.6*10^{10}$ molecules of selected library were taken and 15 PCR cycles were performed; concentration of final agarose-gel purified PCR fragment was 200 ng/µl.

3rd selection round

Third selection round was performed following general setup of 1st selection round of experimental scheme with minor modifications in PCR cycles, emulsified RT reaction temperature and few more details.

All modifications are given below:

- transcription - 5 µl - 200 ng/µl (~1000ng) of agarose-gel purified PCR fragment were taken; final concentration of mRNA used in 3rd slection round was 1.5 µg/µl;
- translation - 0.8 µl - 1.5 µg/µl (~1.2 µg) of mRNA were taken;
- emulsified RT reaction was performed 1 hr at 55°C;
- 1st PCR (RD_Nde//pD_55) - 25 cycles were performed;
- 2nd (nested) PCR (M_Esp//M_Eri) - 22 cycles were performed;
- final concentration of digested PCR fragment - 70 ng/µl;
- ligation - 200 ng (~$0.9*10^{11}$ molecules) of MLV RT library were taken;
- PCR (on ligation mix) - <$0.6*10^{10}$ molecules of selected library were taken and 15 PCR cycles were performed; concentration of final agarose-gel purified PCR fragment was 100 ng/µl.

4th selection round

Fourth selection round was performed following general setup of 1st selection round of experimental scheme with minor modifications in PCR cycles, emulsified RT reaction temperature and few more details.

All modifications are given bellow:

- transcription - 10 µl - 100 ng/µl (~1000ng) of agarose-gel purified PCR fragment were taken; final concentration of mRNA used in 4th slection round was 1.8 µg/µl;
- translation - 0.67 µl - 1.8 µg/µl (~1.2 µg) of mRNA were taken;
- emulsified RT reaction was performed 1 hr at 57.5°C;
- 1st PCR (RD_Nde//pD_55) - 25 cycles were performed;
- 2nd (nested) PCR (M_Esp//M_Eri) - 24 cycles were performed;
- final concentration of digested PCR fragment - 50 ng/µl;
- ligation - 200 ng (~$0.9*10^{11}$ molecules) of MLV RT library were taken;
- PCR (on ligation mix) - <$0.6*10^{10}$ molecules of selected library were taken and 15 PCR cycles were performed; concentration of final agarose-gel purified PCR fragment was 100 ng/µl.

5th selection round

Fith selection round was performed following general setup of 1st selection round of experimental scheme with some modifications in PCR cycles, emulsified RT reaction temperature and final stage of analysis.

All modifications are given bellow:

- transcription - 10 µl - 100 ng/µl (~1000ng) of agarose-gel purified PCR fragment were taken; final concentration of mRNA used in 5th slection round was 1.1 µg/µl;
- translation - 1.1 µl - 1.1 µg/µl (~1.2 µg) of mRNA were taken;
- emulsified RT reaction was performed 1 hr at 60°C;

- 1st PCR (RD_Nde//pD_55) - 25 cycles were performed;
- 2nd (nested) PCR (M_Esp//M_Eri) - 33 cycles were performed;

Nested PCR mixture (500 µl) for full gene amplification was prepared on ice: 50 µl - 10X Taq buffer with KCl (Fermentas); 50 µl - 2 mM of each dNTP (Fermentas); 30 µl - 25 mM $MgCl_2$ (Fermentas); 6.25 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 2.5 µl - 2.5 u/µl Pfu DNA Polymerase (Fermentas); 2.5 µl - 100 µM M_Esp primer (SEQ ID No: 13); 2.5 µl - 100 µM M_Hind3+ primer (SEQ ID No: 18); 50 µl of first PCR (primers set RD_Nde//pD_55); 306 µl water. The cycling protocol for 2077 bp PCR fragment was: initial denaturation step 3 min at 94°C, 22 cycles (45 sec at 94°C, 45 sec at 55°C, and 3 min at 72°C) and final elongation 5 min at 72°C.

Final PCR fragment of selection sample was agarose-gel purified using Qiagen gel extraction kit (elution in 60 µl ~60 ng/µl). Purified PCR fragment was digested with HindIII and Esp3I for 1 hr at 37°C and again agarose-gel purified (elution in 40 µl ~50 ng/µl).

Recovered MLV reverse transcriptase library after 5th selection round was ligated into plasmid vector prepared from pET_his_MLV_pD (SEQ ID No: 1 and FIG 2) digested with NcoI and HindIII, giving new 7474 bp plasmid pET_his_MLV (SEQ ID No: 19) encoding MLV RT with N-terminal his-tag and no pD fusion on C-terminus for fast protein purification using affinity chromatography.

Ligated MLV RT library after 5th selection round was electroporated into T7 expression strain ER2566. Individual clones were sequenced and analyzed. Pool of evolved proteins as well as individual mutants and primary wt M-MuLV reverse transcriptase in the same construction were grown in 200 ml of LB to A590 ~0.7 and purified via his-tag by affinity chromatography using 2 ml of Qiagen - Ni-NTA Superflow resin (all purification was performed under native conditions according to suppliers recommendations). Elution was performed in 1 ml of EB (50 mM - NaH_2PO_4, 300 mM - NaCl, 250 mM - imidazol, pH 8.0, 10 mM - β-mercaptoethanol and 0.1% of triton X-100). All proteins were dialyzed against 50 times excess of storage buffer (50 mM Tris-HCl (pH 8.3 at 25°C), 0.1 M NaCl, 1 mM EDTA, 5 mM DTT, 0.1% (v/v) Triton

X-100 and 50% (v/v) glycerol). Protein purity was checked on SDS-PAGE (usually ~40-80% of target protein). Protein concentration was determined using Bredford method based Bio-Rad protein assay (500-0006).

MLV reverse transcriptase specific activities were measured at 37°C, 50°C (enzyme diluted in special dilution buffer: 30 mM Tris-HCl pH 8.3 at 25°C, 10 mM DTT, 0.5 mg/ml BSA) and rest of activity at 37°C after 5 min enzyme incubation at 50°C (enzyme diluted and its stability measured in 1x RT reaction buffer: 50 mM Tris-HCl pH 8.3 at 25°C, 4 mM $MgCl_2$, 10 mM DTT, 50 mM KCl). Enzyme activity in all cases was assayed in the following final mixture: 50 mM Tris-HCl (pH 8.3 at 25°C), 6 mM $MgCl_2$, 10 mM DTT, 40 mM KCl, 0.5 mM dTTP, 0.4 MBq/ml [3H]-dTTP, 0.4 mM polyA•oligo$(dT)_{12-18}$. Activity units were determined measuring incorporation of dTMP into a polynucleotide fraction (adsorbed on DE-81) in 10 min at particular reaction temperature and comparing to known amounts of commercial enzyme.

Results

During analysis of selection data we have accumulated 104 sequences expressing full length M-MuLV reverse transcriptase. The total CLUSTALW alignment of all proteins (Appendix 2) is compiled without N terminal His tag in order to have the same numeration of amino acids as is usually used in literature. Wild type sequence denoted as MLV is always given as the first sequence. Mutations are marked using white font color in black background (Appendix 2). Amino acids positions, mutations of which somehow improve M-MuLV reverse transcriptase properties and are described in different patent applications are marked in the alignment as columns of amino acids (white font) highlighted in grey. Mutations originating from our selection and located in grey columns serve as proof that our selection procedure precisely targeted the beneficial hot spot or even exact amino acid mutations described elsewhere. Out of 104 sequenced clones we found 98 unique sequences and 1 wt (L5_87) sequence. Five sequences are repeated twice (L5_21 and L5_111; L5_43 and L5_112; L5_49 and L5_63; L5_64 and L5_93; L5_85 and L5_96). In total we had randomly expressed 55 proteins. Out of 55 expressed proteins 40 enzymaticaly active mutant variants (including control wt) of M-MuLV reverse transcriptase were successfully purified to 40-80% homogeneity according to SDS-PAGE. Total protein concentration in purified RT samples was in range of 0.6-5.5 mg/ml. Mutant RT variants were tested for reverse transcriptase activity at 37°C, 50°C

and residual activity at 37°C after 5 min incubation at 50°C (FIG 8). Reverse transcriptase activity at 37°C is normalized to be 100% and is omitted in FIG 8. Thus only two types of columns (percents of RT activity at 50°C and residual RT activity at 37°C after 5 min incubation at 50°C) are shown. As a control wt M-MuLV reverse transcriptase used for mutants library construction is presented. This primary enzyme was expressed in the same vector and purified in the same way as mutant variants of RT. An average value of wt enzyme RT activity at 50°C is about 45% comparing to activity at 37°C. Almost all tested proteins, with few exceptions, have higher than 45% activity at 50°C. An average value of RT activity at 50°C for all tested mutants is about ~92% and is more than 2 times higher comparing to wt enzyme (45%). Some mutants are 100% or even more active at 50°C as they are at 37°C: 20, 23, L5_16, L5_24, L5_30, L5_35, L5_37, L5_43, L5_46, L5_47, L5_49, L5_52, L5_55, L5_64, L5_65, L5_68, L5_72. The best mutants found have RT activity at 50°C about 140% and more (3 times higher than 45% of wt): 20 (165%), L5_37 (162%), L5_43 (156%), L5_46 (135%), L5_47 (179%), L5_52 (137%), L5_64 (142%) and L5_68 (153%).

Even though majority of mutants have very high RT activities at 50°C, they are not as thermostable. Residual RT activity at 37°C after 5 min incubation at 50°C of wt control is ~11%. The same average residual activity of selected enzymes is also similar ~12%. Although some tested RT variants are substantially more thermostable and have residual activity 2-3 times higher than wt enzyme (11%): L5_8 (25%), L5_43 (32%), L5_46 (27%), L5_64 (28%), L5_65 (25%), L5_68 (31%).

Specific activity (u/mg of protein) for partially purified wt enzyme is ~200'000 u/mg (FIG 9). Selected RT variants were expressed and purified in very diverse manner and an average specific activity (~155'000 u/mg) is slightly lower as for wt control (FIG 9). In some cases specific activity is decreased, in some - increased (20 - ~274'000 u/mg; L5_11 - ~273'000 u/mg; L5_28 - ~230'000 u/mg; L5_30 - ~224'000 u/mg; L5_35 - ~316'000 u/mg; L5_43 - ~328'000 u/mg; L5_46 - ~304'000 u/mg; L5_52 - ~310'000 u/mg; L5_64 - ~256'000 u/mg; L5_65 - ~247'000 u/mg).

It is obvious that our selection system works well. Using increased temperature of RT reaction as a selection pressure factor, we have managed to evolve faster (specific RT activity at 50°C

is higher) and more thermostable (residual RT activity at 37°C after 5 min preincubation at 50°C) reverse transcriptases.

The source of valuable information is an alignment of selected protein sequences (Appendix 2). Sequences of analyzed proteins, whose activity at 50°C was substantially better as compared to primary wt M-MuLV (70% and more comparing to 45% of wt activity) are highlighted in grey (Appendix 2). Number of mutagenized amino acids varied in range of 0 (wt or L5_87) to 12 (L5_9). List of mutations found in all selected RT variants is given in Appendix 3. Proteins are sorted by decreasing number of mutations. Most mutants (53 out of 104) had 4-6 mutations per sequence. It is obvious that reverse transcriptase sequence has some hot spots, which are very important and beneficial for the RT reaction in general and for the thermostability of enzyme. Those hot spots can be easily identified in multiple sequence alignment (Appendix 2) as conglomeration of mutations at particular position. Especially important are mutations found in better performing variants of M-MuLV reverse transcriptase (sequences are highlighted in grey – Appendix 2). Summarized information about the most frequent mutations (in decreasing order) is given in Appendix 4. As in previous appendix mutant proteins with substantially higher activity at 50°C are highlighted in grey. If same mutation repeats for many times and tested reverse transcriptases with this mutation are better performing at 50°C, that means this mutation is somehow beneficial for reverse transcription reaction.

According to the frequency with which mutations are found they can be divided into five classes: 21-31 repeats; 14-18 repeats; 4-7 repeats; 2-3 repeats and 1 repeat. The first group of the most frequent mutations comprises four amino acids D524 (31 repeats); D200 (30 repeats); D653 (23 repeats) and D583 (21 repeats). Two amino acids (D524 and D583) are known to complex magnesium ions in active centre of ribonuclease H domain. Mutants D524G, D583N and E562Q are used to turn off RNase H activity of M-MuLV reverse transcriptase (Gerard et al., 2002), what improves synthesis of cDNA. Results of our selection are strikingly similar. Mutation of aspartate 524 is found in 31 sequences out of 98. Moreover, D524N substitution is found once, D524A - 10 times and finally D524G - 20 times. Thus our selection not only precisely targeted important amino acids, but also the same amino acid substitutions, which are known to be the best. Exactly the same situation is with mutation of aspartate 583, which is

repeated in 21 selected proteins out of 104. Substitution D583E is found once, D583A - 3 times, D583G - 7 times and finally D583N - 10 times. Again, same amino acid and same substitution (D583N), which is known to be the best is selected most frequently. Commercial enzyme SUPERSCRIPT II from Invitrogen has three mutations: D524G, D583N and E562Q (WO2004024749). An interesting thing is that mutation of the third amino acid substitution E562 in our selection is found only once (E562K in L5_71). This result suggests that most likely glutamate 562 is not as important as aspartates 524 and 583, or for some reasons exchange of this amino acid can cause some side effects and is not beneficial for RT reactions performed at increased temperatures (>50°C).

Further analysis of selected proteins sequences allowed to identify many more hot amino acids positions, mutations of which are described in other patent applications for improved M-MuLV reverse transcriptase: H204R - 7 repeats (US7078208); H638R - 4 repeats (US20050232934A1); T197A - 2 repeats (US7056716); M289V(L), T306A(M) - 2 repeats (US7078208); E302K, N454K - 2 repeats (WO07022045A2); E69G, L435P - 1 sequence (WO07022045A2); Y64C, Q190R, V223M, F309S - 1 sequence (US7056716); E562K - 1 sequence (US7078208). There are also two selected sequences of reverse transcriptases which have combination of three amino acids substitutions described in literature (30 - D200N, T306M, D524N, D583G; L5_28 - T306A, F309S, D524A, H594R, F625S).

In addition to known mutations we have identified many more amino acids positions which are mutated quite often: D200N(A, G) - 30 repeats; D653N(G, A, H, V) - 23 repeats; L603W(M) - 18 repeats; T330P - 15 repeats; L139P - 14 repeats; Q221R - 6 repeats; T287A - 6 repeats; I49V(T) - 5 repeats; N479D - 5 repeats; H594R(Q) - 5 repeats; F625S(L) - 5 repeats; P65S - 4 repeats; H126S(R) - 4 repeats; L333Q(P) - 4 repeats; A502V - 4 repeats; E607K(G, A) - 4 repeats; K658R(Q) - 4 repeats; H8P(R) - 3 repeats; P130S - 3 repeats; E233K - 3 repeats; Q237R - 3 repeats; N249D - 3 repeats; A283D(T) - 3 repeats; A307V - 3 repeats; Y344H - 3 repeats; P407S(L) - 3 repeats; M428L - 3 repeats; Q430R - 3 repeats; D449G(A) - 3 repeats; A644V(T) - 3 repeats; N649S - 3 repeats; L671P - 3 repeats; E673G(K) - 3 repeats; N678I - 3 repeats (Appendix 4).

Best performing variants of RT usually have mutations of amino acids, which are modified most frequently:

20 (50°C - 123%) - D200N (30 repeats), L603W (18 repeats) and slightly modified C terminus - N678I, S679P, R680A;

L5_35 (50°C - 125%) - D200N (30 repeats), T330P (15 repeats), N479D (5 repeats);

L5_37 (50°C - 162%) - H123S (4 repeats), L149F (1 sequence), D200N (30 repeats), N454K (2 repeats), D583N (21 repeats);

L5_43 (50°C - 160%) - D200N (30 repeats), Q237R (3 repeats), T330P (15 repeats), D524G (31 repeats), F625S (5 repeats), D653N (23 repeats);

L5_46 (50°C - 135%) - D200N (30 repeats), T330P (15 repeats), D583N (21 repeats), T644T (3 repeats);

L5_47 (50°C - 179%) - N107S (1 repeat), H126R (4 repeats), T128A (1 repeat), I179V (2 repeats), D200N (30 repeats), H642Y (2 repeats), D653N (23 repeats);

L5_52 (50°C - 137%) - D200N (30 repeats), T330P (15 repeats), Q374R (2 repeats), (D583N (21 repeats);

L5_64 (50°C - 142%) - D200N (30 repeats), D216G (2 repeats), D524G (31 repeats), E545G (2 repeats);

L5_65 (50°C - 127%) - D200N (30 repeats), Q238H (1 repeat), L570I (1 repeat), L603W (18 repeats);

L5_68 (50°C - 153%) - M39V (2 repeats), I49V (2 repeats), Q91R (2 repeats), H204R (7 repeats), T287A (6 repeats), N454K (2 repeats), F625L (5 repeats), D653H (23 repeats).

The combined data set of measured RT activities and sequence alignment analysis of mutant proteins allowed us to determine many beneficial mutations (and combinations thereof) in the sequence of M-MuLV reverse transcriptase sequence.

Example 3 - *Analysis of Moloney Murine Leukemia Virus reverse transcriptase mutants*

In vitro evolution experiment described in 2nd example was very efficient. Gradually increased temperature of reverse transcription reaction was used as a selection pressure and generated a lot of different mutant variants of M-MuLV RT. Most of them were able to perform better at elevated temperatures comparing to primary enzyme. Sequence analysis of

evolved reverse transcriptases indicates hot spots and most important amino acids positions (replacements), responsible for complex improvement of enzyme properties. In order to elucidate individual impact of different mutations single and multiple mutants of M-MuLV RT were constructed, partially purified and analyzed. Starting point for mutants construction was 7474 bp plasmid pET_his_MLV (SEQ ID No: 19) encoding M-MuLV RT with N-terminal his-tag for fast protein purification using affinity chromatography. M-MuLV reverse transcriptase specific activity at 37°C, relative activity at 50°C and relative residual activity at 37°C after 5 min enzyme incubation at 50°C were determined. In some cases RNase H activity was checked and cDNA synthesis reaction at different temperatures on 1kb or 4.5kb RNA was performed.

Methods and materials

Initial plasmid pET_his_MLV (SEQ ID No: 19) was used as a starting material for mutagenic PCR. Mutations were introduced using mutagenic primers. Individual clones were sequenced and analyzed. M-MuLV RT mutants were expressed in T7 expression strain ER2566. Individual proteins and primary wt M-MuLV reverse transcriptase in the same construction were grown in 200 ml of LB to A590 ~0.7 and purified via his-tag by affinity chromatography using 2 ml of Qiagen - Ni-NTA Superflow resin (all purification was performed under native conditions according to suppliers recommendations). Elution was performed in 1 ml of EB (50 mM - NaH_2PO_4, 300 mM - NaCl, 250 mM - imidazol, pH 8.0, 10 mM - β-mercaptoethanol and 0.1% of triton X-100). All proteins were dialyzed against 50 times excess of storage buffer (50 mM Tris-HCl (pH 8.3 at 25°C), 0.1 M NaCl, 1 mM EDTA, 5 mM DTT, 0.1% (v/v) Triton X-100 and 50% (v/v) glycerol). Protein purity was checked on SDS-PAGE (usually ~40-80% of target protein). Protein concentration was determined using Bradford reagent (Fermentas #R1271).

MLV reverse transcriptase activities were measured at 37°C, 50°C (enzyme diluted in special dilution buffer: 30 mM Tris-HCl pH 8.3 at 25°C, 10 mM DTT, 0.5 mg/ml BSA) and rest of activity at 37°C after 5 min enzyme incubation at 50°C (enzyme diluted and its stability measured in 1x RT reaction buffer: 50 mM Tris-HCl pH 8.3 at 25°C, 4 mM $MgCl_2$, 10 mM DTT, 50 mM KCl). Enzyme activity in all cases was assayed in the following final mixture: 50 mM Tris-HCl (pH 8.3 at 25°C), 6 mM $MgCl_2$, 10 mM DTT, 40 mM KCl, 0.5 mM dTTP, 0.4 MBq/ml [3H]-dTTP, 0.4 mM polyA•oligo$(dT)_{12-18}$. Activity units were determined measuring

incorporation of dTMP into a polynucleotide fraction (adsorbed on DE-81) in 10 min at particular reaction temperature and comparing to known amounts of commercial enzyme.

RNase H activity of M-MuLV reverse transcriptase variants was measured according to US patent US5405776. RNase H activity of purified enzymes was assayed in reaction mixtures (50 µl) containing 50mM Tris-HCl pH 8.3, 2 mM $MnCl_2$, 1 mM DTT and [3H](A)n*(dT)n (5 µM [3H](A)n, 35 cpm/pmol; 20 µM (dT)n). Reactions were incubated at 37°C for 10 min and were stopped by adding 10 µl of tRNA (1 mg/ml) and 20 µl of cold 50% TCA. After 10 minutes on ice, the mixture was centrifuged for 10 minutes in an Eppendorf centrifuge (at 25000g). Forty µl of the supernatant was counted in a LSC-universal cocktail (Roth - Rotiszint eco plus). One unit of RNase H activity is the amount of enzyme required to solubilize one mole of [3H](A)n in [3H](A)n*(dT)n in 10 min at 37°C.

"RevertAid^{TM} First Strand cDNA Synthesis Kit" (#K1622 - Fermentas) and its control 1.1 kb RNA with a 3'-poly(A) tail in combination with oligo$(dT)_{18}$ primer was used to check purified reverse transcriptases for their ability to synthesize cDNA at different temperatures. Alternatively 4.5 kb RNA (synthesized from Eco31I linearized pTZ19R plasmid, which additionally contains piece of phage lambda DNA 5505-8469bp) was used to test reverse transcription reaction. cDNA was synthesized 1 hr in 20 µl reaction volume using kit's components 1 µg of synthetic RNA, following provided protocol with only some minor modifications (without 5min preincubation at 37°C). Reverse transcription reactions were performed in 96 well PCR plate in Eppendorf Mastercycler gradient PCR machine applying corresponding temperature gradient. Synthesized cDNA was analyzed by alkaline agarose gel electrophoresis (staining with ethidium bromide). Samples of cDNA synthesis analysis on alkaline agarose gels are given in FIG 16.

Results

According to the frequency, with which mutations are found during the evolution of M-MuLV reverse transcriptase, they can be divided into five classes: 21-31 repeats; 14-18 repeats; 4-7 repeats; 2-3 repeats and 1 repeat. Construction of individual reverse transcriptase mutants in general was performed according to this information. Most frequently found mutations were tested first. Reverse transcriptase specific activity at 37°C, relative activity at 50°C and relative residual activity at 37°C after 5 min enzyme incubation at 50°C were determined. In some

cases RNase H activity was checked and cDNA synthesis reaction at different temperatures on 1kb or 4.5kb RNA was performed. All experimental data on individual mutants are presented in Appendix 5. Second column ("Selection frequency") indicates the number of sequenced mutants, which had exact mutation and the number in the parentheses indicates total number of particular amino acid mutations found in selection. For example D200N - 25 (30) means, that aspartate 200 replacement to asparagine was found 25 times out of 30 total D200 mutations. Reverse transcriptase specific activity measured at 37°C is given in units per mg of protein. Relative enzyme activity at 50°C and relative residual RT activity at 37°C after 5 min incubation at 50°C are given in percents normalized to specific activity (100%) of the same enzyme measured at 37°C. As a control (first line) is given wt M-MuLV reverse transcriptase used for mutants library construction. This primary enzyme is expressed in the same vector and purified in the same way as mutant variants of RT. Specific activity of wt enzyme is about 200'000 u/mg at 37°C, relative activity at 50°C (comparing to activity at 37°C) - 45-50% (90'000-100'000 u/mg) and relative residual RT activity at 37°C after 5 min incubation at 50°C (comparing to activity at 37°C) is about 11% (~22'000 u/mg). Wild type enzyme has about 160-200 u/mol of RNase H activity and can synthesize full length 1kb cDNA at 48°C. It is known, that M-MuLV reverse transcriptase is protected from thermal inactivation by the binding to template-primer substrate and contrary, enzyme is less thermostable in solution alone (Gerard et al., 2002). Relative residual RT activity at 37°C after 5 min incubation at 50°C directly indicates enzyme thermostability in solution without substrate. Meanwhile relative activity at 50°C represents enzyme thermostability in complex with RNA/DNA substrate and speed of cDNA synthesis. Fast mutant variant of reverse transcriptase will give increased numbers of polymerase units at 50°C, even if it's thermostability will be the same as wild type enzyme. Highest temperature of cDNA synthesis (in our case 1 kb or 4,5 kb) is the most comprehensive parameter, which represents general ability of enzyme to synthesize cDNA at increased temperatures. Reverse transcriptase mutants, which have at least 10% increased: specific activity at 37° (≥220'000 u/mg, 200'000 u/mg - wt), relative activity at 50°C comparing to mutant activity at 37°C (≥54%, 45-50% - wt) or relative residual activity at 37°C after 5 min incubation at 50°C comparing to mutant activity at 37°C (≥13%, 11% - wt),- are shadowed in gray and considered as significantly improved enzymes. Mutants able to synthesize full length 1kb cDNA at temperatures higher than 48°C are also shadowed in gray.

Reverse transcriptase mutants with increased specific activity (≥220'000 u/mg) at 37°C are (Appendix 5):

D200 (D200N - 254'000 u/mg; D200G - 276'000 u/mg; D200H - 234'000 u/mg),

T330 (T330N - 223'000 u/mg; T330D - 240'000 u/mg),

Q221 (Q221R - 268'000 u/mg),

H594 (H594K - 270'000 u/mg; H594Q -231'000 u/mg),

D449 (D449E - 224'000 u/mg; D449N - 221'000 u/mg),

M39 (M39N - 349'000 u/mg),

M66 (M66L - 237'000 u/mg; M66V - 227'000 u/mg; M66I - 240'000 u/mg),

H126 (H126R - 227'000 u/mg),

W388 (W388R - 266'000 u/mg),

I179 (I179V - 251'000 u/mg).

Reverse transcriptase mutants with increased relative activity (≥54%) at 50°C (comparing to activity at 37°C) are (Appendix 5):

D200 (D200N - 84%; D200A - 87%; D200Q - 103%; D200E - 79%; D200V - 131%; D200W - 103%; D200G - 88%; D200K - 102%; D200R - 68%; D200H - 54%),

L603 (L603W - 105%; L603F - 104%; L603Y - 95%; L603M - 77%),

D653 (D653N - 93%; D653K - 106%; D653A - 99%; D653V - 98%; D653Q - 93%; D653L - 83%; D653H - 116%; D653G - 90%; D653W - 93%; D653E - 80%),

T330 (T330P - 80%; T330N - 69%; T330D - 55%; T330V - 65%; T330S - 67%),

Q221 (Q221R - 94%; Q221K - 77%; Q221E - 64%; Q221M - 58%; Q221Y - 77%),

E607 (E607K - 84%; E607A - 98%; E607G - 72%; E607D - 69%),

L139 (L139P - 59%),

T287 (T287S - 68%),

N479 (N479D - 81%),

H594 (H594R - 69%; H594K - 80%; H594Q - 75%; H594N - 61%),

D449 (D449G - 79%; D449E - 77%; D449N - 75%; D449A - 99%; D449V - 83%),

M39 (M39V - 54%; M39N - 71%),

M66 (M66L - 79%; M66V - 73%; M66I - 80%),

L333 (L333Q - 54%),

H126 (H126R - 58%),
P130 (P130S - 70%),
Q91 (Q91R - 56%),
W388 (W388R - 72%),
R390 (R390W - 64%),
Q374 (Q374R - 56%),
E5 (E5K - 67%).

Reverse transcriptase mutants with increased relative residual activity (≥13%) at 37°C after 5 min incubation at 50°C (comparing to activity at 37°C) are (Appendix 5):
D200 (D200N - 15%; D200A - 18%; D200Q - 23%; D200R - 27%; D200H - 27%),
L603 (L603W - 23%; L603Y - 13%; L603P - 15%),
D653 (D653N - 21%; D653K - 15%; D653A - 18%; D653V - 16%; D653Q - 18%; D653H - 13%; D653G - 13%; D653W - 13%; D653E - 19%),
T330 (T330P - 21%; T330N - 13%; T330D - 16%; T330S - 15%),
T287 (T287A - 13%; T287F - 13%),
H594 (H594R - 14%; H594Q - 13%),
D449 (D449G - 13%),
M39 (M39V- 13%),
M66 (M66L - 13%),
Y344 (Y344H - 13%),
Q91 (Q91R - 13%),
N649 (N649S - 16%),
W388 (W388R - 14%).

Mutants able to synthesize full length 1kb cDNA at temperatures higher than 48°C are (Appendix 5):
D200 (D200N - 50.4°C; D200H - 50.4°C),
L603 (L603W - 53.1°C; L603F - 50.4°C; L603Y - 47.8-50.4°C),
D653 (D653N - 50.4-53.1°C; D653K - 50.4-53.1°C; D653A - 50.4°C; D653V - 50.4°C; D653Q - 50.4°C; D653L - 50.4°C; D653H - 50.4-53.1°C; D653G - 50.4°C; D653W - 50.4°C),
Q221 (Q221R - 50.4°C),

E607 (E607K - 47.8-50.4°C),

H594 (H594K- 47.8-50.4°C; H594Q - 47.8-50.4°C).

Samples of 1 kb cDNA synthesis analysis on alkaline agarose gels are given in FIG 16 A-D.

According to collected biochemical data most important positions in M-MuLV reverse transcriptase sequence, which can impact cDNA synthesis at increased temperatures, are: **D200, L603, D653, T330, Q221, E607, L139, T287, N479, H594, D449, M39, M66, L333, H126, Y344, P130, Q91, N649, W388, R390, I179, Q374, E5.**

In general mutations of interest can be combined in between and M-MuLV reverse transcriptase thermostability with and without substrate, velocity, processivity and overall ability to synthesize cDNA at increased temperatures can be further improved. Some data, which illustrates enzyme improvement by combinatorial approach are presented in Appendix 6. Single mutants D200N and L603W have relative activity at 50°C 84% and 105%. Highest temperatures of 1kb cDNA synthesis are 50.4°C and 53.1°C. Double mutant D200N; L603W has relative activity 131% at 50°C and can synthesize 1kb cDNA at 56°C. Triple mutant D200N; L603W; T330P (80% at 50°C; 1kb cDNA at 47.8°C) is improved further and has relative activity 175% at 50°C and can synthesize 1kb cDNA at 56-58°C. Quadruple mutant D200N; L603W; T330P; E607K (84% at 50°C; 1kb cDNA at 47.8-50.1°C) has relative activity 174% at 50°C and can synthesize 1kb cDNA at 60-62°C. Quintuple mutant D200N; L603W; T330P; E607K; L139P (59% at 50°C; 1kb cDNA at 47.8°C) has relative activity 176% at 50°C and can synthesize 1kb cDNA at 62°C and that is about 14°C higher temperature comparing to wild type M-MuLV reverse transcriptase (1kb cDNA at 47.8°C). Additive character of thermostability is also observed in case of:

N479D, H594R mutants (D200N; L603W - 131% at 50°C, 1kb cDNA at 56°C versus D200N; L603W; N479D; H594R - 182% at 50°C, 1kb cDNA at 56-58°C, 4.5kb cDNA at 56-58°C),

T330P mutant (D200N; L603W; D653N; D524G - 155% at 50°C, 1kb cDNA at 58-60°C versus D200N; L603W; D653N; D524G; T330P - 180% at 50°C, 1kb cDNA at 60-62°C).

Samples of 4.5 kb cDNA synthesis analysis on alkaline agarose gels are given in FIG 16 E-G.

Example 4 - *Modification of CRD – selection for DNA dependent DNA polymerase activity using biotin-dUTP (proof of principle).*

This example illustrates activity-based selection strategy of reverse transcriptase as DNA dependent DNA polymerase, which is able to incorporate modified nucleotides into DNA-DNA substrate. The principle scheme of selection is schematically illustrated in FIG 12. Two plasmids pET_his_MLV_D583N_pD (encoding RNase H minus Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase fused to protein D spacer) and its derivative pET_his_del_pD (encoding inactivated reverse transcriptase; 57 amino acids deletion in pol domain and mutation D583N in RNase H domain, Example 1, SEQ ID No: 2) were used as a starting material in this example. Initial DNA fragments encoding active and inactive reverse transcriptases were synthesized in two separate polymerase chain reactions using plasmids pET_his_MLV_D583N_pD and pET_his_del_pD as a target. Synthesized PCR fragments were used in transcription reaction for synthesis of mRNA, which lacks STOP codon at the 3' end. Purified mRNAs were mixed to a ratio of 1:20=MLV (active RT):del (inactive RT). Double stranded DNA adapter (required for selection of DNA dependent DNA polymerase activity) was ligated to 3' mRNA by T4 DNA ligase. This mRNA-dsDNA complex is used for *in vitro* translation reaction. The ribosome moving along the mRNA stops translation at the site of RNA–DNA hybridization (Tabuchi et al., 2001). Ribosome-mRNA/dsDNA-protein complexes were stabilized (as in conventional ribosome display) by dilution of translation mixture with ice-cold buffer containing 50 mM Mg^{2+}. Mixture of ternary complexes (TC) was purified by ultracentrifugation on sucrose cushions. Purified ternary complexes ($<3*10^9$ molecules taken) containing mRNA-dsDNA linked to *in vitro* translated M-MuLV (RNase H-) reverse transcriptases were used to prepare reaction mixture additionally supplemented with biotin-dUTP and reaction buffer. Ice-cold reaction mixture was emulsified yielding $\sim 1*10^{10}$ water in oil compartments ~2μm in size. Emulsified reaction mixture (less than one TC, ribosome-mRNA/dsDNA-protein per compartment) was incubated for 30 min at 37°C. After the temperature of compartmentalized reaction mixture is raised most of TCs dissociate releasing mRNA/dsDNA and reverse transcriptase. Successful incorporation reaction can occur only in compartments containing active M-MuLV (RNase H-) reverse transcriptase resulting in biotinylation of mRNA/dsDNA complex. Biotinylated complex can be selectively immobilized

on streptavidin coated magnetic beads and specifically amplified by RT-PCR. As a result of successful experiment genes encoding active enzyme (in our case RT-PCR fragment of MLV_D583N_pD reverse transcriptase) should be enriched over genes encoding inactive enzyme (del_pD).

Methods and materials

Production of mRNA/dsDNA complex.

(1) Determination of ligation efficiency. Efficiency of ligation reaction was determined by ligation of MLV_pD mRNA (synthesized from pET_his_MLV_pD plasmid, Example 1) with primer Long+Tb (SEQ ID No: 23) using ddC-Long2 primer (SEQ ID No: 22) as a splint. The 5' end of Long+Tb was previously phosphorylated using T4 Polynucleotide Kinase (Fermentas).

Annealing mix of 36µl was prepared by mixing of 8.5 pmol (~10µg) of purified MLV_pD mRNA with 4 times of molar excess of Long+Tb (SEQ ID No: 23) and 4.2 times of molar excess of ddC-Long2 (SEQ ID No: 22) in nuclease free water. The mixture was incubated at 70°C for 5 min and then cooled to room temperature for 20 min. Before adding ligation reaction components mixture was moved to cooling stand for 2 min.

4.5 µl of 10x ligation buffer and 4.5µl of T4 DNA ligase (5v/µl) (Fermentas) were added to 36µl of annealing mixture. Ligation reaction was performed for 30min at 37°C and followed extraction once with equal volume of Roti® Phenol/chloroform (ROTH) and twice with equal volume of chloroform. Assuming that mRNA amount was close to initial amount taken for ligation reaction ~5µg of ligation products mix was diluted to 43µl by nuclease free water. Aliquot of 2µl of resultant mixture was left for analysis on agarose gel before immobilization on Dynabeads M-280 Streptavidin beads (Dynabeads® kilobase BINDER™ Kit (DYNAL Biotech)).

10µl of resuspended Dynabeads was transferred to a 1.5 ml microcentrifuge tube, washed with 50 µl of Binding Solution provided in the kit. The tube was placed on the magnet for 1-2 min until beads settled at the tube and solution was removed. Dynabeads were gently resuspended by pipetting in 38 µl of Binding Solution supplemented with 2µl of aqueous tRNA (tRNA from yeast (Roche)) solution (1µg/µl) to minimize non-specific mRNA binding. 40µl of Dynabeads

in Binding solution was added to solution (~40µl) containing ligation products mix. The tube was incubated shaking at the +22°C in termomixer (Eppendorf) for 60 min. After the binding of ligated mRNA/dsDNA, supernatant was removed and the beads were washed three times with 50 µl of Washing Solution (supplied in the kit). Collected Dynabeads with immobilizes ligated mRNA/dsDNA complex was resuspended in 26µl of nuclease free water and followed extraction once with 40 µl of Roti® Phenol/chloroform (ROTH) and twice with 40µl volume of chloroform to release mRNA/dsDNA complex from magnetic beads. 1; 2 and 5µl of final mix were analysed on agarose gel along with sample (2 µl) left before immobilization and Mass Ruler™High Range DNA ladder. Amount of mRNA in mRNA/DNA complex purified on streptavidin beads was determined comparing mRNA amounts before and after the immobilization. The yield of recovered mRNA was ~60%, what means that at least 60% of mRNA was successfully ligated with DNA duplex, resulting in mRNA/dsDNA complex.

(2) Determination of biotin-dUTP incorporation efficiencies into mRNA/dsDNA complex and into self primed mRNA. As it was shown in first mRNA to dsDNA ligation experiment ligation reaction efficiency is ~60% and more. Free mRNA left in ligation mixture can self prime and participate in extension reaction using M-MuLV reverse transcriptase and biotin-dUTP. This experiment was performed in order to demonstrate that mRNA/dsDNA complex (ligation product) is better substrate than free self primed mRNA.

MLV_D583N_pD mRNA was ligated with long+ oligonucleotide (SEQ ID No: 21) according procedure described above (construction of initial plasmids pET_his_MLV_D583N_pD and pET_his_del_pD in details is described in Example 1). ~12,5 ng of prepared (MLV_D583N_pD mRNA/long+) substrate was combined with ~12,5 ng of del_pD mRNA previously incubated at 70°C for 5 min, then cooled to room temperature for 20 min in nuclease free water, in total volume of 12.5 µl. The second mix for dTTP or biotin-dUTP incorporation by reverse transcriptase was prepared: 8 µl of 5x Reaction buffer for reverse transcriptase; 1 µl - 40 u/µl RiboLock™ RNase inhibitor (Fermentas); 18.6 µl nuclease-free water; 0.4 µl - 200 u/µl RevertAid™ Minus M-MuLV Reverse transcriptase (Fermentas). Prepared mix was divided to two tubes 2x 15 µl and 1 µl of 1 mM dTTP (Fermentas) or 1 µl of 1 mM biotin-dUTP (Fermentas) was added. Subsequentially 5 µl of substrate (from the first mix) was added to dTTP and biotin-dUTP containing mixtures. Reactions were carried out at

37°C for 60 min. After that 1 µl of 0.5M EDTA (pH 8.0) was added to both samples and reaction mixtures were extracted once with equal volume of Roti® Phenol/chloroform (ROTH) and once with equal volume of chloroform followed by purification on G-50 MicroColumns (GE Healthcare). 2µl of resultant reaction products were left for direct RT reaction (without streptavidin beads purification) and the remaining part of solutions were used for biotinylated mRNA-dsDNA complex immobilization on Dynabeads M-280 Streptavidin beads (Dynabeads® kilobase BINDER™ Kit (DYNAL Biotech)). 10 µl of resuspended Dynabeads was transferred to a 1.5 ml microcentrifuge tube, washed with 25 µl of provided Binding Solution. The tube was placed on the magnet for 1-2 min until beads settled at the bottom of the tube and solution was removed. Dynabeads were gently resuspended by pipetting in 90 µl of Binding Solution and 2 µl of aqueous tRNA (tRNA from yeast (Roche)) solution (1µg/µl) was added to minimize non-specific mRNA binding. 40 µl of Dynabeads in Binding solution was added to solution (~40µl) containing the elongated by dTTP and biotin-dUTP RNA-DNA fragments. The tubes were incubated at the +22°C in termomixer (Eppendorf) for 40 min by shaking (1400 rpm). After the binding step, supernatant was removed and the beads were washed three times with 50 µl of Washing Solution (provided in the kit) shaking (1400 rpm) for 5 min at +22°C. Collected Dynabeads with immobilizes elongated mRNA-dsDNA complex was resuspended in reverse transcription reaction mixture. Reverse transcription reaction mixture was prepared on ice: 20 µl - 5X reaction buffer for Reverse Transcriptase (Fermentas); 10 µl – 10 mM dNTP (Fermentas); 2.5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 5 µl - 20 µM pD_42 oligonucleotide (SEQ ID No: 8); 2,5 µl RevertAid™ Minus M-MuLV Reverse transcriptase (200 u/µl)(Fermentas); 55 µl nuclease-free water. Prepared mix was divided to five 19 µl (5x19 µl) aliquots: two were used to resuspend Dynabeads with immobilized elongated mRNA/dsDNA complex or mRNA, the other two were transferred to the tubes with samples left without streptavidin beads purification and to the rest of 19 µl aliquot was added 1µl of nuclease free water – negative reaction control to prove, that reaction mixture is not contaminated by DNA. All reaction mixtures were incubated by shaking (1000 rpm) at the +42°C in termomixer (Eppendorf) for 1 hour until cDNA synthesis reaction was finished.

Amplification of cDNA was performed by nested PCR. Initial PCR mixture was prepared on ice: 14 µl - 10X *Taq* buffer with KCl (Fermentas); 14 µl - 2 mM of each dNTP (Fermentas);

8,4 µl - 25 mM $MgCl_2$ (Fermentas); 2.8 µl - 1 u/µl LC (recombinant) *Taq* DNA Polymerase (Fermentas); 0.7 µl - 100 µM M_F primer (SEQ ID No: 11); 0.7 µl - 100 µM M_2R primer (SEQ ID No: 12); 92.4 µl water - mixture was divided into 7 samples for 19 µl (7x19 µl). To 5x 19 µl of PCR master mix were added 1 µl of cDNA (1-5 RT samples); to the 6^{th} tube of PCR master mix 1 µl of water was added - negative PCR control; to the 7^{th} tube (positive PCR control) - 1 µl of pET_his_MLV_D583N_pD plasmid (~1 ng) was added. The cycling protocol was: initial denaturation step 3 min at 94°C, 25 cycles (45 sec at 94°C, 45 sec at 57°C, and 1 min at 72°C) and final elongation 3 min at 72°C.

Predicted amplicons size 907 bp for MLV_D583N_pD and 736 bp for del_pD cDNA. PCR products were analyzed on 1% agarose gel loading 10 µl of PCR mix per well (FIG 13).

As it was expected efficient cDNA amplification is observed only in elongation reaction with biotin-dUTP. Very faint bands of amplified cDNA can be detected in elongation reaction with dTTP and can be explained by weak nonspecific binding of mRNA to streptavidin beads. After purification on streptavidin beads DNA encoding MLV_D583N_pD gene (907 bp) is enriched over DNA of del_pD (736 bp). That means mRNA/dsDNA complex is elongated by biotin-dUTP much more efficiently than self primed del_pD mRNA.

(3) Preparation of mRNA mixture (MLV_D583N_pD:del_pD=1:20) and RNA/dsDNA complex.

Preparation of PCR fragments for in vitro transcription. PCR mixture was prepared on ice: 20 µl - 10X Taq buffer with KCl (Fermentas); 20 µl - 2 mM of each dNTP (Fermentas); 12 µl - 25 mM $MgCl_2$ (Fermentas); 16 µl - DMSO (D8418 - Sigma); 4 µl - 1 u/µl LC (recombinant) Taq DNA Polymerase (Fermentas); 1 µl - 100 µM pro-pIVEX primer (SEQ ID No: 3); 1 µl - 100 µM pD-ter- primer (SEQ ID No: 20); 122 µl water - mixture divided into two tubes 2x 98 µl. To 2x 98 µl of PCR master mix were added either 2 µl of pET_his_MLV_D583N_pD (diluted to ~1 ng/µl) or 2 µl of pET_his_del_pD (diluted to ~1 ng/µl) (construction of initial plasmids pET_his_MLV_D583N_pD and pET_his_del_pD in details is described in Example 1). The cycling protocol was: initial denaturation step 3 min at 94°C, 30 cycles (45 sec at 94°C, 45 sec at 53°C, and 2 min at 72°C) and final elongation 5 min at 72°C. Amplification efficiency was ~7000 fold from 2 ng of plasmid (7873 bp) target to ~5 µg (50 ng/µl) of amplified product

(2702 bp PCR fragment MLV_D583N_pD from pET_his_MLV_D583N_pD; 2531 bp PCR fragment del_pD from pET_his_del_pD).

Transcription mixture was prepared: 80 µl - 5x T7 transcription buffer (1 M HEPES-KOH pH 7.6; 150 mM Mg acetate; 10 mM spermidin; 0.2 M DTT); 56 µl - 112 mM of each NTP (Fermentas); 16 µl - 20 u/µl T7 RNA polymerase (Fermentas); 8 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 114 µl nuclease-free water - mixture divided into two tubes 2x 165 µl and add 35 µl - 20 ng/µl of PCR fragment MLV_D583N_pD (unpurified PCR mixture) or 35 µl - 20 ng/µl of PCR fragment del_pD PCR (unpurified PCR mixture). Transcription was performed 2 hr at 37°C.

Both transcription mixtures were diluted to 200 µl with ice-cold nuclease-free water and 200 µl of 6 M LiCl solution were added. Mixtures incubated 25 min at +4°C and centrifuged for 25 min at +4°C in cooling centrifuge at max speed (25'000 g). Supernatant was discarded and RNA pellet washed with 500 µl of ice-cold 75% ethanol. Tubes were centrifuged again for 5 min at +4°C at max speed and supernatant was discarded. RNA pellet was dried for 5 min at room temperature and subsequently resuspended in 400 µl of nuclease-free ice-cold water by shaking for 15 min at +4°C and 1400 rpm. Tubes were centrifuged again for 5 min at +4°C at max speed to separate not dissolved RNA. About 380 µl of supernatant were transferred to new tube with 42 µl of 10x DNase I buffer (Mg^{2+}) (Fermentas); 3 µl - 1 u/µl DNaseI (RNase-free) (Fermentas) and incubated for 20 min at +37°C to degrade DNA. Reaction mixtures were extracted once with equal volume of Roti® Phenol/chloroform (ROTH) and twice with equal volume of chloroform to remove DNaseI. 43 µl of 3 M Sodium acetate pH 5.0 solution and 1075 µl of ice-cold 96% ethanol were added to each tube. Finally RNA was precipitated by incubation for 30 min at -20°C and centrifugation for 25 min at +4°C at max speed (25'000 g). Supernatant was discarded and RNA pellet washed with 500 µl of ice-cold 75% ethanol. Tubes were centrifuged again for 4 min at +4°C at max speed and supernatant was discarded. RNA pellet was dried for 5 min at room temperature and subsequently resuspended in 150 µl of nuclease-free ice-cold water by shaking (1400 rpm) for 15 min at +4°C. RNA solution was aliquot for 10 µl and liquid nitrogen frozen. Concentration of mRNA was measured

spectrophotometrically and double-checked on agarose gel along with RiboRuler™ RNA Ladder, High Range (Fermentas).

mRNA/dsDNA complex was produced by ligation of long+ oligodeoxynucleotide to mRNA using ddC-Long2 oligodeoxynucleotide as a splint. The 5' end of long+ was previously phosphorylated using T4 Polynucleotide Kinase (Fermentas). Oligodeoxynucleotide ddC-Long2 has 3' end modification (ddC) in order to prevent possibility of 3' end extension by reverse transcriptase on its natural RNA-DNA substrate.

Annealing mix of 50µl was prepared by mixing of 17 pmol of purified mRNA mixture at ratio of 1:20=MLV_D583N_pD (active RT) : del_pD (inactive RT) with 4.3 time of molar excess of long+ and 4.1 time of molar excess of ddC-Long2 in nuclease free water. The mixture was incubated at 70°C for 5 min and then cooled to room temperature for 20 min. Before adding ligation reaction components mixture was moved to cooling stand for 2 min.

5 µl of 10x ligation buffer and 5µl of T4 DNA ligase (5v/µl) (Fermentas) were added to 40µl of annealing mixture. Negative ligation reaction was carried out using the same annealing mixture in 1x ligation buffer without T4 DNA ligase. Prepared ligation reaction mixtures were incubated at 37°C for 30min. To stop ligation 1µl of 0.5M EDTA (pH 8.0) was added to both tubes and reaction mixtures were extracted once with equal volume of Roti® Phenol/chloroform (ROTH) and twice with equal volume of chloroform followed by concentration of reaction products at 30°C for 10 min in vacuum Concentrator 5301 (Eppendorf). Desalting was performed using illiustra ProbeQuant G-50 MicroColumns (GE Healthcare). Concentrations of ligation products were determined on agarose gel along with RiboRuler™ RNA Ladder, High Range (Fermentas) – mRNA mixture (MLV_D583N_pD:del_pD=1:20) ligated to long+/ddC-Long2 oligodeoxynucleotides ~0.24 µg/µl (sample with T4 DNA ligase) and simple mRNA mixture with long+/ddC-Long2 oligodeoxynucleotides ~0.06 µg/µl (sample without T4 DNA ligase).

(4) General control of mRNA/dsDNA complex by incorporation of [α-P^{33}]dATP. Prepared mRNA/dsDNA complex (substrate) was tested for dTTP (or biotin-dUTP) and subsequent [α-P^{33}]dATP incorporation by reverse transcriptase. Reaction mixture: 16µl of 5x Reaction buffer for reverse transcriptase; 4 µl - 40 u/µl RiboLock ™ RNase inhibitor (Fermentas); 2 µl [α-P^{33}]dATP (10mCi)/ml, SRF-203 (Hartmann Analytic)); 35 µl nuclease free water; 1µl - 200 u/µl RevertAid™ Minus M-MuLV Reverse transcriptase (Fermentas). Prepared mix was

divided to two tubes 2x 28 µl and 2µl of 1mM dTTP (Fermentas) or 2µl of 1mM biotin-dUTP (Fermentas) was added. Resultant mixtures were divided to two tubes 2x 15µl. To first tube 1.25 µl (~0.3µg) of ligation products plus 3.75 µl of nuclease free water were added. To second tube 5 µl (~0.3µg) of negative ligation reaction products were added. Reaction mixtures were incubated at 37°C for 30 min. After that, 1µl of 0.5M EDTA (pH 8.0) was added to all tubes and reaction mixtures were extracted once with equal volume of Roti® Phenol/chloroform (ROTH) and twice with equal volume of chloroform followed by purification on illiustra ProbeQuant G-50 MicroColumns (GE Healthcare). Reactions products were analyzed on agarose gel along with RiboRuler™ RNA Ladder, High Range (Fermentas) FIG 14A. In all samples (with and without ligase) we can see discreet band (~2500b) of del_pD mRNA (20 times smaller amount of MLV_D583N_pD mRNA ~2700b, which was present in mRNA mixture can not be distinguished). Subsequently agarose gel was dried on filter paper and radiolabeled mRNA/dsDNA complex (at the same position as mRNA) was detected only in case of positive ligation samples (with ligase) and not in case of negative ligation samples (without ligase) FIG 14B.

(5) Selection for DNA dependent DNA polymerase activity using biotin-dUTP.

Previously prepared mRNA/dsDNA complex (mRNA mix MLV_D583N_pD (active RT):del_pD (inactive RT)=1:20) was used for in vitro translation employing synthetic WakoPURE system (295-59503 - Wako). Translation mixture for WakoPURE system (25 µl): 12,5 µl - A solution (Wako); 5 µl - B solution (Wako); 0,5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 0.25 µl - 1 M DTT; 1,75 µl nuclease-free water and 5 µl - 0.24 µg/µl mRNA/dsDNA substrate (~1200 ng). In vitro translation was performed for 120 min at 37°C. Translations (~25 µl) was stopped by adding 155 µl of ice-cold stop buffer WBK_{500}+DTT+triton (50 mM Tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - Triton X-100 (T8787 - Sigma)) and centrifuged for 5 min at +4°C and 25'000 g. Very carefully 160 µl of centrifuged translation mixture was transferred on the top of 840 µl 35% (w/v) sucrose solution in WBK_{500}+DTT+Triton X-100 (50 mM Tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - Triton X-100 (T8787 - Sigma); 35% (w/v) - sucrose (84097 - Fluka)) to transparent 1 ml ultracentrifugation tubes (343778 - Beckman). Ternary complexes (TC) consisted of mRNA/dsDNA-ribosome-protein(tRNA) were purified by ultracentrifugation at

TL-100 Beckman ultracentrifuge in TLA100.2 fixed angle rotor (Beckman) at 100'000 rpm for 9 min at +4°C. Initially 750 µl of solution was removed from the very top of the centrifugation tube. Then (very carefully to keep small transparent pellet of TC at the bottom of ultracentrifugation tube intact) tube walls were washed with 750 µl of WBK_{500} (50 mM tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl). Finally all solution was removed starting from the very top of the centrifugation tube and pellet was dissolved in 30 µl of ice-cold stop buffer WBK_{500}+DTT+triton (50 mM Tris-acetate pH 7.5 at 25°C; 50 mM NaCl; 50 mM Mg-acetate; 500 mM KCl; 10 mM DTT; 0.1% (v/v) - Triton X-100 (T8787 - Sigma)).

As it was determined using radioactively labeled mRNA after ultracentrifugation 5% - 30 % of input mRNA is located in ternary complex pellet. Therefore it was predicted to have less than 360 ng (30% from 1200 ng mRNA used in translation reaction) of mRNA in 30 µl of buffer (~12 ng/µl or $9*10^9$ molecules/µl of ternary complex).

Modified nucleotide incorporation reaction mix was prepared on ice by mixing: 5 µl - 5X reaction buffer for Reverse Transcriptase (Fermentas); 1.25 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 2.5 µl – 1mM biotin-dUTP (Fermentas); 40.95 µl nuclease-free water and 0.3 µl of purified (<$2.7*10^9$ molecules) TC. According to the protocol 50 µl of nucleotide incorporation reaction mix contains <$2.7*10^9$ molecules of ternary complex.

Oil-surfactant mixture was prepared by mixing ABIL EM 90 (Goldschmidt) into mineral oil (M5904 - Sigma) to final concentration of 4% (v/v) (Ghadessy and Holliger, 2004; US2005064460). Emulsions were prepared at +4°C in 5 ml cryogenic vials (430492 - Corning) by mixing 950 µl of oil-surfactant mixture with 50 µl of RT mixture. Mixing was performed using MS-3000 magnetic stirrer with speed control (Biosan) at ~2100 rpm; Rotilabo®- (3x8 mm) magnetic bar with centre ring (1489.2 - Roth); water phase was added in 10 µl aliquots every 30 sec, continuing mixing for 2 more minutes (total mixing time - 4 min). According to optical microscopy data compartments in prepared emulsions vary from 0.5 µm to 10 µm in size with average diameter of ~2 µm. Therefore it was expected to have ~$1*10^{10}$ water in oil compartments after the emulsification of 50 µl reverse transcription reaction mixture, which

contains less than $2.7*10^9$ molecules of ternary complex (about one mRNA-dsDNA complex and reverse transcriptase molecules per 3-4 compartments).

Prepared emulsion was incubated 30 min at +37°C.

To recover the reaction mixture from emulsion 20µl of 0.1M EDTA was added to emulsion, stirred for 10 sec, then 50µl phenol/chloroform mix was added and stirred for additional 10 sec. After that, emulsion was transferred to 1.5 ml microcentrifuge tube, 0.5ml of water-saturated ether was added mixed by vortexing and centrifuged for 10 min at room temperature for 16'000 g. Oil-ether phase was removed leaving concentrated (but still intact) emulsion at the bottom of the tube. Finally emulsions were broken by extraction with 0.9 ml water-saturated ether; 0.9 ml water-saturated ethyl-acetate (in order to remove ABIL EM 90 detergent) and twice 0.9 ml water-saturated ether. Water phase was dried for 12 min under vacuum at room temperature followed removal of incorporated nucleotides on illiustra ProbeQuant G-50 MicroColumns (GE Healthcare). 2 µl aliquot of resultant mixture was left for direct RT reaction (without streptavidin beads purification) and the remaining part of solution was used for biotinylated mRNA/dsDNA complex immobilization on Dynabeads M-280 Streptavidin beads (DYNAL Biotech).

Dynabeads® kilobase BINDER™ Kit (DYNAL Biotech) was used for isolation of biotinylated mRNA-dsDNA complex according provided product description. 5µl of resuspended Dynabeads was transferred to a 1.5 ml microcentrifuge tube, washed with 20 µl of provided Binding Solution. The tube was placed on the magnet for 1-2 min until beads settled at the tube and solution was removed. Dynabeads were gently resuspended by pipetting in 50 µl of Binding Solution and 1µl of aqueous tRNA (tRNA from yeast (Roche)) solution (1µg/µl) was added to minimize non-specific mRNA binding. 50µl of Dynabeads in Binding solution were added to solution (~50µl) containing the biotinylated RNA-DNA fragments. The tube was incubated shaking at the +22°C in termomixer (Eppendorf) for 1 hour. After the binding of mRNA/dsDNA, supernatant was removed from the beads, and the beads were washed two times with 50 µl of Washing Solution (provided in the kit) shaking (1400 rpm) for 5 min at +22°C and once with 50 µl of Washing Solution shaking (1400 rpm) for 12 min at +22°C.

Collected Dynabeads with immobilized biotinylated mRNA/dsDNA complex were resuspended in reverse transcription reaction mixture.

Reverse transcription reaction mixture for selected mRNA/dsDNA complex was prepared on ice: 12 µl - 5X reaction buffer for Reverse Transcriptase (Fermentas); 6 µl – 10 mM dNTP (Fermentas); 1.5 µl - 40 u/µl RiboLock RNase inhibitor (Fermentas); 0,3 µl - 20 µM pD_42 oligonucleotide (SEQ ID No: 8); 1,5µl RevertAid™ Minus M-MuLV Reverse transcriptase (200 u/µl)(Fermentas); 35.7 µl nuclease-free water. Prepared mix was divided to three 19 µl aliquots: one was used to resuspend Dynabeads with immobilized biotinylated mRNA/dsDNA complex, the other aliquot of 19 µl was transferred to the tube with sample of elongated mRNA/dsDNA complex left without streptavidin beads purification and to the rest of 19 µl aliquot was added 1µl of nuclease free water (negative reaction control - to prove that reaction mixture is not contaminated). All reaction mixtures were incubated by shaking (1000 rpm) at the +42°C in termomixer (Eppendorf) for 1 hour.

Amplification of cDNA was performed by nested PCR. Initial PCR mixture was prepared on ice: 10 µl - 10X Taq buffer with KCl (Fermentas); 10 µl - 2 mM of each dNTP (Fermentas); 6 µl - 25 mM $MgCl_2$ (Fermentas); 2 µl - 1 u/µl LC (recombinant) *Taq* DNA Polymerase (Fermentas); 1 µl – 2,5 u/µl *Pfu* DNA Polymerase (Fermentas); 0.5 µl - 100 µM RD_Nde primer (SEQ ID No: 9); 0.5 µl - 100 µM pD_55 primer (SEQ ID No: 10); 65 µl water - mixture was divided into 5 samples for 19 µl (5x19 µl). To 3x 19 µl of PCR master mix were added 1 µl of cDNA (1-3 RT samples); to one tube with 19 µl of PCR master mix 1 µl water was added - negative PCR control. For positive PCR control 1 µl of pET_his_MLV_pD plasmid (~1ng) was added. The cycling protocol was: initial denaturation step 3 min at 94°C, 30 cycles (45 sec at 94°C, 45 sec at 58°C, and 3 min at 72°C) and final elongation 5 min at 72°C.

Nested PCR mixture for partial gene amplification (for better resolution of MLV_D583N_pD:del_pD cDNA ratio in RT samples) was prepared on ice: 20 µl - 10X Taq buffer with KCl (Fermentas); 20 µl - 2 mM of each dNTP (Fermentas); 12 µl - 25 mM $MgCl_2$ (Fermentas); 0.9 µl - 5 u/µl *Taq* DNA Polymerase (Fermentas); 1.0 µl - 100 µM M_F primer

(SEQ ID No: 11); 1.0 µl - 100 µM M_2R primer (SEQ ID No: 12); 135.1 µl water - mixture was divided 5x 38 µl. 2 µl of first PCR (primers set RD_Nde//pD_55) product were added to prepared nested PCR mixture. Master mix was mixed again and divided into two tubes (2x 20 µl) for 30 or 35 PCR cycles amplification. The cycling protocol was: initial denaturation step 3 min at 94°C, 30 or 35 cycles (45 sec at 94°C, 45 sec at 57°C, and 1 min at 72°C) and final elongation 3 min at 72°C. Expected length of PCR fragments was 907 bp for MLV_D583N_pD and 736 bp for del_pD. Amplification was analyzed on 1% agarose gel loading 10 µl of PCR mix per well (FIG 15).

Results

1. Double stranded (dsDNA) adaptor was successfully ligated to mRNA using T4 DNA ligase. Ligation efficiency is about 60% as it was determined by ligation of dsDNA-biotin adapter. mRNA/dsDNA complex can be specifically purified on streptavidin beads, providing an opportunity to discriminate between biotin labelled and unlabelled substrates. Free mRNA is much worse substrate for DNA dependent DNA polymerase comparing to mRNA/dsDNA. As a consequence 60% ligation efficiency of dsDNA to mRNA is good enough and such a substrate can be successfully used in evolution scheme.

2. General selection experiment using mRNA/dsDNA complex (mRNA mixture MLV_D583N_pD:del_pD=1:20) was performed. In vitro translation was performed using WakoPURE protein translation system and compartmentalized biotin-dUTP incorporation reaction in to dsDNA was carried out to demonstrate the enrichment of genes encoding active (MLV_D583N_pD) reverse transcriptase over genes encoding inactivated enzyme (del_pD). According to selection scheme (FIG 12) incorporation reaction of biotin-dUTP should occur only in aqueous compartments containing active (MLV_D583N_pD) reverse transcriptase resulting in biotinylation of mRNA/dsDNA complex. DNA dependent DNA polymerase is selected by binding of the biotinylated complex to streptavidin immobilized on magnetic beads, and then the selected gene is amplified by RT-PCR. Genes encoding active enzyme (in our case RT-PCR fragment for MLV_D583N_pD reverse transcriptase) were enriched over genes encoding inactive enzyme (FIG 15). Initial ratio of genes MLV_D583N_pD:del_pD was 1:20 and final ratio (after enrichment) was ~1:1. Respectively an enrichment factor in this particular experiment was ~20 folds. Enrichment factors calculated from different experiments varied in range from 5 to 200. And it was confirmed that DNA dependent DNA polymerase could be selected for modified nucleotide incorporation applying Compartmentalized

Ribosome Display (CRD) method. Selection of conventional DNA dependent DNA polymerases can be performed straight away. Biotin-dUTP can be exchanged to different nucleotide analogues of interest including nucleotide analogues having 3' modifications. After the incorporation of such nucleotide analogues into DNA strand 3' end is blocked, can't be extended and elongation reaction will be terminated. This approach is used in sequencing by synthesis (SBS) scheme and DNA polymerase suitable for SBS can be easily evolved using compartmentalized ribosome display (CRD) technique.

NcoI

```
ccatgggcatgaccctaaatatagaagatgagcatcggctacatgagacctcaaaagagc    60
cagatgtttctctagggtccacatggctgtctgattttcctcaggcctgggcggaaaccg   120
ggggcatgggactggcagttcgccaagctcctctgatcatacctctgaaagcaacctcta   180
ccccccgtgtccataaaacaatacccccatgtcacaagaagccagactggggatcaagcccc  240
                                g                           g
acatacagagactgttggaccagggaatactggtaccctgccagtcccccctggaacacgc   300
                                                    a
ccctgctacccgttaagaaaccagggactaatgattataggcctgtccaggatctgagag    360
                        g                       g
aagtcaacaagcgggtggaagacatccaccccaccgtgcccaacccttacaacctcttga    420
                  c
gcgggctcccaccgtcccaccagtggtacactgtgcttgatttaaaggatgccttttttct   480
          c
gcctgagactccacccccaccagtcagcctctcttcgcctttgagtggagagatccagaga   540
tgggaatctcaggacaattgacctggaccagactcccacagggtttcaaaaacagtccca    600
                        c                               t   t
ccctgtttgatgaggcactgcacagagacctagcagacttccggatccagcaccccagact   660
             g
tgatcctgctacagtacgtggatgacttactgctggccgccacttctgagctagactgcc    720
aacaaggtactcgggccctgttacaaaccctagggaacctcgggtatcgggcctcggcca    780
           g                    c
agaaagcccaaatttgccagaaacaggtcaagtatctggggtatcttctaaaagagggtc    840
agagatggctgactgaggccagaaaagagactgtgatggggcagcctactccgaagaccc    900
                                                    a
ctcgacaactaagggagttcctagggacggcaggcttctgtcgcctctggatccctgggt    960
ttgcagaaatggcagcccccttgtaccctctcaccaaaacggggactctgtttaattggg   1020
gcccagaccaacaaaaggcctatcaagaaatcaagcaagctcttctaactgccccagccc   1080
                                    c
tggggttgccagatttgactaagccctttgaactctttgtcgacgagaagcagggctacg   1140
ccaaaggtgtcctaacgcaaaaactgggaccttggcgtcggccggtggcctacctgtcca   1200
aaaagctagacccagtagcagctgggtggcccccttgcctacggatggtagcagccattg   1260
ccgtactgacaaaggatgcaggcaagctaaccatgggacagccactagtcattctggccc   1320
  -------                           ---------------
cccatgcagtagaggcactagtcaaacaacccccgaccgctggctttccaacgcccgga    1380
tgactcactatcaggccttgcttttggacacggaccgggtccagttcggaccggtggtag   1440
ccctgaacccggctacgctgctcccactgcctgaggaagggctgcaacacaactgccttg   1500
             g
atatcctggccgaagcccacggaacccgacccgacctaacggaccagccgctcccagacg   1560
                                    c       g
ccgaccacacctggtacacggatggaagcagtctcttacaagagggacagcgtaaggcgg   1620
gagctgcggtgaccaccgagaccgaggtaatctgggctaaagccctgccagccgggacat   1680
ccgctcagcgggctgaactgatagcactcacccaggccctaaagatggcagaaggtaaga   1740
                                                        g
agctaaatgtttatactgatagccgttatgcttttgctactgcccatatccatggagaaa   1800
tatacagaaggcgtgggttgctcacatcagaaggcaaagagatcaaaaataaagacgaga   1860
tcttggccctactaaaagccctctttctgcccaaaagacttagcataatccattgtccag   1920
           a                    g
gacatcaaaagggacacagcgccgaggctagaggcaaccggatggctgaccaagcggccc   1980
gaaaggcagccatcacagagactccagacacctctacctcctcatagaaaattcatcac    2040
ccaattcccgcttaattaatgaattc
                    EcoRI
```

APPENDIX 2

```
MLV      1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
1        1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLVI PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
3        1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
5        1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
7        1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
8        1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVSC QSPWNTPLLP
10       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
11       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDLP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
12       1   TLNIEDEHRL HETSKEPDVS LGPTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILAPC QSPWNTPLLP
13       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII SLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
16       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
17       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLVI PLKATSTPVS IKQYSMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
18       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLNQGILVPC QSPWNTPLLP
20       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
21       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
25       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
28       1   TLNIEDERRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
30       1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_1     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_2     1   TLNIKDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAIRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_3     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_4     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_6     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_8     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_9     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLNQGILVPC QSPWNTPLLP
L5_11    1   TLNIEDEHRL HETSTEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_13    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_14    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETWGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_15    1   TLNIEDEHRL HETSKESDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_16    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC LSPWNTPLLP
L5_18    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_20    1   TLNIEDEHRL HATSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGIPVPC QSPWSTPLLP
L5_21    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYSMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_23    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFL QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_24    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_25    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGLG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_28    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_29    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_30    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_32    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQGA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_35    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_37    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_39    1   TLNIEDEHRL HVTSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGVLVPC QSPWNTPLLP
L5_40    1   TLNIEDEHRL HETSKEPDVS LGSAWLSDFP QAWAETGGMG RAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_41    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVA IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_42    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_43    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_44    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QPPWNTPLLP
L5_46    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_47    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_48    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKERIQR LLDQGILVPC QSPWNTPLLP
L5_49    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLTV PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
L5_51    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII PLKATSTPVS IKQYPMSQEA RLGIKPHIQR LLDQGILVPC QSPWNTPLLP
```

APPENDIX 2 - continued

```
L5_52     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_53     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_55     1   TLNIKDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_56     1   TLNIEDEPRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_57     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLVI P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_58     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PLSQ A RLGIKPHIQR LLD GILVPY QSPWNTPLLP
L5_59     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_60     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_61     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQC PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_62     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_63     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLIV P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_64     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_65     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_66     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSLWNTPLLP
L5_68     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGVG LAVRQAPLVL P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC RSPWNTPLLP
L5_69     1   TLNIEDEHRL HETTKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_71     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTSLLP
L5_72     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_73     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMPQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_75     1   TLNIEDEHRL YETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_76     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PLSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_78     1   TLNIEDEHRL RETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_79     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGTKPHIQR LLD GILVPC QSPWNTPLLP
L5_80     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_81     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_82     1   TLNIEDEHRL HETSKEPDVS LGSTWPSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_84     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_85     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_87     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_88     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ V RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_90     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_92     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_93     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_94     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_95     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_96     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_97     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_99     1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_101    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ SMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_103    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNMPLLP
L5_104    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_106    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_107    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_111    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ SMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_112    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_114    1   TLNIEDEHRL HETSKESDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_115    1   TLNIEDEPRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_116    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_117    1   TLNIEDEHRL HETPKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWSTPLLP
L5_118    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
L5_120    1   TLNIEDEHRL HETSKEPDVS LGSTWLSDFP QAWAETGGMG LAVRQAPLII P KATSTPVS IKQ PMSQ A RLGIKPHIQR LLD GILVPC QSPWNTPLLP
```

APPENDIX 2 - continued

MLV	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
1	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
3	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
5	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGPP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
7	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
8	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGPP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
10	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
11	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTMLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
12	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
13	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNIWSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
16	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
17	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
18	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGPP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
20	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
21	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP SPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGTS GQLTWTRLP GFKNSPLFD
23	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
28	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
30	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
L5_1	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_2	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLAWTRLP GFKNSPLFD
L5_3	101	VKKPGTNDYR PVQDLEVNK RVEDVHPTVP NPNLLSRLP PSRQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_4	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_6	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_8	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_9	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPVNPLSGLP PSHQWYTVLD LDAFCLRL HPTSQSLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_11	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
L5_13	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVS NPNLLSGLP PSHQWYTVLD LDAFCLKL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFA
L5_14	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPALFG
L5_15	101	VKKPGTNEYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_16	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGDP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_18	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_20	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFA
L5_21	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_23	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_24	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGPP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_25	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_28	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_29	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRGPEMGIS GQLTWTRLP GFKNSPLFD
L5_30	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFA
L5_32	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNPLSGPP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_35	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
L5_37	101	VKKPGTNDYR PVQDLEVNK RVEDISPTVP NPNLLSGLP PSHQWYTVFD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_39	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
L5_40	101	VKKPGTNDYR PVQDLEVNK RVEDIHSTVP NPNLLSGLP PSHQWYTVLD FDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_41	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGPP PSHQWYTVLD LDAFCLRL HPTSQPLSAF EWRDPEMGIS GQLTWTRLP GFKNSPLYD
L5_42	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GLKNSPLFD
L5_43	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
L5_44	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_46	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFN
L5_47	101	VKKPGTSDYR PVQDLEVNK RVEDIRPAVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGVS GQLTWTRLP GFKNSPLFN
L5_48	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD
L5_49	101	VKKPGTNDYR PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKSSPLFD
L5_51	101	VKKPGTNDYG PVQDLEVNK RVEDIHPTVP NPNLLSGLP PSHQWYTVLD LDAFCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP GFKNSPLFD

APPENDIX 2 - continued

```
L5_52     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_53     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVS NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_55     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_56     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_57     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWKDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_58     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_59     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_60     101  VKKPGTNDYR PAQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_61     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_62     101  VKKPGTNDYR PVQDL EANK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_63     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKSSP LFD
L5_64     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_65     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_66     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_68     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_69     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_71     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLAWTRLP  GFKNSP LFD
L5_72     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_73     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_75     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_76     101  VKKPETNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_78     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTMLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_79     101  VKKRGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_80     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_81     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQSLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LLD
L5_82     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVS NP NLLCGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_84     101  VKKPGTNDYR PVQDL EVNK RVEDIRPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_85     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_87     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_88     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_90     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_92     101  VKKPGTNDYR PVQDL EVNK RVEDISPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_93     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_94     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_95     101  VKKPGTNDYR PVQDL EVNK RVECIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWIGLP  GFKNSP LFD
L5_96     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_97     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_99     101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_101    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_103    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_104    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NS NLLSGLP PSHQWYTVLD L DA FCLRL HPTGQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_106    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_107    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_111    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_112    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFN
L5_114    101  VKKPGTNEYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_115    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSPALFG
L5_116    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_117    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGPP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLPR GFKNSP LFD
L5_118    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
L5_120    101  VKKPGTNDYR PVQDL EVNK RVEDIHPTVP NP NLLSGLP PSHQWYTVLD L DA FCLRL HPTSQPLFAF EWRDPEMGIS GQLTWTRLP  GFKNSP LFD
```

APPENDIX 2 - continued

```
MLV      201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
1        201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
3        201  EALIRDLADF RIQHPDLILL RYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
5        201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEDRKETVMG QPTPKTPRQL
7        201  EAIRRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
8        201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKEAVMG QPTPKTPRQL
10       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
11       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
12       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCRQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
13       201  EALIRDPADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
16       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT QALLQTLGNL GYRASAKKTQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
17       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
18       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
20       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
21       201  EALRRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
23       201  EALIRDLADF RIQHPDLILL RYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
28       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
30       201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_1     201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGDL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_2     201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_3     201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_4     201  EALIRDLVDF RIQHPDLILL QYVDCLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_6     201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKEAVMG QPTPKTPRQL
L5_8     201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_9     201  EALIRDLADF RIQRPDLILL QCVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPAPKTPRQL
L5_11    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_13    201  EALIRDLADF RIQHPDLILL RYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_14    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_15    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGDL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_16    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_18    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_20    201  EALIRDLADF RIQHPDLILL RYVDDLLLAA TSELDCQQGT RTLLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_21    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSKLDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_23    201  EALIRDLADF RIQHPDLILL QYMDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_24    201  EALIRDLADF RIQHPDLILL RYVDDLLLAA TSELDCQQGT RALEQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_25    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_28    201  EALIRDLADF RIQHPDEILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_29    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_30    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_32    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_35    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_37    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_39    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_40    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_41    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_42    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_43    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCRQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_44    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_46    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_47    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_48    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
L5_49    201  EALRRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKEAVMG QPTPKTPRQL
L5_51    201  EALIRDLADF RIQHPDLILL QYVDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVMG QPTPKTPRQL
```

APPENDIX 2 - continued

```
L5_52    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_53    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_55    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGA RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_56    201  EALRDLADF WIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QLTPKTPRQL
L5_57    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_58    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_59    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_60    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWP TEARKETVLG QPTPKTPRQL
L5_61    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_62    201  EALRRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TCARKETVLG QPTPKTPRQL
L5_63    201  EALRRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKEAVLG QPTPKTPRQL
L5_64    201  EALRDLADF RIQHPGLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_65    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQHGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_66    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_68    201  EALRRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKEAVLG QPTPKTPRQL
L5_69    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_71    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNP GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_72    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_73    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_75    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_76    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVVG QPTPKTPRQL
L5_78    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_79    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_80    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_81    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_82    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_84    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_85    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_87    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_88    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_90    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPGQL
L5_92    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_93    201  EALRDLADF RIQHPGLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_94    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_95    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICRKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_96    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_97    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_99    201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_101   201  EALRRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEDRKETVLG QPTPKTPRQL
L5_103   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_104   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_106   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_107   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSKLDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPETPRQL
L5_111   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSKLDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_112   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCRQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_114   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGDL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_115   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_116   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_117   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
L5_118   201  EALRDLADF RIQHPDLILL RYDDLLLAA TSELDCQQGT RALLQTLGNL GYRASAKKAQ ICQKQVKYLG YLLKEGQRWL TERRKEAVLG QPTPKTPRQL
L5_120   201  EALRDLADF RIQHPDLILL QYDDLLLAA TSELDCQQGT RALLQTLGNL GHRASAKKAQ ICQKQVKYLG YLLKEGQRWL TEARKETVLG QPTPKTPRQL
```

MLV	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
1	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
3	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
5	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALVLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
7	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
8	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
10	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
11	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRW PVAYLSKKLD
12	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
13	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
16	301	R FLG AG C HL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
17	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
18	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
20	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
21	301	R FLG AG C RL IPGFAEM AALLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLAAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
23	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
28	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
30	301	R FLGMAG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_1	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_2	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR LVAYLSKKLD
L5_3	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGA LTQKLGPWRR PVAYLSKKLD
L5_4	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_6	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_8	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_9	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAHQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_11	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_13	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_14	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_15	301	R FLG VG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAHQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_16	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_18	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_20	301	R FLG AG C RL IPGFAEM AAPLYPLTKT ETLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_21	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_23	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPRRW PVAYLSKKLD
L5_24	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_25	301	RKFLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_28	301	R FLGAAGS C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_29	301	R FLG AG C RP IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_30	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_32	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_35	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_37	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_39	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_40	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_41	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQDIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_42	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTPFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_43	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_44	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_46	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_47	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_48	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_49	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_51	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD

APPENDIX 2 - continued

L5_52	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKRGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_53	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_55	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_56	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_57	301	RKFLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PAAYLSKKLD
L5_58	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_59	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_60	301	R FLG AG C RL IPGFAEM ATPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKSV LTQKLGPWRR PVAYLSKKLD
L5_61	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA VLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_62	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQDIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_63	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_64	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_65	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_66	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_68	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_69	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_71	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_72	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVTYLSKKLD
L5_73	301	R FLG VG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_75	301	R FLG AG C RL IPGFAEM AAPLHPLTKT GTQFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_76	301	R FLG AG C RL TPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_78	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_79	301	R FLG AG C RL IPGFAEM AAPLHPLTKT GTQFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_80	301	R FLG AG C RL IPGFAEM AAPLYSLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_81	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_82	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKTYQEIKQA LLTAPCLGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_84	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_85	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_87	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_88	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_90	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKRGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_92	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_93	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_94	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_95	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_96	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_97	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_99	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_101	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR SVAYLSKKLD
L5_103	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_104	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPRRR PVAYLSKKLD
L5_106	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_107	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_111	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_112	301	R FLG AG C RL IPGFAEM AAPLYPLTKP GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_114	301	R FLG VG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAHQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_115	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALWLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_116	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD
L5_117	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTQFNWGPGQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGA LTPKLGPWRR PVAYLSKKLD
L5_118	301	R FLG AG C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELIV DEKQGCAKGV LTQKLGPWRR PVAYLSKKLD
L5_120	301	R FLG AS C RL IPGFAEM AAPLYPLTKT GTLFNWGPDQ QKAYQEIKQA LLTAPALGLP DLTKPFELFV DEKQGYAKGV LTQKLGPWRR PVAYLSKKLD

APPENDIX 2 - continued

```
MLV      401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
1        401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
3        401  PVAAGWPPCL RMVAAIAVLT KDAGKLTLGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
5        401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
7        401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLAIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
8        401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
10       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP TTLLPLPEEG LQHNCLDILA
11       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
12       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
13       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTLGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
16       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
17       401  PVAAGWPPRL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRA QFGPVVALNP ATLLPLPEEG LQHNCLDILA
18       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALDP ATLLPLPEEG LQHNCLDILA
20       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
21       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
23       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
28       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
30       401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_1     401  PVAAGWPPRL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRA QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_2     401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_3     401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_4     401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_6     401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_8     401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPGVALNP ATLLPLPEEG LQHNCLDILA
L5_9     401  PVAAGWLPCL RMVAVIAVLT KDAGKLTMGQ PLVITPHAV EAIAKQLPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_11    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_13    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPCR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_14    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PIATAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_15    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_16    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_18    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARTTHY QTLLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_20    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTLGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_21    401  PVAAGWSPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVAPNP ATLLPLPEEG LQHNCLDILA
L5_23    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIPAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_24    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_25    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_28    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_29    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_30    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPGR WLSARMTHY QALLLDTDRV QFGPVVALDP ATLLPLPEEG LQHNCLDILA
L5_32    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ QLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_35    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALDP ATLLPLPEEG LQHNCLDILA
L5_37    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSKARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_39    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_40    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_41    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDH WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_42    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_43    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_44    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGR PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALDP ATLLPLPEEG LQHNCLDILA
L5_46    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_47    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_48    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_49    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_51    401  PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ QLVIAPHAV EALVKQPPDR WLSARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
```

L5_52	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_53	401	PVAAGWPPCL RMVAATVVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_55	401	PVAAGWPPCL RMVAATAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_56	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPPPEEG LQHNCLDILA
L5_57	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_58	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_59	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDH WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_60	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_61	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPAR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_62	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_63	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_64	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_65	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_66	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARRTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_68	401	PVAAGWPPCL RMVAATAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLSKARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCEDILA
L5_69	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_71	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLGILA
L5_72	401	PVAAGWPPCL RMVAATAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_73	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_75	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGR PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_76	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI TPHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG PQHNCLDILA
L5_78	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTRY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_79	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGR PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_80	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_81	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKRPPDR WLS ARMTHY QALLLDTARV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_82	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_84	401	PVAAGWPPCL RMVAAIVVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG PQHNCLDILA
L5_85	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_87	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_88	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALDP ATLLPLPEEG LQHNCLDILA
L5_90	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_92	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_93	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_94	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHDCLDILA
L5_95	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHKCLDILA
L5_96	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_97	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLPPLPEEG LQHNCLDVLA
L5_99	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_101	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_103	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_104	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_106	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_107	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_111	401	PVAAGWSPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVAPNP ATLLPLPEEG LQHNCLDILA
L5_112	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_114	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_115	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_116	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_117	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_118	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVT APHAV EALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA
L5_120	401	PVAAGWPPCL RMVAAIAVLT KDAGKLTMGQ PLVI APHAV GALVKQPPDR WLS ARMTHY QALLLDTDRV QFGPVVALNP ATLLPLPEEG LQHNCLDILA

APPENDIX 2 - continued

MLV	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
1	501	EAHGTRPDLT DQPLPDADHT WYTAGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIRGEIYRR
3	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
5	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
7	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KIAEGKKLNV YTDSRYAFAT AHIHGEIYRR
8	501	EAHGTRPDLT DQPFPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
10	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
11	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
12	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
13	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQW AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
16	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTEIEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
17	501	EVHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTNSRYAFAT AHIHGEIYRR
18	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKQLNV YTGSRYAFAT AHIHGEIYRR
20	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
21	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTASRYAFAT AHIHGEIYRR
23	501	EAHGTRPDLT DQPLPDADHT WYTAGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
28	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
30	501	EAHGTRPDLT DQPLPDADHT WYTNGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTGSRYAFAT AHIHGEIYRR
L5_1	501	EVHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_2	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTKTEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_3	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK APPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_4	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_6	501	EAHGTRPDLT DQPLPDADHT WYTAGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIRGEIYRR
L5_8	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_9	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_11	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGRKLNV YTDSRYAFAT AHIHGEIYRR
L5_13	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTESRYAFAT AHIHGEIYRR
L5_14	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_15	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGRIYRR
L5_16	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTGSRYAFAT AHIHGEIYRK
L5_18	501	EAHRTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSARR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_20	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_21	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_23	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_24	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_25	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_28	501	EAHGTRPDLT DQPLPDADHT WYTAGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIRGEIYRR
L5_29	501	EVHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_30	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTNSRYAFAT AHIHGEIYRR
L5_32	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTGSRYAFAT AHIHGEIYRR
L5_35	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_37	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTNSRYAFAT AHIHGEIYRR
L5_39	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_40	501	EARGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_41	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTGSRYAFAT AHIHGEIYRR
L5_42	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGASAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_43	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTNSRYAFAT AHIHGEIYRR
L5_44	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTNSRYAFAT AHIHGEIYRR
L5_46	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTNSRYAFAT AHIHGEIYRR
L5_47	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_48	501	EAHGTRPDLT DQPLPDADHT WYTGGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_49	501	EAHGTRPDLT DQPLPDADHT WYTAGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR
L5_51	501	EAHGTRPDLT DQPLPDADHT WYTDGSSLLQ EGQRKAGAAV TTETEVIWAK ALPAGTSAQR AELIALTQAL KMAEGKKLNV YTDSRYAFAT AHIHGEIYRR

APPENDIX 2 - continued

APPENDIX 2 - continued

```
MLV       601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
1         601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
3         601  RRWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
5         601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ANQAARKAA ITETPDTSTL LIENSSPNSR LIN
7         601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
8         601  RGLLTSKGKE IKNKDEILAL LKALFLPKRL SIIHSPGIQK GHSAEARGNR ADQAARKAA ITDTPDTSTL LIENSSPNSR LIN
10        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
11        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR AAQAARKAA ITETPDTSTL LIENSSPNSR LIN
12        601  RGLLTSEGKE IKNKDEILAL LKVLFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
13        601  RGWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
16        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQATRKAA ITETPDTSTL LIENSSPNSR LIN
17        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
18        601  RGLLTSEGKE IKNKDEIVAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPIPA
20        601  RGWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPIPA
21        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPIPA
23        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSVEARGNR ADQAARKAA VTETPDTSTL LIENSSPNSR LIN
28        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL STIHCPGIQK GHSVEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
30        601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_1      601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_2      601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_3      601  RGWLTSEGKE IKNKDEILAL LRALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARQAA ITETPDTPTL LIENSSPNSR LIN
L5_4      601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSA
L5_6      601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSP
L5_8      601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNPA
L5_9      601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_11     601  RGLLTSAGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_13     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_14     601  RGWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_15     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR AGQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_16     601  RGMLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_18     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQVARKAA ITETPDTSTL LIENSSPNSR LIN
L5_20     601  RGWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_21     601  RGWLTSEGKE IKNKDEILAL LKALFLPKRLQK GHSAEARGNR ANQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_23     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_24     601  RGWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGSR ADQAARRAA ITETPDTSTL LIENSSPNSR LIN
L5_25     601  RGLLTSEGKE IKNKDEILAL LKALFLPERL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_28     601  RGLLTSEGKE IKNKDEILAL LKALSLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_29     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_30     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRLF SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL PIENSSPNSR LIN
L5_32     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNPR LIN
L5_35     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_37     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_39     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEVRGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_40     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR AGQAARKAA ITETPDTSTL LIENSSPNSR
L5_41     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_42     601  RGWLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_43     601  RGLLTSEGKE IKNKDEILAL LKALSLPKRL SIIHCPGIQK GHSAEARGNR ANQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_44     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_46     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSTEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_47     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GYSAEARGNR ANQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_48     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_49     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ADQAARKAA ITETPDTSTL LIENSSPNSR LIN
L5_51     601  RGLLTSEGKE IKNKDEILAL LKALFLPKRL SIIHCPGIQK GHSAEARGNR ANQAARKAA ITETPDTSTL LIENSSPNSR LIN
```

APPENDIX 2 - continued

L5_9	12 mutations,	83D->N, 135L->P, 166P->S, 214H->R, 222Y->C, 293T->A, 344Y->H, 407P->L, 415A->V, 436A->T, 444V->A, 447P->L
18	11 mutations,	83D->N, 139L->P, 200D->N, 330T->P, 479N->D, 577K->Q, 583D->G, 618L->V, 678N->I, 679S->P, 680R->A
L5_79	10 mutations,	74I->T, 104P->R, 325Y->H, 333L->Q, 430Q->R, 597I->T, 616E->K, 649N->S, 658K->R, 673E->G
L5_82	10 mutations,	26L->P, 130P->S, 137S->G, 343A->T, 356A->G, 524D->G, 539A->T, 603L->W, 653D->N, 676S->P
L5_117	10 mutations,	14S->P, 95N->S, 139L->P, 190Q->R, 333L->Q, 339D->G, 380V->A, 383Q->P, 524D->G, 532G->R
21	9 mutations,	131N->S, 179I->T, 204H->R, 323P->L, 353T->A, 583D->A, 678N->I, 679S->P, 680R->A
L5_3	9 mutations,	125I->V, 138G->R, 143H->R, 380V->A, 552L->P, 603L->W, 622K->R, 658K->Q, 668S->P
L5_20	9 mutations,	12E->A, 87L->P, 95N->S, 200D->A, 221Q->R, 242A->T, 331G->E, 428M->L, 603L->W
L5_75	9 mutations,	11H->Y, 325Y->H, 333L->Q, 430Q->R, 597I->T, 616E->K, 649N->S, 658K->R, 673E->G
L5_76	9 mutations,	66M->L, 105G->E, 200D->N, 289M->V, 314I->T, 436A->T, 491L->P, 573A->T, 583D->N
8	8 mutations,	89P->S, 139L->P, 287T->A, 330T->P, 514L->F, 607E->K, 635C->S, 663E->D
L5_13	8 mutations,	130P->S, 159R->K, 200D->A, 221Q->R, 330T->P, 449D->G, 524D->G, 583D->E
L5_41	8 mutations,	60S->A, 139L->P, 168F->S, 199F->Y, 346E->D, 450R->H, 524D->G, 583D->G
L5_68	8 mutations,	39M->V, 49I->V, 91Q->R, 204H->R, 287T->A, 454N->K, 625F->L, 653D->H
L5_72	8 mutations,	139L->P, 200D->N, 330T->P, 393A->T, 572M->L, 594H->Q, 671L->P, 672I->T
L5_114	8 mutations,	17P->S, 108D->E, 249N->D, 307A->V, 344Y->H, 524D->G, 653D->G, 673E->K
L5_115	8 mutations,	8H->P, 139L->P, 197T->A, 200D->G, 358G->W, 524D->G, 623A->V, 653D->A
17	7 mutations,	49I->V, 65P->S, 200D->N, 409C->R, 470V->A, 502A->V, 583D->N
L5_15	7 mutations,	17P->S, 108D->E, 249N->D, 307A->V, 344Y->H, 524D->G, 653D->G
L5_21, L5_111	7 mutations,	65P->S, 233E->K, 407P->S, 478L->P, 603L->W, 638H->R, 653D->N
L5_40	7 mutations,	24T->A, 41L->R, 127P->S, 151L->F, 330T->P, 503H->R, 653D->G
L5_47	7 mutations,	107N->S, 126H->R, 128T->A, 179I->V, 200D->N, 642H->Y, 653D->N
L5_71	7 mutations,	97P->S, 184T->A, 250L->P, 289M->L, 497D->G, 551A->T, 562E->K
L5_78	7 mutations,	11H->R, 148V->M, 330T->P, 459H->R, 502A->V, 653D->G, 667T->A
L5_81	7 mutations,	166P->S, 199F->L, 446Q->R, 468D->A, 501E->K, 530Q->H, 558A->V
L5_118	7 mutations,	221Q->R, 283A->T, 287T->A, 369F->I, 376Y->C, 434I->T, 603L->W
13	6 mutations,	51P->S, 136L->W, 207L->P, 428M->L, 560R->W, 603L->W
L5_30	6 mutations,	200D->A, 330T->P, 449D->G, 479N->D, 583D->N, 671L->P
L5_32	6 mutations,	69E->G, 135L->P, 139L->P, 431P->Q, 583D->G, 679S->F
L5_43, L5_112	6 mutations,	200D->N, 237Q->R, 330T->P, 524D->G, 625F->S, 653D->N
L5_53	6 mutations,	130P->S, 139L->P, 417A->V, 524D->G, 583D->N, 634H->Y
L5_56	6 mutations,	8H->P, 211R->W, 292P->L, 486L->P, 524D->A, 594H->R
L5_57	6 mutations,	49I->V, 173R->K, 302E->K, 392V->A, 603L->W, 669T->S
L5_60	6 mutations,	112V->A, 200D->N, 280L->P, 322A->T, 379G->S, 653D->N
L5_62	6 mutations,	118V->A, 204H->R, 282E->G, 346E->D, 524D->A, 528L->I
L5_84	6 mutations,	126H->R, 139L->P, 417A->V, 491L->P, 524D->G, 653D->A
L5_95	6 mutations,	124D->G, 187R->G, 263Q->R, 494N->K, 583D->G, 618L->V
L5_107	6 mutations,	139L->P, 233E->K, 295K->E, 633I->T, 642H->R, 643S->G

16	5 mutations,	241R->Q, 259A->T, 311R->H, 544T->I, 656A->T
20	5 mutations,	200D->N, 603L->W, 678N->I, 679S->P, 680R->A
23	5 mutations,	221Q->R, 332T->I, 524D->A, 644A->V, 661I->V
L5_2	5 mutations,	5E->K, 43V->I, 184T->A, 391P->L, 543E->K
L5_14	5 mutations,	37G->W, 197T->A, 200D->G, 433V->A, 603L->W
L5_18	5 mutations,	457M->T, 462A->T, 504G->R, 559Q->R, 655A->V
L5_23	5 mutations,	30P->L, 223V->M, 388W->R, 390R->W, 435L->P
L5_24	5 mutations,	139L->P, 221Q->R, 603L->W, 649N->S, 658K->R
L5_28	5 mutations,	306T->A, 309F->S, 524D->A, 594H->R, 625F->S
L5_37	5 mutations,	126H->S, 149L->F, 200D->N, 454N->K, 583D->N
L5_49, L5_63	5 mutations,	50I->V, 194N->S, 204H->R, 287T->A, 524D->A
L5_55	5 mutations,	5E->K, 200D->N, 240T->A, 653D->H, 671L->P
L5_69	5 mutations,	14S->T, 49I->T, 538A->T, 603L->W, 653D->N
L5_88	5 mutations,	70A->V, 139L->P, 479N->D, 524D->G, 625F->L
L5_101	5 mutations,	65P->S, 204H->R, 283A->D, 391P->S, 583D->N
L5_104	5 mutations,	132P->S, 164S->G, 388W->R, 524D->G, 533Q->K
3	4 mutations,	221Q->R, 428M->L, 602G->R, 603L->W
5	4 mutations,	139L->P, 283A->D, 358G->V, 653D->N
7	4 mutations,	204H->R, 433V->A, 524D->G, 572M->I
11	4 mutations,	29F->L, 148V->M, 390R->W, 653D->A
12	4 mutations,	23S->P, 88V->A, 237Q->R, 623A->V
30	4 mutations,	200D->N, 306T->M, 524D->N, 583D->G
L5_1	4 mutations,	249N->D, 409C->R, 470V->A, 502A->V
L5_6	4 mutations,	287T->A, 524D->A, 594H->R, 680R->P
L5_8	4 mutations,	475V->G, 524D->G, 679S->P, 680R->A
L5_11	4 mutations,	15K->T, 200D->N, 576K->R, 607E->A
L5_16	4 mutations,	91Q->L, 583D->G, 600R->K, 603L->M
L5_29	4 mutations,	174D->G, 312L->P, 502A->V, 524D->G
L5_39	4 mutations,	12E->V, 86I->V, 200D->N, 646A->V
L5_42	4 mutations,	192F->L, 333L->P, 556T->A, 603L->W
L5_44	4 mutations,	92S->P, 430Q->R, 479N->D, 583D->N
L5_46	4 mutations,	200D->N, 330T->P, 583D->N, 644A->T
L5_52	4 mutations,	200D->N, 330T->P, 374Q->R, 583D->N
L5_61	4 mutations,	64Y->C, 351L->V, 449D->A, 530Q->H
L5_64, L5_93	4 mutations,	200D->N, 216D->G, 524D->A, 545E->G
L5_65	4 mutations,	200D->N, 238Q->H, 570L->I, 603L->W
L5_85, L5_96	4 mutations,	200D->N, 330T->P, 583D->A, 638H->R
L5_90	4 mutations,	200D->N, 298R->G, 330T->P, 374Q->R
L5_103	4 mutations,	96T->M, 200D->N, 559Q->P, 607E->G
L5_106	4 mutations,	524D->G, 583D->N, 635C->R, 670L->F
L5_120	4 mutations,	252Y->H, 308G->S, 441E->G, 603L->W
1	3 mutations,	49I->V, 524D->A, 594H->R
28	3 mutations,	8H->R, 632I->T, 644A->V
L5_4	3 mutations,	208A->V, 225D->G, 680R->A
L5_25	3 mutations,	39M->L, 302E->K, 628K->E
L5_35	3 mutations,	200D->N, 330T->P, 479N->D
L5_51	3 mutations,	110R->G, 431P->Q, 653D->N
L5_58	3 mutations,	66M->L, 90C->Y, 653D->V
L5_66	3 mutations,	93P->L, 457M->R, 603L->W
L5_73	3 mutations,	67S->P, 139L->P, 307A->V
L5_80	3 mutations,	326P->S, 583D->G, 676S->P
L5_92	3 mutations,	126H->S, 200D->N, 653D->G
L5_94	3 mutations,	494N->D, 524D->G, 607E->K
L5_97	3 mutations,	484L->P, 498I->V, 653D->A
10	2 mutations,	481A->T, 524D->G
L5_59	2 mutations,	450R->H, 503H->R
L5_99	2 mutations,	36T->I, 524D->G
L5_116	2 mutations,	653D->H, 662T->A
L5_48	1 mutations,	77H->R
WT_MLV, L5_87	0 mutations	

APPENDIX 4

Position 524, total mutations 31, 3 different amino acids
D->G, 20 sequences (7, 10, L5_8, L5_13, L5_15, L5_29, L5_41, L5_43, L5_58, L5_82, L5_84, L5_88, L5_94, L5_99, L5_104, L5_106, L5_112, L5_114, L5_115, L5_117)
D->A, 10 sequences (1, 23, L5_6, L5_28, L5_49, L5_56, L5_62, L5_63, L5_64, L5_93)
D->N, 1 sequences (30)

Position 200, total mutations 30, 3 different amino acids
D->G, 2 sequences (L5_14, L5_115)
D->A, 3 sequences (L5_13, L5_20, L5_30)
D->N, 25 sequences (17, 18, 20, 30, L5_11, L5_35, L5_37, L5_39, L5_43, L5_46, L5_47, L5_52, L5_55, L5_60, L5_64, L5_65, L5_72, L5_76, L5_85, L5_90, L5_92, L5_93, L5_96, L5_103, L5_112)

Position 653, total mutations 23, 5 different amino acids
D->V, 1 sequences (L5_58)
D->G, 5 sequences (L5_15, L5_40, L5_78, L5_92, L5_114)
D->A, 4 sequences (11, L5_84, L5_97, L5_115)
D->N, 10 sequences (5, L5_21, L5_43, L5_47, L5_51, L5_60, L5_69, L5_82, L5_111, L5_112)
D->H, 3 sequences (L5_55, L5_68, L5_116)

Position 583, total mutations 21, 4 different amino acids
D->E, 1 sequences (L5_13)
D->G, 7 sequences (18, 30, L5_16, L5_32, L5_41, L5_80, L5_95)
D->A, 3 sequences (21, L5_85, L5_96)
D->N, 10 sequences (17, L5_30, L5_37, L5_44, L5_46, L5_52, L5_58, L5_76, L5_101, L5_106)

Position 603, total mutations 18, 2 different amino acids
L->W, 17 sequences (3, 13, 20, L5_3, L5_14, L5_20, L5_21, L5_24, L5_42, L5_57, L5_65, L5_66, L5_69, L5_82, L5_111, L5_118, L5_120)
L->M, 1 sequences (L5_116)

Position 330, total mutations 15, 1 different amino acids
T->P, 15 sequences (8, 18, L5_13, L5_30, L5_35, L5_40, L5_48, L5_46, L5_52, L5_72, L5_78, L5_85, L5_90, L5_96, L5_112)

Position 139, total mutations 14, 1 different amino acids
L->P, 14 sequences (5, 8, 18, L5_24, L5_32, L5_41, L5_53, L5_72, L5_73, L5_84, L5_88, L5_107, L5_115, L5_117)

Position 204, total mutations 7, 1 different amino acids
H->R, 7 sequences (7, 21, L5_49, L5_62, L5_63, L5_68, L5_101)

Position 221, total mutations 6, 1 different amino acids
Q->R, 6 sequences (3, 23, L5_13, L5_20, L5_24, L5_118)

Position 287, total mutations 6, 1 different amino acids
T->A, 6 sequences (8, L5_6, L5_49, L5_63, L5_68, L5_118)

Position 680, total mutations 6, 2 different amino acids
R->P, 1 sequences (L5_6)
R->A, 5 sequences (18, 20, 21, L5_4, L5_8)

Position 49, total mutations 5, 2 different amino acids
I->T, 1 sequences (L5_69)
I->V, 4 sequences (1, 17, L5_57, L5_68)

Position 479, total mutations 5, 1 different amino acids
N->D, 5 sequences (18, L5_30, L5_35, L5_44, L5_88)

APPENDIX 4 continued

Position 594, total mutations 5, 2 different amino acids
H->Q, 1 sequences (L5_72)
H->R, 4 sequences (1, L5_6, L5_28, L5_56)

Position 625, total mutations 5, 2 different amino acids
F->S, 3 sequences (L5_28, L5_43, L5_112)
F->L, 2 sequences (L5_68, L5_88)

Position 679, total mutations 5, 2 different amino acids
S->F, 1 sequences (L5_32)
S->P, 4 sequences (18, 20, 21, L5_8)

Position 65, total mutations 4, 1 different amino acids
P->S, 4 sequences (17, L5_21, L5_101, L5_111)

Position 126, total mutations 4, 2 different amino acids
H->S, 2 sequences (L5_37, L5_92)
H->R, 2 sequences (L5_47, L5_84)

Position 333, total mutations 4, 2 different amino acids
L->Q, 3 sequences (L5_75, L5_79, L5_117)
L->P, 1 sequences (L5_42)

Position 502, total mutations 4, 1 different amino acids
A->V, 4 sequences (17, L5_1, L5_29, L5_78)

Position 607, total mutations 4, 3 different amino acids
E->G, 1 sequences (L5_103)
E->A, 1 sequences (L5_11)
E->K, 2 sequences (8, L5_94)

Position 638, total mutations 4, 1 different amino acids
H->R, 4 sequences (L5_21, L5_85, L5_96, L5_111)

Position 658, total mutations 4, 2 different amino acids
K->Q, 1 sequences (L5_3)
K->R, 3 sequences (L5_24, L5_75, L5_79)

Position 8, total mutations 3, 2 different amino acids
H->P, 2 sequences (L5_56, L5_115)
H->R, 1 sequences (28)

Position 130, total mutations 3, 1 different amino acids
P->S, 3 sequences (L5_13, L5_53, L5_82)

Position 233, total mutations 3, 1 different amino acids
E->K, 3 sequences (L5_21, L5_107, L5_111)

Position 237, total mutations 3, 1 different amino acids
Q->R, 3 sequences (12, L5_43, L5_112)

Position 249, total mutations 3, 1 different amino acids
N->D, 3 sequences (L5_1, L5_15, L5_114)

Position 283, total mutations 3, 2 different amino acids
A->D, 2 sequences (5, L5_101)
A->T, 1 sequences (L5_118)

APPENDIX 4 continued

Position 307, total mutations 3, 1 different amino acids
A->V, 3 sequences (L5_15, L5_73, L5_114)

Position 344, total mutations 3, 1 different amino acids
Y->H, 3 sequences (L5_9, L5_15, L5_114)

Position 407, total mutations 3, 2 different amino acids
P->S, 2 sequences (L5_21, L5_111)
P->L, 1 sequences (L5_9)

Position 428, total mutations 3, 1 different amino acids
M->L, 3 sequences (3, 13, L5_20)

Position 430, total mutations 3, 1 different amino acids
Q->R, 3 sequences (L5_44, L5_75, L5_79)

Position 449, total mutations 3, 2 different amino acids
D->G, 2 sequences (L5_13, L5_30)
D->A, 1 sequences (L5_61)

Position 644, total mutations 3, 2 different amino acids
A->T, 1 sequences (L5_46)
A->V, 2 sequences (23, 28)

Position 649, total mutations 3, 1 different amino acids
N->S, 3 sequences (L5_24, L5_75, L5_79)

Position 671, total mutations 3, 1 different amino acids
L->P, 3 sequences (L5_30, L5_55, L5_72)

Position 673, total mutations 3, 2 different amino acids
E->G, 2 sequences (L5_75, L5_79)
E->K, 1 sequences (L5_114)

Position 678, total mutations 3, 1 different amino acids
N->I, 3 sequences (18, 20, 21)

Position 5, total mutations 2, 1 different amino acids
E->K, 2 sequences (L5_2, L5_55)

Position 11, total mutations 2, 2 different amino acids
H->R, 1 sequences (L5_78)
H->Y, 1 sequences (L5_75)

Position 12, total mutations 2, 2 different amino acids
E->V, 1 sequences (L5_39)
E->A, 1 sequences (L5_20)

Position 14, total mutations 2, 2 different amino acids
S->T, 1 sequences (L5_69)
S->P, 1 sequences (L5_117)

Position 17, total mutations 2, 1 different amino acids
P->S, 2 sequences (L5_15, L5_114)

APPENDIX 4 continued

Position 39, total mutations 2, 2 different amino acids
M->V, 1 sequences (L5_68)
M->L, 1 sequences (L5_25)

Position 50, total mutations 2, 1 different amino acids
I->V, 2 sequences (L5_49, L5_63)

Position 66, total mutations 2, 1 different amino acids
M->L, 2 sequences (L5_58, L5_76)

Position 83, total mutations 2, 1 different amino acids
D->N, 2 sequences (18, L5_9)

Position 91, total mutations 2, 2 different amino acids
Q->R, 1 sequences (L5_68)
Q->L, 1 sequences (L5_116)

Position 95, total mutations 2, 1 different amino acids
N->S, 2 sequences (L5_20, L5_117)

Position 108, total mutations 2, 1 different amino acids
D->E, 2 sequences (L5_15, L5_114)

Position 135, total mutations 2, 1 different amino acids
L->P, 2 sequences (L5_9, L5_32)

Position 148, total mutations 2, 1 different amino acids
V->M, 2 sequences (11, L5_78)

Position 166, total mutations 2, 1 different amino acids
P->S, 2 sequences (L5_9, L5_81)

Position 179, total mutations 2, 2 different amino acids
I->T, 1 sequences (21)
I->V, 1 sequences (L5_47)

Position 184, total mutations 2, 1 different amino acids
T->A, 2 sequences (L5_2, L5_71)

Position 194, total mutations 2, 1 different amino acids
N->S, 2 sequences (L5_49, L5_63)

Position 197, total mutations 2, 1 different amino acids
T->A, 2 sequences (L5_14, L5_115)

Position 199, total mutations 2, 2 different amino acids
F->L, 1 sequences (L5_81)
F->Y, 1 sequences (L5_41)

Position 216, total mutations 2, 1 different amino acids
D->G, 2 sequences (L5_64, L5_93)

Position 289, total mutations 2, 2 different amino acids
M->V, 1 sequences (L5_76)
M->L, 1 sequences (L5_71)

APPENDIX 4 continued

Position 302, total mutations 2, 1 different amino acids
E->K, 2 sequences (L5_25, L5_57)

Position 306, total mutations 2, 2 different amino acids
T->A, 1 sequences (L5_28)
T->M, 1 sequences (30)

Position 325, total mutations 2, 1 different amino acids
Y->H, 2 sequences (L5_75, L5_79)

Position 346, total mutations 2, 1 different amino acids
E->D, 2 sequences (L5_41, L5_62)

Position 358, total mutations 2, 2 different amino acids
G->W, 1 sequences (L5_115)
G->V, 1 sequences (5)

Position 374, total mutations 2, 1 different amino acids
Q->R, 2 sequences (L5_52, L5_90)

Position 380, total mutations 2, 1 different amino acids
V->A, 2 sequences (L5_3, L5_117)

Position 388, total mutations 2, 1 different amino acids
W->R, 2 sequences (L5_23, L5_104)

Position 390, total mutations 2, 1 different amino acids
R->W, 2 sequences (11, L5_23)

Position 391, total mutations 2, 2 different amino acids
P->S, 1 sequences (L5_101)
P->L, 1 sequences (L5_2)

Position 409, total mutations 2, 1 different amino acids
C->R, 2 sequences (17, L5_1)

Position 417, total mutations 2, 1 different amino acids
A->V, 2 sequences (L5_53, L5_84)

Position 431, total mutations 2, 1 different amino acids
P->Q, 2 sequences (L5_32, L5_51)

Position 433, total mutations 2, 1 different amino acids
V->A, 2 sequences (7, L5_14)

Position 436, total mutations 2, 1 different amino acids
A->T, 2 sequences (L5_9, L5_76)

Position 450, total mutations 2, 1 different amino acids
R->H, 2 sequences (L5_41, L5_59)

Position 454, total mutations 2, 1 different amino acids
N->K, 2 sequences (L5_37, L5_68)

Position 457, total mutations 2, 2 different amino acids
M->T, 1 sequences (L5_18)
M->R, 1 sequences (L5_66)

APPENDIX 4 continued

Position 470, total mutations 2, 1 different amino acids
V->A, 2 sequences (17, L5_1)

Position 478, total mutations 2, 1 different amino acids
L->P, 2 sequences (L5_21, L5_111)

Position 491, total mutations 2, 1 different amino acids
L->P, 2 sequences (L5_76, L5_84)

Position 494, total mutations 2, 2 different amino acids
N->D, 1 sequences (L5_94)
N->K, 1 sequences (L5_95)

Position 503, total mutations 2, 1 different amino acids
H->R, 2 sequences (L5_40, L5_59)

Position 530, total mutations 2, 1 different amino acids
Q->H, 2 sequences (L5_61, L5_81)

Position 545, total mutations 2, 1 different amino acids
E->G, 2 sequences (L5_64, L5_93)

Position 559, total mutations 2, 2 different amino acids
Q->P, 1 sequences (L5_103)
Q->R, 1 sequences (L5_18)

Position 572, total mutations 2, 2 different amino acids
M->L, 1 sequences (L5_72)
M->I, 1 sequences (7)

Position 597, total mutations 2, 1 different amino acids
I->T, 2 sequences (L5_75, L5_79)

Position 616, total mutations 2, 1 different amino acids
E->K, 2 sequences (L5_75, L5_79)

Position 618, total mutations 2, 1 different amino acids
L->V, 2 sequences (18, L5_95)

Position 623, total mutations 2, 1 different amino acids
A->V, 2 sequences (12, L5_115)

Position 635, total mutations 2, 2 different amino acids
C->S, 1 sequences (8)
C->R, 1 sequences (L5_106)

Position 642, total mutations 2, 2 different amino acids
H->R, 1 sequences (L5_107)
H->Y, 1 sequences (L5_47)

Position 676, total mutations 2, 1 different amino acids
S->P, 2 sequences (L5_80, L5_82)

Position 15, total mutations 1, 1 different amino acids
K->T, 1 sequences (L5_11)

APPENDIX 4 continued

Position 23, total mutations 1, 1 different amino acids
S->P, 1 sequences (12)

Position 24, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_40)

Position 26, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_82)

Position 29, total mutations 1, 1 different amino acids
F->L, 1 sequences (11)

Position 30, total mutations 1, 1 different amino acids
P->L, 1 sequences (L5_23)

Position 36, total mutations 1, 1 different amino acids
T->I, 1 sequences (L5_99)

Position 37, total mutations 1, 1 different amino acids
G->W, 1 sequences (L5_14)

Position 41, total mutations 1, 1 different amino acids
L->R, 1 sequences (L5_40)

Position 43, total mutations 1, 1 different amino acids
V->I, 1 sequences (L5_2)

Position 51, total mutations 1, 1 different amino acids
P->S, 1 sequences (13)

Position 60, total mutations 1, 1 different amino acids
S->A, 1 sequences (L5_41)

Position 64, total mutations 1, 1 different amino acids
Y->C, 1 sequences (L5_61)

Position 67, total mutations 1, 1 different amino acids
S->P, 1 sequences (L5_73)

Position 69, total mutations 1, 1 different amino acids
E->G, 1 sequences (L5_32)

Position 70, total mutations 1, 1 different amino acids
A->V, 1 sequences (L5_88)

Position 74, total mutations 1, 1 different amino acids
I->T, 1 sequences (L5_79)

Position 77, total mutations 1, 1 different amino acids
H->R, 1 sequences (L5_48)

Position 86, total mutations 1, 1 different amino acids
I->V, 1 sequences (L5_39)

Position 87, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_20)

APPENDIX 4 continued

Position 88, total mutations 1, 1 different amino acids
V->A, 1 sequences (12)

Position 89, total mutations 1, 1 different amino acids
P->S, 1 sequences (8)

Position 90, total mutations 1, 1 different amino acids
C->Y, 1 sequences (L5_58)

Position 92, total mutations 1, 1 different amino acids
S->P, 1 sequences (L5_44)

Position 93, total mutations 1, 1 different amino acids
P->L, 1 sequences (L5_66)

Position 96, total mutations 1, 1 different amino acids
T->M, 1 sequences (L5_103)

Position 97, total mutations 1, 1 different amino acids
P->S, 1 sequences (L5_71)

Position 104, total mutations 1, 1 different amino acids
P->R, 1 sequences (L5_79)

Position 105, total mutations 1, 1 different amino acids
G->E, 1 sequences (L5_76)

Position 107, total mutations 1, 1 different amino acids
N->S, 1 sequences (L5_47)

Position 110, total mutations 1, 1 different amino acids
R->G, 1 sequences (L5_51)

Position 112, total mutations 1, 1 different amino acids
V->A, 1 sequences (L5_60)

Position 118, total mutations 1, 1 different amino acids
V->A, 1 sequences (L5_62)

Position 124, total mutations 1, 1 different amino acids
D->G, 1 sequences (L5_95)

Position 125, total mutations 1, 1 different amino acids
I->V, 1 sequences (L5_3)

Position 127, total mutations 1, 1 different amino acids
P->S, 1 sequences (L5_40)

Position 128, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_47)

Position 131, total mutations 1, 1 different amino acids
N->S, 1 sequences (21)

Position 132, total mutations 1, 1 different amino acids
P->S, 1 sequences (L5_104)

APPENDIX 4 continued

Position 136, total mutations 1, 1 different amino acids
L->W, 1 sequences (13)

Position 137, total mutations 1, 1 different amino acids
S->G, 1 sequences (L5_82)

Position 138, total mutations 1, 1 different amino acids
G->R, 1 sequences (L5_3)

Position 143, total mutations 1, 1 different amino acids
H->R, 1 sequences (L5_3)

Position 149, total mutations 1, 1 different amino acids
L->F, 1 sequences (L5_37)

Position 151, total mutations 1, 1 different amino acids
L->F, 1 sequences (L5_40)

Position 159, total mutations 1, 1 different amino acids
R->K, 1 sequences (L5_13)

Position 164, total mutations 1, 1 different amino acids
S->G, 1 sequences (L5_104)

Position 168, total mutations 1, 1 different amino acids
F->S, 1 sequences (L5_41)

Position 173, total mutations 1, 1 different amino acids
R->K, 1 sequences (L5_57)

Position 174, total mutations 1, 1 different amino acids
D->G, 1 sequences (L5_29)

Position 187, total mutations 1, 1 different amino acids
R->G, 1 sequences (L5_95)

Position 190, total mutations 1, 1 different amino acids
Q->R, 1 sequences (L5_117)

Position 192, total mutations 1, 1 different amino acids
F->L, 1 sequences (L5_42)

Position 207, total mutations 1, 1 different amino acids
L->P, 1 sequences (13)

Position 208, total mutations 1, 1 different amino acids
A->V, 1 sequences (L5_4)

Position 211, total mutations 1, 1 different amino acids
R->W, 1 sequences (L5_56)

Position 214, total mutations 1, 1 different amino acids
H->R, 1 sequences (L5_9)

Position 222, total mutations 1, 1 different amino acids
Y->C, 1 sequences (L5_9)

APPENDIX 4 continued

Position 223, total mutations 1, 1 different amino acids
V->M, 1 sequences (L5_23)

Position 225, total mutations 1, 1 different amino acids
D->G, 1 sequences (L5_4)

Position 238, total mutations 1, 1 different amino acids
Q->H, 1 sequences (L5_65)

Position 240, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_55)

Position 241, total mutations 1, 1 different amino acids
R->Q, 1 sequences (16)

Position 242, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_20)

Position 250, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_71)

Position 252, total mutations 1, 1 different amino acids
Y->H, 1 sequences (L5_120)

Position 259, total mutations 1, 1 different amino acids
A->T, 1 sequences (16)

Position 263, total mutations 1, 1 different amino acids
Q->R, 1 sequences (L5_95)

Position 280, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_60)

Position 282, total mutations 1, 1 different amino acids
E->G, 1 sequences (L5_62)

Position 292, total mutations 1, 1 different amino acids
P->L, 1 sequences (L5_56)

Position 293, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_9)

Position 295, total mutations 1, 1 different amino acids
K->E, 1 sequences (L5_107)

Position 298, total mutations 1, 1 different amino acids
R->G, 1 sequences (L5_90)

Position 308, total mutations 1, 1 different amino acids
G->S, 1 sequences (L5_120)

Position 309, total mutations 1, 1 different amino acids
F->S, 1 sequences (L5_28)

Position 311, total mutations 1, 1 different amino acids
R->H, 1 sequences (16)

APPENDIX 4 continued

Position 312, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_29)

Position 314, total mutations 1, 1 different amino acids
I->T, 1 sequences (L5_76)

Position 322, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_60)

Position 323, total mutations 1, 1 different amino acids
P->L, 1 sequences (21)

Position 326, total mutations 1, 1 different amino acids
P->S, 1 sequences (L5_80)

Position 331, total mutations 1, 1 different amino acids
G->E, 1 sequences (L5_20)

Position 332, total mutations 1, 1 different amino acids
T->I, 1 sequences (23)

Position 339, total mutations 1, 1 different amino acids
D->G, 1 sequences (L5_117)

Position 343, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_82)

Position 351, total mutations 1, 1 different amino acids
L->V, 1 sequences (L5_61)

Position 353, total mutations 1, 1 different amino acids
T->A, 1 sequences (21)

Position 356, total mutations 1, 1 different amino acids
A->G, 1 sequences (L5_82)

Position 369, total mutations 1, 1 different amino acids
F->I, 1 sequences (L5_118)

Position 376, total mutations 1, 1 different amino acids
Y->C, 1 sequences (L5_118)

Position 379, total mutations 1, 1 different amino acids
G->S, 1 sequences (L5_60)

Position 383, total mutations 1, 1 different amino acids
Q->P, 1 sequences (L5_117)

Position 392, total mutations 1, 1 different amino acids
V->A, 1 sequences (L5_57)

Position 393, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_72)

Position 415, total mutations 1, 1 different amino acids
A->V, 1 sequences (L5_9)

APPENDIX 4 continued

Position 434, total mutations 1, 1 different amino acids
I->T, 1 sequences (L5_118)

Position 435, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_23)

Position 441, total mutations 1, 1 different amino acids
E->G, 1 sequences (L5_120)

Position 444, total mutations 1, 1 different amino acids
V->A, 1 sequences (L5_9)

Position 446, total mutations 1, 1 different amino acids
Q->R, 1 sequences (L5_81)

Position 447, total mutations 1, 1 different amino acids
P->L, 1 sequences (L5_9)

Position 459, total mutations 1, 1 different amino acids
H->R, 1 sequences (L5_78)

Position 462, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_18)

Position 468, total mutations 1, 1 different amino acids
D->A, 1 sequences (L5_81)

Position 475, total mutations 1, 1 different amino acids
V->G, 1 sequences (L5_8)

Position 481, total mutations 1, 1 different amino acids
A->T, 1 sequences (10)

Position 484, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_97)

Position 486, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_56)

Position 497, total mutations 1, 1 different amino acids
D->G, 1 sequences (L5_71)

Position 498, total mutations 1, 1 different amino acids
I->V, 1 sequences (L5_97)

Position 501, total mutations 1, 1 different amino acids
E->K, 1 sequences (L5_81)

Position 504, total mutations 1, 1 different amino acids
G->R, 1 sequences (L5_18)

Position 514, total mutations 1, 1 different amino acids
L->F, 1 sequences (8)

Position 528, total mutations 1, 1 different amino acids
L->I, 1 sequences (L5_62)

APPENDIX 4 continued

Position 532, total mutations 1, 1 different amino acids
G->R, 1 sequences (L5_117)

Position 533, total mutations 1, 1 different amino acids
Q->K, 1 sequences (L5_104)

Position 538, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_69)

Position 539, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_82)

Position 543, total mutations 1, 1 different amino acids
E->K, 1 sequences (L5_2)

Position 544, total mutations 1, 1 different amino acids
T->I, 1 sequences (16)

Position 551, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_71)

Position 552, total mutations 1, 1 different amino acids
L->P, 1 sequences (L5_3)

Position 556, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_42)

Position 558, total mutations 1, 1 different amino acids
A->V, 1 sequences (L5_81)

Position 560, total mutations 1, 1 different amino acids
R->W, 1 sequences (13)

Position 562, total mutations 1, 1 different amino acids
E->K, 1 sequences (L5_71)

Position 570, total mutations 1, 1 different amino acids
L->I, 1 sequences (L5_65)

Position 573, total mutations 1, 1 different amino acids
A->T, 1 sequences (L5_76)

Position 576, total mutations 1, 1 different amino acids
K->R, 1 sequences (L5_111)

Position 577, total mutations 1, 1 different amino acids
K->Q, 1 sequences (18)

Position 600, total mutations 1, 1 different amino acids
R->K, 1 sequences (L5_16)

Position 602, total mutations 1, 1 different amino acids
G->R, 1 sequences (3)

Position 622, total mutations 1, 1 different amino acids
K->R, 1 sequences (L5_3)

APPENDIX 4 continued

Position 628, total mutations 1, 1 different amino acids
K->E, 1 sequences (L5_25)

Position 632, total mutations 1, 1 different amino acids
I->T, 1 sequences (28)

Position 633, total mutations 1, 1 different amino acids
I->T, 1 sequences (L5_107)

Position 634, total mutations 1, 1 different amino acids
H->Y, 1 sequences (L5_53)

Position 643, total mutations 1, 1 different amino acids
S->G, 1 sequences (L5_107)

Position 646, total mutations 1, 1 different amino acids
A->V, 1 sequences (L5_39)

Position 655, total mutations 1, 1 different amino acids
A->V, 1 sequences (L5_18)

Position 656, total mutations 1, 1 different amino acids
A->T, 1 sequences (16)

Position 661, total mutations 1, 1 different amino acids
I->V, 1 sequences (23)

Position 662, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_116)

Position 663, total mutations 1, 1 different amino acids
E->D, 1 sequences (8)

Position 667, total mutations 1, 1 different amino acids
T->A, 1 sequences (L5_78)

Position 668, total mutations 1, 1 different amino acids
S->P, 1 sequences (L5_3)

Position 669, total mutations 1, 1 different amino acids
T->S, 1 sequences (L5_57)

Position 670, total mutations 1, 1 different amino acids
L->F, 1 sequences (L5_106)

Position 672, total mutations 1, 1 different amino acids
I->T, 1 sequences (L5_72)

APPENDIX 5

	Selection frequency*	Conc.,	Specific activity			RNase H activity		Highest temperature of cDNA synthesis
		mg/ml	37°C (u/mg)	50°C	5 min 50°C => 37°C	u/mol	%	1kb
M-MuLV (wt)		**3.4**	**~200 000**	**45 - 50 %**	**11 %**	**~160-200**	**100 %**	**47.8°C**
D200N	25 (30)	2.91	254 000	84 %	15 %	174	97 %	50.4°C
D200A	3 (30)	2.73	187 000	87 %	18 %	179	99 %	47.8°C
D200Q		3.82	135 000	103 %	23 %	161	89 %	47.8°C
D200E		2.33	71 000	79 %	7 %			47.8°C
D200V		2.04	87 000	131 %	2 %	102	57 %	47.8°C
D200W		2.84	114 000	103 %	4.3 %			47.8°C
D200G	2 (30)	2.68	276 000	88 %	10 %			47.8°C
D200P		1.15	0	0 %	0 %			
D200K		3.14	175 000	102 %	2.5 %			47.8°C
D200R		3.69	52 000	68 %	27 %	171	95 %	45.5°C
D200H		6.23	234 000	54 %	27 %			50.4°C
L603W	17 (18)	2.41	110 000	105 %	23 %	202	112 %	53.1°C
L603F		3.18	109 000	104 %	9 %	112	62 %	50.4°C
L603Y		3.12	139 000	95 %	13 %			47.8 - 50.4°C
L603P		3.24	138 000	20 %	15 %			45.5°C
L603I		2.85	178 000	33 %	7 %	142	79 %	
L603M	1 (18)	3.7	149 000	77 %	9 %	161	89 %	47.8°C
L603V		2.69	190 000	39 %	7 %			45.5°C
L603G		3.42	149 000	36 %	10 %			
D653N	10 (23)	2.39	179 000	93 %	21 %			50.4-53.1°C
D653K		2.38	211 000	106 %	15 %			50.4-53.1°C
D653A	4 (23)	2.98	198 000	99 %	18 %			50.4°C
D653V	1 (23)	2.82	172 000	98 %	16 %			50.4°C
D653Q		2.79	182 000	93 %	18 %			50.4°C
D653L		2.60	212 000	83 %	11 %			50.4°C
D653H	3 (23)	2.41	180 000	116 %	13 %			50.4-53.1°C
D653G	5 (23)	3.05	135 000	90 %	13 %			50.4°C
D653W		2.68	178 000	93 %	13 %			50.4°C
D653E		2.46	200 000	80 %	19 %			47.8°C
T330P	15 (15)	1.76	154 000	80 %	21 %	185	103 %	47.8°C
T330N		2.62	223 000	69 %	13 %			47.8°C

APPENDIX 5 - continued

	Selection frequency*	Conc.,	Specific activity			RNase H activity		Highest temperature of cDNA synthesis
		mg/ml	37°C (u/mg)	50°C	5 min 50°C => 37°C	u/mol	%	1kb
M-MuLV (wt)		**3.4**	**~200 000**	**45 - 50 %**	**11 %**	**~160-200**	**100 %**	**47.8°C**
T330L		3.26	128 000	49 %	9 %			45.5°C
T330D		3.32	240 000	55 %	16 %			47.8°C
T330V		2.56	197 000	65 %	12 %			47.8°C
T330S		2.33	201 000	67 %	15 %			47.8°C
Q221R	6 (6)	2.38	268 000	94 %	0 %	117	65 %	50.4°C
Q221N		3.09	114 000	53 %	10 %			45.5°C
Q221H		2.47	55 000	44 %	8 %			43.6-45.5°C
Q221K		1.85	179 000	77 %	0 %			45.5
Q221E		1.85	83 000	64 %	0 %			43.6-45.5°C
Q221M		1.3	107 000	58 %	2 %			45.5
Q221Y		1.76	60 000	77 %	0 %			43.6-45.5°C
E607K	2 (4)	3.94	182 000	84 %	9 %	260	144 %	47.8-50.4°C
E607A	1 (4)	4.28	204 000	98 %				47.8°C
E607G	1 (4)	4.02	184 000	72 %				47.8°C
E607D		2.24	181 000	69 %				45.5°C
E607Q		1.02	0	0 %				
L139P	14 (14)	1.34	148 000	59 %	7 %	176	98 %	47.8°C
T287A	6 (6)	2.34	204 000	52 %	13 %	174	96 %	45.5-47.8°C
T287S		2.75	213 000	68 %	11 %			45.5-47.8°C
T287L		2.28	179 000	38 %	11 %			45.5°C
T287N		2.46	197 000	50 %	12 %			45.5°C
T287F		2.18	190 000	48 %	13 %			45.5°C
T287P		1.64	134 000	7 %	11 %			41.9°C
N479D	5 (5)	3.15	190 000	81 %	9 %	234	130 %	47.8°C
N479Q		2.82	176 000	33 %				43.6°C
N479E		2.58	179 000	48 %				45.5°C
H594R	4 (5)	2.67	184 000	69 %	14 %	232	129 %	47.8°C
H594K		3.66	270 000	80 %	9 %	170	94 %	47.8-50.4°C
H594Q	1 (5)	3.85	231 000	75 %	13 %			47.8-50.4°C
H594N		4.15	216 000	61 %	11 %			47.8°C
D449G	2 (3)	3.53	123 000	79 %	13 %	162	90 %	45.5-47.8°C

	Selection frequency*	Conc., mg/ml	Specific activity 37°C (u/mg)	50°C	5 min 50°C => 37°C	RNase H activity u/mol	%	Highest temperature of cDNA synthesis 1kb
M-MuLV (wt)		**3.4**	**~200 000**	**45 - 50 %**	**11 %**	**~160-200**	**100 %**	**47.8°C**
D449E		3.25	224 000	77 %				45.5°C
D449N		3.25	221 000	75 %				45.5°C
D449A	1 (3)	4.15	212 000	99 %				45.5-47.8°C
D449V		4.45	216 000	83 %				45.5°C
M39V	1 (2)	3.09	146 000	54 %	13 %	150	83 %	45.5-47.8°C
M39L	1 (2)	2.24	212 000	47 %	9 %			45.5°C
M39F		2.51	141 000	43 %	8 %			45.5°C
M39N		2.38	349 000	71 %	0 %			45.5-47.8°C
M66L	1 (2)	3.35	237 000	79 %	13 %	215	119 %	47.8°C
M66V	1 (2)	2.75	227 000	73 %				45.5-47.8°C
M66I		3.27	240 000	80 %				45.5-47.8°C
L333Q	3 (4)	3.34	177 000	54 %	9 %	233	129 %	47.8°C
H126R	2 (4)	3.58	227 000	58 %	4 %	218	121 %	47.8°C
Y344H	3 (3)	3.29	85 000	30 %	13 %	117	65 %	45.5°C
P130S	3 (3)	2.59	120 000	70 %	10 %	156	87 %	47.8°C
Q91R	1 (2)	2.44	160 000	56 %	13 %	162	90 %	47.8°C
N649S	3 (3)	3.47	157 000	45 %	16 %			47.8°C
W388R	2 (2)	1.75	266 000	72 %	14 %			45.5-47.8°C
R390W	2 (2)	1.98	161 000	64 %	10 %			45.5-47.8°C
I179V	1 (2)	5.44	251 000	52 %				48°C
Q374R	2 (2)	3.17	135 000	56 %	9 %			
E5K	2 (2)	3.2	160 000	67 %	5 %			

APPENDIX 6

	Conc.,	Specific activity		Highest temperature of cDNA synthesis	
	mg/ml	37°C (u/mg)	50°C	1kb	4.5kb
M-MuLV (wt)	**3.4**	**~200 000**	**45 - 50 %**	**47.8°C**	
D200N, L603W (20 mut.)	0.46	260 000	131 %	56°C	
D200N, L603W, T330P (M0_1)	4,17	321 194	175 %	56-58°C	53°C
D200N, L603W, T330P, E607K (M1)	4,27	311 576	174 %	60-62°C	56°C
D200N, L603W, T330P, E607K, L139P (M2)	2,91	348 694	176 %	62°C	61°C
D200N, L603W, N479D, H594R (M3)	5,37	371 510	182 %	56-58°C	56°C
D200N, L603W, D653N, D524G (M4)	4,84	358 112	155 %	58-60°C	58°C
D200N, L603W, D653N, D524G, T330P (M6)	5,23	272 031	180 %	60-62°C	58°C

APPENDIX 7

SEQ ID No: 1

```
LOCUS       pET-his-MLV-pD          7873 bp    DNA     circular      5-JUN-
2007
SOURCE
  ORGANISM
COMMENT     This file is created by Vector NTI
            http://www.invitrogen.com/
COMMENT     ORIGDB|GenBank
COMMENT     VNTDATE|448383004|
COMMENT     VNTDBDATE|448383004|
COMMENT     LSOWNER|
COMMENT     VNTNAME|pET-his-MLV-pD|
COMMENT     VNTAUTHORNAME|Remigijus Skirgaila|
COMMENT     VNTAUTHORTEL|+370-5-2394224|
COMMENT     VNTAUTHOREML|skirgaila@fermentas.lt|
COMMENT     VNTAUTHORWWW|www.fermentas.com|
FEATURES             Location/Qualifiers
     CDS             3534..4391
                     /vntifkey="4"
                     /label=Ap
                     /note="ORF: Frame #2   Start: atg   Stop:  taa"
     CDS             complement(6586..7674)
                     /vntifkey="4"
                     /label=lacI
                     /note="ORF: Frame #3   Start: gtg   Stop:  tga"
     terminator      2864..2910
                     /vntifkey="43"
                     /label=T7\terminator
     rep_origin      2947..3402
                     /vntifkey="33"
                     /label=f1\origin
     rep_origin      complement(2936..2936)
                     /vntifkey="33"
                     /label=ori
     promoter        174..190
                     /vntifkey="30"
                     /label=PT7
     misc_feature    2394..2669
                     /vntifkey="21"
                     /label=pD
     misc_feature    2364..2393
                     /vntifkey="21"
                     /label=gs\linker
     misc_feature    2670..2759
                     /vntifkey="21"
                     /label=gs\linker
     misc_feature    306..2363
                     /vntifkey="21"
                     /label=MLV\H+
     misc_feature    258..305
                     /vntifkey="21"
                     /label=his
BASE COUNT     1873 a      2155 c      2084 g      1761 t
ORIGIN
        1 cacggggcct gccaccatac ccacgccgaa acaagcgctc atgagcccga agtggcgagc
```

```
  61 ccgatcttcc ccatcggtga tgtcggcgat ataggcgcca gcaaccgcac ctgtggcgcc
 121 ggtgatgccg gccacgatgc gtccggcgta gaggatcgag atctatacga aattaatacg
 181 actcactata gggagaccac aacggtttcc ctctagaaat aattttgttt aactttaaga
 241 aagaggagaa attacatatg agaggatcgc atcaccatca ccatcacgga tctggttcca
 301 tgggcatgac cctaaatata gaagatgagc atcggctaca tgagacctca aaagagccag
 361 atgtttctct agggtccaca tggctgtctg attttcctca ggcctgggcg gaaaccgggg
 421 gcatgggact ggcagttcgc caagctcctc tgatcatacc tctgaaagca acctctaccc
 481 ccgtgtccat aaaacaatac cccatgtcac aagaagccag actggggatc aagccccaca
 541 tacagagact gttggaccag ggaatactgg taccctgcca gtccccctgg aacacgcccc
 601 tgctacccgt taagaaacca gggactaatg attataggcc tgtccaggat ctgagagaag
 661 tcaacaagcg ggtggaagac atccacccca ccgtgcccaa cccttacaac ctcttgagcg
 721 ggctcccacc gtcccaccag tggtacactg tgcttgattt aaaggatgcc tttttctgcc
 781 tgagactcca ccccaccagt cagcctctct tcgcctttga gtggagagat ccagagatgg
 841 gaatctcagg acaattgacc tggaccagac tcccacaggg tttcaaaaac agtcccaccc
 901 tgtttgatga ggcactgcac agagacctag cagacttccg gatccagcac ccagacttga
 961 tcctgctaca gtacgtggat gacttactgc tggccgccac ttctgagcta gactgccaac
1021 aaggtactcg ggccctgtta caaaccctag ggaacctcgg gtatcgggcc tcggccaaga
1081 aagcccaaat ttgccagaaa caggtcaagt atctggggta tcttctaaaa gagggtcaga
1141 gatggctgac tgaggccaga aaagagactg tgatggggca gcctactccg aagacccctc
1201 gacaactaag ggagttccta gggacggcag gcttctgtcg cctctggatc cctgggtttg
1261 cagaaatggc agcccccttg taccctctca ccaaaacggg gactctgttt aattggggcc
1321 cagaccaaca aaaggcctat caagaaatca agcaagctct tctaactgcc ccagccctgg
1381 ggttgccaga tttgactaag ccctttgaac tctttgtcga cgagaagcag ggctacgcca
1441 aaggtgtcct aacgcaaaaa ctgggacctt ggcgtcggcc ggtggcctac ctgtccaaaa
1501 agctagaccc agtagcagct gggtggcccc cttgcctacg gatggtagca gccattgccg
1561 tactgacaaa ggatgcaggc aagctaacca tgggacagcc actagtcatt ctggcccccc
1621 atgcagtaga ggcactagtc aaacaacccc ccgaccgctg gctttccaac gcccggatga
1681 ctcactatca ggccttgctt ttggacacgg accgggtcca gttcggaccg gtggtagccc
1741 tgaacccggc tacgctgctc ccactgcctg aggaagggct gcaacacaac tgccttgata
1801 tcctggccga agcccacgga acccgacccg acctaacgga ccagccgctc ccagacgccg
1861 accacacctg gtacacggat ggaagcagtc tcttacaaga gggacagcgt aaggcgggag
1921 ctgcggtgac caccgagacc gaggtaatct gggctaaagc cctgccagcc gggacatccg
1981 ctcagcgggc tgaactgata gcactcaccc aggccctaaa gatggcagaa ggtaagaagc
2041 taaatgttta tactgatagc cgttatgctt ttgctactgc ccatatccat ggagaaatat
2101 acagaaggcg tgggttgctc acatcagaag gcaaagagat caaaaataaa gacgagatct
2161 tggccctact aaaagccctc tttctgccca aaagacttag cataatccat tgtccaggac
2221 atcaaaaggg acacagcgcc gaggctagag gcaaccggat ggctgaccaa gcggcccgaa
2281 aggcagccat cacagagact ccagacacct ctaccctcct catagaaaat tcatcaccca
2341 attcccgctt aattaatgaa ttcggatccg gtggcggttc cggcggtgga tctatgggta
2401 ccgcaaccgc gcccggcgga ttgagtgcga aagcgcctgc aatgaccccg ctgatgctgg
2461 acacctccag ccgtaagctg gttgcgtggg atggcaccac cgacggtgct gccgttggca
2521 ttcttgcggt tgctgctgac cagaccagca ccacgctgac gttctacaag tccggcacgt
2581 tccgttatga ggatgtgctc tggccggagg ctgccagcga cgagacgaaa aaacggaccg
2641 cgtttgccgg aacggcaatc agcatcgttg gatctggtgg cggttccggc ggtggatctg
2701 gtggcggttc cggcggtgga tctggtggcg gttccggcgg tggatcgtgt cttctttaag
2761 cttgcggccg cactcgagca ccaccaccac caccactgag atccggctgc taacaaagcc
2821 cgaaaggaag ctgagttggc tgctgccacc gctgagcaat aactagcata acccccttggg
2881 gcctctaaac gggtcttgag gggtttttg ctgaaaggag gaactatatc cggattggcg
2941 aatgggacgc gccctgtagc ggcgcattaa gcgcggcggg tgtggtggtt acgcgcagcg
3001 tgaccgctac acttgccagc gccctagcgc ccgctccttt cgctttcttc ccttcctttc
3061 tcgccacgtt cgccggcttt ccccgtcaag ctctaaatcg ggggctccct ttagggttcc
3121 gatttagtgc tttacggcac ctcgacccca aaaaacttga ttagggtgat ggttcacgta
3181 gtgggccatc gccctgatag acggtttttc gccctttgac gttggagtcc acgttcttta
3241 atagtggact cttgttccaa actggaacaa cactcaaccc tatctcggtc tattcttttg
3301 atttataagg gattttgccg atttcggcct attggttaaa aaatgagctg atttaacaaa
3361 aatttaacgc gaattttaac aaaatattaa cgtttacaat ttcaggtggc acttttcggg
3421 gaaatgtgcg cggaacccct atttgtttat ttttctaaat acattcaaat atgtatccgc
3481 tcatgagaca ataaccctga taaatgcttc aataatattg aaaaaggaag agtatgagta
3541 ttcaacattt ccgtgtcgcc cttattccct tttttgcggc attttgcctt cctgtttttg
```

3601 ctcacccaga aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtgg
3661 gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac
3721 gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccgtattg
3781 acgccgggca agagcaactc ggtcgccgca tacactattc tcagaatgac ttggttgagt
3841 actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg
3901 ctgccataac catgagtgat aacactgcgg ccaacttact tctgacaacg atcggaggac
3961 cgaaggagct aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt
4021 gggaaccgga gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgcctgcag
4081 caatggcaac aacgttgcgc aaactattaa ctggcgaact acttactcta gcttcccggc
4141 aacaattaat agactggatg gaggcggata aagttgcagg accacttctg cgctcggccc
4201 ttccggctgg ctggtttatt gctgataaat ctggagccgg tgagcgtggg tctcgcggta
4261 tcattgcagc actggggcca gatggtaagc cctcccgtat cgtagttatc tacacgacgg
4321 ggagtcaggc aactatggat gaacgaaata gacagatcgc tgagataggt gcctcactga
4381 ttaagcattg gtaactgtca gaccaagttt actcatatat actttagatt gatttaaaac
4441 ttcatttta atttaaaagg atctaggtga agatcctttt tgataatctc atgaccaaaa
4501 tcccttaacg tgagttttcg ttccactgag cgtcagaccc cgtagaaaag atcaaaggat
4561 cttcttgaga tccttttttt ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc
4621 taccagcggt ggtttgtttg ccggatcaag agctaccaac tctttttccg aaggtaactg
4681 gcttcagcag agcgcagata ccaaatactg tccttctagt gtagccgtag ttaggccacc
4741 acttcaagaa ctctgtagca ccgcctacat acctcgctct gctaatcctg ttaccagtgg
4801 ctgctgccag tggcgataag tcgtgtctta ccgggttgga ctcaagacga tagttaccgg
4861 ataaggcgca gcggtcgggc tgaacggggg gttcgtgcac acagcccagc ttggagcgaa
4921 cgacctacac cgaactgaga tacctacagc gtgagctatg agaaagcgcc acgcttcccg
4981 aagggagaaa ggcggacagg tatccggtaa gcggcagggt cggaacagga gagcgcacga
5041 gggagcttcc agggggaaac gcctggtatc tttatagtcc tgtcgggttt cgccacctct
5101 gacttgagcg tcgatttttg tgatgctcgt caggggggcg gagcctatgg aaaaacgcca
5161 gcaacgcggc ctttttacgg ttcctggcct tttgctggcc ttttgctcac atgttctttc
5221 ctgcgttatc ccctgattct gtggataacc gtattaccgc ctttgagtga gctgataccg
5281 ctcgccgcag ccgaacgacc gagcgcagcg agtcagtgag cgaggaagcg gaagagcgcc
5341 tgatgcggta ttttctcctt acgcatctgt gcggtatttc acaccgcata tatggtgcac
5401 tctcagtaca atctgctctg atgccgcata gttaagccag tatacactcc gctatcgcta
5461 cgtgactggg tcatggctgc gccccgacac ccgccaacac ccgctgacgc gccctgacgg
5521 gcttgtctgc tcccggcatc cgcttacaga caagctgtga ccgtctccgg gagctgcatg
5581 tgtcagaggt tttcaccgtc atcaccgaaa cgcgcgaggc agctgcggta aagctcatca
5641 gcgtggtcgt gaagcgattc acagatgtct gcctgttcat ccgcgtccag ctcgttgagt
5701 ttctccagaa gcgttaatgt ctggcttctg ataaagcggg ccatgttaag ggcggttttt
5761 tcctgtttgg tcactgatgc ctccgtgtaa gggggatttc tgttcatggg ggtaatgata
5821 ccgatgaaac gagagaggat gctcacgata cgggttactg atgatgaaca tgcccggtta
5881 ctggaacgtt gtgagggtaa acaactggcg gtatggatgc ggcgggacca gagaaaaatc
5941 actcagggtc aatgccagcg cttcgttaat acagatgtag gtgttccaca gggtagccag
6001 cagcatcctg cgatgcagat ccggaacata atggtgcagg gcgctgactt ccgcgtttcc
6061 agactttacg aaacacggaa accgaagacc attcatgttg ttgctcaggt cgcagacgtt
6121 ttgcagcagc agtcgcttca cgttcgctcg cgtatcggtg attcattctg ctaaccagta
6181 aggcaacccc gccagcctag ccgggtcctc aacgacagga gcacgatcat gcgcacccgt
6241 ggggccgcca tgccggcgat aatggcctgc ttctcgccga aacgtttggt ggcgggacca
6301 gtgacgaagg cttgagcgag ggcgtgcaag attccgaata ccgcaagcga caggccgatc
6361 atcgtcgcgc tccagcgaaa gcggtcctcg ccgaaaatga cccagagcgc tgccggcacc
6421 tgtcctacga gttgcatgat aaagaagaca gtcataagtg cggcgacgat agtcatgccc
6481 cgcgcccacc ggaaggagct gactgggttg aaggctctca agggcatcgg tcgagatccc
6541 ggtgcctaat gagtgagcta acttacatta attgcgttgc gctcactgcc cgctttccag
6601 tcgggaaacc tgtcgtgcca gctgcattaa tgaatcggcc aacgcgcggg gagaggcggt
6661 ttgcgtattg ggcgccaggg tggtttttct tttcaccagt gagacgggca acagctgatt
6721 gcccttcacc gcctggccct gagagagttg cagcaagcgg tccacgctgg tttgccccag
6781 caggcgaaaa tcctgtttga tggtggttaa cggcgggata taacatgagc tgtcttcggt
6841 atcgtcgtat cccactaccg agatatccgc accaacgcgc agcccggact cggtaatggc
6901 gcgcattgcg cccagcgcca tctgatcgtt ggcaaccagc atcgcagtgg gaacgatgcc
6961 ctcattcagc atttgcatgg tttgttgaaa accggacatg gcactccagt cgccttcccg
7021 ttccgctatc ggctgaattt gattgcgagt gagatattta tgccagccag ccagacgcag
7081 acgcgccgag acagaactta atgggcccgc taacagcgcg atttgctggt gacccaatgc

```
7141 gaccagatgc tccacgccca gtcgcgtacc gtcttcatgg gagaaaataa tactgttgat
7201 gggtgtctgg tcagagacat caagaaataa cgccggaaca ttagtgcagg cagcttccac
7261 agcaatggca tcctggtcat ccagcggata gttaatgatc agcccactga cgcgttgcgc
7321 gagaagattg tgcaccgccg ctttacaggc ttcgacgccg cttcgttcta ccatcgacac
7381 caccacgctg gcacccagtt gatcggcgcg agatttaatc gccgcgacaa tttgcgacgg
7441 cgcgtgcagg gccagactgg aggtggcaac gccaatcagc aacgactgtt tgcccgccag
7501 ttgttgtgcc acgcggttgg gaatgtaatt cagctccgcc atcgccgctt ccacttttte
7561 ccgcgttttc gcagaaacgt ggctggcctg gttcaccacg cgggaaacgg tctgataaga
7621 gacaccggca tactctgcga catcgtataa cgttactggt ttcacattca ccaccctgaa
7681 ttgactctct tccgggcgct atcatgccat accgcgaaag gttttgcgcc attcgatggt
7741 gtccgggatc tcgacgctct cccttatgcg actcctgcat taggaagcag cccagtagta
7801 ggttgaggcc gttgagcacc gccgccgcaa ggaatggtgc atgcaaggag atggcgccca
7861 acagtccccc ggc
```

SEQ ID No: 2

```
LOCUS       pET_his_del_pD          7702 bp    DNA     circular      6-JUN-
2007
SOURCE
  ORGANISM
COMMENT     This file is created by Vector NTI
            http://www.invitrogen.com/
COMMENT     VNTDATE|448455242|
COMMENT     VNTDBDATE|448455855|
COMMENT     LSOWNER|
COMMENT     VNTNAME|pET_his_del_pD|
COMMENT     VNTAUTHORNAME|Remigijus Skirgaila|
COMMENT     VNTAUTHORTEL|+370-5-2394224|
COMMENT     VNTAUTHOREML|skirgaila@fermentas.lt|
COMMENT     VNTAUTHORWWW|www.fermentas.com|
FEATURES             Location/Qualifiers
     misc_feature    2499..2588
                     /vntifkey="21"
                     /label=gs\linker
     misc_feature    2193..2222
                     /vntifkey="21"
                     /label=gs\linker
     misc_feature    2223..2498
                     /vntifkey="21"
                     /label=pD
     promoter        174..190
                     /vntifkey="30"
                     /label=PT7
     rep_origin      complement(2765..2765)
                     /vntifkey="33"
                     /label=ori
     rep_origin      2776..3231
                     /vntifkey="33"
                     /label=f1\origin
     terminator      2693..2739
                     /vntifkey="43"
                     /label=T7\terminator
     CDS             complement(6415..7503)
                     /vntifkey="4"
                     /label=lacI
                     /note="ORF: Frame #3   Start: gtg   Stop:  tga"
     CDS             3363..4220
```

```
                     /vntifkey="4"
                     /label=Ap
                     /note="ORF: Frame #2   Start: atg   Stop:  taa"
     misc_feature    258..305
                     /vntifkey="21"
                     /label=his
     misc_feature    306..2192
                     /vntifkey="21"
                     /label=del
BASE COUNT     1823 a      2115 c      2033 g      1731 t
ORIGIN
        1 cacggggcct gccaccatac ccacgccgaa acaagcgctc atgagcccga agtggcgagc
       61 ccgatcttcc ccatcggtga tgtcggcgat ataggcgcca gcaaccgcac ctgtggcgcc
      121 ggtgatgccg gccacgatgc gtccggcgta gaggatcgag atctatacga aattaatacg
      181 actcactata gggagaccac aacggtttcc ctctagaaat aattttgttt aactttaaga
      241 aagaggagaa attacatatg agaggatcgc atcaccatca ccatcacgga tctggttcca
      301 tgggcatgac cctaaatata gaagatgagc atcggctaca tgagacctca aaagagccag
      361 atgtttctct agggtccaca tggctgtctg attttcctca ggcctgggcg gaaaccgggg
      421 gcatgggact ggcagttcgc caagctcctc tgatcatacc tctgaaagca acctctaccc
      481 ccgtgtccat aaaacaatac cccatgtcac aagaagccag actggggatc aagccccaca
      541 tacagagact gttggaccag ggaatactgg taccctgcca gtccccctgg aacacgcccc
      601 tgctacccgt taagaaacca gggactaatg attataggcc tgtccaggat ctgagagaag
      661 tcaacaagcg ggtggaagac atccacccca ccgtgcccaa cccttacaac ctcttgagcg
      721 ggctcccacc gtcccaccag tggtacactg tgcttgattt aaaggatgcc tttttctgcc
      781 tgagactcca ccccaccagt cagcctctct tcgcctttga gtggagagat ccagagatgg
      841 gaatctcagg acaattgacc tggaccagac tcccacaggg tttcaaaaac agtcccaccc
      901 tgtttgatga ggcactgcac agagacctag cagacttccg gatccagcac ccagacttga
      961 tcctgctaca gtacgtggat gacttactgc tggccgccac ttctgagcta gactgccaac
     1021 aaggtactcg ggccctgtta caaaccctag ggacggcagg cttctgtcgc ctctggatcc
     1081 ctgggtttgc agaaatggca gccccccttgt accctctcac caaaacgggg actctgttta
     1141 attggggccc agaccaacaa aaggcctatc aagaaatcaa gcaagctctt ctaactgccc
     1201 cagccctggg gttgccagat ttgactaagc cctttgaact ctttgtcgac gagaagcagg
     1261 gctacgccaa aggtgtccta acgcaaaaac tgggaccttg gcgtcggccg gtggcctacc
     1321 tgtccaaaaa gctagaccca gtagcagctg ggtggccccc ttgcctacgg atggtagcag
     1381 ccattgccgt actgacaaag gatgcaggca agctaaccat gggacagcca ctagtcattc
     1441 tggcccccca tgcagtagag gcactagtca aacaaccccc cgaccgctgg ctttccaacg
     1501 cccggatgac tcactatcag gccttgcttt tggacacgga ccgggtccag ttcggaccgg
     1561 tggtagccct gaacccggct acgctgctcc cactgcctga ggaagggctg caacacaact
     1621 gccttgatat cctggccgaa gcccacggaa cccgacccga cctaacggac cagccgctcc
     1681 cagacgccga ccacacctgg tacacggatg gaagcagtct cttacaagag ggacagcgta
     1741 aggcgggagc tgcggtgacc accgagaccg aggtaatctg ggctaaagcc ctgccagccg
     1801 ggacatccgc tcagcgggct gaactgatag cactcaccca ggccctaaag atggcagaag
     1861 gtaagaagct aaatgtttat actaatagcc gttatgcttt tgctactgcc catatccatg
     1921 gagaaatata cagaaggcgt gggttgctca catcagaagg caaagagatc aaaaataaag
     1981 acgagatctt ggccctacta aaagccctct ttctgcccaa aagacttagc ataatccatt
     2041 gtccaggaca tcaaaaggga cacagcgccg aggctagagg caaccggatg gctgaccaag
     2101 cggcccgaaa ggcagccatc acagagactc cagacacctc taccctcctc atagaaaatt
     2161 catcacccaa ttcccgctta attaatgaat tcggatccgg tggcggttcc ggcggtggat
     2221 ctatgggtac cgcaaccgcg cccggcggat tgagtgcgaa agcgcctgca atgaccccgc
     2281 tgatgctgga cacctccagc cgtaagctgg ttgcgtggga tggcaccacc gacggtgctg
     2341 ccgttggcat tcttgcggtt gctgctgacc agaccagcac cacgctgacg ttctacaagt
     2401 ccggcacgtt ccgttatgag gatgtgctct ggccggaggc tgccagcgac gagacgaaaa
     2461 aacggaccgc gtttgccgga acggcaatca gcatcgttgg atctggtggc ggttccggcg
     2521 gtggatctgg tggcggttcc ggcggtggat ctggtggcgg ttccggcggt ggatcgtgtc
     2581 ttctttaagc ttgcggccgc actcgagcac caccaccacc accactgaga tccggctgct
     2641 aacaaagccc gaaaggaagc tgagttggct gctgccaccg ctgagcaata actagcataa
     2701 ccccttgggg cctctaaacg ggtcttgagg ggtttttttgc tgaaaggagg aactatatcc
     2761 ggattggcga atgggacgcg ccctgtagcg gcgcattaag cgcggcgggt gtggtggtta
     2821 cgcgcagcgt gaccgctaca cttgccagcg ccctagcgcc cgctcctttc gctttcttcc
```

2881 cttccttttct cgccacgttc gccggctttc cccgtcaagc tctaaatcgg gggctcccttt
2941 tagggttccg atttagtgct ttacggcacc tcgaccccaa aaaacttgat tagggtgatg
3001 gttcacgtag tgggccatcg ccctgataga cggtttttcg ccctttgacg ttggagtcca
3061 cgttctttaa tagtggactc ttgttccaaa ctggaacaac actcaaccct atctcggtct
3121 attcttttga tttataaggg attttgccga tttcggccta ttggttaaaa aatgagctga
3181 tttaacaaaa atttaacgcg aattttaaca aaatattaac gtttacaatt tcaggtggca
3241 cttttcgggg aaatgtgcgc ggaaccccta tttgtttatt tttctaaata cattcaaata
3301 tgtatccgct catgagacaa taaccctgat aaatgcttca ataatattga aaaaggaaga
3361 gtatgagtat tcaacatttc cgtgtcgccc ttattcccttt ttttgcggca ttttgccttc
3421 ctgtttttgc tcacccagaa acgctggtga aagtaaaaga tgctgaagat cagttgggtg
3481 cacgagtggg ttacatcgaa ctggatctca acagcggtaa gatccttgag agttttcgcc
3541 ccgaagaacg ttttccaatg atgagcactt ttaaagttct gctatgtggc gcggtattat
3601 cccgtattga cgccgggcaa gagcaactcg gtcgccgcat acactattct cagaatgact
3661 tggttgagta ctcaccagtc acagaaaagc atcttacgga tggcatgaca gtaagagaat
3721 tatgcagtgc tgccataacc atgagtgata acactgcggc caacttactt ctgacaacga
3781 tcggaggacc gaaggagcta accgcttttt tgcacaacat gggggatcat gtaactcgcc
3841 ttgatcgttg ggaaccggag ctgaatgaag ccataccaaa cgacgagcgt gacaccacga
3901 tgcctgcagc aatggcaaca acgttgcgca aactattaac tggcgaacta cttactctag
3961 cttcccggca acaattaata gactggatgg aggcggataa agttgcagga ccacttctgc
4021 gctcggccct tccggctggc tggtttattg ctgataaatc tggagccggt gagcgtgggt
4081 ctcgcggtat cattgcagca ctggggccag atggtaagcc ctcccgtatc gtagttatct
4141 acacgacggg gagtcaggca actatggatg aacgaaatag acagatcgct gagataggtg
4201 cctcactgat taagcattgg taactgtcag accaagttta ctcatatata ctttagattg
4261 atttaaaact tcatttttaa tttaaaagga tctaggtgaa gatccttttt gataatctca
4321 tgaccaaaat cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga
4381 tcaaaggatc ttcttgagat cctttttttc tgcgcgtaat ctgctgcttg caaacaaaaa
4441 aaccaccgct accagcggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga
4501 aggtaactgg cttcagcaga gcgcagatac caaatactgt ccttctagtg tagccgtagt
4561 taggccacca cttcaagaac tctgtagcac cgcctacata cctcgctctg ctaatcctgt
4621 taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttggac tcaagacgat
4681 agttaccgga taaggcgcag cggtcgggct gaacgggggg ttcgtgcaca cagcccagct
4741 tggagcgaac gacctacacc gaactgagat acctacagcg tgagctatga gaaagcgcca
4801 cgcttcccga agggagaaag gcggacaggt atccggtaag cggcagggtc ggaacaggag
4861 agcgcacgag ggagcttcca gggggaaacg cctggtatct ttatagtcct gtcgggtttc
4921 gccacctctg acttgagcgt cgatttttgt gatgctcgtc aggggggcgg agcctatgga
4981 aaaacgccag caacgcggcc tttttacggt tcctggcctt ttgctggcct tttgctcaca
5041 tgttctttcc tgcgttatcc cctgattctg tggataaccg tattaccgcc tttgagtgag
5101 ctgataccgc tcgccgcagc cgaacgaccg agcgcagcga gtcagtgagc gaggaagcgg
5161 aagagcgcct gatgcggtat tttctcctta cgcatctgtg cggtatttca caccgcatat
5221 atggtgcact ctcagtacaa tctgctctga tgccgcatag ttaagccagt atacactccg
5281 ctatcgctac gtgactgggt catggctgcg ccccgacacc cgccaacacc cgctgacgcg
5341 ccctgacggg cttgtctgct cccggcatcc gcttacagac aagctgtgac cgtctccggg
5401 agctgcatgt gtcagaggtt ttcaccgtca tcaccgaaac gcgcgaggca gctgcggtaa
5461 agctcatcag cgtggtcgtg aagcgattca cagatgtctg cctgttcatc cgcgtccagc
5521 tcgttgagtt tctccagaag cgttaatgtc tggcttctga taaagcgggc catgttaagg
5581 gcggtttttt cctgtttggt cactgatgcc tccgtgtaag ggggatttct gttcatgggg
5641 gtaatgatac cgatgaaacg agagaggatg ctcacgatac gggttactga tgatgaacat
5701 gcccggttac tggaacgttg tgagggtaaa caactggcgg tatggatgcg gcgggaccag
5761 agaaaaatca ctcagggtca atgccagcgc ttcgttaata cagatgtagg tgttccacag
5821 ggtagccagc agcatcctgc gatgcagatc cggaacataa tggtgcaggg cgctgacttc
5881 cgcgtttcca gactttacga aacacggaaa ccgaagacca ttcatgttgt tgctcaggtc
5941 gcagacgttt tgcagcagca gtcgcttcac gttcgctcgc gtatcggtga ttcattctgc
6001 taaccagtaa ggcaaccccg ccagcctagc cgggtcctca acgacaggag cacgatcatg
6061 cgcacccgtg gggccgccat gccggcgata atggcctgct tctcgccgaa acgtttggtg
6121 gcgggaccag tgacgaaggc ttgagcgagg gcgtgcaaga ttccgaatac cgcaagcgac
6181 aggccgatca tcgtcgcgct ccagcgaaag cggtcctcgc cgaaaatgac ccagagcgct
6241 gccggcacct gtcctacgag ttgcatgata aagaagacag tcataagtgc ggcgacgata
6301 gtcatgcccc gcgcccaccg gaaggagctg actgggttga aggctctcaa gggcatcggt
6361 cgagatcccg gtgcctaatg agtgagctaa cttacattaa ttgcgttgcg ctcactgccc

```
6421 gctttccagt cgggaaacct gtcgtgccag ctgcattaat gaatcggcca acgcgcgggg
6481 agaggcggtt tgcgtattgg gcgccagggt ggtttttctt ttcaccagtg agacgggcaa
6541 cagctgattg cccttcaccg cctggccctg agagagttgc agcaagcggt ccacgctggt
6601 ttgccccagc aggcgaaaat cctgtttgat ggtggttaac ggcgggatat aacatgagct
6661 gtcttcggta tcgtcgtatc ccactaccga gatatccgca ccaacgcgca gcccggactc
6721 ggtaatggcg cgcattgcgc ccagcgccat ctgatcgttg gcaaccagca tcgcagtggg
6781 aacgatgccc tcattcagca tttgcatggt ttgttgaaaa ccggacatgg cactccagtc
6841 gccttcccgt tccgctatcg gctgaatttg attgcgagtg agatatttat gccagccagc
6901 cagacgcaga cgcgccgaga cagaacttaa tgggcccgct aacagcgcga tttgctggtg
6961 acccaatgcg accagatgct ccacgcccag tcgcgtaccg tcttcatggg agaaaataat
7021 actgttgatg ggtgtctggt cagagacatc aagaaataac gccggaacat tagtgcaggc
7081 agcttccaca gcaatggcat cctggtcatc cagcggatag ttaatgatca gcccactgac
7141 gcgttgcgcg agaagattgt gcaccgccgc tttacaggct tcgacgccgc ttcgttctac
7201 catcgacacc accacgctgg cacccagttg atcggcgcga gatttaatcg ccgcgacaat
7261 ttgcgacggc gcgtgcaggg ccagactgga ggtggcaacg ccaatcagca acgactgttt
7321 gcccgccagt tgttgtgcca cgcggttggg aatgtaattc agctccgcca tcgccgcttc
7381 cacttttcc cgcgttttcg cagaaacgtg gctggcctgg ttcaccacgc gggaaacggt
7441 ctgataagag acaccggcat actctgcgac atcgtataac gttactggtt tcacattcac
7501 caccctgaat tgactctctt ccgggcgcta tcatgccata ccgcgaaagg ttttgcgcca
7561 ttcgatggtg tccgggatct cgacgctctc ccttatgcga ctcctgcatt aggaagcagc
7621 ccagtagtag gttgaggccg ttgagcaccg ccgccgcaag gaatggtgca tgcaaggaga
7681 tggcgcccaa cagtcccccg gc
```

SEQ ID No: 3

oligonucleotide (22 b)
pro-pIVEX
5′-GCGAGCCCGATCTTCCCCATCG-3′

SEQ ID No: 4

oligonucleotide (21 b)
pD-ter
5′-AAGAAGACACGATCCACCGCC-3′

SEQ ID No: 5
oligonucleotide (38 b)
assrA
5′-TTAAGCTGCTAAAGCGTAGTTTTCGTCGTTTGCGACTA-3′

SEQ ID No: 6

protein (833 aa)
MLV_pD

```
mrgshhhhhhgsgsmgmtlniedehrlhetskepdvslgstwlsdfpqawaetggmglavrqapliiplkatstpvs
ikqypmsqearlgikphiqrlldqgilvpcqspwntpllpvkkpgtndyrpvqdlrevnkrvedihptvpnpynlls
glppshqwytvldlkdaffclrlhptsqplfafewrdpemgisgqltwtrlpqgfknsptlfdealhrdladfriqh
pdlillqyvddlllaatseldcqqgtrallqtlgnlgyrasakkaqicqkqvkylgyllkegqrwltearketvmgq
ptpktprqlreflgtagfcrlwipgfaemaaplypltktgtlfnwgpdqqkayqeikqalltapalglpdltkpfel
fvdekqgyakgvltqklgpwrrpvaylskkldpvaagwppclrmvaaiavltkdagkltmgqplvilaphavealvk
qppdrwlsnarmthyqallldtdrvqfgpvvalnpatllplpeeglqhncldilaeahgtrpdltdqplpdadhtwy
tdgssllqegqrkagaavtteteviwakalpagtsaqraelialtqalkmaegkklnvytdsryafatahihgeiyr
```

rrglltsegkeiknkdeilallkalflpkrlsiihcpghqkghsaeargnrmadqaarkaaitetpdtstlliensspnsrlinefgsgggsgggsmgtatapgglsakapamtplmldtssrklvawdgttdgaavgilavaadqtsttltfyksgtfryedvlwpeaasdetkkrtafagtaisivgsgggsgggsgggsgggsgggsgggscll

SEQ ID No: 7

protein (766 aa)
del_pD
mrgshhhhhhgsgsmgmtlniedehrlhetskepdvslgstwlsdfpqawaetggmglavrqapliiplkatstpvsikqypmsqearlgikphiqrlldqgilvpcqspwntpllpvkkpgtndyrpvqdlrevnkrvedihptvpnpynllsglppshqwytvldlkdaffclrlhptsqplfafewrdpemgisgqltwtrlpqgfknsptlfdealhrdladfriqhpdlillqyvddlllaatseldcqqgtrallqtlgtagfcrlwipgfaemaaplyplтktgtlfnwgpdqqkayqeikqalltapalglpdltkpfelfvdekqgyakgvltqklgpwrrpvaylskkldpvaagwppclrmvaaiavltkdagkltmgqplvilaphavealvkqppdrwlsnarmthyqallldtdrvqfgpvvalnpatllplpeeglqhncldilaeahgtrpdltdqplpdadhtwytdgssllqegqrkagaavtteteviwakalpagtsaqraelialtqalkmaegkklnvytdsryafatahihgeiyrrrglltsegkeiknkdeilallkalflpkrlsiihcpghqkghsaeargnrmadqaarkaaitetpdtstlliensspnsrlinefgsgggsgggsmgtatapgglsakapamtplmldtssrklvawdgttdgaavgilavaadqtsttltfyksgtfryedvlwpeaasdetkkrtafagtaisivgsgggsgggsgggsgggsgggsgggscll

SEQ ID No: 8

oligonucleotide (15 b)
pD_42
5'-TTACGGCTGGAGGTG-3'

SEQ ID No: 9

oligonucleotide (28 b)
RD_Nde
5'- CTTTAAGAAAGAGGAGAAATTACATATG-3'

SEQ ID No: 10

oligonucleotide (14 b)
pD_55
5'-GCCGGGCGCGGTTG-3'

SEQ ID No: 11
oligonucleotide (23 b)
M_F
5'-GATCAAGCCCCACATACAGAGAC-3'

SEQ ID No: 12

oligonucleotide (19 b)
M_2R
5'-GCCCTGCTTCTCGTCGACA-3'

SEQ ID No: 13

oligonucleotide (40 b)
M_Esp
5'-ATCGTCTCCCATGGGCATGACCCTAAATATAGAAGATGAG-3'

SEQ ID No: 14

oligonucleotide (32 b)
M_Eri
5'-AATGAATTCATTAATTAAGCGGGAATTGGGTG-3'

SEQ ID No: 15

oligonucleotide (18 b)
M_1R
5'-CAGGGCCCGAGTACCTTG-3'

SEQ ID No: 16

oligonucleotide (17 b)
M_3F
5'-CCAGTTCGGACCGGTGG-3'

SEQ ID No: 17

oligonucleotide (19 b)
pD_ter-
5'-AAGAAGACACGATCCACCG-3'

SEQ ID No: 18

oligonucleotide (36 b)
M_Hind3+
5'-CGGATCAAGCTTAATTAATTAAGCGGGAATTGGGTG-3'

SEQ ID No: 19

```
LOCUS       pET_his_MLV            7474 bp    DNA     circular      4-JUN-2007
SOURCE
  ORGANISM
COMMENT     This file is created by Vector NTI
            http://www.invitrogen.com/
COMMENT     ORIGDB|GenBank
COMMENT     VNTDATE|446486240|
COMMENT     VNTDBDATE|448293778|
COMMENT     LSOWNER|
COMMENT     VNTNAME|pET_his_MLV|
COMMENT     VNTAUTHORNAME|Remigijus Skirgaila|
COMMENT     VNTAUTHORTEL|+370-5-2394224|
```

```
COMMENT     VNTAUTHOREML|skirgaila@fermentas.lt|
COMMENT     VNTAUTHORWWW|www.fermentas.com|
FEATURES             Location/Qualifiers
     CDS             3135..3992
                     /vntifkey="4"
                     /label=Ap
                     /note="ORF: Frame #2   Start: atg   Stop:  taa"
     CDS             complement(6187..7275)
                     /vntifkey="4"
                     /label=lacI
                     /note="ORF: Frame #3   Start: gtg   Stop:  tga"
     terminator      2465..2511
                     /vntifkey="43"
                     /label=T7\terminator
     rep_origin      2548..3003
                     /vntifkey="33"
                     /label=f1\origin
     rep_origin      complement(2537..2537)
                     /vntifkey="33"
                     /label=ori
     promoter        174..190
                     /vntifkey="30"
                     /label=PT7
     CDS             306..2360
                     /vntifkey="4"
                     /label=MLV
     CDS             258..305
                     /vntifkey="4"
                     /label=his
BASE COUNT     1810 a      2046 c      1942 g      1676 t
ORIGIN
        1 cacggggcct gccaccatac ccacgccgaa acaagcgctc atgagcccga agtggcgagc
       61 ccgatcttcc ccatcggtga tgtcggcgat ataggcgcca gcaaccgcac ctgtggcgcc
      121 ggtgatgccg gccacgatgc gtccggcgta gaggatcgag atctatacga aattaatacg
      181 actcactata gggagaccac aacggtttcc ctctagaaat aattttgttt aactttaaga
      241 aagaggagaa attacatatg agaggatcgc atcaccatca ccatcacgga tctggttcca
      301 tgggcatgac cctaaatata gaagatgagc atcggctaca tgagacctca aaagagccag
      361 atgtttctct agggtccaca tggctgtctg attttcctca ggcctgggcg gaaaccgggg
      421 gcatgggact ggcagttcgc caagctcctc tgatcatacc tctgaaagca acctctaccc
      481 ccgtgtccat aaaacaatac cccatgtcac aagaagccag actggggatc aagccccaca
      541 tacagagact gttggaccag ggaatactgg taccctgcca gtccccctgg aacacgcccc
      601 tgctacccgt taagaaacca gggactaatg attataggcc tgtccaggat ctgagagaag
      661 tcaacaagcg ggtggaagac atccacccca ccgtgcccaa cccttacaac ctcttgagcg
      721 ggctcccacc gtcccaccag tggtacactg tgcttgattt aaaggatgcc tttttctgcc
      781 tgagactcca ccccaccagt cagcctctct tcgcctttga gtggagagat ccagagatgg
      841 gaatctcagg acaattgacc tggaccagac tcccacaggg tttcaaaaac agtcccaccc
      901 tgtttgatga ggcactgcac agagacctag cagacttccg gatccagcac ccagacttga
      961 tcctgctaca gtacgtggat gacttactgc tggccgccac ttctgagcta gactgccaac
     1021 aaggtactcg ggccctgtta caaaccctag ggaacctcgg gtatcgggcc tcggccaaga
     1081 aagcccaaat ttgccagaaa caggtcaagt atctggggta tcttctaaaa gagggtcaga
     1141 gatggctgac tgaggccaga aaagagactg tgatggggca gcctactccg aagacccctc
     1201 gacaactaag ggagttccta gggacggcag gcttctgtcg cctctggatc cctgggtttg
     1261 cagaaatggc agcccccttg taccctctca ccaaaacggg gactctgttt aattggggcc
     1321 cagaccaaca aaaggcctat caagaaatca agcaagctct tctaactgcc ccagccctgg
     1381 ggttgccaga tttgactaag ccctttgaac tctttgtcga cgagaagcag ggctacgcca
     1441 aaggtgtcct aacgcaaaaa ctgggacctt ggcgtcggcc ggtggcctac ctgtccaaaa
     1501 agctagaccc agtagcagct gggtggcccc cttgcctacg gatggtagca gccattgccg
     1561 tactgacaaa ggatgcaggc aagctaacca tgggacagcc actagtcatt ctggcccccc
     1621 atgcagtaga ggcactagtc aaacaacccc ccgaccgctg gctttccaac gcccggatga
```

1681 ctcactatca ggccttgctt ttggacacgg accgggtcca gttcggaccg gtggtagccc
1741 tgaacccggc tacgctgctc ccactgcctg aggaagggct gcaacacaac tgccttgata
1801 tcctggccga agcccacgga acccgacccg acctaacgga ccagccgctc ccagacgccg
1861 accacacctg gtacacggat ggaagcagtc tcttacaaga gggacagcgt aaggcgggag
1921 ctgcggtgac caccgagacc gaggtaatct gggctaaagc cctgccagcc gggacatccg
1981 ctcagcgggc tgaactgata gcactcaccc aggccctaaa gatggcagaa ggtaagaagc
2041 taaatgttta tactgatagc cgttatgctt ttgctactgc ccatatccat ggagaaatat
2101 acagaaggcg tgggttgctc acatcagaag gcaaagagat caaaaataaa gacgagatct
2161 tggccctact aaaagccctc tttctgccca aaagacttag cataatccat tgtccaggac
2221 atcaaaaggg acacagcgcc gaggctagag gcaaccggat ggctgaccaa gcggcccgaa
2281 aggcagccat cacagagact ccagacacct ctaccctcct catagaaaat tcatcaccca
2341 attcccgctt aattaattaa gcttgcggcc gcactcgagc accaccacca ccaccactga
2401 gatccggctg ctaacaaagc ccgaaaggaa gctgagttgg ctgctgccac cgctgagcaa
2461 taactagcat aacccccttgg ggcctctaaa cgggtcttga ggggtttttt gctgaaagga
2521 ggaactatat ccggattggc gaatgggacg cgccctgtag cggcgcatta agcgcggcgg
2581 gtgtggtggt tacgcgcagc gtgaccgcta cacttgccag cgccctagcg cccgctcctt
2641 tcgctttctt cccttccttt ctcgccacgt tcgccggctt tccccgtcaa gctctaaatc
2701 gggggctccc tttagggttc cgatttagtg ctttacggca cctcgacccc aaaaaacttg
2761 attagggtga tggttcacgt agtgggccat cgccctgata gacggttttt cgccctttga
2821 cgttggagtc cacgttcttt aatagtggac tcttgttcca aactggaaca acactcaacc
2881 ctatctcggt ctattctttt gatttataag ggattttgcc gatttcggcc tattggttaa
2941 aaaatgagct gatttaacaa aaatttaacg cgaattttaa caaaatatta acgtttacaa
3001 tttcaggtgg cacttttcgg ggaaatgtgc gcggaacccc tatttgttta tttttctaaa
3061 tacattcaaa tatgtatccg ctcatgagac aataaccctg ataaatgctt caataatatt
3121 gaaaaaggaa gagtatgagt attcaacatt tccgtgtcgc ccttattccc ttttttgcgg
3181 cattttgcct tcctgttttt gctcacccag aaacgctggt gaaagtaaaa gatgctgaag
3241 atcagttggg tgcacgagtg ggttacatcg aactggatct caacagcggt aagatccttg
3301 agagttttcg ccccgaagaa cgttttccaa tgatgagcac ttttaaagtt ctgctatgtg
3361 gcgcggtatt atcccgtatt gacgccgggc aagagcaact cggtcgccgc atacactatt
3421 ctcagaatga cttggttgag tactcaccag tcacagaaaa gcatcttacg gatggcatga
3481 cagtaagaga attatgcagt gctgccataa ccatgagtga taacactgcg gccaacttac
3541 ttctgacaac gatcggagga ccgaaggagc taaccgcttt tttgcacaac atgggggatc
3601 atgtaactcg ccttgatcgt tgggaaccgg agctgaatga agccatacca aacgacgagc
3661 gtgacaccac gatgcctgca gcaatggcaa caacgttgcg caaactatta actggcgaac
3721 tacttactct agcttcccgg caacaattaa tagactggat ggaggcggat aaagttgcag
3781 gaccacttct gcgctcggcc cttccggctg gctggtttat tgctgataaa tctggagccg
3841 gtgagcgtgg gtctcgcggt atcattgcag cactggggcc agatggtaag ccctcccgta
3901 tcgtagttat ctacacgacg gggagtcagg caactatgga tgaacgaaat agacagatcg
3961 ctgagatagg tgcctcactg attaagcatt ggtaactgtc agaccaagtt tactcatata
4021 tactttagat tgatttaaaa cttcattttt aatttaaaag gatctaggtg aagatccttt
4081 ttgataatct catgaccaaa atcccttaac gtgagttttc gttccactga gcgtcagacc
4141 ccgtagaaaa gatcaaagga tcttcttgag atccttttttt tctgcgcgta atctgctgct
4201 tgcaaacaaa aaaaccaccg ctaccagcgg tggtttgttt gccggatcaa gagctaccaa
4261 ctcttttttcc gaaggtaact ggcttcagca gagcgcagat accaaatact gtccttctag
4321 tgtagccgta gttaggccac cacttcaaga actctgtagc accgcctaca tacctcgctc
4381 tgctaatcct gttaccagtg gctgctgcca gtggcgataa gtcgtgtctt accgggttgg
4441 actcaagacg atagttaccg gataaggcgc agcggtcggg ctgaacgggg ggttcgtgca
4501 cacagcccag cttggagcga acgacctaca ccgaactgag atacctacag cgtgagctat
4561 gagaaagcgc cacgcttccc gaagggagaa aggcggacag gtatccggta agcggcaggg
4621 tcggaacagg agagcgcacg agggagcttc cagggggaaa cgcctggtat ctttatagtc
4681 ctgtcgggtt tcgccacctc tgacttgagc gtcgattttt gtgatgctcg tcaggggggc
4741 ggagcctatg gaaaaacgcc agcaacgcgg cctttttacg gttcctggcc ttttgctggc
4801 cttttgctca catgttcttt cctgcgttat cccctgattc tgtggataac cgtattaccg
4861 cctttgagtg agctgatacc gctcgccgca gccgaacgac cgagcgcagc gagtcagtga
4921 gcgaggaagc ggaagagcgc ctgatgcggt attttctcct tacgcatctg tgcggtattt
4981 cacaccgcat atatggtgca ctctcagtac aatctgctct gatgccgcat agttaagcca
5041 gtatacactc cgctatcgct acgtgactgg gtcatggctg cgccccgaca cccgccaaca
5101 cccgctgacg cgccctgacg ggcttgtctg ctcccggcat ccgcttacag acaagctgtg
5161 accgtctccg ggagctgcat gtgtcagagg ttttcaccgt catcaccgaa acgcgcgagg

```
5221 cagctgcggt aaagctcatc agcgtggtcg tgaagcgatt cacagatgtc tgcctgttca
5281 tccgcgtcca gctcgttgag tttctccaga agcgttaatg tctggcttct gataaagcgg
5341 gccatgttaa gggcggtttt ttcctgtttg gtcactgatg cctccgtgta agggggattt
5401 ctgttcatgg gggtaatgat accgatgaaa cgagagagga tgctcacgat acgggttact
5461 gatgatgaac atgcccggtt actggaacgt tgtgagggta aacaactggc ggtatggatg
5521 cggcgggacc agagaaaaat cactcagggt caatgccagc gcttcgttaa tacagatgta
5581 ggtgttccac agggtagcca gcagcatcct gcgatgcaga tccggaacat aatggtgcag
5641 ggcgctgact tccgcgtttc cagactttac gaaacacgga aaccgaagac cattcatgtt
5701 gttgctcagg tcgcagacgt tttgcagcag cagtcgcttc acgttcgctc gcgtatcggt
5761 gattcattct gctaaccagt aaggcaaccc cgccagccta gccgggtcct caacgacagg
5821 agcacgatca tgcgcacccg tggggccgcc atgccggcga taatggcctg cttctcgccg
5881 aaacgtttgg tggcgggacc agtgacgaag gcttgagcga gggcgtgcaa gattccgaat
5941 accgcaagcg acaggccgat catcgtcgcg ctccagcgaa agcggtcctc gccgaaaatg
6001 acccagagcg ctgccggcac ctgtcctacg agttgcatga taaagaagac agtcataagt
6061 gcggcgacga tagtcatgcc ccgcgcccac cggaaggagc tgactgggtt gaaggctctc
6121 aagggcatcg gtcgagatcc cggtgcctaa tgagtgagct aacttacatt aattgcgttg
6181 cgctcactgc ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc
6241 caacgcgcgg ggagaggcgg tttgcgtatt gggcgccagg gtggtttttc ttttcaccag
6301 tgagacgggc aacagctgat tgcccttcac cgcctggccc tgagagagtt gcagcaagcg
6361 gtccacgctg gtttgcccca gcaggcgaaa atcctgtttg atggtggtta acggcgggat
6421 ataacatgag ctgtcttcgg tatcgtcgta tcccactacc gagatatccg caccaacgcg
6481 cagcccggac tcggtaatgg cgcgcattgc gcccagcgcc atctgatcgt tggcaaccag
6541 catcgcagtg ggaacgatgc cctcattcag catttgcatg gtttgttgaa aaccggacat
6601 ggcactccag tcgccttccc gttccgctat cggctgaatt tgattgcgag tgagatattt
6661 atgccagcca gccagacgca gacgcgccga gacagaactt aatgggcccg ctaacagcgc
6721 gatttgctgg tgacccaatg cgaccagatg ctccacgccc agtcgcgtac cgtcttcatg
6781 ggagaaaata atactgttga tgggtgtctg gtcagagaca tcaagaaata acgccggaac
6841 attagtgcag gcagcttcca cagcaatggc atcctggtca tccagcggat agttaatgat
6901 cagcccactg acgcgttgcg cgagaagatt gtgcaccgcc gctttacagg cttcgacgcc
6961 gcttcgttct accatcgaca ccaccacgct ggcacccagt tgatcggcgc gagatttaat
7021 cgccgcgaca atttgcgacg gcgcgtgcag ggccagactg gaggtggcaa cgccaatcag
7081 caacgactgt ttgcccgcca gttgttgtgc cacgcggttg ggaatgtaat tcagctccgc
7141 catcgccgct tccacttttt cccgcgtttt cgcagaaacg tggctggcct ggttcaccac
7201 gcgggaaacg gtctgataag agacaccggc atactctgcg acatcgtata acgttactgg
7261 tttcacattc accaccctga attgactctc ttccgggcgc tatcatgcca taccgcgaaa
7321 ggttttgcgc cattcgatgg tgtccgggat ctcgacgctc tcccttatgc gactcctgca
7381 ttaggaagca gcccagtagt aggttgaggc cgttgagcac cgccgccgca aggaatggtg
7441 catgcaagga gatggcgccc aacagtcccc cggc
```

SEQ ID No: 20

oligonucleotide (19 b)
pD-ter-
5'-AAGAAGACACGATCCACCG-3'

SEQ ID No: 21

oligonucleotide (35 b)
long+
5'- CGAACGTGGCGAGAAAGGAAGGGAAGAAAGAAGTC-3'

SEQ ID No: 22

oligonucleotide (71 b + 3′ modification ddC)
ddC-Long2
5′-TTTTTTTAGACTTCTTTCTTCCCTTCCTTTCTCGCCACGTTCGAAGAAGACACGAT
CCACCGCCGGTTCCG-ddC-3′

SEQ ID No: 23

oligonucleotide (35 b + 3′ Biotin (TEG)
Long+Tb
5′- CGAACGTGGCGAGAAAGGAAGGGAAGAAAGAAGTC-Bio-3′

References

Mattheakis LC, Bhatt RR, Dower WJ. An in vitro polysome display system for identifying ligands from very large peptide libraries. Proc Natl Acad Sci U S A. 1994 Sep 13;91(19):9022-6.

Matsuura T, Pluckthun A. Selection based on the folding properties of proteins with ribosome display. FEBS Lett. 2003 Mar 27;539(1-3):24-8.

Hanes J, Pluckthun A. In vitro selection and evolution of functional proteins by using ribosome display. Proc Natl Acad Sci U S A. 1997 May 13;94(10):4937-42.

He M, Taussig MJ. Antibody-ribosome-mRNA (ARM) complexes as efficient selection particles for in vitro display and evolution of antibody combining sites. Nucleic Acids Res. 1997 Dec 15;25(24):5132-4.

Irving RA, Coia G, Roberts A, Nuttall SD, Hudson PJ. Ribosome display and affinity maturation: from antibodies to single V-domains and steps towards cancer therapeutics. J Immunol Methods. 2001 Feb 1;248(1-2):31-45. Review.

Amstutz P, Pelletier JN, Guggisberg A, Jermutus L, Cesaro-Tadic S, Zahnd C, Pluckthun A. In vitro selection for catalytic activity with ribosome display. J Am Chem Soc. 2002 Aug 14;124(32):9396-403.

Takahashi F, Ebihara T, Mie M, Yanagida Y, Endo Y, Kobatake E, Aizawa M. Ribosome display for selection of active dihydrofolate reductase mutants using immobilized methotrexate on agarose beads. FEBS Lett. 2002 Mar 6;514(1):106-10.

Tawfik DS, Griffiths AD. Man-made cell-like compartments for molecular evolution.
Nat Biotechnol. 1998 Jul;16(7):652-6.

Lee YF, Tawfik DS, Griffiths AD. Investigating the target recognition of DNA cytosine-5 methyltransferase HhaI by library selection using in vitro compartmentalisation. Nucleic Acids Res. 2002 Nov 15;30(22):4937-44.

Cohen HM, Tawfik DS, Griffiths AD. Altering the sequence specificity of HaeIII methyltransferase by directed evolution using in vitro compartmentalization.
Protein Eng Des Sel. 2004 Jan;17(1):3-11.

Ghadessy FJ, Ong JL, Holliger P. Directed evolution of polymerase function by compartmentalized self-replication. Proc Natl Acad Sci U S A. 2001 Apr 10;98(8):4552-7. Epub 2001 Mar 27.

Ghadessy FJ, Ramsay N, Boudsocq F, Loakes D, Brown A, Iwai S, Vaisman A, Woodgate R, Holliger P. Generic expansion of the substrate spectrum of a DNA polymerase by directed evolution. Nat Biotechnol. 2004 Jun;22(6):755-9. Epub 2004 May 23.

Ong JL, Loakes D, Jaroslawski S, Too K, Holliger P. Directed evolution of DNA polymerase, RNA polymerase and reverse transcriptase activity in a single polypeptide. J Mol Biol. 2006 Aug 18;361(3):537-50. Epub 2006 Jul 5.

Bernath K, Hai M, Mastrobattista E, Griffiths AD, Magdassi S, Tawfik DS. In vitro compartmentalization by double emulsions: sorting and gene enrichment by fluorescence activated cell sorting. Anal Biochem. 2004 Feb 1;325(1):151-7.

Mastrobattista E, Taly V, Chanudet E, Treacy P, Kelly BT, Griffiths AD. High-throughput screening of enzyme libraries: in vitro evolution of a beta-galactosidase by fluorescence-activated sorting of double emulsions. Chem Biol. 2005 Dec;12(12):1291-300.

Bertschinger J, Neri D. Covalent DNA display as a novel tool for directed evolution of proteins in vitro. Protein Eng Des Sel. 2004 Sep;17(9):699-707. Epub 2004 Nov 2.

Doi N, Yanagawa H. STABLE: protein-DNA fusion system for screening of combinatorial protein libraries in vitro. FEBS Lett. 1999 Aug 27;457(2):227-30.

Yonezawa M, Doi N, Kawahashi Y, Higashinakagawa T, Yanagawa H. DNA display for in vitro selection of diverse peptide libraries. Nucleic Acids Res. 2003 Oct 1;31(19):e118.

Sepp A, Choo Y. Cell-free selection of zinc finger DNA-binding proteins using in vitro compartmentalization. J Mol Biol. 2005 Nov 25;354(2):212-9. Epub 2005 Oct 3.

Sepp A, Tawfik DS, Griffiths AD. Microbead display by in vitro compartmentalisation: selection for binding using flow cytometry. FEBS Lett. 2002 Dec 18;532(3):455-8.

Griffiths AD, Tawfik DS. Directed evolution of an extremely fast phosphotriesterase by in vitro compartmentalization. EMBO J. 2003 Jan 2;22(1):24-35.

Bernath K, Magdassi S, Tawfik DS. Directed evolution of protein inhibitors of DNA-nucleases by in vitro compartmentalization (IVC) and nano-droplet delivery. J Mol Biol. 2005 Feb 4;345(5):1015-26. Epub 2004 Dec 7.

Thorsen T, Roberts RW, Arnold FH, Quake SR. Dynamic pattern formation in a vesicle-generating microfluidic device. Phys Rev Lett. 2001 Apr 30;86(18):4163-6.

Okushima S, Nisisako T, Torii T, Higuchi T. Controlled production of monodisperse double emulsions by two-step droplet breakup in microfluidic devices. Langmuir. 2004 Nov 9;20(23):9905-8.

Song H, Tice JD, Ismagilov RF. A microfluidic system for controlling reaction networks in time. Angew Chem Int Ed Engl. 2003 Feb 17;42(7):768-72.

Link DR, Grasland-Mongrain E, Duri A, Sarrazin F, Cheng Z, Cristobal G, Marquez M, Weitz DA. Electric control of droplets in microfluidic devices. Angew Chem Int Ed Engl. 2006 Apr 10;45(16):2556-60.

Matsuura T, Yanagida H, Ushioda J, Urabe I, Yomo T. Nascent chain, mRNA, and ribosome complexes generated by a pure translation system. Biochem Biophys Res Commun. 2007 Jan 12;352(2):372-7. Epub 2006 Nov 17.

Gerard GF, D'Alessio JM, Kotewicz ML, Noon MC. Influence on stability in Escherichia coli of the carboxy-terminal structure of cloned Moloney murine leukemia virus reverse transcriptase. DNA. 1986 Aug;5(4):271-9.

Forrer P, Jaussi R. High-level expression of soluble heterologous proteins in the cytoplasm of Escherichia coli by fusion to the bacteriophage lambda head protein D. Gene. 1998 Dec 11;224(1-2):45-52.

Gerard GF, Potter RJ, Smith MD, Rosenthal K, Dhariwal G, Lee J, Chatterjee DK. The role of template-primer in protection of reverse transcriptase from thermal inactivation. Nucleic Acids Res. 2002 Jul 15;30(14):3118-29.

Vichier-Guerre S, Ferris S, Auberger N, Mahiddine K, Jestin JL. A population of thermostable reverse transcriptases evolved from Thermus aquaticus DNA polymerase I by phage display. Angew Chem Int Ed Engl. 2006 Sep 18;45(37):6133-7.

Ghadessy FJ, Holliger P. A novel emulsion mixture for in vitro compartmentalization of transcription and translation in the rabbit reticulocyte system. Protein Eng Des Sel. 2004 Mar;17(3):201-4. Epub 2004 Feb 27.

Roberts RW, Szostak JW. RNA-peptide fusions for the in vitro selection of peptides and proteins. Proc Natl Acad Sci U S A. 1997 Nov 11;94(23):12297-302.

Odegrip R, Coomber D, Eldridge B, Hederer R, Kuhlman PA, Ullman C, FitzGerald K, McGregor D. CIS display: In vitro selection of peptides from libraries of protein-DNA complexes. Proc Natl Acad Sci U S A. 2004 Mar 2;101(9):2806-10. Epub 2004 Feb 23.

Reiersen H, Løbersli I, Løset GA, Hvattum E, Simonsen B, Stacy JE, McGregor D, Fitzgerald K, Welschof M, Brekke OH, Marvik OJ. Covalent antibody display--an in vitro antibody-DNA library selection system. Nucleic Acids Res. 2005 Jan 14;33(1):e10.

Bertschinger J, Neri D. Covalent DNA display as a novel tool for directed evolution of proteins in vitro. Protein Eng Des Sel. 2004 Sep;17(9):699-707. Epub 2004 Nov 2.

Stein V, Sielaff I, Johnsson K, Hollfelder F. A covalent chemical genotype-phenotype linkage for in vitro protein evolution. Chembiochem. 2007 Dec 17;8(18):2191-4.

Tabuchi I, Soramoto S, Nemoto N, Husimi Y. An in vitro DNA virus for in vitro protein evolution. FEBS Lett. 2001 Nov 23;508(3):309-12.

CLAIMS:

1. A reverse transcriptase enzyme comprising a Moloney Murine Leukemia Virus reverse transcriptase amino acid sequence, wherein said enzyme has an optimum activity at a temperature above 42°C, and comprises a mutation at one or more of the following amino acid positions:

D200,	L603,	L139,	T330,	E607,	T287,
Q221,	I49,	N479,	H594,	A502,	
D653,	K658,	P130,	Q237,	H634,	D524
A307,	Y344,	Q430,	D449,	A644,	N649,
L671,	E673,	M39,	Q91,	M66,	W388,
I179	L333	R390	Q374 and	E5	

wherein when the mutation is at position D653 the mutation is not D653N, wherein when the mutation is at position L603 the mutant is not L603A, and wherein when the mutation is at position H594 the mutation is not H594A.

2. A reverse transcriptase enzyme according to claim 1 having an optimum activity at a temperature of at least 50°C.

3. A reverse transcriptase enzyme according to claim 1 or claim 2 having a higher activity at 50°C as compared with the corresponding wild type enzyme.

4. A reverse transcriptase enzyme according to any preceding claim having a specific activity at 37° which is at least 110% of the corresponding wild type enzyme.

5. A reverse transcriptase enzyme according to any preceding claim having a thermostability of at least 1.5 times that of the corresponding wild type enzyme, measured as residual activity at 37°C following treatment at 50°C for 5 minutes.

6. A reverse transcriptase enzyme according to any preceding claim, having one or more of the following mutations:

D200N,A or, G,	L603W or M,	L139P,	T330P,	E607K, G or A,	T287A,
Q221R,	I49V or T,	N479D,	H594K, R or Q,	H634Y	
A502V,	D653G, A, H or V	K658R or Q	P130S,	Q237R,	D524A
A307V,	Y344H,	Q430R,	D449G or A,	A644V or T,	N649S,
L671P,	E673G or K,	M39V or L,	Q91R or L,	M66L,	W388R,
I179T or V	L333Q	R390W	Q374R and	E5K	

7. A reverse transcriptase enzyme according to any preceding claim comprising at least two mutations.

8. A reverse transcriptase enzyme according to claim 7 wherein the at least two mutations are at D200 and at L603.

9. A reverse transcriptase enzyme according to claim 8 wherein the at least two mutations are D200N and at L603W.

10. A reverse transcriptase enzyme according to claim 7 wherein the at least two mutations are at N479 and H594.

11. A reverse transcriptase enzyme according to claim 10 wherein the at least two mutations are N479D and H594K.

12. A polynucleotide encoding the reverse transcriptase enzyme according to any one of claims 1 to 11.