

PYRROLOPYRAZOLES AS N-TYPE CALCIUM CHANNEL BLOCKERS**CROSS REFERENCE TO RELATED APPLICATIONS**

This application claims priority of the benefits of the filing of U.S. Provisional Application Serial No. 61/683,780, filed August 16, 2012. The complete disclosures of the aforementioned related U.S. patent application is/are hereby incorporated herein by reference for all purposes.

**STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR
DEVELOPMENT**

The research and development of the invention described below was not federally sponsored.

BACKGROUND OF THE INVENTION

Calcium ions play a fundamental role in the physiology and biochemistry of organisms and of cells. The entry of calcium into cells through ion channels mediates a variety of cellular and physiological responses, including gene expression, signal transduction, neurotransmitter release, muscle contraction and hormone secretion. Ion channels are classified by gating, or what opens and closes the channel to the flux of ions. Voltage-gated ion channels open or close depending on the voltage gradient across the plasma membrane, whereas ligand-gated ion channels open or close depending on the binding of ligands to the channel. The classification of voltage-gated calcium channels divides them into three groups: (i) high voltage-activated channels, which include L-, N-, P- and Q-type channels; (ii) intermediate voltage-activated R-type channels; and (iii) low voltage-activated T-type channels.


The N-type calcium channel is distributed mainly in central and peripheral neurons, being localized primarily to presynaptic nerve terminals. This channel regulates the calcium flux required for depolarization-evoked release of neurotransmitters from synaptic endings. The transmission of pain signals from the periphery to the central nervous system (CNS) is mediated, inter alia, by N-type calcium channels located in the spinal cord. Inhibition of the N-type calcium channel in the superficial dorsal horn leads to a decrease in membrane excitability and neurotransmitter release, resulting in pain relief. In addition, knock-out mice lacking the N-type calcium channel exhibit reduced nociceptive behaviors in animal models of pain.

N-type calcium channels have been shown to mediate the development and maintenance of the neuronal sensitization processes associated with neuropathic pain and therefore provide attractive targets for the development of analgesic drugs. Three N-type calcium channel modulators are currently approved for the treatment of pain: ω -conotoxin MVIIA (ziconotide), marketed as Prialt®, potently and selectively blocks the N-type calcium channel and is indicated for the management of severe chronic pain; gabapentin, marketed as Neurontin®, and pregabalin, marketed as Lyrica®, bind with high affinity to the α_2 subunit of the N-type calcium channel and are indicated for the treatment of fibromyalgia, diabetic nerve pain and/or post-herpetic neuralgia pain.

It is an object of the present invention to provide N-Type calcium channel blockers. It is also an object of the invention to provide a method of treating, ameliorating or preventing pain by the administration of a compound of Formula (I). And, it is an object of the invention to provide a pharmaceutical composition comprising a compound of Formula (I), useful for treating, ameliorating or preventing pain.

SUMMARY OF THE INVENTION

The present invention is directed to a compound of Formula (I)



Formula (I)

wherein

R^1 is selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, 2,2-difluoroethoxy, di(C_{1-4} alkyl)amino, trifluoromethoxy, fluoro, and trifluoromethyl;


R^2 is hydrogen; or, R^2 may be taken with R^1 and the phenyl ring to which R^1 and R^2 are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R^3 is hydrogen, methyl, methoxy, chloro, or fluoro;


Q is selected from the group consisting of Q1, Q2, and Q3;



Q1



Q2



Q3

wherein

R^4 is selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, cyano, fluoro, and chloro;

R^5 is independently selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, cyano, chloro, and di(C_{1-4} alkyl)amino;

G is selected from the group consisting of 4-methylphenyl, 3-cyanophenyl, 4-cyanophenyl, 4-methoxycarbonylphenyl, 2-chloro-quinolin-6-yl, morpholin-4-ylsulfonyl, morpholin-4-yl(C_{1-4} alkyl)aminosulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C_{6-10} aryl(C_{1-4} alkyl)aminosulfonyl, and 2,6-dimethylmorpholin-4-ylsulfonyl.

₄alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, trifluoromethylsulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isoindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-carbonyl, trifluoromethylsulfonylaminomethylcarbonyl, trifluoromethoxycarbonyl, C₁₋₄alkylsulfonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, pyridinylcarbonyl, furanylcarbonyl, and 5-methylisoxazol-3-ylcarbonyl;

and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

The present invention also provides, *inter alia*, a pharmaceutical composition comprising, consisting of and/or consisting essentially of a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient, and/or a pharmaceutically acceptable diluent, and a compound of Formula (I), or a pharmaceutically acceptable salt form thereof.

Also provided are processes for making a pharmaceutical composition comprising, consisting of, and/or consisting essentially of admixing a compound of Formula (I) and a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient, and/or a pharmaceutically acceptable diluent.

The present invention further provides, *inter alia*, methods for treating or ameliorating a N-Type calcium channel-modulated disorder in a subject, including a human or other mammal in which the disease, syndrome, or condition is affected by the modulation of the N-Type calcium channel, such as pain and the diseases that lead to such pain, using a compound of Formula (I).

The present invention also provides, *inter alia*, methods for producing the instant compounds and pharmaceutical compositions and medicaments thereof.

DETAILED DESCRIPTION OF THE INVENTION

With reference to substituents, the term “independently” refers to the situation where when more than one substituent is possible, the substituents may be the same or different from each other.

The term “alkyl” whether used alone or as part of a substituent group, refers to straight and branched carbon chains having 1 to 8 carbon atoms. Therefore, designated numbers of carbon atoms (e.g. C₁₋₈) refer independently to the number of carbon atoms in an alkyl moiety or to the alkyl portion of a larger alkyl-containing substituent. In substituent groups with multiple alkyl groups such as (C₁₋₆alkyl)₂amino-, the C₁₋₆alkyl groups of the dialkylamino may be the same or different.

The term “alkoxy” refers to an -O-alkyl group, wherein the term “alkyl” is as defined above.

The terms “alkenyl” and “alkynyl” refer to straight and branched carbon chains having 2 or more carbon atoms, wherein an alkenyl chain contains at least one double bond and an alkynyl chain contains at least one triple bond.

The term “cycloalkyl” refers to saturated or partially saturated, monocyclic or polycyclic hydrocarbon rings of 3 to 14 carbon atoms. Examples of such rings include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and adamantyl.

The term “benzo-fused cycloalkyl” refers to a 5- to 8- membered monocyclic cycloalkyl ring fused to a benzene ring. The carbon atom ring members that form the cycloalkyl ring may be fully saturated or partially saturated.

The term “heterocyclyl” refers to a nonaromatic monocyclic or bicyclic ring system having 3 to 10 ring members and which contains carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O, and S. Included within the term heterocyclyl is a nonaromatic cyclic ring of 5 to 7 members in which 1 to 2 members are nitrogen, or a nonaromatic cyclic ring of 5 to 7 members in which 0, 1 or 2 members are nitrogen and up to 2 members are oxygen or sulfur and at least one member must be either nitrogen, oxygen or sulfur; wherein, optionally, the ring contains zero to one unsaturated bonds, and, optionally, when the ring is of 6 or 7 members, it contains up to 2

unsaturated bonds. The carbon atom ring members that form a heterocycle ring may be fully saturated or partially saturated. The term "heterocyclyl" also includes two 5 membered monocyclic heterocycloalkyl groups bridged to form a bicyclic ring. Such groups are not considered to be fully aromatic and are not referred to as heteroaryl groups. When a heterocycle is bicyclic, both rings of the heterocycle are non-aromatic and at least one of the rings contains a heteroatom ring member. Examples of heterocycle groups include, and are not limited to, pyrrolinyl (including 2*H*-pyrrole, 2-pyrrolinyl or 3-pyrrolinyl), pyrrolidinyl, imidazoliny, imidazolidinyl, pyrazoliny, pyrazolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, and piperazinyl. Unless otherwise noted, the heterocycle is attached to its pendant group at any heteroatom or carbon atom that results in a stable structure.

The term "benzo-fused heterocyclyl" refers to a 5 to 7 membered monocyclic heterocycle ring fused to a benzene ring. The heterocycle ring contains carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O, and S. The carbon atom ring members that form the heterocycle ring may be fully saturated or partially saturated. Unless otherwise noted, benzo-fused heterocycle ring is attached to its pendant group at a carbon atom of the benzene ring.

The term "aryl" refers to an unsaturated, aromatic monocyclic or bicyclic ring of 6 to 10 carbon members. Examples of aryl rings include phenyl and naphthalenyl.

The term "heteroaryl" refers to an aromatic monocyclic or bicyclic aromatic ring system having 5 to 10 ring members and which contains carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O, and S. Included within the term heteroaryl are aromatic rings of 5 or 6 members wherein the ring consists of carbon atoms and has at least one heteroatom member. Suitable heteroatoms include nitrogen, oxygen, and sulfur. In the case of 5 membered rings, the heteroaryl ring preferably contains one member of nitrogen, oxygen or sulfur and, in addition, up to 3 additional nitrogens. In the case of 6 membered rings, the heteroaryl ring preferably contains from 1 to 3 nitrogen atoms. For the case wherein the 6 membered ring has 3 nitrogens, at most 2 nitrogen atoms are adjacent. Examples of heteroaryl groups include furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuryl, benzothienyl, indazolyl, benzimidazolyl, benzothiazolyl,

benzoxazolyl, benzisoxazolyl, benzothiadiazolyl, benzotriazolyl, quinolinyl, isoquinolinyl and quiazolinyl. Unless otherwise noted, the heteroaryl is attached to its pendant group at any heteroatom or carbon atom that results in a stable structure.


The term “halogen” or “halo” refers to fluorine, chlorine, bromine and iodine.

The term “formyl” refers to the group —C(=O)H .

The term “oxo” refers to the group $(=\text{O})$.

Whenever the term “alkyl” or “aryl” or either of their prefix roots appear in a name of a substituent (e.g., arylalkyl, alkylamino) the name is to be interpreted as including those limitations given above for “alkyl” and “aryl.” Designated numbers of carbon atoms (e.g., $\text{C}_1\text{—C}_6$) refer independently to the number of carbon atoms in an alkyl moiety, an aryl moiety, or in the alkyl portion of a larger substituent in which alkyl appears as its prefix root. For alkyl and alkoxy substituents, the designated number of carbon atoms includes all of the independent members included within a given range specified. For example C_{1-6} alkyl would include methyl, ethyl, propyl, butyl, pentyl and hexyl individually as well as sub-combinations thereof (e.g., C_{1-2} , C_{1-3} , C_{1-4} , C_{1-5} , C_{2-6} , C_{3-6} , C_{4-6} , C_{5-6} , C_{2-5} , etc.).

In general, under standard nomenclature rules used throughout this disclosure, the terminal portion of the designated side chain is described first followed by the adjacent functionality toward the point of attachment. Thus, for example, a “ $\text{C}_1\text{—C}_6$ alkylcarbonyl” substituent refers to a group of the formula:



The term “R” at a stereocenter designates that the stereocenter is purely of the *R*-configuration as defined in the art; likewise, the term “S” means that the stereocenter is purely of the *S*-configuration. As used herein, the terms “*R” or “*S” at a stereocenter are used to designate that the stereocenter is of pure but unknown configuration. As used herein, the term “RS” refers to a stereocenter that exists as a mixture of the *R*- and *S*-configurations. Similarly, the terms “*RS” or “*SR” refer to a stereocenter that exists as a mixture of the *R*- and *S*-configurations and is of unknown configuration relative to another stereocenter within the molecule.

Compounds containing one stereocenter drawn without a stereo bond designation are a mixture of 2 enantiomers. Compounds containing 2 stereocenters both drawn without stereo bond designations are a mixture of 4 diastereomers. Compounds with 2 stereocenters both labeled “RS” and drawn with stereo bond designations are a 2-component mixture with relative stereochemistry as drawn. Compounds with 2 stereocenters both labeled “*RS” and drawn with stereo bond designations are a 2-component mixture with relative stereochemistry unknown. Unlabeled stereocenters drawn without stereo bond designations are a mixture of the *R*- and *S*-configurations. For unlabeled stereocenters drawn with stereo bond designations, the absolute stereochemistry is as depicted.

Unless otherwise noted, it is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of Formula (I) can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

The term “subject” refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term “therapeutically effective amount” refers to an amount of an active compound or pharmaceutical agent, including a compound of the present invention, which elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation or partial alleviation of the symptoms of the disease, syndrome, condition, or disorder being treated.

The term “composition” refers to a product that includes the specified ingredients in therapeutically effective amounts, as well as any product that results, directly, or indirectly, from combinations of the specified ingredients in the specified amounts.

The term “N-Type calcium channel blocker” is intended to encompass a compound that interacts with the N-Type calcium channel to substantially reduce or eliminate its functional activity, thereby decreasing the flow of calcium ions through the channel and the rise of intracellular calcium concentrations.

The term "N-Type calcium channel-modulated" is used to refer to the condition of being affected by the modulation of the N-Type calcium channel, including the condition of being affected by the inhibition of the N-Type calcium channel, such as, for example, pain, the diseases that lead to such pain and treatments that lead to the reduction of such pain.

As used herein, unless otherwise noted, the term "affect" or "affected" (when referring to a disease, syndrome, condition or disorder that is affected by the inhibition of N-Type calcium channel) shall include a reduction in the frequency and / or severity of one or more symptoms or manifestations of said disease, syndrome, condition or disorder and / or include the prevention of the development of one or more symptoms or manifestations of said disease, syndrome, condition or disorder or the development of the disease, condition, syndrome or disorder.

The compounds of Formula (I) are useful in methods for treating, ameliorating and/or preventing a disease, a syndrome, a condition or a disorder that is affected by the inhibition of N-Type calcium channel. Such methods comprise, consist of and/or consist essentially of administering to a subject, including an animal, a mammal, and a human in need of such treatment, amelioration and / or prevention, a therapeutically effective amount of a compound of Formula (I), or an enantiomer, diastereomer, solvate or pharmaceutically acceptable salt form thereof. In particular, the compounds of Formula (I) are useful for treating, ameliorating and / or preventing pain as well as diseases, syndromes, conditions or disorders causing such pain. More particularly, the compounds of Formula (I) are useful for treating, ameliorating and / or preventing acute pain, inflammatory pain and / or neuropathic pain, comprising administering to a subject in need thereof a therapeutically effective amount of a compound of Formula (I), as herein defined.


Acute pain, as used herein, refers to pain that comes on quickly, can be of varying severity but is self-limiting and of relatively short duration. Examples of acute pain include, but are not limited to, post-operative pain, post-surgical pain, toothache, burn, sunburn, insect/animal bites and stings, headache and/or any pain associated with acute trauma or injury.

Inflammatory pain refers to pain arising from an inflammatory disease, condition, syndrome or disorder, including but not limited to inflammatory bowel disease, irritable bowel syndrome, visceral pain, migraine, post-operative pain, osteoarthritis, rheumatoid

arthritis, back pain, low back pain, joint pain, abdominal pain, chest pain, labor pain, musculoskeletal diseases, skin diseases, toothache, pyresis, burn, sunburn, snake bite, venomous snake bite, spider bite, insect sting, neurogenic or overactive bladder, interstitial cystitis, urinary tract infection, rhinitis, contact dermatitis/hypersensitivity, itch, eczema, pharyngitis, mucositis, enteritis, irritable bowel syndrome, cholecystitis, pancreatitis, post-mastectomy pain syndrome, menstrual pain, endometriosis, pain due to physical trauma, headache, sinus headache, tension headache or arachnoiditis.

A further embodiment of the present invention is directed to a method for treating, ameliorating and / or preventing neuropathic pain. Neuropathic pain refers to a disease, syndrome, condition and/or disorder involving damage to the peripheral or central nervous system, including cancer pain, neurological disorders, spine and peripheral nerve surgery, brain tumor, traumatic brain injury (TBI), chemotherapy-induced pain, pain chronification, radicular pain, HIV pain, spinal cord trauma, chronic pain syndrome, fibromyalgia, chronic fatigue syndrome, lupus, sarcoidosis, peripheral neuropathy, bilateral peripheral neuropathy, diabetic neuropathy, central pain, neuropathies associated with spinal cord injury, stroke, amyotrophic lateral sclerosis (ALS), Parkinson's disease, multiple sclerosis, sciatic neuritis, mandibular joint neuralgia, peripheral neuritis, polyneuritis, stump pain, phantom limb pain, bony fractures, oral neuropathic pain, Charcot's pain, complex regional pain syndrome I and II (CRPS I/II), radiculopathy, Guillain-Barre syndrome, meralgia paresthetica, burning-mouth syndrome, optic neuritis, postfebrile neuritis, migrating neuritis, segmental neuritis, Gombault's neuritis, neuronitis, cervicobrachial neuralgia, cranial neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, migrainous neuralgia, idiopathic neuralgia, intercostals neuralgia, mammary neuralgia, Morton's neuralgia, nasociliary neuralgia, occipital neuralgia, post-herpetic neuralgia, causalgia, red neuralgia, Sluder's neuralgia, splenopalatine neuralgia, supraorbital neuralgia, trigeminal neuralgia, vulvodynia, or vidian neuralgia.


Embodiments of the present invention include a compound of Formula (I)




Formula (I)

wherein


- a) R^1 is selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, 2,2-difluoroethoxy, di(C_{1-4} alkyl)amino, and trifluoromethoxy;
- b) R^1 is selected from the group consisting of ethyl, methoxy, ethoxy, isopropoxy, difluoroethoxy, dimethylamino, and trifluoromethoxy;
- c) R^3 is hydrogen, methyl, chloro, or fluoro;
- d) Q is selected from the group consisting of Q1, Q2, and Q3;



Q1



Q2



Q3

wherein


R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, and chloro;


- e) Q is selected from the group consisting of Q1, Q2, and Q3;



Q1



Q2



Q3

wherein

R^4 is chloro;

R⁵ is independently selected from the group consisting of C₁₋₄alkoxy and chloro;

- f) G is selected from the group consisting of 4-methylphenyl, 2-chloro-quinolin-6-yl, C₆₋₁₀aryl(C₁₋₄)alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, trifluoromethylsulfonyl, C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-carbonyl, C₁₋₄alkylsulfonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, pyridinylcarbonyl, and furanylcarbonyl;
- g) G is selected from the group consisting of methylphenyl, 2-chloro-quinolin-6-yl, C₆₋₁₀aryl(C₁₋₄)alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, 4-C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, trifluoromethylsulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-


carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, and pyridinylcarbonyl;

- h) G is selected from the group consisting of 2-chloro-quinolin-6-yl, C₆₋₁₀aryl(C₁₋₄alkyl)sulfonyl, C₆₋₁₀aryloxy(C₁₋₄alkyl)sulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C₁₋₄alkoxycarbonyl(C₁₋₄alkyl)-sulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄alkyl)-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄alkyl)-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄alkyl)-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C₁₋₄alkyl)-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄alkyl)-carbonyl, and C₁₋₄alkoxycarbonyl;

and any combination of embodiments a) through h) above, provided that it is understood that combinations in which different embodiments of the same substituent would be combined are excluded;

and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)


wherein

R¹ is selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, 2,2-difluoroethoxy, di(C₁₋₄alkyl)amino, trifluoromethoxy, fluoro, and trifluoromethyl;

R^2 is hydrogen; or, R^2 may be taken with R^1 and the phenyl ring to which R^1 and R^2 are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R^3 is hydrogen, methyl, methoxy, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;



wherein


R^4 is selected from the group consisting of chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of 4-methylphenyl, 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4})alkylsulfonyl, C_{6-10} aryloxy(C_{1-4})alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, trifluoromethylsulfonyl, C_{1-4} alkoxycarbonyl(C_{1-4})alkyl-sulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, di(C_{1-4} alkyl)aminosulfonylamino(C_{1-4})alkyl-sulfonyl, C_{1-4} alkylsulfonylamino(C_{1-4})alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C_{1-4})alkyl-sulfonyl, (2H)-1,3-dione-1H-isoindol-1-yl-(C_{1-4})alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C_{1-6} alkylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R^6)-piperidin-1-ylsulfonyl wherein R^6 is C_{1-4} alkoxycarbonyl or C_{1-4} alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C_{1-4} alkoxycarbonylamino(C_{2-4})alkyl-carbonyl, C_{1-4} alkoxycarbonylaminomethyl-carbonyl, C_{1-4} alkylsulfonylaminomethyl-carbonyl, C_{1-4} alkoxycarbonyl, di(C_{1-4} alkyl)aminocarbonyl, C_{1-4} alkoxycarbonyl, pyridinylcarbonyl, and furanylcabonyl;

and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

wherein

R^1 is selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, 2,2-difluoroethoxy, di(C_{1-4} alkyl)amino, trifluoromethoxy, fluoro, and trifluoromethyl;


R^2 is hydrogen; or, R^2 may be taken with R^1 and the phenyl ring to which R^1 and R^2 are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R^3 is hydrogen, methyl, methoxy, chloro, or fluoro;


Q is selected from the group consisting of Q1, Q2, and Q3;



Q1



Q2



Q3

wherein

R^4 is chloro;


R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of methylphenyl, 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4})alkylsulfonyl, C_{6-10} aryloxy(C_{1-4})alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-

sulfonyl, 4- C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, trifluoromethylsulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)- 1,3-dione-1H-isoindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, and pyridinylcarbonyl;

and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)


wherein

R¹ is selected from the group consisting of ethyl, methoxy, ethoxy, isopropoxy, difluoroethoxy, dimethylamino, and trifluoromethoxy;

R² is hydrogen; or, R² may be taken with R¹ and the phenyl ring to which R¹ and R² are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R³ is hydrogen, methyl, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;



wherein


R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4} alkyl)sulfonyl, C_{6-10} aryloxy(C_{1-4} alkyl)sulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C_{1-4} alkoxycarbonyl(C_{1-4} alkyl)-sulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, di(C_{1-4} alkyl)aminosulfonylamino(C_{1-4} alkyl)-sulfonyl, C_{1-4} alkylsulfonylamino(C_{1-4} alkyl)-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C_{1-4} alkyl)-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C_{1-4} alkyl)-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C_{1-6} alkylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R^6)-piperidin-1-ylsulfonyl wherein R^6 is C_{1-4} alkoxycarbonyl or C_{1-4} alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C_{1-4} alkoxycarbonylamino(C_{2-4} alkyl)-carbonyl, and C_{1-4} alkoxycarbonyl;

and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)


wherein

R^1 is selected from the group consisting of ethyl, methoxy, ethoxy, isopropoxy, difluoroethoxy, dimethylamino, and trifluoromethoxy;

R^2 is hydrogen;

R^3 is hydrogen, methyl, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;



wherein


R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4} alkyl)sulfonyl, C_{6-10} aryloxy(C_{1-4} alkyl)sulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C_{1-4} alkoxycarbonyl(C_{1-4} alkyl)-sulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, di(C_{1-4} alkyl)aminosulfonylamino(C_{1-4} alkyl)-sulfonyl, C_{1-4} alkylsulfonylamino(C_{1-4} alkyl)-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C_{1-4} alkyl)-sulfonyl, (2H)-1,3-dione-1H-isoindol-1-yl-(C_{1-4} alkyl)-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C_{1-6} alkylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R^6)-piperidin-1-ylsulfonyl wherein R^6 is C_{1-4} alkoxycarbonyl or C_{1-4} alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C_{1-4} alkoxycarbonylamino(C_{2-4} alkyl)-carbonyl, and C_{1-4} alkoxycarbonyl;

and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

Further embodiments of the present invention are directed to a compound of Formula (I)



Formula (I)

selected from the group consisting of

- Cpd 1**, tert-Butyl 3-(4-chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;
- Cpd 2**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 3**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-[(trifluoromethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 4**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(pyridin-3-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 5**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(pyridin-4-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 6**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(trifluoroacetyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 7**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(pyridin-2-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 8**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(furan-2-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 9**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-[(5-methylisoxazol-3-yl)carbonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 10**, 3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(methoxyacetyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 11**, tert-Butyl {2-[3-(4-chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;

- Cpd 12**, N-{2-[3-(4-Chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl}methanesulfonamide;
- Cpd 13**, N-{2-[3-(4-Chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl}-1,1,1-trifluoromethanesulfonamide;
- Cpd 14**, tert-Butyl 3-(4-chlorophenyl)-2-[2-(trifluoromethyl)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;
- Cpd 15**, 3-(4-Chlorophenyl)-5-(methylsulfonyl)-2-[2-(trifluoromethyl)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 16**, 3-(4-Chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2-[2-(trifluoromethyl)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 17**, 3-(4-Chlorophenyl)-5-[(4-methylphenyl)sulfonyl]-2-[2-(trifluoromethyl)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 18**, tert-Butyl {2-[3-(4-chlorophenyl)-2-[2-(trifluoromethyl)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;
- Cpd 19**, tert-Butyl 3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;
- Cpd 20**, 3-(4-Chlorophenyl)-5-(methylsulfonyl)-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 21**, 3-(4-Chlorophenyl)-5-(trifluoroacetyl)-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 22**, tert-Butyl {2-[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;
- Cpd 23**, 3-[3-(4-Chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]benzonitrile;
- Cpd 24**, Methyl 4-[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]benzoate;
- Cpd 25**, 4-[3-(4-Chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]benzonitrile;
- Cpd 26**, 4-[3-(4-Chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-N-(2-morpholin-4-ylethyl)benzenesulfonamide;
- Cpd 27**, 2-Chloro-6-[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]quinoline;

- Cpd 28**, tert-Butyl 3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;
- Cpd 29**, tert-Butyl {2-[3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;
- Cpd 30**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 31**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 32**, 3-(4-Chlorophenyl)-5-(cyclopropylsulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 33**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 34**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxamide;
- Cpd 35**, 3-(4-Chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 36**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(2,2,2-trifluoroethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 37**, 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 38**, 2-(2- {[3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl} ethyl)-1H-isoindole-1,3(2H)-dione;
- Cpd 39**, 5-(Benzylsulfonyl)-3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 40**, 3-(4-Chlorophenyl)-5-(cyclopentylsulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 41**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(1-methylbutyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 42**, 1-(2- {[3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl} ethyl)pyrrolidine-2,5-dione;
- Cpd 43**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(2-phenoxyethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

- Cpd 44**, N-(2-{{[3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl}ethyl)methanesulfonamide;
- Cpd 45**, N-(2-{{[3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl}ethyl)propane-2-sulfonamide;
- Cpd 46**, N'-(2-{{[3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl}ethyl)-N,N-dimethylsulfamide;
- Cpd 47**, Methyl 3-{{[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl} propanoate;
- Cpd 48**, 3-(4-Chlorophenyl)-5-[(1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl]-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 49**, 3-(4-Chlorophenyl)-2-(2-ethoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 50**, 3-(4-Chlorophenyl)-2-(2-ethoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 51**, 3-(4-Chlorophenyl)-5-(cyclopentylsulfonyl)-2-(2-ethoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 52**, 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(2-ethoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 53**, 5-(Benzylsulfonyl)-3-(4-chlorophenyl)-2-(2-ethoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 54**, 3-(4-Chlorophenyl)-2-(2-ethoxyphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 55**, 5-(Azepan-1-ylsulfonyl)-3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 56**, 3-(4-Chlorophenyl)-N,N-diethyl-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 57**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 58**, Methyl 1-{{[3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl} piperidine-4-carboxylate;
- Cpd 59**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(morpholin-4-ylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

- Cpd 60**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(pyrrolidin-1-ylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 61**, 3-(4-Chlorophenyl)-5-([(2S,6S)-2,6-dimethylmorpholin-4-yl]sulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 62**, 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(piperidin-1-ylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 63**, 3-(4-Chlorophenyl)-N,N-dimethyl-2-[2-(1-methylethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 64**, 2-(4-Chloro-2-methoxyphenyl)-3-(4-chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 65**, 2-(4-Chloro-2-methoxyphenyl)-3-(4-chlorophenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 66**, 3-(4-Chlorophenyl)-2-(4-methoxy-2-methylphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 67**, 3-(4-Chlorophenyl)-2-(4-methoxy-2-methylphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 68**, 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(4-methoxy-2-methylphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 69**, 3-(4-Chlorophenyl)-2-(4-methoxy-2-methylphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 70**, 3-(4-Chlorophenyl)-2-(4-fluoro-2-methoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 71**, 3-(4-Chlorophenyl)-2-(4-fluoro-2-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 72**, 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(4-fluoro-2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 73**, 3-(4-Chlorophenyl)-2-(4-fluoro-2-methoxyphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 74**, 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 75**, 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

- Cpd 76**, 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-[(1-methylbutyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 77**, 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 78**, 3-(4-Chlorophenyl)-2-(2-fluoro-4-methoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 79**, 3-(4-Chlorophenyl)-2-(2-fluoro-4-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 80**, 3-(4-Chlorophenyl)-2-(2-fluoro-4-methoxyphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 81**, 2-(1,3-Benzodioxol-4-yl)-3-(4-chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 82**, 2-(1,3-Benzodioxol-4-yl)-3-(4-chlorophenyl)-5-[(2-methylpropyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 83**, 2-(1,3-Benzodioxol-4-yl)-3-(4-chlorophenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 84**, 3-(4-Chlorophenyl)-2-(2,3-dihydro-1-benzofuran-7-yl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 85**, 3-(4-Chlorophenyl)-2-(2,3-dihydro-1-benzofuran-7-yl)-5-[(2-methylpropyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 86**, 3-(4-Chlorophenyl)-2-(2,3-dihydro-1-benzofuran-7-yl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 87**, 3-(4-Chlorophenyl)-5-[(3,5-dimethylisoxazol-4-yl)sulfonyl]-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 88**, 2-[3-(4-Chlorophenyl)-5-[(1-methylethyl)sulfonyl]-5,6-dihydropyrrolo[3,4-c]pyrazol-2(4H)-yl]-N,N-dimethylaniline;
- Cpd 89**, 2-[3-(4-Chlorophenyl)-5-(ethylsulfonyl)-5,6-dihydropyrrolo[3,4-c]pyrazol-2(4H)-yl]-N,N-dimethylaniline;
- Cpd 90**, 3-(4-Chlorophenyl)-N,N-dimethyl-2-[2-(2,2,2-trifluoroethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 91**, 2-(2-tert-Butoxyphenyl)-3-(4-chlorophenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;

- Cpd 92**, 3-(6-Chloropyridin-3-yl)-2-(2-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 93**, 3-{2-(2-Methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl} quinoline;
- Cpd 94**, 2-(2-Methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-3-(6-methylpyridin-3-yl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 95**, 3-(4-Chlorophenyl)-2-[2-(2,2-difluoroethoxy)phenyl]-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- Cpd 96**, 2-(2-Methoxyphenyl)-3-[6-(1-methylethoxy)pyridin-3-yl]-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Cpd 97**, 2-(2-Methoxyphenyl)-3-(6-methoxypyridin-3-yl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- and pharmaceutically acceptable salt forms thereof.

For use in medicine, salts of compounds of Formula (I) refer to non-toxic “pharmaceutically acceptable salts.” Other salts may, however, be useful in the preparation of compounds of Formula (I) or of their pharmaceutically acceptable salts thereof. Suitable pharmaceutically acceptable salts of compounds of Formula (I) include acid addition salts which can, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of Formula (I) carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, such as sodium or potassium salts; alkaline earth metal salts, such as calcium or magnesium salts; and salts formed with suitable organic ligands, such as quaternary ammonium salts. Thus, representative pharmaceutically acceptable salts include acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate,

mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide and valerate.

Representative acids and bases that may be used in the preparation of pharmaceutically acceptable salts include acids including acetic acid, 2,2-dichloroacetic acid, acylated amino acids, adipic acid, alginic acid, ascorbic acid, L-aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, (+)-camphoric acid, camphorsulfonic acid, (+)-(1S)-camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxy-ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, D-gluconic acid, D-glucuronic acid, L-glutamic acid, α -oxo-glutaric acid, glycolic acid, hippuric acid, hydrobromic acid, hydrochloric acid, (+)-L-lactic acid, ($\pm$)-DL-lactic acid, lactobionic acid, maleic acid, (-)-L-malic acid, malonic acid, ($\pm$)-DL-mandelic acid, methanesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, nitric acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, phosphoric acid, L-pyroglutamic acid, salicylic acid, 4-amino-salicylic acid, sebaic acid, stearic acid, succinic acid, sulfuric acid, tannic acid, (+)-L-tartaric acid, thiocyanic acid, p-toluenesulfonic acid and undecylenic acid; and bases including ammonia, L-arginine, benethamine, benzathine, calcium hydroxide, choline, deanol, diethanolamine, diethylamine, 2-(diethylamino)-ethanol, ethanolamine, ethylenediamine, N-methylglucamine, hydrabamine, 1H-imidazole, L-lysine, magnesium hydroxide, 4-(2-hydroxyethyl)-morpholine, piperazine, potassium hydroxide, 1-(2-hydroxyethyl)-pyrrolidine, sodium hydroxide, triethanolamine, tromethamine and zinc hydroxide.

Embodiments of the present invention include prodrugs of compounds of Formula (I). In general, such prodrugs will be functional derivatives of the compounds that are readily convertible *in vivo* into the required compound. Thus, in the methods of treating or preventing embodiments of the present invention, the term “administering” encompasses the treatment or prevention of the various diseases, conditions, syndromes and disorders described with the compound specifically disclosed or with a compound that may not be

specifically disclosed, but which converts to the specified compound *in vivo* after administration to a patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

Where the compounds according to embodiments of this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention. The skilled artisan will understand that the term compound as used herein, is meant to include solvated compounds of Formula (I).

Where the processes for the preparation of the compounds according to certain embodiments of the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

One embodiment of the present invention is directed to a composition, including a pharmaceutical composition, comprising, consisting of, and/or consisting essentially of the (+)-enantiomer of a compound of Formula (I) wherein said composition is substantially free from the (-)-isomer of said compound. In the present context, substantially free means less than about 25 %, preferably less than about 10 %, more preferably less than about 5 %, even more preferably less than about 2 % and even more preferably less than about 1 %

of the (-)-isomer calculated as.

$$\% (+) - \text{enantiomer} = \frac{(\text{mass}(+) - \text{enantiomer})}{(\text{mass}(+) - \text{enantiomer}) + (\text{mass}(-) - \text{enantiomer})} \times 100$$

Another embodiment of the present invention is a composition, including a pharmaceutical composition, comprising, consisting of, and consisting essentially of the (-)-enantiomer of a compound of Formula (I) wherein said composition is substantially free from the (+)-isomer of said compound. In the present context, substantially free from means less than about 25 %, preferably less than about 10 %, more preferably less than about 5 %, even more preferably less than about 2 % and even more preferably less than about 1 % of the (+)-isomer calculated as

$$\% (-) - \text{enantiomer} = \frac{(\text{mass}(-) - \text{enantiomer})}{(\text{mass}(+) - \text{enantiomer}) + (\text{mass}(-) - \text{enantiomer})} \times 100$$

During any of the processes for preparation of the compounds of the various embodiments of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry, Second Edition*, J.F.W. McOmie, Plenum Press, 1973; T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis, Third Edition*, John Wiley & Sons, 1999. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

Even though the compounds of embodiments of the present invention (including their pharmaceutically acceptable salts and pharmaceutically acceptable solvates) can be administered alone, they will generally be administered in admixture with a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient and/or a pharmaceutically acceptable diluent selected with regard to the intended route of administration and standard pharmaceutical or veterinary practice. Thus, particular

embodiments of the present invention are directed to pharmaceutical and veterinary compositions comprising compounds of Formula (I) and at least one pharmaceutically acceptable carrier, pharmaceutically acceptable excipient, and/or pharmaceutically acceptable diluent.

By way of example, in the pharmaceutical compositions of embodiments of the present invention, the compounds of Formula (I) may be admixed with any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilizing agent(s), and combinations thereof.

Solid oral dosage forms, such as tablets or capsules, containing the compounds of the present invention may be administered in at least one dosage form at a time, as appropriate. It is also possible to administer the compounds in sustained release formulations. Additional oral forms in which the present inventive compounds may be administered include elixirs, solutions, syrups, and suspensions; each optionally containing flavoring agents and coloring agents.

Alternatively, compounds of Formula (I) can be administered by inhalation (intratracheal or intranasal) or in the form of a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. For example, they can be incorporated into a cream comprising, consisting of, and/or consisting essentially of an aqueous emulsion of polyethylene glycols or liquid paraffin. They can also be incorporated, at a concentration of between about 1 % and about 10 % by weight of the cream, into an ointment comprising, consisting of, and/or consisting essentially of a white wax or white soft paraffin base together with any stabilizers and preservatives as may be required. An alternative means of administration includes transdermal administration by using a skin or transdermal patch.

The pharmaceutical compositions of the present invention (as well as the compounds of the present invention alone) can also be injected parenterally, for example intracavernosally, intravenously, intramuscularly, subcutaneously, intradermally or intrathecally. In this case, the compositions will also include at least one of a suitable carrier, a suitable excipient, and a suitable diluent.

For parenteral administration, the pharmaceutical compositions of the present invention are best used in the form of a sterile aqueous solution that may contain other substances, for example, enough salts and monosaccharides to make the solution isotonic

with blood.

For buccal or sublingual administration, the pharmaceutical compositions of the present invention may be administered in the form of tablets or lozenges, which can be formulated in a conventional manner.

By way of further example, pharmaceutical compositions containing at least one of the compounds of Formula (I) as the active ingredient can be prepared by mixing the compound(s) with a pharmaceutically acceptable carrier, a pharmaceutically acceptable diluent, and/or a pharmaceutically acceptable excipient according to conventional pharmaceutical compounding techniques. The carrier, excipient, and diluent may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral, etc.). Thus for liquid oral preparations, such as suspensions, syrups, elixirs and solutions, suitable carriers, excipients and diluents include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers, excipients and diluents include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations also may be optionally coated with substances, such as, sugars, or be enterically-coated so as to modulate the major site of absorption and disintegration. For parenteral administration, the carrier, excipient and diluent will usually include sterile water, and other ingredients may be added to increase solubility and preservation of the composition. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives, such as solubilizers and preservatives.

A therapeutically effective amount of a compound of Formula (I) or a pharmaceutical composition thereof includes a dose range from about 0.1 mg to about 3000 mg, or any particular amount or range therein, in particular from about 1 mg to about 1000 mg, or any particular amount or range therein; or, more particularly, from about 10 mg to about 500 mg, or any particular amount or range therein, of active ingredient in a regimen of about 1 to about 4 times per day for an average (70 kg) human; although, it is apparent to one skilled in the art that the therapeutically effective amount for a compound of Formula (I) will vary as will the diseases, syndromes, conditions, and disorders being treated.

For oral administration, a pharmaceutical composition is preferably provided in the form of tablets containing about 0.01, about 10, about 50, about 100, about 150, about 200, about 250, and about 500 milligrams of a compound of Formula (I).

Advantageously, a compound of Formula (I) may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three and four times daily.

Optimal dosages of a compound of Formula (I) to be administered may be readily determined and will vary with the particular compound used, the mode of administration, the strength of the preparation and the advancement of the disease, syndrome, condition or disorder. In addition, factors associated with the particular subject being treated, including subject gender, age, weight, diet and time of administration, will result in the need to adjust the dose to achieve an appropriate therapeutic level and desired therapeutic effect. The above dosages are thus exemplary of the average case. There can be, of course, individual instances wherein higher or lower dosage ranges are merited, and such are within the scope of this invention.

Compounds of Formula (I) may be administered in any of the foregoing compositions and dosage regimens or by means of those compositions and dosage regimens established in the art whenever use of a compound of Formula (I) is required for a subject in need thereof.

As N-Type calcium channel blockers, the compounds of Formula (I) are useful in methods for treating and/or preventing a disease, a syndrome, a condition or a disorder in a subject, including an animal, a mammal and a human in which the disease, the syndrome, the condition or the disorder is affected by the modulation of the N-Type calcium channel. Such methods comprise, consist of and/or consist essentially of administering to a subject, including an animal, a mammal, and a human in need of such treatment or prevention a therapeutically effective amount of a compound, salt or solvate of Formula (I). In particular, the compounds of Formula (I) are useful for preventing or treating pain, such as inflammatory pain or neuropathic pain, or diseases, syndromes, conditions or disorders causing such pain.

GENERAL SYNTHETIC METHODS

Representative compounds of the present invention can be synthesized in accordance with the general synthetic methods described below and illustrated in the schemes and examples that follow. Since the schemes are an illustration, the invention should not be construed as being limited by the chemical reactions and conditions described in the schemes. The various starting materials used in the schemes and examples are commercially available or may be prepared by methods well within the skill of persons versed in the art. The variables are as defined herein.


Abbreviations used in the instant specification, particularly the schemes and examples, are as follows:

ACN	acetonitrile
AcOH	glacial acetic acid
aq.	aqueous
Bn or Bzl	benzyl
Boc	tert-butyloxycarbonyl
conc.	concentrated
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DCC	<i>N,N'</i> -dicyclohexyl-carbodiimide
DCE	1,2-dichloroethane
DCM	dichloromethane
DIBALH	diisobutylaluminum hydride
DIPEA or DIEA	diisopropyl-ethyl amine
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
EA	ethyl acetate
EDCI	1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
EGTA	ethylene glycol tetraacetic acid
ESI	electrospray ionization
EtOAc or EA	ethyl acetate
EtOH	ethanol
h or hr(s)	hour or hours

HEK	human embryonic kidney
HEPES	(4-(2-hydroxyethyl)-1-piperazineethane sulfonic acid
HPLC	high performance liquid chromatography
LAH	lithium aluminum hydride
LDA	lithium diisopropylamide
LHMDS	lithium bis(trimethylsilyl)amide
mCPBA	<i>meta</i> -chloroperoxybenzoic acid
MeOH	methanol
MHz	megahertz
min	minute or minutes
MS	mass spectrometry
Ms	methanesulfonyl
NMM	N-methylmorpholine
NMR	nuclear magnetic resonance
PCC	pyridinium chlorochromate
RP	reverse-phase
RT	room temperature
R _t	retention time
Sec	second or seconds
TBDMS	t-butyldimethylsilyl
TEA or Et ₃ N	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TIPS	triisopropylsilyl
TLC	thin layer chromatography
TMS	tetramethylsilane

Scheme A illustrates a route for the synthesis of compounds of Formula (I)-A and intermediate A9, wherein Q, R¹, R², and R³ are as defined herein, and G is t-butoxycarbonyl group.


Scheme A



Compound **A1** is either commercially available or may be prepared by methods known in the scientific literature. The compound **A1** may be alkylated with propargyl bromide in the presence of a strong base, such as sodium hydride, to afford compound **A3**. Compound **A3** may be cross-coupled with an aryl or heteroaryl iodide of formula **A4** (wherein Q is as defined herein) in the presence of a palladium catalyst, appropriate ligands, appropriate coupling reagents such as copper iodide, and in the presence of a tertiary amine base such as triethylamine, to afford a compound of formula **A5**. The compound of formula **A5** may be coupled with a substituted hydrazine of formula **A6**, in the presence of an appropriate amide-type coupling reagents such as EDC and HOBt, or with DCC, to afford a compound of formula **A7**. Treatment of a compound of formula **A7** with triphenylphosphine and carbon tetrachloride, or with POCl₃ affords a compound of formula **A8**, which, upon heating to about 100°C in the presence of a tertiary amine base such as triethylamine, affords a compound of formula **(I)-A**. The Boc function group of a compound of formula **(I)-A** may be removed by treatment with a strong acid, such as trifluoroacetic acid, or a mineral acid such as hydrochloric acid to afford an intermediate compound of formula **A9**.

Scheme B illustrates a route for the synthesis of compounds of Formula (I)-B, wherein Q, R¹, R², and R³ are as defined herein, and G is a sulfonyl group, wherein R^B is a substituent as defined by the present invention, including, but not limited to, C₁₋₆alkyl, C₆₋₁₀aryl(C₁₋₄)alkyl, C₆₋₁₀aryloxy(C₁₋₄)alkyl, 3,5-dimethylisoxazol-4-yl, C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl, trifluoromethyl, C₃₋₇cycloalkyl, 2,2,2-trifluoroethyl, di(C₁₋₄alkyl)amino, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl, and 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl.


Scheme B



The compound of formula **A9** may be treated with an R^B-substituted sulfonyl chloride in the presence of an organic amine base, such as triethylamine, to afford a compound of formula **(I)-B**.

Scheme C illustrates a route for the synthesis of compounds of Formula (I)-C, wherein Q, R¹, R², and R³ are as defined herein, and G is a carbonyl group, wherein R^C is a substituent as defined by the present invention, including, but not limited to, methoxymethyl, trifluoromethyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl, C₁₋₄alkoxycarbonylaminomethyl, C₁₋₄alkylsulfonylaminomethyl, C₁₋₄alkoxy, di(C₁₋₄alkyl)amino, and pyridinyl.


Scheme C



The compound of formula **A9** may be coupled with an R^C -substituted carboxylic acid of formula **C1** in the presence of an appropriate amide-type coupling agent such as polystyrene-supported CDI, and in the presence of a coupling activating agent such as HOBt, to afford a compound of formula **(I)-C**.


Scheme D illustrates a route for the synthesis of compounds of Formula **(I)-D**, wherein Q, R^1 , R^2 , and R^3 are as defined herein, and G is a heteroaryl or aryl ring (Ar_D) as defined by G of the present invention.

Scheme D



The compound of formula **A9** may be coupled with an aryl or heteroaryl (Ar_D) iodide of formula **D1** in the presence of a transition metal catalyst such as palladium (II), appropriate ligands, and an inorganic base such as potassium or cesium carbonate, to afford a compound of formula **(I)-D**.


Scheme E illustrates a route for the synthesis of compounds of Formula **(I)-E**, wherein Q, R^1 , R^2 , and R^3 , and G is as defined herein.

Scheme E

The compound **A3** may be brominated in the presence of a brominating reagent such as bromine, to afford compound **E1**. The compound **E1** may be coupled as with a compound of formula **A6**, using the methods described in Scheme A, to afford a compound of formula **E2**. A compound of formula **E2** may be converted to a compound of formula **E3** using the methods described herein, and, upon heating to about 100°C in the presence of a base, the compound of formula **E3** may be cyclized to a compound of formula **E4**. Upon amino deprotection by conventional methods known to one of skill in the art, a G-group of the present invention may be introduced by one of the methods described in the schemes hereinabove to afford a compound of formula **E6**. A compound of formula **E6** may be cross-coupled with an aryl or heteroaryl boronic acid of formula **E7**, in the presence of a transition metal catalyst such as palladium (II) acetate, appropriate ligands, and an inorganic base such as sodium carbonate, to afford a compound of formula (I)-E.

Specific Examples

Example 1



A. To a solution of N-Boc-glycine (**1a**) (3.0 g, 17.1 mmol, 1.0 eq) in DMF (40 mL) at 0°C under an Argon atmosphere was added sodium hydride (60% dispersion in oil, 2.12 g, 53.1 mmol, 3.5 eq) in 2 portions, 10 minutes apart. After stirring 1.5 hrs, the bubbling had slowed and propargyl bromide (**1b**) (80% in toluene, 3.24 mL, 29.1 mmol, 1.7 eq) was added. The grey solution was allowed to warm to rt overnight. Water was added carefully, and the solution brought to pH 3 with 1 N HCl. The aqueous phase was extracted with DCM, dried over MgSO_4 , filtered, and concentrated. Purification by column chromatography (80 g), eluting with 30 to 50% ethyl acetate/hexanes + 0.1% HOAc, afforded compound **1c** (3.17 g, 87%). ^1H NMR (CHLOROFORM- d) δ : 4.06 - 4.28 (m, 4H), 2.27 (t, J = 2.4 Hz, 1H), 1.42 - 1.52 (m, 9H).

B. To a solution of compound **1c** (2.83 g, 13.3 mmol, 1.0 eq) in DMF (30 mL) at rt was added triethylamine (30 mL) followed by 4-chloro-1-iodobenzene (**1d**) (3.49 g, 14.6 mmol, 1.1 eq), $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ (467 mg, 0.67 mmol, 0.05 eq) and copper iodide (253 mg, 1.33 mmol, 0.1 eq). The solution was placed under an Argon atmosphere and heated to 80°C. After 2 hrs, the solution was concentrated, dissolved in DCM, washed sequentially with 1 N HCl and saturated NaHCO_3 , dried over MgSO_4 , filtered, and concentrated.

Purification by column chromatography (150 g), eluting with 20 to 40% ethyl acetate/hexanes, afforded compound **1e** (2.65 g, 62%). ¹H NMR (CHLOROFORM-d) δ: 7.30 - 7.36 (m, 2H), 7.27 - 7.30 (m, 2H), 4.39 (d, J = 30.8 Hz, 2H), 4.19 (d, J = 25.5 Hz, 2H), 1.42 - 1.52 (m, 9H)


C. To a solution of compound **1e** (1.33 g, 4.10 mmol, 1.0 eq) in acetonitrile (30 mL) was added EDCI (1.18 g, 6.15 mmol, 1.5 eq) and hydroxybenzotriazole hydrate (831 mg, 6.15 mmol, 1.5 eq). After 15 min, 2-ethylphenylhydrazine (**1f**) (976 mg, 7.17 mmol, 1.75 eq) was added. After 1 hr, water and 1 N HCl were added, the aqueous phase was extracted with DCM, the combined organic phases were washed with NaHCO₃, dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (115 g), eluting with 10 to 30 to 40% EA/hexanes, gave compound **1g** (1.19 g, 66%). ESI-MS (m/z): Calcd. for C₂₄H₂₈ClN₃O₃: 464.2 (M+23); found: 464.0.

D. To a solution of compound **1g** (1.19 g, 2.69 mmol, 1.0 eq) in acetonitrile (75 mL) was added triphenylphosphine (2.11 g, 8.06 mmol, 3.0 eq) followed by carbon tetrachloride (0.78 mL, 8.06 mmol, 3.0 eq). After 2 hrs, brine was added and the reaction mixture was extracted with EA. The combined organic phases were dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (80 g), eluting with 0 to 10% EA/hexanes, compound **1h** (718 mg, 58%). ¹H NMR (CHLOROFORM-d) δ: 7.74 (br. s., 1H), 7.37 (d, J = 7.8 Hz, 1H), 7.27 - 7.33 (m, 2H), 7.19 - 7.27 (m, 2H), 7.07 - 7.19 (m, 2H), 6.85 - 6.93 (m, 1H), 4.22 - 4.57 (m, 4H), 2.54 (q, J = 7.6 Hz, 2H), 1.42 - 1.58 (m, 9H), 1.25 (t, J = 7.5 Hz, 3H). ESI-MS (m/z): Calcd. for C₂₄H₂₇Cl₂N₃O₂: 482.2 (M+23); found: 481.9.

E. A solution of compound **1h** (718 mg, 1.56 mmol, 1.0 eq) and triethylamine (0.87 mL, 6.25 mmol, 4.0 eq) in toluene (20 mL) were heated to 100 °C overnight. The solution was cooled, water was added, and the aqueous phase was extracted with EA. The organic phases were combined, dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (40 g), eluting with 0 to 15% EA/hexanes, gave compound **1** (600 mg, 91%). ¹H NMR (CHLOROFORM-d) δ: 7.36 - 7.42 (m, 1H), 7.31 - 7.35 (m, 1H), 7.17 - 7.28 (m, 4H), 6.95 - 7.01 (m, 2H), 4.54 - 4.68 (m, 4H), 2.30 - 2.40 (m, 2H), 1.54 (s, 9H), 1.01 (t, J = 7.6 Hz, 3H). ESI-MS (m/z): Calcd. for C₂₄H₂₆ClN₃O₂: 424.2 (M+1); found: 424.0


F. To a solution of compound **1** (600 mg, 1.42 mmol, 1.0 eq) in DCM (20 mL) was added trifluoroacetic acid (2 mL). After 2 hrs, the solution was concentrated. DCM was added, the solution was washed with NaHCO₃, dried over MgSO₄, filtered, and concentrated to give compound **1i** (396 mg, 86%). ¹H NMR (CHLOROFORM-d) δ : 7.35 - 7.41 (m, 1H), 7.32 (d, J = 7.6 Hz, 1H), 7.16 - 7.26 (m, 4H), 6.95 - 7.01 (m, 2H), 4.22 (s, 2H), 4.14 (s, 2H), 2.38 (q, J = 7.6 Hz, 2H), 0.98 - 1.04 (m, 3H). ESI-MS (m/z): Calcd. for C₁₉H₁₈ClN₃: 324.1 (M+1); found: 324.2.

Following the procedure described above for Example 1 and substituting the appropriate reagents, starting materials and purification methods known to those skilled in the art, the following compounds were prepared:




Cpd 19

Cpd **19**: ¹H NMR (CHLOROFORM-d) δ : 7.55 (ddd, J = 10.0, 7.8, 1.9 Hz, 1H), 7.43 - 7.49 (m, 1H), 7.36 - 7.42 (m, 1H), 7.21 - 7.33 (m, 3H), 6.98 - 7.06 (m, 2H), 4.62 (s, 2H), 4.57 (d, J = 4.3 Hz, 2H), 1.53 (d, J = 2.0 Hz, 9H). ESI-MS (m/z): Calcd. for C₂₃H₂₁ClF₃N₃O₃: 480.1 (M+1); found: 480.1.




Cpd 28

Cpd **28**: ¹H NMR (CHLOROFORM-d) δ : 7.34 - 7.43 (m, 2H), 7.23 (dd, J = 8.5, 6.7 Hz, 2H), 7.01 - 7.09 (m, 3H), 6.88 - 6.94 (m, 1H), 4.62 (s, 2H), 4.56 (d, J = 9.6 Hz, 2H), 3.52 (s, 3H), 1.53 (s, 9H). ESI-MS (m/z): Calcd. for C₂₃H₂₄ClN₃O₃: 426.2 (M+1); found: 426.2.

Example 2


To a solution of compound **1i** (18.5 mg, 0.06 mmol, 1.0 eq) in DCM (1.5 mL) was added methanesulfonylchloride (6.7 μ L, 0.09 mmol, 1.5 eq) and triethylamine (15.9 μ L, 0.12 mmol, 2.0 eq). After 1 hr, DCM was added, and the solution was washed with saturated aqueous NaHCO_3 . The organic phase was dried over MgSO_4 , filtered, and concentrated. Purification by column chromatography (8 g), eluting with 20 to 40% EA/hexanes, gave compound **2** (12.8 mg, 56%). ^1H NMR (CHLOROFORM- d) δ : 7.39 - 7.45 (m, 1H), 7.32 - 7.37 (m, 1H), 7.17 - 7.30 (m, 4H), 6.93 - 6.99 (m, 2H), 4.70 (s, 2H), 4.63 (s, 2H), 2.98 (s, 3H), 2.35 (q, J = 7.6 Hz, 2H), 1.02 (t, J = 7.6 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{20}\text{ClN}_3\text{O}_2\text{S}$: 402.1 ($M+1$); found: 402.1.

Following the procedure described above for Example 2 and substituting the appropriate reagents, starting materials and purification methods known to those skilled in the art, the following compounds were prepared:




Cpd 3

Cpd 3: ^1H NMR (CHLOROFORM- d) δ : 7.39 - 7.46 (m, 1H), 7.35 (d, J = 6.6 Hz, 1H), 7.17 - 7.31 (m, 4H), 6.92 - 6.98 (m, 2H), 4.88 (s, 2H), 4.82 (s, 2H), 2.35 (q, J = 7.6 Hz, 2H), 1.03 (t, J = 7.6 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{17}\text{ClF}_3\text{N}_3\text{O}_2\text{S}$: 456.1 ($M+1$); found: 456.1.




Cpd 15

Cpd 15: ^1H NMR (CHLOROFORM- d) δ : 7.84 (dd, J = 5.8, 3.8 Hz, 1H), 7.58 - 7.65 (m, 2H), 7.29 (dd, J = 5.6, 3.5 Hz, 1H), 7.21 - 7.26 (m, 2H), 6.93 - 7.00 (m, 2H), 4.68 (s, 2H), 4.64 (t, J = 1.5 Hz, 2H), 2.98 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{19}\text{H}_{15}\text{ClF}_3\text{N}_3\text{O}_2\text{S}$: 442.1 ($M+1$); found: 442.1.




Cpd 16

Cpd 16: ^1H NMR (CHLOROFORM- d) δ : 7.83 (dd, J = 5.6, 3.8 Hz, 1H), 7.57 - 7.64 (m, 2H), 7.29 (dd, J = 5.4, 3.7 Hz, 1H), 7.20 - 7.25 (m, 2H), 6.93 - 6.99 (m, 2H), 4.75 (br. s., 2H), 4.70 (s, 2H), 3.40 (quin, J = 6.8 Hz, 1H), 1.46 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{19}\text{ClF}_3\text{N}_3\text{O}_2\text{S}$: 470.1 ($M+1$); found: 470.1.




Cpd 17

Cpd 17: ^1H NMR (CHLOROFORM- d) δ : 7.75 - 7.85 (m, 3H), 7.53 - 7.63 (m, 2H), 7.36 (d, J = 8.1 Hz, 2H), 7.17 - 7.25 (m, 3H), 6.87 - 6.95 (m, 2H), 4.59 (s, 2H), 4.56 (s, 2H), 2.44 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{19}\text{ClF}_3\text{N}_3\text{O}_2\text{S}$: 518.1 ($M+1$); found: 518.1.




Cpd 20

Cpd 20: ^1H NMR (CHLOROFORM- d) δ : 7.54 (dd, $J = 7.7, 1.9$ Hz, 1H), 7.48 (td, $J = 7.9, 1.9$ Hz, 1H), 7.38 - 7.45 (m, 1H), 7.23 - 7.33 (m, 3H), 6.97 - 7.03 (m, 2H), 4.65 (d, $J = 4.8$ Hz, 4H), 2.97 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{19}\text{H}_{15}\text{ClF}_3\text{N}_3\text{O}_3\text{S}$: 458.1 ($M+1$); found: 458.1.




Cpd 30

Cpd 30: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.44 (m, 2H), 7.21 - 7.26 (m, 2H), 7.00 - 7.09 (m, 3H), 6.92 (d, $J = 8.1$ Hz, 1H), 4.58 - 4.72 (m, 4H), 3.53 (s, 3H), 2.95 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{19}\text{H}_{18}\text{ClN}_3\text{O}_3\text{S}$: 404.1 ($M+1$); found: 404.1.




Cpd 31

Cpd 31: ^1H NMR (CHLOROFORM- d) δ : 7.35 - 7.44 (m, 2H), 7.19 - 7.25 (m, 2H), 6.99 - 7.09 (m, 3H), 6.88 - 6.95 (m, 1H), 4.63 - 4.79 (m, 4H), 3.54 (s, 3H), 3.38 (quin, $J = 6.8$ Hz, 1H), 1.45 (d, $J = 6.8$ Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{22}\text{ClN}_3\text{O}_3\text{S}$: 432.1 ($M+1$); found: 432.1.




Cpd 32

Cpd 32: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.26 (m, 2H), 7.00 - 7.09 (m, 3H), 6.89 - 6.95 (m, 1H), 4.64 - 4.77 (m, 4H), 3.54 (s, 3H), 2.42 - 2.51 (m, 1H), 1.26 - 1.32 (m, 2H), 0.99 - 1.06 (m, 2H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{20}\text{ClN}_3\text{O}_3\text{S}$: 430.1 ($M+1$); found: 430.1.




Cpd 33

Cpd 33: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 7.00 - 7.08 (m, 3H), 6.89 - 6.94 (m, 1H), 4.63 (s, 2H), 4.59 (s, 2H), 3.53 (s, 3H), 2.90 (s, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{21}\text{ClN}_4\text{O}_3\text{S}$: 433.1 ($M+1$); found: 433.1.




Cpd 35

Cpd 35: ^1H NMR (CHLOROFORM- d) δ : 7.55 (dd, J = 7.6, 1.8 Hz, 1H), 7.48 (td, J = 7.8, 1.8 Hz, 1H), 7.38 - 7.44 (m, 1H), 7.22 - 7.33 (m, 3H), 6.96 - 7.03 (m, 2H), 4.71 (d, J = 10.6 Hz, 4H), 3.39 (quin, J = 6.8 Hz, 1H), 1.46 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{19}\text{ClF}_3\text{N}_3\text{O}_3\text{S}$: 486.1 ($M+1$); found: 486.1.




Cpd 36

Cpd 36: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.44 (m, 2H), 7.21 - 7.26 (m, 2H), 6.99 - 7.09 (m, 3H), 6.89 - 6.95 (m, 1H), 4.62 - 4.81 (m, 4H), 3.88 (q, J = 9.3 Hz, 2H), 3.53 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{17}\text{ClF}_3\text{N}_3\text{O}_3\text{S}$: 472.1 ($M+1$); found: 472.1.




Cpd 37

Cpd 37: ^1H NMR (CHLOROFORM- d) δ : 7.35 - 7.44 (m, 2H), 7.19 - 7.25 (m, 2H), 6.99 - 7.08 (m, 3H), 6.89 - 6.95 (m, 1H), 4.61 - 4.79 (m, 4H), 3.53 (s, 3H), 3.04 - 3.16 (m, 1H), 2.17 - 2.27 (m, 2H), 1.91 (d, J = 12.9 Hz, 2H), 1.59 - 1.76 (m, 3H), 1.17 - 1.37 (m, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{26}\text{ClN}_3\text{O}_3\text{S}$: 472.1 ($M+1$); found: 472.1.




Cpd 38

Cpd 38: ^1H NMR (CHLOROFORM- d) δ : 7.82 - 7.89 (m, 2H), 7.68 - 7.76 (m, 2H), 7.35 - 7.44 (m, 2H), 7.20 - 7.26 (m, 2H), 6.98 - 7.09 (m, 3H), 6.91 (d, J = 8.3 Hz, 1H), 4.60 - 4.76 (m, 4H), 4.23 (t, J = 6.9 Hz, 2H), 3.47 - 3.57 (m, 5H). ESI-MS (m/z): Calcd. for $\text{C}_{28}\text{H}_{23}\text{ClN}_4\text{O}_5\text{S}$: 563.1 ($M+1$); found: 563.1.




Cpd 39

Cpd 39: ^1H NMR (CHLOROFORM- d) δ : 7.30 - 7.44 (m, 7H), 7.17 - 7.23 (m, 2H), 7.02 - 7.08 (m, 1H), 6.89 - 6.95 (m, 3H), 4.47 (s, 2H), 4.37 (s, 4H), 3.54 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{22}\text{ClN}_3\text{O}_3\text{S}$: 480.1 ($M+1$); found: 480.1.




Cpd 40

Cpd 40: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 6.99 - 7.08 (m, 3H), 6.89 - 6.95 (m, 1H), 4.65 - 4.77 (m, 4H), 3.59 - 3.70 (m, 1H), 3.54 (s, 3H), 1.98 - 2.20 (m, 4H), 1.78 - 1.91 (m, 2H), 1.59 - 1.73 (m, 2H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{24}\text{ClN}_3\text{O}_3\text{S}$: 458.1 ($M+1$); found: 458.1.




Cpd 41

Cpd 41: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.44 (m, 2H), 7.20 - 7.25 (m, 2H), 7.00 - 7.08 (m, 3H), 6.92 (dd, $J = 8.7, 1.1$ Hz, 1H), 4.64 - 4.78 (m, 4H), 3.54 (s, 3H), 3.21 (td, $J = 6.7, 3.5$ Hz, 1H), 1.96 - 2.08 (m, 1H), 1.49 - 1.69 (m, 2H), 1.43 (d, $J = 6.8$ Hz, 3H), 1.32 - 1.41 (m, 1H), 0.96 (t, $J = 7.2$ Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{26}\text{ClN}_3\text{O}_3\text{S}$: 460.1 ($M+1$); found: 460.1.




Cpd 42

Cpd 42: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.44 (m, 2H), 7.21 - 7.26 (m, 2H), 7.01 - 7.08 (m, 3H), 6.89 - 6.94 (m, 1H), 4.60 - 4.75 (m, 4H), 4.04 (t, J = 6.6 Hz, 2H), 3.53 (s, 3H), 3.44 (t, J = 6.7 Hz, 2H), 2.73 (s, 4H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{23}\text{ClN}_4\text{O}_5\text{S}$: 515.1 ($M+1$); found: 515.1.




Cpd 43

Cpd 43: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.45 (m, 2H), 7.15 - 7.24 (m, 4H), 7.03 - 7.10 (m, 1H), 6.88 - 6.97 (m, 4H), 6.71 (d, J = 7.8 Hz, 2H), 4.68 (d, J = 12.6 Hz, 4H), 4.44 (t, J = 5.6 Hz, 2H), 3.62 (t, J = 5.6 Hz, 2H), 3.56 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{26}\text{H}_{24}\text{ClN}_3\text{O}_4\text{S}$: 510.1 ($M+1$); found: 510.1.




Cpd 44

Cpd 44: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.45 (m, 2H), 7.24 (d, J = 8.3 Hz, 2H), 7.00 - 7.10 (m, 3H), 6.92 (d, J = 8.3 Hz, 1H), 5.19 - 5.28 (m, 1H), 4.61 - 4.75 (m, 4H), 3.65 - 3.74 (m, 2H), 3.54 (s, 3H), 3.34 (dd, J = 6.6, 4.8 Hz, 2H), 3.01 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{23}\text{ClN}_4\text{O}_5\text{S}_2$: 511.1 ($M+1$); found: 511.1.




Cpd 45

Cpd 45: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.45 (m, 2H), 7.21 - 7.26 (m, 2H), 7.00 - 7.10 (m, 3H), 6.92 (d, J = 8.6 Hz, 1H), 5.02 (br. s., 1H), 4.68 (d, J = 11.9 Hz, 4H), 3.65 - 3.75 (m, 2H), 3.54 (s, 3H), 3.33 (dd, J = 6.6, 4.8 Hz, 2H), 3.15 - 3.25 (m, 1H), 1.40 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{27}\text{ClN}_4\text{O}_5\text{S}_2$: 539.1 ($M+1$); found: 539.1.



Cpd 46


Cpd 46: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.44 (m, 2H), 7.21 - 7.26 (m, 2H), 7.00 - 7.09 (m, 3H), 6.92 (d, J = 7.8 Hz, 1H), 5.06 (t, J = 6.4 Hz, 1H), 4.62 - 4.73 (m, 4H), 3.58 - 3.66 (m, 2H), 3.54 (s, 3H), 3.35 (dd, J = 6.6, 4.8 Hz, 2H), 2.83 (s, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{26}\text{ClN}_5\text{O}_5\text{S}_2$: 540.1 ($M+1$); found: 540.1.



Cpd 47


Cpd 47: ^1H NMR (CHLOROFORM- d) δ : 7.54 (dd, J = 7.8, 1.8 Hz, 1H), 7.48 (td, J = 7.8, 1.8 Hz, 1H), 7.38 - 7.45 (m, 1H), 7.23 - 7.33 (m, 3H), 6.97 - 7.03 (m, 2H), 4.66 (d, J

= 4.5 Hz, 4H), 3.66 (s, 3H), 3.47 (t, $J = 7.3$ Hz, 2H), 2.91 (t, $J = 7.3$ Hz, 2H). ESI-MS (m/z): Calcd. for $C_{22}H_{19}ClF_3N_3O_5S$: 530.1 ($M+1$); found: 530.1.




Cpd 48

Cpd 48: ^1H NMR (CHLOROFORM- d) δ : 7.52 - 7.57 (m, 1H), 7.49 (td, $J = 7.9, 1.9$ Hz, 1H), 7.42 (td, $J = 7.6, 1.4$ Hz, 1H), 7.24 - 7.34 (m, 3H), 6.96 - 7.02 (m, 2H), 4.65 - 4.79 (m, 4H), 4.02 - 4.16 (m, 1H), 3.47 (d, $J = 9.1$ Hz, 2H), 3.36 - 3.45 (m, 1H), 3.16 (dt, $J = 13.4, 8.0$ Hz, 1H), 2.60 - 2.76 (m, 2H). ESI-MS (m/z): Calcd. for $C_{22}H_{19}ClF_3N_3O_5S_2$: 562.0 ($M+1$); found: 562.0.



Cpd 49


Cpd 49: ^1H NMR (CHLOROFORM- d) δ : 7.47 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.34 - 7.41 (m, 1H), 7.23 (d, $J = 8.6$ Hz, 2H), 7.00 - 7.10 (m, 3H), 6.87 (d, $J = 8.3$ Hz, 1H), 4.63 (br. s., 4H), 3.75 (br. s., 2H), 2.96 (s, 3H), 1.01 (t, $J = 6.9$ Hz, 3H). ESI-MS (m/z): Calcd. for $C_{20}H_{20}ClN_3O_3S$: 418.1 ($M+1$); found: 418.1.



Cpd 50


Cpd 50: ^1H NMR (CHLOROFORM- d) δ : 7.47 (dd, $J = 7.7, 1.6$ Hz, 1H), 7.34 - 7.40 (m, 1H), 7.19 - 7.25 (m, 2H), 7.00 - 7.09 (m, 3H), 6.84 - 6.89 (m, 1H), 4.69 (br. s.,

4H), 3.75 (br. s., 2H), 3.39 (quin, $J = 6.8$ Hz, 1H), 1.45 (d, $J = 6.8$ Hz, 6H), 1.02 (t, $J = 7.1$ Hz, 3H). ESI-MS (m/z): Calcd. for $C_{22}H_{24}ClN_3O_3S$: 446.1 ($M+1$); found: 446.1.




Cpd 51

Cpd 51: ^1H NMR (CHLOROFORM- d) δ : 7.47 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.37 (td, $J = 8.0, 1.8$ Hz, 1H), 7.19 - 7.25 (m, 2H), 7.01 - 7.09 (m, 3H), 6.83 - 6.90 (m, 1H), 4.68 (br. s., 4H), 3.59 - 3.91 (m, 3H), 1.99 - 2.21 (m, 4H), 1.78 - 1.92 (m, 2H), 1.59 - 1.71 (m, 2H), 1.01 (t, $J = 6.9$ Hz, 3H). ESI-MS (m/z): Calcd. for $C_{24}H_{26}ClN_3O_3S$: 472.1 ($M+1$); found: 472.1.




Cpd 52

Cpd 52: ^1H NMR (CHLOROFORM- d) δ : 7.47 (dd, $J = 7.7, 1.6$ Hz, 1H), 7.37 (td, $J = 8.0, 1.8$ Hz, 1H), 7.19 - 7.25 (m, 2H), 7.00 - 7.09 (m, 3H), 6.84 - 6.89 (m, 1H), 4.54 - 4.91 (m, 4H), 3.74 (br. s., 2H), 3.11 (tt, $J = 12.1, 3.4$ Hz, 1H), 2.16 - 2.26 (m, 2H), 1.91 (d, $J = 12.9$ Hz, 2H), 1.59 - 1.76 (m, 3H), 1.15 - 1.38 (m, 3H), 1.01 (t, $J = 6.9$ Hz, 3H). ESI-MS (m/z): Calcd. for $C_{25}H_{28}ClN_3O_3S$: 486.2 ($M+1$); found: 486.2.




Cpd 53


Cpd **53**: ^1H NMR (CHLOROFORM- d) δ : 7.46 (dd, $J = 7.7, 1.6$ Hz, 1H), 7.29 - 7.42 (m, 6H), 7.16 - 7.23 (m, 2H), 7.06 (td, $J = 7.6, 1.1$ Hz, 1H), 6.90 - 6.97 (m, 2H), 6.84 - 6.90 (m, 1H), 4.23 - 4.56 (m, 6H), 3.76 (br. s., 2H), 1.04 (t, $J = 7.1$ Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{26}\text{H}_{24}\text{ClN}_3\text{O}_3\text{S}$: 494.1 ($M+1$); found: 494.1.

Cpd **54**

Cpd **54**: ^1H NMR (CHLOROFORM- d) δ : 7.47 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.34 - 7.40 (m, 1H), 7.19 - 7.25 (m, 2H), 7.01 - 7.09 (m, 3H), 6.84 - 6.89 (m, 1H), 4.59 (br. s., 4H), 3.74 (br. s., 2H), 2.90 (s, 6H), 1.02 (t, $J = 7.1$ Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{23}\text{ClN}_4\text{O}_3\text{S}$: 447.1 ($M+1$); found: 447.1.


Cpd **55**

Cpd **55**: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 7.00 - 7.07 (m, 3H), 6.88 - 6.94 (m, 1H), 4.48 - 4.65 (m, 4H), 3.53 (s, 3H), 3.40 - 3.47 (m, 4H), 1.72 - 1.83 (m, 4H), 1.62 - 1.70 (m, 4H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{27}\text{ClN}_4\text{O}_3\text{S}$: 487.2 ($M+1$); found: 487.2.

Cpd **56**


Cpd **56**: ^1H NMR (CHLOROFORM- d) δ : 7.35 - 7.43 (m, 2H), 7.19 - 7.25 (m, 2H), 7.00 - 7.07 (m, 3H), 6.91 (dd, $J = 8.7, 1.1$ Hz, 1H), 4.49 - 4.65 (m, 4H), 3.53 (s, 3H), 3.35

(q, $J = 7.2$ Hz, 4H), 1.23 (t, $J = 7.2$ Hz, 6H). ESI-MS (m/z): Calcd. for $C_{22}H_{25}ClN_4O_3S$: 461.1 ($M+1$); found: 461.1.




Cpd 57

Cpd 57: ^1H NMR (CHLOROFORM- d) δ : 7.35 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 6.99 - 7.08 (m, 3H), 6.89 - 6.94 (m, 1H), 4.52 - 4.68 (m, 4H), 3.70 - 3.80 (m, 2H), 3.53 (s, 3H), 2.78 - 2.90 (m, 2H), 1.68 - 1.77 (m, 2H), 1.43 - 1.57 (m, 1H), 1.20 - 1.34 (m, 2H), 0.97 (d, $J = 6.6$ Hz, 3H). ESI-MS (m/z): Calcd. for $C_{24}H_{27}ClN_4O_3S$: 487.2 ($M+1$); found: 487.2.




Cpd 58

Cpd 58: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 7.00 - 7.08 (m, 3H), 6.92 (dd, $J = 8.7, 1.1$ Hz, 1H), 4.54 - 4.68 (m, 4H), 3.67 - 3.77 (m, 5H), 3.53 (s, 3H), 2.92 - 3.03 (m, 2H), 2.41 - 2.51 (m, 1H), 1.97 - 2.06 (m, 2H), 1.75 - 1.88 (m, 2H). ESI-MS (m/z): Calcd. for $C_{25}H_{27}ClN_4O_5S$: 531.1 ($M+1$); found: 531.1.




Cpd 59


Cpd **59**: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.44 (m, 2H), 7.20 - 7.26 (m, 2H), 6.99 - 7.08 (m, 3H), 6.92 (d, J = 7.8 Hz, 1H), 4.57 - 4.72 (m, 4H), 3.73 - 3.80 (m, 4H), 3.53 (s, 3H), 3.27 - 3.35 (m, 4H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{23}\text{ClN}_4\text{O}_4\text{S}$: 475.1 ($M+1$); found: 475.1.

Cpd **60**

Cpd **60**: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 7.00 - 7.08 (m, 3H), 6.92 (dd, J = 8.7, 1.1 Hz, 1H), 4.52 - 4.70 (m, 4H), 3.53 (s, 3H), 3.36 - 3.43 (m, 4H), 1.91 - 1.99 (m, 4H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{23}\text{ClN}_4\text{O}_3\text{S}$: 459.1 ($M+1$); found: 459.1.


Cpd **61**

Cpd **61**: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.44 (m, 2H), 7.20 - 7.26 (m, 2H), 6.99 - 7.08 (m, 3H), 6.92 (d, J = 7.8 Hz, 1H), 4.53 - 4.70 (m, 4H), 3.70 (ddd, J = 10.4, 6.3, 2.3 Hz, 2H), 3.50 - 3.58 (m, 5H), 2.60 (dd, J = 12.1, 10.6 Hz, 2H), 1.21 (d, J = 6.3 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{27}\text{ClN}_4\text{O}_4\text{S}$: 503.2 ($M+1$); found: 503.2.




Cpd 62

Cpd 62: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 2H), 7.20 - 7.25 (m, 2H), 7.00 - 7.08 (m, 3H), 6.92 (dd, J = 8.7, 1.1 Hz, 1H), 4.53 - 4.69 (m, 4H), 3.53 (s, 3H), 3.24 - 3.33 (m, 4H), 1.52 - 1.71 (m, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{25}\text{ClN}_4\text{O}_3\text{S}$: 473.1 ($M+1$); found: 473.1.




Cpd 63

Cpd 63: ^1H NMR (CHLOROFORM- d) δ : 7.51 (dd, J = 7.8, 1.8 Hz, 1H), 7.35 (td, J = 8.0, 1.8 Hz, 1H), 7.19 - 7.25 (m, 2H), 7.01 - 7.08 (m, 3H), 6.84 (d, J = 7.8 Hz, 1H), 4.59 (br. s., 4H), 4.28 (dt, J = 12.1, 6.1 Hz, 1H), 2.91 (s, 6H), 0.67 - 1.17 (m, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{25}\text{ClN}_4\text{O}_3\text{S}$: 461.1 ($M+1$); found: 461.1.




Cpd 87

Cpd 87: ^1H NMR (CHLOROFORM- d) δ : 7.34 - 7.43 (m, 2H), 7.20 - 7.26 (m, 2H), 6.97 - 7.07 (m, 3H), 6.87 - 6.93 (m, 1H), 4.54 - 4.66 (m, 4H), 3.52 (s, 3H), 2.72 (s, 3H), 2.48 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{21}\text{ClN}_4\text{O}_4\text{S}$: 485.1 ($M+1$); found: 485.1.




Cpd 88


Cpd **88**: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.43 (m, 1H), 7.31 (td, J = 7.8, 1.6 Hz, 1H), 7.16 - 7.22 (m, 2H), 7.02 (td, J = 7.6, 1.3 Hz, 1H), 6.93 - 6.99 (m, 2H), 6.84 (dd, J = 8.2, 1.1 Hz, 1H), 4.72 - 4.92 (m, 2H), 4.62 (d, J = 11.9 Hz, 2H), 3.39 (quin, J = 6.8 Hz, 1H), 2.22 (s, 6H), 1.45 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{25}\text{ClN}_4\text{O}_2\text{S}$: 445.1 ($M+1$); found: 445.1.

Cpd **89**


Cpd **89**: ^1H NMR (CHLOROFORM- d) δ : 7.40 (dd, J = 7.7, 1.6 Hz, 1H), 7.28 - 7.34 (m, 1H), 7.17 - 7.22 (m, 2H), 6.99 - 7.06 (m, 1H), 6.94 - 6.99 (m, 1H), 6.84 (dd, J = 8.2, 1.1 Hz, 1H), 4.68 - 4.86 (m, 2H), 4.52 - 4.65 (m, 2H), 3.15 (q, J = 7.4 Hz, 2H), 2.22 (s, 6H), 1.45 (t, J = 7.5 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{23}\text{ClN}_4\text{O}_2\text{S}$: 431.1 ($M+1$); found: 431.1.

Cpd **90**

Cpd **90**: ^1H NMR (CHLOROFORM- d) δ : 7.52 (dd, J = 7.7, 1.6 Hz, 1H), 7.43 (td, J = 8.0, 1.8 Hz, 1H), 7.17 - 7.25 (m, 3H), 6.98 - 7.04 (m, 2H), 6.93 (d, J = 7.8 Hz, 1H), 4.54 - 4.69 (m, 4H), 4.05 (q, J = 8.2 Hz, 2H), 2.91 (s, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{20}\text{ClF}_3\text{N}_4\text{O}_3\text{S}$: 501.1 ($M+1$); found: 501.1.

Cpd **91**


Cpd 91: ^1H NMR (CHLOROFORM- d) δ : 7.58 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.33 (td, $J = 7.8, 1.8$ Hz, 1H), 7.16 - 7.24 (m, 3H), 6.97 - 7.03 (m, 3H), 4.59 (br. s., 4H), 2.90 (s, 6H), 0.97 - 1.04 (m, 9H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{27}\text{ClN}_4\text{O}_3\text{S}$: 475.2 ($M+1$); found: 475.2.



Cpd 95


Cpd 95: ^1H NMR (CHLOROFORM- d) δ : 7.49 (dd, $J = 7.7, 1.6$ Hz, 1H), 7.38 - 7.44 (m, 1H), 7.20 - 7.25 (m, 2H), 7.16 (td, $J = 7.6, 1.1$ Hz, 1H), 7.00 - 7.05 (m, 2H), 6.91 (d, $J = 8.3$ Hz, 1H), 5.50 - 5.81 (m, 1H), 4.55 - 4.68 (m, 4H), 3.89 (br. s., 2H), 2.91 (s, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{21}\text{ClF}_2\text{N}_4\text{O}_3\text{S}$: 483.1 ($M+1$); found: 483.1.

Example 3




To a solution of compound **1i** (18.5 mg, 0.06 mmol, 1.0 eq) in DCM (1.5 mL) was added nicotinoyl chloride-HCl (15.2 mg, 0.09 mmol, 1.5 eq) and triethylamine (23.8 μL , 0.18 mmol, 3.0 eq). After 1 hr, DCM was added, the solution was washed with saturated aqueous NaHCO_3 , dried over MgSO_4 , filtered, and concentrated. Purification by column chromatography (8 g), eluting with 1 to 3% MeOH/DCM, gave compound **4** (12.2 mg, 50%). ^1H NMR (CHLOROFORM- d) δ : 8.89 (d, $J = 2.3$ Hz, 1H), 8.74 (dd, $J = 4.9, 1.6$ Hz, 1H), 7.89 - 7.97 (m, 1H), 7.31 - 7.48 (m, 3H), 7.13 - 7.30 (m, 4H), 6.97 - 7.06 (m, 1H), 6.85 - 6.94 (m, 1H), 4.99 (d, $J = 24.0$ Hz, 2H), 4.73 (d, $J = 10.6$ Hz, 2H), 2.29 - 2.41 (m, 2H), 1.02 (td, $J = 7.6, 3.0$ Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{21}\text{ClN}_4\text{O}$: 429.1 ($M+1$); found: 429.1.

Example 4




To a solution of 2-furoic acid (7 mg, 0.06 mmol, 1.5 eq) and HOBt (9 mg, 0.07 mmol, 1.7 eq) in DMF (0.33 mL) was added polystyrene supported (PS)-carbodiimide (64 mg, 1.25 mmol/g, 0.08 mmol, 2 eq). After 10 minutes, compound **1i** (13 mg, 0.04 mmol, 1.0 eq) in DCM (1 mL) was added. After 1.5 hr, macroporous (MP)-carbonate resin (65 mg) was added. After 1.5 hrs, the resins were collected by filtration, washed with DCM, and the filtrate was concentrated. Purification by column chromatography (8 g), eluting with 25 to 45% EA/hexanes, gave compound **8** (12.5 mg, 75%). ¹H NMR (CHLOROFORM-d) δ : 7.57 - 7.64 (m, 1H), 7.38 - 7.46 (m, 1H), 7.31 - 7.38 (m, 1H), 7.19 - 7.31 (m, 5H), 6.98 - 7.06 (m, 2H), 6.53 - 6.59 (m, 1H), 5.19 (d, J = 20.5 Hz, 2H), 4.89 - 5.02 (m, 2H), 2.38 (q, J = 7.6 Hz, 2H), 0.98 - 1.08 (m, 3H). ESI-MS (m/z): Calcd. for C₂₄H₂₀ClN₃O₂: 418.1 (M+1); found: 418.1.

Following the procedure described above for Example 4 and substituting the appropriate reagents, starting materials and purification methods known to those skilled in the art, the following compounds were prepared:


Cpd **5**

Cpd **5**: ¹H NMR (CHLOROFORM-d) δ : 8.72 - 8.83 (m, 2H), 7.37 - 7.50 (m, 3H), 7.31 - 7.37 (m, 1H), 7.14 - 7.30 (m, 4H), 6.97 - 7.04 (m, 1H), 6.86 - 6.92 (m, 1H), 5.00 (s, 1H), 4.94 (s, 1H), 4.65 (d, J = 4.0 Hz, 2H), 2.29 - 2.41 (m, 2H), 1.02 (td, J = 7.6, 2.8 Hz, 3H). ESI-MS (m/z): Calcd. for C₂₅H₂₁ClN₄O: 429.1 (M+1); found: 429.1.




Cpd 6

Cpd 6: ^1H NMR (CHLOROFORM- d) δ : 7.39 - 7.46 (m, 1H), 7.32 - 7.38 (m, 1H), 7.18 - 7.30 (m, 5H), 6.95 - 7.01 (m, 2H), 4.98 (d, J = 19.2 Hz, 2H), 4.87 (d, J = 23.0 Hz, 2H), 2.35 (q, J = 7.6 Hz, 2H), 1.03 (td, J = 7.6, 1.0 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{17}\text{ClF}_3\text{N}_3\text{O}$: 420.1 ($M+1$); found: 420.1.




Cpd 10

Cpd 10: ^1H NMR (CHLOROFORM- d) δ : 7.37 - 7.45 (m, 1H), 7.31 - 7.37 (m, 1H), 7.18 - 7.30 (m, 4H), 6.95 - 7.02 (m, 2H), 4.83 (d, J = 8.1 Hz, 2H), 4.75 (d, J = 14.1 Hz, 2H), 4.17 - 4.22 (m, 2H), 3.49 - 3.55 (m, 3H), 2.35 (q, J = 7.4 Hz, 2H), 1.02 (td, J = 7.6, 1.3 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{22}\text{ClN}_3\text{O}_2$: 396.1 ($M+1$); found: 396.1.




Cpd 11

Cpd 11: ^1H NMR (CHLOROFORM- d) δ : 7.38 - 7.46 (m, 1H), 7.32 - 7.37 (m, 1H), 7.24 - 7.30 (m, 2H), 7.19 - 7.24 (m, 2H), 6.93 - 7.01 (m, 2H), 5.53 (br. s., 1H), 4.66 - 4.84 (m, 4H), 4.05 (d, J = 4.5 Hz, 2H), 2.35 (q, J = 7.7 Hz, 2H), 1.47 (s, 9H), 1.01 (t, J = 7.6 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{26}\text{H}_{29}\text{ClN}_4\text{O}_3$: 481.2 ($M+1$); found: 481.2.




Cpd 12

Cpd 12: ^1H NMR (CHLOROFORM- d) δ : 7.39 - 7.46 (m, 1H), 7.32 - 7.38 (m, 1H), 7.18 - 7.31 (m, 4H), 6.94 - 7.02 (m, 2H), 5.50 (q, J = 4.5 Hz, 1H), 4.66 - 4.86 (m, 4H), 4.07 (t, J = 5.4 Hz, 2H), 3.05 (d, J = 2.5 Hz, 3H), 2.35 (q, J = 7.6 Hz, 2H), 1.02 (t, J = 7.6 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{23}\text{ClN}_4\text{O}_3\text{S}$: 459.1 ($M+1$); found: 459.1.




Cpd 18

Cpd 18: ^1H NMR (CHLOROFORM- d) δ : 7.79 - 7.87 (m, 1H), 7.61 (td, J = 4.7, 1.3 Hz, 2H), 7.28 - 7.34 (m, 1H), 7.21 - 7.25 (m, 2H), 6.93 - 7.02 (m, 2H), 5.53 (br. s., 1H), 4.66 - 4.83 (m, 4H), 3.99 - 4.10 (m, 2H), 1.47 (s, 9H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{24}\text{ClF}_3\text{N}_4\text{O}_3$: 521.2 ($M+1$); found: 521.2.




Cpd 21

Cpd 21: ^1H NMR (CHLOROFORM- d) δ : 7.52 - 7.59 (m, 1H), 7.46 - 7.52 (m, 1H), 7.39 - 7.45 (m, 1H), 7.23 - 7.34 (m, 3H), 6.98 - 7.05 (m, 2H), 4.97 (d, J = 3.8 Hz, 2H), 4.85 (d, J = 5.6 Hz, 2H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{12}\text{ClF}_6\text{N}_3\text{O}_2$: 476.1 ($M+1$); found: 476.1.




Cpd 22

Cpd 22: ^1H NMR (CHLOROFORM- d) δ : 7.52 - 7.59 (m, 1H), 7.45 - 7.51 (m, 1H), 7.38 - 7.45 (m, 1H), 7.23 - 7.33 (m, 3H), 6.97 - 7.04 (m, 2H), 5.52 (br. s., 1H), 4.66 - 4.79 (m, 4H), 4.04 (t, J = 4.8 Hz, 2H), 1.47 (s, 9H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{24}\text{ClF}_3\text{N}_4\text{O}_4$: 537.2 ($M+1$); found: 537.2.



Cpd 29


Cpd 29: ^1H NMR (CHLOROFORM- d) δ : 7.36 - 7.44 (m, 2H), 7.20 - 7.28 (m, 2H), 7.00 - 7.09 (m, 3H), 6.92 (dd, J = 8.1, 5.3 Hz, 1H), 5.53 (br. s., 1H), 4.65 - 4.81 (m, 4H), 4.00 - 4.07 (m, 2H), 3.53 (d, J = 3.0 Hz, 3H), 1.47 (s, 9H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{27}\text{ClN}_4\text{O}_4$: 483.2 ($M+1$); found: 483.2.



Cpd 34

Cpd 34: ^1H NMR (CHLOROFORM- d) δ : 7.35 - 7.42 (m, 2H), 7.19 - 7.25 (m, 2H), 7.00 - 7.09 (m, 3H), 6.91 (d, J = 7.6 Hz, 1H), 4.74 (s, 2H), 4.67 (s, 2H), 3.53 (s, 3H), 2.93 (s, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{21}\text{ClN}_4\text{O}_2$: 397.1 ($M+1$); found: 397.1.


Example 5



A. Compound **5a** was prepared according to the procedures described for the synthesis of compound **1i** in Example 1, substituting 2-trifluoromethoxyphenylhydrazine for 2-ethylphenylhydrazine (**1f**) in Step C.

B. To a solution of compound **5a** (18 mg, 0.06 mmol, 1.0 eq) in toluene (1 mL) was added 2-chloro-6-iodoquinoline (29 mg, 0.12 mmol, 2 eq), Pd₂dba₃ (5.5 mg, 0.006 mmol, 0.1 eq), BINAP (11 mg, 0.018 mmol, 0.3 eq) and cesium carbonate (78 mg, 0.24 mmol, 4 eq), and the mixture was heated to 100°C for 3 days. Water was added, and the reaction mixture was extracted with DCM. The combined organic phases were dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (8 g), eluting with 15 to 30% EA/hexanes, gave compound **27** (9.0 mg, 28%). ¹H NMR (CHLOROFORM-*d*) δ: 7.95 (d, *J* = 8.8 Hz, 2H), 7.60 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.45 (dq, *J* = 15.1, 7.7 Hz, 1H), 7.46 (dq, *J* = 15.3, 7.7 Hz, 1H), 7.27 - 7.35 (m, 5H), 7.06 - 7.14 (m, 2H), 6.78 (d, *J* = 2.8 Hz, 1H), 4.70 (d, *J* = 4.3 Hz, 4H). ESI-MS (*m/z*): Calcd. for C₂₇H₁₇Cl₂F₃N₄O: 541.1 (*M*+1); found: 541.1.

Example 6



A. To a solution of KOH (3.27 g, 58.3 mmol, 7 eq) in H_2O (50 mL) at 0°C was added bromine (1.33 g, 8.32 mmol, 1 eq). After 15 min, compound **6a** (1.77 g, 8.32 mmol, 1 eq) in methanol (15 mL) was added dropwise over 2 min, and the reaction mixture was stirred for 1 hr at 0°C . The solution was made acidic with concentrated HCl, then extracted with EA. The combined organic phases were dried over MgSO_4 , filtered, and concentrated. Purification by column chromatography (160 g), eluting with 25 to 50% EA/hexanes + 0.1% HOAc, gave compound **6b** (1207 mg, 47%). ^1H NMR (CHLOROFORM- d) δ : 4.00 - 4.32 (m, 4H), 1.35 - 1.52 (m, 9H).

B. To a solution of compound **6b** (1189 mg, 4.07 mmol, 1 eq) in THF (20 mL) at rt under an Argon atmosphere was added N -methylmorpholine (0.47 mL, 4.27 mmol, 1.05 eq) followed by isobutylchloroformate (0.56 mL, 4.27 mmol, 1.05 eq). After 30 min, a solution of 2-methoxyphenylhydrazine-HCl (746 mg, 4.27 mmol, 1.05 eq) and N -methylmorpholine (0.47 mL, 4.27 mmol, 1.05 eq) in THF (20 mL) (pre-stirred for 20 min) was added and the solution was stirred for 60 min. Saturated aqueous NaHCO_3 and brine were sequentially added, and the aqueous mixture was extracted with EA. The combined organic phases were dried over MgSO_4 , filtered, and concentrated. Purification by column chromatography (40 g), eluting with 25 to 50% EA/hexanes, gave compound **6c** (1.29 g, 77%). ESI-MS (m/z): Calcd. for $\text{C}_{17}\text{H}_{22}\text{BrFN}_3\text{O}_4$: 412.1 ($M+1$); found: 412.1.

C. To a solution of compound **6c** (1.29 g, 3.14 mmol, 1 eq) in acetonitrile (20 mL) at rt was added carbon tetrachloride (0.91 mL, 9.41 mmol, 3 eq) followed by triphenylphosphine (2.47 g, 9.41 mmol, 3 eq). After 2 hrs, brine was added and the aqueous solution was extracted with EA. The combined organic phases were dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (40 g), eluting with 3 to 8% EA/hexanes, gave compound **6d** (604 mg, 45%). ¹H NMR (CHLOROFORM-d) δ: 8.17 (br. s., 1H), 7.34 (d, J = 7.8 Hz, 1H), 6.81 - 6.98 (m, 3H), 4.33 - 4.51 (m, 2H), 4.02 - 4.28 (m, 2H), 3.90 (s, 3H), 1.39 - 1.60 (m, 9H).


D. A solution of compound **6d** (604 mg, 1.4 mmol, 1 eq) and triethylamine (0.78 mL, 5.6 mmol, 4 eq) in toluene (12 mL) was heated to 110 °C for 3 hrs. Water was added to the reaction mixture, and the aqueous phase was extracted with DCM. The combined organics were dried over MgSO₄, filtered, and concentrated. Purification by column chromatography, eluting with 15 to 30% EA/hexanes, gave compound **6e** (411 mg, 74%). ¹H NMR (CHLOROFORM-d) δ: 7.43 - 7.50 (m, 1H), 7.33 (d, J = 7.8 Hz, 1H), 7.02 - 7.10 (m, 2H), 4.59 (s, 1H), 4.53 (s, 1H), 4.47 (s, 1H), 4.44 (s, 1H), 3.83 (s, 3H), 1.52 (d, J = 1.8 Hz, 9H). ESI-MS (m/z): Calcd. for C₁₇H₂₀BrN₃O₃: 394.1 (M+1); found: 394.0.

E. A solution of compound **6e** (411 mg, 1.04 mmol, 1 eq) in trifluoroacetic acid (1 mL) and DCM (5 mL) was stirred 3 hrs and concentrated. The residue was dissolved in DCM and washed with saturated aqueous NaHCO₃. The organic phase was dried over MgSO₄, filtered, and concentrated to give compound **6f** (290 mg, 95%). ¹H NMR (CHLOROFORM-d) δ: 7.45 (td, J = 7.9, 1.6 Hz, 1H), 7.34 (dd, J = 8.1, 1.8 Hz, 1H), 7.01 - 7.09 (m, 2H), 4.11 (s, 2H), 4.04 (s, 2H), 3.83 (s, 3H), 2.97 (br. s., 1H). ESI-MS (m/z): Calcd. for C₁₂H₁₂BrN₃O: 294.0 (M+1); found: 294.0.

F. To compound **6f** (290 mg, 0.99 mmol, 1 eq) in DCM (5 mL) at rt was added diisopropylethylamine (0.52 mL, 2.96 mmol, 3 eq) followed by isopropylsulfonyl chloride (0.22 mL, 1.97 mmol, 2 eq) for approximately 18 hr. Saturated aqueous NaHCO₃ was added, and the mixture was extracted with DCM. The combined organic phases were dried over MgSO₄, filtered, and concentrated. Purification by column chromatography (24 g), eluting with 20 to 40% EA/hexanes, gave compound **6g** (281 mg, 71%). ¹H NMR (CHLOROFORM-d) δ: 7.48 (td, J = 8.0, 1.8 Hz, 1H), 7.30 - 7.36 (m, 1H), 7.02 - 7.10 (m, 2H), 4.67 (s, 2H), 4.58 (d, J = 1.5 Hz, 2H), 3.84 (s, 3H), 3.37 (quin, J = 6.8 Hz, 1H), 1.44


(d, $J = 7.1$ Hz, 6H). ESI-MS (m/z): Calcd. for $C_{15}H_{18}BrN_3O_3S$: 400.0 ($M+1$); found: 400.0.

G. To a solution of compound **6g** (16 mg, 0.04 mmol, 1 eq) in dimethoxyethane (1 mL) and 2 M sodium carbonate (0.25 mL) in a vial was added 2-chloro-5-pyridineboronic acid (19 mg, 0.12 mmol, 3 eq), palladium acetate (1 mg, 0.004 mmol, 0.1 eq) and tri-*o*-tolylphosphine (1.2 mg, 0.004 mmol, 0.1 eq). The mixture was heated to 80 °C for 15 minutes. The mixture was cooled to rt, water was added, and the aqueous phase was extracted with DCM. The combined organic phases were dried over $MgSO_4$, filtered, and concentrated. Purification by column chromatography (8 g), eluting with 20 to 40 to 50% EA/hexanes, gave compound **92** (6.5 mg, 36%). ^1H NMR (CHLOROFORM- d) δ : 8.11 (d, $J = 2.3$ Hz, 1H), 7.31 - 7.39 (m, 2H), 7.27 (dd, $J = 8.3, 2.5$ Hz, 1H), 7.16 (d, $J = 8.3$ Hz, 1H), 6.96 - 7.04 (m, 1H), 6.86 (d, $J = 7.8$ Hz, 1H), 4.65 (d, $J = 14.9$ Hz, 4H), 3.50 (s, 3H), 3.32 (quin, $J = 6.8$ Hz, 1H), 1.38 (d, $J = 6.8$ Hz, 6H). ESI-MS (m/z): Calcd. for $C_{20}H_{21}ClN_4O_3S$: 433.1 ($M+1$); found: 433.1.




Cpd 93

Cpd **93**: ^1H NMR (CHLOROFORM- d) δ : 8.67 (d, $J = 2.3$ Hz, 1H), 8.04 (d, $J = 8.3$ Hz, 1H), 7.86 (d, $J = 2.0$ Hz, 1H), 7.65 - 7.77 (m, 2H), 7.51 - 7.59 (m, 1H), 7.46 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.37 - 7.44 (m, 1H), 7.06 (td, $J = 7.6, 1.1$ Hz, 1H), 6.91 (d, $J = 8.3$ Hz, 1H), 4.84 (s, 2H), 4.75 (s, 2H), 3.50 (s, 3H), 3.41 (quin, $J = 6.8$ Hz, 1H), 1.47 (d, $J = 7.1$ Hz, 6H). ESI-MS (m/z): Calcd. for $C_{24}H_{24}N_4O_3S$: 449.2 ($M+1$); found: 449.2.



Cpd 96


Cpd 96: ^1H NMR (CHLOROFORM- d) δ : 7.96 (d, $J = 2.5$ Hz, 1H), 7.34 - 7.44 (m, 2H), 7.22 (dd, $J = 8.6, 2.5$ Hz, 1H), 7.04 (t, $J = 7.7$ Hz, 1H), 6.94 (d, $J = 8.3$ Hz, 1H), 6.54 (d, $J = 8.6$ Hz, 1H), 5.24 (quin, $J = 6.2$ Hz, 1H), 4.71 (d, $J = 14.1$ Hz, 4H), 3.60 (s, 3H), 3.38 (quin, $J = 6.9$ Hz, 1H), 1.45 (d, $J = 6.8$ Hz, 6H), 1.31 (d, $J = 6.3$ Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{28}\text{N}_4\text{O}_4\text{S}$: 457.2 ($M+1$); found: 457.2.



Cpd 97

Cpd 97: ^1H NMR (CHLOROFORM- d) δ : 7.98 (d, $J = 2.5$ Hz, 1H), 7.33 - 7.45 (m, 2H), 7.21 - 7.27 (m, 1H), 7.04 (t, $J = 7.7$ Hz, 1H), 6.94 (d, $J = 8.1$ Hz, 1H), 6.62 (d, $J = 8.6$ Hz, 1H), 4.71 (d, $J = 12.4$ Hz, 4H), 3.90 (s, 3H), 3.60 (s, 3H), 3.38 (quin, $J = 6.9$ Hz, 1H), 1.45 (d, $J = 6.6$ Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{24}\text{N}_4\text{O}_4\text{S}$: 429.2 ($M+1$); found: 429.2.

Example 7




A. A suspension of 5-chloro-2-nitroanisole (**7a**) (2.67 g, 14.2 mmol) and 5% Pt/C (200 mg) in methanol (60 mL) was placed under a hydrogen gas atmosphere and stirred 3 days. The suspension was filtered through diatomaceous earth and concentrated. Purification by column chromatography (40 g), eluting with 15 to 30% EA/hexanes, gave compound **7b** (1.69 g, 76%). ^1H NMR (CHLOROFORM- d) δ : 6.70 - 6.83 (m, 2H), 6.58 - 6.64 (m, 1H), 3.84 (s, 3H), 3.76 (br. s., 2H). ESI-MS (m/z): Calcd. for $\text{C}_7\text{H}_8\text{ClNO}$: 158.0 ($M+1$); found: 158.0.


B. To a suspension of compound **7b** (1.69 g, 10.7 mmol, 1 eq) in concentrated HCl (25 mL) and acetic acid (10 mL) at 0 °C was added a solution of sodium nitrite (740 mg,

10.7 mmol, 1 eq) in water (10 mL) dropwise. The suspension was warmed to 60 °C for 1.5 hrs. The solution was cooled to 0 °C and a solution of SnCl₂ (5.32 g, 23.6 mmol, 2.2 eq) in concentrated HCl (25 mL) was added. After 30 minutes, a precipitate was collected by filtration to give compound **7c** (2.06 g, 93%). ¹H NMR (DMSO-d₆) δ: 9.94 (br. s., 2H), 7.72 (br. s., 1H), 7.07 (s, 1H), 7.00 (s, 2H), 3.85 (s, 3H).


Following the procedure described above for Example 7 and substituting the appropriate reagents, starting materials and purification methods known to those skilled in the art, the following intermediates to compounds of the present invention were prepared:




Cpd **7d**: ¹H NMR (free base, CHLOROFORM-d) δ: 6.89 (dd, J = 8.7, 5.7 Hz, 1H), 6.60 - 6.69 (m, 1H), 6.52 - 6.60 (m, 1H), 3.0 - 5.0 (brd. s, 3 H), 3.82 (s, 3H).




Cpd **7e**: ¹H NMR (DMSO-d₆) δ: 9.38 (br. s., 2H), 7.36 (br. s., 1H), 6.88 (d, J = 7.8 Hz, 1H), 6.80 - 6.84 (m, 1H), 6.69 - 6.75 (m, 1H), 3.81 (s, 3H), 2.26 (s, 3H).



Cpd **7f**: ¹H NMR (DMSO-d₆) δ: 9.96 (br. s., 2H), 7.82 (br. s., 1H), 7.16 (t, J = 9.3 Hz, 1H), 6.92 (dd, J = 13.1, 2.8 Hz, 1H), 6.79 (dt, J = 8.8, 1.4 Hz, 1H), 3.73 (s, 3H).




Cpd **7g**: ^1H NMR (free base, CHLOROFORM- d) δ : 6.78 (t, J = 8.0 Hz, 1H), 6.56 (d, J = 8.1 Hz, 1H), 6.40 (d, J = 7.8 Hz, 1H), 5.89 (s, 2H).




Cpd **7h**: ^1H NMR (DMSO- d_6) δ : 9.97 (br. s., 2H), 7.75 (br. s., 1H), 6.90 (d, J = 7.1 Hz, 1H), 6.85 (d, J = 8.1 Hz, 1H), 6.77 - 6.82 (m, 1H), 4.58 (t, J = 8.7 Hz, 2H), 3.19 (t, J = 8.8 Hz, 2H).

Using the intermediates prepared by Example 7 and the procedures described in Examples 1 and 2, the following compounds of the present invention were prepared:




Cpd **64**

Cpd **64**: ^1H NMR (CHLOROFORM- d) δ : 7.34 (d, J = 8.3 Hz, 1H), 7.22 - 7.29 (m, 2H), 6.99 - 7.07 (m, 3H), 6.90 (d, J = 2.0 Hz, 1H), 4.70 (d, J = 12.9 Hz, 4H), 3.53 (s, 3H), 3.38 (quin, J = 6.8 Hz, 1H), 1.45 (d, J = 7.1 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{21}\text{Cl}_2\text{N}_3\text{O}_3\text{S}$: 466.1 ($M+1$); found: 466.1.




Cpd 65

Cpd 65: ^1H NMR (CHLOROFORM- d) δ : 7.34 (d, $J = 8.3$ Hz, 1H), 7.24 - 7.26 (m, 2H), 7.00 - 7.07 (m, 3H), 6.90 (d, $J = 2.0$ Hz, 1H), 4.58 (s, 4H), 3.53 (s, 3H), 2.90 (s, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{20}\text{Cl}_2\text{N}_4\text{O}_3\text{S}$: 467.1 ($M+1$); found: 467.1.




Cpd 67

Cpd 67: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.26 (m, 2H), 7.14 - 7.19 (m, 1H), 6.94 - 7.02 (m, 2H), 6.74 - 6.81 (m, 2H), 4.75 (s, 2H), 4.68 (s, 2H), 3.83 (s, 3H), 3.40 (quin, $J = 6.8$ Hz, 1H), 1.96 (s, 3H), 1.46 (d, $J = 6.8$ Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{24}\text{ClN}_3\text{O}_3\text{S}$: 446.1 ($M+1$); found: 446.1.




Cpd 68

Cpd 68: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.25 (m, 2H), 7.14 - 7.20 (m, 1H), 6.94 - 7.00 (m, 2H), 6.75 - 6.80 (m, 2H), 4.74 (s, 2H), 4.67 (s, 2H), 3.83 (s, 3H), 3.05 - 3.17 (m, 1H), 2.17 - 2.28 (m, 2H), 1.96 (s, 3H), 1.91 (d, $J = 13.1$ Hz, 2H), 1.60 - 1.77 (m, 3H), 1.19 - 1.38 (m, 4H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{28}\text{ClN}_3\text{O}_3\text{S}$: 486.2 ($M+1$); found: 486.2.




Cpd 69

Cpd 69: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.26 (m, 2H), 7.14 - 7.19 (m, 1H), 6.94 - 7.01 (m, 2H), 6.74 - 6.80 (m, 2H), 4.66 (s, 2H), 4.58 (s, 2H), 3.83 (s, 3H), 2.91 (s, 6H), 1.96 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{23}\text{ClN}_4\text{O}_3\text{S}$: 447.1 ($M+1$); found: 447.1.



Cpd 70


Cpd 70: ^1H NMR (CHLOROFORM- d) δ : 7.36 (dd, $J = 8.6, 6.1$ Hz, 1H), 7.23 - 7.29 (m, 2H), 7.00 - 7.06 (m, 2H), 6.76 (td, $J = 8.2, 2.5$ Hz, 1H), 6.65 (dd, $J = 10.2, 2.7$ Hz, 1H), 4.56 - 4.72 (m, 4H), 3.55 (s, 3H), 2.96 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{19}\text{H}_{17}\text{ClFN}_3\text{O}_3\text{S}$: 422.1 ($M+1$); found: 422.1.



Cpd 71


Cpd 71: ^1H NMR (CHLOROFORM- d) δ : 7.36 (dd, $J = 8.8, 6.1$ Hz, 1H), 7.22 - 7.29 (m, 2H), 6.98 - 7.06 (m, 2H), 6.75 (td, $J = 8.2, 2.8$ Hz, 1H), 6.65 (dd, $J = 10.2, 2.7$ Hz,

1H), 4.61 - 4.80 (m, 4H), 3.55 (s, 3H), 3.38 (quin, J = 6.9 Hz, 1H), 1.45 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for C₂₁H₂₁ClFN₃O₃S: 450.1 (M+1); found: 450.1.




Cpd 72

Cpd 72: ¹H NMR (CHLOROFORM-d) δ: 7.36 (dd, J = 8.6, 6.1 Hz, 1H), 7.21 - 7.29 (m, 2H), 7.02 (d, J = 8.3 Hz, 2H), 6.75 (td, J = 8.2, 2.5 Hz, 1H), 6.65 (dd, J = 10.4, 2.5 Hz, 1H), 4.59 - 4.80 (m, 4H), 3.55 (s, 3H), 3.04 - 3.16 (m, 1H), 2.22 (d, J = 11.6 Hz, 2H), 1.91 (d, J = 12.9 Hz, 2H), 1.60 - 1.77 (m, 3H), 1.15 - 1.37 (m, 3H). ESI-MS (m/z): Calcd. for C₂₄H₂₅ClFN₃O₃S: 490.1 (M+1); found: 490.1.




Cpd 73

Cpd 73: ¹H NMR (CHLOROFORM-d) δ: 7.36 (dd, J = 8.6, 6.1 Hz, 1H), 7.22 - 7.28 (m, 2H), 7.03 (d, J = 8.6 Hz, 2H), 6.71 - 6.78 (m, 1H), 6.64 (dd, J = 10.1, 2.5 Hz, 1H), 4.50 - 4.69 (m, 4H), 3.75 (d, J = 12.1 Hz, 2H), 3.55 (s, 3H), 2.78 - 2.90 (m, 2H), 1.67 - 1.77 (m, 2H), 1.42 - 1.55 (m, 1H), 1.27 (qd, J = 12.3, 4.0 Hz, 2H), 0.97 (d, J = 6.3 Hz, 3H). ESI-MS (m/z): Calcd. for C₂₄H₂₆ClFN₄O₃S: 505.1 (M+1); found: 505.1.




Cpd 74

Cpd 74: ^1H NMR (CHLOROFORM- d) δ : 7.21 - 7.26 (m, 3H), 7.01 - 7.07 (m, 2H), 6.84 (d, J = 8.1 Hz, 1H), 6.72 (s, 1H), 4.55 - 4.71 (m, 4H), 3.52 (s, 3H), 2.95 (s, 3H), 2.40 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{20}\text{H}_{20}\text{ClN}_3\text{O}_3\text{S}$: 418.1 ($M+1$); found: 418.1.




Cpd 75

Cpd 75: ^1H NMR (CHLOROFORM- d) δ : 7.24 (dd, J = 8.2, 5.9 Hz, 3H), 7.03 (d, J = 8.3 Hz, 2H), 6.84 (d, J = 8.1 Hz, 1H), 6.72 (s, 1H), 4.61 - 4.80 (m, 4H), 3.52 (s, 3H), 3.38 (dt, J = 13.7, 6.9 Hz, 1H), 2.40 (s, 3H), 1.44 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{24}\text{ClN}_3\text{O}_3\text{S}$: 446.1 ($M+1$); found: 446.1.




Cpd 76

Cpd 76: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.26 (m, 3H), 7.01 - 7.07 (m, 2H), 6.84 (d, J = 8.1 Hz, 1H), 6.72 (s, 1H), 4.63 - 4.77 (m, 4H), 3.52 (s, 3H), 3.16 - 3.26 (m, 1H), 2.40 (s, 3H), 1.96 - 2.07 (m, 1H), 1.49 - 1.68 (m, 2H), 1.42 (d, J = 6.8 Hz, 3H), 1.31 - 1.40 (m, 1H), 0.93 - 1.00 (m, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{28}\text{ClN}_3\text{O}_3\text{S}$: 474.2 ($M+1$); found: 474.2.




Cpd 77

Cpd 77: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.26 (m, 3H), 7.00 - 7.07 (m, 2H), 6.83 (d, J = 7.8 Hz, 1H), 6.71 (s, 1H), 4.51 - 4.68 (m, 4H), 3.75 (d, J = 12.4 Hz, 2H), 3.52 (s, 3H), 2.78 - 2.90 (m, 2H), 2.40 (s, 3H), 1.68 - 1.76 (m, 2H), 1.49 (ddd, J = 14.3, 7.2, 3.5 Hz, 1H), 1.20 - 1.34 (m, 2H), 0.97 (d, J = 6.3 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{29}\text{ClN}_4\text{O}_3\text{S}$: 501.2 ($M+1$); found: 501.2.




Cpd 78

Cpd 78: ^1H NMR (CHLOROFORM- d) δ : 7.34 (t, J = 8.7 Hz, 1H), 7.26 - 7.31 (m, 2H), 7.02 - 7.09 (m, 2H), 6.77 (ddd, J = 8.8, 2.7, 1.1 Hz, 1H), 6.66 (dd, J = 11.6, 2.8 Hz, 1H), 4.63 (d, J = 4.8 Hz, 4H), 3.84 (s, 3H), 2.96 (s, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{19}\text{H}_{17}\text{ClFN}_3\text{O}_3\text{S}$: 422.1 ($M+1$); found: 422.1.



Cpd 79


Cpd 79: ^1H NMR (CHLOROFORM- d) δ : 7.34 (t, J = 8.6 Hz, 1H), 7.24 - 7.30 (m, 2H), 7.02 - 7.09 (m, 2H), 6.76 (ddd, J = 8.8, 2.7, 1.1 Hz, 1H), 6.65 (dd, J = 11.6, 2.5 Hz, 1H), 4.70 (dd, J = 10.4, 1.5 Hz, 4H), 3.84 (s, 3H), 3.38 (quin, J = 6.8 Hz, 1H), 1.45 (d, J = 7.1 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{21}\text{ClFN}_3\text{O}_3\text{S}$: 450.1 ($M+1$); found: 450.1.



72


Cpd 80

Cpd 80: ^1H NMR (CHLOROFORM- d) δ : 7.34 (t, J = 8.7 Hz, 1H), 7.24 - 7.30 (m, 2H), 7.02 - 7.09 (m, 2H), 6.76 (ddd, J = 8.8, 2.7, 1.1 Hz, 1H), 6.65 (dd, J = 11.6, 2.8 Hz, 1H), 4.59 (dd, J = 9.9, 1.5 Hz, 4H), 3.83 (s, 3H), 3.75 (d, J = 12.1 Hz, 2H), 2.78 - 2.91 (m, 2H), 1.67 - 1.78 (m, 2H), 1.42 - 1.55 (m, 1H), 1.20 - 1.34 (m, 2H), 0.97 (d, J = 6.6 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{26}\text{ClFN}_4\text{O}_3\text{S}$: 505.1 ($M+1$); found: 505.1.




Cpd 81

Cpd 81: ^1H NMR (CHLOROFORM- d) δ : 7.27 - 7.33 (m, 2H), 7.07 - 7.15 (m, 2H), 6.82 - 6.90 (m, 2H), 6.78 - 6.82 (m, 1H), 5.83 (s, 2H), 4.70 (d, J = 3.5 Hz, 4H), 3.37 (quin, J = 6.8 Hz, 1H), 1.44 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{21}\text{H}_{20}\text{ClN}_3\text{O}_4\text{S}$: 446.1 ($M+1$); found: 446.1.




Cpd 82

Cpd 82: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.25 (m, 2H), 7.01 - 7.07 (m, 2H), 6.75 - 6.82 (m, 2H), 6.73 (dd, J = 6.3, 3.3 Hz, 1H), 5.75 (s, 2H), 4.57 (d, J = 1.8 Hz, 4H), 2.87 (d, J = 6.6 Hz, 2H), 2.28 (dt, J = 13.3, 6.8 Hz, 1H), 1.07 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{22}\text{ClN}_3\text{O}_4\text{S}$: 460.1 ($M+1$); found: 460.1.




Cpd 83

Cpd 83: ^1H NMR (CHLOROFORM- d) δ : 7.27 - 7.32 (m, 2H), 7.08 - 7.15 (m, 2H), 6.83 - 6.89 (m, 2H), 6.77 - 6.83 (m, 1H), 5.82 (s, 2H), 4.59 (d, J = 2.8 Hz, 4H), 3.74 (d, J = 12.1 Hz, 2H), 2.84 (td, J = 12.2, 2.4 Hz, 2H), 1.65 - 1.77 (m, 2H), 1.42 - 1.57 (m, 1H), 1.19 - 1.33 (m, 2H), 0.96 (d, J = 6.6 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{24}\text{H}_{25}\text{ClN}_4\text{O}_4\text{S}$: 501.1 ($M+1$); found: 501.1.




Cpd 84

Cpd 84: ^1H NMR (CHLOROFORM- d) δ : 7.19 - 7.33 (m, 3H), 7.02 - 7.16 (m, 3H), 6.84 - 6.94 (m, 1H), 4.70 (d, J = 9.1 Hz, 4H), 4.44 (t, J = 8.7 Hz, 2H), 3.31 - 3.46 (m, 1H), 3.23 (t, J = 8.6 Hz, 2H), 1.44 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{22}\text{H}_{22}\text{ClN}_3\text{O}_3\text{S}$: 444.1 ($M+1$); found: 444.1.



Cpd 85

Cpd 85: ^1H NMR (CHLOROFORM- d) δ : 7.20 - 7.29 (m, 3H), 7.03 - 7.12 (m, 3H), 6.84 - 6.90 (m, 1H), 4.64 (d, J = 4.8 Hz, 4H), 4.44 (t, J = 8.7 Hz, 2H), 3.23 (t, J = 8.7 Hz, 2H), 2.94 (d, J = 6.3 Hz, 2H), 2.35 (dt, J = 13.3, 6.6 Hz, 1H), 1.14 (d, J = 6.8 Hz, 6H). ESI-MS (m/z): Calcd. for $\text{C}_{23}\text{H}_{24}\text{ClN}_3\text{O}_3\text{S}$: 458.1 ($M+1$); found: 458.1.




Cpd 86

Cpd 86: ^1H NMR (CHLOROFORM- d) δ : 7.19 - 7.29 (m, 3H), 7.02 - 7.13 (m, 3H), 6.83 - 6.91 (m, 1H), 4.59 (d, J = 6.1 Hz, 4H), 4.44 (t, J = 8.7 Hz, 2H), 3.74 (d, J = 12.4 Hz, 2H), 3.23 (t, J = 8.7 Hz, 2H), 2.83 (td, J = 12.2, 2.4 Hz, 2H), 1.67 - 1.75 (m, 2H), 1.41 - 1.57 (m, 1H), 1.19 - 1.33 (m, 2H), 0.96 (d, J = 6.3 Hz, 3H). ESI-MS (m/z): Calcd. for $\text{C}_{25}\text{H}_{27}\text{ClN}_4\text{O}_3\text{S}$: 499.2 ($M+1$); found: 499.2.

The compounds of Table 1, exemplified hereinbelow, were prepared according to the schemes and specific examples described herein.


Table 1. Compounds of Formula (I)

<div style="text-align: center;"> <p>Formula (I)</p> </div>					
Cpd No.	R^1	R^2	R^3	Q	G
1	ethyl	H	H	4-chlorophenyl	<i>t</i> -butoxy carbonyl
2	ethyl	H	H	4-chlorophenyl	methanesulfonyl
3	ethyl	H	H	4-chlorophenyl	trifluoromethyl sulfonyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
4	ethyl	H	H	4-chlorophenyl	pyridin-3-yl carbonyl
5	ethyl	H	H	4-chlorophenyl	pyridin-4-yl carbonyl
6	ethyl	H	H	4-chlorophenyl	trifluoromethyl carbonyl
7	ethyl	H	H	4-chlorophenyl	pyridin-2-yl carbonyl
8	ethyl	H	H	4-chlorophenyl	furan-2-yl carbonyl
9	ethyl	H	H	4-chlorophenyl	5-methyl isoxazol-3-yl carbonyl
10	ethyl	H	H	4-chlorophenyl	methoxymethyl carbonyl
11	ethyl	H	H	4-chlorophenyl	<i>t</i> -butyloxy carbonylamino methylcarbonyl
12	ethyl	H	H	4-chlorophenyl	methanesulfonylamino methyl carbonyl
13	ethyl	H	H	4-chlorophenyl	trifluoromethyl sulfonylamino methylcarbonyl
14	trifluoromethyl	H	H	4-chlorophenyl	<i>t</i> -butyloxy carbonyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
15	trifluoromethyl	H	H	4-chlorophenyl	methanesulfonyl
16	trifluoromethyl	H	H	4-chlorophenyl	isopropyl sulfonyl
17	trifluoromethyl	H	H	4-chlorophenyl	4-methylphenyl sulfonyl
18	trifluoromethyl	H	H	4-chlorophenyl	<i>t</i> -butyloxy carbonylamino methylcarbonyl
19	trifluoro methoxy	H	H	4-chlorophenyl	<i>t</i> -butoxycarbonyl
20	trifluoro methoxy	H	H	4-chlorophenyl	methanesulfonyl
21	trifluoro methoxy	H	H	4-chlorophenyl	trifluoromethyl carbonyl
22	trifluoro methoxy	H	H	4-chlorophenyl	<i>t</i> -butyloxy carbonylamino methylcarbonyl
23	trifluoro methoxy	H	H	4-chlorophenyl	3-cyanophenyl
24	trifluoro methoxy	H	H	4-chlorophenyl	4-methoxy carbonylphenyl
25	trifluoro methoxy	H	H	4-chlorophenyl	4-cyanophenyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
26	trifluoro methoxy	H	H	4-chlorophenyl	4-(2-(morpholin-4- yl)ethyl aminosulfonyl) phenyl
27	trifluoro methoxy	H	H	4-chlorophenyl	2-chloro quinolin-6-yl
28	methoxy	H	H	4-chlorophenyl	<i>t</i> -butoxycarbonyl
29	methoxy	H	H	4-chlorophenyl	<i>t</i> -butoxycarbonyl aminomethyl carbonyl
30	methoxy	H	H	4-chlorophenyl	methanesulfonyl
31	methoxy	H	H	4-chlorophenyl	isopropyl sulfonyl
32	methoxy	H	H	4-chlorophenyl	cyclopropyl sulfonyl
33	methoxy	H	H	4-chlorophenyl	dimethylamino sulfonyl
34	methoxy	H	H	4-chlorophenyl	dimethylamino carbonyl
35	trifluoro methoxy	H	H	4-chlorophenyl	isopropyl sulfonyl
36	methoxy	H	H	4-chlorophenyl	2,2,2-trifluoroethyl sulfonyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
37	methoxy	H	H	4-chlorophenyl	cyclohexyl sulfonyl
38	methoxy	H	H	4-chlorophenyl	2-(1 <i>H</i> -1,3-dioxo-isoindol-2-yl) ethylsulfonyl
39	methoxy	H	H	4-chlorophenyl	phenylmethyl sulfonyl
40	methoxy	H	H	4-chlorophenyl	cyclopentyl sulfonyl
41	methoxy	H	H	4-chlorophenyl	1(R,S)-methylbutyl) sulfonyl
42	methoxy	H	H	4-chlorophenyl	2-(2,5-dioxo-pyrrolidin-1-yl) ethylsulfonyl
43	methoxy	H	H	4-chlorophenyl	2-(phenyloxy)ethyl sulfonyl
44	methoxy	H	H	4-chlorophenyl	2-(methane sulfonylamino) ethylsulfonyl
45	methoxy	H	H	4-chlorophenyl	2-(isopropyl sulfonylamino) ethylsulfonyl
46	methoxy	H	H	4-chlorophenyl	2-(dimethylamino sulfonylamino) ethylsulfonyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
47	trifluoro methoxy	H	H	4-chlorophenyl	2-(methoxy carbonyl)ethyl sulfonyl
48	trifluoro methoxy	H	H	4-chlorophenyl	1,1-dioxido tetrahydro thiophen-3(R,S)- yl)sulfonyl
49	ethoxy	H	H	4-chlorophenyl	methanesulfonyl
50	ethoxy	H	H	4-chlorophenyl	isopropyl sulfonyl
51	ethoxy	H	H	4-chlorophenyl	cyclopentyl sulfonyl
52	ethoxy	H	H	4-chlorophenyl	cyclohexyl sulfonyl
53	ethoxy	H	H	4-chlorophenyl	phenylmethyl sulfonyl
54	ethoxy	H	H	4-chlorophenyl	dimethylamino sulfonyl
55	methoxy	H	H	4-chlorophenyl	azepan-1-yl sulfonyl
56	methoxy	H	H	4-chlorophenyl	diethylamino sulfonyl
57	methoxy	H	H	4-chlorophenyl	4-methyl piperidin-1-yl sulfonyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
58	methoxy	H	H	4-chlorophenyl	4-(methoxy carbonyl) piperidin-1-yl sulfonyl
59	methoxy	H	H	4-chlorophenyl	morpholin-4-yl sulfonyl
60	methoxy	H	H	4-chlorophenyl	pyrrolidin-1-yl sulfonyl
61	methoxy	H	H	4-chlorophenyl	(2S,6S)-2,6-dimethyl morpholin-4-yl sulfonyl
62	methoxy	H	H	4-chlorophenyl	piperidin-1-yl sulfonyl
63	isopropoxy	H	H	4-chlorophenyl	dimethylamino sulfonyl
64	methoxy	H	chloro	4-chlorophenyl	isopropyl sulfonyl
65	methoxy	H	chloro	4-chlorophenyl	dimethylamino sulfonyl
66	methyl	H	methoxy	4-chlorophenyl	methanesulfonyl
67	methyl	H	methoxy	4-chlorophenyl	isopropyl sulfonyl
68	methyl	H	methoxy	4-chlorophenyl	cyclohexyl sulfonyl




Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
69	methyl	H	methoxy	4-chlorophenyl	dimethylamino sulfonyl
70	methoxy	H	fluoro	4-chlorophenyl	methanesulfonyl
71	methoxy	H	fluoro	4-chlorophenyl	isopropyl sulfonyl
72	methoxy	H	fluoro	4-chlorophenyl	cyclohexyl sulfonyl
73	methoxy	H	fluoro	4-chlorophenyl	4-methyl piperidin-1-yl sulfonyl
74	methoxy	H	methyl	4-chlorophenyl	methanesulfonyl
75	methoxy	H	methyl	4-chlorophenyl	isopropyl sulfonyl
76	methoxy	H	methyl	4-chlorophenyl	1(<i>R,S</i>)-methylbutyl sulfonyl
77	methoxy	H	methyl	4-chlorophenyl	4-methyl piperidin-1-yl sulfonyl
78	fluoro	H	methoxy	4-chlorophenyl	methanesulfonyl
79	fluoro	H	methoxy	4-chlorophenyl	isopropyl sulfonyl
80	fluoro	H	methoxy	4-chlorophenyl	4-methylpiperidin-1-ylsulfonyl



Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
81	R ² taken with R ¹ and the phenyl ring to which R ¹ and R ² are both attached to form 1,3-benzodioxol-4-yl		H	4-chlorophenyl	isopropyl sulfonyl
82	R ² taken with R ¹ and the phenyl ring to which R ¹ and R ² are both attached to form 1,3-benzodioxol-4-yl		H	4-chlorophenyl	(2-methyl)propyl sulfonyl
83	R ² taken with R ¹ and the phenyl ring to which R ¹ and R ² are both attached to form 1,3-benzodioxol-4-yl		H	4-chlorophenyl	4-methyl piperidin-1-yl sulfonyl
84	R ² taken with R ¹ and the phenyl ring to which R ¹ and R ² are both attached to form 2,3-dihydrobenzofuran-7-yl		H	4-chlorophenyl	isopropyl sulfonyl
85	R ² taken with R ¹ and the phenyl ring to which R ¹ and R ² are both attached to form 2,3-dihydrobenzofuran-7-yl		H	4-chlorophenyl	(2-methyl) propyl sulfonyl
86	R ² taken with R ¹ and the phenyl ring to which R ¹ and R ² are both attached to form 2,3-dihydrobenzofuran-7-yl		H	4-chlorophenyl	4-methyl piperidin-1-yl sulfonyl



Formula (I)

Cpd No.	R ¹	R ²	R ³	Q	G
87	methoxy	H	H	4-chlorophenyl	(3,5-dimethyl isoxazol-4-yl) sulfonyl
88	dimethylamino	H	H	4-chlorophenyl	isopropyl sulfonyl
89	dimethylamino	H	H	4-chlorophenyl	ethylsulfonyl
90	2,2,2- trifluoroethoxy	H	H	4-chlorophenyl	dimethylaminosulfonyl
91	t-butoxy	H	H	4-chlorophenyl	dimethylamino sulfonyl
92	methoxy	H	H	6-chloro pyridin-3-yl	isopropyl sulfonyl
93	methoxy	H	H	quinolin-3-yl	isopropyl sulfonyl
94	methoxy	H	H	6-methyl pyridin-3-yl	isopropyl sulfonyl
95	2,2- difluoroethoxy	H	H	4-chlorophenyl	dimethylamino sulfonyl
96	methoxy	H	H	6-(isopropyl oxy) pyridin-3-yl	isopropyl sulfonyl
97	methoxy	H	H	6-methoxy pyridin-3-yl	isopropyl sulfonyl

Biological Examples

In Vitro Assays

Example 1

Functional Assay: Antagonism of N-Type Calcium Channel

A stable cell line (HEK parent) co-expressing the α_{1B} (Cav2.2), β_3 and $\alpha_2\delta$ subunits of the N-type calcium channel subunits was used. These cells were routinely grown as monolayers in low glucose-containing Dulbecco's Modified Eagle Medium supplemented with 10% FBS, 2 mM L-glutamine, 100 IU/mL penicillin, 100 μ g/mL streptomycin, 400 μ g/mL G418 and 200 μ g/mL Zeocin (split ratio = 1:5). Cells were maintained in 5% CO₂ at 37 °C. Compounds of Formula (I) were prepared as 10 mM stocks in DMSO from neat compound, if available. Otherwise, the 5 or 10 mM DMSO stock solutions provided in-house were used.

Calcium mobilization responses to KCl depolarization were evaluated by measuring the intensity of calcium-mediated fluorescent signal in the presence of BD Calcium Assay Dye (BD Biosciences, Franklin Lakes, NJ, U.S.A.), utilizing a Functional Drug Screening System (FDSS) by Hamamatsu Corporation (Bridgewater, NJ U.S.A.).

Twenty-four hr prior to assay, cells were seeded in clear-base poly-D-lysine-coated 384-well plates (BD Biosciences) at a density of 5,000 cells per well in culture medium and grown overnight in 5% CO₂ at 37 °C. On the day of assay, growth media were removed, and cells were loaded with BD calcium assay dye (BD Biosciences) for 35 min at 37°C under 5% CO₂ and then for 25 min at room temp. Utilizing the FDSS, cells were exposed to representative compounds of Formula (I) at varying concentrations, and intracellular calcium was measured for 5 min prior to the addition of 50 mM KCl for an additional 3 min of measurement.

Calculations and formulas

IC₅₀ values for representative compounds of Formula (I) were determined from six-point concentration-response experiments and represent the concentration of said compound required to inhibit 50% of the maximal response. Maximal fluorescence

intensity (FI) achieved upon addition of 50 mM KCl was exported from the FDSS software and further analyzed using GraphPad Prism 3.02 (Graph Pad Software Inc., San Diego, CA, U.S.A.). Data were normalized to the maximum average counts from quadruplicate wells for each condition in the presence of 50 mM KCl and to the minimum average counts in the presence of buffer. Theoretical curves were generated using nonlinear regression curve-fitting analysis of either sigmoidal concentration-response or sigmoidal concentration-response (variable slope), and the IC₅₀ values with the best-fit curve determined by GraphPad Prism were reported. Resultant data are shown in Table 2.

Table 2.

Compound No	FDSS IC₅₀ (μM)	%Inhibition (%) at 0.33 μM	%Inhibition (%) at 1 μM
1	0.180		
2	0.085		
3	0.150		
4	0.200		
5	0.200		
6	0.180		
7			58
8	0.600		
9			63
10	0.280		
11	0.100		
12	0.520		
13			59
14			60
15	0.280		
16	0.160		
17	0.300		
18	0.480		
19	0.300		
20	0.040		
21	0.300		
22	0.240		
23			52
24			64
25			57
26			57

Compound No	FDSS IC ₅₀ (μ M)	%Inhibition (%) at 0.33 μ M	%Inhibition (%) at 1 μ M
27	0.093		
28	0.073		
29	0.180		
30	0.021		
31	0.010	94	
32	0.021		
33	0.009		
34	0.170		
35	0.027		
36	0.015		
37	0.016		
38	0.053		
39	0.027		
40	0.012		
41	0.019	90	
42	0.048		
43	0.100		
44	0.009		
45	0.013		
46	0.005		
47	0.031		
48	0.077		
49	0.002		
50	0.003		
51	0.005		
52	0.002		
53	0.007		
54	0.002		
55	0.007		
56	0.003		
57	0.008	95	
58	0.005		
59	0.002		
60	0.002		
61	0.005		
62	0.004		
63	0.003		
64	0.033	86	
65	0.150	74	

Compound No	FDSS IC ₅₀ (μ M)	%Inhibition (%) at 0.33 μ M	%Inhibition (%) at 1 μ M
66		35	
67	0.190	61	
68	0.130	71	
69	0.130	69	
70	0.075	74	
71	0.021	91	
72	0.095	82	
73	0.024	93	
74	0.047	82	
75	0.032	93	
76	0.021	90	
77	0.056	82	
78	0.200	62	
79	0.180	61	
80	0.140	62	
81	0.020	89	
82	0.028	87	
83	0.020	92	
84	0.010	94	
85	0.014	94	
86	0.007	96	
87	0.017	89	
88	0.005	96	
89	0.028	90	
90	0.085	80	
91	0.063	87	
92	0.016	86	
93	0.100		
94		46	
95	0.050	79	
96	0.130	79	
97	0.043	67	

Example 2

Automated Electrophysiology Assay

Cells were grown in T175 flasks to 50%–90% confluence. At the time of use, cells were enzymatically treated with Detachin (Genlantis, San Diego, CA USA), centrifuged, rinsed, and resuspended in 293 SFM II media (Life Technologies, Grand Island, NY U.S.A.) supplemented with 25 mM HEPES (Sigma-Aldrich, St. Louis, MO U.S.A.) to a concentration of $2\text{--}3 \times 10^6$ cells/mL. Cells were added to the automated cell preparation station on the QPatch-HT (Sophion Biosciences, North Brunswick, NJ U.S.A.), and following a 10- to 30-min recovery period with gentle stirring, the assay protocol was initiated. During the automated cell preparation, cells were collected, centrifuged and resuspended in an extracellular (EC) solution containing 132 mM NaCl, 1.8 mM CaCl_2 , 5.4 mM KCl, 0.8 mM MgCl_2 , 10 mM glucose, and 10 mM HEPES (pH = 7.4), adjusted with sucrose to approximately 315 mOsm. The QPlate was primed with an intracellular solution containing 135 mM CsCl, 10 mM EGTA, 4 MgATP, 0.3 NaGTP, and 20 mM HEPES (pH = 7.2), adjusted to approximately 290 mOsm with deionized water and the EC solution. Cells were added to the prepared QPlate wells by robotic pipettes of the QPatch-HT.

For cells determined to be in stable whole-cell patch clamp, the EC solution was replaced with a barium (Ba)/triethylammonium (TEA) solution containing 140 mM TEA-Cl, 10 mM BaCl_2 , 0.8 mM MgCl_2 , 10 mM glucose and 10 mM HEPES (pH = 7.4). High (40 mM) BaCl_2 concentrations were made with adjustments to TEA-Cl (90 mM) to maintain the osmolarity. From a resting potential of -80 mV, a train of depolarizing pulses (15 pulses at 5 Hz, +20 mV) was delivered to the cell once every 30 sec for eight trains (4 min total), and the resulting currents were measured during a control period (no compound). This protocol was repeated for each subsequent addition of control buffer with or without compound (three periods total, each with four trains). The current generated in the 1st and 15th pulses of the last train of each period in the presence of each drug concentration was normalized to the current generated during the control period at the respective pulses (representing low- and high-frequency stimulation, respectively). Data

from both the second and third drug application periods were analyzed for each cell. A final addition of Ba/TEA solution containing 60–100 μM CdCl_2 was made to block all N-type current and to “zero” the currents for each cell. All buffer/ compound additions were made using a “spitting” feature of the QPatch-HT, which added three repetitions of 5 μL solution at the beginning of each recording period.

To examine closed-state inactivation, cells were subjected to a channel-activating 50-msec depolarizing step pulse from -80 to +10 mV, followed by a 5-sec nonactivating step to voltages ranging from -130 to -60 mV in 10 mV increments and then a 50-ms step from -80 to +10 mV to assess the remaining current. Currents from the activating voltage pulse were normalized to the peak value of the test pulse following the -130 mV step and fit to a Boltzman equation to obtain the $V_{1/2}$. Roscovitine (Sigma-Aldrich) was prepared as a 100 mM stock in dimethyl sulfoxide and diluted to the indicated working concentrations. Tetrandrine (Sigma-Aldrich) was prepared as a 4 mM stock in acidic water (pH = 2.0) and then diluted to working concentrations in the external solution. ω -Conotoxin MVIIA (Sigma-Aldrich) was prepared as a 0.3 mg/mL stock solution in water, with 0.1% bovine serum albumin V (Life Technologies). Compounds of Formula (I) were diluted first into dimethyl sulfoxide and then into 10% pluronic F-127 in water (Life Technologies), sonicated for 1 min and diluted into EC buffer. Vehicle controls were run in parallel in all experiments.

Unless otherwise indicated, statistics for comparing among electrophysiological results utilized a one-way analysis of variance with Fisher’s least squares determination test for pair-wise comparison. Resultant data are shown in Tables 3 and 4, below.

Table 3.

QPatch at Low Frequency					
Cpd No.	% Inhibition at Various Concentrations (μM)				
	0.03	0.05	0.1	0.25	1
2					-9
5				0	11
6					16
11				14	37
20					5

QPatch at Low Frequency					
Cpd No.	% Inhibition at Various Concentrations (μ M)				
	0.03	0.05	0.1	0.25	1
22					13
27					-1
28					25
30					28
31			11		
32	-6	10		57	74
33					74
35				22	
36				42	
37				61	
40				57	
41			7		80
44			-2		
45			26		
46			-8		
47			8		
49			18		
50			42		
51			37		
52			40		
54			36		
55			25		
56			28		
57					69
58			53		
59			28		
60			29		
61			53		
62			20		
64			12		
65			-8		
71			-30		
73			-34		
76					-3
81			-19		
82			4		
83			12		
84			-3		

QPatch at Low Frequency					
Cpd No.	% Inhibition at Various Concentrations (μ M)				
	0.03	0.05	0.1	0.25	1
85			13		
86			20		
87			31		
88			26		
89			10		
92			14		
95			-4		
97					40

Table 4.

QPatch at High Frequency					
Cpd No.	% Inhibition at Various Concentrations (μ M)				
	0.03	0.05	0.1	0.25	1
2					22
5				12	37
6					23
11				31	68
20					30
22					28
28					52
30					49
31			47		
32	-5	23		64	83
33					78
35				42	

QPatch at High Frequency					
Cpd No.	% Inhibition at Various Concentrations (μ M)				
	0.03	0.05	0.1	0.25	1
36				58	
37				73	
40				69	
41			40		91
44			-5		
45			26		
46			-6		
47			36		
49			22		
50			50		
51			47		
52			50		
54			46		
55			54		
56			36		
57					82
58			52		
59			51		
60			44		
61			50		


QPatch at High Frequency					
Cpd No.	% Inhibition at Various Concentrations (μ M)				
	0.03	0.05	0.1	0.25	1
62			45		
64			24		
65			1		
71			1		
73			-28		
76					30
82			15		
83			27		
84			7		
85			37		
86			33		
87			49		
88			47		
89			34		
92			10		
95			23		
97					60

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the

practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

We claim:

1. A compound of Formula (I)



Formula (I)


wherein

R^1 is selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, 2,2-difluoroethoxy, di(C_{1-4} alkyl)amino, trifluoromethoxy, fluoro, and trifluoromethyl;

R^2 is hydrogen; or, R^2 may be taken with R^1 and the phenyl ring to which R^1 and R^2 are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R^3 is hydrogen, methyl, methoxy, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;



wherein


R^4 is selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, cyano, fluoro, and chloro;

R^5 is independently selected from the group consisting of hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, cyano, chloro, and di(C_{1-4} alkyl)amino;

G is selected from the group consisting of 4-methylphenyl, 3-cyanophenyl, 4-cyanophenyl, 4-methoxycarbonylphenyl, 2-chloro-quinolin-6-yl, morpholin-4-ylsulfonyl, morpholin-4-yl(C₁₋₄alkyl)aminosulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C₆₋₁₀aryl(C₁₋₄)alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, , C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, , trifluoromethylsulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isoindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethylcarbonyl, trifluoromethylsulfonylaminomethylcarbonyl, trifluoromethoxycarbonyl, C₁₋₄alkylsulfonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, pyridinylcarbonyl, furanylcabonyl, and 5-methylisoxazol-3-ylcarbonyl;

and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof.

2. The compound of claim 1 wherein R¹ is selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, 2,2-difluoroethoxy, di(C₁₋₄alkyl)amino, and trifluoromethoxy.
3. The compound of claim 2 wherein R¹ is selected from the group consisting of ethyl, methoxy, ethoxy, isopropoxy, difluoroethoxy, dimethylamino, and trifluoromethoxy.
4. The compound of claim 1 wherein R³ is hydrogen, methyl, chloro, or fluoro.
5. The compound of claim 1 wherein Q is selected from the group consisting of Q1, Q2, and Q3;




wherein

R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkyl, C_{1-4} alkoxy, and chloro.

6. The compound of claim 5 wherein Q is selected from the group consisting of Q1, Q2, and Q3;



wherein

R^4 is chloro;


R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro.

7. The compound of claim 1 wherein G is selected from the group consisting of 4-methylphenyl, 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4})alkylsulfonyl, C_{6-10} aryloxy(C_{1-4})alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, trifluoromethylsulfonyl, C_{1-4} alkoxycarbonyl(C_{1-4})alkyl-sulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, di(C_{1-4} alkyl)aminosulfonylamino(C_{1-4})alkyl-sulfonyl, C_{1-4} alkylsulfonylamino(C_{1-4})alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C_{1-4})alkyl-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C_{1-4})alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C_{1-6} alkylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R^6)-piperidin-1-ylsulfonyl wherein R^6 is C_{1-4} alkoxycarbonyl or C_{1-4}

- alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-carbonyl, C₁₋₄alkylsulfonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, pyridinylcarbonyl, and furanylcarbonyl.
8. The compound of claim 7 wherein G is selected from the group consisting of methylphenyl, 2-chloro-quinolin-6-yl, C₆₋₁₀aryl(C₁₋₄)alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, 4-C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, trifluoromethylsulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, and pyridinylcarbonyl.
9. The compound of claim 8 wherein G is selected from the group consisting of 2-chloro-quinolin-6-yl, C₆₋₁₀aryl(C₁₋₄)alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl.

alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, and C₁₋₄alkoxycarbonyl.

10. The compound of Formula (I) as in claim 1



Formula (I)


wherein

R¹ is selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, 2,2-difluoroethoxy, di(C₁₋₄alkyl)amino, trifluoromethoxy, fluoro, and trifluoromethyl;

R² is hydrogen; or, R² may be taken with R¹ and the phenyl ring to which R¹ and R² are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R³ is hydrogen, methyl, methoxy, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;




wherein

R⁴ is selected from the group consisting of chloro;

R⁵ is independently selected from the group consisting of C₁₋₄alkoxy and chloro;

G is selected from the group consisting of 4-methylphenyl, 2-chloro-quinolin-6-yl, C₆₋₁₀aryl(C₁₋₄)alkylsulfonyl, C₆₋₁₀aryloxy(C₁₋₄)alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, trifluoromethylsulfonyl, C₁₋₄alkoxycarbonyl(C₁₋₄)alkyl-sulfonyl, C₃₋₇cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄)alkyl-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄)alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄)alkyl-sulfonyl, (2H)-1,3-dione-1H-isoindol-1-yl-(C₁₋₄)alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄)alkyl-carbonyl, C₁₋₄alkoxycarbonylaminomethyl-carbonyl, C₁₋₄alkylsulfonylaminomethyl-carbonyl, C₁₋₄alkoxycarbonyl, di(C₁₋₄alkyl)aminocarbonyl, C₁₋₄alkoxycarbonyl, pyridinylcarbonyl, and furanylcarbonyl; and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof.

11. The compound of Formula (I) as in claim 1



Formula (I)


wherein

R¹ is selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkoxy, 2,2-difluoroethoxy, di(C₁₋₄alkyl)amino, trifluoromethoxy, fluoro, and trifluoromethyl;

R² is hydrogen; or, R² may be taken with R¹ and the phenyl ring to which R¹ and R² are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R^3 is hydrogen, methyl, methoxy, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;




wherein

R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of methylphenyl, 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4})alkylsulfonyl, C_{6-10} aryloxy(C_{1-4})alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, 4- C_{1-4} alkoxycarbonyl(C_{1-4})alkyl-sulfonyl, trifluoromethylsulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, di(C_{1-4} alkyl)aminosulfonylamino(C_{1-4})alkyl-sulfonyl, C_{1-4} alkylsulfonylamino(C_{1-4})alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C_{1-4})alkyl-sulfonyl, (2H)- 1,3-dione-1H-isoindol-1-yl-(C_{1-4})alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C_{1-6} alkylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R^6)-piperidin-1-ylsulfonyl wherein R^6 is C_{1-4} alkoxycarbonyl or C_{1-4} alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, methoxymethylcarbonyl, trifluoromethylcarbonyl, C_{1-4} alkoxycarbonylamino(C_{2-4})alkyl-carbonyl, C_{1-4} alkoxycarbonylaminomethyl-carbonyl, C_{1-4} alkoxycarbonyl, di(C_{1-4} alkyl)aminocarbonyl, C_{1-4} alkoxycarbonyl, and pyridinylcarbonyl; and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof.

12. The compound of Formula (I) as in claim 1



Formula (I)


wherein

R^1 is selected from the group consisting of ethyl, methoxy, ethoxy, isopropoxy, difluoroethoxy, dimethylamino, and trifluoromethoxy;

R^2 is hydrogen; or, R^2 may be taken with R^1 and the phenyl ring to which R^1 and R^2 are both attached to form 2,3-dihydrobenzofuran-7-yl or 1,3-benzodioxol-4-yl;

R^3 is hydrogen, methyl, chloro, or fluoro;

Q is selected from the group consisting of Q1, Q2, and Q3;



wherein


R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4})alkylsulfonyl, C_{6-10} aryloxy(C_{1-4})alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C_{1-4} alkoxycarbonyl(C_{1-4})alkyl-sulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-

trifluoroethylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, di(C₁₋₄alkyl)aminosulfonylamino(C₁₋₄alkyl)-sulfonyl, C₁₋₄alkylsulfonylamino(C₁₋₄alkyl)-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C₁₋₄alkyl)-sulfonyl, (2H)-1,3-dione-1H-isindol-1-yl-(C₁₋₄alkyl)-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C₁₋₆alkylsulfonyl, di(C₁₋₄alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R⁶)-piperidin-1-ylsulfonyl wherein R⁶ is C₁₋₄alkoxycarbonyl or C₁₋₄alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C₁₋₄alkoxycarbonylamino(C₂₋₄alkyl)-carbonyl, and C₁₋₄alkoxycarbonyl; and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof.

13. The compound of Formula (I) as in claim 1



Formula (I)


wherein

R¹ is selected from the group consisting of ethyl, methoxy, ethoxy, isopropoxy, difluoroethoxy, dimethylamino, and trifluoromethoxy;


R² is hydrogen;

R³ is hydrogen, methyl, chloro, or fluoro;


Q is selected from the group consisting of Q1, Q2, and Q3;



Q1



Q2



Q3


wherein

R^4 is chloro;

R^5 is independently selected from the group consisting of C_{1-4} alkoxy and chloro;

G is selected from the group consisting of 2-chloro-quinolin-6-yl, C_{6-10} aryl(C_{1-4})alkylsulfonyl, C_{6-10} aryloxy(C_{1-4})alkylsulfonyl, 3,5-dimethylisoxazol-4-yl-sulfonyl, C_{1-4} alkoxycarbonyl(C_{1-4})alkyl-sulfonyl, C_{3-7} cycloalkylsulfonyl, 2,2,2-trifluoroethylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, di(C_{1-4} alkyl)aminosulfonylamino(C_{1-4})alkyl-sulfonyl, C_{1-4} alkylsulfonylamino(C_{1-4})alkyl-sulfonyl, 2,5-dione-pyrrolidin-1-yl-(C_{1-4})alkyl-sulfonyl, (2H)-1,3-dione-1H-isoindol-1-yl-(C_{1-4})alkyl-sulfonyl, 1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl, C_{1-6} alkylsulfonyl, di(C_{1-4} alkyl)aminosulfonyl, pyrrolidin-1-ylsulfonyl, piperidin-1-ylsulfonyl, 4-(R^6)-piperidin-1-ylsulfonyl wherein R^6 is C_{1-4} alkoxycarbonyl or C_{1-4} alkyl; azepan-1-ylsulfonyl, morpholin-4-ylsulfonyl, 2,6-dimethylmorpholin-4-ylsulfonyl, C_{1-4} alkoxycarbonylamino(C_{2-4})alkyl-carbonyl, and C_{1-4} alkoxycarbonyl; and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof.

14. The compound of Formula (I) as in claim 1



Formula (I)

selected from the group consisting of
tert-Butyl 3-(4-chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;
3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-[(trifluoromethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(pyridin-3-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(pyridin-4-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(trifluoroacetyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(pyridin-2-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(furan-2-ylcarbonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-[(5-methylisoxazol-3-yl)carbonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethylphenyl)-5-(methoxyacetyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

tert-Butyl {2-[3-(4-chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;

N-{2-[3-(4-Chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} methanesulfonamide;

N-{2-[3-(4-Chlorophenyl)-2-(2-ethylphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl}-1,1,1-trifluoromethanesulfonamide;

tert-Butyl 3-(4-chlorophenyl)-2-[2-(trifluoromethyl)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;

3-(4-Chlorophenyl)-5-(methylsulfonyl)-2-[2-(trifluoromethyl)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2-[2-(trifluoromethyl)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-5-[(4-methylphenyl)sulfonyl]-2-[2-(trifluoromethyl)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

tert-Butyl {2-[3-(4-chlorophenyl)-2-[2-(trifluoromethyl)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;

tert-Butyl 3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;

3-(4-Chlorophenyl)-5-(methylsulfonyl)-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-5-(trifluoroacetyl)-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

tert-Butyl {2-[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;

3-[3-(4-Chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]benzonitrile;

Methyl 4-[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]benzoate;

4-[3-(4-Chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]benzonitrile;

4-[3-(4-Chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-N-(2-morpholin-4-ylethyl)benzenesulfonamide;

2-Chloro-6-[3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]quinoline;

tert-Butyl 3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxylate;

tert-Butyl {2-[3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]-2-oxoethyl} carbamate;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-5-(cyclopropylsulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-carboxamide;

3-(4-Chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(2,2,2-trifluoroethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

2-(2-{{3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl}sulfonyl}ethyl)-1H-isoindole-1,3(2H)-dione;

5-(Benzylsulfonyl)-3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-5-(cyclopentylsulfonyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(1-methylbutyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

1-(2-{{3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl}sulfonyl}ethyl)pyrrolidine-2,5-dione;

3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(2-phenoxyethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

N-(2-{{3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl}sulfonyl}ethyl)methanesulfonamide;

N-(2-{{3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl}sulfonyl}ethyl)propane-2-sulfonamide;

N'-(2-{{3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl}sulfonyl}ethyl)-N,N-dimethylsulfamide;

Methyl 3-{{3-(4-chlorophenyl)-2-[2-(trifluoromethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl}sulfonyl}propanoate;

3-(4-Chlorophenyl)-5-[(1,1-dioxidotetrahydrothiophen-3-yl)sulfonyl]-2-[2-(trifluoromethoxy)phenyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2-ethoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

- 3-(4-Chlorophenyl)-5-(cyclopentylsulfonyl)-2-(2-ethoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(2-ethoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 5-(Benzylsulfonyl)-3-(4-chlorophenyl)-2-(2-ethoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-ethoxyphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- 5-(Azepan-1-ylsulfonyl)-3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-N,N-diethyl-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- Methyl 1- {[3-(4-chlorophenyl)-2-(2-methoxyphenyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl]sulfonyl} piperidine-4-carboxylate;
- 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(morpholin-4-ylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(pyrrolidin-1-ylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-5- {[(2S,6S)-2,6-dimethylmorpholin-4-yl]sulfonyl }-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-methoxyphenyl)-5-(piperidin-1-ylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-N,N-dimethyl-2-[2-(1-methylethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- 2-(4-Chloro-2-methoxyphenyl)-3-(4-chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 2-(4-Chloro-2-methoxyphenyl)-3-(4-chlorophenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- 3-(4-Chlorophenyl)-2-(4-methoxy-2-methylphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

- 3-(4-Chlorophenyl)-2-(4-methoxy-2-methylphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(4-methoxy-2-methylphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(4-methoxy-2-methylphenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;
- 3-(4-Chlorophenyl)-2-(4-fluoro-2-methoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(4-fluoro-2-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-5-(cyclohexylsulfonyl)-2-(4-fluoro-2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(4-fluoro-2-methoxyphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-[(1-methylbutyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-methoxy-4-methylphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-fluoro-4-methoxyphenyl)-5-(methylsulfonyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-fluoro-4-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 3-(4-Chlorophenyl)-2-(2-fluoro-4-methoxyphenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 2-(1,3-Benzodioxol-4-yl)-3-(4-chlorophenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;
- 2-(1,3-Benzodioxol-4-yl)-3-(4-chlorophenyl)-5-[(2-methylpropyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

2-(1,3-Benzodioxol-4-yl)-3-(4-chlorophenyl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2,3-dihydro-1-benzofuran-7-yl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2,3-dihydro-1-benzofuran-7-yl)-5-[(2-methylpropyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-(2,3-dihydro-1-benzofuran-7-yl)-5-[(4-methylpiperidin-1-yl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole

3-(4-Chlorophenyl)-5-[(3,5-dimethylisoxazol-4-yl)sulfonyl]-2-(2-methoxyphenyl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

2-[3-(4-Chlorophenyl)-5-[(1-methylethyl)sulfonyl]-5,6-dihydropyrrolo[3,4-c]pyrazol-2(4H)-yl]-N,N-dimethylaniline;

2-[3-(4-Chlorophenyl)-5-(ethylsulfonyl)-5,6-dihydropyrrolo[3,4-c]pyrazol-2(4H)-yl]-N,N-dimethylaniline;

3-(4-Chlorophenyl)-N,N-dimethyl-2-[2-(2,2,2-trifluoroethoxy)phenyl]-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;

2-(2-tert-Butoxyphenyl)-3-(4-chlorophenyl)-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;

3-(6-Chloropyridin-3-yl)-2-(2-methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-{2-(2-Methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazol-3-yl} quinoline;

2-(2-Methoxyphenyl)-5-[(1-methylethyl)sulfonyl]-3-(6-methylpyridin-3-yl)-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

3-(4-Chlorophenyl)-2-[2-(2,2-difluoroethoxy)phenyl]-N,N-dimethyl-2,6-dihydropyrrolo[3,4-c]pyrazole-5(4H)-sulfonamide;

2-(2-Methoxyphenyl)-3-[6-(1-methylethoxy)pyridin-3-yl]-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

2-(2-Methoxyphenyl)-3-(6-methoxypyridin-3-yl)-5-[(1-methylethyl)sulfonyl]-2,4,5,6-tetrahydropyrrolo[3,4-c]pyrazole;

and pharmaceutically acceptable salt forms thereof.

15. A pharmaceutical composition comprising a compound of claim 1 or 14 and at least one of a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient, and a pharmaceutically acceptable diluent.
16. A pharmaceutical composition of claim 15, wherein the composition is a solid oral dosage form.
17. A pharmaceutical composition of claim 15, wherein the composition is a syrup, an elixir or a suspension.