

NOVEL 4-(SUBSTITUTED-AMINO)-7H-PYRROLO[2,3-*d*]PYRIMIDINES AS LRRK2 INHIBITORS

5

Field of the Invention

The present invention relates to small molecule inhibitors of leucine-rich repeat kinase 2 (LRRK2). This invention also relates to methods of inhibiting, in mammals, including humans, LRRK2 by administration of the small molecule LRRK2 inhibitors. The present invention also relates to the treatment of Parkinson's Disease (PD) and other neurodegenerative and/or neurological disorders in mammals, including humans with the LRRK2 inhibitors. More particularly, this invention relates to 4-(substituted-amino)-7H-pyrrolo[2,3-*d*]pyrimidine compounds useful for the treatment of neurodegenerative and/or neurological disorders, such as PD, Alzheimer's Disease (AD) and other LRRK2 associated disorders.

15

Background of the Invention

LRRK2 is a 286 kDa active protein kinase in the ROCO protein family with a complex multidomain structure and which has structural homology to the MAP kinase kinase kinases (MAPKKK). LRRK2 has been shown to phosphorylate moesin (at Thr⁵⁵⁸), ezrin and radixin in vitro. LRRK2 has been found in various regions of the brain as well as in the heart, lung, spleen and kidney. Independent domains that have been established for the LRRK2 protein include an ankyrin-like (ANK) domain, a leucine-rich repeat (LRR) domain, a Ras (renin-angiotensin system) of complex (ROC) domain, a C-terminal of ROC (COR) domain, a kinase (Kinase) domain and a C-terminal WD40 domain. The ROC domain binds guanosine triphosphate (GTP) and the COR domain may be a regulator of the ROC domain's GTPase activity.

25

With multiple domains and both active kinase and guanosine triphosphatase (GTPase) activity, LRRK2 appears to play a complex role in multiple cellular processes. For example, LRRK2 has been associated with NFAT inhibition in the immune system and has been linked to vesicle trafficking, presynaptic homeostasis, mammalian target of rapamycin (mTOR) signaling, signaling through the receptor tyrosine kinase MET in papillary renal and thyroid carcinomas, cytoskeletal dynamics, the mitogen-activated protein kinase (MAPK) pathway, the tumor necrosis factor- α (TNF- α) pathway, the Wnt pathway and autophagy. Recent genome-wide association (GWA) genetic studies have

30

implicated LRRK2 in the pathogenesis of various human diseases such as PD, inflammatory bowel disease (Crohn's disease), cancer and leprosy (Lewis, P.A. and Manzoni, C. Science Signaling 2012, 5(207), pe2).

Parkinson's Disease is a relatively common age-related neurodegenerative disorder resulting from the progressive loss of dopamine-producing neurons and which affects up to 4% of the population over 80. PD is characterized by both motoric and non-motoric symptoms such as tremor at rest, rigidity, akinesia and postural instability as well as non-motor symptoms such as impairment of cognition, sleep and sense of smell. Genome-wide association studies have linked LRRK2 to PD and many patients with point mutations in LRRK2 present symptoms that are indistinguishable from those with idiopathic PD. Over 20 LRRK2 mutations have been associated with autosomal-dominant parkinsonism, and the R1441C, R1441G, R1441H, Y1699C, G2019S, I2020T and N1437H missense mutations are considered to be pathogenic. The LRRK2 R1441G mutation has been shown to increase the release of proinflammatory cytokines (higher levels of TNF- α , IL-1 β , IL-12 and lower levels of IL-10) in microglial cells from transgenic mice and thus may result in direct toxicity to neurons (Gillardon, F. et al. Neuroscience 2012, 208, 41-48). In a murine model of neuroinflammation, induction of LRRK2 in microglia was observed and inhibition of LRRK2 kinase activity with small molecule LRRK2 inhibitors (LRRK2-IN-1 or sunitinib) or LRRK2 knockout resulted in attenuation of TNF- α secretion and nitric oxide synthase (iNOS) induction (Moehle, M. et al. J. Neurosci. 2012, 32(5), 1602-1611). The most common of the LRRK2 mutations, G2019S, is present in more than 85% of PD patients carrying LRRK2 mutations. This mutation, which is present in the LRRK2 kinase domain, leads to an enhancement of LRRK2 kinase activity. In the human brain LRRK2 expression is highest in the same regions of the brain that are impacted by PD, and LRRK2 is found in Lewy Bodies, a hallmark of PD. Recent studies indicate that a potent, selective, brain-penetrant kinase inhibitor for LRRK2 could be a therapeutic treatment for PD.

Dementia results from a wide variety of distinctive pathological processes. The most common pathological processes causing dementia are AD, cerebral amyloid angiopathy (CM) and prion-mediated diseases (see, e.g., Haan et al., Clin. Neurol. Neurosurg. 1990, 92(4):305-310; Glenner et al., J. Neurol. Sci. 1989, 94:1-28). AD is a progressive, neurodegenerative disorder characterized by memory impairment and cognitive dysfunction. AD affects nearly half of all people past the age of 85, the most

rapidly growing portion of the United States population. As such, the number of AD patients in the United States is expected to increase from about 4 million to about 14 million by 2050. LRRK2 mutations have been associated with AD-like pathology, which suggests that there may be a partial overlap between the neurodegenerative pathways in both AD and PD (Zimprach, A. et al. Neuron 2004, 44, 601-607). In addition, the LRRK2 R1628P variant (COR domain) has been associated with an increased incidence of AD in a certain population, perhaps resulting from increased apoptosis and cell death (Zhao, Y. et al.; Neurobiology of Aging 2011, 32, 1990-1993).


An increased incidence of certain non-skin cancers such as renal, breast, lung and prostate cancers, as well as acute myelogenous leukemia (AML), has been reported in Parkinson's disease patients with the LRRK2 G2019S mutation (Saunders-Pullman, R. et al.; Movement Disorders, 2010, 25(15), 2536-2541). Since the G2019S mutation is associated with increased LRRK2 kinase activity, inhibition of this activity may be useful in the treatment of cancer, such as kidney, breast, lung, prostate and blood cancers.

Inflammatory bowel disease (IBD) or Crohn's disease (CD) is a complex disease and is believed to result from an inappropriate immune response to microbiota in the intestinal tract. Genome-wide association studies have recently identified LRRK2 as a major susceptibility gene for Crohn's disease, particularly the M2397T polymorphism in the WD40 domain (Liu, Z. et al. Nat. Immunol. 2011, 12, 1063-1070). In a recent study LRRK2 deficient mice were found to be more susceptible to dextran sodium sulfate induced colitis than their wild-type counterparts, indicating that LRRK2 may play a role in the pathogenesis of IBD (Liu, Z. and Lenardo, M.; Cell Research 2012, 1-3).

Both non-selective and selective small molecule compounds with LRRK2 inhibitory activity such as staurosporine, sunitinib, LRRK2-IN-1, CZC-25146, TAE684 and those in WO 2011/141756, WO 2012/028629 and WO 2012/058193 have been described. It is desirable to provide compounds which are potent and selective inhibitors of LRRK2 with a favorable pharmacokinetic profile and the ability to to traverse the blood brain barrier. Accordingly, the present invention is directed to novel 4-(substituted-amino)-7H-pyrrolo[2,3-d]pyrimidine compounds with LRRK2 inhibitory activity and the use of these compounds in the treatment of diseases associated with LRRK2, such as neurodegenerative diseases, including PD.

Summary of the Invention

A first embodiment of a first aspect of the present invention is a compound of Formula I




5 or a pharmaceutically acceptable salt thereof wherein R^1 and R^2 are each independently hydrogen, C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, a four to seven membered heterocycloalkyl which contains one to three heteroatoms selected from N, O and S; or a five to six membered heteroaryl which contains one to four heteroatoms selected from N, O and S, wherein the C_1 - C_6 alkyl, C_3 - C_7 cycloalkyl, four to seven membered heterocycloalkyl, or five to six membered heteroaryl are optionally substituted with one to three R^6 ; or R^1 and R^2 taken together with the nitrogen to which they are attached are a four to seven membered heterocycloalkyl which optionally contains one to two additional heteroatoms selected from N, O and S, and optionally contains one double bond; a six to eleven membered heterobicycloalkyl which optionally contains one to two additional heteroatoms selected from N, O and S; or a six to twelve membered heterospirocycloalkyl which optionally contains one to two additional heteroatoms selected from N, O and S; and wherein the four to seven membered heterocycloalkyl, six to eleven membered heterobicycloalkyl or six to twelve membered heterospirocycloalkyl is optionally substituted with one to three R^7 ; R^3 is phenyl or a five to ten membered heteroaryl which contains one to four heteroatoms selected from N, O and S; wherein the phenyl and five to ten membered heteroaryl are optionally substituted with one to three R^9 and wherein the phenyl is optionally fused with a C_5 - C_6 cycloalkyl or a five to six membered heterocycloalkyl which contains one to three heteroatoms selected from N, O and S and which is optionally substituted with oxo; R^4 and R^5 are each independently hydrogen or C_1 - C_3 alkyl; R^6 at each occurrence is independently selected from C_1 - C_3 alkyl, C_1 - C_3 alkoxy, hydroxy, halo, $-NR^aR^b$, $-C(O)NR^aR^b$, or a four to seven membered heterocycloalkyl which contains one to three heteroatoms selected from N, O and S; R^7 at each occurrence is independently

selected from halo, hydroxy, cyano, $\text{-NR}^a\text{R}^b$, $\text{-C(O)NR}^a\text{R}^b$, $\text{C}_1\text{-C}_6\text{alkyl}$, $\text{C}_1\text{-C}_6\text{alkoxy}$, phenyl, a five to six membered heteroaryl containing one to four heteroatoms selected from N, O and S, or two R^7 when attached to the same carbon and taken together can be oxo; wherein the $\text{C}_1\text{-C}_6\text{alkyl}$, phenyl and five to six membered heteroaryl are optionally substituted with one to three R^8 ; R^8 at each occurrence is independently hydroxy, halo, cyano, $\text{C}_1\text{-C}_3\text{alkoxy}$, NR^aR^b , $\text{C}_1\text{-C}_3\text{alkyl}$ optionally substituted with one to three halo, $\text{C}_3\text{-C}_7\text{cycloalkyl}$, phenoxy optionally substituted with cyano, or a five to six membered heteroaryloxy containing one to four heteroatoms selected from N, O and S and which is optionally substituted with one or two halo or $\text{C}_1\text{-C}_3\text{alkyl}$; R^9 at each occurrence is independently cyano, halo, hydroxy, $\text{C}_1\text{-C}_3\text{alkyl-S-}$, $\text{-CO}_2\text{H}$, -C(O)NH_2 , $\text{-S(O)}_2\text{NH}_2$, $\text{C}_1\text{-C}_3\text{alkyl}$ optionally substituted with one to three halo or hydroxy, or $\text{C}_1\text{-C}_3\text{alkoxy}$ optionally substituted with one to three halo or hydroxy; and R^a and R^b at each occurrence are each independently hydrogen, $\text{C}_1\text{-C}_6\text{alkyl}$, $\text{C}_3\text{-C}_7\text{cycloalkyl}$ or $\text{-C(O)C}_1\text{-C}_6\text{alkyl}$.

A second embodiment of a first aspect of the present invention is the compound of the first embodiment of the first aspect wherein R^4 and R^5 are each hydrogen; or a pharmaceutically acceptable salt thereof.

A third embodiment of a first aspect of the present invention is the compound of the second embodiment of the first aspect wherein R^1 and R^2 taken together with the nitrogen to which they are attached are azetidin-1-yl, pyrrolidin-1-yl, piperidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, 6-oxa-3-aza-bicyclo[3.1.1]heptan-3-yl,



each of which is optionally substituted with one to three R^7 ; or a pharmaceutically acceptable salt thereof.

A fourth embodiment of a first aspect of the present invention is the compound of the third embodiment of the first aspect wherein R^1 and R^2 taken together with the nitrogen to which they are attached are piperidin-1-yl or morpholin-4-yl; each optionally

substituted with a hydroxy, methyl or 5-methyl-1,2,4-oxadiazol-3-yl; or a pharmaceutically acceptable salt thereof.


A fifth embodiment of a first aspect of the present invention is the compound of the fourth embodiment of the first aspect wherein R^1 and R^2 taken together with the nitrogen to which they are attached are morpholin-4-yl optionally substituted with a methyl or 5-methyl-1,2,4-oxadiazol-3-yl; or a pharmaceutically acceptable salt thereof.

A sixth embodiment of a first aspect of the present invention is the compound of the second embodiment of the first aspect wherein R^1 and R^2 taken together with the nitrogen to which they are attached are methylamino, dimethylamino, diethylamino, N-methylcyclopropylamino or pyrazolylamino; each of which is optionally substituted with one to three R^6 ; or a pharmaceutically acceptable salt thereof.

An seventh embodiment of a first aspect of the present invention is the compound of the third embodiment of the first aspect wherein R^3 is phenyl substituted with one or two R^9 ; and each R^9 is independently selected from cyano, fluoro, chloro or methoxy; or a pharmaceutically acceptable salt thereof.


An eighth embodiment of a first aspect the present invention is the compound of the third embodiment of the first aspect wherein R^3 is pyrazolyl, isothiazolyl, or pyridinyl, each optionally substituted with one R^9 ; and R^9 is cyano or methyl; or a pharmaceutically acceptable salt thereof.

A ninth embodiment of a first aspect of the present invention is the compound of the first embodiment of the first aspect wherein R^1 and R^2 taken together with the nitrogen to which they are attached is selected from the group consisting of:




R³ is 1-methylpyrazol-4-yl, 1*H*-pyrazol-4-yl, 2-cyanopyridin-6-yl, 3-methyl-1,2-thiazol-5-yl, 5-cyano-1-methylpyrrol-3-yl, 3-methylpyridin-5-yl, 3-cyanophenyl, 2-fluorophenyl, 3-fluorophenyl, 2,3-difluorophenyl, 2-fluoro-5-chlorophenyl, 2-fluoro-3-cyanophenyl or 2-methoxy-5-fluorophenyl; and R⁴ and R⁵ are each independently hydrogen or methyl; or a pharmaceutically acceptable salt thereof.

A tenth embodiment of a first aspect of the present invention is the compound of the ninth embodiment wherein R¹ and R² taken together with the nitrogen to which they are attached are a group selected from:



and R^4 and R^5 are each hydrogen; or a pharmaceutically acceptable salt thereof.

An eleventh embodiment of a first aspect of the present invention is the compound of the tenth embodiment of the first aspect wherein R^1 and R^2 taken together
5 with the nitrogen to which they are attached is



or a pharmaceutically acceptable salt thereof.

A twelfth embodiment of a first aspect of the present invention is the compound of the eleventh embodiment of the first aspect wherein R^3 is 3-cyanophenyl, 1-methylpyrazol-4-yl or 5-cyano-1-methylpyrrol-3-yl; or a pharmaceutically acceptable salt
10 thereof.

A thirteenth embodiment of a first aspect of the present invention is the compound of the first embodiment of a first aspect wherein R^1 and R^2 taken together with the nitrogen to which they are attached are piperidinyl or 3-hydroxypiperidinyl; R^3 is
15 1-methylpyrazol-4-yl, 1*H*-pyrazol-4-yl, 2-cyanopyridin-6-yl, 3-methyl-1,2-thiazol-5-yl, 3-methylpyridin-5-yl, 3-cyanophenyl, 2-fluorophenyl, 3-fluorophenyl, 2,3-difluorophenyl, 2-fluoro-5-chlorophenyl, 2-fluoro-3-cyanophenyl or 2-methoxy-5-fluorophenyl; and R^4 and R^5 are each independently hydrogen or methyl; or a pharmaceutically acceptable salt thereof.

A fourteenth embodiment of a first aspect of the invention is a compound of the first embodiment of the first aspect selected from the group consisting of:

5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

3-[6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile;

3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

- 3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 (3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}phenyl)methanol;
 4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[2-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5 {3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 3-[4-(3,3-dimethylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(piperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 1-(5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl)piperidine-3-carboxamide;
 3-{4-[(3*S*)-3-hydroxypiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 10 4-[(3*S*)-3-methylpiperidin-1-yl]-5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-[(3*S*)-3-methylpiperidin-1-yl]-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-phenyl-4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 [1-(5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl)piperidin-4-yl]methanol;
 15 1-{5-[3-(hydroxymethyl)phenyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl}piperidine-3-carbonitrile;
 1-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidine-3-carbonitrile;
 4-(3,5-*cis*-dimethylpiperidin-1-yl)-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-methoxy-3-[4-(3-methylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 20 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-4-methoxybenzonitrile;
 5-(5-chloro-2-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 3-{4-[4-(1*H*-irridazol-2-yl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[3-(methoxymethyl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-[4-(9-methyl-1-oxa-4,9-diazaspiro[5.5]undec-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-
 25 yl]benzonitrile;
 3-[4-(3-methoxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
*N*³-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]-*N,N*-dimethyl-beta-alaninamide;
 3-[4-(4,4-difluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 30 3-[4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[4-(1*H*-pyrazol-3-yl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[(3*S*)-3-fluoropyrrolidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[2-(1*H*-pyrazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

- 3-{4-[3-(1*H*-pyrazol-3-yl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
3-{4-[3-(1*H*-imidazol-2-yl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
3-{4-[(1-methylpiperidin-3-yl)amino]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
3-[4-(3-oxo-2,7-diazaspiro[4.5]dec-7-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
5 3-[4-((3*R*)-3-methylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-(4-{2-(morpholin-4-yl)ethyl}amino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
3-[4-(2-oxa-7-azaspiro[4.5]dec-7-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
N-{1-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-4-yl}acetamide;
5-(1*H*-indazol-5-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
10 3-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzenesulfonamide;
5-(2-fluorophenyl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(1*H*-indazol-4-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(6-fluoro-5-methylpyridin-3-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
15 5-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2,3-dihydro-1*H*-
isoindol-1-one;
4-[(3*S*)-3-methylpiperidin-1-yl]-5-(pyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
4-[(3*S*)-3-methylpiperidin-1-yl]-5-(5-methylpyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
4-[(3*S*)-3-methylpiperidin-1-yl]-5-(7*H*-pyrrolo[2,3-*b*]pyridin-5-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
20 6-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2,3-dihydro-1*H*-
isoindol-1-one;
4-[(3*S*)-3-methylpiperidin-1-yl]-5-(1*H*-pyrrolo[3,2-*b*]pyridin-6-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
25 4-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenol;
4-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzamide;
3-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenol;
5-(2-chloro-5-methylpyridin-3-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
30 3-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzamide;
3-[4-(3-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-[4-(4-methylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzoic acid;
3-[4-(methylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

- 3-[4-(3,5-*cis*-dimethylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-4-methoxybenzonitrile;
 4-methoxy-3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
- 5 *N,N*-dimethyl-5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2-fluorobenzonitrile;
 3-[4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-((3*R*)-3-methylpyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-4-fluorobenzonitrile;
- 10 3-[4-(diethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*R*)-2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-5-fluorobenzonitrile;
 3-{4-[2-(1*H*-pyrazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
- 15 3-[4-((3*S*)-3-methylpyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*R*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-(4-fluoropiperidin-1-yl)-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 20 2-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(3-chlorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluoro-5-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2,5-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 25 5-(2,3-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-chloro-2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 {2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 5-(2,4-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 30 5-(3,5-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(morpholin-4-yl)-5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-chlorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

- 5-(5-fluoro-2-methylphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 {2-fluoro-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 5-(4-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5 {3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 5-(2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-[3-(methylsulfanyl)phenyl]-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(morpholin-4-yl)-5-(pyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]isoquinoline;
 10 5-(5-bromopyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-chloro-5-methylphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-methylphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(1-methyl-1*H*-pyrazol-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[4-(4-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 15 3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-methyl-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 20 3-[4-(6-oxa-3-azabicyclo[3.1.1]hept-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-chloro-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-methoxy-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5-chloro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 6-methyl-5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 25 3-methoxy-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(thiomorpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-[5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 4-[(2*S*)-2-methylmorpholin-4-yl]-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 30 4-[(2*R*)-2-methylmorpholin-4-yl]-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(3-fluoropiperidin-1-yl)-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 {4-[5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]morpholin-2-yl}methanol;

- 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(2-[[6-methylpyridin-3-yl]oxy]methyl)morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- N,N*-dimethyl-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
- N*-cyclopropyl-*N*-methyl-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
- 3-[4-(3,3-difluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-(3-aminopiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-(2-[[5-(difluoromethyl)-1,2,4-oxadiazol-3-yl]methyl]morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-[2-(1,3-thiazol-2-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-(4-oxopiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-[4-fluoro-4-(hydroxymethyl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-[2-(3-hydroxyphenyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-[2-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-(3,3-difluoropyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-(4-{2-[(3-cyanophenoxy)methyl]morpholin-4-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile;
- 3-[4-(3-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 1-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidine-3-carboxamide;
- 3-[4-(2-ethylmorpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-[2-(pyrimidin-4-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-[2-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-(4-{2-[(dimethylamino)methyl]morpholin-4-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile;
- 3-[4-(1*H*-pyrazol-4-ylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-(4-{2-[(5-methyl-1,2,4-oxadiazol-3-yl)methyl]morpholin-4-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile;
- 3-[4-[3-(5-methyl-1,3,4-oxadiazol-2-yl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-5-methoxybenzonitrile;
- 5-(1-ethyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 5-(5-methylpyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

- 5-(5-chloropyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(6-methoxypyrazin-2-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(1,3-dimethyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
3-{4-[(3*S*)-3-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
5 2-fluoro-3-[2-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-{4-[(3*R*)-3-hydroxypiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
1-[5-(3-fluorophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
5-(5-fluoro-2-methoxyphenyl)-4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-
pyrrolo[2,3-*d*]pyrimidine;
10 2-fluoro-3-[4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
4-(4-fluoropiperidin-1-yl)-5-(2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(3-fluoro-5-methoxyphenyl)-4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
3-[4-(4,4-difluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2-fluorobenzonitrile;
1-[5-(5-methylpyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
15 5-(5-fluoro-2-methoxyphenyl)-4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
1-[5-(3-fluoro-5-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
1-[5-(2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
2-fluoro-3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]-
pyrimidin-5-yl}benzonitrile;
20 2-fluoro-3-{4-[2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-
yl}benzonitrile;
4-(azetidin-1-yl)-5-(5-fluoro-2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
4-[(2*S*)-2-methylmorpholin-4-yl]-5-(5-methylpyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(3-fluorophenyl)-4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
25 5-(3-fluorophenyl)-4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]-
pyrimidine;
5-(2-methoxyphenyl)-4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(2-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
2-fluoro-3-{4-[(3*S*)-3-fluoropyrrolidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
30 4-[(3*S*)-3-fluoropyrrolidin-1-yl]-5-(2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(2-methoxyphenyl)-4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
5-(3-fluoro-5-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
2-fluoro-3-[4-(3-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

- 5-(5-fluoro-2-methoxyphenyl)-4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 5-(5-fluoro-2-methoxyphenyl)-4-(3-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 5 5-(3-fluorophenyl)-4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 2-fluoro-3-[4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5-fluoro-2-methoxyphenyl)-4-[(3*S*)-3-fluoropyrrolidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
- 10 5-(4-methyl-1,3-thiazol-2-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(4-methylpyridin-2-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-fluoro-6-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 5-(2-fluoro-6-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 15 5-(2,6-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-methyl-1,2-thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-chloro-3-fluoro-6-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(4-methoxypyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
- 20 3-{4-[2-((5*R*)-5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[2-((5*S*)-5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[(2*S*)-2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
- 25 or a pharmaceutically acceptable salt thereof.

A fifteenth embodiment of a first aspect of the present invention is a compound of the first embodiment of a first aspect selected from the group consisting of:

- 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
- 30 6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile;
 3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[4-(piperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

- 2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2,3-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-chloro-2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5 5-(3-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-[5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 5-(5-methylpyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-[4-(3-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 10 2-fluoro-3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 5-(3-methyl-1,2-thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine; and
 3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-
 15 yl}benzonitrile;
 or a pharmaceutically acceptable salt thereof.

A sixteenth embodiment of a first aspect of the present invention is the compound of the first embodiment of the first aspect selected from the group consisting of:

- 20 3-[6-(difluoromethyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5,6-dihydro-2*H*-pyran-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3,4-dihydro-2*H*-pyran-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(morpholin-4-yl)-5-[3-(1,2,4-oxadiazol-3-yl)phenyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[2-(3-methyl-1,2-oxazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-
 25 yl}benzonitrile;
 2-methyl-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridin-2(1*H*)-one;
 5-(imidazo[2,1-*b*][1,3]thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
rel-3-{4-[(3*aS*,6*aS*)-hexahydro-5*H*-furo[2,3-*c*]pyrrol-5-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-
 30 yl}benzonitrile;
rel-3-{4-[(3*aR*,6*aS*)-tetrahydro-1*H*-furo[3,4-*c*]pyrrol-5(3*H*)-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-
 5-yl}benzonitrile;

- rel*-3-{4-[(4*aR*,7*aS*)-hexahydrocyclopenta[*b*][1,4]oxazin-4(4*aH*)-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]morpholine-2-carbonitrile;
 3-[4-(2,2-dimethylmorpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5 5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]thiophene-2-carbonitrile;
 5-(imidazo[1,2-*b*]pyridazin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-{4-[2(*R*)-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-{4-[2(*S*)-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 10 *d*]pyrimidin-5-yl}benzonitrile;
 6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carboxamide;
 4-(morpholin-4-yl)-5-(pyrazolo[1,5-*a*]pyrimidin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-methyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrrole-2-carbonitrile;
 5-(6-methylimidazo[2,1-*b*][1,3]thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 15 *d*]pyrimidine;
 3-{4-[2-(1,2-oxazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 1-methyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-imidazole-2-carbonitrile;
 4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]thiophene-2-carbonitrile;
 20 4-(morpholin-4-yl)-5-(pyrazolo[1,5-*a*]pyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1,5-dimethyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrrole-2-carbonitrile;
 1-methyl-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrazole-5-carbonitrile; and
 25 3-{4-[2-(cyanomethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 or a pharmaceutically acceptable salt thereof.

A first embodiment of a second aspect of the present invention is a pharmaceutical composition comprising a therapeutically effective amount of a compound according to any one of the first through sixteenth embodiments of the first aspect of the invention, or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier.

A first embodiment of a third aspect of the present invention is a method of treating Parkinson's disease in a patient, the method comprising administering to a patient in need thereof a therapeutically effective amount of a compound or pharmaceutically acceptable salt thereof according to any one of the first through
5 sixteenth embodiments of the first aspect of the invention.

Another aspect of the present invention is the compound or pharmaceutically acceptable salt thereof according to any one of the first through sixteenth embodiments of the first aspect of the invention for use in the treatment of Parkinson's disease.

Another aspect of the present invention is the use of any of the preceding
10 compounds of formula I and their compositions for inhibiting LRRK2 kinase. In a further embodiment, the compounds of formula I or compositions thereof are useful for treating a neurodegenerative disease. In yet another embodiment, the neurodegenerative disease is Parkinson's Disease.

Accordingly, the invention is also directed to methods of treating a patient
15 (preferably a human) for diseases in which the LRRK2 kinase is involved, such as Parkinson's Disease, by administering a therapeutically effective amount of a compound of any of the embodiments of formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The invention is also directed to methods of inhibiting LRRK2 kinase activity, by
20 administering a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, to a mammal or a patient in need thereof. The invention is also directed to methods of treating disorders responsive to the inhibition of LRRK2 kinase activity, such as neurological disorders (particularly Parkinson's disease), certain cancers, and certain
25 immunological disorders (such as Crohn's disease and leprosy) by administering a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, to a mammal or a patient in need thereof.

The invention is also directed to methods for treating conditions or diseases of the
30 central nervous system and neurological disorders in which the LRRK2 kinase is involved, particularly Parkinson's disease (but also including other neurological diseases which may include migraine; epilepsy; Alzheimer's disease; brain injury; stroke; cerebrovascular diseases (including cerebral arteriosclerosis, cerebral amyloid

angiopathy, hereditary cerebral hemorrhage, and brain hypoxia-ischemia); cognitive disorders (including amnesia, senile dementia, HIV-associated dementia, Alzheimer's disease, Huntington's disease, Lewy body dementia, vascular dementia, drug-related dementia, tardive dyskinesia, myoclonus, dystonia, delirium, Pick's disease, Creutzfeldt-Jacob disease, HIV disease, Gilles de la Tourette's syndrome, epilepsy, muscular spasms and disorders associated with muscular spasticity or weakness including tremors, and mild cognitive impairment); mental deficiency (including spasticity, Down syndrome and fragile X syndrome); sleep disorders (including hypersomnia, circadian rhythm sleep disorder, insomnia, parasomnia, and sleep deprivation) and psychiatric disorders such as anxiety (including acute stress disorder, generalized anxiety disorder, social anxiety disorder, panic disorder, post-traumatic stress disorder, agoraphobia, and obsessive-compulsive disorder); factitious disorder (including acute hallucinatory mania); impulse control disorders (including compulsive gambling and intermittent explosive disorder); mood disorders (including bipolar I disorder, bipolar II disorder, mania, mixed affective state, major depression, chronic depression, seasonal depression, psychotic depression, seasonal depression, premenstrual syndrome (PMS) premenstrual dysphoric disorder (PDD), and postpartum depression); psychomotor disorder; psychotic disorders (including schizophrenia, schizoaffective disorder, schizophreniform, and delusional disorder); drug dependence (including narcotic dependence, alcoholism, amphetamine dependence, cocaine addiction, nicotine dependence, and drug withdrawal syndrome); eating disorders (including anorexia, bulimia, binge eating disorder, hyperphagia, obesity, compulsive eating disorders and pagophagia); sexual dysfunction disorders; urinary incontinence; neuronal damage disorders (including ocular damage, retinopathy or macular degeneration of the eye, tinnitus, hearing impairment and loss, and brain edema) and pediatric psychiatric disorders (including attention deficit disorder, attention deficit/hyperactive disorder, conduct disorder, and autism) in a mammal, preferably a human, comprising administering to said mammal a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt thereof.

The text revision of the fourth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR) (2000, American Psychiatric Association, Washington D.C.) provides a diagnostic tool for identifying many of the disorders described herein. The skilled artisan will recognize that there are alternative nomenclatures, nosologies,

and classification systems for disorders described herein, including those as described in the DMS-IV-TR, and that terminology and classification systems evolve with medical scientific progress.

Preferred methods are for treating a neurological disorder, most preferably Parkinson's disease, (but also other neurological disorders such as migraine; epilepsy; Alzheimer's disease; Niemann-Pick type C; brain injury; stroke; cerebrovascular disease; cognitive disorder; sleep disorder) or a psychiatric disorder (such as anxiety; factitious disorder; impulse control disorder; mood disorder; psychomotor disorder; psychotic disorder; drug dependence; eating disorder; and pediatric psychiatric disorder) in a mammal, preferably a human, comprising administering to said mammal a therapeutically effective amount of a compound of Formula I or pharmaceutically acceptable salt thereof. In addition, the compounds of Formula I and pharmaceutically acceptable salts thereof may also be employed in methods of treating other disorders associated with LRRK2 such as Crohn's disease, leprosy and certain cancers, such as kidney, breast, lung, prostate, lung and blood cancer.

Also provided herein are compositions comprising a pharmaceutically effective amount of one or more of the compounds described herein and a pharmaceutically acceptable vehicle, carrier or excipient.

The present invention is also directed to the use of a combination of a LRRK2 inhibitor compound of formula I, and one or more additional pharmaceutically active agent(s).

Other features and advantages of this invention will be apparent from this specification and the appendent claims which describe the invention.

Definitions

The term "alkyl" refers to a linear or branched-chain saturated hydrocarbyl substituent (i.e., a substituent obtained from a hydrocarbon by removal of a hydrogen); in one embodiment from one to six carbon atoms (i.e., C₁-C₆alkyl); in another embodiment, from one to three carbon atoms (i.e., C₁-C₃alkyl). Examples of such substituents include methyl, ethyl, propyl (including *n*-propyl and isopropyl), butyl (including *n*-butyl, isobutyl, *sec*-butyl and *tert*-butyl), pentyl, isoamyl, hexyl and the like.

The term "alkoxy" refers to a linear or branched-chain saturated hydrocarbyl substituent (i.e., a substituent obtained from a hydrocarbon by removal of a hydrogen)

which is in turn attached to an oxygen atom; in one embodiment from one to six carbon atoms (i.e., C₁-C₆alkoxy); in another embodiment, from one to three carbon atoms (i.e., C₁-C₃alkoxy). Examples of such substituents include methoxy, ethoxy, propoxy (including *n*-propoxy and isopropoxy), butoxy (including *n*-butoxy, isobutoxy, *sec*-butoxy and *tert*-butoxy), pentoxy and the like.

The term "cycloalkyl" refers to a carbocyclic substituent obtained by removing a hydrogen from a saturated carbocyclic molecule and having the specified number of carbon atoms. In one embodiment, a cycloalkyl substituent has three to seven carbon atoms (i.e., C₃-C₇cycloalkyl). Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The term "cycloalkyl" includes mono-, bi- and tricyclic saturated carbocycles, as well as bridged and fused ring carbocycles, as well as spiro-fused ring systems.

In some instances, the number of atoms in a cyclic substituent containing one or more heteroatoms (i.e., heteroaryl or heterocycloalkyl) is indicated by the prefix "x to y membered", wherein x is the minimum and y is the maximum number of atoms forming the cyclic moiety of the substituent. The term "heterocycloalkyl" refers to a substituent obtained by removing a hydrogen from a saturated or partially saturated ring structure containing the specified number of ring atoms, wherein at least one of the ring atoms is a heteroatom (i.e., oxygen, nitrogen, or sulfur), with the remaining ring atoms being independently selected from the group consisting of carbon, oxygen, nitrogen, and sulfur. If the heterocycloalkyl substituent is in turn substituted with a group or substituent, the group or substituent may be bound to a nitrogen heteroatom, or it may be bound to a ring carbon atom, as appropriate. As used herein, the term "heterocycloalkyl" as used herein refers to a monocyclic ring system containing the heteroatoms N, O or S as specified. Thus, for example, "four to seven membered heterocycloalkyl" refers to a heterocycloalkyl containing from 4 to 7 atoms, including one or more heteroatoms, in the cyclic moiety of the heterocycloalkyl. The term "heterobicycloalkyl" as used herein refers to a non-spiro bicyclic ring system containing the heteroatoms N, O or S as specified. Thus, for example, "six to twelve membered heterobicycloalkyl" refers to a heterobicycloalkyl containing from 6 to 12 atoms, including one or more heteroatoms, in the cyclic moieties of the heterobicycloalkyl. The term "heterospirocycloalkyl" as used herein refers to a spirocyclic ring system containing the heteroatoms N, O or S as specified. For example, "six to twelve membered

heterospirocycloalkyl” means a six to twelve membered spirocyclic ring system which contains at least one heteroatom as specified.

The term “hydrogen” refers to a hydrogen substituent, and may be depicted as -H.

5 The term “hydroxy” or “hydroxyl” refers to -OH. Compounds bearing a carbon to which one or more hydroxy substituents are attached include, for example, alcohols, enols and phenol.

10 The term “halo” or “halogen” refers to fluoro (which may be depicted as -F), chloro (which may be depicted as -Cl), bromo (which may be depicted as -Br), or iodo (which may be depicted as -I).

15 The term “heteroaryl” refers to an aromatic ring structure containing the specified number of ring atoms in which at least one of the ring atoms is a heteroatom (i.e., oxygen, nitrogen, or sulfur), with the remaining ring atoms being independently selected from the group consisting of carbon, oxygen, nitrogen, and sulfur. A five to six membered heteroaryl is an aromatic ring system which has five or six ring atoms with at least one of the ring atoms being N, O or S. Similarly, a five to ten membered heteroaryl is an aromatic ring system which has five to ten ring atoms with at least one of the ring atoms being N, O or S. A heteroaryl may be a single ring or 2 fused rings. Examples of heteroaryl substituents include 6-membered ring substituents such as
20 pyridyl, pyrazyl, pyrimidinyl, and pyridazinyl; 5-membered ring substituents such as triazolyl, imidazolyl, furanyl, thiophenyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, 1,2,3-, 1,2,4-, 1,2,5-, or 1,3,4-oxadiazolyl and isothiazolyl; 6/5-membered fused ring substituents such as benzothiofuranyl, isobenzothiofuranyl, benzisoxazolyl, benzoxazolyl, purinyl, and anthranilyl; and 6/6-membered fused rings such as quinolinyl, isoquinolinyl, cinnolinyl, quinazolinyl, and 1,4-benzoxazinyl. In a group that has a heteroaryl substituent, the ring atom of the heteroaryl substituent that is bound to the group may be the at least one heteroatom, or it may be a ring carbon atom, where the ring carbon atom may be in the same ring as the at least one heteroatom or where the ring carbon atom may be in a different ring from the at least one heteroatom. Similarly,
25 if the heteroaryl substituent is in turn substituted with a group or substituent, the group or substituent may be bound to the at least one heteroatom, or it may be bound to a ring carbon atom, where the ring carbon atom may be in the same ring as the at least one heteroatom or where the ring carbon atom may be in a different ring from the at least
30

one heteroatom. The term "heteroaryl" also includes pyridyl *N*-oxides and groups containing a pyridine *N*-oxide ring.

Examples of single-ring heterocycloalkyls include azetidiny, oxetanyl, thietanyl, dihydrofuranyl, tetrahydrofuranyl, dihydrothiophenyl, tetrahydrothiophenyl, pyrrolinyl, pyrrolidinyl, imidazolinyl, imidazolidinyl, pyrazolinyl, pyrazolidinyl, thiazolinyl, isothiazolinyl, thiazolidinyl, isothiazolidinyl, dihydropyranyl, piperidinyl, morpholinyl, piperazinyl, azepinyl, oxepinyl, thiepinyl, and diazepinyl.

Examples of 2-fused-ring heteroaryls include, indolizinyl, pyranopyrrolyl, 4*H*-quinolizinyl, purinyl, naphthyridinyl, pyridopyridinyl (including pyrido[3,4-*b*]-pyridinyl, pyrido[3,2-*b*]-pyridinyl, or pyrido[4,3-*b*]-pyridinyl), and pteridinyl, indolyl, isoindolyl, indoleninyl, isoindazolyl, benzazinyl, phthalazinyl, quinoxalinyl, quinazolinyl, benzodiazinyl, benzopyranyl, benzothiopyranyl, benzoxazolyl, indoxazinyl, anthraninyl, benzodioxolyl, benzodioxanyl, benzoxadiazolyl, benzofuranyl, isobenzofuranyl, benzothienyl, isobenzothienyl, benzothiazolyl, benzothiadiazolyl, benzimidazolyl, benzotriazolyl, benzoxazinyl, benzisoxazinyl, pyrrolopyridinyl, pyrazolopyridinyl and imidazothiazolyl.

Other examples of fused-ring heteroaryls include benzo-fused heteroaryls such as indolyl, isoindolyl, indoleninyl, isoindazolyl, benzazinyl (including quinolinyl or isoquinolinyl), phthalazinyl, quinoxalinyl, quinazolinyl, benzodiazinyl (including cinnolinyl or quinazolinyl).

The foregoing groups, as derived from the groups listed above, may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached). Further, a group derived from imidazole may be imidazol-1-yl (N-attached) or imidazol-2-yl (C-attached).

If substituents are described as being "independently selected" from a group, each instance of a substituent is selected independent of the other. Each substituent therefore may be identical to or different from the other substituent(s).

As used herein the term "formula I" or "Formula I" may be referred to as a "compound(s) of the invention." Such terms are also defined to include all forms of the compound of formula I, including hydrates, solvates, isomers, crystalline and non-crystalline forms, isomorphs, polymorphs, and metabolites thereof. For example, the compounds of the invention, or pharmaceutically acceptable salts thereof, may exist in unsolvated and solvated forms. When the solvent or water is tightly bound, the complex

will have a well-defined stoichiometry independent of humidity. When, however, the solvent or water is weakly bound, as in channel solvates and hygroscopic compounds, the water/solvent content will be dependent on humidity and drying conditions. In such cases, non-stoichiometry will be the norm.

5 The compounds of the invention may exist as clathrates or other complexes. Included within the scope of the invention are complexes such as clathrates, drug-host inclusion complexes wherein the drug and host are present in stoichiometric or non-stoichiometric amounts. Also included are complexes of the compounds of the invention containing two or more organic and/or inorganic components which may be in
10 stoichiometric or non-stoichiometric amounts. The resulting complexes may be ionized, partially ionized, or non-ionized. For a review of such complexes, see J. Pharm. Sci., 64 (8), 1269-1288 by Haleblan (August 1975).

 The compounds of the invention may have asymmetric carbon atoms. The carbon-carbon bonds of the compounds of the invention may be depicted herein using a
15 solid line (—), a solid wedge (), or a dotted wedge (). The use of a solid line to depict bonds to asymmetric carbon atoms is meant to indicate that all possible stereoisomers (e.g., specific enantiomers, racemic mixtures, etc.) at that carbon atom are included. The use of either a solid or dotted wedge to depict bonds to asymmetric carbon atoms is meant to indicate that only the stereoisomer shown is
20 meant to be included. It is possible that compounds of Formula I may contain more than one asymmetric carbon atom. In those compounds, the use of a solid line to depict bonds to asymmetric carbon atoms is meant to indicate that all possible stereoisomers are meant to be included. For example, unless stated otherwise, it is intended that the compounds of Formula I can exist as enantiomers and diastereomers or as racemates
25 and mixtures thereof. The use of a solid line to depict bonds to one or more asymmetric carbon atoms in a compound of Formula I and the use of a solid or dotted wedge to depict bonds to other asymmetric carbon atoms in the same compound is meant to indicate that a mixture of diastereomers is present.

 Stereoisomers of Formula I include cis and trans isomers, optical isomers such
30 as R and S enantiomers, diastereomers, geometric isomers, rotational isomers, conformational isomers, and tautomers of the compounds of the invention, including compounds exhibiting more than one type of isomerism; and mixtures thereof (such as racemates and diastereomeric pairs). Also included are acid addition or base addition

salts wherein the counterion is optically active, for example, D-lactate or L-lysine, or racemic, for example, DL-tartrate or DL-arginine.

When any racemate crystallizes, crystals of two different types are possible. The first type is the racemic compound (true racemate) referred to above wherein one homogeneous form of crystal is produced containing both enantiomers in equimolar amounts. The second type is the racemic mixture or conglomerate wherein two forms of crystal are produced in equimolar amounts each comprising a single enantiomer.

The present invention comprises the tautomeric forms of compounds of the invention. Where structural isomers are interconvertible *via* a low energy barrier, tautomeric isomerism ('tautomerism') can occur. This can take the form of proton tautomerism in compounds of the invention containing, for example, an imino, keto, or oxime group, or so-called valence tautomerism in compounds which contain an aromatic moiety. It follows that a single compound may exhibit more than one type of isomerism. The various ratios of the tautomers in solid and liquid form is dependent on the various substituents on the molecule as well as the particular crystallization technique used to isolate a compound.

The compounds of this invention may be used in the form of salts derived from inorganic or organic acids. Depending on the particular compound, a salt of the compound may be advantageous due to one or more of the salt's physical properties, such as enhanced pharmaceutical stability in differing temperatures and humidities, or a desirable solubility in water or oil. In some instances, a salt of a compound also may be used as an aid in the isolation, purification, and/or resolution of the compound.

Where a salt is intended to be administered to a patient (as opposed to, for example, being used in an *in vitro* context), the salt preferably is pharmaceutically acceptable. The term "pharmaceutically acceptable salt" refers to a salt prepared by combining a compound of formula I with an acid whose anion, or a base whose cation, is generally considered suitable for human consumption. Pharmaceutically acceptable salts are particularly useful as products of the methods of the present invention because of their greater aqueous solubility relative to the parent compound. For use in medicine, the salts of the compounds of this invention are non-toxic "pharmaceutically acceptable salts." Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid.

Suitable pharmaceutically acceptable acid addition salts of the compounds of the present invention when possible include those derived from inorganic acids, such as hydrochloric, hydrobromic, hydrofluoric, boric, fluoroboric, phosphoric, metaphosphoric, nitric, carbonic, sulfonic, and sulfuric acids, and organic acids such as acetic, benzenesulfonic, benzoic, citric, ethanesulfonic, fumaric, gluconic, glycolic, isothionic, lactic, lactobionic, maleic, malic, methanesulfonic, trifluoromethanesulfonic, succinic, toluenesulfonic, tartaric, and trifluoroacetic acids. Suitable organic acids generally include, for example, aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic, and sulfonic classes of organic acids.

Specific examples of suitable organic acids include acetate, trifluoroacetate, formate, propionate, succinate, glycolate, gluconate, digluconate, lactate, malate, tartaric acid, citrate, ascorbate, glucuronate, maleate, fumarate, pyruvate, aspartate, glutamate, benzoate, anthranilic acid, stearate, salicylate, *p*-hydroxybenzoate, phenylacetate, mandelate, embonate (pamoate), methanesulfonate, ethanesulfonate, benzenesulfonate, pantothenate, toluenesulfonate, 2-hydroxyethanesulfonate, sufanilate, cyclohexylaminosulfonate, β -hydroxybutyrate, galactarate, galacturonate, adipate, alginate, butyrate, camphorate, camphorsulfonate, cyclopentanepropionate, dodecylsulfate, glycoheptanoate, glycerophosphate, heptanoate, hexanoate, nicotinate, 2-naphthalesulfonate, oxalate, palmoate, pectinate, 3-phenylpropionate, picrate, pivalate, thiocyanate, and undecanoate.

Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, i.e., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts. In another embodiment, base salts are formed from bases which form non-toxic salts, including aluminum, arginine, benzathine, choline, diethylamine, diolamine, glycine, lysine, meglumine, olamine, tromethamine and zinc salts.

Organic salts may be made from secondary, tertiary or quaternary amine salts, such as tromethamine, diethylamine, *N,N'*-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (*N*-methylglucamine), and procaine. Basic nitrogen-containing groups may be quaternized with agents such as lower alkyl (C_1 - C_6) halides (e.g., methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides), dialkyl sulfates (i.e., dimethyl, diethyl, dibutyl, and diamyl sulfates), long

chain halides (e.g., decyl, lauryl, myristyl, and stearyl chlorides, bromides, and iodides), arylalkyl halides (e.g., benzyl and phenethyl bromides), and others.

In one embodiment, hemisalts of acids and bases may also be formed, for example, hemisulfate and hemicalcium salts.

5 Also within the scope of the present invention are so-called "prodrugs" of the compound of the invention. Thus, certain derivatives of the compound of the invention which may have little or no pharmacological activity themselves can, when administered into or onto the body, be converted into the compound of the invention having the desired activity, for example, by hydrolytic cleavage. Such derivatives are referred to as
10 "prodrugs." Further information on the use of prodrugs may be found in "Pro-drugs as Novel Delivery Systems, Vol. 14, ACS Symposium Series (T. Higuchi and V. Stella) and "Bioreversible Carriers in Drug Design," Pergamon Press, 1987 (ed. E. B. Roche, American Pharmaceutical Association). Prodrugs in accordance with the invention can, for example, be produced by replacing appropriate functionalities present in the
15 compounds of any of formula I with certain moieties known to those skilled in the art as "pro-moieties" as described, for example, in "Design of Prodrugs" by H. Bundgaard (Elsevier, 1985).

The present invention also includes isotopically labeled compounds, which are identical to those recited in formula I, but for the fact that one or more atoms are
20 replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the present invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, sulfur, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{11}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds of the present
25 invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically labeled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue
30 distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or

reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labeled compounds of formula I of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples and Preparations below, by substituting a readily available isotopically
5 labeled reagent for a non-isotopically labeled reagent.

DETAILED DESCRIPTION OF THE INVENTION

Typically, a compound of the invention is administered in an amount effective to treat a condition as described herein. The compounds of the invention are administered
10 by any suitable route in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. Therapeutically effective doses of the compounds required to treat the progress of the medical condition are readily ascertained by one of ordinary skill in the art using preclinical and clinical approaches familiar to the medicinal arts.

The term "treating", as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, unless otherwise indicated, refers to the act of treating as "treating" is defined immediately above. The term "treating" also includes adjuvant and
15 neo-adjuvant treatment of a subject.

The compounds of the invention may be administered orally. Oral administration may involve swallowing, so that the compound enters the gastrointestinal tract, or buccal or sublingual administration may be employed, by which the compound enters the blood stream directly from the mouth.

In another embodiment, the compounds of the invention may also be administered directly into the blood stream, into muscle, or into an internal organ. Suitable means for parenteral administration include intravenous, intraarterial, intraperitoneal, intrathecal, intraventricular, intraurethral, intrasternal, intracranial, intramuscular and subcutaneous. Suitable devices for parenteral administration include
25 needle (including microneedle) injectors, needle-free injectors and infusion techniques.

In another embodiment, the compounds of the invention may also be administered topically to the skin or mucosa, that is, dermally or transdermally. In another embodiment, the compounds of the invention can also be administered

intranasally or by inhalation. In another embodiment, the compounds of the invention may be administered rectally or vaginally. In another embodiment, the compounds of the invention may also be administered directly to the eye or ear.

The dosage regimen for the compounds and/or compositions containing the compounds is based on a variety of factors, including the type, age, weight, sex and medical condition of the patient; the severity of the condition; the route of administration; and the activity of the particular compound employed. Thus the dosage regimen may vary widely. Dosage levels of the order from about 0.01 mg to about 100 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions. In one embodiment, the total daily dose of a compound of the invention (administered in single or divided doses) is typically from about 0.01 to about 100 mg/kg. In another embodiment, the total daily dose of the compound of the invention is from about 0.1 to about 50 mg/kg, and in another embodiment, from about 0.5 to about 30 mg/kg (i.e., mg compound of the invention per kg body weight). In one embodiment, dosing is from 0.01 to 10 mg/kg/day. In another embodiment, dosing is from 0.1 to 1.0 mg/kg/day. Dosage unit compositions may contain such amounts or submultiples thereof to make up the daily dose. In many instances, the administration of the compound will be repeated a plurality of times in a day (typically no greater than 4 times). Multiple doses per day typically may be used to increase the total daily dose, if desired.

For oral administration, the compositions may be provided in the form of tablets containing from about 0.01 mg to about 500 mg of the active ingredient, or in another embodiment, from about 1 mg to about 100 mg of active ingredient. Intravenously, doses may range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion.

Suitable subjects according to the present invention include mammalian subjects. Mammals according to the present invention include, but are not limited to, canine, feline, bovine, caprine, equine, ovine, porcine, rodents, lagomorphs, primates, and the like, and encompass mammals in utero. In one embodiment, humans are suitable subjects. Human subjects may be of either gender and at any stage of development.

In another embodiment, the invention comprises the use of one or more compounds of the invention for the preparation of a medicament for the treatment of the conditions recited herein.

For the treatment of the conditions referred to above, the compound of the invention can be administered as compound per se. Alternatively, pharmaceutically acceptable salts are suitable for medical applications because of their greater aqueous solubility relative to the parent compound.

5 In another embodiment, the present invention comprises pharmaceutical compositions. Such pharmaceutical compositions comprise a compound of the invention presented with a pharmaceutically acceptable carrier. The carrier can be a solid, a liquid, or both, and may be formulated with the compound as a unit-dose composition, for example, a tablet, which can contain from 0.05% to 95% by weight of
10 the active compounds. A compound of the invention may be coupled with suitable polymers as targetable drug carriers. Other pharmacologically active substances can also be present.

The compounds of the present invention may be administered by any suitable route, preferably in the form of a pharmaceutical composition adapted to such a route,
15 and in a dose effective for the treatment intended. The active compounds and compositions, for example, may be administered orally, rectally, parenterally, or topically.

Oral administration of a solid dose form may be, for example, presented in discrete units, such as hard or soft capsules, pills, cachets, lozenges, or tablets, each
20 containing a predetermined amount of at least one compound of the present invention. In another embodiment, the oral administration may be in a powder or granule form. In another embodiment, the oral dose form is sub-lingual, such as, for example, a lozenge. In such solid dosage forms, the compounds of formula I are ordinarily combined with one or more adjuvants. Such capsules or tablets may contain a controlled-release
25 formulation. In the case of capsules, tablets, and pills, the dosage forms also may comprise buffering agents or may be prepared with enteric coatings.

In another embodiment, oral administration may be in a liquid dose form. Liquid dosage forms for oral administration include, for example, pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents
30 commonly used in the art (e.g., water). Such compositions also may comprise adjuvants, such as wetting, emulsifying, suspending, flavoring (e.g., sweetening), and/or perfuming agents.

In another embodiment, the present invention comprises a parenteral dose form. "Parenteral administration" includes, for example, subcutaneous injections, intravenous injections, intraperitoneal injections, intramuscular injections, intrasternal injections, and infusion. Injectable preparations (e.g., sterile injectable aqueous or oleaginous suspensions) may be formulated according to the known art using suitable dispersing, wetting agents, and/or suspending agents.

In another embodiment, the present invention comprises a topical dose form. "Topical administration" includes, for example, transdermal administration, such as via transdermal patches or iontophoresis devices, intraocular administration, or intranasal or inhalation administration. Compositions for topical administration also include, for example, topical gels, sprays, ointments, and creams. A topical formulation may include a compound which enhances absorption or penetration of the active ingredient through the skin or other affected areas. When the compounds of this invention are administered by a transdermal device, administration will be accomplished using a patch either of the reservoir and porous membrane type or of a solid matrix variety. Typical formulations for this purpose include gels, hydrogels, lotions, solutions, creams, ointments, dusting powders, dressings, foams, films, skin patches, wafers, implants, sponges, fibers, bandages and microemulsions. Liposomes may also be used. Typical carriers include alcohol, water, mineral oil, liquid petrolatum, white petrolatum, glycerin, polyethylene glycol and propylene glycol. Penetration enhancers may be incorporated; see, for example, J. Pharm. Sci., 88 (10), 955-958, by Finnin and Morgan (October 1999).

Formulations suitable for topical administration to the eye include, for example, eye drops wherein the compound of this invention is dissolved or suspended in a suitable carrier. A typical formulation suitable for ocular or aural administration may be in the form of drops of a micronized suspension or solution in isotonic, pH-adjusted, sterile saline. Other formulations suitable for ocular and aural administration include ointments, biodegradable (e.g., absorbable gel sponges, collagen) and non-biodegradable (e.g., silicone) implants, wafers, lenses and particulate or vesicular systems, such as niosomes or liposomes. A polymer such as cross-linked polyacrylic acid, polyvinyl alcohol, hyaluronic acid, a cellulosic polymer, for example, (hydroxypropyl)methyl cellulose, hydroxyethyl cellulose, or methyl cellulose, or a heteropolysaccharide polymer, for example, gelatin gum, may be incorporated together

with a preservative, such as benzalkonium chloride. Such formulations may also be delivered by iontophoresis.

For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant. Formulations suitable for intranasal administration are typically administered in the form of a dry powder (either alone, as a mixture, for example, in a dry blend with lactose, or as a mixed component particle, for example, mixed with phospholipids, such as phosphatidylcholine) from a dry powder inhaler or as an aerosol spray from a pressurized container, pump, spray, atomizer (preferably an atomizer using electrohydrodynamics to produce a fine mist), or nebulizer, with or without the use of a suitable propellant, such as 1,1,1,2-tetrafluoroethane or 1,1,1,2,3,3,3-heptafluoropropane. For intranasal use, the powder may comprise a bioadhesive agent, for example, chitosan or cyclodextrin.

In another embodiment, the present invention comprises a rectal dose form. Such rectal dose form may be in the form of, for example, a suppository. Cocoa butter is a traditional suppository base, but various alternatives may be used as appropriate.

Other carrier materials and modes of administration known in the pharmaceutical art may also be used. Pharmaceutical compositions of the invention may be prepared by any of the well-known techniques of pharmacy, such as effective formulation and administration procedures. The above considerations in regard to effective formulations and administration procedures are well known in the art and are described in standard textbooks. Formulation of drugs is discussed in, for example, Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania, 1975; Liberman *et al.*, Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Kibbe *et al.*, Eds., Handbook of Pharmaceutical Excipients (3rd Ed.), American Pharmaceutical Association, Washington, 1999.

The compounds of the present invention can be used, alone or in combination with other therapeutic agents, in the treatment of various conditions or disease states. The compound(s) of the present invention and other therapeutic agent(s) may be administered simultaneously (either in the same dosage form or in separate dosage forms) or sequentially.

Two or more compounds may be administered simultaneously, concurrently or sequentially. Additionally, simultaneous administration may be carried out by mixing the compounds prior to administration or by administering the compounds at the same point in time but at different anatomic sites or using different routes of administration.

5 The phrases “concurrent administration,” “co-administration,” “simultaneous administration,” and “administered simultaneously” mean that the compounds are administered in combination.

 The present invention includes the use of a combination of a LRRK2 inhibitor compound as provided in formula I and one or more additional pharmaceutically active agent(s). If a combination of active agents is administered, then they may be administered sequentially or simultaneously, in separate dosage forms or combined in a single dosage form. Accordingly, the present invention also includes pharmaceutical compositions comprising an amount of: (a) a first agent comprising a compound of formula I or a pharmaceutically acceptable salt of the compound; (b) a second
10 pharmaceutically active agent; and (c) a pharmaceutically acceptable carrier, vehicle or diluent.

 Various pharmaceutically active agents may be selected for use in conjunction with the compounds of Formula I, depending on the disease, disorder, or condition to be treated. For example, a pharmaceutical composition for use in treating Parkinson's
20 disease may comprise a compound of formula I or a pharmaceutically acceptable salt thereof together with another agent such as a dopamine (levodopa, either alone or with a DOPA decarboxylase inhibitor), a monoamine oxidase (MAO) inhibitor, a catechol O-methyltransferase (COMT) inhibitor or an anticholinergic agent, or any combination thereof. Particularly preferred agents to combine with the compounds of formula I for
25 use in treating Parkinson's disease include levodopa, carbidopa, tolcapone, entacapone, selegiline, benzotropine and trihexyphenidyl, or any combination thereof. Pharmaceutically active agents that may be used in combination with the compounds of formula I and compositions thereof include, without limitation:

(i) levodopa (or its methyl or ethyl ester), alone or in combination with a DOPA
30 decarboxylase inhibitor (e.g., carbidopa (SINEMET, CARBILEV, PARCOPA), benserazide (MADOPAR), α -methyldopa, monofluoromethyldopa, difluoromethyldopa, brocresine, or *m*-hydroxybenzylhydrazine);

- (ii) anticholinergics, such as amitriptyline (ELAVIL, ENDEP), butriptyline, benzotropine mesylate (COGENTIN), trihexyphenidyl (ARTANE), diphenhydramine (BENADRYL), orphenadrine (NORFLEX), hyoscyamine, atropine (ATROPEN), scopolamine (TRANSDERM-SCOP), scopolamine methylbromide (PARMINE), dicycloverine (BENTYL, BYCLOMINE, DIBENT, DILOMINE), tolterodine (DETROL), oxybutynin (DITROPAN, LYRINEL XL, OXYTROL), penthienate bromide, propantheline (PRO-BANTHINE), cyclizine, imipramine hydrochloride (TOFRANIL), imipramine maleate (SURMONTIL), lofepramine, desipramine (NORPRAMIN), doxepin (SINEQUAN, ZONALON), trimipramine (SURMONTIL), and glycopyrrolate (ROBINUL);
- (iii) catechol O-methyltransferase (COMT) inhibitors, such as nitecapone, tolcapone (TASMAR), entacapone (COMTAN), and tropolone;
- (iv) monoamine oxidase (MAO) inhibitors, such as selegiline (EMSAM), selegiline hydrochloride (l-deprenyl, ELDEPRYL, ZELAPAR), dimethylselegiline, brofaromine, phenelzine (NARDIL), tranlycypromine (PARNATE), moclobemide (AURORIX, MANERIX), biefloxatone, safinamide, isocarboxazid (MARPLAN), nialamide (NIAMID), rasagiline (AZILECT), iproniazide (MARSILID, IPROZID, IPRONID), iproclozide, toloxatone (HUMORYL, PERENUM), bifemelane, desoxypeganine, harmine (also known as telepathine or banasterine), harmaline, linezolid (ZYVOX, ZYVOXID), and pargyline (EUDATIN, SUPIRDYL);
- (v) acetylcholinesterase inhibitors, such as donepezil hydrochloride (ARICEPT®, MEMAC), physostigmine salicylate (ANTILIRIUM®), physostigmine sulfate (ESERINE), ganstigmine, rivastigmine (EXELON®), ladostigil, NP-0361, galantamine hydrobromide (RAZADYNE®, REMINYL®, NIVALIN®), tacrine (COGNEX®), tolserine, memoquin, huperzine A (HUP-A; Neuro-Hitech), phenserine, bisnorcymserine (also known as BNC), and INM-176;
- (vi) amyloid- β (or fragments thereof), such as A β ₁₋₁₅ conjugated to pan HLA DR-binding epitope (PADRE®), ACC-001 (Elan/Wyeth), and Affitope;
- (vii) antibodies to amyloid- β (or fragments thereof), such as ponezumab, solanezumab, bapineuzumab (also known as AAB-001), AAB-002 (Wyeth/Elan), Gantenerumab, intravenous Ig (GAMMAGARD®), LY2062430 (humanized m266; Lilly), and those disclosed in International Patent Publication Nos WO04/032868, WO05/025616, WO06/036291, WO06/069081, WO06/118959, in

US Patent Publication Nos US2003/0073655, US2004/0192898, US2005/0048049, US2005/0019328, in European Patent Publication Nos EP0994728 and 1257584, and in US Patent No 5,750,349;

- (viii) amyloid-lowering or -inhibiting agents (including those that reduce amyloid production, accumulation and fibrillization) such as eprodinate, celecoxib, lovastatin, anapsos, colostrinin, pioglitazone, clioquinol (also known as PBT1), PBT2 (Prana Biotechnology), flurbiprofen (ANSAID®, FROBEN®) and its *R*-enantiomer tarenflurbil (FLURIZAN®), nitroflurbiprofen, fenoprofen (FENOPRON, NALFON®), ibuprofen (ADVIL®, MOTRIN®, NUROFEN®), ibuprofen lysinate, meclofenamic acid, meclofenamate sodium (MECLOMEN®), indomethacin (INDOCIN®), diclofenac sodium (VOLTAREN®), diclofenac potassium, sulindac (CLINORIL®), sulindac sulfide, diflunisal (DOLOBID®), naproxen (NAPROSYN®), naproxen sodium (ANAPROX®, ALEVE®), insulin-degrading enzyme (also known as insulysin), the ginkgo biloba extract EGb-761 (ROKAN®, TEBONIN®), tramiprosate (CEREBRIL®, ALZHEMED®), KIACTA®, neprilysin (also known as neutral endopeptidase (NEP)), scyllo-inositol (also known as scyllitol), atorvastatin (LIPITOR®), simvastatin (ZOCOR®), ibutamoren mesylate, BACE inhibitors such as LY450139 (Lilly), BMS-782450, GSK-188909; gamma secretase modulators and inhibitors such as ELND-007, BMS-708163 (Avagacestat), and DSP8658 (Dainippon); and RAGE (receptor for advanced glycation end-products) inhibitors, such as TTP488 (Transtech) and TTP4000 (Transtech), and those disclosed in US Patent No 7,285,293, including PTI-777;
- (ix) alpha-adrenergic receptor agonists, and beta-adrenergic receptor blocking agents (beta blockers); anticholinergics; anticonvulsants; antipsychotics; calcium channel blockers; catechol O-methyltransferase (COMT) inhibitors; central nervous system stimulants; corticosteroids; dopamine receptor agonists and antagonists; dopamine reuptake inhibitors; gamma-aminobutyric acid (GABA) receptor agonists; immunosuppressants; interferons; muscarinic receptor agonists; neuroprotective drugs; nicotinic receptor agonists; norepinephrine (noradrenaline) reuptake inhibitors; quinolines; and trophic factors;
- (x) histamine 3 (H3) antagonists, such as PF-3654746 and those disclosed in US Patent Publication Nos US2005-0043354, US2005-0267095, US2005-0256135, US2008-0096955, US2007-1079175, and US2008-0176925; International Patent

Publication Nos WO2006/136924, WO2007/063385, WO2007/069053, WO2007/088450, WO2007/099423, WO2007/105053, WO2007/138431, and WO2007/088462; and US Patent No 7,115,600);

- (xi) *N*-methyl-D-aspartate (NMDA) receptor antagonists, such as memantine (NAMENDA, AXURA, EBIXA), amantadine (SYMMETREL), acamprosate (CAMPRAL), besonprodil, ketamine (KETALAR), delucemine, dexanabinol, dexefaroxan, dextromethorphan, dextrorphan, traxoprodil, CP-283097, himantane, idantadol, ipenoxazone, L-701252 (Merck), lancicemine, levorphanol (DROMORAN), methadone, (DOLOPHINE), neramexane, perzinfotel, phencyclidine, tianeptine (STABLON), dizocilpine (also known as MK-801), ibogaine, voacangine, tiletamine, riluzole (RILUTEK), aptiganel (CERESTAT), gavestinel, and remacimide;
- (xii) phosphodiesterase (PDE) inhibitors, including (a) PDE1 inhibitors; (b) PDE2 inhibitors; (c) PDE3 inhibitors; (d) PDE4 inhibitors; (e) PDE5 inhibitors; (f) PDE9 inhibitors (e.g., PF-04447943, BAY 73-6691 (Bayer AG) and those disclosed in US Patent Publication Nos US2003/0195205, US2004/0220186, US2006/0111372, US2006/0106035, and USSN 12/118,062 (filed May 9, 2008)); and (g) PDE10 inhibitors such as 2-({4-[1-methyl-4-(pyridin-4-yl)-1*H*-pyrazol-3-yl]phenoxy}methyl)quinoline (PF-2545920);
- (xiii) serotonin (5-hydroxytryptamine) 1A (5-HT_{1A}) receptor antagonists, such as spiperone, *levo*-pindolol, lecozotan;
- (xiv) serotonin (5-hydroxytryptamine) 2C (5-HT_{2c}) receptor agonists, such as vabicaserin, and zicronapine; serotonin (5-hydroxytryptamine) 4 (5-HT₄) receptor agonists/antagonists, such as PRX-03140 (Epix) and PF-04995274;
- (xv) serotonin (5-hydroxytryptamine) 3C (5-HT_{3c}) receptor antagonists, such as Ondansetron (Zofran);
- (xvi) serotonin (5-hydroxytryptamine) 6 (5-HT₆) receptor antagonists, such as mianserin (TOLVON, BOLVIDON, NORVAL), methiothepin (also known as metitepine), ritanserin, SB-271046, SB-742457 (GlaxoSmithKline), Lu AE58054 (Lundbeck A/S), SAM-760, and PRX-07034 (Epix);
- (xvii) serotonin (5-HT) reuptake inhibitors such as alaproclate, citalopram (CELEXA, CIPRAMIL), escitalopram (LEXAPRO, CIPRALEX), clomipramine (ANAFRANIL), duloxetine (CYMBALTA), femoxetine (MALEXIL), fenfluramine (PONDIMIN),

- norfenfluramine, fluoxetine (PROZAC), fluvoxamine (LUVOX), indalpine, milnacipran (IXEL), paroxetine (PAXIL, SEROXAT), sertraline (ZOLOFT, LUSTRAL), trazodone (DESYREL, MOLIPAXIN), venlafaxine (EFFEXOR), zimelidine (NORMUD, ZELMID), bicifadine, desvenlafaxine (PRISTIQ),
5 brasofensine, vilazodone, cariprazine and tesofensine;
- (xviii) Glycine transporter-1 inhibitors such as paliflutine, ORG-25935, and ORG-26041; and mGluR modulators such as AFQ-059 and amantidine;
- (xix) AMPA-type glutamate receptor modulators such as perampanel, mibampator, selurampanel, GSK-729327, and *N*-{(3*S*,4*S*)-4-[4-(5-cyanothiophen-2-yl)phenoxy]tetrahydrofuran-3-yl}propane-2-sulfonamide;
10
- (xx) P450 inhibitors, such as ritonavir;
- (xxi) tau therapy targets, such as davunetide; and the like.

15 The present invention further comprises kits that are suitable for use in performing the methods of treatment described above. In one embodiment, the kit contains a first dosage form comprising one or more of the compounds of the present invention and a container for the dosage, in quantities sufficient to carry out the methods of the present invention.

20 In another embodiment, the kit of the present invention comprises one or more compounds of the invention.

General Synthetic Schemes

The compounds of Formula I may be prepared by the methods described below,
25 together with synthetic methods known in the art of organic chemistry, or modifications and transformations that are familiar to those of ordinary skill in the art. The starting materials used herein are commercially available or may be prepared by routine methods known in the art [such as those methods disclosed in standard reference books such as the *Compendium of Organic Synthetic Methods*, Vol. I-XII (published by
30 Wiley-Interscience)]. Preferred methods include, but are not limited to, those described below.

During any of the following synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned.

This can be achieved by means of conventional protecting groups, such as those described in T. W. Greene, Protective Groups in Organic Chemistry, John Wiley & Sons, 1981; T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Chemistry, John Wiley & Sons, 1991; and T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Chemistry, John Wiley & Sons, 1999, which are hereby incorporated by reference.

Compounds of Formula I, or their pharmaceutically acceptable salts, can be prepared according to the reaction Schemes discussed herein below. Unless otherwise indicated, the substituents in the Schemes are defined as above. Isolation and purification of the products is accomplished by standard procedures, which are known to a chemist of ordinary skill.

One skilled in the art will recognize that in many cases, the compounds in Schemes 1 through 4 will be generated as a mixture of diastereomers and/or enantiomers; these may be separated at various stages of the synthetic schemes using conventional techniques or a combination of such techniques, such as, but not limited to, crystallization, normal-phase chromatography, reversed phase chromatography and chiral chromatography, to afford the single enantiomers of the invention.

It will be understood by one skilled in the art that the various symbols, superscripts and subscripts used in the schemes, methods and examples are used for convenience of representation and/or to reflect the order in which they are introduced in the schemes, and are not intended to necessarily correspond to the symbols, superscripts or subscripts in the appended claims. The schemes are representative of methods useful in synthesizing the compounds of the present invention. They are not to constrain the scope of the invention in any way.

The reactions for preparing compounds of the invention can be carried out in suitable solvents, which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, e.g., temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.

Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ^1H or ^{13}C), infrared spectroscopy, spectrophotometry (e.g., UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC) or thin layer chromatography (TLC).

DETAILED DESCRIPTION OF THE INVENTION

Compounds of Formula I and intermediates thereof may be prepared according to the following reaction schemes and accompanying discussion. Unless otherwise indicated, R^1 , R^2 , R^3 , R^4 and R^5 in the reaction schemes and discussions that follow are as defined hereinabove. In general, the compounds of this invention may be made by processes which include processes analogous to those known in the chemical arts, particularly in light of the description contained herein. Certain processes for the manufacture of the compounds of this invention and intermediates thereof are provided as further features of the invention and are illustrated by the following reaction schemes. Other processes may be described in the experimental section. The schemes and examples provided herein (including the corresponding description) are for illustration only, and not intended to limit the scope of the present invention.


Scheme 1 refers to preparation of compounds of Formula I. Referring to Scheme 1, compounds of Formula 1-1 and 1-2 [wherein X is a leaving group such as Br or I, and Pg is a suitable protecting group, such as 2-(trimethylsilyl)ethoxymethyl (SEM), *p*-toluenesulfonyl (tosyl) or *tert*-butoxycarbonyl (BOC)] are commercially available or can be made by methods described herein or other methods well known to those skilled in the art.

A compound of Formula 1-3 can be prepared by coupling a compound of Formula 1-1 with a compound of Formula 1-2, for example, by heating a mixture of a compound of Formula 1-1 with a compound of Formula 1-2 in the presence of a base, such as *N,N*-diisopropylethylamine, in an appropriate solvent, such as *n*-butanol, at temperatures ranging between 50 °C and 200 °C. Suitable reaction times are typically from 20 minutes to 48 hours. Alternatively, a metal-catalyzed (such as using a palladium or copper catalyst) coupling may be employed to accomplish the aforesaid coupling. In this variant of the coupling, a mixture of a compound of Formula 1-1 and a

compound of Formula 1-2 can be heated at temperatures ranging between 50 °C and 120 °C in the presence of a base [such as cesium carbonate], a metal catalyst [such as a palladium catalyst, e.g., palladium(II) acetate], and a ligand [such as 1,1'-binaphthalene-2,2'-diylbis(diphenylphosphane) (BINAP)] in an appropriate solvent, such as 1,4-dioxane. Suitable reaction times are typically from 30 minutes to 48 hours.


A compound of Formula 1-3 can subsequently be reacted with a compound of Formula R^3-M [wherein M can be $B(OH)_2$; $B(OR)_2$ wherein each R is independently H or C_{1-6} alkyl, or wherein two (OR) groups, together with the B atom to which they are attached, form a 5- to 10-membered heterocyclic ring optionally substituted with one or more C_{1-6} alkyl; a trialkyltin moiety; or the like] by a metal-catalyzed (such as using a palladium catalyst) coupling reaction to obtain a compound of Formula 1-4. Compounds of Formula R^3-M are commercially available or can be prepared by methods analogous to those described in the chemical art. Alternatively, a compound of Formula 1-3 can be converted to a compound of Formula 1-5 [wherein M is defined as above]. A compound of Formula 1-5 can then be reacted with a compound of Formula R^3-X [wherein X is defined as above] by a metal-catalyzed (such as using a palladium catalyst) coupling reaction to obtain a compound of Formula I. Compounds of Formula R^3-X are commercially available or can be prepared by methods analogous to those described in the chemical art. The type of reaction employed depends on the selection of X and M. For example, when X is halogen or triflate and the R^3-M reagent is a boronic acid or boronic ester, a Suzuki reaction may be used [A. Suzuki, *J. Organomet. Chem.* **1999**, 576, 147-168; N. Miyaura and A. Suzuki, *Chem. Rev.* **1995**, 95, 2457-2483; A. F. Littke et al., *J. Am. Chem. Soc.* **2000**, 122, 4020-4028]. Alternatively, when X is halogen or triflate and M is trialkyltin, a Stille coupling may be employed [V. Farina et al., *Organic Reactions* **1997**, 50, 1-652]. Where X is Br, I or triflate and M is Zn or Mg, a Negishi coupling or Kumada coupling may be used [E. Erdik, *Tetrahedron* **1992**, 48, 9577-9648; T. Banno et al., *J. Organomet. Chem.* **2002**, 653, 288-291]. Removal of the protecting group from compounds of Formula 1-4 under conditions well known to those skilled in the art affords compounds of Formula I.

Scheme 1



Scheme 2 also refers to preparation of compounds of Formula I. Referring to Scheme 2, compounds of Formula I may be prepared utilizing analogous chemical transformations to those described in Scheme 1, but with a different ordering of steps. A compound of Formula 1-1 (as in Scheme 1) can be converted to a compound of Formula 2-1 either directly or after conversion to a compound of Formula 2-2 using methods analogous to those described in Scheme 1. A compound of Formula 2-1 may then be coupled to a compound of Formula 1-2 as in Scheme 1, to produce a compound of Formula 1-4. The coupling conditions employed may be analogous to those described for the preparation of a compound of Formula 1-3 in Scheme 1.


Scheme 2



Scheme 3 refers to a preparation of a compound of Formula 1-1. Referring to Scheme 3, compounds of Formula 3-1 are commercially available or can be made by methods described herein or other methods well known to those skilled in the art. A
5 compound of Formula 3-1 can be treated with a strong base and the intermediate can be subsequently reacted with an electrophile to obtain a compound of Formula 1-1. Examples of suitable reaction conditions for the reaction include mixing a compound of Formula 3-1 with a suitable base, such as lithium diisopropylamide, in a suitable reaction solvent such as tetrahydrofuran. This is followed by addition of an electrophile
10 such as an alkyl iodide or bromide. Suitable temperatures for the aforesaid reaction are typically between -78°C and 30°C . Suitable reaction times typically are from 20 minutes to 48 hours. A compound of Formula 1-1 can be converted to a compound of Formula I using chemistry described in Schemes 1 and 2.

15

Scheme 3



5 Scheme 4 refers to a preparation of a compound of Formula 4-5, wherein X is a leaving group such as Br or I. Referring to Scheme 4, compounds of Formula 4-1 (wherein each R is independently H or C₁₋₆ alkyl) and Formula 4-2 are commercially available or can be made by methods described herein or other methods well known to those skilled in the art.


10 A compound of Formula 4-3 can be prepared by condensation of a compound of Formula 4-1 with a compound of Formula 4-2, for example, through heating a compound of Formula 4-1 and a compound of Formula 4-2 in the presence of a base, such as sodium methoxide, in an appropriate solvent, such as ethanol, at temperatures ranging between 50 °C and 100 °C. Suitable reaction times are typically between 1 and

15 24 hours. A compound of Formula 4-3 can be converted to a compound of Formula 4-4 by treatment with a chlorinating agent, such as phosphorus oxychloride, either in the presence of a suitable solvent or neat, at a temperature ranging between 70 °C and 120 °C and a reaction time of between 1 and 24 hours. A compound of Formula 4-4 may then be treated with an appropriate halogenating agent, such as *N*-iodosuccinimide, in a

20 suitable solvent, such as dichloromethane, to produce a compound of Formula 4-5. Suitable reaction temperatures range from 0 °C to 50 °C, and suitable reaction times are typically from 30 minutes to 24 hours. A compound of Formula 4-5 can be protected (i.e. the pyrrole ring nitrogen is protected) by methods well known in the art and then the corresponding protected compound can be converted to a compound of Formula I using

25 chemistry described in Schemes 1, 2 and 3.

Scheme 4



Experimental Procedures and Working Examples

The following illustrate the synthesis of various compounds of the present invention. Additional compounds within the scope of this invention may be prepared using the methods illustrated in these Examples, either alone or in combination with techniques generally known in the art.


Experiments were generally carried out under inert atmosphere (nitrogen or argon), particularly in cases where oxygen- or moisture-sensitive reagents or intermediates were employed. Commercial solvents and reagents were generally used without further purification, including anhydrous solvents where appropriate. In some cases, Sure-SealTM products from the Aldrich Chemical Company, Milwaukee, Wisconsin were used. In other cases, commercial solvents were passed through columns packed with 4Å molecular sieves, until the following QC standards for water were attained: a) <100 ppm for dichloromethane, toluene, *N,N*-dimethylformamide and tetrahydrofuran; b) <180 ppm for methanol, ethanol, 1,4-dioxane and diisopropylamine. For very sensitive reactions, solvents were further treated with metallic sodium, calcium hydride or molecular sieves, and distilled just prior to use.

Products were generally dried under vacuum before being carried on to further reactions or submitted for biological testing. Mass spectrometry data is reported from either liquid chromatography-mass spectrometry (LCMS), atmospheric pressure chemical ionization (APCI) or gas chromatography-mass spectrometry (GCMS) instrumentation. Chemical shifts for nuclear magnetic resonance (NMR) data are expressed in parts per million (ppm, δ) referenced to residual peaks from the deuterated solvents employed.

For syntheses referencing procedures in other Examples or Methods, reaction conditions (length of reaction and temperature) may vary. In general, reactions were followed by thin layer chromatography or mass spectrometry, and subjected to work-up when appropriate. Purifications may vary between experiments: in general, solvents and the solvent ratios used for eluents/gradients were chosen to provide appropriate *R_f*s or retention times.

Preparation P1

5-Iodo-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine
(**P1**)



- 5 Step 1. Synthesis of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C1**).

A solution of 4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine (9.8 g, 35 mmol) in tetrahydrofuran (250 mL) was cooled to 0 °C and treated with sodium hydride (60% in oil, 1.54 g, 38.5 mmol) in three portions. After the reaction mixture had stirred at 0 °C for
10 1 hour, 2-(trimethylsilyl)ethoxymethyl chloride (6.4 g, 38 mmol) was added drop-wise, and the reaction mixture was warmed to room temperature and allowed to stir for 3 hours. The reaction was quenched with saturated aqueous sodium chloride solution (250 mL), and the organic layer was dried over sodium sulfate, filtered, and concentrated *in vacuo*. Silica gel chromatography (Eluent: 10:1 petroleum ether / ethyl
15 acetate) afforded the product as a white solid. Yield: 8 g, 20 mmol, 57%. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.69 (s, 1H), 8.14 (s, 1H), 5.60 (s, 2H), 3.51 (t, *J*=8 Hz, 2H), 0.82 (t, *J*=8 Hz, 2H), -0.10 (s, 9H).


Step 2. Synthesis of 5-iodo-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**P1**).

20 Morpholine (2.45 g, 28.1 mmol) and *N,N*-diisopropylethylamine (6.63 g, 51.3 mmol) were added to a solution of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C1**) (10.5 g, 25.6 mmol) in *n*-butanol (300 mL), and the reaction mixture was heated at reflux for 18 hours, then concentrated under reduced pressure. Aqueous hydrochloric acid (0.1 M, 100 mL) was added and the resulting solid
25 was collected by filtration, washed with water (20 mL) and dried under vacuum to provide the product as a yellow solid. Yield: 8.0 g, 17 mmol, 66%. LCMS *m/z* 461.2


[M+H⁺]. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.39 (s, 1H), 7.81 (s, 1H), 5.52 (s, 2H), 3.80-3.86 (m, 4H), 3.46-3.53 (m, 6H), 0.77-0.84 (m, 2H), -0.10 (s, 9H).

Preparation P2

5 4-(Morpholin-4-yl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P2**)



To a solution of 5-iodo-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P1**) (500 mg, 1.09 mmol) and 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (543 mg, 4.24 mmol) in 1,4-dioxane (20 mL) was added
 10 tris(dibenzylideneacetone)dipalladium(0) (99.7 mg, 0.109 mmol), triethylamine (439 mg, 4.34 mmol) and 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (XPhos, 51.8 mg, 0.109 mmol), and the reaction mixture was heated at 95 °C for 18 hours. After cooling to room temperature, the reaction mixture was diluted with water (100 mL) and extracted
 15 with ethyl acetate (3 x 50 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo*. Purification via silica gel chromatography (Gradient: 0% to 30% ethyl acetate in petroleum ether) afforded the product as a yellow oil. Yield: 415 mg, 0.901 mmol, 83%. LCMS *m/z* 461.3 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃) δ 8.45
 20 (s, 1H), 7.73 (s, 1H), 5.59 (s, 2H), 3.87-3.93 (m, 4H), 3.68-3.74 (m, 4H), 3.49-3.56 (m, 2H), 1.35 (s, 12H), 0.87-0.93 (m, 2H), -0.06 (s, 9H).


Preparation P35-Bromo-4-chloro-7-[(4-methylphenyl)sulfonyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P3**)

4-Methylbenzenesulfonyl chloride (5.21 g, 27.3 mmol) was added to a suspension of 5-bromo-4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidine (6.34 g, 27.3 mmol) in acetone (70 mL). After addition of an aqueous solution of sodium hydroxide (1.11 g, 27.8 mmol in 15 mL water), the reaction mixture was stirred at room temperature for 16 hours. The mixture was then diluted with water (125 mL), stirred for 10 minutes, and filtered. The collected solids were washed with water to afford the product as a solid.

Yield: 8.88 g, 23.0 mmol, 84%. LCMS m/z 385.9, 387.9, 389.8 $[M+H]^+$. ^1H NMR (400 MHz, DMSO- d_6) δ 8.84 (s, 1H), 8.42 (s, 1H), 8.06 (br d, $J=8.4$ Hz, 2H), 7.46-7.51 (m, 2H), 2.37 (br s, 3H).

Preparation P4

7-[(4-Methylphenyl)sulfonyl]-4-(3-methylpiperidin-1-yl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P4**)




5-Bromo-7-[(4-methylphenyl)sulfonyl]-4-(3-methylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (racemic version of **C15**, see Example 7) (1.44 g, 3.20 mmol) was added to a solution of triethylamine (1.3 g, 13 mmol) in 1,4-dioxane (15 mL). 4,4,5,5-Tetramethyl-1,3,2-dioxaborolane (1.64 g, 12.8 mmol) was then slowly added, and the mixture was

degassed by bubbling nitrogen through it for 5 minutes. 2-Dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl (XPhos, 137 mg, 0.287 mmol) and tris(dibenzylideneacetone)dipalladium(0) (146 mg, 0.160 mmol) were then introduced, and the reaction mixture was heated at 95 °C for 18 hours. After cooling, the mixture was filtered through cotton and the filtrate was concentrated *in vacuo*. Chromatography on silica gel (Gradient: 0% to 75% ethyl acetate in heptane) afforded the product as a solid. Yield: 1.25 g, 2.52 mmol, 79%. ¹H NMR (400 MHz, CDCl₃) δ 8.41 (s, 1H), 8.10 (br d, *J*=8.4 Hz, 2H), 7.95 (s, 1H), 7.30 (br d, *J*=8.6 Hz, 2H), 4.28 (br d, *J*=13 Hz, 1H), 4.18-4.24 (m, 1H), 2.99 (ddd, *J*=13, 11, 4 Hz, 1H), 2.70 (dd, *J*=12.9, 10.9 Hz, 1H), 2.40 (s, 3H), 1.77-1.85 (m, 1H), 1.56-1.70 (m, 4H), 1.35 (s, 12H), 0.88 (d, *J*=6.6 Hz, 3H).

Preparation P5

5-Iodo-4-[(3*S*)-3-methylpiperidin-1-yl]-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P5**)




15

4-Chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C1**) was converted to the product using the method described for synthesis of 5-bromo-7-[(4-methylphenyl)sulfonyl]-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C15**) in Example 7. Yield: 7.38 g, 15.6 mmol, 80%. LCMS *m/z* 473.1 [*M*+*H*⁺]. ¹H NMR (400 MHz, CDCl₃) δ 8.42 (s, 1H), 7.34 (s, 1H), 5.55 (s, 2H), 4.06-4.17 (m, 2H), 3.50-3.57 (m, 2H), 2.91 (ddd, *J*=12.6, 11.5, 3.5 Hz, 1H), 2.58 (dd, *J*=12.6, 10.8 Hz, 1H), 1.95-2.07 (m, 1H), 1.76-1.95 (m, 3H), 1.11-1.23 (m, 1H), 0.96 (d, *J*=6.8 Hz, 3H), 0.88-0.94 (m, 2H), -0.04 (s, 9H).

25

Preparation P6

4-Chloro-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P6**)

Step 1. Synthesis of 4-chloro-5-(1-methyl-1*H*-pyrazol-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C2**).

To a solution of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C1**) (4.1 g, 10 mmol), 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (2.1 g, 10 mmol) and potassium carbonate (2.8 g, 20 mmol) in aqueous 1,4-dioxane was added dichlorobis(triphenylphosphine)palladium(II) (350 mg, 0.50 mmol). The reaction mixture was degassed and purged with nitrogen; this procedure was carried out a total of three times. After heating at reflux for 18 hours, the reaction mixture was cooled to room temperature, diluted with water (200 mL) and extracted with ethyl acetate (3 x 200 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (500 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo*. Purification by chromatography on silica gel (Gradient: 0% to 10% ethyl acetate in petroleum ether) afforded the product as a white solid. Yield: 2.0 g, 5.5 mmol, 55%. ¹H NMR (400 MHz, CDCl₃) δ 8.66 (s, 1H), 7.68 (d, *J*=0.5 Hz, 1H), 7.62 (br s, 1H), 7.36 (s, 1H), 5.68 (s, 2H), 3.99 (s, 3H), 3.54-3.59 (m, 2H), 0.91-0.97 (m, 2H), -0.03 (s, 9H).


Step 2. Synthesis of 4-chloro-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P6**).

4-Chloro-5-(1-methyl-1*H*-pyrazol-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C2**) was converted to the product using the method described for synthesis of 3-[6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**2**) in Example 2. In this case the product, obtained as a yellow solid, was purified by recrystallization from ethyl acetate, rather than via preparative HPLC. Yield: 1.0 g, 4.3 mmol, 43% over two steps. LCMS *m/z* 234.0 [M+H⁺]. ¹H NMR (400 MHz,

DMSO- d_6) δ 12.66 (br s, 1H), 8.58 (s, 1H), 7.91 (s, 1H), 7.69 (d, $J=2.5$ Hz, 1H), 7.63 (d, $J=0.7$ Hz, 1H), 3.89 (s, 3H).

Preparation P7


- 5 *N,N*-Dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine (**P7**)



- 5-iodo-*N,N*-dimethyl-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine [**C3**, prepared from 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C1**) according to the method described for synthesis of 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**5**) in Example 5] was converted to the product according to the method described for synthesis of 4-(morpholin-4-yl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P2**) in Preparation
- 15 **P2**. The product was obtained as a yellow solid. Yield: 342 mg, 0.817 mmol, 69%. LCMS m/z 419.3 [$M+H^+$]. ^1H NMR (400 MHz, $CDCl_3$) δ 8.37 (s, 1H), 7.57 (s, 1H), 5.56 (s, 2H), 3.49-3.57 (m, 2H), 3.25 (s, 6H), 1.36 (s, 12H), 0.86-0.94 (m, 2H), -0.06 (s, 9H).

Example 1

- 20 5-(1-Methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**1**)



Step 1. Synthesis of 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C4**).

To a solution of 5-iodo-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P1**) (500 mg, 1.1 mmol) and 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (272 mg, 1.31 mmol) in a mixture of ethanol and water (4:1, 10 mL) were added dichlorobis(triphenylphosphine)palladium(II) (41 mg, 58 μ mol) and potassium carbonate (447 mg, 3.23 mmol). The reaction mixture was degassed and purged with nitrogen; this procedure was carried out a total of three times. It was then heated at 100 °C for 18 hours. After concentration *in vacuo*, the residue was purified via chromatography on silica gel (Eluent: 1:1 ethyl acetate / petroleum ether) to provide the product as a yellow solid. Yield: 200 mg, 0.48 mmol, 44%. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.39 (s, 1H), 7.86 (s, 1H), 7.59 (s, 1H), 7.52 (s, 1H), 5.57 (s, 2H), 3.90 (s, 3H), 3.50-3.58 (m, 6H), 3.20-3.27 (m, 4H), 0.83 (dd, *J*=8.0, 7.9 Hz, 2H), -0.09 (s, 9H).

Step 2. Synthesis of [5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl]methanol (**C5**).

A solution of 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C4**) (200 mg, 0.48 mmol) in trifluoroacetic acid (5 mL) was stirred at room temperature for 2 hours. The reaction

mixture was concentrated under reduced pressure to afford the product as a yellow oil, which was used for the next step without additional purification.


Step 3. Synthesis of 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**1**).


- 5 A solution of [5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl]methanol (**C5**) (material from the previous step, ≤ 0.48 mmol) in methanol (5 mL) was brought to a pH of >12 via addition of solid potassium carbonate. The reaction mixture was stirred for 30 minutes, filtered, and concentrated *in vacuo*. Purification via preparative HPLC (Column: Agella Venusil ASB C18, 5 μ m; Mobile
- 10 phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile; Eluent: 13% B) provided the product as a yellow solid. Yield over two steps: 90 mg, 0.32 mmol, 67%. LCMS m/z 285.1 $[M+H^+]$. ^1H NMR (400 MHz, DMSO- d_6) δ 12.96 (br s, 1H), 8.47 (s, 1H), 7.85 (s, 1H), 7.57 (s, 1H), 7.52 (d, $J=2.5$ Hz, 1H), 3.90 (s, 3H), 3.53-3.59 (m, 4H), 3.45-3.51 (m, 4H).

15

Example 2

3-[6-Methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**2**)





Step 1. Synthesis of 4-chloro-5-iodo-6-methyl-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidine (**C6**).

To a $-78\text{ }^{\circ}\text{C}$ solution of 4-chloro-5-iodo-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidine (**C1**) (15.0 g, 36.6 mmol) in tetrahydrofuran (500 mL) was added lithium diisopropylamide (2 M solution in heptane / tetrahydrofuran / ethylbenzene, 183 mL, 366 mmol), and the reaction mixture was stirred at $-20\text{ }^{\circ}\text{C}$ for 2 hours, then re-cooled to $-78\text{ }^{\circ}\text{C}$. Iodomethane (52.1 g, 367 mmol) was added at $-78\text{ }^{\circ}\text{C}$, and the reaction mixture was allowed to stir at $-20\text{ }^{\circ}\text{C}$ for 2 hours. Saturated aqueous ammonium chloride solution was added, and the mixture was extracted with ethyl acetate (3 x 500 mL). The combined organic layers were washed with water (100 mL) and with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo*. Purification via chromatography on silica gel (Eluent: 10:1 petroleum ether / ethyl acetate) afforded the product as a yellow oil. Yield: 10 g, 24 mmol, 66%. LCMS m/z 424.1 $[\text{M}+\text{H}^+]$. ^1H NMR (400 MHz, CDCl_3) δ 8.58 (s, 1H), 5.69 (s, 2H), 3.48-3.54 (m, 2H), 2.60 (s, 3H), 0.88-0.95 (m, 2H), -0.05 (s, 9H).

Step 2. Synthesis of 3-(4-chloro-6-methyl-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C7**).

To a solution of 4-chloro-5-iodo-6-methyl-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidine (**C6**) (300 mg, 0.71 mmol) and (3-cyanophenyl)boronic acid (104 mg, 0.708 mmol) in a mixture of 1,2-dimethoxyethane and water (5:1, 10 mL) was added potassium carbonate (193 mg, 1.40 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (17 mg, 23 μmol). The reaction mixture was degassed and purged with nitrogen; this procedure was carried out a total of three times. After heating at reflux for 18 hours, the reaction mixture was cooled to room temperature and extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (50 mL), dried

over sodium sulfate, filtered and concentrated *in vacuo*. Purification via silica gel chromatography (Gradient: 0% to 20% ethyl acetate in petroleum ether) afforded the product as a yellow oil. Yield: 140 mg, 0.351 mmol, 50%. LCMS *m/z* 399.2 [M+H⁺].

Step 3. Synthesis of 3-[6-methyl-4-(morpholin-4-yl)-7-{{2-(trimethylsilyl)ethoxy}methyl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C8**).

To a solution of 3-(4-chloro-6-methyl-7-{{2-(trimethylsilyl)ethoxy}methyl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile (**C7**) (140 mg, 0.351 mmol) in *n*-butanol (5 mL) was added morpholine (30.6 mg, 0.351 mmol) and *N,N*-diisopropylethylamine (90.9 mg, 0.703 mmol). The reaction mixture was heated at 100 °C for 18 hours, then concentrated *in vacuo* to provide the product as a yellow solid. Yield: 125 mg, 0.278 mmol, 79%. LCMS *m/z* 450.3 [M+H⁺].


Step 4. Synthesis of 3-[7-(hydroxymethyl)-6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C9**).

A solution of 3-[6-methyl-4-(morpholin-4-yl)-7-{{2-(trimethylsilyl)ethoxy}methyl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C8**) (125 mg, 0.278 mmol) in trifluoroacetic acid (10 mL) was stirred at room temperature for 2 hours, then concentrated *in vacuo* to provide the product as a yellow solid (125 mg). This was used in the next step without additional purification. LCMS *m/z* 349.9 [M+H⁺].

Step 5. Synthesis of 3-[6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**2**).

A solution of 3-[7-(hydroxymethyl)-6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C9**) (from the previous step, 125 mg, ≤0.278 mmol) in acetonitrile (5 mL) was adjusted to a pH of >11 by addition of solid potassium carbonate. The mixture was filtered and concentrated *in vacuo*; purification via preparative HPLC (Column: Boston Symmetrix ODS-H, 5 μm; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile; Gradient: 19% to 39% B) afforded the product as a yellow solid. Yield: 54 mg, 0.17 mmol, 61% over two steps. LCMS *m/z* 319.9 [M+H⁺]. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.17 (br s, 1H), 8.34 (s, 1H), 7.77-7.83 (m, 2H), 7.65-7.74 (m, 2H), 3.27-3.33 (m, 4H), 2.98-3.06 (m, 4H), 2.36 (s, 3H).

Example 3

6-[4-(Morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile (**3**)

Step 1. Synthesis of 6-[4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile (**C10**).

To a solution of 6-bromopyridine-2-carbonitrile (80 mg, 0.44 mmol) and 4-(morpholin-4-yl)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**P2**) (241 mg, 0.523 mmol) in 1,4-dioxane (2.5 mL) and water (0.5 mL) was added tetrakis(triphenylphosphine)palladium(0) (51 mg, 44 μ mol) and sodium carbonate (140 mg, 1.32 mmol). The reaction mixture was heated at 120 $^{\circ}$ C under microwave irradiation for 15 minutes, then diluted with water (30 mL) and extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo*; purification via preparative thin layer chromatography (Eluent: 1:1 petroleum ether / ethyl acetate) afforded the product as a brown oil. Yield: 110 mg, 0.252 mmol, 57%. ^1H NMR (400 MHz, CDCl_3) δ 8.52 (s, 1H), 7.84-7.93 (m, 2H), 7.74 (s, 1H), 7.59 (dd, $J=7.0, 1.2$ Hz, 1H), 5.66 (s, 2H), 3.56-3.65 (m, 6H), 3.34-3.40 (m, 4H), 0.93 (dd, $J=8.3, 8.0$ Hz, 2H), -0.05 (s, 9H).

Step 2. Synthesis of 6-[7-(hydroxymethyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile (**C11**).


A solution of 6-[4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile (**C10**) (110 mg, 0.252 mmol) in trifluoroacetic acid (3 mL) was stirred at room temperature for 2 hours. The reaction mixture was concentrated *in vacuo* to provide the product as a yellow oil, which was used for the next step without additional purification.

Step 3. Synthesis of 6-[4-(morpholin-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile (**3**).

A solution of 6-[7-(hydroxymethyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile (**C11**) (from the previous step, 85 mg, ≤ 0.25 mmol) in acetonitrile (3 mL) was brought to a pH of >12 via addition of solid potassium carbonate. After 30 minutes at room temperature, the reaction mixture was filtered and concentrated *in vacuo*. Purification via preparative HPLC (Column: Phenomenex Gemini C18, 8 μ m; Mobile phase A: ammonia in water, pH 10; Mobile phase B: acetonitrile; Gradient: 10% to 50% B) afforded the product as a white solid. Yield: 15.2 mg, 49.6 μ mol, 20% over two steps. LCMS m/z 307.2 $[M+H]^+$. ^1H NMR (400 MHz, DMSO- d_6) δ 8.38 (s, 1H), 8.13 (dd, $J=8.1, 7.7$ Hz, 1H), 7.96 (br d, $J=8$ Hz, 1H), 7.89 (br d, $J=8$ Hz, 1H), 7.86 (s, 1H), 3.50-3.55 (m, 4H), 3.19-3.24 (m, 4H).

Example 4

3-[4-(Morpholin-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**4**)



Step 1. Synthesis of 3-(4-chloro-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile (**C12**).

To a stirred mixture of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C1**) (8.2 g, 20 mmol), (3-cyanophenyl)boronic acid (3.2 g, 22 mmol) and potassium carbonate (8.3 g, 60 mmol) in a mixture of 1,2-dimethoxyethane and water (4:1 ratio, 250 mL) was added [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (731 mg, 1.00 mmol). The reaction mixture was degassed and then charged with nitrogen; this procedure was carried out a total of three times. The reaction mixture was heated at reflux for 3 hours, then cooled to room temperature and diluted with saturated aqueous sodium chloride solution (100 mL). The organic layer was dried over sodium sulfate, filtered, and concentrated under reduced pressure. Purification via silica gel column chromatography (Eluent: 10:1 petroleum ether / ethyl acetate) provided the product as a yellow oil. Yield: 5.0 g, 12 mmol, 60%. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.75 (s, 1H), 8.13 (s, 1H), 8.00-8.02 (m, 1H), 7.84-7.92 (m, 2H), 7.68 (dd, *J*=7.8, 7.8 Hz, 1H), 5.70 (s, 2H), 3.60 (dd, *J*=8.0, 8.0 Hz, 2H), 0.86 (dd, *J*=8.0, 8.0 Hz, 2H), -0.08 (s, 9H).

Step 2. Synthesis of 3-[4-chloro-7-(hydroxymethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**C13**).

A solution of 3-(4-chloro-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C12**) (3.8 g, 9.9 mmol) in trifluoroacetic acid (25 mL) was stirred at room temperature for 2 hours. The reaction mixture was concentrated *in vacuo* to afford the product (4 g, >100% mass recovery) as a yellow oil, which was used in the next step without further purification.

Step 3. Synthesis of 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C14**).

A solution of 3-[4-chloro-7-(hydroxymethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**C13**) (4 g from the previous step, ≤9.9 mmol) in methanol (100 mL) was adjusted to pH >12 by addition of solid potassium carbonate. Solvent was removed *in vacuo* and the residue was mixed with water (100 mL). The resulting solid was isolated via filtration and washed with water, providing the product as a white solid. Yield: 1.3 g, 5.1 mmol, 52% over two steps. LCMS *m/z* 255.0 [M+H⁺].


Step 4. Synthesis of 3-[4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**4**).

Morpholine (871 mg, 10 mmol) and *N,N*-diisopropylethylamine (2.6 g, 20 mmol) were added to a solution of 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C14**) (2.5 g, 9.8 mmol) in *n*-butanol (100 mL), and the reaction mixture was heated at

reflux for 3 hours. Solvents were removed *in vacuo* and the residue was purified using chromatography on silica gel (Eluent: 1:1 ethyl acetate / petroleum ether). Subsequent recrystallization from ethyl acetate and *tert*-butyl methyl ether afforded the product as a white solid. Yield: 770 mg, 2.52 mmol, 26%. LCMS *m/z* 306.0 [M+H⁺]. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.34 (br s, 1H), 8.41 (s, 1H), 7.99-8.02 (m, 1H), 7.89 (br d, *J*=8 Hz, 1H), 7.76 (br d, *J*=7.5 Hz, 1H), 7.71 (s, 1H), 7.68 (dd, *J*=7.8, 7.8 Hz, 1H), 3.44-3.50 (m, 4H), 3.11-3.17 (m, 4H).

Example 5


3-[4-(Dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**5**)



A mixture of 3-(4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile (**C14**) (157 mg, 0.616 mmol), dimethylamine (189 mg, 4.19 mmol) and triethylamine (182 mg, 1.80 mmol) in *n*-butanol (12 mL) was heated under microwave irradiation at 150 °C for 25 minutes. After concentration *in vacuo*, the residue was purified by preparative HPLC (Column: Phenomenex Gemini C18, 8 μm; Mobile phase A: ammonia in water, pH 10; Mobile phase B: acetonitrile; Gradient: 25% to 65% B) to give the product as a white solid. Yield: 72.1 mg, 0.274 mmol, 44%. LCMS *m/z* 263.8 [M+H⁺]. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.15 (br s, 1H), 8.28 (s, 1H), 7.93 (br s, 1H), 7.78 (br d, *J*=7.8 Hz, 1H), 7.73 (br d, *J*=7.8 Hz, 1H), 7.62 (br dd, *J*=8.0, 7.8 Hz, 1H), 7.57 (br s, 1H), 2.73 (s, 6H).

Example 6


3-{4-[(2*S*)-2-Methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile (**6**)



To a solution of 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C14**) (100 mg, 0.393 mmol) and (2S)-2-methylmorpholine (54 mg, 0.53 mmol) in *n*-butanol (20 mL) was added *N,N*-diisopropylethylamine (152 mg, 1.18 mol), and the reaction mixture was heated at 100 °C for 18 hours. The reaction mixture was then concentrated *in vacuo*; purification via preparative HPLC (Column: Phenomenex Gemini C18, 8 µm; Mobile phase A: ammonia in water, pH 10; Mobile phase B: acetonitrile; Gradient: 27% to 47% B) provided the product as a white solid. Yield: 17.5 mg, 54.8 µmol, 14%. LCMS *m/z* 320.2 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃) δ 10.71 (br s, 1H), 8.53 (br s, 1H), 7.73-7.89 (m, 2H), 7.51-7.69 (m, 2H), 7.28 (s, 1H, assumed; partially obscured by solvent peak), 3.56-3.77 (m, 3H), 3.39-3.54 (m, 2H), 2.86-2.98 (m, 1H), 2.53-2.65 (m, 1H), 0.93-1.03 (m, 3H).

Example 7

(3-{4-[(3S)-3-Methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}phenyl)methanol (**7**)



15

Step 1. Synthesis of 5-bromo-7-[(4-methylphenyl)sulfonyl]-4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidine (**C15**).

5-Bromo-4-chloro-7-[(4-methylphenyl)sulfonyl]-7H-pyrrolo[2,3-d]pyrimidine (**P3**) was reacted with the hydrochloride salt of (3S)-3-methylpiperidine according to the method described for preparation of 3-{4-[(2S)-2-methylmorpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile (**6**) in Example 6. In this case, purification was effected via

20

silica gel chromatography (Eluent: 5:1 petroleum ether / ethyl acetate), providing the product as a white solid. Yield: 4.3 g, 9.6 mmol, 92%.

Step 2. Synthesis of (3-{7-[(4-methylphenyl)sulfonyl]-4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}phenyl)methanol (**C16**).


5 A mixture of 5-bromo-7-[(4-methylphenyl)sulfonyl]-4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidine (**C15**) (0.225 g, 0.501 mmol), [3-(hydroxymethyl)phenyl]boronic acid (0.104 g, 0.684 mmol), sodium carbonate (0.159 g, 1.50 mmol) and dichlorobis(triphenylphosphine)palladium(II) (36 mg, 51 μ mol) in acetonitrile (2 mL) and water (2 mL) was heated to 150 °C under microwave irradiation
10 for 15 minutes. The reaction was concentrated *in vacuo* to afford the product (0.35 g, >100%), which was used directly in the following step without additional purification.

Step 3. Synthesis of (3-{4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}phenyl)methanol (**7**).

To a stirred solution of (3-{7-[(4-methylphenyl)sulfonyl]-4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}phenyl)methanol (**C16**) from the previous step
15 (<0.50 mmol) in 2-propanol (20 mL) was added lithium hydroxide monohydrate (0.42 g, 10 mmol) and water (3 mL) and the reaction mixture was stirred at room temperature for 18 hours. After concentration *in vacuo*, the residue was purified by preparative HPLC (Column: Waters XBridge; Mobile phase A: 0.1% ammonia in water; Mobile phase B: acetonitrile; Gradient: 44% to 60% B) to provide the product as a white solid. Yield: 130
20 mg, 0.403 mmol, 80% over two steps. LCMS *m/z* 323.4 [$M+H^+$]. ^1H NMR (400 MHz, CD_3OD) δ 8.24 (s, 1H), 7.46 (br s, 1H), 7.35-7.43 (m, 2H), 7.30 (br d, $J=6$ Hz, 1H), 7.23 (s, 1H), 4.67 (s, 2H), 3.89 (br d, $J=13$ Hz, 1H), 3.79 (br d, $J=12$ Hz, 1H), 2.58-2.69 (m, 1H), 2.27 (dd, $J=12.0, 11.0$ Hz, 1H), 1.62-1.71 (m, 1H), 1.30-1.52 (m, 3H), 0.87-1.01 (m,
25 1H), 0.60 (d, $J=6.5$ Hz, 3H).

Example 8

4-(Morpholin-4-yl)-5-(1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine, formate salt (**8**)



Step 1. Synthesis of 1-(tetrahydro-2*H*-pyran-2-yl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (**C17**).

- 5 3,4-Dihydro-2*H*-pyran (5.6 g, 67 mmol) and trifluoroacetic acid (1.17 g, 10.3 mmol) were added to a solution of 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole (10.0 g, 51.5 mmol) in toluene (200 mL), and the reaction mixture was heated to 90 °C for 2 hours. After cooling to room temperature, the reaction mixture was partitioned between ethyl acetate (200 mL) and saturated aqueous sodium bicarbonate solution (100 mL), and the aqueous layer was extracted with ethyl acetate (100 mL).
 10 The combined organic layers were washed with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated *in vacuo*. Silica gel chromatography (Gradient: 10% to 50% ethyl acetate in petroleum ether) provided the product as a white solid. Yield: 13.4 g, 48.2 mmol, 94%. ¹H NMR (400 MHz, CDCl₃)

δ 7.94 (s, 1H), 7.83 (s, 1H), 5.41 (dd, $J=9.5$, 2.5 Hz, 1H), 4.01-4.08 (m, 1H), 3.65-3.74 (m, 1H), 1.98-2.18 (m, 3H), 1.6-1.76 (m, 3H), 1.32 (s, 12H).

Step 2. Synthesis of 4-chloro-5-[1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl]-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C18**).

5 A mixture of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C1**) (2.0 g, 4.9 mmol), 1-(tetrahydro-2H-pyran-2-yl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (**C17**) (1.91 g, 6.87 mmol), potassium phosphate (4.0 g, 19 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.10 g, 87 μ mol) was degassed several times with nitrogen and irradiated in a microwave synthesizer at 130
10 °C for 1.5 hours. The reaction mixture was partitioned between ethyl acetate (400 mL) and water (60 mL), and the aqueous layer was extracted with ethyl acetate (200 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. Chromatography on silica gel (Gradient: 10% to 80% ethyl acetate in
15 petroleum ether) afforded the product as a brown oil. Yield: 1.33 g, 3.06 mmol, 62%. ^1H NMR (400 MHz, CDCl_3) δ 8.67 (s, 1H), 7.87 (s, 1H), 7.76 (s, 1H), 7.38 (s, 1H), 5.68 (s, 2H), 5.46 (dd, $J=9.4$, 2.9 Hz, 1H), 4.08-4.15 (m, 1H), 3.71-3.79 (m, 1H), 3.57 (dd, $J=8.3$, 8.0 Hz, 2H), 2.04-2.24 (m, 3H), 1.61-1.79 (m, 3H), 0.94 (dd, $J=8.3$, 8.3 Hz, 2H), -0.03 (s, 9H).

20 Step 3. Synthesis of 4-(morpholin-4-yl)-5-[1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl]-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C19**).

A solution of 4-chloro-5-[1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl]-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**C18**) (3.3 g, 7.6 mmol), morpholine (0.99 g, 11 mmol) and *N,N*-diisopropylethylamine (6 mL, 34 mmol) in *n*-
25 butanol (15 mL) was irradiated in a microwave synthesizer at 100 °C for 30 minutes. After removal of solvents *in vacuo*, the residue was purified by silica gel chromatography (Gradient: 10% to 50% ethyl acetate in petroleum ether) to provide the product as a brown oil. Yield: 2.6 g, 5.4 mmol, 71%. ^1H NMR (400 MHz, CDCl_3) δ 8.50 (s, 1H), 7.79 (s, 1H), 7.71 (s, 1H), 7.17 (s, 1H), 5.62 (s, 2H), 5.44 (dd, $J=6.5$, 6.0 Hz, 1H), 4.08-4.15 (m, 1H), 3.71-3.79 (m, 1H), 3.62-3.67 (m, 4H), 3.58 (dd, $J=8.5$, 8.0 Hz, 2H), 3.33-3.39 (m, 4H), 2.06-2.19 (m, 3H), 1.62-1.79 (m, 3H), 0.93 (dd, $J=8.5$, 7.5 Hz, 2H), -0.04 (s, 9H).

Step 4. Synthesis of 4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C20**).

To a solution of 4-(morpholin-4-yl)-5-[1-(tetrahydro-2*H*-pyran-2-yl)-1*H*-pyrazol-4-yl]-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C19**) (4.0 g, 8.2 mmol) in dichloromethane (20 mL) was added a solution of hydrogen chloride in 1,4-dioxane (4 M, 100 mL). The reaction mixture was stirred at room temperature for 2 hours, then poured into a mixture of ethyl acetate (300 mL) and saturated aqueous sodium bicarbonate solution (300 mL). The aqueous layer was extracted with ethyl acetate (2 x 300 mL), and the combined organic layers were washed with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. Purification via silica gel chromatography (Gradient: 10% to 80% ethyl acetate in petroleum ether) afforded the product as a yellow oil. Yield: 2.18 g, 5.44 mmol, 66%. LCMS *m/z* 401.3 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃) δ 8.51 (s, 1H), 7.78 (s, 2H), 7.19 (s, 1H), 5.64 (s, 2H), 3.56-3.66 (m, 6H), 3.34-3.40 (m, 4H), 0.91-0.98 (m, 2H), -0.03 (s, 9H).

Step 5. Synthesis of [4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl]methanol (**C21**).

4-(Morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C20**) (0.13 g, 0.32 mmol) was dissolved in trifluoroacetic acid (10 mL) and stirred at room temperature for 2 hours. Removal of solvent *in vacuo* provided the product (100 mg) as a brown solid; this was used in the next step without further purification. LCMS *m/z* 301.2 [M+H⁺].


Step 6. Synthesis of 4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine, formate salt (**8**).

To a solution of [4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-7-yl]methanol (**C21**) (from the previous step, 0.10 g, ≤0.32 mmol) in acetonitrile (10 mL) was added potassium carbonate (0.34 g, 2.5 mmol) and the reaction mixture was stirred at room temperature for 30 minutes. The mixture was filtered and the filter cake was washed with ethyl acetate (3 x 20 mL). The combined filtrates were concentrated *in vacuo*; purification via preparative HPLC (Column: DIKMA Diamonsil(2) C18, 5 μm; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile; Gradient: 0% to 17% B) afforded the product as a pale solid. Yield: 36 mg, 0.11 mmol, 34% over 2 steps. LCMS *m/z* 271.2 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃) δ

10.72 (br s, 1H), 8.47 (s, 1H), 7.78 (s, 2H), 7.19 (s, 1H), 3.60-3.67 (m, 4H), 3.41-3.48 (m, 4H).

Example 9

5 3-[2-Methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**9**)



Step 1. Synthesis of 4-chloro-5-iodo-2-methyl-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C22**).

To a solution of 4-chloro-2-methyl-7*H*-pyrrolo[2,3-*d*]pyrimidine (400 mg, 2.4 mmol) in dichloromethane (10 mL) was added *N*-iodosuccinimide (537 mg, 2.39 mmol). The mixture was stirred at room temperature for 2 hours, then washed with aqueous sodium sulfite solution, dried over sodium sulfate, filtered, and concentrated *in vacuo* to provide the product as a brown solid. Yield: 330 mg, 1.12 mmol, 47%. LCMS *m/z* 293.8 [M+H⁺].

Step 2. Synthesis of 4-chloro-5-iodo-2-methyl-7-([2-(trimethylsilyl)ethoxy]methyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C23**).

4-Chloro-5-iodo-2-methyl-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C22**) was converted to the product according to the method used for synthesis of 4-chloro-5-iodo-7-([2-

(trimethylsilyl)ethoxy)methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C1**) in Preparation P1. The product was obtained as a yellow oil. Yield: 400 mg, 0.94 mmol, 84%. LCMS *m/z* 424.0 [M+H⁺].

Step 3. Synthesis of 5-iodo-2-methyl-4-(morpholin-4-yl)-7-[[2-

5 (trimethylsilyl)ethoxy)methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C24**).

The product was prepared from 4-chloro-5-iodo-2-methyl-7-[[2-(trimethylsilyl)ethoxy)methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C23**) according to the method described for synthesis of 3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile (**6**) in Example 6. In this case, purification was carried out via silica gel
10 chromatography (Gradient: 0% to 50% ethyl acetate in petroleum ether) to provide the product as a yellow oil. Yield: 300 mg, 0.63 mmol, 67%. LCMS *m/z* 475.2 [M+H⁺].

Step 4. Synthesis of 3-[2-methyl-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy)methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C25**).

To a solution of 5-iodo-2-methyl-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy)methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C24**) (100 mg, 0.21 mmol),
15 (3-cyanophenyl)boronic acid (62 mg, 0.42 mmol) and potassium carbonate (100 mg, 0.72 mmol) in a mixture of ethanol and water (4:1, 5 mL) was added dichlorobis(triphenylphosphine)palladium (15 mg, 21 μmol). The reaction mixture was degassed and purged with nitrogen; this procedure was carried out a total of three
20 times. After the reaction mixture had heated at 90 °C for 3 hours, it was allowed to cool and concentrated *in vacuo*. Purification via preparative thin layer chromatography (Eluent: 1:1 petroleum ether / ethyl acetate) provided the product as a yellow oil. Yield: 52 mg, 0.12 mmol, 57%. LCMS *m/z* 451.3 [M+H⁺].


Step 5. Synthesis of 3-[2-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**9**).
25

3-[2-Methyl-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy)methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C25**) was converted to the product using the method described for synthesis of 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**1**) in Example 1. In this case, preparative HPLC purification
30 was carried out using a Phenomenex Gemini C18 column, 8 μm; Mobile phase A: ammonia in water, pH 10; Mobile phase B: acetonitrile; Gradient: 30% to 50% B. The product was obtained as a white solid. Yield: 17.3 mg, 54.2 μmol, 45% over 2 steps. LCMS *m/z* 320.1 [M+H⁺]. ¹H NMR (400 MHz, CD₃OD) δ 7.91-7.93 (m, 1H), 7.85-7.88

(m, 1H), 7.60-7.69 (m, 2H), 7.38 (s, 1H), 3.51-3.55 (m, 4H), 3.24-3.28 (m, 4H), 2.59 (s, 3H).

Example 10

5 {3-[4-(Dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol (**10**)



Step 1. Synthesis of {3-[4-(dimethylamino)-7-([2-(trimethylsilyl)ethoxy]methyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol (**C26**).

A solution of 5-iodo-*N,N*-dimethyl-7-([2-(trimethylsilyl)ethoxy]methyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine (**C3**) (418 mg, 1.00 mmol), [3-(hydroxymethyl)phenyl]boronic acid (228 mg, 1.50 mmol), tetrakis(triphenylphosphine)palladium(0) (115 mg, 0.100 mmol), and cesium carbonate (625 mg, 1.92 mmol) in 1,4-dioxane (6 mL) and water (1.5 mL) was purged with nitrogen, then heated under microwave irradiation at 120 °C for 20 minutes. The reaction mixture was diluted with saturated aqueous sodium chloride solution (50 mL) and extracted with dichloromethane (3 x 60 mL). The combined organic layers were dried over sodium sulfate, filtered, and concentrated *in vacuo*. Chromatography on silica gel (Gradient: 2% to 17% ethyl acetate in petroleum ether) afforded the product as a brown oil. Yield: 369 mg, 0.926 mmol, 93%. LCMS *m/z* 399.0 [*M*+*H*⁺].

Step 2. Synthesis of 3-[4-(dimethylamino)-7-(hydroxymethyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzyl trifluoroacetate (**C27**).

A solution of {3-[4-(dimethylamino)-7-([2-(trimethylsilyl)ethoxy]methyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol (**C26**) (397 mg, 0.996 mmol) in


trifluoroacetic acid (10 mL) was stirred at room temperature for 2 hours. The reaction mixture was concentrated *in vacuo* to provide the product (400 mg) as a brown oil. This material was taken directly into the following step. LCMS *m/z* 395.1 [M+H⁺].

Step 3. Synthesis of {3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol (**10**).

A mixture of 3-[4-(dimethylamino)-7-(hydroxymethyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzyl trifluoroacetate (**C27**) (from the previous step, 400 mg, ≤0.996 mmol) and potassium carbonate (500 mg, 3.6 mmol) in methanol (10 mL) was stirred at 80 °C for 30 minutes. The reaction mixture was diluted with saturated aqueous sodium chloride solution (50 mL) and extracted with dichloromethane (3 x 60 mL); the combined organic layers were dried over sodium sulfate, filtered, and concentrated *in vacuo*. Purification via preparative HPLC (Column: Phenomenex Gemini C18, 8 μm; Mobile phase A: ammonia in water, pH 10; Mobile phase B: acetonitrile; Gradient: 21% to 33% B) afforded the product as a white solid. Yield: 77.7 mg, 0.290 mmol, 29% over two steps. LCMS *m/z* 268.9 [M+H⁺]. ¹H NMR (400 MHz, CDCl₃) δ 10.83 (br s, 1H), 8.39 (s, 1H), 7.50 (br s, 1H), 7.38-7.42 (m, 2H), 7.29-7.34 (m, 1H), 7.11 (s, 1H), 4.78 (s, 2H), 2.84 (s, 6H).

Example 198

3-[6-(Difluoromethyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**198**)



Step 1. Synthesis of 4-chloro-5-iodo-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**C28**).

A solution of diisopropylamine (2.9 g, 29 mmol) in tetrahydrofuran (30 mL) was cooled to $-78\text{ }^{\circ}\text{C}$ and treated drop-wise with *n*-butyllithium (2.5 M, 11.6 mL, 29 mmol). The reaction mixture was stirred at $0\text{ }^{\circ}\text{C}$ for 1 hour, and then cooled to $-78\text{ }^{\circ}\text{C}$. A solution of 4-chloro-5-iodo-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidine (**C1**) (8.0 g, 20 mmol) in tetrahydrofuran (10 mL) was added drop-wise, and stirred was continued at $-78\text{ }^{\circ}\text{C}$ for 1 hour. After drop-wise addition of a solution of ethyl formate (2.6 g, 35 mmol) in tetrahydrofuran (10 mL) to the $-78\text{ }^{\circ}\text{C}$ reaction mixture, it was allowed to warm to room temperature and stir for 18 hours. It was then cooled to $0\text{ }^{\circ}\text{C}$, and the reaction was quenched by addition of saturated aqueous ammonium chloride solution (20 mL), then extracted with ethyl acetate (3 x 50 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (50 mL), dried over sodium sulfate, filtered, and concentrated. Purification via silica gel chromatography (Gradient: 0% to 25% ethyl acetate in petroleum ether) afforded the

product as a yellow oil. Yield: 5.0 g, 11 mmol, 55%. ¹H NMR (400 MHz, CDCl₃) δ 10.11 (s, 1H), 8.80 (s, 1H), 6.05 (s, 2H), 3.55-3.62 (m, 2H), 0.89-0.96 (m, 2H), -0.05 (s, 9H).

Step 2. Synthesis of 5-iodo-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**C29**).

5 Morpholine (1.09 g, 12.5 mmol) and *N,N*-diisopropylethylamine (2.94 g, 22.7 mmol) were added to a solution of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**C28**) (5.0 g, 11 mmol) in acetonitrile (20 mL). The reaction mixture was stirred at reflux for 16 hours, whereupon it was cooled and concentrated *in vacuo*; the residue was purified by chromatography on silica gel
10 (Gradient: 0% to 20% ethyl acetate in petroleum ether) to provide the product as a yellow oil. Yield: 4.0 g, 8.2 mmol, 74%. ¹H NMR (400 MHz, CDCl₃) δ 9.99 (s, 1H), 8.54 (s, 1H), 6.01 (s, 2H), 3.92-3.98 (m, 4H), 3.70-3.75 (m, 4H), 3.58-3.65 (m, 2H), 0.91-0.96 (m, 2H), -0.05 (s, 9H).

Step 3. Synthesis of 3-[6-formyl-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**C30**).

5-Iodo-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**C29**) was converted to the product using the method described for synthesis of 3-(4-chloro-6-methyl-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C7**) in Example 2. The product was obtained as
20 a yellow solid. Yield: 1.5 g, 3.2 mmol, 78%. ¹H NMR (400 MHz, CDCl₃) δ 9.75 (s, 1H), 8.60 (s, 1H), 7.77-7.83 (m, 2H), 7.72-7.76 (m, 1H), 7.67 (br dd, *J*=8, 8 Hz, 1H), 6.09 (s, 2H), 3.66-3.71 (m, 2H), 3.38-3.44 (m, 4H), 3.23-3.28 (m, 4H), 0.94-1.01 (m, 2H), -0.03 (s, 9H).

Step 4. Synthesis of 3-[6-(difluoromethyl)-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**C31**).

A solution of 3-[6-formyl-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**C30**) (200 mg, 0.43 mmol) and (diethylamino)sulfur trifluoride (276 mg, 1.71 mmol) in dichloromethane (10 mL) was stirred at 40 °C for 16 hours. The reaction mixture was poured into saturated aqueous
30 sodium bicarbonate solution (10 mL) and extracted with dichloromethane (3 x 20 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (50 mL), dried over sodium sulfate, filtered, and concentrated; purification by preparative thin-layer chromatography on silica gel (Eluent: 1:1 ethyl acetate /


petroleum ether) afforded the product as a yellow solid. Yield: 150 mg, 0.31 mmol, 72%. ^1H NMR (400 MHz, CDCl_3) δ 8.58 (s, 1H), 7.76 (ddd, $J=7.3, 1.6, 1.5$ Hz, 1H), 7.72-7.74 (m, 1H), 7.61-7.70 (m, 2H), 6.74 (t, $J_{\text{HF}}=52.5$ Hz, 1H), 5.86 (s, 2H), 3.65-3.72 (m, 2H), 3.34-3.42 (m, 4H), 3.15-3.21 (m, 4H), 0.95-1.02 (m, 2H), -0.02 (s, 9H).

- 5 Step 5. Synthesis of 3-[6-(difluoromethyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**198**).

A solution of 3-[6-(difluoromethyl)-4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile (**C31**) (10 mg, 21 μmol) in trifluoroacetic acid (2 mL) was stirred at room temperature for 24 hours. The reaction mixture was concentrated and purified by preparative reversed phase high-performance liquid chromatography (Column: DIKMA Diamonsil(2) C18, 5 μm ; Eluent: 22% acetonitrile in water containing 0.225% formic acid) to provide the product as a white solid. Yield: 2.0 mg, 5.6 μmol , 27%. LCMS m/z 355.9 $[\text{M}+\text{H}^+]$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 13.06 (br s, 1H), 8.47 (s, 1H), 7.92 (ddd, $J=6.7, 2.1, 1.7$ Hz, 1H), 7.83-7.85 (m, 1H), 7.69-7.76 (m, 2H), 7.02 (t, $J_{\text{HF}}=52.5$ Hz, 1H), 3.22-3.30 (m, 4H), 3.03-3.10 (m, 4H).

Example 199 and Example 200

- 5-(5,6-Dihydro-2H-pyran-3-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (**199**) and
20 5-(3,4-Dihydro-2H-pyran-5-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine (**200**)



Step 1. Synthesis of 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)tetrahydro-2H-pyran-3-ol (**C32**).

To a $-78\text{ }^{\circ}\text{C}$ solution of 4-chloro-5-iodo-7*H*-pyrrolo[2,3-*d*]pyrimidine (1.5 g, 5.4 mmol) in tetrahydrofuran (50 mL) was added *n*-butyllithium (2.5 M, 6.4 mL, 16.1 mmol) drop-wise. After the reaction mixture had been stirred at $-78\text{ }^{\circ}\text{C}$ for 2 hours, it was treated with dihydro-2*H*-pyran-3(4*H*)-one (1.61 g, 16.1 mmol), warmed to room temperature, and stirred for 18 hours. The reaction was quenched with water (50 mL), and the aqueous layer was extracted with ethyl acetate (3 x 30 mL); the combined organic layers were washed with saturated aqueous sodium chloride solution (100 mL), dried over sodium sulfate, filtered, and concentrated. Silica gel chromatography (Gradient: 0% to 50% ethyl acetate in petroleum ether) afforded the product as a yellow oil. Yield: 200 mg, 0.79 mmol, 15%. LCMS m/z 254.0 $[\text{M}+\text{H}^+]$.

Step 2. Synthesis of 3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]tetrahydro-2*H*-pyran-3-ol (**C33**).

To a solution of 3-(4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)tetrahydro-2*H*-pyran-3-ol (**C32**) (200 mg, 0.79 mmol) and morpholine (134 mg, 1.54 mol) in *n*-butanol (10 mL) was added *N,N*-diisopropylethylamine (305 mg, 2.36 mol), and the reaction mixture was heated at $70\text{ }^{\circ}\text{C}$ for 2 hours. After concentration *in vacuo*, preparative reversed phase high-performance liquid chromatography (Column: Phenomenex Gemini C18, 8 μm ; Mobile phase A: aqueous ammonia, pH 10; Mobile phase B: acetonitrile; Gradient: 9% to 29% B) provided the product as a white solid. Yield: 180 mg, 0.59 mmol, 75%. LCMS m/z 305.1 $[\text{M}+\text{H}^+]$. ^1H NMR (400 MHz, $\text{CDCl}_3 + \text{D}_2\text{O}$) δ 8.73 (s, 1H), 7.52 (s, 1H), 3.92-4.03 (m, 4H), 3.74-3.88 (m, 4H), 3.32-3.40 (m, 2H), 3.15-3.24 (m, 2H), 2.07-2.16 (m, 1H), 1.88-2.06 (m, 2H), 1.57-1.68 (m, 1H).

Step 3. Synthesis of 5-(5,6-dihydro-2*H*-pyran-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**199**) and 5-(3,4-dihydro-2*H*-pyran-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine (**200**).

Triethylsilane (10 mL) and trifluoroacetic acid (10 mL) were added over 10 minutes to a solution of 3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]tetrahydro-2*H*-pyran-3-ol (**C33**) (40 mg, 0.13 mmol) in dichloromethane (10 mL). The reaction mixture was stirred at room temperature for 2 hours, whereupon it was concentrated *in vacuo*. Preparative reversed phase high-performance liquid chromatography (Column: Boston Symmetrix ODS-H, 5 μm ; Mobile phase A: water containing 0.225% formic acid; Mobile phase B: acetonitrile; Gradient: 11% to 31% B) afforded **199** as a white solid


(Yield: 11.3 mg, 39.5 μ mol, 30%) and **200**, also as a white solid (Yield: 7.0 mg, 24 μ mol, 18%).

199: LCMS m/z 287.0 $[M+H]^+$. ^1H NMR (400 MHz, DMSO- d_6) δ 11.97 (br s, 1H), 8.29 (s, 1H), 7.31 (s, 1H), 5.87-5.93 (m, 1H), 4.30-4.36 (m, 2H), 3.68-3.78 (m, 6H), 3.36-3.43 (m, 4H), 2.21-2.29 (m, 2H).

200: LCMS m/z 287.0 $[M+H]^+$. ^1H NMR (400 MHz, DMSO- d_6) δ 11.83 (br s, 1H), 8.26 (s, 1H), 7.18 (s, 1H), 6.65 (s, 1H), 3.92-3.97 (m, 2H), 3.69-3.76 (m, 4H), 3.40-3.46 (m, 4H), 2.29-2.36 (m, 2H), 1.88-1.96 (m, 2H).

Example 201

4-(Morpholin-4-yl)-5-[3-(1,2,4-oxadiazol-3-yl)phenyl]-7H-pyrrolo[2,3-*d*]pyrimidine (**201**)



Step 1. Synthesis of 3-[4-(morpholin-4-yl)-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C34**).

3-[4-(Morpholin-4-yl)-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**4**) was converted to the product using the method described for synthesis of 4-chloro-5-iodo-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-*d*]pyrimidine (**C1**) in Preparation P1. The product was obtained as a yellow solid. Yield: 0.90 g, 2.1 mmol, 70%. ^1H NMR (400 MHz, $CDCl_3$) δ 8.55 (s, 1H), 7.86 (br s, 1H), 7.79 (br d, $J=7.6$ Hz, 1H), 7.63 (br d, $J=7.6$ Hz, 1H), 7.57 (dd, $J=7.6$, 7.6 Hz, 1H), 7.29 (s, 1H), 5.67 (s, 2H), 3.58-3.64 (m, 2H), 3.52-3.58 (m, 4H), 3.27-3.32 (m, 4H), 0.92-0.98 (m, 2H), -0.03 (s, 9H).

Step 2. Synthesis of *N'*-hydroxy-3-[4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzenecarboximidamide (**C35**).


Hydroxylamine hydrochloride (278 mg, 4.00 mmol) and sodium carbonate (318 mg, 3.00 mmol) were added to a stirred solution of 3-[4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile (**C34**) (436 mg, 1.00 mmol) in ethanol (10 mL) and water (5 mL). The reaction mixture was heated at reflux for 2 hours, then concentrated *in vacuo*, poured into water (25 mL), and extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (25 mL), dried over sodium sulfate, filtered, and concentrated, affording the product as a white solid. Yield: 280 mg, 0.60 mmol, 60%. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.67 (s, 1H), 8.43 (s, 1H), 7.89 (br s, 1H), 7.72 (s, 1H), 7.64 (br d, *J*=7.8 Hz, 1H), 7.53 (br d, *J*=7.8 Hz, 1H), 7.46 (dd, *J*=7.6, 7.6 Hz, 1H), 5.89 (br s, ~1.6H), 5.61 (s, 2H), 3.53-3.61 (m, 2H), 3.39-3.47 (m, 4H), 3.13-3.21 (m, 4H), 0.81-0.88 (m, 2H), -0.09 (s, 9H).

Step 3. Synthesis of 4-(morpholin-4-yl)-5-[3-(1,2,4-oxadiazol-3-yl)phenyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**201**).

Triethyl orthoformate (760 mg, 5.1 mol) and boron trifluoride diethyl etherate (370 mg, 2.6 mol) were added to a 0 °C solution of *N'*-hydroxy-3-[4-(morpholin-4-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzenecarboximidamide (**C35**) (600 mg, 1.3 mmol) in tetrahydrofuran (25 mL). The reaction mixture was heated at 40 °C for 18 hours, then poured into water (50 mL) and extracted with ethyl acetate (3 x 25 mL). The combined organic layers were washed with saturated aqueous sodium chloride solution (50 mL), dried over sodium sulfate, filtered, and concentrated. Purification via preparative reversed phase high-performance liquid chromatography (Column: Phenomenex Gemini C18, 8 μm; Mobile phase A: aqueous ammonia, pH 10; Mobile phase B: acetonitrile; Gradient: 30% to 50% B) provided the product as a yellow solid. Yield: 62 mg, 0.18 mmol, 14%. LCMS *m/z* 349.0 [M+H⁺]. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.77 (s, 1H), 8.40 (s, 1H), 8.24 (br s, 1H), 7.98 (br d, *J*=7.5 Hz, 1H), 7.82 (br d, *J*=7.3 Hz, 1H), 7.63-7.70 (m, 2H), 3.39-3.45 (m, 4H), 3.13-3.20 (m, 4H).

Example 202

3-{4-[2-(3-Methyl-1,2-oxazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile (**202**)



5 Step 1. Synthesis of 4-benzyl-2-(3-methyl-1,2-oxazol-5-yl)morpholin-2-ol (**C36**).

A solution of 2-bromo-1-(3-methyl-1,2-oxazol-5-yl)ethanone (2.50 g, 12.2 mmol) and 2-(benzylamino)ethanol (3.70 g, 24.5 mmol) in acetonitrile (25 mL) was stirred at 35 °C for 18 hours. The reaction mixture was concentrated *in vacuo* and purified using silica gel chromatography (Gradient: 0% to 50% ethyl acetate in petroleum ether) to provide the product as a yellow oil. Yield: 1.2 g, 4.4 mmol, 36%. ¹H NMR (400 MHz, CDCl₃) δ 7.27-7.38 (m, 5H), 6.22 (s, 1H), 4.11-4.19 (m, 1H), 3.76-3.84 (m, 1H), 3.60 (AB quartet, *J*_{AB}=13.0 Hz, Δ*v*_{AB}=13.2 Hz, 2H), 2.99 (br d, *J*=11.0 Hz, 1H), 2.71-2.79 (m, 1H), 2.55 (d, *J*=11.0 Hz, 1H), 2.42 (ddd, *J*=11.7, 11.5, 3.6 Hz, 1H), 2.29 (s, 3H).

Step 2. Synthesis of 4-benzyl-2-(3-methyl-1,2-oxazol-5-yl)morpholine (**C37**).

15 Trimethylsilyl trifluoromethanesulfonate (2.9 g, 13 mmol) and triethylsilane (2.6 g, 22 mmol) were added to a solution of 4-benzyl-2-(3-methyl-1,2-oxazol-5-yl)morpholin-2-ol (**C36**) (1.2 g, 4.4 mmol) in dichloromethane (15 mL), and the reaction mixture was heated at 80 °C for 2 hours. The reaction mixture was then adjusted to a pH of ~5 to 6 via addition of saturated aqueous sodium bicarbonate solution. After extraction with dichloromethane (3 x 20 mL), the combined organic layers were washed with saturated aqueous sodium chloride solution (25 mL), dried over sodium sulfate, filtered, and concentrated. Purification via silica gel chromatography (Gradient: 0% to 50% ethyl acetate in petroleum ether) afforded the product as a yellow oil. Yield: 500 mg, 1.9

mmol, 43%. LCMS m/z 258.9 $[M+H]^+$. ^1H NMR (400 MHz, $CDCl_3$) δ 7.27-7.37 (m, 5H), 6.07 (s, 1H), 4.75 (dd, $J=9.7, 2.3$ Hz, 1H), 3.96 (ddd, $J=11.3, 2.8, 2.4$ Hz, 1H), 3.81 (ddd, $J=11.3, 10.8, 2.5$ Hz, 1H), 3.52-3.61 (m, 2H), 3.02 (br d, $J=11.4$ Hz, 1H), 2.68-2.75 (m, 1H), 2.29 (s, 3H), 2.27-2.37 (m, 2H).

5 Step 3. Synthesis of 2-(3-methyl-1,2-oxazol-5-yl)morpholine (**C38**).

Ammonium cerium(IV) nitrate (442 mg, 0.806 mmol) was added to a solution of 4-benzyl-2-(3-methyl-1,2-oxazol-5-yl)morpholine (**C37**) (100 mg, 0.39 mmol) in acetonitrile / water (5:1 mixture, 6 mL). The reaction mixture was stirred at 25 °C for 4 hours, then concentrated *in vacuo*; preparative thin-layer chromatography on silica gel
10 (Eluent: 10:1 dichloromethane / methanol) provided the product as a yellow oil. Yield: 40 mg, 0.24 mmol, 62%. ^1H NMR (400 MHz, $CDCl_3$) δ 6.11 (s, 1H), 4.79 (dd, $J=10.0, 2.4$ Hz, 1H), 4.01 (ddd, $J=11.8, 2.5, 2.4$ Hz, 1H), 3.81-3.89 (m, 1H), 3.35 (dd, $J=12.7, 2.5$ Hz, 1H), 2.99-3.07 (m, 3H), 2.30 (s, 3H).

Step 4. Synthesis of 3-{4-[2-(3-methyl-1,2-oxazol-5-yl)morpholin-4-yl]-7H-pyrrolo[2,3-
15 d]pyrimidin-5-yl}benzonitrile (**202**).


To a solution of 2-(3-methyl-1,2-oxazol-5-yl)morpholine (**C38**) (20 mg, 0.12 mmol) in *n*-butanol (7 mL) was added 3-(4-chloro-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile (**C14**) (36.2 mg, 0.142 mol), and the reaction mixture was placed in a sealed tube and heated at 150 °C in a microwave reactor for 2 hours. After removal of
20 solvent, the residue was purified by preparative reversed phase high-performance liquid chromatography (Column: YMC-Actus Triart C18, 5 μ m; Mobile phase A: water containing 0.225% formic acid; Mobile phase B: acetonitrile; Gradient: 33% to 53% B) to afford the product as a white solid. Yield: 0.78 mg, 2.0 μ mol, 2%. LCMS m/z 387.0 $[M+H]^+$. ^1H NMR (400 MHz, $DMSO-d_6$), characteristic peaks: δ 12.45 (br s, 1H), 8.45 (s, 1H), 8.00-8.03 (m, 1H), 7.85 (br d, $J=8$ Hz, 1H), 7.76 (br d, $J=8$ Hz, 1H), 7.72 (d, $J=2.8$ Hz, 1H), 7.66 (dd, $J=8, 8$ Hz, 1H), 6.25 (s, 1H), 4.58-4.63 (m, 1H), 3.71-3.83 (m, 2H), 3.01-3.09 (m, 1H), 2.88-2.97 (m, 1H), 2.20 (s, 3H).

Methods

30

Method A


Introduction of a 4-amino substituent onto 3-(4-chloro-7H-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile via chloride displacement



A solution of 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzonitrile (**C14**) in dimethyl sulfoxide (0.1 M, 500 μ L, 50 μ mol) was added to a vial containing the appropriate amine (150 μ mol). After addition of cesium fluoride (23 mg, 150 μ mol), the reaction mixture was shaken at 100 $^{\circ}$ C for 16 hours, then purified via preparative HPLC using an appropriate gradient with one of the following systems: 1) Column: DIKMA Diamonsil(2) C18, 5 μ m; Mobile phase A: 0.1% trifluoroacetic acid in water; Mobile phase B: acetonitrile; 2) Column: Boston Symmetrix ODS-H, 5 μ m or Phenomenex Synergi C18, 4 μ m or Agella Venusil ASB C18, 5 μ m; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile.

Method B

Synthesis of 5-substituted-4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidines via Suzuki reaction



Step 1. Suzuki reaction

A solution of the appropriate boronic ester in aqueous 1,4-dioxane (1:5 v/v water / 1,4-dioxane) (0.3 M, 400 μ L, 120 μ mol) was combined with a solution of 5-iodo-4-[(3S)-3-methylpiperidin-1-yl]-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine (**P5**) (0.25 M in 1:5 water / 1,4-dioxane; 400 μ L, 100 μ mol). The solution was treated with cesium carbonate (65 mg, 200 μ mol) and degassed by bubbling nitrogen through it. [1,1'-Bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II) (3

mg, 5 μ mol) was added to the reaction mixture, which was again degassed with nitrogen, then shaken at 150 °C for 4 hours. After removal of solvent using a Speedvac® concentrator, the products were purified via preparative thin layer chromatography in an appropriate solvent system and taken to the following step.


5 Step 2. Deprotection

The product from the previous step was treated with a solution of trifluoroacetic acid in dichloromethane (1:5 v/v; 2 mL), and the reaction mixture was shaken at 30 °C for 2 hours. After concentration using the Speedvac®, the residue was treated with a solution of ammonium hydroxide in methanol (1:4 v/v; 2 mL) and shaken at 30 °C for 16
10 hours. Solvents were removed on the Speedvac® and the product was purified via preparative HPLC using an appropriate gradient (Column: DIKMA Diamonsil(2) C18, 5 μ m or Agella Venusil ASB C18, 5 μ m; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile).

15

Method C

Synthesis of 5-substituted-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidines via Suzuki
reaction



Step 1. Suzuki reaction

20 A solution of 5-iodo-4-(morpholin-4-yl)-7-([2-(trimethylsilyl)ethoxy]methyl)-7H-pyrrolo[2,3-d]pyrimidine (**P1**) in 1,4-dioxane (0.20 M, 400 μ L, 80 μ mol) was combined with a solution of the appropriate boronic acid in 1,4-dioxane (0.50 M, 200 μ L, 100 μ mol). An aqueous solution of potassium phosphate (0.80 M, 200 μ L, 160 μ mol) was added, followed by [1,1'-bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II) (3
25 mg, 5 μ mol), and the reaction mixture was shaken at 100 °C for 16 hours. The mixture was filtered and the filtrate was concentrated on a Speedvac®. The residue was partitioned between dichloromethane and saturated aqueous sodium carbonate

solution, and the aqueous layer was extracted with dichloromethane (2 x 1 mL). The combined organic layers were dried over sodium sulfate, filtered, and concentrated using the Speedvac® to afford the crude product, which was used directly in the next step.

5 Step 2. Deprotection


The product of the preceding reaction was treated with a solution of concentrated hydrochloric acid in ethanol (1:6 v/v; 1 mL), and the reaction mixture was shaken at 80 °C for 16 hours. After removal of solvents on the Speedvac®, the residue was purified via preparative HPLC using an appropriate gradient with one of the following systems:

- 10 1) Column: Kromasil Eternity-5-C18, 5 µm or Boston Symmetrix ODS-H, 5 µm or Phenomenex Synergi C18, 4 µm; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile; 2) Column: Phenomenex Gemini C18, 8 µm; Mobile phase A: aqueous ammonium hydroxide, pH 10; Mobile phase B: acetonitrile.

15

Method D

Introduction of a 4-amino substituent onto 7*H*-pyrrolo[2,3-*d*]pyrimidines via chloride displacement




- A solution of the 4-chloro-7*H*-pyrrolo[2,3-*d*]pyrimidine substrate in dimethyl sulfoxide (0.5 M, 200 µL, 100 µmol) was added to a solution of the appropriate amine in dimethyl sulfoxide (0.8 M, 200 µL, 160 µmol). *N,N*-Diisopropylethylamine (120 µL, 700 µmol) was added, and the reaction mixture was shaken at 80 °C for 16 hours. After removal of solvents using a Speedvac®, the product was purified via preparative HPLC using an appropriate gradient with one of the following systems: 1) Column: Phenomenex Gemini C18, 8 µm or Agela Durashell C18, 5 µm; Mobile phase A: aqueous ammonium hydroxide, pH 10; Mobile phase B: acetonitrile; or 2) Column: DIKMA Diamonsil(2) C18, 5 µm or Phenomenex Synergi C18, 4 µm; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile.

30

Method E

Synthesis of 4-amino-substituted, 5-substituted-7*H*-pyrrolo[2,3-*d*]pyrimidines from 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine



Step 1. Amine displacement reaction

5 A solution of 4-chloro-5-iodo-7-[[2-(trimethylsilyl)ethoxy]methyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine (**C1**) in *N,N*-dimethylformamide (0.25 M, 500 μ L, 125 μ mol) was added to the appropriate amine (150 μ mol). Triethylamine (35 μ L, 250 μ mol) was added, and the reaction mixture was shaken at 100 $^{\circ}$ C for 16 hours. This solution was used directly in the following step.

10 Step 2. Suzuki reaction

The product solution from the preceding step (≤ 125 μ mol) was mixed with a solution of the appropriate boronic acid in *N,N*-dimethylformamide (0.25 M, 500 μ L, 125 μ mol). An aqueous solution of cesium carbonate (1.25 M, 200 μ L, 250 μ mol) was added, followed by [1,1'-bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II) (1.6 mg, 2.5 μ mol), and the reaction mixture was shaken at 120 $^{\circ}$ C for 4 hours. Removal of solvent using a Speedvac[®] provided a residue, which was used directly in the following step.


Step 3. Deprotection

20 The product from the previous step (≤ 125 μ mol) was treated with a solution of concentrated hydrochloric acid in ethanol (1:6 v/v; 2 mL), and the reaction mixture was shaken at 80 $^{\circ}$ C for 16 hours. After removal of solvent via the Speedvac[®], the residue was taken up in a solution of ammonium hydroxide (30% aqueous) in methanol (1:4 v/v, 2 mL). The mixture was shaken at 30 $^{\circ}$ C for 16 hours, concentrated on the Speedvac[®] and purified via preparative HPLC using an appropriate gradient (Column: DIKMA

Diamonsil(2) C18, 5 μ m or Agella Venusil ASB C18, 5 μ m; Mobile phase A: 0.225% formic acid in water; Mobile phase B: acetonitrile).

Method F

- 5 Introduction of a 5-substituent onto 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidin-4-amines via Suzuki reaction



Step 1. Suzuki reaction

- A solution of the 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7-[[2-(trimethylsilyl)ethoxy]methyl]-7H-pyrrolo[2,3-d]pyrimidine substrate in 1,4-dioxane (0.1 M, 1 mL, 100 μ mol) was added to the appropriate aryl or heteroaryl halide (100 μ mol). An aqueous solution of cesium carbonate (1.0 M, 200 μ L, 200 μ mol) was added, followed by [1,1'-bis(di-*tert*-butylphosphino)ferrocene]dichloropalladium(II) (1.3 mg, 2.0 μ mol), and the reaction mixture was shaken at 120 $^{\circ}$ C for 4 hours. Removal of solvent using a Speedvac[®] provided a residue, which was used directly in the following step.




Step 2. Deprotection





Protecting group removal and purification were carried out as described in the final step of Method E.







- The following table (Table 1) provides Examples 11-197 and Examples 203-225 of the present invention, the structure of the compound, reference to the method of preparation and characterizing data for the compound. Certain of the compounds depicted in the table are racemic and others are depicted as single enantiomers with the absolute stereochemistry as shown. In certain cases the racemic compound has been separated into the individual enantiomers, although the absolute stereochemistry of the single enantiomer(s) may not have been determined. In the table the individual separated enantiomers may be referred to as ENT-1 or ENT-2, which are abbreviations for the separated enantiomer-1 and enantiomer-2, respectively. It is to be understood






that one of the compounds designated as ENT-1 or ENT-2 will have the (*R*) absolute stereochemistry at the chiral center while the other will have the (*S*) absolute stereochemistry.







Table 1








Example #	Structure	Method of Preparation; Non-commercial starting materials	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ (ppm); Mass spectrum, observed ion <i>m/z</i> (M+H ⁺) or HPLC retention time; Mass spectrum <i>m/z</i> (M+H ⁺) (unless otherwise indicated)
11		Ex 6	12.20 (br s, 1H), 8.32 (s, 1H), 7.90 (dd, <i>J</i> =1.8, 1.5 Hz, 1H), 7.83 (ddd, <i>J</i> =8.0, 1.5, 1.3 Hz, 1H), 7.76 (ddd, <i>J</i> =8, 1.5, 1 Hz, 1H), 7.65 (dd, <i>J</i> =7.8, 7.5 Hz, 1H), 7.59 (d, <i>J</i> =2.3 Hz, 1H), 2.96-3.03 (m, 4H), 1.20-1.27 (m, 2H), 1.10-1.19 (m, 2H), 0.87 (s, 6H); 331.9
12		Ex 6	12.21 (br s, 1H), 8.35 (s, 1H), 7.95 (br s, 1H), 7.85 (br d, <i>J</i> =7.5 Hz, 1H), 7.74 (br d, <i>J</i> =7.5 Hz, 1H), 7.61-7.67 (m, 2H), 3.13-3.19 (m, 4H), 1.40-1.48 (m, 2H), 1.29-1.38 (m, 4H); 303.9
13		Ex 6	12.05 (br s, 1H), 8.31 (s, 1H), 7.37-7.50 (m, 5H), 7.24-7.31 (m, 1H), 7.08 (br s, 1H), 6.73 (br s, 1H), 3.93 (br d, <i>J</i> =12.4 Hz, 1H), 3.65 (br d, <i>J</i> =12.4 Hz, 1H), 2.74 (dd, <i>J</i> =12.4, 11.7 Hz, 1H), 2.5-2.58 (m, 1H, assumed; partially obscured by solvent peak), 2.21-2.31 (m, 1H), 1.68-1.78 (m, 1H), 1.26-1.44 (m, 2H), 1.15-1.26 (m, 1H), 322.2







14		Ex 5	¹ H NMR (400 MHz, CDCl ₃), characteristic peaks: δ 10.44 (br s, 1H), 8.46 (s, 1H), 7.82 (dd, J=1.5, 1.3 Hz, 1H), 7.75 (ddd, J=7.7, 1.5, 1.5 Hz, 1H), 7.62 (ddd, J=7.8, 1.5, 1.3 Hz, 1H), 7.54 (dd, J=8.0, 7.5 Hz, 1H), 7.28 (d, J=2.5 Hz, 1H), 4.83 (br s, 1H), 4.14 (br d, J=14 Hz, 1H), 3.95-4.00 (m, 1H), 3.45 (dd, J=13.9, 2.4 Hz, 1H), 3.13-3.21 (m, 1H), 2.82-2.91 (m, 1H), 1.13-1.22 (m, 1H); 320.2
15		Ex 7 ¹	¹ H NMR (400 MHz, CD ₃ OD) δ 8.25 (s, 1H), 7.38-7.49 (m, 4H), 7.27-7.33 (m, 1H), 7.21 (s, 1H), 3.91 (br d, J=13 Hz, 1H), 3.78 (br d, J=13 Hz, 1H), 2.63 (ddd, J=12.6, 12.6, 3.0 Hz, 1H), 2.25 (dd, J=12.6, 11.0 Hz, 1H), 1.62-1.70 (m, 1H), 1.31-1.52 (m, 3H), 0.88-1.00 (m, 1H), 0.59 (d, J=6.5 Hz, 3H); 293.3
16		Ex 6	¹ H NMR (400 MHz, CDCl ₃) δ 10.55 (br s, 1H), 8.49 (s, 1H), 7.83 (br s, 1H), 7.75 (br d, J=7.5 Hz, 1H), 7.61 (br d, J=7.5 Hz, 1H), 7.54 (dd, J=8.0, 7.5 Hz, 1H), 7.23 (s, 1H), 3.84 (br d, J=13 Hz, 1H), 3.77 (br d, J=13 Hz, 1H), 2.61-2.70 (m, 1H), 2.34 (dd, J=12.0, 11.5 Hz, 1H), 1.65-1.77 (m, 1H), 1.29-1.55 (m, 3H), 0.91-1.03 (m, 1H), 0.68 (d, J=6.5 Hz, 3H); 318.3
17		Ex 7 ²	¹ H NMR (400 MHz, CD ₃ OD) δ 8.23 (s, 1H), 7.71 (s, 1H), 7.57 (s, 1H), 7.19 (s, 1H), 3.95 (s, 3H), 3.88-3.95 (m, 1H), 3.83 (br d, J=12 Hz, 1H), 2.66-2.74 (m, 1H), 2.33 (dd, J=12, 11 Hz, 1H), 1.71-1.79 (m, 1H), 1.40-1.62 (m, 3H), 0.96-1.08 (m, 1H), 0.72 (d, J=6.8 Hz, 3H), 297.6








18		Ex 6	3.10 minutes ³ ; 265.1
19		Ex 6	12.02 (br s, 1H), 8.29 (s, 1H), 7.45-7.51 (m, 2H), 7.37-7.45 (m, 3H), 7.24-7.30 (m, 1H), 4.39 (t, <i>J</i> =5.1 Hz, 1H), 3.82 (br d, <i>J</i> =12.4 Hz, 2H), 3.11-3.18 (m, 2H), 2.61 (br dd, <i>J</i> =12, 11 Hz, 2H), 1.35-1.46 (m, 3H), 0.88-1.03 (m, 2H); 309.1
20		Ex 7	¹ H NMR (400 MHz, CD ₃ OD) δ 8.34 (s, 1H), 7.50 (br s, 1H), 7.43-7.46 (m, 2H), 7.32-7.35 (m, 1H), 7.30 (s, 1H), 4.70 (s, 2H), 3.64-3.71 (m, 1H), 3.46-3.54 (m, 1H), 3.21-3.27 (m, 1H), 3.09-3.18 (m, 1H), 2.85-2.92 (m, 1H), 1.82-1.91 (m, 1H), 1.67-1.77 (m, 1H), 1.41-1.51 (m, 1H), 1.19-1.31 (m, 1H); 334.3
21		Ex 7	¹ H NMR (400 MHz, CD ₃ OD) δ 8.39 (s, 1H), 7.92 (br s, 1H), 7.87 (br d, <i>J</i> =7 Hz, 1H), 7.63-7.72 (m, 2H), 7.47 (s, 1H), 3.58-3.70 (m, 2H), 3.06-3.19 (m, 2H), 2.96-3.04 (m, 1H), 1.74-1.92 (m, 2H), 1.38-1.50 (m, 1H), 1.17-1.30 (m, 1H); 329.4
22		Ex 7 ²	¹ H NMR (400 MHz, CD ₃ OD) δ 8.23 (s, 1H), 7.69 (s, 1H), 7.55 (s, 1H), 7.18 (s, 1H), 3.94 (s, 3H), 3.89 (br d, <i>J</i> =12 Hz, 2H), 2.20 (dd, <i>J</i> =12.0, 11.5 Hz, 2H), 1.73 (br d, <i>J</i> =12.6 Hz, 1H), 1.48-1.62 (m, 2H), 0.74 (d, <i>J</i> =7.0 Hz, 6H), 0.65 (dt, <i>J</i> =12, 12 Hz, 1H); 311.5
23		Ex 3 ^{4,5,6} ; P4	¹ H NMR (300 MHz, DMSO- <i>d</i> ₆) characteristic peaks: δ 12.01 (br s, 1H), 8.26 (s, 1H), 7.81 (dd, <i>J</i> =8.4, 2.1 Hz, 1H), 7.68 (d, <i>J</i> =2.1 Hz, 1H), 7.36 (d, <i>J</i> =2.1 Hz, 1H), 7.26 (d, <i>J</i> =9.1 Hz, 1H), 3.82 (s, 3H), 3.74-3.84 (m, 1H), 3.57-3.66 (m, 1H), 2.17 (dd, <i>J</i> =12.5, 11.2 Hz,







			1H), 1.56-1.66 (m, 1H), 1.38-1.50 (m, 1H), 1.05-1.24 (m, 2H), 0.80-0.96 (m, 1H), 0.55 (d, $J=6.3$ Hz, 3H); 348.2
24		Ex 1 ^{1.7}	8.17 (s, 1H), 7.80 (dd, $J=8.5, 2.5$ Hz, 1H), 7.70 (d, $J=2.0$ Hz, 1H), 7.30 (s, 1H), 7.22 (d, $J=9.0$ Hz, 1H), 3.80 (s, 3H), 2.73 (s, 6H); 293.8
25		Ex 7 ^{1.8}	11.88 (br s, 1H), 8.16 (s, 1H), 7.30-7.35 (m, 2H), 7.25 (d, $J=2.6$ Hz, 1H), 7.06 (d, $J=8.5$ Hz, 1H), 3.71 (s, 3H), 2.75 (s, 6H); 303.2
26	 •CF ₃ COOH	Method A	2.07 minutes ⁹ ; 370
27		Method A	2.58 minutes ⁹ ; 348
28		Ex 6	12.29 (br s, 1H), 8.37 (s, 1H), 7.92 (dd, $J=1.5, 1.5$ Hz, 1H), 7.86 (ddd, $J=7.5, 1.5, 1.5$ Hz, 1H), 7.77 (ddd, $J=7.5, 1.5, 1.5$ Hz, 1H), 7.62-7.68 (m, 2H), 3.23-3.28 (m, 2H), 3.19 (s, 2H), 2.95-3.00 (m, 2H), 2.25-2.32 (m, 2H), 2.14-2.23 (m, 2H), 2.13 (s, 3H), 1.74-1.82 (m, 2H), 1.45-1.54 (m, 2H); 389.2






29		Method A	2.46 minutes ⁹ ; 334
30		Method A	2.42 minutes ⁹ ; 350
31		Method A	2.23 minutes ⁹ ; 335
32		Method A	2.78 minutes ⁹ ; 340
33		Method A	2.27 minutes ¹⁰ ; 322
33A		Ex 6	¹ H NMR (400 MHz, CDCl ₃) δ 10.65 (br s, 1H), 8.51 (s, 1H), 7.84 (s, 1H), 7.76 (br d, J=7.5 Hz, 1H), 7.62 (br d, J=8 Hz, 1H), 7.55 (br dd, J=8, 7.5 Hz, 1H), 7.24-7.30 (1H, assumed, obscured by solvent peak), 4.75 (br d, J _{HF} =48 Hz, 1H), 3.32-3.51 (m, 4H), 1.65-1.84 (m, 4H, assumed; partially obscured by water peak); 322.2






34		Method A	2.33 minutes ⁹ ; 370
35		Method A	2.33 minutes ⁹ ; 308
36		Method A	2.33 minutes ⁹ ; 372
37		Method A	2.42 minutes ⁹ ; 370
38		Method A	2.15 minutes ⁹ ; 370
39		Method A	2.08 minutes ⁹ ; 333
40		Method A	2.27 minutes ⁹ ; 373






41	 ENT-2	Ex 6 ^{11,12}	¹ H NMR (400 MHz, CD ₃ OD) δ 8.30 (s, 1H), 7.84 (br s, 1H), 7.81 (br d, J =7.5 Hz, 1H), 7.66 (br d, J =7.5 Hz, 1H), 7.61 (dd, J =8.0, 7.5 Hz, 1H), 7.39 (s, 1H), 3.81 (br d, J =13.0 Hz, 1H), 3.70 (br d, J =12.6 Hz, 1H), 2.65 (ddd, J =12.6, 12.3, 2.8 Hz, 1H), 2.29 (dd, J =12.6, 11.0 Hz, 1H), 1.65-1.74 (m 1H), 1.47-1.56 (m, 1H), 1.31-1.46 (m, 2H), 0.91-1.04 (m, 1H), 0.64 (d, J =6.5 Hz, 3H); 318.0
42		Method A	1.99 minutes ⁹ ; 349
43		Method A	2.47 minutes ⁹ ; 360
44		Method A	2.13 minutes ⁹ ; 361
45		Method B	2.06 minutes ¹⁰ ; 333
46		Method B	2.32 minutes ⁹ ; 372






47		Method B	2.53 minutes ¹⁰ ; 311
48		Method B	2.15 minutes ¹⁰ ; 333
49		Method B	2.39 minutes ¹⁰ ; 326
50		Method B	1.86 minutes ¹⁰ ; 348
51		Method B	2.04 minutes ⁹ ; 294
52		Method B	1.69 minutes ¹⁰ ; 308
53		Method B	2.02 minutes ¹⁰ ; 333








54		Method B	1.96 minutes ¹⁰ ; 348
55		Method B	1.11 minutes ¹⁰ ; 333
56		Method B	2.11 minutes ¹⁰ ; 309
57		Method B	2.17 minutes ⁹ ; 336
58		Method B	2.66 minutes ¹³ ; 309
59		Method B	2.31 minutes ¹⁰ ; 342








60		Ex 1 ^{7,14} ; P5	characteristic peaks, 12.09 (br s, 1H), 8.31 (s, 1H), 7.99-8.03 (m, 2H), 7.79 (br d, $J=7.5$ Hz, 1H), 7.62 (br d, $J=7.5$ Hz, 1H), 7.47-7.52 (m, 2H), 7.36 (br s, 1H), 3.75 (br d, $J=13$ Hz, 1H), 3.68 (br d, $J=12$ Hz, 1H), 2.21 (dd, $J=12, 11$ Hz, 1H), 1.54-1.61 (m, 1H), 1.33-1.43 (m, 2H), 1.19-1.32 (m, 1H), 0.81-0.95 (m, 1H), 0.54 (d, $J=7.0$ Hz, 3H); 336.2
61		Ex 6	8.35 (s, 1H), 7.92 (br s, 1H), 7.82 (br d, $J=8$ Hz, 1H), 7.74 (br d, $J=7.5$ Hz, 1H), 7.60-7.66 (m, 2H), 4.60 (br d, $J_{\text{HF}}=47$ Hz, 1H), 3.26-3.4 (m, 2H, assumed; partially obscured by water peak), 3.16-3.25 (m, 1H), 3.01-3.10 (m, 1H), 1.71-1.87 (m, 1H), 1.59-1.70 (m, 1H), 1.48-1.58 (m, 1H), 1.11-1.21 (m, 1H); 322.2
62		Ex 6	12.83 (br s, 1H), 8.44 (s, 1H), 7.96 (br s, 1H), 7.83 (br d, $J=8$ Hz, 1H), 7.80 (br d, $J=8$ Hz, 1H), 7.75 (d, $J=2.5$ Hz, 1H), 7.67 (dd, $J=8.0, 7.5$ Hz, 1H), 3.75 (br d, $J=13$ Hz, 2H), 2.84 (br dd, $J=13, 12$ Hz, 2H), 1.42-1.55 (m, 3H), 0.91-1.03 (m, 2H), 0.83 (d, $J=6.0$ Hz, 3H); 318.2
63		Ex 1 ^{7,14} ; P5	¹ H NMR (400 MHz, CDCl ₃), characteristic peaks: δ 8.42 (s, 1H), 8.28-8.31 (m, 1H), 8.09 (br d, $J=7$ Hz, 1H), 7.69 (br d, $J=7$ Hz, 1H), 7.52-7.58 (m, 1H), 3.90-4.03 (m, 2H), 2.63-2.73 (m, 1H), 2.32-2.41 (m, 1H), 0.65 (d, $J=6$ Hz, 3H); 337.0
64		Ex 6	11.94 (br s, 1H), 8.22 (s, 1H), 7.86 (dd, $J=1.5, 1.5$ Hz, 1H), 7.73-7.78 (m, 2H), 7.64 (dd, $J=7.8, 7.5$ Hz, 1H), 7.35 (s, 1H), 5.76 (q, $J=4.6$ Hz, 1H), 2.90 (d, $J=4.8$ Hz, 3H); 250.1








65		Ex 4	¹ H NMR (400 MHz, CD ₃ OD) δ 8.24 (s, 1H), 7.72 (dd, <i>J</i> =8.7, 2.1 Hz, 1H), 7.65 (d, <i>J</i> =2.0 Hz, 1H), 7.28 (s, 1H), 7.23 (d, <i>J</i> =8.8 Hz, 1H), 3.87 (s, 3H), 3.83 (br d, <i>J</i> =12 Hz, 2H), 2.13 (dd, <i>J</i> =12.0, 12.0 Hz, 2H), 1.68 (br d, <i>J</i> =13 Hz, 1H), 1.25-1.37 (m, 2H), 0.68 (d, <i>J</i> =6.5 Hz, 6H), 0.60 (dt, <i>J</i> =12, 12 Hz, 1H); 362.1
66		Ex 1 ^{7,14} ; P5	¹ H NMR (400 MHz, CDCl ₃) δ 10.70 (br s, 1H), 8.42 (s, 1H), 7.61-7.67 (m, 2H), 7.25-7.29 (m, 1H, assumed; partially obscured by solvent peak), 7.03 (d, <i>J</i> =8.5 Hz, 1H), 3.89 (s, 3H), 3.87-3.96 (m, 1H), 3.81 (br d, <i>J</i> =13 Hz, 1H), 2.61 (ddd, <i>J</i> =13, 13, 3 Hz, 1H), 2.29 (dd, <i>J</i> =12.6, 11.0 Hz, 1H), 1.66-1.75 (m, 1H), 1.45-1.54 (m 1H), 1.18-1.37 (m, 2H), 0.89-1.00 (m, 1H), 0.65 (d, <i>J</i> =7.0 Hz, 3H); 348.2
67		Ex 1 ^{7,14} ; P5	¹ H NMR (400 MHz, CDCl ₃), characteristic peaks: δ 10.36 (br s, 1H), 8.45 (s, 1H), 7.70 (ddd, <i>J</i> =7.5, 7.5, 1.8 Hz, 1H), 7.59 (ddd, <i>J</i> =7.8, 5.5, 1.8 Hz, 1H), 7.31-7.37 (m, 2H), 3.76-3.88 (m, 2H), 2.66 (ddd, <i>J</i> =13, 12, 2.5 Hz, 1H), 2.36 (dd, <i>J</i> =12.7, 11.2 Hz, 1H), 1.46-1.54 (m, 1H), 1.35-1.45 (m, 1H), 1.22-1.34 (m, 1H), 0.92-1.03 (m, 1H), 0.69 (d, <i>J</i> =6.5 Hz, 3H), 336.2
68		Ex 1 ^{1,7} ; C3	¹ H NMR (400 MHz, CDCl ₃) δ 8.39 (s, 1H), 7.48 (br d, <i>J</i> =8 Hz, 2H), 7.41 (br dd, <i>J</i> =8, 8 Hz, 2H), 7.30-7.34 (m, 1H), 7.09 (s, 1H), 2.84 (s, 6H); 239.1
69		Ex 1 ^{1,7} ; C3	¹ H NMR (400 MHz, CDCl ₃) δ 12.24 (br s, 1H), 8.25 (s, 1H), 7.84-7.91 (m, 1H), 7.75-7.82 (m, 1H), 7.51 (br s, 1H), 7.47 (dd, <i>J</i> =8.0, 7.5 Hz, 1H), 2.73 (s, 6H); 282.2








70		Ex 5	8.35 (s, 1H), 7.92 (s, 1H), 7.85 (d, $J=7.5$ Hz, 1H), 7.76 (d, $J=7.5$ Hz, 1H), 7.63 (dd, $J=8.0$, 7.5 Hz, 1H), 7.57 (br s, 1H), 3.19-3.3 (br m, 4H), 1.70-1.79 (br m, 4H); 290.1
71		Ex 5 ^{15,12}	¹ H NMR (400 MHz, CD ₃ OD) δ 8.15 (s, 1H), 7.78-7.80 (m, 1H), 7.73 (ddd, $J=7.8$, 1.6, 1.3 Hz, 1H), 7.69 (ddd, $J=7.8$, 1.5, 1.2 Hz, 1H), 7.58 (dd, $J=7.8$, 7.8 Hz, 1H), 7.19 (s, 1H), 3.36-3.44 (m, 2H), 3.21-3.30 (m, 1H), 2.75 (dd, $J=10.4$, 8.2 Hz, 1H), 2.05-2.15 (m, 1H), 1.82-1.91 (m, 1H), 1.30-1.41 (m, 1H), 0.90 (d, $J=6.5$ Hz, 3H); 304.1
72		Ex 1 ^{1,7} ; C3	¹ H NMR (400 MHz, CD ₃ OD) δ 8.23 (s, 1H), 7.87 (dd, $J=6.9$, 2.1 Hz, 1H), 7.76 (ddd, $J=8.5$, 4.5, 2.3 Hz, 1H), 7.37-7.42 (m, 2H), 2.85 (s, 6H); 282.1
73		Ex 6	¹ H NMR (400 MHz, DMSO- d_6 + D ₂ O) δ 8.29 (s, 1H) 7.88-7.90 (m, 1H), 7.84 (br d, $J=8.5$ Hz, 1H), 7.71 (br d, $J=8.0$ Hz, 1H), 7.61 (dd, $J=8.0$, 7.5 Hz, 1H), 7.56 (s, 1H), 3.20 (q, $J=7.0$ Hz, 4H), 0.88 (t, $J=7.0$ Hz, 6H); 292.0
74		Ex 6 ^{16,17}	¹ H NMR (400 MHz, CDCl ₃) δ 11.17 (br s, 1H), 8.52 (s, 1H), 7.83 (br s, 1H), 7.76 (br d, $J=8.0$ Hz, 1H), 7.64 (br d, $J=7.5$ Hz, 1H), 7.57 (dd, $J=8.0$, 7.5 Hz, 1H), 7.29 (s, 1H), 3.73-3.84 (m, 2H), 3.63 (br d, $J=13.6$ Hz, 1H), 3.51-3.59 (m, 1H), 3.47 (ddd, $J=12.0$, 11.5, 2.5 Hz, 1H), 3.30 (s, 3H), 3.20-3.30 (m, 2H), 2.96 (ddd, $J=13$, 12, 3.5 Hz, 1H), 2.79 (dd, $J=12.8$, 10.8 Hz, 1H); 350.2






75		Ex 1 ^{1,7} ; C3	8.31 (s, 1H), 7.81-7.83 (m, 1H), 7.71-7.76 (m, 1H), 7.67 (s, 1H), 7.64-7.69 (m, 1H), 2.76 (s, 6H); 282.1
76		Ex 6 ^{18,12}	¹ H NMR (400 MHz, CDCl ₃), characteristic peaks: δ 8.51 (br s, 1H), 7.84 (s, 1H), 7.77 (br d, J=7.5 Hz, 1H), 7.62 (br d, J=8 Hz, 1H), 7.56 (dd, J=8.0, 7.5 Hz, 1H), 7.51 (br s, 1H), 6.06 (br s, 1H), 4.55 (br d, J=10 Hz, 1H), 3.99 (br d, J=13 Hz, 1H), 3.84-3.91 (m, 1H), 3.60-3.75 (m, 2H), 3.02-3.13 (m, 2H); 372.1
77		Ex 5 ^{15,17}	¹ H NMR (400 MHz, CD ₃ OD) δ 8.19 (s, 1H), 7.82 (br s, 1H), 7.71-7.77 (m, 2H), 7.61 (dd, J=7.5, 7.5 Hz, 1H), 7.29 (s, 1H), 3.40-3.49 (m, 2H), 3.24-3.3 (m, 1H, assumed; partially obscured by solvent peak), 2.76 (dd, J=9.5, 9.0 Hz, 1H), 2.10-2.22 (m, 1H), 1.88-1.98 (m, 1H), 1.36-1.48 (m, 1H), 0.91 (d, J=6.5 Hz, 3H), 304.0
78		Ex 1 ⁷	8.38 (s, 1H), 7.80-7.90 (m, 2H), 7.64 (d, J=1.0 Hz, 1H), 7.52 (dd, J=8.0, 7.5 Hz, 1H), 3.35-3.41 (m, 4H), 3.09-3.15 (m, 4H); 323.9
79		Ex 6	¹ H NMR (400 MHz, CDCl ₃) δ 11.02 (br s, 1H), 8.53 (s, 1H), 7.85 (s, 1H), 7.78 (d, J=7.0 Hz, 1H), 7.53-7.68 (m, 2H), 7.30 (s, 1H), 3.57-3.77 (m, 3H), 3.39-3.54 (m, 2H), 2.92 (br dd, J=12, 12 Hz, 1H), 2.59 (dd, J=11.5, 11.0 Hz, 1H), 0.98 (d, J=6.0 Hz, 3H); 319.9







80		Ex 1 ⁷	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆ + D ₂ O) δ 8.35 (s, 1H), 7.97 (dd, <i>J</i> =7.0, 2.0 Hz, 1H), 7.81-7.87 (m, 1H), 7.61 (s, 1H), 7.53 (dd, <i>J</i> =9.0, 9.0 Hz, 1H), 3.36-3.42 (m, 4H), 3.08-3.15 (m, 4H); 324.2
81		Ex 4 ^{2,4}	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆ + D ₂ O) δ 8.36 (s, 1H), 7.81 (s, 1H), 7.54 (s, 1H), 7.46 (s, 1H), 4.84 (br d, <i>J</i> _{HF} =48 Hz, 1H), 3.85 (s, 3H), 3.43-3.58 (m, 4H), 1.58-1.88 (m, 4H); 301.1
82		Method C	2.27 minutes ⁹ ; 306
83		Method C	2.39 minutes ¹⁰ ; 315
84		Method C	2.39 minutes ⁹ ; 299
85		Method C	2.54 minutes ⁹ ; 329
86		Method C	2.46 minutes ⁹ ; 317







87		Method C	2.47 minutes ⁹ ; 317
88		Method C	2.38 minutes ¹⁰ ; 333
89		Method C	2.07 minutes ⁹ ; 329
90		Method C	2.46 minutes ⁹ ; 317
91		Method C	2.46 minutes ⁹ ; 299
92		Method C	2.57 minutes ⁹ ; 317
93		Method C	2.40 minutes ⁹ ; 281








94		Method C	2.41 minutes ⁹ ; 329
95		Method C	2.46 minutes ⁹ ; 315
96		Method C	2.51 minutes ⁹ ; 313
97		Method C	2.42 minutes ⁹ ; 311
98		Method C	2.18 minutes ⁹ ; 329
99		Method C	2.44 minutes ⁹ ; 299
100		Method C	2.11 minutes ⁹ ; 311







101		Method C	2.36 minutes ⁹ ; 311
102		Method C	2.34 minutes ⁹ ; 327
103		Method C	1.44 minutes ⁹ ; 282
104		Method C	1.96 minutes ⁹ ; 332
105		Method C	2.34 minutes ⁹ ; 360
106		Method C	2.14 minutes ¹⁰ ; 329
107		Method C	2.30 minutes ⁹ ; 295








108		Ex 3 ¹⁹	¹ H NMR (400 MHz, CDCl ₃) δ 9.70 (br s, 1H), 8.45 (s, 1H), 7.43 (d, <i>J</i> =2.0 Hz, 1H), 7.33 (d, <i>J</i> =2.0 Hz, 1H), 6.43 (d, <i>J</i> =2.0 Hz, 1H), 3.99 (s, 3H), 3.66 (dd, <i>J</i> =4.8, 4.8 Hz, 4H), 3.42 (dd, <i>J</i> =4.9, 4.4 Hz, 4H); 285.2
109		Ex 5	¹ H NMR (400 MHz, CDCl ₃), characteristic peaks: δ 10.64 (br s, 1H), 8.50 (s, 1H), 7.84-7.86 (m, 1H), 7.77 (ddd, <i>J</i> =7.7, 1.8, 1.4 Hz, 1H), 7.61 (ddd, half of ABXY pattern, <i>J</i> =7.8, 1.5, 1.5 Hz, 1H), 7.55 (dd, half of ABX pattern, <i>J</i> =7.8, 7.8 Hz, 1H), 3.77-3.85 (m, 1H), 3.67-3.76 (m, 2H), 3.03 (ddd, <i>J</i> =13.2, 9.8, 3.0 Hz, 2H), 1.71-1.80 (m, 2H), 1.35-1.46 (m, 2H); 319.8
110		Ex 5	characteristic peaks: 12.35 (br s, 1H), 8.43 (s, 1H), 8.01 (br s, 1H), 7.69-7.81 (m, 3H), 7.59 (dd, <i>J</i> =7.5, 7.5 Hz, 1H), 4.94 (br d, <i>J</i> =8 Hz, 1H), 3.91 (br d, <i>J</i> =13.5 Hz, 1H), 3.62-3.70 (m, 1H), 3.43-3.51 (m, 1H), 2.88-2.97 (m, 1H), 2.35 (s, 3H); 388.2
111		Ex 3	8.39 (s, 1H), 7.81 (br s, 1H), 7.71 (br s, 1H), 7.68 (s, 1H), 7.57 (br s, 1H), 3.46 (dd, <i>J</i> =4.5, 4.5 Hz, 4H), 3.10-3.16 (m, 4H), 2.43 (s, 3H); 320.2
112		Ex 5 ²⁰	characteristic peaks: 8.43 (s, 1H), 8.00 (br s, 1H), 7.85 (br d, <i>J</i> =8.0 Hz, 1H), 7.75 (br d, <i>J</i> =7.5 Hz, 1H), 7.71 (s, 1H), 7.65 (dd, <i>J</i> =7.5, 7.5 Hz, 1H), 4.65-4.70 (m, 1H), 3.82-3.88 (m, 1H), 3.68-3.75 (m, 1H), 3.43-3.51 (m, 1H), 3.19 (dd, <i>J</i> =13, 10 Hz, 1H), 2.85-2.94 (m, 1H), 2.57 (s, 3H); 388.2








113		Ex 5	12.21 (br s, 1H), 8.32 (s, 1H), 7.92-7.94 (m, 1H), 7.81 (br d, $J=8$ Hz, 1H), 7.75 (br d, $J=8$ Hz, 1H), 7.60 (dd, $J=8.0, 7.5$ Hz, 1H), 7.55 (d, $J=2.5$ Hz, 1H), 4.36 (br d, $J=6.0$ Hz, 2H), 3.72 (d, $J=12.6$ Hz, 2H), 3.43 (d, $J=12.6$ Hz, 2H), 2.91-2.98 (m, 1H), 1.89 (d, $J=8.5$ Hz, 1H); 318.1
114		Ex 3	8.41 (s, 1H), 8.04 (br s, 1H), 7.97 (br s, 1H), 7.90 (br s, 1H), 7.83 (s, 1H), 3.49-3.56 (m, 4H), 3.10-3.17 (m, 4H); 340.1
115		Ex 3	8.33 (s, 1H), 7.82 (dd, $J=8.5, 2.0$ Hz, 1H), 7.73 (d, $J=2.0$ Hz, 1H), 7.48 (s, 1H), 7.28 (d, $J=9.0$ Hz, 1H), 3.85 (s, 3H), 3.3-3.38 (m, 4H, assumed; partially obscured by water peak), 3.08-3.15 (m, 4H); 336.2
116		Ex 3	^1H NMR (400 MHz, DMSO- d_6 + D $_2$ O) δ 8.29 (s, 1H), 7.41 (s, 1H), 7.33 (dd, half of ABX pattern, $J=8.5, 2.8$ Hz, 1H), 7.30 (d, half of AB quartet, $J=2.5$ Hz, 1H), 7.10 (d, $J=8.5$ Hz, 1H), 3.74 (s, 3H), 3.33-3.39 (m, 4H), 3.08-3.14 (m, 4H); 345.2
117		Ex 2 ²	11.89 (br s, 1H), 8.25 (s, 1H), 7.74 (s, 1H), 7.48 (s, 1H), 3.90 (s, 3H), 3.43-3.49 (m, 4H), 3.08-3.15 (m, 4H), 2.31 (s, 3H); 299.1
118		Ex 3	^1H NMR (400 MHz, DMSO- d_6 + D $_2$ O) δ 8.36 (s, 1H), 7.67 (s, 1H), 7.54-7.57 (m, 1H), 7.39-7.42 (m, 1H), 7.30-7.33 (m, 1H), 3.86 (s, 3H), 3.43-3.50 (m, 4H), 3.10-3.16 (m, 4H); 335.9







119		Method D ²¹ ; P6	2.06 minutes ⁹ ; 301
120		Method D ²¹ ; P6	1.90 minutes ¹³ ; 299
121		Method D ²¹ ; P6	1.96 minutes ⁹ ; 299
122		Method D ²¹ ; P6	1.96 minutes ⁹ ; 299
123		Method D ²¹ ; P6	2.02 minutes ⁹ ; 301
124		Method D ²¹ ; P6	1.80 minutes ¹³ ; 315
125		Method D ²¹ ; P6 ²²	1.89 minutes ⁹ ; 406







126		Method D ²¹ ; P6	1.76 minutes ⁹ ; 243
127		Method D ²¹ ; P6	1.92 minutes ⁹ ; 269
128		Method D; C14	2.49 minutes ¹³ ; 340
129		Method D; C14	2.05 minutes ¹³ ; 319
130		Method D; C14 ²³	2.43 minutes ¹³ ; 438
131		Method D; C14	2.39 minutes ¹³ ; 389
132		Method D; C14	2.13 minutes ¹³ ; 318








133		Method D; C14	2.03 minutes ⁹ ; 352
134		Method D; C14	2.36 minutes ⁹ ; 398
135		Method D; C14 ²⁴	2.49 minutes ⁹ ; 414
136		Method D; C14	2.08 minutes ¹⁰ ; 326
137		Method D; C14 ²⁵	2.43 minutes ¹⁰ ; 437
138		Method D; C14	2.00 minutes ⁹ ; 320








139		Method D; C14	1.93 minutes ⁹ ; 347
140		Method D; C14	2.39 minutes ⁹ ; 334
141		Method D; C14 ²⁶	2.22 minutes ⁹ ; 384
142		Method D; C14	2.61 minutes ⁹ ; 414
143		Method D; C14	1.87 minutes ⁹ ; 363
144		Method D; C14	1.86 minutes ⁹ ; 302
145		Method D; C14 ²⁷	2.25 minutes ⁹ ; 402








146		Method D; C14	2.17 minutes ⁹ ; 386
147		Ex 3	¹ H NMR (400 MHz, CDCl ₃) δ 11.32 (br s, 1H), 8.46 (s, 1H), 7.39-7.43 (m, 1H), 7.27-7.30 (m, 1H), 7.24 (s, 1H), 7.08-7.11 (m, 1H), 3.88 (s, 3H), 2.88 (s, 6H); 294.1
148		C20 ²⁸	2.07 minutes ⁹ ; 299
149		Ex 3	8.59 (br d, <i>J</i> =1.5 Hz, 1H), 8.39 (s, 1H), 8.33-8.36 (m, 1H), 7.73-7.76 (m, 1H), 7.63 (s, 1H), 3.4-3.47 (m, 4H, assumed; partially obscured by water peak), 3.12-3.17 (m, 4H), 2.38 (s 3H); 296.0
150		Ex 3	8.78 (br s, 1H), 8.54 (br d, <i>J</i> =2 Hz, 1H), 8.42 (s, 1H), 8.07-8.10 (m, 1H), 7.80 (s, 1H), 3.46-3.54 (m, 4H), 3.11-3.18 (m, 4H); 316.1
151		Ex 3 ²⁹	8.48 (s, 1H), 8.36 (s, 1H), 8.13 (s, 1H), 7.87 (s, 1H), 3.99 (s, 3H), 3.49-3.56 (m, 4H), 3.21-3.27 (m, 4H); 313.1
152		Ex 3	8.30 (s, 1H), 7.61 (s, 1H), 7.22 (s, 1H), 3.80 (s, 3H), 3.4-3.48 (m, 4H, assumed; partially obscured by water peak), 3.15-3.21 (m, 4H), 2.13 (s, 3H); 299.2








153		Ex 5	characteristic peaks: 8.40 (s, 1H), 8.00 (br s, 1H), 7.90 (br d, $J=8$ Hz, 1H), 7.76 (br d, $J=7$ Hz, 1H), 7.65-7.70 (m, 2H), 3.60-3.74 (m, 2H), 0.93 (d, $J=6.5$ Hz, 3H); 320.1
154		Ex 1 ⁷	¹ H NMR (400 MHz, CD ₃ OD) δ 7.80-7.86 (m, 1H), 7.70-7.75 (m, 1H), 7.47 (dd, $J=7.8, 7.8$ Hz, 1H), 7.39 (d, $J=1$ Hz, 1H), 3.45-3.50 (m, 4H), 3.24-3.28 (m, 4H), 2.58 (s, 3H); 338.2
155		Ex 6	characteristic peaks: 12.21 (br s, 1H), 8.33 (s, 1H), 7.94 (br s, 1H), 7.82 (br d, $J=8$ Hz, 1H), 7.73 (br d, $J=8$ Hz, 1H), 7.60-7.66 (m, 2H), 4.71 (d, $J=4.5$ Hz, 1H), 3.70-3.79 (m, 1H), 3.44-3.52 (m, 1H), 2.60-2.70 (m, 1H), 1.73-1.82 (m, 1H), 1.38-1.48 (m, 1H), 1.12-1.26 (m, 2H); 320.1
156		Method E	2.48 minutes ⁹ ; 313
157		Method E ²⁰	2.66 minutes ⁹ ; 411
158		Method E	2.41 minutes ¹⁰ ; 340






159		Method E	2.66 minutes ⁹ ; 327
160		Method E	2.60 minutes ¹⁰ ; 345
161		Method E	2.92 minutes ⁹ ; 358
162		Method E	1.88 minutes ⁹ ; 310
163		Method E	2.47 minutes ¹⁰ ; 345
164		Method E	2.54 minutes ⁹ ; 343

165		Method E	2.41 minutes ⁹ ; 325
166		Method E	2.65 minutes ¹³ ; 406
167		Method E	2.58 minutes ⁹ ; 368
168		Method E	2.17 minutes ⁹ ; 299
169		Method E	1.74 minutes ⁹ ; 310
170		Method E	2.02 minutes ¹⁰ ; 283
171		Method E ²⁰	2.43 minutes ⁹ ; 381






172		Method E	2.27 minutes ⁹ ; 325
173		Method E	2.12 minutes ⁹ ; 269
174		Method E	2.13 minutes ⁹ ; 326
175		Method E	2.22 minutes ⁹ ; 313
176		Method E	2.24 minutes ⁹ ; 295
177		Method E	2.24 minutes ⁹ ; 287
178		Method E	2.04 minutes ⁹ ; 338







179		Method E	2.31 minutes ⁹ ; 343
180		Method E	2.27 minutes ⁹ ; 338
181		Method E	2.13 minutes ¹⁰ ; 345
182		Method E	2.14 minutes ¹⁰ ; 313
183		Method E	2.15 minutes ⁹ ; 287
184		Method E	2.12 minutes ⁹ ; 308
185		Method E	2.26 minutes ⁹ ; 331







186		Method E ²⁰	2.36 minutes ⁹ ; 406
187		Method F; P2	2.07 minutes ⁹ ; 302
188		Method F; P2	1.84 minutes ⁹ ; 296
189		Method F; P7	2.29 minutes ⁹ ; 287
190		Method F; P2	2.33 minutes ⁹ ; 329
191		Method F; P2	2.35 minutes ⁹ ; 317
192		Method F; P2	2.18 minutes ⁹ ; 302


193		Method F; P2 ³⁰	2.47 minutes ⁹ ; 363
194		Method F; P2	1.94 minutes ¹³ ; 312
195		Ex 5 ^{20,31,17}	characteristic peaks: 12.37 (br s, 1H), 8.43 (s, 1H), 7.99-8.02 (m, 1H), 7.85 (ddd, $J=7.8$, 1, 1 Hz, 1H), 7.75 (ddd, $J=7.8$, 1, 1 Hz, 1H), 7.71 (s, 1H), 7.65 (dd, $J=7.8$, 7.8 Hz, 1H), 4.67 (dd, $J=9.5$, 2.5 Hz, 1H), 3.85 (br d, $J=13$ Hz, 1H), 3.68-3.74 (m, 1H), 3.18 (dd, $J=13$, 9 Hz, 1H), 2.86-2.94 (m, 1H), 2.57 (s, 3H); 388.1
196		Ex 5 ^{20,31,12}	characteristic peaks: 12.37 (br s, 1H), 8.43 (s, 1H), 8.00 (dd, $J=1.5$, 1.5 Hz, 1H), 7.85 (ddd, $J=7.8$, 1.5, 1.5 Hz, 1H), 7.75 (ddd, $J=7.8$, 1.5, 1.2 Hz, 1H), 7.71 (s, 1H), 7.65 (dd, $J=7.8$, 7.8 Hz, 1H), 4.67 (dd, $J=9.7$, 2.6 Hz, 1H), 3.85 (br d, $J=13$ Hz, 1H), 3.68-3.74 (m, 1H), 3.18 (dd, $J=13.0$, 9.5 Hz, 1H), 2.86-2.94 (m, 1H), 2.57 (s, 3H); 388.1
197		Ex 6 ^{16,12}	¹ H NMR (400 MHz, CDCl ₃) δ 8.48 (s, 1H), 7.75 (br s, 1H), 7.65-7.70 (m, 2H), 7.59 (dd, $J=7.8$, 7.5 Hz, 1H), 7.33 (s, 1H), 4.02 (br d, $J=13.3$ Hz, 1H), 3.89 (br d, $J=13.3$ Hz, 1H), 3.80 (br dd, $J=12.0$, 2.3 Hz, 1H), 3.41-3.54 (m, 2H), 3.29 (s, 3H), 3.25-3.3 (m, 1H), 3.22 (dd, half of ABX pattern, $J=10.2$, 4.6 Hz, 1H), 2.98-3.08 (m, 1H), 2.87 (dd, $J=13.2$, 10.7 Hz, 1H); 350.2

203		Example 3; P2	¹ H NMR (400 MHz, CDCl ₃) δ 10.10 (br s, 1H), 8.50 (s, 1H), 7.67 (br d, <i>J</i> =7.5 Hz, 1H), 7.55 (br d, <i>J</i> =7.5 Hz, 1H), 7.38 (dd, <i>J</i> =7.9, 7.8 Hz, 1H), 7.11-7.13 (m, 1H), 3.35-3.42 (m, 4H), 3.21-3.26 (m, 4H), 2.49 (s, 3H); 319.9
204		Example 4; P2	8.34 (s, 1H), 7.75 (s, 1H), 7.37 (d, <i>J</i> =6.5 Hz, 1H), 6.49 (br s, 1H), 6.44 (dd, <i>J</i> =7.0, 1.5 Hz, 1H), 3.60-3.66 (m, 4H), 3.21-3.27 (m, 4H, assumed; partially obscured by water peak); 297.9
205		Example 3 ³² , P2 ³³	¹ H NMR (400 MHz, CDCl ₃) δ 10.08 (br s, 1H), 8.51 (s, 1H), 7.36 (d, <i>J</i> =0.9 Hz, 1H), 7.29-7.31 (m, 1H), 7.25-7.27 (m, 1H, assumed; partially obscured by solvent peak), 6.87 (dd, <i>J</i> =4.5, 1.0 Hz, 1H), 3.31-3.35 (m, 4H), 3.26-3.31 (m, 4H); 327.1
206		Example 4; C14	¹ H NMR (400 MHz, CDCl ₃), characteristic peaks: δ 11.32 (br s, 1H), 8.42 (s, 1H), 7.78 (br s, 1H), 7.72 (br d, <i>J</i> =7.8 Hz, 1H), 7.62 (br d, <i>J</i> =7.8 Hz, 1H), 7.52 (dd, <i>J</i> =7.8, 7.8 Hz, 1H), 7.18 (s, 1H), 4.34 (br dd, <i>J</i> =5, 5 Hz, 1H), 3.83-3.91 (m, 1H), 3.74 (ddd, <i>J</i> =8.3, 7.8, 5.3 Hz, 1H), 3.46-3.56 (m, 3H), 3.19 (dd, <i>J</i> =12.4, 4.9 Hz, 1H), 2.74-2.84 (m, 1H), 1.99-2.10 (m, 1H); 331.9
207		Example 4; C14	¹ H NMR (400 MHz, CDCl ₃), δ 10.25 (br s, 1H), 8.43 (s, 1H), 7.81 (br s, 1H), 7.73 (br d, <i>J</i> =7.8 Hz, 1H), 7.63 (br d, <i>J</i> =7.8 Hz, 1H), 7.54 (dd, <i>J</i> =7.8, 7.7 Hz, 1H), 7.18 (s, 1H), 3.71 (br dd, <i>J</i> =9.0, 6.2 Hz, 2H), 3.56 (br dd, <i>J</i> =11.3, 7.2 Hz, 2H), 3.47 (dd, <i>J</i> =9.0, 2.6 Hz, 2H), 3.21 (dd, <i>J</i> =11.4, 4.1 Hz, 2H), 2.75-2.81 (m, 2H); 331.9

208		Example 4; C14	8.49 (s, 1H), 8.14 (br s, 1H), 8.03 (br d, $J=7.8$ Hz, 1H), 7.79 (s, 1H), 7.76 (br d, $J=7.5$ Hz, 1H), 7.68 (dd, $J=7.8, 7.8$ Hz, 1H), 3.4-3.50 (m, 2H, assumed; partially obscured by water peak), 2.92-3.08 (m, 2H), 2.78 (br d, $J=12.3$ Hz, 1H), 2.65-2.75 (m, 1H), 2.42-2.5 (m, 1H, assumed; partially obscured by solvent peak), 1.83-1.94 (m, 1H), 1.61-1.83 (m, 2H), 1.47-1.61 (m, 1H), 1.31-1.45 (m, 1H); 345.9
209		Example 4; C14	8.45 (s, 1H), 8.04 (br s, 1H), 7.91 (br d, $J=8$ Hz, 1H), 7.80 (br d, $J=8$ Hz, 1H), 7.76 (s, 1H), 7.66 (dd, $J=8, 8$ Hz, 1H), 5.06-5.09 (m, 1H), 3.79-3.86 (m, 1H), 3.43-3.52 (m, 2H), 3.3-3.41 (m, 1H, assumed; partially obscured by water peak), 3.10-3.18 (m, 1H), 2.82-2.91 (m, 1H); 353.1 [M+Na ⁺]
210		Example 4; C14	8.37 (s, 1H), 7.95 (br s, 1H), 7.87 (br d, $J=7.9$ Hz, 1H), 7.78 (br d, $J=7.6$ Hz, 1H), 7.64-7.69 (m, 2H), 3.21-3.27 (m, 2H), 3.17 (br s, 2H), 2.87-2.93 (m, 2H), 1.19 (s, 6H); 334.0
211		Example 3; P2	12.54 (br s, 1H), 8.41 (s, 1H), 8.00 (d, $J=3.8$ Hz, 1H), 7.78 (d, $J=2.5$ Hz, 1H), 7.28 (d, $J=3.8$ Hz, 1H), 3.52-3.58 (m, 4H), 3.20-3.26 (m, 4H); 311.9
212		Example 3; P2	8.54 (br d, $J=4.3$ Hz, 1H), 8.36 (s, 1H), 8.20 (br d, $J=9.3$ Hz, 1H), 7.94 (s, 1H), 7.72 (s, 1H), 7.28 (dd, $J=9.2, 4.4$ Hz, 1H), 3.01-3.13 (m, 8H); 321.9

213	 <p>ENT-2</p>	Example 4 ¹ ; C1 ³⁴	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆ + D ₂ O) δ 8.41 (s, 1H), 7.80-7.86 (m, 1H), 7.70-7.77 (m, 1H), 7.65 (d, <i>J</i> =1 Hz, 1H), 7.42 (dd, <i>J</i> =7.8, 7.8 Hz, 1H), 4.74-4.80 (m, 1H), 3.78-3.85 (m, 1H), 3.60-3.66 (m, 1H), 3.29-3.45 (m, 3H), 2.88-2.97 (m, 1H), 2.34 (s, 3H); LCMS <i>m/z</i> 428.0 [M+Na ⁺]
214	 <p>ENT-1</p>	Example 4 ¹ ; C1 ³⁴	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆ + D ₂ O) δ 8.38 (s, 1H), 7.75-7.82 (m, 1H), 7.68-7.74 (m, 1H), 7.60 (s, 1H), 7.40 (dd, <i>J</i> =7.8, 7.8 Hz, 1H), 4.68-4.74 (m, 1H), 3.71-3.78 (m, 1H), 3.56-3.64 (m, 1H), 3.29-3.44 (m, 3H), 2.89-2.98 (m, 1H), 2.31 (s, 3H); LCMS <i>m/z</i> 428.0 [M+Na ⁺]
215	 <p>• HCOOH</p>	Example 4; P2	12.46 (br s, 1H), 8.40 (s, 1H), 8.23 (br s, 1H), 8.11 (d, <i>J</i> =2.5 Hz, 1H), 8.06 (dd, <i>J</i> =8.0, 7.5 Hz, 1H), 7.85-7.90 (m, 2H), 7.77 (br s, 1H), 3.49-3.55 (m, 4H), 3.21-3.27 (m, 4H); 347.1 [M+Na ⁺]
216	 <p>• HCOOH</p>	Example 3 ³⁵ ; P2	characteristic peaks: 9.16 (br d, <i>J</i> =7 Hz, 1H), 8.57-8.60 (m, 1H), 8.41 (s, 1H), 8.35 (s, 1H), 7.58 (s, 1H), 7.06-7.11 (m, 1H), 3.14-3.19 (m, 4H); 322.1
217	 <p>• HCOOH</p>	Example 3; P2	12.31 (br s, 1H), 8.38 (s, 1H), 7.42 (d, <i>J</i> =2.3 Hz, 1H), 7.35 (d, <i>J</i> =1.8 Hz, 1H), 7.08 (d, <i>J</i> =1.8 Hz, 1H), 3.82 (s, 3H), 3.54-3.60 (m, 4H), 3.27-3.33 (m, 4H); 308.9
218	 <p>• HCOOH</p>	Example 3; P2	12.42 (br s, 1H), 8.40 (s, 1H), 8.13 (s, ~0.6 H), 7.58 (d, <i>J</i> =2.3 Hz, 1H), 7.53 (d, <i>J</i> =4.5 Hz, 1H), 7.25 (d, <i>J</i> =4.3 Hz, 1H), 3.22-3.31 (m, 2H), 3.06-3.14 (m, 2H), 2.97-3.06 (m, 4H), 2.30 (s, 3H); 340.9

219		Example @109; C14	characteristic peaks: 12.40 (br s, 1H), 8.53 (d, $J=1.5$ Hz, 1H), 8.44 (s, 1H), 8.00-8.02 (m, 1H), 7.86 (br d, $J=8$ Hz, 1H), 7.75 (br d, $J=8$ Hz, 1H), 7.71 (d, $J=2.5$ Hz, 1H), 7.66 (dd, $J=8, 8$ Hz, 1H), 6.39 (d, $J=1.5$ Hz, 1H), 4.66-4.71 (m, 1H), 3.80-3.86 (m, 1H), 3.72-3.78 (m, 1H), 3.04-3.11 (m, 1H), 2.88-2.97 (m, 1H); 394.9 [M+Na ⁺]
220		Example 4; P2 ³⁶	1.43 minutes ³⁷ ; 310.3
221		Method F; P2	1.95 minutes ³⁸ ; 312.2
222		Method F; P2	1.57 minutes ³⁸ ; 321.3
223		Method F; P2	1.86 minutes ³⁸ ; 323.3
224		Example 4; P2 ³⁹	1.63 minutes ³⁷ ; 310.2

225		Example 4; C14 ⁴⁰	characteristic peaks: 12.38 (br s, 1H), 8.42 (s, 1H), 8.00 (dd, <i>J</i> =1.6, 1.4 Hz, 1H), 7.86 (ddd, <i>J</i> =7.7, 1.7, 1.4 Hz, 1H), 7.77 (ddd, <i>J</i> =7.6, 1.5, 1.4 Hz, 1H), 7.71-7.73 (m, 1H), 7.69 (dd, <i>J</i> =8.0, 7.5 Hz, 1H), 3.72-3.79 (m, 1H), 3.60-3.68 (m, 2H), 2.63-2.77 (m, 4H); 344.9
-----	---	--	--

1. In this case, tetrakis(triphenylphosphine)palladium(0) was used in the Suzuki reaction.

2. Boronate ester 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1*H*-pyrazole
5 was used in the Suzuki reaction.

3. Conditions for analytical HPLC. Column: Waters XBridge C18, 2.1 x 50 mm, 5 µm; Mobile phase A: 0.1% trifluoroacetic acid in water; Mobile phase B: 0.1% trifluoroacetic acid in acetonitrile; Gradient: 5% B for 1.0 minute, then 5% to 100% B over 5.75 minutes; Flow rate: 0.5 mL/minute.

10 4. Dichlorobis(triphenylphosphine)palladium(II) was used in the Suzuki reaction.

5. 1 N Sodium hydroxide solution was employed in the Suzuki reaction; this served to remove the (4-methylphenyl)sulfonyl protecting group.

6. In this case, preparative HPLC purification was carried out using a 10 µm Sunfire C-18 column; Eluent: 1:1 acetonitrile / (0.1% formic acid in water).

15 7. A boronic acid was used rather than a boronic ester.

8. The final deprotection was carried out using tetrabutylammonium fluoride in tetrahydrofuran.

9. Conditions for analytical HPLC. Column: Waters XBridge C18, 2.1 x 50 mm, 5 µm; Mobile phase A: 0.0375% trifluoroacetic acid in water; Mobile phase B: 0.01875% trifluoroacetic acid in acetonitrile; Gradient: 1% to 5% B over 0.60 minutes; then 5% to 100% B over 3.40 minutes; Flow rate: 0.8 mL/minute.

10. Conditions for analytical HPLC as in footnote 9, except that the gradient employed was 10% to 100% B over 4.00 minutes.

11. The racemic product was separated into its enantiomers using supercritical fluid chromatography. Column: Chiral Technologies Chiralpak® AD, 5 µm; Eluent: 68:32 carbon dioxide / methanol.

12. This Example was the second-eluting enantiomer from the supercritical fluid chromatographic separation.

13. Conditions for analytical HPLC. Column: Waters XBridge C18, 2.1 x 50 mm, 5 µm; Mobile phase A: 0.05% ammonium hydroxide in water; Mobile phase B: acetonitrile; Gradient: 5% to 100% B over 3.40 minutes; Flow rate: 0.8 mL/minute.

14. [1,1'-Bis(diphenylphosphino)ferrocene]dichloropalladium(II) was used in the Suzuki
5 reaction.

15. The racemic product was separated into its enantiomers using supercritical fluid chromatography. Column: Chiral Technologies Chiralpak® AD, 5 µm; Eluent: 60:40 carbon dioxide / (methanol containing 0.05% diethylamine).

16. The racemic product was separated into its enantiomers using supercritical fluid
10 chromatography. Column: Chiral Technologies Chiralpak® AD, 20 µm; Eluent: 60:40 carbon dioxide / (ethanol containing 0.05% diethylamine).

17. This Example was the first-eluting enantiomer from the supercritical fluid chromatographic separation.

18. The racemic product was separated into its enantiomers using supercritical fluid
15 chromatography. Column: Chiral Technologies Chiralpak® AS, 5 µm; Eluent: 60:40 carbon dioxide / (2-propanol containing 0.05% diethylamine).

19. 3-Iodo-1-methyl-1*H*-pyrazole was used in the Suzuki reaction.

20. The requisite 2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholine may be prepared from 4-benzylmorpholine-2-carbonitrile using chemistry described by D. Sakai and K. Watanabe, *PCT
20 Int. Appl.* **2009**, WO 2009035159 A1 20090319.

21. In this case, the displacement reaction was carried out at 110 °C.

22. The requisite 2-[[6-methylpyridin-3-yl]oxy]methyl)morpholine may be prepared via a Mitsunobu reaction between *tert*-butyl 2-(hydroxymethyl)morpholine-4-carboxylate and 6-methylpyridin-3-ol, followed by acidic removal of the protecting group.

23. 2-[[5-(Difluoromethyl)-1,2,4-oxadiazol-3-yl]methyl)morpholine may be prepared in the
25 following manner: reaction of (4-benzylmorpholin-2-yl)acetonitrile with hydroxylamine hydrochloride and base provides 2-(4-benzylmorpholin-2-yl)-*N'*-hydroxyethanimidamide. Coupling of this compound with difluoroacetic acid may be carried out using any of a number of coupling reagents, such as 1,3-dicyclohexylcarbodiimide. Subsequent cyclization using thermal
30 conditions or tetrabutylammonium fluoride (see A. R. Gangloff et al., *Tetrahedron Lett.* **2001**, 42,1441-1443), followed by debenzylation, provides the requisite amine.

24. Reaction of 4-benzylmorpholine-2-carbonitrile with hydroxylamine hydrochloride and sodium hydroxide provided 4-benzyl-*N'*-hydroxymorpholine-2-carboximidamide, which was condensed with cyclopropanecarboxylic acid using 1,3-dicyclohexylcarbodiimide. Cyclization
35 using tetrabutylammonium fluoride (see A. R. Gangloff et al., *Tetrahedron Lett.* **2001**, 42,1441-1443), followed by debenzylation, afforded 2-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)morpholine.

25. The requisite 3-(morpholin-2-ylmethoxy)benzonitrile may be prepared in a manner analogous to that described in footnote 22.

26. 2-(Pyrimidin-4-yl)morpholine may be prepared from 1-(4-benzylmorpholin-2-yl)ethanone via conversion to the enamine and reaction with 1,3,5-triazine (see D. L. Boger et al., *J. Org. Chem.* **1982**, 47, 2673-2675), followed by debenzylation.

27. 2-[(5-Methyl-1,2,4-oxadiazol-3-yl)methyl]morpholine may be prepared via the chemistry described in footnote 23, by employing acetic acid in place of difluoroacetic acid.

28. Compound **C20** was *N*-alkylated with ethyl iodide and potassium *tert*-butoxide at 30 °C in tetrahydrofuran. Deprotection was carried out using the procedure described in step 2 of Method C.

29. 2-Chloro-6-methoxypyrazine was used in the Suzuki reaction.

30. 2-Chloro-1-fluoro-3-iodo-4-methoxybenzene may be prepared by iodination of 2-chloro-1-fluoro-4-methoxybenzene according to the method of R. Sanz et al., *J. Org. Chem.* **2007**, 72, 5113-5118.

31. The racemic product was separated into its enantiomers using supercritical fluid chromatography. Column: Chiral Technologies Chiralpak® AS, 5 µm; Eluent: 63:37 carbon dioxide / (methanol containing 0.05% ammonium hydroxide).

32. In this case, potassium phosphate and 1,2-dimethoxyethane were used in place of sodium carbonate and 1,4-dioxane.

33. 5-Bromoimidazo[2,1-*b*][1,3]thiazole may be prepared via bromination of imidazo[2,1-*b*][1,3]thiazole using *N*-bromosuccinimide.

34. The two enantiomers were separated by supercritical fluid chromatography. Example 213 was the second-eluting enantiomer, and Example 214 was the first-eluting isomer, using a Chiral Technologies Chiralcel OJ-3 column, and gradient of 5% to 40% methanol in carbon dioxide containing 0.05% diethylamine.

35. In this case, the Suzuki reaction was carried out using bis[di-*tert*-butyl(4-dimethylaminophenyl)phosphine]dichloropalladium(II) and cesium fluoride in a mixture of 1,4-dioxane and water at 100 °C for 18 hours.

36. Treatment of 4-bromo-1-methyl-1*H*-imidazole with lithium diisopropylamide and *N,N*-dimethylformamide provided 4-bromo-1-methyl-1*H*-imidazole-2-carbaldehyde, which was converted to the requisite 4-bromo-1-methyl-1*H*-imidazole-2-carbonitrile via reaction with ammonium hydroxide and iodine, according to the method of J.-J. Shie and J.-M. Fang, *J. Org. Chem.* **2007**, 72, 3141-3144.

37. Conditions for analytical HPLC. Column: Waters Atlantis dC18, 4.6 x 50 mm, 5 µm; Mobile phase A: 0.05% trifluoroacetic acid in water (v/v); Mobile phase B: 0.05% trifluoroacetic

acid in acetonitrile (v/v); Gradient: 5.0% to 95% B, linear over 4.0 minutes; Flow rate: 2 mL/minute.

38. Conditions for analytical HPLC. Column: Waters XBridge C18, 4.6 x 50 mm, 5 μ m; Mobile phase A: 0.03% ammonium hydroxide in water (v/v); Mobile phase B: 0.03% ammonium hydroxide in acetonitrile (v/v); Gradient: 5.0% to 95% B, linear over 4.0 minutes; Flow rate: 2 mL/minute).

39. 3-Bromo-1-methyl-1*H*-pyrazole-5-carbaldehyde was converted to 3-bromo-1-methyl-1*H*-pyrazole-5-carbonitrile using the method of J.-J. Shie and J.-M. Fang; see footnote 36.

40. Treatment of (4-benzylmorpholin-2-yl)acetonitrile with ammonium cerium(IV) nitrate provided the requisite morpholin-2-ylacetonitrile.

Biological Assays

LRRK2 assay

LRRK2 kinase activity was measured using Lantha Screen technology from Invitrogen. GST-tagged truncated LRRK2 from Invitrogen (Cat # PV4874) was incubated with a fluorescein-labeled peptide substrate based upon ezrin/radixin/moesin (ERM), also known as LRRKtide (Invitrogen cat # PR8976A), in the presence of a dose response of compound. Upon completion, the assay was stopped and detected with a terbium labeled anti-phospho-ERM antibody (Invitrogen, cat # PR8975A). The assay was carried out under the following protocol: 3 μ L of a working solution of substrate (233 nM LRRKtide, 117 μ M ATP) prepared in assay buffer (50 mM HEPES, pH 7.5, 3 mM $MgCl_2$, with 2 mM DTT and 0.01% Brij35 added fresh) was added to a low volume Greiner 384-well plate. The compound dose response was prepared by diluting compound to a top concentration of 3.16 mM in 100% DMSO and serially diluted by half-log in DMSO 11 times. Aliquots (3.5 μ L) of the 100% DMSO dose response were mixed with 46.5 μ L water then 1 μ L of this mixture was added to the 3 μ L substrate mix in the 384-well plate. The kinase reaction was started with 3 μ L of a working solution of LRRK2 enzyme at a concentration of 4 μ g/mL. The final reaction concentrations were 100 nM LRRKtide, 50 μ M ATP, 1.7 μ g/mL LRRK2 enzyme and a compound dose response with a top dose of 32 μ M. The reaction was allowed to progress at room temperature for two hours and then stopped with the addition of 7 μ L of detection buffer (20 mM Tris pH 7.6, 0.01% NP-40, 0.02% NaN_3 , 6 mM EDTA with 2 nM terbium labeled anti-phospho-ERM). After an incubation of 1 hour at room temperature, the plate was read on an Envision with an excitation wavelength of 340 nm and a reading emission at

both 520 nm and 495 nm. The ratio of the 520 nm and 495 nm emission was used to analyze the data.

Inhibition of mutant G2019S LRRK2 (Invitrogen cat # PV4881) was measured in the exact same method. All final concentrations of substrate ATP and enzyme were the same. However, since the mutant enzyme is more active the reaction time was reduced to 90 minutes to ensure that inhibition was measured at steady state before any substrate depletion could occur.

Table 2, below, provides the LRRK2 IC₅₀ data for the compounds of the invention.

Table 2

Example Number	IUPAC Name	LRRK2 WT IC ₅₀ (nM)*	LRRK2 G2019S IC ₅₀ (nM)*
1	5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	13 ^a	36 ^a
2	3-[6-methyl-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	12	43
3	6-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]pyridine-2-carbonitrile	9	26
4	3-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	3 ^a	11 ^a
5	3-[4-(dimethylamino)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	5	15
6	3-{4-[(2 <i>S</i>)-2-methylmorpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	7	37
7	(3-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}phenyl)methanol	18 ^a	18 ^a
8	4-(morpholin-4-yl)-5-(1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	25	33
9	3-[2-methyl-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	42	513
10	{3-[4-(dimethylamino)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]phenyl}methanol	34	23

11	3-[4-(3,3-dimethylpiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	90	325
12	3-[4-(piperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	6	28
13	1-(5-phenyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl)piperidine-3-carboxamide	41	39
14	3-{4-[(3 <i>S</i>)-3-hydroxypiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	17 ^b	95 ^b
15	4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-5-phenyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	91	92
16	3-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	16 ^a	45 ^a
17	4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	14	12
18	5-phenyl-4-(pyrrolidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	71	47
19	[1-(5-phenyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl)piperidin-4-yl]methanol	38	29
20	1-{5-[3-(hydroxymethyl)phenyl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl}piperidine-3-carbonitrile	56	31
21	1-[5-(3-cyanophenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidine-3-carbonitrile	57	183
22	4-(3,5- <i>cis</i> -dimethylpiperidin-1-yl)-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	16	16
23	4-methoxy-3-[4-(3-methylpiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	25	46
24	3-[4-(dimethylamino)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-4-methoxybenzonitrile	24	30
25	5-(5-chloro-2-methoxyphenyl)- <i>N,N</i> -dimethyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-amine	22	19
26	3-{4-[4-(1 <i>H</i> -imidazol-2-yl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, trifluoroacetate salt	42	230
27	3-{4-[3-(methoxymethyl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	95	219

28	3-[4-(9-methyl-1-oxa-4,9-diazaspiro[5.5]undec-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	40	201
29	3-[4-(3-methoxypiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	79	154
30	3-{4-[2-(methoxymethyl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	17	121
31	<i>N</i> ³ -[5-(3-cyanophenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]- <i>N,N</i> -dimethyl-beta-alaninamide	54	248
32	3-[4-(4,4-difluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	67	333
33	3-[4-(4-fluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	17	96
33A	3-[4-(4-fluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	17	56 ^b
34	3-{4-[4-(1 <i>H</i> -pyrazol-3-yl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	69	136
35	3-{4-[(3 <i>S</i>)-3-fluoropyrrolidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	30	155
36	3-{4-[2-(1 <i>H</i> -pyrazol-3-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	43	238
37	3-{4-[3-(1 <i>H</i> -pyrazol-3-yl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	84	200
38	3-{4-[3-(1 <i>H</i> -imidazol-2-yl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, formate salt	53	331
39	3-{4-[(1-methylpiperidin-3-yl)amino]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, formate salt	78	243
40	3-[4-(3-oxo-2,7-diazaspiro[4.5]dec-7-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	35	119
41	3-[4-(3-methylpiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile, ENT-2	70	172
42	3-(4-{[2-(morpholin-4-yl)ethyl]amino}-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl)benzonitrile, formate salt	89	487
43	3-[4-(2-oxa-7-azaspiro[4.5]dec-7-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	34	244

44	<i>N</i> -{1-[5-(3-cyanophenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidin-4-yl}acetamide	50	190
45	5-(1 <i>H</i> -indazol-5-yl)-4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	67	66 ^b
46	3-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzenesulfonamide	24	33 ^b
47	5-(2-fluorophenyl)-4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	97	122 ^b
48	5-(1 <i>H</i> -indazol-4-yl)-4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	10 ^a	14
49	5-(6-fluoro-5-methylpyridin-3-yl)-4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, trifluoroacetate salt	64	65 ^b
50	5-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}-2,3-dihydro-1 <i>H</i> -isoindol-1-one	33	33 ^b
51	4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-5-(pyridin-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	56	95 ^b
52	4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-5-(5-methylpyridin-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	15	19 ^b
53	4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-5-(7 <i>H</i> -pyrrolo[2,3- <i>b</i>]pyridin-5-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	74	69 ^b
54	6-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}-2,3-dihydro-1 <i>H</i> -isoindol-1-one	68	59 ^b
55	4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-5-(1 <i>H</i> -pyrrolo[3,2- <i>b</i>]pyridin-6-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	53	50 ^b
56	4-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}phenol	26	28 ^b
57	4-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzamide	13	9 ^b
58	3-{4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}phenol	9	12 ^b
59	5-(2-chloro-5-methylpyridin-3-yl)-4-[(3 <i>S</i>)-3-methylpiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	30	30 ^b

60	3-{4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzamide	86	58
61	3-[4-(3-fluoropiperidin-1-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	12	80
62	3-[4-(4-methylpiperidin-1-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile, hydrochloride salt	91	273
63	3-{4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzoic acid	55	39
64	3-[4-(methylamino)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	45	97
65	3-[4-(3,5- <i>cis</i> -dimethylpiperidin-1-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-4-methoxybenzonitrile	85	176
66	4-methoxy-3-{4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	21	51
67	2-fluoro-3-{4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	11	35
68	<i>N,N</i> -dimethyl-5-phenyl-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-4-amine	63	39
69	3-[4-(dimethylamino)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-2-fluorobenzonitrile	6	17
70	3-[4-(pyrrolidin-1-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	8	64
71	3-[4-(3-methylpyrrolidin-1-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile, ENT-2	21	102
72	3-[4-(dimethylamino)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-4-fluorobenzonitrile	37	190
73	3-[4-(diethylamino)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	13	119
74	3-{4-[2-(methoxymethyl)morpholin-4-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-1	12	100
75	3-[4-(dimethylamino)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-5-fluorobenzonitrile	53	396
76	3-{4-[2-(1 <i>H</i> -pyrazol-3-yl)morpholin-4-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-2, formate salt,	23	208

77	3-[4-(3-methylpyrrolidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile, ENT-1	39	158
78	2-fluoro-3-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	2	7
79	3-{4-[(2 <i>R</i>)-2-methylmorpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	18	69
80	4-fluoro-3-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	49	393
81	4-(4-fluoropiperidin-1-yl)-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	36	50
82	2-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	95	148
83	5-(3-chlorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	8	13
84	5-(2-fluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	22	30
85	5-(3-fluoro-5-methoxyphenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	9	20
86	5-(2,5-difluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	76	145
87	5-(2,3-difluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	12	24
88	5-(5-chloro-2-fluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	9	12
89	{2-fluoro-3-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]phenyl}methanol	32	75
90	5-(2,4-difluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	41	62
91	5-(3-fluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	9	21
92	5-(3,5-difluorophenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	92	238
93	4-(morpholin-4-yl)-5-phenyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	9	14

94	5-(5-fluoro-2-methoxyphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	7	10
95	5-(2-chlorophenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	21	33
96	5-(5-fluoro-2-methylphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	91	223
97	5-(3-methoxyphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	28	46
98	{2-fluoro-5-[4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]phenyl}methanol	70	135
99	5-(4-fluorophenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	95	154
100	{3-[4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]phenyl}methanol	12	19
101	5-(2-methoxyphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	14	23
102	5-[3-(methylsulfanyl)phenyl]-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	52	78
103	4-(morpholin-4-yl)-5-(pyridin-3-yl)-7H-pyrrolo[2,3-d]pyrimidine, formate salt	35	159
104	4-[4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]isoquinoline, formate salt	27	51
105	5-(5-bromopyridin-3-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine, formate salt	6	14
106	5-(2-chloro-5-methylphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	22	44
107	5-(3-methylphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	13	20
108	5-(1-methyl-1H-pyrazol-3-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	53	90
109	3-[4-(4-hydroxypiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile	52	109
110	3-[4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile	20	176

111	3-methyl-5-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	14 ^b	64 ^b
112	3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	10	41
113	3-[4-(6-oxa-3-azabicyclo[3.1.1]hept-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	58 ^b	523 ^b
114	3-chloro-5-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	77 ^b	406 ^b
115	4-methoxy-3-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	48	121
116	5-(5-chloro-2-methoxyphenyl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	85	67
117	6-methyl-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	47	89
118	3-methoxy-5-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	77	177
119	5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-4-(thiomorpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	71	51
120	1-[5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidin-3-ol	44	34
121	4-[(2 <i>S</i>)-2-methylmorpholin-4-yl]-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	57	59
122	4-[(2 <i>R</i>)-2-methylmorpholin-4-yl]-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	67	31
123	4-(3-fluoropiperidin-1-yl)-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	31	27
124	{4-[5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]morpholin-2-yl}methanol	77	80
125	5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-4-(2-[(6-methylpyridin-3-yl)oxy]methyl)morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	54	87
126	<i>N,N</i> -dimethyl-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-amine	55	35
127	<i>N</i> -cyclopropyl- <i>N</i> -methyl-5-(1-methyl-1 <i>H</i> -pyrazol-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-amine	37	37

128	3-[4-(3,3-difluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	79	169
129	3-[4-(3-aminopiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	14	21
130	3-[4-(2-([5-(difluoromethyl)-1,2,4-oxadiazol-3-yl]methyl)morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	44	134
131	3-{4-[2-(1,3-thiazol-2-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	45	181
132	3-[4-(4-oxopiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	66	223
133	3-{4-[4-fluoro-4-(hydroxymethyl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	42	106
134	3-{4-[2-(3-hydroxyphenyl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	91	281
135	3-{4-[2-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	19	47
136	3-[4-(3,3-difluoropyrrolidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	74	194
137	3-(4-{2-[(3-cyanophenoxy)methyl]morpholin-4-yl}-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl)benzonitrile	30	51
138	3-[4-(3-hydroxypiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	10	27
139	1-[5-(3-cyanophenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidine-3-carboxamide	49	136
140	3-[4-(2-ethylmorpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	61	95
141	3-[4-[2-(pyrimidin-4-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	79	193
142	3-{4-[2-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	30	84
143	3-(4-{2-[(dimethylamino)methyl]morpholin-4-yl}-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl)benzonitrile	45	110
144	3-[4-(1 <i>H</i> -pyrazol-4-ylamino)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	69	36

145	3-(4-{2-[(5-methyl-1,2,4-oxadiazol-3-yl)methyl]morpholin-4-yl}-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl)benzonitrile	55	155
146	3-{4-[3-(5-methyl-1,3,4-oxadiazol-2-yl)piperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	96	217
147	3-[4-(dimethylamino)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-5-methoxybenzonitrile	37	93
148	5-(1-ethyl-1 <i>H</i> -pyrazol-4-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	77 ^b	92 ^b
149	5-(5-methylpyridin-3-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	18	30
150	5-(5-chloropyridin-3-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	28	36
151	5-(6-methoxypyrazin-2-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	32	37
152	5-(1,3-dimethyl-1 <i>H</i> -pyrazol-4-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	60	91
153	3-{4-[(3 <i>S</i>)-3-methylmorpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	44	165
154	2-fluoro-3-[2-methyl-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	30	365
155	3-{4-[(3 <i>R</i>)-3-hydroxypiperidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	8 ^b	37 ^b
156	1-[5-(3-fluorophenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidin-3-ol	57	86
157	5-(5-fluoro-2-methoxyphenyl)-4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	44	46
158	2-fluoro-3-[4-(4-fluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	18	86
159	4-(4-fluoropiperidin-1-yl)-5-(2-methoxyphenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	67	74
160	5-(3-fluoro-5-methoxyphenyl)-4-(4-fluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	84	96

161	3-[4-(4,4-difluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-2-fluorobenzonitrile	N.D. ^c	N.D.
162	1-[5-(5-methylpyridin-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidin-3-ol, formate salt	36	67
163	5-(5-fluoro-2-methoxyphenyl)-4-(4-fluoropiperidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	25	29
164	1-[5-(3-fluoro-5-methoxyphenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidin-3-ol	29	38
165	1-[5-(2-methoxyphenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]piperidin-3-ol	58	62
166	2-fluoro-3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	38	272
167	2-fluoro-3-{4-[2-(methoxymethyl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	61	372
168	4-(azetidin-1-yl)-5-(5-fluoro-2-methoxyphenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	85	75
169	4-[(2 <i>S</i>)-2-methylmorpholin-4-yl]-5-(5-methylpyridin-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	99	280
170	5-(3-fluorophenyl)-4-(pyrrolidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	N.D.	N.D.
171	5-(3-fluorophenyl)-4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	N.D.	N.D.
172	5-(2-methoxyphenyl)-4-[(2 <i>S</i>)-2-methylmorpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	N.D.	N.D.
173	5-(2-methoxyphenyl)- <i>N,N</i> -dimethyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-amine	43	29
174	2-fluoro-3-{4-[(3 <i>S</i>)-3-fluoropyrrolidin-1-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	N.D.	N.D.
175	4-[(3 <i>S</i>)-3-fluoropyrrolidin-1-yl]-5-(2-methoxyphenyl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	41	54
176	5-(2-methoxyphenyl)-4-(pyrrolidin-1-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	54	68
177	5-(3-fluoro-5-methoxyphenyl)- <i>N,N</i> -dimethyl-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-4-amine	41	49

178	2-fluoro-3-[4-(3-hydroxypiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile	9	42
179	5-(5-fluoro-2-methoxyphenyl)-4-[(2S)-2-methylmorpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidine	19	30
180	2-fluoro-3-[4-[(2S)-2-methylmorpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile	10	62
181	5-(5-fluoro-2-methoxyphenyl)-4-(3-fluoropiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidine	22	24
182	5-(3-fluorophenyl)-4-[(2S)-2-methylmorpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidine	38	82
183	5-(5-fluoro-2-methoxyphenyl)-N,N-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine	13	11
184	2-fluoro-3-[4-(pyrrolidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile	15	82
185	5-(5-fluoro-2-methoxyphenyl)-4-[(3S)-3-fluoropyrrolidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidine	15	23
186	2-fluoro-3-[4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile	17	90
187	5-(4-methyl-1,3-thiazol-2-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	N.D.	N.D.
188	5-(4-methylpyridin-2-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine, formate salt	43	90
189	5-(2-fluoro-6-methoxyphenyl)-N,N-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine	40	27
190	5-(2-fluoro-6-methoxyphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	25	34
191	5-(2,6-difluorophenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	N.D.	N.D.
192	5-(3-methyl-1,2-thiazol-5-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	23	73
193	5-(2-chloro-3-fluoro-6-methoxyphenyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine	47	51
194	5-(4-methoxypyridin-3-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3-d]pyrimidine, formate salt	N.D.	N.D.

195	3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-1	8	73
196	3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-2	N.D.	N.D.
197	3-{4-[2-(methoxymethyl)morpholin-4-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-2	214	245
198	3-[6-(difluoromethyl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	127	1080
199	5-(5,6-dihydro-2H-pyran-3-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidine	97.8 ^b	351 ^b
200	5-(3,4-dihydro-2H-pyran-5-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidine	76.7 ^b	167 ^b
201	4-(morpholin-4-yl)-5-[3-(1,2,4-oxadiazol-3-yl)phenyl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidine	129 ^b	475 ^b
202	3-{4-[2-(3-methyl-1,2-oxazol-5-yl)morpholin-4-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	64.7	350
203	2-methyl-3-[4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	13.2	90.1
204	4-[4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]pyridin-2(1H)-one	11.7	32.7
205	5-(imidazo[2,1- <i>b</i>][1,3]thiazol-5-yl)-4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidine	15.8	60.5
206	<i>rel</i> -3-{4-[(3a <i>S</i> ,6a <i>S</i>)-hexahydro-5H-furo[2,3- <i>c</i>]pyrrol-5-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	139	591
207	<i>rel</i> -3-{4-[(3a <i>R</i> ,6a <i>S</i>)-tetrahydro-1H-furo[3,4- <i>c</i>]pyrrol-5(3H)-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	223 ^b	687 ^b
208	<i>rel</i> -3-{4-[(4a <i>R</i> ,7a <i>S</i>)-hexahydrocyclopenta[<i>b</i>][1,4]oxazin-4(4aH)-yl]-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	27.2	186
209	4-[5-(3-cyanophenyl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-4-yl]morpholine-2-carbonitrile	42.2	336
210	3-[4-(2,2-dimethylmorpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]benzonitrile	35.2	304
211	5-[4-(morpholin-4-yl)-7H-pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]thiophene-2-carbonitrile	21.9	118

212	5-(imidazo[1,2- <i>b</i>]pyridazin-3-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	6.26	12.3
213	2-fluoro-3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-2	7.71	95.7
214	2-fluoro-3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile, ENT-1	N.D.	N.D.
215	6-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]pyridine-2-carboxamide, formate salt	30.6	79.3
216	4-(morpholin-4-yl)-5-(pyrazolo[1,5- <i>a</i>]pyrimidin-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	35.6	119
217	1-methyl-4-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-1 <i>H</i> -pyrrole-2-carbonitrile	6.37	13
218	5-(6-methylimidazo[2,1- <i>b</i>][1,3]thiazol-5-yl)-4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine, formate salt	37.4 ^b	135 ^b
219	3-{4-[2-(1,2-oxazol-5-yl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	37	234
220	1-methyl-4-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-1 <i>H</i> -imidazole-2-carbonitrile, trifluoroacetate salt	7.47	14.7
221	4-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]thiophene-2-carbonitrile	N.D.	N.D.
222	4-(morpholin-4-yl)-5-(pyrazolo[1,5- <i>a</i>]pyridin-3-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidine	N.D.	N.D.
223	1,5-dimethyl-4-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-1 <i>H</i> -pyrrole-2-carbonitrile	N.D.	N.D.
224	1-methyl-3-[4-(morpholin-4-yl)-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl]-1 <i>H</i> -pyrazole-5-carbonitrile, trifluoroacetate salt	N.D.	N.D.
225	3-{4-[2-(cyanomethyl)morpholin-4-yl]-7 <i>H</i> -pyrrolo[2,3- <i>d</i>]pyrimidin-5-yl}benzonitrile	N.D.	N.D.

* Geometric mean of 2 – 5 determinations unless otherwise indicated

a. IC₅₀ value represents the geometric mean of ≥6 determinations.

b. IC₅₀ value derived from a single determination.

c. N.D. – not determined

Certain of the compounds of the present invention were assessed for kinase selectivity using a commercially available ActiveX targeted kinase probe in which the tissue employed was human peripheral blood mononuclear cells (PBMC). The test compounds were provided to ActivX Biosciences, Inc., 11025 North Torrey Pines Road, Suite 120; LaJolla, CA, USA 92037. The compounds were run in the ActivX proprietary kinase assay and results were obtained at compound test concentrations of 1 μ M (columns 2, 4 and 6) and 0.1 μ M (columns 3, 5 and 7). The results as percent inhibition obtained for the compounds of Examples 1, 4 and 217 are provided below in Table 3.

TABLE 3

Kinase	Example 1		Example 4		Example 217	
	1 μ M	0.1 μ M	1 μ M	0.1 μ M	1 μ M	0.1 μ M
ABL,ARG	22.5	25.8	-0.9	-29.4	-1.6	4.6
ACK	-8	5.2	7.7	0.8	-19.5	-9.8
AKT1	18.8	11.3	-6.3	-8.8	-3.6	5.5
AKT2,AKT3	10.7	1.6	6.8	-2.6	-3.8	-9.7
AMPKa1	27	1.5	16.4	-2.1	48.8	27.2
AMPKa1,AMPKa2	49.7	10.8	32.5	-4.5	16.5	5.7
BRAF	-51.5	-6.5	-1.2	1.2	-10.4	-8.4
BTK	3.7	8.7	9.5	-2.2	8.5	24.3
CaMK1a	6.3	2	1.9	-9.2	-7.1	-5.4
CaMK1d	3.2	8.4	0.9	-12.1	3.1	11.7
CaMK2d	-38.1	-45.3	10.8	-0.4	16.1	9.8
CaMK2g	-15.5	-16.9	2.9	-13.6	18.7	4.3
CDC2	14.7	-4	12.9	-6.7	24.4	28.7
CDK11,CDK8	2.4	4.4	36	5	-5.2	-5.6
CDK2	9.5	-12.1	9.2	11	16.5	3.6
CDK5	-35.1	-6.5	0.1	-12.2	7.9	0.5
CHK2	17.7	0.9	2.2	-6.8	6.4	8.6
DGKA	-16.6	-21	-8.7	-15.9	4.7	-2.8
DNAPK	-30	-4.4	14.7	-19.3	31.5	31.0
eEF2K	-6.5	-9.6	2.8	-27.1	2.2	-12.1
EphA1	19.6	-9.5	-2.5	-17	-16.1	11.1
Erk5	-41.9	-32.8	-6.3	-5	-5.4	-2.6
FAK	20	18	11.1	-8.9	17.4	8.1
FER	-17.9	-7.3	1.6	-26.4	16.3	5.2
FES	-0.9	4.3	6.9	-9.2	6.2	8.4
FGR	4.6	12.9	1.7	-10.8	-0.5	-1.2
FRAP	-1.2	-3.2	9.8	-20.7	0.9	0.7
FYN	-10.1	6.7	-9.1	-1.9	-6.8	-2.1
FYN,SRC,YES	-56.3	-24.1	-8.8	-22.8	-1.4	18.4

GCK	-6.7	4.4	31.2	3.6	27.2	-5.2
GCN2	-25.1	0.1	0.6	-13.6	-3.1	-7.7
GPRK6	23.7	8.9	10.9	-3.9	3.6	-0.8
GSK3A	-11.4	-0.2	7.2	-0.9	8.9	0.6
GSK3B	-5.2	2.4	6.7	-3.7	12.4	-4.3
HPK1	-14.9	-9.7	5.7	-1.5	1.7	-13.7
IKKa	-40.5	-20	-6.6	-12.1	-3.7	-2.5
IKKb	10.8	16.1	-15.8	-15.5	6.5	7.1
IKKe	17.7	2.6	7.4	-7	20.4	-3.0
IKKe,TBK1	23.7	1.5	18.8	-6.7	13.3	2.3
ILK	-56.6	-34	-31.6	-8.9	-3.8	-6.3
IRAK1	-9.1	-13.8	2.1	-2.9	7.5	-1.7
IRAK3	68.7	30.1	33.7	29.8	72.4	16.0
IRAK4	34.7	17.7	10.7	-12.8	34.5	14.7
IRE1	-8	-1.3	1.8	-16	-19.5	3.2
JAK1	54.2	15	5	-9.3	11.4	-3.0
JAK1 domain2	-13.7	-8.1	18.9	-6.2	10.1	7.4
JAK2 domain2	16.7	13.8	16.4	0.7	-6.4	4.8
JAK3 domain2	-5.2	2.7	6.8	6.9	4.9	-7.8
JNK1,JNK2,JNK3	7.7	7.1	5.4	-5.3	-6.4	-18.2
KHS1	24.6	17.8	12.9	-25.9	7.6	10.7
KSR1,KSR2	-14	1.5	11.7	-8.7	-6.8	-6.2
LATS1	-16.2	4.2	5.4	-4.8	1.9	-3.9
LATS2	11	21.2	16.2	-7.2	-23.7	5.8
LKB1	-14	-1.1	16.5	3.5	2.4	-5.3
LOK	22.6	5.7	53.6	11.9	56.1	-6.4
LRRK2	88.1	69	94.1	79.9		88.4
LYN	7.8	3.8	0.8	-5.4	-19.2	8.3
MAP2K1	17	3.5	14	-17.2	-1.5	17.2
MAP2K1,MAP2K2	-2.2	1	6.4	-7.1	-2.6	10.9
MAP2K3	-9	14.4	6.2	-12.5	3.3	2.5
MAP2K4	9.6	10.1	19.2	-20.5	1.6	2.3
MAP2K6	19	17.5	13.5	-15.1	-0.1	11.3
MAP3K1	11.9	-9.4	44.5	6.4	14.8	-17.3
MAP3K3	7.2	-9.2	7.4	-0.5	-6.7	-3.4
MAP3K4	6.9	-14	29.8	9.7	-27.4	-35.5
MAP3K5	2.1	6.2	21.3	3.3	44.5	7.0
MARK2	41.6	10.1	27.3	-4.7	13.1	-5.0
MARK3	29.1	1	23.7	-2.6	18.4	16.2
MAST3	-6.7	9	0.8	2.6	0.4	-11.7
MLK3	4.5	1.4	19.9	-14.1	-9.6	-13.5
MLKL	8.4	1.6	14.5	-10.1	-3.3	2.5
MSK1 domain1	-30.4	-7	-3.8	1	-2.2	-12.1


MST1	50.8	13.2	82.2	41.9	73.5	25.6
MST1,MST2	24.8	-14.5	70.8	34.4	72.6	21.9
MST2	49.4	10.1	82.6	47.3	83.2	34.9
MST3	28.4	10.9	46.2	0.1	10.9	6.7
MST4,YSK1	15.1	-11.4	60.7	11.9	9.9	5.3
NDR1	-7.8	-19.6	1.7	-4.4	-17.7	16.3
NDR2	31.2	22.8	22.1	-18.7	-6.0	28.9
NEK1	-10.1	7	-1.9	-8.9	-5.1	4.8
NEK6,NEK7	-17	-1.7	-4.9	-5.3	6.4	-2.9
NEK7	-18.8	-0.6	7.3	2.2	-2.3	-9.0
NEK9	12.6	-0.9	1	-7.1	1.5	-11.0
NuaK2	8.9	3.6	18.1	-11.9	24.9	1.3
OSR1	-4	-7.8	-55.4	-23.1	-12.3	5.2
p38a	-19.8	-23.5	-1.7	-14.6	-4.6	-1.1
p38d,p38g	-24	-37.3	-5.7	-17.5	1.5	-0.4
p70S6K	18.9	23	11.1	-9	-9.6	3.5
p70S6K,p70S6Kb	-3.5	9.4	-9.6	-10.5	5.4	5.5
p70S6Kb	10.1	15	18.5	11.4	-16.7	-2.2
PAN3	0.8	0.6	15.1	-0.2	-7.5	5.6
PCTAIRE2,PCTAIRE3	-2.5	7.5	24.7	-5.4	15.1	-2.6
PI4KB	16.1	34	4.9	-7.4	-0.5	9.9
PIK3C3	4.8	23.3	13.6	-2.4	10.2	19.9
PIK3CB	-29.8	-2.6	-2.5	0.5	-29.7	-5.3
PIK3CD	-65.4	-101.6	-12.4	-14	10.0	-0.4
PIK3CG	-62.9	-18.6	-5.6	-20.4	-22.0	-3.4
PIP4K2A	6.8	12.1	1.9	-12.1	8.2	7.8
PIP4K2C	-1.4	1.4	58.4	3.1	-11.9	6.2
PIP4K2C	0.9	-48	54.3	9.3	4.0	-4.7
PIP5K3	15	1.9	21.6	-11.2	2.5	16.0
PITSLRE	-3.2	-13	10.7	-12.4	6.6	-6.7
PKD2	20.1	2.7	9.4	0.7	4.5	5.8
PKR	-17.3	-4.3	7.2	-3.3	1.8	-7.7
PRPK	-13.8	-3.4	-3.7	3	12.5	2.6
PYK2	19.5	5.9	2.5	-3.9	8.4	-1.0
RIPK3	-4.2	5.6	8	-3.2	-9.1	-7.3
RSK1 domain1	59.8	16.3	32.9	-26.8	45.6	27.6
RSK1 domain2	18.1	-12.7	55.8	2.7	5.0	-3.4
RSK1,RSK2,RSK3 domain1	18.7	3	13.4	-0.4	48.5	5.0
RSK2 domain1	5.6	-6.8	-11.2	-6	61.6	13.2
RSK2 domain2	9.1	0.6	21.4	-10	2.8	-4.1
RSKL1 domain1	-0.2	5.5	-1.8	1.1	-30.6	-15.9
SGK3	9.9	2.5	-0.8	-5.5	16.9	8.5
SLK	50.6	22.9	60.7	10.5	39.1	22.5

SMG1	-19.2	-1.3	13	-2.5	-6.1	1.5
smMLCK	-1.9	-11.5	16.2	-21.6	3.5	1.7
SRC	9.7	1.5	-2.2	-18.3	1.1	-1.0
STLK5	-46.5	-1.5	1.2	-15.4	-11.5	-6.9
SYK	-6.1	-8.9	7.2	-8.3	-5.1	-2.0
TAO1,TAO3	-14.9	-24	5.9	-6.5	-16.5	-9.7
TAO2	-24.2	-6.3	-2.5	-12.7	-18.4	-20.2
TBK1	13.4	0.1	16.4	7.3	0.1	1.9
TLK1	2.7	7.8	-3.7	-4.6	16.7	7.6
TLK2	-21.1	-5.8	-9.4	-8.4	16.0	11.1
ULK3	8	-1.9	5.2	3.1	17.4	-8.6
Wnk1,Wnk2,Wnk3	-24	-18.8	-6	1.7	-21.1	-6.3
Wnk1,Wnk2,Wnk4	-13.5	-15	-1.3	-1.9	-5.4	2.2
ZAK	23.1	15	-42.2	-20.5	-1.2	-2.9
ZAP70	3.1	-23.2	6.8	-4.9	-6.0	-25.0
ZC1/HGK,ZC2/TNIIK,ZC3/MINK	4.9	-18	14.8	-3.2	7.7	-1.2
ZC2/TNIIK	31.2	-10.6	29.2	-7.9	3.9	-10.7

CLAIMS

We claim:

1. A compound of Formula I



wherein

R^1 and R^2 taken together with the nitrogen to which they are attached are morpholin-4-yl optionally substituted with a methyl or 5-methyl-1,2,4-oxadiazol-3-yl;

R^3 is phenyl or a five to ten membered heteroaryl which contains one to four heteroatoms selected from N, O and S; wherein the phenyl and five to ten membered heteroaryl are optionally substituted with one to three R^9 and wherein the phenyl is optionally fused with a C_5 - C_6 cycloalkyl or a five to six membered heterocycloalkyl which contains one to three heteroatoms selected from N, O and S and which is optionally substituted with oxo;


R^4 and R^5 are each hydrogen; and

R^9 at each occurrence is independently cyano, halo, hydroxy, C_1 - C_3 alkyl-S-, - CO_2H , - $C(O)NH_2$, - $S(O)_2NH_2$, C_1 - C_3 alkyl optionally substituted with one to three halo or hydroxy, or C_1 - C_3 alkoxy optionally substituted with one to three halo or hydroxy; or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1 wherein R^3 is phenyl substituted with one or two R^9 ; and each R^9 is independently selected from cyano, fluoro, chloro or methoxy; or a pharmaceutically acceptable salt thereof.


3. The compound of claim 1 wherein R^3 is pyrazolyl, isothiazolyl, or pyridinyl, each optionally substituted with one R^9 ; and R^9 is cyano or methyl; or a pharmaceutically acceptable salt thereof.

4. A compound of Formula I



wherein


R^1 and R^2 taken together with the nitrogen to which they are attached is selected from the group consisting of:



R³ is 1-methylpyrazol-4-yl, 1*H*-pyrazol-4-yl, 2-cyanopyridin-6-yl, 3-methyl-1,2-thiazol-5-yl, 5-cyano-1-methylpyrrol-3-yl, 3-methylpyridin-5-yl, 3-cyanophenyl, 2-fluorophenyl, 3-fluorophenyl, 2,3-difluorophenyl, 2-fluoro-5-chlorophenyl, 2-fluoro-3-cyanophenyl or 2-methoxy-5-fluorophenyl; and


R⁴ and R⁵ are each independently hydrogen or methyl;
or a pharmaceutically acceptable salt thereof.

5. The compound of claim 4 wherein R¹ and R² taken together with the nitrogen to which they are attached are a group selected from



and R⁴ and R⁵ are each hydrogen; or a pharmaceutically acceptable salt thereof.

6. The compound of claim 5 wherein R¹ and R² taken together with the nitrogen to which they are attached is




or a pharmaceutically acceptable salt thereof.

7. The compound of claim 6 wherein

R³ is 3-cyanophenyl, 1-methylpyrazol-4-yl or 5-cyano-1-methylpyrrol-3-yl;
or a pharmaceutically acceptable salt thereof.

8. A compound of Formula I



or a pharmaceutically acceptable salt thereof wherein

R^1 and R^2 taken together with the nitrogen to which they are attached are piperidiny1 or 3-hydroxypiperidiny1;

R^3 is 1-methylpyrazol-4-yl, 1*H*-pyrazol-4-yl, 2-cyanopyridin-6-yl, 3-methyl-1,2-thiazol-5-yl, 3-methylpyridin-5-yl, 3-cyanophenyl, 2-fluorophenyl, 3-fluorophenyl, 2,3-difluorophenyl, 2-fluoro-5-chlorophenyl, 2-fluoro-3-cyanophenyl or 2-methoxy-5-fluorophenyl; and

R^4 and R^5 are each independently hydrogen or methyl.

9. A compound selected from the group consisting of:

5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
3-[6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile;
3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
{3-[4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
3-[2-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
{3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
3-[4-(3,3-dimethylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
3-[4-(piperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
1-(5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl)piperidine-3-carboxamide;
3-{4-[(3*S*)-3-hydroxypiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

4-[(3S)-3-methylpiperidin-1-yl]-5-phenyl-7H-pyrrolo[2,3-d]pyrimidine;
 3-{4-[(3S)-3-methylpiperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 4-[(3S)-3-methylpiperidin-1-yl]-5-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine;
 5-phenyl-4-(pyrrolidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidine;
 [1-(5-phenyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl)piperidin-4-yl]methanol;
 1-{5-[3-(hydroxymethyl)phenyl]-7H-pyrrolo[2,3-d]pyrimidin-4-yl}piperidine-3-carbonitrile;
 1-[5-(3-cyanophenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]piperidine-3-carbonitrile;
 4-(3,5-cis-dimethylpiperidin-1-yl)-5-(1-methyl-1H-pyrazol-4-yl)-7H-pyrrolo[2,3-d]pyrimidine;
 4-methoxy-3-[4-(3-methylpiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-[4-(dimethylamino)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]-4-methoxybenzonitrile;
 5-(5-chloro-2-methoxyphenyl)-N,N-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine;
 3-{4-[4-(1H-imidazol-2-yl)piperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-{4-[3-(methoxymethyl)piperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-[4-(9-methyl-1-oxa-4,9-diazaspiro[5.5]undec-4-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-[4-(3-methoxypiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-{4-[2-(methoxymethyl)morpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 N³-[5-(3-cyanophenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-N,N-dimethyl-beta-alaninamide;
 3-[4-(4,4-difluoropiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-[4-(4-fluoropiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-{4-[4-(1H-pyrazol-3-yl)piperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-{4-[(3S)-3-fluoropyrrolidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-{4-[2-(1H-pyrazol-3-yl)morpholin-4-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-{4-[3-(1H-pyrazol-3-yl)piperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-{4-[3-(1H-imidazol-2-yl)piperidin-1-yl]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-{4-[(1-methylpiperidin-3-yl)amino]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-[4-(3-oxo-2,7-diazaspiro[4.5]dec-7-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-[4-((3R)-3-methylpiperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;
 3-{4-[[2-(morpholin-4-yl)ethyl]amino]-7H-pyrrolo[2,3-d]pyrimidin-5-yl}benzonitrile;
 3-[4-(2-oxa-7-azaspiro[4.5]dec-7-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl]benzonitrile;

N-{1-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-4-yl}acetamide;
 5-(1*H*-indazol-5-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzenesulfonamide;
 5-(2-fluorophenyl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(1*H*-indazol-4-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(6-fluoro-5-methylpyridin-3-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}-2,3-dihydro-1*H*-isoindol-1-one;
 4-[(3*S*)-3-methylpiperidin-1-yl]-5-(pyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-[(3*S*)-3-methylpiperidin-1-yl]-5-(5-methylpyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-[(3*S*)-3-methylpiperidin-1-yl]-5-{7*H*-pyrrolo[2,3-*b*]pyridin-5-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 6-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}-2,3-dihydro-1*H*-isoindol-1-one;
 4-[(3*S*)-3-methylpiperidin-1-yl]-5-(1*H*-pyrrolo[3,2-*b*]pyridin-6-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}phenol;
 4-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzamide;
 3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}phenol;
 5-(2-chloro-5-methylpyridin-3-yl)-4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzamide;
 3-[4-(3-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(4-methylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzoic acid;
 3-[4-(methylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(3,5-*cis*-dimethylpiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-4-methoxybenzonitrile;
 4-methoxy-3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-{4-[(3*S*)-3-methylpiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
N,N-dimethyl-5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2-fluorobenzonitrile;

3-[4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-((3*R*)-3-methylpyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-4-fluorobenzonitrile;
 3-[4-(diethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*R*)-2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-5-fluorobenzonitrile;
 3-{4-[2-(1*H*-pyrazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-[4-((3*S*)-3-methylpyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*R*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-(4-fluoropiperidin-1-yl)-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(3-chlorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluoro-5-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2,5-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2,3-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-chloro-2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 {2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 5-(2,4-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3,5-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(morpholin-4-yl)-5-phenyl-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-chlorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methylphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 {2-fluoro-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 5-(4-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

{3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]phenyl}methanol;
 5-(2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-[3-(methylsulfonyl)phenyl]-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(morpholin-4-yl)-5-(pyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]isoquinoline;
 5-(5-bromopyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-chloro-5-methylphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-methylphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(1-methyl-1*H*-pyrazol-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[4-(4-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-methyl-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-[4-(6-oxa-3-azabicyclo[3.1.1]hept-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-chloro-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-methoxy-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5-chloro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 6-methyl-5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-methoxy-5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(thiomorpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-[5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 4-[(2*S*)-2-methylmorpholin-4-yl]-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-[(2*R*)-2-methylmorpholin-4-yl]-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(3-fluoropiperidin-1-yl)-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 {4-[5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]morpholin-2-yl}methanol;
 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(2-[[[6-methylpyridin-3-yl]oxy]methyl]morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
N,N-dimethyl-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;

N-cyclopropyl-*N*-methyl-5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;

3-[4-(3,3-difluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-[4-(3-aminopiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-[4-(2-[[5-(difluoromethyl)-1,2,4-oxadiazol-3-yl]methyl]morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-{4-[2-(1,3-thiazol-2-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-[4-(4-oxopiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-{4-[4-fluoro-4-(hydroxymethyl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-{4-[2-(3-hydroxyphenyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-{4-[2-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-[4-(3,3-difluoropyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-(4-{2-[(3-cyanophenoxy)methyl]morpholin-4-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile;

3-[4-(3-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

1-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidine-3-carboxamide;

3-[4-(2-ethylmorpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-{4-[2-(pyrimidin-4-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-{4-[2-(3-cyclopropyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-(4-{2-[(dimethylamino)methyl]morpholin-4-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile;

3-[4-(1*H*-pyrazol-4-ylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

3-(4-{2-[[5-methyl-1,2,4-oxadiazol-3-yl]methyl]morpholin-4-yl}-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)benzonitrile;

3-{4-[3-(5-methyl-1,3,4-oxadiazol-2-yl)piperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-5-methoxybenzonitrile;

5-(1-ethyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

5-(5-methylpyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

5-(5-chloropyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;

5-(6-methoxypyrazin-2-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(1,3-dimethyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[(3*S*)-3-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-[2-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(3*R*)-3-hydroxypiperidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 1-[5-(3-fluorophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 5-(5-fluoro-2-methoxyphenyl)-4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-[4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-(4-fluoropiperidin-1-yl)-5-(2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluoro-5-methoxyphenyl)-4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[4-(4,4-difluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2-fluorobenzonitrile;
 1-[5-(5-methylpyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 5-(5-fluoro-2-methoxyphenyl)-4-(4-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-[5-(3-fluoro-5-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 1-[5-(2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 2-fluoro-3-{4-[2-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-{4-[2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-(azetidin-1-yl)-5-(5-fluoro-2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-[(2*S*)-2-methylmorpholin-4-yl]-5-(5-methylpyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluorophenyl)-4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluorophenyl)-4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-methoxyphenyl)-4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 2-fluoro-3-{4-[(3*S*)-3-fluoropyrrolidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-[(3*S*)-3-fluoropyrrolidin-1-yl]-5-(2-methoxyphenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-methoxyphenyl)-4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluoro-5-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;

2-fluoro-3-[4-(3-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5-fluoro-2-methoxyphenyl)-4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 5-(5-fluoro-2-methoxyphenyl)-4-(3-fluoropiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluorophenyl)-4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 2-fluoro-3-[4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5-fluoro-2-methoxyphenyl)-4-[(3*S*)-3-fluoropyrrolidin-1-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-{4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 5-(4-methyl-1,3-thiazol-2-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(4-methylpyridin-2-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-fluoro-6-methoxyphenyl)-*N,N*-dimethyl-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-amine;
 5-(2-fluoro-6-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2,6-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-methyl-1,2-thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2-chloro-3-fluoro-6-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(4-methoxypyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[2-((5*R*)-5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[2-((5*S*)-5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-{4-[(2*S*)-2-(methoxymethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

or a pharmaceutically acceptable salt thereof.

10. A compound selected from the group consisting of:

5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[6-methyl-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile;
 3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;

4-(morpholin-4-yl)-5-(1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[4-(piperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(2,3-difluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-chloro-2-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3-fluorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-[5-(1-methyl-1*H*-pyrazol-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]piperidin-3-ol;
 5-(5-methylpyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-[4-(3-hydroxypiperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-fluoro-3-[4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-fluoro-3-[4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(3-methyl-1,2-thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine; and
 3-[4-[2-(5-methyl-1,2,4-oxadiazol-3-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 or a pharmaceutically acceptable salt thereof.

11. A compound selected from the group consisting of:

3-[6-(difluoromethyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(5,6-dihydro-2*H*-pyran-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(3,4-dihydro-2*H*-pyran-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 4-(morpholin-4-yl)-5-[3-(1,2,4-oxadiazol-3-yl)phenyl]-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-[4-[2-(3-methyl-1,2-oxazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-methyl-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridin-2(1*H*)-one;
 5-(imidazo[2,1-*b*][1,3]thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
rel-3-[4-[(3*aS*,6*aS*)-hexahydro-5*H*-furo[2,3-*c*]pyrrol-5-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;

rel-3-{4-[(3*aR*,6*aS*)-tetrahydro-1*H*-furo[3,4-*c*]pyrrol-5(3*H*)-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
rel-3-{4-[(4*aR*,7*aS*)-hexahydrocyclopenta[*b*][1,4]oxazin-4(4*aH*)-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 4-[5-(3-cyanophenyl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl]morpholine-2-carbonitrile;
 3-[4-(2,2-dimethylmorpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]thiophene-2-carbonitrile;
 5-(imidazo[1,2-*b*]pyridazin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 2-fluoro-3-{4-[2(*R*)-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 2-fluoro-3-{4-[2(*S*)-(3-methyl-1,2,4-oxadiazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carboxamide;
 4-(morpholin-4-yl)-5-(pyrazolo[1,5-*a*]pyrimidin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1-methyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrrole-2-carbonitrile;
 5-(6-methylimidazo[2,1-*b*][1,3]thiazol-5-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 3-{4-[2-(1,2-oxazol-5-yl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 1-methyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-imidazole-2-carbonitrile;
 4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]thiophene-2-carbonitrile;
 4-(morpholin-4-yl)-5-(pyrazolo[1,5-*a*]pyridin-3-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 1,5-dimethyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrrole-2-carbonitrile;
 1-methyl-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrazole-5-carbonitrile; and
 3-{4-[2-(cyanomethyl)morpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 or a pharmaceutically acceptable salt thereof.

12. A compound selected from the group consisting of
 5-(1-methyl-1*H*-pyrazol-4-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 6-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]pyridine-2-carbonitrile;
 3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-{4-[(2*S*)-2-methylmorpholin-4-yl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl}benzonitrile;
 3-[4-(piperidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 3-[4-(dimethylamino)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-2-fluorobenzonitrile;
 3-[4-(pyrrolidin-1-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 2-fluoro-3-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]benzonitrile;
 5-(3-chlorophenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-fluoro-2-methoxyphenyl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(5-bromopyridin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine;
 5-(imidazo[1,2-*b*]pyridazin-3-yl)-4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidine; and
 1-methyl-4-[4-(morpholin-4-yl)-7*H*-pyrrolo[2,3-*d*]pyrimidin-5-yl]-1*H*-pyrrole-2-carbonitrile;
 or a pharmaceutically acceptable salt thereof.

13. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to any one of claims 1 to 12, or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier.

14. Use of a compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 12 in the manufacture of a medicament for treating Parkinson's disease in a patient.