LIQUID DEPOT FOR NON-INVASIVE SUSTAINED DELIVERY OF AGENTS TO THE EYE

RELATED APPLICATION

[0001] This Application claims priority benefit of U.S. Provisional Application No. 62/665,367, filed May 1, 2018, which is incorporated fully herein for all purposes.

[0002] The present embodiments provide compositions and methods for the treatment of ocular conditions or maladies.

BACKGROUND

[0003] There remains a need for noninvasive, sustained delivery of pharmaceutical agents to tissues and liquid portions of the eye, such as the conjunctiva, cornea, or aqueous humor.

SUMMARY

[0003a] In one aspect, the present disclosure refers to a liquid depot for sustained release of an active agent in the eye, the liquid depot comprising: a tocopherol; an ocular film-forming excipient which are medium-chain triglycerides, wherein the tocopherol:ocular film-forming excipient is at a weight ratio of 90:10 to 60:40; and an active agent, which is a corticosteroid; wherein the liquid depot has a viscosity of 850 cP to 1100 cP, inclusive and water has a viscosity of 1.00 cP.

[0003b] In another aspect, the present disclosure refers to a kit comprising at least one single-use dispenser, wherein the at least single-use dispenser comprises the liquid depot disclosed herein.

[0003c] In yet another aspect, the present disclosure refers to a use of a liquid depot comprising (a) at least one active agent, which is a corticosteroid; and (b) a mixture of tocopheryl acetate:medium-chain triglycerides at a weight ratio of 90:10 to 60:40, in the manufacture of a medicament for treating a malady of the eye in a subject, wherein the medicament is to be topically administered to the eye of the subject, wherein the liquid depot has a viscosity of 850 cP to 1100 cP.

CLAIMS

1. A liquid depot for sustained release of an active agent in the eye, the liquid depot comprising: a tocopherol; an ocular film-forming excipient which are medium-chain triglycerides, wherein the tocopherol:ocular film-forming excipient is at a weight ratio of 90:10 to 60:40; and an active agent, which is a corticosteroid; wherein the liquid depot has a viscosity of 850 cP to 1100 cP, inclusive and water has a viscosity of 1.00 cP.

2. The liquid depot of claim 1, wherein the tocopherol is α-, β-, γ- or δ-tocopherol, or α-, β-, γ- or δ-tocotrienol.

3. The liquid depot of any one of claims 1-2, wherein the tocopherol is an α-, β-, γ-, or δ-tocopherol.

4. The liquid depot of any one of claims 1-3, wherein the tocopherol is tocopheryl acetate.

5. The liquid depot of any one of claims 1-4, wherein the medium-chain triglycerides is a member selected from the group consisting of decanoyl/octanoyl glycerides (MIGLYOL® 810) or caprylic/capric triglyceride (MIGLYOL® 812).

6. The liquid depot of claim 1, wherein the corticosteroid is a member selected from the group consisting of dexamethasone, prednisolone, prednisone, loteprednol, triamcinolone and fluorometholone.

7. The liquid depot of claim 6, wherein the liquid depot comprises (a) 10% dexamethasone; and (b) 90% of a mixture of tocopheryl acetate:medium-chain triglycerides at a ratio of 90:10 to 60:40.

8. The liquid depot of claim 6, wherein the liquid depot comprises, on a wt% basis, (a) 10% prednisolone and (b) 90% of a mixture of tocopheryl acetate:medium-chain triglycerides at wt/wt a ratio of 80:20.

9. The liquid depot of claim 6, wherein the liquid depot comprises, on a wt% basis, (a) 10% loteprednol and (b) 90% of a mixture of tocopheryl acetate:medium-chain triglycerides at wt/wt a ratio of 80:20.

10. The liquid depot of any one claim claims 1-5, comprising dexamethasone and moxifloxacin in a mixture of tocopheryl acetate:medium-chain triglycerides at a ratio of 90:10 to 60:40.

11. The liquid depot of any one of the preceding claims, wherein the liquid depot is disposed within a single-use dispenser.

12. A kit comprising at least one single-use dispenser, wherein the at least single-use dispenser comprises the liquid depot of any one of claims 1-10.

13. Use of a liquid depot comprising (a) at least one active agent, which is a corticosteroid; and (b) a mixture of tocopheryl acetate:medium-chain triglycerides at a weight ratio of 90:10 to 60:40, in the manufacture of a medicament for treating a malady of the eye in a subject, wherein the medicament is to be topically administered to the eye of the subject, wherein the liquid depot has a viscosity of 850 cP to 1100 cP.

14. The use of claim 13, wherein the medium-chain triglycerides is at least one MIGLYOL® neutral oil.

15. The use of claim 13 or 14, wherein the malady of the eye is inflammation, infection, allergic reaction, or glaucoma.

16. The use of claim 13 or 14, wherein the malady of the eye is infection, corneal abrasion or other trauma to the eye surface, blepharitis, inflammatory meibomian gland disease, meibomian gland dysfunction, allergic conjunctivitis, chronic conjunctivitis, Sjögren's syndrome, dry eye (keratoconjunctivitis sicca), aqueous-deficient dry eye, mucin-deficient dry eye (punctate epithelial keratitis), episcleritis, keratitis (corneal ulcers), pterigia, Stevens-Johnson syndrome, ocular citatrical pemphigoid/mucous membrane pemphigoid, irregular cornea condition or other surface abnormalities, epitheliopathy, neurotrophic cornea, corneal

dystrophy such as Fuch's dystrophy, peripheral or marginal degeneration of the cornea, conjunctivochalasis, glaucoma, conjunctival degeneration such as pinguecula, or pingueculitis.

17. The use of any one of claims 13-16, wherein the medicament is to be intermittently administered.

18. The use of claim 17, wherein the medicament is to be intermittently administered for at least 24 hours, 48 hours, 72 hours (3 days), 4 days, 5 days, 6 days, 7 days (1 week), 14 days (2 weeks), or 21 days (3 weeks).

19. The use of claim 17, wherein the medicament is to be intermittently administered for at least 24 hours.