

T CELL WHICH EXPRESSES A GAMMA-DELTA T CELL RECEPTOR (TCR)
AND A CHIMERIC ANTIGEN RECEPTOR (CAR)

FIELD OF THE INVENTION

- 5 The present invention relates to immunotherapeutic T cells. In particular, the invention provides immunotherapeutic gamma-delta T cells comprising a chimeric antigen receptor (CAR).

BACKGROUND TO THE INVENTION

- 10 Chimeric antigen receptors (CARs) developed for cancer immunotherapy combine an extracellular antigen recognition domain with signalling domains specific for effector cells within a single molecule. The most common CAR system involves an antigen recognition domain derived from a monoclonal antibody fused to signalling domains
15 which provide activating signals for T cells.

Typically, the signalling domains of a CAR provides cytotoxicity, proliferation and survival signals to activate the effector cell upon binding of antigen to the antigen recognition domain (Signals 1 and 2).

- 20 A limitation of this technology is potential 'on target-off tumour toxicity'. This toxicity is caused by the recognition of low levels of a cancer-associated antigen recognised by a CAR on normal tissues. For instance GD2 is a target for neuroblastoma but also is expressed on nerves; and PSMA is a target for prostate cancer cells but is also found
25 on normal kidney, liver and colon cells, and brain astrocytes. This problem is more profound in solid tumours where there is a dearth of highly selective targets.

Thus there is a need for cancer immunotherapies which address the above problems.

- 30 SUMMARY OF ASPECTS OF THE INVENTION

- The present inventors have determined a mechanism of reducing 'on target-off tumour toxicity' by using CARs in gamma delta ($\gamma\delta$) T-cells. In the system described herein, a CAR is used to provide a co-stimulatory signal (signal 2) to a $\gamma\delta$ T-cell upon
35 binding of antigen to the antigen recognition domain of the CAR. In this way, signal 2 is only provided to the T-cell upon binding of the CAR to its target antigen (Figure 2A). Signal 1 for $\gamma\delta$ T-cell activation is provided by the endogenous TCR, which is activated by danger signals, such as phosphoantigens.

A $\gamma\delta$ T-cell requires both signal 1 and signal 2 for optimal effector function. Thus, in the present system the $\gamma\delta$ T-cell will only be fully activated for cytotoxicity, proliferation and cytokine secretion if the target cell: (i) expresses the antigen
5 recognised by the CAR; and (ii) expresses danger signals recognised by the endogenous $\gamma\delta$ TCR.

Thus, in a first aspect the present invention provides a T cell which expresses a gamma-delta T cell receptor (TCR) and a chimeric antigen receptor (CAR), wherein
10 the CAR comprises;

- (i) an antigen binding domain;
- (ii) a transmembrane domain; and
- (iii) a co-stimulatory intracellular signalling domain;

wherein the intracellular signalling domain provides a co-stimulatory signal to the T
15 cell following binding of antigen to the antigen binding domain.

As such, binding of a first antigen to the $\gamma\delta$ TCR results in signal 1 production and binding of a second antigen to the antigen binding domain of the CAR results in signal
20 2 production.

The antigen binding domain may be capable of binding to a tumour-associated antigen (TAA).

The antigen binding domain may be capable of binding to GD2, CD33, CD19 or
25 EGFR.

The intracellular signalling domain may comprise the DAP10, CD28, CD27, 41BB, OX40, CD30, IL2-R, IL7-R, IL21-R, NKp30, NKp44 or DNAM-1 (CD226) signalling
30 domain.

The transmembrane domain of the CAR may comprise a CD8 stalk or a CD28 transmembrane domain.

The intracellular signalling domain of the CAR may comprise the DAP10 signalling
35 domain.

The CAR may further comprise a spacer domain between the antigen binding domain and the transmembrane domain.

5 The $\gamma\delta$ TCR may be capable of binding to a phosphoantigen/butyrophilin 3A1 complex; major histocompatibility complex class I chain-related A (MICA); major histocompatibility complex class I chain-related B (MICB); NKG2D ligand 1-6 (ULBP 1-6); CD1c; CD1d; endothelial protein C receptor (EPCR); lipohexapeptides; phycoerythrin or histidyl-tRNA-synthase.

10 The CAR may comprise one of the following amino acid sequences:

SEQ ID NO: 1 (aCD33-Fc-DAP10 CAR)

15 MAVPTQVLGLLLLWLT DARCDIQMTQSPSSLSASVGDRVTITCRASEDIYFNLVWYQ
QKPGKAPKLLIYDTNRLADGVPSRFSGSGSGTQYTLTISSLQPEDFATYYCQHYKNY
PLTFGQGKLEIKRSGGGGSGGGGSGGGGSGGGGSRSEVQLVESGGGLVQPGG
SLRLSCAASGFTLSNYGMHWIRQAPGKGLEWSSISLNGGSTYYRDSVKGRFTISR
DNAKSTLYLQMNSLRAEDTAVYYCAAQDAYTGGYFDYWQGQGLTVTVSSMDPAEPK
SPDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMIARTPEVTCVVVDVSHEDPEVKFN
20 WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI
EKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPE
NNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS
PGKKDKPFWLVVVGGLACYSLLVTVAFIIFWVCARPRRSPAQEDGKVYINMPGR
G

25

SEQ ID NO: 2 (aGD2-Fc-DAP10 CAR)

30 METDTLLLWLLLWPGSTGQVQLQESGPGLVKPSQTLTCTVSGFSLASYNHWW
RQPPGKGLEWLGVIWAGGSTNYNSALMSRLTISKDNSKNQVFLKMSSLTAADTAVY
YCAKRSDDYSWFAYWGQGLTVTVSSGGGGSGGGGSGGGGSENQMTQSPSSLSA
SVGDRVTMTCRASSSVSSSYLHWYQQKSGKAPKVIYSTSNLASGVPSRFSGSGS
GTDYTLTISSLQPEDFATYYCQQYSGYPITFGQGTKVEIKRSDPAEPKSPDKTHTCP
PCPAPPVAGPSVFLFPPKPKDTLMIARTPEVTCVVVDVSHEDPEVKFNWYVDGVEV
HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG
35 QPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPV
LDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKKDKPKF
WLVVVGGLACYSLLVTVAFIIFWVCARPRRSPAQEDGKVYINMPGRG

In a further aspect the present invention provides a CAR comprising; (i) an antigen-binding domain; (ii) a transmembrane domain; and (iii) an intracellular signalling domain; wherein the intracellular signalling domain comprises a co-stimulatory intracellular signalling domain but does not comprise a CD3 endodomain.

5

The co-stimulatory intracellular signalling domain may be selected from a DAP10, CD28, CD27, 41BB, OX40, CD30, IL2-R, IL7-R, IL21-R, NKp30, NKp44 or DNAM-1 (CD226) signalling domain.

10 In a second aspect the present invention provides a CAR comprising, an antigen-binding domain; a transmembrane domain; and an intracellular signalling domain; wherein the intracellular signalling domain comprises a DAP10 signalling domain. The intracellular signalling domain may consist of or consist essentially of a DAP10 signalling domain.

15

In a particular embodiment the intracellular signalling domain of the CAR according to the second aspect of the invention does not comprise a CD3 endodomain.

The CAR according to the second aspect of the invention may be a CAR as defined
20 in the first aspect of the invention.

In a third aspect the present invention provides a nucleic acid sequence encoding a CAR as defined in the first or second aspects of the invention.

25 In a fourth aspect the present invention provides a vector comprising a nucleic acid sequence as defined by the third aspect of the invention.

The vector may be a retroviral vector, a lentiviral vector or a transposon.

30 In a fifth aspect the present invention relates to method for making a cell according to the first aspect of the invention, which comprises the step of introducing; a nucleic acid sequence according to the third aspect of the invention or a vector according to fourth aspect of the invention into a cell.

35 The method may comprise the step of stimulating the cell with a gamma delta T cell stimulating agent.

The $\gamma\delta$ T cell stimulating agent may be selected from, for example, isopentenyl pyrophosphate (IPP); analogs of IPP such as bromohydrin pyrophosphate and (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate; and inhibitors of farnesyl pyrophosphate synthase (FPPS) such as aminobisphosphonates (e.g. zoledronate or pamidronate).

5

The cell may be from a sample isolated from a subject.

In a sixth aspect the present invention provides a pharmaceutical composition comprising a cell according to the first aspect of the present invention.

10

In a seventh aspect the present invention relates to a method for treating a disease, which comprises the step of administering a pharmaceutical composition according to the sixth aspect of the invention to a subject.

15 The method may comprise the step of administering a $\gamma\delta$ T cell stimulating agent to the subject.

The $\gamma\delta$ T cell stimulating agent may be selected from, for example, isopentenyl pyrophosphate (IPP); analogs of IPP such as bromohydrin pyrophosphate and (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate; and inhibitors of farnesyl pyrophosphate synthase (FPPS) such as aminobisphosphonates (e.g. zoledronate or pamidronate).

20 The method may comprise the following steps:

- (i) isolation of a cell-containing sample from a subject;
- 25 (ii) transduction or transfection of cells with: a nucleic acid sample according to the third aspect of the present invention or a vector according to the fourth aspect of the present invention; and
- (iii) administering the cells from (ii) to the subject.

30 In an eighth aspect the present invention relates to a pharmaceutical composition according to the sixth aspect of the present invention for use in treating a disease.

In a ninth aspect the present invention relates to the use of a cell according to the first aspect of the present invention in the manufacture of a medicament for treating and/or preventing a disease.

35

The disease described herein may be cancer, microbial infection or viral infection.

The present invention therefore provides a $\gamma\delta$ T cell which is only fully activated by, and therefore capable of killing, a target cell which expresses a first antigen which is capable of binding to the endogenous $\gamma\delta$ TCR (and thus stimulating productive signal 1) and a second antigen which is capable of binding to the CAR (and thus stimulating productive signal 2).

The $\gamma\delta$ T cells of the invention are therefore useful for reducing unwanted 'on target-off tumour' effects. In particular, a normal cell which expresses low levels of a TAA will not activate the $\gamma\delta$ T cell of the invention as it will not express a danger signal recognised by the endogenous $\gamma\delta$ TCR and thus will not provide signal 1, which is required for full activation of the $\gamma\delta$ T cell.

DESCRIPTION OF THE FIGURES

Figure 1 – Diagram of the signalling required for full activation of a $\gamma\delta$ T cell which results in killing of the target cell. A) and B) Signalling via the $\gamma\delta$ TCR or co-receptors alone does not result in full activation of the $\gamma\delta$ T cell. C) A combination of $\gamma\delta$ TCR and co-receptor signalling results in full activation of the $\gamma\delta$ T cell

Figure 2 – Illustrative diagram of a $\gamma\delta$ T cell of the present invention. A) Normal activation of a $\gamma\delta$ T cell by a target cell. B) Blocking of signal 2 by soluble NKG2D ligands secreted by cancer cells prevents full activation of $\gamma\delta$ T cells. C) Full activation of a $\gamma\delta$ T cell of the present invention by a transformed cell. D) Normal healthy cells do not express danger signals recognised by endogenous $\gamma\delta$ T cell receptors and do not fully activate $\gamma\delta$ T cells of the present invention.

Figure 3 – Examples of illustrative CARs which may be used in the present invention

Figure 4 – Representative flow cytometric dot plots to illustrate co-expression of a $\gamma\delta$ TCR (V δ 2) and GD2-DAP10 CAR (Fc, CD20 marker and CD34 marker) in a $\gamma\delta$ T cell

Figure 5 - Killing of GD2+ cell lines LAN1 and TC71 by V δ 2 $\gamma\delta$ T cells transduced with the aGD2-Fc-DAP10 CAR

(A) Significant killing of GD2+ neuroblastoma cell line LAN1 is only seen when CAR transduced cells are used and not when non-transduced (NT) V δ 2 are used as effectors. (B) Additive effect of aGD2-Fc-DAP10 CAR when combined with 24h

zoledronic acid exposure which increases phosphoantigen production, against the GD2+ Ewing sarcoma cell line TC71. **(C)** Addition of the CAR to $\alpha\beta$ T cells, which lack the signal 1 provided by the $\gamma\delta$ TCR in response to cellular stress, has no effect on cytotoxicity, unlike the effect of the CAR in V δ 2+ $\gamma\delta$ T cells. This indicates that the CAR signal alone is insufficient for T-cell activation. Error bars denote SEM for 3-6 independent donors.

Figure 6 – Killing of GD2+ cell line LAN1 and no killing of GD2- cell line SKNSH. Error bars denote SEM for 3-6 independent donors.

10

Figure 7 - Preservation of CAR expression following prolonged co-culture and GD2 specific expansion

(A) Co-culture was started 24 days after transduction (labelled D0). Serial analyses of cells for presence of CAR (Y axis) and TCRV δ 2 (X axis) were taken in the presence of irradiated GD2+ (LAN1) and GD2- (SK-N-SH) neuroblastoma cells. Representative data from 1 of 3 donors is shown. **(B)** Expansion of aGD2-Fc-DAP10 transduced V δ 2+ cells was only seen in the presence of irradiated GD2+ target cells (graphical representation, n=3 independent donors, error bars denote SEM).

Figure 8 – Flow cytometric staining for CD33 expression of AML cell lines (Nomo1, Sh1 and MV4;11) and freshly isolated monocytes is equivalent.

Figure 9 – A) aCD33-DAP10-transduced V δ 2 cells spare monocytes in the absence of ZOL but aCD33-CD28z-transduced V δ 2 cells do not. B) aCD33-DAP10-transduced V δ 2 cells kill AML better than NT V δ 2 cells, but spare monocytes. Error bars indicate SEM for 3 independent donors.

Figure 10 - Nucleic acid and amino acid sequences of an anti-GD2-Fc-DAP10 CAR

Figure 11 - Nucleic acid and amino acid sequences of an anti-CD33-Fc-DAP10 CAR

Figure 12 - aCD33-DAP10-transduced V δ 2 cells spare haemopoietic stem cells but aCD33-CD28z-transduced V δ 2 cells do not. Normal human bone marrow was cultured overnight with the indicated CAR T cells. Surviving haemopoietic stem cells were assayed by myeloid colony formation in soft agar. Data is derived using transduced V δ 2 cells from three independent donors.

Figure 13 - Differential cross-linking of "costimulation-only" CAR and V γ 9v δ 2 TCR leads to differential cytokine responses. Top; Schematic of experimental design. Biotinylated beads are coated with (A) no/irrelevant antibodies, or (B) antibodies to bind either the TCR (anti-CD3) or the CAR (anti-Ig binding the spacer region of the CAR); C) following cross linking, intracellular cytokine secretion is used to measure activation. As a control, stimulatory anti-CD3/CD28 beads (Miltenyi) are used. Bottom-left: representative FACS plots; bottom-right: cytokine responses to cross linking show that the "costimulation-only" CAR cross linking leads to a TNF- α response but that additional TCR engagement is required for full response comprising both interferon gamma and TNF- α . Data is means $\pm$ SD of 5 donors.

DETAILED DESCRIPTION

$\gamma\delta$ T CELL

T-cells are divided into two groups based on their T-Cell Receptor (TCR) components. The TCR heterodimer consists of an α and β chain in 95% of T cells. These recognise foreign antigens via peptides presented by MHC molecules on antigen presenting cells and are essential for adaptive immunity.

5% of T cells have TCRs consisting of γ and δ chains. $\gamma\delta$ TCRs are MHC independent and detect markers of cellular stress expressed by tumours.

$\gamma\delta$ T cells recognize pathogens and transformed cells in an HLA-unrestricted manner. They respond to markers of cellular stress (e.g. phosphoantigens released by transformed cells as by-products of the mevalonate biosynthetic pathway). $\gamma\delta$ T cells display both innate cytotoxic functions and antigen-presenting capability, particularly in the presence of antibody-opsonized target cells.

$\gamma\delta$ T-cells are responsible for "lymphoid stress surveillance," i.e., sensing and responding immediately to infections or non-microbial stress without the need of clonal expansion or *de novo* differentiation.

The activation of $\gamma\delta$ T cells is regulated by a balance between stimulatory and inhibitory signals. They are activated by $\gamma\delta$ TCR ligands (e.g. phosphoantigens) in combination with MHC-associated ligands of the activatory receptor killer cell lectin-like receptor subfamily K, member 1 (KLRK1), also known as NKG2D, such as MHC

class I polypeptide-related sequence A (MICA), MICB, and various members of the UL16-binding protein (ULBP) family.

5 $\gamma\delta$ cells also express killer-cell immunoglobulin-like receptors (KIRs), which can be either activatory or inhibitory, including killer cell immunoglobulin-like receptor, 2 domains, long cytoplasmic tail, 1 (KIR2DL1) and killer cell immunoglobulin-like receptor, 3 domains, long cytoplasmic tail, 1 (KIR3DL1).

10 Full activation of a $\gamma\delta$ T cell which results in the effective killing of a target cell requires productive signal 1 and signal 2 generation (Figures 1 and 2A).

15 $\gamma\delta$ T-cells derive signal 1 of T cell activation from danger signal antigens present on transformed or infected cells. These danger signal antigens are recognised through the $\gamma\delta$ TCR. Signal 2 of T cell activation for $\gamma\delta$ T-cells is also commonly derived by danger signal molecules (such as MICA) present on transformed or infected cells. Signal 2 may be transduced, for example, through the NKG2D receptor and DAP 10 (Figure 2A).

20 As a means of avoiding immune detection, cancer cells frequently secrete soluble NKG2D ligands effectively blocking signal 2 in $\gamma\delta$ T-cells, thus preventing their activation and facilitating tumour infiltration (Figure 2B).

25 In a first aspect, the present invention provides a T cell which expresses a $\gamma\delta$ TCR and a CAR, wherein the intracellular signalling domain of the CAR provides a co-stimulatory signal to the T cell.

Thus, the arrangement of the $\gamma\delta$ TCR and the CAR is such that the $\gamma\delta$ TCR provides signal 1 and the CAR provides signal 2 upon binding to each receptor, respectively.

30 As used herein, co-stimulatory signal is synonymous with signal 2, which is required for full $\gamma\delta$ T cell activation.

35 Thus, a $\gamma\delta$ T cell according to the first aspect of the present invention will only be fully activated and capable of killing a target cell which expresses a first antigen which is capable of binding to the $\gamma\delta$ TCR (and thus stimulating productive signal 1) and a second antigen which is capable of binding to the CAR (and thus stimulating productive signal 2) (Figure 2C).

In the absence of antigen binding to the $\gamma\delta$ TCR, signal 1 is not generated and full $\gamma\delta$ T cell activation is not achieved. In other words, in the absence of antigen binding to the $\gamma\delta$ TCR, the $\gamma\delta$ T cell is not stimulated to kill the target cell (Figure 2D).

- 5 In the absence of antigen binding to the CAR, signal 2 is not generated and full $\gamma\delta$ T cell activation is not achieved. In other words, in the absence of antigen binding to the CAR, the $\gamma\delta$ T cell is not stimulated to kill the target cell.

The $\gamma\delta$ T cell of the present invention may express any $\gamma\delta$ TCR. Examples of $\gamma\delta$ TCR
10 ligands are known in the art (see Vantourout, P. & Hayday, A. Nat. Rev. Immunol. 13, 88–100 (2013), for example).

By way of example, the $\gamma\delta$ TCR expressed by a cell of the present invention may recognise phosphoantigens (e.g. Isopentenyl pyrophosphate (IPP), Bromohydrin
15 Pyrophosphate (BrHPP) and (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate (HMBPP)); major histocompatibility complex class I chain-related A (MICA); major histocompatibility complex class I chain-related B (MICB); NKG2D ligand 1-6 (ULBP 1-6); CD1c; CD1d; endothelial protein C receptor (EPCR); lipohexapeptides; phycoreythrins or histidyl-tRNA-synthase.

20

One advantage of the cell of the present invention is that it comprises a CAR comprising (i) an antigen binding domain which binds a specific antigen and (ii) a particular co-stimulatory endodomain. As such, the cell of the present invention will have a greater propensity towards activation in an environment comprising an antigen
25 which can be bound by the CAR, as the binding of antigen by the CAR will result in signalling through the co-stimulatory endodomain and signal 2 production. For example, if the antigen-binding domain of the CAR is specific for a TAA, the cell of the present invention will have an increased propensity towards activation in a tumour environment where the TAA is expressed due to the co-stimulatory signal provided by
30 the CAR.

CHIMERIC ANTIGEN RECEPTOR

The T cell according to the present invention expresses a chimeric antigen receptor
35 (CAR).

Chimeric antigen receptors (CARs) are engineered receptors which graft an arbitrary specificity onto an immune effector cell. In a classical CAR, the specificity of a monoclonal antibody is grafted on to a T cell. CAR-encoding nucleic acids may be transferred to T cells using, for example, retroviral vectors. In this way, a large number of cancer-specific T cells can be generated for adoptive cell transfer. Phase I clinical studies of this approach show efficacy.

The target-antigen binding domain of a CAR is commonly fused via a spacer and transmembrane domain to a signaling endodomain. When the CAR binds the target-antigen, this results in the transmission of an activating signal to the T-cell it is expressed on.

Early CAR designs had endodomains derived from the intracellular parts of either the γ chain of the Fc ϵ R1 or CD3 ζ . Consequently, these first generation receptors transmitted immunological signal 1, which was sufficient to trigger T-cell killing of cognate target cells but failed to fully activate the T-cell to proliferate and survive. To overcome this limitation, compound endodomains have been constructed: fusion of the intracellular part of a T-cell co-stimulatory molecule to that of CD3 ζ results in second generation receptors which can transmit an activating and co-stimulatory signal simultaneously after antigen recognition. The co-stimulatory domain most commonly used is that of CD28. This supplies the most potent co-stimulatory signal - namely immunological signal 2, which triggers T-cell proliferation. Some receptors have also been described which include TNF receptor family endodomains, such as the closely related OX40 and 41BB which transmit survival signals. Even more potent third generation CARs have now been described which have endodomains capable of transmitting activation, proliferation and survival signals.

The $\gamma\delta$ T cell of the present invention comprises a CAR which comprises a co-stimulatory signalling endodomain which transmits signal 2 to the $\gamma\delta$ T cell upon the binding of target antigen.

The CARs of the T cell of the present invention may comprise a signal peptide so that when the CAR is expressed inside a cell, such as a T-cell, the nascent protein is directed to the endoplasmic reticulum and subsequently to the cell surface, where it is expressed.

The core of the signal peptide may contain a long stretch of hydrophobic amino acids that has a tendency to form a single alpha-helix. The signal peptide may begin with a short positively charged stretch of amino acids, which helps to enforce proper topology of the polypeptide during translocation. At the end of the signal peptide
5 there is typically a stretch of amino acids that is recognized and cleaved by signal peptidase. Signal peptidase may cleave either during or after completion of translocation to generate a free signal peptide and a mature protein. The free signal peptides are then digested by specific proteases.

The signal peptide may be at the amino terminus of the molecule.

10 The signal peptide may comprise the SEQ ID NO: 6, 7 or 8 or a variant thereof having 5, 4, 3, 2 or 1 amino acid mutations (insertions, substitutions or additions) provided that the signal peptide still functions to cause cell surface expression of the CAR.

SEQ ID NO: 6: MGTSLLCWMALCLLGADHADG

The signal peptide of SEQ ID NO: 6 is compact and highly efficient. It is predicted to
15 give about 95% cleavage after the terminal glycine, giving efficient removal by signal peptidase.

SEQ ID NO: 7: MSLPVTALLPLALLHAARP

The signal peptide of SEQ ID NO: 7 is derived from IgG1.

SEQ ID NO: 8: MAVPTQVLGLLLLWLTDARC

20 The signal peptide of SEQ ID NO: 8 is derived from CD8.

CO-STIMULATORY INTRACELLULAR SIGNALLING DOMAIN

The intracellular domain/endodomain is the signal-transmission portion of a classical CAR.
25

The $\gamma\delta$ T cell of the present invention comprises a CAR which comprises a co-stimulatory signalling endodomain which transmits signal 2 to the $\gamma\delta$ T cell upon the binding of target antigen. Accordingly, $\gamma\delta$ T cell of the present invention comprises a CAR which does not transmit signal 1 to the $\gamma\delta$ T cell upon the binding of target
30 antigen.

T-cell costimulatory receptors are known to induce qualitative and quantitative changes that lower activation thresholds and prevent T cell anergy and enhance T cell function.

- 5 A number of co-receptors for $\gamma\delta$ T cells are known in the art. Productive signalling via one or more of these receptors can result in full activation of the $\gamma\delta$ T cell and target cell killing.

10 The $\gamma\delta$ T cell of the present invention comprises an intracellular signalling domain from a $\gamma\delta$ T cell co-receptor, such that binding of antigen to the antigen-binding domain of the CAR generates productive signal 2 signalling in the $\gamma\delta$ T cell.

The intracellular signalling domain may, for example, comprise the DAP10, CD28, CD27, 41BB, OX40, CD30, IL2-R, IL7-R, IL21-R, NKp30, NKp44 or DNAM-1 (CD226) signalling domain.

The intracellular signalling domain may comprise the DAP10 signalling domain.

20 DAP10 is a signalling subunit which associates with the NKG2D receptor (see Figure 1). It is the exclusive binding partner and signalling intermediate for NKG2D and contains a YxxM activation motif that triggers the lipid kinase cascade.

An example of an amino acid sequence for a DAP10 signalling domain is shown below:

25 SEQ ID NO: 3 – CARPRRSPAQEDGKVYINMPGRG

Further illustrative co-stimulatory domains are shown as SEQ ID NO: 9-19

30 SEQ ID NO: 9 (CD28 endodomain)
KRSRLLHSDYMNMTPRRPGPTRKHYPYAPPRDFAAY

SEQ ID NO: 10 (CD27 endodomain)
QRRKYRSNKGESPVEPAEPCHYSCPREEEGSTIPIQEDYRKPEPACSP

35 SEQ ID NO: 11 (41BB endodomain)
KRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL

SEQ ID NO: 12 (OX40 endodomain)

RRDQRLPPDAHKPPGGGSFRTPIQEEQADAHSTLAKI

SEQ ID NO: 13 (CD30 endodomain)

HRRACRKIRQKLHLCYPVQTSQPKLELVDSRPRRSSTQLRSGASVTEPVAEERGL

5 MSQPLMETCHSVGAAYLESLPLQDASPAGGPSSPRDLPEPRVSTEHTNNKIEKIYIM
KADTVIVGTVKAELPEGRGLAGPAEPELEEELEADHTPHYPEQETEPPLGSCSDVML
SVEEEGKEDPLPTAASGK

SEQ ID NO: 14 (IL2-R endodomain)

10 TWQRRQRKSRRTI

SEQ ID NO: 15 (IL7-R endodomain)

KKRIKPIVWPSLPDHKKTLEHLCKKPRKLNVSFNPESFLDCQIHRVDDIQRDEVEG

FLQDTFPQQLEESEKQRLGGDVQSPNCPSEDVVITPESFGRDSSLTCLAGNVSACD

15 APILSSSRSLDCRESGKNGPHVYQDLLLSLGTNSTLPPPFSLQSGILTLPVAQGGQ
PILTSLGSNQEEAYVTMSSFYQNG

SEQ ID NO: 16 (IL21-R endodomain)

SLKTHPLWRLWKKIWAVSPERFFMPLYKGCSDGDFKKWWGAPFTGSSLELGPWSP

20 EVPSTLEVYSCHPPRSPAKRLQLTELQEPALVESDGVKPSFWPTAQNSGGSAYS
EERDRPYGLVSDTVTVLDAEGPCTWPCSCEDDGYPALDLDAGLEPSPGLEDPLLD
AGTTVLSCGCVSAGSPGLGGPLGSLDLRKPPLADGEDWAGGLPWGGRSPGGVS
ESEAGSPLAGLMDTFD SGFVSGDCSSPVECDFTSPGDEGP PRSYLRQWWIPPP
LSSPGPQAS

25

SEQ ID NO: 17 (NKp30 endodomain)

GSTVYYQGKCLTWKGPRRQLPAVVPAPLPPPCGSSAHLPPVPGG

SEQ ID NO: 18 (NKp44 endodomain)

30 WWGDIWWKTMMELRSLDTQKATCHLQQVTDLPWTSVSSPVEREILYHTVARTKISD
DDDEHTL

SEQ ID NO: 19 (DNAM-1 (CD226) endodomain)

NRRRRRRERRDLFTESWDTQKAPNNYRSPISTSQPTNQSMDDTREDIYVNYPTFSRR

35 PKTRV

The intracellular signalling domain may comprise, consist essentially of or consist of a co-stimulatory signalling domain as described herein.

5 The intracellular signalling domain may comprise a sequence shown as SEQ ID NO: 3 or 9-19 or a variant thereof.

10 The variant may comprise a sequence which shares at least 75% sequence identity with SEQ ID NO: 3 or 9-19 provided that the sequence provides an effective co-stimulatory signaling domain.

The variant may comprise a sequence which shares at least 80% sequence identity with SEQ ID NO: 3 or 9-19 provided that the sequence provides an effective co-stimulatory signaling domain.

15 The variant may comprise a sequence which shares at least 85% sequence identity with SEQ ID NO: 3 or 9-19 provided that the sequence provides an effective co-stimulatory signaling domain.

20 The variant may comprise a sequence which shares at least 90% sequence identity with SEQ ID NO: 3 or 9-19 provided that the sequence provides an effective co-stimulatory signaling domain.

25 The variant may comprise a sequence which shares at least 95% sequence identity with SEQ ID NO: 3 or 9-19 provided that the sequence provides an effective co-stimulatory signaling domain.

30 The variant may comprise a sequence which shares at least 99% sequence identity with SEQ ID NO: 3 or 9-19 provided that the sequence provides an effective co-stimulatory signaling domain.

In one embodiment, the intracellular signalling domain may comprise a sequence shown as SEQ ID NO: 3 or a variant thereof which shares at least 75, 80, 85, 90, 95 or 99% sequence identity with SEQ ID NO: 3, provided that the sequence provides an effective co-stimulatory signaling domain.

35 In one embodiment, the endodomain does not comprise the CD3 endodomain. For example, the endodomain does not comprise the CD3 epsilon chain, the CD3 gamma

chain and/or the CD3 delta chain. In a particular embodiment, the endodomain does not comprise the CD3-zeta endodomain.

An illustrative CD3-zeta endodomain is shown as SEQ ID NO: 26.

5

SEQ ID NO: 26 (CD3 zeta endodomain)

RSRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPRRK
NPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDALHMQ
ALPPR

10

The CD3-zeta endodomain as described herein may comprise or consist of SEQ ID NO: 26 or a variant thereof which has at least 80%, 85%, 90%, 95%, 98% or 99% sequence identity to SEQ ID NO: 26 and provides an effective transmembrane domain/intracellular T cell signaling domain.

15

ANTIGEN BINDING DOMAIN

The antigen binding domain is the portion of the CAR which recognizes antigen. Numerous antigen-binding domains are known in the art, including those based on the antigen binding site of an antibody, antibody mimetics, and T-cell receptors. For example, the antigen-binding domain may comprise: a single-chain variable fragment (scFv) derived from a monoclonal antibody; a natural ligand of the target antigen; a peptide with sufficient affinity for the target; a single domain antibody; an artificial single binder such as a Darpin (designed ankyrin repeat protein); or a single-chain derived from a T-cell receptor.

25

The antigen binding domain may comprise a domain which is not based on the antigen binding site of an antibody. For example the antigen binding domain may comprise a domain based on a protein/peptide which is a soluble ligand for a tumour cell surface receptor (e.g. a soluble peptide such as a cytokine or a chemokine); or an extracellular domain of a membrane anchored ligand or a receptor for which the binding pair counterpart is expressed on the tumour cell.

30

By way of example, the examples described herein relate to CARs which bind GD2 and CD33, respectively.

35

The antigen binding domain may be based on a natural ligand of the antigen.

The antigen binding domain may comprise an affinity peptide from a combinatorial library or a *de novo* designed affinity protein/peptide.

5 TUMOUR-ASSOCIATED ANTIGEN (TAA)

The antigen binding domain may bind to a tumour-associated antigen (TAA).

10 An extensive range of TAAs are known in the art and the CAR used in the present invention may comprise any antigen binding domain which is capable of specifically binding to any TAA.

By way of example, the CAR for use in the present invention may be capable of specifically binding to a TAA listed in Table 1.

15

Table 1

Antigen	Tumour of interest
CD20	B-cell lymphomas, CLL
CD19	Pre-B ALL, B-cell lymphoma, CLL
CD22	Pre-B ALL, B-cell lymphomas, CLL
CD30	Hodgkin's lymphoma, ALCL
CD52	T-cell AML, Pre-B ALL
CD70	Hodgkins Lymphoma, DLCL, Renal cell carcinoma, EBV+ glioblastoma, undifferentiated nasopharyngeal sarcoma
CD33	AML, MDS, APL, CML, JMML, ALL (18% only)
CD47	Pre-B ALL, T cell ALL, AML
IL7 receptor α	Pre-B ALL, B cell lymphomas
TSLPR	Pre-B ALL (7%), Pre-B aLL in Down's syndrome (60%)
ROR1	Pre-B ALL, CLL mantle cell lymphoma
GD2	Neuroblastoma, osteosarcoma, Ewing sarcoma, soft tissue sarcomas, melanoma
IL13R α 2	Glioblastoma, DIPG, melanoma, various carcinomas, mesothelioma
VEGFR2	Tumour vasculature
HER2	Osteosarcoma, colon cancer, breast cancer
ALK	Neuroblastoma, neuroectodermal tumours, glioblastoma, rhabdomyosarcoma, melanoma
EGFRvIII	Glioma
FGFR4	Rhabdomyosarcoma
B7-H3	Neuroblastoma
Glypican-3/Glypican-5	Wilm's tumour, neuroblastoma, rhabdomyosarcoma, hepatic carcinoma, melanoma
FOLR1	Rhabdomyosarcoma, osteosarcoma

A problem associated with the targeting of TAAs in cancer immunotherapy is that low levels of the TAAs may be expressed on normal tissues. For instance GD2 is a neuroblastoma TAA, but it is also expressed on nerves; PSMA is a prostate cancer TAA but also is found on normal kidney, liver and colon cells, and brain astrocytes.

5 This problem is more profound in solid tumours where there is a dearth of highly selective targets.

The expression of TAAs on normal, healthy cells may result in 'on-target, off-tumour' side effects. The present invention mitigates these effects because the $\gamma\delta$ T cell of
10 the present invention is only activated by cells which express a ligand for both the $\gamma\delta$ TCR and the CAR. Normal, healthy cells which express the TAA at low levels will therefore not activate the $\gamma\delta$ T cell of the present invention because they do not express a danger signal antigen capable of binding to the $\gamma\delta$ TCR (Figure 2D).

15 The antigen binding domain of the CAR may be capable of binding GD2, CD33, CD19 or EGFR.

Disialoganglioside (GD2, for example as shown by pubchem: 6450346) is a sialic acid-containing glycosphingolipid expressed primarily on the cell surface. The
20 function of this carbohydrate antigen is not completely understood; however, it is thought to play an important role in the attachment of tumour cells to extracellular matrix proteins. GD2 is densely, homogenously and almost universally expressed on neuroblastoma. In normal tissues, GD2 expression is largely limited to skin melanocytes, and peripheral pain fibre myelin sheaths. Within the CNS, GD2
25 appears to be an embryonic antigen but is found dimly expressed in scattered oligodendrocytes and within the posterior pituitary.

The antigen binding domain may comprise a sequence shown as SEQ ID NO: 20 or a variant thereof, providing that the variant retains the ability to bind to GD2.

30

SEQ ID NO: 20

METDTLLLWLLLWPGSTGQVQLQESGPGLVKPSQTLSTCTVSGFSLASYNHWWRP
KGLEWLGVWAGGSTNYNSALMSRLTISKDNSKNQVFLKMSSLTAADTAVYYCAKRSDDYS
WFAYWGQGTLLTVSSGGGGSGGGGSGGGGSENQMTQSPSSLSASVGDRVTMTCRASSS
35 VSSSYLHWYQQKSGKAPKVWIYSTSNLASGVPSRFSGSGSGTDYTLTISSLQPEDFATYYCQ
QYSGYPITFGQGTVKVEIKRS

The antigen binding domain may comprise a sequence shown as SEQ ID NO: 20 or a variant thereof which shares at least 75, 80, 85, 90, 95 or 99% sequence identity with SEQ ID NO: 20, providing that the variant retains the ability to bind to GD2.

- 5 CD33 (for example as shown by Uniprot accession number P20138) is a putative adhesion molecule of myelomonocytic-derived cells that mediates sialic-acid dependent binding to cells. It is usually considered myeloid-specific, but it can also be found on some lymphoid cells.
- 10 The antigen binding domain may comprise a sequence shown as SEQ ID NO: 21 or a variant thereof, providing that the variant retains the ability to bind to GD2.

SEQ ID NO: 21

MAVPTQVLGLLLLWLTARCDIQMTQSPSSLSASVGDRVITCRASEDIYFNLVWYQQKPGK
 15 APKLLIYDTNRLADGVPSRFSGSGSGTQYTLTISSLQPEDFATYYCQHYKNYPLTFGQGKLE
 IKRSGGGGSGGGGSGGGGSGGGGSRSEVQLVESGGGLVQPGGSLRLSCAASGFTLSNYG
 MHWIRQAPGKLEWVSSISLNGGSTYYRDSVKGRFTISRDNASTLYLQMNSLRRAEDTAVYY
 CAAQDAYTGGYFDYWGGGTLTVSSM

- 20 The antigen binding domain may comprise a sequence shown as SEQ ID NO: 21 or a variant thereof which shares at least 75, 80, 85, 90, 95 or 99% sequence identity with SEQ ID NO: 21, providing that the variant retains the ability to bind to GD2.

The human CD19 antigen is a 95 kd transmembrane glycoprotein belonging to the immunoglobulin superfamily (for example as shown by Uniprot P15391). CD19 is
 25 expressed very early in B-cell differentiation and is only lost at terminal B-cell differentiation into plasma cells. Consequently, CD19 is expressed on all B-cell malignancies apart from multiple myeloma. CD19 is also expressed by the normal B cell compartment.

30

EGFR (for example as shown by Uniprot accession number P00533) is a receptor tyrosine kinase which binds ligands of the EGF family and activates several signaling cascades to convert extracellular cues into appropriate cellular responses. Known
 ligands include EGF, TGFA/TGF- α , amphiregulin, epigen/EPGN,
 35 BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. EGFR is expressed at high levels by many cancer cells. However, it is also expressed by normal, healthy cells.

SPACER DOMAIN

CARs may comprise a spacer sequence to connect the antigen-binding domain with the transmembrane domain and spatially separate the antigen-binding domain from the endodomain. A flexible spacer allows the antigen-binding domain to orient in different directions to facilitate binding.

The spacer sequence may, for example, comprise an IgG1 Fc region, an IgG1 hinge or a human CD8 stalk or the mouse CD8 stalk. The spacer may alternatively comprise an alternative linker sequence which has similar length and/or domain spacing properties as an IgG1 Fc region, an IgG1 hinge or a CD8 stalk. A human IgG1 spacer may be altered to remove Fc binding motifs.

Examples of amino acid sequences for these spacers are given below:

SEQ ID NO: 22 (hinge-CH₂CH₃ of human IgG1)
 AEPKSPDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMIARTPEVTCVVDVSHEDPE
 VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA
 LPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESN
 GQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQK
 SLSLSPGKKD

SEQ ID NO: 23 (human CD8 stalk)
 TTPAPRPPTPAPTASQPLSLRPEACRPAAGGAVHTRGLDFACDI

SEQ ID NO: 24 (human IgG1 hinge)
 AEPKSPDKTHTCPPCPKDPK

The spacer may be a variant of any of SEQ ID NO: 22 to 24 which shares at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99% sequence identity with SEQ ID NO: 22 to 24 and retains the functional activity of the amino acid sequence shown as SEQ ID NO: 9 to 11.

TRANSMEMBRANE DOMAIN

30

The transmembrane domain is the sequence of the CAR that spans the membrane.

A transmembrane domain may be any protein structure which is thermodynamically stable in a membrane. This is typically an alpha helix comprising of several

hydrophobic residues. The transmembrane domain of any transmembrane protein can be used to supply the transmembrane portion of the invention. The presence and span of a transmembrane domain of a protein can be determined by those skilled in the art using the TMHMM algorithm (<http://www.cbs.dtu.dk/services/TMHMM-2.0/>).

5 Further, given that the transmembrane domain of a protein is a relatively simple structure, i.e a polypeptide sequence predicted to form a hydrophobic alpha helix of sufficient length to span the membrane, an artificially designed TM domain may also be used (US 7052906 B1 describes synthetic transmembrane components).

10 The transmembrane domain may be derived from any type I transmembrane protein. The transmembrane domain may be a synthetic sequence predicted to form a hydrophobic helix.

The transmembrane domain may be derived from CD28, which gives good receptor stability.

15

The transmembrane domain may comprise the sequence shown as SEQ ID NO: 25.

SEQ ID NO: 25 (CD28 transmembrane domain)

FWLVVVGGVLACYSLLVTVAFIIFWV

NUCLEIC ACID

20

The present invention further provides a nucleic acid sequence which encodes a CAR as described herein.

25 The nucleic acid sequence may be capable of encoding a CAR having the amino acid sequence shown as SEQ ID NO: 1 or SEQ ID NO: 2.

SEQ ID NO: 4 (aCD33-Fc-DAP10 CAR)

30 ATGGCCGTGCCCACTCAGGTCCTGGGGTTGTTGCTACTGTGGCTTACAGATGCC
AGATGTGACATCCAGATGACACAGTCTCCATCTTCCCTGTCTGCATCTGTCGGA
GATCGCGTCACCATCACCTGTCGAGCAAGTGAGGACATTTATTTTAATTTAGTGT
GGTATCAGCAGAAACCAGGAAAGGCCCTAAGCTCCTGATCTATGATACAAATC
GCTTGGCAGATGGGGTCCCATCACGGTTCAGTGGCTCTGGATCTGGCACACAG
TATACTCTAACCATAAGTAGCCTGCAACCCGAAGATTTGCAACCTATTATTGTC
35 AACACTATAAGAATTATCCGCTCACGTTCCGTCAGGGGACCAAGCTGGAAATCA
AAAGATCTGGTGGCGGAGGGTCAGGAGGCGGAGGCAGCGGAGGCGGTGGCTC

GGGAGGCGGAGGCTCGAGATCTGAGGTGCAGTTGGTGGAGTCTGGGGGCGGC
TTGGTGCAGCCTGGAGGGTCCCTGAGGCTCTCCTGTGCAGCCTCAGGATTAC
TCTCAGTAATTATGGCATGCACTGGATCAGGCAGGCTCCAGGGAAGGGTCTGGA
GTGGGTCTCGTCTATTAGTCTTAATGGTGGTAGCACTTACTATCGAGACTCCGTG
5 AAGGGCCCGATTCACTATCTCCAGGGACAATGCAAAAAGCACCCCTCTACCTTCAA
ATGAATAGTCTGAGGGCCGAGGACACGGCCGTCTATTACTGTGCAGCACAGGA
CGCTTATACGGGAGGTTACTTTGATTACTGGGGCCAAGGAACGCTGGTCACAGT
CTCGTCTATGGATCCCGCCGAGCCCAAATCTCCTGACAAAACCTCACACATGCCC
ACCGTGCCCAGCACCTCCCGTGGCCGGCCCGTCAGTCTTCCTCTTCCCCCAA
10 AACCCAAGGACACCCTCATGATCGCCCGGACCCCTGAGGTCACATGCGTGGTG
GTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGG
CGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCA
CGTACCGTGTGGTCAGCGTCCTACCGTCCTGCACCAGGACTGGCTGAATGGC
AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAA
15 ACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCC
CCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCA
AAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAACCG
GAGAACAACCTACAAGACCACGCCTCCCGTGTGGACTCCGACGGCTCCTTCTTC
CTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTT
20 CTCATGCTCCGTGATGCATGAGGCCCTGCACAATCACTATACCCAGAAATCTCT
GAGTCTGAGCCCAGGCAAGAAGGACCCCAAGTTCTGGGTCTGGTGGTGGTGG
GAGGCGTGCTGGCCTGTTACTCTCTCCTGGTGACCGTGGCCTTCATCATCTTCT
GGGTGTGCGCCAGACCACGGCGGAGCCCAGCCCAGGAGGACGGCAAGGTGTA
CATCAACATGCCCCGGCCGCGGCTGA

25

SEQ ID NO: 5 (aGD2-Fc-DAP10 CAR)

ATGGAGACCGACACCCTGCTGCTGTGGGTGCTGCTGCTGTGGGTGCCAGGCAG
CACCGGCCAGGTGCAGCTGCAGGAGTCTGGCCCAGGCCTGGTGAAGCCCAGC
30 CAGACCCTGAGCATCACCTGCACCGTGAGCGGCTTCAGCCTGGCCAGCTACAA
CATCCACTGGGTGCGGCAGCCCCCAGGCAAGGGCCTGGAGTGGCTGGGCGTG
ATCTGGGCTGGCGGCAGCACCAACTACAACAGCGCCCTGATGAGCCGGCTGAC
CATCAGCAAGGACAACAGCAAGAACCAGGTGTTCTGAAGATGAGCAGCCTGAC
AGCCGCCGACACCGCCGTGTACTACTGCGCCAAGCGGAGCGACGACTACAGCT
35 GGTTGCGCTACTGGGGCCAGGGCACCCCTGGTGACCGTGAGCTCTGGCGGAGG
CGGCTCTGGCGGAGGCGGCTCTGGCGGAGGCGGCAGCGAGAACCAGATGACC
CAGAGCCCCAGCAGCTTGAGCGCCAGCGTGGGCGACCGGGTGACCATGACCT

GCAGAGCCAGCAGCAGCGTGAGCAGCAGCTACCTGCACTGGTACCAGCAGAAG
AGCGGCAAGGCCCCAAAGGTGTGGATCTACAGCACCAGCAACCTGGCCAGCGG
CGTGCCCAGCCGGTTCAGCGGCAGCGGCAGCGGCACCGACTACACCCTGACC
ATCAGCAGCCTGCAGCCCCGAGGACTTCGCCACCTACTACTGCCAGCAGTACAG
5 CGGCTACCCCATCACCTTCGGCCAGGGCACCAAGGTGGAGATCAAGCGGTTCGG
ATCCCGCCGAGCCCAAATCTCCTGACAAAACCTCACACATGCCACCGTGCCAG
CACCTCCCGTGGCCGGCCCGTCAGTCTTCTCTTCCCCC AAAACCCAAGGACA
CCCTCATGATCGCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGC
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCA
10 TAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGG
TCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAG
TGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGAGAAAACCATCTCCAAA
GCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGA
TGAGCTGACCAAGAACCAGGTACGCCTGACCTGCCTGGTCAAAGGCTTCTATCC
15 CAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAACCGGAGAACAACTACA
AGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGC
TCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTG
ATGCATGAGGCCCTGCACAATCACTATACCCAGAAATCTCTGAGTCTGAGCCCA
GGCAAGAAGGACCCCAAGTTCTGGGTCTGGTGGTGGTGGGAGGCGTGCTGG
20 CCTGTTACTCTCTCTGGTGACCGTGGCCTTCATCATCTTCTGGGTGTGCGCCA
GACCACGGCGGAGCCCAGCCCAGGAGGACGGCAAGGTGTACATCAACATGCC
CGGCCGCGGCTGA

The nucleic acid sequence may encode the same amino acid sequence as that
25 encoded by SEQ ID NO: 1 or 2, but may have a different nucleic acid sequence, due
to the degeneracy of the genetic code. The nucleic acid sequence may have at least
80, 85, 90, 95, 98 or 99% identity to the sequence shown as SEQ ID NO: 4 or SEQ ID
NO: 5, provided that it encodes a CAR as defined in the first aspect of the invention.

30 VARIANT

Sequence comparisons can be conducted by eye, or more usually, with the aid of
readily available sequence comparison programs. These publicly and commercially
available computer programs can calculate sequence identity between two or more
sequences.

35 Sequence identity may be calculated over contiguous sequences, i.e. one sequence
is aligned with the other sequence and each amino acid in one sequence directly

compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an “ungapped” alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues (for example less than 50 contiguous amino acids).

- 5 Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to
- 10 produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting “gaps” in the sequence alignment to try to maximise local homology.

However, these more complex methods assign “gap penalties” to each gap that occurs in the alignment so that, for the same number of identical amino acids, a

15 sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. “Affine gap costs” are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of

20 course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use the default values when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

- 25 Calculation of maximum % sequence identity therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A; Devereux *et al.*, 1984, Nucleic Acids Research 12:387). Examples of other software than can perform sequence comparisons
- 30 include, but are not limited to, the BLAST package (see Ausubel *et al.*, 1999 *ibid* – Chapter 18), FASTA (Atschul *et al.*, 1990, J. Mol. Biol., 403-410) and the GENEWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel *et al.*, 1999 *ibid*, pages 7-58 to 7-60). However it is preferred to use the GCG Bestfit program.

Although the final sequence identity can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). It is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

Once the software has produced an optimal alignment, it is possible to calculate % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

The terms "variant" according to the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acids from or to the sequence providing the resultant amino acid sequence retains substantially the same activity as the unmodified sequence.

Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

It will be understood by a skilled person that numerous different polynucleotides and nucleic acids can encode the same polypeptide as a result of the degeneracy of the genetic code. In addition, it is to be understood that skilled persons may, using routine techniques, make nucleotide substitutions that do not affect the polypeptide sequence
5 encoded by the polynucleotides described here to reflect the codon usage of any particular host organism in which the polypeptides are to be expressed.

A nucleic acid sequence or amino acid sequence as described herein may comprise, consist of or consist essentially of a nucleic acid sequence or amino acid sequence as shown herein.

10 VECTOR

The present invention also provides a vector which comprises a nucleic acid sequence according to the present invention. Such a vector may be used to introduce the nucleic acid sequence into a host cell so that it expresses and produces
15 a molecule according to the first aspect of the invention.

The vector may, for example, be a plasmid or a viral vector, such as a retroviral vector or a lentiviral vector.

20 The vector may be capable of transfecting or transducing a T cell.

The vector may also comprise a nucleic acid sequence encoding a suicide gene, such as iCasp9 or RQR8.

25 A suicide-gene is a genetically encoded mechanism which allows selective destruction of adoptively transferred cells, such as T-cells, in the face of unacceptable toxicity.

Activation of Caspase 9 results in cell apoptosis. The activation mechanism behind
30 Caspase 9 was exploited by the iCasp9 molecule. All that is needed for Caspase 9 to become activated, is overcoming the energetic barrier for Caspase 9 to homodimerize. The homodimer undergoes a conformational change and the proteolytic domain of one of a pair of dimers becomes active. Physiologically, this occurs by binding of the CARD domain of Caspase 9 to APAF-1. In iCasp9, the APAF-1 domain is replaced
35 with a modified FKBP12 which has been mutated to selectively bind a chemical inducer of dimerization (CID). Presence of the CID results in homodimerization and

activation. iCasp9 is based on a modified human caspase 9 fused to a human FK506 binding protein (FKBP) (Straathof et al (2005) Blood 105:4247-4254). It enables conditional dimerization in the presence of a small molecule CID, known as AP1903.

- 5 Expression of RQR8 renders T-cells susceptible to anti-CD20 antibody Rituximab but is more compact than the full-length CD20 molecule (Philip, B. et al. (2014) Blood doi:10.1182/blood-2014-01-545020).

PHARMACEUTICAL COMPOSITION

10

The present invention also relates to a pharmaceutical composition containing a vector or a CAR-expressing T cell of the invention together with a pharmaceutically acceptable carrier, diluent or excipient, and optionally one or more further pharmaceutically active polypeptides and/or compounds. Such a formulation may, for
15 example, be in a form suitable for intravenous infusion.

METHOD

- 20 The present invention also relates to a method for making a cell according to the present invention, which comprises the step of introducing a nucleic acid sequence or vector according to the present invention into a cell.

- CAR-expressing cells according to the present invention may either be created *ex vivo* either from a patient's own peripheral blood (1st party), or in the setting of a
25 haematopoietic stem cell transplant from donor peripheral blood (2nd party), or peripheral blood from an unconnected donor (3rd party). Alternatively, CAR T-cells may be derived from *ex-vivo* differentiation of inducible progenitor cells or embryonic progenitor cells to T-cells. In these instances, CAR T-cells are generated by introducing DNA or RNA coding for the CAR by one of many means including
30 transduction with a viral vector, transfection with DNA or RNA.

- The method may further comprise stimulating the cell with a $\gamma\delta$ T cell stimulating agent. As used herein, a ' $\gamma\delta$ T cell stimulating agent' refers to any agent which selectively stimulates the proliferation and/or survival of $\gamma\delta$ T cells from a mixed
35 starting population of cells.

Thus, the resulting cell population is enriched with an increased number of $\gamma\delta$ T cells - for example particular $\gamma\delta$ T cells expressing a particular $\gamma\delta$ TCR receptor - compared with the starting population of cells.

5 $\gamma\delta$ T cell populations produced in accordance with the present invention may be enriched with $\gamma\delta$ T cells, for example particular $\gamma\delta$ T cells expressing a particular $\gamma\delta$ TCR receptor. That is, the $\gamma\delta$ T cell population that is produced in accordance with the present invention will have an increased number of $\gamma\delta$ T cells. For example, the $\gamma\delta$ T cell population of the invention will have an increased number of $\gamma\delta$ T cells
10 expressing a particular $\gamma\delta$ TCR receptor compared with the $\gamma\delta$ T cells in a sample isolated from a subject. That is to say, the composition of the $\gamma\delta$ T cell population will differ from that of a "native" T cell population (i.e. a population that has not undergone expansion steps discussed herein), in that the percentage or proportion of $\gamma\delta$ T cells will be increased.

15 The $\gamma\delta$ T cell population according to the invention may have at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% $\gamma\delta$ T cells.

20 The $\gamma\delta$ T cell population according to the invention may have at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% $\gamma\delta$ T cells expressing a particular $\gamma\delta$ TCR receptor.

By way of example, the $\gamma\delta$ T cell stimulating agent may be isopentenyl pyrophosphate (IPP); an analog of IPP (e.g. bromohydrin pyrophosphate or (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate); an inhibitor of farnesyl pyrophosphate synthase (FPPS) or
25 an aminobisphosphonate such as zoledronate or pamidronate.

The $\gamma\delta$ T cell stimulating agent may be used in combination with a general T cell mitogen, for example a mitogenic cytokine such as IL-2.
30

Additional methods of stimulating $\gamma\delta$ T cells are known in art and include, for example, the use of Concanavalin A (Siegers, G. M. *et al. PLoS ONE* **6**, e16700 (2011)), anti- $\gamma\delta$ TCR antibodies immobilized on plastic; engineered artificial antigen
35 presenting cells as feeders and engineered artificial antigen presenting cells coated in anti- $\gamma\delta$ TCR antibody (Fisher, J. *et al.; Clin. Cancer Res.* (2014)).

METHOD OF TREATMENT

A method for the treatment of disease relates to the therapeutic use of a vector or T cell of the invention. In this respect, the vector or T cell may be administered to a
5 subject having an existing disease or condition in order to lessen, reduce or improve at least one symptom associated with the disease and/or to slow down, reduce or block the progression of the disease.

CAR- expressing T cells may either be created *ex vivo* either from a patient's own
10 peripheral blood (1st party), or in the setting of a haematopoietic stem cell transplant from donor peripheral blood (2nd party), or peripheral blood from an unconnected donor (3rd party). Alternatively, CAR T-cells may be derived from *ex-vivo* differentiation of inducible progenitor cells or embryonic progenitor cells to T-cells. In these instances, CAR T-cells are generated by introducing DNA or RNA coding for
15 the CAR by one of many means including transduction with a viral vector, transfection with DNA or RNA.

In one embodiment, the sample comprising $\gamma\delta$ T cell may have been previously isolated from the subject.

20

A CAR T cell according to the present invention may be generated by a method as described herein. In particular, a CAR- expressing T cell for use in a method for the treatment of a disease may be generated by a method comprising the steps of transduction of the T cell with a viral vector or transfection with DNA or RNA encoded
25 the co-stimulatory CAR as described herein and expansion of $\gamma\delta$ T cells using a $\gamma\delta$ T cell stimulating agent.

The $\gamma\delta$ T cell stimulating agent may be isopentenyl pyrophosphate (IPP); an analog of IPP (e.g. bromohydrin pyrophosphate or (E)-4-Hydroxy-3-methyl-but-2-enyl
30 pyrophosphate); an inhibitor of farnesyl pyrophosphate synthase (FPPS) or aminobisphosphonates such as zoledronate or pamidronate, for example.

T cells expressing a CAR molecule of the present invention may be used for the treatment of a various diseases including, for example, cancer, microbial infection and
35 viral infection.

The cancer may be, for example, bladder cancer, breast cancer, colon cancer, endometrial cancer, kidney cancer (renal cell), lung cancer, brain cancer, melanoma, leukaemia, lymphoma, pancreatic cancer, prostate cancer or thyroid cancer.

5 The methods and uses according to the present invention may be practiced in combination with additional compositions. For example, where the disease to be treated is cancer, the composition of the present invention may be administered in combination with additional cancer therapies such as chemotherapy and/or radiotherapy.

10

A composition of the present invention may be administered in combination with a $\gamma\delta$ T cell stimulating agent such as isopentenyl pyrophosphate (IPP); an analog of IPP (e.g. bromohydrin pyrophosphate or (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate); an inhibitor of farnesyl pyrophosphate synthase (FPPS) or
15 aminobisphosphonates such as zoledronate or pamidronate.

In particular, Zoledronate and Pamidronate can be used for *in vivo* expansion of V δ 2+ $\gamma\delta$ T cells in combination with IL-2. There are a number of Phase I clinical trials that have used this approach (see Fisher *et al.*; OncoImmunology; 3; e27572).

20

'In combination' may refer to administration of the additional therapy or $\gamma\delta$ T cell stimulating agent before, at the same time as or after administration of the composition according to the present invention.

25 The invention will now be further described by way of Examples, which are meant to serve to assist one of ordinary skill in the art in carrying out the invention and are not intended in any way to limit the scope of the invention.

EXAMPLES

30

Example 1 - Generation of $\gamma\delta$ T cells expressing a co-stimulatory CAR

PBMCs were extracted from the blood of healthy donors using Ficoll density gradient separation. They were cultured in RPMI 1640 medium supplemented with 10% FCS, 1% penicillin/streptomycin, 100u/ml human IL-2 and 5 μ M zoledronic acid for 5 days.

35

After 5 days they were transduced with retrovirus containing the CAR construct fused to RQR8, which acts as a marker gene and also provides a Rituximab (α CD20) sensitive suicide gene.

The illustrative CAR described herein includes aGD2-specific scFv, a linker based on the Fc portion of IgG1, a transmembrane domain derived from CD28 and the endodomain of DAP10 (see Figure 10).

5

A second illustrative CAR includes a CD33-specific scFv, a linker based on the Fc portion of IgG1, a transmembrane domain derived from CD28 and the endodomain of DAP10 (see Figure 11).

10 Co-expression of an anti-GD2-Fc-DAP10 CAR with the endogenous TCR of a $\gamma\delta$ T cell was demonstrated (Figure 4).

Example 2 - Killing of GD2+ cell lines LAN1 and TC71 by V δ 2 $\gamma\delta$ T cells transduced with the aGD2-Fc-DAP10 CAR

15

Both the LAN1 and TC71 cells lines are known to express GD2.

Significant killing of GD2+ neuroblastoma cell line LAN1 was only seen when CAR transduced cells were used and not when non-transduced (NT) V δ 2 cells were used as effectors (Figure 5A).

20

There was an additive effect against the GD2+ Ewing sarcoma cell line TC71 when the aGD2-Fc-DAP10 CAR was used in combination with 24h zoledronic acid treatment (Figure 5B).

25

Addition of the CAR to $\alpha\beta$ T cells, which lack the signal 1 provided by the $\gamma\delta$ TCR in response to cellular stress, had no effect on cytotoxicity, unlike the effect of the CAR in V δ 2+ $\gamma\delta$ T cells (Figure 5C). This indicates that the CAR signal alone is insufficient for T-cell activation.

30

Expression of the aGD2-Fc-DAP10 CAR in $\gamma\delta$ T cells did not result in GD2-specific killing of GD2 negative SK-N-SH cells (Figure 6).

Example 3 - Preservation of CAR expression following prolonged co-culture and GD2 specific expansion

35

Co-culture was started 24 days after transduction and serial analyses of cells for the presence of CAR and TCRV δ 2 were taken in the presence of irradiated GD2+ (LAN1) and GD2- (SK-N-SH) neuroblastoma cells (Figure 7A).

- 5 The expansion of aGD2-Fc-DAP10 transduced V δ 2+ cells was only seen in the presence of irradiated GD2+ target cells (Figure 7B).

Example 4 – Specific killing of CD33+ AML cells but not CD33+ monocytes by $\gamma\delta$ T cells expressing an anti-CD33-DAP10 CAR

10

Equivalent levels of CD33 expression were demonstrated in three AML cell lines and monocytes (Figure 8).

- 15 V δ 2 $\gamma\delta$ T cells were transduced with either an anti-CD33-Fc-DAP10 or anti-CD33-Fc-CD28-CD3z CAR construct.

The anti-CD33-Fc-CD28-CD3z CAR construct provides signal 1 and signal 2 in the presence of CD33. The anti-CD33-Fc-DAP10 provides signal 2 in the presence of CD33.

20

Cells transduced with the aCD33-CD28-CD3z CAR killed any CD33 positive cell and did not spare healthy monocytes. Cells transduced with the aCD33-Fc-DAP10 CAR do not kill monocytes (Figure 9A).

- 25 There was significant enhancement of killing of the AML but no enhancement of the killing of monocytes by V δ 2 $\gamma\delta$ T cells transduced with the aCD33-Fc-DAP10 CAR compared to non-transduced controls (Figure 9B).

30 All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments.

35 Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology, cellular immunology or related fields are intended to be within the scope of the following claims.