

FUSED HETEROCYCLIC COMPOUNDS AS PROTEIN KINASE INHIBITORS

INTRODUCTION

[001] Bruton's tyrosine kinase (Btk) belongs to the Tec tyrosine kinase family (Vetrie et al., *Nature* **361**: 226-233, 1993; Bradshaw, *Cell Signal.* **22**: 1175-84, 2010). Btk is primarily expressed in most hematopoietic cells such as B cells, mast cells and macrophages (Smith et al., *J. Immunol.* **152**: 557-565, 1994) and is localized in bone marrow, spleen and lymph node tissue. Btk plays important roles in B-cell receptor (BCR) and FcR signaling pathways, which involve in B-cell development, differentiation (Khan, *Immunol. Res.* **23**: 147, 2001). Btk is activated by upstream Src-family kinases. Once activated, Btk in turn phosphorylates PLC gamma, leading to effects on B-cell function and survival (Humphries et al., *J. Biol.Chem.* **279**: 37651, 2004). These signaling pathways must be precisely regulated. Mutations in the gene encoding Btk cause an inherited B-cell specific immunodeficiency disease in humans, known as X-linked agammaglobulinemia (XLA) (Conley et al., *Annu. Rev. Immunol.* **27**: 199-227, 2009). Aberrant BCR-mediated signaling may result in dysregulated B-cell activation leading to a number of autoimmune and inflammatory diseases. Preclinical studies show that Btk deficient mice are resistant to developing collagen-induced arthritis. Moreover, clinical studies of Rituxan, a CD20 antibody to deplete mature B-cells, reveal the key role of B-cells in a number of inflammatory diseases such as rheumatoid arthritis, systemic lupus erythematosus and multiple sclerosis (Gurcan et al., *Int. Immunopharmacol.* **9**: 10-25, 2009). Therefore, Btk inhibitors can be used to treat autoimmune and/or inflammatory diseases.

[002] In addition, aberrant activating of Btk plays important role in pathogenesis of B-cell lymphomas indicating that inhibition of Btk is useful in the treatment of hematological malignancies (Davis et al., *Nature* **463**: 88-92, 2010). Preliminary clinical trial results showed that the Bruton's tyrosine kinase (Btk) inhibitor PCI-32765 was effective in treatment of several types of B-cell lymphoma (for example, 54th American Society of Hematology (ASH) annual meeting abstract, Dec. 2012: 686 *The Bruton's Tyrosine Kinase (Btk) Inhibitor, Ibrutinib (PCI-32765), Has Preferential Activity in the ABC Subtype of Relapsed/Refractory De Novo Diffuse Large B-Cell Lymphoma (DLBCL): Interim Results of a Multicenter, Open-Label, Phase2 Study*). Because Btk plays a central role as a mediator in multiple signal transduction pathways, inhibitors of Btk are of great interest as anti-inflammatory and/or anti-cancer agents (Mohamed


et al., *Immunol. Rev.* **228**: 58-73, 2009; Pan, *Drug News perspect* **21**: 357-362, 2008; Rokosz et al., *Expert Opin. Ther. Targets* **12**: 883-903, 2008; Uckun et al., *Anti-cancer Agents Med. Chem.* **7**: 624-632, 2007; Lou et al., *J. Med. Chem.* **55**(10): 4539-4550, 2012).

[003] Small molecule inhibitors of Btk are being developed for anti-inflammatory and anticancer therapy. Ibrutinib (PCI-32765, See: US7514444B2 and related documents, for examples, US2012053189A1; WO 2011153514; WO 2011046964; US2010254905A1; WO2010009342; WO2008121742; WO2008054827; US20080139582; US20080076921; US7718662B1; WO2007087068; US20100035841) is a first-in class of Btk inhibitor, currently undergoing multiple clinical trials in relapsed or refractory mantle cell lymphoma (MCL) and chronic lymphocytic leukaemia (CLL). Another Btk inhibitor entered clinical trials is AVL-292 (See, for example, US 20100249092; US20100029610; US2010016296; US20120077832; WO 2011090760; WO 2010028236; WO 2009158571; WO2009051822; WO2010123870). Ono pharmaceuticals and Mannkind Corporation have been doing clinical trials with their small molecular Btk inhibitors, respectively (See, for example, ONO-4059, WO2011152351; WO2007136790A2).

[004] Other Btk inhibitors are also known. See, for example, US2012/0232054 (LOCUS PHARMACEUTICALS, INC.), WO2010126960 (LOCUS PHARMACEUTICALS, INC.), WO 2011/162515 (HANMI HOLDINGS CO. LTD), WO2012135801 (UNIVERSITY OF UTAH RESEARCH FOUNDATION), Kim et al., *Bioorg. Med. Chem. Lett.* **21**: 6258–6263, 2011(Pfizer), US8084620B2 (BMS), WO2002050071; WO2008116064; WO2010011837; WO 2011159857(BMS), US2012058996A1; US2012082702A1; US20100160303 (BMS), US2012129852A1 (BMS), WO 2011019780 (BMS), WO2011029043; WO2011029046 (Biogen Idec), US7393848 (CGI), US20060178367; US20060183746 (CGI), EP2068849 (CGI), WO 2005005429; WO 2005014599; WO 2005047290; WO 2006053121; WO2008033834; WO 2008033858; WO 2006099075; WO 2008033854; WO 2008033857; WO 2009039397 (CGI), WO 2009137596; WO 2010056875; WO 2010068788; WO 2010068806; WO 2010068810 (CGI, GENENTECH), WO 2011140488; WO 2012030990; WO 2012031004 (GILEAD & GENENTECH), US2012040961A1 (DANA-FARBER CANCER INSTITUTE), WO 2005011597; WO 2008045627; WO 2008144253 (IRM LLC), WO 2007140222; WO 2013008095 (NOVARTIS), WO 2012170976A2 (Merck), WO2012135944A1 (PHARMASCIENCE), US2010144705A1; US20120028981A1 (PRINCIPIA BIOPHARMA), WO 2010065898A2; WO 2012158795A1; WO 2012158764A1; WO 2012158810A1 (PRINCIPIA BIOPHARMA), US20090318448A1; US20100016301; US2009105209A1; US20100222325; US20100004231 (ROCHE), WO 2012156334A1; WO 2012020008; WO

2010122038; WO 2010006970; WO 2010006947; WO 2010000633; WO 2009077334; WO 2009098144 (ROCHE), WO 2006065946; WO 2007027594; WO 2007027729 (VERTEX).


[005] WO 2007/026720 A1 discloses that a ring-fused pyrazole compound of formula (A), wherein n represents 2 or 3; **A** represents the formula: -O- or the like; **B** represents a C₁₋₁₀ alkylene group or the like; **C** represents a single bond or the formula: -O-; **R**¹- represents a hydrogen atom, a pyrrolidinyl group or the like; **R**⁴, **R**⁵ and **R**⁶ independently represents a hydrogen atom, a halogen atom or the like; **D**₁ = **D**₂ represents the formula: -CH=CH- or the like; **E** represents the formula: -O- or -NH- or the like; **G** represents a C1-10 alkylene group or the like; and **R**⁷ represents a hydrogen atom, a phenyl group or the like, is useful as an Lck kinase inhibitor:



SUMMARY OF THE INVENTION

[006] The invention provides methods and compositions for inhibiting Btk and treating disease associated with undesirable Btk activity (Btk-related diseases).

[007] In one embodiment the invention provides Btk inhibitors or compounds of formula:



[008] stereoisomers thereof, and pharmaceutically acceptable salts thereof, wherein:

[009] A is a 5- or 6-membered aromatic ring comprising 0-3 heteroatoms of N, S or O;

[010] each W is independently $-(CH_2)-$ or $-C(O)-$;

[011] L is a bond, CH_2 , NR^{12} , O, or S;

[012] S/D is a single or double bond, and when a double bond, R^5 and R^7 are absent;

[013] m is 0, or an integer of 1-4;

[014] n is 0, or an integer of 1-4, wherein when n is more than 1, each R^2 may be different;

[015] p is 0, or an integer of 1-2, wherein when p is 0, m is non-zero, and when p is more than 1, each R^6 and each R^7 may be different;

[016] R^1 , R^4 , R^5 , R^6 , and R^7 are each independently H, halogen, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, alkynyl, $-CN$, $-NR^{13}R^{14}$, $-OR^{13}$, $-COR^{13}$, $-CO_2R^{13}$, $-CONR^{13}R^{14}$, $-C(=NR^{13})NR^{14}R^{15}$, $-NR^{13}COR^{14}$, $-NR^{13}CONR^{14}R^{15}$, $-NR^{13}CO_2R^{14}$, $-SO_2R^{13}$, $-NR^{13}SO_2NR^{14}R^{15}$, or $-NR^{13}SO_2R^{14}$, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R^{16} , wherein (R^4 and R^5), or (R^4 and R^6), or (R^6 and R^7), or (R^6 and R^6 when p is 2), together with the atoms to which they are attached, can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R^{16} ;

[017] R^2 is halogen, alkyl, $-S$ -alkyl, $-CN$, $-NR^{13}R^{14}$, $-OR^{13}$, $-COR^{13}$, $-CO_2R^{13}$, $-CONR^{13}R^{14}$, $-C(=NR^{13})NR^{14}R^{15}$, $-NR^{13}COR^{14}$, $-NR^{13}CONR^{14}R^{15}$, $-NR^{13}CO_2R^{14}$, $-SO_2R^{13}$, $-NR^{13}SO_2NR^{14}R^{15}$, or $-NR^{13}SO_2R^{14}$;

[018] R^{12} is H or lower alkyl;

[019] R^{13} , R^{14} and R^{15} are each independently H, heteroalkyl, alkyl, alkenyl, alkynyl, cycloalkyl, saturated or unsaturated heterocyclyl, aryl, or heteroaryl; wherein (R^{13} and R^{14}), and/or (R^{14} and R^{15}) together with the atom(s) to which they are attached, each can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R^{16} ;

[020] R^{16} is halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocyclyl, oxo, -CN, -OR', -NR'R'', -COR', -CO₂R', -CONR'R'', -C(=NR')NR''R''', -NR'COR'', -NR'CONR'R'', -NR'CO₂R'', -SO₂R', -SO₂aryl, -NR'SO₂NR''R''', or -NR'SO₂R'', wherein R', R'', and R''' are independently hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocyclyl, wherein (R' and R''), and/or (R'' and R''') together with the atoms to which they are attached, can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings.

[021] In exemplary particular embodiments:

[022] (a) S/D is a double bond and R^5 and R^7 are absent;

[023] (b) R^1 is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R^{16} ;


[024] (c) p is 1 and m is 0, 1 or 2, preferably 0 or 1;

[025] (d) A is phenyl;

[026] (e) each R^2 is independently halogen, lower alkyl, or lower alkoxy

[027] (f) R^4 and R^6 , together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R^{16} .

[028] (g) R^4 and R^6 , together with the atoms to which they are attached, form a ring of formula:



Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4;

[029] (h) S/D is a single bond.

[030] (i) p is 0 and R⁶ and R⁷ are absent.

[031] The invention includes all combinations of the recited particular embodiments, such as (a)-(i), supra, as if each combination had been laboriously separately recited.

[032] In exemplary combinations of particular embodiments:

[033] (i) S/D is a double bond and R⁵ and R⁷ are absent; R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶; and R¹⁶ is halogen, lower alkyl, or lower alkoxy;


[034] (ii) S/D is a double bond and R⁵ and R⁷ are absent; p is 1 and m is 0 or 1 (or 2);

[035] (iii) S/D is a double bond and R⁵ and R⁷ are absent; p is 1 and m is 0 or 1 (or 2); A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy (see, formula II);

[036] (iv) S/D is a double bond and R⁵ and R⁷ are absent; and R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶.

[037] (v) S/D is a double bond and R⁵ and R⁷ are absent; R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy.


[038] (vi) S/D is a double bond and R⁵ and R⁷ are absent; R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; p is 1 and m 0 or 1 (or 2); A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy; and the R⁴-R⁶ ring is of formula:



wherein Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4 (see, formula III);

[039] (vii) S/D is a double bond and R⁵ and R⁷ are absent; R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; p

is 1 and m is 0 or 1 (or 2); A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy; and the R⁴-R⁶ ring is of formula:



wherein Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4; and R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶;


[040] (viii) S/D is a single bond; p is 1 and m is 0, 1 or 2; A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy (see, formula IV);

[041] (ix) S/D is a single bond; p is 1 and m is 0, 1 or 2; A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy, and R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶; and R¹⁶ is halogen, lower alkyl, or lower alkoxy;

[042] (x) S/D is a single bond; p is 0 and R⁶ and R⁷ are absent; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy (see, formula V);

[043] (xi) S/D is a single bond; p is 0 and R⁶ and R⁷ are absent; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy, and R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶; and R¹⁶ is halogen, lower alkyl, or lower alkoxy.

[044] In particular embodiments the invention provides compounds of formula II, III, IV and V, stereoisomers thereof, and pharmaceutically acceptable salts thereof, wherein substituents are as defined herein:



[045] The invention also provides compounds of the examples herein, or of Table I, II or III, (below), stereoisomers thereof, and pharmaceutically acceptable salts thereof.

[046] The invention also provides subject compounds having a Btk-inhibiting activity corresponding to a IC50 of 10 uM or less in the BTK KINASE ASSAY.

[047] The invention also provides pharmaceutical compositions comprising a therapeutically effective amount of a subject compound in unit dosage form and one or more pharmaceutically acceptable carriers.

[048] The invention also provides combinations comprising a therapeutically effective amount of a subject compound and a different agent therapeutically active against an autoimmune and/or inflammatory disease.

[049] The invention also provides methods treating a Btk related disease, or disease associated with undesirable Btk activity, particularly an allergic disease, an autoimmune disease (e.g. rheumatoid arthritis), an inflammatory disease, or cancer (e.g. a B-cell proliferative disorder, such as chronic lymphocytic lymphoma, non-Hodgkin's lymphoma, diffuse large B cell lymphoma, mantle cell lymphoma, follicular lymphoma or chronic lymphocytic leukemia), which methods generally comprise administering to a mammal in need thereof an effective amount of a subject compound, an N-oxide thereof or a prodrug thereof, and optionally detecting a resultant amelioration of disease or symptom thereof, or Btk-inhibition.

[050] The invention also provides pharmaceutical compositions comprising a subject compound in unit dosage, administrable form, and methods of inducing autophagy, comprising administering to a person in need thereof an effective amount of a subject compound or composition.

[051] The invention also provides the subject compounds for use as a medicament, and use of the subject compounds in the manufacture of a medicament for the treatment of a Btk related disease.

DESCRIPTION OF PARTICULAR EMBODIMENTS OF THE INVENTION

[052] Disclosed herein are compounds that can inhibit tyrosine kinases, such as Btk, Blk, Bmx, EGFR, ERBB2, ERBB4, Itk, Jak3, Tec and Txk kinases.

[053] The following words, phrases and symbols are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise. The following abbreviations and terms have the indicated meanings throughout.

[054] The term "alkyl" refers to a hydrocarbon group selected from linear and branched saturated hydrocarbon groups of 1-18, or 1-12, or 1-6 carbon atoms. Examples of the alkyl group include methyl, ethyl, 1-propyl or n-propyl ("n-Pr"), 2-propyl or isopropyl ("i-Pr"), 1-butyl or n-butyl ("n-Bu"), 2-methyl-1-propyl or isobutyl ("i-Bu"), 1-methylpropyl or s-butyl ("s-Bu"), and 1,1-dimethylethyl or t-butyl ("t-Bu"). Other examples of the alkyl group include 1-pentyl, 2-pentyl, 3-pentyl, 2-methyl-2-butyl, 3-methyl-2-butyl, 3-methyl-1-butyl, 2-methyl-1-butyl, 1-hexyl, 2-hexyl, 3-hexyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 3-methyl-3-pentyl, 2-methyl-3-pentyl, 2,3-dimethyl-2-butyl and 3,3-dimethyl-2-butyl groups.

[055] Lower alkyl means 1-8, preferably 1-6, more preferably 1-4 carbon atoms; lower alkenyl or alkynyl means 2-8, 2-6 or 2-4 carbon atoms.

[056] The term "alkenyl" refers to a hydrocarbon group selected from linear and branched hydrocarbon groups comprising at least one C=C double bond and of 2-18, or 2-12, or 2-6 carbon atoms. Examples of the alkenyl group may be selected from ethenyl or vinyl, prop-1-enyl, prop-2-enyl, 2-methylprop-1-enyl, but-1-enyl, but-2-enyl, but-3-enyl, buta-1,3-dienyl, 2-methylbuta-1,3-diene, hex-1-enyl, hex-2-enyl, hex-3-enyl, hex-4-enyl, and hexa-1,3-dienyl groups.

[057] The term "alkynyl" refers to a hydrocarbon group selected from linear and branched hydrocarbon group, comprising at least one C≡C triple bond and of 2-18, or 2-12, or 2-6 carbon atoms. Examples of the alkynyl group include ethynyl, 1-propynyl, 2-propynyl (propargyl), 1-butylnyl, 2-butylnyl, and 3-butylnyl groups.

[058] The term "cycloalkyl" refers to a hydrocarbon group selected from saturated and partially unsaturated cyclic hydrocarbon groups, comprising monocyclic and polycyclic (e.g., bicyclic and tricyclic) groups. For example, the cycloalkyl group may be of 3-12, or 3-8, or 3-6 carbon atoms. Even further for example, the cycloalkyl group may be a monocyclic group of 3-12, or 3-8, or 3-6 carbon atoms. Examples of the monocyclic cycloalkyl group include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, cyclohexadienyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, cycloundecyl, and cyclododecyl groups. Examples of the bicyclic cycloalkyl groups include those having 7-12 ring atoms arranged as a

bicycle ring selected from [4,4], [4,5], [5,5], [5,6] and [6,6] ring systems, or as a bridged bicyclic ring selected from bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, and bicyclo[3.2.2]nonane. The ring may be saturated or have at least one double bond (i.e. partially unsaturated), but is not fully conjugated, and is not aromatic, as aromatic is defined herein.

[059] The term "Aryl" herein refers to a group selected from: 5- and 6-membered carbocyclic aromatic rings, for example, phenyl; bicyclic ring systems such as 7-12 membered bicyclic ring systems wherein at least one ring is carbocyclic and aromatic, selected, for example, from naphthalene, indane, and 1,2,3,4-tetrahydroquinoline; and tricyclic ring systems such as 10-15 membered tricyclic ring systems wherein at least one ring is carbocyclic and aromatic, for example, fluorene.

[060] For example, the aryl group is selected from 5- and 6-membered carbocyclic aromatic rings fused to a 5- to 7-membered cycloalkyl or heterocyclic ring optionally comprising at least one heteroatom selected from N, O, and S, provided that the point of attachment is at the carbocyclic aromatic ring when the carbocyclic aromatic ring is fused with a heterocyclic ring, and the point of attachment can be at the carbocyclic aromatic ring or at the cycloalkyl group when the carbocyclic aromatic ring is fused with a cycloalkyl group. Bivalent radicals formed from substituted benzene derivatives and having the free valences at ring atoms are named as substituted phenylene radicals. Bivalent radicals derived from univalent polycyclic hydrocarbon radicals whose names end in "-yl" by removal of one hydrogen atom from the carbon atom with the free valence are named by adding "-idene" to the name of the corresponding univalent radical, e.g., a naphthyl group with two points of attachment is termed naphthylidene. Aryl, however, does not encompass or overlap with heteroaryl, separately defined below. Hence, if one or more carbocyclic aromatic rings are fused with a heterocyclic aromatic ring, the resulting ring system is heteroaryl, not aryl, as defined herein.

[061] The term "halogen" or "halo" refers to F, Cl, Br or I.

[062] The term "heteroalkyl" refers to alkyl comprising at least one heteroatom.

[063] The term "heteroaryl" refers to a group selected from:

[064] 5- to 7-membered aromatic, monocyclic rings comprising 1, 2, 3 or 4 heteroatoms selected from N, O, and S, with the remaining ring atoms being carbon;

[065] 8- to 12-membered bicyclic rings comprising 1, 2, 3 or 4 heteroatoms, selected from N, O, and S, with the remaining ring atoms being carbon and wherein at least one ring is aromatic and at least one heteroatom is present in the aromatic ring; and

[066] 11- to 14-membered tricyclic rings comprising 1, 2, 3 or 4 heteroatoms, selected from N, O, and S, with the remaining ring atoms being carbon and wherein at least one ring is aromatic and at least one heteroatom is present in an aromatic ring.

[067] For example, the heteroaryl group includes a 5- to 7-membered heterocyclic aromatic ring fused to a 5- to 7-membered cycloalkyl ring. For such fused, bicyclic heteroaryl ring systems wherein only one of the rings comprises at least one heteroatom, the point of attachment may be at the heteroaromatic ring or at the cycloalkyl ring.

[068] When the total number of S and O atoms in the heteroaryl group exceeds 1, those heteroatoms are not adjacent to one another. In some embodiments, the total number of S and O atoms in the heteroaryl group is not more than 2. In some embodiments, the total number of S and O atoms in the aromatic heterocycle is not more than 1.

[069] Examples of the heteroaryl group include, but are not limited to, (as numbered from the linkage position assigned priority 1) pyridyl (such as 2-pyridyl, 3-pyridyl, or 4-pyridyl), cinnoliny, pyrazinyl, 2,4-pyrimidinyl, 3,5-pyrimidinyl, 2,4-imidazolyl, imidazopyridinyl, isoxazolyl, oxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, tetrazolyl, thienyl, triazinyl, benzothieryl, furyl, benzofuryl, benzoimidazolyl, indolyl, isoindolyl, indolinyl, phthalazinyl, pyrazinyl, pyridazinyl, pyrrolyl, triazolyl, quinoliny, isoquinoliny, pyrazolyl, pyrrolopyridinyl (such as 1H-pyrrolo[2,3-b]pyridin-5-yl), pyrazolopyridinyl (such as 1H-pyrazolo[3,4-b]pyridin-5-yl), benzoxazolyl (such as benzo[d]oxazol-6-yl), pteridinyl, purinyl, 1-oxa-2,3-diazolyl, 1-oxa-2,4-diazolyl, 1-oxa-2,5-diazolyl, 1-oxa-3,4-diazolyl, 1-thia-2,3-diazolyl, 1-thia-2,4-diazolyl, 1-thia-2,5-diazolyl, 1-thia-3,4-diazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxalinyl, naphthyridinyl, furopyridinyl, benzothiazolyl (such as benzo[d]thiazol-6-yl), indazolyl (such as 1H-indazol-5-yl) and 5,6,7,8-tetrahydroisoquinoline.

[070] The term "heterocyclic" or "heterocycle" or "heterocyclyl" refers to a ring selected from 4- to 12-membered monocyclic, bicyclic and tricyclic, saturated and partially unsaturated rings comprising at least one carbon atoms in addition to 1, 2, 3 or 4 heteroatoms, selected from oxygen, sulfur, and nitrogen. "Heterocycle" also refers to a 5- to 7-membered heterocyclic ring comprising at least one heteroatom selected from N, O, and S fused with 5-, 6-, and/or 7-membered cycloalkyl, carbocyclic aromatic or heteroaromatic ring, provided that the point of attachment is at the heterocyclic ring when the heterocyclic ring is fused with a carbocyclic aromatic or a heteroaromatic ring, and that the point of attachment can be at the cycloalkyl or heterocyclic ring when the heterocyclic ring is fused with cycloalkyl.

[071] "Heterocycle" also refers to an aliphatic spirocyclic ring comprising at least one heteroatom selected from N, O, and S, provided that the point of attachment is at the heterocyclic ring. The rings may be saturated or have at least one double bond (i.e. partially unsaturated). The heterocycle may be substituted with oxo. The point of the attachment may be carbon or heteroatom in the heterocyclic ring. A heterocycle is not a heteroaryl as defined herein.

[072] Examples of the heterocycle include, but not limited to, (as numbered from the linkage position assigned priority 1) 1-pyrrolidinyl, 2-pyrrolidinyl, 2,4-imidazolidinyl, 2,3-pyrazolidinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 2,5-piperazinyl, pyranlyl, 2-morpholinyl, 3-morpholinyl, oxiranyl, aziridinyl, thiranyl, azetidiny, oxetanyl, thietanyl, 1,2-dithietanyl, 1,3-dithietanyl, dihydropyridinyl, tetrahydropyridinyl, thiomorpholinyl, thioxanyl, piperazinyl, homopiperazinyl, homopiperidinyl, azepanyl, oxepanyl, thiepanyl, 1,4-oxathianyl, 1,4-dioxepanyl, 1,4-oxathiepanyl, 1,4-oxaazepanyl, 1,4-dithiepanyl, 1,4-thiazepanyl and 1,4-diazepane 1,4-dithianyl, 1,4-azathianyl, oxazepinyl, diazepinyl, thiazepinyl, dihydrothienyl, dihydropyranlyl, dihydrofuranyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranlyl, tetrahydrothiopyranlyl, 1-pyrrolinyl, 2-pyrrolinyl, 3-pyrrolinyl, indolinyl, 2H-pyranlyl, 4H-pyranlyl, 1,4-dioxanyl, 1,3-dioxolanyl, pyrazolinyl, pyrazolidinyl, dithianyl, dithiolanyl, pyrazolidinylimidazolinyl, pyrimidinonyl, 1,1-dioxo-thiomorpholinyl, 3-azabicyclo[3.1.0]hexanyl, 3-azabicyclo[4.1.0]heptanyl and azabicyclo[2.2.2]hexanyl. Substituted heterocycle also includes ring systems substituted with one or more oxo moieties, such as piperidinyl N-oxide, morpholinyl-N-oxide, 1-oxo-1-thiomorpholinyl and 1, 1-dioxo-1-thiomorpholinyl.

[073] Substituents are selected from: halogen, -R', -OR', =O, =NR', =N-OR', -NR'R'', -SR', -SiR'R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR'-SO₂NR''', -NR''CO₂R', -NH-C(NH₂)=NH, -NR''C(NH₂)=NH, -NH-C(NH₂)=NR', -S(O)R', -SO₂R', -SO₂NR'R'', -NR''SO₂R', -CN and -NO₂, -N₃, -CH(Ph)₂, perfluoro(C1-C4)alkoxy and perfluoro(C1-C4)alkyl, in a number ranging from zero to three, with those groups having zero, one or two substituents being particularly preferred. R', R'' and R''' each independently refer to hydrogen, unsubstituted (C1-C8)alkyl and heteroalkyl, unsubstituted aryl, aryl substituted with one to three halogens, unsubstituted alkyl, alkoxy or thioalkoxy groups, or aryl-(C1-C4)alkyl groups. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6- or 7-membered ring. Hence, -NR'R'' includes 1-pyrrolidinyl and 4-morpholinyl, "alkyl" includes groups such as trihaloalkyl (e.g., -CF₃ and -CH₂CF₃), and when the aryl group is 1,2,3,4-tetrahydronaphthalene, it may be substituted with a substituted or unsubstituted (C3-C7)spirocycloalkyl group. The (C3-C7)spirocycloalkyl group may be substituted in the same manner as defined herein for "cycloalkyl".

[074] Preferred substituents are selected from: halogen, -R', -OR', =O, -NR'R'', -SR', -SiR'R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR''CO₂R', -NR'-SO₂NR''R''', -S(O)R', -SO₂R', -SO₂NR'R'', -NR''SO₂R', -CN and -NO₂, perfluoro(C1-C4)alkoxy and perfluoro(C1-C4)alkyl, where R' and R'' are as defined above.

[075] The term "fused ring" herein refers to a polycyclic ring system, e.g., a bicyclic or tricyclic ring system, in which two rings share only two ring atoms and one bond in common. Examples of fused rings may comprise a fused bicyclic cycloalkyl ring such as those having

from 7 to 12 ring atoms arranged as a bicyclic ring selected from [4,4], [4,5], [5,5], [5,6] and [6,6] ring systems as mentioned above; a fused bicyclic aryl ring such as 7 to 12 membered bicyclic aryl ring systems as mentioned above, a fused tricyclic aryl ring such as 10 to 15 membered tricyclic aryl ring systems mentioned above; a fused bicyclic heteroaryl ring such as 8- to 12-membered bicyclic heteroaryl rings as mentioned above, a fused tricyclic heteroaryl ring such as 11- to 14-membered tricyclic heteroaryl rings as mentioned above; and a fused bicyclic or tricyclic heterocyclyl ring as mentioned above.

[076] The compounds may contain an asymmetric center and may thus exist as enantiomers. Where the compounds possess two or more asymmetric centers, they may additionally exist as diastereomers. Enantiomers and diastereomers fall within the broader class of stereoisomers.

All such possible stereoisomers as substantially pure resolved enantiomers, racemic mixtures thereof, as well as mixtures of diastereomers are intended to be included. All stereoisomers of the compounds and/or pharmaceutically acceptable salts thereof are intended to be included.

Unless specifically mentioned otherwise, reference to one isomer applies to any of the possible isomers. Whenever the isomeric composition is unspecified, all possible isomers are included.

[077] The term "substantially pure" means that the target stereoisomer contains no more than 35%, such as no more than 30%, further such as no more than 25%, even further such as no more than 20%, by weight of any other stereoisomer(s). In some embodiments, the term "substantially pure" means that the target stereoisomer contains no more than 10%, for example, no more than 5%, such as no more than 1%, by weight of any other stereoisomer(s).

[078] When compounds contain olefin double bonds, unless specified otherwise, such double bonds are meant to include both E and Z geometric isomers.

[079] Some of the compounds may exist with different points of attachment of hydrogen, referred to as tautomers. For example, compounds including carbonyl $-\text{CH}_2\text{C}(\text{O})-$ groups (keto forms) may undergo tautomerism to form hydroxyl $-\text{CH}=\text{C}(\text{OH})-$ groups (enol forms). Both keto and enol forms, individually as well as mixtures thereof, are also intended to be included where applicable.

[080] It may be advantageous to separate reaction products from one another and/or from starting materials. The desired products of each step or series of steps is separated and/or purified (hereinafter separated) to the desired degree of homogeneity by the techniques common in the art. Typically such separations involve multiphase extraction, crystallization from a solvent or solvent mixture, distillation, sublimation, or chromatography. Chromatography can involve any number of methods including, for example: reverse-phase and normal phase; size exclusion; ion exchange; high, medium and low pressure liquid chromatography methods and apparatus; small scale analytical; simulated moving bed ("SMB") and preparative thin or thick

layer chromatography, as well as techniques of small scale thin layer and flash chromatography. One skilled in the art will apply techniques most likely to achieve the desired separation.

[081] Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereoisomers to the corresponding pure enantiomers. Enantiomers can also be separated by use of a chiral HPLC column.

[082] A single stereoisomer, e.g., a substantially pure enantiomer, may be obtained by resolution of the racemic mixture using a method such as formation of diastereomers using optically active resolving agents (Eliel, E. and Wilen, S. Stereochemistry of Organic Compounds. New York: John Wiley & Sons, Inc., 1994; Lochmuller, C. H., et al. "Chromatographic resolution of enantiomers: Selective review." J. Chromatogr., 113(3) (1975): pp. 283-302). Racemic mixtures of chiral compounds of the invention can be separated and isolated by any suitable method, including: (1) formation of ionic, diastereomeric salts with chiral compounds and separation by fractional crystallization or other methods, (2) formation of diastereomeric compounds with chiral derivatizing reagents, separation of the diastereomers, and conversion to the pure stereoisomers, and (3) separation of the substantially pure or enriched stereoisomers directly under chiral conditions. See: Wainer, Irving W., Ed. Drug Stereochemistry: Analytical Methods and Pharmacology. New York: Marcel Dekker, Inc., 1993.

[083] "Pharmaceutically acceptable salts" include, but are not limited to salts with inorganic acids, selected, for example, from hydrochlorates, phosphates, diphosphates, hydrobromates, sulfates, sulfinates, and nitrates; as well as salts with organic acids, selected, for example, from malates, maleates, fumarates, tartrates, succinates, citrates, lactates, methanesulfonates, p-toluenesulfonates, 2-hydroxyethylsulfonates, benzoates, salicylates, stearates, alkanoates such as acetate, and salts with $\text{HOOC}-(\text{CH}_2)_n\text{-COOH}$, wherein n is selected from 0 to 4. Similarly, examples of pharmaceutically acceptable cations include, but are not limited to, sodium, potassium, calcium, aluminum, lithium, and ammonium.

[084] In addition, if a compound is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt. Conversely, if the product is a free base, an addition salt, such as a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds. Those


skilled in the art will recognize various synthetic methodologies that may be used without undue experimentation to prepare non-toxic pharmaceutically acceptable addition salts.

[085] "Treating," "treat," or "treatment" refers to administering at least one compound and/or at least one stereoisomer thereof, and/or at least one pharmaceutically acceptable salt thereof to a subject in recognized need thereof that has, for example, cancer.

[086] An "effective amount" refers to an amount of at least one compound and/or at least one stereoisomer thereof, and/or at least one pharmaceutically acceptable salt thereof effective to "treat" a disease or disorder in a subject, and that will elicit, to some significant extent, the biological or medical response of a tissue, system, animal or human that is being sought, such as when administered, is sufficient to prevent development of, or alleviate to some extent, one or more of the symptoms of the condition or disorder being treated. The therapeutically effective amount will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

[087] The term "at least one substituent" includes, for example, from 1 to 4, such as from 1 to 3, further as 1 or 2, substituents. For example, "at least one substituent R¹⁶" herein includes from 1 to 4, such as from 1 to 3, further as 1 or 2, substituents selected from the list of R¹⁶ as described herein.

[088] The invention provides compounds of formula:



[089] stereoisomers thereof, and pharmaceutically acceptable salts thereof.

[090] R¹ (and R⁴, R⁵, R⁶, and R⁷) are each independently H, halogen, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, alkynyl, -CN, -NR¹³R¹⁴, -OR¹³, -COR¹³, -CO₂R¹³, -CONR¹³R¹⁴, -C(=NR¹³)NR¹⁴R¹⁵, -NR¹³COR¹⁴, -NR¹³CONR¹⁴R¹⁵, -NR¹³CO₂R¹⁴, -SO₂R¹³, -NR¹³SO₂NR¹⁴R¹⁵, or -NR¹³SO₂R¹⁴, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶.

[091] R¹³, R¹⁴ and R¹⁵ are each independently H, heteroalkyl, alkyl, alkenyl, alkynyl, cycloalkyl, saturated or unsaturated heterocyclyl, aryl, or heteroaryl; wherein (R¹³ and R¹⁴),

and/or (R^{14} and R^{15}) together with the atom(s) to which they are attached, each can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R^{16} . In particular embodiments R^{13} , R^{14} and R^{15} are each independently H, lower alkyl, or lower alkoxy.

[092] R^{16} is halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocyclyl, oxo, -CN, -OR', -NR'R'', -COR', -CO₂R', -CONR'R'', -C(=NR')NR''R''', -NR'COR'', -NR'CONR'R'', -NR'CO₂R'', -SO₂R', -SO₂aryl, -NR'SO₂NR''R''', or -NR'SO₂R'', wherein R', R'', and R''' are independently hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocyclyl, wherein (R' and R''), and/or (R'' and R''') together with the atoms to which they are attached, can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings.

[093] R^{16} of R^1 in particular embodiments is halogen, lower alkyl, or lower alkoxy.

[094] R^1 in particular embodiments is an optionally hetero-, optionally substituted hydrocarbon selected from heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R^{16} , wherein cyclic structures are preferably 3-8 membered ring structures comprising 0-3 heteroatoms of N, S or O, and the aryl is preferably a 5- or 6-membered aromatic ring comprising 0-3 heteroatoms of N, S or O, wherein the hydrocarbon is preferably a C1-C12 or C1-C8 hydrocarbon.

[095] R^1 in particular embodiments is heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R^{16} .

[096] R^1 in particular embodiments is lower alkyl or alkenyl, optionally cyclic and optionally substituted, particularly with halogen, lower alkyl, or lower alkoxy, and comprising 0-3 heteroatoms of N, S or O. Examples include methylcyclopropyl, cyclohexyl, cyclopentyl, methoxyethyl, halide, methyl, ethyl, propyl and butyl. Other exemplary R^1 moieties include 5-membered aromatic rings like pyrrole, pyrazole, imidazole, furan and hydrogenations thereof (e.g. pyrrolidine, pyrazolidine, imidazolidine, tetrahydrofuran), and 6-membered rings like benzene (phenyl), pyridine, pyran, diazines, triazines and tetrazines, and hydrogenations thereof

(e.g. cyclohexane, di- and tetra-hydropyridine, piperidine, tetrahydropyran, etc.), each of which may be substituted, particularly with halogen, lower alkyl, or lower alkoxy.

[0097] L is a bond, CH₂, NR¹², O, or S, wherein R¹² is H or lower alkyl, e.g. methyl.

[0098] A is a 5- or 6-membered aromatic ring comprising 0-3 heteroatoms of N, S or O. Preferred 5-membered aromatic rings include pyrrole, pyrazole, imidazole, furan and 6-membered rings include benzene (phenyl), pyridine, pyran, diazines, triazines and tetrazines.

[0099] n is 0, 1, 2, 3 or 4, wherein when n is more than 1, each R² may be different. In particular embodiments n is 0 (i.e. A is unsubstituted).

[0100] R² is halogen, alkyl, -S-alkyl, -CN, -NR¹³R¹⁴, -OR¹³, -COR¹³, -CO₂R¹³, -CONR¹³R¹⁴, -C(=NR¹³)NR¹⁴R¹⁵, -NR¹³COR¹⁴, -NR¹³CONR¹⁴R¹⁵, -NR¹³CO₂R¹⁴, -SO₂R¹³, -NR¹³SO₂NR¹⁴R¹⁵, or -NR¹³SO₂R¹⁴; wherein R¹³, R¹⁴ and R¹⁵ are each independently H, heteroalkyl, alkyl, alkenyl, alkynyl, cycloalkyl, saturated or unsaturated heterocyclyl, aryl, or heteroaryl; wherein (R¹³ and R¹⁴), and/or (R¹⁴ and R¹⁵) together with the atom(s) to which they are attached, each can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶.

[0101] R² in particular embodiments is halogen, alkyl, -S-alkyl, -CN, -NR¹³R¹⁴, -OR¹³, -COR¹³, -CO₂R¹³, -CONR¹³R¹⁴, -C(=NR¹³)NR¹⁴R¹⁵, -NR¹³COR¹⁴, -NR¹³CONR¹⁴R¹⁵, -NR¹³CO₂R¹⁴, -SO₂R¹³, -NR¹³SO₂NR¹⁴R¹⁵, or -NR¹³SO₂R¹⁴, preferably wherein the alkyl (including -S-alkyl) is lower alkyl, and R¹³, R¹⁴ and R¹⁵ are each independently H, lower alkyl, or lower alkoxy.

[0102] R² in particular embodiments is halogen, lower alkyl, or lower alkoxy.

[0103] each W is independently -(CH₂)- or -C(O)-, wherein if m is 2, 3 or 4, preferably no more than one W is carbonyl;

[0104] S/D is a single or double bond, and when a double bond, R⁵ and R⁷ are absent;

[0105] m is 0, 1, 2, 3 or 4, preferably 0, 1, 2 or 3, and in particular embodiments 0, 1 or 2, or 0 or 1.

[0106] p is 0 (and R⁶ and R⁷ are absent), or an integer of 1-2, wherein when p is 0, m is non-zero, and when p is more than 1, each R⁶ and each R⁷ may be different; generally p+m is 1, 2, 3 or 4, preferably 1, 2 or 3, and in particular embodiments 1.


[0107] In particular embodiments p is 2 and m is 0 or 1; p is 1 and m is 0 or 1 (or 0, 1 or 2); or p is 0 and m is 1 or 2 (or 1, 2 or 3).

[0108] R⁴ and R⁵, or R⁴ and R⁶, or R⁶ and R⁷, or R⁶ and R⁶ (when p is 2), together with the atoms to which they are attached, can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶. These rings are generally 4-8-membered and include 5-membered aromatic

rings like pyrrole, pyrazole, imidazole, furan and hydrogenations thereof (e.g. pyrrolidine, pyrazolidine, imidazolidine, tetrahydrofuran), and 6-membered rings like benzene (phenyl), pyridine, pyran, diazines, triazines and tetrazines, and hydrogenations thereof (e.g. cyclohexane, di- and tetra-hydropyridine, piperidine, tetrahydropyran, etc.), each of which may be unsubstituted or substituted, particularly with halogen, lower alkyl, lower alkoxy, -COR', or NR'COR'', wherein R', R'' are substituted or unsubstituted alkenyl.

[0109] R⁴ and R⁵ (or R⁶ and R⁷, or R⁶ and R⁶, when p is 2), in particular embodiments form piperidine, azacycloheptanyl, or azetidine, optionally substituted, particularly N-substituted with moieties such as benzyl, acyl, acryloyl, etc.


[0110] R⁴ and R⁶, in particular embodiments, together with the atoms to which they are attached, form a ring of formula:




Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4.

[0111] R⁴ and R⁶ in particular embodiments form phenyl, piperidine, azacycloheptenyl, pyrrolidine, optionally substituted, particularly N-substituted with moieties such as benzyl, acyl, acryloyl, methylamine-acryloyl, etc.

[0112] The invention includes all combinations of the recited particular and preferred embodiments as if each combination had been laboriously separately recited. For example, in particular embodiments, supra, A is phenyl; W is -(CH₂)_n; L is O; S/D is a single bond; m is 1; n is 0; p is 1; R¹ is phenyl; R² is absent; R⁵ is H; and R⁶ and R⁷ are H; yielding the combination:



[0113] R⁴, supra, includes N-containing C1-C8 alkyl, N-containing C3-C8 cycloalkyl and phenyl, for example, methylamine, aniline, azetidine, pyrrolidine, piperidine, azacycloheptenyl, each optionally substituted, particularly N-substituted with moieties such as benzyl, acyl, acryloyl, substituted-acryloyl, propiolyl, substituted-propiolyl, etc., such as structure combinations:



[0114] In particular examples, R⁴ is 1-acryloylpiperidin-4-yl (e.g. compound 27) or 1-(but-2-ynoyl)piperidin-4-yl (e.g. compound 176).

[0115] The invention also provides all the compounds of the examples herein, and of Table I, II and III, stereoisomers thereof, and pharmaceutically acceptable salts thereof.

[0116] The subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof may be employed alone or in combination with at least one other therapeutic agent for treatment. In some embodiments, the compounds, stereoisomers thereof, and pharmaceutically acceptable salts thereof can be used in combination with at least one additional therapeutic agent. The at least one additional therapeutic agent can be, for example, selected from anti-hyperproliferative, anti-cancer, and chemotherapeutic agents. The compound and/or one pharmaceutically acceptable salt disclosed herein may be administered with the at least one other therapeutic agent in a single dosage form or as a separate dosage form. When administered as a separate dosage form, the at least one other therapeutic agent may be administered prior to, at the same time as, or following administration of the compound and/or one pharmaceutically acceptable salt disclosed herein.

[0117] A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer, regardless of mechanism of action. Chemotherapeutic agents include compounds used in "targeted therapy" and conventional chemotherapy. Suitable chemotherapeutic agents can be, for example, selected from: agents that induce apoptosis; polynucleotides (e.g., ribozymes); polypeptides (e.g., enzymes); drugs; biological mimetics; alkaloids; alkylating agents; antitumor antibiotics; antimetabolites; hormones; platinum compounds; monoclonal antibodies conjugated with anticancer drugs, toxins, and/or radionuclides; biological response modifiers (e.g., interferons, such as IFN- α and interleukins, such as IL-2); adoptive immunotherapy agents; hematopoietic growth factors; agents that induce tumor cell differentiation (e.g., all-trans-retinoic acid); gene therapy reagents; antisense therapy reagents and nucleotides; tumor vaccines; and inhibitors of angiogenesis.

[0118] Examples of chemotherapeutic agents include Erlotinib (TARCEVA®, Genentech/OSI Pharm.); Bortezomib (VELCADE®, Millennium Pharm.); Fulvestrant (FASLODEX®, AstraZeneca); Sunitinib (SUTENT®, Pfizer); Letrozole (FEMARA®, Novartis); Imatinib mesylate (GLEEVEC®, Novartis); PTK787/ZK 222584 (Novartis); Oxaliplatin (Eloxatin®, Sanofi); 5-FU (5-fluorouracil); Leucovorin; Rapamycin (Sirolimus, RAPAMUNE®, Wyeth); Lapatinib (TYKERB®, GSK572016, Glaxo Smith Kline); Lonafernib (SCH 66336); Sorafenib (NEXAVAR®, Bayer); Irinotecan (CAMPTOSAR®, Pfizer) and Gefitinib (IRESSA®, AstraZeneca); AG1478, AG1571 (SU 5271, Sugen); alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylenelamines such as altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylmelamine; acetogenins (such as bullatacin and bullatacinone); a camptothecin (such as the synthetic analog topotecan); bryostatin; calystatin; CC-1065 and its adozelesin, carzelesin and bizelesin synthetic analogs; cryptophycins (such as cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin and the synthetic analogs thereof, such as KW-2189 and CB1-TM1; eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, chlorophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, such as calicheamicin gammaII and calicheamicin omegaII (Angew Chem. Intl. Ed. Engl. (1994) 33:183-186); dynemicin, such as dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores, aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, caminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® (doxorubicin), morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, porfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogs such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone,

dromostanolone propionate, epitio stanol, mepitio stanone, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminol evulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diazi quone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, Oreg.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triazi quone; 2,2',2''-trichlorotriethylamine; trichothecenes (such as T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., TAXOL® (paclitaxel; Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE® (Cremophor-free), albumin-engineered nanoparticle formulations of paclitaxel (American Pharmaceutical Partners, Schaumburg, Ill.), and TAXOTERE® (doxetaxel; Rhone-Poulenc Rorer, Antony, France); chloranmbucil; GEMZAR® (gemcitabine); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE® (vinorelbine); novantrone; teniposide; edatrexate; daunomycin; aminopterin; capecitabine (XELODA®); ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; and pharmaceutically acceptable salts, acids and derivatives of any of the above.

[0119] The "chemotherapeutic agent" can also be selected, for example, from: (i) anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX®; tamoxifen citrate), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® (toremifine citrate); (ii) aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® (megestrol acetate), AROMASIN® (exemestane; Pfizer), formestanie, fadrozole, RIVISOR® (vorozole), FEMARA® (letrozole; Novartis), and ARIMIDEX® (anastrozole; AstraZeneca); (iii) anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); (iv) protein kinase inhibitors; (v) lipid kinase inhibitors; (vi) antisense oligonucleotides, such as those which inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC- α , Ral and H-Ras; (vii) ribozymes such as VEGF expression inhibitors (e.g.,

ANGIOZYME®) and HER2 expression inhibitors; (viii) vaccines such as gene therapy vaccines, for example, ALLOVECTIN®, LEUVECTIN®, and VAXID®; PROLEUKIN® rIL-2; a topoisomerase 1 inhibitor such as LURTOTECAN®; ABARELIX® rmRH; (ix) anti-angiogenic agents such as bevacizumab (AVASTIN®, Genentech); and (x) pharmaceutically acceptable salts, acids and derivatives of any of the above.

[0120] The "chemotherapeutic agent" can also be selected, for example, from therapeutic antibodies such as alemtuzumab (Campath), bevacizumab (AVASTIN®, Genentech); cetuximab (ERBITUX®, Imclone); panitumumab (VECTIBIX®, Amgen), rituximab (RITUXAN®, Genentech/Biogen Idec), pertuzumab (OMNITARG®, 2C4, Genentech), trastuzumab (HERCEPTIN®, Genentech), tositumomab (Bexxar, Corixa), and the antibody drug conjugate, gemtuzumab ozogamicin (MYLOTARG®, Wyeth).

[0121] Humanized monoclonal antibodies with therapeutic potential as chemotherapeutic agents in combination with a subject compound and stereoisomers thereof, and pharmaceutically acceptable salt thereof may, for example, be selected from: alemtuzumab, apolizumab, aselizumab, atlizumab, bapineuzumab, bevacizumab, bivatumab mertansine, cantuzumab mertansine, cedelizumab, certolizumab pegol, cidfusituzumab, cidtuzumab, daclizumab, eculizumab, efalizumab, epratuzumab, erlizumab, felvizumab, fontolizumab, gemtuzumab ozogamicin, inotuzumab ozogamicin, ipilimumab, labetuzumab, lintuzumab, matuzumab, mepolizumab, motavizumab, motovizumab, natalizumab, nimotuzumab, nolovizumab, numavizumab, ocrelizumab, omalizumab, palivizumab, pascolizumab, pecfusituzumab, pectuzumab, pertuzumab, pexelizumab, ralivizumab, ranibizumab, reslivizumab, reslizumab, resyvizumab, rovelizumab, ruplizumab, sibrotuzumab, siplizumab, sontuzumab, tacatuzumab tetraxetan, tadocizumab, talizumab, tefibazumab, tocilizumab, toralizumab, trastuzumab, tucotuzumab celmoleukin, tucusituzumab, umavizumab, urtoxazumab, and visilizumab.

[0122] Also provided is a composition comprising a subject compound and stereoisomers thereof, and pharmaceutically acceptable salts thereof, and at least one pharmaceutically acceptable carrier.

[0123] The composition comprising a subject compound and stereoisomers thereof, and pharmaceutically acceptable salts thereof can be administered in various known manners, such as orally, topically, rectally, parenterally, by inhalation spray, or via an implanted reservoir, although the most suitable route in any given case will depend on the particular host, and nature and severity of the conditions for which the active ingredient is being administered. The term "parenteral" as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional and intracranial

injection or infusion techniques. The compositions disclosed herein may be conveniently presented in unit dosage form and prepared by any of the methods well known in the art.

[0124] The subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof can be administered orally in solid dosage forms, such as capsules, tablets, troches, dragées, granules and powders, or in liquid dosage forms, such as elixirs, syrups, emulsions, dispersions, and suspensions. The subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof disclosed herein can also be administered parenterally, in sterile liquid dosage forms, such as dispersions, suspensions or solutions. Other dosage forms that can also be used to administer the subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof disclosed herein as an ointment, cream, drops, transdermal patch or powder for topical administration, as an ophthalmic solution or suspension formation, i.e., eye drops, for ocular administration, as an aerosol spray or powder composition for inhalation or intranasal administration, or as a cream, ointment, spray or suppository for rectal or vaginal administration.

[0125] Gelatin capsules containing the compound and/or the at least one pharmaceutically acceptable salt thereof disclosed herein and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like, can also be used. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of time. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

[0126] Liquid dosage forms for oral administration can further comprise at least one agent selected from coloring and flavoring agents to increase patient acceptance.

[0127] In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols can be examples of suitable carriers for parenteral solutions. Solutions for parenteral administration may comprise a water soluble salt of the at least one compound describe herein, at least one suitable stabilizing agent, and if necessary, at least one buffer substance. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, can be examples of suitable stabilizing agents. Citric acid and its salts and sodium EDTA can also be used as examples of suitable stabilizing agents. In addition, parenteral solutions can further comprise at least one preservative, selected, for example, from benzalkonium chloride, methyl- and propylparaben, and chlorobutanol.

[0128] A pharmaceutically acceptable carrier is, for example, selected from carriers that are compatible with active ingredients of the composition (and in some embodiments, capable of stabilizing the active ingredients) and not deleterious to the subject to be treated. For example, solubilizing agents, such as cyclodextrins (which can form specific, more soluble complexes with the at least one compound and/or at least one pharmaceutically acceptable salt disclosed herein), can be utilized as pharmaceutical excipients for delivery of the active ingredients. Examples of other carriers include colloidal silicon dioxide, magnesium stearate, cellulose, sodium lauryl sulfate, and pigments such as D&C Yellow # 10. Suitable pharmaceutically acceptable carriers are described in *Remington's Pharmaceutical Sciences*, A. Osol, a standard reference text in the art.

[0129] The subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof disclosed herein can further be examined for efficacy in treating Btk related diseases by *in vivo* assays. For example, the compound and/or the at least one pharmaceutically acceptable salts thereof disclosed herein can be administered to an animal (e.g., a mouse model) having Btk related diseases and its therapeutic effects can be accessed. Positive results in one or more of such tests are sufficient to increase the scientific storehouse of knowledge and hence sufficient to demonstrate practical utility of the compounds and/or salts tested. Based on the results, an appropriate dosage range and administration route for animals, such as humans, can also be determined.

[0130] For administration by inhalation, the subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or nebulisers. The subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof may also be delivered as powders, which may be formulated and the powder composition may be inhaled with the aid of an insufflation powder inhaler device. One exemplary delivery system for inhalation can be metered dose inhalation (MDI) aerosol, which may be formulated as a suspension or solution of a subject compound and stereoisomers thereof, and pharmaceutically acceptable salts thereof disclosed herein in at least one suitable propellant, selected, for example, from fluorocarbons and hydrocarbons.

[0131] For ocular administration, an ophthalmic preparation may be formulated with an appropriate weight percentage of a solution or suspension of the subject compound and stereoisomers thereof, and pharmaceutically acceptable salts thereof in an appropriate ophthalmic vehicle, such that the subject compound and stereoisomers thereof, and at least one pharmaceutically acceptable salts thereof is maintained in contact with the ocular surface for a

sufficient time period to allow the compound to penetrate the corneal and internal regions of the eye.

[0132] Useful pharmaceutical dosage-forms for administration of the subject compounds and stereoisomers thereof, and pharmaceutically acceptable salts thereof disclosed herein include, but are not limited to, hard and soft gelatin capsules, tablets, parenteral injectables, and oral suspensions.

[0133] The dosage administered will be dependent on factors, such as the age, health and weight of the recipient, the extent of disease, type of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. In general, a daily dosage of the active ingredient can vary, for example, from 0.1 to 2000 milligrams per day. For example, 10- 500 milligrams once or multiple times per day may be effective to obtain the desired results.

[0134] In some embodiments, a large number of unit capsules can be prepared by filling standard two-piece hard gelatin capsules each with, for example, 100 milligrams of the subject compound and stereoisomers thereof, and pharmaceutically acceptable salt thereof disclosed herein in powder, 150 milligrams of lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

[0135] In some embodiments, a mixture of the compound, stereoisomers thereof, and pharmaceutically acceptable salts thereof a digestible oil such as soybean oil, cottonseed oil or olive oil can be prepared and injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

[0136] In some embodiments, a large number of tablets can be prepared by conventional procedures so that the dosage unit comprises, for example, 100 milligrams of the compound, stereoisomers thereof, and pharmaceutically acceptable salts thereof, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275 milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

[0137] In some embodiments, a parenteral composition suitable for administration by injection can be prepared by stirring 1.5% by weight of the compound and/or at least an enantiomer, a diastereomer, or pharmaceutically acceptable salt thereof disclosed herein in 10% by volume propylene glycol. The solution is made to the expected volume with water for injection and sterilized.

[0138] In some embodiment, an aqueous suspension can be prepared for oral administration. For example, each 5 milliliters of an aqueous suspension comprising 100 milligrams of finely divided compound, stereoisomers thereof, and pharmaceutically acceptable salts thereof, 100

milligrams of sodium carboxymethyl cellulose, 5 milligrams of sodium benzoate, 1.0 grams of sorbitol solution, U.S.P., and 0.025 milliliters of vanillin can be used.

[0139] The same dosage forms can generally be used when the compound, stereoisomers thereof, and pharmaceutically acceptable salts thereof are administered stepwise or in conjunction with at least one other therapeutic agent. When drugs are administered in physical combination, the dosage form and administration route should be selected depending on the compatibility of the combined drugs. Thus the term coadministration is understood to include the administration of at least two agents concomitantly or sequentially, or alternatively as a fixed dose combination of the at least two active components.

[0140] The compounds, stereoisomers thereof, and pharmaceutically acceptable salt thereof disclosed herein can be administered as the sole active ingredient or in combination with at least one second active ingredient, selected, for example, from other active ingredients known to be useful for treating Btk related diseases in a patient.


[0141] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein, including citations therein, are hereby incorporated by reference in their entirety for all purposes.

[0142] GENERAL REACTION SCHEME FOR COMPOUND PREPARATION

[0143] The subject compounds and pharmaceutically acceptable salts thereof, can be prepared from (a) commercially available starting materials (b) known starting materials which may be prepared as described in literature procedures (c) new intermediates described in the schemes and experimental procedures herein. In making the compounds of the invention, the order of synthetic steps may be varied to increase the yield of desired product. Some of compounds in this invention may be generated by the methods as shown in the following reaction schemes and the description thereof.

[0144]


Scheme I



[0145] Scheme I above shows a general synthetic route that is used for preparing the compound S-6 of this invention, where A, R₁, R₂, R₄, R₆, L and n are as described herein. Reaction of methyl ketone S-1 in acetals of *N,N*-dialkylformamides or acetals of *N,N*-dialkylacetamide at reflux temperature for several hours afforded 3-dialkylamino-1-(aryl, heteroaryl or alkyl)-2-propen-1-one S-2. Intermediate S-3 may be prepared by methods substantially similar to those described in International Patent Publication No. WO 2001/019829 and No. WO 2011/046964. The reaction of intermediate S-3 and an appropriately substituted 3-dialkylamino-1-(aryl, heteroaryl or alkyl)-2-propen-1-one S-2 in weak acid such as acetic acid or in an inert solvent such as toluene, acetonitrile or dimethoxyethane with catalytic acid, at 80 °C to reflux temperature for several hours afforded the nitrile compound S-4. Reduction of the pyrimidine ring with reducing agents such as sodium borohydride (NaBH₄), Pd/C or NaBH₄ followed with Pd/C gave tetrahydropyrazolopyrimidine S-5, and subsequent hydrolysis of nitrile under alkaline condition such as NaOH or KOH plus H₂O₂ in alcohol, or under acid condition such as H₃PO₄, H₂SO₄, or BF₃·HOAc, yielded the carboxamide S-6.

[0146]


Scheme II



[0147] Scheme II describes a general synthetic route to prepare carboxamide **S-13**, where A, R₁, R₂, R₄, R₅, L, n and p are as described herein. The protected hydrazine **S-7** known can be conveniently prepared by the methods as described in the literature (J. Med. Chem. 1992, 35, 2392). Deprotection with acid, followed by condensation of this building block with intermediate **S-10** (also described in International Patent Publication No. WO 2001/019829 and No. WO 2011/046964) in alkaline solvent such as TEA/ethanol afforded pyrazole ester **S-9**. The pyrazole alcohol **S-11** can be prepared from an ester **S-9** through a reductive process. The reducing agents which may be used for this process include, but are not limited to LiBH₄, NaBH₄ and Super Hydride. Intramolecular *N*-alkylation or reductive amination gave the nitriles **S-12**, which was converted to carboxamide **S-13** by the same methods as described in Scheme I.

[0148]


Scheme III



[0149] Scheme III describes alternative routes to prepare carboxamide **S-13**.

[0150]


Scheme IV



[0151] Scheme IV above shows general synthetic routes that have been used for preparing compounds **S-19** and **S-20** of this invention, where **A**, **R₁**, **R₂**, **R₄**, **R₆**, **L** and **n** are as described herein. Hydrolysis of nitrile **S-3** afforded carboxamide **S-18** by the same methods as described in Scheme I. The cyclization of pyrazole carboxamide **S-18** or pyrazole nitrile **S-3** with the commercially available or prepared haloketone **S-23** gave regioisomers **S-19** and **S-20**, or **S-21** and **S-22**. The nitriles **S-21** and **S-22** were hydrolyzed to afford carboxamides **S-19** and **S-20** according to Scheme I.

[0152]

Scheme V




[0153] Scheme V above shows general synthetic routes that have been used for preparing compound **S-27** of this invention, where **A**, **R₁**, **R₂**, **R₄**, **R₆**, **L**, **m** and **n** are as described herein. The pyrazole **S-25** was prepared by cyclization of intermediate **S-10** with the commercially available or prepared hydrazine **S-24** in alcohol according to one of steps in Scheme II. The intramolecular cyclization of intermediate **S-25** was performed with the process included, but not limited to Cu catalytic coupling, Buchwald reaction, SN₂ reaction or reductive amination. Finally, nitrile **S-26** was hydrolyzed to afford carboxamide **S-27** according to Scheme I.

Alternatively, nucleophilic substitution, followed by intramolecular cyclization afforded carboxamide **S-27**.

[0154]


Scheme VI



[0155] Scheme VI above shows a general synthetic route that has been used for preparing compound **S-30** of this invention, where A , R_1 , R_2 , R_4 , R_6 , L , m and n are as described herein. The intramolecular amide **S-30** was prepared by heating the mixture of nitrile **S-3** or carboxamide **S-18** and ester **S-29** in inert solvent such as DMF with presence of base such as K_2CO_3 and TEA.

[0156]


Scheme VII



[0157] Scheme VII above shows a general synthetic route that has been used for preparing compound **S-32** of this invention, where A , R_1 , R_2 , R_4 , L and n are as described herein. The amide **S-32** was prepared by heating the mixture of nitrile **S-3** or carboxamide **S-18** and β -ketoester **S-31** in weak acid such as acetic acid or in an inert solvent such as toluene, acetonitrile or dimethoxyethane with catalytic acid.

[0158]

Scheme VIII



[0159] Scheme VIII above shows a general synthetic route that has been used for preparing compound **S-36** of this invention, when A, R₁, R₂, R₆, R₇, L, W, m and n are as described herein. The intermolecular cyclization of nitrile **S-3** or carboxamide **S-18** and commercially available or prepared intermediate **S-33** was performed in inert solvent such as DMF with presence of base such as K₂CO₃ and TEA to afford compound **S-36**. Alternatively, condensation of nitrile **S-3** or carboxamide **S-18** with dialdehyde **S-34**, followed by reduction yielded compound **S-36**.

[0160] EXAMPLES

[0161] The examples below are intended to be purely exemplary and should not be considered to be limiting in any way. Efforts have been made to ensure accuracy with respect to numbers used (for example, amounts, temperature, etc.), but some experimental errors and deviations should be accounted for. Unless indicated otherwise, temperature is in degrees Centigrade. Reagents were purchased from commercial suppliers such as Sigma-Aldrich, Alfa Aesar, or TCI, and were used without further purification unless indicated otherwise.

[0162] Unless indicated otherwise, the reactions set forth below were performed under a positive pressure of nitrogen or argon or with a drying tube in anhydrous solvents; the reaction flasks were fitted with rubber septa for the introduction of substrates and reagents via syringe; and glassware was oven dried and/or heat dried.

[0163] ¹H NMR spectra were recorded on a Agilent instrument operating at 400 MHz. ¹HNMR spectra were obtained using CDCl₃, CD₂Cl₂, CD₃OD, D₂O, d₆-DMSO, d₆-acetone or (CD₃)₂CO as solvent and tetramethylsilane (0.00 ppm) or residual solvent (CDCl₃: 7.25 ppm; CD₃OD: 3.31 ppm; D₂O: 4.79 ppm; d₆-DMSO: 2.50 ppm; d₆-acetone: 2.05; (CD₃)₂CO: 2.05) as the reference standard. When peak multiplicities are reported, the following abbreviations are used: s (singlet), d (doublet), t (triplet), q (quartet), qn (quintuplet), sx (sextuplet), m (multiplet), br (broadened),

dd (doublet of doublets), dt (doublet of triplets). Coupling constants, when given, are reported in Hertz (Hz).

[0164] LC-MS spectrometer (Agilent 1260) Detector: MWD (190-400 nm), Mass detector: 6120 SQ

Mobile phase: A: acetonitrile with 0.1% Formic acid, B: water with 0.1% Formic acid

Column: Poroshell 120 EC-C18, 4.6×50 mm, 2.7µm

Gradient method: Flow: 1.8 mL/min

Time (min) A (%) B (%)

0.00	5	95
1.5	95	5
2.0	95	5
2.1	5	95
3.0	5	95

[0165] Preparative HPLC was conducted on a column (150 × 21.2 mm ID, 5 µm, Gemini NX-C18) at a flow rate of 20 ml/min, injection volume 2 ml, at room temperature and UV Detection at 214 nm and 254 nm.


[0166] In the following examples, the abbreviations below are used:

Brine	Saturated aqueous sodium chloride solution
BnNH ₂	Benzyl amine
CbzCl	Benzyl chloroformate
DCM	Dichloromethane
DCE	1,2-Dichlorethane
DIEA	<i>N,N</i> -diisopropylethylamine
DMF	<i>N,N</i> -Dimethylformamide
DMF-DMA	<i>N,N</i> -dimethylformamide dimethylacetal
DMAP	<i>N,N</i> -dimethylpyridin-4-amine
DCC	<i>N,N</i> -dicyclohexylcarbodiimide
DHP	3,4-Dihydro-2H-pyran
EA	Ethyl acetate
HATU	O-(7-Azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
HOAc	Acetic acid
H ₂ O ₂	Hydrogen peroxide solution 30 % (w/w) in H ₂ O
IPA	Isopropanol
MTBE	Methyl tert-butyl ether


MsCl	Methanesulfonyl chloride
NBS	<i>N</i> -Bromosuccinimide
Pd/C	Palladium on carbon powder
Pd(OH) ₂ /C	Palladium hydroxide on carbon powder
PE	Petroleum ether
PPh ₃	Triphenylphosphine
Pre-TLC	Prepared thin layer chromatography
sat.	Saturated
Tf ₂ O	Trifluoromethanesulfonic anhydride
THF	Tetrahydrofuran
TEA	Triethylamine
TMSCHN ₂	(Trimethylsilyl)diazomethane
TMSCl	Chlorotrimethylsilane

[0167] Example 1: Synthesis of compounds 1-4

[0168] Compound 1: 7-(3-Aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0169] Step 1: 2-(Hydroxy(4-phenoxyphenyl)methylene)malononitrile




[0170] A solution of 4-phenoxybenzoic acid (300 g, 1.4 mol) in SOCl₂ (1.2 L) was stirred at 80 °C under N₂ for 3 hr. The mixture was concentrated in vacuum to give the intermediate (315 g) which was used for next step without further purification.

[0171] To a solution of propanedinitrile (89.5 g, 1355 mmol) and DIEA (350 g, 2710 mmol) in THF (800 mL) was dropwise a solution of the intermediate (315 g) in toluene (800 mL) at 0~5 °C over 2 hr. The resultant mixture was allowed to warm to RT and stirred for 16 hr. The reaction was quenched with water (2.0 L) and extracted with of EA (2.0 L × 3). The combined organic layers were washed with 1000 mL of 3 N HCl aqueous solution, brine (2.0 L × 3), dried


over Na₂SO₄ and concentrated to give the crude product (330 g, 93%). ¹H NMR (DMSO-d₆) δ 7.62 (d, *J* = 8.8 Hz, 2H), 7.46-7.38 (m, 2H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.06 (d, *J* = 8.0 Hz, 2H), 6.94 (d, *J* = 8.8 Hz, 2H). MS (ESI) *m/e* [M+1]⁺ 262.9.

[0172] Step 2: 2-(Methoxy(4-phenoxyphenyl)methylene)malononitrile




[0173] A solution of 2-(hydroxy(4-phenoxyphenyl)methylene)malononitrile (50 g, 190.8 mmol) in CH(OMe)₃ (500 mL) was heated to 75 °C for 16 hr. Then the mixture was concentrated to a residue and washed with MeOH (50 mL) to give 25 g (47.5%) of 2-(methoxy(4-phenoxyphenyl)methylene)malononitrile as a yellow solid. ¹H NMR (DMSO-d₆) δ 7.70 (d, *J* = 8.4 Hz, 2H), 7.52-7.45 (m, 2H), 7.28 (t, *J* = 7.6 Hz, 1H), 7.22-7.06 (m, 4H), 3.93 (s, 3H). MS (ESI) *m/e* [M+1]⁺ 276.9.

[0174] Step 3: 5-Amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile




[0175] To a solution of 2-(methoxy(4-phenoxyphenyl)methylene)malononitrile (80 g, 290 mmol) in ethanol (200 mL) was added hydrazine hydrate (20 mL). The mixture was stirred at RT for 16 hr then was concentrated to give the crude product and washed with MeOH (30 mL) to afford 55 g (68.8%) of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile as a off-white solid. ¹H NMR (DMSO-d₆) δ 12.11 (br s, 1H), 7.80 (d, *J* = 8.8 Hz, 2H), 7.46-7.39 (m, 2H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.12-7.04 (m, 4H), 6.43 (br s, 2H).

[0176] Step 4: (E)-N-(3-(3-(dimethylamino)acryloyl)phenyl)acetamide




[0177] A solution of *N*-(3-acetylphenyl)acetamide (1.77 g, 10.0 mmol) in DMF-DMA (6 mL) with molecular sieve (10 portions) was stirred at 100 °C under N₂ for 2 hr. the mixture was concentrated and washed with MTBE (30 mL) to afford 2.1 g (90%) of (E)-*N*-(3-(3-(dimethylamino)acryloyl)phenyl)acetamide as a yellow solid.

[0178] Step 5: *N*-(3-(3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)phenyl)acetamide




[0179] To a solution of (E)-*N*-(3-(3-(dimethylamino)acryloyl)phenyl)acetamide (46 mg, 0.2 mmol) in HOAc (5 mL) was added 5-amino-3-(4-phenoxyphenyl)-1*H*-pyrazole-4-carbonitrile (55 mg, 0.2 mmol). The mixture was stirred at 118 °C for 4 hr. Then the reaction mixture was concentrated to a residue and partitioned between ethyl acetate (100 mL) and brine (100 mL). Organic layer was separated, washed with brine (2×100 mL), dried over sodium sulfate and concentrated to afford 80 mg (90%) of *N*-(3-(3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-*a*]pyrimidin-7-yl)phenyl)acetamide as a colorless oil. MS (ESI) *m/e* [M+1]⁺ 446.

[0180] Step 6: 7-(3-Aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-*a*]pyrimidine-3-carbonitrile




[0181] To a solution of *N*-(3-(3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-*a*]pyrimidin-7-yl)phenyl)acetamide (285 mg, 0.64 mmol) in ethanol (6 mL) was added HCl (3 mL). The mixture stirred at 75 °C for 3 hr. Concentrated to afford 250 mg (97%) of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-*a*]pyrimidine-3-carbonitrile as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.90 (d, *J* = 4.4 Hz, 1H), 8.67 (br s, 2H), 8.15 (d, *J* = 8.8 Hz, 2H), 8.06 (s, 1H), 7.87 (d, *J* = 8.0 Hz, 1H), 7.64 (t, *J* = 8.0 Hz, 1H), 7.57 (d, *J* = 4.4 Hz, 1H), 7.48-7.44 (m, 3H), 7.25-7.18 (m, 3H), 7.13 (d, *J* = 8.0 Hz, 2H). MS (ESI) *m/e* [M+1]⁺ 404.

[0182] Compound **2**: 7-(3-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-*a*]pyrimidine-3-carboxamide




[0183] Step 1: 7-(3-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-*a*]pyrimidine-3-carbonitrile



[0184] To a solution of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (150 mg, 0.372 mmol) in ethanol (10 mL) was added NaBH₄ (70 mg, 1.86 mmol). The mixture was stirred at rt for 16 hr and 60 °C for 2 hr. Then the reaction mixture was concentrated to a residue and partitioned between EA (50 mL) and brine (40 mL). Organic layer was separated from aqueous layer, washed with brine (50 mL × 2), dried over Na₂SO₄ and concentrated to afford 135 mg (crude) of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile as a yellow solid. MS (ESI) m/e [M+1]⁺ 408.

[0185] Step 2: 7-(3-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0186] To a solution of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (130 mg, 0.32 mmol) in DMSO (2 mL) and ethanol (2 mL) was added a solution of 5 N NaOH aqueous solution (1 mL) and H₂O₂ (1 mL). The mixture was stirred at 60 °C for 30 minutes, concentrated and partitioned between EA (100 mL) and brine (100 mL). Organic layer was separated, washed with brine (3×100 mL), dried over Na₂SO₄ and purified by chromatography column on silica gel eluting with PE/EA to afford 35 mg (26%) of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 7.50 (d, *J* = 8.4 Hz, 2H), 7.40-7.33 (m, 2H), 7.16 (t, *J* = 7.6 Hz, 1H), 7.06 (d, *J* = 8.0 Hz, 2H), 7.03 (d, *J* = 8.4 Hz, 2H), 6.97 (t, *J* = 7.6 Hz, 1H), 6.76 (s, 1H), 6.44 (d, *J* = 8.4 Hz, 1H), 6.23-6.21 (m, 2H), 5.30-5.25 (m, 1H), 5.09 (s, 2H), 3.30-3.28 (m, 1H), 3.12-3.02 (m, 1H), 2.34-2.26 (m, 1H), 2.05-2.01 (m, 1H). MS (ESI) m/e [M+1]⁺ 426.

[0187] Compound **2** was separated into two enantiomeric stereoisomers compound **2a** (peak 1, R or S, retention time at 8.94 min in chiral analysis), and compound **2b** (peak 2, S or R,


retention time at 10.11 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.

Column	CHIRALCEL AS-H
Column size	2 cm × 25 cm
Injection	2 mL
Mobile phase	CO ₂ /MeOH = 70/30
Flow rate	45 mL/min
Wave length	UV 210 nm
Temperature	35 °C
Sample solution	6 mg/mL in mobile phase
Prep-SFC equipment	DAICEL-SFC

[0188] The chiral analysis conditions are shown below.

Column	CHIRALPAK AD-H
Column size	0.46 cm I.D. × 15 cm L, 5 μm
Injection	2 μL
Mobile phase	n-Hexane/EtOH (0.1% triethyl amine) = 50/50 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm

[0189] Compound **3** and **4**: 7-(3-Acrylamidophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide and 7-(3-(3-chloropropanamido)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0190] To a solution of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (30 mg, 0.071 mmol) in DCM (2 mL) was added pyridine (0.2 mL). Then acryloyl chloride (6 mg, 0.084 mmol) was added dropwise. The mixture was stirred at RT for 0.5 hr and partitioned between DCM (20 mL) and brine (20 mL). Organic layer was separated from aqueous layer, washed with brine (2×20 mL), dried over Na₂SO₄ and purified by Pre-TLC (DCM/CH₃OH = 10/1) to afford 1.82 mg (5.38%) of 7-(3-acrylamidophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-

carboxamide as a white solid. ^1H NMR (400 MHz, CD_3OD) δ 7.58 (d, J = 8.8 Hz, 1H), 7.46 (d, J = 8.8 Hz, 2H), 7.34-7.26 (m, 4H), 7.10 (t, J = 7.6 Hz, 1H), 7.04-6.95 (m, 4H), 6.84 (d, J = 7.6 Hz, 1H), 6.39-6.25 (m, 2H), 5.69 (dd, J = 2.4, 9.6 Hz, 1H), 5.47-5.44 (m, 1H), 3.38-3.31 (m, 1H), 3.22-3.12 (m, 1H), 2.52-2.42 (m, 1H), 2.23-2.17 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 480.


[0191] 2.21 mg of byproduct 7-(3-(3-chloropropanamido)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide as a white solid. ^1H NMR (400 MHz, CD_3OD) δ 7.57 (d, J = 8.0 Hz, 1H), 7.52 (d, J = 8.8 Hz, 2H), 7.42-7.30 (m, 4H), 7.16 (t, J = 7.6 Hz, 1H), 7.09-7.01 (m, 4H), 6.89 (d, J = 8.0 Hz, 1H), 5.53-5.48 (m, 1H), 3.85 (t, J = 6.4 Hz, 2H), 3.44-3.37 (m, 1H), 3.26-3.19 (m, 1H), 2.83 (t, J = 6.4 Hz, 2H), 2.60-2.44 (m, 1H), 2.33-2.22 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 516.2.

[0192] Compound **3a** (peak 1, R or S, retention time at 4.45 min) and **3b** (peak 2, R or S, retention time at 7.41 min) was prepared from **2a** and **2b** according to the procedure similar to those for compound **3**.

[0193] The chiral analysis conditions are shown below.

Column	CHIRALPAK AD-H
Column size	0.46 cm I.D. $\times$ 15 cm L, 5 μm
Injection	2 μL
Mobile phase	n-Hexane/EtOH (0.1% triethyl amine) = 50/50 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 254 nm

[0194] Compound **5**: 7-(2-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide



[0195] The desired product was prepared from 1-(2-nitrophenyl)ethanone and 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile according to the procedures similar to those for 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (step 4 to step 5), compound **2** (step 1 and 2) and compound **68** (step 8) under appropriate conditions recognized by one of ordinary skill in the art. ^1H NMR ($\text{DMSO}-d_6$) δ 7.48-7.44 (m, 2H), 7.40-7.33 (m, 2H), 7.15-7.09 (m, 1H), 7.05-6.97 (m, 4H), 6.92 (td, J = 8.0, 1.2 Hz, 1H), 6.75 (br s, 1H), 6.64 (dd, J = 8.0, 1.2 Hz, 1H), 6.46 (td, J = 7.6, 1.2 Hz, 1H), 6.23 (dd, J = 7.6, 1.2 Hz, 1H),

5.57-5.52 (m, 1H), 5.19 (br s, 2H), 3.27-3.17 (m, 1H), 2.93 (td, $J = 2.8, 12.0$ Hz, 1H), 2.21-2.11 (m, 1H), 2.10-2.05 (m, 1H). MS (ESI) m/e $[M+1]^+$ 426.0.


[0196] Compound **5** was separated into two enantiomeric stereoisomers compound **5a** (peak 1, R or S, retention time at 7.30 min in chiral analysis), and compound **5b** (peak 2, S or R, retention time at 9.68 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.

Column	CHIRALCEL OD-H
Column size	3 cm $\times$ 25 cm
Injection	8 mL
Mobile phase	Hexane/IPA = 50/50
Flow rate	20 mL/min
Wave length	UV 254 nm
Temperature	35
Sample solution	4 mg/mL in mobile phase
Prep-SFC equipment	DAICEL-YMC

[0197] The chiral analysis conditions are shown below.

Column	CHIRALPAK IC
Column size	0.46 cm I.D. $\times$ 15 cm L
Injection	2 μ L
Mobile phase	MeOH = 100 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 254 nm

[0198] Compound **6**: 7-(2-Acrylamidophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide



[0199] The desired compound was prepared from compound **5** and acryloyl chloride according to the procedure similar to that for compound **3**. ^1H NMR (DMSO- d_6) δ 9.81 (s, 1H), 7.50-7.32

(m, 5H), 7.26 (td, $J = 7.6, 1.2$ Hz, 1H), 7.21-7.08 (m, 2H), 7.04-6.98 (m, 4H), 6.79 (s, 1H), 6.60 (dd, $J = 7.6, 1.2$ Hz, 1H), 6.50 (dd, $J = 17.0, 10.2$ Hz, 1H), 6.24 (dd, $J = 17.0, 1.9$ Hz, 1H), 5.77-5.74 (m, 2H), 3.26-3.22 (m, 1H), 2.98-2.92 (m, 1H), 2.32-2.25 (m, 1H), 1.96-1.93 (m, 1H). MS (ESI) m/e $[M+1]^+$ 480.


[0200] Compound **6a** (peak 1, R or S, retention time at 4.02 min) and **6b** (peak 2, R or S, retention time at 6.68 min) was prepared from **5a** and **5b** according to the procedure similar to that for compound **3**.

[0201] The chiral analysis conditions are shown below.

Column	CHIRALPAK AD-H
Column size	0.46 cm I.D. $\times$ 15 cm L, 5 μ m
Injection	2 μ L
Mobile phase	n-Hexane/EtOH (0.1% triethyl amine) = 70/30 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm


[0202] Example 2: Synthesis of compounds 7 and 8

[0203] Compound **7**: 7-(3-Aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a] pyrimidine-3-carboxamide



[0204] To a solution of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide (4 mg, 0.01 mmol) in DCE (2 mL) was added active MnO_2 (100 mg, 1.15 mmol). The mixture was stirred at 75 $^{\circ}C$ for 2 hr and filtered. The filtrate was purified by Pre-TLC (DCM/ CH_3OH = 10/1) to afford 2 mg (50%) of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carboxamide as a yellow solid. 1H NMR (400 MHz, CD_3OD-d_4 and $CDCl_3-d_1$) δ 8.62 (d, $J = 4.4$ Hz, 1H), 7.88 (d, $J = 8.8$ Hz, 2H), 7.41-7.39 (m, 2H), 7.31-7.26 (m, 3H), 7.11 (d, $J = 4.4$ Hz, 1H), 7.10-7.04 (m, 1H), 7.01-6.97 (m, 4H), 6.90-6.86 (m, 1H).


[0205] Compound **8**: 7-(3-Acrylamidophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a] pyrimidine-3-carboxamide




[0206] To a solution of 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a] pyrimidine-3-carboxamide (100 mg, 0.24 mmol) in DCM (10 mL) was added TEA (3 drops), followed by acryloyl chloride (32 mg, 0.36 mmol). The mixture was stirred at RT for 1 min and partitioned between DCM (50 mL) and brine (50 mL). Organic layer was separated from aqueous layer, dried over Na₂SO₄, concentrated and purified by Pre-TLC (DCM/CH₃OH = 10/1) to afford 12 mg (11%) of 7-(3-acrylamido phenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carboxamide as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 10.42 (s, 1H), 8.80 (d, *J* = 4.4 Hz, 1H), 8.47 (s, 1H), 8.07 (s, 1H), 7.95 (d, *J* = 8.8 Hz, 2H), 7.89 (d, *J* = 8.4 Hz, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 7.56 (t, *J* = 8.0 Hz, 1H), 7.47 (br s, 1H), 7.42-7.37 (m, 3H), 7.15 (t, *J* = 7.6 Hz, 1H), 7.06 (d, *J* = 8.0 Hz, 2H), 7.01 (d, *J* = 8.8 Hz, 2H), 6.44 (dd, *J* = 10.1, 16.9 Hz, 1H), 6.26 (dd, *J* = 1.6, 16.9 Hz, 1H), 5.76 (dd, *J* = 1.6, 10.1 Hz, 1H). MS (ESI) *m/e* [M+1]⁺ 476.

[0207] **Example 3: Synthesis of compounds 9-10**

[0208] Compound **9**: 7-(2-Aminophenyl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide




[0209] Step 1, 2, 3: 5-Amino-3-(4-(benzyloxy)phenyl)-1H-pyrazole-4-carbonitrile




[0210] The desired product was prepared from 4-(benzyloxy)benzoic acid according to the procedures similar to those (step 1 to 3) for 5-amino-3-(4-phenoxyphenyl)-1H- pyrazole-4-carbonitrile under appropriate conditions recognized by one of ordinary skill in the art. ¹H NMR (400 MHz, DMSO-d₆) δ 12.01 (s, 1H), 7.72 (d, *J* = 8.8 Hz, 2H), 7.50-7.44 (m, 2H), 7.44-7.37 (m, 2H), 7.36-7.32 (m, 1H), 7.11 (d, *J* = 7.2 Hz, 2H), 6.39 (br s, 2H) and 5.16 (s, 2H).

[0211] Step 4, 5: 2-(4-(Benzyloxy)phenyl)-7-(2-nitrophenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0212] The desired product was prepared from 1-(2-nitrophenyl)ethanone and 5-amino-3-(4-(benzyloxy)phenyl)-1H-pyrazole-4-carbonitrile according to the procedures similar to those (step 4 and step 5) for *N*-(3-(3-cyano-2-(4-phenoxyphenyl) pyrazolo[1,5-a]pyrimidin-7-yl)phenyl)acetamide under appropriate conditions recognized by one of ordinary skill in the art.

[0213] Step 6: 7-(2-Aminophenyl)-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carbonitrile



[0214] To a solution of 2-(4-(benzyloxy)phenyl)-7-(2-nitrophenyl)pyrazolo[1,5-a] pyrimidine-3-carbonitrile (2 g, 4.47 mmol) in CH₃OH (20 mL) and DCM (20 mL) was added 10% w/w Pd/C (300 mg). The mixture was stirred at RT under H₂ for 16 hr. Filtered and purified by chromatography column on silica gel (elution with DCM/CH₃OH) to afford 0.92 g (62%) of 7-(2-aminophenyl)-2-(4-hydroxyphenyl)- 4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile as a yellow solid. MS (ESI, m/e) [M+1]⁺ 331.9.


[0215] Step 7: 7-(2-Aminophenyl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carbonitrile



[0216] To a solution of 7-(2-aminophenyl)-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carbonitrile (331 mg, 1.0 mmol) in acetone (10 mL) was added (bromomethyl)benzene (204 mg, 1.2 mmol) and K₂CO₃ (276 mg, 2.0 mmol). The mixtre was stirred at RT for 16 hr. 50 mL of acetone was added and filtered. The filtrate was concentrated to afford 400 mg (95%) of 7-(2-aminophenyl)-2-(4- (benzyloxy)phenyl)-4,5,6,7-


tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile as a yellow solid. MS (ESI) m/e $[M+1]^+$ 421.9.

[0217] Step 8: 7-(2-Aminophenyl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide



[0218] The desired product was prepared from 7-(2-aminophenyl)-2-(4-(benzyloxy) phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound **2**. ^1H NMR (400 MHz, DMSO- d_6) δ 7.47-7.33 (m, 7H), 7.06 (d, J = 8.4 Hz, 2H), 6.96 (t, J = 7.6 Hz, 1H), 6.75 (br s, 1H), 6.68 (d, J = 7.6 Hz, 1H), 6.50 (t, J = 7.6 Hz, 1H), 6.28 (d, J = 7.6 Hz, 1H), 5.59-5.54 (m, 1H), 5.19 (br s, 2H), 5.12 (s, 2H), 3.30-3.20 (m, 1H), 3.02-2.92 (m, 1H), 2.25-2.14 (m, 1H) and 2.13-2.03 (m, 1H). MS (ESI) m/e $[M+1]^+$ 439.9.

[0219] Compound **10**: 7-(2-Acrylamidophenyl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0220] The desired product was prepared from 7-(2-aminophenyl)-2-(4-(benzyloxy)phenyl) -4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 9.84 (s, 1H), 7.46-7.37 (m, 7H), 7.33-7.28 (m, 2H), 7.21 (t, J = 7.6 Hz, 1H), 7.06 (d, J = 8.8 Hz, 2H), 6.81 (br s, 1H), 6.64 (d, J = 7.6 Hz, 1H), 6.53 (dd, J = 10.3, 16.8 Hz, 1H), 6.27 (d, J = 1.8, 16.8 Hz, 1H), 5.59-5.57 (m, 2H), 5.12 (s, 2H), 3.30-3.26 (m, 1H), 3.04-2.92 (m, 1H), 2.35-2.27 (m, 1H) and 1.95-1.97 (m, 1H). MS (ESI) m/e $[M+1]^+$ 493.9.

[0221] Compound **10** was separated into two enantiomeric stereoisomers compound **10a** (peak 1, R or S, retention time at 3.15 min in chiral analysis), and compound **10b** (peak 2, S or R, retention time at 3.91 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.


Column	CHIRALPAK IC-3
Column size	2 cm × 25 cm
Injection	3mL
Mobile phase	MeOH/ACN = 50/50
Flow rate	10 mL/min
Wave length	UV 220 nm
Temperature	35
Sample solution	2.5 mg/mL in mobile phase
Prep-SFC equipment	DAICEL-YMC

[0222] The chiral analysis conditions are shown below.


Column	CHIRALPAK IC
Column size	0.46 cm I.D.× 15 cm L, 5 um
Injection	3 uL
Mobile phase	MeOH/ACN=50/50 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm

[0223] Example 4: Synthesis of compounds 11-12

[0224] Compound 11: 7-(2-Aminophenyl)-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide




[0225] Step 1: Benzyl 2-(3-cyano-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenylcarbamate




[0226] To a solution of 7-(2-aminophenyl)-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carbonitrile (730 mg, 2.21 mmol) in THF (30 mL) was added K_2CO_3 (610 mg, 4.42 mmol), CbzCl (564 mg, 3.32 mmol). After stirring at 65 °C for 16 hr, the mixture was concentrated in vacuum. The residue was partitioned between 150 mL of DCM and 150 mL of brine. Organic layers were separated from aqueous layers, dried over Na_2SO_4 and purified by chromatography column on silica gel eluting with DCM/ CH_3OH to afford 370 mg (62%) of benzyl 2-(3-cyano-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl) phenylcarbamate as a white solid. 1H NMR (400 MHz, $DMSO-d_6$) δ 9.71 (s, 1H), 9.33 (s, 1H), 7.66 (s, 1H), 7.58 (d, J = 8.4 Hz, 2H), 7.44-7.28 (m, 7H), 7.17 (t, J = 7.6 Hz, 1H), 6.79 (d, J = 8.4 Hz, 2H), 6.59 (d, J = 7.6 Hz, 1H), 5.82-5.77 (m, 1H), 5.17 (s, 2H), 3.25-3.18 (m, 1H), 2.97-2.87 (m, 1H), 2.36-2.24 (m, 1H), 2.08-2.00 (m, 1H).

[0227] Step 2: Benzyl 2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidin-7-yl)phenylcarbamate




[0228] To a solution of benzyl 2-(3-cyano-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidin-7-yl)phenylcarbamate (370 mg, 0.8 mmol) in 20 mL of DCM was added 4-fluorophenylboronic acid (167 mg, 1.2 mmol), TEA (162 mg, 1.6 mmol) and $Cu(OAc)_2$ (216 mg, 1.2 mmol). After stirring at RT for 16 hr, 100 mL of DCM, 10 mL of CH_3OH and 100 mL of brine were added to the mixture. Organic layers were separated from aqueous layers, washed with brine (100 mL $\times$ 2), dried over Na_2SO_4 and purified by chromatography column on silica gel eluting with DCM/ CH_3OH to afford 334 mg (75%) of benzyl 2-(3-cyano-2-(4-(4-fluorophenoxy) phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidin-7-yl)phenylcarbamate as a white solid. 1H NMR (400 MHz, $DMSO-d_6$) δ 9.34 (s, 1H), 7.76-7.74 (m, 3H), 7.45-7.10 (m, 12H), 7.02 (d, J = 8.4 Hz, 2H), 6.59 (d, J = 8.0 Hz, 1H), 5.85-5.80 (m, 1H), 5.17 (s, 2H), 3.25-3.18 (m, 1H), 2.97-2.87 (m, 1H), 2.36-2.24 (m, 1H), 2.08-2.00 (m, 1H).

[0229] Step 3: Benzyl 2-(3-carbamoyl-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidin-7-yl)phenylcarbamate




[0230] The desired product was prepared from benzyl 2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenylcarbamate using the procedure similar to step 2 for compound **2**. MS (ESI) m/e $[M+1]^+$ 577.9.

[0231] Step 4: 7-(2-Aminophenyl)-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide



[0232] To a solution of benzyl 2-(3-carbamoyl-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenylcarbamate (100 mg, 0.17 mmol) in 5 mL of DCM and 5 mL of CH₃OH was added 10% w/w Pd/C (50 mg). After stirring at RT under H₂ for 16 hr, the mixture was filtered and the cake was washed with DCM/CH₃OH (1/1, 50 mL). The filtrate was concentrated and purified by chromatography column on silica gel eluting with DCM/CH₃OH to afford 10 mg (13%) of 7-(2-aminophenyl)-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide as a white solid. ¹H NMR (400 MHz, CD₃OD-*d*₄) δ 7.39 (d, J = 8.8 Hz, 2H), 7.02-6.91 (m, 7H), 6.66 (d, J = 7.6 Hz, 1H), 6.53 (t, J = 7.6 Hz, 1H), 6.33 (d, J = 7.6 Hz, 1H), 5.54-5.50 (m, 1H), 3.30-3.24 (m, 1H), 3.12-3.06 (m, 1H) and 2.31-2.20 (m, 2H). MS (ESI) m/e $[M+1]^+$ 443.9.

[0233] Compound **12**: 7-(2-Acrylamidophenyl)-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0234] The desired product was prepared from 7-(2-aminophenyl)-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide and acryloyl chloride


using the procedure similar to that for compound 8. ^1H NMR (400 MHz, $\text{CD}_3\text{OD}-d_4$) δ 7.47 (d, J = 8.4 Hz, 2H), 7.39 (t, J = 7.6 Hz, 1H), 7.33 (t, J = 7.6 Hz, 1H), 7.26 (t, J = 7.6 Hz, 1H), 7.12-7.00 (m, 6H), 6.81 (d, J = 8.0 Hz, 1H), 6.53-6.46 (m, 1H), 6.39-6.35 (m, 1H), 5.87-5.76 (m, 1H), 5.73-5.69 (m, 1H), 3.36-3.30 (m, 1H), 3.22-3.17 (m, 1H), 2.45-2.39 (m, 1H) and 2.17-2.14 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 497.9.

[0235] Example 5: Synthesis of compounds 13-14

[0236] Compound 13: 7-(2-(Methylamino)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0237] Step 1: *N*-(2-(3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide



[0238] To a solution of 7-(2-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (60 mg, 0.15 mmol) in 5 mL of DCM was added three drops of DIEA and three drops of trifluoroacetic anhydride. The reaction mixture was stirred at rt for 2 hr, then partitioned between water (20 mL) and DCM (20 mL). The organic layer was concentrated to give the product as a yellow solid (50 mg, yield: 67%), which was used in the next step without further purification.


[0239] Step 2: 7-(2-(Methylamino)phenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0240] To a solution of *N*-(2-(3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide (50 mg, 0.1 mmol) in 5 mL of acetone was added KOH (11.2 mg, 0.2 mmol) and CH_3I (0.5 mL). The reaction mixture was stirred at rt for 15 hr, then


concentrated to remove acetone. The residue was partitioned between 20 mL of water and 20 mL of EA. The organic layer was concentrated and purified by pre-TLC (PE/EA = 2/1) to give the product as a white solid (15 mg, yield: 37%). ¹H NMR (DMSO-d₆) δ 8.82 (d, *J* = 4.4 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 2H), 7.47-7.38 (m, 4H), 7.31 (dd, *J* = 1.6, 7.2 Hz, 1H), 7.23-7.17 (m, 3H), 7.11 (d, *J* = 8.8 Hz, 2H), 6.77-6.69 (m, 2H), 5.35-5.31 (m, 1H), 2.68 (d, *J* = 4.8 Hz, 3H). MS (ESI) *m/e* [M+1]⁺ 417.9.

[0241] Step 3: 7-(2-(Methylamino)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0242] To a solution of 7-(2-(methylamino)phenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (250 mg, 0.6 mmol) in 10 mL EtOH was added NaBH₄ (100 mg). The reaction mixture was stirred at rt for 15 hr, and then concentrated to remove EtOH. The residue was partitioned between 30 mL of water and 30 mL of EA. The organic layer was concentrated. The residue was dissolved in 10 mL of MeOH, followed by 10% w/w Pd/C (50 mg). The reaction mixture was stirred at rt under 1 atm. of H₂ for 15 hr. Then, the mixture was filtered. The filtrate was concentrated and purified by the flash chromatography (PE/EA = 1/1) to give the product as a white solid (144 mg, yield: 56.5%). MS (ESI) *m/e* [M+1]⁺ 421.9.

[0243] Step 4: 7-(2-(Methylamino)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0244] The desired product was prepared from 7-(2-(methylamino)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound 2. ¹H NMR (DMSO-d₆) δ 7.49 (d, *J* = 8.8 Hz, 2H), 7.44-7.35 (m, 2H), 7.18-7.01 (m, 6H), 6.76 (s, 1H), 6.58 (d, *J* = 7.6 Hz, 2H), 6.54 (t, *J* = 7.6 Hz, 1H), 6.28 (dd, *J* = 1.2, 7.6 Hz, 1H), 5.60-5.57 (m, 1H), 5.50-5.49 (m, 1H), 3.29-3.24 (m, 1H), 2.98-2.93 (m, 1H), 2.76 (d, *J* = 4.8 Hz, 3H), 2.24-2.18 (m, 1H), 2.06-2.03 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 439.8.


[0245] Compound 14: 7-(2-(*N*-methylacrylamido)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0246] The desired product was prepared from 7-(2-(methylamino)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (DMSO- d_6) δ 7.45-7.37 (m, 6H), 7.29-7.24 (m, 1H), 7.15 (t, J = 7.6 Hz, 1H), 7.06-7.00 (m, 5H), 6.81 (d, J = 7.6 Hz, 1H), 6.25-6.16 (m, 2H), 5.94-5.87 (m, 1H), 5.63-5.51 (m, 1H), 5.41-5.35 (m, 1H), 3.41-3.22 (m, 5H), 2.41-1.97 (m, 2H). MS (ESI) m/e $[M+1]^+$ 493.9.

[0247] **Example 6: Synthesis of compounds 15-16**

[0248] Compound **15**: 2-(4-(4-Fluorophenoxy)phenyl)-7-(2-(methylamino)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0249] Step 1: *N*-(2-(3-cyano-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a] pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide




[0250] To a solution of 7-(2-(aminophenyl)-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (33.1 mg, 0.1 mmol) in DCM (10 mL) was added trifluoroacetic anhydride (1 drop) and DIEA (1 drop). After stirring at RT for 1hr, the mixture was partitioned between 10 mL of DCM and 10 mL of brine. Organic layer was separated from aqueous layers, dried over Na_2SO_4 and concentrated to afford 40 mg (93%) of *N*-(2-(3-cyano-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a] pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide as a yellow solid. MS (ESI) m/e $[M+1]^+$ 427.8.

[0251] Step 2: *N*-(2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide




[0252] To a solution of *N*-(2-(3-cyano-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide (42.7 mg, 0.10 mmol) in DCM (3 mL) was added 4-fluorophenylboronic acid (17 mg, 0.12 mmol), TEA (21 mg, 0.2 mmol) and Cu(OAc)₂ (22 mg, 0.12 mmol). After stirring at RT for 16 hr, the mixture was purified by Pre-TLC (DCM/CH₃OH = 20/1) to afford 30 mg (57%) of *N*-(2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide as a colorless oil. ¹H NMR (400 MHz, DMSO-d₆) δ 11.33 (s, 1H), 7.90-7.77 (m, 3H), 7.52-7.37 (m, 3H), 7.34-7.27 (m, 2H), 7.22-7.14 (m, 2H), 7.10 (d, *J* = 8.8 Hz, 2H), 6.80 (d, *J* = 6.4 Hz, 1H), 5.80-5.75 (m, 1H), 3.37-3.28 (m, 1H), 3.09-2.95 (m, 1H), 2.50-2.37 (m, 1H), 2.10-1.95 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 521.8.

[0253] Step 3: *N*-(2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoro-*N*-methylacetamide




[0254] To a solution of *N*-(2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoroacetamide (187 mg, 0.36 mmol) in acetone (10 mL) was added K₂CO₃ (100 mg, 0.72 mmol) and CH₃I (15 drops). After stirring at RT for 2 hr, the mixture was filtered. The filtrate was concentrated to afford 192 mg (100%) of *N*-(2-(3-cyano-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoro-*N*-methylacetamide as yellow solid. MS (ESI) *m/e* [M+1]⁺ 535.8.

[0255] Step 4: 2-(4-(4-Fluorophenoxy)phenyl)-7-(2-(methylamino)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0256] The desired product was prepared from *N*-(2-(3-cyano-2-(4-(4-fluorophenoxy) phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)phenyl)-2,2,2-trifluoro-*N*-methylacetamide using the procedure similar to step 2 for compound 2. ¹H NMR (400 MHz, CD₃OD-*d*₄) δ 7.39 (d, *J* = 8.8 Hz, 2H), 7.09-6.90 (m, 7H), 6.58 (d, *J* = 8.0 Hz, 1H), 6.51 (t, *J* = 7.2 Hz, 1H), 6.30 (d, *J* = 7.2 Hz, 1H), 5.52-5.48 (m, 1H), 3.27-3.24 (m, 1H), 3.11-3.02 (m, 1H), 2.76 (s, 3H), 2.32-2.10 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 457.9.


[0257] Compound 16: 2-(4-(4-Fluorophenoxy)phenyl)-7-(2-(*N*-methylacrylamido)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0258] The desired product was prepared from 2-(4-(4-fluorophenoxy)phenyl)-7-(2-(methylamino)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide and acryloyl chloride using the procedure similar to that for compound 8. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.47-7.37 (m, 4H), 7.30-7.15 (m, 3H), 7.12-7.08 (m, 2H), 6.99 (d, *J* = 8.4 Hz, 2H), 6.81 (d, *J* = 7.2 Hz, 1H), 6.25-5.85 (m, 2H), 5.63-5.49 (m, 1H), 5.42-5.32 (m, 1H), 3.40-3.20 (m, 6H), 2.45-2.20 (m, 1H), 2.15-1.90 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 511.9.

[0259] Example 7: Synthesis of compounds 17-18


[0260] Compound 17: 7-(2-Aminophenyl)-2-(4-(cyclopentyloxy)phenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide



[0261] The desired product was prepared from 7-(2-aminophenyl)-2-(4-hydroxyphenyl) - 4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile and bromocyclopentane using the

procedures similar to those (step 7 and step 8) for compound **9**, under appropriate conditions recognized by one of ordinary skill in the art. ^1H NMR (400 MHz, DMSO- d_6) δ 7.37 (d, J = 8.8 Hz, 2H), 6.97-6.92 (m, 3H), 6.74 (br s, 1H), 6.67 (d, J = 7.6 Hz, 1H), 6.49 (t, J = 7.6 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.58-5.53 (m, 1H), 5.15 (s, 2H), 4.86-4.80 (m, 1H), 3.24-3.27 (m, 1H), 3.02-2.93 (m, 1H), 2.22-2.16 (m, 1H), 2.14-2.08 (m, 1H), 1.98-1.85 (m, 2H), 1.91-1.70 (m, 4H), 1.64-1.54 (m, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 418.0.


[0262] Compound 18: 7-(2-Acrylamidophenyl)-2-(4-(cyclopentyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0263] The desired product was prepared from compound **17** and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 9.83 (s, 1H), 7.45 (d, J = 7.6 Hz, 1H), 7.37 (d, J = 8.8 Hz, 2H), 7.29 (t, J = 7.6 Hz, 1H), 7.21 (t, J = 7.6 Hz, 1H), 6.93 (d, J = 8.8 Hz, 2H), 6.81 (s, 1H), 6.64 (d, J = 7.6 Hz, 1H), 6.53 (dd, J = 10.2, 17.0 Hz, 1H), 6.27 (d, J = 17.0 Hz, 1H), 5.80-5.77 (m, 2H), 4.86-4.79 (m, 1H), 3.27-3.23 (m, 1H), 3.03-2.94 (m, 1H), 2.36-2.25 (m, 1H), 1.99-1.91 (m, 3H) and 1.75-1.53 (m, 6H). MS (ESI) m/e $[\text{M}+1]^+$ 471.9.

[0264] Example 8: Synthesis of compounds 19-20

[0265] Compound 19: 7-(2-Aminophenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0266] Step 1: 7-(2-Aminophenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0267] To a solution of 7-(2-aminophenyl)-2-(4-(hydroxymethyl)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (331 mg, 1.0 mmol) in acetone (15 mL) was added (bromomethyl)cyclopropane (135 mg, 1.0 mmol) and K_2CO_3 (276 mg, 2.0 mmol). After stirring at 56 °C for 16 hr, the mixture was filtered. The cake was washed with acetone (20 mL $\times$ 2). The filtrate was concentrated to afford 300 mg of desired product (78%) as a yellow solid. MS (ESI) m/e $[M+1]^+$ 386.0.

[0268] Step 2: 7-(2-Aminophenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0269] To a solution of 7-(2-aminophenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (350 mg, 0.91 mmol) in EtOH (4 mL) and DMSO (4 mL) was added NaOH aqueous solution (5 N, 2 mL) and H_2O_2 (2 mL). After stirring at 60 °C for 3hr, the mixture was partitioned between 100 mL of H_2O and 100 mL of EA. The organic layer was separated from aqueous layers, washed with saturated brines (100 mL $\times$ 2), dried over Na_2SO_4 and concentrated. The residue was purified by chromatography column on silica gel (elution with DCM/MeOH) to afford 150 mg (41%) of desired product as a white solid. 1H NMR (400 MHz, $DMSO-d_6$) δ 7.37 (d, J = 8.8 Hz, 2H), 6.97-6.92 (m, 3H), 6.75 (br s, 1H), 6.67 (d, J = 8.0 Hz, 1H), 6.49 (t, J = 7.6 Hz, 1H), 6.27 (d, J = 7.6 Hz, 1H), 5.58-5.54 (m, 1H), 5.16 (s, 2H), 3.83 (d, J = 7.2 Hz, 2H), 3.27-3.24 (m, 1H), 3.01-2.92 (m, 1H), 2.23-2.13 (m, 1H), 2.13-2.07 (m, 1H), 1.24-1.18 (m, 1H), 0.59-0.54 (m, 2H) and 0.34-0.30 (m, 2H). MS (ESI) m/e $[M+1]^+$ 404.0.

[0270] Compound **20**: 7-(2-Acrylamidophenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0271] The desired product was prepared from compound **19** and acryloyl chloride according to the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO-d_6) δ 9.83 (s, 1H), 7.45 (d, $J = 8.0$ Hz, 1H), 7.37 (d, $J = 8.4$ Hz, 2H), 7.30 (t, $J = 7.6$ Hz, 1H), 7.21 (t, $J = 7.6$ Hz, 1H), 6.96 (d, $J = 8.4$ Hz, 2H), 6.81 (s, 1H), 6.64 (d, $J = 8.0$ Hz, 1H), 6.53 (dd, $J = 10.2, 16.7$ Hz, 1H), 6.27 (d, $J = 16.7$ Hz, 1H), 5.80-5.77 (m, 2H), 3.83 (d, $J = 7.2$ Hz, 2H), 3.27-3.23 (m, 1H), 3.03-2.93 (m, 1H), 2.36-2.25 (m, 1H), 2.02-1.91 (m, 1H), 1.23-1.14 (m, 1H), 0.59-0.53 (m, 2H) and 0.34-0.29 (m, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 457.9.

[0272] Compound **20** was separated into two enantiomeric stereoisomers compound **20a** (peak 1, R or S, retention time at 3.03 min in chiral analysis), and compound **20b** (peak 2, S or R, retention time at 3.82 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.


Column	CHIRALPAK IC-3
Column size	2 cm $\times$ 25 cm
Injection	3mL
Mobile phase	MeOH/ACN=50/50
Flow rate	10 mL/min
Wave length	UV 254 nm
Temperature	35 $^\circ\text{C}$
Sample solution	2.5 mg/mL in mobile phase
Prep-SFC equipment	DAICEL-YMC

[0273] The chiral analysis conditions are shown below.

Column	CHIRALPAK IC
Column size	0.46 cm I.D. $\times$ 15 cm L, 5 μm
Injection	3 μL
Mobile phase	MeOH/ACN = 50/50 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm


[0274] Example 9: Synthesis of compounds 21-22

[0275] Compound **21**: 7-(2-Aminophenyl)-2-(4-(2-methoxyethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0276] The desired product was prepared from 7-(2-aminophenyl)-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile and 1-bromo-2-methoxyethane using the procedures similar to those (step 7 and step 8) for compound **9**, under appropriate conditions recognized by one of ordinary skill in the art. ^1H NMR (400 MHz, DMSO- d_6) δ 7.38 (d, J = 8.6 Hz, 2H), 7.00-6.94 (m, 3H), 6.75 (br s, 1H), 6.69 (d, J = 7.6 Hz, 1H), 6.51 (t, J = 7.6 Hz, 1H), 6.28 (d, J = 7.6 Hz, 1H), 5.59-5.54 (m, 1H), 5.21 (br s, 1H), 4.11 (t, J = 4.0 Hz, 2H), 3.66 (t, J = 4.0 Hz, 2H), 3.30 (s, 3H), 3.28-3.25 (m, 1H), 3.02-2.92 (m, 1H), 2.24-2.16 (m, 1H) and 2.13-2.05 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 407.9.


[0277] Compound **22**: 7-(2-Acrylamidophenyl)-2-(4-(2-methoxyethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0278] The desired product was prepared from compound **21** and acryloyl chloride according to procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 9.84 (s, 1H), 7.45 (d, J = 8.0 Hz, 1H), 7.37 (d, J = 8.6, 2H), 7.30 (t, J = 7.6 Hz, 1H), 7.21 (t, J = 7.6 Hz, 1H), 6.93 (d, J = 8.6 Hz, 2H), 6.81 (s, 1H), 6.64 (d, J = 8.0 Hz, 1H), 6.53 (dd, J = 10.5, 17.0 Hz, 1H), 6.27 (dd, J = 1.7, 17.0 Hz, 1H), 5.80-5.77 (m, 2H), 4.11 (t, J = 4.4 Hz, 2H), 3.66 (t, J = 4.4 Hz, 2H), 3.30 (s, 3H), 3.27-3.23 (m, 1H), 3.03-2.94 (m, 1H), 2.36-2.25 (m, 1H), 2.01-1.95 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 462.0.

[0279] Example 10: Synthesis of compounds 23-24

[0280] Compound **23**: 7-(2-Aminophenyl)-2-(4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0281] Step 1: 7-(2-Aminophenyl)-2-(4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0282] To a solution of 7-(2-aminophenyl)-2-(4-hydroxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (33 mg, 0.1 mmol) in THF (5 mL) was added PPh₃ (78.6 mg, 0.25 mmol) and tetrahydro-2H-pyran-4-ol (10 mg, 0.1 mmol). Then, DIAD (51 mg, 0.25 mmol) was added dropwise to the mixture at 0 °C and stirred at 0 °C for 10 min under N₂. The mixture was allowed to warm to rt and stirred at RT for 16 hr. The mixture was concentrated in vacuum and partitioned between DCM (20 mL) and brine (20 mL). Organic layer was separated from aqueous layers, dried over sodium sulfate and purified by Pre-TLC (DCM/CH₃OH=10/1) to afford 5 mg (12%) of 7-(2-aminophenyl)-2-(4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile as a colorless oil. ¹H NMR (400 MHz, DMSO-d₆) δ 7.70 (d, *J* = 8.4 Hz, 2H), 7.66-7.58 (m, 2H), 7.58-7.52 (m, 1H), 7.03 (d, *J* = 8.4 Hz, 2H), 6.97 (t, *J* = 7.6 Hz, 1H), 6.69 (d, *J* = 8.0 Hz, 1H), 6.49 (t, *J* = 7.6 Hz, 1H), 6.23 (d, *J* = 8.0 Hz, 1H), 5.63 (s, 1H), 5.21 (s, 2H), 4.66-4.56 (m, 1H), 3.90-3.79 (m, 2H), 3.54-3.43 (m, 2H), 3.25-3.18 (m, 1H), 2.97-2.86 (m, 1H), 2.21-2.07 (m, 2H), 2.02-1.92 (m, 2H), 1.65-1.50 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 415.9.


[0283] Step 2: 7-(2-Aminophenyl)-2-(4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0284] The desired product was prepared from 7-(2-aminophenyl)-2-(4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound 2. ¹H NMR (400 MHz, DMSO-d₆) δ 7.38 (d, *J* = 8.8 Hz, 2H),

7.02 (d, $J = 8.8$ Hz, 2H), 6.95 (t, $J = 7.6$ Hz, 1H), 6.75 (s, 1H), 6.67 (d, $J = 7.8$ Hz, 1H), 6.49 (t, $J = 7.6$ Hz, 1H), 6.26 (d, $J = 7.8$ Hz, 1H), 5.56 (s, 1H), 5.16 (s, 2H), 4.64-4.54 (m, 1H), 3.91-3.79 (m, 2H), 3.53-3.42 (m, 2H), 3.29-3.19 (m, 1H), 3.01-2.92 (m, 1H), 2.25-2.05 (m, 2H), 2.02-1.91 (m, 2H) and 1.64-1.52 (m, 2H). MS (ESI) m/e $[M+1]^+$ 433.9.


[0285] Compound 24: 7-(2-Acrylamidophenyl)-2-(4-(tetrahydro-2H-pyran-4-yloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0286] The desired product was prepared from compound **23** and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 9.83 (s, 1H), 7.45 (d, $J = 8.0$ Hz, 1H), 7.38 (d, $J = 8.6$ Hz, 2H), 7.30 (t, $J = 7.6$ Hz, 1H), 7.21 (t, $J = 7.6$ Hz, 1H), 7.02 (d, $J = 8.6$ Hz, 2H), 6.81 (s, 1H), 6.63 (d, $J = 8.0$ Hz, 1H), 6.53 (dd, $J = 17.0, 10.3$ Hz, 1H), 6.27 (dd, $J = 17.0, 1.6$ Hz, 1H), 5.82-5.74 (m, 2H), 4.66-4.51 (m, 1H), 3.90-3.78 (m, 2H), 3.54-3.40 (m, 2H), 3.31-3.18 (m, 1H), 3.03-2.93 (m, 1H), 2.37-2.24 (m, 1H), 2.04-1.92 (m, 3H) and 1.64-1.51 (m, 2H). MS (ESI) m/e $[M+1]^+$ 487.9.


[0287] Example 11: Synthesis of compounds 25-27

[0288] Compound 25: 7-(1-(Tert-butoxycarbonyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0289] The desired product was prepared from tert-butyl 4-acetylpiperidine-1-carboxylate and 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile according to the procedures for 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (step 4 to 5) and compound **2** (step 1 and 2) under appropriate conditions recognized by one of ordinary skill in the art. ^1H NMR ($\text{CD}_3\text{OD}-d_4$) δ 7.40 (d, $J = 8.4$ Hz, 2H), 7.32-7.25 (m, 2H), 7.06 (t, $J = 7.6$ Hz, 1H), 7.01-6.94 (m, 4H), 4.10-4.00 (m, 2H), 3.98-3.91 (m, 1H), 3.35-3.30 (m, 2H), 2.70-2.58 (m, 2H), 2.18-2.02 (m, 2H), 2.02-1.84 (m, 1H), 1.65-1.45 (m, 2H), 1.39-1.12 (m, 2H), 1.35 (s, 9H). MS (ESI) m/e $[M+1]^+$ 518.0.

[0290] Compound 26: 7-(Piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate



[0291] The desired product was prepared from compound **25** according to the procedure similar to step 2 for compound **38**. ^1H NMR (DMSO- d_6) δ 8.47 (br s, 1H), 8.16 (br s, 1H), 7.50 (d, J = 8.4 Hz, 2H), 7.46-7.38 (m, 2H), 7.18 (t, J = 7.6 Hz, 1H), 7.10-7.04 (m, 4H), 6.72 (br s, 1H), 4.08-4.01 (m, 1H), 3.34-3.26 (m, 4H), 2.94-2.75 (m, 2H), 2.28-2.14 (m, 1H), 2.07-1.88 (m, 2H), 1.87-1.80 (m, 1H), 1.75-1.66 (m, 1H), 1.60-1.43 (m, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 418.0.

[0292] Compound **27**: 7-(1-Acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0293] The desired product was prepared from compound **26** and acryloyl chloride according to the procedure similar to that for compound **8**. ^1H NMR (DMSO- d_6) δ 7.50 (d, J = 8.8 Hz, 2H), 7.46-7.38 (m, 2H), 7.17 (t, J = 7.6 Hz, 1H), 7.08 (d, J = 7.6 Hz, 2H), 7.05 (d, J = 8.8 Hz, 2H), 6.83-6.76 (m, 1H), 6.68 (br s, 1H), 6.07 (d, J = 18.4 Hz, 1H), 5.64 (d, J = 10.4 Hz, 1H), 4.52-4.42 (m, 1H), 4.11-3.98 (m, 2H), 3.33-3.24 (m, 2H), 3.04-2.94 (m, 1H), 2.67-2.55 (m, 1H), 2.33-2.25 (m, 1H), 2.01-1.93 (m, 2H), 1.78-1.66 (m, 1H), 1.61-1.50 (m, 1H), 1.30-1.18 (m, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 471.9.

[0294] Compound **27** was separated into two enantiomeric stereoisomers compound **27a** (peak 1, R or S, retention time at 6.49 min in chiral analysis), and compound **27b** (peak 2, S or R, retention time at 8.03 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.

Column	CHIRALPAK IC
Column size	2 cm $\times$ 25 cm
Injection	2mL
Mobile phase	MeOH/ACN=60/40
Flow rate	15 mL/min
Wave length	UV 254 nm


Temperature	35 °C
Sample solution	0.5 mg/mL in mobile phase
Prep-SFC equipment	DAICEL-YMC

[0295] The chiral analysis conditions are shown below.


Column	CHIRALPAK IC
Column size	0.46 cm I.D.× 15 cm L, 5 μ m
Injection	2 μ L
Mobile phase	MeOH/ACN = 60/40 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm

[0296] Example 12: Synthesis of compounds 28-29

[0297] Compound **28**: 7-(Azetidin-3-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide




[0298] Step 1: Tert-butyl 3-(methoxy(methyl)carbamoyl)azetidine-1-carboxylate




[0299] To a solution of 1-(tert-butoxycarbonyl)azetidine-3-carboxylic acid (5.15 g, 25.6 mmol) in THF (100 mL) was added DCC (7.11 g, 34.5 mmol), Et₃N (5.18 g, 51.2 mmol) and *N,O*-dimethylhydroxylamine hydrochloride (3.44 g, 35.3 mmol), the reaction was stirred at RT for about 16 hr. Concentrated under reduced pressure to remove solvent, the residue was portioned between EA (100 mL) and water (50 mL), the aqueous was further extracted with EA (50 mL × 3). The combined organic phases were washed with brine (20 mL), concentrated under reduced pressure to remove solvent, then purified by column chromatography on silica gel (200-300 mesh, CH₂Cl₂/MeOH = 20/1) to give the crude product (~8.0 g) as a colorless oil. MS (ESI) *m/e* [M+23]⁺ 266.9, [M-55]⁺ 189.0.

[0300] Step 2: Tert-butyl 3-acetylazetidine-1-carboxylate




[0301] To a solution of tert-butyl 3-(methoxy(methyl)carbamoyl)azetidine-1-carboxylate (7.0 g, 28.7 mmol) in THF (150 mL) was added CH_3MgBr (43 mL, 43 mmol) at 0 °C, then slowly warmed to RT for about 2 hr. 10% aqueous of citric acid (30 mL) was added to the mixture, and extracted with EA (50 mL $\times$ 3), the combined organic phases were washed with brine (20 mL), dried over Na_2SO_4 , filtered, concentrated and purified by column chromatography on silica gel (200-300 mesh, PE/EA = 2/1), to give the crude product (4.0 g, 70%) as a colorless oil. MS (ESI) m/e $[\text{M}-55]^+$ 144.0.

[0302] Compound **28**: 7-(Azetidin-3-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide



[0303] The desired product was prepared from tert-butyl 3-acetylazetidine-1-carboxylate and 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile according to the procedures for 7-(3-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine -3-carbonitrile (step 4 to 5), compound **2** (step 1 and 2) and compound **38** (step 2), under appropriate conditions recognized by one of ordinary skill in the art. ^1H NMR ($\text{DMSO}-d_6$) δ 8.70 (br s, 1H), 8.44 (br s, 1H), 7.50 (d, J = 8.6 Hz, 2H), 7.45-7.40 (m, 2H), 7.18 (t, J = 7.6 Hz, 1H), 7.08 (d, J = 7.4 Hz, 2H), 7.06 (d, J = 8.6 Hz, 2H), 6.72 (br s, 1H), 4.46-4.38 (m, 1H), 4.24-4.15 (m, 1H), 4.07-3.94 (m, 3H), 3.29-3.24 (m, 2H), 3.19-3.10 (m, 1H), 2.13-2.04 (m, 1H), 1.78-1.69 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 390.0.

[0304] Compound **29**: 7-(1-Acryloylazetidin-3-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0305] The desired product was prepared from compound **28** and acryloyl chloride according to the procedure similar to that for compound **8**. ^1H NMR ($\text{DMSO}-d_6$) δ 7.50 (d, J = 8.4 Hz, 2H), 7.44-7.40 (m, 2H), 7.20-7.15 (m, 1H), 7.10-7.04 (m, 4H), 6.69 (br s, 1H), 6.37-6.26 (m, 1H), 6.12-6.04 (m, 1H), 5.68-5.60 (m, 1H), 4.43-4.25 (m, 2H), 4.18-4.08 (m, 1H), 4.04-3.96 (m, 1H),

3.86-3.80 (m, 1H), 3.32-3.26 (m, 2H), 3.02-2.92 (m, 1H), 2.14-2.06 (m, 1H), 1.79-1.70 (m, 1H).

MS (ESI) m/e [M+1]⁺ 444.0.

[0306] Compound **29** was separated into two enantiomeric stereoisomers compound **29a** (peak 1, R or S, retention time at 10.54 min in chiral analysis), and compound **29b** (peak 2, S or R, retention time at 13.98 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.


Column	CHIRALCEL OJ-H
Column size	2 cm × 25 cm
Injection	3ml
Mobile phase	CO ₂ /(MeOH70%ACN30%) = 75/25
Flow rate	45 ml/min
Wave length	UV 220 nm
Temperature	35
Sample solution	3 mg/ml in mobile phase
Prep-SFC equipment	DAICEL-SFC

[0307] The chiral analysis conditions are shown below.

Column	CHIRALPAK AD
Column size	0.46 cm I.D.× 15 cm L, 5 μm
Injection	3 μL
Mobile phase	n-Hexane/EtOH (0.1% triethyl amine) = 70/30 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm

[0308] Example 13: Synthesis of compounds 30-31

[0309] Compound **30**: Cis-7-Acryloyl-2-(4-phenoxyphenyl)-4,5,5a,6,7,8,9,9a-octahydro pyrazolo[1,5-a]pyrido[3,4-e]pyrimidine-3-carboxamide and Cis-7-acryloyl-2-(4-phenoxyphenyl)-4,4a,5,6,7,8,8a,9-octahidropyrzazolo[1,5-a]pyrido[4,3-d]pyrimidine-3-carboxamide



[0310] The desired product was prepared from tert-butyl 4-oxopiperidine-1-carboxylate and 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile according to the procedures for compound **27**, under appropriate conditions recognized by one of ordinary skill in the art. ¹H NMR (DMSO-d₆) δ 7.51 (d, *J* = 8.4 Hz, 2H), 7.46-7.38 (m, 2H), 7.17 (t, *J* = 7.6 Hz, 1H), 7.11-7.01 (m, 4H), 6.90-6.78 (m, 1H), 6.65 (s, 1H), 6.18-6.06 (m, 1H), 5.73-5.63 (m, 1H), 4.45-4.33 (m, 1H), 3.84-3.34 (m, 5H), 3.22-3.16 (m, 1H), 2.40-2.32 (m, 1H), 2.21-2.10 (m, 1H), 2.00-1.90 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 443.9.

[0311] Compound **30** was separated into two enantiomeric stereoisomers compound **30a** (peak 1, R or S, retention time at 10.64 min in chiral analysis), and compound **30b** (peak 2, S or R, retention time at 15.18 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.

Column	CHIRALCEL ID
Column size	3 cm × 25 cm
Injection	7 mL
Mobile phase	MeOH/EA = 30/70
Flow rate	25 ml/min
Wave length	UV 254 nm
Temperature	35 °C
Sample solution	2 mg/ml in mobile phase
Prep-SFC equipment	DAICEL-YMC


[0312] The chiral analysis conditions are shown below.

Column	CHIRALPAK AD-H
Column size	0.46 cm I.D. × 15 cm L, 5 μm
Injection	3 μL
Mobile phase	n-Hexane/EtOH (0.1% triethyl amine) = 50/50 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm


[0313] The compound **31** was obtained as one byproduct in the preparation of compound **30**. ^1H NMR (DMSO- d_6 at 80 °C) δ 7.53 (d, J = 8.4 Hz, 2H), 7.46-7.38 (m, 2H), 7.17 (t, J = 7.6 Hz, 1H), 7.11-7.02 (m, 4H), 6.82 (dd, J = 16.8, 10.6 Hz, 1H), 6.42 (br s, 1H), 6.10 (dd, J = 16.8, 2.3 Hz, 1H), 6.01 (br s, 2H), 5.68 (dd, J = 10.6, 2.3 Hz, 1H), 4.17 (dd, J = 5.4, 12.2 Hz, 1H), 3.67 (t, J = 12.2 Hz, 1H), 3.28 (td, J = 4.0, 10.4 Hz, 1H), 2.24-2.17 (m, 1H), 1.93-1.81 (m, 1H), 1.47-1.33 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 443.9.

[0314] Example 14: Synthesis of compounds 32-33

[0315] Compound 32: 7-(Aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide




[0316] Step 1: (*E*)-2-(4-(Dimethylamino)-2-oxobut-3-enyl)isoindoline-1,3-dione



[0317] To a solution of 2-(2-oxopropyl)isoindoline-1,3-dione (1 g, 4.9 mmol) in 20 mL of DMF-DMA was added some of 4A molecular sieve. The reaction mixture was stirred at 100 °C under N_2 for 15 hr. After cooling down to RT, the mixture was filtered and collected 600 mg (47.5%) of crude (*E*)-2-(4-(dimethylamino)-2-oxobut-3-enyl) isoindoline-1,3-dione as a solid. MS (ESI) m/e $[\text{M}+1]^+$ 259.1.


[0318] Step 2: 7-((1,3-Dioxoisindolin-2-yl)methyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0319] A mixture of (*E*)-2-(4-(dimethylamino)-2-oxobut-3-enyl)isoindoline-1,3-dione (600 mg, 2.33 mmol) and 5-amino-3-(4-phenoxy phenyl)-1H-pyrazole-4-carbonitrile (642 mg, 2.33 mmol) in 20 mL of HOAc was stirred and heated to 120 °C for 15 hr. The mixture was concentrated and suspended in 30 mL of solvent (PE/EA = 4/1). The mixture was filtered and the solid was purified by pre-TLC (DCM/EA = 50/1) to give 430 mg (40%) of 7-((1,3-dioxoisindolin-2-yl)methyl)-2-(4-phenoxyphenyl)pyrazolo [1,5-a]pyrimidine-3-carbonitrile as


a solid. ^1H NMR (400 MHz, DMSO- d_6) δ 8.74 (d, J = 4.4 Hz, 1H), 8.05 (d, J = 8.8 Hz, 2H), 7.96-7.90 (m, 1H), 7.88-7.85 (m, 2H), 7.50 (d, J = 4.4 Hz, 1H), 7.46-7.37 (m, 2H), 7.21-7.09 (m, 5H), 5.32 (s, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 472.1.

[0320] Step 3: 7-(Aminomethyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3- carbonitrile




[0321] To a solution of 7-((1,3-dioxoisindolin-2-yl)methyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (290 mg, 0.62 mmol) in 5 mL of MeOH and 5 mL of dioxane was added 12 drops of hydrazine hydrate. The reaction mixture was stirred and heated to 60 °C for 3 hr. The mixture was concentrated and suspended in 20 mL of solvent (DCM/MeOH = 10/1). The mixture was filtered and the filtrate was concentrated and purified by the flash chromatography eluting with EA followed by DCM/MeOH (10/1) to give 150 mg (71%) of 7-(aminomethyl)-2-(4-phenoxyphenyl) pyrazolo[1,5-a]pyrimidine-3-carbonitrile as a solid. MS (ESI) m/e $[\text{M}+1]^+$ 342.1.

[0322] Step 4: 7-(Aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0323] To a solution of 7-(aminomethyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine- 3-carbonitrile (150 mg, 0.44 mmol) in 20 mL of EtOH was added NaBH_4 (200 mg). The reaction mixture was stirred at rt for 15 hr. The mixture was concentrated to remove the solvent. The residue was partitioned between EA (20 mL) and water (20 mL). The organic layers were concentrated to give 150 mg (100%) of crude 7-(aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile as a solid, which was used in the next step without further purification. MS (ESI) m/e $[\text{M}+1]^+$ 346.0.

[0324] Step 5: 7-(Aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine -3-carboxamide



[0325] To a solution of 7-(aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carbonitrile (15 mg, 0.043 mmol) in 2 mL of EtOH was added 1 mL of DMSO, 0.5 mL of NaOH (5N) and 0.5 mL of H₂O₂ (30% aqueous solution). After stirring at 60 °C for 2 hr, the mixture was concentrated to remove EtOH. The residue was partitioned between water (30 mL) and EA (20 mL). The organic phase was concentrated and purified by pre-HPLC eluting from 20% to 40% CH₃CN in 0.1% TFA in H₂O. Fractions containing the desired product were combined and lyophilized overnight to give the product as a TFA salt (10 mg, 64%). ¹H NMR (400 MHz, DMSO-d₆) δ 8.07 (br s, 3H), 7.62 (d, *J* = 8.8 Hz, 2H), 7.43-7.35 (m, 2H), 7.25 (t, *J* = 7.6 Hz, 1H), 7.15-7.12 (m, 4H), 6.77 (br s, 1H), 4.38-4.30 (m, 1H), 3.40-3.20 (m, 4H), 2.16-2.06 (m, 1H), 2.00-1.80 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 364.0.

[0326] Compound **33**: 7-(Acrylamidomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0327] To a solution of 7-(aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide (125 mg, 0.344 mmol) in 5 mL of DCM was added Et₃N (four drops) and acryloyl chloride (46.5 mg, 0.52 mmol). After stirring at rt for 30 mins. The mixture was partitioned between water (10 mL) and DCM (5 mL). The organic layer was concentrated and purified by pre-TLC (DCM/MeOH = 20/1) to give 40 mg (28%) of product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.37 (t, *J* = 6.0 Hz, 1H), 7.59 (d, *J* = 8.8 Hz, 2H), 7.54-7.43 (m, 2H), 7.25 (t, *J* = 7.6 Hz, 1H), 7.16-7.11 (m, 4H), 6.76 (s, 1H), 6.33 (dd, *J* = 10.1, 17.0 Hz, 1H), 6.18 (dd, *J* = 1.9, 17.0 Hz, 1H), 5.69 (dd, *J* = 1.9, 10.1 Hz, 1H), 4.28-4.22 (m, 1H), 3.92-3.80 (m, 1H), 3.50-3.30 (m, 3H), 2.14-1.94 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 417.9.

[0328] Compound **33** was separated into two enantiomeric stereoisomers compound **33a** (peak 1, R or S, retention time at 6.04 min in chiral analysis), and compound **33b** (peak 2, S or R,

retention time at 8.87 min in chiral analysis) by chiral prep-HPLC. The chiral separation conditions are shown below.


Column	CHIRALCEL AD-H
Column size	3cm × 25 cm
Injection	5 mL
Mobile phase	CO ₂ /MeOH = 75/25
Flow rate	65 mL/min
Wave length	UV 220 nm
Temperature	35
Sample solution	4.5 mg/mL in mobile phase
Prep-SFC equipment	DAICEL-SFC

[0329] The chiral analysis conditions are shown below.


Column	CHIRALPAK AD-H
Column size	0.46 cm I.D. × 15 cm L, 5 μm
Injection	1 μL
Mobile phase	n-Hexane/EtOH (0.1% triethyl amine) = 70/30 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm

[0330] Example 15: Synthesis of compounds 34-35

[0331] Compound **34**: 7-((Methylamino)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide




[0332] Step 1: *N*-((3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)methyl)-2,2,2-trifluoroacetamide




[0333] To a solution of 7-(aminomethyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (50 mg, 0.147 mmol) in 5 mL of DCM was added three drops of DIEA and two drops of trifluoroacetic anhydride. After stirring at rt for 2 hr, 10 mL water was added to the mixture and extracted with DCM (5 mL $\times$ 2). The DCM layers were concentrated to give a yellow solid (50 mg, 78%), which was used in the next step without further purification.

[0334] Step 2: *N*-((3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)methyl)-2,2,2-trifluoro-*N*-methylacetamide




[0335] To a solution of *N*-((3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)methyl)-2,2,2-trifluoro-*N*-methylacetamide (40 mg, 0.091 mmol) in 5 mL of acetone was added K_2CO_3 (50 mg) and CH_3I (0.5 mL). After stirring at rt for 4 hr, the mixture was concentrated. The residue was partitioned between 10 mL of water and 10 mL of DCM. The organic layer was concentrated and purified by Pre-TLC (DCM/MeOH = 20/1) to give a yellow solid (30 mg, 73%). MS (ESI) m/e $[M+1]^+$ 451.9.

[0336] Step 3: 7-((Methylamino)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carbonitrile




[0337] To a solution of *N*-((3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidin-7-yl)methyl)-2,2,2-trifluoro-*N*-methylacetamide (40 mg, 0.089 mmol) in 15 mL of EtOH was added $NaBH_4$ (50 mg). After stirring at rt for 30 mins, the mixture was concentrated. The residue was partitioned between 20 mL of water and 20 mL of EA. The EA layer was concentrated and purified by Pre-TLC (DCM/MeOH = 5/1) to give a white solid (20 mg, 63%). MS (ESI) m/e $[M+1]^+$ 359.9.

[0338] Step 4: 7-((Methylamino)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide



[0339] To a solution of 7-((methylamino)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carbonitrile (20 mg, 0.0557 mmol) in 2 mL of EtOH was added 1 mL of DMSO, 0.5 mL of NaOH (5N) and 0.5 mL of H₂O₂ (30% solution). The reaction mixture was stirred at 60 °C for 1 hr. Then the mixture was concentrated to remove EtOH. The residue was partitioned between water (20 mL) and EA (20 mL). The EA layer was concentrated to give the product as a white solid (10 mg, yield: 48%). ¹H NMR (400 MHz, DMSO-d₆) δ 7.51 (d, *J* = 8 Hz, 2H), 7.47-7.40 (m, 2H), 7.19 (t, *J* = 7.6 Hz, 1H), 7.11-7.04 (m, 4H), 6.61 (s, 1H), 4.26-4.16 (m, 1H), 3.34-3.27 (m, 2H), 2.97-2.95 (m, 1H), 2.83-2.77 (m, 1H), 2.33 (s, 3H), 2.08-2.02 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 378.0.


[0340] Compound **35**: 7-((*N*-Methyl(acrylamido)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide




[0341] To a solution of 7-((methylamino)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide (92 mg, 0.244 mmol) in 5 mL of DCM was added Et₃N (three drops) and acryloyl chloride (33 mg, 0.366 mmol). The reaction mixture was stirred at rt for 30 min. Then the mixture was partitioned between water (30 mL) and DCM (20 mL). The organic layer was concentrated and purified by pre-TLC (DCM/MeOH = 15/1) to give the product as a white solid (25 mg, yield: 24%). ¹H NMR (400 MHz, DMSO-d₆ and D₂O at 80 °C) δ 7.51 (d, *J* = 8.4 Hz, 2H), 7.44-7.39 (m, 2H), 7.20-7.14 (m, 1H), 7.10-7.04 (m, 4H), 6.75-6.57 (m, 2H), 6.10-5.85 (m, 1H), 5.67-5.50 (m, 1H), 4.45-4.38 (m, 1H), 4.00-3.70 (m, 2H), 3.40-3.30 (m, 2H), 3.00 (s, 3H), 2.14-1.90 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 432.0.

[0342] Example 16: Synthesis of compounds 36-37

[0343] Compound **36**: 7-(Aminomethyl)-2-(1-benzyl-1H-pyrazol-4-yl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate




[0344] Step 1: Ethyl 1-benzyl-1H-pyrazole-4-carboxylate




[0345] To a mixture of ethyl 1H-pyrazole-4-carboxylate (35.0 g, 250 mmol) and K_2CO_3 (69.0 g, 500 mmol) in CH_3CN (250 mL) was added BnBr (42.7 g, 250 mmol). The mixture was stirred at RT for 18 h and concentrated. The residue was suspended in EA (500 mL), washed with water (200 mL $\times$ 2), dried over Na_2SO_4 and concentrated to give the desired compound as a white solid (53.0 g, 91.2%). 1H NMR ($DMSO-d_6$) δ 8.46 (s, 1H), 7.88 (s, 1H), 7.41-7.19 (m, 5H), 5.37 (s, 2H), 4.21 (q, 2H, $J = 5.4$ Hz), 1.25 (t, 3H, $J = 5.4$ Hz). MS (ESI) m/e $[M+1]^+$ 231.0.

[0346] Step 2: 1-Benzyl-1H-pyrazole-4-carboxylic acid



[0347] A mixture of ethyl 1-benzyl-1H-pyrazole-4-carboxylate (53.0 g, 0.23 mol) and LiOH (19.4 g, 0.46 mol) in THF (100 mL) and H_2O (100 mL) was stirred at reflux for 6 h. Then, THF was removed, the residue was acidified by 6 N HCl, precipitation was formed, filtered and dried to give the desired compound as a white solid (44.0 g, 92.8%). 1H NMR ($DMSO-d_6$) δ 12.36 (s, 1H), 8.39 (s, 1H), 7.85 (s, 1H), 7.38-7.27 (m, 5H), 5.37 (s, 2H). MS (ESI) m/e $[M+1]^+$ 202.9.


[0348] Step 3: 2-((1-Benzyl-1H-pyrazol-4-yl)(hydroxy)methylene)malononitrile



[0349] A solution of 1-benzyl-1H-pyrazole-4-carboxylic acid (25.0 g, 123.8 mmol) in $SOCl_2$ (250 mL) was heated to reflux for 3 hr. The mixture was concentrated in vacuum to give the intermediate, which was used in the next step without further purification. To a solution of


propanedinitrile (8.2 g, 12.8 mmol), DIEA (32.0 g, 247.6 mmol) in THF (250 mL) was added dropwise a solution of intermediate in toluene (250 mL) at 0~5 °C over 1 hr. The resultant mixture was allowed to warm to RT and stirred for 16 hr. The reaction was quenched with water (500 mL) and extracted with EA (500 mL × 3), The organic layers were washed with 3 N HCl (500 mL) , brine (500 mL×3), dried over Na₂SO₄ and concentrated to give the crude product (26.5 g, 85.0%) as a yellow solid. MS (ESI) m/e [M+1]⁺ 250.9.

[0350] Step 4: 2-((1-Benzyl-1H-pyrazol-4-yl)(methoxy)methylene)malononitrile




[0351] A solution of 2-((1-benzyl-1H-pyrazol-4-yl)(hydroxy)methylene)malononitrile (26.5g, 106 mmol) in CH(OMe)₃ (250 mL) was heated to 75 °C for 16 hr. Then the solution was concentrated. The residue was washed with MeOH (50 mL) to give 14.5 g (51.8%) of 2-((1-benzyl-1H-pyrazol-4-yl)(methoxy)methylene)malononitrile as an off-white solid. ¹H NMR (DMSO-d₆) δ 8.71 (s, 1H), 8.08 (s, 1H), 7.42-7.24 (m, 5H), 5.46 (s, 2H), 4.12 (s, 3H). MS (ESI) m/e [M+1]⁺ 264.9.

[0352] Step 5: 5-Amino-2'-benzyl-3,4'-bi(1H-pyrazole)-4-carbonitrile




[0353] A mixture of 2-((1-benzyl-1H-pyrazol-4-yl)(methoxy)methylene)malononitrile (14.5 g, 54.9 mmol) and hydrazine hydrate (10 mL) in EtOH (500 mL) was stirred at RT for 4 hr. Then the mixture was concentrated to give the crude product, washed with MeOH to afford 10 g (69.0%) of 5-amino-2'-benzyl-3,4'-bi(1H-pyrazole)-4- carbonitrile as an off-white solid. ¹H NMR (DMSO-d₆) δ 11.76 (br s, 1H), 8.18 (s, 1H), 7.82 (s, 1H), 7.34-7.26 (m, 5H), 6.11 (br s, 2H), 5.40 (s, 2H). MS (ESI) m/e [M+1]⁺ 264.9.

[0354] Compound **36**: 7-(Aminomethyl)-2-(1-benzyl-1H-pyrazol-4-yl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate



[0355] The desired product was prepared from 5-amino-2'-benzyl-3,4'-bi(1H-pyrazole)-4-carbonitrile and (*E*)-2-(4-(dimethylamino)-2-oxobut-3-enyl)isoindoline-1,3-dione according to the procedures (step 2 to 5) for compound **32**, under appropriate conditions recognized by one of ordinary skill in the art. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.14 (s, 1H), 7.97 (br s, 3H), 7.74 (s, 1H), 7.39-7.24 (m, 5H), 5.37 (s, 2H), 4.40-4.25 (m, 1H), 3.37-3.16 (m, 4H), 2.16-2.08 (m, 1H), 1.99-1.89 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 352.0.


[0356] Compound **37**: 7-(Acrylamidomethyl)-2-(1-benzyl-1H-pyrazol-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-*a*]pyrimidine-3-carboxamide




[0357] The desired product was prepared from compound **36** and acryloyl chloride using the procedure similar to that for compound **8**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.27 (t, *J* = 6.1 Hz, 1H), 8.11 (s, 1H), 7.67 (s, 1H), 7.41-7.21 (m, 5H), 6.58 (s, 1H), 6.25 (br s, 2H), 6.25 (dd, *J* = 17.1, 10.1 Hz, 1H), 6.10 (dd, *J* = 17.1, 2.1 Hz, 1H), 5.62 (dd, *J* = 10.1, 2.1 Hz, 1H), 5.36 (s, 2H), 4.16-4.10 (m, 1H), 3.83-3.72 (m, 1H), 3.42-3.30 (m, 1H), 3.30-3.22 (m, 2H), 2.00-1.96 (m, 1H), 1.95-1.86 (m, 1H). MS (ESI) *m/e* [M+1]⁺ 405.9.

[0358] Example 17: Synthesis of compounds 38-40

[0359] Compound **38** 1'-Benzyl-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-*b*]pyrazole-3,4'-piperidine]-7-carboxamide




[0360] Step 1: di-*tert*-Butyl 1-(1-benzyl-4-(ethoxycarbonyl)piperidin-4-yl)hydrazine-1,2-dicarboxylate




[0361] To a solution of diisopropylamine (153 mg, 1.5 mmol) in THF (20 mL) was added n-BuLi (2.5 M, 0.6 mL) at -75 °C under N₂. After 5 min, ethyl 1-benzylpiperidine-4-carboxylate (247 mg, 1.0 mmol) was added and the resulting mixture then stirred at -70 °C for 10 min, before adding di-tert-butyl azodicarboxylate (345 mg, 1.5 mmol). The reaction was stirred for 30 min, then quenched with aqueous NH₄Cl (10 mL) and extracted with EA (10 mL × 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give the crude product (350 mg, 72%) as an off-white solid. MS (ESI, m/e) [M+1]⁺ 478.3.

[0362] Step 2: Ethyl 1-benzyl-4-hydrazinylpiperidine-4-carboxylate hydrochloride




[0363] A mixture of di-tert-butyl 1-(1-benzyl-4-(ethoxycarbonyl)piperidin-4-yl)hydrazine-1,2-dicarboxylate (1.0 g, 2.09 mmol) and con. HCl (1.0 mL) in MeOH (10 mL) was heated to reflux for 2 hr. The mixture was then concentrated to give the crude product (650 mg, 88.9%) as a yellow solid. MS (ESI, m/e) [M+1]⁺ 278.0.

[0364] Step 3: Ethyl 4-(5-amino-4-cyano-3-(4-phenoxyphenyl)-1H-pyrazol-1-yl)-1-benzylpiperidine-4-carboxylate




[0365] A mixture of ethyl 1-benzyl-4-hydrazinylpiperidine-4-carboxylate hydrochloride (580 mg, 1.57 mmol), 2-(methoxy(4-phenoxyphenyl)methylene)malononitrile (433 mg, 1.57 mmol) and TEA (475 mg, 4.71 mmol) in CHCl₃ (20 mL) was heated to reflux for 16 hr under N₂. The mixture was concentrated and purified by chromatography column on silica gel using 50% of EA in PE as eluant to give the product (280 mg 34.2%) as a yellow solid. MS (ESI, m/e) [M+1]⁺ 521.9.

[0366] Step 4: 5-Amino-1-(1-benzyl-4-(hydroxymethyl)piperidin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile




[0367] To a solution of ethyl 4-(5-amino-4-cyano-3-(4-phenoxyphenyl)-1H-pyrazol-1-yl)-1-benzylpiperidine-4-carboxylate (52 mg, 0.1 mmol) in MeOH (5 mL) was added NaBH₄ (8 mg, 0.2 mmol). After 10 min, the reaction was quenched with water (5 mL) and extracted with EA (10 mL × 3). The organic combined layers were dried over Na₂SO₄ and concentrated to give the crude product (34 mg, 70.9%) as an off-white solid. MS (ESI, m/e) [M+1]⁺ 480.0.

[0368] Step 5: (4-(5-Amino-4-cyano-3-(4-phenoxyphenyl)-1H-pyrazol-1-yl)-1-benzylpiperidin-4-yl) methyl methanesulfonate



[0369] To a solution of 5-amino-1-(1-benzyl-4-(hydroxymethyl)piperidin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (50 mg, 0.1 mmol) and TEA (20 mg, 0.20 mmol) in DCM (5 mL) was added MsCl (14 mg, 0.12 mmol) at 0 °C. After 5 min, the reaction was quenched with water (5 mL) and extracted with DCM (5 mL × 3). The combined organic layers were dried over Na₂SO₄ and concentrated and purified by Prep-TLC (10% of MeOH in DCM) to give the product (35 mg, 62.8%) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 7.80 (d, *J* = 8.8 Hz, 2H), 7.45-7.36 (m, 2H), 7.37-7.22 (m, 5H), 7.17 (t, *J* = 7.6 Hz, 1H), 7.11 (d, *J* = 8.8 Hz, 2H), 7.06 (d, *J* = 8.0 Hz, 2H), 6.49 (br s, 2H), 4.48 (s, 2H), 3.40 (s, 2H), 3.10 (s, 3H), 2.88-2.55 (m, 4H), 2.19-2.16 (m, 2H), 1.92-1.86 (m, 2H). MS (ESI, m/e) [M+1]⁺ 557.9.


[0370] Step 6: 1'-Benzyl-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carbonitrile



[0371] A mixture of (4-(5-amino-4-cyano-3-(4-phenoxyphenyl)-1H-pyrazol-1-yl)-1-benzylpiperidin-4-yl)methyl methanesulfonate (35 mg, 0.06 mmol) and Cs₂CO₃ (31 mg, 0.09 mmol) in DMF (2 mL) was heated to 50 °C for 16 hr. The reaction was quenched with water (5


mL) and extracted with EA (5 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄, concentrated and purified by Prep-TLC (10% of MeOH in DCM) to give the product (12 mg, 43.2%) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 7.79 (d, J = 8.4 Hz, 2H), 7.43-7.40 (m, 3H), 7.38-7.29 (m, 4H), 7.26 (br s, 1H), 7.19 (t, J = 7.6 Hz, 1H), 7.12-7.05 (m, 4H), 3.85 (s, 2H), 3.53 (s, 2H), 2.89-2.80 (m, 2H), 2.22-2.12 (m, 2H), 2.10-1.96 (m, 2H), 1.85-1.75 (m, 2H). MS (ESI, m/e) [M+1]⁺ 462.0.

[0372] Step 7: 1'-Benzyl-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide




[0373] The desired product was prepared from 1'-benzyl-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carbonitrile according to the procedure similar to step 2 for compound **2**. ¹H NMR (400 MHz, DMSO-d₆) δ 7.63 (d, J = 8.4 Hz, 2H), 7.45-7.39 (m, 3H), 7.37-7.31 (m, 4H), 7.16 (t, J = 7.6 Hz, 1H), 7.12-6.94 (m, 4H), 6.43 (s, 1H), 3.78 (s, 2H), 3.55 (br s, 2H), 2.89-2.82 (m, 2H), 2.15-2.11 (m, 2H), 1.84-1.72 (m, 2H). MS (ESI, m/e) [M+1]⁺ 479.9.

[0374] Compound **39**: 6-(4-Phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide



[0375] A mixture of 1'-benzyl-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide (50 mg, 0.10 mmol), 10% w/w Pd(OH)₂/C (5 mg) in MeOH (10 mL) and HOAc (1 drop) was stirred for 16 hr under H₂. The mixture was filtrated and the filtrate was concentrated to give the crude product (20 mg 51.4%) as a yellow solid. MS (ESI, m/e) [M+1]⁺ 390.0.


[0376] Compound **40**: 1'-Acryloyl-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide




[0377] Compound **40** was prepared from compound **39** and acryloyl chloride according to the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 7.64 (d, J = 8.8 Hz, 2H), 7.44-7.37 (m, 2H), 7.16 (t, J = 7.6 Hz, 1H), 7.05 (d, J = 7.6 Hz, 2H), 6.97 (d, J = 8.8 Hz, 2H), 6.86 (dd, J = 10.5, 16.7 Hz, 1H), 6.51 (br s, 1H), 6.12 (dd, J = 2.3, 16.7 Hz, 1H), 5.69 (dd, J = 2.3, 10.5 Hz, 1H), 4.13-3.95 (m, 2H), 3.83 (s, 2H), 3.60-3.38 (m, 2H), 1.99-1.76 (m, 4H). MS (ESI, m/e) $[\text{M}+1]^+$ 443.9.

[0378] **Example 18: Synthesis of compounds 41-43**

[0379] Compound **41**: 1-Benzyl-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine- 4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide




[0380] Step 1: Ethyl 2-(1-benzylpiperidin-4-ylidene)acetate




[0381] To a suspension of NaH (318 mg, 7.94 mmol) in THF (20 mL) was added a solution of ethyl 2-(diethoxyphosphoryl)acetate (1.78 g, 7.94 mmol) in THF (5 mL) dropwise over 30 min at 0 °C. After stirring for 10 min, a solution of 1-benzylpiperidin-4-one (1.0 g, 5.29 mmol) in THF (5 mL) was added dropwise at 0 °C over 20 min. The mixture was allowed to stir for 60 min. Then, the reaction was quenched with water (10 mL). The mixture was extracted with EA (10 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , concentrated and purified by chromatography column on silica gel using 25% of EA in PE as eluant to give the product (1.2 g, 87.3%) as a yellow oil. MS (ESI, m/e) $[\text{M}+1]^+$ 260.0.

[0382] Step 2: 1-Benzyl-5'-oxo-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine- 4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carbonitrile




[0383] A mixture of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (1.0 g, 3.6 mmol), ethyl 2-(1-benzylpiperidin-4-ylidene)acetate (1.1 g, 4.3 mmol) and K_2CO_3 (745 mg, 5.4 mmol) in DMF (20 mL) was heated to 80 °C for 16 hr under N_2 . The reaction was quenched with water (20 mL) and extracted with EA (20 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , concentrated and purified by chromatography column on silica gel using 30% of EA in PE as eluant to give the product (950 mg, 54.9%) as a yellow solid. 1H NMR (400 MHz, DMSO- d_6) δ 11.92 (s, 1H), 7.84 (d, J = 8.8 Hz, 2H), 7.48-7.40 (m, 2H), 7.36-7.29 (m, 4H), 7.27-7.23 (m, 1H), 7.20 (t, J = 7.6 Hz, 1H), 7.15-7.07 (m, 4H), 3.55 (s, 2H), 3.01 (s, 2H), 2.81-2.73 (m, 2H), 2.39-2.27 (m, 2H), 2.26-2.16 (m, 2H), 1.83-1.74 (m, 2H). MS (ESI, m/e) $[M+1]^+$ 489.9.

[0384] Step 3: 1-Benzyl-5'-oxo-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine- 4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide



[0385] A solution of 1-benzyl-5'-oxo-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro [piperidine-4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carbonitrile (500 mg, 1.02 mmol) in H_3PO_4 (5 mL) was heated to 130 °C for 1 hr. The mixture was poured to water (20 mL) and extracted with EA (20 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , concentrated and purified by chromatography column on silica gel using 5% of MeOH in DCM as eluant to afford the product (180 mg, 34.8%) as a yellow solid. MS (ESI, m/e) $[M+1]^+$ 507.9.


[0386] Step 4: 1-Benzyl-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine-4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide



[0387] A solution of 1-benzyl-5'-oxo-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro [piperidine-4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide (180 mg, 0.36 mmol) in BH_3/THF (1N, 20 mL)


was heated to reflux for 3 hr. The reaction was quenched with MeOH (20 mL) and con. HCl (2 mL). The mixture was stirred at 60 °C for 1 hr, then was basified with NaHCO₃ and extracted with EA (20 mL × 3). The combined organic layers were dried over Na₂SO₄, concentrated and purified by chromatography column on silica gel eluting with 5% of MeOH in DCM to give the product (120 mg, 67.6%) as a yellow solid. ¹H NMR (400 MHz, CD₃OD-d₄) δ 7.39-7.22 (m, 9H), 7.09-7.04 (m, 1H), 6.99-6.94 (m, 4H), 3.86 (s, 2H), 3.35 (t, *J* = 5.6 Hz, 2H), 3.19-3.11 (m, 2H), 2.80-2.66 (m, 2H), 2.50-2.40 (m, 2H), 2.10 (t, *J* = 5.6 Hz, 2H), 1.87-1.78 (m, 2H). MS (ESI, m/e) [M+1]⁺ 493.9.

[0388] Compound 42: 2'-(4-Phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine-4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide trifluoroacetate



[0389] Compound **42** was prepared from 1-benzyl-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine-4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide according to the procedure similar to that for compound **39**. ¹H NMR (400 MHz, DMSO-d₆) δ 8.66 (br s, 2H), 7.51 (d, *J* = 8.6 Hz, 2H), 7.46-7.37 (m, 2H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.11-7.04 (m, 4H), 6.79 (s, 1H), 3.40-3.33 (m, 4H), 3.17-3.06 (m, 2H), 2.46-2.35 (m, 2H), 2.17-2.10 (m, 2H), 1.96-1.87 (m, 2H). MS (ESI, m/e) [M+1]⁺ 403.9.


[0390] Compound 43: 1-Acryloyl-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[piperidine-4,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide




[0391] Compound **43** was prepared from compound **42** and acryloyl chloride according to the procedure similar to that for compound **8**. ¹H NMR (400 MHz, DMSO-d₆) δ 7.50 (d, *J* = 8.8 Hz, 2H), 7.45-7.37 (m, 2H), 7.17 (t, *J* = 7.2 Hz, 1H), 7.12-7.01 (m, 4H), 6.85 (dd, *J* = 10.4, 16.7 Hz, 1H), 6.73 (s, 1H), 6.11 (dd, *J* = 2.4, 16.7 Hz, 1H), 5.68 (dd, *J* = 2.4, 10.4 Hz, 1H), 4.23 (d, *J* = 13.2 Hz, 1H), 4.03 (d, *J* = 13.2 Hz, 1H), 3.43 (t, *J* = 12.0, 1H), 3.37-3.33 (m, 2H), 3.16 (t, *J* = 12.0 Hz, 1H), 2.24-2.08 (m, 4H), 1.82-1.73 (m, 2H). MS (ESI, m/e) [M+1]⁺ 457.9.

[0392] Example 19: Synthesis of compounds 44-46

[0393] Compound **44**: 1-Benzyl-6'-(4-phenoxyphenyl)-1',2'-dihydrospiro[azetidine-3,3'-imidazo[1,2-b]pyrazole]-7'-carboxamide




[0394] Step 1: Methyl 1-benzylazetidine-3-carboxylate




[0395] To a solution of methyl azetidine-3-carboxylate (5.0 g, 33.1 mmol) and DIEA (10.7 g, 82.8 mmol) in DMF (50 mL) was added dropwise bromomethyl benzene (5.7 g, 33.1 mmol) at 0 °C over 10 min. After stirring for 2 hr at rt, the mixture was poured to water (50 mL) and extracted with EA (50 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄ and concentrated and purified by chromatography column on silica (EA/PE = 1/4) to give the product (3.5 g, 51.6%) as a light yellow oil. ¹H NMR (DMSO-d₆) δ 7.34-7.20 (m, 5H), 3.62 (s, 3H), 3.53 (s, 2H), 3.41-3.35 (m, 1H), 3.29-3.31 (m, 2H), 3.19-3.22 (m, 2H). MS (ESI, m/e) [M+1]⁺ 206.0.

[0396] Step 2: Di-tert-butyl 1-(1-benzyl-3-(methoxycarbonyl)azetidin-3-yl)hydrazine-1,2-dicarboxylate



[0397] The desired product was prepared from methyl 1-benzylazetidine-3-carboxylate and di-tert-butyl azodicarboxylate using the procedure similar to step 1 for compound **38**. MS (ESI, m/e) [M+1]⁺ 435.9.


[0398] Step 3 to 5: 5-Amino-1-(1-benzyl-3-(hydroxymethyl)azetidin-3-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile



[0399] The desired product was prepared from di-tert-butyl 1-(1-benzyl-3-(methoxycarbonyl)azetidin-3-yl)hydrazine-1,2-dicarboxylate using the procedures similar to those (step 2 to 4) for compound **38**. ¹H NMR (DMSO-d₆) δ 7.78 (d, J = 8.8 Hz, 2H), 7.46-7.38


(m, 2H), 7.36-7.21 (m, 5H), 7.18 (t, $J = 7.6$ Hz, 1H), 7.11-7.04 (m, 4H), 6.17 (s, 2H), 5.50 (t, $J = 5.2$ Hz, 1H), 3.89 (d, $J = 5.2$ Hz, 2H), 3.61-3.63 (m, 4H), 3.38 (d, $J = 6.4$ Hz, 2H). MS (ESI, m/e) $[M+1]^+$ 451.9.

[0400] Step 6: (3-(5-Amino-4-cyano-3-(4-phenoxyphenyl)-1H-pyrazol-1-yl)-1-benzyl azetidin-3-yl)methyl methanesulfonate




[0401] The desired product was prepared from 5-amino-1-(1-benzyl-3-(hydroxymethyl)azetidin-3-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile using the procedure similar to step 5 for compound **38**. ^1H NMR (DMSO- d_6) δ 7.82-7.74 (m, 2H), 7.46-7.38 (m, 2H), 7.36-7.23 (m, 5H), 7.21-7.15 (m, 1H), 7.13-7.04 (m, 4H), 6.52 (s, 2H), 4.69 (s, 2H), 3.70 (d, $J = 8.4$ Hz, 2H), 3.66 (s, 2H), 3.49 (d, $J = 8.4$ Hz, 2H), 3.13 (s, 3H). MS (ESI, m/e) $[M+1]^+$ 529.9.

[0402] Step 7, 8: 1-Benzyl-6'-(4-phenoxyphenyl)-1',2'-dihydrospiro[azetidine-3,3'-imidazo[1,2-b]pyrazole]-7'-carboxamide




[0403] The desired product was prepared from (3-(5-amino-4-cyano-3-(4-phenoxyphenyl)-1H-pyrazol-1-yl)-1-benzyl azetidin-3-yl)methyl methanesulfonate using the procedures similar to those (step 6 and 7) for compound **38**. ^1H NMR (DMSO- d_6) δ 7.66 (d, $J = 8.8$ Hz, 2H), 7.46-7.38 (m, 2H), 7.35-7.28 (m, 4H), 7.28-7.21 (m, 1H), 7.16 (t, $J = 7.6$ Hz, 1H), 7.06 (d, $J = 8.0$ Hz, 2H), 7.01 (d, $J = 8.8$ Hz, 2H), 6.52 (s, 1H), 4.19 (s, 2H), 3.67 (s, 2H), 3.54 (s, 4H). MS (ESI, m/e) $[M+1]^+$ 451.9.

[0404] Compound **45**: 6'-(4-Phenoxyphenyl)-1',2'-dihydrospiro[azetidine-3,3'-imidazo[1,2-b]pyrazole]-7'-carboxamide



[0405] The desired product was prepared from compound **44** using the procedure similar to that for compound **39**. ¹H NMR (DMSO-d₆) δ 7.67 (d, *J* = 8.4 Hz, 2H), 7.46-7.36 (m, 2H), 7.16 (t, *J* = 7.6 Hz, 1H), 7.06 (d, *J* = 7.6 Hz, 2H), 7.00 (d, *J* = 8.4 Hz, 2H), 6.62 (s, 1H), 4.19-4.20 (m, 4H), 3.78 (d, *J* = 9.6 Hz, 2H). MS (ESI, *m/e*) [*M*+1]⁺ 361.9.


[0406] Compound **46**: 1-Acryloyl-6'-(4-phenoxyphenyl)-1',2'-dihydrospiro[azetidine-3,3'-imidazo[1,2-*b*]pyrazole]-7'-carboxamide




[0407] The desired product was prepared from compound **45** and acryloyl chloride using the procedure similar to that for compound **8**. ¹H NMR (DMSO-d₆) δ 7.66 (d, *J* = 8.8 Hz, 2H), 7.45-7.37 (m, 2H), 7.16 (t, *J* = 7.6 Hz, 1H), 7.06 (d, *J* = 7.6 Hz, 2H), 7.00 (d, *J* = 8.8 Hz, 2H), 6.62 (s, 1H), 6.36 (dd, *J* = 17.0, 10.3 Hz, 1H), 6.15 (dd, *J* = 17.0, 2.1 Hz, 1H), 5.72 (dd, *J* = 10.3, 2.1 Hz, 1H), 4.62 (d, *J* = 9.6 Hz, 1H), 4.56 (d, *J* = 9.6 Hz, 1H), 4.32 (d, *J* = 11.2 Hz, 1H), 4.27 (d, *J* = 11.2 Hz, 1H), 4.22 (s, 2H). MS (ESI, *m/e*) [*M*+1]⁺ 415.9.

[0408] Example 20: Synthesis of compounds **47-50**

[0409] Compound **47**: 2-(3-Aminophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-*b*] pyrazole-7-carboxamide




[0410] Step 1: 5-Amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide




[0411] A solution of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (1.0 g, 3.6 mmol) in H₃PO₄ (20 mL) was heated to 120 °C for 4 hr. The mixture was then poured into water (100 mL), extracted with EA (100 mL × 3). The combined organic layers were dried over Na₂SO₄ and concentrated to give the product (850 mg, 77.5%) as yellow solid. MS (ESI, *m/e*) [*M*+1]⁺ 295.1.

[0412] Step 2: 2-(3-Nitrophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide




[0413] A mixture of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (29.4 mg, 0.1 mmol) and 2-bromo-1-(3-nitrophenyl)ethanone (24.4 mg, 0.1 mmol) in EtOH (2 mL) was stirred at 80 °C for 16 hr. The mixture was filtered to afford 5 mg of crude 2-(3-nitrophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide as a yellow solid. MS (ESI) m/e [M+1]⁺ 440.0.

[0414] Step 3: 2-(3-Aminophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide




[0415] To a solution of 2-(3-nitrophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide (600 mg, 1.37 mmol) in 10 mL of MeOH and 10 mL of DCM was added 10% w/w Pd/C (100 mg). After stirring at RT under H₂ for 4 hr, the mixture was filtered. The filtrate was concentrated and purified by Pre-HPLC eluting from 30% to 90% CH₃CN in 0.1% TFA in H₂O. Fractions containing the desired product were combined and lyophilized overnight to afford 73 mg (13%) of 2-(3-aminophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 12.03 (d, *J* = 10.4 Hz, 1H), 8.14 (d, *J* = 8.0 Hz, 1H), 7.73 (d, *J* = 8.4 Hz, 2H), 7.47-7.40 (m, 2H), 7.38-7.26 (m, 3H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.10 (d, *J* = 8.0 Hz, 2H), 7.03 (d, *J* = 8.4 Hz, 2H) and 6.98-6.86 (m, 2H). MS (ESI) m/e [M+1]⁺ 409.9.

[0416] Compound **48**: 2-(3-Acrylamidophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide




[0417] The desired product was prepared from 2-(3-aminophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, $\text{CD}_3\text{OD}-d_4$) δ 8.06 (s, 1H), 7.83 (s, 1H), 7.64 (d, $J = 8.4$ Hz, 1H), 7.52-7.38 (m, 5H), 7.15 (t, $J = 7.6$ Hz, 1H), 7.09-7.05 (m, 4H), 6.49-6.37 (m, 2H) and 5.80 (dd, $J = 4.0, 8.8$ Hz, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 463.9.

[0418] Compound **49**: 3-(3-Aminophenyl)-6-(4-phenoxyphenyl)-1H-pyrazolo[1,5-a]imidazole-7-carboxamide



[0419] The desired compound was separated as another isomer in the step 2 of compound **48**. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.22 (d, $J = 2.4$ Hz, 1H), 8.03 (s, 1H), 7.89 (d, $J = 2.4$ Hz, 1H), 7.81-7.68 (m, 3H), 7.46-7.37 (m, 4H), 7.19 (t, $J = 7.6$ Hz, 1H), 7.13-7.08 (m, 5H) and 6.98 (d, $J = 7.6$ Hz, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 410.1.


[0420] Compound **50**: 3-(3-Acrylamidophenyl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide




[0421] The desired product was prepared from compound **49** and acryloyl chloride using the procedure similar to compound **8**. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.18 (d, $J = 2.4$ Hz, 1H), 10.31 (s, 1H), 8.38 (s, 1H), 7.85 (d, $J = 8.0$ Hz, 1H), 7.80-7.73 (m, 4H), 7.46-7.40 (m, 3H), 7.19 (t, $J = 8.0$ Hz, 1H), 7.13-7.07 (m, 4H), 6.50 (dd, $J = 10.2, 17.0$ Hz, 1H), 6.27 (d, $J = 17.0$ Hz, 1H), 5.76 (d, $J = 10.2$ Hz, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 463.9.

[0422] Example 21: Synthesis of compounds **51** to **60**

[0423] Compound **51**: 2-(4-Phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide trifluoroacetate




[0424] Step 1: tert-Butyl 3-bromo-4-oxopiperidine-1-carboxylate



[0425] To a solution of tert-butyl 4-oxopiperidine-1-carboxylate (5 g, 25 mmol) in 30 mL of DMF at RT was added TEA (7.7 mL, 55 mmol) followed by TMSCl (3.5 mL, 27.6 mmol), then the mixture was stirred at 75 °C overnight. The reaction was cooled to RT, cold *sat. aq.* NaHCO₃ (200 mL) was added followed by cold hexane (200 mL). The organic layer was washed with brine, dried over Na₂SO₄, concentrated to get the crude product directly used in the next step. The residue was dissolved in 15 mL of THF and stirred at 0 °C for 15 min. A solution of NBS (4.47 g, 25 mmol) in 80 mL of THF was added slowly. After addition, the reaction was stirred at RT overnight. Water (200 mL) was added to the reaction followed by 200 mL of hexane. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated to get crude product which was chromatographed on 60 g of silica gel using PE/EA (20/1 to 8/1) as eluant to afford 5.56 g (78%) of tert-butyl 3-bromo-4-oxopiperidine-1-carboxylate as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 4.85-4.70 (m, 1H), 4.20-4.00 (m, 1H), 3.90-3.55 (m, 3H), 2.80-2.68 (m, 1H), 2.54-2.44 (m, 1H), 1.43 (s, 9H). MS (ESI) m/e [M-t-Bu]⁺ 221.9, 224.0.


[0426] Step 2: tert-Butyl 3-cyano-2-(4-phenoxyphenyl)-5,6-dihydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-7(8H)-carboxylate



[0427] A mixture of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (1.5 g, 5.4 mmol) and K₂CO₃ (2.24 g, 16.3 mmol) in 50 mL of DMF at 80 °C was stirred under N₂ for 45 min before tert-butyl 3-bromo-4-oxopiperidine-1-carboxylate (4.5 g, 16.3 mmol) was added in one portion. Then the mixture was stirred at 80 °C for 1 hr. After cooling down to RT, 150 mL of water and 150 mL of EA was added. Aqueous phase was further extracted with EA (100 mL


× 3). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated to get crude product which was chromatographed on 15 g of silica gel using DCM/MeOH (400/1 to 200/1) as eluant to afford 850 mg (35%) of tert-butyl 3-cyano-2-(4-phenoxyphenyl)-5,6-dihydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-7(8H)-carboxylate as an off-white solid. MS (ESI) m/e [M+1]⁺ 455.9.

[0428] Step 3: 2-(4-Phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide trifluoroacetate



[0429] A solution of tert-butyl 3-cyano-2-(4-phenoxyphenyl)-5,6-dihydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-7(8H)-carboxylate (130 mg, 0.28 mmol) in H₃PO₄ (85 wt. % in H₂O, 20 mL) was stirred at 100 °C for 1.5 hr, until TLC and LCMS analysis showed that most of starting material was consumed. The mixture was cooled to room temperature and poured into water (100 mL). The mixture was adjust to PH = 9-10 with solid K₂CO₃. The suspension was extracted with EA (100 mL × 4). The combined organic layers were washed with brine (200 mL), dried over Na₂SO₄ and concentrated to get the crude product which was purified with pre-HPLC eluting from 10% to 90% CH₃CN in 0.1% TFA in H₂O. Fractions containing the desired product were combined and lyophilized overnight to give 15 mg (11%) of 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide trifluoroacetate as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 11.99 (s, 1H), 9.32 (s, 2H), 7.66 (d, *J* = 8.6 Hz, 2H), 7.43 (d, *J* = 7.6 Hz, 1H), 7.41 (d, *J* = 7.6 Hz, 1H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.10-7.06 (m, 4H), 4.44 (s, 2H), 3.49 (m, 2H), 2.95-2.92 (m, 2H). MS (ESI) m/e [M+1]⁺ 373.9.


[0430] Compound **52**: 7-Acryloyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide



[0431] The desired product was prepared from compound **51** and acryloyl chloride using the procedure similar to that for compound **8**. ¹H NMR (400 MHz, DMSO-d₆ at 80 °C) δ 11.55 (s, 1H), 7.71 (d, *J* = 8.6 Hz, 2H), 7.43 (d, *J* = 7.6 Hz, 1H), 7.41 (d, *J* = 7.6 Hz, 1H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.10-7.05 (m, 4H), 6.88 (dd, *J* = 10.6, 17.1 Hz, 1H), 6.24 (s, 2H), 6.15 (d, *J* = 17.1 Hz,


1H), 5.74 (d, J = 10.6 Hz, 1H), 4.78 (s, 2H), 3.94-3.91 (m, 2H), 2.80-2.76 (m, 2H). MS (ESI) m/e $[M+1]^+$ 427.9.

[0432] Compound **53**: 7-(3-Chloropropanoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide



[0433] 7-Acryloyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide (40 mg, 0.09 mmol) was suspended in *Sat.* HCl (gas) /dioxane (50 mL), then the mixture was stirred RT for about 1.5 hr, and concentrated to dryness. The residue was suspended into 2 mL of MeOH and 2 mL of water. The organic layer was discarded, aqueous layer was lyophilized to get 40 mg (90%) of 7-(3-chloropropanoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide as an off-white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 11.79-11.76 (m, 1H), 7.68 (d, J = 8.6 Hz, 2H), 7.43 (d, J = 7.6 Hz, 1H), 7.41 (d, J = 7.6 Hz, 1H), 7.18 (t, J = 7.6 Hz, 1H), 7.10-7.04 (m, 4H), 4.71-4.70 (m, 2H), 3.85-3.79 (m, 4H), 3.02 (t, J = 6.4 Hz, 2H), 2.79-2.69 (m, 2H). MS (ESI) m/e $[M+1]^+$ 463.8, 465.8.

[0434] Compound **54** and **55**: (*E*)-7-(4-(Dimethylamino)but-2-enoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide and 7-Acetyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide




[0435] A mixture of (*E*)-4-(dimethylamino)but-2-enoic acid hydrochloride (147 mg, 0.88 mmol), HATU (611 mg, 1.6 mmol) and TEA (328 mg, 3.2 mmol) in 50 mL of DCM was stirred at RT for about 2 hr before 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide (300 mg, 0.8 mmol) was added. The mixture was stirred at RT overnight. TLC and LCMS analysis showed that starting material was consumed. To the reaction were added 100 mL of water and 50 mL of DCM. Aqueous phase was further extracted with 50 mL of DCM. The combined organic layers were washed with

brine, dried over Na₂SO₄, concentrated to get crude product which was chromatographed on 5 g of silica gel using DCM/MeOH (20/1 to 10/1) as eluant to afford 145 mg (37%) of (*E*)-7-(4-(dimethylamino) but-2-enoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-*c*]pyridine-3-carboxamide, which was dried by lyophilization. ¹H NMR (400 MHz, DMSO-*d*₆ at 80 °C) δ 11.52 (s, 1H), 7.69 (d, *J* = 8.6 Hz, 2H), 7.42 (d, *J* = 7.6 Hz, 1H), 7.41 (d, *J* = 7.6 Hz, 1H), 7.16 (t, *J* = 7.6 Hz, 1H), 7.12-6.99 (m, 4H), 6.78-6.56 (m, 2H), 6.22 (s, 2H), 4.75 (s, 2H), 3.89 (t, *J* = 5.6 Hz, 2H), 3.12 (d, *J* = 5.6 Hz, 2H), 2.80-2.72 (m, 2H), 2.22 (s, 6H). MS (ESI) *m/e* [M+1]⁺ 484.9.


[0436] 7-Acetyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-*c*]pyridine-3-carboxamide was prepared as a byproduct due to some of HOAc residue in the last step. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.77-11.73 (m, 1H), 7.70-7.66 (m, 2H), 7.43 (d, *J* = 7.6 Hz, 1H), 7.41 (d, *J* = 7.6 Hz, 1H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.13-6.94 (m, 4H), 4.68 (s, 2H), 3.83-3.76 (m, 2H), 2.82-2.77 (m, 2H), 2.14 (s, 3H). MS (ESI) *m/e* [M+1]⁺ 416.

[0437] Compound **56**: 7-(2-Cyanoacetyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-*c*]pyridine-3-carboxamide




[0438] The desired compound was prepared from 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-*c*]pyridine-3-carboxamide and 2-cyanoacetic acid according to the procedure similar to that for compound **54**. ¹H NMR (400 MHz, DMSO-*d*₆ at 80 °C) δ 11.55 (s, 1H), 7.68 (d, *J* = 8.6 Hz, 2H), 7.41 (d, *J* = 7.6 Hz, 1H), 7.39 (d, *J* = 7.6 Hz, 1H), 7.16 (t, *J* = 7.6 Hz, 1H), 7.11-6.93 (m, 4H), 6.22 (br s, 2H), 4.68 (s, 2H), 4.14 (s, 2H), 3.86-3.79 (m, 2H), 2.84-2.73 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 440.9.

[0439] Compound **57**: 7-(3-(Dimethylamino)propanoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-*c*]pyridine-3-carboxamide trifluoroacetate




[0440] To a solution of 7-acryloyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide (6 mg, 0.014 mmol) in 5 mL of MeOH at RT was added NaOMe (15 mg, 0.28 mmol) followed by dimethylamine hydrochloride (12 mg, 0.14 mmol), then the mixture was stirred at 50 °C overnight. After cooling down to RT, the mixture was concentrated. The residue was purified by pre-HPLC eluting from 0% to 60% CH₃CN in H₂O. Fractions containing the desired product were combined and lyophilized overnight to give 2.5 mg (35%) of 6-(3-(dimethylamino)propanoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 11.84-11.82 (m, 1H), 9.55 (s, 1H), 7.69-7.66 (m, 2H), 7.44 (d, *J* = 7.6 Hz, 1H), 7.42 (d, *J* = 7.6 Hz, 1H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.10-7.04 (m, 4H), 4.75-4.72 (m, 2H), 3.87-3.82 (m, 2H), 3.02-3.00 (m, 2H), 2.84-2.78 (m, 2H), 2.77 (s, 6H), 2.71-2.68 (m, 2H). MS (ESI) *m/e* [M+1]⁺ 472.9.

[0441] Compound **58**: 7-(But-2-enoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide



[0442] The desired compound was prepared from 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide and but-2-enoic acid according to the procedure similar to that for compound **54**. ¹H NMR (400 MHz, DMSO-d₆) δ 11.78-11.71 (m, 1H), 7.68 (d, *J* = 8.6 Hz, 2H), 7.46-7.39 (m, 2H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.11-7.04 (m, 4H), 6.73-6.63 (m, 2H), 4.80-4.71 (m, 2H), 3.90 (s, 2H), 2.76-2.70 (m, 2H), 1.88-1.86 (m, 3H). MS (ESI) *m/e* [M+1]⁺ 441.9.

[0443] Compound **59** and **60**: (*E*)-7-(3-Cyanoallyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide and (*Z*)-7-(3-Cyanoallyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide




[0444] To a solution of 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide (100 mg, 0.268 mmol) in 10 mL of acetone at RT was added K_2CO_3 (140 mg, 1.07 mmol). After stirring at RT for 2 hr, 4-bromobut-2-enenitrile (40 mg, 0.268 mmol) in 2 mL of acetone was added and stirred at RT overnight. The mixture was then partitioned between EA (50 mL) and water (100 mL). The aqueous phase was further extracted with 50 mL of EA. The combined organic layers were washed with brine, dried over Na_2SO_4 , concentrated to get crude product which was further purified by pre-TLC (DCM/MeOH = 15/1) to afford 7 mg (6%) of (*E*)-7-(3-cyanoallyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide as a light yellow solid. 1H NMR (400 MHz, $CDCl_3$) δ 10.45 (s, 1H), 7.59 (d, J = 8.4 Hz, 2H), 7.42-7.30 (m, 2H), 7.16 (t, J = 7.6 Hz, 1H), 7.10 (d, J = 8.4 Hz, 2H), 7.06 (d, J = 7.6 Hz, 2H), 6.76 (dt, J = 16.3, 4.6 Hz, 1H), 5.67 (d, J = 16.3 Hz, 1H), 5.62 (s, 2H), 3.82 (s, 2H), 3.41 (d, J = 4.6 Hz, 2H), 2.95-2.75 (m, 4H). MS (ESI) m/e $[M+1]^+$ 439.9.


[0445] 4 mg (3.4%) of (*Z*)-7-(3-cyanoallyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide as a light yellow solid. 1H NMR (400 MHz, $CDCl_3$) δ 10.22 (s, 1H), 7.60 (d, J = 8.4 Hz, 2H), 7.41-7.33 (m, 2H), 7.16 (t, J = 7.6 Hz, 1H), 7.10 (d, J = 8.4 Hz, 2H), 7.06 (d, J = 7.6 Hz, 2H), 6.70-6.59 (m, 1H), 5.58 (s, 2H), 5.53 (d, J = 11.2 Hz, 1H), 3.85 (s, 2H), 3.64 (d, J = 6.4 Hz, 2H), 2.99-2.80 (m, 4H). MS (ESI) m/e $[M+1]^+$ 439.9.

[0446] Example 22: Synthesis of compounds **61** to **64**

[0447] Compound **61**: 2-(4-Phenoxyphenyl)-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide




[0448] Step 1: 3-Bromo-4-hydrazinylpyridine




[0449] A mixture of 3-bromo-4-chloropyridine (5 g, 0.026 mol) and hydrazine hydrate (80% in water, 80 mL) in dioxane (100 mL) was stirred at 100 °C overnight. After cooling down to RT, the mixture was concentrated. The residue was partitioned between 300 mL of EA and 300 mL of *aq. sat.* NH₄Cl. The organic layer was washed with brine, dried over Na₂SO₄, filtered and concentrated to get crude product which was suspended in 30 mL of cold isopropyl alcohol and filtered. The collected solid was dried in air to get 4.2 g (87%) of 3-bromo-4-hydrazinylpyridine as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.18 (s, 1H), 8.07 (d, *J* = 5.6 Hz, 1H), 7.37 (s, 1H), 7.01 (d, *J* = 5.6 Hz, 1H), 4.36 (s, 2H).

[0450] Step 2: 5-Amino-1-(3-bromopyridin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile




[0451] A mixture of 2-(methoxy(4-phenoxyphenyl)methylene)malononitrile (7.3 g, 0.026 mol) and 3-bromo-4-hydrazinylpyridine (4.2 g, 0.022 mol) in ethanol (300 mL) was stirred at reflux under N₂ overnight. The reaction was cooled to RT slowly and stirred at RT for about 4 hr till solid precipitated. The solid was filtered, collected and washed with hexane to get 3.38 g (35%) of 5-amino-1-(3-bromopyridin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile as a light yellow solid. MS (ESI) *m/e* [M+1]⁺ 431.8, 433.8.

[0452] Step 3: 5-Amino-1-(3-bromopyridin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide




[0453] The desired compound was prepared from 5-amino-1-(3-bromopyridin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile according to the procedure similar to step 3 for compound **51**. MS (ESI) *m/e* [M+1]⁺ 449.8, 451.8.

[0454] Step 4: 2-(4-Phenoxyphenyl)-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide




[0455] A mixture of 5-amino-1-(3-bromopyridin-4-yl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (3.96 g, 8.8 mmol), CuI (836 mg, 4.4 mmol), *N*¹,*N*²-dimethylethane-1,2-diamine (77 mg, 0.88 mmol), K₃PO₄ (5.59 g, 26.4 mmol) in 100 mL of DMF was stirred at 100 °C under N₂ for 2 hr, until TLC showed that most of starting material was consumed. After cooling down to RT, the mixture was filtered and concentrated. The residue was chromatographed on 30 g of silica gel using DCM/ MeOH (20/1 to 10/1) as eluant to afford 3.2 g (99%) of 2-(4-Phenoxyphenyl)-4H-pyrazolo [1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide as a tan solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.63 (br s, 1H), 8.85 (s, 1H), 8.46 (s, 1H), 7.96 (d, *J* = 4.6 Hz, 1H), 7.81 (d, *J* = 8.6 Hz, 2H), 7.45 (d, *J* = 7.6 Hz, 1H), 7.43 (d, *J* = 7.6 Hz, 1H), 7.20 (t, *J* = 7.6 Hz, 1H), 7.14-7.09 (m, 4H). MS (ESI) *m/e* [M+1]⁺ 369.9.

[0456] Compound **62**: 2-(4-Phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide




[0457] Step 1: 6-Benzyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide



[0458] To a suspension of 2-(4-phenoxyphenyl)-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide (2.46 g, 0.0067 mol) in 150 mL of THF was added benzyl bromide (1.14 g, 0.0067 mol) dropwise, then the mixture was stirred at 65 °C overnight. After cooling down to RT, the mixture was concentrated, the residue was suspended in 150 mL of MeOH, NaBH₄ (10 g, 0.26 mol) was added portionwise. The mixture was stirred at RT overnight. To the reaction


was added 200 mL of water followed by 200 mL of DCM. The aqueous phase was further extracted with 100 mL of DCM. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated to get the crude product which was chromatographed on 10 g of silica gel using DCM/MeOH (200/1 to 80/1) as eluant to afford 0.786 g (26%) of 6-benzyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide as a tan foam. MS (ESI) m/e [M+1]⁺ 463.9.

[0459] Step 2: 2-(4-Phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide




[0460] A mixture of 6-benzyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide (640 mg, 0.00138 mol) and 10% w/w Pd/C (700 mg) in 60 mL of MeOH was stirred at RT under 1 atm of H₂ overnight. TLC and LCMS analysis showed starting material was consumed. The reaction was filtered, filtrate was concentrated to get 397 mg (77%) of 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide. ¹H NMR (400 MHz, CD₃OD-d₄) δ 7.65-7.55 (m, 2H), 7.43-7.33 (m, 2H), 7.18-7.12 (m, 1H), 7.11-6.99 (m, 4H), 4.18 (s, 2H), 3.44 (t, *J* = 5.8 Hz, 2H), 3.01 (t, *J* = 5.8 Hz, 2H). MS (ESI) m/e [M+1]⁺ 373.9.

[0461] Compound **63**: 6-Acryloyl-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide trifluoroacetate



[0462] The desired compound was prepared from 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide and acryloyl chloride according to the procedure similar to that for compound **8**. ¹H NMR (400 MHz, DMSO-d₆) δ 11.79 (s, 1H), 7.73 (d, *J* = 8.6 Hz, 2H), 7.50 (d, *J* = 7.6 Hz, 1H), 7.48 (d, *J* = 7.6 Hz, 1H), 7.24 (t, *J* = 7.6 Hz, 1H), 7.17-7.11 (m, 4H), 7.06-6.98 (m, 1H), 7.22-6.26 (m, 1H), 5.84-5.82 (m, 1H), 4.72 (s, 2H), 4.00-3.97 (m, 2H), 2.94-2.90 (m, 2H). MS (ESI) m/e [M+1]⁺ 427.9.


[0463] Compound **64**: (E)-6-(4-(Dimethylamino)but-2-enoyl)-2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide trifluoroacetate




[0464] The desired compound was prepared from 2-(4-phenoxyphenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[1',5':1,2]imidazo[4,5-c]pyridine-3-carboxamide and (E)-4-(dimethylamino)but-2-enoic acid hydrochloride according to the procedure similar to that for compound **54**. ^1H NMR (400 MHz, DMSO- d_6) δ 11.82-11.74 (m, 1H), 10.06 (br s, 1H), 7.66 (d, J = 8.6 Hz, 2H), 7.43 (d, J = 7.6 Hz, 1H), 7.42 (d, J = 7.6 Hz, 1H), 7.18 (t, J = 7.6 Hz, 1H), 7.10-7.05 (m, 4H), 6.74-6.54 (m, 2H), 4.66 (s, 2H), 3.99-3.86 (m, 4H), 2.88-2.76 (m, 2H), 2.78 (s, 6H). MS (ESI) m/e $[\text{M}+1]^+$ 484.9.

[0465] Example 23: Synthesis of compounds 65-67

[0466] Compound **65**: 7-Nitro-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b]pyrazole-3-carboxamide




[0467] Step 1: (2-Bromo-5-nitrophenyl)hydrazine



[0468] To a suspension of 2-bromo-5-nitroaniline (1 g, 4.6 mmol) in *conc.* HCl (10 mL) at 0 $^{\circ}\text{C}$ was slowly added a solution of NaNO_2 (382 mg, 5.5 mmol) in water (1.5 mL). Then, the mixture was stirred at 0 $^{\circ}\text{C}$ for 3hr until TLC and LCMS analysis showed that most of 2-bromo-5-nitroaniline was consumed. SnCl_2 (1.90 g, 10 mmol) in *conc.* HCl (3 mL) was slowly added. The mixture was then stirred at RT for 2 hr before re-cooled to 0 $^{\circ}\text{C}$. Then, the PH was adjusted with *sat. aq.* NaHCO_3 to 7-8. The mixture was extracted with ethyl acetate (3 $\times$ 50 mL). The


combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated to get crude product which was further chromatographed on 10 g of silica gel using PE/EA (20/1 to 4/1) as eluant to afford 560 mg (51%) of (2-bromo-5-nitrophenyl)hydrazine as a orange solid. ¹H NMR (400 MHz, DMSO-d₆) δ 7.89 (d, *J* = 2.8 Hz, 1H), 7.60 (d, *J* = 8.6 Hz, 1H), 7.29 (dd, *J* = 2.8, 8.6 Hz, 1H), 7.04 (s, 1H), 4.38 (s, 2H). MS (ESI) *m/e* [M+1]⁺ 232, 234.

[0469] Step 2: 5-Amino-1-(2-bromo-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile




[0470] To a solution of 2-(methoxy(4-phenoxyphenyl)methylene)malononitrile (392 mg, 1.42 mmol) in ethanol (30 mL) was added (2-bromo-5-nitrophenyl)hydrazine (300 mg, 1.29 mmol) in one portion, then the mixture was stirred at 70 °C under N₂ overnight. The mixture was concentrated to dryness and chromatographed on 5 g of silica gel using PE/EA (10/1 to 2/1) as eluant to afford 128 mg (21%) of 5-amino-1-(2-bromo-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile as a yellow solid. MS (ESI) *m/e* [M+1]⁺ 476, 478.

[0471] Step 3: 5-Amino-1-(2-bromo-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide




[0472] A mixture of 5-amino-1-(2-bromo-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (137 mg, 0.287 mmol) in phosphorous acid (85 wt. % in H₂O, 10 mL) was stirred at 100 °C for 1 hr, until TLC and LCMS analysis showed that most of starting material was consumed. The reaction was cooled to room temperature and partitioned between water (40 mL) and EA (40 mL). Organic layer was separated from aqueous layer. The aqueous phase was then extracted with EA (20 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄ and concentrated to get the crude product (149 mg) which was used in next step without further purification. MS (ESI) *m/e* [M+1]⁺ 494, 496.

[0473] Step 4: 7-Nitro-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b]pyrazole-3-carboxamide




[0474] A mixture of 5-amino-1-(2-bromo-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (149 mg, 0.3 mmol, crude), CuI (5.7 mg, 0.03 mmol), *N*¹,*N*²-dimethylethane-1,2-diamine (3 mg, 0.03 mmol), K₃PO₄ (64 mg, 0.3 mol) in 15 mL of DMF was stirred at 60 °C under N₂ for 5hr, until TLC analysis showed that most of starting material was consumed. The reaction was cooled to RT. The solvent was removed under reduced pressure. The residue was chromatographed on 5 g of silica gel using DCM/MeOH (200/1 to 20/1) to afford 62 mg (52%) of 7-nitro-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-*b*]pyrazole-3-carboxamide as a tan solid. MS (ESI) *m/e* [M+1]⁺ 414.

[0475] Compound **66**: 7-Amino-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-*b*]pyrazole-3-carboxamide



[0476] To a solution of 7-nitro-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-*b*]pyrazole-3-carboxamide (9 mg, 0.022 mmol) in 3 mL of HOAc was added zinc powder (14 mg, 0.22 mmol). Then the mixture was stirred at RT for 20 min, until TLC and LCMS analysis showed that most of starting material was consumed. The reaction solid was filtered off. The filtrate was concentrated, suspended in 10 mL of EA and filtered. The filtrate was concentrated to get the product as a white solid (4 mg, 50%). ¹H NMR (400 MHz, CD₃OD-*d*₄) δ 7.60 (d, *J* = 8.6 Hz, 2H), 7.32 (d, *J* = 8.0 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 1H), 7.23 (d, *J* = 8.4 Hz, 1H), 7.16 (s, 1H), 7.08 (t, *J* = 8.0 Hz, 1H), 7.05-6.95 (m, 4H), 6.79 (dd, *J* = 1.6, 8.4 Hz, 1H). MS (ESI) *m/e* [M+1]⁺ 384.


[0477] Compound **67**: 7-Acrylamido-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-*b*]pyrazole-3-carboxamide




[0478] To a solution of 7-amino-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b] pyrazole-3-carboxamide (45 mg, 0.11 mmol) in 10 mL of DCM at 0 °C was added TEA (36 mg, 0.35 mmol). Acryloyl chloride (11 mg, 0.12 mmol) in 2 mL of DCM was added dropwise over a period of 20 min. The mixture was stirred until TLC and LCMS analysis showed that most of starting material was consumed. The mixture was then partitioned between water (50 mL) and DCM (20 mL), extracted with additional 20 mL of DCM. The combined organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by Pre-TLC (DCM/MeOH = 20/1) to get 4 mg (7.8%) of 7-acrylamido-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b] pyrazole-3-carboxamide as a grey solid. ¹H NMR (400 MHz, CD₃OD-d₄) δ 8.25 (s, 1H), 7.61 (d, *J* = 8.4 Hz, 2H), 7.44 (d, *J* = 8.8 Hz, 1H), 7.39 (d, *J* = 8.8 Hz, 1H), 7.32 (d, *J* = 7.6 Hz, 1H), 7.30 (d, *J* = 7.6 Hz, 1H), 7.08 (t, *J* = 7.6 Hz, 1H), 7.05-6.90 (m, 4H), 6.42-6.25 (m, 2H), 5.69 (dd, *J* = 9.6, 1.9 Hz, 1H). MS (ESI) *m/e* [M+1]⁺ 438.

[0479] Example 24: Synthesis of compounds 68-69

[0480] Compound **68**: 8-Amino-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a] quinazoline-3-carboxamide




[0481] Step 1: (2-Fluoro-4-nitrophenyl)methanol




[0482] To a solution of 2-fluoro-4-nitrobenzaldehyde (1.0 g, 5.92 mmol) in CH₃OH (10 mL) was added NaBH₄ (814 mg, 22 mmol). After stirring at RT for 15 min, the mixture was concentrated. The residue was partitioned between 100 mL of EA and 100 mL of brine. The combined organic layers were washed with brine (100 mL × 2), dried over Na₂SO₄, and concentrated to afford 1.0 g of (2-fluoro-4-nitrophenyl) methanol (99%) as a red solid. MS (ESI) *m/e* [M+1]⁺ 172.0.

[0483] Step 2: 2-(2-Fluoro-4-nitrobenzyloxy)-tetrahydro-2H-pyran




[0484] To a solution of (2-fluoro-4-nitrophenyl)methanol (755 mg, 4.42 mmol) in 10 mL of DCM was added TsOH (100 mg, 0.13 mmol) and DHP (408 mg, 4.86 mmol). After stirring at RT for 16 hr, the mixture was concentrated. The residue was partitioned between 100 mL of EA and 100 mL of brine. The combined organic layers were washed with brine (100 mL $\times$ 2), dried over Na₂SO₄, concentrated and purified by chromatography column on silica gel (elution with PE/EA) to afford 900 mg (80%) of 2-(2-fluoro-4-nitrobenzyloxy)-tetrahydro-2H-pyran as a colorless oil. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.34 (dd, *J* = 3.0, 6.2 Hz, 1H), 8.26-8.30 (m, 1H), 7.53 (t, *J* = 9.2 Hz, 1H), 4.82-4.76 (m, 2H), 4.62 (d, *J* = 12.0 Hz, 1H), 3.80-3.74 (m, 1H), 3.52-3.47 (m, 1H), 1.76-1.64 (m, 2H) and 1.58-1.45 (m, 4H).

[0485] Step 3: 5-Amino-1-(5-nitro-2-((tetrahydro-2H-pyran-2-yloxy)methyl)phenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide



[0486] To a solution of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (27.6 mg, 0.1 mmol) in DMF (3 mL) and CH₃CN (5 mL) was added 2-(2-fluoro-4-nitrobenzyloxy)-tetrahydro-2H-pyran (25.5 mg, 0.1 mmol) and K₂CO₃ (27.6 mg, 0.2 mmol). After stirring at 80 °C under N₂ for 16 hr, the mixture was concentrated and recrystallized with PE/EA to afford 40 mg (80%) of 5-amino-1-(5-nitro-2-((tetrahydro-2H-pyran-2-yloxy)methyl)phenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide as a yellow solid. MS (ESI) *m/e* [M+1]⁺ 512.2.


[0487] Step 4: 5-Amino-1-(2-(hydroxymethyl)-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide



[0488] To a solution of 5-amino-1-(5-nitro-2-((tetrahydro-2H-pyran-2-yloxy)methyl)phenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (690 mg, 1.3 mmol) in 10 mL of CH₃CN was added hydrochloric acid (3 mL). After stirring at RT for 15 min, the mixture was concentrated to


afford 550 mg (95%) of 5-amino-1-(2-(hydroxyl methyl)-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide as a yellow solid.

[0489] Step 5: 5-Amino-1-(2-formyl-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole -4-carboxamide




[0490] To a solution of 5-amino-1-(2-(hydroxymethyl)-5-nitrophenyl)-3-(4-phenoxy phenyl)-1H-pyrazole-4-carboxamide (550 mg, 1.24 mmol) in 20 mL of DCM was added MnO_2 (500 mg, 5.75 mmol). After stirring at RT for 16 hr, the mixture was filtered. The filtrate was concentrated to afford 400 mg (73%) of 5-amino-1-(2-formyl -5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide as a yellow solid.

[0491] Step 6: 8-Nitro-2-(4-phenoxyphenyl)pyrazolo[1,5-a]quinazoline-3-carboxamide



[0492] To a solution of 5-amino-1-(2-formyl-5-nitrophenyl)-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (400 mg, 0.9 mmol) in 5 mL of CH_3OH and 5 mL of DCM was added HOAc (1 drops). After stirring at RT for 16 hr, the mixture was concentrated and purified by chromatography column on silica gel eluting with PE/EA to afford 240 mg (63%) of 8-nitro-2-(4-phenoxyphenyl)pyrazolo [1,5-a]quinazoline-3-carboxamide as a yellow solid. MS (ESI) m/e $[\text{M}+1]^+$ 425.8.


[0493] Step 7: 8-Nitro-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide



[0494] To a solution of 8-nitro-2-(4-phenoxyphenyl)pyrazolo[1,5-a]quinazoline-3- carboxamide (240 mg, 0.57 mmol) in 10 mL of EtOH and 10 mL of DCM was added NaBH_4 (86 mg, 2.26 mmol) at RT. After stirring at RT for 20 min, 10 mL of water was added. The mixture was concentrated. 5 mL of water was added and filtered. The cake was washed with tert-Butyl


methyl ether (30 mL) and dried to afford 200 mg (83%) of 8-nitro-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide as a yellow solid. MS (ESI) m/e $[M+1]^+$ 427.9.

[0495] Step 8: 8-Amino-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide and 8-amino-2-(4-phenoxyphenyl)pyrazolo[1,5-a]quinazoline-3-carboxamide




[0496] To a solution of 8-nitro-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide (200 mg, 0.47 mmol) in 30 mL of CH_3OH and 30 mL of DCM was added 10% w/w Pd/C (100 mg). After stirring at RT for 1 hr, the mixture was filtered. The filtrate was concentrated to afford 130 mg (70%) of crude 8-amino-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide and 8-amino-2-(4-phenoxyphenyl)pyrazolo[1,5-a]quinazoline-3-carboxamide as a yellow solid. MS (ESI) m/e $[M+1]^+$ 398.0, 395.9.

[0497] Step 9: 8-Amino-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide



[0498] To a solution of the mixture of 8-amino-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide and 8-amino-2-(4-phenoxyphenyl)pyrazolo[1,5-a]quinazoline-3-carboxamide (130 mg, 0.33 mmol) in 10 mL of DCM and 10 mL of CH_3OH was added NaBH_4 (277 mg, 3.3 mmol). After stirring at RT for 15 min, 50 mL of water was added. The mixture was concentrated and filtered. The cake was washed with water (50 mL $\times$ 2) to afford 60 mg (46%) of 8-amino-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.60 (d, J = 8.0 Hz, 2H), 7.44 (d, J = 7.6 Hz, 1H), 7.42 (d, J = 7.6 Hz, 1H), 7.28 (d, J = 7.6 Hz, 1H), 7.18 (t, J = 7.6 Hz, 1H), 7.14-7.16 (m, 4H), 6.81 (s, 1H), 6.52 (d, J = 8.0 Hz, 1H), 6.43 (s, 1H), 5.16 (s, 2H) and 4.37 (s, 2H). MS (ESI) m/e $[M+1]^+$ 397.9.


[0499] Compound **69**: 8-Acrylamido-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide




[0500] The desired product was prepared from compound **68** and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 10.26 (s, 1H), 7.39-7.63 (m, 4H), 7.52 (d, J = 8.8 Hz, 1H), 7.45 (d, J = 8.0 Hz, 1H), 7.42 (d, J = 8.0 Hz, 1H), 7.19 (t, J = 8.0 Hz, 1H), 7.14-7.07 (m, 4H), 7.01 (s, 1H), 6.44 (dd, J = 10.4, 17.0 Hz, 1H), 6.26 (dd, J = 1.6, 17.0 Hz, 1H), 5.77 (dd, J = 1.6, 10.4 Hz, 1H) and 4.51 (s, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 451.9.

[0501] **Example 25: Synthesis of compounds 70-72**

[0502] Compound **70**: 8-Nitro-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo [1,5-a][1,3]diazepine-3-carboxamide




[0503] Step 1: 2-(2-Fluoro-5-nitrophenyl)ethanol




[0504] To a solution of 2-(2-fluoro-5-nitrophenyl)acetic acid (2.0 g, 10 mmol) in THF (50 mL) was added borane dimethyl sulfide complex solution (4.0 g, 25 mmol). The reaction was warmed to 60°C stirred for about 12 hr. After cooling down to RT, CH₃OH (20 mL) was slowly added to the reaction, concentrated under reduced pressure to remove solvent. The residue was purified by column chromatography on silica gel (200-300 mesh, PE/EA = 2/1) to afford the product as a colorless oil (1.6 g, 86.1%). ^1H NMR (400 MHz, DMSO- d_6) δ 8.27 (t, J = 3.2, 6.4 Hz, 1 H), 8.19-8.14 (m, 1 H), 8.45 (t, J = 9.2 Hz, 1 H), 4.80 (t, J = 5.6 Hz, 1 H), 3.66 (dt, J = 5.6, 6.4 Hz, 1 H), 2.86 (t, J = 6.4 Hz, 2 H). MS (ESI) m/e $[\text{M}+1]^+$ 186.

[0505] Step 2: 8-Nitro-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide




[0506] To a solution of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (30 mg, 0.10 mmol) in DMF (5.0 mL) was added K_2CO_3 (28 mg, 0.20 mmol), followed by 2-(2-fluoro-5-nitrophenyl)ethanol (37 mg, 0.20 mmol). The mixture was warmed to 80 °C stirred for about 16 hr. After cooling down to RT, the mixture was concentrated under reduced pressure. The residue was partitioned between ethyl acetate (15 mL) and water (15 mL), the aqueous was extracted with ethyl acetate (3 × 10 mL). The combined organic phases were washed brine (10 mL), dried over Na_2SO_4 , filtered, concentrated and purified by Pre-TLC (DCM/ CH_3OH = 20/1) to afford the product about 5.0 mg (11.1%). 1H NMR (400 MHz, $DMSO-d_6$) δ 8.35 (t, J = 4.0 Hz, 1 H), 8.28 (d, J = 2.8 Hz, 1 H), 8.22 (dd, J = 2.8, 9.2 Hz, 1 H), 8.12 (d, J = 9.2 Hz, 1 H), 7.64-7.60 (m, 2 H), 7.45-7.41 (m, 2 H), 7.22-7.17 (m, 1 H), 7.14-7.09 (m, 4 H), 3.72-3.65 (m, 2 H), 3.29-3.24 (m, 2 H). MS (ESI) m/e $[M+1]^+$ 442.

[0507] Compound **71**: 8-Amino-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo [1,5-a][1,3]diazepine-3-carboxamide



[0508] To the solution of 8-nitro-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo [1,5-a][1,3]diazepine-3-carboxamide (80 mg, 0.18 mmol) in ethanol (20 mL) was added 10% w/w Pd/C (20 mg), the reaction was stirred at RT under H_2 for about 3 hr. Filtered and washed with CH_3OH (20 mL), the filtrate was concentrated under reduced pressure, the residue was purified by Pre-TLC (DCM/ CH_3OH = 20/1) to afford the product as a white solid (20 mg, 26.8%). 1H NMR (400 MHz, $DMSO-d_6$) δ 7.73-7.69 (m, 1H), 7.59-7.55 (m, 2H), 7.45-7.40 (m, 3H), 7.21-7.16 (m, 1H), 7.13-7.07 (m, 4H), 6.59-6.54 (m, 1H), 6.50-6.47 (m, 1H), 5.70-5.40 (br s, 2H), 3.64-3.59 (m, 2H), 2.99-2.94 (m, 2H). MS (ESI) m/e $[M+1]^+$ 412.


[0509] Compound **72**: 8-Acrylamido-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide




[0510] The desired compound was prepared from compound **71** and acryloyl chloride according to the procedure similar to that for compound **8**. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.25 (s, 1H), 7.95 (t, $J = 4.0$ Hz, 1H), 7.76 (d, $J = 8.8$ Hz, 1H), 7.65-7.58 (m, 4H), 7.46-7.41 (m, 2H), 7.21-7.17 (m, 1H), 7.13-7.08 (m, 4H), 6.44 (dd, $J = 10.0, 16.8$ Hz, 1H), 6.27 (dd, $J = 2.0, 16.8$ Hz, 1H), 5.77 (dd, $J = 2.0, 10.0$ Hz, 1H), 3.68-3.65 (m, 2H), 3.05-3.03 (m, 2H). MS (ESI) m/e $[\text{M}+1]^+$ 466.

[0511] **Example 26: Synthesis of compounds 73-75**

[0512] Compound **73**: 8-Nitro-5-oxo-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide




[0513] Step 1: Methyl 2-(2-fluoro-5-nitrophenyl)acetate




[0514] To a solution of 2-(2-fluoro-5-nitrophenyl)acetic acid (1.0 g, 5.0 mmol) in CH_3OH (20 mL) was added con. H_2SO_4 (0.50 mL), the reaction was warmed to 80°C and stirred for about 3 hr. After cooling down to RT, the reaction was poured into water (20 mL) and concentrated to remove CH_3OH . The aqueous was extracted with ethyl acetate (2×20 mL), the combined organic phases were washed brine (10 mL), dried over Na_2SO_4 , filtered and concentrated to afford the product about 1.0 g (93.4%) as a colorless oil. MS (ESI) m/e $[\text{M}+1]^+$ 214.0.

[0515] Step 2: 8-Nitro-5-oxo-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide




[0516] To a solution of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (30 mg, 0.10 mmol) in DMF (5 mL) was added K_2CO_3 (28 mg, 0.20 mmol) and methyl 2-(2-fluoro-5-nitrophenyl)acetate (21 mg, 0.10 mmol). The mixture was warmed to 80 °C stirred for about 16 hr. After cooling down to RT, the mixture was concentrated under reduced pressure to remove solvent. The residue was portioned with DCM (10 mL) and water (10 mL), the aqueous was extracted with DCM (2 × 10 mL), the combined organic phases were washed sat. Sodium chloride (10 mL), dried over anhydrous Sodium sulfate, filtered, concentrated and purified by pre-TLC (DCM/CH₃OH = 20/1) got the product about 10 mg (21.9%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.66 (br s, 1H), 8.51 (d, *J* = 2.6 Hz, 1H), 8.37 (dd, *J* = 2.6, 9.2 Hz, 1H), 8.07 (d, *J* = 9.2 Hz, 1H), 7.85 (d, *J* = 8.8 Hz, 2H), 7.65-7.53 (br s, 2H), 7.45-7.40 (m, 2H), 7.22-7.16 (m, 1H), 7.12-7.05 (m, 4H), 3.92 (s, 2 H). MS (ESI) *m/e* [M+1]⁺ 456.1.

[0517] Compound **74**: 8-Amino-5-oxo-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide



[0518] Compound **74** was prepared from 8-nitro-5-oxo-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide according to the procedure similar to that for compound **71**. ¹H NMR (400 MHz, CD₃OD-*d*₄) δ 7.72-7.67 (m, 2H), 7.52 (d, *J* = 8.8 Hz, 1H), 7.42-7.37 (m, 2H), 7.19-7.14 (m, 1H), 7.11-7.05 (m, 4H), 6.80 (dd, *J* = 2.6, 8.8 Hz, 1H), 6.69 (d, *J* = 2.6 Hz, 1H), 3.59 (s, 2H). MS (ESI) *m/e* [M+1]⁺ 426.1.


[0519] Compound **75**: 8-Acrylamido-5-oxo-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide



[0520] Compound **75** was prepared from 8-amino-5-oxo-2-(4-phenoxyphenyl)-5,6-dihydro-4H-benzo[f]pyrazolo[1,5-a][1,3]diazepine-3-carboxamide and acryloyl chloride according to the procedure similar to that for compound **8**. ¹H NMR (400 MHz, CD₃OD-*d*₄) δ 7.85-7.79 (m, 3H), 7.75-7.71 (m, 2H), 7.43-7.37 (m, 2H), 7.20-7.14 (m, 1H), 7.12-7.06 (m, 4H), 6.50-6.35 (m, 2H), 5.81 (dd, *J* = 2.6, 9.0 Hz, 1H), 3.75 (s, 2H). MS (ESI) *m/e* [M+1]⁺ 480.1.


[0521] **Example 27: Synthesis of compounds 76-79**

[0522] Compound **76**: 7-Nitro-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide




[0523] A mixture of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (220 mg, 0.75 mmol), methyl 2-chloro-5-nitrobenzoate (160 mg, 0.75 mmol) and K₂CO₃ (155 mg, 1.13 mmol) in DMF (10 mL) was heated to 80 °C for 16 hr under N₂. The reaction was poured into water (30 ml), and extracted with ethyl acetate (20 mL × 3). The combined organic layers were dried over Na₂SO₄, concentrated under reduced pressure to a residue, which was purified by a silica gel column eluting with 10% to 50% EA in PE to afford 85 mg (27.3%) of 7-nitro-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide as a yellow solid. MS (ESI, *m/e*) [M+1]⁺ 442.1.

[0524] Compound **77**: 7-Amino-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide




[0525] Compound **77** was prepared from 7-nitro-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide according to the procedure similar to that for compound **71**. ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.88 (br s, 1H), 7.80 (d, *J* = 8.4 Hz, 1H), 7.71 (d, *J* = 6.8 Hz, 2H), 7.36-7.42 (m, 2H), 7.27 (s, 1H), 7.02-7.17 (m, 6H), 5.62 (s, 2H). MS (ESI, *m/e*) [M+1]⁺ 412.1.

[0526] Compound **78**: 7-Acrylamido-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide



[0527] Compound **78** was prepared from 7-amino-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]quinazoline-3-carboxamide and acryloyl chloride according to the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 11.37 (s, 1H), 10.77 (s, 1H), 8.65 (s, 1H), 8.24 (d, J = 8 Hz, 1H), 8.15 (d, J = 8 Hz, 1H), 7.85 (d, J = 7.6 Hz, 2H), 7.48-7.52 (m, 2H), 7.2-7.15 (m, 5H), 6.57 (dd, J = 9.2, 18.0 Hz, 1H), 6.37 (d, J = 18.0 Hz, 1H), 5.87 (d, J = 9.2 Hz, 1H). MS (ESI, m/e) $[\text{M}+1]^+$ 466.1.


[0528] Compound **79**: 8-Amino-5-oxo-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo [1,5-a]quinazoline-3-carboxamide




[0529] The desired product was prepared from methyl 2-chloro-4-nitrobenzoate and 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide using the procedures similar to those for compound **76** and **77**. ^1H NMR (400 MHz, DMSO- d_6) δ 10.61 (s, 1H), 7.77 (d, J = 8.4 Hz, 1H), 7.71 (d, J = 8.0 Hz, 2H), 7.44-7.36 (m, 2H), 7.19-7.02 (m, 6H), 6.64 (d, J = 8.4 Hz, 1H), 6.55 (br s, 2H). MS (ESI, m/e) $[\text{M}+1]^+$ 412.1.

[0530] Example 28: Synthesis of compounds 80-81

[0531] Compound **80**: 5-Oxo-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5-dihydropyrazolo [1,5-a]pyrimidine-3-carboxamide




[0532] Step 1: tert-Butyl 4-(2,2-dimethyl-4,6-dioxo-1,3-dioxane-5-carbonyl)piperidine-1-carboxylate




[0533] To a stirred mixture of 1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid (1.15 g, 5 mmol) and DMAP (61 mg, 0.5 mmol) in DCM (50 mL) was added DCC (1.14 g, 5.5 mmol) and 2,2-dimethyl-1,3-dioxane-4,6-dione (0.8 g, 5.5 mmol). The resulting mixture was stirred at rt for 16 hr and filtered. The filtrate was concentrated under vacuum to afford tert-butyl 4-(2,2-dimethyl-4,6-dioxo-1,3-dioxane-5-carbonyl) piperidine-1-carboxylate 2 g (crude) as a yellow oil, which was used in the next step without further purification. MS (ESI) m/e $[M+23]^+$ 378.1.

[0534] Step 2: tert-Butyl 4-(3-ethoxy-3-oxopropanoyl)piperidine-1-carboxylate



[0535] A solution of tert-butyl 4-(2,2-dimethyl-4,6-dioxo-1,3-dioxane-5-carbonyl) piperidine-1-carboxylate (2 g, 5.63 mmol) in ethanol (50 ml) was refluxed for 20 h, then the solvent was removed under vacuum, and the residue was purified by silica gel chromatography eluted with DCM to afford 0.5 g (30%) of tert-butyl 4-(3-ethoxy-3-oxopropanoyl)piperidine-1-carboxylate as a reddish oil. MS (ESI) m/e $[M+23]^+$ 322.2.

[0536] Step 3: 5-Oxo-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5-dihydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0537] A mixture of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carboxamide (412 mg, 1.4 mmol) and tert-butyl 4-(3-ethoxy-3-oxopropanoyl)piperidine-1-carboxylate (420 mg, 1.4 mmol) in HOAc (20 mL) was stirred at 90 °C for 16 hr. The solvent was removed under vacuum, and the residue was partitioned between aq. NaHCO_3 and ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 and concentrated under vacuum. The residue was purified by Pre-HPLC eluting from 25% to 90% CH_3CN in 0.1% TFA in H_2O . Fractions containing the desired product were combined and lyophilized overnight to afford 5-oxo-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5-dihydropyrazolo[1,5-a]pyrimidine-3-carboxamide (0.3 g, 50%) as a white solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.71 (br s, 1H), 8.68-8.65 (m, 1H), 8.42-8.39 (m, 1H), 7.74 (d, J = 8.4 Hz, 2H), 7.47-7.41 (m, 2H), 7.22-7.18 (m, 1H), 7.14-7.09 (m, 4H), 5.72 (s, 1H),

3.50-3.35 (m, 2H), 3.17-3.06 (m, 1H), 3.01-2.87 (m, 2H), 2.15-2.05 (m, 2H), 1.83-1.72 (m, 2H).

MS (ESI) m/e $[M+1]^+$ 430.1.


[0538] Compound **81**: 5-Oxo-2-(4-phenoxyphenyl)-7-(1-propionylpiperidin-4-yl)-4,5-dihydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0539] The desired product was prepared from compound **80** and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO-d_6) δ 11.74 (s, 1H), 7.75 (d, $J = 8.4$ Hz, 2H), 7.47-7.41 (m, 2H), 7.20 (t, $J = 7.2$ Hz, 1H), 7.12-7.09 (m, 4H), 6.85 (dd, $J = 10.6, 16.6$ Hz, 1H), 6.12 (dd, $J = 2.4, 16.6$ Hz, 1H), 5.76 (s, 1H), 5.69 (dd, $J = 2.4, 10.6$ Hz, 1H), 4.63-4.58 (m, 1H), 4.24-4.20 (m, 1H), 3.15-3.05 (m, 2H), 2.69-2.63 (m, 1H), 1.99-1.91 (m, 2H), 1.61-1.58 (m, 2H). MS (ESI) m/e $[M+1]^+$ 483.9.

[0540] Example 29: Synthesis of compounds 82-83

[0541] Compound **82**: 2-Oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b] pyrazole-3,4'-piperidine]-7-carboxamide




[0542] Step 1: 1'-Benzyl-2-oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b] pyrazole-3,4'-piperidine]-7-carbonitrile




[0543] A mixture of ethyl 1-benzyl-4-hydrazinylpiperidine-4-carboxylate hydrochloride (350 mg, 1.0 mmol), 2-(methoxy(4-phenoxyphenyl)methylene)malononitrile (276 mg, 1.0 mmol) and K_2CO_3 (414 mg, 3.0 mmol) in MeOH (20 mL) was heated to reflux for 16 hr. The mixture was filtered and the filtrate was concentrated to give the crude product (280 mg, 58.9%) as a yellow solid. MS (ESI, m/e) $[M+1]^+$ 475.9.

[0544] Step 2: 1'-Benzyl-2-oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide




[0545] A solution of 1'-benzyl-2-oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide (200 mg, 0.42 mmol) in H_3PO_4 (15 mL) was heated to 120°C for 2 hr. The solution was poured to water (10 mL) and extracted with EA (10 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 and concentrated to give the crude product (120 mg, 58.0%) as an off-white solid. MS (ESI, m/e) $[\text{M}+1]^+$ 493.9.

[0546] Step 3: 2-Oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide



[0547] To a solution of 1'-benzyl-2-oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide (120 mg, 0.24 mmol) in MeOH (10 mL) was added 10% w/w $\text{Pd}(\text{OH})_2/\text{C}$ (5 mg) and stirred for 16 hr under H_2 . The mixture was filtered and the filtrate was concentrated to give the crude product (280 mg, 58.9%) as a yellow solid. MS (ESI, m/e) $[\text{M}+1]^+$ 403.9.

[0548] Compound **83**: 1'-Acryloyl-2-oxo-6-(4-phenoxyphenyl)-1,2-dihydrospiro[imidazo[1,2-b]pyrazole-3,4'-piperidine]-7-carboxamide




[0549] The desired product was prepared from compound **82** and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO-d_6) δ 11.91 (s, 1H), 7.71 (d, J = 8.4 Hz, 2H), 7.47-7.37 (m, 2H), 7.25 (br s, 1H), 7.17 (t, J = 7.6 Hz, 1H), 7.08-7.02 (m, 4H), 6.88 (dd, J = 16.6, 10.4 Hz, 1H), 6.80 (br s, 1H), 6.16 (d, J = 16.6 Hz, 1H), 5.72 (d, J =

10.4 Hz, 1H), 4.32-4.14 (m, 1H), 4.12-3.99 (m, 1H), 3.97-3.81 (m, 1H), 3.76-3.60 (m, 1H), 1.86-1.91 (m, 4H). MS (ESI, m/e) $[M+1]^+$ 457.9.


[0550] Example 30: Synthesis of compounds 84-85

[0551] Compound 84: 6-Amino-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0552]

S Step 1: 6-Nitro-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0553] To a solution of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (83 mg, 0.3 mmol) in HOAc (2 mL) was added sodium 2-nitro-1,3-dioxopropan-2-ide (47 mg, 0.3 mmol). After stirring at RT for 1 hr, water (2 mL) was added. The mixture was partitioned between EA (25 mL) and brine (25 mL). The combined organic layers were washed with brine (25 mL $\times$ 2), dried over Na_2SO_4 and concentrated to afford 90 mg of 6-nitro-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (84%) as a yellow solid. MS (ESI) m/e $[M+1]^+$ 358.2.


[0554] Step 2: 6-Nitro-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0555] To a solution of 6-nitro-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (90 mg, 0.25 mmol) in 2 mL of ethanol and 2 mL of DCM was added NaBH_4 (19 mg, 0.5 mmol) at RT. After stirring at RT for 30 min, 5 mL of water was added. The mixture was concentrated. The residue was partitioned between 50 mL of DCM and 50 mL of brine. The combined organic


layers were washed with brine (50 mL $\times$ 2), dried over Na_2SO_4 and concentrated to afford 50 mg of 6-nitro-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (55%) as a yellow solid. MS (ESI) m/e $[\text{M}+1]^+$ 362.1.

[0556] Step 3: 6-Amino-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0557] To a solution of 6-nitro-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile (600 mg, 1.67 mmol) in 30 mL of methanol and 10 mL of DCM was added 10% w/w Pd/C (100 mg). The mixture was stirred at RT under H_2 for 2 hr and filtered. The filtrate was concentrated and purified by chromatography column on silica gel eluting with PE/EA to afford 200 mg (36%) of 6-amino-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile as a white solid. MS (ESI) m/e $[\text{M}+1]^+$ 332.1.

[0558] Step 4: 6-Amino-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0559] The desired product was prepared from 6-amino-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound **2**. ^1H NMR (400 MHz, DMSO-d_6) δ 7.53-7.48 (m, 2H), 7.46-7.40 (m, 2H), 7.22-7.16 (m, 1H), 7.11-7.03 (m, 4H), 6.58 (br s, 1H), 4.15-4.08 (m, 1H), 3.72-3.67 (m, 1H), 3.40-3.30 (m, 2H) and 3.06-2.98 (m, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 350.2.


[0560] Compound **85**: 6-Acrylamido-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0561] The desired product was prepared from compound **84** and acryloyl chloride using the procedure similar to that for compound **8**. ^1H NMR (400 MHz, DMSO- d_6) δ 8.47 (d, J = 7.2 Hz, 1H), 7.52 (d, J = 8.8 Hz, 2H), 7.46-7.38 (m, 2H), 7.18 (dd, J = 7.2, 7.6 Hz, 1H), 7.09 (d, J = 8.0 Hz, 2H), 7.05 (d, J = 8.8 Hz, 2H), 6.67 (br s, 1H), 6.34 (dd, J = 10.0, 17.2 Hz, 1H), 6.15 (dd, J = 2.0, 17.2 Hz, 1H), 5.63 (dd, J = 2.0, 10.0 Hz, 1H), 4.32-4.40 (m, 1H), 4.22 (dd, J = 4.8, 12.4 Hz, 1H), 3.91 (dd, J = 4.8, 12.4 Hz, 1H), 3.40 (m, 1H) and 3.26 (dd, J = 5.2 Hz, J = 12.0 Hz, 1H). MS (ESI) m/e $[\text{M}+1]^+$ 404.1.

[0562] Example 31: Synthesis of compound 86

[0563] Compound **86**: 6-(Acrylamidomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide




[0564] Step 1: Ethyl 3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-6-carboxylate




[0565] To a solution of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (276 mg, 1.0 mmol) in EtOH (10 mL) was added ethyl 2-formyl-3-oxopropanoate (144 mg, 1.0 mmol) and HOAc (5 drops). After stirring at RT for 16 hr, the mixture was filtered. The cake was washed with H_2O (10 mL $\times$ 2) and dried to afford 250 mg (65%) of ethyl 3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-6-carboxylate as a yellow solid. MS (ESI) m/e $[\text{M}+1]^+$ 384.9.

[0566] Step 2: 6-(Hydroxymethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0567] To a solution of ethyl 3-cyano-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-6-carboxylate (250 mg, 0.65 mmol) in DCM (5 mL) and CH₃OH (5 mL) was added NaBH₄ (250 mg, 6.5 mmol). After stirring at RT for 16 hr, the mixture was partitioned between DCM/CH₃OH (100 mL/5 mL) and brine (100 mL). The organic layer was separated from aqueous layers, dried over Na₂SO₄ and concentrated to afford 250 mg (100%) of 6-(hydroxymethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carbonitrile. MS (ESI) m/e [M+1]⁺ 346.9.

[0568] Step 3: 6-(Hydroxymethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0569] The desired product was prepared from 6-(hydroxymethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound **2**. MS (ESI) m/e [M+1]⁺ 364.9.

[0570] Step 4: 6-((1,3-Dioxoisindolin-2-yl)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro pyrazolo[1,5-a]pyrimidine-3-carboxamide




[0571] To a solution of isoindoline-1,3-dione (74 mg, 0.5 mmol) in THF (20 mL) was added PPh₃ (393 mg, 1.5 mmol) and 6-(hydroxymethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (180 mg, 0.5 mmol). DIAD (253 mg, 1.25 mmol) was added dropwise at 0 °C and stirred for 10 min. The mixture was allowed to warm to rt and stirred for 16 hr. Concentrated and purified by chromatography column on 5 g of silica gel eluting with DCM/CH₃OH to afford 200 mg (62%) of 6-((1,3-dioxoisindolin-2-yl)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide as a yellow solid. MS (ESI) m/e [M+1]⁺ 493.9.

[0572] Step 5: 6-(Aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0573] To a solution of 6-((1,3-dioxoisindolin-2-yl)methyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (200 mg, 0.40 mmol) in CH₃OH (5 mL) was added hydrazine hydrate (1 mL, 80% of aqueous solution). The mixture was stirred at 70 °C under N₂ for 4 hr, concentrated and purified by chromatography column on 5 g of silica gel eluting with DCM/CH₃OH to afford 63 mg (43%) of 6-(aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide as colorless oil. MS (ESI) m/e [M+1]⁺ 363.9.


[0574] Step 6: 6-(Acrylamidomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide




[0575] The desired product was prepared from 6-(aminomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a] pyrimidine-3-carboxamide and acryloyl chloride using the procedure similar to that for compound **8**. ¹H NMR (400 MHz, DMSO-d₆) δ 8.31 (t, *J* = 5.6 Hz, 1H), 7.51 (d, *J* = 8.8 Hz, 2H), 7.46-7.38 (m, 2H), 7.17 (t, *J* = 7.6 Hz, 1H), 7.08 (d, *J* = 7.6 Hz, 2H), 7.04 (d, *J* = 8.8 Hz, 2H), 6.61 (s, 1H), 6.25 (dd, *J* = 17.1, 10.1 Hz, 1H), 6.11 (dd, *J* = 17.1, 2.2 Hz, 1H), 5.62 (dd, *J* = 10.1, 2.2 Hz, 1H), 4.07 (dd, *J* = 12.4, 6.0 Hz, 1H), 3.73 (dd, *J* = 12.4, 8.0 Hz, 1H), 3.41-3.34 (m, 1H), 3.27-3.21 (m, 2H), 3.09-2.96 (m, 1H), 2.37-2.24 (m, 1H). MS (ESI) m/e [M+1]⁺ 417.9.

[0576] Example 32: Synthesis of compounds 87-88

[0577] Compound **87**: 2'-(4-Phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine-3,6'-pyrazolo [1,5-a]pyrimidine]-3'-carboxamide




[0578] Step 1: Diethyl 1-benzylazetidine-3,3-dicarboxylate




[0579] To a solution of diethyl 2,2-bis(hydroxymethyl)malonate (4.4 g, 20 mmol) in CH_3CN (50 mL) was added Tf_2O (7.1 mL, 11.85 g, 42 mmol) at -20°C , followed by two batches of DIEA (6.45 g, 50 mmol). After 0.5 hr, benzylamine (3.21 g, 35 mmol) was added at -20°C . The mixture was stirred at 70°C for 2 hr. 100 mL of EA and 100 mL of brine were added. Organic layers were dried over Na_2SO_4 . Purified by chromatography column on silica gel eluting with PE/EA to afford 4.8 g (82%) of diethyl 1-benzylazetidine-3,3-dicarboxylate as a yellow oil. ^1H NMR (400 MHz, DMSO-d_6) δ 7.29-7.34 (m, 2H), 7.22-7.27 (m, 3H), 4.17 (q, $J = 7.2$ Hz, 4H), 3.56 (s, 2H), 3.51 (s, 4H), 2.39 (s, 3H) and 1.17 (t, $J = 7.2$ Hz, 6H).

[0580] Step 2: (1-Benzylazetidine-3,3-diyl)dimethanol



[0581] To a solution of diethyl 1-benzylazetidine-3,3-dicarboxylate (4.8 g, 16.5 mmol) in CH_3OH (10 mL) was added NaBH_4 (1.25 g, 33 mmol). The mixture was stirred at RT for 1 hr. 100 mL of brine and 200 mL of DCM were added. Organic layers were separated from aqueous layers, dried over Na_2SO_4 , concentrated to afford 2.328 g (68%) of (1-benzylazetidine-3,3-diyl)dimethanol as a yellow oil. ^1H NMR (400 MHz, DMSO-d_6) δ 7.21-7.31 (m, 5 H), 4.15-4.05 (m, 2 H), 3.48 (d, $J = 4.8$ Hz, 2H), 3.17 (d, $J = 4.8$ Hz, 4 H) and 2.89 (s, 2 H).


[0582] Step 3: (1-Benzylazetidine-3,3-diyl)bis(methylene)dimethanesulfonate



[0583] To a solution of (1-benzylazetidine-3,3-diyl)dimethanol (50 mg, 0.24 mmol) in DCM (10 mL) was added TEA (222 mg, 2.2 mmol) and MsCl (249 mg, 2.2 mmol). After stirring for at RT for 4 hr, the mixture was concentrated. The residue was partitioned between brine (100 mL) and EA (100 mL). The organic layer was washed with brine (100 mL $\times$ 2), dried over Na_2SO_4


and concentrated to afford 300 mg (83%) of crude (1-benzylazetidine-3,3-diyl)bis(methylene)dimethanesulfonate as a yellow oil. MS (ESI) m/e $[M+1]^+$ 363.9.

[0584] Step 4: 1-Benzyl-2'-(4-phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine-3,6'-pyrazolo[1,5-a]pyrimidine]-3'-carbonitrile




[0585] To a solution of (1-benzylazetidine-3,3-diyl)bis(methylene)dimethanesulfonate (300 mg, 0.83 mmol) in DMF (10 mL) was added 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (230 mg, 0.83 mmol) and K_2CO_3 (230 mg, 1.66 mmol). The mixture was stirred at 80°C under N_2 for 16 hr. The mixture was concentrated. The residue was washed with H_2O (100 mL $\times$ 2), dried and purified by pre-TLC (DCM/ CH_3OH = 10/1) to afford 30 mg (10%) of desired product as a yellow liquid. MS (ESI) m/e $[M+1]^+$ 447.9.

[0586] Step 5: 1-Benzyl-2'-(4-phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine-3,6'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide




[0587] The desired product was prepared from 1-benzyl-2'-(4-phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine-3,6'-pyrazolo[1,5-a]pyrimidine]-3'-carbonitrile using the procedure similar to step 2 for compound **2**. MS (ESI) m/e $[M+1]^+$ 465.9.

[0588] Step 6: 2'-(4-Phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine-3,6'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide



[0589] To a solution of 1-benzyl-2'-(4-phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine- 3,6'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide (200 mg, 0.43 mmol) in 10 mL of DCM and 10 mL of CH₃OH was added 10% w/w Pd/C (100 mg). After stirring at RT under H₂ for 16 hr, the mixture was filtered and concentrated. The residue was purified by pre-HPLC eluting from 25% to 90% CH₃CN in 0.1% TFA in H₂O. Fractions containing the desired product were combined and lyophilized overnight to afford 30 mg (19%) of desired product as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 8.94 (br s, 1H), 8.84 (br s, 1H), 7.50 (d, *J* = 8.4 Hz, 2H), 7.46-7.38 (m, 2H), 7.18 (t, *J* = 8.0 Hz, 1H), 7.08 (d, *J* = 8.0 Hz, 2H), 7.05 (d, *J* = 8.4 Hz, 2H), 6.81 (br s, 1 H), 4.23 (s, 2 H), 3.90-4.00 (m, 2 H), 3.78-3.87 (m, 2 H) and 3.47 (s, 2H). MS (ESI) *m/e* [M+1]⁺ 375.9.


[0590] Compound **88**: 1-Acryloyl-2'-(4-phenoxyphenyl)-5',7'-dihydro-4'H-spiro[azetidine- 3,6'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide




[0591] The desired product was prepared from compound **87** and acryloyl chloride using the procedure similar to that for compound **8**. ¹H NMR (400 MHz, CD₃OD-d₄) δ 7.49 (d, *J* = 7.6 Hz, 2H), 7.41-7.32 (m, 2H), 7.14 (t, *J* = 8.0 Hz, 1H), 7.05 (d, *J* = 8.0 Hz, 2H), 7.04 (d, *J* = 7.6 Hz, 2H), 6.35 (dd, *J* = 16.8, 10.1 Hz, 1H), 6.25 (dd, *J* = 16.8, 1.6 Hz, 1H), 5.74 (dd, *J* = 10.1, 1.6 Hz, 1H), 4.20-4.27 (m, 4 H), 3.92-3.98 (m, 2 H) and 3.54 (s, 2 H). MS (ESI) *m/e* [M+1]⁺ 429.9.

[0592] Example 33: Synthesis of compounds 89-90

[0593] Compound **89**: 6-(2-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide




[0594] Step 1: 6-Bromo-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0595] A mixture of 5-amino-3-(4-phenoxyphenyl)-1H-pyrazole-4-carbonitrile (28 mg, 0.1 mmol), 2-bromomalonaldehyde (15 mg, 0.1 mmol) in EtOH (5 mL) was stirred at RT for 2 hr. Then, the mixture was filtered to give the crude product (20 mg, 62.9%) as a yellow solid. ^1H NMR (400 MHz, DMSO- d_6) δ 9.89 (d, J = 2.0 Hz, 1H), 8.95 (d, J = 2.0 Hz, 1H), 8.08 (d, J = 8.4 Hz, 2H), 7.52-7.43 (m, 2H), 7.27-7.19 (m, 3H), 7.15 (d, J = 7.6 Hz, 2H). MS (ESI, m/e) $[\text{M}+1]^+$ 391.9.

[0596] Step 2: 6-(2-Aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0597] A mixture of 6-bromo-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (500 mg, 1.28 mmol), 2-aminophenylboronic acid (175 mg, 1.28 mmol), Cs_2CO_3 (623 mg, 1.92 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (74 mg, 0.06 mmol) in 1,4-dioxane (30 mL) and water (1.0 mL) was heated to 80 $^\circ\text{C}$ for 16 hr under N_2 . The mixture was filtered and the filtrate was concentrated and purified by chromatography column on silica gel using 50% of EA in PE as eluant to give the crude product (320 mg, 59.1%) as a yellow solid. MS (ESI, m/e) $[\text{M}+1]^+$ 403.9.


[0598] Step 3: 6-(2-Aminophenyl)-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]pyrimidine-3-carbonitrile



[0599] To a solution of 6-(2-aminophenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (320 mg, 0.79 mmol) in MeOH (20 mL) was added NaBH_4 (86 mg, 2.28 mmol). The solution was stirred at rt for 30 min then was poured into water (50 mL) and extracted with


EA (50 mL $\times$ 3). The organic combined layers were dried over Na_2SO_4 and concentrated to give the crude product (240 mg, 75%) as a yellow solid. MS (ESI, m/e) $[\text{M}+1]^+$ 406.0.

[0600] Step 4: 6-(2-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0601] The desired product was prepared from 6-(2-aminophenyl)-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 8 for compound **68**. MS (ESI, m/e) $[\text{M}+1]^+$ 407.9.

[0602] Step 5: 6-(2-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0603] The desired product was prepared from 6-(2-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound **2**. ^1H NMR (400 MHz, DMSO-d_6) δ 7.58 (d, $J = 8.4$ Hz, 2H), 7.53-7.44 (m, 2H), 7.24 (t, $J = 7.6$ Hz, 1H), 7.19-7.09 (m, 4H), 7.06-7.99 (m, 2H), 6.84 (d, $J = 1.6$ Hz, 1H), 6.74 (d, $J = 7.2$ Hz, 1H), 6.61 (t, $J = 7.6$ Hz, 1H), 5.20 (s, 2H), 4.24 (dd, $J = 4.0, 12.0$, Hz, 1H), 4.07 (dd, $J = 12.0, 12.0$ Hz, 1H), 3.56-3.41 (m, 3H). MS (ESI, m/e) $[\text{M}+1]^+$ 425.9.

[0604] Compound **90**: 6-(3-Aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0605] The desired product was prepared from 6-bromo-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile and 3-aminophenylboronic acid according to the similar procedures


(step 2 to 5) for compound **89** under appropriate conditions recognized by one of ordinary skill in the art. ^1H NMR (400 MHz, DMSO- d_6) δ 7.59 (d, J = 8.4 Hz, 2H), 7.52-7.45 (m, 2H), 7.24 (t, J = 7.6 Hz, 1H), 7.15 (d, J = 8.4 Hz, 2H), 7.12 (d, J = 7.6 Hz, 2H), 7.07 (t, J = 7.6 Hz, 1H), 6.85 (br s, 1H), 6.62-6.54 (m, 3H), 5.23 (br s, 2H), 4.24 (dd, J = 12.0, 4.8 Hz, 1H), 4.09 (t, J = 12.0 Hz, 1H), 3.56-3.50 (m, 1H), 3.40-3.35 (m, 1H), 3.28-3.19 (m, 1H). MS (ESI, m/e) $[\text{M}+1]^+$ 425.9.

[0606] Example 34: Synthesis of compound 91

[0607] Compound 91: 6-(3-(2-(Dimethylamino)ethoxy)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0608] Step 1: 6-(3-Hydroxyphenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0609] To a solution of 6-bromo-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile (782 mg, 2.0 mmol) in dioxane (10 mL) and H₂O (10 mL) was added 3-hydroxyphenylboronic acid (276 mg, 2.0 mmol), Pd(PPh₃)₄ (240 mg, 0.2 mmol) and Na₂CO₃ (424 mg, 4.0 mmol). After stirring at 65 °C under N₂ for 16 hr, the mixture was concentrated and 100 mL of DCM, 10 mL of CH₃OH, 100 mL of H₂O were added. Organic layers were separated from aqueous layers and dried over Na₂SO₄ and purified by chromatography column on silica gel eluting with DCM/CH₃OH to afford 500 mg (62%) of 6-(3-hydroxyphenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile as a yellow solid. MS (ESI) m/e $[\text{M}+1]^+$ 404.9.

[0610] Step 2: 6-(3-Hydroxyphenyl)-2-(4-phenoxyphenyl)-6,7-dihydropyrazolo[1,5-a]pyrimidine-3-carbonitrile and 6-(3-hydroxyphenyl)-2-(4-phenoxyphenyl)-4,5-dihydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0611] The desired product was prepared from 6-(3-hydroxyphenyl)-2-(4-phenoxyphenyl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 3 for compound **89**. MS (ESI) m/e $[M+1]^+$ 406.9.

[0612] Step 3: 6-(3-Hydroxyphenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile




[0613] The desired product was prepared from intermediate in the last step using the procedure similar to step 4 for compound **89**. MS (ESI) m/e $[M+1]^+$ 408.9.

[0614] Step 4: 6-(3-Hydroxyphenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0615] The desired product was prepared from 6-(3-hydroxyphenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carbonitrile using the procedure similar to step 2 for compound **2**. MS (ESI) m/e $[M+1]^+$ 426.9.

[0616] Step 5: 6-(3-(2-(Dimethylamino)ethoxy)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0617] The desired product was prepared from 6-(3-hydroxyphenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a] pyrimidine-3-carboxamide and 2-chloro-*N,N*-dimethylethanamine hydrochloride using the procedure similar to step 7 for compound **9**. ^1H NMR (400 MHz, DMSO- d_6) δ 7.52 (d, J = 8.4 Hz, 2H), 7.46-7.38 (m, 2H), 7.29 (t, J = 7.6 Hz, 1H), 7.18 (t, J = 7.6 Hz, 1H), 7.10-7.05 (m, 4H), 6.98-6.96 (m, 2H), 6.90-6.88 (m, 1H), 6.81 (d, J = 2.4 Hz, 1H), 4.25-4.20 (m, 1H), 4.16-4.11 (m, 3H), 3.50-3.47 (m, 1H), 3.43-3.38 (m, 2H), 2.96-2.84 (m, 2H) and 2.43 (s, 6H). MS (ESI) m/e $[\text{M}+1]^+$ 497.9.


[0618] Compound **178**: *N*1-(2-(4-((*E*)-4-(4-((*S*)-3-carbamoyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)piperidin-1-yl)-4-oxobut-2-en-1-yl)piperazin-1-yl)ethyl)-*N*5-(15-oxo-19-((3*aR*,4*R*,6*aS*)-2-oxohexahydro-1*H*-thieno[3,4-*d*]imidazol-4-yl)-4,7,10-trioxa-14-azanonadecyl)glutaramide trifluoroacetate



[0619] The desired compound was prepared according to the scheme, step and intermediates described below.




[0620] Step 1: (*E*)-4-(4-(2-((*tert*-butoxycarbonyl)amino)ethyl)piperazin-1-yl)but-2-enoic acid



[0621] A mixture of (*E*)-4-bromobut-2-enoic acid (500 mg, 3.03 mmol), *tert*-butyl (2-(piperazin-1-yl)ethyl)carbamate (694 mg, 3.03 mmol) and Et_3N (612 mg, 6.06 mmol) in 20 mL of THF was stirred at RT for 15 h. The mixture was concentrated and used in the next step without further purification. MS (ESI) m/e $[\text{M}+1]^+$ 314.0.

[0622] Step 2: (*S*)-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide




[0623] A solution of 2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo [1,5-a]pyrimidine-3-carboxamide (2.0 g, 4.8 mmol) in MeOH/H₂O = 3/1 (110 mL) was warmed to 50 °C and stirred for about 20 min until all starting material dissolved, then to solution was added a solution of L-DBTA (600 mg, 1.6 mmol) in MeOH/H₂O = 3/1 (10 mL), the solution was stirred at 50 °C for about 30 min, then slowly cooled to 40 °C (about 2 h). To the solution was added crystal seed (10 mg). The mixture was stirred at 40 °C for 2 h, then slowly cooled to ambient temperature and stirred for about 48 h. Filtered, the solid was washed with MeOH/H₂O = 3/1 (5 mL), dried under reduced pressure to give the product as a white solid about 1.1 g (38% yield, 93% ee value). The solid (500 mg) was added to the solvent of THF/H₂O = 1/1 (20 mL), the solution was warmed to 70 °C and stirred for about 1 h until all solid dissolved, then slowly cooled to 40 °C (3 h) and added crystal seed (10 mg), after stirring for about 2 h, the solution was slowly cooled to ambient temperature and stirred for about 48 h. Filtered, solid was washed with water (4 mL), dried under reduced pressure to give the product as a white solid about 330 mg (65% yield, >99.5% ee value) as its L-DBTA salt. Suitable single crystal of this L-DBTA salt was obtained by slow cooling in MeOH/H₂O (1:1, v/v). Configuration of chiral carbon in freebase was determined to be *S*. The DBTA salt was converted to the free base by using aqueous NaOH solution and extracting with DCM.

[0624] The chiral analysis conditions for the chiral resolution are shown below.


Column	CHIRALPAK IC
Column size	0.46 cm I.D.× 15 cm L, 5 μm
Injection	2 μL
Mobile phase	n-Hexane/EtOH(0.1% triethylamine) = 50/50 (v/v)
Flow rate	1.0 mL/min
Wave length	UV 214, 254 nm

[0625] Step 3: (*S,E*)-tert-butyl (2-(4-(4-(4-(3-carbamoyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)piperidin-1-yl)-4-oxobut-2-en-1-yl)piperazin-1-yl)ethyl)carbamate




[0626] A mixture of (*S*)-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (1.26 g, 3.03 mmol), (*E*)-4-(4-(2-((tert-butoxycarbonyl)amino)ethyl)piperazin-1-yl)but-2-enoic acid (948.4 mg, 3.03 mmol), HATU (1.21 g, 3.18 mmol), DIEA (782 mg, 6.06 mmol) in 30 mL of DMF was stirred at RT for 15 hr. The mixture was poured into 300 mL of water and extracted with EA (100 mL). The organic phase was washed with water (100 mL $\times$ 3) and concentrated to give 1.25 g (58%) of product as a yellow solid. ^1H NMR (400 MHz, DMSO- d_6) δ 7.50 (d, J = 8.4 Hz, 2H), 7.42 (t, J = 8.4 Hz, 2H), 7.17 (t, J = 7.4 Hz, 1H), 7.12-7.02 (m, 4H), 6.67 (br s, 1H), 6.64-6.53 (m, 3H), 4.53-4.40 (m, 1H), 4.14-4.05 (m, 1H), 3.32-3.25 (m, 2H), 3.10-2.91 (m, 5H), 2.45-2.15 (m, 11H), 2.10-2.00 (m, 1H), 1.96-1.84 (m, 1H), 1.78-1.65 (m, 1H), 1.62-1.50 (m, 1H), 1.37 (s, 9H), 1.31-1.10 (m, 3H). MS (ESI) m/e $[M+1]^+$ 713.0.

[0627] Step 4: (*S,E*)-7-(1-(4-(4-(2-aminoethyl)piperazin-1-yl)but-2-enoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate




[0628] To a solution of (*S,E*)-tert-butyl (2-(4-(4-(4-(3-carbamoyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)piperidin-1-yl)-4-oxobut-2-en-1-yl)piperazin-1-yl)ethyl)carbamate (150 mg, 0.21 mmol) in 10 mL of DCM was added 2 mL of TFA. The reaction mixture was stirred at RT for 15 hr and concentrated to remove the solvent. The residue was used in the next step without further purification. MS (ESI) m/e $[M+1]^+$ 613.0.

[0629] Compound **178**: *N*1-(2-(4-((*E*)-4-(4-((*S*)-3-carbamoyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidin-7-yl)piperidin-1-yl)-4-oxobut-2-en-1-yl)piperazin-1-yl)ethyl)-*N*5-(15-oxo-19-((3*a*R,4*R*,6*a*S)-2-oxohexahydro-1*H*-thieno[3,4-*d*]imidazol-4-yl)-4,7,10-trioxa-14-azanonadecyl)glutaramide trifluoroacetate




[0630] A mixture of (*S,E*)-7-(1-(4-(4-(2-aminoethyl)piperazin-1-yl)but-2-enoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate (129 mg, 0.21 mmol, crude), N-BIOTINYL-NH-(PEG)2-COOH-DIEA (118 mg, 0.21 mmol), HATU (80 mg, 0.21 mmol), TEA (63.6 mg, 0.63 mmol) in 5 mL of DMF was stirred at 40 °C for 15 hr. The mixture was concentrated and purified by Pre-HPLC to afford 160 mg (60%) of product as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.04 (t, *J* = 5.2 Hz, 1H), 7.80 (t, *J* = 5.6 Hz, 1H), 7.76 (t, *J* = 5.6 Hz, 1H), 7.50 (d, *J* = 8.5 Hz, 2H), 7.42 (t, *J* = 8.5 Hz, 2H), 7.18 (t, *J* = 7.6 Hz, 1H), 7.12-7.03 (m, 4H), 6.89-6.77 (m, 1H), 6.62-6.51 (m, 1H), 6.44 (s, 1H), 4.55-4.40 (m, 1H), 4.35-4.27 (m, 1H), 4.17-3.95 (m, 3H), 3.74-3.60 (m, 2H), 3.55-3.44 (m, 9H), 3.41-3.27 (m, 10H), 3.14-3.02 (m, 9H), 2.82 (dd, *J* = 12.4, 5.0 Hz, 1H), 2.65-2.53 (m, 3H), 2.36-2.18 (m, 1H), 2.13-2.00 (m, 7H), 1.96-1.85 (m, 1H), 1.80-1.67 (m, 3H), 1.66-1.55 (m, 6H), 1.55-1.40 (m, 3H), 1.38-1.21 (m, 4H). MS (ESI) *m/e* [M+1]⁺ 1155.0, [M+23]⁺ 1176.9.

[0631] Compound **181**: (*S*)-7-(1-(2-cyano-3-cyclopropylacryloyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide



[0632] A mixture of compound **180** (60 mg, 0.124 mmol), cyclopropanecarbaldehyde (43.4 mg, 0.62 mmol) and piperidine (52.7 mg, 0.62 mmol) in MeOH (10 mL) was stirred at RT for 15 h. After concentration, to the residue was added EA (50 mL) and water (50 mL). The organic phase was concentrated and purified by chromatography column on silica gel eluting with DCM/MeOH (50/1) to afford 30 mg (45%) of desired compound as a white solid. MS (ESI) *m/e* [M+1]⁺ 537.0.

[0633] Compound **182**: (*S*)-7-(1-cyanopiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide














[0634] To a solution of (S)-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (417 mg, 1 mmol) in DCM (20 mL) was added NaHCO₃ (168 mg, 2 mmol) and water (5 mL), followed by BrCN (127 mg, 1.2 mmol). The mixture was stirred at RT for 16 h. To the mixture was added DCM (50 mL) and brine (20 mL). The organic phase was further washed with brine (100 mL), dried over Na₂SO₄. Concentrated and purified by Pre-TLC (DCM/MeOH, 50/1) to afford 330 mg (75%) of white solid. MS (ESI) m/e [M+1]⁺ 443.0.






[0635] A variety of other compounds have been prepared by methods substantially similar to those of above described Examples. The characterization data for some of these compounds are summarized in Table 1 below and include LC/MS (observed), chiral HPLC and ¹H NMR data.






Table 1. Characterization Data for Selected Compounds






No.	Name	¹ H NMR data, Chiral HPLC and LC/MS m/z (M+1)	Structure
92	(E)-7-(2-(4-(dimethylamino)but-2-enamido)phenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 9.74 (s, 1H), 7.52-7.44 (m, 3H), 7.44-7.36 (m, 2H), 7.29 (t, <i>J</i> = 7.6 Hz, 1H), 7.21-7.04 (m, 2H), 7.11-6.97 (m, 4H), 6.88-6.67 (m, 2H), 6.64 (d, <i>J</i> = 7.6 Hz, 1H), 6.37 (d, <i>J</i> = 15.1 Hz, 1H), 5.81-5.74 (m, 1H), 3.32-3.22 (m, 1H), 3.07 (d, <i>J</i> = 5.6 Hz, 2H), 3.03-2.93 (m, 1H), 2.38-2.26 (m, 1H), 2.19 (s, 6H), 2.05-1.95 (m, 1H). MS (ESI) m/e [M+1] ⁺ 536.9.	
93	7-(2-Aminophenyl)-2-(4-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.40 (d, <i>J</i> = 8.8 Hz, 2H), 6.98 (d, <i>J</i> = 8.8 Hz, 2H), 6.94 (d, <i>J</i> = 8.0 Hz, 1H), 6.75 (s, 1H), 6.67 (d, <i>J</i> = 8.0 Hz, 1H), 6.49 (t, <i>J</i> = 7.6 Hz, 1H), 6.27 (d, <i>J</i> = 7.6 Hz, 1H), 5.57 (s, 1H), 5.16 (s, 2H), 3.78 (s, 3H), 3.28-3.21 (m, 1H), 2.99-2.91 (m, 1H), 2.26-2.04 (m, 2H). MS (ESI) m/e [M+1] ⁺ 363.9.	






94	7-(2-acrylamidophenyl)-2-(4-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 9.84 (s, 1H), 7.45 (d, <i>J</i> = 8.0 Hz, 1H), 7.40 (d, <i>J</i> = 8.8 Hz, 2H), 7.30 (t, <i>J</i> = 7.6 Hz, 1H), 7.21 (t, <i>J</i> = 7.6 Hz, 1H), 6.98 (d, <i>J</i> = 8.8 Hz, 2H), 6.82 (s, 1H), 6.64 (d, <i>J</i> = 8.0 Hz, 1H), 6.53 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 6.27 (dd, <i>J</i> = 17.0, 1.8 Hz, 1H), 5.82-5.74 (m, 2H), 3.77 (s, 3H), 3.31-3.19 (m, 1H), 3.03-2.91 (m, 1H), 2.38-2.24 (m, 1H), 2.02-1.92 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 417.9.	
95	7-(5-amino-2-chlorophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.52-7.44 (m, 2H), 7.40-7.31 (m, 2H), 7.19-7.06 (m, 2H), 7.06-6.98 (m, 4H), 6.58 (dd, <i>J</i> = 8.4, 2.6 Hz, 1H), 6.10 (d, <i>J</i> = 2.6 Hz, 1H), 5.71 (dd, <i>J</i> = 2.0, 5.2 Hz, 1H), 3.37 (dt, <i>J</i> = 12.4, 3.6 Hz, 1H), 3.21 (td, <i>J</i> = 12.4, 3.6 Hz, 1H), 2.50-2.40 (m, 1H), 2.30-2.23 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 460.1.	
96	7-(5-acrylamido-2-chlorophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 10.32 (s, 1H), 8.14 (s, 1H), 7.88 (dd, <i>J</i> = 8.4, 2.0 Hz, 1H), 7.51-7.46 (m, 3H), 7.40 (t, 2H, <i>J</i> = 7.6, 7.6 Hz), 7.16 (t, <i>J</i> = 7.6 Hz, 1H), 7.07-7.01 (m, 5H), 6.95 (br s, 1H), 6.37 (dd, <i>J</i> = 17.0, 10.0 Hz, 1H), 6.23 (dd, <i>J</i> = 17.0, 1.4 Hz, 1H), 5.74 (dd, <i>J</i> = 10.0, 1.4 Hz, 1H), 5.68-5.64 (m, 1H), 3.12-3.02 (m, 1H), 2.46-2.39 (m, 2H), 2.10-2.05 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 513.9.	
97	7-(3-acrylamido-4-chlorophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 9.77 (s, 1H), 7.57 (s, 1H), 7.53-7.46 (m, 3H), 7.44-7.35 (m, 2H), 7.16 (t, <i>J</i> = 7.6 Hz, 1H), 7.08-7.00 (m, 4H), 6.91 (d, <i>J</i> = 7.6 Hz, 1H), 6.83 (br s, 1H), 6.59 (dd, <i>J</i> = 17.1, 10.0 Hz, 1H), 6.27 (d, <i>J</i> = 17.1 Hz, 1H), 5.78 (d, <i>J</i> = 10.0 Hz, 1H), 5.50-5.45 (m, 1H), 3.33 (m, 1H), 3.13-3.03 (m, 1H), 2.42-2.31 (m, 1H), 2.14-2.04 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 513.9.	
98	7-(4-aminophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.50 (d, <i>J</i> = 8.4 Hz, 2H), 7.41-7.32 (m, 2H), 7.11 (t, <i>J</i> = 7.6 Hz, 1H), 7.03-6.98 (m, 4H), 6.69-6.67 (m, 3H), 6.48 (d, <i>J</i> = 8.4 Hz, 2H), 5.24-5.19 (m, 1H), 4.99 (s, 2H), 3.26-3.22 (m, 1H), 3.07-3.00 (m, 1H), 2.28-2.20 (m, 1H), 2.00-1.95 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 426.0.	







99	7-(4-Acrylamidophenyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 10.14 (s, 1H), 7.60 (d, <i>J</i> = 8.4 Hz, 2H), 7.46 (d, <i>J</i> = 8.8 Hz, 2H), 7.40-7.32 (m, 2H), 7.12 (t, <i>J</i> = 7.6 Hz, 1H), 6.94-7.06 (m, 6H), 6.75 (s, 1H), 6.39 (dd, <i>J</i> = 10.1, 17.0 Hz, 1H), 6.21 (dd, <i>J</i> = 1.8, 17.0 Hz, 1H), 5.71 (dd, <i>J</i> = 1.8, 10.1 Hz, 1H), 5.41-5.36 (m, 1H), 3.27-3.22 (m, 1H), 3.07-2.97 (m, 1H), 2.37-2.24 (m, 1H), 2.08-1.95 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 480.	
100	2-(4-phenoxyphenyl)-7-(pyridin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 8.57 (d, <i>J</i> = 6.0 Hz, 2H), 7.50 (d, <i>J</i> = 8.8 Hz, 2H), 7.45-7.37 (m, 2H), 7.22-7.10 (m, 3H), 7.07-7.03 (m, 4H), 6.86-6.83 (m, 1H), 5.56-5.10 (m, 1H), 3.35-3.29 (m, 1H), 3.00-2.94 (m, 1H), 2.48-2.38 (m, 1H), 2.18-2.14 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 412.2.	
101	2-(4-phenoxyphenyl)-7-(pyridin-3-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 8.51 (dd, <i>J</i> = 1.6, 4.8 Hz, 1H), 8.37 (d, <i>J</i> = 1.6 Hz, 1H), 7.52-7.44 (m, 3H), 7.42-7.38 (m, 3H), 7.19-7.13 (m, 1H), 7.07-7.02 (m, 4H), 6.84 (br s, 1H), 5.56-5.51 (m, 1H), 3.34-3.30 (m, 1H), 3.13-3.03 (m, 1H), 2.46-2.40 (m, 1H), 2.21-2.14 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 411.9.	
102	2-(4-phenoxyphenyl)-7-(pyridin-2-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 8.57 (d, <i>J</i> = 4.4 Hz, 1H), 7.79 (td, <i>J</i> = 1.6, 7.6 Hz, 1H), 7.49 (d, <i>J</i> = 8.6 Hz, 2H), 7.40 (t, <i>J</i> = 8.0 Hz, 2H), 7.36-7.28 (m, 1H), 7.16 (t, <i>J</i> = 8.0 Hz, 1H), 7.09-7.00 (m, 5H), 6.81 (br s, 1H), 5.48-5.43 (m, 1H), 3.31-3.25 (m, 1H), 3.10-3.02 (m, 1H), 2.42-2.28 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 411.9.	
103	7-(1-acryloylpyrrolidin-2-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.49 (d, <i>J</i> = 8.6 Hz, 2H), 7.46-7.38 (m, 2H), 7.20-7.15 (m, 1H), 7.09 (d, <i>J</i> = 8.0 Hz, 2H), 7.06 (d, <i>J</i> = 8.6 Hz, 2H), 6.76 (s, 1H), 6.62 (dd, <i>J</i> = 16.8, 10.3 Hz, 1H), 6.16 (dd, <i>J</i> = 16.8, 2.2 Hz, 1H), 5.70 (dd, <i>J</i> = 10.3, 2.2 Hz, 1H), 4.47-4.40 (m, 1H), 4.36-4.28 (m, 1H), 3.65-3.50 (m, 2H), 3.50-3.40 (m, 1H), 3.37-3.33 (m, 1H), 2.06-1.74 (m, 6H). MS (ESI) <i>m/e</i> [M+1] ⁺ 457.9.	
104	7-(1-acryloylazetid-3-yl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-	¹ H NMR (DMSO-d ₆) δ 7.49-7.45 (m, 2H), 7.44-7.37 (m, 4H), 7.37-7.31 (m, 1H), 7.10-7.07 (m, 2H), 6.68 (s, 1H), 6.38-6.25 (m, 1H), 6.12-6.04 (m, 1H), 5.69-5.57 (m, 1H), 5.17-5.12 (m, 2H), 4.42-3.78 (m, 5H), 3.31-3.25 (m, 2H), 3.04-2.88 (m, 1H), 2.15-2.04 (m, 1H), 1.79-1.68 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 457.9.	





	carboxamide		
105	7-(1-acryloylazetidin-3-yl)-2-(4-(pyridin-2-ylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, dmso) δ 8.61-8.57 (m, 1H), 7.90-7.78 (m, 1H), 7.56-7.52 (m, 1H), 7.44-7.39 (m, 2H), 7.38-7.34 (m, 1H), 7.12-7.08 (m, 2H), 6.68 (s, 1H), 6.39-6.23 (m, 1H), 6.12-6.01 (m, 1H), 5.68-5.59 (m, 1H), 5.24-5.21 (m, 2H), 4.44-4.23 (m, 2.5H), 4.19-4.07 (m, 1H), 4.04-3.94 (m, 1H), 3.88-3.78 (m, 0.5H), 3.31-3.24 (m, 2H), 3.02-2.88 (m, 1H), 2.15-2.03 (m, 1H), 1.80-1.68 (m, 1H). MS (ESI) m/e [M+1] ⁺ 458.9	
106	7-(1-acryloylazetidin-3-yl)-2-(4-(4-chlorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.52 (d, <i>J</i> = 8.8 Hz, 2H), 7.46 (d, <i>J</i> = 8.8 Hz, 2H), 7.11 (d, <i>J</i> = 8.8 Hz, 2H), 7.08 (d, <i>J</i> = 8.8 Hz, 2H), 6.69 (s, 1H), 6.38-6.22 (m, 1H), 6.13-6.03 (m, 1H), 5.70-5.60 (m, 1H), 4.43-4.26 (m, 2.5H), 4.19-4.08 (m, 1H), 4.05-3.95 (m, 1H), 3.87-3.79 (m, 0.5H), 3.31-3.25 (m, 2H), 3.04-2.90 (m, 1H), 2.15-2.05 (m, 1H), 1.80-1.66 (m, 1H). MS (ESI) m/e [M+1] ⁺ 477.9.	
107	7-(1-acryloylazetidin-3-yl)-2-(4-(3-chlorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.54 (d, <i>J</i> = 8.4 Hz, 2H), 7.46-7.40 (m, 1H), 7.25-7.19 (m, 1H), 7.18-7.15 (m, 1H), 7.11 (d, <i>J</i> = 8.4 Hz, 2H), 7.07-7.02 (m, 1H), 6.69 (s, 1H), 6.39-6.26 (m, 1H), 6.13-6.01 (m, 1H), 5.69-5.60 (m, 1H), 4.45-4.24 (m, 2.5H), 4.19-4.09 (m, 1H), 4.06-3.94 (m, 1H), 3.88-3.80 (m, 0.5H), 3.32-3.24 (m, 2H), 3.05-2.90 (m, 1H), 2.15-2.04 (m, 1H), 1.80-1.68 (m, 1H). MS (ESI) m/e [M+1] ⁺ 477.9.	
108	7-(1-acrylamidocyclopropyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 8.41 (s, 1H), 7.52 (d, <i>J</i> = 8.6 Hz, 2H), 7.46-7.38 (m, 2H), 7.21-7.15 (m, 1H), 7.11-7.04 (m, 4H), 6.75 (s, 1H), 6.21 (dd, <i>J</i> = 17.1, 9.9 Hz, 1H), 6.09 (dd, <i>J</i> = 17.1, 2.4 Hz, 1H), 5.58 (dd, <i>J</i> = 9.9, 2.4 Hz, 1H), 4.36-4.30 (m, 1H), 3.44-3.35 (m, 2H), 2.31-2.19 (m, 1H), 1.96-1.82 (m, 1H), 1.08-1.00 (m, 1H), 0.81-0.73 (m, 1H), 0.72-0.59 (m, 2H). MS (ESI) m/e [M+1] ⁺ 443.9.	
109	7-(2-acrylamidopropyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 8.06 (s, 1H), 7.53 (d, <i>J</i> = 8.6 Hz, 2H), 7.46-7.37 (m, 2H), 7.17 (t, <i>J</i> = 8.0 Hz, 1H), 7.08 (d, <i>J</i> = 8.0 Hz, 2H), 7.06 (d, <i>J</i> = 8.6 Hz, 2H), 6.77 (s, 1H), 6.28 (dd, <i>J</i> = 17.1, 10.1 Hz, 1H), 6.06 (dd, <i>J</i> = 17.1, 1.6 Hz, 1H), 5.56 (dd, <i>J</i> = 10.1, 1.6 Hz, 1H), 4.80 (t, <i>J</i> = 5.4 Hz, 1H), 3.32-3.26 (m, 2H), 2.11-1.82 (m, 2H). MS (ESI) m/e [M+1] ⁺ 445.9.	

110	7-(aminomethyl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate	¹ H NMR (DMSO-d ₆) δ 7.97 (s, 3H), 7.50-7.44 (m, 4H), 7.43-7.38 (m, 2H), 7.37-7.31 (m, 1H), 7.11 (d, <i>J</i> = 8.8 Hz, 2H), 6.80 (s, 1H), 5.15 (s, 2H), 4.43-4.29 (m, 1H), 3.35-3.21 (m, 4H), 2.20-2.09 (m, 1H), 2.04-1.91 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 337.9.	
111	7-(acrylamidomethyl)-2-(4-(benzyloxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 8.29 (t, <i>J</i> = 6.1 Hz, 1H), 7.51-7.31 (m, 6H), 7.10 (d, <i>J</i> = 8.8 Hz, 1H), 6.68 (s, 1H), 6.26 (dd, <i>J</i> = 17.1, 10.1 Hz, 1H), 6.11 (dd, <i>J</i> = 17.1, 2.1 Hz, 1H), 5.62 (dd, <i>J</i> = 10.1, 2.1 Hz, 1H), 5.15 (s, 2H), 4.21-4.12 (m, 1H), 3.84-3.68 (m, 1H), 3.45-3.36 (m, 1H), 3.32-3.23 (m, 2H), 2.10-1.85 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 431.9.	
112	7-(aminomethyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.39 (d, <i>J</i> = 8.6 Hz, 2H), 6.99 (d, <i>J</i> = 8.6 Hz, 2H), 6.63 (s, 1H), 4.06-3.94 (m, 1H), 3.85 (d, <i>J</i> = 7.0 Hz, 2H), 3.04 (dd, <i>J</i> = 12.8, 6.8 Hz, 1H), 2.87 (dd, <i>J</i> = 12.8, 6.8 Hz, 1H), 2.12-1.91 (m, 2H), 1.30-1.18 (m, 2H), 0.62-0.50 (m, 2H), 0.36-0.28 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 342.0.	
113	7-(acrylamidomethyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (CD ₃ OD-d ₄) δ 7.43 (d, <i>J</i> = 8.4 Hz, 2H), 7.01 (d, <i>J</i> = 8.4 Hz, 2H), 6.32-6.18 (m, 2H), 5.70-5.65 (m, 1H), 4.32-4.22 (m, 1H), 3.87 (d, <i>J</i> = 6.9 Hz, 2H), 3.80-3.66 (m, 2H), 3.52-3.36 (m, 2H), 2.22-1.96 (m, 2H), 1.34-1.20 (m, 1H), 0.66-0.57 (m, 2H), 0.39-0.30 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 395.9.	
114	Cis-7-(1-tert-Butoxycarbonyl)-2-(4-phenoxyphenyl)-5,5a,6,7,8,8a-hexahydro-4H-pyrazolo[1,5-a]pyrrolo[3,4-e]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.52 (d, <i>J</i> = 7.6 Hz, 2H), 7.46-7.38 (m, 2H), 7.18 (t, <i>J</i> = 7.6 Hz, 1H), 7.11-7.03 (m, 4H), 6.71 (br s, 1H), 4.61-4.54 (m, 1H), 3.87 (t, <i>J</i> = 11.6 Hz, 1H), 3.67-3.48 (m, 2H), 3.30-3.12 (m, 3H), 3.06-2.85 (m, 1H), 1.36 (d, 9H). MS (ESI) <i>m/e</i> [M+1] ⁺ 476.0.	






115	Cis-2-(4-phenoxyphenyl)-5,5a,6,7,8,8a-hexahydro-4H-pyrazolo[1,5-a]pyrrolo[3,4-e]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆ and D ₂ O) δ 7.53 (d, <i>J</i> = 8.8 Hz, 2H), 7.44-7.40 (m, 2H), 7.23-7.16 (m, 1H), 7.13-7.04 (m, 4H), 4.79-4.73 (m, 1H), 3.78 (d, <i>J</i> = 12.8 Hz, 1H), 3.57-3.27 (m, 4H), 3.14-3.03 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 376.0.	 Cis
116	Cis-7-acryloyl-2-(4-phenoxyphenyl)-5,5a,6,7,8,8a-hexahydro-4H-pyrazolo[1,5-a]pyrrolo[3,4-e]pyrimidine-3-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.51 (dd, <i>J</i> = 2.0, 8.4 Hz, 2H), 7.45-7.38 (m, 2H), 7.18 (t, <i>J</i> = 7.6 Hz, 1H), 7.08 (d, <i>J</i> = 8.0 Hz, 2H), 7.04 (d, <i>J</i> = 8.4 Hz, 2H), 6.75 (s, 1H), 6.62-6.48 (m, 1H), 6.14-6.04 (m, 1H), 5.69-5.61 (m, 1H), 4.70-4.61 (m, 1H), 4.20-4.10 (m, 1H), 3.98-3.86 (m, 1H), 3.72-3.67 (m, 1H), 3.50-3.40 (m, 1H), 3.42-3.38 (m, 1H), 3.33-3.28 (m, 1H), 3.13-2.94 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 430.0.	 Cis
117	2-(1-acryloylpyrrolidin-3-yl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide	¹ H NMR (DMSO-d ₆) δ 11.76 (s, 1H), 7.67 (d, <i>J</i> = 8.0 Hz, 2H), 7.56 (s, 1H), 7.47-7.38 (m, 2H), 7.17 (t, <i>J</i> = 7.6 Hz, 1H), 7.109 (d, <i>J</i> = 7.6 Hz, 2H), 7.04 (d, <i>J</i> = 8.0 Hz, 2H), 6.60 (ddd, <i>J</i> = 2.2, 10.3, 16.6 Hz, 1H), 6.16 (dd, <i>J</i> = 2.2, 16.6 Hz, 1H), 5.69 (dd, <i>J</i> = 2.2, 10.3 Hz, 1H), 4.09-3.95 (m, 1H), 3.94-3.73 (m, 1H), 3.70-3.37 (m, 3H), 2.42-2.24 (m, 1H), 2.21-2.01 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 441.9.	
118	2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[azetidine-3,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.63 (d, <i>J</i> = 8.4 Hz, 2H), 7.55-7.47 (m, 2H), 7.21-7.14 (m, 4H), 6.88 (s, 1H), 4.47 (d, <i>J</i> = 9.8 Hz, 2H), 3.81 (d, <i>J</i> = 9.8 Hz, 2H), 3.35-3.33 (m, 2H), 2.51-2.44 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 375.9.	
119	1-acryloyl-2'-(4-phenoxyphenyl)-5',6'-dihydro-4'H-spiro[azetidine-3,7'-pyrazolo[1,5-a]pyrimidine]-3'-carboxamide	¹ H NMR (DMSO-d ₆) δ 7.54 (d, <i>J</i> = 8.6 Hz, 2H), 7.46-7.38 (m, 2H), 7.17 (t, <i>J</i> = 7.6 Hz, 1H), 7.11-7.05 (m, 4H), 6.79 (s, 1H), 6.35 (dd, <i>J</i> = 17.0, 10.3 Hz, 1H), 6.14 (dd, <i>J</i> = 17.0, 2.1 Hz, 1H), 5.70 (dd, <i>J</i> = 10.3, 2.1 Hz, 1H), 4.66 (d, <i>J</i> = 8.8 Hz, 1H), 4.38 (d, <i>J</i> = 10.0 Hz, 1H), 4.32 (d, <i>J</i> = 8.8 Hz, 1H), 4.03 (d, <i>J</i> = 10.0 Hz, 1H), 3.31-3.26 (m, 2H), 2.38-2.31 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 429.9.	






120	2-(4-phenoxyphenyl)-4,5,6,7,8,9-hexahydropyrazolo[1',5':1,2]imidazo[4,5-d]azepine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 11.74 (br s, 1H), 9.06 (br s, 2H), 7.65 (d, <i>J</i> = 8.4 Hz, 2H), 7.47-7.39 (m, 2H), 7.18 (t, <i>J</i> = 7.6 Hz, 1H), 7.09 (d, <i>J</i> = 7.6 Hz, 2H), 7.07 (d, <i>J</i> = 8.4 Hz, 2H), 3.19-3.14 (m, 2H), 3.10-3.05 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 387.9.	
121	7-acryloyl-2-(4-phenoxyphenyl)-4,5,6,7,8,9-hexahydropyrazolo[1',5':1,2]imidazo[4,5-d]azepine-3-carboxamide	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.64-7.60 (m, 2H), 7.43-7.37 (m, 2H), 7.19-7.14 (m, 1H), 7.12-7.05 (m, 4H), 6.97-6.85 (m, 1H), 6.32 (dd, <i>J</i> = 2.0, 16.8 Hz, 1H), 5.81 (dd, <i>J</i> = 2.0, 8.4 Hz, 1H), 4.04-3.91 (m, 4H), 3.17-3.09 (m, 2H), 3.07-3.01 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 441.9.	
122	(<i>E</i>)-7-(4-(dimethylamino)but-2-enoyl)-2-(4-phenoxyphenyl)-4,5,6,7,8,9-hexahydropyrazolo[1',5':1,2]imidazo[4,5-d]azepine-3-carboxamide	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.59 (d, <i>J</i> = 8.4 Hz, 2H), 7.41-7.32 (m, 2H), 7.15 (t, <i>J</i> = 7.6 Hz, 1H), 7.10-6.99 (m, 5H), 6.80-6.70 (m, 1H), 4.08-3.84 (m, 6H), 3.15-3.08 (m, 2H), 3.06-3.01 (m, 2H), 2.90 (s, 3H), 2.89 (s, 3H). MS (ESI) <i>m/e</i> [M+1] ⁺ 498.9.	
123	8-acryloyl-2-(4-phenoxyphenyl)-4,5,6,7,8,9-hexahydropyrazolo[5',1':2,3]imidazo[4,5-c]azepine-3-carboxamide As a byproduct in the preparation of compound 121	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.61 (dd, <i>J</i> = 2.4, 8.8 Hz, 2H), 7.41-7.33 (m, 2H), 7.14 (t, <i>J</i> = 8.0 Hz, 1H), 7.11-7.02 (m, 4H), 6.82 (dd, <i>J</i> = 10.6, 16.8 Hz, 1H), 6.19 (dd, <i>J</i> = 1.8, 16.8 Hz, 1H), 5.74 (dd, <i>J</i> = 1.8, 10.6 Hz, 1H), 4.98 (d, <i>J</i> = 12.4 Hz, 2H), 4.06-3.80 (m, 2H), 2.96-2.89 (m, 2H), 2.12-1.93 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 441.9.	
124	(<i>E</i>)-8-(4-(dimethylamino)but-2-enoyl)-2-(4-phenoxyphenyl)-4,5,6,7,8,9-hexahydropyrazolo[5',1':2,3]imidazo[4,5-c]azepine-3-carboxamide As a byproduct in the	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.54 (dd, <i>J</i> = 2.4, 8.8 Hz, 2H), 7.34-7.26 (m, 2H), 7.10-7.04 (m, 1H), 7.03-6.95 (m, 4H), 6.75-6.50 (m, 2H), 4.90 (d, <i>J</i> = 11.6 Hz, 2H), 3.89-3.77 (m, 2H), 3.17-3.12 (m, 2H), 2.88-2.82 (m, 2H), 2.28 (s, 3H), 2.19 (s, 3H), 2.02-1.90 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 498.9.	

	preparation of compound 122.		
125	2-(4-(benzyloxy)phenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide formate	¹ H NMR (DMSO- <i>d</i> ₆) δ 11.94 (s, 1H), 9.78 (s, 1H), 9.33 (s, 2H), 7.30-7.22 (m, 5H), 7.18-7.13 (m, 1H), 6.95-6.85 (m, 1H), 4.41 (s, 2H), 3.91 (s, 2H), 2.94-2.87 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 387.9.	
126	7-Acryloyl-2-(4-(benzyloxy)phenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 9.70 (s, 1H), 7.40 (d, <i>J</i> = 8.2 Hz, 2H), 7.35-7.29 (m, 2H), 7.28-7.23 (m, 1H), 7.19 (d, <i>J</i> = 8.2 Hz, 2H), 6.95-6.87 (m, 1H), 6.86-6.83 (m, 2H), 6.20-6.07 (m, 1H), 5.74-5.73 (m, 3H), 4.80-4.72 (m, 2H), 3.89-3.87 (m, 2H), 2.69-2.67 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 441.9.	
127	7-acryloyl-2-(4-(cyclopropylmethoxy)phenyl)-5,6,7,8-tetrahydro-4H-pyrazolo[5',1':2,3]imidazo[4,5-c]pyridine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 9.67 (s, 1H), 7.39 (d, <i>J</i> = 8.4 Hz, 2H), 7.01-6.89 (m, 1H), 6.84 (d, <i>J</i> = 8.4 Hz, 2H), 6.22-6.11 (m, 1H), 5.80-5.70 (m, 1H), 4.85-4.70 (m, 2H), 4.30-4.26 (m, 2H), 3.99-3.90 (m, 2H), 2.90-2.80 (m, 2H), 1.18-1.06 (m, 1H), 0.44-0.30 (m, 4H). MS (ESI) <i>m/e</i> [M+1] ⁺ 405.9.	
128	6-nitro-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b]pyrazole-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 12.78 (s, 1H), 8.29 (s, 1H), 8.14-8.08 (m, 1H), 8.04-7.96 (m, 1H), 7.91-7.81 (m, 2H), 7.49-7.39 (m, 2H), 7.19 (t, <i>J</i> = 7.6 Hz, 1H), 7.15-7.05 (m, 4H). MS (ESI) <i>m/e</i> [M+1] ⁺ 413.9.	
129	6-amino-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b]pyrazole-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 11.64 (s, 1H), 8.24 (s, 1H), 7.73 (d, <i>J</i> = 8.8 Hz, 2H), 7.47-7.39 (m, 3H), 7.18 (t, <i>J</i> = 7.6 Hz, 1H), 7.12-7.05 (m, 4H), 6.69 (d, <i>J</i> = 1.8 Hz, 1H), 6.50 (dd, <i>J</i> = 1.8, 8.8 Hz, 1H), 5.21 (br s, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 383.9.	
130	6-acrylamido-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b]pyrazole-3-carboxamide	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 8.14 (s, 1H), 7.68 (d, <i>J</i> = 8.8 Hz, 1H), 7.62 (d, <i>J</i> = 8.8 Hz, 2H), 7.38-7.24 (m, 3H), 7.12-6.94 (m, 5H), 6.46-6.25 (m, 2H), 5.71 (dd, <i>J</i> = 2.0, 9.6 Hz, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 437.9.	

131	8-amino-2-(4-phenoxyphenyl)-4H-benzo[4,5]imidazo[1,2-b]pyrazole-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 11.87 (s, 1H), 7.78 (d, <i>J</i> = 8.4 Hz, 2H), 7.48-7.39 (m, 2H), 7.18 (t, <i>J</i> = 7.6 Hz, 1H), 7.13-7.07 (m, 4H), 7.02 (t, <i>J</i> = 8.0 Hz, 1H), 6.70 (d, <i>J</i> = 8.0 Hz, 1H), 6.50 (d, <i>J</i> = 8.0 Hz, 1H), 5.56 (s, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 383.9.	
132	7-(1-acryloylazetidin-3-yl)-2-(4-(3,4-dichlorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 7.61 (dd, <i>J</i> = 8.9, 2.8 Hz, 1H), 7.51 (d, <i>J</i> = 8.4 Hz, 2H), 7.36 (d, <i>J</i> = 2.8 Hz, 1H), 7.10 (d, <i>J</i> = 8.4 Hz, 2H), 7.08-7.03 (m, 1H), 6.66 (s, 1H), 6.35-6.22 (m, 1H), 6.09-6.01 (m, 1H), 5.65-5.55 (m, 1H), 4.40-4.30 (m, 2H), 4.29-4.21 (m, 0.5H), 4.15-4.06 (m, 1H), 4.02-3.92 (m, 1H), 3.84-3.76 (m, 0.5H), 3.28-3.21 (m, 1H), 3.03-2.87 (m, 1H), 2.21-2.00 (m, 1H), 1.77-1.66 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 511.8, 513.8.	
133	7-(2-acrylamidophenyl)-2-(3-methoxy-4-methylphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 9.82 (s, 1H), 7.42 (d, <i>J</i> = 7.4 Hz, 1H), 7.27 (t, <i>J</i> = 7.4 Hz, 1H), 7.19 (d, <i>J</i> = 7.4 Hz, 1H), 7.15 (d, <i>J</i> = 7.4 Hz, 1H), 6.96 (s, 1H), 6.93 (d, <i>J</i> = 7.4 Hz, 1H), 6.83 (s, 1H), 6.61 (d, <i>J</i> = 7.4 Hz, 1H), 6.50 (dd, <i>J</i> = 17.1, 10.0 Hz, 1H), 6.24 (d, <i>J</i> = 17.1 Hz, 1H), 5.80-5.70 (m, 2H), 3.74 (s, 3H), 3.28-3.19 (m, 1H), 2.99-2.87 (m, 1H), 2.33-2.19 (m, 1H), 2.13 (s, 3H), 1.99-1.89 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 431.9.	
133a (peak 1)	(R or S) 7-(2-acrylamidophenyl)-2-(3-methoxy-4-methylphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	Chiral HPLC analysis condition: Instrument: Agilent 1260 HPLC Column: CHIRALCEL OJ-H Column size: 0.46 cm I.D.× 15 cm L, 5 μm Mobile phase: n-Hexane/EtOH(0.1% triethyl amine) =85/15(v/v) Column temperature: 35 °C Flow rate: 1.0 ml/min retention time: 5.86 min	
133b (peak 2)	(S or R) 7-(2-acrylamidophenyl)-2-(3-methoxy-4-methylphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	retention time: 6.64 min	

134	7-(2-acrylamidophenyl)-2-(4-chlorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 9.81 (s, 1H), 7.48 (d, <i>J</i> = 8.6 Hz, 2H), 7.45-7.38 (m, 3H), 7.27 (t, <i>J</i> = 7.2 Hz, 1H), 7.17 (t, <i>J</i> = 7.2 Hz, 1H), 6.76 (s, 1H), 6.59 (d, <i>J</i> = 7.2 Hz, 1H), 6.50 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 6.24 (dd, <i>J</i> = 17.0, 1.8 Hz, 1H), 5.79-5.67 (m, 2H), 3.26-3.18 (m, 1H), 2.99-2.89 (m, 1H), 2.35-2.20 (m, 1H), 1.99-1.88 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 421.8, 423.8.	
134a (peak 1)	(R or S) 7-(2-acrylamidophenyl)-2-(4-chlorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	Chiral HPLC analysis condition: Instrument: Agilent 1260 HPLC Column: CHIRALPAK AD-H Column size: 0.46 cm I.D. × 15 cm L, 5 μm Mobile phase: n-Hexane/EtOH (0.1% triethyl amine)=70/30 (v/v) Column temperature: 35 °C Flow rate: 1.0 ml/min retention time: 4.15 min	
134b (peak 2)	(S or R) 7-(2-acrylamidophenyl)-2-(4-chlorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	retention time: 10.52 min	
135	7-(1-acryloylazetididin-3-yl)-2-(4-(2-cyanophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 7.91-7.85 (m, 1H), 7.71-7.61 (m, 1H), 7.58-7.52 (m, 2H), 7.33-7.24 (m, 1H), 7.20-7.14 (m, 2H), 7.11-7.05 (m, 1H), 6.61 (br s, 1H), 6.36-6.22 (m, 1H), 6.10-6.00 (m, 1H), 5.67-5.54 (m, 1H), 4.42-4.30 (m, 2H), 4.30-4.22 (m, 0.5H), 4.16-4.07 (m, 1H), 4.03-3.92 (m, 1H), 3.87-3.73 (m, 0.5H), 3.30-3.20 (m, 2H), 3.05-2.85 (m, 1H), 2.15-1.97 (m, 1H), 1.78-1.64 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 468.9.	
136	3-(1-acryloylpiperidin-4-yl)-6-(4-phenoxyphenyl)-1H-imidazo[1,2-b]pyrazole-7-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 11.53 (s, 1H), 7.64 (d, <i>J</i> = 8.4 Hz, 2H), 7.40 (d, <i>J</i> = 7.6 Hz, 1H), 7.38 (d, <i>J</i> = 7.6 Hz, 1H), 7.14 (t, <i>J</i> = 7.6 Hz, 1H), 7.05 (d, <i>J</i> = 8.0 Hz, 2H), 7.02 (d, <i>J</i> = 8.0 Hz, 2H), 6.96 (s, 1H), 6.80 (dd, <i>J</i> = 16.8, 10.1 Hz, 1H), 6.40 (brs, 2H), 6.07 (dd, <i>J</i> = 16.8, 2.0 Hz, 1H), 5.64 (dd, <i>J</i> = 10.1, 2.0 Hz, 1H), 4.46 (d, <i>J</i> = 12.8 Hz, 1H), 4.10 (d, <i>J</i> = 12.8 Hz, 1H), 3.24-3.04 (m, 2H), 2.80 (t, <i>J</i> = 12.8 Hz, 1H), 2.20-2.05 (m, 2H), 1.64-1.46 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 455.9.	






137	7-(2-acrylamidophenyl)-2-(3,4-difluorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 9.81 (s, 1H), 7.52-7.36 (m, 3H), 7.34-7.24 (m, 2H), 7.17 (t, <i>J</i> = 7.6 Hz, 1H), 6.74 (br s, 2H), 6.59 (d, <i>J</i> = 7.6 Hz, 1H), 6.50 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 6.24 (dd, <i>J</i> = 17.0, 1.6 Hz, 1H), 5.80-5.70 (m, 2H), 3.28-3.18 (m, 1H), 3.01-2.87 (m, 1H), 2.34-2.20 (m, 1H), 2.01-1.89 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 423.9.	
138	7-(1-acryloylpiperidin-4-yl)-7-methyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 7.50 (d, <i>J</i> = 8.6 Hz, 2H), 7.45-7.37 (m, 2H), 7.17 (t, <i>J</i> = 7.4 Hz, 1H), 7.11-7.03 (m, 4H), 6.85-6.72 (m, 1H), 6.68 (s, 1H), 6.07 (dd, <i>J</i> = 16.7, 2.4 Hz, 1H), 5.64 (d, <i>J</i> = 10.6 Hz, 1H), 4.58-4.41 (m, 1H), 4.19-4.00 (m, 1H), 3.33-3.24 (m, 2H), 3.05-2.85 (m, 1H), 2.60-2.50 (m, 1H), 2.31-2.16 (m, 1H), 2.15-2.00 (m, 1H), 1.80-1.65 (m, 2H), 1.44 (s, 3H), 1.35-1.05 (m, 3H). MS (ESI) <i>m/e</i> [M+1] ⁺ 486.0.	
139	7-(1-acryloylazetidin-3-yl)-2-(3-chloro-4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 7.70-7.66 (m, 1H), 7.52-7.46 (m, 1H), 7.44-7.36 (m, 2H), 7.19-7.07 (m, 2H), 7.04 (d, <i>J</i> = 8.6 Hz, 2H), 6.66 (s, 1H), 6.40-6.25 (m, 1H), 6.13-6.05 (m, 1H), 5.70-5.60 (m, 1H), 4.45-4.36 (m, 2H), 4.33-4.26 (m, 0.5H), 4.21-4.08 (m, 1H), 4.07-3.95 (m, 1H), 3.88-3.80 (m, 0.5H), 3.31-3.24 (m, 2H), 3.04-2.90 (m, 1H), 2.15-2.04 (m, 1H), 1.81-1.67 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 477.9.	
140	7-(1-acryloylazetidin-3-yl)-2-(3-methoxy-4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide formate	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 8.38 (br s, 1H), 7.31-7.19 (m, 3H), 7.15-7.10 (m, 1H), 7.06-6.96 (m, 2H), 6.88 (d, <i>J</i> = 8.1 Hz, 2H), 6.17-6.11 (m, 2H), 5.64-5.54 (m, 1H), 4.24-4.08 (m, 2H), 4.06-3.95 (m, 1H), 3.85-3.70 (m, 4H), 3.55-3.45 (m, 2H), 3.30-3.20 (m, 1H), 3.05-2.90 (m, 1H), 2.48-2.36 (m, 1H), 1.40-1.25 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 473.9.	
141	7-(1-acryloylazetidin-3-yl)-2-(4-phenethoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.41-7.35 (m, 2H), 7.32-7.24 (m, 4H), 7.22-7.15 (m, 1H), 7.03-7.96 (m, 2H), 6.31 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 6.19 (dd, <i>J</i> = 17.0, 2.0 Hz, 1H), 5.74-5.66 (m, 1H), 4.55-4.26 (m, 3H), 4.25-3.94 (m, 4H), 3.48-3.34 (m, 2H), 3.13-3.03 (m, 3H), 2.26-2.14 (m, 1H), 2.04-1.88 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 471.9.	






142	7-(acrylamidomethyl)-2-(4-(benzyloxy)-3-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 8.31 (t, <i>J</i> = 6.0 Hz, 1H), 7.50-7.44 (m, 2H), 7.44-7.31 (m, 3H), 7.17-7.05 (m, 2H), 7.04-6.99 (m, 1H), 6.72 (s, 1H), 6.26 (dd, <i>J</i> = 17.1, 10.1 Hz, 1H), 6.11 (dd, <i>J</i> = 17.1, 1.8 Hz, 1H), 5.62 (dd, <i>J</i> = 10.1, 1.8 Hz, 1H), 5.12 (s, 2H), 4.22-4.12 (m, 1H), 3.85-3.72 (m, 4H), 3.48-3.33 (m, 3H), 2.07-1.87 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 461.9.	
143	7-(acrylamidomethyl)-7-methyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 8.08 (t, <i>J</i> = 6.4 Hz, 1H), 7.53-7.45 (m, 2H), 7.42-7.34 (m, 2H), 7.14 (t, <i>J</i> = 7.4 Hz, 1H), 7.08-7.00 (m, 4H), 6.66 (s, 1H), 6.29 (dd, <i>J</i> = 17.1, 10.2 Hz, 1H), 6.08 (dd, <i>J</i> = 17.1, 2.2 Hz, 1H), 5.58 (dd, <i>J</i> = 10.2, 2.2 Hz, 1H), 3.66 (dd, <i>J</i> = 13.5, 6.8 Hz, 1H), 3.47 (dd, <i>J</i> = 13.5, 6.8 Hz, 1H), 2.05-1.94 (m, 1H), 1.79-1.69 (m, 1H), 1.38 (s, 3H). MS (ESI) <i>m/e</i> [M+1] ⁺ 431.9.	
144	7-(1-acryloylpiperidin-4-yl)-2-phenyl-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide trifluoroacetate	¹ H NMR (CD ₃ OD- <i>d</i> ₄) δ 7.54-7.42 (m, 5H), 6.75 (dd, <i>J</i> = 16.8, 10.7 Hz, 1H), 6.15 (dd, <i>J</i> = 16.8, 1.6 Hz, 1H), 5.70 (dd, <i>J</i> = 10.7, 1.6 Hz, 1H), 4.70-4.55 (m, 1H), 4.25-4.00 (m, 2H), 3.50-3.40 (m, 2H), 3.14-3.03 (m, 1H), 2.73-2.61 (m, 1H), 2.39-2.14 (m, 2H), 2.10-1.94 (m, 1H), 1.84-1.66 (m, 2H), 1.51-1.30 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 380.0.	
145	7-(1-acryloylazetidin-3-yl)-2-(4-chlorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (DMSO- <i>d</i> ₆) δ 7.50 (d, <i>J</i> = 8.6 Hz, 2H), 7.45 (d, <i>J</i> = 8.6 Hz, 2H), 6.62 (s, 1H), 6.35-6.22 (m, 1H), 6.10-6.01 (m, 1H), 5.65-5.55 (m, 1H), 4.40-4.21 (m, 2.5H), 4.14-4.06 (m, 1H), 4.04-3.90 (m, 1H), 3.85-3.75 (m, 0.5H), 3.27-3.21 (m, 2H), 3.00-2.86 (m, 1H), 2.12-1.98 (m, 1H), 1.77-1.63 (m, 1H).	
146	7-(2-Acrylamidophenyl)-2-(3-chloro-4-(pyridin-2-ylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.83 (s, 1H), 8.59 (d, <i>J</i> = 4.2 Hz, 1H), 7.86 (dt, <i>J</i> = 7.6, 2.0 Hz, 1H), 7.56 (d, <i>J</i> = 7.6 Hz, 1H), 7.54 (d, <i>J</i> = 2.0 Hz, 2H), 7.48-7.40 (m, 2H), 7.39-7.33 (m, 1H), 7.33-7.24 (m, 2H), 7.21 (t, <i>J</i> = 7.6 Hz, 1H), 6.77 (s, 1H), 6.63 (d, <i>J</i> = 8.0 Hz, 1H), 6.53 (dd, <i>J</i> = 16.8, 10.2 Hz, 1H), 6.27 (dd, <i>J</i> = 16.8, 1.8 Hz, 1H), 5.82-5.74 (m, 2H), 5.30 (s, 2H), 3.30-3.20 (m, 1H), 3.02-2.92 (m, 1H), 2.37-2.23 (m, 1H), 2.02-1.93 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 528.9.	






147	7-(2-Acrylamido-4-chlorophenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 491.9.	
148	7-(2-Acrylamido-4-methylphenyl)-2-(4-(cyclopropylmethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 471.9.	
149	7-(2-Acrylamidophenyl)-2-(4-(trifluoromethoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 471.9.	
150	7-(2-Acrylamidophenyl)-2-phenyl-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 387.9.	
151	7-(2-Acrylamidophenyl)-2-(4-fluorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 405.9.	






152	7-(2-Acrylamidophenyl)-2-(3-bromo-4-fluorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.83 (s, 1H), 7.77 (dd, <i>J</i> = 6.8, 2.0 Hz, 1H), 7.58-7.51 (m, 1H), 7.45 (d, <i>J</i> = 7.6 Hz, 1H), 7.39 (t, <i>J</i> = 8.6 Hz, 1H), 7.31 (t, <i>J</i> = 7.6 Hz, 1H), 7.21 (t, <i>J</i> = 7.6 Hz, 1H), 6.77 (s, 1H), 6.63 (d, <i>J</i> = 7.6 Hz, 1H), 6.53 (dd, <i>J</i> = 17.0, 10.2 Hz, 1H), 6.28 (dd, <i>J</i> = 17.0, 1.8 Hz, 1H), 5.82-5.75 (m, 2H), 3.31-3.21 (m, 1H), 3.03-2.91 (m, 1H), 2.37-2.22 (m, 1H), 2.03-1.91 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 483.8, 485.8.	
153	(<i>E</i>)-2-(3-Chloro-4-fluorophenyl)-7-(2-(4-(piperidin-1-yl)but-2-enamido)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) <i>m/e</i> [M+1] ⁺ 536.9.	
154	7-(2-Acrylamidophenyl)-2-(4-chloro-3-fluorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.83 (s, 1H), 7.65 (dd, <i>J</i> = 7.3, 2.0 Hz, 1H), 7.54-7.48 (m, 1H), 7.47-7.39 (m, 2H), 7.31 (t, <i>J</i> = 7.6 Hz, 1H), 7.21 (t, <i>J</i> = 7.6 Hz, 1H), 6.77 (s, 1H), 6.63 (d, <i>J</i> = 7.8 Hz, 1H), 6.53 (dd, <i>J</i> = 17.0, 10.3 Hz, 1H), 6.27 (dd, <i>J</i> = 17.0, 1.8 Hz, 1H), 5.83-5.75 (m, 2H), 3.30-3.21 (m, 1H), 3.03-2.97 (m, 1H), 2.36-2.24 (m, 1H), 2.04-1.93 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 439.9.	
155	7-(2-Acrylamidophenyl)-2-(3-chlorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) <i>m/e</i> [M+1] ⁺ 421.9.	
156	7-(2-Acrylamidophenyl)-2-(3,4-dichlorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.83 (s, 1H), 7.71 (d, <i>J</i> = 1.6 Hz, 1H), 7.63 (d, <i>J</i> = 8.4 Hz, 1H), 7.50 (dd, <i>J</i> = 8.4, 1.6 Hz, 1H), 7.45 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (t, <i>J</i> = 7.6 Hz, 1H), 7.21 (t, <i>J</i> = 7.6 Hz, 1H), 6.77 (s, 1H), 6.63 (d, <i>J</i> = 7.6 Hz, 1H), 6.53 (dd, <i>J</i> = 16.8, 10.2 Hz, 1H), 6.28 (d, <i>J</i> = 16.8 Hz, 1H), 5.84-5.75 (m, 2H), 3.31-3.21 (m, 1H), 3.05-2.92 (m, 1H), 2.38-2.23 (m, 1H), 2.03-1.94 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 455.8.	






157	7-(2-Acrylamidophenyl)-2-(4-fluoro-3-methylphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 419.9.	
158	7-(2-Acrylamidophenyl)-2-(3-chloro-4,5-difluorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 457.9.	
159	7-(2-Acrylamidophenyl)-2-(4-fluoro-3-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 435.9.	
160	7-(2-Acrylamidophenyl)-2-(4-chloro-3-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 451.8.	
161	7-(2-Acrylamidophenyl)-2-(4-(trifluoromethyl)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 455.9.	






162	7-(2-Acrylamidophenyl)-2-(3-chloro-4-(trifluoromethyl)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 489.8.	
163	7-(2-Acrylamidophenyl)-2-(4-chloro-3-(trifluoromethyl)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 489.8.	
164	7-(2-Acrylamidophenyl)-2-(p-tolyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 401.9.	
165	(E)-7-(2-(4-(Dimethylamino)but-2-enamido)phenyl)-2-(3-methoxy-4-methylphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 489.0.	
166	7-(2-Acrylamidophenyl)-2-(3-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 417.9.	



167	7-(2-Acrylamidophenyl)-2-(3,4,5-trimethoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 477.9.	
168	7-(2-Acrylamidophenyl)-2-(3,4-dimethoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 447.9.	
169	7-(2-Acrylamidophenyl)-2-(3,5-dichloro-4-methoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 485.8.	
170	(E)-2-(4-Phenoxyphenyl)-7-(1-(4-(piperidin-1-yl)but-2-enoyl)piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 569.0.	
171	(R or S) (E)-7-(1-(4-(Dimethylamino)but-2-enoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 529.0.	

172	7-(1-Acryloylpiperidin-4-yl)-2-(3-methoxy-4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 501.9.	
173	7-(1-Acryloylpiperidin-4-yl)-2-(3-chloro-4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 505.9.	
174	7-(1-Acryloylpiperidin-4-yl)-2-(3-methoxy-4-methylphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 424.0.	
175	7-(1-Acryloylazetidin-3-yl)-7-methyl-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 457.9.	
176	(S)-7-(1-(But-2-ynoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.50 (d, <i>J</i> = 7.2 Hz, 2 H), 7.42 (dd, <i>J</i> = 8.4, 8.4 Hz, 2H), 7.17 (t, <i>J</i> = 7.2 Hz, 1H), 7.11-7.03 (m, 4H), 6.68 (s, 1 H), 4.40-4.24 (m, 2 H), 4.06-3.97 (m, 1 H), 3.33-3.27 (m, 1 H), 3.12-3.00 (m, 1 H), 2.66-2.54 (m, 1H), 2.32-2.20 (m, 1H), 2.07-2.00 (m, 1H), 2.00 (s, 3 H), 1.96-1.86 (m, 1 H), 1.81-1.67 (m, 1 H), 1.63-1.52 (m, 1 H), 1.36-1.08 (m, 2 H). MS (ESI) m/e [M+1] ⁺ 483.9.	

177	(S)-7-(1-(3-chloropropanoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.54-7.47 (m, 2H), 7.45-7.38 (m, 2H), 7.21-7.14 (m, 1H), 7.12-7.02 (m, 4H), 6.69 (s, 1H), 4.45 (d, <i>J</i> = 12.1 Hz, 1H), 4.06-3.97 (m, 1H), 3.90 (t, <i>J</i> = 10.3 Hz, 1H), 3.82-3.72 (m, 2H), 3.33-3.25 (m, 2H), 3.02-2.72 (m, 3H), 2.58-2.43 (m, 1H), 2.35-2.15 (m, 1H), 2.10-1.84 (m, 2H), 1.70 (t, <i>J</i> = 12.1 Hz, 1H), 1.54 (t, <i>J</i> = 12.1 Hz, 1H), 1.41-1.08 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 507.9, 509.9.	
179	7-(1-(But-2-ynoyl)piperidin-4-yl)-2-(4-(2,4-difluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.55-7.45 (m, 3H), 7.41-7.32 (m, 1H), 7.21-7.12 (m, 1H), 7.05-6.99 (m, 2H), 6.67 (s, 1H), 4.39-4.22 (m, 2H), 4.06-3.97 (m, 1H), 3.33-3.25 (m, 2H), 3.12-2.97 (m, 1H), 2.69-2.52 (m, 1H), 2.35-2.17 (m, 1H), 2.08-2.00 (m, 1H), 2.00 (s, 3H), 1.96-1.84 (m, 1H), 1.81-1.65 (m, 1H), 1.63-1.51 (m, 1H), 1.36-1.09 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 519.9.	
180	(S)-7-(1-(2-Cyanoacetyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) <i>m/e</i> [M+1] ⁺ 484.9.	
183	7-(1-(Acryloyl)piperidin-4-yl)-2-(4-(2,4-difluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.58-7.44 (m, 3H), 7.41-7.30 (m, 1H), 7.20-7.10 (m, 1H), 7.02 (d, <i>J</i> = 8.5 Hz, 2H), 6.85-6.73 (m, 1H), 6.67 (br s, 1H), 6.07 (d, <i>J</i> = 16.4 Hz, 1H), 5.64 (d, <i>J</i> = 10.5 Hz, 1H), 4.55-4.39 (m, 1H), 4.17-3.95 (m, 2H), 3.74-3.38 (m, 3H), 3.06-2.89 (m, 1H), 2.35-2.15 (m, 1H), 2.09-1.83 (m, 2H), 1.78-1.65 (m, 1H), 1.62-1.49 (m, 1H), 1.34-1.07 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 507.9.	
184	7-(1-(But-2-ynoyl)azetidin-3-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.53-7.46 (m, 2H), 7.45-7.38 (m, 2H), 7.21-7.13 (m, 1H), 7.12-7.01 (m, 4H), 6.72-6.66 (m, 1H), 4.43-4.34 (m, 1H), 4.33-4.27 (m, 0.5H), 4.26-4.15 (m, 1H), 4.11-4.04 (m, 1H), 4.02-3.92 (m, 1H), 3.85-3.77 (m, 0.5H), 3.30-3.23 (m, 2H), 3.03-2.90 (m, 1H), 2.15-2.04 (m, 1H), 2.00-1.90 (m, 3H), 1.82-1.67 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 455.9.	

185	7-(But-2-ynamidomethyl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.67 (t, <i>J</i> = 6.0 Hz, 1H), 7.56-7.49 (m, 2H), 7.46-7.37 (m, 2H), 7.21-7.13 (m, 1H), 7.12-7.01 (m, 4H), 6.68 (s, 1H), 4.20-4.09 (m, 1H), 3.76-3.65 (m, 1H), 3.36-3.24 (m, 3H), 2.09-1.83 (m, 5H). MS (ESI) <i>m/e</i> [M+1] ⁺ 429.9.	
186	(S)-7-(1-(Pent-2-ynoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.50 (d, <i>J</i> = 8.1 Hz, 2H), 7.45-7.38 (m, 2H), 7.17 (t, <i>J</i> = 7.4 Hz, 1H), 7.12-7.02 (m, 4H), 6.68 (s, 1H), 4.41-4.24 (m, 2H), 4.09-3.97 (m, 1H), 3.33-3.26 (m, 2H), 3.14-2.99 (m, 1H), 2.68-2.53 (m, 1H), 2.43-2.33 (m, 2H), 2.31-2.19 (m, 1H), 2.07-1.83 (m, 2H), 1.82-1.66 (m, 1H), 1.63-1.50 (m, 1H), 1.40-1.13 (m, 2H), 1.11 (t, <i>J</i> = 7.5 Hz, 3H). MS (ESI) <i>m/e</i> [M+1] ⁺ 498.0.	
187	(S)-2-(4-Phenoxyphenyl)-7-(1-(4-(pyrrolidin-1-yl)but-2-ynoyl)piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.50 (d, <i>J</i> = 8.2 Hz, 2H), 7.45-7.37 (m, 2H), 7.17 (t, <i>J</i> = 7.4 Hz, 1H), 7.11-7.02 (m, 4H), 6.67 (s, 1H), 4.41-4.22 (m, 2H), 4.08-3.99 (m, 1H), 3.69 (s, 2H), 3.18-3.04 (m, 1H), 2.75-2.55 (m, 5H), 2.35-2.20 (m, 1H), 2.09-1.85 (m, 2H), 1.83-1.66 (m, 5H), 1.66-1.53 (m, 1H), 1.40-1.11 (m, 4H). MS (ESI) <i>m/e</i> [M+1] ⁺ 553.0.	
188	(S)-7-(1-(4-(Dimethylamino)but-2-ynoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.53-7.47 (m, 2H), 7.46-7.38 (m, 2H), 7.20-7.14 (m, 1H), 7.12-7.02 (m, 4H), 6.69-6.64 (m, 1H), 4.45-4.23 (m, 2H), 4.08-3.99 (m, 1H), 3.52 (s, 2H), 3.19-3.05 (m, 1H), 2.72-2.57 (m, 1H), 2.54-2.49 (m, 1H), 2.36-2.18 (m, 7H), 2.10-1.86 (m, 2H), 1.83-1.68 (m, 1H), 1.67-1.53 (m, 1H), 1.41-1.11 (m, 3H). MS (ESI) <i>m/e</i> [M+1] ⁺ 527.0.	
189	7-(1-(But-2-ynoyl)azetidin-3-yl)-2-(4-(2,4-difluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) <i>m/e</i> [M+1] ⁺ 491.9.	

190	7-(But-2-ynamidomethyl)-2-(4-(2,4-difluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 465.9.	
191	(S)-7-(1-(4-Hydroxybut-2-ynoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	MS (ESI) m/e [M+1] ⁺ 499.9.	
192	(S)-7-(1-(4-Methoxybut-2-ynoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.54-7.47 (m, 2H), 7.45-7.38 (m, 2H), 7.21-7.14 (m, 1H), 7.12-7.03 (m, 4H), 6.69 (s, 1H), 4.41-4.32 (m, 1H), 4.30 (s, 2H), 4.29-4.20 (m, 1H), 4.09-3.98 (m, 1H), 3.33-3.26 (m, 2H), 3.29 (s, 3H), 3.19-3.05 (m, 1H), 2.72-2.57 (m, 1H), 2.36-2.21 (m, 1H), 2.09-1.98 (m, 1H), 2.09-1.97 (m, 1H), 1.84-1.68 (m, 1H), 1.66-1.50 (m, 1H), 1.41-1.10 (m, 2H). MS (ESI) m/e [M+1] ⁺ 514.0.	
193	(S)-7-(1-(Hex-2-ynoyl)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.54-7.48 (m, 2H), 7.45-7.38 (m, 2H), 7.21-7.14 (m, 1H), 7.12-7.02 (m, 4H), 6.68 (s, 1H), 4.42-4.22 (m, 2H), 4.06-3.98 (m, 1H), 3.33-3.25 (m, 2H), 3.15-3.01 (m, 1H), 2.69-2.53 (m, 1H), 2.36 (t, <i>J</i> = 6.9 Hz, 2H), 2.32-2.20 (m, 1H), 2.08-1.97 (m, 1H), 1.97-1.84 (m, 1H), 1.83-1.67 (m, 1H), 1.65-1.46 (m, 3H), 1.38-1.09 (m, 3H), 0.94 (t, <i>J</i> = 7.4 Hz, 3H). MS (ESI) m/e [M+1] ⁺ 512.0.	
194	7-(1-(But-2-ynoyl)piperidin-4-yl)-2-(4-(4-fluorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.53-7.46 (m, 2H), 7.30-7.21 (m, 2H), 7.18-7.11 (m, 2H), 7.07-7.00 (m, 2H), 6.64 (br s, 1H), 4.39-4.23 (m, 2H), 4.06-3.98 (m, 1H), 3.34-3.26 (m, 2H), 3.12-2.99 (m, 1H), 2.69-2.53 (m, 1H), 2.31-2.18 (m, 1H), 2.08-2.01 (m, 1H), 2.00 (s, 3H), 1.97-1.86 (m, 1H), 1.81-1.66 (m, 1H), 1.64-1.51 (m, 1H), 1.36-1.08 (m, 2H). MS (ESI) m/e [M+1] ⁺ 501.9.	

195	7-(1-(But-2-ynoyl)piperidin-4-yl)-2-(4-(3-chlorophenoxy)phenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.57-7.50 (m, 2H), 7.46-7.39 (m, 1H), 7.25-7.20 (m, 1H), 7.17-7.10 (m, 3H), 7.08-7.02 (m, 1H), 6.63 (br s, 1H), 4.40-4.22 (m, 2H), 4.06-3.98 (m, 1H), 3.34-3.26 (m, 2H), 3.13-3.00 (m, 1H), 2.69-2.54 (m, 1H), 2.31-2.19 (m, 1H), 2.08-2.00 (m, 1H), 2.00 (s, 3H), 1.96-1.87 (m, 1H), 1.82-1.66 (m, 1H), 1.64-1.52 (m, 1H), 1.35-1.09 (m, 2H). MS (ESI) <i>m/e</i> [M+1] ⁺ 517.9, 519.9.	
196	7-(2-Acrylamidophenyl)-2-(3-chloro-4-fluorophenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 9.82 (s, 1H), 7.65 (dd, <i>J</i> = 7.4, 2.1 Hz, 1H), 7.55-7.47 (m, 1H), 7.47-7.38 (m, 2H), 7.31 (t, <i>J</i> = 7.1 Hz, 1H), 7.34-7.28 (m, 1H), 6.76 (br s, 1H), 6.63 (d, <i>J</i> = 7.4 Hz, 1H), 6.53 (dd, <i>J</i> = 17.2, 10.3 Hz, 1H), 6.27 (dd, <i>J</i> = 17.2, 1.9 Hz, 1H), 5.83-5.73 (m, 2H), 3.30-3.22 (m, 1H), 3.03-2.93 (m, 1H), 2.37-2.24 (m, 1H), 2.03-1.93 (m, 1H). MS (ESI) <i>m/e</i> [M+1] ⁺ 439.8, 441.8.	

[0636] BTK KINASE ASSAY

[0637] Compounds disclosed herein were tested for inhibition of Btk kinase activity in an assay based on time-resolved fluorescence resonance energy transfer methodology. Recombinant Btk was pre-incubated with the compounds disclosed herein at room temperature for 1 hour in an assay buffer containing 50 mM Tris pH7.4, 10 mM MgCl₂, 2 mM MnCl₂, 0.1 mM EDTA, 1 mM DTT, 20 nM SEB, 0.1% BSA, 0.005% tween-20. The reactions were initiated by the addition of ATP (at the concentration of ATP K_m) and peptide substrate (Biotin-AVLESEEEELYSSARQ-NH₂). After incubating at room temperature for 1 h, an equal volume of stop solution containing 50 mM HEPES pH7.0, 800 mM KF, 20 mM EDTA, 0.1% BSA, Eu cryptate-conjugated p-Tyr66 antibody and streptavidin-labeled XL665 was added to stop the reaction. Plates were further incubated at room temperature for 1 hour, and then the TR-FRET signals (ex337nm, em 620nm/665 nm) were read on BMG PHERAstar FS instrument. The residual enzyme activity in presence of increasing concentrations of compounds was calculated based on the ratio of fluorescence at 615 nm to that at 665nm. The IC₅₀ for each compound was derived from fitting the data to the four-parameter logistic equation by Graphpad Prism software.

[0638] BTKpY223 CELLULAR ASSAY

[0639] Btk pY223 cellular assay is a HTRF based assay intended to determine the endogenous levels of phosphorylated Btk at Tyr223. Phosphorylated Tyr223 is necessary for full activation of Btk. The assay was performed in Ramos cells (CRL-1596, ATCC) with a Btk pY223 assay kit (63IDC000, Cisbio).

[0640] Briefly, Ramos cells were serum starved in 0.5% FBS-containing RPMI1640 for 2 hours. Following starvation, the cells were incubated with compounds to be detected at various concentrations in a CO2 incubator for 1 hour. After incubation, cells were stimulated with 1mM pervanadate (PV) or Na₃VO₄ (OV) for 20 min. Then, the cells were spun down and lysed with 1x lysis buffer at RT for 10 min (4x lysis buffer supplied in the kit). During the incubation, 1x antibody mix was prepared by diluting anti-Btk-d2 and anti-pBtk-K in detection buffer (supplied in the kit). 2 ul/well of 1 x antibody mixture was dispensed into the OptiPlate-384 assay plate (6005620, Perkin Elmer). After that, 18 ul of cell lysate was transferred to the assay plate pre-loaded with antibody solution. After mixing gently and spinning briefly, the plate was sealed up and kept in dark at RT for 18 hours. The fluorescence emission was measured at two different wavelengths (665 nm and 620 nm) on a compatible HTRF reader (PHERAstar FS, BMG). The potency of compounds was calculated basing on the inhibition of ratio between signal intensities at 665nm and 620nm. IC₅₀ values were calculated with GraphPad Prism software using the sigmoidal dose-response function.

[0641] Representative compounds as disclosed herein were tested and found to inhibit Btk and autophosphorylation of Btk at Tyr-223 with IC₅₀ values ranging from subnanomolar to 10 micromolar.

[0642] Table II Assay data for representative compounds

Compound No.	IC ₅₀ (nM)	
	Btk	Btk pY223
1	>10,000	n.d.
2	360	n.d.
2a (peak 1)	270	n.d.
2b (peak 2)	210	n.d.
3	3.6	12.7
3a (peak 1)	2.5	8.7
3b (peak 2)	1.7	2.0
4	26	n.d.
5	200	n.d.
6	0.71	3.1
6a (peak 1)	0.4	3.0
6b (peak 2)	120	n.d.
7	>10,000	n.d.
8	860	n.d.

9	3,700	n.d.
10	1.5	8.0
10a (peak 1)	0.51	2.2
10b (peak 2)	42	n.d.
11	4,500	n.d.
12	1.7	23.6
13	1000	n.d.
14	4.7	232.6
15	2,700	n.d.
16	19	n.d.
17	>10,000	n.d.
18	1.3	26.3
19	5,300	n.d.
20	1.6	12.6
20a (peak 1)	1.2	2.5
20b (peak 2)	110	n.d.
21	>10,000	n.d.
22	2.7	16.7
23	3,200	n.d.
24	39	n.d.
25	3,100	n.d.
26	1700	n.d.
27	1	4.6
27a (peak 1)	0.33	5.7
27b (peak 2)	2	20.0
28	3,400	n.d.
29	0.48	18
29a (peak 1)	0.43	1.6
29b (peak 2)	1.3	4.1
30	1	4.0
30a (peak 1)	0.66	5.6
30b (peak 2)	16	n.d.
31	15	n.d.
32	1,700	n.d.

33	0.9	2.9
33a (peak 1)	0.6	5.9
33b (peak 2)	11	63.3
34	1,900	n.d.
35	1.1	7.9
36	>10,000	n.d.
37	640	n.d.
38	490	n.d.
39	1,400	n.d.
40	13	n.d.
41	180	n.d.
42	330	n.d.
43	3.0	75.8
44	220	n.d.
45	510	n.d.
46	5	22.6
47	110	n.d.
48	150	n.d.
49	29	n.d.
50	2.7	8.0
51	310	n.d.
52	0.14	<0.5
53	7.7	n.d.
54	0.19	6.4
55	82	n.d.
56	110	n.d.
57	37	232.9
58	0.43	4.6
59	43	n.d.
60	40	n.d.
61	140	n.d.
62	240	n.d.
63	0.18	1.5
64	0.17	4.8

65	320	n.d.
66	14	322.2
67	0.32	2.8
68	190	n.d.
69	7.8	34.4
70	19	n.d.
71	55	n.d.
72	4.1	10.5
73	>10,000	n.d.
74	240	n.d.
75	41	222.9
76	5,500	n.d.
77	480	n.d.
78	21	139.6
79	50	n.d.
80	>10,000	n.d.
81	5,900	n.d.
82	3,100	n.d.
83	130	n.d.
84	2,900	n.d.
85	32	73.2
86	2.2	4.4
87	2,000	n.d.
88	16.0	n.d.
89	>10,000	n.d.
90	1,900	n.d.
91	1,900	n.d.
92	1.1	7.0
93	1,500	n.d.
94	1.9	6.7
95	470	n.d.
96	30	n.d.
97	1.1	n.d.
98	77	n.d.


99	9.5	43.4
100	260	n.d.
101	550	n.d.
102	830	n.d.
103	1.1	14
104	2.6	20.6
105	18	n.d.
106	1.4	30.3
107	0.48	11.2
108	3.5	18.9
109	41	n.d.
110	3,000	n.d.
111	24	n.d.
112	>10,000	n.d.
113	94	n.d.
114	6,000	n.d.
115	8,200	n.d.
116	1	3.2
117	34	n.d.
118	410	n.d.
119	40	n.d.
120	180	n.d.
121	0.36	8.2
122	0.78	12.6
123	0.15	1.5
124	0.62	4.6
125	5,000	n.d.
126	20	n.d.
127	300	n.d.
128	930	n.d.
129	23	n.d.
130	14	n.d.
131	170	n.d.
132	1.7	6.2


133	0.53	1.8
133a (peak 1)	0.23	2.9
133b (peak 2)	16	n.d.
134	0.78	4.7
134a (peak 1)	0.64	4.2
134b (peak 2)	1500	n.d.
135	0.38	2.4
136	0.13	0.8
137	3.2	22.7
138	1.0	7.6
139	0.44	3.1
140	3000	n.d.
141	7.0	n.d.
142	54	n.d.
143	0.89	4.1
144	140	n.d.
145	72	n.d.
146	1.3	4.0
147	2.7	n.d.
148	5.6	n.d.
149	4.2	n.d.
150	4.7	n.d.
151	1.9	2.5
152	1.2	1.8
153	6.1	165.2
154	1.2	1.5
155	0.96	n.d.
156	0.55	6.8
157	0.38	20
158	2.9	n.d.
159	1	4.5
160	0.49	1.3
161	4.2	n.d.
162	2.1	3.2


163	7.3	n.d.
164	1.6	3.9
165	1.6	19.1
166	2.7	n.d.
167	2.4	n.d.
168	3.2	n.d.
169	0.15	7
170	3.8	33.7
171	1.4	10.9
172	1.1	0.8
173	0.81	2.1
174	4.8	n.d.
175	1.2	17.6
176	3	15.1
177	5.7	3.9
178	46	>1000
179	48	n.d.
180	87	n.d.
181	25	n.d.
182	124	n.d.
183	1.2	2.9
184	4.7	6.7
185	4.3	4.4
186	1.1	10.3
187	2.1	7.3
188	1.1	8.5
189	42	n.d.
190	38	n.d.
191	0.67	n.d.
192	0.91	n.d.
193	3.9	n.d.
194	45	n.d.
195	8.0	n.d.
196	1.1	2.0


n.d.: No data.

[0643] Table III.


















1. A compound of formula:



A is a 5- or 6-membered aromatic ring comprising 0-3 heteroatoms of N, S or O; each W is independently $-(CH_2)-$ or $-C(O)-$;

S/D is a single or double bond, and when a double bond, R⁵ and R⁷ are absent;

n is 0, or an integer of 1-4, wherein when n is more than 1, each R^2 may be different;

p is 0, or an integer of 1-2, wherein when p is 0, m is non-zero, and when p is more

$R^1, R^4, R^5, R^6,$ and R^7 are each independently H, halogen, heteroalkyl, alkyl, alkenyl,

R¹, R⁴, R⁵, R⁶, and R⁷ are each independently H, halogen, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, alkynyl, -CN, -NR¹³R¹⁴, -OR¹³, -COR¹³, -CO₂R¹³, -CONR¹³R¹⁴, -C(=NR¹³)NR¹⁴R¹⁵, -NR¹³COR¹⁴, -NR¹³CONR¹⁴R¹⁵, -NR¹³CO₂R¹⁴, -SO₂R¹³, -NR¹³SO₂NR¹⁴R¹⁵, or -NR¹³SO₂R¹⁴, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶, wherein (R⁴ and R⁵), or (R⁴ and R⁶), or (R⁶ and R⁷), or (R⁶ and R⁶ when p is 2), together with the atoms to which they are attached, can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶;

R^2 is halogen, alkyl, -S-alkyl, -CN, -NR¹³R¹⁴, -OR¹³, -COR¹³, -CO₂R¹³, -CONR¹³R¹⁴, -C(=NR¹³)NR¹⁴R¹⁵, -NR¹³COR¹⁴, -NR¹³CONR¹⁴R¹⁵, -NR¹³CO₂R¹⁴, -SO₂R¹³, -NR¹³SO₂NR¹⁴R¹⁵, or -NR¹³SO₂R¹⁴;

R¹² is H or lower alkyl;


R¹³, R¹⁴ and R¹⁵ are each independently H, heteroalkyl, alkyl, alkenyl, alkynyl, cycloalkyl, saturated or unsaturated heterocyclyl, aryl, or heteroaryl; wherein (R¹³ and R¹⁴), and/or (R¹⁴ and R¹⁵) together with the atom(s) to which they are attached, each can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶;

R¹⁶ is halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocyclyl, oxo, -CN, -OR', -NR'R'', -COR', -CO₂R', -CONR'R'', -C(=NR'')NR''R''', -NR'COR'', -NR'CONR'R'', -NR'CO₂R'', -SO₂R', -SO₂aryl, -NR'SO₂NR''R''', or -NR'SO₂R'', wherein R', R'', and R''' are independently hydrogen, halogen, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl,

substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted heterocyclyl, wherein (R' and R''), and/or (R''' and R''') together with the atoms to which they are attached, can form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings.

2. The compound of claim 1, comprising one or more embodiments:

- (a) S/D is a double bond and R⁵ and R⁷ are absent;
- (b) R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶;
- (c) p is 1 and m is 0, 1 or 2;
- (d) A is phenyl;
- (e) each R² is independently halogen, lower alkyl, or lower alkoxy;
- (f) R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶;
- (g) R⁴ and R⁶, together with the atoms to which they are attached, form a ring of formula:




Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4;


- (h) S/D is a single bond; and/or
- (i) p is 0 and R⁶ and R⁷ are absent.

3. The compound of claim 1, wherein:

- (i) S/D is a double bond and R⁵ and R⁷ are absent; R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶; and R¹⁶ is halogen, lower alkyl, or lower alkoxy; or
- (ii) S/D is a double bond and R⁵ and R⁷ are absent; p is 1 and m is 0, 1 or 2; or
- (iii) S/D is a double bond and R⁵ and R⁷ are absent; p is 1 and m is 0, 1 or 2; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy; or
- (iv) S/D is a double bond and R⁵ and R⁷ are absent; and R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; or
- (v) S/D is a double bond and R⁵ and R⁷ are absent; R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy; or
- (vi) S/D is a double bond and R⁵ and R⁷ are absent; R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; p is 1 and m is 0, 1 or 2; A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy; and the R⁴-R⁶ ring is of formula:



wherein Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4; or
 (vii) S/D is a double bond and R⁵ and R⁷ are absent; R⁴ and R⁶, together with the atoms to which they are attached, form a ring selected from cycloalkyl, saturated or unsaturated heterocycle, aryl, and heteroaryl rings optionally substituted with at least one substituent R¹⁶; p is 1 and m is 0, 1 or 2; A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy; and the R⁴-R⁶ ring is of formula:



wherein Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4; and R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶; or

(viii) S/D is a single bond; p is 1 and m is 0, 1 or 2; A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy; or

(ix) S/D is a single bond; p is 1 and m is 0, 1 or 2; A is phenyl; each R² is independently halogen, lower alkyl, or lower alkoxy, and R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶; and R¹⁶ is halogen, lower alkyl, or lower alkoxy; or

(x) S/D is a single bond; p is 0 and R⁶ and R⁷ are absent; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy; or


(xi) S/D is a single bond; p is 0 and R⁶ and R⁷ are absent; A is phenyl; and each R² is independently halogen, lower alkyl, or lower alkoxy, and R¹ is H, halogen, alkoxy, heteroalkyl, alkyl, alkenyl, cycloalkyl, aryl, saturated or unsaturated heterocyclyl, heteroaryl, wherein the alkyl, alkenyl, alkynyl, cycloalkyl, heteroaryl, aryl, and saturated or unsaturated heterocyclyl are optionally substituted with at least one substituent R¹⁶.

4. The compound of claim 1, wherein:

S/D is a double bond and R⁵ and R⁷ are absent; p is 1 and m is 0, 1 or 2; A is phenyl;

S/D is a double bond and R⁵ and R⁷ are absent; p is 1 and m is 0, 1 or 2; A is phenyl;

and R⁴ and R⁶, together with the atoms to which they are attached, form a ring of formula:




wherein Q is -CH₂-; J is -CH₂-; and d and b are each independently 0, or an integer of 1-4;

S/D is a single bond; p is 1 and m is 0, 1 or 2; A is phenyl; or

S/D is a single bond; p is 0 and R⁶ and R⁷ are absent; A is phenyl.

5. The compound of claim 1, wherein:

A is phenyl; W is $-(CH_2)-$; L is O; S/D is a single bond; m is 1; n is 0; p is 1; R^1 is phenyl; R^2 is absent; R^5 is H; and R^6 and R^7 are H; yielding the combination structure:




6. The compound of claim 5 wherein R_4 is N-containing C1-C8 alkyl, N-containing C3-C8 cycloalkyl and phenyl, each optionally substituted.


7. The compound of claim 5 wherein R_4 is methylamine, aniline, azetidine, pyrrolidine, piperidine, azacycloheptenyl, each N-substituted with a moiety selected from benzyl, acyl, acryloyl, substituted-acryloyl, propiolyl, and substituted-propiolyl.

8. The compound of claim 5 wherein R_4 is selected from structures:


9. The compound of claim 5 wherein R^4 is 1-acryloylpiperidin-4-yl (compound 27), of formula:




10. The compound of claim 5 wherein R⁴ is 1-acryloylpiperidin-4-yl (compound 27), enantiomeric stereoisomer S of structure:




11. The compound of claim 5 wherein R⁴ is 1-acryloylpiperidin-4-yl (compound 27), enantiomeric stereoisomer R of structure:





12. The compound of claim 5 wherein R⁴ is 1-(but-2-ynoyl)piperidin-4-yl (compound 176):






















13. A compound according to claim 1 selected from the group consisting of, stereoisomers thereof, and pharmaceutically acceptable salts thereof:



























 34	 35	 36
 37	 38	 39
 40	 41	 42
 43	 44	 45
 46	 47	 48
 49	 50	 51
















 67	 68	 69
 70	 71	 72
 73	 74	 75
 76	 77	 78
 79	 80	 81
 82	 83	 84



























14. A compound according to claim 1 selected from the group consisting of, stereoisomers thereof, and pharmaceutically acceptable salts thereof:









































 107	 108	 109
 110	 111	 112
 113	 Cis 114	 Cis 115
 Cis 116	 117	 118













 119	 120	 121
 122	 123	 124
 HCOOH 125	 126	 127
 128	 129	 130








 131	 132	 133
 133a (peak 1)	 133b (peak 2)	 134
 134a (peak 1)	 134b (peak 2)	 135
 136	 137	 138

 139	 140	 141
 142	 143	 144
 145	 146	 147
 148	 149	 150


 151	 152	 153
 154	 155	 156
 157	 158	 159
 160	 161	 162


 163	 164	 165.
 166	 167	 168
 169	 170	 171
 172	 173	 174


 175	 176	 177
 179	 180	 183
 184	 185	 186
 187	 188	 189


 190	 191	 192
 193	 194	 195
 196.		


15. A compound according to claim 1 selected from the group consisting of, stereoisomers thereof, and pharmaceutically acceptable salts thereof:

















16. The compound of any of claims 1 to 15, having a Btk-inhibiting activity corresponding to an IC₅₀ of 10 μ M or less in a Btk Kinase Assay.
17. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any of claims 1 to 15 in unit dosage form and one or more pharmaceutically acceptable carriers.
18. A combination comprising a therapeutically effective amount of a compound of any of claims 1 to 15 and a different agent therapeutically active against an autoimmune and/or inflammatory disease or cancer.
19. A compound for use in treating a disease associated with undesirable Btk activity, which comprises administering to a person in need thereof an effective amount of a compound of any of claims 1-15, wherein the disease is an allergic disease, an autoimmune disease, an inflammatory disease, or cancer.
20. The compound of claim 19 wherein the disease is a B-cell proliferative disorder, selected from chronic lymphocytic lymphoma, non-Hodgkin's lymphoma, diffuse large B cell lymphoma, mantle cell lymphoma, follicular lymphoma and chronic lymphocytic leukemia.