FIG.1

Pure tumor cell culture in flasks
Interferon-gamma treatment for 72 hours
Floating non-autophagic, apoptotic cells
Adherent autophagic, non-apoptotic cells
Tissue culture flask surface
Detached non-autophagic, apoptotic cell
Attached autophagic,non-apoptotic cell
Major histocompatibility complex II-over expressed in autophagic cells
Phosphatidylserine-apoptotic marker



7-AAD-/Annexin-V+(Dying)
7-AAD-/Annexin-V-(Alive)
7-AAD+/Annexin-V+(Dead)
7-AAD+/Annexin-V-(Necrotic)
Percent
100
80
60
40
20
0
IFN-γ - + - + - + - +
2046
2270
2028
2020

FIG. 2A

FIG.2B



B
2048
2270
2028
2020
γIFN
Hsp-60
0.89
0.89
0.88
1.01
Hsp-70
1.26
5.77
1.74
1.28
Hsp-90
0.85
0.50
0.73
1.34
Calreticulin
0.50
1.17
1.44
0.44
HMGB-1
0.84
0.85
2.85
279.0
GADPH

FIG.2C



C
2046
2270
2028
2020
γIFN
ICAM-1
3.68
2.38
2.64
4.80
Mel-4
0.50
0.20
0.34
0.94
Mart-1
0.58
0.25
0.82
0.44
Tyrosinase
0.27
0.88
0.89
1.17
GAPDH

FIG.2D




D
Mean Fluorescence Intensity
6000
5000
4000
3000
2000
1000
0
195 +/- 86
671 +/- 124
77 +/- 199
376 +/- 229
IFN-γ
–
+
–
+
MHC Class I
MHC Class II

FIG.3A






DC+TC
DC+IFN-γ TC
Percent MHC Class II Positive
100
80
60
40
20
0
Isotype
CD80
CD83
CD86

FIG. 3B

FIG.4A







CD86
%
100
80
60
40
20
0
CD86
MFI
5000
4000
3000
2000
1000
0
CD83
%
*
20
15
10
5
0
CD83
MFI
**
300
250
200
150
100
50
0
Pre-ATC Load DC
Post-ATC Load DC
CD80
%
60
50
40
30
20
10
0
CD80
MFI
350
300
250
200
150
100
50
0

FIG. 4B



No IFN-γ IFN-γ

FIG. 5A



No IFN-γ IFN-γ

FIG. 5B

FIG.5C




FIG.6A




A
5% FBS
IFN-γ Adherent Cells
IFN-γ Floating Cells
Serum Free
7-AAD
Annexin-V
R3H

FIG.6B


B
Counts
Autophagy Dye
5% FBS
IFN-γ Adh.
IFN-γ Flts.
SF
0
20
40
60
80
100
10^0
10^1
10^2
10^3
10^4



Fold Change
2.5
2.0
1.5
1.0
0.5
0
IFN-γ Adherent Cells
IFN-γ Floating Cells
Serum Free

FIG. 6C




FIG. 7A





FIG. 7B

FIG.8





Mean Fluorescence Intensity
800
700
600
500
400
300
200
100
0
Non-apoptotic/MHC II+ pre-IFNg
Apoptotic/MHC II+ pre-IFNg
Non-apoptotic/MHC II+ post IFNg
Apoptotic/MHC II+ post IFNg

FIG. 9



Fold change MHC II
16
12
8
4
0
0
5
10
15
20
25
30
35
40
% Annexin-V Positive

FIG.10

FIG.11




Progression Free Survival
IFN-γ resistant
IFN-γ sensitive
1.0
0.8
0.6
0.4
0.2
Proportion
p = 0.003
0
6
12
18
24
30
36
42
48
54
60
Months

FIG.12


Overall Survival
1.0
0.8
0.6
0.4
0.2
Proportion
p = 0.002
0
6
12
18
24
30
36
42
48
54
60
Months



Superimposed survival curves from 3 trials of patient-specific vaccines
DC-54
TC-74
TC-24
DC-18
Percent Surviving
Months
0
10
20
30
40
50
60
70
80
90
100
0
6
12
18
24
30
36
42
48
54
60
66
72

FIG. 13