DESCRIPTION

Title of the Invention

ANTIBODIES FOR DETECTING MICROORGANISMS

DESCRIPTION OF BACKGROUND ART

Field of the Invention

The present invention pertains to antibodies useful in the detection of various microorganisms, particularly bacteria, and a method of detecting microorganisms, reagent kits for detection of microorganisms that use said antibodies, and a method for preparing specific antibodies for detecting microorganisms.

Moreover, the present invention is valuable to the drug industry, particularly for diagnostics for microorganism infections with an emphasis on bacteria.

Prior Art

Diagnosis of microorganism infection is confirmed by detection of the causative pathogen from the infected area or by detection of antibodies to the causative pathogen in serum and body fluids. Detection of the causative pathogen is particularly important in the sense that it makes quick treatment of the patient possible.

Detection of the causative pathogen of infections

This Sheet Has Been Replaced

can be generally classified as cultivation and identification methods, whereby the causative pathogen is Selectively cultivated and then identified based on its biochemical properties; genetic diagnosis, whereby amplification by PCR, etc., is performed based on specific DNA sequence of the causative pathogen and thus the causative pathogen is detected; or immunological methods, whereby the causative pathogen is detected using a specific reaction of antibody with surface antigen marker of the causative pathogen. However, it takes time to obtain results by cultivation and identification methods or genetic diagnosis methods. Therefore, diagnosis by immunological methods is commonly used because the causative pathogen can be detected within a short time with high sensitivity and the patient can be quickly and appropriately treated.

Depending on the bacterial species, a variety of marker antigens and antibodies can be used, alone or in combination to detect the causative pathogen of infections by conventional immunological methods.

For instance, lipopolysaccharide (LPS) represents a genus-specific antigen of *Chlamydia*, (Stephens, R., et al.: *J. Immunol.*, 128:1083-89, 1982, Caldwell, M.D.: *Inf. Immun.*, 44:306-14, 1984), and antibodies to LPS are used as the reagent antibody in a variety

This Sheet Has Been Replaced

of diagnostic kits, particularly for the detection of *Chlamydia trachomatis*.

Moreover, Peterson et al., (*Infection and Immunity*, 59(11), 4147-4153, 1991) and Batteiger et al., (Infection and Immunity, 53(3), 530-533, 1986) have reported monoclonal antibodies to major outer membrane protein (MOMP) of the genus *Chlamydia*.

Publication of unexamined Japanese application No. 298/1988 discusses an immunodetection method based on the western blotting that uses a monoclonal antibody to an approximately 43 kilo Dalton membrane protein antigen of *Mycoplasma pneumoniae*.

A method of preparing monoclonal antibody to *Haemophilus influenzae* and a diagnostic method that uses said antibody are presented in Publication of unexamined Japanese application No. 148859/1987 (Japanese Patent No. 64065/1994).

Proteins of approximately 20 kilo Dalton isolated from the sodium cholate extract of the outer membrane vesicle of *Neisseria gonorrhoeae* strain BS4 (NCTC 11922) are documented and preparation of hybridomas using said substance is disclosed in British Patent Application No. 2,172,704. Moreover, EP 419238A1 describes preparation of a monoclonal antibody, which can bind to a protein of approximately 14 kilo Dalton prepared by using *Neisseria gonorrhoeae* as an immunogen and a method for the preparation of such a

This Sheet Has Been Replaced

monoclonal antibody. Additionally, a method of detecting the same *Neisseria gonorrhoeae* using monoclonal antibody to lipopolysaccharide (LPS) is mentioned in Canadian Patent Application No. 1,220,147.

However, there are problems with said antibodies and detection methods based on those antibodies in that species specificity to microorganisms is insufficient for proper diagnosis. The antibodies do not detect all serum types as different types of surface antigen present within one species.

Marker antigens mentioned in these prior arts are different in terms of their function, structure and cellular locations. Furthermore, they are not evolutionarily conserved and often not resistant to antigenic drift. Antigens like these might have advantage for serotyping but not for routine laboratory diagnosis where a common antibody is needed to detect all of the members within the species. A common antigen that is with common function that is evolutionarily protected but have measurable difference from species to species was not described hitherto.

DISCLOSURE OF THE INVENTION

The present invention strives to make ideal microorganism detection and immunodiagnosis possible

This Sheet Has Been Replaced

and with innovated antibody to the same molecule in different microorganisms. Particularly antibodies to the molecular segment that is a part of same intracellular functional component of all microorganisms. More intensely, it describes an antibody to detect the evolutionary signature within a molecule that is common in all the member of a species but differ form species to species. A method of detecting microorganisms, which is species specific and can cover almost all serum types, a reagent kit for detection of microorganisms that use said antibody, and a method for preparing a specific antibody used for detection of microorganisms.

The present invention describes a protein with the same function, preserved in all microorganisms as a useful antigen protein. Usually, it is expected that such a protein be subjected to a structural change only to a small extent. However, it was found that the antigen epitope of this protein has specificity to certain species or genus of microorganisms, and that the antibody for this protein not only has potentialities of being used for specifically identifying species or genus of microorganisms, but also is capable of detecting all serotypes of the subject microorganisms.

The inventors focused on intracellular molecule that are present in all microbial cells and differ

This Sheet Has Been Replaced

between microorganisms in terms of its amino acid sequence, particularly ribosomal protein L7/L12, which is a member of ribosomal protein. Ribosomal protein L7/L12 is a protein with a molecular weight of approximately 13 kilodaltons and is known to exist as an essential ribosomal protein in protein synthesis. Progress has been made in understanding the complete amino acid sequence of several microorganisms including *Escherichia coli and Baccillus subtilus*, etc., and 50% to 65% homology of the amino acid sequence between the microorganisms has found.

The inventors focused on the fact that even though there are similarities between different microorganisms in terms of said molecule, this molecule also has a structural segment that is unique to each microorganism and discovered that it is possible to detect various microorganisms with species specificity and to detect all serotypes within the same species by using antibody to said protein. As a result of attempting to develop a technology for immunodiagnosis of microorganism species using antibody specific to, for instance, *Haemophilus influenzae, Streptococcus pneumoniae,* and *Neisseria gonorrhoeae*, the inventors completed the present invention upon discovering that antibody specific to said protein of each species of

This Sheet Has Been Replaced

microorganisms can be obtained and species-specific detection of different bacteria is possible using said antibody.

The present invention relates to an antibody used for detecting microorganisms, a method of detecting microorganisms using the antibody, a reagent kit for detecting microorganisms using the antibody, and a method for preparing specific antibodies for detecting microorganisms.

1) Antibodies which are antibodies to ribosomal protein of microorganisms and which react specifically with said microorganisms.

2) The antibodies described in 1) above, where the ribosomal protein of the microorganisms is ribosomal protein L7/L12.

3) The antibodies described in 1) or 2) above, where said microorganisms are microorganisms which cause a sexually transmitted disease (STD).

4) The antibodies described in 1) or 2) above, where said microorganisms are microorganisms which cause respiratory tract infection.

5) The antibody described in 4) above, where the causative microorganisms of respiratory tract infection are microorganisms of *Haemophilus influenzae*.

6) The antibody described in 4) above, where the causative microorganisms of respiratory tract

This Sheet Has Been Replaced

infection are microorganisms of *Streptococcus pneumoniae*.

7) The antibody described in 3) above, where the causative microorganisms of sexually transmitted diseases (STD) are microorganisms of *Neisseria gonorrhoeae*.

8) The antibody described in 7) above, which is the antibody to ribosomal protein L7/L12 of *Neisseria gonorrhoeae* and which identifies a amino acid sequence moiety from 5 to 30 amino acids including the 115th alanine in the amino acid sequence of Sequence ID No. 22 of the Sequence Table.

9) A method of detecting microorganisms, which is characterized by the fact that antibody to intracellular molecules that have the same function for a variety of microorganisms is used.

10) A method of detecting microorganisms, which is characterized by the fact that any antibody described in 1) to 8) above is used.

11) A reagent kit for detecting microorganisms, which is characterized by the fact that antibody to intracellular molecules that have the same function for a variety of microorganisms is used.

12) A reagent kit for detecting microorganisms, which is characterized by the fact that any antibody described in 1) to 8) above is used.

13) A method of preparing any antibody described

This Sheet Has Been Replaced

in 1) to 8) above, characterized by the fact that ribosomal protein L7/L12 of microorganisms obtained by a gene manipulation procedure or by isolation from microorganisms, peptide moiety thereof, or a synthesized peptide corresponding to the peptide moiety is used as an immune source.

The present invention will now be explained in detail.

Sequences No. 1 and No. 2 in the Sequence Table are the DNA sequence of the ribosomal protein L7/L12 gene of *Hemophilus influenzae* and corresponding amino acid sequence. Sequences No. 3 and No. 4 are the DNA sequence of the ribosomal protein L7/L12 gene of *Helicobacter pylori* and the corresponding amino acid sequence. Sequences No. 5 and No. 6 show the DNA sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Streptococcus pneumoniae*. Sequences No. 7 and No. 8 show the DNA sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Neisseria gonorrhoeae*. Sequences No. 9 and No. 10 show the DNA sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Neisseria meningitidis*. Sequence No. 11 and Sequence No. 12 in the Sequence Table are the primers DNA for PCR used to acquire the ribosomal protein L7/L12 gene from *Hemophilus influenzae*. Sequences No. 13 and No. 14

This Sheet Has Been Replaced

in the Sequence Table are the primer DNA for PCR used to acquire the ribosomal protein L7/L12 gene from *Streptococcus pneumoniae*. Sequences No. 15 and No. 16 in the Sequence Table are the primer DNA for PCR used to acquire the ribosomal protein L7/L12 gene from *Neisseria gonorrhoeae*. Sequences No. 17 and No. 18 show the DNA sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Hemophilus influenzae*. Sequences No. 19 and No. 20 show the DNA sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Streptococcus pneumoniae*. Sequences No. 21 and No. 22 show the DNA sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Neisseria gonorrhoeae*.

Furthermore, the left and right terminals of the amino acid sequences entered in the Sequence Table are amino group (referred to below as the N terminal) and carboxyl group terminals (referred to below as the C terminal), respectively, and the left terminal and right terminal of the base sequence is the 5' terminal and the 3' terminal, respectively.

Moreover, the series of biomolecular experiments of gene preparation mentioned in this text can be performed by methods entered in standard experimental manuals. "Molecular cloning: A laboratory manual," Cold Spring Harbor Laboratory Press, Sambrook, J. et

This Sheet Has Been Replaced

al. (1989), is an example of the aforementioned standard experimental manual.

The term microorganism in the present invention refers to all species of microorganisms, including bacteria, yeast, mold, *Actinomyces*, rickettsia, etc., but bacteria in particular pose a problem in terms of diagnosis of microbial infections.

The term "antibody which reacts specifically with microorganisms" as used in the present invention means an antibody, which reacts specifically with a species or group of microorganisms. An antibody, which reacts specifically with a species of microorganisms, is particularly useful for the diagnosis of microbial infection diseases.

In the present invention, causative microorganisms of STD (sexually transmitted disease) include, but are not limited to, *Neisseria gonorrhoeae, Chlamydia trachomatis, Candida albicans, Treponema pallidum,* and *Ureaplasma urealyticum.*

In the present invention, causative microorganisms of respiratory tract infection include, but are not limited to, *Haemophilus influenzae, Streptococcus pneumoniae, Chlamydia pneumoniae, Mycoplasma pneumoniae, Klebsiella pneumoniae, Staphylococcus aureus, Pseudomonas aeruginosa, Streptococcus sp. GroupA, Mycobacterium tuberculosis, Legionella pneumophila,* and *Aspergillus*

This Sheet Has Been Replaced

spp.

The term antibody in the present invention means a polyclonal antibody or monoclonal antibody that can be made using the entire length or only a partial peptide of said ribosomal protein. Although there are no special restrictions to the peptide length for making the antibody, the segment should be of the length characterizing the ribosomal protein L7/L12, and a peptide of 5 amino acids or longer, particularly 8 amino acids or longer, is preferred. Antiserum containing antibody (polyclonal antibody) that identifies ribosomal protein L7/L12 can be obtained by inoculating laboratory animals with adjuvant and a peptide or the full length protein as it is or when necessary, after being crosslinked with a carrier protein such as KLH (keyhole-limpet hemocyanin) and BSA (bovine serum albumin) and recovering the serum. Moreover, the antibody can be used after it has been purified from the antiserum. The laboratory animals that are inoculated include sheep, horses, goats, rabbits, mice, rats, etc., and sheep, rabbits, etc., are particularly preferred for preparation of polyclonal antibody. Moreover, monoclonal antibody can also be obtained by conventional methods of making hybridoma cells, but mice are preferred in this case. The entire length of said protein, or its partial peptide consisting

This Sheet Has Been Replaced

amino acid residues of 5 or more, preferably 8 or more, residues that has been fused with GST (glutathione S-transferase), etc., can be purified and used as antigen, or it can be used as antigen without being purified. The antibody can also be the genetic recombination antibody expressed cellularly using immunoglobulin genes that have been separated by a variety of methods in manuals ("Antibodies: A Laboratory manual," E. Harlow et al., Cold Spring Harbor Laboratory), cloning methods, etc.

Antibody to ribosomal protein L7/L12 that can be employed as the marker antigen of the present invention can be obtained by the following 3 methods, and other similar methods as well; the methods, however, are not limited to these.

a) The desired antibody can be acquired by synthesizing a peptide fragment consisting of 5-30 amino acids for microorganisms with a known ribosomal protein L7/L12 genetic sequence and amino acid sequence using the region least similar to the amino acid sequence of said protein of another microorganisms and making polyclonal antibody, or monoclonal antibody, using this peptide fragment as the immune source.

Moreover, it is possible to acquire the entire sequence of said gene by using a conventional genetic procedure, such as gene amplification by PCR with the

This Sheet Has Been Replaced

DNA sequence at both terminals of said known genetic sequence as the primer, or hybridization using the sequence of a homologous segment as the template probe.

Then, a fused gene with another protein gene is constructed and said fused gene is inserted into the host by conventional gene insertion methods using *Escherichia coli*, etc., as the host and expressed in large quantities. The desired protein antigen can then be acquired by purifying the expressed protein by affinity column methods with antibody to the protein that was used as the fusion protein. In such a case, even if antibody to the amino acid segment retained within the bacteria is acquired, it does not coincide with the purpose of the present invention because the full length of ribosomal protein L7/L12 becomes the antigen. Consequently, hybridoma that produces monoclonal antibody to the antigen that has been obtained by this method is acquired by conventional methods and the desired antibody can be obtained by selecting a clone that produces antibody that will react only with the desired microorganisms.

b) First, since there is 50 to 60% homology between bacteria species in terms of their ribosomal protein L7/L12 amino acid sequence, it is possible to easily acquire said protein genes for microorganisms with unknown L7/L12 amino acid sequence by

This Sheet Has Been Replaced

conventional genetic procedures, such as gene amplification of a specific sequence segment by PCR methods based on the sequence of the homologous segments of the amino acid sequence, or hybridization with the homologous segments as the template probe, using bacteria having a known ribosomal protein L7/L12 amino acid sequence.

Then, a fused gene with another protein gene is constructed and said fused gene is inserted into the host by conventional gene insertion methods using *Escherichia coli*, etc., as the host and expressed in large quantities. The desired protein antigen can then be acquired by purifying the expressed protein by affinity column methods with antibody to the protein that was used as the fusion protein. In such a case, even if antibody to the amino acid segment retained within the bacteria is acquired, it does not coincide with the purpose of the present invention because the full length of ribosomal protein L7/L12 becomes the antigen. Consequently, hybridoma that produces monoclonal antibody to the antigen that has been obtained by this method is acquired by conventional methods and the desired antibody can be obtained by selecting a clone having ability antibody that will react only with the desired microorganisms.

c) Ribosomal protein L7/L12 that has been purified to a high purity can also be obtained by

This Sheet Has Been Replaced

another method, in the case where the amino acid sequence of the ribosomal protein L7/L12 is unknown whereby a peptide of 5 to 30 amino acids corresponding to the common sequence segment retained in the microorganisms is synthesized from the known amino acid sequence of the ribosomal protein L7/L12, and polyclonal antibody or monoclonal antibody to this peptide sequence is made by conventional methods. Then the highly purified Ribosomal protein L7/L12 from disrupted microorganisms is obtained by affinity column chromatography using said antibody.

If purity of the protein is insufficient, it can be purified by conventional methods, such as ion exchange chromatography, hydrophobic chromatography, gel filtration, etc., after which the eluted fraction of ribosomal protein L7/L12 is identified by western blotting, etc., using antibody that was made to obtain the pure fraction. The desired antibody can be obtained by acquiring hybridoma or polyclonal antibody by conventional methods using the pure ribosomal protein L7/L12 antigen that has been obtained and selecting hybridoma or polyclonal antibody that will react specifically with the desired bacteria, as in b).

The antibody of the present invention specific to a variety of microorganisms that has been obtained by the methods in a) through c), etc., can be used in a

This Sheet Has Been Replaced

variety of immunoassay methods to obtain diagnostic reagent kits specific to a variety of microorganisms. For example, this antibody can be used in aggregation reactions where antibody is adsorbed on polystyrene latex particles, ELISA, which is a conventional technology performed in a microtiter plate, conventional immunochromatography methods, sandwich assay, whereby said antibody labeled with colored particles or particles that have coloring capability, or with enzyme or phosphor, and magnetic microparticles coated with capture antibody, etc., are used, etc.

The term detection methods for microorganism using antibody means detection methods that use conventional immunoassay such as aggregation reactions where antibody is adsorbed on polystyrene latex particles, ELISA, which is a conventional technology performed in a microtiter plate, conventional immunochromatography methods, sandwich assay, whereby said antibody labeled with colored particles or particles that have coloring capability, or with enzyme or phosphor, and magnetic microparticles coated with capture antibody, etc., are used, etc.

Moreover, the optical immunoassay (OIA) technology described in International Patent Application Japanese Laid-open (Toku-Hyou) No.

This Sheet Has Been Replaced

509565/1995, in which microorganisms are detected by an optical interference induced by an antibody reaction on the optical thin film which is formed by silicone or silicon nitride, is a useful detection method using an antibody.

Moreover, treatment with an extraction reagent that uses a variety of surfactants, beginning with Triton X-100 and Tween-20, enzyme treatment with an appropriate enzyme, such as protease, etc., methods whereby the known cell structure is disrupted, beginning with disruption of the microorganism by physical methods, can be used to extract the intracellular marker antigen from the necessary microorganism in said detection method. However, it is preferred that the extraction conditions be designed using a combination of surfactants, etc., so that the conditions are optimized for extraction of the bacteria in question with a reagent.

Moreover, the term a reagent kit for detection of microorganisms using antibody means a reagent kit that uses said detection method.

For instance, in the case of obtaining the specific antibody to *Haemophilus influenzae*, which is of extreme diagnostic significance as a causative pathogen of pneumonia, bronchitis, meningitis, etc., the amino acid sequence and DNA sequence of ribosomal protein L7/L12 is entered in data bases, etc..

This Sheet Has Been Replaced

The amino acid and DNA sequence of ribosomal protein L7/L12 of *Haemophilus influenzae* are shown in "Sequences No. 1 and No. 2."

Consequently, in the case of this bacteria, it is possible to similarly compare the amino acid sequence of ribosomal protein L7/L12 with the same protein of, for instance, *Helicobacter pylori*, which is shown in "Sequence No. 3 and No. 4," and synthesize a peptide of 5 to 30 amino acids for the segment of low homology and make polyclonal antibody or monoclonal antibody specific to *Haemophilus influenzae* using this peptide.

In the case of a specific polyclonal antibody, it is preferred that IgG fraction be obtained by purification of the antiserum of immunized laboratory animals with a protein A column, etc., and affinity purification be performed with the synthetic peptide used in immunization of the laboratory animals.

Moreover, PCR primers based on the sequences of N-terminal and C-terminal, for example, the PCR primers shown in Sequences No. 11 and No. 12 in the Sequence Table, were designed from the DNA sequence of ribosomal protein L7/L12 of *Haemophilus influenzae*. Utilizing homology of the PCR primers, DNA fragments amplified by the PCR method or the like using genomic DNA which is extracted from cultivated *Haemophilus influenzae* can be acquired according to a

This Sheet Has Been Replaced

conventional method. The entire length of the gene for ribosomal protein L7/L12 of *Haemophilus influenzae* can be acquired by the analysis of the DNA sequence information of these fragments.

The ribosomal protein L7/L12 gene of *Haemophilus influenzae* thus acquired forms a fusion protein gene with, for instance, GST etc., and expression vector is constructed using the appropriate expression plasmid, *Escherichia coli* is transformed and large quantities of said protein can be expressed. A suitable amount of the transformed *Escherichia coli* is cultivated and the crushed bacterial fluid that is recovered is ~~submitted to~~ purified by an affinity column using GST to obtain the ribosomal protein L7/L12 and GST fusion protein of *Haemophilus influenzae*. It is also possible to acquire the target specific monoclonal antibody by establishing a number of hybridomas using said protein as it is or GST moiety fragments cut from the protein by protease or the like, as an antigen protein and selecting the antibody which exhibits a specific response to *Haemophilus influenzae* bacteria, or a extract of the bacteria, or ribosomal protein L7/L12 of *Haemophilus influenzae*.

Moreover, the amino acid sequence and the DNA sequence of ribosomal protein L7/L12 of *Streptococcus pneumoniae* which is also highly significant as a

This Sheet Has Been Replaced

diagnostic agent for respiratory infection diseases as well as *Haemophilus influenzae*, are known from data bases and the like. The amino acid and DNA sequences of ribosomal protein L7/L12 of *Streptococcus pneumoniae* are shown in Sequences No. 5 and No. 6 of sequence table.

It is therefore possible to acquire a polyclonal antibody or monoclonal antibody which is specific to *Streptococcus pneumoniae* by designing a PCR primer, the PCR primer shown by Sequence ID No. 13 or 14 in the Sequence Table, for example, based on the sequences of N-terminal and C-terminal of DNA sequence of Ribosomal Protein L7/L12s of *Streptococcus pneumoniae* in the same manner as in the case of *Haemophilus influenzae*, and processing thereafter in the same manner as in the case of *Haemophilus influenzae*.

The hybridoma AMSP-2 which produces the monoclonal antibody specific to *Streptococcus pneumoniae* has been deposited with the National Institute of Bioscience and Human-Technology, the Agency of Industrial Science and Technology, the Ministry of International Trade and Industry, Japan, on July 28, 1999, with the deposition number FERM BP-6807.

Moreover, although the DNA and amino acid sequences of the ribosomal protein L7/L12 of, for

This Sheet Has Been Replaced

instance, *Neisseria gonorrhoeae*, which is the causative pathogen of gonorrhea and has been shown to have diagnostic significance as a typical causative pathogen of STD, were unknown, a major part of the DNA sequence and amino acid sequence, which was determined by the Neisseria Gonorrhea Genome Project at Oklahoma University, USA, is disclosed on the Internet.

When part of the known DNA sequence of ribosomal protein L7/L12 was used to probe the existence of DNA fragments with a similar sequence, it was found that DNA sequence corresponding to the ribosomal protein L7/L12 gene is present and it was possible to obtain data on its entire DNA sequence. The entire base sequence and corresponding amino acid sequence of the ribosomal protein L7/L12 gene of this *Neisseria gonorrhoeae* are shown in Sequences No. 7 and No. 8 of the sequence table.

It is therefore possible to acquire the target antibody which is specific to *Neisseria gonorrhoeae* having the entire or partial Ribosomal Protein L7/L12 of *Neisseria gonorrhoeae* as an antigen by designing a PCR primer, the PCR primer shown by Sequence ID No. 15 or 16 in the Sequence Table, for example, based on the sequences of N-terminal and C-terminal of DNA sequence of Ribosomal Protein L7/L12 of *Neisseria gonorrhoeae* in the same manner as in the case of

This Sheet Has Been Replaced

Haemophilus influenzae, and *Streptococcus pneumoniae*, and processing thereafter in exactly the same manner as in the case of *Haemophilus influenzae* or *Streptococcus pneumoniae*.

Particularly, the gene sequence of ribosomal protein L7/L12 of *Neisseria meningitides*, which belongs to the same *Neisseria* genus as *Neisseria gonorrhoeae* is disclosed and readily available on the Internet. The entire base sequence and the corresponding amino acid sequence of the ribosomal protein L7/L12 gene of *Neisseria meningitidis* are shown in "Sequences No. 9 and No. 10." Here, comparing the entire base sequence for ribosomal protein L7/L12 genes of *Neisseria meningitidis* and *Neisseria gonorrhoeae*, only difference in the amino acid sequence is that Neisseria gonorrhoeae has alanine for the 115th amino acid from the N-terminal, whereas *Neisseria meningitidis* has glutamic acid. Therefore, it can be concluded that the antibody for the ribosomal protein L7/L12 of *Neisseria gonorrhoeae* which can specifically detects *Neisseria gonorrhoeae* is the antibody which identifies alanine at 115 from the N-terminal and the amino acid region including the alanine of ribosomal protein L7/L12 as an epitope.

Antibody made based on the present invention can be used in all known types of immunoassay, such as

This Sheet Has Been Replaced

aggregation whereby said antibody is adsorbed on polystyrene latex, ELISA, which is a conventional technology performed in a microtiter plate, conventional immunochromatography, sandwich assay, whereby said antibody labeled with colored particles or particles that have coloring capability, or enzymes or phosphor, and magnetic particles coated with capture antibody are used, etc.

Moreover, antibody that is made based on the present invention can simultaneously function in any of these immunoassay methods as a so-called capture antibody that captures said antigen protein in solid or liquid phase and a so-called enzyme-labeled antibody by modification using an enzyme, such as peroxidase and alkali phosphatase, etc., by conventional methods.

BEST MODE FOR CARRYING OUT THE INVENTION

The following examples are given to explain the present invention in actual terms, the present invention not being restricted to these examples.

Example 1

Cloning of ribosomal protein L7/L12 genes from *Haemophilus influenzae*

After inoculating an appropriate amount of *Haemophilus influenzae* strain ATCC9334 (IID984) (obtained from Tokyo University School of Medicine Laboratories) in a chocolate agar culture medium, the

This Sheet Has Been Replaced

strain was cultivated for 24 hours in a CO_2 incubator under conditions of 37°C and 0.5% CO_2. The colonies that grew were suspended in a TE buffer (manufactured by Wako Pure Chemical Co., Ltd.) to a final concentration of approximately 5 x 10^9 CFU/ml. Approximately 1.5 ml of this suspension was transferred to a microcentrifuge tube and centrifuged for 2 minutes at 10,000 rpm. The supernatant was discarded. The sediment was resuspended in 567 µl TE buffer. Then 30 µl 10% sodium dodecylsulfate (SDS) and 3 µl 20 mg/ml Proteinase K solution were added and thoroughly mixed. The suspension was incubated for another hour at 37°C. Next, after adding 80 µl 10% cetyl trimethyl ammonium bromide/0.7 M NaCl solution and thoroughly mixing the product, it was incubated for 10 minutes at 65°C. Next, 700 µl chloroform-isoamyl alcohol solution at a volume ratio of 24:1 was added and stirred well. The solution was centrifuged for 5 minutes (while being kept at 4°C) at 12,000 rpm using a microcentrifugation device and the aqueous fraction was transferred to a new microtube. Isopropanol was added to the fraction at 0.6-times its volume and the tube was vigorously shaken to form sediment of the DNA. The white DNA sediment was scooped with a glass rod and transferred to a different microcentrifugation tube containing 1 ml 70% ethanol (cooled to -20°C).

This Sheet Has Been Replaced

Next, the product was centrifuged for 5 minutes at 10,000 rpm and the supernatant was gently removed. Then another 1 ml 70% ethanol was added and the product was centrifuged for 5 more minutes.

Once the supernatant had been removed, the sediment was dissolved in 100 µl TE buffer to obtain the DNA solution. The concentration of the genomic DNA solution was determined quantitatively in according to Sectrophotometric determination of the amount of DNA or RNA described else where (Molecular cloning: A laboratory manual," 1989, Eds. Sambrook, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press).

PCR (polymerase chain reaction) was performed using 10 ng of this genomic DNA. Taq polymerase (Takara Co., ltd., code R001A) was employed for PCR. Then 5 µl of buffer, 4 µl dNTP mixture, and 200 pmol each of synthetic oligonucleotide-A, shown in Sequence No. 11 of the Sequence Table and synthetic oligonucleotide-B, shown in Sequence No. 12 of the Sequence table were added to the enzyme to bring the final volume to 50 µl.

This mixture was cycled 5 times with a TAKARA PCR Thermal Cycler 480 for 1 minute at 95°C, 2 minutes at 50°C, and 3 minutes at 72°C and was then cycled 25 times for 1 minute at 95°C, 2 minutes at 60°C, and 3 minutes at 72°C. Electrophoresis was performed in

This Sheet Has Been Replaced

1.5% agarose gel using some of this PCR product. This product was then stained with ethidium bromide (Nihon Gene Co., ltd.) and observed under ultraviolet rays to confirm amplification of approximately 400 bp cDNA. After digestion treatment with restriction endonucleases BamHI and XhoI, electrophoresis was performed in 1.5% agarose gel and staining with ethidium bromide was carried out. An approximately 370 bp band was cut out from the gel. This band was purified with Suprecol (Takara Co., Ltd.) and then inserted into pGEX-4T-1 (Pharmacia), which is a commercial vector. This same vector can function as an expression vector for the desired molecule, which can express fused protein with GST protein, by insertion of the desired gene fragment into the appropriate restriction endonuclease site.

Actually, vector pGEX-4T-1 and the previous DNA were mixed together at a molar ratio of 1:3 and DNA was inserted into the vector with T4 DNA ligase (Invitrogen Co.). Vector pGEX-4T-1 into which DNA had been inserted was genetically introduced to *Escherichia coli* one-shot competent cells (Invitrogen Co., Ltd.) and then inoculated in a plate of LB-Agar (Takara Co., Ltd.) semi-sold culture plate containing 50μg/ml ampicillin (Sigma). The plate was then set-aside at 37°C for 12 hours and the colonies that grew were selected at random and inoculated into 2ml LB-

This Sheet Has Been Replaced

Broth liquid culture medium containing the same concentration of ampicillin. Shake cultivation was performed at 37°C for 7 hours and the bacteria were recovered and the plasmid was separated using Wizard Miniprep (Promega) in accordance with the attached literature. The plasmid was cleaved with restriction endonuclease BamHI/XhoI. Insertion of said PCR product was confirmed by cutting out approximately 370 bp DNA. The base sequence of the DNA that had been inserted was determined using said clone.

Determination of the base sequence of the inserted DNA fragment was performed using the Fluorescence Sequencer of Applied Biosystems. The sequence sample was prepared using PRISM, Ready Reaction Dye Terminator Cycle Sequencing Kit (Applied Biosystems). First, 9.5 µl reaction stock solution, 4.0 µl T7 promoter primer at 0.8 pmol/µl (Gibco BRL) and 6.5 µl of template DNA for sequencing at 0.16 µg/µl were added to a microtube with a capacity of 0.5 ml, mixed and superposed with 100 µl mineral oil. PCR amplification was performed for 25 cycles, where one cycle consisted of 30 seconds at 96°C, 15 seconds at 55°C, and 4 minutes at 60°C. The product was then kept at 4°C for 5 minutes. After the reaction was completed, 80 µl sterilized pure water was added and stirred. The product was centrifuged and the aqueous layer was extracted 3 times with phenol-chloroform.

This Sheet Has Been Replaced

Ten-microliter 3M sodium acetate (pH 5.2) and 300 µl ethanol were added to 100 µl aqueous layer and stirred. The product was then centrifuged for 15 minutes at room temperature and 14,000 rpm and the sediment was recovered. Once the sediment was washed with 75% ethanol, it was dried under a vacuum for 2 minutes to obtain the sequencing sample. The sequencing sample was dissolved in formamide containing 4 µl 10 mM EDTA and denatured for 2 minutes at 90°C. This was then cooled in ice and applied to sequencer.

One of the 5 clones obtained had homology of the sequence with the probe used for PCR. In addition, DNA sequences extremely similar to the gene sequence of ribosomal protein L7/L12 gene of the other microorganisms, for example, *Neisseria gonorrhoeae*, were discovered. The entire base sequence and the corresponding amino acid sequence of the structural gene moiety are as shown in Sequence ID No. 17 and No. 18 of the Sequence Table. This gene fragment clearly codes for *Haemophilus influenzae* ribosomal protein L7/L12.

Example 2

Mass expression in *Escherichia coil* and purification of ribosomal protein L7/L12 from *Haemophilus influenzae*

Fifty milliliters *Escherichia coli* into which

This Sheet Has Been Replaced

expression vector had been inserted were cultivated overnight in LB at 37°C. Then 500 ml YT medium at concentration that was twice that of the aforementioned culture was heated at 37°C for 1 hour. Fifty milliliters of the *Escherichia coli* solution that had been cultivated overnight were introduced to 500 ml of the aforementioned medium. One hour later, 550 µl 100 mM isopropyl β-(D)-thiogalactopyranoside (IPTG) were introduced and cultivated for 4 hours. The product was then recovered and introduced to centrifugation tube at each 250ml and centrifuged for 10 minutes at 7,000 rpm.

The supernatant was discarded and dissolved in 25 ml each 50 mM TrisHCl at a pH of 7.4 and Lysis buffer containing 25% sucrose.

Furthermore, 1.25 ml 10% Nonidet P-40(NP-40) and 125 µl 1 M $MgCl_2$ were added and transferred to a plastic tube. Sonication was performed 1 minute x 5 times while ice cold. The product was centrifuged for 15 minutes at 12,000 rpm and the supernatant was recovered.

Next, the aforementioned supernatant was adsorbed on a glutathione agarose column conditioned with phosphate-buffered saline (PBS).

Then the column was washed with twice the bed volume using a washing solution containing 20 mM Tris buffer at a pH of 7.4, 4.2 mM $MgCl_2$, and 1 mM

This Sheet Has Been Replaced

dithiothreithol (DTT). Elution was performed with 50 mM Tris buffer at a pH of 9.6 containing 5 mM glutathione. The protein content in the fraction was determined by the pigment bonding method (Bradford method; Biorad Co.) and the main fraction was acquired.

Purity of the purified ribosomal protein L7/L12/GST fused protein that was obtained was confirmed by electrophoresis to be approximately 75%, showing that a purity satisfactory for an immunogen could be guaranteed.

Example 3

Preparation of monoclonal antibody to ribosomal protein L7/L12 of *Haemophilus influenzae*

First, 100 μg fused protein antigen of ribosomal protein L7/L12/GST of *Haemophilus influenzae* were dissolved in 200 μl PBS and then 200 μl Freund's complete adjuvant were added and mixed and emulsification was performed. Two-hundred microliters were injected intraperitoneally to immunize mice.

Then the same emulsion antigen was intraperitoneally injected after 2 weeks, after 4 weeks, and after 6 weeks. Two-fold the concentration of antigen emulsion was injected intraperitoneally after 10 weeks and after 14 weeks. The spleen was excised 3 days after the final immunization and

This Sheet Has Been Replaced

submitted to cell fusion.

After thoroughly mixing 2 x 10^7 myeloma cells per 10^8 spleen cells from mice, which had been recovered aseptically, in a glass tube, the mixture was centrifuged for 5 minutes at 1,500 rpm and the supernatant was discarded. The cells were thoroughly mixed.

The myeloma cells used for cell fusion were obtained by cultivation of cell strain NS-1 with an RPMI 1640 culture medium containing 10% FCS, cultivating this product beginning 2 weeks before cell fusion using an RPMI 1640 medium containing 0.13 mM azaguanine, 0.5 µg/ml MC-210, and 10% FCS for 1 weeks, and then further cultivating the cell strain for 1 week with an RPMI 1640 medium containing 10% FCS.

Thirty milliliters of RPMI 1640 culture medium that had been kept at 37°C were added to the mixed cell sample and centrifuged at 1,500 rpm. After removal of the supernatant, 1 ml 50% polyethylene glycol that had been kept at 37°C was added and stirred for 1 minute. 10 ml RPMI 1640 medium kept at 37°C were added and the solution was vigorously mixed for approximately 5 minutes as it was suctioned and evacuated from a sterile pipette.

After centrifugation for 5 minutes at 1,000 rpm and removal of the supernatant, 30 ml HAT medium were

This Sheet Has Been Replaced



added to bring the cell concentration to 5 x 10^6/ml. This mixture was stirred till uniform and then poured, 0.1 ml at a time, into a 96-well culture plate and cultivated at 37°C under 7% CO_2. HAT medium was added, 0.1 ml at a time, on Day 1 and at Week 1 and Week 2.

Then the cells that had produced the desired antibody were screened by ELISA.

Solutions of GST fusion ribosomal protein L7/L12 and GST protein of *Haemophilus influenzae* dissolved in PBS containing 0.05% sodium azide diluted to 10 µg/ml were poured, each 100 µl at a time, into separate 96-well plates and adsorbed overnight at 4°C.

After removal of the supernatant, 200 µl of 1% bovine serum albumin solution (in PBS) were added and reacted and blocked for 1 hour at room temperature. After removal of the supernatant, the product was washed with washing solution (0.02% Tween 20, PBS). One-hundred microliters culture solution of fused cells were added to this and reacted for 2 hours at room temperature. The supernatant was removed and washed with washing solution. Next, 100 µl peroxidase-labeled goat anti-mouse IgG antibody solution at a concentration of 50 ng/ml were added and the solution was reacted for 1 hour at room temperature. The supernatant was removed and the

This Sheet Has Been Replaced

product was again washed with washing solution. Then TMB solution (KPL Co., ltd) was added, 100 µl at a time, and the mixture was reacted for 20 minutes at room temperature. After coloration, 100 µl 1 N sulfuric acid were added to stop the reaction and absorbance at 450 nm was determined.

As a result, positive wells that only reacted with GST fusion ribosomal protein L7/L12 and did not react with GST protein were detected, and it was concluded that antibody to ribosomal protein L7/L12 is present.

Therefore, the cells in the positive wells were recovered and cultivated with HAT medium in a 24-well plastic plate. The fused medium that had been cultivated was diluted with HT medium to a cell count of approximately 20 cells/ml and then mixed with 10^6 six-week-old mouse thyroid cells suspended in HT Cultivation medium in a 96-well culture plate. Culture was performed for 2 weeks at 37°C under conditions of 7% CO_2.

Antibody activity in the culture supernatant was similarly determined by the aforementioned ELISA method and the cells that showed positive reaction with ribosomal protein L7/L12 were recovered.

Furthermore, the same dilution test and cloning procedure was repeated to obtain a total of 5 clones of hybridoma HIRB-1 ~ 5.

This Sheet Has Been Replaced



Example 4

Reaction of monoclonal antibody that reacts with ribosomal protein L7/L12 protein of *Haemophilus influenzae* with Haemophilus influenzae and other bacteria

Monoclonal antibody was produced and recovered in accordance with standard methods using the positive hybridoma cells obtained as previously described.

Basically, 5 x 10^6 cells that had been subcultivated using RPMI 1640 culture medium (containing 10% FCS) were intraperitoneally injected into Balb/C mice that had been intraperitoneally injected with 0.5 ml Pristane 2 weeks earlier. Ascites was recovered 3 weeks later and the centrifugation supernatant was obtained.

The solution containing antibody that was obtained was adsorbed in a Protein A column (5 ml bed, Pharmacia) and rinsed with PBS at 3-times the bed volume. Then elution with citrate buffer at a pH of 3 was performed. The antibody fraction was recovered and the monoclonal antibody that produced by each hybridoma was obtained.

The monoclonal antibody derived from these 5 strains of hybridoma was used in ELISA.

The sandwich assay method was used to assess the monoclonal antibody. The monoclonal antibody that

This Sheet Has Been Replaced

was prepared was used as an enzyme-labeled antibody by being chemically bound to peroxidase.

That is, enzyme labeling was performed in accordance with the method in "Analytical Biochemistry" 132 (1983), 68-73 with the reagent S-acetylthioacetic acid N-hydroxysuccinimide for binding using horseradish peroxidase (Sigma Grade VI). By means of the ELISA reaction, a solution of commercial anti-*Haemophilus influenzae* polyclonal antibody which dissolved in PBS containing 0.05% sodium azide, (Biodesign, rabbit) was diluted to a concentration of 10 µg/ml and poured,100 µl at a time, into a separate 96-well plate and adsorbed overnight at 4°C.

After removal of the supernatant, 200 µl 1% FCS solution (in PBS) were added and reacted and blocked for 1 hour at room temperature. The supernatant was removed and the product was washed with washing solution (0.02% Tween 20, PBS). One-hundred microliters of antigen solution, which had been obtained by adding Triton X-100 to culture solutions of each species of microorganism to a concentration of 0.3% and then extracting the solution for 5 minutes at room temperature, were added to this and reacted for 2 hours at room temperature. The supernatant was removed and the product was further washed with washing solution. Then 100 µl

This Sheet Has Been Replaced

peroxidase-labeled anti-ribosomal protein L7/L12 antibody solution at 5 µg/ml were added and reacted for 1 hour at room temperature. The supernatant was removed and the product was washed with washing solution. TMB (KPL Co.) solution was added, 100 µl at a time, and reacted for 20 minutes at room temperature. After coloration, 100 µl 1 N sulfuric acid were added to stop the reaction. Absorbance at 450 nm was determined.

As a result, as shown in Table 1 it is clear that when monoclonal antibody derived from hybridoma HIRB-2 was used as the enzyme-labeled antibody, all strains of *Haemophilus influenzae* tested were detected at a sensitivity of 10^6 bacteria/ml, while reactivity of other bacteria belonging to the genus *Neisseria* and other microorganisms could not be detected, even at high concentrations of 10^8 bacteria/ml and therefore, antibody with specific reactivity to *Haemophilus influenzae* can be obtained by using monoclonal antibody to ribosomal protein L7/L12.

This Sheet Has Been Replaced

Table 1

		Results of Detection (10^6 cells/ml)
Haemophilus influenzae	ATCC9327	+
Haemophilus influenzae	ATCC9334	+
Haemophilus influenzae	ATCC9007	+
Haemophilus influenzae	ATCC9332	+
Haemophilus influenzae	ATCC8142	+
Haemophilus influenzae	ATCC9833	+
		Results of Detection (10^8 cells/ml)
Neisseria meningitides	ATCC13090	−
Neisseria lactamica	ATCC30011	−
Neisseria mucosa	ATCC35611	−
Neisseria sicca	ATCC9913	−
Branhamella catarrharis	ATCC25240	−
Neisseria gonorrhoeae	ATCC9793	−
Escherichia coli	ATCC25922	−
Klebsiella pneumoniae	ATCC13883	−

(+: Positive; −: Negative)

Example 5

Cloning of ribosomal protein L7/L12 genes from *Streptococcus pneumoniae*, mass expression in Escherichia coli and purification of the same protein and preparation of monoclonal antibody to the same protein

After inoculating an appropriate amount of *Streptococcus pneumoniae* strain IID555 (obtained from Tokyo University School of Medicine Laboratories) in a blood agar culture medium, the strain was cultivated for 48 hours in an incubator at 37°C. The colonies that grew were suspended in a TE buffer to a

This Sheet Has Been Replaced



final concentration of approximately 5 x 10^9 CFU/ml. Approximately 1.5 ml of this suspension was transferred to a microcentrifugation tube and centrifuged for 2 minutes at 10,000 rpm. The supernatant was discarded. The sediment was resuspended in 567 µl TE buffer. Then 30 µl 10% SDS and 3 µl 20 mg/ml Proteinase K solution were added and thoroughly mixed. The suspension was incubated for another hour at 37 °C. Next, after adding 80 µl 10% cetyl trimethyl ammonium bromide/0.7 M NaCl solution and thoroughly mixing the product, it was incubated for 10 minutes at 65°C. Next, 700 µl chloroform-isoamyl alcohol solution at a volume ratio of 24/1 was added and stirred well. The solution was centrifuged for 5 minutes (while being kept at 4°C) at 12,000 rpm using a microcentrifugation device and the aqueous fraction was transferred to a new microtube. Isopropanol was added to the fraction at 0.6-times its volume and the tube was vigorously shaken to form sediment of the DNA. The white DNA sediment was scooped with a glass rod and transferred to a different microcentrifugation tube containing 1 ml 70% ethanol (cooled to -20°C).

Next, the product was centrifuged for 5 minutes at 10,000 rpm and the supernatant was gently removed. Then another 1 ml 70% ethanol was added and the product was centrifuged for 5 more minutes. Once the

This Sheet Has Been Replaced



supernatant had been removed, the sediment was dissolved in 100 µl TE buffer to obtain the DNA solution. The concentration of the genomic DNA solution was determined quantitatively in accordance with E5: Sectrophotometric determination of the amount of DNA or RNA, "Molecular cloning: A laboratory manual," 1989, Eds. Sambrook, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press.

PCR was performed using 10 ng of this genomic DNA. Taq polymerase (Takara Co., ltd., code R001A) was employed for PCR. Then 5 µl of buffer attached to enzyme, 4 µl dNTP mixture attached to enzyme, and 200 pmol each of synthetic oligonucleotide C shown in Sequence No. 13 of the Sequence Table and synthetic oligonucleotide D shown in Sequence No. 14 of the Sequence table were added to the enzyme to bring the final volume to 50 µl.

This mixture was cycled 5 times with a TaKaRa PCR Thermal Cycler 480 for 1 minute at 95°C, 2 minutes at 50°C, and 3 minutes at 72°C and was then cycled 25 times for 1 minute at 95°C, 2 minutes at 60°C, and 3 minutes at 72°C. Electrophoresis was performed in 1.5% agarose gel using some of this PCR product. This product was then stained with ethidium bromide (Nihon Gene Co., ltd.) and observed under ultraviolet rays to confirm amplification of approximately 400 bp

This Sheet Has Been Replaced

cDNA. After digestion treatment with restriction endonucleases BamHI and XhoI, electrophoresis was performed in 1.5% agarose gel and staining with ethidium bromide was carried out. An approximately 370 bp band was cut out from the gel. This band was purified with Suprec01 (Takara Co., Ltd.) and then inserted into pGEX-6P-1 (Pharmacia), which is a commercial vector.

This same vector can function as an expression vector for the desired molecule, which can express fused protein with GST protein, by insertion of the desired gene fragment into the appropriate restriction endonuclease site. Actually, vector pGEX-6P-1 and the previous DNA were mixed together at a molar ratio of 1:5 and DNA was inserted into the vector with T4 DNA ligase (Invitrogen Co.). Vector pGEX-4T-1 into which DNA had been inserted was genetically introduced to *Escherichia coli* One-Shot Competent Cells (Invitrogen Co., Ltd.) and then inoculated in a plate of L-broth (Takara Co., Ltd.) semi-sold culture plate containing 50 µg/ml ampicillin (Sigma). The plate was then set aside at 37°C for 12 hours and the colonies that grew were selected at random and inoculated into 2 ml L-Broth liquid culture medium containing the same concentration of ampicillin. Shake cultivation was performed at 37°C for 8 hours and the bacteria was

This Sheet Has Been Replaced

recovered and the plasmid was separated using Wizard Miniprep in accordance with the attached literature. The plasmid was cleaved with restriction endonuclease BamHI/XhoI. Insertion of said PCR product was confirmed by cutting out approximately 370 bp DNA. The base sequence of the DNA that had been inserted was determined using said clone.

Determination of the base sequence of the inserted DNA fragment was performed using the Fluorescence Sequencer of Applied Biosystems. The sequence sample was prepared using PRISM, Ready Reaction Dye Terminator Cycle Sequencing Kit (Applied Biosystems). First, 9.5 µl reaction stock solution, 4.0 µl T7 promoter primer at 0.8 pmol/µl (Gibco BRL) and 6.5 µl of template DNA for sequencing at 0.16 µg/µl were added to a microtube with a capacity of 0.5 ml, mixed and superposed with 100 µl mineral oil. PCR amplification was performed for 25 cycles, where one cycle consisted of 30 seconds at 96°C, 15 seconds at 55°C, and 4 minutes at 60°C. The product was then kept at 4°C for 4 minutes. After the reaction was completed, 80 µl sterilized pure water was added and stirred. The product was centrifuged and the aqueous layer was extracted 3 times with phenol-chloroform. Ten microliters 3M sodium acetate (pH 5.2) and 300 µl ethanol were added to 100 µl aqueous layer and stirred. The product was then centrifuged for 15

This Sheet Has Been Replaced

minutes at room temperature and 14,000 rpm and the sediment was recovered. Once the sediment was washed with 75% ethanol, it was dried under a vacuum for 2 minutes to obtain the sequencing sample. The sequencing sample was dissolved in formamide containing 4 µl 10 mM EDTA and denatured for 2 minutes at 90°C. This was then cooled in ice and submitted to sequencing.

One of the 7 clones obtained had homology of the sequence with the probe used for PCR. In addition, DNA sequences extremely similar to the gene sequence of ribosomal protein L7/L12 gene of the other microorganisms, for example, *Neisseria gonorrhoeae*, were discovered. The entire base sequence and the corresponding amino acid sequence of the structural gene moiety are as shown in Sequence ID No. 19 and No. 20 of the Sequence Table. This gene fragment clearly codes for *Streptococcus pneumoniae* ribosomal protein L7/L12.

50 ml *Escherichia coli* into which expression vector had been inserted was cultivated overnight in a two-fold concentration YT medium at 37°C. Then, 450ml of the two-fold concentration YT medium was heated at 37°C for 1 hour. 50 ml of the *Escherichia coli* solution that had been cultivated overnight was introduced to 450ml of the aforementioned medium. After cultivation for one hour at 37°C, 100 µl 500m

This Sheet Has Been Replaced

mM IPTG was introduced and cultivated for 4 hours at 25°C. The product was then recovered and introduced to a 250 ml centrifugation tube and centrifuged for 20 minutes at 5000 rpm. The supernatant was discarded and dissolved in 25 ml each 50 mM Tris-HCl at a pH of 7.4 and Lysis buffer containing 25% sucrose.

Furthermore, 1.25 ml 10% NP-40 and 125 µl 1 M $MgCl_2$ were added and transferred to a plastic tube. Sonication was performed 1 minute x 5 times while ice cold. The product was centrifuged for 15 minutes at 12,000 rpm and the supernatant was recovered.

Next, the aforementioned supernatant was adsorbed on a glutathione sepharose column (manufactured by Pharmacia) conditioned with PBS. Then, the column was washed with PBS three times the bed volume. Elution was performed with 50 mM Tris-HCl at a pH of 8.0 containing 10 mM glutathione. The protein content in the fraction was determined by the pigment bonding method (Bradford method; Biorad Co.) and the main fraction was acquired. The main fraction was dialyzed three times against 3L PBS.

1 ml of a cleavage buffer containing 500 mM Tris-HCl (pH 7.0), 1.5 M NaCl, 10 mM EDTA, and 10 mM DTT was added to 10 ml of 1 mg/ml solution of the resulting GST fusion ribosomal protein L7/L12. 100 µl of 2 µg/µl PreScission Protease (manufactured by

This Sheet Has Been Replaced

Pharmacia company) was further added and reacted at 4°C to separate the GST moiety from ribosomal protein L7/L12.

The reaction solution was caused to pass through a glutathione sepharose column which had been conditioned with PBS. The solution coming out from the column was recovered. One-bed volume of PBS was caused to pass through and also recovered. Purity of the purified ribosomal protein L7/L12 that was obtained was confirmed by electrophoresis to be approximately 90%, showing that a purity satisfactory for an immunogen could be guaranteed.

First, 100 µg protein antigen of ribosomal protein L7/L12 of *Streptococcus pneumoniae* was dissolved in 200 µl PBS and then 200 µl Freund's complete adjuvant was added and mixed and emulsification was performed. 200 µl was intraperitoneally injected to immunize mice. Then, the same emulsion antigen was intraperitoneally injected after 2 weeks, after 4 weeks, and after 6 weeks. A two-fold concentration antigen emulsion was further injected intraperitoneally after 10 weeks and after 14 weeks. The spleen was excised 3 days after the final immunization and submitted to cell fusion.

After thoroughly mixing 2 x 10^7 myeloma cells per 10^8 spleen cells from mice, which had been recovered aseptically, in a glass tube, the mixture was

This Sheet Has Been Replaced

centrifuged for 5 minutes at 1500 rpm and the supernatant was discarded. The cells were thoroughly mixed.

The myeloma cells used for cell fusion were obtained by cultivation of cell strain NS-1 with an RPMI 1640 culture medium containing 10% FCS, cultivating this product beginning 2 weeks before cell fusion using an RPMI 1640 medium containing 0.13 mM azaguanine, 0.5 µg/ml MC-210, and 10% FCS for 1 weeks, and then further cultivating the cell strain for 1 week with an RPMI 1640 medium containing 10% FCS. 30 ml of 50 ml RPMI 640 culture medium that had been kept at 37°C was added to the mixed cell sample and centrifuged at 1,500 rpm. After removal of the supernatant, 1 ml 50% polyethylene glycol that had been kept at 37°C was added and stirred for 2 minute. 10 ml RPMI 1640 medium kept at 37°C was added and the solution was vigorously mixed for approximately 5 minutes as it was suctioned and evacuated using a sterile pipette.

After centrifugation for 5 minutes at 1,000 rpm and removal of the supernatant, 30 ml HAT medium were added to bring the cell concentration to 5 x 10^6 cells/ml. This mixture was stirred till uniform and then poured, 0,1ml at a time, into a 96-well culture plate and cultivated at 37°C under 7% CO_2. HAT medium was added, 0.1 ml at a time, on Day 1 and at Week 1

This Sheet Has Been Replaced

and Week 2.

Then, the cells that had produced the desired antibody were screened by ELISA. Solutions of ribosomal protein L7/L12 of *Streptococcus pneumoniae* dissolved in PBS containing 0.05% sodium azide diluted to 10 µg/ml were poured, 100 µl at a time, into separate 96-well plates and adsorbed overnight at 4°. After removal of the supernatant, 200 µl 1% FCS solution (in PBS) were added and reacted and blocked for 1 hour at room temperature. The supernatant was removed and the product was washed with a washing solution (0.02% Tween 20, PBS). 100 µl of a culture solution of fusion cells was added and reacted for two hours at room temperature. The supernatant was removed and the product was further washed with a washing solution. Then, 100 µl of a peroxidase-labeled goat anti-mouse antibody solution at 5 ng/ml was added and reacted for one hour at room temperature. The supernatant was removed and the product was washed with a washing solution. TMB (KPL) solution was added, 100 µl at a time, and reacted for 20 minutes at room temperature. After coloration, 100 µl 1 N sulfuric acid were added to stop the reaction. Absorbance at 450 nm was determined.

As a result, positive wells that reacted with ribosomal protein L7/L12 were detected, confirming

This Sheet Has Been Replaced



presence of the antibody to ribosomal protein L7/L12.

Therefore, the cells in the positive wells were recovered and cultivated with HAT medium in a 24-well plastic plate. The fused medium that had been cultivated was diluted with an HT medium to a cell count of approximately 20 cells/ml and then mixed with 10^6 six-week-old mouse thyroid cells suspended in the HT medium in a 96-well culture plate. The cells were cultivated for 2 weeks at 37°C under the conditions of 7% CO_2. The antibody activity in the culture supernatant was determined by the aforementioned ELISA method and the cells that showed a positive reaction with ribosomal protein L7/L12 were recovered.

Furthermore, the same dilution and cloning procedure was repeated to obtain a total of 4 clones of hybridoma AMSP-1 to 4.

Example 6

Reaction of monoclonal antibody that reacts with ribosomal protein L7/L12 of *Streptococcus pneumoniae* with *Streptococcus pneumoniae* and other microorganisms.

A monoclonal antibody was produced and recovered in accordance with standard methods using the positive hybridoma cells obtained as previously described.

Specifically, cells subcultivated in RPMI 1640

This Sheet Has Been Replaced

medium (containing 10% FCS) was diluted with a serum-free medium to about $2x10^5$cells/ml, $3.3x10^5$cells/ml, and $5x10^5$cells/ml in 25 cm^2 culture flasks, and the total volume was made 5 ml. After cells were grown for 3 to 5 days in 7% CO_2 at 37°C, a flask which contains the least number of original cells was selected among flasks in which cells were grown. The same procedure was repeated until the cells diluted to $2x10^5$cells/ml grow to $2x10^6$cells/ml in 3 to 4 days, thereby acclimatizing the cells with the serum-free medium. Next, cloning was performed in a 96-well plate for bacteria cultivation to select cells exhibiting fastest growth and a highest antibody titer. The selected cells were grown in a 24-well plate and diluted with a serum-free medium in a 25 cm^2 culture flask to a concentration of about $2x10^5$cells/ml and the total volume was made 10 ml. After incubation for 3 to 4 days in 7% CO_2 at 37°C to a concentration of $1x10^6$ cells/ml, the culture broth was transferred to a bottle for mass cultivation together with 100 ml of $1x10^6$ cells/ml cells which were grown in the same manner in a 75 cm^2 flask. 100 ml of a serum-free medium was added to the mixture, which was incubated at 37°C for two days while stirring. 200 ml of the serum-free medium was added again and the mixture was incubated for a further two days. The culture broth was divided into four

This Sheet Has Been Replaced

aliquot, the serum-free medium was added to each portion, followed by incubation for two days. After further addition of 400 ml of the serum-free medium, the culture broth was incubated for 6 days. The culture broth was collected and centrifuged at 10,000 rpm for 15 minutes to obtain a culture supernatant including the target antibody. After the addition of 0.1% sodium azide, the culture supernatant was stored at 4°C. 100 ml of the solution containing the antibody that was obtained was diluted to a 5-fold with PBS and adsorbed in a Protein G column (5 ml bed, Pharmacia) and rinsed with PBS at 3-times the bed volume. Then elution with citrate buffer at a pH of 3 was performed. The antibody fraction was recovered and monoclonal antibody that produced each hybridoma was obtained. The monoclonal antibodies originating from the four hybridoma strains were evaluated according to the OIA method described in Publication of the translation of International Patent Application Japanese Laid-open (Toku-Hyou) No. 509565/1995.

Specifically, the OIA method comprises preparing a reactive substrate by reacting an antibody for capture on a silicon wafer having a thin film layer of silicon nitride, causing this substrate to react with an antigen which is an extract of microorganisms for a prescribed period of time, causing the captured

This Sheet Has Been Replaced

antigen to react with an antibody (an amplification reagent) which is an enzyme-labeled antibody, and finally adding a substrate solution to produce a thin-film precipitate. The antigen-antibody reaction can be judged visually by a degree of light interference color produced in the precipitate.

The monoclonal antibody preparation was used and evaluated as a capture antibody to be immobilized on a silicon wafer having a silicon nitride thin film layer in the OIA method. Moreover, peroxidase-labeled AMGC-1 monoclonal antibody which can non-specifically react with ribosomal proteins L7/L12 protein of a variety of microorganisms described in Reference Example was used as the detect antibody. That is, enzyme labeling was performed in accordance with the method in "Analytical Biochemistry" 132 (1983), 68-73 with the reagent S-acetylthioacetic acid N-hydroxysuccinimide for binding using horseradish peroxidase (Sigma Grade VI).

In the OIA reaction, monoclonal antibody in a PBS containing 0.05% sodium azide was diluted with 0.1 M HEPES (pH 8.0) to a concentration of 10 µg/ml and added onto a silicone wafer which has a thin film layer of silicon nitride, 50 µl at a time, to react for 30 minutes at room temperature, followed by washing with distilled water and use.

15 µl of antigen solution, which had been


This Sheet Has Been Replaced

obtained by adding 0.5% Triton X-100 to culture solutions of various species of microorganisms and then extracting the solution for 5 minutes at room temperature, was added to the specimen obtained in the above-described procedure and reacted for 10 minutes at room temperature. Then, 15 µl of 20 µg/ml peroxidase-labeled AMGC1 anti-body was added and reacted for 10 minutes. After washing with distilled water, a substrate solution (KPL) was added, 15 µl at a time, and reacted for 5 minutes at room temperature. The product was washed with distilled water to judge the concentration of detection signals as an intensity of light interference by necked eyes.

As a result, as shown in Table 2 it is clear that when monoclonal antibody derived from hybridoma AMSP-2 was used as the capture antibody, all strains of *Streptococcus pneumoniae* tested were detected at a sensitivity of 10^6 bacteria/ml, while reactivity of other bacteria could not be detected at a higher concentration of 10^8 bacteria/ml. Thus, the antibody with specific reactivity to *Streptococcus pneumoniae* was confirmed to have been obtained by using the monoclonal antibody to ribosomal protein L7/L12.

The hybridoma AMSP-2 which produces the monoclonal antibody specific to *Streptococcus pneumoniae* has been deposited with National Institute of Bioscience and Human-Technology, the Agency of

This Sheet Has Been Replaced



Industrial Science and Technology, the Ministry of International Trade and Industry, Japan, on July 28, 1999, with the disposition number FERM BP-6870.

5

This Sheet Has Been Replaced




Table 2

	Results of Detection (10^6 cells/ml)	
Streptococcus pneumoniae	ATCC27336	+
Streptococcus pneumoniae	IID554	+
Streptococcus pneumoniae	IID555	+
Streptococcus pneumoniae	IID556	+
Streptococcus pneumoniae	IID557	+
Streptococcus pneumoniae	IID558	+
Streptococcus pneumoniae	IID559	+
Streptococcus pneumoniae	IID1603	+
	Results of Detection (10^8 cells/ml)	
Escherichia coli	ATCC25922	-
Enterococcus faecalis	ATCC19433	-
Haemophilus influenzae	ATCC10211	-
Klebsiella pneumoniae	ATCC13883	-
Neisseria gonorrhoeae	IID821	-
Neisseria lactamica	ATCC23970	-
Neisseria meningitidis	ATCC13090	-
Pseudomonas aeruginosa	ATCC27853	-
GroupB streptococcus	ATCC12386	-
Staphylococcus aureus	ATCC25923	-
Streptococcus pyogenes	ATCC19615	-

(+: Positive; -: Negative)

Example 7

Cloning of ribosomal protein L7/L12 genes from *Neisseria gonorrhoeae*, mass expression in Escherichia coli and purification of the same protein and preparation of monoclonal antibody to the same protein

After inoculating an appropriate amount of *Neisseria gonorrhoeae* strain IID821 (obtained from Tokyo University School of Medicine Laboratories) in a chocolate agar culture medium, the strain was cultivated for 24 hours in a CO_2 incubator under

This Sheet Has Been Replaced



conditions of 37°C and 0.5% CO_2. The colonies that grew were suspended in a TE buffer to a final concentration of approximately 5 x 10^9 CFU/ml. Approximately 1.5 ml of this suspension was transferred to a microcentrifugation tube and centrifuged for 2 minutes at 10,000 rpm. The supernatant was discarded. The sediment was resuspended in 567 µl TE buffer. Then 30 µl 10% SDS and 3 µl 20 mg/ml Proteinase K solution were added and thoroughly mixed. The suspension was incubated for another hour at 37°C.

Next, after adding 80 µl 10% cetyl trimethyl ammonium bromide/0.7 M NaCl solution and thoroughly mixing the product, it was incubated for 10 minutes at 65°C. Next, 700 µl chloroform-isoamyl alcohol solution at a volume ratio of 24/1 was added and stirred well. The solution was centrifuged for 5 minutes (while being kept at 4°C) at 12,000 rpm using a microcentrifugation device and the aqueous fraction was transferred to a new microtube. Isopropanol was added to the fraction at 0.6-times its volume and the tube was vigorously shaken to form sediment of the DNA. The white DNA sediment was scooped with a glass rod and transferred to a different microcentrifugation tube containing 1 ml 70% ethanol (cooled to -20°C).

Next, the product was centrifuged for 5 minutes

This Sheet Has Been Replaced

at 10,000 rpm and the supernatant was gently removed. Then another 1 ml 70% ethanol was added and the product was centrifuged for 5 more minutes.

Once the supernatant had been removed, the sediment was dissolved in 100 µl TE buffer to obtain the DNA solution. The concentration of the genomic DNA solution was determined quantitatively in accordance with E5: Sectrophotometric determination of the amount of DNA or RNA, "Molecular cloning: A laboratory manual," 1989, Eds. Sambrook, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press.

PCR was performed using 10 ng of this genomic DNA. PCR was performed using Taq polymerase (Takara Co., Ltd., code R001A). Then, 5 µl of a buffer attached to enzyme, 4 µl of a dNTP mixture attached to enzyme, and 200 pmol each of synthetic oligonucleotide E shown in Sequence No. 15 of the Sequence Table and synthetic oligonucleotide F shown in Sequence No. 16 of the Sequence Table, which were designed based on the ribosomal protein L7/L12 DNA sequence of *Neisseria gonorrhoeae* acquired from Internet Information (Oklahoma University, *N. Gonorrhoeae* Genome Project, disclosed genomic DNA data) because of the similarity with ribosomal protein L7/L12 DNA sequence of other bacteria, were added to the enzyme to bring the final volume to 50

This Sheet Has Been Replaced

µl.

This mixture was cycled 5 times with a TaKaRa PCR Thermal Cycler 480 for 1 minute at 95°C, 2 minutes at 50°C, and 3 minutes at 72°C and was then cycled 25 times for 1 minute at 95°C, 2 minutes at 60°C, and 3 minutes at 72°C. Electrophoresis was performed in 1.5% agarose gel using some of this PCR product. This product was then stained with ethidium bromide (Nihon Gene Co., ltd.) and observed under ultraviolet rays to confirm amplification of approximately 400 bp cDNA. After digestion treatment with restriction endonucleases BamHI and XhoI, electrophoresis was performed in 1.5% agarose gel and staining with ethidium bromide was carried out. An approximately 370 bp band was cut out from the gel. This band was purified with Suprec01 (Takara Co., Ltd.) and then inserted into pGEX-4T-1 (Pharmacia), which is a commercial vector. Actually, vector pGEX-4T-1 and the previous DNA were mixed together at a molar ratio of 1:3 and DNA was inserted into the vector with T4 DNA ligase (Invitrogen Co.). Vector pGEX-4T-1 into which DNA had been inserted was genetically introduced to *Escherichia coli* One-Shot Competent Cells (Invitrogen Co., Ltd.) and then inoculated in a plate of L-broth (Takara Co., Ltd.) semi-sold culture plate containing 50 µg/ml ampicillin (Sigma). The plate was then set aside at 37°C for 12 hours and the colonies that grew

This Sheet Has Been Replaced

were selected at random and inoculated into 2ml L-Broth liquid culture medium containing the same concentration of ampicillin. Shake cultivation was performed at 37°C for 8 hours and the bacteria was recovered and the plasmid was separated using Wizard Miniprep in accordance with the attached literature. The plasmid was cleaved with restriction endonuclease BamHI/XhoI. Insertion of said PCR product was confirmed by cutting out approximately 370 bp DNA. The base sequence of the DNA that had been inserted was determined using said clone.

Determination of the base sequence of the inserted DNA fragment was performed using the Fluorescence Sequencer of Applied Biosystems. The sequence sample was prepared using PRISM, Ready Reaction Dye Terminator Cycle Sequencing Kit (Applied Biosystems). First, 9.5 µl reaction stock solution, 4.0 µl T7 promoter primer at 0.8 pmol/µl (Gibco BRL) and 6.5 µl template DNA for sequencing at 0.16 µg/µl were added to a microtube with a capacity of 0.5 ml and mixed. After superposing with 100 µl mineral oil, PCR amplification was performed for 25 cycles, where one cycle consisted of 30 seconds at 96°C, 15 seconds at 55°C, and 4 minutes at 60°C. The product was then kept at 4°C for 4 minutes. After the reaction was completed, 80 µl sterilized pure water was added and stirred. The product was centrifuged

This Sheet Has Been Replaced



and the aqueous layer was extracted 3 times with phenol-chloroform. Ten microliters 3M sodium acetate (pH 5.2) and 300 µl ethanol were added to 100 µl aqueous layer and stirred. The product was then centrifuged for 15 minutes at room temperature and 1400 rpm and the sediment was recovered. Once the sediment was washed with 75% ethanol, it was dried under a vacuum for 2 minutes to obtain the sequencing sample. The sequencing sample was dissolved in formamide containing 4 µl 10 mM EDTA and denatured for 2 minutes at 90°C. This was then cooled in ice and submitted to sequencing. One of the 5 clones obtained had homology of the sequence with the probe used for PCR. In addition, DNA sequences extremely similar to the gene sequence of ribosomal protein L7/L12 gene of the other microorganisms, for example, *Haemophilus influenzae*, were discovered. The entire base sequence and the corresponding amino acid sequence of the structural gene moiety are as shown in Sequence No. 21 and No. 22 of the Sequence Table. This gene fragment clearly codes for ribosomal protein L7/L12 gene of *Neisseria gonorrhoeae*.

Neisseria gonorrhoeae fusion GST ribosome protein L7/L12 prepared by the same method as in Example 2 was obtained using the *Neisseria gonorrhoeae* fusion GST ribosomal protein L7/L12 expression vector constructed in this way.

This Sheet Has Been Replaced



Furthermore, hybridoma strain GCRB-3, which produces monoclonal antibody to ribosomal protein L7/L12 of *Neisseria gonorrhoeae*, was obtained in accordance with the method in Example 3.

Example 8

Reaction of monoclonal antibody that reacts with ribosomal protein L7/L12 of *Neisseria gonorrhoeae* with *Neisseria gonorrhoeae* and other microorganism

Monoclonal antibody was produced and recovered in accordance with standard methods using the positive hybridoma cells GCRB-3 obtained as previously described.

Basically, 5 x 10^6 cells (in PBS) that had been subcultivated using RPMI 1640 culture medium (containing 10% FCS) were intraperitoneally injected into Balb/C mice that had been intraperitoneally injected with 0,5ml Pristane 2 weeks earlier. Ascites was recovered 3 weeks later and the centrifugation supernatant was obtained.

The solution containing antibody that was obtained was adsorbed in a Protein A column (5 ml bed, Pharmacia) and rinsed with PBS at 3-times the bed volume. Then elution with citrate buffer at a pH of 3 was performed. The antibody fraction was recovered and the monoclonal antibody that was produced by each hybridoma was obtained. The monoclonal antibody derived from the GCRB-3 hybridoma

This Sheet Has Been Replaced

was used in ELISA.

The sandwich assay method was used to assess the monoclonal antibody. The monoclonal antibody that was prepared was used as an enzyme-labeled antibody by being chemically bound to peroxidase. That is, enzyme labeling was performed in accordance with the method in "Analytical Biochemistry" 132 (1983), 68-73 with the reagent S-acetylthioacetic acid N-hydroxysuccinimide for binding using horseradish peroxidase (Sigma Grade VI). By means of the ELISA reaction, a solution of commercial anti-*Neisseria gonorrhoeae* polyclonal antibody in PBS containing 0.05% sodium azide(Virostat, rabbit) was diluted to a concentration of 10 µg/ml and poured, 100 µl at a time, into a separate 96-well plate and adsorbed overnight at 4°C.

After removal of the supernatant, 200 µl 1% bovine serum albumin solution (in PBS) were added and reacted for 1 hour and blocked at room temperature. The supernatant was removed and the product was washed with washing solution (0.02% Tween 20, PBS). One-hundred microliters of antigen solution, which had been obtained by adding Triton X-100 to culture solutions of each species of microorganisms to a concentration of 0.3% and then extracting the solution for 5 minutes at room temperature, were added to this and reacted for 2 hours at room

This Sheet Has Been Replaced

temperature. The supernatant was removed and the product was further washed with washing solution. Then 100 µl peroxidase-labeled anti-ribosomal protein L7/L12 antibody solution at 5 µg/ml were added and reacted for 1 hour at room temperature. The supernatant was removed and the product was washed with washing solution. TMB (KPL) solution was added, 100 µl at a time, and reacted for 20 minutes at room temperature. After coloration, 100 µl 1 N sulfuric acid were added to stop the reaction. Absorbance at 450 nm was determined.

As a result, as shown in Table 3 it is clear that when monoclonal antibody derived from hybridoma GCRB-3 was used as the enzyme-labeled antibody, all strains of *Neisseria gonorrhoeae* tested were detected at a sensitivity of 10^6cells/ml, while reactivity of other species belonging to the genus *Neisseria* and other microorganisms could not be detected, even at high concentrations of 10^8cells/ml and therefore, antibody with specific reactivity to *Neisseria gonorrhoeae* can be obtained by using monoclonal antibody to ribosomal protein L7/L12.

This Sheet Has Been Replaced

Table 3

		Results of Detection (10^6 cells/ml)
Neisseria gonorrhoeae	ATCC9793	+
Neisseria gonorrhoeae	ATCC19424	+
Neisseria gonorrhoeae	ATCC27628	+
Neisseria gonorrhoeae	ATCC27629	+
Neisseria gonorrhoeae	ATCC27630	+
Neisseria gonorrhoeae	ATCC27631	+
Neisseria gonorrhoeae	ATCC27632	+
Neisseria gonorrhoeae	ATCC27633	+
Neisseria gonorrhoeae	ATCC35541	+
Neisseria gonorrhoeae	ATCC35542	+
Neisseria gonorrhoeae	ATCC43069	+
Neisseria gonorrhoeae	ATCC43070	+
Neisseria gonorrhoeae	ATCC49226	+
		Results of Detection (10^8 cells/ml)
Neisseria meningitidis	ATCC13090	−
Neisseria lactamica	ATCC30011	−
Neisseria mucosa	ATCC35611	−
Neisseria sicca	ATCC9913	−
Branhamella catarrharis	ATCC25240	−
Haemophilus influenzae	ATCC10211	−
Escherichia coli	ATCC25922	−
Klebsiella pneumoniae	ATCC13883	−

(+: Positive; −: Negative)

Example 9

Acquisition of monoclonal antibody to anti-ribosomal protein L7/L12 specific to genus *Neisseria*

After inoculating an appropriate amount of *Neisseria gonorrhoeae* strain IID821 (obtained from Tokyo University School of Medicine Laboratories) in

This Sheet Has Been Replaced

a chocolate agar culture medium, the strain was cultivated for 24 hours in a CO_2 incubator under conditions of 37°C and 0.5% CO_2. The colonies that grew were suspended in a TE buffer to a final concentration of approximately 5 x 10^9 CFU/ml. Approximately 1.5 ml of this suspension was transferred to a microcentrifugation tube and centrifuged for 2 minutes at 10,000 rpm. The supernatant was discarded. The sediment was resuspended in 567 µl TE buffer. Then 30 µl 10% SDS and 3 µl 20 mg/ml Proteinase K solution were added and thoroughly mixed. The suspension was incubated for another hour at 37°C. Next, after adding 80 µl 10% cetyl trimethyl ammonium bromide/0.7 M NaCl solution and thoroughly mixing the product, it was incubated for 10 minutes at 65°C. Next, 700 µl chloroform-isoamyl alcohol solution at a volume ratio of 24/1 was added and stirred well.

The solution was centrifuged for 5 minutes (while being kept at 4°C) at 12,000 rpm using a microcentrifugation device and the aqueous fraction was transferred to a new microtube. Isopropanol was added to the fraction at 0.6-times its volume and the tube was vigorously shaken to form sediment of the DNA. The white DNA sediment was scooped with a glass rod and transferred to a different microcentrifugation tube containing 1 ml 70% ethanol

This Sheet Has Been Replaced



(cooled to -20°C).

Next, the product was centrifuged for 5 minutes at 10,000 rpm and the supernatant was gently removed. Then another 1 ml 70% ethanol was added and the product was centrifuged for 5 more minutes. Once the supernatant had been removed, the sediment was dissolved in 100 µl TE buffer to obtain the DNA solution. The concentration of the genomic DNA solution was determined quantitatively in accordance with E5: Sectrophotometric determination of the amount of DNA or RNA, "Molecular cloning: A laboratory manual," 1989, Eds. Sambrook, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press.

PCR was performed using 10 ng of this genomic DNA. Taq polymerase (Takara Co., Ltd., code R001A) was employed for PCR. Then, 5 µl buffer attached to enzyme, 4 µl dNTP mixture attached to enzyme, and 200 pmol each of synthetic oligonucleotide E shown in Sequence No. 15 of the Sequence Table and synthetic oligonucleotide F shown in Sequence No. 16 of the Sequence Table, which were designed based on the ribosomal protein L12/L12 DNA sequence of *Neisseria gonorrhoeae* acquired from Internet information (Oklahoma University, *N. Gonorrhoeae* Genome Project, disclosed genome DNA data) because of the similarity with ribosomal protein L12/L12 DNA sequence of other

This Sheet Has Been Replaced

bacteria, were added to the enzyme to bring the final volume to 50 µl.

This mixture was cycled 5 times with a TaKaRa PCR Thermal Cycler 480 for 1 minute at 95°C, 2 minutes at 50°C, and 3 minutes at 72°C and was then cycled 25 times for 1 minute at 95°C, 2 minutes at 60°C, and 3 minutes at 72°C. Electrophoresis was performed in 1.5% agarose gel using some of this PCR product. This product was then stained with ethidium bromide (Nihon Gene Co., ltd.) and observed under ultraviolet rays to confirm amplification of approximately 400 bp cDNA. After digestion treatment with restriction endonucleases BamHI and XhoI, electrophoresis was performed in 1.5% agarose gel and staining with ethidium bromide was carried out. An approximately 370 bp band was cut out from the gel. This band was purified with Suprec01 (Takara Co., Ltd.) and then inserted into pGEX-6P-1 (Pharmacia), which is a commercial vector. This same vector can function as an expression vector for the desired molecule, which can express fused protein with GST protein, by insertion of the desired gene fragment into the appropriate restriction endonuclease site. Actually, vector pGEX-6P-1 and the previous DNA were mixed together at a molar ratio of 1:5 and DNA was inserted into the vector with T4 DNA ligase (Invitrogen Co.). Vector pGEX-6P-1 into which DNA had been inserted was

This Sheet Has Been Replaced

genetically introduced to *Escherichia coli* One-Shot Competent Cells (Invitrogen Co., Ltd.) and then inoculated in a plate of L-broth (Tomisaka Co., ltd.) semi-sold culture plate containing 50 µg/ml ampicillin (Sigma). The plate was then set aside at 37°C for 12 hours and the colonies that grew were selected at random and inoculated into 2ml L-Broth liquid culture medium containing the same concentration of ampicillin. Shake cultivation was performed at 37°C for 8 hours and the bacteria was recovered and the plasmid was separated using Wizard Miniprep in accordance with the attached literature. The plasmid was cleaved with restriction endonuclease BamHI/XhoI. Insertion of said PCR product was confirmed by cutting out approximately 370 bp DNA. The base sequence of the DNA that had been inserted was determined using said clone.

Determination of the base sequence of the inserted DNA fragment was performed using the Fluorescence Sequencer of Applied Biosystems. The sequence sample was prepared using PRISM, Ready Reaction Dye Terminator Cycle Sequencing Kit (Applied Biosystems). First, 9.5 µl reaction stock solution, 4.0 µl T7 promoter primer at 0.8 pmol/µl (Gibco BRL) and 6.5 µl template DNA for sequencing at 0.16 µg/µl were added to a microtube with a capacity of 0.5 ml and mixed. After superposing with 100 µl mineral oil,

This Sheet Has Been Replaced

PCR amplification was performed for 25 cycles, where one cycle consisted of 30 seconds at 96°C, 15 seconds at 55°C, and 4 minutes at 60°C. The product was then kept at 4°C for 4 minutes. After the reaction was completed, 80 µl sterilized pure water was added and stirred. The product was centrifuged and the aqueous layer was extracted 3 times with phenol-chloroform. Ten microliters 3M sodium acetate (pH 5.2) and 300 µl ethanol were added to 100 µl the aqueous layer and stirred. The product was then centrifuged for 15 minutes at room temperature and 1400 rpm and the sediment was recovered. Once the sediment was washed with 75% ethanol, it was dried under a vacuum for 2 minutes to obtain the sequencing sample. The sequencing sample was dissolved in formamide containing 4 µl 10 mM EDTA and denatured for 2 minutes at 90°C. This was then cooled in ice and submitted to sequencing.

One of the 4 clones obtained had homology of the sequence with the probe used for PCR. In addition, DNA sequences extremely similar to the gene sequence of ribosomal protein L7/L12 gene of the other microorganisms, for example, *Haemophilus influenzae*, were discovered. The entire base sequence and the corresponding amino acid sequence of the structural gene moiety are as shown in Sequence ID No. 21 and No. 22 of the Sequence Table. This gene fragment clearly

This Sheet Has Been Replaced

codes for *Neisseria gonorrhoeae* ribosomal protein L7/L12.

50 ml *Escherichia coli* into which expression vector had been inserted was cultivated overnight in a two-fold concentration YT medium at 37°C. Then, 450ml of the two-fold concentration YT medium was heated at 37°C for 1 hour. 50 ml of the *Escherichia coli* solution that had been cultivated overnight was introduced to 450ml of the aforementioned medium. After cultivation for one hour at 37°C, 100 µl 500 mM IPTG was introduced and cultivated for 4 hours at 25°C. The product was then recovered and introduced 250 ml to a centrifugation tube and centrifuged for 20 minutes at 5000 rpm. The supernatant was discarded and dissolved in 25 ml each 50 mM Tris-HCl at a pH of 7.4 and Lysis buffer containing 25% sucrose.

Furthermore, 1.25 ml 10% NP-40 and 125 µl 1 M $MgCl_2$ were added and transferred to a plastic tube. Sonication was performed 1 minute x 5 times while ice cold. The product was centrifuged for 15 minutes at 12,000 rpm and the supernatant was recovered.

Next, the aforementioned supernatant was adsorbed on a glutathione sepharose column (manufactured by Pharmacia) conditioned with PBS. Then, the column was washed with PBS three times the bed volume. Elution was performed with 50 mM Tris-HCl at a pH of 8.0 containing 10 mM glutathione. The protein content in

This Sheet Has Been Replaced

the fraction was determined by the pigment bonding method (Bradford method; Biorad Co.) and the main fraction was acquired. The main fraction was dialyzed three times against 3L PBS.

1 ml of a cleavage buffer containing 500 mM Tris-HCl (pH 7.0), 1.5 M NaCl, 10 mM EDTA, and 10 mM DTT was added to 10 ml of 1 mg/ml solution of the resulting GST fusion ribosomal protein L7/L12. 100 µl of 2 u/µl PreScission Protease (manufactured by Pharmacia company) was further added and reacted at 4°C to separate the GST moiety from ribosomal protein L7/L12.

The reaction solution was caused to pass through a glutathione sepharose column which had been conditioned with PBS. The solution coming out from the column was recovered. Onebed volume of PBS was caused to pass through and also recovered. Purity of the purified ribosomal protein L7/L12 that was obtained was confirmed by electrophoresis to be approximately 90%, showing that a purity satisfactory for an immunogen could be guaranteed.

First, 100 µg protein antigen of ribosomal protein L7/L12 of *Neisseria gonorrhoeae* was dissolved in 200 µl PBS and then 200 µl Freund's complete adjuvant was added and mixed and emulsification was performed. 200 µl was intraperitoneally injected to immunize mice. Then, the same emulsion antigen was

This Sheet Has Been Replaced

intraperitoneally injected after 2 weeks, after 4 weeks, and after 6 weeks. A two-fold concentration antigen emulsion was further injected intraperitoneally after 10 weeks and after 14 weeks. The spleen was excised 3 days after the final immunization and submitted to cell fusion.

After thoroughly mixing 2 x 10^7 myeloma cells per 10^8 spleen cells from mice, which had been recovered aseptically, in a glass tube, the mixture was centrifuged for 5 minutes at 1500 rpm and the supernatant was discarded. The cells were thoroughly mixed.

The myeloma cells used for cell fusion were obtained by cultivation of cell strain NS-1 with an RPMI 1640 culture medium containing 10% FCS, cultivating this product beginning 2 weeks before cell fusion using an RPMI 1640 medium containing 0.13 mM azaguanine, 0.5 µg/ml MC-210, and 10% FCS for 1 weeks, and then further cultivating the cell strain for 1 week with an RPMI 1640 medium containing 10% FCS. 30 ml of 50 ml RPMI 1640 culture medium that had been kept at 37°C was added to the mixed cell sample and centrifuged at 1,500 rpm. After removal of the supernatant, 1 ml 50% polyethylene glycol that had been kept at 37°C was added and stirred for 2 minute. 10 ml RPMI 1640 medium kept at 37°C was added and the solution was vigorously mixed for approximately 5

This Sheet Has Been Replaced

minutes as it was suctioned and evacuated from a sterile pipette.

After centrifugation for 5 minutes at 1,000 rpm and removal of the supernatant, 30 ml HAT culture medium were added to bring the cell concentration to 5 x 10^6cells/ml. This mixture was stirred till uniform and then poured, 0,1ml at a time, into a 96-well culture plate and cultivated at 37°C under 7% CO_2. HAT medium was added, 0.1 ml at a time, on Day 1 and at Week 1 and Week 2.

Then the cells that had produced the desired antibody were assessed by ELISA. Solutions of ribosomal protein L7/L12 of *Neisseria gonorrhoeae* dissolved in PBS containing 0.05% sodium azide were diluted to 10 µg/ml and were poured, 100 µl at a time, into separate 96-well plates and adsorbed overnight at 4°C. After removal of the supernatant, 200 µl 1% FCS solution (in PBS) were added and reacted and blocked for 1 hour at room temperature. The supernatant was removed and the product was washed with a washing solution (0.02% Tween 20, PBS). 100 µl of a culture solution of fusion cells was added and reacted for two hours at room temperature. The supernatant was removed and the product was further washed with a washing solution. Then, 100 µl of a peroxidase-labeled goat anti-mouse antibody solution at 5 ng/ml was added and reacted for one hour at room

This Sheet Has Been Replaced

temperature. The supernatant was removed and the product was washed with a washing solution. TMB (KPL) solution was added, 100 µl at a time, and reacted for 20 minutes at room temperature. After coloration, 100 µl 1 N sulfuric acid were added to stop the reaction. Absorbance at 450 nm was determined.

As a result, positive wells that reacted with ribosomal protein L7/L12 were detected, confirming presence of the antibody to ribosomal protein L7/L12.

Therefore, the cells in the positive wells were recovered and cultivated with HAT medium in a 24-well plastic plate. The fused medium that had been cultivated was diluted with an HT medium to a cell count of approximately 20 cells/ml. Then 50 μl the medium was mixed with 10^6 six-week-old mouse thyroid cells suspended in the HT medium in a 96-well culture plate. The cells were cultivated for 2 weeks at 37°C under the conditions of 7% CO_2. The antibody activity in the culture supernatant was determined by the aforementioned ELISA method and the cells that showed a positive reaction with ribosomal protein L7/L12 were recovered.

Furthermore, the same dilution and cloning procedure was repeated to obtain a total of 4 clones of hybridoma AMGC-5 to AMGC-8.

Monoclonal antibody was produced and recovered in

This Sheet Has Been Replaced

accordance with standard methods using the positive hybridoma cells obtained as previously described.

Specifically, cells subcultivated in RPMI 1640 medium (containing 10% FCS) was diluted with a serum-free medium to about $2x10^5$ cells/ml, $3.3x10^5$ cells/ml, and $5x10^5$ cells/ml in $25cm^2$ culture flask, and the total was made 5 ml. After cells were grown for 3 to 5 days in 7% CO_2 at 37°C, a flask which contains the least number of original cells was selected among flasks in which cells were grown. The same procedure was repeated until the cells diluted to $2x10^5$ cells/ml grow to $2x10^6$ cells/ml in 3 to 4 days, thereby acclimatizing the cells with the serum-free medium. Next, cloning was performed in a 96-well plate for bacteria cultivation to select cells exhibiting fastest growth and a highest antibody titer. The selected cells were grown in a 24-well plate and diluted with a serum-free medium in a 25 cm^2 culture flask to a concentration of about $2x10^5$ cells/ml and the total volume was made 10 ml. After incubation for 3 to 4 days in 7% CO_2 at 37°C to a concentration of $1x10^6$ cells/ml, the culture broth was transferred to a bottle for mass cultivation together with 100 ml of $1x10^6$ cells/ml cells which were grown in the same manner in a 75 cm^2 flask. 100 ml of a serum-free medium was added to the mixture, which was incubated at 37°C for two days while

This Sheet Has Been Replaced

stirring. 200 ml of the serum-free medium was added again and the mixture was incubated for a further two days. The culture broth was divided into four aliquot, the serum-free medium was added to each portion, followed by incubation for two days. After further addition of 400 ml of the serum-free medium, the culture broth was incubated for 6 days. The culture broth was collected and centrifuged at 10,000 rpm for 15 minutes to obtain a culture supernatant including the target antibody. After the addition of 0.1% sodium azide, the culture supernatant was stored at 4°C. 100 ml of the solution containing the antibody that was obtained was diluted to a 5-fold with PBS and adsorbed in a protein G column (5 ml bed, Pharmacia) and rinsed with PBS at 3-times the bed volume. Then elution with citrate buffer at a pH of 3 was performed. The antibody fraction was recovered and monoclonal antibody that produced each hybridoma was obtained.

The monoclonal antibodies originating from the four hybridoma strains were evaluated according to the OIA method described in International Patent Application Japanese Laid-open No. 509565/1995.

Specifically, the OIA method comprises preparing a reactive substrate by reacting an antibody for capture on a silicon wafer having a thin film layer of silicon nitride, causing this substrate to react

This Sheet Has Been Replaced



with an antigen which is an extract of microorganisms for a prescribed period of time, causing the captured antigen to react with an antibody (an amplification reagent) which is an enzyme-labeled antibody, and finally adding a substrate solution to produce a thin-film precipitate. The antigen-antibody reaction can be judged visually by a degree of light interference color produced in the precipitate.

The monoclonal antibody preparation was used as a capture antibody to be immobilized on a silicon wafer having a silicon nitride thin film layer in the OIA method. Moreover, peroxidase-labeled AMGC-1 monoclonal antibody which can non-specifically react with ribosomal proteins L7/L12 protein of a variety of microorganisms described in Reference Example was used as the detecte reagent antibody. That is, enzyme labeling was performed in accordance with the method in "Analytical Biochemistry" 132 (1983), 68-73 with the reagent S-acetylthioacetic acid N-hydroxysuccinimide for binding using horseradish peroxidase (Sigma Grade VI).

In the OIA reaction, monoclonal antibody in a PBS containing 0.05% sodium azide was diluted with 0.1 M HEPES (pH 8.0) to a concentration of 10 μg/ml and added onto a silicone wafer which has a thin film layer of silicon nitride, 50 μl at a time, to react for 30 minutes at room temperature, followed by

This Sheet Has Been Replaced

washing with distilled water and use. 15 µl of antigen solution, which had been obtained by adding 0.5% Triton X-100 to culture suspension of various species of microorganisms and then extracting the suspension for 5 minutes at room temperature, was added to the specimen obtained in the above-described procedure and reacted for 10 minutes at room temperature. Then, 15 µl of 20 µg/ml peroxidase-labeled AMGC1 was added and reacted for 10 minutes. After washing with distilled water, a substrate solution (KPL Co.) was added, 15 µl at a time, and reacted for 5 minutes at room temperature. The product was washed with distilled water to judge the concentration of detection signals as an intensity of light interference by necked eyes.

As a result, as shown in Table 4 it is clear that when monoclonal antibody derived from hybridoma AMGC-8 was used as the capture antibody, all strains of *Neisseria* genus tested were detected at a sensitivity of 10^8 cells/ml, while reactivity of other microorganisms could not be detected. Thus, the antibody with specific reactivity to *Neisseria* genus was confirmed to have been obtained by using the monoclonal antibody to ribosomal protein L7/L12.

This Sheet Has Been Replaced



Table 4

		Results of Detection (10^8 cells/ml)
Neisseria gonorrhoeae	IID821	+
Neisseria lactamica	ATCC23970	+
Neisseria meningitidis	ATCC13090	+
Escherichia coli	ATCC25922	−
Enterococcus faecalis	ATCC19433	−
Haemophilus influenzae	ATCC10211	−
Klebsiella pneumoniae	ATCC13883	−
Pseudomonas aeruginosa	ATCC27853	−
GroupB streptococcus	ATCC12386	−
Stauphylococcus aureus	ATCC25923	−
Streptococcus pneumoniae	ATCC27336	−
Streptococcus pyogenes	ATCC19615	−

(+: Positive; −: Negative)

Example 10

Acquisition of a polyclonal antibody which specifically reacts with ribosomal protein L7/L12 of *Haemophilus influenzae* using a ribosomal protein L7/L12protein-immobilized affinity column

A centrifugal supernatant of *Haemophilus influenzae* cell extract which had been treated with 0.5% Triton X-100 was used as an antigen. About 1.2 ml of a physiological saline solution containing 100 µg of antigen was emulsified with the addition of 1.5 ml of Freund's adjuvant. The emulsion was subcutaneously injected into four SPF Japanese White Rabbits to immunize the animals. The rabbits were immunized 5 to 6 times, once every two weeks, and the antibody titer was confirmed.

This Sheet Has Been Replaced

The antibody titer was confirmed by the ELISA method. Solutions of ribosomal protein L7/L12 of *Haemophilus influenzae* dissolved in PBS containing 0.05% sodium azide diluted to 10 µg/ml were poured, 100 µl at a time, into 96-well plates and adsorbed overnight at 4°C. After removal of the supernatant, 200 µl 1% FCS solution (in PBS) were added and reacted and blocked for 1 hour at room temperature. The supernatant was removed and the product was washed with a washing solution (0.02% Tween 20, PBS). 100 µl of a solution obtained by diluting normal rabbit serum and immunized rabbit antiserum was added and reacted for two hours at room temperature. The supernatant was removed and the product was further washed with a washing solution. Then, 100 µl of a peroxidase-labeled goat anti-rabbit IgG antibody solution at 50 ng/ml was added and reacted for one hour at room temperature. The supernatant was removed and the product was washed with a washing solution. OPD solution (Sigma Co.) was added, 100 µl at a time, and reacted for 20 minutes at room temperature. After coloration, 100 µl of 1 N sulfuric acid was added to stop the reaction. Absorbance at 492 nm was determined.

After confirming that the antibody titer had increased, a large quantity of blood was collected. Blood was collected in a centrifugal tube made of

This Sheet Has Been Replaced

glass from the ear artery, allowed to stand for one hour at 37°C, then overnight at 4°C. The mixture was centrifuged at 3000 rpm for 5 minutes and the supernatant was recovered. The resulting anti-serum was stored at 4°C.

Next, an affinity column with immobilized ribosomal protein L7/L12 of *Haemophilus influenzae* and *Neisseria gonorrhoeae* was prepared. HiTrap NHS-activated column (1 ml, manufactured by Pharmacia) was used. Immediately after replacing the column with 6 ml of 1 mM HCl, 1 ml of a PBS solution of ribosomal protein L7/L12 adjusted to 1 mg/ml was charged. The column was allowed to stand for 15 minutes at 25°C. This procedure was repeated 5 times, thereby feeding the total 5 ml of the PBS solution of ribosomal protein L7/L12. Then, 6 ml of Buffer A (0.5M ethanolamine, 0.5 M NaCl, pH 8.3), 6 ml of Buffer B (0.1 M acetic acid, 0.5M NaCl, pH 4), and 6 ml of Buffer A were charged as blocking reagents. After allowing to stand for 15 minutes at 25°C, 6 ml of Buffer B, 6 ml of Buffer A, and 6 ml of Buffer B were further added. The mixture was then equilibrated with 6 ml of PBS.

Using the affinity column with immobilized ribosomal protein L7/L12 of *Haemophilus influenzae*, the polyclonal antibody in the resulting anti-serum was purified using the supernatant of Triton X-100

This Sheet Has Been Replaced

treated bacteria of *Haemophilus influenzae* as an antigen. This antiserum was first diluted with PBS to a volume of 5 times, caused to pass through a 0.45 μm filter, then caused to be adsorbed in the column immobilized with ribosomal protein L7/L12 of *Haemophilus influenzae* at a flow rate of 0.5 ml/min. After elution from the column with 0.1 M glycine (pH 2.1) and immediately neutralizing with 1 M Tris-HCl (pH 9.0), eluted fractions of the target antibody were recovered by the ELISA method, the same as anti-titer measuring method. These fractions were caused to pass through the affinity column immobilized with the ribosomal protein L7/L12 of *Neisseria gonorrhoeae*, which was equilibrated with PBS whereby the antibody that reacts with the ribosomal protein L7/L12 of *Neisseria gonorrhoeae* was adsorbed and fraction which passed through without adsorption was recovered.

The polyclonal antibody purified in this manner was evaluated by the same OIA method as in Example 6.

The purified antibody was used as a capture antibody for the OIA method. Moreover, peroxidase-labeled AMGC-1 monoclonal antibody described in Reference Example was used as the detect antibody. That is, enzyme labeling was performed in accordance with the method in "Analytical Biochemistry" 132 (1983), 68-73 with the reagent S-acetylthioacetic

This Sheet Has Been Replaced

acid N-hydroxysuccinimide for binding using horseradish peroxidase (Sigma Grade VI).

In the OIA reaction, the purified polyclonal antibody in a PBS containing 0.05% sodium azide was diluted with 0.1 M HEPES (pH 8.0) to a concentration of 10 μg/ml and added onto a silicone wafer, 50 μl at a time, to react for 30 minutes at room temperature, followed by washing with distilled water and use.

15 μl of antigen solution, which had been obtained by adding 0.5% Triton X-100 to culture suspension of various species of microorganisms and then extracting the suspension for 5 minutes at room temperature, was added to the specimen obtained in the above-described procedure and reacted for 10 minutes at room temperature. Then, 15 μl of 20 μg/ml peroxidase-labeled AMGC1 was added and reacted for 10 minutes. After washing with distilled water, a substrate solution (KPL) was added, 15 μl at a time, and reacted for 5 minutes at room temperature. The product was washed with distilled water to observe the blue color concentration by the necked eyes.

As a result, as shown in Table 5 it is clear that when the purified polyclonal antibody of APHI2-2 was used as the capture antibody, *Haemophilus influenzae* tested at a sensitivity of 10^8 bacteria/ml was detected, while reactivity of other microorganisms could not be detected. Thus, the antibody with

This Sheet Has Been Replaced

specific reactivity to *Haemophilus influenzae* was confirmed to have been obtained by using the polyclonal antibody purified by an affinity column immobilized with the ribosomal protein L7/L12 of *Haemophilus influenzae*.

This Sheet Has Been Replaced



Table 5

		Results of Detection (10^8 cells/ml)
Haemophilus influenzae	ATCC10211	+
Escherichia coli	ATCC25922	−
Enterococcus faecalis	ATCC19433	−
Klebsiella pneumoniae	ATCC13883	−
Neisseria gonorrhoeae	IID821	−
Neisseria lactamica	ATCC23970	−
Neisseria meningitidis	ATCC13090	−
Pseudomonas aeruginosa	ATCC27853	−
GroupB Streptococcus	ATCC12386	−
Stauphylococcus aureus	ATCC25923	−
Streptococcus pneumoniae	ATCC27336	−
Streptococcus pyogenes	ATCC19615	−

(+: Positive; −: Negative)

Reference Example 1]

Acquisition of monoclonal antibody that reacts non-specifically with ribosomal protein L7/L12 of various bacteria

So called sandwich assay in which an antigen is sandwiched between a capture antibody and a labeled antibody for detection is useful to detect microorganisms by optical immunoassay and the ELISA method because of its high detection sensitivity. In this instance, not only an antibody which specifically reacts with antigens originating from the subject microorganism, but also another antibody which recognizes an antigen epitope are required.

This Sheet Has Been Replaced

Antibodies which react non-specifically with ribosomal protein L7/L12 originating various microorganisms are very useful as an antibody which can constitute a sandwich assay with an antibody which reacts with specifically ribosomal protein L7/L12.

Fortunately, ribosomal proteins L7/L12 protein of a variety of microorganisms have a region wherein the amino acid sequence is homologous. Here, the inventors have been successful in acquiring a monoclonal antibody which exhibits a cross reaction with ribosomal proteins L7/L12 protein of various species of microorganisms from *Neisseria gonorrhoeae*. It has been discovered that an antibody to anti-ribosomal protein L7/L12 having no specificity, which was acquired from one species of microorganisms can be used for sandwich assay of all microorganisms.

Cloning of ribosomal protein L7/L12 genes from *Neisseria gonorrhoeae*, mass expression in Escherichia coli and purification of the same protein, and preparation of monoclonal antibody to the same protein were performed.

After inoculating an appropriate amount of *Neisseria gonorrhoeae* strain IID821 (obtained from Tokyo University School of Medicine Laboratories) in a chocolate agar culture medium, the strain was cultivated for 24 hours in a CO_2 incubator under

This Sheet Has Been Replaced

conditions of 37°C and 0.5% CO_2. The colonies that grew were suspended in a TE buffer to a final concentration of approximately 5 x 10^9 CFU/ml. Approximately 1.5 ml of this suspension was transferred to a microcentrifugation tube and centrifuged for 2 minutes at 10,000 rpm. The supernatant was discarded. The sediment was resuspended in 567 µl TE buffer. Then 30 µl 10% SDS and 3 µl 20 mg/ml Proteinase K solution were added and thoroughly mixed. The suspension was incubated for another hour at 37°C. Next, after adding 80 µl 10% cetyl trimethyl ammonium bromide/0.7 M NaCl solution and thoroughly mixing the product, it was incubated for 10 minutes at 65°C. Next, 700 µl chloroform-isoamyl alcohol solution at a volume ratio of 24/1 was added and stirred well. The solution was centrifuged for 5 minutes (while being kept at 4°C) at 12,000 rpm using a microcentrifugation device and the aqueous fraction was transferred to a new microtube. Isopropanol was added to the fraction at 0.6-times its volume and the tube was vigorously shaken to form sediment of the DNA. The white DNA sediment was scooped with a glass rod and transferred to a different microcentrifugation tube containing 1 ml 70% ethanol (cooled to -20°C).

Next, the product was centrifuged for 5 minutes at 10,000 rpm and the supernatant was gently removed.

This Sheet Has Been Replaced

Then another 1 ml 70% ethanol was added and the product was centrifuged for 5 more minutes. Once the supernatant had been removed, the sediment was dissolved in 100 µl TE buffer to obtain the DNA solution. The concentration of the genomic DNA solution was determined quantitatively in accordance with E5: Sectrophotometric determination of the amount of DNA or RNA, "Molecular cloning: A laboratory manual," 1989, Eds. Sambrook, J., Fritsch, E.F., and Maniatis, T., Cold Spring Harbor Laboratory Press.

PCR was performed using 10 ng of this genomic DNA. Taq polymerase (Takara Co., Ltd., code R001A) was employed for PCR. Then, 5 µl of a buffer attached to enzyme, 4 µl of dNTP mixture attached to enzyme, and 200 pmol of each of synthetic oligonucleotide E shown in Sequence No. 15 of the Sequence Table and synthetic oligonucleotide F shown in Sequence No. 16 of the Sequence Table, which were designed based on the ribosomal protein L7/L12 DNA sequence of *Neisseria gonorrhoeae* acquired from Internet information (Oklahoma University, *N. Gonorrhoeae* Genome Project, disclosed genomic DNA data) because of the similarity with ribosomal protein L7/L12 DNA sequence of other bacteria, were added to the enzyme to bring the final volume to 50 µl.

This Sheet Has Been Replaced

This mixture was cycled 5 times with a TaKaRa PCR Thermal Cycler 480 for 1 minute at 95°C, 2 minutes at 50°C, and 3 minutes at 72°C and was then cycled 25 times for 1 minute at 95°C, 2 minutes at 60°C, and 3 minutes at 72°C. Electrophoresis was performed in 1.5% agarose gel using some of this PCR product. This product was then stained with ethidium bromide (Nihon Gene Co., ltd.) and observed under ultraviolet rays to confirm amplification of approximately 400 bp cDNA. After digestion treatment with restriction endonucleases BamHI and XhoI, electrophoresis was performed in 1.5% agarose gel and staining with ethidium bromide was carried out. An approximately 370 bp band was cut out from the gel. This band was purified with Suprec01 (Takara Co., Ltd.) and then inserted into pGEX-4T-1 (Pharmacia), which is a commercial vector. Actually, vector pGEX-4T-1 and the previous DNA were mixed together at a molar ratio of 1:3 and DNA was inserted into the vector with T4 DNA ligase (Invitrogen Co.). Vector pGEX-4T-1 into which DNA had been inserted was genetically introduced to *Escherichia coli* One-Shot Competent Cells (Invitrogen Co., Ltd.) and then inoculated in a plate of L-broth (Takara Co., Ltd.) semi-sold culture plate containing 50 µg/ml ampicillin (Sigma). The plate was then set aside at 37°C for 12 hours and the colonies that grew were selected at random and

This Sheet Has Been Replaced

inoculated into 2ml L-Broth liquid culture medium containing the same concentration of ampicillin. Shake cultivation was performed at 37°C for 8 hours and the bacteria was recovered and the plasmid was separated using Wizard Miniprep in accordance with the attached literature. The plasmid was cleaved with restriction endonuclease BamHI/XhoI. Insertion of said PCR product was confirmed by cutting out approximately 370 bp DNA. The base sequence of the DNA that had been inserted was determined using said clone.

Determination of the base sequence of the inserted DNA fragment was performed using the Fluorescence Sequencer of Applied Biosystems. The sequence sample was prepared using PRISM, Ready Reaction Dye Terminator Cycle Sequencing Kit (Applied Biosystems). First, 9.5 µl reaction stock solution, 4.0 µl T7 promoter primer at 0.8 pmol/µl (Gibco BRL) and 6.5 µl template DNA for sequencing at 0.16 µg/µl were added to a microtube with a capacity of 0.5 ml and mixed. After superposition with 100 µl mineral oil, PCR amplification was performed for 25 cycles, where one cycle consisted of 30 seconds at 96°C, 15 seconds at 55°C, and 4 minutes at 60°C. The product was then kept at 4°C for 4 minutes. After the reaction was completed, 80 µl sterilized pure water was added and stirred. The product was centrifuged

This Sheet Has Been Replaced

and the aqueous layer was extracted 3 times with phenol-chloroform. Ten microliters 3M sodium acetate (pH 5.2) and 300 µl ethanol were added to 100 µl aqueous layer and stirred. The product was then centrifuged for 15 minutes at room temperature and 14,000 rpm and the sediment was recovered. Once the sediment was washed with 75% ethanol, it was dried under a vacuum for 2 minutes to obtain the sequencing sample. The sequencing sample was dissolved in formamide containing 4 µl 10 mM EDTA and denatured for 2 minutes at 90°C. This was then cooled in ice and submitted to sequencing. One of the 5 clones obtained had homology of the sequence with the probe used for PCR. In addition, DNA sequences extremely similar to the gene sequence of ribosomal protein L7/L12 gene of the other microorganisms, for example, *Haemophilus influenzae*, were discovered. The entire base sequence and the corresponding amino acid sequence of the structural gene moiety are as shown in Sequence No. 21 and No. 22 of the Sequence Table. This gene fragment clearly codes for *Neisseria gonorrhoeae* ribosomal protein L7/L12.

Neisseria gonorrhoeae GST fused ribosome protein L7/L12 prepared by the same method as in Example 2 was obtained using the Neisseria gonorrhoeae GST fusion ribosomal protein L7/L12 expression vector constructed in this way. Furthermore, hybridoma

This Sheet Has Been Replaced



strain AMGC1, which produces monoclonal antibody to ribosomal protein L7/L12 of *Neisseria gonorrhoeae*, was obtained in accordance with the method similar to the method of Example 3. Monoclonal antibody was produced and recovered in accordance with standard methods using the positive hybridoma cells of AMGC1 strain obtained as previously described.

Specifically, cells subcultivated in RPMI 1640 medium (containing 10% FCS) was diluted with a serum-free medium to about $2x10^5$cells/ml, $3.3x10^5$cells/ml, and $5x10^5$ cells/ml in 25 cm^2 culture flasks, and the total was made 5 ml. After cell were grown for 3 to 5 days in 7% CO_2 at 37°C, a flask which contains the least number of original cells was selected among flasks in which cells were grown. The same procedure was repeated until the cells diluted to $2x10^5$ cells/ml grow to $2x10^6$/ml in 3 to 4 days, thereby acclimatizing the cells with the serum-free medium. Next, cloning was performed in a 96-well plate for bacteria cultivation to select cells exhibiting fastest growth and a highest antibody titer. The selected cells were grown in a 24-well plate and diluted with a serum-free medium in a 25 cm^2 culture flask to a concentration of about $2x10^5$ cells/ml and the total volume was made 10 ml. After incubation for 3 to 4 days in 7% CO_2 at 37°C to a concentration of $1x10^6$ cells/ml, the culture broth was transferred

This Sheet Has Been Replaced

to a bottle for mass cultivation together with 100 ml of $1x10^6$ cells/ml cells which were grown in the same manner in a 75 cm^2 flask. 100 ml of a serum-free medium was added to the mixture, which was incubated at 37°C for two days while stirring. 200 ml of the serum-free medium was added again and the mixture was incubated for a further two days. The culture broth was divided into four aliquot, the serum-free medium was added to each portion, followed by incubation for two days. After further addition of 400 ml of the serum-free medium, the culture broth was incubated for 6 days. The culture broth was collected and centrifuged at 10,000 rpm for 15 minutes to obtain a culture supernatant including the target antibody. After the addition of 0.1% sodium azide, the culture supernatant was stored at 4°C. 100 ml of the solution containing the antibody that was obtained was diluted to a 5-fold with PBS and adsorbed in a Protein A column (5 ml bed, Pharmacia) and rinsed with PBS at 3-times the bed volume. Then elution with citrate buffer at a pH of 3 was performed. The antibody fraction was recovered and monoclonal antibody that produced each hybridoma was obtained. The monoclonal antibody derived from the hybridoma was used in ELISA.

To evaluate the antibody, 96 well plates sensitized with ribosomal protein L7/L12s of various

This Sheet Has Been Replaced

microorganisms were used as an antigen. The monoclonal antibody prepared was reacted, followed by the reaction of horseradish peroxidase-labeled anti-mouse IgG (manufactured by MBL, Code 330) as a secondary antigen, and finally the antibody was detected using an enzyme reaction coloring reagent. In the ELISA reaction, solutions of recombinant ribosomal protein L7/L12 of *Neisseria gonorrhoeae*, *Haemophilus influenzae*, and *Streptococcus pneumoniae* dissolved in PBS containing 0.05% sodium azide diluted to 1 μg/ml were poured, 100 μl at a time, into separate 96-well plates and adsorbed overnight at 4°C. After removal of the supernatant, 200 μl 1% bovine serum albumin solution (in PBS) were added and reacted and blocked for 1 hour at room temperature. The supernatant was removed and the product was washed with a washing solution (0.02% Tween 20, PBS). Solutions of AMGC1 antibody, at concentrations of 0.1-1 μg/ml, in an amount of 100 μl each, were added and reacted for two hours at room temperature. The supernatant was removed and the product was further washed with a washing solution. Then, 100 μl of a solution of 5 μg/ml horse radish peroxidase-labeled anti-mouse IgG (manufactured by MBL, Code 330) was added and reacted for one hour at room temperature. The supernatant was removed and the product was washed with a washing solution. TMB (KPL) solution

This Sheet Has Been Replaced

was added, 100 µl at a time, and reacted for 20 minutes at room temperature. After coloration, 100 µl 1 N sulfuric acid were added to stop the reaction. Absorbance at 450 nm was determined.

As a result, as shown in Table 6 it was confirmed that when the monoclonal antibody originating from hybridoma AMGC1 was used, this antibody can react with ribosomal protein L7/L12s of all bacteria such as *Neisseria gonorrhoeae*, *Haemophilus influenzae*, and *Streptococcus pneumoniae*.

Table 6

Results of detection of AMGC1 antibody and Ribosomal Protein L7/L12 of microorganisms	
Neisseria gonorrhoeae	+
Haemophilus influenzae	+
Streptococcus pneumoniae	+

(+: Positive)

The AMGC1 antibody obtained here is very useful as an antibody used for the detection of microorganisms in the so-called sandwich assay by optical immunoassay and ELISA method, in combination with an anti-ribosomal protein L7/L12 antibody which is specific to each microorganism.

Industrial Applicability

According to the present invention, not only microorganisms can be detected specifically according

This Sheet Has Been Replaced

to species, but also microorganisms of all serotypes in the same species can be detected at a high precision, by using antibodies to intracellular molecules of the same function.

By using antibodies to ribosomal proteins L7/L12 of various microorganisms as such antibodies, *Haemophilus influenzae* and *Neisseria gonorrhoeae* can be detected precisely.

Moreover, specific antibodies used for the detection of various kinds of microorganisms can be prepared by using intracellular molecules exhibiting same functions in various microorganisms as an antigen.

This Sheet Has Been Replaced



REMARKS TO DEPOSITED MICROORGANISMS

The organization in which the microorganisms have been deposited:

National Institute of Bioscience and Human Technology, the Agency of Industrial Science and Technology

Address: 1-1-3, Higashi, Yatabe-cho, Tsukuba-shi, Ibaraki-ken, Japan (Postal Code: 305).

Date of deposition: July 28, 1999

Number of deposition given by the deposition organization: FERM PB6807

This Sheet Has Been Replaced

This Sheet Has Been Replaced

CLAIMS:

1. Antibodies which are antibodies to ribosomal protein of microorganisms and which react specifically with said microorganisms.

2. The antibodies according to claim 1, where the ribosomal protein of the microorganisms is ribosomal protein L7/L12.

3. The antibodies according to claim 1 or 2, where said microorganisms are microorganisms which cause a sexually transmitted disease (STD).

4. The antibodies according to claim 1 or 2, where said microorganisms are microorganisms which cause respiratory tract infection.

5. The antibody according to claim 4, where the causative microorganisms of respiratory tract infection are microorganisms of *Haemophilus influenzae*.

6. The antibody according to claim 4, where the causative microorganisms of respiratory tract infection are microorganisms of *Streptococcus pneumoniae*.

7. The antibody according to claim 3, where the causative microorganisms of sexually transmitted diseases (STD) are microorganisms of *Neisseria gonorrhoeae*.

8. The antibody according to claim 3, which is the antibody to ribosomal protein L7/L12 of *Neisseria*

* C L M S *

This Sheet Has Been Replaced

gonorrhoeae and which identifies a continuous amino acid sequence moiety from 5 to 30 amino acids including the 115th alanine in the amino acid sequence of Sequence ID No. 22 of the Sequence Table.

9. A method of detecting microorganisms, which is characterized by the fact that antibody to intracellular molecules that have the same function for a variety of microorganisms is used.

10. A method of detecting microorganisms, which is characterized by the fact that any antibody according to claims 1 to 8 is used.

11. A reagent kit for detecting microorganisms, which is characterized by the fact that antibody to intracellular molecules that have the same function for a variety of microorganisms is used.

12. A reagent kit for detecting microorganisms, which is characterized by the fact that any antibody according to claims 1 to 8 is used.

13. A method of preparing any antibody described any one of claims 1 to 8, characterized by the fact that ribosomal protein L7/L12 of microorganisms obtained by a gene manipulation procedure or by isolation from microorganisms, peptide moiety thereof, or a synthesized peptide corresponding to the peptide moiety is used as an immune source.

* C L M S *