

-1-

A METHOD FOR PRESERVATION OF HUMAN HEMATOPOIETIC STEM OR PROGENITOR CELLS.

Field of invention

5 This invention relates to a method for preservation of Human
hematopoietic stem or progenitor cells.

More particularly, it relates to a method for preservation of Human
hematopoietic stem or progenitor cells with the effect of mannose
binding lectins on human hematopoietic stem/progenitor cells.

Background of the invention

10 Preservation of hematopoietic stem/progenitor cells (HSPC) is
conventionally carried out using low temperature freezing either at -
80°C (mechanical freezers) or at -196°C (liquid nitrogen) and it is
altogether a separate science in itself. Present invention relates to
maintenance of HSC in culture. Maintenance HSPC in vitro without
15 proliferation or differentiation has important clinical applications.
Techniques such as in vitro purging of tumor cells using culture
techniques or by use of chemotherapeutic agents necessitates that the
HSPC remain unaffected in the culture. In genetic engineering
protocols it could be advantageous to eliminate all the short-lived

-2-

progenitors, which are growth factor dependent and then use the surviving primitive cells for transduction purposes. Thus a search for agents which can maintain the HSPC in culture and identification of signals involved in maintenance of the quiescent state is likely to form a fruitful endeavor. Earlier efforts in this direction to maintain HSPC in culture have been made using either specific matrix molecules like denatured collagen (Manney et al, 2004 Biomaterials, 25, 16: 3233-43; Kim et al 2003, Int J Hematol; 78, 2:126-32; Banu et al 2001, Cytokine. 21; 13, 6:349-58) or growth factors like FLT3L (Shapiro et al 1996, Hematother; 5, 6:655-62) or hematopoietic growth factors like Epo (Eridani et al 1998, Biotherapy 10, 4:295-8). These approaches, however, involve use of expensive reagents. The cost factor becomes a major issue when it is to be applied to clinical set ups especially in developing countries. Recently, a lectin named Fli3 receptor-interacting lectin (FRIL) was shown to preserve the quiescent HSPC derived from cord blood (CB) in culture for 30 days without the change of medium and without the exogenous addition of growth factors. (Colucci, G. et al 1999 Proc. Natl. Acad. Sci. USA 96, 646-650; Kollet O. et al 2000 Expt. Hematol. 28, 726-736.) The plant lectin FRIL was shown to support prolonged in vitro maintenance of quiescent human cord blood CD 34 (+) CD38 (-/low)/SCID repopulating stem cells. It was reported that this maintenance of

-3-

quiescent CB progenitors may be through the cell cycle modulation through HTm4 and HTm4S (Xie, X. Y., Xie C., Shi, W., Li, J., Li, Y. H., Wang, D.M., Bai, C.X., Chen, L. & Pei, X.T. 2004 56, 306-312).

Since FRIL is a mannose binding lectin, screening for more mannose
5 binding lectins have been done for their possible HSPC preservation activity so that they can also be used as tools to identify the signals involved in maintenance of quiescent HSPC. Thus the present invention has two distinct advantages over earlier approaches, one of being cost effective and the second of providing useful and specific
10 reagents to study signaling mechanisms.

Maintenance of HSPC in culture has a great clinical utility especially if it can be done without change of medium or use of expensive reagents. The present invention provides a method to achieve this. The mannose specific lectins that have been used in the present study
15 showed HSPC preservation activity up to 30 days without growth factor addition or medium change providing a cost effective method to preserve HSPC in culture.

Objects of the invention

An object of the present invention is to provide a method for
20 preservation

-4-

of Human hematopoietic stem/progenitor cells.

Another object of the present invention is to provide a method for preservation of Human hematopoietic stem/progenitor cells with the effect of mannose binding lectins on human hematopoietic stem/progenitor cells.

Still another object of the present invention is to provide a cost effective method for preservation of Human hematopoietic stem/progenitor cells by using lectins.

Summary of the invention

According to this invention there is provided a method for preservation of Human hematopoietic stem/progenitor cells, wherein the said method comprising the step of:

- a) providing the purified lectins from the group consisting of Dolichos lablab lectin (DL), Garlic lectin (GL), Musa paradisiaca (BL), Artocarpus integrifolia (AL) etc. by dialyzing against phosphate buffered saline (10mM, pH7.4) followed by filtration through 0.2 μ m filters followed by protein determination by using micro BCA kit to check protein loss;

-5-

- b) isolating the CD34⁺ cells from human cord blood sample;
- c) incubating the isolated CD34⁺ cells obtained from step (b) with different concentration of lectins ranges between 100pg/ml to 10ng/ml for a period 10 to 30 days followed by toxicity test;
- 5 d) subjecting incubated CD34⁺ cells obtained from step (c) to CFU assay or seeding on irradiating M210B4 feeders for LTC-IC assays;
- e) interpreting the effect of lectins on human hematopoietic stem/progenitor cells preservation.

Brief description of accompanying figures :

- 10 **Figure 1** represents Mannose specific lectins DL, GL, AL and BL preserve HSPC for 10 days in culture. CD34⁺ cells were incubated with various concentrations of lectins as indicated and the cells were then subjected to CFU assay. The colonies were scored for the presence of CFU-GM, CFU-GEMM and BFU-E. The data represented
15 in this figure are expressed as total progenitors preserved as percent of control cells.

-6-

5 **Figure 2** represents comparative efficacy of GL, BL, AL and DL on CD34⁺ cells isolated from the same sample. A: CFU generated per 1x10⁴ CD34⁺ cells after 10 days incubation with or without lectins B: Total number of progenitors preserved expressed as percent of control.

10 **Figure 3** represents specificity of lectin action: Use of specific sugar Methyl α Mannoside (20mM) in the incubation medium along with lectin abrogates the preservation effect. CD34⁺ cells were incubated with or without lectins for 10 days in the presence or absence of 20mM Methyl α Mannoside. The recovered cells were subjected to standard CFU assay and the colonies were scored at 14 days.
A: Experiment with BL. B: Experiment with GL (* $p \leq 0.05$ ** $p \leq 0.001$)

15 **Figure 4** represents effect of AL, BL, DL and GL in preservation of primitive cells giving rise to LTC-IC. CD34⁺ cells were incubated with 100 and 200 pg/ml of AL, BL, DL and GL independently and then seeded on irradiated M210B4. The cultures were maintained for 5 weeks by weekly demi-defoliation. Both adherent and non-adherent cells were pooled and assayed for colony formation.

-7-

A: Total number of LTC-IC preserved in CD34⁺ cells incubated with AL, BL, DL and GL after 30 days of incubation.

B: Data in A expressed as percent of control.

Detailed description of the invention :

5 Ex vivo preservation of hematopoietic stem and progenitor cells without the addition of growth factors and without the change in medium may be of immense use in various clinical and experimental set ups. Such incubation in culture media may be required for in vitro purging of tumor cells without affecting the quiescent HSPC. It may
10 also help in removal of factor dependent differentiated cells leaving behind primitive HSPC, which can be used for further manipulations including genetic engineering.

In the present invention, some of the mannose binding lectins of defined specificities have been examined for their preservation effect
15 on CD34⁺ cells isolated from cord blood. It has been observed that these lectins indeed have a HSPC preservation activity for up to 30 days without change of medium and without addition of growth factors as assessed by CFU and LTC-IC assays. The optimal concentration for all lectins ranged from 100 – 200 pg/ml. Since the

-8-

binding specificities of these lectins are well characterized (Singh, D.D. and Surolia, A. and Vijayan, M. 2004 Acta Crystallographica D60, 2104 – 2106.) so these lectins can be used as specific and well-defined tools in signal transduction studies.

5 Accordingly the present invention provides a method for preservation of Human hematopoietic stem/progenitor cells, wherein the said method comprising the step of:

a) providing the purified lectins from the group consisting of Dolichos lablab lectin (DL), Garlic lectin (GL), Musa paradisiaca (BL),
10 Artocarpus integrifolia (AL), etc. by dialyzing against phosphate buffered saline (10mM, pH 7.4) followed by filtration through 0.2 μ filters followed by protein determination by using micro BCA kit to check protein loss;

b) isolation of CD34⁺ cells from human cord blood sample;

15 c) incubation of the isolated CD34⁺ cells obtained from step (b) with different concentration of lectins ranges between 100pg/ml to 10ng/ml. for a period 10 to 30 followed by toxicity test;

-9-

- d) subjecting incubated CD34⁺ cells obtained from step (c) to CFU assay or seeding on irradiating M210B4 feeders for LTC-IC assays;
- e) interpreting the effect of lectins on human hematopoietic stem/progenitor cells preservation.

5 In an embodiment of the present invention, lectins used are selected from the group consisting of Dolichos lablab lectin (DL), garlic lectin (GL; *Allium sativum* isoelectin II, viz. ASaII), *Musa Paradisica* (BL) and the mannose binding lectin from *Artocarpus integrifolia* (artocarpin) are purified according to the procedures described in
10 Singh et al. (2004 Acta Crystallographica D60, 2104 – 2106), Mo et al (2001 Eur. J. Biochem. 268, 2609 – 2615), Dam et al (1998 J. Bio. Chem. 273, 5528 – 5535) and Misquith et al (1994 J. Biol. Chem. 269, 30393 – 30401).

15 In another embodiment of the present invention, the homogeneity of purified lectins is confirmed by SDS-polyacrylamide gel electrophoresis.

In yet another embodiment of the present invention, lectins are dialyzed against phosphate buffer saline and filtered through 0.2 to 0.4

-10-

μ filters.

In still another embodiment of the present invention, protein content is determined by using micro BCA kit to check protein loss and aliquots are kept at 2-8°C.

- 5 In yet another embodiment of the present invention, Human cord blood is taken from local hospitals after obtaining informed consent of the mother.

10 In yet another embodiment of the present invention, human cord blood sample (40 to 100 ml depending upon the sample) is collected in sterile bottle containing 40IU of heparin as an anticoagulant per ml of plain culture medium (IMDM, DMEM, RPMI-1640 etc).

15 In yet another embodiment of the present invention, Red blood cells (RBCs) are separated from mononuclear cells (MNC) by mixing HES (1ml; Sigma, hetastarch 6% solution in 0.9% NaCl) and cord blood (1-4ml).

In yet another embodiment of the present invention, suspending the MNC obtained from cord blood by HES separation as described above

-11-

in isolation buffer (0.6% Na-citrate and 2% BSA in 10mM PBS, pH7.4) and isolating the CD34⁺ cells from MNC by immuno-magnetic separation (Dyna) as per the manufacturer's instructions.

5 In yet another embodiment of the present invention, the isolated CD34⁺ cells are incubated with different concentration of lectins ranges between 100pg/ml to 10ng/ml for a period of 10 to 30 days.

In yet another embodiment of the present invention, **concentration range of lectins** used is in the range between 10pg/ml to 80ng/ml for performing toxicity test on TF1 cells using MTT or XTT assays.

10 In yet another embodiment of the present invention, no cytotoxicity is observed in lectin preparations up to 80 ng/ml.

In yet another embodiment of the present invention, CD34⁺ cells were incubated with lectins for various time periods ranging from 10 to 30 days and were subjected to CFU assay or were seeded on irradiated M210B4 feeders for LTC-IC assay.

15

In yet another embodiment of the present invention, the incubated CD34⁺ cells were tested for the number of progenitors preserved by scoring for different types of colonies (such as BFU-E, CFU-GM and

-12-

CFU-GEMM) formed in a semisolid methylcellulose medium (Methyl
cellulose 1%+ Plain IMDM 70 ml+ Human FCS 30% + Na- Pyruvate
10 μ M + Iron saturated Transferrin 0.036% + β Mercapto ethanol
5X10⁻⁵ M + BSA 1%) containing standard growth factors qualified for
5 human clonogenic cells such as hSCF 20ng/ml, GM-CSF- 2ng/ml,
IL3 4ng/ml and Epo 21U/ml. The CFU assay gives an estimate of
different types of progenitors present in a test population.

In yet another embodiment of the present invention, the total numbers
of progenitors preserved by lectins were calculated by normalizing
10 with the number of cells recovered after incubation with lectins for
various time periods.

If "x" is the number of colonies obtained from 1x10⁵ cells (recovered
after incubation with lectins for specified period) in CFU assay and
"y" is the total number of recovered cells then the "absolute number
15 of progenitors preserved" becomes $x y / 1x10^5$

In yet another embodiment of the present invention, the incubated
CD34⁺ cells were optionally seeded on irradiated (8000 rads) M210B4
feeder layer formed on collagen-coated wells using Myelocult
medium (Stem Cell Technology, Canada) followed by maintaining the

-13-

culture for 5-6 weeks by weekly demi-defoliation.

In yet another embodiment of the present invention, the said cultures were harvested by trypsinization.

5 In an embodiment of the present invention, adherent (feeder layer and the hematopoietic cells closely associated with it) and non adherent (floating hematopoietic cells) populations are pooled from harvested cultures to assay for colony formation.

10 The following examples are given by the way of illustration and therefore should not be construed to limit the scope of the present invention.

Example 1 :

Purification of lectins:

15 In the present invention lectins were used namely, Dolichos lablab (DL), Banana lectin (BL) and Artocarpus integrifolia lectin, artocarpin (AL) and garlic lectin (GL). They were purified and characterized as follows.

-14-

Lectins from Dolichos lablab lectin (DL), garlic lectin (GL; Allium sativum isoelectin II, viz. ASII), Musa Paradisica (BL) and the mannose binding lectin from Artocarpus integrifolia; artocarpin (AL) were purified following the procedures described in Singh et al (2004
5 Acta Crystallographica D60, 2104-2106; Mo et al. (2001, Eur. J. Biochem. 268, 2609-2615; Dam et al. (1998, J. Bio. Chem. 273, 5528-5535) and Misquith et al (1994, J. Biol. Chem. 269, 30393-30401) Homogeneity of the purified lectins was confirmed by SDS-polyacrylamide gel electrophoresis (Laemmli, U.K. 1970 Nature 277,
10 680-685).

The lectin preparations were dialyzed extensively against phosphate buffered saline and filtered through 0.2-micron filters. Protein content was determined using micro BCA kit and aliquots were kept at 4°C. The protein determination was frequently done to check protein loss,
15 if any.

Example 2 :

Isolation of CD34⁺ cells from human cord blood:

Cord blood was collected from local hospitals after obtaining informed consent of the mother. The protocols described in these

-15-

studies were approved by the Institutional Ethics Committee. In brief, cord blood was collected in a sterile bottle containing heparin (40IU of heparin/ml of plain medium) and processed soon thereafter by using standard protocols. Red blood cells (RBCs) were separated by
5 HES (4ml of blood: 1 ml of HES) and mononuclear cell (MNC) fraction was isolated by density gradient (Histopaque, Sigma) using standard protocols. MNCs were suspended in isolation buffer and CD34⁺ cells were isolated by immuno-magnetic separation (Dyna) as per the manufacturer's instructions.

10 **Example 3 :**

Functional assays for determination of HSPC activity:

Isolated CD34⁺ cells were incubated with different concentrations (100 pg/ml to 10 ng/ml for all four lectins) of lectins for various time periods (10 to 30 days). This concentration range of lectins used was
15 predetermined by carrying out toxicity tests on TF 1 cells using MTT or XTT assays. The CD34⁺ cells incubated for various time periods were subjected directly to CFU assay or seeded on irradiated M210B4 feeders for

LTC-IC assays as described below. CD34⁺ cells without addition of
20 lectin were kept as control in each experiment performed.

-16-

(a) Colony Forming Unit (CFU) Assay:

CD34⁺ cells incubated for various time periods with the lectins were tested for the number of progenitors preserved by scoring for the different types of colonies (BFU-E, CFU-GM and CFU-GEMM) formed in a semisolid methylcellulose medium containing standard growth factors qualified for human clonogenic cells (hSCF 20ng/ml, GM-CSF-2ng/ml, IL3 4ng/ml and Epo 2IU/ml) using standard criteria (Eaves C and Lambi K (1995) Atlas of Human Hematopoietic colonies. Stem Cell Technologies, Vancouver, Canada). This assay formed a primary screen for the lectins and the optimal concentrations of lectins giving good results in this system were used for long-term assays.

(b) Long Term Culture Initiating Cell (LTC-IC) Assays:

This assay gives an estimate of more primitive HSPC present in the test population than those scored by CFU assays (Sutherland et al 1989 Blood 74: 1563-1570). The CD34⁺ cells incubated with various concentrations and periods as indicated in the results were seeded on irradiated (8000 rads) M210B4 feeder layer formed on collagen coated wells using Myelocult medium. The cultures were maintained for 5-6 weeks by weekly demi-defoliation. The cultures were then harvested by trypsinization. Both adherent and non-adherent

-17-

populations were pooled and assayed for colony formation.

Example 4 :

Determination of optimal concentration of lectins for HSPC preservation:

5 We selected the range of various lectins based on the cytotoxicity assay carried on TF 1 cells. We observed that the lectin preparations did not show any cytotoxicity up to 80ng/ml concentration (data not shown). It was, however, necessary to determine the optimal concentration of the various lectins to be used in the context of their
10 HSPC preservation ability. In order to achieve this we incubated the CD34⁺ cells for a fixed time interval of 10 days with different concentrations of lectins (from 100-400 pg/ml and 10 ng/ml). After incubation cells were subjected to CFU assays as described. Numbers of total progenitors preserved were calculated from the number of
15 colonies formed from recovered cells after incubation with lectin(s). It was observed that all lectins namely AL, BL, DL and AL preserved the progenitors at least up to 10 days of incubation without any medium changes as well as without any cytokine supplementation (Fig 1). Lower concentrations of all lectins (in the range of 100pg/ml
20 and 200pg/ml) were found to be optimal in the context of the

-18-

preservation of HSPC in several experiment (N=10) and, therefore, these concentrations were used in further experiments.

Table 1: Determination of Optimal Concentration of Lectins

Lectin used	Total progenitors preserved after 10 days of incubation by				
	Cells	100pg/ml	200pg/ml	400pg/ml	10ng/ml
AL	357	528	717	489	408
BL	579	1064	734	806	441
DL	3294	7233	5231	4855	3908
GL	495	1925	1808	1562	931

Table 2: Total Progenitors Preserved By Lectins Expressed As Percent of Control

Lectin Used	Total progenitors preserved by lectins Expressed as percent of control				
	Cells	100pg/ml	200pg/ml	400pg/ml	10ng/ml
AL	100	148	201	137	114
BL	100	184	127	139	76
DL	100	220	159	147	119
GL	100	389	365	316	188

-19-

Example 5 :

Comparative efficacy of lectin preparations on HSPC preservation:

The experiments described above were carried out on different
5 samples as the yield of CD34⁺ cells from individual samples is not
enough to carry out experiments with all lectins as a set and thus it
was not possible to compare their efficacy. We therefore carried out
experiment on CD34⁺ cells isolated from single CB samples which
yielded enough CD34⁺ cells for such experiment using the optimal
10 concentrations of 100 and 200 pg/ml. (N=2) It was observed that the
number of CFU per 10⁴ incubated cells was not significantly different
in various sets (Fig 2 a) but when the data were normalized for total
number of progenitors preserved the effect of lectins over control was
evident (Fig 2b). This analysis indicates that probably the lectins
15 preserved more number of HSPC but probably did not affect the
frequency of clonogenic cells. In this particular experiment BL at
100pg/ml and DL at both 100 and 200 pg/ml were found to be more
effective.

-20-

Table 3: Comparative Efficacy of Lectins

	Total progenitors preserved by			
Lectin conc. Used	AL	BL	DL	GL
Cells	337	337	337	337
100pg/ml	407	665	769	462
200pg/ml	382	560	708	585

Table 4: Comparative Efficacy of Lectins Expressed as Percent of Control

	Total progenitors preserved by			
Lectin conc. used	AL	BL	DL	GL
Cells	100	100	100	100
100pg/ml	121	197	228	137
200pg/ml	113	166	210	174

Example 6 :

-21-

Determination of specificity of the lectin action:

We have used mannose-binding lectins in this set of experiments. We therefore set experiments using Methyl α Mannoside in the incubation medium to see if this specific sugar abolishes the HSPC preservation activity of the lectins. CD34⁺ cells were incubated with BL or GL (100 and 200pg/ml) with or without methyl α mannoside sugar (20mM) for 10 days. It was observed that presence of the sugar abrogated the HSPC preservation activity of both BL and GL as seen by the near control colony formation in the sets with sugar while significant preservation activity was observed with lectins. (Fig 3 A and B).

Example 7 :

Effect of lectins on preservation of LTC-IC:

In order to examine whether the lectins also preserved early progenitors we carried out the LTC-IC assay on CD34⁺ cells incubated with the optimal concentrations of lectins (100 and 200 pg/ml) for 30 days. After the incubation with lectins the surviving cells were seeded on irradiated (8000 rads) M210B4 feeder layer using Myelo-cult medium. The cultures were maintained for 5 weeks

-22-

with weekly demi-defoliation. After 5 weeks the cultures were harvested by trypsinization. Both adherent and non-adherent cells were pooled and CFU assay was carried out. It was observed (Fig 4 A and B,) that all lectins could preserve early progenitors up to 30 days of incubation.

Advantages:

1. The invention provides a simple cost effective method for preservation of hematopoietic stem progenitor cells in culture.
2. The invention provides a tool to study signaling mechanisms involved in the maintenance of quiescence in HSPC.
3. The invention provides a method to enrich the HSC for further manipulations.

-23-

We Claim:

1. A method for preservation of Human hematopoietic stem/progenitor cells,
wherein the said method comprising the step of:
 - 5 a. providing the purified lectins from the group consisting of Dolichos lablab lectin (DL), Garlic lectin (GL), Musa paradisiaca (BL), Artocarpus integrifolia (AL) etc. by dialyzing against phosphate buffered saline (10mM, pH7.4) followed by filtration through 0.2 μ m filters followed by protein determination
10 by using micro BCA kit to check protein loss;
 - b. isolating the CD34⁺ cells from human cord blood sample;
 - c. incubating of the isolated CD34⁺ cells obtained from step (b) with different concentration of lectins ranges between 100pg/ml to 10ng/ml for a period 10 to 30 days followed by toxicity test;
 - 15 d. subjecting incubated CD34⁺ cells obtained from step (c) to CFU assay or seeding on irradiating M210B4 feeders for LTC-IC assays;

-24-

- e. interpreting the effect of lectins on human hematopoietic stem/progenitor cells preservation.
2. The method as claimed in claim 1, wherein aliquots of lectins is kept at about 4°C.
 - 5 3. The method as claimed in claim 1, wherein Human cord blood is collected from local hospitals after obtaining informed consent of the mother.
 4. The method as claimed in claim 1, wherein human cord blood sample is collected in sterile bottle containing 40IU of heparin as an anticoagulant per ml to 5 ml of plain medium followed by the
10 separation of RBCs and MNCs fraction by HES in the ratio ranges from 4 ml of cord blood to 1 ml of HES through density gradient.
 5. The method as claimed in claim 1, wherein mononuclear cells
15 (MNC) is suspended in isolation buffer (0.6% Na-citrate and 2% BSA in PBS) and isolating the CD34⁺ cells by immuno-magnetic separation.

-25-

6. The method as claimed in claim 1, wherein toxicity test is carried out on TF1 cells using MTT or XTT assays.
7. The method as claimed in claim 1, wherein incubated CD34⁺ cells is tested for the number of progenitors preserved by scoring for the different types of colonies such as BFU-E, CFU-GM and CFU-GEMM etc. in a semisolid methylcellulose medium (1% Methyl cellulose + Plain IMDM + Human FCS 30% + Na-Pyruvate 10 μ M + Transferrin + β Mercapto ethanol + BSA) containing standard growth factors qualified for human clonogenic cells such as hSCF 20ng/ml, GM-CSF-2ng/ml, IL3 4ng/ml and Epo 2IU/ml by using standard criteria.
8. The method as claimed in claim 1, wherein number of progenitors preserved by lectins is estimated by obtaining CFU from recovered cells after incubation with lectins and calculating the total number of progenitors preserved by lectins by normalizing with the input to provide primary screen for the lectins.
9. The method as claimed in claim 1, wherein seeding the incubated CD34⁺ cells on irradiated (8000 rads) M210B4 feeders layer formed on collagen coated wells using Myelocult medium

-26-

followed by maintaining the culture for 5-6 weeks by weekly demi-defoliation.

10. The method as claimed in claim 1, wherein harvesting of the culture is carried out by trypsinization.
- 5 11. The method as claimed in claim 1, wherein adherent and non-adherent populations from harvested cultures is pooled to assay for colony formation.
12. The method for preservation of Human hematopoietic stem/progenitor cells substantially as herein described with
10 reference to the examples and figures accompanying this specification.