


## PYRIDINONE AND PYRIMIDINONE DERIVATIVES AS FACTOR XIA INHIBITORS

## TECHNICAL FIELD

The present invention relates to a series of pyridinone and pyrimidinone derivatives which are useful as inhibitors of factor XIa.

Thus, the present invention relates to a compound of formula (I):



(wherein all symbols have the same meanings as described hereinafter) or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, use of such compounds in treatment and/or prevention of a thromboembolic disease and processes for the preparation of said compounds.


## BACKGROUND OF THE INVENTION

Thromboembolism is an important cause of morbidity and mortality. It occurs when a blood clot breaks free and is carried by the blood stream to obstruct a blood vessel at another site. Thromboembolic disease includes venous thromboembolism, for example deep vein thrombosis or pulmonary embolism, arterial thrombosis, stroke and myocardial infarction.

Thromboembolic diseases may be treated using anticoagulants. One approach has been to target the inhibition of factor XIa (FXIa). Factor XIa is a plasma serine protease involved in the regulation of blood coagulation. Factor XIa is an activated form of factor XI, which is activated by factor XIIa, thrombin, and it is also autocatalytic. FXIa is a member of the "contact pathway" and activates factor IX by selectively cleaving Arg-Ala and Arg-Val peptide bonds. Factor IXa, in turn, activates factor X. The safety of this target is supported by the observations that FXI deficiency in humans (hemophilia C) results in a mild bleeding disorder. In addition to this, the


efficacy and side effects of this target have been shown using experimental thrombosis and bleeding models in mice lacking FXI, and in baboons and rabbits treated with anti-FXI neutralizing antibodies. These results suggest that FXIa inhibitors will show a potent anti-thrombotic effect without bleeding. Therefore, factor XIa is an attractive target for anti-thrombotic therapy without any bleeding side effect.

It has been described in Patent literature 1 that compounds of formula (A):




wherein  $A^A$  represents a 5- to 12- membered heterocycle, etc.;  $L_{1A}$  represents -CH=CH-, etc.;  $R^{11A}$  represents benzyl, etc.;  $M^A$  represents imidazolyl, etc; are useful as selective inhibitors of factor XIa or dual inhibitors of FXIa and plasma kallikrein.

Furthermore, it has been described in Patent literature 2 that a compound of formula (B-I):




wherein  $A^B$  represents a 5- to 12- membered heterocycle, etc.;  $L_{1B}$  represents -CH=CH-, etc.;  $R^{11B}$  represents benzyl, etc.;  $R^{3B}$  represents phenyl, etc.;  $R^{4B}$  represents chlorine, etc.;  $R^{8aB}$  represents hydrogen, etc; or formula (B-II):




wherein  $M^B$  represents pyridyl, etc.; and the other symbols have the same meanings as described above; inhibit factor XIa and/or plasma kallikrein.

Furthermore, it has been described in Patent literature 3 that compounds of formula (C):




wherein  $W^C$  represents CO, etc.;  $G^C$  represents a direct bond, etc.;  $G^{1C}$ ,  $G^{2C}$ ,  $G^{3C}$  and  $G^{4C}$  each independently represents C or N, etc.;  $R^{9C}$  represents aryl, etc.;  $R^{10C}$  represents heteroaryl, etc.;  $R^{1AC}$  represents heteroarylalkyl, etc.; are useful as gamma secretase modulators, however, it is not reported that the compound represented by formula (C) has factorXIa inhibitory activity.

Furthermore, it has been described in Patent literature 4 that compounds of formula (D):



wherein  $R_{1D}$  represents hydrogen, etc.;  $R_{2D}$  represents aryl, etc.;  $R_{3D}$  represents hydrogen, etc.;  $R_{4D}$  represents hydrogen, etc.;  $R_{5D}$  represents heteroarylalkyl, etc.; is useful as p38 MAP kinase modulator.

Furthermore, it has been described in Patent literature 5 that compounds of formula (E):



wherein  $L^E$  represents a linker providing 0-6 atoms, etc.;  $X^E$  represents heteroaryl, etc.;  $Z^E$  represents halogen, etc.;  $Q^E$  represents CO, etc.;  $R^{2E}$  and  $R^{3E}$  each independently represents hydrogen, aryl, etc.; are useful as dipeptidyl peptidase inhibitors.

[Patent literature 1] WO2007070826

[Patent literature 2] WO2008076805

[Patent literature 3] WO2009076337

[Patent literature 4] WO2003068230

5 [Patent literature 5] EP1506967A1


## DISCLOSURE OF THE INVENTION

It is desirable to find new compounds which may be more effective in treating thromboembolic diseases. Advantageous compounds desirably have good inhibitory  
10 activity and selectivity for factor XIa (and preferably also for plasma kallikrein) with potent anticoagulant activity and/or good oral bioavailability.

The present inventors have made extensive studies to find a compound that can become a therapeutic agent for thromboembolic diseases. As a result, we have found that the object is achieved by a compound represented by formula (I), a salt thereof, an  
15 N-oxide thereof, a solvate thereof, or a prodrug thereof (hereinafter, which may be abbreviated to compounds of the present invention) with good factor XIa inhibitory activity, selectivity for factor XIa, potent anticoagulant activity and/or good oral bioavailability, and then we have completed the present invention.

Namely, the present invention relates to the following aspects.

20 (1) A compound represented by formula (I):



wherein Cyc<sup>1</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

Cyc<sup>2</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-  
25 C10 aryl or 5- or 6-membered heteroaryl;

Cyc<sup>3</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-  
C10 aryl or 5- to 10-membered heteroaryl;



R<sup>1</sup> represents (1) C1-8 alkyl, (2) C2-8 alkenyl, (3) C2-8 alkynyl, (4) halogen, (5) nitro, (6) cyano, (7) oxo, (8) amidino, (9) -C1-8 alkylene -OR<sup>8</sup>, (10) -OR<sup>9</sup>, (11) -COOR<sup>10</sup>, (12) -C1-4 alkylene -COOR<sup>11</sup>, (13) -NHC(O)-C1-4 alkyl, (14) -C1-4 alkylene-O-C(O)-C1-8 alkyl or (15) -NHC(O)O-R<sup>12</sup>,

5 wherein R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup> and R<sup>12</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

10 s represents an integer of 0 to 6,

wherein s represents an integer of 2 to 6, each R<sup>1</sup> may be same or different;

R<sup>2</sup> represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) Cyc<sup>4</sup> or (6) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from halogen, nitro, trifluoromethyl, cyano, Cyc<sup>5</sup>, -NR<sup>13</sup>R<sup>14</sup>, -OR<sup>15</sup>,  
15 -SR<sup>16</sup>, -NHC(O)-Cyc<sup>6</sup>, -NHC(O)-C1-8 alkyl, -NHC(O)O-R<sup>17</sup> and Cyc<sup>5</sup> substituted with 1 to 3 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino and -OR<sup>18</sup>,

wherein R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup> and R<sup>18</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6)  
20 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

Cyc<sup>4</sup>, Cyc<sup>5</sup> and Cyc<sup>6</sup> each independently represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

25 R<sup>3</sup> represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) Cyc<sup>7</sup> or (6) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from halogen, nitro, trifluoromethyl, cyano, Cyc<sup>8</sup>, -NR<sup>19</sup>R<sup>20</sup>, -OR<sup>21</sup>, -SR<sup>22</sup>, -NHC(O)-Cyc<sup>9</sup>, -NHC(O)-C1-8 alkyl, -NHC(O)O-R<sup>23</sup> and Cyc<sup>8</sup> substituted with 1 to 3 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro,  
30 trifluoromethyl, cyano, oxo, amidino and -OR<sup>24</sup>,

wherein R<sup>19</sup>, R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup>, R<sup>23</sup> and R<sup>24</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl

or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

Cyc<sup>7</sup>, Cyc<sup>8</sup> and Cyc<sup>9</sup> each independently represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

5 Y represents N or C(R<sup>5</sup>);

R<sup>4</sup> and R<sup>5</sup> each independently represents (1) hydrogen, (2) halogen, (3) C1-4 alkyl, (4) C3-C8 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl or (8) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and  
10 5- to 10-membered heteroaryl;

R<sup>2</sup> and R<sup>3</sup> may be taken together to form C2-8 alkylene; or

R<sup>3</sup> and R<sup>4</sup> may be taken together to form C2-8 alkylene; wherein one carbon of the alkylene chain may be replaced by oxygen or sulfur;

R<sup>6</sup> represents (1) C1-8 alkyl, (2) C2-8 alkenyl, (3) C2-8 alkynyl, (4) Cyc<sup>10</sup>, (5)  
15 halogen, (6) nitro, (7) cyano, (8) oxo, (9) amidino, (10) -OR<sup>25</sup>, (11) -COOR<sup>26</sup>, (12) -C1-4 alkylene -COOR<sup>27</sup>, (13) -NHC(O)-C1-4 alkyl, (14) -NHC(O)-H, (15) -NHC(O)O-R<sup>28</sup>, (16) trifluoromethyl, (17) -NHC(NH)NH<sub>2</sub>, (18) -C(O)-C1-4 alkyl or (19) Cyc<sup>10</sup> substituted with 1 to 5 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino, -OR<sup>29</sup>, -SR<sup>30</sup>, -NR<sup>31</sup>R<sup>32</sup>, -  
20 NHC(O)NR<sup>33</sup>R<sup>34</sup>, -NHC(O)-C1-4 alkylene -COOH, -NH-S(O)-C1-4 alkyl, -NH-S(O)<sub>2</sub>-C1-4 alkyl, -COOR<sup>35</sup>, -NHC(O)-R<sup>36</sup>, -NHC(O)O-R<sup>37</sup>, -C(O)NH-R<sup>38</sup> and -OC(O)NH-R<sup>39</sup>,

wherein Cyc<sup>10</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

25 R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup>, R<sup>30</sup>, R<sup>31</sup>, R<sup>32</sup>, R<sup>33</sup>, R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup>, R<sup>37</sup>, R<sup>38</sup> and R<sup>39</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and  
30 5- to 10-membered heteroaryl;

m represents an integer of 0 to 6,

wherein m represents an integer of 2 to 6, each R<sup>6</sup> may be same or different;

$R^7$  represents (1) C1-8 alkyl, (2) C2-8 alkenyl, (3) C2-8 alkynyl, (4) halogen, (5) nitro, (6) trifluoromethyl, (7) cyano, (8) oxo, (9) amidino, (10)  $-OR^{40}$ , (11)  $-SR^{41}$ , (12)  $-NR^{42}R^{43}$ , (13)  $-NHC(O)NR^{44}R^{45}$ , (14)  $-NHC(O)-C1-4$  alkylene  $-NR^{46}R^{47}$ , (15)  $-NHC(O)-C1-4$  alkylene  $-COOH$ , (16)  $-NH-S(O)-C1-4$  alkyl, (17)  $-NH-S(O)_2-C1-4$  alkyl, (18)  $-COOR^{48}$ , (19)  $-NHC(O)-R^{49}$ , (20)  $-NHC(O)-C1-4$  alkylene  $-OR^{50}$ , (21)  $-NHC(O)O-R^{51}$ , (22)  $-NHC(O)O-C1-4$  alkylene  $-OR^{52}$ , (23)  $-C(O)NH-R^{53}$ , (24)  $-OC(O)NH-R^{54}$ , (25)  $-OC(O)-R^{55}$ , (26)  $-C(O)-R^{56}$ , (27)  $-CH(OH)-R^{57}$ , (28)  $-C1-4$  alkylene  $-NH_2$ , (29)  $-C1-4$  alkylene  $-OH$ , (30)  $-C1-4$  alkylene  $-OC(O)-C1-4$  alkyl, (31)  $-C1-4$  alkylene  $-NHC(O)-C1-4$  alkyl, (32)  $-C1-4$  alkylene  $-NHC(O)O-C1-4$  alkyl, (33)  $-C1-4$  alkylene  $-NHC(O)-CF_3$ , (34)  $-C1-4$  alkylene  $-NHC(O)NH-C1-4$  alkyl, (35)  $-CH=N-OR^{58}$ , (36)  $-C(O)N(C1-4 \text{ alkyl})_2$ , (37)  $-C(O)NH-R^{63}$ , (38)  $-S(O)_2-NR^{64}R^{65}$ , (39)  $-T-COOR^{66}$ , (40)  $-B(OR^{67})(OR^{68})$ , (41) C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- or 6-membered heteroaryl, which are substituted with 1 to 5 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl,  $-COOH$ ,  $-COO-C1-4$  alkyl,  $OH$ , oxo, and cyano, (42) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from halogen, trifluoromethyl,  $OH$ , oxo,  $-O-C1-4$  alkyl,  $NH_2$ , and cyano, (43)  $-N(OH)C(O)-C1-4$  alkyl, (44)  $-NHC(=N-OR^{69})-C1-4$  alkyl, (45)  $-NHC(S)-C1-4$  alkyl, (46)  $-C(S)-C1-4$  alkyl, (47)  $-S(O)NR^{70}R^{71}$ , (48)  $-C(O)NH(CO)NR^{72}$ , (49)  $-NHC(O)R^{73}$ , (50)  $-NHC(O)O-C1-4$  alkyl substituted with 1 to 5 groups selected from  $-O-C1-4$  alkylene- $O-C1-4$  alkyl,  $NH_2$ , and  $OH$ , (51)  $-NHC(O)-C1-4$  alkyl substituted with 1 to 5 groups selected from  $-O-C1-4$  alkylene- $O-C1-4$  alkyl,  $NR^{74}R^{75}$ , oxo,  $OH$ , halogen and  $-O-C1-4$  alkylene- $O-C1-4$  alkyl substituted with 1 to 2 groups selected from  $OH$ , oxo and halogen, (52)  $-NH-C1-8$  alkyl substituted with 1 to 5 groups selected from  $-O-C1-4$  alkyl, oxo,  $NR^{76}R^{77}$ , C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- or 6-membered heteroaryl, (53) C3-C8 cycloalkyl, (54) 5- to 10-membered heterocycloalkyl, (55) C5-C10 aryl or (56) 5- or 6-membered heteroaryl;

wherein  $R^{40}$ ,  $R^{41}$ ,  $R^{42}$ ,  $R^{43}$ ,  $R^{44}$ ,  $R^{45}$ ,  $R^{46}$ ,  $R^{47}$ ,  $R^{49}$ ,  $R^{50}$ ,  $R^{51}$ ,  $R^{52}$ ,  $R^{53}$ ,  $R^{54}$ ,  $R^{55}$ ,  $R^{56}$ ,  $R^{57}$ ,  $R^{58}$ ,  $R^{67}$ ,  $R^{68}$ ,  $R^{69}$ ,  $R^{70}$ ,  $R^{71}$ ,  $R^{72}$ ,  $R^{73}$ ,  $R^{74}$ ,  $R^{75}$ ,  $R^{76}$  and  $R^{77}$  each independently represents (1) hydrogen, (2) trifluoromethyl, (3) C1-8 alkyl, (4) C2-8 alkenyl, (5) C2-8 alkynyl, (6) C3-C8 cycloalkyl, (7) 5- to 10-membered heterocycloalkyl, (8) C5-C10 aryl, (9) 5- to 10-membered heteroaryl or (10) C1-4 alkyl substituted with 1 to 5 groups

selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

R<sup>63</sup> represents (1) -C1-4 alkylene-O-C1-4 alkyl, (2) -O-C1-4 alkyl, (3) cyano, (4) -C1-4 alkylene-O-C1-4 alkylene-O-C1-4 alkyl or (5) -SO<sub>2</sub>-C1-4 alkyl;

5 R<sup>64</sup> and R<sup>65</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (5) C2-8 alkynyl, (6) -C(O)-C1-4 alkyl or (7) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from OH, oxo, -O-C1-4 alkyl, -O-C1-4 alkylene-O-C1-4 alkyl, halogen, nitro and cyano;

10 T represents (1) C1-4 alkylene, (2) C2-4 alkenylene, (3) -O-C1-4 alkylene-, (4) -O-C2-4 alkenylene-, (5) -S-C1-4 alkylene-, (6) -S-C2-4 alkenylene-, (7) -NH-C1-4 alkylene-, (8) -NH-C2-4 alkenylene-, (9) -NH-C5-C10 aryl- or (10) -NH-5- to 10-membered heteroaryl-;

R<sup>48</sup> and R<sup>66</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from -NH<sub>2</sub>, -NH-C1-4 alkyl, -N(C1-4 alkyl)<sub>2</sub>, OH, oxo, -O-C1-4 alkyl, -O-C1-4 alkylene-O-C1-4 alkyl, halogen, nitro, cyano, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl (6) C3-C10 cycloalkyl, (7) 5- to 10-membered heterocycloalkyl, (8) C5-C10 aryl, (9) 5- to 10-membered heteroaryl, (10) -C1-4 alkylene-C3-C8 cycloalkyl  
20 substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH, trifluoromethyl and halogen, (11) -C1-4 alkylene-C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH, trifluoromethyl and halogen, (12) -C1-4 alkylene-5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH, trifluoromethyl and halogen, (13) -C1-4 alkylene-5- to 10-membered heteroaryl  
25 substituted with 1 to 5 groups selected from C1-4 alkyl, OH, trifluoromethyl and halogen or (14) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 5 groups selected from OH, oxo, -O-C1-4 alkyl, -O-C1-4 alkylene-O-C1-4 alkyl, halogen, nitro, cyano, C3-C8 cycloalkyl, C5-C10 aryl, 5- to 10-membered heterocycloalkyl, -O-5- to 10-membered heteroaryl, -O-C3-C8 cycloalkyl, -O-C5-C10 aryl, -O-5- to 10-membered  
30 heterocycloalkyl, -O-5- to 10-membered heteroaryl, -O-C1-4 alkylene-5- to 10-membered heteroaryl, -O-C1-4 alkylene-C3-C8 cycloalkyl, -O-C1-4 alkylene-C5-C10 aryl, -O-C1-4 alkylene-5- to 10-membered heterocycloalkyl and -O-C1-4 alkylene-5- to 10-membered heteroaryl;

n represents an integer of 0 to 6,


wherein n represents an integer of 2 to 6, each  $R^7$  may be same or different;

and

$R^{62}$  represents hydrogen or halogen,

5 a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.


(2) The compound according to (1), wherein the compound represented by formula (I) represents a compound represented by formula (I-A):



wherein U represents S or  $CH_2$ ; and

10 the other symbols have the same meanings as described above.

(3) The compound according to (1), wherein the compound represented by formula (I) represents a compound represented by formula (I-B):



wherein  $Cyc^{1-B}$  represents 5- to 10-membered heteroaryl;


15  $Cyc^{3-B}$  represents C5-C10 aryl or 5- to 10-membered heteroaryl;

$Cyc^{10-B}$  represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl,  $-COOH$ ,  $-COO-C1-4$  alkyl and  $-CONH_2$ ;

mb represents an integer of 0 to 5; and

the other symbols have the same meanings as described above.

- (4) The compound according to (1), wherein the compound represented by formula (I) represents a compound represented by formula (I-C):



wherein Cyc<sup>1-C</sup> represents 5- to 10-membered heteroaryl;


Cyc<sup>3-C</sup> represents C5-C10 aryl or 5- to 10-membered heteroaryl;

- 10 Cyc<sup>10-C</sup> represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl and -CONH<sub>2</sub>;


mc represents an integer of 0 to 5; and

the other symbols have the same meanings as described above.

- (5) The compound according to (1) or (2), wherein Cyc<sup>1</sup> represents phenyl, imidazolyl, triazolyl, pyrrolyl, pyrazolyl, furanyl, oxazolyl, thiazolyl, isothiazolyl, furazanyl, oxadiazolyl, thiadiazolyl, thienyl, pyridazinyl, indazolyl or benzimidazolyl.
- 15 (6) The compound according to (3), wherein Cyc<sup>1-B</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl.
- (7) The compound according to (4), wherein Cyc<sup>1-C</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl.
- 20 (8) The compound according to (1) or (2), wherein



represents




wherein  $R^{59}$  represents hydrogen, C1-4 alkyl or halogen;

the arrow represents a binding position; and

5 the other symbols have the same meanings as described above.

(9) The compound according to any one of (1) to (8), wherein  $Cyc^2$  represents pyridyl or phenyl.

(10) The compound according to any one of (1) to (9), wherein  $-Cyc^2-(R^6)_m$ ,  $Cyc^2(-R^6)_mCyc^{10-B}$  or  $Cyc^2(-R^6)_{mc}Cyc^{10-C}$  represents



10

wherein  $R^{60}$  represents hydrogen, methyl or halogen;

R<sup>61</sup> represents (1) hydrogen, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) -COOH, (6) -COO-C1-4 alkyl, (7) cyano or (8) -CONH<sub>2</sub>; and the arrow represents a binding position.

(11) The compound according to any one of (1) to (4), wherein Cyc<sup>2</sup> represents pyridyl or phenyl.

(12) The compound according to any one of (1) to (11), wherein R<sup>7</sup> represents (1) C1-8 alkyl, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) cyano, (6) oxo, (7) -OR<sup>40</sup>, (8) -NR<sup>42</sup>R<sup>43</sup>, (9) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (10) -NHC(O)-C1-4 alkylene-NR<sup>46</sup>R<sup>47</sup>, (11) -NHC(O)-C1-4 alkylene-COOH, (12) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (13) -COOR<sup>48</sup>, (14) -NHC(O)-R<sup>49</sup>, (15) -NHC(O)-C1-4 alkylene-OR<sup>50</sup>, (16) -NHC(O)O-R<sup>51</sup>, (17) -NHC(O)O-C1-4 alkylene-OR<sup>52</sup>, (18) -C(O)NH-R<sup>53</sup>, (19) -OC(O)-R<sup>55</sup>, (20) -C(O)-R<sup>56</sup>, (21) -CH(OH)-R<sup>57</sup>, (22) -C1-4 alkylene-NH<sub>2</sub>, (23) -C1-4 alkylene-OH, (24) -C1-4 alkylene-OC(O)-C1-4 alkyl, (25) -C1-4 alkylene-NHC(O)-C1-4 alkyl, (26) -C1-4 alkylene-NHC(O)O-C1-4 alkyl, (27) -C1-4 alkylene-NHC(O)-CF<sub>3</sub>, (28) -C1-4 alkylene-NHC(O)NH-C1-4 alkyl, (29) -CH=N-OR<sup>58</sup> or (30) -T-COOR<sup>66</sup>.

(13) The compound according to (1) or (2), wherein is selected from the group consisting of

- (1) methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl]phenyl]carbamate,
- (2) methyl [4-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl]phenyl]carbamate,
- (3) 4-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl)benzamide,
- (4) 6-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl)-3,4-dihydro-2(1H)-quinolinone,
- (5) methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-5-methyl-1H-imidazol-4-yl]phenyl]carbamate,
- (6) 6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone,
- (7) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzamide,
- (8) 3-[5-(3-amino-1H-indazol-6-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,



- (9) (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (10) (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (11) (3S)-3-[5-(1-amino-6-isoquinolinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (12) 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)benzamide,
- (13) ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,
- 10 (14) methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-methylphenyl]carbamate,
- (15) methyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,
- 15 (16) 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,
- (17) 2-methoxyethyl [4-(2-(7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,
- (18) methyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 20 (19) (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (20) (6S)-6-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,
- 25 (21) 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,
- (22) (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 30 (23) 2-methoxyethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- (24) 2-methoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

(25) 2-methoxyethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

5 (26) 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

(27) 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

(28) 2-methoxyethyl [4-(4-methyl-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

10 (29) 2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

(30) 2-ethoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

15 (31) 2-methoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,

(32) 3-methoxypropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

20 (33) methyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,

(34) 3-oxetanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

25 (35) 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

(36) methyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,

30 (37) 2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,

(38) (2S)-2-methoxypropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

(39) (6S)-6-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

(40) 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

(41) 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(42) 2-(2-methoxyethoxy)ethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(43) 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(44) 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(45) 2-(2-ethoxyethoxy)ethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(46) 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(47) 2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

(48) 2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

(49) 2-ethoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)phenyl]carbamate,

(50) 2-ethoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

5 (51) 2-ethoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

(52) tetrahydro-2-furanylmethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

10 (53) methyl [4-(5-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-2-yl)phenyl]carbamate,

(54) 3-hydroxy-3-methylbutyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(55) 2-(2-ethoxyethoxy)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

15 (56) 2-(2-methoxyethoxy)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(57) 2-fluoroethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

20 (58) 2-hydroxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(59) 2-hydroxy-2-methylpropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(60) (3S)-3-(4-chloro-5-{6-[(1,3-oxazol-2-ylmethyl)amino]-3-pyridinyl})-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

25 (61) 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

(62) (3S)-3-[5-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

30 (63) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]acetamide,

(64) methyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,

- (65) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(2-methoxyethoxy)benzamide,
- (66) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)phenyl]acetamide,
- 5 (67) 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)-3-pyridiny]carbamate,
- (68) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]formamide,
- 10 (69) 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-3-pyridiny]carbamate,
- (70) (1E)-N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-N'-hydroxyethanimidamide,
- 15 (71) (6S)-6-[5-(6-amino-3-pyridiny)-4-fluoro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,
- (72) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-2-fluoroacetamide,
- (73) (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-hydroxyphenyl)-20 1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (74) ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- (75) 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 25 (76) ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- (77) 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (78) 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxamide,
- 30 (79) 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxamide,

- (80) 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-2-yl]phenyl]carbamate,
- (81) 2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-2-yl]phenyl]carbamate,
- 5 (82) 2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-2-yl]phenyl]carbamate,
- (83) 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-2-yl]phenyl]carbamate,
- 10 (84) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (85) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- (86) 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 15 (87) 5-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (88) 4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 20 (89) 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (90) 4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- 25 (91) 2-(2-methoxyethoxy)ethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-2-yl)-2-pyridiny]carbamate,
- (92) 2-(4-morpholinyl)ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 30 (93) 2-(4-morpholinyl)ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(94) 2-methoxyethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

5 (95) (3S)-3-[2-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(96) 4-(2-((3S)-8-bromo-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

(97) ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

10 (98) 2-(4-morpholinyl)ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(99) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

15 (100) isobutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(101) 2-(dimethylamino)-2-oxoethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

20 (102) (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(103) [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(104) methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(105) 3-methylbutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

30 (106) isopropyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(107) 2,3-dihydro-1H-inden-5-yl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(108) phenyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(109) 2-(dimethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(110) methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(111) 2-(diethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(112) 1-(((cyclohexyloxy)carbonyl)oxy)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(113) cyclohexyl 2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-5-((1-(((cyclohexyloxy)carbonyl)oxy)ethoxy)carbonyl)-3-thienyl)-1H-imidazole-1-carboxylate,

(114) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-3-fluoro-2-thiophenecarboxylic acid,

(115) 4-(5-(7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-4-fluoro-1H-imidazol-2-yl)-2-thiophenecarboxylic acid,

(116) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-3-fluoro-2-pyridinecarboxylic acid,

(117) [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(118) [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1-((2,2-dimethylpropanoyl)oxy)methyl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,



(119) 2,3-dihydro-1H-inden-5-yl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(120) ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(121) 2-oxo-2-(1-pyrrolidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(122) 2-oxo-2-(1-piperidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(123) 2-(4-morpholinyl)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(124) isobutyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(125) isopropyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(126) (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate, and

(127) 2-(4-morpholinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate.

(14) A pharmaceutical composition which comprises the compound according to any one of (1) to (13), a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.

(15) The pharmaceutical composition according to (14), which is a factor XIa inhibitor or a factor XIa and plasma kallikrein dual inhibitor.

(16) The pharmaceutical composition according to (15), which is an agent for the treatment or prevention of a thromboembolic disease.

- (17) The compound according to any one of (1) to (13), a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, for use in the treatment of the human or animal body by therapy.
- (18) The compound according to any one of (1) to (13), a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, for use in treating or preventing a thromboembolic disease.
- (19) The compound for use according to (18), wherein the thromboembolic disease is selected from the group consisting of arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, arterial cerebrovascular thromboembolic disorders, venous cerebrovascular thromboembolic disorders and thromboembolic disorders in the chambers of the heart or in the peripheral circulation.
- (20) The compound for use according to (19), wherein the thromboembolic disease is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis.
- (21) A method for treating a patient suffering from or susceptible to a thromboembolic disease, which comprises administering to said patient a therapeutically effective amount of a compound according to any one of (1) to (13), a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.
- (22) Use of a compound according to any one of (1) to (13), a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, in the manufacture of a medicament for use in treating or preventing a thromboembolic disease.
- (23) A method for treating a patient suffering from or susceptible to a thromboembolic disease, which comprises administering to said patient a therapeutically effective amount of a compound according to (1), a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.
- (24) The method according to (23), wherein the thromboembolic disease is selected from the group consisting of arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, arterial cerebrovascular thromboembolic

disorders, venous cerebrovascular thromboembolic disorders and thromboembolic disorders in the chambers of the heart or in the peripheral circulation.

(25) The method according to (24), wherein the thromboembolic disease is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis.

Definitions:

As used herein, a C1-8 alkyl group or moiety is a linear or branched alkyl group or moiety containing from 1 to 8 carbon atoms. Typically a C1-8 alkyl group or moiety is a C1-4 alkyl group or moiety. A C1-4 alkyl group or moiety is a linear or branched alkyl group or moiety containing from 1 to 4 carbon atoms. Examples of C1-8 alkyl groups and moieties include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, t-butyl, 3-methyl-butyl, pentyl, hexyl, heptyl, octyl, and isomers thereof. Examples of C1-4 alkyl groups and moieties include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl and t-butyl. For the avoidance of doubt, where two alkyl moieties are present in a group, the alkyl moieties may be the same or different.

In the present specification, the C2-8 alkenyl includes, for example, ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl and isomers thereof.

In the present specification, the C2-8 alkynyl includes, for example, ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl and isomers thereof.

In the present specification, the C1-4 alkylene includes, for example, methylene, ethylene, propylene, butylene, and the like.

In the present specification, the C1-8 alkylene includes, for example, methylene, ethylene, propylene, isopropylene, butylene, isobutylene, pentamethylene, hexamethylene, heptamethylene and octamethylene and isomers thereof.

5 In the present specification, the C2-4 alkenylene includes, for example, vinylene, propenylene, butenylene and isomers thereof.

In the present specification, the halogen atom includes, for example, fluorine, chlorine, bromine and iodine, and is preferably fluorine, chlorine or bromine.

10

In the present specification, the C3-C10 cycloalkyl includes, for example, cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclooctane, cyclononane, cyclodecane, cyclobutene, cyclopentene, cyclohexene, cycloheptene, cyclooctene, cyclobutadiene, cyclopentadiene, cyclohexadiene, cycloheptadiene,  
15 cyclooctadiene, pentalene, perhydropentalene, perhydroazulene, indene, perhydroindene, indan, dihydronaphthalene, tetrahydronaphthalene, perhydronaphthalene rings and the like.

Cyc<sup>1</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10  
20 aryl or 5- to 10-membered heteroaryl.

“C3-C8 cycloalkyl” refers to a C3-C8 cyclic hydrocarbon. Examples of C3-C8 cycloalkyl include cyclopropane, cyclobutane, cyclopentane, cyclohexane, cycloheptane, cyclooctane, cyclobutene, cyclopentene, cyclohexene, cycloheptene, cyclooctene, cyclobutadiene, cyclopentadiene, cyclohexadiene, cycloheptadiene, cyclooctadiene  
25 rings and the like.

“5- to 10-membered heterocycloalkyl” refers to a “5- to 10-membered mono- or bi- non-aromatic heterocyclic ring having 1 to 4 nitrogen atom(s), 1 or 2 oxygen atom(s) and/or 1 or 2 sulfur atom(s) as a hetero atom(s)”. Examples of 5- to 10-membered heterocycloalkyl include pyrazolidine, dihydropyridine, tetrahydropyridine, piperidine,  
30 dihydropyrazine, tetrahydropyrazine, piperazine, dihydropyrimidine, tetrahydropyrimidine, perhydropyrimidine, dihydropyridazine, tetrahydropyridazine, perhydropyridazine, dihydroazepine, tetrahydroazepine, perhydroazepine, dihydrodiazepine, tetrahydrodiazepine, perhydrodiazepine, dihydrofuran,

tetrahydrofuran, dihydropyran, tetrahydropyran, dihydroxepine, tetrahydroxepine,  
 perhydroxepine, dihydrothiophene, tetrahydrothiophene, dihydrothiopyran,  
 tetrahydrothiopyran, dihydrothiepin, tetrahydrothiepin, perhydrothiepin, dihydroxazole,  
 tetrahydroxazole (oxazolidine), dihydroisoxazole, tetrahydroisoxazole (isoxazolidine),  
 5 dihydrothiazole, tetrahydrothiazole (thiazolidine), dihydroisothiazole,  
 tetrahydroisothiazole (isothiazolidine), dihydrofurazan, tetrahydrofurazan,  
 dihydroxadiazole, tetrahydroxadiazole (oxadiazolidine), dihydroxazine,  
 tetrahydroxazine, dihydroxadiazine, tetrahydroxadiazine, dihydroxazepine,  
 tetrahydroxazepine, perhydroxazepine, dihydroxadiazepine, tetrahydroxadiazepine,  
 10 perhydroxadiazepine, dihydrothiadiazole, tetrahydrothiadiazole (thiadiazolidine),  
 dihydrothiazine, tetrahydrothiazine, dihydrothiadiazine, tetrahydrothiadiazine,  
 dihydrothiazepine, tetrahydrothiazepine, perhydrothiazepine, dihydrothiadiazepine,  
 tetrahydrothiadiazepine, perhydrothiadiazepine, morpholine, thiomorpholine, oxathiane,  
 indoline, isoindoline, dihydrobenzofuran, perhydrobenzofuran, dihydroisobenzofuran,  
 15 perhydroisobenzofuran, dihydrobenzothiophene, perhydrobenzothiophene,  
 dihydroisobenzothiophene, perhydroisobenzothiophene, dihydroindazole,  
 perhydroindazole, dihydroquinoline, tetrahydroquinoline, perhydroquinoline,  
 dihydroisoquinoline, tetrahydroisoquinoline, perhydroisoquinoline, dihydrophthalazine,  
 tetrahydrophthalazine, perhydrophthalazine, dihydronaphthyridine,  
 20 tetrahydronaphthyridine, perhydronaphthyridine, dihydroquinoxaline,  
 tetrahydroquinoxaline, perhydroquinoxaline, dihydroquinazoline, tetrahydroquinazoline,  
 perhydroquinazoline, dihydrocinnoline, tetrahydrocinnoline, perhydrocinnoline,  
 benzoxathiane, dihydrobenzoxazine, dihydrobenzothiazine, pyrazinomorpholine,  
 dihydrobenzoxazole, perhydrobenzoxazole, dihydrobenzothiazole,  
 25 perhydrobenzothiazole, dihydrobenzimidazole, perhydrobenzimidazole, dioxolane, 1,3-  
 dioxole, dioxane, dithiolane, dithiane, dioxaindane, benzodioxane, chroman,  
 benzodithiolane, benzodithiane, 6,7-dihydro-5H-cyclopenta[b]pyrazine, 5H-  
 cyclopenta[b]pyrazine, 2,4-dihydro-1H-benzo[d][1,3]oxazine rings and the like.

“C5-C10 aryl” refers to a “C5-10 mono- or bi- aromatic carbocyclic ring”.

30 Examples of C5-C10 aryl include benzene, azulene, naphthalene rings and the like.  
 Thus the C5-C10 aryl may be, for example, a phenyl ring and the like.

“5- to 10-membered heteroaryl” refers to a “5- to 10-membered mono- or bi-  
 aromatic heterocyclic ring having 1 to 4 nitrogen atom(s), 1 or 2 oxygen atom(s) and/or

1 or 2 sulfur atom(s) as a hetero atom(s)". Examples of 5- to 10-membered heteroaryl include pyrrole, imidazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, furan, thiophene, oxazole, isoxazole, thiazole, isothiazole, furazan, oxadiazole, thiadiazole, indole, isoindole, benzofuran, isobenzofuran, benzothiophene, isobenzothiophene, indazole, quinoline, isoquinoline, purine, phthalazine, pteridine, naphthyridine, quinoxaline, quinazoline, cinnoline, benzoxazole, benzothiazole, benzimidazole, benzofurazan, benzothiadiazole, benzotriazole, isoxazolo[4,5-d]pyridazine rings and the like.

In some embodiments, Cyc<sup>1</sup> also represents Cyc<sup>1-B</sup> or Cyc<sup>1-C</sup>. Cyc<sup>1-B</sup> and Cyc<sup>1-C</sup> independently represent 5- to 10-membered heteroaryl. The "5- to 10-membered heteroaryl" represented by Cyc<sup>1-B</sup> or Cyc<sup>1-C</sup> may be selected from any of the examples provided above for "5- to 10-membered heteroaryl".

Cyc<sup>2</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- or 6-membered heteroaryl.

The "C3-C8 cycloalkyl" represented by Cyc<sup>2</sup> may be selected from any of the examples provided above for "C3-C8 cycloalkyl".

The "5- to 10-membered heterocycloalkyl" represented by Cyc<sup>2</sup> may be selected from any of the examples provided above for "5- to 10-membered heterocycloalkyl".

The "C5-C10 aryl" represented by Cyc<sup>2</sup> may be selected from any of the examples provided above for "C5-C10 aryl".

"5- to 6-membered heteroaryl" refers to a "5- to 6-membered mono-aromatic heterocyclic ring having 1 to 4 nitrogen atom(s), 1 or 2 oxygen atom(s) and/or 1 or 2 sulfur atom(s) as a hetero atom(s)". Examples of 5- to 6-membered heteroaryl include pyrrole, imidazole, triazole, tetrazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, furan, thiophene, oxazole, isoxazole, thiazole, isothiazole, furazan, oxadiazole, thiadiazole, rings and the like.

Cyc<sup>3</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl.

The "C3-C8 cycloalkyl" represented by Cyc<sup>3</sup> may be selected from any of the examples provided above for "C3-C8 cycloalkyl".

The "5- to 10-membered heterocycloalkyl" represented by Cyc<sup>3</sup> may be selected from any of the examples provided above for "5- to 10-membered heterocycloalkyl".

The "C5-C10 aryl" represented by Cyc<sup>3</sup> may be selected from any of the examples provided above for "C5-C10 aryl".

5        The "5- to 10-membered heteroaryl" represented by Cyc<sup>3</sup> may be selected from any of the examples provided above for "5- to 10-membered heteroaryl".

In some embodiments, Cyc<sup>3</sup> also represents Cyc<sup>3-B</sup> or Cyc<sup>3-C</sup>. Cyc<sup>3-B</sup> represents C5-C10 aryl or 5- to 10-membered heteroaryl. The "C5-C10 aryl" represented by Cyc<sup>3-B</sup> may be selected from any of the examples provided above for "C5-C10 aryl". The "5-  
10        to 10-membered heteroaryl" represented by Cyc<sup>3-B</sup> may be selected from any of the examples provided above for "5- to 10-membered heteroaryl".

Cyc<sup>3-C</sup> represents C5-C10 aryl or 5- to 10-membered heteroaryl. The "C5-C10 aryl" represented by Cyc<sup>3-C</sup> may be selected from any of the examples provided above for "C5-C10 aryl". The "5- to 10-membered heteroaryl" represented by Cyc<sup>3-C</sup> may be  
15        selected from any of the examples provided above for "5- to 10-membered heteroaryl".

Cyc<sup>4</sup>, Cyc<sup>6</sup>, Cyc<sup>7</sup> and Cyc<sup>9</sup> each independently represent C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl.

The "C3-C8 cycloalkyl" represented by Cyc<sup>4</sup>, Cyc<sup>6</sup>, Cyc<sup>7</sup> or Cyc<sup>9</sup> may be  
20        selected from any of the examples provided above for "C3-C8 cycloalkyl".

The "5- to 10-membered heterocycloalkyl" represented by Cyc<sup>4</sup>, Cyc<sup>6</sup>, Cyc<sup>7</sup> or Cyc<sup>9</sup> may be selected from any of the examples provided above for "5- to 10-membered heterocycloalkyl".

The "C5-C10 aryl" represented by Cyc<sup>4</sup>, Cyc<sup>6</sup>, Cyc<sup>7</sup> or Cyc<sup>9</sup> may be selected  
25        from any of the examples provided above for "C5-C10 aryl".

The "5- to 10-membered heteroaryl" represented by Cyc<sup>4</sup>, Cyc<sup>6</sup>, Cyc<sup>7</sup> or Cyc<sup>9</sup> may be selected from any of the examples provided above for "5- to 10-membered heteroaryl".

30        Cyc<sup>5</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl, any of which may be optionally substituted with 1 to 3 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino and -OR<sup>18</sup>.

Cyc<sup>8</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl, any of which may be optionally substituted with 1 to 3 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino and -OR<sup>24</sup>.

5           The optionally substituted "C3-C8 cycloalkyl" represented by Cyc<sup>5</sup> or Cyc<sup>8</sup> may be selected from any of the examples provided above for "C3-C8 cycloalkyl".

          The optionally substituted "5- to 10-membered heterocycloalkyl" represented by Cyc<sup>5</sup> or Cyc<sup>8</sup> may be selected from any of the examples provided above for "5- to 10-membered heterocycloalkyl".

10           The optionally substituted "C5-C10 aryl" represented by Cyc<sup>5</sup> or Cyc<sup>8</sup> may be selected from any of the examples provided above for "C5-C10 aryl".

          The optionally substituted "5- to 10-membered heteroaryl" represented by Cyc<sup>5</sup> or Cyc<sup>8</sup> may be selected from any of the examples provided above for "5- to 10-membered heteroaryl".

15

          Cyc<sup>10</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl, any of which may be optionally substituted with 1 to 5 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino, -OR<sup>29</sup>, -SR<sup>30</sup>, -NR<sup>31</sup>R<sup>32</sup>, -NHC(O)NR<sup>33</sup>R<sup>34</sup>, -NHC(O)-C1-4 alkylene -COOH, -NH-S(O)-C1-4 alkyl, -NH-S(O)<sub>2</sub>-C1-4 alkyl, -COOR<sup>35</sup>, -NHC(O)-R<sup>36</sup>, -NHC(O)O-R<sup>37</sup>, -C(O)NH-R<sup>38</sup> and -OC(O)NH-R<sup>39</sup>.

20

          The optionally substituted "C3-C8 cycloalkyl" represented by Cyc<sup>10</sup> may be selected from any of the examples provided above for "C3-C8 cycloalkyl".

          The optionally substituted "5- to 10-membered heterocycloalkyl" represented by Cyc<sup>10</sup> may be selected from any of the examples provided above for "5- to 10-membered heterocycloalkyl".

25

          The optionally substituted "C5-C10 aryl" represented by Cyc<sup>10</sup> may be selected from any of the examples provided above for "C5-C10 aryl".

          The optionally substituted "5- to 10-membered heteroaryl" represented by Cyc<sup>10</sup> may be selected from any of the examples provided above for "5- to 10-membered heteroaryl".

30

          In some embodiments, Cyc<sup>10</sup> also represents Cyc<sup>10-B</sup> or Cyc<sup>10-C</sup>. Cyc<sup>10-B</sup> and Cyc<sup>10-C</sup> each independently represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-



membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl and -CONH<sub>2</sub>. The optionally substituted “5- to 10-membered heteroaryl” in options (1) or (2) of Cyc<sup>10-B</sup> and Cyc<sup>10-C</sup> may be selected from any of the examples provided above for “5- to 10-membered heteroaryl”.

5

Preferably, Cyc<sup>1</sup> represents a C5-C10 aryl or 5- to 10- membered heteroaryl, more preferably phenyl, imidazolyl, triazolyl, tetrazolyl, pyrrolyl, pyrazolyl, furanyl, oxazolyl, thiazolyl, isothiazolyl, furazanyl, oxadiazolyl, thiadiazolyl, thienyl, pyridazinyl, indazolyl or benzimidazolyl, more preferably a 5- to 10- membered heteroaryl such as imidazolyl, triazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl. In one embodiment, Cyc<sup>1</sup> preferably represents imidazolyl or triazolyl.

10

Preferably, Cyc<sup>1-B</sup> or Cyc<sup>1-C</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl, more preferably imidazolyl, triazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl.

15

Preferably, Cyc<sup>2</sup> represents a C5-C10 aryl, preferably a C5-C6 aryl, more preferably phenyl, or a 5- to 6-membered heteroaryl such as pyridyl.

Preferably, Cyc<sup>3</sup> represents (i) C5-C6 cycloalkyl, such as cyclohexane, (ii) 5- to 10-membered heterocycloalkyl such as indoline, isoindoline, dihydroquinoline, dihydroquinazoline, dihydrobenzoxazine or dihydrobenzoxazole, (iii) C5-C10 aryl, for example C5-C7 aryl, such as phenyl, or (iv) 5- to 10-membered heteroaryl, such as pyrazole, pyridine, pyrazine, thiophene, oxazole, thiazole, indazole, quinoline, isoquinoline, quinoxaline or benzimidazole. In particular, the 5- to 10-membered heterocycloalkyl such as indoline, isoindoline, dihydroquinoline, dihydroquinazoline, dihydrobenzoxazine or dihydrobenzoxazole, or 5- to 10-membered heteroaryl such as quinoline or quinoxaline may be substituted with an oxo group to form, for example, indolone, isoindolone, dihydroquinolinone, dihydroquinazolinone, benzoxazinone, benzoxazolone, quinolinone or quinoxalinone. In one embodiment, Cyc<sup>3</sup> preferably represents phenyl or pyridinyl.

20  
25  
30

Preferably, Cyc<sup>3-B</sup> or Cyc<sup>3-C</sup> represents C5-C10 aryl, for example C5-C7 aryl such as phenyl, or 5- to 10-membered heteroaryl, such as pyrazole, pyridine, pyrazine,

thiophene, oxazole, thiazole, indazole, quinoline, isoquinoline, quinoxaline or benzimidazole, more preferably pyrazole or pyridine.

Preferably, Cyc<sup>4</sup>, Cyc<sup>6</sup>, Cyc<sup>7</sup> and Cyc<sup>9</sup> each independently represent C5-C10 aryl, for example C5-C7 aryl such as phenyl.

Preferably, Cyc<sup>5</sup> and Cyc<sup>8</sup> each independently represent C5-C10 aryl, for example C5-C7 aryl such as phenyl, or a 5- to 6-membered heteroaryl such as pyridyl, any of which may be optionally substituted as set out above. Preferably, Cyc<sup>5</sup> and Cyc<sup>8</sup> are unsubstituted.

Preferably, Cyc<sup>10</sup> represents a 5 to 10 membered heteroaryl, preferably imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl, benzimidazolyl, thiazolyl or thiadiazolyl, more preferably thiadiazolyl, triazolyl or tetrazolyl, any of which may be optionally substituted as set out above.

Preferably, Cyc<sup>10-B</sup> or Cyc<sup>10-C</sup> each independently represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl, benzimidazolyl, thiazolyl or thiadiazolyl, more preferably thiadiazolyl, triazolyl or tetrazolyl, any of which may be optionally substituted with 1 to 3 groups selected from halogen, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl, cyano and -CONH<sub>2</sub>.

Preferably, R<sup>1</sup> represents C1-8 alkyl or halogen, more preferably C1-4 alkyl or halogen (preferably chlorine); and  
s represents an integer of 0 to 6,  
wherein s represents an integer of 2 to 6, each R<sup>1</sup> may be same or different.

Preferably, s represents an integer of 0, 1 or 2.

Preferably, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup> and R<sup>11</sup> each independently represents hydrogen or C1-8 alkyl, more preferably C1-4 alkyl.

Preferably, R<sup>12</sup> represents C1-8 alkyl, more preferably C1-4 alkyl.

Preferably, R<sup>2</sup> represents (1) hydrogen, (2) C1-8 alkyl, (3) Cyc<sup>4</sup> or (4) C1-8 alkyl substituted with 1 to 3 groups selected from Cyc<sup>5</sup> and -OR<sup>15</sup>, wherein Cyc<sup>4</sup> and Cyc<sup>5</sup> are preferably as set out above. More preferably, R<sup>2</sup> represents (1) hydrogen, (2) C1-4 alkyl, (3) Cyc<sup>4</sup> or (4) C1-4 alkyl substituted with 1 or 2 groups selected from Cyc<sup>5</sup> and -OR<sup>15</sup>,  
5 wherein Cyc<sup>4</sup> and Cyc<sup>5</sup> are preferably as set out above and R<sup>15</sup> is selected from hydrogen and C1-4 alkyl.

Preferably, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup>, R<sup>16</sup> and R<sup>18</sup> each independently represents hydrogen or C1-8 alkyl, more preferably C1-4 alkyl.

10

Preferably, R<sup>17</sup> represents C1-8 alkyl, more preferably C1-4 alkyl.

When not taken together with R<sup>4</sup> to form a C2-8 alkylene, R<sup>3</sup> preferably represents (1) hydrogen, (2) C1-8 alkyl, (3) Cyc<sup>7</sup> or (4) C1-8 alkyl substituted with 1 to  
15 3 groups selected from Cyc<sup>8</sup> and OR<sup>21</sup>, wherein Cyc<sup>7</sup> and Cyc<sup>8</sup> are preferably as set out above. More preferably, R<sup>3</sup> represents (1) hydrogen, (2) C1-4 alkyl, (3) Cyc<sup>7</sup> or (4) C1-4 alkyl substituted with 1 or 2 groups selected from Cyc<sup>8</sup> and OR<sup>21</sup>, wherein Cyc<sup>7</sup> and Cyc<sup>8</sup> are preferably as set out above and R<sup>21</sup> is selected from hydrogen and C1-4 alkyl. Most preferably, R<sup>3</sup> is hydrogen.

20

Preferably, R<sup>19</sup>, R<sup>20</sup>, R<sup>21</sup>, R<sup>22</sup>, and R<sup>24</sup> each independently represents hydrogen or C1-8 alkyl, more preferably C1-4 alkyl.

Preferably, R<sup>23</sup> represents C1-8 alkyl, more preferably C1-4 alkyl.

25

Preferably, R<sup>4</sup> and R<sup>5</sup> each independently represents hydrogen or C1-4 alkyl, more preferably hydrogen, methyl or ethyl.

In another embodiment, R<sup>3</sup> and R<sup>4</sup> are preferably taken together to form a C2-8  
30 alkylene, preferably a C2-4 alkylene, more preferably ethylene.

Preferably, R<sup>6</sup> represents (1) Cyc<sup>10</sup>, (2) methyl, (3) halogen (preferably chlorine), (4) amidino, or (5) Cyc<sup>10</sup> substituted with 1 to 3 groups selected from halogen

(preferably chlorine), nitro, trifluoromethyl, cyano, -OR<sup>29</sup>, -COOR<sup>35</sup>, -NHC(O)-R<sup>36</sup> and -C(O)NH-R<sup>38</sup>, wherein Cyc<sup>10</sup> is preferably as set out above. More preferably, R<sup>6</sup> represents (1) Cyc<sup>10</sup>, (2) methyl, (3) halogen (preferably chlorine), or (4) Cyc<sup>10</sup> substituted with 1 to 3 groups selected from halogen (preferably chlorine), nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl, cyano or -CONH<sub>2</sub>, wherein Cyc<sup>10</sup> is preferably as set out above; and

m represents an integer of 0 to 6,

wherein m represents an integer of 2 to 6, each R<sup>6</sup> may be same or different.

Preferably, R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>29</sup>, R<sup>30</sup>, R<sup>31</sup>, R<sup>32</sup>, R<sup>33</sup>, R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup>, R<sup>38</sup> and R<sup>39</sup> each independently represents hydrogen or C1-8 alkyl, more preferably C1-4 alkyl.

Preferably, R<sup>28</sup> and R<sup>37</sup> each independently represents C1-8 alkyl, more preferably C1-4 alkyl.

15

Preferably, m represents an integer of 0, 1 or 2.

Preferably, R<sup>7</sup> represents (1) C1-8 alkyl, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) cyano, (6) oxo, (7) -OR<sup>40</sup>, (8) -NR<sup>42</sup>R<sup>43</sup>, (9) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (10) -NHC(O)-C1-4 alkylene-NR<sup>46</sup>R<sup>47</sup>, (11) -NHC(O)-C1-4 alkylene-COOH, (12) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (13) -COOR<sup>48</sup>, (14) -NHC(O)-R<sup>49</sup>, (15) -NHC(O)-C1-4 alkylene-OR<sup>50</sup>, (16) -NHC(O)O-R<sup>51</sup>, (17) -NHC(O)O-C1-4 alkylene-OR<sup>52</sup>, (18) -C(O)NH-R<sup>53</sup>, (19) -OC(O)-R<sup>55</sup>, (20) -C(O)-R<sup>56</sup>, (21) -CH(OH)-R<sup>57</sup>, (22) -C1-4 alkylene-NH<sub>2</sub>, (23) -C1-4 alkylene-OH, (24) -C1-4 alkylene-OC(O)-C1-4 alkyl, (25) -C1-4 alkylene-NHC(O)-C1-4 alkyl, (26) -C1-4 alkylene-NHC(O)O-C1-4 alkyl, (27) -C1-4 alkylene-NHC(O)-CF<sub>3</sub>, (28) -C1-4 alkylene-NHC(O)NH-C1-4 alkyl, (29) -CH=N-OR<sup>58</sup> or (30) -T-COOR<sup>66</sup>. More preferably, R<sup>7</sup> represents (1) methyl, (2) ethyl, (3) fluorine, (4) chlorine, (5) bromine, (6) nitro, (7) trifluoromethyl, (8) cyano, (9) oxo, (10) -OR<sup>40</sup>, (11) -NR<sup>42</sup>R<sup>43</sup>, (12) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (13) -NHC(O)-C1-4 alkylene-NR<sup>46</sup>R<sup>47</sup>, (14) -NHC(O)-C1-4 alkylene-COOH, (15) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (16) -COOR<sup>48</sup>, (17) -NHC(O)-R<sup>49</sup>, (18) -NHC(O)-C1-4 alkylene-OR<sup>50</sup>, (19) -NHC(O)O-R<sup>51</sup>, (20) -NHC(O)O-C1-4 alkylene-OR<sup>52</sup>, (21) -C(O)NH-R<sup>53</sup>, (22) -OC(O)-R<sup>55</sup>, (23) -C(O)-R<sup>56</sup>, (24) -CH(OH)-R<sup>57</sup>, (25) -C1-4 alkylene-NH<sub>2</sub>, (26) -C1-4 alkylene-OH, (27) -C1-4

alkylene -OC(O)-C1-4 alkyl, (28) -C1-4 alkylene -NHC(O)-C1-4 alkyl, (29) -C1-4 alkylene -NHC(O)O-C1-4 alkyl, (30) -C1-4 alkylene -NHC(O)-CF<sub>3</sub>, (31) -C1-4 alkylene -NHC(O)NH-C1-4 alkyl, (32) -CH=N-OR<sup>58</sup> or (33) -T-COOR<sup>66</sup>, wherein R<sup>40</sup>, R<sup>41</sup>, R<sup>42</sup>, R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup>, R<sup>47</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>52</sup>, R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup>, R<sup>57</sup> and R<sup>58</sup> are

5 independently selected from hydrogen, C1-4 alkyl and trifluoromethyl;

R<sup>51</sup> represents C1-8 alkyl, more preferably C1-4 alkyl;

R<sup>48</sup> and R<sup>66</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered  
 10 heteroaryl, (4) C3-C10 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl, (8) -C1-4 alkylene-C3-C8 cycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9) -C1-4 alkylene-C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen, (10) -C1-4 alkylene-5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups  
 15 selected from C1-4 alkyl, oxo, OH and halogen, (11) -C1-4 alkylene-5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen or (12) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 3 groups selected from OH, oxo, -O-C1-4 alkyl, halogen, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, -O-C3-C8 cycloalkyl and -O-5- to 10-membered heterocycloalkyl;

20 T represents C1-4 alkylene or C2-4 alkenylene; and

n represents an integer of 0 to 6,

wherein n represents an integer of 2 to 6, each R<sup>7</sup> may be same or different.

More preferably, R<sup>7</sup> represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl, (3) -NHC(O)O-C1-4 alkylene-O-C1-4 alkyl, (4) halogen, (5) -COOH, (6) -COO-C1-8 alkyl, (7) -COO-C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, (8) -COO- C5-C10 aryl, (9) -COO- C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (10) -COO-C1-4 alkylene-O-C1-8 alkyl  
 25 substituted with 1 to 3 groups selected from OH, oxo, -O-C1-4 alkyl, halogen, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, -O-C3-C8 cycloalkyl and -O-5- to 10-membered heterocycloalkyl, (11) -C1-4 alkylene-COOH, (12) -C1-4 alkylene-COO-

C1-8 alkyl or (13) -C1-4 alkylene-COO- C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl;

n represents an integer of 0 to 3,

wherein n represents an integer of 2 to 3, each R<sup>7</sup> may be same or different.

5

Preferably, R<sup>40</sup>, R<sup>41</sup>, R<sup>42</sup>, R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup>, R<sup>47</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>52</sup>, R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup>, R<sup>57</sup> and R<sup>58</sup> each independently represents hydrogen, trifluoromethyl or C1-8 alkyl, more preferably hydrogen or C1-4 alkyl.

10

Preferably R<sup>51</sup> represents C1-8 alkyl, more preferably C1-4 alkyl.

15

Preferably, R<sup>48</sup> and R<sup>66</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl, (4) C3-C10 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl, (8) -C1-4 alkylene-C3-C8 cycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9) -C1-4 alkylene-C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen, (10) -C1-4 alkylene-5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (11) -C1-4 alkylene-5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen or (12) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 3 groups selected from OH, oxo, -O-C1-4 alkyl, halogen, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, -O-C3-C8 cycloalkyl and -O-5- to 10-membered heterocycloalkyl, more preferably (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and morpholine, (4) phenyl, (5) indane, (6) C1-4 alkylene - 1,3-dioxole substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, or (7) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 3 groups selected from oxo and -O-cyclohexane.

30


Preferably, T represents C1-4 alkylene or C2-4 alkenylene.

Preferably,  $n$  represents an integer of 0, 1, 2 or 3.


Preferably,  $R^{62}$  represents hydrogen or chlorine, more preferably hydrogen.

- 5 In a preferred embodiment,  $Cyc^3$  represents C5-C10 aryl or 5- to 10-membered heteroaryl, more preferably phenyl or thiophene,  $n$  is 1 and  $R^7$  represents (1)  $-NH_2$ , (2)  $-NHC(O)O-C1-4$  alkyl, (3)  $-NHC(O)O-C1-4$  alkylene- $O-C1-4$  alkyl, (4)  $-COOH$ , (5)  $-COO-C1-8$  alkyl, (6)  $-COO-C1-8$  alkyl substituted with 1 to 5 groups selected from  $-N(C1-4$  alkyl) $_2$ , oxo and 5- to 10-membered heterocycloalkyl, (7)  $-COO-$  C5-C10 aryl,
- 10 (8)  $-COO-$  C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9)  $-COO-$  indane, (10)  $-COO-$  C1-4 alkylene - 1,3-dioxole substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (11)  $-COO-C1-4$  alkylene- $O-C1-8$  alkyl substituted with 1 to 3 groups selected from oxo and  $-O$ -cyclohexane, (12)  $-C1-4$  alkylene- $COOH$ , (13)  $-C1-4$
- 15 alkylene- $COO-$  C1-8 alkyl or (14)  $-C1-4$  alkylene- $COO-$  C1-8 alkyl substituted with 1 to 5 groups selected from  $-N(C1-4$  alkyl) $_2$ , oxo and 5- to 10-membered heterocycloalkyl, or  $n$  is 2 and one  $R^7$  represents (1)  $-NH_2$ , (2)  $-NHC(O)O-C1-4$  alkyl or (3)  $-NHC(O)O-C1-4$  alkylene  $-O-C1-4$  alkyl and the other  $R^7$  represents halogen.

- 20 In a preferred embodiment



represents



wherein  $R^{59}$  represents hydrogen, C1-4 alkyl or halogen;


the arrow represents a binding position; and

- 5 the other symbols have the same meanings as described above, preferably wherein  $n$  is 1 and  $R^7$  represents (1)  $-NH_2$ , (2)  $-NHC(O)O-C1-4$  alkyl, (3)  $-NHC(O)O-C1-4$  alkylene  $-O-C1-4$  alkyl, (4)  $-COOH$ , (5)  $-COO-C1-8$  alkyl, (6)  $-COO-C1-8$  alkyl substituted with 1 to 5 groups selected from  $-N(C1-4$  alkyl) $_2$ , oxo and 5- to 10-
- 10 membered heterocycloalkyl, (7)  $-COO-C5-C10$  aryl, (8)  $-COO-C1-4$  alkylene  $-5-$  to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9)  $-COO-$  indane, (10)  $-COO-C1-4$  alkylene  $-1,3$ -dioxole substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (11)  $-COO-C1-4$  alkylene  $-O-C1-8$  alkyl substituted with 1 to 3 groups selected from oxo and
- 15  $-O$ -cyclohexane, (12)  $-C1-4$  alkylene  $-COOH$ , (13)  $-C1-4$  alkylene  $-COO-C1-8$  alkyl or (14)  $-C1-4$  alkylene  $-COO-C1-8$  alkyl substituted with 1 to 5 groups selected from  $-N(C1-4$  alkyl) $_2$ , oxo and 5- to 10-membered heterocycloalkyl, or  $n$  is 2 and one  $R^7$  represents (1)  $-NH_2$ , (2)  $-NHC(O)O-C1-4$  alkyl or (3)  $-NHC(O)O-C1-4$  alkylene  $-O-C1-$
- 20  $-4$  alkyl, (4)  $-COOH$ , (5)  $-COO-C1-8$  alkyl, (6)  $-COO-C1-8$  alkyl substituted with 1 to 5 groups selected from  $-N(C1-4$  alkyl) $_2$ , oxo and 5- to 10-membered heterocycloalkyl, (7)  $-COO-C5-C10$  aryl or (8)  $-COO-C1-4$  alkylene  $-5-$  to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen and the



other  $R^7$  represents halogen.


In a preferred embodiment,  $-\text{Cyc}^2-(R^6)_m$  represents



5 wherein  $R^{60}$  represents hydrogen, methyl or halogen;

$R^{61}$  represents (1) hydrogen, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) -COOH, (6) -COO-C1-4 alkyl, (7) cyano or (8) -CONH<sub>2</sub>; and the arrow represents a binding position.

10 In one embodiment, preferred compounds of the present invention are pyridinone or pyrimidinone derivatives represented by formula (I-A):



wherein U represents S or CH<sub>2</sub>; and

the other symbols have the same meanings as described above. Thus, preferred Cyc<sup>1</sup>, Cyc<sup>2</sup>, Cyc<sup>3</sup>, R<sup>1</sup>, s, R<sup>2</sup>, R<sup>5</sup>, R<sup>6</sup>, m, R<sup>7</sup> and n in the formula (I-A) are the preferred options as described above.

5 Preferred compounds of formula (I-A) are those in which:

Cyc<sup>1</sup> represents a C5-C10 aryl or 5 to 10 membered heteroaryl;

Cyc<sup>2</sup> represents a C5-C10 aryl, preferably a C5-C6 aryl;

10 Cyc<sup>3</sup> represents cyclohexane, indoline, isoindoline, dihydroquinoline, dihydroquinazoline, dihydrobenzoxazine, dihydrobenzoxazole, phenyl, pyrazole, pyridine, pyrazine, thiophene, oxazole, thiazole, indazole, quinoline, isoquinoline, quinoxaline or benzimidazole;

R<sup>1</sup> represents C1-8 alkyl or halogen;

s represents an integer of 0 or 1;

15 R<sup>2</sup> represents (1) hydrogen, (2) C1-8 alkyl, (3) Cyc<sup>4</sup> or (4) C1-8 alkyl substituted with 1 to 3 groups selected from Cyc<sup>5</sup> and -OR<sup>15</sup>;

Cyc<sup>4</sup> represents C5-C10 aryl;

Cyc<sup>5</sup> represents C5-C10 aryl, for example C5-C7 aryl, or a 5- to 6-membered heteroaryl;

R<sup>15</sup> is selected from hydrogen and C1-4 alkyl;

20 Y represents N or =CH-;

U represents S or CH<sub>2</sub>;

R<sup>6</sup> represents (1) methyl, (2) Cyc<sup>10</sup>, (3) halogen, (4) amidino, or (5) Cyc<sup>10</sup> substituted with 1 to 3 groups selected from halogen, trifluoromethyl, cyano, -OR<sup>29</sup>, -COOR<sup>35</sup>, -NHC(O)-R<sup>36</sup> and -C(O)NH-R<sup>38</sup>;

25 R<sup>29</sup>, R<sup>35</sup>, R<sup>36</sup> and R<sup>38</sup> each independently represent hydrogen or C1-4 alkyl;

Cyc<sup>10</sup> represents imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl, benzimidazolyl, thiazolyl or thiadiazolyl;

m represents an integer of 0, 1 or 2,

wherein m represents an integer of 2, each R<sup>6</sup> may be same or different;

30 R<sup>7</sup> represents (1) C1-8 alkyl, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) cyano, (6) oxo, (7) -OR<sup>40</sup>, (8) -NR<sup>42</sup>R<sup>43</sup>, (9) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (10) -NHC(O)-C1-4 alkylene - NR<sup>46</sup>R<sup>47</sup>, (11) -NHC(O)-C1-4 alkylene -COOH, (12) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (13) -COOR<sup>48</sup>, (14) -NHC(O)-R<sup>49</sup>, (15) -NHC(O)-C1-4 alkylene -OR<sup>50</sup>, (16) -NHC(O)O-R<sup>51</sup>,

- (17) -NHC(O)O-C1-4 alkylene -OR<sup>52</sup>, (18) -C(O)NH-R<sup>53</sup>, (19) -OC(O)-R<sup>55</sup>, (20) -C(O)-R<sup>56</sup>, (21) -CH(OH)-R<sup>57</sup>, (22) -C1-4 alkylene -NH<sub>2</sub>, (23) -C1-4 alkylene -OH, (24) -C1-4 alkylene -OC(O)-C1-4 alkyl, (25) -C1-4 alkylene -NHC(O)-C1-4 alkyl, (26) -C1-4 alkylene -NHC(O)O-C1-4 alkyl, (27) -C1-4 alkylene -NHC(O)-CF<sub>3</sub>, (28) -C1-4 alkylene -NHC(O)NH-C1-4 alkyl, (29) -CH=N-OR<sup>58</sup> or (30) -T-COOR<sup>66</sup>;
- 5 R<sup>40</sup>, R<sup>41</sup>, R<sup>42</sup>, R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup>, R<sup>47</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>51</sup>, R<sup>52</sup>, R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup>, R<sup>57</sup> and R<sup>58</sup> each independently represent hydrogen, C1-4 alkyl or trifluoromethyl;
- R<sup>51</sup> represents C1-8 alkyl;
- T represents C1-4 alkylene or C2-4 alkenylene;
- 10 R<sup>48</sup> and R<sup>66</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl which are substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl, (4) C3-C10 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl, (8) -C1-4 alkylene -C3-C8
- 15 cycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9) -C1-4 alkylene -C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen, (10) -C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (11) -C1-4 alkylene -5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from
- 20 C1-4 alkyl, OH and halogen or (12) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 3 groups selected from OH, oxo, -O-C1-4 alkyl, halogen, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, -O-C3-C8 cycloalkyl and -O-5- to 10-membered heterocycloalkyl;
- n represents an integer of 0, 1, 2 or 3,
- 25 wherein n represents an integer of 2 to 3, each R<sup>7</sup> may be same or different; and R<sup>62</sup> represents hydrogen or chlorine.

Preferred compounds of formula (I-A) include those in which:

- Cyc<sup>1</sup> represents phenyl, imidazolyl, triazolyl, tetrazolyl, pyrrolyl, pyrazolyl,
- 30 furanyl, oxazolyl, thiazolyl, isothiazolyl, furazanyl, oxadiazolyl, thiadiazolyl, thienyl, pyridazinyl, indazolyl or benzimidazolyl;
- Cyc<sup>2</sup> represents pyridyl or phenyl;

Cyc<sup>3</sup> represents cyclohexane, indoline, isoindoline, dihydroquinoline, dihydroquinazoline, dihydrobenzoxazine, dihydrobenzoxazole, phenyl, pyrazole, pyridine, pyrazine, thiophene, oxazole, thiazole, indazole, quinoline, isoquinoline, quinoxaline or benzimidazole;

5 R<sup>1</sup> represents methyl, ethyl or chlorine;

s represents an integer of 0 or 1;

R<sup>2</sup> represents hydrogen;

Y represents CH;

U represents CH<sub>2</sub>;

10 R<sup>6</sup> represents (1) methyl, (2) Cyc<sup>10</sup>, (3) halogen (preferably chlorine), or (4) Cyc<sup>10</sup> substituted with 1 to 3 groups selected from halogen (preferably chlorine), nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl, cyano or -CONH<sub>2</sub>;

Cyc<sup>10</sup> represents pyrazolyl, thiadiazolyl, triazolyl or tetrazolyl;

m represents an integer of 0, 1 or 2,

15 wherein m represents an integer of 2, each R<sup>6</sup> may be same or different;

R<sup>7</sup> represents (1) methyl, (2) ethyl, (3) fluorine, (4) chlorine, (5) bromine, (6) nitro, (7) trifluoromethyl, (8) cyano, (9) oxo, (10) -OR<sup>40</sup>, (11) -NR<sup>42</sup>R<sup>43</sup>, (12) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (13) -NHC(O)-C1-4 alkylene -NR<sup>46</sup>R<sup>47</sup>, (14) -NHC(O)-C1-4 alkylene -COOH, (15) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (16) -COOR<sup>48</sup>, (17) -NHC(O)-R<sup>49</sup>, (18) -NHC(O)-C1-4 alkylene -OR<sup>50</sup>, (19) -NHC(O)O-R<sup>51</sup>, (20) -NHC(O)O-C1-4 alkylene -OR<sup>52</sup>, (21) -C(O)NH-R<sup>53</sup>, (22) -OC(O)-R<sup>55</sup>, (23) -C(O)-R<sup>56</sup>, (24) -CH(OH)-R<sup>57</sup>, (25) -C1-4 alkylene -NH<sub>2</sub>, (26) -C1-4 alkylene -OH, (27) -C1-4 alkylene -OC(O)-C1-4 alkyl, (28) -C1-4 alkylene -NHC(O)-C1-4 alkyl, (29) -C1-4 alkylene -NHC(O)O-C1-4 alkyl, (30) -C1-4 alkylene -NHC(O)-CF<sub>3</sub>, (31) -C1-4 alkylene -NHC(O)NH-C1-4 alkyl, (32) -CH=N-OR<sup>58</sup>, (33) -C1-4 alkylene-COOH, (34) -C1-4 alkylene-COO-C1-8 alkyl or (35) -C1-4 alkylene-COO-C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl;

R<sup>40</sup>, R<sup>41</sup>, R<sup>42</sup>, R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup>, R<sup>47</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>52</sup>, R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup>, R<sup>57</sup> and R<sup>58</sup> are independently selected from hydrogen, C1-4 alkyl and trifluoromethyl;

30 R<sup>48</sup> represents (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and morpholine, (4) phenyl, (5) indane, (6) C1-4 alkylene - 1,3-dioxole substituted with 1 to 5 groups selected from C1-4 alkyl, oxo,


OH and halogen or (7) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 3 groups selected from oxo and -O-cyclohexane;

$R^{51}$  represents C1-4 alkyl;


n represents an integer of 0, 1, 2 or 3,

5 wherein n represents an integer of 2 to 3, each  $R^7$  may be same or different; and  $R^{62}$  represents hydrogen.

Further preferred compounds of formula (I-A) include those in which



10 represents




wherein  $R^{59}$  represents hydrogen, C1-4 alkyl or halogen;

the arrow represents a binding position; and

15 the other symbols have the same meanings as described above, preferably wherein n is 1 and  $R^7$  represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl or (3) -NHC(O)O-C1-4 alkylene -O-C1-4 alkyl, (4) -COOH, (5) -COO-C1-8 alkyl, (6) -COO-C1-8 alkyl

substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, (7) -COO- C5-C10 aryl, (8) -COO- C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9) -COO- indane, (10) -COO- C1-4 alkylene - 1,3-dioxole substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (11) -COO-C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 3 groups selected from oxo and -O-cyclohexane, (12) -C1-4 alkylene-COOH, (13) -C1-4 alkylene-COO- C1-8 alkyl or (14) -C1-4 alkylene-COO- C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, or n is 2 and one R<sup>7</sup> represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl or (3) -NHC(O)O-C1-4 alkylene -O-C1-4 alkyl, (4) -COOH, (5) -COO-C1-8 alkyl, (6) -COO-C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, (7) -COO- C5-C10 aryl or (8) -COO- C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen and the other R<sup>7</sup> represents halogen.


Further preferred compounds of formula (I-A) include those in which -Cyc<sup>2</sup>-(R<sup>6</sup>)<sub>m</sub> represents



wherein R<sup>60</sup> represents hydrogen, methyl or halogen;


R<sup>61</sup> represents (1) hydrogen, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) -COOH, (6) -COO-C1-4 alkyl, (7) cyano or (8) -CONH<sub>2</sub>; and the arrow represents a binding position.

Further preferred compounds of formula (I-A) include a compound of (I-A-1):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-1-

5 1):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-1-

10 2):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-2):



5


wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-2-1):



10




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-2-2):




5


wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-3):




10 wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-3-1):



wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-3-2):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-4):




wherein s represents an integer of 0 to 1; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-4-1):




wherein s represents an integer of 0 to 1; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-4-2):



wherein s represents an integer of 0 to 1; and


the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-5):



wherein m represents an integer of 0 to 5; and


the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-5-

5 1):




wherein m represents an integer of 0 to 5; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-5-  
10 2):



wherein m represents an integer of 0 to 5; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-6):




5

wherein s represents an integer 0 to 1;

m represents an integer 0 to 5; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-6-1):


10



wherein  $s$  represents an integer 0 to 1;

$m$  represents an integer 0 to 5; and

- 5 the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-A-6-2):




wherein  $s$  represents an integer 0 to 1;

- 10  $m$  represents an integer 0 to 5; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), and the like.

- In another embodiment, preferred compounds of the present invention are  
 15 pyridinone derivatives represented by formula (I-B):



wherein  $\text{Cyc}^{1-B}$  represents 5- to 10-membered heteroaryl;

$\text{Cyc}^{3-B}$  represents C5-C10 aryl or 5- to 10-membered heteroaryl;

$\text{Cyc}^{10-B}$  represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl,  $-\text{COOH}$ ,  $-\text{COO}-\text{C1-4 alkyl}$  and  $-\text{CONH}_2$ ;

mb represents an integer of 0 to 5; and

the other symbols have the same meanings as described above. Thus, preferred  $\text{Cyc}^{1-B}$ ,  $\text{Cyc}^2$ ,  $\text{Cyc}^{3-B}$ ,  $\text{R}^1$ , s,  $\text{R}^2$ ,  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{Cyc}^{10-B}$ ,  $\text{R}^7$  and n in the formula (I-B) are the preferred options as described above.

Preferred compounds of formula (I-B), are those in which:

$\text{Cyc}^{1-B}$  represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl;

$\text{Cyc}^2$  represents a C5-C10 aryl, preferably a C5-C6 aryl;

$\text{Cyc}^{3-B}$  represents C5-C10 aryl, for example C5-C7 aryl such as phenyl, or 5- to 10-membered heteroaryl, such as pyrazole, pyridine, pyrazine, thiophene, oxazole, thiazole, indazole, quinoline, isoquinoline, quinoxaline or benzimidazole, more preferably pyrazole or pyridine;

$\text{R}^1$  represents halogen or C1-4 alkyl;

s represents an integer of 0 or 1;

$\text{R}^2$  represents (1) hydrogen, (2) C1-8 alkyl, (3)  $\text{Cyc}^4$  or (4) C1-8 alkyl substituted with 1 to 3 groups selected from  $\text{Cyc}^5$  and  $-\text{OR}^{15}$ ;

$\text{Cyc}^4$  represents C5-C10 aryl;

$\text{Cyc}^5$  represents C5-C10 aryl, for example C5-C7 aryl or a 5- to 6-membered heteroaryl;

$\text{R}^{15}$  is selected from hydrogen and C1-4 alkyl;

$R^4$  represents hydrogen or C1-4 alkyl;

$R^5$  represents hydrogen;

$R^6$  represents methyl, halogen or amidino;

mb represents an integer of 0 or 1;

5     Cyc<sup>10-B</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl, benzimidazolyl, thiazolyl or thiadiazolyl, any of which optionally substituted with 1 to 3 groups selected from halogen, cyano, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl and -CONH<sub>2</sub>;

10      $R^7$  represents (1) C1-8 alkyl, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) cyano, (6) oxo, (7) -OR<sup>40</sup>, (8) -NR<sup>42</sup>R<sup>43</sup>, (9) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (10) -NHC(O)-C1-4 alkylene - NR<sup>46</sup>R<sup>47</sup>, (11) -NHC(O)-C1-4 alkylene -COOH, (12) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (13) -COOR<sup>48</sup>, (14) -NHC(O)-R<sup>49</sup>, (15) -NHC(O)-C1-4 alkylene -OR<sup>50</sup>, (16) -NHC(O)O-R<sup>51</sup>, (17) -NHC(O)O-C1-4 alkylene -OR<sup>52</sup>, (18) -C(O)NH-R<sup>53</sup>, (19) -OC(O)-R<sup>55</sup>, (20) -C(O)-R<sup>56</sup>, (21) -CH(OH)-R<sup>57</sup>, (22) -C1-4 alkylene -NH<sub>2</sub>, (23) -C1-4 alkylene -OH, (24) -C1-4 alkylene -OC(O)-C1-4 alkyl, (25) -C1-4 alkylene -NHC(O)-C1-4 alkyl, (26) -C1-4 alkylene -NHC(O)O-C1-4 alkyl, (27) -C1-4 alkylene -NHC(O)-CF<sub>3</sub>, (28) -C1-4 alkylene -NHC(O)NH-C1-4 alkyl, (29) -CH=N-OR<sup>58</sup> or (30) -T-COOR<sup>66</sup>;

15      $R^{40}$ ,  $R^{41}$ ,  $R^{42}$ ,  $R^{43}$ ,  $R^{44}$ ,  $R^{45}$ ,  $R^{46}$ ,  $R^{47}$ ,  $R^{49}$ ,  $R^{50}$ ,  $R^{52}$ ,  $R^{53}$ ,  $R^{54}$ ,  $R^{55}$ ,  $R^{56}$ ,  $R^{57}$  and  $R^{58}$  each independently represent hydrogen, C1-4 alkyl or trifluoromethyl;

20     T represents C1-4 alkylene or C2-4 alkenylene;

25      $R^{48}$  and  $R^{66}$  each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl which are substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl, (4) C3-C10 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl, (8) -C1-4 alkylene -C3-C8 cycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9) -C1-4 alkylene -C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen, (10) -C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen or (11) -C1-4 alkylene -5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen;

30      $R^{51}$  represents C1-8 alkyl; and

n represents an integer of 0, 1, 2 or 3,




wherein n represents an integer of 2 to 3, each R<sup>7</sup> may be same or different.

Preferred compounds of formula (I-B) include those in which:

- Cyc<sup>1-B</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl,  
 5 indazolyl or benzimidazolyl;  
 Cyc<sup>2</sup> represents pyridyl or phenyl;  
 Cyc<sup>3-B</sup> represents C5-C10 aryl, more preferably phenyl;  
 R<sup>1</sup> represents halogen or C1-4 alkyl;  
 s represents an integer of 0 or 1;  
 10 R<sup>2</sup> represents (1) hydrogen, (2) C1-4 alkyl, (3) Cyc<sup>4</sup> or (4) C1-4 alkyl substituted  
 with 1 or 2 groups selected from Cyc<sup>5</sup> and -OR<sup>15</sup>;  
 Cyc<sup>4</sup> represents phenyl;  
 Cyc<sup>5</sup> represents phenyl or pyridyl;  
 R<sup>15</sup> is selected from hydrogen and C1-4 alkyl;  
 15 R<sup>4</sup> represents hydrogen, methyl or ethyl;  
 R<sup>5</sup> represents hydrogen;  
 R<sup>6</sup> represents methyl or halogen (preferably chlorine);  
 mb represents an integer of 0 or 1;  
 Cyc<sup>10-B</sup> represents thiadiazolyl, triazolyl or tetrazolyl, any of which optionally  
 20 substituted with 1 to 3 groups selected from halogen (preferably chlorine), nitro,  
 trifluoromethyl, -COOH, -COO-C1-4 alkyl, cyano or -CONH<sub>2</sub>;  
 R<sup>7</sup> represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl, (3) -NHC(O)O-C1-4  
 alkylene -O-C1-4 alkyl, (4) halogen, (5) -COOH, (6) -COO-C1-8 alkyl, (7) -COO-C1-8  
 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-  
 25 membered heterocycloalkyl, (8) -COO- C5-C10 aryl, (9) -COO- C1-4 alkylene -5- to  
 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl,  
 oxo, OH and halogen, (10) -C1-4 alkylene-COOH or (11) -C1-4 alkylene-COO- C1-8  
 alkyl or (11) -C1-4 alkylene-COO- C1-8 alkyl substituted with 1 to 5 groups selected  
 from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl; and  
 30 n represents an integer of 0, 1, 2 or 3,  
 wherein n represents an integer of 2 to 3, each R<sup>7</sup> may be same or different.

Further preferred compounds of formula (I-B) include those in which  $\text{-Cyc}^{1-B}(\text{-R}^1)_s\text{-Cyc}^{3-B}(\text{-R}^7)_n$  represents



wherein  $\text{R}^{59}$  represents hydrogen, C1-4 alkyl or halogen;

5 the arrow represents a binding position; and


the other symbols have the same meanings as described above, preferably

wherein  $n$  is 1 and  $\text{R}^7$  represents (1)  $\text{-NH}_2$ , (2)  $\text{-NHC(O)O-C1-4 alkyl}$  or (3)  $\text{-NHC(O)O-C1-4 alkylene-O-C1-4 alkyl}$ , (4)  $\text{-COOH}$ , (5)  $\text{-COO-C1-8 alkyl}$ , (6)  $\text{-COO-C1-8 alkyl}$  substituted with 1 to 5 groups selected from  $\text{-N(C1-4 alkyl)}_2$ , oxo and 5- to 10-

10 membered heterocycloalkyl, (7)  $\text{-COO-C5-C10 aryl}$ , (8)  $\text{-COO-C1-4 alkylene-5- to 10-membered heterocycloalkyl}$  substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9)  $\text{-C1-4 alkylene-COOH}$ , (10)  $\text{-C1-4 alkylene-COO-C1-8 alkyl}$  or (11)  $\text{-C1-4 alkylene-COO-C1-8 alkyl}$  substituted with 1 to 5 groups selected from  $\text{-N(C1-4 alkyl)}_2$ , oxo and 5- to 10-membered heterocycloalkyl, or  $n$  is 2 and one  $\text{R}^7$

15 represents (1)  $\text{-NH}_2$ , (2)  $\text{-NHC(O)O-C1-4 alkyl}$ , (3)  $\text{-NHC(O)O-C1-4 alkylene-O-C1-4 alkyl}$ , (4)  $\text{-COOH}$ , (5)  $\text{-COO-C1-8 alkyl}$ , (6)  $\text{-COO-C1-8 alkyl}$  substituted with 1 to 5 groups selected from  $\text{-N(C1-4 alkyl)}_2$ , oxo and 5- to 10-membered heterocycloalkyl, (7)  $\text{-COO-C5-C10 aryl}$  or (8)  $\text{-COO-C1-4 alkylene-5- to 10-membered heterocycloalkyl}$  substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen and the  
20 other  $\text{R}^7$  represents halogen.

Further preferred compounds of formula (I-B) include those in which  $\text{-Cyc}^2(\text{-R}^6)_{mb}\text{Cyc}^{10-B}$  represents




wherein  $R^{60}$  represents hydrogen, methyl or halogen;

$R^{61}$  represents (1) hydrogen, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) -COOH, (6) -COO-C1-4 alkyl, (7) cyano or (8) -CONH<sub>2</sub>; and the arrow represents a binding


5 position.

Further preferred compounds of formula (I-B) include a compound of (I-B-1):




wherein all symbols have the same meanings as described above and the same


10 preferred definitions as set out above (alone or in combination), a compound of (I-B-2):



wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-B-3):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-B-4):




wherein  $s$  represents an integer of 0 to 1; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-B-5):



wherein  $mb$  represents an integer of 0 to 4; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-B-6):




wherein  $s$  represents an integer 0 to 1;

5         $mb$  represents an integer 0 to 4; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), and the like.

10        In another embodiment, preferred compounds of the present invention are pyrimidinone derivatives represented by formula (I-C):



wherein  $Cyc^{1-C}$  represents 5- to 10-membered heteroaryl;

$Cyc^{3-C}$  represents C5-C10 aryl or 5- to 10-membered heteroaryl;

15         $Cyc^{10-C}$  represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl,  $-COOH$ ,  $-COO-C1-4$  alkyl and  $-CONH_2$ ;

$mc$  represents an integer of 0 to 5; and

20        the other symbols have the same meanings as described above. Thus, preferred  $Cyc^{1-C}$ ,  $Cyc^2$ ,  $Cyc^{3-C}$ ,  $R^1$ ,  $s$ ,  $R^2$ ,  $R^4$ ,  $R^6$ ,  $Cyc^{10-C}$ ,  $R^7$  and  $n$  in the formula (I-C) are the preferred options as described above.

Preferred compounds of formula (I-C) are those in which:

Cyc<sup>1-C</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl;

Cyc<sup>2</sup> represents a C5-C10 aryl, preferably a C5-C6 aryl such as phenyl;

5 Cyc<sup>3-C</sup> represents C5-C10 aryl, for example C5-C7 aryl such as phenyl, or 5- to 10-membered heteroaryl, such as pyrazole, pyridine, pyrazine, thiophene, oxazole, thiazole, indazole, quinoline, isoquinoline, quinoxaline or benzimidazole, more preferably pyrazole or pyridine;

R<sup>1</sup> represents halogen or C1-4 alkyl;

10 s represents an integer of 0 or 1;

R<sup>2</sup> represents hydrogen, (1) C1-8 alkyl, (2) Cyc<sup>4</sup> or (3) C1-8 alkyl substituted with 1 to 3 groups selected from Cyc<sup>5</sup> and -OR<sup>15</sup>;

Cyc<sup>4</sup> represents C5-C10 aryl;

15 Cyc<sup>5</sup> represents C5-C10 aryl, for example C5-C7 aryl, or a 5- to 6-membered heteroaryl;

R<sup>15</sup> is selected from hydrogen and C1-4 alkyl;

R<sup>4</sup> represents hydrogen or C1-4 alkyl;

R<sup>6</sup> represents methyl, halogen or amidino;

mc represents an integer of 0 or 1;

20 Cyc<sup>10-C</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl, benzimidazolyl, thiazolyl or thiadiazolyl, any of which optionally substituted with 1 to 3 groups selected from halogen, cyano, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl and -CONH<sub>2</sub>;

R<sup>7</sup> represents (1) C1-8 alkyl, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) cyano, 25 (6) oxo, (7) -OR<sup>40</sup>, (8) -NR<sup>42</sup>R<sup>43</sup>, (9) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (10) -NHC(O)-C1-4 alkylene - NR<sup>46</sup>R<sup>47</sup>, (11) -NHC(O)-C1-4 alkylene -COOH, (12) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (13) -COOR<sup>48</sup>, (14) -NHC(O)-R<sup>49</sup>, (15) -NHC(O)-C1-4 alkylene -OR<sup>50</sup>, (16) -NHC(O)O-R<sup>51</sup>, (17) -NHC(O)O-C1-4 alkylene -OR<sup>52</sup>, (18) -C(O)NH-R<sup>53</sup>, (19) -OC(O)-R<sup>55</sup>, (20) -C(O)-R<sup>56</sup>, (21) -CH(OH)-R<sup>57</sup>, (22) -C1-4 alkylene -NH<sub>2</sub>, (23) -C1-4 alkylene -OH, (24) -C1-4 30 alkylene -OC(O)-C1-4 alkyl, (25) -C1-4 alkylene -NHC(O)-C1-4 alkyl, (26) -C1-4 alkylene -NHC(O)O-C1-4 alkyl, (27) -C1-4 alkylene -NHC(O)-CF<sub>3</sub>, (28) -C1-4 alkylene -NHC(O)NH-C1-4 alkyl, (29) -CH=N-OR<sup>58</sup> or (30) -T-COOR<sup>66</sup>;

$R^{40}, R^{41}, R^{42}, R^{43}, R^{44}, R^{45}, R^{46}, R^{47}, R^{48}, R^{49}, R^{50}, R^{52}, R^{53}, R^{54}, R^{55}, R^{56}, R^{57}$  and  $R^{58}$  each independently represent hydrogen, C1-4 alkyl or trifluoromethyl;

$R^{51}$  represents C1-8 alkyl;

T represents C1-4 alkylene or C2-4 alkenylene;

5  $R^{48}$  and  $R^{66}$  each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C1-8 alkyl which are substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl, (4) C3-C10 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl, (8) -C1-4 alkylene -C3-C8  
10 cycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (9) -C1-4 alkylene -C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH and halogen, (10) -C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen or (11) -C1-4 alkylene -5- to 10-membered heteroaryl substituted with 1 to 5 groups selected  
15 from C1-4 alkyl, OH and halogen; and

n represents an integer of 0, 1, 2 or 3,

wherein n represents an integer of 2 to 3, each  $R^7$  may be same or different.

Preferred compounds of formula (I-C) include those in which:

20  $Cyc^{1-C}$  represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl;

$Cyc^2$  represents pyridyl or phenyl;

$Cyc^{3-C}$  represents C5-C10 aryl, more preferably phenyl;

each  $R^1$  independently represents halogen or C1-4 alkyl;

25 s represents an integer of 0 or 1;

$R^2$  represents (1) hydrogen, (2) C1-4 alkyl, (3)  $Cyc^4$  or (4) C1-4 alkyl substituted with 1 or 2 groups selected from  $Cyc^5$  and -OR<sup>15</sup>;

$Cyc^4$  represents phenyl;

$Cyc^5$  represents phenyl or pyridyl;

30  $R^{15}$  is selected from hydrogen and C1-4 alkyl;

$R^4$  represents hydrogen, methyl or ethyl;

each  $R^6$  independently represents methyl or halogen;


Cyc<sup>10-C</sup> represents thiadiazolyl, triazolyl or tetrazolyl, any of which optionally substituted with 1 to 3 groups selected from halogen (preferably chlorine), nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl, cyano or -CONH<sub>2</sub>;

R<sup>7</sup> represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl, (3) -NHC(O)O-C1-4 alkylene -O-C1-4 alkyl, (4) halogen, (5) -COOH, (6) -COO-C1-8 alkyl, (7) -COO-C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, (8) -COO- C5-C10 aryl, (9) -COO- C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen, (10) -C1-4 alkylene-COOH, (11) -C1-4 alkylene-COO- C1-8 alkyl or (12) -C1-4 alkylene-COO- C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl; and

n represents an integer of 0, 1, 2 or 3,

wherein n represents an integer of 2 to 3, each R<sup>7</sup> may be same or different.

Further preferred compounds of formula (I-C) include those in which -Cyc<sup>1-C</sup>(-R<sup>1</sup>)<sub>s</sub>-Cyc<sup>3-C</sup>(-R<sup>7</sup>)<sub>n</sub> represents



wherein R<sup>59</sup> represents hydrogen, C1-4 alkyl or halogen;

the arrow represents a binding position; and


the other symbols have the same meanings as described above, preferably wherein n is 1 and R<sup>7</sup> represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl, (3) -NHC(O)O-C1-4 alkylene -O-C1-4 alkyl, (4) -COOH, (5) -COO-C1-8 alkyl, (6) -COO-C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, (7) -COO- C5-C10 aryl, (8) -COO- C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl,



oxo, OH and halogen, (9) -C1-4 alkylene-COOH, (10) -C1-4 alkylene-COO- C1-8 alkyl or (11) -C1-4 alkylene-COO- C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, or n is 2 and one R<sup>7</sup> represents (1) -NH<sub>2</sub>, (2) -NHC(O)O-C1-4 alkyl, (3) -NHC(O)O-C1-4 alkylene -O-C1-4 alkyl, (4) -COOH, (5) -COO-C1-8 alkyl, (6) -COO-C1-8 alkyl substituted with 1 to 5 groups selected from -N(C1-4 alkyl)<sub>2</sub>, oxo and 5- to 10-membered heterocycloalkyl, (7) -COO- C5-C10 aryl or (8) -COO- C1-4 alkylene -5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH and halogen and the other R<sup>7</sup> represents halogen.

10


Further preferred compounds of formula (I-C) include those in which -Cyc<sup>2</sup>(-R<sup>6</sup>)<sub>mcCyc</sub><sup>10-C</sup> represents




wherein R<sup>60</sup> represents hydrogen, methyl or halogen;

15 R<sup>61</sup> represents (1) hydrogen, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) -COOH, (6) -COO-C1-4 alkyl, (7) cyano or (8) -CONH<sub>2</sub>; and the arrow represents a binding position.


Further preferred compounds of formula (I-C) include a compound of (I-C-1):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-C-2):



5 wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-C-3):




wherein all symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-C-4):




wherein  $s$  represents an integer of 0 to 1; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-C-5):



wherein  $mc$  represents an integer of 0 to 4; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), a compound of (I-C-6):



wherein  $s$  represents an integer 0 to 1;

$mc$  represents an integer 0 to 4; and

the other symbols have the same meanings as described above and the same preferred definitions as set out above (alone or in combination), and the like.

As used herein, general references to “compounds of formula (I)” include compounds of formula (I-A), (I-B) and (I-C).

Particularly preferred compounds of formula (I) include:

- 5        4-(4-chloro-2-((1S)-1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl)-1H-imidazol-5-yl)benzoic acid,  
       4-(2-((1S)-1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl)-1H-imidazol-5-yl)benzoic acid,  
       3-(4-chloro-2-((1S)-1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl)-1H-imidazol-5-yl)benzoic acid,  
 10       3-{2-[(S)-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl](phenyl)methyl]-1H-imidazol-5-yl}benzoic acid,  
       salts thereof, N-oxides thereof, solvates thereof, and prodrugs thereof.

- 15        Particularly preferred compounds of formula (I-A) include:  
       methyl [4-(2-{2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate,  
       methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-4-yl)phenyl]carbamate,  
 20       methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]carbamate,  
       3-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzoic acid,  
       (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-phenyl-1H-imidazol-2-yl)-  
 25       2,3-dihydro-5(1H)-indolizinone,  
       methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,  
       methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-ethyl-1H-imidazol-5-yl)phenyl]carbamate,  
 30       methyl (4-{2-[7-(5-chloro-2-formamidophenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl]-4-ethyl-1H-imidazol-5-yl}phenyl)carbamate,  
       5-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-4-yl)-1,3-dihydro-2H-indol-2-one,

- 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-4-yl)-3,4-dihydro-2(1H)-quinolinone,
- 6-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone,
- 5 methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-2-fluorophenyl]carbamate,
- methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-2-fluorophenyl]carbamate,
- methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-fluorophenyl]carbamate,
- 10 methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-fluorophenyl]carbamate,
- methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-methylphenyl]carbamate,
- 15 methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-methylphenyl]carbamate,
- methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-ethylphenyl]carbamate,
- 20 methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-ethylphenyl]carbamate,
- methyl [3-chloro-4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]carbamate,
- methyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-methoxyphenyl]carbamate,
- 25 methyl [2-bromo-4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-5-methoxyphenyl]carbamate,
- methyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-(trifluoromethyl)phenyl]carbamate,
- methyl [4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-(trifluoromethyl)phenyl]carbamate,
- 30 1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-(trifluoromethyl)phenyl]carbamate,
- 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-4-yl)benzoic acid,

- 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)benzamide,
- 4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)benzamide,
- 5 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-2-fluorobenzamide,
- 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)-N-methylbenzamide,
- 4-(5-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)-N-methylbenzamide,
- 10 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)-N-ethylbenzamide,
- 4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-N-ethylbenzamide,
- 15 7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-nitrophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 3-[5-(4-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- ethyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]carbamate,
- 20 isopropyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]carbamate,
- 2-methoxyethyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]carbamate,
- 25 N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]acetamide,
- N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]-2-methoxyacetamide,
- N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]-3-methoxypropanamide,
- 30 N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]methanesulfonamide,

- ethyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- isopropyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 5 N-[4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide,
- N-[4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]methanesulfonamide,
- 1-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-3-ethylurea,
- 10 3-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-1,1-dimethylurea,
- N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-2-(dimethylamino)acetamide,
- 15 3-{[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-3-oxopropanoic acid,
- 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-4-hydroxy-2(1H)-quinolinone,
- 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-fluorobenzonitrile,
- 20 3-[5-(3-amino-1H-indazol-6-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 7-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2H-1,4-benzoxazin-3(4H)-one,
- 25 7-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-4-methyl-2H-1,4-benzoxazin-3(4H)-one,
- 7-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-4-methyl-2H-1,4-benzoxazin-3(4H)-one,
- 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1,4-dihydro-2H-3,1-benzoxazin-2-one,
- 30 6-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1,4-dihydro-2H-3,1-benzoxazin-2-one,

- 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-1,3-benzoxazol-2(3H)-one,
- 6-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-1,3-benzoxazol-2(3H)-one,
- 5 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-1-isoindolinone,
- 6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3-methyl-3,4-dihydro-2(1H)-quinazolinone,
- 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-2(1H)-quinoxalinone,
- 10 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1-methyl-1H-benzimidazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 3-[4-chloro-5-(1-methyl-1H-benzimidazol-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 15 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1H-indazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)benzyl acetate,
- 20 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)benzaldehyde oxime,
- 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)benzaldehyde O-methyloxime,
- 25 (3S)-3-[5-(4-acetylphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(1-hydroxyethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- 2-methyl-2-propanyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)benzyl]carbamate,
- 30 (3S)-3-{5-[4-(aminomethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,



- methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]carbamate,
- N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]acetamide,
- 5      N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]-2,2,2-trifluoroacetamide,
- 1-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]-3-ethylurea,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 10      (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(2-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 15      (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(1,3-thiazol-2-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(2-pyrazinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(2-methoxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 20      (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(3-methoxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(4-methoxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 25      (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(3-fluorophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(4-fluorophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 30      (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(2,4-dimethyl-1,3-oxazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(1-methyl-1H-pyrazol-4-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,

- methyl [5-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thienyl]carbamate,
- 2-methyl-2-propanyl [trans-4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)cyclohexyl]carbamate,
- 5 (3S)-3-[5-(trans-4-aminocyclohexyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl [trans-4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)cyclohexyl]carbamate,
- (3S)-3-[5-(trans-4-aminocyclohexyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 10 (3S)-3-[5-(2-amino-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 3-[4-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 15 methyl [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- methyl [5-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- ethyl [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 20 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-3-pyridinyl)-4-methyl-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 25 methyl [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-4-methyl-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-2-methyl-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 30 methyl [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-6-methyl-2-pyridinyl]carbamate,

- (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(6-methyl-3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-3-[4-chloro-5-(6-methyl-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 2-methyl-2-propanyl [6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 3-[5-(5-amino-2-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 10 (3S)-3-[5-(5-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl [6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 15 methyl [6-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- isopropyl [6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- isobutyl [6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 20 N-[6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]acetamide,
- (3S)-3-[5-(5-amino-2-pyridinyl)-4-methyl-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 25 N-[6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-methyl-1H-imidazol-5-yl)-3-pyridinyl]acetamide,
- methyl [6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-methyl-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- (3S)-3-[5-(2-amino-1,3-thiazol-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 30 (3S)-3-[5-(1-amino-6-isoquinolinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- methyl [4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-2-yl)phenyl]carbamate,
- methyl [4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-2-yl)phenyl]carbamate,
- 5 3-[5-(4-aminophenyl)-4H-1,2,4-triazol-3-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl [4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-4H-1,2,4-triazol-3-yl)phenyl]carbamate,
- methyl [4-(6-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-3-oxo-2,3-dihydro-4-pyridaziny)phenyl]carbamate,
- 10 methyl [4-(4-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-1,2,3-triazol-1-yl)phenyl]carbamate,
- methyl [4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1,2-oxazol-3-yl)phenyl]carbamate,
- 15 methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-4-yl)phenyl]carbamate,
- methyl [4-(2-((3R)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-2,3-dihydro-5H-[1,3]thiazolo[3,2-a]pyridin-3-yl)-1H-imidazol-5-yl)phenyl]carbamate,
- methyl [3-chloro-4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-4-yl)phenyl]carbamate,
- 20 6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone,
- ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]carbamate,
- 25 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzamide,
- 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzamide,
- methyl [4-(2-((3R)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]carbamate,
- 30 methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-5-methyl-1H-imidazol-4-yl)phenyl]carbamate,

- methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-5-methyl-1,3-oxazol-4-yl]phenyl]carbamate,  
2-methyl-2-propanyl 1-{4-chloro-2-[(3S)-3-(5-{4-  
[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-  
5 indoliziny]phenyl}-1H-1,2,3-triazole-4-carboxylate,  
methyl 1-{4-chloro-2-[(3S)-3-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl}-1H-1,2,3-triazole-4-carboxylate,  
1-{4-chloro-2-[(3S)-3-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl}-1H-1,2,3-triazole-4-carboxylic acid,  
10 methyl [4-(2-((3S)-7-[2-(4-carbamoyl-1H-1,2,3-triazol-1-yl)-5-chlorophenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
methyl [4-(2-((3S)-7-[5-chloro-2-(4-cyano-1H-1,2,3-triazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
15 methyl [4-(2-((3S)-7-[5-chloro-2-(3-hydroxy-1H-1,2,4-triazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
methyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)-3-pyridiny]l-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
methyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
20 methyl [4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
methyl [4-(2-((3S)-7-[5-fluoro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
25 methyl [4-(2-((3S)-5-oxo-7-[2-(1H-tetrazol-1-yl)-5-(trifluoromethyl)]phenyl]-1,2,3,5-tetrahydro-3-indoliziny]l-1H-imidazol-5-yl)phenyl]carbamate,  
(6S)-6-[5-(6-amino-2-fluoro-3-pyridiny]l-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
(6S)-6-[5-(6-amino-3-pyridiny]l-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
30 (6S)-6-[5-(6-amino-2-chloro-3-pyridiny]l-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzoic acid,

3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzoic acid,

methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]acetate,

[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]acetic acid,

[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]acetic acid,

[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]acetic acid,

[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]acetic acid,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxamide,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-methyl-2-thiophenecarboxamide,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-ethyl-2-thiophenecarboxamide,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(2-methoxyethyl)-2-thiophenecarboxamide,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-[2-(2-methoxyethoxy)ethyl]-2-thiophenecarboxamide,

- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-[2-(4-morpholinyl)ethyl]-2-thiophenecarboxamide,  
5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxamide,  
5 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-methyl-2-thiophenecarboxamide,  
5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-ethyl-2-thiophenecarboxamide,  
5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide,  
10 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(2-methoxyethyl)-2-thiophenecarboxamide,  
5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-[2-(2-methoxyethoxy)ethyl]-2-thiophenecarboxamide,  
5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-[2-(4-morpholinyl)ethyl]-2-thiophenecarboxamide,  
20 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-methoxy-2-thiophenecarboxamide,  
5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-cyano-2-thiophenecarboxamide,  
2-(4-morpholinyl)ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
25 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-methoxy-2-thiophenecarboxamide,  
5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-cyano-2-thiophenecarboxamide,  
30 2-(4-morpholinyl)ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

- methyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetate,  
[5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetic acid,
- 5 methyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetate,  
[5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetic acid,
- 10 2-methyl-2-propanyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,  
4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 15 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,  
ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
isopropyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 20 3-methylbutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
2-(4-morpholinyl)ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 25 (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
phenyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 30 2,3-dihydro-1H-inden-5-yl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
isobutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,



2-(dimethylamino)-2-oxoethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide,

methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[5-(hydroxymethyl)-3-thienyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,

[(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

5-(2-{7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(4-chloro-2-{7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indole-3-carboxylic acid,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indazole-3-carboxylic acid,

4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,

4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,

5-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(4-chloro-2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(4-chloro-2-((3S)-6-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(2-((3S)-6,8-dichloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

- 4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 5 4-(2-((3S)-8-bromo-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (2E)-3-[3-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl}phenyl]acrylic acid,
- (2E)-3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acrylic acid,
- 10 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 15 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1-oxido-4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(3-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 2-methoxyethyl [3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 20 methyl [3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- N-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide,
- 25 methyl 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoate,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[3-(methylsulfonyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[3-(methoxymethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- 30 (3S)-3-[5-(3-acetylphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzaldehyde O-methyloxime,
- 3-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzamide,
- 5 3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzonitrile,
- (3S)-3-[4-chloro-5-(3-hydroxyphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(2-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 10 methyl [2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- N-[2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide,
- 15 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoic acid,
- [3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]boronic acid,
- [3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]boronic acid,
- 20 ethyl [3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetate,
- ethyl [3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetate,
- 25 2-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide,
- 2-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide,
- 2-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-N-methylacetamide,
- 30 2-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-N-methylacetamide,

- 2-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]-2-methylpropanoic acid,
- 2-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]-2-methylpropanoic acid,
- 5 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-methoxybenzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-ethoxybenzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-methoxyethyl)benzamide,
- 10 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(3-methoxypropyl)benzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-methoxyethoxy)benzamide,
- 15 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-fluoroethyl)benzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2,2-difluoroethyl)benzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(3-fluoropropyl)benzamide,
- 20 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-methoxyethyl)benzamide,
- 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-methoxyethoxy)benzamide,
- 25 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-methyl-1H-imidazol-5-yl)benzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-methylbenzamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-fluorobenzamide,
- 30 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxamide,

- 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)benzenecarbothioamide,  
(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
5 (3S)-3-{4-chloro-5-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-{4-chloro-5-[4-(1-hydroxyethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-{5-[2-chloro-4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-10 2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(hydroxymethyl)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-{4-chloro-5-[6-(hydroxymethyl)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
15 (3S)-3-{5-[2-chloro-6-(hydroxymethyl)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-4-methoxybutanamide,  
N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-20 3-indolizinyll}-1H-imidazol-5-yl)phenyl]acetamide,  
N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-2-fluoroacetamide,  
N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-2,2-difluoroacetamide,  
25 N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]tetrahydro-3-furancarboxamide,  
N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]tetrahydro-2H-pyran-4-carboxamide,  
N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-30 3-indolizinyll}-4-fluoro-1H-imidazol-5-yl)phenyl]acetamide,  
N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]ethanethioamide,

N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]formamide,

N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-N-hydroxyacetamide,

5 (1E)-N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-N'-hydroxyethanimidamide,

(1E)-N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-N'-methoxyethanimidamide,

10 N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-3-hydroxy-3-methylbutanamide,

3-{[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-3-oxopropanoic acid,

4-{[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-4-oxobutanoic acid,

15 4-{[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-2,2-dimethyl-4-oxobutanoic acid,

20 4-{[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-2,2-dimethyl-4-oxobutanoic acid,

N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-2-(2-methoxyethoxy)acetamide,

N-[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-2-(2-methoxyethoxy)acetamide,

25 5-{[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-5-oxopentanoic acid,

5-{[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-5-oxopentanoic acid,

30 methyl 5-{[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]amino}-5-oxopentanoate,

N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-N'-methylpentanediamide,

- (2-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2-oxoethoxy)acetic acid,
- (2-{[4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2-oxoethoxy)acetic acid,
- 5 methyl (2-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2-oxoethoxy)acetate,
- methyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)phenyl]carbamate,
- methyl [4-(4-bromo-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 10 2-methoxyethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,
- 2-methoxyethyl [4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 15 2-fluoroethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 2-ethoxyethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 3-methoxypropyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 20 2-(2-methoxyethoxy)ethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 2-(2-ethoxyethoxy)ethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 25 (2S)-2-methoxypropyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 1-methoxy-2-propanyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- tetrahydro-3-furanyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,
- 30 tetrahydro-3-furanylmethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,

- tetrahydro-2-furanylmethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
3-oxetanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
5 2-(dimethylamino)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
3-hydroxy-3-methylbutyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
2-hydroxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
10 2-hydroxy-2-methylpropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
2-hydroxy-2-methylpropyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
15 3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]-1,1-dimethylurea,  
3-[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]-1,1-dimethylurea,  
3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]-1-methoxy-1-methylurea,  
20 3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]-1-(2-methoxyethyl)-1-methylurea,  
3-[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]-1-(2-methoxyethyl)-1-methylurea,  
25 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1,2,3,4-tetrahydro-6-quinolinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,  
6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-methyl-1H-imidazol-5-yl)-3-methyl-3,4-dihydro-2(1H)-quinazolinone,  
2-amino-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)benzoic acid,  
30 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-hydroxybenzoic acid,



- 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-hydroxybenzoic acid,
- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-methylbenzoic acid,
- 5 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-methylbenzoic acid,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(methylsulfonyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{4-[(S)-
- 10 methylsulfinyl]phenyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{4-[(R)-methylsulfinyl]phenyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(S-methylsulfonimidoyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- 15 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(S-methylsulfonimidoyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-2-methoxy-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-
- 20 chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-6-fluoro-2-pyridinyl]carbamate,
- N-[6-chloro-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]acetamide,
- 25 2-methoxyethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 2-methoxyethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-
- 30 1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

- (3S)-3-[5-(6-amino-4-methyl-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-4-methyl-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-3-[5-(6-amino-3-pyridazinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-3-pyridazinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(2-amino-5-pyrimidinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 10 (3S)-3-[5-(2-amino-5-pyrimidinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(5-hydroxy-2-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 15 (3S)-3-[4-chloro-5-(5-hydroxy-2-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(6-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(6-oxo-1,6-dihydro-3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 20 (3S)-3-[4-chloro-5-(6-oxo-1,6-dihydro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 25 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-[2-fluoro-6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-[2-chloro-6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-[6-(dimethylamino)-3-pyridinyl]-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 30 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-[6-(ethylamino)-3-pyridinyl]-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,

- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(2-ethoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(3-methoxypropyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(3-hydroxy-3-methylbutyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(1,3-oxazol-2-ylmethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone,
- 10 3-{[5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-pyridinyl]amino}-N,N-dimethylpropanamide,
- 3-{[5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-pyridinyl]amino}propanamide,
- 15 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(2-oxo-1-pyrrolidinyl)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 20 (3S)-3-{4-chloro-5-[6-(dimethylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-(4-chloro-5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-{4-chloro-5-[6-(ethylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 25 (3S)-3-(4-chloro-5-{6-[(2-ethoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-(4-chloro-5-{6-[(3-methoxypropyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 30 (3S)-3-(4-chloro-5-{6-[(3-hydroxy-3-methylbutyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-(4-chloro-5-{6-[(1,3-oxazol-2-ylmethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

3-{[5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]amino}-N,N-dimethylpropanamide,

(3S)-3-{4-chloro-5-[6-(2-oxo-1-pyrrolidinyl)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(2-amino-1,3-thiazol-5-yl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(2-amino-4-chloro-1,3-thiazol-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

N-[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]acetamide,

methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(3S)-3-[5-(2-amino-3-chloro-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(2-amino-5-chloro-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(2-amino-5-chloro-4-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(2-amino-4-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[2-(methylamino)-4-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-{4-chloro-5-[2-(methylamino)-4-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(3-amino-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-methyl-4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2,6-dimethyl-4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[4-chloro-5-(2,6-dimethyl-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- (3S)-3-[5-(6-amino-2-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-3-[5-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-fluoro-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- (3R)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 10 (3R)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-5-phenyl-1H-imidazole-4-carboxylic acid,
- 15 2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-5-phenyl-1H-imidazole-4-carbonitrile,
- 4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)benzamide,
- 4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)benzamide,
- 20 (3S)-3-[2-(6-amino-3-pyridinyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[2-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 25 methyl [5-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)-2-pyridinyl]carbamate,
- methyl [6-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)-3-pyridinyl]carbamate,
- methyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,
- 30 (3S)-3-[2-(4-aminophenyl)-4-chloro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- (3S)-3-[2-(4-aminophenyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,  
5 2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,  
(1E)-N-[4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]-N'-hydroxyethanimidamide,  
N-[4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-  
10 tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]acetamide,  
2-methoxyethyl [4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,  
3-methoxypropyl [4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,  
15 2-(2-methoxyethoxy)ethyl [4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,  
3-oxetanyl [4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,  
(1E)-N-[4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-  
20 1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]-N'-hydroxyethanimidamide,  
2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,  
2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,  
25 (3S)-3-[2-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-[2-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-[2-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-5-yl]-7-[5-  
30 chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
2-(2-methoxyethoxy)ethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

2-(2-methoxyethoxy)ethyl [5-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4-fluoro-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

(3S)-3-[2-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

methyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

2-fluoroethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

N-[4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4H-1,2,4-triazol-3-yl)phenyl]acetamide,

3-oxetanyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

2-methoxyethyl [3-chloro-4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

(3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-4H-1,2,4-triazol-3-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-phenyl-1H-pyrrol-2-yl)-2,3-dihydro-5(1H)-indolizinone,

(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-nitrophenyl)-1H-pyrrol-2-yl]-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(4-aminophenyl)-1H-pyrrol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

methyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-pyrrol-2-yl)phenyl]carbamate,

- 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-pyrrol-2-yl)phenyl]carbamate,  
(3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-pyrrol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
5 (3S)-3-[1-(4-aminophenyl)-5-oxo-2,5-dihydro-1H-pyrazol-3-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
methyl [4-(3-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-5-oxo-2,5-dihydro-1H-pyrazol-1-yl)phenyl]carbamate,  
2-methoxyethyl [4-(3-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-5-oxo-2,5-dihydro-1H-pyrazol-1-yl)phenyl]carbamate,  
10 2-(2-methoxyethoxy)ethyl [4-(3-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-5-oxo-2,5-dihydro-1H-pyrazol-1-yl)phenyl]carbamate,  
methyl [4-(2-((3S)-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
15 2-methoxyethyl [4-(2-((3S)-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
20 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
methyl [4-(2-((3S)-7-[6-methyl-3-(1H-tetrazol-1-yl)-2-pyridinyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,  
(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-fluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
25 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-fluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[2,3-difluoro-6-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
30 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[2,3-difluoro-6-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[3,5-difluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,



- (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[3,5-difluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-(difluoromethoxy)-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-(difluoromethoxy)-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-(difluoromethoxy)-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 3-{(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyl}-4-(1H-tetrazol-1-yl)benzonitrile,
- 10 3-{(3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyl}-4-(1H-tetrazol-1-yl)benzonitrile,
- 3-{(3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyl}-4-(1H-tetrazol-1-yl)benzonitrile,
- 15 (6S)-6-[5-(4-aminophenyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,
- 3-oxetanyl [4-(2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate,
- 2-methoxyethyl [4-(2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate,
- 20 2-ethoxyethyl [4-(2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate,
- 3-methoxypropyl [4-(2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate,
- 25 methyl [4-(2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,
- 2-methoxyethyl [4-(2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,
- 30 (6S)-6-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,
- (6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

- (6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-(5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
methyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
5 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
methyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
10 methyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,  
2-methoxyethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,  
15 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
20 2-ethoxyethyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
3-methoxypropyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
25 (6S)-6-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
(6S)-6-(4-chloro-5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
30 (6S)-6-[5-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

methyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

5 methyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

10 (6S)-6-[5-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

methyl [4-(5-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-2-yl)phenyl]carbamate,

15 methyl [4-(4-chloro-5-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-2-yl)phenyl]carbamate,

(3S)-3-[5-(4-aminophenyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

2-methoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

20 3-oxetanyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

methyl [4-(4-methyl-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

25 methyl [4-(5-methyl-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1,3-oxazol-4-yl)phenyl]carbamate,

2-methoxyethyl [4-(4-methyl-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

2-ethoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

30 3-methoxypropyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- (3S)-3-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl [6-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 2-methoxyethyl [6-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- 10 6-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone,
- methyl [5-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 15 2-methoxyethyl [5-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 2-ethoxyethyl [4-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,
- 3-methoxypropyl [4-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,
- 20 2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 25 (3S)-3-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl [6-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- methyl [5-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 30 2-methoxyethyl [5-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

- methyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
2-methoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
5 2-ethoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
3-methoxypropyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
methyl [4-(4-methyl-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-10 4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
2-methoxyethyl [4-(4-methyl-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
(6S)-6-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-methyl-2-(1H-tetrazol-15 1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
(6S)-6-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
(6S)-6-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
20 methyl [6-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
2-methoxyethyl [6-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
25 methyl [5-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,  
2-methoxyethyl [5-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,  
30 methyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

5 2-ethoxyethyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

3-methoxypropyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

10 (6S)-6-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

15 methyl [5-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

20 2-methoxyethyl [5-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

methyl [6-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

25 (6S)-6-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate,

(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

30 (3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-3-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
methyl [6-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
5 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
6-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone,  
10 4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzamide,  
methyl [5-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,  
2-methoxyethyl [5-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,  
15 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-3-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
20 methyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
25 methyl [4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
30

methyl [4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

(6S)-6-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-6-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

methyl [6-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)benzamide,

methyl [5-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

methyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,



(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-6-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

5 methyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

10 methyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

15 (6S)-6-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

methyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)phenyl]carbamate,

20 2-methoxyethyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)phenyl]carbamate,

methyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

25 2-methoxyethyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-2,3-dihydro-5(1H)-indolizinone,

30 methyl [5-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-methoxyethyl [5-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

5 methyl [4-(4-chloro-2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

2-methoxyethyl [4-(4-chloro-2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

10 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-2,3-dihydro-5(1H)-indolizinone,

methyl [5-(4-chloro-2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

15 2-methoxyethyl [5-(4-chloro-2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

N-carbamoyl-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxamide,

20 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[5-(methylsulfonyl)-2-thienyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,

(3S)-3-{4-chloro-5-[5-(methylsulfonyl)-2-thienyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

25 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(methylsulfonyl)-2-thiophenecarboxamide,

5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(methylsulfonyl)-2-thiophenecarboxamide,

30 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furoic acid,

5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furamide,

- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-methyl-2-furamide,
- 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furoic acid,
- 5 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furamide,
- 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-methyl-2-furamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furoic acid,
- 10 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furoic acid,
- 2-methyl-2-propenyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-furoate,
- 15 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-furoic acid,
- 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1,3-oxazole-4-carboxylic acid,
- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-pyrrole-2-carboxylic acid,
- 20 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-pyrrole-2-carboxylic acid,
- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-pyrrole-3-carboxylic acid,
- 25 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1-methyl-1H-pyrrol-2-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1-methyl-1H-pyrrole-2-carboxylic acid,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-hydroxy-1-methyl-1H-pyrazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 30 ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thienyl]acetate,

- [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetic acid,  
ethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetate,  
5 [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetic acid,  
4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzenesulfonamide,  
4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-  
10 tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzenesulfonamide,  
3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzenesulfonamide,  
(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-{5-[4-(5-oxo-4,5-dihydro-1,2,4-oxadiazol-3-yl)]phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
15 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-{5-[3-(5-oxo-4,5-dihydro-1,2,4-oxadiazol-3-yl)]phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-{5-[4-(trifluoroacetyl)]phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-{5-[4-(2,2,2-trifluoro-1-  
20 hydroxyethyl)]phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,  
[3-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetic acid,  
2-methyl-2-propanyl {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetate,  
25 {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetic acid,  
{[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetic acid,  
{[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-  
30 tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetic acid,  
2-methyl-2-propanyl {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}acetate,

- {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]amino}acetic acid,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarbonitrile,
- 5 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(methylsulfonyl)-2-thiophenecarboxamide,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-fluoro-2-thiophenecarboxylic acid,
- 10 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-fluoro-2-pyridinecarboxylic acid,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-thiophenecarboxylic acid,
- 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-pyrrol-2-yl)-2-thiophenecarboxylic acid,
- 15 ethyl 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)-2-thiophenecarboxylate,
- ethyl 4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)-2-thiophenecarboxylate,
- 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)-2-thiophenecarboxylic acid,
- 20 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)-2-thiophenecarboxylic acid,
- 4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)-2-thiophenecarboxylic acid,
- 25 4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)-2-thiophenecarboxylic acid,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-pyrimidinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-fluorophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 30 (3S)-3-[5-(4-chlorophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- (3S)-3-[5-(3-chlorophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-3-[5-(2-chlorophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(methylthio)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-methylphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzonitrile,
- 10 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)benzonitrile,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[3-(trifluoromethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone,
- 15 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-hydroxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-nitrophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(4-cyano-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]carbamate,
- 20 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(4-cyano-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)phenyl]carbamate,
- 1-(2-((3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indoliziny)-4-chlorophenyl)-1H-1,2,3-triazole-4-carbonitrile,
- 25 2-hydroxy-2-methylpropyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- 30 3-hydroxy-3-methylbutyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-methoxyethoxy)ethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-ethoxyethoxy)ethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-methoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-ethoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-methoxyethoxy)ethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-ethoxyethoxy)ethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-methoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

2-(2-ethoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(3S)-3-[5-(5-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

- (6S)-6-[5-(5-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,  
methyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,  
5 methyl [4-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,  
methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-methylphenyl]carbamate,  
2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate,  
10 (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-hydroxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,  
(3S)-3-[4-chloro-5-(4-hydroxyphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
15 ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate,  
3-methoxypropyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,  
20 (3S)-3-[5-(6-amino-5-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,  
3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-fluorobenzoic acid,  
5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-fluorobenzoic acid,  
25 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-methylbenzoic acid,  
2-chloro-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoic acid,  
30 1-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]cyclopropanecarboxylic acid,  
1-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]cyclopropanecarboxylic acid,



- 1-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]cyclopropanecarboxylic acid,  
3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2,6-difluorobenzoic acid,  
5 2-chloro-3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)benzoic acid,  
(2E)-3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thienyl]acrylic acid,  
3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-  
10 3-indoliziny])-1H-imidazol-5-yl)-2-thienyl]propanoic acid,  
5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)nicotinic acid,  
2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1,3-thiazole-5-carboxylic acid,  
15 6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,  
4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1,3-thiazole-2-carboxylic acid,  
4-chloro-3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-  
20 tetrahydro-3-indoliziny])-1H-imidazol-5-yl)benzoic acid,  
[5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1-benzothiophen-3-yl]acetic acid,  
[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenoxy]acetic acid,  
25 [3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenoxy]acetic acid,  
2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1,3-thiazole-4-carboxylic acid,  
3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-  
30 indoliziny])-1H-imidazol-5-yl)-4-fluorobenzoic acid,  
[6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1-benzothiophen-3-yl]acetic acid,

- 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)isonicotinic acid,
- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-thiophenecarboxylic acid,
- 5 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-pyrazole-5-carboxylic acid,
- 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indole-2-carboxylic acid,
- 5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-
- 10 tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-1H-indole-2-carboxylic acid,
- [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indol-3-yl]acetic acid,
- [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-
- 15 indoliziny])-1H-imidazol-5-yl)-1H-indol-3-yl]acetic acid,
- [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indazol-3-yl]acetic acid,
- {[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]amino}acetic acid,
- 20 [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indol-1-yl]acetic acid,
- [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-benzimidazol-1-yl]acetic acid,
- 5-(2-((3S)-6-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-
- 25 tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-
- tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- 4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-
- tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- 30 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-
- tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- 4-(4-chloro-2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-
- 1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,

- 4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- 5 (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1,2,3-thiadiazol-4-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 2-oxo-2-(1-pyrrolidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-
- 10 thiophenecarboxylate,
- 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide,
- 15 ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- isobutyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 2,3-dihydro-1H-inden-5-yl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-
- 20 thiophenecarboxylate,
- (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 25 2-(4-morpholinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 2-(dimethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-
- 30 thiophenecarboxylate,
- 2-(diethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

2-oxo-2-(1-piperidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

5 2-(4-morpholinyl)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

isopropyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

10 1-[[cyclohexyloxy]carbonyl]oxyethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

cyclohexyl 2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-5-[[1-  
15 [[cyclohexyloxy]carbonyl]oxy]ethoxy]carbonyl]-3-thienyl)-1H-imidazole-1-carboxylate,

[(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

20 [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1-[[2,2-dimethylpropanoyl]oxy]methyl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-fluoro-2-thiophenecarboxylic acid,

25 5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid, and

30 methyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

salts thereof, N-oxides thereof, solvates thereof, and prodrugs thereof.

Particularly preferred compounds of formula (I-B) include:

- 4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-1-[(5-phenyl-1H-imidazol-2-yl)methyl]-  
2(1H)-pyridinone,
- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
5 pyridinyl}methyl)-1H-imidazol-4-yl]phenyl} carbamate,
- methyl {4-[5-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
pyridinyl}methyl)-1H-imidazol-4-yl]phenyl} carbamate,
- methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
pyridinyl}propyl)-1H-imidazol-4-yl]phenyl} carbamate,
- 10 methyl {4-[5-chloro-2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-  
1(2H)-pyridinyl}propyl)-1H-imidazol-4-yl]phenyl} carbamate,
- methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
pyridinyl}ethyl)-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
15 pyridinyl}cyclopropyl)-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
pyridinyl}-2-phenylethyl)-1H-imidazol-4-yl]phenyl} carbamate,
- methyl [4-(2-{1-[4-(2-carbamimidamido-5-chlorophenyl)-2-oxo-1(2H)-  
pyridinyl]-2-phenylethyl}-1H-imidazol-4-yl)phenyl] carbamate,
- 20 methyl {4-[5-chloro-2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-  
1(2H)-pyridinyl}-2-phenylethyl)-1H-imidazol-4-yl]phenyl} carbamate,
- 4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-1-{2-phenyl-1-[5-(1H-pyrazol-1-yl)-  
1H-benzimidazol-2-yl]ethyl}-2(1H)-pyridinone,
- 1-{1-[5-(4-aminophenyl)-1H-imidazol-2-yl]-2-phenylethyl}-4-(2,5-  
25 dichlorophenyl)-2(1H)-pyridinone,
- methyl [4-(2-{1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-  
phenylethyl}-1H-imidazol-4-yl)phenyl] carbamate,
- methyl [4-(5-chloro-2-{1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-  
phenylethyl}-1H-imidazol-4-yl)phenyl] carbamate,
- 30 methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
pyridinyl}methyl)-4-methyl-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-  
pyridinyl}methyl)-4-ethyl-1H-imidazol-5-yl]phenyl} carbamate,

- 6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]-3,4-dihydro-2(1H)-quinolinone,
- 6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]-3,4-dihydro-2(1H)-quinolinone,
- 5 methyl {3-chloro-4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]benzamide,
- 4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]benzamide,
- 10 6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]-3-methyl-3,4-dihydro-2(1H)-quinazolinone,
- 6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]-3-methyl-3,4-dihydro-2(1H)-quinazolinone,
- 15 1-{{5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl}methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2(1H)-pyridinone,
- methyl {5-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]-2-pyridinyl} carbamate,
- 1-{{5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl}methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2(1H)-pyridinone,
- 20 methyl {6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-1H-imidazol-5-yl]-3-pyridinyl} carbamate,
- methyl {4-[2-(1-{{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}-3-methoxypropyl)-1H-imidazol-5-yl]phenyl} carbamate,
- 25 methyl {4-[4-chloro-2-(1-{{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}-3-methoxypropyl)-1H-imidazol-5-yl]phenyl} carbamate,
- methyl (4-{2-[(1S)-1-{{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}-2-phenylethyl]-1H-imidazol-4-yl}phenyl) carbamate,
- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl)methyl]-4-methyl-1H-imidazol-5-yl]phenyl} carbamate,
- 30 methyl (4-{2-[1-{{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}-2-(3-pyridinyl)ethyl]-1H-imidazol-5-yl}phenyl) carbamate,

- methyl (4-{4-chloro-2-[1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl]-2-(3-pyridinyl)ethyl]-1H-imidazol-5-yl}phenyl)carbamate,
- methyl 1-(4-chloro-2-{1-[(1S)-1-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-2-phenylethyl]-2-oxo-1,2-dihydro-4-pyridinyl}phenyl)-1H-1,2,3-triazole-4-carboxylate,
- 5 methyl [4-(2-{(1S)-1-[4-{5-chloro-2-[4-(trifluoromethyl)-1H-1,2,3-triazol-1-yl]phenyl]-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-4-yl)phenyl]carbamate,
- methyl (4-{2-[(1S)-1-{4-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-5-yl}phenyl)carbamate,
- 10 methyl (4-{2-[(1S)-1-{4-[5-chloro-2-(4-fluoro-1H-pyrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-5-yl}phenyl)carbamate,
- methyl [4-(2-{(1S)-1-[4-{5-chloro-2-[4-(trifluoromethyl)-1H-pyrazol-1-yl]phenyl]-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-4-yl)phenyl]carbamate,
- methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}ethyl)-1H-imidazol-5-yl]phenyl}carbamate,
- 15 methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate,
- methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-ethyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate,
- 20 4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-2-thiophenecarboxylic acid,
- methyl {4-[2-(1-{4-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate,
- methyl {4-[4-chloro-2-(1-{4-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate,
- 25 1-{[5-(4-aminophenyl)-1H-imidazol-2-yl]methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone,
- 2-methoxyethyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate,
- 30 methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-4-methyl-1H-imidazol-5-yl]phenyl}carbamate,
- 2-methoxyethyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-4-methyl-1H-imidazol-5-yl]phenyl}carbamate,

- 1-{{5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl)methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone,  
 4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-1-({5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl)methyl}-2(1H)-pyridinone,  
 5 1-{{5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl)methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone,  
 1-{{5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl)methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone,  
 methyl {6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-3-pyridinyl} carbamate,  
 10 2-methoxyethyl {6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-3-pyridinyl} carbamate,  
 methyl {5-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-2-pyridinyl} carbamate,  
 15 2-methoxyethyl {5-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-2-pyridinyl} carbamate,  
 2-methoxyethyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]phenyl} carbamate,  
 1-({4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl)methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone,  
 20 methyl {6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-3-pyridinyl} carbamate,  
 2-methoxyethyl {6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-3-pyridinyl} carbamate,  
 25 2-methoxyethyl {5-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]-2-pyridinyl} carbamate,  
 1-{{5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl)methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone,  
 1-{{5-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl)methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone, and  
 30 methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl}-1H-imidazol-5-yl]phenyl} carbamate,  
 salts thereof, N-oxides thereof, solvates thereof, and prodrugs thereof.



Particularly preferred compounds of formula (I-C) include:

- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 5 2-methoxyethyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 2-methoxyethyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 3-{[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]methyl}-6-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-4(3H)-pyrimidinone,
- 10 3-{[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]methyl}-6-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-4(3H)-pyrimidinone,
- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-isopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 15 methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-isopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-cyclopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-cyclopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 20 methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-cyclopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-4-fluoro-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-6-oxo-2-(tetrahydro-2H-pyran-4-yl)-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- 25 methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-(methoxymethyl)-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-(methoxymethyl)-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate, and
- 30 methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate,
- salts thereof, N-oxides thereof, solvates thereof, and prodrugs thereof.

Compounds of the present invention containing one or more chiral centres may be used in enantiomerically or diastereoisomerically pure form, or in the form of a mixture of isomers. For the avoidance of doubt, the compounds of the invention may be used in any tautomeric form.

Unless otherwise specifically mentioned, all isomers are included in the present invention. For example, alkyl, alkenyl, alkynyl, alkoxy and alkylthio may be straight chain or branched. Moreover, all isomers due to double bond, ring and fused ring (E-, Z-, cis- and trans-forms), isomers due to the presence of asymmetric carbon(s) etc. (R-, S-,  $\alpha$ - and  $\beta$ -configuration, enantiomer and diastereomer), optically active substances having optical rotation (D-, L-, d- and l-forms), polar compounds by chromatographic separation (more polar compounds and less polar compounds), equilibrium compounds, rotational isomers, a mixture thereof in any proportion and a racemic mixture are included in the present invention.

According to the present invention, symbol  represents  $\alpha$ -configuration, symbol  represents  $\beta$ -configuration and symbol  represents  $\alpha$ -configuration,  $\beta$ -configuration or a mixture of them. There is no particular limitation for the ratio of  $\alpha$ -configuration and  $\beta$ -configuration in the mixture.

20

#### SALTS:

The salt of the compound of formula (I) includes all nontoxic salts or pharmaceutically acceptable salts. With regard to the pharmaceutically acceptable salts, those which are low-toxicity and soluble in water are preferred. Examples of appropriate salts of the compound of formula (I) are salt with alkaline metal (such as potassium, sodium and lithium), salt with alkaline earth metal (such as calcium and magnesium), ammonium salt (such as ammonium salt, tetramethylammonium salt and tetrabutylammonium salt), salt with organic amine (such as triethylamine, methylamine, dimethylamine, cyclopentylamine, benzylamine, phenethylamine, piperidine, monoethanolamine, diethanolamine, tris(hydroxymethyl) methylamine, lysine, arginine and N-methyl-D-glucamine) and acid addition salt (such as inorganic acid salt (e.g., hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate and nitrate) and organic

30

acid salt (e.g., formate, acetate, trifluoroacetate, lactate, tartrate, oxalate, fumarate, maleate, benzoate, citrate, methanesulfonate, ethanesulfonate, benzenesulfonate, toluenesulfonate, isothionate, glucuronate and gluconate), etc.). The salt of the compound of the present invention also includes solvates and also solvates with the above-mentioned alkaline (earth) metal salt, ammonium salt, organic amine salt and acid addition salt. The solvate is preferably low-toxic and water-soluble. Examples of an appropriate solvate are solvates with water and with alcoholic solvent (such as ethanol). The compounds of the present invention are converted to low-toxicity salts or pharmaceutically acceptable salts by known methods.

Moreover, the salt includes a quaternary ammonium salt. The quaternary ammonium salt of the compound represented by formula (I) is the compound where nitrogen of the compounds represented by formula (I) is quarternalized by  $R^0$  ( $R^0$  is C1-8 alkyl or C 1-8 alkyl substituted by phenyl.).

The salt also includes an N-oxide. The compound of the present invention can be converted into an N-oxide by known methods. The N-oxide is the compound where nitrogen of the compound represented by formula (I) is oxidized.

#### Prodrugs:

A prodrug of the compound of formula (I) means a compound which is converted to the compound of formula (I) by reaction with an enzyme, gastric acid or the like in the living body. For example, with regard to a prodrug of the compound of formula (I), when the compound of formula (I) has an amino group, compounds in which the amino group is, for example, acylated, alkylated or phosphorylated (e.g., compounds in which the amino group of the compound of formula (I) is eicosanoylated, alanylated, pentylaminocarbonylated, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylated, tetrahydrofuranylated, pyrrolidylmethylated, pivaloyloxymethylated, acetoxymethylated, tert-butylation, etc.); when the compound of formula (I) has a hydroxyl group, compounds where the hydroxyl group is, for example, acylated, alkylated, phosphorylated or borated (e.g., compounds in which the hydroxyl group of the compound of formula (I) is acetylated, palmitoylated, propanoylated, pivaloylated, succinylated, fumarylated, alanylated or

dimethylaminomethylcarbonylated); and when the compound of formula (I) has a carboxyl group, compounds in which the carboxyl group is, for example, esterified or amidated (e.g., compounds in which the carboxyl group of the compound of formula (I) is made into ethyl ester, phenyl ester, phenylethyl ester, carboxymethyl ester, dimethylaminomethyl ester, pivaloyloxymethyl ester, ethoxycarbonyloxyethyl ester, phthalidyl ester, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester, cyclohexyloxycarbonylethyl ester or methylamide). Those compounds may be produced by a known method per se. The prodrug of the compound of formula (I) may be either a hydrate or a non-hydrate. A prodrug of the compound of formula (I) may also be a compound which is converted to the compound of formula (I) under physiological conditions as described in "*Iyakuhin no kaihatsu*, Vol.7 (Bunshi-sekkei), pp. 163-198 (Hirokawa-Shoten), 1990". Further, the compound of formula (I) may also be labeled by a radio isotope (such as  $^2\text{H}$ ,  $^3\text{H}$ ,  $^{11}\text{C}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$ ,  $^{13}\text{N}$ ,  $^{15}\text{N}$ ,  $^{15}\text{O}$ ,  $^{17}\text{O}$ ,  $^{18}\text{O}$ ,  $^{35}\text{S}$ ,  $^{18}\text{F}$ ,  $^{36}\text{Cl}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ , etc.).

#### PROCESSES FOR THE PREPARATION OF THE COMPOUND OF THE PRESENT INVENTION:


The compounds of the invention can, for example, be prepared according to the following reaction schemes.

The compound of the present invention represented by the formula (I) may be prepared by known methods, for example, a method combining the following methods, the method according to these methods, the methods described in the examples and/or methods described in *Comprehensive Organic Transformations: A Guide to Functional Group Preparations*, 2nd Edition (Richard C. Larock, John Wiley & Sons Inc, 1999), etc., which are appropriately modified in each following method for the preparation. Salts of the starting materials may be used.


It will also be recognized that another major consideration in the planning of any synthetic route in this field is the judicious choice of the protecting group used for protection of the reactive functional groups present in the compounds described in this invention. Protection reactions may be carried out by the methods, for example, described in T. W. Greene, *Protective Groups in Organic Synthesis*, Wiley, New York, 1999.

The compound of formula (I) wherein  $\text{R}^{62}$  represents hydrogen can be prepared

from a compound represented by formula (II):




- wherein  $R^{2-1}$ ,  $R^{3-1}$ ,  $R^{4-1}$  and  $R^{6-1}$  have the same meanings as  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^6$  respectively, with the proviso that carboxyl, hydroxyl, amino or thiol in  $R^{2-1}$ ,  $R^{3-1}$ ,  $R^{4-1}$  and  $R^{6-1}$  may be protected, if necessary, or a compound represented by the formula (III):



wherein Pg represents a protective group of carboxyl, such as C1-4 alkyl, and the other symbols have the same meaning described above, by  $Cyc^1$  ring formation reactions described below.


10

- 1) The compound of formula (I) wherein  $R^{62}$  represent hydrogen,  $Cyc^1$  represents an imidazole ring which is attached to  $Cyc^3$  at the 4-position and has an  $R^{1-1}$ , that is, a compound represented by formula (I-I):




- 15 wherein  $R^{1-1}$  and  $R^{7-1}$  have the same meanings as  $R^1$  and  $R^7$  respectively, with the proviso that, carboxyl, hydroxyl, amino or thiol in  $R^{1-1}$  and  $R^{7-1}$  may be protected, if necessary, and the other symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 1:

## Reaction Scheme 1



In Reaction Scheme 1, the reaction from the compound represented by formula (II) to the compound represented by formula 1b is an alkylation reaction.

- 5 The alkylation reaction is well known. For example, the compound represented by formula (II) with the compound represented by formula 1a:



wherein X represents fluorine, chlorine, bromine or iodine, and the other symbols have the same meaning described above, can be conducted in a solvent such as

- 10 dimethylformamide, tetrahydrofuran, dichloromethane, acetone or acetonitrile in the presence of a base such as sodium carbonate, potassium carbonate, cesium carbonate, sodium bicarbonate potassium bicarbonate, *N,N*-diisopropylethylamine or triethylamine at  $-20^\circ\text{C}$  to reflux temperature to form a compound represented by formula 1b wherein all symbols have the same meaning described above.

- 15 The reaction from the compound represented by formula 1b to the compound represented by formula (I-I) is an imidazole formation reaction.

- The imidazole formation reaction is well known. For example, the compound represented by formula 1b and ammonium acetate or ammonium trifluoroacetate in a suitable solvent such as xylene, toluene or acetic acid, on heating and/or microwave irradiation, can form compounds of formula (I-I).
- 20

Alternatively, the compound represented by formula 1d can be prepared from the compound represented by formula (II). The reaction from the compound represented by formula (II) to the compound represented by formula 1d is an amidation reaction.

The amidation reaction is well known. For example, the reaction of the compound represented by formula (II) with the compound represented by formula 1c wherein all symbols have the same meaning described above is exemplified by:

- (1) A reaction procedure with use of an acid halide,
- (2) A reaction procedure with use of a mixed acid anhydride, and
- (3) A reaction procedure with use of a condensing agent.

Referring specifically to these reaction procedures,

- (1) The reaction procedure employing an acid halide is conducted in practice, for example, by reacting a carboxylic acid with an acid halogenating agent (e.g., oxalyl chloride, thionyl chloride, etc.) in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran, dimethoxyethane, etc.) at a temperature from about -20 °C to the refluxing temperature, followed by reaction of the resultant acid halide with an amine in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran, acetonitrile, ethyl acetate, etc.) or solvent-free in the presence of a base (e.g., pyridine, triethylamine, dimethylaniline, dimethylaminopyridine, diisopropylethylamine, etc.) at a temperature of approximately 0 to 40° C. Alternatively, the procedure can be carried out by reacting the resultant acid halide with an amine in an organic solvent (e.g., dioxane, tetrahydrofuran, dichloromethane, etc.) in the presence or absence of a phase-transfer catalyst (e.g., tetrabutylammonium chloride, triethylbenzylammonium chloride, tri-n-octylmethylammonium chloride, trimethyldecylammonium chloride, tetramethylammonium chloride, trimethyldecylammonium chloride, tetramethylammonium chloride, etc.) at a temperature of about 0 to 40 °C, whilst using an aqueous alkali solution (e.g., an aqueous sodium bicarbonate or sodium hydroxide solution, etc.).
- (2) The reaction procedure employing a mixed acid anhydride is conducted in practice, for example, by reacting a carboxylic acid with an acid halide (pivaloyl chloride, tosyl chloride, mesyl chloride, etc.) or an acid derivative (e.g., ethyl chloroformate, isobutyl chloroformate, etc.) in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran, etc.) or solvent free

in the presence of base (e.g., pyridine, triethylamine, dimethylaniline, dimethylaminopyridine, diisopropylethylamine, etc.) at a temperature of about 0 to 40 °C, followed by reaction of the resultant mixed acid anhydride with an amine in an organic solvent (e.g., chloroform, dichloroethane, diethyl ether, tetrahydrofuran, etc.) at a temperature of about 0 to 40 °C.

- (3) The reaction procedure with use of a condensing agent is carried out, for example, by reacting a carboxylic acid with an amine in an organic solvent (e.g., chloroform, dichloromethane, dimethylformamide, diethyl ether, tetrahydrofuran, etc.) or solvent-free in the presence or absence of a base (e.g., pyridine, triethylamine, dimethylaniline, dimethylaminopyridine, etc.), with use of a condensing agent (e.g., 1,3-dicyclohexylcarbodiimide (DCC), 1-ethyl-3-[3-(dimethylamino)propyl]carbodiimide (EDC), 1,1'-carbonyldiimidazole (CDI), 2-chloro-1-methylpyridinium iodide, 1,1'-propylphosphonic acid anhydride (1-propanephosphonic acid cyclic anhydride, PPA), etc.) and with or without use of 1-hydroxybenzotriazole (HOBt), at a temperature of about 0 to 40 °C.

The reaction from the compound represented by formula 1d to the compound represented by formula (I-I) is an imidazole formation reaction. The imidazole formation reaction can be carried out by the same method as described above in Reaction Scheme 1.

In the course of the synthesis of the compound of the present invention represented by the formula (I-I), the deprotection reaction can be carried out at an appropriate synthetic stage, when the protective groups of carboxy, hydroxy, amino or mercapto group exists.

The deprotection reaction of the protective groups of carboxy, hydroxy, amino or mercapto group is well-known and includes, for example,

- (1) a deprotection reaction by alkali hydrolysis,
- (2) a deprotection under acidic conditions,
- (3) a deprotection reaction by hydrogenolysis,
- (4) a deprotection reaction of silyl group,
- (5) a deprotection reaction using a metal,
- (6) a deprotection reaction using a metal complex, etc.

To explain these methods in detail:

- (1) The deprotection reaction by alkali hydrolysis is carried out, for example, in an



organic solvent (methanol, tetrahydrofuran, 1,4-dioxane, etc.) using a hydroxide of alkali metals (sodium hydroxide, potassium hydroxide, lithium hydroxide, etc.), hydroxide of alkaline earth metals (barium hydroxide, calcium hydroxide, etc.), carbonate (sodium carbonate, potassium carbonate, etc.) or a solution thereof or a mixture thereof at a temperature of 0 to 40 °C.

(2) The deprotection reaction under acidic conditions is carried out, for example, in an organic solvent (dichloromethane, chloroform, dioxane, ethyl acetate, anisole, etc.), in an organic acid (acetic acid, trifluoroacetic acid, methanesulfonic acid, p-toluenesulfonic acid, etc.) or an inorganic acid (hydrochloric acid, sulfuric acid, etc.) or a mixture thereof (hydrobromic acid/acetic acid, etc.) in the presence or absence of 2,2,2-trifluoroethanol at a temperature of 0 to 100 °C.

(3) The deprotection reaction by hydrogenolysis is, for example, carried out in a solvent (e.g. ethers such as tetrahydrofuran, 1,4-dioxane, dimethoxyethane, diethyl ether, etc.; alcohols such as methanol, ethanol, etc.; benzenes such as benzene, toluene, etc.; ketones such as acetone, methyl ethyl ketone, etc.; nitriles such as acetonitrile etc.; amides such as N,N-dimethylformamide, N,N-dimethylacetamide etc.; water, ethyl acetate, acetic acid or a mixture of two or more thereof, etc.) in the presence of a catalyst (palladium-carbon, palladium black, palladium hydroxide, platinum oxide, Raney nickel, etc.) under an atmosphere of hydrogen at normal or increased pressure, or in the presence of ammonium formate at a temperature of 0 to 200 °C.

(4) The deprotection reaction of a silyl group is, for example, carried out in a water-miscible organic solvent (tetrahydrofuran, acetonitrile, etc.) using tetrabutylammonium fluoride at a temperature of 0 to 40 °C.

(5) The deprotection reaction using a metal is carried out, for example, in an acidic solvent (acetic acid, a buffer of pH 4.2 to 7.2 or a mixture of the solution thereof and an organic solvent such as tetrahydrofuran etc.) in the presence of zinc powder at a temperature of 0 to 40 °C optionally under sonication.

(6) The deprotection reaction using a metal complex is carried out, for example, in an organic solvent (dichloromethane, N,N-dimethylformamide, tetrahydrofuran, ethyl acetate, acetonitrile, dioxane, ethanol, etc.), water or a mixture thereof, in the presence of a trapping reagent (tributyltin hydride, triethylsilane, dimedone, morpholine, diethylamine, pyrrolidine, 1,3-dimethylbarbituric acid, etc.), an organic acid (acetic acid, formic acid, 2-ethylhexanecarboxylic acid, etc.) and/or a salt of an organic acid (sodium

2-ethylhexanoate, potassium 2-ethylhexanoate, etc.) in the presence or absence of a phosphine reagent (triphenylphosphine etc.) using a metal complex (tetrakis(triphenylphosphine)palladium (0), palladium(II) bis(triphenylphosphine) dichloride, palladium(II) acetate, rhodium(I) tris(triphenylphosphine) chloride, etc.) at a temperature of 0 to 40 °C.

In addition to the above, deprotection reactions may be carried out by the methods, for example, described in T. W. Greene, *Protective Groups in Organic Synthesis*, Wiley, New York, 1999.

A protective group for carboxy includes, for example, methyl, ethyl, allyl, tert-butyl, trichloroethyl, benzyl (Bn), phenacyl, p-methoxybenzyl, trityl, 2-chlorotrityl or a solid carrier containing these structures, etc.

A protective group for hydroxy includes, for example, methyl, trityl, methoxymethyl (MOM), 1-ethoxyethyl (EE), methoxyethoxymethyl (MEM), 2-tetrahydropyranyl (THP), trimethylsilyl (TMS), triethylsilyl (TES), tert-butyldimethylsilyl (TBDMS), tert-butyldiphenylsilyl (TBDPS), acetyl (Ac), pivaolyl, benzoyl, benzyl (Bn), p-methoxybenzyl, allyloxycarbonyl (Alloc) or 2,2,2-trichloroethoxycarbonyl (Troc), etc.


A protective group for amino includes, for example, benzyloxycarbonyl, tert-butoxycarbonyl, allyloxycarbonyl (Alloc), 1-methyl-1-(4-biphenyl)ethoxycarbonyl (Bpoc), trifluoroacetyl, 9-fluorenylmethoxycarbonyl (FMoc), benzyl (Bn), p-methoxybenzyl, benzyloxymethyl (BOM), 2-(trimethylsilyl)ethoxymethyl (SEM), etc.

A protective group for mercapto includes, for example, benzyl, methoxybenzyl, methoxymethyl (MOM), 2-tetrahydropyranyl (THP), diphenylmethyl, acetyl (Ac), etc.

Protective groups for carboxy, hydroxy, amino or mercapto group are not limited to those described above, but include groups which are easily and selectively deprotected. For example, those groups described in T. W. Greene, *Protective Groups in Organic Synthesis*, Wiley, New York, 1999.

As is easily understood by those skilled in the art, the target compound of the present invention may be prepared easily by selecting these deprotection reactions.


2) The compound of formula (I) wherein  $R^{62}$  represents hydrogen,  $Cyc^1$  represents an imidazole ring which is attached to  $Cyc^3$  at the 4-position and possesses  $R^{1-hal}$ , that is, a compound represented by formula (I-II-b):



wherein  $R^{1-hal}$  represents fluorine, chlorine, bromine or iodine, and the other symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 2:

5

Reaction Scheme 2




wherein all symbols have the same meaning described above.

In Reaction Scheme 2, the reaction from the compound represented by formula (I-II-a) to the compound represented by formula (I-II-b) is a halogenation reaction.

The halogenation reaction is well known. For example, the reaction of the compound represented by formula (I-II-a) with brominating or chlorinating agent, such as N-bromosuccinimide, N-chlorosuccinimide or 1,3-dichloro-5,5-dimethyl-hydantoin in a suitable solvent such as acetonitrile, chloroform or tetrahydrofuran from -20 °C to the refluxing temperature provides the compound represented by formula (I-II-b).

15


3) The compound of formula (I) wherein  $R^{62}$  represents hydrogen,  $Cyc^1$  represents imidazole ring which is attached to  $Cyc^3$  at the 2-position, that is, a compound represented by formula (I-III-a):



20

wherein all symbols have the same meaning described above, and a compound


represented by formula (I-III-b):



wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 3:

5

### Reaction Scheme 3



wherein all symbols have the same meaning described above.

In Reaction Scheme 3, the reaction from the compound represented by formula (II) to the compound represented by formula 3c can be prepared as described below.

10

The compound represented by formula 3b can be prepared by treatment of the appropriately functionalized compound represented by formula (II) with acid halogenating agent (e.g., oxalyl chloride, thionyl chloride, 1-chloro-N,N,2-trimethyl-1-propenylamine etc.) in an organic solvent (e.g., chloroform, dichloromethane, diethyl ether, tetrahydrofuran, dimethoxyethane, etc.) at a temperature from about -20 °C to the refluxing temperature, followed by reaction of the resultant acid halide with diazomethane or (trimethylsilyl)diazomethane in an organic solvent (e.g., chloroform, dichloromethane, hexane, diethyl ether, tetrahydrofuran, acetonitrile, etc.). The compound represented by formula 3c can be prepared by treatment of the compound


15

represented by formula 3b with HX (e.g., hydrochloric acid, hydrobromic acid etc.) in an organic solvent (e.g., chloroform, dichloromethane, hexane, diethyl ether, tetrahydrofuran, acetonitrile, etc.).

The reaction from the compound represented by formula 3c to the compound  
5 represented by formula (I-III-a) is an imidazole formation reaction. The imidazole formation reaction can be carried out by the same method as described in Reaction Scheme 1 above.


The reaction from the compound represented by formula (I-III-a) to the compound represented by formula (I-III-b) is a halogenation reaction. The halogenation  
10 reaction can be carried out by the same method as described in Reaction Scheme 2.

4) The compound of formula (I) wherein  $R^{62}$  represents hydrogen,  $Cyc^1$  represents 1,3,4-triazole ring, that is, a compound represented by formula (I-IV):



wherein all symbols have the same meaning described above, can be prepared as  
15 outlined in Reaction Scheme 4:

Reaction Scheme 4




wherein all symbols have the same meaning described above.

20 In Reaction Scheme 4, the reaction from the compound represented by formula (II) to the compound represented by formula 4b is an amidation reaction. The amidation

reaction can be carried out by the same method as described in Reaction Scheme 1.


The reaction from the compound represented by formula 4b to the compound represented by formula (I-IV) is a 1,3,4-triazole formation reaction. The 1,3,4-triazole formation reaction is well known. For example, the reaction can be carried out by heating the compound represented by 4b in a suitable solvent such as acetic acid, xylene or toluene to give the compound represented by formula (I-IV).

5) The compound of formula (I) wherein  $R^{62}$  represents hydrogen,  $Cyc^1$  represents pyridazinone ring, that is, a compound represented by formula (I-V):



wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 5:

Reaction Scheme 5



wherein all symbols have the same meaning described above.


In Reaction Scheme 5, the reaction from the compound represented by formula (III) to the compound represented by formula 5a can be prepared as described below.

The compound represented by formula 5a can be prepared by treatment of the appropriately functionalized compound represented by formula (III) with dimethylmethylphosphonate in the presence of base such as n-butyllithium in a solvent such as tetrahydrofuran.

The reaction from the compound represented by formula 5a to the compound represented by formula (I-V) is a Horner-Wadsworth-Emmons reaction.

The Horner-Wadsworth-Emmons reaction is well known. For example, the compound represented by formula 5a and the compound represented by formula 5b in the presence of base such as potassium carbonate in a solvent such as ethanol or tetrahydrofuran gives an  $\alpha$ ,  $\beta$ -unsaturated ketone derivative which can then be condensed with a suitably substituted hydrazine derivative represented by  $H_2NNHR^{1-1}$  to give the compound represented by formula (I-V).


6) The compound of formula (I) wherein  $R^{62}$  represents hydrogen,  $Cyc^1$  represents a 1,2,3-triazole ring, that is, a compound represented by formula (I-VI):



wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 6:

15

#### Reaction Scheme 6



wherein all symbols have the same meaning described above.

In Reaction Scheme 6, the reaction from the compound represented by formula (II) to the compound represented by formula 6a is an amidation reaction. The amidation

reaction can be carried out by the same method as described in Reaction Scheme 1 using N,O-dimethylhydroxylamine instead of the compound represented by formula 1c.

The reaction from the compound represented by formula 6a to the compound represented by formula 6b can be prepared as described below.

The compound represented by formula 6b can be prepared by treatment of the appropriately functionalized compound represented by formula 6a with lithium aluminum hydride or diisobutylaluminium hydride in tetrahydrofuran to give the compound represented by formula 6b.


The reaction from the compound represented by formula 6b to the compound represented by formula 6c can be prepared as described below.

The compound represented by formula 6c can be prepared by treatment of the appropriately functionalized compound represented by formula 6b with dimethyl (1-diazo-2-oxopropyl) phosphonate in the presence of base such as potassium carbonate in a solvent such as methanol.

The reaction from the compound represented by formula 6c to the compound represented by formula (I-VI) is a Huisgen reaction. The Huisgen reaction can be prepared as described below.

The compound represented by formula (I-VI) can be prepared by treatment of the appropriately functionalized compound represented by formula 6c and a suitably substituted compound represented by formula 6d in the presence of a copper (II) salt such as copper (II) sulfate, an ascorbate such as sodium ascorbate and a base such as sodium hydroxide which on heating and/or microwave irradiation gives the compound represented by formula (I-VI).

7) The compound of formula (I) wherein  $R^{62}$  represents hydrogen,  $Cyc^1$  represents 1,2-oxazole ring, that is, a compound represented by formula (I-VII):




wherein all symbols have the same meaning described above, can be prepared as



outlined in Reaction Scheme 7:

Reaction Scheme 7



wherein all symbols have the same meaning described above.

- 5 In Reaction Scheme 7, the reaction from the compound represented by formula 7a to the compound represented by formula 7c can be prepared as described below.

The compound represented by formula 7c can be prepared by treatment of an appropriately functionalized compound represented by formula 7a with hydroxylamine hydrochloride in a solvent such as tert-butanol and water to provide the corresponding  
 10 oxime compound represented by formula 7b, which can be converted to a nitrile oxide compound represented by formula 7c by treatment with an oxidant such as chloramine-T trihydrate.

- The reaction from the compound represented by formula (II) to the compound represented by formula 7d can be prepared by the same method as described in Reaction  
 15 Scheme 6.


The reaction from the compound represented by formula 7d to the compound represented by formula (I-VII) can be prepared as described below.

The compound represented by formula (I-VII) can be prepared by combining the

compound represented by formula 7c and the compound represented by formula 7d in the presence of a copper (II) salt such as copper (II) sulfate and copper powder in a solvent such as tert-butanol and water at a temperature from approximately 20 °C to the refluxing temperature with or without microwave irradiation.

5

8) The compound of formula (I-A) wherein  $R^{62}$  represents hydrogen, that is, a compound represented by formula (I-A-H) can be prepared from a compound represented by formula (II-A) or a compound represented by formula (III-A) by  $Cyc^1$  ring formation reactions by the same methods described in Reaction Schemes 1 to 7.



10

wherein all symbols have the same meaning described above.


The compound represented by formula (II-A) and formula (III-A) wherein  $R^{62}$  is hydrogen, Y is C(H) and U is  $CH_2$ , that is, a compound represented by formula (II-A-I) and (III-A-I) respectively, can be prepared as outlined in Reaction Scheme 8:

15

20

25

## Reaction Scheme 8



- 5 wherein Lg represents triflate, tosylate, chlorine or bromine,  $R^{8a}$  represents C1-4 alkyl,  $R^{B-1}$  and  $R^{B-2}$  represents hydrogen, or  $R^{B-1}$  and  $R^{B-2}$  may be taken together to form -C(CH<sub>3</sub>)<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>- and the other symbols have the same meaning described above.

In Reaction Scheme 8, imide formation of an appropriately protected compound represented by formula (IV) can be conducted by using alkylating agent such  
 10 as trimethyloxonium tetrafluoroborate, triethyloxonium tetrafluoroborate or dimethyl sulfate in a solvent such as dichloromethane, acetonitrile or dimethyl carbonate or without a solvent to form an imide compound represented by formula 8a.

Imide compounds represented by formula 8a can be condensed with a suitably protected 1,3-acetonedicarboxylic acid represented by formula 8b in an organic base  
 15 such as triethylamine, N,N-diisopropylethylamine or N-methylmorpholine at a temperature from about 20 °C to the refluxing temperature to provide a diester compound represented by formula 8c.

The diester compound represented by formula 8c can be converted to the

corresponding dicarboxylic acid represented by formula 8d by the same method as described above for the deprotection reaction of carboxyl.

The decarboxylation of a dicarboxylic acid represented by formula 8d can be carried out by treatment with an acid such as hydrochloric acid or 2,4,6-trichlorophenol  
5 at reflux to give the carboxylic acid represented by formula 8e.

The esterification of the compound represented by formula 8e can be conducted in a solvent such as methanol or ethanol in the presence of acid such as hydrochloric acid, sulfuric acid or p-toluenesulfonic acid at refluxing temperature to give the ester compound represented by formula 8f.

10 The ester compound represented by formula 8f can be converted to the compound represented by formula (V-I) by treatment with trifluoromethanesulfonic anhydride, N-phenyltrifluoromethanesulfonimide, 2-[N,N-bis(trifluoromethanesulfonyl)amino]pyridine, p-toluenesulfonyl chloride, phosphorous oxychloride, phosphorous oxybromide at a temperature from about 0 °C to refluxing  
15 temperature in a solvent such as dimethylformamide or tetrahydrofuran or solvent-free in the presence of a base such as triethylamine or diisopropylethylamine or without base.

Suzuki coupling between a compound represented by formula (V-I) and an appropriately functionalized aryl boronic acid or ester compound represented by formula (VI) in the presence of a base such as anhydrous cesium carbonate, cesium  
20 fluoride, sodium carbonate or potassium phosphate in a solvent such as 1,4-dioxane, dimethylformamide or dimethylsulfoxide using a catalyst such as tetrakis(triphenylphosphine)palladium(0), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) chloride, palladium(II) acetate or bis(dibenzylidenacetone)palladium(0), with or without a phosphine ligand such as triphenylphosphine, tri-t-butylphosphine or  
25 1,1'-bis(diphenylphosphino)ferrocene at a temperature from about 70 °C to the refluxing temperature provided the compounds represented by formula (III-A-I).

In cases where suitably substituted boronic acids or esters are not commercially available, the 4,4,5,5-tetramethyl-[1,3,2]dioxaborolane intermediate can be prepared from the corresponding aryl halide or aryl triflate by a palladium mediated coupling  
30 with a diboron species such as bis(pinacolato)diboron using the method of Ishiyama, T. *et al. (J. Org. Chem. 1995, 60(23), 7508)*. Alternatively, the corresponding boronic acid can be prepared by metal-halogen exchange of the aryl/heteroaryl halide, quenching with a trialkoxyborate reagent and aqueous workup to provide the boronic acids

(Miyaura, N.; Suzuki, A. *Chem. Review*, **1995**, *95*, 2457). Alternatively, a compound represented by formula (V-I) can be converted to the corresponding 4,4,5,5-tetramethyl-[1,3,2]dioxaborolane intermediate compound represented by formula (VII-I) by the same methods described above.


- 5 Suzuki coupling between a compound represented by formula (VII-I) and an appropriately functionalized aryl halide or aryl triflate compound represented by formula (VIII) provided the compounds represented by formula (III-A-I).

The compound represented by formula (III-A-I) can be converted to the compound represented by formula (II-A-I) by the deprotection reaction as described  
10 above.

- 9) The compound represented by formula (II-A) and formula (III-A) wherein Y is N and U is CH<sub>2</sub>, that is, a compound represented by formula (II-A-II) and formula (III-A-II) respectively, can be prepared as outlined in Reaction Scheme 9:

15

#### Reaction Scheme 9



wherein X<sub>1</sub> represents hydroxyl, chlorine or -O-C1-4 alkyl and the other symbols have the same meaning described above.

- In Reaction Scheme 9, treatment of an imidate compound represented by  
20 formula 8a with an ammonium salt such as ammonium chloride in a solvent such as

ethanol at a temperature from about 20 °C to the refluxing temperature provides an amidine compound represented by formula 9a.

The amidine compound represented by formula 9a can be condensed with malonic acid derivatives represented by formula 9b such as mono-ethyl malonate or ethyl malonyl chloride by the same methods as described above for the amidation reaction to give the acylated amidine intermediates, which then cyclize to provide a pyrimidinone compound represented by formula 9c.


The pyrimidinone compound represented by formula 9c can be converted to the compound represented by formula (II-A-II) or formula (III-A-II) by the same method described in Reaction Scheme 8.

Alternatively, the amidine compound represented by formula 9a can be condensed with a suitably substituted beta-ketoester compound represented by formula 9d in the presence of base such as triethylamine in a solvent such as toluene at a temperature from about 20 °C to the refluxing temperature to give the compound represented by formula (III-A-II).

The compound represented by formula (III-A-II) can be converted to the compound represented by formula (II-A-II) by the deprotection reaction as described above.

10) The compound represented by formula (II-A) and formula (III-A) wherein Y is C(H) and U is S, that is, a compound represented by formula (II-A-III) and formula (III-A-III) respectively, can be prepared as outlined in Reaction Scheme 10:

#### Reaction Scheme 10




wherein all symbols have the same meaning described above.

In Reaction Scheme 10, the compound represented by formula (III-A-III) can be prepared from acylated Meldrum's acid derivatives represented by formula 10b, which are prepared from the compound represented by formula 10a using the method of Hans, E. *et al.* (*J. Org. Chem.* **2001**, *66*, 6756).

The compound represented by formula (III-A-III) can be converted to the compound represented by formula (II-A-III) by the deprotection reaction as described


above.

11) The compound represented by formula (I-A) wherein  $R^{62}$  is hydrogen, Y is C(H) or N, U is  $CH_2$  or S and at least one of the  $R^6$  is  $Cyc^{10}$ , that is, a compound represented by formula (I-A-IV):



wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 11:

### Reaction Scheme 11



wherein  $R^{6-2}$  represents amine, fluorine, chlorine, bromine or iodine and the other symbols have the same meanings as described above.


In Reaction Scheme 11,  $Cyc^{10}$  can be constructed at the appropriate synthetic

stage by Cyc<sup>10</sup> formation reaction. The Cyc<sup>10</sup> formation reaction can be prepared by the method described in Reaction Schemes 12 to 16 described below.

The reactions such as Suzuki coupling, Cyc<sup>1</sup> formation, etc. described in Reaction Scheme 11 can be prepared by the same methods as described above.


5

12) The compound represented by formula (I-A-IV) wherein Cyc<sup>2</sup> is aryl or heteroaryl and Cyc<sup>10</sup> is tetrazole ring which is attached to Cyc<sup>2</sup> at the 1-position, that is, a compound represented by formula (I-A-IV-I):



10 wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 12:

Reaction Scheme 12



wherein all symbols have the same meaning described above.

15 In Reaction Scheme 12, an appropriately substituted aryl or heteroaryl amine represented by formula (VIII-a-I) can be converted to tetrazole compound represented by formula (VIII-b-I) by treatment with sodium azide and trimethylorthoformate or triethylorthoformate in acetic acid at a temperature from about 0 °C to 95 °C.


20 The compound represented by formula (VIII-b-I) can be converted to the compound represented by formula (I-A-IV-I) by the same method as described in Reaction Scheme 11.

Alternatively, the tetrazole ring formation can be conducted at the appropriate synthetic stage using an intermediate having an aryl or heteroaryl amino group as shown in Reaction Scheme 11.

25




13) The compound represented by formula (I-A-IV) wherein Cyc<sup>2</sup> is aryl or heteroaryl and Cyc<sup>10</sup> is a 1,2,3-triazole ring which is attached to Cyc<sup>2</sup> at the 1-position, that is, a compound represented by formula (I-A-IV-II):



- 5 wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 13:

Reaction Scheme 13



wherein all symbols have the same meaning described above.

- 10 In Reaction Scheme 13, an appropriately substituted aryl or heteroaryl amine compound represented by formula (VIII-a-I) can be treated with sodium nitrite in the presence of acid such as hydrochloric acid in water to produce the corresponding diazonium salt, which can be treated with sodium azide in water to form the corresponding azide compound represented by formula (VIII-a-II). Alternatively, an
- 15 appropriately substituted aryl or heteroaryl amine compound represented by formula (VIII-a-I) can be converted to the corresponding azide compound represented by formula (VIII-a-II) treatment with trimethylsilyl azide and tert-butyl nitrite in acetonitrile at a temperature from about 0 °C to 40 °C.

The azide compound represented by formula (VIII-a-II) can be treated with the


appropriately substituted alkyne in a solvent such as toluene at the refluxing temperature to give 1,2,3-triazole derivatives represented by formula (VIII-b-II).

The compound represented by formula (VIII-b-II) can be converted to the compound represented by formula (I-A-IV-II) by the same method as described in

# 5 Reaction Scheme 11.


Alternatively, 1,2,3-triazole ring formation can be conducted at the appropriate synthetic stage using an intermediate having an aryl or heteroaryl amino group as shown in Reaction Scheme 11.

- 10 14) The compound represented by formula (I-A-IV) wherein  $\text{Cyc}^2$  is aryl or heteroaryl and  $\text{Cyc}^{10}$  is 4-chloro-1,2,3-triazole attached to  $\text{Cyc}^2$  at the 1-position, that is, a compound represented by formula (I-A-IV-III):



- 15 wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 14:

Reaction Scheme 14



wherein all symbols have the same meaning described above.


In Reaction Scheme 14, an appropriately substituted aryl or heteroaryl azide represented by formula (VIII-a-II) can be treated with ethynyl-tri-n-butyltin in a solvent such as toluene at reflux to give the 4-tributylstannyl-1,2,3-triazole represented by formula (VIII-b-III).

The compound represented by formula (VIII-b-III-I) can be prepared from the precursor compound represented by formula (VIII-b-III) by treatment with N-chlorosuccinimide or 1,3-dichloro-5,5-dimethyl-hydantoin in a solvent such as acetonitrile at a temperature from approximately 20 °C to reflux.

The compound represented by formula (VIII-b-III-I) can be converted to the compound represented by formula (I-A-IV-III) by the same method as described in Reaction Scheme 11.


Alternatively, 4-Chloro-1,2,3-triazole ring formation can be conducted at the appropriate synthetic stage using an intermediate having an aryl or heteroaryl amino group as shown in Reaction Scheme 11.

The compound represented by formula (I-A-IV) wherein Cyc<sup>2</sup> is aryl or heteroaryl and Cyc<sup>10</sup> is a 1,2-pyrazole which is attached to Cyc<sup>2</sup> at the 1-position, that is, the compound represented by formula (I-A-IV-IV):



wherein R<sup>A-15</sup> represents hydrogen or C1-4 alkyl and the other symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 15:

#### Reaction Scheme 15




wherein all symbols have the same meaning described above.

In Reaction Scheme 15, an appropriately substituted aryl or heteroaryl fluoride represented by formula (VIII-a-IV) can be converted to the 1,2-pyrazole compound represented by formula (VIII-b-IV) by treatment with the appropriately substituted 1, 2-pyrazole in the presence of a base such as cesium carbonate in a solvent such as N,N-dimethylacetamide at a temperature from approximately 20 °C to 100 °C.


The compound represented by formula (VIII-b-IV) can be converted to the compound represented by formula (I-A-IV-IV) by the same method as described in Reaction Scheme 11.

- 16) The compound represented by formula (I-A-IV) wherein Cyc<sup>10</sup> represents 4-fluoro-1,2-pyrazole, that is, a compound represented by formula (I-A-IV-V):



wherein all symbols have the same meaning described above, can be prepared as outlined in Reaction Scheme 16:

- Reaction Scheme 16



wherein all symbols have the same meaning described above.


- In Reaction Scheme 16, the compound represented by formula (VIII-b-IV-I) can be converted to the 4-fluoro-1,2-pyrazole compound represented by formula (VIII-b-IV-II) by treatment with a fluorinating agent such as 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate) in a solvent such as acetonitrile at a temperature from about 20 °C to 100 °C.

The compound represented by formula (VIII-b-IV-II) can be converted to the

compound represented by formula (I-A-IV-V) by the same method as described in Reaction Scheme 11.

17) The compound of formula (I-B) can be prepared as outlined in Reaction Scheme 17:

### Reaction Scheme 17



wherein all symbols have the same meaning described above.

In Reaction Scheme 17, an appropriately substituted 4-hydroxy-2-pyrone derivative 17a can be treated with a suitably substituted amino acid derivative 17b in the presence of alkali such as sodium hydroxide or potassium hydroxide in water at a temperature from about 20 °C to reflux to give the carboxylic acid compound represented by formula 17c.


The carboxylic acid represented by formula 17c can be converted to the corresponding ester compound represented by formula (V-B-1) by ester formation and leaving group formation as described previously. The ester compound represented by

formula (V-B-1) can be converted to the compound represented by the formula (I-B) by the method as described above.

18) The compound of formula (I-C) can be prepared as outlined in Reaction Scheme 18:

5

### Reaction Scheme 18



wherein all symbols have the same meaning described above.

In Reaction Scheme 18, appropriately substituted boronic ester derivative represented by formula (VI-a), which are commercially available or can be prepared by the method described above, can be converted to a 4-aryl-6-chloropyrimidine compound represented by formula 18a using the 2,4-dichloropyrimidine derivative represented by formula IX by a Suzuki coupling reaction as described above.

4-Aryl-6-chloropyrimidine compounds represented by formula 18a can be hydrolyzed to 4-aryl-pyrimidinone compounds represented by formula 18b by treatment with an acid such as hydrochloric acid at a temperature from approximately 20 °C to reflux.




4-Aryl-pyrimidinone compounds represented by formula 18b can be converted

to the ester compounds represented by formula (III-C-I) by treatment with alkylating agents represented by the formula 18c by the same method as described above.

The ester compound represented by formula (III-C-I) can be converted to the compound represented by the formula (III-C-II) by a  $\text{Cyc}^{10-\text{C}}$  formation reaction. The  $\text{Cyc}^{10-\text{C}}$  formation reaction can be carried out by the same method as described previously.


The compound represented by formula (III-C-II) can be converted to the compound represented by formula (I-C) by the same method as described above for the  $\text{Cyc}^1$  formation reaction.

19) The compound represented by formula (I) wherein Y represents  $\text{C}(\text{R}^5)$  and at least one of the  $\text{R}^5$  or  $\text{R}^{62}$  represent halogen, that is, a compound represented by formula (I-II-1), (I-II-2) or (I-II-3):



wherein  $\text{R}^{5-1}$  and  $\text{R}^{62-1}$  represent chlorine or bromine and the other symbols have the same meanings as described above, can be prepared as outlined in Reaction Scheme 19:

#### Reaction Scheme 19



In Reaction Scheme 19, the reaction from the compound represented by formula (III) to the compound represented by formula (III-II-1), (III-II-2) and (III-II-3) is a halogenation reaction.

- 5 The halogenation reaction is well known. For example, the reaction of the compound represented by formula (III) with brominating or chlorinating agent, such as N-bromosuccinimide, N-chlorosuccinimide or 1,3-dichloro-5,5-dimethyl-hydantoin in a suitable solvent such as acetonitrile, chloroform or tetrahydrofuran from -20 °C to the refluxing temperature provides the compound represented by formula (III-II-1), (III-II-2) and (III-II-3). The carboxylic acid represented by formula (III-II-1), (III-II-2) and (III-II-3) can be converted the compound represented by formula (I-II-1), (I-II-2) and (I-II-3) respectively by the method as described above.

The compounds of the present invention can be prepared by the reactions or modified variants of the reactions described above.

- 15 Other starting compounds or compounds used as reagents are known compounds which can be prepared easily by a combination of known methods, for example, the methods described in Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition (Richard C. Larock, John Wiley & Sons Inc, 1999) or Elmer J. Rauckman et al., J. Org. Chem., vol.41, No.3, 1976, p564-565 etc.

- 20 In each reaction of the specification, the reactions with heating, as will be apparent to those skilled in the art, may be carried out using a water bath, an oil bath, a sand bath, a heating block or by microwave.

In each reaction of the specification, a solid phase reagent may be used which is



supported by a polymer (for example, polystyrene, polyacrylamide, polypropylene or polyethyleneglycol etc.).

In each reaction of the specification, the products obtained may be purified by conventional techniques. For example, the purification may be carried out by distillation at atmospheric or reduced pressure, by high performance liquid chromatography with silica gel or magnesium silicate, by thin layer chromatography, by ion-exchange resin, by scavenger resin, by column chromatography, by washing, trituration or recrystallization. The purification may be carried out after each reaction stage or after several reaction stages.

In a reaction of the specification where polystyrene resin is used, the obtained products may be purified by conventional techniques. For example, the purification may be carried out by multiple washing with a solvent (for example, dimethylformamide, dichloromethane, methanol, tetrahydrofuran, toluene, acetic acid/toluene, etc.).

#### TOXICITY:

The toxicity of the compound represented by formula (I), the salt thereof, the N-oxide thereof or the solvate thereof, or the prodrug thereof is very low and therefore it may be considered safe for pharmaceutical use.

#### APPLICATION TO PHARMACEUTICALS:

The compounds of the present invention are therapeutically useful. The present invention therefore provides a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof, for use in the treatment of the human or animal body by therapy.

Also provided is a pharmaceutical composition comprising a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof, and a pharmaceutically acceptable carrier or diluent.

Said pharmaceutical composition typically contains up to 85 wt% of a compound of the invention. More typically, it contains up to 50 wt% of a compound of the invention. Preferred pharmaceutical compositions are sterile and pyrogen free.

Further, the pharmaceutical compositions provided by the invention typically contain a compound of the invention which is a substantially pure optical isomer.

5 The compounds of the present invention may normally be administered systemically or locally, usually by oral or parenteral administration.

A therapeutically effective amount of a compound of the invention is administered to a patient. The doses to be administered are determined depending upon, for example, age, body weight, symptom, the desired therapeutic effect, the route of  
10 administration, and the duration of the treatment. In the human adult, the doses per person are generally from 1 mg to 1000 mg, by oral administration, up to several times per day, and from 1 mg to 100 mg, by parenteral administration (preferably intravenous administration), up to several times per day, or continuous administration from 1 to 24 hours per day from vein.

15 As mentioned above, the doses to be used depend upon various conditions. Therefore, there are cases in which doses lower than or greater than the ranges specified above may be used.

The compounds or pharmaceutical compositions of the present invention may be administered for example, in the form of a solid for oral administration, liquid forms for  
20 oral administration, injections, liniments or suppositories for parenteral administration. Preferably, the compounds or pharmaceutical compositions of the present invention are administered orally.

Solid forms for oral administration include compressed tablets, pills, capsules, dispersible powders, and granules. Capsules include hard capsules and soft capsules.

25 In such solid forms, one or more of the active compound(s) may be admixed with vehicles (such as lactose, mannitol, glucose, microcrystalline cellulose or starch), binders (such as hydroxypropyl cellulose, polyvinylpyrrolidone or magnesium metasilicate aluminate), disintegrants (such as cellulose calcium glycolate), lubricants (such as magnesium stearate), stabilizing agents, solution adjuvants (such as glutamic  
30 acid or aspartic acid, disaggregating agents, e.g. starch, alginic acid, alginates or sodium starch glycolate; effervescent mixtures, dyestuffs, sweeteners, wetting agents, such as lecithin, polysorbates, laurylsulphates; and, in general, non toxic and pharmacologically inactive substances used in pharmaceutical formulations, and prepared according to

methods well known in normal pharmaceutical practice, for example, by means of mixing, granulating, tableting, sugar coating, or film coating processes. The solid forms may, if desired, be coated with coating agents (such as sugar, gelatin, hydroxypropyl cellulose or hydroxypropylmethyl cellulose phthalate), or be coated with two or more  
5 films. Furthermore, coating may include containment within capsules of absorbable materials such as gelatin.

Liquid forms for oral administration include pharmaceutically acceptable solutions, suspensions, emulsions, syrups and elixirs. In such forms, one or more of the active compound(s) may be dissolved, suspended or emulsified into diluent(s)  
10 commonly used in the art (such as purified water, ethanol or a mixture thereof). Besides such liquid forms may also comprise some additives, such as wetting agents, suspending agents, emulsifying agents, sweetening agents, flavoring agents, aroma, preservative or buffering agent. The syrups may contain as carriers, for example, saccharose or saccharose with glycerine and/or mannitol and/or sorbitol.

Suspensions and emulsions may contain as carrier, for example a natural gum, agar, sodium alginate, pectin, methylcellulose, carboxymethylcellulose, or polyvinyl alcohol. The suspension or solutions for intramuscular injections may contain, together with the active compound, a pharmaceutically acceptable carrier, e.g. sterile water, olive oil, ethyl oleate, glycols, e.g. propylene glycol, and if desired, a suitable amount of  
20 lidocaine hydrochloride.

Solutions for injection or infusion may contain as carrier, for example, sterile water or preferably they may be in the form of sterile, aqueous, isotonic saline solutions.

Injections for parenteral administration include sterile aqueous, suspensions, emulsions and solid forms which are dissolved or suspended into solvent(s) for injection  
25 immediately before use. In injections, one or more of the active compound(s) may be dissolved, suspended or emulsified into solvent(s). The solvents may include distilled water for injection, saline, vegetable oil, propylene glycol, polyethylene glycol, alcohol such as ethanol, or a mixture thereof. Injections may comprise some additives, such as stabilizing agents, solution adjuvants (such as glutamic acid, aspartic acid or  
30 POLYSORBATE80 (registered trade mark)), suspending agents, emulsifying agents, soothing agent, buffering agents, preservative. They may be sterilized at a final step, or may be prepared according to sterile methods. They may also be manufactured in the

form of sterile solid forms such as freeze-dried products, which may be dissolved in sterile water or some other sterile diluent(s) for injection immediately before use.

Other forms for parenteral administration include liquids for external use, ointments and endermic liniments, inhalations, sprays, suppositories and vaginal  
5 suppositories which comprise one or more of the active compound(s) and may be prepared by methods known per se.

Sprays may comprise additional substances other than diluents, used commonly, stabilizers such as sodium hydrogensulfite and buffers capable of imparting isotonicity, for example, isotonic buffers such as sodium chloride, sodium citrate or citric acid.

10

#### EFFECT OF THE INVENTION:

The compounds of the present invention represented by formula (I) act as potent and selective inhibitors of Factor XIa, with potent anticoagulant activity and/or good oral availability. In particular, the compounds of the present invention act as a Factor  
15 XIa inhibitor or a Factor XIa and plasma kallikrein dual inhibitor. Thus the compounds of the present invention are useful in preventing and/or treating thromboembolic diseases. One advantage of the compounds of the present invention is that they can provide high inhibitory activity against FXIa and potent anticoagulant activity and/or high plasma exposure level after oral administration.

20

The present invention therefore provides a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof, for use in treating or preventing a thromboembolic disease. Also provided is a method for treating a patient suffering from or susceptible to a  
25 thromboembolic disease, which method comprises administering to said patient an effective amount of a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof. Further provided is the use of a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a  
30 prodrug thereof, in the manufacture of a medicament for use in treating or preventing a thromboembolic disease.

The thromboembolic disease may be, for example, selected from the group consisting of arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, arterial cerebrovascular thromboembolic disorders, venous cerebrovascular thromboembolic disorders and thromboembolic disorders in the  
5 chambers of the heart or in the peripheral circulation.

More specifically, arterial cardiovascular thromboembolic disorders may be exemplified by coronary artery disease, ischemic cardiomyopathy, acute coronary syndrome, coronary arterial thrombosis, ischemic complications of unstable angina and non-Q-wave myocardial infarction, acute ST-segment elevation myocardial infarction  
10 managed medically or with subsequent percutaneous coronary intervention, angina pectoris such as stable effort angina pectoris, variant angina pectoris, unstable angina pectoris, myocardial infarction (e.g. first myocardial infarction or recurrent myocardial infarction), acute myocardial infarction, reocclusion and restenosis after coronary artery bypass surgery, reocclusion and restenosis after percutaneous transluminal cardiac  
15 angioplasty/ transluminal coronary artery stent placement surgery or after thrombolytic therapy for coronary artery, ischemic sudden death. Venous cardiovascular thromboembolic disorders may be exemplified by deep vein thrombosis (DVT) and/or pulmonary embolism (PE) in major general surgery, abdominal surgery, hip replacement surgery, knee replacement surgery, hip fracture surgery, multiple fracture,  
20 multiple injury, trauma, spinal cord injury, burns, critical care unit, DVT and/or PE in medical patients with severely restricted mobility during acute illness, DVT and/or PE in patients with cancer chemotherapy, DVT and/or PE in patients with stroke, symptomatic or asymptomatic DVT with or without PE, pulmonary embolism. Arterial cerebrovascular thromboembolic disorders may be exemplified by stroke, ischemic  
25 stroke, acute stroke, stroke in patients with non-valvular or valvular atrial fibrillation, cerebral arterial thrombosis, cerebral infarction, transient ischemic attack (TIA), lacuna infarction, atherosclerotic thrombotic cerebral infarction, cerebral artery embolism, cerebral thrombosis, cerebrovascular disorder, asymptomatic cerebral infarction.

Venous cerebrovascular thromboembolic disorders may be exemplified by  
30 intracranial venous thrombosis, cerebral embolism, cerebral thrombosis, sinus thrombosis, intracranial venous sinus thrombosis, cavernous sinus thrombosis. Thromboembolic disorders in the chambers of the heart or in the peripheral circulation may be exemplified by venous thrombosis, systemic venous thromboembolism,

thrombophlebitis, non-valvular or valvular atrial fibrillation, disseminated intravascular coagulopathy (DIC), kidney embolism, atherosclerosis, atherothrombosis, peripheral artery occlusive disease (PAOD), peripheral arterial disease, arterial embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface (such as catheters, stents, artificial heart valves, or hemodialyzer) that promotes thrombosis.

Preferably, the thromboembolic disorder is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial infarction (e.g. first myocardial infarction or recurrent myocardial infarction), ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface (such as catheters, stents or artificial heart valves) that promotes thrombosis.

The compounds of the present invention may also be administered in combination with one or more further therapeutic agents. Thus, in another embodiment, the present invention provides a method for treating a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a first and second therapeutic agent, wherein the first therapeutic agent is a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof, and the second therapeutic agent is at least one agent selected from a second factor XIa inhibitor, an anti-coagulant agent, an anti-platelet agent, a thrombin inhibiting agent, a thrombolytic agent, and a fibrinolytic agent. Preferably, the second therapeutic agent is at least one agent selected from warfarin, unfractionated heparin, low molecular weight heparin (e.g., danaparoid, enoxaparin, bemiparin, dalteparin, nadroparin, parnaparin, reviparin, tinzaparin, ardeparin, adomiparin, semuloparin, RO-14, RO-16, GCC-4401C), synthetic pentasaccharide (e.g., fondaparinux, idraparin, idrabiotaparin, EP-42675, EP-217609, EP-224283), synthetic hexadecasaccharide (e.g., SSR128428), factor Xa inhibitor (e.g., rivaroxaban, edoxaban, apixaban, betrixaban, otamixaban, darexaban, letaxaban, 813893, eribaxaban, AVE-3247, R-1663, BMS-344577, TAK-239),

thrombin inhibitor (argatroban, melagatran, ximelagatran, dabigatran, tanogitrin, dermatan, hirudin, bivalirudin, desirudin, lepirudin, NU-172, DP-4088, RWJ-671818, BL-5030), antithrombin III (e.g., freeze-dried concentrated human antithrombin III, KW-3357), thrombomodulin (e.g., thrombomodulin alfa), non-steroidal anti-

5 inflammatory drugs (e.g., aspirin, acetaminophen, codeine, ibuprofen, naproxen, sulindac, indomethacin, mefenamate, droxicam, indobufen, diclofenac, sulfinpyrazone, piroxicam, fentanyl, ketorolac, mefenamate, morphine, phenacetin, sufentanyl), P2Y<sub>12</sub> receptor antagonist (e.g., ticlopidine, clopidogrel, prasugrel, ticagrelor, elinogrel), phosphodiesterase-III inhibitor (e.g., dipyridamole, cilostazol), phosphodiesterase-V

10 inhibitor (e.g., sildenafil) serotonin 2 antagonist (e.g., sarpogrelate), prostaglandin E<sub>1</sub> agonist (e.g., alprostadil, limaprost, ecraprost), prostaglandin I<sub>2</sub> agonist (e.g., ibudilast, iloprost, beraprost, epoprostenol), thromboxane A synthesis inhibitor (e.g., ozagrel), thromboxane A<sub>2</sub> receptor antagonist (e.g., seratrovast, ramatroban, terutroban, ifetroban), glycoprotein IIb/IIIa blocker (e.g., tirofiban, eptifibatide, abciximab,

15 integrilin), protease-activated receptor-1 antagonist (e.g., vorapaxar) and fibrinolytic agent (e.g., tissue plasminogen activator, anistreplase, alteplase, reteplase, tenecteplase, desmoteplase, amideplase, THR-100, urokinase, ocriplasmin, nasaruplase  $\beta$ , streptokinase). Preferably, the second therapeutic agent is at least one anti-platelet agent. Preferably, the anti-platelet agent(s) are ticlopidine, clopidogrel, prasugrel

20 ticagrelor, elinogrel, cilostazol, sarpogrelate, iloprost, dipyridamole, beraprost, limaprost, ozagrel, vorapaxar and/or aspirin, or a combination thereof. The present invention also provides a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof, in combination with a second therapeutic agent selected from those

25 listed above, for use in treating or preventing a thromboembolic disease. The present invention also provides the use of a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof, in combination with a second therapeutic agent, in the manufacture of a medicament for use in treating or preventing a thromboembolic disease.

30 In another embodiment, the present invention provides a pharmaceutical composition comprising a compound of formula (I), as defined above, or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a solvate thereof or a prodrug thereof and an additional therapeutic agent. Preferably, the further additional

therapeutic agent(s) are selected from potassium channel openers, potassium channel blockers, calcium channel blockers, sodium hydrogen exchanger inhibitors, sodium-potassium pump inhibitors, antiarrhythmic agents, antiatherosclerotic agents, anticoagulants, antiplatelets, antithrombotic agents, prothrombolytic agents, fibrinogen antagonists, diuretics, antihypertensive agents, ATPase inhibitors, mineralocorticoid receptor antagonists, phosphodiesterase inhibitors, antidiabetic agents, anti-inflammatory agents, antioxidants, angiogenesis modulators, antiosteoporosis agents, hormone replacement therapies, hormone receptor modulators, oral contraceptives, antiobesity agents, antidepressants, antianxiety agents, antipsychotic agents, antiproliferative agents, antitumor agents, antiulcer and gastroesophageal reflux disease agents, growth hormone agents and/or growth hormone secretagogues, thyroid mimetics, anti-infective agents, antiviral agents, antibacterial agents, antifungal agents, cholesterol/lipid lowering agents and lipid profile therapies, and agents that mimic ischemic preconditioning and/or myocardial stunning, or a combination thereof.

15 In another embodiment, the present invention provides a pharmaceutical composition further comprising additional therapeutic agent(s) selected from an anti-arrhythmic agent, an anti-hypertensive agent, an antidiabetic agent, an anti-coagulant agent, an anti-platelet agent, a thrombin inhibiting agent, a thrombolytic agent, a fibrinolytic agent, a calcium channel blocker, a potassium channel blocker, a cholesterol/lipid lowering agent, or a combination thereof.

In another embodiment, the present invention provides a pharmaceutical composition further comprising additional therapeutic agent(s) selected from warfarin, unfractionated heparin, low molecular weight heparin (e.g., danaparoid, enoxaparin, bemiparin, dalteparin, nadroparin, parnaparin, reviparin, tinzaparin, ardeparin, adomiparin, semuloparin, RO-14, RO-16, GCC-4401C), synthetic pentasaccharide (e.g., fondaparinux, idraparinux, idrabiotaparinux, EP-42675, EP-217609, EP-224283), synthetic hexadecasaccharide (e.g., SSR128428), factor Xa inhibitor (e.g., rivaroxaban, edoxaban, apixaban, betrixaban, otamixaban, darexaban, letaxaban, 813893, eribaxaban, AVE-3247, R-1663, BMS-344577, TAK-239), thrombin inhibitor (argatroban, melagatran, ximelagatran, dabigatran, tanogitrin, dermatan, hirudin, bivalirudin, desirudin, lepirudin, NU-172, DP-4088, RWJ-671818, BL-5030), antithrombin III (e.g., freeze-dried concentrated human antithrombin III, KW-3357), thrombomodulin (e.g., thrombomodulin alfa), non-steroidal anti-inflammatory drugs (e.g., aspirin,



acetaminophen, codeine, ibuprofen, naproxen, sulindac, indomethacin, mefenamate, droxicam, indobufen, diclofenac, sulfinpyrazone, piroxicam, fentanyl, ketorolac, mefenamate, morphine, phenacetin, sufentanyl), P2Y<sub>12</sub> receptor antagonist (e.g., ticlopidine, clopidogrel, prasugrel, ticagrelor, elinogrel), phosphodiesterase-III inhibitor (e.g., dipyridamole, cilostazol), phosphodiesterase-V inhibitor (e.g., sildenafil) serotonin 2 antagonist (e.g., sarpogrelate), prostaglandin E<sub>1</sub> agonist (e.g., alprostadil, limaprost, ecraprost), prostaglandin I<sub>2</sub> agonist (e.g., ibudilast, iloprost, beraprost, epoprostenol), thromboxane A synthesis inhibitor (e.g., ozagrel), thromboxane A<sub>2</sub> receptor antagonist (e.g., seratrovast, ramatroban, terutroban, ifetroban), glycoprotein IIb/IIIa blocker (e.g., tirofiban, eptifibatide, abciximab, integrilin), protease-activated receptor 1 antagonist (e.g., vorapaxar) and fibrinolytic agent (e.g., tissue plasminogen activator, anistreplase, alteplase, reteplase, tenecteplase, desmoteplase, amideplase, THR-100, urokinase, ocriplasmin, nasaruplase  $\beta$ , streptokinase). Preferably, the second therapeutic agent is at least one anti-platelet agent. Preferably, the anti-platelet agent(s) are ticlopidine, clopidogrel, prasugrel, ticagrelor, elinogrel, cilostazol, sarpogrelate, iloprost, dipyridamole, beraprost, limaprost, ozagrel, vorapaxar and/or aspirin, or a combination thereof.

In a preferred embodiment, the present invention provides a pharmaceutical composition wherein the additional therapeutic agent is an antihypertensive agent selected from angiotensin-converting enzyme inhibitors, angiotensin-1 (AT-1) receptor antagonists, beta- adrenergic receptor antagonists, endothelin A (ETA) receptor antagonists, dual ETA/AT-1 receptor antagonists, and vasopepsidase inhibitors, an antiarrhythmic agent selected from IK<sub>ur</sub> inhibitors, an anticoagulant selected from thrombin inhibitors, antithrombin-III activators, heparin co-factor II activators, other factor XIa inhibitors, other kallikrein inhibitors, plasminogen activator inhibitor (PAI-1) inhibitors, thrombin activatable fibrinolysis inhibitor (TAFI) inhibitors, factor VIIa inhibitors, factor IXa inhibitors, factor XIIIa inhibitors and factor Xa inhibitors, or an anti-platelet agent selected from glycoprotein IIb/IIIa blockers, protease-activated receptor-1 antagonists, protease-activated receptor-4 antagonists, phosphodiesterase-III inhibitors, phosphodiesterase-V inhibitors, prostaglandin E<sub>1</sub> agonists, prostaglandin I<sub>2</sub> agonists, thromboxane A synthesis inhibitors, thromboxane A<sub>2</sub> receptor antagonists, P2Y<sub>1</sub> receptor antagonists, P2Y<sub>12</sub> receptor antagonists, thromboxane receptor antagonists, cyclooxygenase-1 inhibitors, glycoprotein VI antagonists, glycoprotein Ib

antagonists, Growth arrest-specific gene 6 product antagonists and aspirin, or a combination thereof.

In a preferred embodiment, the present invention provides a pharmaceutical composition, wherein the additional therapeutic agent(s) are an anti-platelet agent or a combination thereof.

#### BEST MODE FOR CARRYING OUT THE INVENTION


The present invention is illustrated by the following Examples and biological Examples, but it is not limited thereto.

- 10 The run time, solvents and column conditions used in the LC/MS analysis of the following Examples is reported using a superscript a, b, c, d, e or f appended to the analytical results which corresponds to the following conditions:
- 15 a. 3 minute run time; 0.1% formic acid in water and 0.1% formic acid in acetonitrile as mobile phases, Waters Atlantis® dC18, 2.1 mm x 50 mm, 3 µm column
  - b. 7 minute run time; 0.1% formic acid in water and 0.1% formic acid in acetonitrile as mobile phases, Waters Atlantis® dC18, 2.1 mm x 100 mm, 3 µm column
  - 20 c. 4.5 minute run time; 0.1% formic acid in water and 0.1% formic acid in acetonitrile as mobile phases, Waters Atlantis® dC18, 3 mm x 50 mm, 3 µm column
  - d. 6 minute run time; 0.1% trifluoroacetic acid in water and 0.1% trifluoroacetic acid in acetonitrile as mobile phases, Waters Xterra® MS C18, 3 mm x 50 mm, 5 µm column
  - 25 e. 5.5 minute run time; 0.1% formic acid in water and 0.1% formic acid in acetonitrile as mobile phases, Phenomenex® Kinetex-XB® C18, 2.1 mm x 100 mm, 1.7 µm column
  - f. 1.5 minute run time; 0.1% trifluoroacetic acid in water and 0.1% trifluoroacetic acid in acetonitrile as mobile phases, Waters ACQUITY UPLC®
  - 30 BEH C18, 2.1 mm x 30 mm, 1.7 µm column
  - g. 7 minute run time; 2 mM ammonium bicarbonate in water as mobile phase A and acetonitrile as mobile phase B, Phenomenex® Gemini® C18, 2.0 mm x 100 mm, 3 µm column

The solvents in the parentheses described in chromatographic separation and TLC show the eluting or developing solvents, and the ratios of the solvents used are given as percentage mixtures in chromatographic separations or TLC. Where a compound is described as dried either anhydrous magnesium or sodium sulphate was used. The solvents in the parentheses in NMR show the solvents used in measurement. DMSO represents dimethylsulfoxide;  $\text{CDCl}_3$  represents deuterated chloroform. The following abbreviations are used in reporting the  $^1\text{H}$  NMR spectra: s (singlet), d (doublet), t (triplet), q (quartet), br. (broad), app. (apparent), obs. (obscured).


Including compounds in the following Examples, compounds used in the present specification were commonly named using a computer program capable of naming in accordance with IUPAC rules; ACD/Name® manufactured by Advanced Chemistry Development Inc., JChem for Excel or MarvinSketch manufactured by ChemAxon Ltd., or IUPAC nomenclature. In each of the following Examples, the name of the objective compound of the Example is described subsequently to the number of the Example, and the compound is sometimes referred to as the "title compound".

Example 1: ethyl (2*S*)-5-methoxy-3,4-dihydro-2*H*-pyrrole-2-carboxylate



To a dichloromethane (500 mL) solution of ethyl 5-oxo-L-prolinate (51.2 g) was added trimethyloxonium tetrafluoroborate (50.5 g) and the mixture was stirred at room temperature for 3 hours. To the cooled (0 °C) reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (350 mL) and water (50 mL) was added sequentially followed by extraction with dichloromethane. The combined organic layers were dried and concentrated giving the title compound having the following physical properties (42.2 g). TLC: Rf 0.50 (ethyl acetate).

Example 2: 3-ethyl 8-methyl (3*S*)-7-hydroxy-5-oxo-1,2,3,5-tetrahydroindolizine-3,8-dicarboxylate



The compound prepared in Example 1 (42.2 g) was combined with dimethyl 3-oxopentanedioate (42.9 g) and triethylamine (2.40 mL) and the mixture was stirred at 120 °C for 3 hours. On cooling, this gave the crude title compound having the following physical properties (85.6 g).

LC/MS  $t_R$  1.55 minutes; MS (ES<sup>+</sup>)  $m/z$  282 (M+H)<sup>a</sup>.


Example 3: (3S)-7-hydroxy-5-oxo-1,2,3,5-tetrahydroindolizine-3,8-dicarboxylic acid



To the compound prepared in Example 2 (150.9 g), 2 M sodium hydroxide (1130 mL) was added and the mixture was stirred overnight at room temperature. To the reaction mixture, an aqueous solution of 6 M hydrochloric acid was added and the solid removed by filtration. The filter cake was washed sequentially with water and dichloromethane then dried under vacuum. The filtrate was concentrated to 850 mL and the solid removed by filtration. Washing and drying of the filter cake as detailed above gave the title compound in two batches having the following physical properties (31.7 g).

LC/MS  $t_R$  0.50 minutes; MS (ES<sup>+</sup>)  $m/z$  240 (M+H)<sup>a</sup>.

Example 4: (3S)-7-hydroxy-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid




To the compound prepared in Example 3 (31.7 g), 6 M hydrochloric acid (1140 mL) and 12 M hydrochloric acid (200 mL) were added sequentially and the mixture

stirred at 140 °C for 20 hours. The reaction mixture was concentrated and the solid azeotroped with toluene to dryness giving the title compound having the following physical properties (29.1 g).

LC/MS  $t_R$  0.63 minutes; MS (ES<sup>+</sup>)  $m/z$  196 (M+H)<sup>a</sup>.

5

Example 5: ethyl (3*S*)-7-hydroxy-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate




To an ethanol (115 mL) suspension of the compound prepared in Example 4 (29 g), concentrated sulfuric acid (0.96 mL) was added and the mixture was stirred at 100 °C for 4 hours. The reaction mixture was concentrated to give the title compound having the following physical properties (29.4 g).

10

LC/MS  $t_R$  1.21 minutes; MS (ES<sup>+</sup>)  $m/z$  224 (M+H)<sup>a</sup>.

Example 6: ethyl (3*S*)-5-oxo-7-[[[(trifluoromethyl)sulfonyl]oxy]-1,2,3,5-tetrahydroindolizine-3-carboxylate

15




To an N,N-dimethylformamide (190 mL) solution of the compound prepared in Example 5 (29.4 g), triethylamine (21.2 mL) and 1,1,1-trifluoro-N-phenyl-N-[[[(trifluoromethyl)sulfonyl]methanesulfonamide (49.4 g) were added sequentially and the mixture stirred at room temperature for 1 hour. To the reaction mixture, water (400 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (20 to 50% ethyl acetate in heptanes) to give the title compound having the following physical properties (31.3 g).

20

LC/MS  $t_R$  1.89 minutes; MS (ES<sup>+</sup>)  $m/z$  356 (M+H)<sup>a</sup>.

25


Example 7: ethyl (3*S*)-7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-

3-carboxylate

To a 1,4-dioxane (625 mL) solution of the compound prepared in Example 6 (31.3 g) were added 4-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (22.4 g) and caesium fluoride (33.5 g) under an atmosphere of nitrogen. The mixture was degassed with nitrogen, tetrakis(triphenylphosphine)palladium(0) (2.6 g) added and the reaction mixture stirred at 105 °C for 30 minutes. To the reaction mixture, water (700 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (25% to 100% ethyl acetate in heptanes, then 1 to 5% methanol in ethyl acetate) to give the title compound having the following physical properties (24.3 g).

LC/MS  $t_R$  1.87 minutes; MS ( $ES^+$ )  $m/z$  333 ( $M+H$ )<sup>a</sup>.

15 Example 8: (3S)-7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid




To a methanol (120 mL) solution of the compound prepared in Example 7 (12 g), 2 M sodium hydroxide (72 mL) was added and the mixture stirred at room temperature for 4 hours. The methanol was removed under reduced pressure and 2 M hydrochloric acid added until the solution was pH 4. The solution was extracted with ethyl acetate followed by a 1:1 mixture of 2-propanol and chloroform and the combined organic

layers were dried and concentrated to give the title compound having the following physical properties (10.1 g).

LC/MS  $t_R$  1.58 minutes; MS (ES<sup>+</sup>)  $m/z$  305 (M+H)<sup>a</sup>.

5 Example 9: (3*S*)-7-[5-chloro-2-(1*H*-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid



To a glacial acetic acid (150 mL) solution of the compound prepared in Example 8 (9.3 g) was added trimethyl orthoformate (10 mL) followed after 30 minutes by sodium azide (5.93 g). The mixture was stirred at room temperature for 16 hours. Further trimethyl orthoformate (1.7 mL) and sodium azide (1.0 g) was added and the mixture was stirred at room temperature for a further 16 hours. To the cooled (0 °C) reaction mixture, a solution of sodium nitrite (5.32 g) in water (50 mL) was added dropwise over 30 minutes and the mixture stirred at 0 °C for a further hour followed by extraction into ethyl acetate. The combined organic layers were washed with water and brine, dried and concentrated. The residue was azeotroped with toluene to give the title compound having the following physical properties (10.9 g). LC/MS  $t_R$  1.54 minutes; MS (ES<sup>+</sup>)  $m/z$  358 (M+H)<sup>a</sup>.

20 Example 10: ethyl (3*S*)-7-(2-{bis[(*tert*-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

To a tetrahydrofuran (220 mL) solution of the compound prepared in Example 7 (10 g) was sequentially added di-*tert*-butyl dicarbonate (14.7 g), triethylamine (8.2 mL) and 4-dimethylaminopyridine (0.73 g) and the mixture was stirred at 60 °C for 2 hours. To the reaction mixture, saturated aqueous potassium hydrogen sulfate solution (400 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to obtain the title compound having the

following physical properties (16.8 g).

LC/MS  $t_R$  2.29 minutes; MS ( $ES^+$ )  $m/z$  555 (M+Na), 533 (M+H), 433 (M-CO<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>a</sup>.

5 Example 11: (3*S*)-7-(2-{[(*tert*-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

To a methanol (80 mL) solution of the crude compound prepared in Example 10 (17.6 g) was added 2 M sodium hydroxide (66 mL) and the mixture was stirred at room temperature for 48 hours. The reaction mixture was concentrated and the residue  
10 dissolved in water (200 mL) followed by extraction with dichloromethane. To the aqueous layer, 2 M hydrochloric acid (66 mL) was added followed by extraction into ethyl acetate. The combined ethyl acetate layers were washed with brine, dried and concentrated to obtain the title compound having the following physical properties (12.1 g).

15 LC/MS  $t_R$  1.85 minutes; MS ( $ES^+$ )  $m/z$  405 (M+H)<sup>a</sup>.

Example 12: 7-hydroxy-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

To a stirred melt of 2,4,6-trichlorophenol (98.2 g) at 100 °C was added the compound prepared in Example 3 (29.7 g). The mixture was heated slowly to 180 °C  
20 whilst gas evolution occurred. The reaction mixture was maintained at 180 °C for 48 hours before cooling to room temperature and suspending the residue in dichloromethane (200 mL). The solid was collected by filtration and added to a second melt of 2,4,6-trichlorophenol (98.2 g) at 100 °C. The reaction mixture was stirred at 180 °C for a further 24 hours before cooling to room temperature and suspending the  
25 residue in dichloromethane (200 mL). The solid was collected by filtration to afford the title compound having the following physical properties (24.3 g).

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  10.53 (br. s, 1 H), 5.82 (s, 1 H), 5.43 (d, 1 H), 4.78 (dd, 1 H), 2.98 (dd, 2 H), 2.46 - 2.37 (m, 1 H), 2.16 - 2.05 (m, 1 H).

30 Example 13: ethyl-7-hydroxy-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

The compound prepared in Example 12 (18.9 g) was treated as detailed in Example 5 to give the title compound having the following physical properties (21.5 g).  
LC/MS  $t_R$  1.19 minutes; MS ( $ES^+$ )  $m/z$  224 (M+H)<sup>a</sup>



Example 14: ethyl-5-oxo-7-[[trifluoromethyl)sulfonyl]oxy]-1,2,3,5-tetrahydroindolizine-3-carboxylate

- The compound prepared in Example 13 (20 g) was treated as detailed in Example 6 to give the title compound having the following physical properties (31.8 g). LC/MS  $t_R$  1.89 minutes; MS (ES<sup>+</sup>)  $m/z$  356 (M+H)<sup>a</sup>.

Example 15: ethyl-7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

- The compound prepared in Example 14 (25.1 g) was treated as detailed in Example 7 to give the title compound having the following physical properties (23.8 g). LC/MS  $t_R$  1.81 minutes; MS (ES<sup>+</sup>)  $m/z$  333 (M+H)<sup>a</sup>.

Example 16: 7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

- The compound prepared in Example 15 (2.0 g) was treated as detailed in Example 8 to give the title compound having the following physical properties (1.83 g). LC/MS  $t_R$  1.56 minutes; MS (ES<sup>+</sup>)  $m/z$  305 (M+H)<sup>a</sup>.

Example 17: 7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

- The compound prepared in Example 16 (1.28 g) was treated as detailed in Example 9 to give the title compound having the following physical properties (1.17 g). LC/MS  $t_R$  1.54 minutes; MS (ES<sup>+</sup>)  $m/z$  358 (M+H)<sup>a</sup>.

Example 18: ethyl 7-(2-{bis[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

- The compound prepared in Example 15 (21.8 g) was treated as detailed in Example 10 to give the crude title compound having the following physical properties (32.9 g). LC/MS  $t_R$  2.30 minutes; MS (ES<sup>+</sup>)  $m/z$  555 (M+Na), 533 (M+H), 433 (M-CO<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>a</sup>.

Example 19: 7-(2-[[[(tert-butoxy)carbonyl]amino]-5-chlorophenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

The compound prepared in Example 18 (32.9 g) was treated as detailed in  
5 Example 11 to give the title compound having the following physical properties (20.5 g).  
LC/MS  $t_R$  1.84 minutes; MS ( $ES^+$ )  $m/z$  405 ( $M+Na$ )<sup>a</sup>.

Example 20: tert-butyl 2-[4-(benzyloxy)-2-oxo-1,2-dihydropyridin-1-yl]acetate

To a tetrahydrofuran (700 mL) solution of 4-benzyloxypyridone (25 g),  
10 potassium tert-butoxide (15 g), tetrabutylammonium bromide (2 g) and *tert*-butyl  
bromoacetate (18 mL) was sequentially added and the mixture was stirred at room  
temperature for 48 hours. To the reaction mixture, water (500 mL) was added followed  
by extraction with ethyl acetate. The combined organic layers were washed with brine,  
dried and concentrated to give the title compound having the following physical  
15 properties (35 g).  
LC/MS  $t_R$  1.96 minutes; MS ( $ES^+$ )  $m/z$  316 ( $M+H$ )<sup>a</sup>.

Example 21: tert-butyl 2-[4-hydroxy-2-oxo-1,2-dihydropyridin-1-yl]acetate

To an ethanol (300 mL) solution of the compound prepared in Example 20 (20  
20 g) was added 5% palladium-carbon (4.05 g) and the mixture was stirred under an  
atmosphere of hydrogen for 4 hours. The reaction mixture was then filtered through  
Celite® and the filtrate was concentrated to give the title compound having the  
following physical properties (14.3 g).  
LC/MS  $t_R$  1.42 minutes; MS ( $ES^+$ )  $m/z$  226 ( $M+H$ )<sup>a</sup>.

25

Example 22: tert-butyl 2-{2-oxo-4-[(trifluoromethane)sulfonyloxy]-1,2-dihydropyridin-1-yl}acetate

The compound prepared in Example 21 (14.3 g) was treated as detailed in  
Example 6 to give the title compound having the following physical properties (16.2 g).  
30 LC/MS  $t_R$  1.97 minutes; MS ( $ES^+$ )  $m/z$  301 ( $M-C(CH_3)_3+H$ )<sup>a</sup>.

Example 23: tert-butyl 2-[4-(2-amino-5-chlorophenyl)-2-oxo-1,2-dihydropyridin-1-yl]acetate

The compound prepared in Example 22 (6.4 g) was treated as detailed in Example 7 to give the title compound having the following physical properties (7.0 g). LC/MS  $t_R$  1.95 minutes; MS (ES<sup>+</sup>)  $m/z$  335 (M+H)<sup>a</sup>.

5 Example 24: tert-butyl 2-{4-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-2-oxo-1,2-dihydropyridin-1-yl} acetate

To a glacial acetic acid (100 mL) solution of the compound prepared in Example 23 (7.0 g) was added trimethyl orthoformate (6.9 mL) followed after 30 minutes by sodium azide (4.1 g) and the mixture stirred at room temperature for 16 hours. To the  
10 reaction mixture, water (100 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with water and saturated brine, dried and concentrated. The residue was purified by column chromatography (50 to 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (6.5 g).

15 LC/MS  $t_R$  1.90 minutes; MS (ES<sup>+</sup>)  $m/z$  410 (M+Na)<sup>a</sup>.

Example 25: 2-{4-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-2-oxo-1,2-dihydropyridin-1-yl} acetic acid

To a 1,4-dioxane solution (100 mL) of the compound prepared in Example 24  
20 (6.5) was added 1 M hydrochloric acid (84 mL) and the mixture was heated at 90 °C for 3 hours. The reaction mixture was concentrated to approximately half the original volume and the resultant precipitate collected by filtration to give the title compound having the following physical properties (4.1 g).

LC/MS  $t_R$  1.48 minutes; MS (ES<sup>+</sup>)  $m/z$  332 (M+H)<sup>a</sup>.

25

Example 26: tert-butyl 2-[4-(benzyloxy)-2-oxo-1,2-dihydropyridin-1-yl]-3-phenylpropanoate

To a cooled (-78 °C) tetrahydrofuran (150 mL) solution of the compound prepared in Example 20 (15.3 g) was added benzyl bromide (7.76 mL) followed by a 1  
30 M solution of lithium hexamethyldisilazide in tetrahydrofuran (52.2 mL) and the mixture was stirred at -78 °C for 90 minutes. To the cooled (-78 °C) reaction mixture, a saturated aqueous solution of ammonium chloride (150 mL) was added followed by water (150 mL) and the mixture was allowed to warm to room temperature, then

extracted with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (10 to 30% ethyl acetate in heptanes) to give the title compound having the following physical properties (11.8 g).

5 LC/MS  $t_R$  2.39 minutes; MS (ES<sup>+</sup>)  $m/z$  406 (M+H)<sup>a</sup>.

Example 27: 2-{4-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-2-oxo-1,2-dihydropyridin-1-yl}-3-phenylpropanoic acid

The same operation as in Example 21 → Example 22 → Example 23 →

10 Example 24 → Example 25 was conducted from the compound prepared in Example 26 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.84 minutes; MS (ES<sup>+</sup>)  $m/z$  422 (M+H)<sup>a</sup>.

Example 28: methyl (2S)-2-[4-(benzyloxy)-2-oxopyridin-1(2H)-yl]-3-phenylpropanoate

15 To a tetrahydrofuran (70 mL) solution of 4-benzyloxypyridone (10.8 g), potassium tert-butoxide (6.64 g) and tetrabutylammonium bromide (0.87 g) were added sequentially and the mixture was stirred at room temperature for 30 minutes. To the cooled (0 °C) reaction mixture, a tetrahydrofuran (70 mL) solution of methyl (2R)-3-phenyl-2-[[trifluoromethyl)sulfonyl]oxy]propanoate [Tetrahedron 51(38), 10513  
20 (1995)] (16.8 g) was added over 30 minutes, then the mixture was stirred at 0 °C for 1 hour. To the reaction mixture, water (150 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (50% to 75% ethyl acetate in heptanes) to give the title compound having the following physical properties  
25 (12.7 g).

TLC: R<sub>f</sub> 0.60 (50% ethyl acetate in heptanes).

Example 29: methyl (2S)-2-[4-(2,5-dichlorophenyl)-2-oxopyridin-1(2H)-yl]-3-phenylpropanoate

30 The same operation as in Example 21 → Example 22 → Example 23 was conducted from the compound prepared in Example 28 to give the title compound having the following physical properties. (Note: in the step corresponding to Example

23 in the operation, 2,5-dichlorophenylboronic acid was used).

LC/MS  $t_R$  2.33 minutes; MS ( $ES^+$ )  $m/z$  402 ( $M+H$ )<sup>a</sup>.

Example 30: 2-[4-(2,5-dichlorophenyl)-2-oxopyridin-1(2*H*)-yl]-3-phenylpropanoic acid

5 To a 1: 1 methanol and tetrahydrofuran (18 mL) solution of the compound prepared in Example 29 (0.71 g) was added 1 M sodium hydroxide (4.4 mL) and the mixture was stirred at room temperature for 4 hours. The reaction mixture was concentrated and the residue dissolved in water (50 mL) followed by extraction with *tert*-butyl methyl ether. The aqueous layer was acidified to pH 1 with 1 M hydrochloric  
10 acid and extracted with ethyl acetate. The combined ethyl acetate layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (0.64 g).

LC/MS  $t_R$  4.33 minutes; MS ( $ES^+$ )  $m/z$  388 ( $M+H$ )<sup>b</sup>.

15 Example 31: tert-butyl 2-[4-(2-{bis[(*tert*-butoxy)carbonyl]amino}-5-chlorophenyl)-2-oxo-1,2-dihydropyridin-1-yl]acetate

The compound prepared in Example 23 (4.53 g) was treated as detailed in Example 10 to give the title compound having the following physical properties (4.13 g).  
LC/MS  $t_R$  2.47 minutes; MS ( $ES^+$ )  $m/z$  557 ( $M+Na$ )<sup>a</sup>.

20

Example 32: 2-[4-(2-{bis[(*tert*-butoxy)carbonyl]amino}-5-chlorophenyl)-2-oxo-1,2-dihydropyridin-1-yl]acetic acid

The compound prepared in Example 31 (0.96 g) was treated as detailed in Example 30 to give the title compound having the following physical properties (0.76 g).

25 LC/MS  $t_R$  1.79 minutes; MS ( $ES^+$ )  $m/z$  379 ( $M+H$ )<sup>a</sup>.

Example 33: ethyl 3-(5-chloro-2-nitrophenyl)-3-oxopropanoate

To a toluene (100 mL) solution of 5-chloro-2-nitrobenzoic acid (12 g) was added thionyl chloride (10 mL) and the mixture was stirred at reflux for 4 hours. The reaction  
30 mixture was concentrated and residual thionyl chloride removed by azeotrope with toluene. To a cooled (0 °C) dichloromethane (550 mL) solution of the residue was sequentially added 2,2-dimethyl-1,3-dioxane-4, 6-dione (7.5 g) and 4-dimethylaminopyridine (23 g) and the mixture was stirred at 0 °C for 2 hours. To the

reaction mixture, 1 M hydrochloric acid (100 mL) was added and the dichloromethane layer isolated. The dichloromethane layer was washed, 1 M hydrochloric acid (100 mL x 2), water (100 mL x 2) and brine (100 mL), dried and concentrated. To a toluene (100 mL) solution of the residue was added ethanol (25 mL) and the mixture was stirred in a sealed system at 85 °C for 12 hours. The reaction mixture was concentrated and the residue purified by column chromatography (0 to 30% ethyl acetate in hexanes) to give the title compound having the following physical properties (2.5 g).


LC/MS  $t_R$  2.83 minutes; MS (ES<sup>-</sup>)  $m/z$  270 (M-H)<sup>-</sup>.

10 Example 34: ethyl 2-(5-chloro-2-nitrophenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidine-6-carboxylate

To an *o*-xylene (20 mL) solution of the compound prepared in Example 33 (2 g) was sequentially added 1,8-diazabicyclo[5.4.0]undec-7-ene (4.1 mL) and ethyl (2S)-5-aminopyrrolidine-2-carboxylate [J. Org. Chem. 52(26), 5717 (1987)] (1.2 g) and the mixture was stirred at 130 °C for 1 hour. The reaction mixture was concentrated and the residue purified by column chromatography (0 to 30% ethyl acetate in hexanes) to give the title compound having the following physical properties (0.35 g).

LC/MS  $t_R$  2.71 minutes; MS (ES<sup>+</sup>)  $m/z$  364 (M+H)<sup>+</sup>.

20 Example 35: ethyl 2-(2-amino-5-chlorophenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidine-6-carboxylate



To an ethyl acetate (15 mL) solution of the compound prepared in Example 34 (0.35 g) was sequentially added tin(II) chloride dihydrate (0.73 g) and 6 M HCl (1 mL) and the mixture was stirred at reflux for 2 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (25 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with water and brine, dried and concentrated. The residue was triturated with pentane and the solid

collected by filtration to give the title compound having the following physical properties (0.26 g).

LC/MS  $t_R$  2.53 minutes; MS ( $ES^+$ )  $m/z$  334 ( $M+H$ )<sup>+</sup>.

5 Example 36: tert-butyl N-(4-chloro-2-{6-methyl-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidin-2-yl}phenyl)carbamate

To a *tert*-butanol (10 mL) solution of the compound prepared in Example 35 (0.45 g) was added di-*tert*-butyl dicarbonate (1.17 g) and the mixture was stirred at 80 °C for 8 hours. The reaction mixture was concentrated and the residue obtained  
10 purified by column chromatography (0 to 30% ethyl acetate in hexanes) to give the title compound having the following physical properties (0.40 g).

LC/MS  $t_R$  3.35 minutes; MS ( $ES^+$ )  $m/z$  434 ( $M+H$ )<sup>+</sup>.

15 Example 37: 2-(2-{[(*tert*-butoxy)carbonyl]amino}-5-chlorophenyl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylic acid

To a tetrahydrofuran (5 mL) solution of the compound prepared in Example 36 (0.19 g) was added 2 M sodium hydroxide (1 mL) and the mixture was stirred at room temperature for 12 hours. The reaction mixture was concentrated and the residue obtained dissolved in water (5 mL). To the aqueous solution, potassium hydrogen  
20 sulfate was added to pH of 5 – 6. The mixture was extracted with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (0.13 g).

LC/MS  $t_R$  2.88 minutes; MS ( $ES^+$ )  $m/z$  406 ( $M+H$ )<sup>+</sup>.

25 Example 38: 2-{4-[(methoxycarbonyl)amino]phenyl}-2-oxoethyl 2-(2-{[(*tert*-butoxy)carbonyl]amino}-5-chlorophenyl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylate

To an acetonitrile (15 mL) solution of the compound prepared in Example 37 (0.62 g) and methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10),  
30 2453 (1997)] (0.46 g) was added potassium carbonate (0.53 g) and the mixture was stirred at 50 °C for 8 hours. The reaction mixture was filtered and the filtrate concentrated. The residue was purified by column chromatography (0 – 2% methanol in dichloromethane) to give the title compound having the following physical properties

(0.60 g).

LC/MS  $t_R$  3.40 minutes; MS ( $ES^+$ )  $m/z$  597 ( $M+H$ )<sup>+</sup>

5 Example 39: tert-butyl N-(4-chloro-2-[6-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidin-2-yl]phenyl)carbamate

To a toluene (10 mL) solution of the compound prepared in Example 38 (0.60 g) was added ammonium acetate (0.78 g) and the mixture was heated at reflux for 12 hours. The reaction mixture was concentrated and the residue suspended in water and extracted into ethyl acetate. The combined organic layers were washed with brine, dried and  
10 concentrated. The residue was purified by column chromatography (0 – 45% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.41 g)


LC/MS  $t_R$  2.59 minutes; MS ( $ES^+$ )  $m/z$  577 ( $M+H$ )<sup>+</sup>.

15 Example 40: methyl N-(4-{2-[2-(2-amino-5-chlorophenyl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidin-6-yl]-1H-imidazol-5-yl}phenyl)carbamate

To a dichloromethane (5 mL) solution of the compound prepared in Example 39 (100 mg) was added trifluoroacetic acid (1 mL) and the mixture stirred at room temperature for 2 hours. To the reaction mixture, a saturated aqueous solution of sodium  
20 hydrogen carbonate (15 mL) was added followed by extraction into dichloromethane. The combined organic layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (65 mg).

LC/MS  $t_R$  2.05 minutes; MS ( $ES^+$ )  $m/z$  477 ( $M+H$ )<sup>+</sup>.

25 Example 41: methyl [4-(2-{2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate



To a glacial acetic acid (1 mL) solution of the compound prepared in Example




40 (60 mg) was added triethyl orthoformate (60  $\mu$ L) followed after 30 minutes by sodium azide (25 mg) and the mixture stirred in a sealed tube at 65 °C for 1 hour. The reaction mixture was then concentrated and the residue purified by column chromatography (0 – 5% methanol in dichloromethane) to give the title compound

5 having the following physical properties (20 mg).

LC/MS  $t_R$  3.07 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.45 (s, 1 H), 7.92 (d, 1 H), 7.78 (dd, 1 H), 7.71 (d, 1 H), 7.57 (d, 2 H), 7.44 (d, 2 H), 7.25 (br. s, 1 H), 6.34 (s, 1 H), 5.72 (dd, 1 H), 3.73 (s, 3 H), 3.37 - 3.24 (obs. m, 1 H), 2.97 - 2.87 (m, 1 H), 2.73 - 2.62 (m, 1 H), 2.43 - 2.33 (m, 1 H).

Example 42: methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl)phenyl]carbamate



15 The same operation as in Example 38 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.73 (m, 2 H), 7.70 (d, 1 H), 7.58 (d, 2 H), 7.46 (d, 2 H), 7.24 (br. s, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H), 5.83 - 5.72 (m, 1 H), 3.76 (s, 3 H), 3.52 - 3.41 (m, 1 H), 3.17 - 3.08 (m, 1 H), 2.71 - 2.59 (m, 1 H), 2.54 - 2.44 (m, 1 H).

25 Example 43: tert-butyl N-{4-chloro-2-[3-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl]phenyl}carbamate

The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 19 to give the title compound having the following

physical properties.

LC/MS  $t_R$  1.75 minutes; MS ( $ES^+$ )  $m/z$  576 ( $M+H$ )<sup>a</sup>.

Example 44: tert-butyl N-{4-chloro-2-[3-(4-chloro-5-{4-  
5 [(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-  
tetrahydroindolizin-7-yl]phenyl} carbamate

- A tetrahydrofuran (20 mL) solution of the compound prepared in Example 43 (0.20 g) was cooled to 0 °C and N-chlorosuccinimide (70 mg) was added. The mixture was stirred at room temperature for 16 hours, cooled to 0 °C and further N-  
10 chlorosuccinimide (35 mg) added. The mixture was stirred for a further 24 hours, water (20 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (40 to 50% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.16 g).  
15 TLC:  $R_f$  0.27 (ethyl acetate).

Example 45: methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]carbamate

- The same operation as in Example 40 → Example 41 was conducted from the  
20 compound prepared in Example 44 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.75 minutes; MS ( $ES^+$ )  $m/z$  563 ( $M+H$ )<sup>c</sup>

- <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  11.23 (br. s, 1 H), 8.53 (s, 1 H), 7.68 - 7.28 (m, 5 H),  
7.23 (d, 2 H), 6.77 (s, 1 H), 6.32 (s, 1 H), 5.80 (dd, 1 H), 5.71 (s, 1 H), 3.80 (s, 3 H),  
25 3.52 - 3.35 (m, 1 H), 3.28 - 3.18 (m, 1 H), 3.06 - 2.92 (m, 1 H), 2.53 - 2.35 (m, 1 H).

Example 46: methyl 3-{2-[7-(2-{[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl} benzoate

- The same operation as in Example 38 → Example 39 was conducted from the  
30 compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, methyl 3-(bromoacetyl)benzoate was used in place of methyl [4-(bromoacetyl)-phenyl]carbamate).

LC/MS  $t_R$  2.87 minutes; MS (ES<sup>-</sup>)  $m/z$  559 (M-H)<sup>o</sup>.

Example 47: methyl 3-{2-[7-(2-{[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-chloro-1H-imidazol-5-yl}benzoate

- 5           A dichloromethane (25 mL) solution of the compound prepared in Example 46 (0.25 g) was cooled to 0 °C and N-chlorosuccinimide (72 mg) was added. The mixture was stirred at room temperature for 2 hours. The reaction mixture was then filtered and the filtrate concentrated. The residue was purified by column chromatography (30 to 35% ethyl acetate in heptanes) to give the title compound having the following physical
- 10           properties (0.10 g).

LC/MS  $t_R$  3.44 minutes; MS (ES<sup>+</sup>)  $m/z$  595 (M+H)<sup>o</sup>.

Example 48: methyl 3-{2-[7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-chloro-1H-imidazol-5-yl}benzoate

- 15           The compound prepared in Example 47 (0.25 g) was treated as detailed in Example 40 to give the title compound having the following physical properties (0.20 g). LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  495 (M+H)<sup>o</sup>.

Example 49: 3-{2-[7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-chloro-1H-imidazol-5-yl}benzoic acid

- 20           To a solution of the compound prepared in Example 48 (0.14 g) in a 1 : 1 mixture of tetrahydrofuran and water (4 mL) was added lithium hydroxide monohydrate (24 mg) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated to approximately half its original volume and the aqueous residue
- 25           acidified to pH 2-3 by portion-wise addition of potassium hydrogen sulfate followed by extraction into dichloromethane. The combined organic layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (0.12 g).

LC/MS  $t_R$  2.71 minutes; MS (ES<sup>+</sup>)  $m/z$  481 (M+H)<sup>o</sup>.

30

Example 50: 3-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzoic acid

The compound prepared in Example 49 (75 mg) was treated as detailed in

Example 41 to give the title compound having the following physical properties (60 mg).

LC/MS  $t_R$  2.65 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H), 506 (M-N<sub>2</sub>+H)<sup>c</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.20 (br. s, 1 H), 9.71 (s, 1 H), 8.33 (s, 1 H), 7.97 (d, 1 H), 7.88 (d, 1 H), 7.89 - 7.77 (m, 3 H), 7.60 (t, 1 H), 5.99 (s, 1 H), 5.95 (s, 1 H), 5.59 (dd, 1 H), 3.34 - 3.23 (obs. m, 1 H), 3.07 - 2.90 (m, 1 H), 2.56 - 2.40 (m, 1 H), 2.30 - 2.20 (m, 1 H).

Example 51: 2-oxo-2-phenylethyl (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylate

To an N,N-dimethylformamide (3 mL) solution of the compound prepared in Example 9 (100 mg) was added potassium carbonate (151 mg) and the mixture stirred at room temperature for 30 minutes. To the reaction mixture, a solution of 2-bromo-1-phenylethan-1-one (72 mg) in N,N-dimethylformamide (3 mL) was added and the mixture stirred at room temperature for 2 hours. The reaction mixture was then diluted with ethyl acetate (30 mL), filtered and the filtrate concentrated to give the crude title compound having the following physical properties (193 mg).  
LC/MS  $t_R$  1.97 minutes; MS (ES<sup>+</sup>)  $m/z$  476 (M+H)<sup>a</sup>.

Example 52: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-phenyl-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

To a toluene (5 mL) suspension of the compound prepared in Example 51 (193 mg) was sequentially added glacial acetic acid (0.50 mL) and ammonium acetate (231 mg) and the mixture stirred at reflux for 2 hours. The reaction mixture was concentrated and the residue suspended in a saturated aqueous solution of sodium hydrogen carbonate (10 mL) followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (5% methanol in ethyl acetate) followed by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (54.5 mg).  
LC/MS  $t_R$  3.20 minutes; MS (ES<sup>+</sup>)  $m/z$  456 (M+H)<sup>b</sup>  
NMR analysis showed a 2: 1 ratio of tautomers.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.19 and 10.77 (br. s, 1 H), 8.54 (s, 1 H), 7.78 - 7.73 (m, 1 H), 7.61 (dd, 1 H), 7.57 - 7.52 (m, 1 H), 7.52 - 7.43 (m, 2 H), 7.42 - 7.31 (m, 2 H), 7.27 - 7.15 (m, 2 H), 6.32 (s, 1 H), 5.92 - 5.79 (m, 1 H), 5.70 (s, 1 H), 3.54 - 3.25 (m, 2 H), 3.01 (dd, 1 H), 2.53 - 2.39 (m, 1 H).

5

Example 53: 4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-1-[(5-phenyl-1H-imidazol-2-yl)methyl]-2(1H)-pyridinone

The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 25 to give the title compound having the following physical properties.

10

LC/MS *t<sub>R</sub>* 3.16 minutes; MS (ES<sup>+</sup>) *m/z* 430 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-*d*<sub>6</sub>) δ 9.69 (s, 1 H), 7.80 (app. br. s, 3 H), 7.75 - 7.62 (m, 3 H), 7.50 (br. s, 1 H), 7.32 (t, 2 H), 7.17 (t, 1 H), 6.27 (d, 1 H), 5.86 (dd, 1 H), 5.09 (s, 2 H).

15

Example 54: tert-butyl N-(4-chloro-2-{1-[(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)methyl]-2-oxo-1,2-dihydropyridin-4-yl}phenyl)carbamate

The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 32 to give the title compound having the following physical properties.

20

LC/MS *t<sub>R</sub>* 1.69 minutes; MS (ES<sup>+</sup>) *m/z* 550 (M+H)<sup>a</sup>.

Example 55: methyl N-[4-(2-{[4-(2-amino-5-chlorophenyl)-2-oxo-1,2-dihydropyridin-1-yl]methyl}-1H-imidazol-5-yl)phenyl]carbamate

25

To a 1, 4-dioxane (0.75 mL) solution of the compound prepared in Example 54 (109 mg) was added 4 M hydrochloric acid in dioxane (0.75 mL) and the mixture was stirred at room temperature for 16 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford the title compound having the following physical properties (78 mg).

30


LC/MS *t<sub>R</sub>* 1.54 minutes; MS (ES<sup>+</sup>) *m/z* 450 (M+H)<sup>a</sup>

Example 56: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-4-yl]phenyl}carbamate

The compound prepared in Example 55 (78 mg) was treated as detailed in Example 41 to give the title compound having the following physical properties (40 mg).

- 5 LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  503 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.70 (s, 1 H), 9.68 (s, 1 H), 7.83 - 7.78 (m, 3 H), 7.70 (d, 1 H), 7.62 (d, 2 H), 7.57 - 7.35 (m, 4 H), 6.29 (s, 1 H), 5.89 (d, 1 H), 5.13 (s, 2 H), 3.66 (s, 3 H).

- 10 Example 57: methyl {4-[5-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-4-yl]phenyl}carbamate



The compound prepared in Example 56 (72 mg) was treated as detailed in Example 47 to give the title compound having the following physical properties (16.2 mg).

- 15 LC/MS  $t_R$  3.96 minutes; MS (ES<sup>+</sup>)  $m/z$  537 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.76 (s, 1 H), 9.71 (s, 1 H), 7.84 - 7.79 (m, 3 H), 7.68 (d, 1 H), 7.63 (br. s, 1 H), 7.62 (d, 2 H), 7.52 (d, 2 H), 6.26 (d, 1 H), 5.88 (dd, 1 H), 5.04 (s, 2 H), 3.69 (s, 3 H).

20

Example 58: tert-butyl 2-[4-(2-{bis[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-2-oxo-1,2-dihydropyridin-1-yl]butanoate

To a cooled (-78°C) tetrahydrofuran (25 mL) solution of the compound prepared in Example 31 (2.5 g) was added 1 M lithium hexamethyldisilazide in tetrahydrofuran (5.61 mL) and the mixture was stirred at -78°C for 30 minutes. To the cooled (-78°C) reaction mixture, iodoethane (0.45 mL) was added and the mixture was stirred at -78°C for 2 hours. To the cooled (-78°C) reaction mixture, a saturated aqueous solution of

25

ammonium chloride was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried (MgSO<sub>4</sub>) and concentrated. The residue was purified by column chromatography (10% to 20% ethyl acetate in heptanes) to give the title compound having the following physical properties (2.37 g).

- 5 LC/MS *t*<sub>R</sub> 2.62 minutes; MS (ES<sup>+</sup>) *m/z* 585 (M+Na)<sup>a</sup>.

Example 59: methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}propyl)-1H-imidazol-4-yl]phenyl}carbamate

- The same operation as in Example 25 → Example 9 → Example 38 → Example  
10 39 was conducted from the compound prepared in Example 58 to give the title compound having the following physical properties.

LC/MS *t*<sub>R</sub> 3.54 minutes; MS (ES<sup>+</sup>) *m/z* 531 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.32 (s, 1 H), 9.68 (s, 1 H), 9.62 (s, 1 H), 7.84 - 7.77 (m, 3 H), 7.71 (d, 1 H), 7.68 - 7.39 (m, 5 H), 6.24 (s, 1 H), 5.96 (s, 1 H), 5.92 (dd, 1 H),  
15 3.68 (s, 3 H), 2.28 - 1.85 (m, 2 H), 0.79 (t, 3 H).

Example 60: methyl {4-[5-chloro-2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}propyl)-1H-imidazol-4-yl]phenyl}carbamate

- The compound prepared in Example 59 (200 mg) was treated as detailed in  
20 Example 47 to give the title compound having the following physical properties (42.5 mg).

LC/MS *t*<sub>R</sub> 4.22 minutes; MS (ES<sup>+</sup>) *m/z* 565 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.69 (s, 1 H), 9.79 (s, 1 H), 9.69 (s, 1 H), 7.85 - 7.78 (m, 3 H), 7.67 (d, 1 H), 7.62 (d, 2 H), 7.54 (d, 2 H), 6.24 (d, 1 H), 6.04 - 5.86 (m, 2 H),  
25 3.69 (s, 3 H), 2.23 - 1.83 (m, 2 H), 0.79 (t, 3 H).

Example 61: methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}ethyl)-1H-imidazol-5-yl]phenyl}carbamate

- The same operation as in Example 58 → Example 25 → Example 9 → Example  
30 38 → Example 39 was conducted from the compound prepared in Example 31 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 58 in the operation, iodomethane was used).

LC/MS *t*<sub>R</sub> 3.16 minutes; MS (ES<sup>+</sup>) *m/z* 517 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.29 (br. s, 1 H), 9.68 (s, 1 H), 9.60 (br. s, 1 H), 7.87 - 7.76 (m, 3 H), 7.67 (d, 2 H), 7.60 - 7.38 (m, 4 H), 6.25 (s, 1 H), 6.12 (q, 1 H), 5.90 (d, 1 H), 3.66 (s, 3 H), 1.64 (d, 3 H).

5 Example 62: ethyl 2-[4-(benzyloxy)-2-oxo-1,2-dihydropyridin-1-yl]prop-2-enoate

To a dichloromethane (25 mL) solution of ethyl prop-2-ynoate (0.61 mL), 4-(benzyloxy)pyridin-2(1H)-one (1.0 g) and triphenylphosphine (0.20 g) were sequentially added at 0 °C and the mixture was stirred at room temperature for 16 hours. The reaction mixture was concentrated under reduced pressure and the residue purified  
10 by column chromatography to give the title compound having the following physical properties (1.0 g).

LC/MS *t<sub>R</sub>* 1.87 minutes; MS (ES<sup>+</sup>) *m/z* 300 (M+H)<sup>a</sup>.

15 Example 63: ethyl 1-[4-(benzyloxy)-2-oxo-1,2-dihydropyridin-1-yl]cyclopropane-1-carboxylate

To a dimethylsulfoxide (8 mL) solution of trimethylsulphoxonium iodide (0.91 g), sodium hydride (0.16 g, 60% dispersion in mineral oil) was added and the mixture was stirred at room temperature for 15 minutes. A dimethylsulfoxide (6 mL) solution of the compound prepared in Example 62 (1.0 g) was added and the mixture was stirred at  
20 room temperature for 1.5 hours. To the reaction mixture, water (50 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by flash chromatography to give the title compound having the following physical properties (0.39 g).

LC/MS *t<sub>R</sub>* 1.88 minutes; MS (ES<sup>+</sup>) *m/z* 314 (M+H)<sup>a</sup>.

25

Example 64: 1-[4-(benzyloxy)-2-oxo-1,2-dihydropyridin-1-yl]cyclopropane-1-carboxylic acid

The compound prepared in Example 63 (0.39 g) was treated as detailed in Example 8 to give the title compound having the following physical properties (0.27 g).

30 LC/MS *t<sub>R</sub>* 1.64 minutes; MS (ES<sup>-</sup>) *m/z* 284 (M-H)<sup>a</sup>.

Example 65: methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-oxo-1(2H)-pyridinyl}cyclopropyl)-1H-imidazol-5-yl]phenyl}carbamate



The same operation as in Example 51 → Example 52 → Example 21 → Example 6 → Example 7 → Example 41 was conducted from the compound prepared in Example 64 to give the title compound having the following physical properties.

(Note: in the step corresponding to Example 51 in the operation, methyl [4-

5 (bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).

LC/MS  $t_R$  3.10 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.72 (br. s, 1 H), 9.69 (s, 1 H), 9.59 (br. s, 1 H), 7.84 - 7.80 (m, 2 H), 7.79 (d, 1 H), 7.71 (d, 1 H), 7.58 (d, 2 H), 7.40 (br. d, 2 H), 7.35 (d, 1 H), 6.29 (d, 1 H), 5.79 (dd, 1 H), 3.64 (s, 3 H), 1.63 (br. s, 2 H), 1.38 (br. s, 2 H).

10

Examples 66(1) and 66(2): methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2-oxo-1(2H)-pyridinyl}-2-phenylethyl)-1H-imidazol-4-yl]phenyl}carbamate and methyl [4-(2-{1-[4-(2-carbamimidamido-5-chlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-4-yl)phenyl]carbamate acetate

15 The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 27 to obtain the title products in a 5: 2 ratio having the following physical properties.

Example 66(1):

20 LC/MS  $t_R$  3.63 minutes; MS (ES<sup>+</sup>)  $m/z$  593 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.25 (s, 1 H), 9.64 (s, 1 H), 9.61 (s, 1 H), 7.85 - 7.78 (m, 3 H), 7.76 (d, 1 H), 7.69 (d, 2 H), 7.48 - 7.38 (m, 3 H), 7.25 (t, 2 H), 7.17 (t, 1 H), 7.11 (d, 2 H), 6.33 - 6.21 (m, 1 H), 6.20 (s, 1 H), 5.90 (d, 1 H), 3.67 (s, 3 H), 3.56 - 3.46 (m, 2 H).

25

Example 66(2):

LC/MS  $t_R$  3.50 minutes; MS (ES<sup>+</sup>)  $m/z$  582 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.24 (s, 1 H), 9.61 (s, 1 H), 9.49 (s, 1 H), 7.99 (d, 1 H), 7.71 (d, 2 H), 7.52 (d, 2 H), 7.48 - 7.42 (m, 5 H), 7.37 (s, 1 H), 7.30 - 7.17 (m, 5 H), 7.19 - 7.13 (m, 1 H), 6.40 (s, 1 H), 6.35 (s, 1 H), 6.30 (d, 1 H), 3.66 (s, 3 H), 3.79 - 3.53 (m, 2 H), 1.88 (s, 3 H).

30

Example 67: methyl {4-[5-chloro-2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}-2-phenylethyl)-1H-imidazol-4-yl]phenyl}carbamate

The compound prepared in Example 66(1) (38 mg) was treated as detailed in Example 47 to give the title compound having the following physical properties (16.8 mg).

LC/MS  $t_R$  4.51 minutes; MS (ES<sup>+</sup>)  $m/z$  627 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.77 (s, 1 H), 9.78 (s, 1 H), 9.64 (s, 1 H), 7.84 - 7.78 (m, 3 H), 7.77 (d, 1 H), 7.60 (d, 2 H), 7.53 (d, 2 H), 7.27 (t, 2 H), 7.20 (t, 1 H), 7.09 (d, 2 H), 6.25 (t, 1 H), 6.18 (d, 1 H), 5.93 (dd, 1 H), 3.68 (s, 3 H), 3.56 - 3.35 (m, 2 H).

Example 68: 2-nitro-5-(1H-pyrazol-1-yl)aniline

To an N,N-dimethylformamide (4 mL) solution of 5-chloro-2-nitroaniline (200 mg) was added 1H-pyrazole (316 mg) and potassium hydroxide (98 mg) and the mixture stirred at 100 °C for 20 hours. To the reaction mixture at room temperature, water (10 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 to 50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (52 mg).

LC/MS  $t_R$  1.71 minutes; MS (ES<sup>+</sup>)  $m/z$  205 (M+H)<sup>a</sup>.

Example 69: 4-(1H-pyrazol-1-yl)benzene-1,2-diamine

To a 4 : 1 ethanol: water (2 mL) solution of the compound prepared in Example 68 (52 mg) was added a saturated aqueous solution of ammonium chloride solution (0.5 mL) and iron powder (113 mg) and the mixture stirred at room temperature for 1.5 hours. The reaction mixture was filtered through Celite® and the filtrate concentrated. The residue was triturated with dichloromethane and the precipitate formed was isolated by filtration giving the title compound with the following physical properties (49 mg).

LC/MS  $t_R$  0.70 minutes; MS (ES<sup>+</sup>)  $m/z$  175 (M+H)<sup>a</sup>.

Example 70: N-[2-amino-4-(1H-pyrazol-1-yl)phenyl]-2-{4-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-2-oxo-1,2-dihydropyridin-1-yl}-3-phenylpropanamide

To an N,N-dimethylformamide (1 mL) solution of the compound prepared in

Example 69 (49 mg) and the compound prepared in Example 27 (55 mg) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (59.67 mg) and diisopropylethylamine (40  $\mu$ L) and the mixture stirred at room temperature for 16 hours. To the reaction mixture, water (5 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and then concentrated. The residue was purified by column chromatography (0 to 50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (29 mg).

LC/MS  $t_R$  1.98 minutes; MS ( $ES^+$ )  $m/z$  578 (M+H)<sup>a</sup>.

Example 71: 4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-1-{2-phenyl-1-[5-(1H-pyrazol-1-yl)-1H-benzimidazol-2-yl]ethyl}-2(1H)-pyridinone

A glacial acetic acid (1 mL) solution of the compound prepared in Example 70 (29 mg) was stirred at 60 °C for 1 hour. To the cooled (0 °C) reaction mixture, 2 M sodium hydroxide (2 mL) and water (2 mL) was sequentially added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (5 mg).

LC/MS  $t_R$  4.07 minutes; MS ( $ES^+$ )  $m/z$  560 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.20 (s, 1 H), 8.48 (d, 1 H), 7.96 (br. s, 1 H), 7.86 (d, 1 H), 7.84 - 7.68 (m, 5 H), 7.65 (br. s, 1 H), 7.31 (t, 2 H), 7.23 (t, 1 H), 7.12 (d, 2 H), 6.53 (t, 1 H), 6.47 - 6.36 (m, 1 H), 6.22 (d, 1 H), 5.90 (dd, 1 H), 3.68 (dd, 1 H), 3.46 (dd, 1 H).

Example 72: 2-[4-(2,5-dichlorophenyl)-2-oxo-1,2-dihydropyridin-1-yl]-N-[2-(4-nitrophenyl)-2-oxoethyl]-3-phenylpropanamide

To a pyridine (8 mL) solution of the compound prepared in Example 30 (0.63 g) and 2-amino-1-(4-nitrophenyl)ethanone hydrochloride hydrate (0.46 g) was added N-[3-(dimethylamino)propyl]-N'-ethylcarbodiimide hydrochloride (0.40 g) and the mixture stirred at room temperature for 20 hours. To the reaction mixture, 0.5 M hydrochloric acid (50 mL) was added followed by extraction with ethyl acetate. The combined

organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (5% to 50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.64 g).

TLC: Rf 0.13 (50% ethyl acetate in heptanes).

5

Example 73: 4-(2,5-dichlorophenyl)-1-{1-[4-(4-nitrophenyl)-1H-imidazol-2-yl]-2-phenylethyl}-1,2-dihydropyridin-2-one

To an *o*-xylene (50 mL) solution of the compound prepared in Example 72 (0.62 g) was added ammonium acetate (0.87 g) and the mixture stirred at 150 °C for 1 hour.

10 The reaction mixture was concentrated and the residue suspended in a saturated aqueous solution of sodium hydrogen carbonate (100 mL) followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (10% to 50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.61 g).

15 TLC: Rf 0.11 (50% ethyl acetate in heptanes).

Example 74: 1-{1-[5-(4-aminophenyl)-1H-imidazol-2-yl]-2-phenylethyl}-4-(2,5-dichlorophenyl)-2(1H)-pyridinone

To a solution of the compound prepared in Example 73 (0.30 g) in an 8:1:1  
20 mixture of ethanol, saturated aqueous ammonium chloride solution and water (7.5 mL) was added iron powder (0.25 g) and the mixture stirred at 50 °C for 3 hours. The reaction mixture was filtered through Celite® and the filtrate concentrated. The residue obtained was suspended in water and extracted into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by  
25 column chromatography (50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.27 g).

LC/MS  $t_R$  3.63 minutes; MS (ES<sup>+</sup>)  $m/z$  501 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 7.89 (d, 1 H), 7.48 (d, 1 H), 7.40 (dd, 1 H), 7.37 (d, 1 H), 7.31 (app. br. s, 2 H), 7.25 - 7.19 (m, 4 H), 7.18 (s, 1 H), 7.18 - 7.13 (m, 1 H), 6.73  
30 (d, 2 H), 6.48 (d, 1 H), 6.43 (d, 1 H), 6.42 (br. s, 1 H), 3.78 - 3.59 (m, 1 H), 3.59 - 3.37 (m, 1 H).

Example 75: methyl [4-(2-{1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-

phenylethyl}-1H-imidazol-4-yl)phenyl]carbamate

To a dichloromethane (2.5 mL) solution of the compound prepared in Example 74 (0.26 g) was sequentially added methyl chloroformate (50  $\mu$ l) and N,N-diisopropylethylamine (135  $\mu$ L) and the mixture stirred at room temperature for 1 hour.

- 5 To the reaction mixture, water (20 mL) was added followed by extraction with dichloromethane. The combined organic layers were washed with brine, dried and concentrated. To a methanol (2.5 mL) solution of the residue, 1 M sodium hydroxide (0.51 mL) was added and the mixture was stirred at room temperature for 30 minutes. To the reaction mixture, water (30 mL) was added followed by extraction with ethyl
- 10 acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (10% to 50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.22 g). LC/MS  $t_R$  4.18 minutes; MS (ES<sup>+</sup>)  $m/z$  559 (M+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  7.93 (br. s, 1 H), 7.70 (br. s, 1 H), 7.49 (d, 2 H), 7.46 (d, 2 H), 7.41 (dd, 1 H), 7.38 (d, 1 H), 7.33 (s, 1 H), 7.27 - 7.19 (m, 4 H), 7.19 - 7.12 (m, 1 H), 6.49 (d, 1 H), 6.45 (d, 1 H), 6.55 - 6.30 (m, 1 H), 3.74 (s, 3 H), 3.80 - 3.62 (m, 1 H), 3.53 (app. br. s, 1 H).
- 15

Example 76: methyl [4-(5-chloro-2-{1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-4-yl)phenyl]carbamate

- 20 To an acetonitrile (3.5 mL) solution of the compound prepared in Example 75 (0.195 g) was added 1-chloropyrrolidine-2,5-dione (50 mg) and the mixture stirred at 60 °C for 5.5 hours. The reaction mixture was concentrated and the residue purified by column chromatography (5% to 50% ethyl acetate in heptanes) to obtain the title
- 25 compound having the following physical properties (52 mg). LC/MS  $t_R$  5.09 minutes; MS (ES<sup>+</sup>)  $m/z$  593 (M+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  7.94 (d, 1 H), 7.57 (d, 2 H), 7.50 (d, 2 H), 7.49 (d, 1 H), 7.41 (dd, 1 H), 7.38 (d, 1 H), 7.26 - 7.21 (m, 4 H), 7.21 - 7.15 (m, 1 H), 6.48 (s, 1 H), 6.47 (dd, 1 H), 6.42 (t, 1 H), 3.74 (s, 3 H), 3.65 (dd, 1 H), 3.45 (dd, 1 H).
- 30

Example 77: methyl N-(4-propanoylphenyl)carbamate

To a dichloromethane (10 mL) solution of 1-(4-aminophenyl)propan-1-one (0.25 g), pyridine (0.27 mL) and methyl chloroformate (0.15 mL) were added sequentially

and the mixture stirred overnight at room temperature. To the reaction mixture, water (25 mL) was added followed by extraction with dichloromethane. The combined organic layers were washed sequentially with 1 M hydrochloric acid, water and brine, dried and concentrated to obtain the title compound having the following physical properties (0.23 g).


LC/MS  $t_R$  1.68 minutes; MS ( $ES^+$ )  $m/z$  208 ( $M+H$ )<sup>a</sup>.

Example 78: methyl N-[4-(2-bromopropanoyl)phenyl]carbamate

To a glacial acetic acid (10 mL) solution of the compound prepared in Example 77 (0.23 g) was added pyridinium tribromide (0.35 g) followed by a solution of 33 wt. % hydrogen bromide in acetic acid (0.15 mL) and the mixture stirred at room temperature overnight. The reaction mixture was concentrated and a saturated aqueous solution of sodium hydrogen carbonate (25 mL) was added to the residue followed by extraction with ethyl acetate. The combined organic layers were washed with water, brine, dried and concentrated to obtain the title compound having the following physical properties (0.29 g).

LC/MS  $t_R$  1.87 minutes; MS ( $ES^+$ )  $m/z$  286 and 288 ( $M+H$ )<sup>a</sup>.

Example 79: formic acid - methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -4-methyl-1H-imidazol-5-yl)phenyl]carbamate (1:1)



The same operation as in Example 38 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 78

was used in the step corresponding to Example 41 in the operation, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt).

- 5 LC/MS  $t_R$  3.00 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.92 (s, 1 H), 9.69 (s, 1 H), 9.59 (s, 1 H), 8.51 (br. s, 1 H), 7.80 - 7.79 (m, 3 H), 7.52 - 7.39 (m, 4 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.57 - 5.55 (m, 1 H), 3.66 (s, 3 H), 3.36 - 3.34 (obs. m, 1 H), 3.00 - 2.96 (m, 1 H), 2.51 - 2.49 (obs. m, 1 H), 2.33 (s, 3 H), 2.19 (app. br. s, 1 H).


10

Example 80: methyl N-[4-(2-bromobutanoyl)phenyl]carbamate

The same operation as in Example 77 → Example 78 was conducted from 1-(4-aminophenyl)butan-1-one to obtain the title compound having the following physical properties.

- 15 LC/MS  $t_R$  2.00 minutes; MS (ES<sup>+</sup>)  $m/z$  300 and 302 (M+H)<sup>a</sup>.

Examples 81(1) and 81(2): formic acid - methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-4-ethyl-1H-imidazol-5-yl)phenyl]carbamate (1:1) and formic acid - methyl (4-{2-[7-(5-chloro-2-  
 20 formamidophenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll]-4-ethyl-1H-imidazol-5-yl}phenyl]carbamate (1:1)



- The same operation as in Example 38 → Example 39 → Example 40 →  
 Example 41 was conducted from the compound prepared in Example 19 to obtain the  
 25 title compounds having the following physical properties. (Note: in the step  
 corresponding to Example 38 in the operation, the compound prepared in Example 80  
 was used. In the step corresponding to Example 41 in the operation, high performance  
 liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in

mobile phase A (0.1% aqueous formic acid)] was used to give the title products in a 4 : 1 ratio as their formic acid salts).

Example 81(1):

- 5 LC/MS  $t_R$  3.10 minutes; MS (ES<sup>+</sup>)  $m/z$  557 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.93 (s, 1 H), 9.69 (s, 1 H), 9.60 (s, 1 H), 8.51 (br. s, 1 H), 7.80 - 7.78 (m, 3 H), 7.50 - 7.37 (m, 4 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.57 (d, 1 H), 3.66 (s, 3 H), 3.36 - 3.34 (obs. m, 2 H), 3.00 - 2.95 (m, 1 H), 2.73 - 2.71 (m, 1 H), 2.51 - 2.49 (obs. m, 1 H), 2.31 - 2.27 (m, 1 H), 1.19 (t, 3 H).


10

Example 81(2):

- LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  532 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.93 (s, 1 H), 9.73 (s, 1 H), 9.63 (s, 1 H), 8.21 (d, 1 H), 8.33 (s, 1 H), 7.99 (d, 1 H), 7.49 - 7.41 (m, 6 H), 6.27 (s, 1 H), 6.19 (s, 1 H), 5.65 (d, 1 H), 3.67 (s, 3 H), 3.36 - 3.34 (obs. m, 3 H), 3.17 - 3.12 (m, 1 H), 2.72 - 2.70 (m, 2 H), 1.18 (t, 3 H).

15

Example 82: formic acid - methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-4-methyl-1H-imidazol-5-yl]phenyl}carbamate (1:1)



20

- The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 25 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 78 was used. In the step corresponding to Example 39 in the operation, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt).


25

LC/MS  $t_R$  3.03 minutes; MS (ES<sup>+</sup>)  $m/z$  517 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.98 (br. s), 9.70 (s, 1 H), 9.63 (br. s, 1 H), 8.25 (s, 1 H), 7.86 - 7.73 (m, 3 H), 7.68 (d, 1 H), 7.46 (app. br. s, 4 H), 6.27 (d, 1 H), 5.88 (dd, 1 H), 5.03 (s, 2 H), 3.67 (s, 3 H), 2.30 (s, 3 H).

5 Example 83: formic acid - methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-4-ethyl-1H-imidazol-5-yl]phenyl} carbamate (1:1)



The same operation as in Example 38 → Example 52 was conducted from the compound prepared in Example 25 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 80 was used. In the step corresponding to Example 39 in the operation, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt).

15 LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  531 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.99 (br. s, 1 H), 9.70 (s, 1 H), 9.63 (br. s, 1 H), 8.26 (br. s, 1 H), 7.88 - 7.76 (m, 3 H), 7.68 (d, 1 H), 7.45 (app. br. d, 4 H), 6.28 (d, 1 H), 5.87 (dd, 1 H), 5.04 (s, 2 H), 3.67 (s, 3 H), 2.69 (app. br. s, 2 H), 1.17 (t, 3 H).

20 Example 84: 2-oxo-2-(2-oxo-2,3-dihydro-1H-indol-5-yl)ethyl 7-(2-{{(tert-butoxy)carbonyl}amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

To an acetonitrile (4 mL) solution of the compound prepared in Example 19 (0.35 g) was added 5-(bromoacetyl)-1,3-dihydro-2H-indol-2-one (0.25 g) and N,N-diisopropylethylamine (0.22 mL) and the mixture was stirred at 50 °C for 16 hours. To the reaction mixture, water was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to give the crude title compound having the following physical properties (0.67 g).

LC/MS  $t_R$  1.99 minutes; MS ( $ES^+$ )  $m/z$  578 ( $M+H$ )<sup>a</sup>.

Example 85: 5-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)-1,3-dihydro-2H-indol-2-one



5

The same operation as in Example 39 → Example 55 → Example 41 was conducted from the compound prepared in Example 84 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.80 minutes; MS ( $ES^+$ )  $m/z$  511 ( $M+H$ )<sup>b</sup>

- 10 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.05 (s, 1 H), 10.36 (s, 1 H), 9.70 (s, 1 H), 8.48 (br. s, 1 H), 7.95 - 7.63 (m, 3 H), 7.55 (s, 1 H), 7.50 (br. s, 1 H), 7.35 (s, 1 H), 6.78 (d, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H), 3.50 (s, 2 H), 3.45 - 3.30 (obs. m, 1 H), 2.99 (ddd, 1 H), 2.54 - 2.43 (obs. m, 1 H), 2.33 (app. br. s, 1 H).

- 15 Example 86: 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)-3,4-dihydro-2(1H)-quinolinone

The same operation as in Example 84 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to

20 Example 84 in the operation, 6-(bromoacetyl)-3,4-dihydroquinolin-2(1H)-one was used).

LC/MS  $t_R$  2.88 minutes; MS ( $ES^+$ )  $m/z$  525 ( $M+H$ )<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.09 (br. s, 1 H), 10.08 (br. s, 1 H), 9.69 (s, 1 H), 7.84 - 7.75 (m, 3 H), 7.51 (s, 1 H), 7.46 (d, 1 H), 7.38 (br. s, 1 H), 6.83 (d, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.63 (d, 1 H), 3.42 - 3.34 (obs. m, 1 H), 3.01 (ddd, 1 H), 2.91 (t, 2 H),
- 25 2.56 - 2.49 (obs. m, 1 H), 2.46 (t, 2 H), 2.35 - 2.26 (m, 1 H).

Example 87: 6-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone

The compound prepared in Example 86 (0.10 g) was treated as detailed in Example 44 to give the title compound having the following physical properties (0.11 g).

- 5 LC/MS  $t_R$  3.84 minutes; MS (ES<sup>+</sup>)  $m/z$  559 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.78 (br. s, 1 H), 10.20 (s, 1 H), 9.71 (s, 1 H), 7.90 - 7.76 (m, 3 H), 7.50 (s, 1 H), 7.48 (d, 1 H), 6.93 (d, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.56 (dd 1 H), 3.35 - 3.23 (obs. m, 1 H), 2.99 (dd, 1 H), 2.93 (t, 2 H), 2.62 - 2.42 (obs. m, 3 H), 2.23 - 2.14 (m, 1 H).

10

Example 88: 6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3,4-dihydro-2(1H)-quinolinone

The same operation as in example Example 51 → Example 52 was conducted from the compound prepared in Example 25 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, 6-(bromoacetyl)-3,4-dihydroquinolin-2(1H)-one was used).

15

- LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.78 - 7.74 (m, 2 H), 7.70 (d, 1 H), 7.68 (d, 1 H), 7.54 (br. s, 1 H), 7.50 (br. d, 1 H), 7.33 (br. s, 1 H), 6.90 (d, 1 H), 6.42 (d, 1 H), 6.05 (dd, 1 H), 5.20 (s, 2 H), 3.01 (t, 2 H), 2.61 (t, 2 H).

20

Example 89: 6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3,4-dihydro-2(1H)-quinolinone

The compound prepared in Example 88 (100 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (38 mg).

25

- LC/MS  $t_R$  3.69 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.86 (s, 1 H), 10.22 (s, 1 H), 9.72 (s, 1 H), 7.87 - 7.77 (m, 3 H), 7.69 (d, 1 H), 7.49 (s 1 H), 7.47 (d, 1 H), 6.92 (d, 1 H), 6.28 (s, 1 H), 5.88 (dd, 1 H), 5.05 (s, 2 H), 2.96 - 2.87 (m, 2 H), 2.57 - 2.47 (obs. m, 2 H).

30

Example 90: methyl N-(4-acetyl-2-fluorophenyl)carbamate

To an N,N-dimethylformamide (9 mL) solution of methyl N-(2-fluoro-4-iodophenyl)carbamate [J. Med. Chem. 48(18), 5823 (2005)] (1.20 g) was added

propane-1,3-diylbis(diphenylphosphane) (0.10 g), palladium(II) diacetate (28 mg) and 3 M aqueous potassium carbonate solution (2.28 mL). The mixture was partitioned equally across four microwavable vessels and all were degassed with nitrogen for 5 minutes. To each reaction mixture, *n*-butyl vinyl ether (0.66 mL) was added and the reaction mixtures were irradiated under microwave conditions (100 W) at 100 °C for 30 minutes. The reaction mixtures were combined and concentrated. The residue was suspended in tetrahydrofuran (12 mL) and 1 M hydrochloric acid (12 mL), stirred for 2 hours at room temperature and extracted with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (2% to 100% ethyl acetate in heptanes) to afford the title compound having the following physical properties (0.49 g). LC/MS  $t_R$  1.57 minutes; MS (ES<sup>+</sup>)  $m/z$  212 (M+H)<sup>a</sup>.

Example 91: methyl N-[4-(2-bromoacetyl)-2-fluorophenyl]carbamate

To a cooled (5 °C) glacial acetic acid (16 mL) solution of the compound prepared in Example 90 (0.49 g) was added a solution of 33 wt. % hydrogen bromide in acetic acid (0.32 mL) followed by pyridinium tribromide (0.73 g) and the mixture stirred at room temperature for 5 hours. To the reaction mixture, a saturated solution of aqueous sodium bicarbonate (100 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and then concentrated. The residue was triturated with a 1 : 1 mixture of ethyl acetate and heptane and the precipitate isolated by filtration to give the title compound having the following physical properties (0.44 g). LC/MS  $t_R$  1.79 minutes; MS (ES<sup>+</sup>)  $m/z$  290 and 292 (M+H)<sup>a</sup>.

Example 92: methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]-2-fluorophenyl]carbamate

The same operation as in Example 84 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 91 was used). LC/MS  $t_R$  3.31 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 9.37 (s, 1 H), 7.81 (br. s, 1 H), 7.77 - 7.73 (m, 2 H),

7.72 - 7.68 (m, 1 H), 7.53 - 7.43 (m, 2 H), 7.42 - 7.34 (m, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H), 5.79 (dd, 1 H), 3.78 (s, 3 H), 3.52 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.52 (app. br. s, 1 H).

5 Example 93: methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-fluorophenyl]carbamate

The compound prepared in Example 92 (160 mg) was treated as detailed in Example 47 to give the title compound having the following physical properties (40 mg). LC/MS  $t_R$  4.19 minutes; MS (ES<sup>+</sup>)  $m/z$  581 (M+H)<sup>b</sup>

10 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.03 (br. s, 1 H), 9.71 (s, 1 H), 9.48 (br. s, 1 H), 7.86 - 7.80 (m, 2 H), 7.80 - 7.72 (m, 2 H), 7.59 - 7.48 (m, 2 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.57 (dd, 1 H), 3.69 (s, 3 H), 3.31 - 3.24 (obs. m, 1 H), 3.00 (ddd, 1 H), 2.58 - 2.54 (obs. m, 1 H), 2.24 - 2.17 (m, 1 H).

15 Example 94: methyl N-[4-(2-bromoacetyl)-3-fluorophenyl]carbamate

The same operation as in Example 77 → Example 90 → Example 91 was conducted from 4-bromo-3-fluoroaniline to give the title compound having the following physical properties.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.86 (t, 1 H), 7.50 (dd, 1 H), 6.97 (dd, 1 H), 6.80 (br. s, 1 H), 4.42 (d, 2 H), 3.75 (s, 3 H).

Example 95: methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-fluorophenyl]carbamate

The same operation as in Example 84 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 94 was used).

LC/MS  $t_R$  3.36 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.25 (s, 1 H), 7.72 (app. br. s, 1 H), 7.65 - 7.61 (m, 2 H), 7.60 - 7.56 (m, 1 H), 7.34 (d, 1 H), 7.19 (br. s, 1 H), 7.06 (app. br. s, 1 H), 6.04 (s, 1 H), 5.98 (s, 1 H), 5.70 (dd, 1 H), 3.65 (s, 3 H), 3.39 - 3.30 (m, 1 H), 3.01 (ddd, 1 H), 2.53 (qd, 1 H), 2.41 (br. s, 1 H).

Example 96: methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-fluorophenyl]carbamate

The compound prepared in Example 95 (125 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (25 mg).

- 5 LC/MS  $t_R$  4.19 minutes; MS (ES<sup>+</sup>)  $m/z$  581 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.81 (br. s, 1 H), 10.09 (br. s, 1 H), 9.72 (s, 1 H), 7.88 - 7.78 (m, 3 H), 7.60 - 7.41 (m, 2 H), 7.31 (d, 1 H), 5.99 (s, 1 H), 5.95 (s, 1 H), 5.59 (dd, 1 H), 3.71 (s, 3 H), 3.30 - 3.23 (m, 1 H), 2.98 (ddd, 1 H), 2.55 - 2.44 (obs. m, 1 H), 2.24 - 2.16 (m, 1 H).

10

Example 97: methyl N-[4-(2-bromoacetyl)-3-methylphenyl]carbamate

The same operation as in Example 77 → Example 90 → Example 91 was conducted from 4-bromo-3-methylaniline to give the title compound having the following physical properties.

- 15 LC/MS  $t_R$  1.83 minutes; MS (ES<sup>+</sup>)  $m/z$  286 and 288 (M+H)<sup>a</sup>.

Example 98: methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-methylphenyl]carbamate

- 20 The same operation as in Example 84 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 97 was used).

- LC/MS  $t_R$  3.07 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.25 (s, 1 H), 7.67 - 7.61 (m, 2 H), 7.58 (d, 1 H),  
25 7.22 (m, 3 H), 6.91 (br. s, 1 H), 6.02 (s, 1 H), 5.99 (s, 1 H), 5.70 (dd, 1 H), 3.64 (s, 3 H), 3.39 - 3.29 (m, 1 H), 3.00 (ddd, 1 H), 2.58 - 2.48 (m, 1 H), 2.38 (app. br. s, 1 H), 2.25 (s, 3 H).

- 30 Example 99: methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-methylphenyl]carbamate

The compound prepared in Example 98 (90 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (48 mg).

LC/MS  $t_R$  3.18 minutes; MS (ES<sup>+</sup>)  $m/z$  577 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.26 (s, 1 H), 7.66 - 7.62 (m, 2 H), 7.58 (d, 1 H), 7.30 (app. s, 1 H), 7.26 (d, 1 H), 7.12 (d, 1 H), 6.03 (s, 1 H), 5.99 (s, 1 H), 5.60 (dd, 1 H), 3.65 (s, 3 H), 3.40 - 3.31 (m, 1 H), 3.01 (ddd, 1 H), 2.54 (qd, 1 H), 2.37 - 2.30 (m, 1 H), 2.15 (s, 3 H).

5

Example 100: methyl N-(3-ethylphenyl)carbamate

3-Ethylaniline (4.0 g) was treated as detailed in Example 77 to give the title compound having the following physical properties (5.9 g).

LC/MS *t*<sub>R</sub> 1.90 minutes; MS (ES<sup>+</sup>) *m/z* 180 (M+H)<sup>a</sup>.

10

Example 101: methyl N-(4-acetyl-3-ethylphenyl)carbamate

To a dichloroethane (7 mL) solution of the compound prepared in Example 100 (0.50 g) was added acetyl chloride (0.21 mL) and aluminium trichloride (0.93 g) and the mixture stirred at room temperature overnight. To the reaction mixture, ice-water (5 mL) was cautiously added followed by extraction into dichloromethane. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (7% to 60% ethyl acetate in heptanes) to afford the title compound having the following physical properties (0.43 g).

15

LC/MS *t*<sub>R</sub> 2.13 minutes; MS (ES<sup>+</sup>) *m/z* 222 (M+H)<sup>a</sup>.

20

Example 102: methyl N-[4-(2-bromoacetyl)-3-ethylphenyl]carbamate

The compound prepared in Example 101 (0.41 g) was treated as detailed in Example 91 to give the title compound having the following physical properties (0.41 g). LC/MS *t*<sub>R</sub> 1.96 minutes; MS (ES<sup>+</sup>) *m/z* 300 and 302 (M+H)<sup>a</sup>.

25

Example 103: methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-ethylphenyl]carbamate

The same operation as in Example 84 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 102 was used).

30

LC/MS *t*<sub>R</sub> 3.18 minutes; MS (ES<sup>+</sup>) *m/z* 557 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.77 - 7.72 (m, 2 H), 7.69 (d, 1 H), 7.45 - 7.26 (m, 3 H), 7.11 - 6.84 (m, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.82 (dd, 1 H), 3.76 (s, 3 H), 3.51 - 3.40 (m, 1 H), 3.12 (dd, 1 H), 2.72 (q, 2 H), 2.69 - 2.59 (m, 1 H), 2.51 (app. br. s, 1 H), 1.15 (app. br. s, 3 H).

5

Example 104: methyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])}-1H-imidazol-5-yl)-3-ethylphenyl]carbamate

The compound prepared in Example 103 (40 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (26.7 mg).

10

LC/MS *t<sub>R</sub>* 4.21 minutes; MS (ES<sup>+</sup>) *m/z* 591 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 7.78 - 7.74 (m, 2 H), 7.70 (d, 1 H), 7.45 (app. s, 1 H), 7.39 (dd, 1 H), 7.19 (d, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.70 (dd, 1 H), 3.77 (s, 3 H), 3.51 - 3.42 (m, 1 H), 3.12 (ddd, 1 H), 2.71 - 2.62 (m, 1 H), 2.59 (q, 2 H), 2.47 - 2.40 (m, 1 H), 1.10 (t, 3 H).

15

Example 105: methyl N-[4-(2-bromoacetyl)-3-chlorophenyl]carbamate

The same operation as in Example 77 → Example 90 → Example 91 was conducted from 4-bromo-3-chloroaniline to give the title compound having the following physical properties.

20

LC/MS *t<sub>R</sub>* 1.89 minutes; MS (ES<sup>+</sup>) *m/z* 306 (M+H)<sup>a</sup>.

Example 106: methyl [3-chloro-4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])}-1H-imidazol-5-yl)phenyl]carbamate

25

The same operation as in Example 84 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 105 was used).

LC/MS *t<sub>R</sub>* 3.38 minutes; MS (ES<sup>+</sup>) *m/z* 563 (M+H)<sup>b</sup>

30

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 7.77 - 7.73 (m, 3 H), 7.73 - 7.66 (m, 2 H), 7.53 (app. br. s, 1 H), 7.36 (app. br. s, 1 H), 6.16 (s, 1 H), 6.10 (s, 1 H), 5.83 (dd, 1 H), 3.77 (s, 3 H), 3.53 - 3.42 (m, 1 H), 3.12 (ddd, 1 H), 2.70 - 2.60 (m, 1 H), 2.54 (app. br. s, 1 H).



Example 107: methyl {3-chloro-4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 25 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 105 was used).

LC/MS  $t_R$  3.03 minutes; MS (ES<sup>+</sup>)  $m/z$  537 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.28 (br. s, 1 H), 9.85 (br. s, 1 H), 9.71 (s, 1 H), 7.91 (d, 1 H), 7.86 - 7.77 (m, 3 H), 7.73 (d, 1 H), 7.63 (app. br. s, 1 H), 7.58 (s, 1 H), 7.41 (dd, 1 H), 6.30 (app. br. s, 1 H), 5.87 (dd, 1 H), 5.11 (s, 2 H), 3.68 (s, 3 H).

Example 108(1) and Example 108(2): methyl N-[4-(2-bromoacetyl)-3-methoxyphenyl]carbamate and methyl N-[2-bromo-4-(2-bromoacetyl)-5-methoxyphenyl]carbamate

The same operation as in Example 77 → Example 90 → Example 91 was conducted from 4-bromo-3-methoxyaniline to give a 3 : 5 mixture of the title compounds having the following physical properties.

Example 108(1):

LC/MS  $t_R$  1.82 minutes; MS (ES<sup>+</sup>)  $m/z$  302 and 304 (M+H)<sup>a</sup>.

Example 108(2):

LC/MS  $t_R$  2.10 minutes; MS (ES<sup>+</sup>)  $m/z$  379, 381, 383 (M+H)<sup>a</sup>.

Example 109(1) and Example 109(2): methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-3-methoxyphenyl]carbamate and methyl [2-bromo-4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])1H-imidazol-5-yl)-5-methoxyphenyl]carbamate

The same operation as in Example 84 → Example 39 was conducted from the compound prepared in Example 17 to give the title compounds in a 2 : 1 ratio having

the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the 3: 5 mixture of the compounds prepared in Example 108 was used).

Example 109(1):

- 5 LC/MS  $t_R$  3.12 minutes; MS (ES<sup>+</sup>)  $m/z$  559 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.97 (s, 1 H), 9.70 (s, 1 H), 9.69 (s, 1 H), 7.88 - 7.78 (m, 4 H), 7.31 (d, 1 H), 7.27 (s, 1 H), 7.04 (dd, 1 H), 5.99 (s, 1 H), 5.96 (s, 1 H), 5.63 (dd, 1 H), 3.83 (s, 3 H), 3.66 (s, 3 H), 3.30 - 3.21 (m, 1 H), 3.00 (dd, 1 H), 2.48 - 2.41 (m, 1 H), 2.39 - 2.32 (m, 1 H).

10

Example 109(2):

- LC/MS  $t_R$  3.59 minutes; MS (ES<sup>+</sup>)  $m/z$  637 and 639 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.15 (s, 1 H), 9.70 (s, 1 H), 8.89 (s, 1 H), 8.06 (s, 1 H), 7.85 - 7.77 (m, 3 H), 7.45 (d, 1 H), 7.25 (s, 1 H), 5.99 (s, 1 H), 5.95 (s, 1 H), 5.64 (dd, 1 H), 3.87 (s, 3 H), 3.66 (s, 3 H), 3.45 - 3.38 (m, 1 H), 3.02 (dd, 1 H), 2.54 - 2.46 (obs. m, 1 H), 2.43 - 2.35 (m, 1 H).

15

Example 110: methyl N-[4-(2-bromoacetyl)-3-(trifluoromethyl)phenyl]carbamate

- The same operation as in Example 77 → Example 90 → Example 91 was  
20 conducted from 4-bromo-3-trifluoromethylaniline to give the title compound having the following physical properties.

LC/MS  $t_R$  1.38 minutes; MS (ES<sup>+</sup>)  $m/z$  340 and 342 (M+H)<sup>a</sup>.

- Example 111: methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-(trifluoromethyl)phenyl]carbamate

25

- The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 110 was used).

30

LC/MS  $t_R$  3.59 minutes; MS (ES<sup>+</sup>)  $m/z$  597 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.95 (br. s, 1 H), 7.78 - 7.72 (m, 2 H), 7.73 - 7.66 (m, 2 H), 7.62 - 7.50 (m, 1 H), 7.13 (br. s, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H),


5.82 (dd, 1 H), 3.78 (s, 3 H), 3.53 - 3.40 (m, 1 H), 3.16 - 3.07 (m, 1 H), 2.70 - 2.59 (m, 1 H), 2.53 (app. br. s, 1 H).

Example 112: methyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-(trifluoromethyl)phenyl]carbamate

The compound prepared in Example 111 (50 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (25 mg). LC/MS  $t_R$  4.29 minutes; MS (ES<sup>+</sup>)  $m/z$  631 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.03 (s, 1 H), 7.78 - 7.74 (m, 3 H), 7.70 (d, 1 H), 7.43 (d, 1 H), 6.16 (s, 1 H), 6.11 (s, 1 H), 5.72 (dd, 1 H), 3.80 (s, 3 H), 3.48 - 3.39 (m, 1 H), 3.12 (ddd, 1 H), 2.72 - 2.60 (m, 1 H), 2.45 - 2.36 (m, 1 H).

Example 113: 4-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-4-yl)benzoic acid ammoniate



The same operation as in Example 84 → Example 39 → Example 37 → Example 55 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, methyl 4-(2-bromoacetyl)benzoate [J. Am. Chem. Soc. 122(39), 9361 (2000)] was used. In the step corresponding to Example 41 in the operation, high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) was used to give the title product as the ammonium salt.

LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.70 (s, 1 H), 7.89 (d, 2 H), 7.85 - 7.72 (m, 5 H), 7.64

(s, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.64 (dd, 1 H), 3.45 - 3.30 (obs. m, 1 H), 3.00 (dd, 1 H), 2.58 - 2.45 (obs. m, 1 H), 2.34 (app. br. s, 1 H).

5 Example 114: 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)benzamide

To an N,N-dimethylformamide (2 mL) solution of the compound prepared in Example 113 (88 mg) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (87 mg), ammonium chloride (47 mg) and N,N-diisopropylethylamine (145  $\mu$ L) and the mixture stirred at room temperature for 2  
10 hours. To the reaction mixture, water was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue obtained on concentration was purified by high performance liquid chromatography (5 to 100% acetonitrile in water) to give the title compound having the following physical properties (33.2 mg).

15 LC/MS  $t_R$  2.89 minutes; MS ( $ES^+$ )  $m/z$  499 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.22 (s, 1 H), 9.68 (s, 1 H), 7.90 (br. s, 1 H), 7.84 (d, 2 H), 7.81 - 7.77 (m, 3 H), 7.76 (d, 2 H), 7.63 (s, 1 H), 7.24 (br. s, 1 H), 5.96 (s, 1 H), 5.95 (s, 1 H), 5.64 (d, 1 H), 3.48 - 3.33 (m, 1 H), 3.00 (dd, 1 H), 2.56 - 2.47 (obs. m, 1 H), 2.39 - 2.30 (m, 1 H).

20

Example 115: 4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)benzamide

The compound prepared in Example 114 (88 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (9.9  
25 mg).

LC/MS  $t_R$  3.55 minutes; MS ( $ES^+$ )  $m/z$  533 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.12 (s, 1 H), 9.71 (s, 1 H), 8.01 (s, 1 H), 7.97 (d, 2 H), 7.83 - 7.78 (m, 5 H), 7.41 (s, 1 H), 6.00 (s, 1 H), 5.95 (s, 1 H), 5.59 (dd, 1 H), 3.35 - 3.22 (m, 1 H), 3.00 (dd, 1 H), 2.60 - 2.48 (obs. m, 1 H), 2.21 (t, 1 H).

30

Example 116: tert-butyl N-(4-chloro-2-{3-[4-(4-cyano-3-fluorophenyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, 4-(2-bromoacetyl)-2-fluorobenzonitrile [WO2007/070826, page 121] was used).

5 LC/MS  $t_R$  2.27 minutes; MS ( $ES^+$ )  $m/z$  546 (M+H) <sup>a</sup>.

Example 117: tert-butyl N-(2-{3-[4-(4-carbamoylphenyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}-4-chlorophenyl)carbamate

To a dimethylsulfoxide (6 mL) solution of the compound prepared in Example  
10 116 (0.106 g) was sequentially added potassium carbonate (13 mg) and 30% aqueous hydrogen peroxide solution (0.13 mL) and the mixture stirred at 50 °C for 16 hours. To the reaction mixture, a 10% aqueous solution of lithium chloride (3 mL) was added and the mixture stirred at room temperature for 15 minutes followed by extraction into ethyl acetate. The combined organic layers were sequentially washed with a saturated  
15 aqueous solution of sodium hydrogen carbonate and brine, dried and concentrated to obtain the crude title compound having the following physical properties (0.156 g).  
LC/MS  $t_R$  1.83 minutes; MS ( $ES^+$ )  $m/z$  564 (M+H) <sup>a</sup>.

Example 118: 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-2-fluorobenzamide

The same operation as in Example 40 → Example 41 was conducted from the compound prepared in Example 117 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.30 minutes; MS ( $ES^+$ )  $m/z$  517 (M+H) <sup>b</sup>  
25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.85 (t, 1 H), 7.78 - 7.73 (m, 2 H), 7.70 (d, 1 H), 7.66 - 7.52 (m, 3 H), 6.15 (s, 1 H), 6.11 (s, 1 H), 5.80 (dd, 1 H), 3.56 - 3.42 (m, 1 H), 3.14 (ddd, 1 H), 2.73 - 2.61 (m, 1 H), 2.58 - 2.44 (m, 1 H).

Example 119: 4-(2-bromoacetyl)benzamide

30 4-Acetylbenzamide [J. Org. Chem., 65(26), 9103 (2000)] (0.25 g) was treated as detailed in Example 91 to give the title compound having the following physical properties (0.37 g).

LC/MS  $t_R$  1.33 minutes; MS ( $ES^+$ )  $m/z$  242 and 244 (M+H) <sup>a</sup>.

Example 120: 4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]benzamide

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 25 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 119 was used).

LC/MS  $t_R$  2.88 minutes; MS (ES<sup>+</sup>)  $m/z$  473 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.90 (d, 2 H), 7.80 (d, 2 H), 7.77 - 7.72 (m, 2 H), 7.70 (d, 1 H), 7.68 (d, 1 H), 7.55 (s, 1 H), 6.41 (d, 1 H), 6.04 (dd, 1 H), 5.22 (s, 2 H).

Example 121: 4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]benzamide

The compound prepared in Example 120 (50 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (16 mg).

LC/MS  $t_R$  3.52 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.13 (br. s, 1 H), 9.72 (s, 1 H), 7.99 (br. s, 1 H), 7.93 (d, 2 H), 7.85 - 7.76 (m, 5 H), 7.69 (d, 1 H), 7.38 (br. s, 1 H), 6.26 (app. s, 1 H), 5.88 (dd, 1 H), 5.05 (s, 2 H).

Example 122: 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-4-yl)-N-methylbenzamide

The compound prepared in Example 113 (88 mg) was treated as detailed in Example 114 using methylamine hydrochloride instead of ammonium chloride to give the title compound having the following physical properties (46.8 mg).

LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.23 (s, 1 H), 9.69 (s, 1 H), 8.36 (q, 1 H), 7.91 - 7.68 (m, 7 H), 7.63 (s, 1 H), 5.98 (s, 1 H), 5.97 (s, 1 H), 5.64 (dd, 1 H), 3.44 - 3.33 (m, 1 H), 3.02 (dd, 1 H), 2.78 (d, 3 H), 2.57 - 2.48 (obs. m, 1 H), 2.38 - 2.31 (m, 1 H).

Example 123: 4-(5-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-4-yl)-N-methylbenzamide

The compound prepared in Example 122 (90 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (11.1 mg).

LC/MS  $t_R$  3.72 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.10 (s, 1 H), 9.71 (s, 1 H), 8.48 (q, 1 H), 7.93 (d, 2 H), 7.87 - 7.72 (m, 5 H), 6.00 (s, 1 H), 5.95 (s, 1 H), 5.59 (dd, 1 H), 3.36 - 3.22 (obs. m, 1 H), 2.80 (d, 3 H), 2.59 - 2.50 (obs. m, 1 H), 2.39 - 2.12 (m, 2 H).

10 Example 124: 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-4-yl)-N-ethylbenzamide

The compound prepared in Example 113 (88 mg) was treated as detailed in Example 114 using ethylamine hydrochloride instead of ammonium chloride to give the title compound having the following physical properties (48.6 mg).

LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  527 (M+H)<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.23 (s, 1 H), 9.70 (s, 1 H), 8.40 (t, 1 H), 7.90 - 7.69 (m, 7 H), 7.64 (s, 1 H), 5.98 (s, 1 H); 5.97 (s, 1 H), 5.64 (dd, 1 H), 3.45 - 3.33 (m, 1 H), 3.32 - 3.23 (m, 2 H), 3.01 (dd, 1 H), 2.56 - 2.48 (obs. m, 1 H), 2.38 - 2.30 (m, 1 H), 1.13 (t, 3 H).

20 Example 125: 4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-N-ethylbenzamide

The compound prepared in Example 124 (93 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (24.9 mg).

- 25 LC/MS  $t_R$  3.96 minutes; MS (ES<sup>+</sup>)  $m/z$  561 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.11 (s, 1 H), 9.71 (s, 1 H), 8.51 (t, 1 H), 7.94 (d, 2 H), 7.87 - 7.68 (m, 5 H), 6.00 (s, 1 H), 5.95 (s, 1 H), 5.59 (dd, 1 H), 3.40 - 3.23 (obs. m, 3 H), 3.00 (dd, 1 H), 2.58 - 2.46 (obs. m, 1 H), 2.20 (t, 1 H), 1.14 (t, 3 H).

30 Example 126: 7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-nitrophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizine

The same operational sequence as in Example 38 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the

title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, 2-bromo-1-(4-nitrophenyl)-ethanone was used).

LC/MS  $t_R$  4.03 minutes; MS (ES<sup>+</sup>)  $m/z$  501 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.49 (br. s, 1 H), 9.70 (s, 1 H), 8.21 (d, 2 H), 7.95 (d, 2 H), 7.89 - 7.77 (m, 4 H), 5.99 (s, 1 H), 5.97 (s, 1 H), 5.65 (dd, 1 H), 3.44 - 3.25 (obs. m, 1 H), 3.02 (dd, 1 H), 2.58 - 2.42 (obs. m, 1 H), 2.36 - 2.25 (m, 1 H).

10 Example 127: 3-[5-(4-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 126 (1.10 g) was treated as detailed in Example 74 to give the title compound having the following physical properties (0.86 g).

LC/MS  $t_R$  2.74 minutes; MS (ES<sup>+</sup>)  $m/z$  471 (M+H)<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.73 (m, 2 H), 7.70 (d, 1 H), 7.39 (app. br. s, 2 H), 7.06 (br. s, 1 H), 6.75 (d, 2 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.79 (dd, 1 H), 3.52 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.69 - 2.58 (m, 1 H), 2.47 (app. br. s, 1 H).

20 Example 128: ethyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]carbamate

- To a cooled (0 °C) dichloromethane (5 mL) solution of the compound prepared in Example 127 (130 mg), pyridine (45  $\mu$ L) and ethyl chloroformate (26  $\mu$ L) were added sequentially and the mixture stirred overnight at room temperature. The reaction mixture was concentrated and dissolved in methanol (5 mL), concentrated aqueous ammonia solution (0.21 mL) was added and the mixture stirred 3 hours at room temperature. The reaction mixture was concentrated, the residue suspended in water (3 mL) and extracted into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (25% to 100% ethyl acetate in heptanes followed by 1% to 3% methanol in ethyl acetate) to afford the title compound having the following physical properties (76 mg).

LC/MS  $t_R$  3.21 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.72 (m, 2 H), 7.70 (d, 1 H), 7.65 - 7.40 (m, 4 H), 7.36 - 7.10 (m, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H), 5.80 (dd, 1 H),




4.20 (q, 2 H), 3.52 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.50 (app. br. s, 1 H), 1.33 (t, 3 H).

Example 129(1) to Example 129(6)

- 5 The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 127 and the corresponding acid chlorides, chloroformates, sulfonyl chlorides or acetic anhydride using the method as detailed in Example 128.

10 Example 129(1): isopropyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate



LC/MS  $t_R$  3.36 minutes; MS (ES<sup>+</sup>)  $m/z$  557 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.73 (m, 2 H), 7.70 (d, 1 H),  
 15 7.65 - 7.38 (m, 4 H), 7.30 - 7.10 (m, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H), 5.80 (dd, 1 H),  
 5.02 - 4.94 (m, 1 H), 3.51 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.71 - 2.61 (m, 1 H), 2.52 (br.  
 s, 1 H), 1.32 (d, 6 H).

20 Example 129(2): 2-methoxyethyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  573 (M+H), 545 (M-N<sub>2</sub>+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.79 - 7.72 (m, 2 H), 7.72 - 7.67 (m, 1  
 H), 7.59 (app. br. s, 2 H), 7.46 (app. br. s, 2 H), 7.37 - 7.07 (m, 1 H), 6.14 (s, 1 H), 6.10  
 (s, 1 H), 5.80 (d, 1 H), 4.36 - 4.19 (m, 2 H), 3.72 - 3.61 (m, 2 H), 3.52 - 3.43 (m, 1 H),  
 25 3.41 (s, 3 H), 3.17 - 3.05 (m, 1 H), 2.71 - 2.59 (m, 1 H), 2.49 (app. br. s, 1 H).

Example 129(3): N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-

tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]acetamide

LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.77 - 7.73 (m, 2 H), 7.70 (d, 1 H),  
7.64 - 7.55 (m, 4 H), 7.27 (br. s, 1 H), 6.15 (s, 1 H), 6.11 (s, 1 H), 5.80 (dd, 1 H), 3.52 -  
5 3.43 (m, 1 H), 3.12 (ddd, 1 H), 2.71 - 2.59 (m, 1 H), 2.53 - 2.44 (m, 1 H), 2.14 (s, 3 H).

Example 129(4): N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-2-methoxyacetamide

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>b</sup>

10 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.04 (s, 1 H), 9.72 (s, 1 H), 9.68 (s, 1 H), 7.84 - 7.77  
(m, 3 H), 7.68 (d, 2 H), 7.58 (d, 2 H), 7.43 (s, 1 H), 5.99 (s, 1 H), 5.96 (s, 1 H), 5.62 (dd,  
1 H), 3.99 (s, 2 H), 3.38 (s, 3 H), 3.38 - 3.30 (obs. m, 1 H), 3.00 (dd, 1 H), 2.55 - 2.44  
(obs. m, 1 H), 2.29 - 2.39 (m, 1 H).

15 Example 129(5): N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-3-methoxypropanamide

LC/MS  $t_R$  3.03 minutes; MS (ES<sup>+</sup>)  $m/z$  579 (M+Na), 557 (M+H), 529 (M-N<sub>2</sub>+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.78 - 7.72 (m, 2 H), 7.71 - 7.66 (m, 1  
H), 7.59 (app. br. s, 4 H), 7.27 (br. s, 1 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.79 (dd, 1 H),  
20 3.74 (t, 2 H), 3.53 - 3.40 (m, 1 H), 3.37 (s, 3 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 3 H),  
2.49 (app. br. s, 1 H).

Example 129(6): N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]methanesulfonamide

25 LC/MS  $t_R$  2.98 minutes; MS (ES<sup>+</sup>)  $m/z$  549 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.25 (s, 1 H), 7.66 - 7.61 (m, 2 H), 7.60 - 7.44 (m, 3  
H), 7.22 (br. s, 1 H), 7.15 (d, 2 H), 6.03 (s, 1 H), 5.99 (s, 1 H), 5.68 (dd, 1 H), 3.40 -  
3.29 (m, 1 H), 3.01 (ddd, 1 H), 2.86 (s, 3 H), 2.59 - 2.49 (m, 1 H), 2.38 (app. br. s, 1 H).

30 Example 130(1) to Example 130(4)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 128, 129(1), 129(3) or 129(6)

using the method as detailed in Example 44.

Example 130(1): ethyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate



5

LC/MS  $t_R$  4.28 minutes; MS ( $ES^+$ )  $m/z$  577 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.86 (br. s, 1 H), 9.76 (s, 1 H), 9.71 (s, 1 H), 7.85 - 7.80 (m, 3 H), 7.60 (d, 2 H), 7.55 (d, 2 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.57 (dd, 1 H), 4.15 (q, 2 H), 3.30 - 3.24 (obs. m, 1 H), 3.03 - 2.95 (m, 1 H), 2.58 - 2.54 (obs. m, 1 H), 2.23 - 2.17 (m, 1 H), 1.26 (t, 3 H).

10

Example 130(2): isopropyl [4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  4.46 minutes; MS ( $ES^+$ )  $m/z$  591 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.86 (br. s, 1 H), 9.71 (app. s, 2 H), 7.86 - 7.78 (m, 3 H), 7.64 - 7.52 (m, 4 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.57 (dd, 1 H), 4.91 (quintet, 1 H), 3.30 - 3.25 (m, 1 H), 2.99 (ddd, 1 H), 2.58 - 2.54 (obs. m, 1 H), 2.20 (t, 1 H), 1.27 (d, 6 H).

15

Example 130(3): N-[4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide

LC/MS  $t_R$  3.85 minutes; MS ( $ES^+$ )  $m/z$  547 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.88 (br. s, 1 H), 10.06 (s, 1 H), 9.71 (s, 1 H), 7.89 - 7.75 (m, 3 H), 7.72 - 7.58 (m, 4 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.57 (dd, 1 H), 3.30 - 3.24 (m, 1 H), 2.99 (dd, 1 H), 2.59 - 2.55 (obs. m, 1 H), 2.25 - 2.16 (m, 1 H), 2.06 (s, 3 H).

25

Example 130(4): N-[4-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -1H-imidazol-5-yl)phenyl]methanesulfonamide

LC/MS  $t_R$  3.93 minutes; MS (ES<sup>+</sup>)  $m/z$  583 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.92 (br. s, 1 H), 9.91 (br. s, 1 H), 9.71 (s, 1 H), 7.88  
5 - 7.77 (m, 3 H), 7.70 - 7.62 (m, 2 H), 7.29 (d, 2 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.58 (dd,  
1 H), 3.30 - 3.25 (m, 1 H), 3.04 (s, 3 H), 3.03 - 2.96 (m, 1 H), 2.57 - 2.54 (obs. m, 1 H),  
2.21 (dd, 1 H).

Example 131: 1-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -1H-imidazol-5-yl)phenyl]-3-ethylurea

To a dichloromethane (7 mL) solution of the compound prepared in Example  
127 (130 mg) was added ethyl isocyanate (20  $\mu$ L) and the mixture stirred at room  
temperature for 3 hours. To the reaction mixture, further ethyl isocyanate (20  $\mu$ L) was  
added and the mixture stirred at room temperature for 72 hours. One further treatment  
15 with ethyl isocyanate (20  $\mu$ L) was required, and the mixture stirred at room temperature  
for 4 hours. The precipitate was collected by filtration and purified by high performance  
liquid chromatography (5 to 100% acetonitrile in water) to give the title compound  
having the following physical properties (28 mg).

LC/MS  $t_R$  2.95 minutes; MS (ES<sup>+</sup>)  $m/z$  542 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.77 - 7.73 (m, 2 H), 7.70 (d, 1 H),  
20 7.55 (app. br. s, 2 H), 7.39 (d, 2 H), 7.21 (br. s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.79  
(dd, 1 H), 3.52 - 3.41 (m, 1 H), 3.25 (q, 2 H), 3.12 (ddd, 1 H), 2.72 - 2.59 (m, 1 H), 2.52  
- 2.45 (m, 1 H), 1.17 (t, 3 H).

Example 132: 3-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -1H-imidazol-5-yl)phenyl]-1,1-dimethylurea

To a cooled (0 °C) dichloromethane (7 mL) solution of the compound prepared in  
Example 127 (130 mg) was sequentially added pyridine (45  $\mu$ L) and dimethylcarbamic  
chloride (21  $\mu$ L) and the mixture stirred at room temperature for 3 hours. To the  
30 reaction mixture, further dimethylcarbamic chloride (21  $\mu$ L) was added and the mixture  
stirred at room temperature for 72 hours. One further treatment with dimethylcarbamic  
chloride (21  $\mu$ L) was required, and the mixture stirred at room temperature for 4 hours.  
The reaction mixture was concentrated and purified by high performance liquid

chromatography (5 to 100% acetonitrile in water) to give the title compound having the following physical properties (32 mg).

LC/MS  $t_R$  2.93 minutes; MS (ES<sup>+</sup>)  $m/z$  542 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.75 - 7.70 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.54 (d, 2 H), 7.38 (d, 2 H), 7.20 (br. s, 1 H), 6.13 (s, 1 H), 6.08 (s, 1 H), 5.78 (d, 1 H), 3.49 - 3.39 (m, 1 H), 3.14 - 3.05 (m, 1 H), 3.02 (s, 6 H), 2.68 - 2.57 (m, 1 H), 2.51 - 2.42 (m, 1 H).

Example 133: N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-2-(dimethylamino)acetamide

To an N,N-dimethylformamide (6 mL) and pyridine (2 mL) solution of the compound prepared in Example 127 (100 mg) and N,N-dimethylglycine (26 mg) was added 1-[3-(dimethylaminopropyl)]-3-ethylcarbodiimide hydrochloride (49 mg) and the mixture stirred at room temperature for 16 hours. The reaction was concentrated and the residue suspended in water (15 mL) followed by extraction into a 3: 1 mixture of ethyl acetate and propan-2-ol. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (21 mg).

LC/MS  $t_R$  2.52 minutes; MS (ES<sup>+</sup>)  $m/z$  279 [(M+2H)/2]<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.77 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.63 (app. br. s, 4 H), 7.29 (br. s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.80 (dd, 1 H), 3.53 - 3.38 (m, 1 H), 3.16 (s, 2 H), 3.14 - 3.08 (m, 1 H), 2.70 - 2.59 (m, 1 H), 2.55 - 2.43 (m, 1 H), 2.39 (s, 6 H).

25

Example 134: 3-{[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-3-oxopropanoic acid methyl ester

To a cooled (0°C) dichloromethane (1 mL) solution of the compound prepared in Example 127 (54 mg) was sequentially added pyridine (11  $\mu$ L) and methyl-3-chloro-3-oxopropanoate (20  $\mu$ L) and the mixture stirred for 3 hours at room temperature. To the reaction mixture, water (5 mL) was added followed by extraction into dichloromethane. The combined organic layers were sequentially washed with 1 M hydrochloric acid, a

30

saturated aqueous solution of sodium hydrogen carbonate and brine, dried and concentrated. The residue thus obtained was triturated with a 1: 4 mixture of ethyl acetate in heptanes, the resultant precipitate being isolated by filtration to give the title compound having the following physical properties (56 mg).

- 5 LC/MS  $t_R$  1.21 minutes; MS (ES<sup>+</sup>)  $m/z$  571 (M+H)<sup>a</sup>.

Example 135: 3-{[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-3-oxopropanoic acid

To a methanol (0.5 mL) solution of the compound prepared in Example 134 (56  
10 mg) was added a solution of sodium carbonate (10.4 mg) in water (0.5 mL) and the mixture stirred at room temperature for 16 hours. Further sodium carbonate (10.4 mg) was then added and the mixture stirred at room temperature for 2 hours. The reaction mixture was concentrated and the residue suspended in 1 M hydrochloric acid (0.4 mL) followed by extraction into ethyl acetate. The combined organic layers were washed  
15 with brine, dried and concentrated. The residue thus obtained was triturated with a 1: 4 mixture of ethyl acetate in heptanes, the resultant precipitate being isolated by filtration to give the title compound having the following physical properties (53 mg).

LC/MS  $t_R$  2.84 minutes; MS (ES<sup>+</sup>)  $m/z$  557 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.59 (s, 1 H), 12.05 (s, 1 H), 9.69 (s, 1 H), 7.80 (app.  
20 s, 3 H), 7.59 (d, 2 H), 7.50 (d, 2 H), 7.40 (s, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.62 (d, 1 H), 3.44 - 3.30 (obs. m, 1 H), 3.05 - 2.92 (m, 1 H), 2.83 (s, 2 H), 2.49 - 2.42 (obs. m, 1 H), 2.38 - 2.32 (m, 1 H).

Example 136: 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-4-hydroxy-2(1H)-quinolinone

The compound prepared in Example 135 (31 mg), was suspended in Eaton's reagent (0.21 mL) and the mixture stirred at 100 °C for 2 hours. To the cooled (0 °C) reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (5 mL) was carefully added followed by extraction into ethyl acetate. The combined organic  
30 layers were washed with brine, dried and concentrated and the residue purified by high performance liquid chromatography (5 to 100% acetonitrile in water). The residue was purified further by trituration from a boiling 9 : 1 solution of dichloromethane and methanol. The resultant precipitate was isolated by filtration to give the title compound

having the following physical properties (2.8 mg).

LC/MS  $t_R$  2.88 minutes; MS (ES<sup>+</sup>)  $m/z$  539 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.30 (br. s, 1 H), 11.14 (br. s, 1 H), 9.70 (s, 1 H),  
8.08 (s, 1 H), 7.87 - 7.76 (m, 4 H), 7.51 (br. s, 1 H), 7.24 (br. s, 1 H), 6.53 (br. s, 1 H),  
5 5.99 (s, 1 H), 5.96 (s, 1 H), 5.74 (br. s, 1 H), 5.64 (d, 1 H), 3.36 - 3.27 (obs. m, 1 H),  
3.02 (dd, 1 H), 2.54 - 2.45 (obs. m, 1 H), 2.40 - 2.32 (m, 1 H).

Example 137: 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-2-fluorobenzonitrile

10 The same operation as in Example 40 → Example 41 was conducted from the compound prepared in Example 116 to give the title compound having the following physical properties.

LC/MS  $t_R$  4.12 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.43 (s, 1 H), 7.60 - 7.38 (m, 6 H), 7.29 (s, 1 H), 6.21 (s,  
15 1 H), 5.72 (d, 1 H), 5.65 (s, 1 H), 3.41 - 3.24 (m, 2 H), 3.02 - 2.91 (m, 1 H), 2.49 - 2.31 (m, 1 H).

Example 138: 3-[5-(3-amino-1H-indazol-6-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

20 To a methanol (1 mL) solution of the compound prepared in Example 137 (125 mg) was added hydrazine hydrate (105  $\mu$ L) and the mixture stirred at 80 °C for 20 hours. To the reaction mixture, further hydrazine hydrate (105  $\mu$ L) was added and the mixture stirred at 80 °C for 24 hours. To the reaction mixture, water (5 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine,  
25 dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (6.5 mg).

LC/MS  $t_R$  2.29 minutes; MS (ES<sup>+</sup>)  $m/z$  511 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.39 (s, 1 H), 7.79 - 7.56 (m, 5 H), 7.38 (br. s, 1 H),  
30 7.32 (d, 1 H), 6.16 (s, 1 H), 6.12 (s, 1 H), 5.81 (dd, 1 H), 3.55 - 3.42 (m, 1 H), 3.14 (ddd, 1 H), 2.74 - 2.59 (m, 1 H), 2.52 - 2.49 (m, 1 H).

Example 139: 7-acetyl-3,4-dihydro-2H-1,4-benzoxazin-3-one

To an N,N-dimethylformamide (5 mL) solution of 7-bromo-2H-1,4-benzoxazin-3(4H)-one (0.50 g) was added tributyl(1-ethoxyethenyl)stannane (0.81 mL). The mixture was degassed with nitrogen for 5 minutes and tetrakis(triphenylphosphine)palladium(0) (51 mg) was added. The mixture was stirred at 120 °C in a pressure tube for 3 hours. To the reaction mixture, 1 M hydrochloric acid was added and the mixture was stirred for 2 hours then extracted into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was triturated with hot cyclohexane and the precipitate isolated by filtration to give the title compound having the following physical properties (0.35 g).

LC/MS  $t_R$  1.30 minutes; MS ( $ES^+$ )  $m/z$  192 (M+H)<sup>a</sup>.

Example 140: 7-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-2H-1,4-benzoxazin-3(4H)-one

The same operation as in Example 91 → Example 84 → Example 39 was conducted from the compound prepared in Example 139 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 17 was used).

LC/MS  $t_R$  2.90 minutes; MS ( $ES^+$ )  $m/z$  527 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.07 (s, 1 H), 10.66 (s, 1 H), 9.68 (s, 1 H), 7.85 - 7.75 (m, 3 H), 7.44 (br. s, 1 H), 7.29 (d, 1 H), 7.25 (s, 1 H), 6.85 (d, 1 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.60 (dd, 1 H), 4.55 (s, 2 H), 3.45 - 3.34 (obs. m, 1 H), 2.99 (dd, 1 H), 2.53 - 2.43 (obs. m, 1 H), 2.37 - 2.29 (m, 1 H).

Example 141: 7-bromo-4-methyl-3,4-dihydro-2H-1,4-benzoxazin-3-one

To an N,N-dimethylformamide (15 mL) solution of 7-bromo-2H-1,4-benzoxazin-3(4H)-one (1.0 g) was sequentially added potassium carbonate (1.21 g) and iodomethane (1.0 mL) and the mixture stirred under nitrogen for 3 hours. To the reaction mixture, water (25 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to afford the title compound having the following physical properties (1.1 g).

LC/MS  $t_R$  1.93 minutes; MS ( $ES^+$ )  $m/z$  242 and 244 (M+H)<sup>a</sup>.



Example 142: 7-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-4-methyl-2H-1,4-benzoxazin-3(4H)-one

The same operation as in Example 139 → Example 91 → Example 84 → Example 52 was conducted from the compound prepared in Example 141 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 9 was used).

LC/MS  $t_R$  3.15 minutes; MS (ES<sup>+</sup>)  $m/z$  541 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.12 (br. s, 1 H), 9.69 (s, 1 H), 7.83 - 7.77 (m, 3 H), 7.50 (br. s, 1 H), 7.40 (d, 1 H), 7.31 (d, 1 H), 7.11 (br. s, 1 H), 5.97 (s, 1 H), 5.93 (s, 1 H), 5.60 (dd, 1 H), 4.64 (s, 2 H), 3.27 (s, 3 H), 3.48 - 3.36 (obs. m, 1 H), 2.99 (dd, 1 H), 2.58 - 2.46 (obs. m, 1 H), 2.38 - 2.25 (m, 1 H).

Example 143: 7-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-4-methyl-2H-1,4-benzoxazin-3(4H)-one

The compound prepared in Example 142 (100 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (28 mg).

LC/MS  $t_R$  4.09 minutes; MS (ES<sup>+</sup>)  $m/z$  575 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.99 (s, 1 H), 9.71 (s, 1 H), 7.88 - 7.73 (m, 3 H), 7.43 (dd, 1 H), 7.34 (d, 1 H), 7.27 (d, 1 H), 5.99 (s, 1 H), 5.93 (s, 1 H), 5.56 (dd, 1 H), 4.70 (s, 2 H), 3.35 - 3.21 (m, 1 H), 3.30 (s, 3 H), 2.99 (dd, 1 H), 2.58 - 2.48 (m, 1 H), 2.20 (t, 1 H).

Example 144: 6-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1,4-dihydro-2H-3,1-benzoxazin-2-one

The same operation as in Example 139 → Example 91 → Example 51 → Example 39 was conducted from 6-bromo-1,4-dihydro-2H-3,1-benzoxazin-2-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 17 was used).

LC/MS  $t_R$  2.91 minutes; MS (ES<sup>+</sup>)  $m/z$  527 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.09 (s, 1 H), 10.12 (s, 1 H), 9.70 (s, 1 H), 7.82 -

7.77 (m, 3 H), 7.57 (d, 1 H), 7.54 (s, 1 H), 7.41 (d, 1 H), 6.84 (d, 1 H), 5.95 (app. s, 2 H), 5.61 (d, 1 H), 5.31 (s, 2 H), 3.38 - 3.26 (obs. m, 1 H), 3.00 (dd, 1 H), 2.54 - 2.46 (obs. m, 1 H), 2.20 (t, 1 H).

5    Example 145: 6-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyI}-1H-imidazol-5-yl)-1,4-dihydro-2H-3,1-benzoxazin-2-one

The compound prepared in Example 144 (40 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (10.6 mg).

10    LC/MS  $t_R$  3.79 minutes; MS (ES<sup>+</sup>)  $m/z$  561 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.75 (s, 1 H), 10.30 (s, 1 H), 9.71 (s, 1 H), 7.90 - 7.74 (m, 3 H), 7.57 (d, 1 H), 7.52 (app. s, 1 H), 6.97 (d, 1 H), 6.00 (s, 1 H), 5.94 (s, 1 H), 5.54 (dd, 1 H), 5.35 (s, 2 H), 3.38 - 3.28 (obs. m, 1 H), 2.99 (dd, 1 H), 2.59 - 2.48 (obs. m, 1 H), 2.21 (t, 1 H).

15

Example 146: 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyI}-1H-imidazol-5-yl)-1,3-benzoxazol-2(3H)-one

The same operation as in Example 84 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, 6-(2-bromoacetyl)benzoxazol-2(3H)-one [J. Med. Chem., 34(6), 1860 (1991)] was used).

20    LC/MS  $t_R$  2.96 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.10 (s, 1 H), 11.58 (br. s, 1 H), 9.68 (s, 1 H), 7.87 - 7.74 (m, 3 H), 7.57 (s, 1 H), 7.52 (d, 1 H), 7.48 (s, 1 H), 7.02 (d, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.62 (dd, 1 H), 3.45 - 3.36 (obs. m, 1 H), 3.11 - 2.88 (m, 1 H), 2.50 (obs. m, 1 H), 2.40 - 2.09 (m, 1 H).

25    Example 147: 6-(4-chloro-2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyI}-1H-imidazol-5-yl)-1,3-benzoxazol-2(3H)-one

The compound prepared in Example 146 (105 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (4 mg).

LC/MS  $t_R$  3.85 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.74 (s, 1 H), 7.73 (dd, 1 H), 7.68 (d, 1 H), 7.55 (d, 1 H), 7.52 (dd, 1 H), 7.16 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.71 (dd, 1 H), 3.49 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.65 (ddd, 1 H), 2.40 (ddt, 1 H).

5 Example 148: 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-1-isoindolinone

The same operation as in Example 84 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to  
10 Example 84 in the operation, 6-(bromoacetyl)isoindolin-1-one [Bioorg. Med. Chem., 16(6), 3091 (2008)] was used).

LC/MS *t*<sub>R</sub> 2.98 minutes; MS (ES<sup>+</sup>) *m/z* 511 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.93 (br. s, 1 H), 7.83 (br. s, 1 H), 7.75 (br. s, 1 H), 7.75 - 7.70 (m, 2 H), 7.68 (d, 1 H), 7.54 (br. s, 1 H), 6.14 (s, 1 H), 6.10  
15 (s, 1 H), 5.80 (d, 1 H), 4.49 (s, 2 H), 3.56 - 3.39 (m, 1 H), 3.36 - 3.26 (obs. m, 1 H), 3.12 (ddd, 1 H), 2.71 - 2.58 (m, 1 H).

Example 149: 6-(2-bromoacetyl)-3-methyl-1,2,3,4-tetrahydroquinazolin-2-one

To a cooled (0 °C) suspension of aluminium trichloride (2.06 g) in 1,2-  
20 dichloroethane (5 mL) was added bromoacetyl bromide (1.1 mL) and the mixture stirred at 0 °C for 30 minutes. To the reaction mixture, a solution of 3-methyl-3,4-dihydroquinazolin-2(1H)-one (1.0 g) in 1,2-dichloroethane (5 mL) was added and the mixture stirred at 50 °C for 3 hours. The reaction mixture was concentrated and the residue suspended in iced water (20 mL). The resultant precipitate was isolated by  
25 filtration to give the title compound having the following physical properties (1.51 g).  
LC/MS *t*<sub>R</sub> 1.53 minutes; MS (ES<sup>+</sup>) *m/z* 283 and 285 (M+H)<sup>a</sup>

Example 150: 6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-3-methyl-3,4-dihydro-2(1H)-  
30 quinazolinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the

compound prepared in Example 149 was used).

LC/MS  $t_R$  2.94 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.16 (s, 1 H), 8.25 (s, 1 H), 8.23 (s, 1 H), 8.03 (d, 1 H), 7.98 – 7.89 (m, 2 H), 7.47 (d, 1 H), 7.33 (dd, 1 H), 7.14 (d, 1 H), 6.47 (s, 1 H), 6.40 (br. s, 1 H), 6.22 (s, 1 H), 5.95 (d, 1 H), 3.65 – 3.46 (m, 2 H), 3.24 (dd, 1 H), 2.65 – 2.50 (m, 1 H), 2.05 (s, 2 H), 1.57 (s, 3 H).

Example 151: 6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-methyl-3,4-dihydro-2(1H)-quinazolinone

10 The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 25 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 149 was used).

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.08 (br. s, 1 H), 9.70 (s, 1 H), 9.17 (br. s, 1 H), 7.88 - 7.76 (m, 3 H), 7.69 (d, 1 H), 7.53 - 7.44 (m, 2 H), 7.39 (s, 1 H), 6.72 (d, 1 H), 6.29 (s, 1 H), 5.87 (dd, 1 H), 5.07 (s, 2 H), 4.43 (s, 2 H), 2.87 (s, 3 H).

20 Example 152: 6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-methyl-3,4-dihydro-2(1H)-quinazolinone

The compound prepared in Example 151 (70 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (6.0 mg).

LC/MS  $t_R$  3.68 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.82 (br. s, 1 H), 9.71 (s, 1 H), 9.36 (s, 1 H), 7.88 - 7.77 (m, 3 H), 7.69 (d, 1 H), 7.45 (d, 1 H), 7.41 (s, 1 H), 6.84 (d, 1 H), 6.28 (d, 1 H), 5.89 (dd, 1 H), 5.06 (s, 2 H), 4.46 (s, 2 H), 2.88 (s, 3 H).

Example 153: 2-bromo-1-(2-chloroquinoxalin-6-yl)ethan-1-one

30 To an acetonitrile (10 mL) suspension of 2-oxo-1,2-dihydroquinoxaline-6-carboxylic acid [patent WO2006/040568, page 75, 82] (0.38 g) was sequentially added pyridine (8  $\mu$ L), N,N-dimethylformamide (7.5  $\mu$ L) and thionyl chloride (2.2 mL) and the mixture stirred at 90 °C for 1 hour. The reaction mixture was concentrated and the

residue dissolved in acetonitrile (10 mL). To the cooled (0 °C) acetonitrile solution, a 2 M solution of (trimethylsilyl)diazomethane in diethyl ether (2.5 mL) was added and the mixture stirred at 0 °C for 2 hours. To the reaction mixture, a solution of 33 wt. % hydrogen bromide in acetic acid (0.86 mL) was added and the mixture stirred at 0 °C for a further 30 minutes. A saturated aqueous solution of sodium hydrogen carbonate (50 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to obtain the title compound having the following physical properties (0.35 g).

LC/MS  $t_R$  1.96 minutes; MS (ES<sup>+</sup>)  $m/z$  285 and 287 (M+H)<sup>a</sup>.

Example 154: 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2(1H)-quinoxalinone

The same operation as in Example 51 → Example 52 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 153 was used).

LC/MS  $t_R$  2.95 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.17 (br. s, 1 H), 9.69 (s, 1 H), 8.15 (s, 1 H), 8.06 (d, 1 H), 7.92 (dd, 1 H), 7.82 - 7.76 (m, 3 H), 7.61 (s, 1 H), 7.28 (d, 1 H), 5.98 (s, 1 H), 5.95 (s, 1 H), 5.63 (dd, 1 H), 3.45 - 3.37 (m, 2 H), 3.02 (dd, 1 H), 2.41 - 2.31 (m, 1 H).

Example 155: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(1-methyl-1H-benzimidazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 39 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, 2-bromo-1-(1-methyl-1H-benzimidazol-5-yl)ethanone hydrobromide was used).

LC/MS  $t_R$  2.75 minutes; MS (ES<sup>+</sup>)  $m/z$  510 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 8.13 (s, 1 H), 7.96 (s, 1 H), 7.76 - 7.74 (m, 2 H), 7.71 - 7.69 (m, 2 H), 7.58 - 7.56 (m, 1 H), 7.32 (s, 1 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.84 - 5.82 (m, 1 H), 3.92 (s, 3 H), 3.53 - 3.46 (m, 1 H), 3.16 - 3.11 (m, 1 H), 2.69 - 2.63 (m, 1 H), 2.55 - 2.50 (m, 1 H).

Example 156: 3-[4-chloro-5-(1-methyl-1H-benzimidazol-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 84 → Example 39 → Example 47 → Example 40 → Example 41 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, 2-bromo-1-(1-methyl-1H-benzimidazol-5-yl)ethanone hydrobromide was used).

LC/MS  $t_R$  3.16 minutes; MS ( $ES^+$ )  $m/z$  544 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.90 (br. s, 1 H), 9.71 (s, 1 H), 8.24 (s, 1 H), 7.97 (s, 1 H), 7.84 - 7.78 (m, 3 H), 7.68 (d, 1 H), 7.63 (d, 1 H), 6.00 (s, 1 H), 5.94 (s, 1 H), 5.59 (dd, 1 H), 3.86 (s, 3 H), 3.32 - 3.25 (m, 1 H), 3.00 (dd, 1 H), 2.60 - 2.52 (m, 1 H), 2.29 - 2.16 (m, 1 H).

Example 157: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1H-indazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 39 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, 2-bromo-1-(1H-indazol-5-yl)ethan-1-one [Bioorg. Med. Chem. Lett. 21(5), 1480 (2011)] was used).

LC/MS  $t_R$  2.92 minutes; MS ( $ES^+$ )  $m/z$  496 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.96 (s, 1 H), 12.04 (s, 1 H), 9.69 (s, 1 H), 8.07 (d, 1 H), 8.01 (d, 1 H), 7.84 - 7.77 (m, 3 H), 7.73 (d, 1 H), 7.50 - 7.43 (m, 2 H), 5.99 (s, 1 H), 5.96 (s, 1 H), 5.64 (d, 1 H), 3.38 - 3.27 (obs. m, 1 H), 3.02 (dd, 1 H), 2.56 - 2.44 (obs. m, 1 H), 2.40 - 2.32 (m, 1 H).

Example 158: ethyl 4-{2-[7-(2-{[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-4-yl}benzoate

The same operation as in Example 84 → Example 39 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, ethyl 4-acetylbenzoate [Bioorg. Med. Chem. Lett. 18, 2886 (2008)] was used).

LC/MS  $t_R$  2.16 minutes; MS ( $ES^+$ )  $m/z$  575 (M+H)<sup>a</sup>.

Example 159: tert-butyl N-[4-chloro-2-(3-{4-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl)phenyl]carbamate

To a cooled (0 °C) tetrahydrofuran (20 mL) solution of the compound prepared in Example 158 (0.50 g) was added a 1 M solution of lithium aluminium hydride in tetrahydrofuran (1.39 mL) and the mixture stirred at 0 °C for 1 hour. To the cooled (0 °C) reaction mixture, a saturated aqueous solution of ammonium chloride (10 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (0.30 g).  
LC/MS  $t_R$  1.71 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>a</sup>.

Example 160(1) and Example 160(2): 7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone and 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzyl acetate

The same operation as in Example 40 → Example 41 was conducted from Example 159 to give the title compounds in a 2: 1 ratio having the following physical properties.

20

Example 160(1):

LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  486 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (250 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.78 - 7.58 (m, 5 H), 7.42 - 7.25 (m, 3 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.59 (s, 2 H), 3.43 (dd, 1 H), 3.13 (dd, 1 H), 2.74 - 2.36 (m, 2 H).

25

Example 160(2):

LC/MS  $t_R$  3.31 minutes; MS (ES<sup>+</sup>)  $m/z$  528 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.64 (m, 5 H), 7.38 (d, 2 H), 7.37 (br. s, 1 H), 6.15 (s, 1 H), 6.11 (s, 1 H), 5.80 (dd, 1 H), 5.12 (s, 2 H), 3.53 - 3.41 (m, 1 H), 3.22 - 3.07 (m, 1 H), 2.73 - 2.61 (m, 1 H), 2.55 - 2.40 (m, 1 H), 2.10 (s, 3 H).

30

Example 161: tert-butyl N-(4-chloro-2-{3-[4-(4-formylphenyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

To a dichloromethane (5 mL) solution of the compound prepared in Example 159 (0.39 g) was added Dess-Martin periodinane (1,1,1-Triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one) (0.33 g) and the mixture stirred at room temperature for 16 hours. To the reaction mixture, a 1:1 mixture of a saturated aqueous solution of sodium hydrogen carbonate and a saturated aqueous solution of sodium sulfite was added (10 mL) and the mixture stirred at room temperature for 15 minutes followed by extraction into dichloromethane. The combined organic layers were washed with a saturated aqueous solution of sodium hydrogen carbonate, brine, dried and concentrated to obtain the title compound having the following physical properties (0.30 g).  
LC/MS  $t_R$  2.04 minutes; MS ( $ES^+$ )  $m/z$  531 (M+H) <sup>a</sup>.

Example 162: tert-butyl N-{4-chloro-2-[3-(4-{4-[(hydroxyimino)methyl]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl]phenyl}carbamate

To an ethanol (2 mL) solution of the compound prepared in Example 161 (0.15 g) was added hydroxylamine hydrochloride (15.1 mg) and 2 M sodium hydroxide (124  $\mu$ L) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated and the residue suspended in water (5 mL) and extracted into dichloromethane. The combined organic layers were washed with brine, dried and concentrated to obtain the title compound having the following physical properties (0.15 g).  
LC/MS  $t_R$  1.84 minutes; MS ( $ES^+$ )  $m/z$  546 (M+H) <sup>a</sup>.

Example 163: 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)benzaldehyde oxime

The same operation as in Example 40  $\rightarrow$  Example 41 was conducted from the compound prepared in Example 162 to give the title compound having the following physical properties.  
LC/MS  $t_R$  3.10 minutes; MS ( $ES^+$ )  $m/z$  499 (M+H) <sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 8.09 (s, 1 H), 7.78 - 7.67 (m, 5 H), 7.60 (d, 2 H), 7.40 (br. s, 1 H), 6.15 (s, 1 H), 6.11 (s, 1 H), 5.80 (dd, 1 H), 3.47 (td, 1 H), 3.13 (ddd, 1 H), 2.71 - 2.62 (m, 1 H), 2.49 (br. s, 1 H).



Example 164: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)benzaldehyde O-methyloxime

The same operation as in Example 162 → Example 40 → Example 41 was  
5 conducted from the compound prepared in Example 161 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 162 in the operation, methoxyamine hydrochloride was used).

LC/MS  $t_R$  3.61 minutes; MS ( $ES^+$ )  $m/z$  513 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 8.09 (s, 1 H), 7.78 - 7.68 (m, 5 H),  
10 7.61 (d, 2 H), 7.42 (s, 1 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.81 (dd, 1 H), 3.95 (s, 3 H),  
3.52 - 3.43 (m, 1 H), 3.20 - 3.07 (m, 1 H), 2.71 - 2.62 (m, 1 H), 2.53 - 2.46 (m, 1 H).

Example 165: (3S)-3-[5-(4-acetylphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

15 The same operation as in Example 84 → Example 39 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, 1-(4-acetylphenyl)-2-bromoethan-1-one [Chem. Pharm. Bull. 40 (5), 1170 (1992)] was used).

20 LC/MS  $t_R$  3.48 minutes; MS ( $ES^+$ )  $m/z$  498 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 8.01 (d, 2 H), 7.85 (app. br. s, 2 H),  
7.77 - 7.68 (m, 3 H), 7.55 (br. s, 1 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.81 (dd, 1 H), 3.53 -  
3.43 (m, 1 H), 3.21 - 3.08 (m, 1 H), 2.72 - 2.65 (m, 1 H), 2.62 (s, 3 H), 2.57 - 2.46 (m, 1  
H).

25

Example 166: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(1-hydroxyethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

To a cooled (0 °C) tetrahydrofuran (2 mL) solution of the compound prepared in  
Example 165 (30 mg) was added sodium borohydride (1.1 mg) and the mixture stirred  
30 at room temperature for 25 minutes. To the cooled (0 °C) reaction mixture, water (10  
mL) was added followed by extraction with ethyl acetate. The combined organic layers  
were washed with brine, dried and concentrated. The residue was purified by column  
chromatography (0 to 15% methanol in dichloromethane) to obtain the title compound

having the following physical properties (5.5 mg)

LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.26 (s, 1 H), 7.68 - 7.60 (m, 2 H), 7.58 (d, 1 H), 7.50 (app. br. s, 2 H), 7.28 (d, 2 H), 7.17 (br. s, 1 H), 6.04 (s, 1 H), 6.00 (s, 1 H), 5.68 (dd, 1 H), 4.70 (q, 1 H), 3.41 - 3.30 (m, 1 H), 3.07 - 2.96 (m, 1 H), 2.59 - 2.45 (m, 1 H), 2.41 - 2.25 (m, 1 H), 1.34 (d, 3 H).

Example 167: 2-methyl-2-propanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]carbamate

10 The same operation as in Example 51 → Example 39 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, tert-butyl N-[[4-(2-bromoacetyl)phenyl]methyl] carbamate [Bioorg. Med. Chem. Lett. 13(20), 3557 (2003)] was used).

15 LC/MS  $t_R$  3.51 minutes; MS (ES<sup>+</sup>)  $m/z$  585 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.58 (s, 1 H), 7.62 (dd, 1 H), 7.54 (d, 1 H), 7.52 (d, 1 H), 7.28 (s, 1 H), 7.27 (app. s, 4 H), 7.22 (s, 1 H), 6.31 (s, 1 H), 5.86 (d, 1 H), 5.75 (s, 1 H), 4.88 (br. s, 1 H), 4.31 (br. d, 2 H), 3.53 - 3.42 (m, 1 H), 3.42 - 3.31 (m, 1 H), 3.03 (dd, 1 H), 2.49 (quintet, 1 H), 1.47 (s, 9 H).

20

Example 168: (3S)-3-{5-[4-(aminomethyl)phenyl]-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone dihydrochloride

The compound prepared in Example 167 (0.19 g) was treated in accordance with Example 55 to give the title compound having the following physical properties (0.13 g).

25 LC/MS  $t_R$  2.46 minutes; MS (ES<sup>+</sup>)  $m/z$  485 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.71 (s, 1 H), 8.39 (br. s, 3 H), 8.10 (br. s, 1 H), 7.88 (d, 2 H), 7.84 (dd, 1 H), 7.82 (d, 1 H), 7.72 (d, 1 H), 7.62 (d, 2 H), 6.10 (s, 1 H), 6.00 (s, 1 H), 5.84 (dd, 1 H), 4.14 - 4.02 (q, 2 H), 3.34 - 3.24 (m, 1 H), 3.15 - 3.05 (m, 1 H), 2.75 - 2.67 (m, 1 H), 2.41 - 2.30 (m, 1 H).

30

Example 169: methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]carbamate

The compound prepared in Example 168 (100 mg) was treated in accordance

with Example 77 to give the title compound having the following physical properties (30 mg).

LC/MS  $t_R$  3.07 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.80 - 7.59 (m, 5 H), 7.31 - 7.26 (m, 3 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.80 (dd, 1 H), 4.29 (s, 2 H), 3.68 (s, 3 H), 3.51 - 3.41 (m, 1 H), 3.18 - 3.07 (m, 1 H), 2.65 - 2.60 (m, 1 H), 2.52 - 2.41 (m, 1 H).

Example 170: N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]acetamide

10 To a dichloromethane (3 mL) solution of the compound prepared in Example 168 (100 mg) was sequentially added pyridine (29  $\mu$ L) and acetic anhydride (13.6  $\mu$ L) and the mixture stirred at room temperature for 4 hours. To the cooled (0 °C) reaction mixture, 1 M hydrochloric acid (6.60 mL) was added followed by extraction into dichloromethane. The combined organic layers were washed with brine, dried and  
15 concentrated. The residue was purified by column chromatography (0 to 15% methanol in dichloromethane) to obtain the title compound having the following physical properties (27.4 mg).

LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  527 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.39 (s, 1 H), 7.77 - 7.68 (m, 3 H), 7.63 (d, 2 H),  
20 7.31 - 7.27 (m, 3 H), 6.15 (s, 1 H), 6.14 (s, 1 H), 5.81 (dd, 1 H), 4.37 (s, 2 H), 3.54 - 3.37 (m, 1 H), 3.22 - 3.07 (m, 1 H), 2.74 - 2.59 (m, 1 H), 2.55 - 2.38 (m, 1 H), 2.02 (s, 3 H).

Example 171: N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzyl]-2,2,2-trifluoroacetamide

25 To a cooled (0 °C) dichloromethane (3 mL) solution of the compound prepared in Example 168 (100 mg) was sequentially added triethylamine (57  $\mu$ L) and trifluoroacetic anhydride (59  $\mu$ L) and the mixture stirred at 0 °C for 3 hours. To the reaction mixture water (5 mL) was added followed by extraction with dichloromethane.  
30 The combined organic layers were washed with brine, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate), then triturated with a 1:1 mixture of dichloromethane and heptanes. The precipitate was collected by filtration to

give the title compound having the following physical properties (2 mg).

LC/MS  $t_R$  3.35 minutes; MS (ES<sup>+</sup>)  $m/z$  581 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.78 - 7.73 (m, 2 H), 7.69 (d, 1 H), 7.67 (app. br. s, 2 H), 7.36 (br. s, 1 H), 7.33 (d, 2 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.80 (dd, 1 H), 4.47 (s, 2 H), 3.55 - 3.39 (m, 1 H), 3.14 (dd, 1 H), 2.66 (dd, 1 H), 2.52 - 2.36 (m, 1 H).

Example 172: 1-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]benzyl]-3-ethylurea

10 To a dichloromethane (3 mL) solution of the compound prepared in Example 168 (100 mg) was added ethyl isocyanate (29  $\mu$ L) and the mixture stirred at room temperature for 3 hours. To the reaction mixture water (5 mL) was added followed by extraction with dichloromethane. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by high performance liquid  
15 chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (8.9 mg).  
LC/MS  $t_R$  2.99 minutes; MS (ES<sup>+</sup>)  $m/z$  556 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.27 (s, 1 H), 7.68 - 7.62 (m, 2 H), 7.58 (d, 1 H), 7.57 - 7.44 (m, 2 H), 7.28 - 7.11 (m, 3 H), 6.04 (s, 1 H), 6.01 (s, 1 H), 5.69 (d, 1 H),  
20 4.22 (s, 2 H), 3.40 - 3.29 (m, 1 H), 3.07 (q, 2 H), 3.05 - 2.97 (m, 1 H), 2.61 - 2.42 (m, 1 H), 2.34 - 2.19 (m, 1 H), 1.02 (t, 3 H).

Example 173(1) to Example 173(23)

The compounds of the present invention having the following physical data were  
25 prepared using the corresponding alpha-bromoketones from the compound prepared in Example 9 in the process of Example 51  $\rightarrow$  Example 52.

Example 173(1): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

30 LC/MS  $t_R$  2.81 minutes; MS (ES<sup>+</sup>)  $m/z$  457 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.39 (br. s, 1 H), 9.70 (s, 1 H), 8.47 (d, 2 H), 7.94 - 7.73 (m, 4 H), 7.64 (d, 2 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.63 (dd, 1 H), 3.44 - 3.37 (m, 1 H), 3.01 (dd, 1 H), 2.55 - 2.52 (m, 1 H), 2.38 - 2.26 (m, 1 H).

Example 173(2): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  2.85 minutes; MS (ES<sup>+</sup>)  $m/z$  457 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.94 (br. s, 1 H), 9.70 (s, 1 H), 8.93 (s, 1 H), 8.37 (d, 1 H), 8.03 (d, 1 H), 7.84 - 7.77 (m, 3 H), 7.67 (br. s, 1 H), 7.34 (s, 1 H), 5.99 (s, 1 H), 5.97 (s, 1 H), 5.62 (d, 1 H), 3.45 - 3.36 (obs. m, 1 H), 3.00 (dd, 1 H), 2.58 - 2.45 (obs. m, 1 H), 2.38 - 2.25 (m, 1 H).

10 Example 173(3): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  2.91 minutes; MS (ES<sup>+</sup>)  $m/z$  457 (M+H)<sup>b</sup>

NMR analysis showed a 2 : 1 ratio of tautomers.

- Major tautomer: <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.28 (br. s, 1 H), 9.70 (s, 1 H), 8.46  
15 (d, 1 H), 7.84 - 7.70 (m, 5 H), 7.59 (s, 1 H), 7.16 (td, 1 H), 5.97 (app. s, 2 H), 5.65 (d, 1 H), 3.41 - 3.34 (m, 1 H), 3.01 (dd, 1 H), 2.57 - 2.44 (obs. m, 1 H), 2.38 - 2.31 (m, 1 H).  
Minor tautomer: <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.60 (br. s, 1 H), 9.71 (s, 1 H), 8.55 (d, 1 H), 7.84 - 7.70 (m, 5 H), 7.46 (s, 1 H), 7.23 (dd, 1 H), 5.97 (s, 2 H), 5.95 (s, 1 H), 5.73 (d, 1 H), 3.31 - 3.21 (m, 1 H), 2.98 (dd, 1 H), 2.57 - 2.44 (obs. m, 1 H), 2.28 - 2.21  
20 (m, 1 H).

Example 173(4): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1,3-thiazol-2-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.62 minutes; MS (ES<sup>+</sup>)  $m/z$  463 (M+H)<sup>b</sup>

- 25 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.47 (br. s, 1 H), 9.69 (s, 1 H), 7.84 - 7.77 (m, 3 H), 7.69 (s, 1 H), 7.65 (s, 1 H), 5.99 (s, 1 H), 5.97 (s, 1 H), 5.62 (dd, 1 H), 3.32 - 3.24 (obs. m, 1 H), 3.01 (dd, 1 H), 2.59 - 2.52 (obs. m, 1 H), 2.37 - 2.25 (m, 1 H).

30 Example 173(5): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-pyrazinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, 2-bromo-1-(pyrazin-2-yl)ethan-1-one [J. Med. Chem. 41(13), 2436 (1998)] was used.

LC/MS  $t_R$  3.29 minutes; MS (ES<sup>+</sup>)  $m/z$  458 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.47 (br. s, 1 H), 9.69 (s, 1 H), 8.98 (app. br. s, 1 H), 8.52 (app. br. s, 1 H), 8.42 (app. br. s, 1 H), 7.82 - 7.78 (m, 3 H), 7.73 (br. s, 1 H), 5.97 (s, 1 H), 5.96 (s, 1 H), 5.66 (d, 1 H), 3.45 - 3.34 (obs. m, 1 H), 3.01 (dd, 1 H), 2.59 - 2.45 (obs. m, 1 H), 2.38 - 2.32 (m, 1 H).

5

Example 173(6): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-methoxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS *t<sub>R</sub>* 3.23 minutes; MS (ES<sup>+</sup>) *m/z* 486 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.71 (br. s, 1 H), 8.53 (s, 1 H), 7.60 (dd, 1 H), 7.59 (app. br. s, 1 H), 7.54 (d, 1 H), 7.50 (d, 1 H), 7.30 (br. s, 1 H), 7.26 - 7.18 (m, 1 H), 7.05 - 6.90 (m, 2 H), 6.31 (s, 1 H), 5.88 (d, 1 H), 5.68 (s, 1 H), 4.02 (s, 3 H), 3.54 - 3.26 (m, 2 H), 3.00 (dd, 1 H), 2.45 (quintet, 1 H).

Example 173(7): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-methoxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS *t<sub>R</sub>* 3.31 minutes; MS (ES<sup>+</sup>) *m/z* 486 (M+H)<sup>b</sup>

NMR analysis showed a 2 : 1 ratio of tautomers.

Major tautomer: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.77 (br. s, 1 H), 8.54 (s, 1 H), 7.61 (d, 1 H), 7.54 (d, 1 H), 7.51 (br. s, 1 H), 7.37 - 6.96 (m, 4 H), 6.79 (d, 1 H), 6.32 (s, 1 H), 5.83 (d, 1 H), 5.70 (s, 1 H), 3.86 (s, 3 H), 3.53 - 3.25 (m, 2 H), 3.06 - 2.95 (m, 1 H), 2.53 - 2.40 (m, 1 H).

Minor tautomer: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.16 (br. s, 1 H), 8.54 (s, 1 H), 7.61 (d, 1 H), 7.54 (d, 1 H), 7.51 (br. s, 1 H), 7.37 - 6.96 (m, 4 H), 6.82 (d, 1 H), 6.36 (s, 1 H), 5.87 (d, 1 H), 5.70 (s, 1 H), 3.84 (s, 3 H), 3.53 - 3.25 (m, 2 H), 3.06 - 2.95 (m, 1 H), 2.53 - 2.40 (m, 1 H).

Example 173(8): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-methoxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS *t<sub>R</sub>* 3.17 minutes; MS (ES<sup>+</sup>) *m/z* 486 (M+H)<sup>b</sup>

NMR analysis showed a 3 : 2 ratio of tautomers.

Major tautomer: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.68 (br. s, 1 H), 8.53 (s, 1 H), 7.68 (d, 2 H), 7.62 (d, 1 H), 7.54 (d, 1 H), 7.51 (s, 1 H), 7.17 (s, 1 H), 6.96 - 6.86 (m, 2 H), 6.32

(s, 1 H), 5.90 - 5.80 (m, 1 H), 5.69 (s, 1 H), 3.83 (s, 3 H), 3.54 - 3.27 (m, 2 H), 3.07 - 2.96 (m, 1 H), 2.46 (quintet, 1 H).

Minor tautomer: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.02 (br. s, 1 H), 8.53 (s, 1 H), 7.62 (d, 1 H), 7.54 (d, 1 H), 7.51 (s, 1 H), 7.42 (d, 1 H), 7.11 (s, 1 H), 6.96 - 6.86 (m, 2 H), 6.36 (s, 1 H), 5.90 - 5.80 (m, 1 H), 5.69 (s, 1 H), 3.83 (s, 3 H), 3.54 - 3.27 (m, 2 H), 3.07 - 2.96 (m, 1 H), 2.46 (quintet, 1 H).

Example 173(9): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-fluorophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

10 LC/MS *t*<sub>R</sub> 3.59 minutes; MS (ES<sup>+</sup>) *m/z* 474 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.21 (br. s, 1 H), 9.68 (s, 1 H), 7.84 - 7.74 (m, 3 H), 7.60 (s, 1 H), 7.54 (d, 1 H), 7.48 (dd, 1 H), 7.35 (dd, 1 H), 6.96 (td, 1 H), 5.97 (s, 1 H), 5.96 (s, 1 H), 5.62 (dd, 1 H), 3.44 - 3.36 (m, 1 H), 3.00 (dd, 1 H), 2.60 - 2.45 (obs. m, 1 H), 2.38 - 2.30 (m, 1 H).

15

Example 173(10): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-fluorophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS *t*<sub>R</sub> 3.38 minutes; MS (ES<sup>+</sup>) *m/z* 474 (M+H)<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.11 (br. s, 1 H), 9.69 (s, 1 H), 7.82 - 7.76 (m, 3 H), 7.72 (dd, 2 H), 7.49 (s, 1 H), 7.14 (t, 2 H), 5.97 (s, 1 H), 5.96 (s, 1 H), 5.63 (dd, 1 H), 3.47 - 3.35 (obs. m, 1 H), 3.01 (dd, 1 H), 2.57 - 2.44 (obs. m, 1 H), 2.40 - 2.28 (m, 1 H).

Example 173(11): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2,4-dimethyl-1,3-oxazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

25 LC/MS *t*<sub>R</sub> 3.36 minutes; MS (ES<sup>+</sup>) *m/z* 475 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.97 (br. s, 1 H), 8.57 (s, 1 H), 7.60 (dd, 1 H), 7.53 (s, 1 H), 7.49 (d, 1 H), 7.09 (d, 1 H), 6.29 (s, 1 H), 5.81 (d, 1 H), 5.72 (s, 1 H), 3.50 - 3.37 (m, 1 H), 3.34 (dd, 1 H), 3.00 (dd, 1 H), 2.51 - 2.44 (m, 1 H), 2.43 (s, 3 H), 2.33 (s, 3 H).

30 Example 173(12): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-pyrimidinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, 2-bromo-1-(pyrimidin-4-yl)ethan-1-one [J. Med. Chem., 51(3), 487 (2008)] was used.

LC/MS  $t_R$  3.33 minutes; MS (ES<sup>+</sup>)  $m/z$  458 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, acetone- $d_6$ )  $\delta$  11.59 (br. s, 1 H), 9.30 (s, 1 H), 8.98 (br. s, 1 H), 8.68 (d, 1 H), 7.88 (s, 1 H), 7.91 (s, 1 H), 7.78 (s, 2 H), 7.72 (s, 1 H), 6.12 (br. s, 1 H), 5.94 (s, 1 H), 5.83 (d, 1 H), 3.51 - 3.35 (m, 1 H), 3.12 (m, 1 H), 3.00 (m, 1 H), 2.62 - 2.53 (m, 1 H).

Example 173(13): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-fluorophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.58 minutes; MS (ES<sup>+</sup>)  $m/z$  475 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, acetone- $d_6$ )  $\delta$  11.35 (br. s, 1 H), 9.30 (s, 1 H), 8.17 (m, 1 H), 7.77 (s, 2 H), 7.73 (s, 1 H), 7.52 (m, 1 H), 7.25 (m, 2 H), 7.15 (m, 1 H), 6.12 (s, 1 H), 5.95 (s, 1 H), 5.85 (m, 1 H), 3.47 (m, 1 H), 3.15 - 3.10 (m, 1 H), 3.10 - 3.05 (m, 1 H), 2.6 - 2.5 (m, 1 H).

Example 173 (14): (3S)-3-[5-(4-chlorophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.71 minutes; MS (ES<sup>+</sup>)  $m/z$  492 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, acetone- $d_6$ )  $\delta$  11.25 (br. s, 1 H), 9.30 (s, 1 H), 7.85 (s, 1 H), 7.80 (d, 2 H), 7.75 (s, 2 H), 7.70 (s, 1 H), 7.55 (s, 1 H), 7.35 (d, 2 H), 6.10 (s, 1 H), 5.95 (s, 1 H), 5.78 (m, 1 H), 3.50 - 3.45 (m, 1 H), 3.15 - 3.10 (m, 1 H), 3.10 - 3.05 (m, 1 H), 2.6 - 2.5 (m, 1 H).

Example 173(15): (3S)-3-[5-(3-chlorophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.81 minutes; MS (ES<sup>+</sup>)  $m/z$  492 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, acetone- $d_6$ )  $\delta$  11.32 (br. s, 1 H), 9.30 (s, 1 H), 7.85 (s, 1 H), 7.78 - 7.70 (m, 3 H), 7.73 (s, 1 H), 7.62 (s, 1 H), 7.35 (m, 1 H), 7.18 (m, 1 H), 6.10 (s, 1 H), 5.95 (s, 1 H), 5.78 (m, 1 H), 3.50 - 3.45 (m, 1 H), 3.15 - 3.10 (m, 1 H), 3.10 - 3.05 (m, 1 H), 2.6 - 2.5 (m, 1 H).

Example 173(16): (3S)-3-[5-(2-chlorophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.64 minutes; MS (ES<sup>+</sup>)  $m/z$  492 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 11.30 (br. s, 1 H), 9.25 (s, 1 H), 8.20 (d, 1 H), 7.80 (s, 1 H), 7.75 (s, 2 H), 7.70 (s, 1 H), 7.38 (d, 1 H), 7.35 (t, 1 H), 7.18 (t, 1 H), 6.10 (s, 1 H), 5.92 (s, 1 H), 5.88 (d, 1 H), 3.50 – 3.45 (m, 1 H), 3.15 – 3.10 (m, 1 H), 3.10 – 3.05 (m, 1 H), 2.6 – 2.5 (m, 1 H).

5

Example 173(17): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(methylthio)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

LC/MS *t<sub>R</sub>* 3.33 minutes; MS (ES<sup>+</sup>) *m/z* 502 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.13 (br. s, 1 H), 9.69 (s, 1 H), 7.84 – 7.75 (m, 3 H),  
10 7.63 (d, 2 H), 7.46 (br. s, 1 H), 7.23 (d, 2 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H),  
3.43 – 3.27 (obs. m, 1 H), 3.00 (dd, 1 H), 2.56 – 2.48 (obs. m, 1 H), 2.47 (s, 3 H), 2.38 –  
2.22 (m, 1 H).

Example 173(18): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-methylphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

15 LC/MS *t<sub>R</sub>* 3.36 minutes; MS (ES<sup>+</sup>) *m/z* 471 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.07 (br. s, 0.45 H), 10.70 (br. s, 0.55 H), 8.53 (br. s, 1 H), 7.59 – 7.66 (m, 2 H), 7.53 – 7.56 (m, 1 H), 7.51 (m, 1 H), 7.38 (d, 1 H), 7.15 – 7.23 (m, 3 H), 6.37 (s, 0.45 H), 6.32 (s, 0.55 H), 5.87 (d, 0.45 H), 5.84 (d, 0.55 H), 5.69 (br. s,  
20 1 H), 3.29 – 3.52 (m, 2 H), 3.01 (dd, 1 H), 2.51 – 2.41 (m, 1 H), 2.36 (br. s, 1.35 H), 2.35 (s, 1.65 H).

Example 173(19): 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzonitrile

25 LC/MS *t<sub>R</sub>* 3.71 minutes; MS (ES<sup>+</sup>) *m/z* 481 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>) δ 11.40 (br. s, 1 H), 9.30 (s, 1 H), 8.00 (d, 2 H), 7.75 (m, 2 H), 7.70 (m, 2 H), 6.10 (s, 1 H), 5.95 (s, 1 H), 5.80 (d, 1 H), 3.50 – 3.45 (m, 1 H), 3.15 – 3.10 (m, 1 H), 3.05 – 2.55 (m, 1 H), 2.6 – 2.5 (m, 1 H).

30 Example 173(20): 3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzonitrile

LC/MS *t<sub>R</sub>* 3.61 minutes; MS (ES<sup>+</sup>) *m/z* 481 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.06 (br. s, 1 H), 8.01 (d, 1 H), 7.75 - 7.72 (m, 2 H), 7.70 - 7.67 (m, 1 H), 7.59 - 7.49 (m, 3 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.52 - 3.43 (m, 1 H), 3.12 (ddd, 1 H), 2.69 - 2.60 (m, 1 H), 2.52 (br. s, 1 H).

5

Example 173(21): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[3-(trifluoromethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

LC/MS *t<sub>R</sub>* 4.12 minutes; MS (ES<sup>+</sup>) *m/z* 525 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.86 (br. s, 1 H), 8.53 (s, 1 H), 8.03 (s, 1 H), 7.86 - 7.96 (m, 1 H), 7.44 - 7.74 (m, 5 H), 7.33 (d, 1 H), 6.33 (s, 1 H), 5.85 (d, 1 H), 5.72 (s, 1 H), 3.35 - 3.56 (m, 2 H), 2.93 - 3.08 (m, 1 H), 2.40 - 2.57 (m, 1 H).

10

Example 173(22): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-hydroxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

15 LC/MS *t<sub>R</sub>* 2.96 minutes; MS (ES<sup>+</sup>) *m/z* 472 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.24 (s, 1 H), 7.20 - 7.15 (m, 1 H), 7.14 - 7.06 (m, 2 H), 6.68 (td, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.79 (dd, 1 H), 3.50 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.51 - 2.41 (m, 1 H).

20

Example 173(23): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-nitrophenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, 2-bromo-1-(3-nitrophenyl)ethan-1-one [patent W2010/032010] was used.

25 LC/MS *t<sub>R</sub>* 4.03 minutes; MS (ES<sup>+</sup>) *m/z* 501 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.95 (br. s, 1 H), 8.60 (s, 1 H), 8.55 (s, 1 H), 8.05 (d, 2 H), 7.65 (m, 1 H), 7.60 - 7.50 (m, 3 H), 7.48 (s, 1 H), 6.35 (s, 1 H), 5.85 (d, 1 H), 5.72 (s, 1 H), 3.52 - 3.42 (m, 2 H), 3.10-3.02 (dd, 1 H), 2.55 - 2.45 (m, 1 H).

30 Example 174: 2-bromo-1-(1-methyl-1H-pyrazol-4-yl)ethan-1-one

A suspension of 1-methyl-1H-pyrazole-4-carboxylic acid (0.25 g) in thionyl chloride (4 mL) was stirred at reflux for 30 minutes. The reaction was concentrated, the residue azeotroped with toluene and dissolved in acetonitrile (5 mL). To the cooled

(0 °C) solution, a 2 M solution of (trimethylsilyl)diazomethane in hexanes (1.5 mL) was added and the mixture stirred at 0 °C for 1 hour. A solution of 33 wt. % hydrogen bromide in acetic acid (1.0 mL) was added and the mixture stirred at room temperature for a further 80 minutes. To the reaction mixture, tert-butyl methyl ether (20 mL) was added and the resultant precipitate isolated by filtration to give the title product having the following physical properties (0.22 g).  
LC/MS  $t_R$  1.23 minutes; MS ( $ES^+$ )  $m/z$  203 and 205 ( $M+H$ )<sup>a</sup>.

10 Example 175: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1-methyl-1H-pyrazol-4-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 174 was used).  
15 LC/MS  $t_R$  2.79 minutes; MS ( $ES^+$ )  $m/z$  460 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.94 (br. s, 1 H), 9.74 (s, 1 H), 7.89 - 7.82 (m, 4 H), 7.62 (s, 1 H), 7.17 (s, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.58 (dd, 1 H), 3.87 (s, 3 H), 3.40 - 3.27 (obs. m, 1 H), 3.04 (dd, 1 H), 2.64 - 2.46 (obs. m, 1 H), 2.45 - 2.34 (m, 1 H).

20 Example 176: methyl N-[3-bromo-5-(2-bromoacetyl)thiophen-2-yl]carbamate

To a dichloromethane (15 mL) suspension of 5-acetylthiophene-2-carboxylic acid (1.0 g) was added sequentially oxalyl chloride (0.32 mL) and N,N-dimethylformamide (15  $\mu$ L) and the mixture stirred at room temperature for 2 hours. The reaction mixture was concentrated and the residue dissolved in acetone (6 mL). To the mixture a solution of sodium azide (0.50 g) in water (2 mL) was added and the reaction stirred at room temperature for 30 minutes. The resultant precipitate was isolated by filtration, dissolved in toluene (80 mL) and the mixture refluxed for 1 hour. Methanol (10 mL) was added at room temperature and the mixture stirred at room temperature for a further hour, then concentrated to approximately 30 mL. The resultant precipitate was collected by filtration to give the title compound having the following physical properties (0.56 g).  
30 LC/MS  $t_R$  1.38 minutes; MS ( $ES^+$ )  $m/z$  200 ( $M+H$ )<sup>a</sup>.

Example 177: methyl N-[3-bromo-5-(2-bromoacetyl)thiophen-2-yl]carbamate

The compound prepared in Example 176 (0.46 g) was treated as detailed in Example 91 to give the title compound having the following physical properties (0.68 g). LC/MS  $t_R$  1.84 minutes; MS ( $ES^+$ )  $m/z$  356, 358, 360 ( $M+H$ )<sup>a</sup>.

5

Example 178: methyl N-[5-(2-azidoacetyl)-3-bromothiophen-2-yl]carbamate

To an N,N-dimethylformamide (15 mL) solution of the compound prepared in Example 177 (0.68 g) was added sodium azide (0.25 g) and the mixture stirred at room temperature for 1 hour. To the reaction mixture, water (100 mL) was added followed by  
10 extraction with ethyl acetate. The combined organic layers were dried and concentrated to give the title compound having the following physical properties (0.51 g). LC/MS  $t_R$  1.81 minutes; MS ( $ES^+$ )  $m/z$  319 and 321 ( $M+H$ )<sup>a</sup>.

Example 179: methyl N-[5-(2-aminoacetyl)thiophen-2-yl]carbamate hydrochloride

To an ethanol (25 mL) solution of the compound prepared in Example 178 (0.51 g) was sequentially added 2 M hydrochloric acid (1.60 mL) and 5% palladium-carbon (0.10 g) and the mixture was stirred under an atmosphere of hydrogen for 16 hours. The reaction mixture was filtered through Celite® and the filtrate concentrated. The residue was dissolved in water (15 mL) and washed with ethyl acetate. The aqueous layer was  
20 concentrated to give the title compound as the hydrochloride salt having the following physical properties (0.28 g).  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.52 (br. s, 1 H), 7.88 (d, 1 H), 6.69 (d, 1 H), 4.39 (s, 2 H), 3.77 (s, 3 H).

25 Example 180: methyl N-{5-[2-({7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl}formamido)acetyl]thiophen-2-yl}carbamate

To an N,N-dimethylformamide (1 mL) solution of the compound prepared in Example 179 (0.15 g) and the compound prepared in Example 17 (0.20 g) was sequentially added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium  
30 hexafluorophosphate (0.23 g) and diisopropylethylamine (0.29 mL) and the mixture stirred at room temperature for 3 hours. To the reaction mixture, water (15 mL) was added followed by extraction into ethyl acetate. The combined organic layers were

washed with brine, dried and concentrated. The residue was triturated in ethyl acetate and the precipitate collected by filtration to give the title compound having the following physical properties (0.095 g).

LC/MS  $t_R$  1.70 minutes; MS (ES<sup>+</sup>)  $m/z$  554 (M+H)<sup>a</sup>.

5

Example 181: methyl [5-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl]-2-thienyl]carbamate

To a toluene (3 mL) suspension of the compound prepared in Example 180 (95 mg) was sequentially added glacial acetic acid (0.30 mL) and ammonium acetate (93 mg) and the mixture stirred at reflux for 40 minutes. Further ammonium acetate (62 mg) was added and the mixture stirred at reflux for 50 minutes. To the cooled (room temperature) reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (15 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in water) to give the title compound having the following physical properties (14 mg).

LC/MS  $t_R$  3.18 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.78 - 7.73 (m, 2 H), 7.70 (d, 1 H), 7.09 (br. s, 1 H), 6.96 (d, 1 H), 6.50 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.76 (dd, 1 H), 3.78 (s, 3 H), 3.50 - 3.41 (m, 1 H), 3.11 (ddd, 1 H), 2.67 - 2.59 (m, 1 H), 2.48 (app. br. s, 1 H).

20

Example 182(1) and 182(2): 2-({[4-(bromoacetyl)cyclohexyl]carbamoyl}oxy)-2-methylpropylidene and 2-({[4-(chloroacetyl)cyclohexyl]carbamoyl}oxy)-2-methylpropylidene

25

To a dichloromethane (100 mL) solution of 4-{{tert-butoxy}carbonyl}amino}cyclohexane-1-carboxylic acid [Eur. J. Med. Chem. 36(3), 265 (2001)] (2.80 g) was added 1-chloro-N,N,2-trimethyl-1-propenylamine (3.05 mL) and the mixture stirred at room temperature for 1 hour. To the cooled (0 °C) reaction mixture, a 2 M solution of (trimethylsilyl)diazomethane in hexanes (11.5 mL) was added and the mixture stirred at room temperature for 16 hours. The reaction was cooled (0 °C) and a solution of 48 wt. % hydrogen bromide in water (0.86 mL) was added and the mixture stirred at room temperature for 1 hour. A saturated aqueous

30

solution of sodium hydrogen carbonate (50 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was triturated with a 1: 2 mixture of ethyl acetate and heptanes and the resultant solid isolated via filtration to give the title compounds in a 1 :  
5 1 ratio having the following physical properties (2.13 g).

Example 182(1):

LC/MS  $t_R$  1.99 minutes; MS (ES<sup>+</sup>)  $m/z$  264 and 266 [M-C(CH<sub>3</sub>)<sub>3</sub>+H]<sup>a</sup>.

10 Example 182(2):

LC/MS  $t_R$  1.95 minutes; MS (ES<sup>+</sup>)  $m/z$  220 [M-C(CH<sub>3</sub>)<sub>3</sub>+H]<sup>a</sup>.

Example 183: 2-methyl-2-propanyl [trans-4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)cyclohexyl]carbamate

15 The same operation as in Example 178 → Example 179 → Example 180 → Example 181 was conducted from the 1: 1 mixture of compounds prepared in Example 182 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 180 in the operation, the compound prepared in  
20 Example 9 was used).

LC/MS  $t_R$  3.27 minutes; MS (ES<sup>+</sup>)  $m/z$  577 (M+H)<sup>b</sup>

NMR analysis showed a 2: 1 ratio of tautomers.

Major tautomer: <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.78 - 7.70 (m, 2 H), 7.68 (d, 1 H), 6.65 (s, 1 H), 6.58 (br. s, 1 H), 6.10 (s, 1 H), 6.07 (s, 1 H), 5.71 (app. d, 1  
25 H), 3.59 (br. s, 1 H), 3.43 - 3.31 (obs. m, 1 H), 3.12 - 3.02 (m, 1 H), 2.64 - 2.53 (m, 1 H), 2.49 (t, 1 H), 2.44 - 2.33 (m, 1 H), 2.07 - 1.92 (m, 4 H), 1.82 - 1.62 (m, 2 H), 1.44 (s, 9 H), 1.36 - 1.25 (m, 2 H).

Minor tautomer: <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.78 - 7.70 (m, 2 H), 7.68 (d, 1 H), 6.71 (s, 1 H), 6.58 (br. s, 1 H), 6.10 (s, 1 H), 6.09 (s, 1 H), 5.71 (app. d, 1  
30 H), 3.65 (br. s, 1 H), 3.43 - 3.31 (obs. m, 1 H), 3.12 - 3.02 (m, 1 H), 2.75 - 2.64 (m, 1 H), 2.64 - 2.53 (m, 1 H), 2.44 - 2.33 (m, 1 H), 2.07 - 1.92 (m, 4 H), 1.82 - 1.62 (m, 2 H), 1.44 (s, 9 H), 1.36 - 1.25 (m, 2 H).

Example 184: (3S)-3-[5-(trans-4-aminocyclohexyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 183 (0.32 g) was treated as detailed in Example 40 to give the title compound having the following physical properties (0.15 g).

- 5 LC/MS  $t_R$  2.28 minutes; MS ( $ES^+$ )  $m/z$  477 (M+H), 239 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.77 - 7.70 (m, 2 H), 7.67 (d, 1 H), 6.64 (br. s, 1 H), 6.10 (s, 1 H), 6.08 (s, 1 H), 5.70 (dd, 1 H), 3.42 - 3.32 (m, 1 H), 3.08 (dd, 1 H), 2.70 (tt, 1 H), 2.62 - 2.54 (m, 1 H), 2.54 - 2.45 (m, 1 H), 2.43 - 2.34 (m, 1 H), 2.09 - 1.91 (m, 4 H), 1.43 (ddt, 2 H), 1.28 (ddd, 2 H).

10

Example 185: methyl [trans-4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)cyclohexyl]carbamate

The compound prepared in Example 184 (78 mg) was treated as detailed in Example 128 to give the title compound having the following physical properties (19 mg). (Note: in the step corresponding to Example 128 in the operation, methyl chloroformate was used).

15

- LC/MS  $t_R$  2.92 minutes; MS ( $ES^+$ )  $m/z$  535 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.79 - 7.72 (m, 2 H), 7.70 (d, 1 H), 6.68 (br. s, 1 H), 6.12 (s, 1 H), 6.10 (s, 1 H), 5.72 (dt, 1 H), 3.71 (br. s, 1 H), 3.63 (s, 3  
20 H), 3.45 - 3.35 (m, 2 H), 3.13 - 3.04 (m, 1 H), 2.64 - 2.54 (m, 1 H), 2.41 (br. s, 1 H), 2.10 - 1.97 (m, 2 H), 1.80 (br. s, 2 H), 1.71 (br. s, 2 H), 1.40 - 1.29 (m, 2 H).

Example 186: (3S)-3-[5-(trans-4-aminocyclohexyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

- 25 The same operation as in Example 44 → Example 40 was conducted from the compound prepared in Example 183 to give the title compound having the following physical properties.

- LC/MS  $t_R$  2.94 minutes; MS ( $ES^+$ )  $m/z$  511 (M+H), 255.5 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.78 - 7.72 (m, 2 H), 7.70 (d, 1 H),  
30 6.11 (s, 1 H), 6.10 (s, 1 H), 5.65 (dd, 1 H), 3.48 - 3.35 (m, 1 H), 3.08 (ddd, 1 H), 2.75 (tt, 1 H), 2.68 (tt, 1 H), 2.64 - 2.54 (m, 1 H), 2.33 (ddt, 1 H), 2.01 (br. d, 2 H), 1.93 - 1.84 (m, 2 H), 1.68 - 1.56 (m, 2 H), 1.37 - 1.24 (m, 2 H).

Example 187: 2-([(4-methoxyphenyl)methyl]amino}pyridine-4-carbonitrile

A mixture of 4-cyano-2-fluoropyridine (1.0 g) and 4-methoxybenzylamine (2.25 g) was stirred at 100 °C for 1 hour and purified by column chromatography (0 to 50% ethyl acetate in heptanes) to give the title compound having the following physical

5 properties (1.69 g).

LC/MS  $t_R$  1.81 minutes; MS (ES<sup>+</sup>)  $m/z$  240 (M+H)<sup>a</sup>.

Example 188: 1-(2-([(4-methoxyphenyl)methyl]amino}pyridin-4-yl)ethan-1-one

To a diethyl ether (25 mL) solution of the compound prepared in Example 187  
10 (0.8 g) was added a 1.4 M solution of methyl magnesium bromide in 3: 1 toluene / tetrahydrofuran (7.2 mL) and the mixture stirred at room temperature for 16 hours. To the reaction mixture, a saturated aqueous solution of ammonium chloride (10 mL) was added followed by 6 M hydrochloric acid (5 mL) and the mixture stirred at room temperature for 1 hour. The mixture was then basified to pH 12 by addition of 2 M  
15 sodium hydroxide and extracted into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (0.47 g).

LC/MS  $t_R$  1.35 minutes; MS (ES<sup>+</sup>)  $m/z$  257 (M+H)<sup>a</sup>.

20 Example 189: 1-(2-aminopyridin-4-yl)-2-bromoethan-1-one dihydrobromide

To a glacial acetic acid (7 mL) solution of the compound prepared in Example 188 (0.38 g) was sequentially added 33 wt. % hydrogen bromide in acetic acid (1.28 mL) and bromine (76 µL) and the mixture stirred at 70 °C for 2 hours. The reaction mixture was concentrated and the residue triturated with dichloromethane (10 mL). The  
25 precipitate formed was isolated by filtration to give the title compound having the following physical properties (0.38 g).

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 7.88 - 7.83 (m, 1 H), 7.24 (d, 1 H), 7.01 (dd, 1 H), 3.71 (d, 1H), 3.64 (d, 1H).

30 Example 190: (3S)-3-[5-(2-amino-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the



compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 189 was used).

LC/MS  $t_R$  2.85 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>b</sup>


- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.39 (s, 1 H), 7.85 (d, 1 H), 7.78 - 7.74 (m, 2 H), 7.73 - 7.69 (m, 1 H), 7.52 (br. s, 1 H), 6.96 - 6.88 (m, 2 H), 6.17 (s, 1 H), 6.11 (s, 1 H), 5.80 (dd, 1 H), 3.50 - 3.41 (m, 1 H), 3.13 (ddd, 1 H), 2.70 - 2.61 (m, 1 H), 2.51 - 2.43 (m, 1 H).

10 Example 191: 2-bromo-1-(6-nitropyridin-3-yl)ethanone

To a stirred solution of 1-(6-nitropyridin-3-yl)ethan-1-one [patent WO2010/089292, page 71] (0.83 g) in tetrahydrofuran (10 mL) was added N-bromosuccinimide (1.78 g) and the mixture stirred at 50 °C for 16 hours. The reaction mixture was concentrated and the residue purified by column chromatography (20 to 100% ethyl acetate in dichloromethane) giving the title compound having the following physical properties.

LC/MS  $t_R$  1.53 minutes; MS (ES<sup>+</sup>)  $m/z$  245 and 247 (M+H)<sup>a</sup>.

20 Example 192: 3-[4-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone



- The same operation as in Example 84 → Example 39 → Example 40 → Example 41 → Example 74 was conducted from the compound prepared in Example 19 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 191 was used).

LC/MS  $t_R$  2.66 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.97 (br. s, 1 H), 9.68 (s, 1 H), 8.25 (d, 1 H), 7.83 - 7.75 (m, 3 H), 7.66 (dd, 1H), 7.28 (s, 1 H), 6.43 (d, 1 H), 5.96 (s, 1 H), 5.95 (s, 1 H), 5.81 (br. s, 2 H), 5.60 (d, 1 H), 3.43 - 3.31 (obs. m, 1 H), 3.05 - 2.92 (m, 1 H), 2.52 - 2.43 (m, 1 H), 2.39 - 2.29 (m, 1 H).

5

Example 193: 1-(6-aminopyridin-3-yl)-2-bromoethan-1-one dihydrobromide

1-(6-aminopyridin-3-yl)ethan-1-one [patent US2007/027184, page 17] (0.46 g) was treated as detailed in Example 91 to give the title compound having the following physical properties (0.70 g).

10 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.65 (d, 1 H), 8.01 (dd, 1 H), 6.60 (d, 1 H), 4.51 (s, 2 H).

Example 194: (3S)-3-[4-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

15 The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used).

LC/MS *t*<sub>R</sub> 1.38 minutes; MS (ES<sup>+</sup>) *m/z* 472 (M+H)<sup>a</sup>.

20

Example 195: methyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 194 (80 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the title compound having the following physical properties (66 mg).

25

LC/MS *t*<sub>R</sub> 3.11 minutes; MS (ES<sup>+</sup>) *m/z* 530 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 8.56 (app. br. s, 1 H), 8.02 (app. br. s, 1 H), 7.88 (app. br. s, 1 H), 7.76 - 7.70 (m, 2 H), 7.69 (d, 1 H), 7.41 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.77 (s, 3 H), 3.52 - 3.40 (m, 1 H), 3.16 - 3.06 (m, 1 H), 2.71 - 2.59 (m, 1 H), 2.52 (app. br. s, 1 H).

30

Example 196: methyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 195 (46 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (20 mg).

- 5 LC/MS  $t_R$  3.92 minutes; MS (ES<sup>+</sup>)  $m/z$  564 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.59 (d, 1 H), 8.07 (d, 1 H), 8.00 (d, 1 H), 7.78 - 7.73 (m, 2 H), 7.20 (d, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H), 5.73 (dd, 1 H), 3.52 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.72 - 2.61 (m, 1 H), 2.46 - 2.36 (m, 1 H).


10 Example 197: ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 194 (50 mg) was treated as detailed in Example 128 to give the title compound having the following physical properties (21 mg).

- 15 LC/MS  $t_R$  3.29 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.56 (br. s, 1 H), 8.07 - 8.00 (m, 1 H), 7.91 - 7.84 (m, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.42 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.22 (q, 2 H), 3.50 - 3.41 (m, 1 H), 3.15 - 3.06 (m, 1 H), 2.69 - 2.60 (m, 1 H), 2.55 - 2.46 (m, 1 H), 1.32 (t, 3 H).

20

Example 198: (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone



The compound prepared in Example 194 (50 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (13 mg).

- 25 LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.21 (s, 1 H), 7.78 - 7.71 (m, 3 H),

7.70 - 7.66 (m, 1 H), 6.65 (d, 1 H), 6.13 (s, 1 H), 6.08 (s, 1 H), 5.69 (d, 1 H), 3.49 - 3.39 (m, 1 H), 3.10 (dd, 1 H), 2.69 - 2.58 (m, 1 H), 2.42 - 2.34 (m, 1 H).

5 Example 199: 1-([5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2(1H)-pyridinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 25 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used).

- 10 LC/MS  $t_R$  2.61 minutes MS (ES<sup>+</sup>)  $m/z$  446 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 8.21 (br. s, 1 H), 7.81 - 7.72 (m, 3 H), 7.70 - 7.63 (m, 2 H), 7.26 (br. s, 1 H), 6.66 - 6.62 (m, 1 H), 6.42 - 6.38 (m, 1 H), 6.05 - 6.00 (m, 1 H), 5.17 (s, 2 H).

15 Example 200: methyl {5-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-2-pyridinyl}carbamate

The compound prepared in Example 199 (200 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the title compound having the following physical properties (9 mg).

- 20 LC/MS  $t_R$  3.06 minutes; MS (ES<sup>+</sup>)  $m/z$  504 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.27 (br. s, 1 H), 10.22 (br. s, 1 H), 9.72 (s, 1 H), 8.63 (s, 1 H), 8.01 (d, 1 H), 7.92 - 7.75 (m, 4 H), 7.68 (d, 1 H), 7.55 (br. s, 1 H), 6.26 (s, 1 H), 5.85 (dd, 1 H), 5.09 (s, 2 H), 3.64 (s, 3 H).

25 Example 201: 6-amino-N-methoxy-N-methylpyridine-3-carboxamide

- To an N,N-dimethylformamide (50 mL) solution of 6-aminopyridine-3-carboxylic acid (2.0 g) was added N,O-dimethylhydroxylamine hydrochloride (1.69 g), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (6.06 g) and diisopropylethylamine (7.6 mL) and the mixture stirred at room temperature for  
30 16 hours. To the reaction mixture, water (250 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated and the residue purified by column chromatography (0 to 15% methanol in dichloromethane) to give the title compound having the following physical properties

(0.61 g).

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 8.31 (d, 1 H), 7.69 (dd, 1 H), 6.42 (d, 1 H), 3.57 (s, 3 H), 3.21 (s, 3 H).

5    Example 202: tert-butyl N-[(tert-butoxy)carbonyl]-N-{5-[methoxy(methyl)carbamoyl]pyridin-2-yl}carbamate

The compound prepared in Example 201 (0.61 g) was treated as detailed in Example 10 to give the title compound having the following physical properties (0.58 g). LC/MS *t<sub>R</sub>* 1.99 minutes; MS (ES<sup>+</sup>) *m/z* 382 (M+H)<sup>a</sup>.

10

Example 203: tert-butyl N-(5-propanoylpyridin-2-yl)carbamate

To a cooled (0 °C) tetrahydrofuran (10 mL) solution of the compound prepared in Example 202 (0.53 g) was added a 1 M solution of ethylmagnesium bromide in tetrahydrofuran (1.73 mL) and the mixture stirred at room temperature for 16 hours. To  
15 the cooled (0 °C) reaction mixture, a saturated aqueous solution of ammonium chloride (1.5 mL) and water (30 mL) was sequentially added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (0.43 g). LC/MS *t<sub>R</sub>* 2.03 minutes; MS (ES<sup>+</sup>) *m/z* 251 (M+H)<sup>a</sup>.

20

Example 204: tert-butyl N-[5-(2-bromopropanoyl)pyridin-2-yl]carbamate

To a glacial acetic acid (5 mL) solution of the compound prepared in Example 203 (0.20 g) was added a solution of 33 wt. % hydrogen bromide in acetic acid (0.28 mL) followed by bromine (41 µL) and the mixture stirred at room temperature for 1  
25 hour. To the reaction mixture, a saturated solution of aqueous sodium bicarbonate was added (20 mL) followed by extraction into ethyl acetate. The combined organic layers were dried, filtered and concentrated and the residue purified by column chromatography (0 to 50% ethyl acetate in heptane) to give the title compound having the following physical properties (0.20 g).  
30 LC/MS *t<sub>R</sub>* 2.23 minutes; MS (ES<sup>+</sup>) *m/z* 329 and 331 (M+H)<sup>a</sup>.

Example 205: tert-butyl N-(5-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-methyl-1H-imidazol-5-yl}pyridin-2-yl)carbamate

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 204 was used).

LC/MS  $t_R$  1.66 minutes; MS (ES<sup>+</sup>)  $m/z$  586 (M+H)<sup>a</sup>.

Example 206: (3S)-3-[5-(6-amino-3-pyridinyl)-4-methyl-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

To a 1,4-dioxane (0.5 mL) solution of the compound prepared in Example 205 (42 mg) was added 1 M hydrochloric acid (0.4 mL) and the mixture heated at 90 °C for 3 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (40 mL) was added followed by extraction into ethyl acetate. The combined organic layers were dried, filtered and concentrated to give the title compound having the following physical properties (25 mg).

LC/MS  $t_R$  2.53 minutes; MS (ES<sup>+</sup>)  $m/z$  486 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.25 (s, 1 H), 7.91 (br. s, 1 H), 7.65 - 7.61 (m, 2 H), 7.60 - 7.55 (m, 1 H), 7.51 (d, 1 H), 6.54 (d, 1 H), 6.02 (s, 1 H), 5.98 (s, 1 H), 5.60 (dd, 1 H), 3.39 - 3.29 (m, 1 H), 2.98 (ddd, 1 H), 2.51 (qd, 1 H), 2.37 - 2.28 (m, 1 H), 2.18 (br. s, 3 H).

Example 207: methyl [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-methyl-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 206 (100 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the title compound having the following physical properties (28 mg).

LC/MS  $t_R$  3.00 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 8.41 (s, 1 H), 7.96 - 7.88 (m, 2 H), 7.77 - 7.73 (m, 2 H), 7.71 - 7.68 (m, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.74 (dd, 1 H),

3.79 (s, 3 H), 3.51 - 3.42 (m, 1 H), 3.16 - 3.07 (m, 1 H), 2.69 - 2.60 (m, 1 H), 2.51 - 2.42 (m, 1 H), 2.36 (s, 3 H).

Example 208: 6-chloro-5-iodopyridin-2-amine

- 5 To an N,N-dimethylformamide (30 mL) solution of 2-amino-6-chloropyridine (1.0 g) was added N-iodosuccinimide (1.75 g) and the mixture stirred at room temperature for 16 hours. To the reaction mixture, water (200 mL) was added followed by extraction into ethyl acetate. The combined organic layers were dried, filtered and concentrated. The residue was triturated with dichloromethane and the resultant
- 10 precipitate collected by filtration to give the title compound having the following physical properties (0.49 g).


LC/MS  $t_R$  1.72 minutes; MS ( $ES^+$ )  $m/z$  254 ( $M+H$ )<sup>a</sup>.

Example 209: 1-(6-amino-2-chloropyridin-3-yl)-2-bromoethan-1-one dihydrobromide

- 15 The same operation as in Example 90 → Example 91 was conducted from the compound prepared in Example 208 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.52 minutes; MS ( $ES^+$ )  $m/z$  249 and 251 ( $M+H$ )<sup>a</sup>.

- 20 Example 210: (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone



- The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following
- 25 physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 209 was used).

LC/MS  $t_R$  2.98 minutes; MS ( $ES^+$ )  $m/z$  506 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.81 (app. br. s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.32 (app. br. s, 1 H), 6.56 (d, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.79 (dd, 1 H), 3.50 - 3.38 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.57 (m, 1 H), 2.50 (tdd, 1 H).

5

Example 211: 1-([5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2(1H)-pyridinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 25 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 209 was used).

10

LC/MS *t<sub>R</sub>* 2.92 minutes; MS (ES<sup>+</sup>) *m/z* 481 (M+H)<sup>b</sup>

15

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.86 (br. s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.64 (m, 2 H), 7.39 (br. s, 1 H), 6.56 (d, 1 H), 6.40 (s, 1 H), 6.02 (dd, 1 H), 5.18 (s, 2 H).

Example 212: 1-(6-amino-2-methylpyridin-3-yl)-2-bromoethan-1-one dihydrobromide

The same operation as in Example 90 → Example 189 was conducted from 5-bromo-6-methylpyridin-2-amine to give the title compound having the following physical properties.

20

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.39 (d, 1 H), 6.96 (d, 1 H), 4.57 (s, 2 H), 2.75 (s, 3 H).

Example 213: (3S)-3-[5-(6-amino-2-methyl-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

25

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 212 was used).

30

LC/MS *t<sub>R</sub>* 2.62 minutes; MS (ES<sup>+</sup>) *m/z* 486 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 7.77 - 7.73 (m, 2 H), 7.72 - 7.68 (m, 1 H), 7.54 (d, 1 H), 6.99 (s, 1 H), 6.48 (d, 1 H), 6.14 (s, 1 H), 6.12 (s, 1 H), 5.80 (dd, 1 H), 3.51 - 3.39 (m, 1 H), 3.16 - 3.08 (m, 1 H), 2.69 - 2.59 (m, 1 H), 2.52 - 2.46 (m, 1 H),



2.41 (s, 3 H).

Example 214: methyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-6-methyl-2-pyridinyl]carbamate

5 The compound prepared in Example 213 (33 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the title compound having the following physical properties (21 mg).

LC/MS  $t_R$  3.07 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.89 (br. d, 1 H), 7.77 - 7.70 (m, 3 H),  
10 7.68 (d, 1 H), 7.19 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.80 (dd, 1 H), 3.51 - 3.41 (m, 1 H), 3.15 - 3.07 (m, 1 H), 2.68 - 2.60 (m, 1 H), 2.56 - 2.40 (obs. m, 1 H), 2.50 (s, 3 H).

Example 215: 1-(6-amino-2-fluoropyridin-3-yl)-2-bromoethan-1-one dihydrobromide

The same operation as in Example 90 → Example 189 was conducted from 2-  
15 amino-6-fluoro-5-iodopyridine [J. Org. Chem. 71(8), 2922 (2006)] to give the title compound having the following physical properties.

LC/MS  $t_R$  1.50 minutes; MS (ES<sup>+</sup>)  $m/z$  233 and 235 (M+H)<sup>a</sup>.

Example 216: (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone



The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the  
25 compound prepared in Example 215 was used).

LC/MS  $t_R$  2.94 minutes; MS (ES<sup>+</sup>)  $m/z$  490 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.97 (app. br. s, 1 H), 7.75 - 7.70 (m,

2 H), 7.68 (d, 1 H), 7.15 (br. s, 1 H), 6.45 (dd, 1 H), 6.14 (s, 1 H), 6.07 (s, 1 H), 5.79 (d, 1 H), 3.50 - 3.38 (m, 1 H), 3.14 - 3.05 (m, 1 H), 2.67 - 2.56 (m, 1 H), 2.52 - 2.44 (m, 1 H).

5 Example 217: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(6-methyl-3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 204 → Example 51 → Example 52 was conducted from 2-methyl-5-acetyl-pyridine to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the

10 operation, the compound prepared in Example 9 was used).

LC/MS  $t_R$  2.83 minutes; MS (ES<sup>+</sup>)  $m/z$  471 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.73 (app. s, 1 H), 8.01 (d, 1 H), 7.75 - 7.71 (m, 2 H), 7.68 (d, 1 H), 7.46 (br. s, 1 H), 7.30 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.51 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.70 - 2.59 (m, 2 H), 2.52 (s, 3 H).

15

Example 218: (3S)-3-[4-chloro-5-(6-methyl-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 217 (45 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (19 mg).

20 LC/MS  $t_R$  3.18 minutes; MS (ES<sup>+</sup>)  $m/z$  505 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.75 (d, 1 H), 8.05 (dd, 1 H), 7.76 - 7.72 (m, 2 H), 7.68 (d, 1 H), 7.40 (d, 1 H), 6.13 (s, 1 H), 6.10 (d, 1 H), 5.72 (dd, 1 H), 3.45 (td, 1 H), 3.12 (ddd, 1 H), 2.66 (qd, 1 H), 2.57 (s, 3 H), 2.44 - 2.36 (m, 1 H).

25 Example 219: tert-butyl N-[6-(2-bromoacetyl)pyridin-3-yl]carbamate

The same operation as in Example 10 → Example 188 → Example 91 was conducted from 5-amino-2-cyanopyridine to give the title compound having the following physical properties.

LC/MS  $t_R$  2.07 minutes; MS (ES<sup>+</sup>)  $m/z$  315 and 317 (M+H)<sup>a</sup>.

30

Example 220: 2-methyl-2-propanyl [6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The same operation as in Example 84 → Example 39 was conducted from the

compound prepared in Example 17 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 84 in the operation, the compound prepared in Example 219 was used).

LC/MS  $t_R$  3.37 minutes; MS (ES<sup>+</sup>)  $m/z$  572 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.54 (d, 1 H), 7.92 (dd, 1 H), 7.79 - 7.64 (m, 4 H), 7.48 (s, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.80 (dd, 1 H), 3.54 - 3.37 (m, 1 H), 3.18 - 3.03 (m, 1 H), 2.72 - 2.43 (m, 2 H), 1.53 (s, 9 H).

10 Example 221: 3-[5-(5-amino-2-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 220 (130 mg) was treated as detailed in Example 206 to give the title compound having the following physical properties (104 mg).

LC/MS  $t_R$  2.79 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.95 (br. s, 1 H), 7.75 - 7.71 (m, 2 H), 7.67 (d, 1 H), 7.53 (br. s, 1 H), 7.32 (br. s, 1 H), 7.10 (dd, 1 H), 6.14 (s, 1 H), 6.07 (s, 1 H), 5.79 (d, 1 H), 3.51 - 3.39 (m, 1 H), 3.09 (dd, 1 H), 2.65 - 2.56 (m, 1 H), 2.54 - 2.45 (m, 1 H).

20 Example 222: 3-[5-(5-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 44 → Example 206 was conducted from the compound prepared in Example 220 to give the title compound having the following physical properties.

- 25 LC/MS  $t_R$  3.38 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.04 (s, 1 H), 7.75 - 7.70 (m, 3 H), 7.69 - 7.66 (m, 1 H), 7.09 (dd, 1 H), 6.15 (s, 1 H), 6.07 (s, 1 H), 5.77 (dd, 1 H), 3.43 - 3.35 (m, 1 H), 3.12 - 3.05 (m, 1 H), 2.60 (qd, 1 H), 2.50 - 2.41 (m, 1 H).

30 Example 223: methyl [6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The compound prepared in Example 221 (60 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the

title compound having the following physical properties (30.2 mg).

LC/MS  $t_R$  3.00 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>

NMR analysis showed a 2: 1 ratio of tautomers.

Major tautomer: <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.18 (s, 1 H), 9.79 (br. s, 1 H), 9.69 (s, 1 H), 8.53 (d, 1 H), 7.91 - 7.81 (m, 1 H), 7.81 - 7.74 (m, 3 H), 7.69 (d, 1 H), 7.48 (d, 1 H), 5.97 (app. s, 2 H), 5.64 (dd, 1 H), 3.68 (s, 3 H), 3.40 - 3.30 (obs. m, 1 H), 3.07 - 2.90 (m, 1 H), 2.53 - 2.44 (obs. m, 1 H), 2.38 - 2.30 (m, 1 H).

Minor tautomer: <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.49 (s, 1 H), 9.89 (br. s, 1 H), 9.70 (s, 1 H), 8.63 (d, 1 H), 7.91 - 7.81 (m, 1 H), 7.81 - 7.74 (m, 3 H), 7.68 (d, 1 H), 7.33 (d, 1 H), 5.96 (s, 1 H), 5.95 (s, 1 H), 5.71 (dd, 1 H), 3.70 (s, 3 H), 3.29 - 3.21 (m, 1 H), 3.07 - 2.90 (m, 1 H), 2.53 - 2.44 (obs. m, 1 H), 2.29 - 2.21 (m, 1 H).

Example 224: 2-(5-aminopyridin-2-yl)-2-oxoethyl 2-{4-[5-chloro-2-(1,2,3,4-tetrazol-1-yl)phenyl]-2-oxopyridin-1-yl}acetate

The compound prepared in Example 25 (0.55 g) was treated as detailed in Example 51 using the compound prepared in Example 219 in place of 2-bromo-1-phenylethan-1-one to give the title compound having the following physical properties (0.26 g).

LC/MS  $t_R$  1.66 minutes; MS (ES<sup>+</sup>)  $m/z$  466 (M+H)<sup>a</sup>.

Example 225: 2-{5-[(methoxycarbonyl)amino]pyridin-2-yl}-2-oxoethyl 2-{4-[5-chloro-2-(1,2,3,4-tetrazol-1-yl)phenyl]-2-oxopyridin-1-yl}acetate

To a dichloromethane (25 mL) solution of the compound prepared in Example 224 (0.24 g) was added diisopropylethylamine (0.10 g) and methyl chloroformate (44  $\mu$ L) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated and the residue purified by column chromatography (0 to 80% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.10 g).

LC/MS  $t_R$  3.06 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>a</sup>.

Example 226: methyl {6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-pyridinyl}carbamate

The compound prepared in Example 225 (102 mg) was treated as detailed in

Example 52 to give the title compound having the following physical properties (17 mg).

LC/MS  $t_R$  2.99 minutes; MS (ES<sup>+</sup>)  $m/z$  504 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.59 (br. s, 1 H), 7.97 (br. d, 1 H),  
7.77 (app. br. s, 1 H), 7.77 - 7.71 (m, 2 H), 7.71 - 7.63 (m, 2 H), 7.55 (app. br. s, 1 H),  
5 6.42 (s, 1 H), 6.03 (d, 1 H), 5.22 (s, 2 H), 3.80 (s, 3 H).

Example 227: 2-(5-aminopyridin-2-yl)-2-oxoethyl (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

The compound prepared in Example 9 (0.39 g) was treated as detailed in

- 10 Example 51 using the compound prepared in Example 219 in place of 2-bromo-1-phenylethan-1-one to obtain the title compound having the following physical properties (0.54 g).

LC/MS  $t_R$  1.73 minutes; MS (ES<sup>+</sup>)  $m/z$  492 (M+H)<sup>a</sup>.

- 15 Examples 228(1) and 228(2): 2-(5-{[(tert-butoxy)carbonyl]amino}pyridin-2-yl)-2-oxoethyl (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate and 2-(5-{bis[(tert-butoxy)carbonyl]amino}pyridin-2-yl)-2-oxoethyl (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

- 20 To a dichloromethane (50 mL) solution of the compound prepared in Example 227 (0.54 g) was added di-*tert*-butyl dicarbonate (1.19 g) and 4-dimethylaminopyridine (0.27 g) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated and the residue purified by column chromatography (0 to 20% methanol in dichloromethane) to obtain the title compounds in a 2 : 1 ratio having the  
25 following physical properties (0.65 g).

Example 228(1):

LC/MS  $t_R$  2.09 minutes; MS (ES<sup>+</sup>)  $m/z$  592 (M+H)<sup>a</sup>

- 30 Example 228(2):

LC/MS  $t_R$  2.27 minutes; MS (ES<sup>+</sup>)  $m/z$  692 (M+H)<sup>a</sup>

Example 229: (3S)-3-[4-(5-aminopyridin-2-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 52 → Example 55 was conducted from the 2:1 ratio of compounds prepared in Example 228 (0.65 g) to obtain the title compound  
5 having the following physical properties (0.27 g).

LC/MS  $t_R$  1.54 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>a</sup>.

Example 230: methyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

10 The same operation as in Example 128 → Example 44 was conducted from the compound prepared in Example 229 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 128 in the operation, methyl chloroformate was used).

LC/MS  $t_R$  3.99 minutes; MS (ES<sup>+</sup>)  $m/z$  564 (M+H)<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.97 (br. s, 1 H), 10.00 (br. s, 1 H), 9.70 (s, 1 H), 8.73 (s, 1 H), 7.96 (dd, 1 H), 7.90 (d, 1 H), 7.85 - 7.77 (m, 3 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.67 (dd, 1 H), 3.70 (s, 3 H), 3.23 (td, 1 H), 3.02 - 2.92 (m, 1 H), 2.55 - 2.44 (obs. m, 1 H), 2.18 (t, 1 H).

20 Example 231(1) to Example 231(4)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 229 and the corresponding chloroformates or acetic anhydride using the method as detailed in Example 128.

25 Example 231(1): isopropyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS  $t_R$  3.32 minutes; MS (ES<sup>+</sup>)  $m/z$  558 (M+H)<sup>b</sup>

NMR analysis showed a 2: 1 ratio of tautomers.

Major tautomer: <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.19 (br. s, 1 H), 9.72 (br. s, 1 H),  
30 9.69 (s, 1 H), 8.52 (d, 1 H), 7.91 - 7.83 (m, 1 H), 7.82 - 7.77 (m, 3 H), 7.67 (d, 1 H), 7.48 (s, 1 H), 5.99 - 5.93 (m, 2 H), 5.63 (dd, 1 H), 4.90 (septuplet, 1 H), 3.30 - 3.21 (m, 1 H), 3.04 - 2.93 (m, 1 H), 2.56 - 2.42 (obs. m, 1 H), 2.38 - 2.29 (m, 1 H), 1.25 (d, 6 H).

Minor tautomer:  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.50 (br. s, 1 H), 9.82 (br. s, 1 H), 9.70 (s, 1 H), 8.64 (d, 1 H), 7.91 - 7.83 (m, 1 H), 7.82 - 7.77 (m, 3 H), 7.66 (d, 1 H), 7.32 (s, 1 H), 5.99 - 5.93 (m, 2 H), 5.70 (dd, 1 H), 4.90 (septuplet, 1 H), 3.30 - 3.21 (m, 1 H), 3.04 - 2.93 (m, 1 H), 2.56 - 2.42 (obs. m, 1 H), 2.28 - 2.20 (m, 1 H), 1.27 (d, 6 H).

5

Example 231(2): isobutyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS  $t_R$  3.52 minutes; MS ( $\text{ES}^+$ )  $m/z$  572 ( $\text{M}+\text{H}$ )<sup>b</sup>

NMR analysis showed a 1: 1 ratio of tautomers.

10  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  11.40 and 10.88 (br. s, 1 H), 8.55 (s, 1 H), 8.41 (br. s, 1 H), 8.13 - 7.92 (m, 1 H), 7.87 (d, 0.5 H), 7.61 (dd, 1 H), 7.58 - 7.43 (m, 3 H), 7.32 (br. s, 0.5 H), 7.14 - 7.02 and 6.77 (m and br. s, 1 H), 6.40 and 6.34 (br. s, 1 H), 5.86 (t, 1 H), 5.68 and 5.64 (br. s, 1 H), 3.97 (d, 2 H), 3.52 - 3.17 (m, 2 H), 3.00 (dd, 1 H), 2.53 - 2.39 (m, 1 H), 1.98 (dqintet, 1 H), 0.97 (d, 6 H).

15

Example 231(3): N-[6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]acetamide

LC/MS  $t_R$  2.94 minutes; MS ( $\text{ES}^+$ )  $m/z$  514 ( $\text{M}+\text{H}$ )<sup>b</sup>

NMR analysis showed a 2: 1 ratio of tautomers.

20 Major Tautomer:  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.21 (br. s, 1 H), 10.12 (s, 1 H), 9.69 (s, 1 H), 8.61 (d, 1 H), 8.00 (dd, 1 H), 7.82 - 7.77 (m, 3 H), 7.69 (d, 1 H), 7.50 (s, 1 H), 5.99 - 5.93 (m, 2 H), 5.63 (dd, 1 H), 3.33 - 3.21 (m, 1 H), 3.05 - 2.92 (m, 1 H), 2.56 - 2.42 (obs. m, 1 H), 2.38 - 2.29 (m, 1 H), 2.06 (s, 3 H).

Minor Tautomer:  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.52 (br. s, 1 H), 10.20 (s, 1 H),  
25 9.70 (s, 1 H), 8.70 (d, 1 H), 8.03 (dd, 1 H), 7.82 - 7.77 (m, 3 H), 7.68 (d, 1 H), 7.34 (s, 1 H), 5.99 - 5.93 (m, 2 H), 5.70 (d, 1 H), 3.33 - 3.21 (m, 1 H), 3.05 - 2.92 (m, 1 H), 2.56 - 2.42 (obs. m, 1 H), 2.28 - 2.20 (m, 1 H), 2.07 (s, 3 H).

Example 231(4): methyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

30

LC/MS  $t_R$  0.65 minutes; MS ( $\text{ES}^+$ )  $m/z$  530 ( $\text{M}+\text{H}$ )<sup>f</sup>

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.5 (s, 0.3 H), 12.2 (s, 0.7 H), 9.88 (s, 0.3 H), 9.77 (s, 0.7 H), 9.69 (s, 0.3 H), 9.67 (s, 0.7 H), 8.61 (d, 0.3 H), 8.51 (d, 0.7 H), 7.79 - 7.63 (m, 5

H), 7.47 (d, 0.7 H), 7.31 (d, 0.3 H), 5.97 - 5.92 (m, 2 H), 5.74 - 5.60 (m, 1 H), 3.69 (s, 0.9 H), 3.67 (s, 2.1 H), 3.41 - 3.17 (m, 1 H), 3.09 - 2.90 (m, 1 H), 2.61 - 2.22 (m, 2 H).

5 Examples 232(1) and 232(2): (3S)-3-[5-(5-amino-2-pyridinyl)-4-methyl-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone and N-[6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-methyl-1H-imidazol-5-yl)-3-pyridinyl]acetamide

The same operation as in Example 188 → Example 189 → Example 51 → Example 52 was conducted from 5-amino-2-cyanopyridine to obtain the title  
10 compounds in a 4 : 1 ratio having the following physical properties. (Note: in the step corresponding to Example 188 in the operation, a 3 M solution of ethyl magnesium bromide in tetrahydrofuran was used. In the step corresponding to Example 51 in the operation, the compound prepared in Example 9 was used).

15 Example 232(1):

LC/MS  $t_R$  2.88 minutes; MS (ES<sup>+</sup>)  $m/z$  508 (M+Na), 486 (M+H), 243 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.00 (br. s, 1 H), 7.75 - 7.70 (m, 2 H),  
7.70 - 7.66 (m, 1 H), 7.35 (app. br. s, 1 H), 7.11 (dd, 1 H), 6.13 (s, 1 H), 6.07 (s, 1 H),  
5.74 (app. br. s, 1 H), 3.08 (dd, 1 H), 2.65 - 2.55 (m, 1 H), 2.50 - 2.43 (m, 2 H), 2.36 (br.  
20 s, 3 H).

Example 232(2):

LC/MS  $t_R$  2.92 minutes; MS (ES<sup>+</sup>)  $m/z$  528 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.71 (d, 1 H), 8.05 (br. s, 1 H), 7.73  
25 (app. s, 2 H), 7.71 - 7.56 (m, 2 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.75 (d, 1 H), 3.50 - 3.40  
(m, 1 H), 3.14 - 3.05 (m, 1 H), 2.67 - 2.58 (m, 1 H), 2.52 - 2.43 (obs. m, 1 H), 2.46 (br. s,  
3 H), 2.16 (s, 3 H).

Example 233: methyl [6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-methyl-1H-imidazol-5-yl)-3-pyridinyl]carbamate  
30

The compound prepared in Example 232(1, 80 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the



title compound having the following physical properties (22.9 mg).

LC/MS  $t_R$  3.04 minutes; MS ( $ES^+$ )  $m/z$  544 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.60 (br. s, 1 H), 7.95 (app. br. s, 1 H), 7.72 (app. s, 2 H), 7.70 - 7.67 (m, 1 H), 7.62 (app. br. s, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H),  
5 5.75 (d, 1 H), 3.77 (s, 3 H), 3.51 - 3.39 (m, 1 H), 3.13 - 3.05 (m, 1 H), 2.67 - 2.57 (m, 1 H), 2.52 - 2.41 (obs. m, 1 H), 2.47 (s, 3 H).

Example 234: tert-butyl N-[5-(2-bromoacetyl)-1,3-thiazol-2-yl]carbamate

To an cooled (0 °C) dichloromethane (4 mL) solution of 2-[(tert-  
10 butoxycarbonyl)amino]-1,3-thiazole-5-carboxylic acid (0.30 g) was added a solution of oxalyl chloride (0.13 mL) in dichloromethane (1 mL). One drop of N,N-dimethylformamide was added and the mixture stirred at room temperature for 1 hour. The reaction mixture was concentrated and the residue dissolved in acetonitrile (5 mL). To the cooled (0 °C) acetonitrile solution, a 2 M solution of  
15 (trimethylsilyl)diazomethane in hexanes (1.23 mL) was added and the mixture stirred at 0 °C for 1 hour. To the reaction mixture, a solution of 33 wt. % hydrogen bromide in acetic acid (0.20 mL) was added and the mixture stirred at 0 °C for a further 20 minutes. A saturated aqueous solution of sodium hydrogen carbonate (50 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed  
20 with brine, dried and concentrated to give the crude title compound having the following physical properties (0.37 g).  
LC/MS  $t_R$  1.92 minutes; MS ( $ES^+$ )  $m/z$  265 and 267 ( $M-C(CH_3)_3+H$ )<sup>a</sup>.

Example 235: (3S)-3-[5-(2-amino-1,3-thiazol-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 40 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 234 was used).  
30 LC/MS  $t_R$  2.79 minutes; MS ( $ES^+$ )  $m/z$  478 ( $M+H$ ), 239 [ $(M+2H)/2$ ]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.76 - 7.70 (m, 2 H), 7.68 (d, 1 H), 7.10 (s, 1 H), 7.09 (br. s, 1 H), 6.13 (s, 1 H), 6.07 (s, 1 H), 5.72 (dd, 1 H), 3.46 - 3.36 (m, 1 H), 3.09 (ddd, 1 H), 2.66 - 2.56 (m, 1 H), 2.46 (app. br. s, 1 H).

Example 236: (3S)-3-[5-(1-amino-6-isoquinolinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone diacetate

The same operation as in Example 90 → Example 189 → Example 51 →

- 5 Example 52 was conducted from 6-bromoisoquinolin-1-amine to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 9 was used).

LC/MS  $t_R$  2.94 minutes; MS (ES<sup>+</sup>)  $m/z$  522 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 8.24 (d, 1 H), 8.11 (br. s, 1 H), 7.99  
10 (d, 1 H), 7.77 - 7.73 (m, 2 H), 7.70 (d, 1 H), 7.66 (br. s, 1 H), 7.58 (d, 1 H), 7.15 (d, 1 H),  
6.16 (s, 1 H), 6.12 (s, 1 H), 5.82 (dd, 1 H), 3.53 - 3.43 (m, 1 H), 3.14 (ddd, 1 H), 2.67  
(ddd, 1 H), 2.54 - 2.44 (m, 1 H), 1.98 (s, 6 H).

Example 237: methyl N-(4-carbamimidoylphenyl)carbamate

- 15 To a cooled (0 °C) ethanol (10 mL) suspension of methyl N-(4-cyanophenyl)carbamate [J. Chem. Soc., Perkin Trans. 1, 2587 (1984)] was added acetyl chloride (4 mL) and the mixture stirred at room temperature for 16 hours. Further acetyl chloride (1 mL) was added and the mixture stirred at 50 °C for 3 hours. The reaction mixture was concentrated and the residue dissolved in ethanol (10 mL). To the solution,  
20 ammonium carbonate (2.73 g) was added and the mixture stirred at room temperature for 6 hours. The solids were removed by filtration and the filtrate concentrated. The residue was triturated with dichloromethane (10 mL) and the precipitate isolated by filtration to give the title compound having the following physical properties.

LC/MS  $t_R$  0.66 minutes; MS (ES<sup>+</sup>)  $m/z$  194 (M+H)<sup>a</sup>.

25

Examples 238(1) and 238(2): tert-butyl N-{2-[3-(2-bromoacetyl)-5-oxo-2,3-dihydro-1H-indolizin-7-yl]-4-chlorophenyl}carbamate and tert-butyl N-{4-chloro-2-[3-(2-chloroacetyl)-5-oxo-2,3-dihydro-1H-indolizin-7-yl]phenyl}carbamate

- The compound prepared in Example 19 (0.50 g) was treated as detailed in  
30 Example 182 to give a 2: 7 mixture of the title compounds having the following physical properties (22.9 mg).

Example 238(1):

LC/MS  $t_R$  2.10 minutes; MS ( $ES^+$ )  $m/z$  503 and 505 ( $M+Na$ ), 481 and 483 ( $M+H$ )<sup>a</sup>.

Example 238(2):

LC/MS  $t_R$  2.07 minutes; MS ( $ES^+$ )  $m/z$  459 ( $M+Na$ ), 437 ( $M+H$ )<sup>a</sup>.

5

Example 239: tert-butyl N-{4-chloro-2-[3-(2-{4-[(methoxycarbonyl)amino]phenyl}-3H-imidazol-4-yl)-5-oxo-2,3-dihydro-1H-indolizin-7-yl]phenyl}carbamate

To an acetonitrile (10 mL) suspension of the compound prepared in Example 237 (0.125 g) and the 2 : 7 ratio of compounds prepared in Examples 238(1) and 238(2) (0.26 g) was added potassium carbonate (0.16 g) and the mixture stirred at reflux for 4 hours. Water (50 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated and the residue purified by column chromatography (0 to 5% methanol in dichloromethane) to give the title compound having the following physical properties (0.21 g).

15 LC/MS  $t_R$  1.78 minutes; MS ( $ES^+$ )  $m/z$  576 ( $M+H$ )<sup>a</sup>.

Example 240: methyl [4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl]phenyl]carbamate

The same operation as in Example 40 → Example 41 was conducted from the compound prepared in Example 239 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.99 minutes; MS ( $ES^+$ )  $m/z$  529 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.40 (s, 1 H), 7.80 - 7.74 (m, 4 H), 7.71 (d, 1 H), 7.54 (d, 2 H), 6.77 (s, 1 H), 6.18 (s, 1 H), 6.09 (s, 1 H), 5.82 (d, 1 H), 3.77 (s, 3 H), 3.40 - 3.36 (m, 1 H), 3.06 (dd, 1 H), 2.64 - 2.54 (m, 1 H), 2.42 (dd, 1 H).

25

Example 241: methyl [4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl]phenyl]carbamate

The compound prepared in Example 240 (80 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (21 mg).

LC/MS  $t_R$  3.79 minutes; MS ( $ES^+$ )  $m/z$  563 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.78 - 7.72 (m, 4 H), 7.72 - 7.69 (m, 1 H), 7.55 (d, 2 H), 6.21 (s, 1 H), 6.05 (s, 1 H), 5.81 (dd, 1 H), 3.77 (s, 3 H), 3.40 - 3.34

30

(m, 1 H), 3.14 (ddd, 1 H), 2.75 - 2.65 (m, 1 H), 2.33 (tdd, 1 H).

Example 242: tert-butyl N-[4-chloro-2-(3-{N'-[(4-nitrophenyl)carboximidoyl]hydrazinecarbonyl}-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl)phenyl]carbamate

To a cooled (0 °C) tetrahydrofuran (5 mL) solution of the compound prepared in Example 19 (0.40 g) was added ethyl chloroformate (0.11 mL) and 4-methylmorpholine (0.16 mL) and the mixture stirred at 0 °C for 25 minutes. A solution of N-amino-4-nitrobenzenecarboximidamide [Justus Liebigs Ann. Chem. 298, 51 (1897)] in tetrahydrofuran (5 mL) was added and the mixture stirred at room temperature for 16 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (25 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 to 10% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.23 g).  
LC/MS  $t_R$  1.79 minutes; MS (ES<sup>+</sup>)  $m/z$  567 (M+H)<sup>a</sup>.

Example 243: 7-(2-amino-5-chlorophenyl)-3-[5-(4-nitrophenyl)-4H-1,2,4-triazol-3-yl]-1,2,3,5-tetrahydroindolizin-5-one

A glacial acetic acid (2.1 mL) solution of the compound prepared in Example 242 (0.23 g) was stirred at 110 °C for 3 hours. To the mixture, a saturated aqueous solution of sodium hydrogen carbonate (30 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 to 20% methanol in ethyl acetate) to obtain the title compound having the following physical properties (0.086 g).  
LC/MS  $t_R$  1.97 minutes; MS (ES<sup>+</sup>)  $m/z$  449 (M+H)<sup>a</sup>.

Example 244: 3-[5-(4-aminophenyl)-4H-1,2,4-triazol-3-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 41 → Example 74 was conducted from the compound prepared in Example 243 (110 mg) to obtain the title compound having the following physical properties (77 mg).

LC/MS  $t_R$  3.31 minutes; MS ( $ES^+$ )  $m/z$  472 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.68 (br. s, 1 H), 9.68 (s, 1 H), 7.80 (app. s, 3 H), 7.58 (d, 2 H), 6.61 (d, 2 H), 5.95 (s, 1 H), 5.93 (s, 1 H), 5.62 (d, 1 H), 5.57 (br. s, 2 H), 3.27 - 3.08 (m, 1 H), 2.95 (dd, 1 H), 2.58 - 2.44 (obs. m, 1 H), 2.15 - 2.00 (m, 1 H).

5

Example 245: methyl [4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4H-1,2,4-triazol-3-yl)phenyl]carbamate

The compound prepared in Example 244 (65 mg) was treated as detailed in Example 128 using methyl chloroformate in place of ethyl chloroformate to give the title compound having the following physical properties (29.6 mg).

10

LC/MS  $t_R$  3.67 minutes; MS ( $ES^+$ )  $m/z$  530 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  12.89 (br. s, 1 H), 8.71 (br. s, 1 H), 7.77 (app. br. s, 2 H), 7.65 - 7.53 (m, 2 H), 7.53 - 7.45 (m, 1 H), 7.34 (app. br. s, 2 H), 6.31 (br. s, 1 H), 5.91 (br. s, 1 H), 5.76 (app. br. s, 1 H), 3.75 (br. s, 3 H), 3.46 - 3.28 (m, 1 H), 3.07 - 2.91 (m, 1 H), 2.79 - 2.46 (m, 2 H).

15

Example 246: methyl 7-(2-{[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

To a dichloromethane (7.5 mL) and methanol (2.5 mL) solution of the compound prepared in Example 19 (1.0 g) was added a 2 M solution of (trimethylsilyl)diazomethane in diethyl ether (2.2 mL) at 0 °C and the mixture stirred at room temperature for 16 hours. To the reaction mixture, acetic acid (0.5 mL) and a saturated aqueous solution of sodium hydrogen carbonate (30 mL) were sequentially added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (10% to 100% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.83 g).

20

25

LC/MS  $t_R$  2.01 minutes; MS ( $ES^+$ )  $m/z$  419 ( $M+H$ )<sup>a</sup>.

30

Example 247: tert-butyl N-(4-chloro-2-{3-[2-(dimethoxyphosphoryl)acetyl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

To a cooled (-78 °C) tetrahydrofuran (20 mL) solution of dimethyl methylphosphonate (1.1 mL) was added a 2.5 M solution of *n*-butyllithium in hexanes

- (4.0 mL) and the mixture stirred at -78 °C for 30 minutes. A solution of the compound prepared in Example 246 (0.83 g) in tetrahydrofuran (10 mL) was added and the mixture stirred at -78 °C for 1 hour. To the cooled (-78 °C) reaction mixture, a saturated aqueous solution of ammonium chloride (5 mL) was added followed by water (30 mL) and the mixture was allowed to warm to room temperature then extracted with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (20% to 100% ethyl acetate in heptanes, then 0 to 20% methanol in ethyl acetate) to obtain the title compound having the following physical properties (0.67 g).
- LC/MS  $t_R$  1.92 minutes; MS (ES<sup>+</sup>)  $m/z$  511 (M+H)<sup>a</sup>.

Example 248: tert-butyl N-(4-chloro-2-{3-[5-(4-nitrophenyl)-6-oxo-1,6-dihydropyridazin-3-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

- To a cooled (0 °C) ethanol (7 mL) suspension of the compound prepared in Example 247 (0.67 g) and potassium carbonate (0.27 g) was added ethyl (4-nitrophenyl)(oxo)acetate (0.31 g) and the mixture stirred at 0 °C for 90 minutes then at room temperature for 2 hours. Hydrazine hydrate (0.6 mL) was added and the mixture stirred at room temperature for 3 hours. To the reaction mixture, 0.5 M hydrochloric acid (25 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 to 100% ethyl acetate in heptanes). The product was triturated with a 1: 2 mixture of dichloromethane and heptanes and the solid collected by filtration to give the title compound having the following physical properties (0.24 g) LC/MS  $t_R$  2.08 minutes; MS (ES<sup>+</sup>)  $m/z$  576 (M+H)<sup>a</sup>.

Example 249: methyl [4-(6-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-3-oxo-2,3-dihydro-4-pyridazinyl)phenyl]carbamate

- The same operation as in Example 74 → Example 128 → Example 40 → Example 41 was conducted from the compound prepared in Example 248 (0.24 g) to obtain the title compound having the following physical properties (0.075 g). (Note: in the step corresponding to Example 128 in the operation, methyl chloroformate was used).

LC/MS  $t_R$  3.69 minutes; MS (ES<sup>+</sup>)  $m/z$  557 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.98 (br. s, 1 H), 9.89 (s, 1 H), 9.71 (s, 1 H), 7.88 (d, 2 H), 7.82 - 7.78 (m, 3 H), 7.61 (s, 1 H), 7.56 (d, 2 H), 5.96 (s, 1 H), 5.94 (s, 1 H), 5.55 (dd, 1 H), 3.68 (s, 3 H), 3.16 - 3.05 (m, 1 H), 2.97 (ddd, 1 H), 2.59 - 2.46 (obs. m, 1 H), 2.25 - 2.15 (m, 1 H).

5

Example 250: tert-butyl N-(4-chloro-2-{3-[methoxy(methyl)carbamoyl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

To an N,N-dimethylformamide (12.5 mL) solution of the compound prepared in Example 19 (1.0 g) was added N,O-dimethylhydroxylamine hydrochloride (0.36 g), O-  
10 (7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (1.0 g) and diisopropylethylamine (1.2 mL) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated and the residue suspended in a saturated aqueous solution of sodium hydrogen carbonate (30 mL) followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and  
15 concentrated. The residue was purified by column chromatography (0% to 20% methanol in ethyl acetate) to obtain the title compound having the following physical properties (0.73 g).

LC/MS *t*<sub>R</sub> 1.99 minutes; MS (ES<sup>+</sup>) *m/z* 448 (M+H)<sup>a</sup>.

20 Example 251: tert-butyl N-[4-chloro-2-(3-formyl-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl)phenyl]carbamate

To a cooled (-45 °C) tetrahydrofuran (6.5 mL) solution of the compound prepared in Example 250 (0.60 g) was added a 1 M solution of lithium aluminium hydride in tetrahydrofuran (1.6 mL) over 10 minutes. The reaction mixture was stirred  
25 at -45 °C for 30 minutes, 0 °C for 90 minutes then cooled to -45 °C and a solution of potassium hydrogen sulfate (0.36 g) in water (1.0 mL) added. The mixture was warmed to room temperature and extracted into ethyl acetate. The combined organic layers were washed sequentially with 1 M hydrochloric acid (25 mL), a saturated aqueous solution of sodium hydrogen carbonate (30 mL) and brine (25 mL), dried and concentrated to  
30 give the title compound having the following physical properties (0.45 g).

LC/MS *t*<sub>R</sub> 1.84 minutes; MS (ES<sup>+</sup>) *m/z* 389 (M+H)<sup>a</sup>.

Example 252: tert-butyl N-[4-chloro-2-(3-ethynyl-5-oxo-1,2,3,5-tetrahydroindolizin-7-

yl)phenyl]carbamate

- To a methanol (15 mL) suspension of the compound prepared in Example 251 (0.45 g) and potassium carbonate (0.32 g) was added dimethyl (1-diazo-2-oxopropyl)phosphonate (0.2 mL) and the mixture stirred at room temperature for 4 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (25 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 to 100% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.29 g).
- LC/MS  $t_R$  2.02 minutes; MS ( $ES^+$ )  $m/z$  385 (M+H)<sup>a</sup>.

Example 253: tert-butyl N-(2-{3-[1-(4-aminophenyl)-1H-1,2,3-triazol-4-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}-4-chlorophenyl)carbamate

- To an acetonitrile (1.2 mL) solution of the compound prepared in Example 252 (0.14 g) was added 4-azidoaniline hydrochloride (59 mg), copper(II) sulfate pentahydrate (8 mg), a solution of sodium ascorbate (31 mg) in water (0.60 mL) and 1 M sodium hydroxide (0.35 mL). The reaction mixture was irradiated under microwave conditions (120 W) at 80 °C for 1 hour and a saturated aqueous solution of sodium hydrogen carbonate (30 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography ((0 to 100% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (0.16 g).
- LC/MS  $t_R$  2.01 minutes; MS ( $ES^+$ )  $m/z$  519 (M+H)<sup>a</sup>.

25 Example 254: methyl [4-(4-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-1,2,3-triazol-1-yl)phenyl]carbamate

- The same operation as in Example 128 → Example 40 → Example 41 was conducted from the compound prepared in Example 253 (0.16 g) to obtain the title compound having the following physical properties (0.11 g). (Note: in the step corresponding to Example 128 in the operation, methyl chloroformate was used).
- LC/MS  $t_R$  3.88 minutes; MS ( $ES^+$ )  $m/z$  530 (M+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.94 (s, 1 H), 9.69 (s, 1 H), 8.55 (s, 1 H), 7.83 - 7.74 (m, 5 H), 7.64 (d, 2 H), 5.96 (s, 1 H), 5.95 (s, 1 H), 5.79 (d, 1 H), 3.69 (s, 3 H), 3.39 -



3.33 (m, 1 H), 3.04 (dd, 1 H), 2.61 - 2.52 (m, 1 H), 2.32 (dd, 1 H).

Example 255: tert-butyl N-(4-chloro-2-{3-[3-(4-nitrophenyl)-1,2-oxazol-5-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

- 5 To a *tert*-butanol (1.0 mL) and water (1.0 mL) suspension of 4-nitrobenzaldehyde (63 mg) was added hydroxylamine hydrochloride (29 mg) and the mixture stirred at room temperature for 1 hour. Chloramine-T trihydrate (117 mg) was added and the mixture stirred at room temperature for 5 minutes to afford the intermediate nitrile oxide.
- 10 To the suspension, copper powder (2.4 mg), copper(II) sulfate pentahydrate (9.4 mg) and a solution of the compound prepared in Example 252 (145 mg) in *tert*-butanol (1.0 mL) was sequentially added and the mixture stirred at room temperature for 16 hours. The nitrile oxide was re-prepared as detailed above, added to the reaction and the mixture stirred at room temperature for 16 hours then at 80 °C for 20 hours. To the
- 15 reaction mixture, a 10% aqueous solution of ammonia (20 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue obtained on concentration was purified by column chromatography (12% to 80% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (49.5 mg).
- 20 LC/MS  $t_R$  2.33 minutes; MS (ES<sup>+</sup>)  $m/z$  549 (M+H)<sup>a</sup>.

Example 256: methyl [4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1,2-oxazol-3-yl)phenyl]carbamate

- The same operation as in Example 40 → Example 41 → Example 74 →
- 25 Example 128 was conducted from the compound prepared in Example 255 (49.5 mg) to obtain the title compound having the following physical properties (8.4 mg). (Note: in the step corresponding to Example 128 in the operation, methyl chloroformate was used).
- LC/MS  $t_R$  4.12 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>
- 30 <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 9.91 (s, 1 H), 9.71 (s, 1 H), 7.83 - 7.79 (m, 3 H), 7.77 (d, 2 H), 7.58 (d, 2 H), 6.75 (s, 1 H), 6.02 (s, 1 H), 6.00 (s, 1 H), 5.86 (dd, 1 H), 3.69 (s, 3 H), 3.18 (td, 1 H), 3.09 - 2.99 (m, 1 H), 2.61 - 2.53 (m, 1 H), 2.27 - 2.19 (m, 1 H).

Example 257: 2-methyl-2-propanyl (2S)-2-[2-oxo-4-{{(trifluoromethyl)sulfonyl}oxy}-1(2H)-pyridinyl]-3-phenylpropanoate

The same operation as in Example 20 → Example 21 → Example 22 was conducted from 2-methyl-2-propanyl (2R)-3-phenyl-2-

- 5    {{(trifluoromethyl)sulfonyl}oxy}propanoate (WO2006/57961) to give the title compound having the following physical properties.

TLC: Rf 0.33 (20% ethyl acetate in hexane).

Example 258: (2S)-2-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-3-phenylpropanoic acid

10

The same operation as in Example 23 → Example 25 was conducted from the compound prepared in Example 257 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 23 in the operation, 2,5-dichlorophenylboronic acid was used).

- 15    TLC: Rf 0.18 (10% methanol in dichloromethane).

Example 259: 4-(4-chloro-2-{{(1S)-1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-5-yl})benzoic acid

- 20    The same operation as in Example 72 → Example 73 → Example 76 → Example 49 was conducted from the compound prepared in Example 258 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 72 in the operation, methyl 4-(bromoacetyl)benzoate was used).

TLC: Rf 0.41 (10% methanol in dichloromethane)

- 25    <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.21 (br. s, 1 H), 8.10 - 7.92 (m, 3 H), 7.85 (s, 2 H), 7.66 - 7.44 (m, 3 H), 7.34 - 7.11 (m, 5 H), 6.51 - 6.31 (m, 3 H), 3.60 (dd, 1 H), 3.39 (dd, 1 H).

Example 260: 4-(2-{{(1S)-1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-5-yl})benzoic acid

30

The same operation as in Example 72 → Example 73 → Example 49 was conducted from the compound prepared in Example 258 to give the title compound having the following physical properties. (Note: in the step corresponding to Example

72 in the operation, methyl 4-(bromoacetyl)benzoate was used).

TLC: Rf 0.47 (10% methanol in dichloromethane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.98 (br. s, 1 H), 8.08 - 7.91 (m, 5 H), 7.86 (d, 1 H), 7.68 - 7.40 (m, 3 H), 7.36 - 7.03 (m, 5 H), 6.45 (s, 1 H), 6.38 - 6.32 (m, 1 H), 6.32 - 6.23 (m, 1 H), 3.86 - 3.72 (m, 1 H), 3.65 - 3.48 (m, 1 H).

Example 261: 3-(4-chloro-2-((1S)-1-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl]-2-phenylethyl)-1H-imidazol-5-yl)benzoic acid

The same operation as in Example 72 → Example 73 → Example 76 →

Example 49 was conducted from the compound prepared in Example 258 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 72 in the operation, methyl 3-(bromoacetyl)benzoate was used).

TLC: Rf 0.45 (10% methanol in dichloromethane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.29 (br. s, 1 H), 12.98 (br. s, 1 H), 8.31 (s, 1 H), 8.03 - 7.94 (m, 2 H), 7.89 (d, 1 H), 7.66 - 7.45 (m, 4 H), 7.31 - 7.11 (m, 5 H), 6.50 - 6.32 (m, 3 H), 3.60 (dd, 1 H), 3.47 - 3.36 (m, 1 H).

Example 262: 3-{2-[(S)-[4-(2,5-dichlorophenyl)-2-oxo-1(2H)-pyridinyl](phenyl)methyl]-1H-imidazol-5-yl}benzoic acid

The same operation as in Example 20 → Example 21 → Example 22 → Example 23 → Example 30 → Example 72 → Example 73 → Example 49 was conducted from methyl (2R)-chloro(phenyl)acetate [Justus Liebigs Annalen der Chemie, 501, 208, (1933)] to give the title compound having the following physical properties. (Note: in the step corresponding to Example 23 in the operation, 2,5-dichlorophenylboronic acid was used. In the step corresponding to Example 72 in the operation, methyl 3-(bromoacetyl)benzoate was used).

TLC: Rf 0.57 (10% methanol in dichloromethane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.21 (br. s, 1 H), 12.71 (br. s, 1 H), 8.32 (br. s, 1 H), 7.97 (d, 2 H), 7.77 (d, 2 H), 7.65 - 7.23 (m, 10 H), 6.54 (d, 1 H), 6.42 (d, 1 H).

Example 263: methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl]-3-methoxypropyl)-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 26 → Example 21 → Example 22 → Example 23 → Example 24 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 20 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 26 in the operation, 2-chloroethyl methyl ether was used).

TLC: Rf 0.26 (ethyl acetate)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 11.17 (br. s, 1 H), 8.67 - 8.51 (m, 1 H), 7.82 - 7.33 (m, 8 H), 7.20 (br. s, 1 H), 6.53 - 5.99 (m, 2 H), 5.84 - 5.64 (m, 1 H), 3.89 - 3.68 (m, 3 H), 3.50 - 3.30 (m, 2 H), 3.28 - 3.08 (m, 3 H), 2.71 - 2.33 (m, 2 H).

Example 264: methyl {4-[4-chloro-2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}-3-methoxypropyl)-1H-imidazol-5-yl]phenyl} carbamate

The compound prepared in Example 263 (40 mg) was treated as detailed in Example 47 to give the title compound having the following physical properties (18 mg).

TLC: Rf 0.57 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 11.78 (br. s, 1 H), 8.59 (s, 1 H), 7.79 - 7.39 (m, 8 H), 6.77 (s, 1 H), 6.40 (s, 1 H), 6.19 (br. s, 1 H), 5.79 (d, 1 H), 3.79 (s, 3 H), 3.53 - 3.32 (m, 2 H), 3.32 - 3.06 (m, 3 H), 2.59 (d, 1 H), 2.45 (br. s, 1 H).


Example 265: methyl (4-{2-[(1S)-1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}-2-phenylethyl]-1H-imidazol-4-yl}phenyl)carbamate

The same operation as in Example 23 → Example 24 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 257 to give the title compound having the following physical properties.

TLC: Rf 0.57 (80% ethyl acetate in hexane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.30 (s, 1 H), 9.76 - 9.49 (m, 2 H), 7.91 - 7.61 (m, 5 H), 7.61 - 7.35 (m, 4 H), 7.34 - 6.99 (m, 5 H), 6.40 - 6.10 (m, 2 H), 5.89 (dd, 1 H), 3.66 (s, 3 H), 3.58 - 3.43 (m, 1 H), 3.27 (br. s, 1 H).

Example 266: methyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-4-yl)phenyl]carbamate



The same operation as in Example 38 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 11 to give the title compound having the following physical properties.

5 TLC: R<sub>f</sub> 0.29 (ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (s, 1 H), 9.75 - 9.63 (m, 1 H), 9.58 (s, 1 H), 7.87 - 7.72 (m, 3 H), 7.66 - 7.31 (m, 5 H), 6.01 - 5.88 (m, 2 H), 5.68 - 5.53 (m, 1 H), 3.72 - 3.59 (m, 3 H), 3.48 - 3.34 (m, 1 H), 2.99 (dd, 1 H), 2.46 - 2.10 (m, 2 H).

10 Example 267: 5-[(5-chloro-2-nitrophenyl)(hydroxy)methylene]-2,2-dimethyl-1,3-dioxane-4,6-dione:

A mixture of 5-chloro-2-nitrobenzoic acid (11.0 g) and thionyl chloride (11.8 mL) was stirred at 90 °C for 2 hours then concentrated. The resulting residue was dissolved in dichloromethane (50 mL) and added to a stirred solution of 2,2-dimethyl-1,3-dioxane-4,6-dione (7.86 g) and 4-(dimethylamino)pyridine (13.3 g) in dichloromethane (200 mL) at -12 °C. After being stirred for 2 hours, the reaction mixture was washed with 1 M hydrochloric acid (100 mL x 3) and water, dried and concentrated. The residue thus obtained was triturated with acetonitrile followed by acetone to afford the title compound having the following physical properties (6.0 g).

20 TLC: R<sub>f</sub> 0.43 (ethyl acetate).

Example 268: methyl (3R)-7-(5-chloro-2-nitrophenyl)-5-oxo-2,3-dihydro-5H-[1,3]thiazolo[3,2-a]pyridine-3-carboxylate

To a dichloromethane (80 mL) solution of the compound prepared in Example 267 (7.67 g) was added (4S)-methyl 2-methyl-4,5-dihydro-1,3-thiazole-4-carboxylate [Journal of Organic Chemistry, 66(20), 6756, (2001)] (1.57 g) and the stirred mixture treated with gaseous hydrogen chloride at 0 °C for 15 minutes. The reaction mixture

was then heated for 24 hours at 64 °C for before allowing to cool to room temperature. To the reaction mixture, water was added followed by extraction with dichloromethane. The combined organic layers were washed sequentially with a saturated aqueous solution of sodium hydrogen carbonate and brine, dried and concentrated. The residue  
5 was purified by column chromatography (30 – 55% ethyl acetate in hexanes) to give the title compound having the following physical properties (310 mg).  
TLC: Rf 0.35 (50% ethyl acetate in hexane).

10 Example 269: methyl (3R)-7-[2-(bis{[(2-methyl-2-propanyl)oxy]carbonyl}amino)-5-chlorophenyl]-5-oxo-2,3-dihydro-5H-[1,3]thiazolo[3,2-a]pyridine-3-carboxylate

To a mixture of ammonium chloride (250 mg) and zinc powder (341 mg) in water (2.5 mL) was added a solution of the compound prepared in Example 268 (234 mg) in ethyl acetate (12.5 mL) and the mixture stirred for 3 hours at room temperature. The reaction mixture was filtered through Celite® and the filtrate was washed with  
15 brine, dried and concentrated. The residue thus obtained was dissolved in tetrahydrofuran (5 mL) and to this solution triethylamine (0.16 mL), 4-(dimethylamino)pyridine (13 mg) and di-*tert*-butyl dicarbonate (247 mg) were sequentially added. The reaction mixture was warmed to 60 °C for 1 hour before allowing to cool to room temperature. To the reaction mixture, a saturated aqueous  
20 solution of potassium bisulfate was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (70% ethyl acetate in hexanes) to give the title compound having the following physical properties (189 mg).  
TLC: Rf 0.63 (80% ethyl acetate in hexane).

25

Example 270: methyl [4-(2-{(3R)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-2,3-dihydro-5H-[1,3]thiazolo[3,2-a]pyridin-3-yl}-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 11 → Example 38 → Example 39 →  
Example 40 → Example 41 was conducted from the compound prepared in Example  
30 269 to give the title compound having the following physical properties.  
TLC: Rf 0.49 (5% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.58 (br. s, 1 H), 8.58 (s, 1 H), 7.72 (d, 2 H), 7.67 - 7.28 (m, 4 H), 7.26 - 7.06 (m, 2 H), 6.80 - 6.45 (m, 1 H), 6.23 (d, 1 H), 6.17 - 5.97 (m, 1 H),

5.80 (s, 1 H), 4.71 (d, 1 H), 4.00 - 3.65 (m, 4 H).

Example 271(1) to Example 271(4)

The compounds of the present invention having the following physical data were  
5 prepared using the corresponding alpha-bromoketones from the compound prepared in  
Example 11 in the process of Example 38 → Example 39 → Example 40 → Example  
41

Example 271(1): methyl [3-chloro-4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-  
10 5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl)phenyl]carbamate  
TLC 0.55 (5% methanol in ethyl acetate, NH Silica)  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.20 (s, 1 H), 9.82 (s, 1 H), 9.74 - 9.58 (m, 1 H),  
7.93 - 7.84 (m, 1 H), 7.84 - 7.72 (m, 3 H), 7.62 (d, 1 H), 7.58 - 7.36 (m, 2 H), 5.96 (s, 2  
H), 5.73 - 5.55 (m, 1 H), 3.73 - 3.61 (m, 3 H), 3.46 - 3.33 (m, 1 H), 3.10 - 2.87 (m, 1 H),  
15 2.47 - 2.19 (m, 2 H).

Example 271(2): 6-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-  
tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone  
TLC: R<sub>f</sub> 0.28 (10% methanol in ethyl acetate, NH Silica)  
20 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.01 (s, 1 H), 10.03 (s, 1 H), 9.79 - 9.50 (m, 1 H),  
7.85 - 7.70 (m, 3 H), 7.60 - 7.28 (m, 3 H), 6.98 - 6.61 (m, 1 H), 6.13 - 5.84 (m, 2 H),  
5.60 (d, 1 H), 3.54 - 3.19 (m, 3 H), 3.10 - 2.77 (m, 3 H), 2.45 - 2.07 (m, 2 H).

Example 271(3): ethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-  
25 1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate  
TLC: R<sub>f</sub> 0.52 (10% methanol in ethyl acetate)  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.02 (s, 1 H), 9.79 - 9.60 (m, 1 H), 9.54 (s, 1 H),  
7.95 - 7.68 (m, 3 H), 7.68 - 7.34 (m, 5 H), 6.09 - 5.80 (m, 2 H), 5.73 - 5.48 (m, 1 H),  
4.19 - 4.06 (m, 2 H), 3.46 - 3.35 (m, 1 H), 2.99 (dd, 1 H), 2.61 - 2.53 (m, 1 H), 2.40 -  
30 2.24 (m, 1 H), 1.24 (t, 3 H).

Example 271 (4): 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-  
tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzamide

TLC: Rf 0.55 (10% methanol in dichloromethane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.22 (s, 1 H), 9.72 - 9.66 (m, 1 H), 7.99 - 7.60 (m, 9 H), 7.25 (br. s, 1 H), 6.07 - 5.82 (m, 2 H), 5.67 - 5.53 (m, 1 H), 3.52 - 3.35 (m, 1 H), 3.01 (dd, 1 H), 2.46 - 2.24 (m, 2 H).

5

Example 272: 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzamide

The compound prepared in Example 271(4) (250 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (152 mg).

10

TLC: Rf 0.64 (6% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.11 (s, 1 H), 9.70 (s, 1 H), 8.12 - 7.74 (m, 8 H), 7.40 (br. s, 1 H), 6.06 - 5.82 (m, 2 H), 5.58 (dd, 1 H), 3.28 - 3.15 (m, 1 H), 3.00 (dd, 1 H), 2.65 - 2.54 (m, 1 H), 2.33 - 2.11 (m, 1 H).

15

Example 273: methyl [4-(2-((3R)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in example 42 was separated by preparative chiral HPLC (CHIRALPAK IB 50 mm x 250 mm, hexane/tetrahydrofuran/diethylamine = 20/80/0.1) to give the title compound having the following physical properties.

20

TLC: Rf 0.50 (ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (s, 1 H), 9.81 - 9.63 (m, 1 H), 9.58 (s, 1 H), 7.87 - 7.73 (m, 3 H), 7.68 - 7.29 (m, 5 H), 6.04 - 5.89 (m, 2 H), 5.68 - 5.54 (m, 1 H), 3.71 - 3.59 (m, 3 H), 3.47 - 3.35 (m, 1 H), 2.99 (dd, 1 H), 2.43 - 2.14 (m, 2 H).

25

Example 274(1) and Example 274(2): 2-methyl-2-propanyl {4-chloro-2-[(3S)-3-(5-{4-[(methoxycarbonyl)amino]phenyl}-4-methyl-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl} carbamate and 2-methyl-2-propanyl {4-chloro-2-[(3S)-3-(4-{4-[(methoxycarbonyl)amino]phenyl}-5-methyl-1,3-oxazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl} carbamate

30

The same operation as in Example 38 → Example 39 was conducted from the compound prepared in Example 11 to obtain the title products in a 1: 1 ratio having the following physical properties. (Note: in the step corresponding to Example 38 in the



operation, the compound prepared in Example 78 was used)

Example 274(1):

LC/MS  $t_R$  4.13 minutes; MS (ES<sup>+</sup>)  $m/z$  589 (M+H)<sup>d</sup>.

5 Example 274(2):

LC/MS  $t_R$  4.59 minutes; MS (ES<sup>+</sup>)  $m/z$  590 (M+H)<sup>d</sup>.

Example 275: methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-5-methyl-1H-imidazol-4-yl)phenyl]carbamate

10 The same operation as in Example 40 → Example 41 was conducted from the compound prepared in Example 274(1) to give the title compound having the following physical properties.

TLC: R<sub>f</sub> 0.54 (5% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.87 (s, 1 H), 9.81 - 9.62 (m, 1 H), 9.58 (s, 1 H),  
15 7.92 - 7.68 (m, 3 H), 7.54 - 7.36 (m, 4 H), 6.06 - 5.84 (m, 2 H), 5.63 - 5.40 (m, 1 H),  
3.72 - 3.60 (m, 3 H), 3.47 - 3.35 (m, 1 H), 3.10 - 2.86 (m, 1 H), 2.47 - 2.39 (m, 2 H),  
2.32 (s, 3 H).

Example 276: methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-5-methyl-1,3-oxazol-4-yl)phenyl]carbamate

20 The same operation as in Example 40 → Example 41 was conducted from the compound prepared in Example 274(2) to give the title compound having the following physical properties.

TLC: R<sub>f</sub> 0.45 (5% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.86 - 9.51 (m, 2 H), 7.81 (app. s, 3 H), 7.52 (app. s,  
25 4 H), 6.00 - 5.97 (m, 2 H), 5.66 (dd, 1 H), 3.67 (s, 3 H), 3.25 - 3.09 (m, 1 H), 3.08 - 2.94  
(m, 1 H), 2.48 (s, 3H), 2.31 - 2.10 (m, 2 H).

Example 277: methyl {4-[2-((4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl)methyl)-4-methyl-1H-imidazol-5-yl]phenyl}carbamate

30 The same operation as in Example 38 → Example 39 → Example 40 → Example 41 was conducted from the compound prepared in Example 32 to obtain the title compound having the following physical properties. (Note: in the step

corresponding to Example 38 in the operation, the compound prepared in Example 78 was used)

TLC: Rf 0.32 (5% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.01 (br. s, 1 H), 8.53 (s, 1 H), 7.68 - 7.59 (m, 1 H),  
5 7.59 - 7.32 (m, 7 H), 6.63 (br. s, 1 H), 6.54 (d, 1 H), 5.73 (d, 1 H), 5.08 (s, 2 H), 3.80 (s, 3 H), 2.40 (br. s, 3 H).

Example 278: methyl (4-{2-[1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl]-2-(3-pyridinyl)ethyl]-1H-imidazol-5-yl}phenyl)carbamate

10 The same operation as in Example 26 → Example 21 → Example 22 → Example 23 → Example 24 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 20 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 26 in the operation, 3-(chloromethyl)pyridine hydrochloride was used).

15 TLC: Rf 0.30 (5% methanol in ethyl acetate, NH Silica)  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.32 (s, 1 H), 9.80 - 9.59 (m, 2 H), 8.47 - 8.29 (m, 2 H), 7.87 - 7.61 (m, 6 H), 7.61 - 7.35 (m, 4 H), 7.34 - 7.19 (m, 1 H), 6.44 - 6.07 (m, 2 H), 5.92 (d, 1 H), 3.66 (s, 3 H), 3.62 - 3.40 (m, 1 H), 3.33 - 3.16 (m, 1 H).

20 Example 279: methyl (4-{4-chloro-2-[1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl]-2-(3-pyridinyl)ethyl]-1H-imidazol-5-yl}phenyl)carbamate

The compound prepared in Example 278 (126 mg) was treated as detailed in Example 47 to give the title compound having the following physical properties (60 mg).

TLC: Rf 0.21 (5% methanol in ethyl acetate)

25 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.94 (s, 1 H), 9.80 (s, 1 H), 9.64 (s, 1 H), 8.56 - 8.24 (m, 2 H), 7.95 - 7.68 (m, 4 H), 7.56 (q, 4 H), 7.45 - 7.12 (m, 2 H), 6.34 - 6.20 (m, 1 H), 6.16 (d, 1 H), 5.95 (dd, 1 H), 3.74 - 3.62 (m, 3 H), 3.54 (dd, 1 H), 3.28 (br. s, 1 H).

Example 280: 1-azido-2-bromo-4-chlorobenzene

30 To a solution of concentrated hydrochloric acid (100 mL) in water (300 mL) was added 2-bromo-4-chloroaniline (10.0 g). To this stirred suspension, a solution of sodium nitrite (3.68 g) in water (150 mL) was added dropwise at -5 °C. After being stirred at -5 °C for 1 hour, a solution of sodium azide (3.46 g) in water (15 mL) was

added dropwise and the resulting suspension stirred at -5 °C for a further hour. The reaction mixture was then extracted with ethyl acetate and the organic layer washed with brine, dried and concentrated to obtain the title compound having the following physical properties (10.7 g).

5 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.56 (d, 1 H), 7.32 (dd, 1 H), 7.09 (d, 1 H).

Example 281: 2-methyl-2-propanyl 1-(2-bromo-4-chlorophenyl)-1H-1,2,3-triazole-4-carboxylate

To a toluene (50 mL) solution of the compound prepared in Example 280 (2.4 g),  
10 tert-butyl propiolate (3 mL) was added and the mixture stirred at reflux for 6 hours. The reaction mixture was concentrated and the residue purified by column chromatography (7 to 20% ethyl acetate in hexane) to give the title compound having the following physical properties (2.3 g).

LC/MS *t*<sub>R</sub> 4.51 minutes; MS (ES<sup>+</sup>) *m/z* 358 and 360 (M+H)<sup>d</sup>.

15

Example 282: ethyl (3S)-5-oxo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,5-tetrahydro-3-indolizinecarboxylate

To a 1,4-dioxane (7 mL) solution of the compound prepared in Example 6 (500 mg) was sequentially added bis(pinacolate)diboron (393 mg) and potassium acetate  
20 (415 mg). The mixture was degassed with argon then 1,1'-bis(diphenylphosphino)ferrocene palladium (II) dichloride dichloromethane complex (115 mg) was added and the reaction mixture stirred at 100 °C for 3 hours. To the reaction mixture, ethyl acetate was added followed by filtration with Celite. The filtrate was washed with water and brine, dried and concentrated to obtain the title compound  
25 having the following physical properties (412 mg).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 6.86 (br s, 1 H), 6.40 (br s, 1 H), 5.11 (dd, 1 H), 4.39 - 4.17 (m, 2 H), 3.25 - 3.10 (m, 1 H), 3.09 - 2.96 (m, 1 H), 2.55 - 2.39 (m, 1 H), 2.32 - 2.20 (m, 1 H), 1.39 - 1.21 (m, 15 H).

30 Example 283: ethyl (3S)-7-[5-chloro-2-(4-{[(2-methyl-2-propanyl)oxy]carbonyl}-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylate

To a 1,4-dioxane (30 mL) solution of the compound prepared in Example 281 (1.01 g) was added the compound prepared in Example 282 (1.10 g) and a solution of

sodium carbonate (1.20 g) in water (6.0 mL). To the reaction mixture, tetrakis(triphenylphosphine)palladium(0) (487 mg) was added then the mixture degassed with argon and stirred for 16 h at 100 °C. The reaction mixture was concentrated and the residue thus obtained suspended in ethyl acetate and washed with water and brine. The organic layer was dried and concentrated and the residue purified by column chromatography (15% methanol in ethyl acetate) to give the title compound having the following physical properties (613 mg).  
LC/MS  $t_R$  4.30 minutes; MS (ES<sup>+</sup>)  $m/z$  485 (M+H)<sup>d</sup>.

10 Example 284: 2-methyl-2-propanyl 1-{4-chloro-2-[(3S)-3-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl}-1H-1,2,3-triazole-4-carboxylate

The same operation as in Example 8 → Example 38 → Example 39 was conducted from the compound prepared in Example 283 to give the title compound having the following physical properties.

TLC: R<sub>f</sub> 0.72 (6% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.02 (s, 1 H), 9.57 (s, 1 H), 8.99 (s, 1 H), 7.80 - 7.70 (m, 3 H), 7.62 - 7.37 (m, 5 H), 6.07 (s, 1 H), 5.93 - 5.89 (m, 1 H), 5.60 (d, 1 H), 3.70 - 3.63 (m, 3 H), 3.52 - 3.36 (m, 1 H), 3.03 (dd, 1 H), 2.47 - 2.33 (m, 2 H), 1.59 - 1.47 (m, 9 H).

Example 285: methyl 1-{4-chloro-2-[(3S)-3-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl}-1H-1,2,3-triazole-4-carboxylate

25 To a methanol (1 mL) solution of the compound prepared in Example 284 (91 mg) was added a 20% solution of sodium methoxide in methanol (50 µL) and the mixture stirred for 2 hours at 40 °C. To the cooled (0 °C) reaction mixture, a saturated aqueous solution of sodium bisulfate was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated and the residue purified by column chromatography (20% ethyl acetate in methanol) to give the title compound having the following physical properties (52 mg).

TLC: R<sub>f</sub> 0.55 (6% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.02 (s, 1 H), 9.57 (s, 1 H), 9.11 (s, 1 H), 7.79 - 7.70

(m, 3 H), 7.61 - 7.36 (m, 5 H), 6.01 (s, 1 H), 5.92 (s, 1 H), 5.59 (d, 1 H), 3.87 - 3.80 (m, 3 H), 3.65 (s, 3 H), 3.51 - 3.35 (m, 1 H), 3.06 - 2.95 (m, 1 H), 2.47 - 2.25 (m, 2 H).

5 Example 286: 1-{4-chloro-2-[(3S)-3-(5-{4-[(methoxycarbonyl)amino]phenyl})-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indoliziny]phenyl}-1H-1,2,3-triazole-4-carboxylic acid

To a dichloromethane (1.5 mL) solution of the compound prepared in Example 284 (51 mg) was added trifluoroacetic acid (0.5 mL). After stirring for 5 hours at 45 °C, the reaction mixture was concentrated and the resulting residue triturated with ethyl acetate to give the title compound having the following physical properties (27 mg).

10 TLC: R<sub>f</sub> 0.41 (ethyl acetate/methanol/acetic acid, 10/5/2)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.84 (s, 1 H), 9.03 (s, 1 H), 7.94 - 7.60 (m, 7 H), 7.60 - 7.45 (m, 2 H), 6.12 (s, 1 H), 5.98 (d, 1 H), 5.75 (dd, 1 H), 3.68 (s, 3 H), 3.30 (dt, 1 H), 3.20 - 2.94 (m, 1 H), 2.76 - 2.56 (m, 1 H), 2.40 - 2.16 (m, 1 H).

15

Example 287: 1-(2-bromo-4-chlorophenyl)-1H-1,2,3-triazole-4-carboxylic acid

To a dichloromethane (10 mL) solution of the compound prepared in Example 281 (2.3 g), was added trifluoroacetic acid (10 mL). After being stirred at room temperature for 1 hour, the reaction mixture was concentrated and the resulting residue triturated with dichloromethane to give the title compound having the following physical properties (1.5 g).

20

LC/MS t<sub>R</sub> 4.07 minutes; MS (ES<sup>+</sup>) m/z 302, 304 (M+H)<sup>d</sup>.

Example 288: 1-(2-bromo-4-chlorophenyl)-1H-1,2,3-triazole-4-carboxamide

25 To an N,N-dimethylformamide (3 mL) solution of the compound prepared in Example 287 (500 mg) was sequentially added triethylamine (0.69 mL) and *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (753 mg). After being stirred at room temperature for 15 minutes, a 28% aqueous solution of ammonia (0.7 mL) was added and the mixture stirred at room temperature for a further 15 minutes. To the reaction mixture, water was added and the resulting precipitate collected by filtration. The precipitate was washed with water and dried to give the title compound having the following physical properties (463 mg).

30

LC/MS t<sub>R</sub> 4.02 minutes; MS (ES<sup>+</sup>) m/z 301, 303 (M+H)<sup>d</sup>.

Example 289: methyl [4-(2-((3S)-7-[2-(4-carbamoyl-1H-1,2,3-triazol-1-yl)-5-chlorophenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]carbamate

5           The same operation as in Example 283 → Example 8 → Example 38 → Example 39 was conducted from the compound prepared in Example 288 to give the title compound having the following physical properties.

TLC: R<sub>f</sub> 0.34 (10% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.10 - 11.89 (m, 1 H), 9.57 (s, 1 H), 8.94 - 8.69 (m, 1 H), 7.97 (s, 1 H), 7.83 - 7.64 (m, 3 H), 7.64 - 7.26 (m, 6 H), 6.08 - 5.99 (m, 1 H), 5.93 (s, 1 H), 5.61 (d, 1 H), 3.73 - 3.59 (m, 3 H), 3.50 - 3.40 (m, 1 H), 3.09 - 2.93 (m, 1 H), 2.43-2.27 (m, 2 H).

Example 290: methyl [4-(2-((3S)-7-[5-chloro-2-(4-cyano-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]carbamate

15           To a dichloromethane (50 mL) solution of the compound prepared in Example 289 (70 mg) was added triethylamine (0.17 mL) and trifluoroacetic anhydride (0.17 mL) and the reaction stirred at room temperature for 1 hour. The reaction mixture was concentrated and the resulting residue purified by column chromatography (5% ethyl acetate in methanol, NH silica) to give the title compound having the following physical properties (55 mg).

TLC: R<sub>f</sub> 0.53 (10% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.04 (s, 1 H), 9.57 (s, 1 H), 9.37 (d, 1 H), 7.84 - 7.76 (m, 3 H), 7.62 - 7.51 (m, 2 H), 7.49 - 7.37 (m, 3 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.65- 5.57 (m, 1 H), 3.68 - 3.63 (m, 3 H), 3.47 - 3.36 (m, 1 H), 3.01 (dd, 1 H), 2.41 - 2.08 (m, 2 H).

Example 291: methyl 1-(2-bromo-4-chlorophenyl)-1H-1,2,3-triazole-4-carboxylate

30           To a methanol (5 mL) solution of the compound prepared in Example 287 was added concentrated sulfuric acid (0.2 mL) and the mixture heated at reflux for 6 hours before allowing to cool to room temperature. The reaction mixture was concentrated, diluted with water (5 mL) and neutralized with a saturated aqueous solution of sodium hydrogen carbonate followed by extraction with ethyl acetate. The combined organic

layers were washed with brine, dried and concentrated to give the title compound having the following physical properties (132 mg).

TLC: R<sub>f</sub> 0.50 (50% ethyl acetate in hexane).

5 Example 292: 2-methyl-2-propanyl (2S)-2-[2-oxo-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1(2H)-pyridinyl]-3-phenylpropanoate

The compound prepared in Example 257 (1.88 g) was treated as detailed in Example 282 to give the title compound having the following physical properties (1.35 g).

10 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.30 - 7.05 (m, 6 H), 6.93 (br s, 1 H), 6.27 (dd, 1 H), 5.47 (dd, 1 H), 3.43 (dd, 1 H), 3.25 (dd, 1 H), 1.40 (s, 9 H), 1.30 (s, 12 H).

15 Example 293: methyl 1-(4-chloro-2-{1-[(1S)-1-(5-{4-[(methoxycarbonyl)amino]phenyl}-1H-imidazol-2-yl)-2-phenylethyl]-2-oxo-1,2-dihydro-4-pyridinyl}phenyl)-1H-1,2,3-triazole-4-carboxylate

The same operation as in Example 283 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 292 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 283 in the operation, the compound prepared in Example 291 was used)

20 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.29 (s, 1 H), 9.61 (s, 1 H), 9.20 - 8.90 (m, 1 H), 7.92 - 7.60 (m, 6 H), 7.57 - 7.35 (m, 3 H), 7.30 - 7.06 (m, 5 H), 6.39 - 6.08 (m, 2 H), 6.03 - 5.84 (m, 1 H), 3.85 (s, 3 H), 3.73 - 3.61 (m, 3 H), 3.59 - 3.40 (m, 1 H), 3.31 - 3.09 (m, 1 H).

25

Example 294: 1-(2-bromo-4-chlorophenyl)-4-(trifluoromethyl)-1H-1,2,3-triazole

The compound prepared in Example 280 (2.0 g) was treated as detailed in Example 281 using 3,3,3-trifluoropropyne in place of tert-butyl propiolate to give the title compound having the following physical properties (100 mg)

30 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.25 (m, 1 H), 7.81 (m, 1 H), 7.50 - 7.55 (m, 2 H).

Example 295: methyl [4-(2-{(1S)-1-[4-{5-chloro-2-[4-(trifluoromethyl)-1H-1,2,3-triazol-1-yl]phenyl}-2-oxo-1(2H)-pyridinyl]-2-phenylethyl}-1H-imidazol-4-

yl)phenyl]carbamate

The same operation as in Example 283 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 292 to give the title compound having the following physical properties. (Note: in the step

- 5 corresponding to Example 283 in the operation, the compound prepared in Example 294 was used)

TLC: R<sub>f</sub> 0.53 (66% ethyl acetate in hexane)

- <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.00 (br. s, 1 H), 9.19 (s, 1 H), 8.99 (s, 1 H), 7.87 - 7.60 (m, 5 H), 7.58 - 7.38 (m, 3 H), 7.38 - 7.31 (m, 1 H), 7.30 - 6.98 (m, 5 H), 6.43 - 10 6.12 (m, 2 H), 5.88 (dd, 1 H), 3.69 (s, 3 H), 3.57 (dd, 1 H), 3.27 (dd, 1 H).

Example 296: 1-(2-bromo-4-chlorophenyl)-4-(tributylstannyl)-1H-1,2,3-triazole

The compound prepared in Example 280 (1.86 g) was treated as detailed in Example 281 using ethynyltri-n-butyltin in place of tert-butyl propiolate to give the title

- 15 compound having the following physical properties (5.03 g)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.83 (m, 1 H), 7.75 (dd, 1 H), 7.50 (dd, 1 H), 7.45 (dd, 1 H), 1.64 - 1.50 (m, 6 H), 1.42 - 1.24 (m, 6 H), 1.22 - 1.12 (m, 6 H), 0.89 (t, 9H).

Example 297: 1-(2-bromo-4-chlorophenyl)-4-chloro-1H-1,2,3-triazole

- 20 To an acetonitrile (72 mL) solution of the compound prepared in Example 296 (2.42 g) was added N-chlorosuccinimide (885 mg) and the mixture stirred for 15 hours at 60 °C before cooling to room temperature. The reaction mixture was concentrated and the resulting residue purified by column chromatography (20% ethyl acetate in hexane) to give the title compound having the following physical properties (1.0 g).

- 25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.91 (s, 1 H), 7.79 (t, 1 H), 7.48 – 7.51 (m, 2 H).

Example 298: methyl (4-{2-[(1S)-1-{4-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}-2-phenylethyl]-1H-imidazol-5-yl}phenyl)carbamate

- The same operation as in Example 283 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 292 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 283 in the operation, the compound prepared in Example 297 was used)
- 30



TLC: Rf 0.62 (66% ethyl acetate in hexane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.30 (br. s, 1 H), 9.61 (s, 1 H), 8.65 (s, 1 H), 7.96 - 7.63 (m, 5 H), 7.63 - 7.35 (m, 4 H), 7.35 - 7.05 (m, 5 H), 6.47 - 6.09 (m, 2 H), 5.89 (d, 1 H), 3.77 - 3.62 (m, 3 H), 3.52 (dd, 1 H), 3.26 (d, 1 H).

5

Example 299: 1-(2-bromo-4-chlorophenyl)-1H-pyrazole

To a N,N-dimethylacetamide (20 mL) solution of 2-bromo-4-chloro-1-fluorobenzene (2.0 g) was added cesium carbonate (5.39 g) and pyrazole (780 mg) and the reaction mixture stirred for 3 hours at 100 °C before allowing to cool to room  
10 temperature. The reaction mixture was diluted with ethyl acetate, washed with water and brine, dried and concentrated to give the title compound having the following physical properties (1.52 g).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.81 (d, 1 H), 7.73 (d, 1 H), 7.71 (d, 1 H), 7.46 (d, 1 H), 7.40 (dd, 1 H), 6.47 (t, 1 H).

15

Example 300: 1-(2-bromo-4-chlorophenyl)-4-fluoro-1H-pyrazole

To an acetonitrile (4.5 mL) solution of the compound prepared in Example 299 (300 mg) was added 1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate) (619 mg) and the reaction mixture stirred for 18 hours at 70 °C  
20 before allowing to cool to room temperature. The reaction mixture was diluted with ethyl acetate, washed with a saturated aqueous solution of sodium hydrogen carbonate and brine, dried and concentrated. The residue thus obtained was purified by column chromatography (20% ethyl acetate in hexane) to obtain the title compound having the following physical properties (131 mg).

25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.75 - 7.69 (m, 2 H), 7.60 (dd, 1 H), 7.44 (dd, 1 H), 7.40 (dd, 1 H).

Example 301: methyl (4-{2-[(1S)-1-{4-[5-chloro-2-(4-fluoro-1H-pyrazol-1-yl)phenyl]-2-oxo-1(2H)-pyridinyl}-2-phenylethyl]-1H-imidazol-5-yl}phenyl)carbamate

30 The same operation as in Example 283 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 292 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 283 in the operation, the compound prepared in Example 300

was used)

TLC: Rf 0.40 (50% ethyl acetate in hexane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.30 (s, 1 H), 9.61 (s, 1 H), 8.05 - 7.92 (m, 1 H), 7.81 (d, 1 H), 7.76 - 7.38 (m, 9 H), 7.38 - 7.03 (m, 5 H), 6.47 - 6.10 (m, 2 H), 5.99 - 5.64 (m, 1 H), 3.69 - 3.61 (m, 3 H), 3.61 - 3.43 (m, 1 H), 3.30 - 3.16 (m, 1 H).

Example 302: 1-(2-bromo-4-chlorophenyl)-4-(trifluoromethyl)-1H-pyrazole

2-bromo-4-chloro-1-fluorobenzene (40 mg) was treated with 4-(trifluoromethyl)-1H-pyrazole as detailed in Example 299 to give the title compound having the following physical properties (18 mg).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.07 (t, 1 H), 7.93 (s, 1 H), 7.74 (dd, 1 H), 7.47 (dd, 1 H), 7.44 (dd, 1 H).

Example 303: methyl [4-(2-((1S)-1-[4-{5-chloro-2-[4-(trifluoromethyl)-1H-pyrazol-1-yl]phenyl}-2-oxo-1(2H)-pyridinyl]-2-phenylethyl)-1H-imidazol-4-yl)phenyl]carbamate

The same operation as in Example 283 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 292 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 283 in the operation, the compound prepared in Example 302 was used)

TLC: Rf 0.72 (66% ethyl acetate in hexane)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.29 (s, 1 H), 9.61 (s, 1 H), 8.70 - 8.42 (m, 1 H), 8.14 - 7.95 (m, 1 H), 7.90 - 7.59 (m, 6 H), 7.59 - 7.37 (m, 3 H), 7.33 - 6.98 (m, 5 H), 6.43 - 6.06 (m, 2 H), 5.96 - 5.68 (m, 1 H), 3.66 (s, 3 H), 3.56 - 3.42 (m, 1 H), 3.29 - 3.06 (m, 1 H).

Example 304: 1-(2-bromo-4-chlorophenyl)-1H-1,2,4-triazol-3-ol

2-Bromo-4-chloro-phenylhydrazine [Journal of the American Chemical Society, 79, 934, (1957)] (800 mg) and urea (317 mg) were suspended in xylene (12 mL) and the suspension treated with concentrated sulfuric acid (266 mg) with vigorous stirring. The reaction mixture was heated for 6 hours at 135 °C then allowed to cool to 90 °C, whereupon formic acid (420 mg) and concentrated sulfuric acid (46 mg) was added sequentially. The mixture was heated for 15 hours at 90 °C at which juncture additional

formic acid (1.26 g) and concentrated sulfuric acid (549 mg) were added. The mixture was stirred for a further 3 hours at 90 °C then allowed to cool to room temperature. The resulting precipitate was collected by filtration, washed with water and dried in vacuo to give the title compound having the following physical properties (520 mg).

- 5 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 10.4 (br. s, 1 H), 8.48 (s, 1 H), 8.01 (d, 1 H), 7.66 - 7.58 (m, 2 H).

Example 305: 1-(2-bromo-4-chlorophenyl)-3-[(4-methoxybenzyl)oxy]-1H-1,2,4-triazole

- To a N,N-dimethylformamide (5 mL) solution of the compound prepared in  
10 Example 304 was sequentially added potassium carbonate (452 mg) and 4-methoxybenzylchloride (256 mg) and the mixture stirred for 3 days at room temperature. To the reaction mixture, water was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated and the residue purified by column chromatography (30% ethyl acetate in hexane) to obtain the  
15 title compound having the following physical properties (120 mg).  
TLC: R<sub>f</sub> 0.87 (66% ethyl acetate in hexane).

Example 306: methyl [4-(2-{(3S)-7-[5-chloro-2-(3-hydroxy-1H-1,2,4-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -1H-imidazol-5-yl)phenyl]carbamate

- 20 The same operation as in Example 283 → Example 8 → Example 38 → Example 39 → Example 40 was conducted from the compound prepared in Example 305 to give the title compound having the following physical properties.  
TLC: R<sub>f</sub> 0.43 (10% methanol in dichloromethane)  
<sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>) δ 8.12 (s, 1 H), 7.73 - 7.50 (m, 5 H), 7.43 (d, 2 H),  
25 7.23 (s, 1 H), 6.26 (s, 1 H), 6.17 (s, 1 H), 5.80 (dd, 1 H), 3.82 - 3.69 (m, 3 H), 3.59 - 3.39 (m, 1 H), 3.24 - 3.03 (m, 1 H), 2.73 - 2.58 (m, 1 H), 2.56 - 2.45 (m, 1 H).

Example 307: ethyl (3S)-7-(2-amino-5-methyl-3-pyridinyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylate

- 30 To a 1,4-dioxane (20 mL) solution of 2-amino-3-bromo-5-methylpyridine (1.0 g) was added bis(pinacolate)diboron (1.63 g) and potassium acetate (788 mg) and the mixture degassed with argon. 1,1'-Bis(diphenylphosphino)ferrocene palladium (II) dichloride dichloromethane complex (218 mg) was then added and the reaction mixture

stirred for 3 hours at 110 °C before allowing to cool to room temperature. The reaction mixture was diluted with ethyl acetate, filtered and the filtrate washed with brine, dried and concentrated. The resulting residue was treated with the compound prepared in

5 following physical properties (290 mg).

LC/MS  $t_R$  3.32 minutes; MS (ES<sup>+</sup>)  $m/z$  314 (M+H)<sup>d</sup>.

Example 308: methyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)-3-pyridinyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]carbamate

10 The same operation as in Example 8 → Example 38 → Example 39 → Example 41 was conducted from the compound prepared in Example 307 to give the title compound having the following physical properties.

TLC: R<sub>f</sub> 0.38 (5% methanol in ethyl acetate, NH Silica)

15 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.11 - 11.96 (m, 1 H), 9.95 - 9.80 (m, 1 H), 9.58 (s, 1 H), 8.58 (d, 1 H), 8.07 (d, 1 H), 7.63 - 7.51 (m, 2 H), 7.49 - 7.37 (m, 3 H), 6.18 - 5.93 (m, 2 H), 5.63 (d, 1 H), 3.76 - 3.56 (m, 3 H), 3.49 - 3.30 (m, 1 H), 3.20 - 2.90 (m, 1 H), 2.65 - 2.54 (m, 1 H), 2.47 (br. s, 3 H), 2.40 - 2.27 (m, 1 H).

20 Example 309: methyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]carbamate

The same operation as in Example 7 → Example 8 → Example 38 → Example 39 → Example 41 was conducted from the compound prepared in Example 6 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 7 in the operation, 2-amino-5-methylphenylboronic acid

25 pinacol ester was used)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (s, 1 H), 9.65 (s, 1 H), 9.58 (s, 1 H), 7.62 - 7.54 (m, 3 H), 7.50 (d, 3 H), 7.45 - 7.37 (m, 2 H), 5.93 (s, 1 H), 5.86 (s, 1 H), 5.60 (d, 1 H), 3.65 (s, 3 H), 3.45 - 3.35 (m, 1 H), 3.07 - 2.89 (m, 1 H), 2.46 (s, 3 H), 2.40 - 2.23 (m, 2 H).

30

Example 310: methyl [4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]carbamate

The same operation as in Example 283 → Example 8 → Example 38 →

Example 39 was conducted from the compound prepared in Example 282 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 283 in the operation, the compound prepared in Example 297 was used)

- 5 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (s, 1 H), 9.57 (s, 1 H), 8.79 - 8.58 (m, 1 H), 7.83 - 7.65 (m, 3 H), 7.65 - 7.30 (m, 5 H), 6.04 - 5.81 (m, 2 H), 5.61 (dd, 1 H), 3.74 - 3.59 (m, 3 H), 3.49 - 3.36 (m, 1 H), 3.01 (dd, 1 H), 2.55 (br. s, 1 H), 2.43 - 2.13 (m, 1 H).

Example 311: 1-(2-bromo-4-fluorophenyl)-1H-tetrazole

- 10 To a glacial acetic acid (10 mL) solution of 2-bromo-4-fluoroaniline (570 mg) was added trimethyl orthoformate (1 mL) followed after 30 minutes by sodium azide (5.93 g) and the mixture stirred for 7 hours at 90 °C. The reaction mixture was concentrated and the resulting residue suspended in ethyl acetate. The suspension thus formed was washed sequentially with water, a saturated aqueous solution of sodium  
15 hydrogen carbonate and brine, dried and concentrated. The residue thus obtained was purified by column chromatography (60% ethyl acetate in hexane) to obtained the title compound having the following physical properties (700 mg).  
TLC R<sub>f</sub> 0.26 (25% ethyl acetate in hexane).

- 20 Example 312: methyl [4-(2-((3S)-7-[5-fluoro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl]phenyl]carbamate

- The same operation as in Example 283 → Example 8 → Example 38 → Example 39 was conducted from the compound prepared in Example 282 to give the title compound having the following physical properties. (Note: in the step  
25 corresponding to Example 283 in the operation, the compound prepared in Example 311 was used)

TLC R<sub>f</sub> 0.33 (5% methanol in ethyl acetate, NH Silica)

- <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.04 (s, 1 H), 9.68 (s, 1 H), 9.58 (s, 1 H), 7.87 - 7.78 (m, 1 H), 7.63 - 7.51 (m, 4 H), 7.49 - 7.37 (m, 3 H), 5.94 (app. s, 2 H), 5.60 (d, 1 H),  
30 3.68 - 3.63 (m, 3 H), 3.30 - 3.21 (m, 1 H), 3.05 - 2.92 (m, 1 H), 2.46 - 2.23 (m, 2 H).

Example 313: 1-[2-iodo-4-(trifluoromethyl)phenyl]-1H-tetrazole

2-iodo-4-trifluoromethylaniline (861 mg) was treated as detailed in Example 311

to give the title compound having the following physical properties (873 mg).  
TLC Rf 0.24 (25% ethyl acetate in hexane).

Example 314: methyl [4-(2-((3S)-5-oxo-7-[2-(1H-tetrazol-1-yl)-5-(trifluoromethyl)phenyl]-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 283 → Example 8 → Example 38 → Example 39 was conducted from the compound prepared in Example 282 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 283 in the operation, the compound prepared in Example 313 was used)

TLC Rf 0.55 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.03 (s, 1 H), 9.75 - 9.73 (m, 1 H), 9.58 (s, 1 H), 8.15 - 8.10 (m, 1 H), 8.04 - 8.01 (m, 2 H), 7.66 - 7.35 (m, 5 H), 6.04 - 5.96 (m, 2 H), 5.62 (d, 1 H), 3.67 - 3.64 (m, 3 H), 3.46 - 3.35 (m, 1 H), 3.12 - 2.90 (m, 1 H), 2.46 - 2.28 (m, 2 H).

Example 315: 2-(4-hydroxy-6-methyl-2-oxo-1(2H)-pyridinyl)propanoic acid

A suspension of 4-hydroxy-6-methyl-2-pyrone (2.52 g) and DL-alanine (1.78 g) in 1 M sodium hydroxide (20 mL) was heated at reflux for 15 hours. To the reaction mixture, an aqueous solution of 1 M hydrochloric acid (22 mL) was added and the resulting precipitate collected by filtration. The filter cake was washed with 1 M hydrochloric acid and dried in vacuo to obtain the title compound having the following physical properties (2.3 g).


LC/MS *t<sub>R</sub>* 2.90 minutes; MS (ES<sup>+</sup>) *m/z* 198 (M+H)<sup>d</sup>.

Example 316: ethyl 2-(4-hydroxy-6-methyl-2-oxo-1(2H)-pyridinyl)propanoate

An ethanol (15 mL) suspension of the compound prepared in Example 315 (2.3 g) was treated with concentrated sulphuric acid (43 µL) and the mixture stirred for 15 hours at 90 °C. The forthcoming precipitate was collected by filtration, washed with ethanol and dried to obtain the title compound having the following physical properties (0.78 g).

LC/MS *t<sub>R</sub>* 3.47 minutes; MS (ES<sup>+</sup>) *m/z* 226 (M+H)<sup>d</sup>.

Example 317: methyl {4-[2-(1-{4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}ethyl)-1H-imidazol-5-yl]phenyl}carbamate



5           The same operation as in Example 6 → Example 7 → Example 9 → Example  
11 → Example 38 → Example 39 was conducted from the compound prepared in  
Example 316 to give the title compound having the following physical properties.

TLC Rf 0.58 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 11.01 (br. s, 1 H), 8.69 - 8.37 (m, 1 H), 7.76 - 7.45 (m, 5 H), 7.39 (d, 2 H), 7.18 (s, 1 H), 6.69 (br. s, 1 H), 6.36 (s, 1 H), 5.73 (br. s, 1 H), 5.62 - 5.46 (m, 1 H), 3.85 - 3.70 (m, 3 H), 2.50 (br. s, 3 H), 1.99 (d, 3 H).

Example 318: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

15           The same operation as in Example 315 → Example 316 → Example 6 →  
Example 7 → Example 9 → Example 11 → Example 38 → Example 39 was conducted  
to give the title compound having the following physical properties. (Note: in the step  
corresponding to Example 315 in the operation, glycine was used in place of DL-  
alanine.)

20 TLC Rf 0.42 (5% methanol in ethyl acetate, NH Silica)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.97 (s, 1 H), 9.75 - 9.64 (m, 1 H), 9.58 (s, 1 H), 7.86 - 7.73 (m, 3 H), 7.63 - 7.50 (m, 2 H), 7.49 - 7.36 (m, 3 H), 6.12 - 5.95 (m, 1 H), 5.91 (d, 1 H), 5.25 - 5.08 (m, 2 H), 3.70 - 3.60 (m, 3 H), 2.55 - 2.53 (m, 3 H).

25 Example 319: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-6-ethyl-2-oxo-  
1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl} carbamate

The same operation as in Example 315 → Example 316 → Example 6 → Example 7 → Example 9 → Example 11 → Example 38 → Example 39 was conducted

from 6-ethyl-4-hydroxy-2-pyrone to give the title compound having the following physical properties. (Note: in the step corresponding to Example 315 in the operation, glycine was used in place of DL- alanine.)

TLC Rf 0.50 (ethyl acetate)

- 5 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.93 (s, 1 H), 9.77 - 9.63 (m, 1 H), 9.58 (s, 1 H), 7.85 - 7.80 (m, 3 H), 7.65 - 7.36 (m, 5 H), 6.33 - 6.04 (m, 1 H), 5.72 - 5.49 (m, 1 H), 5.17 (s, 2 H), 3.74 - 3.61 (m, 3 H), 3.00 - 2.75 (m, 2 H), 1.10 - 0.91 (m, 3 H).

Example 320: 4-chloro-2-(6-chloro-2-methyl-4-pyrimidinyl)aniline

- 10 To a 1,4-dioxane (150 mL) solution of 4,6-dichloro-2-methylpyrimidine (2.0 g) stirring under an atmosphere of argon was added 4-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (3.11 g) and 2 M aqueous sodium carbonate solution (18.5 mL). The mixture was degassed with argon then tetrakis(triphenylphosphine) palladium(0) (0.71 g) added and the mixture stirred for 2 hours at 100 °C. To the  
15 reaction mixture, ethyl acetate was added and the organic layer washed with a saturated aqueous solution of sodium hydrogen carbonate and brine, dried and concentrated. The residue thus obtained was triturated with dichloromethane to give the title compound having the following physical properties (1.6 g).  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.54 (d, 1 H), 7.48 (s, 1 H), 7.18 (dd, 1 H), 6.69 (d, 1 H),  
20 6.13 (br. s, 2 H), 2.75 (s, 3 H).

Example 321: 6-(2-amino-5-chlorophenyl)-2-methyl-4(3H)-pyrimidinone

- A suspension of the compound prepared in Example 320 (1.2 g) in 5 M hydrochloric acid (30 mL) was refluxed for 2 hours. The reaction mixture was  
25 concentrated and the residue azeotroped with ethanol and toluene to obtain the title compound having the following physical properties (1.7 g).  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 7.38 (d, 1 H), 7.17 (dd, 1 H), 6.80 (d, 1 H), 6.46 (s, 1 H), 2.37 (s, 3 H), 4.50-3.50 (obs. m, 3 H).

- 30 Example 322: 2-methyl-2-propanyl [4-(2-amino-5-chlorophenyl)-2-methyl-6-oxo-1(6H)-pyrimidinyl]acetate


To an N,N-dimethylformamide (30 mL) solution of the compound prepared in Example 321 (0.69 g) was added *tert*-butyl bromoacetate (0.45 mL) and potassium



carbonate (1.06 g) and the mixture stirred at room temperature for 3 hours. The reaction mixture was then diluted with ethyl acetate, washed with water and brine, dried and concentrated. The residue thus obtained was purified by column chromatography (15 to 50% ethyl acetate in hexane) to give the title compound having the following physical properties (370 mg).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.37 (d, 1 H), 7.12 (dd, 1 H), 6.62 (d, 1 H), 6.60 (s, 1 H), 5.38 (br. s, 2 H), 4.74 (s, 2 H), 2.51 (s, 3 H), 1.50 (s, 9 H).

Example 323: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-methyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate



The same operation as in Example 24 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 322 to give the title compound having the following physical properties.

TLC R<sub>f</sub> 0.58 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.10 (br. s, 1 H), 9.73 (s, 1 H), 9.57 (s, 1 H), 7.98 (s, 1 H), 7.92 - 7.70 (m, 2 H), 7.60 (d, 2 H), 7.55 - 7.26 (m, 3 H), 6.53 (s, 1 H), 5.18 (s, 2 H), 3.65 (s, 3 H), 2.55 - 2.44 (obs. m, 3 H).

Example 324: 2-methoxyethyl [4-(bromoacetyl)phenyl]carbamate

The same operation as in Example 77 → Example 78 was conducted from 4-aminoacetophenone to give the title compound having the following physical properties. (Note: in the step corresponding to Example 77 in the operation, 2-methoxyethyl chloroformate was used)

TLC R<sub>f</sub> 0.40 (50% ethyl acetate in hexane).

Example 325: 2-methoxyethyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2-methyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl} carbamate

The same operation as in Example 24 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 322 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 324 was used)

TLC Rf 0.52 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.10 (s, 1 H), 9.75 - 9.63 (m, 2 H), 8.04 - 7.92 (m, 1 H), 7.92 - 7.69 (m, 2 H), 7.62 - 7.34 (m, 5 H), 6.70 - 6.33 (m, 1 H), 5.18 (s, 2 H), 4.23 - 4.13 (m, 2 H), 3.71 - 3.46 (m, 2 H), 3.28 - 3.26 (m, 3 H), 2.55 - 2.44 (obs. m, 3 H).

Example 326: 2-methoxyethyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-methyl-6-oxo-1(6H)-pyrimidinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The compound prepared in Example 325 (39 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (11 mg).

TLC Rf 0.83 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.84 (s, 1 H), 9.88 (s, 1 H), 9.74 (s, 1 H), 8.01 (d, 1 H), 7.91 - 7.71 (m, 2 H), 7.69 - 7.34 (m, 4 H), 6.51 (s, 1 H), 5.17 (s, 2 H), 4.28 - 4.13 (m, 2 H), 3.67 - 3.49 (m, 2 H), 3.29 (s, 3 H), 2.39 (s, 3 H).

Example 327: 3-{[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]methyl}-6-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-methyl-4(3H)-pyrimidinone

The same operation as in Example 24 → Example 25 → Example 38 → Example 39 was conducted from the compound prepared in Example 322 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 193 was used)

TLC Rf 0.20 (5% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.07 (s, 1 H), 9.73 (s, 1 H), 8.29 - 8.17 (m, 1 H), 7.98 (s, 1 H), 7.92 - 7.73 (m, 2 H), 7.65 (d, 2 H), 7.31 (s, 1 H), 6.52 (s, 1 H), 6.42 (d, 1 H), 5.78 (d, 1 H), 5.17 (s, 2 H), 2.55 - 2.44 (obs. m, 3 H).


Example 328: 3-{[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]methyl}-6-[5-

chloro-2-(1H-tetrazol-1-yl)phenyl]-2-methyl-4(3H)-pyrimidinone

The compound prepared in Example 327 (30 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (7 mg).

TLC Rf 0.62 (5% methanol in ethyl acetate)

- 5 <sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>) δ 9.44 (s, 1 H), 8.18 (d, 1 H), 8.03 - 7.85 (m, 1 H), 7.84 - 7.56 (m, 3 H), 6.64 (dd, 1 H), 6.50 (s, 1 H), 5.25 (s, 2 H), 2.46 (s, 3 H).

Example 329: (6S)-6-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one


10

The same operation as in Example 8 → Example 9 → Example 51 → Example 52 was conducted from the compound prepared in Example 35 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 215 was used).

- 15 LC/MS *t<sub>R</sub>* 2.86 minutes; MS (ES<sup>+</sup>) *m/z* 491 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.22 (br. s, 1 H), 9.73 (s, 1 H), 7.99 (dd, 1 H), 7.96 (d, 1 H), 7.85 (dd, 1 H), 7.82 (d, 1 H), 7.11 (d, 1 H), 6.36 (dd, 1 H), 6.35 (s, 1 H), 6.26 (br. s, 2 H), 5.62 (dd, 1 H), 3.18 (ddd, 1 H), 2.76 (ddd, 1 H), 2.59 – 2.52 (obs. m, 1 H), 2.24 – 2.16 (m, 1 H).

20

Example 330: (6S)-6-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one




The same operation as in Example 8 → Example 9 → Example 51 → Example 52 → Example 44 was conducted from the compound prepared in Example 35 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used).

LC/MS  $t_R$  2.78 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (300 MHz, methanol- $d_4$ )  $\delta$  9.45 (s, 1 H), 8.20 (d, 1 H), 7.92 (d, 1 H), 7.80 – 7.73 (m, 2 H), 7.70 (d, 1 H), 6.65 (d, 1 H), 6.36 (s, 1 H), 5.65 (dd, 1 H), 3.30 – 3.22 (m, 1 H), 2.91 (ddd, 1 H), 2.67 (dq, 1 H), 2.34 (ddt, 1 H).

Example 331: (6S)-6-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one




The same operation as in Example 8 → Example 9 → Example 51 → Example 52 was conducted from the compound prepared in Example 35 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 209 was used).

TLC R<sub>f</sub> 0.40 (10% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.23 (s, 1 H), 9.71 (s, 1 H), 8.02 - 7.91 (m, 2 H), 7.88 - 7.72 (m, 2 H), 7.42 (d, 1 H), 6.55 - 6.24 (m, 4 H), 5.62 (dd, 1 H), 3.25 - 3.07 (m, 1 H), 2.83 - 2.66 (m, 1 H), 2.59 – 2.52 (obs. m, 1 H), 2.31 - 2.13 (m, 1 H).

Example 332: 2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate



5

The same operation as Example 8 → Example 9 → Example 51 → Example 52 → Example 55 → Example 128 → Example 44 was conducted from the compound prepared in Example 35 to obtain the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 219 was used. In the step corresponding to Example 128 in the operation, 2-methoxyethyl chloroformate was used).


10

LC/MS  $t_R$  3.87 minutes; MS (ES<sup>+</sup>)  $m/z$  609 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.09 (br. s, 1 H), 10.12 (br. s, 1 H), 9.73 (s, 1 H), 8.76 (d, 1 H), 8.00 (d, 1 H), 7.96 (dd, 1 H), 7.91 (d, 1 H), 7.86 (dd, 1 H), 7.82 (d, 1 H), 6.33 (s, 1 H), 5.65 (dd, 1 H), 4.25 (dd, 2 H), 3.61 - 3.57 (m, 2 H), 3.30 (s, 3 H), 3.08 (td, 1 H), 2.77 (ddd, 1 H), 2.60 - 2.53 (m, 1 H), 2.18 - 2.08 (m, 1 H).

15

Example 333: ethyl (6S)-2-[(4-methylbenzenesulfonyl)oxy]-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylate



20

To a stirred suspension of ethyl (2S)-5-aminopyrrolidine-2-carboxylate [J. Org. Chem. 52(26), 5717 (1987)] (300 mg), malonic acid monoethyl ester potassium salt (531 mg) and EDC.HCl (598 mg) in N,N-dimethylformamide (3 mL) was added diisopropylethylamine (0.326 mL). The suspension was stirred at 60 °C for 16 hours.

25


The mixture was cooled to 5 °C then treated with diisopropylethylamine (0.326 mL)

and *p*-toluenesulfonyl chloride (357 mg). The mixture was stirred at room temperature for 2 hours. Water was poured into the reaction mixture. The aqueous mixture was extracted with ethyl acetate. The combined organic solution was washed with brine and concentrated under reduced pressure. The residue was purified by column

5 chromatography (70% ethyl acetate in heptanes) to give the title compound having the following physical properties (362 mg)

LC/MS  $t_R$  1.93 minutes; MS ( $ES^+$ )  $m/z$  401 ( $M+Na$ ), 379 ( $M+H$ )<sup>a</sup>.


10 Example 334: ethyl (6S)-2-(2-amino-5-chlorophenyl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylate



To a mixture of the compound prepared in Example 333 (30 g), 4-chloro-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (24.1 g) and triphenylphosphine (2.08 g) in toluene (360 mL), were added sodium carbonate (25.2 g) in water (120 ml) and palladium(II) bis(triphenylphosphosphine) dichloride (2.78 g). The mixture was stirred at 100 °C for 6 hours under an atmosphere of argon. The insoluble materials were removed by filtration. The resulting residue was washed with ethyl acetate (150 mL) and water (150 mL). The organic layer was separated and the aqueous layer was extracted with ethyl acetate (150 mL). The combined organic layers were washed with brine (90 mL), dried and concentrated. The resulting residue was purified by column chromatography (70% ethyl acetate in heptanes) to give the title compound having the following physical properties (19.3 g).


LC/MS  $t_R$  1.81 minutes; MS ( $ES^+$ )  $m/z$  334 ( $M+H$ )<sup>a</sup>.

25 Example 335: (6S)-2-(2-amino-5-chlorophenyl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylic acid




- A suspension of the compound prepared in Example 334 (17.5 g) in 2 M hydrochloric acid (160 mL) and 1,4-dioxane (40 mL) was stirred at 90 °C for 8 hours. The reaction mixture was cooled to room temperature. The resulting precipitate was collected by filtration, washed with 1,4-dioxane and dried to give the title compound (10.4 g) as the hydrochloride salt. The filtrate was cooled on an ice bath and treated with 5 M sodium hydroxide to pH 3-4. The aqueous mixture was saturated with ammonium sulfate and extracted with ethyl acetate / methanol (85/15). The combined organic layers were dried and concentrated to give the title compound (7.2 g) as the free base.
- LC/MS  $t_R$  1.52 minutes; MS ( $ES^+$ )  $m/z$  306 ( $M+H$ )<sup>a</sup>.

Example 336: (6S)-2-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylic acid



- The compound prepared in Example 335 (4.34 g) was treated as detailed in Example 9 to give the title compound having the following physical properties (3.24 g). LC/MS  $t_R$  1.40 minutes; MS ( $ES^+$ )  $m/z$  739 ( $2M+Na$ ), 381 ( $M+Na$ ), 359 ( $M+H$ ), 331 ( $M-N_2+H$ )<sup>a</sup>.

- Example 337: 3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoic acid




The same operation as in Example 51 → Example 52 → Example 37 →

Example 55 → Example 24 was conducted from the compound prepared in Example 11 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, methyl 3-(2-bromoacetyl)benzoate [J. Med. Chem., 13(4), 674 (1970)] was used).

LC/MS  $t_R$  3.08 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 8.32 (app. s, 1 H), 7.93 – 7.84 (m, 2 H), 7.78 - 7.72 (m, 2 H), 7.70 (d, 1 H), 7.47 (t, 1 H), 7.41 (s, 1 H), 6.15 (s, 1 H), 6.11 (s, 1 H), 5.81 (dd, 1 H), 3.48 (td, 1 H), 3.13 (ddd, 1 H), 2.72 - 2.62 (m, 1 H), 2.55 - 2.44 (m, 1 H).

Example 338: 3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)benzoic acid



15

To a cooled (0 °C) tetrahydrofuran (5 mL) solution of the compound prepared in Example 337 (90 mg) was added a solution of 1,3-dichloro-5,5-dimethyl imidazolidine-2,4-dione (25 mg) in tetrahydrofuran (0.5 mL) and the reaction mixture stirred at 0 °C for 0.5 hours. To the reaction mixture, a 20% aqueous solution of sodium thiosulfate (1 mL) was added and the mixture stirred for 5 minutes before diluting with water and extracting into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue obtained was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile

20




phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (18 mg).

LC/MS  $t_R$  3.88 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.40 (app. s, 1 H), 8.04 - 7.92 (m, 2 H), 7.79 - 7.68 (m, 3 H), 7.58 (t, 1 H), 6.18 (s, 1 H), 6.08 (s, 1 H), 5.75 (dd, 1 H), 3.53 - 3.41 (m, 1 H), 3.13 (ddd, 1 H), 2.72 - 2.61 (m, 1 H), 2.46 - 2.34 (m, 1 H).


Example 339: methyl 2-(4-acetylphenyl)acetate



10 Methyl (4-bromophenyl)acetate (0.50 g) was treated as detailed in Example 139 to give the title compound having the following physical properties (0.26 g).

LC/MS  $t_R$  1.60 minutes; MS (ES<sup>+</sup>)  $m/z$  234 (M+H+MeCN), 193 (M+H)<sup>a</sup>.

Example 340: methyl 2-[4-(2-bromoacetyl)phenyl]acetate



15

To a dichloromethane (4 mL) solution of the compound prepared in Example 339 (0.21 g) was added bromine (53  $\mu$ L) and the mixture stirred at room temperature for 3 hours. The reaction mixture was concentrated to give the crude title product having the following physical properties (0.28 g).


20 LC/MS  $t_R$  1.82 minutes; MS (ES<sup>+</sup>)  $m/z$  271 and 273 (M+H)<sup>a</sup>.

Example 341: methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]acetate

25 The same operation as in Example 51  $\rightarrow$  Example 52 was conducted from the compound prepared in Example 340 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.54 minutes; MS (ES<sup>+</sup>)  $m/z$  528 (M+H)<sup>a</sup>.

30 Example 342: [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]acetic acid




To a 1,4-dioxane (2 mL) solution of the compound prepared in Example 341 (90 mg) was added 1 M hydrochloric acid (0.85 mL) and the mixture warmed at 90 °C for 3 hours. The solvents were removed *in vacuo* and the residue purified by high

5 performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (21 mg).

LC/MS  $t_R$  2.96 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.39 (br. s, 1 H), 9.70 (s, 1 H), 8.05 (br. s, 1 H), 7.86  
10 - 7.80 (m, 2 H), 7.77 - 7.69 (m, 3 H), 7.41 (d, 2 H), 6.07 (s, 1 H), 6.02 (s, 1 H), 5.81 (dd, 1 H), 3.64 (s, 2 H), 3.27 - 3.24 (m, 1 H), 3.14 - 3.05 (m, 1 H), 2.75 - 2.65 (m, 1 H), 2.38 - 2.30 (m, 1 H).

15 Example 343: [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizine-1-yl)phenyl]acetic acid



The same operation as in Example 338 → Example 342 was conducted from the compound prepared in Example 341 to give the title compound having the following physical properties.

20 LC/MS  $t_R$  3.83 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.99 (br. s, 1 H), 9.71 (s, 1 H), 7.88 - 7.78 (m, 3 H), 7.63 (d, 2 H), 7.35 (d, 2 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.58 (dd, 1 H), 3.59 (s, 2 H), 3.40 - 3.20 (obs. m, 1 H), 2.99 (dd, 1 H), 2.60 - 2.50 (obs. m, 1 H), 2.25 - 2.15 (m, 1 H).

Example 344: methyl 2-(3-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}phenyl)acetate

The same operation as in Example 139 → Example 340 → Example 51 → Example 52 was conducted from methyl (3-bromophenyl)acetate to give the title compound having the following physical properties.

LC/MS  $t_R$  1.57 minutes; MS (ES<sup>+</sup>)  $m/z$  528 (M+H)<sup>a</sup>.


Example 345: [3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl}phenyl)acetic acid



The compound prepared in Example 344 (270 mg) was treated using the method as detailed in Example 342 to give the title compound having the following physical properties (260 mg).

LC/MS  $t_R$  3.00 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.34 (br. s, 1 H), 12.15 (br. s, 1 H), 9.70 (s, 1 H), 7.87 - 7.74 (m, 3 H), 7.62 - 7.54 (m, 2 H), 7.51 (br. s, 1 H), 7.30 (m, 1 H), 7.10 (m, 1 H), 5.99 (s, 1 H), 5.96 (s, 1 H), 5.65 (d, 1 H), 3.40 - 3.30 (obs. m, 1 H), 3.59 (s, 2 H), 3.02 (dd, 1 H), 2.55 - 2.45 (obs. m, 1 H), 2.35 (m, 1 H).

Example 346: [3-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl}phenyl)acetic acid




The same operation as in Example 338 → Example 342 was conducted from the compound prepared in Example 344 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.84 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.02 (br. s, 1 H), 12.42 (br. s, 1 H), 9.71 (s, 1 H), 7.89 - 7.75 (m, 3 H), 7.65 - 7.56 (m, 2 H), 7.42 (t, 1 H), 7.23 (d, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.58 (dd, 1 H), 3.62 (s, 2 H), 3.30 - 3.23 (m, 1 H), 2.99 (dd, 1 H), 2.58 - 2.53 (m, 1 H), 2.20 (app. t, 1 H).

- 10 Example 347: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid




The same operation as in Example 234 → Example 51 → Example 52 → Example 8 → Example 55 → Example 24 was conducted from 5-

- 15 (methoxycarbonyl)thiophene- 2-carboxylic acid to give the title compound having the following physical properties.(Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 11 was used).  
LC/MS  $t_R$  3.38 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.73 (m, 2 H), 7.70 - 7.65 (m, 2 H),  
20 7.41 (s, 1 H), 7.27 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.50 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.49 (tdd, 1 H).

Example 348: 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid

25




The compound prepared in Example 347 (95 mg) was treated using the method as detailed in Example 338 to give the title compound having the following physical properties (17 mg).

LC/MS  $t_R$  3.81 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.65 (m, 4 H), 7.38 (d, 1 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.70 (dd, 1 H), 3.50 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.66 (qd, 1 H), 2.43 - 2.33 (m, 1 H).

Example 349(1) to Example 349(4)



10

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 347 using the corresponding amine hydrochloride salts, employing the method as detailed in Example 114.

- 15 Example 349(1): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy)-1H-imidazol-5-yl)-2-thiophenecarboxamide

LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  505 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.77 - 7.67 (m, 3 H), 7.62 (d, 1 H), 7.39 (s, 1 H), 7.26 (d, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.77 (dd, 1 H), 3.46 (td, 1 H),  
20 3.12 (ddd, 1 H), 2.72 - 2.58 (m, 1 H), 2.54 - 2.44 (m, 1 H).

Example 349(2): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy)-1H-imidazol-5-yl)-N-methyl-2-thiophenecarboxamide

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  519 (M+H)<sup>b</sup>

- 25 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.78 - 7.64 (m, 3 H), 7.53 (d, 1 H), 7.37 (s, 1 H), 7.23 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.75 (dd, 1 H), 3.44 (td, 1 H), 3.11 (ddd, 1 H), 2.89 (s, 3 H), 2.68 - 2.57 (m, 1 H), 2.53 - 2.41 (m, 1 H).

Example 349(3): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy)-1H-imidazol-5-yl)-N-ethyl-2-thiophenecarboxamide

30

LC/MS  $t_R$  3.40 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.78 – 7.63 (m, 3 H), 7.57 (d, 1 H), 7.41 (s, 1 H), 7.25 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.49 – 3.33 (m, 3 H), 3.12 (ddd, 1 H), 2.72 – 2.59 (m, 1 H), 2.46 (tt, 1 H), 1.21 (t, 3 H).

5


Example 349(4): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide

LC/MS  $t_R$  3.36 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 – 7.64 (m, 3 H), 7.42 (app. s, 2 H), 7.26 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.41 (dd, 1 H), 3.21 (br. s, 6 H), 3.11 (ddd, 1 H), 2.74 – 2.60 (m, 1 H), 2.52 – 2.40 (m, 1 H).

10

Example 350(1) to Example 350(3)



15 The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 347 using the corresponding amines, employing the method as detailed in Example 70.

Example 350(1): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(2-methoxyethyl)-2-thiophenecarboxamide

20

LC/MS  $t_R$  3.29 minutes; MS (ES<sup>+</sup>)  $m/z$  563 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.58 (d, 1 H), 7.37 (br. s, 1 H), 7.23 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.75 (dd, 1 H), 3.56 - 3.50 (m, 4 H), 3.49 - 3.40 (m, 1 H), 3.37 (s, 3 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.48 (app. br. s, 1 H).

25

Example 350(2): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-[2-(2-methoxyethoxy)ethyl]-2-thiophenecarboxamide

30


LC/MS  $t_R$  3.34 minutes; MS (ES<sup>+</sup>)  $m/z$  607 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.60 (d, 1 H), 7.37 (br. s, 1 H), 7.25 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.76 (dd, 1 H), 3.66 - 3.62 (m, 4 H), 3.58 - 3.52 (m, 4 H), 3.49 - 3.40 (m, 1 H), 3.37 (s, 3 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.53 - 2.44 (m, 1 H).

Example 350(3): 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-N-[2-(4-morpholinyl)ethyl]-2-thiophenecarboxamide

LC/MS  $t_R$  2.70 minutes; MS (ES<sup>+</sup>)  $m/z$  618 (M+H), 309.5 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.56 (d, 1 H), 7.39 (br. s, 1 H), 7.24 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.75 (dd, 1 H), 3.70 (app. t, 4 H), 3.51 (t, 2 H), 3.48 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.69 - 2.43 (m, 8 H).

Example 351(1) to Example 351(4)



The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 348 using the corresponding amine hydrochloride salts, employing the method as detailed in Example 114.

Example 351(1): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-thiophenecarboxamide

LC/MS  $t_R$  3.57 minutes; MS (ES<sup>+</sup>)  $m/z$  539 and 541 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.31 (br. s, 1 H), 9.71 (s, 1 H), 8.00 (s, 1 H), 7.89 - 7.76 (m, 3 H), 7.72 (d, 1 H), 7.54 - 7.19 (m, 2 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.54 (d, 1 H), 3.33 - 3.21 (obs. m, 1 H), 2.98 (dd, 1 H), 2.60 - 2.52 (obs. m, 1 H), 2.25 - 2.12 (m, 1 H).

Example 351(2): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-methyl-2-thiophenecarboxamide

LC/MS  $t_R$  3.71 minutes; MS (ES<sup>+</sup>)  $m/z$  553 and 555 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.29 (br. s, 1 H), 9.71 (s, 1 H), 8.50 (app. s, 1 H),  
 5 7.88 - 7.76 (m, 3 H), 7.68 (d, 1 H), 7.39 (d, 1 H), 5.99 (s, 1 H), 5.93 (s, 1 H), 5.54 (d, 1 H), 3.31 - 3.20 (obs. m, 1 H), 2.99 (dd, 1 H), 2.76 (d, 3 H), 2.59 - 2.52 (obs. m, 1 H), 2.27 - 2.14 (m, 1 H).

Example 351(3): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-ethyl-2-thiophenecarboxamide

10 LC/MS  $t_R$  3.87 minutes; MS (ES<sup>+</sup>)  $m/z$  567 and 569 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.28 (br. s, 1 H), 9.71 (s, 1 H), 8.52 (app. s, 1 H),  
 7.90 - 7.75 (m, 3 H), 7.70 (d, 1 H), 7.39 (d, 1 H), 5.99 (s, 1 H), 5.93 (s, 1 H), 5.54 (dd, 1 H), 3.31 - 3.19 (m, 3 H), 2.98 (dd, 1 H), 2.60 - 2.50 (obs. m, 1 H), 2.23 - 2.13 (m, 1 H),  
 15 1.12 (t, 3 H).

Example 351(4): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide

20 LC/MS  $t_R$  3.87 minutes; MS (ES<sup>+</sup>)  $m/z$  567 and 569 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.32 (br. s, 1 H), 9.71 (s, 1 H), 7.93 - 7.70 (m, 3 H),  
 7.53 (d, 1 H), 7.40 (d, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.55 (dd, 1 H), 3.35 - 3.22 (obs. m, 1 H), 3.12 (br. s, 6 H), 2.99 (dd, 1 H), 2.59 - 2.50 (obs. m, 1 H), 2.25 - 2.12 (m, 1 H).

25 Example 352(1) to Example 352(3)



The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 350(1), 350(2) and 350(3) using the method as detailed in Example 338.



Example 352(1): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-methoxyethyl)-2-thiophenecarboxamide

LC/MS  $t_R$  3.80 minutes; MS ( $ES^+$ )  $m/z$  597 and 599 ( $M+H$ )<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.97 (br. s, 1 H), 8.70 (br. s, 1 H), 7.63 (dd, 1 H), 7.58 (d, 1 H), 7.55 (d, 1 H), 7.35 (d, 1 H), 6.99 (br. s, 1 H), 6.57 (app. br. s, 1 H), 6.32 (s, 1 H), 5.84 (s, 1 H), 5.81 (d, 1 H), 3.67 - 3.47 (m, 5 H), 3.40 (s, 3 H), 3.04 (dd, 1 H), 2.95 - 2.86 (m, 1 H), 2.59 - 2.46 (m, 1 H).

10 Example 352(2): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-[2-(2-methoxyethoxy)ethyl]-2-thiophenecarboxamide

LC/MS  $t_R$  3.76 minutes; MS ( $ES^+$ )  $m/z$  641 and 643 ( $M+H$ )<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.77 - 7.72 (m, 2 H), 7.71 - 7.65 (m, 2 H), 7.37 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.71 (dd, 1 H), 3.67 - 3.62 (m, 4 H), 3.59 - 3.54 (m, 4 H), 3.49 - 3.40 (m, 1 H), 3.38 (s, 3 H), 3.12 (ddd, 1 H), 2.72 - 2.60 (m, 1 H), 2.42 - 2.32 (m, 1 H).


20 Example 352(3): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-[2-(4-morpholinyl)ethyl]-2-thiophenecarboxamide

LC/MS  $t_R$  2.96 minutes; MS ( $ES^+$ )  $m/z$  652 and 654 ( $M+H$ ), 326.5 and 327.5

$[(M+2H)/2]$ <sup>b</sup>

- 25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.77 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.64 (d, 1 H), 7.36 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.70 (dd, 1 H), 3.76 - 3.66 (m, 4 H), 3.53 (t, 2 H), 3.43 (td, 1 H), 3.11 (ddd, 1 H), 2.65 (qd, 1 H), 2.59 (t, 2 H), 2.55 (app. br. s, 4 H), 2.41 - 2.34 (m, 1 H).

30 Example 353: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-methoxy-2-thiophenecarboxamide




To an N,N-dimethylformamide (6 mL) solution of the compound prepared in Example 347 (150 mg) was added 1-[3-(dimethylaminopropyl)]-3-ethylcarbodiimide hydrochloride (170 mg), 4-dimethylaminopyridine (18 mg) and methoxamide hydrochloride (136 mg). Triethylamine (270  $\mu$ L) was added and the mixture stirred at room temperature for 16 hours. A saturated aqueous solution of sodium hydrogen carbonate was added followed by extraction with ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 to 15% methanol in dichloromethane, NH silica) to give the title compound having the following physical properties (21.6 mg).

LC/MS  $t_R$  3.23 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.76 - 7.70 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.51 (br. s, 1 H), 7.38 (br. s, 1 H), 7.24 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.75 (dd, 1 H), 3.79 (s, 3 H), 3.49 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.53 - 2.42 (m, 1 H).

Example 354: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-N-cyano-2-thiophenecarboxamide




To an N,N-dimethylformamide (4 mL) solution of the compound prepared in Example 347 (100 mg) was added (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (105 mg), cyanamide (12.5 mg) and diisopropylethylamine (98  $\mu$ L) and the mixture stirred at room temperature for 16 hours. Water was added followed by extraction with a 2: 1 mixture of dichloromethane and propan-2-ol. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column

chromatography (0 to 40% methanol in dichloromethane) followed by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (32 mg).

LC/MS  $t_R$  3.24 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>


- 5 <sup>1</sup>H NMR (500 MHz, 2:1 methanol-d<sub>4</sub> and D<sub>2</sub>O)  $\delta$  9.38 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.53 (d, 1 H), 7.32 (br. s, 1 H), 7.19 (d, 1 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.77 (d, 1 H), 3.47 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.71 - 2.59 (m, 1 H), 2.50 - 2.39 (m, 1 H).

- 10 Example 355: 2-(4-morpholinyl)ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate



- To a cooled (0 °C) dichloromethane (15 mL) suspension of the compound prepared in Example 347 (150 mg) was added oxalyl chloride (75.3  $\mu$ L) followed by one drop of N,N-dimethylformamide and the mixture stirred at room temperature for 3 hours. The reaction mixture was concentrated and the residue re-dissolved in dichloromethane (15 mL). To this cooled (0 °C) solution, N-(2-hydroxyethyl)morpholine (108  $\mu$ L) and triethylamine (124  $\mu$ L) were added and the mixture stirred at room temperature for 1 hour. Water was added followed by extraction with dichloromethane. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 to 5% methanol in ethyl acetate) to give the title compound having the following physical properties (60 mg).

- 25 LC/MS  $t_R$  1.72 minutes; MS (ES<sup>+</sup>)  $m/z$  619 (M+H), 310 [(M+2H)/2]<sup>c</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.78 - 7.71 (m, 3 H), 7.70 - 7.65 (m, 1 H), 7.45 (br. s, 1 H), 7.28 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.75 (dd, 1 H), 4.43 (t, 2 H), 3.72 - 3.68 (m, 4 H), 3.50 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.77 (t, 2 H), 2.69 - 2.61 (m, 1 H), 2.62 - 2.56 (m, 4 H), 2.50 (app. br. s, 1 H).

Example 356(1) to Example 356(3)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 353, 354, and 355 using the method as detailed in Example 338.


Example 356(1): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy)-1H-imidazol-5-yl)-N-methoxy-2-thiophenecarboxamide

- LC/MS  $t_R$  2.68 minutes; MS (ES<sup>+</sup>)  $m/z$  569 and 571 (M+H)<sup>+</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.78 - 7.73 (m, 2 H), 7.70 - 7.64 (m, 1 H), 7.59 (app. br. s, 1 H), 7.37 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.70 (dd, 1 H), 3.81 (s, 3 H), 3.44 (td, 1 H), 3.16 - 3.06 (m, 1 H), 2.72 - 2.59 (m, 1 H), 2.41 - 2.32 (m, 1 H).

- Example 356(2): 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy)-1H-imidazol-5-yl)-N-cyano-2-thiophenecarboxamide


- LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  564 and 566 (M+H)<sup>+</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.68 (d, 1 H), 7.59 (d, 1 H), 7.32 (d, 1 H), 6.13 (s, 1 H), 6.08 (s, 1 H), 5.69 (dd, 1 H), 3.42 (td, 1 H), 3.10 (ddd, 1 H), 2.71 - 2.60 (m, 1 H), 2.41 - 2.33 (m, 1 H).

Example 356(3): 2-(4-morpholinyl)ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy)-1H-imidazol-5-yl)-2-thiophenecarboxylate



LC/MS  $t_R$  3.11 minutes; MS ( $ES^+$ )  $m/z$  653 and 655 ( $M+H$ ), 327 and 328 [ $(M+2H)/2$ ]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.80 (d, 1 H), 7.75 - 7.72 (m, 2 H),  
 7.70 - 7.66 (m, 1 H), 7.39 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.70 (dd, 1 H), 4.47 (t, 2  
 H), 3.73 - 3.69 (m, 4 H), 3.49 - 3.39 (m, 1 H), 3.12 (ddd, 1 H), 2.82 (t, 2 H), 2.69 - 2.61  
 5 (m, 5 H), 2.42 - 2.33 (m, 1 H).


Example 357: methyl [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]-2-thienyl]acetate



10 The same operation as in Example 51  $\rightarrow$  Example 52 was conducted from methyl 2-[5-(2-chloroacetyl)thiophen-2-yl]acetate to give the title compound having the following physical properties.

LC/MS  $t_R$  3.46 minutes; MS ( $ES^+$ )  $m/z$  534 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1  
 15 H), 7.19 (s, 1 H), 7.08 (d, 1 H), 6.85 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H),  
 3.86 (s, 2 H), 3.72 (s, 3 H), 3.43 (td, 1 H), 3.10 (ddd, 1 H), 2.64 (qd, 1 H), 2.46 (tdd, 1  
 H).

Example 358: [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]-2-thienyl]acetic acid hydrochloride




The compound prepared in Example 357 (18.2 mg) was treated as detailed in Example 342 to give, on trituration with a 9: 1 mixture of dichloromethane and acetonitrile, the title compound as the hydrochloride salt having the following physical  
 25 properties (14.5 mg).

LC/MS  $t_R$  3.08 minutes; MS ( $ES^+$ )  $m/z$  520 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.68 (br. s, 1 H), 9.70 (s, 1 H), 7.94 - 7.55 (m, 4 H), 7.29 (br. s, 1 H), 6.96 (app. s, 1 H), 6.04 (s, 1 H), 6.01 (s, 1 H), 5.72 (dd, 1 H), 3.87 (s, 2 H), 3.31 - 3.24 (obs. m, 1 H), 3.11 - 3.01 (m, 1 H), 2.70 - 2.58 (m, 1 H), 2.37 - 2.26 (m, 1 H).

5

Example 359: methyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-thienyl]acetate




The compound prepared in Example 357 (69 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (35 mg).

LC/MS *t<sub>R</sub>* 4.18 minutes; MS (ES<sup>+</sup>) *m/z* 568 and 570 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 11.37 (br. s, 1 H), 8.59 (s, 1 H), 7.64 (dd, 1 H), 7.59 - 7.53 (m, 2 H), 7.09 (d, 1 H), 6.90 (d, 1 H), 6.36 (s, 1 H), 5.81 (d, 1 H), 5.76 (s, 1 H), 3.85 (s, 2 H), 3.76 (s, 3 H), 3.50 - 3.38 (m, 1 H), 3.17 (dd, 1 H), 3.02 (dd, 1 H), 2.54 - 2.41 (m, 1 H).

15

Example 360: [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-thienyl]acetic acid hydrochloride



20

The compound prepared in Example 359 (20 mg) was treated as detailed in Example 342 to give, on trituration with a 9: 1 mixture of dichloromethane and acetonitrile, the title compound as the hydrochloride salt having the following physical properties (11.5 mg).


LC/MS *t<sub>R</sub>* 3.08 minutes; MS (ES<sup>+</sup>) *m/z* 554 and 556 (M+H)<sup>b</sup>

25

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.11 (br. s, 1 H), 12.61 (s, 1 H), 9.71 (s, 1 H), 7.86 - 7.76 (m, 3 H), 7.25 (d, 1 H), 6.96 (d, 1 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.54 (dd, 1 H), 3.85 (s, 2 H), 3.33 - 3.22 (m, 1 H), 2.98 (dd, 1 H), 2.58 - 2.49 (obs. m, 1 H), 2.24 - 2.11 (m, 1 H).

5

Example 361: tert-butyl 4-(2-bromoacetyl)thiophene-2-carboxylate



To a tetrahydrofuran (100 mL) solution of tert-butyl 4-acetylthiophene-2-carboxylate [patent EP1357123, 2003] (7.91 g) was added a solution of  
 10 phenyltrimethylammonium tribromide (12.5 g) in tetrahydrofuran (20 mL) and the mixture stirred at room temperature for 3 hours. To the reaction mixture, water (100 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 – 20% ethyl acetate in heptanes) to give the title product having the  
 15 following physical properties (7.73 g).

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.29 (d, 1 H), 8.11 (d, 1 H), 4.31 (s, 2 H), 1.60 (s, 9 H).

Example 362: 2-methyl-2-propanyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate


20 The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 361 to give the title compound having the following physical properties.

LC/MS *t*<sub>R</sub> 1.92 minutes; MS (ES<sup>+</sup>) *m/z* 562 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.95 (s, 1 H), 7.77 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.34 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.47 (app. br. s, 1 H), 1.59 (s, 9 H).

Example 363: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride

30



To a 1,4-dioxane (60 mL) solution of the compound prepared in Example 362 (3.0 g) was added 1 M hydrochloric acid (26.7 mL) and the mixture heated at 90 °C for 5 hours. On cooling the solvent was reduced in volume *in vacuo* to give a precipitate which was isolated by filtration and washed with water. Drying of the filter cake *in vacuo* gave the title compound having the following physical properties (2.28 g).

LC/MS  $t_R$  1.47 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  14.63 (br. s, 1 H), 13.42 (br. s, 1 H), 9.70 (s, 1 H), 8.25 (br. s, 1 H), 8.17 (s, 1 H), 8.01 (br. s, 1 H), 7.88 - 7.77 (m, 2 H), 7.72 (d, 1 H), 6.05 (s, 1 H), 6.02 (s, 1 H), 5.77 (dd, 1 H), 3.32 - 3.28 (m, 1 H), 3.15 - 3.04 (m, 1 H), 2.76 - 2.60 (m, 1 H), 2.39 - 2.30 (m, 1 H).

Example 364: tert-butyl 4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-fluoro-1H-imidazol-5-yl}thiophene-2-carboxylate


To an N,N-dimethylformamide (2 mL) solution of the compound prepared in Example 362 (100 mg) was added sodium carbonate (24.5 mg) and Accufluor® (126 mg) and the suspension stirred at room temperature for 1 hour. Water (10 mL) was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (25 – 100% ethyl acetate in heptanes) to give the title product having the following physical properties (29 mg).

LC/MS  $t_R$  2.31 minutes; MS (ES<sup>+</sup>)  $m/z$  602 (M+Na), 524 [M-C(CH<sub>3</sub>)<sub>3</sub>+H]<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.86 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.64 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.66 (dd, 1 H), 3.48 - 3.38 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.40 - 2.33 (m, 1 H), 1.59 (s, 9 H).



Example 365: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride




5 The compound prepared in Example 364 (26 mg) was treated as detailed in Example 363 to give, on concentration of the reaction mixture, the title compound as the hydrochloride salt having the following physical properties (25 mg).

LC/MS  $t_R$  3.79 minutes; MS ( $ES^+$ )  $m/z$  524 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.12 (very br. s, 1 H), 12.97 (br. s, 1 H), 9.70 (s, 1 H), 7.93 (s, 1 H), 7.84 (d, 1 H), 7.81 - 7.77 (m, 3 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.51 (d, 1 H), 3.33 - 3.23 (m, 1 H), 2.98 (dd, 1 H), 2.58 - 2.54 (m, 1 H), 2.22 - 2.13 (m, 1 H).

Example 366: 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride




The same operation as in Example 338 → Example 363 was conducted from the compound prepared in Example 362 to give the title compound having the following physical properties.

20 LC/MS  $t_R$  3.84 minutes; MS ( $ES^+$ )  $m/z$  540 and 542 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.24 (br. s, 1 H), 9.71 (s, 1 H), 8.14 (d, 1 H), 8.08 (d, 1 H), 7.83 - 7.79 (m, 3 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.54 (dd, 1 H), 3.32 - 3.24 (m, 1 H), 2.99 (dd, 1 H), 2.59 - 2.53 (m, 1 H), 2.19 (t, 1 H).

Example 367: ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate




To an N,N-dimethylformamide (2 mL) suspension of the compound prepared in  
 5 Example 363 (80 mg) was added ethanol (92  $\mu$ L), (1H-benzotriazol-1-yloxy)  
 [tris(dimethylamino)]phosphonium hexafluorophosphate (77 mg) and N,N-  
 diisopropylethylamine (68  $\mu$ L) and the mixture stirred at room temperature for 16 hours.  
 Water (10 mL) was added followed by extraction into ethyl acetate. The combined  
 organic layers were washed with brine, dried and concentrated. The residue was purified  
 10 by column chromatography (50 – 100% ethyl acetate in heptanes, then 0 – 5% methanol  
 in ethyl acetate) to give the title product having the following physical properties (59  
 mg).

LC/MS  $t_R$  3.64 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.04 (d, 1 H), 7.82 (d, 1 H), 7.77 -  
 15 7.72 (m, 2 H), 7.71 - 7.66 (m, 1 H), 7.33 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76  
 (dd, 1 H), 4.34 (q, 2 H), 3.44 (td, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.46 (app. br. s,  
 1 H), 1.37 (t, 3 H).

Example 368(1) to Example 368(7)



20

The compounds of the present invention having the following physical data were  
 synthesised from the compound prepared in Example 363 using the corresponding  
 commercially available alcohols, employing the method as detailed in Example 367.

Example 368(1): isopropyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate

Note: in this instance, the reaction was heated at 50 °C for an additional 24 hours to achieve sufficient conversion to the ester.

- 5 LC/MS  $t_R$  3.84 minutes; MS ( $ES^+$ )  $m/z$  548 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.02 (d, 1 H), 7.81 (d, 1 H), 7.76 - 7.72 (m, 2 H), 7.70 - 7.67 (m, 1 H), 7.32 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 5.19 (td, 1 H), 3.50 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.52 - 2.39 (m, 1 H), 1.37 (d, 6 H).

10


Example 368(2): 3-methylbutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate

LC/MS  $t_R$  4.32 minutes; MS ( $ES^+$ )  $m/z$  576 ( $M+H$ )<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.03 (d, 1 H), 7.83 (d, 1 H), 7.77 - 7.72 (m, 2 H), 7.72 - 7.65 (m, 1 H), 7.33 (br. s, 1 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 4.35 (t, 2 H), 3.52 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.65 (qd, 1 H), 2.47 (app. br. s, 1 H), 1.85 - 1.72 (m, 1 H), 1.66 (app. q, 2 H), 0.99 (d, 6 H).

Example 368(3): 2-(4-morpholinyl)ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate


20



LC/MS  $t_R$  2.73 minutes; MS ( $ES^+$ )  $m/z$  619 ( $M+H$ ), 310 [ $(M+2H)/2$ ]<sup>b</sup>

- 25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.07 (d, 1 H), 7.84 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.35 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 4.45 (t, 2 H), 3.72 - 3.67 (m, 4 H), 3.49 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.78 (t, 2 H), 2.68 - 2.54 (m, 5 H), 2.46 (app. br. s, 1 H).

Example 368(4): (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate



- 5 LC/MS  $t_R$  3.68 minutes; MS ( $ES^+$ )  $m/z$  618 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 8.10 (d, 1 H), 7.88 (d, 1 H), 7.79 - 7.65 (m, 3 H), 7.34 (br. s, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.77 (dd, 1 H), 5.15 (s, 2 H), 3.54 - 3.37 (m, 1 H), 3.10 (dd, 1 H), 2.72 - 2.55 (m, 1 H), 2.54 - 2.31 (m, 1 H), 2.24 (s, 3 H).

10

Example 368(5): phenyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate

LC/MS  $t_R$  4.10 minutes; MS ( $ES^+$ )  $m/z$  582 ( $M+H$ )<sup>b</sup>


- 15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.24 (d, 1 H), 7.96 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.71 - 7.65 (m, 1 H), 7.49 - 7.42 (m, 2 H), 7.39 (br. s, 1 H), 7.32 - 7.27 (m, 1 H), 7.24 (d, 2 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.46 (td, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.48 (app. br. s, 1 H).

20 Example 368(6): 2,3-dihydro-1H-inden-5-yl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate

LC/MS  $t_R$  4.55 minutes; MS ( $ES^+$ )  $m/z$  622 ( $M+H$ )<sup>b</sup>

- 25 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.17 (br. s, 1 H), 9.69 (s, 1 H), 8.23 (s, 1 H), 7.98 (s, 1 H), 7.84 - 7.75 (m, 3 H), 7.58 (br. s, 1 H), 7.28 (d, 1 H), 7.13 (s, 1 H), 7.01 (d, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.62 (d, 1 H), 3.45 - 3.34 (obs. m, 1 H), 3.00 (dd, 1 H), 2.88 (q, 4 H), 2.55 - 2.44 (obs. m, 1 H), 2.38 - 2.28 (m, 1 H), 2.06 (quintet, 2 H).


Example 368(7): isobutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate



LC/MS  $t_R$  4.06 minutes; MS (ES<sup>+</sup>)  $m/z$  562 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.04 (d, 1 H), 7.83 (s, 1 H), 7.77 - 7.72 (m, 2 H), 7.71 - 7.65 (m, 1 H), 7.36 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 4.09 (d, 2 H), 3.49 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.47 (app. br. s, 1 H), 2.06 (td, 1 H), 1.02 (d, 6 H).

- 10 Example 369(1) and 369(2): 2-(dimethylamino)-2-oxoethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate and 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide



15

The compound prepared in Example 363 (100 mg) was treated as detailed in Example 367 to give the two title compounds having the following physical properties.

Example 369(1): 20 mg


- 20 LC/MS  $t_R$  3.15 minutes; MS (ES<sup>+</sup>)  $m/z$  591 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.13 (d, 1 H), 7.87 (d, 1 H), 7.78 - 7.71 (m, 2 H), 7.71 - 7.63 (m, 1 H), 7.32 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1

H), 5.03 (s, 2 H), 3.52 - 3.40 (m, 1 H), 3.10 (ddd, 1 H), 3.08 (s, 3 H), 2.97 (s, 3 H), 2.71 - 2.56 (m, 1 H), 2.51 - 2.40 (m, 1 H).

Example 369(2): 12 mg

- 5 LC/MS  $t_R$  2.99 minutes; MS ( $ES^+$ )  $m/z$  533 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.76 - 7.72 (m, 4 H), 7.70 (d, 1 H), 7.31 (s, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.77 (dd, 1 H), 3.45 (td, 1 H), 3.25 (very br. s, 6 H), 3.13 (dd, 1 H), 2.71 - 2.59 (m, 1 H), 2.51 - 2.39 (m, 1 H).

- 10 Example 370: methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate




- To a solution of the compound prepared in Example 363 (100 mg) in a 4:1 mixture of dichloromethane and methanol (2.5 mL) was added a 2 M solution of (trimethylsilyl)diazomethane in diethyl ether (296  $\mu$ L) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated and the residue purified by column chromatography (50 - 100% ethyl acetate in heptanes, then 0 - 10% methanol in ethyl acetate) to give the title product having the following physical properties (72 mg).

- 20 LC/MS  $t_R$  3.46 minutes; MS ( $ES^+$ )  $m/z$  520 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.05 (d, 1 H), 7.83 (d, 1 H), 7.77 - 7.71 (m, 2 H), 7.72 - 7.65 (m, 1 H), 7.32 (br. s, 1 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.89 (s, 3 H), 3.52 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.47 (app. br. s, 1 H).

25

Example 371: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[5-(hydroxymethyl)-3-thienyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone



To a cooled (0 °C) tetrahydrofuran (6 mL) solution of the compound prepared in Example 370 (66 mg) was added a 1.2 M solution of diisobutylaluminium hydride in toluene (0.13 mL). The mixture was allowed to warm to room temperature and stirred 1.5 hours. The reaction mixture was cooled to 0 °C, a 1.2 M solution of diisobutylaluminium hydride in toluene (0.13 mL) added then the mixture warmed to room temperature and stirred 16 hours. To the reaction mixture, a 10% aqueous solution of potassium sodium tartrate was added (10 mL) followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 – 10% methanol in dichloromethane) to give the title product having the following physical properties (36 mg).

LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  492 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.78 - 7.71 (m, 2 H), 7.70 - 7.64 (m, 1 H), 7.44 (s, 1 H), 7.24 (s, 1 H), 7.17 (br. s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 4.73 (s, 2 H), 3.49 - 3.37 (m, 1 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.44 (app. br. s, 1 H).


Example 372: 4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1-[(prop-2-en-1-yloxy)carbonyl]-1H-imidazol-5-yl}thiophene-2-carboxylic acid

To an acetonitrile (8 mL) suspension of the compound prepared in Example 363 (400 mg) was added N,N-diisopropylethylamine (0.34 mL). The solution was cooled to 0 °C and a solution of allyl chloroformate (84  $\mu$ L) in acetonitrile (0.5 mL) added. The mixture was warmed to room temperature and stirred 16 hours before diluting with ethyl acetate (20 mL) and water (10 mL). The layers were separated and 1 M hydrochloric acid was added to the aqueous layer until pH 2 was attained. The aqueous layer was washed with dichloromethane, then a 9: 1 mixture of dichloromethane and propan-2-ol. The combined dichloromethane layers were dried and concentrated and the

residue triturated with ethyl acetate. Isolation of the solids by filtration gave the title product having the following physical properties (335 mg).

LC/MS  $t_R$  1.97 minutes; MS (ES<sup>+</sup>)  $m/z$  590 (M+H)<sup>a</sup>.

- 5 Example 373: [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate




- To an N,N-dimethylformamide (4 mL) solution of the compound prepared in  
 10 Example 372 (330 mg) was added potassium carbonate (116 mg) and the suspension stirred at room temperature for 10 minutes. A solution of chloromethyl pivalate (89  $\mu$ L) in N,N-dimethylformamide (1 mL) was added and the mixture stirred at room temperature for 16 hours. Water (10 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated.  
 15 The residue was purified by column chromatography (20 – 100% ethyl acetate in heptanes, then 0 – 10% methanol in ethyl acetate) to give the title product having the following physical properties (131 mg).

LC/MS  $t_R$  4.17 minutes; MS (ES<sup>+</sup>)  $m/z$  620 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.11 (d, 1 H), 7.90 (d, 1 H), 7.79 -  
 20 7.64 (m, 3 H), 7.36 (br. s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.95 (s, 2 H), 5.76 (dd, 1 H), 3.54 - 3.35 (m, 1 H), 3.19 - 3.01 (m, 1 H), 2.78 - 2.53 (m, 1 H), 2.45 (app. br. s, 1 H), 1.21 (s, 9 H).


Example 374: 2-methyl-2-propanyl 5-(bromoacetyl)-2-thiophenecarboxylate





tert-butyl 5-acetylthiophene-2-carboxylate (4.5 g) was treated as detailed in Example 361 to give the title compound having the following physical properties (3.9 g). TLC Rf 0.25 (10% ethyl acetate in hexane)

- 5 Example 375: 7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylic acid




- The same operation as in Example 7 → Example 8 → Example 9 was conducted from the compound prepared in Example 6 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 7 in the operation, 2-amino-5-methylphenylboronic acid pinacol ester and 1,1'-bis(diphenylphosphino)ferrocene palladium (II) dichloride dichloromethane complex were used). TLC Rf 0.18 (25% methanol in dichloromethane).

- 15 Example 376: 5-(2-{7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyI}-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride

- The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 375 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 374 was used). LC/MS  $t_R$  0.67 minutes; MS (ESI)  $m/z$  484 (M-H)<sup>+</sup> <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.65 (s, 1 H), 7.88 (m, 1 H), 7.69 (m, 1 H), 7.60 (d, 1 H), 7.55-7.42 (m, 3 H), 5.98 (s, 1 H), 5.89 (s, 1 H), 5.71 (m, 1 H), 3.30 (m, 1 H), 3.05 (m, 1 H), 2.50 - 2.02 (m, 2 H), 2.46 (s, 3 H).

Example 377: 5-(4-chloro-2-{7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyI}-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride




The same operation as in Example 51 → Example 52 → Example 338 → Example 363 was conducted from the compound prepared in Example 375 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 374 was used).

LC/MS  $t_R$  0.79 minutes; MS (ESI<sup>+</sup>)  $m/z$  518 (M-H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.66 (s, 1 H), 7.72 (d, 1 H), 7.58 (d, 1 H), 7.56-7.42 (m, 3 H), 5.94 (s, 1 H), 5.85 (s, 1 H), 5.55 (d, 1 H), 3.25 (m, 1 H), 3.00 (m, 1 H), 2.60 - 2.10 (m, 2 H), 2.46 (s, 3 H).


Example 378: 2-methyl-2-propanyl 5-bromo-1H-indole-3-carboxylate



A suspension of 5-bromo-1H-indole-3-carboxylic acid (500 mg) in N,N-dimethylformamide di-tert-butyl acetal (2.5 mL) was stirred at 100 °C for 17 hours. The reaction mixture was cooled to room temperature. Water was added followed by extraction with ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (5 – 40% ethyl acetate in hexane) to give the title product having the following physical properties (173 mg).

LC/MS  $t_R$  1.07 minutes; MS (ESI<sup>+</sup>)  $m/z$  297 (M+H)<sup>+</sup>.


Example 379: bis(2-methyl-2-propanyl) 5-bromo-1H-indole-1,3-dicarboxylate



To a dichloromethane (2 mL) solution of compound prepared in Example 378 were added di-*tert*-butyl dicarbonate (153 mg), triethylamine (0.12 mL) and 4-dimethylaminopyridine (7.1 mg) and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated and the residue was purified by column chromatography (5 – 20% ethyl acetate in hexane) to give the title product having the following physical properties (204 mg).

TLC R<sub>f</sub> 0.70 (25% ethyl acetate in hexane).

- 10 Example 380: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-1H-indole-3-carboxylic acid hydrochloride




The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 379 to give the title compound having the following physical properties.

LC/MS *t<sub>R</sub>* 0.61 minutes; MS (ES<sup>+</sup>) *m/z* 539 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 12.05 (s, 1 H), 9.69 (s, 1 H), 8.39 (s, 1 H), 8.09 (d, 1 H), 7.98 (d, 1 H), 7.86 – 7.82 (m, 2 H), 7.71 (s, 1 H), 7.61 – 7.58 (m, 2 H), 6.09 – 6.03 (m, 2 H), 5.87 – 5.76 (m, 1 H), 3.50 – 3.10 (m, 2 H), 2.65 – 2.30 (m, 2 H).


- 20 Example 381: 2-methyl-2-propanyl 5-bromo-1-(2-methyl-2-propanyl)-1H-indazole-3-carboxylate



5-bromo-1H-indazole-3-carboxylic acid (395 mg) was treated as detailed in Example 378 to give the title compound having the following physical properties (484 mg).

- 5 TLC Rf 0.91 (33% Ethyl acetate in hexane).

Example 382: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-1H-indazole-3-carboxylic acid




- 10 The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 381 to give the title compound having the following physical properties.

LC/MS  $t_R$  0.59 minutes; MS (ESI<sup>+</sup>)  $m/z$  540 (M+H)<sup>+</sup>

- <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 8.18 (s, 1 H), 7.82 - 7.62 (m, 4 H), 7.59 - 7.40 (m, 2 H), 5.98 (s, 1 H), 5.92 (s, 1 H), 5.62 (m, 1 H), 3.52 - 2.90 (m, 2 H), 2.60 - 2.30 (m, 2 H).

- Example 383(1), (2) and (3): 8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylic acid, 6-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylic acid and 6,8-dichloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylic acid



- To a tetrahydrofuran (5 mL) solution of the compound prepared in Example 9 (500 mg) was added 1,3-dichloro-5,5-dimethyl imidazolidine-2,4-dione (206 mg) and the reaction mixture stirred at 40 °C for 0.5 hours. The reaction mixture was concentrated. The residue obtained was purified by high performance liquid chromatography [5 to 95% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] to give the title compounds having the following physical properties.

Example 383(1): 143 mg

- 10 LC/MS  $t_R$  0.71 minutes; MS ( $ES^+$ )  $m/z$  392 and 394 ( $M+H$ )<sup>f</sup>.


Example 383(2): 121 mg

LC/MS  $t_R$  0.69 minutes; MS ( $ES^+$ )  $m/z$  392 and 394 ( $M+H$ )<sup>f</sup>.

- 15 Example 383(3): 75 mg

LC/MS  $t_R$  0.73 minutes; MS ( $ES^+$ )  $m/z$  426, 428 and 430 ( $M+H$ )<sup>f</sup>.

Example 384: 2-methyl-2-propanyl 4-(bromoacetyl)-2-pyridinecarboxylate




- 20 The same operation as in Example 378 → Example 139 → Example 361 was conducted from 4-bromo-2-pyridinecarboxylic acid to give the title compound having the following physical properties.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.99 (d, 1 H), 8.50 – 8.46 (m, 1 H), 7.91 (dd, 1 H), 4.47 (s, 2 H), 1.68 (s, 9 H).

25

Example 385: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid trifluoroacetate



The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 384 to give the title compound having the following physical properties. (Note: high performance liquid


5 chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt).

LC/MS  $t_R$  0.58 minutes; MS (ES<sup>+</sup>)  $m/z$  501 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.68 (s, 1 H), 8.70 (d, 1 H), 8.48 (s, 1 H), 8.36 (s, 1  
10 H), 8.12 (d, 1 H), 7.90 – 7.79 (m, 2 H), 7.74 (s, 1 H), 6.01 (s, 1 H), 5.97 (s, 1 H), 5.75 – 5.67 (m, 1 H), 3.40 – 3.23 (m, 1 H), 3.14 – 2.97 (m, 1 H), 2.70 – 2.25 (m, 2 H).

Example 386: 4-(2-{(3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-  
1,2,3,5-tetrahydro-3-indolizinyloxy}-1H-imidazol-5-yl)-2-pyridinecarboxylic acid

15 hydrochloride




The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 384 to give the title compound having the following physical properties. (Note: in the step corresponding to Example

20 51 in the operation, the compound prepared in Example 383(1) was used).

LC/MS  $t_R$  0.64 minutes; MS (ES<sup>+</sup>)  $m/z$  535 and 537 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.79 – 9.70 (m, 1 H), 8.34 (d, 1 H), 8.59 – 8.47 (m, 1  
H), 8.43 (s, 1 H), 8.27 – 8.17 (m, 1 H), 7.95 – 7.84 (m, 2 H), 7.74 – 7.67 (m, 1 H), 6.40  
– 6.31 (m, 1 H), 5.94 – 5.77 (m, 1 H), 3.46 – 3.25 (m, 1 H), 3.24 – 3.10 (m, 1 H), 2.77 –  
25 2.32 (m, 2 H).

Example 387: 5-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride



5 The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 374 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 383(1) was used).

LC/MS  $t_R$  0.75 minutes; MS (ES<sup>+</sup>)  $m/z$  540 and 542 (M+H)<sup>+</sup>

10 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.77 – 9.65 (m, 1 H), 7.89 – 7.84 (m, 2 H), 7.75 – 7.60 (m, 3 H), 7.45 – 7.20 (m, 1 H), 6.36 – 6.30 (m, 1 H), 5.80 – 5.69 (m, 1 H), 3.50 – 3.25 (m, 1 H), 3.22 – 3.05 (m, 1 H), 2.75 – 2.20 (m, 2 H).

Example 388: 5-(4-chloro-2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid trifluoroacetate

The same operation as in Example 51 → Example 52 → Example 338 → Example 363 was conducted from the compound prepared in Example 374 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 383(1) was used, high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt).

LC/MS  $t_R$  0.86 minutes; MS (ES<sup>+</sup>)  $m/z$  574, 576 and 578 (M+H)<sup>+</sup>

25 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.75 – 9.69 (m, 1 H), 7.91 – 7.79 (m, 2 H), 7.77 – 7.66 (m, 2 H), 7.46 (d, 1 H), 6.34 – 6.28 (m, 1 H), 5.72 – 5.59 (m, 1 H), 3.40 – 3.22 (m, 1 H), 3.20 – 3.04 (m, 1 H), 2.75 – 2.15 (m, 2 H).

Example 389: 5-(4-chloro-2-((3S)-6-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid trifluoroacetate

The same operation as in Example 51 → Example 52 → Example 338 →

- 5 Example 363 was conducted from the compound prepared in Example 374 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 383(2) was used, high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous
- 10 trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt). LC/MS  $t_R$  0.84 minutes; MS ( $ES^+$ )  $m/z$  574, 576 and 578 ( $M+H$ )<sup>f</sup> <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.70 (s, 1 H), 7.91 - 7.81 (m, 2 H), 7.77 - 7.65 (m, 2 H), 7.47 (d, 1 H), 6.35 (s, 1 H), 5.71 - 5.57 (m, 1 H), 3.45 - 3.25 (m, 1 H), 3.20 - 3.03 (m, 1 H), 2.70 - 2.15 (m, 2 H).


Example 390: 5-(2-((3S)-6,8-dichloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid trifluoroacetate

The same operation as in Example 51 → Example 52 → Example 363 was

- 20 conducted from the compound prepared in Example 374 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 383(3) was used, high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to
- 25 give the title product as the trifluoroacetic acid salt). LC/MS  $t_R$  0.78 minutes; MS ( $ES^+$ )  $m/z$  574, 576 and 578 ( $M+H$ )<sup>f</sup> <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.84 (s, 1 H), 7.99 - 7.88 (m, 2 H), 7.81 (d, 1 H), 7.69 - 7.61 (m, 2 H), 7.30 (d, 1 H), 5.88 - 5.82 (m, 1 H), 3.53 - 3.35 (m, 1 H), 3.30 - 3.16 (m, 1 H), 2.75 - 2.30 (m, 2 H).

Example 391: 4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid trifluoroacetate






The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 361 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 381(1) was used, high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt).

LC/MS  $t_R$  0.68 minutes; MS (ES<sup>+</sup>)  $m/z$  540 and 542 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.77 – 9.67 (m, 1 H), 8.22 – 8.11 (m, 2 H), 8.00 – 7.85 (m, 3 H), 7.98 – 7.86 (m, 1 H), 6.37 (s, 1 H), 5.94 – 5.77 (m, 1 H), 3.45 – 3.10 (m, 2 H), 2.80 – 2.30 (m, 2 H).

Example 392: 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride




The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 361 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 336 was used).

LC/MS  $t_R$  0.60 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.71 (s, 1 H), 8.20 (s, 1 H), 8.13 (s, 1 H), 7.99 (s, 1 H), 7.93 (d, 1 H), 7.90 – 7.80 (m, 2 H), 6.40 (s, 1 H), 5.83 – 5.74 (m, 1 H), 3.21 – 3.04 (m, 1 H), 2.96 – 2.81 (m, 1 H), 2.75 – 2.20 (m, 2 H).

Example 393(1) and 393(2): 8-bromo-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylic acid and 6-bromo-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylic acid



5

To a cooled (0 °C) tetrahydrofuran (5 mL) solution of the compound prepared in Example 9 (300 mg) was added N-bromosuccinimide (117 mg) and the reaction mixture stirred at 0 °C for 0.5 hours. To the reaction mixture, sodium thiosulfate was added and the mixture stirred for 0.5 hours before concentration. The residue obtained was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] to give the title compounds having the following physical properties.

10


Example 393(1): 145 mg

15 LC/MS  $t_R$  0.93 minutes; MS (ES<sup>+</sup>)  $m/z$  436 and 438 (M+H)<sup>+</sup>.

Example 393(2): 100 mg

LC/MS  $t_R$  0.88 minutes; MS (ES<sup>+</sup>)  $m/z$  436 and 438 (M+H)<sup>+</sup>.

20 Example 394: 4-(2-((3S)-8-bromo-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride



The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 361 to give the title compound


25

having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 393(1) was used)

LC/MS  $t_R$  0.69 minutes; MS ( $ES^+$ )  $m/z$  584 and 586 ( $M+H$ )<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.73 – 9.64 (m, 1 H), 8.25 – 8.08 (m, 2 H), 7.96 – 7.82 (m, 3 H), 7.65 – 7.56 (m, 1 H), 6.37 – 6.33 (m, 1 H), 5.96 – 5.81 (m, 1 H), 3.46 – 3.25 (m, 1 H), 3.23 – 3.08 (m, 1 H), 2.65 – 2.30 (m, 2 H).

Example 395: 3-[5-(3-bromophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone



The same operation as in Example 51 → Example 52 was conducted from 2-bromo-1-(3-bromophenyl)ethanone to give the title compound having the following physical properties.


LC/MS  $t_R$  0.85 minutes; MS ( $ES^+$ )  $m/z$  534 and 536 ( $M+H$ )<sup>f</sup>.

Example 396: 2-methyl-2-propanyl (2E)-3-[3-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]acrylate

To a 1,4-dioxane (1 mL) solution of the compound prepared in Example 395 (30 mg) was sequentially added N,N-dicyclohexylmethylamine (18  $\mu$ L) and tert-butyl acrylate (11 mg). The mixture was degassed with argon then bis(tri-tert-butylphosphine) palladium (1.4 mg) added and the reaction mixture irradiated under microwave conditions (100 W) at 110 °C for 10 minutes before concentration. The residue obtained was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] to give the title compound having the following physical properties (15 mg).

LC/MS  $t_R$  0.92 minutes; MS ( $ES^+$ )  $m/z$  582 ( $M+H$ )<sup>f</sup>.

Example 397: (2E)-3-[3-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acrylic acid hydrochloride




The compound prepared in Example 396 (22 mg) was treated as detailed in Example 363 to give the title compound having the following physical properties (20 mg).

LC/MS  $t_R$  0.66 minutes; MS (ES<sup>+</sup>)  $m/z$  526 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.69 (s, 1 H), 8.22 – 8.16 (m, 1 H), 7.90 – 7.79 (m, 3 H), 7.80 – 7.50 (m, 5 H), 6.67 (d, 1 H), 6.05 (s, 1 H), 6.02 (s, 1 H), 5.86 – 5.77 (m, 1 H), 3.40 - 3.20 (m, 1 H), 3.19 – 3.02 (m, 1 H), 2.80 – 2.20 (m, 2 H).

Example 398: (2E)-3-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acrylic acid trifluoroacetate




The same operation as in Example 51 → Example 52 → Example 396 → Example 363 was conducted from 2-bromo-1-(4-bromophenyl)ethanone to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt).

LC/MS  $t_R$  0.67 minutes; MS (ES<sup>+</sup>)  $m/z$  526 (M+H)<sup>+</sup>

$^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 8.15 - 7.70 (m, 8 H), 7.58 (d, 1 H), 6.57 (d, 1 H), 6.03 (s, 1 H), 6.00 (s, 1 H), 5.80 - 5.75 (m, 1 H), 3.50 - 3.20 (m, 1 H), 3.15 - 3.00 (m, 1 H), 2.70 - 2.20 (m, 2 H).


- 5 Example 399: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid trifluoroacetate



The same operation as in Example 51  $\rightarrow$  Example 52  $\rightarrow$  Example 364  $\rightarrow$

- 10 Example 363 was conducted from the compound prepared in Example 374 to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt).
- 15 LC/MS  $t_R$  0.80 minutes; MS (ESI $^-$ )  $m/z$  522 (M-H) $^+$
- $^1\text{H}$  NMR (300 MHz, methanol- $d_4$ )  $\delta$  9.34 (s, 1 H), 7.80 - 7.62 (m, 4 H), 7.19 (d, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.65 (m, 1 H), 3.40 (m, 1 H), 3.08 (m, 1 H), 2.62 (m, 1 H), 2.37 (m, 1 H).

- 20 Example 400: 4-[2-((4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl)methyl)-1H-imidazol-5-yl]-2-thiophenecarboxylic acid hydrochloride




- The same operation as in Example 6  $\rightarrow$  Example 7  $\rightarrow$  Example 8  $\rightarrow$  Example 9  $\rightarrow$  Example 51  $\rightarrow$  Example 52  $\rightarrow$  Example 363 was conducted from methyl 2-(4-hydroxy-6-methyl-2-oxo-1,2-dihydropyridin-1-yl)acetate [J. Het. Chem., 27(5), 1401
- 25

(1990)] to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 361 was used).

TLC Rf 0.65 (10% methanol, 1% acetic acid in dichloromethane)

- 5  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 8.17 (s, 1 H), 8.12 (d, 1 H), 7.90 (br. s, 1 H), 7.81 (app. s, 2 H), 7.72 (s, 1 H), 6.10 (d, 1 H), 6.01 (d, 1 H), 5.31 (s, 2 H), 2.40 (s, 3 H).

- 10 Example 401: ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylate



The same operation as in Example 361  $\rightarrow$  Example 51  $\rightarrow$  Example 52 was conducted from 5-acetyl-thiophene-2-carboxylic acid ethyl ester to give the title compound having the following physical properties.

- 15 TLC Rf 0.28 (ethyl acetate)  
 $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  10.88 (s, 1 H), 8.53 (s, 1 H), 7.69 (d, 1 H), 7.61 (dd, 1 H), 7.53 (d, 1 H), 7.49 (d, 1 H), 7.25 (m, 1 H), 7.19 (d, 1 H), 6.30 (d, 1 H), 5.80 (d, 1 H), 5.72 (s, 1 H), 4.33 (q, 2 H), 3.50 - 3.30 (m, 2 H), 3.10 - 2.95 (m, 1 H), 2.55 - 2.40 (m, 1 H), 1.37 (t, 3 H).

20

Example 402: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(1-oxido-4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operational sequence as in Example 78  $\rightarrow$  Example 51  $\rightarrow$  Example 52 was conducted from 4-acetylpyridine-N-oxide [patent WO9509853] to give the title

- 25 compound having the following physical properties.

LC/MS  $t_R$  3.19 minutes; MS ( $\text{ES}^+$ )  $m/z$  472 ( $\text{M}+\text{H}$ )<sup>b</sup>

$^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.44 (br. s, 0.7 H), 9.69 (s, 1 H), 8.29 (s, 0.3 H), 8.12 (d, 2 H), 7.75 - 7.89 (m, 3 H), 7.73 (br. s, 1 H), 7.66 (d, 2 H), 5.96 (d, 2 H), 5.62 (d, 1 H), 3.29 - 3.39 (m, 1 H), 3.00 (dd, 1 H), 2.54 (br. s, 1 H), 2.29 (t, 1 H).

30

Example 403: (3S)-3-[5-(3-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 173(21) (350 mg) was treated as detailed in Example 74 to give the title compound having the following physical properties (288 mg).

LC/MS  $t_R$  2.98 minutes; MS (ES<sup>+</sup>)  $m/z$  471 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.70 - 7.75 (m, 2 H), 7.64 - 7.69 (m, 1 H), 6.82 - 7.35 (m, 4 H), 6.62 (d, 1 H), 6.13 (s, 1 H), 6.07 (s, 1 H), 5.78 (dd, 1 H), 4.35 - 4.75 (m, 1 H), 3.36 - 3.49 (m, 1 H), 3.02 - 3.14 (m, 1 H), 2.61 (qd, 1 H), 2.44 (m, 1 H).

Example 404(1) to Example 404(3)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 403 and the corresponding acid chlorides or chloroformates using the method as detailed in Example 128.

Example 404(1): 2-methoxyethyl [3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.25 minutes; MS (ES<sup>+</sup>)  $m/z$  573 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.10 (br. s, 1 H), 9.51 - 9.86 (m, 2 H), 7.68 - 7.93 (m, 4 H), 7.03 - 7.45 (m, 4 H), 5.82 - 6.11 (2 x s, 2 H), 5.51 - 5.75 (m, 1 H), 4.12 - 4.25 (m, 2 H), 3.56 (m, 2 H), 3.31 (s, 2 H), 3.28 (s, 3 H), 3.01 (dd, 1 H), 2.30 - 2.42 (m, 1 H).

Example 404(2): methyl [3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.91 (d, 0.66 H), 8.62 - 8.49 (m, 0.34 H), 8.09 (t, 0.66 H), 7.79 - 7.60 (m, 4.33 H), 7.45 - 7.10 (br. m, 4 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.79 (d, 1 H), 3.74 (s, 3 H), 3.51 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.70 - 2.57 (m, 1 H), 2.56 - 2.31 (m, 1 H).

Example 404(3): N-[3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetamide

LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 8.36 (s, 1 H), 7.79 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.71 - 7.66 (m, 1 H), 7.49 (d, 1 H), 7.38 (d, 1 H), 7.33 - 7.26 (m, 2 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.79 (dd, 1 H), 3.48 - 3.40 (m, 1 H), 3.11 (dd, 1 H), 2.68 - 2.59 (m, 1 H), 2.50 - 2.41 (m, 1 H), 2.14 (s, 3 H).

#### Example 405(1) to Example 405(2)

The compounds of the present invention having the following physical data were prepared using the corresponding carboxylic acids from the compound prepared in Example 9 in the process of Example 234 → Example 51 → Example 52.

#### Example 405(1): methyl 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoate

LC/MS  $t_R$  3.58 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.32 (s, 1 H), 7.87 (d, 1 H), 7.90 (d, 1 H), 7.70 - 7.75 (m, 2 H), 7.65 - 7.70 (m, 1 H), 7.47 (t, 1 H), 7.42 (s, 1 H), 6.09 (s, 1 H), 6.13 (s, 1 H), 5.79 (dd, 1 H), 3.92 (s, 3 H), 3.41 - 3.52 (m, 1 H), 3.11 (ddd, 1 H), 2.60 - 2.69 (m, 1 H), 2.45 - 2.54 (m, 1 H).

#### Example 405(2): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-{5-[3-(methylsulfonyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.41 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.25 (br. s, 1 H), 8.01 (d, 1 H), 7.78 (d, 1 H), 7.69 - 7.75 (m, 2 H), 7.64 - 7.69 (m, 1 H), 7.60 (t, 1 H), 7.53 (br. s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.79 (dd, 1 H), 3.47 (td, 1 H), 3.14 - 3.16 (s, 3 H), 3.08 - 3.19 (m, 1 H), 2.59 - 2.70 (m, 1 H), 2.51 (br. s, 1 H).

#### Example 406: 3-(methoxymethyl)benzoic acid



A mixture of 3-bromomethyl benzoic acid methyl ester (0.573 g) and potassium carbonate (691 mg) in methanol (8 mL) and tetrahydrofuran (8 mL) was heated and



stirred under nitrogen at 55 °C for 2 h. The reaction mixture was filtered and the filtrate containing 3-methoxymethylbenzoic acid methyl ester used directly in the next stage. 1 M sodium hydroxide (3.75 ml) was added and the reaction was stirred at 70 °C for 30 minutes. The solvent was evaporated and the residue dissolved water and acidified (pH~ 2) by the addition of 1 N HCl. The aqueous layer was extracted with ethyl acetate and the organic layer washed with water, saturated brine and dried to afford the required product as a white solid (395 mg) having the following physical properties.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.10 (s, 1 H), 8.06 (d, 1 H), 7.62 (d, 1 H), 7.49 (t, 1 H), 4.54 (s, 2 H), 3.44 (s, 3 H).

Example 407: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[3-(methoxymethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operational sequence as in Example 234 → Example 51 → Example 52 was conducted from the compound prepared in Example 406 to give the title compound having the following physical properties.

LC/MS *t<sub>R</sub>* 3.28 minutes; MS (ES<sup>+</sup>) *m/z* 500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 9.36 (s, 1 H), 7.70 - 7.76 (m, 2 H), 7.66 - 7.70 (m, 1 H), 7.64 (s, 1 H), 7.58 (d, 1 H), 7.29 - 7.38 (m, 2 H), 7.21 (d, 1 H), 6.09 (s, 1 H), 6.12 (s, 1 H), 5.79 (dd, 1 H), 4.48 (s, 2 H), 3.41 - 3.49 (m, 1 H), 3.39 (s, 3 H), 3.10 (ddd, 1 H), 2.64 (qd, 1 H), 2.42 - 2.51 (m, 1 H).

Example 408: (3S)-3-[5-(3-acetylphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operational sequence as in Example 340 → Example 51 → Example 52 was conducted from 1,3-diacetylbenzene to give the title compound having the following physical properties.

LC/MS *t<sub>R</sub>* 3.40 minutes; MS (ES<sup>+</sup>) *m/z* 498 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 9.37 (s, 1 H), 8.31 (br. s, 1 H), 7.92 (br. s, 1 H), 7.86 (m, 2 H), 7.76 - 7.71 (m, 2 H), 7.71 - 7.62 (m, 2 H), 7.61 - 7.34 (m, 3 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.81 (dd, 1 H), 3.54 - 3.41 (m, 1 H), 3.13 (ddd, 1 H), 2.73 - 2.60 (m, 4 H), 2.52 (br. s, 1 H).

Example 409: (3S)-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}benzaldehyde

The same operational sequence as in Example 51 → Example 52 was conducted from 3-(2-bromoacetyl)benzaldehyde [J. Med. Chem., 50(18), 4405 (2007)] to give the  
5 title compound having the following physical properties.

LC/MS  $t_R$  1.64 minutes; MS (ES<sup>+</sup>)  $m/z$  484 (M+H)<sup>a</sup>.

Example 410: 3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzaldehyde O-methyloxime

10 The compound prepared in Example 409 (100 mg) was vigorously stirred with hydroxylamine hydrochloride (35 mg) and sodium acetate (34 mg) in ethanol (2.5 mL) overnight. The reaction was stirred for a further 6 days, ethanol evaporated under a stream of nitrogen and further hydroxylamine hydrochloride (70 mg) and sodium acetate (68 mg) added in 2 mL of a 4: 1 mixture of acetic acid and water. The reaction  
15 was heated at for a further 16 hours at 100 °C. The solvent was evaporated and purified by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (6.5 mg).

LC/MS  $t_R$  3.60 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.69 (s, 1 H), 8.26 (s, 1 H), 7.91 (s, 1 H), 7.80 (m, 3 H), 7.72 (d, 1 H), 7.50 (br. s, 1 H), 7.44 - 7.40 (m, 1 H), 7.39 - 7.35 (m, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.63 (d, 1 H), 4.08 (br. s, 1 H), 3.90 (s, 3 H), 3.00 (dd, 1 H), 2.49 - 2.45 (m, 1 H), 2.35 - 2.26 (m, 1 H).

25 Example 411: 3-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)benzamide

The same operational sequence as in Example 51 → Example 52 → Example 117 → Example 55 → Example 24 was conducted from 3-(bromoacetyl)benzonitrile to give the title compound having the following physical properties.

30 LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 8.20 (s, 1 H), 7.85 (d, 1 H), 7.76 (m, 3 H), 7.70 (d, 1 H), 7.49 (t, 1 H), 7.46 (s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.80 (dd, 1 H), 3.51 - 3.41 (m, 1 H), 3.11 (ddd, 1 H), 2.65 (ddd, 1 H), 2.0 (m, 1 H).

Example 412: 3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]benzonitrile

The compound prepared in Example 173(20) (91 mg) was treated using the  
5 method as detailed in Example 338 to give the title compound having the following physical properties (32 mg).

LC/MS  $t_R$  4.12 minutes; MS (ES<sup>+</sup>)  $m/z$  515 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 8.06 - 7.99 (m, 2 H), 7.76 - 7.72 (m, 2 H), 7.70 (s, 2 H), 7.67 - 7.61 (m, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.73 (dd, 1 H), 3.50 -  
10 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.71 - 2.61 (m, 1 H), 2.44 - 2.36 (m, 1 H).

Example 413: (3S)-3-[4-chloro-5-(3-hydroxyphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 173(22) (84 mg) was treated using the  
15 method as detailed in Example 338 to give the title compound having the following physical properties (57 mg).

LC/MS  $t_R$  3.81 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.65 (m, 1 H), 7.28 - 7.23 (m, 1 H), 7.18 (t, 1 H), 7.14 (d, 1 H), 6.76 (dd, 1 H), 6.13 (s, 1 H), 6.09  
20 (s, 1 H), 5.72 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.64 (qd, 1 H), 2.43 - 2.34 (m, 1 H).

Example 414: (3S)-3-[5-(2-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 74 was  
25 conducted from 2-bromo-1-(2-nitrophenyl)ethanone to give the title compound having the following physical properties.

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  471 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.39 - 7.20 (m, 2 H), 7.04 - 6.93 (m, 1 H), 6.78 (d, 1 H), 6.68 (br. s, 1 H), 6.16 (s, 1 H), 6.05 (s, 1 H), 5.82 (dd, 1 H), 3.49 - 3.38 (m, 1 H), 3.10 (dd, 1 H), 2.69 - 2.47 (m, 2 H).  
30

Example 415: methyl [2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 414 (50 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (30 mg).

LC/MS  $t_R$  4.09 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.65 (br. s, 1 H), 11.88 (br. s, 1 H), 9.68 (s, 1 H), 8.11 (d, 1 H), 7.81 (m, 2 H), 7.75 (s, 1 H), 7.68 - 7.63 (m, 2 H), 7.18 (t, 1 H), 6.99 (t, 1 H), 6.00 (s, 1 H), 5.99 (s, 1 H), 5.70 (d, 1 H), 3.63 (s, 3 H), 3.53 - 3.45 (m, 1 H), 3.07 (dd, 1 H), 2.62 - 2.54 (m, 1 H), 2.39 - 2.26 (m, 1 H).

Example 416: N-[2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]acetamide

The compound prepared in Example 414 (20 mg) was treated with acetyl chloride following the method of Example 128 to give the title compound having the following physical properties (12.7 mg).

LC/MS  $t_R$  3.45 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.66 (br. s, 1 H), 11.96 (s, 1 H), 9.71 (s, 1 H), 8.38 (d, 1 H), 7.86 - 7.78 (m, 2 H), 7.74 (d, 1 H), 7.70 - 7.64 (m, 2 H), 7.16 (t, 1 H), 7.02 (t, 1 H), 6.00 (s, 1 H), 5.99 (s, 1 H), 5.71 (d, 1 H), 3.33 - 3.29 (m, 1 H), 3.06 (dd, 1 H), 2.62 - 2.55 (m, 1 H), 2.30 (dd, 1 H), 1.93 (s, 3 H).

Example 417: prop-2-en-1-yl 2-acetylbenzoate

To an N,N-dimethylformamide (60 mL) suspension of 2-acetylbenzoic acid (5 g) was added potassium carbonate (5 g) and the mixture stirred at room temperature for 30 minutes. To the reaction mixture, allyl bromide (2.8 mL) was added and the mixture stirred at room temperature for 18 hours. The reaction mixture was concentrated and the residue suspended in water (50 mL) followed by extraction into ethyl acetate. The combined organic layers were washed with water, saturated saline, dried and concentrated to give the title compound having the following physical properties (6.04 g).

LC/MS  $t_R$  1.82 minutes; MS (ES<sup>+</sup>)  $m/z$  205 (M+H)<sup>a</sup>.

Example 418: prop-2-en-1-yl 2-{1-[(trimethylsilyl)oxy]ethenyl}benzoate

To a cooled (0 °C) dichloromethane (50 mL) solution of the compound prepared in Example 417 (2 g), triethylamine (3.06 mL) and bromo(trimethyl)silane (2.33 mL) were added sequentially and the mixture stirred at room temperature for 60 hours. To the reaction mixture, water (30 mL) was added followed by extraction into dichloromethane. The combined organic layers were washed with saturated saline, dried and concentrated to give the title compound having the following physical properties (2.53 g).

<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ 7.64 - 7.57 (m, 1 H), 7.50 - 7.45 (m, 1 H), 7.41 (dt, 1 H), 7.34 (dt, 1 H), 6.03 (tdd, 1 H), 5.41 (qd, 1 H), 5.27 (qd, 1 H), 4.78 (td, 2 H), 4.63 (d, 1 H), 4.49 (d, 1 H), 0.18 (s, 9 H).

Example 419: prop-2-en-1-yl 2-(2-bromoacetyl)benzoate

To a tetrahydrofuran (50 mL) solution of the compound prepared in Example 418 (2.53 g) was added N-bromosuccinimide (1.63 g) and the reaction stirred at room temperature for 1 hour. To the reaction mixture, water (25 mL) and saturated aqueous solution of sodium carbonate (25 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with water, saturated saline, dried and concentrated to give the title compound having the following physical properties (2.33 g).

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.05 (dd, 1 H), 7.65 (dt, 1 H), 7.56 (dt, 1 H), 7.42 (dd, 1 H), 6.02 (tdd, 1 H), 5.43 (qd, 1 H), 5.33 (dd, 1 H), 4.83 (td, 2 H), 4.34 (s, 2 H).

Example 420: 2-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy}-1H-imidazol-5-yl)benzoic acid

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 419 to give the title compound having the following physical properties. Hydrolysis of the allyl ester occurred under the conditions employed in the step corresponding to Example 52 in the operation.

LC/MS *t*<sub>R</sub> 3.32 minutes; MS (ES<sup>+</sup>) *m/z* 500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 10.72 (br. s, 1 H), 9.98 (br. s, 1 H), 8.71 (s, 1 H), 8.28 (d, 1 H), 7.80 - 7.64 (m, 3 H), 7.63 - 7.56 (m, 1 H), 7.56 - 7.49 (m, 2 H), 7.43 (t, 1 H), 6.36 (s, 1 H), 5.78 (s, 1 H), 5.50 (d, 1 H), 3.34 - 3.19 (m, 1 H), 2.93 (dd, 1 H), 2.86 - 2.69 (m,

1 H), 2.36 (td, 1 H).

Example 421: 2-bromo-1-[3-(tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]ethan-1-one

To a dichloromethane (3 mL) and glacial acetic acid (0.06 mL) solution of 1-[3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]ethanone (300 mg) was added  
5 bromine (62.64  $\mu$ L) and the mixture stirred at room temperature for 2 hours. The reaction mixture was concentrated and a saturated aqueous solution of sodium hydrogen carbonate (25 mL) was added to the residue followed by extraction with dichloromethane. The combined organic layers were washed with water, saturated  
10 saline, dried and concentrated to obtain the title compound having the following physical properties (0.37 g).

LC/MS  $t_R$  2.34 minutes; MS (ES<sup>+</sup>)  $m/z$  325 (M+H)<sup>a</sup>.

Example 422: (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-3-{5-[3-(tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1H-imidazol-2-yl}-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 51  $\rightarrow$  Example 52 was conducted from Example 421 to give the title compound having the following physical properties.  
LC/MS  $t_R$  1.83 minutes; MS (ES<sup>+</sup>)  $m/z$  582 (M+H)<sup>a</sup>.

20

Example 423: [3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]boronic acid

To a 1, 4-dioxane (1 mL) solution of the compound prepared in Example 422 (40 mg) was added 1 M hydrochloric acid in water (0.34 mL) and the mixture was  
25 stirred overnight at 90 °C. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate was added followed by extraction into dichloromethane. The combined organic layers were washed with saturated saline, dried (Na<sub>2</sub>SO<sub>4</sub>) and then concentrated. The residue was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile  
30 phase A (0.1% aqueous formic acid)] was used to give the title compound having the following physical properties (10 mg).

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.08 (s, 1 H), 9.70 (s, 1 H), 8.27 (s, 1 H), 8.16 - 8.07

(m, 2 H), 7.82 - 7.77 (m, 3 H), 7.72 (d, 1 H), 7.58 (d, 1 H), 7.43 (d, 1 H), 7.28 (t, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.68 - 5.58 (m, 1 H), 3.48 - 3.42 (m, 1 H), 3.01 (dd, 1 H), 2.54 - 2.53 (m, 1 H), 2.42 - 2.33 (m, 1 H).

5 Example 424: [3-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]boronic acid

The same operation as in Example 338 → Example 423 was conducted from the compound prepared in Example 422 to give the title compound having the following physical properties.

10 LC/MS  $t_R$  3.69 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.00 (s, 1 H), 9.72 (s, 1 H), 8.30 (br. s, 2 H), 8.07 (s, 1 H), 7.85 - 7.78 (m, 3 H), 7.73 (t, 2 H), 7.44 (t, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.59 (dd, 1 H), 3.28 - 3.25 (m, 1 H), 3.03 - 2.94 (m, 1 H), 2.59 - 2.54 (m, 1 H), 2.23 - 2.14 (m, 1 H).

15

Example 425: ethyl [3-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetate

To an ethanol (1 mL) solution of the compound prepared in Example 345 (39 mg) was added 2 M hydrochloric acid in ether (18.97  $\mu$ L) and the mixture was stirred at  
20 90 °C for 2 hours. The reaction mixture was concentrated and water was added to the residue and then neutralized with a saturated aqueous solution of sodium hydrogen carbonate followed by extraction with dichloromethane. The combined organic layers were washed with water, saturated saline, dried and then concentrated. The residue was purified by column chromatography (eluent: 50% to 100% ethyl acetate in heptanes and  
25 then 1% methanol in ethyl acetate) to afford the title compound having the following physical properties (30 mg).

LC/MS  $t_R$  3.47 minutes; MS (ES<sup>+</sup>)  $m/z$  542 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.71 - 7.66 (m, 1 H), 7.64 - 7.46 (m, 2 H), 7.42 - 7.26 (m, 2 H), 7.15 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.79 (d, 1 H), 4.14 (q, 2 H), 3.67 (s, 2 H), 3.51 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.70 - 2.58 (m, 1 H), 2.57 - 2.44 (m, 1 H), 1.24 (t, 3 H).  
30

Example 426: ethyl [3-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-

oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]acetate

The compound prepared in Example 346 (50 mg) was treated using the method as detailed in Example 425 to give the title compound having the following physical properties (28 mg).

- 5 LC/MS  $t_R$  4.07 minutes; MS (ES<sup>+</sup>)  $m/z$  576 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.77 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.65 - 7.60 (m, 2 H), 7.42 (t, 1 H), 7.27 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.73 (dd, 1 H), 4.16 (q, 2 H), 3.70 (s, 2 H), 3.50 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.65 (qd, 1 H), 2.45 - 2.35 (m, 1 H), 1.26 (t, 3 H).

10

Example 427: 2-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]acetamide

The compound prepared in Example 345 (45 mg) was treated using the method as detailed in Example 114 to give the title compound having the following physical properties (18 mg).

15

- LC/MS  $t_R$  2.86 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.10 (br. s, 1 H), 9.71 (s, 1 H), 7.85 - 7.75 (m, 3 H), 7.59 (s, 1 H), 7.54 - 7.46 (m, 3 H), 7.23 (t, 1 H), 7.06 (d, 1 H), 6.88 (br. s, 1 H), 5.98 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H), 3.42 - 3.39 (m, 3 H), 3.01 (dd, 1 H), 2.55 - 2.45 (m, 20 obs. 1 H), 2.40 - 2.33 (m, 1 H).

Example 428: 2-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]acetamide

The compound prepared in Example 346 (50 mg) was treated using the method as detailed in Example 114 to give the title compound having the following physical properties (21 mg).

25

- LC/MS  $t_R$  3.65 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.01 (br. s, 1 H), 9.71 (s, 1 H), 7.87 - 7.77 (m, 3 H), 7.63 - 7.48 (m, 3 H), 7.40 (t, 1 H), 7.22 (d, 1 H), 6.93 (br. s, 1 H), 5.99 (s, 1 H), 5.94 (s, 30 1 H), 5.58 (dd, 1 H), 3.42 (s, 2 H), 3.30 - 3.23 (m, 1 H), 3.07 - 2.87 (m, 1 H), 2.59 - 2.54 (m, 1 H), 2.19 (t, 1 H).

Example 429: 2-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-



tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-N-methylacetamide

The compound prepared in Example 345 (45 mg) was treated using the method as detailed in Example 114 using methylamine hydrochloride to give the title compound having the following physical properties (17 mg).

- 5 LC/MS  $t_R$  2.95 minutes; MS (ES<sup>+</sup>)  $m/z$  527 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.11 (br. s, 1 H), 9.70 (s, 1 H), 7.94 (m, 1 H), 7.85 - 7.75 (m, 3 H), 7.58 (s, 1 H), 7.54 (m, 2 H), 7.48 - 7.45 (m, 1 H), 7.23 (t, 1 H), 5.98 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H), 3.44 - 3.40 (m, 1 H), 3.39 (s, 2 H), 3.01 (dd, 1 H), 2.57 (s, 3 H), 2.55 - 2.45 (m, obs. 1 H), 2.38 - 2.35 (m, 1 H).

10

Example 430: 2-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-N-methylacetamide

The compound prepared in Example 346 (50 mg) was treated using the method as detailed in Example 114 using methylamine hydrochloride to give the title compound

- 15 having the following physical properties (20 mg).

LC/MS  $t_R$  3.77 minutes; MS (ES<sup>+</sup>)  $m/z$  561 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.01 (br. s, 1 H), 9.71 (s, 1 H), 8.00 (d, 1 H), 7.87 - 7.75 (m, 3 H), 7.63 - 7.51 (m, 2 H), 7.39 (t, 1 H), 7.21 (d, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.58 (dd, 1 H), 3.44 (s, 2 H), 3.31 - 3.19 (m, 1 H), 3.04 - 2.90 (m, 1 H), 2.58 (s, 3 H),  
20 2.55 - 2.45 (m, obs. 1 H), 2.19 (t, 1 H).

Example 431: 2-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)phenyl]-2-methylpropanoic acid - formic acid (1:1)

- 25 The same operation as in Example 139 → Example 340 → Example 51 → Example 52 → Example 342 was conducted from methyl 2-(3-bromophenyl)-2-methylpropanoate to give the title compound having the following physical properties.

LC/MS  $t_R$  3.23 minutes; MS (ES<sup>+</sup>)  $m/z$  542 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.71 (br. s, 1 H), 12.11 (br. s, 1 H), 9.69 (s, 1 H),  
30 8.17 (s, 1 H), 7.84 - 7.76 (m, 3 H), 7.70 - 7.61 (m, 1 H), 7.58 - 7.41 (m, 2 H), 7.35 - 7.24 (m, 1 H), 7.19 - 7.09 (m, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H), 3.50 - 3.20 (obs. m, 1H), 3.00 (dd, 1 H), 2.53 - 2.47 (obs. m, 1 H), 2.40 - 2.30 (m, 1 H), 1.49 (s, 6 H).

Example 432: 2-[3-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-2-methylpropanoic acid

The same operation as in Example 139 → Example 340 → Example 51 →

- 5 Example 52 → Example 338 → Example 342 was conducted from methyl 2-(3-bromophenyl)-2-methylpropanoate to give the title compound having the following physical properties.

LC/MS  $t_R$  3.82 minutes; MS (ES<sup>+</sup>)  $m/z$  576 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.07 (br. s, 1 H), 9.72 (s, 1 H), 7.87 - 7.77 (m, 3 H),  
10 7.69 (s, 1 H), 7.59 (d, 1 H), 7.43 (t, 1 H), 7.30 (d, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.58 (dd, 1 H), 3.28 - 3.21 (m, 1 H), 3.06 - 2.91 (m, 1 H), 2.60 - 2.55 (m, 1 H), 2.26 - 2.16 (m, 1 H), 1.52 (s, 6 H).

Example 433: 4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro indolizin-3-yl]-1H-imidazol-4-yl}benzoic acid

- 15 The same operation as in Example 51 → Example 52 → Example 37 → Example 55 → Example 24 was conducted from ethyl 4-(2-bromoacetyl)benzoate [Chem. Pharm. Bull., 54(9), 1318 (2006)] to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the  
20 operation, the compound prepared in Example 11 was used).

LC/MS  $t_R$  3.14 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.30 (s, 1 H), 9.70 (s, 1 H), 7.96 - 7.74 (m, 7 H),  
25 7.69 (s, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.66 (m, 1 H), 3.35 (obs. m, 1 H), 3.01 (dd, 1 H), 2.50 (obs. m, 1 H), 2.40 - 2.20 (m, 1 H).

Example 434(1) to Example 434(8)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 433 and the corresponding amines or amine HCl salts using the method as detailed in Example 114.

- 30 Example 434(1): formic acid - 4-(2-[(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-methoxybenzamide (1:1)

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 8.19 (br. s, 1 H), 7.83 - 7.67 (m, 7 H), 7.49 (s, 1 H), 6.14 (s, 1 H), 6.12 (s, 1 H), 5.81 (dd, 1 H), 3.83 (s, 3 H), 3.53 - 3.43 (m, 1 H), 3.22 - 3.09 (m, 1 H), 2.74 - 2.61 (m, 1 H), 2.55 - 2.43 (m, 1 H).

5 Example 434(2): formic acid - 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-ethoxybenzamide (1:1)

LC/MS *t<sub>R</sub>* 3.11 minutes; MS (ES<sup>+</sup>) *m/z* 543 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 7.84 - 7.67 (m, 7 H), 7.48 (s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.80 (dd, 1 H), 4.04 (q, 2 H), 3.54 - 3.40 (m, 1 H), 3.13 (ddd, 1 H), 2.71 - 2.58 (m, 1 H), 2.54 - 2.44 (m, 1 H), 1.33 (t, 3 H).

Example 434(3): 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(2-methoxyethyl)benzamide

LC/MS *t<sub>R</sub>* 3.09 minutes; MS (ES<sup>+</sup>) *m/z* 557 (M+H)<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 7.86 - 7.68 (m, 7 H), 7.54 - 7.35 (m, 1 H), 6.18 (s, 1 H), 6.12 (s, 1 H), 5.80 (d, 1 H), 3.59 (s, 4 H), 3.52 - 3.44 (m, 1 H), 3.40 (s, 3 H), 3.17 - 3.09 (m, 1 H), 2.71 - 2.61 (m, 1 H), 2.56 - 2.48 (m, 1 H).

Example 434(4): formic acid - 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-

20 oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(3-methoxypropyl)benzamide (1:1)

LC/MS *t<sub>R</sub>* 3.17 minutes; MS (ES<sup>+</sup>) *m/z* 571 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 7.87 - 7.68 (m, 7 H), 7.47 (s, 1 H), 6.13 (2 x s, 2 H), 5.81 (dd, 1 H), 3.50 (td, 5 H), 3.38 (s, 3 H), 3.20 - 3.10 (m, 1 H), 2.73 - 2.60 (m, 1 H), 2.54 - 2.45 (m, 1 H), 1.90 (m, 2 H).

Example 434(5): formic acid - 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(2-methoxyethoxy)benzamide (1:1)

30 LC/MS *t<sub>R</sub>* 3.05 minutes; MS (ES<sup>+</sup>) *m/z* 573 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 7.83 - 7.67 (m, 7 H), 7.49 (s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.81 (dd, 1 H), 4.13 (dd, 2 H), 3.75 - 3.68 (m, 2 H), 3.52 - 3.42 (m, 1 H), 3.41 (s, 3 H), 3.18 - 3.07 (m, 1 H), 2.68 (s, 1 H), 2.56 - 2.44 (m, 1 H).

Example 434(6): formic acid - 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(2-fluoroethyl)benzamide (1:1)

- 5 LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  545 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.91 - 7.66 (m, 7 H), 7.48 (s, 1 H), 6.16 (s, 1 H), 6.11 (s, 1 H), 5.81 (dd, 1 H), 4.63 (t, 1 H), 4.53 (t, 1 H), 3.76 - 3.62 (m, 2 H), 3.53 - 3.41 (m, 1 H), 3.13 (m, 1 H), 2.74 - 2.60 (m, 1 H), 2.54 - 2.42 (m, 1 H).

- 10 Example 434(7): 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(2,2-difluoroethyl)benzamide

- LC/MS  $t_R$  3.28 minutes; MS (ES<sup>+</sup>)  $m/z$  563 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.93 - 7.80 (m, 4 H), 7.79 - 7.66 (m, 4 H), 7.53 (br. s, 1 H), 6.20 - 6.09 (2 x s, 2 H), 6.06 - 5.89 (m, 1 H), 5.81 (m, 1 H), 3.75  
15 (m, 2 H), 3.47 (br. s, 1 H), 3.13 (m, 1 H), 2.71 - 2.58 (m, 1 H), 2.56 - 2.47 (m, 1 H).

Example 434(8): 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(3-fluoropropyl)benzamide

- LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  559 (M+H)<sup>b</sup>  
20 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.42 (s, 1 H), 7.87 - 7.66 (m, 7 H), 7.47 (s, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.81 (m, 1 H), 4.60 (t, 1 H), 4.50 (t, 1 H), 3.56 - 3.43 (m, 3 H), 3.18 - 3.09 (m, 1 H), 2.73 - 2.59 (m, 1 H), 2.54 - 2.43 (m, 1 H), 2.09 - 1.92 (m, 2 H).

- Example 435: 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(2-methoxyethyl)benzamide

- 25 The compound prepared in Example 434(3) (20 mg) was treated using the method as detailed in Example 338 to give the title compound having the following physical properties (2 mg).  
LC/MS  $t_R$  3.84 minutes; MS (ES<sup>+</sup>)  $m/z$  591 (M+H)<sup>b</sup>  
30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.39 (s, 1 H), 7.93 (d, 2 H), 7.84 (d, 2 H), 7.78 - 7.68 (m, 3 H), 6.15 (s, 1 H), 6.13 (s, 1 H), 5.81 - 5.69 (m, 1 H), 3.60 (s, 4 H), 3.46 (m, 1 H), 3.41 (s, 3 H), 3.19 - 3.09 (m, 1 H), 2.71 - 2.63 (m, 1 H), 2.46 - 2.38 (m, 1 H).

Example 436: formic acid - 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-(2-methoxyethoxy)benzamide (1:1)

The compound prepared in Example 434(5) (50 mg) was treated using the method as detailed in Example 338 to give the title compound having the following physical properties (7 mg).

LC/MS  $t_R$  3.77 minutes; MS ( $ES^+$ )  $m/z$  607 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.39 (s, 1 H), 7.88 - 7.81 (m, 4 H), 7.78 - 7.68 (m, 3 H), 6.15 (s, 1 H), 6.13 (s, 1 H), 5.75 (dd, 1 H), 4.18 - 4.12 (m, 2 H), 3.72 (m, 2 H), 3.43 (m, s, 4 H), 3.18 - 3.09 (m, 1 H), 2.68 (m, 1 H), 2.46 - 2.38 (m, 1 H).

Example 437: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-methyl-1H-imidazol-5-yl)benzamide

The same operation as in Example 114 → Example 78 → Example 51 → Example 52 was conducted from 4-propanoylbenzoic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  2.84 minutes; MS ( $ES^+$ )  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.41 (s, 1 H), 7.97 - 7.89 (m, 2 H), 7.74 (m, 6 H), 6.16 - 6.08 (m, 2 H), 5.78 - 5.72 (m, 1 H), 3.52 - 3.42 (m, 1 H), 3.17 - 3.08 (m, 1 H), 2.71 - 2.61 (m, 1 H), 2.52 - 2.36 (m, 4 H).

Example 438: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-methylbenzamide

The same operation as in Example 139 → Example 78 → Example 51 → Example 52 was conducted from 4-bromo-3-methylbenzamide to give the title compound having the following physical properties.

LC/MS  $t_R$  2.93 minutes; MS ( $ES^+$ )  $m/z$  514 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.29 (s, 1 H), 9.70 (s, 1 H), 7.90 (s, 1 H), 7.82 (m, 4 H), 7.72 (s, 1 H), 7.70 (d, 1 H), 7.39 (s, 1 H), 7.25 (s, 1 H), 5.98 (s, 2 H), 3.41 - 3.3 (obs, 2 H), 3.04 - 2.95 (m, 1 H), 2.45 (s, 3 H), 2.35 (m, 1 H).

Example 439: 4-(2-((7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-fluorobenzamide

The same operation as in Example 234 → Example 84 → Example 52 → Example 117 → Example 55 → Example 24 was conducted from 4-cyano-2-fluorobenzoic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  3.23 minutes; MS (ES<sup>+</sup>)  $m/z$  517 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 8.10 (br. s, 1 H), 7.80 - 7.64 (m, 5 H), 7.56 - 7.46 (m, 1 H), 6.17 (s, 1 H), 6.10 (s, 1 H), 5.89 - 5.76 (m, 1 H), 3.56 - 3.43 (m, 1 H), 3.19 - 3.07 (m, 1 H), 2.71 - 2.61 (m, 1 H), 2.58 - 2.47 (m, 1 H).

Example 440: 5-(2-bromoacetyl)pyridine-2-carbonitrile

- 10 6-cyanopyridine-3-carboxylic acid (750 mg) was treated using the method as detailed in Example 234 to give the title compound having the following physical properties (655 mg).

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 9.14 - 9.03 (m, 1 H), 8.24 (dd, 1 H), 7.69 (dd, 1 H), 4.25 (s, 2 H).

15

Example 441: 2-(6-cyanopyridin-3-yl)-2-oxoethyl 7-(2-([(tert-butoxy)carbonyl]amino})-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

- 20 The compound prepared in Example 19 (400 mg) was dissolved in acetonitrile (4 mL). N,N-diisopropylethylamine (64.1 μl) was added and the reaction stirred at room temperature for 10 minutes. Tetrabutyl ammonium iodide (164 mg) and 6-cyanopyridine-3-carboxylic acid (67.9 mg) was added and the reaction stirred at 50 °C for 1 h. The reaction mixture was diluted with water and extracted into ethyl acetate. The organic layer was washed with brine, dried and concentrated *in vacuo* to give the title compound having the following physical properties (196 mg).

- 25 LC/MS  $t_R$  2.14 minutes; MS (ES<sup>+</sup>)  $m/z$  493 (M+H)<sup>a</sup>.

Example 442: 5-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy}-1H-imidazol-5-yl)-2-pyridinecarboxamide

- 30 The same operation as in Example 52 → Example 117 → Example 55 → Example 24 was conducted from the compound prepared in Example 441 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.16 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.40 (s, 1 H), 9.69 (s, 1 H), 8.95 (d, 1 H), 8.26 - 8.15

(m, 1 H), 8.01 (m, 2 H), 7.87 - 7.74 (m, 4 H), 7.61 - 7.50 (m, 1 H), 5.98 (s, 1 H), 5.97 (s, 1 H), 5.65 (dd, 1 H), 3.35 (obs. m, 1 H), 3.17 (d, 1 H), 3.08 - 2.98 (m, 1 H), 2.40 - 2.30 (m, 1 H).

5 Example 443: 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny})-1H-imidazol-5-yl)benzenecarbothioamide

To a tetrahydrofuran (5 mL) solution of Example 114 (125 mg) was added Lawesson's reagent (71 mg) and the reaction heated at 70 °C overnight. Further Lawesson's reagent (70 mg) was added and the reaction heated at 70 °C for 3 h. On  
10 concentration, the residue was suspended in a saturated aqueous solution of sodium hydrogen carbonate (5 mL) and extracted into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0% - 20% methanol in dichloromethane) followed by preparative TLC (10% methanol in dichloromethane) to give the title product (4.4 mg)  
15 having the following physical properties.

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  515 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.28 (s, 1 H), 7.87 (d, 2 H), 7.64 - 7.56 (m, 5 H), 7.49 (s, 1 H), 6.02 (s, 1 H), 6.00 (s, 1 H), 5.71 (m, 1 H), 3.39 - 3.31 (m, 1 H), 3.09 - 3.02 (m, 1 H), 2.62 - 2.55 (m, 1 H), 2.41 - 2.33 (m, 1 H).

20

Example 444: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 371 was conducted from ethyl 4-(2-bromoacetyl)benzoate [Chem. Pharm. Bull., 54(9), 1318  
25 (2006)] to give the title compound having the following physical properties.

LC/MS  $t_R$  2.84 minutes; MS (ES<sup>+</sup>)  $m/z$  486 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.80 - 7.56 (m, 5 H), 7.37 (br. s, 3 H), 6.15 (s, 1 H), 6.13 (s, 1 H), 5.86 - 5.75 (m, 1 H), 4.62 (s, 2 H), 3.54 - 3.40 (m, 1 H), 3.13 (ddd, 1 H), 2.72 - 2.60 (m, 1 H), 2.48 - 2.31 (m, 1 H).

30

Example 445: (3S)-3-{4-chloro-5-[4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 444 (70 mg) was treated using the method

as detailed in Example 338 to give the title compound having the following physical properties (47.7 mg).

LC/MS  $t_R$  3.81 minutes; MS (ES<sup>+</sup>)  $m/z$  520 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.79 - 7.67 (m, 5 H), 7.46 (d, 2 H),  
5 6.13 (d, 2 H), 5.74 (dd, 1 H), 4.66 (s, 2 H), 3.52 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.66  
(m, 1 H), 2.46 - 2.36 (m, 1 H).

Example 446: (3S)-3-{4-chloro-5-[4-(1-hydroxyethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

10 The compound prepared in Example 166 (73 mg) was treated using the method as detailed in Example 338 to give the title compound having the following physical properties (24.5 mg).

LC/MS  $t_R$  3.93 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.39 (s, 1 H), 7.78 - 7.74 (m, 2 H), 7.72 - 7.68 (m, 3  
15 H), 7.48 (d, 2 H), 6.13 (d, 2 H), 5.74 (dd, 1 H), 4.89 - 4.85 (m, 1 H), 3.52 - 3.42 (m, 1  
H), 3.12 (ddd, 1 H), 2.71 - 2.62 (m, 1 H), 2.41 (dd, 1 H), 1.48 (d, 3 H).

Example 447: (3S)-3-{5-[2-chloro-4-(hydroxymethyl)phenyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

20 The same operation as in Example 234 → Example 51 → Example 52 → Example 371 was conducted from 2-chloro-4-(methoxycarbonyl)benzoic acid [Bioorg. Med. Chem. Lett., 20(22), 6748 (2010)] to give the title compound having the following physical properties.

LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  521 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.85 (br. s, 1 H), 7.78 - 7.69 (m, 3 H),  
7.62 - 7.50 (m, 2 H), 7.33 (d, 1 H), 6.18 (s, 1 H), 6.12 (s, 1 H), 5.83 (m, 1 H), 4.62 (s, 2  
H), 3.54 - 3.42 (m, 1 H), 3.18 - 3.08 (m, 1 H), 2.71 - 2.61 (m, 1 H), 2.54 (br. s, 1 H).

Example 448: methyl 5-(1-ethoxyethenyl)pyridine-2-carboxylate

30 To an N,N-dimethylformamide (10 mL) solution of methyl 5-bromopyridine-2-carboxylate (1.0 g) was added tributyl(1-ethoxyethenyl)stannane (1.72 mL). The mixture was degassed with nitrogen for 5 minutes, tetrakis(triphenylphosphine)palladium(0) (107 mg) added and the mixture heated at 120




°C for 3 hours. On cooling to room temperature, water (30 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with brine, dried and concentrated. The residue was purified by column chromatography (0 - 70% ethyl acetate in heptanes) to give the title product having the following physical

5 properties (618 mg).

LC/MS  $t_R$  1.80 minutes; MS (ES<sup>+</sup>)  $m/z$  208 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  8.96 (d, 1 H), 8.16 (dd, 1 H), 8.06 (d, 1 H), 5.08 (d, 1 H), 4.56 (d, 1 H), 3.95 (q, 2 H), 3.89 (s, 3 H), 1.37 (t, 3 H).

10 Example 449: methyl 5-(2-bromoacetyl)pyridine-2-carboxylate



To a tetrahydrofuran (28.7 ml) solution of the compound prepared in Example 448 (594 mg) was added water (1.9 ml) followed by N-bromosuccinimide (510 mg) and the mixture stirred at room temperature for 20 minutes. On concentration, the residue

15 was purified by column chromatography (10 - 70% ethyl acetate in heptanes) to give the title product having the following physical properties (568 mg).

LC/MS  $t_R$  1.48 minutes; MS (ES<sup>+</sup>)  $m/z$  258 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.24 (d, 1 H), 8.52 (dd, 1 H), 8.21 (d, 1 H), 5.07 (s, 2 H), 3.97 - 3.88 (m, 3 H).

20

Example 450: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(hydroxymethyl)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 371 was conducted from the compound prepared in Example 449 to give the title compound

25 having the following physical properties.

LC/MS  $t_R$  2.78 minutes; MS (ES<sup>+</sup>)  $m/z$  487 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.26 (m, 1 H), 8.77 - 8.64 (m, 1 H), 8.09 - 7.98 (m, 1 H), 7.68 (m, 2 H), 7.55 (d, 1 H), 7.49 - 7.39 (m, 2 H), 6.08 (s, 1 H), 5.95 (s, 1 H), 5.69 (dd, 1 H), 4.62 - 4.57 (m, 2 H), 3.41 - 3.30 (m, 1 H), 3.09 - 2.97 (m, 1 H), 2.61 - 2.43 (m,

1 H), 2.41 - 2.30 (m, 1 H).

Example 451: (3S)-3-{4-chloro-5-[6-(hydroxymethyl)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

5        The compound prepared in Example 450 (15 mg) was treated using the method as detailed in Example 338 to give the title compound having the following physical properties (7.4 mg).

LC/MS  $t_R$  3.24 minutes; MS (ES<sup>+</sup>)  $m/z$  521 (M+H)<sup>b</sup>

10        <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.40 (s, 1 H), 8.88 (m, 1 H), 8.19 (m, 1 H), 7.76 (m, 2 H), 7.71 - 7.62 (m, 2 H), 6.15 (s, 1 H), 6.11 (s, 1 H), 5.72 (m, 1 H), 4.75 (s, 2 H), 3.52 - 3.40 (m, 1 H), 3.21 - 3.10 (m, 1 H), 2.75 - 2.61 (m, 1 H), 2.47 - 2.36 (m, 1 H).

Example 452: methyl 5-bromo-6-chloropyridine-2-carboxylate

15        To a stirred solution of 5-bromo-2-(methoxycarbonyl)pyridine-N-oxide [Bioorg. Med. Chem. Lett., 18(4), 1407 (2008)] (2.1 g) in toluene (40 ml) was added phosphorus oxychloride (0.76 mL) and the mixture stirred at 80 °C for 16 h. To the reaction mixture, water (50 mL) was added followed by extraction into ethyl acetate. The combined organic layers were sequentially washed with a saturated aqueous solution of sodium hydrogen carbonate, water and brine then dried and concentrated. The residue was  
20        purified by column chromatography (50% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.65 g).

LC/MS  $t_R$  1.81 minutes; MS (ES<sup>+</sup>)  $m/z$  250 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  8.34 (d, 1 H), 7.98 (d, 1 H), 4.02 - 3.93 (m, 3 H).

25        Example 453: (3S)-3-{5-[2-chloro-6-(hydroxymethyl)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

      The same operation as in Example 448 → Example 449 → Example 51 → Example 52 → Example 371 was conducted from the compound prepared in Example 452 to give the title compound having the following physical properties.

30        LC/MS  $t_R$  3.24 minutes; MS (ES<sup>+</sup>)  $m/z$  521 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.26 (s, 1 H), 8.31 - 8.22 (m, 1 H), 7.63 - 7.55 (m, 4 H), 7.46 (d, 1 H), 6.09 (s, 1 H), 5.91 (s, 1 H), 5.71 (m, 1 H), 4.55 (s, 2 H), 3.43 - 3.31 (m, 1 H), 3.09 - 2.96 (m, 1 H), 2.62 - 2.48 (m, 1 H), 2.47 - 2.36 (m, 1 H).

Example 454: N-[4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -1H-imidazol-5-yl)phenyl]-4-methoxybutanamide

The compound prepared in Example 127 (105 mg) was treated with 4-methoxybutanoic acid (33  $\mu$ L) following the method of Example 133 to give the title compound having the following physical properties (37.3 mg).

LC/MS  $t_R$  3.24 minutes; MS (ES<sup>+</sup>)  $m/z$  571 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.75 - 7.69 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.62 - 7.53 (m, 4 H), 7.25 (s, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.77 (dd, 1 H), 3.46 (t, 2 H), 3.48 - 3.39 (m, 1 H), 3.34 (s, 3 H), 3.09 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.44 (t, 2 H), 2.50 - 2.41 (m, 1 H), 1.94 (quintet, 1 H).

Example 455: (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-3-[5-(4-nitrophenyl)-1H-imidazol-2-yl]-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 51  $\rightarrow$  Example 52 was conducted from 2-bromo-1-(4-nitrophenyl)ethanone to give the title compound having the following physical properties.

LC/MS  $t_R$  1.90 minute; MS (ES<sup>+</sup>)  $m/z$  520 (M+H)<sup>a</sup>.

Example 456: (3S)-3-[5-(4-aminophenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

The compound prepared in Example 455 (4.26 g) was treated as detailed in Example 74 to give the title compound having the following physical properties (1.76 g).

LC/MS  $t_R$  1.36 minutes; MS (ES<sup>+</sup>)  $m/z$  471 (M+H)<sup>a</sup>.

Example 457(1) to Example 457(5)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 456 and the corresponding acid chlorides or anhydrides using the method as detailed in Example 128.

Example 457(1): N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -1H-imidazol-5-yl)phenyl]acetamide

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  513 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.66 - 7.49 (m, 4 H), 7.36 - 7.14 (m, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.59 (m, 1 H), 2.47 (br. s, 1 H), 2.13 (s, 3 H).

5

Example 457(2): N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]-2-fluoroacetamide

LC/MS *t<sub>R</sub>* 2.99 minutes; MS (ES<sup>+</sup>) *m/z* 531 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.09 (s, 1 H), 10.06 (s, 1 H), 9.69 (s, 1 H), 7.83 - 7.76 (m, 3 H), 7.70 - 7.56 (m, 4 H), 7.44 (d, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H), 4.96 (d, 2 H), 3.47 - 3.36 (m, 2 H), 3.04 - 2.94 (m, 1 H), 2.38 - 2.30 (m, 1 H).

10

Example 457(3): N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]-2,2-difluoroacetamide

LC/MS *t<sub>R</sub>* 3.15 minutes; MS (ES<sup>+</sup>) *m/z* 549 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.75 - 7.60 (m, 7 H), 7.33 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 6.16 (t, 1 H), 5.78 (dd, 1 H), 3.49 - 3.40 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.59 (m, 1 H), 2.48 (br. s, 1 H).

15

Example 457(4): N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]tetrahydro-3-furancarboxamide

LC/MS *t<sub>R</sub>* 2.98 minutes; MS (ES<sup>+</sup>) *m/z* 569 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.72 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.58 (br. s, 4 H), 7.29 (br. s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.06 - 4.00 (m, 1 H), 3.96 - 3.87 (m, 2 H), 3.86 - 3.80 (m, 1 H), 3.49 - 3.39 (m, 1 H), 3.23 - 3.16 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.46 (br. s, 1 H), 2.20 (m, 2 H).

25

Example 457(5): N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]tetrahydro-2H-pyran-4-carboxamide

LC/MS *t<sub>R</sub>* 3.03 minutes; MS (ES<sup>+</sup>) *m/z* 583 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.76 - 7.70 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.65 - 7.48 (m, 4 H), 7.35 - 7.10 (m, 1 H), 6.12 (br. s, 1 H), 6.08 (br. s, 1 H), 5.78 (d,

30

1 H), 4.05 – 3.97 (m, 2 H), 3.53 - 3.39 (m, 3 H), 3.14 - 3.05 (m, 1 H), 2.69 - 2.58 (m, 2 H), 2.55 - 2.37 (m, 1 H), 1.91 - 1.80 (m, 2 H), 1.80 - 1.73 (m, 2 H).

5 Example 458: formic acid - N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4-fluoro-1H-imidazol-5-yl)phenyl]acetamide (1:1)

The compound prepared in Example 457(1) (70 mg) was treated as detailed in Example 364 to give the title compound having the following physical properties (14 mg).

- 10 LC/MS  $t_R$  3.79 minutes; MS (ES<sup>+</sup>)  $m/z$  531 (M+H), 553 (M+Na) <sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.61 (d, 2 H), 7.50 (d, 2 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.69 (dd, 1 H), 3.47 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.67 - 2.58 (m, 1 H), 2.40 - 2.34 (m, 1 H), 2.13 (s, 3 H).

15 Example 459: formic acid - N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]ethanethioamide (1:1)

- To a tetrahydrofuran solution (2 mL) of the compound prepared in Example 457(1) (70 mg) was added Lawesson's reagent (55 mg) and the mixture heated at 50 °C for 2 hours, then warmed to 70 °C and stirred 3 hours. Further Lawesson's reagent (55 mg) was added and the mixture cooled to 50 °C and stirred 16 hours. An additional portion of Lawesson's reagent (55 mg) was added, the mixture stirred 50 °C for 6 hours, then further Lawesson's reagent (55 mg) added and the mixture stirred at 50 °C for 16 hours. On cooling to room temperature, the mixture was diluted with a saturated aqueous solution of sodium hydrogen carbonate (20 mL) and extracted into ethyl acetate.
- 25 The combined organic layers were dried and concentrated and the residue purified by column chromatography (0 – 6% methanol in dichloromethane) followed by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (17 mg).
- 30 LC/MS  $t_R$  3.17 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H) <sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.16 (br. s, 1 H), 11.64 - 11.51 (m, 1 H), 9.70 (s, 1 H), 7.84 - 7.76 (m, 5 H), 7.74 - 7.66 (m, 2 H), 7.51 (br. s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1

H), 5.62 (d, 1 H), 3.56 - 3.24 (m, 1 H), 3.04 - 2.96 (m, 1 H), 2.59 (s, 3 H), 2.53 - 2.50 (m, 1 H), 2.35 - 2.28 (m, 1 H).

5 Example 460: N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]formamide

A mixture of acetic anhydride (28  $\mu$ L) and formic acid (14  $\mu$ L) was stirred at 60 °C for 2 hours. On cooling to room temperature, this solution was added to a stirred tetrahydrofuran solution (2 mL) of the compound prepared in Example 456 (80 mg) and this mixture stirred at room temperature for 1 hour 30 minutes. To the reaction mixture,  
10 water (20 mL) was added followed by extraction into ethyl acetate. The combined organic layers were dried and concentrated and the residue purified by column chromatography (0 - 8% methanol in dichloromethane) to give the title compound having the following physical properties (53 mg).

LC/MS  $t_R$  2.87 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.26 (s, 1 H), 7.79 - 7.52 (m, 7 H), 7.36 - 7.16 (m, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.50 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.70 - 2.58 (m, 1 H), 2.55 - 2.39 (m, 1 H).

20 Example 461: N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]-N-hydroxyacetamide

To an N,N-dimethylformamide solution (30 mL) of the compound prepared in Example 455 (150 mg) was added water (1 mL), ammonium chloride (64 mg) and zinc dust (78 mg) and the mixture stirred at room temperature for 1 hour and 30 minutes. To the reaction mixture, water (15 mL) was added followed by extraction into  
25 dichloromethane. The combined organic layers were dried and concentrated. To a dichloromethane (10 mL) solution of this crude residue was added N,N-diisopropylethylamine (53  $\mu$ L) and acetyl chloride (21  $\mu$ L) and the mixture stirred at room temperature for 45 minutes. An additional aliquot of acetyl chloride (21  $\mu$ L) was added and the mixture stirred at room temperature for a further 2 hours before diluting  
30 with water (30 mL) and extraction into ethyl acetate. The combined organic layers were dried and concentrated and the residue purified by column chromatography (0 - 10% methanol in dichloromethane) to give the title compound having the following physical properties (33 mg).

LC/MS  $t_R$  2.93 minutes; MS ( $ES^+$ )  $m/z$  529 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.78 - 7.52 (m, 7 H), 7.31 (br. s, 1 H), 6.11 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.09 (ddd, 1 H), 2.68 - 2.57 (m, 1 H), 2.50 - 2.42 (m, 1 H), 2.29 (br. s, 3 H).

5

Example 462: (1E)-N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-N'-hydroxyethanimidamide

To an ethanol suspension (1 mL) of the compound prepared in Example 456 (80 mg) was added ethyl N-hydroxyethenecarboximide (35 mg) and the mixture stirred at 80 °C for 3 days. On concentration the residue was purified by column chromatography (0 – 6% methanol in dichloromethane) to give the title compound having the following physical properties (13 mg).

LC/MS  $t_R$  2.72 minutes; MS ( $ES^+$ )  $m/z$  528 ( $M+H$ ), 264.5 ( $M/2+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.73 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.66 - 7.56 (m, 2 H), 7.27 (br. s, 1 H), 7.13 (d, 2 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.50 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.47 (br. s, 1 H), 1.93 (s, 3 H).

Example 463: formic acid - (1E)-N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-N'-methoxyethanimidamide (2:1)

The compound prepared in Example 456 (80 mg) was treated as detailed in Example 462 using ethyl N-methoxyethenecarboximide [Tetrahedron, 40(19), 3769 (1984)] in place of N-hydroxyacetimide to give the title compound having the following physical properties (2.4 mg).

LC/MS  $t_R$  2.99 minutes; MS ( $ES^+$ )  $m/z$  542 ( $M+H$ ), 271.5 ( $M/2+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.62 (d, 2 H), 7.28 (s, 1 H), 7.12 (d, 2 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.78 (s, 3 H), 3.49 - 3.40 (m, 1 H), 3.15 - 3.07 (m, 1 H), 2.67 - 2.59 (m, 1 H), 2.51 - 2.43 (m, 1 H), 1.89 (s, 2 H).

Example 464: N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]-3-hydroxy-3-methylbutanamide

The compound prepared in Example 456 (80 mg) was treated with 3-hydroxy-3-methyl butyric acid (22 mg) following the method of Example 133 to give the title compound having the following physical properties (26 mg).

LC/MS  $t_R$  2.99 minutes; MS ( $ES^+$ )  $m/z$  571 (M+H) <sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.66 - 7.52 (m, 4 H), 7.36 - 7.16 (m, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.50 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.52 (s, 2 H), 2.53 - 2.40 (m, 1 H), 1.33 (s, 6 H).

- 10 Example 465: 3-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-3-oxopropanoic acid

The same operation as in Example 134 → Example 135 was conducted from the compound prepared in Example 456 to give the title compound having the following physical properties.

- 15 LC/MS  $t_R$  2.84 minutes; MS ( $ES^+$ )  $m/z$  557 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.08 (br. s, 1 H), 10.53 (br. s, 1 H), 9.69 (s, 1 H), 7.93 - 7.74 (m, 3 H), 7.68 - 7.49 (m, 4 H), 7.41 (br. s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.61 (dd, 1 H), 3.51 - 3.27 (m, 1 H), 3.24 (s, 2 H), 3.06 - 2.94 (m, 1 H), 2.52 - 2.44 (m, 1 H), 2.37 - 2.27 (m, 1 H).

20

Example 466: 4-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-4-oxobutanoic acid

To a stirred dichloromethane solution (2 mL) of the compound prepared in Example 456 (60 mg) was added triethylamine (27  $\mu$ L) and succinic anhydride (19 mg) and the mixture stirred at room temperature for 16 hours. To the reaction mixture, water (10 mL) and dichloromethane (10 mL) were added and the resultant precipitate isolated by filtration to give the title compound having the following physical properties (24 mg).

LC/MS  $t_R$  2.84 minutes; MS ( $ES^+$ )  $m/z$  570 (M+H) <sup>b</sup>

- 30 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.06 (br. s, 1 H), 9.94 (s, 1 H), 9.69 (s, 1 H), 7.82 - 7.77 (m, 3 H), 7.63 - 7.52 (m, 4 H), 7.41 (d, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.61 (d, 1 H), 3.43 - 3.37 (m, 1 H), 3.03 - 2.96 (m, 1 H), 2.57 - 2.50 (m, 5 H), 2.38 - 2.30 (m, 1 H).



Example 467: 4-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2,2-dimethyl-4-oxobutanoic acid

The compound prepared in Example 456 (50 mg) was treated with 2,2-dimethylsuccinic anhydride (18  $\mu$ L) following the method of Example 466 to give the title compound having the following physical properties (13 mg).  
LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  599 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.39 (s, 1 H), 7.75 (dd, 1 H), 7.72 (d, 1 H), 7.69 (d, 1 H), 7.68 - 7.60 (m, 5 H), 6.16 (s, 1 H), 6.14 (s, 1 H), 5.86 (dd, 1 H), 3.43 - 3.36 (m, 1 H), 3.35 (s, 2 H), 3.26 - 3.15 (m, 1 H), 2.87 - 2.77 (m, 1 H), 2.69 (s, 2 H), 2.49 - 2.39 (m, 1 H), 1.31 (s, 6 H).

Example 468: 4-{[4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2,2-dimethyl-4-oxobutanoic acid

The compound prepared in Example 467 (17 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (6.3 mg).  
LC/MS  $t_R$  3.87 minutes; MS (ES<sup>+</sup>)  $m/z$  633 (M+H)<sup>b</sup>  
NMR analysis showed a 2:1 ratio of tautomers.  
Major tautomer: <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.90 (s, 1 H), 12.05 (br. s, 1 H), 10.02 (s, 1 H), 9.71 (s, 1 H), 7.84 - 7.77 (m, 3 H), 7.66 (d, 2 H), 7.61 (d, 2 H), 5.99 (s, 1 H), 5.93 (s, 1 H), 5.56 (dd, 1 H), 3.30 - 3.23 (m, 1 H), 3.04 - 2.95 (m, 1 H), 2.59 (s, 2 H), 2.55 - 2.46 (obs. m, 1 H), 2.23 - 2.16 (m, 1 H), 1.21 (s, 6 H).

Minor tautomer: <sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.90 (s, 1 H), 12.05 (br. s, 1 H), 9.88 (br. s, 1 H), 9.69 (s, 1 H), 7.84 - 7.77 (m, 3 H), 7.61 - 7.30 (m, 4 H), 5.99 (s, 1 H), 5.96 (s, 1 H), 5.67 - 5.59 (m, 1 H), 3.30 - 3.23 (m, 1 H), 3.04 - 2.95 (m, 1 H), 2.57 (s, 2 H), 2.55 - 2.46 (obs. m, 1 H), 2.35 - 2.25 (m, 1 H), 1.20 (s, 6 H).

Example 469: N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-2-(2-methoxyethoxy)acetamide

The compound prepared in Example 456 (80 mg) was treated with (2-

methoxyethoxy)acetic acid (21  $\mu$ L) following the method of Example 133 to give the title compound having the following physical properties (18.4 mg).

LC/MS  $t_R$  3.07 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.09 (br. s, 1 H), 9.69 (s, 1 H), 9.60 (s, 1 H), 7.84 -  
5 7.76 (m, 3 H), 7.64 (d, 2 H), 7.59 (d, 2 H), 7.43 (s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H),  
5.61 (d, 1 H), 4.06 (s, 2 H), 3.70 - 3.64 (m, 2 H), 3.56 - 3.50 (m, 2 H), 3.46 - 3.28 (obs.  
m, 1 H), 3.30 (s, 3 H), 3.00 (dd, 1 H), 2.58 - 2.44 (obs. m, 1 H), 2.38 - 2.28 (m, 1 H).

10 Example 470: (3S)-3-[5-(4-aminophenyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-  
(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 51  $\rightarrow$  Example 52  $\rightarrow$  Example 338  $\rightarrow$   
Example 74 was conducted from 2-bromo-1-(4-nitrophenyl)ethanone to give the title  
compound having the following physical properties.

LC/MS  $t_R$  2.07 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>a</sup>.

15

Example 471: N-[4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-  
1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-2-(2-  
methoxyethoxy)acetamide

20 The compound prepared in Example 470 (50 mg) was treated with (2-  
methoxyethoxy)acetic acid (8.4  $\mu$ L) following the method of Example 133 to give the  
title compound having the following physical properties (13.8 mg).

LC/MS  $t_R$  3.95 minutes; MS (ES<sup>+</sup>)  $m/z$  621 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.93 (br. s, 1 H), 9.78 (s, 1 H), 9.71 (s, 1 H), 7.85 -  
7.78 (m, 3 H), 7.73 (d, 2 H), 7.65 (d, 2 H), 5.99 (s, 1 H), 5.93 (s, 1 H), 5.57 (dd, 1 H),  
25 4.09 (s, 2 H), 3.68 (dd, 2 H), 3.54 (dd, 2 H), 3.30 (s, 3 H), 3.29 - 3.22 (m, 1 H), 2.99 (dd,  
1 H), 2.58 - 2.43 (obs. m, 1 H), 2.19 (t, 1 H).

30 Example 472: 5-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-  
tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-5-oxopentanoic acid -  
formic acid (1:1)

The compound prepared in Example 456 (170 mg) was treated with glutaric  
anhydride (62 mg) following the method of Example 466 to give the title compound  
after purification by high performance liquid chromatography [5 to 100% mobile phase

B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (19 mg).

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  585 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.07 (br. s, 1 H), 9.89 (br. s, 1 H), 9.69 (s, 1 H), 8.19 (s, 1 H), 7.84 - 7.77 (m, 3 H), 7.59 (d, 2 H), 7.56 (d, 2 H), 7.38 (br. s, 1 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.61 (dd, 1 H), 3.42 - 3.32 (obs. m, 1 H), 2.99 (dd, 1 H), 2.55 - 2.44 (obs. m, 1 H), 2.34 (t, 2 H), 2.38 - 2.24 (obs. m, 1 H), 2.27 (t, 2 H), 1.80 (quintet, 2 H).

Example 473: 5-{[4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-5-oxopentanoic acid

The compound prepared in Example 470 (100 mg) was treated with glutaric anhydride (23 mg) following the method of Example 466 to give the title compound having the following physical properties (30.8 mg).

LC/MS  $t_R$  3.70 minutes; MS (ES<sup>+</sup>)  $m/z$  619 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.91 (br. s, 1 H), 10.10 (s, 1 H), 9.71 (s, 1 H), 7.84 - 7.77 (m, 3 H), 7.68 (d, 2 H), 7.62 (d, 2 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.56 (dd, 1 H), 3.29 - 3.23 (m, 1 H), 2.98 (ddd, 1 H), 2.54 (s, 3 H), 2.53 - 2.47 (obs. m, 1 H), 2.36 (t, 2 H), 2.26 (t, 2 H), 2.22 - 2.16 (m, 1 H), 1.80 (quintet, 1 H).

Example 474: methyl 5-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-5-oxopentanoate

A methanol (5 mL) solution of the compound prepared in Example 472 (125 mg) was stirred at room temperature for 2 days. The reaction mixture was concentrated and the residue purified by column chromatography (0% to 20% methanol in dichloromethane) to afford the title compound having the following physical properties (9 mg).

LC/MS  $t_R$  3.15 minutes; MS (ES<sup>+</sup>)  $m/z$  599 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.75 - 7.70 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.59 (d, 2 H), 7.56 (d, 2 H), 7.25 (br. s, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 3.66 (s, 3 H), 3.49 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.43 (t, 4 H), 2.50 - 2.38 (obs. m, 1 H), 1.99 (quintet, 2 H).

Example 475: N-[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-N'-methylpentanediamide

The compound prepared in Example 472 (98 mg) was treated with methylamine hydrochloride (19 mg) following the method of Example 114 to give the title compound having the following physical properties (2.0 mg).

LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  598 (M+H), 300 (M+2H)/2 <sup>b</sup>

5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.06 (s, 1 H), 9.85 (s, 1 H), 9.69 (s, 1 H), 7.83 - 7.78 (m, 3 H), 7.77 - 7.71 (m, 1 H), 7.63 - 7.58 (m, 2 H), 7.57 - 7.52 (m, 2 H), 7.40 (d, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.61 (dd, 1 H), 3.44 - 3.35 (m, 1 H), 3.00 (dd, 1 H), 2.56 (d, 3 H), 2.54 - 2.45 (obs. m, 1 H), 2.28 (t, 2 H), 2.35 - 2.25 (m, 1 H), 2.10 (t, 2 H), 1.84 - 1.75 (m, 2 H).

10

Example 476: (2-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2-oxoethoxy)acetic acid

The compound prepared in Example 456 (170 mg) was treated with diglycolic anhydride (63 mg) following the method of Example 466 to give the title compound

15 having the following physical properties (87 mg).

LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.54 (br. s, 1 H), 12.08 (br. s, 1 H), 10.03 (br. s, 1 H), 9.69 (s, 1 H), 7.84 - 7.74 (m, 3 H), 7.62 (br. s, 4 H), 7.43 (br. s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.62 (d, 1 H), 4.19 (s, 2 H), 4.16 (s, 2 H), 3.52 - 3.17 (obs. m, 1 H), 2.99 (dd, 1 H), 2.57 - 2.46 (obs. m, 1 H), 2.38 - 2.21 (m, 1 H).

20

Example 477: (2-{[4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2-oxoethoxy)acetic acid

25 The compound prepared in Example 470 (100 mg) was treated with diglycolic anhydride (23 mg) following the method of Example 466 to give the title compound having the following physical properties (25.2 mg).

LC/MS  $t_R$  3.68 minutes; MS (ES<sup>+</sup>)  $m/z$  621 (M+H) <sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.93 (s, 1 H), 10.88 (br. s, 1 H), 9.71 (s, 1 H), 7.84 - 7.78 (m, 3 H), 7.77 (d, 2 H), 7.64 (d, 2 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.57 (dd, 1 H), 4.16 (s, 2 H), 4.12 (s, 2 H), 3.32 - 3.23 (obs. m, 1 H), 2.98 (ddd, 1 H), 2.56 - 2.45 (obs. m, 1 H), 2.24 - 2.16 (m, 1 H).

Example 478: methyl (2-{[4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]amino}-2-oxoethoxy)acetate

The compound prepared in Example 476 (100 mg) was treated as detailed in Example 474 to give the title compound having the following physical properties (17 mg).

LC/MS  $t_R$  3.10 minutes; MS (ES<sup>+</sup>)  $m/z$  601 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.54 (m, 5 H), 7.34 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.34 (s, 2 H), 4.22 (s, 2 H), 3.79 (s, 3 H), 3.51 - 3.38 (m, 1 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.57 - 2.31 (m, 1 H).

Example 479: methyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 266 (200 mg) was treated as detailed in Example 364 to give the title compound having the following physical properties (23 mg).

LC/MS  $t_R$  4.02 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.75 (s, 1 H), 9.75 (s, 1 H), 9.71 (s, 1 H), 7.88 - 7.76 (m, 3 H), 7.56 - 7.44 (m, 4 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.54 (d, 1 H), 3.68 (s, 3 H), 3.32 - 3.23 (m, 1 H), 3.05 - 2.93 (m, 1 H), 2.59 - 2.52 (m, 1 H), 2.23 - 2.12 (m, 1 H).

Example 480: methyl [4-(4-bromo-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

To a cooled (0 °C) dichloromethane (3 mL) solution of the compound prepared in Example 266 (200 mg) was added N-bromosuccinimide (67.3 mg) and the mixture stirred at 0 °C for 75 minutes. The reaction mixture was then diluted with dichloromethane and washed with water. The aqueous layer was extracted twice more with dichloromethane and the combined organic layers washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 to 10% methanol in dichloromethane) to give the title compound having the following physical properties (54.2 mg).

LC/MS  $t_R$  4.08 minutes; MS (ES<sup>+</sup>)  $m/z$  607 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.74 - 7.69 (m, 2 H), 7.69 - 7.65 (m,

1 H), 7.63 – 7.58 (m, 2 H), 7.54 – 7.47 (m, 2 H), 6.12 (s, 1 H), 6.07 (s, 1 H), 5.72 (dd, 1 H), 3.75 (s, 3 H), 3.49 – 3.36 (m, 1 H), 3.16 – 3.00 (m, 1 H), 2.70 – 2.54 (m, 1 H), 2.47 – 2.29 (m, 1 H).

5 Example 481: 2-methoxyethyl N-[4-(2-bromopropanoyl)phenyl]carbamate

The same operation as in Example 77 → Example 78 was conducted from 1(4-aminophenyl)propan-1-one to give the title compound having the following physical properties (Note: in the step corresponding to Example 77 in the operation, 2-methoxyethyl chloroformate was used).

10 LC/MS  $t_R$  1.91 minutes; MS (ES<sup>+</sup>)  $m/z$  330 and 332 (M+H)<sup>a</sup>.

Example 482: 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate

15 The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 481 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.77 - 7.73 (m, 2 H), 7.72 - 7.67 (m, 1 H), 7.49 (br. s, 2 H), 7.47 - 7.39 (m, 2 H), 6.12 (2 x s, 2 H), 5.74 (dd, 1 H), 4.34 - 4.24 (m, 2 H), 3.69 - 3.63 (m, 2 H), 3.46 (td, 1 H), 3.42 - 3.40 (m, 3 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.45 (br. s, 1 H), 2.33 (br. s, 3 H).

Example 483: 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate

25 The same operation as in Example 128 → Example 44 was conducted from the compound prepared in Example 456 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 128 in the operation, 2-methoxyethyl chloroformate was used).

LC/MS  $t_R$  4.07 minutes; MS (ES<sup>+</sup>)  $m/z$  607 (M+H)<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.77 - 7.63 (m, 3 H), 7.60 (d, 2 H), 7.50 (d, 2 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.71 (dd, 1 H), 4.33 - 4.21 (m, 2 H), 3.70 - 3.60 (m, 2 H), 3.53 - 3.35 (obs. m, 1 H), 3.39 (s, 3 H), 3.17 - 2.99 (m, 1 H), 2.72 - 2.53 (m, 1 H), 2.46 - 2.29 (m, 1 H).

Example 484: formic acid - 2-fluoroethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate (1:1)

5 To a 1,4-dioxane (1 mL) solution of the compound prepared in Example 456 (100 mg) was added N,N-diisopropylethylamine (106  $\mu$ L) followed by a solution of 2-fluoroethyl chloroformate (50  $\mu$ L) in 1,4-dioxane (1 mL) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated, the residue re-dissolved in methanol (2 mL) and concentrated ammonia solution (59  $\mu$ L) added. The  
10 mixture was stirred for 1 hour at room temperature then concentrated. The residue was suspended in water (25 mL) and extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 – 10% methanol in dichloromethane) followed by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] to give the title  
15 compound as the formic acid salt having the following physical properties (61.1 mg). LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  561 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.17 (s, 0.5 formate salt), 7.74 - 7.72 (m, 2 H), 7.69 - 7.67 (m, 1 H), 7.57 (d, 2 H), 7.45 (d, 2 H), 7.25 (s, 1 H), 6.13 (s, 1 H),  
20 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.71 - 4.66 (m, 1 H), 4.61 - 4.56 (m, 1 H), 4.42 - 4.38 (m, 1 H), 4.36 - 4.32 (m, 1 H), 3.48 - 3.41 (m, 1 H), 3.11 (ddd, 1 H), 2.68 - 2.60 (m, 1 H), 2.49 - 2.43 (m, 1 H).

Example 485: 2-ethoxyethyl chloroformate

25 To a cooled (0 °C) tetrahydrofuran (50 mL) solution of triphosgene (4.11 g) was added N,N-diisopropylethylamine (0.11 mL) and the mixture stirred for 5 minutes. A solution of 2-ethoxyethanol (2.68 mL) in tetrahydrofuran (15 mL) was added slowly and the mixture stirred at room temperature for 18 hours. The reaction mixture was concentrated and the residue dissolved in ethyl acetate (100 mL) and washed  
30 sequentially with 0.2 M hydrochloric acid and saturated saline, dried and concentrated to afford the title compound having the following physical properties (2.77 g).  
<sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>)  $\delta$  4.49 - 4.41 (m, 2 H), 3.74 - 3.67 (m, 2 H), 3.56 (q, 2 H), 1.23 (t, 3 H).

Example 486: 2-ethoxyethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 456 (100 mg) was treated with 2-ethoxyethyl chloroformate (100 mg) as detailed in Example 484 to give the title compound having the following physical properties (48.8 mg).

LC/MS  $t_R$  3.26 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.34 (s, 1 H), 7.75 - 7.69 (m, 2 H), 7.68 - 7.63 (m, 1 H), 7.55 (app. br. s, 2 H), 7.42 (d, 2 H), 7.22 (br. s, 1 H), 6.11 (s, 1 H), 6.06 (s, 1 H), 5.77 (dd, 1 H), 4.30 - 4.18 (m, 2 H), 3.72 - 3.64 (m, 2 H), 3.56 (q, 2 H), 3.43 (td, 1 H), 3.08 (ddd, 1 H), 2.61 (qd, 1 H), 2.45 (app. br. s, 1 H), 1.20 (t, 3 H).

Example 487: 3-methoxypropyl chloroformate

2-Methoxypropan-1-ol (2.49 g) was treated as detailed in Example 485 to give the title compound having the following physical properties (3.97 g).

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  4.43 (t, 2 H), 3.48 (t, 2 H), 3.35 (s, 3 H), 2.00 (quintet, 2 H).


Example 488: 3-methoxypropyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 456 (100 mg) was treated with 3-methoxypropyl chloroformate (81 mg) as detailed in Example 484 to give the title compound having the following physical properties (95 mg).

LC/MS  $t_R$  3.19 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.75 - 7.70 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.56 (app. br. s, 2 H), 7.44 (app. br. s, 2 H), 7.25 (br. s, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.77 (dd, 1 H), 4.20 (t, 2 H), 3.52 (t, 2 H), 3.49 - 3.38 (m, 1 H), 3.34 (s, 3 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.47 (app. br. s, 1 H), 1.94 (quintet, 2 H).

Example 489: 2-(2-methoxyethoxy)-ethyl chloroformate




2-(2-Methoxyethoxy)-ethan-1-ol (1 g) was treated as detailed in Example 485 to



give the title compound having the following physical properties (1.20 g).

$^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  4.52 - 4.41 (m, 2 H), 3.83 - 3.73 (m, 2 H), 3.70 - 3.62 (m, 2 H), 3.61 - 3.52 (m, 2 H), 3.39 (s, 3 H).

- 5 Example 490: formic acid - 2-(2-methoxyethoxy)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate (1:1)




- 10 The compound prepared in Example 456 (100 mg) was treated with 2-(2-methoxyethoxy)-ethyl chloroformate (96.9 mg) as detailed in Example 484 to give the title compound having the following physical properties (95 mg).

LC/MS  $t_R$  3.09 minutes; MS ( $\text{ES}^+$ )  $m/z$  617 ( $\text{M}+\text{H}$ )<sup>b</sup>

- $^1\text{H}$  NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.57 (d, 2 H), 7.44 (d, 2 H), 7.23 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H),  
15 4.30 - 4.24 (m, 2 H), 3.76 - 3.71 (m, 2 H), 3.68 - 3.64 (m, 2 H), 3.58 - 3.53 (m, 2 H), 3.49 - 3.39 (m, 1 H), 3.36 (s, 3 H), 3.10 (ddd, 1 H), 2.64 (qd, 1 H), 2.50 - 2.42 (m, 1 H).


Example 491: 2-(2-ethoxyethoxy)-ethyl chloroformate



- 20 2-(2-Ethoxyethoxy)-ethan-1-ol (1 g) was treated as detailed in Example 485 to give the title compound having the following physical properties (1.26 g).

$^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  4.48 - 4.39 (m, 2 H), 3.79 - 3.71 (m, 2 H), 3.67 - 3.60 (m, 2 H), 3.60 - 3.54 (m, 2 H), 3.51 (q, 2 H), 1.19 (t, 3 H).

- 25 Example 492: formic acid - 2-(2-ethoxyethoxy)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate (1:1)



The compound prepared in Example 456 (100 mg) was treated with the chloroformate prepared in 2-(2ethoxyethoxy)chloroformate (104 mg) as detailed in Example 484 to give the title compound having the following physical properties (95 mg).

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  631 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.57 (d, 2 H), 7.45 (d, 2 H), 7.23 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.30 - 4.26 (m, 2 H), 3.77 - 3.72 (m, 2 H), 3.68 - 3.64 (m, 2 H), 3.61 - 3.58 (m, 2 H), 3.53 (q, 2 H), 3.49 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.64 (qd, 1 H), 2.50 - 2.41 (m, 1 H), 1.18 (t, 3 H).

#### Example 493(1) to Example 493(5)

The compounds of the present invention having the following physical data were prepared from the compound prepared in Example 456 using the corresponding alcohols in the process of Example 485 → Example 484.

Example 493(1): (2S)-2-methoxypropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.16 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.34 (s, 1 H), 7.74 - 7.69 (m, 2 H), 7.68 - 7.64 (m, 1 H), 7.63 - 7.35 (m, 4 H), 7.34 - 7.00 (m, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.77 (dd, 1 H), 4.15 (dd, 1 H), 4.07 (dd, 1 H), 3.68 - 3.59 (m, 1 H), 3.40 (s, 3 H), 3.50 - 3.36 (m, 1 H), 3.09 (ddd, 1 H), 2.62 (qd, 1 H), 2.46 (app. br. s, 1 H), 1.20 (d, 3 H).

Example 493(2): 1-methoxy-2-propanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.20 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.56 (d, 2 H), 7.44 (d, 2 H), 7.22 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1

H), 5.05 - 4.98 (m, 1 H), 3.51 (dd, 1 H), 3.49 - 3.39 (m, 2 H), 3.38 (s, 3 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.47 (app. br. s, 1 H), 1.27 (d, 3 H).

Example 493(3): tetrahydro-3-furanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-

5 yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  585 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.56 (app. br. s, 2 H), 7.44 (app. br. s, 2 H), 7.23 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 5.31 (dd, 1 H), 3.97 - 3.83 (m, 4 H), 3.44 (td, 1 H), 3.10 (ddd, 1 H),  
10 2.63 (qd, 1 H), 2.46 (app. br. s, 1 H), 2.23 (dtd, 1 H), 2.14 - 2.04 (m, 1 H).

Example 493(4): formic acid - tetrahydro-3-furanylmethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate (1:1)

15 LC/MS  $t_R$  3.14 minutes; MS (ES<sup>+</sup>)  $m/z$  599 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.57 (d, 2 H), 7.45 (d, 2 H), 7.25 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.16 (dd, 1 H), 4.05 (dd, 1 H), 3.92 - 3.83 (m, 2 H), 3.76 (q, 1 H), 3.64 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.11 (ddd, 1 H), 2.70 - 2.59 (m, 2 H), 2.50 - 2.41 (m, 1 H), 2.14 - 2.05 (m,  
20 1 H), 1.72 (dt, 1 H).

Example 493(5): tetrahydro-2-furanylmethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.16 minutes; MS (ES<sup>+</sup>)  $m/z$  599 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (250 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 - 7.70 (m, 2 H), 7.70 - 7.64 (m, 1 H), 7.56 (app. br. s, 2 H), 7.46 (app. br. s, 2 H), 7.27 (br. s, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 4.23 - 4.02 (m, 3 H), 3.95 - 3.73 (m, 2 H), 3.44 (td, 1 H), 3.10 (ddd, 1 H), 2.73 - 2.55 (m, 1 H), 2.46 (app. br. s, 1 H), 2.13 - 1.84 (m, 3 H), 1.80 - 1.63 (m, 1 H).

30

Example 494: 3-oxetanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

To an acetonitrile (26 mL) suspension of N,N'-disuccinimidyl carbonate (2.05 g)

was added oxetan-3-ol (0.44 mL) and the mixture stirred at room temperature for 5 minutes. To the reaction mixture, N,N-diisopropylethylamine (3.3 mL) was added and the mixture stirred at room temperature for 20 hours to form a 0.25 M solution of the mixed carbonate product.

- 5 To a 2.5 M solution in acetonitrile (4 mL) of the above prepared mixed carbonate, the compound prepared in Example 456 (100 mg) and N,N-diisopropylethylamine (105  $\mu$ L) were added sequentially and the mixture stirred at room temperature for 18 hours. On concentration, the residue was dissolved in methanol (2 mL) whereupon concentrated aqueous ammonia (60  $\mu$ L) was added and the mixture stirred at room temperature for 2
- 10 hours. The reaction mixture was concentrated, the residue suspended in saturated sodium carbonate solution (30 mL) and extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in water) to afford the title compound having the following physical properties (72 mg).
- 15 LC/MS  $t_R$  2.98 minutes; MS ( $ES^+$ )  $m/z$  571 (M+H) <sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.34 (s, 1 H), 7.74 - 7.69 (m, 2 H), 7.68 - 7.64 (m, 1 H), 7.56 (app. br. s, 2 H), 7.43 (d, 2 H), 7.23 (br. s, 1 H), 6.12 (s, 1 H), 6.07 (s, 1 H), 5.77 (dd, 1 H), 5.51 - 5.42 (m, 1 H), 4.92 (t, 2 H), 4.68 (dd, 2 H), 3.49 - 3.36 (m, 1 H), 3.09 (ddd, 1 H), 2.62 (q, 1 H), 2.45 (app. br. s, 1 H).

20

#### Example 495(1) to Example 495(3)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 456 and the corresponding alcohols using the method as detailed in Example 494.

25

Example 495(1): 2-(dimethylamino)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  2.50 minutes; MS ( $ES^+$ )  $m/z$  586 (M+H), 293 (M+2H)/2 <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.76 - 7.69 (m, 2 H), 7.69 - 7.64 (m, 1 H), 7.56 (app. br. s, 2 H), 7.45 (app. br. s, 2 H), 7.23 (br. s, 1 H), 6.12 (s, 1 H), 6.07 (s, 1 H), 5.77 (dd, 1 H), 4.26 (t, 2 H), 3.50 - 3.37 (m, 1 H), 3.09 (ddd, 1 H), 2.67 (t, 2 H), 2.65 - 2.56 (m, 1 H), 2.55 - 2.38 (m, 1 H), 2.33 (s, 6 H).

30

Example 495(2): 2-(oxan-2-yloxy)ethyl N-(4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}phenyl)carbamate

LC/MS  $t_R$  1.70 minutes; MS ( $ES^+$ )  $m/z$  643 (M+H), 559 (M-C<sub>5</sub>H<sub>7</sub>O+2H)/2 <sup>a</sup>.

5

Example 495(3): 3-hydroxy-3-methylbutyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.08 minutes; MS ( $ES^+$ )  $m/z$  601 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.75 - 7.70 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.65 - 7.36 (m, 4 H), 7.35 - 7.00 (m, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.77 (d, 1 H), 4.28 (t, 2 H), 3.51 - 3.35 (m, 1 H), 3.10 (ddd, 1 H), 2.66 - 2.57 (m, 1 H), 2.57 - 2.28 (m, 1 H), 1.88 (t, 2 H), 1.26 (s, 6 H).

10

Example 496: 2-hydroxyethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]carbamate

15

To a 1,4-dioxane (4 mL) solution of the compound prepared in Example 495(2) (137 mg) was added 1 M hydrochloric acid (2.1 mL) and the mixture stirred at room temperature for 4 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (25 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in water) to afford the title compound having the following physical properties (24 mg).

20

LC/MS  $t_R$  2.84 minutes; MS ( $ES^+$ )  $m/z$  559 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.04 (br. s, 1 H), 9.68 (s, 1 H), 9.63 (br. s, 1 H), 7.83 - 7.75 (m, 3 H), 7.59 (d, 2 H), 7.42 (d, 2 H), 7.39 (s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.61 (d, 1 H), 4.81 (t, 1 H), 4.09 (t, 2 H), 3.61 (q, 2 H), 3.44 - 3.37 (m, 1 H), 2.99 (dd, 1 H), 2.56 - 2.43 (obs. m, 1 H), 2.38 - 2.29 (m, 1 H).

25

Example 497: 2-hydroxy-2-methylpropyl N-(4-acetylphenyl)carbamate

30

To a dichloromethane (62 mL) solution of 4-acetylphenyl isocyanate (1 g) and 4-dimethylaminopyridine (76 mg) was added 2-methylpropane-1,2-diol [patent US2010/249087] (0.58 g) and the mixture stirred at room temperature for 3 hours. The reaction mixture was concentrated and the residue purified by column chromatography

(0 to 100% EtOAc in heptanes) to give the title compound having the following physical properties (0.92 g).

LC/MS  $t_R$  1.49 minutes; MS (ES<sup>+</sup>)  $m/z$  252 (M+H)<sup>a</sup>.

5    Example 498: 2-hydroxy-2-methylpropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 204 → Example 51 → Example 52 was conducted from the compound prepared in Example 497 to give the title compound having the following physical properties.

10    LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  587 (M+H)<sup>b</sup>  
      <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.76 - 7.69 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.64 - 7.36 (m, 4 H), 7.36 - 7.01 (m, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 3.99 (s, 2 H), 3.51 - 3.35 (m, 1 H), 3.09 (ddd, 1 H), 2.68 - 2.56 (m, 1 H), 2.56 - 2.28 (m, 1 H), 1.26 (s, 6 H).

15

Example 499: 2-hydroxy-2-methylpropyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

20    The compound prepared in Example 498 (104 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (59.7 mg).

LC/MS  $t_R$  3.94 minutes; MS (ES<sup>+</sup>)  $m/z$  621 (M+H)<sup>b</sup>  
      <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.75 - 7.70 (m, 2 H), 7.69 - 7.65 (m, 1 H), 7.61 (d, 2 H), 7.53 (d, 2 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.71 (dd, 1 H), 4.00 (s, 2 H),  
25    3.44 (td, 1 H), 3.09 (ddd, 1 H), 2.63 (qd, 1 H), 2.43 - 2.32 (m, 1 H), 1.27 (s, 6 H).

Example 500: 3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]-1,1-dimethylurea

30    The compound prepared in Example 456 (400 mg) was treated as detailed in Example 132 to give the title compound having the following physical properties (128 mg).

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  542 (M+H)<sup>b</sup>  
      <sup>1</sup>H NMR (250 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.77 - 7.65 (m, 3 H), 7.56 (d, 2 H),

7.39 (d, 2 H), 7.22 (s, 1 H), 6.12 (s, 1 H), 6.09 (d, 1 H), 5.78 (dd, 1 H), 3.53 - 3.36 (m, 1 H), 3.18 - 3.07 (m, 1 H), 3.02 (s, 6 H), 2.70 - 2.54 (m, 1 H), 2.53 - 2.36 (m, 1 H).

Example 501: 3-[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]-1,1-dimethylurea

The compound prepared in Example 500 (48 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (28.6 mg).

LC/MS  $t_R$  3.85 minutes; MS (ES<sup>+</sup>)  $m/z$  576 and 578 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.73 (app. s, 1 H), 7.72 (dd, 1 H), 7.68 (d, 1 H), 7.59 (d, 2 H), 7.48 (d, 2 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.70 (dd, 1 H), 3.43 (td, 1 H), 3.09 (ddd, 1 H), 3.03 (s, 6 H), 2.68 - 2.58 (m, 1 H), 2.43 - 2.33 (m, 1 H).

Example 502: formic acid - 3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]-1-methoxy-1-methylurea (1:1)

The compound prepared in Example 456 (100 mg) was treated as detailed in Example 484 using N-methoxy-N-methylcarbamoyl chloride [Chem. Pharm. Bull., 55(2), 328 (2007)] in place of 2-fluoroethyl chloroformate to give the title compound having the following physical properties (13.9 mg).

LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  558 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.59 (d, 2 H), 7.51 (d, 2 H), 7.25 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.76 (s, 3 H), 3.44 (td, 1 H), 3.15 (s, 3 H), 3.10 (ddd, 1 H), 2.69 - 2.58 (m, 1 H), 2.51 - 2.41 (m, 1 H).

Example 503: 3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]-1-(2-methoxyethyl)-1-methylurea

The compound prepared in Example 456 (50 mg) was treated as detailed in Example 484 using N-(2-methoxyethyl)-N-methylcarbamoyl chloride [Chem. Pharm. Bull., 55(2), 328 (2007)] in place of 2-fluoroethyl chloroformate to give the title compound having the following physical properties (10.4 mg).

LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  586 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.56 (d, 2 H), 7.35 (d, 2 H), 7.22 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.63 - 3.59 (m, 2 H), 3.58 - 3.54 (m, 2 H), 3.49 - 3.39 (m, 1 H), 3.43 (s, 3 H), 3.10 (ddd, 1 H), 3.05 (s, 3 H), 2.69 - 2.58 (m, 1 H), 2.50 - 2.42 (m, 1 H).

5

Example 504: 3-[4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]-1-(2-methoxyethyl)-1-methylurea

The compound prepared in Example 503 (43.5 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (17.4 mg).

LC/MS *t<sub>R</sub>* 4.01 minutes; MS (ES<sup>+</sup>) *m/z* 620 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 7.77 - 7.71 (m, 2 H), 7.68 (dd, 1 H), 7.61 (d, 2 H), 7.43 (d, 2 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.71 (dd, 1 H), 3.64 - 3.60 (m, 2 H), 3.59 - 3.55 (m, 2 H), 3.50 - 3.38 (m, 1 H), 3.43 (s, 3 H), 3.14 - 3.07 (m, 1 H), 3.06 (s, 3 H), 2.69 - 2.59 (m, 1 H), 2.43 - 2.34 (m, 1 H).

15

Example 505: 1-(1,2,3,4-tetrahydroquinolin-1-yl)ethan-1-one

To a dichloromethane (25 mL) solution of 1,2,3,4-tetrahydroquinoline (1.89 mL) was added N,N-diisopropylethylamine (3.12 mL). To the cooled (0 °C) reaction mixture, acetyl chloride (1.18 mL) was added and the mixture stirred at room temperature for 2 hours. To the reaction mixture, dichloromethane was added and the organic layer washed with water. The aqueous layer was extracted with dichloromethane and the combined organic layers washed with saturated saline, dried and concentrated to give the title compound having the following physical properties (2.97 g).

25

LC/MS *t<sub>R</sub>* 1.66 minutes; MS (ES<sup>+</sup>) *m/z* 176 (M+H)<sup>a</sup>.

Example 506: 1-(1-acetyl-1,2,3,4-tetrahydroquinolin-6-yl)-2-bromoethan-1-one

To a cooled (0 °C) 1,2-dichloroethane (15 mL) solution of aluminum trichloride (5.65 g) was added bromoacetyl bromide (2.96 mL) and the mixture was stirred at 0 °C for 30 minutes. The compound prepared in Example 505 (2.97 g) was added and the mixture stirred at room temperature for 2 hours. On concentration, the residue was suspended in water followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated to obtain the title

30



compound having the following physical properties (6.17 g).

LC/MS  $t_R$  1.75 minutes; MS ( $ES^+$ )  $m/z$  296 (M+H) <sup>a</sup>.

Example 507: 2-chloro-1-(1,2,3,4-tetrahydroquinolin-6-yl)ethan-1-one hydrochloride

5        The compound prepared in Example 506 (6.17 g) was suspended in concentrated hydrochloric acid (100 mL) and the mixture stirred at 80 °C for 2 hours. On concentration, the residue was triturated with a 1:1 mixture of ethyl acetate and dichloromethane, the resultant precipitate being collected by filtration to give the title compound having the following physical properties (1.08 g).

10      LC/MS  $t_R$  1.78 minutes; MS ( $ES^+$ )  $m/z$  210 (M+H) <sup>a</sup>.

Example 508: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1,2,3,4-tetrahydro-6-quinolinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

15        The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 507 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.05 minutes; MS ( $ES^+$ )  $m/z$  511 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.76 (s, 1 H), 9.69 (s, 1 H), 7.95 – 7.58 (m, 3 H), 7.23 – 7.06 (m, 3 H), 6.38 (d, 1 H), 6.00 – 5.90 (m, 2 H), 5.64 – 5.51 (m, 2 H), 3.52 –  
20      3.48 (m, 1 H), 3.22 – 3.11 (m, 2 H), 3.04 – 2.91 (m, 1 H), 2.71 – 2.66 (m, 2 H), 2.51 – 2.48 (m, 1 H), 2.38 – 2.31 (m, 1 H), 1.87 – 1.74 (m, 2 H).

Example 509: 6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-methyl-1H-imidazol-5-yl)-3-methyl-3,4-dihydro-2(1H)-quinazolinone

25        The same operation as in Example 506 → Example 51 → Example 52 was conducted from 3-methyl-3,4-dihydroquinazolin-2(1H)-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 506 in the operation, 2-bromopropanoyl bromide was used).

30      LC/MS  $t_R$  2.93 minutes; MS ( $ES^+$ )  $m/z$  554 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.86 (s, 1 H), 9.69 (s, 1 H), 9.15 (s, 1 H), 7.92 – 7.67 (m, 3 H), 7.36 – 7.12 (m, 2 H), 6.76 (d, 1 H), 5.96 (s, 1 H), 5.94 (s, 1 H), 5.63 – 5.46 (m, 1 H), 4.44 (s, 2 H), 3.43 – 3.36 (m, 1 H), 3.02 – 2.92 (m, 1 H), 2.87 (s, 3 H), 2.48 – 2.42 (m, 1 H), 2.30 (s, 3 H), 2.24 – 2.16 (m, 1 H).

Example 510: 2-amino-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)benzoic acid

The same operation as in Example 448 → Example 449 → Example 51 →

- 5 Example 52 was conducted from 6-bromo-2H-3,1-benzoxazine-2,4(1H)-dione to give the title compound having the following physical properties. (Note: in the step corresponding to Example 52 in the operation, the deprotection of the amino acid is also accomplished).


LC/MS  $t_R$  2.80 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>a</sup>

- 10 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.85 (s, 1 H), 7.78 - 7.66 (m, 3 H), 7.50 (d, 1 H), 7.18 (s, 1 H), 6.79 (d, 1 H), 6.12 (d, 2 H), 5.79 (dd, 1 H), 3.52 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.50 - 2.44 (m, 1 H).

Example 511: 6-acetyl-2,2-dimethyl-2,4-dihydro-1,3-benzodioxin-4-one

- 15 To a solution of 5-acetyl-2-hydroxybenzoic acid (500 mg) in trifluoroacetic acid (4 ml) and acetone (1.2 ml) was added trifluoroacetic anhydride (1.17 ml) and the mixture heated at 90 °C for 16 h. The reaction mixture was concentrated, the residue dissolved in ethyl acetate and washed sequentially with a saturated solution of sodium hydrogen carbonate and saturated saline, dried and concentrated. The residue was
- 20 purified by column chromatography (0 - 60% ethyl acetate in heptanes) to afford the title compound having the following physical properties (432 mg).
- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.55 (d, 1 H), 8.27 (dd, 1 H), 7.17 (d, 1 H), 2.63 (s, 3 H), 1.77 (s, 6 H).

- 25 Example 512: 6-(2-bromoacetyl)-2,2-dimethyl-2,4-dihydro-1,3-benzodioxin-4-one



- To a dichloromethane (6 mL) solution of the compound prepared in Example 511 (362 mg) was added acetic acid (0.2 ml) followed by bromine (93  $\mu$ l) and the mixture stirred at room temperature for 2 hrs. The reaction mixture was concentrated to give the
- 30 crude title compound having the following physical properties (534 mg).

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.52 - 8.45 (m, 1 H), 8.21 (dd, 1 H), 7.11 (d, 1 H), 4.58 (s, 2 H), 1.73 - 1.64 (s, 6 H).

5 Example 513: (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-3-[5-(2,2-dimethyl-4-oxo-2,4-dihydro-1,3-benzodioxin-6-yl)-1H-imidazol-2-yl]-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 512 to give the title compound having the following physical properties.

10 LC/MS *t<sub>R</sub>* 1.73 minutes; MS (ES<sup>+</sup>) *m/z* 556 (M+H)<sup>a</sup>.

Example 514: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-hydroxybenzoic acid

To a 1,4-dioxane (2 mL) solution of the compound prepared in Example 513 (100  
15 mg) was added a 4 M solution of hydrogen chloride in 1,4-dioxane (2 mL) followed by water (1 mL) and the mixture heated at 60 °C for 3 hours. On cooling to room temperature, the reaction mixture was diluted with ethyl acetate (30 mL) and washed with a saturated aqueous solution of sodium hydrogen carbonate. The combined  
aqueous layers were treated with concentrated hydrochloric acid until pH 2 – 3 was  
20 attained, extracted once into ethyl acetate then washed twice more with a 1: 3 mixture of propan-2-ol and chloroform. The combined organic phases were dried and concentrated and the residue triturated with dichloromethane. The resultant precipitate was collected by filtration to give the title compound having the following physical properties (29.6  
mg).

25 LC/MS *t<sub>R</sub>* 3.06 minutes; MS (ES<sup>+</sup>) *m/z* 516 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.43 (s, 1 H), 8.29 (d, 1 H), 7.89 - 7.82 (m, 2 H), 7.81 - 7.68 (m, 3 H), 7.12 (d, 1 H), 6.21 (s, 1 H), 6.19 (s, 1 H), 5.91 (dd, 1 H), 3.43 - 3.36 (m, 1 H), 3.31 - 3.22 (m, 1 H), 2.94 - 2.81 (m, 1 H), 2.51 - 2.39 (m, 1 H).

30 Example 515: 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-hydroxybenzoic acid

The same operation as in Example 338 → Example 514 was conducted from the compound prepared in Example 513 to give the title compound having the following

physical properties.

LC/MS  $t_R$  3.92 minutes; MS (ES<sup>+</sup>)  $m/z$  550 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.25 - 8.17 (m, 1 H), 7.81 - 7.64 (m, 4 H), 7.01 - 6.95 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.73 (dd, 1 H), 3.50 - 3.41 (m, 1 H), 3.11 (ddd, 1 H), 2.66 - 2.60 (m, 1 H), 2.43 - 2.33 (m, 1 H).

Example 516: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-methylbenzoic acid

The same operation as in Example 139 → Example 78 → Example 51 (with example 11) → Example 52 → Example 8 → Example 55 → Example 24 was conducted from methyl 5-bromo-2-methylbenzoate to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 11 was used).

LC/MS  $t_R$  3.15 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.20 (m, 1 H), 7.79 - 7.65 (m, 4 H), 7.35 (s, 1 H), 7.30 (d, 1 H), 6.14 (s, 1 H), 6.14 (s, 1 H), 5.81 (dd, 1 H), 3.55 - 3.41 (m, 1 H), 3.19 - 3.08 (m, 1 H), 2.74 - 2.62 (m, 1 H), 2.58 (s, 3 H), 2.53 - 2.45 (m, 1 H).

Example 517: 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-methylbenzoic acid

The compound prepared in Example 516 (90 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (26 mg).

LC/MS  $t_R$  4.00 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.39 (s, 1 H), 8.26 (s, 1 H), 7.82 - 7.67 (m, 4 H), 7.39 (d, 1 H), 6.18 (s, 1 H), 6.08 (s, 1 H), 5.74 (dd, 1 H), 3.51 - 3.40 (m, 1 H), 3.17 - 3.07 (m, 1 H), 2.67 (m, 1 H), 2.62 (s, 3 H), 2.40 (d, 1 H).

Example 518: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-{5-[4-(methylsulfonyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

To a dichloromethane (1 mL) solution of the compound prepared in Example 173(17) (50 mg) was added a solution of *meta*-chloroperbenzoic acid (45 mg, 77% in water) in dichloromethane (1 mL) and the mixture stirred at room temperature for 45

minutes. To the reaction mixture, water (10 mL) was added followed by extraction into dichloromethane. The combined organic layers were washed with saturated sodium hydrogen carbonate, saturated saline, dried and concentrated. The residue was purified by column chromatography (0 - 100% ethyl acetate in heptanes, then 0 - 10% methanol in ethyl acetate) to give the title compound having the following physical properties (53 mg).

LC/MS  $t_R$  3.33 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.37 (br. s, 1 H), 9.69 (s, 1 H), 7.94 (d, 2 H), 7.86 (d, 2 H), 7.82 - 7.74 (m, 4 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.64 (dd, 1 H), 3.43 - 3.27 (obs. m, 1 H), 3.19 (s, 3 H), 3.01 (dd 1 H), 2.55 - 2.45 (obs. m, 1 H), 2.38 - 2.29 (m, 1 H).

Example 519: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{4-[(S)-methylsulfinyl]phenyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

To a dichloromethane (8 mL) solution of titanium(IV) isopropoxide (0.3 mL) and diethyl (2S,3S)-2,3-dihydroxybutanedioate (0.34 mL) was added water (18  $\mu$ L) and the mixture stirred at room temperature for 20 minutes. A dichloromethane (2 mL) solution of the compound prepared in Example 173(17) (0.50 g) was added and the reaction mixture cooled to -20 °C. A 2.84 M solution of tert-butyl hydroperoxide in toluene (0.39 mL) [Org. Synth. Coll., 7, 461, 1990] was added and the reaction stood at -18 °C for 20 hours. To the chilled (-20 °C) reaction mixture, water (0.18 mL) was added and the mixture allowed to warm to room temperature over 1.5 hours. Alumina was added and the solids removed by filtration through Celite®, washing the filter cake with dichloromethane. The filtrate was then washed with a saturated aqueous solution of sodium sulphite (30 mL) then saturated saline, dried and concentrated. The residue was purified by column chromatography (50 - 100% ethyl acetate in heptanes, then 0 - 20% methanol in ethyl acetate) to give the title compound having the following physical properties (321 mg).

LC/MS  $t_R$  1.81 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>e</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.32 (br. s, 1 H), 9.69 (s, 1 H), 7.88 (d, 2 H), 7.83 - 7.75 (m, 3 H), 7.74 - 7.55 (m, 3 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.64 (dd, 1 H), 3.42 - 3.36 (obs. m, 1 H), 3.01 (ddd 1 H), 2.73 (s, 3 H), 2.57 - 2.51 (obs. m, 1 H), 2.38 - 2.28 (m, 1 H).

Example 520: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{4-[(R)-methylsulfinyl]phenyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 173(17) (0.50 g) was treated as detailed in  
5 Example 519 using diethyl (2R,3R)-2,3-dihydroxybutanedioate in place of diethyl (2S,3S)-2,3-dihydroxybutanedioate to give the title compound having the following physical properties (350 mg).

LC/MS  $t_R$  1.81 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>e</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.32 (br. s, 1 H), 9.69 (s, 1 H), 7.88 (d, 2 H), 7.84 -  
10 7.76 (m, 3 H), 7.74 - 7.57 (m, 3 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.65 (dd, 1 H), 3.42 -  
3.37 (obs. m, 1 H), 3.01 (dd, 1 H), 2.73 (s, 3 H), 2.57 - 2.51 (obs. m, 1 H), 2.38 - 2.27  
(m, 1 H).

Example 521: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(S-methylsulfonylimidoyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

To a dichloromethane (5 mL) solution of the compound prepared in Example  
519 (261 mg), rhodium(II)acetate dimer (5.7 mg), magnesium oxide (81 mg), 2,2,2-  
trifluoroacetamide (114 mg) and (diacetoxyiodo)benzene (243 mg) were added  
sequentially and the mixture stirred at room temperature for 16 hours. To the reaction  
20 mixture, dichloromethane (30 mL) was added and the solids removed by filtration  
through Celite®, washing the filter cake with dichloromethane. The filtrate was  
concentrated and the residue dissolved in methanol (5 mL) whereupon potassium  
carbonate (348 mg) was added and the mixture stirred at room temperature for 2 hours.  
To the reaction mixture, dichloromethane (30 mL) was added and the solids removed by  
25 filtration through Celite®, washing the filter cake with dichloromethane. The filtrate  
was concentrated and the residue purified by column chromatography (0% to 10%  
methanol in dichloromethane) followed by high performance liquid chromatography (5  
to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title  
compound having the following physical properties (29.2 mg).

30 LC/MS  $t_R$  3.49 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>g</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 8.03 (d, 2 H), 7.98 (d, 2 H), 7.76 -  
7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.64 (br. s, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.80  
(dd, 1 H), 3.46 (td, 1 H), 3.33 (s, 3 H), 3.14 (ddd, 1 H), 2.72 - 2.61 (m, 1 H), 2.49 (tt, 1

H).

Example 522: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(S-methylsulfonimidoyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

5           The compound prepared in Example 520 (296 mg) was treated as detailed in Example 521 to give the title compound having the following physical properties (3.4 mg).

LC/MS  $t_R$  3.47 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>g</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.36 (s, 1 H), 8.06 - 7.84 (m, 4 H), 7.76 - 7.71 (m, 2  
10 H), 7.71 - 7.66 (m, 1 H), 7.59 (br. s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.79 (dd, 1 H),  
3.47 (td, 1 H), 3.16 (s, 3 H), 3.12 (ddd, 1 H), 2.70 - 2.60 (m, 1 H), 2.51 (app. br. s, 1 H).

Example 523: 1-(6-amino-2-fluoropyridin-3-yl)ethan-1-one

2-amino-6-fluoro-5-iodopyridine [J. Org. Chem. 71(8), 2922 (2006)] (1.5 g) was  
15 treated as detailed in Example 90 to give the title compound having the following physical properties (0.46 g).

LC/MS  $t_R$  1.19 minutes; MS (ES<sup>+</sup>)  $m/z$  155 (M+H)<sup>a</sup>.

Example 524: 1-(6-amino-2-methoxypyridin-3-yl)ethan-1-one

20           A methanol (3 mL) suspension of the compound prepared in Example 523 (150 mg) and potassium tert-butoxide (546 mg) was heated at 60 °C for 3 hours. On cooling to room temperature, the reaction mixture was diluted with water (20 mL) and extracted into ethyl acetate. The combined organic layers were dried and concentrated to give the title compound having the following properties (138 mg).

25           LC/MS  $t_R$  1.35 minutes; MS (ES<sup>+</sup>)  $m/z$  167 (M+H)<sup>a</sup>.

Example 525: (3S)-3-[5-(6-amino-2-methoxy-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

30           The same operation as in Example 204 → Example 51 → Example 52 was conducted from the compound prepared in Example 524 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.99 minutes; MS (ES<sup>+</sup>)  $m/z$  502 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.79 - 7.60 (m, 4 H), 7.12 (br. s, 1 H),

6.21 - 6.15 (m, 2 H), 6.12 - 6.07 (m, 1 H), 5.86 (br. s, 1 H), 3.99 (br. s, 3 H), 3.48 - 3.37 (m, 1 H), 3.16 - 3.08 (m, 1 H), 2.66 - 2.55 (m, 2 H).

Example 526: (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 216 (94 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (38 mg). LC/MS  $t_R$  3.75 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.78 - 7.71 (m, 3 H), 7.70 - 7.66 (m, 1 H), 6.46 (dd, 1 H), 6.14 (s, 1 H), 6.07 (s, 1 H), 5.72 (dd, 1 H), 3.46 - 3.37 (m, 1 H), 3.13 - 3.05 (m, 1 H), 2.67 - 2.57 (m, 1 H), 2.45 - 2.38 (m, 1 H).

Example 527: methyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-6-fluoro-2-pyridinyl]carbamate

The compound prepared in Example 216 (60 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (24 mg).

LC/MS  $t_R$  3.54 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 8.30 (br. s, 1 H), 7.82 (d, 1 H), 7.76 - 7.70 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.32 (br. s, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.80 (dd, 1 H), 3.77 (s, 3 H), 3.51 - 3.40 (m, 1 H), 3.11 (dd, 1 H), 2.68 - 2.58 (m, 1 H), 2.55 - 2.47 (m, 1 H).

Example 528: N-[6-chloro-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinyl]acetamide

To a dichloromethane (5 mL) solution of the compound prepared in Example 210 (150 mg) was added triethylamine (83  $\mu$ L) followed by acetyl chloride (42  $\mu$ L) and the mixture stirred at room temperature for 2 hours. To the reaction mixture, further triethylamine (83  $\mu$ L) and acetyl chloride (42  $\mu$ L) was added and the mixture stirred at room temperature for 16 hours. At this juncture, an additional aliquot of acetyl chloride (42  $\mu$ L) was added and the mixture stirred at room temperature for 16 hours. A final aliquot of acetyl chloride (42  $\mu$ L) was then added and the mixture stirred at room temperature for 72 hours.



To the reaction mixture, water (10 mL) was added followed by extraction with dichloromethane. The combined organic layers were dried and concentrated and the residue dissolved in methanol (10 mL). To this methanolic solution, concentrated ammonia solution (0.10 mL) was added and the mixture stirred at room temperature for 1 hour. On concentration, the residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in 2 mM aqueous ammonium hydrogen carbonate) to give the title compound having the following physical properties (28 mg).

LC/MS  $t_R$  3.41 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 8.26 (br. s, 1 H), 8.11 (d, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.61 (br. s, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.81 (d, 1 H), 3.51 - 3.42 (m, 1 H), 3.11 (dd, 1 H), 2.68 - 2.59 (m, 1 H), 2.57 - 2.49 (m, 1 H), 2.16 (s, 3 H).

Example 529: 2-methoxyethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The same operation as in Example 51 → Example 52 → Example 128 was conducted from the compound prepared in Example 193 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 128 in the operation, 2-methoxyethyl chloroformate was used).

LC/MS  $t_R$  3.19 minutes; MS (ES<sup>+</sup>)  $m/z$  574 (M+H), 288 (M/2+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.56 (br. s, 1 H), 8.01 (br. s, 1 H), 7.88 (d, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.40 (br. s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 4.33 - 4.28 (m, 2 H), 3.68 - 3.64 (m, 2 H), 3.50 - 3.41 (m, 1 H), 3.39 (s, 3 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.49 (br. s, 1 H).

Example 530: 2-methoxyethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 529 (71 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (27 mg).

LC/MS  $t_R$  3.93 minutes; MS (ES<sup>+</sup>)  $m/z$  608 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.57 (s, 1 H), 8.03 (dd, 1 H), 7.98 (d, 1 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.71 (dd, 1

H), 4.34 - 4.30 (m, 2 H), 3.69 - 3.64 (m, 2 H), 3.49 - 3.41 (m, 1 H), 3.39 (s, 3 H), 3.11 (ddd, 1 H), 2.70 - 2.60 (m, 1 H), 2.43 - 2.35 (m, 1 H).

5 Example 531: 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The compound prepared in Example 229 (160 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (133 mg).

LC/MS  $t_R$  3.15 minutes; MS (ES<sup>+</sup>)  $m/z$  574 (M+H)<sup>b</sup>

10 <sup>1</sup>H NMR (250 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.56 (d, 1 H), 7.94 (dd, 1 H), 7.85 - 7.62 (m, 4 H), 7.49 (br. s, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.80 (dd, 1 H), 4.35 - 4.24 (m, 2 H), 3.72 - 3.60 (m, 2 H), 3.56 - 3.42 (m, 1 H), 3.39 (s, 3 H), 3.19 - 2.99 (m, 1 H), 2.75 - 2.38 (m, 2 H).

15 Example 532: 2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The compound prepared in Example 531 (89 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (94 mg).

20 LC/MS  $t_R$  4.00 minutes; MS (ES<sup>+</sup>)  $m/z$  608 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.95 (br. s, 1 H), 10.10 (br. s, 1 H), 9.70 (s, 1 H), 8.74 (d, 1 H), 7.96 (dd, 1 H), 7.89 (d, 1 H), 7.84 - 7.77 (m, 3 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.67 (dd, 1 H), 4.27 - 4.21 (m, 2 H), 3.60 - 3.56 (m, 2 H), 3.34 (s, 3 H), 3.27 - 3.18 (m, 1 H), 2.97 (ddd, 1 H), 2.54 - 2.43 (obs. m, 1 H), 2.24 - 2.12 (m, 1 H).

25

Example 533: tert-butyl N-(5-acetyl-4-methylpyridin-2-yl)carbamate

tert-Butyl-(5-bromo-4-methylpyridin-2-yl)-N-[(tert-butoxy)carbonyl]carbamate [Bioorg. Med. Chem., 12(5), 1151 (2004)] (3.23 g) was treated as detailed in Example 90 to give the title compound having the following physical properties (0.22 g).

30 LC/MS  $t_R$  1.93 minutes; MS (ES<sup>+</sup>)  $m/z$  251 (M+H), 195 (M-t-Bu+H)<sup>a</sup>.

Example 534: tert-butyl N-[5-(2-bromoacetyl)-4-methylpyridin-2-yl]carbamate hydrobromide

To a tetrahydrofuran solution (5 mL) of the compound prepared in Example 533 (236 mg) was added a 33% weight solution of hydrogen bromide in acetic acid (0.23 mL) followed by bromine (23  $\mu$ L) and the mixture stirred at room temperature for 1 hour. Concentration of the reaction mixture gave the crude title compound having the following physical properties (0.28 g).  
LC/MS  $t_R$  2.14 minutes; MS ( $ES^+$ )  $m/z$  329 and 331 ( $M+H$ ), 273 and 275 ( $M-t-Bu+H$ )<sup>a</sup>.

Example 535: tert-butyl N-(5-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}-4-methylpyridin-2-yl)carbamate

The same operation as in Example 51  $\rightarrow$  Example 52 was conducted from the compound prepared in Example 534 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.71 minutes; MS ( $ES^+$ )  $m/z$  586 ( $M+H$ ), 265.5 ( $M/2+H$ )<sup>a</sup>.

Example 536: (3S)-3-[5-(6-amino-4-methyl-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 535 (35 mg) was treated as detailed in Example 55 to give the title compound having the following physical properties (15 mg).

LC/MS  $t_R$  2.73 minutes; MS ( $ES^+$ )  $m/z$  286 ( $M+H$ ), 244 ( $M-t-Bu+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.91 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.00 (br. s, 1 H), 6.49 (s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.78 (dd, 1 H), 3.49 - 3.38 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.57 (m, 1 H), 2.54 - 2.43 (m, 1 H), 2.26 (s, 3 H).

Example 537: (3S)-3-[5-(6-amino-4-methyl-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 44  $\rightarrow$  Example 55 was conducted from the compound prepared in Example 535 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.97 minutes; MS ( $ES^+$ )  $m/z$  520 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.78 (s, 1 H), 7.75 - 7.72 (m, 2 H), 7.70 - 7.66 (m, 1 H), 6.52 (s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.68 (dd, 1 H), 3.49 -

3.40 (m, 1 H), 3.10 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.47 - 2.38 (m, 1 H), 2.16 (s, 3 H).

Example 538: 1-(6-aminopyridazin-3-yl)ethan-1-one

A solution of 1-(6-chloropyridazin-3-yl)ethan-1-one [Bioorg. Med. Chem.,  
5 13(11), 3705 (2005)] (1.30 g) in concentrated aqueous ammonia (1.5 mL) was heated  
for 50 minutes at 120 °C by microwave irradiation. The reaction mixture was then  
extracted into ethyl acetate. The combined organic layers were dried and concentrated  
and the residue purified by column chromatography (0 – 10% methanol in  
dichloromethane) to give the title compound having the following physical properties  
10 (0.42 g).

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 7.85 (d, 1 H), 6.93 (d, 1 H), 2.67 (s, 3 H).

Example 539: (3S)-3-[5-(6-amino-3-pyridazinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

15 The same operation as in Example 204 → Example 51 → Example 52 was  
conducted from the compound prepared in Example 538 to give the title compound  
having the following physical properties.

LC/MS *t*<sub>R</sub> 2.79 minutes; MS (ES<sup>+</sup>) *m/z* 473 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.83 - 7.71 (m, 3 H), 7.70 - 7.66 (m, 1  
20 H), 7.56 (br. s, 1 H), 6.97 (d, 1 H), 6.15 (s, 1 H), 6.08 (s, 1 H), 5.81 (d, 1 H), 3.50 - 3.37  
(m, 1 H), 3.10 (ddd, 1 H), 2.69 - 2.58 (m, 1 H), 2.55 - 2.42 (m, 1 H).

Example 540: (3S)-3-[5-(6-amino-3-pyridazinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

25 The same operation as in Example 228 → Example 534 → Example 51 →  
Example 52 → Example 44 → Example 55 was conducted from the compound  
prepared in Example 538 to give the title compound having the following physical  
properties.

LC/MS *t*<sub>R</sub> 3.06 minutes; MS (ES<sup>+</sup>) *m/z* 507 (M+H)<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.98 (d, 1 H), 7.75 - 7.72 (m, 2 H),  
7.70 - 7.67 (m, 1 H), 6.99 (d, 1 H), 6.15 (s, 1 H), 6.08 (s, 1 H), 5.77 (dd, 1 H), 3.45 -  
3.36 (m, 1 H), 3.10 (ddd, 1 H), 2.69 - 2.60 (m, 1 H), 2.44 - 2.37 (m, 1 H).

Example 541: (3S)-3-[5-(2-{bis[(4-methoxyphenyl)methyl]amino}pyrimidin-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

5 The same operation as in Example 51 → Example 52 was conducted from 1-(2-{bis[(4-methoxyphenyl)methyl]amino}pyrimidin-5-yl)-2-bromoethan-1-one [patent WO2011/141713] to give the title compound having the following physical properties. LC/MS  $t_R$  2.04 minutes; MS ( $ES^+$ )  $m/z$  713 (M+H) <sup>a</sup>.

10 Example 542: (3S)-3-[5-(2-amino-5-pyrimidinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

To a cooled (0 °C) 1,2-dichloroethane (2 mL) solution of the compound prepared in Example 541 (0.13 g) was added trifluoroacetic acid (1 mL) followed by concentrated sulfuric acid (2 drops) and the mixture stirred for 25 minutes at 0 °C then warmed to room temperature and stirred a further 72 hours. On concentration, the residue was  
15 dissolved in water (2 mL) and treated with a saturated aqueous solution of sodium hydrogen carbonate until pH 8 was attained followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 – 10% methanol in dichloromethane then 100% methanol) to afford the title compound having the  
20 following physical properties (38 mg).

LC/MS  $t_R$  2.78 minutes; MS ( $ES^+$ )  $m/z$  473 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.56 (s, 2 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.31 (br. s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.49 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.51 - 2.42 (m, 1 H).

25

Example 543: (3S)-3-[5-(2-amino-5-pyrimidinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 338 → Example 542 was conducted from the compound prepared in Example 541 to give the title compound having the following  
30 physical properties.

LC/MS  $t_R$  3.42 minutes; MS ( $ES^+$ )  $m/z$  507 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.92 (s, 1 H), 9.71 (s, 1 H), 8.49 (s, 2 H), 7.85 - 7.78 (m, 3 H), 6.94 (s, 2 H), 5.99 (s, 1 H), 5.93 (s, 1 H), 5.54 (dd, 1 H), 3.31 - 3.24 (m, 1 H),

2.98 (ddd, 1 H), 2.53 - 2.51 (m, 1 H), 2.23 - 2.15 (m, 1 H).

Example 544: 1-(5-hydroxypyridin-2-yl)ethan-1-one

To a cooled (0 °C) tetrahydrofuran (15 mL) solution of 5-hydroxypyridine-2-carbonitrile (0.92 g) was added a 1.4 M solution of methyl magnesium bromide in a 3:1 mixture of toluene and tetrahydrofuran (16.5 mL) and the mixture warmed to room temperature and stirred 16 hours. At this juncture, a 1.4 M solution of methyl magnesium bromide in a 3:1 mixture of toluene and tetrahydrofuran (8.25 mL) was added and the reaction mixture stirred a further 2 hours at room temperature. To the cooled (0 °C) reaction mixture, a saturated aqueous solution of ammonium chloride (5 mL) was added followed by concentrated sulfuric acid (1.85 mL). To this mixture, a 4 M solution of sodium hydroxide was added until pH 5 was attained, whereupon the mixture was extracted into ethyl acetate. The combined organic layers were dried and concentrated to give the title compound having the following physical properties (1 g).  
LC/MS  $t_R$  0.96 minutes; MS (ES<sup>+</sup>)  $m/z$  138 (M+H)<sup>a</sup>.

Example 545: formic acid - (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(5-hydroxy-2-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone (2:1)

The same operation as in Example 204 → Example 51 → Example 52 was conducted from the compound prepared in Example 544 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties.

LC/MS  $t_R$  2.88 minutes; MS (ES<sup>+</sup>)  $m/z$  473 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.42 (s, 2 H), 8.05 (d, 1 H), 7.75 - 7.72 (m, 2 H), 7.70 - 7.64 (m, 2 H), 7.40 (s, 1 H), 7.24 (dd, 1 H), 6.14 (s, 1 H), 6.09 (s, 1 H), 5.81 (dd, 1 H), 3.49 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.54 - 2.47 (m, 1 H).

Example 546: (3S)-3-[4-chloro-5-(5-hydroxy-2-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 545 (19 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (6.8

mg).

LC/MS  $t_R$  4.12 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.16 (d, 1 H), 7.87 (d, 1 H), 7.77 - 7.70 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.27 (dd, 1 H), 6.15 (s, 1 H), 6.08 (s, 1 H), 5.77 (dd, 1 H), 3.46 - 3.34 (m, 1 H), 3.09 (ddd, 1 H), 2.62 (qd, 1 H), 2.50 - 2.39 (m, 1 H).

Example 547: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(6-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 204 → Example 51 → Example 52 was conducted from 1-(6-fluoropyridin-3-yl)ethan-1-one [patent WO2006/082392] to give the title compound having the following physical properties.

LC/MS  $t_R$  3.39 minutes; MS (ES<sup>+</sup>)  $m/z$  475 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.29 (br. s, 1 H), 9.69 (s, 1 H), 8.54 (d, 1 H), 8.25 - 8.19 (m, 1 H), 7.82 - 7.76 (m, 3 H), 7.64 (d, 1 H), 7.14 (dd, 1 H), 5.97 (br. s, 1 H), 5.96 (s, 1 H), 5.63 (dd, 1 H), 3.42 - 3.35 (m, 1 H), 3.00 (dd, 1 H), 2.55 - 2.51 (m, 1 H), 2.37 - 2.30 (m, 1 H).

Example 548: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(6-oxo-1,6-dihydro-3-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

To a 1,2-dimethoxyethane (3 mL) suspension of the compound prepared in Example 547 (80 mg) was added 3 M hydrochloric acid and the mixture stirred at 80 °C for 2 hours 30 minutes. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (12 mL) was added followed by extraction into ethyl acetate. The combined organic layers were dried and concentrated and the residue triturated in dichloromethane (3 mL), the resultant precipitate being collected by filtration to give the title compound having the following physical properties (61 mg).

LC/MS  $t_R$  2.73 minutes; MS (ES<sup>+</sup>)  $m/z$  473 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.92 (dd, 1 H), 7.78 - 7.71 (m, 3 H), 7.71 - 7.67 (m, 1 H), 7.29 (s, 1 H), 6.59 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.47 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.68 - 2.59 (m, 1 H), 2.48 - 2.41 (m, 1 H).

Example 549: (3S)-3-[4-chloro-5-(6-oxo-1,6-dihydro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 548 (49 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (15 mg).

LC/MS  $t_R$  3.35 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H) <sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.89 (d, 1 H), 7.79 - 7.71 (m, 3 H), 7.71 - 7.66 (m, 1 H), 6.64 (d, 1 H), 6.12 (s, 1 H), 6.10 (s, 1 H), 5.68 (dd, 1 H), 3.48 - 3.40 (m, 1 H), 3.11 (ddd, 1 H), 2.70 - 2.58 (m, 1 H), 2.42 - 2.34 (m, 1 H).

Example 550: 1-[6-(methylamino)pyridin-3-yl]ethan-1-one

- 10 A suspension of 1-(6-chloropyridin-3-yl)ethanone (0.30 g) in a 40% weight solution of methylamine in water (4.5 mL) was heated at 145 °C for 50 minutes by microwave irradiation. On cooling to room temperature, the reaction mixture was extracted into ethyl acetate. The combined organic layers were dried and concentrated and the residue purified by column chromatography (0 – 70% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.19 g).

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.73 (d, 1 H), 8.03 (dd, 1 H), 6.40 (d, 1 H), 3.01 (d, 3 H), 2.51 (s, 3 H).

Example 551: 2-bromo-1-[6-(methylamino)pyridin-3-yl]ethan-1-one dihydrobromide

- 20 The compound prepared in Example 550 (0.17 g) was treated as detailed in Example 204 to give the title compound having the following physical properties (0.20 g).


LC/MS  $t_R$  0.89 minute; MS (ES<sup>+</sup>)  $m/z$  229 and 231 (M+H) <sup>a</sup>.

- 25 Example 552: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 551 to give the title compound having the following physical properties.

- 30 LC/MS  $t_R$  2.73 minutes; MS (ES<sup>+</sup>)  $m/z$  486 (M+H), 244 (M/2+H) <sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.24 (d, 1 H), 7.72 (s, 3 H), 7.69 (s, 1 H), 7.17 (br. s, 1 H), 6.56 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.77 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.88 (s, 3 H), 2.68 - 2.58 (m, 1 H), 2.50 - 2.42 (m, 1 H).



Example 553: tert-butyl N-(5-acetyl-6-fluoropyridin-2-yl)carbamate

The same operation as in Example 90 → Example 228 was conducted from 2-amino-6-fluoro-5-iodopyridine [J. Org. Chem. 71(8), 2922 (2006)] to give the title compound having the following physical properties.

LC/MS  $t_R$  2.01 minute; MS ( $ES^+$ )  $m/z$  199 ( $M-t-Bu+H$ )<sup>a</sup>.

Example 554: tert-butyl N-(5-acetyl-6-fluoropyridin-2-yl)-N-methylcarbamate

To an N,N-dimethylformamide (2 mL) suspension of sodium hydride (46 mg, 60% dispersion in mineral oil) was added a solution of the compound prepared in Example 553 (314 mg) in N,N-dimethylformamide (5 mL) and the mixture stirred at room temperature for 10 minutes. At this juncture, iodomethane (72  $\mu$ L) was added and the mixture stirred at room temperature for 16 hours. To the reaction mixture, water (10 mL) was added followed by extraction into ethyl acetate. The organic layer was washed sequentially with water and saturated saline, dried and concentrated. The residue was purified by column chromatography (0 - 40% ethyl acetate in heptanes) to give the title compound having the following physical properties (146 mg).

LC/MS  $t_R$  2.24 minute; MS ( $ES^+$ )  $m/z$  213 ( $M+H$ )<sup>a</sup>.

Example 555: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[2-fluoro-6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 204 → Example 51 → Example 52 was conducted from the compound prepared in Example 554 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.13 minutes; MS ( $ES^+$ )  $m/z$  504 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  10.72 (br. s, 1 H), 8.53 (s, 1 H), 8.26 (dd, 1 H), 7.64 - 7.60 (m, 1 H), 7.56 - 7.53 (m, 1 H), 7.51 (d, 1 H), 6.35 (s, 1 H), 6.30 (dd, 1 H), 5.83 (d,

1 H), 5.65 (s, 1 H), 4.62 - 4.53 (m, 1 H), 3.50 - 3.23 (m, 2 H), 2.99 (dd, 1 H), 2.94 (s, 3 H), 2.52 - 2.40 (m, 1 H).

5 Example 556: (3S)-3-{5-[2-chloro-6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 550 → Example 90 → Example 204 → Example 51 → Example 52 was conducted from 2,6-dichloro-3-iodopyridine to give the title compound having the following physical properties.

LC/MS  $t_R$  3.20 minutes; MS (ES<sup>+</sup>)  $m/z$  520 (M+H)<sup>b</sup>

10 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.82 - 7.71 (m, 3 H), 7.71 - 7.66 (m, 1 H), 7.30 (br. s, 1 H), 6.48 (d, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.79 (dd, 1 H), 3.48 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.87 (s, 3 H), 2.67 - 2.57 (m, 1 H), 2.53 - 2.46 (m, 1 H).

15 Example 557: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(dimethylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 550 → Example 204 → Example 51 → Example 52 was conducted from 1-(6-chloropyridin-3-yl)ethan-1-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 550 in the operation, a 40% weight solution of dimethylamine in water was used).

20 LC/MS  $t_R$  2.89 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.34 (br. s, 1 H), 7.80 (br. s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.22 (br. s, 1 H), 6.70 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.14 - 3.10 (m, 1 H), 3.09 (s, 6 H),  
25 2.68 - 2.59 (m, 1 H), 2.55 - 2.41 (m, 1 H).

Example 558: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(ethylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 550 → Example 204 → Example 51 → Example 52 was conducted from 1-(6-chloropyridin-3-yl)ethan-1-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 550 in the operation, a 70% weight solution of ethylamine in water was used).

30 LC/MS  $t_R$  2.78 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 8.24 (d, 1 H), 7.77 - 7.68 (m, 4 H), 7.18 (br. s, 1 H), 6.57 (d, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.78 (dd, 1 H), 3.35 - 3.31 (m, 2 H), 3.50 - 3.41 (m, 1 H), 3.12 (ddd, 1 H), 2.65 (ddd, 1 H), 2.51 - 2.44 (m, 1 H), 1.25 (t, 3 H).

5

Example 559: 1-{6-[(2-methoxyethyl)amino]pyridin-3-yl}ethan-1-one

To a propan-2-ol (7 mL) solution of 1-(6-chloropyridin-3-yl)ethanone (0.70 g) was added 2-methoxyethylamine (1.57 mL) and the mixture was heated at 180 °C for 50 minutes by microwave irradiation. On cooling to room temperature, the reaction mixture was diluted with water (40 mL) and extracted into ethyl acetate. The combined organic layers were dried and concentrated and the residue purified by column chromatography (0 - 80% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.55 g).

10

LC/MS *t<sub>R</sub>* 0.80 minute; MS (ES<sup>+</sup>) *m/z* 195 (M+H)<sup>a</sup>.

15

Example 560: 2-bromo-1-{6-[(2-methoxyethyl)amino]pyridin-3-yl}ethan-1-one dihydrobromide

20

The compound prepared in Example 559 (0.30 g) was treated as detailed in Example 204 to give the title compound having the following physical properties (0.67 g).

LC/MS *t<sub>R</sub>* 1.25 minute; MS (ES<sup>+</sup>) *m/z* 273 and 275 (M+H)<sup>a</sup>.

Example 561: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

25

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 560 to give the title compound having the following physical properties.

LC/MS *t<sub>R</sub>* 3.06 minutes; MS (ES<sup>+</sup>) *m/z* 530 (M+H), 265.5 (M/2+H)<sup>b</sup>

30

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.23 (d, 1 H), 7.75 - 7.70 (m, 3 H), 7.69 (s, 1 H), 7.17 (s, 1 H), 6.61 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.60 - 3.55 (m, 2 H), 3.51 - 3.46 (m, 2 H), 3.46 - 3.39 (m, 1 H), 3.38 (s, 3 H), 3.10 (ddd, 1 H), 2.69 - 2.58 (m, 1 H), 2.50 - 2.42 (m, 1 H).

Example 562 : (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(2-ethoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 559 → Example 204 → Example 51 → Example 52 was conducted from 1-(6-chloropyridin-3-yl)ethan-1-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 559 in the operation, 2-ethoxyethylamine was used).

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H), 272.5 (M/2+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.23 (d, 1 H), 7.75 - 7.71 (m, 3 H), 7.69 - 7.65 (m, 1 H), 7.18 (s, 1 H), 6.62 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.62 (t, 1 H), 3.55 (q, 2 H), 3.48 (t, 2 H), 3.46 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.49 - 2.42 (m, 1 H), 1.20 (t, 3 H).

Example 563 : (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(3-methoxypropyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 559 → Example 204 → Example 51 → Example 52 was conducted from 1-(6-chloropyridin-3-yl)ethan-1-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 559 in the operation, 3-methoxypropylamine was used).

LC/MS  $t_R$  2.85 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.22 (br. s, 1 H), 7.76 - 7.66 (m, 4 H), 7.16 (br. s, 1 H), 6.56 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.51 (t, 2 H), 3.42 - 3.46 (m, 1 H), 3.37 (t, 2 H), 3.35 (s, 3 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.52 - 2.41 (m, 1 H), 1.87 (quintet, 2 H).

Example 564 : (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(3-hydroxy-3-methylbutyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 559 → Example 204 → Example 51 → Example 52 was conducted from 1-(6-chloropyridin-3-yl)ethan-1-one to give the title compound having the following physical properties. (Note: in the step corresponding to Example 559 in the operation, 4-amino-2-methylbutan-2-ol was used).

LC/MS  $t_R$  2.85 minutes; MS (ES<sup>+</sup>)  $m/z$  558 (M+H), 279.5 (M/2+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.23 (d, 1 H), 7.76 - 7.70 (m, 3 H), 7.70 - 7.66 (m, 1 H), 7.17 (s, 1 H), 6.57 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1

H), 3.49 - 3.42 (m, 1 H), 3.41 - 3.37 (m, 2 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.50 - 2.42 (m, 1 H), 1.83 - 1.77 (m, 2 H), 1.26 (s, 6 H).

Example 565: 1-{6-[(1,3-oxazol-2-yl)methyl]amino}pyridin-3-yl}ethan-1-one

- 5 To a propan-2-ol (10 mL) solution of 1-(6-fluoropyridin-3-yl)ethanone [patent WO2006/082392] (0.50 g) was added 1-(1,3-oxazol-2-yl)methylamine hydrochloride (0.73 g) and N,N-diisopropylethylamine (1.25 mL) and the mixture was heated at 110 °C for 30 minutes by microwave irradiation. On cooling to room temperature, the reaction mixture was diluted with water (20 mL) and extracted into ethyl acetate. The  
10 combined organic layers were dried and concentrated and the residue purified by column chromatography (0 – 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.22 g).  
LC/MS  $t_R$  1.04 minute; MS (ES<sup>+</sup>)  $m/z$  218 (M+H)<sup>a</sup>.

- 15 Example 566: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-{6-[(1,3-oxazol-2-yl)methyl]amino}-3-pyridinyl)-1H-imidazol-2-yl)-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 204 → Example 51 → Example 52 was conducted from the compound prepared in Example 565 to give the title compound having the following physical properties.

- 20 LC/MS  $t_R$  2.83 minutes; MS (ES<sup>+</sup>)  $m/z$  553 (M+H), 277 (M/2+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.26 (d, 1 H), 7.84 (s, 1 H), 7.76 - 7.71 (m, 3 H), 7.69-7.66 (m, 1 H), 7.17 (br. s, 1 H), 7.09 (s, 1 H), 6.65 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.76 (dd, 1 H), 4.67 (s, 2 H), 3.48 - 3.39 (m, 1 H), 3.09 (ddd, 1 H), 2.67 - 2.58 (m, 1 H), 2.49 - 2.42 (m, 1 H).

25

Example 567(1) and 567(2): 3-{[5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]amino}-N,N-dimethylpropanamide and 3-{[5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]amino}propanamide

30

The same operation as in Example 565 → Example 55 → Example 250 → Example 204 → Example 51 → Example 52 was conducted from 1-(6-fluoropyridin-3-yl)ethan-1-one [patent WO2006/082392] to give the title compounds in a 1:1 ratio

having the following physical properties (Note: in the step corresponding to Example 565 in the operation,  $\beta$ -alanine hydrochloride tert-butyl ester was used. In the step corresponding to Example 250 in the operation, dimethylamine hydrochloride was used).

5 Example 567(1):

LC/MS  $t_R$  2.82 minutes; MS ( $ES^+$ )  $m/z$  571 (M+H), 286 (M/2+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.25 (s, 1 H), 7.73 (d, 4 H), 7.17 (br. s, 1 H), 6.57 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.60 (t, 2 H), 3.48 - 3.40 (m, 1 H), 3.10 (ddd, 1 H), 3.04 (s, 3 H), 2.94 (s, 3 H), 2.69 (t, 2 H), 2.66 - 2.60 (m, 1 H),  
10 2.50 - 2.42 (m, 1 H).

Example 567(2):

LC/MS  $t_R$  2.67 minutes; MS ( $ES^+$ )  $m/z$  543 (M+H), 272 (M/2+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.25 (d, 1 H), 7.77 - 7.65 (m, 4 H),  
15 7.17 (br. s, 1 H), 6.58 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.77 (dd, 1 H), 3.59 (t, 2 H), 3.50 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.66 - 2.58 (m, 1 H), 2.52 (t, 2 H), 2.49 - 2.41 (m, 1 H).

Example 568: trifluoroacetic acid - 4-[(5-acetylpyridin-2-yl)amino]butanoic acid (1: 1)

20 The same operation as in Example 565  $\rightarrow$  Example 40 was conducted from 1-(6-fluoropyridin-3-yl)ethan-1-one [patent WO2006/082392] to give the title compound having the following physical properties. (Note: in the step corresponding to Example 565 in the operation, tert-butyl 4-aminobutanoate was used).

LC/MS  $t_R$  0.63 minutes; MS ( $ES^+$ )  $m/z$  223 (M+H) <sup>a</sup>.

25

Example 569: 1-(5-acetylpyridin-2-yl)pyrrolidin-2-one

The compound prepared in Example 568 (360 mg) was treated as detailed in Example 250 to give the title compound having the following physical properties (65 mg).

30 LC/MS  $t_R$  1.40 minute; MS ( $ES^+$ )  $m/z$  205 (M+H) <sup>a</sup>.

Example 570: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[6-(2-oxo-1-pyrrolidinyl)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

35 The same operation as in Example 204  $\rightarrow$  Example 51  $\rightarrow$  Example 52 was conducted from the compound prepared in Example 569 to give the title compound

having the following physical properties.

LC/MS  $t_R$  3.21 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H), 562 (M+Na), 270.5 (M/2+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.65 (s, 1 H), 8.61 (s, 1 H), 8.36 (d, 1 H), 7.93 (d, 1 H),  
7.62 (dd, 1 H), 7.55 (m, 2 H), 7.29 - 7.24 (m, 1 H), 6.27 (s, 1 H), 5.89 (d, 1 H), 5.83 (s,  
5 1 H), 4.10 (t, 2 H), 3.53 - 3.44 (m, 1 H), 3.25 - 3.17 (m, 1 H), 3.06 (dd, 1 H), 2.67 (t, 2  
H), 2.60 - 2.51 (m, 1 H), 2.15 (quintet, 2 H).

Example 571(1) to Example 571(3)

The compounds of the present invention having the following physical data were  
10 synthesised from the compounds prepared in Examples 552, 557 and 561 using the  
method as detailed in Example 44.

Example 571(1): (3S)-3-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-  
7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

15 LC/MS  $t_R$  3.03 minutes; MS (ES<sup>+</sup>)  $m/z$  520 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.27 (d, 1 H), 7.76 - 7.71 (m, 3 H),  
7.70 - 7.66 (m, 1 H), 6.58 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.69 (dd, 2 H), 3.49 -  
3.38 (m, 1 H), 3.09 (ddd, 1 H), 2.89 (s, 3 H), 2.68 - 2.58 (m, 1 H), 2.42 - 2.34 (m, 1 H).

20 Example 571(2): (3S)-3-{4-chloro-5-[6-(dimethylamino)-3-pyridinyl]-1H-imidazol-2-  
yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.21 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.39 (d, 1 H), 7.83 (dd, 1 H), 7.78 -  
7.73 (m, 2 H), 7.72 - 7.68 (m, 1 H), 6.76 (d, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.72 (dd,  
25 1 H), 3.51 - 3.42 (m, 1 H), 3.16 - 3.08 (m, 7 H), 2.65 (qd, 1 H), 2.44 - 2.36 (m, 1 H).

Example 571(3): (3S)-3-(4-chloro-5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-  
imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.21 minutes; MS (ES<sup>+</sup>)  $m/z$  564 (M+H)<sup>b</sup>  
30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.26 (d, 1 H), 7.75 - 7.70 (m, 3 H),  
7.70 - 7.66 (m, 1 H), 6.63 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.69 (dd, 1 H), 3.58 (t, 2  
H), 3.51 (t, 2 H), 3.48 - 3.40 (m, 1 H), 3.38 (s, 3 H), 3.10 (ddd, 1 H), 2.68 - 2.59 (m, 1  
H), 2.42 - 2.34 (m, 1 H).

Example 572(1) to Example 572(7)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 558, 562, 563, 564, 566, 567(1) and 570 using the method as detailed in Example 338.

Example 572(1): (3S)-3-{4-chloro-5-[6-(ethylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.24 (d, 1 H), 7.79 - 7.64 (m, 4 H), 6.57 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.70 (dd, 1 H), 3.48 - 3.39 (m, 1 H), 3.35 - 3.31 (m, 2 H), 3.09 (ddd, 1 H), 2.57 - 2.68 (m, 1 H), 2.41 - 2.33 (m, 1 H), 1.24 (t, 3 H).

Example 572(2): (3S)-3-(4-chloro-5-{6-[(2-ethoxyethyl)amino]-3-pyridinyl}-1H-

imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.16 minutes; MS (ES<sup>+</sup>)  $m/z$  578 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.26 (d, 1 H), 7.75 - 7.71 (m, 3 H), 7.70 - 7.67 (m, 1 H), 6.64 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.69 (dd, 1 H), 3.63 (t, 2 H), 3.55 (q, 2 H), 3.51 (t, 2 H), 3.48 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.42 - 2.34 (m, 1 H), 1.20 (t, 3 H).

Example 572(3): (3S)-3-(4-chloro-5-{6-[(3-methoxypropyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  2.20 minutes; MS (ES<sup>+</sup>)  $m/z$  578 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.24 (d, 1 H), 7.74 - 7.70 (m, 2 H), 7.70 - 7.66 (m, 2 H), 6.57 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.70 (dd, 1 H), 3.50 (t, 2 H), 3.48 - 3.40 (m, 1 H), 3.38 (t, 2 H), 3.34 (s, 3 H), 3.09 (ddd, 1 H), 2.67 - 2.58 (m, 1 H), 2.40 - 2.33 (m, 1 H), 1.87 (quintet, 2 H).

Example 572(4): (3S)-3-(4-chloro-5-{6-[(3-hydroxy-3-methylbutyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  592 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.26 (d, 1 H), 7.75 - 7.71 (m, 3 H), 7.70 - 7.67 (m, 1 H), 6.59 (d, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.69 (dd, 1 H), 3.48 - 3.39 (m, 3 H), 3.10 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.42 - 2.34 (m, 1 H), 1.83 - 1.78 (m, 2 H), 1.27 (s, 6 H).

5

Example 572(5): (3S)-3-(4-chloro-5-{6-[(1,3-oxazol-2-ylmethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS *t<sub>R</sub>* 3.33 minutes; MS (ES<sup>+</sup>) *m/z* 587 (M+H)<sup>b</sup>

10 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 - 9.34 (m, 1 H), 8.27 (br. s, 1 H), 7.85 (s, 1 H), 7.78 - 7.71 (m, 3 H), 7.70 - 7.66 (m, 1 H), 7.10 (s, 1 H), 6.72 - 6.67 (m, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.69 (dd, 1 H), 4.70 (s, 2 H), 3.47 - 3.39 (m, 1 H), 3.13 - 3.05 (m, 1 H), 2.68 - 2.58 (m, 1 H), 2.41 - 2.34 (m, 1 H).

15 Example 572(6): 3-{[5-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-pyridinyl]amino}-N,N-dimethylpropanamide

LC/MS *t<sub>R</sub>* 3.06 minutes; MS (ES<sup>+</sup>) *m/z* 605 (M+H)<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 8.28 (br. s, 1 H), 7.76 - 7.66 (m, 4 H), 6.62 - 6.57 (m, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.70 (dd, 1 H), 3.65 - 3.60 (m, 2 H), 3.48 - 3.39 (m, 1 H), 3.13 - 3.06 (m, 1 H), 3.05 (s, 3 H), 2.94 (s, 3 H), 2.70 (t, 2 H), 2.67 - 2.58 (m, 1 H), 2.42 - 2.34 (m, 1 H).

Example 572(7): formic acid - (3S)-3-{4-chloro-5-[6-(2-oxo-1-pyrrolidinyl)-3-

25 pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone (1:1)

LC/MS *t<sub>R</sub>* 3.99 minutes; MS (ES<sup>+</sup>) *m/z* 574 (M+H)<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 8.70 (d, 1 H), 8.41 (d, 1 H), 8.08 (dd, 1 H), 7.74 (s, 2 H), 7.70 - 7.67 (m, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.72 (dd, 1 H), 4.14 (t, 2 H), 3.50 - 3.40 (m, 1 H), 3.12 (ddd, 1 H), 2.68 (t, 2 H), 2.66 - 2.61 (m, 1 H), 2.43 - 2.36 (m, 1 H), 2.17 (quintet, 2 H).

Example 573: (3S)-3-[5-(2-amino-1,3-thiazol-5-yl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 235 (67 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (1.8 mg).

LC/MS  $t_R$  3.18 minutes; MS (ES<sup>+</sup>)  $m/z$  512 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.77 - 7.71 (m, 2 H), 7.71 - 7.66 (m, 1 H), 7.22 (s, 1 H), 6.13 (s, 1 H), 6.08 (s, 1 H), 5.65 (dd, 1 H), 3.41 (td, 1 H), 3.09 (ddd, 1 H), 2.70 - 2.56 (m, 1 H), 2.39 - 2.30 (m, 1 H).

Example 574: (3S)-3-[5-(2-amino-4-chloro-1,3-thiazol-5-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 201 → Example 203 → Example 204 → Example 51 → Example 52 was conducted from 2-{[(tert-butoxy)carbonyl]amino}-4-chloro-1,3-thiazole-5-carboxylic acid [Bioorg. Med. Chem., 12(23), 6171 (2004)] to give the title compound having the following physical properties. (Note: in the step corresponding to Example 203 in the operation, methyl magnesium bromide was used. In the step corresponding to Example 204 in the operation, removal of the tert-butoxycarbonyl group was also accomplished).

LC/MS  $t_R$  3.38 minutes; MS (ES<sup>+</sup>)  $m/z$  512 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.77 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.37 (br. s, 1 H), 6.14 (s, 1 H), 6.06 (s, 1 H), 5.74 (d, 1 H), 3.47 - 3.36 (m, 1 H), 3.14 - 3.02 (m, 1 H), 2.68 - 2.54 (m, 1 H), 2.52 - 2.41 (m, 1 H).

Example 575: N-[4-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-pyridinyl]acetamide

The compound prepared in Example 190(2) (50 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (14 mg).

LC/MS  $t_R$  3.49 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 8.52 (app. br. s, 1 H), 8.32 (d, 1 H), 7.78 - 7.72 (m, 2 H), 7.70 - 7.66 (m, 1 H), 7.43 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.74 (dd, 1 H), 3.51 - 3.38 (m, 1 H), 3.19 - 3.07 (m, 1 H), 2.74 - 2.62 (m, 1 H), 2.44 - 2.35 (m, 1 H), 2.20 (s, 3 H).

Example 576: methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 190(1) (67 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (40 mg).

LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>

<sup>1</sup>H-NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.97 (br. s, 1 H), 9.69 (s, 1 H), 8.14 - 8.02 (m, 2 H), 7.83 - 7.70 (m, 3 H), 7.58 (br. s, 1 H), 7.29 (d, 1 H), 5.98 (s, 1 H), 5.92 (s, 1 H), 5.61 (d, 1 H), 3.67 (s, 3 H), 3.35 (obs. m, 1 H), 2.98 (dd, 1 H), 2.50 (obs. m, 1 H), 2.33 - 2.22 (m, 1 H).

Examples 577(1), 577(2) and 577(3): (3S)-3-[5-(2-amino-3-chloro-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone, (3S)-3-[5-(2-amino-5-chloro-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone and (3S)-3-[5-(2-amino-5-chloro-4-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 190(1) (97 mg) was treated as detailed in Example 44 to give the title products having the following physical properties.

Example 577(1): 9.7 mg

LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 7.82 - 7.74 (m, 4 H), 7.70 (s, 1 H), 7.16 (d, 1 H), 6.00 (s, 2 H), 5.94 (s, 1 H), 5.92 (s, 1 H), 5.63 (dd, 1 H), 3.65 - 3.07 (obs. m, 1 H), 2.92 - 2.92 (m, 1 H), 2.94 (dd, 1 H), 2.48 - 2.39 (m, 1 H), 2.29 - 2.20 (m, 1 H).

Example 577(2): 12.6 mg

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.70 (s, 1 H), 7.88 (s, 1 H), 7.82 - 7.79 (m, 2 H), 7.79 - 7.76 (m, 1 H), 7.74 (s, 1 H), 7.05 (s, 1 H), 6.04 (s, 2 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.65 (dd, 1 H), 3.52 - 3.18 (obs. m, 1 H), 3.05 - 2.95 (m, 1 H), 2.58 - 2.43 (obs. m, 1 H), 2.35 - 2.27 (m, 1 H).

Example 577(3): 8.6 mg

LC/MS  $t_R$  3.40 minutes; MS ( $ES^+$ )  $m/z$  540 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.67 (s, 1 H), 7.97 (s, 1 H), 7.84 - 7.73 (m, 3 H), 7.23 (br. s, 1 H), 6.44 (br. s, 2 H), 5.96 (s, 2 H), 5.67 (dd, 1 H), 3.49 - 3.19 (obs. m, 1 H),  
5 2.97 (dd, 1 H), 2.57 - 2.43 (obs. m, 1 H), 2.39 - 2.25 (m, 1 H).

Example 578(1) and 578(2): tert-butyl N-(4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}pyridin-2-yl)carbamate and tert-butyl N-[(tert-butoxy)carbonyl]-N-(4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}pyridin-2-yl)carbamate  
10

To a *tert*-butanol (1.3 mL) solution of the compound prepared in Example 190(1) (123 mg) was added di-*tert*-butyl dicarbonate (250 mg). The mixture was stirred at room temperature for 36 hours then at 50 °C for a further 3 hours. To the ambient  
15 reaction mixture, concentrated ammonia solution (0.15 mL) was added and the mixture stirred at room temperature for 6 hours then at 50 °C for a further hour. The reaction mixture was concentrated and the residue purified by column chromatography (0 - 15% methanol in ethyl acetate) to give the title products in a 4:1 ratio.

20 Example 578(1):

LC/MS  $t_R$  1.69 minutes; MS ( $ES^+$ )  $m/z$  572 (M+H)<sup>a</sup>.

Example 578(2):

LC/MS  $t_R$  2.10 minutes; MS ( $ES^+$ )  $m/z$  672 (M+H)<sup>a</sup>.

25

Example 579: (3S)-3-[5-(2-amino-4-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 44 → Example 206 was conducted from the 4: 1 ratio of compounds prepared in Example 578 to give the title compound having the  
30 following physical properties.

LC/MS  $t_R$  3.04 minutes; MS ( $ES^+$ )  $m/z$  506 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.09 (br. s, 1 H), 9.71 (s, 1 H), 7.96 (d, 1 H), 7.85 - 7.76 (m, 3 H), 6.85 - 6.75 (m, 2 H), 6.06 (br. s, 2 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.57

(dd, 1 H), 3.29 - 3.21 (m, 1 H), 3.04 - 2.93 (m, 1 H), 2.58 - 2.52 (obs m, 1 H), 2.23 - 2.13 (m, 1 H).

Example 580: tert-butyl N-(4-acetylpyridin-2-yl)-N-methylcarbamate

- 5        The same operation as in Example 550 → Example 228 was conducted from 1-(2-chloropyridin-4-yl)ethanone to give the title compound having the following physical properties.

LC/MS  $t_R$  2.05 minutes; MS (ES<sup>+</sup>)  $m/z$  251 (M+H)<sup>a</sup>.

10    Example 581: 2-bromo-1-[2-(methylamino)pyridin-4-yl]ethan-1-one

To a tetrahydrofuran (6 mL) solution of the compound prepared in Example 580 (161 mg) was added a 33% weight solution of hydrogen bromide in acetic acid (222  $\mu$ L) followed by bromine (33  $\mu$ L) and the mixture was stirred at room temperature for 1 hour. The reaction mixture was concentrated and the residue triturated with a 1:3 mixture of dichloromethane and *tert*-butyl methyl ether (20 mL), the resultant precipitate being isolated by filtration to give the title compound having the following physical properties.

LC/MS  $t_R$  0.71 minutes; MS (ES<sup>+</sup>)  $m/z$  229 (M+H)<sup>a</sup>.

20    Example 582: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[2-(methylamino)-4-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 581 to give the title compound having the following physical properties.

- 25    LC/MS  $t_R$  2.93 minutes; MS (ES<sup>+</sup>)  $m/z$  486 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.87 (d, 1 H), 7.72 (m, 2 H), 7.68 (dd, 1 H), 7.49 (br. s, 1 H), 6.84 (app. d, 2 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 3.43 (td, 1 H), 3.11 (ddd, 1 H), 2.89 (s, 3 H), 2.64 (qd, 1 H), 2.51 - 2.38 (m, 1 H).

30    Example 583: prop-2-en-1-yl N-(4-{4-chloro-2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}pyridin-2-yl)-N-methylcarbamate

The same operation as in Example 484 → Example 44 was conducted from the compound prepared in Example 582 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 484 in the operation, allyl chloroformate was used).

- 5 LC/MS  $t_R$  2.04 minutes; MS ( $ES^+$ )  $m/z$  604 ( $M+H$ )<sup>a</sup>.

Example 584: (3S)-3-{4-chloro-5-[2-(methylamino)-4-pyridinyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

- To a nitrogen degassed tetrahydrofuran (1.5 mL) solution of the compound prepared in Example 583 (93 mg) and N,N'-dimethylbarbituric acid (29 mg) was added tetrakis(triphenylphosphine)palladium(0) (9 mg) and the mixture stirred at room temperature for 2.5 hours. To the reaction mixture, saturated sodium hydrogen carbonate solution (30 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by high performance liquid chromatography (5 to 100% acetonitrile in water) to give the title compound having the following physical properties (36 mg).
- 10  
15

LC/MS  $t_R$  3.10 minutes; MS ( $ES^+$ )  $m/z$  520 ( $M+H$ )<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.96 (dd, 1 H), 7.79 - 7.63 (m, 3 H), 6.91 - 6.82 (m, 2 H), 6.12 (s, 1 H), 6.11 (s, 1 H), 5.72 (dd, 1 H), 3.53 - 3.35 (m, 1 H), 3.19 - 3.02 (m, 1 H), 2.89 (s, 3 H), 2.65 (qd, 1 H), 2.45 - 2.29 (m, 1 H).
- 20

Example 585: (3S)-3-[5-(3-amino-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

- The same operation as in Example 51 → Example 52 → Example 74 was conducted from 2-bromo-1-(3-nitropyridin-4-yl)ethan-1-one [J. Med. Chem., 53(2), 787 (2010)] to give the title compound having the following physical properties.
- 25

LC/MS  $t_R$  2.95 minutes; MS ( $ES^+$ )  $m/z$  472 ( $M+H$ )<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1H), 8.00 (s, 1 H), 7.78 - 7.64 (m, 4 H), 7.59 (s, 1 H), 7.39 (d, 1 H), 6.18 (s, 1 H), 6.05 (s, 1 H), 5.83 (dd, 1 H), 3.48 - 3.37 (m, 1 H), 3.11 (dd, 1 H), 2.65 - 2.56 (m, 1 H), 2.56 - 2.47 (m, 1 H).
- 30

Example 586: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2-methyl-4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone acetate

The same operation as in Example 51 → Example 52 was conducted from 2-bromo-1-(2-methylpyridin-4-yl)ethan-1-one [J. Med. Chem., 53(2), 787 (2010)] to give the title compound having the following physical properties.

LC/MS  $t_R$  2.91 minutes; MS (ES<sup>+</sup>)  $m/z$  471 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.33 (d, 1 H), 7.77 - 7.62 (m, 5 H), 7.56 (br. s, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.79 (dd, 1 H), 3.51 - 3.39 (m, 1 H), 3.12 (ddd, 1 H), 2.70 - 2.59 (m, 1 H), 2.55 (s, 3 H), 2.52 - 2.43 (m, 1 H).

Example 587: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(2,6-dimethyl-4-pyridinyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 139 → Example 204 → Example 51 → Example 52 was conducted from 4-bromo-2,6-dimethylpyridine to give the title compound having the following physical properties.

LC/MS  $t_R$  2.99 minutes; MS (ES<sup>+</sup>)  $m/z$  485 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.67 (m, 1 H), 7.61 (br. s, 1 H), 7.40 (br. s, 2 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.51 - 3.40 (m, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.52 - 2.43 (obs. m, 1 H), 2.50 (s, 6 H).

Example 588: (3S)-3-[4-chloro-5-(2,6-dimethyl-4-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 587 (69 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (3 mg).

LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  519 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.46 (s, 2 H), 6.13 (s, 1 H), 6.11 (s, 1 H), 5.72 (dd, 1 H), 3.47 - 3.39 (m, 1 H), 3.12 (ddd, 1 H), 2.71 - 2.61 (m, 1 H), 2.54 (s, 6 H), 2.42 - 2.34 (m, 1 H).

Example 589: 1-(6-aminopyridin-2-yl)ethan-1-one dihydrobromide

The same operation as in Example 187 → Example 188 → Example 189 was conducted from 6-fluoropyridine-2-carbonitrile to give the title compound having the following physical properties.

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 7.96 (t, 1 H), 7.59 (d, 1 H), 7.15 (d, 1 H), 4.95 (s, 2 H), 4.12 (br. s, 3 H).

5 Example 590: prop-2-en-1-yl N-(6-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}pyridin-2-yl)carbamate

The same operation as in Example 51 → Example 484 → Example 52 was conducted from the compound prepared in Example 589 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 10 484 in the operation, allyl chloroformate was used. Additionally, in the same step in the operation, the stage employing methanol and concentrated ammonia solution was omitted and the crude product extracted into ethyl acetate from water).  
LC/MS *t<sub>R</sub>* 1.73 minutes; MS (ES<sup>+</sup>) *m/z* 556 (M+H)<sup>a</sup>.

15 Example 591: (3S)-3-[5-(6-amino-2-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 590 (75 mg) was treated as detailed in Example 584 to give the title compound having the following physical properties (11.5 mg).  
20 LC/MS *t<sub>R</sub>* 2.83 minutes; MS (ES<sup>+</sup>) *m/z* 472 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.76 - 7.70 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.51 - 7.36 (m, 2 H), 7.01 (br. s, 1 H), 6.44 (d, 1 H), 6.14 (s, 1 H), 6.08 (s, 1 H), 5.81 (dd, 1 H), 3.49 - 3.37 (m, 1 H), 3.10 (ddd, 1 H), 2.68 - 2.57 (m, 1 H), 2.55 - 2.46 (m, 1 H).

25 Example 592: (3S)-3-[5-(6-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 338 → Example 584 was conducted from the compound prepared in Example 590 to give the title compound having the following  
30 physical properties.  
LC/MS *t<sub>R</sub>* 3.19 minutes; MS (ES<sup>+</sup>) *m/z* 506 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.75 - 7.70 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.47 (t, 1 H), 7.25 (d, 1 H), 6.46 (d, 1 H), 6.16 (s, 1 H), 6.08 (s, 1 H), 5.80 (dd, 1 H), 3.39 (td, 1 H), 3.09 (ddd, 1 H), 2.67 - 2.56 (m, 1 H), 2.53 - 2.44 (m, 1 H).

5 Example 593: formic acid - (3S)-3-[5-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone (1:1)

The same operation as in Example 484 → Example 364 → Example 584 was conducted from the compound prepared in Example 194 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties. (Note: in the step corresponding to Example 484 in the operation, allyl chloroformate was used).

LC/MS *t<sub>R</sub>* 2.93 minutes; MS (ES<sup>+</sup>) *m/z* 490 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.38 (s, 1 H), 8.19 (br. s, 1 H), 8.09 (br. s, 1 H), 7.78 - 7.67 (m, 4 H), 6.73 (d, 1 H), 6.13 (s, 1 H), 6.11 (s, 1 H), 5.68 (dd, 1 H), 3.45 (td, 1 H), 3.12 (ddd, 1 H), 2.70 - 2.59 (m, 1 H), 2.43 - 2.34 (m, 1 H).

20 Example 594: 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-fluoro-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The compound prepared in Example 531 (200 mg) was treated as detailed in Example 364 to give the title product having the following physical properties (10.5 mg).

LC/MS *t<sub>R</sub>* 3.95 minutes; MS (ES<sup>+</sup>) *m/z* 592 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 8.58 (s, 1 H), 7.93 (d, 1 H), 7.75 - 7.70 (m, 2 H), 7.67 (d, 1 H), 7.49 (d, 1 H), 6.15 (s, 1 H), 6.07 (s, 1 H), 5.75 (d, 1 H), 4.31 - 4.27 (m, 2 H), 3.68 - 3.63 (m, 2 H), 3.45 - 3.35 (m, 1 H), 3.39 (s, 3 H), 3.09 (ddd, 1 H), 2.61 (qd, 1 H), 2.45 - 2.38 (m, 1 H).

30 Example 595: ethyl (3R)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

The same operation as in Example 1 → Example 2 → Example 3 → Example 4 → Example 5 → Example 6 → Example 7 → Example 9 was conducted from ethyl 5-oxo-D-prolinate to give the title compound having the following physical properties.

LC/MS  $t_R$  3.70 minutes; MS ( $ES^+$ )  $m/z$  793 (2M+Na), 771 (2M+H), 408 (M+Na), 386 (M+H), 358 (M-N<sub>2</sub>+H)<sup>b</sup>.

Example 596: (3R)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

To a 1,4-dioxane (3.8 mL) solution of the compound prepared in Example 595 (1.28 g) was added 2 M hydrochloric acid (6.4 mL) and the reaction stirred at 70 °C for 2 hours. A second aliquot of 2 M hydrochloric acid (6.4 mL) was added and the mixture stirred at the same temperature a further 2 hours. A third aliquot of 2 M hydrochloric acid (6.4 mL) was added and the reaction stirred at 70 °C for a further 2 hours before cooling to room temperature. To the reaction mixture, ammonium sulfate (13.1 g) was added and the resultant suspension extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was triturated with a 1: 1 mixture of ethyl acetate and heptane to give the title compound having the following physical properties (1.15 g).

LC/MS  $t_R$  3.19 minutes; MS ( $ES^+$ )  $m/z$  737 (2M+Na), 715 (2M+H), 380 (M+Na), 358 (M+H), 330 (M-N<sub>2</sub>+H)<sup>b</sup>.

Example 597: (3R)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 596 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used)

LC/MS  $t_R$  2.68 minutes; MS ( $ES^+$ )  $m/z$  472 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 8.20 (app. br. s, 1 H), 7.79 - 7.70 (m, 3 H), 7.67 (d, 1 H), 7.18 (br. s, 1 H), 6.60 (d, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 3.44 (td, 1 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.46 (app. br. s, 1 H).

Example 598: formic acid - (3R)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone (1:1)

The compound prepared in Example 597 (250 mg) was treated as detailed in Example 44 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (86 mg).

LC/MS  $t_R$  2.94 minutes; MS ( $ES^+$ )  $m/z$  506 and 508 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 8.21 (d, 1 H), 8.17 (br. s, 1 H), 7.85 (d, 1 H), 7.79 - 7.73 (m, 2 H), 7.72 - 7.69 (m, 1 H), 6.74 (d, 1 H), 6.14 (s, 1 H), 6.12 (s, 1 H), 5.71 (dd, 1 H), 3.46 (td, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.44 - 2.36 (m, 1 H).

Example 599: prop-2-en-1-yl 2-[(diphenylmethylidene)amino]acetate

To a dichloromethane (16 mL) solution of glycine allyl ester [J. Org. Chem., 66(15), 5241 (2001)] (5.05 g) was added benzophenone imine (5.6 mL) and the reaction stirred at room temperature for 22 hours. To the reaction mixture, dichloromethane (25 mL) was added and the suspension filtered through Celite®. The filtrate was concentrated to half the original volume and *tert*-butyl methyl ether added. The suspension was filtered once more through Celite® and the filtrate concentrated to give the title compound having the following physical properties (8.56 g).

<sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  7.71 - 7.64 (m, 2 H), 7.53 - 7.45 (m, 3 H), 7.44 - 7.39 (m, 1 H), 7.35 (app. t, 2 H), 7.19 (dd, 2 H), 6.00 - 5.87 (m, 1 H), 5.37 - 5.29 (m, 1 H), 5.25 (dd, 1 H), 4.66 (d, 2 H), 4.25 (s, 2 H).

Example 600: prop-2-en-1-yl 2-amino-3-oxo-3-phenylpropanoate hydrochloride

To a cooled (-78 °C) tetrahydrofuran (7.2 mL) solution of the compound prepared in Example 599 (1.0 g) was added a solution of 1 M lithium bis(trimethylsilyl)amide in tetrahydrofuran (3.6 mL) and the mixture stirred at -78 °C for 30 minutes. The resultant solution was added to a cooled (-78 °C) tetrahydrofuran (3.6 mL) solution of benzoyl chloride (0.42 mL) and the mixture stirred at -78 °C for 2 hours. To the cooled (-78 °C) reaction mixture, a 1 M aqueous solution of hydrochloric acid (7.2 mL) was added and the reaction stirred at room temperature for 1 hour. The reaction mixture was concentrated and the residue triturated with *tert*-butyl methyl ether.

The resultant precipitate was collected by filtration to give the title compound having the following physical properties (1.13 g).

LC/MS  $t_R$  1.13 minutes; MS (ES<sup>+</sup>)  $m/z$  220 (M+H)<sup>a</sup>.

5 Example 601: prop-2-en-1-yl 2-([(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]formamido)-3-oxo-3-phenylpropanoate

To a cooled (-20 °C) tetrahydrofuran (6 mL) solution of the compound prepared in Example 9 (1.1 g) and N-methylmorpholine (0.85 mL) was added isobutyl chloroformate (0.41 mL) and the mixture stirred at -20 °C for 40 minutes then at room  
10 temperature for 5 minutes. To the reaction mixture, an N,N-dimethylformamide (7 mL) suspension of the compound prepared in Example 600 (1.01 g) and N-methylmorpholine (0.85 mL) was added and the mixture stirred at room temperature for 1 hour. To the reaction mixture, water (20 mL) and a saturated aqueous solution of sodium hydrogen carbonate (20 mL) were added followed by extraction into ethyl  
15 acetate. The combined organic layers were washed sequentially with 1 M hydrochloric acid and saturated saline, dried and concentrated. The residue was purified by column chromatography (20% - 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (1.17 g).

LC/MS  $t_R$  1.99 minutes; MS (ES<sup>+</sup>)  $m/z$  559 (M+H)<sup>a</sup>.

20

Example 602: prop-2-en-1-yl 2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-5-phenyl-1H-imidazole-4-carboxylate

To an acetic acid (8.1 mL) suspension of the compound prepared in Example 601 (0.9 g) was added ammonium acetate (1.84 g) and the mixture stirred at reflux for 2  
25 hours. To the ambient reaction mixture, saturated sodium hydrogen carbonate was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (20% - 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (358 mg).

30 LC/MS  $t_R$  2.03 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>a</sup>.

Example 603: 2-[(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-5-phenyl-1H-imidazole-4-carboxylic acid

The compound prepared in Example 602 (100 mg) was treated as detailed in Example 584 to give the title compound having the following physical properties (38 mg).

LC/MS  $t_R$  3.62 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.77 - 7.70 (m, 4 H), 7.70 - 7.65 (m, 1 H), 7.42 - 7.32 (m, 3 H), 6.13 (s, 1 H), 6.08 (s, 1 H), 5.76 (dd, 1 H), 3.47 (td, 1 H), 3.11 (ddd, 1 H), 2.65 (qd, 1 H), 2.52 - 2.42 (m, 1 H).

10 Example 604: 2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-5-phenyl-1H-imidazole-4-carbonitrile

- To an N,N-dimethylformamide (2 mL) solution of the compound prepared in Example 603 (50 mg) was added 1.1'-carbonyldiimidazole (14 mg) and the mixture stirred at 100 °C for 16 hours. Further 1.1'-carbonyldiimidazole (14 mg) was added and the mixture stirred a further 24 hours at 100 °C. The mixture was then allowed to cool
- 15 to room temperature whereupon concentrated ammonia solution (67  $\mu$ L) was added. The mixture was transferred to a sealed tube and heated at 50 °C for 16 hours. To the ambient reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate was added followed by extraction into ethyl acetate. The combined organic layers were washed with water and saturated saline, dried and concentrated. The residue was
- 20 purified by column chromatography (25% - 100% ethyl acetate in heptanes) to afford the title compound having the following physical properties (6 mg).
- LC/MS  $t_R$  1.67 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>a</sup>.

25 Example 605: 2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-5-phenyl-1H-imidazole-4-carbonitrile

- To a dichloromethane (0.5 mL) solution of the compound prepared in Example 604 (6 mg) was added (methoxycarbonylsulfamoyl)triethylammonium hydroxide, inner salt (8.6 mg) and the mixture stirred at room temperature for 2 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate was added
- 30 followed by extraction into dichloromethane. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (80% - 100% ethyl acetate in heptanes) to afford the title compound having the following physical properties (3.5 mg).

LC/MS  $t_R$  4.06 minutes; MS (ES<sup>+</sup>)  $m/z$  481 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 7.80 (d, 2 H), 7.78 - 7.72 (m, 2 H), 7.71 - 7.68 (m, 1 H), 7.54 - 7.49 (m, 2 H), 7.47 - 7.42 (m, 1 H), 6.15 (s, 1 H), 6.10 (s, 1 H), 5.77 (dd, 1 H), 3.48 - 3.38 (m, 1 H), 3.12 (ddd, 1 H), 2.72 - 2.61 (m, 1 H), 2.45 - 2.35 (m, 1 H).

Example 606(1) and 606(2): 3-(2-chloroacetyl)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one and 3-(2-bromoacetyl)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

The compound prepared in Example 9 (0.50 g) was treated as detailed in Example 234 to give a 2: 1 ratio of the title compounds having the following physical properties (0.65 g).

Example 606(1):

LC/MS  $t_R$  1.73 minutes; MS (ES<sup>+</sup>)  $m/z$  412 and 414 (M+Na); 390 and 392 (M+H)<sup>a</sup>.

Example 606(2):

LC/MS  $t_R$  1.76 minutes; MS (ES<sup>+</sup>)  $m/z$  456 and 458 (M+Na); 434 and 436 (M+H)<sup>a</sup>.

Example 607: 4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)benzamide

The 2: 1 mixture of compounds prepared in Example 606 (32 mg) were treated with 4-amidinobenzamide hydrochloride following the method of Example 239 to give the title compound having the following physical properties (3.9 mg).

LC/MS  $t_R$  2.98 minutes; MS (ES<sup>+</sup>)  $m/z$  499 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.61 (s, 1 H), 9.71 (s, 1 H), 8.01 (s, 1 H), 7.97 - 7.91 (m, 4 H), 7.84 - 7.77 (m, 3 H), 7.40 (s, 1 H), 6.99 (s, 1 H), 5.92 (s, 2 H), 5.62 - 5.60 (m, 1 H), 3.02 - 2.90 (m, 1 H), 2.54 - 2.51 (m, 1 H), 2.47 - 2.35 (m, 1 H), 2.33 - 2.21 (m, 1 H).

Example 608: 4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)benzamide

The compound prepared in Example 607 (60 mg) was treated using the method

as detailed in Example 44 to give the title compound having the following physical properties (6.0 mg).

LC/MS  $t_R$  3.64 minutes; MS (ES<sup>+</sup>)  $m/z$  533 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.04 (s, 1 H), 9.66 (s, 1 H), 8.03 (s, 1 H), 7.96 (d, 2 H), 7.92 (d, 2 H), 7.82 (s, 2 H), 7.76 (s, 1 H), 7.42 (s, 1 H), 6.00 (s, 1 H), 5.94 (s, 1 H), 5.66 – 5.63 (m, 1 H), 3.24 – 3.10 (m, 1 H), 3.09 – 2.98 (m, 1 H), 2.64 – 2.55 (m, 1 H), 2.20 – 2.09 (m, 1 H).

Example 609: tert-butyl N-[5-(N-hydroxycarbamimidoyl)pyridin-2-yl]carbamate

To an ethanol (5 mL) solution of tert-butyl 5-cyano-2-pyridinylcarbamate (0.30 g) was added hydroxylamine hydrochloride (0.95 g) followed by N,N-diisopropylethylamine (244  $\mu$ L) and the mixture stirred at 80 °C for 1.5 hours. The reaction mixture was concentrated and azeotroped with chloroform to give the title compound having the following physical properties (0.345 g).

LC/MS  $t_R$  1.21 minutes; MS (ES<sup>+</sup>)  $m/z$  253 (M+H)<sup>a</sup>.

Example 610: (6-[(tert-butoxy)carbonyl]amino}pyridin-3-yl)methanimidamido acetate

To an acetic acid (10 mL) solution of the compound prepared in Example 609 (0.345 g) was added acetic anhydride (3.85 mL) and the mixture stirred at room temperature for 30 minutes. The reaction mixture was concentrated, the residue dissolved in ethyl acetate and washed sequentially with a saturated aqueous solution of sodium hydrogen carbonate and saturated saline, dried and concentrated to obtain the title compound having the following physical properties (0.244 g).

LC/MS  $t_R$  1.60 minutes; MS (ES<sup>+</sup>)  $m/z$  295 (M+H)<sup>a</sup>.

Example 611: acetic acid - tert-butyl N-(5-carbamimidoylpyridin-2-yl)carbamate (1 : 1)

To an ethanol (3 mL) solution of the compound prepared in Example 610 (0.244 g) was added palladium on carbon (50 mg) and the mixture stirred at room temperature under an atmosphere of hydrogen for 1 hour. The reaction mixture was filtered through Celite® and the filtrate concentrated to give the title compound having the following physical properties (0.174 g).

LC/MS  $t_R$  1.15 minutes; MS (ES<sup>+</sup>)  $m/z$  237 (M+H)<sup>a</sup>.

Example 612: tert-butyl N-(5-{5-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl}-1H-imidazol-2-yl}pyridin-2-yl)carbamate

To a tetrahydrofuran (2 mL) and water (0.6 mL) solution of the 2: 1 mixture of compounds prepared in Example 606 (109 mg) was added sodium hydrogen carbonate (85 mg) and the mixture warmed to 70 °C for 30 minutes. The compound prepared in Example 611 (109 mg) was added and the mixture stirred at 70 °C for 3 hours. The reaction mixture was diluted with ethyl acetate and washed sequentially with water and saturated saline, dried and concentrated. The residue was purified by column chromatography (0 - 100% methanol in dichloromethane) to give the title compound having the following physical properties (18.5 mg).  
LC/MS  $t_R$  1.68 minutes; MS (ES<sup>+</sup>)  $m/z$  572 (M+H)<sup>a</sup>.

Example 613: (3S)-3-[2-(6-amino-3-pyridinyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone hydrochloride

The compound prepared in Example 612 (18.5 mg) was treated using the method as detailed in Example 55 to give the title compound having the following physical properties (6.8 mg).  
LC/MS  $t_R$  2.79 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.73 (s, 1 H), 8.79 (s, 1 H), 8.46 (d, 2 H), 7.83 (s, 2 H), 7.77 (s, 1 H), 7.37 (s, 1 H), 7.09 (d, 1 H), 6.00 (s, 2 H), 5.79 – 5.60 (m, 1 H), 3.31 – 3.20 (m, 1 H), 3.06 – 2.94 (m, 1 H), 2.61 – 2.55 (m, 1 H), 2.33 – 2.24 (m, 1 H).

Example 614: (3S)-3-[2-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone hydrochloride

The same operation as in Example 44 → Example 55 was conducted from the compound prepared in Example 612 to give the title compound having the following physical properties.  
LC/MS  $t_R$  3.03 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.08 (s, 1 H), 9.67 (s, 1 H), 8.51 – 8.10 (m, 4 H), 7.84 – 7.79 (m, 2 H), 7.74 (d, 1 H), 7.08 (d, 1 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.63 (dd, 1 H), 3.21 – 2.97 (m, 2 H), 2.62 – 2.06 (m, 2 H).



Example 615: methyl [5-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)-2-pyridinyl]carbamate

The compound prepared in Example 614 (25 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (3.8 mg).

LC/MS  $t_R$  3.70 minutes; MS ( $ES^+$ )  $m/z$  564 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 8.69 (d, 1 H), 8.13 (dd, 1 H), 7.99 (d, 1 H), 7.76 - 7.71 (m, 2 H), 7.71 - 7.66 (m, 1 H), 6.20 (s, 1 H), 6.04 (s, 1 H), 5.82 - 5.77 (m, 1 H), 3.79 (s, 3 H), 3.42 - 3.38 (m, 1 H), 3.15 - 3.10 (m, 1 H), 2.75 - 2.62 (m, 1 H), 2.31 - 2.28 (m, 1 H).


Example 616: methyl [6-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)-3-pyridinyl]carbamate

The same operation as in Example 77 → Example 609 → Example 610 → Example 611 → Example 612 → Example 44 was conducted from 5-aminopyridine-2-carbonitrile to give the title compound having the following physical properties.

LC/MS  $t_R$  3.82 minutes; MS ( $ES^+$ )  $m/z$  564 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.08 (s, 1 H), 10.05 (s, 1 H), 9.64 (s, 1 H), 8.73 (s, 1 H), 7.95 (d, 1 H), 7.88 (d, 1 H), 7.81 (s, 2 H), 7.75 (s, 1 H), 5.99 (s, 1 H), 5.91 (s, 1 H), 5.73 - 5.66 (m, 1 H), 3.72 (s, 3 H), 3.18 - 2.94 (m, 2 H), 2.56 - 2.07 (m, 2 H).

Example 617: tert-butyl N-{4-chloro-2-[3-(2-chloroacetyl)-5-oxo-2,3-dihydro-1H-indolizin-7-yl]phenyl}carbamate



To a dichloromethane (100 mL) solution of the compound prepared in Example 19 (2.5 g) was added 1-chloro-N,N,2-trimethylprop-1-en-1-amine (1.80 ml) and the mixture stirred at room temperature for 1 hour. A 2 M solution of (trimethylsilyl)diazomethane in diethyl ether (6.2 mL) was added and the mixture

stirred at room temperature for 45 minutes. Concentrated hydrochloric acid (0.53 mL) was then added and the mixture stirred vigorously at room temperature for 30 minutes. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (50 mL) was added followed by extraction into dichloromethane. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (20% - 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (1.79 g).  
LC/MS  $t_R$  2.09 minutes; MS (ES<sup>+</sup>)  $m/z$  459 (M+Na), 437 (M+H)<sup>a</sup>.

10 Example 618: methyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate

The same operation as in Example 239 → Example 40 → Example 24 → Example 364 was conducted from the compound prepared in Example 617 to give the title compound having the following physical properties.

15 LC/MS  $t_R$  3.89 minutes; MS (ES<sup>+</sup>)  $m/z$  547 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  9.33 (s, 1 H), 7.80 - 7.63 (m, 5 H), 7.52 (d, 2 H), 6.17 (s, 1 H), 6.04 (s, 1 H), 5.79 (dd, 1 H), 3.75 (s, 3 H), 3.39 - 3.32 (obs. m, 1 H), 3.18 - 3.07 (m, 1 H), 2.65 (qd, 1 H), 2.48 - 2.37 (m, 1 H).

20 Example 619: prop-2-en-1-yl N-(4-carbamimidoylphenyl)carbamate

The same operation as in Example 225 → Example 237 was conducted from 4-aminobenzonitrile to give the title compound having the following physical properties. (Note: in the step corresponding to Example 225 in the operation, allyl chloroformate was used in place of methyl chloroformate).

25 LC/MS  $t_R$  1.05 minutes; MS (ES<sup>+</sup>)  $m/z$  220 (M+H)<sup>a</sup>.

Example 620: tert-butyl N-(4-chloro-2-{5-oxo-3-[2-(4-{[(prop-2-en-1-yloxy)carbonyl]amino}phenyl)-1H-imidazol-5-yl]-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

30 To a dimethylsulfoxide (3.5 mL) solution of the compounds prepared in Example 617 (200 mg) and Example 619 (83 mg) was added potassium carbonate (135 mg). The mixture was stirred at room temperature for 2.5 hours then at 90 °C for 1 hour. To the reaction mixture, water (30 mL) was added followed by extraction into ethyl

acetate. The combined organic layers were washed with water and saturated saline, dried and concentrated. The residue was purified by column chromatography (0% to 10% methanol in dichloromethane) to give the title compound having the following physical properties (84.9 mg).

- 5 LC/MS  $t_R$  1.75 minutes; MS ( $ES^+$ )  $m/z$  602 ( $M+H$ )<sup>a</sup>.

Example 621: (3S)-3-[2-(4-aminophenyl)-4-chloro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 40 → Example 24 → Example 44 →

- 10 Example 584 was conducted from the compound prepared in Example 620 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.29 minutes; MS ( $ES^+$ )  $m/z$  505 ( $M+H$ )<sup>b</sup>


- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.34 (br. s, 1 H), 9.65 (s, 1 H), 7.84 - 7.78 (m, 3 H), 7.77 - 7.73 (m, 1 H), 7.51 (d, 2 H), 6.60 (d, 2 H), 5.98 (s, 1 H), 5.91 (s, 1 H), 5.60 (d, 1  
15 H), 5.66 - 5.40 (m, 1 H), 3.20 - 3.09 (m, 1 H), 3.04 - 2.94 (m, 1 H), 2.60 - 2.49 (m, 1 H), 2.14 - 2.05 (m, 1 H).

Example 622: (3S)-7-(2-{bis[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

- 20 To a methanol solution (130 mL) of the compound prepared in Example 10 (17.54 g) was added 2 M sodium hydroxide (54.6 mL) and the mixture stirred at room temperature for 1 hour 30 minutes. The methanol was removed *in vacuo* and the residual aqueous solution diluted with water (200 mL) and washed with dichloromethane. The aqueous layer was then treated with 2 M hydrochloric acid (30  
25 mL) and extracted into ethyl acetate. The combined organic layers were dried and concentrated to give the title compound having the following physical properties (14.75 g).

LC/MS  $t_R$  2.05 minutes; MS ( $ES^+$ )  $m/z$  405 ( $M-Boc+H$ ), 527 ( $M+Na$ )<sup>a</sup>.

- 30 Example 623: tert-butyl N-[(tert-butoxy)carbonyl]-N-{4-chloro-2-[3-(2-chloroacetyl)-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl]phenyl}carbamate



To a dichloromethane (160 mL) solution of the compound prepared in Example 617 (6.0 g) was added 1-chloro-N,N,2-trimethylprop-1-en-1-amine (3.46 mL) and the mixture stirred at room temperature for 1 hour. A 2 M solution of

- 5 (trimethylsilyl)diazomethane in diethyl ether (11.9 mL) was added and the reaction mixture stirred at room temperature for 45 minutes. Concentrated hydrochloric acid (1.0 mL) was then added dropwise and the reaction stirred vigorously at room temperature for 30 minutes. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (250 mL) was added followed by extraction into dichloromethane.
- 10 The combined organic layers were dried and concentrated to afford the title compound having the following physical properties (8.87 g).

LC/MS  $t_R$  2.28 minutes; MS ( $ES^+$ )  $m/z$  437 (M-Boc+H), 559 (M+Na) <sup>a</sup>.

Example 624: (3S)-3-[2-(4-aminophenyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

- 15

The same operation as in Example 620 → Example 40 → Example 24 → Example 584 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.88 minutes; MS ( $ES^+$ )  $m/z$  471 (M+H) <sup>b</sup>

- 20 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.93 (br. s, 1 H), 9.69 (s, 1 H), 7.82 - 7.77 (m, 3 H), 7.53 (d, 2 H), 6.73 (br. s, 1 H), 6.57 (d, 2 H), 5.93 - 5.88 (m, 2 H), 5.56 (d, 1 H), 5.31 (br. s, 2 H), 3.38 - 3.27 (m, 1 H), 2.95 - 2.87 (m, 1 H), 2.43 - 2.34 (m, 1 H), 2.27 - 2.20 (m, 1 H).

- 25 Example 625: 2-methoxyethyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-2-yl)phenyl]carbamate

To a dichloromethane (1 mL) solution of the compound prepared in Example 624 (30 mg) was added N,N-diisopropylethylamine (23  $\mu$ L) followed by 2-

methoxyethyl chloroformate (7.5  $\mu$ L) and the reaction mixture stirred at room temperature for 16 hours. Further N,N-diisopropylethylamine (13  $\mu$ L) and 2-methoxyethyl chloroformate (7.5  $\mu$ L) was added and the mixture stirred at room temperature for 2 hours. To the reaction mixture, water (10 mL) was added followed by  
5 extraction into dichloromethane. The combined organic layers were dried and concentrated, the residue dissolved in methanol (1 mL) and treated with concentrated ammonia solution (0.20 mL). After stirring 1 hour at room temperature the mixture was concentrated and the residue purified by column chromatography (0 to 6% methanol in dichloromethane) to give the title compound having the following physical properties  
10 (22 mg).  
LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  573 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.39 (s, 1 H), 7.78 - 7.72 (m, 4 H), 7.71 - 7.67 (m, 1 H), 7.53 (d, 2 H), 6.74 (s, 1 H), 6.15 (s, 1 H), 6.07 (s, 1 H), 5.80 (d, 1 H), 4.31 - 4.25 (m, 2 H), 3.68 - 3.63 (m, 2 H), 3.39 (s, 3 H), 3.36 - 3.31 (m, 1 H), 3.08 - 3.00 (m, 1 H), 2.62  
15 - 2.51 (m, 1 H), 2.44 - 2.36 (m, 1 H).

Example 626: 2-(2-methoxyethoxy)ethyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-2-yl)phenyl]carbamate

The compound prepared in Example 624 (30 mg) was treated as detailed in  
20 Example 625 using the chloroformate prepared in Example 489 to give the title compound having the following physical properties (15 mg).  
LC/MS  $t_R$  3.09 minutes; MS (ES<sup>+</sup>)  $m/z$  617 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.39 (s, 1 H), 7.77 - 7.72 (m, 4 H), 7.71 - 7.67 (m, 1 H), 7.53 (d, 2 H), 6.74 (s, 1 H), 6.15 (s, 1 H), 6.07 (s, 1 H), 5.80 (d, 1 H), 4.31 - 4.27 (m,  
25 2 H), 3.76 - 3.72 (m, 2 H), 3.68 - 3.64 (m, 2 H), 3.57 - 3.54 (m, 2 H), 3.36 (s, 3 H), 3.34 - 3.29 (m, 1 H), 3.08 - 3.00 (m, 1 H), 2.61 - 2.51 (m, 1 H), 2.43 - 2.36 (m, 1 H).

Example 627: (1E)-N-[4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-2-yl)phenyl]-N'-hydroxyethanimidamide

The compound prepared in Example 624 (35 mg) was treated as detailed in  
30 Example 462 to give the title compound having the following physical properties (1.4 mg).

LC/MS  $t_R$  2.67 minutes; MS (ES<sup>+</sup>)  $m/z$  528 (M+H), 264.5 (M/2+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.34 (br. s, 0.3 H), 12.20 (br. s, 0.4 H), 12.07 (br. s, 0.3 H), 9.71 - 9.68 (m, 1.3 H), 9.62 (s, 0.3 H), 9.27 (s, 0.4 H), 8.21 (s, 0.4 H), 8.01 (s, 0.3 H), 7.79 (d, 3.3 H), 7.74 (d, 0.6 H), 7.69 (d, 0.8 H), 7.54 (d, 0.8 H), 7.52 - 7.47 (m, 0.6 H), 7.14 (d, 0.6 H), 6.86 (d, 0.3 H), 6.81 - 6.75 (m, 1.3 H), 5.93 - 5.86 (m, 2 H), 5.60 - 5.52 (m, 1 H), 3.42 - 3.29 (m, 1 H), 2.97 - 2.86 (m, 1 H), 2.44 - 2.37 (m, 1 H), 2.30 - 2.21 (m, 1 H), 1.98 (s, 1.2 H), 1.92 (s, 1.8 H).

Example 628: N-[4-(4-chloro-5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-2-yl)phenyl]acetamide

10        The compound prepared in Example 621 (25 mg) was treated with acetyl chloride following the method of Example 484 to give the title compound having the following physical properties (18.8 mg).

LC/MS *t<sub>R</sub>* 3.63 minutes; MS (ES<sup>+</sup>) *m/z* 547 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.34 (s, 1 H), 7.77 - 7.70 (m, 4 H), 7.68 (d, 1 H),  
15    7.65 (d, 2 H), 6.19 (s, 1 H), 6.03 (s, 1 H), 5.79 (dd, 1 H), 3.38 - 3.32 (obs. m, 1 H), 3.12 (ddd, 1 H), 2.74 - 2.62 (m, 1 H), 2.36 - 2.26 (m, 1 H), 2.14 (s, 3 H).

Example 629: 2-methoxyethyl [4-(4-chloro-5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-2-yl)phenyl]carbamate

20        The compound prepared in Example 621 (25 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (17.1 mg).

LC/MS *t<sub>R</sub>* 3.91 minutes; MS (ES<sup>+</sup>) *m/z* 607 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.34 (s, 1 H), 7.76 - 7.70 (m, 4 H), 7.68 (d, 1 H),  
25    7.53 (d, 2 H), 6.20 (br. s, 1 H), 6.03 (s, 1 H), 5.79 (dd, 1 H), 4.31 - 4.25 (m, 2 H), 3.68 - 3.62 (m, 2 H), 3.39 (s, 3 H), 3.43 - 3.33 (obs. m, 1 H), 3.18 - 3.06 (m, 1 H), 2.74 - 2.61 (m, 1 H), 2.37 - 2.25 (m, 1 H).

Example 630: 3-methoxypropyl [4-(4-chloro-5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-

30    yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-2-yl)phenyl]carbamate

The compound prepared in Example 621 (30 mg) was treated with the chloroformate prepared in Example 487 following the method of Example 625 to give the title compound having the following physical properties (12 mg).

LC/MS  $t_R$  3.87 minutes; MS ( $ES^+$ )  $m/z$  621 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.69 (br. s, 1 H), 9.83 (br. s, 1 H), 9.65 (s, 1 H), 7.81 (s, 2 H), 7.77 - 7.72 (m, 3 H), 7.53 (d, 2 H), 5.99 (br. s, 1 H), 5.91 (s, 1 H), 5.62 (d, 1 H), 4.14 (t, 2 H), 3.43 (t, 2 H), 3.25 (s, 3 H), 3.19 - 3.10 (m, 1 H), 3.05 - 2.96 (m, 1 H), 2.61 - 2.55 (m, 1 H), 2.16 - 2.08 (m, 1 H), 1.86 (quintet, 2 H).

Example 631: 2-(2-methoxyethoxy)ethyl [4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate

10 The compound prepared in Example 621 (30 mg) was treated with the chloroformate prepared in Example 489 following the method of Example 625 to give the title compound having the following physical properties (15 mg).

LC/MS  $t_R$  3.76 minutes; MS ( $ES^+$ )  $m/z$  651 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.69 (br. s, 1 H), 9.95 (br. s, 1 H), 9.65 (s, 1 H), 7.85 - 7.72 (m, 5 H), 7.54 (d, 2 H), 5.99 (s, 1 H), 5.91 (s, 1 H), 5.63 (dd, 1 H), 4.23 - 4.19 (m, 2 H), 3.67 - 3.63 (m, 2 H), 3.58 - 3.54 (m, 2 H), 3.47 - 3.43 (m, 2 H), 3.24 (s, 3 H), 3.20 - 3.10 (m, 1 H), 3.05 - 2.96 (m, 1 H), 2.62 - 2.54 (m, 1 H), 2.17 - 2.08 (m, 1 H).

Example 632: 3-oxetanyl [4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate

20 The compound prepared in Example 621 (30 mg) was treated as detailed in Example 494 to give the title compound having the following physical properties (15 mg).

LC/MS  $t_R$  3.67 minutes; MS ( $ES^+$ )  $m/z$  605 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.72 (s, 1 H), 10.11 (br. s, 1 H), 9.64 (s, 1 H), 7.83 - 7.80 (m, 2 H), 7.79 - 7.72 (m, 3 H), 7.53 (d, 2 H), 5.99 (s, 1 H), 5.91 (s, 1 H), 5.63 (dd, 1 H), 5.43 (quintet, 1 H), 4.83 (t, 2 H), 4.56 (dd, 2 H), 3.19 - 3.10 (m, 1 H), 3.01 (ddd, 1 H), 2.60 - 2.55 (m, 1 H), 2.16 - 2.08 (m, 1 H).

30 Example 633: (1E)-N-[4-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]-N'-hydroxyethanimidamide

The compound prepared in Example 621 (46 mg) was treated as detailed in Example 462 to give the title compound having the following physical properties (2.5 mg).

LC/MS  $t_R$  3.09 minutes; MS ( $ES^+$ )  $m/z$  562 (M+H), 281.5 (M/2+H) <sup>b</sup>

5 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 7.79 - 7.62 (m, 6 H), 7.19 (d, 2 H), 6.20 (s, 1 H), 6.04 (s, 1 H), 5.80 (ddd, 1 H), 3.36 - 3.30 (m, 1 H), 3.18 - 3.08 (m, 1 H), 2.73 - 2.65 (m, 1 H), 2.36 - 2.27 (m, 1 H), 2.00 (s, 3 H).

10 Example 634: 3-[2-(4-aminophenyl)-4-fluoro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 620 → Example 40 → Example 24 → Example 364 → Example 584 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.65 minutes; MS ( $ES^+$ )  $m/z$  489 (M+H) <sup>a</sup>.

15

Example 635: 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate

The compound prepared in Example 634 (28 mg) was treated with as detailed in Example 625 to give the title compound having the following physical properties (11 mg).

20

LC/MS  $t_R$  3.86 minutes; MS ( $ES^+$ )  $m/z$  591 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.33 (s, 1 H), 7.76 - 7.66 (m, 5 H), 7.53 (d, 2 H), 6.18 (s, 1 H), 6.04 (s, 1 H), 5.79 (dd, 1 H), 4.30 - 4.26 (m, 2 H), 3.67 - 3.63 (m, 2 H), 3.39 (s, 3 H), 3.34 - 3.30 (m, 1 H), 3.16 - 3.09 (m, 1 H), 2.70 - 2.60 (m, 1 H), 2.46 - 2.39 (m, 1 H).

25

Example 636: 2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate

30

The compound prepared in Example 634 (28 mg) was treated with the chloroformate prepared in Example 489 following the method of Example 625 to give the title compound having the following physical properties (12 mg).

LC/MS  $t_R$  3.86 minutes; MS ( $ES^+$ )  $m/z$  635 (M+H) <sup>b</sup>



<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.78 - 7.67 (m, 5 H), 7.54 (d, 2 H), 6.19 (s, 1 H), 6.06 (s, 1 H), 5.81 (dd, 1 H), 4.33 - 4.28 (m, 2 H), 3.78 - 3.74 (m, 2 H), 3.70 - 3.66 (m, 2 H), 3.59 - 3.55 (m, 2 H), 3.38 (s, 3 H), 3.35 - 3.32 (m, 1 H), 3.18 - 3.10 (m, 1 H), 2.71 - 2.61 (m, 1 H), 2.48 - 2.40 (m, 1 H).

5

Example 637: prop-2-en-1-yl N-(6-chloro-5-cyanopyridin-2-yl)carbamate

To an acetonitrile (40 mL) solution of 6-amino-2-chloropyridine-3-carbonitrile [Bioorg. Med. Chem. Lett., 20(5), 1697 (2010)] (1.56 g) was added allyl chloroformate (2.71 mL) followed by N,N-diisopropylethylamine (5.0 mL) and 4-  
10 dimethylaminopyridine (0.12 g) and the mixture stirred at room temperature for 72 hours. Further aliquots of allyl chloroformate (1.1 mL) and N,N-diisopropylethylamine (1.67 mL) were then added and the reaction mixture stirred at room temperature for 29 hours. The mixture was concentrated and the residue purified by column chromatography (0 – 50% ethyl acetate in heptanes) to give the title compound having  
15 the following physical properties (1.31 g).  
LC/MS *t<sub>R</sub>* 1.90 minute; MS (ES<sup>+</sup>) *m/z* 238 (M+H)<sup>+</sup>.

Example 638: prop-2-en-1-yl N-(5-carbamimidoyl-6-chloropyridin-2-yl)carbamate

The same operation as in Example 609 → Example 610 → Example 611 was  
20 conducted from the compound prepared in Example 619 to give the title compound having the following physical properties.  
LC/MS *t<sub>R</sub>* 0.92 minute; MS (ES<sup>+</sup>) *m/z* 255 (M+H)<sup>+</sup>.

Example 639: (3S)-3-[2-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 620 → Example 40 → Example 24 →  
Example 584 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties. (Note: in the step  
corresponding to Example 620 in the operation, the compound prepared in Example 638  
30 was used).  
LC/MS *t<sub>R</sub>* 2.90 minutes; MS (ES<sup>+</sup>) *m/z* 506 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.86 (s, 1 H), 9.69 (s, 1 H), 7.82 - 7.77 (m, 3 H), 7.75 (d, 1 H), 6.87 (d, 1 H), 6.65 (s, 1 H), 6.47 (d, 1 H), 5.91 (s, 1 H), 5.89 (s, 1 H), 5.58

(d, 1 H), 3.31 - 3.26 (m, 1 H), 2.95 - 2.88 (m, 1 H), 2.42 - 2.37 (m, 1 H), 2.30 - 2.23 (m, 1 H).

Example 640: 6-amino-2-fluoropyridine-3-carbonitrile

5 To an N,N-dimethylformamide (60 mL) suspension of 6-fluoro-5-iodopyridin-2-amine [J. Org. Chem., 71(7), 2922 (2006)] (8.0 g) was added zinc cyanide (4.15 g) and the mixture degassed with nitrogen for 10 minutes. Tetrakis(triphenylphosphine)palladium(0) (1.94 g) was then added and the reaction heated at 90 °C for 2 hours 30 minutes. To the ambient reaction mixture, water (300 mL) was added followed  
10 by extraction into ethyl acetate. The combined organic layers were dried and concentrated and the residue purified by column chromatography (0 to 50% ethyl acetate in heptanes) to give the title compound having the following physical properties (3.42 g).

LC/MS  $t_R$  1.18 minute; MS ( $ES^+$ )  $m/z$  138 (M+H)<sup>a</sup>.

15

Example 641: prop-2-en-1-yl N-(5-carbamimidoyl-6-fluoropyridin-2-yl)carbamate

The same operation as in Example 637 → Example 609 → Example 610 → Example 611 was conducted from the compound prepared in Example 640 to give the title compound having the following physical properties.

20 LC/MS  $t_R$  0.93 minute; MS ( $ES^+$ )  $m/z$  239 (M+H)<sup>a</sup>.

Example 642: (3S)-3-[2-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 620 → Example 40 → Example 24 →  
25 Example 584 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 620 in the operation, the compound prepared in Example 641 was used).

LC/MS  $t_R$  2.87 minutes; MS ( $ES^+$ )  $m/z$  490 (M+H)<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.73 (s, 1 H), 9.70 (s, 1 H), 7.96 (dd, 1 H), 7.83 - 7.77 (m, 3 H), 6.81 (d, 1 H), 6.65 (s, 1 H), 6.41 (dd, 1 H), 5.92 (s, 1 H), 5.89 (s, 1 H), 5.58 (d, 1 H), 3.32 - 3.27 (m, 1 H), 2.95 - 2.88 (m, 1 H), 2.43 - 2.34 (m, 1 H), 2.29 - 2.21 (m, 1 H).

Example 643: (3S)-3-[2-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 620 → Example 40 → Example 24 →

- 5 Example 44 → Example 584 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 620 in the operation, the compound prepared in Example 641 was used).

LC/MS  $t_R$  3.58 minutes; MS ( $ES^+$ )  $m/z$  524 ( $M+H$ )<sup>b</sup>

- 10 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.18 (br. s, 1 H), 9.65 (s, 1 H), 7.95 - 7.88 (m, 1 H), 7.83 - 7.80 (m, 2 H), 7.75 (s, 1 H), 6.76 (s, 2 H), 6.40 (dd, 1 H), 6.00 (s, 1 H), 5.91 (s, 1 H), 5.72 (d, 1 H), 3.15 - 3.04 (m, 1 H), 3.03 - 2.95 (m, 1 H), 2.56 - 2.47 (m, 1 H), 2.17 - 2.09 (m, 1 H).

- 15 Example 644: 2-(2-methoxyethoxy)ethyl N-(5-cyanopyridin-2-yl)carbamate

- To an acetonitrile (150 mL) solution of 6-aminopyridine-3-carbonitrile (3.70 g) was added the chloroformate prepared in Example 489 (7.94 g) followed by N,N-diisopropylethylamine (7.72 mL) and the mixture stirred at room temperature for 16 hours. The precipitate formed in the reaction mixture was isolated by filtration to give the title compound (2.12 g). To the filtrate, water (300 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was suspended in acetonitrile (25 mL), concentrated ammonia solution (5 mL) added and the mixture stirred at room temperature for 16 hours. The resultant precipitate was isolated by filtration to give a second batch of the title compound (2.52 g) having the following physical properties. LC/MS  $t_R$  1.42 minutes; MS ( $ES^+$ )  $m/z$  288 ( $M+Na$ ), 266 ( $M+H$ )<sup>a</sup>.

Example 645: 2-(2-methoxyethoxy)ethyl N-(5-carbamimidoylpyridin-2-yl)carbamate

- The same operation as in Example 609 → Example 610 → Example 611 was conducted from the compound prepared in Example 644 to give the title compound having the following physical properties.

LC/MS  $t_R$  0.84 minutes; MS ( $ES^+$ )  $m/z$  283 ( $M+H$ )<sup>a</sup>.

Example 646: 2-(2-methoxyethoxy)ethyl [5-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l}-4-fluoro-1H-imidazol-2-yl)-2-pyridinyl]carbamate

The same operation as in Example 620 → Example 40 → Example 24 →

- 5 Example 364 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 620, the compound prepared in Example 645 was used).

LC/MS  $t_R$  3.74 minutes; MS (ES<sup>+</sup>)  $m/z$  636 (M+H)<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.75 (br. s, 1 H), 10.41 (s, 1 H), 9.66 (s, 1 H), 8.76 -  
10 8.66 (m, 1 H), 8.15 (dd, 1 H), 7.89 (d, 1 H), 7.83 - 7.78 (m, 2 H), 7.78 - 7.75 (m, 1 H),  
5.97 (s, 1 H), 5.93 (s, 1 H), 5.63 (dd, 1 H), 4.26 - 4.20 (m, 2 H), 3.67 - 3.62 (m, 2 H),  
3.58 - 3.54 (m, 2 H), 3.47 - 3.43 (m, 2 H), 3.24 (s, 3 H), 3.24 - 3.16 (m, 1 H), 3.00 (ddd,  
1 H), 2.62 - 2.53 (m, 1 H), 2.23 - 2.16 (m, 1 H).

- 15 Example 647: 2-(2-methoxyethoxy)ethyl [5-(4-chloro-5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l}-1H-imidazol-2-yl)-2-pyridinyl]carbamate

The same operation as in Example 620 → Example 40 → Example 24 →

- Example 44 was conducted from the compound prepared in Example 623 to give the  
20 title compound having the following physical properties. (Note: in the step corresponding to Example 620, the compound prepared in Example 645 was used).

LC/MS  $t_R$  3.67 minutes; MS (ES<sup>+</sup>)  $m/z$  652 and 654 (M+H), 326.5 and 327.5

[(M+2H)/2]<sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.00 (br. s, 1 H), 10.43 (s, 1 H), 9.65 (s, 1 H), 8.72  
25 (d, 1 H), 8.16 (dd, 1 H), 7.90 (d, 1 H), 7.85 - 7.78 (m, 2 H), 7.75 (app. s, 1 H), 5.98 (s, 1  
H), 5.92 (s, 1 H), 5.63 (dd, 1 H), 4.25 - 4.21 (m, 2 H), 3.67 - 3.63 (m, 2 H), 3.58 - 3.54  
(m, 2 H), 3.46 - 3.43 (m, 2 H), 3.24 (s, 3 H), 3.22 - 3.12 (m, 1 H), 3.05 - 2.97 (m, 1 H),  
2.61 - 2.53 (m, 1 H), 2.17 - 2.08 (m, 1 H).

- 30 Example 648: 2-methoxyethyl N-(5-carbamimidoylpyridin-2-yl)carbamate

The same operation as in Example 644 → Example 609 → Example 610 →

Example 611 was conducted from 6-aminopyridine-3-carbonitrile to give the title

compound having the following physical properties. (Note: in the step corresponding to Example 644, 2-methoxyethyl chloroformate was used).

LC/MS  $t_R$  0.66 minutes; MS (ES<sup>+</sup>)  $m/z$  239 (M+H)<sup>a</sup>.

- 5    Example 649: tert-butyl N-[(tert-butoxy)carbonyl]-N-(4-chloro-2-{3-[2-(6-[(2-methoxyethoxy)carbonyl]amino}pyridin-3-yl)-1H-imidazol-5-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}phenyl)carbamate

10    The compound prepared in Example 623 (3.55 g) was treated with the compound prepared in Example 648 following the method of Example 620 to give the title compound having the following physical properties (1.09 g).

LC/MS  $t_R$  1.91 minutes; MS (ES<sup>+</sup>)  $m/z$  721 (M+H), 621 {M-[CO<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>]+H}<sup>a</sup>.

Example 650: 2-methoxyethyl [5-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-2-yl)-2-pyridinyl]carbamate

- 15    The same operation as in Example 40 → Example 24 was conducted from the compound prepared in Example 649 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.96 minutes; MS (ES<sup>+</sup>)  $m/z$  574 (M+H), 287.5 [(M+2H)/2]<sup>b</sup>

- 20    <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.50 (s, 1 H), 10.36 (s, 1 H), 9.69 (s, 1 H), 8.74 (d, 1 H), 8.18 (dd, 1 H), 7.88 (d, 1 H), 7.82 - 7.77 (m, 3 H), 6.93 (d, 1 H), 5.92 (s, 1 H), 5.90 (s, 1 H), 5.60 (d, 1 H), 4.26 - 4.21 (m, 2 H), 3.59 - 3.55 (m, 2 H), 3.28 (s, 3 H), 3.31 - 3.26 (obs. m, 1 H), 2.93 (dd, 1 H), 2.45 - 2.39 (m, 1 H), 2.29 - 2.22 (m, 1 H).

- 25    Example 651: 2-methoxyethyl [5-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4-fluoro-1H-imidazol-2-yl)-2-pyridinyl]carbamate

The compound prepared in Example 650 (380 mg) was treated as detailed in Example 364 to give the title compound having the following physical properties (21.3 mg).

- 30    LC/MS  $t_R$  3.75 minutes; MS (ES<sup>+</sup>)  $m/z$  592 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.75 (br. s, 1 H), 10.40 (s, 1 H), 9.67 (s, 1 H), 8.71 (d, 1 H), 8.15 (dd, 1 H), 7.89 (d, 1 H), 7.83 - 7.77 (m, 2 H), 7.77 - 7.74 (m, 1 H), 5.97 (s,

1 H), 5.93 (s, 1 H), 5.63 (dd, 1 H), 4.24 (dd, 2 H), 3.57 (dd, 2 H), 3.28 (s, 3 H), 3.21 (td, 1 H), 3.00 (ddd, 1 H), 2.62 - 2.53 (m, 1 H), 2.24 - 2.16 (m, 1 H).

Example 652: 2-methoxyethyl [5-(4-chloro-5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-2-yl)-2-pyridinyl]carbamate

The compound prepared in Example 650 (100 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (40.4 mg).

LC/MS  $t_R$  3.68 minutes; MS ( $ES^+$ )  $m/z$  608 and 610 (M+H), 304.5 and 305.5 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.02 (br. s, 1 H), 10.42 (s, 1 H), 9.65 (s, 1 H), 8.72 (d, 1 H), 8.16 (dd, 1 H), 7.90 (d, 1 H), 7.84 - 7.78 (m, 2 H), 7.75 (app. s, 1 H), 5.98 (s, 1 H), 5.92 (s, 1 H), 5.63 (dd, 1 H), 4.27 - 4.21 (m, 2 H), 3.60 - 3.55 (m, 2 H), 3.28 (s, 3 H), 3.22 - 3.13 (m, 1 H), 3.01 (ddd, 1 H), 2.63 - 2.54 (m, 1 H), 2.17 - 2.09 (m, 1 H).

Example 653: tert-butyl N-(2-{3-[2-(6-aminopyridin-3-yl)-1H-imidazol-5-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl}-4-chlorophenyl)carbamate

To a methanol (5.5 mL) solution of the compound prepared in Example 649 (185 mg) was added 6 M sodium hydroxide (5.5 mL) and the mixture stirred at room temperature for 24 hours. The methanol was removed *in vacuo* and the aqueous residue treated with 1 M hydrochloric acid (11 mL) and water (20 mL) followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue obtained on concentration was purified by column chromatography (0 to 10% methanol in dichloromethane) to obtain the title compound having the following physical properties (107 mg).

LC/MS  $t_R$  1.51 minutes; MS ( $ES^+$ )  $m/z$  519 (M+H), 463 {M-[C(CH<sub>3</sub>)<sub>3</sub>]+H}<sup>a</sup>.

Example 654: (3S)-3-[2-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 637 → Example 40 → Example 24 → Example 364 → Example 584 was conducted from the compound prepared in Example 653 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  490 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.17 (s, 1 H), 9.66 (s, 1 H), 8.38 (d, 1 H), 7.83 – 7.78 (m, 2 H), 7.78 – 7.73 (m, 2 H), 6.47 (d, 1 H), 6.26 (br. s, 2 H), 5.96 (s, 1 H), 5.91 (s, 1 H), 5.59 (dd, 1 H), 3.20 (td, 1 H), 2.98 (ddd, 1 H), 2.59 – 2.52 (m, 1 H), 2.21 – 2.14 (m, 1 H).

Example 655: (3S)-3-[5-(4-aminophenyl)-4H-1,2,4-triazol-3-yl]-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 242 → Example 243 → Example 74 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.54 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>a</sup>.

Example 656(1) to 656(4)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 655 and the corresponding acid chlorides and chloroformates using the method as detailed in Example 484.

Example 656(1): methyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4H-1,2,4-triazol-3-yl)phenyl]carbamate

LC/MS  $t_R$  3.61 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.85 (d, 2 H), 7.77 – 7.72 (m, 2 H), 7.69 (d, 1 H), 7.58 (d, 2 H), 6.17 (s, 1 H), 6.09 (s, 1 H), 5.85 (dd, 1 H), 3.76 (s, 3 H), 3.40 – 3.33 (obs. m, 1 H), 3.10 (ddd, 1 H), 2.73 – 2.62 (m, 1 H), 2.37 – 2.27 (m, 1 H).

Example 656(2): formic acid - 2-fluoroethyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-4H-1,2,4-triazol-3-yl)phenyl]carbamate (1:1)

LC/MS  $t_R$  3.72 minutes; MS (ES<sup>+</sup>)  $m/z$  562 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.86 (d, 2 H), 7.78 – 7.72 (m, 2 H), 7.70 (d, 1 H), 7.60 (d, 2 H), 6.17 (s, 1 H), 6.09 (s, 1 H), 5.85 (dd, 1 H), 4.71 – 4.66 (m, 1 H), 4.62 – 4.56 (m, 1 H), 4.44 – 4.39 (m, 1 H), 4.38 – 4.33 (m, 1 H), 3.41 – 3.23 (m, 1 H), 3.10 (dd, 1 H), 2.73 – 2.62 (m, 1 H), 2.32 (app. br. s, 1 H).

Example 656(3): formic acid - 2-methoxyethyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4H-1,2,4-triazol-3-yl)phenyl]carbamate (1:1)

- 5 LC/MS  $t_R$  3.64 minutes; MS ( $ES^+$ )  $m/z$  574 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  14.12 (br. s, 1 H), 9.99 (br. s, 1 H), 9.69 (s, 1 H), 7.84 (d, 2 H), 7.82 - 7.75 (m, 3 H), 7.58 (d, 2 H), 5.97 (s, 1 H), 5.94 (s, 1 H), 5.67 (dd, 1 H), 4.25 - 4.18 (m, 2 H), 3.60 - 3.53 (m, 2 H), 3.28 (s, 3 H), 3.27 - 3.18 (m, 1 H), 2.98 (dd, 1 H), 2.59 - 2.52 (obs. m, 1 H), 2.16 - 2.05 (m, 1 H).

10

Example 656(4): N-[4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4H-1,2,4-triazol-3-yl)phenyl]acetamide

- LC/MS  $t_R$  3.51 minutes; MS ( $ES^+$ )  $m/z$  514 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.88 (d, 2 H), 7.76 - 7.72 (m, 2 H),  
15 7.72 - 7.67 (m, 3 H), 6.16 (s, 1 H), 6.09 (s, 1 H), 5.85 (dd, 1 H), 3.40 - 3.32 (obs. m, 1 H), 3.10 (ddd, 1 H), 2.73 - 2.62 (m, 1 H), 2.37 - 2.28 (m, 1 H), 2.15 (s, 3 H).

Example 657: formic acid - 3-oxetanyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-4H-1,2,4-triazol-3-yl)phenyl]carbamate (1:1)

- 20 The compound prepared in Example 655 (75 mg) was treated as detailed in Example 494 to give the title compound having the following physical properties (12.9 mg).

- LC/MS  $t_R$  3.65 minutes; MS ( $ES^+$ )  $m/z$  572 ( $M+H$ )<sup>b</sup>  
25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.86 (d, 2 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.58 (d, 2 H), 6.16 (s, 1 H), 6.09 (s, 1 H), 5.85 (dd, 1 H), 5.53 - 5.46 (m, 1 H), 4.93 (t, 2 H), 4.69 (dd, 2 H), 3.39 - 3.32 (obs. m, 1 H), 3.10 (ddd, 1 H), 2.72 - 2.62 (m, 1 H), 2.37 - 2.28 (m, 1 H).

- 30 Example 658: (3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carbonitrile



The same operation as in Example 114 → Example 605 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.60 minutes; MS ( $ES^+$ )  $m/z$  339 (M+H) 361 (M+Na)<sup>a</sup>.

5

Example 659: (3S)-N-amino-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboximidamide

To a methanol (2 mL) suspension of the compound prepared in Example 658 (90 mg) was added hydrazine hydrate (120  $\mu$ L) and the mixture stirred at 60 °C for 1.5 hours. The reaction mixture was concentrated to give the crude title compound having the following physical properties (95 mg).

10

LC/MS  $t_R$  1.16 minutes; MS ( $ES^+$ )  $m/z$  371 (M+H)<sup>a</sup>.

Example 660: 2-methoxyethyl [3-chloro-4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}] -4H-1,2,4-triazol-3-yl)phenyl]carbamate

15

The same operation as in Example 242 → Example 243 → Example 74 → Example 484 was conducted from the compound prepared in Example 659 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 242 in the operation, 2-chloro-4-nitrobenzoic acid was used. In the step corresponding to Example 484 in the operation, 2-methoxyethyl chloroformate was used).

20

LC/MS  $t_R$  3.76 minutes; MS ( $ES^+$ )  $m/z$  608 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  14.02 (br. s, 1 H), 10.20 (br. s, 1 H), 9.69 (s, 1 H), 7.82 - 7.78 (m, 3 H), 7.76 - 7.68 (m, 2 H), 7.51 (d, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.71 (d, 1 H), 4.26 - 4.21 (m, 2 H), 3.58 (dd, 2 H), 3.28 (s, 3 H), 3.21 - 3.13 (m, 1 H), 3.03 - 2.94 (m, 1 H), 2.59 - 2.54 (m, 1 H), 2.19 - 2.09 (m, 1 H).

25

Example 661: tert-butyl N-[(tert-butoxy)carbonyl]-N-(6-fluoro-5-iodopyridin-2-yl)carbamate

30

The compound prepared in Example 640 (5.0 g) was treated as detailed in Example 228 to give the title compound having the following physical properties (5.26 g).

LC/MS  $t_R$  2.47 minutes; MS ( $ES^+$ )  $m/z$  461 ( $M+Na$ ) 283 [ $M-boc-tBu$ ] $^{+a}$ .

Example 662(1) and 662(2): 6-{bis[(tert-butoxy)carbonyl]amino}-2-fluoropyridine-3-carboxylic acid and 6-[[[(tert-butoxy)carbonyl]amino]-2-fluoropyridine-3-carboxylic acid

To a cooled (0 °C) tetrahydrofuran (40 mL) solution of the compound prepared in Example 661 (2.0 g) was added a 2 M solution of isopropylmagnesium chloride in tetrahydrofuran (2.28 mL) and the mixture stirred at 0 °C for 30 minutes. The cooled (0 °C) reaction mixture was treated with an excess of solid carbon dioxide and slowly allowed to warm to room temperature. To the reaction mixture, a saturated aqueous solution of ammonium chloride was added followed by a saturated aqueous solution of sodium hydrogen carbonate and this mixture was washed with ethyl acetate. The aqueous layer was treated with 2 M hydrochloric acid until pH 1 had been attained then extracted into dichloromethane. The combined organic layers were washed with saturated saline, dried and concentrated to give the title compounds in a 9: 1 ratio having the following physical properties (0.73 g).

Example 662(1):

LC/MS  $t_R$  2.05 minutes; MS ( $ES^+$ )  $m/z$  356 ( $M+Na$ ) 201 [ $M-boc-tBu$ ] $^{+a}$ .

Example 662(2):

LC/MS  $t_R$  1.74 minutes; MS ( $ES^+$ )  $m/z$  279 ( $M+Na$ ) 201 [ $M-tBu$ ] $^{+a}$ .

Example 663(1) and 663(2): tert-butyl N-[(tert-butoxy)carbonyl]-N-(5-{N'-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboximidoyl]hydrazinecarbonyl}-6-fluoropyridin-2-yl)carbamate and tert-butyl N-(5-{N'-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboximidoyl]hydrazinecarbonyl}-6-fluoropyridin-2-yl)carbamate

To a cooled (0 °C) dichloromethane (2 mL) suspension of the 9: 1 mixture of compounds prepared in Example 662 (52 mg) was added oxalyl chloride (13.5  $\mu$ L) followed by 1 drop of N,N-dimethylformamide and the mixture stirred at 0 °C for 1 hour. The reaction mixture was concentrated, the residue dissolved in tetrahydrofuran (2

mL) and added to a cooled (0 °C) tetrahydrofuran (2 mL) solution of the compound prepared in Example 659 (70 mg). The mixture was stirred 18 hours at room temperature then concentrated and the residue suspended in a saturated aqueous solution of sodium hydrogen carbonate, followed by extraction into ethyl acetate. The combined organic layers were washed with water and saturated saline, dried and concentrated to obtain the title compounds in a 3: 2 ratio having the following physical properties (120 mg).

Example 663(1):

10 LC/MS  $t_R$  1.85 minutes; MS ( $ES^+$ )  $m/z$  709 (M+H) 609 [M-boc]<sup>+</sup><sup>a</sup>.

Example 663(2):

LC/MS  $t_R$  1.59 minutes; MS ( $ES^+$ )  $m/z$  609 (M+H)<sup>a</sup>.

15 Example 664: (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-4H-1,2,4-triazol-3-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The 3: 2 mixture of compounds prepared in Example 663 (160 mg) was treated as detailed in Example 243 to give the title compound having the following physical properties (25 mg).

20 LC/MS  $t_R$  3.35 minutes; MS ( $ES^+$ )  $m/z$  491 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 8.07 (t, 1 H), 7.77 - 7.72 (m, 2 H), 7.71 - 7.68 (m, 1 H), 6.50 (dd, 1 H), 6.16 (s, 1 H), 6.08 (s, 1 H), 5.84 (d, 1 H), 3.37 - 3.34 (m, 1 H), 3.08 (dd, 1 H), 2.73 - 2.59 (m, 1 H), 2.36 - 2.22 (m, 1 H).

25 Example 665: ethyl 2-benzoyl-4-[7-(2-[[tert-butoxy]carbonyl]amino)-5-chlorophenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-oxobutanoate

To a cooled (0 °C) tetrahydrofuran (5 mL) suspension of sodium hydride (11 mg, 60% dispersion in mineral oil) and sodium iodide (34 mg), ethyl benzoylacetate (48  $\mu$ L) was added and the mixture was stirred at 0 °C for 30 minutes. A tetrahydrofuran (1.2 mL) solution of the compound prepared in Example 617 (100 mg) was added and the mixture was stirred at room temperature for 16 hours. To the reaction mixture, 0.5 M hydrochloric acid in water (30 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated.

The residue was purified by column chromatography (20 to 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (112 mg). LC/MS  $t_R$  2.33 minutes; MS ( $ES^+$ )  $m/z$  593 ( $M+H$ )<sup>a</sup>.

5 Example 666: 1-[7-(2-amino-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-4-phenylbutane-1,4-dione

To a 1,4-dioxane (3 mL) solution of the compound prepared in Example 665 was added 6 M hydrochloric acid in water (3 mL) and the mixture stirred at 100 °C for 3 hours. The mixture was then diluted with water, basified to pH 8 by addition of  
10 sodium hydrogen carbonate and extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (20% to 100% ethyl acetate in heptanes, then 0 to 15% methanol in ethyl acetate) to give the title compound having the following physical properties (29.4 mg).

15 LC/MS  $t_R$  2.00 minutes; MS ( $ES^+$ )  $m/z$  421 ( $M+H$ )<sup>a</sup>.

Example 667: 7-(2-amino-5-chlorophenyl)-3-(5-phenyl-1H-pyrrol-2-yl)-1,2,3,5-tetrahydroindolizin-5-one

To an acetic acid (1 mL) solution of the compound prepared in Example 666 (39  
20 mg) was added ammonium acetate (100 mg) and the mixture was stirred at 90 °C for 1 hour. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (30 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated to give the title compound having the following physical properties (50 mg).

25 LC/MS  $t_R$  2.26 minutes; MS ( $ES^+$ )  $m/z$  402 ( $M+H$ )<sup>a</sup>.

Example 668: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-(5-phenyl-1H-pyrrol-2-yl)-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 667 (37 mg) was treated as detailed in  
30 Example 24 to give the title compound having the following physical properties (20 mg). LC/MS  $t_R$  4.51 minutes; MS ( $ES^+$ )  $m/z$  455 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.47 (br. s, 1 H), 8.52 (s, 1 H), 7.61 (dd, 1 H), 7.54 (d, 1 H), 7.52 - 7.48 (m, 3 H), 7.35 (t, 2 H), 7.23 - 7.17 (m, 1 H), 6.41 (s, 1 H), 6.40 (t, 1 H),

6.18 (t, 1 H), 5.96 (d, 1 H), 5.58 (s, 1 H), 3.28 (ddd, 1 H), 2.98 (dd, 1 H), 2.69 (dd, 1 H), 2.49 (tt, 1 H).

5 Example 669: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(4-nitrophenyl)-1H-pyrrol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 665 → Example 666 → Example 667 → Example 24 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 665 in the operation, ethyl 4-nitrobenzoylacetate was used).

- 10 LC/MS  $t_R$  4.51 minutes; MS (ES<sup>+</sup>)  $m/z$  500 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.69 (br. s, 1 H), 9.71 (s, 1 H), 8.22 (d, 2 H), 7.86 - 7.77 (m, 5 H), 6.74 (t, 1 H), 6.01 (s, 1 H), 5.95 (s, 1 H), 5.76 - 5.69 (m, 2 H), 3.17 - 3.06 (m, 1 H), 2.96 (dd, 1 H), 2.57 - 2.51 (obs. m, 1 H), 2.18 (dd, 1 H).

15 Example 670: (3S)-3-[5-(4-aminophenyl)-1H-pyrrol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 669 (608 mg) was treated as detailed in Example 74 to give the title compound having the following physical properties (161 mg).

- 20 LC/MS  $t_R$  3.40 minutes; MS (ES<sup>+</sup>)  $m/z$  470 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.03 (br. s, 1 H), 9.70 (s, 1 H), 8.05 (br. s, 2 H), 7.83 - 7.77 (m, 3 H), 7.45 (d, 2 H), 6.95 (d, 2 H), 6.25 (t, 1 H), 6.01 (s, 1 H), 5.93 (s, 1 H), 5.71 (d, 1 H), 5.66 (t, 1 H), 3.17 - 3.07 (m, 1 H), 2.95 (dd, 1 H), 2.47 - 2.42 (obs. m, 1 H), 2.22 (dd, 1 H).

25

Example 671: methyl [4-(5-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-pyrrol-2-yl)phenyl]carbamate

The compound prepared in Example 670 (65 mg) was treated with methyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (45.2 mg).

- 30 LC/MS  $t_R$  4.19 minutes; MS (ES<sup>+</sup>)  $m/z$  528 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.40 (br. s, 1 H), 8.53 (s, 1 H), 7.60 (dd, 1 H), 7.54 (d, 1 H), 7.50 (d, 1 H), 7.44 (d, 2 H), 7.38 (d, 2 H), 6.77 (br. s, 1 H), 6.41 (s, 1 H), 6.33 (t, 1

H), 6.15 (t, 1 H), 5.95 (d, 1 H), 5.59 (s, 1 H), 3.28 (ddd, 1 H), 2.97 (dd, 1 H), 2.68 (dd, 1 H), 2.48 (tt, 1 H).

Example 672: 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-pyrrol-2-yl)phenyl]carbamate

The compound prepared in Example 670 (65 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (32.2 mg).

LC/MS  $t_R$  4.20 minutes; MS (ES<sup>+</sup>)  $m/z$  572 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.40 (br. s, 1 H), 8.53 (s, 1 H), 7.61 (dd, 1 H), 7.54 (d, 1 H), 7.50 (d, 1 H), 7.44 (d, 2 H), 7.37 (d, 2 H), 6.80 (br. s, 1 H), 6.42 (s, 1 H), 6.33 (t, 1 H), 6.16 (app. br. s, 1 H), 5.95 (d, 1 H), 5.58 (s, 1 H), 4.37 - 4.31 (m, 2 H), 3.69 - 3.62 (m, 2 H), 3.42 (s, 3 H), 3.33 - 3.22 (m, 1 H), 2.97 (dd, 1 H), 2.68 (dd, 1 H), 2.48 (tt, 1 H).

Example 673: methyl 6-amino-2-fluoropyridine-3-carboxylate

To a methanol (20 mL) solution of 2-amino-6-fluoro-5-iodopyridine (1.0 g), triethylamine (0.76 mL), palladium(II) dichloride (12 mg) and 1,1'-binaphthalene-2,2'-diylbis(diphenylphosphane) (45 mg) were added and the mixture heated at 100 °C under an atmosphere of carbon monoxide (3.8 bar) for 20 hours. The reaction mixture was concentrated and the residue purified by column chromatography (10% to 80% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (615 mg).

LC/MS  $t_R$  1.24 minutes; MS (ES<sup>+</sup>)  $m/z$  171 (M+H)<sup>a</sup>.

Example 674: methyl 6-{bis[(prop-2-en-1-yloxy)carbonyl]amino}-2-fluoropyridine-3-carboxylate

To an acetonitrile (3.2 mL) solution of the compound prepared in Example 673 (276 mg), 4-dimethylaminopyridine (20 mg), N,N-diisopropylethylamine (2.0 mL) and allyl chloroformate (2.6 mL) were added and the mixture stirred at room temperature for 5 days. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (40 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue

was purified by column chromatography (0% to 60% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (134 mg).

LC/MS  $t_R$  1.98 minutes; MS ( $ES^+$ )  $m/z$  339 ( $M+H$ )<sup>a</sup>.

5 Example 675(1) and 675(2): 2-fluoro-6-{[(prop-2-en-1-yloxy)carbonyl]amino}pyridine-3-carboxylic acid and 2-fluoro-6-[(methoxycarbonyl)amino]pyridine-3-carboxylic acid

To a methanol (3.5 mL) and tetrahydrofuran (3.5 mL) solution of the compound prepared in Example 674 was added 1 M sodium hydroxide (3.4 mL) and the mixture stirred at room temperature for 16 hours. The reaction mixture was concentrated and the  
10 residue diluted with water (20 mL) and washed with ethyl acetate. The aqueous layer was acidified with 1 M hydrochloric acid to attain pH 3 – 4 then extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated to obtain the title products in a 2: 1 ratio.

15 Example 675(1)

LC/MS  $t_R$  1.53 minutes; MS ( $ES^+$ )  $m/z$  241 ( $M+H$ )<sup>a</sup>.

Example 675(2)

LC/MS  $t_R$  1.23 minutes; MS ( $ES^+$ )  $m/z$  215 ( $M+H$ )<sup>a</sup>.

20

Example 676: ethyl 3-(2-fluoro-6-{[(prop-2-en-1-yloxy)carbonyl]amino}pyridin-3-yl)-3-oxopropanoate

To a cooled (0 °C) dichloromethane (12 mL) solution of the 2: 1 mixture of compounds prepared in Example 675 (287 mg) was added oxalyl chloride (145  $\mu$ L) and  
25 N,N-dimethylformamide (5  $\mu$ L) and the mixture stirred at room temperature for 2 hours. The reaction mixture was then concentrated and the residue dissolved in acetonitrile (5 mL).

Separately, to an acetonitrile (5 mL) suspension of ethyl potassium malonate (488 mg) was added sequentially triethylamine (620  $\mu$ L) and magnesium chloride (310 mg) and  
30 the mixture stirred at room temperature for 4 hours. To this reaction mixture, the above prepared acid chloride solution and triethylamine (167  $\mu$ L) were added sequentially and the mixture stirred at room temperature for 2 days. To the reaction mixture, 0.5 M hydrochloric acid (30 mL) was added and the mixture stirred at room temperature for 10

minutes. The mixture was then extracted into ethyl acetate and the combined organic layers washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0% to 50% ethyl acetate in heptanes) to obtain the title compound having the following physical properties (208 mg).

- 5 LC/MS  $t_R$  1.90 minutes; MS (ES<sup>+</sup>)  $m/z$  311 (M+H)<sup>a</sup>.

Example 677: ethyl 4-{7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl}-2-(2-fluoro-6-[[prop-2-en-1-yloxy]carbonyl]amino}pyridine-3-carbonyl)-4-oxobutanoate

- 10 The same operation as in Example 665 → Example 40 → Example 24 was conducted from the compound prepared in Example 623 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 665 in the operation, the compound prepared in Example 676 was used).  
LC/MS  $t_R$  2.09 minutes; MS (ES<sup>+</sup>)  $m/z$  664 (M+H)<sup>a</sup>.


15

Example 678: prop-2-en-1-yl N-[5-(4-{7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl}-4-oxobutanoyl)-6-fluoropyridin-2-yl]carbamate

- To a 1,4-dioxane (2 mL) solution of the compound prepared in Example 677 (164 mg) was added 2 M hydrochloric acid (1 mL) and the mixture stirred at 100 °C for  
20 5.5 hours. To the reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate (30 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (50 to 100% ethyl acetate in heptanes, then 0 to 10% methanol in ethyl acetate) to give the title compound having the following  
25 physical properties (37.2 mg).  
LC/MS  $t_R$  2.00 minutes; MS (ES<sup>+</sup>)  $m/z$  592 (M+H)<sup>a</sup>.

Example 679: (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-pyrrol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone





The same operation as in Example 667 → Example 584 was conducted from the compound prepared in Example 678 to give the title compound having the following physical properties.

- 5 LC/MS  $t_R$  3.97 minutes; MS (ES<sup>+</sup>)  $m/z$  489 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  10.78 (br. s, 1 H), 9.70 (s, 1 H), 7.82 - 7.78 (m, 3 H), 7.74 (dd, 1 H), 6.40 - 6.31 (m, 3 H), 6.17 (app. br. s, 1 H), 6.05 (s, 1 H), 5.93 (s, 1 H), 5.81 (t, 1 H), 5.74 (d, 1 H), 3.17 - 3.06 (m, 1 H), 2.96 (dd, 1 H), 2.46 - 2.40 (m, 1 H), 2.30 (dd, 1 H).

10

Example 680(1) and 680(2): ethyl 3-[7-(2-{bis[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-3-oxopropanoate and ethyl 3-[7-(2-{[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-3-oxopropanoate

- 15 The compound prepared in Example 622 (3.0 g) was treated as detailed in Example 676 to give the title products in a 6: 5 ratio having the following physical properties.

Example 680(1):

- 20 LC/MS  $t_R$  2.26 minutes; MS (ES<sup>+</sup>)  $m/z$  597 (M+Na)<sup>a</sup>.

Example 680(2):

LC/MS  $t_R$  2.05 minutes; MS (ES<sup>+</sup>)  $m/z$  475 (M+H)<sup>a</sup>.

- 25 Example 681(1) and 681(2): tert-butyl N-[(tert-butoxy)carbonyl]-N-{4-chloro-2-[5-oxo-3-(5-oxo-2,5-dihydro-1H-pyrazol-3-yl)-1,2,3,5-tetrahydroindolizin-7-yl]phenyl} carbamate and tert-butyl N-{4-chloro-2-[5-oxo-3-(5-oxo-2,5-dihydro-1H-pyrazol-3-yl)-1,2,3,5-tetrahydroindolizin-7-yl]phenyl} carbamate

To an ethanol (12 mL) solution of the 6: 5 mixture of compounds prepared in Example 680 (2.15 g) was added a 35% weight solution of hydrazine hydrate in water (0.65 mL) and the mixture stirred at 50 °C for 3 hours. The reaction mixture was concentrated and the residue purified by column chromatography (0% to 10% methanol in dichloromethane) to give the title products in a 1 : 2 ratio.

Example 681(1):

LC/MS  $t_R$  1.97 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H), 443 (M-CO<sub>2</sub>C(CH<sub>3</sub>)<sub>3</sub>+2H) <sup>a</sup>.

10 Example 681(2):

LC/MS  $t_R$  1.76 minutes; MS (ES<sup>+</sup>)  $m/z$  443 (M+H) <sup>a</sup>.

Example 682(1) and 682(2): tert-butyl N-[(tert-butoxy)carbonyl]-N-{4-chloro-2-[(3S)-3-[1-(4-nitrophenyl)-5-oxo-2,5-dihydro-1H-pyrazol-3-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl]phenyl} carbamate and tert-butyl N-{4-chloro-2-[(3S)-3-[1-(4-nitrophenyl)-5-oxo-2,5-dihydro-1H-pyrazol-3-yl]-5-oxo-1,2,3,5-tetrahydroindolizin-7-yl]phenyl} carbamate

To a dimethylsulfoxide (12 mL) solution of the 1: 2 mixture of compounds prepared in Example 681 (1.03 g) was added 1-fluoro-4-nitrobenzene (0.32 g) and potassium carbonate (0.48 g) and the mixture stirred at 50 °C for 3 hours. To the reaction mixture, water (40 mL) was added followed by extraction into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by flash chromatography (30% to 100% ethyl acetate in heptanes) to give the title products in a 1 : 2 ratio.

25 Example 682(1):

LC/MS  $t_R$  2.39 minutes; MS (ES<sup>+</sup>)  $m/z$  686 (M+Na) <sup>a</sup>.

Example 682(2):

30 LC/MS  $t_R$  2.20 minutes; MS (ES<sup>+</sup>)  $m/z$  564 (M+H) <sup>a</sup>.

Example 683: (3S)-3-[1-(4-aminophenyl)-5-oxo-2,5-dihydro-1H-pyrazol-3-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 40 → Example 24 → Example 74 was conducted from the 1: 2 mixture of compounds prepared in Example 682 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.90 minutes; MS ( $ES^+$ )  $m/z$  487 (M+H) <sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.17 (br. s, 1 H), 9.68 (s, 1 H), 7.86 - 7.70 (m, 3 H), 6.82 (d, 2 H), 6.62 (d, 2 H), 5.97 (br. s, 1 H), 5.91 (br. s, 1 H), 5.75 (br. s, 2 H), 5.60 (d, 1 H), 5.43 (br. s, 1 H), 3.21 - 3.06 (m, 1 H), 2.94 (dd, 1 H), 2.59 - 2.42 (obs. m, 1 H), 2.21 - 2.09 (m, 1 H).

10 Example 684: methyl [4-(3-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-5-oxo-2,5-dihydro-1H-pyrazol-1-yl]phenyl]carbamate

The compound prepared in Example 683 (110 mg) was treated with methyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (19 mg).

- 15 LC/MS  $t_R$  3.71 minutes; MS ( $ES^+$ )  $m/z$  545 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.34 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.40 (d, 2 H), 7.03 - 6.98 (m, 2 H), 6.17 (s, 1 H), 6.03 (s, 1 H), 5.76 (d, 1 H), 5.59 (s, 1 H), 3.72 (s, 3 H), 3.29 - 3.20 (m, 1 H), 3.06 (dd, 1 H), 2.65 - 2.54 (m, 1 H), 2.33 (dd, 1 H).

20

Example 685: 2-methoxyethyl [4-(3-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-5-oxo-2,5-dihydro-1H-pyrazol-1-yl]phenyl]carbamate

The compound prepared in Example 683 (110 mg) was treated with 2-

- 25 methoxyethyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (39 mg).

LC/MS  $t_R$  3.79 minutes; MS ( $ES^+$ )  $m/z$  589 (M+H) <sup>b</sup>

- <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.34 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.70 - 7.65 (m, 1 H), 7.40 (d, 2 H), 7.03 - 6.97 (m, 2 H), 6.16 (s, 1 H), 6.03 (s, 1 H), 5.76 (d, 1 H), 5.60 (s, 30 1 H), 4.28 - 4.22 (m, 2 H), 3.67 - 3.60 (m, 2 H), 3.38 (s, 3 H), 3.29 - 3.20 (m, 1 H), 3.05 (dd, 1 H), 2.65 - 2.54 (m, 1 H), 2.32 (dd, 1 H).

Example 686: 2-(2-methoxyethoxy)ethyl [4-(3-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l}-5-oxo-2,5-dihydro-1H-pyrazol-1-yl)phenyl]carbamate

The compound prepared in Example 683 (100 mg) was treated with the  
5 chloroformate as prepared in Example 489 following the method of Example 484 to give the title compound having the following physical properties (51.6 mg).

LC/MS  $t_R$  3.78 minutes; MS ( $ES^+$ )  $m/z$  633 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.34 (s, 1 H), 7.76 - 7.70 (m, 2 H), 7.69 - 7.64 (m, 1 H), 7.39 (d, 2 H), 7.03 - 6.97 (m, 2 H), 6.16 (s, 1 H), 6.02 (s, 1 H), 5.76 (d, 1 H), 5.59 (s,  
10 1 H), 4.27 - 4.22 (m, 2 H), 3.74 - 3.68 (m, 2 H), 3.66 - 3.61 (m, 2 H), 3.57 - 3.51 (m, 2 H), 3.35 (s, 3 H), 3.29 - 3.18 (m, 1 H), 3.05 (dd, 1 H), 2.65 - 2.53 (m, 1 H), 2.32 (dd, 1 H).


Example 687: tert-butyl 2-[2-oxo-4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2-dihydropyridin-1-yl]acetate

The compound prepared in Example 22 (0.50 g) was treated as detailed in Example 282 to give the crude title compound having the following physical properties (0.50 g). (Note: the reaction time was limited to 40 minutes).

LC/MS  $t_R$  1.35 minutes; MS ( $ES^+$ )  $m/z$  254 (M-C<sub>6</sub>H<sub>10</sub>+H), 198 (M-C<sub>6</sub>H<sub>10</sub>-C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>a</sup>.

20

Example 688: tert-butyl 2-[4-(2-amino-5-methylphenyl)-2-oxo-1,2-dihydropyridin-1-yl]acetate



To a 1,4-dioxane (4.5 mL) solution of the compound prepared in Example 687  
25 (150 mg) was added 2-bromo-4-methylaniline (83 mg) and caesium fluoride (154 mg) under an atmosphere of nitrogen. The mixture was degassed with nitrogen, tetrakis(triphenylphosphine)palladium(0) (23.5 mg) added and the mixture stirred at 90 °C for 1.5 hours. On cooling to room temperature, ethyl acetate (18 mL) was added, the resultant suspension filtered through Celite® and the filtrate concentrated. The

residue was purified by column chromatography (0% to 10% methanol in ethyl acetate) to give the title compound having the following physical properties (100 mg).

LC/MS  $t_R$  1.78 minutes; MS ( $ES^+$ )  $m/z$  315 ( $M+H$ ), 259 ( $M-C(CH_3)_3+H$ )<sup>a</sup>.

5 Example 689: methyl {4-[2-({4-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 24 → Example 25 → Example 51 → Example 52 was conducted from the compound prepared in Example 688 to give the title compound having the following physical properties. (Note: in the step

10 corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).

LC/MS  $t_R$  3.03 minutes; MS ( $ES^+$ )  $m/z$  483 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.11 (br. s, 1 H), 9.67 (s, 1 H), 9.61 (br. s, 1 H), 7.68 (d, 1 H), 7.63 – 7.60 (m, 3 H), 7.52 (d, 1 H), 7.51 (s, 1 H), 7.44 (d, 1 H), 7.41 (d, 2 H),

15 6.20 (d, 1 H), 5.83 (dd, 1 H), 5.07 (s, 2 H), 3.65 (s, 3 H), 2.45 (s, 3 H).


Example 690: methyl {4-[4-chloro-2-({4-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl]-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The compound prepared in Example 689 (35 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (9 mg).

LC/MS  $t_R$  3.86 minutes; MS ( $ES^+$ )  $m/z$  517 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.87 (br. s, 1 H), 9.80 (br. s, 1 H), 9.69 (s, 1 H), 7.67 (d, 1 H), 7.64 – 7.57 (m, 3 H), 7.57 – 7.47 (m, 4 H), 6.19 (d, 1 H), 5.85 (dd, 1 H), 5.06 (s, 2 H), 3.68 (s, 3 H), 2.46 (s, 3 H).

25 Example 691: methyl (3S)-5-oxo-7-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,5-tetrahydroindolizine-3-carboxylate



The same operation as in Example 5 → Example 6 → Example 282 was conducted from the compound prepared in Example 4 to give the title compound having

30

the following physical properties. (Note: in the step corresponding to Example 5 in the operation, methanol was used in place of ethanol).

LC/MS  $t_R$  1.05 minutes; MS ( $ES^+$ )  $m/z$  238 ( $M-C_4H_{10}+H$ )<sup>a</sup>.

5 Example 692: (3S)-7-[5-methoxy-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

The same operation as in Example 688 → Example 8 → Example 9 was conducted from the compound prepared in Example 691 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 10 688 in the operation, 2-bromo-4-methoxyaniline was used).

LC/MS  $t_R$  1.44 minutes; MS ( $ES^+$ )  $m/z$  354 ( $M+H$ )<sup>a</sup>.

Example 693: methyl [4-(2-[(3S)-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

15 The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 692 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).

20 LC/MS  $t_R$  3.03 minutes; MS ( $ES^+$ )  $m/z$  525 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.04 (br. s, 1 H), 9.63 (s, 1 H), 9.59 (br. s, 1 H), 7.63 (d, 1 H), 7.59 (d, 2 H), 7.44 - 7.35 (m, 3 H), 7.22 (dd, 1 H), 7.16 (d, 1 H), 5.94 (s, 1 H), 5.92 (s, 1 H), 5.60 (d, 1 H), 3.89 (s, 3 H), 3.65 (s, 3 H), 3.43 - 3.36 (m, 1 H), 2.99 (dd, 1 H), 2.57 - 2.42 (obs. m, 1 H), 2.39 - 2.29 (m, 1 H).

25

Example 694: 2-methoxyethyl [4-(2-[(3S)-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 692 to give the title compound having the following 30 physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 324 was used).

LC/MS  $t_R$  2.94 minutes; MS ( $ES^+$ )  $m/z$  569 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.27 (s, 1 H), 7.54 (d, 1 H), 7.54 (app. br. s, 2 H), 7.43 (app. br. s, 2 H), 7.20 (dd, 1 H), 7.14 (d, 1 H), 7.26 (br. s, 1 H), 6.10 (s, 1 H), 6.07 (s, 1 H), 5.76 (dd, 1 H), 4.29 - 4.22 (m, 2 H), 3.91 (s, 3 H), 3.67 - 3.61 (m, 2 H), 3.48 - 3.39 (m, 1 H), 3.38 (s, 3 H), 3.07 (ddd, 1 H), 2.60 (qd, 1 H), 2.46 (app. br. s, 1 H).

5

Example 695: 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 694 (83 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (59.5 mg).

10

LC/MS *t<sub>R</sub>* 3.90 minutes; MS (ES<sup>+</sup>) *m/z* 603 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.29 (s, 1 H), 7.64 - 7.58 (m, 2 H), 7.55 (d, 1 H), 7.52 (d, 2 H), 7.22 (dd, 1 H), 7.17 (d, 1 H), 6.11 (s, 1 H), 6.09 (s, 1 H), 5.70 (dd, 1 H), 4.30 - 4.25 (m, 2 H), 3.93 (s, 3 H), 3.68 - 3.62 (m, 2 H), 3.48 - 3.41 (m, 1 H), 3.39 (s, 3 H), 3.09 (ddd, 1 H), 2.62 (qd, 1 H), 2.42 - 2.34 (m, 1 H).

15

Example 696: (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-methoxy-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 484 → Example 338 → Example 584 was conducted from the compound prepared in Example 692 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used. In the step corresponding to Example 484 in the operation, allyl chloroformate was used).

20

LC/MS *t<sub>R</sub>* 2.82 minutes; MS (ES<sup>+</sup>) *m/z* 502 (M+H)<sup>b</sup>

25

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.29 (s, 1 H), 8.21 (s, 1 H), 7.75 (dd, 1 H), 7.56 (d, 1 H), 7.22 (dd, 1 H), 7.17 (d, 1 H), 6.65 (d, 1 H), 6.11 (s, 1 H), 6.09 (s, 1 H), 5.68 (dd, 1 H), 3.93 (s, 3 H), 3.50 - 3.36 (m, 1 H), 3.09 (ddd, 1 H), 2.62 (qd, 1 H), 2.43 - 2.30 (m, 1 H).

30

Example 697: methyl [4-(2-((3S)-7-[6-methyl-3-(1H-tetrazol-1-yl)-2-pyridinyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 688 → Example 8 → Example 24 → Example 51 → Example 52 was conducted from the compound prepared in Example 691 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 688 in the operation, 2-bromo-6-methylpyridin-3-amine was used. In the step corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).  
LC/MS  $t_R$  2.90 minutes; MS ( $ES^+$ )  $m/z$  510 (M+H), 482 (M-N<sub>2</sub>+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.05 (br. s, 1 H), 9.74 (s, 1 H), 9.58 (br. s, 1 H), 8.12 (d, 1 H), 7.63 (d, 1 H), 7.59 (d, 2 H), 7.40 (d, 2 H), 7.38 (s, 1 H), 6.19 (s, 1 H), 5.78 (s, 1 H), 5.61 (d, 1 H), 3.71 – 3.62 (obs. m, 1 H), 3.65 (s, 3 H), 3.46 – 3.36 (obs. m, 1 H), 3.07 (dd, 1 H), 2.65 (s, 3 H), 2.34 – 2.27 (m, 1 H).

Example 698: (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-fluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 7 → Example 9 → Example 596 → Example 51 → Example 52 was conducted from the compound prepared in Example 6 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 7 in the operation, 4-fluoro-2-(tetramethyl-1,3,2-dioxaborolan-2-yl)aniline [patent EP1719773 (2006)] was used. In the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used).

LC/MS  $t_R$  2.50 minutes; MS ( $ES^+$ )  $m/z$  456 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.35 (s, 1 H), 8.19 (d, 1 H), 7.77 (dd, 1 H), 7.74 - 7.69 (m, 1 H), 7.51 - 7.44 (m, 2 H), 7.20 (s, 1 H), 6.63 (d, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.50 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.62 (qd, 1 H), 2.50 - 2.41 (m, 1 H).

Example 699: (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-fluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 698 (103 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (51 mg).

LC/MS  $t_R$  2.78 minutes; MS ( $ES^+$ )  $m/z$  490 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 8.21 (app. s, 1 H), 7.83 - 7.64 (m, 2 H), 7.58 - 7.39 (m, 2 H), 6.65 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.71 (dd, 1 H), 3.52 - 3.38 (m, 1 H), 3.11 (td, 1 H), 2.74 - 2.54 (m, 1 H), 2.45 - 2.30 (m, 1 H).

5 Example 700: (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[2,3-difluoro-6-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 688 → Example 9 → Example 596 → Example 51 → Example 52 was conducted from the compound prepared in Example 282 to give the title compound having the following physical properties. (Note: in the  
10 step corresponding to Example 688 in the operation, 2-bromo-3,4-difluoroaniline was used. In the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used)

LC/MS *t<sub>R</sub>* 2.57 minutes; MS (ES<sup>+</sup>) *m/z* 474 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 8.20 (app. br. s, 1 H), 7.75 (d, 1 H),  
15 7.65 (q, 1 H), 7.60 - 7.54 (m, 1 H), 7.19 (br. s, 1 H), 6.61 (d, 1 H), 6.21 (s, 1 H), 6.18 (s, 1 H), 5.77 (dd, 1 H), 3.49 (td, 1 H), 3.14 (ddd, 1 H), 2.64 (qd, 1 H), 2.52 - 2.42 (m, 1 H).

Example 701: (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[2,3-difluoro-6-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

20 The compound prepared in Example 700 (75 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (28.9 mg).

LC/MS *t<sub>R</sub>* 2.85 minutes; MS (ES<sup>+</sup>) *m/z* 508 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.37 (s, 1 H), 8.22 (d, 1 H), 7.77 (dd, 1 H), 7.67 (q,  
25 1 H), 7.61 - 7.55 (m, 1 H), 6.66 (d, 1 H), 6.23 (s, 1 H), 6.20 (s, 1 H), 5.71 (dd, 1 H), 3.51 (td, 1 H), 3.15 (ddd, 1 H), 2.66 (qd, 1 H), 2.41 (tdd, 1 H).

Example 702: formic acid - (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[3,5-difluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone (2:1)

30 The same operation as in Example 688 → Example 9 → Example 596 → Example 51 → Example 52 was conducted from the compound prepared in Example 282 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile

phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties. (Note: in the step corresponding to Example 688 in the operation, 2-bromo-4,6-difluoroaniline was used. In the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used).

- 5 LC/MS  $t_R$  2.60 minutes; MS (ES<sup>+</sup>)  $m/z$  474 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.47 (s, 1 H), 8.29 (br. s, 2 H), 8.15 (d, 1 H), 7.96 (dd, 1 H), 7.56 - 7.47 (m, 1 H), 7.41 - 7.35 (m, 1 H), 7.31 (s, 1 H), 6.80 (d, 1 H), 6.14 (s, 1 H), 6.13 (s, 1 H), 5.75 (dd, 1 H), 3.51 - 3.38 (m, 1 H), 3.11 (ddd, 1 H), 2.70 - 2.52 (m, 1 H), 2.48 - 2.38 (m, 1 H).

10

Example 703: formic acid - (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[3,5-difluoro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone (1:1)

The compound prepared in Example 702 (40 mg) was treated as detailed in Example 338 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (8.7 mg).

- 15 LC/MS  $t_R$  2.87 minutes; MS (ES<sup>+</sup>)  $m/z$  508 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.47 (s, 1 H), 8.30 (br. s, 1 H), 8.20 (d, 1 H), 7.78 (dd, 1 H), 7.57 - 7.48 (m, 1 H), 7.39 (dd, 1 H), 6.68 (d, 1 H), 6.14 (s, 1 H), 6.13 (s, 1 H), 5.67 (dd, 1 H), 3.44 (td, 1 H), 3.10 (ddd, 1 H), 2.69 - 2.55 (m, 1 H), 2.37 (tdd, 1 H).

20

Example 704: (3S)-7-[5-(difluoromethoxy)-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

- 25 The same operation as in Example 688 → Example 9 → Example 596 was conducted from the compound prepared in Example 282 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 688 in the operation, 2-bromo-4-(difluoromethoxy)aniline [patent WO2009/144632] was used).

- 30 LC/MS  $t_R$  1.51 minutes; MS (ES<sup>+</sup>)  $m/z$  390 (M+H), 362 (M-N<sub>2</sub>+H)<sup>a</sup>.

Example 705: (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-(difluoromethoxy)-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 704 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used)

- 5 LC/MS  $t_R$  2.71 minutes; MS (ES<sup>+</sup>)  $m/z$  504 (M+H), 252.5 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 8.19 (d, 1 H), 7.75 (dd, 1 H), 7.72 (d, 1 H), 7.47 (dd, 1 H), 7.43 (d, 1 H), 7.19 (s, 1 H), 7.08 (t, 1 H), 6.62 (d, 1 H), 6.12 (s, 1 H), 6.10 (s, 1 H), 5.76 (dd, 1 H), 3.44 (td, 1 H), 3.10 (ddd, 1 H), 2.63 (qd, 1 H), 2.49 - 2.41 (m, 1 H).

10

Example 706: (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-(difluoromethoxy)-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 705 (90 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (50 mg).

15

- LC/MS  $t_R$  2.95 minutes; MS (ES<sup>+</sup>)  $m/z$  538 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.38 (s, 1 H), 8.22 (d, 1 H), 7.77 (dd, 1 H), 7.74 (d, 1 H), 7.51 - 7.48 (m, 1 H), 7.46 (d, 1 H), 7.11 (t, 1 H), 6.67 (d, 1 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.71 (dd, 1 H), 3.44 (td, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.44 - 2.34 (m, 1 H).

20

Example 707: (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-(difluoromethoxy)-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 704 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 215 was used).

25

- LC/MS  $t_R$  2.94 minutes; MS (ES<sup>+</sup>)  $m/z$  522 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.98 (app. br. s, 1 H), 7.74 (d, 1 H), 7.49 (dd, 1 H), 7.45 (d, 1 H), 7.16 (br. s, 1 H), 7.09 (t, 1 H), 6.47 (dd, 1 H), 6.16 (s, 1 H), 6.10 (s, 1 H), 5.80 (dd, 1 H), 3.45 (td, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.52 - 2.45 (m, 1 H).

30

Example 708: (3S)-7-[5-cyano-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid

- The same operation as in Example 688 → Example 9 → Example 596 was conducted from the compound prepared in Example 282 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 688 in the operation, 2-bromo-4-cyanoaniline was used).
- LC/MS  $t_R$  1.51 minutes; MS ( $ES^+$ )  $m/z$  719 (2M+Na), 697 (2M+H)<sup>a</sup>.

Example 709: 3-{(3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyll}-4-(1H-tetrazol-1-yl)benzonitrile

- The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 708 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used).
- LC/MS  $t_R$  2.42 minutes; MS ( $ES^+$ )  $m/z$  463 (M+H), 435 (M-N<sub>2</sub>+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.41 (s, 1 H), 8.20 (app. s, 1 H), 8.13 – 8.05 (m, 2 H), 7.90 (d, 1 H), 7.75 (dd, 1 H), 7.20 (br. s, 1 H), 6.62 (d, 1 H), 6.17 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.45 (td, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.52 - 2.40 (m, 1 H).

Example 710: 3-{(3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyll}-4-(1H-tetrazol-1-yl)benzonitrile

- The compound prepared in Example 709 (70 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (22.9 mg).
- LC/MS  $t_R$  2.73 minutes; MS ( $ES^+$ )  $m/z$  497 (M+H), 469 (M-N<sub>2</sub>+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.44 (s, 1 H), 8.23 (d, 1 H), 8.15 - 8.07 (m, 2 H), 7.93 (d, 1 H), 7.79 (dd, 1 H), 6.68 (d, 1 H), 6.19 (s, 1 H), 6.12 (s, 1 H), 5.72 (dd, 1 H), 3.47 (td, 1 H), 3.13 (ddd, 1 H), 2.67 (qd, 1 H), 2.45 - 2.37 (m, 1 H).

Example 711: 3-{(3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyll}-4-(1H-tetrazol-1-yl)benzonitrile


The same operation as in Example 51 → Example 52 was conducted from the compound prepared in Example 708 to give the title compound having the following

physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 215 was used).

LC/MS  $t_R$  2.66 minutes; MS ( $ES^+$ )  $m/z$  481 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.42 (s, 1 H), 8.12 - 8.08 (m, 2 H), 7.98 (br. s, 1 H),  
5 7.92 (d, 1 H), 7.17 (d, 1 H), 6.47 (dd, 1 H), 6.20 (s, 1 H), 6.11 (s, 1 H), 5.82 (dd, 1 H),  
3.46 (td, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.54 - 2.47 (m, 1 H).

Example 712: methyl N-[6-(2-bromoacetyl)pyridin-3-yl]carbamate




10 The same operation as in Example 10 → Example 188 → Example 55 →  
Example 128 → Example 204 was conducted from 5-amino-2-cyanopyridine to give the  
title compound having the following physical properties. (Note: in the step  
corresponding to Example 128 in the operation, methyl chloroformate was used).

LC/MS  $t_R$  1.68 minutes; MS ( $ES^+$ )  $m/z$  273 and 275 ( $M+H$ )<sup>a</sup>.

15


Example 713: 2-methoxyethyl N-[6-(2-bromoacetyl)pyridin-3-yl]carbamate



The same operation as in Example 128 → Example 188 → Example 204 was  
conducted from 5-amino-2-cyanopyridine to give the title compound having the  
20 following physical properties. (Note: in the step corresponding to Example 128 in the  
operation, 2-methoxyethyl chloroformate was used)

LC/MS  $t_R$  1.69 minutes; MS ( $ES^+$ )  $m/z$  317 and 319 ( $M+H$ )<sup>a</sup>.

Example 714: 2-ethoxyethyl N-[4-(2-bromoacetyl)phenyl]carbamate




25

The same operation as in Example 128 → Example 78 was conducted from 4-aminoacetophenone to give the title compound having the following physical properties. (Note: in the step corresponding to Example 128 in the operation, the chloroformate prepared in Example 485 was used)

- 5 LC/MS  $t_R$  1.85 minutes; MS ( $ES^+$ )  $m/z$  330 and 332 ( $M+H$ )<sup>a</sup>.

Example 715: 3-methoxypropyl N-[4-(2-bromoacetyl)phenyl]carbamate



- 10 The same operation as in Example 128 → Example 78 was conducted from 4-aminoacetophenone to give the title compound having the following physical properties. (Note: in the step corresponding to Example 128 in the operation, the chloroformate prepared in Example 487 was used)

LC/MS  $t_R$  1.83 minutes; MS ( $ES^+$ )  $m/z$  330 and 332 ( $M+H$ )<sup>a</sup>.

- 15 Example 716: (6S)-6-[5-(4-aminophenyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

- The same operation as in Example 51 → Example 52 → Example 74 was conducted from the compound prepared in Example 336 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 20 51 in the operation, 2-bromo-1-(4-nitrophenyl)ethan-1-one was used). LC/MS  $t_R$  2.64 minutes; MS ( $ES^+$ )  $m/z$  472 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.45 (s, 1 H), 7.92 (d, 1 H) 7.80 - 7.66 (m, 2 H), 7.44 - 7.36 (m, 2 H), 7.10 (s, 1 H), 6.78 - 6.71 (m, 2 H), 6.35 (s, 1 H), 5.73 (dd, 1 H), 3.29 - 3.25 (m, 1 H), 2.92 - 2.88 (m, 1 H), 2.67 - 2.63 (m, 1 H), 2.38 - 2.35 (m, 1 H).

25

Example 717: 3-oxetanyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

- The compound prepared in Example 716 (100 mg) was treated as detailed in Example 494 to give the title compound having the following physical properties (13.4 30 mg).

LC/MS  $t_R$  2.96 minutes; MS ( $ES^+$ )  $m/z$  572 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.13 (s, 1 H), 9.74 (s, 1 H) 7.97 (d, 1 H) 7.89 - 7.79 (m, 2 H) 7.61 (br. s, 1 H) 7.43 (d, 2 H) 6.34 (s, 1 H) 5.61 (dd, 1 H) 5.41 (t, 1 H) 4.82 - 4.78 (m, 2 H) 4.59 - 4.51 (m, 2 H) 3.20 - 3.18 (m, 1 H) 2.81 - 2.72 (m, 1 H) 2.59 - 2.54 (m, 1 H) 2.26 - 2.18 (m, 1 H).

Example 718(1) to Example 718(13)

The compounds of the present invention having the following physical data were prepared using the corresponding alpha-bromoketones from the compound prepared in Example 336 in the process of Example 51 → Example 52.

Example 718(1): 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl])-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 324 was used.

LC/MS  $t_R$  3.06 minutes; MS ( $ES^+$ )  $m/z$  574 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.46 (s, 1 H), 7.92 (d, 1 H), 7.81 - 7.67 (m, 2 H), 7.59 (d, 2 H), 7.46 (d, 2 H), 7.27 (s, 1 H), 6.36 (s, 1 H), 5.74 (dd, 1 H), 4.32 - 4.24 (m, 2 H), 3.66 (dd, 2 H), 3.41 (s, 3 H), 3.30 - 3.24 (m, 1 H), 3.01 - 2.87 (m, 1 H), 2.77 - 2.59 (m, 1 H), 2.45 - 2.34 (m, 1 H).

Example 718(2): 2-ethoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl])-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 714 was used.

LC/MS  $t_R$  2.93 minutes; MS ( $ES^+$ )  $m/z$  588 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H-NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.15 (s, 1 H), 9.68 (s, 2 H), 7.99 (s, 1 H), 7.88 - 7.80 (m, 2 H), 7.64 - 7.40 (m, 5 H), 6.36 (s, 1 H), 5.64 - 5.59 (m, 1 H), 4.22 - 4.16 (m, 2 H), 3.63 - 3.57 (m, 2 H), 3.51 - 3.43 (m, 2 H), 3.26 - 3.12 (m, 1 H), 2.80 - 2.72 (m, 1 H), 2.60 - 2.54 (m, 1 H), 2.26 - 2.12 (m, 1 H) 1.10 (t, 3H).

Example 718(3): 3-methoxypropyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

5 In the step corresponding to Example 51 in the process, the compound prepared in Example 715 was used.

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  589 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.44 (s, 1 H), 7.90 (d, 1 H), 7.76 (dd, 1 H), 7.69 (d, 1 H), 7.59 - 7.54 (bd, 2 H), 7.46 - 7.41 (bd, 2 H), 7.28 - 7.22 (bs, 1 H), 6.34 (s, 1 H), 5.72 (dd, 10.0 Hz, 1 H), 4.20 (t, 2 H), 3.51 (t, 2 H), 3.34 (s, 3 H), 3.29 - 3.25 (m, 1 H),  
10 2.94 - 2.98 (m, 1 H), 2.70 - 2.62 (m, 1 H), 2.41 - 2.34 (m, 1 H), 1.93 (quintet, 2 H).

Example 718(4): methyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate

15 In the step corresponding to Example 51 in the process, the compound prepared in Example 78 was used.

LC/MS  $t_R$  3.00 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H)<sup>b</sup>

<sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.99 (s, 1 H), 9.77 (s, 1 H), 9.57 (s, 1 H), 8.02 - 7.94 (m, 1 H), 7.89 - 7.77 (m, 2 H), 7.57 - 7.35 (m, 4 H), 6.36 (s, 1 H), 5.59 - 5.50 (m, 1 H),  
20 3.72 (s, 3 H), 3.20 (s, 1 H), 2.76 - 2.71 (m, 1 H), 2.55 - 2.50 (m, 1 H), 2.35 - 2.15 (m, 4 H).

Example 718(5): 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate

25 In the step corresponding to Example 51 in the process, the compound prepared in Example 481 was used.

LC/MS  $t_R$  3.01 minutes; MS (ES<sup>+</sup>)  $m/z$  588 (M+H)<sup>b</sup>

<sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  11.98 (s, 1 H), 9.85 (s, 2 H), 8.02 (d, 1 H), 7.89 - 7.78 (m, 2 H), 7.56 - 7.36 (m, 4 H), 6.36 (s, 1 H), 5.61 - 5.48 (m, 1 H), 4.25 (t, 2 H),  
30 3.50 (t, 2 H), 3.25 (s, 3 H), 3.25 - 3.13 (m, 1 H), 2.81 - 2.78 (m, 1 H), 2.69 - 2.65 (m, 1 H), 2.35 - 2.16 (m, 4 H).



Example 718(6): (6S)-6-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

- 5 LC/MS  $t_R$  2.52 minutes; MS ( $ES^+$ )  $m/z$  473 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.47 (s, 1 H), 8.22 (s, 1 H), 7.93 (d, 1 H), 7.82 - 7.69 (m, 3 H), 7.24 (br. s., 1 H), 6.65 (d, 1 H), 6.36 (s, 1 H), 5.74 (dd, 1 H), 3.31 - 3.26 (m, 1 H), 2.94 (ddd, 1 H), 2.75 - 2.62 (m, 1 H), 2.44 - 2.34 (m, 1 H).

- 10 Example 718(7): (6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

In the step corresponding to Example 51 in the process, the compound prepared in Example 551 was used.

- LC/MS  $t_R$  2.59 minutes; MS ( $ES^+$ )  $m/z$  487 ( $M+H$ )<sup>b</sup>  
15 <sup>1</sup>H-NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.08 (s, 1 H), 9.67 (s, 1 H), 8.36 (s, 1 H), 7.99 (s, 1 H), 7.89 - 7.79 (m, 2 H), 7.71 - 7.57 (m, 1 H), 7.34 (s, 1 H), 6.63 - 6.29 (m, 3 H), 5.64 - 5.53 (m, 1 H), 3.21 (dd, 1 H), 2.83 - 2.73 (m, 4 H), 2.62 - 2.55 (m, 1 H), 2.32 - 2.13 (m, 1 H).

- 20 Example 718(8): (6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-(5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

In the step corresponding to Example 51 in the process, the compound prepared in Example 560 was used.

- 25 LC/MS  $t_R$  2.68 minutes; MS ( $ES^+$ )  $m/z$  531 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.34 (s, 1 H), 8.09 (d, 1 H), 7.81 - 7.73 (m, 2 H), 7.69 - 7.64 (m, 1 H), 7.62 - 7.55 (m, 1 H), 7.18 (s, 1 H), 6.64 (d, 1 H), 6.24 (s, 1 H), 5.62 (dd, 1 H), 3.54 - 3.45 (m, 2 H), 3.43 - 3.36 (m, 2 H), 3.29 (s, 3 H), 3.20 - 3.12 (m, 1 H), 2.82 (ddd, 1 H), 2.56 (dd, 1 H), 2.32 - 2.21 (m, 1 H).

30

Example 718(9): (6S)-6-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one (enantiopure)

In the step corresponding to Example 51 in the process, the compound prepared in Example 215 was used.

LC/MS  $t_R$  2.86 minutes; MS (ES<sup>+</sup>)  $m/z$  491 (M+H)<sup>b</sup>

<sup>1</sup>H-NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.22 (s, 1 H), 9.74 (s, 1 H), 8.02 – 7.95 (m, 2 H),  
5 7.86 (dd, 1 H), 7.83 (d, 1 H), 7.12 (d, 1 H), 6.42 – 6.34 (m, 2 H), 6.27 (s, 2 H), 5.62 (dd,  
1 H), 3.19 (td, 1 H), 2.81 – 2.71 (m, 1 H), 2.60 – 2.55 (m, 1 H), 2.26 – 2.16 (m, 1 H).

Example 718(10): (6S)-6-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

10 In the step corresponding to Example 51 in the process, the compound prepared in Example 209 was used.

TLC R<sub>f</sub> 0.40 (10% methanol in ethyl acetate)

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.23 (s, 1 H), 9.71 (s, 1 H), 8.02 - 7.91 (m, 2 H), 7.88  
- 7.72 (m, 2 H), 7.42 (d, 1 H), 6.55 - 6.24 (m, 4 H), 5.62 (dd, 1 H), 3.25 - 3.07 (m, 1 H),  
15 2.83 - 2.66 (m, 1 H), 2.59 – 2.52 (obs. m, 1 H), 2.31 - 2.13 (m, 1 H).

Example 718(11): methyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-3-pyridinyl]carbamate

20 In the step corresponding to Example 51 in the process, the compound prepared in Example 712 was used.

LC/MS  $t_R$  2.94 minutes; MS (ES<sup>+</sup>)  $m/z$  531 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.50 (s, 1 H), 8.55 (s, 1 H), 8.00 - 7.88 (m, 3 H),  
7.83 - 7.49 (m, 3 H), 6.40 (s, 1 H), 5.81 - 5.70 (m, 1 H), 3.78 (s, 3 H), 3.41 - 3.34 (m, 1  
25 H), 2.99 - 2.83 (m, 1 H), 2.74 - 2.66 (m, 1 H), 2.51 - 2.40 (m, 1 H).

Example 718(12): 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-3-pyridinyl]carbamate


30 In the step corresponding to Example 51 in the process, the compound prepared in Example 713 was used.

LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  575 (M+H)<sup>b</sup>

<sup>1</sup>H-NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.26 (s, 1 H), 9.87 (s, 1 H), 9.78 (s, 1 H), 8.39 (s, 1 H), 8.00 - 7.64 (m, 4 H), 7.55 - 7.30 (m, 1 H), 6.39 - 6.27 (m, 1 H), 5.74 - 5.46 (m, 1 H), 4.31 - 4.10 (m, 2 H), 3.65 - 3.53 (m, 2 H), 3.29 (s, 3 H), 3.23 - 3.04 (m, 1 H), 2.83 - 2.68 (m, 1 H), 2.62 - 2.56 (m, 1 H), 2.28 - 2.10 (m, 1 H).

5

Example 718(13): methyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl]phenyl]carbamate



LC/MS *t<sub>R</sub>* 0.68 minutes; MS (ES<sup>+</sup>) *m/z* 530 (M+H)<sup>+</sup>

10 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.5 (s, 0.3 H), 12.2 (s, 0.7 H), 9.75 - 9.65 (s, 1.3 H), 9.57 (s, 0.7 H), 7.98 - 7.92 (m, 1 H), 7.91 - 7.77 (m, 2 H), 7.65 - 7.35 (m, 4.7 H), 7.12 (s, 0.3 H), 6.33 (s, 0.7 H), 6.30 (s, 0.3 H), 5.65 - 5.54 (m, 1 H), 3.66 (s, 0.9 H), 3.64 (s, 2.1 H), 3.39 - 3.10 (m, 1 H), 2.80 - 2.68 (m, 1 H), 2.58 - 2.13 (m, 2 H).

15 Example 719(1) to 719(2)

The compounds of the present invention having the following physical data were synthesised from the compound prepared in Example 718(6) and the corresponding chloroformates using the method as detailed in Example 128.

20 Example 719(1): methyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS *t<sub>R</sub>* 1.48 minutes; MS (ES<sup>+</sup>) *m/z* 531 (M+H)<sup>+</sup>

25 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.3 (s, 1 H), 10.1 (s, 1 H), 9.71 (s, 1 H), 8.59 (d, 1 H), 8.06 - 7.75 (m, 5 H), 7.56 (d, 1 H), 6.34 (s, 1 H), 5.65 - 5.57 (m, 1 H), 3.65 (s, 3 H), 3.26 - 3.17 (m, 1 H), 2.82 - 2.71 (m, 1 H), 2.71 - 2.38 (m, 1 H), 2.32 - 2.17 (m, 1 H).

Example 719(2): 2-methoxyethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.10 minutes; MS ( $ES^+$ )  $m/z$  575 ( $M+H$ )<sup>b</sup>

5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.35 (s, 1 H), 10.16 (s, 1 H), 9.79 (s, 1 H), 8.70 (s, 1 H), 8.13 - 8.01 (m, 2 H), 7.95 - 7.81 (m, 3 H), 7.64 (s, 1 H), 6.45 - 6.35 (m, 1 H), 5.69 (d, 1 H), 4.33 - 4.23 (m, 2 H), 3.67 - 3.58 (m, 2 H), 3.34 (s, 3 H), 3.32 - 3.19 (m, 1 H), 2.90 - 2.79 (m, 1 H), 2.67 - 2.62 (m, 1 H), 2.33 - 2.22 (m, 1 H).

10 Example 720: 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The compound prepared in Example 718(12) (267 mg) was treated as detailed in Example 364 to give the title product having the following physical properties (11.4 mg).

LC/MS  $t_R$  3.79 minutes; MS ( $ES^+$ )  $m/z$  593 ( $M+H$ )<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.44 (s, 1 H), 8.62 (br. s, 1 H), 7.97 - 7.87 (m, 2 H), 7.77 (dd, 1 H), 7.70 (d, 1 H), 7.52 (d, 1 H), 6.37 (s, 1 H), 5.69 (dd, 1 H), 4.33 - 4.27 (m, 2 H), 3.69 - 3.62 (m, 2 H), 3.39 (s, 3 H), 3.25 (td, 1 H), 2.90 (ddd, 1 H), 2.65 (qd, 1 H),  
20 2.39 - 2.30 (m, 1 H).

Example 721(1) to Example 721(10)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 718(1), 718(2), 718(3), 718(7),  
25 718(8), 718(9), 718(11), 718(12), 719(1), 719(2) using the method as detailed in Example 338.

Example 721(1): 2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl]phenyl]carbamate

LC/MS  $t_R$  3.94 minutes; MS ( $ES^+$ )  $m/z$  608 ( $M+H$ )<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.92 (s, 1 H), 9.91 (s, 1 H), 9.76 (s, 1 H), 8.00 (d, 1 H), 7.86 (dd, 1 H), 7.83 (d, 1 H), 7.60 (d, 2 H), 7.56 (d, 2 H), 6.34 (s, 1 H), 5.57 (dd, 1

H), 4.24 – 4.20 (m, 2 H), 3.60 – 3.55 (m, 2 H), 3.29 (s, 3 H), 3.17 – 3.07 (m, 1 H), 2.82 – 2.72 (m, 1 H), 2.62 – 2.55 (m, 1 H), 2.22 – 2.12 (m, 1 H).

Example 721(2): 2-ethoxyethyl [4-(4-chloro-2-{{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}}-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  4.12 minutes; MS (ES<sup>+</sup>)  $m/z$  623 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.92 (s, 1 H), 9.92 (br. s, 1 H), 9.74 (s, 1 H), 8.00 (d, 1 H), 7.88 - 7.81 (m, 2 H), 7.63 - 7.54 (m, 4 H), 6.34 (s, 1 H), 5.57 (dd, 1 H), 4.23 - 4.19 (m, 2 H), 3.63 - 3.59 (m, 2 H), 3.48 (q, 2 H), 3.17 - 3.08 (m, 1 H), 2.77 (d, 1 H), 2.61 - 2.56 (m, 1 H), 2.17 (d, 1 H), 1.13 (t, 3 H).

Example 721(3): 3-methoxypropyl [4-(4-chloro-2-{{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}}-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  4.05 minutes; MS (ES<sup>+</sup>)  $m/z$  622 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.45 (s, 1 H), 7.91 (d, 1 H), 7.76 (dd, 1 H), 7.70 (d, 1 H), 7.61 (d, 2 H), 7.52 (d, 2 H), 6.35 (s, 1 H), 5.66 (dd, 1 H), 4.21 (t, 2 H), 3.52 (t, 2 H), 3.34 (s, 3 H), 3.29 - 3.25 (m, 1 H), 2.94 - 2.87 (m, 1 H), 2.70 - 2.62 (m, 1 H), 2.36 - 2.30 (m, 1 H), 1.94 (quintet, 2 H).

Example 721(4): (6S)-6-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

LC/MS  $t_R$  2.82 minutes; MS (ES<sup>+</sup>)  $m/z$  521 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.81 (s, 1 H), 9.76 (s, 1 H), 8.29 (d, 1 H), 8.00 (d, 1 H), 7.86 (dd, Hz, 1 H), 7.83 (d, 1 H), 7.65 (dd, 1 H), 6.76 (d, 1 H), 6.54 (d, 1 H), 6.34 (s, 1 H), 5.55 (dd, 1 H), 3.17 – 3.08 (m, 1 H), 2.80 (s, 3 H), 2.78 – 2.72 (m, 1 H), 2.61 – 2.54 (m, 1 H), 2.22 – 2.12 (m, 1 H).

Example 721(5): (6S)-6-(4-chloro-5-{6-[(2-methoxyethyl)amino]-3-pyridinyl}-1H-imidazol-2-yl)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

LC/MS  $t_R$  2.94 minutes; MS (ES<sup>+</sup>)  $m/z$  565 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.46 (s, 1 H), 8.27 (d, 1 H), 7.92 (d, 1 H), 7.80 - 7.68 (m, 3 H), 6.65 (d, 1 H), 6.37 (s, 1 H), 5.67 (dd, 1 H), 3.65 - 3.57 (m, 2 H), 3.55 - 3.50 (m, 2 H), 3.40 (s, 3 H), 3.30 - 3.25 (m, 1 H), 2.97 - 2.86 (m, 1 H), 2.68 (dd, 1 H), 2.39 - 2.30 (m, 1 H).

5

Example 721(6): (6S)-6-[5-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one  
LC/MS *t<sub>R</sub>* 3.63 minutes; MS (ES<sup>+</sup>) *m/z* 525 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.71 (s, 1 H), 9.75 (s, 1 H), 8.00 (d, 1 H), 7.86 (dd, 1 H), 7.82 (d, 1 H), 7.61 (dd, 1 H), 6.66 (s, 2 H), 6.41 (dd, 1 H), 6.34 (s, 1 H), 5.57 (dd, 1 H), 3.16 - 3.06 (m, 1 H), 2.80 - 2.70 (m, 1 H), 2.60 - 2.54 (m, 1 H), 2.20 - 2.10 (m, 1 H).

Example 721(7): methyl [6-(4-chloro-2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS *t<sub>R</sub>* 3.83 minutes; MS (ES<sup>+</sup>) *m/z* 565 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.15 (s, 1 H), 10.06 (s, 1 H), 9.80 (s, 1 H), 8.80 (s, 1 H), 8.06 (d, 1 H), 8.03 (dd, 1 H), 7.97 (d, 1 H), 7.92 (dd, 1 H), 7.88 (d, 1 H), 6.38 (s, 1 H), 5.71 (dd, 1 H), 3.77 (s, 3 H), 3.20 - 3.08 (m, 1 H), 2.89 - 2.77 (m, 1 H), 2.66 - 2.58 (m, 1 H), 2.24 - 2.14 (m, 1 H).

Example 721(8): 2-methoxyethyl [6-(4-chloro-2-{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS *t<sub>R</sub>* 3.87 minutes; MS (ES<sup>+</sup>) *m/z* 609 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.09 (s, 1 H), 10.12 (s, 1 H), 9.74 (s, 1 H), 8.76 (d, 1 H), 8.00 (d, 1 H), 7.97 (d, 1 H), 7.91 (d, 1 H), 7.86 (dd, 1 H), 7.82 (s, 1 H), 6.33 (s, 1 H), 5.65 (dd, 1 H), 4.29 - 4.21 (m, 2 H), 3.64 - 3.53 (m, 2 H), 3.30 (s, 3 H), 3.13 - 3.01 (m, 1 H), 2.83 - 2.70 (m, 1 H), 2.60 - 2.53 (m, 1 H), 2.18 - 2.07 (m, 1 H).

Example 721(9): methyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.77 minutes; MS ( $ES^+$ )  $m/z$  565 ( $M+H$ )<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.20 (s, 1 H), 10.41 (s, 1 H), 9.81 (s, 1 H), 8.62 (s, 1 H), 8.09 - 7.90 (m, 3 H), 7.88 - 7.80 (m, 2 H), 6.36 (s, 1 H), 5.61 - 5.53 (m, 1 H), 3.73 - 3.64 (s, 3 H), 3.21 - 3.08 (m, 1 H), 2.83 - 2.69 (m, 1 H), 2.61 - 2.56 (m, 1 H), 2.21 - 2.14 (m, 1 H).

- 10 Example 721(10): 2-methoxyethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.77 minutes; MS ( $ES^+$ )  $m/z$  609 ( $M+H$ )<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.46 (s, 1 H), 8.60 (d, 1 H), 8.08 (dd, 1 H), 8.00 (d, 1 H), 7.95 (d, 1 H), 7.79 (dd, 1 H), 7.72 (d, 1 H), 6.39 (s, 1 H), 5.70 (dd, 1 H), 4.34 (t, 2 H), 3.69 (t, 2 H), 3.42 (s, 3 H), 3.30 - 3.27 (m, 1 H), 2.98 - 2.90 (m, 1 H), 2.75 - 2.65 (m, 1 H), 2.41 - 2.33 (m, 1 H).

- 20 Example 722: ((6S)-6-[5-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

The same operation as in Example 484 → Example 364 → Example 584 was conducted from the compound prepared in Example 718(6) to give the title compound having the following physical properties. (Note: in the step corresponding to Example 484 in the operation, allyl chloroformate was used).

- 25 LC/MS  $t_R$  2.80 minutes; MS ( $ES^+$ )  $m/z$  491 ( $M+H$ )<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.43 (s, 1 H), 8.09 (s, 1 H), 7.89 (d, 1 H), 7.75 (dd, 1 H), 7.68 (d, 1 H), 7.62 (dd, 1 H), 6.63 (d, 1 H), 6.34 (s, 1 H), 5.62 (dd, 1 H), 3.29 - 3.21 (m, 1 H), 2.89 (ddd, 1 H), 2.65 (qd, 1 H), 2.37 - 2.28 (m, 1 H).

- 30 Example 723: ((6S)-6-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one (enantiopure)

The same operation as in Example 128 → Example 338 → Example 584 was conducted from the compound prepared in Example 718(6) to give the title compound having the following physical properties. (Note: in the step corresponding to Example 128 in the operation, allyl chloroformate was used).

- 5 LC/MS  $t_R$  2.78 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.46 (s, 1 H), 8.24 - 8.20 (m, 1 H), 7.93 (d, 1 H), 7.81 - 7.76 (m, 2 H), 7.74 - 7.69 (m, 1 H), 6.67 (d, 1 H), 6.38 (s, 1 H), 5.69 - 5.63 (m, 1 H), 3.31 - 3.25 (m, 1 H), 2.97 - 2.87 (m, 1 H), 2.73 - 2.63 (m, 1 H), 2.40 - 2.30 (m, 1 H).

- 10 Example 724: methyl [4-(5-{{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}}-1H-imidazol-2-yl)phenyl]carbamate

The same operation as in Example 234 → Example 239 was conducted from the compound prepared in Example 336 to give the title compound having the following physical properties.

- 15 LC/MS  $t_R$  2.93 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.82 (s, 1 H), 7.73 - 7.65 (m, 3 H), 7.62 (dd, 1 H), 7.50 (d, 1 H), 7.38 (d, 2 H), 7.00 (br. s, 1 H), 6.94 (s, 1 H), 6.34 (s, 1 H), 5.75 (dd, 1 H), 3.79 (s, 3 H), 3.48 - 3.36 (m, 1 H), 2.88 (ddd, 1 H), 2.60 - 2.47 (m, 2 H).

- 20 Example 725: methyl [4-(4-chloro-5-{{(6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}}-1H-imidazol-2-yl)phenyl]carbamate

The compound prepared in Example 724 (55 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (5.6 mg).

- 25 LC/MS  $t_R$  3.70 minutes; MS (ES<sup>+</sup>)  $m/z$  564 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.43 (s, 1 H), 7.90 (d, 1 H), 7.77 (dd, 1 H), 7.73 (d, 2 H), 7.71 (d, 1 H), 7.54 (d, 2 H), 6.42 (s, 1 H), 5.76 (dd, 1 H), 3.75 (s, 3 H), 3.26 - 3.14 (m, 1 H), 2.96 (ddd, 1 H), 2.76 - 2.64 (m, 1 H), 2.28 (tdd, 1 H).

30

- Example 726: 6-(5-chloro-2-nitrophenyl)-2-(propan-2-yl)-3,4-dihydropyrimidin-4-one

To an N,N-dimethylformamide (3 mL) solution of the compound prepared in Example 33 (0.60 g) was added potassium carbonate (0.46 g) and 2-



methylpropanimidamide hydrochloride (0.30 g) and the mixture stirred at 100 °C for 5 hours. The reaction mixture was then concentrated and the residue purified by column chromatography (10 to 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (0.31 g).

- 5 <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 12.29 (br. s, 1 H), 7.86 (d, 1 H), 7.61 - 7.52 (m, 2 H), 6.56 (s, 1 H), 2.93 (td, 1 H), 1.32 (d, 6 H).

Example 727: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-isopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl}carbamate

- 10 The same operation as in Example 322 → Example 74 → Example 9 → Example 25 → Example 51 → Example 52 was conducted from the compound prepared in Example 726 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).
- 15 LC/MS *t<sub>R</sub>* 3.40 minutes; MS (ES<sup>+</sup>) *m/z* 546 (M+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.16 (br. s, 1 H), 9.77 (s, 1 H), 9.60 (br. s, 1 H), 7.97 (d, 1 H), 7.89 - 7.84 (m, 1 H), 7.83 - 7.78 (m, 1 H), 7.59 (d, 2 H), 7.44 (s, 1 H), 7.40 (d, 2 H), 6.71 (s, 1 H), 5.25 (s, 2 H), 3.65 (s, 3 H), 3.32 - 3.30 (obs. m, 1 H), 0.87 (d, 6 H).

- 20 Example 728: methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-isopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl}carbamate

The compound prepared in Example 727 (55 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (17 mg).

- 25 LC/MS *t<sub>R</sub>* 4.32 minutes; MS (ES<sup>+</sup>) *m/z* 580 (M+H)<sup>b</sup>
- <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.87 (br. s, 1 H), 9.80 (br. s, 1 H), 9.79 (s, 1 H), 8.00 (d, 1 H), 7.88 - 7.84 (m, 1H), 7.84 - 7.79 (m, 1 H), 7.62 - 7.57 (m, 2 H), 7.57 - 7.49 (m, 2 H), 6.71 (s, 1 H), 5.23 (s, 2 H), 3.68 (s, 3 H), 3.16 (td, 1 H), 0.84 (d, 6 H).

- 30 Example 729: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-cyclopropyl-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 726 → Example 20 → Example 74 → Example 9 → Example 25 → Example 51 → Example 52 was conducted from

cyclopropanecarboximidamide hydrochloride to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).

- 5 LC/MS  $t_R$  3.30 minutes; MS (ES<sup>+</sup>)  $m/z$  544 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.41 (s, 1 H), 7.87 (d, 1 H), 7.78 - 7.72 (m, 1 H), 7.67 (d, 1 H), 7.60 (m, 2 H), 7.44 (m, 2 H), 7.37 - 7.25 (br. s, 1 H), 6.61 (s, 1 H), 5.52 (s, 2 H), 3.74 (s, 3 H), 2.22 - 2.10 (m, 1 H), 0.89 (m, 2 H), 0.63 - 0.56 (m, 2 H).

10 Example 730: methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-cyclopropyl-6-oxo-1(6H)-pyrimidinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The compound prepared in Example 729 (70 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (40 mg).

- 15 LC/MS  $t_R$  4.21 minutes; MS (ES<sup>+</sup>)  $m/z$  578 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.87 (br. s, 1 H), 9.81 (br. s, 1 H), 9.74 (s, 1 H), 7.97 (d, 1 H), 7.88 - 7.77 (m, 2 H), 7.64 - 7.50 (m, 4 H), 6.64 (s, 1 H), 5.36 (s, 2 H), 3.68 (s, 3 H), 2.20 - 2.09 (m, 1 H), 0.81 (m, 2 H), 0.44 (m, 2 H).

20 Example 731: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-cyclopropyl-6-oxo-1(6H)-pyrimidinyl}methyl)-4-fluoro-1H-imidazol-5-yl]phenyl}carbamate

The compound prepared in Example 729 (70 mg) was treated as detailed in Example 364 to give the title compound having the following physical properties (20 mg).

- 25 LC/MS  $t_R$  4.18 minutes; MS (ES<sup>+</sup>)  $m/z$  562 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.71 (br. s, 1 H), 9.74 (m, 2 H), 7.95 (s, 1 H), 7.87 - 7.75 (m, 2 H), 7.57 - 7.40 (m, 4 H), 6.63 (s, 1 H), 5.34 (s, 2 H), 3.67 (s, 3 H), 2.23 - 2.04 (m, 1 H), 0.88 - 0.72 (m, 2 H), 0.44 (m, 2 H).

30 Example 732: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-oxo-2-(tetrahydro-2H-pyran-4-yl)-1(6H)-pyrimidinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 726 → Example 20 → Example 74 → Example 9 → Example 25 → Example 51 → Example 52 was conducted from tetrahydro-2H-pyran-4-carboximidamide hydrochloride to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).

LC/MS  $t_R$  3.27 minutes; MS (ES<sup>+</sup>)  $m/z$  588 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.20 (br. s, 1 H), 9.79 (s, 1 H), 9.60 (br. s, 1 H), 7.96 (d, 1 H), 7.85 (d, 1 H), 7.82 (s, 1 H), 7.58 (d, 2 H), 7.44 (s, 1 H), 7.39 (d, 2 H), 6.68 (s, 1 H), 5.31 (s, 2 H), 3.77 (m, 2 H), 3.65 (s, 3 H), 3.41 (m, 1 H), 3.29 (t, 2 H), 1.44 (m, 2 H), 1.28 - 1.18 (m, 2 H).

Example 733: methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-(methoxymethyl)-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl}phenyl}carbamate

The same operation as in Example 726 → Example 20 → Example 74 → Example 9 → Example 25 → Example 51 → Example 52 was conducted from methoxyacetamide hydrochloride to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used).

LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.13 (br. s, 1 H), 9.78 (s, 1 H), 9.60 (br. s, 1 H), 7.99 (d, 1 H), 7.89 - 7.85 (m, 1 H), 7.84 - 7.80 (m, 1 H), 7.59 (d, 2 H), 7.44 - 7.36 (m, 3 H), 6.70 (s, 1 H), 5.24 (s, 2 H), 4.49 (s, 2 H), 3.65 (s, 3 H), 3.15 (s, 3 H).

Example 734: methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl}-2-(methoxymethyl)-6-oxo-1(6H)-pyrimidinyl)methyl]-1H-imidazol-5-yl}phenyl}carbamate

The compound prepared in Example 733 (85 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (26 mg).

LC/MS  $t_R$  4.04 minutes; MS (ES<sup>+</sup>)  $m/z$  582 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.84 (br. s, 1 H), 9.86 - 9.73 (m, 2 H), 8.03 (s, 1 H), 7.92 - 7.78 (m, 2 H), 7.63 - 7.47 (m, 4 H), 6.71 (s, 1 H), 5.21 (s, 2 H), 4.35 (s, 2 H), 3.67 (s, 3 H), 3.12 (s, 3 H).

5 Example 735: (3S)-3-[5-(4-aminophenyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 74 was conducted from the compound prepared in Example 375 to give the title compound having the following physical properties. (Note: in the step corresponding to Example

10 51 in the operation, 2-bromo-1-(4-nitrophenyl)ethanone was used).

LC/MS *t<sub>R</sub>* 2.72 minutes; MS (ES<sup>+</sup>) *m/z* 451 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.32 (s, 1 H), 7.53 (m, 2 H), 7.49 (s, 1 H), 7.37 (m, 2 H), 7.04 (br. s, 1 H), 6.74 (d, 2 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.77 (dd, 1 H), 3.48 - 3.38 (m, 1 H), 3.13 - 3.04 (m, 1 H), 2.65 - 2.56 (m, 1 H), 2.52 (s, 3 H), 2.51 - 2.41 (m, 1

15 H).

Example 736: 2-methoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 735 (40 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (36 mg).

LC/MS *t<sub>R</sub>* 3.12 minutes; MS (ES<sup>+</sup>) *m/z* 553 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.30 (s, 1 H), 7.55 (d, 2 H), 7.52 - 7.45 (m, 3 H), 7.43 (d, 2 H), 7.21 (s, 1 H), 6.09 (s, 1 H), 6.05 (s, 1 H), 5.77 (dd, 1 H), 4.28 - 4.22 (m, 2

25 H), 3.67 - 3.61 (m, 2 H), 3.47 - 3.40 (m, 1 H), 3.38 (s, 3 H), 3.12 - 3.02 (m, 1 H), 2.66 - 2.55 (m, 1 H), 2.50 (s, 3 H), 2.48 - 2.40 (m, 1 H).

Example 737: 3-oxetanyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate

The compound prepared in Example 735 (100 mg) was treated as detailed in Example 494 to give the title compound having the following physical properties (15 mg).

LC/MS *t<sub>R</sub>* 2.93 minutes; MS (ES<sup>+</sup>) *m/z* 551 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.31 (s, 1 H), 7.65 - 7.54 (m, 2 H), 7.54 - 7.46 (m, 3 H), 7.46 - 7.36 (m, 2 H), 7.36 - 7.20 (m, 1 H), 6.10 (s, 1 H), 6.06 (s, 1 H), 5.77 (d, 1 H), 5.47 (quintet, 1 H), 4.92 (t, 2 H), 4.68 (dd, 2 H), 3.50 - 3.37 (m, 1 H), 3.08 (ddd, 1 H), 2.62 (qd, 1 H), 2.51 (s, 3 H), 2.49 - 2.39 (m, 1 H).

5

Example 738(1) to Example 738(12)

The compounds of the present invention having the following physical data were prepared using the corresponding alpha-bromoketones from the compound prepared in Example 375 in the process of Example 51 → Example 52.

10

Example 738(1) and 738(2): methyl [4-(4-methyl-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]carbamate and methyl [4-(5-methyl-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1,3-oxazol-4-yl)phenyl]carbamate

15

The title compounds were obtained in a (1.9 : 1) ratio and separated by column chromatography. (Note: in the step corresponding to Example 51 in the process, the compound prepared in Example 78 was used).

Example 738(1):

20 LC/MS *t<sub>R</sub>* 2.94 minutes; MS (ES<sup>+</sup>) *m/z* 523 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.88 (s, 1 H), 9.65 (s, 1 H), 9.59 (br. s, 1 H), 7.59 (d, 1 H), 7.50 (m, 2 H), 7.45 (m, 3 H), 7.38 (d, 1 H), 5.93 (s, 1 H), 5.85 (s, 1 H), 5.59 - 5.49 (m, 1 H), 3.65 (s, 3 H), 3.40 - 3.38 (m, 1 H), 3.36 - 3.33 (obs. m, 1 H), 3.02 - 2.90 (m, 1 H), 2.45 (s, 3 H), 2.35 - 2.26 (s, 3 H), 2.23 - 2.16 (m, 1 H).

25

Example 738(2):

LC/MS *t<sub>R</sub>* 4.02 minutes; MS (ES<sup>+</sup>) *m/z* 524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.31 (s, 1 H), 7.54 - 7.46 (m, 7 H), 6.13 (s, 1 H), 6.05 (s, 1 H), 5.76 (dd, 1 H), 3.73 (s, 3 H), 3.34 (s, 3 H), 3.32 - 3.27 (obs. m, 1 H), 3.17 - 3.01 (m, 1 H), 2.70 - 2.58 (m, 1 H), 2.51 (s, 3 H), 2.48 (s, 3 H), 2.43 - 2.24 (m, 1 H).

30

Example 738(3): 2-methoxyethyl [4-(4-methyl-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 481 was used.

LC/MS  $t_R$  3.11 minutes; MS ( $ES^+$ )  $m/z$  567 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.31 (s, 1 H), 7.54 - 7.49 (m, 3 H), 7.49 - 7.45 (m, 2 H), 7.45 - 7.37 (m, 2 H), 6.10 (s, 1 H), 6.06 (s, 1 H), 5.71 (d, 1 H), 4.28 - 4.25 (m, 2 H), 3.66 - 3.62 (m, 2 H), 3.48 - 3.41 (m, 1 H), 3.39 (s, 3 H), 3.07 (ddd, 1 H), 2.61 (qd, 1 H), 2.51 (s, 3 H), 2.47 - 2.37 (m, 1 H), 2.31 (app. br. s, 3 H).

10 Example 738(4): 2-ethoxyethyl [4-(2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l}-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 714 was used.

LC/MS  $t_R$  3.13 minutes; MS ( $ES^+$ )  $m/z$  567 ( $M+H$ )<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.32 (s, 1 H), 7.57 (d, 2 H), 7.54 - 7.47 (m, 3 H), 7.45 (d, 2 H), 7.23 (s, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.78 (dd, 1 H), 4.28 - 4.25 (m, 2 H), 3.71 - 3.67 (m, 2 H), 3.60 - 3.54 (m, 2 H), 3.48 - 3.39 (m, 1 H), 3.13 - 3.04 (m, 1 H), 2.63 (qd, 1 H), 2.52 (s, 3 H), 2.50 - 2.43 (m, 1 H), 1.20 (t, 3 H).

20 Example 738(5): 3-methoxypropyl [4-(2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]l}-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 715 was used.

LC/MS  $t_R$  3.11 minutes; MS ( $ES^+$ )  $m/z$  567 ( $M+H$ )<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.32 (s, 1 H), 7.64 - 7.56 (m, 1 H), 7.55 - 7.47 (m, 4 H), 7.46 - 7.38 (m, 2 H), 7.29 (br. s, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.78 (d, 1 H), 4.21 (t, 2 H), 3.53 (t, 2 H), 3.48 - 3.39 (m, 1 H), 3.35 (s, 3 H), 3.09 (ddd, 1 H), 2.68 - 2.57 (m, 1 H), 2.52 (s, 3 H), 2.50 - 2.39 (m, 1 H), 1.94 (quintet, 2 H).

30 Example 738(6): (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

LC/MS  $t_R$  2.67 minutes; MS ( $ES^+$ )  $m/z$  452 ( $M+H$ ) 227 ( $M+H$ )/2<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.32 (s, 1 H), 8.29 - 8.08 (m, 1 H), 7.81 - 7.67 (m, 1 H), 7.57 - 7.50 (m, 2 H), 7.49 (s, 1 H), 7.36 - 6.99 (m, 1 H), 6.61 (d, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.76 (dd, 1 H), 3.51 - 3.38 (m, 1 H), 3.15 - 3.01 (dd, 1 H), 2.70 - 2.56 (m, 1 H), 2.52 (s, 3 H), 2.50 - 2.32 (m, 1 H).

5

Example 738(7): (3S)-3-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 551 was used.

10 LC/MS *t<sub>R</sub>* 2.76 minutes; MS (ES<sup>+</sup>) *m/z* 466 (M+H) 234 (M+H)/2 <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.32 (s, 1 H), 8.33 - 8.17 (m, 1 H), 7.79 - 7.64 (m, 1 H), 7.57 - 7.51 (m, 2 H), 7.49 (s, 1 H), 7.21 (br. s, 1 H), 6.54 (d, 1 H), 6.10 (s, 1 H), 6.07 (s, 1 H), 5.76 (dd, 1 H), 3.53 - 3.38 (m, 1 H), 3.09 (ddd, 1 H), 2.90 (s, 3 H), 2.69 - 2.58 (m, 1 H), 2.52 (s, 3 H), 2.49 - 2.37 (m, 1 H).

15

Example 738(8): (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 215 was used.

20 LC/MS *t<sub>R</sub>* 3.06 minutes; MS (ES<sup>+</sup>) *m/z* 470 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.31 (s, 1 H), 8.16 - 7.76 (m, 1 H), 7.55 - 7.49 (m, 2 H), 7.47 (s, 1 H), 7.14 (br. s, 1 H), 6.45 (d, 1 H), 6.12 (s, 1 H), 6.05 (s, 1 H), 5.78 (d, 1 H), 3.48 - 3.37 (m, 1 H), 3.08 (dd, 1 H), 2.68 - 2.55 (m, 1 H), 2.53 - 2.43 (m, 4 H).

25 Example 738(9): (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 209 was used.

LC/MS *t<sub>R</sub>* 3.11 minutes; MS (ES<sup>+</sup>) *m/z* 486 (M+H) <sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.31 (s, 1 H), 7.99 - 7.65 (m, 1 H), 7.55 - 7.50 (m, 2 H), 7.48 (s, 1 H), 7.37 (br. s, 1 H), 6.56 (d, 1 H), 6.12 (s, 1 H), 5.79 (dd, 1 H), 3.49 - 3.38 (m, 1 H), 3.09 (dd, 1 H), 2.65 - 2.57 (m, 1 H), 2.55 - 2.44 (m, 4 H).

Example 738(10): methyl [6-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 712 was used.

5 LC/MS  $t_R$  3.07 minutes; MS (ES<sup>+</sup>)  $m/z$  510 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.32 (s, 1 H), 8.64 - 8.46 (m, 1 H), 7.95 (d, 1 H), 7.81 (br. s, 1 H), 7.58 - 7.30 (m, 4 H), 6.13 (s, 1 H), 6.06 (s, 1 H), 5.80 (d, 1 H), 3.77 (s, 3 H), 3.50 - 3.40 (m, 1 H), 3.10 (ddd, Hz, 1 H), 2.68 - 2.56 (m, 1 H), 2.52 (m, 4 H).

10 Example 738(11): formic acid - 2-methoxyethyl [6-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 713 was used. In the step corresponding to Example 52 in the process, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt.

15 LC/MS  $t_R$  2.98 minutes; MS (ES<sup>+</sup>)  $m/z$  554 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.32 (s, 1 H), 8.58 (s, 1 H), 8.17 (s, 1 H), 7.96 (d, 1 H), 7.76 (d, 1 H), 7.56 - 7.46 (m, 4 H), 6.13 (s, 1 H), 6.07 (s, 1 H), 5.81 (dd, 1 H), 4.33 - 4.25 (m, 2 H), 3.69 - 3.62 (m, 2 H), 3.52 - 3.41 (m, 1 H), 3.40 (s, 3 H), 3.10 (ddd, 1 H), 2.69 - 2.57 (m, 1 H), 2.55 - 2.45 (m, 4 H).

25 Example 738(12): 6-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone

In the step corresponding to Example 51 in the process, 6-(bromoacetyl)-3,4-dihydroquinolin-2(1H)-one was used.

LC/MS  $t_R$  2.97 minutes; MS (ES<sup>+</sup>)  $m/z$  505 (M+H)<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.32 (s, 1 H), 7.55 - 7.43 (m, 5 H), 7.24 (br. s, 1 H), 6.86 (d, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.77 (dd, 1 H), 3.44 (td, 1 H), 3.09 (ddd, 1 H), 2.98 (t, 2 H), 2.68 - 2.60 (m, 1 H), 2.60 - 2.55 (m, 2 H), 2.52 (s, 3 H), 2.50 - 2.42 (m, 1 H).



Example 739: methyl [5-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 738(6) (200 mg) was treated with methyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (213 mg).

LC/MS  $t_R$  3.12 minutes; MS ( $ES^+$ )  $m/z$  510 (M+H) 256 (M+H)/2<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.32 (s, 1 H), 8.62 - 8.48 (m, 1 H), 8.10 - 7.97 (m, 1 H), 7.88 (d, 1 H), 7.55 - 7.47 (m, 3 H), 7.40 (br. s, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.78 (dd, 1 H), 3.77 (s, 3 H), 3.51 - 3.39 (m, 1 H), 3.10 (ddd, 1 H), 2.70 - 2.58 (m, 1 H), 2.52 (s, 4 H).

Example 740: 2-methoxyethyl [5-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 738(6) (124 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 484 to give the title compound having the following physical properties (81 mg).

LC/MS  $t_R$  3.13 minutes; MS ( $ES^+$ )  $m/z$  554 (M+H) 278 (M+H)/2<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.32 (s, 1 H), 8.62 - 8.49 (m, 1 H), 8.03 (d, 1 H), 7.89 (d, 1 H), 7.55 - 7.47 (m, 3 H), 7.40 (br. s, 1 H), 6.11 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 4.31 (dd, 2 H), 3.69 - 3.64 (m, 2 H), 3.51 - 3.43 (m, 1 H), 3.40 (s, 3 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.52 (s, 4 H).

Example 741(1) to Example 741(3):

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 738(4), 738(5) and 738(11) using the method as detailed in Example 338.

Example 741(1): 2-ethoxyethyl [4-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  4.14 minutes; MS ( $ES^+$ )  $m/z$  601 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.32 (s, 1 H), 7.62 (d, 2 H), 7.56 - 7.51 (m, 4 H), 7.50 (s, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.71 (dd, 1 H), 4.30 - 4.25 (m, 2 H), 3.72 -

3.67 (m, 2 H), 3.57 (q, 2 H), 3.48 - 3.38 (m, 1 H), 3.09 (ddd, 1 H), 2.69 - 2.57 (m, 1 H), 2.52 (s, 3 H), 2.44 - 2.35 (m, 1 H), 1.21 (t, 3 H).

Example 741(2): 3-methoxypropyl [4-(4-chloro-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate  
LC/MS  $t_R$  4.10 minutes; MS (ES<sup>+</sup>)  $m/z$  601 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.33 (s, 1 H), 7.62 (d, 2 H), 7.56 - 7.51 (m, 4 H), 7.50 (s, 1 H), 6.11 (s, 1 H), 6.07 (s, 1 H), 5.71 (dd, 1 H), 4.25 - 4.20 (m, 2 H), 3.53 (t, 2 H), 3.44 (td, 1 H), 3.36 (s, 3 H), 3.09 (ddd, 1 H), 2.64 (qd, 1 H), 2.52 (s, 3 H), 2.44 - 2.35 (m, 1 H), 1.95 (quintet, 2 H).

Example 741(3): 2-methoxyethyl [6-(4-chloro-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS  $t_R$  3.96 minutes; MS (ES<sup>+</sup>)  $m/z$  588 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.32 (s, 1 H), 8.68 - 8.61 (m, 1 H), 8.01 - 7.89 (m, 2 H), 7.56 - 7.45 (m, 3 H), 6.13 (s, 1 H), 6.06 (s, 1 H), 5.78 (dd, 1 H), 4.33 - 4.26 (m, 2 H), 3.71 - 3.62 (m, 2 H), 3.44 - 3.35 (m, 4 H), 3.14 - 3.04 (m, 1 H), 2.66 - 2.55 (m, 1 H), 2.51 (s, 3 H), 2.48 - 2.41 (m, 1 H).

Example 742(1) to Example 742(5):

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 738(6), 738(7), 738(10), 739 and 740 using the method as detailed in Example 44.

Example 742(1): (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  2.96 minutes; MS (ES<sup>+</sup>)  $m/z$  486 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.32 (s, 1 H), 8.21 (d, 1 H), 7.76 (dd, 1 H), 7.55 - 7.51 (m, 2 H), 7.51 - 7.48 (m, 1 H), 6.65 (d, 1 H), 6.10 (s, 1 H), 6.07 (s, 1 H), 5.69 (dd, 1 H), 3.48 - 3.39 (m, 1 H), 3.09 (ddd, 1 H), 2.69 - 2.57 (m, 1 H), 2.52 (s, 3 H), 2.42 - 2.33 (m, 1 H).

Example 742(2): formic acid - (3S)-3-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone (1:1)

Purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] gave the title compound as the formic acid salt.

LC/MS  $t_R$  3.00 minutes; MS ( $ES^+$ )  $m/z$  500 (M+H) 250 (M+H)/2 <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.32 (s, 1 H), 8.26 (d, 1 H), 8.15 (s, 1 H), 7.78 (dd, 1 H), 7.55 - 7.51 (m, 2 H), 7.50 (s, 1 H), 6.64 (d, 1 H), 6.10 (s, 1 H), 6.07 (s, 1 H), 5.69 (dd, 1 H), 3.48 - 3.38 (m, 1 H), 3.09 (ddd, 1 H), 2.91 (s, 3 H), 2.63 (qd, 1 H), 2.52 (s, 3 H), 2.42 - 2.33 (m, 1 H).

Example 742(3): formic acid - methyl [6-(4-chloro-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-

pyridinyl]carbamate (1:1)

LC/MS  $t_R$  3.95 minutes; MS ( $ES^+$ )  $m/z$  544 (M+H) 250 (M+H)/2 <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.96 (br. s, 1 H), 10.00 (br. s, 1 H), 9.67 (s, 1 H), 8.73 (s, 1 H), 8.02 - 7.93 (m, 1 H), 7.92 - 7.86 (m, 1 H), 7.60 (d, 1 H), 7.55 - 7.47 (m, 2 H), 5.95 (s, 1 H), 5.86 (s, 1 H), 5.66 (dd, 1 H), 3.70 (s, 3 H), 3.21 - 3.16 (m, 1 H), 3.02 - 2.92 (m, 1 H), 2.55 - 2.48 (obs. m, 1 H), 2.46 (s, 3 H), 2.26 - 2.10 (m, 1 H).

Example 742(4): methyl [5-(4-chloro-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.88 minutes; MS ( $ES^+$ )  $m/z$  544 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.33 (s, 1 H), 8.58 (d, 1 H), 8.07 - 8.02 (m, 1 H), 8.02 - 7.96 (m, 1 H), 7.54 - 7.52 (m, 2 H), 7.50 (s, 1 H), 6.11 (s, 1 H), 6.08 (s, 1 H), 5.71 (dd, 1 H), 3.79 (s, 3 H), 3.45 (td, 1 H), 3.10 (ddd, 1 H), 2.65 (qd, 1 H), 2.53 (s, 3 H), 2.44 - 2.35 (m, 1 H).


Example 742(5): 2-methoxyethyl [5-(4-chloro-2-{(3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.87 minutes; MS ( $ES^+$ )  $m/z$  588 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.09 (br. s, 1 H), 10.37 (s, 1 H), 9.68 (s, 1 H), 8.57 (d, 1 H), 8.04 (dd, 1 H), 7.91 (d, 1 H), 7.60 (d, 1 H), 7.55 - 7.49 (m, 2 H), 5.96 (s, 1 H), 5.86 (s, 1 H), 5.57 (dd, 1 H), 4.24 (dd, 2 H), 3.58 (dd, 2 H), 3.29 (s, 3 H), 3.27 - 3.22 (m, 1 H), 3.00 (dd, 1 H), 2.55 - 2.53 (m, 1 H), 2.47 (s, 3 H), 2.24 - 2.14 (m, 1 H).

5

Example 743: ethyl (6S)-2-hydroxy-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylate



To a 1,4-dioxane (570 mL) suspension of ethyl (2S)-5-aminopyrrolidine-2-carboxylate hydrochloride [J. Org. Chem. 52(26), 5717 (1987)] (28.3 g) was sequentially added triethylamine (51.2 mL) and ethyl malonyl chloride (20.8 mL) and the mixture stirred at 80 °C for 1 hour. The reaction mixture was cooled to room temperature, diluted with ethyl acetate (500 mL), filtered and the filtrate concentrated to give the crude title compound having the following physical properties (42.6 g).

LC/MS *t<sub>R</sub>* 0.80 minutes; MS (ES<sup>+</sup>) *m/z* 225 (M+H)<sup>+</sup>.

15

Example 744: ethyl (6S)-2-chloro-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylate

The compound prepared in Example 743 (30.2 g) was suspended in phosphorous(V) oxychloride (125 mL) and the mixture stirred at 100 °C for 1 hour. The reaction mixture was cooled to room temperature, concentrated, the residue suspended in a 1 M aqueous solution of sodium hydrogen carbonate (100 mL) and extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 to 70% ethyl acetate in heptanes) to give the title compound having the following physical properties (5.31 g).

25


LC/MS *t<sub>R</sub>* 1.43 minutes; MS (ES<sup>+</sup>) *m/z* 265 (M+Na), 243 (M+H)<sup>+</sup>.

30

Example 745: ethyl (6S)-2-(2-amino-5-methylphenyl)-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylate

To a 1,4-dioxane (290 mL) solution of the compound prepared in Example 744 (7.32 g) was added 4-methyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline hydrochloride (9.50 g) and a 2 M aqueous solution of sodium carbonate (45.2 mL) under an atmosphere of nitrogen. The mixture was degassed with nitrogen, 1,1'-bis(diphenylphosphino)ferrocene palladium (II) dichloride dichloromethane complex (1.31 g) added and the mixture stirred at 100 °C for 4 hours before cooling to room temperature. The reaction mixture was diluted with ethyl acetate (300 mL) and the organic layer isolated, dried and concentrated. The residue was purified by column chromatography (10% to 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (6.50 g).  
LC/MS  $t_R$  1.52 minutes; MS ( $ES^+$ )  $m/z$  314 ( $M+H$ )<sup>a</sup>.

Example 746: (6S)-2-[5-methyl-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylic acid



15

The same operation as in Example 8 → Example 9 was conducted from the compound prepared in Example 745 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.37 minutes; MS ( $ES^+$ )  $m/z$  699 ( $2M+Na$ ), 677 ( $2M+H$ ), 361 ( $M+Na$ ), 339 ( $M+H$ ), 311 ( $M-N_2+H$ )<sup>a</sup>.

20

Example 747(1) to Example 747(11)

The compounds of the present invention having the following physical data were prepared using the corresponding alpha-bromoketones from the compound prepared in Example 746 in the process of Example 51 → Example 52.

25

Example 747(1): methyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used.

LC/MS  $t_R$  2.99 minutes; MS ( $ES^+$ )  $m/z$  510 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.40 (s, 1 H), 7.67 (s, 1 H), 7.63 - 7.51 (m, 4 H),  
5 7.44 (br. s, 2 H), 7.29 (br. s, 1 H), 6.33 (s, 1 H), 5.72 (dd, 1 H), 3.74 (s, 3 H), 3.33 (br. s, 1 H), 3.30 - 3.23 (m, 1 H), 2.96 - 2.84 (m, 1 H), 2.66 (qd, 1 H), 2.52 (s, 3 H), 2.39 (d, 1 H).

10 Example 747(2): 2-methoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl]phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 324 was used.

LC/MS  $t_R$  3.04 minutes; MS ( $ES^+$ )  $m/z$  554 ( $M+H$ )<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.41 (s, 1 H), 7.68 (s, 1 H), 7.58 (app. br. s, 2 H), 7.56 (app. s, 2 H), 7.46 (app. br. s, 2 H), 7.28 (br. s, 1 H), 6.35 (s, 1 H), 5.74 (dd, 1 H), 4.28 (dd, 2 H), 3.66 (dd, 2 H), 3.41 (s, 3 H), 3.35 - 3.25 (m, 1 H), 2.91 (ddd, 1 H), 2.68 (qd, 1 H), 2.54 (s, 3 H), 2.40 (app. br. s, 1 H).

20 Example 747(3): 2-ethoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl]phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 714 was used.

25 LC/MS  $t_R$  3.11 minutes; MS ( $ES^+$ )  $m/z$  568 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.40 (s, 1 H), 7.70 - 7.13 (m, 8 H), 6.33 (s, 1 H), 5.72 (dd, 1 H), 4.33 - 4.19 (m, 2 H), 3.73 - 3.65 (m, 2 H), 3.57 (q, 2 H), 3.33 (m, 1 H), 2.90 (m, 1 H), 2.71 - 2.61 (m, 1 H), 2.53 (s, 3 H), 2.40 (br. s, 1 H), 1.20 (t, 3 H).

30 Example 747(4): 3-methoxypropyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl]phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 715 was used.

LC/MS  $t_R$  1.54 minutes; MS (ES<sup>+</sup>)  $m/z$  568 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.39 (s, 1 H), 7.84 - 6.91 (m, 9 H), 6.33 (s, 1 H),  
5 5.72 (dd, 1 H), 4.62 (br. s, 1 H), 4.20 (t, 2 H), 3.52 (t, 2 H), 3.34 (s, 3 H), 3.29 - 3.20 (m,  
1 H), 2.90 (ddd, 1 H), 2.65 (m, 1 H), 2.52 (s, 3 H), 2.39 (br. s, 1 H), 1.93 (quintet, 2 H).

Example 747(5): formic acid - methyl [4-(4-methyl-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 78 was used. In the step corresponding to Example 52 in the process, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title  
15 product as the formic acid salt

LC/MS  $t_R$  2.93 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.03 (br. s, 1 H), 9.69 (br. s, 1 H), 9.64 (br. s, 1 H),  
8.21 (br. s, 1 H), 7.83 - 7.24 (m, 7 H), 6.18 (br. s, 1 H), 5.55 (d, 1 H), 3.66 (s, 3 H), 3.68  
- 3.56 (obs. m, 1 H), 3.28 - 3.10 (m, 1 H), 2.84 - 2.68 (m, 1 H), 2.47 (s, 3 H), 2.29 (br. s,  
20 3 H), 2.19 (br. s, 1 H).

Example 747(6): 2-methoxyethyl [4-(4-methyl-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 481 was used.

LC/MS  $t_R$  3.03 minutes; MS (ES<sup>+</sup>)  $m/z$  568 (M+H)<sup>b</sup>

NMR analysis showed a 3: 2 ratio of tautomers.

Major tautomer: <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.00 (s, 1 H), 9.70 (br. s, 1 H), 9.69  
30 (s, 1 H), 7.70 (br. s, 1 H), 7.61 (s, 1 H), 7.59 - 7.35 (m, 5 H), 6.19 (s, 1 H), 5.55 (dd, 1  
H), 4.21 - 4.17 (m, 2 H), 3.58 - 3.54 (m, 2 H), 3.28 (s, 3 H), 3.26 - 3.13 (m, 1 H), 2.81 -  
2.70 (m, 1 H), 2.58 - 2.48 (m, 1 H), 2.47 (s, 3 H), 2.33 (s, 3 H), 2.26 - 2.17 (m, 1 H).

Minor tautomer: <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.20 (s, 1 H), 9.82 (br. s, 1 H), 9.71 (s, 1 H), 7.72 (br. s, 1 H), 7.63 (s, 1 H), 7.59 - 7.35 (m, 5 H), 6.18 (s, 1 H), 5.56 (dd, 1 H), 4.24 - 4.20 (m, 2 H), 3.60 - 3.56 (m, 2 H), 3.29 (s, 3 H), 3.26 - 3.13 (m, 1 H), 2.81 - 2.70 (m, 1 H), 2.58 - 2.48 (m, 1 H), 2.48 (s, 3 H), 2.19 (s, 3 H), 2.19 - 2.11 (m, 1 H).

5

Example 747(7): (6S)-6-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

10 LC/MS *t<sub>R</sub>* 2.54 minutes; MS (ES<sup>+</sup>) *m/z* 453 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 (s, 1 H), 8.25 - 8.15 (br. s, 1 H), 7.80 - 7.70 (br. s, 1 H), 7.65 (s, 1 H), 7.55 (s, 2 H), 7.30 - 7.20 (br. s, 1 H), 6.58 (d, 1 H), 6.32 (s, 1 H), 5.70 (m, 1 H), 3.35 (m, 1 H), 2.92 - 2.85 (m, 1 H), 2.70 - 2.60 (m, 1 H), 2.52 (s, 3 H), 2.43 - 2.30 (br. s, 1 H).

15

Example 747(8): formic acid - (6S)-6-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 551 was used.

20

LC/MS *t<sub>R</sub>* 2.51 minutes; MS (ES<sup>+</sup>) *m/z* 467 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 (s, 1 H), 8.25 (s, 1 H), 8.20 (s, 1 H), 7.87 (d, 1 H), 7.67 (s, 1 H), 7.56 (m, 2 H), 7.28 (s, 1 H), 6.70 (d, 1 H), 6.34 (s, 1 H), 5.72 (dd, 1 H), 3.30 - 3.23 (m, 1 H), 3.01 - 2.85 (m, 1 H), 2.93 (s, 3 H), 2.67 (dt, 1 H), 2.53 (s, 3 H), 2.38 (td, 1 H).

25

Example 747(9): (6S)-6-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

In the step corresponding to Example 51 in the process, the compound prepared in Example 209 was used.

30

LC/MS *t<sub>R</sub>* 2.20 minutes; MS (ES<sup>+</sup>) *m/z* 487 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.41 (s, 1 H), 7.91 (br. s, 1 H), 7.68 (s, 1 H), 7.57 (app. s, 2 H), 7.39 (br. s, 1 H), 6.57 (d, 1 H), 6.36 (s, 1 H), 5.76 (ddd, 1 H), 3.36 - 3.26 (obs. m, 1 H), 2.91 (ddd, 1 H), 2.67 (qd, 1 H), 2.54 (s, 3 H), 2.48 - 2.36 (m, 1 H).

5 Example 747(10): methyl [6-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 712 was used.

10 LC/MS *t*<sub>R</sub> 2.94 minutes; MS (ES<sup>+</sup>) *m/z* 511 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 (s, 1 H), 8.56 (br. s, 1 H), 7.94 (d, 1 H), 7.76 (br. s, 1 H), 7.66 (s, 1 H), 7.55 (app. s, 2 H), 7.51 (br. s, 1 H), 6.34 (s, 1 H), 5.75 (dd, 1 H), 3.76 (s, 3 H), 3.36 - 3.25 (obs. m, 1 H), 2.90 (ddd, 1 H), 2.66 (qd, 1 H), 2.52 (s, 3 H), 2.47 - 2.37 (m, 1 H).

15

Example 747(11): 2-methoxyethyl [6-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 713 was used.

20 LC/MS *t*<sub>R</sub> 2.88 minutes; MS (ES<sup>+</sup>) *m/z* 555 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 (s, 1 H), 8.57 (br. s, 1 H), 7.94 (d, 1 H), 7.87 - 7.31 (m, 5 H), 6.35 (s, 1 H), 5.75 (d, 1 H), 4.60 (br. s, 1 H), 4.34 - 4.23 (m, 2 H), 3.71 - 3.61 (m, 2 H), 3.43 - 3.37 (m, 3 H), 3.28 - 3.36 (m, 1 H), 2.91 (ddd, 1 H), 2.72 - 2.61 (m, 1 H), 2.53 (s, 3 H), 2.35 - 2.47 (m, 1 H).

25

Example 748: methyl [5-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

30 The compound prepared in Example 747(7) (212 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (181 mg).

LC/MS *t*<sub>R</sub> 2.87 minutes; MS (ES<sup>+</sup>) *m/z* 511 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 (s, 1 H), 8.55 (s, 1 H), 8.02 (d, 1 H), 7.89 (d, 1 H), 7.67 (s, 1 H), 7.56 (s, 2 H), 7.40 (br. s, 1 H), 6.34 (s, 1 H), 5.73 (dd, 1 H), 3.77 (s, 3 H), 3.25 - 3.38 (m, 1 H), 2.91 (ddd, 1 H), 2.67 (qd, 1 H), 2.53 (s, 3 H), 2.35 - 2.46 (m, 1 H).

5

Example 749: 2-methoxyethyl [5-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 747(7) (212 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (235 mg).

LC/MS *t*<sub>R</sub> 2.98 minutes; MS (ES<sup>+</sup>) *m/z* 555 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.40 (s, 1 H), 8.56 (br. s, 1 H), 8.03 (d, 1 H), 7.88 (d, 1 H), 7.67 (s, 1 H), 7.56 (s, 2 H), 7.41 (br. s, 1 H), 6.34 (s, 1 H), 5.73 (dd, 1 H), 4.62 (br. s, 1 H), 4.25 - 4.35 (m, 2 H), 3.61 - 3.70 (m, 2 H), 3.39 (s, 3 H), 3.32 - 3.35 (m, 1 H), 2.91 (ddd, 1 H), 2.67 (qd, 1 H), 2.53 (s, 3 H), 2.35 - 2.46 (m, 1 H).

15

Example 750(1) to Example 750(8)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 747(1), 747(2), 747(3), 747(4), 747(8), 747(11), 748 and 749 using the method as detailed in Example 338.

20

Example 750(1): methyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

25

LC/MS *t*<sub>R</sub> 3.82 minutes; MS (ES<sup>+</sup>) *m/z* 544 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.90 (br. s, 1 H), 9.80 (s, 1 H), 9.71 (s, 1 H), 7.72 (s, 1 H), 7.60 (m, 3 H), 7.52 (m, 3 H), 5.75 (m, 1 H), 3.75 (s, 3 H), 3.20 - 3.05 (m, 1 H), 2.8 - 2.7 (m, 1 H), 2.62 - 2.50 (obs. m, 2 H), 2.45 (s, 3 H).

30

Example 750(2): 2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.82 minutes; MS ( $ES^+$ )  $m/z$  588 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.91 (br. s, 1 H), 9.91 (br. s, 1 H), 9.71 (s, 1 H), 7.73 (s, 1 H), 7.66 - 7.58 (m, 3 H), 7.58 - 7.52 (m, 3 H), 6.21 (s, 1 H), 5.57 (dd, 1 H), 4.22 (dd, 2 H), 3.58 (dd, 2 H), 3.29 (s, 3 H), 3.18 - 3.06 (m, 1 H), 2.77 (ddd, 1 H), 2.63 - 2.53 (m, 1 H), 2.48 (s, 3 H), 2.21 - 2.12 (m, 1 H).

Example 750(3): 2-ethoxyethyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

10 LC/MS  $t_R$  4.00 minutes; MS ( $ES^+$ )  $m/z$  602 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.40 (s, 1 H), 7.68 (s, 1 H), 7.64 (s, 1 H), 7.62 (s, 1 H), 7.57 - 7.48 (m, 3 H), 6.35 (s, 1 H), 5.66 (dd, 1 H), 4.36 - 4.17 (m, 2 H), 3.78 - 3.63 (m, 2 H), 3.57 (q, 2 H), 3.29 - 3.22 (m, 1 H), 2.90 (ddd, 1 H), 2.75 - 2.60 (m, 2 H), 2.53 (s, 3 H), 2.40 - 2.24 (m, 1 H), 1.21 (t, 3 H).

15

Example 750(4): 3-methoxypropyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  3.95 minutes; MS ( $ES^+$ )  $m/z$  602 ( $M+H$ )<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.40 (s, 1 H), 7.68 (s, 1 H), 7.62 (d, 2 H), 7.59 - 7.43 (m, 4 H), 6.35 (s, 1 H), 5.66 (dd, 1 H), 4.28 - 4.16 (m, 2 H), 3.53 (t, 2 H), 3.35 (s, 3 H), 3.30 - 3.22 (m, 1 H), 2.90 (ddd, 1 H), 2.67 (qd, 1 H), 2.53 (s, 3 H), 2.41 - 2.28 (m, 1 H), 1.95 (quintet, 2 H).

25 Example 750(5): formic acid - (6S)-6-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one (1:1)

LC/MS  $t_R$  2.75 minutes; MS ( $ES^+$ )  $m/z$  501 ( $M+H$ )<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.81 (br. s, 1 H), 9.70 (s, 1 H), 8.29 (d, 1 H), 8.16 (s, 2 H), 7.72 (s, 1 H), 7.64 (dd, 1 H), 7.62 (d, 1 H), 7.58 - 7.52 (m, 1 H), 6.72 - 6.78 (m, 1 H), 6.53 (d, 1 H), 6.20 (s, 1 H), 5.55 (dd, 1 H), 3.12 (td, 1 H), 2.83 - 2.71 (m, 4 H), 2.62 - 2.52 (m, 1 H), 2.48 (s, 3 H), 2.16 (tdd, 1 H).

Example 750(6): 2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS  $t_R$  3.76 minutes; MS ( $ES^+$ )  $m/z$  589 ( $M+H$ )<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.08 (br. s, 1 H), 10.11 (br. s, 1 H), 9.69 (s, 1 H), 8.75 (s, 1 H), 8.00 - 7.93 (m, 1 H), 7.93 - 7.87 (m, 1 H), 7.73 (s, 1 H), 7.62 (d, 1 H), 7.55 (d, 1 H), 6.19 (s, 1 H), 5.64 (dd, 1 H), 4.27 - 4.19 (m, 2 H), 3.66 - 3.51 (m, 2 H), 3.29 (s, 3 H), 3.07 (td, 1 H), 2.77 (ddd, 1 H), 2.61 - 2.52 (m, 1 H), 2.48 (s, 3 H), 2.16 - 2.07 (m, 1 H).

10

Example 750(7): methyl [5-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.63 minutes; MS ( $ES^+$ )  $m/z$  545 ( $M+H$ )<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.42 (s, 1 H), 8.60 (d, 1 H), 8.04 - 8.10 (m, 1 H), 7.98 - 8.03 (m, 1 H), 7.70 (s, 1 H), 7.55 - 7.60 (m, 2 H), 6.37 (s, 1 H), 5.69 (dd, 1 H), 3.80 (s, 3 H), 3.35 (d, 1 H), 2.92 (ddd, 1 H), 2.69 (qd, 1 H), 2.55 (s, 3 H), 2.32 - 2.41 (m, 1 H).

- 20 Example 750(8): 2-methoxyethyl [5-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  3.66 minutes; MS ( $ES^+$ )  $m/z$  589 ( $M+H$ )<sup>b</sup>

- 25 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.40 (s, 1 H), 8.58 (d, 1 H), 8.02 - 8.11 (m, 1 H), 7.92 - 8.03 (m, 1 H), 7.68 (s, 1 H), 7.50 - 7.62 (m, 2 H), 6.35 (s, 1 H), 5.67 (dd, 1 H), 4.23 - 4.41 (m, 2 H), 3.60 - 3.75 (m, 2 H), 3.40 (s, 3 H), 3.36 (br. s, 1 H), 2.90 (ddd, 1 H), 2.68 (qd, 1 H), 2.53 (s, 3 H), 2.35 (qd, 1 H).

- 30 Example 751: methyl [6-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

The compound prepared in Example 747(10) (106 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (20 mg).

LC/MS  $t_R$  3.78 minutes; MS (ES<sup>+</sup>)  $m/z$  545 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.08 (br. s, 1 H), 10.00 (br. s, 1 H), 9.69 (s, 1 H), 8.73 (s, 1 H), 7.96 (dd, 1 H), 7.90 (d, 1 H), 7.73 (s, 1 H), 7.61 (d, 1 H), 7.55 (dd, 1 H), 6.19 (s, 1 H), 5.64 (dd, 1 H), 3.70 (s, 3 H), 3.07 (td, 1 H), 2.76 (ddd, 1 H), 2.60 – 2.52 (obs. m, 1 H), 2.48 (3H, s), 2.16 – 2.07 (m, 1 H).

Example 752: (6S)-6-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

The same operation as in Example 228 → Example 338 → Example 55 was conducted from the compound prepared in Example 747(7) to give the title compound having the following physical properties.

LC/MS  $t_R$  2.70 minutes; MS (ES<sup>+</sup>)  $m/z$  487 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.40 (s, 1 H), 8.21 (d, 1 H), 7.76 (dd, 1 H), 7.67 (s, 1 H), 7.59 - 7.51 (m, 2 H), 6.65 (d, 1 H), 6.34 (s, 1 H), 5.64 (dd, 1 H), 3.30 - 3.22 (m, 1 H), 2.89 (ddd, 1 H), 2.71 - 2.60 (m, 3 H), 2.53 (s, 3 H), 2.33 (tdd, 1 H).

Example 753: ethyl (3S)-7-(2-azido-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

To a cooled (0 °C) acetonitrile (100 mL) solution of the compound prepared in Example 7 (8.95 g) was sequentially added azidotrimethylsilane (4.24 mL) and *tert*-butyl nitrite (4.84 mL) and the mixture stirred at 0 °C for 1 hour. The reaction mixture was concentrated and the residue purified by column chromatography (10% to 75% ethyl acetate in heptanes) to give the title compound having the following physical properties (7.0 g).

LC/MS  $t_R$  2.10 minutes; MS (ES<sup>+</sup>)  $m/z$  359 (M+H)<sup>a</sup>.

Example 754: ethyl (3S)-7-{5-chloro-2-[4-(tributylstannyl)-1H-1,2,3-triazol-1-yl]phenyl}-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

To a toluene (50 mL) solution of the compound prepared in Example 753 (5.0 g) was added tributyl(ethynyl)stannane (6.1 mL) and the mixture stirred at reflux for 3 hours. The reaction mixture was concentrated and the residue purified by column chromatography (5% to 75% ethyl acetate in heptanes) to give the title compound having the following physical properties (6.42 g).

LC/MS  $t_R$  2.95 minutes; MS ( $ES^+$ )  $m/z$  671, 673 and 675 ( $M+H$ )<sup>a</sup>.


Example 755: ethyl (3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylate

5 To a cooled (0 °C) THF (540 mL) solution of copper(II) chloride (1.51 g) was added a solution of the compound prepared in Example 754 (3.4 g) in THF (36 mL) dropwise over 15 minutes. The mixture was stirred at 0 °C a further 15 minutes then concentrated. The residue was dissolved in acetonitrile, washed with hexane and the acetonitrile layer concentrated. The residue was suspended in water and extracted into  
10 ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (50 to 100% ethyl acetate in heptanes) to give the title compound having the following physical properties (1.12 g).

LC/MS  $t_R$  1.92 minutes; MS ( $ES^+$ )  $m/z$  441 and 443 ( $M+Na$ ), 419 and 421 ( $M+H$ )<sup>a</sup>.

15

Example 756: (3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid



The compound prepared in Example 755 (0.90 g) was treated as detailed in  
20 Example 8 to give the title compound having the following physical properties (0.81 g).  
LC/MS  $t_R$  1.67 minutes; MS ( $ES^+$ )  $m/z$  391 and 393 ( $M+H$ )<sup>a</sup>.

Example 757(1) to Example 757(9)

The compounds of the present invention having the following physical data were  
25 prepared using the corresponding alpha-bromoketones from the compound prepared in  
Example 756 in the process of Example 51 → Example 52.

Example 757(1): 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]carbamate

5 In the step corresponding to Example 51 in the process, the compound prepared in Example 324 was used.

LC/MS  $t_R$  3.38 minutes; MS ( $ES^+$ )  $m/z$  606 and 608 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.30 (s, 1 H), 7.73 – 7.69 (m, 2 H), 7.65 (d, 1 H), 7.59 (app. br. s, 2 H), 7.45 (app. br. s, 2 H), 7.35 – 7.12 (m, 1 H), 6.16 (s, 1 H), 6.15 (s, 1 H), 5.81 (d, 1 H), 4.31 – 4.26 (m, 2 H), 3.69 – 3.63 (m, 2 H), 3.52 – 3.45 (m, 1 H), 3.41  
10 (s, 3 H), 3.14 (ddd, 1 H), 2.66 (qd, 1 H), 2.51 (app. br. s, 1 H).

Example 757(2): (3S)-3-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone

15 In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

LC/MS  $t_R$  2.92 minutes; MS ( $ES^+$ )  $m/z$  505 and 507 ( $M+H$ ), 253 and 254 [ $(M+2H)/2$ ]<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.30 (s, 1 H), 8.22 (d, 1 H), 7.78 (dd, 1 H), 7.74 – 7.68 (m, 2 H), 7.65 (d, 1 H), 7.22 (br. s, 1 H), 6.64 (d, 1 H), 6.16 (s, 1 H), 6.14 (s, 1 H), 5.80 (dd, 1 H), 3.48 (td, 1 H), 3.14 (ddd, 1 H), 2.66 (qd, 1 H), 2.53 – 2.45 (m, 1 H).  
20

Example 757(3): (3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)]phenyl]-3-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 551 was used.

25 LC/MS  $t_R$  2.98 minutes; MS ( $ES^+$ )  $m/z$  519 and 521 ( $M+H$ ), 260 and 261 [ $(M+2H)/2$ ]<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.28 (s, 1 H), 8.25 (br. s, 1 H), 7.73 (br. s, 1 H), 7.72 – 7.67 (m, 2 H), 7.63 (d, 1 H), 7.17 (br. s, 1 H), 6.54 (d, 1H), 6.14 (s, 1 H), 6.12 (s, 1 H), 5.78 (dd, 1H), 3.47 (td, 1 H), 3.13 (ddd, 1 H), 2.88 (s, 3 H), 2.64 (qd, 1 H), 2.49 (app. br. s, 1 H).  
30

Example 757(4): (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)]phenyl]-2,3-dihydro-5(1H)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 215 was used.

LC/MS  $t_R$  3.22 minutes; MS ( $ES^+$ )  $m/z$  523 and 525 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.28 (s, 1 H), 8.04 (app. br. s, 1 H), 7.72 - 7.67 (m, 2 H), 7.64 (d, 1 H), 7.18 (br. s, 1 H), 6.44 (d, 1 H), 6.14 (s, 1 H), 6.13 (s, 1 H), 5.84 - 5.76 (m, 1 H), 3.52 - 3.40 (m, 1 H), 3.12 (ddd, 1 H), 2.63 (qd, 1 H), 2.55 - 2.46 (m, 1 H).

Example 757(5): (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

10 In the step corresponding to Example 51 in the process, the compound prepared in Example 209 was used.

LC/MS  $t_R$  3.23 minutes; MS ( $ES^+$ )  $m/z$  539, 541 and 543 ( $M+H$ ), 270, 271 and 272 [ $(M+2H)/2$ ]<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.27 (s, 1 H), 7.88 (app. br. s, 1 H), 7.72 - 7.65 (m, 2 H), 7.62 (d, 1 H), 7.38 (br. s, 1 H), 6.53 (d, 1 H), 6.13 (s, 1 H), 6.11 (s, 1 H), 5.80 (d, 1 H), 3.54 - 3.39 (m, 1 H), 3.11 (dd, 1 H), 2.68 - 2.56 (m, 1 H), 2.55 - 2.47 (m, 1 H).

Example 757(6): methyl [6-(2-{(3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate

20 In the step corresponding to Example 51 in the process, the compound prepared in Example 712 was used.

LC/MS  $t_R$  3.31 minutes; MS ( $ES^+$ )  $m/z$  585 and 587 ( $M+Na$ ), 563 and 565 ( $M+H$ ), 282 and 283 [ $(M+2H)/2$ ]<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.56 (app. br. s, 1 H), 8.28 (s, 1 H), 7.94 (d, 1 H), 7.81 (app. br. s, 1 H), 7.72 - 7.67 (m, 2 H), 7.66 - 7.61 (m, 1 H), 7.54 (br. s, 1 H), 6.14 (app. s, 2 H), 5.82 (app. br. s, 1 H), 3.77 (s, 3 H), 3.49 (app. br. s, 1 H), 3.13 (ddd, 1 H), 2.64 (qd, 1 H), 2.54 (app. br. s, 1 H).

30 Example 757(7): formic acid - 2-methoxyethyl [6-(2-{(3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-pyridinyl]carbamate (1:1)



In the step corresponding to Example 51 in the process, the compound prepared in Example 713 was used. In the step corresponding to Example 52 in the process, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt.

LC/MS  $t_R$  3.27 minutes; MS ( $ES^+$ )  $m/z$  607 and 609 (M+H), 304 and 305 [(M+2H)/2]<sup>b</sup> NMR analysis showed a 2: 1 ratio of tautomers.

Major tautomer: <sup>1</sup>H-NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.21 (br. s, 1 H), 9.90 (br. s, 1 H), 8.70 (s, 1 H), 8.52 (br. s, 1 H), 8.23 (br. s, 0.5 H), 7.90 – 7.82 (m, 1 H), 7.79 – 7.71 (m, 3 H), 7.68 (d, 1 H), 7.48 (br. s, 1 H), 5.98 (s, 1 H), 5.96 (br. s, 1 H), 5.64 (d, 1 H), 4.24 – 4.19 (m, 2 H), 3.59 – 3.54 (m, 2 H), 3.42 – 3.34 (obs. m, 1 H), 3.28 (s, 3 H), 3.06 – 2.95 (m, 1 H), 2.55 – 2.50 (obs. m, 1 H), 2.37 – 2.29 (m, 1 H).

Minor tautomer: <sup>1</sup>H-NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.50 (br. s, 1 H), 10.00 (br. s, 1 H), 8.70 (s, 1 H), 8.63 (br. s, 1 H), 8.23 (br. s, 0.5 H), 7.90 – 7.82 (m, 1 H), 7.79 – 7.71 (m, 3 H), 7.68 (d, 1 H), 7.33 (br. s, 1 H), 5.98 (s, 1 H), 5.95 (br. s, 1 H), 5.71 (app. br. s, 1 H), 4.24 – 4.19 (m, 2 H), 3.59 – 3.54 (m, 2 H), 3.42 – 3.34 (obs. m, 1 H), 3.28 (s, 3 H), 3.06 – 2.95 (m, 1 H), 2.55 – 2.50 (obs. m, 1 H), 2.31 – 2.21 (app. br. s, 1 H).

Example 757(8): 6-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone

In the step corresponding to Example 51 in the process, 6-(bromoacetyl)-3,4-dihydroquinolin-2(1H)-one was used.

LC/MS  $t_R$  3.16 minutes; MS ( $ES^+$ )  $m/z$  558 and 560 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.29 (s, 1 H), 7.72 – 7.67 (m, 2 H), 7.66 – 7.60 (m, 1 H), 7.50 (br. s, 1 H), 7.46 (d, 1 H), 7.24 (br. s, 1 H), 6.86 (d, 1 H), 6.15 (s, 1 H), 6.13 (s, 1 H), 5.79 (dd, 1 H), 3.47 (td, 1 H), 3.13 (ddd, 1 H), 2.98 (t, 2 H), 2.65 (qd, 1 H), 2.58 (t, 2 H), 2.54 – 2.43 (m, 1 H).

Example 757(9): formic acid - 4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzamide (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 119 was used. In the step corresponding to Example 52 in the process, high

performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt.

LC/MS  $t_R$  3.12 minutes; MS ( $ES^+$ )  $m/z$  532 and 534 ( $M+H$ )<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.26 (br. s, 1 H), 8.69 (s, 1 H), 8.22 (br. s, 1 H), 7.93 (br. s, 1 H), 7.84 (d, 2 H), 7.79 - 7.67 (m, 5 H), 7.62 (br. s, 1 H), 7.28 (br. s, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.64 (d, 1 H), 3.42 - 3.34 (obs. m, 1 H), 3.02 (dd, 1 H), 2.58 - 2.50 (obs. m, 1 H), 2.32 (app. br. s, 1 H).

10 Example 758: methyl [5-(2-{(3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 757(2) (50 mg) was treated with methyl chloroformate following the method of Example 484 to give the title compound having  
15 the following physical properties (27 mg).

LC/MS  $t_R$  3.42 minutes; MS ( $ES^+$ )  $m/z$  563 and 565 ( $M+H$ ), 282 and 283 [ $(M+2H)/2$ ]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.56 (app. br. s, 1 H), 8.28 (s, 1 H), 8.07 - 7.97 (m, 1 H), 7.93 - 7.83 (m, 1 H), 7.73 - 7.67 (m, 2 H), 7.66 - 7.62 (m, 1 H), 7.41 (br. s, 1 H), 6.14 (s, 1 H), 6.13 (s, 1 H), 5.80 (dd, 1 H), 3.77 (s, 3 H), 3.48 (td, 1 H), 3.14 (ddd, 1 H),  
20 2.65 (qd, 1 H), 2.51 (app. br. s, 1 H).

Example 759: 2-methoxyethyl [5-(2-{(3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyll}-1H-imidazol-5-yl)-2-pyridinyl]carbamate

- 25 The compound prepared in Example 757(2) (50 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (30.7 mg).

LC/MS  $t_R$  3.47 minutes; MS ( $ES^+$ )  $m/z$  607 and 609 ( $M+H$ ), 304 and 305 [ $(M+2H)/2$ ]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.57 (app. br. s, 1 H), 8.28 (s, 1 H), 8.09 - 7.98 (m, 1 H), 7.92 - 7.82 (m, 1 H), 7.72 - 7.67 (m, 2 H), 7.66 - 7.61 (m, 1 H), 7.42 (br. s, 1 H),  
30 6.13 (app. s, 2 H), 5.80 (dd, 1 H), 4.34 - 4.27 (m, 2 H), 3.69 - 3.63 (m, 2 H), 3.49 (td, 1 H), 3.39 (s, 3 H), 3.14 (ddd, 1 H), 2.65 (qd, 1 H), 2.52 (app. br. s, 1 H).

Example 760(1) to Example 760(3)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 757(2), 757(3) and 757(6) using the method as detailed in Example 44.

5

Example 760(1): (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  3.18 minutes; MS ( $ES^+$ )  $m/z$  539, 541 and 543 ( $M+H$ ), 270, 271 and 272 [ $(M+2H)/2$ ]<sup>b</sup>

10  $^1H$  NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.28 (s, 1 H), 8.21 (d, 1 H), 7.76 (dd, 1 H), 7.72 – 7.67 (m, 2 H), 7.63 (d, 1 H), 6.65 (d, 1 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.71 (dd, 1 H), 3.47 (td, 1 H), 3.12 (ddd, 1 H), 2.65 (qd, 1 H), 2.44 - 2.36 (m, 1 H).

Example 760(2): (3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-3-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

15 LC/MS  $t_R$  3.18 minutes; MS ( $ES^+$ )  $m/z$  553, 555 and 557 ( $M+H$ ), 277, 278 and 279 [ $(M+2H)/2$ ]<sup>b</sup>

20  $^1H$  NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.28 (s, 1 H), 8.27 (d, 1 H), 7.74 (dd, 1 H), 7.72 – 7.67 (m, 2 H), 7.63 (d, 1 H), 6.59 (d, 1 H), 6.15 (s, 1 H), 6.12 (s, 1 H), 5.71 (dd, 1 H), 3.47 (td, 1 H), 3.12 (ddd, 1 H), 2.90 (s, 3 H), 2.65 (qd, 1 H), 2.44 - 2.36 (m, 1 H).

Example 760(3): methyl [6-(4-chloro-2-{(3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-3-pyridinyl]carbamate

25 LC/MS  $t_R$  4.27 minutes; MS ( $ES^+$ )  $m/z$  597, 599 and 601 ( $M+H$ ), 299, 300 and 301 [ $(M+2H)/2$ ]<sup>b</sup>


30  $^1H$  NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.64 (d, 1 H), 8.28 (s, 1 H), 8.01 - 7.92 (m, 2 H), 7.72 - 7.67 (m, 2 H), 7.63 (d, 1 H), 6.15 (app. s, 2 H), 5.81 (dd, 1 H), 3.77 (s, 3 H), 3.44 (td, 1 H), 3.12 (ddd, 1 H), 2.63 (qd, 1 H), 2.50 - 2.42 (m, 1 H).

Example 761: formic acid - 2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-3-pyridinyl]carbamate (1:1)

The compound prepared in Example 757(7) (54 mg) was treated as detailed in Example 338 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (4 mg).

LC/MS  $t_R$  4.33 minutes; MS (ES<sup>+</sup>)  $m/z$  641, 643 and 645 (M+H), 321, 322 and 323 [(M+2H)/2]<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.98 (br. s, 1 H), 10.12 (br. s, 1 H), 8.75 (app. br. s, 1 H), 8.72 (s, 1 H), 8.51 (s, 0.5 H), 7.97 (d, 1 H), 7.90 (d, 1 H), 7.81 – 7.76 (m, 2 H), 7.74 (d, 1 H), 6.01 (s, 1 H), 5.94 (s, 1 H), 5.69 (dd, 1 H), 4.27 – 4.22 (m, 2 H), 3.62 – 3.56 (m, 2 H), 3.30 (s, 3 H), 3.25 (td, 1 H), 3.00 (dd, 1 H), 2.54 – 2.45 (obs. m, 1 H), 2.25 – 2.16 (m, 1 H).

Example 762: (6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4H,6H,7H,8H-pyrrolo[1,2-a]pyrimidine-6-carboxylic acid



The same operation as in Example 753 → Example 754 → Example 755 → Example 8 was conducted from the compound prepared in Example 334 to give the title compound having the following physical properties.

LC/MS  $t_R$  1.59 minutes; MS (ES<sup>+</sup>)  $m/z$  392 and 394 (M+H)<sup>a</sup>.

Example 763(1) to Example 763(9)

The compounds of the present invention having the following physical data were prepared using the corresponding alpha-bromoketones from the compound prepared in Example 762 in the process of Example 51 → Example 52.

5    Example 763(1): methyl [4-(2-{{(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used.

10    LC/MS  $t_R$  3.24 minutes; MS (ES<sup>+</sup>)  $m/z$  563 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  8.36 (s, 1 H), 7.88 (d, 1 H), 7.72 (dd, 1 H), 7.64 (d, 1 H), 7.61 - 7.51 (m, 2 H), 7.50 - 7.37 (m, 2 H), 7.26 (br. s, 1 H), 6.16 (s, 1 H), 5.74 (dd, 1 H), 3.73 (s, 3 H), 3.39 (td, 1 H), 3.01 (ddd, 1 H), 2.69 (qd, 1 H), 2.51 - 2.35 (m, 1 H).

15    Example 763(2): 2-methoxyethyl [4-(2-{{(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 324 was used.

20    LC/MS  $t_R$  3.29 minutes; MS (ES<sup>+</sup>)  $m/z$  607 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  8.38 (s, 1 H), 7.89 (d, 1 H), 7.73 (dd, 1 H), 7.65 (d, 1 H), 7.63 - 7.53 (m, 2 H), 7.49 - 7.39 (m, 2 H), 7.29 (br. s, 1 H), 6.17 (s, 1 H), 5.75 (dd, 1 H), 4.30 - 4.21 (m, 2 H), 3.70 - 3.61 (m, 2 H), 3.48 - 3.35 (m, 4 H), 3.02 (ddd, 1 H), 2.71 (qd, 1 H), 2.51 - 2.33 (m, 1 H).

25

Example 763(3): methyl [4-(2-{{(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-4-methyl-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 78 was used.

30

LC/MS  $t_R$  3.19 minutes; MS (ES<sup>+</sup>)  $m/z$  577 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.38 (s, 1 H), 7.89 (d, 1 H), 7.73 (dd, 1 H), 7.65 (d, 1 H), 7.55 - 7.39 (m, 4 H), 6.16 (s, 1 H), 5.69 (dd, 1 H), 3.78 - 3.71 (s, 3 H), 3.47 - 3.35 (m, 1 H), 3.01 (ddd, 1 H), 2.69 (qd, 1 H), 2.46 - 2.20 (m, 4 H).

5 Example 763(4): 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl]-4-methyl-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 481 was used.

10 LC/MS *t*<sub>R</sub> 3.20 minutes; MS (ES<sup>+</sup>) *m/z* 621 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.39 (s, 1 H), 7.90 (d, 1 H), 7.74 (dd, 1 H), 7.66 (d, 1 H), 7.58 - 7.36 (m, 4 H), 6.17 (s, 1 H), 5.71 (dd, 1 H), 4.35 - 4.23 (m, 2 H), 3.72 - 3.61 (m, 2 H), 3.49 - 3.36 (m, 4 H), 3.02 (ddd, 1 H), 2.80 - 2.60 (m, 1 H), 2.52 - 2.23 (m, 4 H).

15

Example 763(5): (6S)-6-[5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

20 LC/MS *t*<sub>R</sub> 2.86 minutes; MS (ES<sup>+</sup>) *m/z* 506 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.27 (s, 1 H), 8.10 (s, 1 H), 7.79 (d, 1 H), 7.70 - 7.60 (m, 2 H), 7.55 (d, 1 H), 7.12 (br. s, 1 H), 6.52 (d, 1 H), 6.06 (s, 1 H), 5.63 (dd, 1 H), 3.29 (td, 1 H), 2.92 (ddd, 1 H), 2.65 - 2.53 (m, 1 H), 2.36 - 2.26 (m, 1 H).

25 Example 763(6): formic acid - (6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-6-{5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 551 was used. In the step corresponding to Example 52 in the process, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt.

30

LC/MS *t*<sub>R</sub> 2.81 minutes; MS (ES<sup>+</sup>) *m/z* 520 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.38 (s, 1 H), 8.22 (s, 1 H), 8.20 (s, 1 H), 7.94 - 7.87 (m, 2 H), 7.73 (dd, 1 H), 7.65 (d, 1 H), 7.31 (s, 1 H), 6.73 (d, 1 H), 6.17 (s, 1 H), 5.74 (dd, 1 H), 3.46 - 3.35 (m, 1 H), 3.03 (ddd, 1 H), 2.94 (s, 3 H), 2.77 - 2.64 (m, 1 H), 2.48 - 2.33 (m, 1 H).

5

Example 763(7): formic acid - methyl [6-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 712 was used. In the step corresponding to Example 52 in the process, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt.

LC/MS *t<sub>R</sub>* 3.16 minutes; MS (ES<sup>+</sup>) *m/z* 564 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.58 - 8.53 (m, 1 H), 8.36 (s, 1 H), 8.18 (s, 1 H), 7.96 - 7.90 (m, 1 H), 7.88 (d, 1 H), 7.78 - 7.68 (m, 2 H), 7.66 - 7.61 (m, 1 H), 7.51 (s, 1 H), 6.18 (s, 1 H), 5.77 (dd, 1 H), 3.76 (s, 3 H), 3.41 (td, 1 H), 3.01 (ddd, 1 H), 2.70 (qd, 1 H), 2.45 (tdd, 1 H).

Example 763(8): 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 713 was used.

LC/MS *t<sub>R</sub>* 3.16 minutes; MS (ES<sup>+</sup>) *m/z* 608 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.58 (app. br. s, 1 H), 8.39 (s, 1 H), 7.96 (d, 1 H), 7.90 (d, 1 H), 7.82 - 7.72 (m, 2 H), 7.67 (d, 1 H), 7.53 (br. s, 1 H), 6.20 (s, 1 H), 5.79 (dd, 1 H), 4.33 - 4.29 (m, 2 H), 3.69 - 3.65 (m, 2 H), 3.48 - 3.38 (m, 4 H), 3.04 (ddd, 1 H), 2.77 - 2.67 (m, 1 H), 2.51 - 2.42 (m, 1 H).

30

Example 763(9): formic acid - 4-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)benzamide (1:1)

In the step corresponding to Example 51 in the process, the compound prepared in Example 119 was used. In the step corresponding to Example 52 in the process, high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] was used to give the title product as the formic acid salt.

LC/MS  $t_R$  3.09 minutes; MS ( $ES^+$ )  $m/z$  533 and 535 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.37 (s, 1 H), 8.35 (br. s, 1 H), 7.92 - 7.83 (m, 3 H), 7.77 (d, 2 H), 7.72 (dd, 1 H), 7.64 (d, 1 H), 7.49 (s, 1 H), 6.17 (s, 1 H), 5.76 (dd, 1 H), 3.41 (td, 1 H), 3.03 (ddd, 1 H), 2.71 (qd, 1 H), 2.49 - 2.39 (m, 1 H).

Example 764: methyl [5-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 763(5) (175 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (162 mg).

LC/MS  $t_R$  3.33 minutes; MS ( $ES^+$ )  $m/z$  565 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.58 (br. s, 1 H), 8.39 (s, 1 H), 8.04 (d, 1 H), 7.93 - 7.88 (m, 2 H), 7.75 (dd, 1 H), 7.69 - 7.64 (m, 1 H), 7.44 (br. s, 1 H), 6.19 (s, 1 H), 5.77 (dd, 1 H), 3.79 (s, 3 H), 3.48 - 3.38 (m, 1 H), 3.05 (ddd, 1 H), 2.79 - 2.68 (m, 1 H), 2.51 - 2.41 (m, 1 H).

Example 765: 2-methoxyethyl [5-(2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 763(5) (175 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (116 mg).

LC/MS  $t_R$  3.38 minutes; MS ( $ES^+$ )  $m/z$  608 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.58 (br. s, 1 H), 8.39 (s, 1 H), 8.04 (d, 1 H), 7.94 - 7.88 (m, 2 H), 7.75 (dd, 1 H), 7.69 - 7.65 (m, 1 H), 7.44 (br. s, 1 H), 6.19 (s, 1 H), 5.77 (dd, 1 H), 4.34 - 4.31 (m, 2 H), 3.69 - 3.67 (m, 2 H), 3.48 - 3.38 (m, 4 H), 3.05 (ddd, 1 H), 2.73 (qd, 1 H), 2.46 (m, 1 H).



Example 766(1) to Example 766(7):

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 763(1), 763(2), 763(6), 763(7),

5 763(8), 764 and 765 using the method as detailed in Example 338.

Example 766(1): methyl [4-(4-chloro-2-{(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate

10 LC/MS  $t_R$  4.22 minutes; MS (ES<sup>+</sup>)  $m/z$  597 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.91 (s, 1 H), 9.81 (s, 1 H), 8.77 (s, 1 H), 7.96 (d, 1 H), 7.85 - 7.80 (m, 1 H), 7.77 - 7.74 (m, 1 H), 7.62 - 7.58 (m, 2 H), 7.57 - 7.52 (m, 2 H), 6.15 (s, 1 H), 5.58 (dd, 1 H), 3.68 (s, 3 H), 3.24 - 3.15 (m, 1 H), 2.86 (ddd, 1 H), 2.66 - 2.57 (m, 1 H), 2.23 - 2.15 (m, 1 H).

15

Example 766(2): 2-methoxyethyl [4-(4-chloro-2-{(6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl}-1H-imidazol-5-yl)phenyl]carbamate

LC/MS  $t_R$  4.22 minutes; MS (ES<sup>+</sup>)  $m/z$  641 (M+H)<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  8.39 (s, 1 H), 7.91 (d, 1 H), 7.75 (dd, 1 H), 7.69 - 7.63 (m, 3 H), 7.55 (d, 2 H), 6.20 (s, 1 H), 5.70 (dd, 1 H), 4.32 - 4.28 (m, 2 H), 3.69 - 3.66 (m, 2 H), 3.45 - 3.37 (m, 4 H), 3.03 (ddd, 1 H), 2.72 (qd, 1H), 2.39 (tdd, 1 H).

Example 766(3): (6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-6-{4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

25 LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  554 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  8.36 (s, 1 H), 8.26 (d, 1 H), 7.89 (d, 1 H), 7.77 - 7.69 (m, 2 H), 7.65 (d, 1 H), 6.59 (d, 1 H), 6.17 (s, 1 H), 5.66 (dd, 1 H), 3.37 (qd, 1 H),  
30 3.00 (ddd, 1 H), 2.89 (s, 3 H), 2.69 (qd, 1 H), 2.37 (tdd, 1 H).

Example 766(4): formic acid - methyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate (1:1)

Purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] gave the title product as the formic acid salt.

LC/MS  $t_R$  4.19 minutes; MS ( $ES^+$ )  $m/z$  598 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.69 (s, 1 H), 8.39 (s, 1 H), 8.37 (s, 1 H), 8.02 - 7.94 (m, 2 H), 7.90 (d, 1 H), 7.77 - 7.71 (m, 1 H), 7.66 (d, 1 H), 6.19 (s, 1 H), 5.75 (dd, 1 H), 3.78 (s, 3 H), 3.41 - 3.34 (m, 1 H), 3.01 (ddd, 1 H), 2.75 - 2.65 (m, 1 H), 2.44 - 2.34 (m, 1 H).

Example 766(5): 2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate

LC/MS  $t_R$  4.17 minutes; MS ( $ES^+$ )  $m/z$  642 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.11 (br. s, 1 H), 10.12 (br. s, 1 H), 8.77 (s, 1 H), 8.76 (d, 1 H), 7.99 - 7.96 (m, 2 H), 7.93 - 7.90 (m, 1 H), 7.85 - 7.81 (m, 1 H), 7.78 - 7.74 (m, 1 H), 6.13 (s, 1 H), 5.67 (dd, 1 H), 4.26 - 4.23 (m, 2 H), 3.60 - 3.58 (m, 2 H), 3.30 (s, 3 H), 3.21 - 3.13 (m, 1 H), 2.90 - 2.82 (m, 1 H), 2.61 - 2.55 (m, 1 H), 2.19 - 2.12 (m, 1 H).

Example 766(6): methyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  4.05 minutes; MS ( $ES^+$ )  $m/z$  598 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.11 (br. s, 1 H), 10.38 (s, 1 H), 8.77 (s, 1 H), 8.57 (s, 1 H), 8.08 - 8.01 (m, 1 H), 7.98 - 7.89 (m, 2 H), 7.86 - 7.79 (m, 1 H), 7.78 - 7.71 (m, 1 H), 6.15 (s, 1 H), 5.59 (dd, 1 H), 3.69 (s, 3 H), 3.21 (td, 1 H), 2.87 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.30 - 2.12 (m, 1 H).

Example 766(7): 2-methoxyethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  4.07 minutes; MS ( $ES^+$ )  $m/z$  642 (M+H)<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.12 (br. s, 1 H), 10.39 (s, 1 H), 8.80 - 8.76 (s, 1 H), 8.60 - 8.54 (m, 1 H), 8.07 - 8.01 (m, 1 H), 7.98 - 7.89 (m, 2 H), 7.86 - 7.80 (m, 1 H), 7.78 - 7.72 (m, 1 H), 6.18 - 6.12 (s, 1 H), 5.59 (dd, 1 H), 4.30 - 4.17 (m, 2 H), 3.62 - 3.53 (m, 2 H), 3.30 - 3.27 (s, 3 H), 3.25 - 3.14 (m, 1 H), 2.87 (ddd, 1 H), 2.61 (qd, 1 H), 2.27 - 2.13 (m, 1 H).

10

Example 767: (6S)-6-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one


- 15 The compound prepared in Example 763(5) (160 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (60 mg).

LC/MS  $t_R$  3.11 minutes; MS ( $ES^+$ )  $m/z$  541 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.39 (s, 1 H), 8.23 (d, 1 H), 7.91 (d, 1 H), 7.82 - 7.73 (m, 2 H), 7.67 (d, 1 H), 6.67 (dd, 1 H), 6.20 (s, 1 H), 5.69 (dd, 1 H), 3.45 - 3.36 (m, 1 H), 3.03 (ddd, 1 H), 2.75 - 2.67 (m, 1 H), 2.44 - 2.35 (m, 1 H).

20

Example 768: (3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydroindolizine-3-carboxylic acid



- 25 The same operation as in Example 7 → Example 753 → Example 754 → Example 755 → Example 8 was conducted from the compound prepared in Example 6 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 7 in the operation, 2-amino-5-methylphenylboronic acid

pinacol ester and 1,1'-bis(diphenylphosphino)ferrocene palladium (II) dichloride dichloromethane complex were used).

LC/MS  $t_R$  1.63 minutes; MS (ES<sup>+</sup>)  $m/z$  371 and 373 (M+H)<sup>a</sup>.

5 Example 769(1) to Example 769(5)

The compounds of the present invention having the following physical data were prepared using the corresponding alpha-bromoketones from the compound prepared in Example 768 in the process of Example 51 → Example 52.

10 Example 769(1): methyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, methyl [4-(bromoacetyl)-phenyl]carbamate [J. Am. Chem. Soc. 119(10), 2453 (1997)] was used.

15 LC/MS  $t_R$  3.19 minutes; MS (ES<sup>+</sup>)  $m/z$  542 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.23 (s, 1 H), 7.57 (d, 2 H), 7.49 - 7.47 (m, 2 H), 7.47 - 7.42 (m, 3 H), 7.23 (s, 1 H), 6.13 - 6.10 (m, 2 H), 5.79 (dd, 1 H), 3.74 (s, 3 H), 3.51 - 3.42 (m, 1 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.52 - 2.47 (m, 4 H).

20 Example 769(2): 2-methoxyethyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 324 was used.

25 LC/MS  $t_R$  3.24 minutes; MS (ES<sup>+</sup>)  $m/z$  586 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 8.23 (s, 1 H), 7.57 (br. s, 2 H), 7.49 - 7.38 (m, 5 H), 7.33 - 7.13 (m, 1 H), 6.11 (s, 2 H), 5.79 (dd, 1 H), 4.29 - 4.24 (m, 2 H), 3.67 - 3.62 (m, 2 H), 3.51 - 3.41 (m, 1 H), 3.39 (s, 3 H), 3.11 (ddd, 1 H), 2.68 - 2.58 (m, 1 H), 2.55 - 2.42 (m, 4 H).

30

Example 769(3): methyl [4-(2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-methyl-1H-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 78 was used.

LC/MS *t*R 3.18 minutes; MS (ES+) *m/z* 556 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.24 (s, 1 H), 7.51 - 7.44 (m, 5 H), 7.44 - 7.40 (m, 2 H), 6.12 (s, 1 H), 6.11 (s, 1 H), 5.74 (dd, 1 H), 3.74 (s, 3 H), 3.51 - 3.42 (m, 1 H), 3.10 (ddd, 1 H), 2.67 - 2.58 (m, 1 H), 2.50 (s, 3 H), 2.49 - 2.42 (m, 1 H), 2.31 (s, 3 H).

Example 769(4): 2-methoxyethyl [4-(2-((3*S*)-7-[2-(4-chloro-1*H*-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4-methyl-1*H*-imidazol-5-yl)phenyl]carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 481 was used.

LC/MS *t*R 3.19 minutes; MS (ES+) *m/z* 600 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.23 (s, 1 H), 7.52 - 7.38 (m, 7 H), 6.14 - 6.08 (m, 2 H), 5.77 - 5.71 (m, 1 H), 4.29 - 4.25 (m, 2 H), 3.67 - 3.62 (m, 2 H), 3.52 - 3.42 (m, 1 H), 3.39 (s, 3 H), 3.10 (ddd, 1 H), 2.65 - 2.57 (m, 1 H), 2.50 (s, 3 H), 2.49 - 2.41 (m, 1 H), 2.31 (br. s, 3 H).

Example 769(5): (3*S*)-3-[5-(6-amino-3-pyridinyl)-1*H*-imidazol-2-yl]-7-[2-(4-chloro-1*H*-1,2,3-triazol-1-yl)-5-methylphenyl]-2,3-dihydro-5(1*H*)-indolizinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

LC/MS *t*R 2.87 minutes; MS (ES+) *m/z* 485 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.23 (s, 1 H), 8.21 (br. s, 1 H), 7.75 (d, 1 H), 7.50 - 7.48 (m, 2 H), 7.46 (s, 1 H), 7.20 (br. s, 1 H), 6.61 (d, 1 H), 6.12 (s, 1 H), 6.11 (s, 1 H), 5.78 (dd, 1 H), 3.52 - 3.41 (m, 1 H), 3.18 - 3.08 (m, 1 H), 2.69 - 2.59 (m, 1 H), 2.50 (s, 3 H), 2.55 - 2.45 (m, 1 H).

Example 770: methyl [5-(2-((3*S*)-7-[2-(4-chloro-1*H*-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1*H*-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 769(5) (140 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (84 mg).

LC/MS *t*R 3.32 minutes; MS (ES+) *m/z* 543 (M+H), 272 (M/2+H) <sup>b</sup>

5 <sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.55 (br. s, 1 H), 8.22 (s, 1 H), 8.01 (d, 1 H), 7.87 (d, 1 H), 7.46 - 7.49 (m, 2 H), 7.45 (br. s, 1 H), 7.39 (br. s, 1 H), 6.11 (s, 2 H), 5.79 (dd, 1 H), 3.77 (s, 3 H), 3.41 - 3.53 (m, 1 H), 3.12 (ddd, 1 H), 2.64 (qd, 1 H), 2.46 - 2.55 (m, 4 H).

10 Example 771: 2-methoxyethyl [5-(2-((3*S*)-7-[2-(4-chloro-1*H*-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1*H*-imidazol-5-yl)-2-pyridinyl]carbamate

The compound prepared in Example 769(5) (140 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 128 to give the title  
15 compound having the following physical properties (98 mg).

LC/MS *t*R 3.32 minutes; MS (ES+) *m/z* 587 (M+H), 294 (M/2+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.57 (br. s, 1 H), 8.24 (s, 1 H), 8.03 (br. s, 1 H), 7.88 (d, 1 H), 7.51 - 7.48 (m, 2 H), 7.46 (s, 1 H), 7.42 (br. s, 1 H), 6.15 - 6.10 (m, 2 H), 5.80 (dd, 1 H), 4.34 - 4.29 (m, 2 H), 3.69 - 3.65 (m, 2 H), 3.53 - 3.44 (m, 1 H), 3.40 (s, 3  
20 H), 3.13 (ddd, 1 H), 2.70 - 2.60 (m, 1 H), 2.51 (s, 3 H), 2.57 - 2.47 (m, 1 H).

Example 772(1) to Example 772(2)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 769(1) and 769(2) using the  
25 method as detailed in Example 338.

Example 772(1): methyl [4-(4-chloro-2-((3*S*)-7-[2-(4-chloro-1*H*-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1*H*-imidazol-5-yl)phenyl]carbamate

30 LC/MS *t*R 4.29 minutes; MS (ES+) *m/z* 576 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 12.85 (br. s, 1 H), 9.80 (s, 1 H), 8.68 (s, 1 H), 7.63 - 7.59 (m, 2 H), 7.57 - 7.52 (m, 3 H), 7.52 - 7.46 (m, 2 H), 5.98 (s, 1 H), 5.86 (s, 1 H),

5.57 (dd, 1 H), 3.68 (s, 3 H), 3.31 - 3.25 (m, 1 H), 3.04 - 2.96 (m, 1 H), 2.57 - 2.53 (m, 1 H), 2.45 (s, 3 H), 2.23 - 2.16 (m, 1 H).

Example 772(2): 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

LC/MS *t*R 4.31 minutes; MS (ES+) *m/z* 620 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.23 (s, 1 H), 7.63 (d, 2 H), 7.53 (d, 2 H), 7.50 - 7.48 (m, 2 H), 7.47 (s, 1 H), 6.13 (s, 1 H), 6.11 (s, 1 H), 5.73 (dd, 1 H), 4.30 - 4.27 (m, 2 H), 3.67 - 3.64 (m, 2 H), 3.51 - 3.42 (m, 1 H), 3.40 (s, 3 H), 3.12 (ddd, 1 H), 2.69 - 2.60 (m, 1 H), 2.51 (s, 3 H), 2.44 - 2.37 (m, 1 H).

Example 773(1) to Example 773(3)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 769(5), 770 and 771 using the method as detailed in Example 44

Example 773(1): (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS *t*R 3.05 minutes; MS (ES+) *m/z* 519 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.23 (s, 1 H), 8.22 (d, 1 H), 7.77 (dd, 1 H), 7.50 - 7.48 (m, 2 H), 7.47 (s, 1 H), 6.65 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.70 (dd, 1 H), 3.51 - 3.42 (m, 1 H), 3.11 (ddd, 1 H), 2.69 - 2.59 (m, 1 H), 2.51 (s, 3 H), 2.43 - 2.36 (m, 1 H).

Example 773(2): methyl [5-(4-chloro-2-((3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS *t*R 4.14 minutes; MS (ES+) *m/z* 577 (M+H)<sup>b</sup>


<sup>1</sup>H NMR (500 MHz, methanol-*d*<sub>4</sub>) δ 8.58 (d, 1 H), 8.23 (s, 1 H), 8.05 - 8.02 (m, 1 H), 8.00 - 7.96 (m, 1 H), 7.49 - 7.47 (m, 2 H), 7.46 (s, 1 H), 6.13 (s, 1 H), 6.11 (s, 1 H), 5.73 (dd, 1 H), 3.78 (s, 3 H), 3.52 - 3.43 (m, 1 H), 3.12 (ddd, 1 H), 2.70 - 2.61 (m, 1 H), 2.50 (s, 3 H), 2.44 - 2.37 (m, 1 H).

Example 773(3): 2-methoxyethyl [5-(4-chloro-2-{(3S)-7-[2-(4-chloro-1H-1,2,3-triazol-1-yl)-5-methylphenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate

- 5 LC/MS  $t_R$  4.15 minutes; MS (ES<sup>+</sup>)  $m/z$  621 (M+H), 311 (M/2+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  8.58 (d, 1 H), 8.23 (s, 1 H), 8.07 - 8.03 (m, 1 H), 8.00 - 7.97 (m, 1 H), 7.50 - 7.48 (m, 2 H), 7.47 (s, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.72 (dd, 1 H), 4.34 - 4.30 (m, 2 H), 3.69 - 3.65 (m, 2 H), 3.52 - 3.43 (m, 1 H), 3.40 (s, 3 H), 3.12 (ddd, 1 H), 2.70 - 2.61 (m, 1 H), 2.51 (s, 3 H), 2.45 - 2.37 (m, 1 H).

10

Example 774: 2-{4-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1,2-dihydropyridin-1-yl}acetic acid



- The same operation as in Example 6 → Example 7 → Example 8 → Example 9  
15 was conducted from methyl 2-(4-hydroxy-6-methyl-2-oxo-1,2-dihydropyridin-1-yl)acetate [J. Het. Chem., 27(5), 1401 (1990)] to give the title compound having the following physical properties.

LC/MS  $t_R$  1.53 minutes; MS (ES<sup>+</sup>)  $m/z$  713 (2M+Na), 691 (2M+H), 368 (M+Na), 346 (M+H), 318 (M-N<sub>2</sub>+H)<sup>a</sup>.

20

Example 775: 1-{[5-(4-aminophenyl)-1H-imidazol-2-yl]methyl}-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

- The same operation as in Example 51 → Example 52 → Example 74 was  
conducted the compound prepared in Example 774 to give the title compound having  
25 the following physical properties. (Note: in the step corresponding to Example 51 in the process, 2-bromo-1-(4-nitrophenyl)ethan-1-one was used).

LC/MS  $t_R$  2.87 minutes; MS (ES<sup>+</sup>)  $m/z$  459 (M+H)<sup>b</sup>



<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.75 (m, 2 H), 7.70 (d, 1 H), 7.45 – 7.35 (br. s, 2 H), 7.20 – 7.00 (br. s, 1 H), 6.72 (d, 2 H), 6.25 (s, 1 H), 5.95 (s, 1 H), 5.32 (s, 2 H), 2.67 (s, 2 H), 2.47 (s, 3 H).

5 Example 776: 2-methoxyethyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The compound prepared in Example 775 (232 mg) was treated with 2-methoxyethyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (70 mg).

10 LC/MS *t<sub>R</sub>* 3.22 minutes; MS (ES<sup>+</sup>) *m/z* 561 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.75 – 7.65 (m, 3 H), 7.50 – 7.40 (br. d, 2 H), 7.30 – 7.20 (br. d, 2 H), 7.17 – 7.05 (br. s, 1 H), 6.27 (s, 1 H), 5.95 (s, 1 H), 5.35 (s, 2 H), 4.25 (m, 2 H), 3.65 (m, 2 H), 3.4 (s, 3 H), 2.45 (s, 3 H).

15 Example 777(1) to Example 777(8)

The compounds of the present invention having the following physical data were prepared using the corresponding alpha-bromoketones from the compound prepared in Example 774 in the process of Example 51 → Example 52

20 Example 777(1): methyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-4-methyl-1H-imidazol-5-yl]phenyl}carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 78 was used.

LC/MS *t<sub>R</sub>* 3.06 minutes; MS (ES<sup>+</sup>) *m/z* 531 (M+H)<sup>b</sup>

25 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.80 (s, 1 H), 9.70 (m, 1 H), 9.60 (br. s, 1 H), 7.82 (m, 3 H), 7.50 – 7.35 (m, 4 H), 6.05 (s, 1 H), 5.95 (s, 1 H), 5.10 (s, 2 H), 3.15 (s, 3 H), 2.55 (s, 3 H), 2.30 (s, 3 H).

30 Example 777(2): 2-methoxyethyl {4-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-4-methyl-1H-imidazol-5-yl]phenyl}carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 481 was used.

LC/MS *t<sub>R</sub>* 3.09 minutes; MS (ES<sup>+</sup>) *m/z* 575 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.76 (s, 1 H), 9.72 (m, 2 H), 7.80 (m, 3 H), 7.60 – 7.40 (m, 4 H), 6.27 (s, 1 H), 5.95 (s, 1 H), 5.10 (s, 2 H), 4.24 (m, 2 H), 3.60 (m, 2 H), 3.30 (s, 3 H), 2.5 (s, 3 H), 2.30 (s, 3 H).

5 Example 777(3): 1-([5-(6-amino-3-pyridinyl)-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used.

LC/MS *t<sub>R</sub>* 2.68 minutes; MS (ES<sup>+</sup>) *m/z* 460 (M+H)<sup>b</sup>

10 <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 11.89 (s, 1 H), 9.68 (s, 1 H), 8.24 (s, 1 H), 8.07 – 7.72 (m, 3 H), 7.65 (d, 1 H), 7.26 (s, 1 H), 6.41 (d, 1 H), 6.02 (s, 2 H), 5.96 – 5.70 (m, 2 H), 5.13 (s, 2 H), 3.32 (d, 3 H).

15 Example 777(4): 4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-1-([5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl]methyl)-2(1H)-pyridinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 551 was used.

LC/MS *t<sub>R</sub>* 2.73 minutes; MS (ES<sup>+</sup>) *m/z* 474 (M+H)<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 8.32 – 8.22 (br. s., 1 H), 7.80 – 7.73 (m, 3 H), 7.70 (d, 1 H), 7.3 – 7.2 (br. s, 1 H), 6.55 (d, 1 H), 6.29 (s, 1 H), 6.00 (s, 1 H), 5.34 (s, 2 H), 2.89 (s, 3 H), 2.50 (s, 3 H).

25 Example 777(5): 1-([5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 215 was used.

LC/MS *t<sub>R</sub>* 3.04 minutes; MS (ES<sup>+</sup>) *m/z* 478 (M+H), 500 (M+Na)<sup>b</sup>

30 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 11.95 (s, 1 H), 9.70 (s, 1 H), 7.95 (dd, 1 H), 7.80 (m, 3 H), 7.05 (s, 1 H), 6.35 (d, 1 H), 6.27 (s, 2 H), 6.05 (s, 1 H), 5.70 (s, 1 H), 5.16 (s, 2 H), 3.30 (s, 3 H).

Example 777(6): 1-([5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

In the step corresponding to Example 51 in the process, the compound prepared in Example 209 was used.

LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  494 (M+H), 516 (M+Na)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.05 (s, 1 H), 9.70 (s, 1 H), 7.95 (d, 1 H), 7.80 (m, 3 H), 7.40 (s, 1 H), 6.50 (d, 1 H), 6.33 (s, 2 H), 6.08 (s, 1 H), 5.90 (s, 1 H), 5.17 (s, 2 H), 3.30 (s, 3 H).

Example 777(7): methyl {6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-pyridinyl}carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 712 was used.

LC/MS  $t_R$  1.52 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>a</sup>

<sup>1</sup>H NMR (250 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.56 (s, 1 H), 7.95 (d, 1 H), 7.73 (m, 4 H), 7.50 (s, 1 H), 6.26 (s, 1 H), 5.97 (s, 1 H), 5.49 (s, 1 H), 5.34 (s, 2 H), 3.76 (s, 3 H), 2.49 (s, 3 H).

Example 777(8): 2-methoxyethyl {6-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-pyridinyl}carbamate

In the step corresponding to Example 51 in the process, the compound prepared in Example 713 was used.

LC/MS  $t_R$  3.04 minutes; MS (ES<sup>+</sup>)  $m/z$  562 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 8.56 (s, 1 H), 8.03 – 7.86 (m, 1 H), 7.87 – 7.60 (m, 4 H), 7.51 (s, 1 H), 6.27 (s, 1 H), 5.98 (s, 1 H), 5.35 (s, 2 H), 4.37 – 4.16 (m, 2 H), 3.82 – 3.62 (m, 2 H), 3.39 (s, 3 H), 2.49 (s, 3 H).

Example 778: methyl {5-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-2-pyridinyl}carbamate

The compound prepared in Example 777(3) (30 mg) was treated with methyl chloroformate following the method of Example 128 to give the title compound having the following physical properties (7 mg).

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 12.10 (s, 1 H), 10.13 (s, 1 H), 9.68 (s, 1 H), 8.58 (s, 1 H), 8.01 (d, 1 H), 7.78 (d, 4 H), 7.52 (s, 1 H), 6.04 (s, 1 H), 5.91 (s, 1 H), 5.16 (s, 2 H), 3.65 (s, 3 H), 2.50 (obs. s, 3 H).

5 Example 779: 2-methoxyethyl {5-[2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-2-pyridinyl} carbamate

The same operation as in Example 51 → Example 128 → Example 52 was conducted the compound prepared in Example 774 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the process, the compound prepared in Example 193 was used. In the step corresponding to

10 Example 128 in the process, 2-methoxyethyl chloroformate was used).

LC/MS *t<sub>R</sub>* 3.19 minutes; MS (ES<sup>+</sup>) *m/z* 562 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 10.15 (br. s, 2 H), 9.70 (s, 1 H), 8.60 (s, 1 H), 8.02 (d, 1 H), 7.79 (m, 4 H), 7.52 (br. s, 1 H), 6.08 (s, 1 H), 5.92 (s, 1 H), 5.20 (s, 2 H), 4.22 (m,

15 2 H), 3.58 (m, 2 H), 3.30 (s, 3 H), 2.50 (obs. s, 3 H).

Example 780(1) to Example 780(5)

The compounds of the present invention having the following physical data were synthesised from the compounds prepared in Examples 776, 777(4), 777(7), 777(8) and

20 779 using the method as detailed in Example 338.

Example 780(1): 2-methoxyethyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl} carbamate

25 LC/MS *t<sub>R</sub>* 4.08 minutes; MS (ES<sup>+</sup>) *m/z* 595 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 12.80 (br. s, 1 H), 9.86 (s, 1 H), 9.70 (s, 1 H), 7.80 (m, 3 H), 7.55 (dd, 4 H), 6.00 (s, 1 H), 5.90 (s, 1 H), 5.12 (s, 2 H), 4.20 (m, 2 H), 3.55 (m, 2 H), 3.30 (s, 3 H), 2.40 (s, 3 H).

30 Example 780(2): 1-({4-chloro-5-[6-(methylamino)-3-pyridinyl]-1H-imidazol-2-yl}methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

LC/MS *t<sub>R</sub>* 3.00 minutes; MS (ES<sup>+</sup>) *m/z* 508 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.71 (s, 1 H), 8.29 (s, 1 H), 7.80 (m, 3 H), 7.63 (m, 1 H), 6.71 (m, 1 H), 6.52 (d, 1 H), 6.03 (s, 1 H), 5.94 (s, 1 H), 5.15 (s, 2 H), 2.80 (s, 3 H), 2.45 (s, 3 H).

5 Example 780(3): methyl {6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-pyridinyl} carbamate

LC/MS *t<sub>R</sub>* 4.01 minutes; MS (ES<sup>+</sup>) *m/z* 552 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 12.92 (s, 1 H), 9.98 (s, 1 H), 9.70 (s, 1 H), 8.71 (s, 1 H), 7.98 – 7.91 (m, 1 H), 7.87 (d, 1 H), 7.81 (m, 3 H), 6.00 (s, 1 H), 5.93 (s, 1 H), 5.17  
10 (s, 2 H), 3.69 (s, 3 H), 2.38 (s, 3 H).

Example 780(4): 2-methoxyethyl {6-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-3-pyridinyl} carbamate

15 LC/MS *t<sub>R</sub>* 4.03 minutes; MS (ES<sup>+</sup>) *m/z* 596 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.92 (s, 1 H), 10.09 (s, 1 H), 9.70 (s, 1 H), 8.73 (d, 1 H), 7.99 – 7.91 (m, 1 H), 7.87 (d, 1 H), 7.81 (m, 3 H), 6.00 (s, 1 H), 5.93 (s, 1 H), 5.17 (s, 2 H), 4.30 – 4.16 (m, 2 H), 3.65 – 3.52 (m, 2 H), 3.28 (s, 3 H), 2.38 (s, 3 H).

20 Example 780(5): 2-methoxyethyl {5-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]-2-pyridinyl} carbamate

LC/MS *t<sub>R</sub>* 3.94 minutes; MS (ES<sup>+</sup>) *m/z* 596 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.72 (s, 1 H), 8.58 (s, 1 H), 8.03 (m, 1 H), 7.90 (m, 1  
25 H), 7.82 (m, 4 H), 6.04 (s, 1 H), 5.95 (s, 1 H), 5.18 (s, 2 H), 4.25 (m, 2 H), 3.59 (m, 2 H), 3.30 (s, 3 H), 2.48 (s, 3 H).

Example 781(1) to Example 781(2)

The compounds of the present invention having the following physical data were  
30 synthesised from the compounds prepared in Examples 777(3) and 777(5) using the method as detailed in Example 44.

Example 781(1): 1-([5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

LC/MS  $t_R$  3.05 minutes; MS ( $ES^+$ )  $m/z$  494 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO- $d_6$ )  $\delta$  12.71 (s, 1 H), 9.70 (s, 1 H), 8.19 (d, 1 H), 7.97 – 7.70 (m, 3 H), 7.62 (d, 1 H), 6.49 (d, 1 H), 6.17 (s, 2 H), 6.01 (d, 1 H), 5.93 (s, 1 H), 5.11 (s, 2 H), 2.45 (m, 3 H).

Example 781(2): 1-([5-(6-amino-2-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]methyl)-4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2(1H)-pyridinone

LC/MS  $t_R$  3.84 minutes; MS ( $ES^+$ )  $m/z$  512 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.79 – 7.64 (m, 4 H), 6.45 (m, 1 H), 6.23 (s, 1 H), 5.97 (s, 1 H), 5.28 (s, 2 H), 2.69 (s, 2 H), 2.45 (s, 3 H).

Example 782: N-carbamoyl-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxamide

To an N,N-dimethylformamide (1 mL) solution of the compound prepared in Example 347 (25 mg) was added (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (26.2 mg), cyanamide (3.1 mg) and diisopropylethylamine (24.5  $\mu$ L) and the mixture stirred at room temperature for 16 hours. To the reaction mixture, concentrated hydrochloric acid (1 mL) was added and the mixture stirred at room temperature for 24 hours then warmed to 50 °C and stirred a further 16 hours. The reaction mixture was cooled to room temperature, diluted with water (6 mL) and sufficient solid sodium hydrogen carbonate added to basify the supernatant to pH 9. The resultant suspension was extracted into a 1:1 mixture of ethyl acetate and propan-2-ol, the combined organic layers washed with saturated saline, dried and concentrated. The residue obtained on concentration was triturated with dichloromethane, the resultant precipitate being isolated by filtration to obtain the title compound having the following physical properties (5.3 mg).

LC/MS  $t_R$  3.29 minutes; MS ( $ES^+$ )  $m/z$  548 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.40 (br. s, 1 H), 10.50 (br. s, 1 H), 9.69 (s, 1 H), 8.09 (d, 1 H), 7.91 (br. s, 1 H), 7.84 – 7.75 (m, 3 H), 7.60 (s, 1 H), 7.33 (br. s, 1 H), 7.28 (d, 1 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.60 (d, 1 H), 3.36 – 3.26 (obs. m, 1 H), 3.00 (dd, 1 H), 2.57 – 2.49 (obs. m, 1 H), 2.39 – 2.27 (m, 1 H).

Example 783: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[5-(methylsulfonyl)-2-thienyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 234 → Example 51 → Example 52 was  
5 conducted from 5-methanesulfonylthiophene-2-carboxylic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  3.61 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.21 (br. s, 1 H), 8.62 (s, 1 H), 7.62 (dd, 1 H), 7.59 (d, 1 H), 7.55 (d, 1 H), 7.52 (d, 1 H), 7.27 (obs. s, 1 H), 7.20 (d, 1 H), 6.28 (s, 1 H), 5.83 (d, 1 H), 5.81 (br. s, 1 H), 3.51 - 3.40 (m, 1 H), 3.31 - 3.22 (m, 1 H), 3.18 (s, 3 H), 3.05 (dd, 1 H), 2.59 - 2.45 (m, 1 H).

Example 784: (3S)-3-{4-chloro-5-[5-(methylsulfonyl)-2-thienyl]-1H-imidazol-2-yl}-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

15 The compound prepared in Example 783 (100 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (60 mg).

LC/MS  $t_R$  3.91 minutes; MS (ES<sup>+</sup>)  $m/z$  596 and 598 (M+Na), 574 and 576 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  12.35 (br. s, 1 H), 8.71 (s, 1 H), 7.68 - 7.60 (m, 2 H), 7.57 (d, 1 H), 7.56 - 7.52 (m, 1 H), 7.05 (app. br. s, 1 H), 6.32 (s, 1 H), 5.95 (s, 1 H), 5.84 (d, 1 H), 3.64 - 3.52 (m, 1 H), 3.22 (s, 3 H), 3.10 (dd, 1 H), 2.85 (app. br. s, 1 H), 2.65 - 2.52 (m, 1 H).

Example 785: 5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-N-(methylsulfonyl)-2-thiophenecarboxamide

The compound prepared in Example 347 (62 mg) was treated as detailed in Example 355 using methanesulfonamide in place of N-(2-hydroxyethyl)morpholine to give, on purification by high performance liquid chromatography [5 to 100% mobile  
30 phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)], the title compound having the following physical properties (11 mg).

LC/MS  $t_R$  3.47 minutes; MS (ES<sup>+</sup>)  $m/z$  583 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.36 (s, 1 H), 7.78 - 7.72 (m, 3 H), 7.71 - 7.67 (m, 1 H), 7.45 (s, 1 H), 7.29 (d, 1 H), 6.13 (s, 1 H), 6.10 (s, 1 H), 5.76 (dd, 1 H), 3.50 - 3.40 (m, 1 H), 3.33 (s, 3 H), 3.12 (ddd, 1 H), 2.69 - 2.60 (m, 1 H), 2.53 - 2.45 (m, 1 H).

5 Example 786: 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(methylsulfonyl)-2-thiophenecarboxamide

The compound prepared in Example 785 (26 mg) was treated as detailed in Example 338 to give, on purification by high performance liquid chromatography [5 to 10 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)], the title compound having the following physical properties (4.5 mg).

LC/MS *t<sub>R</sub>* 3.91 minutes; MS (ES<sup>+</sup>) *m/z* 617 and 619 (M+H)<sup>b</sup>

15 <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 9.37 (s, 1 H), 7.82 (d, 1 H), 7.78 - 7.72 (m, 2 H), 7.71 - 7.66 (m, 1 H), 7.40 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.70 (dd, 1 H), 3.49 - 3.39 (m, 1 H), 3.33 (s, 3 H), 3.12 (ddd, 1 H), 2.71 - 2.61 (m, 1 H), 2.42 - 2.34 (m, 1 H).

Example 787: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-furoic acid

20 The same operation as in Example 51 → Example 52 → Example 8 → Example 55 → Example 24 was conducted from the compound prepared in Example 662 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the process, methyl 5-(2-bromoacetyl)furan-2-carboxylate [patent US2007/265265] was used).

25 LC/MS *t<sub>R</sub>* 3.20 minutes; MS (ES<sup>+</sup>) *m/z* 490 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.02 (br. s, 1 H), 12.44 (br. s, 1 H), 9.69 (s, 1 H), 7.84 - 7.76 (m, 3 H), 7.50 (d, 1 H), 7.21 (d, 1 H), 6.63 (d, 1 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.62 (dd, 1 H), 3.31 - 3.28 (m, 1 H), 3.00 (dd, 1 H), 2.56 - 2.52 (m, 1 H), 2.33 - 2.24 (m, 1 H).

30

Example 788: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-furamide



The compound prepared in Example 787 (40 mg) was treated as detailed in Example 114 to give the title compound having the following physical properties (16 mg).

LC/MS  $t_R$  3.07 minutes; MS ( $ES^+$ )  $m/z$  489 ( $M+H$ )<sup>b</sup>

- 5 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.66 (m, 1 H), 7.53 (br. s, 1 H), 7.16 (d, 1 H), 6.70 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.48 - 3.39 (m, 1 H), 3.12 (ddd, 1 H), 2.66 (qd, 1 H), 2.51 - 2.40 (m, 1 H).

10 Example 789: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-methyl-2-furamide

The compound prepared in Example 787 (40 mg) was treated as detailed in Example 114 using methylamine hydrochloride in place of ammonium chloride to give the title compound having the following physical properties (19 mg).

LC/MS  $t_R$  3.18 minutes; MS ( $ES^+$ )  $m/z$  502 ( $M+H$ )<sup>b</sup>

- 15 <sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.38 (s, 1 H), 7.77 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.52 (br. s, 1 H), 7.10 (d, 1 H), 6.68 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.78 (dd, 1 H), 3.48 - 3.39 (m, 1 H), 3.13 (ddd, 1 H), 2.92 (s, 3 H), 2.66 (qd, 1 H), 2.52 - 2.40 (m, 1 H).

20 Example 790: 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-furoic acid

The compound prepared in Example 787 (50 mg) was treated using the method as detailed in Example 338 to give the title compound having the following physical properties (30 mg).

- 25 LC/MS  $t_R$  3.81 minutes; MS ( $ES^+$ )  $m/z$  524 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.53 (br. s, 1 H), 9.71 (s, 1 H), 7.82 (d, 1 H), 7.81 - 7.80 (m, 2 H), 7.25 (d, 1 H), 6.82 (d, 1 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.60 (dd, 1 H), 3.27 - 3.23 (m, 1 H), 2.99 (ddd, 1 H), 2.56 - 2.52 (m, 1 H), 2.17 (dd, 1 H).

30 Example 791: 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-furamide

The compound prepared in Example 790 (60 mg) was treated as detailed in Example 114 to give the title compound having the following physical properties (9 mg).

LC/MS  $t_R$  3.64 minutes; MS (ES<sup>+</sup>)  $m/z$  523 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 7.76 - 7.72 (m, 2 H), 7.71 - 7.67 (m, 1 H), 7.23 (d, 1 H), 6.84 (d, 1 H), 6.14 (s, 1 H), 6.10 (s, 1 H), 5.72 (dd, 1 H), 3.45 (td, 1 H), 3.12 (ddd, 1 H), 2.70 - 2.61 (m, 1 H), 2.42 (tdd, 1 H).

5

Example 792: formic acid - 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-N-methyl-2-furamide (1:1)

The compound prepared in Example 790 (60 mg) was treated using the method as detailed in Example 114 using methylamine hydrochloride in place of ammonium chloride, to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (12 mg).

LC/MS  $t_R$  3.76 minutes; MS (ES<sup>+</sup>)  $m/z$  537 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.37 (s, 1 H), 8.13 (br. s, 1 H), 7.77 - 7.72 (m, 2 H), 7.72 - 7.66 (m, 1 H), 7.17 (d, 1 H), 6.83 (d, 1 H), 6.14 (s, 1 H), 6.11 (s, 1 H), 5.72 (dd, 1 H), 3.45 (td, 1 H), 3.20 - 3.07 (m, 1 H), 2.94 (s, 3 H), 2.73 - 2.59 (m, 1 H), 2.43 (tdd, 1 H).

20

Example 793: tert-butyl 4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}furan-2-carboxylate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 was conducted from tert-butyl 4-bromofuran-2-carboxylate [patent EP1489077, 2004] to give the title compound having the following physical properties.

25

LC/MS  $t_R$  1.81 minutes; MS (ES<sup>+</sup>)  $m/z$  546 (M+H), 490 (M-C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>a</sup>.

Example 794: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-furoic acid hydrochloride

The compound prepared in Example 793 (100 mg) was treated as detailed in Example 363 to give the title compound as the hydrochloride salt having the following physical properties (17.7 mg).

30

LC/MS  $t_R$  2.85 minutes; MS (ES<sup>+</sup>)  $m/z$  490 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.46 (br. s, 1 H), 9.71 (s, 1 H), 8.40 (br. s, 1 H), 7.89 (br. s, 1 H), 7.86 - 7.80 (m, 2 H), 7.72 (d, 1 H), 7.68 (s, 1 H), 6.05 (s, 1 H), 6.01 (s, 1 H), 5.77 (dd, 1 H), 3.34 - 3.28 (m, 1 H), 3.12 - 3.04 (m, 1 H), 2.72 - 2.62 (m, 1 H), 2.38 - 2.29 (m, 1 H).

Example 795: 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-furoic acid - formic acid (1:1)

The same operation as in Example 338 → Example 363 was conducted from the compound prepared in Example 793 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties.

LC/MS  $t_R$  3.70 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.09 (br. s, 1 H), 9.71 (s, 1 H), 8.19 (s, 1 H), 8.18 (s, 0.5 H), 7.83 - 7.77 (m, 3 H), 7.48 (s, 1 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.54 (dd, 1 H), 3.34 - 3.23 (m, 1 H), 2.98 (dd, 1 H), 2.58 - 2.48 (m, 1 H), 2.23 - 2.14 (m, 1 H).

Example 796: 2-methyl-2-propanyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-furoate

The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 was conducted from 5-bromofuran-3-carboxylic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  4.22 minutes; MS (ES<sup>+</sup>)  $m/z$  546 (M+H), 490 (M-C(CH<sub>3</sub>)<sub>3</sub>+H), 462 (M-N<sub>2</sub>-C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, methanol- $d_4$ )  $\delta$  9.36 (s, 1 H), 7.98 (s, 1 H), 7.76 - 7.68 (m, 3 H), 7.30 (s, 1 H), 6.80 (s, 1 H), 6.12 (s, 1 H), 6.09 (s, 1 H), 5.76 (dd, 1 H), 3.51 - 3.38 (m, 1 H), 3.17 - 3.04 (m, 1 H), 2.70 - 2.57 (m, 1 H), 2.50 - 2.40 (m, 1 H), 1.55 (s, 9 H).

Example 797: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-furoic acid - formic acid (1:1)

The compound prepared in Example 796 (135 mg) was treated as detailed in Example 363 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (62.6 mg).

LC/MS  $t_R$  3.22 minutes; MS (ES<sup>+</sup>)  $m/z$  490 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.32 (br. s, 1 H), 9.69 (s, 1 H), 8.18 (app. s, 2 H), 7.83 - 7.78 (m, 3 H), 7.36 (br. s, 1 H), 6.67 (br. s, 1 H), 5.98 (s, 1 H), 5.95 (s, 1 H), 5.61 (dd, 1 H), 3.39 - 3.30 (m, 1 H), 3.00 (dd, 1 H), 2.54 - 2.44 (obs. m, 1 H), 2.32 - 2.25 (m, 1 H).

Example 798: 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-1,3-oxazole-4-carboxylic acid

The same operation as in Example 361 → Example 51 → Example 52 → Example 8 → Example 55 → Example 24 was conducted from ethyl 2-acetyl-1,3-oxazole-4-carboxylate [patent WO2010/143116] to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 11 was used).

LC/MS  $t_R$  3.16 minutes; MS (ES<sup>+</sup>)  $m/z$  491 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.73 (br. s, 1 H), 9.71 (s, 1 H), 8.56 (br. s, 1 H), 8.20 (s, 0.5 H), 7.84 - 7.72 (m, 4 H), 6.00 (s, 1 H), 5.92 (s, 1 H), 5.64 (d, 1 H), 3.40 - 3.29 (obs. m, 1 H), 3.01 (dd, 1 H), 2.58 - 2.47 (obs. m, 1 H), 2.32 - 2.22 (m, 1 H).

Example 799: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-1H-pyrrole-2-carboxylic acid

The same operation as in Example 51 → Example 52 → Example 8 → Example 55 → Example 24 was conducted from methyl 5-(2-bromoacetyl)-1H-pyrrole-2-carboxylate [Chem. Pharm. Bull., 46(4), 559 (1998)] to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the

formic acid salt having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 622 was used).

LC/MS  $t_R$  3.02 minutes; MS (ES<sup>+</sup>)  $m/z$  489 (M+H)<sup>b</sup>

5 <sup>1</sup>H NMR (250 MHz, DMSO- $d_6$ )  $\delta$  12.04 (br. s, 1 H), 11.57 (br. s, 1 H), 9.69 (s, 1 H), 8.19 (s, 0.5 H), 7.85 - 7.75 (m, 3 H), 7.50 (br. s, 1 H), 6.71 (dd, 1 H), 6.33 (br. s, 1 H), 5.96 (s, 1 H), 5.94 (s, 1 H), 5.60 (dd, 1 H), 3.47 - 3.28 (obs. m, 1 H), 3.05 - 2.91 (m, 1 H), 2.56 - 2.39 (obs. m, 1 H), 2.38 - 2.20 (m, 1 H).

10 Example 800: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1H-pyrrole-2-carboxylic acid

The same operation as in Example 51 → Example 52 → Example 8 → Example 55 → Example 24 was conducted from ethyl 4-(2-chloroacetyl)-1H-pyrrole-2-carboxylate [Chem. Pharm. Bull., 44(1), 48 (1996)] to give the title compound after purification by  
15 high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 11 was used).

LC/MS  $t_R$  2.74 minutes; MS (ES<sup>+</sup>)  $m/z$  489 (M+H)<sup>b</sup>

20 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.53 - 11.33 (m, 2 H), 9.69 (s, 1 H), 7.83 - 7.74 (m, 3 H), 7.28 - 6.77 (m, 3 H), 5.95 (s, 1 H), 5.93 (s, 1 H), 5.57 (d, 1 H), 3.42 - 3.31 (m, 1 H), 2.97 (dd, 1 H), 2.51 - 2.42 (m, 1 H), 2.26 (app. br. s, 1 H).

25 Example 801: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1H-pyrrole-3-carboxylic acid - formic acid (1:1)

The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 5-bromo-1H-pyrrole-3-carboxylic acid to give the title compound after purification by high performance  
30 liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties.

LC/MS  $t_R$  2.85 minutes; MS (ES<sup>+</sup>)  $m/z$  489 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.02 (br. s, 1 H), 11.40 (br. s, 1 H), 9.70 (s, 1 H), 8.24 (s, 0.5 H), 7.83 - 7.80 (m, 3 H), 7.31 - 7.18 (m, 2 H), 6.50 (br. s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.60 (dd, 1 H), 3.40 - 3.30 (obs. m, 1 H), 2.98 (dd, 1 H), 2.52 - 2.43 (obs. m, 1 H), 2.33 (app. br. s, 1 H).

Example 802: tert-butyl 5-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}-1-methyl-1H-pyrrole-2-carboxylate

The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 was conducted from 5-bromo-1-methyl-1H-pyrrole-2-carboxylic acid to give the title compound having the following physical properties. LC/MS  $t_R$  4.08 minutes; MS (ES<sup>+</sup>)  $m/z$  559 (M+H), 503 (M-C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>b</sup>

Example 803: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(1-methyl-1H-pyrrol-2-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 802 (135 mg) was treated as detailed in Example 363 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] having the following physical properties (42.9 mg).

LC/MS  $t_R$  3.02 minutes; MS (ES<sup>+</sup>)  $m/z$  459 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.36 (s, 1 H), 7.76 - 7.71 (m, 2 H), 7.68 (d, 1 H), 7.12 (s, 1 H), 6.73 (t, 1 H), 6.28 (dd, 1 H), 6.13 (s, 1 H), 6.10 (d, 1 H), 6.07 (dd, 1 H), 5.79 (dd, 1 H), 3.67 (s, 3 H), 3.43 - 3.36 (m, 1 H), 3.13 (ddd, 1 H), 2.72 - 2.63 (m, 1 H), 2.45 (tdd, 1 H).

Example 804: 5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-1-methyl-1H-pyrrole-2-carboxylic acid

To a cooled (0 °C) dichloromethane (12.6 mL) solution of the compound prepared in Example 802 (35 mg) was added a 1 M solution of titanium(IV) chloride in dichloromethane (0.63 mL), the mixture warmed to room temperature and stirred 1 hour.

The reaction mixture was cooled to 0 °C, 1 M hydrochloric acid (12.6 mL) added and the mixture stirred vigorously for 5 minutes before filtering through a pad of Celite®. The filter cake was sequentially washed with water (30 mL), propan-2-ol (50 mL) and dichloromethane (50 mL). The organic layer of the biphasic filtrate was isolated and the pH of the aqueous phase adjusted to 2 – 3 by addition of solid sodium hydrogen carbonate. The aqueous was extracted with a 1: 1 mixture of propan-2-ol and dichloromethane, the organic layers combined, washed with saturated saline, dried and concentrated. The residue was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (14.9 mg).

LC/MS  $t_R$  3.11 minutes; MS (ES<sup>+</sup>)  $m/z$  503 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.30 (br. s, 1 H), 9.67 (s, 1 H), 7.86 - 7.73 (m, 3 H), 7.32 (br. s, 1 H), 6.79 (d, 1 H), 6.21 (app. br. s, 1 H), 5.95 (app. s, 2 H), 5.63 (d, 1 H), 3.98 (br. s, 3 H), 3.51 - 3.17 (obs. m, 1 H), 2.99 (dd, 1 H), 2.55 - 2.44 (obs. m, 1 H), 2.35 - 2.26 (m, 1 H).

Example 805: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-[5-(3-hydroxy-1-methyl-1H-pyrazol-5-yl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 234 → Example 51 → Example 52 was conducted from 3-hydroxy-1-methyl-1H-pyrazole-5-carboxylic acid [Chem. Ber., 109, 268 (1976)] to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] having the following physical properties.

LC/MS  $t_R$  3.02 minutes; MS (ES<sup>+</sup>)  $m/z$  476 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.34 (s, 1 H), 7.76 – 7.70 (m, 2 H), 7.68 (d, 1 H), 7.26 (br. s, 1 H), 6.13 (s, 1 H), 6.08 (s, 1 H), 5.78 (dd, 1 H), 5.69 (s, 1 H), 3.75 (s, 3 H), 3.43 (td, 1 H), 3.11 (ddd, 1 H), 2.68 – 2.57 (m, 1 H), 2.52 – 2.42 (m, 1 H).

Example 806: ethyl [4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-thienyl]acetate

The same operation as in Example 139 → Example 361 → Example 51 → Example

52 was conducted from ethyl 2-(4-bromothiophen-2-yl)acetate [patent US6184245, 2001] to give the title compound having the following physical properties.

LC/MS  $t_R$  3.33 minutes; MS (ES<sup>+</sup>)  $m/z$  548 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.05 (br. s, 1 H), 8.62 (s, 1 H), 7.63 (dd, 1 H), 7.56 (d, 1 H), 7.54 (d, 1 H), 7.35 (br. s, 1 H), 7.14 (s, 1 H), 7.08 (s, 1 H), 6.31 (s, 1 H), 5.86 (d, 1 H), 5.78 (s, 1 H), 4.21 (q, 2 H), 3.83 (s, 2 H), 3.55 - 3.45 (m, 1 H), 3.35 - 3.26 (m, 1 H), 3.04 (dd, 1 H), 2.56 - 2.45 (m, 1 H), 1.30 (t, 3 H).

Example 807: [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetic acid - formic acid (1:1)

The compound prepared in Example 806 (75 mg) was treated as detailed in Example 342 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt having the following physical properties (52.7 mg).

LC/MS  $t_R$  2.90 minutes; MS (ES<sup>+</sup>)  $m/z$  520 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.61 (br. s, 1 H), 12.01 (br. s, 1 H), 9.70 (s, 1 H), 8.15 (s, 1 H), 7.84 - 7.76 (m, 3 H), 7.34 (br. s, 1 H), 7.30 (br. s, 1 H), 7.20 (br. s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.60 (d, 1 H), 3.80 (s, 2 H), 3.40 - 3.20 (obs. m, 1 H), 2.99 (dd, 1 H), 2.59 - 2.48 (obs. m, 1 H), 2.33 (app. br. s, 1 H).

Example 808: ethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetate

The compound prepared in Example 806 (153 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (92 mg).

LC/MS  $t_R$  4.44 minutes; MS (ES<sup>+</sup>)  $m/z$  604 and 606 (M+Na), 582 and 584 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  11.23 (br. s, 1 H), 8.49 (s, 1 H), 7.55 (dd, 1 H), 7.48 - 7.44 (m, 2 H), 7.35 (s, 1 H), 7.11 (s, 1 H), 6.26 (s, 1 H), 5.73 (d, 1 H), 5.67 (s, 1 H), 4.13 (q, 2 H), 3.75 (s, 2 H), 3.40 - 3.31 (m, 1 H), 3.10 (dd, 1 H), 2.93 (dd, 1 H), 2.44 - 2.33 (m, 1 H), 1.23 (t, 3 H).



Example 809: [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thienyl]acetic acid hydrochloride

The compound prepared in Example 808 (50 mg) was treated as detailed in Example 342 to give, on concentration of the reaction mixture, the title compound as the

5 hydrochloride salt having the following physical properties (48.5 mg).

LC/MS  $t_R$  3.86 minutes; MS (ES<sup>+</sup>)  $m/z$  554 and 556 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.00 (br. s, 1 H), 9.71 (s, 1 H), 7.91 - 7.74 (m, 3 H), 7.64 (s, 1 H), 7.39 (s, 1 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.55 (dd, 1 H), 3.87 (s, 2 H), 3.34 - 3.21 (m, 1 H), 2.98 (dd, 1 H), 2.58 - 2.47 (obs. m, 1 H), 2.18 (app. t, 1 H).

10

Example 810: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzenesulfonamide

The same operation as in Example 51 → Example 52 was conducted from 4-(2-bromoacetyl)benzene-1-sulfonamide [patent WO2009/009122] to give the title

15 compound having the following physical properties.

LC/MS  $t_R$  3.05 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.32 (br. s, 1 H), 9.70 (s, 1 H), 7.88 (d, 2 H), 7.84 - 7.78 (m, 3 H), 7.77 (d, 2 H), 7.70 (d, 1 H), 7.30 (br. s, 2 H), 5.99 (s, 1 H), 5.97 (s, 1 H), 5.64 (dd, 1 H), 3.42 - 3.32 (m, 1 H), 3.02 (dd, 1 H), 2.58 - 2.49 (m, 1 H), 2.38 - 2.31 (m, 1 H).

20

Example 811: 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzenesulfonamide

The compound prepared in Example 810 (70 mg) was treated as detailed in Example 338 to give the title compound having the following physical properties (31 mg).

25

LC/MS  $t_R$  3.69 minutes; MS (ES<sup>+</sup>)  $m/z$  569 and 571 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol- $d_4$ )  $\delta$  9.37 (s, 1 H), 7.96 (d, 2 H), 7.86 (d, 2 H), 7.75 - 7.71 (m, 2 H), 7.70 - 7.66 (m, 1 H), 6.13 (s, 1 H), 6.11 (s, 1 H), 5.73 (dd, 1 H), 3.50 - 3.40 (m, 1 H), 3.12 (ddd, 1 H), 2.66 (qd, 1 H), 2.43 - 2.36 (m, 1 H).

30

Example 812: 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)benzenesulfonamide

The same operation as in Example 51 → Example 52 was conducted from 3-(2-bromoacetyl)benzene-1-sulfonamide [Chem. Pharm. Bull., 30(11), 4092 (1982)] to give the title compound having the following physical properties.

LC/MS  $t_R$  3.09 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO- $d_6$ )  $\delta$  12.27 (br. s, 1 H), 9.69 (s, 1 H), 8.14 (s, 1 H), 7.88 (d, 1 H), 7.82 - 7.73 (m, 3 H), 7.65 - 7.43 (m, 3 H), 7.35 (s, 2 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.62 (dd, 1 H), 3.49 - 3.38 (m, 1 H), 3.09 - 2.92 (m, 1 H), 2.62 - 2.52 (m, 1 H), 2.43 - 2.28 (m, 1 H).

Example 813: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(5-oxo-4,5-dihydro-1,2,4-oxadiazol-3-yl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 was conducted from 3-(4-bromophenyl)-4,5-dihydro-1,2,4-oxadiazol-5-one to give the title compound having the following physical properties.

LC/MS  $t_R$  3.29 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.91 (br. s, 1 H), 12.32 (br. s, 1 H), 9.69 (s, 1 H), 7.89 (d, 2 H), 7.84 - 7.75 (m, 5 H), 7.71 (s, 1 H), 5.98 (s, 1 H), 5.96 (s, 1 H), 5.63 (dd, 1 H), 3.44 - 3.33 (obs. m, 1 H), 3.01 (dd, 1 H), 2.59 - 2.48 (obs. m, 1 H), 2.38 - 2.30 (m, 1 H).

Example 814: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[3-(5-oxo-4,5-dihydro-1,2,4-oxadiazol-3-yl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 was conducted from 3-(3-bromophenyl)-4,5-dihydro-1,2,4-oxadiazol-5-one to give the title compound having the following physical properties.

LC/MS  $t_R$  3.34 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 12.94 (br. s, 1 H), 12.25 (br. s, 1 H), 9.70 (s, 1 H), 8.14 (s, 1 H), 7.93 (d, 1 H), 7.85 - 7.74 (m, 3 H), 7.66 - 7.46 (m, 3 H), 5.99 (s, 1 H), 5.95 (s, 1 H), 5.64 (dd, 1 H), 3.53 - 3.28 (obs. m, 1 H), 3.02 (dd, 1 H), 2.59 - 2.28 (obs. m, 2 H).

5

Example 815: 2-(4-bromophenyl)-2-(trifluoromethyl)-1,3-dioxolane

A vigorously stirred solution of 1-(4-bromophenyl)-2,2,2-trifluoroethanone (2.0 g) and 2-chloroethanol (0.79 mL) in N,N-dimethylformamide (4 mL) and tetrahydrofuran (2 mL) was cooled to -60 °C and a solution of potassium *tert*-butoxide (1.33 g) in N,N-dimethylformamide (3 mL) added dropwise over 20 minutes. The reaction mixture was stirred at -60 °C for 90 minutes whereupon a saturated aqueous solution of ammonium chloride (10 mL) was added. On warming to room temperature, water (5 mL) was added followed by saturated saline (5 mL) and the mixture extracted into ethyl acetate. The combined organic layers were washed twice with water and once with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 - 40% ethyl acetate in heptanes) to give the title compound having the following physical properties.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.54 (d, 2 H), 7.49 (d, 2 H), 4.33 - 4.24 (m, 2 H), 4.13 - 4.04 (m, 2 H).

20

Example 816: (3S)-7-(2-amino-5-chlorophenyl)-3-(5-{4-[2-(trifluoromethyl)-1,3-dioxolan-2-yl]phenyl}-1H-imidazol-2-yl)-1,2,3,5-tetrahydroindolizin-5-one

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 40 was conducted from the compound prepared in Example 815 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 11 was used).

LC/MS *t*<sub>R</sub> 1.89 minutes; MS (ES<sup>+</sup>) *m/z* 543 (M+H)<sup>+</sup> <sup>a</sup>

Example 817: (3S)-7-(2-amino-5-chlorophenyl)-3-{5-[4-(trifluoroacetyl)phenyl]-1H-imidazol-2-yl}-1,2,3,5-tetrahydroindolizin-5-one

To a cooled (0 °C) dichloromethane (6.8 mL) solution of the compound prepared in

Example 816 (131 mg) was added a 1 M solution of boron tribromide in dichloromethane (1.33 mL) and the mixture stirred at 0 °C for 1 hour then warmed to room temperature and stirred 16 hours. To the cooled (0 °C) reaction mixture, a saturated aqueous solution of sodium hydrogen carbonate was added followed by extraction with dichloromethane. The combined dichloromethane layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 – 15% methanol in ethyl acetate) to give the title compound having the following physical properties (50 mg).

LC/MS  $t_R$  1.57 minutes; MS (ES<sup>+</sup>)  $m/z$  517 (M+H<sub>2</sub>O+H) <sup>a</sup>

Example 818: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(trifluoroacetyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

The compound prepared in Example 817 (67.9 mg) was treated as detailed in Example 24 to give the title compound having the following physical properties (60.1 mg).

LC/MS  $t_R$  4.31 minutes; MS (ES<sup>+</sup>)  $m/z$  570 (M+H<sub>2</sub>O+H), 552 (M+H) <sup>g</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>)  $\delta$  9.35 (s, 1 H), 7.74 – 7.66 (m, 5 H), 7.59 (d, 2 H), 7.38 (s, 1 H), 6.13 (s, 1 H), 6.09 (s, 1 H), 5.79 (dd, 1 H), 3.45 (td, 1 H), 3.11 (ddd, 1 H), 2.64 (qd, 1 H), 2.53 – 2.44 (m, 1 H).

Example 819: (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-3-{5-[4-(2,2,2-trifluoro-1-hydroxyethyl)phenyl]-1H-imidazol-2-yl}-2,3-dihydro-5(1H)-indolizinone

To a methanol (3.2 mL) solution of the compound prepared in Example 818 (32 mg) was added sodium borohydride (5.1 mg) and the mixture stirred one hour at room temperature. On concentration, the residue was suspended in a saturated aqueous solution of ammonium chloride (5 mL) and extracted into ethyl acetate. The combined organic layers were washed with saturated saline, dried and concentrated. The residue was purified by column chromatography (0 – 10% methanol in dichloromethane) to give the title compound having the following physical properties (25.3 mg).

LC/MS  $t_R$  3.30 minutes; MS (ES<sup>+</sup>)  $m/z$  554 (M+H) <sup>b</sup>

<sup>1</sup>H NMR (500 MHz, methanol-d<sub>4</sub>) δ 9.35 (s, 1 H), 7.79 – 7.57 (m, 5 H), 7.47 (app. br. s, 2 H), 7.43 – 7.20 (m, 1 H), 6.12 (s, 1 H), 6.08 (s, 1 H), 5.79 (d, 1 H), 5.07 – 4.96 (m, 1 H), 3.51 – 3.38 (m, 1 H), 3.10 (ddd, 1 H), 2.64 (qd, 1 H), 2.56 – 2.34 (m, 1 H).

5 Example 823: [3-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]acetic acid

The compound prepared in Example 345 (150 mg) was treated using the method as detailed in Example 338 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] as the formic acid salt  
10 having the following physical properties (10 mg).

LC/MS *t<sub>R</sub>* 3.22 minutes; MS (ES<sup>+</sup>) *m/z* 548 and 550 (M+H)<sup>b</sup>

NMR analysis showed a 2: 1 ratio of atropisomers

Major atropisomer: <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.20 (br. s, 1 H), 9.76 (s, 1 H),  
15 8.21 (app. s, 1 H), 7.91 – 7.82 (m, 2 H), 7.79 (s, 1 H), 7.66 – 7.39 (m, 3 H), 7.35 – 7.20 (m, 1 H), 7.13 – 7.03 (m, 1 H), 6.29 (s, 1 H), 5.74 (dd, 1 H), 3.55 (s, 2 H), 3.36 – 3.25 (m, 1 H), 3.18 – 3.05 (m, 1 H), 2.66 – 2.54 (m, 1 H), 2.35 (app. br. s, 1 H).

Minor atropisomer: <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 12.67 (very br. s, 1 H), 9.70 (s, 1 H), 8.22 (app. s, 1 H), 7.91 – 7.82 (m, 2 H), 7.72 (s, 1 H), 7.66 – 7.39 (m, 3 H), 7.35 –  
20 7.20 (m, 1 H), 7.13 – 7.03 (m, 1 H), 6.34 (s, 1 H), 5.72 (dd, 1 H), 3.65 (s, 2 H), 3.36 – 3.25 (m, 1 H), 3.18 – 3.05 (m, 1 H), 2.66 – 2.54 (m, 1 H), 2.35 (app. br. s, 1 H).

25 Example 824: 2-methyl-2-propanyl {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 was conducted from tert-butyl 2-[(4-bromophenyl)sulfanyl]acetate [patent US2004/010019] to give the title compound having the following physical properties.

LC/MS *t<sub>R</sub>* 3.89 minutes; MS (ES<sup>+</sup>) *m/z* 602 (M+H)<sup>b</sup>

30 <sup>1</sup>H NMR (250 MHz, DMSO-d<sub>6</sub>) δ 12.15 (br. s, 1 H), 9.69 (s, 1 H), 7.78 – 7.74 (m, 3 H), 7.66 (d, 2 H), 7.53 (s, 1 H), 7.31 (d, 2 H), 5.97 (s, 1 H), 5.96 (s, 1 H), 5.61 (d, 1 H), 3.71

(s, 2 H), 3.49 – 3.31 (obs. m, 1 H), 3.08 – 2.89 (m, 1 H), 2.60 – 2.41 (obs. m, 1 H), 2.42 – 2.25 (m, 1 H), 1.33 (s, 9 H).

5 Example 825: {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetic acid

The compound prepared in Example 824 (235 mg) was treated as detailed in Example 363 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] having the following physical properties (104 mg).  
10 LC/MS  $t_R$  3.13 minutes; MS (ES<sup>+</sup>)  $m/z$  546 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.12 (br. s, 1 H), 9.68 (s, 1 H), 7.83 – 7.76 (m, 3 H), 7.64 (d, 2 H), 7.50 (s, 1 H), 7.29 (d, 2 H), 5.96 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd, 1 H), 3.74 (s, 2 H), 3.44 – 3.30 (obs. m, 1 H), 2.99 (dd, 1 H), 2.53 – 2.44 (obs. m, 1 H), 2.38 – 2.30 (m, 1 H).

15

Example 826: tert-butyl 2-[(3-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl}-1H-imidazol-5-yl)phenyl)sulfanyl]acetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 was conducted from tert-butyl 2-[(3-bromophenyl)sulfanyl]acetate [J. Med. Chem.,  
20 47(1), 18 (2004)] to give the title compound having the following physical properties.  
LC/MS  $t_R$  1.92 minutes; MS (ES<sup>+</sup>)  $m/z$  602 (M+H)<sup>a</sup>.

Example 827: {[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]thio}acetic acid

25 The compound prepared in Example 826 (80 mg) was treated as detailed in Example 363 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] having the following physical properties (40 mg).  
LC/MS  $t_R$  3.14 minutes; MS (ES<sup>+</sup>)  $m/z$  546 (M+H)<sup>b</sup>  
30 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.16 (br. s, 1 H), 9.69 (s, 1 H), 7.85 – 7.76 (m, 3 H), 7.64 (app. s, 1 H), 7.58 – 7.44 (m, 2 H), 7.33 – 7.23 (m, 1 H), 7.18 – 7.10 (m, 1 H), 5.97

(s, 1 H), 5.94 (s, 1 H), 5.62 (dd, 1 H), 3.79 (s, 2 H), 3.43 – 3.30 (obs. m, 1 H), 3.00 (dd, 1 H), 2.57 – 2.49 (obs. m, 1 H), 2.41 – 2.29 (m, 1 H).

5 Example 828: {[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]thio}acetic acid

The same operation as in Example 338 → Example 363 was conducted from the compound prepared in Example 826 to give the title compound after purification by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] having the following  
10 physical properties.

LC/MS  $t_R$  4.02 minutes; MS (ES<sup>+</sup>)  $m/z$  580 and 582 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.08 (br. s, 1 H), 9.71 (s, 1 H), 7.85 – 7.77 (m, 3 H), 7.64 (app. s, 1 H), 7.53 (d, 1 H), 7.41 (t, 1 H), 7.28 (d, 1 H), 5.98 (s, 1 H), 5.94 (s, 1 H), 5.57 (dd, 1 H), 3.83 (s, 2 H), 3.35 – 3.23 (obs. m, 1 H), 2.99 (dd, 1 H), 2.59 – 2.48 (obs.  
15 m, 1 H), 2.24 – 2.17 (m, 1 H).

Example 829: 2-methyl-2-propanyl {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]amino}acetate

20 The same operation as in Example 90 → Example 361 → Example 51 → Example 52 was conducted from tert-butyl 2-[(4-bromophenyl)amino]acetate [patent WO2004/113279] to give the title compound having the following physical properties.

LC/MS  $t_R$  3.40 minutes; MS (ES<sup>+</sup>)  $m/z$  585 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  10.96 (br. s, 1 H), 8.57 (s, 1 H), 7.61 (dd, 1 H), 7.54 (d, 1  
25 H), 7.51 (d, 1 H), 7.42 (app. br. s, 2 H), 7.07 (s, 1 H), 6.60 (d, 2 H), 6.32 (s, 1 H), 5.85 (d, 1 H), 5.71 (s, 1 H), 4.36 (br. s, 1 H), 3.82 (s, 2 H), 3.53 – 3.40 (m, 1 H), 3.39 – 3.28 (m, 1 H), 3.01 (dd, 1 H), 2.52 – 2.40 (m, 1 H), 1.50 (s, 9 H).

Example 830: {[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]amino}acetic acid hydrochloride  
30


To a 1,4-dioxane (1.5 mL) solution of the compound prepared in Example 829 (25.6 mg) was added concentrated hydrochloric acid (40  $\mu$ L) and the mixture stirred at room

temperature for 7 hours. On concentration, the residue was triturated with dichloromethane and the resultant precipitate collected by filtration to give the title compound having the following physical properties (20 mg).

LC/MS  $t_R$  1.66 minutes; MS ( $ES^+$ )  $m/z$  529 ( $M+H$ )<sup>e</sup>

- 5 <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  14.68 (br. s, 1 H), 14.51 (br. s, 1 H), 9.70 (s, 1 H), 7.84 - 7.79 (m, 3 H), 7.71 (d, 1 H), 7.53 (d, 2 H), 6.68 (d, 2 H), 6.06 (d, 1 H), 6.01 (s, 1 H), 5.80 (dd, 1 H), 3.87 (s, 2 H), 3.32 - 3.23 (m, 1 H), 3.14 - 3.05 (m, 1 H), 2.74 - 2.65 (m, 1 H), 2.38 - 2.29 (m, 1 H).

- 10 Example 831: 4-{2-[(3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-5-yl}thiophene-2-carboxamide



The compound prepared in Example 363 (20 mg) was treated as detailed in Example 114 to give the title compound having the following physical properties (16 mg).

- 15 LC/MS  $t_R$  1.36 minutes; MS ( $ES^+$ )  $m/z$  505 ( $M+H$ )<sup>a</sup>.

Example 832: 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-5-yl)-2-thiophenecarbonitrile

- To a pyridine (0.5 mL) solution of the compound prepared in Example 831 (16 mg) was added methanesulfonyl chloride (2.45  $\mu$ L). The mixture was stirred at room temperature for 20 hours, then warmed to 50 °C and stirred 7 hours. The reaction was cooled to room temperature, further methanesulfonyl chloride (4.90  $\mu$ L) added and the mixture stirred at room temperature for 16 hours. On concentration, the residue was dissolved in methanol (0.5 mL), treated with concentrated aqueous ammonia solution (12.3  $\mu$ L) and stirred at room temperature for 5 days. On concentration, the residue was suspended in saturated aqueous solution of sodium hydrogen carbonate and extracted into a 9:1 mixture of dichloromethane and methanol. The combined organic phases



were dried and concentrated and the residue purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (8 mg).

- 5 LC/MS  $t_R$  3.60 minutes; MS (ES<sup>+</sup>)  $m/z$  487 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.24 (br. s, 1 H), 9.68 (s, 1 H), 8.20 (s, 1 H), 7.97 (s, 1 H), 7.85 - 7.73 (m, 3 H), 7.48 (s, 1 H), 5.96 (s, 1 H), 5.95 (s, 1 H), 5.61 (d, 1 H), 3.38 - 3.28 (obs. m, 1 H), 2.99 (dd, 1 H), 2.53 - 2.44 (obs. m, 1 H), 2.35 - 2.25 (m, 1 H).

10 Example 833: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-N-(methylsulfonyl)-2-thiophenecarboxamide

- To a cooled (0 °C) tetrahydrofuran (2.5 mL) suspension of the compound prepared in Example 363 (170 mg) was sequentially added triethylamine (159  $\mu$ L)  
15 followed by isobutyl chloroformate (95.9  $\mu$ L) and the mixture stirred at room temperature for 2 hours. On cooling to 0 °C, a solution of methanesulfonamide (164 mg) and triethylamine (93.7  $\mu$ L) in tetrahydrofuran (2.5 mL) was added. 4-Dimethylaminopyridine (40.8 mg) was then added and stirring continued at room temperature for 16 hours. To the reaction mixture, concentrated aqueous ammonia  
20 solution (125  $\mu$ L) was added and the mixture stirred at room temperature a further hour then concentrated. The residue was suspended in 1 M hydrochloric acid (5 mL) and extracted into a 9:1 mixture of dichloromethane and methanol. The aqueous layer was extracted thrice more with a 9:1 mixture of chloroform and propan-2-ol, the combined organic layers dried and concentrated. The residue was purified by high performance  
25 liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] then further purified by trituration from dichloromethane, the resultant precipitate being isolated by filtration to obtain the title compound (11.5 mg). The filtrate was concentrated and residual 2-methyl-propan-1-ol removed by azeotrope with deuterated chloroform to give a second batch of the title  
30 compound having the following physical properties (13 mg).

LC/MS  $t_R$  3.15 minutes; MS (ES<sup>+</sup>)  $m/z$  583 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  13.03 - 11.93 (m, 2 H), 9.69 (s, 1 H), 8.22 (s, 1 H), 7.87 (s, 1 H), 7.82 - 7.74 (m, 3 H), 7.31 (br. s, 1 H), 5.97 (s, 1 H), 5.95 (s, 1 H), 5.62 (dd,

1 H), 3.44 - 3.28 (obs. m, 1 H), 3.27 (s, 3 H), 3.01 (dd, 1 H), 2.56 - 2.48 (obs. m, 1 H), 2.34 (app. br. s, 1 H).

5 Example 834: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3-fluoro-2-thiophenecarboxylic acid

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 8 → Example 55 → Example 24 was conducted from methyl 4-bromo-3-fluorothiophene-2-carboxylate [Tetrahedron Lett., 42(50), 8797 (2001)] to give, after trituration from dichloromethane, the title compound having the following physical  
10 properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 11 was used).

LC/MS  $t_R$  3.34 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.30 (br. s, 1 H), 12.26 (br. s, 1 H), 9.68 (s, 1 H), 7.85 - 7.78 (m, 3 H), 7.77 (s, 1 H), 7.29 (s, 1 H), 5.96 (app. s, 2 H), 5.63 (dd, 1 H), 3.38  
15 - 3.27 (obs. m, 1 H), 2.99 (dd, 1 H), 2.53 - 2.47 (obs. m, 1 H), 2.35 - 2.28 (m, 1 H).

Example 835: tert-butyl 4-(2-((3S)-7-[5-chloro-2-(1H-1,2,3,4-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl)-1H-imidazol-5-yl)-3-fluoropyridine-2-carboxylate

20 The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 was conducted from 4-chloro-3-fluoropyridine-2-carboxylic acid [Eur. J. Org. Chem., 10, 2116 (2005)] to give the title compound having the following physical properties.

LC/MS  $t_R$  2.00 minutes; MS (ES<sup>+</sup>)  $m/z$  575 (M+H), 519 (M-C(CH<sub>3</sub>)<sub>3</sub>+H)<sup>+</sup>; 447 (M-CO<sub>2</sub>  
25 C(CH<sub>3</sub>)<sub>3</sub>-N<sub>2</sub>+H)<sup>a</sup>.

Example 836: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-3-fluoro-2-pyridinecarboxylic acid

To a 1,4-dioxane (4.2 mL) solution of the compound prepared in Example 835 (102  
30 mg) was added 6 M hydrochloric acid (4.2 mL) and the mixture heated at 60 °C for 1 hour. On concentration the residue was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile

phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (33.1 mg).

LC/MS  $t_R$  3.16 minutes; MS (ES<sup>+</sup>)  $m/z$  519 (M+H), 447 (M-CO<sub>2</sub>-N<sub>2</sub>+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.52 (br. s, 1 H), 12.63 (br. s, 1 H), 9.69 (s, 1 H),  
5 8.41 (d, 1 H), 8.04 (t, 1 H), 7.86 - 7.75 (m, 3 H), 7.69 (d, 1 H), 5.98 (app. s, 2 H), 5.77 -  
5.58 (m, 1 H), 3.37 - 3.25 (obs. m, 1 H), 3.02 (dd, 1 H), 2.59 - 2.45 (m, 1 H), 2.39 - 2.28  
(m, 1 H).

10 Example 837: 4-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)-3-thiophenecarboxylic acid hydrochloride

The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 4-bromothiophene-3-carboxylic acid to give the title compound as the hydrochloride salt having the following physical properties.

15 LC/MS  $t_R$  3.37 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>)  $\delta$  14.34 (br. s, 1 H), 9.69 (s, 1 H), 8.46 (d, 1 H), 7.96 (d, 1 H), 7.84 (dd, 1 H), 7.82 (d, 1 H), 7.79 (s, 1 H), 7.71 (d, 1 H), 6.06 - 6.04 (m, 1 H),  
6.03 - 6.01 (m, 1 H), 5.82 (dd, 1 H), 3.26 (td, 1 H), 3.14 - 3.05 (m, 1 H), 2.76 - 2.66 (m, 1 H), 2.33 - 2.21 (m, 1 H).

20

Example 838(1) and 838(2): 2-tert-butyl 4-methyl thiophene-2,4-dicarboxylate and 4-(methoxycarbonyl)thiophene-2-carboxylic acid

tert-Butyl 4-bromothiophene-2-carboxylate [J. Med. Chem., 55(12), 5982, supporting information (2012)] (1.0 g) was treated as detailed in Example 673 to give  
25 the title compounds having the following physical properties.

Example 838(1): 309 mg

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.22 (d, 1 H), 8.07 (d, 1 H), 3.89 (s, 3 H), 1.59 (s, 9 H).

30 Example 838(2): 363 mg

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.35 (d, 1 H), 8.27 (d, 1 H), 3.91 (s, 3 H).

Example 839: 5-[(tert-butoxy)carbonyl]thiophene-3-carboxylic acid

To a stirred solution of the compound prepared in Example 838(1) (305 mg) in tetrahydrofuran (12.8 mL) and water (12.8 mL) was added lithium hydroxide monohydrate (63.1 mg) and the mixture stirred at room temperature for 5 hours. To the reaction mixture, water (20 mL) and tert-butyl methyl ether (30 mL) were added and the phases separated. The organic layer was washed with a saturated aqueous solution of sodium hydrogen carbonate (30 mL), the aqueous layers then combined and treated with 2 M hydrochloric acid until pH 1 - 2 was attained. This aqueous suspension was extracted with ethyl acetate, the combined organic layers washed with saturated saline, dried and concentrated to give the title compound having the following physical properties (269 mg).  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.04 (br. s, 1 H), 8.51 (d, 1 H), 7.87 (d, 1 H), 1.53 (s, 9 H).

Example 840: 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-pyrrol-2-yl)-2-thiophenecarboxylic acid

The same operation as in Example 676 → Example 665 → Example 666 → Example 667 → Example 24 was conducted from the compound prepared in Example 839 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 665 in the operation, the compound prepared in Example 617 was used).  
LC/MS *t<sub>R</sub>* 3.94 minutes; MS (ES<sup>+</sup>) *m/z* 505 (M+H)<sup>b</sup>  
<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.08 (br. s, 1 H), 11.22 (br. s, 1 H), 9.70 (s, 1 H), 7.97 (d, 1 H), 7.84 - 7.73 (m, 4 H), 6.32 (t, 1 H), 6.00 (s, 1 H), 5.92 (s, 1 H), 5.67 (d, 1 H), 5.62 (t, 1 H), 3.13 (td, 1 H), 2.94 (dd, 1 H), 2.55 - 2.41 (obs. m, 1 H), 2.18 (dd, 1 H).


Example 841: ethyl 4-cyanothiophene-2-carboxylate

To an N,N-dimethylformamide (75 mL) solution of ethyl 4-bromothiophene-2-carboxylate [patent US2012/022123] (4.98 g) was added copper(I) cyanide (2.09g) and the mixture heated at 150 °C for 9.5 hours. The reaction mixture was cooled to room temperature, diluted with ethyl acetate (350 mL) and washed with water (200 mL) and 2 M aqueous ammonia solution (200 mL). The organic layer was dried, concentrated and

the residue purified by column chromatography (0 – 20% ethyl acetate in heptanes) to give the title compound having the following physical properties (2.39 g).

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  8.07 (d, 1 H), 7.93 (d, 1 H), 4.39 (q, 2 H), 1.39 (t, 3 H).

5 Example 842: ethyl 4-carbamimidoylthiophene-2-carboxylate




The compound prepared in Example 841 (1.79 g) was treated as detailed in Example 237 to give, after trituration from a mixture of water (15 mL) and a saturated aqueous solution of sodium hydrogen carbonate (15 mL), the title compound having the

10 following physical properties (1.08 g).

$^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ )  $\delta$  8.37 (d, 1 H), 8.18 (d, 1 H), 7.36 (br. s, 3 H), 4.31 (q, 2 H), 1.31 (t, 3 H).

15 Example 843: ethyl 4-{5-[7-(2-{bis[(tert-butoxy)carbonyl]amino}-5-chlorophenyl)-5-oxo-1,2,3,5-tetrahydroindolizin-3-yl]-1H-imidazol-2-yl}thiophene-2-carboxylate



The compound prepared in Example 842 (0.57 g) was treated with the compound prepared in Example 623 (1.60 g) as detailed in Example 620 to give the title compound having the following physical properties (1.44 g).

20 LC/MS  $t_R$  2.08 minutes; MS ( $\text{ES}^+$ )  $m/z$  704 ( $\text{M}+\text{Na}$ ), 681 ( $\text{M}+\text{H}$ )<sup>a</sup>.

Example 844: ethyl 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl}-1H-imidazol-2-yl)-2-thiophenecarboxylate

25 The same operation as in Example 40 → Example 24 was conducted from the compound prepared in Example 843 to give the title compound having the following physical properties.

LC/MS  $t_R$  3.46 minutes; MS ( $ES^+$ )  $m/z$  534 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.48 (br. s, 1 H), 9.68 (s, 1 H), 8.18 (d, 1 H), 8.15 (d, 1 H), 7.85 - 7.74 (m, 3 H), 6.89 (s, 1 H), 5.91 (app. s, 2 H), 5.58 (d, 1 H), 4.32 (q, 2 H), 3.36 - 3.25 (m, 1 H), 2.93 (dd, 1 H), 2.47 - 2.38 (m, 1 H), 2.26 (dd, 1 H), 1.32 (t, 3 H).

5

Example 845: ethyl 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-4-fluoro-1H-imidazol-2-yl)-2-thiophenecarboxylate

The compound prepared in Example 844 (296 mg) was treated as detailed in Example 364 to give the title compound having the following physical properties (20.4 mg).

10

LC/MS  $t_R$  4.25 minutes; MS ( $ES^+$ )  $m/z$  552 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz,  $CDCl_3$ )  $\delta$  11.26 (br. s, 1 H), 8.62 (s, 1 H), 8.07 (s, 1 H), 7.93 (s, 1 H), 7.62 (dd, 1 H), 7.58 - 7.51 (m, 2 H), 6.36 (s, 1 H), 5.85 (d, 1 H), 5.79 (s, 1 H), 4.36 (q, 2 H), 3.42 - 3.27 (m, 1 H), 3.06 (dd, 1 H), 2.84 (dd, 1 H), 2.61 - 2.46 (m, 1 H), 1.39 (t, 3 H).

15

Example 846: ethyl 4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-2-yl)-2-thiophenecarboxylate

The compound prepared in Example 844 (90 mg) was treated as detailed in Example 44 to give the title compound having the following physical properties (54.9 mg).

20

LC/MS  $t_R$  4.10 minutes; MS ( $ES^+$ )  $m/z$  568 and 570 ( $M+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.90 (br. s, 1 H), 9.64 (s, 1 H), 8.21 (d, 1 H), 8.17 (d, 1 H), 7.84 - 7.78 (m, 2 H), 7.76 - 7.72 (m, 1 H), 5.98 (s, 1 H), 5.92 (s, 1 H), 5.60 (dd, 1 H), 4.32 (q, 2 H), 3.18 (td, 1 H), 3.02 (ddd, 1 H), 2.63 - 2.53 (m, 1 H), 2.16 - 2.08 (m, 1 H), 1.32 (t, 3 H).

25

Example 847: 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-2-yl)-2-thiophenecarboxylic acid

The same operation as in Example 8  $\rightarrow$  Example 40  $\rightarrow$  Example 24 was conducted from the compound prepared in Example 843 to give the title compound having the following physical properties.

30

LC/MS  $t_R$  2.96 minutes; MS ( $ES^+$ )  $m/z$  506 ( $M+H$ ), 478 ( $M-N_2+H$ )<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.25 (br. s, 1 H), 12.46 (br. s, 1 H), 9.69 (s, 1 H), 8.18 - 8.06 (m, 2 H), 7.84 - 7.76 (m, 3 H), 6.88 (br. s, 1 H), 5.92 (app. s, 2 H), 5.60 (d, 1 H), 3.40 - 3.26 (obs. m, 1 H), 2.94 (dd, 1 H), 2.48 - 2.38 (m, 1 H), 2.26 (dd, 1 H).

5 Example 848: 4-(5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-4-fluoro-1H-imidazol-2-yl)-2-thiophenecarboxylic acid

The same operation as in Example 364 → Example 8 → Example 40 → Example 24 was conducted from the compound prepared in Example 843 to give the title compound having the following physical properties.

10 LC/MS *t*<sub>R</sub> 3.66 minutes; MS (ES<sup>+</sup>) *m/z* 524 (M+H)<sup>b</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.32 (br. s, 1 H), 12.57 (s, 1 H), 9.66 (s, 1 H), 8.12 - 8.03 (m, 2 H), 7.85 - 7.77 (m, 2 H), 7.76 (d, 1 H), 5.97 (s, 1 H), 5.91 (s, 1 H), 5.60 (dd, 1 H), 3.25 - 3.15 (m, 1 H), 3.05 - 2.95 (m, 1 H), 2.62 - 2.53 (m, 1 H), 2.24 - 2.13 (m, 1 H).

15

Example 849: 4-(4-chloro-5-{7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-2-yl)-2-thiophenecarboxylic acid

The compound prepared in Example 847 (70 mg) was treated as detailed in Example 338. The reaction mixture was concentrated directly and purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% formic acid in acetonitrile) in mobile phase A (0.1% aqueous formic acid)] to give the title compound having the following physical properties (18.1 mg).

20 LC/MS *t*<sub>R</sub> 2.57 minutes; MS (ES<sup>+</sup>) *m/z* 540 and 542 (M+H)<sup>c</sup>

<sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>) δ 13.30 (br. s, 1 H), 12.85 (br. s, 1 H), 9.64 (s, 1 H), 8.13 (s, 1 H), 8.09 (s, 1 H), 7.83 - 7.78 (m, 2 H), 7.74 (app. s, 1 H), 5.99 (s, 1 H), 5.91 (s, 1 H), 5.61 (dd, 1 H), 3.21 - 3.12 (m, 1 H), 3.06 - 2.97 (m, 1 H), 2.62 - 2.53 (m, 1 H), 2.16 - 2.07 (m, 1 H).

30 Example 850: 2-methoxyethyl [4-(2-{(3S)-7-[5-chloro-2-(4-cyano-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny}-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 283 → Example 8 → Example 38 → Example 39 → Example 290 was conducted from the compound prepared in Example 288 to give

the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 324 was used)

LC/MS  $t_R$  0.72 minutes; MS (ES<sup>+</sup>)  $m/z$  597 (M+H)<sup>f</sup>

- 5 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.6 – 11.8 (m, 1H), 9.85 – 9.55 (m, 1H), 9.35 (s, 1 H), 7.85 – 7.73 (m, 3 H), 7.64 – 7.02 (m, 5 H), 5.95 (s, 1 H), 5.94 (s, 1 H), 5.60 (d, 1 H), 4.18 (dd, 2 H), 3.56 (dd, 2 H), 3.46 – 3.30 (m, 1 H), 3.27 (s, 3H), 2.99 (dd, 1 H), 2.60-2.45 (m, 1 H), 2.40 – 2.25 (m, 1 H).

- 10 Example 851: 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(4-cyano-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]phenyl]carbamate

The same operation as in Example 44 was conducted from the compound prepared in Example 850 to give the title compound having the following physical properties.

- 15 TLC: R<sub>f</sub> 0.51 (CH<sub>2</sub>Cl<sub>2</sub>/AcOEt/MeOH, 8/4/1)  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.8 (s, 1H), 9.89 (s, 1H), 9.36 (s, 1 H), 7.85 – 7.77 (m, 3 H), 7.63 – 7.48 (m, 4 H), 5.96 (s, 1 H), 5.94 (s, 1 H), 5.56 (dd, 1 H), 4.20 (dd, 2 H), 3.56 (dd, 2 H), 3.28 (s, 3H), 3.25 – 3.17 (m, 1 H), 3.06-2.90 (m, 1 H), 2.60-2.45 (m, 1 H), 2.28 – 2.14 (m, 1 H).

Example 852: 1-(2-((3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-5-oxo-1,2,3,5-tetrahydro-7-indolizinyl)-4-chlorophenyl)-1H-1,2,3-triazole-4-carbonitrile

- 25 The same operation as in Example 283 → Example 8 → Example 38 → Example 39 → Example 10 → Example 290 → Example 338 → Example 287 was conducted from the compound prepared in Example 288 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 38 in the operation, the compound prepared in Example 193 was used)

LC/MS  $t_R$  0.70 minutes; MS (ES<sup>+</sup>)  $m/z$  530 (M+H)<sup>f</sup>


- 30 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.7 (s, 1H), 9.36 (s, 1 H), 8.19 (d, 1H), 7.83 – 7.72 (m, 3 H), 7.63 (dd, 1 H), 6.50 (d, 1H), 6.17 (s, 2 H), 5.95 (s, 1 H), 5.93 (s, 1 H), 5.53 (dd, 1 H), 3.30 – 3.06 (m, 1 H), 3.05 – 2.92 (m, 1 H), 2.60-2.45 (m, 1 H), 2.24 – 2.12 (m, 1 H).




Example 853(1) to Example 853(2)

The same operation as in Example 485 was conducted from corresponding alcohols to give the title compounds having the following physical properties.

5

Example 853(1): 3-hydroxy-3-methylbutyl carbonochloridate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  4.51 (t, 2 H), 1.94 (t, 2 H), 1.29 (s, 6 H).

10 Example 853(2): 2-hydroxy-2-methylpropyl carbonochloridate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  4.18 (s, 2 H), 1.30 (s, 6 H).

Example 854(1) to Example 854(3)


15 The compounds of the present invention having the following physical data were prepared using the compound prepared in Example 194 and the corresponding chloroformates using the method as detailed in Example 128.

20 Example 854(1): 2-hydroxy-2-methylpropyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  0.66 minutes; MS ( $\text{ES}^+$ )  $m/z$  588 ( $\text{M}+\text{H}$ )<sup>f</sup>


$^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  12.7-12.1 (m, 1 H), 10.1-9.98 (m, 1 H), 9.70-9.60 (m, 1 H), 8.62-8.52 (m, 1 H), 8.05-7.95 (m, 1 H), 7.86-7.72 (m, 4 H), 7.52-7.20 (m, 1 H),  
 25 5.97-5.88 (m, 2 H), 5.60 (d, 1 H), 4.54 (m, 1 H), 3.90-3.83 (m, 2 H), 3.38 (m, 1 H), 2.98 (m, 1 H), 2.60-2.20 (m, 2 H), 1.13 (s, 6 H).

Example 854(2): 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-pyridinyl]carbamate



- 5 LC/MS  $t_R$  0.69 minutes; MS (ES<sup>+</sup>)  $m/z$  618 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.7 (s, 0.2 H), 12.2 (s, 0.8 H), 10.3 (s, 0.2 H), 10.1 (s, 0.8 H), 9.69 (s, 0.2 H), 9.67 (s, 0.8 H), 8.60 - 8.51 (m, 1 H), 8.03 - 7.95 (m, 1 H), 7.85 - 7.75 (m, 4 H), 7.52 (d, 0.8 H), 7.21 (d, 0.2 H), 5.97 - 5.90 (m, 2 H), 5.64 - 5.58 (m, 1 H), 4.23 - 4.16 (m, 2 H), 3.66 - 3.59 (m, 2 H), 3.58 - 3.52 (m, 2 H), 3.50 - 3.32 (m, 5 H),  
 10 3.05 - 2.92 (m, 1 H), 2.62 - 2.18 (m, 2 H), 1.07 (t, 3 H).

Example 854(3): 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-pyridinyl]carbamate



- 15 LC/MS  $t_R$  0.72 minutes; MS (ES<sup>+</sup>)  $m/z$  632 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.7 (s, 0.2 H), 12.2 (s, 0.8 H), 10.3 (s, 0.2 H), 10.1 (s, 0.8 H), 9.69 (s, 0.2 H), 9.67 (s, 0.8 H), 8.59 - 8.51 (m, 1 H), 8.03 - 7.95 (m, 1 H), 7.82 - 7.75 (m, 4 H), 7.53 (d, 0.8 H), 7.22 (d, 0.2 H), 5.97 - 5.92 (m, 2 H), 5.64 - 5.58 (m, 1 H),  
 20 4.24 - 4.16 (m, 2 H), 3.66 - 3.56 (m, 2 H), 3.56 - 3.51 (m, 2 H), 3.46 - 3.28 (m, 3 H), 3.24 (s, 3 H), 3.05 - 2.92 (m, 1 H), 2.60 - 2.18 (m, 2 H).

Example 855(1) to Example 855(3)


The compounds of the present invention having the following physical data were prepared using the compound prepared in Example 198 and the corresponding chloroformates using the method as detailed in Example 128.

5 Example 855(1): 3-hydroxy-3-methylbutyl [5-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  0.82 minutes; MS (ES<sup>+</sup>)  $m/z$  636 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.0 (brs, 1 H), 10.2 (s, 1 H), 9.69 (s, 1 H), 8.55 (d, 1 H), 8.04 (dd, 1 H), 7.92 (d, 1 H), 7.86-7.75 (m, 3 H), 5.97 (s, 1 H), 5.92 (s, 1 H), 5.55 (dd, 1 H), 4.35 (s, 1 H), 4.20 (t, 2 H), 3.28 (m, 1 H), 2.98 (m, 1 H), 2.52 (m, 1 H), 2.20 (m, 1 H), 1.74 (t, 2 H), 1.13 (s, 6 H).

15 Example 855(2): 2-(2-methoxyethoxy)ethyl [5-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate



LC/MS  $t_R$  0.82 minutes; MS (ES<sup>+</sup>)  $m/z$  652 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  13.1 (s, 1 H), 10.4 (s, 1 H), 9.70 (s, 1 H), 8.56 (d, 1 H), 8.02 (dd, 1 H), 7.91 (d, 1 H), 7.84-7.77 (m, 3 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.59-5.52 (m, 1 H), 4.26-4.19 (m, 2 H), 3.67-3.61 (m, 2 H), 3.59-3.52 (m, 2 H), 3.47-3.40 (m, 2 H), 3.37-3.24 (m, 1 H), 3.23 (s, 3 H), 3.04-2.92 (m, 1 H), 2.60-2.28 (m, 1 H), 2.27-2.13 (m, 1 H).

25 Example 855(3): 2-(2-ethoxyethoxy)ethyl [5-(4-chloro-2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinyl]carbamate


LC/MS  $t_R$  0.86 minutes; MS (ES<sup>+</sup>)  $m/z$  666 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 13.1 (s, 1 H), 10.4 (s, 1 H), 9.70 (s, 1 H), 8.56 (d, 1 H), 8.02 (dd, 1 H), 7.91 (d, 1 H), 7.84 – 7.77 (m, 3 H), 5.98 (s, 1 H), 5.93 (s, 1 H), 5.59 – 5.52 (m, 1 H), 4.26 – 4.19 (m, 2 H), 3.68 – 3.61 (m, 2 H), 3.59 – 3.52 (m, 2 H), 3.51 – 3.20 (m, 5 H), 3.04 – 2.92 (m, 1 H), 2.60 – 2.28 (m, 1 H), 2.27 – 2.13 (m, 1 H), 1.08 (t, 3H).

#### Example 856(1) to Example 856(2)

The compounds of the present invention having the following physical data were prepared using the compound prepared in Example 593 and the corresponding chloroformates using the method as detailed in Example 128.

#### Example 856(1): 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate



15

LC/MS *t<sub>R</sub>* 0.82 minutes; MS (ES<sup>+</sup>) *m/z* 636 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1 H), 10.3 (s, 1 H), 9.69 (s, 1 H), 8.46 (s, 1 H), 7.92 – 7.77 (m, 5 H), 5.96 (s, 1 H), 5.93 (s, 1 H), 5.53 (d, 1 H), 4.25 – 4.18 (m, 2 H), 3.67 – 3.60 (m, 2 H), 3.59 – 3.52 (m, 2 H), 3.47 – 3.40 (m, 2 H), 3.37 – 3.24 (m, 1 H), 3.23 (s, 3 H), 3.03 – 2.92 (m, 1 H), 2.60 – 2.28 (m, 1 H), 2.27 – 2.13 (m, 1 H).

20

#### Example 856(2): 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate

25 LC/MS *t<sub>R</sub>* 0.85 minutes; MS (ES<sup>+</sup>) *m/z* 650 (M+H)<sup>f</sup>


<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1 H), 10.3 (s, 1 H), 9.69 (s, 1 H), 8.45 (s, 1 H), 7.91 – 7.77 (m, 5 H), 5.96 (s, 1 H), 5.93 (s, 1 H), 5.52 (d, 1 H), 4.25 – 4.18 (m, 2 H), 3.67 – 3.60 (m, 2 H), 3.58 – 3.52 (m, 2 H), 3.50 – 3.19 (m, 5 H), 3.03 – 2.92 (m, 1 H),

2.60 - 2.28 (m, 1 H), 2.27 - 2.13 (m, 1 H), 1.08 (t, 3 H).

Example 857(1) to Example 857(2)

The compounds of the present invention having the following physical data were prepared using the compound prepared in Example 718(6) and the corresponding chloroformates using the method as detailed in Example 128.

Example 857(1): 2-(2-methoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate



LC/MS  $t_R$  0.67 minutes; MS (ES<sup>+</sup>)  $m/z$  619 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>)  $\delta$  9.43 (s, 1 H), 8.63 - 8.50 (m, 1 H), 8.10 - 7.97 (m, 1 H), 7.97 - 7.83 (m, 2 H), 7.77 (dd, 1 H), 7.70 (d, 1 H), 7.49 - 7.39 (m, 1 H), 6.34 (s, 1 H), 5.78 - 5.69 (m, 1 H), 4.31 (t, 2 H), 3.77 (t, 2 H), 3.69 - 3.63 (m, 2 H), 3.58 - 3.52 (m, 2 H), 3.36 (s, 3 H), 3.28 - 3.21 (m, 1 H), 3.01 - 2.85 (m, 1 H), 2.76 - 2.60 (m, 1 H), 2.50 - 2.32 (m, 1 H).

Example 857(2): 2-(2-ethoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  0.70 minutes; MS (ES<sup>+</sup>)  $m/z$  633 (M+H)<sup>+</sup>


<sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>)  $\delta$  9.43 (s, 1 H), 8.60 - 8.49 (m, 1 H), 8.08 - 7.98 (m, 1 H), 7.93 - 7.83 (m, 2 H), 7.77 (dd, 1 H), 7.69 (d, 1 H), 7.49 - 7.39 (m, 1 H), 6.35 (s, 1 H), 5.78 - 5.70 (m, 1 H), 4.32 (t, 2 H), 3.76 (t, 2 H), 3.69 - 3.63 (m, 2 H), 3.62 - 3.57 (m, 2 H), 3.53 (q, 2 H), 3.28 - 3.22 (m, 1 H), 3.01 - 2.86 (m, 1 H), 2.77 - 2.59 (m, 1 H), 2.51 - 2.32 (m, 1 H), 1.17 (t, 3 H).

Example 858(1) to Example 858(2)

The compounds of the present invention having the following physical data were prepared using the compound prepared in Example 330 and the corresponding chloroformates using the method as detailed in Example 128.

5

Example 858(1): 2-(2-methoxyethoxy)ethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-pyridinyl]carbamate



10 LC/MS  $t_R$  0.80 minutes; MS ( $ES^+$ )  $m/z$  653 ( $M+H$ )<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, methanol- $d_4$ )  $\delta$  9.44 (s, 1 H), 8.58 (d, 1 H), 8.05 (dd, 1 H), 7.99 (d, 1 H), 7.92 (d, 1 H), 7.77 (dd, 1 H), 7.70 (d, 1 H), 6.37 (s, 1 H), 5.71 – 5.64 (m, 1 H), 4.33 (t, 2 H), 3.76 (t, 2 H), 3.72 – 3.62 (m, 2 H), 3.59 – 3.52 (m, 2 H), 3.36 (s, 3 H), 3.26 – 3.20 (m, 1 H), 2.99 – 2.85 (m, 1 H), 2.76 – 2.60 (m, 1 H), 2.43 – 2.29 (m, 1 H).

15

Example 858(2): 2-(2-ethoxyethoxy)ethyl [5-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  0.83 minutes; MS (ES<sup>+</sup>)  $m/z$  667 (M+H)<sup>f</sup>

20 <sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>) δ 9.44 (s, 1 H), 8.62 – 8.55 (m, 1 H), 8.09 – 7.96 (m, 2 H), 7.92 (d, 1 H), 7.78 (dd, 1 H), 7.70 (d, 1 H), 6.36 (s, 1 H), 5.71 – 5.63 (m, 1 H), 4.33 (t, 2 H), 3.77 (t, 2 H), 3.69 – 3.63 (m, 2 H), 3.62 – 3.57 (m, 2 H), 3.53 (q, 2 H), 3.28 – 3.20 (m, 1 H), 3.00 – 2.85 (m, 1 H), 2.77 – 2.61 (m, 1 H), 2.43 – 2.29 (m, 1 H), 1.17 (t, 3 H).


25

Example 859(1) to Example 859(2)

The compounds of the present invention having the following physical data were prepared using the compound prepared in Example 722 and the corresponding

chloroformates using the method as detailed in Example 128.

Example 859(1): 2-(2-methoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate



LC/MS  $t_R$  0.79 minutes; MS ( $ES^+$ )  $m/z$  637 ( $M+H$ )<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, methanol- $d_4$ )  $\delta$  9.43 (s, 1 H), 8.46 (d, 1 H), 7.99 - 7.87 (m, 3 H), 7.77 (dd, 1 H), 7.70 (d, 1 H), 6.36 (s, 1 H), 5.67 - 5.61 (m, 1 H), 4.32 (t, 2 H), 3.78 (t, 2 H), 3.69 - 3.61 (m, 2 H), 3.59 - 3.52 (m, 2 H), 3.37 - 3.34 (m, 3 H), 3.26 - 3.20 (m, 1 H), 2.99 - 2.85 (m, 1 H), 2.76 - 2.62 (m, 1 H), 2.41 - 2.29 (m, 1 H).

Example 859(2): 2-(2-ethoxyethoxy)ethyl [5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate

LC/MS  $t_R$  0.83 minutes; MS ( $ES^+$ )  $m/z$  651 ( $M+H$ )<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, methanol- $d_4$ )  $\delta$  9.43 (s, 1 H), 8.45 (d, 1 H), 7.99 - 7.88 (m, 3 H), 7.77 (dd, 1 H), 7.70 (d, 1 H), 6.35 (s, 1 H), 5.67 - 5.61 (m, 1 H), 4.33 (t, 2 H), 3.76 (t, 2 H), 3.69 - 3.63 (m, 2 H), 3.62 - 3.57 (m, 2 H), 3.53 (q, 2 H), 3.28 - 3.20 (m, 1 H), 3.00 - 2.85 (m, 1 H), 2.75 - 2.58 (m, 1 H), 2.42 - 2.27 (m, 1 H), 1.17 (t, 3 H).

Example 860: (3S)-3-[5-(5-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 52 → Example 44 → Example 55 was conducted from the compound prepared in Example 228 to give the title compound having the following physical properties.

LC/MS  $t_R$  0.71 minutes; MS ( $ES^+$ )  $m/z$  506 ( $M+H$ )<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.6 (brs, 1 H), 9.68 (s, 1 H), 8.02-7.70 (m, 4 H),

7.65 (d, 1 H), 7.00 (d, 1 H), 5.96 (s, 1 H), 5.92 (s, 1 H), 5.63 (d, 1 H), 5.54 (brs, 2 H), 3.22 (m, 1 H), 2.98 (m, 1 H), 2.45 (m, 1 H), 2.20 (m, 1 H).

5 Example 861: (6S)-6-[5-(5-amino-2-pyridinyl)-4-chloro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one

The same operation as in Example 51 → Example 52 → Example 44 → Example 55 was conducted from the compound prepared in Example 336 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 219 was used)

10 LC/MS  $t_R$  0.68 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.77 (s, 1 H), 8.07 (d, 1 H), 7.84 (d, 1 H), 7.67 - 7.60 (m, 2 H), 7.46 (d, 1 H), 7.01 (dd, 1 H), 6.41 (s, 1 H), 5.74 (dd, 1 H), 3.79 (s, 2 H), 3.46 - 3.15 (m, 2 H), 2.95 - 2.80 (m, 1 H), 2.55 - 2.39 (m, 1 H).

15 Example 862: methyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl])-1H-imidazol-5-yl)phenyl]carbamate

The same operation as in Example 44 was conducted from the compound prepared in Example 718(13) to give the title compound having the following physical properties.

20 LC/MS  $t_R$  0.84 minutes; MS (ES<sup>+</sup>)  $m/z$  564 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.9 (s, 1 H), 9.79 (s, 1 H), 9.74 (s, 1 H), 7.98 (d, 1 H), 7.88 - 7.78 (m, 2 H), 7.62 - 7.50 (m, 4 H), 6.32 (s, 1 H), 5.59 - 5.52 (m, 1 H), 3.67 (s, 3 H), 3.20 - 3.02 (m, 1 H), 2.81 - 2.71 (m, 1 H), 2.62 - 2.39 (m, 1 H), 2.18 - 2.09 (m, 25 1 H).

Example 863: methyl [4-(4-chloro-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate

30 The same operation as in Example 44 was conducted from the compound prepared in Example 309 to give the title compound having the following physical properties.

LC/MS  $t_R$  2.18 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>d</sup>  
<sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>)  $\delta$  9.31 (s, 1 H), 7.66-7.46 (m, 7 H), 6.10 (s, 1 H),



6.06 (s, 1 H), 5.69 (dd, 1 H), 3.74 (s, 3 H), 3.42 (m, 1 H), 3.08 (m, 1 H), 2.62 (m, 1 H), 2.51 (s, 3 H), 2.38 (m, 1 H).

Example 864(1) to Example 864(3)

- 5           The compounds of the present invention having the following physical data were prepared from the compound prepared in Example 9 using the corresponding alpha-bromoketones in the process of Example 51 → Example 52.

Example 864(1): Methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3-methylphenyl]carbamate

- 10           LC/MS  $t_R$  3.73 minutes; MS (ES<sup>+</sup>)  $m/z$  543 (M+H)<sup>d</sup>  
              <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.26 – 12.10 (m, 1 H), 9.74 – 9.57 (m, 2 H), 7.84 – 7.80 (m, 3 H), 7.62 (d, 1 H), 7.39 – 7.30 (m, 2 H), 7.19 (d, 1 H), 5.99 (s, 2 H), 5.69 – 5.66 (m, 1 H), 3.73 – 3.69 (m, 3 H), 3.48 – 3.30 (m, 1 H), 3.08 – 3.00 (m, 1 H), 2.56 –  
15           2.53 (m, 2 H), 2.38 – 2.30 (m, 3 H).

Example 864(2): 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)phenyl]carbamate

- LC/MS  $t_R$  0.66 minutes; MS (ES<sup>+</sup>)  $m/z$  573 (M+H)<sup>f</sup>  
20           <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.02 (brs, 1 H), 9.69 (s, 1 H), 9.68 (s, 1 H), 7.85 – 7.74 (m, 3 H), 7.59 (d, 2 H), 7.41 (d, 2 H), 7.39 (d, 1 H), 6.02 – 5.90 (m, 2 H), 5.61 (d, 1 H), 4.28 – 4.12 (m, 2 H), 3.62 – 3.50 (m, 2 H), 3.45 – 3.21 (m, 1 H), 3.32 (s, 3 H), 3.06 – 2.93 (m, 1 H), 2.57 – 2.25 (m, 2 H).

- 25           Example 864(3): (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(4-hydroxyphenyl)-1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone

- LC/MS  $t_R$  0.64 minutes; MS (ES<sup>+</sup>)  $m/z$  472 (M+H)<sup>f</sup>  
              <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.3 (s, 0.3 H), 11.9 (s, 0.7 H), 9.69 (s, 0.3 H), 9.67 (s, 0.7 H), 9.47 (s, 0.3 H), 9.24 (s, 0.7 H), 7.80 – 7.75 (m, 3 H), 7.51 – 7.37 (m, 2 H), 7.26  
30           (d, 0.7 H), 6.98 (d, 0.3 H), 6.79 – 6.67 (m, 2 H), 5.97 – 5.90 (m, 2 H), 5.62 – 5.55 (m, 1 H), 3.43 – 3.21 (m, 1 H), 3.02 – 2.91 (m, 1 H), 2.63 – 2.13 (m, 2 H).

Example 865(1) to Example 865(2)


The compounds of the present invention having the following physical data were prepared from the compound prepared in Example 9 using the corresponding alpha-bromoketones in the process of Example 51 → Example 52 → Example 44.

5 Example 865(1): (3S)-3-[4-chloro-5-(4-hydroxyphenyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

LC/MS  $t_R$  0.82 minutes; MS (ES<sup>+</sup>)  $m/z$  506 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.7 (s, 1 H), 9.70 (s, 1 H), 9.66 (s, 1 H), 7.85 - 7.77 (m, 3 H), 7.48 (d, 2 H), 6.84 (d, 2 H), 5.97 (s, 1 H), 5.92 (s, 1 H), 5.54 (dd, 1 H), 3.40 - 3.20 (m, 1 H), 3.02 - 2.89 (m, 1 H), 2.62 - 2.30 (m, 1 H), 2.24 - 2.14 (m, 1 H).


Example 865(2): ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-thiophenecarboxylate



15 TLC R<sub>f</sub> 0.52 (ethyl acetate)

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  9.34 (s, 1 H), 7.75 - 7.65 (m, 4 H), 7.37 (d, 1 H), 6.12 (d, 1 H), 6.08 (d, 1 H), 5.68 (dd, 1 H), 4.34 (q, 2 H), 3.50 - 3.30 (m, 1 H), 3.08 - 3.10 (m, 1 H), 2.70 - 2.58 (m, 1 H), 2.43 - 2.30 (m, 1 H), 1.37 (t, 3 H).

20 Example 867: 3-methoxypropyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate



The same operation as in Example 128 → Example 188 → Example 204 →

Example 51 → Example 52 → Example 338 was conducted from 5-amino-2-cyanopyridine to give the title compound having the following physical properties.

(Note: in the step corresponding to Example 128 in the operation, 3-methoxypropyl chloroformate was used. In the step corresponding to Example 51 in the operation, the compound prepared in Example 336 was used)

LC/MS  $t_R$  0.85 minutes; MS (ES<sup>+</sup>)  $m/z$  623 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 11.45 – 11.25 (m, 1 H), 8.82 (s, 1 H), 8.20 (d, 1 H), 8.13 – 8.03 (m, 1 H), 7.90 (d, 1 H), 7.70 – 7.60 (m, 2 H), 7.48 (d, 1 H), 7.20 – 7.12 (m, 1 H), 6.48 (s, 1 H), 5.85 – 5.79 (m, 1 H), 4.30 (t, 2 H), 3.57 – 3.40 (m, 3 H), 3.36 (s, 3 H), 3.16 – 3.03 (m, 1 H), 2.96 – 2.82 (m, 1 H), 2.60 – 2.42 (m, 1 H), 2.03 – 1.92 (m, 2 H).

Example 868: bis(2-methyl-2-propanyl) [5-(bromoacetyl)-3-fluoro-2-pyridinyl]imidodicarbonate

The same operation as in Example 90 → Example 228 → Example 361 was conducted from (5-iodo-3-fluoropyridin-2-yl)-amine to give the title compound having the following physical properties.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.87 (dd, 1H), 8.03 (dd, 1H), 4.42 (s, 2 H), 1.44 (s, 18 H).

Example 869: (3S)-3-[5-(6-amino-5-fluoro-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 51 → Example 52 → Example 338 → Example 55 was conducted from the compound prepared in Example 9 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 868 was used)

LC/MS  $t_R$  0.75 minutes; MS (ES<sup>+</sup>)  $m/z$  524 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.9 (s, 1H), 9.70 (s, 1 H), 8.09 (s, 1H), 7.86 – 7.74 (m, 3 H), 7.60 (dd, 1 H), 6.49 (s, 2 H), 5.97 (s, 1 H), 5.92 (s, 1 H), 5.52 (dd, 1 H), 3.30 – 3.17 (m, 1 H), 3.03 – 2.92 (m, 1 H), 2.60–2.45 (m, 1 H), 2.24 – 2.12 (m, 1 H).

Example 870: 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-fluorobenzoic acid hydrochloride

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 3-bromo-2-fluorobenzoic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.68 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>f</sup>

- 5 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 8.12 – 8.04 (m, 1 H), 7.88 – 7.71 (m, 5 H), 7.43 – 7.36 (m, 1 H), 6.02 (s, 1 H), 6.00 (s, 1 H), 5.80 – 5.72 (m, 1 H), 3.45 – 3.21 (m, 1 H), 3.17 – 2.98 (m, 1 H), 2.72 – 2.25 (m, 2 H).

10 Example 871: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-fluorobenzoic acid

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 5-bromo-2-fluorobenzoic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.65 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>f</sup>

- 15 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.69 (s, 1 H), 8.29 – 8.21 (m, 1 H), 8.07 – 7.93 (m, 2 H), 7.85 – 7.81 (m, 2 H), 7.76 – 7.72 (m, 1 H), 7.50 – 7.40 (m, 1 H), 6.03 (s, 1 H), 6.00 (s, 1 H), 5.78 – 5.70 (m, 1 H), 3.37 – 3.20 (m, 1 H), 3.15 – 3.00 (m, 1 H), 2.71 – 2.59 (m, 1 H), 2.42 – 2.28 (m, 1 H).

20 Example 872: 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-methylbenzoic acid hydrochloride

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 3-bromo-2-methylbenzoic acid to give the title compound having the following physical properties.

- 25 LC/MS  $t_R$  0.60 minutes; MS (ES<sup>+</sup>)  $m/z$  514 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.67 (s, 1 H), 7.86 – 7.76 (m, 4 H), 7.70 (d, 1 H), 7.56 (dd, 1 H), 7.42 (dd, 1 H), 6.04 (s, 1 H), 6.03 (s, 1 H), 5.84 – 5.75 (m, 1 H), 3.42 – 3.00 (m, 2 H), 3.79 – 2.23 (m, 5 H).

30 Example 873: 2-chloro-5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)benzoic acid hydrochloride

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-chloro-5-iodobenzoic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.69 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>f</sup>

- 5 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.68 (s, 1 H), 8.18 (d, 1 H), 8.13 (s, 1 H), 7.92 (dd, 1 H), 7.83 – 7.79 (m, 2 H), 7.73 – 7.65 (m, 2 H), 6.04 (s, 1 H), 6.01 (s, 1 H), 5.80 – 5.72 (m, 1 H), 3.39 – 3.21 (m, 1 H), 3.18 – 3.00 (m, 1 H), 2.74 – 2.23 (m, 2 H).

10 Example 874: 1-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)]-1H-imidazol-5-yl]phenyl]cyclopropanecarboxylic acid

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 1-(4-bromophenyl)cyclopropanecarboxylic acid to give the title compound having the following physical properties.

- 15 LC/MS  $t_R$  0.68 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>) δ 9.39 (s, 1 H), 7.83 (s, 1 H), 7.77 (dd, 1 H), 7.74 – 7.71 (m, 1 H), 7.79 – 7.63 (m, 3 H), 7.52 (d, 2 H), 6.18 (s, 1 H), 6.15 (s, 1 H), 5.94 – 5.85 (m, 1 H), 2.97 – 2.79 (m, 2 H), 2.52 – 2.36 (m, 2 H), 1.66 – 1.61 (m, 2 H), 1.27 – 1.21 (m, 2 H).

20

Example 875: 1-[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)]-1H-imidazol-5-yl]phenyl]cyclopropanecarboxylic acid

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 1-(3-bromophenyl)cyclopropanecarboxylic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.69 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>f</sup>

- 25 <sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>) δ 9.39 (s, 1 H), 7.87 (s, 1 H), 7.80 – 7.67 (m, 4 H), 7.63 – 7.57 (m, 1 H), 7.51 – 7.44 (m, 2 H), 6.18 (s, 1 H), 6.15 (s, 1 H), 5.93 – 5.86 (m, 1 H), 2.95 – 2.80 (m, 2 H), 2.55 – 2.38 (m, 2 H), 1.68 – 1.62 (m, 2 H), 1.31 – 1.24 (m, 2 H).
- 30

Example 876: 1-[3-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]phenyl]cyclopropanecarboxylic acid

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 338 → Example 363 was conducted from 1-(3-bromophenyl)cyclopropanecarboxylic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.87 minutes; MS (ES<sup>+</sup>)  $m/z$  574 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.69 – 8.56 (m, 1 H), 7.67 – 7.44 (m, 5 H), 7.39 – 7.19 (m, 2 H), 6.30 – 6.10 (m, 1 H), 5.93 – 5.73 (m, 2 H), 3.67 – 3.45 (m, 1 H), 3.12 – 2.85 (m, 1 H), 2.65 – 2.40 (m, 2 H), 1.80 – 1.60 (m, 2 H), 1.40 – 1.20 (m, 2 H).

Example 877: 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2,6-difluorobenzoic acid hydrochloride

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 3-bromo-2,6-difluorobenzoic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.65 minutes; MS (ES<sup>+</sup>)  $m/z$  536 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.67 (s, 1 H), 8.06 – 7.94 (m, 1 H), 7.82 – 7.71 (m, 3 H), 7.65 – 7.56 (m, 1 H), 7.34 – 7.24 (m, 1 H), 5.99 (s, 1 H), 5.98 (s, 1 H), 5.74 – 5.67 (m, 1 H), 3.39 – 3.22 (m, 1 H), 3.08 – 2.96 (m, 1 H), 2.75 – 2.25 (m, 2 H).

Example 878: 2-chloro-3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoic acid trifluoroacetate

The same operation as in Example 378 → Example 90 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 3-bromo-2-chlorobenzoic acid to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.64 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.68 (s, 1 H), 8.03 – 7.93 (m, 1 H), 7.85 – 7.74 (m, 4 H), 7.61 – 7.54 (m, 1 H), 7.47 (t, 1 H), 6.02 – 5.97 (m, 2 H), 5.75 – 5.67 (m, 1 H), 3.43 – 3.23 (m, 2 H), 3.11 – 2.97 (m, 2 H).

Example 879: (2E)-3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thienyl]acrylic acid hydrochloride

- 5        The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from (2E)-3-(4-bromo-2-thienyl)acrylic acid to give the title compound having the following physical properties.  
LC/MS  $t_R$  0.63 minutes; MS ( $ES^+$ )  $m/z$  532(M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.70 (s, 1 H), 8.23 (s, 1H), 7.97 (s, 1H), 7.95 (d, 1H),  
10    7.82 (s, 2H), 7.78 – 7.66 (m, 3H), 6.27 (d, 1H), 6.05 (s, 1 H), 5.99 (s, 1H), 5.83 – 5.79 (m, 1H), 3.39 – 3.24 (m, 1H), 3.18 – 3.04 (m, 1H), 2.79 – 2.63 (m, 1 H), 2.46 – 2.26 (m, 1 H).

Example 880: 3-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thienyl]propanoic acid hydrochloride

- 15        The same operation as in Example 378 → Example 139 → Example 21 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from (2E)-3-(4-bromo-2-thienyl)acrylic acid to give the title compound having the following physical properties.  
20    LC/MS  $t_R$  0.58 minutes; MS ( $ES^+$ )  $m/z$  535(M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 7.84 – 7.76 (m, 3H), 7.69 (s, 1H), 7.28 (s, 1H), 6.04 (s, 1H), 6.00 (s, 1 H), 5.79 – 5.75 (m, 1H), 3.35 – 3.20 (m, 1H), 3.18 – 3.04 (m, 1H), 3.04 (t, 1H), 2.76 – 2.63 (m, 1 H), 2.61 (t, 1H), 2.40 – 2.19 (m, 1 H).

25    Example 881: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)nicotinic acid dihydrochloride

- The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 5-bromonicotinic acid to give the title compound having the following physical properties.  
30    LC/MS  $t_R$  0.59 minutes; MS ( $ES^+$ )  $m/z$  501 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 9.21 (d, 1 H), 9.01 (d, 1 H), 8.65 (t, 1 H), 8.21 – 8.15 (m, 1 H), 7.84 – 7.80 (m, 2 H), 7.73 (s, 1 H), 6.03 (s, 1 H), 6.00 (s, 1 H), 5.81 – 5.73 (m, 1 H), 3.41 – 3.25 (m, 1 H), 3.18 – 3.01 (m, 2 H), 2.71 – 2.61 (m, 1 H).


Example 882: 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-1,3-thiazole-5-carboxylic acid trifluoroacetate

5 The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-bromo-1,3-thiazole-5-carboxylic acid to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)]) was used to give the title product as the trifluoroacetic acid salt)

10 LC/MS  $t_R$  0.67 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.34 (s, 1 H), 8.27 (s, 1H), 7.81 (s, 1H), 7.74 – 7.65 (m, 3 H), 6.14 (s, 1 H), 6.10 (s, 1H), 5.82 – 5.77 (m, 1 H), 3.51 – 3.40 (m, 1 H), 3.20 – 3.08 (m, 1H), 2.76 – 2.61 (m, 1 H), 2.60 – 2.46 (m, 1 H).

15 Example 883: 6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid trifluoroacetate



The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 6-bromo-2-pyridinecarboxylic acid to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)]) was used to give the title product as the trifluoroacetic acid salt)

20 LC/MS  $t_R$  0.62 minutes; MS (ES<sup>+</sup>)  $m/z$  501 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 8.13 – 8.01 (m, 3 H), 8.00 – 7.90 (m, 1 H), 7.86 – 7.72 (m, 3 H), 6.05 – 5.98 (m, 2 H), 5.84 – 5.72 (m, 1 H), 3.40 – 3.20 (m, 2



H), 3.15 – 2.98 (m, 2 H).

Example 884: 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1,3-thiazole-2-carboxylic acid

5 hydrochloride

The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 4-bromo-1,3-thiazole-2-carboxylic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.58 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>+</sup>

10 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.69 (s, 1 H), 8.46 (s, 1 H), 8.10 (s, 1 H), 7.87 – 7.78 (m, 2 H), 7.71 (s, 1 H), 6.03 (s, 1 H), 6.01 (s, 1 H), 5.86 – 5.75 (m, 1 H), 3.35 – 3.05 (m, 4 H).

Example 885: 4-chloro-3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)benzoic acid trifluoroacetate

15 The same operation as in Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 3-bromo-4-chlorobenzoic acid to give the title compound having the following physical properties.

(Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.70 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>+</sup>

20 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.37 (s, 1 H), 8.31 (s, 1H), 8.07 – 8.04 (m, 1H), 7.88 (s, 1H), 7.76 – 7.66 (m, 4H), 6.16 (s, 1 H), 6.14 (s, 1H), 5.94 – 5.89 (m, 1 H), 3.42 –  
25 3.10 (m, 2 H), 3.90 – 2.79 (m, 1H), 2.53 – 2.41 (m, 1 H).

Example 886: [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1-benzothiophen-3-yl]acetic acid trifluoroacetate

30 The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-methyl-2-propenyl (5-chloro-1-benzothiophen-3-yl)acetate to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B

(0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.67 minutes; MS ( $ES^+$ )  $m/z$  570(M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 8.19 (s, 1H), 8.14 – 8.09 (m, 2H), 7.82 (s, 2H), 7.79 – 7.70 (m, 3H), 6.06 (s, 1 H), 6.03 (s, 1H), 5.84 – 5.79 (m, 1 H), 3.91 (s, 2H), 3.36 – 3.22 (m, 1H), 3.18 – 3.05 (m, 1H), 2.78 – 2.63 (m, 1 H), 2.43– 2.29 (m, 1 H).

10 Example 887: [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenoxy]acetic acid trifluoroacetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-methyl-2-propanyl (3-iodophenoxy)acetate [patent EP1386913] to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 15 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.59 minutes; MS ( $ES^+$ )  $m/z$  530 (M+H)<sup>f</sup>

20 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.38 (s, 1 H), 7.76 – 7.70 (m, 4 H), 7.63 (d, 2H), 7.07 (d, 2H), 6.16 (s, 1 H), 6.14 (s, 1H), 5.90 – 5.85 (m, 1 H), 3.35 - 3.17 (m, 2 H), 2.94 – 2.79 (m, 1 H), 2.49 – 2.35 (m, 1 H).

Example 888: [3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenoxy]acetic acid trifluoroacetate

25 The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-methyl-2-propanyl (4-iodophenoxy)acetate [patent EP1386913] to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A 30 (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.62 minutes; MS ( $ES^+$ )  $m/z$  530 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.38 (s, 1 H), 7.85 (s, 1H), 7.76 – 7.67 (m, 3 H), 7.42

(t, 1H), 7.30 (d, 1H), 7.29 (d, H), 7.04 (dd, 1H), 6.16 (s, 1 H), 6.15 (s, 1H), 5.92 – 5.86 (m, 1 H), 3.41 - 3.17 (m, 2 H), 2.94 – 2.79 (m, 1 H), 2.49 – 2.39 (m, 1 H).

5 Example 889: 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1,3-thiazole-4-carboxylic acid trifluoroacetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-methyl-2-propanyl 2-bromo-1,3-thiazole-4-carboxylate [patent WO2008/54701] to give the title compound having the  
10 following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.69 minutes; MS (ES<sup>+</sup>)  $m/z$  507 (M+H)<sup>f</sup>  
15 <sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.35 (s, 1 H), 8.31 (s, 1H), 7.90 (s, 1H), 7.75 – 7.65 (m, 3 H), 6.15 (s, 1 H), 6.09 (s, 1H), 5.83 – 5.79 (m, 1 H), 3.48 – 3.37 (m, 1 H), 3.20 – 3.08 (m, 1H), 2.78 – 2.62 (m, 1 H), 2.57 – 2.43 (m, 1 H).

20 Example 890: 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-4-fluorobenzoic acid trifluoroacetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 2-methyl-2-propanyl 3-bromo-4-fluorobenzoate [patent WO2010/37210] to give the title compound having the following  
25 physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.69 minutes; MS (ES<sup>+</sup>)  $m/z$  518 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.66 (s, 1 H), 8.51 – 8.47 (m, 1H), 7.90 – 7.82 (m, 1H), 7.78 (s, 1H), 7.75 (s, 1 H), 7.63 – 7.59 (m, 1H), 7.43 – 7.37 (m, 1H), 6.01 (s, 1 H),  
30 5.96 (s, 1H), 5.71 – 5.68 (m, 1 H), 3.40 – 3.27 (m, 1 H), 3.13 – 3.02 (m, 1H), 2.71 – 2.24 (m, 2 H).

Example 891: methyl (6-(((trifluoromethyl)sulfonyl)oxy)-1-benzothiophen-3-yl)acetate

To a solution of methyl (6-hydroxy-1-benzothiophen-3-yl)acetate (883 mg) and triethylamine (1.1 mL) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added trifluoromethanesulfonic anhydride (1.23 g) at 0 °C. After being stirred for 25 min at 0 °C, the reaction mixture was poured into a saturated aqueous solution of ammonium chloride. The mixture was extracted with dichloromethane. The organic layers were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified by column chromatography (10 – 20% ethyl acetate in hexane) to give the title compound having the following physical properties (1.16 g). TLC: R<sub>f</sub> 0.76(30% ethyl acetate in hexane).

10 Example 892: [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1-benzothiophen-3-yl]acetic acid trifluoroacetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 342 was conducted from the compound prepared in Example 891 to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS *t<sub>R</sub>* 0.64 minutes; MS (ES<sup>+</sup>) *m/z* 570(M+H)<sup>+</sup>

20 <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.68 (s, 1 H), 8.36 (s, 1H), 8.07 (s, 1H), 7.89 (d, 1H), 7.84 – 7.71 (m, 3H), 7.70 (d, 1H), 7.67 (s, 1H), 6.05 (s, 1 H), 6.02 (s, 1H), 5.83 – 5.78 (m, 1 H), 3.88 (s, 2H), 3.37 – 3.22 (m, 1H), 3.18 – 3.06 (m, 1H), 2.78 – 2.63 (m, 1 H), 2.43– 2.24 (m, 1 H).


25 Example 893: 2-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)isonicotinic acid trifluoroacetate

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 342 was conducted from methyl 2-bromoisonicotinate to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS *t<sub>R</sub>* 0.60 minutes; MS (ES<sup>+</sup>) *m/z* 501 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.68 (s, 1 H), 8.75 (d, 1 H), 8.28 (s, 1 H), 8.17 – 8.05 (m, 1 H), 7.91 – 7.69 (m, 4 H), 6.01 (s, 1 H), 5.98 (s, 1 H), 5.79 – 5.70 (m, 1 H), 3.40 – 3.23 (m, 1 H), 3.14 – 2.98 (m, 2 H), 2.65 – 2.30 (m, 1 H).

- 5 Example 894: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-3-thiophenecarboxylic acid trifluoroacetate



The same operation as in Example 90 → Example 361 → Example 51 → Example 52 → Example 342 was conducted from methyl 5-iodo-3-

- 10 thiophenecarboxylate to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt) LC/MS *t<sub>R</sub>* 0.73 minutes; MS (ES<sup>+</sup>) *m/z* 506 (M+H)<sup>+</sup>
- 15 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.68 (s, 1 H), 8.25 (s, 1 H), 7.92 (s, 1 H), 7.83 – 7.75 (m, 3 H), 7.71 (s, 1 H), 6.03 (s, 1 H), 5.99 (s, 1 H), 5.73 (dd, 1 H), 3.40 – 3.20 (m, 1 H), 3.14 – 2.98 (m, 1 H), 2.65 – 2.25 (m, 2 H).

Example 895: 2-methyl-2-propanyl 3-acetyl-1H-pyrazole-5-carboxylate

- 20 To a stirred solution of indium (III) chloride (22 mg) in water (3 mL) was added 3-butyne-2-one (340 mg) and tert-butyl diazoacetate (710 mg) at room temperature. After being stirred at room temperature for 1 hour, the resulting precipitate was collected by filtration, washed with water and dried in vacuo to give the title compound having the following physical properties (1.05 g).
- 25 TLC: R<sub>f</sub> 0.74 (50% ethyl acetate in hexane).

Example 896: 3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-1H-pyrazole-5-carboxylic acid

The same operation as in Example 361 → Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 895 to give the title compound having the following physical properties.

LC/MS  $t_R$  0.59 minutes; MS (ES<sup>+</sup>)  $m/z$  490 (M+H)<sup>f</sup>

- 5 <sup>1</sup>H NMR (300 MHz, methanol-d<sub>4</sub>) δ 9.34 (s, 1 H), 7.78-7.62 (m, 3 H), 7.30 (s, 1 H), 6.87 (s, 1 H), 6.13 (s, 1 H), 6.07 (s, 1 H), 5.78 (dd, 1 H), 3.40 - 3.02 (m, 2 H), 2.72 - 2.40 (m, 2 H).

10 Example 897: 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-indole-2-carboxylic acid trifluoroacetate

The same operation as in Example 378 → Example 139 → Example 379 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 5-bromo-1H-indole-2-carboxylic acid to give the title compound having the following

15 physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.59 minutes; MS (ES<sup>+</sup>)  $m/z$  539(M+H)<sup>f</sup>

- 20 <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 11.56 (s, 1H), 9.39 (s, 1 H), 7.96 (d, 1H), 7.99 (s, 1H), 7.76 - 7.66 (m, 4H), 7.57 (s, 2H), 7.21 (s, 1H), 6.17 (s, 1 H), 6.15 (s, 1H), 5.94 - 5.88 (m, 1 H), 3.43 - 3.18 (m, 2 H), 2.94 - 2.80 (m, 1 H), 2.52 - 2.39 (m, 1 H).

25 Example 898: 5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo [1,2-a]pyrimidin-6-yl])-1H-imidazol-5-yl)-1H-indole-2-carboxylic acid hydrochloride

The same operation as in Example 378 → Example 139 → Example 379 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from 5-bromo-1H-indole-2-carboxylic acid to give the title compound having the following

30 physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 336 was used)

LC/MS  $t_R$  0.58 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 12.05 (s, 1 H), 9.73 (s, 1 H), 8.09 (s, 1 H), 8.04 (s, 1 H), 7.94 (d, 1 H), 7.91 - 7.79 (m, 2 H), 7.69 - 7.62 (m, 1 H), 7.58 - 7.51 (m, 1 H), 7.17 (s,

1 H), 6.41 (s, 1 H), 5.87 (dd, 1 H), 3.21 - 3.04 (m, 1 H), 3.01 - 2.83 (m, 1 H), 2.81 - 2.61 (m, 1 H) 2.42 - 2.28 (m, 1 H).

Example 899: [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1H-indol-3-yl]acetic acid hydrochloride

The same operation as in Example 378 → Example 379 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from (5-bromo-1H-indol-3-yl)acetic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.60 minutes; MS (ES<sup>+</sup>)  $m/z$  553 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  11.23 (s, 1H), 9.69 (s, 1 H), 7.96 (d, 1H), 7.95 (s, 1H), 7.82 (s, 2H), 7.71 (s, 1H), 7.48 (s, 2H), 7.34 (s, 1H), 6.06 (s, 1 H), 6.03 (s, 1H), 5.84 – 5.79 (m, 1 H), 3.70 (s, 2H), 3.39 – 3.22 (m, 1 H), 3.18 – 3.05 (m, 1H), 2.80 – 2.64 (m, 1 H), 2.43 – 2.28 (m, 1 H).

Example 900: [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1H-indol-3-yl]acetic acid hydrochloride

The same operation as in Example 378 → Example 379 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from (6-bromo-1H-indol-3-yl)acetic acid to give the title compound having the following physical properties.

LC/MS  $t_R$  0.62 minutes; MS (ES<sup>+</sup>)  $m/z$  553 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  11.3 (s, 1 H), 9.69 (s, 1 H), 8.03 (s, 1 H), 7.86-7.75 (m, 3 H), 7.71 (d, 1 H), 7.62 (d, 1 H), 7.40 (d, 1 H), 7.36 (d, 1 H), 6.06 (s, 1 H), 6.04 (s, 1 H), 5.82 (dd, 1 H), 3.67 (s, 2 H), 3.28 (m, 1 H), 3.14 (m, 1 H), 2.72 (m, 1 H), 2.38 (m, 1 H).

Example 901: [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-1H-indazol-3-yl]acetic acid

dihydrochloride

The same operation as in Example 379 → Example 378 → Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from (5-

bromo-1H-indazol-3-yl)acetic acid to give the title compound having the following physical properties.


LC/MS  $t_R$  0.56 minutes; MS (ES<sup>+</sup>)  $m/z$  554 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  14.66 (br s, 1 H), 13.2 (br s, 1 H), 9.70 (s, 1 H), 8.18 – 8.16 (m, 1 H), 8.05 (s, 1 H), 7.86 – 7.78 (m, 2 H), 7.75 (dd, 1 H), 7.71 (dd, 1 H), 7.64 (dd, 1 H), 6.06 (s, 1 H), 6.03 (s, 1 H), 5.82 (dd, 1 H), 3.95 (s, 2 H), 3.39 – 3.22 (m, 1 H), 3.20 – 3.03 (m, 1 H), 2.80 – 2.62 (m, 1 H), 2.44 – 2.31 (m, 1 H).

Example 902: tert-butyl 2-(3-acetylanilino)acetate

To a solution of 1-(3-aminophenyl)ethanone (600 mg) in acetone (20 mL) were added potassium carbonate (1.23 g) and tert-butyl 2-bromoacetate (1.73 g). The mixture was refluxed overnight then cooled to room temperature. The insoluble material was removed by filtration. The filtrate was concentrated. The residue was purified by column chromatography (10 – 20% ethyl acetate in hexane) to give the title compound having the following physical properties (1.10 g).  
TLC: R<sub>f</sub> 0.62 (20% ethyl acetate in hexane).

Example 903: {[3-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)phenyl]amino}acetic acid dihydrochloride



The same operation as in Example 379 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 902 to give the title compound having the following physical properties.

LC/MS  $t_R$  0.57 minutes; MS (ES<sup>+</sup>)  $m/z$  529 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.68 (s, 1 H), 7.98 (s, 1 H), 7.81 (s, 2 H), 7.69 (d, 1 H), 7.63 – 7.45 (m, 1 H), 7.18 (d, 1 H), 7.06 – 6.88 (m, 2 H), 6.67 (d, 1 H), 6.04 (s, 1 H), 6.00 (s, 1 H), 5.88 – 5.72 (m, 1 H), 3.88 (s, 2 H), 3.34 – 3.18 (m, 1 H), 3.16 – 2.98 (m, 1 H).



H), 2.77 - 2.60 (m, 1 H) 2.43 - 2.28 (m, 1 H).

Example 904: 2-(1H-indol-5-yl)-2-oxoethyl (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylate

- 5        The same operation as in Example 361 → Example 51 → Example 363 was conducted from 2-methyl-2-propanyl 5-acetyl-1H-indole-1-carboxylate [patent WO2005/26175] to give the title compound having the following physical properties. TLC: R<sub>f</sub> 0.47 (20% ethyl acetate in hexane).

- 10    Example 905: 2-(1-{2-[(2-methyl-2-propanyl)oxy]-2-oxoethyl}-1H-indol-5-yl)-2-oxoethyl (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinecarboxylate

- 15        To a solution of the compound prepared in Example 904 (140 mg) in N,N-dimethylformamide (3 mL) were added potassium carbonate (792 mg) and tert-butyl 2-bromoacetate (395 mg) at room temperature. After being stirred for 1 hour at room temperature, the reaction mixture was poured into a saturated aqueous solution of ammonium chloride. The mixture was extracted with ethyl acetate. The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified by column chromatography (0 – 10% methanol in ethyl acetate) to give the title compound  
20    having the following physical properties (99 mg).  
TLC: R<sub>f</sub> 0.53 (ethyl acetate).

Example 906: [5-(2-{(3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyloxy}-1H-imidazol-5-yl)-1H-indol-1-yl]acetic acid

- 25        The same operation as in Example 52 → Example 363 was conducted from the compound prepared in Example 905 to give the title compound having the following physical properties.  
LC/MS *t<sub>R</sub>* 0.63 minutes; MS (ES<sup>+</sup>) *m/z* 553 (M+H)<sup>+</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.69 (s, 1 H), 7.88-7.02 (m, 8 H), 6.35 (brs, 1 H),  
30    5.97 (brs, 1 H), 5.94 (s, 1 H), 5.63 (d, 1 H), 4.57 (s, 2 H), 3.53-2.85 (m, 2 H), 2.62-2.38 (m, 2 H).

Example 907: 2-methyl-2-propanyl (5-bromo-1H-benzimidazol-1-yl)acetate

To a stirred suspension of sodium hydride (487 mg) in N,N-dimethylformamide (10 mL) was added a solution of 5-bromo-1H-benzimidazole (2 g) in N,N-dimethylformamide (5 mL) at room temperature. To the resulting mixture was added tert-butyl 2-bromoacetate (2.38 g) at room temperature. After being stirred for 20 minutes, the reaction mixture was poured into a saturated aqueous solution of ammonium chloride. The mixture was extracted with ethyl acetate. The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified by column chromatography (0 – 50% ethyl acetate in hexane) to give the title compound having the following physical properties (3.16 g).

TLC: R<sub>f</sub> 0.59 (ethyl acetate).


Example 908: [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-1H-benzimidazol-1-yl]acetic acid dihydrochloride

The same operation as in Example 139 → Example 361 → Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 907 to give the title compound having the following physical properties.

LC/MS *t*<sub>R</sub> 0.55 minutes; MS (ES<sup>+</sup>) *m/z* 554 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.70 (s, 1 H), 9.22 (brs, 1 H), 8.31 (s, 1 H), 8.17 (s, 1 H), 8.06-7.92 (m, 2 H), 7.85-7.78 (m, 2 H), 7.71 (s, 1 H), 6.07 (s, 1 H), 6.01 (s, 1 H), 5.86 (dd, 1 H), 5.38 (s, 2 H), 3.32 (m, 1 H), 3.12 (m, 1 H), 2.72 (m, 1 H), 2.42 (m, 1 H).

Example 909: 5-(2-((3S)-6-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride



The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 383(2) to give the title compound

having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 374 was used).

LC/MS  $t_R$  0.64 minutes; MS (ES<sup>+</sup>)  $m/z$  540 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.73–9.63 (m, 1 H), 7.91–7.85 (m, 2 H), 7.78–

5 7.61 (m, 3 H), 7.41–7.31 (m, 1 H), 6.43–6.35 (m, 1 H), 5.79–5.68 (m, 1 H), 3.80–3.05 (m, 4 H).


Example 910: 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid hydrochloride

The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 336 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 384 was used)

15 LC/MS  $t_R$  0.55 minutes; MS (ES<sup>+</sup>)  $m/z$  502 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.73 (s, 1 H), 8.78 – 8.71 (m, 1 H), 8.53 (s, 1 H), 8.47 (s, 1 H), 8.21 (d, 1 H), 7.95 (s, 1 H), 7.91 – 7.80 (m, 2 H), 6.36 (s, 1 H), 5.82 – 5.71 (m, 1 H), 3.24 – 3.05 (m, 2 H), 2.95 – 2.78 (m, 2 H).

20 Example 911: 4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-  
4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl})-1H-imidazol-5-yl)-2-  
pyridinecarboxylic acid hydrochloride




The same operation as in Example 51 → Example 52 → Example 338 →  
25 Example 363 was conducted from the compound prepared in Example 336 to give the  
title compound having the following physical properties. (Note: in the step  
corresponding to Example 51 in the operation, the compound prepared in Example 384  
was used)

LC/MS  $t_R$  0.59 minutes; MS (ES<sup>+</sup>)  $m/z$  536 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.74 (s, 1 H), 8.78 (d, 1 H), 8.53 (d, 1 H), 8.13 (dd, 1 H), 7.98 (d, 1 H), 7.84 (dd, 1 H), 7.81 (d, 1 H), 6.34 (s, 1 H), 5.63 (dd, 1 H), 3.13 – 3.03 (m, 1 H), 2.85 – 2.73 (m, 1 H), 2.65 – 2.52 (m, 1 H), 2.26 – 2.15 (m, 1 H).

5

Example 912: 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid trifluoroacetate



10

The same operation as in Example 51 → Example 52 → Example 338 →

Example 363 was conducted from the compound prepared in Example 384 to give the title compound having the following physical properties. (Note: high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

15

LC/MS  $t_R$  0.63 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.70 (s, 1 H), 8.75 (d, 1 H), 8.46 (d, 1 H), 7.98 (dd, 1 H), 7.83 – 7.76 (m, 3 H), 5.99 (s, 1 H), 5.94 (s, 1 H), 5.59 (dd, 1 H), 3.38 – 3.20 (m, 1 H), 3.07 – 2.91 (m, 1 H), 2.50 – 2.30 (m, 1 H), 2.28 – 2.16 (m, 1 H).

20

Example 913: 4-(4-chloro-2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl)-1H-imidazol-5-yl)-2-pyridinecarboxylic acid trifluoroacetate

The same operation as in Example 51 → Example 52 → Example 338 →

25


Example 363 was conducted from the compound prepared in Example 383(1) to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 384 was used. High performance liquid chromatography [5 to 100% mobile phase B (0.1%

trifluoroacetic acid in acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] was used to give the title product as the trifluoroacetic acid salt)

LC/MS  $t_R$  0.69 minutes; MS (ES<sup>+</sup>)  $m/z$  569 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.74 – 9.72 (m, 1 H), 8.77 – 8.75 (m, 1 H), 8.45 – 8.43 (m, 1 H), 8.00 – 7.95 (m, 1 H), 7.89 – 7.68 (m, 3 H), 6.32 – 6.25 (m, 1 H), 5.74 – 5.68 (m, 1 H), 3.40 – 3.21 (m, 1 H), 3.19 – 3.03 (m, 1 H), 2.77 – 2.60 (m, 1 H), 2.42 – 2.39 (m, 1 H).

Example 914: 4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride



The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 756 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 361 was used)

LC/MS  $t_R$  0.68 minutes; MS (ES<sup>+</sup>)  $m/z$  539 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.70 (s, 1 H), 8.29 (s, 1 H), 8.17 (s, 1 H), 8.05 (s, 1 H), 7.78 (dd, 1 H), 7.73 (d, 1 H), 7.68 (d, 1 H), 6.10 (s, 1 H), 5.82 (s, 1 H), 5.79 (dd, 1 H), 3.30 (m, 1 H), 3.14 (m, 1 H), 2.72 (m, 1 H), 2.36 (m, 1 H).

Example 915: 4-(2-((3S)-7-[5-chloro-2-(4-chloro-1H-1,2,3-triazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-2-pyridinecarboxylic acid hydrochloride

The same operation as in Example 51 → Example 52 → Example 363 was conducted from the compound prepared in Example 756 to give the title compound having the following physical properties. (Note: in the step corresponding to Example

51 in the operation, the compound prepared in Example 384 was used)

LC/MS  $t_R$  0.65 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.73 (d, 1 H), 8.70 (s, 1 H), 8.52 (d, 1 H), 8.44 (s, 1 H), 8.18 (dd, 1 H), 7.78 (dd, 1 H), 7.77-7.68 (m, 2 H), 6.04 (s, 1 H), 5.96 (s, 1 H), 5.73 (dd, 1 H), 3.38 (m, 1 H), 3.08 (m, 1 H), 2.68-2.25 (m, 2 H).

Example 916: 5-chloro-2-(1,2,3-thiadiazol-4-yl)phenol

The same operation as in the literature [Tetrahedron 56 (24), 3933 (2000)] was conducted from 1-(4-chloro-2-hydroxyphenyl)ethanone to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 384 was used) TLC: R<sub>f</sub> 0.74 (50% ethyl acetate in hexane).


Example 917: (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1,2,3-thiadiazol-4-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone

The same operation as in Example 6 → Example 283 → Example 8 → Example 51 → Example 52 → Example 10 → Example 338 → Example 363 was conducted from the compound prepared in Example 916 to give the title compound having the following physical properties. (Note: in the step corresponding to Example 51 in the operation, the compound prepared in Example 193 was used)

LC/MS  $t_R$  0.71 minutes; MS (ES<sup>+</sup>)  $m/z$  522 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.72 (brs, 1 H), 9.04 (d, 1 H), 8.20 (d, 1 H), 7.86 (d, 1 H), 7.73 – 7.57 (m, 3 H), 6.51 (d, 1 H), 6.18 (brs, 2 H), 5.98 (d, 1 H), 5.56 (d, 1 H), 3.45 - 3.14 (m, 1 H), 3.04 - 2.88 (m, 1 H), 2.60 - 2.34 (m, 1 H), 2.29 - 2.12 (m, 1 H).

Example 918: methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate




The same operation as in Example 370 was conducted from the compound prepared in Example 392 to give the title compound having the following physical properties.

- 5 LC/MS  $t_R$  0.71 minutes; MS ( $ES^+$ )  $m/z$  521 ( $M+H$ )<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, Methanol- $d_4$ )  $\delta$  9.44 (s, 1 H), 8.04 (d, 1 H), 7.91 (d, 1 H), 7.82 (d, 1 H) 7.79 - 7.73 (m, 1 H) 7.72 - 7.66 (m, 1 H) 7.40 - 7.34 (m, 1 H), 6.34 (s, 1 H), 5.71 (dd, 1 H), 3.87 (s, 3 H), 3.27 - 3.20 (m, 1 H), 3.02 - 2.84 (m, 1 H), 2.77 - 2.55 (m, 1 H), 2.49 - 2.29 (m, 1 H).

10

Example 919(1) and Example 919(2) : 2-oxo-2-(1-pyrrolidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate trifluoroacetate and 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-N,N-dimethyl-2-thiophenecarboxamide

15



To a solution of the compound prepared in Example 392 (100 mg), 2-hydroxy-1-pyrrolidin-1-yl-ethanone (48 mg), and N,N-diisopropylethylamine (0.080 mL) in N,N-dimethylformamide (2 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-

- 20 tetramethyluronium hexafluorophosphate (84 mg) at room temperature. After being stirred at 40 °C for 15 hours, the reaction mixture was poured into saturated aqueous solution of sodium hydrogen carbonate and extracted with ethyl acetate. The combined organic layer was dried and concentrated. The residue was purified by high performance liquid chromatography [5 to 100% mobile phase B (0.1% trifluoroacetic acid in

acetonitrile) in mobile phase A (0.1% aqueous trifluoroacetic acid)] to give the title compounds having the following physical properties.

Example 919(1): 48 mg


- 5 LC/MS  $t_R$  0.71 minutes; MS (ES<sup>+</sup>)  $m/z$  618 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  9.72 (s, 1 H), 8.16 (s, 1 H), 8.07 (s, 1 H), 7.95 (d, 1 H), 7.92-7.79 (m, 2 H), 7.76 (brs, 1H), 6.37 (s, 1 H), 5.68 (dd, 1 H), 4.91 (s, 2 H), 3.50-3.02 (m, 5 H), 2.82 (m, 1 H), 2.60 (m, 1 H), 2.24 (m, 1 H), 1.96-1.92 (m, 2 H), 1.82-1.70 (m, 2 H).

10

Example 919(2): 20 mg.

- LC/MS  $t_R$  0.61 minutes; MS (ES<sup>+</sup>)  $m/z$  534 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.7-10.4 (m, 1 H), 8.78 (m, 1 H), 7.72-7.35 (m, 5 H), 7.18-7.08 (m 1 H), 6.42-6.37 (m, 1 H), 5.75 (m, 1 H), 3.48-3.00 (m, 8 H), 2.92 (m, 1 H),  
 15 2.48(m, 1 H).

Example 920(1) to Example 920(9)



- The compounds of the present invention having the following physical data were  
 20 synthesised from the compound prepared in Example 392 using the corresponding  
 alcohols, employing the method as detailed in Example 367.

Example 920(1): Ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate

25

- LC/MS  $t_R$  0.77 minutes; MS (ES<sup>+</sup>)  $m/z$  535 (M+H)<sup>f</sup>  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.78 (s, 1 H), 8.00 (s, 1 H), 7.72 – 7.62 (m, 3 H), 7.47 (d,




1 H), 7.18 (s, 1 H), 6.39 (s, 1 H), 5.76 (d, 1 H), 4.35 (q, 2 H), 3.41 – 3.35 (m, 2 H), 2.96 – 2.87 (m, 1 H), 2.53 – 2.47 (m, 1 H), 1.39 (t, 3 H).

Example 920(2): Isobutyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-  
 5 4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-  
thiophenecarboxylate

LC/MS  $t_R$  0.88 minutes; MS (ES<sup>+</sup>)  $m/z$  563 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.78 (s, 1 H), 7.98 (s, 1 H), 7.68 – 7.61 (m, 3 H), 7.47 (d, 1 H), 7.18 (s, 1 H), 6.39 (s, 1 H), 5.76 (d, 1 H), 4.09 (d, 2 H), 3.43 – 3.35 (m, 2 H), 2.96 – 2.91 (m, 1 H), 2.54 – 2.47 (m, 1 H), 2.12 – 2.07 (m, 1 H), 1.01 (d, 6 H).


Example 920(3): 2,3-dihydro-1H-inden-5-yl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-  
yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-  
thiophenecarboxylate



LC/MS  $t_R$  0.99 minutes; MS (ES<sup>+</sup>)  $m/z$  623 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.39 (s, 1H), 8.79 (s, 1 H), 8.16 (s, 1H), 7.79 (s, 1H), 7.68 (d, 1H), 7.67 – 7.62 (m, 1H), 7.47 (d, 1H), 7.21 (s, 1H), 7.06 (s, 1H), 6.97 – 6.94 (m, 1 H), 6.39 (s, 1H), 5.76 (d, 1H), 3.47 – 3.35 (m, 2H), 2.92 (q, 4H), 2.58 – 2.43 (m, 1H), 2.18 – 2.06 (m, 3H).

Example 920(4): (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((6S)-2-[5-chloro-2-  
(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-  
imidazol-5-yl)-2-thiophenecarboxylate



LC/MS  $t_R$  0.76 minutes; MS (ES<sup>+</sup>)  $m/z$  619 (M+H)<sup>f</sup>


<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.7-10.3 (m, 1 H), 8.79 (m, 1 H), 8.05-7.90 (m, 1 H), 7.81-7.43 (m, 4 H), 7.22-7.14 (m, 1 H), 6.42-6.37 (m, 1 H), 5.80-5.70 (m, 1 H), 5.06 (s, 2 H), 3.46-3.25 (m, 2 H), 2.93 (m, 1 H), 2.50 (m, 1 H), 2.25 (m, 3 H).

Example 920(5): 2-(4-morpholinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate

LC/MS  $t_R$  0.56 minutes; MS (ES<sup>+</sup>)  $m/z$  620 (M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.7-10.4 (m, 1 H), 8.78 (m, 1 H), 8.01-7.88 (m, 1 H), 7.75-7.59 (m, 3 H), 7.47 (m, 1 H), 7.21-7.12 (m, 1 H), 6.42-6.37 (m, 1 H), 5.80-5.70 (m, 1 H), 4.46-4.40 (m, 2 H), 3.72 (t, 4 H), 3.48-3.18 (m, 2 H), 2.92 (m, 1 H), 2.76 (t, 2 H), 2.57 (t, 4 H), 2.51 (m, 1 H).

Example 920(6): 2-(dimethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate



LC/MS  $t_R$  0.66 minutes; MS (ES<sup>+</sup>)  $m/z$  592 (M+H)<sup>f</sup>


<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.4 (brs, 1 H), 8.79 (m, 1 H), 8.04 (brs, 1 H), 7.75 (brs, 1 H), 7.67 (d, 1 H), 7.62 (dd, 1 H), 7.47 (d, 1 H), 7.14 (s, 1 H), 6.38 (s, 1 H), 5.75 (d, 1 H), 4.94 (s, 2 H), 3.39 (m, 1 H), 3.21 (m, 1 H), 3.04 (s, 3 H), 3.00 (s, 3 H), 2.93 (m, 1 H), 2.51 (m, 1 H).

Example 920(7): 2-(diethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate

TLC R<sub>f</sub> 0.42 (10% methanol in ethyl acetate)

- 5 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.36 (br. s, 1 H), 8.78 (s, 1 H), 8.16 – 7.93 (m, 1 H), 7.83 – 7.55 (m, 3 H), 7.47 (d, 1 H), 7.21 – 7.09 (m, 1 H), 6.50 – 6.29 (m, 1 H), 5.74 (d, 1 H), 4.93 (s, 2 H), 3.65 – 3.22 (m, 6 H), 3.22 – 3.65 (m, 6 H), 3.01 – 2.83 (m, 1 H), 2.64 – 2.37 (m, 1 H), 1.26 (t, 3 H), 1.15 (t, 3 H).

- 10 Example 920(8): 2-oxo-2-(1-piperidiny)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate trifluoroacetate



LC/MS *t*<sub>R</sub> 0.76 minutes; MS (ES<sup>+</sup>) *m/z* 632 (M+H)<sup>+</sup>

- 15 <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.72 (s, 1 H), 8.20 (s, 1 H), 8.15 (s, 1 H), 7.98-7.80 (m, 4 H), 6.41 (s, 1 H), 5.73 (dd, 1 H), 5.03 (s, 2 H), 3.46-3.25 (m, 4 H), 3.12 (m, 1 H), 2.83 (m, 1 H), 2.65 (m, 1 H), 2.28 (m, 1 H), 1.65-1.12 (m, 6 H).

- 20 Example 920(9): 2-(4-morpholinyl)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate

LC/MS *t*<sub>R</sub> 0.65 minutes; MS (ES<sup>+</sup>) *m/z* 634 (M+H)<sup>+</sup>

- <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>) δ 9.72 (s, 1 H), 8.11 (s, 1H), 7.98 (s, 1H), 7.95 (d, 1H), 7.86 – 7.78 (m, 2H), 7.60 (s, 1H), 6.35 (s, 1H), 5.65 – 5.61 (m, 1H), 5.03 (s, 2H), 3.62 – 3.54 (m, 2H), 3.44 – 3.40 (m, 2H), 3.21 – 3.10 (m, 1H), 2.82 – 2.71 (m, 1H), 2.62 – 2.50 (m, 1H), 2.26 – 2.19 (m, 1H).
- 25


Example 921: Isopropyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate

The same operation as in Example 367 was conducted from the compound prepared in Example 392 and 2-propanol to give the title compound having the following physical properties.

LC/MS  $t_R$  0.82 minutes; MS (ES<sup>+</sup>)  $m/z$  549 (M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.78 (s, 1 H), 7.96 (s, 1 H), 7.68 – 7.61 (m, 3 H), 7.47 (d, 1 H), 7.17 (s, 1 H), 6.39 (s, 1 H), 5.27 – 5.18 (m, 1 H), 3.42 – 3.35 (m, 2 H), 2.96 – 2.87 (m, 1 H), 2.54 – 2.47 (m, 1 H), 1.37 (d, 6 H).

Example 922(1) and Example 922(2): 1-(((cyclohexyloxy)carbonyl)oxy)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate and cyclohexyl 2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-5-{5-[(1-(((cyclohexyloxy)carbonyl)oxy)ethoxy)carbonyl]-3-thienyl}-1H-imidazole-1-carboxylate



The same operation as in Example 373 was conducted from the compound prepared in Example 392 and 1-chloroethyl cyclohexyl carbonate to give the title compounds having the following physical properties.

Example 922(1):


LC/MS  $t_R$  0.99 minutes; MS (ES<sup>+</sup>)  $m/z$  677(M+H)<sup>+</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.41 (s, 1 H), 8.79 (s, 1 H), 8.02 (s, 1 H), 7.75 (s, 1 H), 7.67 (d, 1 H), 7.65 – 7.61 (m, 1 H), 7.47 (d, 1 H), 7.17 (s, 1 H), 6.98 (q, 1 H), 6.38 (s, 1 H), 5.75 (d, 1 H), 4.69 – 4.60 (m, 1 H), 3.43 – 3.33 (m, 2 H), 2.96 – 2.85 (m, 1 H), 2.60 – 2.43 (m, 1 H), 1.98 – 1.90 (m, 2 H), 1.81 – 1.71 (m, 2 H), 1.64 (d, 3 H), 1.60 – 1.21 (m, 6 H).

Example 922(2):LC/MS  $t_R$  1.24 minutes; MS ( $ES^+$ )  $m/z$  803(M+H)<sup>f</sup>

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  8.78 (s, 1 H), 7.96 (d, 1H), 7.88 (d, 1H), 7.74 (d, 1H),  
 7.66 – 7.62 (m, 1H), 7.55 (s, 1H), 7.52 (d, 1H), 7.52 (s, 1H), 6.97 (q, 1 H), 6.42 (s, 1H),  
 5 6.35 – 6.32 (m, 1H), 5.08 – 4.98 (m, 1H), 4.69 – 4.60 (m, 1H), 3.20 – 3.08 (m, 1H), 2.80  
 – 2.77 (m, 1H), 2.69 – 2.55 (m, 1H), 2.31 – 2.21 (m, 1H), 2.08 – 1.21 (m, 20H), 1.63 (d,  
 3H).

Example 923(1) and Example 923(2): [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((6S)-  
 10 2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-  
a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate and [(2,2-  
dimethylpropanoyl)oxy]methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-  
oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1-[(2,2-  
dimethylpropanoyl)oxy]methyl)-1H-imidazol-5-yl)-2-thiophenecarboxylate



The same operation as in Example 373 was conducted from the compound prepared in Example 392 and chloromethyl 2,2-dimethylpropanoate to give the title compounds having the following physical properties.

20 Example 923(1):LC/MS  $t_R$  0.90 minutes; MS ( $ES^+$ )  $m/z$  621(M+H)<sup>f</sup>


<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.43 (s, 1H), 8.79 (s, 1 H), 8.05 (s, 1H), 7.68 (s, 1H),  
 7.66 (d, 1H), 7.64 – 7.62 (m, 1H), 7.47 (d, 1H), 7.17 (s, 1H), 6.38 (s, 1 H), 5.96 (s, 2H),  
 5.75 (d, 1 H), 3.48 – 3.32 (m, 2H), 2.97 – 2.85 (m, 1H), 2.60 – 2.43 (m, 1 H), 1.23 (s,  
 25 9H).

Example 923(2):LC/MS  $t_R$  1.13 minutes; MS ( $ES^+$ )  $m/z$  735(M+H)<sup>f</sup>

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.78 (s, 1 H), 8.01 (d, 1H), 7.79 (d, 1H), 7.68 (d, 1H), 7.64 – 7.60 (m, 1H), 7.49 (d, 1H), 7.21 (s, 1H), 6.45 (d, 1H), 6.27 (s, 1 H), 5.94 (s, 2H), 5.91 (d, 1H), 5.87 – 5.83 (m, 1 H), 3.77 – 3.62 (m, 1H), 2.98 – 2.85 (m, 1H), 2.65 – 2.49 (m, 2 H), 1.21 (s, 9H), 1.18 (s, 9H).


5

Example 924: 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-fluoro-2-thiophenecarboxylic acid




- 10 LC/MS  $t_R$  0.63 minutes; MS ( $\text{ES}^+$ )  $m/z$  525 ( $\text{M}+\text{H}$ )<sup>f</sup>  
 $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.71 (s, 1 H), 7.94 (d, 1 H), 7.89 (d, 1 H), 7.87 - 7.78 (m, 2 H), 7.35 (d, 1 H), 6.34 (s, 1 H), 5.63 (dd, 1 H), 3.21 - 3.04 (m, 1 H), 2.84 - 2.68 (m, 1 H), 2.63 - 2.51 (m, 1 H) 2.28 - 2.11 (m, 1 H).

- 15 Example 925: 5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride



- LC/MS  $t_R$  0.65 minutes; MS ( $\text{ES}^+$ )  $m/z$  507( $\text{M}+\text{H}$ )<sup>f</sup>  
 20  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.72 (s, 1 H), 7.97 – 7.90 (m, 2H), 7.88 – 7.80 (m, 3H), 7.70 (d, 1H), 7.54 (d, 1H), 6.36 (s, 1 H), 5.80 – 5.74 (m, 1 H), 3.20 – 3.06 (m, 1H), 2.92 – 2.80 (m, 1H), 2.70 – 2.56 (m, 1 H), 2.36– 2.23 (m, 1 H).

Example 926: 5-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid hydrochloride



- 5 LC/MS  $t_R$  0.75 minutes; MS ( $ES^+$ )  $m/z$  525(M+H)<sup>+</sup>  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  13.28 (s, 1H), 9.74 (s, 1 H), 7.97 (d, 1H), 7.87 – 7.79 (m, 2H), 7.69 (d, 1H), 7.26 (d, 1H), 6.35 (s, 1 H), 5.57 – 5.50 (m, 1 H), 3.17 – 3.01 (m, 1H), 2.83 – 2.70 (m, 1H), 2.63 – 2.47 (m, 1 H), 2.21– 2.10 (m, 1 H).

- 10 Example 927: methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-6-methyl-2-oxo-1(2H)-pyridinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 44 was conducted from the compound prepared in Example 318 to give the title compounds having the following physical properties.

- 15 TLC Rf 0.80 (10% methanol in ethyl acetate)  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.81 (s, 1 H), 9.86 - 9.59 (m, 2 H), 7.90 - 7.70 (m, 3 H), 7.68 - 7.44 (m, 4 H), 6.02 (s, 1 H), 5.94 (s, 1 H), 5.15 (s, 2 H), 3.67 (s, 3 H), 2.43 (s, 3 H).

- 20 Example 928: methyl {4-[4-chloro-2-({4-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2-methyl-6-oxo-1(6H)-pyrimidinyl}methyl)-1H-imidazol-5-yl]phenyl}carbamate

The same operation as in Example 44 was conducted from the compound prepared in Example 323 to give the title compounds having the following physical properties.

- 25 TLC Rf 0.63 (1% methanol in ethyl acetate)  
<sup>1</sup>H NMR (300 MHz, DMSO- $d_6$ )  $\delta$  12.84 (s, 1 H), 9.78 (s, 2 H), 8.01 (d, 1 H), 7.90 - 7.75 (m, 2 H), 7.67 - 7.41 (m, 4 H), 6.51 (s, 1 H), 5.17 (s, 2 H), 3.68 (s, 3 H), 2.39 (s, 3 H).

## PHARMACOLOGICAL ACTIVITIES

The compounds of the present invention possess Factor XIa inhibitory activity and oral bioavailability, for example, such an effect of the compounds of the present invention was confirmed by the following tests.

5 All the procedures were conducted by conventionally used techniques on the basis of basic biological methods. Furthermore, the measuring method of the present invention was modified to improve the accuracy and/or sensitivity of measurement for evaluating the compound of the present invention. The detailed experimental method was as follows.

## 10 EXPERIMENTAL METHOD

### (1) In Vitro Assay

Inhibitory activities of compounds of the present invention against factor XIa, Xa, XIIa, IXa, VIIa, plasma kallikrein or thrombin were evaluated using appropriate purified proteases and synthetic substrates. The rate of hydrolysis of the chromogenic  
15 substrate by the relevant protease was continuously measured at 405 nm.

Inhibitory activity against each enzyme was calculated as % inhibition using the equation described below.

% Inhibition =  $\frac{[(\text{rate without compound}) - (\text{rate with compound})]}{(\text{rate without compound})} \times 100\%$ .

20 Each half maximal inhibitory concentration (IC<sub>50</sub>) value was determined by plotting the concentration of compound of the invention against the % inhibition.

#### (1-1) Factor XIa enzyme activity

Human Factor XIa (Haematologic Technologies Inc.) activity was measured at  
25 an enzyme concentration of 0.1 U/mL in 150 mM NaCl, 5 mM KCl, 1 mg/mL PEG6000, 50 mM HEPES-NaOH (pH7.4) with 300 μM S-2366 (pyroGlu-Pro-Arg-pNA, Chromogenix).

#### (1-2) Plasma kallikrein enzyme activity

30 Human plasma kallikrein (Enzyme Research Laboratories Ltd) activity was measured at an enzyme concentration of 0.605 mU/mL in 200 mM NaCl, 5 mg/mL PEG6000, 100 mM Phosphate-NaOH (pH7.4) with 150 μM S-2302 (H-D-Pro-Phe-Arg-pNA, Chromogenix).



(1-3) Factor Xa and thrombin enzyme activity

Human Factor Xa (American Diagnostica Inc.) and human thrombin (Sigma) activities were measured at the enzyme concentrations of 0.18 U/mL and 0.12 U/mL, respectively in the same buffer containing 150 mM NaCl, 2 mg/mL PEG6000, 50 mM Tris-HCl (pH7.4), except that the reactions were started with 300  $\mu$ M S-2222 (phenyl-Ile-Glu-Gly-Arg-pNA, Chromogenix) and 300  $\mu$ M S-2366, respectively.

(1-4) Factor XIIa enzyme activity

Human Factor  $\alpha$ -XIIa (Enzyme Research Laboratories Ltd) activity was measured at an enzyme concentration of 0.17 U/mL in 150 mM NaCl, 50 mM Tris-HCl (pH7.4) with 300  $\mu$ M S-2302 (H-D-Pro-Phe-Arg-pNA, Chromogenix).

(1-5) Factor IXa enzyme activity

Human Factor IXa (American Diagnostica Inc.) activity was measured at an enzyme concentration of 13 U/mL in 100 mM NaCl, 5 mM  $\text{CaCl}_2$ , 30% ethylene glycol, 50 mM Tris-HCl (pH7.4) with 3 mM Pefachrome IXa 3960 (Leu-Ph'Gly-Arg-pNA, Pentapharm).

(1-6) Factor VIIa enzyme activity

Human Factor VIIa activity was measured using recombinant human Factor VIIa (American Diagnostica Inc.) in the presence of recombinant human tissue factor which was produced according to the method described in the literature (Protein expression and purification, 3, 453-460 (1992) in a buffer containing 150 mM NaCl, 5 mM  $\text{CaCl}_2$ , 0.5 mg/mL PEG6000, 50 mM HEPES-NaCl (pH7.4) with 3 mM S-2288 (Ile-Pro-Arg-pNA, Chromogenix).

(1-7) APTT, PT measurement

Activated partial thromboplastin time (APTT) and prothrombin time (PT) were measured using automatic coagulation analyzer (CA-1500, Sysmex Corporation). For the APTT or PT measurement, standard human plasma (Siemens Healthcare Diagnostics GmbH) were mixed with each compound dilutions followed by the automatic addition of APTT reagent (Siemens Healthcare Diagnostics GmbH) and 0.02

M calcium chloride or PT reagent (Siemens Healthcare Diagnostics GmbH) to start clot formation. The anticoagulant activities (APTT2 or PT2) of the compounds of the invention were expressed as the concentrations necessary to double the clotting time in vehicle (1% DMSO) group. APTT2 or PT2 was determined by plotting the  
5 concentration of compound of the invention against the fold increase of clotting time.

(2) Rat oral administration test

Each compound of the present invention in a solution of 20% wellsolve (celeste) was given to fasted male Crj:CD(SD)IGS rats as a single 3 mg/kg, p.o. dose by gavage.  
10 Blood samples were drawn from jugular vein into syringes containing 3.2% sodium citrate (the volume ratio of blood to anticoagulant = 9:1) after oral administration at 0.5, 1, 2, 4 and 6 hours. Plasma was harvested by centrifugation and stored at -20°C until measurement of plasma concentration.

To measure plasma concentrations of the compounds of the present invention,  
15 plasma samples were deproteinized with acetonitrile, followed by evaporation of the acetonitrile to dryness. The residual was then re-dissolved with DMSO and its FXIa inhibitory activity was measured by enzyme assay described above. Plasma concentrations of each compound of the present invention were quantitated by generating a standard curve of known compound concentration in rat plasma.

20

The compounds of the present invention were tested in the Factor XIa assay and the rat oral administration test described above, and found to have a good Factor XIa inhibitory activity and good oral bioavailability. Table 1 described below lists Factor XIa IC<sub>50</sub> values measured for the following examples.

25








Table 1

| Example No | In vitro FXIa inhibitory activity<br>IC <sub>50</sub> (μM) |
|------------|------------------------------------------------------------|
| 41         | 0.0065                                                     |
| 45         | 0.015                                                      |
| 57         | 0.0024                                                     |
| 60         | 0.0072                                                     |
| 61         | 0.014                                                      |
| 86         | 0.0039                                                     |
| 129(2)     | 0.0046                                                     |
| 181        | 0.047                                                      |
| 198        | 0.0015                                                     |
| 210        | 0.0028                                                     |
| 216        | 0.0038                                                     |
| 230        | 0.011                                                      |
| 241        | 0.0048                                                     |
| 245        | 0.017                                                      |
| 264        | 0.0061                                                     |
| 270        | 0.011                                                      |
| 278        | 0.0057                                                     |
| 290        | 0.0025                                                     |
| 309        | 0.0083                                                     |
| 310        | 0.0049                                                     |
| 347        | 0.0042                                                     |
| 363        | 0.0014                                                     |
| 387        | 0.0016                                                     |
| 391        | 0.0011                                                     |
| 392        | 0.0021                                                     |
| 399        | 0.0011                                                     |
| 400        | 0.013                                                      |
| 834        | 0.0016                                                     |
| 848        | 0.0017                                                     |

#### Comparative experiments

CompEx1, Ex2, Ex3 and Ex4 as comparative compounds are prepared according to the reported preparation in WO2007/070826 (Ex.64 and Ex.254), WO2008/076805 (Ex.1) and WO2009/076337 (Ex.1). The comparative compounds were also tested in the Factor XIa assay and the rat oral administration test described above. Table 2 described below lists Factor XIa IC<sub>50</sub> values and plasma concentrations measured for the comparative compounds and the typical compound of the present invention.

Table 2

| Example No | Structure                                                                           | In vitro FXIa inhibitory activity<br>IC <sub>50</sub> (μM) | rat plasma concentration<br>after po administration<br>C <sub>max</sub> (μM) at 3mpk |
|------------|-------------------------------------------------------------------------------------|------------------------------------------------------------|--------------------------------------------------------------------------------------|
| 45         |    | 0.015                                                      | 1.25                                                                                 |
| 41         |    | 0.0065                                                     | 2.28                                                                                 |
| 57         |   | 0.0024                                                     | 0.51                                                                                 |
| Comp Ex 1  |  | 0.0036                                                     | 0.015                                                                                |
| Comp Ex 2  |  | 0.040                                                      | 0.025                                                                                |
| Comp Ex 3  |  | 0.130                                                      | 0.088                                                                                |
| Comp. Ex 4 |  | >10                                                        | Not tested                                                                           |

As a result, the compounds of the present invention showed good oral bioavailability compared with the comparative compounds described above.

Therefore, the results indicated that the compounds of the present invention possess Factor XIa inhibitory activity, with good oral bioavailability.

Additionally, the good oral bioavailability of compounds of the present invention can be determined using the following experimental methods.

10 (3-1) Pharmacokinetic (PK) study in rat

Each compound of the present invention in a solution of 20% wellsolve (celeste) was given to fasted male Crj:CD(SD) rats as a single 3 mg/kg, p.o. dose by gavage. Blood samples were drawn from jugular vein into syringes containing 3.2% sodium citrate (the volume ratio of blood to anticoagulant = 9:1) or heparinized syringes at 0.5, 1, 3, 7 hours after oral administration. Plasma was obtained by centrifugation and stored at -20°C until measurement of plasma concentration.

To measure plasma concentrations of the compounds of the present invention, plasma samples were deproteinized with acetonitrile, followed by evaporation of the acetonitrile to dryness. Then the sample was reconstituted in the mobile phase and analyzed by LC/MS/MS. An analytical column (Shim-pack XR-ODS II, 2.0 mm × 75 mm, 2.2 μm) and mobile phase (0.1% formic acid in water and 0.1% formic acid in acetonitrile, flow rate of 0.5 mL/min) were used. The system was used in multiple reaction monitoring (MRM) mode with positive ion detection.

25 (3-2) Pharmacokinetic (PK) study of the compound which has an ester group(s) in rat

Each compound of the present invention in a solution of 20% wellsolve (celeste) was given to fasted male Crj:CD(SD) rats as a single 3 mg/kg, p.o. dose by gavage. Blood samples were drawn from jugular vein into syringes treated with heparin-diisopropyl fluorophosphate mixture (500:1) at 0.5, 1, 3, 7 hours after oral administration. Plasma was obtained by centrifugation and stored at -20°C until measurement of plasma concentration.

To measure plasma concentrations of the compounds of the present invention, plasma samples were deproteinized with acetonitrile, followed by evaporation of the

acetonitrile to dryness. Then the sample was reconstituted in the mobile phase and analyzed by LC/MS/MS. An analytical column (Shim-pack XR-ODS II, 2.0 mm × 75 mm, 2.2 μm) and mobile phase (0.1% formic acid in water and 0.1% formic acid in acetonitrile, flow rate of 0.5 mL/min) were used. The system was used in multiple  
5 reaction monitoring (MRM) mode with positive ion detection.

Table 3

| Example No. of administered compound | rat plasma concentration of ester compound after po administration Cmax (μM) at 3mpk | rat plasma concentration of corresponding carboxylic acid after po administration Cmax (μM) at 3mpk |
|--------------------------------------|--------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|
| 401                                  | 0.96                                                                                 | 5.6                                                                                                 |

Additionally, enzymatic hydrolysis of an ester group in the compound of the  
10 present invention can be determined using the following experimental methods.

(4-1) Analysis of enzymatic hydrolysis of an ester group(s) in the compounds of the present invention using hepatocytes prepared from various species (rat, dog, monkey, human)

15 A typical assay procedure was conducted by using cryopreserved hepatocytes prepared from various species. A mixture of hepatocytes, buffer (pH 7.4), and each test compound were incubated. The final test compound concentration was typically 100 ng/mL, with a usual cell density of 1,000,000 cells/ml for all species. The incubation was at 37°C, with time-points taken over 120 minutes. Reaction termination was  
20 achieved by addition of an aliquot of the hepatocyte / test compound mixture to acetonitrile/ethanol (7/3) to effect protein precipitation, followed by centrifugation. Then the sample was diluted with distilled water and analyzed by LC/MS/MS. An analytical column (Shim-pack XR-ODS II, 2.0 mm × 75 mm, 2.2 μm) and mobile phase (0.1% formic acid in water and 0.1% formic acid in acetonitrile, flow rate of 0.5  
25 mL/min) were used. The system was used in multiple reaction monitoring (MRM) mode with positive ion detection.

(4-2) Analysis of enzymatic hydrolysis of the compound which has an ester group(s) using blood from various species (rat, dog, monkey, human)

Each compound of the present invention in a solution of acetonitrile were incubated in blood from various species. The incubation was typically performed at a concentration of 100ng/mL of test compound at 37°C, with time points taken over 60 minutes. The reaction was terminated by addition of an aliquot of blood / test compound mixture to acetonitrile/ethanol (7/3) to effect protein precipitation, followed by centrifugation. Then the sample was diluted with distilled water and analyzed by LC/MS/MS. An analytical column (Shim-pack XR-ODS II, 2.0 mm × 75 mm, 2.2 μm) and mobile phase (0.1% formic acid in water and 0.1% formic acid in acetonitrile, flow rate of 0.5 mL/min) were used. The system was used in multiple reaction monitoring (MRM) mode with positive ion detection.

Formulation example 1

The following components were admixed in conventional method and punched out to obtain 10,000 tablets each containing 10 mg of active ingredient.

- methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-5-methyl-1H-imidazol-4-yl)phenyl]carbamate 100 g
- Carboxymethylcellulose calcium (disintegrating agent) 20 g
- Magnesium stearate (lubricating agent) 10 g
- Microcrystalline cellulose 870 g

Formulation example 2

The following components were admixed in conventional method. The solution was sterilized in conventional manner, filtered through dust removal equipment, placed 5 mL portions into ampoules and sterilized by autoclave to obtain 10,000 ampoules each containing 20 mg of the active ingredient.

- methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-5-methyl-1H-imidazol-4-yl)phenyl]carbamate 200 g
- mannitol 20 g
- distilled water 50 L


## INDUSTRIAL APPLICABILITY

The compounds of the present invention represented by formula (I) act as potent and selective inhibitors of Factor XIa, with potent anticoagulant activity and/or good oral availability. In particular, the compounds of the present invention act as a Factor XIa inhibitor or a Factor XIa and plasma kallikrein dual inhibitor. Thus the compounds of the present invention are useful in preventing and/or treating thromboembolic diseases, for example arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, arterial cerebrovascular thromboembolic disorders, venous cerebrovascular thromboembolic disorders and thromboembolic disorders in the chambers of the heart or in the peripheral circulation. The compound of the present invention is therefore useful as a medicament.



## CLAIMS

1. A compound represented by formula (I):



5            wherein Cyc<sup>1</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

             Cyc<sup>2</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- or 6-membered heteroaryl;

10            Cyc<sup>3</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

             R<sup>1</sup> represents (1) C1-8 alkyl, (2) C2-8 alkenyl, (3) C2-8 alkynyl, (4) halogen, (5) nitro, (6) cyano, (7) oxo, (8) amidino, (9) -C1-8 alkylene -OR<sup>8</sup>, (10) -OR<sup>9</sup>, (11) -COOR<sup>10</sup>, (12) -C1-4 alkylene -COOR<sup>11</sup>, (13) -NHC(O)-C1-4 alkyl, (14) -C1-4 alkylene-O-C(O)-C1-8 alkyl or (15) -NHC(O)O-R<sup>12</sup>,

15            wherein R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup> and R<sup>12</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

20            s represents an integer of 0 to 6,

             wherein s represents an integer of 2 to 6, each R<sup>1</sup> may be same or different;

             R<sup>2</sup> represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) Cyc<sup>4</sup> or (6) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from halogen, nitro, trifluoromethyl, cyano, Cyc<sup>5</sup>, -NR<sup>13</sup>R<sup>14</sup>, -OR<sup>15</sup>, -SR<sup>16</sup>, -NHC(O)-Cyc<sup>6</sup>, -NHC(O)-C1-8 alkyl, -NHC(O)O-R<sup>17</sup> and Cyc<sup>5</sup> substituted with  
25            1 to 3 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino and -OR<sup>18</sup>,

wherein  $R^{13}$ ,  $R^{14}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{17}$  and  $R^{18}$  each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

$Cyc^4$ ,  $Cyc^5$  and  $Cyc^6$  each independently represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

$R^3$  represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5)  $Cyc^7$  or (6) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from halogen, nitro, trifluoromethyl, cyano,  $Cyc^8$ ,  $-NR^{19}R^{20}$ ,  $-OR^{21}$ ,  $-SR^{22}$ ,  $-NHC(O)-Cyc^9$ ,  $-NHC(O)-C1-8$  alkyl,  $-NHC(O)O-R^{23}$  and  $Cyc^8$  substituted with 1 to 3 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino and  $-OR^{24}$ ,

wherein  $R^{19}$ ,  $R^{20}$ ,  $R^{21}$ ,  $R^{22}$ ,  $R^{23}$  and  $R^{24}$  each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

$Cyc^7$ ,  $Cyc^8$  and  $Cyc^9$  each independently represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

Y represents N or  $C(R^5)$ ;

$R^4$  and  $R^5$  each independently represents (1) hydrogen, (2) halogen, (3) C1-4 alkyl, (4) C3-C8 cycloalkyl, (5) 5- to 10-membered heterocycloalkyl, (6) C5-C10 aryl, (7) 5- to 10-membered heteroaryl or (8) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

$R^2$  and  $R^3$  may be taken together to form C2-8 alkylene; or

$R^3$  and  $R^4$  may be taken together to form C2-8 alkylene; wherein one carbon of the alkylene chain may be replaced by oxygen or sulfur;

$R^6$  represents (1) C1-8 alkyl, (2) C2-8 alkenyl, (3) C2-8 alkynyl, (4)  $Cyc^{10}$ , (5) halogen, (6) nitro, (7) cyano, (8) oxo, (9) amidino, (10)  $-OR^{25}$ , (11)  $-COOR^{26}$ , (12)  $-C1-4$  alkylene- $COOR^{27}$ , (13)  $-NHC(O)-C1-4$  alkyl, (14)  $-NHC(O)-H$ , (15)  $-NHC(O)O-R^{28}$ , (16) trifluoromethyl, (17)  $-NHC(NH)NH_2$ , (18)  $-C(O)-C1-4$  alkyl or (19)  $Cyc^{10}$

substituted with 1 to 5 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, cyano, oxo, amidino, -OR<sup>29</sup>, -SR<sup>30</sup>, -NR<sup>31</sup>R<sup>32</sup>, -NHC(O)NR<sup>33</sup>R<sup>34</sup>, -NHC(O)-C1-4 alkylene -COOH, -NH-S(O)-C1-4 alkyl, -NH-S(O)<sub>2</sub>-C1-4 alkyl, -COOR<sup>35</sup>, -NHC(O)-R<sup>36</sup>, -NHC(O)O-R<sup>37</sup>, -C(O)NH-R<sup>38</sup> and -OC(O)NH-R<sup>39</sup>,

5 wherein Cyc<sup>10</sup> represents C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- to 10-membered heteroaryl;

R<sup>25</sup>, R<sup>26</sup>, R<sup>27</sup>, R<sup>28</sup>, R<sup>29</sup>, R<sup>30</sup>, R<sup>31</sup>, R<sup>32</sup>, R<sup>33</sup>, R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup>, R<sup>37</sup>, R<sup>38</sup> and R<sup>39</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C3-C8 cycloalkyl, (6) 5- to 10-membered heterocycloalkyl, (7) C5-C10 aryl, 10 (8) 5- to 10-membered heteroaryl or (9) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

m represents an integer of 0 to 6,

wherein m represents an integer of 2 to 6, each R<sup>6</sup> may be same or different;

15 R<sup>7</sup> represents (1) C1-8 alkyl, (2) C2-8 alkenyl, (3) C2-8 alkynyl, (4) halogen, (5) nitro, (6) trifluoromethyl, (7) cyano, (8) oxo, (9) amidino, (10) -OR<sup>40</sup>, (11) -SR<sup>41</sup>, (12) -NR<sup>42</sup>R<sup>43</sup>, (13) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (14) -NHC(O)-C1-4 alkylene-NR<sup>46</sup>R<sup>47</sup>, (15) -NHC(O)-C1-4 alkylene-COOH, (16) -NH-S(O)-C1-4 alkyl, (17) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (18) -COOR<sup>48</sup>, (19) -NHC(O)-R<sup>49</sup>, (20) -NHC(O)-C1-4 alkylene-OR<sup>50</sup>, (21) -NHC(O)O-R<sup>51</sup>, (22) -NHC(O)O-C1-4 alkylene -OR<sup>52</sup>, (23) -C(O)NH-R<sup>53</sup>, (24) -OC(O)NH-R<sup>54</sup>, (25) -OC(O)-R<sup>55</sup>, (26) -C(O)-R<sup>56</sup>, (27) -CH(OH)-R<sup>57</sup>, (28) -C1-4 alkylene-NH<sub>2</sub>, (29) -C1-4 alkylene-OH, (30) -C1-4 alkylene-OC(O)-C1-4 alkyl, (31) -C1-4 alkylene-NHC(O)-C1-4 alkyl, (32) -C1-4 alkylene-NHC(O)O-C1-4 alkyl, (33) -C1-4 alkylene-NHC(O)-CF<sub>3</sub>, (34) -C1-4 alkylene-NHC(O)NH-C1-4 alkyl, (35) -CH=N-OR<sup>58</sup>, (36) -C(O)N(C1-4 alkyl)<sub>2</sub>, (37) -C(O)NH-R<sup>63</sup>, (38) -S(O)<sub>2</sub>-NR<sup>64</sup>R<sup>65</sup>, (39) -T-COOR<sup>66</sup>, (40) -B(OR<sup>67</sup>)(OR<sup>68</sup>), (41) C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl or 5- or 6-membered heteroaryl, which are substituted with 1 to 5 groups selected from C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, halogen, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl, OH, oxo, and cyano, (42) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from halogen, trifluoromethyl, OH, oxo, -O-C1-4 alkyl, NH<sub>2</sub>, and cyano, (43) -N(OH)C(O)-C1-4 alkyl, (44) -NHC(=N-OR<sup>69</sup>)-C1-4 alkyl, (45) -NHC(S)-C1-4 alkyl, (46) -C(S)-C1-4 alkyl, (47) -S(O)NR<sup>70</sup>R<sup>71</sup>, (48) -C(O)NH(CO)NR<sup>72</sup>, (49) -NHC(O)R<sup>73</sup>, (50) -NHC(O)O-C1-4 alkyl

substituted with 1 to 5 groups selected from -O-C1-4 alkylene-O-C1-4 alkyl, NH<sub>2</sub>, and OH, (51) -NHC(O)-C1-4 alkyl substituted with 1 to 5 groups selected from -O-C1-4 alkylene-O-C1-4 alkyl, NR<sup>74</sup>R<sup>75</sup>, oxo, OH, halogen and -O-C1-4 alkylene-O-C1-4 alkyl substituted with 1 to 2 groups selected from OH, oxo and halogen, (52) -NH-C1-8 alkyl substituted with 1 to 5 groups selected from -O-C1-4 alkyl, oxo, NR<sup>76</sup>R<sup>77</sup>, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- or 6-membered heteroaryl, (53) C3-C8 cycloalkyl, (54) 5- to 10-membered heterocycloalkyl, (55) C5-C10 aryl, or (56) 5- or 6-membered heteroaryl;

wherein R<sup>40</sup>, R<sup>41</sup>, R<sup>42</sup>, R<sup>43</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup>, R<sup>47</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>51</sup>, R<sup>52</sup>, R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup>, R<sup>57</sup>, R<sup>58</sup>, R<sup>67</sup>, R<sup>68</sup>, R<sup>69</sup>, R<sup>70</sup>, R<sup>71</sup>, R<sup>72</sup>, R<sup>73</sup>, R<sup>74</sup>, R<sup>75</sup>, R<sup>76</sup> and R<sup>77</sup> each independently represents (1) hydrogen, (2) trifluoromethyl, (3) C1-8 alkyl, (4) C2-8 alkenyl, (5) C2-8 alkynyl, (6) C3-C8 cycloalkyl, (7) 5- to 10-membered heterocycloalkyl, (8) C5-C10 aryl, (9) 5- to 10-membered heteroaryl or (10) C1-4 alkyl substituted with 1 to 5 groups selected from C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl;

R<sup>63</sup> represents (1) -C1-4 alkylene-O-C1-4 alkyl, (2) -O-C1-4 alkyl, (3) cyano, (4) -C1-4 alkylene-O-C1-4 alkylene-O-C1-4 alkyl or (5) -SO<sub>2</sub>-C1-4 alkyl;

R<sup>64</sup> and R<sup>65</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (5) C2-8 alkynyl, (6) -C(O)-C1-4 alkyl or (7) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from OH, oxo, -O-C1-4 alkyl, -O-C1-4 alkylene-O-C1-4 alkyl, halogen, nitro and cyano;

T represents (1) C1-4 alkylene, (2) C2-4 alkenylene, (3) -O-C1-4 alkylene-, (4) -O-C2-4 alkenylene-, (5) -S-C1-4 alkylene-, (6) -S-C2-4 alkenylene-, (7) -NH-C1-4 alkylene-, (8) -NH-C2-4 alkenylene-, (9) -NH-C5-C10 aryl- or (10) -NH-5- to 10-membered heteroaryl-;

R<sup>48</sup> and R<sup>66</sup> each independently represents (1) hydrogen, (2) C1-8 alkyl, (3) C2-8 alkenyl, (4) C2-8 alkynyl, (5) C1-8 alkyl, C2-8 alkenyl or C2-8 alkynyl, which are substituted with 1 to 5 groups selected from -NH<sub>2</sub>, -NH-C1-4 alkyl, -N(C1-4 alkyl)<sub>2</sub>, OH, oxo, -O-C1-4 alkyl, -O-C1-4 alkylene-O-C1-4 alkyl, halogen, nitro, cyano, C3-C8 cycloalkyl, 5- to 10-membered heterocycloalkyl, C5-C10 aryl and 5- to 10-membered heteroaryl (6) C3-C10 cycloalkyl, (7) 5- to 10-membered heterocycloalkyl, (8) C5-C10 aryl, (9) 5- to 10-membered heteroaryl, (10) -C1-4 alkylene-C3-C8 cycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH, trifluoromethyl and

halogen, (11) -C1-4 alkylene-C5-C10 aryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH, trifluoromethyl and halogen, (12) -C1-4 alkylene-5- to 10-membered heterocycloalkyl substituted with 1 to 5 groups selected from C1-4 alkyl, oxo, OH, trifluoromethyl and halogen, (13) -C1-4 alkylene-5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from C1-4 alkyl, OH, trifluoromethyl and halogen or (14) -C1-4 alkylene-O-C1-8 alkyl substituted with 1 to 5 groups selected from OH, oxo, -O-C1-4 alkyl, -O-C1-4 alkylene-O-C1-4 alkyl, halogen, nitro, cyano, C3-C8 cycloalkyl, C5-C10 aryl, 5- to 10-membered heterocycloalkyl, -O-5- to 10-membered heteroaryl, -O-C3-C8 cycloalkyl, -O-C5-C10 aryl, -O-5- to 10-membered heterocycloalkyl, -O-5- to 10-membered heteroaryl, -O-C1-4 alkylene-5- to 10-membered heteroaryl, -O-C1-4 alkylene-C3-C8 cycloalkyl, -O-C1-4 alkylene-C5-C10 aryl, -O-C1-4 alkylene-5- to 10-membered heterocycloalkyl and -O-C1-4 alkylene-5- to 10-membered heteroaryl;


n represents an integer of 0 to 6,

wherein n represents an integer of 2 to 6, each  $R^7$  may be same or different; and

$R^{62}$  represents hydrogen or halogen,

a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.


2. The compound according to claim 1, wherein the compound represented by formula (I) represents a compound represented by formula (I-A):



wherein U represents S or CH<sub>2</sub>; and

the other symbols have the same meanings as described in claim 1.

3. The compound according to claim 1, wherein the compound represented by formula (I) represents a compound represented by formula (I-B):



wherein Cyc<sup>1-B</sup> represents 5- to 10-membered heteroaryl;


5 Cyc<sup>3-B</sup> represents C5-C10 aryl or 5- to 10-membered heteroaryl;

Cyc<sup>10-B</sup> represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl and -CONH<sub>2</sub>;

mb represents an integer of 0 to 5; and

10 the other symbols have the same meanings as described in claim 1.

4. The compound according to claim 1, wherein the compound represented by formula (I) represents a compound represented by formula (I-C):



15 wherein Cyc<sup>1-C</sup> represents 5- to 10-membered heteroaryl;

Cyc<sup>3-C</sup> represents C5-C10 aryl or 5- to 10-membered heteroaryl;

Cyc<sup>10-C</sup> represents (1) 5- to 10-membered heteroaryl or (2) 5- to 10-membered heteroaryl substituted with 1 to 5 groups selected from halogen, cyano, nitro, trifluoromethyl, -COOH, -COO-C1-4 alkyl and -CONH<sub>2</sub>;

20 mc represents an integer of 0 to 5; and


the other symbols have the same meanings as described in claim 1.

5. The compound according to claim 1 or 2, wherein Cyc<sup>1</sup> represents phenyl, imidazolyl, triazolyl, pyrrolyl, pyrazolyl, furanyl, oxazolyl, thiazolyl, isothiazolyl, furazanyl, oxadiazolyl, thiadiazolyl, thienyl, pyridazinyl, indazolyl or benzimidazolyl.


6. The compound according to claim 3, wherein Cyc<sup>1-B</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl.

7. The compound according to claim 4, wherein Cyc<sup>1-C</sup> represents imidazolyl, triazolyl, tetrazolyl, oxazolyl, pyridazinyl, indazolyl or benzimidazolyl.

8. The compound according to claim 1 or claim 2, wherein



represents




wherein  $R^{59}$  represents hydrogen, C1-4 alkyl or halogen;

the arrow represents a binding position; and

5 the other symbols have the same meanings as described in claim 1.

9. The compound according to any one of claims 1 to 8, wherein  $Cyc^2$  represents pyridyl or phenyl.

10. The compound according to any one of claims 1 to 9, wherein  $-Cyc^2-$  ( $R^6$ )<sub>m</sub>,  $Cyc^2(-R^6)$ <sub>mb</sub> $Cyc^{10-B}$  or  $Cyc^2(-R^6)$ <sub>mc</sub> $Cyc^{10-C}$  represents





wherein R<sup>60</sup> represents hydrogen, methyl or halogen;

R<sup>61</sup> represents (1) hydrogen, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) -COOH, (6) -COO-C1-4 alkyl, (7) cyano or (8) -CONH<sub>2</sub>; and the arrow represents a binding position.

5

11. The compound according to any one of claims 1 to 10, wherein R<sup>7</sup> represents (1) C1-8 alkyl, (2) halogen, (3) nitro, (4) trifluoromethyl, (5) cyano, (6) oxo, (7) -OR<sup>40</sup>, (8) -NR<sup>42</sup>R<sup>43</sup>, (9) -NHC(O)NR<sup>44</sup>R<sup>45</sup>, (10) -NHC(O)-C1-4 alkylene-NR<sup>46</sup>R<sup>47</sup>, (11) -NHC(O)-C1-4 alkylene-COOH, (12) -NH-S(O)<sub>2</sub>-C1-4 alkyl, (13) -COOR<sup>48</sup>, (14) -NHC(O)-R<sup>49</sup>, (15) -NHC(O)-C1-4 alkylene-OR<sup>50</sup>, (16) -NHC(O)O-R<sup>51</sup>, (17) -NHC(O)O-C1-4 alkylene -OR<sup>52</sup>, (18) -C(O)NH-R<sup>53</sup>, (19) -OC(O)-R<sup>55</sup>, (20) -C(O)-R<sup>56</sup>, (21) -CH(OH)-R<sup>57</sup>, (22) -C1-4 alkylene -NH<sub>2</sub>, (23) -C1-4 alkylene-OH, (24) -C1-4 alkylene-OC(O)-C1-4 alkyl, (25) -C1-4 alkylene-NHC(O)-C1-4 alkyl, (26) -C1-4 alkylene-NHC(O)O-C1-4 alkyl, (27) -C1-4 alkylene-NHC(O)-CF<sub>3</sub>, (28) -C1-4 alkylene-NHC(O)NH-C1-4 alkyl, (29) -CH=N-OR<sup>58</sup> or (30) -T-COOR<sup>66</sup>.

12. The compound according to claim 1 or claim 2, wherein is selected from the group consisting of

- (1) methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl]phenyl]carbamate,
- (2) methyl [4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl]phenyl]carbamate,
- (3) 4-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl]benzamide,
- (4) 6-(2-{7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-4-yl)-3,4-dihydro-2(1H)-quinolinone,
- (5) methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-5-methyl-1H-imidazol-4-yl]phenyl]carbamate,
- (6) 6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl)-3,4-dihydro-2(1H)-quinolinone,
- (7) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny]-1H-imidazol-5-yl]benzamide,

- (8) 3-[5-(3-amino-1H-indazol-6-yl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (9) (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 5 (10) (3S)-3-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (11) (3S)-3-[5-(1-amino-6-isoquinolinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (12) 4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl)benzamide,
- 10 (13) ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl)phenyl]carbamate,
- (14) methyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]-3-methylphenyl]carbamate,
- 15 (15) methyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,
- (16) 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl)phenyl]carbamate,
- (17) 2-methoxyethyl [4-(2-(7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl)phenyl]carbamate,
- 20 (18) methyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]-3-pyridinyl]carbamate,
- (19) (3S)-3-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- 25 (20) (6S)-6-[5-(6-amino-2-chloro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,
- (21) 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,
- 30 (22) (3S)-3-[5-(6-amino-3-pyridinyl)-4-chloro-1H-imidazol-2-yl]-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,
- (23) 2-methoxyethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indolizinyl))-1H-imidazol-5-yl]-2-pyridinyl]carbamate,

- (24) 2-methoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,
- (25) 2-methoxyethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-2-pyridinyl]carbamate,
- (26) 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- (27) 2-methoxyethyl [4-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,
- (28) 2-methoxyethyl [4-(4-methyl-2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,
- (29) 2-methoxyethyl [6-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)-3-pyridinyl]carbamate,
- (30) 2-ethoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,
- (31) 2-methoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,
- (32) 3-methoxypropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,
- (33) methyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,
- (34) 3-oxetanyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])]-1H-imidazol-5-yl)phenyl]carbamate,
- (35) 2-methoxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,
- (36) methyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,
- (37) 2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)phenyl]carbamate,

(38) (2S)-2-methoxypropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

(39) (6S)-6-[5-(6-amino-2-fluoro-3-pyridinyl)-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,

5 (40) 2-methoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-methyl-1H-imidazol-5-yl)phenyl]carbamate,

(41) 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(42) 2-(2-methoxyethoxy)ethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

15 (43) 2-(2-methoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(44) 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

20 (45) 2-(2-ethoxyethoxy)ethyl [5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(46) 2-(2-ethoxyethoxy)ethyl [5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-5-yl)-2-pyridinyl]carbamate,

(47) 2-methoxyethyl [4-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

25 (48) 2-methoxyethyl [6-(4-chloro-2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-3-pyridinyl]carbamate,

30 (49) 2-ethoxyethyl [4-(2-((3S)-7-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)phenyl]carbamate,

(50) 2-ethoxyethyl [4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

5 (51) 2-ethoxyethyl [4-(2-((6S)-2-[5-methyl-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)phenyl]carbamate,

(52) tetrahydro-2-furanylmethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

10 (53) methyl [4-(5-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-2-yl)phenyl]carbamate,

(54) 3-hydroxy-3-methylbutyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(55) 2-(2-ethoxyethoxy)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

15 (56) 2-(2-methoxyethoxy)ethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(57) 2-fluoroethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

20 (58) 2-hydroxyethyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(59) 2-hydroxy-2-methylpropyl [4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]carbamate,

(60) (3S)-3-(4-chloro-5-{6-[(1,3-oxazol-2-ylmethyl)amino]-3-pyridinyl})-1H-imidazol-2-yl)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

25 (61) 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4H-1,2,4-triazol-3-yl)phenyl]carbamate,

(62) (3S)-3-[5-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-2-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

30 (63) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])phenyl]acetamide,

(64) methyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,

- (65) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-N-(2-methoxyethoxy)benzamide,
- (66) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-4-fluoro-1H-imidazol-5-yl]phenyl]acetamide,
- 5 (67) 2-methoxyethyl [6-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-4-fluoro-1H-imidazol-5-yl]-3-pyridiny]carbamate,
- (68) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]phenyl]formamide,
- 10 (69) 2-methoxyethyl [6-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-4-fluoro-1H-imidazol-5-yl]-3-pyridiny]carbamate,
- (70) (1E)-N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]phenyl]-N'-hydroxyethanimidamide,
- 15 (71) (6S)-6-[5-(6-amino-3-pyridiny)-4-fluoro-1H-imidazol-2-yl]-2-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-7,8-dihydropyrrolo[1,2-a]pyrimidin-4(6H)-one,
- (72) N-[4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl]phenyl]-2-fluoroacetamide,
- (73) (3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl]-3-[5-(4-hydroxyphenyl)-
- 20 1H-imidazol-2-yl]-2,3-dihydro-5(1H)-indolizinone,
- (74) ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- (75) 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 25 (76) ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- (77) 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (78) 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-
- 30 tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxamide,
- (79) 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)]phenyl)-5-oxo-1,2,3,5-tetrahydro-3-indoliziny)-1H-imidazol-5-yl)-2-thiophenecarboxamide,

- (80) 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,
- (81) 2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-2-yl)phenyl]carbamate,
- 5 (82) 2-(2-methoxyethoxy)ethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,
- (83) 2-methoxyethyl [4-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)phenyl]carbamate,
- 10 (84) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (85) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- (86) 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 15 (87) 5-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (88) 4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- 20 (89) 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,
- (90) 4-(2-((3S)-8-chloro-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-pyridinecarboxylic acid,
- 25 (91) 2-(2-methoxyethoxy)ethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)-2-pyridiny]carbamate,
- (92) 2-(4-morpholinyl)ethyl 5-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,
- 30 (93) 2-(4-morpholinyl)ethyl 5-(4-chloro-2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(94) 2-methoxyethyl [5-(5-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-4-fluoro-1H-imidazol-2-yl)-2-pyridinyl]carbamate,

5 (95) (3S)-3-[2-(6-amino-3-pyridinyl)-4-fluoro-1H-imidazol-5-yl]-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-2,3-dihydro-5(1H)-indolizinone,

(96) 4-(2-((3S)-8-bromo-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

(97) ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

10 (98) 2-(4-morpholinyl)ethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(99) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-4-fluoro-1H-imidazol-5-yl)-2-thiophenecarboxylic acid,

15 (100) isobutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(101) 2-(dimethylamino)-2-oxoethyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

20 (102) (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(103) [(2,2-dimethylpropanoyl)oxy]methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(104) methyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(105) 3-methylbutyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

30 (106) isopropyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-2-thiophenecarboxylate,



(107) 2,3-dihydro-1H-inden-5-yl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(108) phenyl 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny])-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(109) 2-(dimethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(110) methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(111) 2-(diethylamino)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(112) 1-(((cyclohexyloxy)carbonyl)oxy)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(113) cyclohexyl 2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-5-((1-(((cyclohexyloxy)carbonyl)oxy)ethoxy)carbonyl)-3-thienyl))-1H-imidazole-1-carboxylate,

(114) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-3-fluoro-2-thiophenecarboxylic acid,

(115) 4-(5-(7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-4-fluoro-1H-imidazol-2-yl)-2-thiophenecarboxylic acid,

(116) 4-(2-((3S)-7-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-5-oxo-1,2,3,5-tetrahydro-3-indoliziny))-1H-imidazol-5-yl)-3-fluoro-2-pyridinecarboxylic acid,

(117) ((2,2-dimethylpropanoyl)oxy)methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(118) ((2,2-dimethylpropanoyl)oxy)methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl))-1-(((2,2-dimethylpropanoyl)oxy)methyl))-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(119) 2,3-dihydro-1H-inden-5-yl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(120) ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(121) 2-oxo-2-(1-pyrrolidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(122) 2-oxo-2-(1-piperidinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(123) 2-(4-morpholinyl)-2-oxoethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(124) isobutyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(125) isopropyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate,

(126) (5-methyl-2-oxo-1,3-dioxol-4-yl)methyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate, and

(127) 2-(4-morpholinyl)ethyl 4-(2-((6S)-2-[5-chloro-2-(1H-tetrazol-1-yl)phenyl]-4-oxo-4,6,7,8-tetrahydropyrrolo[1,2-a]pyrimidin-6-yl)-1H-imidazol-5-yl)-2-thiophenecarboxylate.

13. A pharmaceutical composition which comprises the compound according to any one of claims 1 to 12, a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.

14. The pharmaceutical composition according to claim 13, which is a factor XIa inhibitor or a factor XIa and plasma kallikrein dual inhibitor.

15. The pharmaceutical composition according to claim 14, which is an agent for the treatment or prevention of a thromboembolic disease.

5           16. The compound according to any one of claims 1 to 12, a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, for use in the treatment of the human or animal body by therapy.

10           17. The compound according to any one of claims 1 to 12, a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, for use in treating or preventing a thromboembolic disease.

15           18. The compound for use according to claim 17, wherein the thromboembolic disease is selected from the group consisting of arterial cardiovascular thromboembolic disorders, venous cardiovascular thromboembolic disorders, arterial cerebrovascular thromboembolic disorders, venous cerebrovascular thromboembolic disorders and thromboembolic disorders in the chambers of the heart or in the peripheral circulation.

20           19. The compound for use according to claim 18, wherein the thromboembolic disease is selected from unstable angina, an acute coronary syndrome, atrial fibrillation, myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, peripheral occlusive arterial disease, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary arterial thrombosis, cerebral arterial thrombosis, cerebral embolism, kidney embolism, pulmonary embolism, and thrombosis resulting from medical implants, devices, or procedures in which blood is exposed to an artificial surface that promotes thrombosis.

30           20. A method for treating a patient suffering from or susceptible to a thromboembolic disease, which comprises administering to said patient a therapeutically effective amount of a compound according to any one of claims 1 to 12, a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof.

21. Use of a compound according to any one of claims 1 to 12, a salt thereof, an N-oxide thereof, a solvate thereof, or a prodrug thereof, in the manufacture of a medicament for use in treating or preventing a thromboembolic disease.