



(51) International Patent Classification:

C12N 15/11 (2006.01) C12Q 1/70 (2006.01)
C12Q 1/68 (2006.01)

(21) International Application Number:

PCT/US2014/061127

(22) International Filing Date:

17 October 2014 (17.10.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/892,796 18 October 2013 (18.10.2013) US

(71) Applicants: NATIONAL TAIWAN UNIVERSITY;

No.1, Sec. 4, Roosevelt Road, Taipei, 10617 (TW).
CHUNG WO, Andrew Man [US/US]; 25 Dodgers Hole
Road, Edgartown, Massachusetts 02539 (US).

(72) Inventors: **CHEN, Pei-Jer**; No. 7, Chung-Shan South

Road, Taipei, 10002 (TW). **YEH, Shiou-Hwei**; No. 7,
Chung-Shan South Road, Taipei, 10002 (TW). **LI, Chiao-**

Ling; No. 7, Chung-Shan South Road, Taipei, 10002
(TW). **CHEN, Ding-Shinn**; No. 7, Chung-Shan South
Road, Taipei, 10002 (TW).

(74) Agents: **WEAVER, Andrew Z.** et al.; 1000 Louisiana

Street, Fifty-Third Floor, Houston, Texas 77002 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,

[Continued on next page]

(54) Title: CIRCULATING CANCER BIOMARKER AND ITS USE




FIG.2

(57) Abstract: A chimera nucleic acid can be obtained from circulatory system for monitoring tumor status. The nucleic acid comprises partial sequence derived from host genome and partial sequence derived from non-host genome. The partial sequence derived from host genome and the partial sequence derived from non-host genome form a chimera junction. The chimera junction is obtained from cell-free nucleic acids and is indicative of disease status.



DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, **Published:**

LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE,

SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,

GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

— *with international search report (Art. 21(3))*