



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C08G 63/668, 18/42	A1	(11) International Publication Number: WO 99/42512 (43) International Publication Date: 26 August 1999 (26.08.99)
(21) International Application Number: PCT/EP99/00837 (22) International Filing Date: 9 February 1999 (09.02.99) (30) Priority Data: 09/025,608 18 February 1998 (18.02.98) US 09/116,112 15 July 1998 (15.07.98) US (71) Applicant (for all designated States except MN): ARCO CHEMICAL TECHNOLOGY, L.P. [US/US]; Suite 238, Two Greenville Crossing, 4001 Kennett Pike, Greenville, DE 19807 (US). (71) Applicant (for MN only): ARCO CHEMIE TECHNOLOGIE NEDERLAND B.V. [NL/NL]; Theemsweg 14, NL-3197 KM Botlek Rotterdam (NL). (72) Inventors: YANG, Lau, S.; 2604 North Gate Road, Wilming- ton, DE 19810 (US). CAI, Gangfeng; 335 Glasgow Circle, Danville, CA 94526 (US). ARMSTEAD, Diandre; 654 Park Lane, Philadelphia, PA 19144 (US). (74) Agent: CROPP, John, Anthony, David; Mathys & Squire, 100 Gray's Inn Road, London WC1X 8AL (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: PROCESS FOR MAKING POLYETHERESTER RESINS WITH HIGH AROMATIC ESTER CONTENT (57) Abstract A process for making polyetherester resins is disclosed, wherein a glycol ester of an aromatic diacid such as terephthalic acid is reacted with a polyether and a source of a dicarboxylic acid such as maleic anhydride to produce a polyetherester resin, wherein the aromatic diester content may be conveniently adjusted as desired. The process permits the preparation of unsaturated polyetherester resins with relatively high levels of terephthalate repeating units while avoiding the difficulties associated with the direct use of terephthalic acid in previously known procedures. Such unsaturated polyetheresters are useful for making thermoset resins with excellent mechanical and physical properties. New glycol esters based on 2-methyl-1,3-propanediol, compositions of such glycol esters and 2-methyl-1,3-propanediol, and polyester resins and thermosets made from the glycol esters, are also disclosed.		