

Description

BENZOTHAIAZOL-2-ON DERIVATIVES AS LIPASE AND PHOSPHOLIPASE INHIBITORS

5

The present invention relates to benzothiazol-2-one derivatives of the general formula I, to their pharmaceutically useful salts and to their use as medicinal substances.


10 Benzothiazolone derivatives of similar structure are described in EP 0334134, EP 0391186 and DE 2101150. However, these are used as pesticides and fungicides.

Compounds having an inhibitory effect on hormone-sensitive lipase are described in
15 the prior art for example in WO2004/035550, WO2005/073199 or WO03/051842.
Compounds with an inhibitory effect on endothelial lipase are described in the prior art for example in WO2004/094394, WO2004/094393.

It is an aim of the present disclosure to provide compounds which have an inhibitory
20 effect on hormone-sensitive lipase or endothelial lipase.

Summary of the Invention

According to a first aspect, there is provided a compound of the formula I



25

wherein the meanings are:

- R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl or aminosulfonyl;
- 5 R2 hydrogen, (C₁-C₆)-alkyl; or
- 10 R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -C(=O)-, -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;
- 15 R3 hydrogen;
- 20 R4, R5 identically or differently hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, (C₁-C₆)-alkylcarbonyl, CO-OR₇, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 25
 30

R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

with the proviso that not more than one of the substituents R₃, R₄, R₅ is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

R7 hydrogen, (C₁-C₆)-alkyl, benzyl;

R8, R8a identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR₇, cyclopropyl, cyclopropylene;

R9, R10 identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;

the tautomeric forms of the compounds and the physiologically tolerated salts thereof.

According to a second aspect, there is provided a medicament comprising one or more compounds of the formula I, in which the meanings are:

R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-

alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl, aminosulfonyl;

5

R2 hydrogen, (C₁-C₆)-alkyl; or

10

R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -C(=O)-, -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;

15

20

R3, R4, R5 identically or differently hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, (C₁-C₆)-alkylcarbonyl, CO-OR₇, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

25

30

R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-

(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

with the proviso that not more than one of the substituents R₃, R₄, R₅ is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

R₇ hydrogen, (C₁-C₆)-alkyl, benzyl;

R₈, R_{8a} identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR₇, cyclopropyl, cyclopropylene;

R₉, R₁₀ identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;

the tautomeric forms of the compounds and the physiologically tolerated salts thereof or one or more compounds of the formula I as defined above.

According to a third aspect, there is provided the use of the compounds of the formula I, in which the meanings are:

R₁ (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl, aminosulfonyl;

- R2 hydrogen, (C₁-C₆)-alkyl; or
- 5 R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -C(=O)-, -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;
- 10 R3, R4, R5 identically or differently hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, (C₁-C₆)-alkylcarbonyl, CO-OR₇, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 15 20 R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 25 R7 hydrogen, (C₁-C₆)-alkyl, benzyl;

1f

R8, R8a identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR7, cyclopropyl, cyclopropylene;

5 R9, R10 identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;

10 the tautomeric forms of the compounds and the physiologically tolerated salts thereof or the compounds of the formula I as defined above for producing a medicament for the treatment and/or prevention of diabetes mellitus .

15 According to a fourth aspect, there is provided the use of the compounds of the formula I as defined above for producing a medicament for the treatment and/or prevention of dyslipidemias .

20 According to a fifth aspect, there is provided the use of the compounds of the formula I as defined above for producing a medicament for the treatment and/or prevention of atherosclerotic disorders.

25 According to a sixth aspect, there is provided a process for producing a medicament comprising one or more of the compounds of the formula I as defined above, which comprises mixing the latter with a pharmaceutically suitable carrier and converting this mixture into a form suitable for administration.



According to a seventh aspect, there is provided a process for preparing compounds of the general formula I as defined above, which comprises benzothiazol-2-one derivatives of the formula II

a) being acylated with carbamoyl chlorides of the formula III;


or

b) in two stages being reacted first with phosgene or equivalents such as trichloromethyl chlorocarbonate, ditrichloromethyl carbonate or 4-nitrophenyl chloroformate and in a second step with amines of the formula IV,

in which the substituents have the meanings indicated above.




10 The disclosure relates to benzothiazol-2-one derivatives of the general formula I



in which the meanings are:

R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where

According to an eighth aspect, there is provided a process for preparing compounds of the general formula I with R2 hydrogen as defined above, which comprises reacting benzothiazol-2-one derivatives of the formula II with isocyanates of the formula V: $O=C=N-R_1$.



- aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl or aminosulfonyl;
- R2 hydrogen, (C₁-C₆)-alkyl; or
- 10 R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series
- 15 -CHR8-, -CR8R8a-, -(C=R8)-, -NR9-, -C(=O)-, -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;
- 20 R3, R4, R5 hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, (C₁-C₆)-alkylcarbonyl, CO-OR7, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR9R10, O-CO-NR9R10, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH,
- 25 O-CO-(C₁-C₆)-alkylene-CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 30 R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR9R10, O-CO-NR9R10, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-

C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

with the proviso that not more than one of the substituents R₃, R₄, R₅ is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

R₇ hydrogen, (C₁-C₆)-alkyl, benzyl;

R₈, R_{8a} identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR₇, cyclopropyl, cyclopropylene;

R₉, R₁₀ identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;

the tautomeric forms of the compounds and the physiologically tolerated salts thereof.

Preferred compounds of the formula I are those in which

R₁ is (C₆-C₁₂)-alkyl, (C₁-C₃)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₄-C₁₂)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, amino, (C₁-C₆)-alkylamino, trifluoromethyl;

R₂ is hydrogen, (C₁-C₆)-alkyl; or

R₁ and R₂ together with the nitrogen atom bearing them are a monocyclic, saturated 5- to 6-membered ring system or a bicyclic saturated or

partially unsaturated 9- to 10-membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR8-, -CR8R8a-, -(C=R8)-, -NR9-, -O-, -S-, with the proviso that two units from the series -O-, -S- may not be adjacent;

5

R3, R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, amino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, COOR7, (C₁-C₆)-alkylsulfonyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₁-C₆)-alkylcarbonyl, CO-NR9R10, O-CO-NR9R10, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

10

15

R6 is hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

20

with the proviso that not more than one of the substituents R3, R4, R5 is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

25 R7 is hydrogen, (C₁-C₆)-alkyl, benzyl;

R8, R8a are identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, COOR7, cyclopropyl, cyclopropylene;

30 R9, R10 are identically or differently hydrogen, (C₁-C₆)-alkyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl.

Preferred compounds of the formula I are also those in which

- 5 R1 is (C₆-C₁₂)-alkyl, (C₁-C₃)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₄-C₁₂)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, amino, (C₁-C₆)-alkylamino, trifluoromethyl;
- 10 R2 is hydrogen, (C₁-C₆)-alkyl; or
- R1 and R2 together with the nitrogen atom bearing them are a monocyclic, saturated 5- to 6-membered ring system or a bicyclic saturated or partially unsaturated 9- to 10 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, (C=R₈)-, -NR₉-, -O-, -S-, with the proviso that two units from the series -O-, -S- may not be adjacent;
- 20 R3, R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, amino, COOR₇, (C₁-C₆)-alkylsulfonyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₁-C₆)-alkylcarbonyl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 25 R6 is hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylamino carbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 30

with the proviso that not more than one of the substituents R³, R⁴, R⁵ is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

- | | | |
|----|---------|--|
| 5 | R7 | is hydrogen, (C ₁ -C ₆)-alkyl, benzyl; |
| | R8, R8a | are identically or differently (C ₁ -C ₆)-alkyl, halogen, trifluoromethyl, COOR7, cyclopropyl, cyclopropylene; |
| 10 | R9, R10 | are identically or differently hydrogen, (C ₁ -C ₆)-alkyl, (C ₆ -C ₁₀)-aryl, (C ₁ -C ₄)-alkylene-(C ₆ -C ₁₀)-aryl. |

Particularly preferred compounds of the formula I are those in which

- 15 R1 is (C₆-C₁₂)-alkyl, (C₁-C₃)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₄-C₁₂)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, amino, (C₁-C₆)-alkylamino, trifluoromethyl;
- 20 R2 is hydrogen, (C₁-C₆)-alkyl; or
- R1 and R2 together with the nitrogen atom bearing them are a monocyclic, saturated 5- to 6-membered ring system or a bicyclic saturated or partially unsaturated 9- to 10 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -O-, -S-, with the proviso that two units from the series -O-, -S- may not be adjacent;
- 30 R3, R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl, (C₁-C₆)-alkyl, hydroxy, amino, COOR₇, (C₁-C₆)-alkylsulfonyl, aminosulfonyl,

pentafluorosulfanyl, (C₁-C₆)-alkylcarbonyl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

5

R₆ is hydrogen, hydroxy, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylamino carbonyl, di-(C₂-C₈)-alkylaminocarbonyl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

10

with the proviso that not more than one of the substituents R₃, R₄, R₅ is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

R₇ is hydrogen, (C₁-C₆)-alkyl;

15

R₈, R_{8a} are identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, cyclopropyl;

R₉, R₁₀ are identically or differently hydrogen, (C₁-C₆)-alkyl.

20

Further particularly preferred compounds of the formula I are those in which

R₁ is (C₆-C₁₂)-alkyl, benzyl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycyl, where benzyl, heteroaryl, cycloalkyl or bicycyl may be substituted by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy or trifluoromethyl;

25

R₂ is hydrogen; or

30 R₁ and R₂ together with the nitrogen atom bearing them form a monocyclic, saturated 5- to 6-membered ring system, of which individual members of the ring systems may be replaced by one to two atoms or atomic

groups from the series -CHR₈-, -NR₉-, in which R₈ is as defined above, and R₉ is (C₁-C₆)-alkyl or cyclopropyl.

Further particularly preferred compounds of the formula I are also those in which

R₁ is (C₆-C₁₂)-alkyl, benzyl, (C₁-C₂)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₂)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycyl, where benzyl, heteroaryl, cycloalkyl or bicycyl may be substituted by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkoxy or trifluoromethyl;

R₂ is hydrogen; or

R₁ and R₂ together with the nitrogen atom bearing them form a monocyclic, saturated 5- to 6-membered ring system, of which individual members of the ring systems may be replaced by one to two atoms or atomic groups from the series -CHR₈-, -NR₉-, in which R₈ is as defined above, and R₉ is (C₁-C₆)-alkyl or cyclopropyl.

Very particularly preferred compounds of the formula I are those in which

R₁ is (C₇-C₁₂)-alkyl, benzyl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl or (C₈-C₁₄)-bicycyl, where benzyl, heteroaryl, cycloalkyl or bicycyl may be substituted by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy or trifluoromethyl;

R₂ is hydrogen;

R₃, R₄, R₅ are identically or differently hydrogen, halogen, hydroxy, (C₁-C₆)-alkyl, trifluoromethyl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy.

Very particularly preferred compounds of the formula I are also those in which

R1 is (C₆-C₁₂)-alkyl, benzyl, CH₂-(C₅-C₁₂)-heteroaryl or (C₈-C₁₄)-bicycle, where benzyl, heteroaryl or bicycle may be substituted by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy or trifluoromethyl;

5 R2 is hydrogen;

R3, R4, R5 are identically or differently hydrogen, halogen, hydroxy, (C₁-C₆)-alkyl, trifluoromethyl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

10

R6 is hydrogen.

Further very particularly preferred compounds of the formula I are those in which

15

R1 is (C₆-C₁₂)-alkyl, benzyl, where benzyl may be substituted by methyl;

R2 is hydrogen;

20 R3 is hydrogen;

R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl;

R6 is hydrogen.

25

Compounds of the formula I in which R3 is hydrogen are a preferred embodiment of the disclosure.

Further preferred compounds of the formula I are those in which R3, R6 is hydrogen, and R4 or R5 are not hydrogen.

30

The invention relates to compounds of the formula I in the form of their salts, racemates, racemic mixtures and pure enantiomers, and to their diastereomers and mixtures thereof.

5 The alkyl radicals in the substituents R1, R2, R3, R4, R5, R6, R7, R8, R8a, R9 and R10 may be either straight-chain or branched. Halogen is fluorine, chlorine, bromine or iodine, in particular fluorine or chlorine.

Aryl means an aromatic carbocyclic mono- or bicyclic ring system which comprises 6 to 10 atoms in the ring or in the rings.

10

Heteroaryl is a mono- or bicyclic aromatic ring system having 5 to 12 ring members, in which at least one atom in the ring system is a heteroatom from the series N, O and S.

Suitable "heteroaryl rings" or "heteroaryl radicals" are, for example,


15 benzimidazolyl, benzofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, quinolinyl, furyl, furazanyl, imidazolyl, 1H-indazolyl, indolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, pyrimidinyl, pyrazinyl, pyrazolyl, pyridyl, pyrrolyl, thiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl,
20 thiophenyl.

Preferred heteroaryl radicals are thiophenyl, pyridyl and pyrazolyl. The heteroaryl rings or heteroaryl radicals may be substituted one or more times by suitable groups.

25 A cycloalkyl radical is a ring system which comprises one or more rings and which is in saturated or partially unsaturated (with one or two double bonds) form and which is composed exclusively of carbon atoms. The cycloalkyl radicals may be substituted one or more times by suitable groups.

30 Bicycle means bicyclic ring systems which are in saturated or partially unsaturated form and which, apart from carbon, may also comprise one or more heteroatoms such as, for example, nitrogen, oxygen or sulfur. This definition also includes ring systems which comprise a fused benzene nucleus. Examples which may be

mentioned are the tetrahydronaphthyl, alpha- or beta-tetralon-, indanyl- or indan-1-on-yl radical. Preferred bicyclic radicals are those of formula Ic



- 5
- Pharmaceutically acceptable salts are, because their solubility in water is greater than that of the initial or basic compounds, particularly suitable for medical applications. These salts must have a pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the
- 10 invention are salts of inorganic acids such as hydrochloric acid, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acid, and of organic acids such as, for example, acetic acid, benzenesulfonic, benzoic, citric, ethanesulfonic, frumaric, gluconic, glycolic, isethionic, lactic, lactobionic, maleic, malic, methanesulfonic, succinic, p-toluenesulfonic and tartaric acid. Suitable pharmaceutically acceptable
- 15 basic salts are ammonium salts, alkali metal salts (such as sodium and potassium salts) and alkaline earth metal salts (such as magnesium and calcium salts) and salts of trometamol (2-amino-2-hydroxymethyl-1,3-propanediol), diethanolamine, lysine or ethylenediamine.
- 20 Salts with a pharmaceutically unacceptable anion such as, for example, trifluoroacetate likewise belong within the framework of the disclosure as useful intermediates for the preparation or purification of pharmaceutically acceptable salts and/or for use in nontherapeutic, for example in vitro, applications.
- 25 The term "physiologically functional derivative" used herein refers to any physiologically tolerated derivative of a compound of the disclosure of the formula I, for example an ester, which on administration to a mammal such as, for example, a human is able to form (directly or indirectly) a compound of the formula I or an active metabolite thereof.

Physiologically functional derivatives also include prodrugs of the compounds of the invention as, for example, described in H. Okada et al., Chem. Pharm. Bull. 1994, 42, 57-61. Such prodrugs can be metabolized in vivo to a compound of the disclosure. These prodrugs may themselves be active or not.

5

The compounds of the disclosure may also exist in various polymorphous forms, for example as amorphous and crystalline polymorphous forms. All polymorphous forms of the compounds of the disclosure belong within the framework of the disclosure and are a further embodiment of the disclosure.

10

All references to "compound(s) of formula I" hereinafter refer to compound(s) of the formula I as described above, and their salts, solvates and physiologically functional derivatives as described herein.

15

Use

The compounds of the invention of the general formula I in which the meanings are:

20 R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, 25 mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl, aminosulfonyl;

R2 hydrogen, (C₁-C₆)-alkyl; or

30

R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a

bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR8-, -CR8R8a-, -(C=R8)-, -NR9-, -C(=O)-, -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;

- 5 R3, R4, R5 identically or differently hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, -(C₁-C₆)-alkylcarbonyl, CO-OR7, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR9R10, O-CO-NR9R10, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 10
- 15
- R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR9R10, O-CO-NR9R10, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 20
- 25 R7 hydrogen, (C₁-C₆)-alkyl, benzyl;
- R8, R8a identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR7, cyclopropyl, cyclopropylene;
- 30 R9, R10 identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-

alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;

the tautomeric forms of the compounds and the physiologically tolerated salts thereof
 5 have a surprising inhibitory effect on hormone sensitive lipase, HSL, an allosteric enzyme in adipocytes which is inhibited by insulin and is responsible for the breakdown of fats in fat cells and thus for transferring fat constituents into the blood stream. Inhibition of this enzyme is therefore equivalent to an insulin-like effect of the compounds of the invention, eventually leading to a reduction of free fatty acids in
 10 the blood and of blood glucose. They can therefore be employed for metabolic derangements such as, for example, for non-insulin-dependent diabetes mellitus, for diabetic syndrome and for direct pancreatic damage.

The compounds of the invention of the general formula I, especially those in which
 15 R₂ is hydrogen, may additionally have an inhibitory effect on endothelial lipase (EL). The preferred substrate for EL is HDL, which has antiatherosclerotic activity. A reduction in the HDL level leads to progression of atherosclerosis and its sequelae such as coronary heart disease and moreover favors development of the metabolic syndrome and its sequelae diabetes. An inhibition of EL should thus generally lead
 20 to prevention of atherosclerotic disorders and indirectly reduce the probability of people with an increased risk for diabetes becoming ill.

It has further been found that the inhibitory effect of the compounds of the invention of the general formula I is selective in relation to other lipases.

25 The compounds of the invention of the formula I may also have an inhibitory effect on triglyceride lipase.

Compounds of this type are particularly suitable for the treatment and/or prevention
 30 of.

1. - Disorders of fatty acid metabolism and glucose utilization disorders
2. - Disorders of the insulin sensitivity of myo-, adipo- and hepatocytes (insulin resistance) - metabolic syndrome
3. Diabetes mellitus, especially type 2 diabetes, including the prevention of the
5 sequelae associated therewith.
Particular aspects in this connection are
 - hyperglycemia,
 - improvement in insulin resistance,
 - improvement in glucose tolerance,
 - 10 - protection of the pancreatic β cells
 - prevention of macro- and microvascular disorders
4. Dyslipidemias and the sequelae thereof such as, for example, atherosclerosis,
15 coronary heart disease, cerebrovascular disorders etc., especially those (but not
restricted thereto) which are characterized by one or more of the following
factors:
 - high plasma triglyceride concentrations, high postprandial plasma triglyceride
concentrations
 - low HDL cholesterol concentration
 - 20 - low apoA lipoprotein concentrations
 - high LDL cholesterol concentrations
 - small dense LDL cholesterol particles
 - high apoB lipoprotein concentrations
- 25 5. Various other conditions which may be associated with the metabolic syndrome,
such as:
 - obesity (excess weight), including central obesity
 - thromboses, hypercoagulable and prothrombotic stages (arterial and venous)
 - high blood pressure
 - 30 - heart failure such as, for example (but not restricted thereto), following
myocardial infarction, hypertensive heart disease or cardiomyopathy

6. Other disorders or conditions in which inflammatory reactions or cell differentiation may for example be involved are:

- atherosclerosis such as, for example (but not restricted thereto), coronary sclerosis including angina pectoris or myocardial infarction, stroke
- 5 - vascular restenosis or reocclusion
- chronic inflammatory bowel diseases such as, for example, Crohn's disease and ulcerative colitis
- pancreatitis
- other inflammatory states
- 10 - retinopathy
- adipose cell tumors
- adipose cell carcinomas such as, for example, liposarcoma
- solid tumors and neoplasms such as, for example (but not restricted thereto), carcinomas of the gastrointestinal tract, of the liver, of the biliary tract and of
- 15 the pancreas, endocrine tumors, carcinomas of the lungs, of the kidneys and of the urinary tract, of the genital tract, prostate carcinomas etc.
- acute and chronic myeloproliferative disorders and lymphomas
- angiogenesis
- neurodegenerative disorders
- 20 - Alzheimer's disease
- multiple sclerosis
- Parkinson's disease
- erythemato-squamous dermatoses such as, for example, psoriasis
- acne vulgaris
- 25 - other skin disorders and dermatological conditions which are modulated by PPAR
- eczemas and neurodermatitis
- dermatitis such as, for example, seborrheic dermatitis or photodermatitis
- keratitis and keratosis such as, for example, seborrheic keratoses, senile
- 30 keratoses, actinic keratosis, photo-induced keratoses or keratosis follicularis
- keloids and keloid prophylaxis
- warts, including condylomata or condylomata acuminata

- human papilloma viral (HPV) infections, such as, for example, venereal papillomata, viral warts such as, for example, molluscum contagiosum, leukoplakia
- papular dermatoses such as, for example, lichen planus
- 5 - skin cancer such as, for example, basal cell carcinoma, melanomas or cutaneous T-cell lymphomas
- localized benign epidermal tumors such as, for example, keratoderma, epidermal naevi
- chilblains
- 10 - high blood pressure
- syndrome X
- polycystic ovary syndrome (PCOS)
- asthma
- osteoarthritis
- 15 - lupus erythematosus (LE) or inflammatory rheumatic disorders such as, for example, rheumatoid arthritis
- vasculitis
- wasting (cachexia)
- gout
- 20 - ischemia/reperfusion syndrome
- acute respiratory distress syndrome (ARDS)

Compounds which inhibit endothelial lipase are particularly suitable for the treatment
25 and/or prevention of

1. Dyslipidemias and general impairments of lipid metabolism and their sequelae such as, for example, atherosclerosis, coronary heart disease, cerebrovascular disorders etc., especially those (but not restricted thereto) which are characterized by one or more of the following factors:
- 30 - high plasma triglyceride concentrations, high postprandial plasma triglyceride concentrations,

- low HDL cholesterol concentration
 - low ApoA lipoprotein concentrations
 - high LDL cholesterol concentrations
 - small dense LDL cholesterol particles
 - 5 - high ApoB lipoprotein concentrations
2. Various other conditions which may be associated with the metabolic syndrome, such as:
- obesity (excess weight), including central obesity
 - 10 - thromboses, hypercoagulable and prothrombotic stages (arterial and venous)
 - high blood pressure
 - heart failure such as, for example (but not restricted thereto), following myocardial infarction, hypertensive heart disease or cardiomyopathy
 - diabetes mellitus, in particular type 2 diabetes including the prevention of the
 - 15 sequelae associated therewith (hyperglycemia, glucose intolerance, loss of pancreatic β cells, macro- and microvascular disorders
3. Other disorders or conditions in which inflammatory reactions or cell differentiation may for example be involved are:
- 20 - atherosclerosis such as, for example (but not restricted thereto), coronary sclerosis including angina pectoris or myocardial infarction, stroke
 - vascular restenosis or reocclusion
 - chronic inflammatory bowel diseases such as, for example, Crohn's disease and ulcerative colitis
 - 25 - pancreatitis
 - other inflammatory states
 - retinopathy
 - adipose cell tumors
 - adipose cell carcinomas such as, for example, liposarcomas
 - 30 - solid tumors and neoplasms such as, for example (but not restricted thereto), carcinomas of the gastrointestinal tract, of the liver, of the biliary tract and of the pancreas, endocrine tumors, carcinomas of the lungs, of the kidneys and

the urinary tract, of the genital tract, prostate carcinomas etc.

- acute and chronic myeloproliferative disorders and lymphomas
- angiogenesis
- neurodegenerative disorders
- 5 - Alzheimer's disease
- multiple sclerosis
- Parkinson's disease
- erythemato-squamous dermatoses such as, for example, psoriasis
- acne vulgaris
- 10 - other skin disorders and dermatological conditions which are modulated by PPAR
- eczemas and neurodermatitis
- dermatitis such as, for example, seborrheic dermatitis or photodermatitis
- keratitis and keratoses such as, for example, seborrheic keratoses, senile
- 15 keratoses, actinic keratosis, photo-induced keratoses or keratosis follicularis
- keloids and keloid prophylaxis
- warts, including condylomata or condylomata acuminata
- human papilloma viral (HPV) infections such as, for example, venereal papillomata, viral warts such as, for example, molluscum contagiosum,
- 20 leukoplakia
- papular dermatoses such as, for example, lichen planus
- skin cancer such as, for example, basal-cell carcinomas, melanomas or cutaneous T-cell lymphomas
- localized benign epidermal tumors such as, for example, keratoderma,
- 25 epidermal naevi
- chilblains
- high blood pressure
- syndrome X
- polycystic ovary syndrome (PCOS)
- 30 - asthma
- osteoarthritis
- lupus erythematosus (LE) or inflammatory rheumatic disorders such as, for

example, rheumatoid arthritis

- vasculitis
- wasting (cachexia)
- gout
- 5 - ischemia/reperfusion syndrome
- acute respiratory distress syndrome (ARDS)

Formulations

- 10 The amount of a compound of the invention necessary to achieve the desired biological effect depends on a number of factors, for example the specific compound chosen, the intended use, the mode of administration and the clinical condition of the patient. The daily dose is generally in the range from 0.3 mg to 100 mg (typically from 3 mg to 50 mg) per day and per kilogram of body weight, for example
- 15 3-10 mg/kg/day. An intravenous dose may be, for example, in the range from 0.3 mg to 1.0 mg/kg, which can suitably be administered as infusion of 10 ng to 100 ng per kilogram and per minute. Suitable infusion solutions for these purposes may contain, for example, from 0.1 ng to 10 mg, typically from 1 ng to 10 mg, per milliliter. Single
- 20 doses may contain, for example, from 1 mg to 10 g of the active ingredient. Thus, ampoules for injections may contain, for example, from 1 mg to 100 mg, and single-dose formulations which can be administered orally, such as, for example, tablets or capsules, may contain, for example, from 0.05 to 1000 mg, typically from 0.5 to 600 mg. For the therapy of the abovementioned conditions, the compounds of
- 25 formula I may be used as the compound itself, but they are preferably in the form of a pharmaceutical composition with an acceptable carrier. The carrier must, of course, be acceptable in the sense that it is compatible with the other ingredients of the composition and is not harmful for the patient's health. The carrier may be a solid or a liquid or both and is preferably formulated with the compound as a single dose, for example as a tablet, which may contain from 0.05% to 95% by weight of the
- 30 active ingredient. Other pharmaceutically active substances may likewise be present, including other compounds of the invention. The pharmaceutical compositions of the invention can be produced by one of the known pharmaceutical methods, which

essentially consist of mixing the ingredients with pharmacologically acceptable carriers and/or excipients.

Pharmaceutical compositions of the invention are those suitable for oral, rectal, topical, peroral (for example sublingual) and parenteral (for example subcutaneous, intramuscular, intradermal or intravenous) administration, although the most suitable mode of administration depends in each individual case on the nature and severity of the condition to be treated and on the nature of the compound of formula I used in each case. Coated formulations and coated slow-release formulations also belong within the framework of the invention. Preference is given to acid- and gastric juice-resistant formulations. Suitable coatings resistant to gastric juice comprise cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropylmethylcellulose phthalate and anionic polymers of methacrylic acid and methyl methacrylate.

Suitable pharmaceutical preparations for oral administration may be in the form of separate units such as, for example, capsules, cachets, suckable tablets or tablets, each of which contain a defined amount of the compound of formula I; as powders or granules; as solution or suspension in an aqueous or nonaqueous liquid; or as an oil-in-water or water-in-oil emulsion. These compositions may, as already mentioned, be prepared by any suitable pharmaceutical method which includes a step in which the active ingredient and the carrier (which may consist of one or more additional ingredients) are brought into contact. The compositions are generally produced by uniform and homogeneous mixing of the active ingredient with a liquid and/or finely divided solid carrier, after which the product is shaped if necessary. Thus, for example, a tablet can be produced by compressing or molding a powder or granules of the compound, where appropriate with one or more additional ingredients. Compressed tablets can be produced by tableting the compound in free-flowing form such as, for example, a powder or granules, where appropriate mixed with a binder, glidant, inert diluent and/or one or more surface-active/dispersing agent(s) in a suitable machine. Molded tablets can be produced by molding the compound, which is in powder form and is moistened with an inert liquid diluent, in a suitable machine.

Pharmaceutical compositions which are suitable for peroral (sublingual) administration comprise suckable tablets which contain a compound of formula I with a flavoring, normally sucrose and gum arabic or tragacanth, and pastilles which comprise the compound in an inert base such as gelatin and glycerol or sucrose and gum arabic.

Pharmaceutical compositions suitable for parenteral administration comprise preferably sterile aqueous preparations of a compound of formula I, which are preferably isotonic with the blood of the intended recipient. These preparations are preferably administered intravenously, although administration may also take place by subcutaneous, intramuscular or intradermal injection. These preparations can preferably be produced by mixing the compound with water and making the resulting solution sterile and isotonic with blood. Injectable compositions of the invention generally contain from 0.1 to 5% by weight of the active compound.

Pharmaceutical compositions suitable for rectal administration are preferably in the form of single-dose suppositories. These can be produced by mixing a compound of the formula I with one or more conventional solid carriers, for example cocoa butter, and shaping the resulting mixture.

Pharmaceutical compositions suitable for topical use on the skin are preferably in the form of ointment, cream, lotion, paste, spray, aerosol or oil. Carriers which can be used are petrolatum, lanolin, polyethylene glycols, alcohols and combinations of two or more of these substances. The active ingredient is generally present in a concentration of from 0.1 to 15% by weight of the composition, for example from 0.5 to 2%.

Transdermal administration is also possible. Pharmaceutical compositions suitable for transdermal uses can be in the form of single patches which are suitable for long-term close contact with the patient's epidermis. Such patches suitably contain the active ingredient in an aqueous solution which is buffered where appropriate, dissolved and/or dispersed in an adhesive or dispersed in a polymer. A suitable

active ingredient concentration is about 1% to 35%, preferably about 3% to 15%. A particular possibility is for the active ingredient to be released by electrotransport or iontophoresis as described, for example, in *Pharmaceutical Research*, 2(6): 318 (1986).

5

The compounds of the formula I with R² hydrogen are distinguished by favorable effects on disorders of lipid metabolism. They beneficially influence the HDL to LDL ratio and increase in particular the HDL level and are suitable for the prevention and treatment of dyslipidemias and metabolic syndrome and their diverse sequelae such as atherosclerosis, coronary heart disease, heart failure, obesity and diabetes.

10

Combinations with other medicaments

The compounds of the invention can be administered alone or in combination with one or more further pharmacologically active substances which have, for example, favorable effects on metabolic disturbances or disorders frequently associated therewith. Examples of such medicaments are

15

1. medicaments which lower blood glucose, antidiabetics,
2. active ingredients for the treatment of dyslipidemias,
- 20 3. antiatherosclerotic medicaments,
4. antiobesity agents,
5. antiinflammatory active ingredients
6. active ingredients for the treatment of malignant tumors
7. antithrombotic active ingredients
- 25 8. active ingredients for the treatment of high blood pressure
9. active ingredients for the treatment of heart failure and
10. active ingredients for the treatment and/or prevention of complications caused by diabetes or associated with diabetes.

25

30 They can be combined with the compounds of the invention of the formula I in particular for a synergistic improvement in the effect. Administration of the active ingredient combination can take place either by separate administration of the active

ingredients to the patient or in the form of combination products in which a plurality of active ingredients are present in one pharmaceutical preparation.

Examples which may be mentioned are:

5

Antidiabetics

Suitable antidiabetics are disclosed for example in the Rote Liste 2001, chapter 12 or in the USP Dictionary of USAN and International Drug Names, US Pharmacopeia, 10 Rockville 2003. Antidiabetics include all insulins and insulin derivatives such as, for example, Lantus® (see www.lantus.com) or Apidra®, and other fast-acting insulins (see US 6,221,633), GLP-1 receptor modulators as described in WO 01/04146 or else, for example, those disclosed in WO 98/08871 of Novo Nordisk A/S.

The orally effective hypoglycemic active ingredients include, preferably, the 15 sulfonylureas which act on the ATP-dependent potassium channel of the beta cells (e.g. disclosed in WO 97/26265 and WO 99/03861), biguanides, meglitinides, glucagon antagonists, oral GLP-1 agonists, DPP-IV inhibitors, insulin sensitizers, e.g. PPAR and PXR modulators and active ingredients such as, for example, oxadiazolidinediones, thiazolidinediones, inhibitors of liver enzymes which are 20 involved in stimulating gluconeogenesis and/or glycogenolysis, modulators of glucose uptake such as, for example, glucosidase inhibitors, compounds which alter lipid metabolism and lead to a change in the blood lipid composition, compounds which reduce food intake or food uptake.

25 In one embodiment of the invention, the compounds of the formula I are administered in combination with insulin.

In one embodiment of the invention, the compounds of the formula I are administered in combination with substances which influence hepatic glucose production such as, for example, glycogen phosphorylase inhibitors (see: 30 WO 01/94300, WO 02/096864, WO 03/084923, WO 03/084922, WO 03/104188).

In one embodiment, the compounds of the formula I are administered in combination

with an active ingredient which acts on the ATP-dependent potassium channel of the beta cells, such as, for example, sulfonylureas (e.g. tolbutamide, glibenclamide, glipizide, glimepiride) or glinides (e.g. repaglinide).

5 In one embodiment, the compounds of the formula I are administered in combination with a biguanide such as, for example, metformin.

In one embodiment, the compounds of the formula I are administered in combination with a PPARgamma agonist or thiazolidinedione such as, for example, ciglitazone, pioglitazone, rosiglitazone or the compounds disclosed in WO 97/41097 of
10 Dr. Reddy's Research Foundation, in particular 5-[[4-[(3,4-dihydro-3-methyl-4-oxo-2-quinazolinylmethoxy)phenyl]methyl]-2,4-thiazolidinedione.

In one embodiment, the compounds of the formula I are administered in combination with a DPPIV inhibitor as described, for example, in WO98/19998, WO99/61431,
15 WO99/67278, WO99/67279, WO01/72290, WO 02/38541, WO03/040174, in particular P 93/01 (1-cyclopentyl-3-methyl-1-oxo-2-pentan ammonium chloride), P-31/98, LAF237 (1-[2-[3-hydroxyadamant-1-ylamino)acetyl]pyrrolidine-2-(S)-carbonitrile), TS021 ((2S, 4S)-4-fluoro-1-[[2-hydroxy-1,1-dimethylethyl)amino]-acetyl]pyrrolidine-2-carbonitrile monobenzenesulfonate).

20 In one embodiment, the compounds of the formula I are administered in combination with compounds with an inhibitory effect on SGLT-1 and/or 2, as disclosed directly or indirectly for example in WO2004/007517, WO2004/052902, WO2004/052903 and WO2005/121161.

25 In one embodiment, the compounds of the formula I are administered in combination with an α -glucosidase inhibitor such as, for example, miglitol or acarbose.

In one embodiment, the compounds of the formula I are administered in combination with more than one of the aforementioned compounds, e.g. in combination with a
30 sulfonylurea and metformin, a sulfonylurea and acarbose, repaglinide and metformin, insulin and a sulfonylurea, insulin and metformin, insulin and troglitazone, insulin and lovastatin, etc.

Lipid modulators

In one embodiment of the invention, the compounds of the formula I are
5 administered in combination with an HMGCoA reductase inhibitor such as lovastatin, fluvastatin, pravastatin, simvastatin, ivastatin, itavastatin, atorvastatin, rosuvastatin.

In one embodiment of the invention, the compounds of the formula I are
administered in combination with a bile acid absorption inhibitor (see, for example,
10 US 6,245,744, US 6,221,897, US 6,277,831, EP 0683 773, EP 0683 774).

In one embodiment of the invention, the compounds of the formula I are
administered in combination with a polymeric bile acid adsorbent such as, for
example, cholestyramine, colesevelam.

15 In one embodiment of the invention, the compounds of the formula I are
administered in combination with a cholesterol resorption inhibitor as described for
example in WO 02/50027, or ezetimibe, tiqueside, pamaqueside.

20 In one embodiment of the invention, the compounds of the formula I are
administered in combination with an LDL receptor inducer (see, for example,
US 6,342,512).

In one embodiment, the compounds of the formula I are administered in combination
25 with bulking agents, preferably insoluble bulking agents (see, for example,
carob/Caromax® (Zunft H J; et al., Carob pulp preparation for treatment of
hypercholesterolemia, ADVANCES IN THERAPY (2001 Sep-Oct), 18(5), 230-6)).
Caromax is a carob-containing product from Nutrinova, Nutrition Specialties & Food
Ingredients GmbH, Industriepark Hoechst, 65926 Frankfurt/Main). Combination with
30 Caromax® is possible in one preparation or by separate administration of
compounds of the formula I and Caromax®. Caromax® can in this connection also
be administered in the form of food products such as, for example, in bakery

products or muesli bars.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a PPARalpha agonist.

5

In one embodiment of the invention, the compounds of the formula I are administered in combination with a mixed PPAR alpha/gamma agonist such as, for example, AZ 242 (tesaglitazar, (S)-3-(4-[2-(4-methanesulfonyloxyphenyl)ethoxy]-phenyl)-2-ethoxypropionic acid), BMS 298585 (N-[(4-methoxyphenoxy)carbonyl]-N-
10 [[4-[2-(5-methyl-2-phenyl-4-oxazolyl)ethoxy]phenyl]methyl]glycine) or as described in WO 99/62872, WO 99/62871, WO 01/40171, WO 01/40169, WO96/38428, WO 01/81327, WO 01/21602, WO 03/020269, WO 00/64888 or WO 00/64876.

In one embodiment of the invention, the compounds of the formula I are
15 administered in combination with a fibrate such as, for example, fenofibrate, gemfibrozil, clofibrate, bezafibrate.

In one embodiment of the invention, the compounds of the formula I are administered in combination with nicotinic acid or niacin.

20

In one embodiment of the invention, the compounds of the formula I are administered in combination with a CETP inhibitor, e.g. CP- 529, 414 (torcetrapib).

In one embodiment of the invention, the compounds of the formula I are
25 administered in combination with an ACAT inhibitor.

In one embodiment of the invention, the compounds of the formula I are administered in combination with an MTP inhibitor such as, for example, implitapide.

30 In one embodiment of the invention, the compounds of the formula I are administered in combination with an antioxidant.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a lipoprotein lipase inhibitor.

5 In one embodiment of the invention, the compounds of the formula I are administered in combination with an ATP citrate lyase inhibitor.

In one embodiment of the invention, the compounds of the formula I are administered in combination with a squalene synthetase inhibitor.

10 In one embodiment of the invention, the compounds of the formula I are administered in combination with a lipoprotein(a) antagonist.

Antiobesity agents

15 In one embodiment of the invention, the compounds of the formula I are administered in combination with a lipase inhibitor such as, for example, orlistat.

In one embodiment, the further active ingredient is fenfluramine or dexfenfluramine.
In another embodiment, the further active ingredient is sibutramine.

20 In a further embodiment, the compounds of the formula I are administered in combination with CART modulators (see "Cocaine-amphetamine-regulated transcript influences energy metabolism, anxiety and gastric emptying in mice" Asakawa, A, et al., M.: Hormone and Metabolic Research (2001), 33(9), 554-558),

25 NPY antagonists, e.g. naphthalene-1-sulfonic acid {4-[(4-aminoquinazolin-2-ylamino)methyl]cyclohexylmethyl}amide; hydrochloride (CGP 71683A)), MC4 agonists (e.g. 1-amino-1,2,3,4-tetrahydronaphthalene-2-carboxylic acid [2-(3a-benzyl-2-methyl-3-oxo-2,3,3a,4,6,7-hexahydropyrazolo[4,3-c]pyridin-5-yl)-1-(4-chlorophenyl)-2-oxoethyl]amide; (WO 01/91752)), orexin antagonists (e.g. 1-(2-

30 methylbenzoxazol-6-yl)-3-[1,5]naphthyridin-4-ylurea; hydrochloride (SB-334867-A)), H3 agonists (3-cyclohexyl-1-(4,4-dimethyl-1,4,6,7-tetrahydroimidazo[4,5-c]pyridin-5-yl)propan-1-one oxalic acid salt (WO 00/63208)); TNF agonists, CRF antagonists

(e.g. [2-methyl-9-(2,4,6-trimethylphenyl)-9H-1,3,9-triazafluoren-4-yl]dipropylamine (WO 00/66585)), CRF BP antagonists (e.g. urocortin), urocortin agonists, $\beta 3$ agonists (e.g. 1-(4-chloro-3-methanesulfonylmethylphenyl)-2-[2-(2,3-dimethyl-1H-indol-6-yloxy)ethylamino]ethanol; hydrochloride (WO 01/83451)), MSH (melanocyte-stimulating hormone) agonists, CCK-A agonists (e.g. {2-[4-(4-chloro-2,5-dimethoxyphenyl)-5-(2-cyclohexylethyl)thiazol-2-ylcarbamoyl]-5,7-dimethylindol-1-yl}acetic acid trifluoroacetic acid salt (WO 99/15525)), serotonin reuptake inhibitors (e.g. dexfenfluramine), mixed serotoninergic and noradrenergic compounds (e.g. WO 00/71549), 5HT agonists e.g. 1-(3-ethylbenzofuran-7-yl)piperazine oxalic acid salt (WO 01/09111), bombesin agonists, galanin antagonists, growth hormone (e.g. human growth hormone), growth hormone-releasing compounds (6-benzyloxy-1-(2-diisopropylaminoethylcarbamoyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tertiary butyl ester (WO 01/85695)), TRH agonists (see, for example, EP 0 462 884), uncoupling protein 2 or 3 modulators, leptin agonists (see, for example, Lee, Daniel W.; Leinung, Matthew C.; Rozhavskaya-Arena, Marina; Grasso, Patricia. Leptin agonists as a potential approach to the treatment of obesity. *Drugs of the Future* (2001), 26(9), 873-881), DA agonists (bromocriptine, Doprexin), lipase/amylase inhibitors (e.g. WO 00/40569), PPAR modulators (e.g. WO 00/78312), RXR modulators or TR- β agonists.

In another embodiment, the further active ingredient is a cannabinoid receptor 1 antagonist (such as, for example, rimonabant, SR147778 or those as are described in, for example, EP 0656354, WO 00/15609, WO 02/076949, WO2005/080345, WO2005/080328, WO2005/080343, WO2005/075450, WO2005/080357, WO2001/70700, WO2003/026647-48, WO2003/02776, WO2003/040107, WO2003/007887, WO2003/027069, US6,509,367, WO2001/32663, WO2003/086288, WO2003/087037, WO2004/048317, WO2004/058145, WO2003/084930, WO2003/084943, WO2004/058744, WO2004/013120, WO2004/029204, WO2004/035566, WO2004/058249, WO2004/058255, WO2004/058727, WO2004/069838, US2004/0214837, US2004/0214855, US2004/0214856, WO2004/096209, WO2004/096763, WO2004/096794, WO2005/000809, WO2004/099157, US2004/0266845, WO2004/110453,

WO2004/108728, WO2004/000817, WO2005/000820, US2005/0009870, WO2005/00974, WO2004/111033-34, WO2004/11038-39, WO2005/016286, WO2005/007111, WO2005/007628, US2005/0054679, WO2005/027837, WO2005/028456, WO2005/063761-62, WO2005/061509, WO2005/077897).

5

In one embodiment of the invention, the further active ingredient is leptin.

In one embodiment, the further active ingredient is dexamphetamine, amphetamine, mazindole or phentermine.

10 In one embodiment, the compounds of the formula I are administered in combination with medicaments having effects on the coronary circulation and the vascular system, such as, for example, ACE inhibitors (e.g. ramipril), medicaments which act on the angiotensin-renaline system, calcium antagonists, beta blockers etc.

15 In one embodiment, the compounds of the formula I are administered in combination with medicaments having an antiinflammatory effect.

In one embodiment, the compounds of the formula I are administered in combination with medicaments which are employed for cancer therapy and cancer prevention.

20

It will be appreciated that every suitable combination of the compounds of the invention with one or more of the aforementioned compounds and optionally one or more other pharmacologically active substances is regarded as falling within the protection conferred by the present invention.

25

The activity of the compounds of the invention of the formula I was tested in the following enzyme assay systems:

30 1. HSL inhibition assay

1.1. Preparation of the partially purified HSL:

Isolated rat fat cells are obtained from epididymal adipose tissue from untreated male rats (Wistar, 220-250 g) by collagenase treatment in accordance with published methods (e.g. S. Nilsson et al., *Anal. Biochem.* 158, 1986, 399-407; G. Fredrikson et al., *J. Biol. Chem.* 256, 1981, 6311-6320; H. Tornquist et al., *J. Biol. Chem.* 251, 1976, 813-819). The fat cells from 10 rats are washed three times by flotation with 50 ml of homogenization buffer (25 mM Tris/HCl, pH 7.4, 0.25 M sucrose, 1 mM ETDA, 1 mM DTT, 10 µg/ml leupeptin, 10 µg/ml antipain, 20 µg/ml pepstatin) each time and finally taken up in 10 ml of homogenization buffer. The fat cells are homogenized in a Teflon-in-glass homogenizer (Braun-Melsungen) by 10 strokes at 1500 rpm and 15°C. The homogenate is centrifuged (Sorvall SM24 tubes, 5000 rpm, 10 min, 4°C). The subnatant between the layer of fat at the top and the pellet is removed and the centrifugation is repeated. The subnatant resulting therefrom is centrifuged again (Sorvall SM24 tubes, 20 000 rpm, 45 min, 4°C). The subnatant is removed, and 1 g of heparin-Sepharose (Pharmacia-Biotech, CL-6B, washed 5× with 25 mM Tris/HCl, pH 7.4, 150 mM NaCl) is added. After incubation at 4°C for 60 min (shaking at intervals of 15 min), the mixture is centrifuged (Sorvall SM24 tubes, 3000 rpm, 10 min, 4°C). The supernatant is adjusted to pH 5.2 by adding glacial acetic acid and is incubated at 4°C for 30 min. The precipitates are collected by centrifugation (Sorvall SS34, 12 000 rpm, 10 min, 4°C) and suspended in 2.5 ml of 20 mM Tris/HCl, pH 7.0, 1 mM EDTA, 65 mM NaCl, 13% sucrose, 1 mM DTT, 10 µg/ml leupeptin/pepstatin/antipain. The suspension is dialyzed against 25 mM Tris/HCl, pH 7.4, 50% glycerol, 1 mM DTT, 10 µg/ml leupeptin, pepstatin, antipain at 4°C overnight and then loaded onto a hydroxiapatite column (0.1 g per 1 ml of suspension, equilibrated with 10 mM potassium phosphate, pH 7.0, 30% glycerol, 1 mM DTT). The column is washed with four volumes of equilibration buffer at a flow rate of 20 to 30 ml/h. The HSL is eluted with one volume of equilibration buffer containing 0.5 M potassium phosphate and then dialyzed (see above) and concentrated 5- to 10-fold by ultrafiltration (Amicon Diaflo PM 10 Filter) at 4°C. The partially purified HSL can be stored at -70°C for 4 to 6 weeks.

1.2 HSL activity assay:

To prepare the substrate, 25-50 μCi of $[^3\text{H}]$ trioleoylglycerol (in toluene), 6.8 μmol of unlabeled trioleoylglycerol and 0.6 mg of phospholipids (phosphatidylcholine/phosphatidylinositol 3:1 w/v) are mixed, dried with N_2 and then taken up in 2 ml of 0.1 M KPi (pH 7.0) by ultrasound treatment (Branson 250, microtip, setting 1-2, 2×1 min with an interval of 1 min). After addition of 1 ml of KPi and renewed ultrasound treatment (4×30 sec on ice with intervals of 30 sec), 1 ml of 20% BSA (in KPi) is added (final concentration of trioleoylglycerol 1.7 mM). For the reaction, 100 μl of substrate solution are pipetted into 100 μl of HSL solution (HSL prepared as above, diluted in 20 mM KPi, pH 7.0, 1 mM EDTA, 1 mM DTT, 0.02% BSA, 20 $\mu\text{g}/\text{ml}$ pepstatin, 10 $\mu\text{g}/\text{ml}$ leupeptin) and incubated at 37°C for 30 min. Addition of 3.25 ml of methanol/chloroform/heptane (10:9:7) and of 1.05 ml of 0.1 M K_2CO_3 , 0.1 M boric acid (pH 10.5) is followed by thorough mixing and finally centrifugation ($800 \times g$, 20 min). After phase separation, one equivalent of the upper phase (1 ml) is removed and the radioactivity is determined by liquid scintillation measurement.

1.3 Evaluation of the HSL-inhibitory effect:

Substances are normally tested in four independent mixtures. The inhibition of the HSL enzymatic activity by a test substance is determined by comparing with an uninhibited control reaction. The IC_{50} is calculated from an inhibition plot with at least 10 concentrations of the test substance. The GRAPHIT, Elsevier-BIOSOFT software package is used to analyze the data.

2. EL inhibition assay:

2.1. Preparation of EL

EL is released as secretory protein in high concentration into cell culture medium (conditioned medium) by recombinant cell lines (CHO, HEK293). This is employed

as enzyme solution after concentration.

2.2. EL activity assay

- 5 The phospholipase-specific substrate 1,2-bis(4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-3-undecanoyl)-*sn*-glycero-3-phosphocholine, (manufacturer Molecular Probes) is used to characterize the enzymatic activity of endothelial lipase and the effect of inhibitors. Hydrolysis of the A1 ester linkage of this phospholipid by the enzyme liberates the fluorescent dye Bodipy which can be detected after
10 separation by thin-layer chromatography on an HPTLC plate (silica gel 60, Merck) or directly in the reaction vessel by measuring the fluorescence.
The substrate solution is prepared by taking up 100 µg of 1,2-bis(4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-3-undecanoyl)-*sn*-glycero-3-phospho-
choline (manufacturer Molecular Probes), 2.4 mg of tripalmitin (Sigma) and 7.9 mg of
15 DOP – choline (1,2-dioleoyl-*sn*-glycero-3-phosphocholine) in 393 µl of chloroform and then transferring 157 µl into a fresh reaction vessel. After evaporation of the solvent, the lipid mixture is dissolved in 4 ml of 200 mM TRIS-HCl, 150 mM sodium chloride, pH = 7.4, by sonication twice. The subsequent enzymic reaction takes place at 37°C for 60 minutes. For this purpose, 45 µl of the substrate solution are
20 incubated with 1 µl of inhibitor of appropriate concentration (dissolved in DMSO, pure DMSO solution is used as control) and 5 µl of enzyme solution (conditioned medium). Then 3 µl of the assay mixture are loaded onto an HPTLC plate (silica gel 60, Merck), and the liberated fluorescent dye is separated for detection with an eluent (diethyl ether:petroleum benzine:acetic acid [78:22:1]). After evaporation of
25 the eluent, the plate is read in a fluorescence scanner. An increased liberation of the fluorescent dye in the uninhibited reaction is to be observed as a measure of the enzymic activity.

2.3. Evaluation of the EL-inhibitory effect:

The enzymatic activity is reduced as a function of the inhibitor concentration used, and the inhibitor concentration at which a half-maximum enzymatic activity is observed is called IC₅₀.

2.4. Further EL inhibition assay:

The phospholipase-specific substrate 1,2-bis(4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-3-undecanoyl)-*sn*-glycero-3-phosphocholine, (manufacturer Molecular Probes) is used to characterize the enzymatic activity of endothelial lipase and the effect of inhibitors. Hydrolysis of the A1 ester linkage of this phospholipid by the enzyme liberates the fluorescent dye Bodipy which can be detected after separation by thin-layer chromatography on an HPTLC plate (silica gel 60, Merck) or directly in the reaction vessel by measuring the fluorescence.

The substrate solution is prepared by dissolving 100 µg of 1,2-bis(4,4-difluoro-5,7-dimethyl-4-bora-3a,4a-diaza-s-indacene-3-undecanoyl)-*sn*-glycero-3-phosphocholine (manufacturer Molecular Probes) in 100 µl of DMSO and taking up in 2.4 mg of tripalmitin (Sigma) in 393 µl of chloroform which comprises 20 mg/ml DOP–choline (1,2-dioleoyl-*sn*-glycero-3-phosphocholine). 39.3 µl of this lipid mixture are transferred into a fresh reaction vessel, and the solvent is evaporated off. The lipid mixture is dissolved in 4 ml of 200 mM TRIS-HCl, 150 mM sodium chloride, pH = 7.4, by sonication twice. The subsequent enzymic reaction takes place at 37°C for 90 minutes. For this purpose, 20 µl of the substrate solution are incubated with 2 µl of inhibitor of appropriate concentration (dissolved in 10% DMSO, using 10% strength DMSO solution for control) and 2 µl of enzyme solution (conditioned medium). The 4 µl of the assay mixture are loaded onto an HPTLC plate (silica gel 60, Merck), and the liberated fluorescent dye is separated for detection with an eluent (diethyl ether:petroleum benzine:acetic acid [78:22:1]). After evaporation of the eluent, the plate is read in a fluorescence scanner. An increased liberation of the fluorescent dye in the uninhibited reaction is to be observed as a measure of the enzymic activity.

In this assay, the compounds of the examples showed the following IC₅₀ values:

Example	IC ₅₀ [μM] EL
1	0.16
2	0.051
3	0.052
4	0.215
5	0.435
6	0.014
7	0.12


Preparation processes



5

The compounds of the invention of the general formula I are prepared by methods known per se, e.g. by acylation of substituted or unsubstituted benzothiazol-2-one derivatives II with carbamoyl chlorides III (method A), or in two stages by reacting benzothiazol-2-one derivatives II with phosgene or equivalents such as

10 trichloromethyl chlorocarbonate, ditrichloromethyl carbonate or 4-nitrophenyl chloroformate and further reaction of the resulting benzothiazol-2-onecarboxylic acid derivative with amines IV (method B). For compounds in which R₂ is hydrogen, the benzothiazol-2-one derivatives II can also be reacted with the appropriate

15 isocyanates V R₁-N=C=O (method C).





Since acids are usually liberated in these reactions, it is advisable to add bases such as pyridine, triethylamine, sodium hydroxide solution or alkali metal carbonates for expedition. The reactions can be carried out in wide temperature ranges. It has usually proved to be advantageous to operate at from 0°C to the boiling point of the solvent used. Examples of solvents employed are methylene chloride, THF, DMF, toluene, ethyl acetate, n-heptane, dioxane, diethyl ether or pyridine. If anhydrous conditions are used, strong bases such as lithium hydride, sodium hydride or potassium tert-butoxide in aprotic solvents such as THF or DMF have also proved suitable.

The benzothiazol-2-one derivatives employed as starting compounds II are commercially available or can be prepared by processes known from the literature (e.g. C. Flouzat, Y. Bresson, A. Mattio, J. Bonnet, G. Guillaumet *J. Med. Chem.* 1993, 36, 497-503; F. Mutterer, C.D. Weis, *J. Het. Chem.* 1976, 13, 1103-1104; K. Takeda; H. Ogura, *Synthetic Communications* (1982), 12(3), 213-17; K. Konishi, I. Nishiguchi; T. Hirashima, N. Sonoda; S. Murai, *Synthesis* (1984), (3), 254-5).

The examples detailed below serve to illustrate the invention without, however, restricting it.

Examples

5

Example 1:

2-Oxobenzothiazole-3-hexylcarboxamide

- 75 mg (0.45 mmol) of 3H-benzothiazol-2-one were dissolved in 10 ml of pyridine.
10 Addition of 76 mg (0.59 mmol) of 1-isocyanatohexane was followed by stirring at 100°C for 6 h, addition once again of the same amount of isocyanatohexane and stirring at 100°C for 6 h. The reaction mixture was concentrated, and the residue was dissolved in water and extracted with ethyl acetate. The organic phase was concentrated and purified by preparative HPLC (PR18, acetonitrile/water 0.1% TFA).
15 Yield: 38 mg (28%), M+H+: 279.15.

Example 2:

2-Oxobenzothiazole-3-heptylcarboxamide

- 20 300 mg (1.98 mmol) of 3H-benzothiazol-2-one were reacted in analogy to Example 1 with 336 mg (2.38 mmol) of 1-isocyanatoheptane in dioxane at 80°C. Yield: 125 mg (22%), M+H+: 293.11.

Example 3:

- 25 2-Oxobenzothiazole-3-(2-methylbenzyl)carboxamide

300 mg (1.98 mmol) of 3H-benzothiazol-2-one were reacted in analogy to Example 1 with 350 mg (2.38 mmol) of 1-isocyanatomethyl-2-methylbenzene. Yield: 129 mg (22%), M+H+: 299.05.

30

Example 4:

2-Oxobenzothiazole-3-(3-methylbenzyl)carboxamide

300 mg (1.98 mmol) of 3H-benzothiazol-2-one were reacted in analogy to Example 1 with 350 mg (2.38 mmol) of 1-isocyanatomethyl-3-methylbenzene in dioxane at 80°C. Yield: 130 mg (22%), M+H+: 299.05.

5

Example 5:

6-Bromo-2-oxobenzothiazole-3-hexylcarboxamide

200 mg (0.869 mmol) of 6-bromo-3H-benzothiazol-2-one were reacted in analogy to Example 1 with 132.7 mg (1.04 mmol) of 1-isocyanatohexane in dioxane at 80°C. Yield: 185 mg (59%), M+H+: 357.1.

10

Example 6:

5-Chloro-2-oxobenzothiazole-3-heptylcarboxamide

15

300 mg (1.37 mmol) of 5-chloro-3H-benzothiazol-2-one were reacted in analogy to Example 1 with 174 mg (1.37 mmol) of 1-isocyanatoheptane in dioxane at 80°C. Yield: 100 mg (19%), M+H+: 327.38.

20

Example 7:


2-Oxo-5-trifluoromethylbenzothiazole-3-hexylcarboxamide

300 mg (1.37 mmol) of 5-trifluoromethyl-3H-benzothiazol-2-one were reacted in analogy to Example 1 with 202 mg (1.37 mmol) of 1-isocyanatohexane in dioxane at 80°C. Yield: 125 mg (26%), M+H+: 347.05.

25

Claims

1. A compound of the formula I



(I)

wherein the meanings are:

R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl or aminosulfonyl;

R2 hydrogen, (C₁-C₆)-alkyl; or

R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -C(=O)-, -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;

R3 hydrogen;

R4, R5 identically or differently hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, (C₁-C₆)-alkylcarbonyl, CO-OR₇, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

with the proviso that not more than one of the substituents R₃, R₄, R₅ is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

R7 hydrogen, (C₁-C₆)-alkyl, benzyl;

R8, R8a identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR₇, cyclopropyl, cyclopropylene;

R9, R10 identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;

the tautomeric forms of the compounds and the physiologically tolerated salts thereof.

2. A compound of the general formula I as claimed in claim 1, wherein

5 R1 is (C₆-C₁₂)-alkyl, (C₁-C₃)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₄-C₁₂)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, amino, (C₁-C₆)-alkylamino, trifluoromethyl;

10

R2 is hydrogen, (C₁-C₆)-alkyl; or

R1 and R2 together with the nitrogen atom bearing them are a monocyclic, saturated 5- to 6-membered ring system or a bicyclic saturated or partially unsaturated 9- to 10 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -O-, -S-, with the proviso that two units from the series -O-, -S- may not be adjacent;

20

R3 hydrogen;

R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, amino, COOR₇, (C₁-C₆)-alkylsulfonyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₁-C₆)-alkylcarbonyl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

30

- 5
10
15
20
25
30
- R6 is hydrogen, hydroxy, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylamino carbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- with the proviso that not more than one of the substituents R3, R4, R5 is halogen, trifluoromethyl or (C₁-C₆)-alkyl;
- R7 is hydrogen, (C₁-C₆)-alkyl, benzyl;
- R8, R8a are identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, COOR7, cyclopropyl, cyclopropylene;
- R9, R10 are identically or differently hydrogen, (C₁-C₆)-alkyl, (C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl.
3. A compound of the general formula I as claimed in claim 1 or 2, wherein
- R1 is (C₆-C₁₂)-alkyl, (C₁-C₃)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₃)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₃)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₄-C₁₂)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, amino, (C₁-C₆)-alkylamino, trifluoromethyl;
- R2 is hydrogen, (C₁-C₆)-alkyl; or
- R1 and R2 together with the nitrogen atom bearing them are a monocyclic, saturated 5- to 6-membered ring system or a bicyclic saturated or partially unsaturated 9- to 10 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR8-, -CR8R8a-,

-(C=R8)-, -NR9-, -O-, -S-, with the proviso that two units from the series -O-, -S- may not be adjacent;

R3 hydrogen;

5

R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl, (C₁-C₆)-alkyl, hydroxy, amino, COOR7, (C₁-C₆)-alkylsulfonyl, aminosulfonyl, pentafluorosulfonyl, (C₁-C₆)-alkylcarbonyl, CO-NR9R10, O-CO-NR9R10, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

10

R6 is hydrogen, hydroxy, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylamino carbonyl, di-(C₂-C₈)-alkylaminocarbonyl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

15

with the proviso that not more than one of the substituents R3, R4, R5 is halogen, trifluoromethyl or (C₁-C₆)-alkyl;

20

R7 is hydrogen, (C₁-C₆)-alkyl;

R8, R8a are identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, cyclopropyl;

25

R9, R10 are identically or differently hydrogen, (C₁-C₆)-alkyl.

4. A compound of the formula I as claimed in claims 1, 2 or 3, wherein

30

R1 is (C₆-C₁₂)-alkyl, benzyl, (C₁-C₂)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₂)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where benzyl, heteroaryl,

cycloalkyl or bicycle may be substituted by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy or trifluoromethyl;

R₂ is hydrogen; or

5

R₁ and R₂ together with the nitrogen atom bearing them are a monocyclic, saturated 5- to 6-membered ring system, of which individual members of the ring systems may be replaced by one to two atoms or atomic groups from the series -CHR₈-, -NR₉-, where R₈ is as defined above, and R₉ is (C₁-C₆)-alkyl or cyclopropyl.

10

5. A compound of the formula I as claimed in claims 1 to 4, in which

R₁ is (C₆-C₁₂)-alkyl, benzyl, CH₂-(C₅-C₁₂)-heteroaryl or (C₈-C₁₄)-bicycyl, where benzyl, heteroaryl or bicycyl may be substituted by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy or trifluoromethyl;

15

R₂ is hydrogen;

20 R₃ hydrogen;

R₄, R₅ are identically or differently hydrogen, halogen, hydroxy, (C₁-C₆)-alkyl, trifluoromethyl or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

25

R₆ is hydrogen.

6. A compound of the formula I as claimed in claims 1 to 5, wherein

30 R₁ is (C₆-C₁₂)-alkyl, benzyl, where benzyl may be substituted by methyl;

R₂ is hydrogen;

R3 is hydrogen;

R4, R5 are identically or differently hydrogen, halogen, trifluoromethyl;

R6 is hydrogen.

7. A compound of the formula I as claimed in claims 1 to 6, wherein

R3, R6 are hydrogen and R4 or R5 are not hydrogen.

8. A medicament comprising one or more compounds of the formula I, in which the meanings are:

R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl, aminosulfonyl;

R2 hydrogen, (C₁-C₆)-alkyl; or

R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -C(=O)-, -O-, -S-, -SO-, -SO₂-,

with the proviso that two units from the series $-O-$, $-S-$, $-SO-$, $-SO_2-$ may not be adjacent;

- 5 R3, R4, R5 identically or differently hydrogen, halogen, (C_1-C_6) -alkyl, (C_1-C_3) -alkyloxy- (C_1-C_3) -alkylene, hydroxy, (C_1-C_6) -alkylmercapto, amino, (C_1-C_6) -alkylamino, di- (C_2-C_{12}) -alkylamino, (C_1-C_6) -alkylcarbonyl, CO-OR7, trifluoromethyl, (C_1-C_6) -alkylsulfonyl, (C_1-C_6) -alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C_6-C_{10}) -aryl, (C_5-C_{12}) -heteroaryl, CO-NR9R10, O-CO-NR9R10, O-CO- (C_1-C_6) -alkylene-CO-O- (C_1-C_6) -alkyl, O-CO- (C_1-C_6) -alkylene-CO-OH, O-CO- (C_1-C_6) -alkylene-CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C_1-C_6) -alkyloxy;
- 10
- 15 R6 hydrogen, (C_1-C_3) -alkyloxy- (C_1-C_3) -alkylene, hydroxy, (C_1-C_6) -alkylmercapto, amino, (C_1-C_6) -alkylamino, di- (C_2-C_{12}) -alkylamino, pentafluorosulfanyl, (C_6-C_{10}) -aryl, (C_5-C_{12}) -heteroaryl, CO-NR9R10, O-CO-NR9R10, O-CO- (C_1-C_6) -alkylene-CO-O- (C_1-C_6) -alkyl, O-CO- (C_1-C_6) -alkylene-CO-OH, O-CO- (C_1-C_6) -alkylene-CO-NR9R10 or unsubstituted or mono- or poly-F-substituted (C_1-C_6) -alkyloxy;
- 20
- with the proviso that not more than one of the substituents R3, R4, R5 is halogen, trifluoromethyl or (C_1-C_6) -alkyl;
- 25 R7 hydrogen, (C_1-C_6) -alkyl, benzyl;
- R8, R8a identically or differently (C_1-C_6) -alkyl, halogen, trifluoromethyl, CO-OR7, cyclopropyl, cyclopropylene;
- 30 R9, R10 identically or differently hydrogen, (C_1-C_6) -alkyl, $-(C_6-C_{10})$ -aryl, (C_5-C_{12}) -heteroaryl, (C_3-C_{12}) -cycloalkyl, (C_1-C_4) -alkylene- (C_6-C_{10}) -aryl, (C_1-C_4) -alkylene- (C_5-C_{12}) -heteroaryl, (C_1-C_4) -alkylene- (C_3-C_{12}) -cycloalkyl, (C_8-C_{14}) -bicycle;

the tautomeric forms of the compounds and the physiologically tolerated salts thereof or one or more compounds of the formula I as claimed in claims 1 to 7.

- 9 The use of the compounds of the formula I, in which the meanings are:
 10 R1 (C₅-C₁₆)-alkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-
 5 heteroaryl, (C₁-C₄)-alkylene-(C₄-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle, where aryl, heteroaryl, cycloalkyl or bicycle may be substituted one or more times by halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, mono-(C₁-C₆)-alkylaminocarbonyl, di-(C₂-C₈)-alkylaminocarbonyl, (C₁-C₆)-alkoxycarbonyl, 10 (C₁-C₆)-alkylcarbonyl, cyano, trifluoromethyl, trifluoromethyloxy, (C₁-C₆)-alkylsulfonyl, aminosulfonyl;
- R2 hydrogen, (C₁-C₆)-alkyl; or
- 15 R1 and R2 form together with the nitrogen atom bearing them a monocyclic, saturated or partially unsaturated 4- to 7-membered ring system or a bicyclic saturated or partially unsaturated 8- to 14 membered ring system, of which individual members of the ring systems may be replaced by one to three atoms or atomic groups from the series -CHR₈-, -CR₈R_{8a}-, -(C=R₈)-, -NR₉-, -C(=O)-, 20 -O-, -S-, -SO-, -SO₂-, with the proviso that two units from the series -O-, -S-, -SO-, -SO₂- may not be adjacent;
- R3, R4, R5 identically or differently hydrogen, halogen, (C₁-C₆)-alkyl, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, (C₁-C₆)-alkylcarbonyl, CO-OR₇, trifluoromethyl, (C₁-C₆)-alkylsulfonyl, (C₁-C₆)-alkylsulfinyl, aminosulfonyl, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-

30

35

alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;

- 5 R6 hydrogen, (C₁-C₃)-alkyloxy-(C₁-C₃)-alkylene, hydroxy, (C₁-C₆)-alkylmercapto, amino, (C₁-C₆)-alkylamino, di-(C₂-C₁₂)-alkylamino, pentafluorosulfanyl, (C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, CO-NR₉R₁₀, O-CO-NR₉R₁₀, O-CO-(C₁-C₆)-alkylene-CO-O-(C₁-C₆)-alkyl, O-CO-(C₁-C₆)-alkylene-CO-OH, O-CO-(C₁-C₆)-alkylene-CO-NR₉R₁₀ or unsubstituted or mono- or poly-F-substituted (C₁-C₆)-alkyloxy;
- 10 R7 hydrogen, (C₁-C₆)-alkyl, benzyl;
- R8, R8a identically or differently (C₁-C₆)-alkyl, halogen, trifluoromethyl, CO-OR₇, cyclopropyl, cyclopropylene;
- 15 R9, R10 identically or differently hydrogen, (C₁-C₆)-alkyl, -(C₆-C₁₀)-aryl, (C₅-C₁₂)-heteroaryl, (C₃-C₁₂)-cycloalkyl, (C₁-C₄)-alkylene-(C₆-C₁₀)-aryl, (C₁-C₄)-alkylene-(C₅-C₁₂)-heteroaryl, (C₁-C₄)-alkylene-(C₃-C₁₂)-cycloalkyl, (C₈-C₁₄)-bicycle;
- 20

the tautomeric forms of the compounds and the physiologically tolerated salts thereof or the compounds of the formula I as claimed in claims 1 to 7 for producing a medicament for the treatment and/or prevention of diabetes mellitus .

- 25 10. The use of the compounds of the formula I as claimed in claims 1 to 9 for producing a medicament for the treatment and/or prevention of dyslipidemias.

11. The use of the compounds of the formula I as claimed in claims 1 to 9 for producing a medicament for the treatment and/or prevention of atherosclerotic disorders.

5

12. A process for producing a medicament comprising one or more of the compounds of the formula I as claimed in claims 1 to 9, which comprises mixing the latter with a pharmaceutically suitable carrier and converting this mixture into a form suitable for administration.

10

13. A process for preparing compounds of the general formula I as claimed in claims 1 to 7, which comprises benzothiazol-2-one derivatives of the formula II

15



20

25

30

35

- a) being acylated with carbamoyl chlorides of the formula III;
 or
 b) in two stages being reacted first with phosgene or equivalents such as trichloromethyl chlorocarbonate, ditrichloromethyl carbonate or 4-nitrophenyl chloroformate and in a second step with amines of the formula IV, in which the substituents have the meanings indicated above.



14. A process for preparing compounds of the general formula I with R2 hydrogen as claimed in claims 1 to 7, which comprises reacting benzothiazol-2-one derivatives of the formula II with isocyanates of the formula V: $O=C=N-R1$.

