

PATENT SPECIFICATION

(11) 1 463 885

1 463 885

- (21) Application No. 19093/74 (22) Filed 1 May 1974
 (31) Convention Application No. 160 (32) Filed 3 May 1973 in
 (33) India (IN)
 (44) Complete Specification published 9 Feb. 1977
 (51) INT CL³ C07D 521/00 A61K 31/395 (C07D 521/00 209/04 239/10)
 (52) Index at acceptance


C2C 1343 1354 1410 1412 1602 1745 213 214 215 247 250
 251 252 25Y 28X 29X 29Y 305 30Y 314 31Y 337
 360 362 364 365 366 367 36Y 371 373 37Y 551 552
 614 623 624 628 672 678 775 790 79Y QL QM QS



(54) CONDENSED PYRROLE MERCAPTO COMPOUNDS

(71) We, CIBA-GEIGY AG, a Body Corporate organised according to the laws of Switzerland, of Basle, Switzerland, do hereby declare the invention for which we pray that a patent may be granted to us, and the method by which it is to be performed, to be particularly described in and by the following statement:—

The subject of this invention is the preparation of a compound of the formula I



where Ar is a grouping which along with the adjacent carbon atoms form an o-arylene or an o-heteroarylene residue, each of R₁ and R₂ independently is hydrogen or a radical of aliphatic character, R₂ is a hydrogen atom, a radical of aliphatic character or a free carboxyl group or functional derivative thereof and Alk represents a substituted or unsubstituted alkylene or alkenylene radical contributing 2—5 carbon atoms to the ring, their tautomers and salts.

The residue Ar forms together with the two adjacent carbon atoms of the pyrrole ring a carbocyclic or monoheterocyclic radical of aromatic character. A carbocyclic radical of aromatic character is especially an o-phenylene radical which may be substituted by one, two or more identical or different substituents attached to any one of the positions available for substitution. A monoheterocyclic residue of aromatic character is primarily a monoaza, monooxa, or monothia-cyclic group of aromatic character which, in accordance to the o-phenylene residue, may be substituted. A monoazacyclic residue of aromatic character is e.g. an o-pyridylene, while monooxacyclic and monothiacyclic residues of aromatic character are o-furylene and o-thienylene groups, respectively.

Substituents of the carbocyclic and mono-

heterocyclic radicals of aromatic character are optionally substituted hydrocarbon radicals, e.g. optionally substituted aliphatic, cycloaliphatic, cycloaliphatic-aliphatic, aromatic or araliphatic hydrocarbon radicals, with aliphatic or cycloaliphatic radicals preferably being saturated. In cycloaliphatic hydrocarbon radicals the cyclic moiety is preferably monocyclic, but it can also be bicyclic or polycyclic.

An aliphatic hydrocarbon radical which can optionally be substituted is in particular an alkyl, alkenyl or alkynyl radical, in particular a straight-chain or branched lower alkyl, lower alkenyl or lower alkynyl radical. Examples of substituents of aliphatic hydrocarbons are free, esterified or etherified hydroxy groups, such as lower alkoxy, lower alkenyloxy groups or halogen atoms or functional derivatives of a carboxyl group, e.g. lower alkoxy-carbonyl groups.

The term "lower" as used hereinbefore or hereinafter in connection with the definition of organic compounds, groups and radicals, signifies that such compounds, groups and radicals contain up to and including 7 carbon atoms.

Examples of lower alkyl groups are preferably methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec. butyl, tert. butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, isohexyl or n-heptyl groups. Lower alkenyl is e.g. the allyl or 2-methylallyl group and lower alkynyl groups are preferably propargyl groups. Substituted lower alkyl groups are e.g. the trifluoromethyl group or an optionally esterified carboxymethyl group e.g. the alkoxy-carbonylmethyl group.

Further substituents of the carbocyclic or monoheterocyclic radicals of aromatic character are free, etherified or esterified hydroxy and etherified mercapto groups, e.g. alkoxy groups and halogen atoms, in particular the methoxy, ethoxy, n-propoxy, isopropoxy, n-butyloxy, isobutyloxy, tert. butyloxy, sec. butyloxy groups and alkylmercapto groups, e.g. methylmercapto or ethylmercapto groups.

Also possible are the lower alkoxy groups with 2 or 3 carbon atoms in the alkoxy moiety which are substituted by a free, etherified or esterified hydroxyl group, e.g. as hydroxy-lower alkoxy radical the 2-hydroxyethoxy radical, as lower alkoxy-alkoxy radical the 2-methoxyethoxy radical, and as halo-lower alkoxy radical the 2-chloroethoxy radical, and as optionally substituted aryl-lower alkoxy radical, for example a phenyl-lower alkoxy radical, in particular the benzyloxy radical, and as optionally substituted aryloxy radical the phenyloxy radical.

Halogen atoms are primarily fluorine, chlorine or bromine atoms, but can also be iodine atoms.

The aromatic radicals mentioned in the last cited groups can in their turn be substituted by the substituents and functional groups defined hereinbefore as in the enumeration of the individual hydrocarbon groups. The aromatic radicals of the substituents can also be substituted by an acyloxy group, for example by a lower alkanoyloxy group, in particular by the acetyloxy or propionyloxy group.

As further substituents of the carbocyclic and monoheterocyclic radicals of aromatic character are to be mentioned the nitro group, the free or substituted amino group, e.g. a lower alkylamino group, in particular the methylamino or ethylamino group or an acylamino group, e.g. a lower alkanylamino group, in particular the acetylamino or the propionylamino group. As disubstituted amino group there may be mentioned the N,N - di-lower alkylamino group, e.g. the N,N - dimethylamino or the N,N - diethylamino groups.

Examples of further secondary and tertiary amino groups are: N - ethyl - N - methylamino, n-propylamino, di - n - propylamino, isopropylamino, diisopropylamino or di - n - butylamino groups.

Further substituents of the carboaromatic or heteroaromatic radicals are acyl groups, free or functionally modified carboxy groups, in particular esterified carboxy groups, e.g. the methoxycarbonyl or ethoxycarbonyl groups or lower alkanoyl groups, for example the acetyl or the propionyl groups.

Radicals of aliphatic character of the substituents R₁, R₂ and R₃ are optionally substituted hydrocarbon radicals, e.g. aliphatic, cycloaliphatic, cycloaliphatic-aliphatic, aromatic or araliphatic hydrocarbon radicals.

An aliphatic hydrocarbon radical preferably has a maximum of 7 carbon atoms and is e.g. lower alkyl, lower alkenyl or lower alkynyl. A lower alkyl radical is especially one mentioned above. A lower alkenyl group preferably is allyl or methallyl, and a lower alkynyl group propargyl. A cycloaliphatic hydrocarbon residue preferably has 3-8 ring carbon atoms, and represents, for example, a

cycloalkyl, e.g. cyclopentyl or cyclohexyl group. A cycloaliphatic-aliphatic hydrocarbon radical preferably contains as a cycloaliphatic residue one of those mentioned above and represents preferably a cycloalkyl-lower alkyl radical, e.g. cyclopentyl-methyl, cyclohexyl-methyl, or cyclohexylethyl. The above mentioned hydrocarbon radicals may be mono-, di- or poly-substituted aliphatic hydrocarbon radicals.

Aliphatic hydrocarbon radicals can be substituted by a free hydroxy, etherified or esterified hydroxy group, free or substituted amino group or nitro group, a lower alkylenedioxy group or by substituents which have already been defined hereinbefore.

Cycloaliphatic hydrocarbon radicals can be substituted by lower alkyl radicals, in particular by the lower alkyl radicals which have already been defined hereinbefore in detail or by the substituents which have already been mentioned in the enumeration of the aliphatic hydrocarbon radicals.

The aliphatic radical of a cycloaliphatic-aliphatic or araliphatic group can be substituted principally by one of the lower alkyl radicals already defined hereinbefore. The aromatic portion of the araliphatic group may be substituted for example by substituents mentioned above for an o-phenylene radical.

Aliphatic hydrocarbon radicals of the substituents R₁, R₂ and R₃ are in particular alkyl radicals, particularly however, lower alkyl radicals.

The bivalent alkylene and alkenylene radical "Alk" can be monosubstituted or polysubstituted, in particular by free or substituted hydroxy or mercapto groups such as etherified hydroxy or mercapto groups, e.g. lower alkoxy groups, in particular methoxy and ethoxy groups, or e.g. by lower alkylmercapto groups, in particular methylmercapto or ethylmercapto groups. Also possible are esterified hydroxy groups, e.g. lower alkanoyloxy groups, in particular the acetyloxy group, halogen atoms, especially fluorine, chlorine or bromine atoms, the oxo group or functionally modified oxo groups, e.g. the hydroxyimino, hydrazone or semicarbazone radical. A lower alkylene or alkenylene radical "Alk" can be, for example, the 1,2-ethylene, 1,2- or 1,3-propylene, 2,3- or 2,4- or 1,5-pentylene, vinylene or 1,3-propenylene radical. Besides representing an alkyl group, the substituent R₄ represents above all a carboxy group, but it can also represent a functional derivative of a carboxy group, e.g. a lower alkoxy-carbonyl group, in particular the methoxycarbonyl, n-propoxycarbonyl or n-butoxycarbonyl group. Also possible are nitrogen-containing functional derivatives of the carboxy group, e.g. the carbamoyl group, which can be optionally substituted at the nitrogen atom, for example a carbamoyl group which is substituted by N-lower alkyl, in particu-

70

75

80

85

90

95

100

105

110

115

120


125

130

lar the N-methylcarbamoyl, N,N-dimethylcarbamoyl, N-ethylcarbamoyl or the N,N-diethylcarbamoyl groups. As further nitrogen-containing functional derivatives of the carboxy group mention may also be made of the nitriles.

The new compounds exhibit valuable pharmacological properties. In addition to ophthalmological and hypotensive effects, they primarily show vasoconstrictor, especially decongestant, e.g. nasal decongestant properties. The new compounds are, therefore, useful as pharmacologically effective compounds, primarily as nasal decongestant agents for which purpose daily doses of 0.1 to 50 mg may be used. As anti-hypertensive agents they are useful at a daily dose of 50—500 mg. They are also useful as intermediate products in the preparation of other valuable compounds, especially of pharmacologically active substances.

The invention concerns primarily compounds of the formula (Ia)



wherein X is nitrogen or represents the group CH, and each of Ra and Rb is hydrogen or any of the substituents of the carbocyclic or heterocyclic aromatic radicals mentioned earlier, each of R₁ and R₂ independently is hydrogen or a lower alkyl group, R₃ is hydrogen, lower alkyl or a free or esterified carboxy group as for example an alkoxy carbonyl group and n is an integer denoting 2—5, their tautomers and acid addition salts.

Of special interest in this invention are compounds of formula Ia in which X represents the group CH, each of R₁ and R₂ independently is hydrogen or lower alkyl and R₃ is hydrogen, lower alkyl, a free carboxy or methoxycarbonyl or ethoxycarbonyl group, and each of R₄ and R₅ is hydrogen, lower alkyl, halogen or a lower alkoxy group and n is an integer denoting 2—4, their tautomers and acid addition salts.

Especially the invention concerns compounds of formula Ia in which X represents the group CH, R₁ and R₂ independently is hydrogen or lower alkyl and R₃ is hydrogen, lower alkyl or a free carboxy group, and each of R₄ and R₅ is hydrogen, halogen or a lower alkoxy group and n is an integer denoting 2—4, their tautomers and acid addition salts.


Furthermore the invention concerns in particular compounds of formula Ia in which X represents the group CH, R₁ and R₂ are hydrogen or methyl and R₃ is hydrogen, methyl or a free carboxy group, and each of R₄ and R₅ is hydrogen, chlorine, bromine

or methoxy, and n is an integer denoting 2—4, their tautomers and acid addition salts.


Of the new compounds of formula Ia is especially named the 3 - (2 - imidazolin - 2 - ylthio) - indole hydrochloride and hydroiodide.

The new compounds of the general formula I are obtained by methods which are in themselves known, as for example

a) by reacting a compound of the formula II wherein R₁ and "Alk" have the meanings defined in formula I




or a tautomer thereof with a compound of the formula III



wherein R₁ and R₂ have the meanings given under formula I, in the presence of a suitable oxidising agent and if desired converting the resulting compound to an acid addition salt thereof.

This process is performed with compounds of the formula II and III in the presence of an oxidising agent, especially halogen, particularly, iodine or bromine in the presence of suitable solvents and other additives such as an alkali metal salt of a hydrohalic acid especially a potassium salt of hydroiodic or hydrochloric acid. Other oxidising agents may also be used such as ferric chloride or ferricyanide salts, advantageously, in the presence of a suitable solvent.

b) In a second process a compound of the formula IV




wherein R₁, R₂ and Ar have the meanings given under I, R₄ is a free or functionally converted derivative of a hydroxy group, a free mercapto group or a modified mercapto group or amino group, R₅ is an oxygen or sulphur atom, or an imino group or R₄ and R₅ taken together represents a triple bonded nitrogen atom (nitrilo residue) is reacted with a compound of the general formula V,



(14)

in which R_6 and R_7 , each depending on the meaning of R_4 and R_5 , represents an amino, imino, hydroxy or functionally converted hydroxy, whereby the substituents R_4 — R_7 contain at least 2 nitrogen atoms in order to obtain the ring of formula VI



and if desired, converting a resulting compound to an acid addition salt thereof.


The above defined reaction results with compounds of formula IV, which, depending on the meaning of the substituents R_4 and R_5 , are primarily derivatives of the thiocarbonic, dithiocarbonic or trithiocarbonic acid, such as amide, imides, nitriles, halides or intramolecular anhydrides.

The substituents R_6 and R_7 of the general formula V are amino, imino, hydroxy or a reactive esterified hydroxy group.

For example in compounds of the general formula IV the substituent R_4 is an NHR_5 group and R_5 an unsubstituted amino group and in compounds of formula V each of the substituent R_6 and R_7 is a hydroxy, especially, a reactive esterified hydroxy group, particularly a halogen atom.

As a further example, in a compound of formula IV the substituent R_4 is an NHR_5 group and R_5 an oxo or a thioxo group and R_6 in the compound of formula V a hydroxy or reactive esterified hydroxy group as mentioned before and R_7 an amino group. Furthermore in a compound of formula IV R_4 is a hydroxy, mercapto or a reactive esterified hydroxy group and R_5 an amino group and in the compound of formula V the substituent R_6 is an NHR_5 group and R_7 a hydroxy or a reactive esterified hydroxy group as described before.

In a further example, the substituent R_4 of compound of formula IV is a hydroxy, reactive esterified hydroxy or mercapto group and R_5 an oxo, thioxo group or R_4 and R_5 taken together in the residue




as a nitrilo residue form a nitrile and in the compound V R_4 is an NHR_5 group and R_5 an amino group.

Preferably in the above process as a compound of formula IV a pseudothiurea is used in which the substituent R_4 is an NHR_5 group and R_5 an amino group and as compound of formula V a reactive diester in


which R_6 and R_7 , each represent a hydroxy group is used.

The reactor of a pseudothiurea compound of formula IV with a reactive diester of an alkanediol of formula V is carried out by using a diester with an above defined strong acid, in particular with a hydrohalic acid. The reaction is carried out in a suitable solvent, in particular a lower alcohol e.g. ethanol and, if desired, in the presence of an acid-binding medium, e.g. an alkali-metal or alkaline earth metal hydroxide, carbonate or hydrogen carbonate.

c) According to a third process compounds of formula I may be prepared by reacting a compound of the general formula VII



in which R_1 , R_2 and Ar have the meanings defined under formula I and Y is an anion of an inorganic acid with a compound of the general formula II



wherein R_3 has the same meaning as defined under formula I. The reaction is carried out by reacting a diazonium salt of formula VII, in which Y^- is an anion of an inorganic acid, for example hydrochloric acid, sulphuric acid, fluoboric acid or phosphoric acid with a compound of formula II, wherein R_3 has the above defined meaning.

When a compound of formula II is added to a diazonium salt solution of formula VII in an acidic or neutral solution, an isothio-uronium salt is formed which is substituted at the sulphur atom.

Resulting compounds may be converted into others of the formula I, for example, by modifying, introducing or splitting off substituents according to methods known per se. Substituents attached to aromatic rings may be converted to other substituents, for example, nitro groups into amino groups by reduction in the usual manner, for example by treatment with catalytically activated hydrogen.

The reactions of this invention are carried out in the usual manner, at room temperature or with cooling or heating, under atmospheric or superatmospheric pressure, if necessary, in the presence or absence of diluents, catalysts and condensing agents, and/or in the atmosphere of an inert gas, e.g. nitrogen.

The starting materials to be used in the present process are advantageously those which give rise to the compounds referred to above as being especially valuable.

The starting materials are known or, in so far as they are new, they can be prepared by known methods. Thus, compounds of the formula II are prepared by reacting an alkylenediamine of the formula



with carbon disulphide or with thiophosgene. Compounds of the formula III are well-known in the literature and may be prepared apart from other methods by the Fischer indole cyclization of suitable hydrazone compounds.

Starting materials of the formula IV are prepared for example by reacting an optionally substituted thiourea with compounds of the formula III in the presence of a suitable oxidising agent as described under the first process for the preparation of pseudo thiourea compounds. Other starting materials of the formula IV may be obtained by reacting a suitable thiocyanate, thiocarbonic acid or one of its derivatives with compounds of the formula III.

Compounds of the formula VII of the third process can be prepared by the conventional diazotization of 3-amino derivatives of the formula III in the presence of a nitrite such as an alkali metal nitrite, for example sodium nitrite or an organic nitrite for example amyl nitrite in the presence of a suitable acid such as inorganic acid, for example hydrochloric, sulphuric, phosphoric or fluoboric acids. The diazonium salts are usually prepared and reacted in solution.

Depending on the reaction conditions employed, the new compounds are obtained in the free form or in the form of their salts. The latter are acid addition salts such, for example, as pharmaceutically acceptable, non-toxic acid addition salts, for example, salts with inorganic acids such as hydrochloric, hydrobromic, nitric, sulphuric or phosphoric acids, or with organic acids such as organic carboxylic acids, for example acetic, propionic, glycollic, malonic, succinic, maleic, hydroxymaleic, fumaric, malic, tartaric, citric, benzoic, cinnamic, mandelic, salicylic, 4-aminosalicylic, 2-phenoxybenzoic, 2-acetoxybenzoic, embonic, glucuronic, nicotinic or isonicotinic acid, or with organic sulphonic acids, for example, methanesulphonic, ethanesulphonic, 2-hydroxyethanesulphonic, 1,2-ethanedisulphonic, benzenesulphonic, paratoluenesulphonic or 2-naphthalenesulphonic acid. These and other acid addition salts may also be used as intermediate products, for example, for purifying the free compounds or in the manufacture of other salts, as well as for identification

purposes. Salts particularly suitable for the last mentioned purpose are, for example, those with acid organic nitro compounds, for example, picric, picrolonic or flavianic acid, or with metallic complex acids, for example, phosphotungstic, phosphomolybdic, chloroplatinic acid or Reinecke acid, either mono-salts or poly-salts may be formed.

Free compounds may be converted into their acid addition salts, for example, by treatment with acids, such as the acids mentioned above, for instance by treating a solution of the base in a suitable inert solvent or solvent mixture with an acid or a solution thereof, or with a suitable ion exchange resin. Salts may also be obtained in the form of their hydrates, or they may contain solvent of crystallization.

A resulting salt may be converted into the free compound, for example, by treatment with a base, such as an alkali or alkaline earth metal hydroxide or carbonate, or with ammonia or with a suitable ion exchange resin.

A resulting salt may also be converted into another salt, for example, by treatment with an ion exchange resin or by reacting a salt with an inorganic acid with a metal salt, for example, a sodium, barium or silver salt, of an acid in a suitable solvent in which the resulting inorganic compound is insoluble.

In view of the close relationship between the new compounds in the free form and in the form of their salts, whenever a free compound or a salt is referred to in this context, the corresponding salt and free compound, respectively, is also intended, provided such is possible or appropriate under the circumstances.

Mixtures of isomers may be resolved into their constituents by known methods. Thus, for example, resulting racemates may be resolved into the optically active d-forms and l-forms by crystallization from optically active solvents, or by treating the racemic compounds with an optically active form of an acid, containing an asymmetric carbon atom, preferably in the presence of a suitable solvent. Particularly suitable optically active forms of acids are d-tartaric acid and l-tartaric acid, as well as the optically active forms of maleic, mandelic, camphor-10-sulphonic or quinic acid. Resulting salts may be converted into other salts or into the free and optically active bases, and an optically active base may be converted into an acid addition salt by the methods referred to above.

The invention includes also any modification of the present process in which a starting material is formed under the reaction conditions or used in the form of a derivative, for example, a salt thereof.

The invention also includes pharmaceutical preparations which contain one of the new compounds or a salt thereof in admixture

(16)

with a pharmaceutical organic or inorganic solid or liquid carrier suitable for enteral, as for example oral, parenteral or topical administration.

- 5 Suitable dosage units are dragées tablets, capsules, suppositories or ampoules. These pharmaceutical preparations contain per dosage unit, e.g. 0.01—100 mg, preferably 10 0.1—50 mg of the new compound or a pharmaceutically acceptable acid addition salt thereof. It is moreover possible to use corresponding amounts of preparations, solutions, suspensions and emulsions which are not administered as individual doses, such as 15 in the case of drops, syrups and elixirs.

- In dosage units for oral administration, the content of active substance is preferably between 10% and 90%. Such dosage units are produced by the combination of the active substance with, e.g. solid pulverulent carriers such as lactose, saccharose, sorbitol, mannitol; starches such as potato starch, maize starch or amylopectin, also laminaria powder or citrus pulp powder; cellulose derivatives or gelatine, optionally with the addition of 25 lubricants such as magnesium or calcium stearate, or polyethylene glycols, to form tablets or dragée cores.

- The latter are coated, for example, with 30 concentrated sugar solutions which may also contain, e.g. gum arabic, talcum and/or titanium dioxide; or with a lacquer dissolved in readily volatile organic solvents or solvent mixtures. Dyestuffs can be added to these 35 coatings; e.g., for identification of the various doses of active substance. Further suitable oral dosage units are hard gelatine capsules, as well as soft closed capsules made from gelatine and a softener such as glycerin. The 40 hard capsules contain the active substance preferably as a granulate in admixture with lubricants such as talcum or magnesium, and, optionally, stabilisers such as sodium metabisulphite ($\text{Na}_2\text{S}_2\text{O}_5$) or ascorbic acid. In soft 45 capsules the active substance is preferably dissolved or suspended in suitable liquids such as liquid polyethylene glycols, to which likewise stabilisers may be added.

- Suitable dosage units for rectal administration are, e.g. suppositories consisting of a combination of an active substance with a suppository mixture based on natural or synthetic triglycerides (e.g. cocoa butter), polyethylene glycols or suitable higher fatty 55 alcohols; and gelatine rectal capsules containing a combination of the active substance with polyethylene glycols.

- Ampoule solutions for parenteral administration, particularly intramuscular or intravenous administration, contain, for example, 60 a compound of the general formula I, in a concentration of preferably 0.5—5%, as an aqueous dispersion prepared with the aid of conventional solubility-promoting agents and/or emulsifiers; or preferably an aqueous 65

solution of a pharmaceutically acceptable, water-soluble acid addition salt of a compound of the general formula I.

In the case of liquids to be taken orally, such as syrups and elixirs, the concentration of the active substance is designed to enable 70 single doses to be easily measured out, e.g. as the content of a tea-spoon or of a measuring spoon containing, e.g. 4 ml, or as a multiple of this volume. Suitable syrups are, 75 for example, solutions of water-soluble acid addition salts, or suspensions of insoluble but absorbable acid addition salts, in aqueous solutions of sugars and/or alkanepolyols, such as unrefined sugar or sorbitol or glycerin, 80 flavourings and aromatics, as well as, optionally, preservatives and stabilisers. Elixirs are aqueous-alcoholic solutions of a compound of the general formula I, or of pharmaceutically acceptable salts thereof, which 85 may likewise contain the additives mentioned in the case of syrups. Further oral preparations to be mentioned are dropping-solutions which usually have a higher alcohol content and, at the same time, a higher content of 90 active substance, so that a single dose can be measured out, e.g. as 10 to 40 drops. Furthermore compounds of formula I, especially its pharmaceutically acceptable salts can 95 be used therapeutically in the form of drops in a solution.

The following Examples a) to g) illustrate the production of some typical preparations, but these examples, however, in no way represent the only embodiments of such 100 preparations.

a) An amount of 250.0 g of active substance is mixed with 550.0 g of lactose and 292.0 g of potato starch; the mixture is moistened with an alcoholic solution of 8 g of 105 gelatine, and granulated through a sieve. The obtained granulate is dried and 60.0 g of potato starch, 60.0 g of talcum, 10.0 g of magnesium stearate and 20.0 g of colloidal silicon dioxide are mixed in; the mixture is then pressed into 10,000 tablets each weighing 125 mg and each containing 25 mg of 110 active substance. The tablets can optionally be provided with grooves for a more precise adjustment of the dosage amount. 115

b) A granulate is prepared from 100.0 g of active substance, 379.0 g of lactose, and the alcoholic solution of 6.0 g of gelatine; the granulate is then dried and mixed with 10.0 g of colloidal silicon dioxide, 40.0 g of talcum, 6.0 g of potato starch and 5.0 g of magnesium stearate; and the mixture finally 120 pressed to obtain 10,000 dragée cores. These are subsequently coated with a concentrated syrup made from 533.5 g of crystallised saccharose, 20.0 g of shellac, 75.0 g of gum arabic, 250.0 g of talcum, 20.0 g of colloidal silicon dioxide and 1.5 g of dyestuff. After 125 drying, the obtained dragées each weight

150 mg and each contain 10 mg of active substance.

c) An amount of 20.0 g of active substance is dissolved in 1500 ml of boiled pyrogen-free water, the solution being then made up to 2000 ml with similarly treated water. The solution is filtered off and filled into 1000 ampoules each containing 2 ml, these being finally sterilised. A 2 ml ampoule contains 20 mg or 1.0% of active substance.

d) An amount of 25.0 g of active substance and 1975 g of finely ground suppository foundation substance (e.g. cocoa butter) is thoroughly mixed and then melted. The melt is maintained homogeneous by stirring whilst being poured out to form 1000 suppositories each weighing 2.0 g. They each contain 25 mg of active substance.

e) For the preparation of a syrup containing 0.25% of active substance, 1.5 litres of glycerin, 42 g of *p*-hydrobenzoic acid methyl ester, 18 g of *p*-hydroxybenzoic acid-*n*-propyl ester and, with slight heating, 25.0 g of active substance are dissolved in 3 litres of distilled water; additions are then made to the solution of 4 litres of 70% sorbitol solution, 1000 g of cryst. saccharise, 250 g of glucose and an aromatic, e.g. 250 g of Orange Peel Soluble Fluid of the firm Eli Lilly and Co., Indianapolis, or 5 g each of natural lemon aroma and "Halb und Halb" essence, both manufactured by the firm Haarmann und Reimer, Halzminden, Germany; the obtained solution is filtered, and the filtrate made up with distilled water to 10 litres.

f) In order to prepare a dropping solution containing 1.5% of active substance. 150.9 g of active substance and 30 g of sodium cyclamate are dissolved in a mixture of 4 litres of ethanol (96%) and 1 litre of propylene glycol. A mixture is then prepared separately of 3.5 litres of 70% sorbitol solution with 1 litre of water; this mixture is subsequently added to the above mentioned active substance solution. An addition is thereupon made of an aromatic, e.g. 5 g of cough-sweet aroma or 30 g of grapefruit essence, both from the firm Haarmann and Reimer, Halzminder, Germany; and the whole is afterwards well mixed, filtered and made up with distilled water to 10 litres.

g) In order to prepare a solution for using therapeutically in the form of drops, the following compositions are added:


active substance (compound of formula I)	1 g
para-hydroxybenzoic acid propylester	0.8 g
para-hydroxybenzoic acid methylester	0.4 g
de-ionized water, to make up	1000 cc.

The following examples illustrate the pro-

duction of the new compounds of the general formula I; these examples, however, are not intended in any way to limit the scope of the invention. The temperatures are expressed in degrees Centigrade.

Example 1

A mixture of 11.7 g indole in 100 ml methanol and 10.2 g 2-imidazolidinethione in 150 ml methanol is added to a well-stirred solution of 25.4 g iodine and 50 g potassium iodide in 100 ml water. The mixture is stirred for 2 hours at room temperature. The clear solution is concentrated in vacuo to one third of the original volume and cooled. The solid which separates is filtered off, dried and recrystallised from alcohol-ether to give 3 - (2 - imidazolin - 2 - ylthio) - indole hydroiodide of the formula



melting at 210—211°.

By passing a solution of 10 g of above hydroiodide in 50 ml methanol through a column of 100 g Amberlite (Registered Trade Mark) IRA-400 (chloride form) resin in methanol, the hydrochloride is obtained. After recrystallization from alcohol-ether, it melts at 240—241°.

Example 2

A mixture of 11.7 g indole in 50 ml methanol and 11.6 g 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 100 ml methanol is added to a well-stirred solution of 5.4 g iodine and 50 g potassium iodide in 100 ml water. The mixture is stirred for 12 hours at room temperature and cooled. The crystalline product is filtered off. The filtrate is concentrated to one third of the original volume, and cooled to yield more of the resulting product. The combined product is recrystallised from alcohol to give 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydroiodide, melting at 251°.

By passing a solution of 7 g of above hydroiodide in 35 ml methanol through a column of 70 g Amberlite IRA-400 (chloride form) resin in methanol, the hydrochloride is obtained. Upon recrystallization from alcohol-ether, it is obtained as a monohydrate melting at 125—126°.

Example 3

A mixture of 3.2 g indole-2-carboxylic acid in 50 ml methanol and 2 g 2-amidazolidinethione in 100 ml methanol is added to a well-stirred solution of 3 g iodine and 25 g potassium iodide in 50 ml water. After being stirred for 2 hours at room temperature, the mixture is filtered to remove the

product. The filtrate is concentrated and cooled to give further yield. The combined product is recrystallised from alcohol-ether to give 3 - (2 - imidazolin - 2 - ylthio) - indole - 2 - carboxylic acid hydriodide melting above 310°.

By passing a solution of 2 g of the above hydriodide in 10 ml methanol through a column of 20 g Amberlite IRA-400 (chloride form) resin in methanol, the hydrochloride is obtained. Upon recrystallization from methanol-ether, it is obtained as a monohydrate melting at 187—188° (with decomposition).

The free base is obtained by treatment of an aqueous suspension of the hydriodide with sodium bicarbonate. Upon crystallization from aqueous alcohol it melts at 204—206°.

Example 4

A mixture of 6.4 g indole - 2 - carboxylic acid in 25 ml methanol and 4.6 g 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 50 ml methanol is added to a well-stirred solution of 10.1 g iodine and 16.3 potassium iodide in 50 ml water. The mixture is stirred at room temperature for 1½ hours, and cooled. The crystalline product is filtered off. A second yield is obtained by concentrating the filtrate to a small volume and cooling. The combined product is recrystallised from hot water to give 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid hydriodide melting at 252°.

Example 5

A mixture of 2.8 g 7 - azaindole in 25 ml methanol and 3.5 g 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 50 ml methanol is added to a well-stirred solution of 7.6 g iodine and 15 g potassium iodide in 50 ml water. The mixture is stirred at room temperature for 14 hours and filtered. The precipitate is washed with water and ether and then recrystallised from water to give 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - 7 - aza - indole hydriodide melting at 294—295°.

Example 6

A solution of 19.6 g of 5 - bromo - indole in 50 ml of methanol is added to a stirred solution of 50 g potassium iodide and 25.4 g of iodine in 150 ml of water at room temperature. The mixture is further treated with a solution of 11.6 g of 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 270 ml of warm methanol. The reaction mixture is then stirred at room temperature for 2 hours, filtered and the filtrate is concentrated to one-fifth of the original volume and cooled. A crystalline precipitate is formed. This is filtered and recrystallised from a mixture of methanol, ethylacetate and ether to yield 5 -

bromo - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 241—242°.

Example 7

A solution of 11.7 g of indole in 100 ml of methanol is added portionwise to a stirred solution of 50 g potassium iodide and 25.4 g of iodine in 150 ml of water at room temperature. This mixture is further treated with a solution of 13 g of 1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 150 ml of methanol. The reaction mixture is then stirred at room temperature for 2 hours, filtered and the filtrate concentrated to one-fourth of the original volume and cooled. A crystalline precipitate is formed. This is filtered and recrystallised from a mixture of methanol, ethyl acetate and ether to yield 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 207°.

The starting material is prepared as follows:

To a solution of 162 g 3 - methylamino-propylamine in 600 ml of 50% aqueous ethanol, 121 ml carbon disulphide is added dropwise so that the temperature does not exceed 60°. The reaction mixture is heated at 60° for 2 hours and 15 ml concentrated hydrochloric acid is added and then the reaction mixture is heated under reflux for 16 hours at 100°. On cooling to -5°, a crystalline precipitate is formed which is filtered and recrystallised from isopropanol to afford 1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol which melts at 125°.

Example 8

A solution of 9.36 g of indole in 60 ml methanol is added dropwise to a well-stirred solution of 40 g potassium iodide and 20.3 g of iodine in 120 ml of water at room temperature. This is reacted with a solution of 10.6 g of 2 - mercapto - 1,4,5,6 - tetrahydro - 5 - pyrimidinol in 1800 ml of 50% aqueous methanol. The reaction mixture is stirred at room temperature for 2 hours and worked up in the manner, described under Example 1 to yield 3 - (5 - hydroxy - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 228—229° (with decomposition) after recrystallization from a mixture of methanol, ethyl acetate and ether.

The starting material is prepared as follows:

A mixture of 42 g 1,3 - diamino - 2 - propanol and 150 ml of 50% ethanol is treated dropwise with 31 ml carbon disulphide and the reaction mixture heated at 60° for 2 hours, 4 ml of concentrated hydrochloric acid is added and the reaction mixture is heated for 10 hours at 100°. It is cooled to 5° and the crystalline precipitate

formed, is filtered off and recrystallized from 50% aqueous ethanol to yield 2 - mercapto - 1,4,5,6 - tetrahydro - 5 - pyrimidinol which melts at 236°.

Example 9

5 A solution of 28.98 g of indole - 2 - carboxylic acid in 300 ml of methanol is added to a well-stirred solution of 90 g of potassium iodide and 45.6 g of iodine in 270 ml of water at room temperature. This mixture is further treated with a solution of 23.4 g of 1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 420 ml of warm methanol. The reaction mixture is stirred for 2 hours at room temperature and worked up in the manner described under Example 1 to yield 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid hydriodide which melts at 230—231° after recrystallization from a mixture of methanol, ethyl acetate and ether.

Example 10

25 A solution of 5.7 g of 5 - methoxy - indole - 2 - carboxylic acid in 70 ml of hot methanol is added portionwise to a solution of 15 g potassium iodide and 7.6 g of iodine in 45 ml of water. This is further treated with a solution of 3.48 g of 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 70 ml of hot methanol, stirred at room temperature for 2 hours and filtered. The filtrate is worked up in the manner described under Example 1 to yield 5 - methoxy - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid hydriodide which melts at 239—240° after recrystallization from a mixture of methanol, ethyl acetate and ether.

Example 11

40 A solution of 19.65 g of 2 - methylindole in 250 ml of methanol is added to a well-stirred solution of 75 g of potassium iodide and 38 g of iodine in 225 ml of water at room temperature. This mixture is further treated with a solution of 17.4 g of 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 350 ml of warm methanol, stirred for 2 hours and worked up in the manner described under Example 1 to yield 2 - methyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 242° after recrystallization from a mixture of methanol and ethyl acetate.

Example 12

55 A solution of 8.6 g of 1 - methylindole - 2 - carboxylic acid in 250 ml of warm methanol is added to a well-stirred solution of 25 g of potassium iodide and 12.2 g of iodine in 75 ml of water and treated with a solution of 5.2 g of 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol. The reaction is carried out and then worked up in the manner described

under Example 1 to yield 1 - methyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid hydriodide which melts at 227° (with decomposition) after recrystallization from a mixture of methanol and ethyl acetate.

Example 13

70 A solution of 13.1 g of 2 - methylindole in 175 ml of warm methanol is added to a well-stirred solution of 50 g of potassium iodide and 24.5 g of iodine in 150 ml of water. This mixture is further treated with a hot solution of 10.2 g of 2 - imidazolidine-thione in 150 ml of warm methanol. The reaction is carried out and worked up in the manner described under Example 1 to yield 2 - methyl - 3 - (2 - imidazolin - 2 - ylthio) - indole hydriodide which melts at 227—228° (with decomposition) after recrystallization from a mixture of methanol, ethyl acetate and ether.

Example 14

85 A solution of 3.45 g of 5 - methoxy - indole - 2 - carboxylic acid in 50 ml methanol is added to a solution of 4.57 g of iodine and 9 g of potassium iodide in 27 ml of water. This is treated with a solution of 2.34 g of 1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 30 ml of water. The reaction is carried out and worked up as described under Example 1 to yield 5 - methoxy - 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid hydriodide which melts at 256° (with decomposition) after recrystallization from a mixture of methanol and ethyl acetate.

Example 15

100 A mixture of 5.85 g indole in 50 ml methanol and 5.8 g 1 - methyl - 2 - imidazolidine-thione in 100 ml methanol is treated with a solution of 12.7 g of iodine and 25 g potassium iodide in 100 ml water. The mixture is stirred for 2 hours at room temperature, the clear solution is concentrated in vacuo to one third of the original volume and cooled. The solid which separates is filtered off and recrystallized from a mixture of methanol and ethyl acetate to give 3 - (1 - methyl - 2 - imidazolin - 2 - ylthio) - indole hydriodide melting at 215—216°.

Example 16

115 A solution of 12.7 g of iodine and 25 g of potassium iodide in 100 ml of water is added dropwise to a mixture of 7.25 g of 1,2 - dimethylindole in 60 ml methanol and 5.8 g of 1 - methyl - 2 - imidazolidine-thione in 100 ml methanol. The reaction mixture is stirred for 2 hours at room temperature. The crystalline precipitate formed was filtered and recrystallized from a mixture of

methanol, ethyl acetate and ether to yield 1,2 - dimethyl - 2 - (1 - methyl - 2 - imidazolin - 2 - ylthio) - indole hydriodide which melts at 250—251° (with decomposition).

Example 17


A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml water was added dropwise to a mixture of 7.25 g of 1,2 - dimethylindole and 6.5 g of 1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 120 ml of methanol. The reaction mixture is stirred at room temperature for 2 hours. The crystalline precipitate formed was filtered and recrystallized from a mixture of methanol and ethyl acetate to yield 1,2 - dimethyl - 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 239—240° (with decomposition).

Example 18

A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml of water is added dropwise to a mixture of 7.25 g of 1,2 - dimethylindole in 30 ml of methanol and 5.8 g of 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 120 ml of warm methanol. The reaction mixture is stirred for 2 hours at room temperature. The crystalline precipitate formed was filtered and recrystallized from a mixture of methanol and ethyl acetate to yield 1,2 - dimethyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 246° (with decomposition).

Example 19

A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml of water is added dropwise to a solution of 7.25 g indole in 30 ml methanol and 6.5 g of 4,5,6,7 - tetrahydro - 1H - 1,3 - diazepine - 2 - thiol in 100 ml of methanol. The reaction mixture is stirred for 2 hours at room temperature. The crystalline precipitate formed was filtered off and recrystallized from a mixture of methanol and ethyl acetate to afford 1,2 - dimethyl - 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole hydriodide of the formula



which melts at 245—246° (with decomposition).

Example 20

A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml of water is added dropwise to a well-stirred mixture of 5.85 g of indole in 20 ml methanol and

6.5 g of 4,5,6,7 - tetrahydro - 1H - 1,3 - diazepine - 2 - thiol in 100 ml methanol. The reaction mixture is stirred at room temperature for 2 hours. The crystalline precipitate formed was filtered off and recrystallized from a mixture of methanol, ethyl acetate and ether to afford 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole hydriodide which melts at 236° (with decomposition).

Example 21

To a stirred solution of 3.6 g 1,2 - dimethylindole and 2.55 g 2 - imidazolidinethione in 80 ml methanol is added a solution of 6.35 g iodine and 1 g potassium iodide in 50 ml water. After 1 hour at room temperature, the solution is concentrated in vacuo until crystals separate, then cooled and filtered. The precipitate is washed with water and ether and is recrystallized from alcohol to give 1,2 - dimethyl - 3 - (2 - imidazolin - 2 - ylthio) - indole hydriodide melting at 203—205° (with decomposition).

Example 22


To a stirred solution of 1.76 g indole and 1.5 g imidazole 2 - thiol in 50 ml methanol is added a solution of .81 g iodine and 8 g potassium iodide in 30 ml water. After stirring for 3 hours at room temperature, the solution is concentrated in vacuo. An oil separates which crystallizes upon addition of ether. The crystals are filtered off and washed with ethyl acetate; and finally recrystallized from a mixture of ethanol and ethyl acetate to give 3 - (2 - imidazol - 2 - ylthio) - indole hydriodide melting at 199—201°.

Example 23

A solution of 6.35 g of iodine and 11 g of potassium iodide in 50 ml of water is added dropwise to a mixture of 4.9 g of 5 - bromoindole in 35 ml methanol and 3.25 g of 4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - thione in 50 ml of methanol. The reaction mixture is allowed to be stirred at room temperature for 2 hours. The crystalline precipitate formed is filtered off and recrystallized from a mixture of methanol and ethyl acetate to yield 5 - bromo - 3 - (4,5,6,7 - tetrahydro - (1H - 1,3 - diazepin - 2 - ylthio) - indole hydriodide which melts at 243—246° (with decomposition).

WHAT WE CLAIM IS:—

1. Compounds having the general formula I



where Ar is a grouping which along with the



methanol, ethyl acetate and ether to yield 1,2 - dimethyl - 2 - (1 - methyl - 2 - imidazolin - 2 - ylthio) - indole hydriodide which melts at 250—251° (with decomposition).

Example 17


A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml water was added dropwise to a mixture of 7.25 g of 1,2 - dimethylindole and 6.5 g of 1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 120 ml of methanol. The reaction mixture is stirred at room temperature for 2 hours. The crystalline precipitate formed was filtered and recrystallized from a mixture of methanol and ethyl acetate to yield 1,2 - dimethyl - 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 239—240° (with decomposition).

Example 18

A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml of water is added dropwise to a mixture of 7.25 g of 1,2 - dimethylindole in 30 ml of methanol and 5.8 g of 1,4,5,6 - tetrahydro - 2 - pyrimidinethiol in 120 ml of warm methanol. The reaction mixture is stirred for 2 hours at room temperature. The crystalline precipitate formed was filtered and recrystallized from a mixture of methanol and ethyl acetate to yield 1,2 - dimethyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole hydriodide which melts at 246° (with decomposition).

Example 19

A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml of water is added dropwise to a solution of 7.25 g indole in 30 ml methanol and 6.5 g of 4,5,6,7 - tetrahydro - 1H - 1,3 - diazepine - 2 - thiol in 100 ml of methanol. The reaction mixture is stirred for 2 hours at room temperature. The crystalline precipitate formed was filtered off and recrystallized from a mixture of methanol and ethyl acetate to afford 1,2 - dimethyl - 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole hydriodide of the formula



which melts at 245—246° (with decomposition).

Example 20

A solution of 12.7 g of iodine and 20 g of potassium iodide in 100 ml of water is added dropwise to a well-stirred mixture of 5.85 g of indole in 20 ml methanol and

6.5 g of 4,5,6,7 - tetrahydro - 1H - 1,3 - diazepine - 2 - thiol in 100 ml methanol. The reaction mixture is stirred at room temperature for 2 hours. The crystalline precipitate formed was filtered off and recrystallized from a mixture of methanol, ethyl acetate and ether to afford 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole hydriodide which melts at 236° (with decomposition).

Example 21

To a stirred solution of 3.6 g 1,2 - dimethylindole and 2.55 g 2 - imidazolidinethione in 80 ml methanol is added a solution of 6.35 g iodine and 1 g potassium iodide in 50 ml water. After 1 hour at room temperature, the solution is concentrated in vacuo until crystals separate, then cooled and filtered. The precipitate is washed with water and ether and is recrystallized from alcohol to give 1,2 - dimethyl - 3 - (2 - imidazolin - 2 - ylthio) - indole hydriodide melting at 203—205° (with decomposition).

Example 22


To a stirred solution of 1.76 g indole and 1.5 g imidazole 2 - thiol in 50 ml methanol is added a solution of .81 g iodine and 8 g potassium iodide in 30 ml water. After stirring for 3 hours at room temperature, the solution is concentrated in vacuo. An oil separates which crystallises upon addition of ether. The crystals are filtered off and washed with ethyl acetate; and finally recrystallized from a mixture of ethanol and ethyl acetate to give 3 - (2 - imidazol - 2 - ylthio) - indole hydriodide melting at 199—201°.

Example 23

A solution of 6.35 g of iodine and 11 g of potassium iodide in 50 ml of water is added dropwise to a mixture of 4.9 g of 5 - bromoindole in 35 ml methanol and 3.25 g of 4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - thione in 50 ml of methanol. The reaction mixture is allowed to be stirred at room temperature for 2 hours. The crystalline precipitate formed is filtered off and recrystallized from a mixture of methanol and ethyl acetate to yield 5 - bromo - 3 - (4,5,6,7 - tetrahydro - (1H - 1,3 - diazepin - 2 - ylthio) - indole hydriodide which melts at 243—246° (with decomposition).

WHAT WE CLAIM IS:—

1. Compounds having the general formula I



where Ar is a grouping which along with the



* C L M S *

21


11

1,463,885

1

adjacent carbon atoms form an o-arylene or an o-heteroarylene residue, each of R_1 and R_2 independently is hydrogen or a radical of aliphatic character, R_3 is a hydrogen atom, a radical of aliphatic character or a free carboxyl group or functional derivative thereof and Alk represents a substituted or unsubstituted alkylene or alkenylene radical contributing 2-5 carbon atoms to the ring, their tautomers and salts.

2. Compounds having the general formula Ia



wherein X is nitrogen or represents the group CH, and each of R_1 and R_2 is hydrogen, an optionally substituted hydrocarbon radical, a free, etherified or esterified hydroxy or etherified mercapto group, nitro, a free or substituted amino group, an acyl radical, a free or functionally modified carboxy group, each of R_1 and R_2 independently is hydrogen or a lower alkyl group, R_3 is hydrogen, lower alkyl, a free carboxy group or an alkoxy carbonyl group and n is an integer denoting 2-5, their tautomers and acid addition salts.

3. Compounds of the general formula Ia as illustrated in claim 2 in which X represents the group CH, each of R_1 and R_2 independently is hydrogen or lower alkyl and R_3 is hydrogen, lower alkyl, a free carboxy or methoxycarbonyl or ethoxycarbonyl group, and each of R_1 and R_2 is hydrogen, lower alkyl, halogen or a lower alkoxy group and n is an integer denoting 2-4, their tautomers and acid addition salts.

4. Compounds of the general formula Ia as illustrated in claim 2 in which X represents the group CH, R_1 and R_2 independently is hydrogen or lower alkyl and R_3 is hydrogen, lower alkyl or a free carboxy group, and each of R_1 and R_2 is hydrogen, halogen or a lower alkoxy group and n is an integer denoting 2-4, their tautomers and acid addition salts.

5. Compounds of the general formula Ia as illustrated in claim 2 in which X represents the group CH, R_1 and R_2 are hydrogen or methyl and R_3 is hydrogen, methyl or a free carboxy group, and each of R_1 and R_2 is hydrogen, chlorine, bromine or methoxy, and n is an integer denoting 2-4, their tautomers and acid addition salts.

6. The pharmaceutically acceptable acid addition salts of a compound as claimed in claim 1.

7. The pharmaceutically acceptable acid addition salts of a compound as claimed in claim 2.

8. The pharmaceutically acceptable acid

addition salts of a compound as claimed in claim 3. 60

9. The pharmaceutically acceptable acid addition salts of a compound as claimed in claim 4.

10. The pharmaceutically acceptable acid addition salts of a compound as claimed in claim 5. 65

11. 3 - (2 - imidazolin - 2 - ylthio) - indole .

12. 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole. 70

13. 3 - (2 - imidazolin - 2 - ylthio) - indole - 2 - carboxylic acid.

14. 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid. 75

15. 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) 7 - aza - indole.

16. 5 - bromo - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole.

17. 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole. 80

18. 3 - (5 - hydroxy - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole.

19. 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid . 85

20. 5 - methoxy - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid.

21. 2 - methyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole. 90

22. 1 - methyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid.

23. 2 - methyl - 3 - (2 - imidazolin - 2 - ylthio) - indole. 95

24. 5 - methoxy - 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole - 2 - carboxylic acid.

25. 3 - (1 - methyl - 2 - imidazolin - 2 - ylthio) - indole. 100

26. 1,2 - dimethyl - 3 - (1 - methyl - 2 - imidazolin - 2 - ylthio) - indole.

27. 1,2 - dimethyl - 3 - (1 - methyl - 1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole. 105

28. 1,2 - dimethyl - 3 - (1,4,5,6 - tetrahydro - 2 - pyrimidinylthio) - indole.

29. 1,2 - dimethyl - 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole. 110

30. 1,2 - dimethyl - 3 - (2 - imidazolin - 2 - ylthio) - indole .

31. 3 - (2 - imidazol - 2 - ylthio) - indole. 115


31a. 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole.

32. 5 - bromo - 3 - (4,5,6,7 - tetrahydro - 1H - 1,3 - diazepin - 2 - ylthio) - indole.


33. The acid addition salts of a compound as claimed in any one of claims 11 to 32. 120

34. The pharmaceutically acceptable acid addition salts of a compound as claimed in any one of claims 11 to 32.

35. Process for the production of compounds having the general formula I as defined in claim 1 which comprises reacting a compound of the formula II



wherein R_3 and "Alk" have the meanings defined in formula I or a tautomer thereof with a compound of the formula III



wherein R_1 and R_2 have the meanings given under formula I, in the presence of a suitable oxidising agent and if desired converting the resulting compound to an acid addition salt thereof.

36. Process as defined in claim 35 wherein the reaction of compounds of formula II with compounds of formula III is performed in the presence of a halogen as an oxidising agent and a suitable alkali metal salt of a hydrohalic acid.


37. Process as defined in claims 35 and 36 wherein the reaction of compounds of formula II with compounds of formula III is performed in the presence of iodine or bromine as an oxidising agent and a suitable alkali metal salt of a hydrohalic acid.

38. Process as defined in claims 35, 36 and 37 wherein the reaction of compounds of formula II with compounds of formula III is performed in the presence of iodine and potassium iodide.

39. Process as defined in claim 35 wherein the reaction of a compound of formula II with compounds of formula III is performed in the presence of ferric chloride or ferricyanide salts as oxidising agent and a suitable solvent.

40. Process as defined in claims 35-39 wherein in the compounds of formula II and formula III X represents the group CH_2 , R_1 and R_2 are hydrogen or methyl and R_3 is hydrogen, methyl or a free carboxy group, and each of R_4 and R_5 is hydrogen, chlorine, bromine or methoxy, and n is an integer denoting 2-4.


41. Process for the production of compounds having the general formula I as illustrated in claim 1 which comprises reacting a compound of formula IV



wherein R_1 , R_2 and Ar have the meanings given under formula I, R_4 is a free or functionally converted derivative of a hydroxy group, a free mercapto group or a modified mercapto group or amino group, R_5 is an oxygen or sulphur atom, or an imino group or R_4 and R_5 taken together represents a triple bonded nitrogen atom (nitrilo residue) with a compound of the general formula V



in which R_6 and R_7 each depending on the meaning of R_4 and R_5 represents an amino, imino, hydroxy or functionally converted hydroxy, whereby the substituents R_4 - R_7 contain at least 2 nitrogen atoms in order to obtain the ring of formula VI



and if desired, converting a resulting compound of formula I to an acid addition salt thereof.

42. Process as defined in claim 41 wherein in compound of general formula IV the substituents R_4 and R_5 are derivatives of the thiocarbonic, dithiocarbonic or trithiocarbonic acid and the substituents R_6 and R_7 of compounds of formula V are an amino, imino, hydroxy or a reactive esterified hydroxy group.

43. Process as defined in claims 41 and 42 wherein in compounds of general formula IV the substituents R_4 and R_5 are amides, imides, nitriles, halides or intramolecular anhydrides of the thiocarbonic, dithiocarbonic or trithiocarbonic acid and the substituents R_6 and R_7 of compounds of formula V are an amino, imino, hydroxy or a reactive esterified hydroxy group.

44. Process as defined in claims 41, 42 and 43 wherein as a compound of formula IV a pseudothiurea is used in which the substituent R_4 is an NHR_8 group and R_5 an amino group and as a compound of formula V a reactive diester of an alkanediol in which R_6 and R_7 each represent a hydroxy group is used.

45. Process as defined in claim 44, wherein the reaction of a pseudothiurea compound of formula IV with a reactive diester



1,463,885


13

of an alkanediol is carried out in a suitable solvent, and if desired, in the presence of an acid binding agent.


46. Process as claimed in claim 45, wherein the solvent referred to is a lower alcohol.

47. Process as claimed in either of claims 45 or 46, wherein the acid binding agent is an alkali metal or alkaline earth metal hydroxide, carbonate, or hydrogen carbonate.

48. Process for the production of compounds having the general formula I as defined in claim 1, which comprises reacting a compound of general formula VII



15 in which R_1 , R_2 and Ar have the meanings defined under formula I and Y is an anion of an inorganic acid with a compound of formula II



20 wherein R_3 has the same meaning as defined under formula I and if desired, converting a resulting compound of formula I to an acid addition salt thereof.

25 49. Process as defined in claim 48, wherein a compound of formula VII is used in which Y is an anion of hydrochloric, sulphuric, fluoroboric or phosphoric acid.

30 50. Process as claimed in any one of claims 35 to 49, wherein a compound of the formula I thus obtained is converted into a pharmaceutically acceptable acid addition salt.

35 51. Process for the production of compounds having the general formula I as defined in claim 1 substantially as herein described with reference to any one of the foregoing Examples 1 to 14.

40 52. Process for the production of compounds having the general formula I as defined in claim 1 substantially as herein described with reference to any one of the foregoing Examples 15 to 23.

53. A therapeutic composition comprising a compound of formula I as defined in claim 1 and a pharmaceutically acceptable solid carrier or diluent therefore.

54. A vasoconstrictor active composition useful as nasal decongestant agent comprising a compound having the general formula I as defined in claim 1 or a pharmaceutically acceptable acid addition salt of such a compound and a pharmaceutically acceptable diluent or carrier therefore.

55. A vasoconstrictor active composition useful as nasal decongestant agent comprising a compound having the general formula Ia as defined in claim 2 or a pharmaceutically acceptable acid addition salt of such a compound and a pharmaceutically acceptable diluent or carrier therefore.

56. A vaso constrictor active composition useful as nasal decongestant agent comprising a compound having the general formula Ia as defined in claim 3 or a pharmaceutically acceptable acid addition salt of such a compound and a pharmaceutically acceptable diluent or carrier therefore.

57. A vasoconstrictor active composition useful as nasal decongestant agent comprising a compound having the general formula Ia as defined in claim 4 or a pharmaceutically acceptable acid addition salt of such a compound and a pharmaceutically acceptable diluent or carrier therefore.

58. A vasoconstrictor active composition useful as nasal decongestant agent comprising a compound having the general formula Ia as defined in claim 5 or a pharmaceutically acceptable acid addition salt of such a compound and a pharmaceutically acceptable diluent or carrier therefore.

59. A vasoconstrictor active composition useful as nasal decongestant agent comprising the compound 3 - (2 - imidazolin - 2 - ylthio) - indole or a pharmaceutically acceptable acid addition salt of such a compound and a pharmaceutically acceptable diluent or carrier therefore.

60. A hypotensive effective composition useful as antihypertensive agent comprising a compound having the general formula I as defined in claim 1 or a pharmaceutically acceptable acid addition salt of such a compound and pharmaceutically acceptable diluent or carrier therefore.

61. A hypertensive effective composition useful as antihypertensive agent comprising

28



14

1,463,885

14

- 5 a compound having the general formula I as defined in claim 2 or a pharmaceutically acceptable acid addition salt of such a compound or a pharmaceutically acceptable diluent or carrier therefore.

ABEL & IMRAY,
Chartered Patent Agents,
Northumberland House,
303-306, High Holborn,
London, WC1V 7LH.

Printed for Her Majesty's Stationery Office, by the Courier Press, Leamington Spa, 1977
Published by The Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from
which copies may be obtained.