

4-Haloalkyl-3-heterocyclylpyridines and 4-haloalkyl-5-heterocyclyl-pyrimidines, processes for their preparation, compositions comprising them, and their use as pesticides

- 5 The present invention relates to 4-haloalkyl-3-heterocyclylpyridines and 4-haloalkyl-5-heterocyclylpyrimidines, to processes for their preparation, to compositions comprising them and to the use of novel and known 4-haloalkyl-3-heterocyclylpyridines and 4-haloalkyl-5-heterocyclylpyrimidines for controlling animal pests, in particular insects, spider mites,
10 ectoparasites and helminths.

It is already known that appropriately substituted pyridines or pyrimidines have acaricidal and insecticidal activity. Thus, WO 95/07891 describes pyridines which carry a cycloalkyl radical in position 4 which is linked via a
15 hetero atom and a group of various substituents in position 3. WO 93/19050 discloses 4-cycloalkylamino- and 4-cycloalkoxypyrimidines which carry in position 5 inter alia alkyl, alkoxy or haloalkoxy groups. However, the desired activity against the harmful organisms is not always sufficient. Additionally, these compounds often have undesirable toxicologic
20 properties toward mammals and aquatic living beings.

Pyridyl-1,2,4-thiadiazoles having fungicidal properties are described in DE-A 42 39 727. The compounds disclosed therein carry the thiadiazole ring in position 2, 3 or 4 of the unsubstituted pyridine.

25 EP-A 0 371 925 discloses some 1,3,4-oxadiazolyl- and 1,3,4-thiadiazolyl-pyrimidines having nematicidal and fungicidal properties. In the biologically effective compounds disclosed in this publication, the pyrimidine carries the oxadiazolyl or thiadiazolyl ring either

- 30 a) in position 5 and is optionally substituted by a thiomethyl group in position 2, or

- b) in position 2 and is optionally substituted in position 4 and 6, in each case by a methyl group.

Aryltriazole derivatives for use as pesticides are known from EP-A 0 185 256. In addition to the phenyltriazoles, which are particularly preferred, three haloalkyl-3-pyridyltriazoles are disclosed :

- 3-(2-chlorophenyl)-1-methyl-5-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole
- 10 • 3-(2,6-difluorophenyl)-1-methyl-5-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole and
- 3-(2-chloro-4-fluorophenyl)-1-methyl-5-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole,

their desired activity at low application rates, however, is not always satisfactory, especially when controlling insects and spider mites.

Some commercially available 4-haloalkyl-3-heterocyclylpyridines are known from the Maybridge Catalogue 1996/1997, Maybridge Chemical CO. LTD., Trevillet Tintagel, GB:

20

- 3-(3,5-dichlorophenyl)- 5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(4-trifluoromethyl-3-pyridyl)- 3-phenyl-1,2,4-oxadiazole
- 3-(4-trifluoromethyl-3-pyridyl)- 5-phenyl-1,2,4-oxadiazole
- 5-(2-chlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 25 • 5-(3-chlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(4-chlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(2-fluorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(4-fluorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(2,4-dichlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 30 • 5-(3,4-dichlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(3,5-dichlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(2,6-dichloro-4-pyridyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole


- 5-(3,5-bis(trifluoromethyl)phenyl)-3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 2-(2-chlorophenyl)-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 2-(3-chlorophenyl)-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 5 • 2-(4-chlorophenyl)-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 2-(2-trifluoromethoxyphenyl)-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 2-(4-trifluoromethoxyphenyl)-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 10 • 2-(4-trifluoromethyl-3-pyridyl)-5-phenyl-1,3,4-oxadiazole
- 2-(4-trifluoromethyl-3-pyridyl)-4-methylthiazolecarbohydrazide
- ethyl 2-(4-trifluoromethyl-3-pyridyl)-4-methylthiazolecarboxylate
- N-(4-chlorophenyl)carbonyl-N'-[2-(4-trifluoromethyl-3-pyridyl)-4-methyl-5-thiazolyl]carbonylhydrazine
- 15 • 2-(4-trifluoromethyl-3-pyridyl)-4-thiazolecarbohydrazide
- 4-(4-chlorophenyl)-2-(4-trifluoromethyl-3-pyridyl)thiazole
- 4-(4-cyanophenyl)-2-(4-trifluoromethyl-3-pyridyl)thiazole
- N-(4-trifluoromethylphenyl)carbonyl-N'-[2-(4-trifluoromethyl-3-pyridyl)-4-thiazolyl]carbonylhydrazine
- 20 • 2-(2-(4-trifluoromethyl-3-pyridyl)thiazolyl)-5-chloro-3-methylbenzo[b]thiophene
- 2-(4-chlorophenylmethylthio)-5-(4-trifluoromethyl-3-pyridyl)-1-methyl-1H-1,3,4-triazole
- 2-(4-chlorophenylcarbonylmethylthio)-5-(4-trifluoromethyl-3-pyridyl)-1-methyl-1H-1,3,4-triazole and
- 25 • 2-ethoxycarbonylmethylthio-5-(4-trifluoromethyl-3-pyridyl)-1-methyl-1H-1,3,4-triazole.

However, a biological activity toward harmful organisms has hitherto not
 30 been disclosed.

It is an object of the present invention to provide compounds having good insecticidal and acaricidal properties and simultaneously low toxicity toward mammals and aquatic living beings.

- 5 It has now been found that compounds of the formula I, optionally as salts, have a wider activity spectrum against animal pests and simultaneously more favorable toxicologic properties toward mammals and aquatic living beings than the prior art compounds.


10



15

In the formula (I):

- Y is halo-C₁-C₆-alkyl;
 X is CH or N;
 20 m is 0 or 1;
 Q is a 5-membered heterocyclic group



in which

- a) $X^1 = W$, $X^2 = NR^a$, $X^3 = CR^bR^1$ or
 25 b) $X^1 = NR^a$, $X^2 = CR^bR^1$, $X^3 = W$ or
 c) $X^1 = V$, $X^2 = CR^aR^1$, $X^3 = NR^b$ or
 d) $X^1 = V$, $X^2 = CR^aR^2$, $X^3 = CR^bR^3$ or
 e) $X^1 = V$, $X^2 = CR^4R^5$, $X^3 = CR^6R^7$ or
 f) $X^1 = NR^a$, $X^2 = CR^bR^1$, $X^3 = NR^8$;

30

R^a and R^b together are a bond

This Sheet Has Been Replaced





5

V is oxygen, sulfur or NR^9 ;

W is oxygen or sulfur;

R^1 is hydrogen,

(C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₃-C₈)-

5

cycloalkyl,

(C₄-C₈)-cycloalkenyl, (C₆-C₈)-cycloalkynyl,

where the six last-mentioned radicals are optionally substituted
by one or more radicals from the group

halogen, cyano, nitro, hydroxyl, $-\text{C}(=\text{W})\text{R}^{10}$, $-\text{C}(=\text{NOR}^{10})\text{R}^{10}$,

10

$-\text{C}(=\text{NNR}^{10}_2)\text{R}^{10}$, $-\text{C}(=\text{W})\text{OR}^{10}$, $-\text{C}(=\text{W})\text{NR}^{10}_2$, -

$\text{OC}(=\text{W})\text{R}^{10}$, $-\text{OC}(=\text{W})\text{OR}^{10}$, $-\text{NR}^{10}\text{C}(=\text{W})\text{R}^{10}$, -

$\text{N}[\text{C}(=\text{W})\text{R}^{10}]_2$, $-\text{NR}^{10}\text{C}(=\text{W})\text{OR}^{10}$, $-\text{C}(=\text{W})\text{NR}^{10}-\text{NR}^{10}_2$,

$-\text{C}(=\text{W})\text{NR}^{10}-\text{NR}^{10}[\text{C}(=\text{W})\text{R}^{10}]$, $-\text{NR}^{10}-\text{C}(=\text{W})\text{NR}^{10}_2$,

$-\text{NR}^{10}-\text{NR}^{10}\text{C}(=\text{W})\text{R}^{10}$, $-\text{NR}^{10}-\text{N}[\text{C}(=\text{W})\text{R}^{10}]_2$,

15

$-\text{N}[(\text{C}=\text{W})\text{R}^{10}]-\text{NR}^{10}_2$,

$-\text{NR}^{10}-\text{NR}^{10}[(\text{C}=\text{W})\text{R}^{10}]$, $-\text{NR}^{10}-\text{NR}^{10}[(\text{C}=\text{W})\text{WR}^{10}]$,

$-\text{NR}^{10}-\text{R}^{10}[(\text{C}=\text{W})\text{NR}^{10}_2]$, $-\text{NR}^{10}(\text{C}=\text{NR}^{10})\text{R}^{10}$,

$-\text{NR}^{10}(\text{C}=\text{NR}^{10})\text{NR}^{10}_2$,

$-\text{O}-\text{NR}^{10}_2$, $-\text{O}-\text{NR}^{10}(\text{C}=\text{W})\text{R}^{10}$, $-\text{SO}_2\text{NR}^{10}_2$, $-\text{NR}^{10}\text{SO}_2\text{R}^{10}$,

20

$-\text{SO}_2\text{OR}^{10}$, $-\text{OSO}_2\text{R}^{10}$, $-\text{OR}^{10}$, $-\text{NR}^{10}_2$, $-\text{SR}^{10}$, $-\text{SiR}^{10}_3$,

$-\text{SeR}^{10}$, $-\text{PR}^{10}_2$, $-\text{P}(=\text{W})\text{R}^{10}_2$,

$-\text{SOR}^{10}$, $-\text{SO}_2\text{R}^{10}$, $-\text{PW}_2\text{R}^{10}_2$, $-\text{PW}_3\text{R}^{10}_2$, aryl and

heterocyclyl,

the two last-mentioned radicals optionally being substituted
by one or more radicals from the group

25

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl,

(C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₆-C₈)-

cycloalkynyl, (C₁-C₆)-haloalkyl, (C₂-C₆)-haloalkenyl, (C₂-C₆)-haloalkynyl, halogen, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃, -C(=W)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -SOR¹⁰, -SO₂R¹⁰, nitro, cyano and hydroxyl,

5

aryl,

which is optionally substituted by one or more radicals from the group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl,

10

where these six abovementioned radicals are optionally substituted by one or more radicals from the group

halogen, cyano, nitro, halogen, -C(=W)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SOR¹⁰ and -SO₂R¹⁰,

15

halogen, cyano, nitro, -C(=W)R¹⁰, -C(=NOR¹⁰)R¹⁰, -C(=NNR¹⁰₂)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -OC(=W)R¹⁰, -OC(=W)OR¹⁰, -NR¹⁰C(=W)R¹⁰, -N[C(=W)R¹⁰]₂, -NR¹⁰C(=W)OR¹⁰, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃, -PR¹⁰₂, -SOR¹⁰, -SO₂R¹⁰, -PW₂R¹⁰₂ and -PW₃R¹⁰₂,

20

heterocyclyl,

which is optionally substituted by one or more radicals from the group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl,

25

where the six abovementioned radicals are optionally substituted by one or more radicals from the group

- cyano, nitro, halogen, $-C(=W)R^{10}$, $-C(=W)OR^{10}$,
 $-C(=W)NR^{10}_2$, $-NR^{10}C(=W)R^{10}$, $-N[C(=W)R^{10}]_2$,
 $-OC(=W)R^{10}$, $-OC(=W)OR^{10}$, $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$,
 $-SOR^{10}$ and $-SO_2R^{10}$,
- 5 halogen, cyano, nitro, $-C(=W)R^{10}$, $-C(=W)OR^{10}$,
 $-C(=W)NR^{10}_2$, $-OC(=W)R^{10}$, $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$, $-SOR^{10}$
and $-SO_2R^{10}$,
 $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$, $-SOR^{10}$, $-SO_2R^{10}$, $-C(=W)R^{10}$,
 $-C(=NOR^{10})R^{10}$, $-C(=NNR^{10}_2)R^{10}$, $-C(=W)OR^{10}$,
10 $-C(=W)NR^{10}_2$, $-OC(=W)R^{10}$, $-OC(=W)OR^{10}$, $-NR^{10}C(=W)R^{10}$,
 $-N[C(=W)R^{10}]_2$, $-NR^{10}C(=W)OR^{10}$, $-C(=W)NR^{10}-NR^{10}_2$,
 $-C(=W)NR^{10}-NR^{10}[C(=W)R^{10}]$, $-NR^{10}-C(=W)NR^{10}_2$,
 $-NR^{10}-NR^{10}C(=W)R^{10}$, $-NR^{10}-NC(=W)R^{10}_2$, $-N(C=W)R^{10}$
 $-NR^{10}_2$, $-NR^{10}-NR^{10}[(C=W)R^{10}]$, $-NR^{10}-NR^{10}[(C=W)WR^{10}]$,
15 $-NR^{10}-NR^{10}[(C=W)NR^{10}_2]$, $-NR^{10}(C=NR^{10})R^{10}$,
 $-NR^{10}(C=NR^{10})NR^{10}_2$, $-O-NR^{10}_2$, $-O-NR^{10}(C=W)R^{10}$,
 $-SO_2NR^{10}_2$, $-NR^{10}SO_2R^{10}$, $-SO_2OR^{10}$, $-OSO_2R^{10}$,
 $-SC(=W)R^{10}$, $-SC(=W)OR^{10}$, $-SC(=W)R^{10}$, $-PR^{10}_2$, $-PW_2R^{10}_2$,
 $-PW_3R^{10}_2$, SiR^{10}_3 or halogen;

20

R^2 and R^3 independently of one another have the definitions given in R^1 ;

- R^2 and R^3 together form a 5- to 7-membered ring which may be partially
or fully unsaturated and may be interrupted by one or more atoms from the
25 group nitrogen, oxygen and sulfur, the oxygen atoms not being directly
adjacent to one another, and the ring optionally being substituted by one or
more, but at most 5, radicals R^1 ;

R^4 and R^6 independently of one another have the definitions given in R^1 ;

R^4 and R^6 together form a 4- to 7-membered ring which may be partially or fully unsaturated and may be interrupted by one or more atoms from the group nitrogen, oxygen and sulfur, the oxygen atoms not being directly adjacent to one another, and the ring optionally being substituted by one or more, but at most 5, radicals R^1 ;

R^5 and R^7 independently of one another are hydrogen,

(C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₆-C₈)-cycloalkynyl, where the six last-mentioned radicals are optionally substituted by one or more radicals from the group

halogen, cyano, nitro, hydroxyl, $-C(=W)R^{10}$, $-C(=NOR^{10})R^{10}$, $-C(=NNR^{10}_2)R^{10}$, $-C(=W)OR^{10}$, $-C(=W)NR^{10}_2$, $-OC(=W)R^{10}$, $-OC(=W)OR^{10}$, $-NR^{10}C(=W)R^{10}$, $-N[C(=W)R^{10}]_2$, $-NR^{10}C(=W)OR^{10}$, $-C(=W)NR^{10}-NR^{10}_2$, $-C(=W)NR^{10}-NR^{10}[C(=W)R^{10}]$, $-NR^{10}-C(=W)NR^{10}_2$, $-NR^{10}-NR^{10}C(=W)R^{10}$, $-NR^{10}-N[C(=W)R^{10}]_2$, $-N[(C=W)R^{10}]-NR^{10}_2$, $-NR^{10}-NR^{10}[(C=W)R^{10}]$, $-NR^{10}-NR^{10}[(C=W)WR^{10}]$, $-NR^{10}-NR^{10}[(C=W)NR^{10}_2]$, $-NR^{10}(C=NR^{10})R^{10}$, $-NR^{10}(C=NR^{10})NR^{10}_2$, $-O-NR^{10}_2$, $-O-NR^{10}(C=W)R^{10}$, $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$, $-SiR^{10}_3$, $-SeR^{10}$, $-PR^{10}_2$, $-P(=W)R^{10}_2$, $-SOR^{10}$, $-SO_2R^{10}$, $-PW_2R^{10}_2$, $-PW_3R^{10}_2$, aryl and heterocyclyl, of which the two mentioned last are optionally substituted by one or more radicals from the group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
 cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₆-C₈)-cycloalkynyl, (C₁-
 C₆)-haloalkyl, (C₂-C₆)-haloalkenyl, (C₂-C₆)-haloalkynyl,
 halogen, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃, -C(=W)R¹⁰,
 5 -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -SOR¹⁰, -SO₂R¹⁰, nitro,
 cyano and hydroxyl,

aryl,

which is optionally substituted by one or more radicals from the
 group

10 (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
 cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl,

where these six abovementioned radicals are optionally
 substituted by one or more radicals from the group

15 halogen, cyano, nitro, halogen, -C(=W)R¹⁰, -C(=W)OR¹⁰,
 -C(=W)NR¹⁰₂, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SOR¹⁰ and
 -SO₂R¹⁰,

halogen, cyano, nitro, -C(=W)R¹⁰, -C(=NOR¹⁰)R¹⁰,
 -C(=NNR¹⁰₂)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -OC(=W)R¹⁰,
 20 -OC(=W)OR¹⁰, -NR¹⁰C(=W)R¹⁰, -N[C(=W)R¹⁰]₂,
 -NR¹⁰C(=W)OR¹⁰, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃, -PR¹⁰₂,
 -SOR¹⁰, -SO₂R¹⁰, -PW₂R¹⁰₂ and -PW₃R¹⁰₂,

pyridyl,

25 which is optionally substituted by one or more radicals from the
 group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
 cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl,

where the six abovementioned radicals are optionally substituted by one or more radicals from the group

5 cyano, nitro, halogen, $-C(=W)R^{10}$, $-C(=W)OR^{10}$,
 $-C(=W)NR^{10}_2$, $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$, $-SOR^{10}$ and
 $-SO_2R^{10}$,

10 halogen, cyano, nitro, $-C(=W)R^{10}$, $-C(=W)OR^{10}$,
 $-C(=W)NR^{10}_2$, $-OC(=W)R^{10}$, $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$, $-SOR^{10}$
and $-SO_2R^{10}$,

$-C(=W)R^{10}$, $-C(=NOR^{10})R^{10}$, $-C(=NNR^{10}_2)R^{10}$, $-C(=W)OR^{10}$,
 $-C(=W)NR^{10}_2$ or halogen;

15 R^4 and R^5 together form a 4- to 7-membered ring which may be partially unsaturated and may be interrupted by one or more atoms from the group nitrogen, oxygen and sulfur, oxygen atoms not being directly adjacent to one another, and the ring optionally being substituted by one or more, but at most 5, radicals R^1 ;

20 R^4 and R^5 together form one of the groups $=O$, $=S$ or $=N-R^9$;

R^6 and R^7 together form a 5- to 7-membered ring which may be partially unsaturated and may be interrupted by one or more atoms from the group nitrogen, oxygen and sulfur, oxygen atoms not being directly adjacent to one another, and the ring optionally being substituted by one or more, but at most 5, radicals R^1 ;

R^6 and R^7 together form one of the groups $=O$, $=S$ or $=N-R^9$;



R^8 is hydrogen,

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
 cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-
 alkyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkyl, (C₃-C₈)-cycloalkyl-(C₂-
 5 C₄)-alkenyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyl, (C₁-C₆)-alkyl-
 (C₃-C₈)-cycloalkyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkyl, (C₂-C₆)-
 alkynyl-(C₃-C₈)-cycloalkyl, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenyl,
 (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyl,

10 where the fourteen last-mentioned radicals are optionally
 substituted by one or more radicals from the group

halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-
 alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-
 15 haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy,
 (C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-
 halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, (C₃-C₈)-
 cycloalkyl-(C₁-C₄)-alkoxy, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-
 alkoxy, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenyloxy, (C₄-C₈)-
 20 cycloalkenyl-(C₁-C₄)-alkenyloxy, (C₁-C₆)-alkyl-(C₃-C₈)-
 cycloalkoxy, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkoxy, (C₂-C₆)-
 alkynyl-(C₃-C₈)-cycloalkoxy, (C₁-C₆)-alkyl-(C₄-C₈)-
 cycloalkenyloxy, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyloxy,
 (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, (C₁-C₄)-alkoxy-(C₂-C₆)-
 25 alkenyloxy, carbamoyl, (C₁-C₆)-mono- or dialkylcarbamoyl,
 (C₁-C₆)-mono- or dihaloalkylcarbamoyl, (C₃-C₈)-mono- or



- dicycloalkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-
 cycloalkoxycarbonyl, (C₁-C₆)-alkanoyloxy, (C₃-C₈)-
 cycloalkanoyloxy, (C₁-C₆)-haloalkoxycarbonyl, (C₁-C₆)-
 haloalkanoyloxy, (C₁-C₆)-alkaneamido, (C₁-C₆)-
 5 haloalkaneamido, (C₂-C₆)-alkeneamido, (C₃-C₈)-
 cycloalkaneamido, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkaneamido,
 (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-alkynylthio,
 (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio, (C₂-C₆)-
 haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-
 10 cycloalkenylthio, (C₃-C₈)-halocycloalkthio, (C₄-C₈)-
 halocycloalkenylthio, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylthio,
 (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylthio, (C₃-C₈)-cycloalkyl-
 (C₂-C₄)-alkenylthio, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylthio,
 (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylthio, (C₂-C₆)-alkenyl-(C₃-C₈)-
 15 cycloalkylthio, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylthio, (C₁-
 C₆)-alkyl-(C₄-C₈)-cycloalkenylthio, (C₂-C₆)-alkenyl-(C₄-C₈)-
 cycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-alkenylsulfinyl,
 (C₂-C₆)-alkynylsulfinyl, (C₁-C₆)-haloalkylsulfinyl, (C₂-C₆)-
 haloalkenylsulfinyl, (C₂-C₆)-haloalkynylsulfinyl, (C₃-C₈)-
 20 cycloalkylsulfinyl, (C₄-C₈)-cycloalkenylsulfinyl, (C₃-C₈)-
 halocycloalksulfinyl, (C₄-C₈)-halocycloalkenylsulfinyl, (C₃-C₈)-
 cycloalkyl-(C₁-C₄)-alkylsulfinyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-
 alkylsulfinyl, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylsulfinyl, (C₄-
 C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfinyl, (C₁-C₆)-alkyl-(C₃-
 25 C₈)-cycloalkylsulfinyl, (C₂-C₆)-alkenyl-(C₃-C₈)-
 cycloalkylsulfinyl, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylsulfinyl,




- (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₁-C₆)-alkylsulfonyl, (C₂-C₆)-alkenylsulfonyl, (C₂-C₆)-alkynylsulfonyl, (C₁-C₆)-haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl, (C₂-C₆)-haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl, (C₄-C₈)-cycloalkenylsulfonyl, (C₃-C₈)-halocycloalkylsulfonyl, (C₄-C₈)-halocycloalkenylsulfonyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylsulfonyl, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfonyl, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylsulfonyl, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₁-C₆)-alkylamino, (C₂-C₆)-alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-haloalkylamino, (C₂-C₆)-haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-C₈)-cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-halocycloalkylamino, (C₄-C₈)-halocycloalkenylamino, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylamino, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylamino, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylamino, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylamino, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylamino, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylamino, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylamino, (C₁-C₆)-trialkylsilyl, aryl, arylloxy, arylthio, arylamino, arylcarbamoyl, aroyl, aroyloxy,



- aryloxycarbonyl, aryl-(C₁-C₄)-alkoxy, aryl-(C₂-C₄)-alkenyloxy,
 aryl-(C₁-C₄)-alkylthio, aryl-(C₂-C₄)-alkenylthio, aryl-(C₁-C₄)-
 alkylamino, aryl-(C₂-C₄)-alkenylamino, aryl-(C₁-C₆)-
 dialkylsilyl, diaryl-(C₁-C₆)-alkylsilyl, triarylsilyl and 5- or 6-
 5 membered heterocyclyl,
 of which the nineteen last-mentioned radicals are optionally
 substituted in their cyclic moiety by one or more substituents
 from the group
 halogen, cyano, nitro, amino, hydroxyl, thio,
 10 (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy,
 (C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-
 haloalkylthio, (C₁-C₄)-alkylamino, (C₁-C₄)-haloalkylamino,
 formyl and (C₁-C₄)-alkanoyl,
 aryl, which is optionally substituted by one or more radicals from
 15 the group
 halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-
 alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-
 haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy,
 (C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-
 20 halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, carbamoyl,
 (C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl,
 (C₁-C₆)-alkanoyloxy, (C₁-C₆)-mono- or dihaloalkylcarbamoyl,
 (C₁-C₆)-haloalkoxycarbonyl, (C₁-C₆)-haloalkanoyloxy, (C₁-
 C₆)-alkaneamido, (C₁-C₆)-haloalkaneamido, (C₂-C₆)-
 25 alkeneamido, (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-
 alkynylthio, (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio,
 (C₂-C₆)-haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-



- cycloalkenylthio, (C₃-C₈)-halocycloalkthio, (C₃-C₈)-
 halocycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-
 alkenylsulfinyl, (C₂-C₆)-alkynylsulfinyl, (C₁-C₆)-
 haloalkylsulfinyl, (C₂-C₆)-haloalkenylsulfinyl, (C₂-C₆)-
 haloalkynylsulfinyl, (C₃-C₈)-cycloalkylsulfinyl, (C₄-C₈)-
 cycloalkenylsulfinyl, (C₃-C₈)-halocycloalksulfinyl, (C₄-C₈)-
 halocycloalkenylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₂-C₆)-
 alkenylsulfonyl, (C₂-C₆)-alkynylsulfonyl, (C₁-C₆)-
 haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl, (C₂-C₆)-
 haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl, (C₄-C₈)-
 cycloalkenylsulfonyl, (C₃-C₈)-halocycloalksulfonyl, (C₄-C₈)-
 halocycloalkenylsulfonyl, (C₁-C₆)-alkylamino, (C₂-C₆)-
 alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-haloalkylamino,
 (C₂-C₆)-haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-
 C₈)-cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-
 halocycloalkamino and (C₄-C₈)-halocycloalkenylamino,
 -C(=W)R¹¹, OR¹¹ or NR¹¹₂;
- R⁹ is (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
 cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-
 alkyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkyl, (C₃-C₈)-cycloalkyl-
 (C₂-C₄)-alkenyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyl,
 where the nine last-mentioned radicals are optionally substituted
 by one or more radicals from the group
- halogen, cyano, (C₁-C₆)-alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-
 alkynyloxy and (C₁-C₆)-haloalkyloxy;

R¹⁰

is hydrogen,

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
 cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-
 alkyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkyl, (C₃-C₈)-cycloalkyl-
 (C₂-C₄)-alkenyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyl, (C₁-C₆)-
 alkyl-(C₃-C₈)-cycloalkyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkyl,
 (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkyl, (C₁-C₆)-alkyl-(C₄-C₈)-
 cycloalkenyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyl,

where the fourteen last-mentioned radicals are optionally
 substituted by one or more radicals from the group


halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-
 alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-
 haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy,
 (C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-
 halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, (C₃-C₈)-
 cycloalkyl-(C₁-C₄)-alkoxy, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-
 alkoxy, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenyloxy, (C₄-C₈)-
 cycloalkenyl-(C₁-C₄)-alkenyloxy, (C₁-C₆)-alkyl-(C₃-C₈)-
 cycloalkoxy, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkoxy, (C₂-C₆)-
 alkynyl-(C₃-C₈)-cycloalkoxy, (C₁-C₆)-alkyl-(C₄-C₈)-
 cycloalkenyloxy, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyloxy,
 (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, (C₁-C₄)-alkoxy-(C₂-C₆)-
 alkenyloxy, carbamoyl,

(C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-mono- or
 dihaloalkylcarbamoyl, (C₃-C₈)-mono- or

dicycloalkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-



- cycloalkoxycarbonyl, (C₁-C₆)-alkanoyloxy, (C₃-C₈)-
 cycloalkanoyloxy, (C₁-C₆)-haloalkoxycarbonyl, (C₁-C₆)-
 haloalkanoyloxy, (C₁-C₆)-alkaneamido, (C₁-C₆)-
 haloalkaneamido, (C₂-C₆)-alkeneamido, (C₃-C₈)-
 5 cycloalkaneamido, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkaneamido,
 (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-alkynylthio,
 (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio, (C₂-C₆)-
 haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-
 cycloalkenylthio, (C₃-C₈)-halocycloalkthio, (C₄-C₈)-
 10 halocycloalkenylthio, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylthio,
 (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylthio, (C₃-C₈)-cycloalkyl-
 (C₂-C₄)-alkenylthio, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylthio,
 (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylthio, (C₂-C₆)-alkenyl-(C₃-C₈)-
 cycloalkylthio, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylthio, (C₁-C₆)-
 15 alkyl-(C₄-C₈)-cycloalkenylthio, (C₂-C₆)-alkenyl-(C₄-C₈)-
 cycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-alkenylsulfinyl,
 (C₂-C₆)-alkynylsulfinyl, (C₁-C₆)-haloalkylsulfinyl, (C₂-C₆)-
 haloalkenylsulfinyl, (C₂-C₆)-haloalkynylsulfinyl, (C₃-C₈)-
 cycloalkylsulfinyl, (C₄-C₈)-cycloalkenylsulfinyl, (C₃-C₈)-
 20 halocycloalksulfinyl, (C₄-C₈)-halocycloalkenylsulfinyl, (C₃-C₈)-
 cycloalkyl-(C₁-C₄)-alkylsulfinyl,
 (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylsulfinyl, (C₃-C₈)-cycloalkyl-
 (C₂-C₄)-alkenylsulfinyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-
 alkenylsulfinyl, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylsulfinyl, (C₂-
 25 C₆)-alkenyl-(C₃-C₈)-cycloalkylsulfinyl, (C₂-C₆)-alkynyl-(C₃-
 C₈)-cycloalkylsulfinyl, (C₁-C₆)-alkyl-(C₄-C₈)-



- cycloalkenylsulfinyl, (C₂-C₆)-alkenyl-(C₄-C₈)-
 cycloalkenylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₂-C₆)-
 alkenylsulfonyl,
 (C₂-C₆)-alkynylsulfonyl, (C₁-C₆)-haloalkylsulfonyl, (C₂-C₆)-
 5 haloalkenylsulfonyl, (C₂-C₆)-haloalkynylsulfonyl, (C₃-C₈)-
 cycloalkylsulfonyl, (C₄-C₈)-cycloalkenylsulfonyl, (C₃-C₈)-
 halocycloalksulfonyl, (C₄-C₈)-halocycloalkenylsulfonyl,
 (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylsulfonyl, (C₄-C₈)-cycloalkenyl-
 (C₁-C₄)-alkylsulfonyl, (C₃-C₈)-cycloalkyl-(C₂-C₄)-
 10 alkenylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfonyl,
 (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkenyl-
 (C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkynyl-(C₃-C₈)-
 cycloalkylsulfonyl, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylsulfonyl,
 (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylsulfonyl,
 15 (C₁-C₆)-alkylamino, (C₂-C₆)-alkenylamino, (C₂-C₆)-
 alkynylamino, (C₁-C₆)-haloalkylamino, (C₂-C₆)-
 haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-C₈)-
 cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-
 halocycloalkamino, (C₄-C₈)-halocycloalkenylamino, (C₃-C₈)-
 20 cycloalkyl-(C₁-C₄)-alkylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-
 alkylamino, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylamino,
 (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylamino, (C₁-C₆)-alkyl-
 (C₃-C₈)-cycloalkylamino, (C₂-C₆)-alkenyl-(C₃-C₈)-
 cycloalkylamino, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylamino,
 25 (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylamino, (C₂-C₆)-alkenyl-
 (C₄-C₈)-cycloalkenylamino, (C₁-C₆)-trialkylsilyl, aryl, aryloxy,



5

arylthio, arylamino, aryl-(C₁-C₄)-alkoxy, aryl-(C₂-C₄)-
alkenyloxy, aryl-(C₁-C₄)-alkylthio, aryl-(C₂-C₄)-alkenylthio,
aryl-(C₁-C₄)-alkylamino, aryl-(C₂-C₄)-alkenylamino, aryl-
(C₁-C₆)-dialkylsilyl, diaryl-(C₁-C₆)-alkylsilyl, triarylsilyl and 5-
or 6-membered heterocyclyl,
where the cyclic moiety of the fourteen last-mentioned
radicals is optionally substituted by one or more radicals from
the group

10

halogen, cyano, nitro, amino, hydroxyl, thio, (C₁-C₄)-alkyl,
(C₁-C₄)-haloalkyl, (C₃-C₈)-cycloalkyl, (C₁-C₄)-alkoxy,
(C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-halo-
alkylthio, (C₁-C₄)-alkylamino, (C₁-C₄)-haloalkylamino,
formyl and (C₁-C₄)-alkanoyl,

15

aryl, 5- or 6-membered heteroaromatic,
where the two last-mentioned radicals are optionally substituted
by one or more radicals from the group

20

halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-
alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-
haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy,
(C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-
halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, carbamoyl,
(C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl,
(C₁-C₆)-alkanoyloxy,

25

(C₁-C₆)-mono- or dihaloalkylcarbamoyl, (C₁-C₆)-
haloalkoxycarbonyl, (C₁-C₆)-haloalkanoyloxy, (C₁-C₆)-
alkaneamido, (C₁-C₆)-haloalkaneamido, (C₂-C₆)-
alkeneamido, (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-



alkynylthio, (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio,
 (C₂-C₆)-haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-
 cycloalkenylthio,
 (C₃-C₈)-halocycloalkthio, (C₄-C₈)-halocycloalkenylthio,
 5 (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-alkenylsulfinyl, (C₂-C₆)-
 alkynylsulfinyl, (C₁-C₆)-haloalkylsulfinyl, (C₂-C₆)-
 haloalkenylsulfinyl, (C₂-C₆)-haloalkynylsulfinyl, (C₃-C₈)-
 cycloalkylsulfinyl, (C₄-C₈)-cycloalkenylsulfinyl, (C₃-C₈)-
 halocycloalksulfinyl, (C₄-C₈)-halocycloalkenylsulfinyl, (C₁-C₆)-
 10 alkylsulfonyl, (C₂-C₆)-alkenylsulfonyl, (C₂-C₆)-alkynylsulfonyl,
 (C₁-C₆)-haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl,
 (C₂-C₆)-haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl,
 (C₄-C₈)-cycloalkenylsulfonyl, (C₃-C₈)-halocycloalksulfonyl,
 (C₄-C₈)-halocycloalkenylsulfonyl, (C₁-C₆)-alkylamino,
 15 (C₂-C₆)-alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-
 haloalkylamino, (C₂-C₆)-haloalkenylamino, (C₂-C₆)-
 haloalkynylamino, (C₃-C₈)-cycloalkylamino, (C₄-C₈)-
 cycloalkenylamino, (C₃-C₈)-halocycloalkamino and (C₄-C₈)-
 halocycloalkenylamino;

20

R¹¹is (C₁-C₁₀)-alkyl, haloalkyl, aryl,which is optionally substituted by one or more radicals from the
grouphalogen, cyano, nitro, (C₁-C₄)-alkoxy, (C₁-C₄)-alkyl, amino,

25

(C₁-C₄)-monoalkylamino and (C₁-C₄)-dialkylamino,NR¹⁰₂, OR¹⁰ or SR¹⁰.

The term "halogen" includes fluorine, chlorine, bromine and iodine.

The term "(C₁-C₄)-alkyl" is to be understood as a straight-chain or branched hydrocarbon radical having 1, 2, 3 or 4 carbon atoms, such as, for example, the methyl, ethyl, propyl, isopropyl, 1-butyl, 2-butyl, 2-methylpropyl or tert-butyl radical. Correspondingly, alkyl radicals having a greater range of carbon atoms are to be understood as straight-chain or branched saturated hydrocarbon radicals which contain a number of carbon atoms which corresponds to the range stated. Thus, the term "(C₁-C₆)-alkyl" includes the abovementioned alkyl radicals, and, for example, the pentyl, 2-methylbutyl, 1,1-dimethylpropyl, hexyl radical. The term "(C₁-C₁₀)-alkyl" is to be understood as the abovementioned alkyl radicals, and, for example, the nonyl, 1-decyl or 2-decyl radical and the term "(C₁-C₂₀)-alkyl" is to be understood as the abovementioned alkyl radicals, and, for example, the undecyl, dodecyl, pentadecyl or eicosyl radical.

"(C₁-C₄)-Haloalkyl" is to be understood as an alkyl group mentioned under the term "(C₁-C₄)-alkyl" in which one or more hydrogen atoms are replaced by the same number of identical or different halogen atoms, preferably by fluorine or chlorine, such as the trifluoromethyl, the 1-fluoroethyl, the 2,2,2-trifluoroethyl, the chloromethyl, fluoromethyl, the difluoromethyl and the 1,1,2,2-tetrafluoroethyl group.

"(C₁-C₄)-Alkoxy" is to be understood as an alkoxy group whose hydrocarbon radical has the meaning given under the term "(C₁-C₄)-alkyl". Alkoxy groups embracing a greater range of carbon atoms are to be understood correspondingly.

The terms "alkenyl" and "alkynyl" having a prefix stating the range of carbon atoms denote a straight-chain or branched hydrocarbon radical having a number of carbon atoms corresponding to the range stated which comprises at least one multiple bond which may be in any position of the

unsaturated radical in question. "(C₂-C₄)-Alkenyl" is thus, for example, the vinyl, allyl, 2-methyl-2-propene or 2-butenyl group; "(C₂-C₆)-alkenyl" denotes the abovementioned radicals and, for example, the pentenyl, 2-methylpentenyl or the hexenyl group. The term "(C₂-C₂₀)-alkenyl" is to be understood as the abovementioned radicals and, for example, the 2-decenyl or the 2-eicosenyl group. "(C₂-C₄)-Alkynyl" is, for example, the ethynyl, propargyl, 2-methyl-2-propyne or 2-butyne group. "(C₂-C₆)-Alkynyl" is to be understood as the abovementioned radicals and, for example, the 2-pentynyl- or the 2-hexynyl group and "(C₂-C₂₀)-alkynyl" is to be understood as the abovementioned radicals and, for example, the 2-octynyl or the 2-decynyl group.

"(C₃-C₈)-Cycloalkyl" denotes monocyclic alkyl radicals, such as the cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl radical and bicyclic alkyl radicals, such as the norbornyl radical.

The term "(C₃-C₈)-cycloalkyl-(C₁-C₄)-alkyl" is to be understood as, for example, the cyclopropylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclohexylethyl and cyclohexylbutyl radical, and the term "(C₁-C₆)-alkyl-(C₃-C₈)-cycloalkyl is to be understood as, for example, the 1-methylcyclopropyl, 1-methylcyclopentyl, 1-methylcyclohexyl, 3-hexylcyclobutyl and 4-tert-butyl-cyclohexyl radical.

"(C₁-C₄)-Alkoxy-(C₁-C₆)-alkyloxy" is an alkoxy group as defined above which is substituted by a further alkoxy group, such as, for example, 1-ethoxyethoxy.

"(C₃-C₈)-Cycloalkoxy" or "(C₃-C₈)-cycloalkylthio" is to be understood as one of the abovementioned (C₃-C₈)-cycloalkyl radicals which is linked via an oxygen or sulfur atom.

"(C₃-C₈)-Cycloalkyl-(C₁-C₆)-alkoxy" is, for example, the cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy, cyclohexylethoxy or the cyclohexylbutoxy group;

- 5 The term "(C₁-C₄)-alkyl-(C₃-C₈)-cycloalkoxy" is, for example, the methylcyclopropyloxy, methylcyclobutyloxy or the butylcyclohexyloxy group.

"(C₁-C₆)-Alkylthio" is an alkylthio group whose hydrocarbon radical has the meaning given under the term "(C₁-C₆)-alkyl".

- 10 Correspondingly, "(C₁-C₆)-alkylsulfinyl" is, for example, the methyl-, ethyl-, propyl-, isopropyl-, butyl-, isobutyl-, sec-butyl- or tert-butylsulfinyl group and "(C₁-C₆)-alkylsulfonyl" is, for example, the methyl-, ethyl-, propyl-, isopropyl-, butyl-, isobutyl-, sec-butyl- or tert-butylsulfonyl group.

- 15 "(C₁-C₆)-Alkylamino" is a nitrogen atom which is substituted by one or two identical or different alkyl radicals of the above definition.

- The term "(C₁-C₆)-mono- or dialkylcarbamoyl" is a carbamoyl group having one or two hydrocarbon radicals which have the meaning given under the
20 term "(C₁-C₆)-alkyl" and which, in the case of two hydrocarbon radicals, may be identical or different.

- Correspondingly, "(C₁-C₆)-dihaloalkylcarbamoyl" is a carbamoyl group which carries two (C₁-C₆)-haloalkyl radicals in accordance with the above
25 definition or one (C₁-C₆)-haloalkyl radical and one (C₁-C₆)-alkyl radical in accordance with the above definition.

"(C₁-C₆)-Alkanoyl" is, for example, the acetyl, propionyl, butyryl or 2-methylbutyryl group.

The term "aryl" is to be understood as an isocyclic aromatic radical preferably having 6 to 14, in particular 6 to 12, carbon atoms, such as, for example, phenyl, naphthyl or biphenyl, preferably phenyl. "Aroyl" is thus an aryl radical as defined above which is attached via a carbonyl group, such as, for example, the benzoyl group.

The term "heterocyclyl" denotes a cyclic radical which may be fully saturated, partially unsaturated or fully unsaturated and which may be interrupted by at least one or more identical atoms from the group nitrogen, sulfur or oxygen, oxygen atoms, however, not being directly adjacent to one another and at least one carbon atom being present in the ring, such as, for example, a thiophene, furan, pyrrole, thiazole, oxazole, imidazole, isothiazole, isoxazole, pyrazole, 1,3,4-oxadiazole, 1,3,4-thiadiazole, 1,3,4-triazole, 1,2,4-oxadiazole, 1,2,4-thiadiazole, 1,2,4-triazole, 1,2,3-triazole, 1,2,3,4-tetrazole, benzo[b]thiophene, benzo[b]furan, indole, benzo[c]thiophene, benzo[c]furan, isoindole, benzoxazole, benzothiazole, benzimidazole, benzisoxazole, benzisothiazole, benzopyrazole, benzothiadiazole, benzotriazole, dibenzofuran, dibenzothiophene, carbazole, pyridine, pyrazine, pyrimidine, pyridazine, 1,3,5-triazine, 1,2,4-triazine, 1,2,4,5-triazine, quinoline, isoquinoline, quinoxaline, quinazoline, cinnoline, 1,8-naphthyridine, 1,5-naphthyridine, 1,6-naphthyridine, 1,7-naphthyridine, phthalazine, pyridopyrimidine, purine, pteridine 4H-quinolizine; piperidine, pyrrolidine, oxazoline, tetrahydrofuran, tetrahydropyran, isoxazolidine or thiazolidine radical. The term "heteroaromatic" thus embraces, from among the meanings mentioned above under "heterocyclyl", in each case the fully unsaturated aromatic heterocyclic compounds.

"Aryl-(C₁-C₄)-alkoxy" is an aryl radical which is attached via a (C₁-C₄)-alkoxy group, for example the benzyloxy, phenylethoxy, phenylbutoxy or naphthylmethoxy radical.

"Arylthio" is an aryl radical attached via a sulfur atom, for example the phenylthio or the 1- or 2-naphthylthio radical. Correspondingly, "aryloxy" is, for example, the phenoxy or 1- or 2-naphthyloxy radical.


- 5 "Aryl-(C₁-C₄)-alkylthio" is an aryl radical which is attached via an alkylthio radical, for example the benzylthio, naphthylmethylthio or the phenylethylthio radical.

The term "(C₁-C₆)-trialkylsilyl" denotes a silicon atom which carries three
 10 identical or different alkyl radicals in accordance with the above definition. Correspondingly "aryl-(C₁-C₆)-dialkylsilyl" is a silicon atom which carries one aryl radical and two identical or different radicals in accordance with the above definition, "diaryl-(C₁-C₆)-alkylsilyl" is a silicon atom which carries
 15 one alkyl radical and two identical or different aryl radicals in accordance with the above definition, and "triarylsilyl" is a silicon atom which carries three identical or different aryl radicals in accordance with the above definition.


In cases where two or more radicals R¹⁰ are present in a substituent, such
 20 as, for example, in -C(=W)NR¹⁰₂, these radicals may be identical or different.

Preference is given to those compounds of the formula I in which

- Y is C₁-C₆-alkyl which is mono- or polysubstituted by chlorine
 25 and/or fluorine;
 m is zero;
 Q is a 5-membered heterocyclic group



in which



5 R^a and R^b together are a bond ;

R^1, R^2, R^3, R^4 and R^6 are each independently of one another

hydrogen, halogen, C₁-C₁₂-alkyl, C₃-C₈-cycloalkyl, C₂-C₈-

alkenyl, C₂-C₈-alkynyl, where the four last-mentioned

hydrocarbon radicals are optionally mono- or polysubstituted

10 by identical or different radicals from a group A1 consisting of

C₁-C₆-alkylcarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-

alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, C₁-C₆-

alkylcarbonylamino, C₁-C₆-alkylsulfonylamino, phenyl, furyl,

pyrryl, thienyl, halogen, cyano, phenyloxy, phenylthio and

15 phenylamino, where the eleven first-mentioned radicals of

group A1 are each optionally mono- or polysubstituted by

identical or different radicals from a group B1 consisting of

halogen, cyano, C₁-C₃-alkoxy and phenyl which is optionally

mono- or polysubstituted by one or more halogen atoms and

20 where the three last-mentioned radicals of group A1 are each

optionally mono- or polysubstituted by identical or different

radicals from a group B2 consisting of H halogen, cyano,

nitro, C₁-C₃-alkyl and C₁-C₃-alkoxy, or are C₁-C₆-

alkylcarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-

25 alkoxycarbonyl, phenyl, pyridyl, furyl, thienyl, pyrryl, where the

eight last-mentioned radicals are optionally mono- or

polysubstituted by identical or different radicals from group

B1, or are OR^{10} , SR^{10} or $N(R^{10})_2$;

- R^5 and R^7 are each independently of one another hydrogen, halogen, C₁-C₁₂-alkyl, C₃-C₈-cycloalkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, where the four last-mentioned hydrocarbon radicals are optionally mono- or polysubstituted by identical or different radicals from a group A2 consisting of C₁-C₆-alkylcarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, C₁-C₆-alkylcarbonylamino, phenyl, furyl, pyrrol, thienyl, halogen, cyano, phenyloxy, phenylthio and phenylamino, where the ten first-mentioned radicals of group A2 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A2 are each optionally mono- or polysubstituted by identical or different radicals from the group B2, or are C₁-C₆-alkylcarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-alkoxycarbonyl, phenyl, pyridyl, furyl, thienyl, pyrrol, where the eight last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from the group B1, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;
- R^{10} is hydrogen, benzyl, C₁-C₆-alkyl, C₁-C₆-cycloalkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, phenyl, C₁-C₆-alkylcarbonyl or C₁-C₆-alkylsulfonyl, where the eight last-mentioned radicals are optionally mono- or polysubstituted by identical or different halogen atoms.

Particular preference is given to compounds of the formula I in which Y is trifluoromethyl;

R^1 , R^2 , R^3 , R^4 and R^6 are each independently of one another halogen, C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or

different radicals from a group A3 consisting of C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, C₁-C₄-alkylsulfonylamino, phenyl, furyl, pyrrol, thienyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the eleven first-mentioned radicals of group A3 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A3 are each optionally mono- or polysubstituted by identical or different radicals from the group B2, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;

R⁵ and R⁷ are each independently of one another halogen, C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A4 consisting of C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, phenyl, furyl, pyrrol, thienyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the ten first-mentioned radicals of group A4 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A4 are each optionally mono- or polysubstituted by identical or different radicals from the group B2, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;

R¹⁰ is hydrogen, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, phenyl, C₁-C₄-alkylcarbonyl or C₁-C₄-alkylsulfonyl, where the six last-mentioned radicals are optionally mono- or polysubstituted by identical or different halogen atoms.

Very particular preference is given to compounds of the formula I in which R^1 , R^2 , R^3 , R^4 and R^6 are each independently of one another C₁-C₁₀-

5 alkyl, C₂-C₁₀-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A5 consisting of C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, C₁-C₄-alkylsulfonylamino, phenyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the
10 eight first-mentioned radicals of group A5 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A5 are each optionally mono- or polysubstituted by identical or different radicals from the group B2;

15 R^5 and R^7 are each independently of one another C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A6 consisting of

20 C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, phenyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the seven first-mentioned radicals of group A6 are each optionally mono- or polysubstituted by identical or different
25 radicals from the group B1 and the three last-mentioned radicals of group A6 are each optionally mono- or polysubstituted by identical or different radicals from the group B2.

30 Depending on the nature of the substituents defined above, the compounds of the formula (I) have acidic or basic properties and can form salts. If the compounds of the formula (I) carry, for example, groups such as hydroxyl, carboxyl and other groups inducing acidic properties, these

compounds can be reacted with bases to give salts. Suitable bases are, for example, hydroxides, carbonates, bicarbonates of the alkali metals and alkaline earth metals, in particular those of sodium, potassium, magnesium and calcium, further ammonia, primary, secondary and tertiary amines

- 5 having (C₁-C₄)-alkyl radicals and also mono-, di- and trialkanolamines of (C₁-C₄)-alkanols. If the compounds of the formula (I) carry, for example, groups such as amino, alkylamino and other groups inducing basic properties, these compounds can be reacted with acids to give salts. Suitable acids are, for example, mineral acids, such as hydrochloric acid, sulfuric acid and phosphoric acid, organic acids, such as acetic acid, oxalic acid and acidic salts, such as NaHSO₄ and KHSO₄. The salts which can be obtained in this manner likewise have insecticidal, acaricidal and nematicidal properties.


- 15 The compounds of the formula (I) may have one or more asymmetric carbon atoms or stereoisomers on double bonds. Enantiomers or diastereomers may therefore be present. The invention embraces both the pure isomers and mixtures thereof. The mixtures of diastereomers can be separated into the isomers by customary methods, for example by selective crystallization from suitable solvents or by chromatography. Racemates can be separated into the enantiomers by customary methods.

The present invention also provides processes for preparing compounds of the formula I:


- 25 To prepare compounds of the formula (I) in which

$$a) \quad X^1 = W, X^2 = NR^a, X^3 = CR^bR^1$$

and R^a, R^b and R¹ are as defined above and W is oxygen, activated derivatives of the acid of the formula (II)



where X and Y are as defined above, are reacted in the presence of a base with a compound of the formula (III)




in which the radical R^1 is as defined in formula (I). Suitable activated

- 5 derivatives are, for example, acyl halides, esters and anhydrides. Suitable bases are amines, such as triethylamine, diisopropylethylamine, pyridine or lutidine, alkali metal hydroxides, alkali metal alkoxides, such as sodium ethoxide or potassium tert-butoxide, or alkylmetal compounds, such as butyllithium.

10

Depending on the conditions, the reaction described above can be carried out as a one-step process or as a two-step process via intermediates of the formula (IV):



15


Compounds of the formula (IV) can be cyclized to the 1,2,4-oxadiazoles by heating in an inert solvent at temperatures of up to 180°C.

- Compounds of the formula (IV) are also directly obtainable from the acid of
 20 the formula (II) and amidoximes of the formula (III) by using a dehydrating reagent such as dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide or N,N'-carbonyldiimidazole.


- Both acids of the formula (II) and amidoximes of the formula (III) are
 25 commercially available or can be prepared by methods known from the literature (see, for example: Houben-Weyl, Methoden der organischen

Chemie, Volume X/4, pages 209-212; EP-A 0 580 374; G.F. Holland, J.N.Pereira, J. Med. Chem., 1967, 10, 149).


- In the abovementioned case a) where W is sulfur, the compounds of the formula (I) can be obtained in a manner known from the literature by reaction of a compound of the formula (VII) with an electrophilic amination reagent, such as hydroxylamine-O-sulfonic acid (Y. Lin, S.A. Lang, S.R. Petty, J. Org. Chem. 1980, 45, 3750).




- The compounds of the formula (VII) required as starting materials for this reaction can be prepared by reacting the thioamides of the formula (VIII) with dialkylamide dialkyl acetals, formula (IX), where R^1 is as defined above and R^{12} and R^{13} are each C_1 - C_4 -alkyl.



To prepare compounds of the formula (I) in which




- 20 and R^a , R^b and R^1 are as defined above, and W is oxygen, amidoximes of the formula (V) can be reacted with activated derivatives of the acids of the formula (VI) or with the acids of the formula (VI) themselves.



To prepare compounds of the formula (I) in which

c) $X^1 = V$, $X^2 = CR^aR^1$, $X^3 = NR^b$


- 5 and R^a , R^b and R^1 are as defined above and V is sulfur, N,N'-diacylhydrazines of the formula (XIII) can be cyclized with a thiolation reagent, such as Lawesson's reagent (A.A. El-Barbary, S. Scheibyl, S.O. Lawesson, H. Fritz, Acta Chem. Scand. 1980, 597), in an inert solvent, such as toluene.




10

In the abovementioned case b) where W is oxygen, the compounds of the formula (I) can be prepared by reaction of acids of the formula (II) with hydrazines of the formula (X), in which R^1 is as defined above, using an activating reagent, such as phosphorus oxychloride or phosphorus pentachloride.

15




It is also possible to react acid hydrazides of the formula (XI) with ortho esters of the formula (XII) where R^1 is as defined above, and R^{12} is (C₁-C₄)-alkyl.




The reaction can be carried out with or without solvent and with or without an activating reagent. Suitable solvents are hydrocarbons, such as toluene, or ethers, such as 1,2-dimethoxyethane. A suitable activating reagent is, for example, phosphorus oxychloride. The reaction temperature is generally the reflux temperature of the solvent.

10 To prepare compounds of the formula (I) in which




and R^a , R^b and R^3 are as defined above and V is oxygen, compounds of the formula (XIV) are reacted with a dehydrating reagent.



- 15 Suitable dehydrating reagents are inorganic acyl chlorides, such as phosphorus oxychloride or thionyl chloride, inorganic acids, such as sulfuric acid or polyphosphoric acid, or a mixture of phosphoric acid and acetic anhydride (Houben-Weyl, Methoden der organischen Chemie, Volume E8a, pages 935 - 941).
- 20 The reaction can be carried out with or without a solvent. Suitable solvents are inert solvents, such as toluene, benzene, dimethoxyethane,

dimethylformamide, dimethylacetamide and chlorobenzene. The reaction temperature is advantageously in a range between 50°C and 150°C .


- Compounds of the formula (XIV) can be obtained, for example, by oxidation of the corresponding hydroxyl compound of the formula (XV), it being possible to employ all reagents which are customarily used for this purpose in organic chemistry. (Milos Hudlický, "Oxidations in Organic Chemistry", ACS Monograph 186, American Chemical Society, Washington, DC, 1990)



10

In the abovementioned case d) where V is sulfur, the compounds of the formula (I) can be prepared by condensation of thioamides of the formula (XVII) with carbonyl derivatives of the formula (XVIII), where Z is halogen, in particular chlorine or bromine, acyloxy or sulfonyloxy, in particular methanesulfonyloxy or tolylsulfonyloxy.

15




To prepare compounds of the formula (I) in which


e) $X^1 = V$, $X^2 = CR^4R^5$, $X^3 = CR^6R^7$

- and R^4 , R^5 , R^6 and R^7 are as defined above and V is oxygen, compounds of the formula (XV) are reacted with cyclization reagents, such as Burgess' reagent (G.M. Atkins, E.M. Burgess, J. Am. Chem. Soc. 1968, 90, 4744.), in a solvent such as tetrahydrofuran and 1,4-dioxane, at a temperature

which is in a range between room temperature and the reflux temperature of the solvent.




Compounds of the formula (XV) can be obtained by reacting activated derivatives of the acid in formula (II) with β -aminoalcohols of the formula (XVI), if appropriate in the presence of a base, such as, for example, triethylamine, in an inert solvent, such as, for example, dichloromethane.



An acyl halide or an anhydride can be used as activated derivative of the acid.


A number of β -aminoalcohols of the formula (XVI) are commercially available. For others, there is a large number of preparation procedures in the literature, for example a reduction of α -amino acids (B.M. Trost "Comprehensive Organic Synthesis, Reduction", Volume 8, Pergamon Press, Oxford, 1991).

In the abovementioned case e) where V is sulfur, the compounds of the formula (I) can be prepared by reaction of thioamides of the formula (XVII) with compounds of the formula (XIX), the two substituents Z being as defined above and either identical or different (A.R. Katritzky "Comprehensive Heterocyclic Chemistry", Volume 6, pages 306 - 312, Pergamon Press, Oxford).




Thioamides of the formula (XVII) are either commercially available or can be obtained by addition of hydrogen sulfide to the corresponding carbonitriles in the presence of a base (A.E.S. Fairfull, J.L. Lowe, D.A. Peak, J. Chem. Soc. 1952, 742).



For preparing compounds of the formula (I) in which



and R^a , R^b , R^1 and R^8 are as defined above, hydrazides of the formula (XX)



are reacted with a compound of the formula (XXI) or with thioamides of the formula (XXII) (Houben-Weyl, Methoden der organischen Chemie, Volume E8d, pages 510 - 512).



This reaction can be carried out with or without using a solvent, suitable solvents being alcohols, such as ethanol and propanol, or aromatic hydrocarbons, such as toluene and xylene. If the reaction is carried out in a solvent, the reaction temperature to be chosen is advantageously the reflux

temperature of the solvent. If, on the other hand, the reaction is carried out without a solvent, it is possible to heat up to 200°C, if appropriate.

Once the group Q has been assembled, for example by condensation,
 5 cyclization or cycloaddition reactions, the radicals R^1 to R^9 may be derivatized further, if desired, employing the extensive arsenal of methods of organochemical synthesis.

To assemble compounds of the formula (I), in which m is 1, compounds of
 10 the formula (I) in which m is 0 can be treated with an oxidizing agent, such as, for example, meta-chloroperoxybenzoic acid.

The compounds of the formula (I) (also referred to as "active compounds" hereinbelow) have good plant tolerance, favorable homotherm toxicity and
 15 advantageous properties with respect to aquatic organisms and are suitable for controlling animal pests, in particular insects, arachnids (Acarina), helminths and molluscs, especially preferably for controlling insects and arachnids which are encountered in agriculture, in animal husbandry, in forests, in the preservation of stored products and materials
 20 and in the hygiene sector. They are active against normally sensitive and resistant species and all or individual stages of development. The abovementioned pests include:

From the order of the Acarina, for example, *Acarus siro*, *Argas* spp., *Ornithodoros* spp., *Dermanyssus gallinae*, *Eriophyes ribis*, *Phyllocoptruta*
 25 *oleivora*, *Boophilus* spp., *Rhipicephalus* spp., *Amblyomma* spp., *Hyalomma* spp., *Ixodes* spp., *Psoroptes* spp., *Chorioptes* spp., *Sarcoptes* spp., *Tarsonemus* spp., *Bryobia praetiosa*, *Panonychus* spp., *Tetranychus* spp., *Eotetranychus* spp., *Oligonychus* spp. and *Eutetranychus* spp.

30 From the order of the Isopoda, for example, *Oniscus asellus*, *Armadium vulgare* and *Porcellio scaber*.

From the order of the Diplopoda, for example, *Blaniulus guttulatus*.

From the order of the Chilopoda, for example, *Geophilus carpophagus* and *Scutigera* spp.

From the order of the Symphyla, for example, *Scutigerella immaculata*.

From the order of the Thysanura, for example, *Lepisma saccharina*.

- 5 From the order of the Collembola, for example, *Onychiurus armatus*.

From the order of the Orthoptera, for example, *Blatta orientalis*, *Periplaneta americana*, *Leucophaea madeirae*, *Blatella germanica*, *Acheta domesticus*, *Gryllotalpa* spp., *Locusta migratoria migratorioides*, *Melanoplus differentialis* and *Schistocerca gregaria*.

10

From the order of the Isoptera, for example, *Reticulitermes* spp.

From the order of the Anoplura, for example, *Phylloera vastatrix*, *Pemphigus* spp., *Pediculus humanus corporis*, *Haematopinus* spp. and *Linognathus* spp.

15

From the order of the Mallophaga, for example, *Trichodectes* spp. and *Damalinea* spp.

From the order of the Thysanoptera, for example, *Hercinothrips femoralis*, *Thrips tabaci* and *Frankliniella* spp.

20

From the order of the Heteroptera, for example, *Eurygaster* spp., *Dysdercus intermedius*, *Piesma quadrata*, *Cimex lectularius*, *Rhodnius prolixus* and *Triatoma* spp.

- 25 From the order of the Homoptera, for example, *Aleurodes brassicae*, *Bemisia tabaci*, *Trialeurodes vaporariorum*, *Aphis* spp., *Brevicoryne brassicae*, *Cryptomyzus ribis*, *Doralis fabae*, *Doralis pomi*, *Eriosoma lanigerum*, *Hyalopterus arundinis*, *Macrosiphum avenae*, *Myzus* spp., *Phorodon humuli*, *Rhopalosiphum padi*, *Empoasca* spp., *Euscelus*
- 30 *bilobatus*, *Nephotettix cincticeps*, *Lecanium corni*, *Saissetia oleae*, *Laodelphax striatellus*, *Nilaparvata lugens*, *Aonidiella aurantii*, *Aspidiotus hederae*, *Pseudococcus* spp. and *Psylla* spp.

- From the order of the Lepidoptera, for example, *Pectinophora gossypiella*,
Bupalus piniarius, *Cheimatobia brumata*, *Lithocolletis blancardella*,
Hyponomeuta padella, *Plutella maculipennis*, *Malacosoma neustria*,
Euproctis chrysorrhoea, *Lymantria* spp., *Bucculatrix thurberiella*,
5 *Phyllocnistis citrella*, *Agrotis* spp., *Euxoa* spp., *Feltia* spp., *Earias insulana*,
Heliothis spp., *Laphygma exigua*, *Mamestra brassicae*, *Panolis flammea*,
Prodenia litura, *Spodoptera* spp., *Trichoplusia ni*, *Carpocapsa pomonella*,
Pieris spp., *Chilo* spp., *Pyrausta nubilalis*, *Ephestia kuehniella*, *Galleria*
mellonella, *Cacoecia podana*, *Capua reticulana*, *Choristoneura fumiferana*,
10 *Clysia ambiguella*, *Homona magnanima*, *Tortrix viridana*, *Cuaphalocrocis*
spp. and *Manduca* spp.

- From the order of the Coleoptera, for example, *Anobium punctatum*,
Rhizopertha dominica, *Bruchidius obtectus*, *Acanthoscelides obtectus*,
15 *Hylotrupes bajulus*, *Agelastica alni*, *Leptinotarsa decemlineata*, *Phaedon*
cochleariae, *Diabrotica* spp., *Psylloides chrysocephala*, *Epilachna*
varivestis, *Atomaria* spp., *Oryzaephilus surinamensis*, *Anthonomus* spp.,
Sitophilus spp., *Otiorrhynchus sulcatus*, *Cosmopolites sordidus*,
Ceuthorrhynchus assimilis, *Hypera postica*, *Dermestes* spp., *Trogoderma*,
20 *Anthrenus* spp., *Attagenus* spp., *Lyctus* spp., *Meligethes aeneus*, *Ptinus*
spp., *Niptus hololeucus*, *Gibbium psyllodes*, *Tribolium* spp., *Tenebrio*
molitor, *Agriotes* spp., *Conoderus* spp., *Melolontha melolontha*,
Amphimallon solstitialis, *Costelytra zealandica* and *Lissorhoptus* spp.

- 25 From the order of the Hymenoptera, for example, *Diprion* spp.,
Hoplocampa spp., *Lasius* spp., *Monomorium pharaonis* and *Vespa* spp.

- From the order of the Diptera, for example, *Aedes* spp., *Anopheles* spp.,
Culex spp., *Drosophila melanogaster*, *Musca* spp., *Fannia* spp., *Calliphora*
30 *erythrocephala*, *Lucilia* spp., *Chrysomyia* spp., *Cuterebra* spp.,
Gastrophilus spp., *Hypobosca* spp., *Stomoxys* spp., *Oestrus* spp.,
Hypoderma spp., *Tabanus* spp., *Tannia* spp., *Bibio hortulanus*, *Oscinella*
frit, *Phorbia* spp., *Pegomyia hyoscyami*, *Ceratitis capitata*, *Dacus oleae*
and *Tipula paludosa*.

From the order of the Siphonaptera, for example, *Xenopsylla cheopsis* and *Ceratophyllus* spp.

From the order of the Arachnida, for example, *Scorpio maurus* and

- 5 *Latrodectus mactans*.

From the class of helminths, for example, *Haemonchus*, *Trichostrongylus*, *Ostertagia*, *Cooperia*, *Chabertia*, *Strongyloides*, *Oesophagostomum*, *Hyostrogylus*, *Ancylostoma*, *Ascaris* and *Heterakis*, as well as *Fasciola*.

- From the class of Gastropoda, for example, *Deroceras* spp., *Arion* spp.,
10 *Lymnaea* spp., *Galba* spp., *Succinea* spp., *Biomphalaria* spp., *Bulinus* spp. and *Oncomelania* spp.

From the class of Bivalva, for example, *Dreissena* spp.

- The phytoparasitic nematodes which can be controlled according to the
15 invention include, for example, the root-parasitic soil nematodes, such as, for example, those of the genera *Meloidogyne* (root gall nematodes, such as *Meloidogyne incognita*, *Meloidogyne hapla* and *Meloidogyne javanica*), *Heterodera* and *Globodera* (cyst-forming nematodes, such as *Globodera rostochiensis*, *Globodera pallida* and *Heterodera trifolii*) and of the genera
20 *Radopholus* (such as *Radopholus similis*), *Pratylenchus* (such as *Pratylenchus neglectus*, *Pratylenchus penetrans* and *Pratylenchus curvatus*), *Tylenchulus* (such as *Tylenchulus semipenetrans*), *Tylenchorhynchus* (such as *Tylenchorhynchus dubius* and *Tylenchorhynchus claytoni*), *Rotylenchus* (such as *Rotylenchus robustus*),
25 *Heliocotylenchus* (such as *Heliocotylenchus multicinctus*), *Belonoaimus* (such as *Belonoaimus longicaudatus*), *Longidorus* (such as *Longidorus elongatus*), *Trichodorus* (such as *Trichodorus primitivus*) and *Xiphinema* (such as *Xiphinema index*).

- 30 The nematode genera *Ditylenchus* (stem parasites, such as *Ditylenchus dipsaci* and *Ditylenchus destructor*), *Aphelenchoides* (leaf nematodes, such as *Aphelenchoides ritzemabosi*) and *Anguina* (blossom nematodes, such as *Anguina tritici*) can furthermore be controlled with the compounds according to the invention.

The invention also relates to compositions, in particular insecticidal and acaricidal compositions, which comprise the compounds of the formula (I) in addition to suitable formulation auxiliaries.

5

The compositions according to the invention in general comprise the active compounds of the formula (I) to the extent of 1 to 95% by weight. They can be formulated in various ways, depending on how this is determined by the biological and/or chemico-physical parameters. Suitable formulation

10 possibilities are therefore:

Wettable powders (WP), emulsifiable concentrates (EC), aqueous solutions (SL), emulsions, sprayable solutions, oil- or water-based dispersions, suspension concentrates (SC), suspoemulsions (SE), dusting powders (DP), seed dressings, granules in the form of microgranules,
15 sprayed granules, absorption granules and adsorption granules, water-dispersible granules (WG), ULV formulations, microcapsules, waxes or baits.

These individual types of formulation are known in principle and are
20 described, for example, in: Winnacker-Küchler, "Chemische Technologie" [Chemical Technology], Volume 7, C. Hauser Verlag Munich, 4th Edition 1986; van Falkenberg, "Pesticides Formulations", Marcel Dekker N.Y., 2nd Edition 1972-73; K. Martens, "Spray Drying Handbook", 3rd Edition 1979, G. Goodwin Ltd. London.

25

The necessary formulation auxiliaries, i.e. carrier substances and surface-acting substances, such as inert materials, surfactants, solvents and further additives, are likewise known and are described, for example, in: Watkins, "Handbook of Insecticide Dust Diluents and Carriers",
30 2nd Edition, Darland Books, Caldwell N.J.; H. v. Olphen, "Introduction to Clay Colloid Chemistry", 2nd Edition, J. Wiley & Sons, N.Y.; Marsden, "Solvents Guide", 2nd Edition, Interscience, N.Y. 1950; McCutcheon's, "Detergents and Emulsifiers Annual", MC Publ. Corp., Ridgewood N.J.; Sisley and Wood, "Encyclopedia of Surface Active Agents", Chem. Publ.

Co. Inc., N.Y. 1964; Schönfeldt, "Grenzflächenaktive Äthylenoxidaddukte" [Surface-active ethylene oxide adducts], Wiss. Verlagsgesell., Stuttgart 1967; Winnacker-Küchler, "Chemische Technologie" [Chemical Technology], Volume 7, C. Hauser Verlag Munich, 4th Edition 1986.

5

Combinations with other substances having a pesticidal action, fertilizers and/or growth regulators can be prepared on the basis of these formulations, for example in the form of a ready-to-use formulation or as a tank mix. Wettable powders are preparations which are uniformly

10 dispersible in water and which, alongside the active compound, and in addition to a diluent or inert substance, also comprise wetting agents, for example polyethoxylated alkylphenols, polyethoxylated fatty alcohols or alkyl- or alkylphenol-sulfonates, and dispersing agents, for example sodium ligninsulfonate or sodium 2,2'-dinaphthylmethane-6,6'-disulfonate.

15 Emulsifiable concentrates are prepared by dissolving the active compound in an organic solvent, for example butanol, cyclohexanone, dimethylformamide, xylene or also higher-boiling aromatics or hydrocarbons, with the addition of one or more emulsifiers. Emulsifiers which can be used are, for example: calcium alkylarylsulfonates, such as
20 Ca dodecylbenzenesulfonate, or nonionic emulsifiers, such as fatty acid polyglycol esters, alkylaryl polyglycol ethers, fatty alcohol polyglycol ethers, propylene oxide/ethylene oxide condensation products, alkyl polyethers, sorbitan fatty acid esters, polyoxyethylene sorbitan fatty acid esters or polyoxyethylene sorbitol esters.

25

Dusting powders are obtained by grinding the active compound with finely divided solid substances, for example talc, naturally occurring clays, such as kaolin, bentonite and pyrophyllite, or diatomaceous earth. Granules can be prepared either by spraying the active compound onto granular inert

30 material capable of adsorption or by applying active compound concentrates to the surface of carrier substances, such as sand, kaolinites or granular inert material, by means of adhesives, for example polyvinyl alcohol, sodium polyacrylate or mineral oils. Suitable active compounds

can also be granulated in the manner customary for the preparation of fertilizer granules - if desired as a mixture with fertilizers.

5 In wettable powders, the active compound concentration is generally about 10 to 90% by weight, the remainder to make up 100% by weight comprising customary formulation constituents. In emulsifiable concentrates, the active compound concentration can be about 5 to 80% by weight. Dust-like formulations usually comprise 5 to 20% by weight of active compound, and sprayable solutions about 2 to 20% by weight. In 10 granules, the content of active compound partly depends on whether the active compound is present in liquid or solid form and what granulating auxiliaries, fillers and the like are used.

15 In addition, the active compound formulations mentioned comprise, if appropriate, the particular customary tackifiers, wetting agents, dispersing agents, emulsifiers, penetration agents, solvents, fillers or carriers.

20 For use, the concentrates in the commercially available form are diluted in the customary manner, if appropriate, for example by means of water in the case of wettable powders, emulsifiable concentrates, dispersions and in some cases also microgranules. Dust-like and granular formulations as well as sprayable solutions are usually not diluted further with additional inert substances before use.

25 The required amount applied varies with external conditions, such as temperature, humidity and the like. It can vary within wide limits, for example between 0.0005 and 10.0 kg/ha or more of active substance, but is preferably between 0.001 and 5 kg/ha.

30 The active compounds according to the invention can be present in their commercially available formulations and in the use forms prepared from these formulations as mixtures with other active compounds, such as insecticides, attractants, sterilizing agents, acaricides, nematocides, fungicides, growth-regulating substances or herbicides.

The pest control agents include, for example, phosphoric acid esters, carbamates, carboxylic acid esters, formamidines, tin compounds, substances produced by microorganisms and the like.

5 Preferred partners for the mixtures are

1. from the group of phosphorus compounds
 acephate, azamethiphos, azinphos-ethyl-, azinphosmethyl, bromophos, bromophos-ethyl, chlorfenvinphos, chlormephos, chlorpyrifos, chlorpyrifos-
 10 methyl, demeton, demeton-S-methyl, demeton-S-methyl sulfone, dialifos, diazinon, dichlorvos, dicrotophos, O,O-1,2,2,2-tetrachloroethyl phosphorthioate (SD 208 304), dimethoate, disulfoton, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitriothion, fensulfothion, fenthion, fonofos, formothion, heptenophos, isozophos, isothioate,
 15 isoxathion, malathion, methacrifos, methamidophos, methidathion, salithion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, parathion, parathion-methyl, phenthoate, phorate, phosalone, phosfolan, phosmet, phosphamidon, phoxim, pirimiphos, primiphos-ethyl, pirimiphos-methyl, profenofos, propaphos, proetamphos, prothiofos,
 20 pyraclofos, pyridapenthion, quinalphos, sulprofos, temephos, terbufos, tetrachlorvinphos, thiometon, triazophos, trichlorphon, vamidothion;
2. from the group of carbamates
 aldicarb, 2-sec-butylphenyl methylcarbamate (BPMC), carbaryl,
 25 carbofuran, carbosulfan, cloethocarb, benfuracarb, ethiofencarb, furathiocarb, isoprocarb, methomyl, 5-methyl-m-cumenyl butyryl(methyl)carbamate, oxamyl, pirimicarb, propoxur, thiodicarb, thiofanox, ethyl 4,6,9-triaza-4-benzyl-6, 10-dimethyl-8-oxa-7-oxo-5,11-dithia-9-dodecenoate (OK 135), 1-methylthio(ethylideneamino) N-methyl-
 30 N-(morpholiniothio)carbamate (UC 51717);
3. from the group of carboxylic acid esters
 allethrin, alphametrin, 5-benzyl-3-furylmethyl (E)-(1R)-cis, 2,2-di-methyl-3-(2-oxothiolan-3-ylidenemethyl)cyclopropanecarboxylate, bioallethrin,

- bioallethrin ((S)-cyclopentyl isomer), bioresmethrin, biphenate, (RS)-1-cyano-1-(6-phenoxy-2-pyridyl)methyl (1RS)-trans-3-(4-tert-butylphenyl)-2,2-dimethylcyclopropanecarboxylate (NCI 85193), cycloprothrin, cyhalothrin, cythithrin, cypermethrin, cyphenothrin, deltamethrin, empenethrin,
- 5 esfenvalerate, fenfluthrin, fenpropathrin, fenvalerate, flucythrinate, flumethrin, fluvalinate (D isomer), permethrin, pheothrin ((R) isomer), d-pralethrin, pyrethrins (naturally occurring products), resmethrin, tefluthrin, tetramethrin and tralomethrin;
- 10 4. from the group of amidines
amitraz, chlordimeform;
5. from the group of tin compounds
cyhexatin, fenbutatin oxide;
- 15 6. others
abamectin, *Bacillus thuringiensis*, bensultap, binapacryl, bromopropylate, buprofezin, camphechlor, cartap, chlorobenzilate, chlorfluazuron, 2-(4-chlorophenyl)-4,5-diphenylthiophene (UBI-T 930), chlorfentezine, 2-
- 20 naphthylmethyl cyclopropanecarboxylate (Ro 12-0470), cyromazin, N-(3,5-dichloro-4-(1,1,2,3,3,3-hexafluoro-1-propyloxy)phenyl)carbamoyl)-2-chlorobenzocarboximide acid ethyl ester, dicofol, N-(N-(3,5-di-chloro-4-(1,1,2,2-tetrafluoroethoxy)phenylamino)carbonyl)-2,6-difluorobenzamide (XRD 473), diflubenzuron, N-(2,3-dihydro-3-methyl-1,3-thiazol-2-ylidene)-
- 25 2,4-xylylidene, dinobuton, dinocap, endosulfan, ethofenprox, (4-ethoxyphenyl)(dimethyl)(3-(3-phenoxyphenyl)propyl)silane, (4-ethoxyphenyl)(3-(4-fluoro-3-phenoxyphenyl)propyl)dimethylsilane, fenoxycarb, 2-fluoro-5-(4-(4-ethoxyphenyl)-4-methyl-1-pentyl)diphenyl ether (MTI 800), granulosin and nuclear polyhedrosis viruses, fenthion, flubenzimine, flucycloxuron, flufenoxuron, gamma-HCH, hexythiazox,
- 30 hydramethylnon (AC 217300), ivermectin, 2-nitromethyl-4,5-dihydro-6H-thiazine (DS 52618), 2-nitromethyl-3,4-dihydrothiazole (SD 35651), 2-nitromethylene-1,2-thiazinan-3-ylcarbamide (WL 108477),

propargite, teflubenzuron, tetradifon, tetrasul, thiocyclam, trifumuron, imidacloprid.

5 The abovementioned combination partners are known active compounds, and most of them are described in Ch. R. Worthing, S.B. Walker, The Pesticide Manual, 7th Edition (1983), British Crop Protection Council.

10 The active compound content of the use forms prepared from the commercially available formulations can be from 0.00000001 to 95% by weight of active compound, preferably between 0.00001 and 1% by weight.

The active compounds are used in a customary manner appropriate for the use forms.

15 The active compounds according to the invention are also suitable for controlling endo- and ectoparasites in the veterinary medicine field and in the field of animal husbandry. The active compounds according to the invention are used here in a known manner, such as by oral use in the form of, for example, tablets, capsules, potions or granules, by means of
20 dermal use in the form of, for example, dipping, spraying, pouring-on, spotting-on and dusting, and by parenteral use in the form of, for example, injection.

25 The novel compounds of the formula (I) can accordingly also particularly advantageously be used in livestock husbandry (for example cattle, sheep, pigs and poultry, such as chickens, geese and the like). In a preferred embodiment of the invention, the compounds are administered orally to the animals, if appropriate in suitable formulations (cf. above) and if appropriate with the drinking water or feed. Since excretion in the faeces
30 takes place in an active manner, the development of insects in the faeces of the animals can be prevented very easily in this way. The dosages and formulations suitable in each case depend in particular on the species and the development stage of the stock animals and also on the pressure of infestation, and can easily be determined and specified by the customary

methods. The novel compounds can be employed in cattle, for example, in dosages of 0.01 to 1 mg/kg of body weight.

In addition to the application methods mentioned hereinabove, the active compounds of the formula I according to the invention also have excellent systemic action. The active compounds can therefore also be introduced into the plants via below-ground and above-ground parts of plants (root, stem, leaf), when the active compounds are applied in liquid or solid form to the immediate surroundings of the plants (for example granules in soil application, application in flooded rice fields).

Furthermore, the active compounds according to the invention are particularly useful for treating vegetative and generative propagation stock, such as, for example, seed of, for example, cereals, vegetables, cotton, rice, sugar beet and other crops and ornamentals, of bulbs, cuttings and tubers of other vegetatively propagated crops and ornamentals. To this end, treatment can be carried out prior to sowing or planting (for example by special seed coating techniques, by seed dressings in liquid or solid form or by seed box treatment), during sowing or planting or after sowing or planting by special application techniques (for example seed row treatment). Depending on the application, the amount of active compound applied can vary within a relatively wide range. In general, the application rates are between 1 g and 10 kg of active compound per hectare of soil area.

25

The examples below serve to illustrate the invention.

A. Formulation examples

- 5 a) A dusting powder is obtained by mixing 10 parts by weight of active compound and 90 parts by weight of talc, as the inert substance, and comminuting the mixture in an impact mill.
- 10 b) A wettable powder which is readily dispersible in water is obtained by mixing 25 parts by weight of active compound, 65 parts by weight of kaolin-containing quartz, as the inert substance, 10 parts by weight of potassium ligninsulfonate and 1 part by weight of sodium oleoylmethyltauride, as wetting and dispersing agent and grinding the mixture in a pinned disk mill.
- 15 c) A dispersion concentrate which is readily dispersible in water is prepared by mixing 40 parts by weight of active compound with 7 parts by weight of a sulfosuccinic monoester, 2 parts by weight of a sodium ligninsulfonate and 51 parts by weight of water and grinding the mixture to a fineness of below 5 microns in a grinding bead mill.
- 20 d) An emulsifiable concentrate can be prepared from 15 parts by weight of active compound, 75 parts by weight of cyclohexane, as the solvent, and 10 parts by weight of ethoxylated nonylphenol (10 EO), as the emulsifier.
- 25 e) Granules can be prepared from 2 to 15 parts by weight of active compound and an inert granule carrier material, such as attapulgite, pumice granules and/or quartz sand. A suspension of the wettable powder from Example b) having a solids content of 30% is expediently used, and this is sprayed onto the surface of attapulgite granules and the
- 30 components are dried and mixed intimately. The weight content of the wettable powder is approximately 5% and that of the inert carrier material is approximately 95% of the finished granules.

B. Chemical examples**Example No. 1**

3-Isopropyl-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole (Table 1, No. 81)

- 5 2 g of methyl 4-trifluoromethylnicotinate and 1.56 g of isobutyramide oxime were initially charged in 15 ml of ethanol and cooled to 0°C. 10 ml of a 1.2 molar sodium ethoxide solution were added dropwise to this solution. The mixture was allowed to warm to room temperature over a period of two hours and stirring was then continued at this temperature until the reaction, according to TLC, had ended.

The reaction mixture was concentrated and the residue was taken up in saturated ammonium chloride solution and extracted with diethyl ether. Chromatographic purification of the crude product gave the desired compound as a yellowish oil.

- 15 ¹H-NMR (CDCl₃, 300 MHz): δ = 1.41 (d, J = 6.9 Hz, 6H), 3.22 (m, 1H), 7.78 (d, J = 5 Hz, 1H), 9.02 (d, J = 5 Hz, 1H), 9.34 (s, 1H) ppm.

Example No. 2

- 20 *3-Isopropyl-5-(4-trifluoromethyl-5-pyrimidyl)-1,2,4-oxadiazole* (Table 1, No. 189)

- 25 2 g of ethyl 4-trifluoromethylpyrimidine-5-carboxylate and 1.56 g of isobutyramide oxime were initially charged in 15 ml of ethanol and cooled to 0°C. 10 ml of a 1.2 molar sodium ethoxide solution were added dropwise to this solution. The mixture was allowed to warm to room temperature over a period of one hour and then heated under reflux until the reaction, according to TLC, had ended. The reaction mixture was concentrated and the residue was taken up in saturated ammonium chloride solution and extracted with diethyl ether. Chromatographic purification of the crude product gave the desired compound as a yellowish oil.

¹H-NMR (CDCl₃, 300 MHz): d = 1.43 (d, J = 7 Hz, 6H), 3.22 (hept., J = 7 Hz, 1H), 9.52 (s, 1H), 9.58 (s, 1H) ppm.

Example No. 3

5 *2-Methyl-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole (Table 3, No. 549)*

500 mg of 4-trifluoromethylnicotinic hydrazide were heated under reflux in 3.5 ml of triethyl orthoacetate for 2 hours. The reaction mixture was subsequently concentrated and the residue was carefully admixed with 2 ml of phosphorus oxychloride. The mixture was stirred at room

10 temperature for 1 hour and then poured on ice and extracted with ethyl acetate. Chromatographic purification of the crude product obtained after drying and concentrating gave the desired compound as a yellowish oil.

¹H-NMR (CDCl₃, 300 MHz): d = 2.67 (s, 3H), 7.75 (d, J = 5 Hz, 1H), 8.99 (d, J = 5 Hz, 1H), 9.34 (s, 1H) ppm.

15

Example No. 4

4-(Ethoxycarbonylmethyl)-2-(4-trifluoromethyl-3-pyridyl)thiazole (Table 4, No. 688)

20

500 mg of 4-trifluoromethylpyridine-3-thiocarboxamide and 440 mg of ethyl 4-chloroacetate were dissolved in 5 ml of dimethylformamide and heated at 100°C for 4 hours. After cooling, the reaction mixture was poured onto ice-water and extracted with diethyl ether. The diethyl ether phase was dried

25 (MgSO₄), filtered and concentrated and the residue was purified by chromatography. This gave the desired product in pure form as a colorless oil.

¹H-NMR (CDCl₃, 300 MHz): d = 1.28 (t, J = 7.5 Hz, 3H), 3.92 (s, 2H), 4.22 (q, J = 7.5 Hz, 2H), 7.43 (s, 1H), 7.68 (d, J = 5 Hz, 1H), 8.86 (d, J = 5 Hz, 1H), 8.97 (s, 1H) ppm.

30

Example No. 5:*4-Ethyl-2-(4-trifluoromethyl-3-pyridyl)oxazole (Table 4, No. 762)*

5

2.6 g of 4-trifluoromethylnicotinic acid were admixed with 20 ml of thionyl chloride and heated at reflux temperature for 1 hour. After cooling, excess thionyl chloride was distilled off and the acyl chloride which remained as a pale yellow oil was taken up in 30 ml of dichloromethane. This solution was subsequently added dropwise to a solution of 2.4 g of 2-amino-1-butanol and 2.75 g of triethylamine in 30 ml of dichloromethane cooled in an ice bath. After the addition had ended, stirring was continued at room temperature for approximately 2 hours. The mixture was poured into ammonium chloride solution and extracted with ethyl acetate. The crude N-(1-hydroxy-2-butyl)-4-trifluoromethylnicotinamide (2.3 g) obtained after drying and concentrating the ethyl acetate phase was dissolved at room temperature in 100 ml dichloromethane and mixed with 4.6 g of periodinan (Dess-Martin reagent). After the reaction had ended, according to TLC, the reaction mixture was concentrated and purified by column chromatography. The resulting 2-(trifluoromethylpyridin-3-amido)butanal (1.5 g) was dissolved in 30 ml of dimethylformamide, admixed with 2.72 g of phosphorus oxychloride and heated at 90°C for 15 minutes. The solution was then poured onto ice and extracted with diethyl ether. Drying and concentration of the diethyl ether phase and chromatographic purification of the residue gave the product as a brownish oil.

25

¹H-NMR (CDCl₃, 300 MHz): δ = 1.3 (t, J = 7.4 Hz, 3H), 2.66 (qd, J = 7.4 Hz, J < 1 Hz, 2H), 7.58 (t, J < 1 Hz, 1H), 7.65 (d, J = 5 Hz, 1H), 8.83 (d, J = 5 Hz, 1H), 9.33 (d, J = 5 Hz, 1H) ppm.

30

Example No. 6*4-Ethyl-2-(4-trifluoromethyl-3-pyridyl)-4,5-dihydrooxazole (Table 5, No 876)*

5 1 g of 4-trifluoromethylnicotinic acid was admixed with 8 ml of thionyl chloride and heated at reflux temperature for 1 hour. After cooling, excess thionyl chloride was distilled off and the acyl chloride which remained as a pale yellow oil was taken up in 10 ml of dichloromethane. This solution was subsequently added dropwise to a solution of 930 mg of 2-amino-1-butanol
10 and 1.06 g of triethylamine in 10 ml of dichloromethane cooled in an ice bath. After the addition had ended, stirring was continued for approximately 2 hours at room temperature. The mixture was poured into an ammonium chloride solution and extracted with ethyl acetate. The crude N-(1-hydroxy-2-butyl)-4-trifluoromethylnicotinamide (1.03 g) obtained after drying and
15 concentration of the ethyl acetate phase was dissolved at room temperature in 6 ml of tetrahydrofuran and admixed with 1.09 g of N-[(triethylammonio)sulfonyl]-methylcarbamate (Burgess' reagent). The mixture was stirred at 60°C for 3 hours. After cooling, the batch was concentrated and the residue was taken up in water and extracted with
20 ethyl acetate. Chromatographic purification of the crude product gave the product as a colorless oil.

¹H-NMR (CDCl₃, 200 MHz): δ = 1.03 (t, J = 7.6 Hz, 3H), 1.72 (m, 2H), 4.15 (t, J = 7.5 Hz, 1H), 4.32 (m, 1H), 4.58 (t, J = 7.5 Hz, 1H), 7.6 (d, J = 5 Hz, 1H), 8.87 (d, J = 5 Hz, 1H), 9.06 (s, 1H) ppm.

25

Example No. 7*2-(3-Thienylmethyl)-5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole (Table 3, No. 572)*

30

880 mg of thiophene-3-acetic hydrazide were added to a solution of 960 mg of 4-trifluoromethylpyridine-3-carboxylic acid in 5 ml of phosphorus oxychloride, and the mixture was heated at reflux for 2 hours. The reaction

mixture was subsequently added dropwise to ice-water, made neutral using concentrated ammonia solution and extracted with ethyl acetate. Drying (Na_2SO_4), concentration and chromatographic purification gave 624 mg of the desired product as a slightly brown oil.

- 5 $^1\text{H-NMR}$ (CDCl_3 , 200 MHz): δ = 4.38 (s, 2H), 7.1 (d, J = 5 Hz, 1H), 7.23 (s, 1H), 7.37 (dd, J = 5 Hz, J = 3 Hz, 1H), 7.75 (d, J = 6 Hz, 1H), 8.98 (d, J = 6 Hz, 1H), 9.36 (s, 1H) ppm.

Example No. 8

10

5-Methyl-3-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole (Table 6, No. 947)

- 15 A mixture of 290 mg of ethylacetimidate hydrochloride and 100 mg of sodium hydroxide in 2 ml of ethanol was filtered and added to 500 mg of 4-trifluoromethyl-3-pyridinecarbohydrazide, and the mixture was heated under reflux for 3 hours. The reaction mixture was concentrated and the residue was suspended in xylene and refluxed for 4 hours. For work-up, the batch was diluted with ethyl acetate and washed with water.

Chromatographic purification gave the pure product as a colorless solid.

- 20 $^1\text{H-NMR}$ (CDCl_3 , 300 MHz): δ = 2.58 (s, 3H), 7.64 (d, J = 5 Hz, 1H), 8.85 (d, J = 5 Hz, 1H), 9.19 (s, 1H) ppm.

Example 9

- 25 *3-(N-Isopropylcarbamoylmethyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole*

Step 1: Tert-butyl 3-amino-3-(4-trifluoromethyl-3-pyridinecarbonyloxy-imino)propionate

30

30 g of 4-trifluoromethyl-3-pyridinecarboxylic acid is initially charged in 150 ml of dry THF and, a little at a time, admixed with 25.3 g of carbonyl-

diimidazole. The mixture is stirred at room temperature for 30 min. 27.2 g of tert-butoxycarbonylacetamide oxime dissolved in 150 ml of THF are then added dropwise. The mixture is stirred overnight, the solvent is evaporated and the residue is taken up in ethyl acetate, washed three times with 1 M sulfuric acid and once with saturated sodium bicarbonate solution. Concentration of the ethyl acetate phase gives 28 g of the product as a pale yellow solid.

¹H-NMR (CDCl₃, 300MHz): d = 1.5 (s, 9H), 3.3 (s, 2H), 5.55 (br.s, 2H), 7.83 (d, J = 5 Hz, 1H), 8.97 (d, J = 5Hz, 1H), 9.13 (s, 1H) ppm.

Step 2: 3-(Tert-butoxycarbonylmethyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole

28 g of tert-butyl 3-amino-3-(4-trifluoromethyl-3-pyridinecarbonyloxy-imino)propionate are dissolved in 380 ml of toluene and heated under reflux for 17 hours. Concentration and chromatographic purification of the residue over silica gel gives 14.4 g of the product as a pale brown oil.

¹H-NMR (CDCl₃, 300MHz): d = 1.5 (s, 9H), 3.88 (s, 2H), 7.79 (d, J = 5 Hz, 1H), 9.02 (d, J = 5Hz, 1H), 9.33 (s, 1H) ppm.

Step 3: 3-(Hydroxycarbonylmethyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole

12.4 g of 3-(tert-butoxycarbonylmethyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole are dissolved in 110 ml of dichloromethane and admixed with 57 ml of trifluoroacetic acid. The reaction mixture is stirred at room temperature for 1.5 hours and subsequently concentrated under reduced pressure. The residue is repeatedly taken up in dichloromethane and reconcentrated to remove any remaining trifluoroacetic acid. The mixture is

finally triturated with diethyl ether, giving 8.1 g of the product as a white solid.

5 **Step 4: 3-(N-Isopropylcarbamoylmethyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole**

1 g of the product of the previous step are dissolved in 10 ml of THF and is mixed with 0.59 g of carbonyldiimidazole. The mixture is stirred at room temperature for 10 minutes, 0.22 g of isopropylamine are added dropwise
 10 and the mixture is allowed to react for a further 1.5 hours at room temperature with stirring. The reaction mixture is subsequently concentrated and the residue is taken up in ethyl acetate and washed three times with 1 M sulfuric acid and once with saturated sodium bicarbonate solution. The solid residue obtained after drying and
 15 concentrating the ethyl acetate phase is recrystallized from tert-butyl methyl ether, giving 0.46 g of the pure product as a pale yellow solid.

¹H-NMR (CDCl₃, 300MHz): d = 1.20 (d, J = 7.6 Hz, 6H), 3.82 (s, 2H), 4.12 (m, 1H), 6.50 (br.s, 1H), 7.81 (d, J = 5 Hz, 1H), 9.02 (d, J = 5Hz, 1H), 9.37
 20 (s, 1H) ppm.

Example No. 10

25 **3-(N,N-Dimethylaminocarbamoyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole**

Step 1: Ethyl 2-amino-2-(4-trifluoromethyl-3-pyridinecarbonyloxyimino)acetate
 30

17.3 g of carbonyldiimidazole are initially charged in 200 ml of 1,4-dioxane and, a little at a time, admixed with 20 g of 4-trifluoromethyl-3-pyridinecarboxylic acid. The mixture is stirred at room temperature for 1 h

and subsequently heated to 45°C for 2 h. After cooling to 30°C, 14.5 g of ethoxycarbonylformamide oxime are added and the mixture is stirred at 45°C for 3 h. The precipitated solid is filtered off with suction and the filtrate is concentrated to 50 ml and, together with the solid, added to 250 ml of ice-water. The solid is filtered off with suction and dried at 50°C under reduced pressure. This gives 28.7 g of the product as a white solid of mp. 172-174°C.

Step 2: 3-Ethoxycarbonyl-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole

20 g of ethyl 2-amino-2-(4-trifluoromethyl-3-pyridinecarbonyloxyimino)-acetate are dissolved in 200 ml of a mixture of xylene and toluene and admixed with 5 g of Amberlyst 15. The mixture is boiled at 125-130°C for 6 h using a Dean-Stark apparatus. After the reaction has ended, the mixture is cooled and admixed with a small amount of diethyl ether. The mixture is filtered with suction through a glass filter frit, and the solution is then concentrated. This gives 15.8 g of the product as a yellow oil.

Step 3: 5-(4-Trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole-3-carboxylic acid

15.8 g of 3-ethoxycarbonyl-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole are initially charged in 13 ml of methanol, and, with ice-cooling at 0°C, a solution of 2.8 g of lithium hydroxide in 50 ml of water is added dropwise. The mixture is stirred at room temperature for 2 h, 20 ml of ice-water are added and the mixture is extracted with 200 ml of diethyl ether. The aqueous phase is adjusted to pH=2 using dil. HCl, and the precipitated product is filtered off with suction. After drying, 13.8 g of 5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole-3-carboxylic acid are obtained as a white solid of mp. 157-159°C.

Step 4: N,N-Dimethyl-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole-3-carboxamide

5.8 g of carbonyldiimidazole are initially charged in 90 ml of tetrahydrofuran and, a little at a time, admixed with 9 g of 5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole-3-carboxylic acid. The mixture is stirred at room temperature for 15 min and then heated at 50°C for 2 h. After cooling to room temperature, 2.3 g of dimethylamine are introduced in a very gentle gas stream over a period of 2 h. After a reaction time of 12 h, the mixture is concentrated and taken up in 200 ml of diethyl ether. The mixture is washed with ice-cold half conc. hydrochloric acid solution, washed neutral with sat. sodium bicarbonate sol., dried over magnesium sulfate and concentrated under reduced pressure. This gives a slightly yellow oil which [lacuna] after a number of days to a solid of mp. 52-54°C.


In a similar manner, it is possible to prepare the compounds shown in Tables 1 to 6 below.

The abbreviations used denote

Ph: phenyl

THP: 2-tetrahydropyranyl

Table 1



No.	X	Y	m	W	R ¹	m.p. [°C]
1	N	CCl ₃	0	O	CH ₃	
2	N	CCl ₃	0	O	CH ₂ CH ₃	
3	N	CCl ₃	0	O	COOCH ₂ CH ₃	
4	CH	CCl ₃	0	O	CH ₃	
5	CH	CCl ₃	0	O	COOCH ₂ CH ₃	
6	N	(CF ₂) ₃ CHF ₂	0	O	CH ₃	
7	N	(CF ₂) ₃ CHF ₂	0	O	COOCH ₂ CH ₃	
8	CH	(CF ₂) ₃ CHF ₂	0	O	CH ₃	
9	CH	(CF ₂) ₃ CHF ₂	0	O	COOCH ₂ CH ₃	
10	N	(CF ₂) ₃ CHF ₂	0	S	CH ₂ COOC(CH ₃) ₃	
11	N	(CF ₂) ₃ CHF ₂	0	S	CH ₂ CONHCH ₃	
12	CH	(CF ₂) ₃ CHF ₂	0	S	(CH ₂) ₂ CH ₃	
13	CH	(CF ₂) ₃ CHF ₂	0	S	COOCH ₂ CH ₃	
14	N	(CF ₂) ₂ CHF ₂	0	O	CH ₂ CH ₃	
15	N	(CF ₂) ₂ CHF ₂	0	O	COOCH ₂ CH ₃	
16	N	(CF ₂) ₂ CHF ₂	0	O	OH	
17	N	(CF ₂) ₂ CHF ₂	0	O	OCH ₃	
18	CH	(CF ₂) ₂ CHF ₂	0	O	CH ₃	
19	CH	(CF ₂) ₂ CHF ₂	0	O	COOCH ₂ CH ₃	
20	CH	(CF ₂) ₂ CHF ₂	0	O	OH	

No.	X	Y	m	W	R ¹	m.p. [°C]
21	CH	(CF ₂) ₂ CHF ₂	0	O	NHCH ₃	
22	N	CF ₂ CF ₃	0	O	CH ₃	
23	N	CF ₂ CF ₃	0	O	CH ₂ CH ₃	
24	N	CF ₂ CF ₃	0	O	(CH ₂) ₂ CH ₃	
25	N	CF ₂ CF ₃	0	O	CH(CH ₃) ₂	
26	N	CF ₂ CF ₃	0	O	Cyclo-C ₆ H ₁₁	
27	N	CF ₂ CF ₃	0	O	CH ₂ C=CH ₂	
28	N	CF ₂ CF ₃	0	O	CH ₂ C≡CH	
29	N	CF ₂ CF ₃	0	O	CH ₂ CH ₂ C≡CH	
30	N	CF ₂ CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₃	
31	N	CF ₂ CF ₃	0	O	(CH ₂) ₄ C≡CH	
32	N	CF ₂ CF ₃	0	O	CHFCF ₃	
33	N	CF ₂ CF ₃	0	O	COOCH ₂ CH ₃	
34	N	CF ₂ CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	
35	N	CF ₂ CF ₃	0	O	CH ₂ CONHCH ₃	
36	N	CF ₂ CF ₃	0	O	NH ₂	
37	N	CF ₂ CF ₃	0	O	NHCH ₂ CH ₃	
38	CH	CF ₂ CF ₃	0	O	CH ₃	
39	CH	CF ₂ CF ₃	0	O	CH ₂ CH ₃	
40	CH	CF ₂ CF ₃	0	O	(CH ₂) ₂ CH ₃	
41	CH	CF ₂ CF ₃	0	O	CH(CH ₃) ₂	
42	CH	CF ₂ CF ₃	0	O	Cyclo-C ₆ H ₁₁	
43	CH	CF ₂ CF ₃	0	O	CH ₂ C=CH ₂	
44	CH	CF ₂ CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	
45	CH	CF ₂ CF ₃	0	O	NH ₂	
46	CH	CF ₂ CF ₃	0	O	NHCOCH ₃	
47	CH	CF ₂ CF ₃	0	O	NHCOCH ₂ CH ₃	

No.	X	Y	m	W	R ¹	m.p. [°C]
48	N	CF ₂ CF ₃	0	S	CH ₃	
49	N	CF ₂ CF ₃	0	S	CH ₂ CH ₃	
50	N	CF ₂ CF ₃	0	S	(CH ₂) ₂ CH ₃	
51	N	CF ₂ Cl	0	O	CH ₃	
52	N	CF ₂ Cl	0	O	CH ₂ CH ₃	
53	N	CF ₂ Cl	0	O	(CH ₂) ₂ CH ₃	
54	N	CF ₂ Cl	0	O	CH(CH ₃) ₂	
55	N	CF ₂ Cl	0	O	CH ₂ COOC(CH ₃) ₃	
56	N	CF ₂ Cl	0	O	CH ₂ CONHCH ₃	
57	N	CF ₂ Cl	0	O	OH	
58	N	CF ₂ Cl	0	O	OCH ₃	
59	N	CF ₂ Cl	0	O	OCH ₂ CH ₃	
60	N	CF ₂ Cl	0	O	NHCH ₃	
61	CH	CF ₂ Cl	0	O	CH ₃	
62	CH	CF ₂ Cl	0	O	CH ₂ CH ₃	
63	CH	CF ₂ Cl	0	O	(CH ₂) ₂ CH ₃	
64	CH	CF ₂ Cl	0	O	CH(CH ₃) ₂	
65	CH	CF ₂ Cl	0	O	CH ₂ COOC(CH ₃) ₃	
66	CH	CF ₂ Cl	0	O	CH ₂ CONHCH ₃	
67	CH	CF ₂ Cl	0	O	OH	
68	CH	CF ₂ Cl	0	O	OCH ₃	
69	CH	CF ₂ Cl	0	O	OCH ₂ CH ₃	
70	CH	CF ₂ Cl	0	O	NHCH ₃	
71	CH	CF ₂ Cl	0	O	Cyclo-C ₆ H ₁₁	
72	CH	CF ₂ Cl	0	O	CH ₂ C=CH ₂	
73	CH	CF ₂ Cl	0	O	COOCH ₂ CH ₃	
74	CH	CF ₂ Cl	0	O	CH ₂ COOC(CH ₃) ₃	

No.	X	Y	m	W	R ¹	m.p. [°C]
75	CH	CF ₂ Cl	0	O	CH ₂ CONHCH ₃	
76	CH	CF ₂ Cl	0	O	OCH ₃	
77	CH	CF ₂ Cl	0	O	NHCH ₃	
78	CH	CF ₃	0	O	CH ₃	oil
79	CH	CF ₃	0	O	CH ₂ CH ₃	oil
80	CH	CF ₃	0	O	(CH ₂) ₂ CH ₃	oil
81	CH	CF ₃	0	O	CH(CH ₃) ₂	oil
82	CH	CF ₃	0	O	Cyclo-C ₃ H ₅	oil
83	CH	CF ₃	0	O	(CH ₂) ₃ CH ₃	oil
84	CH	CF ₃	0	O	CH(CH ₃)CH ₂ CH ₃	oil
85	CH	CF ₃	0	O	CH ₂ CH(CH ₃) ₂	oil
86	CH	CF ₃	0	O	C(CH ₃) ₃	oil
87	CH	CF ₃	0	O	Cyclo-C ₄ H ₇	
88	CH	CF ₃	0	O	(CH ₂) ₄ CH ₃	oil
89	CH	CF ₃	0	O	CH(CH ₃)(CH ₂) ₂ CH ₃	
90	CH	CF ₃	0	O	(CH ₂) ₂ CH(CH ₃) ₂	
91	CH	CF ₃	0	O	CH ₂ C(CH ₃) ₃	
92	CH	CF ₃	0	O	Cyclo-C ₅ H ₉	oil
93	CH	CF ₃	0	O	(CH ₂) ₅ CH ₃	
94	CH	CF ₃	0	O	C(CH ₂ CH ₃) ₂ CH ₃	
95	CH	CF ₃	0	O	Cyclo-C ₆ H ₁₁	
96	CH	CF ₃	0	O	(CH ₂) ₆ CH ₃	
97	CH	CF ₃	0	O	CH(CH ₃)(CH ₂) ₄ CH ₃	
98	CH	CF ₃	0	O	Cyclo-C ₇ H ₁₃	
99	CH	CF ₃	0	O	CH ₂ -cyclo-C ₆ H ₁₁	
100	CH	CF ₃	0	O	2-Norbornyl	
101	CH	CF ₃	0	O	(CH ₂) ₇ CH ₃	

No.	X	Y	m	W	R ¹	m.p. [°C]
102	CH	CF ₃	0	O	CH(CH ₂ CH ₃)(CH ₂) ₅ CH ₃	
103	CH	CF ₃	0	O	(CH ₂) ₈ CH ₃	
104	CH	CF ₃	0	O	(CH ₂) ₃ -cyclo-C ₈ H ₁₁	
105	CH	CF ₃	0	O	(CH ₂) ₉ CH ₃	
106	CH	CF ₃	0	O	1-Adamantyl	
107	CH	CF ₃	0	O	(CH ₂) ₁₀ CH ₃	
108	CH	CF ₃	0	O	(CH ₂) ₁₁ CH ₃	
109	CH	CF ₃	0	O	CH(CH ₃)(CH ₂) ₉ CH ₃	
110	CH	CF ₃	0	O	(CH ₂) ₁₂ CH ₃	
111	CH	CF ₃	0	O	(CH ₂) ₁₃ CH ₃	
112	CH	CF ₃	0	O	(CH ₂) ₁₄ CH ₃	
113	CH	CF ₃	0	O	(CH ₂) ₁₅ CH ₃	
114	CH	CF ₃	0	O	(CH ₂) ₁₇ CH ₃	
115	CH	CF ₃	0	O	(CH ₂) ₁₉ CH ₃	
116	CH	CF ₃	0	O	CHO	
117	CH	CF ₃	0	O	CH=CH ₂	oil
118	CH	CF ₃	0	O	CH ₂ C=C(CH ₃) ₂	
119	CH	CF ₃	0	O	CH ₂ CH ₂ C=CH ₂	
120	CH	CF ₃	0	O	CH ₂ C=CH ₂	
121	CH	CF ₃	0	O	C(CH ₃)=CH ₂	
122	CH	CF ₃	0	O	(E)-CH ₂ CH=CHCH ₂ CH ₃	
123	CH	CF ₃	0	O	(Z)-CH ₂ CH=CHCH ₂ CH ₃	
124	CH	CF ₃	0	O	(CH ₂) ₅ C=CH ₂	
125	CH	CF ₃	0	O	C(=CHCH ₃)CH ₃	62-64
126	CH	CF ₃	0	O	Geranyl	
127	CH	CF ₃	0	O	3-Menthyl	
128	CH	CF ₃	0	O	C≡CH	

No.	X	Y	m	W	R ¹	m.p. [°C]
129	CH	CF ₃	0	O	CH ₂ C≡CH	
130	CH	CF ₃	0	O	CH ₂ CH ₂ C≡CH	
131	CH	CF ₃	0	O	CH ₂ CH ₂ C≡CH	
132	CH	CF ₃	0	O	(CH ₂) ₄ C≡CH	
133	CH	CF ₃	0	O	CHFCF ₃	oil
134	CH	CF ₃	0	O	COOCH ₂ CH ₃	oil
135	CH	CF ₃	0	O	CH ₂ CH ₂ OH	oil
136	CH	CF ₃	0	O	CH ₂ CH ₂ OCH ₃	oil
137	CH	CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	oil
138	CH	CF ₃	0	O	CH ₂ SC ₆ H ₅	oil
139	CH	CF ₃	0	O	CH ₂ CONHCH ₃	109-111
140	CH	CF ₃	0	O	CH ₂ CH(OH)CH ₂ OH	
141	CH	CF ₃	0	O	CH ₂ COCH ₃	
142	CH	CF ₃	0	O	COCH ₃	
143	CH	CF ₃	0	O	CH ₂ OC ₆ H ₅	
144	CH	CF ₃	0	O	COC ₆ H ₅	
145	CH	CF ₃	0	O	CO(4-Cl)-C ₆ H ₄	
146	CH	CF ₃	0	O	CF ₂ CH ₃	
147	CH	CF ₃	0	O	CH ₂ CN	
148	CH	CF ₃	0	O	CH ₂ CH ₂ CN	
149	CH	CF ₃	0	O	CH ₂ CH(-O-)CH ₂	
150	CH	CF ₃	0	O	CH ₂ (4-OCH ₃)C ₆ H ₅	
151	CH	CF ₃	0	O	CH ₂ -cyclo-(4-Oxo)-C ₆ H ₈	
152	CH	CF ₃	0	O	CH ₂ CH(OH)CH ₂ SC ₆ H ₅	
153	CH	CF ₃	0	O	CH ₂ CH ₂ Si(CH ₃) ₃	
154	CH	CF ₃	0	O	CH=CF ₂	
155	CH	CF ₃	0	O	CCl=CHCl	

No.	X	Y	m	W	R ¹	m.p. [°C]
156	CH	CF ₃	0	O	2-Pyridyl	99 - 101
157	CH	CF ₃	0	O	2-Furyl	
158	CH	CF ₃	0	O	2-Thienyl	106 -108
159	CH	CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₂ OTHP	
160	CH	CF ₃	0	O	CH ₂ CH ₂ Cl	oil
161	CH	CF ₃	0	O	Si(CH ₃) ₃	
162	CH	CF ₃	0	O	OC ₆ H ₅	
163	CH	CF ₃	0	O	OH	
164	CH	CF ₃	0	O	OCH ₃	
165	CH	CF ₃	0	O	OCH ₂ CH ₃	
166	CH	CF ₃	0	O	OCHF ₂	
167	CH	CF ₃	0	O	OCH ₂ C ₆ H ₅	
168	CH	CF ₃	0	O	CH ₂ SCH ₃	48-49
169	CH	CF ₃	0	O	SC ₆ H ₅	
170	CH	CF ₃	0	O	SeC ₆ H ₅	
171	CH	CF ₃	0	O	NH ₂	116 -118
172	CH	CF ₃	0	O	NHCH ₃	
173	CH	CF ₃	0	O	NHCH ₂ CH ₃	
174	CH	CF ₃	0	O	N(CH ₂ CH ₃) ₂	
175	CH	CF ₃	0	O	CONHCH ₂ C=CH ₂	105 -
176	CH	CF ₃	0	O	Cl	
177	CH	CF ₃	0	O	Br	
178	CH	CF ₃	0	O	CONH ₂	206 - 208
179	CH	CF ₃	0	O	NHCOCH ₃	129-131
180	CH	CF ₃	0	O	NHCOCH ₂ CH ₃	
181	CH	CF ₃	0	O	OSO ₂ CH ₃	
182	CH	CF ₃	0	O	SOCH ₂ (4-Br)-C ₆ H ₄	



No.	X	Y	m	W	R ¹	m.p. [°C]
183	CH	CF ₃	0	O	N(CH ₃)COOCH ₂ C ₆ H ₅	
184	CH	CF ₃	0	O	NHNH ₂	
185	CH	CF ₃	0	O	NHN(CH ₃) ₂	
186	N	CF ₃	0	O	CH ₃	
187	N	CF ₃	0	O	CH ₂ CH ₃	oil
188	N	CF ₃	0	O	(CH ₂) ₂ CH ₃	oil
189	N	CF ₃	0	O	CH(CH ₃) ₂	oil
190	N	CF ₃	0	O	(CH ₂) ₃ CH ₃	oil
191	N	CF ₃	0	O	CH ₂ CH(CH ₃) ₂	oil
192	N	CF ₃	0	O	C(CH ₃) ₃	
193	N	CF ₃	0	O	(CH ₂) ₄ CH ₃	oil
194	N	CF ₃	0	O	CH(CH ₃)(CH ₂) ₂ CH ₃	
195	N	CF ₃	0	O	CH ₂ C(CH ₃) ₃	
196	N	CF ₃	0	O	Cyclo-C ₅ H ₉	
197	N	CF ₃	0	O	(CH ₂) ₅ CH ₃	
198	N	CF ₃	0	O	Cyclo-C ₆ H ₁₁	
199	N	CF ₃	0	O	CH(CH ₃)(CH ₂) ₄ CH ₃	
200	N	CF ₃	0	O	CH ₂ -cyclo-C ₆ H ₁₁	
201	N	CF ₃	0	O	(CH ₂) ₇ CH ₃	
202	N	CF ₃	0	O	(CH ₂) ₈ CH ₃	
203	N	CF ₃	0	O	(CH ₂) ₉ CH ₃	
204	N	CF ₃	0	O	CH(CH ₃)(CH ₂) ₉ CH ₃	
205	N	CF ₃	0	O	(CH ₂) ₁₅ CH ₃	
206	N	CF ₃	0	O	(CH ₂) ₁₇ CH ₃	
207	N	CF ₃	0	O	(CH ₂) ₁₉ CH ₃	
208	N	CF ₃	0	O	CH ₂ CH=C(CH ₃) ₂	
209	N	CF ₃	0	O	CH ₂ CH ₂ CH=CH ₂	




No.	X	Y	m	W	R ¹	m.p. [°C]
210	N	CF ₃	0	O	CH ₂ CH=CH ₂	
211	N	CF ₃	0	O	(Z)-CH ₂ CH=CHCH ₂ CH ₃	
212	N	CF ₃	0	O	(CH ₂) ₅ CH=CH ₂	
213	N	CF ₃	0	O	CH ₂ C≡CH	
214	N	CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₃	
215	N	CF ₃	0	O	CHFCF ₃	
216	N	CF ₃	0	O	COOCH ₂ CH ₃	
217	N	CF ₃	0	O	CH ₂ CH ₂ OH	
218	N	CF ₃	0	O	CH ₂ CH ₂ OCH ₃	
219	N	CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	
220	N	CF ₃	0	O	CH ₂ SC ₆ H ₅	
221	N	CF ₃	0	O	CH ₂ CONHCH ₃	
222	N	CF ₃	0	O	CH ₂ CH(OH)CH ₂ OH	
223	N	CF ₃	0	O	CHO	
224	N	CF ₃	0	O	COCH ₃	
225	N	CF ₃	0	O	CH ₂ OC ₆ H ₅	
226	N	CF ₃	0	O	COC ₆ H ₅	
227	N	CF ₃	0	O	CF ₂ CH ₃	
228	N	CF ₃	0	O	CH ₂ CN	
229	N	CF ₃	0	O	CH ₂ CH ₂ CN	
230	N	CF ₃	0	O	CH=CF ₂	
231	N	CF ₃	0	O	2-Furyl	
232	N	CF ₃	0	O	CH ₂ C≡C-I	
233	N	CF ₃	0	O	OH	
234	N	CF ₃	0	O	OCH ₃	
235	N	CF ₃	0	O	OCH ₂ CH ₃	
236	N	CF ₃	0	O	OCHF ₂	



No.	X	Y	m	W	R ¹	m.p. [°C]
237	N	CF ₃	0	O	OCH ₂ C ₆ H ₅	
238	N	CF ₃	0	O	SC ₆ H ₅	
239	N	CF ₃	0	O	NH ₂	
240	N	CF ₃	0	O	NHCH ₃	
241	N	CF ₃	0	O	NHCH ₂ CH ₃	
242	N	CF ₃	0	O	N(CH ₂ CH ₃) ₂	
243	N	CF ₃	0	O	N(CH ₂ CN) ₂	
244	N	CF ₃	0	O	N(CH ₃) ₂	
245	N	CF ₃	0	O	NHCOCH ₃	
246	N	CF ₃	0	O	NHCOCH ₂ CH ₃	
247	N	CF ₃	0	O	OSO ₂ CH ₃	
248	N	CF ₃	0	O	NHNH ₂	
249	CH	CF ₃	0	S	CH ₃	
250	CH	CF ₃	0	S	CH ₂ CH ₃	
251	CH	CF ₃	0	S	(CH ₂) ₂ CH ₃	
252	CH	CF ₃	0	S	CHO	
253	CH	CF ₃	0	S	CHFCF ₃	
254	CH	CF ₃	0	S	CH ₂ C≡CH	
255	CH	CF ₃	0	S	COOCH ₂ CH ₃	
256	CH	CF ₃	0	S	CH ₂ COOC(CH ₃) ₃	
257	CH	CF ₃	0	S	CH ₂ CN	
258	CH	CF ₃	0	S	SeC ₆ H ₅	
259	N	CF ₃	0	S	CH ₃	
260	N	CF ₃	0	S	CH ₂ CH ₃	
261	N	CF ₃	0	S	(CH ₂) ₂ CH ₃	
262	N	CF ₃	0	S	CHFCF ₃	
263	N	CF ₃	0	S	CH ₂ CH ₂ OH	


No.	X	Y	m	W	R ¹	m.p. [°C]
264	N	CF ₃	0	S	CH ₂ COOC(CH ₃) ₃	
265	CH	CH ₂ CH ₂ Cl	0	O	CH ₃	
266	CH	CH ₂ CH ₂ Cl	0	O	CH ₂ CH ₃	
267	CH	CH ₂ CH ₂ Cl	0	O	(CH ₂) ₂ CH ₃	
268	CH	CH ₂ CH ₂ Cl	0	O	CH(CH ₃) ₂	
269	CH	CH ₂ CH ₂ Cl	0	O	CH ₂ SC ₆ H ₅	
270	CH	CH ₂ CH ₂ Cl	0	O	CH ₂ CONHCH ₃	
271	CH	CH ₂ CH ₂ Cl	0	O	NH ₂	
272	CH	CH ₂ CH ₂ Cl	0	O	NHCH ₂ CH ₃	
273	N	CH ₂ CH ₂ Cl	0	O	CH ₂ CH ₃	
274	N	CH ₂ CH ₂ Cl	0	O	NH ₂	
275	N	CH ₂ Cl	0	O	CH ₃	
276	CH	CH ₂ Cl	0	O	CH ₃	
277	CH	CHF ₂	0	O	CH ₃	
278	CH	CHF ₂	0	O	CH ₂ CH ₃	
279	CH	CHF ₂	0	O	(CH ₂) ₂ CH ₃	
280	CH	CHF ₂	0	O	CH ₂ CH=CH ₂	
281	CH	CHF ₂	0	O	C(CH ₃)=CH ₂	
282	CH	CHF ₂	0	O	COOCH ₂ CH ₃	
283	CH	CHF ₂	0	O	CH ₂ CONHCH ₃	
284	CH	CHF ₂	0	O	CF ₂ CH ₃	
285	CH	CHF ₂	0	O	CHO	
286	CH	CHF ₂	0	O	NH ₂	
287	CH	CHF ₂	0	O	Cl	
288	CH	CHF ₂	0	O	NHCOCH ₃	
289	CH	CHF ₂	0	O	NHNH ₂	
290	N	CHF ₂	0	O	CH ₃	




No.	X	Y	m	W	R ¹	m.p. [°C]
291	N	CHF ₂	0	O	CH ₂ CH ₃	
292	N	CHF ₂	0	O	CH(CH ₃)(CH ₂) ₄ CH ₃	
293	N	CHF ₂	0	O	CH ₂ CH=CH ₂	
294	N	CHF ₂	0	O	COOCH ₂ CH ₃	
295	N	CHF ₂	0	O	NH ₂	
296	CH	CF ₃	1	O	CH ₃	
297	CH	CF ₃	1	O	COOCH ₂ CH ₃	
298	CH	CF ₃	1	O	CH ₂ COOC(CH ₃) ₃	
299	CH	CF ₃	1	O	CHFCH ₃	
300	N	CF ₃	0	O	CH ₂ NHSO ₂ CH ₃	
301	N	CF ₃	0	O	(CH ₂) ₂ NHSO ₂ CH ₃	
302	N	CF ₃	0	O	CH ₂ NHSO ₂ CH ₂ CH ₃	
303	N	CF ₃	0	O	CH ₂ NHSO ₂ CH ₂ C ₆ H ₅	
304	CH	CF ₃	0	O	(CH ₂) ₄ NHSO ₂ CF ₃	
305	CH	CF ₃	0	O	(CH ₂) ₂ S(CH ₂) ₂ CH ₃	
306	CH	CF ₃	0	O	(CH ₂) ₄ S(CH ₂) ₄ OCH ₃	
307	CH	CF ₃	0	S	(CH ₂) ₂ S(CH ₂) ₂ CN	
308	CH	CF ₃	0	S	CH ₂ NHSO ₂ CH ₂ CH ₃	
309	CH	CF ₃	0	S	CH ₂ NHSO ₂ CH ₂ C ₆ H ₅	
310	CH	CF ₃	0	S	(CH ₂) ₂ NHSO ₂ CH ₃	
311	CH	CF ₃	0	S	CH ₂ NHSO ₂ CH ₃	
312	CH	CF ₃	0	S	CH(CH ₃)CH ₂ NHC ₆ H ₅	
313	CH	CF ₃	0	S	(CH ₂) ₂ S(2-F)-C ₆ H ₄	
314	CH	CF ₃	0	S	(CH ₂) ₆ NHCH ₂) ₆ OCH ₃	
315	CH	CF ₃	0	S	(CH ₂) ₂ NH-(2-F)-C ₆ H ₄	
316	CH	CF ₃	0	S	(CH ₂) ₃ NHCH ₂ CN	
317	CH	CF ₃	0	S	(CH ₂) ₂ O(3-Cl)-C ₆ H ₄	




No.	X	Y	m	W	R ¹	m.p. [°C]
318	CH	CF ₃	0	S	(CH ₂) ₆ NHCH ₂ CF ₃	
319	CH	CF ₃	0	S	(CH ₂) ₂ O(3-CH ₃)-C ₆ H ₄	
320	CH	CF ₃	0	O	CH ₂ NHC ₆ H ₅	
321	CH	CF ₃	0	O	(CH ₂) ₄ S(2-Br)-C ₆ H ₄	
322	CH	CF ₃	0	O	(CH ₂) ₆ NH(CH ₂) ₂ OCH ₃	
323	CH	CF ₃	0	O	(CH ₂) ₂ NH(CH ₂) ₄ OCH ₃	
324	CH	CF ₃	0	O	(CH ₂) ₃ NH-(4-CN)-C ₆ H ₄	
325	CH	CF ₃	0	O	(CH ₂) ₂ O(3-CH ₃)-C ₆ H ₄	
326	CH	CF ₃	0	O	(CH ₂) ₄ NHCH ₂ CF ₃	
327	CH	CF ₃	0	O	(CH ₂) ₄ NHCH ₂ CN	
328	CH	CF ₃	0	O	(CH ₂) ₃ O(4-OCH ₃)-C ₆ H ₄	
329	CH	CF ₃	0	O	CH ₂ SO ₂ -tert-C ₄ H ₉	oil
330	CH	CF ₃	0	O	CH ₂ SO ₂ -(4-F)-C ₆ H ₄	oil
331	CH	CF ₃	0	O	CH ₂ SO ₂ -C ₆ H ₅	oil
332	CH	CF ₃	0	O	CH ₂ SOCH ₃	63
333	CH	CF ₃	0	O	CH ₂ SO-C ₆ H ₅	oil
334	CH	CF ₃	0	O	CH ₂ CONH(CH ₂) ₂ CH ₃	80 – 82
335	CH	CF ₃	0	O	(4-OCF ₃)-C ₆ H ₄	57 – 59
336	CH	CF ₃	0	O	CH ₂ OCH ₃	oil
337	CH	CF ₃	0	O		53 – 54
338	CH	CF ₃	0	O		oil
339	CH	CF ₃	0	O	CH ₂ CH ₂ OCH ₂ CH ₃	oil
340	CH	CF ₃	0	O	CH ₂ CH ₂ NC ₆ H ₅	80-83


No.	X	Y	m	W	R ¹	m.p. [°C]
341	CH	CF ₃	0	O		80 – 81
342	CH	CF ₃	0	O		110 - 111
343	CH	CF ₃	0	O	CH ₂ CH ₂ O(CO)-(4-Cl)-C ₆ H ₄	80 – 82
344	CH	CF ₃	0	O	CH ₂ -(4-OCH ₃)-C ₆ H ₄	54 – 55
345	CH	CF ₃	0	O	CH ₂ -(3-Cl)-C ₆ H ₄	51 – 52
346	CH	CF ₃	0	O	CH ₂ -cyclo-C ₃ H ₅	oil
347	CH	CF ₃	0	O	CH ₂ -(4-C ₆ H ₅)-C ₆ H ₄	oil
348	CH	CF ₃	0	O		143 - 144
349	CH	CF ₃	0	O	CH ₂ CH ₂ O(CO)-(2,6-F ₂)-C ₆ H ₃	57 - 58
350	CH	CF ₃	0	O	CH ₂ CH ₂ O(CO)-(4-NO ₂)-C ₆ H ₄	80 - 81
351	CH	CF ₃	0	O	CH ₂ -(2,6-Cl ₂)-C ₆ H ₃	91 - 92
352	CH	CF ₃	0	O	CH ₂ CH ₂ OSO ₂ CH ₃	oil
353	CH	CF ₃	0	O	CH ₂ CH ₂ O(CO)-tert-C ₄ H ₉	oil
354	CH	CF ₃	0	O	CH ₂ -(3-F)-C ₆ H ₄	50 - 51
355	CH	CF ₃	0	O	CH ₂ CONCH ₂ C≡CH	129 - 131


No.	X	Y	m	W	R ¹	m.p. [°C]
356	CH	CF ₃	0	O	CH ₂ CH ₂ O(CO)-cyclo-C ₃ H ₇	oil
357	CH	CF ₃	0	O	CH ₂ CH ₂ O(CO)CH ₃	oil
358	CH	CF ₃	0	O	CH ₂ -[2,4-(CH ₃) ₂]-C ₆ H ₃	85 - 86
359	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH=CH ₂	210 - 212
360	CH	CF ₃	0	O	CH ₂ CON(CH ₂ CH ₃) ₂	oil
361	CH	CF ₃	0	O	CH ₂ CON(CH ₂) ₃ CH ₃	77 - 79
362	CH	CF ₃	0	O	CH ₂ CONCH ₂ -(2-furyl)	139 - 141
363	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃) ₂	112 - 114
364	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)[(CH ₂) ₄ CH ₃]	73 - 75
365	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ C ₆ H ₅	120 - 122
366	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ OCH ₂ CH ₃	78
367	CH	CF ₃	0	O	CH ₂ CONCH ₂ CF ₃	176 - 178
368	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)[(CH ₂) ₅ CH ₃]	85 - 86
369	CH	CF ₃	0	O		oil
370	CH	CF ₃	0	O		oil
371	CH	CF ₃	0	O	CH ₂ CH ₂ -(1-pyrrol)	oil
372	CH	CF ₃	0	O	CH ₂ CH ₂ C ₆ H ₅	oil





No.	X	Y	m	W	R ¹	m.p. [°C]
373	CH	CF ₃	0	O	CH ₂ Cl	53 - 54
374	CH	CF ₃	0	O	(CH ₂) ₃ OH	38 - 39
375	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)[(CH ₂) ₂]CH ₃	68 - 69
376	CH	CF ₃	0	O	CH ₂ CH(OCH ₃) ₂	oil
377	CH	CF ₃	0	O	CH ₂ CONCH ₂ C(CH ₃) ₃	oil
378	CH	CF ₃	0	O	CH ₂ CONC(CH ₃) ₂ (CH ₂ CH ₃)	oil
379	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ -cyclo-C ₆ H ₁₁	82 - 85
380	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)(1-naphthyl)	142 - 146
381	CH	CF ₃	0	O	(CH ₂) ₃ Cl	oil
382	CH	CF ₃	0	O	CH ₂ CON-tert-C ₄ H ₉	oil
383	CH	CF ₃	0	O	CH ₂ CON(iso-C ₃ H ₇) ₂	70 - 72
384	CH	CF ₃	0	O	CH ₂ CON(CH ₂) ₇ CH ₃	79 - 81
385	CH	CF ₃	0	O	CH ₂ CON-cyclo-C ₆ H ₁₁	119 - 121
386	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ -(4-Cl)-C ₆ H ₄	120 - 121
387	CH	CF ₃	0	O	CH ₂ CONCH ₂ -(2-thienyl)	137 - 139
388	CH	CF ₃	0	O		151 - 153
389	CH	CF ₃	0	O	CH ₂ CONHCH(CH ₃)(CH ₂ CH ₃)	87-89
390	CH	CF ₃	0	O	(CH ₂) ₃ SCH ₃	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
391	CH	CF ₃	0	O	(CH ₂) ₃ SOCH ₃	oil
392	CH	CF ₃	0	O	CH ₂ CONC(CH ₃) ₂ (C≡CH)	111-113
393	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)CH ₂ CH ₂ CH(CH ₃) ₂	72-74
394	CH	CF ₃	0	O		oil
395	CH	CF ₃	0	O	CH ₂ CON-cyclo-C ₅ H ₉	110 - 112
396	CH	CF ₃	0	O	CH ₂ CON(CH ₂) ₄ CH ₃	75 - 77
397	CH	CF ₃	0	O		190 - 192
398	CH	CF ₃	0	O	CH ₂ CON(3-CF ₃)C ₆ H ₄	136 - 138
399	CH	CF ₃	0	O	CH ₂ CON-cyclo-C ₈ H ₁₇	115 - 117
400	CH	CF ₃	0	O		oil
401	CH	CF ₃	0	O	CH ₂ CON-Adamantyl	oil
402	CH	CF ₃	0	O	CH ₂ CON(CH ₂ CH ₂ CH ₃) ₂	oil
403	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)[(4-F)-C ₆ H ₄]	111 - 113
404	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH(CH ₃) ₂	91 - 93






No.	X	Y	m	W	R ¹	m.p. [°C]
405	CH	CF ₃	0	O		Oil
406	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ OC ₆ H ₅	99 - 101
407	CH	CF ₃	0	O	CH ₂ CH=NOCH ₃	oil
408	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ -[3,4-(OCH ₃) ₂]C ₆ H ₃	123-125
409	CH	CF ₃	0	O	CH ₂ CON-(2-Cl)C ₆ H ₄	138 - 140
410	CH	CF ₃	0	O	CH ₂ CON-(2-SCH ₃)C ₆ H ₄	136 - 138
411	CH	CF ₃	0	O		222 - 225
412	CH	CF ₃	0	O		207 - 209
413	CH	CF ₃	0	O	CH ₂ CON-(3-Br)C ₆ H ₄	129 - 131
414	CH	CF ₃	0	O	CH ₂ CON-N-(2,4,6-Cl ₃)C ₆ H ₂	153 - 155
415	CH	CF ₃	0	O	CH ₂ CON-(4-I)C ₆ H ₄	143 - 145
416	CH	CF ₃	0	O	CH ₂ CON-NCOCH ₂ (3-Thienyl)	185 - 187
417	CH	CF ₃	0	O	CH ₂ CH ₂ CHO	oil
418	CH	CF ₃	0	O	CH ₂ CON(CH ₃)[(CH ₂) ₃ CH ₃]	oil
419	CH	CF ₃	0	O	CH ₂ CON-(3,5-Cl ₂ -2,4-F ₂)C ₆ H	166 - 167





No.	X	Y	m	W	R ¹	m.p. [°C]
420	CH	CF ₃	0	O	CH ₂ CON-C ₆ H ₅	215 – 217
421	CH	CF ₃	0	O	CH ₂ CON(CH ₃)(C ₆ H ₁₁)	oil
422	CH	CF ₃	0	O	CH ₂ CON(CH ₂ CH ₃)(CH ₂ CH=CH ₂)	oil
423	CH	CF ₃	0	O	CH ₂ CON(CH ₂ CH ₃)[CH(CH ₃) ₂]	oil
424	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)[(CH ₃) ₂]	108 – 110
425	CH	CF ₃	0	O	CH ₂ CON(CH ₂ CH ₃)[CH ₂ C(=CH ₂)(CH ₃)]	oil
426	CH	CF ₃	0	O	CH ₂ CONCH ₂ (4-tert-C ₄ H ₉)C ₆ H ₄	oil
427	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)(tert-C ₄ H ₉)	oil
428	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)[CH ₂ CH(CH ₃)(CH ₂ CH ₃)]	oil
429	CH	CF ₃	0	O	CH ₂ CONCH ₂ COOCH ₂ CH ₃	103 - 105
430	CH	CF ₃	0	O	CH ₂ CON[(CH ₂) ₂ CH ₃](CH ₂ -cyclo-C ₃ H ₇)	oil
431	CH	CF ₃	0	O	CH ₂ CONCH(CH ₃)CH ₂ CH ₂ CH(CH ₃) ₂	80 - 82
432	CH	CF ₃	0	O	CH ₂ CONCH(CH ₂ CH ₃)[CH ₂ CH(CH ₃) ₂]	oil
433	CH	CF ₃	0	O	CH ₂ C=O-(1-PiperidinyI)	oil
434	CH	CF ₃	0	O		180 - 182
435	CH	CF ₃	0	O	CH ₂ CONCH ₂ C(=CH ₂)(CH ₃)	86 - 87
436	CH	CF ₃	0	O	CH ₂ CONCH[CH(CH ₃) ₂](COOCH ₃)	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
437	CH	CF ₃	0	O	CH ₂ CONCH ₂ -cyclo-C ₃ H ₇	oil
438	CH	CF ₃	0	O	CH ₂ CON(CH ₂) ₅ OH	oil
439	CH	CF ₃	0	O	CH ₂ CON(CH ₃)(CH ₂ CO ₂ CH ₃)	oil
440	CH	CF ₃	0	O	CH ₂ CON(CH ₃)(CH ₂ CN)	oil
441	CH	CF ₃	0	O	CH ₂ CONCH[CH ₂ CH(CH ₃) ₂](CO ₂ CH ₃)	oil
442	CH	CF ₃	0	O	CH ₂ CON-(1-PiperidinyI)	oil
443	CH	CF ₃	0	O	CH ₂ CONCH ₂ CH ₂ OCH ₃	97 - 99
444	CH	CF ₃	0	O	CH ₂ CH ₂ SC ₆ H ₅	oil
445	CH	CF ₃	0	O	CH ₂ CH ₂ SCH ₃	oil
446	CH	CF ₃	0	O	CH ₂ CH ₂ SCH ₂ C ₆ H ₅	oil
447	CH	CF ₃	0	O		oil
448	CH	CF ₃	0	O	CH ₂ CON-(2-OH)C ₆ H ₄	162 - 164
449	CH	CF ₃	0	O	CH ₂ CON-(3-OH)C ₆ H ₄	oil
450	CH	CF ₃	0	O	CH ₂ CON-(2-CH ₃)C ₆ H ₄	163 - 164
451	CH	CF ₃	0	O	CH ₂ CON-(3-NO ₂)C ₆ H ₄	176 - 178
452	CH	CF ₃	0	O	CH ₂ CON-(3-OCF ₂ CHFCI)C ₆ H ₄	120 - 121
453	CH	CF ₃	0	O	CH ₂ CON-(3-CF ₃ -4-F)C ₆ H ₃	168 - 170
454	CH	CF ₃	0	O	CH ₂ CON-(2,4-Cl ₂)C ₆ H ₃	120 - 122





No.	X	Y	m	W	R ¹	m.p. [°C]
455	CH	CF ₃	0	O	CH ₂ CON-(2-F-4-Cl)C ₆ H ₃	148 - 151
456	CH	CF ₃	0	O	CH ₂ CON-[2,4-(CH ₃) ₂]C ₆ H ₃	123 - 125
457	CH	CF ₃	0	O	CH ₂ CON-[2,3-(CH ₃) ₂]C ₆ H ₃	waxy
458	CH	CF ₃	0	O		waxy
459	CH	CF ₃	0	O	CH ₂ CON-(2-CH ₃ -3-Cl)C ₆ H ₃	160 - 162
460	CH	CF ₃	0	O	CH ₂ CON(CH ₂ CH ₃)(C ₆ H ₅)	oil
461	CH	CF ₃	0	O		124 - 126
462	CH	CF ₃	0	O	CH ₂ CON(2-OCH ₃ -5-Ph)C ₆ H ₃	167 - 169
463	CH	CF ₃	0	O		157 - 158
464	CH	CF ₃	0	O	CH ₂ CON-(3-NO ₂ -4-Cl)C ₆ H ₃	oil
465	CH	CF ₃	0	O	CH ₂ CON-(2-Cl-4-CH ₃)C ₆ H ₃	106 - 108
466	CH	CF ₃	0	O	CH ₂ CON-(3-OCH ₂ CH ₃)C ₆ H ₄	waxy
467	CH	CF ₃	0	O		169 - 171



No.	X	Y	m	W	R ¹	m.p. [°C]
468	CH	CF ₃	0	O	CH ₂ CON-(4-CH ₃)C ₆ H ₄	139 - 141
469	CH	CF ₃	0	O	CH ₂ CON-(1-Naphthyl)	155 - 157
470	CH	CF ₃	0	O	CH ₂ CON-(3-I)C ₆ H ₄	135 - 137
471	CH	CF ₃	0	O	CH ₂ CON-(2-OCH ₂ CH ₃)C ₆ H ₄	138
472	CH	CF ₃	0	O	CH ₂ CON-(2-OCH ₃)C ₆ H ₄	130 - 132
473	CH	CF ₃	0	O	CH ₂ CON-[3,5-(OCH ₃) ₂]C ₆ H ₃	130 - 132
474	CH	CF ₃	0	O	CH ₂ CON-(4-Cl)C ₆ H ₄	139 - 141
475	CH	CF ₃	0	O	CH ₂ CON-(3-CH ₃)C ₆ H ₄	oil
476	CH	CF ₃	0	O	CH ₂ CON-(3-OCH ₃)C ₆ H ₄	oil
477	CH	CF ₃	0	O	CH ₂ CON-(4-CH ₂ CH ₃)C ₆ H ₄	122 - 123
478	CH	CF ₃	0	O	CH ₂ CON-(4-CF ₃)C ₆ H ₄	151 - 152
479	CH	CF ₃	0	O	CH ₂ CON-(2-CH ₃ -4-Cl)C ₆ H ₃	165 - 167
480	CH	CF ₃	0	O	CH ₂ CH ₂ NCH ₂ C ₆ H ₅	oil
481	CH	CF ₃	0	O	CH ₂ CH ₂ NCH ₂ -(3-Pyridyl)	oil
482	CH	CF ₃	0	O	CH ₂ CH=NOCH ₂ CH ₃	oil
483	CH	CF ₃	0	O	CH ₂ CH=NOC ₆ H ₅	oil
484	CH	CF ₃	0	O	CH ₂ CON-(4-NO ₂)C ₆ H ₄	181 - 183
485	CH	CF ₃	0	O	CH ₂ CON-(2-CH ₃ -4-NO ₂)C ₆ H ₃	129 - 131
486	CH	CF ₃	0	O	CH ₂ CON-(2-Cl-3-CF ₃)C ₆ H ₃	136


No.	X	Y	m	W	R ¹	m.p. [°C]
487	CH	CF ₃	0	O	CH ₂ CON-(2-CN-4-Cl)C ₆ H ₃	157 - 159
488	CH	CF ₃	0	O	CH ₂ CON-(3,5-Cl ₂)C ₆ H ₃	167 - 169
489	CH	CF ₃	0	O	CH ₂ CON-(3,5-Cl ₂ -4-OCF ₂ CHF ₂)C ₆ H ₂	132 - 134
490	CH	CF ₃	0	O	CH ₂ CON-(2,4,5-Cl ₃)C ₆ H ₂	146
491	CH	CF ₃	0	O	CH ₂ CON-(3,5-Cl ₂ -4-OCF ₂ CHFCF ₃)C ₆ H ₂	124 - 126
492	CH	CF ₃	0	O	CH ₂ CON-(2-CF ₃ -4-Cl)C ₆ H ₃	136
493	CH	CF ₃	0	O		oil
494	CH	CF ₃	0	O		91-93
495	CH	CF ₃	0	O		123-125
496	CH	CF ₃	0	O		81-83
497	CH	CF ₃	0	O		113-115

No.	X	Y	m	W	R ¹	m.p. [°C]
498	CH	CF ₃	0	O	COOH	155-157
499	CH	CF ₃	0	O	4-F-C ₆ H ₄	104-106
500	CH	CF ₃	0	O	CON(C ₂ H ₅) ₂	oil
501	CH	CF ₃	0	O	CONCH(CH ₃) ₂	oil
502	CH	CF ₃	0	O	CON(CH ₃) ₂	52-54
503	CH	CF ₃	0	O	CONHCH ₂ CCH	105-107
504	CH	CF ₃	0	O	CONH-cyclo-C ₃ H ₅	101-103
505	CH	CF ₃	0	O	CONH ₂	206-208
506	CH	CF ₃	0	O		72-74
507	CH	CF ₃	0	O		98-100
508	CH	CF ₃	0	O		108-110
509	CH	CF ₃	0	O		140-142






No.	X	Y	m	W	R ¹	m.p. [°C]
510	CH	CF ₃	0	O	CONHCH ₃	127-129
511	CH	CF ₃	0	O	CONHCH ₂ CH=CH ₂	oil
512	CH	CF ₃	0	O	CON(CH ₂ CN) ₂	90-92
513	CH	CF ₃	0	O	4-(t-C ₄ H ₉)-C ₆ H ₄	64-66
514	CH	CF ₃	0	O	4-CF ₃ -C ₆ H ₄	89-91
515	CH	CF ₃	0	O	4-CH ₃ -3-F-C ₆ H ₃	104-106
516	CH	CF ₃	0	O	2,4-di-Cl-C ₆ H ₃	70-72
517	CH	CF ₃	0	O	4-(NHSO ₂ CH ₃)-C ₆ H ₄	204-206
518	CH	CF ₃	0	O	2,6-di-Cl-C ₆ H ₃	139-141
519	CH	CF ₃	0	O	COOCH ₂ C ₆ H ₅	83-85
520	CH	CF ₃	0	O	CONHC ₃ H ₇	oil
521	CH	CF ₃	0	O	3,5-di-Br-4-(OCH ₃)-C ₆ H ₂	132-134
522	CH	CF ₃	0	O	CHCl ₂	oil
523	CH	CF ₃	0	O	CCl ₃	oil
524	CH	CF ₃	0	O	CH(OCH ₃) ₂	oil
525	CH	CF ₃	0	O	3-CF ₃ -C ₆ H ₄	57-59
526	CH	CF ₃	0	O	CON(CH ₂) ₅	oil
527	CH	CF ₃	0	O	CON(CH ₃)CH ₂ C ₆ H ₅	oil
528	CH	CF ₃	0	O	CONHCH ₂ C ₆ H ₅	96-98






No.	X	Y	m	W	R ¹	m.p. [°C]
529	CH	CF ₃	0	O		oil
530	CH	CF ₃	0	O	CONH-n-C ₈ H ₁₃	oil
531	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ C ₆ H ₅	oil
532	CH	CF ₃	0	O	CONH-c-C ₆ H ₁₁	115-117
533	CH	CF ₃	0	O	CON(n-C ₄ H ₉) ₂	oil
534	CH	CF ₃	0	O		oil
535	CH	CF ₃	0	O	CONH-i-C ₄ H ₉	oil
536	CH	CF ₃	0	O		oil
537	CH	CF ₃	0	O	CON(CH ₂) ₄	68-70
538	CH	CF ₃	0	O	CON(CH ₃)-n-C ₆ H ₁₃	oil
539	CH	CF ₃	0	O		oil






No.	X	Y	m	W	R ¹	m.p. [°C]
540	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₃	oil
541	CH	CF ₃	0	O	CONHOCH ₃	oil
542	CH	CF ₃	0	O		oil
543	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ CH ₃	oil
544	CH	CF ₃	0	O	CONHCH ₂ CH(OCH ₃) ₂	oil
545	CH	CF ₃	0	O	CONH-t-C ₄ H ₉	113-115
546	CH	CF ₃	0	O	CONHCH ₂ -4-Cl-C ₆ H ₄	oil
547	CH	CF ₃	0	O	CONHCH(CH ₃)C ₆ H ₅	oil
548	CH	CF ₃	0	O	CONHCH ₂ CH ₂ OCH ₃	92-94
549	CH	CF ₃	0	O		190-192
550	CH	CF ₃	0	O	CONHC(CH ₃) ₂ CCH	90-92
551	CH	CF ₃	0	O	CONHCH ₂ -2-Furyl	93-95
552	CH	CF ₃	0	O	CON(CH ₂) ₃	91-93
553	CH	CF ₃	0	O	CONHCH ₂ -c-C ₃ H ₅	oil
554	CH	CF ₃	0	O	CONHC(CH ₃) ₂ CH ₂ CH ₃	oil



No.	X	Y	m	W	R ¹	m.p. [°C]
555	CH	CF ₃	0	O	CONH(CH ₂) ₃ C ₆ H ₅	oil
556	CH	CF ₃	0	O	CONHCH ₂ -3-Pyridyl	132-134
557	CH	CF ₃	0	O	CON(CH ₃)-n-C ₄ H ₉	oil
558	CH	CF ₃	0	O	CON(CH ₂ CH ₃)-i-C ₃ H ₇	oil
559	CH	CF ₃	0	O		oil
560	CH	CF ₃	0	O	CONHCH ₂ CH ₂ Cl	oil
561	CH	CF ₃	0	O	CONHCH ₂ CN	152-157
562	CH	CF ₃	0	O	CON(CH ₃)OCH ₃	oil
563	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH=CH ₂	oil
564	CH	CF ₃	0	O	CONHCH ₂ COOCH ₃	oil
565	CH	CF ₃	0	O	CON(CH ₃)-i-C ₃ H ₇	oil
566	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ CN	oil
567	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH(OCH ₃) ₂	oil
568	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH(-CH ₂ CH ₂ O-)	oil
569	CH	CF ₃	0	O	CONHCH ₂ C(=CH ₂)CHH ₃	oil
570	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ CH=CH ₂	oil
571	CH	CF ₃	0	O	CONHC ₆ H ₅	83-85





No.	X	Y	m	W	R ¹	m.p. [°C]
572	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CCH	oil
573	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CN	oil
574	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ N(CH ₃) ₂	oil
575	CH	CF ₃	0	O	CONHOCH ₂ CH ₃	114-116
576	CH	CF ₃	0	O	CONHCH ₂ CF ₃	74-76
577	CH	CF ₃	0	O	CON(CH ₂ CH ₂ Cl) ₂	oil
578	CH	CF ₃	0	O	CONH-C-C ₄ H ₇	oil
579	CH	CF ₃	0	O	CON(CH ₂ CH ₂ CH ₃)CH ₂ -C-C ₃ H ₅	oil
580	CH	CF ₃	0	O	CON(CH ₃)-C-C ₆ H ₁₁	oil
581	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ C(=CH ₂)CH ₃	oil
582	CH	CF ₃	0	O	CONHOCH ₂ CH=CH ₂	90-92
583	CH	CF ₃	0	O	CONHOCH ₂ C ₆ H ₅	126-128
584	CH	CF ₃	0	O	CON(CH ₃)CH ₂ COOCH ₃	oil
585	CH	CF ₃	0	O	COONHCH ₃	230-232
586	CH	CF ₃	0	O	CONHCH ₂ CH ₃	83-85
587	CH	CF ₃	0	O	CONHCH(CH ₃)COOCH ₃	104-106
588	CH	CF ₃	0	O	CONHCH(i-C ₃ H ₇)COOCH ₃	oil
589	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CON(CH ₃) ₂	oil
590	CH	CF ₃	0	O	CON(CH ₃)-t-C ₄ H ₉	oil




No.	X	Y	m	W	R ¹	m.p. [°C]
591	CH	CF ₃	0	O	CONHO-t-C ₄ H ₉	103-105
592	CH	CF ₃	0	O	CON(CH ₃)CH(i-C ₃ H ₇)COOCH ₃	oil
593	CH	CF ₃	0	O	CH(OCH ₂ CH ₃) ₂	oil
594	CH	CF ₃	0	O		oil
595	CH	CF ₃	0	O		oil
596	CH	CF ₃	0	O		oil
597	CH	CF ₃	0	O		oil
598	CH	CF ₃	0	O		oil
599	CH	CF ₃	0	O	CONHCH ₂ CONHCH ₃	101-103
600	CH	CF ₃	0	O	CON(CH ₂) ₇	oil
601	CH	CF ₃	0	O	CON(CH ₂) ₆	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
602	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ CH ₂ OCH ₃	oil
603	CH	CF ₃	0	O		oil
604	CH	CF ₃	0	O		oil
605	CH	CF ₃	0	O		oil
606	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ CH ₂ CN	oil
607	CH	CF ₃	0	O		oil
608	CH	CF ₃	0	O	CON(CH ₂ CH ₃)-n-C ₄ H ₉	oil
609	CH	CF ₃	0	O		179-181
610	CH	CF ₃	0	O	CONHCH(CH ₃)CONHCH ₃	136-138
611	CH	CF ₃	0	O	COON(CH ₂) ₄	64-66
612	CH	CF ₃	0	O	CONHCH ₂ CON(CH ₃) ₂	107-109
613	CH	CF ₃	0	O	CON(CH ₂ COOCH ₂ CH ₃) ₂	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
614	CH	CF ₃	0	O		180-182
615	CH	CF ₃	0	O		221-223
616	CH	CF ₃	0	O		234-236
617	CH	CF ₃	0	O		oil
618	CH	CF ₃	0	O	CON(CH ₃)CH ₂ -6-Cl-3-pyridyl	oil
619	CH	CF ₃	0	O		105-107
620	CH	CF ₃	0	O	CONHCH(CH ₃)CH(OCH ₃) ₂	oil
621	CH	CF ₃	0	O	CONHCH ₂ CH ₂ SCH ₃	oil
622	CH	CF ₃	0	O	CONHCH(CH ₃)CH ₂ OCH ₃	70-72
623	CH	CF ₃	0	O	CONHCH ₂ CH ₂ NHCOCH ₃	124-126
624	CH	CF ₃	0	O	CONH(CH ₂) ₃ OCH ₂ CH ₃	oil
625	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ CH ₂ CH ₃	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
626	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ OCH ₃	oil
627	CH	CF ₃	0	O	CONHCH ₂ CH ₂ SCH ₂ CH ₃	oil
628	CH	CF ₃	0	O	CONHCH ₂ CH ₂ OCH ₂ CH ₃	59-61
629	CH	CF ₃	0	O		oil
630	CH	CF ₃	0	O		174-176
631	CH	CF ₃	0	O	CONHCH(CH ₃)CH(OCH ₃) ₂	oil
632	CH	CF ₃	0	O	CONHCH ₂ CH ₂ SCH ₃	oil
633	CH	CF ₃	0	O	CONHCH(CH ₃)CH ₂ OCH ₃	70-72
634	CH	CF ₃	0	O	CONHCH ₂ CH ₂ NHCOCH ₃	124-126
635	CH	CF ₃	0	O	CONH(CH ₂) ₃ OCH ₂ CH ₃	oil
636	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ CH ₂ CH ₃	oil
637	CH	CF ₃	0	O	CON(CH ₂ CH ₃)CH ₂ OCH ₃	oil
638	CH	CF ₃	0	O	CONHCH ₂ CH ₂ SCH ₂ CH ₃	oil
639	CH	CF ₃	0	O	CONHCH(CH ₃)CH ₂ COOCH ₂ CH ₃	oil
640	CH	CF ₃	0	O	CONH-4-COOCH ₃ -C ₆ H ₄	189-191
641	CH	CF ₃	0	O	CONH-4-CONH ₂ -C ₆ H ₄	265-267
642	CH	CF ₃	0	O	CONHCH ₂ CH ₂ Br	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
643	CH	CF ₃	0	O	CONHCH ₂ CH=CHCH ₂ Cl	oil
644	CH	CF ₃	0	O	CONH-4-CONHCH ₃ -C ₆ H ₄	219-221
645	CH	CF ₃	0	O	CONHCH ₂ CH ₂ CH ₂ Br	oil
646	CH	CF ₃	0	O	CONHCH ₂ CH ₂ CH ₂ OCH ₃	oil
647	CH	CF ₃	0	O	CONH-4-CH ₂ CH ₃ -C ₆ H ₄	97-99
648	CH	CF ₃	0	O	CONHCH ₂ CH ₂ OCH(CH ₃) ₂	oil
649	CH	CF ₃	0	O	CONHCH ₂ CH ₂ CH ₂ OCH ₂ CH ₃	oil
650	CH	CF ₃	0	O		oil
651	CH	CF ₃	0	O		64-66
652	CH	CF ₃	0	O		oil
653	CH	CF ₃	0	O		oil
654	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH ₃	oil

No.	X	Y	m	W	R ¹	m.p. [°C]
655	CH	CF ₃	0	O	CH ₂ CON(CH ₃) ₂	58-60
656	CH	CF ₃	0	O	CH ₂ CON(CH ₂) ₄	101-103
657	CH	CF ₃	0	O		oil
658	CH	CF ₃	0	O		90-92
659	CH	CF ₃	0	O	CH ₂ CONHCH ₂ CH ₃	104-106
660	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH ₂ OH	oil
661	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH ₂ CH ₃	oil
662	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH(-OCH ₂ CH ₂ O-)	oil
663	CH	CF ₃	0	O	CH ₂ CONHCH ₂ CH ₃	104-106
664	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH ₂ OH	oil
665	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH ₂ CH ₃	oil
667	CH	CF ₃	0	O	CH ₂ CON(CH ₃)CH ₂ CH(-OCH ₂ CH ₂ O-)	oil
668	CH	CF ₃	0	O		79-81
669	CH	CF ₃	0	O	CH ₂ CONHCH ₂ CH ₂ SCH ₃	65-67
670	CH	CF ₃	0	O	CH ₂ CONHCH(CH ₃)CH ₂ OCH ₃	86-88




No.	X	Y	m	W	R ¹	m.p. [°C]
671	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ OCO-c-C ₄ H ₇	oil
672	CH	CF ₃	0	O	CH ₂ CONHCH ₂ CH ₂ Br	87-89
673	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ OCOC ₆ H ₅	oil
674	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ OCO-c-C ₃ H ₅	oil
675	CH	CF ₃	0	O	CONH-2-CH ₃ -C ₆ H ₄	104-106
676	CH	CF ₃	0	O	CH ₂ CON(i-C ₃ H ₇)-4-F-C ₆ H ₄	102-104
677	CH	CF ₃	0	O		oil
678	CH	CF ₃	0	O		oil
679	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ OCONHC ₆ H ₅	100-102
680	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ OCONHCH ₂ CH ₃	oil
681	CH	CF ₃	0	O	CON(CH ₃)CH ₂ CH ₂ OSO ₂ CH ₃	oil
682	CH	CF ₃	0	O	CH ₂ CONH-c-C ₄ H ₇	133-135
683	CH	CF ₃	0	O	CH ₂ CONHCH ₂ CN	158-160

Table 2



No.	X	Y	W	R ¹	m.p. [°C]
684	N	(CF ₂) ₃ CHF ₂	O	CH ₃	
685	N	(CF ₂) ₂ CF ₃	O	CH ₂ CH ₃	
686	N	(CF ₂) ₂ CF ₃	O	COOCH ₂ CH ₃	
687	N	(CF ₂) ₂ CF ₃	O	OH	
688	N	(CF ₂) ₂ CF ₃	O	OCH ₃	
689	N	CF ₂ CF ₃	O	CH ₃	
690	N	CF ₂ CF ₃	O	CH ₂ CH ₃	
691	N	CF ₂ CF ₃	S	CH ₃	
692	N	CF ₂ CF ₃	S	CH ₂ CH ₃	
693	N	CF ₂ CF ₃	S	(CH ₂) ₂ CH ₃	
694	CH	CF ₃	O	CH ₃	oil
695	CH	CF ₃	O	CH ₂ CH ₃	
696	CH	CF ₃	O	(CH ₂) ₂ CH ₃	
697	CH	CF ₃	O	CH(CH ₃) ₂	
698	CH	CF ₃	O	(CH ₂) ₃ CH ₃	
699	CH	CF ₃	O	CH(CH ₃)CH ₂ CH ₃	
700	CH	CF ₃	O	CH ₂ CH(CH ₃) ₂	
701	CH	CF ₃	O	C(CH ₃) ₃	oil
702	CH	CF ₃	O	(CH ₂) ₄ CH ₃	
703	CH	CF ₃	O	CH(CH ₃)(CH ₂) ₂ CH ₃	

No.	X	Y	W	R ¹	m.p. [°C]
704	CH	CF ₃	O	(CH ₂) ₂ CH(CH ₃) ₂	
705	CH	CF ₃	O	CH ₂ C(CH ₃) ₃	
706	CH	CF ₃	O	cyclo-C ₅ H ₉	
707	CH	CF ₃	O	cyclo-C ₆ H ₁₁	
708	CH	CF ₃	O	CHO	
709	CH	CF ₃	O	CH=CH ₂	
710	CH	CF ₃	O	CH ₂ CH=C(CH ₃) ₂	
711	CH	CF ₃	O	CH ₂ CH=CH ₂	
712	CH	CF ₃	O	C(CH ₃)=CH ₂	
713	CH	CF ₃	O	(CH ₂) ₅ C=CH ₂	
714	CH	CF ₃	O	C(=CHCH ₃)CH ₃	
715	CH	CF ₃	O	CH ₂ C≡CH	
716	CH	CF ₃	O	CH ₂ CH ₂ C≡CH	
717	CH	CF ₃	O	CH ₂ C≡CCH ₂ CH ₃	
718	CH	CF ₃	O	(CH ₂) ₄ C≡CH	
719	CH	CF ₃	O	CHFCF ₃	
720	CH	CF ₃	O	COOCH ₂ CH ₃	
721	CH	CF ₃	O	CH ₂ CH ₂ OH	
722	CH	CF ₃	O	CH ₂ CH ₂ OCH ₃	
723	CH	CF ₃	O	CH ₂ COOC(CH ₃) ₃	
724	CH	CF ₃	O	CH ₂ SC ₆ H ₅	
725	CH	CF ₃	O	CH ₂ CONHCH ₃	
726	CH	CF ₃	O	CH ₂ CH(OH)CH ₂ OH	
727	CH	CF ₃	O	CH ₂ COCH ₃	
728	CH	CF ₃	O	COCH ₃	
729	CH	CF ₃	O	CH ₂ OC ₆ H ₅	
730	CH	CF ₃	O	COC ₆ H ₅	
731	CH	CF ₃	O	CF ₂ CH ₃	


No.	X	Y	W	R ¹	m.p. [°C]
732	CH	CF ₃	O	CH ₂ CN	
733	CH	CF ₃	O	CH ₂ CH(-O-)CH ₂	
734	CH	CF ₃	O	CH ₂ (4-OCH ₃)C ₆ H ₅	
735	CH	CF ₃	O	CH ₂ CH(OH)CH ₂ SC ₆ H ₅	
736	CH	CF ₃	O	CH=CF ₂	
737	CH	CF ₃	O	CCl=CHCl	
738	CH	CF ₃	O	2-Pyridyl	
739	CH	CF ₃	O	OC ₆ H ₅	
740	CH	CF ₃	O	OH	
741	CH	CF ₃	O	OCH ₃	
742	CH	CF ₃	O	OCH ₂ CH ₃	
743	CH	CF ₃	O	OCHF ₂	
744	CH	CF ₃	O	OCH ₂ C ₆ H ₅	
745	CH	CF ₃	O	SCH ₃	
746	CH	CF ₃	O	SC ₆ H ₅	
747	CH	CF ₃	O	NH ₂	
748	CH	CF ₃	O	NHCH ₃	
749	CH	CF ₃	O	NHCH ₂ CH ₃	
750	CH	CF ₃	O	N(CH ₂ CH ₃) ₂	
751	CH	CF ₃	O	N(CH ₂ CN) ₂	
752	CH	CF ₃	O	N(CH ₃) ₂	
753	CH	CF ₃	O	NHCOCH ₃	
754	CH	CF ₃	O	NHCOCH ₂ CH ₃	
755	CH	CF ₃	O	OSO ₂ CH ₃	
756	CH	CF ₃	O	SOCH ₂ (4-Br)-C ₆ H ₄	
757	CH	CF ₃	O	N(CH ₃)COOCH ₂ C ₆ H ₅	
758	N	CF ₃	O	CH ₃	
759	N	CF ₃	O	CH ₂ CH ₃	

No.	X	Y	W	R ¹	m.p. [°C]
760	N	CF ₃	O	(CH ₂) ₂ CH ₃	
761	N	CF ₃	O	CH(CH ₃) ₂	
762	N	CF ₃	O	(CH ₂) ₃ CH ₃	
763	N	CF ₃	O	CH ₂ CH(CH ₃) ₂	
764	N	CF ₃	O	C(CH ₃) ₃	
765	N	CF ₃	O	CH ₂ C(CH ₃) ₃	
766	N	CF ₃	O	cyclo-C ₅ H ₉	
767	N	CF ₃	O	cyclo-C ₆ H ₁₁	
768	N	CF ₃	O	CH ₂ C=C(CH ₃) ₂	
769	N	CF ₃	O	CH ₂ CH ₂ C=CH ₂	
770	N	CF ₃	O	CH ₂ CH=CH ₂	
771	N	CF ₃	O	(CH ₂) ₅ CH=CH ₂	
772	N	CF ₃	O	CH ₂ C≡CH	
773	N	CF ₃	O	CH ₂ C≡CCH ₂ CH ₃	
774	N	CF ₃	O	CHFCF ₃	
775	N	CF ₃	O	COOCH ₂ CH ₃	
776	N	CF ₃	O	CH ₂ CH ₂ OH	
777	N	CF ₃	O	CH ₂ CH ₂ OCH ₃	
778	N	CF ₃	O	CH ₂ COOC(CH ₃) ₃	
779	N	CF ₃	O	CH ₂ SC ₆ H ₅	
780	N	CF ₃	O	CH ₂ CONHCH ₃	
781	N	CF ₃	O	CH ₂ CH(OH)CH ₂ OH	
782	N	CF ₃	O	CHO	
783	N	CF ₃	O	COCH ₃	
784	N	CF ₃	O	CH ₂ OC ₆ H ₅	
785	N	CF ₃	O	COC ₆ H ₅	
786	N	CF ₃	O	CF ₂ CH ₃	
787	N	CF ₃	O	CH ₂ CN	

No.	X	Y	W	R ¹	m.p. [°C]
788	N	CF ₃	O	CH ₂ CH ₂ CN	
789	N	CF ₃	O	CH=CF ₂	
790	N	CF ₃	O	2-Furyl	
791	N	CF ₃	O	OH	
792	N	CF ₃	O	OCH ₃	
793	N	CF ₃	O	OCH ₂ CH ₃	
794	N	CF ₃	O	OCHF ₂	
795	N	CF ₃	O	OCH ₂ C ₆ H ₅	
796	N	CF ₃	O	NH ₂	
797	N	CF ₃	O	NHCH ₃	
798	N	CF ₃	O	NHCH ₂ CH ₃	
799	N	CF ₃	O	N(CH ₂ CH ₃) ₂	
800	N	CF ₃	O	N(CH ₂ CN) ₂	
801	N	CF ₃	O	N(CH ₃) ₂	
802	N	CF ₃	O	NHCOCH ₃	
803	N	CF ₃	O	NHCOCH ₂ CH ₃	
804	N	CF ₃	O	OSO ₂ CH ₃	
805	CH	CF ₃	S	CH ₃	
806	CH	CF ₃	S	CH ₂ CH ₃	
807	CH	CF ₃	S	(CH ₂) ₂ CH ₃	
808	CH	CF ₃	S	CHO	
809	CH	CF ₃	S	CHFCH ₃	
810	CH	CF ₃	S	CH ₂ C≡CH	
811	CH	CF ₃	S	COOCH ₂ CH ₃	
812	CH	CF ₃	S	CH ₂ COOC(CH ₃) ₃	
813	CH	CF ₃	S	CH ₂ CN	
814	N	CF ₃	S	CH ₃	
815	N	CF ₃	S	CH ₂ CH ₃	

No.	X	Y	W	R ¹	m.p. [°C]
816	N	CF ₃	S	(CH ₂) ₂ CH ₃	
817	N	CF ₃	S	CHF ₂ CF ₃	
818	N	CF ₃	S	CH ₂ CH ₂ OH	
819	N	CF ₃	S	CH ₂ COOC(CH ₃) ₃	
820	N	CH ₂ CH ₂ Cl	O	CH ₂ CH ₃	
821	N	CH ₂ CH ₂ Cl	O	NH ₂	
822	N	CH ₂ Cl	O	CH ₃	
823	CH	CHF ₂	O	CH ₃	
824	CH	CHF ₂	O	CH ₂ CH ₃	
825	CH	CHF ₂	O	(CH ₂) ₂ CH ₃	
826	CH	CHF ₂	O	CH ₂ C=CH ₂	
827	CH	CHF ₂	O	C(CH ₃)=CH ₂	
828	CH	CHF ₂	O	COOCH ₂ CH ₃	
829	CH	CHF ₂	O	CH ₂ CONHCH ₃	
830	CH	CHF ₂	O	CF ₂ CH ₃	
831	CH	CHF ₂	O	CHO	
832	CH	CHF ₂	O	NH ₂	
833	CH	CHF ₂	O	NHCOCH ₃	
834	N	CHF ₂	O	CH ₃	
835	N	CHF ₂	O	CH ₂ CH ₃	
836	N	CHF ₂	O	CH(CH ₃)(CH ₂) ₄ CH ₃	
837	N	CHF ₂	O	CH ₂ CH=CH ₂	
838	N	CHF ₂	O	COOCH ₂ CH ₃	
839	N	CHF ₂	O	NH ₂	

Table 3



No.	X	Y	m	V	R'	m.p. [°C]
840	N	(CF ₂) ₃ CHF ₂	0	O	CH ₃	
841	N	(CF ₂) ₂ CF ₃	0	O	CH ₂ CH ₃	
842	N	(CF ₂) ₂ CF ₃	0	O	COOCH ₂ CH ₃	
843	N	(CF ₂) ₂ CF ₃	0	O	SH	
844	N	(CF ₂) ₂ CF ₃	0	O	SCH ₃	
845	N	(CF ₂) ₂ CF ₃	0	O	SCH ₂ C≡CH	
846	N	CF ₂ CF ₃	0	O	CH ₃	
847	N	CF ₂ CF ₃	0	O	CH ₂ CH ₃	
848	N	CF ₃	0	O	CH ₃	
849	N	CF ₃	0	O	CH ₂ CH ₃	
850	N	CF ₃	0	O	(CH ₂) ₂ CH ₃	
851	N	CF ₃	0	O	CH(CH ₃) ₂	
852	N	CF ₃	0	O	(CH ₂) ₃ CH ₃	
853	N	CF ₃	0	O	CH ₂ CH(CH ₃) ₂	
854	N	CF ₃	0	O	C(CH ₃) ₃	
855	N	CF ₃	0	O	CH ₂ C(CH ₃) ₃	
856	N	CF ₃	0	O	Cyclo-C ₆ H ₉	
857	N	CF ₃	0	O	Cyclo-C ₆ H ₁₁	
858	N	CF ₃	0	O	CH ₂ CH=C(CH ₃) ₂	
859	N	CF ₃	0	O	CH ₂ CH ₂ CH=CH ₂	

No.	X	Y	m	V	R ¹	m.p. [°C]
860	N	CF ₃	0	O	CH ₂ CH=CH ₂	
861	N	CF ₃	0	O	(CH ₂) ₅ CH=CH ₂	
862	N	CF ₃	0	O	CH ₂ C≡CH	
863	N	CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₃	
864	N	CF ₃	0	O	CHFCF ₃	
865	N	CF ₃	0	O	COOCH ₂ CH ₃	
866	N	CF ₃	0	O	CH ₂ CH ₂ OH	
867	N	CF ₃	0	O	CH ₂ CH ₂ OCH ₃	
868	N	CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	
869	N	CF ₃	0	O	CH ₂ SPh	
870	N	CF ₃	0	O	CH ₂ CONHCH ₃	
871	N	CF ₃	0	O	CH ₂ CH(OH)CH ₂ OH	
872	N	CF ₃	0	O	CHO	
873	N	CF ₃	0	O	COCH ₃	
874	N	CF ₃	0	O	CH ₂ OC ₆ H ₅	
875	N	CF ₃	0	O	COPh	
876	N	CF ₃	0	O	CF ₂ CH ₃	
877	N	CF ₃	0	O	CH ₂ CN	
878	N	CF ₃	0	O	CH ₂ CH ₂ CN	
879	N	CF ₃	0	O	CH=CF ₂	
880	N	CF ₃	0	O	2-Furyl	
881	N	CF ₃	0	O	OH	
882	N	CF ₃	0	O	OCH ₃	
883	N	CF ₃	0	O	OCH ₂ CH ₃	
884	N	CF ₃	0	O	OCHF ₂	
885	N	CF ₃	0	O	OCH ₂ Ph	
886	N	CF ₃	0	O	NH ₂	
887	N	CF ₃	0	O	NHCH ₃	

No.	X	Y	m	V	R ¹	m.p. [°C]
888	N	CF ₃	0	O	NHCH ₂ CH ₃	
889	N	CF ₃	0	O	N(CH ₂ CH ₃) ₂	
890	N	CF ₃	0	O	N(CH ₂ CN) ₂	
891	N	CF ₃	0	O	N(CH ₃) ₂	
892	N	CF ₃	0	O	NHCOCH ₃	
893	N	CF ₃	0	O	NHCOCH ₂ CH ₃	
894	N	CF ₃	0	O	OSO ₂ CH ₃	
895	N	CH ₂ CH ₂ Cl	0	O	CH ₂ CH ₃	
896	N	CH ₂ CH ₂ Cl	0	O	NH ₂	
897	N	CH ₂ Cl	0	O	CH ₃	
898	N	CHF ₂	0	O	CH ₃	
899	N	CHF ₂	0	O	CH ₂ CH ₃	
900	N	CHF ₂	0	O	CH(CH ₃)(CH ₂) ₄ CH ₃	
901	N	CHF ₂	0	O	CH ₂ CH=CH ₂	
902	N	CHF ₂	0	O	COOCH ₂ CH ₃	
903	N	CHF ₂	0	O	NH ₂	
904	CH	CF ₃	0	O	CH ₃	60-61
905	CH	CF ₃	1	O	CH ₃	
906	CH	CF ₃	0	O	CH ₂ CH ₃	oil
907	CH	CF ₃	1	O	CH ₂ CH ₃	oil
908	CH	CF ₃	0	O	(CH ₂) ₂ CH ₃	oil
909	CH	CF ₃	1	O	(CH ₂) ₂ CH ₃	oil
910	CH	CF ₃	0	O	CH(CH ₃) ₂	
911	CH	CF ₃	1	O	CH(CH ₃) ₂	
912	CH	CF ₃	0	O	(CH ₂) ₃ CH ₃	
913	CH	CF ₃	1	O	(CH ₂) ₃ CH ₃	
914	CH	CF ₃	0	O	CH(CH ₃)CH ₂ CH ₃	
915	CH	CF ₃	1	O	CH(CH ₃)CH ₂ CH ₃	


No.	X	Y	m	V	R ¹	m.p. [°C]
916	CH	CF ₃	0	O	CH ₂ CH(CH ₃) ₂	
917	CH	CF ₃	1	O	CH ₂ CH(CH ₃) ₂	
918	CH	CF ₃	0	O	C(CH ₃) ₃	
919	CH	CF ₃	1	O	C(CH ₃) ₃	
920	CH	CF ₃	0	O	(CH ₂) ₄ CH ₃	
921	CH	CF ₃	1	O	(CH ₂) ₄ CH ₃	
922	CH	CF ₃	0	O	CH(CH ₃)(CH ₂) ₂ CH ₃	
923	CH	CF ₃	0	O	(CH ₂) ₂ CH(CH ₃) ₂	
924	CH	CF ₃	0	O	CH ₂ C(CH ₃) ₃	
925	CH	CF ₃	0	O	cyclo-C ₅ H ₉	
926	CH	CF ₃	0	O	cyclo-C ₆ H ₁₁	
927	CH	CF ₃	0	O	CH ₂ (3-Thienyl)	oil
928	CH	CF ₃	0	O	CHO	
929	CH	CF ₃	0	O	CH=CH ₂	
930	CH	CF ₃	0	O	CH ₂ Ph	61-63
931	CH	CF ₃	0	O	CH ₂ CH=C(CH ₃) ₂	
932	CH	CF ₃	0	O	CH ₂ CH=CH ₂	
933	CH	CF ₃	0	O	C(CH ₃)=CH ₂	
934	CH	CF ₃	0	O	(CH ₂) ₅ C=CH ₂	
935	CH	CF ₃	0	O	C(=CHCH ₃)CH ₃	
936	CH	CF ₃	0	O	CH ₂ C≡CH	
937	CH	CF ₃	0	O	CH ₂ CH ₂ C≡CH ₂	
938	CH	CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₃	
939	CH	CF ₃	0	O	(CH ₂) ₄ C≡CH	
940	CH	CF ₃	0	O	CHFCF ₃	
941	CH	CF ₃	0	O	COOCH ₂ CH ₃	
942	CH	CF ₃	0	O	CH ₂ CH ₂ OH	
943	CH	CF ₃	0	O	CH ₂ CH ₂ OCH ₃	

No.	X	Y	m	V	R ¹	m.p. [°C]
944	CH	CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	
945	CH	CF ₃	0	O	CH ₂ SPh	
946	CH	CF ₃	0	O	CH ₂ CONHCH ₃	
947	CH	CF ₃	0	O	CH ₂ CH(OH)CH ₂ OH	
948	CH	CF ₃	0	O	CH ₂ COCH ₃	
949	CH	CF ₃	0	O	COCH ₃	
950	CH	CF ₃	0	O	CH ₂ Oph	
951	CH	CF ₃	0	O	COPh	
952	CH	CF ₃	0	O	CF ₂ CH ₃	
953	CH	CF ₃	0	O	CH ₂ CN	oil
954	CH	CF ₃	0	O	CH ₂ CH(-O-)CH ₂	
955	CH	CF ₃	0	O	CH ₂ (4-OCH ₃)Ph	
956	CH	CF ₃	0	O	CH ₂ CH(OH)CH ₂ SPh	
957	CH	CF ₃	0	O	CH=CF ₂	
958	CH	CF ₃	0	O	CCl=CHCl	
959	CH	CF ₃	0	O	Ph	120-121
960	CH	CF ₃	0	O	2-Thienyl	87-89
961	CH	CF ₃	0	O	Oph	
962	CH	CF ₃	0	O	OH	
963	CH	CF ₃	0	O	OCH ₃	
964	CH	CF ₃	0	O	OCH ₂ CH ₃	
965	CH	CF ₃	0	O	OCHF ₂	
966	CH	CF ₃	0	O	OCH ₂ Ph	
967	CH	CF ₃	0	O	SCH ₃	
968	CH	CF ₃	0	O	SPh	
969	CH	CF ₃	0	O	NH ₂	190-191
970	CH	CF ₃	0	O	NHCH ₃	
971	CH	CF ₃	0	O	NHCH ₂ CH ₃	

No.	X	Y	m	V	R ¹	m.p. [°C]
972	CH	CF ₃	0	O	N(CH ₂ CH ₃) ₂	
973	CH	CF ₃	0	O	N(CH ₂ CN) ₂	
974	CH	CF ₃	0	O	N(CH ₃) ₂	
975	CH	CF ₃	0	O	NHCOCH ₃	
976	CH	CF ₃	0	O	NHCOCH ₂ CH ₃	
977	CH	CF ₃	0	O	OSO ₂ CH ₃	
978	CH	CF ₃	0	O	SOCH ₂ (4-Br)-C ₆ H ₄	
979	CH	CF ₃	0	O	N(CH ₃)COOCH ₂ Ph	
980	CH	CF ₃	0	NCH ₃	CH ₃	
981	CH	CF ₃	0	NCH ₂ CH ₃	CH ₃	
982	CH	CF ₃	0	NCH ₂ CH ₃	CH ₂ CH ₃	
983	CH	CF ₃	0	NCH ₂ CN	CH ₂ CH ₃	
984	CH	CF ₃	0	NCH ₂ OCH ₃	NHCH ₃	
985	CH	CF ₃	0	NCH ₂ OCH ₂ CH ₃	CN	
986	CH	CF ₃	0	NCH ₂ CH=CH ₂	CH ₃	
987	CH	CF ₃	0	NCH ₂ CH=CF ₂	SCH ₃	
988	CH	CF ₃	0	NCH ₂ OCH ₃	SCH ₂ CH ₃	
989	CH	CF ₃	0	NCH ₂ OCH ₃	SCH ₂ Ph	
990	CH	CHF ₂	0	O	CH ₃	
991	CH	CHF ₂	0	O	CH ₂ CH ₃	
992	CH	CHF ₂	0	O	(CH ₂) ₂ CH ₃	
993	CH	CHF ₂	0	O	CH ₂ CH=CH ₂	
994	CH	CHF ₂	0	O	C(CH ₃)=CH ₂	
995	CH	CHF ₂	0	O	COOCH ₂ CH ₃	
996	CH	CHF ₂	0	O	CH ₂ CONHCH ₃	
997	CH	CHF ₂	0	O	CF ₂ CH ₃	
998	CH	CHF ₂	0	O	CHO	
999	CH	CHF ₂	0	O	NH ₂	

No.	X	Y	m	V	R ¹	m.p. [°C]
1000	CH	CHF ₂	0	O	NHCOCH ₃	
1001	N	CF ₂ CF ₃	0	S	CH ₃	
1002	N	CF ₂ CF ₃	0	S	CH ₂ CH ₃	
1003	N	CF ₂ CF ₃	0	S	(CH ₂) ₂ CH ₃	
1004	N	CF ₃	0	S	CH ₃	
1005	N	CF ₃	0	S	CH ₂ CH ₃	
1006	N	CF ₃	0	S	(CH ₂) ₂ CH ₃	
1007	N	CF ₃	0	S	CHFCF ₃	
1008	N	CF ₃	0	S	CH ₂ CH ₂ OH	
1009	N	CF ₃	0	S	CH ₂ COOC(CH ₃) ₃	
1010	CH	CF ₃	0	S	CH ₃	
1011	CH	CF ₃	0	S	CH ₂ CH ₃	
1012	CH	CF ₃	0	S	(CH ₂) ₂ CH ₃	
1013	CH	CF ₃	0	S	CHO	
1014	CH	CF ₃	0	S	CHFCF ₃	
1015	CH	CF ₃	0	S	CH ₂ C≡CH	
1016	CH	CF ₃	0	S	COOCH ₂ CH ₃	
1017	CH	CF ₃	0	S	CH ₂ COOC(CH ₃) ₃	
1018	CH	CF ₃	0	S	CH ₂ CN	

Table 4



No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1019	N	(CF ₂) ₃ CHF ₂	0	S	H	CH ₂ CH ₃	
1020	N	CF ₂ CF ₂ CF ₃	0	S	H	CH ₂ CH ₃	
1021	N	CF ₂ CF ₃	0	S	H	CH ₂ CH ₃	
1022	N	CH ₂ CH ₂ Cl	0	S	H	CH ₂ CH ₃	
1023	N	CH ₂ Cl	0	S	H	CH ₂ CH ₃	
1024	N	CF ₃	0	S	CH ₂ CH ₃	CH ₂ CH ₃	
1025	N	CF ₃	0	S	(CH ₂) ₂ CH ₃	H	
1026	N	CF ₃	0	S	CH(CH ₃) ₂	H	
1027	N	CF ₃	0	S	CH ₂ CH(CH ₃) ₂	H	
1028	N	CF ₃	0	S	C(CH ₃) ₃	H	
1029	CH	CF ₃	0	S	H	CH ₃	oil
1030	CH	CF ₃	0	S	H	CH ₂ CH ₃	oil
1031	CH	CF ₃	0	S	H	C(CH ₃) ₃	oil
1032	CH	CF ₃	0	S	CH ₂ CH ₃	COOCH ₂ CH ₃	
1033	CH	CF ₃	0	S	(CH ₂) ₂ CH ₃	COOCH ₂ CH ₃	
1034	CH	CF ₃	0	S	CH(CH ₃) ₂	COOCH ₂ CH ₃	
1035	CH	CF ₃	0	S	CH(CH ₃) ₂	CONHCH ₂ CH ₃	
1036	CH	CF ₃	0	S	CH(CH ₃) ₂	CONHCH ₂ CH ₃	
1037	CH	CF ₃	0	S	CH(CH ₃) ₂	CON(CH ₂ CH ₃) ₂	
1038	CH	CF ₃	0	S	CH(CH ₃) ₂	CONH-cyclo-C ₃ H ₇	

No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1039	CH	CF ₃	0	S	C(CH ₃) ₃	COOCH ₂ CH ₃	
1040	CH	CF ₃	0	S	H	CONHCH ₂ CH ₃	
1041	CH	CF ₃	0	S	H	CON(CH ₂ CH ₃) ₂	
1042	CH	CF ₃	0	S	H	COOCH ₂ CH ₃	oil
1043	CH	CF ₃	0	S	H	CH ₂ COOCH ₂ CH ₃	oil
1044	CH	CF ₃	0	S	H	CH ₂ CHO	
1045	CH	CF ₃	0	S	H	CH ₂ OCH ₃	
1046	CH	CF ₃	0	S	H	CH ₂ OCH ₂ Ph	
1047	CH	CF ₃	0	S	H	H	
1048	CH	CF ₃	0	S	cyclo-C ₅ H ₉	H	
1049	CH	CF ₃	0	S	CON(CH ₃) ₂	CH ₃	oil
1050	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ OH	
1051	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ OCH ₃	
1052	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ OCH ₂ Ph	
1053	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ SPh	
1054	CH	CF ₃	0	S	CH ₃	CH ₃	oil
1055	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ CHO	
1055	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ CHNPh	
1057	CH	CF ₃	0	S	CH ₃	CH ₂ CH ₂ CONH ₂	
1058	CH	CF ₃	0	S	H	(4-CF ₃ O)C ₆ H ₄	120-121
1059	CH	CF ₃	0	S	CH ₂ C≡CH	H	
1060	CH	CF ₃	0	S	CH ₂ CH ₂ C≡CH	H	
1061	CH	CF ₃	0	S	CH ₂ C≡CCH ₂ CH ₃	H	
1062	CH	CF ₃	0	S	CH ₂ CH=C(CH ₃) ₂	H	
1063	CH	CF ₃	0	S	CH ₂ CH ₂ CH=CH ₂	H	
1064	CH	CF ₃	0	S	CH ₂ CH=CH ₂	H	
1065	CH	CF ₃	0	S	C(CH ₃)=CH ₂	H	
1066	CH	CF ₃	0	S	CHFCF ₃	H	

No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1067	CH	CF ₃	0	S	COOCH ₂ CH ₃	H	
1068	CH	CF ₃	0	S	CH ₂ CH ₂ OH	H	
1069	CH	CF ₃	0	S	CH ₂ CH ₂ OCH ₃	H	
1070	CH	CF ₃	0	S	CH ₂ COOC(CH ₃) ₃	H	
1071	CH	CF ₃	0	S	CH ₂ COCH ₃	H	
1072	CH	CF ₃	0	S	COCH ₃	H	
1073	CH	CF ₃	0	S	CH ₂ Oph	H	
1074	CH	CF ₃	0	S	COPh	H	
1075	CH	CF ₃	0	S	CO(4-Cl)-C ₆ H ₄	H	
1076	CH	CF ₃	0	S	CF ₂ CH ₃	H	
1077	CH	CF ₃	0	S	CH ₂ CN	H	
1078	CH	CF ₃	0	S	CH ₂ CH ₂ CN	H	
1079	N	CF ₃	0	S	H	H	
1080	N	CF ₃	0	S	H	CH ₂ CH ₂ CN	
1081	N	CF ₃	0	S	H	CH ₂ CO ₂ C(CH ₃) ₃	
1082	N	CF ₃	0	S	H	CH ₂ CHO	
1083	N	CF ₃	0	S	H	CH ₂ CH ₂ OH	
1084	N	CF ₃	0	S	H	CH ₂ CH ₂ OCH ₃	
1085	N	CF ₃	0	S	cyclo-C ₅ H ₉	H	
1086	N	CF ₃	0	S	CH ₃	COOCH ₂ CH ₃	
1087	N	CF ₃	0	S	CH ₃	COOH	
1088	N	CF ₃	0	S	CH ₃	CONH ₂	
1089	N	CF ₃	0	S	CH ₃	CONHCH ₂ CH ₃	
1090	N	CF ₃	0	S	CH ₃	CON(CH ₂ CH ₃) ₂	
1091	N	CF ₃	0	S	CH ₃	CONHCH ₃	
1092	N	CF ₃	0	S	CH ₃	CONHCH ₂ CN	
1093	N	CF ₃	0	S	CH ₃	CON(CH ₂ CN) ₂	
1094	N	CF ₃	0	S	CH ₃	CON(CH ₃) ₂	

No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1095	N	CF ₃	0	S	CH ₂ C≡CH	OCH ₂ CH ₃	
1096	N	CF ₃	0	S	CH ₂ CH ₂ C≡CH	OCH ₂ CH ₃	
1097	N	CF ₃	0	S	CH ₂ C≡CCH ₂ CH ₃	OCH ₂ CH ₃	
1098	N	CF ₃	0	S	CH ₂ CH=C(CH ₃) ₂	OCH ₂ CH ₃	
1099	N	CF ₃	0	S	CH ₂ CH ₂ CH=CH ₂	OCH ₂ CH ₃	
1100	N	CF ₃	0	S	CH ₂ CH=CH ₂	OCH ₂ CH ₃	
1101	N	CF ₃	0	S	C(CH ₃)=CH ₂	OCH ₂ CH ₃	
1102	N	CF ₃	0	S	CHFCF ₃	OCH ₂ CH ₃	
1103	N	CF ₃	0	S	COOCH ₂ CH ₃	OCH ₂ CH ₃	
1104	N	CF ₃	0	S	CH ₂ CH ₂ OH	OCH ₂ CH ₃	
1105	N	CF ₃	0	S	CH ₂ CH ₂ OCH ₃	OCH ₂ CH ₃	
1106	N	CF ₃	0	S	CH ₂ COOC(CH ₃) ₃	OCH ₂ CH ₃	
1107	N	CF ₃	0	S	CH ₂ COCH ₃	H	
1108	N	CF ₃	0	S	COCH ₃	H	
1109	N	CF ₃	0	S	CH ₂ Oph	H	
1110	N	CF ₃	0	S	COPh	H	
1111	N	CF ₃	0	S	CO(4-Cl)-C ₆ H ₄	H	
1112	N	CF ₃	0	S	CF ₂ CH ₃	H	
1113	N	CF ₃	0	S	CH ₂ CN	H	
1114	N	CF ₃	0	S	CH ₂ CH ₂ CN	H	
1115	CH	CF ₃	0	O	CH ₂ CH ₃	CH ₂ CH ₃	
1116	CH	CF ₃	0	O	(CH ₂) ₂ CH ₃	H	
1117	CH	CF ₃	0	O	H	CH ₂ CH ₃	oil
1118	CH	CF ₃	0	O	CH(CH ₃) ₂	COOCH ₂ CH ₃	
1119	CH	CF ₃	0	O	CH(CH ₃) ₂	COOH	
1120	CH	CF ₃	0	O	CH(CH ₃) ₂	CONH ₂	
1121	CH	CF ₃	0	O	CH(CH ₃) ₂	CH ₃	
1122	CH	CF ₃	0	O	C(CH ₃) ₃	H	


No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1123	CH	CF ₃	0	O	H	CH ₃	
1124	CH	CF ₃	0	O	H	cyclo-C ₅ H ₉	
1125	CH	CF ₃	0	O	H	CH ₂ CH ₂ CH ₃	
1126	CH	CF ₃	0	O	H	Ph	103-104
1127	CH	CF ₃	0	O	H	2-Pyridyl	
1128	CH	CF ₃	0	O	H	2-Furyl	
1129	CH	CF ₃	0	O	cyclo-C ₅ H ₉	H	
1130	CH	CF ₃	0	O	CH ₃	COOCH ₂ CH ₃	
1131	CH	CF ₃	0	O	CH ₃	COOH	
1132	CH	CF ₃	0	O	CH ₃	CONH ₂	
1133	CH	CF ₃	0	O	CH ₃	CONHCH ₂ CH ₃	
1134	CH	CF ₃	0	O	CH ₃	CON(CH ₂ CH ₃) ₂	
1135	CH	CF ₃	0	O	CH ₃	CONHCH ₃	
1136	CH	CF ₃	0	O	CH ₃	CONHCH ₂ CN	
1137	CH	CF ₃	0	O	CH ₃	CON(CH ₂ CN) ₂	
1138	CH	CF ₃	0	O	CH ₃	CON(CH ₃) ₂	
1139	CH	CF ₃	0	O	CH ₂ C≡CH	H	
1140	CH	CF ₃	0	O	CH ₂ CH ₂ C≡CH	H	
1141	CH	CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₃	H	
1142	CH	CF ₃	0	O	CH ₂ CH=C(CH ₃) ₂	H	
1143	CH	CF ₃	0	O	CH ₂ CH ₂ C=CH	H	
1144	CH	CF ₃	0	O	CH ₂ CH=CH ₂	H	
1145	CH	CF ₃	0	O	C(CH ₃)=CH ₂	H	
1146	CH	CF ₃	0	O	CHFCF ₃	H	
1147	CH	CF ₃	0	O	COOCH ₂ CH ₃	H	
1148	CH	CF ₃	0	O	CH ₂ CH ₂ OH	H	
1149	CH	CF ₃	0	O	CH ₂ CH ₂ OCH ₃	H	
1150	CH	CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	H	

No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1151	CH	CF ₃	0	O	CH ₂ COCH ₃	H	
1152	CH	CF ₃	0	O	COCH ₃	H	
1153	CH	CF ₃	0	O	CH ₂ Oph	H	
1154	CH	CF ₃	0	O	COPh	H	
1155	CH	CF ₃	0	O	CO(4-Cl)-C ₆ H ₄	H	
1156	CH	CF ₃	0	O	CF ₂ CH ₃	H	
1157	CH	CF ₃	0	O	CH ₂ CN	H	
1158	CH	CF ₃	0	O	CH ₂ CH ₂ CN	H	
1159	N	CF ₃	0	O	CH ₂ CH ₃	CH ₂ CH ₃	
1160	N	CF ₃	0	O	(CH ₂) ₂ CH ₃	H	
1161	N	CF ₃	0	O	CH(CH ₃) ₂	CONH ₂	
1162	N	CF ₃	0	O	CH(CH ₃) ₂	CH ₃	
1163	N	CF ₃	0	O	C(CH ₃) ₃	H	
1164	N	CF ₃	0	O	H	CH ₃	
1165	N	CF ₃	0	O	H	CH ₂ CH ₃	
1166	N	CF ₃	0	O	H	CH ₂ CH ₂ CH ₃	
1167	N	CF ₃	0	O	H	Ph	
1168	N	CF ₃	0	O	H	2-Pyridyl	
1169	N	CF ₃	0	O	H	2-Furyl	
1170	N	CF ₃	0	O	cyclo-C ₅ H ₉	H	
1171	N	CF ₃	0	O	CH ₃	COOCH ₂ CH ₃	
1172	N	CF ₃	0	O	CH ₃	COOH	
1173	N	CF ₃	0	O	CH ₃	CONH ₂	
1174	N	CF ₃	0	O	CH ₃	CONHCH ₂ CH ₃	
1175	N	CF ₃	0	O	CH ₃	CON(CH ₂ CH ₃) ₂	
1176	N	CF ₃	0	O	CH ₃	CONHCH ₃	
1177	N	CF ₃	0	O	CH ₃	CONHCH ₂ CN	
1178	N	CF ₃	0	O	CH ₃	CON(CH ₂ CN) ₂	

No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1179	N	CF ₃	0	O	CH ₃	CON(CH ₃) ₂	
1180	N	CF ₃	0	O	CH ₂ C≡CH	H	
1181	N	CF ₃	0	O	CH ₂ CH ₂ C≡CH	H	
1182	N	CF ₃	0	O	CH ₂ C≡CCH ₂ CH ₃	H	
1183	N	CF ₃	0	O	CH ₂ CH=C(CH ₃) ₂	H	
1184	N	CF ₃	0	O	CH ₂ CH ₂ CH=CH ₂	H	
1185	N	CF ₃	0	O	CH ₂ CH=CH ₂	H	
1186	N	CF ₃	0	O	C(CH ₃)=CH ₂	H	
1187	N	CF ₃	0	O	CHFCF ₃	H	
1188	N	CF ₃	0	O	COOCH ₂ CH ₃	H	
1189	N	CF ₃	0	O	CH ₂ CH ₂ OH	H	
1190	N	CF ₃	0	O	CH ₂ CH ₂ OCH ₃	H	
1191	N	CF ₃	0	O	CH ₂ COOC(CH ₃) ₃	H	
1192	N	CF ₃	0	O	CH ₂ COCH ₃	H	
1193	N	CF ₃	0	O	COCH ₃	H	
1194	N	CF ₃	0	O	CH ₂ Oph	H	
1195	N	CF ₃	0	O	COPh	H	
1196	N	CF ₃	0	O	CO(4-Cl)-C ₆ H ₄	H	
1197	N	CF ₃	0	O	CF ₂ CH ₃	H	
1198	N	CF ₃	0	O	CH ₂ CN	H	
1199	N	CF ₃	0	O	CH ₂ CH ₂ CN	H	
1200	N	CF ₃	0	O	CH ₂ NHSO ₂ CH ₃	CH ₃	
1201	N	CF ₃	0	O	(CH ₂) ₂ NHSO ₂ CH ₃	CH ₃	
1202	N	CF ₃	0	O	CH ₂ NHSO ₂ CH ₂ CH ₃	CH ₃	
1203	N	CF ₃	0	O	H	CH ₂ NHSO ₂ CH ₂ Ph	
1204	CH	CF ₃	0	O	(CH ₂) ₄ NHSO ₂ CF ₃	CH ₃	
1205	CH	CF ₃	0	O	(CH ₂) ₂ S(CH ₂) ₂ CH ₃	CH ₂ CH ₂ CH ₃	
1206	CH	CF ₃	0	O	(CH ₂) ₄ S(CH ₂) ₄ OCH ₃	CH ₃	

No.	X	Y	m	V	R ²	R ³	m.p. [°C]
1207	CH	CF ₃	0	S	CH ₃	(CH ₂) ₂ S(CH ₂) ₂ CN	
1208	CH	CF ₃	0	S	CH ₂ NHSO ₂ CH ₂ CH ₃	CH ₃	
1209	CH	CF ₃	0	S	CH ₂ NHSO ₂ CH ₂ Ph	CH ₂ CH ₂ CH ₃	
1210	CH	CF ₃	0	S	(CH ₂) ₂ NHSO ₂ CH ₃	CF ₃	
1211	CH	CF ₃	0	S	H	CH ₂ NHSO ₂ CH ₃	
1212	CH	CF ₃	0	S	CH(CH ₃)CH ₂ NHPh	CF ₃	
1213	CH	CF ₃	0	S	(CH ₂) ₂ S(2-F)-C ₆ H ₄	CH ₂ CH ₂ CH ₃	
1214	CH	CF ₃	0	S	(CH ₂) ₆ NHCH ₂) ₆ OCH ₃	CF ₃	
1215	CH	CF ₃	0	S	H	(CH ₂) ₂ NH-(2-F)-C ₆ H ₄	
1216	CH	CF ₃	0	S	(CH ₂) ₃ NHCH ₂ CN	H	
1217	CH	CF ₃	0	S	(CH ₂) ₂ O(3-Cl)-C ₆ H ₄	CH ₃	
1218	CH	CF ₃	0	S	CF ₃	(CH ₂) ₆ NHCH ₂ CF ₃	
1219	CH	CF ₃	0	S	CH ₃	(CH ₂) ₂ O(3-CH ₃)-C ₆ H ₄	
1220	CH	CF ₃	0	O	H	CH ₂ NHPh	
1221	CH	CF ₃	0	O	CH ₃	(CH ₂) ₄ S(2-Br)-C ₆ H ₄	
1222	CH	CF ₃	0	O	(CH ₂) ₆ NH(CH ₂) ₂ OCH ₃	CH ₃	
1223	CH	CF ₃	0	O	(CH ₂) ₂ NH(CH ₂) ₄ OCH ₃	H	
1224	CH	CF ₃	0	O	CF ₃	(CH ₂) ₃ NH-(4-CN)-C ₆ H ₄	
1225	CH	CF ₃	0	O	(CH ₂) ₄ NHCH ₂ CF ₃	CH ₃	
1226	CH	CF ₃	0	O	C ₂ F ₅	(CH ₂) ₂ O(3-CH ₃)-C ₆ H ₄	
1227	CH	CF ₃	0	O	(CH ₂) ₄ NHCH ₂ CN	H	
1228	CH	CF ₃	0	O	(CH ₂) ₃ O(4-Cl)-C ₆ H ₄	C ₂ F ₅	


Table 5



No.	X	Y	V	R ⁴	R ⁵	R ⁶	R ⁷	m.p. [°C]
1229	CH	CF ₃	O	H	H	H	H	oil
1230	CH	CF ₃	O	H	H	CH ₃	H	oil
1231	CH	CF ₃	O	H	H	CH ₂ CH ₃	H	oil
1232	CH	CF ₃	O	H	H	CH(CH ₃) ₂	H	
1233	CH	CF ₃	O	H	H	CH ₂ CH(CH ₃) ₂	H	
1234	CH	CF ₃	O	H	H	CH(CH ₃)CH ₂ CH ₃	H	
1235	CH	CF ₃	O	H	H	CH ₂ OH	H	
1236	CH	CF ₃	O	H	H	CH(OH)CH ₃	H	
1237	CH	CF ₃	O	H	H	CH ₂ SH	H	
1238	CH	CF ₃	O	H	H	CH ₂ CH ₂ SCH ₃	H	
1239	CH	CF ₃	O	H	H	(CH ₂) ₃ NH ₂	H	
1240	CH	CF ₃	O	H	H	(CH ₂) ₄ NH ₂	H	
1241	CH	CF ₃	O	H	H	CH=CH ₂	H	
1242	CH	CF ₃	O	H	H	(CH ₂) ₂ COOCH ₃	H	
1243	CH	CF ₃	O	H	H	(CH ₂) ₂ COOH	H	
1244	CH	CF ₃	O	H	H	(CH ₂) ₂ CONH ₂	H	
1245	CH	CF ₃	S	CH ₃	CH ₃	H	H	
1246	CH	CF ₃	O	H	H	CH ₃	CH ₃	oil
1247	CH	CF ₃	O	H	H	CH ₂ COOCH ₃	H	
1248	CH	CF ₃	O	H	H	CH ₂ COOH	H	

No.	X	Y	V	R ⁴	R ⁵	R ⁶	R ⁷	m.p. [°C]
1249	CH	CF ₃	O	H	H	CH ₂ CONH ₂	H	
1250	CH	CF ₃	O	H	H	CH ₂ Ph	H	
1251	CH	CF ₃	O	H	H	CH ₂ -(4-OH)-C ₆ H ₄	H	
1252	CH	CF ₃	O	H	H	CH ₂ -(3-indolyl)	H	
1253	CH	CF ₃	O	CH ₃	CH ₃	H	H	oil
1254	CH	CF ₃	O	CH ₃	H	H	H	oil
1255	CH	CF ₃	O	CH ₃	H	H	Ph	
1256	CH	CF ₃	O	H		(CH ₂) ₄	H	
1257	CH	CF ₃	NH	H		(CH ₂) ₄	H	
1258	CH	CF ₃	NCH ₃	H		(CH ₂) ₄	H	
1259	CH	CF ₃	NCH ₂ C ₆ H ₄	H		(CH ₂) ₄	H	
1260	CH	CF ₃	NCH(CH ₃) ₂	H		(CH ₂) ₄	H	
1261	CH	CF ₃	O	Ph	H	Ph	H	
1262	CH	CF ₃	NH	Ph	H	Ph	H	
1263	CH	CF ₃	NCH ₃	Ph	H	Ph	H	
1264	CH	CF ₃	NCH ₂ C ₆ H ₄	Ph	H	Ph	H	
1265	N	CF ₃	O	H	H	CH ₂ CH ₃	H	oil
1266	N	CF ₃	O	H	H	CH(CH ₃) ₂	H	
1267	N	CF ₃	O	H	H	CH ₂ CH(CH ₃) ₂	H	
1268	N	CF ₃	O	H	H	CH ₂ COOH	H	
1269	N	CF ₃	O	H	H	CH ₂ COOCH ₃	H	
1270	N	CF ₃	O	H	H	CH ₂ CONH ₂	H	
1271	N	CF ₃	O	CH ₃	CH ₃	H	H	
1272	N	CF ₃	O	H	(CH ₂) ₄		H	
1273	N	CF ₃	O	H	H	CH ₂ CH ₂ SCH ₃	H	
1274	CH	CF ₃	S	H	H	H	H	oil

Table 6



No.	X	Y	R ⁸	R ¹	m.p. [°C]
1275	CH	CF ₃	CH ₃	SH	209-210
1276	CH	CF ₃	CH ₃	SCH ₃	
1277	CH	CF ₃	CH ₃	SCH ₂ CH ₃	
1278	CH	CF ₃	CH ₃	S(CH ₂) ₂ CH ₃	
1279	CH	CF ₃	CH ₃	SCH(CH ₃) ₂	
1280	CH	CF ₃	CH ₃	SPh	
1281	CH	CF ₃	CH ₃	S(CH ₂) ₃ CH ₃	
1282	CH	CF ₃	CH ₃	SCH(CH ₃)CH ₂ CH ₃	
1283	CH	CF ₃	CH ₃	SCH ₂ CH(CH ₃) ₂	
1284	CH	CF ₃	CH ₃	OH	119-120
1285	CH	CF ₃	CH ₃	OCH ₃	
1286	CH	CF ₃	CH ₃	OCH ₂ CH ₃	
1287	CH	CF ₃	CH ₃	OCHF ₂	
1288	CH	CF ₃	CH ₃	OCH ₂ Ph	
1289	CH	CF ₃	CH ₃	OCONHPh	
1290	CH	CF ₃	CH ₃	OCONH-(4-F)-C ₆ H ₄	
1291	CH	CF ₃	CH ₃	OCONH-(3,5-di-Cl)-C ₆ H ₃	
1292	CH	CF ₃	CH ₂ CN	OCH ₃	
1293	CH	CF ₃	CH ₂ CN	OCH ₂ CH ₃	
1294	CH	CF ₃	CH ₂ CN	OCHF ₂	

No.	X	Y	R ⁸	R ¹	m.p. [°C]
1295	CH	CF ₃	CH ₂ CN	OCH ₂ Ph	
1296	CH	CF ₃	CH ₂ CN	OCONHPh	
1297	CH	CF ₃	CH ₂ CN	OCONH-(4-F)-C ₆ H ₄	
1298	CH	CF ₃	CH ₂ OCH ₂ CH ₃	OCH ₃	
1299	CH	CF ₃	CH ₂ OCH ₂ CH ₃	OCH ₂ CH ₃	
1300	CH	CF ₃	CH ₂ OCH ₂ CH ₃	OCHF ₂	
1301	CH	CF ₃	CH ₂ OCH ₂ CH ₃	OCH ₂ Ph	
1302	CH	CF ₃	CH ₂ OCH ₂ CH ₃	OCONHPh	
1303	CH	CF ₃	H	CH ₃	203-204
1304	CH	CF ₃	H	CH ₂ CH ₃	134-135
1305	CH	CF ₃	H	(CH ₂) ₂ CH ₃	
1306	CH	CF ₃	H	CH(CH ₃) ₂	
1307	CH	CF ₃	H	Cyclo-C ₃ H ₅	
1308	CH	CF ₃	H	(CH ₂) ₃ CH ₃	
1309	CH	CF ₃	H	CH(CH ₃)CH ₂ CH ₃	
1310	CH	CF ₃	H	CH ₂ CH(CH ₃) ₂	
1311	CH	CF ₃	H	CH=CH ₂	
1312	CH	CF ₃	H	CH ₂ CH=C(CH ₃) ₂	
1313	CH	CF ₃	H	CH ₂ CH ₂ CH=CH ₂	
1314	CH	CF ₃	H	CH ₂ CH=CH ₂	
1315	CH	CF ₃	H	C(CH ₃)=CH ₂	
1316	CH	CF ₃	H	CHFCF ₃	
1317	CH	CF ₃	H	COOCH ₂ CH ₃	
1318	CH	CF ₃	H	CH ₂ CH ₂ OH	
1319	CH	CF ₃	H	CH ₂ CH ₂ OCH ₃	
1320	CH	CF ₃	H	CH ₂ COOC(CH ₃) ₃	
1321	CH	CF ₃	CH ₃	CH ₂ COOC(CH ₃) ₃	
1322	CH	CF ₃	CH ₂ CN	CH ₂ COOC(CH ₃) ₃	

No.	X	Y	R ⁸	R ¹	m.p. [°C]
1323	CH	CF ₃	CH ₂ OCH ₂ CH ₃	CH ₂ COOC(CH ₃) ₃	
1324	CH	CF ₃	H	CH ₂ SPh	
1325	CH	CF ₃	H	CH ₂ CONHCH ₃	
1326	CH	CF ₃	H	CH ₂ COCH ₃	
1327	CH	CF ₃	H	COCH ₃	
1328	CH	CF ₃	H	CH ₂ Oph	
1329	CH	CF ₃	H	COPh	
1330	CH	CF ₃	H	CO(3-Cl)-C ₆ H ₄	
1331	CH	CF ₃	H	CF ₂ CH ₃	
1332	CH	CF ₃	H	CH ₂ CN	
1333	CH	CF ₃	H	CH ₂ CH ₂ CN	
1334	CH	CF ₃	H	CH ₂ CH(-O-)CH ₂	
1336	CH	CF ₃	H	CH ₂ (4-OCH ₃)Ph	
1337	N	CF ₃	CH ₃	SH	
1338	N	CF ₃	CH ₃	SCH ₃	
1339	N	CF ₃	CH ₃	SCH ₂ CH ₃	
1340	N	CF ₃	CH ₃	SPh	
1341	N	CF ₃	CH ₃	SCH ₂ CH(CH ₃) ₂	
1342	N	CF ₃	CH ₃	OH	
1343	N	CF ₃	CH ₃	OCH ₃	
1344	N	CF ₃	CH ₃	OCH ₂ CH ₃	
1345	N	CF ₃	CH ₃	OCH ₂ Ph	
1346	N	CF ₃	CH ₃	OCONHPh	
1347	N	CF ₃	CH ₂ CN	OCH ₃	
1348	N	CF ₃	CH ₂ CN	OCH ₂ CH ₃	
1349	N	CF ₃	CH ₂ CN	OCH ₂ Ph	
1350	N	CF ₃	CH ₂ CN	OCONHPh	
1351	N	CF ₃	CH ₂ OCH ₂ CH ₃	OCH ₃	

No.	X	Y	R ⁸	R ¹	m.p. [°C]
1352	N	CF ₃	CH ₂ OCH ₂ CH ₃	OCH ₂ Ph	
1353	N	CF ₃	CH ₂ OCH ₂ CH ₃	OCONHPh	
1354	N	CF ₃	H	CH ₃	
1355	N	CF ₃	H	CH ₂ CH ₃	
1356	N	CF ₃	H	(CH ₂) ₂ CH ₃	
1357	N	CF ₃	H	CH(CH ₃) ₂	
1358	N	CF ₃	H	(CH ₂) ₃ CH ₃	
1359	N	CF ₃	H	CH(CH ₃)CH ₂ CH ₃	
1360	N	CF ₃	H	CH ₂ CH(CH ₃) ₂	
1361	N	CF ₃	H	CH ₂ C=C(CH ₃) ₂	
1362	N	CF ₃	H	CH ₂ CH=CH ₂	
1363	N	CF ₃	H	C(CH ₃)H=CH ₂	
1364	N	CF ₃	H	COOCH ₂ CH ₃	
1365	N	CF ₃	H	CH ₂ CH ₂ OH	
1366	N	CF ₃	H	CH ₂ CH ₂ OCH ₃	
1367	N	CF ₃	H	CH ₂ COOC(CH ₃) ₃	
1368	N	CF ₃	H	CH ₂ SPh	
1369	N	CF ₃	H	CH ₂ CONHCH ₃	
1370	N	CF ₃	H	CH ₂ COCH ₃	
1371	N	CF ₃	H	COCH ₃	
1372	N	CF ₃	H	CH ₂ Oph	
1373	N	CF ₃	H	COPh	
1374	N	CF ₃	H	CH ₂ CN	
1375	N	CF ₃	H	CH ₂ CH ₂ CN	
1376	CH	CF ₃	CH ₃	CH ₂ CH ₃	oil

C. Biological examples

Example 1

5

- A Petri dish whose bottom is covered with filter paper and which contains about 5 ml of culture medium is prepared. Pieces of filter paper with about 30, 24-hour-old eggs of the American tobacco budworm (*Heliothis virescens*) are dipped into an aqueous solution of the formulated preparation to be examined for 5 seconds and subsequently placed in the Petri dish. A further 200 μ l of the aqueous solution are spread over the culture medium. The Petri dish is closed and then kept at about 25°C in a climatized chamber. After 6 days' storage, the effect of the preparation on the eggs and the larvae which may have hatched from these is determined.
- 15 At a concentration of 300 ppm (based on the content of active compound), the preparations of Example Nos. 79 and 88 effect a mortality of 90-100%.

Example 2

- 20 Germinated broad bean seeds (*Vicia faba*) with radicles are transferred into brown glass bottles filled with tap water and subsequently populated with approximately 100 black bean aphids (*Aphis fabae*) belegt. Plants and aphids are then dipped for 5 seconds into an aqueous solution of the formulated preparation to be examined. After the solution has dripped off,
- 25 plant and animals are kept in a climatized chamber (16 hours of light/day, 25°C, 40-60% relative atmospheric humidity. After 3 and 6 days' storage, the effect of the preparation on the aphids is determined. At a concentration of 300 ppm (based on the content of active compound), the preparations of Example Nos. 79, 78, 80, 81, 83, 84, 88, 133, 135, 136,
- 30 137, 138, 139, 1117, 1229, 1230, 1231, 1246 and 1254 effect a mortality of 90-100% among the aphids.

Example 3

The leaves of 12 rice plants having a stem length of 8 cm are dipped for 5 seconds into an aqueous solution of the formulated preparation to be examined. After the solution has dripped off, the rice plants treated in this manner are placed in a Petri dish and populated with approximately 20 larvae (L3 stage) of the rice leaf hopper species *Nilaparvata lugens*. The Petri dish is closed and stored in a climatized chamber (16 hours of light/day, 25°C, 40-60% relative atmospheric humidity). After 6 days' storage, the mortality among the leaf hopper larvae is determined. At a concentration of 300 ppm (based on the content of active compound), the preparations of Example Nos. 88, 139 and 927 effect a mortality of 90-100%.

15 Example 4


Germinated broad bean seeds (*Vicia faba*) with radicles are transferred into brown glass bottles filled with tap water. Four milliliters of an aqueous solution of the formulated preparation to be examined are pipetted into the brown glass bottle. The broad bean is subsequently heavily populated with approximately 100 black bean aphids (*Aphis fabae*). Plant and animals are then stored in a climatized chamber (16 hours of light/day, 25°C, 40-60% relative atmospheric humidity). After 3 and 6 days' storage, the root-systemic activity of the preparation on the aphids is determined. At a concentration of of 30 ppm (based on the content of active compound), the Preparations of Example Nos. 78, 79, 80, 81, 83, 84, 88, 133, 135, 136, 137, 138, 139, 187, 1117, 1229, 1230, 1231, 1246 and 1254 effect a mortality of 90-100% among the aphids by root-systemic action.



Claims

1. A 4-haloalkyl-3-heterocyclylpyridine or 4-haloalkyl-5-heterocyclylpyrimidine of the formula I if desired in the form of its salt

5



10


wherein

Y is halo-C₁-C₆-alkyl;

X is CH or N;

15 m is 0 or 1;

Q is a 5-membered heterocyclic group



in which

- 20 a) $X^1 = W$, $X^2 = NR^a$, $X^3 = CR^bR^1$ or
 b) $X^1 = NR^a$, $X^2 = CR^bR^1$, $X^3 = W$ or
 c) $X^1 = V$, $X^2 = CR^aR^1$, $X^3 = NR^b$ or
 d) $X^1 = V$, $X^2 = CR^aR^2$, $X^3 = CR^bR^3$ or
 e) $X^1 = V$, $X^2 = CR^4R^5$, $X^3 = CR^6R^7$ or
 25 f) $X^1 = NR^a$, $X^2 = CR^bR^1$, $X^3 = NR^8$;

 R^a and R^b together are a bondV is oxygen, sulfur or NR^9 ;

W is oxygen or sulfur;

This Sheet Has Been Replaced



R¹

is hydrogen,

(C₁-C₂₀)-alkyl, (C₂-C₂₀)-alkenyl, (C₂-C₂₀)-alkynyl, (C₃-C₈)-cycloalkyl,

(C₄-C₈)-cycloalkenyl, (C₆-C₈)-cycloalkynyl,

5 where the six last-mentioned radicals are optionally substituted by one or more radicals from the group

halogen, cyano, nitro, hydroxyl, -C(=W)R¹⁰, -C(=NOR¹⁰)R¹⁰,
 -C(=NNR¹⁰₂)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -
 OC(=W)R¹⁰, -OC(=W)OR¹⁰, -NR¹⁰C(=W)R¹⁰, -
 10 N[C(=W)R¹⁰]₂, -NR¹⁰C(=W)OR¹⁰, -C(=W)NR¹⁰-NR¹⁰₂,
 -C(=W)NR¹⁰-NR¹⁰[C(=W)R¹⁰], -NR¹⁰-C(=W)NR¹⁰₂,
 -NR¹⁰-NR¹⁰C(=W)R¹⁰, -NR¹⁰-N[C(=W)R¹⁰]₂,
 -N[(C=W)R¹⁰]-NR¹⁰₂,
 -NR¹⁰-NR¹⁰[(C=W)R¹⁰], -NR¹⁰-NR¹⁰[(C=W)WR¹⁰],
 15 -NR¹⁰-R¹⁰[(C=W)NR¹⁰₂], -NR¹⁰(C=NR¹⁰)R¹⁰,
 -NR¹⁰(C=NR¹⁰)NR¹⁰₂,
 -O-NR¹⁰₂, -O-NR¹⁰(C=W)R¹⁰, -SO₂NR¹⁰₂, -NR¹⁰SO₂R¹⁰,
 -SO₂OR¹⁰, -OSO₂R¹⁰, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃,
 -SeR¹⁰, -PR¹⁰₂, -P(=W)R¹⁰₂,
 20 -SOR¹⁰, -SO₂R¹⁰, -PW₂R¹⁰₂, -PW₃R¹⁰₂, aryl and
 heterocyclyl,


the two last-mentioned radicals optionally being substituted by one or more radicals from the group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl,

25 (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₆-C₈)-cycloalkynyl, (C₁-C₆)-haloalkyl, (C₂-C₆)-haloalkenyl, (C₂-C₆)-haloalkynyl, halogen, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃,



126




aryl,

which is optionally substituted by one or more radicals from the group


5

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl, where these six abovementioned radicals are optionally substituted by one or more radicals from the group

10



15




heterocyclyl,

20

which is optionally substituted by one or more radicals from the group

25

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl, where the six abovementioned radicals are optionally substituted by one or more radicals from the group





127

- $-\text{OC}(=\text{W})\text{R}^{10}$, $-\text{OC}(=\text{W})\text{OR}^{10}$, $-\text{OR}^{10}$, $-\text{NR}^{10}_2$, $-\text{SR}^{10}$,
 $-\text{SOR}^{10}$ and $-\text{SO}_2\text{R}^{10}$,
halogen, cyano, nitro, $-\text{C}(=\text{W})\text{R}^{10}$, $-\text{C}(=\text{W})\text{OR}^{10}$,
 $-\text{C}(=\text{W})\text{NR}^{10}_2$, $-\text{OC}(=\text{W})\text{R}^{10}$, $-\text{OR}^{10}$, $-\text{NR}^{10}_2$, $-\text{SR}^{10}$, $-\text{SOR}^{10}$
5 and $-\text{SO}_2\text{R}^{10}$,
 $-\text{OR}^{10}$, $-\text{NR}^{10}_2$, $-\text{SR}^{10}$, $-\text{SOR}^{10}$, $-\text{SO}_2\text{R}^{10}$, $-\text{C}(=\text{W})\text{R}^{10}$,
 $-\text{C}(=\text{NOR}^{10})\text{R}^{10}$, $-\text{C}(=\text{NNR}^{10}_2)\text{R}^{10}$, $-\text{C}(=\text{W})\text{OR}^{10}$,
 $-\text{C}(=\text{W})\text{NR}^{10}_2$, $-\text{OC}(=\text{W})\text{R}^{10}$, $-\text{OC}(=\text{W})\text{OR}^{10}$, $-\text{NR}^{10}\text{C}(=\text{W})\text{R}^{10}$,
 $-\text{N}[\text{C}(=\text{W})\text{R}^{10}]_2$, $-\text{NR}^{10}\text{C}(=\text{W})\text{OR}^{10}$, $-\text{C}(=\text{W})\text{NR}^{10}-\text{NR}^{10}_2$,
10 $-\text{C}(=\text{W})\text{NR}^{10}-\text{NR}^{10}[\text{C}(=\text{W})\text{R}^{10}]$, $-\text{NR}^{10}-\text{C}(=\text{W})\text{NR}^{10}_2$,
 $-\text{NR}^{10}-\text{NR}^{10}\text{C}(=\text{W})\text{R}^{10}$, $-\text{NR}^{10}-\text{NC}(=\text{W})\text{R}^{10}_2$, $-\text{N}(\text{C}=\text{W})\text{R}^{10}$
 $-\text{NR}^{10}_2$, $-\text{NR}^{10}-\text{NR}^{10}[(\text{C}=\text{W})\text{R}^{10}]$, $-\text{NR}^{10}-\text{NR}^{10}[(\text{C}=\text{W})\text{WR}^{10}]$,
 $-\text{NR}^{10}-\text{NR}^{10}[(\text{C}=\text{W})\text{NR}^{10}_2]$, $-\text{NR}^{10}(\text{C}=\text{NR}^{10})\text{R}^{10}$,
 $-\text{NR}^{10}(\text{C}=\text{NR}^{10})\text{NR}^{10}_2$, $-\text{O}-\text{NR}^{10}_2$, $-\text{O}-\text{NR}^{10}(\text{C}=\text{W})\text{R}^{10}$,
15 $-\text{SO}_2\text{NR}^{10}_2$, $-\text{NR}^{10}\text{SO}_2\text{R}^{10}$, $-\text{SO}_2\text{OR}^{10}$, $-\text{OSO}_2\text{R}^{10}$,
 $-\text{SC}(=\text{W})\text{R}^{10}$, $-\text{SC}(=\text{W})\text{OR}^{10}$, $-\text{SC}(=\text{W})\text{R}^{10}$, $-\text{PR}^{10}_2$, $-\text{PW}_2\text{R}^{10}_2$,
 $-\text{PW}_3\text{R}^{10}_2$, SiR^{10}_3 or halogen;

- R^2 and R^3 independently of one another have the definitions given in R^1 ;
20 R^2 and R^3 together form a 5- to 7-membered ring which may be partially
or fully unsaturated and may be interrupted by one or more atoms from the
group nitrogen, oxygen and sulfur, the oxygen atoms not being directly
adjacent to one another, and the ring optionally being substituted by one or
more, but at most 5, radicals R^1 ;
25 R^4 and R^6 independently of one another have the definitions given in R^1 ;



* C L M S *

R^4 and R^6 together form a 4- to 7-membered ring which may be partially or fully unsaturated and may be interrupted by one or more atoms from the group nitrogen, oxygen and sulfur, the oxygen atoms not being directly adjacent to one another, and the ring optionally being substituted by one or more, but at most 5, radicals R^1 ;

R^5 and R^7 independently of one another are hydrogen,

(C_1 - C_{20})-alkyl, (C_2 - C_{20})-alkenyl, (C_2 - C_{20})-alkynyl, (C_3 - C_8)-cycloalkyl, (C_4 - C_8)-cycloalkenyl, (C_6 - C_8)-cycloalkynyl,

where the six last-mentioned radicals are optionally substituted by one or more radicals from the group

halogen, cyano, nitro, hydroxyl, $-C(=W)R^{10}$, $-C(=NOR^{10})R^{10}$,
 $-C(=NNR^{10}_2)R^{10}$, $-C(=W)OR^{10}$, $-C(=W)NR^{10}_2$,
 $-OC(=W)R^{10}$, $-OC(=W)OR^{10}$, $-NR^{10}C(=W)R^{10}$, -
 $N[C(=W)R^{10}]_2$, $-NR^{10}C(=W)OR^{10}$, $-C(=W)NR^{10}-NR^{10}_2$,
 $-C(=W)NR^{10}-NR^{10}[C(=W)R^{10}]$, $-NR^{10}-C(=W)NR^{10}_2$,
 $-NR^{10}-NR^{10}C(=W)R^{10}$, $-NR^{10}-N[C(=W)R^{10}]_2$,
 $-N[(C=W)R^{10}]-NR^{10}_2$, $-NR^{10}-NR^{10}[(C=W)R^{10}]$, $-NR^{10}$
 $-NR^{10}[(C=W)WR^{10}]$, $-NR^{10}-NR^{10}[(C=W)NR^{10}_2]$,
 $-NR^{10}(C=NR^{10})R^{10}$, $-NR^{10}(C=NR^{10})NR^{10}_2$, $-O-NR^{10}_2$,
 $-O-NR^{10}(C=W)R^{10}$, $-OR^{10}$, $-NR^{10}_2$, $-SR^{10}$, $-SiR^{10}_3$, $-SeR^{10}$,
 $-PR^{10}_2$, $-P(=W)R^{10}_2$, $-SOR^{10}$, $-SO_2R^{10}$, $-PW_2R^{10}_2$,
 $-PW_3R^{10}_2$, aryl and heterocyclyl,

of which the two mentioned last are optionally substituted by one or more radicals from the group

(C_1 - C_6)-alkyl, (C_2 - C_6)-alkenyl, (C_2 - C_6)-alkynyl, (C_3 - C_8)-cycloalkyl, (C_4 - C_8)-cycloalkenyl, (C_6 - C_8)-cycloalkynyl,



(C₁-C₆)-haloalkyl, (C₂-C₆)-haloalkenyl, (C₂-C₆)-haloalkynyl, halogen, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃, -C(=W)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -SOR¹⁰, -SO₂R¹⁰, nitro, cyano and hydroxyl,

5 aryl,

which is optionally substituted by one or more radicals from the group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl,

10 where these six abovementioned radicals are optionally substituted by one or more radicals from the group

halogen, cyano, nitro, halogen, -C(=W)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SOR¹⁰ and -SO₂R¹⁰,

15 halogen, cyano, nitro, -C(=W)R¹⁰, -C(=NOR¹⁰)R¹⁰, -C(=NNR¹⁰₂)R¹⁰, -C(=W)OR¹⁰, -C(=W)NR¹⁰₂, -OC(=W)R¹⁰, -OC(=W)OR¹⁰, -NR¹⁰C(=W)R¹⁰, -N[C(=W)R¹⁰]₂, -NR¹⁰C(=W)OR¹⁰, -OR¹⁰, -NR¹⁰₂, -SR¹⁰, -SiR¹⁰₃, -PR¹⁰₂, -SOR¹⁰, -SO₂R¹⁰, -PW₂R¹⁰₂ and -PW₃R¹⁰₂,




20 pyridyl,

which is optionally substituted by one or more radicals from the group

(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-cycloalkyl, (C₄-C₈)-cycloalkenyl and (C₆-C₈)-cycloalkynyl,

25 where the six abovementioned radicals are optionally substituted by one or more radicals from the group

cyano, nitro, halogen, -C(=W)R¹⁰, -C(=W)OR¹⁰,



R^4 and R^5 together form a 4- to 7-membered ring which may be partially
unsaturated and may be interrupted by one or more atoms from the group
nitrogen, oxygen and sulfur, oxygen atoms not being directly adjacent to
one another, and the ring optionally being substituted by one or more, but
at most 5, radicals R^1 ;

R^4 and R^5 together form one of the groups $=\text{O}$, $=\text{S}$ or $=\text{N}-\text{R}^9$;

R^6 and R^7 together form a 5- to 7-membered ring which may be partially
unsaturated and may be interrupted by one or more atoms from the group
nitrogen, oxygen and sulfur, oxygen atoms not being directly adjacent to
one another, and the ring optionally being substituted by one or more, but
at most 5, radicals R^1 ;

R^6 and R^7 together form one of the groups $=\text{O}$, $=\text{S}$ or $=\text{N}-\text{R}^9$;

R^8 is hydrogen,

(C_1-C_6)-alkyl, (C_2-C_6)-alkenyl, (C_2-C_6)-alkynyl, (C_3-C_8)-
cycloalkyl, (C_4-C_8)-cycloalkenyl, (C_3-C_8)-cycloalkyl-(C_1-C_4)-
alkyl, (C_4-C_8)-cycloalkenyl-(C_1-C_4)-alkyl, (C_3-C_8)-cycloalkyl-(C_2-
 C_4)-alkenyl, (C_4-C_8)-cycloalkenyl-(C_1-C_4)-alkenyl, (C_1-C_6)-alkyl-
(C_3-C_8)-cycloalkyl, (C_2-C_6)-alkenyl-(C_3-C_8)-cycloalkyl, (C_2-C_6)-



alkynyl-(C₃-C₈)-cycloalkyl, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenyl,

(C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyl,

where the fourteen last-mentioned radicals are optionally substituted by one or more radicals from the group

- 5 halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy, (C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, (C₃-C₈)-
- 10 cycloalkyl-(C₁-C₄)-alkoxy, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkoxy, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenyloxy, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyloxy, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkoxy, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkoxy, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkoxy, (C₁-C₆)-alkyl-(C₄-C₈)-
- 15 cycloalkenyloxy, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyloxy, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, (C₁-C₄)-alkoxy-(C₂-C₆)-alkenyloxy, carbamoyl, (C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-mono- or dihaloalkylcarbamoyl, (C₃-C₈)-mono- or dicycloalkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-
- 20 cycloalkoxycarbonyl, (C₁-C₆)-alkanoyloxy, (C₃-C₈)-cycloalkanoyloxy, (C₁-C₆)-haloalkoxycarbonyl, (C₁-C₆)-haloalkanoyloxy, (C₁-C₆)-alkaneamido, (C₁-C₆)-haloalkaneamido, (C₂-C₆)-alkeneamido, (C₃-C₈)-cycloalkaneamido, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkaneamido,
- 25 (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-alkynylthio, (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio, (C₂-C₆)-




- haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-
cycloalkenylthio, (C₃-C₈)-halocycloalkthio, (C₄-C₈)-
halocycloalkenylthio, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylthio,
(C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylthio, (C₃-C₈)-cycloalkyl-
5 (C₂-C₄)-alkenylthio, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylthio,
(C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylthio, (C₂-C₆)-alkenyl-(C₃-C₈)-
cycloalkylthio, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylthio, (C₁-
C₆)-alkyl-(C₄-C₈)-cycloalkenylthio, (C₂-C₆)-alkenyl-(C₄-C₈)-
cycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-alkenylsulfinyl,
10 (C₂-C₆)-alkynylsulfinyl, (C₁-C₆)-haloalkylsulfinyl, (C₂-C₆)-
haloalkenylsulfinyl, (C₂-C₆)-haloalkynylsulfinyl, (C₃-C₈)-
cycloalkylsulfinyl, (C₄-C₈)-cycloalkenylsulfinyl, (C₃-C₈)-
halocycloalksulfinyl, (C₄-C₈)-halocycloalkenylsulfinyl, (C₃-C₈)-
cycloalkyl-(C₁-C₄)-alkylsulfinyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-
15 alkylsulfinyl, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylsulfinyl, (C₄-
C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfinyl, (C₁-C₆)-alkyl-(C₃-
C₈)-cycloalkylsulfinyl, (C₂-C₆)-alkenyl-(C₃-C₈)-
cycloalkylsulfinyl, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylsulfinyl,
(C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylsulfinyl, (C₂-C₆)-alkenyl-
20 (C₄-C₈)-cycloalkenylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₂-C₆)-
alkenylsulfonyl, (C₂-C₆)-alkynylsulfonyl, (C₁-C₆)-
haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl, (C₂-C₆)-
haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl, (C₄-C₈)-
cycloalkenylsulfonyl, (C₃-C₈)-halocycloalksulfonyl, (C₄-C₈)-
25 halocycloalkenylsulfonyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-
alkylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylsulfonyl, (C₃-



- (C₈)-cycloalkyl-(C₂-C₄)-alkenylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfonyl, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylsulfonyl, (C₁-C₆)-alkyl-
- 5 (C₄-C₈)-cycloalkenylsulfonyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₁-C₆)-alkylamino, (C₂-C₆)-alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-haloalkylamino, (C₂-C₆)-haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-C₈)-cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-
- 10 halocycloalkamino, (C₄-C₈)-halocycloalkenylamino, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylamino, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylamino, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylamino, (C₂-C₆)-alkenyl-(C₃-C₈)-
- 15 cycloalkylamino, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylamino, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylamino, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylamino, (C₁-C₆)-trialkylsilyl, aryl, aryloxy, arylthio, arylamino, arylcarbonyl, aroyl, aroyloxy, aryloxycarbonyl, aryl-(C₁-C₄)-alkoxy, aryl-(C₂-C₄)-alkenyloxy,
- 20 aryl-(C₁-C₄)-alkylthio, aryl-(C₂-C₄)-alkenylthio, aryl-(C₁-C₄)-alkylamino, aryl-(C₂-C₄)-alkenylamino, aryl-(C₁-C₆)-dialkylsilyl, diaryl-(C₁-C₆)-alkylsilyl, triarylsilyl and 5- or 6-membered heterocyclyl,
- 25 of which the nineteen last-mentioned radicals are optionally substituted in their cyclic moiety by one or more substituents from the group
- halogen, cyano, nitro, amino, hydroxyl, thio,



- (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₁-C₄)-alkoxy,
(C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-
haloalkylthio, (C₁-C₄)-alkylamino, (C₁-C₄)-haloalkylamino,
formyl and (C₁-C₄)-alkanoyl,
- 5 aryl, which is optionally substituted by one or more radicals from
the group
- halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-
alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-
haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy,
- 10 (C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-
halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, carbamoyl,
(C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl,
(C₁-C₆)-alkanoyloxy, (C₁-C₆)-mono- or dihaloalkylcarbamoyl,
(C₁-C₆)-haloalkoxycarbonyl, (C₁-C₆)-haloalkanoyloxy,
- 15 (C₁-C₆)-alkaneamido, (C₁-C₆)-haloalkaneamido, (C₂-C₆)-
alkeneamido, (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-
alkynylthio, (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio,
(C₂-C₆)-haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-
cycloalkenylthio, (C₃-C₈)-halocycloalkthio, (C₃-C₈)-
- 20 halocycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-
alkenylsulfinyl, (C₂-C₆)-alkynylsulfinyl, (C₁-C₆)-
haloalkylsulfinyl, (C₂-C₆)-haloalkenylsulfinyl, (C₂-C₆)-
haloalkynylsulfinyl, (C₃-C₈)-cycloalkylsulfinyl, (C₄-C₈)-
cycloalkenylsulfinyl, (C₃-C₈)-halocycloalksulfinyl, (C₄-C₈)-
- 25 halocycloalkenylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₂-C₆)-
alkenylsulfonyl, (C₂-C₆)-alkynylsulfonyl, (C₁-C₆)-



haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl, (C₂-C₆)-
haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl, (C₄-C₈)-
cycloalkenylsulfonyl, (C₃-C₈)-halocycloalkylsulfonyl, (C₄-C₈)-
halocycloalkenylsulfonyl, (C₁-C₆)-alkylamino, (C₂-C₆)-
5 alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-haloalkylamino,
(C₂-C₆)-haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-
C₈)-cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-
halocycloalkylamino and (C₄-C₈)-halocycloalkenylamino,
-C(=W)R¹¹, OR¹¹ or NR¹¹₂;

10

R⁹ is (C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-
alkyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkyl, (C₃-C₈)-cycloalkyl-
(C₂-C₄)-alkenyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyl,

15

where the nine last-mentioned radicals are optionally substituted
by one or more radicals from the group

halogen, cyano, (C₁-C₆)-alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-
alkynyloxy and (C₁-C₆)-haloalkyloxy;

20 R¹⁰

is hydrogen,
(C₁-C₆)-alkyl, (C₂-C₆)-alkenyl, (C₂-C₆)-alkynyl, (C₃-C₈)-
cycloalkyl, (C₄-C₈)-cycloalkenyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-
alkyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkyl, (C₃-C₈)-cycloalkyl-
(C₂-C₄)-alkenyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyl, (C₁-C₆)-
25 alkyl-(C₃-C₈)-cycloalkyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkyl,



(C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkyl, (C₁-C₆)-alkyl-(C₄-C₈)-

cycloalkenyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyl,

where the fourteen last-mentioned radicals are optionally substituted by one or more radicals from the group

5 halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy,

(C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkoxy, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkoxy, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenyloxy, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenyloxy, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkoxy, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkoxy, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkoxy, (C₁-C₆)-alkyl-(C₄-C₈)-

15 cycloalkenyloxy, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenyloxy, (C₁-C₄)-alkoxy-(C₁-C₆)-alkoxy, (C₁-C₄)-alkoxy-(C₂-C₆)-alkenyloxy, carbamoyl,

(C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-mono- or dihaloalkylcarbamoyl, (C₃-C₈)-mono- or

20 dicycloalkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl, (C₃-C₈)-cycloalkoxycarbonyl, (C₁-C₆)-alkanoyloxy, (C₃-C₈)-cycloalkanoyloxy, (C₁-C₆)-haloalkoxycarbonyl, (C₁-C₆)-haloalkanoyloxy, (C₁-C₆)-alkaneamido, (C₁-C₆)-haloalkaneamido, (C₂-C₆)-alkeneamido, (C₃-C₈)-

25 cycloalkaneamido, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkaneamido, (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-alkynylthio,



- (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio, (C₂-C₆)-haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-cycloalkenylthio, (C₃-C₈)-halocycloalkthio, (C₄-C₈)-halocycloalkenylthio, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylthio,
- 5 (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylthio, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylthio, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylthio, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylthio, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylthio, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylthio, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylthio, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-alkenylsulfinyl,
- 10 (C₂-C₆)-alkynylsulfinyl, (C₁-C₆)-haloalkylsulfinyl, (C₂-C₆)-haloalkenylsulfinyl, (C₂-C₆)-haloalkynylsulfinyl, (C₃-C₈)-cycloalkylsulfinyl, (C₄-C₈)-cycloalkenylsulfinyl, (C₃-C₈)-halocycloalksulfinyl, (C₄-C₈)-halocycloalkenylsulfinyl, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylsulfinyl,
- 15 (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylsulfinyl, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylsulfinyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfinyl, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylsulfinyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylsulfinyl, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylsulfinyl, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylsulfinyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylsulfinyl, (C₁-C₆)-alkylsulfonyl, (C₂-C₆)-alkenylsulfonyl,
- 20 (C₂-C₆)-alkynylsulfonyl, (C₁-C₆)-haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl, (C₂-C₆)-haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl, (C₄-C₈)-cycloalkenylsulfonyl, (C₃-C₈)-halocycloalksulfonyl, (C₄-C₈)-halocycloalkenylsulfonyl, (C₃-
- 25



This Sheet Has Been Replaced



138

- 5 C₈)-cycloalkyl-(C₁-C₄)-alkylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylsulfonyl, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylsulfonyl, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylsulfonyl, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylsulfonyl, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylsulfonyl, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylsulfonyl, (C₁-C₆)-alkylamino, (C₂-C₆)-alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-haloalkylamino, (C₂-C₆)-haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-C₈)-cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-halocycloalkamino; (C₄-C₈)-halocycloalkenylamino, (C₃-C₈)-cycloalkyl-(C₁-C₄)-alkylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkylamino, (C₃-C₈)-cycloalkyl-(C₂-C₄)-alkenylamino, (C₄-C₈)-cycloalkenyl-(C₁-C₄)-alkenylamino, (C₁-C₆)-alkyl-(C₃-C₈)-cycloalkylamino, (C₂-C₆)-alkenyl-(C₃-C₈)-cycloalkylamino, (C₂-C₆)-alkynyl-(C₃-C₈)-cycloalkylamino, (C₁-C₆)-alkyl-(C₄-C₈)-cycloalkenylamino, (C₂-C₆)-alkenyl-(C₄-C₈)-cycloalkenylamino, (C₁-C₆)-trialkylsilyl, aryl, aryloxy, arylthio, arylamino, aryl-(C₁-C₄)-alkoxy, aryl-(C₂-C₄)-alkenyloxy, aryl-(C₁-C₄)-alkylthio, aryl-(C₂-C₄)-alkenylthio, aryl-(C₁-C₄)-alkylamino, aryl-(C₂-C₄)-alkenylamino, aryl-(C₁-C₆)-dialkylsilyl, diaryl-(C₁-C₆)-alkylsilyl, triarylsilyl and 5- or 6-membered heterocyclyl,



where the cyclic moiety of the fourteen last-mentioned radicals is optionally substituted by one or more radicals from the group

5 halogen, cyano, nitro, amino, hydroxyl, thio, (C₁-C₄)-alkyl, (C₁-C₄)-haloalkyl, (C₃-C₈)-cycloalkyl, (C₁-C₄)-alkoxy, (C₁-C₄)-haloalkoxy, (C₁-C₄)-alkylthio, (C₁-C₄)-haloalkylthio, (C₁-C₄)-alkylamino, (C₁-C₄)-haloalkylamino, formyl and (C₁-C₄)-alkanoyl,

10 aryl, 5- or 6-membered heteroaromatic, where the two last-mentioned radicals are optionally substituted by one or more radicals from the group

halogen, cyano, nitro, hydroxyl, thio, amino, formyl, (C₁-C₆)-alkoxy, (C₂-C₆)-alkenyloxy, (C₂-C₆)-alkynyloxy, (C₁-C₆)-haloalkyloxy, (C₂-C₆)-haloalkenyloxy, (C₂-C₆)-haloalkynyloxy, 15 (C₃-C₈)-cycloalkoxy, (C₄-C₈)-cycloalkenyloxy, (C₃-C₈)-halocycloalkoxy, (C₄-C₈)-halocycloalkenyloxy, carbamoyl, (C₁-C₆)-mono- or dialkylcarbamoyl, (C₁-C₆)-alkoxycarbonyl, (C₁-C₆)-alkanoyloxy, (C₁-C₆)-mono- or dihaloalkylcarbamoyl, (C₁-C₆)-haloalkoxycarbonyl, 20 (C₁-C₆)-haloalkanoyloxy, (C₁-C₆)-alkaneamido, (C₁-C₆)-haloalkaneamido, (C₂-C₆)-alkeneamido, (C₁-C₆)-alkylthio, (C₂-C₆)-alkenylthio, (C₂-C₆)-alkynylthio, (C₁-C₆)-haloalkylthio, (C₂-C₆)-haloalkenylthio, (C₂-C₆)-haloalkynylthio, (C₃-C₈)-cycloalkylthio, (C₄-C₈)-cycloalkenylthio, 25 (C₃-C₈)-halocycloalkthio, (C₄-C₈)-halocycloalkenylthio, (C₁-C₆)-alkylsulfinyl, (C₂-C₆)-alkenylsulfinyl, (C₂-C₆)-



- alkynylsulfinyl, (C₁-C₆)-haloalkylsulfinyl, (C₂-C₆)-
haloalkenylsulfinyl, (C₂-C₆)-haloalkynylsulfinyl, (C₃-C₈)-
cycloalkylsulfinyl, (C₄-C₈)-cycloalkenylsulfinyl, (C₃-C₈)-
halocycloalksulfinyl, (C₄-C₈)-halocycloalkenylsulfinyl, (C₁-C₆)-
5 alkylsulfonyl, (C₂-C₆)-alkenylsulfonyl, (C₂-C₆)-alkynylsulfonyl,
(C₁-C₆)-haloalkylsulfonyl, (C₂-C₆)-haloalkenylsulfonyl, (C₂-
C₆)-haloalkynylsulfonyl, (C₃-C₈)-cycloalkylsulfonyl, (C₄-C₈)-
cycloalkenylsulfonyl, (C₃-C₈)-halocycloalksulfonyl, (C₄-C₈)-
halocycloalkenylsulfonyl, (C₁-C₆)-alkylamino, (C₂-C₆)-
10 alkenylamino, (C₂-C₆)-alkynylamino, (C₁-C₆)-haloalkylamino,
(C₂-C₆)-haloalkenylamino, (C₂-C₆)-haloalkynylamino, (C₃-
C₈)-cycloalkylamino, (C₄-C₈)-cycloalkenylamino, (C₃-C₈)-
halocycloalkamino and (C₄-C₈)-halocycloalkenylamino;
- 15 R¹¹ is (C₁-C₁₀)-alkyl, haloalkyl, aryl,
which is optionally substituted by one or more radicals from the
group
halogen, cyano, nitro, (C₁-C₄)-alkoxy, (C₁-C₄)-alkyl, amino,
(C₁-C₄)-monoalkylamino and (C₁-C₄)-dialkylamino,
20 NR¹⁰₂, OR¹⁰ or SR¹⁰
with the proviso that the compounds listed below are not included:
- 3-(2-chlorophenyl)-1-methyl-5-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole
 - 3-(2,6-difluorophenyl)-1-methyl-5-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole
 - 25 • 3-(2-chloro-4-fluorophenyl)-1-methyl-5-(4-trifluoromethyl-3-pyridyl)-1H-1,2,4-triazole
 - 3-(3,5-dichlorophenyl)-5-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole




- 5-(4-trifluoromethyl-3-pyridyl)- 3-phenyl-1,2,4-oxadiazole
- 3-(4-trifluoromethyl-3-pyridyl)- 5-phenyl-1,2,4-oxadiazole
- 5-(2-chlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(3-chlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5 • 5-(4-chlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(2-fluorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(4-fluorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(2,4-dichlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(3,4-dichlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 10 • 5-(3,5-dichlorophenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(2,6-dichloro-4-pyridyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 5-(3,5-bistrifluoromethylphenyl)- 3-(4-trifluoromethyl-3-pyridyl)-1,2,4-oxadiazole
- 2-(2-chlorophenyl)- 5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 15 • 2-(3-chlorophenyl)- 5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 2-(4-chlorophenyl)- 5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 2-(2-trifluoromethoxyphenyl)- 5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 2-(4-trifluoromethoxyphenyl)- 5-(4-trifluoromethyl-3-pyridyl)-1,3,4-oxadiazole
- 20 • 2-(4-trifluoromethyl-3-pyridyl)-5-phenyl-1,3,4-oxadiazole
- 2-(4-trifluoromethyl-3-pyridyl)-4-methylthiazolecarbohydrazide
- ethyl 2-(4-trifluoromethyl-3-pyridyl)-4-methylthiazolecarboxylate
- N-(4-chlorophenyl)carbonyl-N'-[2-(4-trifluoromethyl-3-pyridyl)-4-methyl-5-thiazolyl]carbonylhydrazine
- 25 • 2-(4-trifluoromethyl-3-pyridyl)-4-thiazolecarbohydrazide
- 4-(4-chlorophenyl)-2-(4-trifluoromethyl-3-pyridyl)thiazole
- 4-(4-cyanophenyl)-2-(4-trifluoromethyl-3-pyridyl)thiazole
- N-(4-trifluoromethylphenyl)carbonyl-N'-[2-(4-trifluoromethyl-3-pyridyl)-4-thiazolyl]carbonylhydrazine
- 30 • 2-(2-(4-trifluoromethyl-3-pyridyl)thiazolyl)-5-chloro-3-methylbenzo[b]thiophene



- 2-(4-chlorophenylmethylthio)-5-(4-trifluoromethyl-3-pyridyl)-1-methyl-1H-1,3,4-triazole
 - 2-(4-chlorophenylcarbonylmethylthio)-5-(4-trifluoromethyl-3-pyridyl)-1-methyl-1H-1,3,4-triazole and
- 5 • 2-ethoxycarbonylmethylthio-5-(4-trifluoromethyl-3-pyridyl)-1-methyl-1H-1,3,4-triazole.

2. A 4-haloalkyl-3-heterocyclylpyridine or 4-haloalkyl-5-heterocyclylpyrimidine as claimed in claim 1, wherein

- 10 Y is C₁-C₆-alkyl which is mono- or polysubstituted by chlorine and/or fluorine;
- m is zero;
- Q is a 5-membered heterocyclic group



15

in which

- a) $X^2 = NR^a$ and $X^3 = CR^bR^1$ or
- b) $X^2 = CR^aR^2$ and $X^3 = CR^bR^3$ or
- c) $X^2 = CR^4R^5$ and $X^3 = CR^6R^7$;
- 20 R^a and R^b together are a bond ;
- R^1, R^2, R^3, R^4 and R^6 are each independently of one another
- hydrogen, halogen, C₁-C₁₂-alkyl, C₃-C₈-cycloalkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, where the four last-mentioned
- hydrocarbon radicals are optionally mono- or polysubstituted
- 25 by identical or different radicals from a group A1 consisting of
- C₁-C₆-alkylcarbonyl, C₁-C₆-alkylaminocarbonyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, C₁-C₆-alkylcarbonylamino, C₁-C₆-alkylsulfonylamino, phenyl, furyl, pyrrolyl, thienyl, halogen, cyano, phenyloxy, phenylthio and



- phenylamino, where the eleven first-mentioned radicals of group A1 are each optionally mono- or polysubstituted by identical or different radicals from a group B1 consisting of halogen, cyano, C₁-C₃-alkoxy and phenyl which is optionally
- 5 mono- or polysubstituted by one or more halogen atoms and where the three last-mentioned radicals of group A1 are each optionally mono- or polysubstituted by identical or different radicals from a group B2 consisting of halogen, cyano, nitro, C₁-C₃-alkyl and C₁-C₃-alkoxy, or are C₁-C₆-alkylcarbonyl,
- 10 C₁-C₆-alkylaminocarbonyl, C₁-C₆-alkoxycarbonyl, phenyl, pyridyl, furyl, thienyl, pyrrol, where the eight last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from group B1, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;
- 15 R⁵ and R⁷ are each independently of one another hydrogen, halogen, C₁-C₁₂-alkyl, C₃-C₈-cycloalkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl, where the four last-mentioned hydrocarbon radicals are optionally mono- or polysubstituted by identical or different radicals from a group A2 consisting of C₁-C₆-alkylcarbonyl,
- 20 C₁-C₆-alkylaminocarbonyl, C₁-C₆-alkoxy, C₁-C₆-alkylthio, C₁-C₆-alkylamino, C₁-C₆-alkylcarbonylamino, phenyl, furyl, pyrrol, thienyl, halogen, cyano, phenyloxy, phenylthio and phenylamino, where the ten first-mentioned radicals of group A2 are each optionally mono- or polysubstituted by identical
- 25 or different radicals from the group B1 and the three last-mentioned radicals of group A2 are each optionally mono- or polysubstituted by identical or different radicals from the group B2, or are C₁-C₆-alkylcarbonyl, C₁-C₆-alkylaminocarbonyl,



5 C₁-C₆-alkoxycarbonyl, phenyl, pyridyl, furyl, thienyl, fluorine, chlorine, bromine, pyrrol, where the eight last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from the group B1, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;

R¹⁰ is hydrogen, benzyl, C₁-C₆-alkyl, C₁-C₆-cycloalkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, phenyl, C₁-C₆-alkylcarbonyl or C₁-C₆-alkylsulfonyl, where the eight last-mentioned radicals are optionally mono- or polysubstituted by identical or different halogen atoms.

3. A 4-haloalkyl-3-heterocyclylpyridine or 4-haloalkyl-5-heterocyclylpyrimidine as claimed in claim 2, wherein

Y is trifluoromethyl;

15 R¹, R², R³, R⁴ and R⁶ are each independently of one another halogen, C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A3 consisting of C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, C₁-C₄-alkylsulfonylamino, phenyl, furyl, pyrrol, thienyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the eleven first-mentioned radicals of group A3 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A3 are each optionally mono- or polysubstituted by identical or different radicals from the group B2, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;




- R^5 and R^7 are each independently of one another halogen, C₁-C₁₂-alkyl, C₂-C₁₂-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A4 consisting of C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, phenyl, furyl, pyrrol, thienyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the ten first-mentioned radicals of group A4 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A4 are each optionally mono- or polysubstituted by identical or different radicals from the group B2, or are OR¹⁰, SR¹⁰ or N(R¹⁰)₂;
- R^{10} is hydrogen, C₁-C₆-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, phenyl, C₁-C₄-alkylcarbonyl or C₁-C₄-alkylsulfonyl, where the six last-mentioned radicals are optionally mono- or polysubstituted by identical or different halogen atoms.
4. A 4-haloalkyl-3-heterocyclylpyridine or 4-haloalkyl-5-heterocyclyl-pyrimidine as claimed in claim 3, wherein R^1 , R^2 , R^3 , R^4 and R^6 are each independently of one another C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A5 consisting of C₁-C₄-alkylcarbonyl, C₁-C₄-alkylaminocarbonyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, C₁-C₄-alkylsulfonylamino, phenyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the




5 eight first-mentioned radicals of group A5 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A5 are each optionally mono- or polysubstituted by identical or different radicals from the group B2;

10 R^5 and R^7 are each independently of one another C_1 - C_{10} -alkyl, C_2 - C_{10} -alkenyl, where the two last-mentioned radicals are optionally mono- or polysubstituted by identical or different radicals from a group A6 consisting of C_1 - C_4 -alkylcarbonyl, C_1 - C_4 -alkylaminocarbonyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylamino, C_1 - C_4 -alkylcarbonylamino, phenyl, fluorine, chlorine, bromine, cyano, phenyloxy, phenylthio and phenylamino, where the
15 seven first-mentioned radicals of group A6 are each optionally mono- or polysubstituted by identical or different radicals from the group B1 and the three last-mentioned radicals of group A6 are each optionally mono- or polysubstituted by identical or different radicals from the group B2.


20 5. A process for preparing compounds of the formula I as claimed in any of claims 1 to 4, which comprises reacting, if



activated derivatives of the acid of the formula (II),




where X and Y are as defined above, in the presence of a base with a compound of the formula (III),




in which the radical R^1 is as defined in formula (I),

5


or, if




compounds of the formula (VII) with an electrophilic amination agent



or, if




amidoximes of the formula (V) with activated derivatives of acids of the
15 formula (VI) or with the acids of the formula (VI) themselves






or, if




N,N'-diacylhydrazines of the formula (XIII) with a thiolation agent, such as Lawessons reagent, in an inert solvent


5



or, if



acids of the formula (II) with hydrazides of the formula (X), in which R^1 is as defined above using an activating reagent, such as phosphorus oxychloride or phosphorus pentachloride,




15


or acid hydrazides of the formula (XI) with orthoesters of the formula (XII), where R^1 is as defined above, and R^{12} is (C₁-C₄)-alkyl




149




or, if



5 and R^a , R^b and R^3 are as defined above, compounds of the formula (XIV) with a dehydrating reagent, such as inorganic acid chlorides, inorganic acids and anhydrides,




10 or, if




thioamides of the formula (XVII) with carbonyl derivatives of the formula (XVIII) where Z is halogen, in particular chlorine or bromine, acyloxy or sulfonyloxy, in particular methanesulfonyloxy or tolylsulfonyloxy

15







or, if




- 5 thioamides of the formula (XVII) with compounds of the formula (XIX) where the two substituents Z are as defined above and may either be identical or different



or, if





and R^a , R^b , R^1 and R^8 are as defined above, hydrazides of the formula (XX)



15

with a compound of the formula (XXI) or with thioamides of the formula (XXII),





6. A composition having insecticidal, acaricidal and/or nematocidal action, which comprises at least one compound as claimed in any of claims 1 to 4.
- 5
7. A composition having insecticidal, acaricidal and nematocidal action as claimed in claim 6 in a mixture with carriers and/or surfactants.
8. A composition as claimed in claim 6 or 7, which contains a
- 10 further active compound from the group acaricides, fungicides, herbicides, insecticides, nematocides or growth-regulating substances.
9. A compound as claimed in any of claims 1 to 4 or a composition as claimed in claim 6 or 7, for use as veterinary medicament,
- 15 preferably for controlling endo- and ectoparasites.
10. A method for controlling harmful insects, acarina and nematodes, which comprises applying an effective amount of a compound as claimed in any of claims 1 to 4 or of a composition as claimed in any of
- 20 claims 6 to 8 to the area where the action is desired.
11. A process for protecting useful plants against the undesired action of harmful insects, acarina and nematodes, which comprises using at least one of the compounds as claimed in any of claims 1 to 4 for
- 25 treating the seed of the useful plants.
12. The use of compounds as claimed in any of claims 1 to 4 or of a composition as claimed in any of claims 6 to 8 for controlling harmful insects, acarina and nematodes.