

**NEURAL REGENERATING CELLS
WITH ALTERATIONS IN DNA METHYLATION**

STATEMENT REGARDING FEDERAL SUPPORT

5 [0001] Not applicable.

FIELD

[0002] The present disclosure is in the fields of cellular therapy for neural disorders and epigenetic regulation of gene expression and differentiation.

10

BACKGROUND

[0003] Cellular differentiation is controlled, in part, by regulation of gene expression. Regulation of transcription; *i.e.*, use of DNA as a template for the synthesis of a mRNA molecule; is one of the mechanisms by which gene expression is regulated. Transcriptional regulation of gene expression can result, for example, from alteration of chromatin structure and/or binding of transcriptional regulatory proteins to specific DNA sequences in or near the gene.

15

[0004] Another means by which transcriptional control of gene expression is effected is by chemical alteration of DNA. The most widely-studied aspect of this form of regulation is DNA methylation. In eukaryotic genomes, the primary form of DNA methylation is the conversion of cytosine to 5-methyl-cytosine, through the action of one of a number of cellular methyltransferases. In most cases, methylated C residues are located directly upstream of a G residue. In general, methylation of C residues in or near a gene is correlated with reduced expression of the gene. In most cases, CpG methylation is not itself the proximate cause of transcriptional repression of a gene, but appears to be a mechanism to perpetuate transcriptional repression initially mediated by gene regulatory proteins.

20

25

[0005] The frequency of CG dinucleotide sequences in the upstream regions of certain non-cell-type-specific vertebrate genes (*i.e.*, housekeeping genes) is much higher than would be expected based on the GC content of the genome; such regions are known as CpG Islands. CpG Islands are sites at which the methylation state of the C residues can affect transcription of the associated gene. Conversely, the methylation state of C residues in a CpG island or other region associated with a particular gene can be used as a potential indicator of the transcriptional state of that

30

gene and/or as a diagnostic marker to characterize a particular cell type. See, for example, WO 2006/094836.

SUMMARY

5 [0006] Disclosed herein are cells that are capable of stimulating neural recovery and/or neural regeneration after transplantation to sites of nervous system injury or disease.

[0006a] In a first aspect, the present invention refers to an *in vitro* method for altering the methylation state of one or more genes in a progenitor cell, wherein the
10 progenitor cell is a marrow adherent stromal cell (MASC), by the process of:

transfecting the progenitor cell with a polynucleotide comprising sequences encoding a Notch intracellular domain;

wherein the alterations in methylation state comprise:

(a) increased methylation of the PITX2, DNMT3b, Igf2R and SDF4 genes,
15 and

(b) decreased methylation of the ROPN1L and TMEM179 genes.

[0006b] In a second aspect, the present invention refers to an *in vitro* method for converting a progenitor cell to a cell that stimulates neural recovery and/or neural regeneration after transplantation to a site of nervous
20 system injury or disease, the method comprising altering the methylation state of one or more genes in the progenitor cell, wherein the progenitor cell is a marrow adherent stromal cell;

further wherein the alterations in methylation state comprise:

(i) increased methylation of the PITX2, DNMT3b, Igf2R and SDF4 genes,
25 and

(ii) decreased methylation of the ROPN1L and TMEM179 genes;

further wherein neither the progenitor cell nor any of its descendents were transfected with a polynucleotide comprising sequences encoding a Notch intracellular domain (NICD).

30 [0006c] In a third aspect, the present invention refers to a method for identifying a cell, descended from a marrow adherent stromal cell, that stimulates neural recovery and/or neural regeneration after transplantation to a site of nervous system injury or disease, the method comprising assaying the methylation state of one or more genes in the descendent cell;

wherein the one or more genes is selected from the group consisting of PITX2, DNMT3b, IGF2R, SDF4, ROPN1L and TMEM179;

further wherein:

(i) methylation of the PITX2, DNMT3b, IGF2R and SDF4 genes is increased, and

(ii) methylation of the ROPN1L and TMEM179 genes is decreased.

[0006d] In certain exemplifications, the cells are descended from marrow adherent stem cells (MASCs), but have undergone alterations in the methylation status of certain genes after treatment and culture *in vitro*. Thus, the inventor has discovered that alteration of the methylation state of one or more genes can convert a progenitor cell into a descendent cell having neural regenerative properties not possessed by the progenitor cell.

[0007] As a result of this discovery, the present disclosure encompasses, *inter alia*, the following exemplifications:

1. A method for altering the methylation state of a gene in a cell, the method comprising:

(a) transfecting the cell with a polynucleotide comprising sequences encoding a Notch intracellular domain; and

(b) culturing the transfected cell such that the methylation state of a gene in the cell or one or more descendents of the cell is altered as compared to the gene in an untransfected cell, thereby altering the methylation state of the gene.

2. The method of exemplification 1, wherein the gene is the PITX2 gene.

3. The method of exemplification 1, wherein the gene is the DNMT3b gene.

4. The method of exemplification 1, wherein the gene is the IGF2R gene.

5. The method of exemplification 1, wherein the gene is the SDF4 gene.

6. The method of exemplification 1, wherein the gene is the ROPN1L gene.

7. The method of exemplification 1, wherein the gene is the TMEM179 gene.

8. The method of any of exemplifications 1-5, wherein methylation of the gene is increased in the descendent cell.

9. The method of any of exemplifications 1 6 or 7, wherein methylation of the gene is decreased in the descendent cell.

10. The method of exemplification 9, wherein the sequence C-A-T-C^{me}-G-C-C-C is converted to C-A-T-C-G-C-C-C.

11. The method of any of exemplifications 1-10, wherein the cell is a marrow adherent stromal cell (MASC).

12. A method for making a descendent cell in which the methylation state of a gene is altered, the method comprising:

- 5 (a) transfecting a progenitor cell with a polynucleotide comprising sequences encoding a Notch intracellular domain;
- (b) culturing the transfected cell; and
- (c) obtaining, among the progeny of the transfected cell, one or more descendent cells in which the methylation state of the gene is altered.

10 13. The method of exemplification 12, wherein the gene is the PITX2 gene.

14. The method of exemplification 12, wherein the gene is the DNMT3b gene.

15. The method of exemplification 12, wherein the gene is the IGF2R gene.

16. The method of exemplification 12, wherein the gene is the SDF4 gene.

17. The method of exemplification 12, wherein the gene is the ROPN1L gene.

15 18. The method of exemplification 12, wherein the gene is the TMEM179 gene.

19. The method of any of exemplifications 12-16, wherein methylation of the gene is increased in the descendent cell compared to the progenitor cell.

20. The method of any of exemplifications 12 or 17 or 18, wherein methylation of the gene is decreased in the descendent cell compared to the progenitor cell.

20 21. The method of exemplification 20, wherein the sequence C-A-T-C^{me}-G-C-C-C is converted to C-A-T-C-G-C-C-C.

22. The method of any of exemplifications 12-21, wherein the progenitor cell is a marrow adherent stromal cell (MASC).

23. A method for converting a progenitor cell to a neural regenerating cell, the method comprising altering the methylation state of one or more genes in the progenitor cell.

24. The method of exemplification 23, wherein the gene is the PITX2 gene.

25. The method of exemplification 23, wherein the gene is the DNMT3b gene.

26. The method of exemplification 23, wherein the gene is the IGF2R gene.

27. The method of exemplification 23, wherein the gene is the SDF4 gene.

28. The method of exemplification 23, wherein the gene is the ROPN1L gene.

29. The method of exemplification 23, wherein the gene is the TMEM179 gene.

30. The method of any of exemplifications 23-27, wherein methylation of the gene is increased in the neural regenerating cell, compared to the progenitor cell.

31. The method of any of exemplifications 23, 28 or 29, wherein methylation of the gene is decreased in the neural regenerating cell, compared to the progenitor cell.

32. The method of exemplification 31, wherein the sequence C-A-T-C^{me}-G-C-C-C is converted to C-A-T-C-G-C-C-C.

5 33. The method of any of exemplifications 23-32, wherein the progenitor cell is a marrow adherent stromal cell (MASC).

34. The method of exemplification 23, wherein the methylation state of the gene is altered by:

(a) transfecting the progenitor cell with a polynucleotide comprising
10 sequences encoding a Notch intracellular domain;

(b) culturing the transfected cell; and

(c) obtaining, among the progeny of the transfected cell, one or more
descendent cells in which the methylation state of the gene is altered;

wherein said descendent cell in which the methylation state of the gene is
15 altered is a neural regenerating cell.

35. The method of exemplification 30, wherein the methylation state of the gene is altered by contacting the progenitor cell with a fusion protein comprising a methylation domain and a DNA-binding domain, or with a nucleic acid encoding a fusion protein comprising a methylation domain and a DNA-binding domain, wherein
20 the DNA binding domain is engineered to bind to one or more sequences in the gene.

36. The method of exemplification 31, wherein the methylation state of the gene is altered by contacting the progenitor cell with a fusion protein comprising a demethylation domain and a DNA-binding domain, or with a nucleic acid encoding a fusion protein comprising a demethylation domain and a DNA-binding domain,
25 wherein the DNA binding domain is engineered to bind to one or more sequences in the gene.

37. A cell that is descended from a progenitor cell through culture *in vitro*, wherein:

(a) the cell supports the growth and/or regeneration of neural tissue;

30 (b) the methylation state of one or more genes in the cell is altered compared to the progenitor cell; and

(c) during the culture *in vitro*, neither the progenitor cell nor any of its descendents were transfected with a polynucleotide comprising sequences encoding the Notch intracellular domain (NICD).

38. The cell of exemplification 37, wherein the gene is the PITX2 gene.

39. The cell of exemplification 37, wherein the gene is the DNMT3b gene.

40. The cell of exemplification 37, wherein the gene is the IGF2R gene.

41. The cell of exemplification 37, wherein the gene is the SDF4 gene.

5 42. The cell of exemplification 37, wherein the gene is the ROPN1L gene.

43. The cell of exemplification 37, wherein the gene is the TMEM179 gene.

44. The cell of any of exemplifications 37-41, wherein methylation of the gene is increased in the neural regenerating cell, compared to the progenitor cell.

10 45. The cell of any of exemplifications 37, 42 or 43, wherein methylation of the gene is decreased in the neural regenerating cell compared to the progenitor cell.

46. The cell of exemplification 45, wherein the sequence C-A-T-C^{me}-G-C-C-C is converted to C-A-T-C-G-C-C-C.

47. The cell of any of exemplifications 37-46, wherein the progenitor cell is a marrow adherent stromal cell (MASC).

15 48. The cell of exemplification 37, wherein the methylation state of the gene is altered by contacting the progenitor cell with a fusion protein comprising a methylation domain and a DNA-binding domain, or with a nucleic acid encoding a fusion protein comprising a methylation domain and a DNA-binding domain, wherein the DNA binding domain is engineered to bind to one or more sequences in the gene.

20 49. The cell of exemplification 37, wherein the methylation state of the gene is altered by contacting the progenitor cell with a fusion protein comprising a demethylation domain and a DNA-binding domain, or with a nucleic acid encoding a fusion protein comprising a demethylation domain and a DNA-binding domain, wherein the DNA binding domain is engineered to bind to one or more sequences in
25 the gene.

50. A method for identifying a neural regenerating cell, the method comprising assaying the methylation state of one or more genes in the cell, wherein a change in the methylation state of the assayed genes is indicative of a neural regenerating cells.

30 51. The method of exemplification 50, wherein the assay is for increased methylation of the one or more genes.

52. The method of exemplification 50, wherein the assay is for decreased methylation of the one or more genes.

53. The method of exemplification 50, wherein the assay is for increased methylation of one or more first genes and for decreased methylation of one or more second genes.

54. The method of exemplification 50, wherein the one or more genes is/are selected from the group consisting of PITX2, ROPN1L, DNMT3b, IGF2R, TMEM179 and SDF4.

BRIEF DESCRIPTION OF THE DRAWINGS

[0008] Not applicable.

DETAILED DESCRIPTION

Methylation State as a Diagnostic

[0009] Analysis of changes in the DNA methylation state of specific CpG sequences in or near one or more genes of interest can be used to identify a cell and to distinguish it from other cells with different DNA methylation patterns. For example, if, in a stem or other type of progenitor cell, a particular CpG sequence is methylated on its C residue, and upon further differentiation, the C residue becomes demethylated; the demethylation of that C residue can be used as a marker for that differentiation step. Conversely, methylation of a C residue may serve as a marker for differentiation. Total, all-or-none changes in methylation state are not required; a change in methylation frequency at a particular CpG sequence can also be diagnostic.

[0010] A number of methods, known in the art, can be used to distinguish methylated from unmethylated cytosine residues. These include, but are not limited to, treatment of DNA with bisulfite, and assay of DNA cleavage with methylation-sensitive and methylation-dependent restriction enzymes. Bisulfite (SO_3^-) treatment deaminates unmethylated cytosine, converting it to deoxyuridine which, upon replication, templates an adenosine residue in the nascent DNA strand. Thus, bisulfite treatment results in eventual conversion of a C-G base pair into a T-A base pair; and such changes can be detected by standard DNA sequencing methods. Methylated C residues are unaffected by bisulfite treatment; therefore C^{me} -G base pairs remain unchanged.

[0011] For assay of methylation status using restriction enzymes, enzymes with the sequence CG in their recognition site can be used. For certain recognition sites which contain the sequence CG, an enzyme which recognizes the site will fail to

cleave it if the C residue is methylated, but an isoschizomer of that enzyme (*i.e.*, an enzyme that recognizes the same sequence) will cleave the site whether or not the C residue is methylated. For example, both *HpaII* and *MspI* recognize the sequence CCGG. *MspI* cleaves the site regardless of whether the second C residue is

5 methylated. However, *HpaII* will cleave the site only if the second C residue is unmethylated. Thus, cleavage of a CCGG sequence by both enzymes indicates that the second C residue in the site is unmethylated (*i.e.*, the site has the sequence C-C-G-G); while cleavage by *MspI* only indicates that the second C residue is methylated (*i.e.*, the site has the sequence C-C^{me}-G-G).

10 [0012] In practice, analysis of the methylation state of a particular CpG sequence involves identification of a longer sequence that includes the CpG of interest. This sequence, often denoted an amplicon, is generally chosen so that it includes one or more CpG dinucleotide sequences (at which the methylation state may differ in different cell types) and is suitable for amplification; *e.g.*, by polymerase

15 chain reaction. Such amplicon sequences are generally long enough to be unique in a mammalian-sized genome.

[0013] Additional details and other information relating to methylation analysis and exemplary amplicons that can be used for analysis of DNA methylation are found in WO 2006/094836 (Sept. 14, 2006), the disclosure of which is

20 incorporated by reference for the purposes of providing additional details and other information relating to methylation analysis and exemplary amplicons that can be used for analysis of DNA methylation.

Progenitor Cells

25 [0014] Progenitor cells, which can be converted to neural regenerating cells by altering the methylation status of certain genes, can be any type of non-terminally differentiated cell. For example, totipotent stem cells as disclosed for example, in U.S. Patent Nos. 5,843,780; 6,200,806 and 7,029,913 can be used as progenitor cells. Totipotent stem cells can be cultured (*e.g.*, U.S. Patent Nos. 6,602,711 and 7,005,252)

30 and differentiated into various types of pluripotent cells (*e.g.*, U.S. Patent Nos. 6,280,718; 6,613,568 and 6,887,706), which can also be used as progenitor cells in the practice of the disclosed methods.

[0015] Another exemplary type of progenitor cells are marrow adherent stromal cells (MASCs), also known as bone marrow stromal cells (BMSCs), marrow

- adherent stem cells and mesenchymal stem cells. Exemplary disclosures of MSCs are provided in U.S. patent application publication No. 2003/0003090; Prockop (1997) *Science* **276**:71-74 and Jiang (2002) *Nature* **418**:41-49. Methods for the isolation and purification of MSCs can be found, for example, in U.S. Patent No. 5,486,359; Pittenger *et al.* (1999) *Science* **284**:143-147 and Dezawa *et al.* (2001) *Eur. J. Neurosci.* **14**:1771-1776. Human MSCs are commercially available (*e.g.*, BioWhittaker, Walkersville, MD) or can be obtained from donors by, *e.g.*, bone marrow aspiration, followed by selection for adherent bone marrow cells. See, *e.g.*, WO 2005/100552.
- 10 [0016] MSCs can also be isolated from umbilical cord blood. See, for example, Campagnoli *et al.* (2001) *Blood* **98**:2396-2402; Erices *et al.* (2000) *Br. J. Haematol.* **109**:235-242 and Hou *et al.* (2003) *Int. J. Hematol.* **78**:256-261.

Notch Intracellular Domain

- 15 [0017] The Notch protein is a transmembrane receptor, found in all metazoans, that influences cell differentiation through intracellular signaling. Contact of the Notch extracellular domain with a Notch ligand (*e.g.*, Delta, Serrate, Jagged) results in two proteolytic cleavages of the Notch protein, the second of which is catalyzed by a γ -secretase and releases the Notch intracellular domain (NICD) into the
- 20 cytoplasm. In the mouse Notch protein, this cleavage occurs between amino acids gly1743 and val1744. The NICD translocates to the nucleus, where it acts as a transcription factor, recruiting additional transcriptional regulatory proteins (*e.g.*, MAM, histone acetylases) to relieve transcriptional repression of various target genes (*e.g.*, Hes 1).
- 25 [0018] Additional details and information regarding Notch signaling are found, for example in Artavanis-Tsakonas *et al.* (1995) *Science* **268**:225-232; Mumm and Kopan (2000) *Develop. Biol.* **228**:151-165 and Ehebauer *et al.* (2006) *Sci. STKE* **2006** (364), cm7. [DOI: 10.1126/stke.3642006cm7].
- [0019] Transfection of progenitor cells (*e.g.*, MSCs) with a nucleic acid
- 30 encoding the human Notch intracellular domain, followed by enrichment of transfected cells by drug selection and further culture, results in the production of neural regenerating cells with altered DNA methylation in their genomes. See Example 2, *infra*, for additional details.

Cell Culture and Transfection

[0020] Standard methods for cell culture are known in the art. See, for example, R. I. Freshney "Culture of Animal Cells: A Manual of Basic Technique," Fifth Edition, Wiley, New York, 2005.

5 [0021] Methods for introduction of exogenous DNA into cells (*i.e.*, transfection) are also well-known in the art. See, for example, Sambrook *et al.* "Molecular Cloning: A Laboratory Manual," Third Edition, Cold Spring Harbor Laboratory Press, 2001; Ausubel *et al.*, "Current Protocols in Molecular Biology," John Wiley & Sons, New York, 1987 and periodic updates.

10 [0022] Exemplary methods for transfection and culture are provided in Examples 1 and 2, *infra*.

Methods for Targeted Alteration of DNA Methylation

[0023] Because conversion of progenitor cells to neural regenerating cells is
15 accompanied by changes in the methylation state of certain genes; targeted alteration of methylation state can be used to convert a progenitor cell to a neural regenerating cell.

[0024] Methods for altering the methylation state at a particular C residue are known in the art. To increase the methylation of a particular sequence, fusion
20 proteins comprising a DNA-binding domain and a methylation domain can be used. See, for example, Bestor U.S. 2002/0188103 (Dec. 12, 2002) and WO 97/11972 (April 3, 1997). Exemplary DNA methyltransferase enzymes, which can serve as a source of methylation domains, are disclosed in the aforementioned references. A DNA methyltransferase is a protein which is capable of methylating a particular DNA
25 sequence, which particular DNA sequence may be CpG. This protein may be a mutated DNA methyltransferase, a wild type DNA methyltransferase, a naturally occurring DNA methyltransferase, a variant of a naturally occurring DNA methyltransferase, a truncated DNA methyltransferase, or a segment of a DNA methyltransferase which is capable of methylating DNA. The DNA
30 methyltransferase may include mammalian DNA methyltransferase, bacterial DNA methyltransferase, M.SssI DNA methyltransferase and other proteins or polypeptides that have the capability of methylating DNA.

[0025] Exemplary DNA methyltransferases that can serve as a source of methylation domains for the construction of fusion proteins include, but are not

limited to, cytosine DNA methyltransferases, *dam* methyltransferase, *dcm* methyltransferase, DNMT1, DNMT2, DNMT3a, DNMT3b, CpG methylases, M.SssI, M.CviPI, *HhaI* methyltransferase, *HpaII* methyltransferase, *MspI* methyltransferase, *TaqI* methyltransferase, *BamHI* methyltransferase, *EcoRI* methyltransferase, *HaeIII* methyltransferase, *AluI* methyltransferase, and *SssI* methyltransferase.

[0026] For reducing the degree of methylation of a particular DNA sequence, fusions between a DNA-binding domain and a demethylating domain can be used. Exemplary DNA demethylating domains have been described. See, for example, Bhattacharya *et al.* (1999) *Nature (London)* 397:579-583; Cervoni *et al.* (1999) *J. Biol. Chem.* 274:8363-8366.

[0027] Another exemplary method for reducing the degree of methylation of a sequence of interest is to express, in the cell, a fusion between a DNA binding domain (that binds to the sequence of interest) and a 5-methylcytosine DNA-glycosylase. The fusion protein removes the methylated cytosine base from the DNA sugar-phosphate backbone, to be replaced with cytosine by cellular DNA repair enzymes.

[0028] Demethylation of a DNA sequence of interest can also be achieved by blocking access of maintenance methylases to that sequence during replication; thereby preventing methylation of the unmethylated strand of newly-replicated hemimethylated DNA. A further round of replication will result in daughter DNA duplexes that are unmethylated at the sequence of interest. Such blockage can be achieved by expression in the cell of a zinc finger DNA-binding domain that is engineered to bind to the sequence of interest (see below).

[0029] The activity of a methylation domain or demethylation domain can be targeted to a particular C residue by constructing a fusion protein (or a nucleic acid encoding the fusion protein) comprising a methylation domain and a DNA binding domain, wherein the DNA-binding domain either binds naturally to a sequence at or near the chosen C residue or has been engineered to bind to a sequence at or near the chosen C residue. The DNA-binding domain can be a naturally-occurring DNA-binding domain or a non-naturally-occurring, engineered DNA-binding domain.

[0030] In this regard, the zinc finger DNA-binding domain is useful, inasmuch as it is possible to engineer zinc finger proteins to bind to any DNA sequence of choice. A zinc finger binding domain comprises one or more zinc finger structures. Miller *et al.* (1985) *EMBO J* 4:1609-1614; Rhodes (1993) *Scientific American* February: 56-65; U.S. Patent No. 6,453,242. Typically, a single zinc finger is about

30 amino acids in length and contains four zinc-coordinating amino acid residues. Structural studies have demonstrated that the canonical (C₂H₂) zinc finger motif contains two beta sheets (held in a beta turn which generally contains two zinc-coordinating cysteine residues) and an alpha helix (generally containing two zinc
5 coordinating histidine residues).

[0031] Zinc fingers include both canonical C₂H₂ zinc fingers (*i.e.*, those in which the zinc ion is coordinated by two cysteine and two histidine residues) and non-canonical zinc fingers such as, for example, C₃H zinc fingers (those in which the zinc ion is coordinated by three cysteine residues and one histidine residue) and C₄ zinc
10 fingers (those in which the zinc ion is coordinated by four cysteine residues). Non-canonical zinc fingers can also include those in which an amino acid other than cysteine or histidine is substituted for one of these zinc-coordinating residues. See *e.g.*, WO 02/057293 (July 25, 2002) and US 2003/0108880 (June 12, 2003).

[0032] Zinc finger binding domains can be engineered to bind to a sequence
15 of choice. See, for example, Beerli *et al.* (2002) *Nature Biotechnol.* 20:135-141; Pabo *et al.* (2001) *Ann. Rev. Biochem.* 70:313-340; Isalan *et al.* (2001) *Nature Biotechnol.* 19:656-660; Segal *et al.* (2001) *Curr. Opin. Biotechnol.* 12:632-637; Choo *et al.* (2000) *Curr. Opin. Struct. Biol.* 10:411-416. Zinc finger binding domain are engineered to have a novel binding specificity, compared to a naturally-occurring zinc
20 finger protein. Engineering methods include, but are not limited to, rational design and various types of empirical selection methods. Rational design includes, for example, using databases comprising triplet (or quadruplet) nucleotide sequences and individual zinc finger amino acid sequences, in which each triplet or quadruplet nucleotide sequence is associated with one or more amino acid sequences of zinc
25 fingers which bind the particular triplet or quadruplet sequence. See, for example, U.S. Patent Nos. 6,140,081; 6,453,242; 6,534,261; 6,610,512; 6,746,838; 6,866,997; 7,067,617; U.S. Patent Application Publication Nos. 2002/0165356; 2004/0197892; 2007/0154989; 2007/0213269; and International Patent Application Publication Nos. WO 98/53059 and WO 2003/016496.

30 [0033] Exemplary selection methods, including phage display, interaction trap, hybrid selection and two-hybrid systems, are disclosed in U.S. Patent Nos. 5,789,538; 5,925,523; 6,007,988; 6,013,453; 6,140,466; 6,200,759; 6,242,568; 6,410,248; 6,733,970; 6,790,941; 7,029,847 and 7,297,491; as well as U.S. Patent

Application Publication Nos. 2007/0009948 and 2007/0009962; WO 98/37186; WO 01/60970 and GB 2,338,237.

[0034] Enhancement of binding specificity for zinc finger binding domains has been described, for example, in U.S. Patent No. 6,794,136 (Sept. 21, 2004).

5 Additional aspects of zinc finger engineering, with respect to inter-finger linker sequences, are disclosed in U.S. Patent No. 6,479,626 and U.S. Patent Application Publication No. 2003/0119023. See also Moore *et al.* (2001a) *Proc. Natl. Acad. Sci. USA* 98:1432-1436; Moore *et al.* (2001b) *Proc. Natl. Acad. Sci. USA* 98:1437-1441 and WO 01/53480.

10 [0035] All of the references cited in this section, entitled "Methods for targeted alteration of DNA Methylation," are hereby incorporated by reference in their entireties for the purposes of disclosing exemplary methylation domains and demethylation domains (wild-type and mutant), art-recognized methods for the design, selection and engineering of zinc finger DNA-binding domains, and
15 construction of fusion proteins comprising methylation domains and/or zinc finger DNA-binding domains.

EXAMPLES

Example 1: Preparation of Marrow Adherent Stromal Cells (MASCs)

20 [0036] Bone marrow aspirates, obtained from human donors, were divided into 12.5 ml aliquots in 50 ml tubes, and 12.5 ml of growth medium (10% FBS in α MEM, supplemented with penicillin/streptomycin and 2 mM L-glutamine) was added to each tube. The contents of the tubes were mixed by inversion and the tubes were centrifuged at 200 x g for 8 minutes. The upper, clear phase was discarded, the
25 volume of the lower phase was adjusted to 25 ml with fresh growth medium, and the tubes were again mixed and centrifuged. The upper layer was again removed. The volume of the lower phase in each tube was again adjusted to 25 ml and the contents of all tubes was pooled in a 250 ml tube. After determination of cell concentration by Trypan Blue exclusion and determination of nucleated cell count, cells were plated in
30 T225 flasks, in 40 ml per flask of growth medium at a density of 100×10^6 total nucleated cells per flask. The flasks were incubated at 37°C for 3 days in a CO₂ incubator, during which time the MASCs attached to the flask.

[0037] After 3 days, unattached cells were removed by rocking the flasks and withdrawing the culture medium. Each flask was washed three times with 40 ml of

α MEM supplemented with penicillin/streptomycin; then 40 ml of prewarmed (37°C) growth medium was added to each flask and the cells were cultured at 37°C in a CO₂ incubator. During this time, the medium was replaced with 40 ml of fresh growth medium every 3-4 days, and cells were monitored for growth of colonies and cell density.

[0038] When the cultures achieved 25-30% confluence (usually 10,000-20,000 cells per colony and within 10-14 days), the MASCs (passage M0) were harvested for further passage. MASCs were harvested from up to 10 T-225 flasks at a time. Medium was removed from the flasks and the adherent cells were rinsed with 20ml of DPBS w/o Ca/Mg (DPBS -/-, HyClone) 2 times. Ten ml of 0.25% Trypsin/EDTA (Invitrogen, Carlsbad, CA) was added to each flask and flasks were incubated for approximately 5 min at room temperature. When cells had detached and the colonies had dispersed into single cells, the trypsin was inactivated by addition of 10ml of growth medium followed by gentle mixing. The cell suspensions were withdrawn from the flasks, and pooled in 250ml tubes. The tubes were subjected to centrifugation at 200 x g for 8 minutes. The supernatants were carefully removed and the wet cell pellets were resuspended in growth medium to an estimated cell concentration of approximately 1×10^6 cells/ml. Viable cell count was determined and cells were plated in T225 flasks at a concentration of 2×10^6 cells per flask in growth medium (passage M1). Cells were grown for 3-5 days, or until 85-90% confluent, changing medium every 2 to 3 days. At 85-90% confluence, passage M1 cells were harvested by trypsinization and replated at 2×10^6 cells per T225 flask as described above, to generate passage M2 cultures. M2 cultures were fed fresh medium every three days, if necessary. When passage M2 cultures reached 85-90% confluence (usually within 3-5 days), they were either harvested for transfection to generate NRCs (Example 2 below) or frozen for future use.

Example 2: Preparation of Neural Regenerating Cells (NRCs)

[0039] Neural regenerating cells, also known as NRCs or SB623 cells, were prepared from MASCs harvested from passage M2 cultures, as follows.

A. Preparation of transfection mixture

[0040] Neural regenerating cells were made by transfection of passage M2 MASCs with a plasmid encoding the Notch intracellular domain. The plasmid (pN2) comprised a pCI-neo backbone (Promega, Madison, WI) in which sequences

encoding amino acids 1703-2504 of the human Notch-1 protein, which encode the intracellular domain, were introduced into the multiple cloning site. For each flask of MASCs, 5 ml of transfection mixture, containing 40 μ g of plasmid and 0.2 ml of Fugene 6[®] solution, was used. To make the transfection mixture, the appropriate amount of Fugene[®] solution (depending on the number of flasks of cells to be transfected) was added to α MEM in a sterile 250ml tube, using a glass pipette. The solution was mixed gently and incubated for 5 min at room temperature. The appropriate amount of plasmid DNA was then added dropwise to the Fugene[®]/ α MEM mixture, gently mixed, and incubated for 30 min at room temperature.

- 10 [0041] Prior to the addition of pN2 DNA to the Fugene[®]/MEM mixture, 5ml was removed and placed into a 15ml tube to which was added 40ug of pEGFP plasmid. This solution was used to transfect one flask of cells, as a control for transfection efficiency.

B. Transfection

- 15 [0042] For transfection, passage M2 MASCs were harvested by trypsinization (as described in Example 1) and plated at a density of 2.5×10^6 cells in 40 ml of growth medium per T225 flask. When the cells reached 50-70% confluence (usually within 18-24 hours) they were prepared for transfection, by replacing their growth medium with 35 ml per flask of transfection medium (α MEM +10% FBS without penicillin/streptomycin).

- 20 [0043] Three hours after introduction of transfection medium, 5ml of the transfection mixture (Section A above) was added to each T-225 flask by pipetting directly into the medium, without contacting the growth surface, followed by gentle mixing. A control T-225 flask was transfected with 40 μ g of pEGFP plasmid, for
25 determination of transfection efficiency.

[0044] After incubating cultures at 37°C in transfection medium for 24 hours, the transfection medium was replaced with α MEM +10% FBS + penicillin/streptomycin.

C. Selection of Transfected Cells

- 30 [0045] Cells that had incorporated plasmid DNA were selected 48 hrs after transfection by replacing the medium with 40 ml per flask of selection medium (growth medium containing 100 μ g/ml G-418). Fresh selection medium was provided 3 days, and again 5 days after selection was begun. After 7 days, selection medium was removed and the cells were fed with 40 ml of growth medium. The

cultures were then grown for about 3 weeks (range 18 to 21 days), being re-fed with fresh growth medium every 2-3 days.

[0046] Approximately 3 weeks after selection was begun, when surviving cells began to form colonies, cells were harvested. Medium was removed from the flasks using an aspirating pipette and 20 ml of DPBS without $\text{Ca}^{2+}/\text{Mg}^{2+}$, at room temperature, was added to each flask. The culture surface was gently rinsed, the wash solution was removed by aspiration and the rinse step was repeated. Then 10 ml of prewarmed (37°C) 0.25% Trypsin/EDTA was added to each flask, rinsed over the growth surface, and the flasks were incubated for 5-10 min. at room temperature. Cultures were monitored with a microscope to ensure complete detachment of cells. When detachment was complete, trypsin was inactivated by addition of 10ml of growth medium per flask. The mixture was rinsed over the culture surface, mixed by pipetting 4-5 times with a 10ml pipette, and the suspension was transferred into a sterile 50 ml conical centrifuge tube. Cells harvested from several flasks could be pooled in a single tube. If any clumps were present, they were allowed to settle and the suspension was removed to a fresh tube.

[0047] The cell suspensions were centrifuged at 800 rpm (200 x g) for 8 min at room temperature. Supernatants were removed by aspiration. Cell pellets were loosened by tapping the tube, about 10 ml of DPBS without $\text{Ca}^{2+}/\text{Mg}^{2+}$ was added to each tube and cells were resuspended by gently pipetting 4-5 times with a 10ml pipette to obtain a uniform suspension.

D. Expansion of transfected cells

[0048] Cell number was determined for the suspension of transformed, selected cells and the cells were plated in T-225 flasks at 2×10^6 cells per flask (providing approximately 30% seeding of viable cells). This culture is denoted M2P1 (passage #1). M2P1 cultures were fed with fresh medium every 2-3 days, and when cells reached 90-95% confluence (usually 4-7 days after passage), they were harvested and replated at 2×10^6 cells per flask to generate passage M2P2. When M2P2 cultures reached 90-95% confluence, they were harvested for further assay.

30

Example 3: Comparison of methylation patterns between MASCs and NRCs

[0049] MASCS were prepared from each of three independent human donors (denoted D33, D39 and D41), as described in Example 1 above. A portion of each

preparation of MASCs was used to prepare neural regenerating cells, as described in Example 2, above. Genomic DNA was isolated from each of these six preparations of cells, and for each of the three donors, the methylation state of DNA from neural regenerating cells was compared to that of DNA from their MASC progenitor cells.

5 [0050] Genes whose methylation state were analyzed were selected according to three criteria:

1. known DNA methylation markers for MASCs and mesenchymal cell lines;
2. genes identified as methylation markers for MASCs in a genome-wide screen using differential methylation hybridization; and

10 3. genes reported in the literature to have an effect on embryonic stem cell differentiation.

[0051] For analysis of methylation status, bisulfite sequencing was performed on selected portions (amplicons) of genes selected according to the criteria listed above. Certain genes showed no significant differences in methylation status between
 15 MASCs and NRCs. These genes are listed in Table 1. A number of genes contained amplicons that exhibited differences in methylation status between MASCs and NRCs. These are listed in Table 2. Among these were five genes whose methylation differences were sufficiently significant to be useful in distinguishing NRCs from MASCs. These were PITX2 (also known as Pituitary homeobox 2; RIEG bicoid-
 20 related homeobox transcription factor), ROPN1L (Ropporin 1-like protein; AKAP-associated sperm protein), DNMT3b (DNA C5-N-Methyl Transferase 3b), IGF2R (Insulin-like growth factor 2 receptor) and SDF4 (Stromal cell-derived factor 4). Details of the methylation differences for selected amplicons in these five genes are provided in Tables 3-7.

25 [0052] Tables 3-7 show methylation state at a number of CpG sequences within each amplicon. "Control Cells" refer to MASCs, and "Target Cells" refer to NRCs. Cells were obtained from three different donors, and both MASCs and NRCs were prepared from each donor. SB101 MASCs and SB102 NRCs were from the same donor; SB103 MASCs and SB104 NRCs were obtained from a second donor,
 30 and SB105 MASCs and SB106 NRCs were obtained from a third donor. Each table show results obtained for a different amplicon. Columns 2-7 in each table show methylation levels for particular CpG sites within the amplicon (identified by the number following the colon in Column 1) in MASCs (columns 2-4) and NRCs (columns 5-7). Mean methylation level for each CpG assayed is provided in column

8 for MASCs and in column 9 for NRCs, and the difference in mean methylation level between MASCs and NRCs is shown in column 10.

[0053] Column 11 shows the "Fisher Score" for each CpG sequence assayed.

The Fisher Score is calculated as follows:

5

$$\frac{[\text{Mean methylation value (MASCs)} - \text{Mean methylation value (NRCs)}]^2}{[\text{Standard deviation (MASCs)}]^2 + [\text{Standard deviation (NRCs)}]^2}$$

10 [0054] The Fisher criterion indicates the variability in methylation levels at a particular CpG site. Fisher scores above 1 are considered significant.

[0055] These data show that methylation of CpG sequences in the PITX2, DNMT3b, IGF2R and SDF4 genes is increased in NRCs, compared to MASCs. In contrast, methylation of CpG sequences in the RPON1L gene is reduced in NRCs, compared to MASCs. Similarly, methylation of TMEM179 is decreased. In particular, demethylation of a methylated C residue at position 292 in amplicon 549 represents a significant difference in NRCs, compared to their MASC progenitor cells.

[0056] Accordingly, these methylation changes are diagnostic for NRCs; moreover, achieving the same methylation changes by other means is also useful for preparing NRCs.

Table 1: Amplicons showing no methylation differences between MASCs and NRCs

Amplicon No.	Gene
32	CD4 precursor
38	Collagen α 3 (VI)
105	Collagen α 1 (II)
112	Prolargin precursor
121	Osteopontin
127	BMP4
135	Anexin 6
179	HIF 1A
208	GLI3
308	Keratin 8
475	LRRK 1
488	KCTD 5
509	Frizzled 1 precursor
510	HAND 2
514	ZNF 74
522	PKNOX 2
528	GENSCAN 00000032124
532	Q9C015
537	C15orf27
546	GENSCAN 00000000442
563	GENSCAN 00000003261

Amplicon No.	Gene
824	LIF
827	OCT4
830	DNMT3B
831	IGF2R
832	IGF2R
835	SLUG
836	SLUG
837	PTN
838	PTN
839	ID3
840	ID3
841	ID4
842	ID4
843	SDF4
845	KLF2
846	KLF2
847	P107/RBL1
848	P107/RBL1
849	RELN
850	RELN
851	SST
852	SST

Table 2: Amplicons that are differentially methylated in MASCs and NRCs

Amplicon ID	Gene
18	GDF5
164	FGFR1
227	LPIN1
825	NANOG
826	NANOG
834	NNAT
497	PITX2
549	ROPN1L
829	DNMT3b
833	IGF2R
844	SDF4
1303	TMEM179

Table 3: Methylation changes in the PITX2 Gene

CpG IDs	Control Cells			Target Cells			Mean Methylation		Methylation Difference	Fisher Criterion
	SBI01	SBI03	SBI05	SBI02	SBI04	SBI06	Control	Target		
AMP497:62	1.00	1.00	1.00	0.97	0.97	1.00	1.00	0.98	0.02	1.33
AMP497:73	0.60	0.58	0.64	0.56	0.66	0.94	0.61	0.72	-0.11	0.32
AMP497:82	0.80	0.79	0.80	0.67	0.82	0.98	0.80	0.82	-0.03	0.03
AMP497:105	0.81	0.89	0.85	0.94	0.93	1.00	0.85	0.96	-0.11	3.75
AMP497:109	0.64	0.69	0.77	0.66	0.87	0.96	0.70	0.83	-0.13	0.60
AMP497:135	0.69	0.66	0.78	0.75	0.84	0.82	0.71	0.80	-0.09	1.52
AMP497:139	0.76	0.77	0.83	0.96	0.89	1.00	0.79	0.95	-0.16	5.88
AMP497:142	0.83	0.83	0.84	1.00	0.94	1.00	0.83	0.98	-0.15	17.44
AMP497:153	0.76	0.76	0.88	0.92	1.00	0.97	0.80	0.96	-0.16	4.15
AMP497:172	0.95	0.88	0.96	1.00	1.00	1.00	0.93	1.00	-0.07	2.92
AMP497:184	0.84	0.77	0.92	1.00	1.00	1.00	0.84	1.00	-0.16	4.77

20

Table 4: Methylation changes in the ROPN1L gene

CpG IDs	Control Cells			Target Cells			Mean Methylation		Methylation Difference	Fisher Criterion
	SBI01	SBI03	SBI05	SBI02	SBI04	SBI06	Control	Target		
AMP549:148	0.47	1.00	1.00	0.34	0.72	0.90	0.82	0.65	0.17	0.16
AMP549:167	0.39	0.83	0.83	0.02	0.28	0.80	0.68	0.37	0.32	0.45
AMP549:185	0.09	0.66	0.80	0.00	0.23	0.53	0.52	0.25	0.26	0.33
AMP549:190	0.12	0.66	0.66	0.00	0.43	0.47	0.48	0.30	0.18	0.20
AMP549:249	0.32	0.97	1.00	0.00	0.37	0.65	0.76	0.34	0.43	0.71
AMP549:292	1.00	1.00	1.00	0.38	0.55	1.00	1.00	0.64	0.36	1.25
AMP549:359	1.00	1.00	NA	1.00	1.00	NA	1.00	1.00	0.00	NA

Table 5: Methylation changes in the DNMT3b gene

CpG IDs	Control Cells			Target Cells			Mean Methylation		Methylation Difference	Fisher Criterion
	SBI01	SBI03	SBI05	SBI02	SBI04	SBI06	Control	Target		
AMP829:63	1.00	0.56	0.84	NA	1.00	0.92	0.80	0.96	-0.16	NA
AMP829:67	0.56	0.52	0.63	NA	0.81	0.61	0.57	0.71	-0.14	NA
AMP829:111	0.79	0.50	0.76	0.50	1.00	0.78	0.68	0.76	-0.08	0.07
AMP829:116	0.80	0.58	0.75	0.39	1.00	0.72	0.71	0.70	0.01	0.00
AMP829:127	0.73	0.71	0.76	0.62	1.00	0.70	0.73	0.77	-0.04	0.04
AMP829:134	1.00	1.00	0.76	0.93	0.84	1.00	0.92	0.92	0.00	0.00
AMP829:137	0.51	0.21	0.73	0.73	0.73	0.56	0.48	0.67	-0.19	0.46
AMP829:140	1.00	0.58	0.45	0.60	0.38	0.74	0.68	0.57	0.10	0.09
AMP829:154	0.46	0.21	0.21	0.36	0.23	0.30	0.29	0.30	0.00	0.00
AMP829:158	0.67	0.54	0.65	0.56	0.91	0.74	0.62	0.74	-0.12	0.38
AMP829:186	0.21	0.16	0.14	0.20	0.26	0.26	0.17	0.24	-0.07	1.96
AMP829:216	0.33	0.69	0.00	1.00	1.00	0.80	0.34	0.93	-0.59	2.66
AMP829:298	0.08	0.06	0.00	0.53	0.12	0.00	0.05	0.22	-0.17	0.37
AMP829:337	0.20	0.05	0.00	0.00	0.00	0.00	0.08	0.00	0.08	0.64
AMP829:358	0.03	0.01	0.00	0.17	0.00	0.00	0.01	0.06	-0.04	0.19
AMP829:369	0.34	0.07	0.00	0.08	0.00	0.00	0.14	0.03	0.11	0.35

Table 6: Methylation changes in the IGF2R gene

CpG IDs	Control Cells			Target Cells			Mean Methylation		Methylation Difference	Fisher Criterion
	SBI01	SBI03	SBI05	SBI02	SBI04	SBI06	Control	Target		
AMP833:33	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:44	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:58	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:82	1.00	1.00	0.83	1.00	NA	1.00	0.94	1.00	-0.06	NA
AMP833:102	1.00	0.91	1.00	1.00	1.00	1.00	0.97	1.00	-0.03	0.33
AMP833:104	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:116	1.00	0.94	0.84	1.00	1.00	1.00	0.93	1.00	-0.07	0.82
AMP833:118	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:146	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:155	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:162	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:164	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	NA
AMP833:176	0.82	0.83	0.87	1.00	1.00	1.00	0.84	1.00	-0.16	36.57
AMP833:178	0.72	0.77	0.92	1.00	1.00	1.00	0.80	1.00	-0.20	3.57
AMP833:191	0.77	0.85	0.85	0.97	1.00	0.96	0.82	0.98	-0.15	9.16
AMP833:193	0.87	0.92	0.88	0.97	1.00	1.00	0.89	0.99	-0.10	10.00
AMP833:217	0.76	0.74	0.77	0.86	0.90	0.88	0.76	0.88	-0.12	24.02
AMP833:230	0.74	0.83	0.82	0.94	0.90	0.93	0.80	0.92	-0.13	5.60
AMP833:239	0.59	0.80	0.81	0.92	0.84	0.89	0.73	0.88	-0.15	1.32
AMP833:246	0.64	0.79	0.78	0.89	0.87	0.87	0.74	0.88	-0.14	2.73
AMP833:248	0.66	0.80	0.76	0.83	0.85	0.84	0.74	0.84	-0.10	1.80
AMP833:265	0.65	0.76	0.75	0.87	0.86	0.85	0.72	0.86	-0.14	5.09
AMP833:277	0.65	0.73	0.73	0.87	0.84	0.84	0.70	0.85	-0.15	8.98
AMP833:291	0.52	0.75	0.49	0.71	0.74	0.70	0.59	0.72	-0.13	0.82
AMP833:306	0.48	0.66	0.60	0.76	0.69	0.67	0.58	0.71	-0.13	1.51
AMP833:321	0.46	0.56	0.50	0.70	0.67	0.67	0.51	0.68	-0.17	10.60
AMP833:347	0.47	0.64	0.53	0.48	0.70	0.63	0.55	0.60	-0.06	0.15

AMP833:356	0.45	0.53	0.34	0.61	0.51	0.59	0.44	0.57	-0.13	1.41
AMP833:360	0.41	0.47	0.41	0.61	0.51	0.51	0.43	0.54	-0.11	2.96
AMP833:383	0.40	0.48	0.40	0.64	0.60	0.57	0.43	0.60	-0.18	9.60
AMP833:385	0.34	0.51	0.44	0.53	0.62	0.56	0.43	0.57	-0.14	2.09
AMP833:399	0.53	0.49	0.35	0.62	0.66	0.52	0.46	0.60	-0.14	1.43
AMP833:401	NA	0.42	0.32	0.49	0.61	0.47	0.37	0.52	-0.15	NA
AMP833:416	NA	0.42	0.33	0.60	0.76	0.32	0.38	0.56	-0.19	NA
AMP833:418	NA	0.29	0.33	0.48	0.45	0.21	0.31	0.38	-0.07	NA

Table 6 (continued)

Table 7: Methylation changes in the SDF4 gene

CpG IDs	Control Cells			Target Cells			Mean Methylation		Methylation Difference	Fisher Criterion
	SBI01	SBI03	SBI05	SBI02	SBI04	SBI06	Control	Target		
AMP844:215	0.29	0.00	0.59	0.48	NA	0.68	0.29	0.58	-0.29	NA
AMP844:221	0.52	0.00	0.80	0.72	NA	1.00	0.44	0.86	-0.42	NA
AMP844:224	0.40	0.00	0.51	0.89	NA	0.66	0.30	0.77	-0.47	NA
AMP844:235	0.67	0.00	0.77	0.49	0.59	1.00	0.48	0.69	-0.21	0.18
AMP844:237	0.43	0.00	0.79	0.91	0.66	0.93	0.41	0.83	-0.43	1.03
AMP844:240	0.74	0.00	0.71	1.00	0.55	0.97	0.48	0.84	-0.36	0.53
AMP844:243	0.47	0.00	0.87	0.58	0.15	0.89	0.45	0.54	-0.09	0.03
AMP844:272	0.32	0.13	0.72	0.64	0.38	0.79	0.39	0.60	-0.22	0.35
AMP844:275	0.40	0.07	0.60	0.60	0.06	0.45	0.36	0.37	-0.01	0.00
AMP844:278	0.35	0.12	0.77	0.77	0.57	0.77	0.41	0.70	-0.29	0.71
AMP844:303	0.43	0.30	0.56	1.00	0.31	0.82	0.43	0.71	-0.28	0.54
AMP844:340	0.65	0.42	0.74	0.71	0.27	0.88	0.60	0.62	-0.01	0.00
AMP844:342	1.00	0.77	1.00	0.72	0.29	1.00	0.92	0.67	0.25	0.44
AMP844:375	0.48	0.10	0.41	0.43	0.65	0.61	0.33	0.56	-0.23	1.01
AMP844:393	0.32	0.07	0.65	0.21	0.49	0.51	0.35	0.40	-0.06	0.03
AMP844:417	0.43	0.23	NA	0.64	0.43	0.61	0.33	0.56	-0.23	NA
AMP844:426	0.26	0.12	NA	0.56	0.27	0.58	0.19	0.47	-0.28	NA

Example 4: Changes in methylation state of the ROPN1L gene

[0057] The nucleotide sequences of the amplicons containing changes in methylation status in the ROPN1L gene in NRCs, compared to MASCs, were analyzed to identify precisely the nucleotides whose methylation state was altered. The results of this analysis are presented in Table 8.

Table 8: Methylation changes at specific cytosine residues in the ROPN1L gene

Amplicon number	Sequence in:	
	MASCs	NRCs
549:148	T-T-A-C ^{me} -G-C-C-T	T-T-A-C-G-C-C-T
549:167	T-C-T-C ^{me} -G-G-A-G	T-C-T-C-G-G-A-G
549:185	C-C-T-C ^{me} -G-G-G-G	C-C-T-C-G-G-G-G
549:190	G-G-A-C ^{me} -G-A-T-C	G-G-A-C-G-A-T-C
549:249	C-C-T-C ^{me} -G-G-C-C	C-C-T-C-G-G-C-C
549:292	C-A-T-C ^{me} -G-C-C-C	C-A-T-C-G-C-C-C
549:359	G-T-C-C ^{me} -G-A-T-G	G-T-C-C-G-A-T-G

Example 5: Neural regenerating properties of NRCs with altered DNA methylation

[0058] Neural regenerating cells prepared as described in Example 2, which have the methylation changes described in Examples 3 and 4, are useful in the treatment of various disorders of the central and peripheral nervous systems. See, for example, co-owned WO 2009/023251 (Feb. 19, 2009); the disclosures of which is incorporated by reference in its entirety for all purposes.

[0059] The cells described and characterized in the present disclosure can also be converted, after further treatments, into cells that have the properties of neural cells and neural precursor cells. See, for example, US Patent Application Publication No. 2006/0166362 (July 27, 2006), the disclosure of which is incorporated by reference, which discloses such exemplary treatments, and the properties of the cells so treated. See also US Patent Application Publication No. 2006/0216276 (Sept. 28, 2006), the disclosure of which is incorporated by reference, which discloses additional properties of cells so treated.

CLAIMS

1. An *in vitro* method for altering the methylation state of one or more genes in a progenitor cell, wherein the progenitor cell is a marrow adherent stromal cell (MASC),
5 by the process of:

transfecting the progenitor cell with a polynucleotide comprising sequences encoding a Notch intracellular domain;

wherein the alterations in methylation state comprise:

(a) increased methylation of the PITX2, DNMT3b, Igf2R and SDF4 genes,
10 and

(b) decreased methylation of the ROPN1L and TMEM179 genes.

2. An *in vitro* method for converting a progenitor cell to a cell that stimulates neural recovery and/or neural regeneration after transplantation to a site of nervous
15 system injury or disease, the method comprising altering the methylation state of one or more genes in the progenitor cell, wherein the progenitor cell is a marrow adherent stromal cell;

further wherein the alterations in methylation state comprise:

(i) increased methylation of the PITX2, DNMT3b, Igf2R and SDF4 genes,
20 and

(ii) decreased methylation of the ROPN1L and TMEM179 genes;

further wherein neither the progenitor cell nor any of its descendents were transfecting with a polynucleotide comprising sequences encoding a Notch intracellular domain (NICD).

25 3. The method of claim 2, wherein the methylation state of the PITX2, DNMT3b, IGF2R and SDF4 genes is increased by contacting the progenitor cell with one or more fusion proteins comprising a methylation domain and a DNA-binding domain, or with one or more nucleic acids encoding a fusion protein comprising a methylation domain
30 and a DNA-binding domain, wherein the DNA binding domains are engineered to bind to one or more sequences in each of the PITX2, DNMT3b, IGF2R and SDF4 genes.

4. The method of claim 2, wherein the methylation state of the ROPN1L and TMEM 179 genes is decreased by contacting the progenitor cell with one or more fusion proteins comprising a demethylation domain and a DNA-binding domain, or with one or more nucleic acids encoding a fusion protein comprising a demethylation domain and a DNA-binding domain, wherein the DNA binding domains are engineered to bind to one or more sequences in each of the ROPN1L and TMEM 179 genes.

5. A method for identifying a cell, descended from a marrow adherent stromal cell, that stimulates neural recovery and/or neural regeneration after transplantation to a site of nervous system injury or disease, the method comprising assaying the methylation state of one or more genes in the descendent cell;

wherein the one or more genes is selected from the group consisting of PITX2, DNMT3b, IGF2R, SDF4, ROPN1L and TMEM179;

further wherein:

(i) methylation of the PITX2, DNMT3b, IGF2R and SDF4 genes is increased, and

(ii) methylation of the ROPN1L and TMEM179 genes is decreased.