

**Test methods for determining the intracellular concentration of cyclic
nucleotides**

The invention relates to processes for the quantitative optical analysis of the intracellular concentration of cyclic guanosine 3',5'-monophosphate (cGMP) or cyclic adenosine 3',5'-monophosphate (cAMP). These processes may be useful for identifying and evaluating the pharmacological properties of test substances which influence the activity of receptors or enzymes involved in the synthesis or degradation of the cyclic nucleotides cGMP and cAMP. As a result, the process is suitable, for example, for finding modulators of soluble and membrane-bound guanylate cyclases, phosphodiesterases (PDEs), NO synthases and G-protein-coupled receptors (GPCRs).

The invention describes a process in which the intracellular concentrations of cGMP or cAMP are measured with the aid of recombinant cell lines. These cell lines express recombinantly a combination of particular ion channels activated by cyclic nucleotides (CNG channels), a calcium-sensitive photoprotein and a receptor or an enzyme for which modulators are to be found. The process is suitable for automation and for high-throughput screening (HTS) and ultra high-throughput screening (uHTS) for modulators.

Current processes for measuring the intracellular concentration of cGMP or cAMP have the disadvantage of being very expensive, due to the use of radioactivity, being only partially automatable, if at all, and being very complicated.

Conventional commercial systems for determining cGMP and cAMP are radioactive techniques such as the radioimmunoassay (RIA; e.g. from IBL, Hamburg, Germany), the scintillation proximity assay (SPA; Amersham Bioscience, UK) and the non-radioactive enzymed-linked immunosorbent assay (ELISA; for example from Amersham Bioscience, UK).

The above-described processes cannot be used for measuring the intracellular cGMP and cAMP concentrations in living cells "in situ". To carry out the measurements, the cells need to be disrupted and the cyclic nucleotides then be extracted.

- 5 Another process for measuring intracellular cAMP is the measurement with the aid of inducible promoter systems ("cAMP response element", CRE) and the enzyme luciferase (Goetz et al., J. Biomol. Screen. (2000) 5 377-384). The process has the disadvantage, however, that both transcription and translation need to take place in order for the signal to form, causing the measurement to last for hours. This method
10 is thus not suitable for rapid measurement of the actual cAMP concentrations in cells.

It is the object of the invention to find an improved, non-radioactive process for measuring the intracellular concentrations of cGMP and cAMP. The intention here is
15 to be able to carry out the measurements very rapidly and also with high sensitivity. The process should be automatable and suitable for high-throughput screening of samples or test compounds (HTS and uHTS).

Test compounds in accordance with the invention are preferably small-molecule
20 chemical compounds having a molecular weight of from 100 to 500 or 100 to 1000 or else higher than that. These compounds may be used individually or else as a mixture in the processes according to the invention. Test compounds in accordance with the invention also include antibodies, natural substances, extracts of natural substances, peptides, proteins and nucleic acids. This list is to be regarded only by
25 way of example and not as finite.

In order to establish a screening system for enzymes and receptors which influence the intracellular cAMP or cGMP level, various ion channels activated by cyclic nucleotides (CNG channels) are transfected, alone or in various combinations, into
30 cells which additionally express a calcium-sensitive photoprotein in a recombinant manner. These cells are subsequently transfected with various enzymes or receptors in order to test the suitability of the cells for high-throughput screening. This involves using, for example, soluble guanylate cyclase, since it is capable of

producing intracellular cGMP after stimulation. An example of a receptor which may be used is the β -adrenergic receptor which, after an activation, induces an increase in intracellular cAMP concentration.

- 5 CNG channels are integral membrane proteins which are opened by the cyclic nucleotides cGMP and cAMP. This family of ion channels are non-selective cation channels which are permeable to sodium, potassium and calcium ions. The functional channels are composed of four identical or else different subunits (homo- or heterotetramers). The sensitivities to cGMP and cAMP can be influenced by
10 combining various subunits (Biel et al. TCM (1996) 6, 274-280; Kaupp and Seifert, Physiol. Rev. (2002) 82, 769-824). The channel subunits CNG1-CNG6 which, according to a more recent nomenclature, are also referred to as CNGA1-CNB3, have been cloned and published (Bradley et al., Science (2001) 294, 2095-2096). There exist also splice variants such as CNG4.3, for example (Sauter et al., PNAS
15 (1998) 95, 4696-4701).

- The increase in intracellular cGMP or cAMP concentration results in the opening of the CNG channels and thereby in an influx of calcium ions. The calcium flowing in may then be detected with the aid of calcium-sensitive fluorescent indicators, such
20 as, for example, FURA, Fluo-3 etc., or with the aid of calcium-sensitive photoproteins such as aequorin or obelin.

- The photoprotein aequorin which is derived from the jellyfish *Aequorea victoria* is a highly sensitive calcium indicator in cells. The aequorin complex consists of the
25 protein apoaequorin, molecular oxygen and the luminophore coelenterazine. This complex emits blue light in the presence of calcium (Ca^{2+} ions), with maximum emission at 466 nm (Jones et al., Trends Biotechnol. (1999) 17, 477-481).

- Another photoprotein is obelin which has been cloned from various hydrozoa such as
30 *Obelia longissima*, for example. It consists of apoobelin, O_2 and the luminophore coelenterazine. Likewise, blue light is emitted in the presence of calcium (Illarionov et al., Methods Enzymol. (2000) 305, 223-49).

Guanylate cyclases are responsible for producing cGMP from GTP. A distinction is made between soluble and membrane-bound guanylate cyclases. Soluble guanylate cyclase is a heterodimer composed of an α 1 subunit and a β 1 subunit. It is the natural receptor for nitrogen monoxide (NO) produced by the endothelial NO synthase and can therefore also be activated pharmacologically by NO-releasing substances such as, for example, SIN-1. Since the cGMP produced by guanylate cyclase induces relaxation of blood vessels, the enzyme is a highly interesting pharmacological target (Hobbs AJ., TIPS (1997) 18, 484-491). The membrane-bound guanylate cyclases possess a transmembrane segment and are activated by various agonists such as ANP, BNP, CNP or guanylin (Wedel and Garbers, Annual Rev. Physiol. (2001) 63, 215-233).

Another pharmacologically interesting protein family are the "G-protein-coupled receptors" (GPCRs). These are integral membrane proteins which can transduce the action of an extracellular hormone into the cell interior. The intracellular levels of cAMP or calcium are influenced via coupling of these receptors to various G proteins. The activation of "Gq-coupled receptors" results in a release of calcium from internal stores and thus in an increase in cytoplasmic calcium concentration. The activation of "Gs-coupled receptors" results in an increase in cAMP concentration, whereas activation of the "Gi-coupled receptors" reduces the intracellular cAMP content (Gurrath M., Curr. Med. Chem. (2001) 8, 1605-1548). An example of Gs-coupled receptors are the " β -adrenergic receptors" which can be stimulated with the aid of agonists such as isoprenaline (Dzimiri N., Pharmacol. Rev. (1999) 51, 465-501).

25

The cyclic nucleotides cGMP and cAMP produced by guanylate cyclases and GPCRs, respectively, are hydrolyzed by "phosphodiesterases" (PDEs) to give GMP and AMP, respectively. The phosphodiesterases are an enzyme family whose members are distinctly different with regard to substrate specificity, regulation and expression patterns in the body (Francis et al., Prog. Nucleic Acid Res. Mol. Biol. (2001) 65, 1-52). PDEs play a decisive part in controlling the intracellular cAMP and cGMP levels and are therefore likewise important targets for pharmacological intervention.

30

Surprisingly, combining the CNG3 channel (SEQ ID NO: 5, Acc. No. X 89600) or, in particular, the CNG2 channel (SEQ ID NO: 1, Acc. No. X55010) with a photoprotein proved to be particularly suitable for detecting changes in intracellular cGMP concentrations (Figure 1).

A similar surprise was a combination of CNG2/CNG4.3/CNG5 (SEQ ID NO: 1/ SEQ ID NO: 2, Acc. No. AJ000515/SEQ ID NO: 3, Acc. No. U12623) and a photoprotein, which proved to be very suitable for detecting changes in intracellular cAMP concentration (Figure 3). The combination of CNG2(T537A)/CNG4.3/CNG5 (SEQ ID NO: 4/ SEQ ID NO: 2/ SEQ ID NO: 3) with a photoprotein proved to be particularly suitable for cAMP measurement. In the CNG2 mutant CNG2(T537A) (SEQ ID NO: 4), the amino acid threonine-537 has been replaced with alanine (Altenhofen et al., PNAS (1991) 88, 9868-9872).

The recombinant cell lines may be prepared using common vectors such as pcDNA1.1 Amp or pcDNA3 (Invitrogen Life Technologies). Common transfection reagents such as, for example, lipofectamine (Invitrogen Life Technologies) may be used for transfecting the corresponding plasmid constructs.

The cells are seeded for the measurements on 384-well microtitre plates (MTPs) at 1500 cells per well and on 1536-well MTPs at 250 cells per well. After 1-2 days of growth at 37°C/5% CO₂, the cell culture medium (DMEM/F12 containing 10% foetal calf serum) is removed and the cells are charged with coelenterazine (0.8 µg/ml) in calcium-free Tyrode's at 37°C/5% CO₂ for 3 h. Subsequently, test substances and appropriate control substances (e.g. SIN-1 or isoprenaline) are added in calcium-free Tyrode's and incubated on the cells for 5-10 min. The extra measurement is then carried out in a light-tight box by adding calcium-containing Tyrode's via a comb (final calcium concentration: 3 mM), with the aid of a charge-coupled device camera.

To measure the effects of substances directly, the cells are charged with coelenterazine in calcium-containing Tyrode's. The substances are then added within

the light-tight box and measurement commences immediately with addition of the substances (Figure 5).

5 The advantages of this process are the high sensitivity of the measurement, the low costs per measuring point and the suitability for HTS and uHTS. The process has an excellent signal-to-noise ratio, and it is possible to achieve stimulation factors of 50-150. It is moreover possible to apply test substances for very short periods of time, since a signal can already be observed after a few seconds (Figure 5). This has the advantage of unspecific effects of test substances (e.g. due to cytotoxicity) on the
10 cells used being minimized. The process is moreover suitable for characterizing "orphan receptors" (receptors whose natural ligand is unknown), since it is possible to observe both changes in the cAMP level (Gs coupling and Gi coupling) and changes in intracellular calcium concentration (Gq coupling) by means of luminescence or fluorescence measurement.

15

The invention relates to processes for determining the intracellular concentration of cyclic nucleotides, characterized in that a cell expressing a CNG channel together with a photoprotein is prepared and used, and the intracellular concentration of cyclic nucleotides is determined by means of the luminescence signal of the photoprotein.

20

The invention likewise relates to processes as described above, wherein the CNG channel is the CNG2 or the CNG3 channel.

25

The invention comprises processes as described above, wherein the cyclic nucleotide is cGMP.

30

The invention likewise relates to processes as described above, wherein the cyclic nucleotide is cAMP and a cell expressing a combination of the CNG2/CNG4.3/CNG5 channels is prepared and used.

The invention also relates to the above-described processes, wherein the cyclic nucleotide is cAMP and a cell expressing a combination of the CNG2(T537A)/CNG4.3/CNG5 channels is prepared and used.

The invention relates to the processes as described above, wherein the photoprotein is aequorin.

- 5 The invention also relates to processes as described above, wherein the photoprotein is obelin.

The invention relates to a method of screening test compounds for identifying receptor ligands, wherein any of the above-described processes is used in which the
10 cell used expresses the receptor and possesses an intracellular messenger system which allows a receptor-ligand bond to cause measurable modulation of the ion flow through the ion channel, the said cell is incubated with test substances of which those modulating the luminescence are selected.

- 15 The invention also relates to a method of this type, wherein the receptor is a G-protein-coupled receptor.

The invention also relates to a corresponding method, wherein the G-protein-coupled receptor is an orphan receptor.

20

The invention furthermore relates to a method of screening test compounds for identifying modulators of phosphodiesterases, wherein any of the above-described processes is used in which the cell used expresses the phosphodiesterase, the said cell is incubated with test substances of which those modulating the luminescence are
25 selected.

The invention likewise relates to a method of screening test compounds for identifying modulators of guanylate cyclases, wherein any of the above-described processes is used in which the cell used expresses the guanylate cyclase, the said cell
30 is incubated with test substances of which those modulating the luminescence are selected.

The invention also comprises a method of screening test compounds for identifying modulators of NO synthase, wherein any of the above-described processes is used in which the cell used expresses the NO synthase and soluble guanylate cyclase, the said cell is incubated with test substances of which those modulating the luminescence are selected.

The invention also relates to a corresponding method, wherein the NO synthase and the soluble guanylate cyclase are expressed in different cells.

The invention also relates to a cell prepared by any of the processes illustrated above.

Description of the figures

Figure 1: Schematic representation of the process for determining intracellular cGMP concentration.

Figure 2: Luminescence measurement after activation of soluble guanylate cyclase with SIN-1. The "relative light units" (RLUs) were measured for 1 min. The amount of light measured at the SIN-1 concentration indicated over this period is shown.

Figure 3: Schematic representation of the process for determining intracellular cAMP concentration.

Figure 4: Luminescence measurement after activation of the β -adrenergic receptor with isoprenaline. The incubation time with isoprenaline is 10 min in calcium-free Tyrode's. Measurement is started by adding calcium. The relative light units (RLUs) are measured in second intervals. The time course of the luminescence signal after addition of calcium and previous incubation with 0 M (stars), 10^{-7} M (squares) and 10^{-6} M (triangles) isoprenaline is shown.

Figure 5: Kinetics of the luminescence signal after activation of the β -adrenergic receptor with isoprenaline in calcium-containing Tyrode's. The measurement is started with the addition of the agonist isoprenaline (10^{-6} M) within the light-tight box.

5

Examples

10 If, for example, soluble guanylate cyclase is additionally expressed in a cell line which expresses, as a cGMP measuring system, the CNG2 channel (SEQ ID NO: 1) together with a calcium-sensitive photoprotein, it is possible to increase the intercellular cGMP concentration with the aid of the guanylate cyclase stimulator SIN-1 in a dose-dependent manner and to detect an enhanced luminescence signal (Figure 2).

15

If, for example, the β -adrenergic receptor is additionally expressed in cells which, as a cAMP-measuring system, express a calcium-sensitive photoprotein together with the CNG-channel subunits CNG2(T537A)/CNG4.3/CNG5 (SEQ ID NO: 4/SEQ ID NO: 2/ SEQ ID NO: 3), activation of this receptor by means of isoprenaline results in
20 a dose-dependent manner in an increased cAMP concentration and an enhanced luminescence signal (Figure 4). If the cells are charged with coelenterazine in calcium-containing Tyrode's, the agonist isoprenaline may also be added directly to the cells within the light-tight box. A luminescence signal will then be observed within a few seconds (Figure 5).

Claims

1. Use of CNG2 and a photoprotein for the measurement of cGMP.
2. Use of claim 1, wherein Aequorin is used as a reporter gene.

SEQUENCE LISTING

<110> Bayer AG, BHC

<120> Test methods for determining the intracellular concentration of cyclic nucleotides

<130> Le A 36 621

<160> 5

<170> PatentIn version 3.1

<210> 1

<211> 3166

<212> DNA

<213> Bos taurus

<400> 1

gctcacctac	tggcaggctg	gggtgtgcag	gccccacagt	gagagggtca	gacatccagc	60
cagtgggcag	aactgtttct	ccagtgggca	agtgtgccc	tgggccgagg	cctcgacctc	120
gcctcagctg	taggcatggg	ctcctctgag	tactgccc	cctccccctgc	tggcctcagg	180
cagggcgggc	agctctttga	ctgagaggct	gagaagctgc	gtggggattc	ggggatcccc	240
ctgggtctggg	gacaggagct	ttgggaggtc	tctactgtcc	cttggtgctg	atgagctccc	300
gagggttctg	ctctgactgg	aaagcttctg	gacagacctc	tagcctgtgg	ccaagggaca	360
ccctgcctcg	gaatccgtct	ggtgaggaaa	ggtgagggtc	ctggttgtac	atggaggatg	420
acagaaaaag	ccaatggcgt	gaagagctcc	ccagccaata	accacaacca	ccatgccctc	480
cctgccatca	aggccagtgg	caaagatgac	cacagggcc	gcagccggcc	acagtctgct	540
gctgtgatg	acacctctc	agagctacag	caactggcag	agatggatgc	ccccagcag	600
aggaggggtg	gcttcgcag	gattgccgc	ctgggtgggg	tcctcagaga	gtgggcttac	660
aggaacttcc	gtgaggagga	gcctagacct	gactcattcc	ttgagcggtt	ccgggggcct	720
gagctccaca	ccgtgacaac	acaacaagga	gacggcaaag	gcgacaagga	cggcgagggc	780
aagggcacca	agaagaagtt	tgaactcttt	gtcttgacc	cagccgggga	ctggtactac	840
cgctggcttt	ttctcattgc	cttgcccgtc	ctctacaact	ggtgcctatt	ggtggccaga	900
gcctgcttca	gtgacctgca	gaaaggctac	tacatagtgt	ggctgggtct	ggattacgtc	960
tcagatgtgg	tctacatcgc	agacctctc	atccgactgc	gcacaggttt	cttgagcag	1020
gggctactgg	tgaagacac	caagaagttg	cgggacaact	acatccacac	catgcagttt	1080
aaagtgttgg	tggcctccat	catcccctaca	gacctgatct	atcttgctgt	gggatccat	1140
aaccctgagg	tgcgcttcaa	ccgctgcta	cactttgccc	gcattgttga	gttctttgac	1200
cgcactgaga	cacgcaccag	ctaccccaac	atcttccgaa	taagcaacct	gatectctac	1260
atcttgatca	tcattcactg	gaatgcctgc	atctactatg	ccatctccaa	gtccatcggc	1320
tttggggtag	acacctgggt	ttaccccaac	atcactgacc	ctgagtatgg	ctacctgtct	1380
aggaggtaca	tctattgcct	ttactggtct	acactgacct	tcaccaccat	tggggagaca	1440
ccacccctg	taaaggatga	ggagtacctg	tttgtcatct	ttgacttcct	gattggtgtc	1500
ctcatccttg	ccaccatcgt	gggaaatgtg	ggctccatga	tctccaacat	gaatgccacc	1560
cgggtgagt	tccaggccaa	gattgatgct	gtcaaacatt	atatgcagtt	ccgaaaggtc	1620
agcaaggaga	tggaaagcaa	ggtcattagg	tggtttgact	acttgtggac	caataagaag	1680
agtgtagatg	agcgagaagt	cctcaaaaac	ctgccagcaa	agctcagggc	tgagatagcc	1740
atcaacgtcc	acctgtccac	actcaagaaa	gtgcgcattc	ttcaggactg	tgaggctggc	1800
ctgctggtgg	aactggtatt	aaagctccgg	cctcaggctc	ttagccctgg	ggactacatt	1860
tgcgcgaagg	gggatattgg	gaaggagatg	tacataatca	aggagggaaa	attggcagtg	1920
atggctgatg	acggtgtcac	tcagtatgcc	ctgctctcgg	ctgggagttg	ctttggagag	1980
gcagtatcc	ttaatatata	gggcagcaaa	atgggcaatc	ggcgacagc	caacatccgc	2040
agtcttggtc	actctgatct	gttctgcttg	tccaaggatg	atcttatgga	agctgtgact	2100
gagtaccctg	atgccaagag	ggtcttgagg	gagagaggcc	gggagattct	gatgaaggag	2160
ggcttggttg	atgagaatga	ggtggcagcc	agcatggagg	tagatgtgca	ggaaaagcta	2220
gaacagctgg	agaccaacat	ggacaccttg	tacactcggt	ttgccgcct	gctggccgag	2280
tacacgggag	cccagcagaa	gctcaagcag	cgcatacacag	ttttggaaac	gaagatgaag	2340
cagaataatg	aggatgactc	cctgtcagat	gggatgaaca	gcccagagcc	acctgccgag	2400
aagccataat	ggcttgcccc	aatgccctc	cagccttg	ctttgacccc	agggttgga	2460
gagctgtgta	ggtccccaca	tatatatgca	ttaccacatc	cccttgaatt	ctcccagaag	2520
cctctctgct	ggaaggttta	gggtctgac	atccagaagc	cctcctccaa	gtccgactaa	2580
cagctaattct	tgtgcagggc	atagactgtg	cttagctcgg	cttcagaag	cttcagcctg	2640
tctaagtttg	aggaagaaa	aaaagaggag	catctctcca	ggctcttttg	catctagtta	2700
cctcctactt	gattcttttc	taatattgtg	tctgaatatt	tccatttccc	tgcagcagta	2760
tgtagttaga	acactggctg	cagacaccca	gcactggctc	cagtgtcctt	ttcccaaggc	2820
aggcaaaagg	tgtggagggg	ggcaaggagg	agatgtcac	tccaagtctc	gccgtgctga	2880
ttcctccgcc	tgtctgcca	ccagagtg	gagccttg	gtcttttctg	aaccagggg	2940
aggaggtatg	tcctggtctc	caatccatcc	caggacatgg	agtgaggaa	tagcagttgg	3000
ccagcaggca	aggcacctgg	agaagggtgg	gggcaggagc	ctggccatca	cccctctatg	3060

cagagtgttt	ctcaggaggc	cctgaggctg	atgggtgggtg	gatgactctt	caagttaaca	3120
tctgcagtag	agacacttac	aagttaataa	attcctctga	actttt		3166

<210> 2
 <211> 3236
 <212> DNA
 <213> Rattus norvegicus

<400> 2						
caccagctcc	atgtgtgtgcc	aatcacagcg	tcttagaagc	tgctgtcctg	gctgacccag	60
ggcttctccc	ggctccactc	atatgtcca	gtcccaggct	tggtatcact	gtcacagccc	120
caggctgacc	tactgcaaca	tggaccttcc	taccagcaga	gcccgaaga	cccaatgtcc	180
tgctggatag	ctgtttggtg	gtgcaggctg	atgtggacca	gtgccagcta	gaaagggcac	240
agccagagac	agcatcgatc	caggagttac	cagaagaaga	agaggagaag	gaggagaaga	300
agaaggagga	ggaagaggag	aaggaggagg	aggaagagaa	ggaagaggaa	gaggagaagg	360
aggaggaggg	ggaggccaca	aactcaacag	tccccaggat	cacgcctctg	cctgacacct	420
acgggaccca	gtacatggc	taaaccaggc	ctgaacatgg	agctgaatcg	cctgggtccag	480
gaccagccac	ctggccagga	gggccccaga	cctggccccc	caaaccacgc	cgagcacctc	540
cccaatgtgc	ccagctaccg	tcccgcaccc	acccgcaccc	ccgtcctcgt	ctcccgagg	600
acagccttgt	ccaactccaa	cttcaccaag	gagatcagga	gctccatccg	tcgcctggtg	660
ccagccacga	aagagcaccc	ggagctccag	gtggaagaca	cagatgccga	agctggcccc	720
ctcatcccag	aggagacgat	cccggcacct	gagagaccac	cagtgtctcc	cgccaagtct	780
gacacccctg	cggttcccag	cgcagcaacc	cacaggaaga	agctaccttc	tcaggatgat	840
gaggctgaag	aactcaaggc	cctgtcaccc	cagtgggtcc	cgtgtgtg	ctgggtccag	900
cccacccccc	cacaggaggc	tgatggcgag	gaccgtgcgg	cctccacagc	cagccagaac	960
agtgccatca	tcaacgaccg	gctccaggag	ctggtgaaga	tggtcaagga	gcggaacagag	1020
aagggtgaag	agaagctcat	tgaccctgac	gtcacctccg	atgaggagag	ccccaaagccc	1080
tccccagcca	agaaggcccc	agactcagcc	ccagcccgga	agccggcgga	ggcagaggcg	1140
gcagaggagg	agcactactg	tgacatgtct	tgctgcaagt	tcaagcgag	gccctggaag	1200
atgtaccagt	tccccagag	catcgaccca	ctgaccaacc	tcagtgtacat	cctgtggtg	1260
ttcttcgtg	tgctggcctg	gaactgggaa	tgctggctga	ttcctgtg	ctgggccttc	1320
ccgtaccagc	gggcagacaa	catccacctc	tggtgtctca	tggtactact	gtgcgacttc	1380
atctacctcc	tgacatcac	cgtgttccag	atgcgtctcc	agtttgtcaa	aggcggggac	1440
atcattacag	ataagaagga	gatgcgtaat	aattacctga	agtctcaacg	atttaagatg	1500
gacttgctct	gccttttgcc	tttggtattt	ctctacttga	aacttgccgt	gaacccccct	1560
cttcgcctgc	cccgtgcct	gaagtacatg	gccttctttg	agtttaataa	ccgtctggaa	1620
gccatccctca	gcaaagccta	cgtttacagg	gttatcagga	ccaccgccta	cctgtctgtat	1680
agcttgcatc	tcaactcctg	tctttactac	tgggcgctcg	ccttccaggg	catcggttcc	1740
actcactggg	tttatgacgg	cgtggggaac	agctacattc	gatgtactac	ctgggctgtg	1800
aaaactctca	tcaccatcgg	aggactgccc	gacccccaga	cgctctttga	gatcgtcttc	1860
cagctgtctga	attattttac	agggtgtctc	gctttctctg	tgatgattgg	acagatgaga	1920
gatgtggtgg	gggcccgcac	ggcagggcag	acgtactacc	gcagctgcat	ggacagcacg	1980
gtgaagtaca	tgaacttcta	caagatcccc	aggtctgtgc	agaaccgcgt	caagacctgg	2040
tatgagtaca	cctggcattc	acaaggcatg	ctggatgagt	cagagctgat	ggttcagctt	2100
ccggacaaga	tgctgtctga	cctggccatt	gacgtaaact	acaacattgt	cagcaaatgt	2160
gcgtctcttc	agggtcgcga	ccggcagatg	atcttcgaca	tgctcaagcg	acttcgtcca	2220
gtcgtctacc	tacccaatga	ctatgtgtgc	aagaaggggg	agattggccg	agagatgtat	2280
attatccagg	cggggcaggt	gcaggtgtct	ggcgccccag	atggaaaggg	tgctcctggt	2340
acactcaaag	ccggtatcgt	gtttggagag	ataagcttgc	tggtgtcg	ggcggtaac	2400
aggcgacag	ccaatgtggt	ggcccacggc	ttcaccaatc	tcttcattct	ggataagaag	2460
gacttgaatg	agatttttgt	gcattaccct	gaatctcaga	agctgtccg	gaagaaggcc	2520
aggcgcatgc	tcagaaacaa	caacaaaccc	aaggaggaga	agagtgtgct	catcctgccc	2580
ccagctgcgg	gcaccccga	gctcttcaat	gctgcctgg	ctgcagcagg	aaagatgggc	2640
cccaggggag	ccaaggcg	caagctcgcc	cacctgagag	ccaggctcaa	agaactggct	2700
gcactggagg	cagccgcacg	acagcagcag	ctgctggaac	aggccaagag	ctcgcaagaa	2760
gccgggggag	aggagggtc	tgagccaca	gaccaacctg	caccccagga	gccgtcagag	2820
cccaaggagc	ccccggagcc	cccagccccg	agttctccac	cgccagcctc	agcaaagccc	2880
gagggaagca	cggaggaggc	cgcaggggcg	ccggagcctt	cagtgaggat	ccgtgtgagt	2940
ccaggccctg	atcccgggg	acagacacta	tcggtggaga	tgctggaaga	gaagaaggag	3000
gaggtggagt	gaggtgcatt	cagcacctg	atcccatccc	taggtgcgga	ccagaccttt	3060
gaaccacctg	cctcaccaag	catcttagcc	taaaggcgcc	ttctcaccgg	cttatagaaa	3120
ctctcaaatt	tcacaaagta	gtggtggtga	cctcgagctc	ctaacattcc	cggagcaccc	3180
tgctccccctc	tcacagttgc	tattttctag	aagaaaaccg	ccttgacctt	taaaaa	3236

<210> 3
 <211> 2351
 <212> DNA
 <213> Rattus norvegicus

<400> 3
 cgctggagcc cgagggcagc ctcttcacag aagccaggca ctttcccaag ccacctataa 60
 tcggagaatc actggatagt ctacagacta gaaccacact acggaaacat gagccaggac 120
 ggcaaaagtga agaccacaga gtccacgccc ccagcccca ccaaagccag gaagtggctg 180
 ccagtcctag acccatctgg ggattactac tactgggtgg tgaacacaat ggtcttccca 240
 atcatgtaca acctcatcat cgttgatgac agggcctgct ttcttgactt gcagcacagt 300
 tacctggtag cctggttcgt gctggactac acgagtgacc tgctgtacct actagatatt 360
 ggggtacgct tccacacagg attcctagag cagggcatcc tgggtgtaga caaaggcatg 420
 atcgccagtc gctacgtccg cacctggagc ttctgtttgg acctggcgct cctggtcccc 480
 acagatgcgg cctatgtgca gctgggcccc cacatcccta cactccggct aaaccgcttt 540
 ctccgagtcg cccgcctctt cgaggctttt gatcgtagag agaccgcac ggcttaccca 600
 aatgccttcc gcatagccaa cctgatgctt tacatttttg ttgtcatcca ttggaacagt 660
 tgcttatact tcgccctgtc caggtaacct ggctttggac gggatgcgtg ggtataccca 720
 gacctgcgc aacctggctt tgagcgcttg cggcgccagt atctctacag cttctacttc 780
 tccactctga tctgaccac agtgggtgac acgcccgtgc cagaccgaga ggaagagtac 840
 ctcttcatgg tgggtgactt cctgctggcc gtcattgggt tcgccaccat catgggtagc 900
 atgagctctg tcatttaca catgaacact gcagatgcgg ccttctaacc agaccatgcg 960
 ctggttaaaga agtacatgaa gctgcagcat gtcaacaaga ggctggagcg gcgagttatt 1020
 gactggtacc agcatcttca gatcaacaag aagatgacca acgagtagc catcttgacg 1080
 cacctgcctg agcggctgcg ggccggaggt gctgtgtccg tgcacctgtc taccctgagc 1140
 cgagtacaga tcttcagaa ctgtgaagcc agcctgctgg aagagctggg gctgaagcta 1200
 cagccccaga cctactcgcc aggcgaatat gtgtgccgca aaggggacat tggccgagag 1260
 atgtacatca tccgtgaggg ccagctggct gtggtggccg atgatggtg cacacagtat 1320
 gctgtgcttg gtgcagggtc ctactttggg gagatcagta tcatcaacat caaagggaac 1380
 atgtctggaa accgacgaac agccaacatc aagagcctag gttattcaga cctgttctgc 1440
 ctacgaagg aggatctgcg ggaggtactg agtgagtacc cacaggccca ggccgtcatg 1500
 gaggagaagg gccgagaaat cttgctcaaa atgaataagt tggatgtgaa tgcgtgagca 1560
 gctgagatcg cctccagga ggccacagag tctcggctca aaggcctcga ccagcagctt 1620
 gatgatctgc agaccaagt tgctcgcta ctggctgagc tggagtccag tgcactgaag 1680
 atagcttacc gcatcgaag gctggagtgg cagactcgag agtggccaa gccagaggac 1740
 atgggtgagg ctgatgatga ggctgagcct ggagaaggga cgtccaagga tggagaggga 1800
 aaggctggcc aggggggacc ctacggcata gaatgacccc gtcctgaccc taggactccc 1860
 agctcaaata aatccagagt ggtgggaaa tctgcctgca ggaactgtca tctctttgc 1920
 taggttacag aacgtagta aattggtcta tagatgccta gctaggatg tgggtcacag 1980
 gatccattag tccatactc accagcaaat gtacacacac acacacacac aaacatgcgc 2040
 acgcacacgc tcaagactga gttctacata atgttctctg cttctgcaca catgtgcacc 2100
 atgatcacag agaacaattg aatgcactgg gactcttgag gtaagcttta cacatcttaa 2160
 aggaggtctg ctggttttag gcaaggacag tgttgagct acagggaaa tagacagata 2220
 caattcctga actctaggtc accacttcag ctgcttcagg gtatggccc acttcagggtc 2280
 tgagctggga caggcggtat tttgaaatga acctgatgtg tgatttatta ataaaatata 2340
 aggttcataa c 2351

<210> 4
 <211> 1992
 <212> DNA
 <213> mutated CNG2

<400> 4
 atgacagaaa aagccaatgg cgtgaagagc tccccagcca ataaccacaa ccaccatgcc 60
 cctcctgcca tcaaggccag tggcaaagat gaccacaggg ccagcagccg gccacagtct 120
 gctgctgctg atgacacctc ctacagacta cagcaactgg cagagatgga tgccccccag 180
 cagaggaggg gtggcttccg caggattgcc cgcctgggtg gggctcctcag agagtgggct 240
 tacaggaact tccgtgagga ggagcctaga cctgactcat tccctgagcg tttccggggg 300
 cctgagctcc acaccgtgac aacacaacaa ggagacggca aaggcgacaa ggacggcgag 360
 ggcaagggca ccaagaagaa gtttgaactc tttgtcttgg accagccgg ggactggtag 420
 taccgctggc tttttctcat tgccttgccc ctgctctaca actgggtgct attgtgggcc 480
 agagcctgct tcagtgcact gcagaaaggc tactacatag tgtggctggg gctggattac 540
 gtctcagatg tgggtctacat cgcagacctc ttcatccgac tgcgcacagg tttcttgagg 600
 caggggctac tgggtgaaaga caccaagaag ttgcgggaca actacatcca caccatgcag 660
 tttagctgg atgtggcctc catcatccct acagacctga tctattttgc tgtggggatc 720
 cataaccctg aggtgcgctt caaccgcctg ctacactttg cccgcatggt tgagttcttt 780
 gaccgcactg agacacgcac cagctacccc aacatcttcc gaataagcaa cctgatcctc 840
 tacatcttga tcatcattca ctggaatgcc tgcatctact atgccatctc caagtccatc 900
 ggctttgggg tagacacctg ggtttacccc aacatcactg accctgagta tggctacctg 960
 tctaggagat acatctattg cctttactgg tctacactga ccctcaccac cattggggag 1020

acaccacccc	ctgtaaagga	tgaggagtag	ctgtttgtca	tctttgactt	cctgattggt	1080
gtcctcatct	ttgccacat	cgtgggaaat	gtgggctcca	tgatctccaa	catgaatgcc	1140
acccgggctg	agttccaggc	caagattgat	gctgtcaaac	attatatgca	gttccgaaaag	1200
gtcagcaagg	agatggaagc	caaggtcatt	aggtgggttg	actacttggt	gaccaataag	1260
aagagtgtag	atgagcgaga	agtccctcaa	aacctgccag	caaagctcag	ggctgagata	1320
gccatcaacg	tccacctgtc	cacactcaag	aaagtgcgca	tctttcagga	ctgtgagggt	1380
ggcctgctgg	tggaaactgg	attaaagctc	cggcctcagg	tcttttagccc	tggggactac	1440
atttgccgca	agggggatat	tgggaaggag	atgtacataa	tcaaggaggg	aaaattggca	1500
gtggtggctg	atgacgggtg	cactcagtat	gccctgctct	cggctgggag	ttgctttgga	1560
gagatcagta	tccttaatat	taagggcagc	aaaatgggca	atcggcgcgc	agccaacatc	1620
cgcagtcttg	gctactctga	tctgttctgc	ttgtccaagg	atgatcttat	ggaagctgtg	1680
actgagtacc	ctgatgccaa	gaggttcttg	gaggagagag	gccgggagat	tctgatgaag	1740
gagggcttgt	tggatgagaa	tgagggtggc	gccagcatgg	aggtagatgt	gcaggaaaag	1800
ctagaacagc	tggagaccaa	catggacacc	ttgtacactc	gttttgcccg	cctgctggcc	1860
gagtacacgg	gagcccagca	gaagctcaag	cagcgcatca	cagttttgga	aacgaagatg	1920
aagcagaata	atgaggatga	ctccctgtca	gatgggatga	acagcccaga	gccacctgcc	1980
gagaagccat	aa					1992

<210> 5
 <211> 2641
 <212> DNA
 <213> Bos taurus

<400> 5						
gtcgtcatgg	cggaggatgc	tgtgcgaggt	ggatggcagc	cttctcccca	ggaccctcca	60
ggtccgagtc	tgtggaggcc	tcagcaactg	gagtagtggg	acactgccat	tcagcagcca	120
ccttgggggtg	gaatgtgaca	cagcaagaag	atggccaaga	ttagcaccca	atactccac	180
cccaccagga	cacacccctc	ggtcaggacc	atggacagag	atctggattg	cattgaaaac	240
ggtctcagca	ggaccacatt	gccatgcgag	gagacatcgt	cagaactgca	ggaaggcatt	300
gccatggaga	ctagaggact	ggctgaatcc	aggcaaagct	ccttcaccag	ccagggtccc	360
accaggttgt	cacgcctcat	catctcgctc	cgtgcctgga	gcgccaggca	cttacaccag	420
gaggaccaga	ggcccgaactc	tttcttgag	cgtttccgcg	gagctgagct	ccaagagggtg	480
tctagccgag	aaagccatgt	ccagttcaat	gtgggcagcc	aggagccacc	agacagaggg	540
agaagtgcct	ggcccctggc	cagaaacaac	accaacacct	gcaacaactc	ggagaaggat	600
gacaaggcga	aaaaggagga	gaaagagaaa	aaggaagaga	aaaaggagaa	cccaagaaa	660
gaggagaaga	agaaggacag	cgttgtgatg	gaccttcca	gcaacatgta	ctaccactgg	720
ctgactgtca	tcgccgtgcc	tgtcttctac	aactgggtgc	tgctcgtgtg	cagggcctgt	780
ttcgatgagc	tccagtcoga	gcacctgatg	ccttggctgg	tcctggacta	ctcagcagac	840
atcctctatg	gcattggatat	gctggtccga	gcccgacag	gcttcttgga	gcaaggcctg	900
atggtcatgg	acgccagcga	gctgtggaag	cactacacac	agaccttgca	cttcaagctg	960
gacgtgttgt	ccctgggtgcc	cacagacctg	gcttatttta	agctgggcat	gaactaccca	1020
gaactgaggt	tcaaccgcct	cctgaagttg	gcccggtctt	tcgaattctt	tgaccgcacg	1080
gagacaagga	ccaactaccc	caacatgttc	aggatcgga	acttggctct	gtacatcctc	1140
atcatcatcc	actggaatgc	ctgcatctac	tttgccattt	ccaagttcat	tggtttcggg	1200
accgactcct	gggtctaccc	aaacgtctcc	aaccagagt	atgggcgcct	ctccagaaaag	1260
tacatttaca	gtctctactg	gtccaccttg	accctgacca	ccattgggga	gaccccgccc	1320
cccgtaaaag	acgaggaata	tctctttgtg	gtcatcgact	tcctgggtgg	cgtcctgatt	1380
tttgccacca	tcgtgggcaa	cgtgggtccc	atgatctcaa	acatgaatgc	ttcacgggcc	1440
gagttccagg	ccaagatcga	ttccatcaag	cagtacatgc	agttccgcaa	ggtgaccaag	1500
gacttgagga	cacgggtgat	ccgctggttc	gactacctgt	gggccaataa	gaagacagtg	1560
gatgagaagg	aggtgctcaa	gagcctcccc	gacaagctga	aggccgagat	cgccatcaac	1620
gtgcacctgg	acaccctgag	gaaggtccga	atcttccagg	actgcgaggc	ggggctgctg	1680
gtggagctgg	tgctgaagct	gcggccggca	gtgttcagcc	ccggggacta	catctgataag	1740
aagggggaca	tcgggaggga	gatgtacatt	atcaaggagg	gcaagctggc	cgtggtggcc	1800
gaggacggca	tcaccagtt	cgtggtcctc	ggcgacggga	gttatttcgg	ggagatcagc	1860
atcttgaaca	tcaaggggag	caagtccggg	aaccgcccga	cggccaacat	caggagcatc	1920
ggctactcgg	acctgttctg	cctctccaag	gacgacctga	tggaggcgct	caccgagtac	1980
cccagggcca	agaaggcgct	ggaggagaaa	gggcggcaga	tcctcatgaa	ggacaacctg	2040
atcgacgagg	agctggccaa	ggccggggca	gaccccaaag	acatcgagga	gaagtgagg	2100
cacctcgaga	cctccctgga	ctccctgcag	accaggtttg	cgcggctcct	gagtgagtac	2160
aacgccaccc	agatgaaggt	gaagcagcgg	ctcagccagt	tggaaagcca	ggtgaagatg	2220
ggcctcccgc	ctgatggcga	tgctccgcaa	actgaggcca	gtcagccctg	aagacacagg	2280
tgccctctcc	tgccctcccg	gggcggcggt	cagtgcgacg	ctgcgccgca	cggggctcgg	2340
ccgggaccga	attctagctt	tccccacct	ctgtgcgctg	cgtggccttg	ggagagagcc	2400
ttggtttccc	tcacttagat	aacaggactc	tttatgtctg	tcccagttaa	gtgacaggtt	2460
gctgtgagct	ccacaagaaa	cgcttcgtga	ggcagggttt	tgtaagtgt	gagatgtttc	2520
taggccaaga	gtataaaaat	gtgagcacag	aagttgtttt	tttttttttt	ttttaatccg	2580
tggggaatat	ttagactcct	gaacttcatt	ttttttgtaa	atgggaggtt	atttacttac	2640
c						2641