

ENDOGLYCOSIDASE FROM STREPTOCOCCUS PYOGENES AND METHODS USING IT

Field of the Invention

The present invention relates to a novel endoglycosidase, mutants thereof
5 lacking glycan hydrolyzing activity, and its use in methods of hydrolyzing the glycan
of glycoproteins.

Background of the Invention

Endoglycosidase S (EndoS) is secreted by a number of serotypes of
10 *Streptococcus pyogenes* and has a specific endoglycosidase activity on native IgG by
hydrolyzing the conserved glycans attached to the asparagine 297 residue on the
heavy chains of IgG, Collin and Olsen, The EMBO Journal, 2001, 20 3046-3055.
EndoS is the first known bacterial enzyme with a unique specificity for native IgG. In
contrast, the activities of other known endoglycosidases require or are enhanced by
15 denaturation of the glycoprotein substrate.

Antibodies such as IgG have many applications in basic research as well as in
diagnostics and drug development. In some of these applications, such as
immunohistochemistry, immunoassays, tumour detection, radiotherapy,
crystallographic studies of antibody binding sites and immunotargeting, it is more
20 convenient to use Fab fragments than whole IgG molecules. Some of the advantages
of using Fab fragments are that they will not be affected by Fc receptors on cells or
precipitate antigen, they display a reduced immunogenicity and are less susceptible to
phagocytosis, and that radiolabelled Fab fragments are more rapidly cleared from
tissue than whole IgG molecules. For other applications, it is desirable to use Fc
25 fragments of IgG. In further applications, it may be desirable to use deglycosylated
versions of the antibodies or other glycoproteins.

The cleavage of IgG into Fab and Fc fragments is most often carried out using
proteolytic enzymes such as pepsin or papain. These enzymes often cleave other
proteins, so the cleavage reaction generally has to be performed on a purified IgG
30 fraction. Furthermore, pepsin and papain typically cleave IgG in more than one place.
This means that the fragments obtained often do not correspond to whole Fab or Fc
fragments, and even if cleavage does result in Fab and Fc fragments, they are
typically susceptible to further cleavage into smaller fragments. The isolation of Fc
fragments from Fab fragments is most often carried out using protein A or G affinity

separation columns, which utilise the Fc-binding properties of the bacterial proteins A and G.

Many different glycoproteins have utility in therapeutic applications. Methods to analyse the glycosylation of such proteins have utility in the research and development of the proteins as therapeutics. It may also be desirable to provide deglycosylated versions of these proteins.

Summary

The inventors have identified a novel endoglycosidase from serotype M49 *Streptococcus pyogenes*, referred to herein as EndoS49. EndoS49 was isolated from strain NZ131, a nephritogenic and highly transformable strain of serotype M49. NZ131 strain is a clinical isolate from a case of acute post-streptococcal glomerulonephritis in New Zealand. At a protein level, EndoS49 has less than 40% identity to EndoS, and is a 90kDa protein, compared to the 108kDa of EndoS. EndoS49 has deglycosylation activity for a broader range of proteins than EndoS.

The enzyme is a 90kDa enzyme, having a family 18 glycoside hydrolase catalytic domain. EndoS49 hydrolyzes glycan on human glycoproteins, and in particular IgG1-4, and alpha-1-microglobulin. EndoS49 can be used in the hydrolysis of glycans on human glycoproteins including IgG and alpha-1-microglobulin. EndoS49 can thus be used in glycoprofiling analysis in which the enzyme is contacted with a glycoprotein, and the products produced are separated for analysis of the glycans and the protein. EndoS49 can also be used to prepare deglycosylated proteins. The enzyme can be modified to reduce or remove endoglycosidase activity.

The modified EndoS49 polypeptide which lacks endoglycosidase activity can be used in methods for isolating glycosylated and/or functionally active IgG. By using such a modified EndoS49 polypeptide in combination with an additional IgG-binding reagent which is capable of binding denatured and/or deglycosylated IgG, the inventors have also identified a method for assessing the glycosylation state or functional quality of an IgG-containing sample.

In accordance with a first aspect, there is thus provided a polypeptide comprising

- (a) the amino acid sequence of SEQ ID NO: 1;
- (b) a variant thereof having at least 50% identity to the amino acid sequence of SEQ ID NO: 1 and having endoglycosidase activity; or

(c) a fragment of either thereof having endoglycosidase activity. In a second aspect, there is provided a polypeptide capable of binding to IgG and which does not have endoglycosidase activity comprising

(a) the amino acid sequence of SEQ ID NO: 2;

5 (b) a variant thereof having at least 50% identity to the amino acid sequence of SEQ ID NO: 1, in which the amino acid equivalent to glutamic acid at position 186 is substituted; or

(c) a fragment of either thereof.

The disclosure also provides polynucleotides, expression vectors and host cells
10 encoding or expressing the polypeptides of the disclosure. The disclosure also relates to the use of the polypeptides of the disclosure in a method of determining or analysing the glycosylation state of the protein, and in particular of an antibody, in particular an IgG antibody.

In a third aspect, there is provided a polynucleotide comprising a sequence which
15 encodes a polypeptide as defined herein.

In a fourth aspect, there is provided an expression vector, comprising a polynucleotide as defined herein.

In a fifth aspect, there is provided a method for hydrolyzing glycan of a glycoprotein comprising incubating the glycoprotein with a polypeptide as defined herein.

20 In a sixth aspect, a method for assessing the glycosylation status of a glycoprotein comprising incubating a glycoprotein with a polypeptide as defined herein, and analysing the products produced.

The disclosure also provides a method for isolating IgG from an IgG-containing sample, which method comprises:

25 (a) contacting said IgG-containing sample with a modified EndoS49 polypeptide which lacks IgG endoglycosidase activity

(b) separating said EndoS49 from the contacted sample;
thereby obtaining isolated IgG.

In seventh aspect, there is provided a method for isolating IgG from an IgG-containing sample, which method comprises
30

(a) contacting said IgG-containing sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, to thereby allow formation of a IgG-EndoS49 polypeptide complex;

(b) separating said EndoS49 from the contacted sample; and thereby obtaining
35 isolated IgG.

In an eighth aspect, there is provided a method for determining the presence or absence of IgG in a sample, which method comprises

- (a) contacting said sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, to thereby allow formation of a IgG-EndoS49 polypeptide complex;
- 5 (b) optionally separating said EndoS49 from the contacted sample; and
- (c) determining whether the separated EndoS49 is bound to IgG; and thereby determining the presence or absence of IgG in the sample.

Additionally there is provided method of assessing the glycosylation state or functional quality of an IgG-containing sample, which method comprises taking a first and a second sub-sample of the IgG-containing sample, and wherein steps (a) and (b) according to the method above are applied to the first sub-sample, and wherein steps (a) and (b) as above are applied to the second sub-sample except the EndoS49 polypeptide is substituted with an alternative IgG-binding reagent which is capable of binding denatured and/or deglycosylated IgG, and further comprising:

- 15 (c) quantifying the amount of IgG bound to the EndoS49 polypeptide in the first sub-sample, and the amount of IgG bound to the alternative IgG-binding reagent in the second sub-sample; and
- (d) comparing both the amounts of bound IgG determined in (c); and thereby assessing the glycosylation state or functional quality of an IgG containing sample.

In a ninth aspect, there is provided a method of assessing the glycosylation state or functional quality of an IgG containing sample, which method comprises taking a first and a second sub-sample of the IgG-containing sample, and wherein steps (a) and (b) according to the seventh aspect are applied to the first sub-sample, and steps (a) and (b) according to the seventh aspect are applied to the second sub-sample except the EndoS49 polypeptide is substituted with an alternative IgG-binding reagent which is capable of binding unglycosylated and/or denatured, inactive IgG, and further comprising

- 25 (a) quantifying the amount of IgG bound to the EndoS49 polypeptide in the first sub-sample, and the amount of IgG bound to the alternative IgG-binding reagent in the second sub-sample; and
- 30 (b) comparing both of the amounts of bound IgG determined in (c); and thereby assessing the glycosylation state or functional quality of an IgG containing sample.

The modified enzyme of the present disclosure may also be used in methods for isolating Fab or Fc fragments of IgG. The methods of the disclosure make use of a

highly specific IgG cleaving enzyme from *S. pyogenes*, IdeS (Immunoglobulin G-degrading enzyme of *S. pyogenes*), and an EndoS49 polypeptide.

In an tenth aspect, there is provided a method for isolating Fab fragments from IgG-containing samples, which method comprises (a) contacting said Ig-G containing sample with IdeS, and an EndoS49 polypeptide; (b) separating said IdeS and said EndoS49 polypeptide from the contacted sample; and thereby isolating Fab fragments.

In a eleventh aspect, there is provided a method for isolating Fc fragments from IgG-containing samples, which method comprises (a) contacting said IgG-containing sample with IdeS; (b) separating IdeS from the mixture obtained in step (a), thereby isolating Fab and Fc fragments; (c) contacting said Fab and Fc fragments with an EndoS49 polypeptide to thereby allow formation of a Fc fragment-EndoS49 polypeptide complex; (d) separating the Fc fragment-EndoS49 polypeptide complexes from the mixture obtained in step (c); and (e) isolating Fc fragments from the Fc fragment-EndoS49 polypeptide complexes obtained in step (d).

In one method of the disclosure, a sample containing IgG is contacted with IdeS and an EndoS49 polypeptide, which is a modified EndoS49 polypeptide which lacks endoglycosidase activity as described above.

In the methods of the disclosure, typically IdeS cleaves the IgG into Fab and Fc fragments and the EndoS49 polypeptide binds to the Fc fragments. The Fc fragments are then separated from the Fab fragments.

This method is particularly useful for isolating Fab or Fc fragments from samples comprising purified IgG. More specifically, it is useful for isolating Fab or Fc fragments from a sample comprising IgG purified using the modified EndoS49 polypeptide of the disclosure. However, the method can also be adapted for use on samples containing unpurified IgG, such as serum, cell lysate or cell culture medium.

Also provided are kits for carrying out the methods according to the disclosure.

In a twelfth aspect, there is provided a kit for isolating IgG from an IgG-containing sample, comprising (a) an EndoS49 polypeptide as defined herein; and optionally (b) means for separating said Endo S49 polypeptide from a sample.

In a thirteenth aspect, there is provided a kit for determining the presence or absence of IgG in a sample, comprising (a) an EndoS49 polypeptide as defined herein; and optionally (b) means for separating said Endo S49 polypeptide from a sample.

In a fourteenth aspect, there is provided a kit for assessing the glycosylation

state or functional quality of an IgG containing sample, comprising (a) an EndoS49 polypeptide as defined herein; (b) an alternative IgG-binding reagent which is capable of binding denatured and/or deglycosylated IgG; (c) means for separating said EndoS49 polypeptide and said alternative igG-binding reagent from a sample.

5 In a fifteenth aspect, there is provided kit for isolating Fab or Fc fragments of IgG comprising (a) IdeS; (b) an EndoS49 polypeptide; and (c) means for separating said IdeS and said EndoS49 polypeptide from a sample.

Brief Description of the Figures

10 **Figure 1.** ClustalW alignment of EndoS49 and EndoS reveals two different proteins. EndoS49 and EndoS was aligned using ClustalW in the software MacVector. GH18 catalytic motif (D**D*D*E) is present at position 179-186 with Glu186 as the catalytic residue.

15 **Figure 2.** EndoS49 has activity on glycoproteins. A. 1 µg of EndoS49, its catalytic mutant and the truncated versions was incubated with 3 µg human IgG in PBS overnight at 37°C and analyzed on a SDS-PAGE gel and with LCAlectin blot. B. 1 µg EndoS49 and EndoS49(E186L) was incubated with 3 µg of subclasses 1-4 of human IgG and analyzed as above. C. 1 µg EndoS49 and its mutants were incubated with 3 µg Alpha-1-microglobulin and analyzed on a 10 % SDS-PAGE gel.

20 **Figure 3.** EndoS49 binds to IgG. 4, 2 and 1 µg of EndoS49 and its mutants were immobilized on a PVDF membrane and incubated with human IgG and later with protein-G coupled to HRP.

25 **Figure 4.** The genomic context of *ndoS49* and *ndoS*. A comparison of the genes surrounding *ndoS49* and *ndoS* in GAS strains NZ131 (M49) and 5005 (M1) was carried out in MacVector.

Figure 5. Phylogenetic analysis of EndoS49 and other bacterial endoglycosidases.

Figure 6. SDS Page gel of Avastin and Erbitux after digestion with EndoS or EndoS49 followed by IdeS digestion.

Brief Description of the Sequences

- 5 SEQ ID NO: 1 is an amino acid sequence of an EndoS49 polypeptide isolated from *S. pyogenes* M49 serotype NZ131.
- SEQ ID NO: 2 is an amino acid sequence of a modified EndoS49 polypeptide (E186L) derived from the sequence of SEQ ID NO: 1.
- SEQ ID NO: 3 is a nucleotide sequence encoding EndoS49 polypeptide
- 10 SEQ ID NO: 4 is an amino acid sequence of IdeS isolated from *S. pyogenes* API.

Detailed Description of the Invention

General polypeptide features

- 15 The present invention relates to a novel polypeptide EndoS49. The invention also provides various methods which utilize the bacterial proteins EndoS49 and IdeS, as well as other proteins. The terms protein, peptide and polypeptide are used interchangeably herein. It will be understood that certain polypeptides and methods of the invention require an EndoS49 polypeptide having endoglycosidase activity,
- 20 whereas other polypeptides and methods of the invention require a modified EndoS49 polypeptide lacking said activity.

- The following section relates to general features of all polypeptides of the invention, and in particular to variations, alterations, modifications or derivatisations of amino acid sequence which are included within the polypeptides of the invention.
- 25 It will be understood that such variations, alterations, modifications or derivatisations of polypeptides as are described herein are subject to the requirement that the polypeptides retain any further required activity or characteristic as may be specified subsequent sections of this disclosure.

- Variants of polypeptides of the invention may be defined by particular levels
- 30 of amino acid identity which are described in more detail in subsequent sections of this disclosure. Amino acid identity may be calculated using any suitable algorithm. For example the PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (such as identifying equivalent or corresponding sequences (typically on their default settings), for example as described in Altschul S. F. (1993) J

Mol Evol 36:290-300; Altschul, S, F *et al* (1990) J Mol Biol 215:403-10. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two polynucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001. Alternatively, the UWGCG Package provides the BESTFIT program which can be used to calculate homology (for example used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, 387-395).

It will be understood that variants of polypeptides of the invention also includes substitution variants. Substitution variants preferably involve the replacement of one or more amino acids with the same number of amino acids and making

conservative amino acid substitutions. For example, an amino acid may be substituted with an alternative amino acid having similar properties, for example, another basic amino acid, another acidic amino acid, another neutral amino acid, another charged amino acid, another hydrophilic amino acid, another hydrophobic amino acid, another polar amino acid, another aromatic amino acid or another aliphatic amino acid. Some properties of the 20 main amino acids which can be used to select suitable substituents are as follows:

Ala	aliphatic, hydrophobic, neutral	Met	hydrophobic, neutral
Cys	polar, hydrophobic, neutral	Asn	polar, hydrophilic, neutral
Asp	polar, hydrophilic, charged (-)	Pro	hydrophobic, neutral
Glu	polar, hydrophilic, charged (-)	Gln	polar, hydrophilic, neutral
Phe	aromatic, hydrophobic, neutral	Arg	polar, hydrophilic, charged (+)
Gly	aliphatic, neutral	Ser	polar, hydrophilic, neutral
His	aromatic, polar, hydrophilic, charged (+)	Thr	polar, hydrophilic, neutral
Ile	aliphatic, hydrophobic, neutral	Val	aliphatic, hydrophobic, neutral
Lys	polar, hydrophilic, charged(+)	Trp	aromatic, hydrophobic, neutral
Leu	aliphatic, hydrophobic, neutral	Tyr	aromatic, polar, hydrophobic

The polypeptides of the invention and for use in the invention may be in a substantially isolated form. It will be understood that the polypeptide may be mixed with carriers or diluents which will not interfere with the intended purpose of the polypeptide and still be regarded as substantially isolated. A polypeptide for use in the invention may also be in a substantially purified form, in which case it will generally comprise the polypeptide in a preparation in which more than 50%, e.g. more than 80%, 90%, 95% or 99%, by weight of the polypeptide in the preparation is a polypeptide of the invention.

The amino acid sequence of polypeptides of the invention and for use in the invention may be modified to include non-naturally occurring amino acids or to increase the stability of the compound. When the polypeptides are produced by synthetic means, such amino acids may be introduced during production. The

polypeptides may also be modified following either synthetic or recombinant production.

Polypeptides of the invention or for use in the invention may also be produced using D-amino acids. In such cases the amino acids will be linked in reverse sequence in the C to N orientation. This is conventional in the art for producing such polypeptides.

A number of side chain modifications are known in the art and may be made to the side chains of the polypeptides, subject to the polypeptides retaining any further required activity or characteristic as may be specified herein.

It will also be understood that the polypeptides of the invention and used in the invention may be chemically modified, e.g. post-translationally modified. For example, they may be glycosylated, phosphorylated or comprise modified amino acid residues. They may be modified by the addition of a signal sequence to promote insertion into the cell membrane.

The polypeptides of the invention may also be derivatised or modified to assist with their isolation or purification. Thus, in one embodiment of the invention, the polypeptide for use in the invention is derivatised or modified by addition of a ligand which is capable of binding directly and specifically to a separation means.

Alternatively, the polypeptide is derivatised or modified by addition of one member of a binding pair and the separation means comprises a reagent that is derivatised or modified by addition of the other member of a binding pair. Any suitable binding pair can be used. In a preferred embodiment where the polypeptide for use in the invention is derivatised or modified by addition of one member of a binding pair, the polypeptide is preferably histidine-tagged or biotin-tagged. Typically the amino acid coding sequence of the histidine or biotin tag is included at the gene level and the proteins are expressed recombinantly in *E. coli*. The histidine or biotin tag is typically present at one end of the polypeptide, either at the N-terminus or at the C-terminus.

The histidine tag typically consists of six histidine residues, although it can be longer than this, typically up to 7, 8, 9, 10 or 20 amino acids or shorter, for example 5, 4, 3, 2 or 1 amino acids. Furthermore, the histidine tag may contain one or more amino acid substitutions, preferably conservative substitutions as defined above.

EndoS49 polypeptides having endoglycosidase activity

The EndoS49 polypeptide in this instance is preferably *S. pyogenes* EndoS49, or a variant or fragment of *S. pyogenes* EndoS49 which retains endoglycosidase activity. The variant may be an EndoS49 polypeptide from another *Streptococcus equi*, *Streptococcus zooepidemicus* or, preferably, *Streptococcus pyogenes*

The EndoS49 polypeptide may comprise:

- (a) the amino acid sequence of SEQ ID NO: 1;
- (b) a variant thereof having at least 50% identity to the amino acid sequence of SEQ ID NO: 1 and having endoglycosidase activity; or
- (c) a fragment of either thereof having endoglycosidase activity.

Preferably, the polypeptide comprises, or consists of, the sequence of SEQ ID NO: 1. SEQ ID NO: 1 is the sequence of EndoS49 from *S. pyogenes*. The EndoS49 polypeptide of the invention may additionally not comprise a signal sequence.

Variant polypeptides as described in this section are those for which the amino acid sequence varies from that in SEQ ID NO: 1, but which retain the endoglycosidase activity of EndoS49. Such variants may include allelic variants and the deletion, modification or addition of single amino acids or groups of amino acids within the protein sequence, as long as the peptide maintains IgG endoglycosidase activity.

The variant sequences typically differ by at least 1, 2, 3, 5, 10, 20, 30, 50, 100 or more mutations (which may be substitutions, deletions or insertions of amino acids). For example, from 1 to 100, 2 to 50, 3 to 30 or 5 to 20 amino acid substitutions, deletions or insertions may be made, provided the modified polypeptide retains activity as an IgG-specific endoglycosidase.

Variants of the amino acid sequence of SEQ ID NO: 1 preferably contain residues 179 to 186 of SEQ ID NO: 1, and in particular include the motif D**D*D*E. These amino acids constitute a family 18 glycoside hydrolase catalytic domain. The glutamic acid at position 186 is essential for enzymatic activity. Most preferably, therefore, the variant of SEQ ID NO: 1 contains glutamic acid at the position equivalent to position 186 of SEQ ID NO: 1. The variant of SEQ ID NO: 1 may contain residues 179 to 186 of SEQ ID NO: 1 having one or more conservative substitutions, provided that the variant contains glutamic acid at the position equivalent to position 186 of SEQ ID NO: 1.

Typically, polypeptides which display the endoglycosidase activity of EndoS49 with more than about 50%, 55% or 65% identity, preferably at least 70%, at least 80%, at least 90% and particularly preferably at least 95%, at least 97% or at least 99% identity, with the amino acid sequence of SEQ ID NO: 1 are considered variants of the protein. The identity of variants of SEQ ID NO: 1 may be measured over a region of at least 100, at least 250, at least 500, at least 750, at least 800, at least 810, at least 820, at least 930, at least 940 or more contiguous amino acids of the sequence shown in SEQ ID NO: 1, or more preferably over the full length of SEQ ID NO: 1.

The fragment of the EndoS49 polypeptide used in the invention is typically at least 400, 500, 600, 700, 750, 800, or 825 amino acids in length, as long as it retains the IgG endoglycosidase activity of EndoS. Preferably, the fragment of the EndoS49 polypeptide used in the invention encompasses residues 179 to 186 of SEQ ID NO: 1.

Polypeptides for use in the present invention may be isolated from any suitable organism that expresses an EndoS49 polypeptide or a variant of an EndoS49 polypeptide. Typically, the EndoS49 polypeptide is isolated from suitable EndoS49 expressing strains of *Streptococcus*, preferably strains of *S. pyogenes*, and in particular those of serotype M49.

Isolation and purification of EndoS49 from an expressing *S. pyogenes* culture, or from cultures of other cells expressing EndoS49 is typically on the basis of endoglycosidase activity. Preferably the purification method involves an ammonium sulphate precipitation step and an ion exchange chromatography step. According to one method, the culture medium is fractionated by adding increasing amounts of ammonium sulphate. The amounts of ammonium sulphate may be 10 to 80%.

Preferably the culture medium is fractionated with 50% ammonium sulphate, and the resulting supernatant is further precipitated with 70% ammonium sulphate. Pelleted polypeptides may then be subjected to ion exchange chromatography, for example by FPLC on a Mono Q column. Eluted fractions may be assayed for endoglycosidase activity and peak activity fractions may be pooled. Fractions may be analysed by SDS PAGE. Fractions may be stored at -80°C.

Polypeptides for use in the invention may also be prepared as fragments of such isolated polypeptides. Further, the EndoS49 polypeptides may also be made synthetically or by recombinant means. For example, a recombinant EndoS49

polypeptide may be produced by transfecting mammalian cells in culture with an expression vector comprising a nucleotide sequence encoding the polypeptide operably linked to suitable control sequences, culturing the cells, extracting and purifying the EndoS49 polypeptide produced by the cells.

5 The EndoS49 polypeptides of invention described in this section display endoglycosidase activity. Preferably, the polypeptide hydrolyses IgG or IgG Fc fragments by hydrolysing glycan linked of a full-length IgG heavy chain polypeptide. Preferably the EndoS49 polypeptide of the invention also has endoglycosidase activity, and is capable of glycan hydrolysis of alpha-1-microglobulin.

10 The endoglycosidase activity may be determined by means of a suitable assay. For example, a test polypeptide may be incubated with glycoprotein such as IgG or alpha-1-microglobulin at a suitable temperature, such as 37°C. The starting materials and the reaction products may then be analysed by SDS PAGE. Typically, the molecular mass of the IgG heavy chain is reduced by approximately 3kDa to 4kDa if
15 the test polypeptide has IgG endoglycosidase activity. Another assay for determining whether a test polypeptide has IgG endoglycosidase activity is by detection of glycosylated IgG using *Lens culinaris* agglutinin lectin (LCA), optionally using horseradish peroxidase and peroxidase substrate. Typically, the carbohydrate signal is reduced if the test polypeptide has IgG endoglycosidase activity. Another assay for
20 determining whether a test polypeptide has IgG endoglycosidase activity is by incubation of a test polypeptide with purified IgG Fc fragments followed by reduction of the sample with 10 mM dithiotreitol and mass spectroscopy (MALDI-TOF) analysis. Endoglycosidase activity can also be measured for EndoS49 polypeptides by using alpha-1-microglobulin in place of IgG in the assays mentioned above.

25 The endoglycosidase activity of the polypeptides can be further characterised by inhibition studies.

 The EndoS49 polypeptide is capable of hydrolyzing glycoprotein molecules present in a sample taken from a subject. Thus, where the subject is a human, the EndoS49 polypeptide is capable of hydrolyzing the glycans in glycoproteins of a
30 subject, such as on the heavy chains of human IgG or alpha-1-microglobulin. EndoS49 is capable of hydrolyzing human IgG of all four subclasses (IgG₁₋₄). In preferred embodiments, the EndoS49 polypeptide has the ability to hydrolyze human IgG and alpha-1-microglobulin.

EndoS49 polypeptides which lack endoglycosidase activity

The EndoS49 polypeptide in this instance may also be modified *S. pyogenes* EndoS49, which has been engineered to lack endoglycosidase activity but which possesses IgG binding activity. Such modified EndoS49 is particularly useful in the methods described herein. By IgG binding activity it will be understood that the modified EndoS49 binds to IgG, or a variant or fragment thereof, in particular the Fc fragment thereof, which is normally glycosylated. By “normally glycosylated” it will be understood that the IgG molecule, or variant of fragment thereof, is a glycoprotein comprising at least the IgG polypeptide heavy chain (or variant of fragment thereof) coupled to at least one carbohydrate group as found coupled to naturally occurring IgG molecules. In particular, the at least one carbohydrate group is a glycan linked to the asparagine residue corresponding to residue 297 of a full-length IgG heavy chain polypeptide.

The EndoS49 polypeptide is preferably engineered by site-directed mutagenesis. By IgG binding activity it will be understood that the EndoS49 polypeptides described in this section bind at least one, preferably two, three or all four of the IgG subclasses, IgG₁₋₄. Preferably the at least one IgG subclass is bound with high affinity and/or high specificity.

By high affinity it is meant that the binding affinity constant (K_D) for the interaction of the modified EndoS49 with an IgG subclass is greater than 0.05 μM , preferably greater than 0.06 μM , 0.07 μM or 0.08 μM . Binding activity may be determined, and binding affinity may be assessed by any suitable means. For example, by surface plasmon resonance interaction analysis, equilibrium dialysis analysis, or any standard biochemical methods in conjunction with, for example, Scatchard analysis.

The variant may be derived from an EndoS49 polypeptide from another organism, such as another bacterium, as is described in the preceding section with the exception that the variant in this instance lacks endoglycosidase activity but possesses IgG binding activity. The modified EndoS49 polypeptide may comprise:

- (a) the amino acid sequence of SEQ ID NO: 2;
- (b) a variant thereof having at least 50% identity to the amino acid sequence of SEQ ID NO: 2 which lacks endoglycosidase activity; or
- (c) a fragment of either thereof which lacks endoglycosidase activity.

Preferably, the polypeptide comprises, or consists of, the sequence of SEQ ID NO: 2. SEQ ID NO: 2 is derived from the sequence of SEQ ID NO: 1, but has been engineered to lack endoglycosidase activity by the substitution of glutamic acid (E) for leucine (L) at position 186 of SEQ ID NO: 1. Such polypeptides typically possess IgG-binding activity as described above.

Variant polypeptides as described in this section are those for which the amino acid sequence varies from that in SEQ ID NO: 2, but which lack endoglycosidase activity and retain IgG-binding activity. Such variants may include allelic variants and the deletion, modification or addition of single amino acids or groups of amino acids within the protein sequence, as long as the peptide maintains the above characteristics.

The variant sequences typically differ by at least 1, 2, 3, 5, 10, 20, 30, 50, 100 or more mutations (which may be substitutions, deletions or insertions of amino acids). For example, from 1 to 100, 2 to 50, 3 to 30 or 5 to 20 amino acid substitutions, deletions or insertions may be made, provided the modified polypeptide lacks endoglycosidase activity and retains IgG-binding activity.

Typically, polypeptides which lack endoglycosidase activity and retain IgG-binding activity with more than about 50%, 55% or 65% identity, preferably at least 70%, at least 80%, at least 90% and particularly preferably at least 95%, at least 97% or at least 99% identity, with the amino acid sequence of SEQ ID NO: 2 are considered variants of the protein. The identity of variants of SEQ ID NO: 2 may be measured over a region of at least 100, at least 250, at least 500, at least 750, at least 800, at least 820, at least 830 or more contiguous amino acids of the sequence shown in SEQ ID NO: 2, or more preferably over the full length of SEQ ID NO: 2.

The fragment of the EndoS49 polypeptide used in the invention is typically at least 300, 400, 500, 600, 700, 750, 800 or 830 amino acids in length, as long as it lacks endoglycosidase activity and retains IgG-binding activity.

In an alternative method, an EndoS49 protein with the desired characteristics can be produced by altering a nucleotide encoding an EndoS49 protein, and then expressing said nucleotide in a suitable system. Suitable methods include site-directed mutagenesis of the nucleotide encoding the protein. This technique has been widely used in the study of protein functions. The technique is typically oligonucleotide-based and involves the following steps:

(1) Cloning the DNA encoding the protein of interest into a plasmid vector.

(2) Denaturing the plasmid DNA to produce single strands.

(3) Contacting the denatured DNA with a synthetic oligonucleotide (or
5 oligonucleotides) complementary to the target sequence but incorporating the desired mutation(s) (point mutation, deletion, or insertion), such that the synthetic oligonucleotide anneals to the target region.

(4) Extending the mutated strand by a DNA-polymerase using the plasmid DNA strand as the template.

10 (5) Propagating the heteroduplex (mutated/non-mutated strand) by transformation in *E. coli*.

After propagation, about 50% of the produced heteroduplexes are mutants and the other 50% are "wild type" (no mutation). Selection and enrichment methods are used to favor the production of mutants. For example, the parental non-mutated
15 strand can be digested with a restriction enzyme that only digests methylated DNA (DpnI). This allows removal of the parental strand from the reaction before transformation of *E. coli* by since the newly synthesized strands are un-methylated while the parental strand (if purified from the correct *E. coli* background) is methylated.

20 Alternatives to site-directed mutagenesis include:

(1) Polymerase chain reaction (PCR) based methods using specific mutagenic primers, or error-prone PCR with subsequent screening for desired mutations or loss/gain of protein function.

(2) Introduction of a plasmid harboring the gene of interest into an *E. coli*
25 mutator strain (deficient in DNA proofreading systems) and subsequent screening for desired mutations or loss/gain of protein function.

(3) Chemical synthesis of partial or whole genes containing the desired mutations and subsequent introduction into an appropriate protein expression system.

Alternatively, an EndoS49 protein with the desired characteristics can be
30 produced by DNA-independent methods, which include chemical synthesis of parts of a polypeptide with the desired mutation.

Polypeptides for use in the invention may also be prepared as fragments of such isolated polypeptides. Further, the EndoS49 polypeptides may also be made synthetically or by recombinant means. For example, a recombinant EndoS49

polypeptide may be produced by transfecting mammalian cells in culture with an expression vector comprising a nucleotide sequence encoding the polypeptide operably linked to suitable control sequences, culturing the cells, extracting and purifying the EndoS49 polypeptide produced by the cells.

5 The EndoS49 polypeptide is capable of binding to IgG molecules present in a sample taken from a subject. Thus, where the subject is a human, the EndoS49 polypeptide is capable of binding human IgG. EndoS49 is capable of binding human IgG of all four subclasses (IgG₁₋₄).

10 *Polynucleotides, vectors and host cells*

 The invention also relates to polynucleotides encoding the above polypeptides, and their use in medicine. In particular the invention relates to polynucleotides comprising or consisting of (a) the coding sequence of SEQ ID NO:3 or a complementary sequence thereto; (b) sequence which hybridises under stringent
15 conditions to the sequences defined in (a); (c) sequence which is degenerate as a result of the genetic code to sequence as defined in (a) or (b); (d) sequence having at least 60% identity to sequences defined in (a) (b) or (c); and (e) fragments of the above sequences.

 Typically the polynucleotide is DNA. However, the invention may
20 comprise RNA polynucleotides. The polynucleotides may be single or double stranded, and may include within them synthetic or modified nucleotides.

 A polynucleotide of the invention can hybridize to the coding sequence or the complement of the coding sequence of SEQ ID NO: 3 at a level significantly above background. Background hybridization may occur, for example, because of other
25 DNAs present in a DNA library. The signal level generated by the interaction between a polynucleotide of the invention and the coding sequence or complement of the coding sequence of SEQ ID NO: 3 is typically at least 10 fold, preferably at least 100 fold, as intense as interactions between other polynucleotides and the coding sequence of SEQ ID NO: 3. The intensity of interaction may be measured, for
30 example, by radiolabelling the probe, e.g. with ³²P. Selective hybridisation may typically be achieved using conditions of medium to high stringency. However, such hybridisation may be carried out under any suitable conditions known in the art (see Sambrook *et al*, 1989. For example, if high stringency is required suitable conditions

include from 0.1 to 0.2 x SSC at 60°C up to 65°C. If lower stringency is required suitable conditions include 2 x SSC at 60°C.

The coding sequence of SEQ ID NO: 3 may be modified by nucleotide substitutions, for example from 1, 2 or 3 to 10, 25, 50 or 100 substitutions. The polynucleotide of SEQ ID NO: 3 may alternatively or additionally be modified by one or more insertions and/or deletions and/or by an extension at either or both ends. Additional sequences such as signal sequences may also be included. The modified polynucleotide generally encodes a polypeptide which has endoglycosidase activity. Alternatively, a polynucleotide encodes an epitope portion of an EndoS49 polypeptide. Degenerate substitutions may be made and/or substitutions may be made which would result in a conservative amino acid substitution when the modified sequence is translated, for example as shown in the Table above.

A nucleotide sequence which is capable of selectively hybridizing to the complement of the DNA coding sequence of SEQ ID NO: 3 will generally have at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 98% or at least 99% sequence identity to the coding sequence of SEQ ID NO: 3 over a region of at least 20, preferably at least 30, for instance at least 40, at least 60, more preferably at least 100 contiguous nucleotides or most preferably over the full length of SEQ ID NO: 3. Methods for the calculation of sequence identity or similarity are discussed in more detail above in relation to the polypeptides of the invention.

Any combination of the above mentioned degrees of sequence identity and minimum sizes may be used to define polynucleotides of the invention, with the more stringent combinations (i.e. higher sequence identity over longer lengths) being preferred. Thus, for example a polynucleotide which has at least 90% sequence identity over 25, preferably over 30 nucleotides forms one aspect of the invention, as does a polynucleotide which has at least 95% sequence identity over 40 nucleotides.

Polynucleotide fragments, such as those suitable for use as probes or primers will preferably be at least 10, preferably at least 15 or at least 20, for example at least 25, at least 30 or at least 40 nucleotides in length. They will typically be up to 40, 50, 60, 70, 100 or 150 nucleotides in length. Probes and fragments can be longer than 150 nucleotides in length, for example up to 200, 300, 400, 500, 600, 700 nucleotides in length, or even up to a few nucleotides, such as five or ten nucleotides, short of the coding sequence of SEQ ID NO: 3.

Polynucleotides according to the invention may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques. The polynucleotides are typically provided in isolated and/or purified form.

5 In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

Longer polynucleotides will generally be produced using recombinant means,
10 for example using PCR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15-30 nucleotides) to a region of the *ndoS49* gene which it is desired to clone, bringing the primers into contact with DNA obtained from a bacterial cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the
15 amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector.

Although in general the techniques mentioned herein are well known in the
20 art, reference may be made in particular to Sambrook *et al*, Molecular Cloning: A Laboratory Manual, 1989.

The polynucleotides according to the invention have utility in production of the polypeptides according to the invention, which may take place *in vitro*. Polynucleotides of the invention may be used as a primer, e.g. a PCR primer, a primer
25 for an alternative amplification reaction, a probe e.g. labelled with a revealing label by conventional means using radioactive or non-radioactive labels, or the polynucleotides may be cloned into vectors.

Polynucleotides or primers of the invention may carry a revealing label. Suitable labels include radioisotopes such as ³²P or ³⁵S, enzyme labels, or other
30 protein labels such as biotin. Such labels may be added to polynucleotides or primers of the invention and may be detected using techniques known *per se*.

Polynucleotides or primers of the invention or fragments thereof, labelled or unlabelled, may be used by a person skilled in the art in nucleic acid-based tests for detecting or sequencing *ndoS49* in a sample.

Such tests for detecting generally comprise bringing a sample containing DNA or RNA into contact with a probe comprising a polynucleotide or primer of the invention under hybridizing conditions and detecting any duplex formed between the probe and nucleic acid in the sample. Such detection may be achieved using techniques such as PCR or by immobilizing the probe on a solid support, removing nucleic acid in the sample which is not hybridized to the probe, and then detecting nucleic acid which has hybridized to the probe. Alternatively, the sample nucleic acid may be immobilized on a solid support, and the amount of probe bound to such a support can be detected.

The probes of the invention may conveniently be packaged in the form of a test kit in a suitable container. In such kits the probe may be bound to a solid support where the assay formats for which the kit is designed requires such binding. The kit may also contain suitable reagents for treating the sample to be probed, hybridizing the probe to nucleic acid in the sample, control reagents, instructions, and the like.

The polynucleotides of the invention may be incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Therefore, polynucleotides of the invention may be made by introducing a polynucleotide of the invention into a replicable vector, introducing the vector into a compatible host cell and growing the host cell under conditions which bring about replication of the vector.

Preferably the vector is an expression vector comprising a nucleic acid sequence that encodes a polypeptide of the invention. Such expression vectors are routinely constructed in the art of molecular biology and may for example involve the use of plasmid DNA and appropriate initiators, promoters, enhancers and other elements, which may be necessary, and which are positioned in the correct orientation, in order to allow for protein expression. Other suitable vectors would be apparent to persons skilled in the art. By way of further example in this regard we refer to Sambrook *et al.* 1989.

Polynucleotides according to the invention may also be inserted into the vectors described above in an antisense orientation in order to provide for the production of antisense RNA. Antisense RNA or other antisense polynucleotides or interfering RNA, iRNA may also be produced by synthetic means. Such antisense polynucleotides or iRNA may be used as test compounds in the assays of the

invention or may be useful in a method of treatment of the human or animal body by therapy.

Preferably, a polynucleotide of the invention or for use in the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A regulatory sequence, such as a promoter, "operably linked" to a coding sequence is positioned in such a way that expression of the coding sequence is achieved under conditions compatible with the regulatory sequence.

The vectors may be for example, plasmid, virus or phage vectors provided with a origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a resistance gene for a fungal vector.

Promoters and other expression regulation signals may be selected to be compatible with the host cell for which expression is designed. For example, yeast promoters include *S. cerevisiae* GAL4 and ADH promoters, *S. pombe* *nmt1* and *adh* promoter. Mammalian promoters include the metallothionein promoter which can be induced in response to heavy metals such as cadmium. Viral promoters such as the SV40 large T antigen promoter or adenovirus promoters may also be used. All these promoters are readily available in the art.

Mammalian promoters, such as β -actin promoters, may be used. Tissue-specific promoters are especially preferred. Viral promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR), the rous sarcoma virus (RSV) LTR promoter, the SV40 promoter, the human cytomegalovirus (CMV) IE promoter, adenovirus, HSV promoters (such as the HSV IE promoters), or HPV promoters, particularly the HPV upstream regulatory region (URR). Viral promoters are readily available in the art.

Expression vectors may be transformed into a suitable host cell to provide for expression of a polypeptide or polypeptide fragment of the invention. The host cell, transformed or transfected with an expression vector as described above, is cultivated under conditions to allow for expression of the polypeptide or fragment, and the

expressed polypeptide or fragment is recovered. Isolation and purification may be carried out as described above. Host cells will be chosen to be compatible with the vector and will preferably be bacterial. Host cells may also be cells of a non-human animal, or a plant transformed with a polynucleotide of the invention.

5

IdeS

IdeS is an extracellular cysteine protease produced by the human pathogen *S. pyogenes* and is described in WO 03/051914. IdeS was originally isolated from a group A streptococcal strain of serotype M1, but the *ides* gene has now been identified in all tested group A streptococcal strains. IdeS has an extraordinarily high degree of substrate specificity, with its only identified substrate being IgG. IdeS catalyses a single proteolytic cleavage in the lower hinge region of human IgG. This proteolytic degradation promotes inhibition of opsonophagocytosis and interferes with the killing of group A *Streptococcus*. IdeS also cleaves some subclasses of IgG in various animals and efficiently converts IgG into Fc and Fab fragments. The *ides* gene has been cloned and expressed in *E. coli* as a GST fusion protein.

The IdeS polypeptide for use in the methods of the invention is preferably *S. pyogenes* IdeS, or a variant or fragment of *S. pyogenes* IdeS which retains cysteine protease activity. The variant may be an IdeS polypeptide from another organism, such as another bacterium. The bacterium is preferably a *Streptococcus*. The *Streptococcus* is preferably a group A *Streptococcus*, a group C *Streptococcus* or a group G *Streptococcus*. In particular, the variant may be an IdeS polypeptide from a group C *Streptococcus* such as *S. equi* or *S. zooepidemicus*. Alternatively, the variant may be from *Pseudomonas putida*.

25 The IdeS polypeptide may comprise:

- (a) the amino acid sequence of SEQ ID NO: 4;
- (b) a variant thereof having at least 50% identity to the amino acid sequence of SEQ ID NO: 4 and having IgG cysteine protease activity; or
- (c) a fragment of either thereof having IgG cysteine protease activity.

30 Preferably, the IdeS polypeptide comprises, or consists of, the sequence of SEQ ID NO: 4. SEQ ID NO: 4 is the sequence of the mature form of IdeS, without the signal sequence.

Variant IdeS polypeptides are those for which the amino acid sequence varies from that in SEQ ID NO: 4, but which display the same IgG cysteine protease activity

as IdeS. Typically, polypeptides with more than about 50%, 55% or 65% identity, preferably at least 70%, at least 80%, at least 90% and particularly preferably at least 95%, at least 97% or at least 99% identity, with the amino acid sequence of SEQ ID NO: 4 are considered variants of the protein. Such variants may include allelic variants and the deletion, modification or addition of single amino acids or groups of amino acids within the protein sequence, as long as the peptide maintains the basic functionality of IdeS. The identity of variants of SEQ ID NO: 4 may be measured over a region of at least 50, at least 75, at least 100, at least 150, at least 200, at least 250, at least 275, at least 300 or more contiguous amino acids of the sequence shown in SEQ ID NO: 4, or more preferably over the full length of SEQ ID NO: 4.

Variants of the amino acid sequence of SEQ ID NO: 4 preferably contain residues Lys-55 and/or Cys-65 and/or His-233 and/or Asp-255 and/or Asp-257 of SEQ ID NO: 4. Most preferably, the variant of SEQ ID NO: 4 contains each of residues Lys-55, Cys-65, His-233, Asp-255 and Asp-257 of SEQ ID NO: 4.

The variant sequences typically differ by at least 1, 2, 5, 10, 20, 30, 50 or more mutations (which may be substitutions, deletions or insertions of amino acids). For example, from 1 to 50, 2 to 30, 3 to 20 or 5 to 10 amino acid substitutions, deletions or insertions may be made. The modified polypeptide retains activity as an IgG-specific cysteine protease. Preferably the variant polypeptides comprise a cysteine residue and a histidine residue at a spacing typically found in cysteine proteases. For example, in SEQ ID NO: 4, these residues are found at a spacing of about 130 amino acids, as is typically found in cysteine proteases.

The fragment of the IdeS polypeptide used in the invention is typically at least 10, for example at least 15, 20, 25, 30, 40, 50 or more amino acids in length, up to 100, 150, 200, 250 or 300 amino acids in length, as long as it retains the IgG cysteine protease activity of IdeS. Preferably, the fragment of the IdeS polypeptide used in the invention encompasses residues Lys-55 and/or Cys-65 and/or His-233 and/or Asp-255 and/or Asp-257 of SEQ ID NO: 4. Most preferably, the fragment encompasses each of residues Lys-55, Cys-65, His-233, Asp-255 and Asp-257 of SEQ ID NO: 4.

IdeS polypeptides for use in accordance with the invention display immunoglobulin cysteine protease activity, and in particular IgG cysteine protease activity. Preferably, the polypeptide cleaves IgG in the hinge region and more particularly in the hinge region of the heavy chain. Cleavage results in production of Fc and Fab fragments of IgG. Preferably the activity is specific for IgG. The cysteine

protease activity may be determined by means of a suitable assay. For example, a test polypeptide may be incubated with IgG at a suitable temperature, such as 37°C. The starting materials and the reaction products may then be analysed by SDS PAGE to determine whether the desired IgG cleavage product is present. Typically this

5 cleavage product is a 31kDa fragment. Typically there is no further degradation of IgG after this first cleavage. The cleavage product may be subjected to N-terminal sequencing to verify that cleavage has occurred in the hinge region of IgG. Preferably the N-terminal sequence comprises the sequence GPSVFLFP.

The cysteine protease activity of the polypeptides can be further characterised

10 by inhibition studies. Preferably, the activity is inhibited by the peptide derivate Z-LVG-CHN₂ and/or by iodoacetic acid both of which are protease inhibitors. However, the activity is generally not inhibited by E64.

The cysteine protease activity of the polypeptides is generally IgG-specific in that the polypeptides may not degrade the other classes of Ig, namely IgM, IgA, IgD

15 and IgE, when incubated with these immunoglobulins under conditions that permit cleavage of IgG. The IdeS polypeptide is capable of cleaving human IgG. In preferred embodiments the polypeptide has the ability to cleave human, rabbit, mouse or goat IgG.

IdeS polypeptides for use in the present invention may be isolated from any

20 suitable organism that expresses an IdeS polypeptide. Typically, the IdeS polypeptide is isolated from suitable IdeS expressing strains of *S. pyogenes*. Suitable organisms and strains may be identified by a number of techniques. For example, *S. pyogenes* strains may initially be tested for the presence an *ides* gene. The presence of the *ides* gene can then be verified by PCR using the primers or by hybridisation of the probes

25 to genomic DNA of the *S. pyogenes* strain.

S. pyogenes strains expressing active IdeS can be identified by assaying for IgG cysteine protease activity in the culture supernatant. Preferably inhibitor E64 is added to the supernatant to inhibit any SpeB cysteine protease activity. At least five strains express active IdeS: strains AP1, AP12, AP55, KTL3 and SF370. Preferably

30 the expressing strain is selected from AP1, AP12 and AP55.

Isolation and purification of IdeS from an expressing *S. pyogenes* culture, or from cultures of other cells expressing IdeS is typically on the basis of IgG cysteine protease activity. Preferably the purification method involves an ammonium sulphate

precipitation step and an ion exchange chromatography step. According to one method, the culture medium is fractionated by adding increasing amounts of ammonium sulphate. The amounts of ammonium sulphate may be 10 to 80%. Preferably the culture medium is fractionated with 50% ammonium sulphate, and the resulting supernatant is further precipitated with 70% ammonium sulphate. Pelleted polypeptides may then be subjected to ion exchange chromatography, for example by FPLC on a Mono Q column. Eluted fractions may be assayed for IgG cysteine protease activity and peak activity fractions may be pooled. Fractions may be analysed by SDS PAGE. For example, an N-terminal sequence can be obtained from the SDS PAGE protein band. Fractions may be stored at -20°C.

Methods using the endoglycosidase activity of EndoS49

As described herein, EndoS49 has endoglycosidase activity and is able to hydrolyse the glycan of glycoproteins including IgG and alpha-1-microglobulin. The present invention thus provides methods for deglycosylation of glycoproteins, and in particular, hydrolysis of glycan from glycoproteins, and in particular, from IgG and alpha-1-microglobulin. Typically, such a method includes incubating a sample containing glycoprotein with EndoS49 with a glycoprotein under conditions which allow the endoglycosidase activity. Suitable conditions include use of EndoS49 at a concentration of at least 1 µg/ml, 2 µg/ml, 4 µg/ml, 6 µg/ml, 8 µg/ml, 10 µg/ml, 12 µg/ml, 15 µg/ml or 20 µg/ml, preferably at least 10 µg/ml. Suitable conditions also include incubation of the sample with EndoS49 for at least 20 minutes, 30 minutes, 40 minutes, 50 minutes, 60 minutes, 70 minutes, 80 minutes, 90 minutes or 120 minutes, preferably at least 60 minutes. Incubation preferably takes place at room temperature, more preferably at approximately 20°C, 25°C, 30°C, 35°C, 40°C or 45°C, and most preferably at approximately 37°C.

These methods may be used to provide deglycosylated glycoproteins, which may themselves be useful in research or therapy. These methods may also be used to characterise glycans on glycoproteins, for example, in glycomapping or glycoprofiling. Such glycomapping and glycoprofiling is particularly useful for antibody molecules, such as IgG molecules, for example, in the analysis of monoclonal IgG molecules. Typically, the methods involved incubating the protein with EndoS49 to hydrolyse the glycans of the protein. Subsequently, the glycans and

the protein or polypeptide are separated, for example, using any suitable technique such as HPLC or gel chromatography. The separated moieties can then be analysed using any suitable analytical method, such as mass spectrometry, HPLC, gel chromatography, gel electrophoresis, spectrometry, capillary electrophoresis and other standard laboratory techniques for the analysis of glycans and/or proteins.

In accordance with additional methods of the present invention, the methods may also comprise utilising additional enzymes such as IdeS so that the glycans on the Fc portion of the antibody can be analysed in more details using the methods and techniques described herein.

One example is to analyze the fucosylation of an immunoglobulin. The degree of fucosylation on the Fc glycans on an IgG molecule is important for the therapeutic potential of an IgG drug candidate. Afucosylated IgG molecules increase the ADCC (nn) effect of the therapeutic IgG molecule. Thus, in accordance with the present invention, there is provided a method for analyzing the amount of fucose in the Fc glycans of an IgG, using EndoS49.

Typically, such a method includes incubating an glycoprotein, in this case an immunoglobulin, with EndoS49 under conditions which allow the endoglycosidase activity of EndoS49. Suitable conditions include use of EndoS49 at a concentration of at least 1 µg/ml, 2 µg/ml, 4 µg/ml, 6 µg/ml, 8 µg/ml, 10 µg/ml, 12 µg/ml, 15 µg/ml or 20 µg/ml, preferably at least 10 µg/ml. Suitable conditions also include incubation of the sample with EndoS49 for at least 20 minutes, 30 minutes, 40 minutes, 50 minutes, 60 minutes, 70 minutes, 80 minutes, 90 minutes or 120 minutes, preferably at least 60 minutes. Incubation preferably takes place at room temperature, more preferably at approximately 20°C, 25°C, 30°C, 35°C, 40°C or 45°C, and most preferably at approximately 37°C. IdeS may be added after the reaction with EndoS49, or in the same reaction mixture, to induce proteolysis, dividing the immunoglobulin molecule into F(ab')₂ and Fc. The two fragments are separated using a separation method before injection into a mass spectrometer. After glycan cleavage, a GlcNAc and core Fuc residue remain attached to Asn at the consensus Fc/2 glycosylation site. Since an afucosylated immunoglobulin is not core fucosylated, some Fc/2 will contain only a GlcNAc after digestion. The characteristic mass difference (-146 Da) resulting from the absence of fucose is readily apparent in the deconvoluted mass spectrum. Use of EndoS49 therefore, facilitates the direct estimation of the degree of core afucosylation of IgG.

Method for determining the presence or absence of IgG in a sample, or for isolating IgG from an IgG-containing sample

The isolation and/or detection of IgG is typically carried out in the art using such agents as Protein G or Protein A. These bacterial proteins interact well with IgG. However, Protein A does not bind to all four IgG subclasses (IgG₁₋₄), and both Protein A and Protein G are unable to discriminate between unglycosylated and/or denatured, inactive IgG and glycosylated and/or native, functionally active IgG. By contrast, the present inventors have identified that EndoS49 polypeptides which lack IgG endoglycosidase activity typically bind all four IgG subclasses with high affinity, and are selective for normally glycosylated IgG, i.e. IgG in its native, functionally active form.

Accordingly, the present invention provides an improved method for determining the presence or absence of IgG in a sample, which method comprises contacting said sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, separating said EndoS49 from the contacted sample, and thereby determining the presence or absence of IgG and, optionally, where IgG is present, obtaining isolated IgG. The invention therefore also provides a method for isolating IgG from an IgG-containing sample, which method comprises contacting said sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, separating said EndoS49 from the contacted sample, and thereby obtaining isolated IgG.

The above samples are contacted with EndoS49 polypeptide under conditions suitable for interaction between the polypeptide and the sample to take place and IgG binding activity to occur, i.e. to allow formation of a IgG-EndoS49 polypeptide complex. Suitable conditions include use of EndoS49 at a concentration of at least 1 µg/ml, 2 µg/ml, 4 µg/ml, 6 µg/ml, 8 µg/ml, 10 µg/ml, 12 µg/ml, 15 µg/ml or 20 µg/ml, preferably at least 10 µg/ml. Suitable conditions also include incubation of the sample with EndoS49 for at least 20 minutes, 30 minutes, 40 minutes, 50 minutes, 60 minutes, 70 minutes, 80 minutes, 90 minutes or 120 minutes, preferably at least 60 minutes. Incubation preferably takes place at room temperature, more preferably at approximately 20°C, 25°C, 30°C, 35°C, 40°C or 45°C, and most preferably at approximately 37°C.

A particular advantage of EndoS49 in these methods is that EndoS49 specifically binds to normally glycosylated IgG. The IgG binding activities of other

IgG binding agents typically require or are enhanced by denaturation of the IgG glycoprotein. This is typically achieved by treating an IgG-containing sample with acid. Such treatment may damage or denature some antibodies (acid-sensitive antibodies). Since the method of the invention requires no such treatment, the method
5 is particularly suitable for isolating acid-sensitive IgG in its native form from a sample.

The EndoS49 may be separated from the contacted sample by any suitable method. A preferred method for removal of the EndoS49 from a sample comprises using an EndoS49 which is derivatised or modified as described above.

10 A preferred modification comprises the addition of a histidine tag. The presence of a histidine tag means that the polypeptide binds with a high affinity to a reagent or separating means containing chelating groups on its surface which carry a nickel, copper or zinc ion. The histidine tag binds strongly to these metal ions. Such a reagent can therefore be used to separate EndoS49 from a sample.

15 Another preferred modification comprises the addition of a biotin tag. The presence of a biotin tag means that the polypeptide binds with a high affinity to a reagent or separating means comprising streptavidin. The biotin tag binds strongly to streptavidin. Such a reagent can therefore be used to separate EndoS49 from a sample.

20 Preferred reagents or separating means are populations of magnetic particles capable of binding to the EndoS49 polypeptide. For example, where the EndoS49 polypeptide is derivatised with a histidine tag, the magnetic particles contain on their surface chelating groups which carry a nickel, copper or zinc ion. Alternatively, where the EndoS49 polypeptide is derivatised with a biotin tag, the magnetic particles
25 contain on their surface streptavidin.

Accordingly, a preferred method of removing EndoS49 from a sample comprises using a population of magnetic particles as described above and carrying out magnetic field separation on the sample. The magnetic particles are preferably magnetic nanoparticles, and the magnetic field separation is preferably high-gradient
30 magnetic field separation.

It will be understood that any suitable separation means may be used. For example, the alternative means described in the preceding section.

The EndoS49 of the contacted sample may be assessed for the presence or absence of bound IgG by any suitable means.

For example, the molecular weight of the EndoS49 may be analysed. EndoS49 bound to IgG will have a higher molecular weight than EndoS49 not bound to IgG. Accordingly suitable methods include any method able to discriminate protein species by weight, for example SDS-PAGE and Western Blot, Mass spectrometry etc. Alternatively, the above Western Blot may be directly analysed for the presence of IgG by using IgG-specific antibodies or antibodies specific to a particular IgG sub-class. Detection of proteins in a blot in this manner is a widely used technique in the art.

Other suitable means for detecting the presence or absence of IgG bound to EndoS49 include incubating the EndoS49 with antibodies to IgG or IgG-binding proteins with coupled enzymes (e.g. horseradish peroxidase, alkaline phosphatase) followed by addition of fluorogenic/chromogenic substrates. In this instance, the development of a colour signal indicates the presence of IgG, with the quantity of IgG being proportional to the strength of the signal. Detection of proteins in this manner is a widely used technique in the art.

Further suitable means for detecting the presence or absence of IgG bound to EndoS49 comprise first separating bound IgG from EndoS49 so that it can be analysed/detected independently of EndoS49 by any of the above methods or any suitable method. IgG may be separated from EndoS49 by any suitable means. Suitable means include the elution of IgG from EndoS49 by contacting the EndoS49 from the contacted sample with a suitable elution buffer. The choice of elution buffer will typically depend on whether or not the IgG bound to EndoS49 is known or suspected to be acid-sensitive, i.e. denatured/inactivated by contact with acids.

Where the antibody is not acid-sensitive, an elution protocol using a low pH elution buffer may typically be employed. Elution protocols of this type are well known in the art. Such elution buffers have a pH typically below about pH 3, most preferably below about pH 2. Preferred examples include 0.1 M Glycine at pH2. In addition, or optionally, such elution buffers may typically comprise at least one of the following:

- Sodium or potassium salts, preferably at a concentration of about 0.5M to about 1M;
 - Mono-, di-, or polysaccharides with structures similar to the glycan associated with Asn-297 on native IgG;
- or any combination thereof.

However, as outlined above, the methods of the invention are particularly suitable for detection/isolation of acid-sensitive antibodies. Where the IgG bound to EndoS49 is known or suspected to be acid-sensitive, it is therefore preferable to use elution buffers and protocols which do not require a low pH. Such protocols are also known in the art and are based on the principle of providing a buffer comprising a molecule which competes with the bound IgG for binding to EndoS49, thus leading to release of the bound IgG. Suitable competition elution buffers therefore typically comprise one or more mono-, di-, or polysaccharides with structures similar to the glycan associated with Asn-297 on native IgG. Particularly preferred elution buffers comprise sucrose at about 0.25M to about 0.5M, preferably with pH from about 5.3 to about 8.3. Examples of specific preferred elution buffers include, for example: Sucrose 0.25M, in PBS pH7.4; Sucrose 0.5M, in PBS pH7.4; Sucrose 0.25M, in PBS pH5.3; Sucrose 0.25M, in PBS pH8.5 and Sucrose 0.25M, Maltose 0.25M, in PBS pH7.4.

In addition, or optionally, such competition elution buffers may typically comprise Sodium or potassium salts, preferably at a concentration of about 0.5M to about 1M.

The means for separating bound IgG from EndoS49 as described above may also be used to obtain isolated IgG.

Method of assessing the glycosylation state and/or functional quality of an IgG containing sample

The EndoS49 polypeptides of the invention in unmodified form have the ability to hydrolyse glycan of IgG. Also, as described above, the EndoS49 polypeptides of the invention lacking IgG endoglycosidase activity and having IgG binding activity are specific for glycosylated and/or native, functionally active IgG. Thus, the EndoS49 polypeptides can be used for analysing the glycosylation state of a glycoprotein, and in particular an IgG antibody.

In accordance with one aspect of the invention, an IgG antibody is incubated with an EndoS49 polypeptide of the invention which has endoglycosidase activity. The products obtained can be analysed by any suitable techniques, including HPLC, mass spec, gel chromatography, gel electrophoresis, spectrophotometer, capillary electrophoresis. Such methods of analysis can be conducted at any stage in the preparation of a protein, for example during screening of drug candidates, during

development of production processes of biologic drugs as well as a quality control in release assays and during production.

The EndoS49 polypeptides which have been modified to remove or reduce the endoglycosidase activity, or which retain the ability to bind to IgG in its native form
5 can also be useful in glycomapping, and in particular the analysis of glycoprotein, and in particular IgG structure.

For example, by using said EndoS49 polypeptides, optionally in combination with an alternative IgG-binding reagent, the present invention provides a method of assessing the glycosylation state or functional quality of an IgG containing sample,
10 which method comprises taking a first and a second sub-sample from the IgG-containing sample, contacting the first sub-sample with an EndoS49 polypeptide as described in the preceding section and the second sub-sample with an alternative IgG-binding reagent which is capable of binding unglycosylated and/or denatured, inactive IgG, and then quantifying the amount of IgG bound to the EndoS49 polypeptide in the
15 first sub-sample, and the amount of IgG bound to the alternative IgG-binding reagent in the second sub-sample. Finally, by comparing both of the amounts of bound IgG determined in the first and second sub-samples, the glycosylation state or functional quality of an IgG containing sample can be assessed.

The alternative IgG-binding reagent is typically Protein A or Protein G, which
20 bind to all forms (native or denatured) of IgG. In this instance, the amount of IgG bound to said reagent therefore represents the total IgG present in the second sub-sample. The EndoS49 polypeptide binds only to glycosylated and/or native, functionally active IgG, and therefore the amount of IgG bound to EndoS49 represents only the glycosylated and/or native, functionally active IgG present in the
25 first sub-sample. By comparing the concentration of total IgG from the second sub-sample to the concentration of native IgG in the first sample, the skilled person will recognise that one obtains a ratio which reflects the proportion of IgG in the original sample which is present in its glycosylated and/or native, functionally active form.

In another embodiment, the alternative IgG-binding reagent could be specific
30 for unglycosylated and/or denatured IgG. Such a reagent could be, for example, an antibody. Accordingly, in this embodiment the proportion of IgG in the original sample which is present in its glycosylated and/or native, functionally active form can be assessed by the formula:

Amount of IgG in first sub-sample / (Amount of IgG in first sub-sample + Amount of IgG in second sub-sample)

The samples in the above methods are contacted with EndoS49 polypeptide or alternative IgG-binding reagent under conditions suitable for interaction between the polypeptide or reagent and the sample to take place and IgG binding activity to occur. Suitable conditions are, for example, equivalent to those set out in the preceding section.

Method for isolating IgG Fab or Fc fragments from an IgG-containing sample

The methods of the present invention can be used for isolating Fab fragments from IgG-containing samples. In one embodiment, the present invention provides a method for isolating Fab fragments of IgG from an IgG-containing sample, which method comprises:

- (a) contacting said IgG-containing sample with IdeS, and an EndoS49 polypeptide;
- (b) separating said IdeS and said EndoS49 polypeptide from the contacted sample; and

thereby isolating Fab fragments.

Preferred methods for separating IdeS and EndoS49 polypeptide from a sample comprises using an IdeS and/or EndoS49 polypeptide which is derivatised or modified as described above. The same or a different modification may be used on each of IdeS and the EndoS49 polypeptide.

A preferred modification comprises the addition of a histidine tag. The presence of a histidine tag means that the polypeptide binds with a high affinity to a reagent or separating means containing chelating groups on its surface which carry a nickel, copper or zinc ion. The histidine tag binds strongly to these metal ions. Such a reagent can therefore be used to separate IdeS and/or EndoS49 polypeptide from a sample.

Another preferred modification comprises the addition of a biotin tag. The presence of a biotin tag means that the polypeptide binds with a high affinity to a reagent or separating means comprising streptavidin. The biotin tag binds strongly to streptavidin. Such a reagent can therefore be used to separate IdeS and/or EndoS49 polypeptide from a sample.

Preferred reagents or separating means are populations of magnetic particles capable of binding to the EndoS49 polypeptide. For example, where the IdeS and/or EndoS49 polypeptide polypeptide is derivatised with a histidine tag, the magnetic particles contain on their surface chelating groups which carry a nickel, copper or zinc ion. Alternatively, where the IdeS and/or EndoS49 polypeptide polypeptide is derivatised with a biotin tag, the magnetic particles contain on their surface streptavidin.

Accordingly, a preferred method of removing EndoS49 from a sample comprises using a population of magnetic particles as described above and carrying out magnetic field separation on the sample. The magnetic particles are preferably magnetic nanoparticles, and the magnetic field separation is preferably high-gradient magnetic field separation.

Thus, step (a) of the above method preferably additionally comprises contacting the sample with a population of magnetic nanoparticles capable of binding to IdeS and the EndoS49 polypeptide, and wherein step (b) comprises carrying out magnetic field separation on the sample.

The EndoS49 polypeptide is preferably a modified EndoS49 polypeptide lacking endoglycosidase activity.

In the above embodiment of the invention, the IgG-containing sample typically comprises purified or isolated IgG. By "purified or isolated IgG" is meant an IgG fraction with a purity of normal commercial grade. IgG is typically isolated from a sample such as serum or, in the case of recombinant IgG, from cell lysate. Isolation may be carried out according to any suitable method, preferably according to the method described above for the isolation of IgG using a modified EndoS49 polypeptide lacking endoglycosidase activity. Thus, one embodiment of the invention encompasses the method set out above comprising, prior to step (a):

- (i) contacting said IgG-containing sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, to thereby allow formation of a IgG-EndoS49 polypeptide complex;
- (ii) separating said IgG-EndoS49 polypeptide complex from the contacted sample;
- (iii) eluting IgG from the IgG-EndoS49 polypeptide complex thereby obtaining an IgG-containing sample;

and wherein steps (a) and (b) are carried out the IgG-containing sample obtained in step (iii). Separation of the IgG-EndoS49 polypeptide complex from a sample is preferably carried out according to the methods set out in the section above relating to methods for determining the presence or absence of IgG in a sample, or for isolating
5 IgG from an IgG-containing sample.

In an alternative embodiment of the invention, the methods are adapted to isolate Fab fragments from IgG-contained samples without the need to purify the IgG before carrying out the method. These methods can be carried out on a sample containing unpurified IgG, for example, whole serum, cell lysate or cell culture
10 medium. In this embodiment of the invention, the method comprises:

- (a) contacting said IgG-containing sample with an EndoS49 polypeptide to thereby allow formation of a IgG-EndoS49 polypeptide complex;
- (b) separating said IgG-EndoS49 polypeptide complex from the contacted sample;
- 15 (c) adding to IgG-EndoS49 polypeptide complexes obtained in step (b) IdeS; and (d) separating said IdeS and said EndoS49 polypeptide from the mixture obtained in (c);
and thereby isolating Fab fragments.

The methods for separating IdeS and/or EndoS49 polypeptide from the
20 samples/mixtures above preferably comprise using an IdeS and/or EndoS49 polypeptide which is derivatised or modified as described above. The same or a different modification may be used on each of IdeS and the EndoS49 polypeptide. Preferred reagents or separating means are populations of magnetic particles capable of binding to the IdeS and/or EndoS49 polypeptide. For example, where the IdeS
25 and/or EndoS49 polypeptide polypeptide is derivatised with a histidine tag, the magnetic particles contain on their surface chelating groups which carry a nickel, copper or zinc ion. Alternatively, where the IdeS and/or EndoS49 polypeptide polypeptide is derivatised with a biotin tag, the magnetic particles contain on their surface streptavidin.

Thus, step (a) of the above method preferably additionally comprises
30 contacting the sample with a population of magnetic nanoparticles capable of binding to the EndoS49 polypeptide, step (c) additionally comprises contacting the IgG-EndoS49 polypeptide complexes obtained in step (b) with a population of magnetic nanoparticles capable of binding to IdeS and the EndoS49 polypeptide, and wherein

steps (b) and (d) comprise carrying out magnetic field separation on the sample of (a) and mixture obtained in (c), respectively.

It will be understood that any suitable separation means may be used. For example, the alternative means described in the section relating to methods for
5 isolating a population of cells which are substantially free of IgG molecules bound to FcγRs could be adapted for separation of IdeS and/or EndoS49 polypeptide.

The EndoS49 polypeptide in the above embodiment is preferably a modified EndoS49 polypeptide lacking endoglycosidase activity.

The above methods of the invention can also be used for isolating Fc
10 fragments from IgG-containing samples. In one such embodiment of the invention, the method comprises:

- (a) contacting said IgG-containing sample with IdeS;
- (b) separating IdeS from the mixture obtained in step (a), thereby isolating Fab and Fc fragments;
- 15 (c) contacting said Fab and Fc fragments with an EndoS49 polypeptide to thereby allow formation of a Fc fragment-EndoS49 polypeptide complex;
- (d) separating the Fc fragment-EndoS49 polypeptide complexes from the mixture obtained in step (c); and
- (e) isolating Fc fragments from the Fc fragment-EndoS49 polypeptide
20 complexes obtained in step (d).

It will be understood that any suitable separation means may be used as described above, however, the methods for separating IdeS and/or EndoS49 polypeptide from the samples/mixtures above preferably comprise using an IdeS and/or EndoS49 polypeptide which is derivatised or modified as described above.

25 Preferably, step (a) of the above method additionally comprises contacting the sample with a population of magnetic nanoparticles capable of binding to IdeS, step (c) additionally comprises contacting the Fab and Fc fragments obtained in step (b) with a population of magnetic nanoparticles capable of binding to the EndoS49 polypeptide, and wherein steps (b) and (d) comprise carrying out magnetic field
30 separation on the sample of (a) and mixture obtained in (c), respectively. The EndoS49 polypeptide is preferably a modified EndoS49 polypeptide lacking endoglycosidase activity.

In an alternative embodiment, Fc fragments may be isolated from an IgG-containing sample by a method comprising:

(a) contacting said IgG-containing sample with IdeS and an EndoS49 polypeptide

(b) separating the EndoS49 polypeptide from the mixture obtained in (a); thereby isolating Fc fragments.

5 It will be understood that any suitable separation means may be used as described above. However, preferably step (a) of the above method additionally comprises contacting the sample with a population of magnetic nanoparticles capable of binding to the EndoS49 polypeptide but not to IdeS, and wherein step (b) comprises carrying out magnetic field separation on the mixture obtained in (a).

10 Preferably, the IdeS and/or the EndoS49 polypeptide are derivatised or modified as described above, with the proviso that a different modification is applied to each. For example, where the IdeS is modified by addition of a histidine tag such that it binds to a population of magnetic particles containing on their surface chelating groups which carry a nickel, copper or zinc ion, the EndoS49 polypeptide might be modified by
15 addition of a biotin tag such that it binds to a population of magnetic particles containing on their surface streptavidin. The EndoS49 polypeptide is preferably a modified EndoS49 polypeptide lacking endoglycosidase activity.

Similar to the methods for isolating Fab fragments, it will be appreciated that in the methods for separating Fc fragments the IgG-containing sample typically
20 comprises purified or isolated IgG. A preferred method of isolating IgG is described in steps (i) to (iii) as set out above.

Kits

The present invention provides

- 25 - a kit for isolating IgG from an IgG-containing sample, comprising:
- (a) an EndoS49 polypeptide according to the invention which lacks endoglycosidase activity; and optionally
 - (b) means for separating said EndoS49 polypeptide from a sample.
- 30 - a kit for determining the presence or absence of IgG in a sample, comprising:
- (a) an EndoS49 polypeptide according to the invention which lacks endoglycosidase activity; and optionally
 - (b) means for separating said EndoS49 polypeptide from a sample.

- a kit for assessing the glycosylation state and/or functional quality of an IgG containing sample, comprising:

- (a) an EndoS49 polypeptide according to the invention which lacks endoglycosidase activity; and optionally;
- 5 (b) an alternative IgG-binding reagent which is capable of binding denatured and/or deglycosylated IgG;
- (c) means for separating said EndoS49 polypeptide from a sample; and
- (d) means for separating said alternative IgG-binding reagent from a sample.

10 The alternative IgG-binding reagent comprises Protein G and/or Protein A and/or Protein A/G.

The present invention also provides:

- a kit for isolating Fab or Fc fragments of IgG comprising:

- (a) IdeS;
- 15 (b) an EndoS49 polypeptide; and
- (c) means for separating said IdeS and said EndoS49 polypeptide from a sample.

In a preferred embodiment, the kit additionally comprises an EndoS49 polypeptide according to the invention which lacks endoglycosidase activity and a
20 means for separating said EndoS49 polypeptide from a sample. The EndoS49 polypeptide is preferably an EndoS49 polypeptide according to the invention which lacks endoglycosidase activity.

Preferred embodiments of the above kits further comprise instructions for using the kit in a method of the invention. Further preferred embodiments include
25 those wherein the means for separating an EndoS49 polypeptide, an alternative IgG-binding reagent, an IdeS polypeptide, or an EndoS49 polypeptide from a sample are populations of magnetic nanoparticles, wherein each population is capable of binding to at least one of the indicated polypeptides/reagents/proteins. In this embodiment the kit typically additionally comprises instructions to perform magnetic field separation
30 on the sample.

In preferred embodiments of the above methods and kits, the polypeptides/proteins/reagents used are derivatised with an affinity tag, preferably a histidine tag, to assist with separation of said polypeptides.

The following Examples illustrate the invention:

Example 1

5 MATERIALS AND METHODS

Bacterial strains and growth

The genome of GAS strain NZ131 of serotype M49 has been sequenced and this strain was therefore selected as reference strain in this work (McShan *et al.*, 2008) (Chaussee *et al.*, 1999). GAS strain NZ131 was propagated on blood agar
10 and *Escherichia coli* strains Top10 (Invitrogen) and BL21 pLysS (Invitrogen) were propagated on lysogeny broth (LB) agar. For selection in *E. coli* Top10 cells, carbenicillin was used at 100 µg/mL and for *E. coli* BL21 pLysS, 100 µg/mL carbenicillin and 34 µg/mL chloramphenicol were used. Overnight cultures of *E. coli*
15 strain NZ131 was performed using Puregene DNA Purification Kit (Qiagen). Transformation was carried out using heat-shock at 42°C for 30 s. Plasmid preparations from *E. coli* were performed using Plasmid Miniprep Kit I (E.Z.N.A.). All primers used in this work are listed in Table 2.

20 **Recombinant expression of EndoS49**

Recombinant expression of EndoS49 in *E. coli* was established by PCR amplification of the *ndoS49* gene from group A *Streptococcus* strain NZ131, serotype M49 with the primers *ndoS49*-F-BamHI, CTGTAAGGATCCAGGAGAAGACTG, and *ndoS49*-R-XhoI, GAAACCTCGAGTCTTTGTAATCGTAGGACTT. The
25 *ndoS49* gene fragment was digested with restriction enzymes *Bam*HI and *Xho*I and ligated into the expression vector pGEX-5X-3 (Amersham Biosciences) using DNA ligase T4 (Fermentas) creating the plasmid pGEX-*ndoS49*. The expression vector was transformed into *E. coli* Top10 chemically competent cells and recombinant cells were grown on 100 µg/mL carbenicillin plates and screened with PCR using primers
30 *ndoS49*-F-BamHI and *ndoS49*-R-XhoI. Positive clones were isolated and the pGEX-*ndoS49* plasmid was purified and transformed into the *E. coli* expression strain BL21 pLysS as described above.

One recombinant clone was grown overnight at 37°C with antibiotics and diluted 1:20 in LB medium with antibiotics and grown for 3 h. The expression of the
35 protein EndoS49 was induced with 0.1 mM IPTG for 3 h. The cells were harvested

and lysed with BugBuster Protein Extraction Reagent (Novagen). Recombinant GST-EndoS49 was purified on column with Glutathione Sepharose 4B (GE Healthcare) and eluted with reduced glutathione.

5 Mutagenesis of EndoS49

Site-directed mutagenesis of the glutamic acid 186(Glu-186) to leucine (E186L) was carried out using QuickChange II Site-Directed Mutagenesis Kit (Stratagene) according to the manufacturer's instructions. The mutagenesis primer used was CTAGATATTGATATTCTTCACGAATTTACGAAC in combination with
10 the antisense of the sequence above and the plasmid pGEX-*ndoS49*(mutation underlined). This generated the plasmids pGEX-*ndoS49*(E186L) and, after sequencing, recombinant EndoS49(E186L) was expressed and purified as described for EndoS49. The truncated versions of EndoS49 were constructed by amplifying parts of the *ndoS49* gene from GAS NZ131 with primers *ndoS49*(trunc1-5) containing
15 restrictions sites BamHI and XhoI (Table 2). The fragments were digested and ligated into the pGEX vector as above, and transformed into *E. coli* Top10 and subsequently to BL21 pLysS and grown with antibiotics. The proteins were produced as above and the proteins EndoS49(trunc1) 80 kDa, EndoS49(trunc2) 70 kDa, EndoS49(trunc3) 60 kDa, EndoS49(trunc4) 50 kDa, EndoS49(trunc5) 42 kDa were purified.

20

Glycoprotein glycan hydrolysis assay

1 µg recombinant EndoS49 and its mutants were incubated with 3 µg of each glycoprotein in 20 µL PBS overnight at 37°C. Glycan hydrolysis was analyzed on a 10% SDS-PAGE gel and subsequently analyzed with LCA lectin blot as previously
25 described (Collin and Olsén, 2001a).

Slot-blot analysis

EndoS49 and its mutants were immobilized on a methanol activated PVDF membrane at 4, 2, 1 µg in PBS per slot using Millipore slot blot equipment. The
30 membrane was blocked with 5% skim milk (Difco) for 1 h at room temperature. Washing was consistently carried out for 3x10 minutes in PBST. The membrane was incubated with 10 µg human IgG (Sigma) in 0,5% skim milk for 1 h at 37°C and then washed. 5 µg horseradish peroxidase conjugated with protein G (Invitrogen) was added to the membrane and incubated for 1 h at 37°C. After washing the membrane

was developed with Supersignal West Pico Chemiluminiscent Substrate (Thermo Scientific).

Bioinformatic analysis

5 The genes *ndoS49* and *ndoS* were translated into EndoS49 and EndoS and compared using the ClustalW algorithm within the software MacVector (MacVector Inc.). The phylogenetic tree was constructed with MacVector using protein sequences from NCBI PubMed with the following accession numbers: EndoS (AF296340), EndoE (AAR20477), EndoH (NP_631673), EndoC (ADC53484), EndoF2 (P36912),
10 EndoF3 (P36913) (Collin and Olsén, 2001b) (Collin and Fischetti, 2004) (Tarentino and Plummer, 1974) (Tarentino *et al.*, 1993).

PCR screening for *ndoS49*

 Primers amplifying the *ndoS49* gene from GAS were designed and denoted
15 ndoS49-F (AAAACGCGGACCACTATATGC) and ndoS49-R (AAACGTTGTCCGAGGATTTG). 42 GAS strains were propagated on blood agar and grown overnight at 37°C with 5% CO₂. Single colonies were picked and lysed in 20 µL sterile H₂O at 99°C for 10 minutes. These lysates were used as template for a stringent PCR reaction to detect *ndoS49* in the 42 GAS strains. As a positive control,
20 primers for the amplification of the gene *recA* were designed, recA-F (AGCCCTTGATGATGCTTTG) and recA-R (AACAATTCTGGGTGATCGG). As positive controls, both PCR reactions used genomic DNA from GAS strain NZ131 (M49) and AP1 (M1) as template.

25 RESULTS

ClustalW analysis reveals two different enzymes: EndoS49 and EndoS

 The genes *ndoS49* and *ndoS* were *in silico* translated into proteins and compared using the ClustalW algorithm. On the gene level the identity is 50% and 37% on the protein level. The ClustalW analysis revealed a (nearly) identical signal
30 peptide sequence and a conserved family 18 glycoside hydrolase catalytic domain (DGLDIDIE) (Figure 1). Experimental analysis of EndoS has shown that tryptophans are essential for the glycan-hydrolyzing activity (Allhorn *et al.*, 2008). These tryptophans are also conserved in EndoS49.

Recombinant EndoS49 show glycan hydrolyzing activity on human glycoproteins

The 90 kDa EndoS49 was successfully recombinantly expressed in *E. coli* BL21 and purified from the soluble fraction using the GST-tag. EndoS49(E186L), a catalytic mutant with the glutamic acid of the GH18 motif (E186) substituted for a leucine (L), was constructed and purified in the same way. To map the activity of the protein, 5 carboxy-terminally truncated versions of the enzymes were constructed and denoted EndoS49(trunc1) 80 kDa, EndoS49(trunc2) 70 kDa, EndoS49(trunc3) 60 kDa, EndoS49(trunc4) 50 kDa and EndoS49(trunc5) 42 kDa. This collection of enzymes was utilized to analyze the glycan hydrolyzing activity of EndoS49 on human glycoproteins. First, the enzymes were incubated with human IgG overnight and analyzed on a SDS-PAGE gel and with LCA lectin blot, detecting the mannose structures in the glycan of IgG. The gel revealed a shift of 4 kDa of the IgG heavy chain treated with EndoS49 and the LCA lectin blot confirmed this shift as a lack of the *N*-linked glycan (Figure 2A). EndoS49(E186L) showed no shift and no change in glycan composition suggesting that E186 plays a crucial role in the catalytic activity of EndoS49. Concerning the truncated enzymes, EndoS49(trunc1-4) showed activity on the glycan of IgG but EndoS49(trunc5), the smallest of the enzymes (42 kDa), showed no glycan-hydrolyzing activity (Figure 2A).

Further analysis of the IgG deglycosylation by EndoS49 was carried out by incubating IgG₁₋₄ with EndoS49 and EndoS49(E186L), overnight. The IgG subclasses were analyzed as above and showed that EndoS49 has activity on all four subclasses of IgG, and in line with previous result, the catalytic mutant showed no activity (Figure 2B). Incubating the collection of enzymes with alpha-1-microglobulin, a heavy glycosylated human serum protein, and analysis on SDS-PAGE showed glycan hydrolysis activity of EndoS49 on this glycoprotein (Figure 2C). To further elucidate the specificity of EndoS49, a model substrate consisting of a *N*-acetyl-beta-D-glucosaminide coupled to the fluorescent 4-methylumbelliferyl was incubated with EndoS49 for 1, 2, 3, 4 and 16 h and the fluorescence measured. The fluorescence will increase if the sugar is cleaved but no such increase in intensity was observed (data not shown) suggesting that EndoS49 has activity only on glycoprotein substrates.

EndoS49 binding to IgG

The finding that EndoS49 has glycan hydrolyzing activity on IgG led us to believe that the enzyme binds IgG. This was evaluated with slot blot analysis where

EndoS49 and its catalytic mutant and truncated versions were immobilized on a PVDF membrane and incubated with IgG and the binding detected with protein G coupled to Horseradish-peroxidase (HRP). The slot blot show an increased binding to IgG by the catalytic mutant EndoS49(E186L) (Figure 3).

5

The gene *ndoS49* is present in GAS serotype M49

To elucidate whether *ndoS49* is present in any other serotypes than M49 a stringent PCR was deployed to analyze the presence of the *ndoS49* gene in a selection of GAS strains. The primers *ndoS49*-F and *ndoS49*-R was used in a PCR on lysates
 10 from GAS colonies together with the positive control amplifying the *recA* gene, present in all GAS strains. The *ndoS49* gene was amplified in all selected GAS M49 serotypes and also in serotype M60, whereas no other serotype gave a PCR product (Table 1).

In the sequenced genomes of GAS strains NZ131 (M49) and MGAS5005
 15 (M1) the genes surrounding *ndoS49* and *ndoS* were compared revealing that the genes are located in the same genomic context and that the surrounding genes are highly conserved (Figure 4). The full length EndoS49 was compared to a selection of previously described endoglycosidases, EndoS, EndoC, EndoH, EndoE, EndoF2, EndoF3) and a phylogenetic tree was reconstructed (Figure 5). This revealed that
 20 EndoS and EndoC are more closely related than EndoS and EndoS49 and that endoglycosidases from *Streptococcus* are close related compared to enzymes from other bacteria.

Table 1. The presence of *ndoS49* in a selection of GAS serotypes

Strain	Serotype	<i>ndoS49</i>	<i>recA</i>
5448	M1	-	+
SF370	M1	-	+
ACN1	M1	-	+
ACN2	M2	-	+
ANC3	M3	-	+
20224	M3	-	+
ACN4	M4	-	+
ACN5	M5	-	+
Manfredo	M5	-	+
ACN6	M6	-	+
AP6	M6	-	+
ACN9	M9	-	+
ACN11	M11	-	+

ACN12	M12	-	+
ACN18	M18	-	+
ACN19	M19	-	+
ACN22	M22	-	+
ACN24	M24	-	+
ACN28	M28	-	+
NZ131	M49	+	+
3487-05	M49	+	+
AW1	M49	+	+
AW2	M49	+	+
AW3	M49	+	+
AW4	M49	+	+
AW6	M49	+	+
AW7	M49	+	+
AW8	M49	+	+
AW9	M49	+	+
AW10	M49	+	+
AW11	M49	+	+
AW12	M49	+	+
AW13	M49	+	+
ACN49	M49	+	+
AP49	M49	+	+
CS101	M49	+	+
AP53	M53	-	+
ALAB49	M53	-	+
ACN55	M55	-	+
ACN57	M57	-	+
ACN60	M60	+	+
AP74	M74	-	+

Table 2. Primers used in this work

Primer name	Sequence (5'-3')
ndoS49-F-BamHI	CTGTAAGGATCCAGGAGAAGACTG
ndoS49-R-XhoI	GAAACCTCGAGTCTTTGTAATCGTAGGACTT
ndoS49(E186L)-F	CTAGATATTGATATTCTTCACGAATTTACGAAC
ndoS49(E186L)-R	GTTCGTAAATTCGTGAAGAATATCAATATCTAG
ndoS49(trunc1)-R-XhoI	ATTTCTCGAGCTGAAGACGTCCTTTAGCCACG
ndoS49(trunc2)-R-XhoI	TAAACTCGAGCCCCATCAGAAACATCTACTAAG
ndoS49(trunc3)-R-XhoI	ATTTTCTCGAGGCATTATCAACATCATAATGACC
ndoS49(trunc4)-R-XhoI	TAAACTCGAGCCAGTCATGCCTACCATAACAAGCTCAGC

ndoS49(trunc5)-R-XhoI	ATTTCTCGAGCTGTCCAACCTTGTTGAATG
ndoS49-F	AAAACGCGGACCACTATATGC
ndoS49-R	AAACGTTGTCCGAGGATTTG
recA-F	AGCCCTTGATGATGCTTTG
recA-R	AACAATTCTGGGTGATCGG

The protein sequence of EndoS49

NCBIReferenceSequences

NZ131 genome: NC_011375.1

5 *ndoS49* gene sequence: NC_011375.1

EndoS49 protein sequence: YP_002286383.1

EndoS49 Protein Sequence:

MDKHL LVKRTL GCVCAATLMGAALATHHDSLNTVKAEEKTVQTG
10 KTDQQVGAKLVQEIREGKRGPYAGYFRTWHDRASTGIDGKQHPENTMAEVPKEVDI
LFVFHDHTASDSPFWSELKDSYVHKLHQQTALVQTIGVNELNGRTGLSKDYPTPEG
NKALAAIVKAFVTDGRVDGLDIDIEHEFTNKRTPPEEDARALNVFKEIAQLIGKNGSD
KSKLLIMDTTSLVENNPIFKGIAEDLDYLLRQYYGSQGGEAEVDTINSDWNQYQNYID
15 ASQFMIGSFEEESASKGNLWFDVNEYDPNNPEKGKDIEGTRAKKYAEWQPSTGGLKA
GIFSIAIDRDGVAHV PSTYKNRTSTNLQRHEVDNISHTDYTVSRKLKTLMTEDKRYDV
IDQKDIPDPALREQIIQQVGQYKGDLERYNKTLLVLTGDKIQNLKGLEKLSKLQKLELR
QLSNVKEITPELLPESMKKDAELVMVGMTGLEKLNLSGLNRQTLGDGIDVNSITHLTSF
DISHNSLDLSEKSEDRKLLMTLMEQVSNHQKITVKN TAFENQKPKGYYPQTYDTKEGH
YDVDNAEHDILTDFVFGTVTKRNTFIGDEEAFAIYKEGAVDGRQYVSKDYTYEAFRKD
20 YKGYKVHLTASN LGETVTSKVTATTD ETYLVDSVDGEKV VHHMKNLIGSGAIMMENLA
KGAKVIGTSGDFEQAKKIFDGEKSDRFFTWGQTNWIAFDLGEINLAKEWRLFNAETNT
EIKTDSSSLNVAKGRQLILKDTTIDLEKMDIKNRKEYLSNDENWTDVAQMDDAKAIFNS
KLSNVLSRYWRFCDVGGASSYYPQYTELQILGQRLSNDVANTLKD

25 Example 2

Monoclonal Ig molecules can show minor variations in the Fc glycans and as a consequence the Fc glycans can appear as an inhomogeneous pool of Fc glycans. The vast majority of the glycans are identical but a minority can show variable carbohydrate structure or composition. The variety arises both from the origin of the Fc part, which can be human, humanized or from another species, and from the choice of production cell-line and cell culture conditions. In this example two well-known IgG based drugs, Avastin and Erbitux, were deglycosylated both with EndoS and with EndoS49. The samples were incubated with EndoS or EndoS49 as set out in the Table below.

Lane	Sample	Erbtux 100 µg	Avastin 100 µg	EndoS 0.1 mg/ml (µl)	EndoS49 0.1 mg/ml (µl)	ideS 1 µg
1	MW standard	-	-	-	-	-
2		-	+	5	-	+
3		-	+	50	-	+
4		-	+	-	5	+
5		-	+	-	50	+
6		-	+	-	-	+
7		+	-	5	-	+
8		+	-	50	-	+
9		+	-	-	5	+
10		+	-	-	50	+
11		+	-	-	-	+
12	MW standard	-	-	-	-	-

The results are presented in Figure 6. It was found that EndoS49 has the potential to cleave a larger variety of Fc glycans than EndoS. Even if incubation was left over night in order to minimize the effect of potential differences in the enzymatic activities the EndoS enzyme could not fully deglycosylate Erbtux Fc glycans. However EndoS49 shows a complete deglycosylation profile and is hence the more favourable enzyme when it comes to glycan profiling of immunoglobulins.

CLAIMS

1. An isolated polypeptide comprising
 - (a) the amino acid sequence of SEQ ID NO: 1;
 - (b) a variant thereof having at least 95% identity to the amino acid sequence of SEQ ID NO: 1 over at least 810 contiguous amino acids of SEQ ID NO: 1 and having the endoglycosidase activity of a polypeptide consisting of the amino acid sequence of SEQ ID NO: 1.
2. A polypeptide according to claim 1 which consists of the amino acid sequence of SEQ ID NO: 1.
3. An isolated polypeptide capable of binding to IgG and which does not have endoglycosidase activity comprising
 - (a) the amino acid sequence of SEQ ID NO: 2;
 - (b) a variant thereof having at least 95% identity to the amino acid sequence of SEQ ID NO: 1 over at least 810 contiguous amino acids of SEQ ID NO: 1, in which the amino acid equivalent to glutamic acid at position 186 is substituted.
4. A polynucleotide comprising a sequence which encodes a polypeptide as defined in claim 1, 2 or 3.
5. A polynucleotide according to claim 4 which comprises
 - (i) SEQ ID NO: 3 or a complementary sequence thereto;
 - (ii) a sequence which hybridises under stringent conditions to the sequence defined in (i);
 - (iii) a sequence which is degenerate as a result of the genetic code to a sequence as defined in (i) or (ii);
 - (iv) a sequence having at least 60% identity to a sequence as defined in (i), (ii) or (iii);
 - (v) a fragment of any of the sequences (i), (ii), (iii) or (iv), and which encodes a polypeptide having the endoglycosidase activity of the polypeptide of SEQ ID NO: 1; or
 - (vi) a variant of any of the sequences of (i), (ii), (iii), (iv) or (v), which comprises a substitution at the encoded amino acid for glutamic acid at position 186 or an equivalent position thereof, which encodes a polypeptide which does not have endoglycosidase activity, but is capable of binding to IgG.

6. An expression vector, comprising a polynucleotide as defined in claim 4 or 5.
7. A method for hydrolysing glycan of a glycoprotein comprising incubating the glycoprotein with a polypeptide as defined in claim 1 or 2.
8. A method for assessing the glycosylation status of a glycoprotein comprising incubating a glycoprotein with a polypeptide according to claim 1 or 2, and analysing the products produced.
9. The method according to claim 8 which comprises:
 - (a) contacting the glycoprotein with the said polypeptide to hydrolyse glycan of the glycoprotein;
 - (b) separating the glycan from the deglycosylated protein;
 - (c) analysing the glycan and/or deglycosylated protein so produced.
10. The method according to any one of claims 7 to 9 wherein the glycoprotein comprises an IgG antibody or a monoclonal IgG antibody.
11. A method for isolating IgG from an IgG-containing sample, which method comprises:
 - (a) contacting said IgG-containing sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, to thereby allow formation of a IgG-EndoS49 polypeptide complex;
 - (b) separating said EndoS49 from the contacted sample; andthereby obtaining isolated IgG.
12. A method for determining the presence or absence of IgG in a sample, which method comprises:
 - (a) contacting said sample with an EndoS49 polypeptide which lacks IgG endoglycosidase activity, to thereby allow formation of a IgG-EndoS49 polypeptide complex;
 - (b) optionally separating said EndoS49 from the contacted sample; and
 - (c) determining whether the separated EndoS49 is bound to IgG; andthereby determining the presence or absence of IgG in the sample.

13. Method according to claim 11 or 12, wherein said EndoS49 polypeptide comprises:
 - (a) the amino acid sequence of SEQ ID NO: 2;
 - (b) a variant thereof having at least 95% identity to the amino acid sequence of SEQ ID NO: 2 over at least 820 contiguous amino acids of SEQ ID NO: 2 and having IgG-binding activity.
14. Method according to any one of claims 11 to 13, wherein the isolated or detected IgG is in its native, functionally active form.
15. Method according to any one of claims 11 to 14, wherein the isolated or detected IgG comprises at least one, preferably two, three or all four of the IgG subclasses, IgG₁, IgG₂, IgG₃ and IgG₄.
16. Method according to any one of claims 11 to 15 wherein step(a) additionally comprises contacting the sample with a population of magnetic nanoparticles capable of binding to the EndoS49 polypeptide, and wherein step (b) comprises carrying out magnetic field separation on the contacted sample.
17. A method of assessing the glycosylation state or functional quality of an IgG containing sample, which method comprises taking a first and a second sub-sample of the IgG-containing sample, and wherein steps (a) and (b) according to any one of claims 11 or 13 to 15 are applied to the first sub-sample, and steps (a) and (b) according to any one of claims 11 or 13 to 15 are applied to the second sub-sample except the EndoS49 polypeptide is substituted with an alternative IgG-binding reagent which is capable of binding unglycosylated and/or denatured, inactive IgG, and further comprising:
 - (a) quantifying the amount of IgG bound to the EndoS49 polypeptide in the first sub-sample, and the amount of IgG bound to the alternative IgG-binding reagent in the second sub-sample; and
 - (b) comparing both of the amounts of bound IgG determined in (c);and thereby assessing the glycosylation state or functional quality of an IgG containing sample.

18. Method according to claim 17 wherein the alternative IgG-binding reagent comprises Protein G and/or Protein A.

19. A method for isolating Fab fragments from IgG-containing samples, which method comprises:

- (a) contacting said IgG-containing sample with IdeS, and an EndoS49 polypeptide;
- (b) separating said IdeS and said EndoS49 polypeptide from the contacted sample; and

thereby isolating Fab fragments.

20. Method according to claims 19 wherein step(a) additionally comprises contacting the sample with a population of magnetic nanoparticles capable of binding to the IdeS polypeptide and the EndoS49 polypeptide, and wherein step (b) comprises carrying out magnetic field separation on the contacted sample, and/or wherein the EndoS49 polypeptide is an EndoS49 polypeptide as defined in claim 13.

21. A method for isolating Fc fragments from IgG-containing samples, which method comprises:

- (a) contacting said IgG-containing sample with IdeS;
- (b) separating IdeS from the mixture obtained in step (a), thereby isolating Fab and Fc fragments;
- (c) contacting said Fab and Fc fragments with an EndoS49 polypeptide to thereby allow formation of a Fc fragment-EndoS49 polypeptide complex;
- (d) separating the Fc fragment-EndoS49 polypeptide complexes from the mixture obtained in step (c); and
- (e) isolating Fc fragments from the Fc fragment-EndoS49 polypeptide complexes obtained in step (d).

22. Method according to claim 21 wherein step (a) additionally comprises contacting the sample with a population of magnetic nanoparticles capable of binding to IdeS, step (c) additionally comprises contacting the isolated Fab and Fc fragments obtained in step (b) with a population of magnetic nanoparticles capable of binding to the EndoS49 polypeptide, and wherein steps (b) and (d) comprise carrying out

magnetic field separation on the contacted sample or isolated Fab and Fc fragments, respectively and/or wherein the EndoS49 polypeptide is an EndoS49 polypeptide as defined in claim 13.

23. Method according to any one of claims 19 to 22, which method comprises prior to step (a):

- (i) contacting said IgG-containing sample with an EndoS49 polypeptide as defined in claim 13, to thereby allow formation of a IgG-EndoS49 polypeptide complex;
- (ii) separating said IgG-EndoS49 polypeptide complex from the contacted sample;
- (iii) eluting IgG from the IgG-EndoS49 polypeptide complex thereby obtaining an IgG-containing sample;

and wherein steps (a) and (b) are carried out the IgG-containing sample obtained in step (iii).

24. A kit for isolating IgG from an IgG-containing sample, comprising:

- (a) an EndoS49 polypeptide as defined in claim 13; and optionally
- (b) means for separating said EndoS49 polypeptide from a sample.

25. A kit for determining the presence or absence of IgG in a sample, comprising:

- (a) an EndoS49 polypeptide as defined in claim 13; and optionally
- (b) means for separating said EndoS49 polypeptide from a sample.

26. A kit for assessing the glycosylation state or functional quality of an IgG containing sample, comprising:

- (a) an EndoS49 polypeptide as defined in claim 13;
- (b) an alternative IgG-binding reagent which is capable of binding denatured and/or deglycosylated IgG;
- (c) means for separating said EndoS49 polypeptide and said alternative IgG-binding reagent from a sample.

27. Kit according to claim 26 wherein the alternative IgG-binding reagent comprises Protein G and/or Protein A and/or Protein A/G.

28. Kit for isolating Fab or Fc fragments of IgG comprising:
- (a) IdeS;
 - (b) an EndoS49 polypeptide; and
 - (c) means for separating said IdeS and said EndoS49 polypeptide from a sample.
29. Kit according to claim 28 wherein the EndoS49 polypeptide is said EndoS49 polypeptide as defined in claim 13.
30. Kit according to any one of claims 24 to 29, wherein the means for separating said EndoS49 polypeptide, said alternative IgG-binding reagent, or said IdeS from a sample are populations of magnetic nanoparticles, wherein each population is capable of binding to at least one of the indicated polypeptides/reagents/proteins.