

New compositions for treating CMT and related disorders

The present invention relates to compositions and methods for the treatment of the Charcot-Marie-Tooth disease and related disorders.

5

Charcot-Marie-Tooth disease ("CMT") is an orphan genetic peripheral poly neuropathy. Affecting approximately 1 in 2,500 individuals, this disease is the most common inherited disorder of the peripheral nervous system. Its onset typically occurs during the first or second decade of life, although it may be detected in infancy. Course of disease is chronic with gradual neuromuscular degeneration. The disease is invalidating with cases of accompanying neurological pain and extreme muscular disability. CMT is one of the best studied genetic pathologies with approximately 30,000 cases in France. While a majority of CMT patients harbour a duplication of a chromosome 17 fragment containing a myelin gene: PMP22 (form CMT1A), two dozens of genes have been implicated in different forms of CMT. Accordingly, although monogenic in origin, this pathology manifests clinical heterogeneity due to possible modulator genes. The genes mutated in CMT patients are clustering around tightly connected molecular pathways affecting differentiation of Schwann cells or neurons or changing interplay of these cells in peripheral nerves.

20 Mining of publicly available data, describing molecular mechanisms and pathological manifestations of the CMT1A disease, allowed us to prioritize a few functional cellular modules - transcriptional regulation of PMP22 gene, PMP22 protein folding/degradation, Schwann cell proliferation and apoptosis, death of neurons, extra-cellular matrix deposition and remodelling, immune response - as potential legitimate targets for CMT-relevant therapeutic interventions. The combined impact of these deregulated functional modules on onset and progression of pathological manifestations of Charcot-Marie-Tooth justifies a potential efficacy of combinatorial CMT treatment.

25 International patent application n° PCT/EP2008/066457 describes a method of identifying drug candidates for the treatment of the Charcot-Marie-Tooth disease by building a dynamic model of the pathology and targeting functional cellular pathways which are relevant in the regulation of CMT disease.

30

International patent application n° PCT/EP2008/066468 describes compositions for the treatment of the Charcot-Marie-Tooth disease which comprise at least two compounds selected from the group of multiple drug candidates.

5

Summary of invention

The purpose of the present invention is to provide new therapeutic combinations for treating CMT and related disorders. The disclosure thus relates to compositions and methods for treating CMT and related disorders, in particular toxic neuropathy and
10 amyotrophic lateral sclerosis, using particular drug combinations.

An embodiment of this disclosure more specifically relates to a composition comprising Baclofen, Sorbitol and a compound selected from Pilocarpine, Methimazole, Mifepristone, Naltrexone, Rapamycine, Flurbiprofen and Ketoprofen, salts or prodrugs
15 thereof, for simultaneous, separate or sequential administration to a mammalian subject.

A particular embodiment of the present disclosure relates to a composition comprising Baclofen, Sorbitol and Naltrexone, for simultaneous, separate or sequential administration to a mammalian subject.
20

Another embodiment relates to a composition comprising (a) rapamycin, (b) mifepristone or naltrexone, and (c) a PMP22 modulator, for simultaneous, separate or sequential administration to a mammalian subject.

25 In a particular embodiment, the PMP22 modulator is selected from Acetazolamide, Albuterol, Amiloride, Aminoglutethimide, Amiodarone, Aztreonam, Baclofen, Balsalazide, Betaine, Bethanechol, Bicalutamide, Bromocriptine, Bumetanide, Buspirone, Carbachol, Carbamazepine, Carbimazole, Cevimeline, Ciprofloxacin, Clonidine, Curcumin, Cyclosporine A, Diazepam, Diclofenac, Dinoprostone,
30 Disulfiram, D-Sorbitol, Dutasteride, Estradiol, Exemestane, Felbamate, Fenofibrate, Finasteride, Flumazenil, Flunitrazepam, Flurbiprofen, Furosemide, Gabapentin, Galantamine, Haloperidol, Ibuprofen, Isoproterenol, Ketoconazole, Ketoprofen, L-

carnitine, Liothyronine (T3), Lithium, Losartan, Loxapine, Meloxicam, Metaproterenol, Metaraminol, Metformin, Methacholine, Methimazole, Methylergonovine, Metoprolol, Metyrapone, Miconazole, Mifepristone, Nadolol, Naloxone, Naltrexone; Norfloxacin, Pentazocine, Phenoxybenzamine, Phenylbutyrate, Pilocarpine, Pioglitazone, Prazosin, 5 Propylthiouracil, Raloxifene, Rapamycin, Rifampin, Simvastatin, Spironolactone, Tacrolimus, Tamoxifen, Trehalose, Trilostane, Valproic acid, salts or prodrugs thereof.

Another embodiment is a composition comprising Rapamycin and mifepristone, for simultaneous, separate or sequential administration to a mammalian subject.

10

A further embodiment is a composition as disclosed above further comprising one or several pharmaceutically acceptable excipients or carriers (i.e., a pharmaceutical composition).

15 Another embodiment relates to a composition as disclosed above for treating CMT or a related disorder.

A further embodiment relates to the use of a combination of compounds as disclosed above for the manufacture of a medicament for the treatment of CMT or a 20 related disorder.

A further embodiment is a method for treating CMT or a related disorder, the method comprising administering to a subject in need thereof an effective amount of a composition as defined above.

25

A further embodiment is a method of preparing a pharmaceutical composition, the method comprising mixing the above compounds in an appropriate excipient or carrier.

A more specific embodiment is a method of treating CMT1a in a subject, the method comprising administering to the subject in need thereof an effective amount of a compound or combination of compounds as disclosed above.

- 5 A further specific embodiment is a method of treating toxic neuropathy in a subject, the method comprising administering to the subject in need thereof an effective amount of a compound or combination of compounds as disclosed above.

- 10 A further specific embodiment is a method of treating ALS in a subject, the method comprising administering to the subject in need thereof an effective amount of a compound or combination of compounds as disclosed above.

- 15 Any of the various uses or methods of treatment disclosed herein can also include an optional step of diagnosing a patient as having CMT or a related disorder, particularly CMT1A, or identifying an individual as at risk of developing CMT or a related disorder, particularly CMT1A.

- 20 In this regard, a further embodiment is a method of treating CMT, particularly CMT1a, the method comprising (1) assessing whether a subject has CMT, particularly CMT1a and (2) treating the subject having CMT, particularly CMT1a with an effective amount of a combination of compounds as described above. Determining whether a subject has CMT, particularly CMT1a, can be done by various tests known per se in the art, such as DNA assays.

- 25 The disclosure may be used for treating CMT or a related disorder in any mammalian subject, particularly human subjects, more preferably CMT1a.

- 30 In one aspect of the disclosure, there is provided a composition comprising baclofen, D-sorbitol and naltrexone, salts, enantiomers, or racemates thereof, for simultaneous, separate or sequential administration to a mammalian subject.

In another aspect, there is provided the use of the composition as defined above, in the manufacture of a medicament for the treatment of Charcot Marie Tooth (“CMT”) disease, a toxic neuropathy or Amyotrophic Lateral Sclerosis (“ALS”) in a mammalian subject.

5

In another aspect, there is provided the use of Baclofen, D-sorbitol and naltrexone, salts, enantiomers, or racemates thereof in the manufacture of a medicament for the treatment of CMT disease, a toxic neuropathy or ALS in a mammalian subject, said medicament being formulated for simultaneous, separate or sequential administration.

10

Legend to the figures

Figure 1. Synergistic effect of drug combination, dose 1: effect of **A)** Mix7 (dose 1, day 10), **B)** d-Sorbitol (SRB, 500 μ M, day 10), **C)** (R/S)-Baclofen (BCL, 5 μ M, day 10) and **D)** Naltrexone (NTX, 5 μ M, day 10) on MBP expression. *:p<0.05:

15

significantly different from control (=ascorbic acid) (One-Way ANOVA followed by Fisher Post-hoc test); ns: not statistically different

Figure 2. Synergistic effect of drug combination, dose 6 **A)** Mix7 (dose 6, day 10), **B)** SRB (160 nM, day 10), **C)** BCL (1.6 nM, day 10) and **D)** NTX (1.6 nM, day 10) on MBP expression. *:p<0.05: significantly different from control (=ascorbic acid) (One-Way ANOVA followed by Fisher Post-hoc test); ns: not statistically different

Figure 3. Positive effect of Mix7 (7 doses) **A)** on day 10 and **B)** on day 11 in co-incubation with ascorbic acid in PMP22 TG co-cultures on MBP expression in percentage of control (= ascorbic acid). One-Way Anova followed by Fisher post-hoc test.

Figure 4. Positive effect on male rats of the 3 and 6 weeks treatment with Mix1 measured using bar test. Latencies were measured as the mean of two first assays of the tests (white bars represent control rats treated with placebo; black bars represent transgenic rats treated with placebo ; grey bars represent transgenic rats treated with Mix1. ** p<0,01. Statistics are realised with the Student bilateral test).

Figure 5. Positive effect on gait of male rats of the 3 and 6-week (respectively left and right graph treatment with Mix1 composition (white bars represent fluid gait ; grey bars represent not fluid gait ; black bars represent rats with a severe incapacity to walk. Statistics are realised with the Student bilateral test).

Figure 6. Positive effect on male rats of the Mix1 composition in rats using inclined plane test (25°). Rats were examined after 3, 6, 9 and 12 weeks of treatment (white bars represent control rats treated with placebo ; black bars represent transgenic rats treated with placebo ; grey bars represent transgenic rats treated with Mix1. * p<0,05. Statistics are realised with the Student bilateral test).

Figure 7. Positive effect on female rats of the 3 weeks treatment with the Mix2 composition in rats, using an inclined plane test (white bars represent control rats treated with placebo ; black bars represent transgenic rats treated with placebo ; grey bars represent transgenic rats treated with Mix2. ** p<0,01. Statistics are realised with the Student bilateral test).

Figure 8. Protective effect on male rats of Mix1 on oxaliplatin-induced neuropathy (white bars represent wild type rats treated with placebo ; black bars represent wild type

rats treated with reference product gabapentin ; grey bars represent wild type rats treated with Mix1. * $p < 0,05$; ** $p < 0,01$. Statistics are realised with the Student bilateral test).

Figure 9. Significant decrease of pmp22 RNA expression in treated transgenic animals compared to PMP22 transgenic rats, observed after 9 weeks of treatment with the Mix7-dose 3 (MPZ as reference gene, Sereda et al, 1996) ($p = 0.0015$). The transgene integration and the overexpression of pmp22 gene have also been confirmed; pmp22 RNA in transgenic PMP22 rats was 1,8 fold overexpressed compared to their wild type littermates controls ($p < 1.10^{-4}$). Extraction of pmp22 RNA was performed on sciatic nerves of 16 weeks old male rats ($n=18$ for the Wild Type, $n=20$ for the transgenic rats and $n=18$ for TG treated with Mix7-dose3). Statistical analysis was performed by using the Welch t-test.

Figure 10. A clustering analysis was performed on the inclined plane test score at 35° (to distribute in the poor, intermediate and good performance classes at all time points of evaluation (3, 6 and 9 weeks of treatment analyzed together). A significant difference was observed between WT and TG placebo: 68% of WT belonged to the good performances group and only 5% of TG placebo belonged to this group ($p=0,0003$). Mix7-dose 2 and dose 3 improved the performances of TG rats. Statistical analysis were performed by applying a trend-test at the 5% significance level ($n=18$ for WT placebo rats, $n=20$ for TG placebo rats, $n=17$ for TG treated with Mix7-dose 2, $n=18$ for TG treated with Mix7-dose3).

Figure 11. The fall latencies of TG rats in the bar test after 9 weeks of treatment with Mix7-dose3 were analyzed using a Cox model with a sandwich variance estimator, and compared to the reference TG placebo by applying a log rank-test at the 5% significance level. Mix7-dose 3 significantly increased the fall latency of TG rats after 9 weeks of treatment.

Figure 12. The grip strength of groups of wild type, transgenic placebo and transgenic animals treated with Mix7-dose 3 daily for 9 weeks was modeled using a Cox model with a sandwich variance estimator over all the times after treatment (3, 6 and 9 weeks) and compared to the reference TG placebo by applying a log rank-test at the 5% significance level. The corresponding p-values were presented on Kaplan-Meier curves A significant decrease of the fore paws grip strength of transgenic placebo rats was observed (black plain line, $n=21$) compared to WT rats (grey plain line, $p=1.45.10^{-}$

5, n=19). The treatment with Mix7-dose 3 significantly increased the strength of the fore paws (black dashed line; $p=0.03$, $n=18$).

Figure 13. A Pearson correlation test showed a significant correlation between the fall latency time in the bar test (after 9 weeks of treatment) and the pmp22 RNA expression level: $p=1,6 \cdot 10^{-4}$ (WT, TG placebo and TG treated with the Mix7-dose 3 analysed together); $p=0,07$ (TG placebo and TG treated with the Mix7-dose 3 analysed together).

The lower the pmp22 RNA expression was, the better the bar test performances were. Male rats were 16 weeks old ($n=18$ for WT rats, white circles; $n=20$ for the TG placebo, black circles and $n=18$ for TG treated with the Mix7-dose3, white triangles).

Figure 14. A Pearson correlation test showed a significant correlation between the fall latency time in the bar test (after 9 weeks of treatment) and the conduction velocity of the sensitive nerve (NCV): $p=1,34 \cdot 10^{-6}$ (WT, TG placebo and TG treated with Mix7-dose3 analysed together) and $p=0,04$ (TG placebo and TG treated with Mix7-dose3 analysed together). The higher the conduction velocity was, the better the performances in bar test were. Male rats were 16 weeks old ($n=18$ for WT rats, white circles; $n=20$ for the TG placebo, black circles and $n=18$ for TG treated with the Mix7-dose3, white triangles).

20 Detailed description of the invention

The present invention provides new therapeutic approaches for treating CMT or related disorders. The invention discloses novel drug combinations which allow an effective correction of such diseases and may be used in any mammalian subject.

25

Within the context of this invention, CMT includes CMT1A, CMT1B, CMT1C, CMT1D, CMT1X, CMT2A, CMT2B, CMT2D, CMT2E, CMT2-P0, CMT4A, CMT4B1, CMT4B2, CMT4D, CMT4F, CMT4, or AR-CMT2A, more preferably CMT1a.

30

Within the context of the present invention, the term "CMT related disorder" designates other diseases associated with abnormal expression of PMP22 leading to

abnormal myelination and loss of neurons. The term "CMT related disorder" more particularly includes Alzheimer's disease (AD), senile dementia of AD type (SDAT), Parkinson's disease, Lewis body dementia, vascular dementia, autism, mild cognitive impairment (MCI), age-associated memory impairment (AAMI) and problem associated
5 with ageing, post-encephalitic Parkinsonism, schizophrenia, depression, bipolar disease and other mood disorders, Huntington's disease, motor neurone diseases including amyotrophic lateral sclerosis (ALS), multiple sclerosis, idiopathic neuropathies, diabetic neuropathy, toxic neuropathy including neuropathy induced by drug treatments, neuropathies provoked by HIV, radiation, heavy metals and vitamin deficiency states,
10 prion-based neurodegeneration, including Creutzfeld-Jakob disease (CJD), bovine spongiform encephalopathy (BSE), GSS, FFI, Kuru and Alper's syndrome.

In a preferred embodiment, "CMT related disorder" designates a toxic neuropathy, particularly drug-induced neuropathies, or ALS.

15

As used herein, "treatment" of a disorder includes the therapy, prevention, prophylaxis, retardation or reduction of pain provoked by the disorder. The term treatment includes in particular the control of disease progression and associated symptoms.

20

Also, the term "compound" designates the chemical compounds as specifically named in the application, as well as any pharmaceutically composition with acceptable salt, hydrate, ester, ether, isomers, racemate, conjugates, pro-drugs thereof. The compounds listed in this application may also be identified with its corresponding CAS
25 number.

Thus, the preferred compounds used in the invention are Baclofen (CAS 1134-47-0) and its possible salts, enantiomers, racemates, prodrugs and derivatives; Sorbitol (CAS 50-70-4) and its possible salts, enantiomers, racemates, prodrugs and derivatives; Naltrexone (CAS 16590-41-3) and its possible salts, enantiomers, racemates, prodrugs
30 and derivatives; Mifepristone (CAS 84371-65-3) and its possible salts, enantiomers, racemates, prodrugs and derivatives; Pilocarpine (CAS 54-71-7) and its possible salts, enantiomers, racemates, prodrugs and derivatives; Methimazole (CAS 60-56-0) and its

possible salts, enantiomers, racemates, prodrugs and derivatives; Ketoprofen (CAS 22071-15-4) and its possible salts, enantiomers, racemates, prodrugs and derivatives; Flurbiprofen (5104-49-4) and its possible salts, enantiomers, racemates, prodrugs and derivatives and Rapamycin (CAS 53123-88-9) and its possible salts, enantiomers, racemates, prodrugs and derivatives.

Further compounds used in the invention are Acetazolamide (CAS 59-66-5) and its possible salts, enantiomers, prodrugs and derivatives; Albuterol (CAS 18559-94-9) and its possible salts, enantiomers, prodrugs and derivatives; Amiloride (CAS 2016-88-8) and its possible salts, enantiomers, prodrugs and derivatives; Aminoglutethimide (CAS 125-84-8) and its possible salts, enantiomers, prodrugs and derivatives; Amiodarone (CAS 1951-25-3) and its possible salts, enantiomers, prodrugs and derivatives; Aztreonam (CAS 78110-38-0) and its possible salts, enantiomers, prodrugs and derivatives; Baclofen (CAS 1134-47-0) and its possible salts, enantiomers, prodrugs and derivatives; Balsalazide (CAS 80573-04-2) and its possible salts, enantiomers, prodrugs and derivatives; Betaine (CAS 107-43-7) and its possible salts, enantiomers, prodrugs and derivatives; Bethanechol (CAS 674-38-4) and its possible salts, enantiomers, prodrugs and derivatives; Bicalutamide (CAS 90357-06-5) and its possible salts, enantiomers, prodrugs and derivatives; Bromocriptine (CAS 25614-03-3) and its possible salts, enantiomers, prodrugs and derivatives; Bumetanide (CAS 28395-03-1) and its possible salts, enantiomers, prodrugs and derivatives; Buspirone (CAS 36505-84-7) and its possible salts, enantiomers, prodrugs and derivatives; Carbachol (CAS 51-83-2) and its possible salts, enantiomers, prodrugs and derivatives; Carbamazepine (CAS 298-46-4) and its possible salts, enantiomers, prodrugs and derivatives; Carbimazole (CAS 22232-54-8) and its possible salts, enantiomers, prodrugs and derivatives; Cevimeline (CAS 107233-08-9) and its possible salts, enantiomers, prodrugs and derivatives; Ciprofloxacin (CAS 85721-33-1) and its possible salts, enantiomers, prodrugs and derivatives; Clonidine (CAS 4205-90-7) and its possible salts, enantiomers, prodrugs and derivatives; Curcumin (CAS 458-37-7) and its possible salts, enantiomers, prodrugs and derivatives; Cyclosporine A (CAS 59865-13-3) and its possible salts, enantiomers, prodrugs and derivatives; Diazepam (CAS 439-14-5) and its possible salts, enantiomers, prodrugs and derivatives; Diclofenac (CAS 15307-86-5) and its possible salts, enantiomers, prodrugs and derivatives; Dinoprostone (CAS 363-

24-6) and its possible salts, enantiomers, prodrugs and derivatives; Disulfiram (CAS 97-77-8) and its possible salts, enantiomers, prodrugs and derivatives; D-Sorbitol (CAS 50-70-4) and its possible salts, enantiomers, prodrugs and derivatives; Dutasteride (CAS 164656-23-9) and its possible salts, enantiomers, prodrugs and derivatives; Estradiol
5 (CAS 50-28-2) and its possible salts, enantiomers, prodrugs and derivatives; Exemestane (CAS 107868-30-4) and its possible salts, enantiomers, prodrugs and derivatives; Felbamate (CAS 25451-15-4) and its possible salts, enantiomers, prodrugs and derivatives; Fenofibrate (CAS 49562-28-9) and its possible salts, enantiomers, prodrugs and derivatives; Finasteride (CAS 98319-26-7) and its possible salts,
10 enantiomers, prodrugs and derivatives; Flumazenil (CAS 78755-81-4) and its possible salts, enantiomers, prodrugs and derivatives; Flunitrazepam (CAS 1622-62-4) and its possible salts, enantiomers, prodrugs and derivatives; Flurbiprofen (CAS 5104-49-4) and its possible salts, enantiomers, prodrugs and derivatives; Furosemide (CAS 54-31-9) and its possible salts, enantiomers, prodrugs and derivatives; Gabapentin (CAS
15 60142-96-3) and its possible salts, enantiomers, prodrugs and derivatives; Galantamine (CAS 357-70-0) and its possible salts, enantiomers, prodrugs and derivatives; Haloperidol (CAS 52-86-8) and its possible salts, enantiomers, prodrugs and derivatives; Ibuprofen (CAS 15687-27-1) and its possible salts, enantiomers, prodrugs and derivatives; Isoproterenol (CAS 7683-59-2) and its possible salts, enantiomers,
20 prodrugs and derivatives; Ketoconazole (CAS 65277-42-1) and its possible salts, enantiomers, prodrugs and derivatives; Ketoprofen (CAS 22071-15-4) and its possible salts, enantiomers, prodrugs and derivatives; L-carnitine (CAS 541-15-1) and its possible salts, enantiomers, prodrugs and derivatives; Liothyronine (T3) (CAS 6893-02-3) and its possible salts, enantiomers, prodrugs and derivatives; Lithium (CAS 7439-93-2) and its possible salts, enantiomers, prodrugs and derivatives; Losartan (CAS 114798-26-4) and its possible salts, enantiomers, prodrugs and derivatives; Loxapine (CAS 1977-10-2) and its possible salts, enantiomers, prodrugs and derivatives; Meloxicam (CAS 71125-38-7) and its possible salts, enantiomers, prodrugs and derivatives; Metaproterenol (CAS 586-06-1) and its possible salts, enantiomers, prodrugs and
30 derivatives; Metaraminol (CAS 54-49-9) and its possible salts, enantiomers, prodrugs and derivatives; Metformin (CAS 657-24-9) and its possible salts, enantiomers, prodrugs and derivatives; Methacholine (CAS 55-92-5) and its possible salts,

enantiomers, prodrugs and derivatives; Methimazole (CAS 60-56-0) and its possible salts, enantiomers, prodrugs and derivatives; Methylergonovine (CAS 113-42-8) and its possible salts, enantiomers, prodrugs and derivatives; Metoprolol (CAS 37350-58-6) and its possible salts, enantiomers, prodrugs and derivatives; Metyrapone (CAS 54-36-4) and its possible salts, enantiomers, prodrugs and derivatives; Miconazole (CAS 22916-47-8) and its possible salts, enantiomers, prodrugs and derivatives; Mifepristone (CAS 84371-65-3) and its possible salts, enantiomers, prodrugs and derivatives; Nadolol (CAS 42200-33-9) and its possible salts, enantiomers, prodrugs and derivatives; Naloxone (CAS 465-65-6) and its possible salts, enantiomers, prodrugs and derivatives; Naltrexone (CAS 16590-41-3) and its possible salts, enantiomers, prodrugs and derivatives; Norfloxacin (CAS 70458-96-7) and its possible salts, enantiomers, prodrugs and derivatives; Pentazocine (CAS 359-83-1) and its possible salts, enantiomers, prodrugs and derivatives; Phenoxybenzamine (CAS 59-96-1) and its possible salts, enantiomers, prodrugs and derivatives; Phenylbutyrate (CAS 1821-12-1) and its possible salts, enantiomers, prodrugs and derivatives; Pilocarpine (CAS 54-71-7) and its possible salts, enantiomers, prodrugs and derivatives; Pioglitazone (CAS 111025-46-8) and its possible salts, enantiomers, prodrugs and derivatives; Prazosin (CAS 19216-56-9) and its possible salts, enantiomers, prodrugs and derivatives; Propylthiouracil (CAS 51-52-5) and its possible salts, enantiomers, prodrugs and derivatives; Raloxifene (CAS 84449-90-1) and its possible salts, enantiomers, prodrugs and derivatives; Rapamycin (CAS 53123-88-9) and its possible salts, enantiomers, prodrugs and derivatives; Rifampin (CAS 13292-46-1) and its possible salts, enantiomers, prodrugs and derivatives; Simvastatin (CAS 79902-63-9) and its possible salts, enantiomers, prodrugs and derivatives; Spironolactone (CAS 52-01-7) and its possible salts, enantiomers, prodrugs and derivatives; Tacrolimus (CAS 104987-11-3) and its possible salts, enantiomers, prodrugs and derivatives; Tamoxifen (CAS 10540-29-1) and its possible salts, enantiomers, prodrugs and derivatives; Trehalose (CAS 99-20-7) and its possible salts, enantiomers, prodrugs and derivatives; Trilostane (CAS 13647-35-3) and its possible salts, enantiomers, prodrugs and derivatives; Valproic acid (CAS 99-66-1) and its possible salts, enantiomers, prodrugs and derivatives.

The term "combination" designates a treatment wherein several drugs are co-administered to a subject to cause a biological effect. In a combined therapy, the drugs may be administered together or separately, at the same time or sequentially. Also, the drugs may be administered through different routes and protocols.

5

The invention now discloses the identification and activities of particular drug combinations which provide an efficient treatment for CMT. More specifically, the invention discloses novel ternary combinations which provide a significant effect in vitro and in vivo on CMT or related disorders.

10

In this regard, the invention relates to a composition comprising Baclofen, Sorbitol and a compound selected from Pilocarpine, Methimazole, Mifepristone, Naltrexone, Rapamycine, Flurbiprofen and Ketoprofen, salts, enantiomers, racemates, or prodrugs thereof.

15

More preferably, the invention relates to a composition comprising Baclofen, Sorbitol and a compound selected from Pilocarpine, Methimazole, Mifepristone, Naltrexone, and Ketoprofen.

20

In the most preferred embodiment, the present invention relates to a composition comprising Naltrexone, Baclofen and Sorbitol, for simultaneous, separate or sequential administration to a mammalian subject.

25

Preferably, in the above compositions, Sorbitol is D-Sorbitol and Baclofen is RS-Baclofen or S-Baclofen, more preferably RS-baclofen.

30

Another preferred object of the invention relates to a composition comprising:

- (a) rapamycin,
- (b) mifepristone or naltrexone, and
- (c) a PMP22 modulator,

for simultaneous, separate or sequential administration to a mammalian subject.

Another preferred object of this invention is a composition comprising:

- (a) rapamycin,
- (b) mifepristone, and
- (c) a PMP22 modulator,

for simultaneous, separate or sequential administration to a mammalian subject.

5

The PMP22 modulator may be any compound that modulates PMP22 pathway in a cell and essentially causes or contributes to normalization of myelin organization and/or inhibition of neuron loss. The PMP22 modulator may be selected from Acetazolamide, Albuterol, Amiloride, Aminoglutethimide, Amiodarone, Aztreonam, Baclofen, Balsalazide, Betaine, Bethanechol, Bicalutamide, Bromocriptine, Bumetanide, Buspirone, Carbachol, Carbamazepine, Carbimazole, Cevimeline, Ciprofloxacin, Clonidine, Curcumin, Cyclosporine A, Diazepam, Diclofenac, Dinoprostone, Disulfiram, D-Sorbitol, Dutasteride, Estradiol, Exemestane, Felbamate, Fenofibrate, Finasteride, Flumazenil, Flunitrazepam, Flurbiprofen, Furosemide, Gabapentin, Galantamine, Haloperidol, Ibuprofen, Isoproterenol, Ketoconazole, Ketoprofen, L-carnitine, Liothyronine (T3), Lithium, Losartan, Loxapine, Meloxicam, Metaproterenol, Metaraminol, Metformin, Methacholine, Methimazole, Methylergonovine, Metoprolol, Metyrapone, Miconazole, Mifepristone, Nadolol, Naloxone, Naltrexone; Norfloxacin, Pentazocine, Phenoxybenzamine, Phenylbutyrate, Pilocarpine, Pioglitazone, Prazosin, Propylthiouracil, Raloxifene, Rapamycin, Rifampin, Simvastatin, Spironolactone, Tacrolimus, Tamoxifen, Trehalose, Trilostane, Valproic acid salts or prodrugs thereof.

In a preferred embodiment, compound (c) is selected from pilocarpin, methimazole and baclofen. In this regard, a most preferred composition of this invention comprises:

- (a) rapamycin,
- (b) mifepristone, and
- (c) a compound selected from pilocarpin, methimazole and baclofen,

for simultaneous, separate or sequential administration to a mammalian subject.

Specific examples of such compositions include compositions comprising:

- Rapamycin ; mifepristone and pilocarpin;
- Rapamycin ; mifepristone and Baclofen;

- Rapamycin ; mifepristone and methimazole; or
- Rapamycin ; Naltrexone and methimazole.

The experimental section shows these particular drug combinations are able to efficiently correct PMP22 expression in vitro, to restore normal myelination and neuron integrity, and thus to ameliorate CMT in animals in vivo. The results also show these combinations can protect animals from chemotherapy-induced neuropathy. As a result, these compositions may be used to prevent or reduce chemotherapy-induced neuropathy, thereby allowing patients to receive chemotherapy for longer periods.

Another object of this invention is a composition comprising Naltrexone, Baclofen and a further distinct PMP22 inhibitor as defined above.

A further object of this invention is a composition as disclosed above further comprising one or several pharmaceutically acceptable excipients or carriers (i.e., a pharmaceutical composition).

Another object of the present invention relates to a composition as disclosed above for treating CMT or a related disorder.

A further object of this invention relates to the use of a combination of compounds as disclosed above for the manufacture of a medicament for the treatment of CMT or a related disorder.

A further object of this invention is a method for treating CMT or a related disorder, the method comprising administering to a subject in need thereof an effective amount of a composition as defined above.

A further object of this invention is a method of preparing a pharmaceutical composition, the method comprising mixing the above compounds in an appropriate excipient or carrier.

A more specific object of this invention is a method of treating CMT1a in a subject, the method comprising administering to the subject in need thereof an effective amount of a compound or combination of compounds as disclosed above.

5 A further specific object of this invention is a method of treating toxic neuropathy in a subject, the method comprising administering to the subject in need thereof an effective amount of a compound or combination of compounds as disclosed above.

10 A further specific object of this invention is a method of treating ALS in a subject, the method comprising administering to the subject in need thereof an effective amount of a compound or combination of compounds as disclosed above.

Therapy according to the invention may be performed as drug combination and/or in conjunction with any other therapy. It and may be provided at home, the doctor's
15 office, a clinic, a hospital's outpatient department, or a hospital, so that the doctor can observe the therapy's effects closely and make any adjustments that are needed.

The duration of the therapy depends on the stage of the disease being treated, the age and condition of the patient, and how the patient responds to the treatment.

20

Additionally, a person having a greater risk of developing an additional neuropathic disorder (e.g., a person who is genetically predisposed to or have, for example, diabetes, or is being under treatment for an oncological condition, etc.) may receive prophylactic treatment to alleviate or to delay eventual neuropathic response.

25

The dosage, frequency and mode of administration of each component of the combination can be controlled independently. For example, one drug may be administered orally while the second drug may be administered intramuscularly. Combination therapy may be given in on-and-off cycles that include rest periods so that
30 the patient's body has a chance to recovery from any as yet unforeseen side-effects. The drugs may also be formulated together such that one administration delivers both drugs.

Formulation of Pharmaceutical Compositions

The administration of each drug of the combination may be by any suitable means that results in a concentration of the drug that, combined with the other component, is able to ameliorate the patient condition (which may be determined e.g., in vitro by an effect on elevated expression of PMP22 upon reaching the peripheral nerves).

While it is possible for the active ingredients of the combination to be administered as the pure chemical it is preferable to present them as a pharmaceutical composition, also referred to in this context as pharmaceutical formulation. Possible compositions include those suitable for oral, rectal, topical (including transdermal, buccal and sublingual), or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration.

More commonly these pharmaceutical formulations are prescribed to the patient in "patient packs" containing a number dosing units or other means for administration of metered unit doses for use during a distinct treatment period in a single package, usually a blister pack. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions. Thus, the invention further includes a pharmaceutical formulation, as herein before described, in combination with packaging material suitable for said formulations. In such a patient pack the intended use of a formulation for the combination treatment can be inferred by instructions, facilities, provisions, adaptations and/or other means to help using the formulation most suitably for the treatment. Such measures make a patient pack specifically suitable for and adapted for use for treatment with the combination of the present invention.

30

The drug may be contained in any appropriate amount in any suitable carrier substance, and is may be present in an amount of 1-99% by weight of the total weight of

the composition. The composition may be provided in a dosage form that is suitable for the oral, parenteral (e.g., intravenously, intramuscularly), rectal, cutaneous, nasal, vaginal, inhalant, skin (patch), or ocular administration route. Thus, the composition may be in the form of, e.g., tablets, capsules, pills, powders, granulates, suspensions, emulsions, solutions, gels including hydrogels, pastes, ointments, creams, plasters, drenches, osmotic delivery devices, suppositories, enemas, injectables, implants, sprays, or aerosols.

The pharmaceutical compositions may be formulated according to conventional pharmaceutical practice (see, e.g., Remington: The Science and Practice of Pharmacy (20th ed.), ed. A. R. Gennaro, Lippincott Williams & Wilkins, 2000 and Encyclopedia of Pharmaceutical Technology, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York).

Pharmaceutical compositions according to the invention may be formulated to release the active drug substantially immediately upon administration or at any predetermined time or time period after administration.

The controlled release formulations include (i) formulations that create a substantially constant concentration of the drug within the body over an extended period of time; (ii) formulations that after a predetermined lag time create a substantially constant concentration of the drug within the body over an extended period of time; (iii) formulations that sustain drug action during a predetermined time period by maintaining a relatively, constant, effective drug level in the body with concomitant minimization of undesirable side effects associated with fluctuations in the plasma level of the active drug substance; (iv) formulations that localize drug action by, e.g., spatial placement of a controlled release composition adjacent to or in the diseased tissue or organ; and (v) formulations that target drug action by using carriers or chemical derivatives to deliver the drug to a particular target cell type.

30

Administration of drugs in the form of a controlled release formulation is especially preferred in cases in which the drug in combination, has (i) a narrow therapeutic index

(i.e., the difference between the plasma concentration leading to harmful side effects or toxic reactions and the plasma concentration leading to a therapeutic effect is small; in general, the therapeutic index, TI, is defined as the ratio of median lethal dose (LD50) to median effective dose (ED50)); (ii) a narrow absorption window in the gastro-intestinal tract; or (iii) a very short biological half-life so that frequent dosing during a day is required in order to sustain the plasma level at a therapeutic level.

Any of a number of strategies can be pursued in order to obtain controlled release in which the rate of release outweighs the rate of metabolism of the drug in question. Controlled release may be obtained by appropriate selection of various formulation parameters and ingredients, including, e.g., various types of controlled release compositions and coatings. Thus, the drug is formulated with appropriate excipients into a pharmaceutical composition that, upon administration, releases the drug in a controlled manner (single or multiple unit tablet or capsule compositions, oil solutions, suspensions, emulsions, microcapsules, microspheres, nanoparticles, patches, and liposomes).

Solid Dosage Forms for Oral Use

Formulations for oral use include tablets containing the active ingredient(s) in a mixture with non-toxic pharmaceutically acceptable excipients. These excipients may be, for example, inert diluents or fillers (e.g., sucrose, microcrystalline cellulose, starches including potato starch, calcium carbonate, sodium chloride, calcium phosphate, calcium sulfate, or sodium phosphate); granulating and disintegrating agents (e.g., cellulose derivatives including microcrystalline cellulose, starches including potato starch, croscarmellose sodium, alginates, or alginic acid); binding agents (e.g., acacia, alginic acid, sodium alginate, gelatin, starch, pregelatinized starch, microcrystalline cellulose, carboxymethylcellulose sodium, methylcellulose, hydroxypropyl methylcellulose, ethylcellulose, polyvinylpyrrolidone, or polyethylene glycol); and lubricating agents, glidants, and antiadhesives (e.g., stearic acid, silicas, or talc). Other pharmaceutically acceptable excipients can be colorants, flavoring agents, plasticizers, humectants, buffering agents, and the like.

The tablets may be uncoated or they may be coated by known techniques, optionally to delay disintegration and absorption in the gastrointestinal tract and thereby providing a sustained action over a longer period. The coating may be adapted to release the active
5 drug substance in a predetermined pattern (e.g., in order to achieve a controlled release formulation) or it may be adapted not to release the active drug substance until after passage of the stomach (enteric coating). The coating may be a sugar coating, a film coating (e.g., based on hydroxypropyl methylcellulose, methylcellulose, methyl hydroxyethylcellulose, hydroxypropylcellulose, carboxymethylcellulose, acrylate
10 copolymers, polyethylene glycols and/or polyvinylpyrrolidone), or an enteric coating (e.g., based on methacrylic acid copolymer, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, shellac, and/or ethylcellulose). A time delay material such as, e.g., glyceryl monostearate or glyceryl distearate may be employed.

15

The solid tablet compositions may include a coating adapted to protect the composition from unwanted chemical changes, (e.g., chemical degradation prior to the release of the active drug substance). The coating may be applied on the solid dosage form in a similar manner as that described in Encyclopedia of Pharmaceutical
20 Technology.

The drugs may be mixed together in the tablet, or may be partitioned. For example, a first drug is contained on the inside of the tablet, and a second drug is on the outside, such that a substantial portion of the second drug is released prior to the release of the
25 first drug.

Formulations for oral use may also be presented as chewable tablets, or as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent (e.g., potato starch, microcrystalline cellulose, calcium carbonate, calcium phosphate or
30 kaolin), or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, liquid paraffin, or olive oil. Powders and granulates may be

prepared using the ingredients mentioned above under tablets and capsules in a conventional manner.

Controlled release compositions for oral use may, e.g., be constructed to release the
5 active drug by controlling the dissolution and/or the diffusion of the active drug substance.

Dissolution or diffusion controlled release can be achieved by appropriate coating
of a tablet, capsule, pellet, or granulate formulation of drugs, or by incorporating the
10 drug into an appropriate matrix. A controlled release coating may include one or more
of the coating substances mentioned above and/or, e.g., shellac, beeswax, glycowax,
castor wax, carnauba wax, stearyl alcohol, glyceryl monostearate, glyceryl distearate,
glycerol palmitostearate, ethylcellulose, acrylic resins, dl-polylactic acid, cellulose
acetate butyrate, polyvinyl chloride, polyvinyl acetate, vinyl pyrrolidone, polyethylene,
15 polymethacrylate, methylmethacrylate, 2-hydroxymethacrylate, methacrylate hydrogels,
1,3 butylene glycol, ethylene glycol methacrylate, and/or polyethylene glycols. In a
controlled release matrix formulation, the matrix material may also include, e.g.,
hydrated methylcellulose, carnauba wax and stearyl alcohol, carbopol 934, silicone,
glyceryl tristearate, methyl acrylate-methyl methacrylate, polyvinyl chloride,
20 polyethylene, and/or halogenated fluorocarbon.

A controlled release composition containing one or more of the drugs of the
claimed combinations may also be in the form of a buoyant tablet or capsule (i.e., a
tablet or capsule that, upon oral administration, floats on top of the gastric content for a
25 certain period of time). A buoyant tablet formulation of the drug(s) can be prepared by
granulating a mixture of the drug(s) with excipients and 20-75% w/w of hydrocolloids,
such as hydroxyethylcellulose, hydroxypropylcellulose, or
hydroxypropylmethylcellulose. The obtained granules can then be compressed into
tablets. On contact with the gastric juice, the tablet forms a substantially water-
30 impermeable gel barrier around its surface. This gel barrier takes part in maintaining a
density of less than one, thereby allowing the tablet to remain buoyant in the gastric
juice.

Liquids for Oral Administration

Powders, dispersible powders, or granules suitable for preparation of an aqueous suspension by addition of water are convenient dosage forms for oral administration. Formulation as a suspension provides the active ingredient in a mixture with a dispersing or wetting agent, suspending agent, and one or more preservatives. Suitable suspending agents are, for example, sodium carboxymethylcellulose, methylcellulose, sodium alginate, and the like.

Parenteral Compositions

The pharmaceutical composition may also be administered parenterally by injection, infusion or implantation (intravenous, intramuscular, subcutaneous, or the like) in dosage forms, formulations, or via suitable delivery devices or implants containing conventional, non-toxic pharmaceutically acceptable carriers and adjuvants. The formulation and preparation of such compositions are well known to those skilled in the art of pharmaceutical formulation.

Compositions for parenteral use may be provided in unit dosage forms (e.g., in single-dose ampoules), or in vials containing several doses and in which a suitable preservative may be added (see below). The composition may be in form of a solution, a suspension, an emulsion, an infusion device, or a delivery device for implantation or it may be presented as a dry powder to be reconstituted with water or another suitable vehicle before use. Apart from the active drug(s), the composition may include suitable parenterally acceptable carriers and/or excipients. The active drug(s) may be incorporated into microspheres, microcapsules, nanoparticles, liposomes, or the like for controlled release. The composition may include suspending, solubilizing, stabilizing, pH-adjusting agents, and/or dispersing agents.

The pharmaceutical compositions according to the invention may be in the form suitable for sterile injection. To prepare such a composition, the suitable active drug(s) are dissolved or suspended in a parenterally acceptable liquid vehicle. Among

acceptable vehicles and solvents that may be employed are water, water adjusted to a suitable pH by addition of an appropriate amount of hydrochloric acid, sodium hydroxide or a suitable buffer, 1,3-butanediol, Ringer's solution, and isotonic sodium chloride solution. The aqueous formulation may also contain one or more preservatives
5 (e.g., methyl, ethyl or n-propyl p-hydroxybenzoate). In cases where one of the drugs is only sparingly or slightly soluble in water, a dissolution enhancing or solubilizing agent can be added, or the solvent may include 10-60% w/w of propylene glycol or the like.

Controlled release parenteral compositions may be in form of aqueous suspensions,
10 microspheres, microcapsules, magnetic microspheres, oil solutions, oil suspensions, or emulsions. Alternatively, the active drug(s) may be incorporated in biocompatible carriers, liposomes, nanoparticles, implants, or infusion devices. Materials for use in the preparation of microspheres and/or microcapsules are, e.g., biodegradable/bioerodible polymers such as polygalactin, poly-(isobutyl cyanoacrylate), poly(2-hydroxyethyl-L-
15 glutamine). Biocompatible carriers that may be used when formulating a controlled release parenteral formulation are carbohydrates (e.g., dextrans), proteins (e.g., albumin), lipoproteins, or antibodies. Materials for use in implants can be non-biodegradable (e.g., polydimethyl siloxane) or biodegradable (e.g., poly(caprolactone), poly(glycolic acid) or poly(ortho esters)).

20

Rectal Compositions

For rectal application, suitable dosage forms for a composition include suppositories (emulsion or suspension type), and rectal gelatin capsules (solutions or
25 suspensions). In a typical suppository formulation, the active drug(s) are combined with an appropriate pharmaceutically acceptable suppository base such as cocoa butter, esterified fatty acids, glycerinated gelatin, and various water-soluble or dispersible bases like polyethylene glycols. Various additives, enhancers, or surfactants may be incorporated.

30

Percutaneous and Topical Compositions

The pharmaceutical compositions may also be administered topically on the skin for percutaneous absorption in dosage forms or formulations containing conventionally non-toxic pharmaceutical acceptable carriers and excipients including microspheres and liposomes. The formulations include creams, ointments, lotions, liniments, gels, hydrogels, solutions, suspensions, sticks, sprays, pastes, plasters, and other kinds of transdermal drug delivery systems. The pharmaceutically acceptable carriers or excipients may include emulsifying agents, antioxidants, buffering agents, preservatives, humectants, penetration enhancers, chelating agents, gel-forming agents, ointment bases, perfumes, and skin protective agents.

The emulsifying agents may be naturally occurring gums (e.g., gum acacia or gum tragacanth)

The preservatives, humectants, penetration enhancers may be parabens, such as methyl or propyl p-hydroxybenzoate, and benzalkonium chloride, glycerin, propylene glycol, urea, etc.

The pharmaceutical compositions described above for topical administration on the skin may also be used in connection with topical administration onto or close to the part of the body that is to be treated. The compositions may be adapted for direct application or for application by means of special drug delivery devices such as dressings or alternatively plasters, pads, sponges, strips, or other forms of suitable flexible material.

Dosages and duration of the treatment

It will be appreciated that the drugs of the combination may be administered concomitantly, either in the same or different pharmaceutical formulation or sequentially. If there is sequential administration, the delay in administering one of the active ingredients should not be such as to lose the benefit of the efficacious effect of the combination of the active ingredients. A minimum requirement for a combination according to this description is that the combination should be intended for combined

use with the benefit of the efficacious effect of the combination of the active ingredients. The intended use of a combination can be inferred by facilities, provisions, adaptations and/or other means to help using the combination according to the invention.

5 Therapeutically effective amounts of the drugs that are subject of this invention can be used together for the preparation of a medicament useful for reducing the effect of increased expression of PMP22 gene; restoration of normal myelination and nerve integrity, preventing or reducing the risk of developing CMT disease, halting or slowing the progression of CMT disease once it has become clinically manifest, and preventing
10 or reducing the risk of a first or subsequent occurrence of an neuropathic event.

Although the active drugs of the present invention may be administered in divided doses, for example two or three times daily, a single daily dose of each drug in the combination is preferred, with a single daily dose of all drugs in a single pharmaceutical composition (unit dosage form) being most preferred.

15 Administration can be one to several times daily for several days to several years, and may even be for the life of the patient. Chronic or at least periodically repeated long-term administration will be indicated in most cases.

• The term “unit dosage form” refers to physically discrete units (such as capsules, tablets, or loaded syringe cylinders) suitable as unitary dosages for human
20 subjects, each unit containing a predetermined quantity of active material or materials calculated to produce the desired therapeutic effect, in association with the required pharmaceutical carrier.

The amount of each drug in the combination preferred for a unit dosage will depend
25 upon several factors including the administration method, the body weight and the age of the patient, the severity of the neuropathic damage caused by CMT disease or risk of potential side effects considering the general health status of the person to be treated.

Additionally, pharmacogenomic (the effect of genotype on the pharmacokinetic, pharmacodynamic or efficacy profile of a therapeutic) information about a particular
30 patient may affect the dosage used.

Except when responding to especially impairing CMT disease cases when higher dosages may be required, or when treating children when lower dosages should be

chosen, the preferred dosage of each drug in the combination will usually lie within the range of doses not above the usually prescribed for long-term maintenance treatment or proven to be safe in the large phase 3 clinical studies.

5 For example,

- for Rapamycin, from about 1 to about 100 $\mu\text{g/kg}$ per day, typically from 1 to 50 $\mu\text{g/kg}$, for instance between 5 and 30 $\mu\text{g/kg/day}$.
- for Mifepristone, from about 1 to about 300 $\mu\text{g/kg}$ per day, typically from 10 to 200 $\mu\text{g/kg}$, for instance between 10 and 80 $\mu\text{g/kg/day}$.
- 10 • for Naltrexone, from about 1 to about 100 $\mu\text{g/kg}$ per day, typically from 1 to 50 $\mu\text{g/Kg}$, for instance between 1 and 20 $\mu\text{g/kg/day}$.
- for Pilocarpin, from about 1 to about 100 $\mu\text{g/kg}$ per day, typically from 1 to 50 $\mu\text{g/Kg}$, for instance between 1 and 20 $\mu\text{g/kg/day}$.
- for Baclofen, from about 1 to about 300 $\mu\text{g/kg}$ per day, typically from 10
15 to 200 $\mu\text{g/kg}$, for instance between 20 and 100 $\mu\text{g/kg/day}$.
- for Methimazole, from about 1 to about 100 $\mu\text{g/kg}$ per day, typically from 1 to 50 $\mu\text{g/kg}$, for instance between 1 and 20 $\mu\text{g/kg/day}$.

The most preferred dosage will correspond to amounts from 1% up to 10% of those usually prescribed for long-term maintenance treatment.

20 It will be understood that the amount of the drug actually administered will be determined by a physician, in the light of the relevant circumstances including the condition or conditions to be treated, the exact composition to be administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration. Therefore, the above dosage ranges are intended
25 to provide general guidance and support for the teachings herein, but are not intended to limit the scope of the invention.

The following examples are given for purposes of illustration and not by way of limitation.

30

Examples

A. Preparation of drug combinations

5

The following drug combinations were prepared:

	Molecule	dose
Mix1	Sorbitol	2,1 mg/kg/day
	S-Baclofen (-)	60 µg/kg/day
	Naltrexone	7 µg/kg/day

10

	Molecule	dose
Mix2	Rapamycin	15 µg/kg/day
	Mifepristone	40 µg/kg/day

	Molecule	dose
Mix3	Rapamycin	15 µg/kg/day
	Mifepristone	40 µg/kg/day
	Pilocarpin	7 µg/kg/day

15

	Molecule	dose
Mix4	Rapamycin	15 µg/kg/day
	Mifepristone	40 µg/kg/day
	Baclofen	60 µg/kg/day

	Molecule	dose
Mix5	Rapamycin	15 µg/kg/day
	Mifepristone	40 µg/kg/day
	Methimazole	4,2 µg/kg/day

5

	Molecule	dose
Mix6	Rapamycin	15 µg/kg/day
	Naltrexone	7 µg/kg/day
	Methimazole	4,2 µg/kg/day

	Molecule	dose 1	dose 2	dose 3
Mix7	Sorbitol	10,5 mg/kg/day	2,1 mg/kg/day	1,05 mg/kg/day
	(RS) Baclofen	0,3 mg/kg/day	60 µg/kg/day	30 µg/kg/day
	Naltrexone	35 µg/kg/day	7 µg/kg/day	3,5 µg/kg/day

10

B. In vitro Experiments

1. PMP22 expression assays on Schwann cells treated with Mix1-6

5 1.1 Cell culture

1.1.1: Commercially available rat primary Schwann cells

Vials of rat Schwann cells (SC) primary culture (Sciencell # R1700) are defrost and seeded at the density of 10 000 cells/cm² in “Sciencell Schwann cell medium” (basal medium from Sciencell # R1701) in poly-L-lysine pre-coated 75 cm² flasks. The culture medium is composed of basal medium, 5% Fetal Bovine Serum (3H-Biomedical AB #1701-0025), 1% Schwann cell growth supplement (3H Biomedical AB #1701-1752), 1% Gentamicin (Sigma #G1397) and 10µM of Forskolin (Sigma # F6886) to promote their proliferation.

15 After reaching confluency (4 to 10 days depending on cell batch), Schwann cells are purified by gentle agitation or by thy1.1 immunopanning that allow SC isolation from adherent fibroblasts, to produce cultures that are at least 95% pure. SC are then counted (Tryptan blue method) and seeded in poly-L-lysine pre-coated 75 cm² flask in the same SC medium. At confluency, cells are rinsed, trypsinized (trypsin-EDTA 1x diluted from
20 Invitrogen #1540054), diluted in PBS without calcium and magnesium) counted and plated in 12 well-dishes (140 000 cells/well) in Sciencell Schwann cell medium with 5% of FBS, 1% of cell growth supplement (CGS), 40µg/ml of gentamicin and 4µM Forskolin.

25 *1.1.2 Custom -made rat primary Schwann cells*

Primary Schwann cell cultures (SC) are established from Sprague-Dawley newborn rats (between P0 and P2) sciatic nerves. All newborn rats are sacrificed and isolated in a Petri dish. Dissection is performed under sterile conditions.

The dorsal skin is removed from the hind paw and the lower torso. The sciatic nerve is isolated and transferred to a culture dish containing ice-cold Leibovitz (L15, Invitrogen
30 #11415) supplemented with 1% penicillin/streptomycin solution (50UI/ml and 50µg/ml,

respectively; Invitrogen #15070) and 1% of bovine serum albumin (BSA, Sigma A6003). Both nerves per rats are transferred in a 15ml tube containing ice-cold L15. The L15 medium is then removed and replaced by 2.4ml of DMEM (Invitrogen #21969035) with 10mg/ml of collagenase (Sigma #A6003). Nerves are incubated in this
5 medium for 30 minutes at 37°C. The medium is then removed and both nerves are dissociated by trypsin (10% trypsin EDTA 10x, Invitrogen #15400054) diluted in PBS without calcium and magnesium (Invitrogen # 2007-03) for 20min at 37°C. The reaction is stopped by addition of DMEM containing DNase I grade II (0.1mg/ml Roche diagnostic #104159) and foetal calf serum (FCS 10%, Invitrogen #10270). The cell
10 suspension was triturated with a 10ml pipette and passed through a filter in a 50ml tube (Swinnex 13mm filter units, Millipore, with 20µm nylon-mesh filters, Fisher). The cell suspension is centrifuged at 350g for 10min at room temperature (RT) and the pellets are suspended in DMEM with 10% FCS and 1% penicillin/streptomycin. Cells are counted (Trypan blue method) and seeded in Falcon 100mm Primaria tissue culture
15 plates at the density of $5 \cdot 10^5$ to 10^6 cells/dish.

After one day of culture, the medium is changed with DMEM, 10% FCS, 1% penicillin/streptomycin and 10µM of cytosine b-D-arabinofuranoside (Sigma #C1768). 48hrs later, medium is eliminated and cells are washed three times with DMEM. The SC growth medium is then added, composed of DMEM, 10% FCS, 1%
20 penicillin/streptomycin, 2µM of Forskolin (Sigma #F6886), 10µg/ml of bovine pituitary extract (PEX, Invitrogen #13028). The medium is replaced every 2-3 days.

After 8 days of culture (4 to 10 days depending on cell batches), Schwann cells reach confluency and the culture, containing a large amount of contaminating fibroblasts, is purified by the thy1.1 immunopanning method. After this purification, cells are
25 suspended in growth medium at 10 000 cells/cm² in poly-L-lysine pre-coated 75 cm² flasks. Once they reach confluency, cells are rinsed, trypsinized (trypsin-EDTA), counted and plated in 12 well-dishes (100 000 cells/well).

1.1.3 Drug incubation

30 After cells being plated in 12well-dishes, the medium is replaced by a defined medium consisting in a mix of DMEM-F12 (Invitrogen # 21331020) complemented by 1% of

N2 supplement (Invitrogen # 17502), 1% L-Glutamine (Invitrogen #25030024) 2.5% FBS (Sciencell #0025), 0.02 µg/ml of corticosterone (Sigma # C2505), 4µM Forskolin and 50µg/ml of gentamycin. Growth factors are not added to this medium, to promote SC differentiation

- 5 24 hours later, the medium is replaced by a defined medium (DMEM-F12) complemented with 1 % Insulin-Transferrin-Selenium – X (ITS, Invitrogen # 51300), 16µg/ml of Putrescine (Sigma # P5780), 0.02 µg/ml of corticosterone and 50µg/ml of gentamicin. At this step, neither progesterone nor forskolin are present in the medium.

One day later, Schwann cells are stimulated by combinations of drugs during 24hrs (3
10 wells/condition). The preparation of each compound is performed just prior to its addition to the cell culture medium.

Drugs are added to a defined medium composed of DMEM-F12, with 1 % Insulin-Transferrin-Selenium – X (ITS, Invitrogen # 51300), 16µg/ml of Putrescine, 0.02 µg/ml of corticosterone, 10nM Progesterone and 50µg/ml of gentamicin. The absence of
15 Forskolin during drug stimulation avoids adenylate cyclase saturation.

1.2. Schwann cells purification by Thy1.1 immunopanning

To prevent fibroblast culture contamination, Schwann cells are purified using the clone
20 Thy1.1 (ATCC TIB-103TM) immunopanning protocole.

Antibody pre-coated 100mm bacteria Petri dishes are prepared as follows: these dishes are washed three times with PBS and treated by 20ml of Tris HCl solution 50 mM, pH 9.5, with 10 µg/ml of goat Anti-Mouse IgM MU antibody (Jackson ImmunoResearch #115-005-020) overnight at 4°C; then rinsed 3 times with PBS and treated by a solution
25 of PBS with 0.02% of BSA and supernatant obtained from T11D7e2 hybridoma culture (ATCC #TIB-103) containing the Thy1.1 IgM antibody for 2hours at room temperature. Finally, the plates are washed three times with PBS before the cell suspensions are added.

SC are detached with trypsin EDTA. As soon as the majority of cells are in suspension,
30 the trypsin is neutralized with DMEM-10% FBS and the cells are centrifuged. The

pellet of dissociated cells is resuspended in 15ml of medium with 0.02% BSA at the density of 0.66×10^6 cells per ml (maximum) and transferred to Petri dish (about 6.6 million of cells/10ml/dish of 100mm).

The cell suspension is incubated in the Thy 1.1 coated Petri dish during 45 min at 37°C with gentle agitation every 15 min to prevent non-specific binding. The majority of fibroblast cells expressing Thy1.1 adhere on the dish. At the end of the incubation, the cell suspension is recovered and centrifuged. This cell suspension contains in theory only Schwann cells. Cells are centrifuged and cell pellet is suspended in growth medium with 10μM of Forskolin at 16 000 cells/cm² in T75 cm² flask Poly-L-Lysine treated.

1.3 Quantitative reverse transcriptase polymerase chain reaction (Q- RT-PCR)

Quantitative RT-PCR is used to compare the levels of PMP22 mRNA after drug stimulation, relative with housekeeping Ribosomal L13A mRNA in rat Schwann cell primary culture.

After rinsing with cold sterilized PBS, total RNAs from each cell sample are extracted and purified from SC using the Qiagen RNeasy micro kit (Qiagen #74004). Nucleic acids are quantified by Nanodrop spectrophotometer using 1μl of RNA sample. The RNA integrity is determined through a BioAnalyzer (Agilent) apparatus.

RNAs are reverse-transcribed into cDNA according to standard protocol. cDNA templates for PCR amplification are synthesized from 200ng of total RNA using SuperScript II reverse-transcriptase (Invitrogen # 18064-014) for 60 min at 42°C in the presence of oligo(dT), in a final volume of 20μl.

cDNAs are subjected to PCR amplification using the «LightCycler® 480» system (Roche Molecular Systems Inc.) Each cDNA are diluted five times before being used for PCR amplification. 2.5μl of this cDNAs enters the PCR reaction solution (final volume of 10μl). Preliminary experiments ensured that quantitation was done in the exponential phase of the amplification process for both sequences and that expression of the reference gene was uniform in the different culture conditions.

PCR reaction is performed by amplification of 500nM of forward primer of rat PMP22 (NM_017037), 5-GGAAACGCGAATGAGGC-3 (SEQ ID NO: 1), and 500nM of reverse primer 5-GTTCTGTTTGGTTTGGCTT-3 (SEQ ID NO: 2) (amplification of 148-bp). A 152-bp fragment of the RPL13A ribosomal (NM_173340) RNA is amplified
5 in parallel in separate reactions for normalization of the results by using a 500nM of forward primer 5-CTGCCCTCAAGGTTGTG-3 (SEQ ID NO: 3), and a 500nM of reverse primer 5- CTTCTTCTTCCGGTAATGGAT-3 (SEQ ID NO: 4).

We used FRET chemistry to perform RT-Q-PCR analysis. FRET probes are composed of 0.3μM of Pmp22-FL-5-GCTCTGAGCGTGCATAGGGTAC (SEQ ID NO: 5) or
10 Rpl13A-FL- 5-TCGGGTGGAAGTACCAGCC (SEQ ID NO: 6), labelled at their 3' end with a donor fluorophore dye (Fluorescein). 0.15μM Red640 probes are defined as follows: Pmp22-red-5'-AGGGAGGGAGGAAGGAAACCAGAAA (SEQ ID NO: 7) or Rpl13A-red-5'-TGACAGCTACTCTGGAGGAGAAACGGAA (SEQ ID NO: 8), labelled at their 5' end with an acceptor fluorophore dye (Rhodamine Red 640).

15 Each PCR reaction contained 2.5μl cDNA template in a final volume of 10μl of master mix kit (Roche #04-887301001).

The following PCR conditions are used: 10sec at 95°C, 10sec at 63°C and 12 sec at 72°C and 30sec at 40°C (Forty amplification cycles). The relative levels of PMP22 gene expression is measured by determining the ratio between the products generated from
20 the target gene PMP22 and the endogenous internal standard RPL13A.

1.4 PMP22 protein expression analysis by flow cytometry (FACS)

8hrs, 24hrs and 48hrs after drugs incubation, supernatants are recovered, centrifuged
25 and frozen. SC are detached with trypsin-EDTA. As soon as the majority of cells are in suspension, the trypsin is neutralised using DMEM with 10% FCS.

Supernatants with cells are recovered and centrifuged. The pellets of cells are transferred in micro tubes, washed in PBS once and fixed with a specific solution (AbCys #Reagent A BUF09B). 10 minutes later, cells are rinsed once with PBS and
30 kept at 4°C.

Five days after cell fixation, all cell preparations with different incubation times are labelled using the following protocol.

Cells are centrifuged at 7000 rpm for 5 minutes and the pellets are suspended in a solution of permeabilization (AbCys #Reagent B BUF09B) and labelled with primary
5 PMP22 antibody (Abcam #ab61220, 1/50) for 1hr room at temperature. Cells are then centrifuged at 7000 rpm for 5 minutes and cell pellets are rinsed once in PBS. A secondary antibody is added, coupled to Alexa Fluor 488 (goat anti-rabbit IgG, Molecular Probes #A11008, 1/100), for one hour at room temperature. Cells are then
10 centrifuged at 7000 rpm for 5 minutes and cell pellets are rinsed once in PBS. The labelling is increased adding a tertiary antibody coupled to Alexa Fluor 488 (chicken anti-goat IgG, Molecular Probes #A21467, 1/100) for one hour incubation, at room temperature. Cells are then rinsed once in PBS. Control without any antibody (unlabelled cells) is performed to determine the level of autofluorescence and adapted the sensitivity of the photomultipliers. Control with both secondary and tertiary
15 antibodies but without primary antibody, is performed to assess non specific binding of antibodies.

Data acquisition and analysis are performed with a FACS Array cytometer and FACS Array software (Becton Dickinson) on 5000 cells. Forward Scatter (FSC) correlated with cell volume (size) and Side Scatter (SSC) depending on inner complexity of cells
20 (granularity) are analysed. For expression of PMP22, analysis is performed within the total cells and percent of positive cells is calculated. Positive cells are cells with fluorescence intensity higher than the control with secondary antibody.

In order to quantify the number of SC, cells in control medium are analysed using antibodies anti-S100 Protein.

25 Cells are prepared according to the following protocol: Schwann cells are stained with antibody anti-S100 Protein (Dako #S0311, 1/100) for 1hr room at temperature. This antibody is labelled according to protocol described above for PMP22 immunostaining but without incubation with tertiary antibody.

30 1.5. Drug incubation and activity

Drugs are incubated for 24hrs or 48hrs in the same defined medium than described above (3 wells/condition) in absence of Forskolin to avoid adenylate cyclase stimulation saturation, but in presence of 10nM of progesterone. After drug incubation, supernatants are recovered and Schwann cells are frozen for RT-Q-PCR analysis.

5

These experiments are summarized in Table 1.

Table 1

Combination	PMP22 expression
Mix1	down regulation
Mix2	down regulation
Mix3	down regulation
Mix4	down regulation
Mix5	down regulation
Mix6	down regulation

10

2. Assessment of synergistic effect of compounds in Mix7 in a co-culture model for CMT

15

A model of co-culture was used as an *in vitro* model of CMT1A. This model of myelination consists in co-culturing sensory neurons and Schwann cells from male PMP22 Transgenic (TG) dissociated Dorsal Root Ganglia (DRG).

The aim of this study is to assess the effect of 3 test compounds (+/-Baclofen, Naltrexone and Sorbitol) and Mix7 (a mixture of these 3 drugs) on myelination process. The effect of the 3 test compounds, and their mixture on myelination, are assessed by evaluating Myelin Basic Protein (MBP) expression in presence of ascorbic acid.

20

2.1 Materials and Methods

25

15 days gestation pregnant female rats are killed by cervical dislocation. The embryos are removed from the uterus and are at similar fetal stage of development.

2.1.1 Genotyping

- A piece of each embryo head (3 mm^3) is placed in a 2 ml tube DNase free. The DNA is extracted with the SYBR Green Extract-N-Amp tissue PCR kit (Sigma, ref XNATG-1KT). 120 μl of extraction solution (Kit Sigma, ref XNATG-1KT) was put on each piece of embryo head. The heads are incubated for 10 min at room temperature. At the end of this incubation, the heads are incubated for 5 min at 95°C in the extraction solution. Immediately after this last incubation, 100 μl of neutralizing solution are added, each DNA extract is diluted at 1/40 with sterile ultrapure water (Biosolve, ref: 91589) and stored at $+4^\circ\text{C}$ until use. The genotyping of female (F) and male (M) embryos is performed during the dissection of the DRG, with the kit Fast SYBR Green Master Mix (Applied Biosystem, 4385612). The gender of each embryo is determined using the male SRY gene. The SRY primers are supplied by Pharnext (SRY-F (SEQ ID NO:9): 5'-GAGAGAGGCACAAGTTGGC-3'; SRY-R (SEQ ID NO:10): 5'-GCCTCCTGGAAAAAGGGCC-3'). SRY primers are diluted at $3\mu\text{M}$ in sterile ultrapure water (Biosolve, ref: 91589). A mix for PCR is prepared with ultrapure water (4 μl per sample), primer at $3\mu\text{M}$ (2 μl per sample) and Master Mix (10 μl per sample). In a PCR 96 wells plate, 16 μl of PCR mix is deposited in each well. 4 μl of each diluted DNA is added according to a plan deposit. The PCR is run using the 7500 fast RT-PCR system (Applied Biosystem), with the following program:
- Beginning: 95°C – 20 sec
- 45 cycles: 95°C – 10 sec, 65°C – 10 sec, 72°C – 30 sec (data acquisition).
- Melt curve: 95°C – 15 sec, 64°C – 1min, 90°C – 30 sec (continuous data acquisition), 60°C 15 sec. The amplification plots and melt curves are analyzed with the 7500 software (Applied Biosystems).
- The results for each sample are compared to negative control (ultrapure water) and to the positive control (TG/Male and WT/Female), to conclude on the genotype of each embryo.

2.1.2 Sensory neurons and Schwann cells co-cultures

- Rat Dorsal root ganglions are cultured as previously described by Cosgaya et al., 2002 and Rangaraju et al., 2008.

Each embryo is dispatched on numerating petri dish (35 mm of diameter). The head of embryo is cut, placed on 1.5 ml tube DNAase free; the ADN is extracted with the Extract-N-Amp Tissue Kit (Sigma Aldrich). The genotyping (Male (M) and female (F), wild type and PMP22 transgenic) is performed with the kit Fast SYBR Green Master Mix (Applied Biosystem). This genotyping is performed in parallel of the dissection of dorsal root ganglia (DRG), so that at the end of the dissection, only one type of culture (DRG from transgenic male) is done. DRG of each embryo is collected, placed in ice-cold medium of Leibovitz (L15, Invitrogen). At the end of the dissection, DRG of TGM are pooled and dissociated by trypsinization (trypsin EDTA, 0.05%; Invitrogen) for 20 min at 37°C. The reaction is stopped by addition of DMEM containing 10% of fetal bovine serum (FBS) in the presence of DNAase I (Roche). The suspension is triturated with a 10 ml pipette. Cells are then centrifuged at 350 x g for 10 min at room temperature. The pellet of dissociated cells is resuspended in neurobasal medium (Invitrogen) containing 2 % of B27 (Invitrogen), 1% of penicillin-streptomycin (Invitrogen), 1 % de L glutamine and 50 ng/ml NGF (Sigma). This medium is the neuronal medium. Viable cells are counted in a Neubauer cytometer using the trypan blue exclusion test (Sigma) and seeded on the basis of 10 000 cells per well in 96 well-plates (Greiner) treated with poly-L-lysine. The plates are maintained at 37°C in a humidified incubator, in an atmosphere of air (95%)-CO₂ (5%). Half of the standard neuronal culture medium is changed every other day. The cultures are maintained in standard neurobasal medium for 7 days to allow Schwann cells to populate the sensory neuron neurites. On day 7, the cultures are fed with standard neuronal medium supplemented or not with 50 µg/ml ascorbic acid in order to initiate basal lamina formation and myelination.

25

2.1.3. Drug incubation

On day 7, the following test compounds (alone or in combination) are added in the medium with 50 µg/ml ascorbic acid:

30

- (RS)-Baclofen
- Natrexone
- d-Sorbitol
- Mix7= the combination of the 3 individual compounds.

These compounds or compound combination are tested at the following concentrations (Table 2):

		Dose 1	Dose 2	Dose 3	Dose 4	Dose 5	Dose 6	Dose 7
Individual drugs	Naltrexone	5 μ M	1 μ M	200nM	40nM	8nM	1.6nM	320pM
	d-Sorbitol	500 μ M	100 μ M	20 μ M	4 μ M	800nM	160nM	32nM
	(RS)- Baclofen	5 μ M	1 μ M	200nM	40nM	8nM	1.6nM	320pM
Mix7	Naltrexone	5 μ M	1 μ M	200nM	40nM	8nM	1.6nM	320pM
	d-Sorbitol	500 μ M	100 μ M	20 μ M	4 μ M	800nM	160nM	32nM
	(RS)- Baclofen	5 μ M	1 μ M	200nM	40nM	8nM	1.6nM	320pM

- 5 **Table 2:** Concentration of individual drugs or in combination used for in vitro studies of MBP expression in TG DRG/SC co-cultures.

The test compounds are incubated for 5 different times: 5, 9, 10, 11 and 13 days.

- 10 Three separate and independent cultures of DRG (from TG embryos male rats) are done. These conditions are assessed in presence of ascorbic acid, 6 wells per condition. The solution ready to use of all test compounds are extemporaneously prepared from a stock solution, stored at -20°C. This solution is prepared once a week. Half of the standard neuronal medium supplemented with test compounds and ascorbic acid (each at the concentration 1X) are changed every other day.

15

2.1.4 Staining protocol

After 5, 9, 10, 11 and 13 days of incubation, cells are fixed by a cold solution of ethanol (95%) and acetic acid (5%) for 10 min. The cells are permeabilized and blocked with PBS containing 0.1% saponin and 10% goat serum for 15 min. Then, the cells are

incubated with a specific marker of myelin: polyclonal antibody anti-myelin basic protein (MBP) antibody (Sigma 118K0431).

This antibody is revealed with Alexa Fluor 568 goat anti-rabbit IgG and Alexa Fluor 488 goat anti-mouse IgG (Molecular probe 687621, 623962). Nuclei of neurons are
5 labeled by a fluorescent marker (Hoechst solution, SIGMA ref B1155).

2.1.5. Data processing

Per well, 20 pictures are taken using InCell AnalyzerTM 1000 (GE Healthcare) with 20x magnification. All images are taken in the same conditions. Analysis of total length
10 of myelinated axons was automatically done (length and area around axons) using Developer software (GE Healthcare). All values will be expressed as mean $\pm$ s.e.mean. Statistic analyses are done on different conditions (ANOVA followed by Fisher's PLSD test when allowed).

15 2.2. Results

Synergistic effect of drugs in the efficacy of Mix7

An important synergistic effect of drugs composing the Mix7 combination is observed on MBP expression. Indeed, on day 10 (=17 days of culture), combination of (RS)-Baclofen, Naltrexone and d-Sorbitol significantly increases MBP expression at doses 1
20 and 6 as shown in Figures 1A and 2A. By contrast, the above drugs used individually have no substantial effect compared to control (Figures 1B-D and 2B-D). A significant effect on MBP expression is also recorded after 10 days of incubation at doses 2, 3, 4, 5 and 7 of Mix7 (Figure 3A).

25 This effect is still observed on day 11 with doses 2-7 (Figure 3B) in form of a clear bell shape curve.

C. Experiments in vivo in CMT animal model

30 We tested the compounds for therapeutic effect in a rat model.

The experimental groups are formed with young rats of both genders separately. The rats are assigned to the groups following randomization schedule based on the body

weight. In some experiments the randomization is based on the performances of the rats in the bar test. Both genders are represented by separate control groups that are numerically equal or bigger than the treatment groups.

- 5 The rats are treated chronically with drugs - force fed or injected by Alzet osmotic subcutaneous pump (DURECT Corporation Cupertino, CA), depending on each drug bioavailability during 3 or 6 weeks. In all the *in vivo* experiments performed, the Mix7 is administered by gavage.
- 10 The animals are weighted twice a week in order to adjust the doses to growing body weight. If the osmotic pump is chosen for the treatment administration, the doses of the drug are calculated on the basis of the estimated mean body weight of the animals expected for their age over the period of the pump duration (6 weeks). The pumps are re-implanted if necessary, with the appropriated anesthesia protocol.

15

Behavioural tests

- Each three or four weeks the animals are subjected to a behavioural test. Each test is conducted by the same investigator in the same room and at the same time of the day; this homogeneity is maintained throughout entire experiment. All treatments and
- 20 genotype determination are blinded for the investigator. "Bar test" and "Grip strength" has been mainly used to access the performance throughout study. The schedule of the bar test may change as the animal growth (in order to avoid the bias due to the learning, for example).

- 25 The assay of the grip strength allows detection of subtle differences in the grip performance that seems to be composed of the muscle force, sensitivity status (for instance, painful tactile feelings may change measured values of the force), behavioural component ("motivation"). The values differ between fore and hind limbs and greatly depend on the age of the animals.

30

The grip strength test measures the strength with which an animal holds on to a grip with its forepaws or its hindpaws separately. A dynamometer is placed with a grip to

measure the strength (Force Gauge FG-5000A). The rat is held by the experimenter in a way that it grasps the grip either with its forepaws or with its hind paws and pulls gently the rat backwards until it releases the grip. The force measured when the animal releases the grip is recorded.

5

Two successive trials measuring the forepaws and two successive trials measuring the hindpaws strength per animal are processed ; only the maximum score (one for forepaws and one for hindpaws) is noticed (in N).

10 The Bar Test

The bar test evaluates rats' ability to hold on a fix rod. Pmp22 rats which display muscular weakness, exhibit a performance deficit in this test (Sereda et al, 1996). The rat is placed on its four paws on the middle of the rod (diameter: 2.5 cm ; length : 50 cm; 30 cm above the table). Trials are performed consecutively; the number and the
15 duration of trials in our experiments have been depending on batches of the animals. This variability in the testing has been introduced in order to determine the schedule appropriated to the best detection of the motor deficiency in the CMT rats in the course of the experiments.

20 Performance indices are recorded on each session:

- The number of trials needed to hold for 60 sec (or 30 sec for batch 1, session 1 and 2) on the rod.
- The time spent on the bar (i.e. the fall latency) in each trial and the average on the session. In the experimental procedures where the session ends after the rat
25 has stayed for a cut-off time, i.e. 30 or 60 s, on the bar, a performance of the cut-off time (30 s or 60 s) is assigned to trials not completed (eg: for batch 8, for an animal which stays on the bar less than 10 sec on trials 1, 2 and 3, then for 60 sec on trials 4 and 5, 60 s is assigned to trials 6 to 10).
- The number of falls.

30

General health assessment

Body weights, overt signs (coat appearance, body posture, gait, tremor etc.) of the animals are monitored throughout the experiment. The rating scale is used for recording: 0= normal, 1=abnormal.

5 The gait

Each rat is observed in a novel rat cage (dimensions 55x33x18cm) without litter for five minutes. The gait of rats is evaluated with 4 parameters:

- Score 0: normal gait (fluid)
- Score 1: abnormal gait (not fluid or the rat has a slight limp)
- 10 -Score 2: moderate incapacity (the rat drags one's leg and is able to put it right and walk)
- Score 3: serious incapacity (the rat drags its one's or both hindpaws but is unable to put it/them right).

Inclined plane test

- 15 The sliding apparatus had a 30x50cm Plexiglas plane that could be inclined at an angle of 0° (horizontal) to 60°. Each rat was initially placed on the 25°-angled inclined plane in the up-headed position (head-up orientation); two trials separated by 1 min are performed. 30 min later, the same experiment is realized on a 35°-angled inclined plane then on 40°-angled inclined plane. During this time the rat was returned to its cage. The
- 20 plane is cleaned after each trial.

The performances of rats are evaluated by 4 different scores :

- Score 0: no slide
- Score 1: a little slide (one or two paws)
- Score 2: a moderate slide (4 paws) but not until the end of the plane
- 25 -Score 3: the rat is sliding until the very bottom of the plane.

Further tests

When appropriate, the rats are subjected to electrophysiological evaluation, histological measurement and the pmp22 RNA expression level in the sciatic nerve is quantified.

Quantification of pmp22 RNA in sciatic nerve by quantitative RT- PCR

Total RNA was isolated from left sciatic nerves using Qiazol (ref N°79306, Qiagen GmbH, Germany) followed by the single-step purification method with RNeasy Mini Kit (ref N° 74106, Qiagen GmbH, Germany) described by the manufacturer's protocol
5 (Qiagen-RNeasy Fibrous tissue Handbook). DNA contamination was removed by digestion with RNase-free DNase I by use of the DNA-free kit (Qiagen-Rnase-free dnase set 1500 Kunits, ref N° 1023460).

RNA concentrations are estimated by NanoDrop ND-1000 and a test of quality control was done by Agilent RNA 6000 nano chips on Agilent 2100 Bioanalyzer.

- 10 Reverse transcription and real-time PCR: Quantitative RT-PCR (RT-Q-PCR) was performed as follows: 80 ng of total RNA was reverse transcribed using SuperScript™ II Reverse Transcriptase (Invitrogen, Carlsbad, CA) with Oligo(dT)12-18 (Invitrogen, Carlsbad, CA) in a 20-μl reaction volume.

- Real-time PCR was performed with a rapid thermal cycler system (LightCycler® 480
15 II, 384-Well, Roche, Switzerland). Amplifications are performed in a 10μl total volume with primers concentration optimized between 130nM and 1μM. Primers and template are supplemented with LightCycler® 480 SYBR Green I Master (2× conc. Roche, Cat. Ref N° 04 887 352 001). Nucleotides, MgCl₂, Taq DNA polymerase and buffer are included in the mix. An amplification protocol incorporated an initial incubation at
20 95°C for 10min for the activation of Taq DNA polymerase followed by 45 cycles, with a 95°C denaturation for 10s, 60°C annealing for 40s and 72°C extension for 10s (detection of the fluorescent product was performed at the end of the 72°C extension period by a single acquisition mode) and ended by a cycle of melting curve with 95°C
25 is 0.11°C/s and detection of the fluorescent product was continuous). To confirm the amplification specificity, the PCR product from each primer pair was subjected to a melting curve analysis. Relative quantification was performed based on the crossing point (Cp value) for each of the PCR samples. The point at which the fluorescence of a sample rises above the background fluorescence is called the "crossing point (Cp) of the
30 sample. Rattus norvegicus Myelin Protein Zero (MPZ) gene was used for normalization (Sereda et al, 2006). The sequences of the primers (synthesized by Eurofins MWG Operon, Germany) used for the RT-Q-PCR analysis are:

PMP22 – forward: 5'-TGTACCACATCCGCCTTGG-3' (SEQ ID NO: 11) and
PMP22 – reverse: 5'-GAGCTGGCAGAAGAACAGGAAC-3' (SEQ ID NO: 12).
MPZ-forward: 5'-TGTTGCTGCTGTTGCTCTTC-3' (SEQ ID NO: 13) and
MPZ-reverse: 5'-TTGTGAAATTTCCCCTTCTCC-3' (SEQ ID NO: 14).

5

Results

Mix1 composition improves bar test performances throughout the treatment procedure (Fig 4).

10

Mix1 improves the gait score of transgenic rats after 3 and 6 weeks of treatment as shown in figure 5.

Mix1 increases the performances of transgenic rats after 3, 6, 9 and 12 weeks of treatment in the inclined plane test at 25° described in the figure 6.

15

Figure 7 illustrates the positive effect of Mix2 on gait score of transgenic rats at 25, 35 and 40° in the inclined plane test.

Mix7 (dose 3) significantly decreases the pmp22 RNA gene expression in the sciatic nerve of pmp22 transgenic rats (Figure 9).

20

The performances of pmp22 rats treated with Mix7 (dose 2 and dose 3) are improved in the inclined plane test at 35° (Figure 10). More specifically, 29 and 33% of rats belong to the good performance group, compared to 5% for TG placebo group, and 29 and 11% of rats belong to the poor performance group compared to 60% for the TG placebo group. P-value (versus the TG placebo) is equal to 0,0152 for the TG rats treated with Mix7-dose 2 and p-value is equal to 0,002 for the TG rats treated with Mix7-dose 3 versus the TG placebo).

25

30

Mix7-dose 3 significantly increases the fall latency time of pmp22 rats in bar test after 9 weeks of treatment (Figure 11): black dashed line, $p=4,56.10^{-2}$, $n=18$. Significant

difference between TG placebo rats (black plain line, n=20) and WT placebo rats (grey plain line, $p=3,82 \cdot 10^{-7}$, n=18) is also observed.

The Figure 12 illustrates the improvement of grip strength of the pmp22 rats treated
5 with the Mix7-dose 3.

The Figure 13 shows the significant correlation between the bar test latency time (after 9 weeks of treatment with the Mix7-dose 3) and the expression level of pmp22 RNA.

10 The Figure 14 displays the significant correlation between the bar test latency time after 9 weeks of treatment with Mix7-dose 3 and the conduction velocity of the sensitive nerve (tail).

Similar results are obtained for other combinations (see Table 3).

15

Table 3

Combination	PMP22 rat disease phenotype
Mix1	improvement
Mix2	improvement
Mix3	improvement
Mix4	improvement
Mix5	improvement
Mix6	improvement

These data show that, in vivo, the combinations and regimens of this invention allow effective treatment of CMT.

20

D. In vivo effect in a model of toxic neuropathy

The drug treatments or regimens are orally administered from the day before the first intraperitoneal injection of Oxaliplatin 3mg/kg (D -1) until the day before the last
25 testing day (D16). Animals belonging to the Oxaliplatin-treated group are dosed daily with distilled water (10 ml/kg). Animals are dosed with the tested treatment and distilled water daily during the morning whereas Oxaliplatin is administered on the afternoon.

During the testing days (i.e. D1, D4, D10), the treatment and distilled water are administered after the test. Regarding the testing day (D4), including compounds and vehicle administrations and Oxaliplatin injection, the treatment and distilled water are administered prior to the injection of Oxaliplatin after the test. Animals from the reference-treated group are dosed only during the testing days (i.e. D1, D4, D10 and D17).

Cold allodynia is assessed by measuring the responses to thermal non-nociceptive stimulation (acetone test) on D1 (around 24h after the first injection of Oxaliplatin 3 mg/kg (acute effect of Oxaliplatin), on D4, D10 and (chronic effect of Oxaliplatin) and on D17 (residual effect of Oxaliplatin one week after completion of treatment).

Testing is done using the acetone test 2h post-administration of the reference. The reference substance is Gabapentin, 100 mg/kg, per os (once a day x 4 testing days).

Acetone test

Cold allodynia is assessed using the acetone test. In this test, latency of hindpaw withdrawal is measured after application of a drop of acetone to the plantar surface of both hindpaws (reaction time) and the intensity of the response is scored (cold score).

Reaction time to the cooling effect of acetone is measured within 20 sec (cut-off) after acetone application. Responses to acetone are also graded to the following 4-point scale: 0 (no response); 1 (quick withdrawal, flick of the paw); 2 (prolonged withdrawal or marked flicking of the paw); 3 (repeated flicking of the paw with licking or biting).

Six trials by rat are performed. For each experimental group, the results are expressed as the cumulative cold score defined as the sum of the 6 scores for each rat together $\pm$ SEM. The minimum score being 0 (no response to any of the 6 trials) and the maximum possible score being 18 (repeated flicking and licking or biting of paws on each of the six trials).

Gabapentin source: Zhejiang Chiral Medicine Chemicals, China

Oxaliplatin source: Sigma, France

The results are depicted on Figure 8. They clearly show a protective effect of the
5 composition of this invention on oxaliplatin-induced neuropathy.

E. In vivo effect in a model of ALS

Animal model

10 We have chosen the SOD1^{G93A} rat model (generated by Howland et al) to mimic the Amyotrophic Lateral Sclerosis pathology. This model overexpresses the mutated SOD1 gene in spinal cord, many brain regions as well as peripheral tissues. The onset of the motor neuron disease of this model is about at 115 days; it appears as hind limb abnormal gait. In few days, the paralysis of hind limb arises.

15

Experimental procedures

We obtained colonies by crossing breeder SOD1^{G93A} rats with Sprague Dawley female rats. Heterozygous SOD1^{G93A} rats are identified with polymerase chain reaction (PCR) of tail DNA with primers specific for hSOD1 [1]. Animals are maintained in a room
20 with controlled illumination (lights on 0500–1900 h) and temperature (23±1°C), and given free access to food and water. All the animal procedures in the present study are carried out in accordance with the guidelines standards of animal care.

Body weight measurement was performed every week and behavioural tests began at an age of 60 days and continued until endpoint. The treatments are administered every day
25 per oral or subcutaneous way from the age of 5 weeks.

1. Observation test : characterization of the general aspect

Each rat was observed in a novel rat cage (dimensions 55x33x18cm) without litter for five minutes. 5 different parameters are recorded:

30 **The gait**

- score 0 : normal gait (fluid)
- score 1 : abnormal gait (not fluid or the rat has a slight limp)

- score 2 : moderate incapacity (the rat drags one's leg and is able to put it right and walk)
- score 3 : serious incapacity (the rat drags its one's or both hindpaws but is unable to put it/them right)

5 **The coat aspect**

- score 0 : clean and silky coat
- score 1 : piloerection or dirty coat

The tremor

- score 0 : no tremor
- 10 - score 1 : tremor

The body position

- score 0 : normal
- score 1: abnormal (flattened or arching its back)

The hindpaws position

- 15 - score 0 : normal
- score 1 : spread hindpaws

2. The motor score test : characterization of the motor deficit

This test evaluates the ability of rats to right themselves within 30 sec of being turned
20 on either side (righting reflex) (Gale et al) .

A non-parametrical scoring system was used following these criteria (Matsumoto et al, Thonhoff et al):

- score 0: the rat is unable to right itself from either side within 30 sec
- score 1: the rat is unable to right itself from only one side within 30 sec
- 25 - score 2: the rat is able to right itself from both sides within 30 sec but is unable to stand in the cage; it is always dragging some parts of body
- score 3: the rat is able to right itself from both sides within 30 sec, is unable to stand in the cage but is not dragging some parts of body
- score 4: the rat is able to right itself from both sides within 30 sec, is able to stand in
30 the cage but has visible functional deficits
- score 5: the rat is able to right itself from both sides within 30 sec, is able to stand in the cage and no visible functional deficits.

The end-point of disease is fixed at score 0; the rat is then euthanized.

3. Inclined plane test : characterization of the motor deficit

- 5 The sliding apparatus had a 30x50cm plexiglas plane that could be inclined at an angle of 0° (horizontal) to 60°. Each rat was initially placed on the 25°-angled inclined plane in the up-headed position (head-up orientation); two trials separated by 1 min are performed. 30 min later, the same experiment is realized on a 35°-angled inclined plane then on 40°-angled inclined plane. During this time the rat was returned to its cage. The
10 plane is cleaned after each trial.

The performances of rats are evaluated by 4 different scores:

- score 0: no slide
- score 1: a little slide (one or two paws)
- score 2: a moderate slide (4 paws) but not until the end of the plane
- 15 -score 3: the rat is sliding until the very bottom of the plane.

4. The wire mesh test : characterization of the motor ability in difficult situation

- A wire mesh was placed in contact with a box at the top (at an angle of 70°) and the edge of a table at the bottom. Each rat was placed on the bottom of the wire mesh and
20 motivated to ascend by placing their littermates in the box at the top. Each rat was trained once a week (3 trials).

The recorded parameter was the latency time to reach the top of the wire mesh.

5. The open field test : characterization of the locomotor activity

- 25 The locomotor activity was measured in a Plexiglas box (45x45x30 cm, Acti-Track by BIOSEB, Lyon, France) with 16 photo-cell beams following the two axes, 1 and 5 cm above the floor.

The spontaneous and exploratory activity of each rat was evaluated during 3 hours.

- 4 parameters are recorded (the total travelled distance, the number of rearings, the
30 percentage of travelled distance and of time spent in the center of the openfield).

Bibliography

- Amici SA, Dunn WA Jr, Murphy AJ, Adams NC, Gale NW, Valenzuela DM,
5 Yancopoulos GD, Notterpek L. Peripheral myelin protein 22 is in complex with
alpha6beta4 integrin, and its absence alters the Schwann cell basal lamina. *J Neurosci.*
2006; 26(4):1179-1189.
- Amici SA, Dunn WA Jr, Notterpek L. Developmental abnormalities in the nerves
of peripheral myelin protein 22-deficient mice. *J Neurosci Res.* 2007; 85(2): 238-249.
- 10 Atanasoski S, Scherer SS, Nave K-A, Suter U. Proliferation of Schwann Cells and
Regulation of Cyclin D1 Expression in an Animal Model of Charcot-Marie-Tooth
Disease Type 1A. *J Neurosci Res.* 2002; 67(4):443-449.
- Basta-Kaim A, Budziszewska B, Jaworska-Feil L, Tetich M, Leśkiewicz M,
Kubera M, Lasoń W. Chlorpromazine inhibits the glucocorticoid receptor-mediated
15 gene transcription in a calcium-dependent manner. *Neuropharmacology.*
2002;43(6):1035-1043
- Batty IH, Fleming IN, Downes CP. Muscarinic-receptor-mediated inhibition of
insulin-like growth factor-1 receptor-stimulated phosphoinositide 3-kinase signalling in
1321N1 astrocytoma cells. *Biochem J.* 2004; 379(Pt 3):641-651.
- 20 Bogoyevitch MA, Ketterman AJ, Sugden PH. Cellular stresses differentially
activate c-Jun N-terminal protein kinases and extracellular signal-regulated protein
kinases in cultured ventricular myocytes. *J Biol Chem.* 1995;270(50):29710-29717.
- Brancolini C, Marzinotto S, Edomi P, Agostoni E, Fiorentini C, Müller HW,
Schneider C. Rho-dependent regulation of cell spreading by the tetraspan membrane
25 protein Gas3/PMP22. *Mol. Biol. Cell* 1999; 10: 2441-2459.
- Castellone MD, Teramoto H, Gutkind JS. Cyclooxygenase-2 and Colorectal Cancer
Chemoprevention: The β -Catenin Connection. *Cancer Res.* 2006; 66(23):11085-11088.
- Chen XR, Besson VC, Palmier B, Garcia Y, Plotkine M, Marchand-Leroux C.
Neurological recovery-promoting, anti-inflammatory, and anti-oxidative effects
30 afforded by fenofibrate, a PPAR alpha agonist, in traumatic brain injury. *J Neurotrauma*
2007; 24 (7): 1119-1131.

Chies R, Nobbio L, Edomi P, Schenone A, Schneider C, Brancolini C. Alterations in the Arf6-regulated plasma membrane endosomal recycling pathway in cells overexpressing the tetraspan protein Gas3/PMP22. *J Cell Sci.* 2003; 116(Pt 6): 987-999.

Constable AL, Armati PJ. DMSO induction of the leukotriene LTC₄ by Lewis rat Schwann cells. *J Neurol Sci* 1999; 162(2): 120-126.

Cosgaya J. M., Chan J. R., Shooter E. M. The Neurotrophin Receptor p75^{NTR} as a Positive Modulator of Myelination. *Science.* 2002; 298; 1245-1248.

Devaux JJ, Scherer SS. Altered ion channels in an animal model of Charcot-Marie-Tooth disease type IA. *J Neurosci.* 2005; 25(6): 1470-1480.

10 Diep QN, Benkirane K, Amiri F, Cohn JS, Endemann D, Schiffrin EL. PPAR alpha activator fenofibrate inhibits myocardial inflammation and fibrosis in angiotensin II-infused rats. *J Mol Cell Cardiol.* 2004; 36 (2): 295-304.

Dracheva S, Davis KL, Chin B, Woo DA, Schmeidler J, Haroutunian V. Myelin-associated mRNA and protein expression deficits in the anterior cingulate cortex and hippocampus in elderly schizophrenia patients. *Neurobiol Dis.* 2006 Mar;21(3):531-540.

D'Urso D, Ehrhardt P, Müller HW. Peripheral myelin protein 22 and protein zero: a novel association in peripheral nervous system myelin. *J Neurosci.* 1999; 19(9):3396-3403.

20 Fortun J, Dunn WA Jr, Joy S, Li J, Notterpek L. Emerging role for autophagy in the removal of aggresomes in Schwann cells. *J Neurosci.* 2003; 23(33): 10672-10680.

Fortun J, Li J, Go J, Fenstermaker A, Fletcher BS, Notterpek L. Impaired proteasome activity and accumulation of ubiquitinated substrates in a hereditary neuropathy model. *J Neurochem* 2005; 92:1531-1541.

25 Fortun J, Go JC, Li J, Amici SA, Dunn WA Jr, Notterpek L. Alterations in degradative pathways and protein aggregation in a neuropathy model based on PMP22 overexpression. *Neurobiol Dis.* 2006; 22(1):153-164.

Fortun J, Verrier JD, Go JC, Madorsky I, Dunn WA, Notterpek L. The formation of peripheral myelin protein 22 aggregates is hindered by the enhancement of autophagy and expression of cytoplasmic chaperones. *Neurobiol Dis.* 2007; 25(2): 252-265.

30 Gale K, Kerasidis H, Wrathall JR. Spinal cord contusion in the rat: behavioral analysis of functional neurologic impairment. *Exp Neurol.* 1985 Apr; 88(1):123-34.

- Galvez AS, Ulloa JA, Chiong M, Criollo A, Eisner V, Barros LF, Lavandero S. Aldose reductase induced by hyperosmotic stress mediates cardiomyocyte apoptosis: differential effects of sorbitol and mannitol. *J Biol Chem*. 2003; 278(40):38484-38494.
- Groyer G, Eychenne B, Girard C, Rajkowski K, Schumacher M, Cadepond F.
5 Expression and functional state of the corticosteroid receptors and 11 beta-hydroxysteroid dehydrogenase type 2 in Schwann cells. *Endocrinology*. 2006; 147(9):4339-4350.
- Howland DS, Liu J, She Y, Goad B, Maragakis NJ, Kim B, Erickson J, Kulik J, DeVito L, Psaltis G, DeGennaro LJ, Cleveland DW, Rothstein JD. Focal loss of the
10 glutamate transporter EAAT2 in a transgenic rat model of SOD1 mutant-mediated amyotrophic lateral sclerosis (ALS). *Proc Natl Acad Sci U S A*. 2002 Feb 5; 99(3):1604-9. Epub 2002 Jan 29.
- Kantamneni S, Corrêa SA, Hodgkinson GK, Meyer G, Vinh NN, Henley JM, Nishimune A. GISP: a novel brain-specific protein that promotes surface expression and
15 function of GABA(B) receptors. *J Neurochem*. 2007;100(4):1003-17.
- Khajavi M, Shiga K, Wiszniewski W, He F, Shaw CA, Yan J, Wensel TG, Snipes GJ, Lupski JR. Oral curcumin mitigates the clinical and neuropathologic phenotype of the Trembler-J mouse: a potential therapy for inherited neuropathy. *Am J Hum Genet*. 2007; 81(3): 438-453.
- 20 Kobsar I, Hasenpusch-Theil K, Wessig C, Müller HW, Martini R. Evidence for Macrophage-Mediated Myelin Disruption in an Animal Model for Charcot-Marie-Tooth Neuropathy Type 1A. *J. Neurosci Res* 2005; 81:857-864.
- Lange CA, Shen T et al. Phosphorylation of human progesterone receptors at serine-294 by mitogen-activated protein kinase signals their degradation by the 26S
25 proteasome. *PNAS USA*. 2000; 97: 1032-1037.
- Le-Niculescu H, Kurian SM, Yehyaw N, Dike C, Patel SD, Edenberg HJ, Tsuang MT, Salomon DR, Nurnberger JI Jr, Niculescu AB. Identifying blood biomarkers for mood disorders using convergent functional genomics. *Mol Psychiatry*. 2008 Feb 26. [Epub ahead of print].
- 30 Li WW, Le Goascogne C, Ramaugé M, Schumacher M, Pierre M, Courtin F. Induction of type 3 iodothyronine deiodinase by nerve injury in the rat peripheral nervous system. *Endocrinology*. 2001; 142(12):5190-5197.

- Lupski JR, Wise CA, Kuwano A, Pentao L, Parke JT, Glaze DG, Ledbetter DH, Greenberg F, Patel PI. Gene dosage is a mechanism for Charcot-Marie-Tooth disease type 1A. *Nat Genet.* 1992 ;1(1): 29-33.
- Matsumoto A, Okada Y, Nakamichi M, Nakamura M, Toyama Y, Sobue G, Nagai M, Aoki M, Itoyama Y, Okano H. Disease progression of human SOD1 (G93A) transgenic ALS model rats.. *J Neurosci Res.* 2006 Jan; 83(1):119-33.
- Mäurer M, Kobsar I, Berghoff M, Schmid CD, Carenini S, Martini R. Role of immune cells in animal models for inherited neuropathies: facts and visions. *J Anat.* 2002; 200(4): 405-414.
- 10 Melcangi RC, Cavarretta IT, Ballabio M, Leonelli E, Schenone A, Azcoitia I, Miguel Garcia-Segura L, Magnaghi V. Peripheral nerves: a target for the action of neuroactive steroids. *Brain Res Rev.* 2005; 48(2): 328-338.
- Mercier G, Turque N, Schumacher M. Rapid effects of triiodothyronine on immediate-early gene expression in Schwann cells. *Glia.* 2001; 35(2):81-89.
- 15 Meyer Zu Horste G., Nave K-A. Animal models of inherited neuropathies. *Curr. Opin. Neurol.* 2006; 19(5): 464-473.
- Meyer zu Horste G, Prukop T, Liebetanz D, Mobius W, Nave KA, Sereda MW. Antiprogestosterone therapy uncouples axonal loss from demyelination in a transgenic rat model of CMT1A neuropathy. *Ann Neurol.* 2007; 61 (1): 61-72.
- 20 Miller AL, Garza AS, Johnson BH, Thompson EB. Pathway interactions between MAPKs, mTOR, PKA, and the glucocorticoid receptor in lymphoid cells. *Cancer Cell Int.* 2007; 28:7:3
- Muja N, Blackman SC, Le Breton GC, DeVries GH. Identification and functional characterization of thromboxane A2 receptors in Schwann cells. *J Neurochem.* 2001; 25 78(3):446-456.
- Muller DL, Unterwald EM. In Vivo Regulation of Extracellular Signal-Regulated Protein Kinase (ERK) and Protein Kinase B (Akt) Phosphorylation by Acute and Chronic Morphine. *JPET* 2004; 310:774–782.
- Nambu H, Kubo E, Takamura Y, Tsuzuki S, Tamura M, Akagi Y. Attenuation of aldose reductase gene suppresses high-glucose-induced apoptosis and oxidative stress in rat lens epithelial cells. *Diabetes Res Clin Pract.* 2008; 82(1):18-24.
- 30

- Nave KA, Sereda MW, Ehrenreich H. Mechanisms of disease: inherited demyelinating neuropathies--from basic to clinical research. *Nat Clin Pract Neurol*. 2007; 3(8): 453-464.
- Niemann S., Sereda M.W., Rossner M., Stewart H., Suter U., Meinck H.M., Griffiths I.R., Nave K-A. The "CMT rat": peripheral neuropathy and dysmyelination caused by transgenic overexpression of PMP22. *Ann. N.- Y. Acad. Sci.* 1999; 883:254-261.
- Notterpek L, Shooter EM, Snipes GJ. Upregulation of the endosomal-lysosomal pathway in the trembler-J neuropathy. *J Neurosci*. 1997;17(11): 4190-4200.
- Obrietan K, van den Pol AN. GABAB receptor-mediated inhibition of GABAA receptor calcium elevations in developing hypothalamic neurons. *J Neurophysiol*. 1998; 79(3):1360-1370.
- Ogata T, Iijima S, Hoshikawa S, Miura T, Yamamoto S, Oda H, Nakamura K, Tanaka S Opposing extracellular signal-regulated kinase and Akt pathways control Schwann cell myelination. *J Neurosci*. 2004; 24(30):6724-6732.
- Ohsawa Y, Murakami T, Miyazaki Y, Shirabe T, Sunada Y. Peripheral myelin protein 22 is expressed in human central nervous system. *J Neurol Sci*. 2006; 247(1):11-15.
- Passage E, Norreel JC, Noack-Fraissignes P, Sanguedolce V, Pizant J, Thirion X, Robaglia-Schlupp A, Pellissier JF, Fontes M. Ascorbic acid treatment corrects the phenotype of a mouse model of Charcot-Marie-Tooth disease. *Nature Med*. 2004; 10(4): 396-401.
- Perea J, Robertson A, Tolmachova T, Muddle J, King RH, Ponsford S, Thomas PK, Huxley C. Induced myelination and demyelination in a conditional mouse model of Charcot-Marie-Tooth disease type 1A. *Hum Mol Genet*. 2001; 10(10):1007-1018.
- Rangaraju S, Madorsky I, Pileggi JG, Kamal A, Notterpek L. Pharmacological induction of the heat shock response improves myelination in a neuropathic model. *Neurobiology of Disease*. 2008; 32(105-115).
- Roa BB, Garcia CA, Suter U, Kulpa DA, Wise CA, Mueller J, Welcher AA, Snipes GJ, Shooter EM, Patel PI, Lupski JR. Charcot-Marie-Tooth disease type 1A. Association with a spontaneous point mutation in the PMP22 gene. *N Engl J Med*. 1993; 329(2): 96-101.

- Robaglia-Schlupp A, Pizant J, Norreel JC, Passage E, Saberan-Djoneidi D, Ansaldi JL, Vinay L, Figarella-Branger D, Levy N, Clarac F, Cau P, Pellissier JF, Fontes M. PMP22 overexpression causes dysmyelination in mice. *Brain* 2002; 125(Pt 10): 2213-2221.
- 5 Robert F, Guennoun R, Désarnaud F, Do-Thi A, Benmessahel Y, Baulieu EE, Schumacher M. Synthesis of progesterone in Schwann cells: regulation by sensory neurons. *Eur J Neurosci.* 2001; 13(5): 916-924.
- Roux KJ, Amici SA, Notterpek L. The temporospatial expression of peripheral myelin protein 22 at the developing blood-nerve and blood-brain barriers. *J Comp*
10 *Neurol.* 2004; 474(4):578-588.
- Sancho S, Young P, Suter U. Regulation of Schwann cell proliferation and apoptosis in PMP22-deficient mice and mouse models of Charcot-Marie-Tooth disease type 1A. *Brain* 2001; 124(Pt 11): 2177-2187.
- Schumacher M, Guennoun R, Mercier G, Désarnaud F, Lacor P, Bénavides J,
15 Ferzaz B, Robert F, Baulieu EE. Progesterone synthesis and myelin formation in peripheral nerves. *Brain Res Rev.* 2001; 37(1-3): 343-359.
- Sereda MW, Meyer zu Horste G, Suter U, et al. Therapeutic administration of progesterone antagonist in a model of Charcot-Marie-Tooth disease (CMT-1A). *Nat Med* 2003; 9: 1533-1537.
- 20 Sereda MW, Nave KA. Animal models of Charcot-Marie-Tooth disease type 1A (CMT1A). *Neuromol Med* 2006; 8: 205-215.
- Stirnweiss J, Valkova C, Ziesché E, Drube S, Liebmann C. Muscarinic M2 receptors mediate transactivation of EGF receptor through Fyn kinase and without matrix metalloproteases. *Cell Signal.* 2006; 18(8):1338-1349.
- 25 Suter U, Scherer SS. Disease mechanisms in inherited neuropathies. *Nat. Rev. Neurosci.* 2003; 4: 714-726.
- Suter U, Welcher AA, Ozcelik T, Snipes GJ, Kosaras B, Francke U, Billings-Gagliardi S, Sidman RL, Shooter EM. Trembler mouse carries a point mutation in a myelin gene. *Nature.* 1992; 356(6366): 241-244.
- 30 Thonhoff JR, Jordan PM, Karam JR, Bassett BL, Wu P. Identification of early disease progression in an ALS rat model. *Neurosci Lett.* 2007 Mar 30; 415(3):264-8. Epub 2007 Jan 14.

Thomas PK, Marques W Jr, Davis MB, Sweeney MG, King RH, Bradley JL, Muddle JR, Tyson J, Malcolm S, Harding AE. The phenotypic manifestations of chromosome 17p11.2 duplication. *Brain* 1997; 120 (Pt 3): 465-478.

- 5 Tobler AR, Liu N, Mueller L, Shooter EM. Differential aggregation of the Trembler and Trembler J mutants of peripheral myelin protein 22. *PNAS U S A.* 2002; 99(1):483-488.

Tu H, Rondard P, Xu C, Bertaso F, Cao F, Zhang X, Pin JP, Liu J. Dominant role of GABAB2 and Gbetagamma for GABAB receptor-mediated-ERK1/2/CREB pathway in cerebellar neurons. *Cell Signal.* 2007; 19(9):1996-2002.

Uht RM, Anderson CM, Webb P, Kushner PJ. Transcriptional activities of estrogen and glucocorticoid receptors are functionally integrated at the AP-1 response element. *Endocrinology.* 1997 Jul;138(7):2900-2908.

- 15 Ulzheimer JC, Peles E, Levinson SR, Martini R. Altered expression of ion channel isoforms at the node of Ranvier in P0-deficient myelin mutants. *Mol Cell Neurosci.* 2004; 25(1): 83-94.

Vallat JM, Sindou P, Preux PM, Tabaraud F, Milor AM, Couratier P, LeGuern E, Brice A. Ultrastructural PMP22 expression in inherited demyelinating neuropathies. *Ann Neurol.* 1996; 39(6): 813-817.

- 20 Walter IB. Nuclear triiodothyronine receptor expression is regulated by axon-Schwann cell contact. *Neuroreport.* 1993; 5(2):137-140.

Walter IB, Deruaz JP, de Tribolet N. Differential expression of triiodothyronine receptors in schwannoma and neurofibroma: role of Schwann cell-axon interaction. *Acta Neuropathol (Berl).* 1995; 90(2):142-149.

- 25 Welch WJ, Brown CR. Influence of molecular and chemical chaperones on protein folding. *Cell Stress Chaperones.* 1996;1(2):109-115.

Woodhams PL, MacDonald RE, Collins SD, Chessell IP, Day NC. Localisation and modulation of prostanoid receptors EP1 and EP4 in the rat chronic constriction injury model of neuropathic pain. *Eur J Pain.* 2007; 11(6):605-613.

Claims

1. A composition comprising baclofen, D-sorbitol and naltrexone, salts, enantiomers, or racemates thereof, for simultaneous, separate or sequential administration to a mammalian subject.

2. The composition of claim 1, further comprising a pharmaceutically suitable excipient or carrier.

3. The composition according to claim 1 or 2, wherein said compounds are formulated with a biomolecule, a micelle or liposome-forming lipids or oil in water emulsions, or pegylated or solid nanoparticles or microparticles for oral or parenteral or intrathecal administration.

4. The composition of any one of the preceding claims, which is a liquid formulation suitable for oral administration.

5. Use of the composition of any one of claims 1 to 4, in the manufacture of a medicament for the treatment of Charcot Marie Tooth ("CMT") disease, a toxic neuropathy or Amyotrophic Lateral Sclerosis ("ALS") in a mammalian subject.

6. The use of claim 5, for the treatment of CMT1A in said mammalian subject.

7. The use of claim 5 or 6, wherein the mammalian subject is a human subject.

8. Use of Baclofen, D-sorbitol and naltrexone, salts, enantiomers, or racemates thereof in the manufacture of a medicament for the treatment of CMT disease, a toxic neuropathy or ALS in a mammalian subject, said medicament being formulated for simultaneous, separate or sequential administration.