

TREATMENT OF CLUSTER HEADACHE USING ANTI-CGRP ANTIBODIES

PRIOR APPLICATION INFORMATION

[0001] This application claims priority to U.S. Provisional Application No. 63/237,639 filed August 27, 2021, and is herein incorporated by reference in its entirety.

SEQUENCE LISTING DISCLOSURE

[0002] The contents of the electronic sequence listing (1267 SEQ listing-2022-08-03.xml; Size: 771,633 bytes; and Date of Creation: August 3, 2022) is herein incorporated by reference in its entirety.

SEQUENCES NOT PERMITTED TO BE ENTERED IN ST.26 XML FILE DUE TO SEQUENCE LENGTH

[0003] Table A below lists sequences present in the priority application U.S. Provisional Application No. 63/237,639 (identified above, which is herein incorporated by reference in its entirety), but cannot be included in the 1267 SEQ listing-2022-08-03.xml file submitted herewith due to the length of the sequences.

TABLE A										
Item No.	Residues	Length	Molecule Type	Organism	Contains DNA and RNA fragments	Sequence Skipped in XML file	Feature Key	Feature Location	Qualifiers	Previously Presented In Priority Application As SEQ ID
1	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 8: "1267-US-PSP_seq_8"
							source	1..3	mol_type = protein	
2	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 18: "1267-US-PSP_seq_18"
							source	1..9	mol_type = other DNA organism = synthetic construct	
3	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 48: "1267-US-PSP_seq_48"

TABLE A										
Item No.	Residues	Length	Molecule Type	Organism	Contains DNA and RNA fragments	Sequence Skipped in XML file	Feature Key	Feature Location	Qualifiers	Previously Presented In Priority Application As SEQ ID
							source	1..3	mol_type = protein organism = synthetic construct	
4	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 58: "1267-US-PSP_seq_58"
							source	1..9	mol_type = other DNA organism = synthetic construct	
5	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 88: "1267-US-PSP_seq_88"
							source	1..3	mol_type = protein	
6	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 98: "1267-US-PSP_seq_98"
							source	1..9	mol_type = other DNA organism = synthetic construct	
7	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 128: "1267-US-PSP_seq_128"
							source	1..3	mol_type = protein organism = synthetic construct	
8	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 138: "1267-US-PSP_seq_138"
							source	1..9	mol_type = other DNA organism = synthetic construct	
9	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 168: "1267-US-PSP_seq_168"
							source	1..3	mol_type = protein organism = synthetic construct	

TABLE A										
Item No.	Residues	Length	Molecule Type	Organism	Contains DNA and RNA fragments	Sequence Skipped in XML file	Feature Key	Feature Location	Qualifiers	Previously Presented In Priority Application As SEQ ID
10	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 178: "1267-US-PSP_seq_178"
							source	1..9	mol_type = other DNA organism = synthetic construct	
11	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 208: "1267-US-PSP_seq_208"
							source	1..3	mol_type = protein organism = synthetic construct	
12	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 218: "1267-US-PSP_seq_218"
							source	1..9	mol_type = other DNA organism = synthetic construct	
13	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 248: "1267-US-PSP_seq_248"
							source	1..3	mol_type = protein organism = synthetic construct	
14	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 258: "1267-US-PSP_seq_258"
							source	1..9	mol_type = other DNA organism = synthetic construct	
15	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 288: "1267-US-PSP_seq_288"
							source	1..3	mol_type = protein organism = synthetic construct	

TABLE A										
Item No.	Residues	Length	Molecule Type	Organism	Contains DNA and RNA fragments	Sequence Skipped in XML file	Feature Key	Feature Location	Qualifiers	Previously Presented In Priority Application As SEQ ID
16	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 298: "1267-US-PSP_seq_298"
							source	1..9	mol_type = other DNA organism = synthetic construct	
17	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 328: "1267-US-PSP_seq_328"
							source	1..3	mol_type = protein organism = synthetic construct	
18	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 338: "1267-US-PSP_seq_338"
							source	1..9	mol_type = other DNA organism = synthetic construct	
19	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 368: "1267-US-PSP_seq_368"
							source	1..3	mol_type = protein organism = synthetic construct	
20	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 378: "1267-US-PSP_seq_378"
							source	1..9	mol_type = other DNA organism = synthetic construct	
21	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 408: "1267-US-PSP_seq_408"
							source	1..3	mol_type = protein organism = synthetic construct	

TABLE A										
Item No.	Residues	Length	Molecule Type	Organism	Contains DNA and RNA fragments	Sequence Skipped in XML file	Feature Key	Feature Location	Qualifiers	Previously Presented In Priority Application As SEQ ID
22	ggcgacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 418: "1267-US-PSP_seq_418"
							source	1..9	mol_type = other DNA organism = synthetic construct	
23	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 448: "1267-US-PSP_seq_448"
							source	1..3	mol_type = protein organism = synthetic construct	
24	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 458: "1267-US-PSP_seq_458"
							source	1..9	mol_type = other DNA organism = synthetic construct	
25	GDI	3	AA	synthetic construct	No	Yes	REGION	1..3	note = Engineered antibody sequence	Sequence 528: "1267-US-PSP_seq_528"
							source	1..3	mol_type = protein organism = synthetic construct	
26	ggggacatc	9	DNA	synthetic construct	No	Yes	misc_feature	1..9	note = Engineered antibody sequence	Sequence 538: "1267-US-PSP_seq_538"
							source	1..9	mol_type = other DNA organism = synthetic construct	

BACKGROUND

[0004] Field

[0005] This invention pertains to methods of treatment of headache disorders, such as Cluster headache, using antibodies and fragments thereof (including Fab fragments) that specifically bind to human Calcitonin Gene Related Peptide (hereinafter "CGRP"). The invention also pertains to immediate treatment of headache, e.g., cluster headache, using antibodies and fragments thereof (including Fab fragments) that specifically bind to human Calcitonin Gene Related Peptide (hereinafter "CGRP").

[0006] Description of Related Art

[0007] Calcitonin Gene Related Peptide (CGRP) is produced as a multifunctional neuropeptide of 37 amino acids in length. Two forms of CGRP, the CGRP-alpha and CGRP-beta forms, exist in humans and have similar activities. CGRP-alpha and CGRP-beta differ by three amino acids in humans, and are derived from different genes. CGRP is released from numerous tissues such as trigeminal nerves, which when activated release neuropeptides within the meninges, mediating neurogenic inflammation that is characterized by vasodilation, vessel leakage, and mast-cell degradation. Durham, P.L., *New Eng. J. Med.*, 350 (11):1073-75 (2004). Biological effects of CGRP are mediated via the CGRP receptor (CGRP-R), which consists of a seven-transmembrane component, in conjunction with receptor-associated membrane protein (RAMP). CGRP-R further requires the activity of the receptor component protein (RCP), which is essential for an efficient coupling to adenylate cyclase through G proteins and the production of cAMP. Doods, H., *Curr. Op. Invest. Drugs*, 2(9):1261-68 (2001).

[0008] Migraines are neurovascular disorder affecting approximately 10% of the adult population in the U.S., and are typically accompanied by intense headaches. CGRP is believed to play a prominent role in the development of migraines. In fact several companies, i.e., Amgen, Eli Lilly, Teva and Alder Biopharmaceuticals (recently acquired by Lundbeck A/S) have developed anti-CGRP and anti-CGRP-R antibodies for use in treating or preventing migraine headaches. The present assignee has previously filed patent applications related to anti-CGRP antibodies and uses thereof including published PCT Application WO/2012/162243 filed May 21, 2012 entitled "ANTI-CGRP COMPOSITIONS AND USE THEREOF", published PCT Application WO/2012/162257 filed May 21, 2012, entitled "USE OF ANTI-CGRP ANTIBODIES AND ANTIBODY FRAGMENTS TO PREVENT OR INHIBIT PHOTOPHOBIA OR LIGHT AVERSION IN SUBJECTS IN NEED THEREOF, ESPECIALLY MIGRAINE SUFFERERS" published PCT Application WO/2012/162253, filed May 21, 2012, entitled "USE OF ANTI-CGRP OR ANTI-CGRP-R ANTIBODIES OR ANTIBODY FRAGMENTS TO TREAT OR PREVENT CHRONIC AND ACUTE FORMS OF DIARRHEA" and published PCT Application WO/2015/003122, filed July 3, 2014, entitled "REGULATION OF GLUCOSE METABOLISM USING ANTI-CGRP ANTIBODIES" all of which applications are incorporated by reference in their entirety.

[0009] Cluster headache (such as chronic or episodic cluster headache) is a rare but disabling primary headache disorder characterized by attacks of intense unilateral headache, frequently

associated with autonomic symptoms such as lacrimation, conjunctival injection, and nasal congestion (International Headache Society International Classification of Headache Disorders third edition [HIS ICHD-3]). The diagnosis of Cluster headache is distinctly recognized and defined by the ICHD-3.

[0010] The social impact of cluster headache is considerable and is associated with considerable direct and indirect economic consequences. Cluster headache has a prevalence of 0.1% with a 2 to 6 times higher average incidence rate for males compared to females. However, the ratio might be lower due to misdiagnosis of cluster headache in females compared to males.

[0011] The lifetime prevalence of cluster headache, based on a meta-analysis, showed a mean prevalence of 124 per 100,000, where episodic form was 6 times more prevalent than the chronic form. There are significant unmet needs for just about every clinical aspect of cluster headache, particularly related to the severity of the disease, the diagnostic challenges and the available treatment options. Most patients experiencing cluster headache attacks rate their pain intensity as near to or at the worst pain imaginable (using a 10-cm Visual Analog Scale [VAS]).

[0012] The pharmacological armamentarium in cluster headache consists of acute / abortive therapies, transitional therapies and preventive treatments. First line of acute treatment is sumatriptan administered subcutaneously and inhalation of 100% oxygen. Transitional therapies are often used to relieve the patient until the preventive agents are adequately titrated, and consists of oral steroids or greater occipital nerve (GON) blocks. The currently available preventive pharmacological treatments are unspecific, insufficient, and hampered by side-effects. Preventive treatment aims to reduce attack frequency with verapamil being first-choice, but only 50 to 80% of cluster headache patients are responders; and its use is hampered by side-effects since many patients need high doses. Other preventive treatments are less attractive due to their side-effect profile, the scarcity of evidence and high cost. In the clinic, several types of treatment are combined in the effort to provide relief to the patients and improve the quality of life.

[0013] Increased plasma or serum levels of calcitonin gene-related peptide (CGRP) have been associated with painful syndromes such as migraine and cluster headache. Cluster headache patients have higher CGRP levels compared to migraine patients and healthy controls. As in migraine, CGRP levels are altered during attacks.

BRIEF SUMMARY

[0014] In one aspect, the present disclosure provides a method for treatment of cluster headache, either episodic or chronic, in a patient in the need of immediate relief of symptoms or for prevention of cluster headache in a patient in need of immediate preventative treatment, comprising intravenous administering to a patient in need 400 mg of an anti-CGRP antibody comprising the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID

NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively. According to one embodiment the antibody may be of a human IgG1 type.

[0015] In some aspects, said patient may exhibit at least one cluster headache symptom at the time of administration.

[0016] In some aspects, said patient may have head pain.

[0017] In some aspects, the pain may include one or more of the following symptoms migraine-like nausea and aura. Common signs and symptoms during a cluster headache attack include: pain in, behind or around one eye, one-sided pain, restlessness, tearing or excessive tearing, eye redness e.g. redness eye on the affected side of pain, stuffy or runny nose e.g. only on the on the affected side, forehead or facial sweating e.g. on the affected side, pale skin (pallor) or flushing, swelling around the eye e.g. on the affected side, drooping eyelid e.g. on the affected side

[0018] In some aspects, one or more of said signs or symptoms may be alleviated after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

[0019] In some aspects, said patient may no longer have a cluster headache after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

[0020] In some aspects, said anti-CGRP antibody may comprise the light chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12), respectively.

[0021] In some aspects, said anti-CGRP antibody may comprise the variable light chain polypeptide of SEQ ID NO: 222.

[0022] In some aspects, said anti-CGRP antibody may comprise the variable light chain polypeptide encoded by SEQ ID NO: 232.

[0023] In some aspects, said anti-CGRP antibody may comprise the variable heavy chain polypeptide of SEQ ID NO: 202.

[0024] In some aspects, said anti-CGRP antibody may comprise the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

[0025] In some aspects, said anti-CGRP antibody may comprise the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202.

[0026] In some aspects, said anti-CGRP antibody may comprise the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

[0027] In some aspects, said anti-CGRP antibody may comprise the light chain polypeptide of SEQ ID NO: 221.

[0028] In some aspects, said anti-CGRP antibody may comprise the light chain polypeptide encoded by SEQ ID NO: 231.

[0029] In some aspects, said anti-CGRP antibody may comprise the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

[0030] In some aspects, said anti-CGRP antibody may comprise the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

[0031] In some aspects, said anti-CGRP antibody may comprise the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

[0032] In some aspects, said anti-CGRP antibody may comprise the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

[0033] In some aspects, said intravenous administration may be infused over a period of approximately 30 min to 60 minutes.

[0034] In some aspects, the cluster headache symptoms may decline or may be abolished immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

[0035] In some aspects, said patient may be symptom free 2 hours post-completion of infusion.

[0036] In some aspects, said method may further comprise intravenously administering 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

[0037] In some aspects, said method may further comprise intravenously administering 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

[0038] In some aspects, said anti-CGRP antibody may be comprised in a formulation comprising or consisting of histidine (L-histidine), sorbitol, polysorbate 80, and water.

[0039] In some aspects, said formulation may comprise or may consist of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or

having amounts of each constituent within +/- 10% of said values, and having a pH of 5.8 or within +/- 10% of said value.

[0040] In some aspects, said formulation may comprise or may consist of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 5% of said values, and/or having a pH of 5.8 or within +/- 5% of said value.

[0041] In some aspects, said formulation may comprise or may consist of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 1% of said values, and/or having a pH of 5.8 or within +/- 1% of said value.

[0042] In some aspects, said formulation may comprise or may consist of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.5% of said values, and/or having a pH of 5.8 or within +/- 0.5% of said value.

[0043] In some aspects, said formulation may comprise or may consist of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.1% of said values, and/or having a pH of 5.8 or within +/- 0.1% of said value.

[0044] In some aspects, said L- Histidine in said formulation comprises a mixture of L-Histidine and L-Histidine monohydrate. Said 3.1 mg of histidine in said formulation may comprise a mixture of L-Histidine (1 mg) and L-Histidine monohydrate (2.8 mg), which in the final formulation sums up to 3.1 mg L-histidine free base.

[0045] In some aspects, said formulation may be comprised in a 100 mg/mL single-dose vial wherein each mL contains 100 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

[0046] In some aspects, said formulation may be comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

[0047] In some aspects, the patient may be receiving or has received additional migraine or headache medication.

[0048] In some aspects, the patient may receive additional migraine or headache medication prior, concurrent or after administration of the anti-CGRP antibody.

[0049] In some aspects, the patient may receive additional migraine or headache medication within a period of time before and after said anti-CGRP antibody administration, such as within 15

minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said anti-CGRP antibody administration.

[0050] In some aspects, said additional migraine medication may comprise an acute and/or a chronic migraine medication.

[0051] In some aspects, said additional migraine or headache medication may comprise a triptan, an analgesic such as non-opioid or opioid/narcotic, acetaminophen, an NSAID, a combination medication such as EXCEDRIN® or EXCEDRIN MIGRAINE®, antiemetic medication, an ergotamine, or an ergot derivative.

[0052] In some aspects, said additional migraine or headache medication may comprise non-opioid analgesic such as paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two drugs with analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol, butorphanol, morphine, codeine, hydrocodone, or any combination thereof), barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

[0053] In some aspects, said migraine or headache may be selected from the group comprising acute migraine or headache, migraines with or without aura, chronic migraine, episodic migraine, chronic/episodic migraine, hemiplagic migraines, cluster headaches, migrainous neuralgia, chronic headaches, tension headaches, general headaches, headaches due to an underlying structural problem in the head or neck, sinus headaches (such as for example associated with sinusitis), and allergy-induced headaches or migraines.

[0054] In some aspects, said anti-CGRP antibody may be expressed in or obtained by expression in *Pichia pastoris*.

[0055] In some aspects, said anti-CGRP antibody may be expressed in or obtained by expression in CHO cells.

[0056] In some aspects, said patient may be administered 400 mg of said anti-CGRP antibody every three months.

[0057] In some aspects, said method of treatment may result in immediate relief of cluster headache symptoms.

[0058] In some aspects, said method of treatment may result in immediate preventative treatment of cluster headache.

[0059] The present disclosure further provides methods of immediate treatment of cluster headache (such as chronic or episodic headache), comprising administering to a patient in need an

effective amount of at least one anti-CGRP antibody or antibody fragment or an anti-CGRP-R antibody or antibody fragment or one or more formulations comprising said antibody or antibody fragment as disclosed herein,

[0060] In some aspects, said antibody may be administered while said patient has a cluster headache.

[0061] In some aspects, said antibody administration may be initiated within 1-6 hours of the onset of said cluster headache. In some aspects, said cluster headache may comprise episodic cluster headache or chronic cluster headache. In some aspects, said anti-CGRP antibody or antibody fragment Ab6 or a Fab fragment thereof, having the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and the heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11); or having the light chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12), respectively. According to an embodiment the antibody may be of a human IgG1 type. In some aspects, said anti-CGRP antibody may comprise the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202. Said anti-CGRP antibody may comprise the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212. Said anti-CGRP antibody may comprise the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566. In some aspects, said anti-CGRP antibody may comprise the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567. In some aspects, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202, which polypeptides optionally are respectively linked to human light and heavy constant region polypeptides, e.g., human IgG1, IgG2, IgG3 or IgG4 constant regions, which constant regions optionally may be modified to alter glycosylation or proteolysis, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells. In some aspects, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the light chain of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells, wherein the constant regions thereof optionally may be modified to alter glycosylation or proteolysis or other effector functions. In some aspects, any of the aforementioned anti-CGRP antibodies or antibody fragments, preferably Ab6, may be optionally comprised in a formulation as disclosed herein, e.g.,

comprising histidine (L-histidine), sorbitol, polysorbate 80, such as, per 1 mL volume, about 100 mg anti-CGRP antibody, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, having a pH of about 5.8. In some aspects, said administered dosage of said antibody may be 400 mg. In some aspects, said dosage may be administered intravenously, e.g., in a saline solution such as 0.9% sodium chloride in a suitable volume, such as 100 mL.

[0062] In some aspect, said patient fulfill the ICHD-3 diagnosis criteria for cluster headache, such as chronic cluster headache or episodic cluster headache

[0063] In some aspects, said patient may exhibit at least five attacks according to the blow list A to C

A. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes (when untreated)

B. 1 or 2 of the following:

(1) at least one of the following symptoms or signs, ipsilateral to the headache:

i. conjunctival injection and/or lacrimation

ii. nasal congestion and/or rhinorrhoea

iii. eyelid oedema

iv. forehead and facial sweating

v. miosis and/or ptosis

(2) a sense of restlessness or agitation

C. Occurring with a frequency between one every other day and eight per day

[0064] In some aspects the cluster headache symptom is selected from the list consisting of or comprising severe or very severe unilateral orbital, supraorbital and/or temporal pain e.g. lasting 15–180 minutes (when untreated), and the one or more (such as 1, 2, 3, 4, 5, or all) of the following symptoms or signs, ipsilateral to the headache: conjunctival injection and/or lacrimation; nasal congestion and/or rhinorrhea; eyelid oedema; forehead and facial sweating; miosis and/or ptosis; a sense of restlessness or agitation.

[0065] In some aspects, said patient may exhibit the attacks according to the above without a remission period, or with remissions lasting less than 3 months, for at least 1 year.

[0066] The present disclosure provides methods of treating or preventing cluster headache, comprising administering to a patient in need an effective amount of an anti-CGRP antibody or anti-CGRP antibody fragment or one or more formulations comprising said anti-CGRP antibody or anti-CGRP antibody fragment as disclosed herein. In some aspects, said anti-CGRP antibody Ab6, having the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and the heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11); or having the light chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences

encoded by SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12), respectively. In some aspects, said anti-CGRP antibody may comprise the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202. Said anti-CGRP antibody may comprise the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212. Said anti-CGRP antibody may comprise the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566. In some aspects, said anti-CGRP antibody may comprise the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567. In some aspects, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202, which polypeptides optionally are respectively linked to human light and heavy constant region polypeptides, e.g., human IgG1, IgG2, IgG3 or IgG4 constant regions, which constant regions optionally may be modified to alter glycosylation or proteolysis, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells. Said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the light chain of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells, wherein the constant regions thereof optionally may be modified to alter glycosylation or proteolysis or other effector functions. Any of the aforementioned anti-CGRP antibodies or antibody fragments, preferably Ab6, may be optionally comprised in a formulation as disclosed herein, e.g., comprising histidine (L-histidine), sorbitol, polysorbate 80, such as, per 1 mL volume, about 100 mg anti-CGRP antibody, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, having a pH of about 5.8. The administered dosage of said antibody may be between about 100 mg and about 400 mg, such as 400 mg. The dosage may be administered by different means, e.g., intravenously, e.g., in a saline solution such as 0.9% sodium chloride in a suitable volume, such as 100 mL.

[0067] The present disclosure also provides methods of treating cluster headache, comprising intravenously administering to a patient in need thereof a first dosage comprising 400 mg of an anti-CGRP antibody, wherein said anti-CGRP antibody preferably comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or 566, wherein in the first 24 hours (such as within $\frac{1}{2}$ hour, 1 hour, 2 hour or 12 hours) after administration of said first dosage the patient exhibits relief in pain or other signs or symptoms related to cluster headache.

[0068] In another aspect, the disclosure provides methods of treating cluster headache, comprising intravenously administering to a patient in need thereof a first dosage comprising between about 100 mg and about 300 mg of an anti-CGRP antibody, wherein said anti-CGRP antibody

preferably comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or 566, wherein on the first day following the day of administration the patient exhibits relief in pain or symptoms related to cluster headache.

[0069] In some exemplary embodiments, the dosage, e.g., the first dosage, of said anti-CGRP antibody may be 400 mg.

[0070] In other exemplary embodiments, the method of treating cluster headache may comprise intravenously administering 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

[0071] The antibody may be provided or administered in a formulation as disclosed herein, e.g., comprising histidine (L-histidine), sorbitol, polysorbate 80, such as, per 1 mL volume, about 100 mg anti-CGRP antibody, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, having a pH of about 5.8.

[0072] In some embodiments, the patient to be treated may be a chronic cluster headache patient or episodic cluster headache patient

[0073] In some embodiments, said anti-CGRP antibody may be aglycosylated or if glycosylated only may contain only mannose residues.

[0074] In some embodiments, said anti-CGRP antibody may consist of the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566. Said anti-CGRP antibody may consist of the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

[0075] In some embodiments, said anti-human CGRP antibody or antibody fragment comprises the variable light chain of SEQ ID NO: 222 and/or the variable heavy chain of SEQ ID NO: 202. In some embodiments, said anti-human CGRP antibody or antibody fragment comprises the variable light chain encoded by SEQ ID NO: 232 and/or the variable heavy chain encoded by SEQ ID NO: 212.

[0076] In some embodiments, said anti-human CGRP antibody or antibody fragment comprises the light chain of SEQ ID NO: 221 and/or the heavy chain of SEQ ID NO: 201 or SEQ ID NO: 566. In some embodiments, said anti-human CGRP antibody or antibody fragment comprises the light chain encoded by SEQ ID NO: 231 and/or the heavy chain encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

[0077] In some embodiments, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the VL polypeptide of SEQ ID NO: 222 and the VH polypeptide of SEQ ID NO: 202, which polypeptides optionally are respectively linked to human light and heavy constant region polypeptides, e.g., human IgG1, IgG2, IgG3 or IgG4 constant regions, which constant regions optionally may be modified to alter glycosylation or proteolysis, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells.

[0078] In some embodiments, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the light chain of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells, wherein the constant regions thereof optionally may be modified to alter glycosylation or proteolysis or other effector functions.

[0079] In some embodiments, any of the aforementioned anti-CGRP antibodies or antibody fragments may be comprised in a formulation as disclosed herein, e.g., comprising histidine (L-histidine), sorbitol, polysorbate 80, such as, per 1 mL volume, about 100 mg anti-CGRP antibody, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, having a pH of about 5.8. The antibody or fragment may be administered by different means, e.g., intravenously, e.g., in a saline solution such as 0.9% sodium chloride in a suitable volume, such as 100 mL.

[0080] In other embodiments, about 400 mg of said anti-CGRP antibody or antibody fragment is administered, e.g., intravenously.

[0081] In exemplary embodiments, the anti-human CGRP antibody or antibody fragment is administered, e.g., intravenously at a frequency which is at most every 10-14 weeks, preferably every 11-13 weeks, more preferably every 3 months or every 12 weeks, wherein the antibody dosage is administered in a single formulation or divided into different formulations which are administered at a frequency of approximately every 10-14 weeks, preferably every 11-13 weeks, more preferably every 3 months or every 12 weeks. The phrase "the antibody dosage is administered in a single formulation or divided into different formulations" refers to the administration of the recited amount of antibody within a relatively short period of time, e.g., within a period of several hours, e.g., 1 to 8 hours, about one day, within about two days, or within about one week, which may be by the same or different routes (e.g., i.v., i.m., and/or s.c.), sites of administration.

[0082] In other exemplary embodiments, the anti-human CGRP antibody used in the aforementioned methods has an *in vivo* half-life of at least 10 days.

[0083] In other exemplary embodiments, the anti-human CGRP antibody has an *in vivo* half-life of at least 15 days.

[0084] In other exemplary embodiments, the anti-human CGRP antibody used in the aforementioned methods has an *in vivo* half-life of at least 20 days.

[0085] In other exemplary embodiments, the anti-human CGRP antibody used in the aforementioned methods has an *in vivo* half-life of at least 20-30 days.

[0086] In other exemplary embodiments, the anti-human CGRP antibody used in the aforementioned methods binds to human α - and β -CGRP.

[0087] In other exemplary embodiments, the administered anti-human CGRP antibody dosage results in sustained pharmacodynamic (PK) activity, within 5% of the maximal response (I_{max}) (as compared to lower antibody doses).

[0088] In other exemplary embodiments, the administered anti-human CGRP antibody dosage results in sustained pharmacodynamic (PK) activity which is maintained for at least 2-3 months after antibody administration, wherein PK analysis of the anti-human CGRP antibody is derived from plasma concentrations.

[0089] In other exemplary embodiments, the administered anti-human CGRP antibody dosage is 400 mg which is administered no more frequently than every 2 months.

[0090] The present invention is additionally directed to the use of specific antibodies and fragments thereof having binding specificity for CGRP, in particular antibodies having desired epitopic specificity, high affinity or avidity and/or functional properties. A preferred embodiment of the invention is directed to usage of chimeric or humanized antibodies and fragments thereof (including Fab fragments) capable of binding to CGRP and/or inhibiting the biological activities mediated by the binding of CGRP to the CGRP receptor ("CGRP-R") e.g., wherein such antibodies optionally are derived from recombinant cells engineered to express same, optionally yeast or mammalian cells, further optionally *Pichia pastoris* and CHO cells.

[0091] High affinity for a specific and useful epitope is particularly desirable for the avoidance of side effects, since higher affinity of the therapeutic antibody to the desirable epitope can lead to a reduction in dose or a reduction of frequency of administration of the antibody and thus the reduction of side effects. The stronger and more robust the effect triggered by the high affinity binding of the therapeutic antibody is, the less frequent the antibody needs to be administered, and/or can be administered at a reduced dose. As frequent administration and high doses carry a higher risk of eliciting side effects, an antibody that provides robust and prolonged effects by high affinity binding to a specific epitope is highly advantageous. The claimed invention provides such antibodies.

[0092] The data in the application as filed demonstrates that the listed antibodies effectively bind to CGRP, prevent binding of the peptide hormone to its cellular receptor and thereby inhibit downstream cellular signaling. Ab6 (also known as ALD403 or Eptinezumab) is disclosed in the application as filed and is exemplified as a useful antibody, showing the desired epitopic specificity, high affinity and functional properties which enable it for the treatment of CGRP related diseases. Ab6 is further characterized by associating quickly with its ligand and releases the ligand only extremely slowly. As presented at the American Headache Society 61st Annual Scientific Meeting July 2019 Ab6 (also named ALD403 or Eptinezumab), the Eptinezumab:CGRP ligand binding interface has been characterized structurally, and the CDRs forms extensive contact to the ligand CGRP. Analysis of the atomic resolution 3-dimensional complex structure reveals that the C-terminal end of CGRP binds in a deep, narrow pocket formed by the heavy and light chains demonstrating that the binding is via a deep positive charged pocket and a large hydrophobic surface to bind CGRP, with both surfaces revealing strong shape and charge complementarity, which contribute to the binding specificity and affinity. The pocket is uniquely formed in the interface between the Ab6 VH and VL domains and intimately accepts the C-terminal part of the CGRP peptide in the activated form with

the terminal acylation of Phe37. The conclusions state that the unique attributes of the Fab region of the Eptinezumab molecular structure demonstrate the specificity and strength of binding to α -CGRP. Clinical trials have confirmed Ab6's particular suitability to treat CGRP related diseases, demonstrating prolonged efficacy over six months for the treatment of migraines (Press Release: "Alder Presents Positive Clinical Data for ALD403 at the 17th Congress of the International Headache Society", May 15, 2015) and Annex 4 (Phase 2 Human Clinical Trial Results: "Randomized, double-blind, placebo-controlled trial of ALD403, an anti-CGRP antibody in the prevention of frequent episodic migraine" ("Trial Results")). Further results from clinical trials with Ab6 have demonstrated that following administration of a single intravenous dose of Ab6(ALD403) (1000 mg) to individuals with a diagnosis of migraine, patients reported a significant reduction in the number of migraine days. In particular, over 12 weeks following treatment, 61% of patients receiving ALD403 reported a 50% reduction in migraine days (as compared to 33% of patients receiving placebo), 33% of patients receiving ALD403 reported a 75% reduction in migraine days (as compared to 9% of patients receiving placebo), and, strikingly, 16% of patients receiving ALD403 reported a 100% reduction in migraine days (as compared to 0% of patients receiving placebo). The same trend was observed after 24 weeks following treatment with ALD403. In particular, 53% of patients receiving ALD403 reported a 50% reduction in migraine days (as compared to 28% of patients receiving placebo), 26% of patients receiving ALD403 reported a 75% in migraine (as compared to 7% of patients receiving placebo), and 11% of patients receiving ALD403 reported a 100% reduction in migraine days (as compared to 0% of patients receiving placebo).

[0093] In another preferred embodiment of the invention, full length antibodies and Fab fragments thereof are contemplated that inhibit the CGRP-alpha-, CGRP-beta-, and rat CGRP-driven production of cAMP. In a further preferred embodiment of the invention, full length and Fab fragments thereof are contemplated that reduce vasodilation in a recipient following administration.

[0094] The invention also contemplates usage of conjugates of anti-CGRP antibodies and binding fragments thereof conjugated to one or more functional or detectable moieties. The invention also contemplates usage of chimeric or humanized anti-CGRP or anti-CGRP/CGRP-R complex antibodies and binding fragments thereof. In one embodiment, binding fragments include, but are not limited to, Fab, Fab', F(ab')₂, Fv, scFv fragments, SMIPs (small molecule immunopharmaceuticals), camelbodies, nanobodies, and IgNAR.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

[0095] **FIGs. 1A-1F** provide the polypeptide sequences of the full-length heavy chain for antibodies Ab1-Ab14 with their framework regions (FR), complementarity determining regions (CDRs), and constant region sequences delimited.

[0096] **FIGs. 2A-2D** provide the polypeptide sequences of the full-length light chain for antibodies Ab1-Ab14 with their framework regions (FR), complementarity determining regions (CDRs), and constant region sequences delimited.

[0097] **FIGs. 3A-3P** provide exemplary polynucleotide sequences encoding the full-length heavy chain for antibodies Ab1-Ab14 with their framework regions (FR), complementarity determining regions (CDRs), and constant region coding sequences delimited.

[0098] **FIGs. 4A-4I** provide exemplary polynucleotide sequences encoding the full-length light chain for antibodies Ab1-Ab14 with their framework regions (FR), complementarity determining regions (CDRs), and constant region coding sequences delimited.

[0099] **FIG. 5** provides the polypeptide sequence coordinates within the full-length heavy chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the variable region and complementarity determining regions (CDRs), and the SEQ ID NO of each individual feature.

[0100] **FIG. 6** provides the polypeptide sequence coordinates within the full-length heavy chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the framework regions (FRs) and constant region, and the SEQ ID NO of each individual feature.

[0101] **FIG. 7** provides the polypeptide sequence coordinates within the full-length light chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the variable region and complementarity determining regions (CDRs), and the SEQ ID NO of each individual feature.

[0102] **FIG. 8** provides the polypeptide sequence coordinates within the full-length light chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the framework regions (FRs) and constant region, and the SEQ ID NO of each individual feature.

[0103] **FIG. 9** provides the polynucleotide sequence coordinates within the exemplary polynucleotide sequences encoding the full-length heavy chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the variable region and complementarity determining regions (CDRs), and the SEQ ID NO of each individual feature.

[0104] **FIG. 10** provides the polynucleotide sequence coordinates within the exemplary polynucleotide sequences encoding the full-length heavy chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the framework regions (FRs) and constant region, and the SEQ ID NO of each individual feature.

[0105] **FIG. 11** provides the polynucleotide sequence coordinates within the exemplary polynucleotide sequences encoding the full-length light chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the variable region and complementarity determining regions (CDRs), and the SEQ ID NO of each individual feature.

[0106] **FIG. 12** provides the polynucleotide sequence coordinates within the exemplary polynucleotide sequences encoding the full-length light chain polypeptide sequences of antibodies Ab1-Ab14 of sequence features including the framework regions (FRs) and constant region, and the SEQ ID NO of each individual feature.

[0107] **FIG. 13** shows the number of subjects in a human clinical trial described in Example 2 who were either treated with Ab6 (treatment group) or placebo groups who showed a 50, 75 or 100% reduction in migraines at each monitoring point throughout the period. The right bar in each group corresponds to patients receiving 1000 mg Ab6 and the left bar in each group corresponds to matched placebo controls. In each response rate group the patients receiving Ab6 had a significantly greater response rate than placebo-treated controls, with p values of 0.0155, 0.0034, and 0.0006 in each respective group as indicated. The administered antibody was produced in *P. pastoris* and consisted of the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201.

[0108] **FIG. 14** shows the median ($\pm$ QR) % change from baseline in the number of migraine days per month in the placebo and Ab6 –treated group over the 12 weeks post-treatment. (p=0.0078). The upper (red) line and lower (blue) line show results for placebo-treated controls and patients administered 1000 mg Ab6, respectively.

[0109] **FIG. 15** shows the median ($\pm$ QR) % change from baseline in the number of migraine episodes per month in the placebo and Ab6 –treated group over the 12 weeks post-treatment. The upper (red) line and lower (blue) line show results for placebo-treated controls and patients administered 1000 mg Ab6, respectively.

[0110] **FIG. 16** shows the median ($\pm$ QR) % change from baseline in the number of migraine hours per month in the placebo and Ab6 –treated group over the 12 weeks post-treatment. The upper (red) line and lower (blue) line show results for placebo-treated controls and patients administered 1000 mg Ab6, respectively.

[0111] **FIG. 17** summarizes the screening of patients, allocation into the treatment and control groups, and loss of patients through follow-up.

[0112] **FIG. 18** compares the HIT-6 responder analysis for the Ab6-treated and placebo groups at baseline, week 4 after treatment, week 8 after treatment and week 12 after treatment.

[0113] **FIG. 19** shows the percentage of patients for whom the HIT-6 analysis indicated that the effect of headaches was only “some” or “little/none” at baseline and after Ab6 administration. At baseline most patients had either “substantial” or “severe” impact from migraines. At each subsequent time point, a significantly greater percentage of patients administered 1000 mg Ab6 had only “some” or “little/none” HIT-6 impact (left bar in each group, colored blue) as compared to placebo controls (right bar in each group, colored red).

[0114] **FIG. 20** contains the pharmacokinetic (PK) profile for Ab6 administered intravenously at a single dosage of 1000 mg.

[0115] **FIG. 21** contains plasma-free pharmacokinetic (PK) parameters N (number of patients), mean, and standard deviation (SD) for a single 1000 mg intravenous dosage of Ab6. The parameters shown in the table and the units are $C_{\max}$ ($\mu\text{g/mL}$), $\text{AUC}_{0-\infty}$ ($\text{mg}\cdot\text{hr/mL}$), half-life (days), V_z (L) and CL (mL/hr).

[0116] **FIG. 22** shows the change (mean $\pm$ SEM) change from baseline in migraine days per month for Ab6 (1000 mg i.v.) versus placebo as a single dose for the study described in Example 2.

[0117] **FIG. 23** shows the average migraine days ($\pm$ SD) over time for the full analysis population for the study described in Example 2. Normalization was applied to visit intervals where eDiaries were completed for 21-27 days by multiplying the observed frequency by the inverse of the completion rate.

[0118] **FIG. 24** shows the distribution of migraine days actual and change for the Ab6 treatment group during weeks 1-4 for the study described in Example 2.

[0119] **FIG. 25** shows the distribution of migraine days actual and change for the placebo group during weeks 1-4 for the study described in Example 2.

[0120] **FIG. 26** shows the distribution of migraine days actual and change for the Ab6 treatment group during weeks 5-8 for the study described in Example 2.

[0121] **FIG. 27** shows the distribution of migraine days actual and change for the placebo group during weeks 5-8 for the study described in Example 2.

[0122] **FIG. 28** shows the distribution of migraine days actual and change for the Ab6 treatment group during weeks 9-12 for the study described in Example 2.

[0123] **FIG. 29** shows the distribution of migraine days actual and change for the placebo group during weeks 9-12 for the study described in Example 2.

[0124] **FIG. 30** shows the 50% responder rate for the Ab6 and placebo treatment groups for the study described in Example 2. Subjects with $\geq 50\%$ reduction in migraine frequency were considered to be a 50% responder. Normalization was applied to visit intervals where eDiary was completed for 21-27 days by multiplying the observed frequency by the inverse of the completion rate.

[0125] **FIG. 31** shows the 75% responder rate for the Ab6 and placebo treatment groups for the study described in Example 2. Subjects with $\geq 75\%$ reduction in migraine frequency were considered to be a 75% responder. Normalization was applied as described with **FIG. 30**.

[0126] **FIG. 32** shows the 100% responder rate for the Ab6 and placebo treatment group for the study described in Example 2. Subjects with 100% reduction in migraine frequency were considered to be a 100% responder. Normalization was applied as described with **FIG. 30**.

[0127] **FIG. 33** shows the mean migraine severity over time for the full analysis population for the study described in Example 2. On the scale used, a mean migraine score of 3 represents "moderate pain."

[0128] **FIG. 34** summarizes the change from baseline in measured attributes for the placebo and treatment groups in the study described in Example 2.

[0129] **FIG. 35** shows the percentages of patients with migraine in the 300 mg, 100 mg, and placebo treatment groups at days 1, 7, 14, 21, and 28 in the clinical trial described in Example 3. The uppermost line shows results for placebo, the lowest line shows results for the 300 mg dosage, and the middle line shows results for the 100 mg dosage.

[0130] **FIG. 36** show the percentage of patients in the 300 mg and 100 mg treatment groups achieving a 50% reduction in migraine days in month 1, over months 1-3 (after the 1st infusion), and over months 4-5 (after the 2nd infusion) in the clinical trial described in Example 3. In each graph, the data bars, from left to right, show results for the 100 mg, 300 mg, and placebo groups. Statistical significance is as shown. ++ indicates a statistically significant difference from placebo; + indicates a statistically significant difference from placebo (unadjusted); and § indicates a statistically significant difference from placebo (post hoc).

[0131] **FIG. 37** show the percentage of patients in the 300 mg and 100 mg treatment groups achieving a 75% reduction in migraine days in month 1, over months 1-3 (after the 1st infusion), and over months 4-5 (after the 2nd infusion) in the clinical trial described in Example 3. Data order and statistical significance labels are as indicated with **FIG. 36**.

[0132] **FIG. 38** show the percentage of patients in the 300 mg and 100 mg treatment groups achieving a 100% reduction in migraine days in month 1, over months 1-3 (after the 1st infusion), and over months 4-5 (after the 2nd infusion) in the clinical trial described in Example 3. Data order and statistical significance labels are as indicated with **FIG. 36**.

[0133] **FIG. 39** summarizes the characteristics of patients in each treatment group in the clinical trial described in Example 3. * According to the American Academy of Neurology/American Headache Society guidelines for migraine preventative treatment (medications identified by clinical review of coded medical data); SD, standard deviation; BMI, body mass index.

[0134] **FIG. 40.** Difference from placebo in change from baseline in mean migraine days (MMD) over months 1-3 by baseline subgroup for a human clinical trial of chronic migraine patients. In the graph, the data point refers to the mean value and the line shows the 95% confidence interval (CI) of the change from placebo for the 100 mg (upper line) or 300 mg (lower line) treatment group, for each subgroup as labeled at the far left.

[0135] **FIG. 41.** Difference from placebo in change from baseline in mean migraine days (MMD) over months 1-3 by baseline subgroup for a human clinical trial of episodic migraine patients. The graph is labeled as in **FIG. 40**.

[0136] **FIG. 42.** Change from baseline in mean migraine days (MMDs) across 2 dose intervals in chronic migraine patients with at least 1 day of acute medication use per month at baseline. Triangle: placebo (n=366). Circle: 100 mg Ab6 per dose (n=356). Square: 300 mg Ab6 per dose (n=350).

[0137] **FIG. 43.** Mean days with acute medication use in chronic migraine patients with at least one day per month of acute medication use at baseline. Triangle: placebo (n=366). Circle: 100 mg Ab6 per dose (n=356). Square: 300 mg Ab6 per dose (n=350).

[0138] **FIG. 44.** Change from baseline in acute medication use by subgroups of chronic migraine patients with differing baseline days of acute medication use. Solid lines: patients with 10 or more days of acute medication use per month at baseline. Dashed lines: patients with at least 1 and less

than 10 days of acute medication use per month at baseline. Triangle: placebo. Circle: 100 mg Ab6 per dose. Square: 300 mg Ab6 per dose.

[0139] **FIG. 45.** Summary of Acute Medication Days by Subgroups of Chronic Migraine Patients with Baseline Acute Medication Use.

[0140] **FIG. 46.** Change from baseline in mean migraine days (MMDs) across 2 dose intervals in episodic migraine patients with at least 1 day of acute medication use per month at baseline. Triangle: placebo (n=222). Circle: 100 mg Ab6 per dose (n=221). Square: 300 mg Ab6 per dose (n=222).

[0141] **FIG. 47.** Mean days with acute medication use in episodic migraine patients with at least one day per month of acute medication use at baseline. Triangle: placebo (n=222). Circle: 100 mg Ab6 per dose (n=221). Square: 300 mg Ab6 per dose (n=222).

[0142] **FIG. 48.** Change from baseline in acute medication use by subgroups of episodic migraine patients with differing baseline days of acute medication use. Solid lines: patients with 10 or more days of acute medication use per month at baseline. Dashed lines: patients with at least 1 and less than 10 days of acute medication use per month at baseline. Triangle: placebo. Circle: 100 mg Ab6 per dose. Square: 300 mg Ab6 per dose.

[0143] **FIG. 49.** Summary of Acute Medication Days by Subgroups of Episodic Migraine Patients with Baseline Acute Medication Use.

[0144] **FIG. 50.** Inclusion of Day -1 in the Migraine Data. Day 0 is defined as the day of the infusion. Thus, the data on Day 0 are indicative of the treatment effect post-infusion.

DETAILED DESCRIPTION

[0145] Use of anti-CGRP antibodies for treatment of cluster headache is described herein. Additionally, anti-CGRP antibodies are believed to be effective for treatment of chronic and episodic cluster headache. The treatment was shown to have a very rapid onset of efficacy, with relief on the first day following administration, and the effect on symptoms and pain even within the first ½ hour (such as within the first hour, 2 hours, 3 hours or e.g. 12 hours).

[0146] Definitions

[0147] It is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, animal species or genera, and reagents described, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments, only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. As used herein the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the protein" includes reference to one or more proteins and equivalents thereof known to those skilled in the art, and so forth. All technical and

scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs unless clearly indicated otherwise.

[0148] As used herein, the term "cluster headache" and "chronic" or "episodic" cluster headache refers to a cluster headache that meets the criteria for that condition specified in ICHD-3 (Headache Classification Committee of the International Headache Society (IHS), The International Classification of Headache Disorders, 3rd edition, Cephalalgia. 2018 Jan;38(1):1-211). Cluster period generally lasts for several weeks to months. The starting date and the duration of each cluster period might be consistent from period to period. For example, cluster periods can occur seasonally, such as every spring or every fall. In episodic cluster headaches, the headaches occur for one week to e.g 12 weeks, separated by a pain-free remission period that can last as long e.g as 12 months before another cluster period starts. Chronic cluster periods might continue for more than a year, or pain-free periods might last less than one month.

[0149] A cluster headache strikes quickly, usually without warning, although a migraine-like nausea and aura may be early signs. Common signs and symptoms during a headache include:

- Excruciating pain that is generally situated in, behind or around one eye, but may radiate to other areas of your face, head and neck
- One-sided pain
- Restlessness
- Excessive tearing
- Redness of the eye on the affected side
- Stuffy or runny nose on the affected side
- Forehead or facial sweating on the affected side
- Pale skin (pallor) or flushing face
- Swelling around the eye on the affected side
- Drooping eyelid on the affected side

[0150] During a cluster period, headaches usually occur every day, sometimes several times a day. A single attack can last from 15 minutes to three hours and the attacks often occur at the same time each day. Most attacks occur at night, usually one to two hours after you go to bed

[0151] A subject experiencing cluster headache will experience attacks of severe, strictly unilateral pain which is orbital, supraorbital, temporal or in any combination of these sites, lasting 15-180 minutes and occurring from once every other day to eight times a day. The pain is associated with ipsilateral conjunctival injection, lacrimation, nasal congestion, rhinorrhoea, forehead and facial sweating, miosis, ptosis and/or eyelid oedema, and/or with restlessness or agitation. The diagnosis criteria include a subject which exhibit at least five attacks according to the below list A to C

A. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes (when untreated)

B. 1 or 2 of the following

(1) at least one of the following symptoms or signs, ipsilateral to the headache:

- i. conjunctival injection and/or lacrimation
- ii. nasal congestion and/or rhinorrhoea
- iii. eyelid oedema
- iv. forehead and facial sweating
- v. miosis and/or ptosis

(2) a sense of restlessness or agitation

C. Occurring with a frequency between one every other day and eight per day

[0152] In case of chronic cluster headache the subject may exhibit the attacks according to the above without a remission period, or with remissions lasting less than 3 months, for at least 1 year. In case of episodic cluster headache the subject may exhibit the attacks according to the above occurring in periods lasting from 7 days to one year, separated by pain-free periods lasting at least 3 months.

[0153] The pain usually ends as suddenly as it began, with rapidly decreasing intensity. After attacks, most people are pain-free but exhausted.

[0154] As used herein, the term "reduction in reduction in weekly attacks " refers to a reduction (e.g., a stated percentage reduction, such as 50%) in the likelihood of a patient having attacks of intense unilateral headache and/or a reduction in the likelihood of patients rating their attacks pain intensity as near to or at the worst pain imaginable (using a 10-cm Visual Analog Scale [VAS]) in the stated period, such as the 18 hour, 20 hour, 24 hour, 28 hour, or 30 hour period, preferably the 24 hour period, after a first dosage of an antibody, or on the first day following the day of antibody administration (i.e., on the first full day following the day on which the antibody administration is completed).

[0155] As used herein, the term "diagnosed with cluster headache" refers to a patient meeting the clinical criteria for cluster headache, such as the clinical criteria for chronic cluster headache or episodic cluster headache, whether or not a formal diagnosis of that patient was performed.

[0156] As used herein, the term "intravenously administering" refers to a mode of administration wherein a substance, e.g., an antibody, is introduced directly into the circulation of that patient, most typically into the venous circulation. The substance may be introduced in a carrier fluid, such as an aqueous solution, e.g., normal saline. The substance may be administered in a single formulation or in multiple formulations, as long as the administration is completed over a short period of time (e.g., within 1 day, preferably within 12 hours, more preferably within 6 hours, and most preferably within 1-2 hours).

[0157] As used herein, the term "immediate relief" is intended to mean a relief in symptoms in a patient, e.g., cluster headache signs or symptoms associated with an chronic/episodic cluster headache, wherein said relief of signs or symptoms is experienced rapidly or immediately after anti-CGRP antibody treatment, e.g., relief of one or more signs or symptoms is experienced by the patient

within a short time period post-infusion with Ab6, such as within minutes or a few hours, such as within 10 minutes, 20 minutes, 30 minutes, 60 minutes, 1 hour, 2 hours or 6 hours, up to *e.g.* a day.

[0158] As used herein, the term “immediate preventative treatment” is intended to mean prevention of cluster headache symptoms in a patient, *e.g.*, prevention of cluster headache associated with chronic/episodic cluster headache. In this context, “immediate preventative treatment” refers to the treatment of a subject or a prophylactic treatment of a subject who is at risk of developing cluster headache, resulting in a decrease in the probability that the subject will develop cluster headache. Typically the prevention of symptoms is experienced rapidly or immediately after anti-CGRP antibody treatment, *e.g.*, prevention of one or more symptoms is experienced by the patient within a short time period post-infusion with Ab6, such as within minutes or a few hours, such as within 10 minutes, 20 minutes, 30 minutes, 60 minutes, 1 hour, 2 hours or 6 hours, up to *e.g.* a day.

[0159] As used herein, the terms “4-point scale” or “4 point pain scale” or “VRS” or “VRS-4” refer to the 4-point verbal rating scale (VRS) used to measure pain (VRS-4) (*see* “The International Classification of Headache Disorders, 3rd edition”, Cephalalgia, 2018, Vol. 38(1) 1–211, at pg. 210 (“intensity of pain”)). In the VRS the patient is asked to rate the pain verbally on a 4 point scale (between 0 and 3), with 3 being severe, 2 being moderate, 1 being mild, and 0 being no pain. It may also be scored on a verbal rating scale expressed in terms of its functional consequence: 0, no pain; 1, mild pain, does not interfere with usual activities; 2, moderate pain, inhibits but does not wholly prevent usual activities; 3, severe pain, prevents all activities.

[0160] *Calcitonin Gene Related Peptide (CGRP)*: As used herein, CGRP encompasses not only the following *Homo sapiens* CGRP-alpha and *Homo sapiens* CGRP-beta amino acid sequences available from American Peptides (Sunnyvale CA) and Bachem (Torrance, CA):

[0161] CGRP-alpha: ACDTATCVTHRLAGLLSRSGGVVKNNFVPTNVGSKAF-NH₂ (SEQ ID NO: 561), wherein the terminal phenylalanine is amidated;

[0162] CGRP-beta: ACNTATCVTHRLAGLLSRSGGMVKSNFVPTNVGSKAF-NH₂ (SEQ ID NO: 562), wherein the terminal phenylalanine is amidated; but also any membrane-bound forms of these CGRP amino acid sequences, as well as mutants (mutiens), splice variants, isoforms, orthologs, homologues and variants of this sequence.

[0163] *Expression Vector*: These DNA vectors contain elements that facilitate manipulation for the expression of a foreign protein within the target host cell, *e.g.*, a yeast or mammalian cell such as *Pichia pastoris* or CHO cells. Conveniently, manipulation of sequences and production of DNA for transformation is first performed in a bacterial host, *e.g.* *E. coli*, and usually vectors will include sequences to facilitate such manipulations, including a bacterial origin of replication and appropriate bacterial selection marker. Selection markers encode proteins necessary for the survival or growth of transformed host cells grown in a selective culture medium. Host cells not transformed with the vector containing the selection gene will not survive in the culture medium. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, (b) complement auxotrophic

deficiencies, or (c) supply critical nutrients not available from complex media. Exemplary vectors and methods for transformation of yeast are described, for example, in Burke, D., Dawson, D., & Stearns, T. (2000). *Methods in yeast genetics: a Cold Spring Harbor Laboratory course manual*. Plainview, N.Y.: Cold Spring Harbor Laboratory Press.

[0164] Expression vectors for use in yeast or mammalian cells will generally further include yeast or mammalian specific sequences, including a selectable auxotrophic or drug marker for identifying transformed yeast strains or transformed mammalian cells. A drug marker may further be used to amplify copy number of the vector in the host cell.

[0165] The polypeptide coding sequence of interest is operably linked to transcriptional and translational regulatory sequences that provide for expression of the polypeptide in host cells, e.g., *Pichia pastoris* or CHO cells. These vector components may include, but are not limited to, one or more of the following: an enhancer element, a promoter, and a transcription termination sequence. Sequences for the secretion of the polypeptide may also be included, e.g. a signal sequence, and the like. A yeast or mammalian origin of replication is optional, as expression vectors are often integrated into the host cell genome. In one embodiment of the invention, the polypeptide of interest is operably linked, or fused, to sequences providing for optimized secretion of the polypeptide from yeast diploid cells.

[0166] Nucleic acids are "operably linked" when placed into a functional relationship with another nucleic acid sequence. For example, DNA for a signal sequence is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites or alternatively via a PCR/recombination method familiar to those skilled in the art (Gateway[®] Technology; Invitrogen, Carlsbad California). If such sites do not exist, the synthetic oligonucleotide adapters or linkers are used in accordance with conventional practice.

[0167] Promoters are untranslated sequences located upstream (5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of particular nucleic acid sequences to which they are operably linked. Such promoters fall into several classes: inducible, constitutive, and repressible promoters (that increase levels of transcription in response to absence of a repressor). Inducible promoters may initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, e.g., the presence or absence of a nutrient or a change in temperature.

[0168] The promoter fragment may also serve as the site for homologous recombination and integration of the expression vector into the same site in the host genome; alternatively a selectable marker is used as the site for homologous recombination. Examples of suitable promoters from

Pichia include the AOX1 and promoter (Cregg *et al.* (1989) Mol. Cell. Biol. 9:1316-1323); ICL1 promoter (Menendez *et al.* (2003) Yeast 20(13):1097-108); glyceraldehyde-3-phosphate dehydrogenase promoter (GAP) (Waterham *et al.* (1997) Gene 186(1):37-44); and FLD1 promoter (Shen *et al.* (1998) Gene 216(1):93-102). The *GAP* promoter is a strong constitutive promoter and the AOX and FLD1 promoters are inducible.

[0169] Other yeast promoters include ADH1, alcohol dehydrogenase II, GAL4, PHO3, PHO5, Pyk, and chimeric promoters derived therefrom. Additionally, non-yeast promoters may be used in the invention such as mammalian, insect, plant, reptile, amphibian, viral, and avian promoters. Most typically the promoter will comprise a mammalian promoter (potentially endogenous to the expressed genes) or will comprise a yeast or viral promoter that provides for efficient transcription in yeast systems.

[0170] Examples of mammalian promoters include cytomegalovirus (CMV) derived promoters, chicken 3-actin (CBM) derived promoters, adenomatous polyposis coli (APC) derived promoters, leucine-rich repeat containing G protein-coupled receptor 5 (LGR5) promoters, CAG promoter, Beta actin promoter, elongation factor-1 (EF1) promoter, early growth response 1 (EGR-1) promoter, eukaryotic initiation factor 4A (EIF4A1) promoter, simian virus 40 (SV40) early promoter, mouse mammary tumor virus (MMTV), human immunodeficiency virus (HIV) long terminal repeat (LTR) promoter, MoMuLV promoter, an avian leukemia virus promoter, an Epstein-Barr virus immediate early promoter, a Rous sarcoma virus promoter, as well as human gene promoters such as, but not limited to, the actin promoter, the myosin promoter, the hemoglobin promoter, and the creatine kinase promoter, among others. Combinations of two or more of the foregoing promoters may also be used. Further, inducible promoters may be used. The use of an inducible promoter provides a molecular switch capable of turning on expression of the polynucleotide sequence which it is operatively linked when such expression is desired, or turning off the expression when expression is not desired. Examples of inducible promoters include, but are not limited to a metallothionine promoter, a glucocorticoid promoter, a progesterone promoter, and a tetracycline promoter.

[0171] The polypeptides of interest may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, *e.g.* a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the polypeptide coding sequence that is inserted into the vector. The heterologous signal sequence selected preferably is one that is recognized and processed through one of the standard pathways available within the host cell. The *S. cerevisiae* alpha factor pre-pro signal has proven effective in the secretion of a variety of recombinant proteins from *P. pastoris*. Other yeast signal sequences include the alpha mating factor signal sequence, the invertase signal sequence, and signal sequences derived from other secreted yeast polypeptides. Additionally, these signal peptide sequences may be engineered to provide for enhanced secretion in diploid yeast expression systems. Secretion signals for use in mammalian as

well as yeast cells include mammalian signal sequences, which may be heterologous to the protein being secreted, or may be a native sequence for the protein being secreted. Signal sequences include pre-peptide sequences, and in some instances may include propeptide sequences. Many such signal sequences are known in the art, including the signal sequences found on immunoglobulin chains, *e.g.*, K28 preprotoxin sequence, PHA-E, FACE, human MCP-1, human serum albumin signal sequences, human Ig heavy chain, human Ig light chain, and the like. For example, see Hashimoto *et. al.* Protein Eng 11(2) 75 (1998); and Kobayashi *et. al.* Therapeutic Apheresis 2(4) 257 (1998).

[0172] Transcription may be increased by inserting a transcriptional activator sequence into the vector. These activators are cis-acting elements of DNA, usually about from 10 to 300 bp, which act on a promoter to increase its transcription. Transcriptional enhancers are relatively orientation and position independent, having been found 5' and 3' to the transcription unit, within an intron, as well as within the coding sequence itself. The enhancer may be spliced into the expression vector at a position 5' or 3' to the coding sequence, but is preferably located at a site 5' from the promoter.

[0173] Expression vectors used in eukaryotic host cells may also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from 3' to the translation termination codon, in untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA.

[0174] Construction of suitable vectors containing one or more of the above-listed components employs standard ligation techniques or PCR/recombination methods. Isolated plasmids or DNA fragments are cleaved, tailored, and re-ligated in the form desired to generate the plasmids required or via recombination methods. For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform host cells, and successful transformants selected by antibiotic resistance (*e.g.* ampicillin or Zeocin) where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion and/or sequenced.

[0175] As an alternative to restriction and ligation of fragments, recombination methods based on att sites and recombination enzymes may be used to insert DNA sequences into a vector. Such methods are described, for example, by Landy (1989) *Ann.Rev.Biochem.* 58:913-949; and are known to those of skill in the art. Such methods utilize intermolecular DNA recombination that is mediated by a mixture of lambda and *E. coli* –encoded recombination proteins. Recombination occurs between specific attachment (*att*) sites on the interacting DNA molecules. For a description of att sites see Weisberg and Landy (1983) Site-Specific Recombination in Phage Lambda, in *Lambda II*, Weisberg, ed.(Cold Spring Harbor, NY:Cold Spring Harbor Press), pp. 211-250. The DNA segments flanking the recombination sites are switched, such that after recombination, the *att* sites are hybrid sequences comprised of sequences donated by each parental vector. The recombination can occur between DNAs of any topology.

[0176] *Att* sites may be introduced into a sequence of interest by ligating the sequence of interest into an appropriate vector; generating a PCR product containing *att* B sites through the use of specific primers; generating a cDNA library cloned into an appropriate vector containing *att* sites; and the like.

[0177] *Folding*, as used herein, refers to the three-dimensional structure of polypeptides and proteins, where interactions between amino acid residues act to stabilize the structure. Proper folding is typically the arrangement of a polypeptide that results in optimal biological activity, and in the case of antibodies can conveniently be monitored by assays for activity, *e.g.* antigen binding.

[0178] The expression host may be further modified by the introduction of sequences encoding one or more enzymes that enhance folding and disulfide bond formation, *i.e.* foldases, chaperonins, *etc.* Such sequences may be constitutively or inducibly expressed in the yeast host cell, using vectors, markers, *etc.* as known in the art. Preferably the sequences, including transcriptional regulatory elements sufficient for the desired pattern of expression, are stably integrated in the yeast genome through a targeted methodology.

[0179] For example, the eukaryotic PDI is not only an efficient catalyst of protein cysteine oxidation and disulfide bond isomerization, but also exhibits chaperone activity. Co-expression of PDI can facilitate the production of active proteins having multiple disulfide bonds. Also of interest is the expression of BIP (immunoglobulin heavy chain binding protein); cyclophilin; and the like. In one embodiment of the invention, each of the haploid parental strains expresses a distinct folding enzyme, *e.g.* one strain may express BIP, and the other strain may express PDI or combinations thereof.

[0180] The terms "*desired protein*" or "*desired antibody*" are used interchangeably and refer generally to a parent antibody specific to a target, *i.e.*, CGRP or a chimeric or humanized antibody or a binding portion thereof derived therefrom as described herein. The term "antibody" is intended to include any polypeptide chain-containing molecular structure with a specific shape that fits to and recognizes an epitope, where one or more non-covalent binding interactions stabilize the complex between the molecular structure and the epitope. The archetypal antibody molecule is the immunoglobulin, and all types of immunoglobulins, IgG, IgM, IgA, IgE, IgD, *etc.*, from all sources, *e.g.* human, rodent, rabbit, cow, sheep, pig, dog, other mammals, chicken, other avians, *etc.*, are considered to be "antibodies." A preferred source for producing antibodies useful as starting material according to the invention is rabbits. Numerous antibody coding sequences have been described; and others may be raised by methods well-known in the art. Examples thereof include chimeric antibodies, human antibodies and other non-human mammalian antibodies, humanized antibodies, single chain antibodies (such as scFvs), camelbodies, nanobodies, IgNAR (single-chain antibodies derived from sharks), small-modular immunopharmaceuticals (SMIPs), and antibody fragments such as Fabs, Fab', F(ab')₂ and the like. See Streltsov VA, *et al.*, Structure of a shark IgNAR antibody variable domain and modeling of an early-developmental isotype, *Protein Sci.* 2005 Nov;14(11):2901-9. Epub 2005 Sep 30; Greenberg AS, *et al.*, A new antigen receptor gene family that undergoes

rearrangement and extensive somatic diversification in sharks, *Nature*. 1995 Mar 9;374(6518):168-73; Nuttall SD, et al., Isolation of the new antigen receptor from wobbegong sharks, and use as a scaffold for the display of protein loop libraries, *Mol Immunol*. 2001 Aug;38(4):313-26; Hamers-Casterman C, et al., Naturally occurring antibodies devoid of light chains, *Nature*. 1993 Jun 3;363(6428):446-8; Gill DS, et al., Biopharmaceutical drug discovery using novel protein scaffolds, *Curr Opin Biotechnol*. 2006 Dec; 17(6):653-8. Epub 2006 Oct 19.

[0181] For example, antibodies or antigen binding fragments may be produced by genetic engineering. In this technique, as with other methods, antibody-producing cells are sensitized to the desired antigen or immunogen. The messenger RNA isolated from antibody producing cells is used as a template to make cDNA using PCR amplification. A library of vectors, each containing one heavy chain gene and one light chain gene retaining the initial antigen specificity, is produced by insertion of appropriate sections of the amplified immunoglobulin cDNA into the expression vectors. A combinatorial library is constructed by combining the heavy chain gene library with the light chain gene library. This results in a library of clones which co-express a heavy and light chain (resembling the Fab fragment or antigen binding fragment of an antibody molecule). The vectors that carry these genes are co-transfected into a host cell. When antibody gene synthesis is induced in the transfected host, the heavy and light chain proteins self-assemble to produce active antibodies that can be detected by screening with the antigen or immunogen.

[0182] Antibody coding sequences of interest include those encoded by native sequences, as well as nucleic acids that, by virtue of the degeneracy of the genetic code, are not identical in sequence to the disclosed nucleic acids, and variants thereof. Variant polypeptides can include amino acid (aa) substitutions, additions or deletions. The amino acid substitutions can be conservative amino acid substitutions or substitutions to eliminate non-essential amino acids, such as to alter a glycosylation site, or to minimize misfolding by substitution or deletion of one or more cysteine residues that are not necessary for function. Variants can be designed so as to retain or have enhanced biological activity of a particular region of the protein (*e.g.*, a functional domain, catalytic amino acid residues, *etc.*). Variants also include fragments of the polypeptides disclosed herein, particularly biologically active fragments and/or fragments corresponding to functional domains. Techniques for in vitro mutagenesis of cloned genes are known. Also included in the subject invention are polypeptides that have been modified using ordinary molecular biological techniques so as to improve their resistance to proteolytic degradation or to optimize solubility properties or to render them more suitable as a therapeutic agent.

[0183] Chimeric antibodies may be made by recombinant means by combining the variable light and heavy chain regions (V_L and V_H), obtained from antibody producing cells of one species with the constant light and heavy chain regions from another. Typically chimeric antibodies utilize rodent or rabbit variable regions and human constant regions, in order to produce an antibody with predominantly human domains. The production of such chimeric antibodies is well known in the art,

and may be achieved by standard means (as described, e.g., in U.S. Patent No. 5,624,659, incorporated herein by reference in its entirety). It is further contemplated that the human constant regions of chimeric antibodies of the invention may be selected from IgG1, IgG2, IgG3, and IgG4 constant regions.

[0184] Humanized antibodies are engineered to contain even more human-like immunoglobulin domains, and incorporate only the complementarity-determining regions of the animal-derived antibody. This is accomplished by carefully examining the sequence of the hyper-variable loops of the variable regions of the monoclonal antibody, and fitting them to the structure of the human antibody chains. Although facially complex, the process is straightforward in practice. See, e.g., U.S. Patent No. 6,187,287, incorporated fully herein by reference.

[0185] In addition to entire immunoglobulins (or their recombinant counterparts), immunoglobulin fragments comprising the epitope binding site (e.g., Fab', F(ab')₂, or other fragments) may be synthesized. "Fragment," or minimal immunoglobulins may be designed utilizing recombinant immunoglobulin techniques. For instance "Fv" immunoglobulins for use in the present invention may be produced by synthesizing a fused variable light chain region and a variable heavy chain region. Combinations of antibodies are also of interest, e.g. diabodies, which comprise two distinct Fv specificities. In another embodiment of the invention, SMIPs (small molecule immunopharmaceuticals), camelbodies, nanobodies, and IgNAR are encompassed by immunoglobulin fragments.

[0186] Immunoglobulins and fragments thereof may be modified post-translationally, e.g. to add effector moieties such as chemical linkers, detectable moieties, such as fluorescent dyes, enzymes, toxins, substrates, bioluminescent materials, radioactive materials, chemiluminescent moieties and the like, or specific binding moieties, such as streptavidin, avidin, or biotin, and the like may be utilized in the methods and compositions of the present invention. Examples of additional effector molecules are provided *infra*.

[0187] A polynucleotide sequence "corresponds" to a polypeptide sequence if translation of the polynucleotide sequence in accordance with the genetic code yields the polypeptide sequence (i.e., the polynucleotide sequence "encodes" the polypeptide sequence), one polynucleotide sequence "corresponds" to another polynucleotide sequence if the two sequences encode the same polypeptide sequence.

[0188] A "heterologous" region or domain of a DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. Another example of a heterologous region is a construct where the coding sequence itself is not found in nature (e.g., a cDNA where the genomic coding sequence contains introns, or synthetic

sequences having codons different than the native gene). Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

[0189] A "coding sequence" is an in-frame sequence of codons that (in view of the genetic code) correspond to or encode a protein or peptide sequence. Two coding sequences correspond to each other if the sequences or their complementary sequences encode the same amino acid sequences. A coding sequence in association with appropriate regulatory sequences may be transcribed and translated into a polypeptide. A polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence. A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. Promoter sequences typically contain additional sites for binding of regulatory molecules (e.g., transcription factors) which affect the transcription of the coding sequence. A coding sequence is "under the control" of the promoter sequence or "operatively linked" to the promoter when RNA polymerase binds the promoter sequence in a cell and transcribes the coding sequence into mRNA, which is then in turn translated into the protein encoded by the coding sequence.

[0190] Vectors are used to introduce a foreign substance, such as DNA, RNA or protein, into an organism or host cell. Typical vectors include recombinant viruses (for polynucleotides) and liposomes (for polypeptides). A "DNA vector" is a replicon, such as plasmid, phage or cosmid, to which another polynucleotide segment may be attached so as to bring about the replication of the attached segment. An "expression vector" is a DNA vector which contains regulatory sequences which will direct polypeptide synthesis by an appropriate host cell. This usually means a promoter to bind RNA polymerase and initiate transcription of mRNA, as well as ribosome binding sites and initiation signals to direct translation of the mRNA into a polypeptide(s). Incorporation of a polynucleotide sequence into an expression vector at the proper site and in correct reading frame, followed by transformation of an appropriate host cell by the vector, enables the production of a polypeptide encoded by said polynucleotide sequence.

[0191] "Amplification" of polynucleotide sequences is the *in vitro* production of multiple copies of a particular nucleic acid sequence. The amplified sequence is usually in the form of DNA. A variety of techniques for carrying out such amplification are described in a review article by Van Brunt (1990, Bio/Technol., 8(4):291-294). Polymerase chain reaction or PCR is a prototype of nucleic acid amplification, and use of PCR herein should be considered exemplary of other suitable amplification techniques.

[0192] The general structure of antibodies in vertebrates now is well understood (Edelman, G. M., Ann. N.Y. Acad. Sci., 190: 5 (1971)). Antibodies consist of two identical light polypeptide chains of molecular weight approximately 23,000 daltons (the "light chain"), and two identical heavy chains of molecular weight 53,000-70,000 (the "heavy chain"). The four chains are joined by disulfide bonds in a "Y" configuration wherein the light chains bracket the heavy chains starting at the mouth

of the “Y” configuration. The “branch” portion of the “Y” configuration is designated the F_{ab} region; the stem portion of the “Y” configuration is designated the F_c region. The amino acid sequence orientation runs from the N-terminal end at the top of the “Y” configuration to the C-terminal end at the bottom of each chain. The N-terminal end possesses the variable region having specificity for the antigen that elicited it, and is approximately 100 amino acids in length, there being slight variations between light and heavy chain and from antibody to antibody.

[0193] The variable region is linked in each chain to a constant region that extends the remaining length of the chain and that within a particular class of antibody does not vary with the specificity of the antibody (i.e., the antigen eliciting it). There are five known major classes of constant regions that determine the class of the immunoglobulin molecule (IgG, IgM, IgA, IgD, and IgE corresponding to γ , μ , α , δ , and ϵ (gamma, mu, alpha, delta, or epsilon) heavy chain constant regions). The constant region or class determines subsequent effector function of the antibody, including activation of complement (Kabat, E. A., *Structural Concepts in Immunology and Immunochemistry*, 2nd Ed., p. 413-436, Holt, Rinehart, Winston (1976)), and other cellular responses (Andrews, D. W., *et al.*, *Clinical Immunobiology*, pp 1-18, W. B. Sanders (1980); Kohl, S., *et al.*, *Immunology*, 48: 187 (1983)); while the variable region determines the antigen with which it will react. Light chains are classified as either κ (kappa) or λ (lambda). Each heavy chain class can be prepared with either kappa or lambda light chain. The light and heavy chains are covalently bonded to each other, and the “tail” portions of the two heavy chains are bonded to each other by covalent disulfide linkages when the immunoglobulins are generated either by hybridomas or by B cells.

[0194] The expression “variable region” or “VR” refers to the domains within each pair of light and heavy chains in an antibody that are involved directly in binding the antibody to the antigen. Each heavy chain has at one end a variable domain (V_H) followed by a number of constant domains. Each light chain has a variable domain (V_L) at one end and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain.

[0195] The expressions “complementarity determining region,” “hypervariable region,” or “CDR” refer to one or more of the hyper-variable or complementarity determining regions (CDRs) found in the variable regions of light or heavy chains of an antibody (*See* Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest*, National Institutes of Health, Bethesda, Md., (1987)). These expressions include the hypervariable regions as defined by Kabat *et al.* (“Sequences of Proteins of Immunological Interest,” Kabat E., *et al.*, US Dept. of Health and Human Services, 1983) or the hypervariable loops in 3-dimensional structures of antibodies (Chothia and Lesk, *J Mol. Biol.* 196 901-917 (1987)). The CDRs in each chain are held in close proximity by framework regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site. Within the CDRs there are select amino acids that have been described as the selectivity determining regions (SDRs) which represent the critical contact residues used by the CDR in the

antibody-antigen interaction (Kashmiri, S., *Methods*, 36:25-34 (2005)). In the present invention when specific antibody amino acid or nucleic acid residues are referenced by number this generally refers to its position within a specified amino acid or nucleic acid sequence (i.e., particular sequence identifier) and/or in accordance with Kabat *et al* numbering.

[0196] The expressions “framework region” or “FR” refer to one or more of the framework regions within the variable regions of the light and heavy chains of an antibody (*See* Kabat, E. A. *et al.*, *Sequences of Proteins of Immunological Interest*, National Institutes of Health, Bethesda, Md., (1987)). These expressions include those amino acid sequence regions interposed between the CDRs within the variable regions of the light and heavy chains of an antibody.

[0197] “C_{max}” refers to the maximum (or peak) concentration that an antibody or other compound achieves in tested area (e.g., in the serum or another compartment such as cerebrospinal fluid) after the drug has been administered. For example, serum C_{max} may be measured from serum, e.g., prepared by collecting a blood sample, allowing it to clot and separating solid components by centrifugation or other means to yield serum (blood containing neither blood cells nor clotting factors), and then detecting the concentration of the analyte in the serum by ELISA or other means known in the art.

[0198] “AUC” refers to the area under the concentration-time curve which is expressed in units of mg/mL * hr (or equivalently mg*hr/ml) unless otherwise specified. “AUC_{0-t}” refers to the area under the concentration-time curve from time=0 to last quantifiable concentration. “AUC_{0-inf}” refers to the area under the concentration-time curve from time=0 extrapolated to infinity.

[0199] “I_{max}” refers to the maximal pharmacodynamic response elicited by an anti-CGRP antibody dosage, preferably a dosage of 350 mg or more, more typically at least 750 or 1000 mg, as compared to the response elicited by a lower anti-CGRP antibody doses, e.g., wherein such response may be detected by the inhibition of vasodilation after topical application of capsaicin.

[0200] Anti-CGRP Antibodies and Binding Fragments Thereof Having Binding Specificity for CGRP

[0201] The invention specifically includes the use of specific anti-CGRP antibodies and antibody fragments referred to herein as Ab1-Ab14 which comprise or consist of the CDR, VL, VH, CL, CH polypeptides sequences identified in **FIGs. 1A-12**. The polypeptides comprised in an especially preferred anti-CGRP antibody, Ab6 is further described below.

[0202] Antibody Ab6

[0203] In a preferred exemplary embodiment, the invention includes humanized antibodies having binding specificity to CGRP and possessing a variable light chain sequence comprising the sequence set forth below:

QVLTQSPSSLSASVGDRVTINCQASQSVYHNTYLAWYQQKPGKVPKQLIYDASTLASGVPSR
FSGSGSGTDFTLTISSLQPEDVATYYCLGSYDCTNGDCFVFGGGTKVEIKR (SEQ ID NO: 222).

[0204] The invention also includes humanized antibodies having binding specificity to CGRP and possessing a light chain sequence comprising the sequence set forth below:

QVLTQSPSSLSASVGDRVTINCQASQSVYHNTYLAWYQQKPGKVPKQLIYDASTLASGVPSR
FSGSGSGTDFLTISLQPEDVATYYCLGSYDCTNGDCFVFGGGTKVEIKRTVAAPSVFIFPPS
DEQLKSGTASVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYLSSTLTLSK
ADYEEKHKVYACEVTHQGLSPVTKSFNRGEC (SEQ ID NO: 221).

[0205] The invention further includes humanized antibodies having binding specificity to CGRP and possessing a variable heavy chain sequence comprising the sequence set forth below:

EVQLVESGGGLVQPGGSLRLSCAVSGIDLSGYMNWVRQAPGKGLEWVGIVINGATYYAS
WAKGRFTISRDNKSTTVYLQMNSLRAEDTAVYFCARGDIWGQGTLLTVSS (SEQ ID NO:
202).

[0206] The invention also includes humanized antibodies having binding specificity to CGRP and possessing a heavy chain sequence comprising the sequence set forth below:

EVQLVESGGGLVQPGGSLRLSCAVSGIDLSGYMNWVRQAPGKGLEWVGIVINGATYYAS
WAKGRFTISRDNKSTTVYLQMNSLRAEDTAVYFCARGDIWGQGTLLTVSSASTKGPSVFPLA
PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
LGTQTYICNVNHKPSNTKVDARVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISR
TPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLN
GKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA
VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQ
KSLSLSPGK (SEQ ID NO: 201).

[0207] Alternatively, the heavy chain of Ab6 may lack the C-terminal lysine of SEQ ID NO: 201, i.e., a heavy chain sequence comprising the sequence set forth below:

EVQLVESGGGLVQPGGSLRLSCAVSGIDLSGYMNWVRQAPGKGLEWVGIVINGATYYAS
WAKGRFTISRDNKSTTVYLQMNSLRAEDTAVYFCARGDIWGQGTLLTVSSASTKGPSVFPLA
PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS
LGTQTYICNVNHKPSNTKVDARVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISR
TPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLN
GKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIA
VEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQ
KSLSLSPG (SEQ ID NO: 566).

[0208] The invention further contemplates antibodies comprising one or more of the polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228 which correspond to the complementarity-determining regions (CDRs, or hypervariable regions) of the variable light chain sequence of SEQ ID NO: 222 or the light chain sequence of SEQ ID NO: 221, and/or one or more of the polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11) which correspond to the complementarity-determining regions (CDRs, or

hypervariable regions) of the variable heavy chain sequence of SEQ ID NO: 202 or the heavy chain sequence of SEQ ID NO: 201 or SEQ ID NO: 566, or combinations of these polypeptide sequences. In another embodiment of the invention, the antibodies of the invention or fragments thereof comprise, or alternatively consist of, combinations of one or more of the CDRs, the variable heavy and variable light chain sequences, and the heavy and light chain sequences set forth above, including all of them.

[0209] The invention also contemplates fragments of the antibody having binding specificity to CGRP. In one embodiment of the invention, antibody fragments of the invention comprise, or alternatively consist of, the polypeptide sequence of SEQ ID NO: 222 or SEQ ID NO: 221. In another embodiment of the invention, antibody fragments of the invention comprise, or alternatively consist of, the polypeptide sequence of SEQ ID NO: 202 or SEQ ID NO: 201 or SEQ ID NO: 566.

[0210] In a further embodiment of the invention, fragments of the antibody having binding specificity to CGRP comprise, or alternatively consist of, one or more of the polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228 which correspond to the complementarity-determining regions (CDRs, or hypervariable regions) of the variable light chain sequence of SEQ ID NO: 222 or the light chain sequence of SEQ ID NO: 221.

[0211] In a further embodiment of the invention, fragments of the antibody having binding specificity to CGRP comprise, or alternatively consist of, one or more of the polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11) which correspond to the complementarity-determining regions (CDRs, or hypervariable regions) of the variable heavy chain sequence of SEQ ID NO: 202 or the heavy chain sequence of SEQ ID NO: 201 or SEQ ID NO: 566.

[0212] The invention also contemplates antibody fragments which include one or more of the antibody fragments described herein. In one embodiment of the invention, fragments of the antibodies having binding specificity to CGRP comprise, or alternatively consist of, one, two, three or more, including all of the following antibody fragments: the variable light chain region of SEQ ID NO: 222; the variable heavy chain region of SEQ ID NO: 202; the complementarity-determining regions (SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228) of the variable light chain region of SEQ ID NO: 222; and the complementarity-determining regions (SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11)) of the variable heavy chain region of SEQ ID NO: 202.

[0213] In a particularly preferred embodiment of the invention, the humanized anti- CGRP antibody is Ab6, comprising, or alternatively consisting of, SEQ ID NO: 221 and SEQ ID NO: 201 or SEQ ID NO: 566, and having at least one of the biological activities set forth herein.

[0214] In a further particularly preferred embodiment of the invention, antibody fragments comprise, or alternatively consist of, Fab (fragment antigen binding) fragments having binding specificity for CGRP. With respect to antibody Ab6, the Fab fragment includes the variable light

chain sequence of SEQ ID NO: 222 and the variable heavy chain sequence of SEQ ID NO: 202. This embodiment of the invention further contemplates additions, deletions, and variants of SEQ ID NO: 222 and/or SEQ ID NO: 202 in said Fab while retaining binding specificity for CGRP.

[0215] In another particularly preferred embodiment of the invention, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202, which polypeptides optionally are respectively linked to human light and heavy constant region polypeptides, e.g., human IgG1, IgG2, IgG3 or IgG4 constant regions, which constant regions optionally may be modified to alter glycosylation or proteolysis, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells.

[0216] In another particularly preferred embodiment of the invention, said anti-CGRP antibody may comprise the antibody expression product isolated from recombinant cells which express nucleic acid sequences encoding the light chain of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566, wherein said recombinant cells optionally comprise yeast or mammalian cells, e.g., *Pichia pastoris* or CHO cells, wherein the constant regions thereof optionally may be modified to alter glycosylation or proteolysis or other effector functions.

[0217] In another particularly preferred embodiment of the invention, any of the aforementioned anti-CGRP antibodies or antibody fragments may be optionally comprised in a formulation as disclosed herein, e.g., comprising histidine (L-histidine), sorbitol, polysorbate 80, such as, per 1 mL volume, about 100 mg anti-CGRP antibody, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, having a pH of about 5.8.

[0218] In one embodiment of the invention described herein (infra), Fab fragments may be produced by enzymatic digestion (e.g., papain) of Ab6. In another embodiment of the invention, anti-CGRP antibodies such as Ab6 or Fab fragments thereof may be produced via expression in mammalian cells such as CHO, NSO or HEK 293 cells, fungal, insect, or microbial systems such as yeast cells (for example diploid yeast such as diploid *Pichia*) and other yeast strains. Suitable *Pichia* species include, but are not limited to, *Pichia pastoris*.

[0219] In another embodiment, antibody fragments may be present in one or more of the following non-limiting forms: Fab, Fab', F(ab')₂, Fv and single chain Fv antibody forms. In a preferred embodiment, the anti-CGRP antibodies described herein further comprises the kappa constant light chain sequence comprising the sequence set forth below:

[0220] TVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 563).

[0221] In another preferred embodiment, the anti-CGRP antibodies described herein further comprises the gamma-1 constant heavy chain polypeptide sequence comprising the sequence set forth

below or the same sequence lacking the carboxy terminal lysine residue (SEQ ID NO: 564 and SEQ ID NO: 565, respectively):

[0222] ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELL
GGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY
ASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM
TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQG
NVFSCSVMHREALHNHYTQKSLSLSPGK (SEQ ID NO: 564).

[0223] ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPA
VLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELL
GGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY
ASTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM
TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQG
NVFSCSVMHREALHNHYTQKSLSLSPG (SEQ ID NO: 565).

[0224] For clarity, any antibody disclosed herein is intended to include any variant of the disclosed constant region variant sequences, e.g., Ab6 may comprise the constant region of SEQ ID NO: 564 containing the C-terminal lysine or may comprise the constant region of SEQ ID NO: 565 lacking the C-terminal lysine. Thus, every disclosure herein of the heavy chain of SEQ ID NO: 201 also includes a variant lacking the C-terminal lysine residue thereof, *i.e.*, having the heavy chain variable region sequence of Ab6 (SEQ ID NO: 202) and the constant region sequence of SEQ ID NO: 565. For example, the sequence encoding an antibody comprising a C-terminal lysine in the heavy chain may, when expressed in cell lines such as CHO cells, produce an antibody lacking said C-terminal lysine due to proteolysis, or a mixture of heavy chains containing or lacking said C-terminal lysine.

[0225] In another embodiment, the invention contemplates use of an isolated anti-CGRP antibody comprising a V_H polypeptide sequence selected from: SEQ ID NO: 2, SEQ ID NO: 42, SEQ ID NO: 82, SEQ ID NO: 122, SEQ ID NO: 162, SEQ ID NO: 202, SEQ ID NO: 242, SEQ ID NO: 282, SEQ ID NO: 322, SEQ ID NO: 362, SEQ ID NO: 402, SEQ ID NO: 442, SEQ ID NO: 482, or SEQ ID NO: 522, or a variant thereof; and further comprising a V_L polypeptide sequence selected from: SEQ ID NO: 22, SEQ ID NO: 62, SEQ ID NO: 102, SEQ ID NO: 142, SEQ ID NO: 182, SEQ ID NO: 222, SEQ ID NO: 262, SEQ ID NO: 302, SEQ ID NO: 342, SEQ ID NO: 382, SEQ ID NO: 422, SEQ ID NO: 462, SEQ ID NO: 502, or SEQ ID NO: 542, or a variant thereof, wherein one or more of the framework residues (FR residues) in said V_H or V_L polypeptide has been substituted with another amino acid residue resulting in an anti-CGRP antibody that specifically binds CGRP. The invention contemplates humanized and chimeric forms of these antibodies. The chimeric antibodies may include an Fc derived from IgG1, IgG2, IgG3, or IgG4 constant regions.

[0226] In one embodiment of the invention, the antibodies or V_H or V_L polypeptides originate or are selected from one or more rabbit B cell populations prior to initiation of the humanization process referenced herein.

[0227] In another embodiment of the invention, the anti-CGRP antibodies and fragments thereof do not have binding specificity for CGRP-R. In a further embodiment of the invention, the anti-CGRP antibodies and fragments thereof inhibit the association of CGRP with CGRP-R. In another embodiment of the invention, the anti-CGRP antibodies and fragments thereof inhibit the association of CGRP with CGRP-R and/or additional proteins and/or multimers thereof, and/or antagonizes the biological effects thereof.

[0228] As stated herein, antibodies and fragments thereof may be modified post-translationally to add effector moieties such as chemical linkers, detectable moieties such as for example fluorescent dyes, enzymes, substrates, bioluminescent materials, radioactive materials, and chemiluminescent moieties, or functional moieties such as for example streptavidin, avidin, biotin, a cytotoxin, a cytotoxic agent, and radioactive materials.

[0229] Antibodies or fragments thereof may also be chemically modified to provide additional advantages such as increased solubility, stability and circulating time (*in vivo* half-life) of the polypeptide, or decreased immunogenicity (See U.S. Pat. No. 4,179,337). The chemical moieties for derivatization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The antibodies and fragments thereof may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

[0230] The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog). For example, the polyethylene glycol may have an average molecular weight of about 200, 500, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10,000, 10,500, 11,000, 11,500, 12,000, 12,500, 13,000, 13,500, 14,000, 14,500, 15,000, 15,500, 16,000, 16,500, 17,000, 17,500, 18,000, 18,500, 19,000, 19,500, 20,000, 25,000, 30,000, 35,000, 40,000, 50,000, 55,000, 60,000, 65,000, 70,000, 75,000, 80,000, 85,000, 90,000, 95,000, or 100,000 kDa. Branched polyethylene glycols are described, for example, in U.S. Pat. No. 5,643,575; Morpurgo et al., *Appl. Biochem. Biotechnol.* 56:59-72 (1996); Vorobjev et al., *Nucleosides Nucleotides* 18:2745-

2750 (1999); and Caliceti et al., *Bioconjug. Chem.* 10:638-646 (1999), the disclosures of each of which are incorporated herein by reference.

[0231] There are a number of attachment methods available to those skilled in the art, See e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), See also Malik et al., *Exp. Hematol.* 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

[0232] As suggested above, polyethylene glycol may be attached to proteins via linkage to any of a number of amino acid residues. For example, polyethylene glycol can be linked to polypeptides via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (e.g., lysine, histidine, aspartic acid, glutamic acid, or cysteine) or to more than one type of amino acid residue (e.g., lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof).

[0233] Alternatively, antibodies or fragments thereof may have increased *in vivo* half-lives via fusion with albumin (including but not limited to recombinant human serum albumin or fragments or variants thereof (See, e.g., U.S. Pat. No. 5,876,969, issued Mar. 2, 1999, EP Patent 0 413 622, and U.S. Pat. No. 5,766,883, issued Jun. 16, 1998, herein incorporated by reference in their entirety)) or other circulating blood proteins such as transferrin or ferritin. In a preferred embodiment, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) are fused with the mature form of human serum albumin (i.e., amino acids 1-585 of human serum albumin as shown in FIGS. 1 and 2 of EP Patent 0 322 094) which is herein incorporated by reference in its entirety. Polynucleotides encoding fusion proteins of the invention are also encompassed by the invention.

[0234] Regarding detectable moieties, further exemplary enzymes include, but are not limited to, horseradish peroxidase, acetylcholinesterase, alkaline phosphatase, beta-galactosidase and luciferase. Further exemplary fluorescent materials include, but are not limited to, rhodamine, fluorescein, fluorescein isothiocyanate, umbelliferone, dichlorotriazinylamine, phycoerythrin and dansyl chloride. Further exemplary chemiluminescent moieties include, but are not limited to, luminol. Further exemplary bioluminescent materials include, but are not limited to, luciferin and aequorin. Further exemplary radioactive materials include, but are not limited to, Iodine 125 (¹²⁵I), Carbon 14 (¹⁴C), Sulfur 35 (³⁵S), Tritium (³H) and Phosphorus 32 (³²P).

[0235] Regarding functional moieties, exemplary cytotoxic agents include, but are not limited to, methotrexate, aminopterin, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine; alkylating agents such as mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU), mitomycin C, lomustine (CCNU), 1-methylnitrosourea, cyclophosphamide, mechlorethamine, busulfan, dibromomannitol, streptozotocin, mitomycin C, cis-dichlorodiamine platinum (II) (DDP) cisplatin and carboplatin (paraplatin); anthracyclines include daunorubicin (formerly daunomycin), doxorubicin (adriamycin), detorubicin, carminomycin, idarubicin, epirubicin, mitoxantrone and bisantrene; antibiotics include dactinomycin (actinomycin D), bleomycin, calicheamicin, mithramycin, and anthramycin (AMC); and antimetabolic agents such as the vinca alkaloids, vincristine and vinblastine. Other cytotoxic agents include paclitaxel (taxol), ricin, pseudomonas exotoxin, gemcitabine, cytochalasin B, gramicidin D, ethidium bromide, emetine, etoposide, teniposide, colchicin, dihydroxy anthracin dione, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, procarbazine, hydroxyurea, asparaginase, corticosteroids, mytotate (O,P⁺-(DDD)), interferons, and mixtures of these cytotoxic agents.

[0236] Further cytotoxic agents include, but are not limited to, chemotherapeutic agents such as carboplatin, cisplatin, paclitaxel, gemcitabine, calicheamicin, doxorubicin, 5-fluorouracil, mitomycin C, actinomycin D, cyclophosphamide, vincristine and bleomycin. Toxic enzymes from plants and bacteria such as ricin, diphtheria toxin and Pseudomonas toxin may be conjugated to the humanized or chimeric antibodies, or binding fragments thereof, to generate cell-type-specific-killing reagents (Youle, et al., Proc. Nat'l Acad. Sci. USA 77:5483 (1980); Gilliland, et al., Proc. Nat'l Acad. Sci. USA 77:4539 (1980); Krock, et al., Proc. Nat'l Acad. Sci. USA 77:5419 (1980)).

[0237] Other cytotoxic agents include cytotoxic ribonucleases as described by Goldenberg in U.S. Pat. No. 6,653,104. Embodiments of the invention also relate to radioimmunoconjugates where a radionuclide that emits alpha or beta particles is stably coupled to the antibody, or binding fragments thereof, with or without the use of a complex-forming agent. Such radionuclides include beta-emitters such as Phosphorus-32 (³²P), Scandium-47 (⁴⁷Sc), Copper-67 (⁶⁷Cu), Gallium-67 (⁶⁷Ga), Yttrium-88 (⁸⁸Y), Yttrium-90 (⁹⁰Y), Iodine-125 (¹²⁵I), Iodine-131 (¹³¹I), Samarium-153 (¹⁵³Sm), Lutetium-177 (¹⁷⁷Lu), Rhenium-186 (¹⁸⁶Re) or Rhenium-188 (¹⁸⁸Re), and alpha-emitters such as Astatine-211 (²¹¹At), Lead-212 (²¹²Pb), Bismuth-212 (²¹²Bi) or -213 (²¹³Bi) or Actinium-225 (²²⁵Ac).

[0238] Methods are known in the art for conjugating an antibody or binding fragment thereof to a detectable moiety and the like, such as for example those methods described by Hunter et al, Nature 144:945 (1962); David et al, Biochemistry 13:1014 (1974); Pain et al, J. Immunol. Meth. 40:219 (1981); and Nygren, J., Histochem. and Cytochem. 30:407 (1982).

[0239] Embodiments described herein further include variants and equivalents that are substantially homologous to the antibodies, antibody fragments, diabodies, SMIPs, camelbodies, nanobodies, IgNAR, polypeptides, variable regions and CDRs set forth herein. These may contain, e.g., conservative substitution mutations, (i.e., the substitution of one or more amino acids by similar

amino acids). For example, conservative substitution refers to the substitution of an amino acid with another within the same general class, e.g., one acidic amino acid with another acidic amino acid, one basic amino acid with another basic amino acid, or one neutral amino acid by another neutral amino acid. What is intended by a conservative amino acid substitution is well known in the art.

[0240] In another embodiment, the invention contemplates polypeptide sequences having at least 90% or greater sequence homology to any one or more of the polypeptide sequences of antibody fragments, variable regions and CDRs set forth herein. More preferably, the invention contemplates polypeptide sequences having at least 95% or greater sequence homology, even more preferably at least 98% or greater sequence homology, and still more preferably at least 99% or greater sequence homology to any one or more of the polypeptide sequences of antibody fragments, variable regions and CDRs set forth herein. Methods for determining homology between nucleic acid and amino acid sequences are well known to those of ordinary skill in the art.

[0241] In another embodiment, the invention further contemplates the above-recited polypeptide homologs of the antibody fragments, variable regions and CDRs set forth herein further having anti-CGRP activity. Non-limiting examples of anti-CGRP activity are set forth herein.

[0242] The present invention also contemplates anti-CGRP antibodies comprising any of the polypeptide or polynucleotide sequences described herein substituted for any of the other polynucleotide sequences described herein. For example, without limitation thereto, the present invention contemplates antibodies comprising the combination of any of the variable light chain and variable heavy chain sequences described herein, and further contemplates antibodies resulting from substitution of any of the CDR sequences described herein for any of the other CDR sequences described herein.

[0243] Additional Exemplary Embodiments of the Invention

[0244] In another embodiment, the invention contemplates treatment methods using one or more anti-human CGRP antibodies or antibody fragments thereof which specifically bind to the same overlapping linear or conformational epitope(s) and/or competes for binding to the same overlapping linear or conformational epitope(s) on an intact human CGRP polypeptide or fragment thereof as an anti-human CGRP antibody selected from Ab1, Ab2, Ab3, Ab4, Ab5, Ab6, Ab7, Ab8, Ab9, Ab10, Ab11, Ab12, Ab13, or Ab14. In a preferred embodiment, the anti-human CGRP antibody or fragment thereof specifically binds to the same overlapping linear or conformational epitope(s) and/or competes for binding to the same overlapping linear or conformational epitope(s) on an intact human CGRP polypeptide or a fragment thereof as Ab3, Ab6, Ab13, or Ab14.

[0245] A preferred embodiment of the invention is directed to treatment methods using chimeric or humanized antibodies and fragments thereof (including Fab fragments) having binding specificity for CGRP and inhibiting biological activities mediated by the binding of CGRP to the CGRP receptor. In a particularly preferred embodiment of the invention, the chimeric or humanized anti-CGRP antibodies are selected from Ab3, Ab6, Ab13, or Ab14.

[0246] In another embodiment of the invention, the anti-human CGRP antibody used in the described treatment methods is an antibody which specifically binds to the same overlapping linear or conformational epitopes on an intact CGRP polypeptide or fragment thereof that is (are) specifically bound by Ab3, Ab6, Ab13, or Ab14 as ascertained by epitopic mapping using overlapping linear peptide fragments which span the full length of the native human CGRP polypeptide.

[0247] In another embodiment, the invention is also directed to treatment methods using an isolated anti-CGRP antibody or antibody fragment comprising one or more of the CDRs contained in the V_H polypeptide sequences selected from: 3, 13, 23, 33, 43, 53, 63, 73, 83, 93, 103, 113, 123, or 133, or a variant thereof, and/or one or more of the CDRs contained in the V_L polypeptide sequences selected from: 1, 11, 21, 31, 41, 51, 61, 71, 81, 91, 101, 111, 121, or 131, or a variant thereof.

[0248] In one embodiment of the invention, the anti-human CGRP antibody discussed in the two prior paragraphs comprises at least 2 complementarity determining regions (CDRs) in each the variable light and the variable heavy regions which are identical to those contained in an anti-human CGRP antibody selected from Ab1, Ab2, Ab3, Ab4, Ab5, Ab6, Ab7, Ab8, Ab9, Ab10, Ab11, Ab12, Ab13, or Ab14.

[0249] In a preferred embodiment, the anti-human CGRP antibody used in the described treatment methods comprises at least 2 complementarity determining regions (CDRs) in each the variable light and the variable heavy regions which are identical to those contained in Ab3 or Ab6. In another embodiment, all of the CDRs of the anti-human CGRP antibody discussed above are identical to the CDRs contained in an anti-human CGRP antibody selected from Ab1, Ab2, Ab3, Ab4, Ab5, Ab6, Ab7, Ab8, Ab9, Ab10, Ab11, Ab12, Ab13, or Ab14. In a preferred embodiment of the invention, all of the CDRs of the anti-human CGRP antibody discussed above are identical to the CDRs contained in an anti-human CGRP antibody selected from Ab3 or Ab6.

[0250] The invention further contemplates treatment methods wherein the one or more anti-human CGRP antibodies discussed above are aglycosylated or if glycosylated are only mannosylated; that contain an Fc region that has been modified to alter effector function, half-life, proteolysis, and/or glycosylation; are human, humanized, single chain or chimeric; and are a humanized antibody derived from a rabbit (parent) anti-human CGRP antibody. An exemplary mutation which impairs glycosylation comprises the mutation of the Asn residue at position 297 of an IgG heavy chain constant region such as IgG1 to another amino acid, such as Ala as described in U.S. Pat. No. 5,624,821, which is incorporated by reference in its entirety.

[0251] The invention further contemplates one or more anti-human CGRP antibodies wherein the framework regions (FRs) in the variable light region and the variable heavy regions of said antibody respectively are human FRs which are unmodified or which have been modified by the substitution of one or more human FR residues in the variable light or heavy chain region with the corresponding FR residues of the parent rabbit antibody, and wherein said human FRs have been derived from human variable heavy and light chain antibody sequences which have been selected

from a library of human germline antibody sequences based on their high level of homology to the corresponding rabbit variable heavy or light chain regions relative to other human germline antibody sequences contained in the library.

[0252] The invention also contemplates a method of treating or preventing medication overuse headache, e.g., associated with the overuse of anti-migraine drugs and/or associated with triptan and/or ergot and/or analgesic overuse, comprising administering to a patient exhibiting medication overuse headache or at risk of developing medication overuse headache a therapeutically effective amount of at least one anti-human CGRP antibody or fragment described herein. The invention also contemplates that the treatment method may involve the administration of two or more anti-CGRP antibodies or fragments thereof and disclosed herein. If more than one antibody is administered to the patient, the multiple antibodies may be administered simultaneously or concurrently, or may be staggered in their administration. The anti-CGRP activity of the anti-CGRP antibodies of the present invention, and fragments thereof having binding specificity to CGRP, may also be described by their strength of binding or their affinity for CGRP. In one embodiment of the invention, the anti-CGRP antibodies of the present invention, and fragments thereof having binding specificity to CGRP, bind to CGRP with a dissociation constant (K_D) of less than or equal to 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, or 10^{-13} M. Preferably, the anti-CGRP antibodies and fragments thereof bind CGRP with a dissociation constant of less than or equal to 10^{-11} M, 5×10^{-12} M, or 10^{-12} M. In another embodiment of the invention, the anti-CGRP antibodies of the present invention, and fragments thereof having binding specificity to CGRP, bind to a linear or conformational CGRP epitope.

[0253] In another embodiment of the invention, the anti-CGRP activity of the anti-CGRP antibodies of the present invention, and fragments thereof having binding specificity to CGRP, bind to CGRP with an off-rate of less than or equal to 10^{-4} S⁻¹, 5×10^{-5} S⁻¹, 10^{-5} S⁻¹, 5×10^{-6} S⁻¹, 10^{-6} S⁻¹, 5×10^{-7} S⁻¹, or 10^{-7} S⁻¹.

[0254] In a further embodiment of the invention, the anti-CGRP activity of the anti-CGRP antibodies of the present invention, and fragments thereof having binding specificity to CGRP, exhibit anti-CGRP activity by preventing, ameliorating or reducing the symptoms of, or alternatively treating, diseases and disorders associated with CGRP. Non-limiting examples of diseases and disorders associated with CGRP are set forth herein and include headache and migraine disorders.

[0255] Polynucleotides Encoding Anti-CGRP Antibody Polypeptides

[0256] As aforementioned the invention specifically includes the use of specific anti-CGRP antibodies and antibody fragments referred to herein as Ab1-Ab14 which comprise or consist of the CDR, VL, VH, CL, and CH polypeptides having the sequences identified in **FIGs. 1A-12**. The nucleic acid sequences encoding the foregoing VL, VH, CL, and CH polypeptides comprised in Ab1-Ab14 are also comprised in **FIGs. 1A-12**. The nucleic acid sequences which encode the CDR, VL,

VH, CL, and CH polypeptides of an especially preferred anti-CGRP antibody, Ab6, are further described below.

[0257] Antibody Ab6

[0258] The invention is further directed to polynucleotides encoding antibody polypeptides having binding specificity to CGRP. In one embodiment of the invention, polynucleotides of the invention comprise, or alternatively consist of, the following polynucleotide sequence encoding the variable light chain polypeptide sequence of SEQ ID NO: 222:

[0259] CAAGTGCTGaccagtcctccatcctccctgtctgcatctgtaggagacagagtcaccatcAATtgcCAGGCCA GTCAGAGTGTTTATCATAACACCTACCTGGCCtggtatcagcagaaaccaggaaagttcctaaCAActgatcta tGATGCATCCACTCTGGCATCTggggccccatctcgtttcagtgccagtggtatctgggacagatttcactctcaccatcagcgc ctgcagcctgaagatgttgaacttattactgtCTGGGCAGTTATGATTGTACTAATGGTGATTGTTTTGTTtgcg gcggaggaaccaaggtggaatcaaactgt (SEQ ID NO: 232).

[0260] In one embodiment of the invention, polynucleotides of the invention comprise, or alternatively consist of, the following polynucleotide sequence encoding the light chain polypeptide sequence of SEQ ID NO: 221:

[0261] CAAGTGCTGaccagtcctccatcctccctgtctgcatctgtaggagacagagtcaccatcAATtgcCAGGCCA GTCAGAGTGTTTATCATAACACCTACCTGGCCtggtatcagcagaaaccaggaaagttcctaaCAActgatcta tGATGCATCCACTCTGGCATCTggggccccatctcgtttcagtgccagtggtatctgggacagatttcactctcaccatcagcgc ctgcagcctgaagatgttgaacttattactgtCTGGGCAGTTATGATTGTACTAATGGTGATTGTTTTGTTtgcg gcggaggaaccaaggtggaatcaaactgtACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGAT GAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGA GAGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAG TGTCACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGCTGA GCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTG AGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTAG (SEQ ID NO: 231).

[0262] In another embodiment of the invention, polynucleotides of the invention comprise, or alternatively consist of, the following polynucleotide sequence encoding the variable heavy chain polypeptide sequence of SEQ ID NO: 202:

[0263] gagggtgcagctTgtggagtctgggggaggcttggtccagcctggggggtccctgagactctctgtgcaGTCtctggaAT CGACCTCagtGGCTACTACATGAACtgggtccgtcaggctccagggaaggggctggagtgggtcGGAGTCATTGG TATTAATGGTGCCACATACTACGCGAGCTGGGCGAAAGGCcgattcaccatctccagagacaattccaagA CCACGGTGtatcttcaaatgaacagcctgagagctgaggacactgctgtgtatTTCtgtGCTAGAGGGGACATCtggggcc aagggaacctcgtcaccgtcTCGAGC (SEQ ID NO: 212).

[0264] In one embodiment of the invention, polynucleotides of the invention comprise, or alternatively consist of, the following polynucleotide sequence encoding the heavy chain polypeptide sequence of SEQ ID NO: 201:

[0265] gaggtgcagctTgtggagtctgggggaggcttggtccagcctggggggtccctgagactctcctgtgcaGTCtctggaATCGACCTCagtGGCTACTACATGAACtgggtccgtcaggctccagggaaggggctggagtgggtcGGAGTCATTGGTATTAATGGTGCCACATACTACGCGAGCTGGGCGAAAGGCcgattcaccatctccagagacaattccaagAACACGGTGtatcttcaaatgaacagcctgagagctgaggacactgctgtgtatTTCtgtGCTAGAGGGGACATCtggggccaaaggaccctcgtcaccgtcTCGAGCGCCTCCACCAAGGGGCCATCGGTCTTCCCCCTGGCAcCCTCTCCaAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCCGGCTGTCTACAGTCTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGACGCGAGAGTTGAGCCCAAATCTTGTGACAAAACCTCACACATGCCACCGTGCCCAGCACCTGAACCTCTGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGACACCTCATGaTCTCCCgGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAA GCCGCGGGAGGAGCAGTACGCCAGCACGTACCGTGTGGTCAGCGTCTCACCCTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA GCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTA CACCTGCCCCCATCCCCGGGAGGAGATGACCAAGAACCAGGTACGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACCTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGC AAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGAT GCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATG A (SEQ ID NO: 211).

[0266] In one embodiment of the invention, polynucleotides of the invention comprise, or alternatively consist of, the following polynucleotide sequence encoding the heavy chain polypeptide sequence of SEQ ID NO: 566:

gaggtgcagctTgtggagtctgggggaggcttggtccagcctggggggtccctgagactctcctgtgcaGTCtctggaATCGACCTCagtGGCTACTACATGAACtgggtccgtcaggctccagggaaggggctggagtgggtcGGAGTCATTGGTATTAATGGTGCCACATACTACGCGAGCTGGGCGAAAGGCcgattcaccatctccagagacaattccaagAACACGGTGtatcttcaaatgaacagcctgagagctgaggacactgctgtgtatTTCtgtGCTAGAGGGGACATCtggggccaaggaccctcgtcaccgtcTCGAGCGCCTCCACCAAGGGGCCATCGGTCTTCCCCCTGGCAcCCTCTCCaAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCCGGCTGCTACAGTCTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGACCCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACGCGAGAGTTGAGCCCAAATCTTGTGACAAAACCTCACACATGCCACCGTGCCAGCACCTGAACCTCTGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGACACCTCATGaTCTCCCgGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGG

TCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGG
 GAGGAGCAGTACGCCAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGA
 CTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCA
 TCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTG
 CCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGG
 CTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCT
 ACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCA
 CCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAG
 GCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTTGA (SEQ ID NO:
 567).

[0267] In a further embodiment of the invention, polynucleotides encoding antibody fragments having binding specificity to CGRP comprise, or alternatively consist of, one or more of the polynucleotide sequences of SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238 which correspond to polynucleotides encoding the complementarity-determining regions (CDRs, or hypervariable regions) of the light chain variable sequence of SEQ ID NO: 222 or the light chain sequence of SEQ ID NO: 221.

[0268] In a further embodiment of the invention, polynucleotides encoding antibody fragments having binding specificity to CGRP comprise, or alternatively consist of, one or more of the polynucleotide sequences of SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12) which correspond to polynucleotides encoding the complementarity-determining regions (CDRs, or hypervariable regions) of the heavy chain variable sequence of SEQ ID NO: 202 or the heavy chain sequence of SEQ ID NO: 201 or SEQ ID NO: 566.

[0269] The invention also contemplates polynucleotide sequences including one or more of the polynucleotide sequences encoding antibody fragments described herein. In one embodiment of the invention, polynucleotides encoding antibody fragments having binding specificity to CGRP comprise, or alternatively consist of, one, two, three or more, including all of the following polynucleotides encoding antibody fragments: the polynucleotide SEQ ID NO: 232 encoding the light chain variable sequence of SEQ ID NO: 222; the polynucleotide SEQ ID NO: 231 encoding the light chain sequence of SEQ ID NO: 221; the polynucleotide SEQ ID NO: 212 encoding the heavy chain variable sequence of SEQ ID NO: 202; the polynucleotide SEQ ID NO: 211 encoding the heavy chain sequence of SEQ ID NO: 201; the polynucleotide SEQ ID NO: 567 encoding the heavy chain sequence of SEQ ID NO: 566; polynucleotides encoding the complementarity-determining regions (SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238) of the light chain variable sequence of SEQ ID NO: 222 or the light chain sequence of SEQ ID NO: 221; and polynucleotides encoding the complementarity-determining regions (SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12)) of the heavy chain variable sequence of SEQ ID NO: 202 or the heavy chain sequence of SEQ ID NO: 201 or SEQ ID NO: 566.

[0270] In a preferred embodiment of the invention, polynucleotides of the invention comprise, or alternatively consist of, polynucleotides encoding Fab (fragment antigen binding) fragments having binding specificity for CGRP. With respect to antibody Ab6, the polynucleotides encoding the full length Ab6 antibody comprise, or alternatively consist of, the polynucleotide SEQ ID NO: 231 encoding the light chain sequence of SEQ ID NO: 221 and the polynucleotide SEQ ID NO: 211 encoding the heavy chain sequence of SEQ ID NO: 201 or the polynucleotide SEQ ID NO: 567 encoding the heavy chain sequence of SEQ ID NO: 566.

[0271] Another embodiment of the invention contemplates these polynucleotides incorporated into an expression vector for expression in mammalian cells such as CHO, NSO, HEK-293, or in fungal, insect, or microbial systems such as yeast cells such as the yeast *Pichia*. Suitable *Pichia* species include, but are not limited to, *Pichia pastoris*. In one embodiment of the invention described herein (infra), Fab fragments may be produced by enzymatic digestion (e.g., papain) of Ab6 following expression of the full-length polynucleotides in a suitable host. In another embodiment of the invention, anti-CGRP antibodies such as Ab6 or Fab fragments thereof may be produced via expression of Ab6 polynucleotides in mammalian cells such as CHO, NSO or HEK 293 cells, fungal, insect, or microbial systems such as yeast cells (for example diploid yeast such as diploid *Pichia*) and other yeast strains. Suitable *Pichia* species include, but are not limited to, *Pichia pastoris*.

[0272] In one embodiment, the invention is directed to an isolated polynucleotide comprising a polynucleotide encoding an anti-CGRP V_H antibody amino acid sequence selected from SEQ ID NO: 2, SEQ ID NO: 42, SEQ ID NO: 82, SEQ ID NO: 122, SEQ ID NO: 162, SEQ ID NO: 202, SEQ ID NO: 242, SEQ ID NO: 282, SEQ ID NO: 322, SEQ ID NO: 362, SEQ ID NO: 402, SEQ ID NO: 442, SEQ ID NO: 482, or SEQ ID NO: 522 or encoding a variant thereof wherein at least one framework residue (FR residue) has been substituted with an amino acid present at the corresponding position in a rabbit anti-CGRP antibody V_H polypeptide or a conservative amino acid substitution.

[0273] In another embodiment, the invention is directed to an isolated polynucleotide comprising the polynucleotide sequence encoding an anti-CGRP V_L antibody amino acid sequence of SEQ ID NO: 22, SEQ ID NO: 62, SEQ ID NO: 102, SEQ ID NO: 142, SEQ ID NO: 182, SEQ ID NO: 222, SEQ ID NO: 262, SEQ ID NO: 302, SEQ ID NO: 342, SEQ ID NO: 382, SEQ ID NO: 422, SEQ ID NO: 462, SEQ ID NO: 502, or SEQ ID NO: 542, or encoding a variant thereof wherein at least one framework residue (FR residue) has been substituted with an amino acid present at the corresponding position in a rabbit anti-CGRP antibody V_L polypeptide or a conservative amino acid substitution.

[0274] In yet another embodiment, the invention is directed to one or more heterologous polynucleotides comprising a sequence encoding the polypeptides contained in SEQ ID NO: 22 and SEQ ID NO: 2; SEQ ID NO: 62 and SEQ ID NO: 42; SEQ ID NO: 102 and SEQ ID NO: 82; SEQ ID NO: 142 and SEQ ID NO: 122; SEQ ID NO: 182 and SEQ ID NO: 162; SEQ ID NO: 222 and SEQ ID NO: 202; SEQ ID NO: 262 and SEQ ID NO: 242; SEQ ID NO: 302 and SEQ ID NO: 282; SEQ ID NO: 342 and SEQ ID NO: 322; SEQ ID NO: 382 and SEQ ID NO: 362; SEQ ID NO: 422 and

SEQ ID NO: 402; SEQ ID NO: 462 and SEQ ID NO: 442; SEQ ID NO: 502 and SEQ ID NO: 482; or SEQ ID NO: 542 and SEQ ID NO: 522.

[0275] In another embodiment, the invention is directed to an isolated polynucleotide that expresses a polypeptide containing at least one CDR polypeptide derived from an anti-CGRP antibody wherein said expressed polypeptide alone specifically binds CGRP or specifically binds CGRP when expressed in association with another polynucleotide sequence that expresses a polypeptide containing at least one CDR polypeptide derived from an anti-CGRP antibody wherein said at least one CDR is selected from those contained in the V_L or V_H polypeptides of SEQ ID NO: 22, SEQ ID NO: 2, SEQ ID NO: 62, SEQ ID NO: 42, SEQ ID NO: 102, SEQ ID NO: 82, SEQ ID NO: 142, SEQ ID NO: 122, SEQ ID NO: 182, SEQ ID NO: 162, SEQ ID NO: 222, SEQ ID NO: 202, SEQ ID NO: 262, SEQ ID NO: 242, SEQ ID NO: 302, SEQ ID NO: 282, SEQ ID NO: 342, SEQ ID NO: 322, SEQ ID NO: 382, SEQ ID NO: 362, SEQ ID NO: 422, SEQ ID NO: 402, SEQ ID NO: 462, SEQ ID NO: 442, SEQ ID NO: 502, SEQ ID NO: 482, SEQ ID NO: 542, or SEQ ID NO: 522.

[0276] Host cells and vectors comprising said polynucleotides are also contemplated.

[0277] The invention further contemplates vectors comprising the polynucleotide sequences encoding the variable heavy and light chain polypeptide sequences, as well as the individual complementarity-determining regions (CDRs, or hypervariable regions), as set forth herein, as well as host cells comprising said vector sequences. In one embodiment of the invention, the host cell is a yeast cell. In another embodiment of the invention, the yeast host cell belongs to the genus *Pichia*.

[0278] Methods of Producing Antibodies and Fragments thereof

[0279] In another embodiment, the present invention contemplates methods for producing anti-CGRP antibodies and fragments thereof. Methods for producing antibodies and fragments thereof secreted from polyploid, preferably diploid or tetraploid strains of mating competent yeast are taught, for example, in U.S. patent application publication no. US 2009/0022659 to Olson et al., and in U.S. patent no. 7,935,340 to Garcia-Martinez et al., the disclosures of each of which are herein incorporated by reference in their entireties. Methods for producing antibodies and fragments thereof in mammalian cells, e.g., CHO cells are further well known in the art.

[0280] Other methods of producing antibodies are also well known to those of ordinary skill in the art. For example, methods of producing chimeric antibodies are now well known in the art (See, for example, U.S. Patent No. 4,816,567 to Cabilly et al.; Morrison et al., *P.N.A.S. USA*, 81:8651-55 (1984); Neuberger, M.S. et al., *Nature*, 314:268-270 (1985); Boulianne, G.L. et al., *Nature*, 312:643-46 (1984), the disclosures of each of which are herein incorporated by reference in their entireties).

[0281] Likewise, other methods of producing humanized antibodies are now well known in the art (See, for example, U.S. Patent Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370 to Queen et al; U.S. Patent Nos. 5,225,539 and 6,548,640 to Winter; U.S. Patent Nos. 6,054,297, 6,407,213 and 6,639,055 to Carter et al; U.S. Patent No. 6,632,927 to Adair; Jones, P.T. et al, *Nature*, 321:522-525

(1986); Reichmann, L., et al, *Nature*, 332:323-327 (1988); Verhoeyen, M, et al, *Science*, 239:1534-36 (1988), the disclosures of each of which are herein incorporated by reference in their entireties).

[0282] The term "opioid analgesic" herein refers to all drugs, natural or synthetic, with morphine-like actions. The synthetic and semi-synthetic opioid analgesics are derivatives of five chemical classes of compound: phenanthrenes; phenylheptylamines; phenylpiperidines; morphinans; and benzomorphans, all of which are within the scope of the term. Exemplary opioid analgesics include codeine, dihydrocodeine, diacetylmorphine, hydrocodone, hydromorphone, levorphanol, oxycodone, alfentanil, buprenorphine, butorphanol, fentanyl, sufentanil, meperidine, methadone, nalbuphine, propoxyphene and pentazocine or pharmaceutically acceptable salts thereof.

[0283] The term "NSAID" refers to a non-steroidal anti-inflammatory compound. NSAIDs are categorized by virtue of their ability to inhibit cyclooxygenase. Cyclooxygenase 1 and cyclooxygenase 2 are two major isoforms of cyclooxygenase and most standard NSAIDs are mixed inhibitors of the two isoforms. Most standard NSAIDs fall within one of the following five structural categories: (1) propionic acid derivatives, such as ibuprofen, naproxen, naprosyn, diclofenac, and ketoprofen; (2) acetic acid derivatives, such as tolmetin and slindac; (3) fenamic acid derivatives, such as mefenamic acid and meclofenamic acid; (4) biphenylcarboxylic acid derivatives, such as diflunisal and flufenisal; and (5) oxicams, such as piroxim, sudoxicam, and isoxicam. Another class of NSAID has been described which selectively inhibit cyclooxygenase 2. Cox-2 inhibitors have been described, e.g., in U.S. Pat. Nos. 5,616,601; 5,604,260; 5,593,994; 5,550,142; 5,536,752; 5,521,213; 5,475,995; 5,639,780; 5,604,253; 5,552,422; 5,510,368; 5,436,265; 5,409,944; and 5,130,311, all of which are hereby incorporated by reference. Certain exemplary COX-2 inhibitors include celecoxib (SC-58635), DUP-697, flosulide (CGP-28238), meloxicam, 6-methoxy-2 naphthylacetic acid (6-MNA), rofecoxib, MK-966, nabumetone (prodrug for 6-MNA), nimesulide, NS-398, SC-5766, SC-58215, T-614; or combinations thereof.

[0284] In some embodiments, aspirin and/or acetaminophen may be taken in conjunction with the subject CGRP antibody or fragment. Aspirin is another type of non-steroidal anti-inflammatory compound.

[0285] The subject to which the pharmaceutical formulation is administered can be, e.g., any human or non-human animal that is in need of such treatment, prevention and/or amelioration, or who would otherwise benefit from the inhibition or attenuation of medication overuse headache. For example, the subject can be an individual that is diagnosed with, or who is deemed to be at risk of being afflicted by medication overuse headache. The present invention further includes the use of any of the pharmaceutical formulations disclosed herein in the manufacture of a medicament for the treatment, prevention and/or amelioration of medication overuse headache.

[0286] Administration

[0287] In an embodiment of the invention, the anti-CGRP antibodies described herein, or CGRP binding fragments thereof, are administered to a subject at a dosage of 400 mg. In a preferred

embodiment of the invention, the anti-CGRP antibodies described herein, or CGRP binding fragments thereof, as well as combinations of said antibodies or antibody fragments, are administered to a recipient subject with a frequency of once every twenty-six weeks or six months or less, such as once every sixteen weeks or four months or less, once every eight weeks or two months or less, once every four weeks or monthly or less, once every two weeks or bimonthly or less, once every week or less, or once daily or less. In general the administration of sequential doses may vary by plus or minus a few days from the aforementioned schedule, e.g., administration every 3 months or every 12 weeks includes administration of a dose varying from the schedule day by plus or minus 1, 2, 3, 4, 5, 5, or 7 days.

[0288] In another embodiment of the invention, the anti-CGRP antibodies described herein, or CGRP binding fragments thereof, as well as combinations of said antibodies or antibody fragments, are administered to a subject in a pharmaceutical formulation.

[0289] A “pharmaceutical composition” refers to a chemical or biological composition suitable for administration to a mammal. Such compositions may be specifically formulated for administration via one or more of a number of routes, including but not limited to buccal, epicutaneous, epidural, inhalation, intraarterial, intracardial, intracerebroventricular, intradermal, intramuscular, intranasal, intraocular, intraperitoneal, intraspinal, intrathecal, intravenous, oral, parenteral, rectally via an enema or suppository, subcutaneous, subdermal, sublingual, transdermal, and transmucosal, preferably intravenous. In addition, administration can occur by means of injection, powder, liquid, gel, drops, or other means of administration.

[0290] A “pharmaceutical excipient” or a “pharmaceutically acceptable excipient” is a carrier, usually a liquid, in which an active therapeutic agent is formulated. In one embodiment of the invention, the active therapeutic agent is a humanized antibody described herein, or one or more fragments thereof. The excipient generally does not provide any pharmacological activity to the formulation, though it may provide chemical and/or biological stability, and release characteristics. Exemplary formulations can be found, for example, in Remington’s Pharmaceutical Sciences, 19th Ed., Grennaro, A., Ed., 1995 which is incorporated by reference.

[0291] As used herein “pharmaceutically acceptable carrier” or “excipient” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents that are physiologically compatible. In one embodiment, the carrier is suitable for parenteral administration. Alternatively, the carrier can be suitable for intravenous, intraperitoneal, intramuscular, or sublingual administration. Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions of the

invention is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0292] Pharmaceutical compositions typically must be sterile and stable under the conditions of manufacture and storage. The invention contemplates that the pharmaceutical composition is present in lyophilized form. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol), and suitable mixtures thereof. The invention further contemplates the inclusion of a stabilizer in the pharmaceutical composition. The proper fluidity can be maintained, for example, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants.

[0293] In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, monostearate salts and gelatin. Moreover, the alkaline polypeptide can be formulated in a time release formulation, for example in a composition which includes a slow release polymer. The active compounds can be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, polylactic acid and polylactic, polyglycolic copolymers (PLG). Many methods for the preparation of such formulations are known to those skilled in the art.

[0294] An exemplary composition comprises, consists essentially of, or consists of an anti-CGRP antibody or fragment thereof (e.g., Ab6), an excipient such as histidine, an isotonic agent such as sorbitol, and a surfactant such as polysorbate 80 in an aqueous solution. For example, the composition may comprise, consist essentially of, or consist of histidine (L-histidine), sorbitol, polysorbate 80, such as, per 1 mL volume, about 100 mg anti-CGRP antibody (e.g., Ab6), about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, having a pH of about 5.8, or approximately that constitution, e.g., within 10% of those values, within 5% of those values, within 1% of those values, within 0.5% of those values, or within 0.1% of those values, and water. For example, the pH value may be within 10% of 5.8, i.e., between 5.22 and 6.38. The Ab6 antibody may comprise or consist of the variable light and heavy chain polypeptides of SEQ ID NO: 222 and SEQ ID NO: 202 respectively, or the light and heavy chain polypeptides of SEQ ID NO: 221 and SEQ ID NO: 201 respectively, or the light and heavy chain polypeptides of SEQ ID NO: 221 and SEQ ID NO: 566 respectively. The composition may be in the form of an aqueous solution, or a concentrate (e.g., lyophilized) which when reconstituted, e.g., by addition of water, yields the aforementioned constitution. An exemplary composition consists of, per mL, 100 mg of the light and heavy chain

polypeptides of SEQ ID NO: 221 and SEQ ID NO: 201 respectively, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, and water Q.S. or approximately that constitution, e.g., within 10% of those quantities, within 5% of those quantities, within 1% of those quantities, within 0.5% of those quantities, or within 0.1% of those quantities. Another exemplary composition consists of, per mL, 100 mg of the light and heavy chain polypeptides of SEQ ID NO: 221 and SEQ ID NO: 566 respectively, about 3.1 mg L-Histidine, about 40.5 mg Sorbitol, and about 0.15 mg Polysorbate 80, and water Q.S. or approximately that constitution, e.g., within 10% of those quantities, within 5% of those quantities, within 1% of those quantities, within 0.5% of those quantities, or within 0.1% of those quantities. The composition may be suitable for intravenous or subcutaneous administration, preferably intravenous administration. For example, the composition may be suitable for mixing with an intravenous solution (such as 0.9% sodium chloride) at an amount of between about 100 mg and about 300 mg antibody added to 100 mL of intravenous solution. Preferably the composition may be shelf-stable for at least 1, 3, 6, 12, 18, or 24 months, e.g., showing formation of aggregates of no more than 5% or no more than 10% of the antibody or fragment after storage at room temperature or when refrigerated at 4°C for the specified duration, or in an accelerated aging test that simulates storage for that duration.

[0295] For each of the recited embodiments, the compounds can be administered by a variety of dosage forms. Any biologically-acceptable dosage form known to persons of ordinary skill in the art, and combinations thereof, are contemplated. Examples of such dosage forms include, without limitation, reconstitutable powders, elixirs, liquids, solutions, suspensions, emulsions, powders, granules, particles, microparticles, dispersible granules, cachets, inhalants, aerosol inhalants, patches, particle inhalants, implants, depot implants, injectables (including subcutaneous, intramuscular, intravenous, and intradermal, preferably intravenous), infusions, and combinations thereof.

[0296] The above description of various illustrated embodiments, of the invention is not intended to be exhaustive or to limit the invention to the precise form disclosed. While specific embodiments, of, and examples for, the invention are described herein for illustrative purposes, various equivalent modifications are possible within the scope of the invention, as those skilled in the relevant art will recognize. The teachings provided herein of the invention can be applied to other purposes, other than the examples described above.

[0297] These and other changes can be made to the invention in light of the above detailed description. In general, in the following claims, the terms used should not be construed to limit the invention to the specific embodiments, disclosed in the specification and the claims. Accordingly, the invention is not limited by the disclosure, but instead the scope of the invention is to be determined entirely by the following claims.

[0298] The invention may be practiced in ways other than those particularly described in the foregoing description and examples. Numerous modifications and variations of the invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

[0299] Certain CGRP antibody polynucleotides and polypeptides are disclosed in the sequence listing accompanying this patent application filing, and the disclosure of said sequence listing is herein incorporated by reference in its entirety.

[0300] The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is herein incorporated by reference in their entireties.

[0301] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the subject invention, and are not intended to limit the scope of what is regarded as the invention. Efforts have been made to ensure accuracy with respect to the numbers used (e.g. amounts, temperature, concentrations, etc.) but some experimental errors and deviations should be allowed for. Unless otherwise indicated, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees centigrade; and pressure is at or near atmospheric.

ADDITIONAL EXEMPLARY EMBODIMENTS

S1. Use of an anti-CGRP antibody for the manufacture of a medicament for treating cluster headache (e.g. chronic cluster headache and/or episodic headache) in a patient in the need of immediate relief of cluster headache symptoms or for prevention of cluster headache (e.g. chronic cluster headache and/or episodic headache) in a patient in need of immediate preventative treatment of cluster headache, wherein said medicament is for intravenous infusion in a dosage of 400 mg of said anti-CGRP antibody, wherein said anti-CGRP antibody comprises the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively.

S2. Use of the anti-CGRP antibody of embodiment S1, wherein said medicament is for use in a patient that exhibits at least one cluster headache symptom at the time of administration. S3. Use of the anti-CGRP antibody of embodiment S2, wherein said at least one cluster headache symptom comprises at least five attacks according to the blow list A to C

A. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes (when untreated)

B. 1 or 2 or more of the following

(1) at least one of the following symptoms or signs, ipsilateral to the headache:

i. conjunctival injection and/or lacrimation

ii. nasal congestion and/or rhinorrhoea

iii. eyelid oedema

iv. forehead and facial sweating

v. miosis and/or ptosis

(2) a sense of restlessness or agitation

C. Occurring with a frequency between one every other day and eight per day

S4. Use of the anti-CGRP antibody of embodiment S3, wherein the subject experience head pain.

S5. Use of the anti-CGRP antibody of any one of embodiments S2-S4, wherein the symptom is alleviated after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

S6. Use of the anti-CGRP antibody of any one of embodiments S2-S5, wherein said patient no longer has a cluster headache after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

S7. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12), respectively.

S8. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222.

S9. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232.

S10. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable heavy chain polypeptide of SEQ ID NO: 202.

S11. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

- S12. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202.
- S13. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212.
- S14. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221.
- S15. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231.
- S16. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.
- S17. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.
- S18. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.
- S19. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.
- S20. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said intravenous administration is infused over a period of approximately 30 min to 60 minutes.
- S21. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein the cluster headache symptoms decline or are abolished immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

S22. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said patient is cluster headache free 2 hours post-completion of infusion.

S23. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody is for intravenous administration in a dosage of about 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

S24. Use of the anti-CGRP antibody of any one of embodiments S1-S22, wherein said anti-CGRP antibody is for intravenous administration in a dosage of 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

S25. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said anti-CGRP antibody is comprised in a formulation comprising or consisting of histidine (L-histidine), sorbitol, polysorbate 80, and water.

S26. Use of the anti-CGRP antibody of embodiment S25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 10% of said values, and having a pH of 5.8 or within +/- 10% of said value.

S27. Use of the anti-CGRP antibody of embodiment S25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 5% of said values, and/or having a pH of 5.8 or within +/- 5% of said value.

S28. Use of the anti-CGRP antibody of embodiment S25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 1% of said values, and/or having a pH of 5.8 or within +/- 1% of said value.

S29. Use of the anti-CGRP antibody of embodiment S25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.5% of said values, and/or having a pH of 5.8 or within +/- 0.5% of said value.

S30. Use of the anti-CGRP antibody of embodiment S25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.1% of said values, and/or having a pH of 5.8 or within +/- 0.1% of said value.

S31. Use of the anti-CGRP antibody of any one of embodiments S25-S30, wherein said L-Histidine in said formulation comprises a mixture of L-Histidine and L-Histidine monohydrate.

S32. Use of the anti-CGRP antibody of any one of embodiments S25-S30, wherein said 3.1 mg of histidine in said formulation comprises a mixture of L-Histidine (1 mg) and L-Histidine monohydrate (2.8 mg), which in the final formulation sums up to 3.1 mg L-histidine free base.

S33. Use of the anti-CGRP antibody of any one of embodiments S26-S32, wherein said formulation is comprised in a 100 mg/mL single-400 vial wherein each mL contains 100 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

S34. Use of the anti-CGRP antibody of any one of embodiments S26-S32, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

S35. Use of the anti-CGRP antibody of any one of embodiments S26-S32, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine such as L-histidine hydrochloride monohydrate, polysorbate 80, sorbitol, and Water for Injection, USP, at a pH of or about 5.8.

S36. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said medicament is for administration to a patient that exhibits a pain level of at least 2 on the VRS-4 at the time of administration of said antibody.

S37. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said medicament is for administration to a patient that exhibits a pain level of at least 3 on the VRS-4 at the time of administration of said antibody.

S38. Use of the anti-CGRP antibody of any one of embodiments S1-S37, wherein said medicament is for administration to a patient that exhibits a pain level of at most 2 on the VRS-4 immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

S38. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said medicament is for administration to a patient that exhibits a pain level at most 1 on the VRS-4

immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

S40. Use of the anti-CGRP antibody of any one of the foregoing embodiments, wherein said medicament is for administration to a patient that is not administered any acute migraine medication within a period of time before and after said administration, such as within 15 minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said administration.

S41. Use of the anti-CGRP antibody of embodiment S40, wherein said acute migraine medication comprises a triptan, an analgesic such as non-opioids or opioids/narcotics, acetaminophen, an NSAID, a combination medication, an ergotamine, or an ergot derivative.

S42. Use of the anti-CGRP antibody of embodiment S41, wherein said non-opioid analgesic comprises paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two drugs with analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol, butorphanol, morphine, codeine, hydrocodone, or any combination thereof), barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

S43. Use of the anti-CGRP antibody of any one of embodiments S1-S39, wherein the patient is receiving or has received additional migraine medication.

S44. Use of the anti-CGRP antibody of any one of embodiments S1-S39 or S43, wherein the patient receives migraine medication prior, concurrent or after administration of the anti-CGRP antibody.

S45. Use of the anti-CGRP antibody of any one of embodiments S1-S39 or S43-S44, wherein the patient receives migraine medication within a period of time before and after said anti-CGRP antibody administration, such as within 15 minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said anti-CGRP antibody administration.

S46. Use of the anti-CGRP antibody of any one of embodiments S44 or S45, wherein said migraine medication comprises an acute and/or a chronic migraine medication.

S47. Use of the anti-CGRP antibody of any one of embodiments S44-S46, wherein said migraine medication comprises a triptan, an analgesic such as non-opioid or opioid/narcotic, acetaminophen, an NSAID, a combination medication, an ergotamine, or an ergot derivative.

S48. Use of the anti-CGRP antibody of embodiment S47, wherein said non-opioid analgesic comprises paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two drugs with analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol, butorphanol, morphine, codeine, hydrocodone, or any combination thereof), barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

S49. Use of the anti-CGRP antibody of any of any one of the foregoing embodiments, wherein said anti-CGRP antibody is expressed in or obtained by expression in *Pichia pastoris*.

S50. Use of the anti-CGRP antibody of any of any one of embodiments S1-S48, wherein said anti-CGRP antibody is expressed in or obtained by expression in CHO cells.

S51. Use of the anti-CGRP antibody of any of any one of the foregoing embodiments, wherein said patient is administered 400 mg of said anti-CGRP antibody every three months.

S52. Use of the anti-CGRP antibody of any of any one of the foregoing embodiments, wherein said method results in immediate relief of cluster headache symptoms.

S53. Use of the anti-CGRP antibody of any of any one of the foregoing embodiments, wherein said method results in immediate preventative treatment of cluster headache.

S54. Use of the anti-CGRP antibody of any of any one of the foregoing embodiments, wherein the cluster headache symptom is selected from the list consisting of or comprising severe or very severe unilateral orbital, supraorbital and/or temporal pain e.g. lasting 15–180 minutes (when untreated), and the one or more (such as 1, 2, 3, 4, 5, or all) of the following symptoms or signs, ipsilateral to the headache:

conjunctival injection and/or lacrimation; nasal congestion and/or rhinorrhea; eyelid oedema; forehead and facial sweating; miosis and/or ptosis; a sense of restlessness or agitation.

FURTHER EXEMPLARY EMBODIMENTS

E1. An anti-CGRP antibody for use in treating cluster headache (e.g. chronic or episodic cluster headache) in a patient in the need of immediate relief of symptoms or for use in preventing cluster headache (e.g. chronic or episodic cluster headache) in a patient in need of immediate preventative treatment of cluster headache, wherein said anti-CGRP antibody is for intravenous infusion in a dosage of 400 mg of said anti-CGRP antibody, wherein said anti-CGRP antibody comprises the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively.

E2. The anti-CGRP antibody for use according to embodiment E1, wherein said anti-CGRP antibody is for use in a patient that exhibits at least one cluster headache symptom at the time of administration.

[0302] E3. The anti-CGRP antibody for use according to embodiment E2, wherein said at least one cluster headache symptom comprises at least five attacks according to the blow list A to C

A. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes (when untreated)

B. 1 or 2 or more of the following

(1) at least one of the following symptoms or signs, ipsilateral to the headache:

- i. conjunctival injection and/or lacrimation
- ii. nasal congestion and/or rhinorrhoea
- iii. eyelid oedema
- iv. forehead and facial sweating
- v. miosis and/or ptosis

(2) a sense of restlessness or agitation

C. Occurring with a frequency between one every other day and eight per day

E4. The anti-CGRP antibody for use according to embodiment E3, wherein the subject experience head pain.

E5. The anti-CGRP antibody for use according to any one of embodiments, E2-E4, wherein the most cluster headache is alleviated after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours

after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

E6. The anti-CGRP antibody for use according to any one of embodiments E2-E5, wherein said patient no longer has a cluster headache after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

E7. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12), respectively.

E8. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222.

E9. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232.

E10. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable heavy chain polypeptide of SEQ ID NO: 202.

E11. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

E12. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202.

E13. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

E14. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221.

E15. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231.

E16. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

E17. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

E18. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

E19. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

E20. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said intravenous administration is infused over a period of approximately 30 min to 60 minutes.

E21. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein the cluster headache symptoms decline or are abolished immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

E22. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said patient is cluster headache free 2 hours post-completion of infusion.

E23. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is for intravenous administration in a dosage of about 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

E24. The anti-CGRP antibody for use according to any one of embodiments E1-E22, wherein said anti-CGRP antibody is for intravenous administration in a dosage of 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

E25. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is comprised in a formulation comprising or consisting of histidine (L-histidine), sorbitol, polysorbate 80, and water.

E26. The anti-CGRP antibody for use according to embodiment E25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 10% of said values, and having a pH of 5.8 or within +/- 10% of said value.

E27. The anti-CGRP antibody for use according to embodiment E25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 5% of said values, and/or having a pH of 5.8 or within +/- 5% of said value.

E28. The anti-CGRP antibody for use according to embodiment E25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 1% of said values, and/or having a pH of 5.8 or within +/- 1% of said value.

E29. The anti-CGRP antibody for use according to embodiment E25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.5% of said values, and/or having a pH of 5.8 or within +/- 0.5% of said value.

E30. The anti-CGRP antibody for use according to embodiment E25, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.1% of said values, and/or having a pH of 5.8 or within +/- 0.1% of said value.

E31. The anti-CGRP antibody for use according to of any one of embodiments E25-E30, wherein said L- Histidine in said formulation comprises a mixture of L-Histidine and L-Histidine monohydrate.

E32. The anti-CGRP antibody for use according to any one of embodiments E25-E30, wherein said 3.1 mg of histidine in said formulation comprises a mixture of L-Histidine (1 mg) and L-Histidine monohydrate (2.8 mg), which in the final formulation sums up to 3.1 mg L-histidine free base.

E33. The anti-CGRP antibody for use according to any one of embodiments E26-E32, wherein said formulation is comprised in a 100 mg/mL single-dose vial wherein each mL contains 100 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

E34. The anti-CGRP antibody for use according to any one of embodiments E26-E32, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

E35. A formulation comprising a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody according to E1-E35, L-histidine or L-histidine hydrochloride monohydrate, polysorbate 80, sorbitol and Water for Injection, USP, at a pH of about or at 5.8.

E36. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is for administration to a patient that exhibits a pain level of at least 2 on the VRS-4 at the time of administration of said antibody.

E37. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is for administration to a patient that exhibits a pain level of at least 3 on the VRS-4 at the time of administration of said antibody.

E38. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is for administration to a patient that exhibits a pain level of at most 2 or 3 on the VRS-4 immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

E39. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is for administration to a patient that exhibits a pain level at most 1 on the VRS-4 immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

E40. The anti-CGRP antibody for use according to any one of the foregoing embodiments, wherein said anti-CGRP antibody is for administration to a patient that is not administered any acute migraine medication within a period of time before and after said administration, such as within 15 minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said administration.

E41. The anti-CGRP antibody for use according to embodiment E40, wherein said acute migraine medication comprises a triptan, an analgesic such as non-opioids or opioids/narcotics, acetaminophen, an NSAID, a combination medication, an ergotamine, or an ergot derivative.

E42. The anti-CGRP antibody for use according to embodiment E41, wherein said non-opioid analgesic comprises paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two drugs with analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol, butorphanol, morphine, codeine, hydrocodone, or any combination thereof), barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

E43. The anti-CGRP antibody for use of any one of embodiments E1-E39, wherein the patient is receiving or has received migraine medication.

E44. The anti-CGRP antibody for use of any one of embodiments E1-E39 or E43, wherein the patient receives migraine medication prior, concurrent or after administration of the anti-CGRP antibody.

E45. The anti-CGRP antibody for use of embodiments E1-S39 or E43-E44, wherein the patient receives migraine medication within a period of time before and after said anti-CGRP antibody administration, such as within 15 minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said anti-CGRP antibody administration.

E46. The anti-CGRP antibody for use of any one of embodiments E44 or E45, wherein said migraine medication comprises an acute and/or a chronic migraine medication.

E47. The anti-CGRP antibody for use of embodiments E44-E46, wherein said migraine medication comprises a triptan, an analgesic such as non-opioid or opioid/narcotic, acetaminophen, an NSAID, a combination medication, an ergotamine, or an ergot derivative.

E48. The anti-CGRP antibody for use of embodiment E47, wherein said non-opioid analgesic comprises paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two drugs with analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol, butorphanol, morphine, codeine, hydrocodone, or any combination thereof), barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

E49. The anti-CGRP antibody for use according to any of any one of the foregoing embodiments, wherein said anti-CGRP antibody is expressed in or obtained by expression in *Pichia pastoris*.

E50. The anti-CGRP antibody for use according to any of any one of embodiments E1-E39, wherein said anti-CGRP antibody is expressed in or obtained by expression in CHO cells.

E51. The anti-CGRP antibody for use of any of any one of the foregoing embodiments, wherein said patient is administered 400 mg of said anti-CGRP antibody every three months.

E52. The anti-CGRP antibody for use of any of any one of the foregoing embodiments, wherein said method results in immediate relief of cluster headache symptoms.

E53. The anti-CGRP antibody for use of any of any one of the foregoing embodiments, wherein said method results in immediate preventative treatment of cluster headache.

E.54 The anti-CGRP antibody for use of any of any one of the foregoing embodiments, wherein the cluster headache symptom is selected from the list consisting of or comprising severe or very severe unilateral orbital, supraorbital and/or temporal pain e.g. lasting 15–180 minutes (when untreated), and the one or more (such as 1, 2, 3, 4, 5, or all) of the following symptoms or signs, ipsilateral to the headache:

conjunctival injection and/or lacrimation; nasal congestion and/or rhinorrhea; eyelid oedema; forehead and facial sweating; miosis and/or ptosis; a sense of restlessness or agitation.

Further Embodiments

EE1. A method for treatment of cluster headache (such as chronic cluster headache or episodic cluster headache) in a patient with cluster headache symptoms or for prevention of cluster headache in a patient in need of preventative treatment of cluster headache (such as chronic cluster headache or episodic cluster headache), comprising intravenous administering to a patient in need 400 mg of an anti-CGRP antibody comprising the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively.

EE2. The method of embodiment 1, wherein said patient exhibits at least one cluster headache symptom at the time of administration.

EE3. The method of embodiment 1 or 2, wherein the patient experiences at least five attacks according to the blow list A to C:

A. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes (when untreated)

B. 1 and/or 2 of the following:

(1) at least one of the following symptoms or signs, ipsilateral to the headache:

- i. conjunctival injection and/or lacrimation
- ii. nasal congestion and/or rhinorrhea
- iii. eyelid oedema
- iv. forehead and facial sweating
- v. miosis and/or ptosis

(2) a sense of restlessness or agitation

C. Occurring with a frequency between one every other day and eight per day.

EE4. The method of any one of embodiments 1-3, wherein said patient has had or been diagnosed with cluster headache for at least 3 months, at least 6 months, at least 9 months, for at least a year, for at least 2 years, for at least 3 years, or more than 3 years.

EE5. The method of any one of embodiments 1-4, wherein said patient classifies according to the following one or more criteria: is male, is a smoker, is between the age of about 20 to about 50 years, has a family history of cluster headache, is using a medication such as nitroglycerin, or other drug used to treat heart disease, has been diagnosed with abnormalities in the body's biological clock (hypothalamus), the cluster headache is typically not associated with triggers, such as foods, hormonal changes or stress, or upon the onset of a cluster period alcohol use exacerbates or triggers more headaches or increased head pain.

- EE6. The method of any one of embodiments 1-5, wherein said patient experiences head pain.
- EE7. The method of any one of embodiments 1-6, wherein the cluster headache sign or symptom is alleviated immediately after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.
- EE8. The method of any one of embodiments 1-7, wherein said patient no longer has a cluster headache after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.
- EE9. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 234; SEQ ID NO: 236; and SEQ ID NO: 238, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences encoded by SEQ ID NO: 214; SEQ ID NO: 216; and RESIDUE GGGGACATC (SEE PAGE 2, TABLE A, ITEM NO. 12), respectively.
- EE10. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222.
- EE11. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232.
- EE12. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable heavy chain polypeptide of SEQ ID NO: 202.
- EE13. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable heavy chain polypeptide encoded by SEQ ID NO: 212.
- EE14. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202.

EE15. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

EE16. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221.

EE17. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231.

EE18. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

EE19. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

EE20. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

EE21. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

EE22. The method of any one of the foregoing embodiments, wherein said intravenous administration is infused over a period of approximately 30 min to 60 minutes.

EE23. The method of any one of the foregoing embodiments, wherein the cluster headache signs or symptoms decline or are abolished immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

EE24. The method of any one of the foregoing embodiments, wherein said patient is cluster headache free 2 hours post-completion of infusion.

EE25. The method of any one of embodiments 1-23, comprising intravenously administering 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.

EE26. The method of any one of the foregoing embodiments, wherein said anti-CGRP antibody is comprised in a formulation comprising or consisting of histidine (L-histidine), sorbitol, polysorbate 80, and water.

EE27. The method of embodiment 26, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 10% of said values, and having a pH of 5.8 or within +/- 10% of said value.

EE28. The method of embodiment 26, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 5% of said values, and/or having a pH of 5.8 or within +/- 5% of said value.

EE29. The method of embodiment 26, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 1% of said values, and/or having a pH of 5.8 or within +/- 1% of said value.

EE30. The method of embodiment 26, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.5% of said values, and/or having a pH of 5.8 or within +/- 0.5% of said value.

EE31. The method of embodiment 26, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.1% of said values, and/or having a pH of 5.8 or within +/- 0.1% of said value.

EE32. The method of any one of embodiments 26-31 wherein said L- Histidine in said formulation comprises a mixture of L-Histidine and L-Histidine monohydrate.

EE33. The method of any one of embodiments 26-31, wherein said 3.1 mg of histidine in said formulation comprises a mixture of L-Histidine (1 mg) and L-Histidine hydrochloride monohydrate (2.8 mg), which in the final formulation sums up to 3.1 mg L-histidine free base.

EE34. The method of any one of embodiments 26-33, wherein said formulation is comprised in a 100 mg/mL single-dose vial wherein each mL contains 100 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

EE35. The method of any one of embodiments 26-33, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

EE36. The method of any one of embodiments 26-33, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine, L-histidine hydrochloride monohydrate, polysorbate 80, sorbitol, and Water for Injection, USP, at a pH at or about 5.8.

EE37. The method of any one of the foregoing embodiments, wherein the patient exhibits a pain level of at least 2 on the VRS-4 at the time of administration of said antibody.

EE38. The method of any one of the foregoing embodiments, wherein the patient exhibits a pain level of at least 3 on the VRS-4 at the time of administration of said antibody.

EE39. The method of any one of embodiments 1-38, wherein the patient exhibits a pain level of at most 2 on the VRS-4 immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

EE40. The method of any one of the foregoing embodiments, wherein the patient exhibits a pain level at most 1 on the VRS-4 immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

EE41. The method of any one of the foregoing embodiments, wherein the patient is not administered any acute migraine medication within a period of time before and after said administration, such as within 15 minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said administration.

EE42. The method of embodiment 41, wherein said acute migraine medication comprises a triptan, an analgesic such as non-opioids or opioids/narcotics, acetaminophen, an NSAID, a combination medication, an antiemetic, a barbiturate, an ergotamine, or an ergot derivative.

EE43. The method of embodiment 42, wherein said non-opioid analgesic comprises paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two or more drugs, optionally of different drug classes including wherein at least one or two of said drugs elicit analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol, butorphanol, morphine, codeine, hydrocodone, or any combination thereof), a barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

EE44. The method of any one of embodiments 1-40, wherein the patient is receiving or has received migraine or headache medication.

EE45. The method of any one of embodiments 1-40 or 44, wherein the patient receives migraine or headache medication prior, concurrent or after administration of the anti-CGRP antibody.

EE46. The method of any one embodiment 1-40 or 45-45, wherein the patient receives migraine or headache medication within a period of time before and after said anti-CGRP antibody administration, such as within 15 minutes, within 30 minutes, within 1 hour, within 2 hours, within 3 hours, within 4 hours, within 5 hours, or within 6 hours before and after said anti-CGRP antibody administration.

EE47. The method of embodiment 45 or 46, wherein said migraine medication comprises an acute and/or a chronic migraine medication.

EE48. The method of any of embodiments 45-47, wherein said migraine or headache medication comprises a triptan, an analgesic such as non-opioid or opioid/narcotic, acetaminophen, an NSAID, a combination medication, an antiemetic, a barbiturate, an ergotamine, or an ergot derivative.

EE49. The method of embodiment 48, wherein said non-opioid analgesic comprises paracetamol (acetaminophen), acetylsalicylic acid (aspirin), another NSAID, or another non-opioid analgesic; said triptan comprises use of one or more of sumatriptan, zolmitriptan, naratriptan, rizatriptan, eletriptan, almotriptan, or frovatriptan; said opioid comprises use of one or more of oxycodone, tramadol, butorphanol, morphine, codeine, and hydrocodone; said combination medication comprises two or more drugs optionally of different drug classes, wherein at least one or two elicit analgesic effects (for example, paracetamol and codeine), an analgesic and an adjuvant (for example, paracetamol and caffeine) and/or said combination-analgesics comprises at least one opioid (such as tramadol,

butorphanol, morphine, codeine, hydrocodone, or any combination thereof), barbiturate such as butalbital, and/or caffeine, and/or said combination-analgesic comprises acetylsalicylic acid (aspirin), paracetamol and caffeine (EXCEDRIN®, EXCEDRIN MIGRAINE®).

EE50. The method of any one of any one of the foregoing embodiments, wherein said anti-CGRP antibody is expressed in or obtained by expression in *Pichia pastoris*.

EE51. The method of any of any one of embodiments 1-49, wherein said anti-CGRP antibody is expressed in or obtained by expression in CHO cells.

EE52. The method of any of any one of the foregoing embodiments, wherein said patient is administered 400 mg of said anti-CGRP antibody every three months.

EE53. The method of any one of the foregoing embodiments, wherein said method results in immediate relief of cluster headache symptoms.

EE54. The method of any one of the foregoing embodiments, wherein said method results in immediate preventative treatment of cluster headache.

EE55. The method of any one of the foregoing embodiments, wherein the cluster headache symptom is selected from the list consisting of or comprising; severe or very severe unilateral orbital, supraorbital and/or temporal pain e.g. lasting 15–180 minutes (when untreated), and one or more (such as 1, 2, 3, 4, 5, or all) of the following symptoms or signs, ipsilateral to the headache:

conjunctival injection and/or lacrimation; nasal congestion and/or rhinorrhea; eyelid oedema; forehead and facial sweating; miosis and/or ptosis; a sense of restlessness or agitation.

EE56. A dosage formulation for treatment or prevention of cluster headache comprising 400 mg of an anti-CGRP antibody comprising the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively and at least one pharmaceutically acceptable carrier.

EE57. A kit for treatment or prevention of cluster headache comprising (i) one or more single dose dosage formulations each comprising 400 mg of an anti-CGRP antibody comprising the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively and at

least one pharmaceutically acceptable carrier and (ii) a label or other documentation providing directions for *in vivo* administration.

EE58. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the variable light chain polypeptide of SEQ ID NO: 222 and the variable heavy chain polypeptide of SEQ ID NO: 202.

EE59. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the variable light chain polypeptide encoded by SEQ ID NO: 232 and the variable heavy chain polypeptide encoded by SEQ ID NO: 212.

EE60. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221.

EE61. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231.

EE62. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

EE63. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

EE64. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

EE65. The dosage formulation or kit of embodiment 56 or 57, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

EE66. The dosage formulation or kit of any one of embodiments 56-65, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 10% of said values, and having a pH of 5.8 or within +/- 10% of said value.

EE67. The dosage formulation or kit of any one of embodiments 56-65, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 5% of said values, and/or having a pH of 5.8 or within +/- 5% of said value.

EE68. The dosage formulation or kit of any one of embodiments 56-65, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 1% of said values, and/or having a pH of 5.8 or within +/- 1% of said value.

EE69. The dosage formulation or kit of any one of embodiments 56-65, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.5% of said values, and/or having a pH of 5.8 or within +/- 0.5% of said value.

EE70. The dosage formulation or kit of any one of embodiments 56-6, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 0.1% of said values, and/or having a pH of 5.8 or within +/- 0.1% of said value.

EE71. The dosage formulation or kit of any one of embodiments 66-70, wherein said L- Histidine in said formulation comprises a mixture of L-Histidine and L-Histidine monohydrate.

EE72. The dosage formulation or kit of any of embodiments 66-71, wherein said 3.1 mg of histidine in said formulation comprises a mixture of L-Histidine (1 mg) and L-Histidine hydrochloride monohydrate (2.8 mg), which in the final formulation method sums up to 3.1 mg L-histidine free base.

EE73. The dosage formulation or kit of any of embodiments 56-72, wherein said formulation is comprised in a 100 mg/mL single-dose vial wherein each mL contains 100 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

EE74. The dosage formulation or kit of any of embodiments 56-73, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine (1 mg), L-histidine hydrochloride monohydrate (2.8 mg), polysorbate 80 (0.15 mg), sorbitol (40.5 mg), and Water for Injection, USP, at a pH of 5.8.

EE75. The dosage formulation or kit of any of embodiments 56-74, wherein said formulation is comprised in a 300 mg/mL single-dose vial wherein each mL contains 300 mg anti-CGRP antibody, L-histidine, L-histidine hydrochloride monohydrate, polysorbate 80, sorbitol, and Water for Injection, USP, at a pH at or about 5.8.

[0303] EXAMPLES

[0304] The following examples are provided in order to illustrate the invention, but are not to be construed as limiting the scope of the claims in any way.

[0305] EXAMPLE 1

[0306] Preparation of Antibodies that Bind CGRP

[0307] The preparation of exemplary anti-CGRP antibodies Ab1-Ab14 having the sequences in **FIGs. 1A-12** is disclosed in owned PCT Application WO/2012/162243, published on November 29, 2012, the contents of which are incorporated by reference herein. This application exemplifies synthesis of these antibodies in *Pichia pastoris* cells. The present Applicant further contemplates synthesis of anti-CGRP antibodies Ab1-Ab14, and Ab6 in particular in CHO cells.

[0308] EXAMPLE 2

[0309] Human Clinical Study Evaluating the Safety and Efficacy of an Anti-CGRP Antibody According to the Invention

[0310] CLINICAL TREATMENT PROTOCOL

[0311] The humanized anti-CGRP IgG1 antibody identified herein as Ab6 was assessed in human subjects for its ability to inhibit, alleviate or prevent the number of, duration, and/or the intensity of migraine episodes. The Ab6 antibody contains the V_L and light chain polypeptides respectively in SEQ ID NO: 222 and SEQ ID NO: 221, and contains the V_H and heavy chain polypeptides respectively in SEQ ID NO: 202 and SEQ ID NO: 201. This antibody comprises an IgG1 constant region that contains a mutation in the heavy chain constant region (replacement of asparagine residue at position 297 with an alanine residue which substantially eliminates glycosylation and lytic activity (*see* US Patent No. 5,624,821).

[0312] Specifically, the clinical efficacy of the Ab6 antibody was tested in a placebo controlled double-blind, randomized study. The individuals in the study were all selected based on specific criteria. Particularly all were diagnosed as migraine sufferers at ≤ 50 years of age (ICHD-II, 2004 Section 1), and further had a history of migraine ≥ 12 months with ≥ 5 and ≤ 14 migraine days in each 28 day period in the 3 months prior to screening.

[0313] Further, all of the individuals in the study used acute migraine medications ≤ 14 days per 28 day period and, within those days, ≤ 10 days of triptan use per 28 day period in the 3 months prior to screening and the 28 day period of completion of eDiary prior to randomization.

[0314] Table 1 summarizes the demographic characteristics of the study population.

Table 1: Baseline Demographics and Clinical Characteristics		
Characteristic	Placebo iv (n=82)	Ab6 1000mg iv (n=81)
Mean $\pm$SD Age (years)	39.0 (9.6)	38.6 (10.8)

Mean ±SD Weight (kg)	75.4 (14.4)	75.0 (16.5)
Female Gender	66 (80%)	67 (83%)
Race:		
Caucasian	66 (80.5%)	66 (81.5%)
African American	9 (11.0%)	10 (12.4%)
Asian	3 (3.7%)	4 (5.0%)
Other	4 (4.8%)	1 (1.1%)
Baseline (per 28 days):		
Mean ±SD Migraine Days	8.8 (2.7)	8.4 (2.1)
Mean ±SD Migraine Episodes	6.7 (2.4)	6.0 (2.2)
Mean ±SD Headache Frequency	9.6 (2.8)	9.2 (2.6)
Mean ±SD Migraine Hours	72.2 (51.0)	80.1 (49.1)
Mean ±SD HIT-6 Score	64.5 (4.44)	63.8 (5.21)
Mean ±SD MSQ RFP Score	49.0 (17.9)	49.5 (21.2)
Mean ±SD MSQ RFR Score	61.9 (22.7)	63.9 (24.0)
Mean ±SD MSQ EF Score	59.5 (22.9)	59.8 (27.0)

[0315] Throughout the study all of the individuals were required to record their migraine status daily using an e-diary. In the e-diary the subjects in the study were required to record the number of migraine days/month, migraine episodes/month, migraine hours/month, migraine severity, and the use of any abortive medicine such as triptans.

[0316] In addition, the study participants were required to use the e-diary to record their migraine status in the 28 day period prior to treatment with antibody or placebo in order to establish a migraine day/hour/episode baseline per month. Also, this allowed the subjects in the study to become familiar with the use of the e-diary.

[0317] After the 28-day run-in the subjects in the study were broken into two groups, each including 80 subjects (**FIG. 17**). In the first group, i.e., the antibody treatment group, (n=80) each subject in the group was administered intravenously a single 1000 mg dose of Ab6. In the second group (n=80), i.e., the placebo group, each of the subjects was given an intravenous injection containing only the aqueous antibody carrier solution.

[0318] The individuals in the treated and placebo groups were assessed in the 24 weeks post-dose administration. Initially, a 12 week interim analysis was conducted. Subsequent to the 12 week interim analysis, a refined analysis was conducted. This refined analysis potentially included, for example, addition or removal of patient data in accord with the study protocol, e.g., updating data that had not been fully loaded from the e-diaries. This refinement resulted in slight changes but did not alter the overall conclusions.

[0319] The efficacy of the antibody versus the placebo was assessed in part based on the recorded data in the e-diary entries. For example, this analysis included a comparison of the number of recorded migraine days/month, migraine episodes/month, migraine hours/month in the subjects in the treated versus the placebo group. The percentage of responders in each group (i.e., the subjects with 50%, 75%, and 100% reduction in migraine days) in both groups was also compared.

[0320] In addition, the responses of the Ab6- and placebo-treated subjects in both groups to MSQ and HIT-6 questionnaires are to be evaluated and compared. MSQ is a frequently utilized disease-specific tool to assess the impact of migraine on health-related quality of life (HRQL). MSQ comprises a 16-item Migraine-Specific Quality-of-Life Questionnaire (Version 1.0), which was developed by Glaxo Wellcome Inc. MSQ is hypothesized to measure 3 parameters: (i) Role Function-Restrictive; (ii) Role Function-Preventive; and (iii) Emotional Function.

[0321] The HIT-6 or functional impact (also called the Headache Impact Test or HIT-6) similarly is a well known tool for assessing migraine intensity. This test uses six questions to capture the impact of headache and its treatment on an individual's functional health and well-being.

[0322] Also, the pharmacokinetic (PK) properties of the CGRP antibody and immunogenicity are to be assessed in the Ab6 antibody treated subjects.

[0323] CLINICAL RESULTS AND ANALYSIS

[0324] The results of this human clinical trial and analysis through week 12 in the treated subjects are summarized in the Table 2 below.

[0325] Table 2. Responder analysis for migraine days

Time period	% reduction migraine days	Placebo iv	Ab6 1000mg iv	P value
Week 1-4		n=80	n=75	
	50	40 (50.0)	58 (77.3)	p=0.0005
	75	19 (23.8)	39 (52.0)	p=0.0005

	100	4 (5.0)	21 (28.0)	p=0.0001
Week 5-8		n=80	n=78	
	50	43 (53.8)	59 (75.6)	p=0.0048
	75	28 (35.0)	35 (44.9)	p=0.2555
	100	12 (15.0)	21 (26.9)	p=0.0791
Week 9-12		n=77	n=72	
	50	51 (66.2)	54 (75.0)	p=0.2827
	75	24 (31.2)	38 (52.8)	p=0.0083
	100	13 (16.9)	29 (40.3)	p=0.0019

[0326] In addition, the results of the clinical study were compared based on the number of responders in the treatment and placebo groups. As shown in **FIG. 13** the number of subjects who showed a 50, 75 or 100% reduction in migraine days for each month of the interim period were compared in the treatment and placebo groups. As shown in the figure, 60% of the Ab6- treated group had at least 50 % reduction in headache days, 31% of the Ab6- treated group had at least 75 % reduction in headache days and 15 % of the Ab6 treated group had 100 % reduction in headache days.

[0327] By contrast, 33% of the placebo-treated group had at least 50 % reduction in headache days, 9% of the placebo-treated group had at least 75 % reduction in headache days, and 0 % (none) of the placebo- treated group had 100 % reduction in headache days.

[0328] These results clearly show that the reduction in the number of migraine days was much greater in the Ab6-treated group. But for the significant placebo effect, the difference in these numbers would have been more pronounced. (Elevated placebo effect is not surprising as the phenomenon is often very high for migraine and other neurological drugs).

[0329] In addition, the % change from baseline in the number of migraine days per month in the placebo and Ab6 –treated group was compared. As shown in **FIG. 14**, the median ($\pm$ QR) % change from baseline in the number of migraine days per month in the placebo and Ab6 –treated group was compared for the 2 groups during the 12 weeks post-treatment. These results which are statistically

significant ($p=0.0078$) clearly show the Ab6-treated group had a much greater reduction in the number of headache days per month compared to baseline than the placebo-treated group.

[0330] Also, the % change from baseline in the number of migraine episodes per month in the placebo and Ab6 –treated group was compared. As shown in **FIG. 15** the median ($\pm$ QR) % change from baseline in the number of migraine episodes per month in the placebo and Ab6 –treated group was compared during the 12 weeks post-treatment. These results indicate that the Ab6-treated group had a significantly greater reduction in the number of migraine episodes per month compared to baseline than the placebo-treated group.

[0331] Further, the % change from baseline in the number of migraine hours per month in the placebo and Ab6-treated group was compared. As shown in **FIG. 16**, the median ($\pm$ QR) % change from baseline in the number of migraine hours per month in the placebo and Ab6 –treated group was compared for the 2 groups during the 12 weeks post-treatment. These results clearly show the Ab6-treated group had a greater reduction in the number of migraine hours per month compared to baseline than the placebo-treated group.

[0332] In addition, the HIT-6 results were compared for both groups. As noted, this questionnaire finds well accepted usage in assessing the migraine status of individuals with frequent/chronic migraine. **FIG. 18** compares the HIT-6 responder analysis for the Ab6-treated and placebo groups at baseline, week 4 after treatment, week 8 after treatment and week 12 after treatment. The results at each time point reveal that the Ab6-treated group had a statistically significant improvement in the HIT-6 scores relative to the placebo group, i.e., 54.4% for the Ab6-treated compared to 30% for the placebo at week 4 ($p=0.0023$), 51.3% for the Ab6-treated compared to 38.0% for the placebo at week 8 ($p=0.1094$) and 61.1% for the Ab6-treated compared to 33.3% for the placebo at week 12 ($p=0.0007$). **FIG. 19** shows the percentage of patients having a HIT-6 score of some or little/none over time in the placebo and Ab6 treatment groups (statistical significance a shown).

[0333] In addition, **FIG. 20** contains the pharmacokinetic (PK) profile for Ab6 administered intravenously at a single dosage of 1000 mg in mg/mL over the 24 week period following Ab6 administration.

[0334] **FIG. 21** contains plasma-free pharmacokinetic (PK) parameters N (number of patients), mean, and standard deviation (SD) for a single 1000 mg intravenous dosage of Ab6. The parameters shown in the table and the units are $C_{\max}$ ($\mu\text{g/mL}$), $\text{AUC}_{0-\infty}$ ($\text{mg}\cdot\text{hr/mL}$), half-life (days), V_z (L) and CL (mL/hr).

[0335] Further analysis was conducted for patient data between 12-weeks and 24-weeks. The treatment group continued to exhibit decreased migraine days relative to the control group, however, the magnitude of the difference decreased over time. Additionally, the control group exhibited fewer migraine days per month than at baseline. This was thought to result at least in part from “diary fatigue” wherein patients potentially report no migraine on a day in which a migraine actually

occurred, in order to avoid the time and effort of answering further queries about the migraine that would result from giving an affirmative answer to the question of whether they had a migraine on a given day.

[0336] Further analysis of the study results are shown in **FIGs. 22-33**. These result include analysis of the change (mean $\pm$ SEM) from baseline in migraine days per month for Ab6 (1000 mg i.v.) versus placebo (**FIG. 22**), change in average migraine days ($\pm$ SD) over time for the full analysis population (**FIG. 23**). Additionally, shown are the distribution of migraine days actual and change for the Ab6 treatment group during weeks 1-4 (**FIG. 24**), distribution of migraine days actual and change for the placebo group during weeks 1-4 (**FIG. 25**), distribution of migraine days actual and change for the Ab6 treatment group during weeks 5-8 (**FIG. 26**), distribution of migraine days actual and change for the placebo group during weeks 5-8 (**FIG. 27**), distribution of migraine days actual and change for the Ab6 treatment group during weeks 9-12 (**FIG. 28**), and distribution of migraine days actual and change for the placebo group during weeks 9-12 (**FIG. 29**).

[0337] Responder rate analysis was also performed (**FIGs. 30-32**). These figures respectively show the 50%, 75%, and 100% responder rate for the Ab6 and placebo treatment groups. Subjects with $\geq 50\%$ reduction in migraine frequency were considered to be a 50% responder. Subjects with $\geq 75\%$ reduction in migraine frequency were considered to be a 75% responder. Likewise, subjects with 100% reduction in migraine frequency were considered to be a 100% responder.

[0338] In **FIGs. 22 and 30-32**, normalization was applied to visit intervals where eDiaries were completed for 21-27 days by multiplying the observed frequency by the inverse of the completion rate.

[0339] Migraine severity was also analyzed. **FIG. 33** shows the mean migraine severity over time for the full analysis population. On the scale used, a mean migraine score of 3 represents "moderate pain."

[0340] **FIG. 34** summarizes the change from baseline in migraine days, migraine episodes, migraine hours, average migraine severity, headache frequency, and outcome measures including the HIT-6 score, MSQ (Migraine Specific Quality of Life Questionnaire) RFP (Role Function-Preventative), MSQ RFR (Role Function-Restrictive), and MSQ EF (Emotional Function).

[0341] EXAMPLE 3

[0342] Human Clinical Study Evaluating the Safety and Efficacy of an Anti-CGRP Antibody in Chronic Migraine Patients

[0343] This example describes a randomized, double-blind, placebo-controlled clinical trial evaluating the safety and efficacy of Ab6 for chronic migraine prevention. In the study, 1,072 patients were randomized to receive Ab6 (300 mg or 100 mg), or placebo administered by infusion once every 12 weeks. To be eligible for the trial, patients must have experienced at least 15 headache days per month, of which at least eight met criteria for migraine. Patients that participated in the trial had an average of 16.1 migraine days per month at baseline. Study endpoints included the mean

change from baseline in monthly migraine days, reduction in migraine prevalence at day 1 and over days 1-28, and reduction of at least 50%, 75%, and 100% from baseline in mean monthly migraine days, change from baseline in mean monthly acute migraine-specific medication days, and reductions from baseline in patient-reported impact scores on the Headache Impact Test (HIT-6). The administered antibody, Ab6, is an anti-CGRP antibody consisting of the light chain polypeptide of SEQ ID NO: 221 and heavy chain polypeptide of SEQ ID NO: 201.

[0344] Patient characteristics are summarized in **FIG. 39**, with separate columns for patients receiving placebo, 100 mg of the antibody, or 300 mg of the antibody. Patients had a mean number of years from migraine diagnosis of between 17.0 and 19.0 years, a mean duration of suffering from chronic migraine of between 11.5 and 12.4 years, and between 44.3% and 45.2% of patients utilized at least one prophylactic medication. At baseline, in both antibody treatment groups the mean number of migraine days per month was 16.1, while for the placebo group, the mean number of migraine days per month was 16.2.

[0345] The reduction in a specified percentage (50%, 75%, or 100%) from baseline in mean monthly migraine days refers to the number or percentage of patients in a treatment group that exhibited the given percentage reduction in the number of migraine days per month. For example, a patient exhibiting 16 migraine days per month at baseline would be a 75% responder if the number of migraine days per month was decreased by at least 12 days per month over specified period.

[0346] The results are shown in **FIGs. 35-39**. **FIG. 35** shows the percentages of patients with migraine in the 300 mg, 100 mg, and placebo treatment groups at days 1, 7, 14, 21, and 28. The uppermost line shows results for placebo, the lowest line shows results for the 300 mg dosage, and the middle line shows results for the 100 mg dosage.

[0347] As shown in **FIG. 35**, at day 1 the percentage reduction in migraine prevalence was 52% for the 300 mg dosage, 50% at the 100 mg dosage, and 27% for placebo. The decrease was statistically significant compared to the placebo group for both the 100 mg and 300 mg treatment groups.

[0348] **FIGs. 36-38** show the percentage of patients in the 300 mg and 100 mg treatment groups achieving, respectively, 50%, 75%, and 100% reduction in migraine days in month 1, over months 1-3 (after the 1st infusion), and over months 4-5 (after the 2nd infusion). In each graph, the data bars, from left to right, show results for the 100 mg, 300 mg, and placebo groups. Statistical significance is as shown. ++ indicates a statistically significant difference from placebo; + indicates a statistically significant difference from placebo (unadjusted); and § indicates a statistically significant difference from placebo (post hoc).

[0349] EXAMPLE 4

[0350] Baseline Subgroup Analysis for Human Clinical Studies Evaluating the Safety and Efficacy of an Anti-CGRP Antibody in Chronic or Episodic Migraine Patients

[0351] In the study of Chronic Migraine described in Example 3, at intake, each patient was assessed for potential medication overuse headache (MOH). MOH was present in 39.9% (139 patients) in the 100 mg treatment group, 42.0% (147 patients) in the 300 mg treatment group, and 39.6% (145 patients) in the placebo group. Assessment of the treatment outcomes in this patient subset indicated that treatment with the anti-CGRP antibody was efficacious for MOH (**FIG. 41**). Specifically, in the 100 mg treatment group, mean migraine days per month changed by -3.0 days (95% CI, -4.56 to -1.52 days) in the patients having MOH at baseline, compared to MOH patients receiving placebo. Similarly, in the 300 mg treatment group, mean migraine days per month changed by -3.2 days (95% CI, -4.66 to -1.78 days) in the patients having MOH at baseline, compared to MOH patients receiving placebo. By contrast, for patients without MOH at baseline, in the 100 mg treatment group, mean migraine days per month changed by -1.3 days (95% CI, -2.43 to -0.16 days), compared to patients without MOH at baseline receiving placebo. Likewise, for patients without MOH at baseline in the 300 mg treatment group, mean migraine days per month changed by -2.1 days (95% CI, -3.24 to -0.88 days), compared to patients without MOH at baseline receiving placebo. Efficacy for other subgroups was shown as well, including efficacy for patients with mean migraine day (MMD) frequency less than 17 days or greater than or equal to 17 days, patients with an age at diagnosis of less than or equal to 21 years or greater than 21 years, patients having a duration of migraine of less than or equal to 15 year or greater than 15 years, patients suffering from migraine with aura or migraine with no aura, patients with prior prophylactic medication use or no prior prophylactic medication use, patients with concomitant prophylactic medication use or no concomitant prophylactic medication use, ant patients with triptan use on greater than or equal to 33% of days, or less than 33% of days. In each case, efficacy for each subgroup was shown (**FIG. 41**).

[0352] In another human clinical trial of patients with episodic migraine, patients were randomized to receive Ab6 100 mg (n=221), 300 mg (n=222), or placebo (n=222) in a double blind, parallel study. After a 28 day screening period, patients were administered the drug or placebo intravenously every 3 months for 4 total infusions (**FIG. 40**). Efficacy was shown over months 1-3 for both the 100 mg and 300 mg treatment groups, with a mean change in migraine days of -3.9 for the 100 mg treatment group and -4.3 days for the 300 mg treatment group, compared to -3.2 days for the placebo group. Efficacy for subgroups of patients was also shown, including efficacy for patients with mean migraine day (MMD) frequency less than or equal to 9 days or greater than 9 days, patients with an age at diagnosis of less than or equal to 21 years or greater than 21 years, patients having a duration of migraine of less than or equal to 15 year or greater than 15 years, and patients suffering from migraine with aura or migraine with no aura.

[0353] EXAMPLE 5

[0354] Effects of Ab6 treatment on medication use in chronic and episodic migraine patients

[0355] During the studies of chronic migraine patients described in Example 3 and episodic migraine patients described in Example 4, patients also recorded use of acute medication in a daily

eDiary and were allowed to use acute medication at their own discretion. Acute medications for migraine included ergots, triptans, and analgesics (e.g., NSAIDs, opioids, and caffeine-containing combination analgesics).

[0356] For further analysis, patients were stratified by the number of days with acute medication use during the 28-day screening period (1-9 or ≥ 10 days; “baseline”). Acute medication days were calculated for individual types of acute medications and combined, meaning that if 2 or more types of medications were used on the same calendar days, they were counted as separate medication use days. For example, if a patient took an opioid and a triptan on the same day, it counted as 2 days of acute medication use. These analyses included patients with at least 1 acute medication use day during the 28-day baseline screening period.

[0357] In both chronic migraine and episodic migraine patients who used acute medication during the 28-day baseline period, Ab6 treatment resulted in greater average reductions in monthly migraine days and acute medication days than placebo as early as Month 1 after dosing, with similar results across 2 dose intervals over 6 months.

[0358] Ab6 consistently demonstrated greater reductions in mean monthly migraine days over 6 months of treatment than placebo in chronic migraine patients taking ≥ 1 day of acute medication use during baseline (**FIG. 42**). Chronic migraine patients who had at least one day of acute medication use per month during baseline demonstrated greater decreases in acute medication use than placebo as early as month 1 after treatment and across the entire 6 month treatment period (**FIG. 43**). In the subgroup of chronic migraine patients who were taking 1–9 days of acute medication during baseline, the change from baseline in days of acute medication use was greater in the 300 mg Ab6 group than placebo across 6 months of treatment (**FIG. 44**). A clear decrease in medication days per month was observed for patients with at least 10 days of medication use per month at baseline for both Ab6 treatment group compared to placebo over the entire 6 month period. **FIG. 45** shows the changes in medication use days at Month 1 and Month 6 in the subgroups of chronic migraine patients with ≥ 1 , 1–9, and ≥ 10 days of acute medication use at baseline. With the exception of Ab6 100 mg at month 6 in patients with 1–9 days/month of use at baseline, Ab6 demonstrated a greater treatment effect in reducing acute medication use than placebo.

[0359] Similarly, across 2 dose intervals over 6 months, episodic migraine patients with one or more days of acute medication use during baseline experienced greater reductions in mean monthly migraine days with Ab6 than Placebo (**FIG. 46**). Episodic migraine patients who had at least one day of acute medication use per month during baseline demonstrated greater decreases in acute medication use than placebo as early as month 1 after treatment and across the entire 6 month treatment period (**FIG. 47**). In the subgroup of episodic migraine patients who were taking 1–9 days of acute medication during baseline, the change from baseline in days of acute medication use was greater with Ab6 than placebo across 6 months of treatment (**FIG. 48**). A similar pattern was observed in the subgroup of patients who were taking ≥ 10 days of acute medication during baseline, though smaller

sample sizes may have contributed to the less consistent pattern over time. **FIG. 49** shows the changes in medication use days at Month 1 and Month 6 in the subgroups of episodic migraine patients with ≥ 1 , 1–9, and ≥ 10 days of acute medication use at baseline. With the exception of Ab6 100 mg at Month 6 in patients with ≥ 10 days/month of use at baseline, the reduction in acute medication use was greater in the Ab6 treatment groups than placebo.

[0360] The results show that both episodic migraine and chronic migraine patients who were at risk for medication-overuse headache (≥ 10 days/month of acute medication use) demonstrated the greatest reductions in acute medication use, with Ab6 treatment generally resulting in larger decreases in medication use days than placebo.

[0361] The most frequently reported acute headache medications in $> 10\%$ of subjects included Thomapyrin N (44.5%) (a combination of paracetamol, aspirin, and caffeine), ibuprofen (40.6%), sumatriptan (33.6%), paracetamol (acetaminophen) (20.3%), and naproxen sodium (10.2%). The most frequently reported preventive headache medication in $> 10\%$ of subjects was topiramate (12.5%).

[0362] Example 6

[0363] Efficacy of anti-CGRP Antibodies in Subjects Experiencing an Acute Attack of Migraine

[0364] This example describes a randomized, double-blind, placebo-controlled clinical trial evaluating the safety and efficacy of Ab6 for the acute treatment of migraine. In the study, approximately 450 patients are randomized 1:1 to receive either 100 mg Ab6 or placebo. During a screening period (approx. 1-8 weeks) patients are assessed for migraine frequency and medication use frequency. Eligible patients have a migraine attack frequency of about 4-15 migraine days per month in the 3 months prior to screening. By history, the subject's typical migraine attack, if untreated, would be associated with headache pain of moderate to severe intensity and a most bothersome symptom of nausea, photophobia, or phonophobia. Subjects must be headache free for at least 24 hours prior to onset of a qualifying migraine in order to participate in the trial. On the day of treatment, the patient will travel to the study site and intravenous infusion of 100 mg Ab6 or placebo will commence between about 1-6 hours from the start of the attack. Patients will not have received any other monoclonal antibody (e.g., any CGRP antagonist antibody) within the 6 month period prior to screening.

[0365] Co-Primary Endpoints are time to headache pain freedom and time to absence of most bothersome symptom. Co-Key secondary are headache pain freedom at 2 hours and absence of most bothersome symptom at 2 hours. Secondary endpoints are time to headache pain relief, headache pain freedom at 2 hours with sustained headache pain freedom for 24 and 48 hours, use of rescue medication by 24 hours and by 48 hours, absence of photophobia at 2 hours, absence of phonophobia at 2 hours, absence of nausea at 2 hours, change from Baseline in Headache Impact Test (HIT 6) at Week 4, and change from Baseline in Migraine Treatment Optimization Questionnaire-6 (mTOQ-6) at Week 4. Exploratory Endpoints are absence of headache pain at all timepoints other than 2 hours, absence of photophobia at all timepoints other than 2 hours, absence of phonophobia at all timepoints

other than 2 hours, absence of nausea at all timepoints other than 2 hours, pain relapse when the subject was headache pain-free at 2 hours, patient Global Impression of Change (PGIC) at Week 4, and time to next migraine. Headache pain is collected on a 4-point scale with 3 being severe, 2 being moderate, 1 being mild, and 0 being no pain. Pain freedom is no pain (0) with the absence of rescue medication (note that in the trial rescue medication is not to be used for 2 hours post completion of infusion in order to separate the effects of the antibody from the rescue medication, however, in the course of normal use, rescue medication optionally may be used; any use of rescue medication is collected as data).

[0366] Statistical analysis is performed to determine significance of the difference in endpoints between patients receiving Ab6 or placebo, including the time to pain freedom and time to absence of most bothersome symptom, and each of the other aforementioned endpoints.

[0367] Use of rescue medication refers to any intervention (medical or device) provided to the subject to provide relief of migraine. In the study this should not be provided sooner than 2 hours following completion of the study drug administration in order to separate the effects of the antibody from the effects of said rescue medication, however, rescue medication is not contraindicated. The proportion of subjects requiring rescue medication use is summarized in the study. Acute rescue medication includes any medication to treat migraine or migraine associated symptoms, e.g., triptans, analgesics such as non-opioids or opioids/narcotics, acetaminophen, NSAIDS, combination medications such as EXCEDRIN® or EXCEDRIN MIGRAINE®, antiemetic medications, ergotamines, ergot derivatives, etc.

[0368] Absence of Migraine-Associated Symptoms (Photophobia, Phonophobia and Nausea) refers to the absence or presence of each of the aforementioned migraine-associated symptoms, as reported by the subject. The proportion of subjects absent the symptoms, with no administration of rescue medication, is summarized in the study.

[0369] Headache Impact Test (HIT-6) is assessed as the change from baseline of the total score, and is summarized and compared between treatment groups in the study.

[0370] Migraine Treatment Optimization Questionnaire-6 (mTOQ-6) is assessed as the change from baseline of the total score and is summarized and compared between the treatment groups in the study.

[0371] Time to Headache Pain Relief is assessed as the first time point post completion of infusion at which the subject reports relief of pain meaning their headache pain has gone from moderate or severe (2 or 3) to mild or no pain (1 or 0) with no administration of rescue medication.

[0372] Pain Relapse is assessed as the occurrence of headache of any severity within 48 hours of drug administration for a patient who has no headache pain (0) at 2 hours. The proportion of subjects with recurrence of headache pain of any severity is summarized in the study.

[0373] The study shows that Ab6 is effective and safe for acute migraine treatment.

[0374] Example 7

[0375] In the pivotal clinical studies the patients received Ab6 as 100 mg or 300 mg dosages, as described in Example 3. Including day -1 (post infusion of Ab6) in the statistical analysis shows that an apparent treatment effect is present immediately after infusion when the treatment effect is assessed (**FIG. 50**). In the Figure Day 0 is defined as the day of the infusion and Day -1 data represent the pre-infusion condition. A substantial decrease in the percentage of migraines from Day -1 (baseline, the day prior to infusion) to Day 0 is apparent. Moreover, the magnitude of the effect is greater with the 300 mg dosage than the 100 mg dosage, and both show a greater effect than the placebo group.

[0376] Example 8

[0377] Interventional, randomized, double-blind, parallel-group, placebo-controlled delayed-start study to evaluate the efficacy and safety of eptinezumab in patients with episodic Cluster Headache.

[0378] In a clinical study the inventors of the present invention is evaluating the efficacy of Eptinezumab (Ab6) in patients with episodic cluster headache (eCH). The target population for this study is defined as patients with eCH, based on the IHS ICHD-3 classification, with documented evidence of eCH prior to a screening visit. The study includes about 300 patients that will be randomly allocated via a randomization system to one of the two initial treatment groups: eptinezumab 400 mg, or placebo, in a ratio of 1:1. The total study duration from Screening Visit to the Safety Follow-up Visit is approximately 77 weeks and includes Screening Period (approximately 12 months / 52 weeks), Screening Period 2 (7 days), a Placebo controlled Period (4 weeks), an active Treatment Period (4 weeks), a Post-treatment Observational Period (8 weeks), and a Safety Follow-up Period (8 weeks).

[0379] The patients will enter Screening Period 2 as soon as possible after they experience the beginning of a cluster headache bout, which is characterized by the presence of at least one typical cluster headache attack, and not later than 1 week after the start of the first attack. Under exceptional circumstances, when a patient is able to attend Screening Visit 2 only during the second week after the first typical cluster headache attack, the possibility to enroll this patient in the study will be discussed with the investigator and the decision will be taken in the context of known history of typical duration of the bout for the individual patient.

[0380] Eligible patients will be randomly assigned to receive at the Baseline Visit (Day 0/Visit 3) either Eptinezumab 400 mg or placebo, administered as an IV infusion over 45 minutes (+15 minutes). Preferably the infusion should be administered in the morning. All patients will continue in the Active Treatment Period of the study and will receive a second IMP infusion (eptinezumab 400 mg or placebo), administered over 45 minutes (+15 minutes), at the end of

Week 4 (Visit 6) in a blinded manner, so that patients previously exposed to eptinezumab will receive placebo and patients

randomized to placebo will receive eptinezumab 400mg.

Evaluation (or Endpoints) will include the measurements of response by: 75% reduction in number of weekly attacks (Weeks 1-2); Change from baseline in weekly number of times an abortive therapy was used (Weeks 1-4). Time to resolution of cluster headache bout within 16 weeks, Use of abortive medication (triptans and O2) per week after the second infusion and Incidence and prevalence of preventive medication by week within 16 weeks

CLAIMS

What is claimed is:

1. A method for treatment of cluster headache (such as chronic cluster headache or episodic cluster headache) in a patient with cluster headache symptoms or for prevention of cluster headache in a patient in need of preventative treatment of cluster headache (such as chronic cluster headache or episodic cluster headache), comprising intravenous administering to a patient in need 400 mg of an anti-CGRP antibody comprising the light chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 224; SEQ ID NO: 226; and SEQ ID NO: 228, respectively and heavy chain CDR 1, 2, and 3 polypeptide sequences of SEQ ID NO: 204; SEQ ID NO: 206; and RESIDUE GDI (SEE PAGE 2, TABLE A, ITEM NO. 11), respectively.
2. The method of claim 1, wherein said patient exhibits at least one cluster headache symptom at the time of administration.
3. The method of claim 1 or 2, wherein the patient experiences at least five attacks according to the blow list A to C:
 - A. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15–180 minutes (when untreated)
 - B. 1 and/or 2 of the following:
 - (1) at least one of the following symptoms or signs, ipsilateral to the headache:
 - i. conjunctival injection and/or lacrimation
 - ii. nasal congestion and/or rhinorrhea
 - iii. eyelid oedema
 - iv. forehead and facial sweating
 - v. miosis and/or ptosis
 - (2) a sense of restlessness or agitation
 - C. Occurring with a frequency between one every other day and eight per day.
4. The method of any one of claims 1-3, wherein said patient has had or been diagnosed with cluster headache for at least 3 months, at least 6 months, at least 9 months, for at least a year, for at least 2 years, for at least 3 years, or more than 3 years.
5. The method of any one of claims 1-4, wherein said patient classifies according to the following one or more criteria: is male, is a smoker, is between the age of about 20 to about 50 years, has a family history of cluster headache, is using a medication such as nitroglycerin, or other drug used to treat heart disease, has been diagnosed with abnormalities in the body's biological clock

(hypothalamus), the cluster headache is typically not associated with triggers, such as foods, hormonal changes or stress, or upon the onset of a cluster period alcohol use exacerbates or triggers more headaches or increased head pain.

6. The method of any one of claims 1-5, wherein said patient experiences head pain.

7. The method of any one of claims 1-6, wherein the cluster headache sign or symptom is alleviated immediately after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

8. The method of any one of claims 1-7, wherein said patient no longer has a cluster headache after said administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

9. The method of any one of the foregoing claims, wherein said anti-CGRP antibody comprises the light chain polypeptide of SEQ ID NO: 221 and the heavy chain polypeptide of SEQ ID NO: 201 or SEQ ID NO: 566.

10. The method of any one of the foregoing claims, wherein said anti-CGRP antibody comprises the light chain polypeptide encoded by SEQ ID NO: 231 and the heavy chain polypeptide encoded by SEQ ID NO: 211 or SEQ ID NO: 567.

11. The method of any one of the foregoing claims, wherein the cluster headache signs or symptoms decline or are abolished immediately after administration, such as within the first day after administration, within 12 hours after administration, within 6 hours after administration within 5 hours after administration, within 4 hours after administration, within 3 hours after administration, within 2 hours after administration, or within 1 hour of after administration, within 30 minutes after administration, or such as between 1-6 hours after administration.

12. The method of any one of the foregoing claims, wherein said patient is cluster headache free 2 hours post-completion of infusion.

13. The method of any one of claims 1-23, comprising intravenously administering 400 mg of said anti-CGRP antibody every 10-14 weeks, preferably every 11-13 weeks, more preferably every 12 weeks.
14. The method of any one of the foregoing claims, wherein said anti-CGRP antibody is comprised in a formulation comprising or consisting of histidine (L-histidine), sorbitol, polysorbate 80, and water.
15. The method of claim 26, wherein said formulation comprises or consists of, per 1 mL volume, 100 mg anti-CGRP antibody, 3.1 mg L-Histidine, 40.5 mg Sorbitol, and 0.15 mg Polysorbate 80, or having amounts of each constituent within +/- 10% of said values, and having a pH of 5.8 or within +/- 10% of said value.