


AMINO QUINAZOLINES DERIVATIVES WITH BLOOD PLATELET REDUCING PROPERTIES

FIELD OF THE INVENTION

This invention relates to the discovery of prodrugs of 3-or 5-substituted analogues of the selective platelet lowering agent anagrelide which have reduced potential for cardiovascular side-effects and which should therefore lead to improved patient compliance and safety in the treatment of myeloproliferative diseases. More specifically, the present invention relates to prodrugs of certain imidazoquinazoline derivatives which have utility as platelet lowering agents in humans. The compounds of the present invention function by inhibiting the formation of blood platelets.

BACKGROUND OF THE INVENTION

Anagrelide hydrochloride (Agrylin®, Xagrid®) is a novel orally administered imidazoquinazoline which selectively reduces platelet count in humans and is used for such purposes in the treatment of myeloproliferative diseases (MPDs), such as essential thrombocythemia (ET), where an elevated platelet count may put the patient at increased thrombotic risk. The chemical structure of anagrelide, 6,7-dichloro-1,5-dihydroimidazo[2,1-*b*]-quinazolin-2(3*H*)-one (hydrochloride monohydrate), is shown in the following formula:



6,7-Dichloro-1,5-dihydroimidazo[2,1-*b*]-quinazolin-2(3*H*)-one hydrochloride monohydrate

The principal side effects of anagrelide are cardiovascular in nature, tachycardia, palpitations etc and limit the utility of the drug. These are largely attributed to its metabolism to 3-hydroxyanagrelide. This compound was surprisingly found to be some 40-fold more potent as an inhibitor of PDEIII, and therefore potential inotropic agent, than anagrelide itself. Furthermore plasma exposure to this metabolite after treatment with anagrelide is typically three times greater

than to the drug itself confirming its pivotal role. Consequently a series of 3-substituted anagrelide analogues has been investigated which has shown that it is possible to introduce metabolism blocking groups at that position and yet still retain the anti-megakaryocytic actions of the drug. Furthermore, these compounds have considerably less potential for cardiovascular effects than 3-hydroxyanagrelide. Indirect steric hindrance to the formation of the 3-hydroxy metabolite may also be achieved by substitution at the 5-position. The 3-substituted compounds typified, by the dimethyl or spirocyclopropyl analogues, are notably less soluble at physiological pH (~7) than the parent compound which presents a significant challenge to their efficient absorption.

Anagrelide HCl itself is a poorly soluble drug substance. In the pH range of 4 – 8, the solubility is less than 10 µg/mL. The solubility increases at pH values above and below this range; for example in 0.1M HCl the solubility is ~170 µg/mL and at pH 11.4 approaches 1mg/mL. The dissociation constants (pKa1 and pKa2) of 2.9 and 9.8 were estimated from the solubility/pH profile of anagrelide HCl. Thus over much of the physiological pH range the drug has very poor aqueous solubility. Even material dissolving in the stomach at pH 1-2 may precipitate in the duodenum at pH 5-6. This presents potential problems for the quantitative absorption from the most likely site for absorption, namely the upper small intestine. As a consequence anagrelide is micronised prior to filling into capsules for clinical use to ensure maximal absorption. This and anagrelide's inherent potency as an anti-megakaryocytic agent – *in vitro* IC₅₀ ~27 nM and *in vivo* doses of just 1-2 mg – serve to limit the potential problem of incomplete absorption. Indeed, a radiolabelling study in humans showed that following the oral administration of 1 mg ¹⁴C-labelled drug >75% of the administered radioactivity was recovered in the urine implying that at least after this dose absorption was >75%. However for those patients requiring larger doses either as the result of relative insensitivity to the drug, higher first-pass pre-systemic metabolism or greater body weight, the possibility exists for incomplete absorption. This would be expected to lead to increased variability in attained plasma drug concentrations and consequential variability in patient response.

Furthermore for less potent analogues of anagrelide, even though they may have better cardiovascular profiles, problems of incomplete absorption may be encountered. Improved water solubility though the use appropriate open ring pro-drugs of anagrelide may therefore offer significant advantage in minimizing this risk.

5

WO2004/063172 relates to the use of 2-amino-2H-quinazoline derivatives for producing therapeutic agents for the treatment of myeloproliferative diseases, high blood pressure and bronchodilation.

10

SUMMARY


This disclosure provides for prodrugs of anagrelide derivatives substituted at either the 3- or 5-position. In these anagrelide derivatives, metabolism to an analogue of the cardioactive 3-hydroxyanagrelide is blocked either directly (3-substitution) or indirectly (5-substitution). The prodrugs are notably more soluble *in vitro* (and under anticipated *in vivo* conditions) than their ring closed analogues offering the potential for better absorption from the GI tract. Such compounds would spontaneously and completely ring close at pH 7 or above thus offering a convenient means of delivering these ring closed anti-megakaryocytic (platelet lowering) agents to the systemic circulation. Since the preferred site of metabolism of anagrelide is the 3-position, such compounds are likely to present improved pharmacokinetic profile and hence improve patient compliance and convenience enabling a broader spectrum of patients to be effectively treated. In the case of the 5-substituted compounds it is expected that a bulky group is more effective than a smaller group when cyclised to the 'closed ring' anagrelide analogue. Groups such as t-butyl and other bulky blocking groups are thus expected to be of most utility when substituted at the 5-position. A substituent comprising a large group at the 5-position is expected to sterically hinder access to the 3-position by the metabolising cytochrome's active site. This should inhibit formation of the cardioactive metabolite, 3-hydroxyanagrelide.

The ring closed compounds of the present invention are especially beneficial because surprisingly they have dramatically lower PDE III inhibitory activity (and hence lower

30

cardioactive potential) than the active metabolite of anagrelide, 3-hydroxyanagrelide and yet also surprisingly retain their anti-megakaryocytic activity. Indeed these compounds have therapeutic indices which are much more favourable than that for anagrelide itself.

In a first aspect, there is provided a compound of Formula (I) or a pharmaceutically acceptable salt or solvate thereof



(I)

wherein:

R^1 and R^2 , are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} alkylsulphonyl and COOH; C_{1-6} hydroxyalkyl; C_{1-6} carboxyalkyl; and sulphide;

or wherein R_1 and R_2 together with the carbon to which they are attached form a C_{3-8} carbocyclic ring may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} haloalkyl, C_{1-4} alkylsulphonyl and COOH;

or wherein R_1 and R_2 together with the carbon to which they are attached represent a C_{2-6} alkenyl or C_{2-6} alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C_{1-4} haloalkyl and COOH, provided always that one of R^1 and R^2 is not hydroxyl when the other is methyl;

R^3 and R^4 are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} alkylsulphonyl and COOH; C_{1-6} hydroxyalkyl; C_{1-6} carboxyalkyl; and sulphide;

or R^3 and R^4 together with the carbon to which they are attached form a C_{3-8} carbocyclic ring which may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} haloalkyl, C_{1-4} alkylsulphonyl and COOH;

or R^3 and R^4 together represent a C_{2-6} alkenyl or C_{2-6} alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C_{1-4} haloalkyl and COOH;

R^5 is selected from the group comprising: fluoro, chloro, bromo and iodo;

R^6 is selected from the group comprising: fluoro, chloro, bromo and iodo; and

R^7 and R^8 are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

R^9 is H or C_{1-6} alkyl,

R^{10} is selected from the group comprising: hydrogen; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl and C_{3-8} cycloalkyl wherein each of the foregoing groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} alkylsulphonyl and $COOH$; or R^{10} is a pharmaceutically acceptable counterion, and

X is O or S,

provided always that R^1 , R^2 , R^3 and R^4 are not all hydrogen.

In a second aspect, there is provided a pharmaceutical composition comprising a compound of formula (I) as defined above, or a pharmaceutically acceptable salt or solvate thereof, together with a pharmaceutically acceptable diluent or carrier, which may be adapted for oral, parenteral or topical administration.


In a third aspect, there is provided a compound of formula (I) as defined above, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition containing any of the foregoing, for use as a medicament.

In a fourth aspect, there is provided the use of a compound of formula (I) as defined above, or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment of a disease selected from: myeloproliferic diseases and generalized thrombotic diseases.

In a fifth aspect, there is provided the use of a compound of formula (I) as defined above in the manufacture of a medicament for the reduction of platelet count.

In one embodiment, the present disclosure comprises a prodrug of an anagrelide analogue comprising a 3-, 5-, 3, 3- or 5,5-substituted anagrelide compound. Thus, for example, in the 3-substituted derivatives, first pass metabolism (of the rapidly ring closed analogue) to 3-hydroxyanagrelide is directly blocked. In particular, the disclosure relates to prodrugs of an anagrelide analogue wherein first pass metabolism to the corresponding analogue of 3-hydroxyanagrelide is effectively blocked.

According to the present disclosure, there is provided a compound of Formula (I) or a pharmaceutically acceptable salt or solvate thereof



wherein:

R^1 , R^2 , R^3 and R^4 independently represent hydrogen or a blocking group which functions to prevent metabolic reaction either directly or indirectly at the 3- position;

or R^1 and R^2 , and/or R^3 and R^4 together with the carbon to which they are attached form a blocking group which functions to prevent metabolic reaction either directly or indirectly at the 3- or 5-position of substitution, the remainder of groups R^1 to R^4 being hydrogen;

R^5 is selected from the group comprising: fluoro, chloro, bromo and iodo;

R⁶ is selected from the group comprising: fluoro, chloro, bromo and iodo; and

R⁷ and R⁸ are independently selected from the group comprising: H; halo; cyano; C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₁₋₆ alkoxy, and C₁₋₆ haloalkoxy; and

5

R⁹ is H or C₁₋₆ alkyl,

R¹⁰ is selected from the group comprising: hydrogen; C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl and C₃₋₈ cycloalkyl wherein each of the foregoing groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C₁₋₄ alkylsulphonyl and COOH; or R¹⁰ is a pharmaceutically acceptable cation, and

10

X is O or S,

15 provided always that R¹, R², R³ and R⁴ are not all hydrogen.

In an embodiment when one of R¹ and R² is methyl and R³ and R⁴ are hydrogen then other of R¹ and R² is not hydrogen.

20 In one embodiment:

R¹ and R², are independently selected from the group comprising: H; halo; cyano; C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₃₋₈ cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C₁₋₄ alkylsulphonyl and COOH; C₁₋₆ hydroxyalkyl; C₁₋₆ carboxyalkyl; and sulphide;

25

or R¹ and R² together with the carbon to which they are attached form a C₃₋₈ carbocyclic ring may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C₁₋₄ haloalkyl, C₁₋₄ alkylsulphonyl and COOH;

30

or R^1 and R^2 together with the carbon to which they are attached represent a C_{2-6} alkenyl or C_{2-6} alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C_{1-4} haloalkyl and $COOH$, provided always that one of R^1 and R^2 is not hydroxyl when the other is methyl.

In a preferred set of compounds, R^1 is an optionally substituted C_{1-4} alkyl or C_{3-8} cycloalkyl group.

10 In a preferred set of compounds, R^2 is an optionally substituted C_{1-4} alkyl or C_{3-8} cycloalkyl group.

Other preferred compounds are those in which at least one of R^1 and R^2 is $-C(H)_n(F)_m$ or $-C(H)_n(F)_m-C(H)_p(F)_q$, where $m = 2$ or 3 , and $n = (3-m)$; and $p = 2$ or 3 , and $q = (3-p)$.

15 More preferably at least one of R^1 and R^2 is CF_3 or CHF_2 . Most preferably, at least one of R^1 and R^2 is CF_3 .

20 In an embodiment, R^1 is preferably methyl, cyclopropyl, CF_3 or CHF_2 . Most preferably, R^1 is methyl.

In an embodiment, R^2 is preferably methyl, cyclopropyl, CF_3 or CHF_2 . Most preferably R^2 is methyl.

25 In another preferred set of compounds, R^1 and R^2 together form an optionally substituted C_{3-8} cycloalkyl group. Most preferably this is a cyclopropyl group.

In an embodiment:

30 R^3 and R^4 are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups

may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C₁₋₄ alkylsulphonyl and COOH; C₁₋₆ hydroxyalkyl; C₁₋₆ carboxyalkyl; and sulphide;

- 5 or R³ and R⁴ together with the carbon to which they are attached form a C₃₋₈ carbocyclic ring may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C₁₋₄ haloalkyl, C₁₋₄ alkylsulphonyl and COOH;

- 10 or R³ and R⁴ together represent a C₂₋₆ alkenyl or C₂₋₆ alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C₁₋₄ haloalkyl and COOH.

- 15 In an embodiment, R³ is H or C₁₋₆ alkyl. Preferably, R³ is H.

In an embodiment, R⁴ is H or C₁₋₆ alkyl. Preferably, R⁴ is H.

In an embodiment, R⁵ is chloro.

- 20 In an embodiment, R⁶ is chloro.

In an embodiment, R⁷ is H.

In an embodiment, R⁸ is H.

- 25 In an embodiment, R⁹ is H or Me. In one embodiment, R⁹ is H and compounds in which R⁹ is H enjoy good solubility. When R⁹ is a C₁₋₆ alkyl group, such as Me, the PDE III inhibiting activity is effectively eliminated. Me represents a particularly preferred alkyl substituent.

- 30 In an embodiment, R¹⁰ is H or optionally substituted C₁₋₆ alkyl. Most preferably, R¹⁰ is C₁₋₆ alkyl. In an alternative embodiment, R¹⁰ is Na or K, with Na being preferred.

In an embodiment, X is O.

In a further embodiment,

5

R^1 and R^2 are independently selected from the group comprising: H; cyano; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} alkylsulphonyl and COOH; C_{1-6} hydroxyalkyl; C_{1-6} carboxyalkyl; and sulphide;

10

or R^1 and R^2 together with the carbon to which they are attached form a C_{3-8} carbocyclic ring may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} haloalkyl, C_{1-4} alkylsulphonyl and COOH;

15

or R^1 and R^2 together represent a C_{2-6} alkenyl or C_{2-6} alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C_{1-4} haloalkyl and COOH;

20

R^3 and R^4 are hydrogen;

R^5 is selected from the group comprising: fluoro, chloro, bromo and iodo;

25 R^6 is selected from the group comprising: fluoro, chloro, bromo and iodo; and

R^7 , R^8 , R^9 and R^{10} are hydrogen.

Another preferred group of compounds is those in which neither R^1 nor R^2 is hydrogen.

30 Amongst these, it is preferred when R^1 and R^2 are both independently selected from the group


comprising: cyano, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, in which the alkyl, alkenyl, and alkynyl groups may be optionally substituted;

or wherein R₁ and R₂ together with the carbon to which they are attached form an optionally substituted C₃₋₈ carbocyclic ring


or wherein R₁ and R₂ together represent an optionally substituted C₂₋₆ alkenyl or C₂₋₆ alkynyl group.

Particularly preferred individual compounds of the disclosure include:


The 3-methylanagrelide derivative



3,3-dimethylanagrelide derivative



and the corresponding spiro 3-cyclopropyl derivative



Regarding the use of the compounds of the disclosure in humans, there is provided:

- a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, together with a pharmaceutically acceptable diluent or carrier,
5 which may be adapted for oral, parenteral or topical administration;
- a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition containing any of the foregoing, for use as a medicament;
- 10 the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment of a disease selected from: myeloproliferic diseases and generalised thrombotic diseases.

- a method of treating a disease selected from: myeloproliferic diseases and generalised thrombotic
15 diseases in a human, which comprises treating said human with an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, or with a pharmaceutical composition containing any of the foregoing.

- The present disclosure also encompasses a method of treating a patient having essential
20 thrombocythemia or other myeloproliferative disease or thrombotic cardiovascular disease or high blood platelet count, which method comprises administering to the patient a therapeutically effective amount of a compound of the present disclosure.

- Another embodiment of the present disclosure includes a method of reducing blood platelet count
25 within a patient, which method comprises administering to the patient a therapeutically effective amount of a compound of the present disclosure.

- The present disclosure encompasses providing the compounds of the present disclosure for the methods listed above, among others, wherein cardiotoxicity is reduced compared to using
30 anagrelide.

The present disclosure also encompasses pharmaceutical compositions comprising a compound or pharmaceutically acceptable salt of a compound of the present disclosure and a pharmaceutically acceptable carrier.

- 5 Pharmaceutically acceptable salts of the compounds of Formula (I) include acid addition salts. Examples include hydrochloric and hydrobromide salts.

DETAILED DESCRIPTION OF THE INVENTION

10

The present invention is directed to new prodrugs of 3-or 5-substituted analogues of the established platelet lowering agent anagrelide. These compounds spontaneously ring close at pH's 7 and above to yield 3-or 5-substituted anagrelides that retain the anti-megakaryocytic properties (hence platelet lowering activity) of anagrelide but have reduced PDEIII inhibitory properties and hence lower potential for unwanted cardiovascular and anti-aggregatory side-effects.

15

Appropriate substitution at the 3-position of the anagrelide molecule effectively blocks the principal site of metabolism and thus precludes the formation of the highly potent PDEIII inhibitor 3-OH anagrelide. The 5-substituted analogues have the potential to indirectly sterically hinder metabolism at the preferred 3-position. These 3-or 5-substituted analogues of anagrelide also have the potential for improved pharmacokinetic characteristics since the 3-position in the anagrelide molecule is known to be the major site of metabolism which is the principal mechanism of drug clearance.

20

25

Use of the corresponding "open ring" prodrugs of these 3- or 5-substituted analogues could offer the added value of improved rates of dissolution and water solubility, allowing easier formulation. For example the aqueous solubility of anagrelide at pH 7 is <10ug/ml. For ethyl-5,6-dichloro-3,4-dihydro-2-(1H)-iminoquinazoline-3-acetate HBr - an unsubstituted but representative example of these ring open prodrugs - the solubility is ~ 5.5mg/ml in distilled water.

30

Such prodrugs are likely to be extremely rapidly and completely cyclised in plasma to the closed ring 3-alkylanagrelide analogues. For example the rapid and quantitative conversion of ethyl-5,6-dichloro-3,4-dihydro-2-(1H)-iminoquinazoline-3-acetate HBr - an unsubstituted but representative example of these ring open prodrugs - to anagrelide was demonstrated in human plasma using LC/MS-MS analytical techniques. Human plasma was spiked with anagrelide prodrug (final concentration 100 ng/mL). Immediately after mixing, and at 15, 30, 45 and 60 minutes afterward samples were analysed for anagrelide prodrug and anagrelide. Even at the first point of measurement no prodrug could be found demonstrating the rapid and complete conversion to anagrelide itself. Figure 1 shows the levels of anagrelide prodrug, ethyl-5,6-dichloro-3,4-dihydro-2-(1H)-iminoquinazoline-3 acetate and anagrelide observed in samples of human plasma, incubated at room temperature over one hour.

The potential benefit of improved water solubility on the absorption of these open-ring analogues was shown in a comparative bioavailability study in the dog. Using the unsubstituted ethyl-5,6-dichloro-3,4-dihydro-2-(1H)-iminoquinazoline-3-acetate HBr as a model compound, a comparison was made of the systemic availability of anagrelide when given as this compound or as anagrelide itself in equimolar doses (7.7 & 6.1mg/kg respectively). Examination of pharmacokinetic parameters for the prodrug showed an approximately 17-fold higher C_{max} , and a mean 16-fold higher AUC for anagrelide than when the drug itself was administered.

20

These results implied that the inherent absorption of anagrelide at this dose (6.1mg/kg, albeit 200 fold above the clinical dose) was comparatively poor (< 6.25 %) since there was little evidence for marked changes in metabolism, the likely alternative explanation. The metabolite-to-drug exposure ratio after anagrelide was 1.5 compared to 0.9 after the prodrug.

25

This study (see tables below) also showed that there was also considerably less variability in C_{max} and AUC after the prodrug. For example C_{max} for anagrelide after the prodrug ranged from 170 – 418 ngmL⁻¹ (relative standard deviation, RSD, 26%) compared to 9.5 to 44.3 ngmL⁻¹ after anagrelide itself (RSD 62.5%). Similarly the AUC for anagrelide after the prodrug ranged from 367 to 1470 ng.hmL⁻¹ (RSD 34%) compared to 21.6 to 188ng.hmL⁻¹(RSD 71%) after anagrelide

30

itself. The lesser variability was consistent with more efficient absorption. This study illustrated the potential benefits of the open-ring prodrugs to improve absorption.

Table 1

5 Pharmacokinetic parameters of anagrelide following a single oral (capsule) administration of anagrelide or an ester open ring prodrug of anagrelide to male dogs at equivalent molar doses

Anagrelide (6.1 mg/kg)

Dog ID number	C _{max} (ng/mL)	T _{max} (hours)	AUC _{0-t} (ng.h/mL)	AUC _{inf} (ng.h/mL)	k (hours ⁻¹)	t _{1/2} (hours)
1	15.7	16	141	-	e	-
3	14.8	1.5	42.0	42.3 ^c	0.4459 ^c	1.6 ^c
11	25.0	2	188	193 ^c	0.3119 ^c	2.2 ^c
23	9.50	1.5	21.6	23.1 ^d	0.1953 ^d	3.5 ^d
29	44.3	1	88.9	89.3	0.3031	2.3
Mean	21.9	1.5 ^b	96.3	-	-	-
SD	13.7		68.9	-	-	

Ester prodrug of anagrelide (7.5 mg/kg)

Dog ID number	C _{max} (ng/mL)	T _{max} (hours)	AUC _{0-t} (ng.h/mL)	AUC _{inf} (ng.h/mL)	k (hours ⁻¹)	t _{1/2} (hours)
1	213	3	678	679	0.1969	3.5
3 ^a	170	1	367	369	0.3071	2.3
11	418	4	1440	1440 ^d	0.1789 ^d	3.9 ^d
23	334	3	951	952	0.4941	1.4
29	353	6	1470	1470	0.4857	1.4
Mean	330	3.5 ^b	1130	1030	0.3922	1.8 ^f
SD	86		390	400	0.1692	

^a Animal vomited *ca* 1 hour post-dose, excluded from calculation of mean

^b Median

^c Estimate based on two data points only, therefore did not meet acceptance criteria, excluded from calculation of mean

^d Could not be estimated in accordance with all acceptance criteria, excluded from calculation of mean

^e Could not be estimated from the available data

^f Calculated as $\ln 2 / (\text{mean rate constant})$

Table 2

Pharmacokinetic parameters of 3-hydroxy anagrelide, following a single oral (capsule) administration of anagrelide or an ester prodrug of anagrelide to male dogs at equivalent molar doses

Anagrelide (6.1 mg/kg)

Dog ID number	C _{max} (ng/mL)	T _{max} (hours)	AUC _{0-t} (ng.h/mL)	AUC _{infin} (ng.h/mL)	k (hours ⁻¹)	t _{1/2} (hours)
1	14.1	16	131	-	d	-
3	18.0	1.5	64.6	65.0	0.2854	2.4
11	29.9	16	274	-	d	-
23	19.4	1.5	50.7	51.8	0.2314	3.0
29	43.0	1.5	122	123	0.2966	2.3
Mean	24.9	1.5 ^b	128	79.9	0.2711	2.6 ^e
SD	11.7		89	37.9	0.0349	

Ester prodrug of anagrelide (7.5 mg/kg)

Dog ID number	C _{max} (ng/mL)	T _{max} (hours)	AUC _{0-t} (ng.h/mL)	AUC _{infin} (ng.h/mL)	k (hours ⁻¹)	t _{1/2} (hours)
1	185	3	564	566	0.1569	4.4
3 ^a	106	1.5	303	303	0.2510	2.8
11	347	4	1280	1290 ^c	0.1235 ^c	5.6 ^c
23	269	3	876	878	0.4425	1.6
29	241	6	1240	1240	0.3776	1.8
Mean	261	3.5 ^b	990	895	0.3257	2.1 ^e
SD	67		337	337	0.1497	

^a Animal vomited *ca* 1 hour post-dose, excluded from calculation of mean

^b Median

^c Could not be estimated in accordance with all acceptance criteria (excluded from calculation of mean)

^d Could not be estimated from the available data

^e Calculated as $\ln 2 / (\text{mean rate constant})$

For those 3- or 5-substituted anagrelide analogues which have a lower therapeutic potency (but not inherent activity) than anagrelide itself, a potentially higher absolute dose may be needed which could present problems for absorption. For example 3,3, dimethyl anagrelide (anti-megakaryocytic IC₅₀ ~160nM cf 27nM for anagrelide) may need to be given at 6 times the current clinical dose of anagrelide. In this situation absorption may be less than complete and a prodrug may be needed to ensure efficient absorption from the GI tract.

The compounds of Formula I can be prepared in an analogous manner to those described in US 4256748 and US 6388073. The disclosures of the synthetic procedures used in each of these documents is intended specifically to be incorporated into this disclosure and forms part of the disclosure of this invention. The contents are not presented here in full for the purposes of brevity but the skilled person is specifically directed to these documents.

A person skilled in the art will be aware of variations of, and alternatives to, the processes referred to in US 4256748 which allow the individual compounds defined by formula (I) to be obtained having been now revealed as desirable targets. The present invention thus further encompasses methods of manufacturing a compound of the present invention to the extent that such processes produce novel intermediates and / or employ novel process features.

It will also be appreciated by a person skilled in the art that the compounds of the invention could be made by adaptation of the methods herein described and/or adaptation of methods known in the art, for example the art described herein, or using standard textbooks such as "Comprehensive Organic Transformations - A Guide to Functional Group Transformations", RC Larock, Wiley-VCH (1999 or later editions), "March's Advanced Organic Chemistry - Reactions, Mechanisms and Structure", MB Smith, J. March, Wiley, (5th edition or later) "Advanced Organic Chemistry, Part B, Reactions and Synthesis", FA Carey, RJ Sundberg, Kluwer Academic/Plenum Publications, (2001 or later editions), "Organic Synthesis - The Disconnection Approach", S Warren (Wiley), (1982 or later editions), "Designing Organic Syntheses" S Warren (Wiley) (1983 or later editions), "Guidebook To Organic Synthesis" RK Mackie and DM Smith (Longman) (1982 or later editions), etc., and the references therein as a guide.

It will also be apparent to a person skilled in the art that sensitive functional groups may need to be protected and deprotected during synthesis of a compound of the invention. This may be achieved by conventional methods, for example as described in "Protective Groups in Organic Synthesis" by TW Greene and PGM Wuts, John Wiley & Sons Inc (1999), and references therein.

Definitions:

Halo means a group selected from: fluoro, chloro, bromo or iodo.

5

The term "alkyl" as used herein as a group or a part of a group refers to a straight or branched hydrocarbon chain containing the specified number of carbon atoms. For example, C₁₋₁₀ alkyl means a straight or branched alkyl containing at least 1 and at most 10 carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, isobutyl, isopropyl, t-butyl, hexyl, heptyl, octyl, nonyl and decyl. A C₁₋₄ alkyl group is one embodiment, for example methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl or t-butyl.

10

The term "cycloalkyl" as used herein refers to a non-aromatic monocyclic hydrocarbon ring of 3 to 8 carbon atoms such as, for example, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

15

The term "spirocyclic" as used herein refers to a ring system joined to a second ring system at one carbon atom.

The term "alkoxy" as used herein refers to a straight or branched hydrocarbon chain group containing oxygen and the specified number of carbon atoms. For example, C₁₋₆ alkoxy means a straight or branched alkoxy containing at least 1 and at most 6 carbon atoms. Examples of "alkoxy" as used herein include, but are not limited to, methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy, 2-methylprop-1-oxy, 2-methylprop-2-oxy, pentoxy and hexyloxy. A C₁₋₄ alkoxy group is one embodiment, for example methoxy, ethoxy, propoxy, prop-2-oxy, butoxy, but-2-oxy or 2-methylprop-2-oxy.

20

25

The term "hydroxyalkyl" as used herein as a group refers to a straight or branched hydrocarbon chain containing the specified number of carbon atoms, which is substituted by 1-3 hydroxyl groups. For example, C₁₋₄ hydroxyalkyl means a straight or branched alkyl chain containing from

30

1 to 4 carbon atoms and at least one hydroxyl group; examples of such group include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxyisopropyl, hydroxybutyl and the like.

5 The term "alkenyl" as used herein as a group or a part of a group refers to a straight or branched hydrocarbon chain containing the specified number of carbon atoms and containing at least one double bond. For example, the term " C_{2-6} alkenyl" means a straight or branched alkenyl containing at least 2 and at most 6 carbon atoms and containing at least one double bond. Examples of "alkenyl" as used herein include, but are not limited to, ethenyl, 2-propenyl, 3-butenyl, 2-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, 3-methylbut-2-enyl, 3-hexenyl
10 and 1,1-dimethylbut-2-enyl. It will be appreciated that in groups of the form $-O-C_{2-6}$ alkenyl, the double bond is preferably not adjacent to the oxygen.

The term "alkynyl" as used herein as a group or a part of a group refers to a straight or branched hydrocarbon chain containing the specified number of carbon atoms and containing at least one
15 triple bond. For example, the term " C_{2-6} alkynyl" means a straight or branched alkynyl containing at least 2 and at most 6 carbon atoms and containing at least one triple bond. Examples of "alkynyl" as used herein include, but are not limited to, ethynyl, 2-propynyl, 3-butyne, 2-butyne, 2-pentyne, 3-pentyne, 3-methyl-2-butyne, 3-methylbut-2-ynyl, 3-hexynyl and 1,1-dimethylbut-2-ynyl. It will be appreciated that in groups of the form
20 $-O-C_{2-6}$ alkynyl, the triple bond is preferably not adjacent to the oxygen. The term "halo" refers to halogens such as fluorine, chlorine, bromine or iodine atoms.

The term "sulfide" refers to a radical of R_a-S-R_b , wherein a sulfur atom is covalently attached to two hydrocarbon chains, R_a and R_b , wherein the two hydrocarbon chains may be, for example,
25 but not limited to, any discussed above.

The compounds of the invention, i.e. those of formula (I), when cyclised may possess antimegakaryocytic activity in humans. Such activity may be assessed using a well established model. Assessment of the *in vitro* anti-megakaryocytic activity - and potentially therefore the
30 platelet lowering capability - of the anagrelide prodrugs can be determined using the model of megakaryocytopoiesis (Cohen-Solal *et al.*, Thromb. Haemost. 1997, 78:37-41 and Cramer *et al.*,

Blood, 1997, 89:2336-46). This involves examining the differentiation of human CD34⁺ stem cells into megakaryocytes which ultimately give rise to blood platelets.

5 The compounds of the invention may be particularly useful in the treatment of myeloproliferative diseases. The compounds may also find utility in the treatment of generalised thrombotic diseases.

10 It is to be appreciated that references to treatment include prophylaxis as well as the alleviation and/or cure of established symptoms of a condition. "Treating" or "treatment" of a state, disorder or condition includes: (1) preventing or delaying the appearance of clinical symptoms of the state, disorder or condition developing in a human that may be afflicted with or predisposed to the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition, (2) inhibiting the state, disorder or condition, *i.e.*, arresting, reducing or delaying the development of the disease or a relapse thereof (in case of
15 maintenance treatment) or at least one clinical or subclinical symptom thereof, or (3) relieving or attenuating the disease, *i.e.*, causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms.

20 Myeloproliferative diseases which may be treatable with the compounds of the present invention include: essential thrombocythemia, polycythemia vera, chronic idiopathic myelofibrosis, chronic myeloid leukaemia with residual thrombocytosis, reactive thrombocytosis immediately preceding a surgical procedures, as an immediate or post operative preventative measures to minimise the risk of thrombus formation during or post surgery.

25 Thrombotic cardiovascular diseases (TCVD) (*i.e.* patients at increased generalised thrombotic risk) which may be treatable with the compounds of the present invention include: myocardial infarct (heart attack) thrombotic stroke, patients having undergone coronary stent placement.


30 The compounds of the present invention may also find utility in indicated for the reduction of atherothrombotic events as follows: recent MI, recent stroke or established peripheral arterial

disease, acute coronary syndrome (unstable angina/non-Qwave MI), cardiovascular death, MI, stroke, and refractory ischemia.

Example 1

- 5 The spirocyclopropylanagrelide derivative can be prepared using established chemistry as described previously according to the following reaction scheme. Other derivatives can be made in a likewise manner. The final product is obtained by cyclisation of the cyclopropyl precursor in the presence of cyanogen bromide and ethanol.

10








Example 2

- 15 The solubilities of various salts was investigated in three solvents [distilled water (pH 5.0), 50 mM ammonium formate (pH 7.9) and 0.1 M hydrochloric acid (pH 0.6)]. An excess of the salt was suspended in the chosen medium and sonicated for 3 minutes. The mixture was centrifuged for 10 minutes, the supernatant liquid was collected and the solubility was measured. Table 3 shows the results of the measurements.

20

25


Solubilities of 2,2-Disubstituted 2-(2-Amino-5,6-dichloro-4*H*-quinazolin-3-yl)-propionates ($\mu\text{g mL}^{-1}$ at ca. 20 °C)

Compound	Ammonium formate (50 mM, pH 7.9)	Water (pH 5.0)	Hydrochloric acid (0.1 M, pH 0.6)
	4	20	215
	16	7340	46
	1	2	35
	unstable	7060	6430
	11	10	169

- 5 The figures show that the compounds of the invention have substantially improved solubility over their corresponding closed ring counterparts. This means that the compounds can be formulated and administered to patients in higher dosages than is possible for the closed ring versions. This represents a major clinical advantage.

CLAIMS

1. A compound of Formula (I) or a pharmaceutically acceptable salt or solvate thereof



(I)

wherein:

R^1 and R^2 , are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} alkylsulphonyl and COOH; C_{1-6} hydroxyalkyl; C_{1-6} carboxyalkyl; and sulphide;

or wherein R_1 and R_2 together with the carbon to which they are attached form a C_{3-8} carbocyclic ring may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} haloalkyl, C_{1-4} alkylsulphonyl and COOH;

or wherein R_1 and R_2 together with the carbon to which they are attached represent a C_{2-6} alkenyl or C_{2-6} alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C_{1-4} haloalkyl and COOH, provided always that one of R^1 and R^2 is not hydroxyl when the other is methyl;

R^3 and R^4 are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-8} cycloalkyl wherein said alkyl, alkenyl, alkynyl or cycloalkyl groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} alkylsulphonyl and COOH; C_{1-6} hydroxyalkyl; C_{1-6} carboxyalkyl; and sulphide;

or R^3 and R^4 together with the carbon to which they are attached form a C_{3-8} carbocyclic ring which may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C_{1-4} haloalkyl, C_{1-4} alkylsulphonyl and COOH;

or R^3 and R^4 together represent a C_{2-6} alkenyl or C_{2-6} alkynyl group bound through a double bond to the ring to which it is attached and being optionally substituted by one to three groups independently selected from the group comprising: halo, hydroxyl, cyano, C_{1-4} haloalkyl and COOH;

R^5 is selected from the group comprising: fluoro, chloro, bromo and iodo;

R^6 is selected from the group comprising: fluoro, chloro, bromo and iodo; and

R^7 and R^8 are independently selected from the group comprising: H; halo; cyano; C_{1-6} alkyl, C_{1-6} haloalkyl, C_{1-6} alkoxy, and C_{1-6} haloalkoxy;

R^9 is H or C_{1-6} alkyl,

R¹⁰ is selected from the group comprising: hydrogen; C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl and C₃₋₈ cycloalkyl wherein each of the foregoing groups may be optionally substituted by 1 to 5 groups chosen independently from the group comprising: halo, hydroxyl, cyano, nitro, C₁₋₄ alkylsulphonyl and COOH; or R¹⁰ is a pharmaceutically acceptable counterion, and

X is O or S,

provided always that R¹, R², R³ and R⁴ are not all hydrogen.

2. A compound as claimed in claim 1, wherein R^3 is H or C_{1-6} alkyl.
3. A compound as claimed in claim 1 or 2, wherein R^4 is H or C_{1-6} alkyl.
4. A compound as claimed in any preceding claim, wherein R^5 is chloro.
5. A compound as claimed in any preceding claim, wherein R^6 is chloro.
6. A compound as claimed in any preceding claim, wherein R^7 is H.
7. A compound as claimed in any preceding claim, wherein R^8 is H.
8. A compound as claimed in any preceding claim, wherein R^9 is H or methyl.
9. A compound as claimed in any preceding claim, wherein R^{10} is C_{1-6} alkyl.

10. A compound as claimed in any preceding claim, wherein R^1 is an optionally substituted C_{1-4} alkyl or C_{3-8} cycloalkyl group, preferably wherein R^1 is methyl, cyclopropyl, CF_3 or CHF_2 .
11. A compound as claimed in any preceding claim, wherein R^2 is an optionally substituted C_{1-4} alkyl or C_{3-8} cycloalkyl group, preferably wherein R^2 is methyl, cyclopropyl, CF_3 or CHF_2 .
12. A compound as claimed in any of claims 1 to 9, wherein R^1 and R^2 together form an optionally substituted C_{3-8} cycloalkyl group.
13. A pharmaceutical composition comprising a compound of formula (I) as defined in any of claims 1 to 12, or a pharmaceutically acceptable salt or solvate thereof, together with a pharmaceutically acceptable diluent or carrier, which may be adapted for oral, parenteral or topical administration.
14. A compound of formula (I) as defined in any of claims 1 to 12, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition containing any of the foregoing, for use as a medicament.
15. Use of a compound of formula (I) as defined in any of claims 1 to 12, or a pharmaceutically acceptable salt or solvate thereof in the manufacture of a medicament for the treatment of a disease selected from: myeloproliferic diseases and generalised thrombotic diseases.

16. Use of a compound of formula (I) as defined in any of claims 1 to 12 in the manufacture of a medicament for the reduction of platelet count.