

Helicobacter pylori antigens and vaccine compositions.

TECHNICAL FIELD

5 The present invention provides recombinant polypeptides which constitute *Helicobacter pylori* antigens, said antigens being expressed on the surface of both dividing (bacillary) forms as well as resting (coccoid) forms of *H. pylori*, and giving rise to both systemic and local (mucosal) production of antibodies. The invention furthermore provides nucleic acid molecules
10 coding for the said polypeptides, as well as vectors and host cells comprising such nucleic acid molecules. The said recombinant polypeptides are useful for the diagnosis of *H. pylori* infections and for the manufacture of vaccine compositions which will elicit a protective immune response against such infections, said vaccine compositions being suitable
15 for both therapeutic and prophylactic use.

BACKGROUND ART

20 The gram-negative bacterium *Helicobacter pylori* is an important human pathogen, involved in several gastroduodenal diseases. Colonization of gastric epithelium by the bacterium leads to active inflammation and progressive chronic gastritis, with a greatly enhanced risk of progression to peptic ulcer disease.

25 In order to colonize the gastric mucosa, *H. pylori* uses a number of virulence factors. Such virulence factors comprise several adhesins, with which the bacterium associates with the mucus and/or binds to epithelial cells; ureases which helps to neutralize the acid environment; and
30 proteolytic enzymes which makes the mucus more fluid.

REPLACED SHEETS SG

☐ Description : paged to paged
☒ Claim(s) : paged 40 to paged 43
☐ Drawing(s) : paged to paged
☐ Abstract : paged to paged

Despite a strong apparent host immune response to *H. pylori*, with production of both local (mucosal) as well as systemic antibodies, the pathogen persists in the gastric mucosa, normally for the life of the host. The reason for this is probably that the spontaneously induced immune-
5 response is inadequate or directed towards the wrong epitopes of the antigens.

In order to understand the pathogenesis and immunology of *H. pylori* infections, it is of great importance to define the antigenic structure of this
10 bacterium. In particular, there is a need for characterization of surface-exposed (like adhesins) and secreted proteins which, in many bacterial pathogens, have been shown to constitute the main virulence factors, and which can be useful for the diagnosis of *H. Pylori* and in the manufacture of vaccine compositions.

15 Cloning of the gene *hpaA*, which codes for a 20 kDa receptor-binding subunit of the N-acetylneuraminyllactose-binding fibrillar hemagglutinin (NLBH) of *H. pylori*, has been disclosed by Evans et al. (1993) J. Bacteriol. 175, 674-683.

20 Monoclonal antibodies (MAbs) against membrane preparations of *H. pylori* have been disclosed by Bölin et al. (1995) J. Clin. Microbiol. 33, 381-384. One of these MAbs, designated HP30-1:1:6, reacted with a 30 kDa protein which was shown to be exposed on the surface of intact bacteria and to
25 have properties like that of an adhesin.

Whenever stressed or threatened, the *H. pylori* cell transforms from a bacillary to a coccoid form. In the coccoid form, the *H. pylori* cell is much less sensitive to antibiotics and other anti-bacterial agents. Circumstantial
30 evidence indicate the *H. pylori* might be transmitted between individuals in this form, possibly via water or direct contact. An efficient vaccine composition should therefore elicit an immune response towards both the

coccoid and the bacillary form of *H. pylori*. Since systemic immunity probably only plays a limited role in protection against mucosal infections, it is also important that the vaccine composition will enhance protective immune mechanisms locally in the stomach.

5

PURPOSE OF THE INVENTION

10 The purpose of this invention is to provide an antigenic *H. pylori* polypeptide which can be useful i.a. for eliciting a protective immune response against, and for diagnosis of, *H. pylori* infection. This purpose has been achieved by the recombinant cloning of a *H. pylori* gene which encodes a surface-exposed protein. The nucleic acid sequence of this gene is similar to the sequence of the *hpaA* gene as published by Evans et al.
15 (1993) in the Journal of Bacteriology, vol. 175, 674-683. However, while the *hpaA* gene was reported to code for a 20 kDa protein, it has surprisingly been found that the DNA molecule according to the invention encodes a polypeptide with a molecular weight of 29 kDa.

20 The 29 kDa polypeptide is shown to be an antigenic protein which is expressed in all strains of *H. pylori*, also in coccoid forms of the bacterium, and which is able to induce a mucosal as well as a systemic immune-response in a host measured as antibody production. The 29 kDa polypeptide is expressed by all *H. pylori* strains tested and antibodies
25 created towards this protein do not cross-react with common endogenous human bacteria of other species or with selected human tissues including the gastric mucosa. Thus being an essential, well conserved adhesin with immunogenic properties, the 29 kDa polypeptide will be useful both for the detection of *H. pylori* infections as well as for the manufacture of
30 vaccine compositions, which when given in an appropriate pharmaceutical formulation will elicit a protective or therapeutic immune response against such infections.

The experimental data below thus indicates that the 29 kDa *H. pylori* protein is important for *H. pylori* colonization and/or persistence of infection, since binding of a monoclonal antibody for the 29 kDa protein results in complete inhibition of colonisation of *H. pylori* in mice. Furthermore, the 29 kDa *H. pylori* protein, when used as an oral immunogen, acts as a stimulator of an immune response leading to a significant reduction of colonisation of *H. pylori* in mice which were infected with *H. pylori* 1 month prior to immunization.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1: Restriction enzyme map of plasmid pAE1 containing the 1.7 kb fragment of *H. pylori* encoding the 29 kDa polypeptide. Hatched bar indicates the position of the structural gene. The location of the T3 and T7 promoter sequences are shown above the black bars indicating the vector.

Fig. 2: Plasmid maps of pS860, pS861, pS862 and pS863.

Filled arrows: *lac* operon promoter (Plac) or bacteriophage T7 RNA Polymerase promoter (T7promoter). Grey fill: PCR generated 5'-end or 3'-end of the 29 kDa gene. Terminator: T7 transcription terminator. Ori: pBR322 plasmid replication origin.

Fig. 3: Effect of monoclonal antibodies on the colonisation of *H. pylori* in BALB/c mice.

Fig. 4: Therapeutic oral immunization of *H. pylori* infected BALB/c mice.

DISCLOSURE OF THE INVENTION

Throughout this description and in particular in the following examples, the terms "standard protocols" and "standard procedures", when used in
5 the context of molecular cloning techniques, are to be understood as protocols and procedures found in an ordinary laboratory manual such as: Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) Molecular Cloning: A laboratory manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

10

In a first important aspect, this invention provides a recombinant polypeptide which has an amino acid sequence identical with, or substantially similar to, a *Helicobacter pylori* surface-exposed antigen with an approximate molecular weight of 29 kDa.

15

The said surface-exposed antigen according to the invention has i.a. the following important properties:

- It is an adhesin, which is important for the colonization of the gastric mucosa;
- 20 • It is expressed on the surface of both dividing (bacillary) forms as well as resting (coccoid) forms of *H. pylori*;
- It is a strong antigen giving rise to both systemic and local (mucosal) production of antibodies;
- It is conserved in all tested strains of *H. pylori*;
- 25 • Antibodies to the 29 kDa polypeptide do not cross-react with a number of different non-helicobacter bacteria, or with selected human tissues, including the gastric mucosa;
- The 29 kDa polypeptide is lipidated and thus post-translationally modified. This feature of the polypeptide may be of importance for
30 its immunogenicity and for its proper exposure on the surface of *H. pylori*. It is known in the art that lipid modification can be essential

for the immunological properties of bacterial lipoproteins (see Weis, J.J. et al. (1994) *Infection and Immunity*, vol. 62, 4632-4636).

- It is a putative virulence-factor, whereby the term "virulence factor" is to be understood a molecule specifically involved in adherence of *H. pylori* to the epithelial surface of the gastric mucosa and / or in the establishment and maintenance of *H. pylori* infection.

In a preferred form, the said polypeptide has an amino acid sequence according to positions 1-260, or 28-260, in SEQ ID NO: 2 or 4 of the Sequence Listing. As further described in the Experimental Section, it is believed that positions 1-260 in SEQ ID NO: 2 and 4 represent the uncleaved protein, while positions 1-27 represent a signal sequence and positions 28-260 represent the mature polypeptide. The only difference between SEQ ID NO: 2 and SEQ ID NO: 4 is that SEQ ID NO: 2 has a Ser residue in position 222, while SEQ ID NO: 4 has an Arg residue in the same position.

However, the polypeptide according to the invention is not to be limited strictly to a polypeptide with an amino acid sequence identical with the above mentioned positions in SEQ ID NO: 2 or 4 in the Sequence Listing. Rather the invention encompasses polypeptides carrying modifications like substitutions, small deletions, insertions or inversions, which polypeptides nevertheless have substantially the properties of the 29 kDa polypeptide according to the invention. Such properties include the ability to elicit a mucosal as well as systemic immune-response against *H. pylori* in a mammal host; the ability to work as an adhesin; and the presence of the polypeptide in both bacillary and coccoid forms of *H. pylori*.

Included in the invention are consequently polypeptides, the amino acid sequence of which is at least 90% homologous, preferably at least 95% homologous, with the amino acid sequence shown as positions 1-260, or positions 28-260, in SEQ ID NO: 2 or 4, in the Sequence Listing, which

polypeptides nevertheless have substantially the biological activities of the 29 kDa polypeptide according to the invention.

5 Included in the invention are also peptides, with a length of at least 5 amino acids, which comprise an immunogenic epitope of the 29 kDa polypeptide according to the invention and retains the ability to elicit an immune response against *H. pylori* bacteria in a mammal host. Such epitope(s) can be presented alone or in the form of fusion proteins, where the epitope is fused to an inert or immunologically active carrier
10 polypeptide. The identification of these epitopes will be based on the presence of host-generated antibodies towards different segments of the 29 kDa polypeptide.

15 One way of obtaining structural information on the epitopes of the 29 kDa polypeptide is the production and characterisation of monoclonal antibodies binding to the polypeptide, followed by mapping of epitopes by e.g. Pepscan analysis. Monoclonal antibodies can be produced by standard methods, such as those described by De St. Groth (1980) in J. Immunol. Methods, vol. 35, 1-21.

20 In another aspect, the invention provides an isolated and purified nucleic acid molecule which has a nucleotide sequence coding for a polypeptide as defined above. In a preferred form of the invention, the said nucleic acid molecule is a DNA molecule which has a nucleotide sequence identical
25 with SEQ ID NO: 1 or 3 of the Sequence Listing. However, the DNA molecule according to the invention is not to be limited strictly to the sequence shown as SEQ ID NO: 1 or 3. Rather the invention encompasses DNA molecules carrying modifications like substitutions, small deletions, insertions or inversions, which nevertheless encode polypeptides having
30 substantially the biochemical activity of the 29 kDa polypeptide according to the invention. It will be known to the skilled person that A $\leftrightarrow$ G and T $\leftrightarrow$ C substitutions, with no effect on the amino acid sequence, are not

unusual in *H. pylori*. The only difference between SEQ ID NO: 1 and SEQ ID NO: 3 is that SEQ ID NO: 1 has an A residue in position 1458, while SEQ ID NO: 3 has a C residue in the same position.

- 5 Included in the invention are also DNA molecules which nucleotide sequences are degenerate, because of the genetic code, to the nucleotide sequence shown as SEQ ID NO: 1 or 3. Since there are 64 possible codons, but only 20 natural amino acids, most amino acids are coded for by more than one codon. This natural "degeneracy", or "redundancy", of the genetic
- 10 code is well known in the art. It will thus be appreciated that the DNA sequence shown in the Sequence Listing is only an example within a large but definite group of DNA sequences which will encode the polypeptide as described above.
- 15 Consequently, the inventions includes an isolated nucleic acid molecule selected from:
- (a) nucleic acid molecules comprising a nucleotide sequence which is identical with, or substantially similar to, positions 796-1572 or 874-1572 in SEQ ID NO: 1 or 3 in the Sequence Listing;
- 20 (b) nucleic acid molecules comprising a nucleotide sequence capable of hybridizing to a nucleotide sequence complementary the polypeptide coding region of a DNA molecule as defined in (a) and which codes for a polypeptide according to the invention, or a functionally equivalent modified form thereof; and (c) nucleic acid molecules comprising a nucleic
- 25 acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence as defined in (a) or (b) and which codes for a polypeptide according to the invention, or a functionally equivalent modified form thereof.
- 30 A further aspect of the invention is a vector which comprises the nucleic acid molecule according to the invention. Such a vector can preferably be

the plasmid vector pAE1 (Deposited under the Budapest Treaty under accession No. NCIMB 40732).

5 A vector according to the invention can also be a replicable expression vector which carries and is capable of mediating the expression of a nucleic acid molecule according to the invention. In the present context the term "replicable" means that the vector is able to replicate in a given type of host cell into which it has been introduced. Examples of vectors are viruses such as bacteriophages, cosmids, plasmids and other recombination
10 vectors. Nucleic acid molecules are inserted into vector genomes by standard methods known in the art. An expression vector according to the invention can preferably be any one of the vectors pAL30:1, pAL30:2, pAL30:3, pAL30:4 or, more preferably, pS863.

15 Included in the invention is also a host cell harbouring a vector according to the invention. Such a host cell can be a prokaryotic cell, a unicellular eukaryotic cell or a cell derived from a multicellular organism. The host cell can thus e.g. be a bacterial cell such as an *E. coli* cell; a cell from a yeast such as *Saccharomyces cerevisiae* or *Pichia pastoris*, or a mammalian cell.
20 The methods employed to effect introduction of the vector into the host cell are standard methods well known to a person familiar with recombinant DNA methods.

In another aspect, the invention provides a process for production of a
25 polypeptide as defined above, said method comprising culturing a host cell transformed with an expression vector as defined above, under conditions whereby said polypeptide is produced, and recovering said polypeptide.

The medium used to grow the cells may be any conventional medium
30 suitable for the purpose. A suitable vector may be any of the vectors described above, and an appropriate host cell may be any of the cell types listed above. The methods employed to construct the vector and effect

introduction thereof into the host cell may be any methods known for such purposes within the field of recombinant DNA. The recombinant polypeptide expressed by the cells may be secreted, i.e. exported through the cell membrane, dependent on the type of cell and the composition of the vector.

If the polypeptide is produced intracellularly by the recombinant host, i.e. is not secreted by the cell, it may be recovered by standard procedures comprising cell disruption by mechanical means, e.g. sonication or homogenization, or by enzymatic or chemical means followed by purification. In order to be secreted, the DNA sequence encoding the polypeptide should be preceded by a sequence coding for a signal peptide, the presence of which ensures secretion of the polypeptide from the cells so that at least a significant proportion of the polypeptide expressed is secreted into the culture medium and recovered.

A further aspect of the invention is a polypeptide according to the invention for use in therapy, for use in the diagnosis of *Helicobacter pylori* infection in a mammal, including man, and for use as a therapeutic or prophylactic vaccine.

Another important aspect of the invention is a vaccine composition for inducing a protective immune response in a mammal, including humans, against the bacillary and/or coccoid form of *Helicobacter pylori*. Such a vaccine composition comprises an immunogenically effective amount of a polypeptide as defined above, including at least a part of the 29 kDa polypeptide comprising an immunogenic epitope, or a modified form of said polypeptide which retains the capability to induce protective immunity against *Helicobacter pylori* infection. The term "modified form" includes, but is not restricted to, forms of the polypeptide which are post-translationally modified, e.g. lipidated. It is believed that the 29 kDa protein is lipidated, cf. Example 4 below.

The vaccine composition comprises optionally also a pharmaceutically acceptable carrier or diluent, or other immunologically active antigens for prophylactic or therapeutic use. Physiologically acceptable carriers and diluents are well known to those skilled in the art and include e.g.
5 phosphate buffered saline (PBS), or, in the case of oral vaccines, HCO_3^- based formulations or enterically coated powder formulations.

The vaccine composition can optionally include or be administered together with acid secretion inhibitors, preferably proton pump inhibitors (PPIs), e.g. omeprazole. The vaccine can be formulated in known delivery
10 systems such as liposomes, ISCOMs, cochleates, etc. (see e.g. Rabinovich et al. (1994) Science 265, 1401-1404) or be attached to or included into polymer microspheres of degradable or non-degradable nature. The antigens could be associated with live attenuated bacteria, viruses or
15 phages or with killed vectors of the same kind.

As will be demonstrated in the Experimental Section below, a vaccine composition according to the invention can be used for both therapeutic and prophylactic purposes. The vaccine composition according to the
20 invention is preferably administered to any mammalian mucosa exemplified by the buccal, the nasal, the tonsillar, the gastric, the intestinal (small and large intestine), the rectal and the vaginal mucosa. The mucosal vaccines can be given together with for the purpose appropriate adjuvants. The vaccine can also be given parenterally, by subcutaneous,
25 intracutaneous or intramuscular route, optionally together with the appropriate adjuvant.

An alternative approach for creating an immune response against the 29 kDa polypeptide is to use the approach known as "nucleic acid
30 vaccination" or "naked DNA" vaccination. It is known in the art that injection into muscle of plasmid DNA encoding an antigen of interest can result in sustained expression of the antigen and generation of an immune

response (see e.g. Rabinovich et al. *supra*). Several routes of administration are possible, such as parental, mucosal or via a "gene-gun" that delivers tiny amounts of DNA-coated gold beads (Fynan et al. (1993) Proc. Natl. Acad. Sci. U.S.A. 90, 11478-11482).

5

Thus, a nucleic acid molecule according to the invention can be expressed in plasmid comprising a suitable eukaryotic promoter. This "naked DNA" can then be injected intramuscularly or given intradermally via a "gene-gun". Epitopes of the expressed protein will be expressed by MHC

10

molecules on the surface of the cells and trigger an immune response.

Consequently, nucleic acid molecules and vectors as disclosed in the previous paragraphs for use in therapy, in particular for use as a vaccine, are further aspects of the invention. The use of such nucleic acid molecules and vectors in the manufacture of compositions for treatment, prophylaxis or diagnosis of *Helicobacter pylori* infection are also further aspects of the invention.

15

Yet another aspect of the invention is the use of a polypeptide as defined above, or a modified form of said polypeptide which retains the capability to induce protective immunity against *Helicobacter pylori* infection, in the manufacture of a compositions for the treatment, prophylaxis or diagnosis of *Helicobacter pylori* infection. Such compositions include in particular a vaccine composition eliciting a protective immune response against the bacillary and/or coccoid form of *Helicobacter pylori*. Included in the invention is also the said use in the manufacture of a diagnostic kit for diagnosis of *Helicobacter pylori* infection. Such a diagnostic kit is further described below.

20

25

In a further aspect, the invention provides a method of eliciting in a mammal, including man, a protective immune response against *Helicobacter pylori* infection, said method comprising the step of administering to the said mammal an immunologically effective amount of a vaccine

30

composition as defined above. The term "immunologically effective amount" is intended to mean an amount which elicit a significant protective *Helicobacter pylori* response, which will eradicate a *H. pylori* infection in an infected mammal or prevent the infection in a susceptible mammal. Typically an immunologically effective amount would comprise approximately 1 µg to 100 mg, preferably approximately 10 µg to 10 mg, of *H. pylori* antigen for oral administration, or approximately less than 100 µg for parenteral administration.

Another aspect of the invention is a method of *in vitro* diagnosis of *Helicobacter pylori* infection, comprising at least one step wherein a polypeptide as defined above, including a part of the 29 kDa polypeptide which part comprises an immunogenic epitope, is used. The polypeptide can optionally be labelled and/or coupled to a solid support. A method of diagnosis can e.g. comprise the steps (a) contacting a said polypeptide, optionally bound to a solid support, with a body fluid taken from a mammal; and (b) detecting antibodies from the said body fluid binding to the said polypeptide. Preferred methods of detecting antibodies are ELISA (Enzyme linked immunoabsorbent assay) methods which are well known in the art.

In yet another aspect, the invention provides a diagnostic kit for the detection of *Helicobacter pylori* infection in a mammal, including man, comprising components which enable a diagnosis method as exemplified above to be carried out.

EXAMPLES

EXAMPLE 1: Cloning and expression of a 29 kDa polypeptide from *H. pylori*

5

1.1. Bacterial strains, vectors and growth conditions

H. pylori CCUG 17874 (= NTCC 11637) was grown on horse blood agar plates in an microaerophilic atmosphere. *E. coli* strains XL1-Blue MRF⁺ and XLOLR (Stratagene, La Jolla, California) were used as host strains for cloning experiments and were grown in Luria-Bertani broth (LB) or NZY medium supplemented with 0.2% maltose and 10 mM MgSO₄ when used for lambda infection. The lambda expression vector ZAP ExpressTM and its phagemid derivative pBK-CMV were obtained from Stratagene.

15

1.2. DNA techniques

Chromosomal DNA from *H. pylori* 17874 was prepared by suspending bacteria from plates incubated for 48 h in 50 mM Tris-Cl, pH 8.0, 25% sucrose, 50 mM EDTA containing 10 mg/ml lysozyme, and 5 ng/ml DNase-free RNase (Boehringer Mannheim Scandinavia AB, Bromma, Sweden). The suspension was incubated for 10 min at +37°C. An equal volume of lysis buffer (0.4% Triton X100 in 50 mM Tris-Cl, pH 8.0; and 62.5 mM EDTA) was added and the suspension was incubated at room temperature until a noticeable lysis of the bacteria occurred. The suspension was then extracted in three steps, with buffered phenol (pH 8.0), phenol/chloroform and chloroform, respectively. The DNA was precipitated from the aqueous phase and dissolved in TE-buffer (10 mM Tris-Cl, pH 8.0; and 1 mM EDTA).

30

Restriction enzymes were purchased from Boehringer Mannheim Scandinavia AB and used according to the manufacturers instructions.

Plasmids and lambda DNA were purified with Wizard kits (Promega, Madison, Wisconsin). Sequencing was performed using the Sequenase 2.0 kit (Amersham Sweden AB, Solna, Sweden). Oligonucleotides were purchased from Innovagen, Lund, Sweden. PCR was performed using Taq DNA polymerase (Boehringer-Mannheim Scandinavia AB).

1.3. Construction of a *H. pylori* genomic library

Chromosomal DNA fragments in the size range 2-12 kb were purified from partially *Sau*3A-cleaved *H. pylori* 17874 DNA and cloned into *Bam*HI digested ZAP Express™ vector as described in the Stratagene protocol. Following *in vitro* packaging, the library was titrated by infecting strain XL-1 Blue MRF and plated onto indicator plates containing isopropyl-β-D-thiogalactopyranoside (IPTG) and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranosid (X-Gal). The titer of the library was 1.2×10^6 PFU/ml with 85% recombinants.

Plaques expressing the 29 kDa polypeptide were detected by immunological screening using MAb HP30-1:1:6 (Bölin et al. (1995) J. Clin. Microbiol. 33, 381-384) according to standard methods. Positive plaques were isolated and the plating and screening with the MAb was repeated until plaque purity was obtained. The conversion to the phagemid form of the ZAP Express clones was accomplished using the ExAssist protocol (Stratagene).

1.4. Immunoblotting and dot blot test

Overnight cultures of *E. coli* XL0LR containing plasmids with cloned inserts from *H. pylori* 17874 depicted in Fig. 1, were diluted 1:100 in 5 ml of LB medium with 50 mg/ml kanamycin. The cultures were incubated at +37°C until the OD at 600 nm was 0.7. IPTG was added to a final concentration of 1 mM and the bacteria were grown for additional 2 h.

Cultures without IPTG were grown similarly. The cultures were centrifuged and resuspended in 1/10 of the volume. Ten µl of the suspension were mixed with an equal volume of 2X sample buffer, boiled and analysed by SDS-PAGE. Strain XL0LR, grown in the same way but without kanamycin, served as a negative control. A suspension of *H. pylori* 17874 in PBS (OD at 600 nm = 1.0) was used as a positive control.

After immobilization of the protein profiles on nitro-cellulose sheets, reaction with the 29 kDa polypeptide-specific MAb HP30-1:1:6 diluted 1:10 was carried out as described previously (Bölin et al., 1995) and bound antibodies were detected by using anti-mouse IgG labelled with peroxidase. Filters were developed with hydrogen peroxide substrate and 4-chloronaphthol chromogen (BioRad Svenska AB).

The dot blot test was performed using overnight cultures of the above strains. Two µl of a suspension were spotted on nitrocellulose filters, air-dried and incubated with MAb HP30-1:1:6 diluted 1:10 for one h. Subsequent steps were carried out as described for immunoblotting.

1.5. Molecular cloning

Partially digested chromosomal DNA from *H. pylori* strain 17874 were cloned into a lambda expression vector (ZAP Express™). Four plaques expressing the 29 kDa polypeptide were detected after screening of 24 000 plaques for reaction with the 29 kDa polypeptide-specific MAb. The positive plaques were purified and the size of the cloned inserts were examined by digestion of DNA-preparations with *Xba*I and *Sall*. The inserts were from 3.7 to 1.78 kb in size. After *in vivo* excision of the pBK-CMV phagemids from the four positive plaques, restriction enzyme maps were constructed and compared with the inserts in the lambda vector. The phagemids were found to contain overlapping DNA-fragments with the

same size as in the lambda vector. Most of the restriction enzymes tested, except for *SmaI* and *NheI*, did not cleave the cloned fragments.

5 The restriction map of the smallest cloned 1.7 fragment (pAE1) that were further analyzed is shown in Fig. 1. One of the cloned inserts were in the opposite direction with regard to the vector promoter. When whole cell extracts of the *E. coli* strains containing these plasmids were analysed in immunoblotting with MAb HP30-1:1:6, they were all found to express a polypeptide with the same molecular weight as *H. pylori* 17874. No
10 difference in expression of the 29 kDa polypeptide was seen when the vector promoter was induced with IPTG. This indicated that the gene was transcribed from its own promoter. Three subclones containing the DNA fragments indicated in Fig. 1 were constructed and examined for expression of the 29 kDa polypeptide. None of the clones expressed the
15 polypeptide. When XL0LR (pAE1) were tested in the dot blot assay (Bölin et al., 1995) and compared with *H. pylori*, it was found to be weakly positive indicating that some of the expressed polypeptide may be exposed on the surface.

20 1.6. Analysis of the DNA sequence

Both strands of the 1.7 kb insert of pAE1 and the subclones were sequenced using T3- and T7-specific primers and, when necessary, supplemented with specific primers to cover regions of the sequence not
25 available with the standard primers. The computer analysis showed that the sequence (SEQ ID NO: 1) contained an open reading frame (ORF) of 780 bp on one strand, spanning the restriction enzyme sites used for subcloning (Fig. 1). A putative ribosome binding site could be identified (positions 782-785 of SEQ ID NO: 1). The ORF coded for 260 amino acids
30 of a polypeptide of a molecular weight of 29,126 Da (SEQ ID NO: 2).

The amino acid sequence was found to contain a possible signal sequence of 27 amino acids. The sequence Leu-Val-Gly-Cys (positions 25 to 28 in SEQ ID NO: 2 and 4) is one of the consensus sequences (Leu-X-Y-Cys) assigned as a recognition site for the enzyme signal peptidase II. The signal peptidase II cleaves the signal sequences before the cysteine residue in prolipoproteins. The characteristics of the signal sequence thus suggest that the 29 kDa protein is a lipoprotein and that the mature protein comprises amino acids 28 to 260.

1.7. Expression of the recombinant 29 kDa polypeptide in *E. coli*

The recombinant 29 kDa polypeptide was produced in high concentration in *E. coli* N4830-1 from the expression vector construct pAL30, which contains the entire gene of the 29 kDa polypeptide (positions 771-1667 in SEQ ID NO: 1 and 3) The vector used for the construct was pML-LCTB λ 7 (obtained from Michael Lebens, University of Gothenburg, Sweden) which contains a strong λP_L promoter. The vector also comprises a β -lactamase gene giving ampicillin resistance. The LCTB gene (encoding the cholera toxin and its signal peptide), which is inserted between the λP_L promoter and a terminator region in the vector, was excised from the vector by cleaving with the restriction enzymes *Sma*I and *Hind*III.

The structural gene encoding the 29 kDa polypeptide, including its signal sequence, was amplified by Polymerase Chain Reaction (PCR). The primers used were HP30N (GGC GTA GAA ATG GAA GCG C; corresponding to positions 522 to 540 in SEQ ID NO: 1 and 3) which binds 271 bp upstream of the ATG start codon and HP30C (CCC AAG ATT CAT CAG CCC TTA AAT ACA CG) which recognizes a DNA fragment 855 bp downstream the start codon (corresponding to positions 1648 to 1667 in SEQ ID NO: 1 and 3). The HP30C primer contained a *Hind*III cleavage site which by the PCR reaction was added to the sequence of the 29 kDa polypeptide gene. The resulting PCR product was 1.1 kb. This DNA fragment was cleaved by *Ssp*I

and *HindIII* which gave a fragment of 0.9 kb which was ligated to the vector fragment (2.7 kb). The vector construct now called pAL30 (3.6 kb) was transformed into *E. coli* N4830-1 by electroporation. Four positive clones were found (pAL30:1, 2, 3, 4).

5

To induce expression of the recombinant polypeptide the N4830-1 cells containing the pAL30: 1 to 4 were grown over night at +30°C (the lambda cI repressor inhibits the transcription at this temperature) in 1 x LB with ampicillin (100 µg/ml). A small part of this over night culture was
10 inoculated in 5 ml 1 x LB with ampicillin and the cells were grown at +30°C until the O.D. at 600 nm was about 0.7. The temperature was then raised to +42°C, whereby the repressor was inactivated, and incubated for two additional hours.

15

Samples taken before and after induction was analysed on 14% SDS-PAGE and by immunoblotting, using the monoclonal antibody HP30-1:1:6 which is specific for the 29 kDa polypeptide. All three induced clones used in immunoblotting (pAL30: 1, 3 and 4) expressed a large amount of the recombinant polypeptide after induction. The suspension from the non-
20 induced cells contained only a low amount of the 29 kDa polypeptide.

25

The clone pAL30:1 was chosen for further analysis. In order to verify that the clone really contained the gene encoding the 29 kDa polypeptide, the ends of the fragment inserted in the vector was sequenced. It was verified
that the sequence inserted into the expression vector corresponded to the expected sequence of the cloned PCR fragment.

30

EXAMPLE 2: Kinetics of expression of the 29 kDa polypeptide during various culture conditions

Two strains of *H. pylori* were used, namely CCUG 17874 (a laboratory strain) and Hel 73 (recently isolated from a patient suffering from duodenal ulcer). Cultivation was performed on blood agar plates, as well as in Brucella Broth supplemented with cyclodextrin. All cultures were
5 incubated in a microaerophilic atmosphere consisting of 5% O₂, 10% CO₂ and 85% N₂. Bacteria were harvested after 2, 4 and 7 days, washed once in PBS and kept at -20°C for subsequent analysis. The expression of the 29 kDa surface polypeptide was analysed by inhibition-ELISA employing specific monoclonal antibodies as previously used for detection of *E. coli*
10 surface antigens (Lopez-Vidal, Y and Svennerholm, A-M., J. Clin. Microbiol. 28, 1906-) against the polypeptide. These antibodies were also used in immunoelectron microscopy.

When CCUG 17874 had been cultivated for 7 days, on blood agar plates as well as in brucella broth, approximately 70% of the bacteria had converted
15 from the spiral form to the coccoid form. This conversion occurred already after 3 days in Hel 73. The inhibition-ELISA showed a fairly constant concentration of the 29 kDa polypeptide in samples from both plate and broth cultures, during the 7 days. The presence of the polypeptide was
20 confirmed by immunoelectron microscopy. The 29 kDa polypeptide was found to be well preserved in coccoid forms of *H. pylori*. The 29 kDa polypeptide was found to be more abundant in Hel 73 than in CCUG 17874.

25

EXAMPLE 3: Antibody responses against the 29 kDa polypeptide

Antibody responses against the 29 kDa polypeptide were determined in sera and gastric aspirates from patients with duodenal ulcers (n=19), in
30 asymptotic *H. pylori* carriers (n=18) and in non-infected age-matched controls (n=20).

Antibody levels against the 29 kDa polypeptide were tested in gastric aspirates and in sera from the three groups of subjects, by means of different ELISA methods. A majority of the infected subjects had significantly higher levels, compared with the healthy control persons, of specific antibodies against the 29 kDa polypeptide both in serum and in gastric aspirates. Antibody titers in asymptomatic carriers were comparable to those of the symptomatic patients.

10 **EXAMPLE 4: Labelling of polypeptides with [³H]palmitate**

Since the amino acid sequence of the 29 kDa polypeptide contained a possible signal peptide typical for lipoproteins, the labelling of the protein with radioactive palmitic acid was investigated:

15

E. coli N4830-1, either lacking or carrying pAL30:1, were grown at +30°C in LB-broth supplemented with 50 µg of carbencillin / ml. At a cell density of 10⁸ bacteria / ml, [³H]palmitic acid (5 mCi/ml; DuPont NEN, Boston, MA) was added to a final concentration of 50 µCi/ml. The temperature was raised to +42°C and the cultures were incubated for another 12 h. The cells were collected by centrifugation and lysed in SDS-PAGE lysis buffer. After electrophoresis, the gel was processed for flougraphy by immersing the gel in Amplify™ (Amersham International, UK) for 30 min, drying it between cellophane sheets and exposing the gel to X-ray film at -70°C for 36 h.

25

The results indicated that the 29 kDa polypeptide is lipidated and thus post-translationally modified.

EXAMPLE 5: Triton X-114 partitioning of *E. coli* expressing the recombinant 29 kDa polypeptide

E. coli cells carrying pAL30:1 were grown at +30°C in LB-broth supplemented with 50 µg carbencillin / ml. At a cell density of 10⁸ bacteria / ml, the temperature was raised to +42°C and the cultures were incubated for another 3 h. The cells were collected by centrifugation (11.300 × g, 10 min, +4°C) and resuspended in 25 ml of PBS per gram of cell pellet. The suspension was frozen and then thawed at room temperature, and 25 µl DNase I (10 µg/µl) was added. The sample was gently shaken by inversion for 30 min at room temperature and chilled to 8-12°C followed by the addition of Triton X-114 (final concentration 0.3%). After incubation by gentle inversion at +4°C for 3 h the insoluble material was collected by centrifugation (18.900 × g, 10 min, +25°C).

The phases were analysed by SDS-PAGE and the identity of the 29 kDa polypeptide was verified by Western blotting using MAb HP30-1:1:6. The results indicated that the 29 kDa polypeptide appeared in the detergent phase, which confirmed that it is a lipoprotein. It is known in the art that integral membrane proteins are normally recovered in the detergent phase (Bordier, C. (1981) J. Biol. Chem., vol. 256, 1604-1607).

This experiment also verified that a plasmid inserted into *E. coli* could express and produce the 29 kDa protein. This is important for the future production of a vaccine in larger scale, since *H. pylori* does not grow very well or fast.

EXAMPLE 6: Construction of expression vector pS863 for production of high levels of *H. pylori* 29 kDa protein

6.1. Preparation of pS860

To generate convenient restriction sites for the 5'-end of the 29 kDa gene, two synthetic oligonucleotides for PCR amplification were synthesized. The
 5 plasmid pS852 (identical to the plasmid pAL30:1 described in Example 1.7) was used as a template for the PCR amplification. The sequences of these two oligonucleotides are listed below:

EcoRI NdeI
 10 5'-CGGAATTCCATATGAGAGCAAATAATCATTTTAAAG-3'

BamHI XmaI NheI
 5'-GCGGATCCCCCGGGCTAGCTGGATGGTAATTCATTTC-3'

15 PCR amplification was performed and the 169 bp amplified fragment was ligated into the TA vector (Mead, D.A. et al. (1991) Bio/Technology 9, 657-663). The constructed plasmid was designated pS860 (Fig. 2). The sequence of the construct was confirmed by dideoxy sequencing (Sanger et al. (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467).

20

6.2. Preparation of pS861

In order to change restriction sites in the 3'-end of the 29 kDa gene two synthetic oligonucleotides for PCR amplification were synthesized. The
 25 plasmid pS852 (pAL30:1) was used as a template for the PCR amplification. The sequences of the two oligonucleotides are listed below:

EcoRI XmaI
 30 5'-CGGAATTCCCCGGGTTATTATTCTCCACCGG-3'

PstI BamHI
 5'-CGCTGCAGGGATCCTTATTATCGGTTTCTTTTGCCTTTTAA-3'

35 PCR amplification were performed and the amplified fragment was digested with *XmaI* and *BamHI* generating a 357 bp fragment. This fragment was cloned into pUC19, the constructed plasmid was designated

pS861 (Fig. 2). The sequence of the construct was confirmed by dideoxy sequencing (Sanger et al. (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467).

6.3. Preparation of plasmid pS862

5

The cDNA encoding the middle part of the 29 kDa gene was isolated by gel electrophoresis as a 280 bp *NheI/XmaI* fragment from the plasmid pS852 (pAL30:1). This fragment was ligated together with a 357 bp *XmaI/BamHI* fragment from pS861 and a 4061 bp *NheI/BamHI* fragment from pS861. The generated plasmid was designated pS862 (Fig. 2).

10

6.4. Preparation of plasmid pS863

Thereafter, a 795 bp *NdeI* and *BamHI* restriction fragment was isolated from pS862 and ligated to a 4 kb *NdeI/BamHI* fragment from T7 vector pS637(pET-3a) (Studier, F.W. et al. (1990) Methods Enzymol. 185, 60-89). The resulting expression vector was designated pS863 (Fig. 2).

15

20 EXAMPLE 7: Purification of recombinant *H. pylori* 29 kDa lipoprotein

7.1. Host strains and bacterial cultures

The expression vector pS863 was transformed into the following *E.coli* host strains; BL21(DE3); BL21(DE3)pLysS; and BL21(DE3)pLysE. The expression experiments were carried out essentially as described by Studier et al. (Methods Enzymol. 185, 60-89, 1990). The bacteria were grown in LB medium (Ausubel, F.M. et al. (eds.) Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1992) containing 50 µg/ml carbencillin. In addition, when BL21(DE3)pLysS and BL21(DE3)pLysE were used, the medium was supplemented with 30 µg/ml chloramphenicol. For

25

30

induction of the T7 expression system, the cultures were grown to a density of approximately $OD_{600} = 0.5$, and then supplemented with 0.4 mM IPTG for induction. The cells were harvested about 180 minutes after induction. The host strain that gave the highest expression level was

5 BL21(DE3)pLysS.

7.2. Purification of the *H. pylori* 29 kDa lipoprotein

Cultures of *E. coli* BL21(DE3)/pLysS transformed with plasmid pS863 were

10 grown as described above and the cells were collected by centrifugation and resuspended in cold buffer (50 mM Tris-HCl, 2 mM EDTA, 10 mM NaCl, pH 8.0). For each gram of pellet (wet weight) 35 ml of buffer was added.

15 7.2.1. Triton X-114 extraction

To extract the lipoprotein, Triton X-114 (TX-114) was added to a final concentration of 1.5% (v/v) and the suspension was stirred for one hour at 0°C. The Triton-insoluble material was pelleted by centrifugation at 18,900

20 x g for 10 min. In some cases was the pellet extracted once more but with half the volume of TX-114 containing buffer. After the second TX-114 extraction the pellet was discarded.

Phase partitioning of the supernatant from the TX-114 extraction was

25 obtained by incubating it for 15 min at +30°C with occasional mixing. The turbid solution was centrifuged at 31,300 x g for 30 min at +30°C. The lower detergent phase was collected and diluted to 1% TX-114 with cold buffer (50 mM Tris-HCl, 2 mM EDTA, 10 mM NaCl, pH 8.0).

7.2.2. *Q-sepharose, pH 8.0*

The diluted TX-114-phase was applied to a Q-sepharose column (Pharmacia) (20 ml/3 g cell pellet) equilibrated with buffer (50 mM Tris-HCl, 2 mM EDTA, 10 mM NaCl, 0.1% Triton X-100, pH 8.0). The 29 kDa lipoprotein was collected as the non-binding fraction. This fraction was phase partitioned by incubating it at +30°C with occasional mixing until the solution was turbid. The two phases were separated by centrifugation at 31,300 x g for 30 min at +30°C. The lower detergent phase was collected and diluted to 1% TX-114 with cold buffer (10 mM Tris-HCl, 2 mM EDTA, pH 8.6).

7.2.3. *Q-sepharose, pH 8.6*

The diluted TX-114-phase was applied to a 100 ml Q-sepharose column (Pharmacia) equilibrated with buffer (10 mM Tris-HCl, 2 mM EDTA, pH 8.6). The non-binding fraction contained TX-114. The column was washed with buffer A (10 mM Tris-HCl, 2 mM EDTA, 0.1% Triton X-100, pH 8.6). The 29 kDa lipoprotein was collected by a salt gradient with buffer B (10 mM Tris-HCl, 2 mM EDTA, 0.1% Triton X-100, 1 M NaCl, pH 8.6). The gradient was as follows; 0-50%B, 40 ml; 50-100%B, 100 ml. The 29 kDa lipoprotein eluted between 60-70% B.

7.2.4. *SDS-PAGE and protein electroblotting*

25

Protein samples from the different purification steps were solubilized in sample buffer (50 mM Tris-HCl, pH 6.8, 8% glycerol, 1.6% SDS, 4% β -mercaptoethanol, 0.02% bromphenol blue) and separated on Novex precast gradient gels (4-20% polyacrylamide) or BioRad precast gradient gels (10-20% polyacrylamide). The electrophoresis running buffer contained 25 mM

30

-27-

Tris, 192 mM glycine, 0.5% SDS, pH 8.3. Gels were stained with 0.1% Coomassie Brilliant Blue R-250 in 40% methanol, 10% acetic acid and destained with 10% methanol, 10% acetic acid.

5 Gels intended for Semi-Dry-electroblotting were not stained but soaked in Transfer buffer (48 mM Tris, 38 mM glycine, 0.075 % SDS, 20 % MeOH) and proteins were transferred onto PVDF membranes (Immobilon[®], Millipore, USA) by a SemiDry electroblotting apparatus (BioRad). Immunodetection was accomplished by first blocking the PVDF membrane for one hour in
10 2% BSA in TBS (50 mM Tris-HCl, 2.5 M NaCl, pH 8.2) and thereafter the membrane was incubated for one hour with a specific monoclonal antibody (IgG1) against the 29 kDa lipoprotein diluted 1:10 with 1% BSA in TBS. After a washing step with TBS the membrane was incubated for one hour with an alkaline phosphatase-conjugated anti-mouse IgG antibody
15 (Dakopatts, Denmark). After an additional wash the membrane was developed with appropriate substrates (5-bromo-4-chloro-3-indolyl phosphate (BCIP) and nitroblue tetrazolium (NBT) (Sigma)).

7.2.5. Protein concentration and pyrogenicity

20

Total protein concentration was determined by the bicinchoninic acid method (BCA Protein Assay. Pierce Chemical Company, USA).

25 The endotoxin content was assayed by a chromogenic *Limulus* amoebocyte lysate (LAL) test (LAL COAMATIC[®] Endotoxin. Endosafe Inc. USA)

Stained SDS-gels were scanned (BioRad Imager GS-) to determine the relative amount of protein contaminants in the final preparations. The preparations contained < 10% protein contaminants.

30

EXAMPLE 8: Analysis of the *H. pylori* 29 kDa protein for use as a vaccine**8.1. Materials & Methods****5 8.1.1. Animals**

Female SPF BALB/c mice were purchased from Bomholt Breeding centre (Denmark). They were kept in ordinary makrolon cages with free supply of water and food. The animals were 4-6 weeks old at arrival.

10

8.1.2. Infection

After a minimum of one week of acclimatization, the animals were infected with a type 2 strain of *H. pylori* (strain 244, originally isolated from an ulcer patient). This strain has earlier proven to be a good colonizer of the mouse stomach. The bacteria were grown overnight in Brucella broth supplemented with 10% fetal calf serum, at +37°C in a microaerophilic atmosphere (10% CO₂, 5% O₂). The animals were given an oral dose of omeprazole (400 mmol/kg) and after 3-5 h an oral inoculation of *H. pylori* (approximately 10⁸ cfu/animal). Infection was checked in control animals 2-3 weeks after the inoculation.

15

20

8.1.3. Immunizations

25 The animals were immunized 4 times over a 34 day period (day 1, 15, 25 and 35). Purified antigen was given at a dose of 100 µg/mouse and membrane proteins (MP) at a dose of 0.5 mg/dose. Membrane proteins were prepared by sonication of bacteria in PBS. Debris was removed by spinning the sonicate at +4°C, 2000 rpm for 5 min. The supernatant was

-29-

transferred to a new tube and spun at +4°C, 15,000 rpm for 20 min. The pellet was recovered and stored at -70°C until use.

5 As an adjuvant, the animals were also given 10 µg/mouse of cholera toxin (CT) with each immunization. Omeprazole (400 µmol/kg) was given orally to the animals 3-5 h prior to immunization as a way of protecting the antigens from acid degradation. Animals were sacrificed 4 weeks after final immunization.

10 8.1.4. *Passive protection*

To analyze the effect of monoclonal antibodies (MAbs) on the ability of *H. pylori* to colonize the mouse stomach, MAbs with different specificities were mixed with *H. pylori* 10 min prior to inoculation as described above. 15 MAbs raised against the 29 kDa protein (HP30-1:1:6), against urease (Ure 8:1); and against the *E. coli* heat-stable protein (ST 1:3) were used. The MAbs were titrated in an ELISA to allow for equal amounts of each MAb to be used in the experiment. A number of 10^7 bacteria per mouse were used for inoculation. The mice were sacrificed 2 weeks post inoculation.

20

8.1.5. *Analysis of infection*

The mice were sacrificed by CO₂ and cervical dislocation. The abdomen was opened and the stomach removed. After cutting the stomach along the greater curvature, it was rinsed in saline. The mucosa from the antrum and 25 corpus of an area of 25 mm² was scraped separately with a surgical scalpel. The mucosa scraping was suspended in Brucella broth and plated onto Blood Skirrow plates. The plates were incubated under microaerophilic conditions for 3-5 days and the number of colonies was counted. The identity of *H. pylori* was ascertained by urease and catalase 30 test and by direct microscopy or Gram staining.

8.2. Results

8.2.1 Passive protection

5 Three groups with 10 animals in each were given a mixture of *H. pylori*
strain 244 and a MAb, and one group was given only *H. pylori*. The
mixture of MAb and bacteria was allowed to react for 10 min before being
inoculated into the mice. None of the MAbs used had any clear effect on
the bacteria *in vitro*. Two weeks after inoculation, the mice were sacrificed
10 and the infection rate was determined for each group (Fig. 3). All of the
mice in the control group and those inoculated with the ST Mab were
infected. In the urease MAb group all mice were infected, but to a
significantly lower degree compared with the controls. In the group
inoculated with the MAb against the 29 kDa protein, none of the mice
15 were infected.

8.2.2. Therapeutic immunization

20 The animals in this study were infected with *H. pylori* strain 244 one month
prior to immunizations. Mice in groups of ten were then immunized with
either cholera toxin (CT) or CT together with membrane proteins, urease or
the 29 kDa protein. Control animals received vehicle only (PBS). One
month after the final immunization, the animals were sacrificed and CFU
was determined (Fig. 4). All control animals, as well as those immunized
25 with only CT, were infected. Animals actively immunized with urease and
CT, or with 29 kDa protein and CT, had significantly decreased CFU
values compared with the controls. Only one animal in the urease-
immunized group was completely cured from the infection.

8.3. Conclusions

5 The results above indicate that the 29 kDa *H. pylori* protein is important for the colonization and/or persistence of an infection, since binding of a MAb to this structure result in complete inhibition of colonisation.

10 Furthermore, the 29 kDa *H. pylori* protein, when used as an oral immunogen in conjunction with cholera toxin as an oral adjuvant, acts as a stimulator of an immune response leading to a significant reduction of the degree of colonisation of *H. pylori* in the used animal model.

15 Taken together, these results strongly support the use of the 29 kDa *H. pylori* protein in an oral vaccine formulation for the use in humans to treat and prevent *H. pylori* infections.

DEPOSIT OF MICROORGANISMS

20 The plasmid pAE1 has been deposited under the Budapest Treaty at the National Collections of Industrial and Marine Bacteria (NCIMB), Aberdeen, Scotland, UK, and under accession number NCIMB 40732. The date of deposit is 16 May 1995.

-32-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: ASTRA AB
(B) STREET: Västra Mälarehamnen 9
(C) CITY: Södertälje
(E) COUNTRY: Sweden
(F) POSTAL CODE (ZIP): S-151 85
(G) TELEPHONE: +46 8 553 260 00
(H) TELEFAX: +46 8 553 288 20
(I) TELEX: 19237 astra s

(ii) TITLE OF INVENTION: Bacterial Antigens and Vaccine Compositions

(iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1670 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 793..1575

(ix) FEATURE:

(A) NAME/KEY: mat_peptide
(B) LOCATION: 793..1572

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GATCCTATCG CGCCAAAGGT GGTATTAGGA ATAAGAGCTT GATTATTAAT CTCCTGGTA	60
AGTCCAAAAA GTATTAGAGA ATGCTTAGAG GCGGTTTTTC CAGCGATTCC TTATTGCGTG	120
GATTTGATTT TAGGGAATTA CATGCAAGTG AATGAAAAAA ACATTCAAGC GTTTGCCCCC	180
AAACAATAAG GTAAAAAATG CCACTCACTC ATTTGAATGA AGAAAAATCAA CCTAAAATGG	240
TGGATATAGG GGATAAGAA ACCACTGAAA GAATCGCTCT AGCAAGCGGT CGTATCAGCA	300

TGAATAAAGA GGCTTATGAC GCTATTATCA ATCATGGCGT CAAAAAGGGT CCGGTATTAC	360
AAACTGCTAT TATTGCTGGG ATTATGGGGG CTAAAAAGAC AAGCGAACTC ATTCCCATGT	420
GCCATCCAAT CATGCTCAAT GGGGTGGATA TTGATATTTT AGAAGAAAAA GAGACTTGTA	480
GTTTTAAACT CTATGCGAGA GTCAAAACTC AAGCTAAAAC GGGCGTAGAA ATGGAAGCGC	540
TAATGAGTGT GAGCGTAGGG CTTTAAACCA TTTATGACAT GGTGAAAGCC ATTGATAAGA	600
GCATGACAAT TAGCGGTGTG ATGCTGGAAT ATAAAAGTGG AGGCAAAAGT GGGGATTATA	660
ACGCTAAAAA ATAGAAAAAG ACTGATAATC TAAAGATATT AGGGTAAAAT AACATTTTGA	720
CAACAAAAGC GTGTTGGTTG CTTGGGATTT GTTGTATAG AAGTCTAAAA TATTACAATC	780
AAGGATAGAA CG ATG AGA GCA AAT AAT CAT TTT AAA GAT TTT GCA TGG	828
Met Arg Ala Asn Asn His Phe Lys Asp Phe Ala Trp	
1 5 10	
AAA AAA TGC CTT TTA GGC GCG AGC GTG GTG GCT TTA TTA GTG GGA TGC	876
Lys Lys Cys Leu Leu Gly Ala Ser Val Val Ala Leu Leu Val Gly Cys	
15 20 25	
AGC CCG CAT ATT ATT GAA ACC AAT GAA GTC GCT TTG AAA TTG AAT TAC	924
Ser Pro His Ile Ile Glu Thr Asn Glu Val Ala Leu Lys Leu Asn Tyr	
30 35 40	
CAT CCA GCT AGC GAG AAA GTT CAA GCG TTA GAT GAA AAG ATT TTG CTT	972
His Pro Ala Ser Glu Lys Val Gln Ala Leu Asp Glu Lys Ile Leu Leu	
45 50 55 60	
TTA AGG CCA GCT TTC CAA TAT AGC GAT AAT ATC GCT AAA GAG TAT GAA	1020
Leu Arg Pro Ala Phe Gln Tyr Ser Asp Asn Ile Ala Lys Glu Tyr Glu	
65 70 75	
AAC AAA TTC AAG AAT CAA ACC GCG CTC AAG GTT GAA CAG ATT TTG CAA	1068
Asn Lys Phe Lys Asn Gln Thr Ala Leu Lys Val Glu Gln Ile Leu Gln	
80 85 90	
AAT CAA GGC TAT AAG GTT ATT AGC GTA GAT AGC AGC GAT AAA GAC GAT	1116
Asn Gln Gly Tyr Lys Val Ile Ser Val Asp Ser Ser Asp Lys Asp Asp	
95 100 105	
TTT TCT TTT GCA CAA AAA AAA GAA GGG TAT TTG GCG GTT GCT ATG AAT	1164
Phe Ser Phe Ala Gln Lys Lys Glu Gly Tyr Leu Ala Val Ala Met Asn	
110 115 120	
GGC GAA ATT GTT TTA CGC CCC GAT CCT AAA AGG ACC ATA CAG AAA AAA	1212
Gly Glu Ile Val Leu Arg Pro Asp Pro Lys Arg Thr Ile Gln Lys Lys	
125 130 135 140	
TCA GAA CCC GGG TTA TTA TTC TCC ACC GGT TTG GAC AAA ATG GAA GCG	1260
Ser Glu Pro Gly Leu Leu Phe Ser Thr Gly Leu Asp Lys Met Glu Gly	
145 150 155	

-34-

GTT TTA ATC CCG GCT GGG TTT ATT AAG GTT ACC ATA CTA GAG CCT ATG Val Leu Ile Pro Ala Gly Phe Ile Lys Val Thr Ile Leu Glu Pro Met 160 165 170	1308
AGT GGG GAA TCT TTG GAT TCT TTT ACG ATG GAT TTG AGC GAG TTG GAC Ser Gly Glu Ser Leu Asp Ser Phe Thr Met Asp Leu Ser Glu Leu Asp 175 180 185	1356
ATT CAA GAA AAA TTC TTA AAA ACC ACC CAT TCA AGC CAT AGC GGG GGG Ile Gln Glu Lys Phe Leu Lys Thr Thr His Ser Ser His Ser Gly Gly 190 195 200	1404
TTA GTT AGC ACT ATG GTT AAG GGA ACG GAT AAT TCT AAT GAC GCG ATC Leu Val Ser Thr Met Val Lys Gly Thr Asp Asn Ser Asn Asp Ala Ile 205 210 215 220	1452
AAG AGC GCT TTG AAT AAG ATT TTT GCA AAT ATC ATG CAA GAA ATA GAC Lys Ser Ala Leu Asn Lys Ile Phe Ala Asn Ile Met Gln Glu Ile Asp 225 230 235	1500
AAA AAA CTC ACT CAA AAG AAT TTA GAA TCT TAT CAA AAA GAC GCC AAA Lys Lys Leu Thr Gln Lys Asn Leu Glu Ser Tyr Gln Lys Asp Ala Lys 240 245 250	1548
GAA TTA AAA GGC AAA AGA AAC CGA TAA AAACAAATAA CGCATAAGAA Glu Leu Lys Gly Lys Arg Asn Arg * 255 260	1595
AAGAACGCTT GAATAAACTG CTTAAAAAGG GTTTTTFAGC GTTCTTTTTC AGCGTGTATT	1655
TAAGGGCTGA TGATC	1670

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 261 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Arg Ala Asn Asn His Phe Lys Asp Phe Ala Trp Lys Lys Cys Leu 1 5 10 15
Leu Gly Ala Ser Val Val Ala Leu Leu Val Gly Cys Ser Pro His Ile 20 25 30
Ile Glu Thr Asn Glu Val Ala Leu Lys Leu Asn Tyr His Pro Ala Ser 35 40 45
Glu Lys Val Gln Ala Leu Asp Glu Lys Ile Leu Leu Leu Arg Pro Ala 50 55 60
Phe Gln Tyr Ser Asp Asn Ile Ala Lys Glu Tyr Glu Asn Lys Phe Lys 65 70 75 80

-35-

Asn	Gln	Thr	Ala	Leu	Lys	Val	Glu	Gln	Ile	Leu	Gln	Asn	Gln	Gly	Tyr	85	90	95	
Lys	Val	Ile	Ser	Val	Asp	Ser	Ser	Asp	Lys	Asp	Asp	Phe	Ser	Phe	Ala	100	105	110	
Gln	Lys	Lys	Glu	Gly	Tyr	Leu	Ala	Val	Ala	Met	Asn	Gly	Glu	Ile	Val	115	120	125	
Leu	Arg	Pro	Asp	Pro	Lys	Arg	Thr	Ile	Gln	Lys	Lys	Ser	Glu	Pro	Gly	130	135	140	
Leu	Leu	Phe	Ser	Thr	Gly	Leu	Asp	Lys	Met	Glu	Gly	Val	Leu	Ile	Pro	145	150	155	160
Ala	Gly	Phe	Ile	Lys	Val	Thr	Ile	Leu	Glu	Pro	Met	Ser	Gly	Glu	Ser	165	170	175	
Leu	Asp	Ser	Phe	Thr	Met	Asp	Leu	Ser	Glu	Leu	Asp	Ile	Gln	Glu	Lys	180	185	190	
Phe	Leu	Lys	Thr	Thr	His	Ser	Ser	His	Ser	Gly	Gly	Leu	Val	Ser	Thr	195	200	205	
Met	Val	Lys	Gly	Thr	Asp	Asn	Ser	Asn	Asp	Ala	Ile	Lys	Ser	Ala	Leu	210	215	220	
Asn	Lys	Ile	Phe	Ala	Asn	Ile	Met	Gln	Glu	Ile	Asp	Lys	Lys	Leu	Thr	225	230	235	240
Gln	Lys	Asn	Leu	Glu	Ser	Tyr	Gln	Lys	Asp	Ala	Lys	Glu	Leu	Lys	Gly	245	250	255	
Lys	Arg	Asn	Arg	*												260			

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1670 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 793..1575

(ix) FEATURE:

- (A) NAME/KEY: mat_peptide
- (B) LOCATION: 793..1572

-36-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GATCCTATCG CGCCAAAGGT GGTATTAGGA ATAAGAGCTT GATTATTAAT CTCCCTGGTA	60
AGTCCAAAAA GTATTAGAGA ATGCTTAGAG GCGGTTTTTC CAGCGATTCC TTATTGCGTG	120
GATTTGATTT TAGGGAATTA CATGCAAGTG AATGAAAAAA ACATTCAAGC GTTTGCCCCC	180
AAACAATAAG GTAAAAATG CCACTCACTC ATTTGAATGA AGAAAATCAA CCTAAAATGG	240
TGGATATAGG GGATAAAGAA ACCACTGAAA GAATCGCTCT AGCAAGCGGT CGTATCAGCA	300
TGAATAAAGA GGCTTATGAC GCTATTATCA ATCATGGCGT CAAAAGGGT CCGGTATTAC	360
AAACTGCTAT TATTGCTGGG ATTATGGGGG CTAAAAAGAC AAGCGAAGTC ATTCCCATGT	420
GCCATCCAAT CATGCTCAAT GGGGTGGATA TTGATATTTT AGAAGAAAAA GAGACTTGTA	480
GTTTTAAACT CTATGCGAGA GTCAAACTC AAGCTAAAAC GGGCGTAGAA ATGGAAGCGC	540
TAATGAGTGT GAGCGTAGGG CTTTAAACCA TTTATGACAT GGTGAAAGCC ATTGATAAGA	600
GCATGACAAT TAGCGGTGTG ATGCTGGAAT ATAAAAGTGG AGGCAAAAGT GGGGATTATA	660
ACGCTAAAAA ATAGAAAAAG ACTGATAATC TAAAGATATT AGGGTAAAAT AACATTTTGA	720
CAACAAAAGC GTGTTGGTTG CTTCCGATTT GTTGTATAG AAGTCTAAAA TATTACAATC	780
AAGGATAGAA CG ATG AGA GCA AAT AAT CAT TTT AAA GAT TTT GCA TGG	828
Met Arg Ala Asn Asn His Phe Lys Asp Phe Ala Trp	
1 5 10	
AAA AAA TGC CTT TTA GGC GCG AGC GTG GTG GCT TTA TTA GTG GGA TGC	876
Lys Lys Cys Leu Leu Gly Ala Ser Val Val Ala Leu Leu Val Gly Cys	
15 20 25	
AGC CCG CAT ATT ATT GAA ACC AAT GAA GTC GCT TTG AAA TTG AAT TAC	924
Ser Pro His Ile Ile Glu Thr Asn Glu Val Ala Leu Lys Leu Asn Tyr	
30 35 40	
CAT CCA GCT AGC GAG AAA GTT CAA GCG TTA GAT GAA AAG ATT TTG CTT	972
His Pro Ala Ser Glu Lys Val Gln Ala Leu Asp Glu Lys Ile Leu Leu	
45 50 55 60	
TTA AGG CCA GCT TTC CAA TAT AGC GAT AAT ATC GCT AAA GAG TAT GAA	1020
Leu Arg Pro Ala Phe Gln Tyr Ser Asp Asn Ile Ala Lys Glu Tyr Glu	
65 70 75	
AAC AAA TTC AAG AAT CAA ACC GCG CTC AAG GTT GAA CAG ATT TTG CAA	1068
Asn Lys Phe Lys Asn Gln Thr Ala Leu Lys Val Glu Gln Ile Leu Gln	
80 85 90	
AAT CAA GGC TAT AAG GTT ATT AGC GTA GAT AGC AGC GAT AAA GAC GAT	1116
Asn Gln Gly Tyr Lys Val Ile Ser Val Asp Ser Ser Asp Lys Asp Asp	
95 100 105	

-37-

TTT TCT TTT GCA CAA AAA AAA GAA GGG TAT TTG GCG GTT GCT ATG AAT Phe Ser Phe Ala Gln Lys Lys Glu Gly Tyr Leu Ala Val Ala Met Asn 110 115 120	1164
GGC GAA ATT GTT TTA CGC CCC GAT CCT AAA AGG ACC ATA CAG AAA AAA Gly Glu Ile Val Leu Arg Pro Asp Pro Lys Arg Thr Ile Gln Lys Lys 125 130 135 140	1212
TCA GAA CCC GGG TTA TTA TTC TCC ACC GGT TTG GAC AAA ATG GAA GGG Ser Glu Pro Gly Leu Leu Phe Ser Thr Gly Leu Asp Lys Met Glu Gly 145 150 155	1260
GTT TTA ATC CCG GCT GGG TTT ATT AAG GTT ACC ATA CTA GAG CCT ATG Val Leu Ile Pro Ala Gly Phe Ile Lys Val Thr Ile Leu Glu Pro Met 160 165 170	1308
AGT GGG GAA TCT TTG GAT TCT TTT ACG ATG GAT TTG AGC GAG TTG GAC Ser Gly Glu Ser Leu Asp Ser Phe Thr Met Asp Leu Ser Glu Leu Asp 175 180 185	1356
ATT CAA GAA AAA TTC TTA AAA ACC ACC CAT TCA AGC CAT AGC GGG GGG Ile Gln Glu Lys Phe Leu Lys Thr Thr His Ser Ser His Ser Gly Gly 190 195 200	1404
TTA GTT AGC ACT ATG GTT AAG GGA ACG GAT AAT TCT AAT GAC GCG ATC Leu Val Ser Thr Met Val Lys Gly Thr Asp Asn Ser Asn Asp Ala Ile 205 210 215 220	1452
AAG AGA GCT TTG AAT AAG ATT TTT GCA AAT ATC ATG CAA GAA ATA GAC Lys Arg Ala Leu Asn Lys Ile Phe Ala Asn Ile Met Gln Glu Ile Asp 225 230 235	1500
AAA AAA CTC ACT CAA AAG AAT TTA GAA TCT TAT CAA AAG GAC GCC AAA Lys Lys Leu Thr Gln Lys Asn Leu Glu Ser Tyr Gln Lys Asp Ala Lys 240 245 250	1548
GAA TTA AAA GGC AAA AGA AAC CGA TAA AAACAAATAA CGCATAAGAA Glu Leu Lys Gly Lys Arg Asn Arg * 255 260	1595
AAGAACGCTT GAATAAACTG CTTAAAAAGG GTTTTTTACG GTTCTTTTTG AGCGTGATT	1655
TAAGGGCTGA TGATC	1670

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 261 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Arg Ala Asn Asn His Phe Lys Asp Phe Ala Trp Lys Lys Cys Leu
1 5 10 15

-38-

Leu Gly Ala Ser Val Val Ala Leu Leu Val Gly Cys Ser Pro His Ile
 20 25 30

Ile Glu Thr Asn Glu Val Ala Leu Lys Leu Asn Tyr His Pro Ala Ser
 35 40 45

Glu Lys Val Gln Ala Leu Asp Glu Lys Ile Leu Leu Leu Arg Pro Ala
 50 55 60

Phe Gln Tyr Ser Asp Asn Ile Ala Lys Glu Tyr Glu Asn Lys Phe Lys
 65 70 75 80

Asn Gln Thr Ala Leu Lys Val Glu Gln Ile Leu Gln Asn Gln Gly Tyr
 85 90 95

Lys Val Ile Ser Val Asp Ser Ser Asp Lys Asp Asp Phe Ser Phe Ala
 100 105 110

Gln Lys Lys Glu Gly Tyr Leu Ala Val Ala Met Asn Gly Glu Ile Val
 115 120 125

Leu Arg Pro Asp Pro Lys Arg Thr Ile Gln Lys Lys Ser Glu Pro Gly
 130 135 140

Leu Leu Phe Ser Thr Gly Leu Asp Lys Met Glu Gly Val Leu Ile Pro
 145 150 155 160

Ala Gly Phe Ile Lys Val Thr Ile Leu Glu Pro Met Ser Gly Glu Ser
 165 170 175

Leu Asp Ser Phe Thr Met Asp Leu Ser Glu Leu Asp Ile Gln Glu Lys
 180 185 190

Phe Leu Lys Thr Thr His Ser Ser His Ser Gly Gly Leu Val Ser Thr
 195 200 205

Met Val Lys Gly Thr Asp Asn Ser Asn Asp Ala Ile Lys Arg Ala Leu
 210 215 220

Asn Lys Ile Phe Ala Asn Ile Met Gln Glu Ile Asp Lys Lys Leu Thr
 225 230 235 240

Gln Lys Asn Leu Glu Ser Tyr Gln Lys Asp Ala Lys Glu Leu Lys Gly
 245 250 255

Lys Arg Asn Arg *
 260

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>30</u> , line <u>17-22</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet ☐	
Name of depositary institution The National Collections of Industrial and Marine Bacteria Limited (NCIMB)	
Address of depositary institution (including postal code and country) 23 St Machar Drive Aberdeen AB2 1RY Scotland, UK	
Date of deposit 16 May 1995	Accession Number NCIMB 40732
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet ☐	
In respect of all designated states in which such action is possible and to the extent that it is legally permissible under the law of the designated state, it is requested that a sample of the deposited micro-organism be made available only by the issue thereof to an independent expert, in accordance with the relevant patent legislation, e.g. Rule 28(4) EPC, and generally similar provisions <i>mutatis mutandis</i> for any other designated state.	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
☒ This sheet was received with the international application	☐ This sheet was received by the International Bureau on:
Authorized officer 	Authorized officer



CLAIMS

1. A recombinant polypeptide which has an amino acid sequence identical with, or substantially similar to, a *Helicobacter pylori* surface-exposed antigen with an approximate molecular weight of 29 kDa.
2. A polypeptide according to claim 1 which amino acid sequence is identical with, or substantially similar to, positions 1-260 or 28-260 in SEQ ID NO: 2 or SEQ ID NO: 4 in the Sequence Listing.
3. A peptide with a length of at least 5 amino acids comprising an immunogenic epitope of a polypeptide according to claim 1.
4. An isolated nucleic acid molecule which has a nucleotide sequence coding for a polypeptide according to claim 1 or 2.
5. An isolated nucleic acid molecule selected from:
 - (a) nucleic acid molecules comprising a nucleotide sequence which is identical with, or substantially similar to, positions 796-1572 or 874-1572 in SEQ ID NO: 1 or SEQ ID NO: 3 in the Sequence Listing;
 - (b) nucleic acid molecules comprising a nucleotide sequence capable of hybridizing to a nucleotide sequence complementary to the polypeptide coding region of a nucleic acid molecule as defined in (a) and which codes for a polypeptide according to claim 1 or 2, or a functionally equivalent modified form thereof; and
 - (c) nucleic acid molecules comprising a nucleic acid sequence which is degenerate as a result of the genetic code to a nucleotide sequence as defined in (a) or (b) and which codes for a polypeptide according to claim 1 or 2, or a functionally equivalent modified form thereof.





6. A vector which comprises the nucleic acid molecule according to claim 4 or 5.
- 5 7. A vector according to claim 6 which is the plasmid vector pAE1 (NCIMB 40732).
8. A vector according to claim 6 which is an expression vector capable of mediating the expression of a DNA molecule according to claim 4 or 5.
- 10 9. A vector according to claim 8 which is the plasmid vector pS863.
10. A host cell harbouring a vector according to any one of claims 6 to 9.
- 15 11. A process for production of a polypeptide which is a *Helicobacter pylori* surface-exposed 29 kDa antigen, which comprises culturing a host cell transformed with an expression vector according to claim 8 or 9 under conditions whereby said polypeptide is produced, and recovering said polypeptide.
- 20 12. A polypeptide or peptide according to any one of claims 1 to 3 for use in therapy.
13. A polypeptide or peptide according to any one of claims 1 to 3 for use in the diagnosis of *Helicobacter pylori* infection.
- 25 14. A polypeptide or peptide according to any one of claims 1 to 3 for use as a vaccine.





15. A vaccine composition for inducing a protective immune response to *Helicobacter pylori* infection, comprising an immunogenically effective amount of a polypeptide according to any one of claims 1 to 3, or a modified form of said polypeptide which retains the capability to induce protective immunity against *Helicobacter pylori* infection, optionally together with a pharmaceutically acceptable carrier or diluent.
16. A vaccine composition according to claim 15 for use as a therapeutic vaccine in a mammal, including man, which is infected by *Helicobacter pylori*.
17. A vaccine composition according to claim 15 for use as a prophylactic vaccine to protect a mammal, including man, from infection by *Helicobacter pylori*.
18. Use of a polypeptide according to any one of claims 1 to 3, or a modified form of said antigen which retains the capability to induce protective immunity against *Helicobacter pylori* infection, in the manufacture of a composition for the treatment, prophylaxis or diagnosis of *Helicobacter pylori* infection.
19. Use of a polypeptide according to any one of claims 1 to 3, or a modified form of said antigen which retains the capability to induce protective immunity against *Helicobacter pylori* infection, in the manufacture of a diagnostic kit for diagnosis of *Helicobacter pylori* infection.
20. Use of a polypeptide according to any one of claims 1 to 3, or a modified form of said polypeptide which retains the capability to induce protective immunity against *Helicobacter pylori* infection, in





the manufacture of a vaccine for use in eliciting a protective immune response against *Helicobacter pylori*.

- 5 21. A method of eliciting in a mammal a protective immune response against *Helicobacter pylori* infection, said method comprising the step of administering to the said mammal an immunologically effective amount of a vaccine composition according to any one of claims 15 to 17.
- 10 22. A method according to claim 21 wherein the said mammal is a human.
- 15 23. A method of *in vitro* diagnosis of *Helicobacter pylori* infection comprising at least one step wherein a polypeptide according to any one of claims 1 to 3, optionally labelled or coupled to a solid support, is used.
- 20 24. A method according to claim 23 comprising the steps
(a) contacting a said polypeptide, optionally bound to a solid support, with a body fluid taken from a mammal; and
(b) detecting antibodies from the said body fluid binding to the said polypeptide.
- 25 25. A diagnostic kit for the detection of *Helicobacter pylori* infection in a mammal, including man, comprising components which enable the method according to claim 23 or 24 to be carried out.

