

Description


This invention relates to a process for preparing monoacylated hydrazines. More particularly, this invention relates to a process which comprises reacting a trichloromethyl aryl ketone with a monoalkylated hydrazine to obtain a monoacyl-monoalkylhydrazine. The monoacyl-monoalkylhydrazines are useful intermediates in the process of preparing 1-alkyl-1,2-diacylhydrazines which are known to have insecticidal activity against Coleoptera and Lepidoptera.

Synthetic Communications, 17(14), 1741-1748 (1987) discusses the acylation of various nitrogen nucleophiles with 2,2,2-trichloro-1-(p-substituted phenyl)ethanones where two examples illustrate the acylation of unsubstituted hydrazine and of phenylhydrazine.

Selectivity is a problem when acylating hydrazines since hydrazine is a difunctional molecule. There are two types of selectivity problems. One is the selectivity to monoacylate hydrazine while preventing or minimizing diacylation. The other is control of the regioselectivity of acylation for monosubstituted hydrazines. The process of the present invention provides an efficient method of selectively monoacylating, and causing acylation on the unsubstituted nitrogen, of monoalkylated hydrazines.

In the process of the present invention, the monoalkylated hydrazine is reacted with a trichloromethyl aryl ketone to obtain the desired monoacylated hydrazine derivative.


Accordingly the present invention provides a process for preparing a monoacylated hydrazine of the formula



which comprises reacting a hydrazine of the formula



or the corresponding hydrazine hydrate or hydrazine salt with a trichloromethyl aryl ketone of the formula



wherein:

R is (C₁-C₆)alkyl; and

Aryl is (i) phenyl optionally substituted with one to three substituents which may be the same or different and which are selected from halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl or halo(C₁-C₆)alkoxy; or (ii) naphthyl.

(C₁-C₆)alkyl includes straight or branched alkyl groups, for example, methyl, ethyl, *n*-propyl, *n*-butyl, isopropyl, *t*-butyl, or neopentyl. (C₁-C₆)alkoxy includes, for example, methoxy. Halo means bromo, chloro, fluoro and iodo. Halo(C₁-C₆)alkyl includes, for example, trifluoromethyl. Halo(C₁-C₆)alkoxy includes, for example, trifluoromethoxy.

In one preferred embodiment, R is *t*-butyl or methyl.

In another preferred embodiment, Aryl is 4-((C₁-C₆)alkyl)phenyl, 2,3-di(C₁-C₆)alkylphenyl, 4-halophenyl, 2-(C₁-C₆)alkyl-3-(C₁-C₆)alkoxyphenyl, 2-(C₁-C₆)alkyl-3-halophenyl, 2,3-dihalophenyl, 2-halo-3-(C₁-C₆)alkylphenyl, 2,3-di(C₁-C₆)alkoxyphenyl, 2-halo-3-(C₁-C₆)alkoxyphenyl or 2,3,5-tri(C₁-C₆)alkylphenyl.

More preferably, Aryl is 4-ethylphenyl, 4-chlorophenyl, 2,3-dimethylphenyl, 2-methyl-3-methoxyphenyl, 4-methylphenyl, 4-*n*-propylphenyl, 2-methyl-3-chlorophenyl, 2,3-dimethoxyphenyl, 2-methyl-3-bromophenyl, 2-methyl-3-fluorophenyl, 2,3-dichlorophenyl, 2-fluoro-3-chlorophenyl, 2,3-difluorophenyl, 4-*isopropylphenyl*, 2-chloro-3-methylphenyl, 2-bromo-3-methylphenyl, 2-chloro-3-methoxyphenyl, 2-ethyl-3-chlorophenyl, 2-fluoro-3-methylphenyl or 2,3-dimethyl-5-*isopropylphenyl*.

The hydrazine used in the process can be a hydrate, the neat hydrazine such as methylhydrazine, or a hydrazine salt. For example, the hydrazine salt may be a hydrazine hydrohalide such as *t*-butylhydrazine hydrochloride. In the case where a hydrazine salt is used, an equivalent of base is added to the reaction to produce the free hydrazine. Examples of bases include potassium carbonate, sodium acetate, sodium

methoxide, sodium ethoxide, triethylamine, and sodium hydroxide. The preferred base is sodium hydroxide.

The reaction process may be carried out in a variety of solvents such as methanol, ethanol, isopropanol, xylene, water, ethyl acetate, toluene and methylene chloride. Preferred solvents are aprotic solvents such as xylene, toluene and methylene chloride.

5 The reaction process is carried out preferably at atmospheric pressure.

The process may be carried out between about 10 °C and about 100 °C, preferably between about 20 °C and about 50 °C.

The following Examples further illustrate the invention but are not intended to limit it in any way.

10 EXAMPLES

Example 1: 1-(4-Ethylbenzoyl)-2-*t*-butylhydrazine

To a mixture of *t*-butylhydrazine hydrochloride (6.36 grams (g), 50 millimoles (mmol)) 5.3 g of water, 20 milliliters (ml) of methylene chloride and 4.0 g of 50% sodium hydroxide solution (50 mmol) under nitrogen there was added dropwise at room temperature over 12 minutes α,α,α -trichloro-4-ethylacetophenone (13.88 g, 95.2% purity, 52.5 mmol). The reaction mixture was stirred at room temperature for 5.75 hours, then 0.4 g of additional sodium hydroxide solution was added and the reaction mixture was stirred overnight. The reaction was quenched with water and the phases were separated. The methylene chloride phase was 20 washed twice with water, dried over magnesium sulfate, filtered and evaporated *in vacuo* to yield 10.24 g (93% yield) of a pale yellow solid. This was found to contain 92.5% of 1-(4-ethyl)benzoyl-2-*t*-butylhydrazine, less than 0.3% of the isomer 1-(4-ethyl)benzoyl-1-*t*-butylhydrazine, and less than 0.3% of the 1,2-di-(4-ethyl)benzoyl-1-*t*-butylhydrazine.

25 Example 2: 1-(4-Chlorobenzoyl)-2-methylhydrazine

To a mixture of methylhydrazine (1.63 g, 34.6 mmol) and 55 ml of toluene under nitrogen, α,α,α -trichloro-4-chloroacetophenone (8.55 g, 97.9% purity, 34.6 mmol) was added dropwise and the reaction mixture was stirred at room temperature for 2.75 hours. The resulting slurry was filtered and the 30 solids were washed with toluene and dried to yield 3.82 g (70% yield) of 1-(4-chlorobenzoyl)-2-methylhydrazine, mp 129.5-131 °C (literature mp 132-133 °C, Meyer, R. F., J. Heterocyclic Chem., 1965, 2, 305).

Using essentially the reaction conditions described in Examples 1 and 2 the compounds listed in Table I were prepared. Where reaction conditions differed, they are noted in the Table. The products of the reactions listed in this Table were isolated either by filtering from the reaction solvent, or by extraction into 35 acid, followed by neutralization and extraction of the product from the aqueous phase. The reported melting points are for the isolated material which had not been further purified by recrystallization.

TABLE I

Ex. No.	Aryl	R	Solvent Temp.	Product Isolation	Yield	mp (°C)	lit mp (°C)
3.	Phenyl	CH ₃	Toluene RT ²	Extraction	73	79.5-82	85-86 ¹ 86-88 ³
4.	Phenyl	C(CH ₃) ₃	Toluene 40 °C	Extraction	75	93.5-94.5	94-95 ⁴
5.	4-Ethylphenyl	CH ₃	Toluene RT	Extraction	81	65-71	
45 6.	4-Ethylphenyl	C(CH ₃) ₃	Toluene 40 °C	Filtration	76		

1. Meyer, R. F., J. Heterocyclic Chem., 1965, 2, 305.

2. RT = Room Temperature

3. Smith, P. A. S., "Derivatives of Hydrazine and other Hydronitrogens having N-N bonds", Benjamin/Cummings Publishing Company, Reading, MA, 1983, p 83.

50 4. MacLeay et al, U.S. Patent 4,008,273.

By using essentially the reaction conditions described in Examples 1 and 2, Examples 8-25, listed in Table II, are prepared.

Table II

<u>Ex. No.</u>	<u>Aryl</u>	<u>R</u>	<u>Solvent Temp.</u>
7.	Phenyl	CH ₃	Xylene RT
8.	2,3-Dimethyl- phenyl	C(CH ₃) ₃	Toluene 40°C
9.	2-Methyl-3- methoxyphenyl	C(CH ₃) ₃	Toluene 40°C

Table II cont'd


Ex. No.	Aryl	R	Solvent Temp.
10.	4-Methylphenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
11.	4- <i>n</i> -Propylphenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
12.	2-Methyl-3-chloro- phenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
13.	2,3-Dimethoxy- phenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
14.	2-Methyl-3-bromo- phenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
15.	2-Methyl-3-fluoro- phenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
16.	2,3-Dichloro- phenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
17.	2-Fluoro-3-chloro- phenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT
18.	2,3-Difluorophenyl	C(CH ₃) ₃	Toluene RT
19.	4-Isopropylphenyl	C(CH ₃) ₃	Toluene RT
20.	2-Chloro-3-methyl- phenyl	C(CH ₃) ₃	Toluene RT
21.	2-Bromo-3-methyl- phenyl	C(CH ₃) ₃	Toluene RT
22.	2-Chloro-3-methoxy- phenyl	C(CH ₃) ₃	Toluene RT
23.	2-Ethyl-3-chloro- phenyl	C(CH ₃) ₃	Toluene RT
24.	2-Fluoro-3-methyl- phenyl	C(CH ₃) ₃	Toluene RT

Table II cont'd

Ex. No.	Aryl	R	Solvent Temp.
25.	2,3-Dimethyl- 5-isopropylphenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT

Claims


1. A process for preparing a monoacylated hydrazine of the formula



which comprises reacting a hydrazine of the formula



or the corresponding hydrazine hydrate or hydrazine salt with a trichloromethyl aryl ketone of the formula



wherein:

R is (C₁-C₆)alkyl; and

Aryl is (i) phenyl optionally substituted with one to three substituents which may be the same or different and which are selected from halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl or halo(C₁-C₆)alkoxy, or (ii) naphthyl.

2. A process as claimed in claim 1, wherein R is *t*-butyl or methyl.
3. A process as claimed in claim 1 or 2, wherein Aryl is 4-(C₁-C₆)alkylphenyl, 2,3-di(C₁-C₆)alkylphenyl, 2-(C₁-C₆)alkyl-3-(C₁-C₆)alkoxyphenyl, 4-halophenyl, 2,3-dihalophenyl, 2-(C₁-C₆)alkyl-3-halophenyl, 2,3-di(C₁-C₆)alkoxyphenyl, 2-halo-3-(C₁-C₆)alkylphenyl, 2-halo-3-(C₁-C₆)alkoxyphenyl or 2,3,5-tri(C₁-C₆)alkylphenyl.
4. A process as claimed in claim 3, wherein Aryl is 4-ethylphenyl, 4-methylphenyl, 4-*n*-propylphenyl, 4-isopropylphenyl, 2,3-dimethylphenyl, 2-methyl-3-methoxyphenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,3-difluorophenyl, 2-fluoro-3-chlorophenyl, 2-methyl-3-chlorophenyl, 2-methyl-3-bromophenyl, 2-methyl-3-fluorophenyl, 2-ethyl-3-chlorophenyl, 2,3-dimethoxyphenyl, 2-chloro-3-methylphenyl, 2-bromo-3-methylphenyl, 2-fluoro-3-methylphenyl, 2-chloro-3-methoxyphenyl or 2,3-dimethyl-5-isopropylphenyl.
5. A process as claimed in any preceding claim, wherein said hydrazine is methyl hydrazine or a hydrazine hydrate.

Table II cont'd

Ex. No.	Aryl	R	Solvent Temp.
25.	2,3-Dimethyl- 5-isopropylphenyl	C(CH ₃) ₃	CH ₂ Cl ₂ /NaOH RT

Claims


1. A process for preparing a monoacylated hydrazine of the formula



which comprises reacting a hydrazine of the formula



or the corresponding hydrazine hydrate or hydrazine salt with a trichloromethyl aryl ketone of the formula



wherein:

R is (C₁-C₆)alkyl; and

Aryl is (i) phenyl optionally substituted with one to three substituents which may be the same or different and which are selected from halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl or halo(C₁-C₆)alkoxy, or (ii) naphthyl.

2. A process as claimed in claim 1, wherein R is *t*-butyl or methyl.

3. A process as claimed in claim 1 or 2, wherein Aryl is 4-(C₁-C₆)alkylphenyl, 2,3-di(C₁-C₆)alkylphenyl, 2-(C₁-C₆)alkyl-3-(C₁-C₆)alkoxyphenyl, 4-halophenyl, 2,3-dihalophenyl, 2-(C₁-C₆)alkyl-3-halophenyl, 2,3-di(C₁-C₆)alkoxyphenyl, 2-halo-3-(C₁-C₆)alkylphenyl, 2-halo-3-(C₁-C₆)alkoxyphenyl or 2,3,5-tri(C₁-C₆)alkylphenyl.

4. A process as claimed in claim 3, wherein Aryl is 4-ethylphenyl, 4-methylphenyl, 4-*n*-propylphenyl, 4-isopropylphenyl, 2,3-dimethylphenyl, 2-methyl-3-methoxyphenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,3-difluorophenyl, 2-fluoro-3-chlorophenyl, 2-methyl-3-chlorophenyl, 2-methyl-3-bromophenyl, 2-methyl-3-fluorophenyl, 2-ethyl-3-chlorophenyl, 2,3-dimethoxyphenyl, 2-chloro-3-methylphenyl, 2-bromo-3-methylphenyl, 2-fluoro-3-methylphenyl, 2-chloro-3-methoxyphenyl or 2,3-dimethyl-5-isopropylphenyl.

5. A process as claimed in any preceding claim, wherein said hydrazine is methyl hydrazine or a hydrazine hydrate.



6. A process as claimed in any of claims 1 to 4, wherein the hydrazine is a hydrazine salt, and an equivalent of base, preferably potassium carbonate, sodium acetate, sodium methoxide, sodium ethoxide, sodium hydroxide or triethylamine, and more preferably sodium hydroxide, is added to the reaction mixture to produce free hydrazine.
7. A process as claimed in claim 6, wherein said hydrazine salt is a hydrazine hydrohalide, preferably *t*-butylhydrazine hydrochloride.
8. A process as claimed in any preceding claim, which is carried out in a solvent, preferably methanol, ethanol, isopropanol, water, xylene, toluene, methylene chloride or ethyl acetate, more preferably xylene, toluene or methylene chloride.
9. A process as claimed in any preceding claim, which is carried out at a temperature between 10 °C and 100 °C, preferably between 20 °C and 50 °C.

Patentansprüche


1. Verfahren zur Herstellung eines monoacylierten Hydrazins der Formel



welches Verfahren ein Umsetzen eines Hydrazins der Formel



oder des korrespondierenden Hydrazinhydrates oder Hydrazinsalzes mit einem Trichlormethylarylketon der Formel



worin;

R(C₁-C₆)Alkyl ist, und

Aryl (i) ein Phenyl, welches gegebenenfalls mit einem bis drei Substituenten substituiert sein kann, welche gleich oder verschieden sein können und welche aus Halogen, (C₁-C₆)Alkyl, (C₁-C₆)Alkoxy, Halogen(C₁-C₆)alkyl oder Halogen(C₁-C₆)alkoxy gewählt sind, oder (ii) Naphthyl ist, umfaßt.


2. Verfahren, wie in Anspruch 1 beansprucht, wobei R tert. Butyl oder Methyl ist.
3. Verfahren, wie in Anspruch 1 oder 2 beansprucht, wobei Aryl ist: 4-(C₁-C₆)Alkylphenyl, 2,3-Di(C₁-C₆)alkylphenyl, 2-(C₁-C₆)Alkyl-3-(C₁-C₆)alkoxyphenyl, 4-Halogenphenyl, 2,3-Dihalogenphenyl, 2-(C₁-C₆)Alkyl-3-halogenphenyl, 2,3-Di(C₁-C₆)alkoxyphenyl, 2-Halogen-3-(C₁-C₆)alkylphenyl, 2-Halogen-3-(C₁-C₆)alkoxyphenyl oder 2,3,5-Tri(C₁-C₆)alkylphenyl.
4. Verfahren, wie in Anspruch 3 beansprucht, wobei Aryl ist: 4-Ethylphenyl, 4-Methylphenyl, 4-n-Propylphenyl, 4-Isopropylphenyl, 2,3-Dimethylphenyl, 2-Methyl-3-methoxyphenyl, 4-Chlorphenyl, 2,3-Dichlorphenyl, 2,3-Difluorphenyl, 2-Fluor-3-chlorphenyl, 2-Methyl-3-chlorphenyl, 2-Methyl-3-bromphenyl, 2-Methyl-3-fluorphenyl, 2-Ethyl-3-chlorphenyl, 2,3-Dimethoxyphenyl, 2-Chlor-3-methylphenyl, 2-Brom-3-methylphenyl, 2-Fluor-3-methylphenyl, 2-Chlor-3-methoxyphenyl oder 2,3-Dimethyl-5-isopropylphenyl.



5. Verfahren, wie in einem vorhergehenden Anspruch beansprucht, wobei das Hydrazin Methylhydrazin oder ein Hydrazinhydrat ist.
6. Verfahren, wie in einem der Anspruch 1 bis 4 beansprucht, wobei das Hydrazin ein Hydrazinsalz ist, und ein Äquivalent Base, vorzugsweise Kaliumcarbonat, Natriumacetat, Natriummethoxid, Natriummethoxid, Natriumhydroxid oder Triethylamin, und mehr bevorzugt Natriumhydroxid, dem Reaktionsgemisch zugegeben wird, um freies Hydrazin herzustellen.
7. Verfahren, wie in Anspruch 6 beansprucht, wobei das Hydrazinsalz ein Hydrazinhydrohalogenid, vorzugsweise tert. Butylhydrazinhydrochlorid, ist.
8. Verfahren, wie in einem vorhergehenden Anspruch beansprucht, welches in einem Lösungsmittel, vorzugsweise Methanol, Ethanol, Isopropanol, Wasser, Xylol, Toluol, Methylenchlorid oder Ethylacetat, mehr bevorzugt Xylol, Toluol oder Methylenchlorid, ausgeführt wird.
9. Verfahren, wie in einem vorhergehenden Anspruch beansprucht, welches bei einer Temperatur zwischen 10 °C und 100 °C, vorzugsweise zwischen 20 °C und 50 °C, ausgeführt wird.

Revendications


1. Procédé pour préparer une hydrazine monoacylée de formule



qui consiste à faire réagir une hydrazine de formule



ou le sel d'hydrazine ou l'hydrate d'hydrazine correspondant avec une trichlorométhylarylcétone de formule



dans laquelle:

R est un groupe alkyle en C₁-C₆; et

le groupe aryle est (i) un groupe phényle éventuellement substitué avec un à trois substituants qui peuvent être identiques ou différents et qui sont choisis parmi Les groupes halogéno, alkyle en C₁-C₆, alcoxy en C₁-C₆, halogénoalkyle en C₁ - C₆ ou halogénoalcoxy en C₁-C₆, ou (ii) un groupe naphthyle.

2. Procédé selon la revendication 1, dans lequel R est un groupe t-butyle ou méthyle.

3. Procédé selon la revendication 1 ou 2, dans lequel le groupe aryle est un 4-alkyl(C₁-C₆)phényle, un 2,3-dialkyl(C₁-C₆)phényle, un 2-alkyl(C₁-C₆)-3-alcoxy(C₁-C₆) phényle, 4-halogénophényle, un 2,3-dihalogénophényle, un 2-alkyl(C₁-C₆)-3-halogénophényle, un 2,3-dialcoxy(C₁-C₆)phényle, un 2-halogéno-3-alkyl(C₁-C₆)phényle, un 2-halogéno-3-alcoxy(C₁-C₆)phényle ou un 2,3,5-trialkyl(C₁-C₆)phényle.

4. Procédé selon la revendication 3, dans lequel le groupe aryle est le 4-éthylphényle, le 4-méthylphényle, le 4-n-propylphényle, le 4-isopropylphényle, le 2,3-diméthylphényle, le 2-méthyl-3-méthoxyphényle, le 4-chlorophényle, le 2,3-dichlorophényle, le 2,3-difluorophényle, le 2-fluoro-3-chlorophényle, le 2-méthyl-3-chlorophényle, le 2-méthyl-3-bromophényle, le 2-méthyl-3-fluorophényle, le 2-éthyl-3-chloro-



phényle, le 2,3-diméthoxyphényle, le 2-chloro-3-méthylphényle, le 2-bromo-3-méthylphényle, le 2-fluoro-3-méthylphényle, le 2-chloro-3-méthoxyphényle ou le 2,3-diméthyl-5-isopropylphényle.

5. Procédé selon l'une quelconque des revendications précédentes, dans lequel ladite hydrazine est la méthylhydrazine ou un hydrate d'hydrazine.
6. Procédé selon l'une quelconque des revendications 1 à 4, dans lequel l'hydrazine est un sel d'hydrazine, et un équivalent de base, de préférence le carbonate de potassium, l'acétate de sodium, le méthylate de sodium, l'éthylate de sodium, l'hydroxyde de sodium ou la triéthylamine, plus particulièrement, l'hydroxyde de sodium, est ajouté au mélange réactionnel pour produire de l'hydrazine libre.
7. Procédé selon la revendication 6, dans lequel ledit sel d'hydrazine est un hydrohalogénure d'hydrazine, de préférence le chlorhydrate de t-butylhydrazine.
8. Procédé selon l'une quelconque des revendications précédentes, qui est effectué dans un solvant, de préférence le méthanol, l'éthanol, l'isopropanol, l'eau, le xylène, le toluène, le chlorure de méthylène ou l'acétate d'éthyle, plus particulièrement, le xylène, le toluène ou le chlorure de méthylène.
9. Procédé selon l'une quelconque des revendications précédentes, qui est effectué à une température comprise entre 10 °C et 100 °C, de préférence entre 20 °C et 50 °C.

