

(51) International Patent Classification:
C12Q 1/68 (2006.01)(21) International Application Number:
PCT/US2012/037155(22) International Filing Date:
9 May 2012 (09.05.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/484,198 9 May 2011 (09.05.2011) US(71) Applicant (for all designated States except US): **FLUIDIGM CORPORATION** [US/US]; 7000 Shoreline Court, Suite 100, South San Francisco, California 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LIVAK, Kenneth J.** [US/US]; 7000 Shoreline Court, Suite 100, South San Francisco, California 94080 (US). **MYERS, Stacey** [US/US]; 7000 Shoreline Court, Suite 100, South SanFrancisco, California 94020 (US). **WANG, Xiaohui** [US/US]; 7000 Shoreline Court, Suite 100, South San Francisco, California 94080 (US). **WANG, Jun** [CA/US]; 7000 Shoreline Court, Suite 100, South San Francisco, California 94080 (US).(74) Agents: **APPLE, Randolph T.** et al.; Kilpatrick Townsend & Stockton LLP, Two Embarcadero Center, 8th Floor, San Francisco, California 94111-3834 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: PROBE BASED NUCLEIC ACID DETECTION




FIG. 1

(57) Abstract: The invention provides a method for detecting a target nucleotide sequence by tagging the nucleotide sequence with a nucleotide tag, providing a probe oligonucleotide with a melting temperature T_{m1} , comprising a regulatory sequence and a nucleotide tag recognition sequence; incorporating the probe oligonucleotide into the tagged polynucleotide in a polynucleotide amplification reaction, providing a regulatory oligonucleotide with a melting temperature T_{m2} , comprising a sequence segment that is at least partially complementary to the regulatory sequence, amplifying the tagged target nucleic acid sequence in a PCR amplification reaction using the probe oligonucleotide as a primer, and detecting the amplification product; wherein T_{m1} and T_{m2} are higher than the annealing temperature associated with the polynucleotide amplification reaction.



GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*