

CYCLIN-DEPENDENT KINASE 7 INHIBITORS

Field of the invention

The present invention relates to novel inhibitors of cyclin-dependent protein kinases 7 (CDK7), to pharmaceutically acceptable salts, solvates or stereoisomers thereof, to pharmaceutical compositions comprising the present compounds, to methods for treating diseases or disorders and use of the present compounds as pharmaceutical products for treating diseases or disorders.

Background of the invention

Cyclin-dependent kinase 7 (CDK7) is an enzyme ubiquitously expressed in diverse tissues of the body; it is a member of the cyclin-dependent protein kinase (CDK) family. CDK7, along with cyclin H and MAT1, forms the CDK-activating complex, which regulates progression of cell cycle phases via phosphorylation of other CDKs. CDK7 is a mandatory component of the transcription factor, TFIIH. The latter is involved in transcription initiation and DNA repair processes. Thus, CDK7 plays a crucial role in the regulation of the cell cycle and gene transcription (Ganuza et al. Genetic inactivation of Cdk7 leads to cell cycle arrest and induces premature aging due to adult stem cell exhaustion. *EMBO J.* 2012;31(11):2498-510; Larochelle et al. Cyclin-dependent kinase control of the initiation-to-elongation switch of RNA polymerase II. *Nat Struct Mol Biol.* 2012;19(11):1108-15).

Uncontrolled proliferation and changes in transcription processes are properties characteristic of malignant tumor cells. Excessive activation of CDK7 has been observed in many cancer types, and correlates with an aggressive course of the disease and a poor prognosis to a patient, such as, for example, stomach cancer, squamous cell carcinoma of the esophagus, ER+ and triple negative breast cancer, hepatocellular carcinoma (Liu et al. Cyclin-dependent kinase 7 (CDK7) expression in human hepatocellular carcinoma: association with HCC progression, prognosis and cell proliferative capacity. *Transl Cancer Res.* 2018;7(3):472-479; Wang et al. Upregulation of CDK7 in gastric cancer cell promotes tumor cell proliferation and predicts poor prognosis. *Exp Mol Pathol.* 2016;100(3):514-21; Patel et al. Expression of CDK7, Cyclin H, and MAT1 Is Elevated in Breast Cancer and Is Prognostic in Estrogen Receptor-Positive Breast Cancer. *Clin Cancer Res.* 2016;22(23):5929-5938; Li et al. Therapeutic Rationale to Target Highly Expressed CDK7 Conferring Poor Outcomes in Triple-Negative Breast Cancer. *Cancer Res.* 2017;77(14):3834-3845; Zhang et al. Low expression of cyclinH and cyclin-dependent kinase 7 can decrease the proliferation of human esophageal squamous cell carcinoma. *Dig Dis Sci.* 2013;58(7):2028-37). Inhibition of CDK7 can have an antitumor effect by means of reduced rate of cell proliferation, as well as by affecting the transcription of genes involved in the malignant transformation of cells, such as cells of ovarian cancer, T-cell acute lymphoblastic leukemia, neuroblastoma, glioma, small cell lung cancer and others (Chipumuro et al. CDK7 inhibition suppresses super-enhancer-linked oncogenic transcription in MYCN-driven cancer. *Cell.* 2014;159(5):1126-1139, Kwiatkowski et al. Targeting transcription regulation in cancer with a covalent CDK7 inhibitor. *Nature.* 2014;511(7511):616-20, Greenall et al. Cyclin-dependent kinase 7 is a therapeutic target in high-grade glioma. *Oncogenesis.* 2017;6(5):e336, Christensen et al. Targeting transcriptional addictions in small cell lung cancer with a covalent CDK7 inhibitor. *Cancer Cell.* 2014 Dec 8;26(6):909-922). Furthermore, protein inhibition potentiates genome instability and reduces PD-L1 expression; it triggers anti-tumor immune responses (Zhang et al. CDK7 Inhibition Potentiates Genome Instability Triggering Anti-tumor Immunity in Small Cell Lung Cancer. *Cancer Cell* 2020;37(1):37-54.e9, Wang et al. CDK7 inhibitor THZ1 enhances

antiPD-1 therapy efficacy via the p38 α /MYC/PD-L1 signaling in non-small cell lung cancer. J Hematol Oncol. 2020;13(1):99).

Thus, there is a need to create novel compounds that preferably target CDK7 and can provide for a better risk–benefit ratio in the therapy of diseases associated with increased CDK7 activity.

Description of the invention

The terms used in the description of this invention appear below.

Optionally substituted in one, two, three, four or several positions means the specified group can be substituted by a radical or any combination of radicals in one, two, three, four or from one to six positions.

An "annelated compound" or "condensed compound" is a polycyclic compound that has two neighboring atoms in common. Examples of condensed compounds include, but are not limited to, naphthalene, quinoline, tetrahydroquinoline, isoquinoline, quinoxaline, 1,2,3,4-tetrahydroquinoxaline, 1,2-dihydroquinoline, indole, dihydroindole, 1H-pyrrolo[2,3-b]pyridine, 1H,6H,7H-pyrrolo[2,3-c]pyridine, 1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridine, 1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepine, 1H,6H,7H-pyrrolo[2,3-c]pyridine, 1H,4H,7H,8H-pyrrolo[2,3-c]azepine, 1H,6H,7H,8H-pyrrolo[2,3-c]azepine, 1H,8H-pyrrolo[2,3-c]azepine, 7H,8H-pyrrolo[2,3-c]azepine, 1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocine, 1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,8H,9H-pyrrolo[2,3-c]azocine, 9H-pyrrolo[2,3-c]azocine, 1H,4H,5H,6H-pyrrolo[2,3-c]pyrrole, 1H-1,3-benzodiazole, azaspiro[3.5]nonan, azaspiro[2.5]octane, 1H,2H,3H,4H,5H,6H-pyrrolo[2,3-f][1,4]oxazepine, 3,4,6,13-tetraazatricyclo[8.3.0.0²,⁶]trideca-1(10),2,4,11-tetraen-11-yl, 1,4-diazatricyclo[7.1.1.0³,⁷]undeca-3(7),5-dienyl.

“Alkyl” means an aliphatic straight chain or branched chain hydrocarbon group having from 1 to 12 carbon atoms, more preferably from 1 to 6 carbon atoms. Branched chain means alkyl chain having one or more “lower alkyl” substituents. Examples of alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, 2-pentyl, 3-pentyl, neopentyl, n-hexyl. Alkyl may have substituents which may be same or different structure.

“Cycloalkyl” means a fully saturated carbocyclic ring that contains from 3 to 10 carbon ring atoms. Cycloalkyl may have substituents which may be same or different structure. Cycloalkyl may be annelated with aryl, heteroaryl, heterocyclyl. Examples of cycloalkyl groups include, but are not limited to, monocyclic groups, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl or cyclodecyl, bicyclic groups, such as bicycloheptyl or bicyclooctyl.

“Alkenyl” means a straight chain or branched chain hydrocarbon group having from 2 to 12 carbon atoms, more preferably from 2 to 6 carbon atoms that contains one or more carbon-carbon double bond. Alkenyl may have substituents which may be same or different structure.

“Aryl” means an aromatic monocyclic or polycyclic system having from 6 to 14 carbon atoms, more preferably from 6 to 10 carbon atoms. Examples of aryl groups include, but are not limited to, phenyl, phenylene, benzenetriyl, indanyl, naphthyl, naphthylene, naphthalenetriyl and anthrylene. Aryl may have substituents which may be same or different structure. Aryl can be annelated with a heterocycle, cycloalkyl, or heteroaryl.

“Alkyloxy”, “Alkoxy” or “alkyloxy group” means an alkyl-O- group, wherein alkyl is defined in this section. Examples of alkoxy groups include, but are not limited to, methoxy, ethoxy, n-propoxy, iso-propoxy, n-butoxy, tert-butoxy, iso-butoxy.

“Amino group” means R'R"N-group.

“Aminocarbonyl” means -C(=O)NR'''R''' group.

Examples of R', R'', R''', R''' include, but are not limited to, substituents selected from the group comprising hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, heteroalkyl or R' and R'' together with the carbon atom they are attached to, can form 4-7-membered heterocyclyl or heteroaryl.

"Alkylsulfonyl" (-S(O)₂-(C₁-C₆)alkyl) means "alkyl" or "cycloalkyl", as defined above, linked to the corresponding fragment of a molecule via a sulfonyl group, -SO₂-. Examples of alkylsulfonyls include, but are not limited to, methylsulfonyl, ethylsulfonyl, propylsulfonyl, cyclopropylsulfonyl, etc.

"Alkylsulfonamide" (-NH-S(O)₂-(C₁-C₆)alkyl) means "alkyl" or "cycloalkyl", as defined above, linked to the corresponding fragment of a molecule via a sulfonamide group, -NH-SO₂-. Examples of alkylsulfonamides include, but are not limited to, methylsulfonamide, ethylsulfonamide, propylsulfonamide, cyclopropylsulfonamide, etc.

"Dialkylphosphine oxide" or "dialkylphosphoryl" (-P(O)((C₁-C₆)alkyl)₂) means "alkyl" or "cycloalkyl" as defined above, linked to the corresponding fragment of the molecule via a phosphoryl group. Examples of dialkylphosphine oxide include, but are not limited to, dimethylphosphine oxide, diethylphosphine oxide, methylethylphosphine oxide, dipropylphosphine oxide, dicyclopropylphosphine oxide, etc.

The term “oxo” as used herein relates to the radical =O.

“Lower alkyl” means a straight chain or branched chain alkyl having from 1 to 4 carbon atoms.

“Halo” or “Halogen” (Hal) means fluoro, chloro, bromo and iodo.

“Heterocycle”, “heterocyclyl”, “heterocyclic ring” means a monocyclic or polycyclic system having from 3 to 11 carbon atoms, of which one or more carbon atoms are substituted by a heteroatom, such as nitrogen, oxygen, sulfur. Heterocycle may have one or more substituents which may be same or different structure. Nitrogen and sulfur atoms of heterocycle could be oxidized to N-oxide, S-oxide or S-dioxide. Heterocycle may be saturated, partially saturated or unsaturated. A heterocyclyl may be represented by a bridged-ring compound with one, two or three -CH₂-groups; or it may be annelated with a 4-5-membered heteroaryl, wherein one, two or three carbon atoms are substituted by nitrogen atoms. Heterocycle may be annelated with cycloalkyl, aryl or heteroaryl. Examples of heterocycles include, but are not limited to, azetidine, pyrrolidine, piperidine, 2,8-diazaspiro[4.5]decane, piperazine, morpholine, diazepan, azepan, azabicycloheptan, oxane, 1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,4H,5H,6H-pyrrolo[2,3-c]pyrrole, 1,2,3,4-tetrahydroquinoxaline, tetrahydroquinoline, 1,2-dihydroquinoline, dihydroindole, azaspiro[3.5]nonan, azaspiro[2.5]octane, 5H,6H,7H,8H,9H-[1,2,4]triazolo[4,3-a]azepino, 5H,6H,7H-[1,2,4]triazolo[4,3-a]azepino, 1-azabicyclo[4.1.1]octano, 1-azabicyclo[4.1.1]oct-3-enyl, etc.

“Heteroaryl”, "heteroaryl", "heteroaryl ring" means an aromatic monocyclic or polycyclic system having from 5 to 11 carbon atoms, preferably from 5 to 10, of which one or more carbon atoms are substituted by a heteroatom, such as nitrogen, sulfur or oxygen. Nitrogen atom of heterocycle could be oxidized to N-oxide. Heteroaryl may have one or more substituents which may be same or different structure. Heteroaryl may be annelated with cycloalkyl, aryl or

heterocycle. Examples of heteroaryls include, but are not limited to, 1H-pyrrolo[2,3-b]pyridine, 7H-pyrrolo[2,3-d]pyrimidine, pyrrole, furan, pyridine, pyrazine, pyrimidine, pyridazine, isoxazole, isothiazole, tetrazole, oxazole, 1,2,4-oxadiazole, thiazole, pyrazole, furazan, 1,2,4-triazole, 1,2,3-triazole, 1,2,4-thiadiazole, 2H-1,2,3,4-tetrazole, quinoxaline, imidazo[1,2-a]pyridine, indole, benzimidazole, quinoline, imidazole, thienopyridine, quinazoline, naphthyridine, thienopyrimidine, imidazopyridine, quinoline, isoquinoline, quinoxaline, indole, 1H-pyrrolo[2,3-b]pyridine, 1H,6H,7H-pyrrolo[2,3-c]pyridine, 1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridine, 1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepine, 1H,6H,7H-pyrrolo[2,3-c]pyridine, 1H,4H,7H,8H-pyrrolo[2,3-c]azepine, 1H,6H,7H,8H-pyrrolo[2,3-c]azepine, 1H,8H-pyrrolo[2,3-c]azepine, 7H,8H-pyrrolo[2,3-c]azepine, 1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocine, 1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocine, 1H,8H,9H-pyrrolo[2,3-c]azocine, 9H-pyrrolo[2,3-c]azocine, 1H,4H,5H,6H-pyrrolo[2,3-c]pyrrole, 1H-1,3-benzodiazole, 1H,2H,3H,4H,5H,6H-pyrrolo[2,3-f][1,4]oxazepine, 3,4,6,13-tetraazatricyclo[8.3.0.0^{2,6}]trideka-1(10),2,4,11-tetraenyl, 1,4-diazatricyclo[7.1.1.0^{3,7}]undeca-3(7),5-dienyl, and others.

“Substituent” means a chemical radical attached to a scaffold (fragment).

“Solvate” is a molecular aggregate that consists of the compound of the present invention, including its pharmaceutically acceptable salts, with one or more solvent molecules. The solvent molecules are molecules of common pharmaceutical solvents, known to be safe for recipients, e.g. water, ethanol, ethylene glycol, and the like. Other solvents, such as MeOH, methyl-tert-butyl ether, ethyl acetate, methyl acetate, (S)-propylene glycol, (R)-propylene glycol, 1,4-butanediol, and the like, can be used as intermediate solvates for obtaining more desirable solvates.

The term “hydrate” refers to a complex, wherein the solvent molecule is water.

Solvates and/or hydrates preferably exist in crystalline form.

The terms “bond”, “chemical bond”, or “single bond” refer to a chemical bond of two atoms or two moieties (i.e. groups, fragments) when the atoms joined by the bond are considered to be part of larger substructure.

A chemical bond in the compounds of the present invention shown as " $\equiv$ " means a double bond, wherein the configuration of substituents corresponds to the formula.

A chemical bond in the compounds of the present invention shown as " $\times$ " means a double bond which may have both *cis*- and *trans*-configurations.

The term “stereoisomers” refers to compounds that have identical chemical composition and the same structure, but differ in the spatial arrangement of atoms or groups. Stereoisomers may include geometric isomers, enantiomers, diastereomers.

The term "standard treatment" includes extraction, processing with solutions of acids, alkalis and/or salts to neutralize or otherwise alter pH and/or remove interferents; filtration through Celite, silica gel and other materials; addition of an inert carrier (celite, silica gel); removal of volatile components and solvents in vacuo.

The term "excipient" is used herein to describe any ingredient other than the compound(s) of the invention.

“Pharmaceutical composition” means a composition comprising a compound of the invention and at least one excipient. The excipient may be selected from a group consisting of pharmaceutically acceptable and pharmacologically compatible fillers, solvents, diluents, carriers, auxiliary, distributing and sensing agents, delivery agents, such as preservatives, stabilizers, fillers, disintegrators, moisteners, emulsifiers, suspending agents, thickeners, sweeteners, flavouring

agents, aromatizing agents, antibacterial agents, fungicides, lubricants, and prolonged delivery controllers, the choice and proportions of which depend on the nature and route of administration and dosage. Examples of suspending agents are ethoxylated isostearyl alcohol, polyoxyethene, sorbitol and sorbitol ether, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacant and mixtures thereof as well. Protection against action of microorganisms can be provided by various antibacterial and antifungal agents, such as, for example, parabens, chlorobutanol, sorbic acid, and similar compounds. The composition may also contain isotonic agents, for example, sugars, sodium chloride, and similar compounds. Prolonged action of the composition may be achieved by agents slowing down absorption of active ingredient, for example, aluminum monostearate and gelatine. Examples of suitable carriers, solvents, diluents and delivery agents are water, ethanol, polyalcohols and mixtures thereof, plant oils (such as olive oil) and organic esters (such as ethyl oleate) for injections. Examples of fillers are lactose, milk sugar, sodium citrate, calcium carbonate, calcium phosphate, and the like. Examples of disintegrators and distributors are starch, alginic acid and its salts, silicates, and the like. Examples of lubricants are magnesium stearate, sodium lauryl sulfate, talc, and polyethylene glycol of high molecular weight as well. The pharmaceutical composition for peroral, sublingual, transdermal, intramuscular, intravenous, subcutaneous, local or rectal administration of active ingredient, alone or in combination with another active ingredient may be administered to animals and human in a standard administration form, in a mixture with traditional pharmaceutical carriers. Suitable standard administration forms include peroral forms such as tablets, gelatin capsules, pills, powders, granules, chewing gums and peroral solutions or suspensions; sublingual and transbuccal administration forms; aerosols; implants; local, transdermal, subcutaneous, intramuscular, intravenous, intranasal or intraocular administration forms and rectal administration forms.

“Pharmaceutically acceptable salt” means relatively nontoxic both organic and inorganic salts of acids and bases disclosed in the present invention. These salts may be prepared in situ in the processes of synthesis, isolation or purification of compounds or they may be prepared specially. In particular, salts of bases may be prepared specially from purified base of the disclosed compound and suitable organic or inorganic acid. Examples of salts prepared in this manner are hydrochlorides, hydrobromides, sulfates, bisulfates, phosphates, nitrates, acetates, oxalates, valerates, oleates, palmitates, stearates, laurates, borates, benzoates, lactates, tosylates, citrates, maleates, fumarates, succinates, tartrates, mesitates, malonates, salicylates, propionates, ethane sulphonates, benzene sulfonates, sulfamates and the like (Detailed description of such salts properties is given in: Berge S.M., et al., “Pharmaceutical Salts” J. Pharm. Sci. 1977, 66: 1-19). Salts of disclosed acids may be prepared by reaction of a purified acid with suitable base; furthermore, metal salts and amine salts may be synthesized as well. Metal salts are salts of sodium, potassium, calcium, barium, zinc, magnesium, lithium and aluminum; sodium and potassium salts are most preferred. Suitable inorganic bases from which metal salts may be prepared are: sodium hydroxide, carbonate, bicarbonate and hydride; potassium hydroxide and bicarbonate, lithium hydroxide, calcium hydroxide, magnesium hydroxide, zinc hydroxide. Organic bases from which salts of disclosed acids may be prepared are amines and amino acids, the basicity of which is sufficient enough to produce a stable salt, and which are suitable for use in medical purposes (in particular, they must have low toxicity). Such amines include ammonia, methylamine, dimethylamine, trimethylamine, ethylamine, diethylamine, triethylamine, benzylamine, dibenzylamine, dicyclohexylamine, piperazine, ethylpiperidine, tris(hydroxymethyl)aminomethane and the like. Besides, salts can be prepared using

tetraalkylammonium hydroxides, such as holine, tetramethylammonium, tetraethylammonium, and the like. Basic aminoacids, i.e. lysine, ornithine and arginine, may be used as aminoacids.

“Medicinal product (drug product, medicament)” is a compound (or a mixture of compounds as a pharmaceutical composition) in the form of tablets, granules, capsules, injections, ointments and other ready forms intended for restoration, improvement or modification of physiological functions in humans and animals, as well as for treatment and prophylaxis of diseases, for diagnostics, anesthesia, contraception, cosmetology and others.

"Treat," "treatment," and "therapy" refer to a method of mitigating or eliminating a biological disorder and/or at least one of its accompanying symptoms. The term “to alleviate” a disease, disorder or condition means reducing the severity and/or occurrence frequency of the symptoms of the disease, disorder, or condition. Further, references herein to "treatment" include references to curative, palliative and prophylactic treatment.

“Prophylaxis”, “prophylactic therapy” refers to a set of measures aimed at preventing the onset, eliminating risk factors, or early detecting a disease or disorder, its progression, worsening of the course, exacerbation, relapse, complications or other consequences.

In one aspect, the subject of prophylaxis or treatment, or patient, is a mammal, preferably a human subject. Said subject may be either male or female, of any age.

The term "disorder" or "disease" means any condition, the manifestations of which may benefit or subject's prognosis with that condition may benefit from treatment according to the present invention. The definition of the term includes chronic and acute disorders or diseases including those pathological conditions that predispose a mammal to the disorder in question. Non-limiting examples of diseases to be treated include an oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma;

an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angiitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjogren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis,

osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory disease or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, otitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis.

“Therapeutically effective amount” refers to that amount of the therapeutic agent being administered in the course of treatment which will relieve the severity or eliminate the symptoms of the disease being treated.

As used in the present description and claims that follow, unless otherwise dictated by the context, the words "have", "include," and "comprise" or variations thereof such as "has", "having," "includes", "including", "comprises," or "comprising," will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.


Description of drawings.

Figure 1 illustrates the dynamics of tumor growth of NCI-H69 xenografts in the control group and in the group receiving compound **CDK7_1187** at a dose of 6 mg/kg QD for 21 days. Data are shown as arithmetic means of tumor volumes $\pm$ standard error of the mean.

Note: *- $p < 0.05$; statistically significant difference from the control group (Mann-Whitney test).

Detailed description of the invention

In one embodiment, the present invention relates to a compound of Formula **I**:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

R₁ is -C(Hal)₃, -CH(Hal)₂, -CH₂Hal, -Hal, -H, -NO₂, -(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -O-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -S-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal;

R₂, R₃ are each independently H, -(C₁-C₆)alkyl;

or R₂, R₃ together form -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several -(C₁-C₆)alkyls;

L₁, L₂ are each independently a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃-O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃-NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

A is 6-10 membered aryl unsubstituted or substituted by one or several R₇; 6-10 membered aryl unsubstituted or substituted by one or several R_{7a}, fused with -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8a}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7b}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8b}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7c}, fused with 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{8c}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{7d}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7e}, fused with (C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8e}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7f}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8f}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7g}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7h}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8h};

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}, -P(O)(C₁-C₆ alkyl)₂, -S(O)₂NH₂, -S(O)₂C₁-C₆ alkyl, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one

or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH, -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;

R₇, R_{7a}, R_{7b}, R_{7c}, R_{7d}, R_{7e}, R_{7f}, R_{7g}, R_{7h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8a}, R_{8b}, R_{8c}, R_{8d}, R_{8e}, R_{8f}, R_{8h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

In one embodiment, the present invention relates to a compound of Formula I:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

R₁ is -C(Hal)₃, -CH(Hal)₂, -CH₂Hal, -Hal, -H, -NO₂, -(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -O-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -S-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal;

R₂, R₃ are each independently H, -(C₁-C₆)alkyl;

or R₂, R₃ together form -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several -(C₁-C₆)alkyls;

L₁, L₂ are each independently a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃-O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃-NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

A is 6-10 membered aryl unsubstituted or substituted by one or several R₇; 6-10 membered aryl unsubstituted or substituted by one or several R_{7a}, fused with -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8a}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7b}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N,

S or O, unsubstituted or substituted by one or several R_{8b}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7c}, fused with 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{8c}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{7d}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7e}, fused with (C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8e}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7f}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8f}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7g}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7h}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8h};

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}, -P(O)(C₁-C₆ alkyl)₂, -S(O)₂NH₂, -S(O)₂C₁-C₆ alkyl, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;

R₇, R_{7a}, R_{7b}, R_{7c}, R_{7d}, R_{7e}, R_{7f}, R_{7g}, R_{7h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8a}, R_{8b}, R_{8c}, R_{8d}, R_{8e}, R_{8f}, R_{8h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

In one embodiment, the present invention relates to a compound of Formula **I.b**:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

R₁ is -C(Hal)₃, -CH(Hal)₂, -CH₂Hal, -Hal, -H, -NO₂, -(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -O-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -S-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal;

R₂, R₃ are each independently H, -(C₁-C₆)alkyl;

or R₂, R₃ together form -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several -(C₁-C₆)alkyls;

L₁, L₂ are each independently a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃-O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃-NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

A is 6-10 membered aryl unsubstituted or substituted by one or several R₇; 6-10 membered aryl unsubstituted or substituted by one or several R_{7a}, fused with -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8a}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7b}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8b}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7c}, fused with 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{8c}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{7d}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7e}, fused with (C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8e}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7f}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8f}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7g}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7h}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8h};

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}, -P(O)(C₁-C₆ alkyl)₂, -S(O)₂NH₂, -S(O)₂C₁-C₆ alkyl, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered

heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j} , fused with phenyl unsubstituted or substituted by one or several R_{4k} ; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l} , fused with phenyl unsubstituted or substituted by one or several R_{4m} ;

R_{4a} , R_{4b} , R_{4c} , R_{4d} , R_{4e} , R_{4f} , R_{4g} , R_{4h} , R_{4i} , R_{4j} , R_{4k} , R_{4l} , R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R_5 , R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a} , R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,

or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;


R_7 , R_{7a} , R_{7b} , R_{7c} , R_{7d} , R_{7e} , R_{7f} , R_{7g} , R_{7h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;


R_{8a} , R_{8b} , R_{8c} , R_{8d} , R_{8e} , R_{8f} , R_{8g} , R_{8h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;


Hal is an atom of F, Cl, Br, I.

In one embodiment, the present invention relates to the S-isomer of the compound of formula **I**.

In one embodiment, the present invention relates to a compound, wherein -A-L₂-R₄ is:







wherein k,m are each independently 0, 1, 2 or 3,

X₁, X₂, X₃, X₄, X₅, X₆ are each independently an atom of C, an atom of N, CH, CH₂, NH, S or O;

Y₁, Y₂, Y₃, Y₄, Y₅ are each independently an atom of C, an atom of N or CH;

L₂ is a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}, -P(O)(C₁-C₆ alkyl)₂, -S(O)₂NH₂, -S(O)₂C₁-C₆ alkyl, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k} 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;


R₇, R_{7b}, R_{7d}, R_{7f}, are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;


R_{8b} , R_{8d} , R_{8f} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.


In one embodiment, the present invention relates to a compound, wherein -A-L₂-R₄ is:

















where p is each independently 0 or 1 and q is each independently 0, 1 or 2, wherein the total number of substituents in each cycle is no more than 3;

X₁, X₂, X₃, X₄, X₅, X₆ are each independently an atom of C, an atom of N, CH, CH₂, NH, S or O; Y₁, Y₂, Y₃, Y₄, Y₅ are each independently an atom of C, an atom of N or CH;

L₂ is a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃-O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃-NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

R₄ is -H, -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}-, -P(O)(C₁-C₆ alkyl)₂-, -S(O)₂NH₂-, -S(O)₂C₁-C₆ alkyl-, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1, 2, 3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1, 2, 3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k}; 5-6 membered heteroaryl with 1, 2, 3 or 4 heteroatoms selected from N, O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH, -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;


R₇, R_{7b}, R_{7d}, R_{7f}, are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8b}, R_{8d}, R_{8f} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

In one embodiment, the present invention relates to a compound, wherein -A-L₂-R₄ is





wherein k,m are each independently 0, 1,2 or 3,

L₂ is a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, - (CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃-O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃-NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}-, -P(O)(C₁-C₆ alkyl)₂-, -S(O)₂NH₂-, -S(O)₂C₁-C₆ alkyl, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k} 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or

substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;


R₇, R_{7b}, R_{7d}, R_{7f}, are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8b}, R_{8d}, R_{8f} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.


In one embodiment, the present invention relates to a compound, wherein -A-L₂-R₄ is

















membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl,

R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,

or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;

R₇, R_{7b}, R_{7d}, R_{7f}, are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8b}, R_{8d}, R_{8f} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

In one embodiment, the present invention relates to a compound, wherein L₁, L₂ are each independently a chemical bond, -C(O)-, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -C(CH₂)₂-, -CF₂-, -S-, -S(O)-, -S(O)₂-, -O-, -NH-, -NCH₃-, -P(O)(CH₃)-, -C(O)O-, -C(O)-NH- or -NH-C(O)-.

In one embodiment, the present invention relates to a compound, wherein R₁ is -H, -CF₃, -CCl₃, -CHF₂, -CH₂F, -CHCl₂, -CH₂Cl, -NO₂.

In one embodiment, the present invention relates to a compound, wherein R₂, R₃ are each independently -H, methyl, ethyl, propyl, or R₂ and R₃ together with the C atom they are attached to form cyclopropyl, cyclobutyl or cyclopentyl.

In one embodiment, the present invention relates to a compound, wherein R₄ is -H, (=O), -OH, -F, -Cl, -Br, -P(O)(C₁-C₆ alkyl)₂; -S(O)₂NH₂; -S(O)₂-(C₁-C₆)alkyl; -(C₁-C₆)alkyl unsubstituted or substituted by one or several (=O), -Hal, -CN, -S(O)₂-(C₁-C₆)alkyl; -O-(C₁-C₆)alkyl unsubstituted or substituted by one or several (=O), -CN, -Hal; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several (=O), -OH, -Hal, -CN; phenyl unsubstituted or substituted by one or several -Hal, -CN, morpholine, -P(O)(CH₃)₂; 5-6 membered heterocyclyl with 1,2 or 3 heteroatoms selected from N or O unsubstituted or substituted by -(C₁-C₆)alkyl, (=O), -Hal, -CN; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by (=O), -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -Hal, -CN; 5-6 membered heteroaryl with 1 or 2 atoms N, unsubstituted or substituted by one or several (=O), -(C₁-C₆)alkyl, fused with phenyl; -NR_{4a}R_{4b},

R_{4a}, R_{4b} are each independently -H, (=O), morpholinyl, -F, -S(O)₂CH₃, -CN, -one, cyclopropyl, thiazolyl, -C₁-C₃ alkyl unsubstituted or substituted with phenyl, -P(O)(CH₃)₂.

In one embodiment, the present invention relates to a compound, wherein R₄ is -H, -OH, (=O), -P(O)(CH₃)₂, -CH₃, -CH₂CH₃, tert-butyl, -CH₂CF₃, -CH₂CH₂S(O)₂CH₃, -CF₃, -Cl, -OCF₃, -S(O)₂NH₂, -S(O)₂CH₃, phenyl unsubstituted or substituted by one or several -Hal, -CN; methylpiperazinyl, piperazinyl, imidazolyl, thiazolyl, methylpyrazolyl, pyrazolyl, pyridinyl, pyrimidinyl, morpholine, oxanyl, oxolanyl, cyclohexyl, cyclopentyl, cyclopropyl, oxazolyl, methyloxazolyl, dimethyloxazolyl, dihydropyridinyl, methyl dihydropyridinyl, -NR_{4a}R_{4b}, R_{4a}, R_{4b} are each independently -H, (=O), morpholinyl, -F, -S(O)₂CH₃, -CN, -one, cyclopropyl, thiazolyl, -C₁-C₃ alkyl unsubstituted or substituted with phenyl, -P(O)(CH₃)₂.

In one embodiment, the present invention relates to a compound, wherein R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O) -CH₃, morpholinyl, -F, -S(O)₂CH₃, -CN, -OH, cyclopropyl, thiazolyl, -C₁-C₃ alkyl unsubstituted or substituted with phenyl, -P(O)(CH₃)₂.

In one embodiment, the present invention relates to a compound, wherein R₅ is -H, -CH₃.

In one embodiment, the present invention relates to a compound, wherein R_{6a} and R_{6b} are each independently -H, -F, -CH₃, or R_{6a} and R_{6b} together with the C atom they are attached to are cyclopropyl.

In one embodiment, the present invention relates to a compound, wherein R₇, R_{7a}, R_{7b}, R_{7c}, R_{7d}, R_{7e}, R_{7f}, R_{7g}, R_{7h} are each independently (=O), -Hal, -CH₃; -OCH₃; -CHAl₃; -OCHAl₃.

In one embodiment, the present invention relates to a compound, wherein R_{8a}, R_{8b}, R_{8c}, R_{8d}, R_{8e}, R_{8f}, R_{8g}, R_{8h} are each independently (=O), -Hal, -CH₃; -OCH₃; -CHAl₃; -OCHAl₃.

Compounds, described in the present invention, may be obtained as, and/or used as, pharmaceutically acceptable salts. The type of pharmaceutical acceptable salts, include, but are not limited to: acid salts formed by reacting the free base form of the compound with a pharmaceutically acceptable inorganic acid such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, metaphosphoric acid, and the like; or with an organic acid such as formic acid, acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, trifluoroacetic acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanedisulfonic acid, benzenesulfonic acid, toluenesulfonic acid, 2-naphthalenesulfonic acid, 4-methylbicyclo-[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-3-hydroxy-2-ene-1-carboxylic acid, 3-phenylpropionic acid, trimethylacetic acid, tert-butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like.

The corresponding counterions of the pharmaceutically acceptable salts may be analyzed and identified using various methods including, but not limited to, ion exchange chromatography, ion chromatography, capillary electrophoresis, inductively coupled plasma, atomic absorption spectroscopy, mass spectrometry, or any combination thereof.



The salts are recovered by using at least one of the following techniques: filtration, precipitation with a non-solvent followed by filtration, evaporation of the solvent, or, in the case of aqueous solutions, lyophilization. It should be understood that a reference to a pharmaceutically acceptable salt includes the solvent addition forms or crystal forms thereof, particularly solvates or polymorphs. Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and may be formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of compounds described in the present patent can









be conveniently prepared or formed during the processes described in the present invention. In addition, the compounds provided in the present invention can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided in the present invention.









Compounds described in the present invention may be provided in various forms, including but not limited to, amorphous forms, milled forms and nano-particulate forms. In addition, compounds described in the present invention include crystalline forms, also known as polymorphs. Polymorphs include different crystal packing arrangements of the same elemental composition of a compound. Polymorphs typically have different X-ray diffraction patterns, infrared spectra, melting points, different density, hardness, crystal shape, optical and electrical properties, stability, and solubility. Various factors such as the recrystallization solvent, rate of crystallization, and storage temperature may cause one crystal form to dominate.











The screening and characterization of the pharmaceutically acceptable salts, polymorphs and/or solvates may be accomplished using a variety of techniques including, but not limited to, thermal analysis, x-ray diffraction, spectroscopy, vapor sorption, and microscopy. Thermal analysis methods address to analysis of thermo chemical degradation or thermo physical processes including, but not limited to, polymorphic transitions, and such methods are used to analyze the relationships between polymorphic forms, to determine weight loss, to find the glass transition temperature, or for excipient compatibility studies. Such methods include, but are not limited to, differential scanning calorimetry (DSC), modulated differential scanning calorimetry (MDCS), thermogravimetric analysis (TGA), thermogravi-metric and infrared analysis (TG/IR). Crystallographic methods include, but are not limited to, single crystal and powder diffractometers and synchrotron sources. The various spectroscopic techniques used include, but are not limited to, Raman (combinational scattering), FTIR, UVIS, and NMR (liquid and solid state). The various microscopy techniques include, but are not limited to, polarized light microscopy, scanning electron microscopy (SEM) with energy dispersive x-ray analysis (EDX), environmental scanning electron microscopy with EDX (in gas or water vapor atmosphere), IR microscopy, and Raman microscopy.









In another embodiment, the present invention relates to compounds selected from the group including:









Code Name	Structure	Code Name	Structure
CDK7_461 4-(2-aminobenzoyl)-N-[(3S)-6,6-dimethylpiperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1000 4-[5-(2-methyl-2H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	









CDK7_461_a 4-(2-aminobenzoyl)-N-(6,6-dimethylpiperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1000_a 4-[5-(2-methyl-2H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_549 4-(2-aminobenzoyl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1060 4-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_549_a 4-(2-aminobenzoyl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1060_a 4-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_813 4-benzoyl-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1102 4-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	









CDK7_813_a 4-benzoyl-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1102_a 4-[5-(1-cyclopropyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_540 4-[(2-aminophenyl)methyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1103 4-[5-(1-cyclopropyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_540_a 4-[(2-aminophenyl)methyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1103_a 4-[5-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_541 4-[1-(2-aminophenyl)ethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1124 4-[5-(1-methyl-1H-1,3-benzodiazol-2-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	









CDK7_541_a 4-[1-(2-aminophenyl)ethyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1124a 4-[5-(1-methyl-1H-1,3-benzodiazol-2-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_541_b 4-[(1S)-1-(2-aminophenyl)ethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1020 ethyl 4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrole-2-carboxylate	
CDK7_541_c 4-[(1R)-1-(2-aminophenyl)ethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1020_a ethyl 4-{2-[[[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrole-2-carboxylate	
CDK7_542 4-[2-(2-aminophenyl)propan-2-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1017 N,N,1-trimethyl-4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrole-2-carboxamide	
CDK7_542_a 4-[2-(2-aminophenyl)propan-2-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1017_a N,N,1-trimethyl-4-(2-[[[piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrole-2-carboxamide	









CDK7_543 4-[1-(2-aminophenyl)cyclopropyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1061 N-[(3S)-piperidin-3-yl]-4-[5-(pyridin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_543_a 4-[1-(2-aminophenyl)cyclopropyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1061_a N-(piperidin-3-yl)-4-[5-(pyridin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_544 4-[(2-aminophenyl)difluoromethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1062 N-[(3S)-piperidin-3-yl]-4-[5-(pyrimidin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_544_a 4-[(2-aminophenyl)difluoromethyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1062_a N-(piperidin-3-yl)-4-[5-(pyrimidin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	









CDK7_545 4-[(2-aminophenyl)sulfanyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1057 4-{5-[2-(dimethylphosphoryl)phenyl]-1H-pyrrol-3-yl}-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_545_a 4-[(2-aminophenyl)sulfanyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1057_a 4-{5-[2-(dimethylphosphoryl)phenyl]-1H-pyrrol-3-yl}-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_546 4-(2-aminobenzenesulfinyl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_955 7-(1-methyl-1H-pyrazol-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_546_a 4-(2-aminobenzenesulfinyl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_955_a 7-(1-methyl-1H-pyrazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_546_b 4-[(R)-2-aminobenzenesulfinyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1122 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_546_c 4-[(S)-2-aminobenzenesulfinyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1122_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_547 4-(2-aminobenzenesulfonyl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1123 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1-methyl-1H-pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_547_a 4-(2-aminobenzenesulfonyl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1123_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(1-methyl-1H-pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_548 4-(2-aminophenoxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1136 3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_548_a 4-(2-aminophenoxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1136_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_550 N4-(2-aminophenyl)-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1137 3-(2-{[(7S)-5-azaspiro[3.5]nonan-7-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_550_a N4-(2-aminophenyl)-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1137_a 3-[2-({5-azaspiro[3.5]nonan-7-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_551 N4-(2-aminophenyl)-N4-methyl-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1138 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1,3-thiazol-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_551_a N4-(2-aminophenyl)-N4-methyl-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1138_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(1,3-thiazol-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_812 N4-phenyl-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1139 7-(3,5-dimethyl-1,2-oxazol-4-yl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_812_a N4-phenyl-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1139_a 7-(3,5-dimethyl-1,2-oxazol-4-yl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_754 4-phenoxy-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1140 5-[3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl]-1-methyl-1,2-dihydropyridin-2-one	
CDK7_754_a 4-phenoxy-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1140_a 5-(3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl)-1-methyl-1,2-dihydropyridin-2-one	
CDK7_766 4-benzyl-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1141 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(pyridin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_766_a 4-benzyl-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1141_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_552 4-[(2-aminophenyl)(methyl)phosphoro]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1142 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(pyridin-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_552_a 4-[(2-aminophenyl)(methyl)phosphoro]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1142_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_552_b 4-[(R)-(2-aminophenyl)(methyl)phosphoro]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1143 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(pyridin-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_552_c 4-[(S)-(2-aminophenyl)(methyl)phosphoro]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1143_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_735 N4-[2-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1144 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(1-methyl-1H-pyrazol-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_735_a N4-[2-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1144_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(1-methyl-1H-pyrazol-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_897 4-[2-(dimethylphosphoryl)phenoxy]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1145 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(1,3-thiazol-2-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_897_a 4-[2-(dimethylphosphoryl)phenoxy]-N-[piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1145_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(1,3-thiazol-2-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_892 4-[3-(dimethylphosphoryl)phenoxy]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1146 7-[(3,5-dimethyl-1,2-oxazol-4-yl)methyl]-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_892_a 4-[3-(dimethylphosphoryl)phenoxy]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1146_a 7-[(3,5-dimethyl-1,2-oxazol-4-yl)methyl]-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_888 N4-[3-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1147 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[1-(1-methyl-1H-pyrazol-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_888_a N4-[3-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1147_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(1-methyl-1H-pyrazol-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_903 N4-[2-(dimethylphosphoryl)-5-methylphenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1148 3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-7-[1-(1,3-thiazol-2-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_903_a N4-[2-(dimethylphosphoryl)-5-methylphenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1148_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(1,3-thiazol-2-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_904 N4-[2-(dimethylphosphoryl)-4-methylphenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1149 7-[1-(3,5-dimethyl-1,2-oxazol-4-yl)ethyl]-3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_904_a N4-[2-(dimethylphosphoryl)-4-methylphenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1149_a 7-[1-(3,5-dimethyl-1,2-oxazol-4-yl)ethyl]-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_905 N4-[2-(dimethylphosphoryl)-5-(trifluoromethyl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1150 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(oxan-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_905_a N4-[2-(dimethylphosphoryl)-5-(trifluoromethyl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1150_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxan-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_906 N4-[2-(dimethylphosphoryl)-4-(trifluoromethyl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1151 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(oxan-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_906_a N4-[2-(dimethylphosphoryl)-4-(trifluoromethyl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1151_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxan-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_907 N4-[5-chloro-2-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1152 3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-(oxolan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_907_a N4-[5-chloro-2-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1152_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxolan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_908 N4-[4-chloro-2-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1153 3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-[(oxolan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_908_a N4-[4-chloro-2-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1153_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxolan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_915 N4-[2-(dimethylphosphoryl)-4-(trifluoromethoxy)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1154 7-cyclopentyl-3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_915_a N4-[2-(dimethylphosphoryl)-4-(trifluoromethoxy)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1154_a 7-cyclopentyl-3-(2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_889 2-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]amino]benzene-1-sulfonamide		CDK7_1155 7-(cyclopentylmethyl)-3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_889_a 2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}amino)benzene-1-sulfonamide		CDK7_1155_a 7-(cyclopentylmethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_890 N4-(3- N4-(3-methanesulfonylphenyl)-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1156 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[1-(oxan-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_890_a N4-(3-methanesulfonylphenyl)-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine		CDK7_1156_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(oxan-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_891 3-[(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)amino]benzene-1-sulfonamide		CDK7_1157 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[1-(oxolan-3-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_891_a 3-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}amino)benzene-1-sulfonamide		CDK7_1157_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(oxolan-3-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_893 2-[(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)oxy]benzenesulfonamide		CDK7_1158 7-(1-cyclopentylethyl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_893_a 2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}oxy)benzenesulfonamide		CDK7_1158_a 7-(1-cyclopentylethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_894 4-(3-methanesulfonylphenoxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1159 7-tert-butyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_894_a 4-(3-methanesulfonylphenoxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1159_a 7-tert-butyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_895 3-[(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)oxy]benzenesulfonamide		CDK7_1160 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2,2-dimethylpropyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_895_a 3-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}oxy)benzenesulfonamide		CDK7_1160_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2,2-dimethylpropyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_827 4-(2,3-dihydro-1H-indol-7-yloxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1161 7-(3,3-dimethylbutan-2-yl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_827_a 4-(2,3-dihydro-1H-indol-7-yloxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1161_a 7-(3,3-dimethylbutan-2-yl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_828 4-(1H-indol-7-yloxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1162 8-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-4-(1-methyl-1H-pyrazol-4-yl)-2H,3H,4H,5H,6H-pyrrolo[2,3-f][1,4]oxazepin-5-one	
CDK7_828_a 4-(1H-indol-7-yloxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1162_a 8-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-4-(1-methyl-1H-pyrazol-4-yl)-2H,3H,4H,5H,6H-pyrrolo[2,3-f][1,4]oxazepin-5-one	
CDK7_829 N-[(3S)-piperidin-3-yl]-4-(1,2,3,4-tetrahydroquinolin-8-yloxy)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1163 (6S)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_829_a 4 N-(piperidin-3-yl)-4-(1,2,3,4-tetrahydroquinolin-8-yloxy)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1163_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_830 N-[(3S)-piperidin-3-yl]-4-(quinolin-8-yloxy)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1164 (6R)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_830_a N-(piperidin-3-yl)-4-(quinolin-8-yloxy)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1164_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_898 N-{2-[(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)amino]phenyl}acetamide		CDK7_1165 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,6-dimethyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_898_a N-[2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}amino)phenyl]acetamide		CDK7_1165_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,6-dimethyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	











CDK7_899 N-{2-[(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)oxy]phenyl}acetamide		CDK7_1105 6-cyclopropyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_899_a N-[2-(2-{[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}oxy)phenyl]acetamide		CDK7_1105_a 6-cyclopropyl-3-(2-{[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_460 N-[(3S)-6,6-dimethylpiperidin-3-yl]-4-(5-phenyl-1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1058 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1,3-thiazol-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_460_a N-(6,6-dimethylpiperidin-3-yl)-4-(5-phenyl-1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1058_a 3-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1,3-thiazol-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_533 4-(5-phenyl-1H-pyrrol-3-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1052 1-methyl-5-[7-oxo-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-6-yl]-1,2-dihydropyridin-2-one	
CDK7_533_a 4-(5-phenyl-1H-pyrrol-3-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1052_a 1-methyl-5-[7-oxo-3-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-6-yl]-1,2-dihydropyridin-2-one	
CDK7_534 4-(4-phenyl-1H-pyrrol-3-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_972 N-[(3S)-piperidin-3-yl]-4-(1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_534_a 4-(4-phenyl-1H-pyrrol-3-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_972_a N-[piperidin-3-yl]-4-(1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	









CDK7_535 4-(5-benzoyl-1H-pyrrol-3-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1021 7-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepine-4,8-dione	
CDK7_535_a 4-(5-benzoyl-1H-pyrrol-3-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1021_a 7-methyl-3-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepine-4,8-dione	
CDK7_536 N,N-dimethyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1121 6-(oxan-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_536_a N,N-dimethyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1121_a 6-(oxan-4-yl)-3-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_537 4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_809 4-(1H-indol-1-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_537_a 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_809_a 4-(1H-indol-1-yl)-N[piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_538 N-phenyl-4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_810 4-(1H-1,3-benzodiazol-1-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_538_a N-phenyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_810_a 4-(1H-1,3-benzodiazol-1-yl)-N[piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	











CDK7_539 4-[5-(4-methylpiperazine-1-carbonyl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_811 4-(1H-1,2,3-benzotriazol-1-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_539_a 4-[5-(4-methylpiperazine-1-carbonyl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_811_a 4-(1H-1,2,3-benzotriazol-1-yl)-N-[piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_922 4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrole-2-carboxylic acid		CDK7_953 7-methyl-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_922_a 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxylic acid		CDK7_953_a 7-methyl-3-(2-[[[piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

CDK7_695 4-[4-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-yl]benzonitrile		CDK7_959 6-(1-methyl-1H-pyrazol-4-yl)-3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_695_a 4-(4-{2-[[piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-yl)benzonitrile		CDK7_959_a 6-(1-methyl-1H-pyrazol-4-yl)-3-(2-{{piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_696 3-[4-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-yl]benzonitrile		CDK7_961 6-benzyl-3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_696_a 3-(4-{2-[[piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-yl)benzonitrile		CDK7_961_a 6-benzyl-3-{2-[[piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_814 3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_966 N,N-dimethyl-4-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-imidazole-2-carboxamide	











CDK7_814_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_966_a N,N-dimethyl-4-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-imidazole-2-carboxamide	
CDK7_815 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[3,2-c]pyridin-4-one		CDK7_991 N,N-dimethyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-3-carboxamide	
CDK7_815_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[3,2-c]pyridin-4-one		CDK7_991_a N,N-dimethyl-4-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-3-carboxamide	
CDK7_843 6-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1176 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(pyridin-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	











CDK7_843_a 6-methyl-3-(2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1176_a 3-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(pyridin-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_844 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-4-one		CDK7_1187 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_844_a 3-(2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-4-one		CDK7_1187_a 3-(2-{[6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_845 6-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1199 3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1-methyl-1H-pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_845_a 6-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1199_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-6-(1-methyl-1H-pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_833 N-(1H-imidazol-2-yl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1200 3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_833_a N-(1H-imidazol-2-yl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1200_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-6-methyl-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_834 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-N-(1,3-thiazol-2-yl)-1H-pyrrole-2-carboxamide		CDK7_1201 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_834_a 4-{2- [(piperidin-3- yl)amino]-5- (trifluoromethy l)pyrimidin-4- yl}-N-(1,3- thiazol-2-yl)- 1H-pyrrole-2- carboxamide		CDK7_1201_a 3-(2-{[(3S)- piperidin-3- yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)-6- (1,3-thiazol-4-yl)- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	
CDK7_835 N-(1-methyl- 1H-pyrazol-4- yl)-4-(2- {[(3S)- piperidin-3- yl]amino}-5- (trifluoromethy l)pyrimidin-4- yl)-1H- pyrrole-2- carboxamide		CDK7_1202 3-(2-{[(3S)- piperidin-3- yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)-6- (1,3-thiazol-4-yl)- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	
CDK7_835_a N-(1-methyl- 1H-pyrazol-4- yl)-4-{2- [(piperidin-3- yl)amino]-5- (trifluoromethy l)pyrimidin-4- yl}-1H- pyrrole-2- carboxamide		CDK7_1202_a 3-{2-[(piperidin- 3-yl)amino]-5- (trifluoromethyl)p yrimidin-4-yl}-6- (1,3-thiazol-4-yl)- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	
CDK7_836 4-(2-{[(3S)- piperidin-3- yl]amino}-5- (trifluoromethy l)pyrimidin-4- yl)-N-(pyridin- 4-yl)-1H- pyrrole-2- carboxamide		CDK7_1203 3-(2-{[(3S)- piperidin-3- yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)-6- (1,3-thiazol-5-yl)- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	
CDK7_836_a 4-{2- [(piperidin-3- yl)amino]-5- (trifluoromethy l)pyrimidin-4- yl}-N- (pyridin-4-yl)-		CDK7_1203_a 3-{2-[(piperidin- 3-yl)amino]-5- (trifluoromethyl)p yrimidin-4-yl}-6- (1,3-thiazol-5-yl)- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	









1H-pyrrole-2-carboxamide			
CDK7_837 N-(4-cyanophenyl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1204 6-(2-methyl-1,3-thiazol-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_837_a N-(4-cyanophenyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1204_a 6-(2-methyl-1,3-thiazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_838 N-[4-(morpholin-4-yl)phenyl]-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1205 6-(2-methyl-1,3-thiazol-5-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_838_a N-[4-(morpholin-4-yl)phenyl]-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1205_a 6-(2-methyl-1,3-thiazol-5-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_846 N-methyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1206 6-(4-methyl-1,3-thiazol-2-yl)-3-{2-[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_846_a N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1206_a 6-(4-methyl-1,3-thiazol-2-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_847 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-N-(2,2,2-trifluoroethyl)-1H-pyrrole-2-carboxamide		CDK7_1207 6-(5-methyl-1,3-thiazol-2-yl)-3-{2-[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_847_a 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-N-(2,2,2-trifluoroethyl)-1H-pyrrole-2-carboxamide		CDK7_1207_a 6-(5-methyl-1,3-thiazol-2-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_848 N-(2-methanesulfonyl-ethyl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1208 6-(cyclopropylmethyl)-3-{2-[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_848_a N-(2-methanesulfonyl-ethyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1208_a 6-(cyclopropylmethyl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_849 4-[5-(morpholine-4-carbonyl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1209 6-cyclobutyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_849_a 4-[5-(morpholine-4-carbonyl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1209_a 6-cyclobutyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_850 N-(oxan-4-yl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1210 6-cyclopentyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_850_a N-(oxan-4-yl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1210_a 6-cyclopentyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_851 N-(4-hydroxycyclohexyl)-4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1211 6-(1-methylcyclopropyl)-3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_851_a N-(4-hydroxycyclohexyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1211_a 6-(1-methylcyclopropyl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_852 N-(4-cyanocyclohexyl)-4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1212 6-cyclohexyl-3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_852_a N-(4-cyanocyclohexyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1212_a 6-cyclohexyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_951 N-cyclopropyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1213 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(pyridin-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_951_a N-cyclopropyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1213_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(pyridin-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_923 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1214 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(pyridin-3-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_923_a 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1214_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(pyridin-3-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_924 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1215 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(pyridin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_924_a 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one		CDK7_1215_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(pyridin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_925 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,7H,8H-pyrrolo[2,3-c]azepin-8-one		CDK7_1216 6-(3-methylpyridin-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_925_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H-pyrrolo[2,3-c]azepin-8-one		CDK7_1216_a 6-(3-methylpyridin-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_926 3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one		CDK7_1217 6-(2-methylpyridin-3-yl)-3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_926_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one		CDK7_1217_a 6-(2-methylpyridin-3-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_927 3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H,8H-pyrrolo[2,3-c]azepin-8-one		CDK7_1218 6-(4-methylpyridin-3-yl)-3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_927_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,8H-pyrrolo[2,3-c]azepin-8-one		CDK7_1218_a 6-(4-methylpyridin-3-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	









CDK7_928 7-methyl-3-(2- {[(3S)- piperidin-3- yl]amino}-5- (trifluorometh- yl)pyrimidin-4- yl)- 1H,4H,7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1219 6-(3- methylpyridin-2- yl)-3-(2-{[(3S)- piperidin-3- yl]amino}-5- (trifluoromethyl)p- yrimidin-4-yl)- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	
CDK7_928_a 7-methyl-3-{2- [(piperidin-3- yl)amino]-5- (trifluorometh- yl)pyrimidin-4- yl}- 1H,4H,7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1219_a 6-(3- methylpyridin-2- yl)-3-{2- [(piperidin-3- yl)amino]-5- (trifluoromethyl)p- yrimidin-4-yl}- 1H,6H,7H- pyrrolo[2,3- c]pyridin-7-one	
CDK7_929 7-methyl-3-{2- [[(3S)-6,6- dimethylpiperidin- 3-yl]amino]-5- (trifluorometh- yl)pyrimidin-4- yl}- 1H,4H,7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1225 7-cyclopropyl-3- (2-{[(3S)-6,6- dimethylpiperidin- 3-yl]amino}-5- (trifluoromethyl)p- yrimidin-4-yl)- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_929_a 7-methyl-3-{2- [(piperidin-3- yl)amino]-5- (trifluorometh- yl)pyrimidin-4- yl}- 1H,4H,7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1225_a 7-cyclopropyl-3- {2-[(6,6- dimethylpiperidin- 3-yl)amino]-5- (trifluoromethyl)p- yrimidin-4-yl}- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	









CDK7_930 7-methyl-3-(2- {[(3S)- piperidin-3- yl]amino}-5- (trifluorometh- yl)pyrimidin-4- yl)-7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1226 7-cyclobutyl-3- (2-{[(3S)-6,6- dimethylpiperidin -3-yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_930_a 7-methyl-3-{2- [(piperidin-3- yl)amino]-5- (trifluorometh- yl)pyrimidin-4- yl}-7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1226_a 7-cyclobutyl-3- (2-{[(6,6- dimethylpiperidin -3-yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_931 3-(2-{[(3S)- piperidin-3- yl]amino}-5- (trifluorometh- yl)pyrimidin-4- yl)-7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1227 7- (cyclopropylmeth yl)-3-(2-{[(3S)- 6,6- dimethylpiperidin -3-yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_931_a 3-{2- [(piperidin-3- yl)amino]-5- (trifluorometh- yl)pyrimidin-4- yl}-7H,8H- pyrrolo[2,3- c]azepin-8-one		CDK7_1227_a 7- (cyclopropylmeth yl)-3-{2-[(6,6- dimethylpiperidin -3-yl)amino]-5- (trifluoromethyl)p yrimidin-4-yl}- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	









CDK7_932 3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1228 7-(cyclobutylmethyl)-3-(2-{{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_932_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1228_a 7-(cyclobutylmethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_933 8-methyl-3-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1229 3-(2-{{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_933_a 8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1229_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl)-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	









CDK7_934 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1230 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2,2,2-trifluoroethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_934_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1230_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2,2,2-trifluoroethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_935 8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1236 4-{5-methyl-3,4,6,13-tetraazatricyclo[8.3.0.0^{2,6}]trideca-1(10),2,4,11-tetraen-11-yl}-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_935_a 8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1236_a 4-{5-methyl-3,4,6,13-tetraazatricyclo[8.3.0.0^{2,6}]trideca-1(10),2,4,11-tetraen-11-yl}-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	











CDK7_936 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1237 4-{5-cyclopropyl-3,4,6,13-tetraazatricyclo[8.3.0.0^{2,6}]trideca-1(10),2,4,11-tetraen-11-yl}-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_936_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1237_a 4-{5-cyclopropyl-3,4,6,13-tetraazatricyclo[8.3.0.0^{2,6}]trideca-1(10),2,4,11-tetraen-11-yl}-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine	
CDK7_937 8-methyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1238 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,6,7-trimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_937_a 8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1238_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,6,7-trimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	











CDK7_938 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1239 (6S)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_938_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1239_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_939 8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1240 (6R)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_939_a 8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1240_a (6R)-3-(2-{[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

CDK7_940 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1241 7-cyclopropyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,6-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_940_a 3-(2-{[piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1241_a 7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,6-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_941 8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1242 (6S)-7-cyclopropyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_941_a 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1242_a 7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	











CDK7_942 8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1243 (6R)-7-cyclopropyl-3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_942_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-9H-pyrrolo[2,3-c]azocin-9-one		CDK7_1243_a 7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_943 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H-pyrrolo[2,3-c]pyrrol-6-one		CDK7_1244 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-5,5,6-trimethyl-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	
CDK7_943_a 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H-pyrrolo[2,3-c]pyrrol-6-one		CDK7_1244_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-5,5,6-trimethyl-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one	

CDK7_944 5-methyl-3-(2- {[(3S)- piperidin-3- yl]amino}-5- (trifluoromethy l)pyrimidin-4- yl)- 1H,4H,5H,6H- pyrrolo[2,3- c]pyrrol-6-one		CDK7_1254 6,7-dimethyl-3- (2-{[(3S)- piperidin-3- yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_944_a 5-methyl-3-{2- [(piperidin-3- yl)amino]-5- (trifluoromethy l)pyrimidin-4- yl}- 1H,4H,5H,6H- pyrrolo[2,3- c]pyrrol-6-one		CDK7_1254_a 6,7-dimethyl-3- {2-[(piperidin-3- yl)amino]-5- (trifluoromethyl)p yrimidin-4-yl}- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_945 N-[(3S)- piperidin-3- yl]-4-(1H- pyrazol-4-yl)- 5- (trifluoromethy l)pyrimidin-2- amine		CDK7_1259 3-(2-{[(3S)-6,6- dimethylpiperidin -3-yl]amino}-5- (trifluoromethyl)p yrimidin-4-yl)-7- ethyl- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one	
CDK7_945_a N-(piperidin-3- yl)-4-(1H- pyrazol-4-yl)- 5- (trifluoromethy l)pyrimidin-2- amine		CDK7_1259_a 3-{2-[(6,6- dimethylpiperidin -3-yl)amino]-5- (trifluoromethyl)p yrimidin-4-yl}-7- ethyl- 1H,4H,5H,6H,7H, 8H-pyrrolo[2,3- c]azepin-8-one7	







CDK7_946 4-(1H-imidazol-4-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1261 6-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,4-diazatricyclo[7.1.1.0 ^{3,7}]undeca-3(7),5-dien-2-one	
CDK7_946_a 4-(1H-imidazol-4-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1261_a 6-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,4-diazatricyclo[7.1.1.0 ^{3,7}]undeca-3(7),5-dien-2-one	
CDK7_947 N-[(3S)-piperidin-3-yl]-4-(1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1264 7-methyl-3-(2-{[(3S,6S)-6-methylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_947_a N-[piperidin-3-yl]-4-(1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1264_a 7-methyl-3-{2-[(6-methylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_948 N-[(3S)-piperidin-3-yl]-4-(1H-1,2,3-triazol-4-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1265 3-(2-{[(3S,6S)-6-ethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

CDK7_948_a N-(piperidin-3-yl)-4-(1H-1,2,3-triazol-4-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1265_a 3-{2-[(6-ethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_949 N-[(3S)-piperidin-3-yl]-4-(1H-1,2,4-triazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1268 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_949_a N-(piperidin-3-yl)-4-(1H-1,2,4-triazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1268_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_950 N-[(3S)-piperidin-3-yl]-4-(2H-1,2,3,4-tetrazol-5-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1269 2-[3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl]acetonitrile	
CDK7_950_a N-(piperidin-3-yl)-4-(2H-1,2,3,4-tetrazol-5-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1269_a 2-(3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl)acetonitrile	

		c]azepin-7-yl)acetonitrile	
CDK7_900 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,2,3,4-tetrahydroquinolin-2-one		CDK7_1270 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_900_a 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,2,3,4-tetrahydroquinolin-2-one		CDK7_1270_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_901 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,2-dihydroquinolin-2-one		CDK7_1271 3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_901_a 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,2-dihydroquinolin-2-one		CDK7_1271_a 3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

CDK7_921 4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1,2-dihydroisoquinolin-1-one		CDK7_1272 3-(2-{{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_921_a 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,2-dihydroisoquinolin-1-one		CDK7_1272_a 3-[2-{{4-azaspiro[2.5]octan-6-yl}amino}-5-(trifluoromethyl)pyrimidin-4-yl]-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_1001 N-cyclopropyl-N-methyl-4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1273 2-[3-(2-{{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl}acetonitrile	
CDK7_1001_a N-cyclopropyl-N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1273_a 2-{3-[2-{{4-azaspiro[2.5]octan-6-yl}amino}-5-(trifluoromethyl)pyrimidin-4-yl]-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl}acetonitrile	
CDK7_1003 N-methyl-4-(2-{{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-N-(1,3-thiazol-2-yl)-		CDK7_1274 3-(2-{{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

1H-pyrrole-2-carboxamide			
CDK7_1003_a N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-N-(1,3-thiazol-2-yl)-1H-pyrrole-2-carboxamide		CDK7_1274_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_997 N-benzyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1275 3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_997_a N-benzyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1275_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_998 N-benzyl-N-methyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrole-2-carboxamide		CDK7_1276 3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-cyclopropyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

CDK7_998_a N-benzyl-N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrole-2-carboxamide		CDK7_1276_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-cyclopropyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_999 4-[5-(1-methyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1277 3-(2-{{(6S)-4-azaspiro[2.5]octan-6-yl}amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	
CDK7_999_a 4-[5-(1-methyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine		CDK7_1277_a 3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one	

The present invention also relates to a method for inhibiting the biological activity of cyclin-dependent protein kinases CDK7 in a subject, consisting in contacting cyclin-dependent protein kinases CDK7 with a compound according to the present invention.

CDK7-inhibiting compounds may be used to manufacture medicinal products intended for treating any of the pathological conditions described herein, for example, compounds of formula I, pharmaceutically acceptable salts, solvates or stereoisomers will be useful in the prophylaxis or treatment of diseases or medical conditions associated, alone or partially, with CDK7 activity, for example, oncological, infectious inflammatory or non-infectious inflammatory, auto-inflammatory, autoimmune disease. Exemplary oncological disease which may be amenable to treatment using the above compounds is selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma,

oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma, but not limited thereto.

Exemplary autoimmune disease is selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angiitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjogren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis, but not limited thereto.

Exemplary autoinflammatory disease is selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS), but not limited thereto.

Exemplary infectious inflammatory disease or non-infectious inflammatory disease is selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, ostitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis,

omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis, but not limited thereto.

In one embodiment, the present invention relates to a pharmaceutical composition that comprises a therapeutically effective amount of the compound described herein, or pharmaceutically acceptable salt, solvate, or stereoisomer thereof, and one or more pharmaceutically acceptable excipients.

In one embodiment, the present invention relates to a pharmaceutical composition comprising a therapeutically effective amount of at least one of the compounds described herein, or pharmaceutically acceptable salt, solvate, stereoisomer thereof, and one or more pharmaceutically acceptable excipients.

In another one embodiment, the pharmaceutical composition of the present invention is intended to treat or prevent a disease or disorder associated with increased activity of cyclin-dependent protein kinases CDK7. In another embodiment of the invention, the pharmaceutical composition according to the present invention is intended for the prophylaxis and treatment of a disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinases CDK7 is an oncological, infectious inflammatory or non-infectious inflammatory, auto-inflammatory, autoimmune disease.

In another one embodiment of the present invention, the pharmaceutical composition comprising compounds of the present invention is intended to prevent or treat

an oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma;

an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angiitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjogren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-

associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, otitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvovaginitis.

In another embodiment of the present invention, the pharmaceutical composition comprising the compounds according to the present invention is intended for the prophylaxis or treatment of an oncological disease being breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer.

The pharmaceutical composition of the present invention may comprise, for example, from about 5% to about 100% active ingredients by weight, from about 10% to about 100% active ingredients, from about 10% to about 60% active ingredients. It is to be understood that each dosage unit may not comprise an effective amount of an active ingredient or ingredients, because the sufficient effective amount may be achieved by administering multiple dosage unit forms.

A typical composition is prepared by mixing the compound of the present invention with a carrier, diluent or excipient. Suitable carriers, diluents and fillers are well known to those skilled in the art and include materials such as carbohydrates, waxes, water soluble and/or swellable polymers, hydrophilic or hydrophobic materials, gelatin, oils, solvents, water, and the like. The particular carrier, diluent or filler used will depend upon the means and purpose for which compound of the present invention is being applied. Solvents are generally selected based on solvents recognized by those skilled in the art as safe to be administered to a mammal. In general,

safe solvents are aqueous solvents such as water and other solvents that are soluble or miscible in water. Suitable aqueous solvents include water, as the main ingredient, ethanol, propylene glycol, polyethylene glycols (e.g., PEG400, PEG300), etc. and mixtures thereof. The compositions may also include one or more buffers, stabilizing agents, surfactants, wetting agents, lubricating agents, emulsifiers, suspending agents, preservatives, antioxidants, opaquing agents, glidants, processing aids, colorants, sweeteners, perfuming agents, flavoring agents and other known additives to provide an elegant presentation of the drug product (i.e., compound of the present invention or pharmaceutical composition thereof) or aid in the manufacturing of the pharmaceutical product (i.e. medicinal product). The pharmaceutical compositions should preferably be manufactured in compliance with the GMP (Good Manufacturing Practice) requirements.

The pharmaceutical compositions may also include salts, solvates and hydrates of compounds of the present invention, or stabilized form of the compound (e.g., complex with a cyclodextrin derivative or other known complexation agent).

The pharmaceutical compositions of the present invention are typically suitable for oral administration. Oral administration may involve swallowing, so that the compound enters the gastrointestinal tract, and/or buccal, lingual, or sublingual administration by which the compound enters the blood stream directly from the mouth.

Formulations suitable for oral administration include solid, semi-solid and liquid systems such as tablets; granules; soft or hard capsules containing multi- or nano-particulates, liquids, or powders; lozenges (including liquid-filled); chews; gels; fast dispersing dosage forms; films; ovules; sprays; and buccal/mucoadhesive patches. More preferred formulations for oral administration are tablets, granules, and capsules.

Liquid formulations include suspensions, solutions, syrups and elixirs. Such formulations may be employed as fillers in soft or hard capsules (made, for example, made of gelatin or hydroxypropylmethylcellulose) and typically comprise a carrier, for example, water, ethanol, polyethylene glycol, propylene glycol, methylcellulose, or a suitable oil, and one or more emulsifying agents and/or suspending agents. Liquid formulations may also be prepared by the reconstitution of a solid, for example, from a sachet.

The pharmaceutical compositions of the invention may also be administered parenterally. As used herein, "parenteral administration" of a pharmaceutical composition includes any route of administration characterized by physical breaching of a tissue of a subject and administration of the pharmaceutical composition through the breach in the tissue, thus generally resulting in the direct administration into the blood stream, into muscle, or into an internal organ. Parenteral administration thus includes, inter alia, administration of a pharmaceutical composition by injection of the composition, by application of the composition through a surgical incision, by application of the composition through a tissue-penetrating non-surgical wound, and the like. In particular, parenteral administration is contemplated to include, inter alia, subcutaneous, intraperitoneal, intramuscular, intravenous, intraarterial, intrathecal, intraventricular, intraurethral, intracranial, intrasynovial injection or infusions; and kidney dialytic infusion techniques. Intratumoral delivery, e.g. intratumoral injection, may also be advantageous. Regional perfusion is also provided.

Formulations of pharmaceutical compositions suitable for parenteral administration typically comprise the active ingredient combined with a pharmaceutically acceptable carrier, such as sterile water or sterile isotonic saline. Such formulations may be prepared, packaged in a form suitable for bolus administration or for continuous administration. Injectable formulations may be prepared, packaged in unit dosage form, such as in ampoules or in multi-dose containers containing

a preservative. Formulations for parenteral administration include, inter alia, suspensions, solutions, emulsions in oily or aqueous vehicles, pastes, and the like.

The compounds of the invention can also be administered intranasally or by inhalation, typically in the form of a dry powder (either alone, as a mixture, or as a mixed component particle, for example, mixed with a suitable pharmaceutically acceptable excipient) from a dry powder inhaler, as an aerosol pressurised container, pump, spray, atomiser (preferably an atomiser using electrohydrodynamics to produce a fine mist), or nebuliser, with or without the use of a suitable propellant, or as nasal drops.

The pressurised container, pump, spray, atomizer, or nebuliser typically contains a solution or suspension of the compound of the present invention comprising, for example, a suitable agent for dispersing, solubilising, or extending release of the active ingredient, a propellant as solvent.

Prior to use as dry powder or suspension, the medicinal product is typically micronised to a size suitable for delivery by inhalation (typically less than 5 microns). This may be achieved by any appropriate comminuting method, such as spiral jet milling, fluid bed jet milling, supercritical fluid processing to form nanoparticles, high pressure homogenisation, or spray drying.

Capsules, blisters and cartridges for use in an inhaler or insufflator may be formulated to contain a powder mix of the compound of the present invention, a suitable powder base and a performance modifier.

A suitable solution formulation for use in an atomiser using electrohydrodynamics to produce a fine mist may contain a suitable dose of the compound of the present invention per actuation and the actuation volume may, for example, vary from 1 μL to 100 μL .

Suitable flavours, such as menthol and levomenthol, or sweeteners, such as saccharin or sodium saccharin, may be added to those formulations of the present invention intended for inhaled/intranasal administration.

Formulations may be formulated to be immediate and/or modified release. Modified release formulations include delayed, sustained, pulsed, controlled, targeted and programmed release.

In one embodiment, the present invention relates to the prophylaxis or method of treatment of a disease or disorder associated with increased activity of cyclin-dependent protein kinases CDK7, which comprises administering in a therapeutically effective amount the compound of the present invention or a pharmacologically acceptable salt, solvate or stereoisomer thereof, or the pharmaceutical composition of the present invention to a subject in need of such treatment or prophylaxis.

In another embodiment, the present invention relates to the prophylaxis or method of treatment of a disease or disorder, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinases CDK7 is an oncological, infectious inflammatory or non-infectious inflammatory, autoinflammatory, autoimmune disease.

In another embodiment, the present invention relates to the prophylaxis or method of treatment of a disease or disorder, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinases CDK7 is an oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal

cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma;

an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angiitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjogren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, otitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse

myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis.

In another embodiment, the present invention relates to the prophylaxis or method of treatment of a disease or disorder, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinases CDK7 is an oncological disease selected from a group comprising breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer.

It is meant that the compounds of the present invention may be used in the prophylaxis, methods of treatment as described above, may be used in the treatment as described above, and/or may be used in the manufacture of therapeutical products (medication) for treatment as described above.

In one embodiment, the present invention relates to the use of the compound according to the present invention or a pharmacologically acceptable salt, solvate or stereoisomer thereof, or the pharmaceutical composition described above for the treatment of a disease or disorder associated with increased activity of CDK7 in a subject in need of such treatment.

In one embodiment, the present invention relates to the use of the compound according to the present invention or a pharmaceutically acceptable salt, solvate or stereoisomer thereof, or the pharmaceutical composition described above for the treatment of a disease or disorder associated with increased activity of CDK7, wherein the disease or disorder associated with increased activity of CDK7 is an oncological, infectious inflammatory or non-infectious inflammatory, autoinflammatory, autoimmune disease.

In another one embodiment, the present invention relates to the use described above, wherein the disease or disorder associated with increased activity of CDK7 is an oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma; an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angiitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjogren's syndrome,

systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, otitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis.

In another embodiment, the invention relates to the above-described use, wherein the oncological disease is selected from a group comprising breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer.

The compounds of the invention may be administered alone or in combination with one or more other drug products or monoclonal antibody-based drug products (or in any combination thereof). The pharmaceutical compositions, methods and use of the invention thus also encompass embodiments of combinations (co-administration) with other active agents.

As used herein, the terms “co-administration”, “co-administered” or “in combination with” referring to the compounds with one or more other therapeutic agents, are intended to mean, refer to or include the following:

- simultaneous administration to said patient of such combination of the compound of the invention and therapeutic agent to a patient in need of treatment, when such components are formulated together into a single dosage form which releases said components at substantially the same time,
- simultaneous administration of such combination of the compound of the invention and therapeutic agent to a patient in need of treatment, when such components are formulated apart from each other into separate dosage forms which are administered at substantially the same time to said patient, whereupon said components are released at substantially the same time in said patient,
- sequential administration of such combination of the compound of the invention and therapeutic agent to a patient in need of treatment, when such components are formulated apart from each other into separate dosage forms which are taken at consecutive times by said patient with a significant time interval between each administration, whereupon said components are released at different times in said patient; as well as
- sequential administration of such combination of the compound of the invention and therapeutic agent to a patient in need of treatment, when such components are formulated together into a single dosage form which releases said components in a controlled manner whereupon they are concurrently, consecutively, and/or overlappingly released at the same and/or different times to said patient, wherein each part may be administered by either the same or a different route.

As well known to those skilled in the art, therapeutically effective dosages may vary when the drug products are used in combination treatment. Methods for experimentally determining therapeutically effective dosages of drug products and other agents for use in combination treatment regimens have been described in the literature. For example, the use of metronomic dosing, i.e., providing more frequent, lower doses in order to minimize toxic side effects, has been described in the literature. Combination treatment further includes periodic treatments that start and stop at various times in accordance with the patient treatment plan. For combination therapy described in the present patent, dosages of co-administered compounds may vary depending on the type of co-drug employed, on the specific drug employed, on the condition or disorder being treated and so forth.

The antitumor treatment described above can be used either as a stand-alone therapy, or in combination with surgery, or radiotherapy, or drug therapy. Such therapy may be administered concurrently, simultaneously, sequentially or separately with the treatment of the compound according to the invention and may comprise one or more agents from the following categories of antitumor agents: antiproliferative/antitumor therapeutical products and combinations thereof used in medical oncology, such as alkylating agents, alkylated sulfonates, nitrosoureas or triazenes; antimetabolites; hormonal agents or hormone antagonists; platinum compounds; antitumor antibiotics; topoisomerase inhibitors.

Examples of antimetabolites include, without limitation, folic acid antagonists (e.g., methotrexate, trimetrexate, pemetrexed, pralatrexate, raltitrexed, calcium levofolinate) or pyrimidine antagonists (e.g., cytarabine, tegafur, fluorouracil, capecitabine, floxouridine, azacitidine, enocitabine, karmofur, gemcitabine, sapacitabine, elacitarabine, doxifluridine), or purine antagonists (for example, mercaptopurine, thioguanine, pentostatin, fludarabine, cladribine, nelarabine, azathioprine, clofarabine), or asparaginase.

Examples of alkylating agents include, without limitation, mechlorethamine, cyclophosphamide, chlorambucil, melphalan, bendamustine, hexamethylmelamine, thiotepa,

busulfan, carmustine, lomustine, larmustine, semustine, streptozocin, dacarbazine, ifosfamide, improsulfan, mitobronitol, mitolactol, nimustine, ranimustine, temozolomide, treosulfan, carboquinone, apaziquinone, fotemustine, altretamine, gluphosphamide, pipobroman, trophosphamide, uramustine, evophosphamide, VAL-083.

Examples of hormonal agents and hormone antagonists include, without limitation, hydrocortisone, dexamethasone, prednisone, prednisone, hydroxyprogesterone caproate, megestrol acetate, medroxyprogesterone acetate, diethylstilbestrol, estradiol, tamoxifen, testosterone propionate, fluoxymesterone, flutamide, leuprolide, abarelix, abiraterone, bicalutamide, buserelin, calusteron, chlorotrianizene, degarelix, dexamethasone, fluocortolone, fulvestrant, goserelin, histrelin, leuprorelin, mitotane, nafarelin, nandrolone, nilutamide, octreotide, raloxifene, thyrotropin alpha, toremifene, triptorelin, diethylstilbestrol, acolbifene, danazol, deslorelin, epitostanol, orteronel, enzalutamide, aminoglutethimide, anastrozole, exemestane, fadrozole, letrozole, testolactone, formestane.

Examples of platinum compounds include, without limitation, cisplatin, carboplatin, oxaliplatin, eptaplatin, miriplatin hydrate, lobaplatin, nedaplatin, picoplatin, satraplatin.

Examples of antitumor antibiotics include, without limitation, doxorubicin, daunorubicin, idarubicin, carubicin, valrubicin, zorubicin, aclarubicin, pirarubicin, nemorubicin, amrubicin, epirubicin, bleomycin, dactinomycin, plicamycin, peplomycin, mitomycin-C, zinostatin, streptozotocin.

Examples of topoisomerase inhibitors include, without limitation, irinotecan, topotecan, belotecan, teniposide, etoposide, voreloxin, amonafide.

Examples of antitumor agents include, without limitation, any one of the following agents: microtubule targeting drugs, such as taxanes (e.g., paclitaxel, nab-paclitaxel, docetaxel, cabazitaxel, tesetaxel), Vinca alkaloids (e.g., vinorelbine, vinblastine, vincristine, vindesine, vinflunine); mitogen-activated protein kinases inhibitors (e.g., U0126, PD98059, PD184352, PD0325901, ARRY-142886, SB239063, SP600125, BAY 43-9006, wortmannin or LY294002); mTOR inhibitors (e.g. sirolimus, temsirolimus, everolimus, ridaforolimus); antibodies (for example, prolgolimab, rituximab, trastuzumab, alemtuzumab, besilesomab, cetuximab, denosumab, ipilimumab, bevacizumab, pertuzumab, pembrolizumab, nivolumab, cemiplimab, durvalumab, atezolizumab, avelumab, ofatumumab, panitumumab, tozitinomab, catumaxomab, elotuzumab, epratuzumab, farletuzumab, mogamulizumab, necitumumab, nimotuzumab, obinutuzumab, okaratumumab, oregovomab, ramucirumab, rilotumumab, siltuximab, tocilizumab, zalutuzumab, zanolimumab, matuzumab, dalotuzumab, onartuzumab, rakotumomab, tabalumab, abituzumab); kinase inhibitors (fosmatanib, entospletinib, erlotinib, imatinib, lapatinib, nilotinib, pazopanib, vemurafenib, gefitinib, crizotinib, dazatinib, regorafenib, ruxolitinib, sorafenib, sunitinib, vandetanib, bozutinib, axitinib, afatinib, alisertib, dabrafenib, dacomitinib, dinaciclib, dovitinib, nintedanib, lenvatinib, linifanib, linsitinib, masitinib, motesanib, neratinib, orantinib, ponatinib, radotinib, tipifarnib, tivantinib, tivozanib, trametinib, apatinib, ibrutinib, acalabrutinib, cobimetinib, fedratinib, brivanib alaninate, cediranib, cabozantinib, icotinib, cipatinib, rigosertib, pimasertib, buparlisib, idelalisib, midostaurin, perifosine, tesevatinib); photosensitizers (e.g., talaporfin, temoporfin, porfimer sodium); cytokines (e.g., aldesleukin, interferon alfa, interferon alfa-2a, interferon alfa-2b, celmoleukin, tasonermin, recombinant interleukin-2, oprelvekin, recombinant interferon beta-1a); vaccines (e.g., picibanil, sipuleucel-T, vitespen, emepepimut-S, oncoVAX, rindopepimut, troVAX, MGN-1601, MGN-1703); bisantrene, decitabine, mitoxantrone, procarbazine, trabectedin, amsacrin, brostallicin, miltefosin, romidepsin, plitidepsin, eribulin, ixabepilone, fosbretabulin, denileukin diftitox, ibritumomab tiuxetan,

prednimustine, trastuzumab emtansine, estramustin, gemtuzumab ozogamicin, aflibercept, oportuzumab monatox, cintredekin besudotox, edotreotide, inotuzumab ozogamicin, naptumomab estafenatox, vintafolide, brentuximab vedotin, bortezomib, ixazomib, carfilzomib, lenalidomide, thalidomide, pomalidomide, zoledronic acid, ibandronic acid, pamidronic acid, alitretinoin, tretinoin, peretinoin, bexarotene, tamibarotene, imiquimod, lentinan, mifamurtide, romurtide, pegaspargase, pentostatin, endostatin, sizofiran, vismodegib, vorinostat, entinostat, panobinostat, celecoxib, cilengitide, etanidazole, ganetespib, idronoxil, iniparib, lonidamine, nimorazole, procodazole, tasquinimod, telotristat, belinostat, thymalfasin, tirapazamine, tosedostat, trabedersen, ubenimex, valspodar, gendicine, reolisin, retaspimicin, trebananib, virulisin.

Thus, in another embodiment of the invention, there is described a pharmaceutical product comprising a compound of formula (I) or pharmaceutically acceptable salt, solvate or stereoisomer thereof as defined hereinbefore, in combination with an anti-tumour agent as defined hereinbefore, intended for the conjoint treatment of cancer.

Dosage regimens may be adjusted to provide the optimum desired response. For example, a single dose may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate peroral compositions in a unit dosage form for ease of administration and uniformity of dosage. A unit dosage form as used herein refers to physically discrete units suited as unitary dosages for patients/subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the desired pharmaceutical carrier. Specification for the unit dosage forms of the invention is typically dictated by and directly dependent on (a) the unique characteristics of a therapeutic agent and particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in subjects.

Thus, a skilled artisan would appreciate, based upon the disclosure provided herein, that the doses and dosage regimen are adjusted in accordance with methods well-known in the therapeutic arts. That is, the maximum tolerable dose can be readily established, and the effective amount providing a detectable therapeutic effect to a patient may also be determined, as can the temporal requirements for administering each agent to provide a detectable therapeutic effect to a patient. Thus, while certain doses and administration regimens are exemplified herein, these examples in no way limit the doses and administration regimens that may be provided to a patient in practicing the embodiments of the invention.

It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated, and may include single or multiple doses. Furthermore, it is to be understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the judgment of a medical professional administering or supervising the administration of the compositions, and that dosage ranges set forth in the present description are exemplary only and are not intended to limit the scope or practice of the claimed compositions. Furthermore, the dosage regimen with the compositions of the present invention may be based on a variety of factors, including the type of disease, age, weight, sex, medical condition of the patient, the severity of the condition, the route of administration, and the particular compound of the present invention employed. Thus, the dosage regimen may vary widely, but can be determined routinely using standard methods. For example, doses may be adjusted based on pharmacokinetic and pharmacodynamic parameters, which may include clinical effects such as toxic effects or laboratory values. Thus, the present invention encompasses intra-patient dose-escalation as

determined by one skilled in the art. Methods for determining appropriate dosage and regimens are well-known in the art and would be understood by a skilled artisan once provided the ideas disclosed herein.

Typically, doses used to treat an adult are usually in the range of 0.01-5000 mg per day, or from 0.1-500 mg per day or from 0.1 to 100 mg per day.

Once improvement of the patient's conditions has occurred, a maintenance dose is administered, if necessary. Subsequently, the dosage or the frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. Patients may be however required periodic treatment on a long-term basis upon any relapse of symptoms.

The foregoing ranges are merely suggestive, as the number of variables in regard to an individual treatment regime is large, and considerable excursions from these recommended values are not uncommon. These dosages may be altered depending on a number of variables, not limited to the activity of the compound used, the disorder or condition to be treated, the method of administration, the requirements of the individual subject, the severity of the disorder or condition being treated, and the judgment of the physician.

The following examples are provided for better understanding of the invention. These examples are for purposes of illustration only and are not to be construed as limiting the scope of the invention in any manner.

The compounds and processes of the present invention will be better understood in connection with the following synthetic schemes which illustrate the methods by which the compounds of the present invention can be prepared. The initial products can be obtained from commercial sources or prepared using conventional prior art methods known to one of ordinary skill in the art. One of ordinary skill in the art will also readily appreciate that the steps of selective protection and deprotection, as well as the order of these steps, may be carried out in different order, depending on the nature of the substituents, for the successful completion of the synthesis described below.

The abbreviations used in this description, including those given in the illustrative diagrams and examples that follow, are well known to one of ordinary skill in the art.

DIPEA – diisopropylethylamine;

TBAF – tetrabutylammonium fluoride;

DMF – N,N-dimethylformamide;

THF – tetrahydrofuran;

BINAP – (±)-2,2'-bis(diphenylphosphino)-1,1'-dinaphthalene;

MTBE – methyl *tert*-butyl ether;

TMEDA - *N,N,N',N'*-tetramethylethane-1,2-diamine;

HATU - 1-[bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxide hexafluorophosphate;

NBS – *N*-bromosuccinimide;

TEA – triethylamine;

DMAP - 4-dimethylaminopyridine;

TFA – trifluoroacetic acid;

TIPS – triisopropyl silane;

dppf – 1,1'-bis(diphenylphosphino)ferrocene;


ac – acetyl;

boc – *tert*-butoxycarbonyl;

pin – pinacolato;
 HOBT – 1-hydroxybenzotriazole;
 EDC – 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide;
 PPA – polyphosphoric acid;
 HPLC – high performance liquid chromatography;
 Ph – phenyl;
 HMDS – hexamethyldisilazane.

Examples

Example 1. Synthesis of compound 1.6.



Synthesis of compound 1.2

TEA (4.28 g, 42.2 mmol), DMAP (0.679 g, 5.50 mmol) and benzenesulfonyl chloride (3.64 g, 21.7 mmol) were added to solution of **1.1** (4.00 g, 18.34 mmol) in 40 ml of dichloromethane. The reaction mass was stirred for 20 h at room temperature. After standard processing, the product was isolated using silica gel column chromatography. 5.63 g (86%) of compound **1.2** were yielded.

Synthesis of compound 1.3

Compound **1.2** (2.5 g, 6.98 mmol), (BPin)₂ (3.62 g, 13.96 mmol), KOAc (1.77 g, 17.45 mmol), Pd(dppf)Cl₂ (0.417 g, 0.56 mmol) in 50 ml of 1,4-dioxane were stirred in nitrogen atmosphere for 10 h at 100°C. After standard treatment, the reaction product was isolated using silica gel column chromatography. 2.16 g (76%) of compound **1.3** were yielded.


Synthesis of compound 1.5

Compound **1.3** (0.90 g, 2.0 mmol), pyrimidine **1.4** (0.692 g, 1.82 mmol), Na₂CO₃ (0.448 g, 4.18 mmol), Pd(dppf)Cl₂ (0.136 g, 0.18 mmol) in a mixture of 18 ml of 1,4-dioxane and 6 ml of water were stirred in nitrogen atmosphere for 4 h at 100°C. After standard treatment, the product was isolated using silica gel column chromatography. 0.95 g (84%) of compound **1.5** were yielded.

Synthesis of compound 1.6

Compound **1.5** (0.90 g, 1.44 mmol) was dissolved in 9 ml of ethanol; NaOH (0.35 g, 8.64 mmol) and 9 ml of water were added. The reaction mass was stirred at 60°C for 6 h, the solvent was removed, water was added to the residue until it completely dissolved (10 ml), pH of the mixture was brought to 3 with a 10% citric acid solution. The precipitate was filtered, the residual solvents were removed in vacuo. 0.6 g (92%) of product **1.6** was yielded.

Example 2. Synthesis of compound 2.4.



Synthesis of compound 2.1

Compound **1.1** (9.08 g, 41.6 mmol) was dissolved in 90 ml of ethanol; NaOH (6.73 g, 167 mmol) and 45 ml of water were added. The reaction mass was stirred at 70°C for 6 h, ethanol was removed in vacuo, water was added to the residue until it completely dissolved, pH of the mixture was brought to 3 with HCl_{conc.}, the precipitate was filtered off, washed with water, residual solvents were removed in vacuo. 7.1 g (90%) of compound **2.1** were yielded.

Synthesis of compound 2.2

Compound **2.1** (3.52 g, 18.5 mmol) was dissolved in 40 ml of dichloromethane; oxalyl chloride (9.41 g, 74.1 mmol) and 0.3 ml of DMF were added. The reaction mass was stirred for 2 h, volatile components were removed in vacuo, the residue was dissolved in 20 ml of dichloromethane and added while being cooled on an ice bath to a solution of *tert*-butanol (2.77 g, 37.1 mmol) and TEA (15.0 g, 148 mmol) in 10 ml of dichloromethane, stirred for 20 h at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 3.7 g (81%) of compound **2.2** were yielded.


Synthesis of compound 2.3

TEA (4.28 g, 42.2 mmol), DMAP (0.22 g, 1.79 mmol) and benzenesulfonyl chloride (3.24 g, 20.2 mmol) were added to a solution of **2.2** (4.5 g, 18.3 mmol) in 50 ml of dichloromethane. The reaction mass was stirred for 20 h at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 6.73 g (80%) of product **2.3** were yielded.

Synthesis of compound 2.4

TFA (1.37 g, 12.0 mmol) was added to a solution of **2.3** (0.579 g, 1.50 mmol) in 12 ml of dichloromethane, the reaction mass was stirred for 20 h at room temperature. Volatile components were removed in vacuo. 0.43 g (95%) of product **2.4** was yielded.

Example 3. Synthesis of compounds CDK7_536, CDK7_537, CDK7_833, CDK7_835, CDK7_846, CDK7_847, CDK7_848, CDK7_849, CDK7_850, CDK7_851, CDK7_951, CDK7_997, CDK7_998, CDK7_1001, CDK7_1003.














Synthesis of compound 3.1

Compound **1.6** (0.065 g, 0.14 mmol), HATU (0.108 g, 0.28 mmol), cyclopropylamine (0.012 g, 0.21 mmol), DIPEA (0.073 g, 0.56 mmol) were dissolved in 0.6 ml DMF and stirred for 20 h at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.035 g (51%) of compound **3.1** was yielded.


Synthesis of compound CDK7_951

Compound **3.1** (0.035 g, 0.07 mmol) was dissolved in TFA (0.403 g, 3.5 mmol) and stirred at 20 °C for 20 h. TFA was removed at reduced pressure, product was isolated using preparative HPLC. 0.015 g (54%) of compound **CDK7_951** was yielded.

The following compounds were produced analogously from compound **1.6** and corresponding amines:

Item no.	Compound	Original amine	Yield at stage 1, %	Yield at stage 2, %
1	CDK7_536		99	79
2	CDK7_833		39	3
3	CDK7_835		57	39
4	CDK7_846		66	58
5	CDK7_847		34	77
6	CDK7_848		69	54
7	CDK7_849		75	48
8	CDK7_850		50	96
9	CDK7_851		8 (total yield for two stages)	
10	CDK7_537		69	36
11	CDK7_1001		53	48
12	CDK7_1003		77	56
13	CDK7_997		99	19
14	CDK7_998		50	50

Example 4. Synthesis of compounds **CDK7_538**, **CDK7_539**, **CDK7_834**, **CDK7_836**, **CDK7_837**, **CDK7_838**



Synthesis of compound 4.2

Compound **2.4** (0.3 g, 0.91 mmol) was dissolved in 5 ml of dichloromethane; oxalyl chloride (0.289 g, 2.28 mmol) and 0.1 ml of DMF were added. The reaction mass was stirred for 3 h. Volatile components were removed in vacuo, the residue was dissolved in 5 ml of dichloromethane and added dropwise while stirring and cooling on an ice bath to a solution of 4-aminopyridine (0.087 g, 0.91 mmol) and TEA (0.386 g, 3.64 mmol) in 5 ml of dichloromethane. The reaction mass was stirred at room temperature for 20 h. After standard treatment, the product was isolated using column chromatography on silica gel. 0.3 g (80%) of compound **4.2** was yielded.

Synthesis of compound 4.3

Compound **4.2** (0.10 g, 0.25 mmol), (BPin)₂ (0.13 g, 0.50 mmol), KOAc (0.063 g, 0.63 mmol), Pd(dppf)Cl₂ (0.021 g, 0.03 mmol) were dissolved in 2 ml of 1,4-dioxane and heated in a sealed vessel while stirring in nitrogen atmosphere for 6 h at 100 °C. The reaction mixture was filtered off through a Celite layer, and the solvent was removed. Compound **1.4** (0.088 g, 0.33 mmol), Na₂CO₃ (0.088 g, 0.82 mmol), Pd(dppf)Cl₂ (0.025 g, 0.03 mmol), 5 ml of 1,4-dioxane and 1.7 ml of water were added to the residue. The reaction mass was heated in a sealed vessel in nitrogen atmosphere for 3 h at 100 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.075 g (44%) of compound **4.3** was yielded.

Synthesis of compound 4.4




An aqueous solution of NaOH (0.089 g, 2.2 mmol) was added to a solution of **4.3** (0.075 g, 0.11 mmol) in ethanol (1.5 ml) and stirred for 3 h at 50 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.042 g (72%) of compound **4.4** was yielded.

Synthesis of compound CDK7_836


Compound **4.4** (0.042 g, 0.08 mmol) was dissolved in 2 ml of dichloromethane; TFA (0.092 g, 0.8 mmol) was added, the reaction mass was stirred for 20 h at 20 °C, the solvent was removed. The product was isolated by HPLC. 0.012 g (34%) of compound **CDK7_836** was yielded.

The following compounds shown in the table below were produced analogously.

N	Product	Original amine	Yield, %, stage			
			1	2+3	4	5
1	CDK7_538		58	79	64	85
2	CDK7_539		**	19	42	77

3	CDK7_834		86	52	80	27
4	CDK7_837		39	23	*	9
5	CDK7_838		71	55	82	44

*Intermediate was not isolated



Synthesis of compound 4.5

Compound **2.1** (0.6 g, 3.16 mmol), *N*-methylpiperazine (0.425 ml, 3.79 mmol), HATU (1.58 g, 4.11 mmol), DIPEA (1.11 ml, 6.32 mmol) were dissolved in 6 ml of DMF and stirred at room temperature for 20 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.86 g (99%) of compound **4.5** was yielded.

Synthesis of compound 4.6

TEA (0.769 ml, 5.52 mmol), DMAP (0.023 g, 0.18 mmol) and benzenesulfonyl chloride (0.36 g, 2.02 mmol) were added to a solution of **4.5** (0.5 g, 1.84 mmol) in 15 ml of dichloromethane. The reaction mass was stirred for 20 h at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.56 g (74%) of product **4.6** was yielded.

Synthesis of compound 4.7

Compound **4.6** (0.56 g, 1.36 mmol), (BPin)₂ (0.698 g, 2.72 mmol), KOAc (0.413 g, 4.08 mmol), Pd(dppf)Cl₂ (0.112 g, 0.14 mmol) in 12 ml of 1,4-dioxane were stirred in nitrogen atmosphere for 10 h at 100°C. After standard treatment, the reaction product was isolated using silica gel column chromatography. 0.52 g (83%) of compound **4.7** was yielded.

Synthesis of compound 4.8

Boronic ether **4.7** (0.149 g, 0.32 mmol), pyrimidine **1.4** (0.1 g, 0.26 mmol), Na₂CO₃ (0.07 g, 0.65 mmol), Pd(dppf)Cl₂ (0.024 g, 0.03 mmol) in 2 ml of 1,4-dioxane and 0.7 ml of water were stirred in nitrogen atmosphere for 4 h at 100°C. After standard treatment, the reaction product was isolated using column chromatography on silica gel. 0.05 g (23%) of compound **4.8** was yielded.


Synthesis of compound 4.9

Compound **4.8** (0.05 g, 0.07 mmol), NaOH (0.065 g, 1.61 mmol) in 0.5 ml of ethanol and 0.5 ml of H₂O were stirred at 50°C for 2 h. 2 ml of 10% solution of citric acid was added, extraction was performed with ethyl acetate, and, following removal of the solvent in vacuo, 0.016 g (42%) of compound **4.9** was yielded.

Synthesis of compound **CDK7_539**

Compound **4.9** (0.016 g, 0.03 mmol) was dissolved in 1 ml of dichloromethane, 0.2 ml of 4 H HCl solution in 1,4-dioxane was added, the reaction mass was stirred at 20 °C for 18 h, volatile components were removed in vacuo, the product was isolated by preparative HPLC. 0.010 g (77%) of compound **CDK7_539** was yielded.

Example 5. Synthesis of compound **CDK7_535**



Synthesis of compound **5.2**

0.75 ml (1.84 mmol) 2.5 M of *n*-butyllithium solution in hexane was added while stirring in nitrogen atmosphere at -78 °C to a solution of compound **5.1** (0.45 g, 1.23 mmol) in 50 ml of THF, stirred for 30 min, benzoyl chloride (0.262 g, 1.84 mmol) was then added, and stirred for another 30 min. After standard treatment, the product was isolated using silica gel column chromatography. 0.3 g (62%) of compound **5.2** was yielded.

Synthesis of compound **5.3**

Pyrrole **5.2** (0.126 g, 0.35 mmol), (BPin)₂ (0.124 g, 0.48 mmol), KOAc (0.063 g, 0.63 mmol), Pd(dppf)Cl₂ (0.012 g, 0.02 mmol) were dissolved in 5 ml of 1,4-dioxane, heated in a sealed vessel for 7 h at 100°C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.05 g (36%) of compound **5.3** was yielded.

Synthesis of compound **5.4**

Boronic ether **5.3** (0.05 g, 0.1 mmol), pyrimidine **1.4** (0.027 g, 0.07 mmol), Na₂CO₃ (0.021 g, 0.2 mmol), Pd(dppf)Cl₂ (0.007 g, 0.01 mmol) in 1.5 ml of 1,4-dioxane and 0.5 ml of water were stirred in nitrogen atmosphere for 3 h at 90 °C. After standard treatment, the reaction product was isolated using column chromatography on silica gel. 0.04 g (62%) of compound **5.4** was yielded.


Synthesis of compound **5.5**

Compound **5.5** was produced analogously to compound **4.9** with a yield of 87%.

Synthesis of compound **CDK7_535**

Compound **CDK7_535** was produced analogously to compound **539** with a yield of 38%.

Example 6. Synthesis of compounds **CDK7_845**, **CDK7_924**, **CDK7_959**, **CDK7_961**, **CDK7_1052**.



Synthesis of compound **6.2**

Compound was produced analogously to compound **4.2** with a yield of 93%.

Synthesis of compound **6.3**

Amide **6.2** (0.29 g, 1.00 mmol) and 0.97 g of methanesulfonic acid (10 mmol) were stirred for 72 h at 20 °C, 5 ml of water was added while cooling to the reaction mass, stirred for 1 hour, the precipitate was filtered off, washed with water, residual solvents were removed in vacuo. 0.19 g (84%) of compound **6.3** was yielded.

Synthesis of compound **6.4**

TEA (0.26 g, 2.60 mmol), DMAP (0.01 g, 0.08 mmol) and benzenesulfonyl chloride (0.165 g, 0.92 mmol) were added to a solution of pyrrolopyridone **6.3** (0.19 g, 0.84 mmol) in 6 ml of dichloromethane, the reaction mass was stirred for 18 h at 20 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.26 g (73%) of compound **6.4** was yielded.

Synthesis of compound **6.5**

Compound **6.4** (0.22 g, 0.60 mmol), (BPin)₂ (0.31 g, 1.2 mmol), KOAc (0.153 g, 1.50 mmol), Pd(dppf)Cl₂ (0.045 g, 0.06 mmol) in 5 ml of 1,4-dioxane was heated while stirring in nitrogen atmosphere for 7 h at 100°C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.14 g (59%) of compound **6.5** was yielded.

Synthesis of compound **6.6**

Compound **6.5** (0.54 g, 1.30 mmol), pyrimidine **1.4** (0.495 g, 1.30 mmol), Na₂CO₃ (0.348 g, 3.25 mmol), Pd(dppf)Cl₂ (0.97 g, 0.13 mmol) in a mixture of 12 ml of 1,4-dioxane and 4 ml of water were stirred in nitrogen atmosphere for 4 h at 100 °C. After standard treatment, the reaction product was isolated using column chromatography on silica gel. 0.26 g (32%) of compound **6.6** was yielded.

Synthesis of compound **6.7**





The compound was produced analogously to compound **4.9** with a yield of 95%.

Synthesis of compound **CDK7_845**


The compound was produced analogously to compound **CDK7_539** with a yield of 38%.

The following compounds were analogously produced:

N	Compound	Initial compound	Stage, yield, %						
			1	2	3	4+5	6	7	

1	CDK7_959		53	48	81	41	77	35
2	CDK7_961		46	93	67	18	99	63
3	CDK7_924		54	50	92	65	91	38
4	CDK7_1052		25	62	76	70	80 (total yield for two stages)	

Example 7. Synthesis of compound CDK7_944



Synthesis of compound 7.3

Compound **7.1** (2.30 g, 12.9 mmol), compound **7.2** (1.49 g, 12.9 mmol) and Ag_2CO_3 (1.87 g, 0.065 mmol) in 23 ml of 1,4-dioxane and stirred for 5 h at 100 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 3.00 g (82%) of compound **7.3** were yielded.

Synthesis of compound 7.4

Compound **7.3** (3.00 g, 8.5 mmol) and LiOH (0.630 g, 25.5 mmol) in a mixture of 15 ml of water and 30 ml of methanol and stirred for 8 h at 55 °C. Methanol was removed at reduced pressure, pH of the mixture was adjusted to 2, precipitate was filtered off and residual solvents were removed in vacuo. 2.10 g (97%) of compound **7.4** were yielded.

Synthesis of compound 7.5

Compound **7.4** (1.070 g, 4.0 mmol), SOCl_2 (0.980 g, 8.0 mmol) and DMF (0.015 g, 0.2 mmol) in 10 ml of dichloromethane and stirred for 2 h at 40 °C. 5 ml of 4 N solution of HCl in 1,4-dioxane was added to the mixture and stirred for 16 h at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.540 g (99%) of compound **7.5** was yielded.

Synthesis of compound 7.6

Compound **7.5** (0.30 g, 2.09 mmol) was dissolved in 30 ml of THF, added at 0 °C to the resulting solution of NaH (0.125 g, 3.13 mmol), and the mixture was stirred for 30 minutes at 0 °C. TIPSCl (0.540 g, 2.72 mmol) was then added dropwise and stirred for 30 min at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.611 g (99%) of compound **7.6** was yielded.

Synthesis of compound 7.7

Compound **7.6** (0.611 g, 1.98 mmol) was dissolved in 20 ml of THF, a solution of NBS (0.356 g, 1.98 mmol) was added at -78 °C in 10 ml of THF, and the mixture was stirred for 30 minutes at -78 °C and for 30 minutes at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.735 g (100%) of compound **7.7** was yielded.

Synthesis of compound 7.8

Compound **7.7** (0.625 g, 1.60 mmol) and 30% TBAF solution in tetrahydrofuran (1.59 g, 1.76 mmol) were mixed in 20 ml of THF. The mixture was stirred for 30 minutes at 25 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.240 g (70%) of compound **7.8** was yielded.

Synthesis of compound 7.9

Compound **7.8** (0.240 g, 1.06 mmol) was dissolved in 30 ml of THF; NaH (0.064 g, 1.59 mmol) was added at 0 °C to the resulting solution and the mixture was stirred for 30 minutes, benzenesulfonyl chloride (0.227 g, 1.27 mmol) was added and stirred for 30 minutes at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.31 g (82%) of compound **7.9** was yielded.

Synthesis of compound 7.10

Compound **7.10** was produced analogously to compound **1.3** with a yield of 80%.

Synthesis of compound 7.11

Compound **7.11** was produced analogously to compound **1.5** with a yield of 65%.


Synthesis of compound 7.12

Compound **7.12** was produced analogously to compound **1.6** with a yield of 97%.

Synthesis of compound CDK7_944

Compound **CDK7_944** was produced analogously to compound **CDK7_951** with a yield of 60%.

Example 8. Synthesis of compound CDK7_929.



Synthesis of compound 8.2

A solution of 4.6 g (0.026 mol) of compound **8.1** in 50 ml of THF was slowly added to a suspension of 2.00 g (0.053 mol) of LiAlH₄ in 100 ml of THF while stirring and cooling. The reaction mass was stirred for 5 h at 100 °C, cooled to 0 °C, and 50 ml of water was added. After standard processing, the product was isolated using silica gel column chromatography. 0.5 g (12%) of compound **8.2** was yielded.

Synthesis of compound **8.3**

Compound **8.3** was produced analogously to compound **6.2** with a yield of 70%.

Synthesis of compound **8.4**

Compound **8.4** was produced analogously to compound **6.3** with a yield of 13%.

Synthesis of compound **8.5**

Compound **8.5** was produced analogously to compound **6.4** with a yield of 56%.

Synthesis of compound **8.6**

Compound **8.6** was produced analogously to compound **6.5** with a yield of 64%.

Synthesis of compound **8.7**

Compound **8.7** was produced analogously to compound **6.6** with a yield of 31%.


Synthesis of compound **8.8**

Compound **8.8** was produced analogously to compound **6.7** with a yield of 95%.

Synthesis of compound **CDK7_929**

Compound **CDK7_929** was produced analogously to compound **CDK7_845** with a yield of 90%.

Example 9. Synthesis of compound **CDK7_966**



Synthesis of compound **9.2**

Compound **9.1** (2.4 g, 8.87 mmol) was dissolved in 30 ml of methanol. 50 ml (44.5 g, 0.395 mmol) of a 40% aqueous solution of dimethylamine was added. The reaction mass was stirred for 24 h, the solvent was removed, the product was isolated using column chromatography on silica gel. 0.5 g (21%) of compound **9.2** was yielded.

Synthesis of compound **9.3**

Compound **9.2** (0.40 g, 1.48 mmol) was dissolved in 10 ml of dichloromethane, a solution of NBS (0.38 g, 2.13 mol) in 10 ml of dichloromethane was added. The reaction mass was stirred for 1 h. After standard treatment, the product was isolated using column chromatography on silica gel. 0.16 g (31%) of compound **9.3** was yielded.

Synthesis of compound **9.4**

The compound **9.3** (0.05 g, 0.14 mmol), (BPin)₂ (0.047 g, 0.18 mmol), KOAc (0.042 g, 0.42 mmol), Pd(dppf)Cl₂ (0.01 g, 0.01 mmol) was dissolved in 1 ml of 1,4-dioxane, heated in a sealed vessel for 1 h at 100° C. The reaction mixture was filtered off through Celite and the solvent was removed in vacuo. 0.055 g (99%) of product in the form of brown oil was yielded.


Synthesis of compound **9.5**

Compound **9.5** was produced analogously to compound **1.5** with a yield of 93%.

Synthesis of compound **CDK7_966**

Compound **CDK7_966** was produced analogously to compound **CDK7_951** with a yield of 69%.

Example 10. Synthesis of compound **CDK7_972**



Synthesis of compound **10.2**

Compound **10.2** was derived from compound **10.1** analogously to compound **1.3** with a yield of 98%.


Synthesis of compound **10.3**

Compound **10.3** was produced analogously to compound **1.5** with a yield of 3%.

Synthesis of compound **CDK7_972**

Compound **CDK7_972** was produced analogously to compound **CDK7_951** with a yield of 33%.

Example 11. Synthesis of compound **CDK7_921**



Synthesis of compound **11.2**

Compound **11.2** was produced analogously to compound **1.3** with a yield of 62%.


Synthesis of compound **11.3**

Compound **11.3** was produced analogously to compound **1.5** with a yield of 34%.

Synthesis of compound **CDK7_921**

Compound **CDK7_921** was produced analogously to compound **CDK7_951** with a yield of 34%.

Example 12. Synthesis of compound **CDK7_901**



Synthesis of compound **12.2**

Compound **12.2** was produced analogously to compound **1.3** with a yield of 87%.


Synthesis of compound **12.3**

Compound **12.3** was produced analogously to compound **1.5** with a yield of 46%.

Synthesis of compound **CDK7_901**

Compound **CDK7_901** was produced analogously to compound **CDK7_951** with a yield of 89%.

Example 13. Synthesis of compound CDK7_766.




Synthesis of compound 13.2

Boronic ether **13.1** (0.04 g, 0.17 mmol), pyrimidine **1.4** (0.059 g, 0.15 mmol), Pd(OAc)₂ (0.003 g, 0.015 mmol), PPh₃ (0.008 g, 0.03 mmol), Na₂CO₃ (0.05 g, 0.46 mmol) in 0.37 ml of 1,4-dioxane and 0.04 ml of water were stirred at 100 °C in nitrogen atmosphere for 8 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.046 g (68%) of compound **13.2** was yielded.

Synthesis of compound CDK7_766

Compound **13.2** (0.025 g, 0.05 mmol) was dissolved in 3 ml of dichloromethane, 0.2 ml of TFA was added, stirred for 18 h at 20 °C, the solvent was removed in vacuo, the product was isolated by preparative HPLC. 0.008 g (40%) of compound **CDK7_766** was yielded.

Example 14. Synthesis of compounds CDK7_809, CDK7_810, CDK7_811, CDK7_829, CDK7_830, CDK7_897



Synthesis of compound 14.1

Method a:

NaH (0.014 g, 0.302 mmol) was added to a solution of indole (0.031 g, 0.276 mmol) in 2 ml of DMF, stirred for 30 min, a solution of pyrimidine **1.4** (0.100 g, 0.263 mmol) in 1 ml of DMF was added, the reaction mass was stirred for 20 h, a saturated solution of NaHCO₃ was added, extraction was performed with ethyl acetate. The product was isolated using column chromatography on silica gel using a mixture of hexane-ethyl acetate (92:8) as an eluent. 0.081 mg (67%) of compound **14.1** was yielded.

Method b is analogous to method A, K₂CO₃ was used instead of NaH, and MeCN was used instead of DMF; instead of stirring at room temperature, the reaction mixture was heated while boiling until the initial reagents disappeared.






Synthesis of compound CDK7_809

Method a:


Compound **14.1** (0.063 g, 0.137 mmol) was dissolved in 1.5 ml of 7.65 M solution of HCl in 1,4-dioxane. The reaction mass was stirred for 18 h at 20 °C. The solvent was removed in vacuo, the residue was treated with ether, the precipitate was filtered off and the residual solvents were removed in vacuo. 0.014 g (24%) of compound **CDK7_809** in the form of hydrochloride was yielded.

Method b is analogous to method A, TFA was used instead of 7.65 M solution of HCl in 1,4-dioxane, and dichloromethane was used as a solvent.

The following compounds were produced analogously:

Item no.	Compound	Initial compound	Reaction conditions, yield at stage 1, %	Reaction conditions, yield at stage 2, %
1	CDK7_810		a , 75%	b , 52%
2	CDK7_811		a , 43%	a , 79%
3	CDK7_829		b , 90%	b , 78%
4	CDK7_830		a , 178%	b , 67%
5	CDK7_897		b , 92%	b , 43%

Example 15. Synthesis of compounds **CDK7_548**, **CDK7_550**, **CDK7_735**, **CDK7_754**, **CDK7_888**, **CDK7_891**, **CDK7_898**, **CDK7_900**, **CDK7_903**, **CDK7_904**



Synthesis of compound **15.2**

DIPEA (0.214 g, 1.66 mmol) was added to a solution of tert-butyl N-(2-hydroxyphenyl)carbamate of compound (0.289 g, 1.38 mmol) in 3 ml of absolute ethanol. A solution of compound **15.1** (0.300 g, 1.38 mmol) in 2 ml of absolute ethanol was added to the resulting mixture at 0°C. The mixture was stirred at 0°C for 30 minutes, then at room temperature for 16 h. After standard treatment, the product was isolated using silica gel column chromatography. Product **15.2** (0.45 g, 84%) was yielded.

Synthesis of compound **15.3**

Compound **15.2** (0.445 g, 1.14 mmol), tert-butyl-3(S)-3-aminopiperidine carboxylate (0.229 g, 1.14 mmol) and DIPEA (0.221 mg, 1.71 mmol) in 2 ml of *i*-PrOH were stirred at room temperature for 16 h. After standard processing, the product was isolated using silica gel column chromatography, and then by preparative TLC. 0.057 g (9%) of compound **15.3** was yielded.

Synthesis of compound **CDK7_548**

1 ml of 7.65 M solution of HCl in 1,4-dioxane was added to a solution of compound **15.3** (0.057 g, 0.107 mmol) in 1 ml of ethanol. The mixture was stirred at room temperature for 12 h, volatile components were removed, the residue was recrystallized from a mixture of MTBE:acetone:*i*-PrOH (4:1:1). 0.015 g (yield 31%) of compound **CDK7_548** in the form of hydrochloride was yielded.

The following compounds were produced analogously:


Item no.	Compound	Initial compound	Reaction conditions, yield at stage 1, %	Reaction conditions, yield at stage 2, %	Reaction conditions, yield at stage 3, %
1	CDK7_550		a , 16%	b , 98%	a , 83%
2	CDK7_735		a , 21%	b , 98%	a , 92%
3	CDK7_754		a , 25%	a , 98%	a , 32%
4	CDK7_888		a , 78%	b , 66%	b , 47%
5	CDK7_891		a , 68%	c , 39%	b , 9%
6	CDK7_898		a , 37%	b , 98%	b , 62%
7	CDK7_900		a , 27%	b , 43%	b , 43%
8	CDK7_903		b , 9%	d , 94%	b , 86%
9	CDK7_904		b , 45%	d , 97%	b , 51%

Reaction conditions: Stage 1 **a**: DIPEA, EtOH, r. t.; **b**: DIPEA, MeCN, r. t. or 40 °C

Stage 2 **a**: **a**: DIPEA, *i*-PrOH, r.t.; **b**: DIPEA, *i*-PrOH, boiling;
c: DIPEA, MeCN, r. t.; **d**: DIPEA, MeCN, boiling.

Stage 3: **a**: HCl, 1,4-dioxane; **b**: TFA, dichloromethane.

Example 16. Preparation of compound **CDK7_991**



Synthesis of compound **16.2**

It was produced analogously to compound **4.2** from compound **16.1**. Yield: 64%.

Synthesis of compound **16.3**

Compound **16.3** was produced analogously to compound **1.2** with a yield of 92%.

Synthesis of compound **16.4**

Compound **16.4** was produced analogously to compound **1.3** with a yield of 38%.

Synthesis of compound **16.5**

Compound **16.5** was produced analogously to compound **1.5** with a yield of 32%.


Synthesis of compound **16.6**

Compound **16.6** was produced analogously to compound **4.5** with a yield of 99%.

Synthesis of compound **CDK7_991**

Compound **CDK7_991** was produced analogously to compound **CDK7_951** with a yield of 26%.

Example 17. Preparation of compound **CDK7_1017**



Synthesis of compound **17.1**

NaH (60% suspension in paraffin oil) (0.087 g, 2.17 mmol) at 0 °C was added to a solution of 4-bromo-N,N-dimethyl-1H-pyrrol-2-carboxamide (0.332 g, 1.45 mmol) in 15 ml of THF, the reaction mass was stirred for 30 min, CH₃I (0.270 g, 1.88 mmol) was added. The reaction mixture was heated to room temperature and stirred for 24 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.33 g (99%) of compound **17.1** was yielded.

Synthesis of compound **17.2**

Compound **17.2** was produced analogously to compound **1.3** with a yield of 98%.


Synthesis of compound **17.3**

Compound **17.3** was produced analogously to compound **1.5** with a yield of 40%.

Synthesis of compound **CDK7_1017**

Compound **CDK7_1017** was produced analogously to compound **CDK7_951** with a yield of 10%.

Example 18. Synthesis of compound **CDK7_953**.



Synthesis of compound **18.2**

Compound **18.2** was produced analogously to compound **8.3** from acid **2.1** and amine **18.1** with a yield of 71%

Synthesis of compound **18.3**

TFA (4.17 g, 36.2 mmol) was added to a solution of amide **18.2** (0.60 g, 1.81 mmol) in 5 ml of dichloromethane, the reaction mass was stirred for 18 h at 20 °C, volatile components were removed in vacuo, polyphosphoric acid (PPA) (9.34 g, 36.2 mmol) and phosphoric anhydride (1.05 g, 3.62 mmol) were added to the residue, heated while stirring at 110 °C for 3 h, 10 g of ice was added, stirred for 2 h, precipitate was filtered off, washed with water, residual solvents were removed in vacuo. 0.32 g (69%) of compound **18.3** was yielded.

Synthesis of compound **18.4**

NaBH₄ (0.053 g, 1.32 mmol) was added while cooling on an ice bath to a solution of compound **18.3** (0.100 g, 0.33 mmol) in 3 ml of ethanol, the reaction mass was stirred for 1 hour at the same temperature, then for 72 h at 20 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.076 g (88%) of compound **18.4** was yielded.

Synthesis of compound **18.5**

A 4 N solution of HCl in 1,4-dioxane (0.19 ml, 0.76 mmol) was added to a solution of compound **18.4** (0.22 g, 0.76 mmol) in 11 ml of chloroform, the reaction mass was stirred for 3 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.14 g (77%) of product **18.5** was yielded.

Synthesis of compound **18.6**

TEA (0.52 g, 5.00 mmol), DMAP (0.024 g, 0.2 mmol) and benzenesulfonyl chloride (0.21 g, 1.18 mmol) were added to a solution of compound **18.5** (0.14 g, 0.49 mmol) in 6 ml of dichloromethane, the reaction mass was stirred for 40 h at 20 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.15 g (78%) of product **18.6** was yielded.

Synthesis of compound **18.7**

Compound **18.7** was synthesized analogously to **6.5** from 0.15 g of compound **18.6**. Following borylation, the reaction mass was filtered off through a silica gel layer, the solvent was removed, yield was 0.54 g of the product in the form of a dark red oil comprising 30% of the target compound **18.7**, the product was used at the next stage without additional purification.

Synthesis of compound **18.8**

Compound **18.7** (0.54 g, 0.38 mmol), pyrimidine **1.4** (0.111 g, 0.29 mmol), Na₂CO₃ (0.094 g, 0.88 mmol), Pd(dppf)Cl₂ (0.022 g, 0.03 mmol) in a mixture of 6 ml of 1,4-dioxane and 2 ml of water were mixed in nitrogen atmosphere for 3 h at 100 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.081 g (43%) of compound **18.8** was yielded.

Synthesis of compound **18.9**

Compound **18.9** was produced analogously to compound **6.7** from 0.081 g of compound **18.8** with a yield of 91%.


Synthesis of compound **18.10**

Pd/C 5% (0.25 g, 0.12 mmol) was added to a solution of compound **18.9** (0.06 g, 0.12 mmol) in 6 ml of methanol. Hydrogenation was performed at atmospheric pressure for 3 h, the reaction mass was filtered off through a Celite layer, the solvent was removed in vacuo. 0.033 g (54%) of compound **18.10** was yielded.

Synthesis of compound **CDK7_953**

TFA (0.346 g, 3 mmol) was added to a solution of compound **18.10** (0.033 g, 0.06 mmol) in 2 ml of dichloromethane, the reaction mass was stirred at 20 °C for 18 h, volatile components were removed in vacuo, the product was isolated using preparative HPLC. 0.009 g (36%) of compound **CDK7_953** was yielded.

Example 19. Synthesis of compound **19.6**.



Synthesis of compound **19.2**

Aldehyde **19.1** (5.00 g, 34.9 mmol), 2,2-dimethoxyethane-1-amine (3.87 ml, 34.9 mmol), sodium triacetoxyborohydride (23.24 g, 105 mmol) were suspended in 150 ml of dichloromethane, stirred at room temperature for 24 h. After standard treatment, the product was isolated using silica gel column chromatography. 6.31 g (80%) of compound **19.2** were yielded.

Synthesis of compound **19.3**

Oxalyl chloride (0.451 ml, 5.26 mmol) and a drop of DMF were added to a suspension of 4-bromo-1H-pyrrole-2-carboxylic acid (**0.5** g, 2.63 mmol) in 5 ml of dichloromethane, stirred for 2 h at room temperature, volatile components were removed in vacuo, added to a solution of compound **19.2** (0.592 g, 2.63 mmol) and TEA in 5 ml of dichloromethane at 0 °C, stirred for 1 h. After standard treatment, the product was isolated using silica gel column chromatography. 1.00 g (95%) of compound **19.3** was yielded.

Synthesis of compound **19.4**

Compound **19.3** (2.68 g, 6.41 mmol) was dissolved in 15 ml of TFA, stirred for 6 days. Volatile components were removed in vacuo and the product was isolated using column chromatography. 1.78 g (83%) of compound **19.4** were yielded.


Synthesis of compound **19.5**

Compound **19.5** was produced analogously to compound **6.4** with a yield of 83%.

Synthesis of compound **19.6**

Compound **19.5** (0.440 g, 0.880 mmol) was dissolved in 2 ml of TFA in a sealed vessel, heated while stirring at 100 °C for 30 h. Volatile components were removed, the product was isolated using column chromatography. 0.22 g (71%) of compound **19.6** was yielded.

Example 20. Synthesis of compound **CDK7_1105**



Synthesis of compound **20.1**

Compound **19.6** (0.040 g, 0.110 mmol), cyclopropylboronic acid (0.029 g, 0.330 mmol), pyridine (0.018 ml, 0.220 mmol), triethylamine (0.081 ml, 0.550 mmol) in 4 ml of tetrahydrofuran was heated in a microwave reactor for 1 hour at 150 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.042 g (98%) of compound **20.1** was yielded.

Synthesis of compound **20.2**

Compound **20.1** (0.042 g, 0.100 mmol), (BPin)₂ (0.039 g, 0.150 mmol), KOAc (0.051 g, 0.500 mmol), Pd(dppf)Cl₂ (0.015 g, 0.020 mmol) was dissolved in 2 ml of 1,4-dioxane, heated in a sealed vessel for 1 h at 100° C. The reaction mixture was filtered off through Celite and the solvent was removed in vacuo. 0.04 g (91%) of product in the form of brown oil was yielded.

Synthesis of compound **20.3**

Compound **20.2** (0.04 g, 0.090 mmol), pyrimidine **1.4** (0.034 g, 0.090 mmol), Na₂CO₃ (0.019 g, 0.180 mmol), Pd(dppf)Cl₂ (0.010 g, 0.010 mmol) in a mixture of 2 ml of 1,4-dioxane and 0.5 ml of water were stirred in nitrogen atmosphere for 1 hour at 100 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.026 g (44%) of compound **20.3** was yielded.


Synthesis of compound **20.4**

Compound **20.3** (0.026 g, 0.04 mmol), NaOH (0.105 g, 2.60 mmol) in 5 ml of methanol and 0.5 ml of water were stirred at room temperature for 3 h, filtered off through Celite, the solvent was removed in vacuo. 0.02 g (95%) of compound **20.4** was yielded.

Synthesis of compound **CDK7_1105**

Compound **20.4** (0.020 g, 0.04 mmol) was dissolved in 2 ml of TFA, the reaction mass was stirred at 20 °C for 18 h, volatile components were removed in vacuo, re-evaporated with dichloromethane, volatile components were removed in vacuo, the product was isolated using preparative HPLC. 0.017 g (47%) of compound **CDK7_1105** was yielded.

Example 21. Synthesis of compound **CDK7_1062**.



Synthesis of compound **21.1**

2.23 ml (5.56 mmol) of 2.5M solution of *n*-butyllithium in hexane was added while stirring in nitrogen atmosphere at -78 °C to a solution of 2-chloropyrimidine (0.5 g, 4.28 mmol) in 10 ml of THF, stirred for 30 minutes, tributyltin hydride (1.42 ml, 5.14 mmol) was then added and stirred for another 30 minutes, temperature was adjusted to room temperature, stirred for 12 h at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.55 g (35%) of compound **21.1** was yielded.

Synthesis of compound **21.2**

Compound **21.1** (0.089 g, 0.240 mmol), 1-benzenesulfonyl-4-bromo-2-iodo-1*H*-pyrrol (0.1 g, 0.240 mmol), copper(I) iodide (0.014 g, 0.070 mmol), Pd(PPh₃)₄ (0.028 g, 0.020 mmol) in 20.1 ml of DMF were heated in a sealed vessel in an inert atmosphere for 4 h at 120 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.035 g (40%) of compound **21.2** was yielded.

Synthesis of compound **21.3**

Compound **21.2** (0.051 g, 0.140 mmol), (BPin)₂ (0.073 g, 0.280 mmol), KOAc (0.035 g, 0.350 mmol), Pd(dppf)Cl₂ (0.010 g, 0.010 mmol) was dissolved in 2 ml of 1,4-dioxane, heated in a sealed vessel for 10 h at 110 °C, the reaction mixture was filtered off through Celite and the solvent was removed in vacuo. 0.058 g (99%) of compound **21.3** in the form of brown oil was yielded.

Synthesis of compound **21.4**

Compound **21.3** (0.058 g, 0.140 mmol), pyrimidine **1.4** (0.053 g, 0.140 mmol), Na₂CO₃ (0.019 g, 0.018 mmol), Pd(dppf)Cl₂ (0.031 g, 0.04 mmol) in a mixture of 2 ml of 1,4-dioxane and 1 ml of water were stirred in nitrogen atmosphere for 2 h at 110 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.044 g (50%) of compound **21.4** was yielded.


Synthesis of compound **21.5**

Compound **21.5** was produced analogously to compound **6.7** with a yield of 99%.

Synthesis of compound **CDK7_1062**

Compound **CDK7_1062** was produced analogously to compound **6.7** with a yield of 37%.

Example 22. Synthesis of compound **CDK7_1058**




Synthesis of compound **22.2**

Compound **19.6** (0.12 g, 0.34 mmol), 2-bromothiazole (0.1 g, 0.6 mmol), CuI (0.06 g, 0.3 mmol), K₂CO₃ (0.09 g, 0.65 mmol) in 2 ml of DMF were heated while stirring in a sealed vessel at 130 °C for 2 h, cooled to room temperature, 3M solution of NaOH was added to pH 8, stirred at room temperature for 2 h, the solvent was removed in vacuo, the product was isolated using column chromatography. 0.1 g of a mixture in the form of oil with a content of compound **22.1** of 60% according to HPLC (0.06 g, 60%) was yielded. 0.056 g (0.32 mmol) of benzenesulfonyl chloride, 0.008 g (0.06 mmol) of DMAP, 0.064 g (0.64 mmol) of triethylamine and 5 ml of dichloromethane were added to this mixture. The reaction mixture was stirred at room temperature for 18 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.015 g (16%) of compound **22.2** was yielded.

Synthesis of compound **CDK7_1058**

0.015 g (0.21 mmol) of compound **22.2**, (BPin)₂ (0.012 g, 0.05 mmol), KOAc (0.01 g, 0.1 mmol), Pd(dppf)Cl₂ (0.01 g, 0.01 mmol) and 1 ml of 1,4-dioxane were mixed, the mixture was heated in a sealed vessel for 1 h at 100 °C. After cooling, the reaction mixture was filtered off through Celite, and the solvent was removed in vacuo. Pyrimidine **1.4** (0.011 g, 0.03 mmol), Na₂CO₃ (0.006 g, 0.06 mmol), Pd(dppf)Cl₂ (0.01 g, 0.01 mmol) in a mixture of 1 ml of 1,4-dioxane and 0.5 ml of water were added to the residue comprising **22.3**. The resulting mixture was stirred in nitrogen atmosphere for 2 h at 110 °C, the reaction mixture was filtered off through celite, and the solvent was removed in vacuo. The residue comprising compound **22.4** was dissolved in 1.5 ml of ethanol, NaOH (0.12 g, 3.00 mmol) and 1 ml of water were added. The reaction mass was stirred at 60 °C for 6 h, solvents were removed in vacuo, water was added to the residue until completely dissolved (1 ml), acidified with 10% solution of citric acid to pH 4, extraction was performed with ethyl acetate, volatile components were removed in vacuo. The residue was dissolved in 1 ml of TFA, the reaction mass was stirred at 20 °C for 18 h, the volatile components were removed in vacuo, the product was isolated using preparative HPLC. 1.3 mg (total yield: 7 %) of compound **CDK7_1058** were yielded.

Example 23. Synthesis of compounds **CDK7_1060**, **CDK7_1103**.



Synthesis of compound **23.2**

A mixture of 4-bromo-1H-pyrrole-2-carbonitrile (0.40 g, 2.29 mmol) (**23.1**), $\text{NH}_2\text{OH}\cdot\text{HCl}$ (0.804 g, 11.45 mmol) and TEA (1.6 ml, 11.45 mmol) in 20 ml of ethanol was boiled while stirring for 1 hour. After standard treatment and without additional purification, 0.465 g (99%) was produced and used in the next step.

Synthesis of compound **23.3**

EDC (0.137 g, 0.71 mmol), HOBt (0.073 g, 0.47 mmol) were suspended in 3 ml of 1,4-dioxane; 0.027 ml (0.47 mmol) of acetic acid was added and stirred at room temperature in nitrogen atmosphere for 1 h. Compound **23.2** (0.100 g, 0.47 mmol) was then added to 3 ml of 1,4-dioxane and stirred at room temperature for 12 h, the volatile components were removed in vacuo. The intermediate was produced by column chromatography, dissolved in 2 ml of 1,4-dioxane, heated in a microwave reactor at 130 °C for 2 h, volatile components were removed in vacuo. The product was isolated using column chromatography on silica gel. 0.025 g (23%) of compound **23.3** was yielded.

Synthesis of compound **23.4**

Compound **23.4** was produced analogously to compound **1.2** with a yield of 71%.

Synthesis of compound **23.5**

Compound **23.4** (0.044 g, 0.120 mmol), $(\text{BPin})_2$ (0.04 g, 0.16 mmol), KOAc (0.036 g, 0.36 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.009 g, 0.01 mmol) in 1 ml of 1,4-dioxane was heated while stirring in nitrogen atmosphere at 100 °C for 4 h. After standard treatment, the product was isolated using column chromatography on silica gel. 0.034 g (68%) of compound **23.5** was yielded.

Synthesis of compound **23.6**


Compound **23.6** was produced analogously to compound **20.3** with a yield of 41%.

Synthesis of compound **CDK7_1060**

Compound **23.6** (0.016 g, 0.02 mmol), NaOH (0.016 g, 0.40 mmol) in 0.5 ml of ethanol and 0.4 ml of water were stirred at room temperature for 4 h. After standard treatment, the residue comprising compound **23.7** was dissolved in 1 ml of dichloromethane, 0.058 ml of TFA was added, the reaction mass was stirred at 20 °C for 18 h, volatile components were removed in vacuo, the product was isolated by preparative HPLC. 0.008 g (66%) of compound **CDK7_1060** was yielded.

Compound **CDK7_1103** was produced analogously using cyclopropyl carboxylic acid with a total yield of 1%.

Example 24. Synthesis of compounds **CDK7_1061**, **CDK7_533**.



Synthesis of compound **24.2**

A mixture of pyridine of **24.1** (0.304 g, 0.82 mmol), 1-(benzenesulfonyl)-4-bromo-2-iodine-1 *H*-pyrrole (0.228 g, 0.55 mmol), CuI (0.032 g, 0.16 mmol), Pd(PPh₃)₄ (0.064 g, 0.06 mmol) in 5 ml of 1,4-dioxane was heated while stirring in nitrogen atmosphere at 100 °C for 4 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.054 g (32%) of compound **24.2** was yielded.

Synthesis of compound **24.3**

Compound **24.2** (0.064 g, 0.18 mmol), (BPin)₂ (0.093 g, 0.36 mmol), KOAc (0.055 g, 0.54 mmol), Pd(dppf)Cl₂ (0.04 g, 0.05 mmol) in 2 ml of 1,4-dioxane were heated while stirring in nitrogen atmosphere at 100 °C for 4 h. After standard treatment, the product was isolated using column chromatography on silica gel. 0.074 g (99%) of compound **24.3** was yielded.

Synthesis of compound **24.4**

Compound **3.3** (0.074 g, 0.18 mmol), pyrimidine of **1.4** (0.069 g, 0.18 mmol), Na₂CO₃ (0.058 g, 0.54 mmol), Pd(dppf)Cl₂ (0.04 g, 0.05 mmol) in a mixture of 2 ml of 1,4-dioxane and 1 ml of water were stirred in nitrogen atmosphere at 100 °C for 2 h. After standard treatment, the reaction product was isolated using column chromatography on silica gel. 0.04 g (35%) of compound **24.4** was yielded.

Synthesis of compound **24.5**


Compound **24.4** (0.04 g, 0.06 mmol), NaOH (0.024 g, 0.6 mmol) in a mixture of 3 ml of ethanol and 3 ml of water were stirred at room temperature for 1 h. After standard treatment, 0.029 g (99%) of compound **24.5** was yielded.

Synthesis of compound **CDK7_1061**

Compound **CDK7_1061** was produced analogously to compound **CDK7_1060** with a yield of 44%.

Compound **CDK7_533** was produced analogously using the corresponding boronic acid (conditions were analogous to the synthesis of compound **21.4**) with a total yield of 22%.


Example 25. Synthesis of compounds **CDK7_1000**




Synthesis of compound 25.2

NaH (0.045 g, 1.13 mmol) was added at 0 °C to a solution of 4-bromo-1H-pyrrol-2-carbonitrile **25.1** (0.130 g, 0.750 mmol) in 3 ml of tetrahydrofuran, stirred for 30 min, 0.217 ml (1.2 mmol) of SEMCl was added, stirred at room temperature for 1 h. After standard treatment, volatile components were removed in vacuo. The residue was dissolved in 2.5 ml of DMF, NaN₃ (0.058 g, 0.88 mmol), NH₄Cl (0.047 g, 0.88 mmol) were added, heated while stirring in a microwave synthesizer vial at 160 °C for 2 h. 0.069 ml (1.10 mmol) of methyl iodide was added to the reaction mixture, stirred at room temperature for 12 h. Volatile components were removed and the product was isolated using column chromatography. (BPin)₂ (0.168 g, 0.66 mmol), KOAc (0.134 g, 1.32 mmol), Pd(dppf)Cl₂ (0.025 g, 0.03 mmol) in 2 ml of 1,4-dioxane were added to the residue comprising compound **25.2** and heated while stirring in nitrogen atmosphere at 100 °C for 4 h. After standard treatment, the reaction mixture was filtered off through a silica gel layer and volatile components were removed. The residue comprising compound **25.3** was dissolved in a mixture of 3 ml of 1,4-dioxane and 1.5 ml of water, pyrimidine **1.4** (0.118 g, 0.310 mmol), Na₂CO₃ (0.100 g, 0.930 mmol), Pd(dppf)Cl₂ (0.069 g, 0.090 mmol) were added and stirred in nitrogen atmosphere for 1 hour at 100 °C. After standard treatment, the reaction mixture was filtered off through a silica gel layer and volatile components were removed. 1 ml of dichloromethane, **0.07 ml of TFA were added to the residue comprising compound 25.4**, the reaction mass was stirred at 20 °C for 18 h, volatile components were removed in vacuo. The residue was dissolved in 1 ml of ethanol; NaOH (0.036 g, 0.90 mmol) and 1 ml of water were added, stirred at 20 °C for 18 h. After standard treatment, the reaction product was isolated using preparative HPLC **CDK7_1000** was yielded.

Example 26. Synthesis of compound CDK7_1124





26.7

CDK7 1124

Synthesis of compound 26.2

Synthesis of compound 26.3

Synthesis of compound 26.4

Synthesis of compound 26.5


Synthesis of compound 26.6

Synthesis of compound 26.7

Synthesis of compound **CDK7_1124**

were removed in vacuo. The product was isolated using preparative HPLC. 0.008 g (36%) of compound **CDK7_1124** was yielded.

Example 27. Synthesis of compound CDK7_1021



Synthesis of compound 27.1

Compound **18.3** (0.250 g, 0.830 mmol) was dissolved in 15 ml of THF, NaH (0.037 g, 0.910 mmol) was added at 0 °C to the resulting solution, and the mixture was stirred for 10 minutes, benzenesulfonyl chloride (0.139 ml, 1.08 mmol) was added and stirred for 1 hour at room temperature. After standard treatment, the product was isolated using silica gel column chromatography. 0.131 g (40%) of compound **27.1** was yielded.

Synthesis of compound 27.2

Compound **27.1** (0.131 g, 0.310 mmol), (BPin)₂ (0.157 g, 0.620 mmol), KOAc (0.094 g, 0.930 mmol), Pd(dppf)Cl₂ (0.023 g, 0.030 mmol) in 5 ml of 1,4-dioxane were stirred in nitrogen atmosphere for 8 h at 100°C. After standard treatment, the reaction product was isolated using column chromatography on silica gel. 0.06 g (43%) of compound **27.2** was yielded.


Synthesis of compound 27.3

Compound **27.3** was produced analogously to compound **4.3** with a yield of 23%.

Synthesis of compound CDK7_1021

Compound **27.3** (0.021 g, 0.03 mmol), NaOH (0.012 g, 0.03 mmol) in 0.5 ml of ethanol and 0.5 ml of H₂O were stirred at room temperature for 2 h. After standard treatment, the residue containing compound **27.4** was dissolved in 1 ml of dichloromethane, 0.093 ml of TFA was added and stirred at room temperature for 20 h. After standard treatment, the product was isolated using preparative HPLC. 0.006 g (35%) of compound **CDK7_1021** was yielded.

Example 28. Synthesis of compound CDK7_1121



Synthesis of compound **28.1**

Compound **19.6** (0.071 g, 0.190 mmol), oxane-4-yl methanesulfonate (0.070 g, 0.380 mmol), Cs₂CO₃ (0.125 g, 0.380 mmol) in a mixture of 1 ml of DMF and 1 ml of dimethoxyethane were heated while stirring at 90°C for 40 h. After standard treatment, the product was isolated using column chromatography on silica gel. 0.058 g (70%) of compound **28.1** was yielded.

Synthesis of compound **28.2**

Compound **28.1** (0.058 g, 0.120 mmol), (BPin)₂ (0.050 g, 0.190 mmol), KOAc (0.039 g, 0.380 mmol), Pd(dppf)Cl₂ (0.009 g, 0.010 mmol) was dissolved in 1 ml of 1,4-dioxane, heated in a sealed vessel for 4 h at 110°C. After standard treatment, the product was isolated using column chromatography. 0.049 g (85%) of compound **28.2** was yielded.


Synthesis of compound **28.3**

Compound **28.2** (0.049 g, 0.100 mmol), pyrimidine **1.4** (0.046 g, 0.120 mmol), Na₂CO₃ (0.040 g, 0.370 mmol), Pd(dppf)Cl₂ (0.030 g, 0.040 mmol) in a mixture of 1 ml of 1,4-dioxane and 0.5 ml of water were stirred in nitrogen atmosphere for 1 hour at 110 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.040 g (57%) of compound **28.3** was yielded.

Synthesis of compound **CDK7_1121**

Compound **28.3** (0.040 g, 0.060 mmol), NaOH (0.121 g, 3.00 mmol) in a mixture of 1 ml of ethanol and 1 ml of H₂O was stirred at room temperature for 2 h. After standard treatment, the residue containing compound **28.4** was dissolved in 3 ml of dichloromethane, 0.696 ml of TFA was added and stirred at room temperature for 18 h. After standard treatment, the product was isolated using preparative HPLC. 0.003 g (11%) of compound **CDK7_1121** was yielded.

Example 29. Synthesis of compound **29.5**.



Synthesis of compound 29.1

Oxalyl chloride (5.94 ml, 67.9 mmol) and a drop of DMF were added to acid **2.1** (4.30 g, 22.6 mmol) in 60 ml of dichloromethane and stirred at room temperature for 1 hour. Volatile components were then removed, the residue was dissolved in 30 ml of dichloromethane and added dropwise to a solution of methyl 3-aminopropanoate hydrochloride (4.11 g, 29.4 mmol) and TEA (16.6 ml, 113 mmol) in 30 ml of dichloromethane at 0 °C. The mixture was stirred at room temperature for 1 h. After standard treatment, the product was isolated using column chromatography on silica gel. 5.70 g (92%) of compound **29.1** was yielded.

Synthesis of compound 29.2

Compound **29.1** (3.90 g, 12.8 mmol) was dissolved in 40 ml of ethanol, NaOH (1.08 g, 26.8 mmol) and 20 ml of water were added. The reaction mass was stirred at room temperature for 18 h. After standard treatment, 2.8 g (84%) of product **29.2** was yielded.

Synthesis of compound 29.3

A mixture of compound **29.2** (4.07 g, 14.8 mmol), PPA (38.2 g, 148 mmol), P₂O₅ (4.29 g, 14.8 mmol) was stirred at 110 °C for 2 h. After standard treatment, 3.51 g (97%) of compound **29.3** were yielded.


Synthesis of compound 29.4

Compound **29.3** (2.04 g, 6.72 mmol), triethylsilane (3.16 g, 26.9 mmol) in 31 ml of TFA were stirred at room temperature for 20 h. After standard treatment, 1.22 g (79%) of compound **29.4** was obtained.

Synthesis of compound 29.5

Compound **29.5** was produced analogously to compound **2.3** with a yield of 50%.

Example 30. Synthesis of compound **CDK7_955**.



Synthesis of compound **30.1**

Compound **29.5** (0.346 g, 0.920 mmol), 4-Iodo-1-methyl-1H-pyrazole (1.17 g, 5.52 mmol), CuI (0.088 g, 0.460 mmol), (1*R*, 2*R*)-cyclohexane-1,2-diamine (0.107 g, 0.920 mmol) in 14 ml of 1,4-dioxane were heated while stirring at 120 °C for 7 h. After standard treatment, the product was isolated using column chromatography. 0.247 g (60%) of compound **30.1** was yielded.

Synthesis of compound **30.2**

Compound **30.1** (0.130 g, 0.290 mmol), (BPin)₂ (0.221 g, 0.870 mmol), KOAc (0.132 g, 1.30 mmol), Pd(dppf)Cl₂ (0.065 g, 0.09 mmol) was dissolved in 9 ml of 1,4-dioxane, heated in a sealed vessel for 12 h at 110° C. After standard treatment, the product was isolated by column chromatography. 0.141 g (98%) of compound **34.2** was yielded.

Synthesis of compound **30.3**

Compound **30.2** (0.225 g, 0.12 mmol), pyrimidine **1.4** (0.059 g, 0.160 mmol), Na₂CO₃ (0.039 g, 0.360 mmol), Pd(dppf)Cl₂ (0.009 g, 0.01 mmol) in a mixture of 3 ml of 1,4-dioxane and 1 ml of water were stirred in nitrogen atmosphere for 10 h at 100 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.060 g (70%) of compound **30.3** was yielded.

Synthesis of compound **30.4**

Compound **30.3** (0.026 g, 0.04 mmol), NaOH (0.112 g, 2.80 mmol) in 5 ml of ethanol and 5 ml of water were stirred at room temperature for 1 hour, filtered off through celite, the solvent was removed in vacuo. 0.08 g (99%) of compound **30.4** was yielded.

Synthesis of compound **CDK7_955**

Compound **30.4** (0.080 g, 0.14 mmol) was dissolved in 2 ml of dichloromethane, 0.217 ml of TFA was added, the reaction mass was stirred at 20 °C for 20 h. After standard treatment, the product was isolated using preparative HPLC. 0.043 g (65%) of compound **CDK7_955** was yielded.

Synthesis of compound **30.5**


Compound **29.5** (0.201 g, 0.53 mmol), cyclopropylboronic acid (0.141 g, 1.59 mmol), copper(II) acetate (0.147 g, 0.079 mmol), TEA (0.339 g, 3.18 mmol), DMAP (0.196 g, 1.59 mmol) in 5 ml

of toluene were stirred at 100 °C in for 12 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.098 g (44%) of product **30.5** was yielded.

Synthesis of compound **30.6**

Compound **30.5** (1.00 g, 2.32 mmol), (BPin)₂ (1.80 g, 6.96 mmol), KOAc (0.704 g, 6.96 mmol), Pd(dppf)Cl₂ (0.139 g, 0.190 mmol) was dissolved in 30 ml of 1,4-dioxane, heated in a sealed vessel for 24 h at 100° C. After standard treatment, the product was isolated using column chromatography. 0.700 g (66%) of compound **30.6** was yielded.

Example 31. Synthesis of compound **CDK7_1122**, **CDK7_1225**.



Synthesis of compound **31.2**


Compound **30.2** (0.216 g, 0.440 mmol), pyrimidine **31.1** (0.125 g, 0.390 mmol), Na₂CO₃ (0.124 g, 1.16 mmol), Pd(dppf)Cl₂ (0.058 g, 0.08 mmol) in a mixture of 6 ml of 1,4-dioxane and 2 ml of water were stirred in nitrogen atmosphere for 4 h at 120 °C. After standard treatment, the product was isolated using column chromatography on silica gel. 0.114 g (46%) of compound **31.2** was yielded.

Synthesis of compound **CDK7_1122**

Compound **31.2** (0.171 g, 0.270 mmol), NaOH (0.109 g, 2.70 mmol) in 2 ml of ethanol and 1 ml of water were stirred at room temperature for 4 h. After standard treatment, the product was isolated using preparative HPLC. 0.028 g (21%) of compound **CDK7_1122** was yielded.

Compound **CDK7_1225** was produced analogously with a total yield of 69%.

Example 32. Synthesis of compound **CDK7_1123**.



Synthesis of compound **32.2**

Compound **32.2** was produced analogously to compound **4.2** with a yield of 79%.

Synthesis of compound **32.3**

Amide **32.2** (7.34 g, 20.5 mmol) and 27 ml of methanesulfonic acid (410 mmol) were stirred for 16 h at 20 °C. After standard treatment, 4.98 g (83%) of compound **32.3** were yielded.

Synthesis of compound **32.4**

DIPEA (6.68 g, 51.2 mmol), DMAP (1.05 g, 8.53 mmol) and benzenesulfonyl chloride (4.87 g, 27.3 mmol) were added to a solution of pyrrolopyridone **32.3** (5.00 g, 17.1 mmol) in 100 ml of dichloromethane, the reaction mass was stirred at room temperature for 18 h. After standard treatment, the product was isolated using silica gel column chromatography. 3.00 g (41%) of compound **32.4** was yielded.

Synthesis of compound **32.5**

Compound **32.4** (1.50 g, 3.46 mmol), (BPin)₂ (1.76 g, 6.92 mmol), KOAc (1.05 g, 10.4 mmol), Pd(dppf)Cl₂ (0.258 g, 0.350 mmol) in 20 ml of 1,4-dioxane was heated while stirring in nitrogen atmosphere for 3 h at 100°C. After standard treatment, the product was isolated using column chromatography on silica gel. 1.60 g (96%) of compound **32.5** was yielded.


Synthesis of compound **32.6**

Compound **32.5** (0.117 g, 0.230 mmol), pyrimidine **31.1** (0.110 g, 0.210 mmol), Na₂CO₃ (0.067 g, 0.630 mmol), Pd(dppf)Cl₂ (0.031 g, 0.040 mmol) in a mixture of 3 ml of 1,4-dioxane and 1 ml of water were stirred in nitrogen atmosphere for 18 h at 100 °C. After standard treatment, 0.130 g (98%) of compound **32.6** was yielded.

Synthesis of compound **CDK7_1123**

Compound **32.6** (0.600 g, 0.960 mmol), NaOH (0.384 g, 9.60 mmol) in 18 ml of ethanol and 3 ml of water were stirred at 45 °C for 3 h. After standard treatment, the product was isolated using preparative HPLC. 0.106 g (23%) of compound **CDK7_1123** was yielded.

Example 33. Synthesis of compound **CDK7_1176**, **CDK7_1172**, **CDK7_1177**, **CDK7_1188**.



Synthesis of compound **33.1**

Compound **33.1** was produced analogously to compound **30.1** with a yield of 83%.

Synthesis of compound **33.2**

Compound **33.2** was produced analogously to compound **30.2** with a yield of 99%.

Synthesis of compound **33.3**

Compound **33.3** was produced analogously to compound **30.3** with a yield of 53%.


Synthesis of compound **33.4**

Compound **33.4** was produced analogously to compound **30.4** with a yield of 99%.

Synthesis of compound **CDK7_1176**


Compound **CDK7_1176** was produced analogously to compound **CDK7_955** with a yield of 70%.

The following compounds were produced in an analogous manner:

No.	Compound	Initial compound	Stage, yield, %				
			1	2	3	4	5
1	CDK7_1172		37	86			13
2	CDK7_1177		38	4			
3	CDK7_1188		18*	84	27		

*Conditions and methods of the stage are described in the synthesis of compound **30.5**

Example 34. Synthesis of compound CDK7_1187.



Synthesis of compound 34.1

Method A:

Compound **34.1** was produced analogously to compound **2630.51** with a yield of 79%.

Method B:

Compound **34.1** was produced analogously to **44.3** from compound **43.1** and MeI with a total yield of 70.0 %


Synthesis of compound 34.2

Compound **34.2** was produced analogously to compound **30.2** with a yield of 99%.

Synthesis of compound CDK7_1187

Compound **34.2** (0.069 g, 0.120 mmol), pyrimidine **31.1** (0.062 g, 0.120 mmol), Na₂CO₃ (0.039 g, 0.360 mmol), Pd(dppf)Cl₂ (0.013 g, 0.020 mmol) in a mixture of 2 ml of 1,4-dioxane and 1 ml of water were stirred in nitrogen atmosphere for 6 h at 100 °C. After standard treatment, the product was isolated using preparative HPLC. 0.05 g (10%) of compound **CDK7_1187** was yielded.

Example 35. Synthesis of compounds CDK7_999, CDK7_1102.



Synthesis of compound 35.1

Compound **35.1** was produced analogously to compound **4.2** with a yield of 96%.

Synthesis of compound **35.2**

Compound **35.1** was dissolved in 10 ml of acetonitrile; NaN₃ (0.136 g, 2.07 mmol) was added. A solution of trifluoromethanesulfonic anhydride (0.570 g, 1.88 mmol) in 5 ml of acetonitrile was then added and stirred at room temperature for 18 h. After standard treatment, the product was isolated using column chromatography. 0.162 g (76%) of compound **35.2** was yielded.

Synthesis of compound **35.3**

Compound **35.3** was produced analogously to compound **26.3** with a yield of 49%.

Synthesis of compound **35.4**

Compound **35.4** was produced analogously to compound **20.2** with a yield of 50%.

Synthesis of compound **35.5**

Compound **35.5** was produced analogously to compound **20.3** with a yield of 84%.

Synthesis of compound **35.6**


Compound **35.6** was produced analogously to compound **24.5** with a yield of 74%.

Synthesis of compound **CDK7_999**

Compound **CDK7_999** was produced analogously to compound **CDK7_951** with a yield of 58%.

Compound **CDK7_1102** was produced analogously using cyclopropylamine with a total yield of 1%.


Example 36. Synthesis of compound **CDK7_814**.



Synthesis of compound **CDK7_814**

Compound **36.1** (0.152 g, 0.210 mmol), pyrimidine **1.4** (0.080 g, 0.210 mmol), Na₂CO₃ (0.067 g, 0.630 mmol), Pd(dppf)Cl₂ (0.047 g, 0.060 mmol) in a mixture of 3 ml of 1,4-dioxane and 1.5 ml of water were stirred in nitrogen atmosphere for 1 hour at 100 °C. Volatile components were removed, and the product was isolated using column chromatography. 3 ml of TFA was added to the residue containing compound **36.2** and stirred at room temperature for 12 h. After standard treatment, the product was isolated using preparative chromatography. 0.023 g (30%) of compound **CDK7_814** was yielded.

Example 37. Synthesis of compound **CDK7_843**.



Synthesis of compound **37.2**


1M solution of LiHMDS in tetrahydrofuran (0.405 ml, 0.410 mmol) was added at -10 °C to a solution of compound **37.1** (0.082 g, 0.340 mmol) in 4 ml of THF and stirred for 1 hour; methyl iodide (0.161 g, 1.12 mmol) was added and stirred at 0 °C for 10 minutes, then stirred at room

temperature for 2 h. After standard treatment, the product was isolated using silica gel column chromatography. 0.067 g (79%) of compound **37.2** was yielded.

Synthesis of compound **CDK7_843**

A mixture of compound **37.2** (0.085 g, 0.330 mmol), (BPin)₂ (0.128 g, 0.490 mmol), [Ir(OMe)cod]₂ (0.016 g, 0.020 mmol), 4,4'-di-tert-butyl-2,2'-bipyridine (0.013 g, 0.050 mmol) was mixed in 2 ml of THF and stirred at room temperature for 10 minutes, a solution of compound **37.2** (0.085 g, 0.330 mmol in 1 ml of THF) was then added and heated in a sealed vessel at 70 °C for 2 h, then at room temperature for 12 h. After cooling, the mixture was filtered off through a silica gel layer, the solvent was removed in vacuo. **Pyrimidine 1.4** (0.122 g, 0.320 mmol), Na₂CO₃ (0.103 g, 0.960 mmol), Pd(dppf)Cl₂ (0.072 g, 0.100 mmol) in a mixture of 3 ml of 1,4-dioxane and 1.5 ml of water were added to the residue containing compound **37.3**. The resulting mixture was stirred in nitrogen atmosphere for 4 h at 100 °C, the reaction mixture was filtered off through a silica gel layer, and the solvent was removed in vacuo. The product was isolated using column chromatography on silica gel. The residue containing compound **37.4** was then dissolved in 3 ml of TFA, stirred at room temperature for 12 h. After standard treatment, the product was isolated using preparative HPLC. 0.006 g (5%) of compound **CDK7_843** was yielded.

Example 38. Synthesis of compound **CDK7_1019**.




Synthesis of compound **38.1**

Compound **16.5** (0.220 g, 0.350 mmol) was dissolved in 1 ml of TFA, stirred at room temperature for 32 h, the solvent was removed in vacuo. The dry residue was then dissolved in 15 ml of dichloromethane, methyl iodide (0.024 ml, 0.390 mmol) and DIPEA (0.311 ml, 1.75 mmol) were added, stirred at room temperature for 12 h. After standard treatment, the product was isolated using column chromatography. 0.160 g (85%) of compound **38.1** was yielded.

Synthesis of compound **CDK7_1019**

Compound **38.1** (0.160 g, 0.280 mmol), NaOH (0.226 g, 5.60 mmol) in 6 ml of ethanol and 6 ml of water were stirred at room temperature for 1 h. The solvent was removed in vacuo, the product was isolated using preparative HPLC. 0.035 g (32%) of compound **CDK7_1019** was yielded.


Example 39. Synthesis of compound **CDK7_1020**.



Synthesis of compound **CDK7_1020**

Compound **1.5** (0.050 g, 0.080 mmol), NaOH (0.065 g, 1.60 mmol) in 2 ml of ethanol and 2 ml of water were stirred at room temperature for 1 h. After standard treatment, the residue was dissolved in 1.5 ml of TFA and stirred at room temperature for 16 h. After standard treatment, the product was isolated using preparative HPLC. 0.015 g (48%) of compound **CDK7_1020** was yielded.


Example 40. Synthesis of compound 40.2.



Synthesis of compound 40.2

Compound 40.2 was produced analogously to compound 15.3 with a yield of 48%.


Example 41. Synthesis of compound CDK7_1199.



Synthesis of compound CDK7_1199

Compound 32.5 (0.053 g, 0.100 mmol), pyrimidine 40.2 (0.028 g, 0.090 mmol), Na_2CO_3 (0.106 g, 0.270 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.020 g, 0.027 mmol) in a mixture of 1 ml of 1,4-dioxane and 0.5 ml of water were stirred in nitrogen atmosphere at 100 °C for 2 h. After standard treatment, intermediate 41.1 was used in the next stage without additional purification. NaOH (0.032 g, 0.800 mmol), 4 ml of ethanol and 4 ml of water were added to the residue containing compound 41.1 and stirred at room temperature for 1 h. After standard treatment, the product was isolated using preparative HPLC. 0.001 g (3%) of compound CDK7_1199 was yielded.

Example 42. Synthesis of compound CDK7 1254.



Synthesis of compound 42.2

Compound 42.2 was produced analogously to compound 29.1 with a yield of 97%.

Synthesis of compound 42.3

Compound 42.3 was produced analogously to compound 29.2 with a yield of 62%.

Synthesis of compound **42.4**

Compound **42.4** was produced analogously to compound **29.3** with a yield of 72%.

Synthesis of compound **42.6**

NaBH₄ (0.062 g, 1.64 mmol) was added while cooling on an ice bath to a solution of compound **42.4** (0.220 g, 0.410 mmol) in 7 ml of ethanol, the reaction mass was stirred at 20 °C for 24 h. After standard processing, the product was isolated using silica gel column chromatography. The residue comprising compound **42.5** was dissolved in 4 ml of dichloromethane, and 0.57 ml of 4N solution of HCl in 1,4-dioxane was added and stirred at 20 °C for 20 h. After standard treatment, the product was isolated using column chromatography. 0.077 g (79%) of compound **42.6** was yielded.

Synthesis of compound **42.7**

Compound **42.7** was produced analogously to compound **2.3** with a yield of 87%.

Synthesis of compound **42.8**

Compound **42.8** was produced analogously to compound **27.2** with a yield of 83%.

Synthesis of compound **42.9**

Compound **42.9** was produced analogously to compound **1.5** with a yield of 41%.


Synthesis of compound **42.10**

Pd/C 5% (0.170 g, 0.080 mmol) was added to a solution of compound **42.9** (0.045 g, 0.080 mmol) in 5 ml of methanol. Hydrogenation was performed at atmospheric pressure for 20 h, the reaction mass was filtered off through a Celite layer, the solvent was removed in vacuo. 0.023 g (55%) of compound **42.10** was yielded.

Synthesis of compound **CDK7_1254**

Compound **CDK7_1254** was produced analogously to compound **CDK7_836** with a yield of 29%.


Example 43. Synthesis of compound **43.1**.



Synthesis of compound **43.1**

NaH (0.227 g, 5.68 mmol) at 0 °C was added to a solution of compound **29.4** (1.00 g, 4.37 mmol) in 50 ml of tetrahydrofuran, stirred for 30 min, 0.789 ml (4.37 mmol) of SEMCl was added, stirred at room temperature for 3 h. After standard treatment, the reaction product was isolated by column chromatography. 0.927 g (59%) of compound **43.1** was yielded.

Example 44. Synthesis of compounds **CDK7_1154**, **CDK7_1155**, **CDK7_1226**, **CDK7_1227**, **CDK7_1229**, **CDK7_1259**, **CDK7_1268**.



Synthesis of compound **44.1**

NaH (0.948 g, 23.7 mmol) at 0 °C was added to a solution of compound **43.1** (0.300 g, 0.790 mmol) in 15 ml of DMF, stirred for 10 minutes, bromocyclopentane (3.72 g, 23.7 mmol) was added, stirred at room temperature for 28 h. After standard treatment, the reaction product was isolated by column chromatography. 0.154 g (46%) of compound **44.1** was yielded.

Synthesis of compound **44.3**

Compound **44.1** was dissolved in 5 ml of TFA and stirred at room temperature for 1 h. Volatile components were then removed in vacuo and the dry residue was dissolved in 5 ml of ethanol. 3 ml of 3M solution of NaOH was added, stirred at room temperature for 1 hour, volatile components were removed. After standard treatment, the product was purified by column chromatography. The residue containing compound **44.2** was then dissolved in 5 ml of DMF; NaH (0.029 g, 0.720 mmol) was added at 0 °C, stirred for 30 minutes; benzenesulfonyl chloride (0.134 g, 0.720 mmol) was then added and stirred at room temperature for 16 h. After standard treatment, the product was isolated using column chromatography. 0.054 g (34%) of compound **44.3** was yielded.

Synthesis of compound **44.4**

Compound **44.4** was produced analogously to compound **30.2**.

Synthesis of compound **44.5**

Compound **44.5** was produced analogously to compound **32.6** using pyrimidine **31.1** with a yield of 54%.

Synthesis of compound **CDK7_1154**


Compound **CDK7_1154** was produced analogously to compound **CDK7_1122** with a yield of 12%.

The following compounds were produced in an analogous manner:

No.	Compound	Initial compound	Stage, yield, %					
			1	2	3	4	5	6
1	CDK7_1155		42	51	67	80	32	56
2	CDK7_1226		25	46			4	
3	CDK7_1227		91	38	72	84		14
4	CDK7_1229		74	92	96	60		36

5	CDK7_1259	<chem>CCCCI</chem>	63			15
6	CDK7_1268	<chem>BrCCOC</chem>	68	43	78	15

Example 45. Synthesis of compound **CDK7_1201**, **CDK7_1277**.



Synthesis of compound **45.1**

Compound **45.1** was produced analogously to compound **44.1** using 2-iodopropane with a yield of 74%.

Synthesis of compound **45.2**

Compound **45.2** was produced analogously to compound **44.2** with a yield of 92%.

Synthesis of compound **45.3**

Compound **45.3** was produced analogously to compound **44.3** with a yield of 96%.

Synthesis of compound **45.4**

Compound **45.4** was produced analogously to compound **44.4** with a yield of 60%.

Synthesis of compound **45.6**

Compound **45.6** was produced analogously to compound **18.8** with a yield of 85%.

Synthesis of compound **45.7**

Compound **45.6** (1.55 g, 2.65 mmol) was dissolved in 15 ml of 4 N solution of HCl in 1,4-dioxane, the reaction mass was stirred at 60 °C for 30 min. After standard treatment, the product was filtered off and volatile components were removed in vacuo. 1.15 g (88%) of compound **45.7** were yielded.

Synthesis of compound **45.8**


Compound **45.7** (0.470 g, 0.900 mmol) was dissolved in 6 ml of POCl₃ and stirred at 90 °C for 30 min. After standard treatment, the product was isolated using column chromatography. 0.367 g (79%) of compound **45.8** was yielded.

Synthesis of compound **CDK7_1277**

A suspension of compound **45.8** (0.534 g, 1.04 mmol), amine **40.1** (0.171 g, 1.35 mmol) and TEA (0.916 ml, 6.24 mmol) in 30 ml of absolute ethanol was stirred at room temperature for 60 h. 2 ml of 3 M solution of NaOH was added to the resulting mixture and mixed at room temperature for 2 h. After standard treatment, the product was isolated using preparative HPLC. 0.170 g (35%) of product **CDK7_1277** was yielded.

Compound **CDK7_1201** was produced analogously with a total yield of 3%.

Example 46. Synthesis of compound **CDK7_1276**.



Synthesis of compound **46.1**

Compound **46.1** was produced analogously to compound **18.8** using pyrimidine **45.5** with a yield of 75%.

Synthesis of compound **46.2**

Compound **46.2** was produced analogously to compound **45.7** using pyrimidine **45.5** with a yield of 75%.

Synthesis of compound **CDK7_1276**

Compound **46.2** (0.075 g, 0.140 mmol) was dissolved in 0.5 ml of POCl₃ and stirred at 90 °C for 30 min. After standard treatment, the product was isolated using column chromatography. The residue containing compound **46.3**, amine **40.1** (0.083 g, 1.70 mmol) and TEA (0.117 ml, 0.840 mmol) in 8 ml of absolute ethanol was mixed at room temperature for 72 h. 2 ml of 3 M solution of NaOH was added to the resulting mixture and stirred at room temperature for 4 h. After standard treatment, the product was isolated using preparative HPLC. 0.012 g of product **CDK7_1276** (15%) was yielded.

Example 47. Analysis of resulting compounds.

The purity and structure of the resulting compounds was confirmed by chromatography/mass spectrometry LC/MS-ESI and ¹H NMR spectroscopy (**Tables 3 and 4**).

Equipment data:

Table 1. Chromatography/mass spectrometry

Name	Manufacturer, country
Agilent Triple Quad LC/MS system:	Agilent, USA
Agilent 1200 autosampler	
Agilent 1200 thermostatted column	
Agilent 1200 degasser	
Agilent 1200 autosampler thermostat	
Agilent 6410 QQQ MS detector	
Agilent 1200 UV Detector	
Agilent 1200 pump	

Table 2. NMR spectrometer

Name	Manufacturer, country	Model, main characteristics

NMR spectrometer	Germany	AVANCE III, 400 MHz
------------------	---------	---------------------

Table 3. Analytical data for exemplary compounds

Code	¹ H NMR (400 MHz, DMSO-d ₆), δ
CDK7_533	¹ H NMR (400 MHz, DMSO) δ 11.89 (s, 1H), 8.62 – 8.30 (m, 1H), 7.74 – 7.32 (m, 6H), 7.31 – 7.14 (m, 1H), 7.03 (s, 1H), 4.00 – 3.71 (m, 1H), 3.14 – 2.94 (m, 1H), 2.87 – 2.70 (m, 1H), 2.47 – 2.24 (m, 2H), 1.99 – 1.81 (m, 1H), 1.70 – 1.58 (m, 1H), 1.57 – 1.36 (m, 1H).
CDK7_535	¹ H NMR (400 MHz, DMSO) δ 12.86 – 12.57 (m, 1H), 8.96 – 8.74 (m, 2H), 8.71 – 8.49 (m, 1H), 8.06 – 7.94 (m, 1H), 7.94 – 7.79 (m, 2H), 7.71 – 7.46 (m, 2H), 7.62 – 7.54 (m, 2H), 7.33 – 7.21 (m, 1H), 4.32 – 4.11 (m, 1H), 3.44 – 3.12 (m, 3H), 2.91 – 2.73 (m, 2H), 2.05 – 1.83 (m, 2H), 1.78 – 1.50 (m, 2H).
CDK7_536	¹ H NMR (400 MHz, DMSO) δ 12.16 – 11.80 (m, 1H), 8.53 and 8.47 (2s, 1H, rotamers), 7.68 – 7.55 (m, 1H), 7.48 and 7.42 (2s, 1H, rotamers), 7.05 (s, 1H), 3.91 – 3.78 (m, 1H), 3.22 – 2.94 (m, 6H), 2.85 – 2.71 (m, 1H), 2.47 – 2.36 (m, 2H), 1.89 (s, 2H), 1.69 – 1.55 (m, 1H), 1.53 – 1.41 (m, 2H).
CDK7_537	¹ H NMR (400 MHz, DMSO) δ 12.28 – 11.80 (m, 1H), 8.64 – 8.43 (m, 1H), 8.34 (s, 1H), 8.15 – 7.58 (m, 2H), 7.54 – 7.24 (m, 2H), 7.20 – 6.94 (m, 1H), 4.10 – 3.95 (m, 1H), 3.72 – 3.52 (m, 1H), 3.30 – 2.86 (m, 2H), 2.68 – 2.58 (m, 1H), 1.97 – 1.84 (m, 1H), 1.79 – 1.45 (m, 3H).
CDK7_538	¹ H NMR (400 MHz, DMSO) δ 12.52 – 12.09 (m, 1H), 10.47 and 10.07 (2s, 1H, rotamers), 8.61 – 8.45 (m, 1H), 8.33 (s, 1H), 7.83 – 7.56 (m, 5H), 7.38 – 7.31 (m, 2H), 7.11 – 7.03 (m, 1H), 4.20 – 4.05 (m, 1H), 3.34 – 3.07 (m, 1H), 3.09 – 2.85 (m, 1H), 2.71 – 2.58 (m, 2H), 2.00 – 1.86 (m, 1H), 1.80 – 1.45 (m, 3H).
CDK7_539	¹ H NMR (400 MHz, DMSO) δ 12.37 – 12.12 (m, 1H), 11.24 – 10.88 (m, 1H), 9.65 – 8.85 (m, 2H), 8.66 – 8.50 (m, 1H), 8.07 – 7.89 (m, 1H), 7.65 – 7.58 and 7.51 – 7.44 (2m, 1H, rotamers), 7.15 – 7.01 (m, 1H), 4.55 – 4.40 (m, 2H), 4.40 – 4.17 (m, 1H), 3.54 – 3.31 (m, 5H), 3.25 – 3.04 (m, 3H), 2.98 – 2.69 (m, 5H), 2.04 – 1.51 (m, 4H).
CDK7_548	¹ H NMR (400 MHz, DMSO) δ 10.49 – 9.94 (m, 1H), 9.33 – 8.84 (m, 2H), 8.80 – 8.44 (m, 1H), 8.43 – 8.24 (m, 1H), 8.00 – 7.73 (m, 1H), 7.22 – 6.71 (m, 3H), 3.96 – 3.81 (m, 1H), 3.36 – 2.73 (m, 4H), 2.01 – 1.43 (m, 4H).
CDK7_735	¹ H NMR (400 MHz, DMSO) δ 11.25 and 10.73 (2s, 1H, rotamers), 9.49 – 8.94 (m, 2H), 8.58 – 8.27 (m, 2H), 7.86 – 7.48 (m, 2H), 7.34 – 7.13 (m, 1H), 4.27 – 4.01 (m, 1H), 3.74 – 3.43 (m, 1H), 3.37 – 3.22 (m, 1H), 3.19 – 3.07 (m, 1H), 3.00 – 2.73 (m, 2H), 2.05 – 1.91 (m, 1H), 1.91 – 1.82 (m, 1H), 1.82 – 1.62 (m, 7H), 1.61 – 1.43 (m, 1H).
CDK7_754	¹ H NMR (400 MHz, DMSO) δ 9.36 – 8.71 (m, 2H), 8.63 – 8.46 (m, 1H), 8.35 – 8.23 and 8.01 – 7.82 (2m, 1H, rotamers), 7.54 – 7.39 (m, 2H), 7.35 – 7.16 (m, 3H), 4.28 – 4.13 (m, 1H), 3.33 – 2.93 (m, 2H), 2.98 – 2.61 (m, 2H), 1.97 – 1.28 (m, 4H).
CDK7_766	¹ H NMR (400 MHz, DMSO) δ 8.56 and 8.47 (2s, 1H, rotamers), 7.88 – 7.80 (m, 1H), 7.33 – 7.18 (m, 5H), 4.03 – 3.95 (m, 2H), 3.92 – 3.79 (m, 1H), 3.06 – 2.94 (m, 1H), 2.86 – 2.77 (m, 1H), 2.46 – 2.36 (m, 2H), 1.91 – 1.76 (m, 1H), 1.68 – 1.55 (m, 1H), 1.50 – 1.32 (m, 2H).
CDK7_809	¹ H NMR (400 MHz, DMSO) δ 9.30 – 9.09 (m, 1H), 8.93 – 8.81 (m, 1H), 8.56 and 8.44 (2d, <i>J</i> = 7.7 Hz, 1H, rotamers), 7.82 – 7.75 (m, 1H), 7.69 – 7.61 (m, 1H), 7.56 – 7.45 (m, 1H), 7.32 – 7.15 (m, 2H), 6.77 (s, 1H), 4.35 – 4.06 (m, 1H), 3.42 – 3.30 (m, 1H), 3.22 – 3.07 (m, 1H), 2.92 – 2.76 (m, 2H), 2.05 – 1.97 (m, 1H), 1.92 – 1.86 (m, 1H), 1.79 – 1.54 (m, 2H).
CDK7_810	¹ H NMR (400 MHz, DMSO) δ 9.04 – 8.93 (2s, 1H, rotamers), 8.70 – 8.60 (m, 2H), 8.49 – 8.41 (m, 1H), 7.82 – 7.73 (m, 1H), 7.73 – 7.64 (m, 1H), 7.41 – 7.30

	(m, 2H), 4.34 – 4.98 (m, 1H), 3.43 – 3.35 (m, 2H), 3.15 (br s, 1H), 2.84 (br s, 2H), 2.07 – 1.84 (m, 2H), 1.74 – 1.54 (m, 2H).
CDK7_811	¹ H NMR (400 MHz, DMSO) δ 8.96 and 8.92 (2s, 1H, rotamers), 8.56 and 8.47 (2d, <i>J</i> = 8.0 Hz, 1H, rotamers), 8.39 and 8.18 (2d, <i>J</i> = 8.4, 1H), 8.28 – 8.22 (m, 2H), 7.75 (q, <i>J</i> = 7.0 Hz, 1H), 7.64 – 7.58 (m, 1H), 4.09 – 3.96 (m, 1H), 3.19 – 3.10 (m, 2H), 2.94 – 2.82 (m, 2H), 2.03 – 1.95 (m, 1H), 1.78 – 1.64 (m, 1H), 1.60 – 1.38 (m, 2H).
CDK7_814	¹ H NMR (400 MHz, DMSO) δ 12.12 (br.s, 1H), 8.59 – 8.44 (m, 1H), 7.80 – 7.62 (m, 1H), 7.34 – 7.27 (m, 1H), 7.24 – 7.15 (m, 1H), 3.97 – 3.86 (m, 1H), 3.42 – 3.34 (m, 2H), 3.13 – 3.06 (m, 1H), 2.99 – 2.93 (m, 2H), 2.92 – 2.85 (m, 1H), 2.57 – 2.51 (m, 2H), 1.95 – 1.86 (m, 1H), 1.73 – 1.64 (m, 1H), 1.54 – 1.38 (m, 2H).
CDK7_829	¹ H NMR (400 MHz, DMSO) δ 8.45 and 8.38 (2s, 1H, rotamers), 7.78 and 7.64 (2d, <i>J</i> = 7.3 Hz, 1H, rotamers), 6.84 – 6.74 (m, 1H), 6.74 – 6.65 (m, 1H), 6.45 (t, <i>J</i> = 7.7 Hz, 1H), 4.92 and 4.80 (2s, 1H, rotamers), 3.18 – 3.11 (m, 2H), 2.97 – 2.88 (m, 1H), 2.85 – 2.56 (m, 4H), 2.39 – 2.23 (m, 2H), 1.83 – 1.76 (m, 2H), 1.75 – 1.51 (m, 1H), 1.49 – 0.82 (m, 4H).
CDK7_830	¹ H NMR (400 MHz, DMSO) δ 8.88 – 8.80 (m, 1H), 8.54 and 8.47 (2s, 1H, rotamers), 8.46 – 8.42 (m, 1H), 8.27 (s, 1H), 7.98 – 7.90 (m, 1H), 7.74 and 7.43 (2s, 1H, rotamers), 7.69 – 7.60 (m, 2H), 7.60 – 7.52 (m, 1H), 3.87 – 3.79 (m, 1H), 2.99 – 2.76 (m, 1H), 2.41 – 2.25 (m, 2H), 2.19 – 2.07 (m, 1H), 1.78 and 1.51 (m, 1H), 1.40 – 1.23 (m, 1H), 1.18 – 0.88 (m, 1H), 0.83 – 0.50 (m, 1H).
CDK7_834	¹ H NMR (400 MHz, DMSO) δ 12.47 (br s, 1H), 8.66 – 8.50 (m, 1H), 8.38 (s, 1H), 8.05 – 7.70 (m, 2H), 7.68 – 7.57 (m, 1H), 7.52 (d, <i>J</i> = 3.6 Hz, 1H), 7.22 (d, <i>J</i> = 3.6 Hz, 1H), 4.22 and 3.95 (m, 1H), 3.45 – 2.87 (m, 2H), 2.73 – 2.55 (m, 2H), 2.04 – 1.85 (m, 1H), 1.84 – 1.64 (m, 2H), 1.64 – 1.44 (m, 1H).
CDK7_835	¹ H NMR (400 MHz, DMSO) δ 12.46 – 12.04 (m, 1H), 10.72 – 10.27 (2s, 1H, rotamers), 8.61 – 8.49 (m, 1H), 8.36 (s, 1H), 7.96 (s, 1H), 7.84 – 7.63 (m, 1H), 7.63 – 7.43 (m, 3H), 4.19 – 4.05 (m, 1H), 4.04 – 3.93 (m, 1H), 3.83 (s, 3H), 3.17 – 3.09 (m, 1H), 3.08 – 3.00 (m, 1H), 2.98 – 2.85 (m, 1H), 2.03 – 1.85 (m, 1H), 1.83 – 1.45 (m, 3H).
CDK7_836	¹ H NMR (400 MHz, DMSO) δ 12.74 – 12.07 (m, 1H), 10.82 and 10.39 (2s, 1H, rotamers), 8.62 – 8.51 (m, 1H), 8.49 – 8.44 (m, 2H), 8.33 (s, 1H), 7.89 – 7.49 (m, 5H), 4.22 – 4.02 (m, 1H), 3.39 – 3.15 (m, 1H), 3.12 – 2.96 (m, 1H), 2.75 – 2.59 (m, 2H), 2.00 – 1.88 (m, 1H), 1.86 – 1.47 (m, 3H).
CDK7_838	¹ H NMR (400 MHz, DMSO) δ 12.23 (m, 1H), 10.31 and 9.91 (2s, 1H, rotamers), 8.65 – 8.46 (m, 1H), 7.81 – 7.44 (m, 5H), 7.02 – 6.88 (m, 2H), 4.16 – 3.90 (m, 1H), 3.77 – 3.71 (m, 4H), 3.37 – 3.26 (m, 1H), 3.10 – 3.02 (m, 4H), 3.02 and 2.90 (2br s, 1H, rotamers), 2.70 – 2.53 (m, 2H), 2.00 – 1.87 (m, 1H), 1.81 – 1.44 (m, 3H).
CDK7_843	¹ H NMR (400 MHz, DMSO) δ 8.60 – 8.47 (m, 1H), 7.86 – 7.62 (m, 1H), 7.39 – 7.32 (m, 1H), 7.30 – 7.22 (m, 1H), 3.99 – 3.87 (m, 4H), 3.40 – 3.31 (m, 2H), 3.13 – 3.05 (m, 1H), 2.97 – 2.84 (m, 3H), 2.57 – 2.51 (m, 2H), 1.95 – 1.86 (m, 1H), 1.75 – 1.64 (m, 1H), 1.54 – 1.40 (m, 2H).
CDK7_845	¹ H NMR (400 MHz, DMSO) δ 8.68 – 8.46 (m, 1H), 7.96 – 7.75 (m, 1H), 7.62 (s, 1H), 7.37 – 7.25 (m, 1H), 7.22 – 6.94 (m, 1H), 4.03 – 3.93 (m, 1H), 3.55 (s, 3H), 3.17 – 3.08 (m, 1H), 2.95 – 2.83 (m, 1H), 2.58 – 2.52 (m, 4H), 2.08 – 1.87 (m, 1H), 1.81 – 1.62 (m, 1H), 1.60 – 1.40 (m, 2H).
CDK7_846	¹ H NMR (400 MHz, DMSO) δ 12.25 – 11.89 (m, 1H), 8.54 (s, 1H), 8.49 and 8.22 (2s, 1H, rotamers), 7.81 – 7.60 (m, 1H), 7.48 and 7.24 (2s, 1H, rotamers), 7.40 (s, 1H), 4.16 – 3.93 (m, 1H), 3.32 – 3.25 and 3.17 – 3.10 (2m, 1H,

	rotamers), 3.08 – 2.89 (m, 1H), 2.84 – 2.69 (m, 4H), 2.69 – 2.56 (m, 2H), 2.03 – 1.83 (m, 1H), 1.81 – 1.43 (m, 3H).
CDK7_847	¹ H NMR (400 MHz, DMSO) δ 12.27 (s, 1H), 9.37 and 8.92 (2s, 1H, rotamers), 8.64 – 8.45 (m, 1H), 7.87 – 7.65 (m, 1H), 7.62 – 7.36 (m, 2H), 4.06 (m, 3H), 3.35 – 3.27 and 3.18 – 3.09 (2m, 1H, rotamers), 3.08 – 2.99 and 2.96 – 2.89 (2m, 1H, rotamers), 2.71 – 2.53 (m, 3H), 1.97 – 1.88 (m, 1H), 1.81 – 1.46 (m, 3H).
CDK7_848	¹ H NMR (400 MHz, DMSO) δ 12.96 – 12.21 (m, 2H), 8.67 – 8.49 (m, 1H), 8.08 – 7.48 (m, 3H), 7.37 – 7.22 (m, 1H), 3.93 – 3.73 (m, 1H), 3.07 – 2.82 (m, 1H), 2.71 – 2.54 (m, 1H), 2.02 – 1.71 (m, 2H), 1.68 – 1.52 (m, 1H), 1.50 – 1.10 (m, 10H).
CDK7_849	¹ H NMR (400 MHz, DMSO) δ 12.26 – 11.93 (m, 1H), 8.63 – 8.45 (m, 1H), 7.85 – 7.67 (m, 1H), 7.57 and 7.43 (2s, 1H, rotamers), 7.02 (s, 1H), 4.07 – 3.93 (m, 2H), 3.75 – 3.69 (m, 4H), 3.67 – 3.62 (m, 4H), 3.25 – 3.09 (m, 2H), 3.03 – 2.87 (m, 1H), 2.01 – 1.86 (m, 1H), 1.82 – 1.66 (m, 1H), 1.63 – 1.42 (m, 2H).
CDK7_850	¹ H NMR (400 MHz, DMSO) δ 12.21 – 11.97 (m, 1H), 8.68 – 8.52 (m, 2H), 8.20 – 8.10 (m, 1H), 7.94 – 7.84 (m, 1H), 7.62 and 7.43 (2s, 1H, rotamers), 7.39 – 7.28 (m, 1H), 4.28 – 4.15 (m, 2H), 4.06 – 3.96 (m, 2H), 3.45 – 3.33 (m, 3H), 3.28 – 3.17 (m, 1H), 2.95 – 2.76 (m, 2H), 2.03 – 1.86 (m, 2H), 1.81 – 1.41 (m, 6H).
CDK7_851	¹ H NMR (400 MHz, DMSO) δ 12.13 – 11.85 (m, 1H), 8.58 – 8.45 (m, 1H), 8.31 (s, 1H), 8.30 – 8.25 and 8.03 – 7.96 (2m, 1H, rotamers), 7.75 – 7.68 and 7.68 – 7.58 (2m, 1H, rotamers), 7.51 – 7.27 (m, 2H), 4.09 – 3.91 (m, 1H), 3.74 – 3.65 (m, 1H), 3.32 – 3.05 (m, 1H), 3.04 – 2.84 (m, 1H), 1.95 – 1.61 (m, 8H), 1.60 – 1.44 (m, 2H), 1.43 – 1.15 (m, 5H).
CDK7_888	¹ H NMR (400 MHz, DMSO) δ 8.51 (d, J = 8 Hz 1H), 8.47 – 8.41 (m, 1H), 8.27 – 8.23 (m, 1H), 7.71 (m, 1H), 7.62 (d, J = 8 Hz, 1H), 7.51 – 7.43 (2H, m), 4.02 – 3.90 (m, 1H), 3.22 – 3.14 (m, 1H), 3.01 – 2.93 (m, 1H), 2.88 – 2.77 (m, 1H), 2.61 – 2.55 (m, 1H), 1.88 – 1.80 (m, 1H), 1.79 – 1.70 (m, 6H), 1.68 – 1.62 (m, 1H), 1.57 – 1.39 (m, 2H).
CDK7_891	¹ H NMR (400 MHz, DMSO) δ 8.64 (s, 1H), 8.39 (s, 1H), 8.29 and 8.24 (2s, 1H, rotamers), 8.26 (s, 1H), 7.99 – 7.83 and 7.41 – 7.24 (2m, 1H, rotamers), 7.74 – 7.54 (m, 2H), 7.54 – 7.44 (m, 2H), 4.03 – 3.92 and 3.85 – 3.70 (2m, 1H, rotamers), 3.14 – 3.00 (m, 1H), 2.96 – 2.78 (m, 1H), 2.48 – 2.31 (m, 2H), 1.92 – 1.76 (m, 1H), 1.75 – 1.56 (m, 1H), 1.55 – 1.34 (m, 2H).
CDK7_897	¹ H NMR (400 MHz, DMSO) δ 8.57 and 8.51 (2s, 1H, rotamers), 8.04 and 7.80 (2d, J = 7.4 Hz, 1H, rotamers), 7.94 – 7.84 (m, 1H), 7.64 (q, 9.0 Hz, 1H), 7.49 – 7.39 (m, 1H), 7.37 – 7.28 (m, 1H), 3.84 – 3.74 and 3.13 – 3.04 (2m, 1H, rotamers), 2.97 – 2.89 and 2.68 – 2.61 (m, 1H, rotamers), 2.78 – 2.70 (m, 1H), 2.36 – 2.25 (m, 2H), 1.86 – 1.63 (m, 1H), 1.63 – 1.50 (m, 7H), 1.50 – 1.27 (m, 2H), 1.27 – 0.99 (m, 1H).
CDK7_898	¹ H NMR (400 MHz, DMSO) δ 10.13 – 9.96 (m, 1H), 8.21 and 8.15 (2s, 1H, rotamers), 8.09 and 7.94 (2s, 1H, rotamers), 7.82 – 7.67 (m, 1H), 7.35 and 7.01 (2d, J = 7.6 Hz, 1H, rotamers), 7.28 – 7.21 (m, 1H), 7.21 – 7.13 (m, 2H), 3.90 – 3.68 and 3.44 – 3.40 (2m, 1H, rotamers), 3.00 – 2.82 (m, 1H), 2.80 – 2.68 (m, 1H), 2.42 – 2.27 (m, 2H), 2.09 – 2.01 (m, 3H), 1.83 – 1.73 (m, 1H), 1.62 – 1.51 (m, 1H), 1.44 – 1.31 (m, 1H), 1.44 – 1.31 (m, 1H), 1.31 – 1.25 (m, 1H), 1.25 – 1.17 and 1.06 – 1.00 (2m, 1H, rotamers).
CDK7_901	¹ H NMR (400 MHz, DMSO) δ 12.01 (br s, 1H), 8.90 – 8.72 (m, 1H), 8.42 – 8.27 (m, 1H), 7.62 – 7.48 (m, 1H), 7.45 – 7.32 (m, 1H), 7.21 – 7.08 (m, 2H), 6.49 (s, 1H), 4.17 – 4.04 (m, 1H), 3.98 – 3.88 (m, 1H), 3.34 – 3.22 (m, 1H),

	3.09 – 2.93 (m, 1H), 2.80 – 2.57 (m, 2H), 2.04 – 1.71 (m, 2H), 1.67 – 1.42 (m, 2H).
CDK7_903	¹ H NMR (400 MHz, DMSO) δ 10.65 and 10.39 (2s, 1H, rotamers), 8.32 – 8.11 (m, 2H), 7.54 and 7.32 (2d, <i>J</i> = 8.0 Hz, 1H, rotamers), 7.49 – 7.38 (m, 1H), 6.99 (d, <i>J</i> = 7.9 Hz, 1H), 4.00 – 3.66 (m, 1H), 3.13 – 2.99 (m, 1H), 2.93 – 2.77 (m, 1H), 2.45 (s, 2H), 2.38 (s, 3H), 1.96 – 1.86 (m, 1H), 1.79 – 1.60 (m, 7H), 1.48 – 1.38 (m, 2H).
CDK7_904	¹ H NMR (400 MHz, DMSO) δ 10.58 and 10.34 (2s, 1H, rotamers), 8.45 – 8.11 (m, 2H), 7.48 and 7.22 (2d, <i>J</i> = 8.0 Hz, 1H, rotamers), 7.41 – 7.29 (m, 2H), 3.86 – 3.54 (m, 1H), 3.02 – 2.93 (m, 1H), 2.80 – 2.73 (m, 1H), 2.44 – 2.34 (m, 2H), 2.31 (s, 3H), 1.93 – 1.80 (m, 1H), 1.78 – 1.68 (m, 6H), 1.63 – 1.56 (m, 1H), 1.42 – 1.31 (m, 2H).
CDK7_921	¹ H NMR (400 MHz, DMSO) δ 12.01 (br s, 1H), 8.79 – 8.64 (m, 1H), 8.33 – 8.20 (m, 1H), 8.10 – 7.94 (m, 1H), 7.73 – 7.64 (m, 1H), 7.58 – 7.47 (m, 1H), 7.33 – 7.17 (m, 2H), 3.92 – 3.64 (m, 1H), 3.11 – 2.94 (m, 1H), 2.83 – 2.66 (m, 1H), 2.45 – 2.32 (m, 2H), 1.92 – 1.89 (m, 2H), 1.69 – 1.52 (m, 1H), 1.52 – 1.21 (m, 2H).
CDK7_929	¹ H NMR (400 MHz, DMSO) δ 12.15 (br s, 1H), 8.63 – 8.48 (m, 1H), 7.88 – 7.72 (m, 1H), 7.29 – 7.16 (m, 1H), 7.10 – 6.92 (m, 1H), 6.08 – 5.92 (m, 1H), 3.95 – 3.78 (m, 2H), 3.77 – 3.62 (m, 2H), 3.12 – 2.95 (m, 4H), 2.90 – 2.75 (m, 1H), 2.49 – 2.41 (m, 2H), 1.96 – 1.80 (m, 1H), 1.74 – 1.58 (m, 1H), 1.52 – 1.31 (m, 2H).
CDK7_944	¹ H NMR (400 MHz, DMSO) δ 12.31 (br s, 1H), 8.58 – 8.46 (m, 1H), 7.80 and 7.60 (2d, 8.0 Hz, 1H, rotamers), 7.54 – 7.42 (m, 1H), 4.61 – 4.35 (m, 2H), 4.03 – 3.85 (m, 1H), 3.18 – 3.06 (m, 1H), 3.04 (s, 3H), 2.96 – 2.83 (m, 1H), 2.62 – 2.52 (m, 2H), 2.07 – 1.86 (m, 1H), 1.80 – 1.63 (m, 1H), 1.62 – 1.42 (m, 2H).
CDK7_951	¹ H NMR (400 MHz, DMSO) δ 12.20 – 11.90 (m, 1H), 8.68 and 8.53 (2s, 1H, rotamers), 8.48 and 8.28 (2s, 1H, rotamers), 7.75 and 7.63 (2d, <i>J</i> = 7.0 Hz, 1H, rotamers), 7.47 and 7.27 (2s, 1H, rotamers), 7.41 (s, 1H), 4.14 – 3.91 (m, 1H), 3.36 – 3.24 and 3.18 – 3.06 (2m, 1H, rotamers), 3.06 – 2.86 (m, 1H), 2.85 – 2.72 (m, 1H), 2.69 – 2.51 (m, 2H), 1.99 – 1.82 (m, 1H), 1.81 – 1.44 (m, 3H), 0.74 – 0.62 (m, 2H), 0.55 (s, 2H).
CDK7_953	¹ H NMR (400 MHz, DMSO) δ 11.64 – 11.41 (m, 1H), 8.62 – 8.46 (m, 1H), 7.87 – 7.70 (m, 1H), 7.12 – 6.99 (m, 1H), 3.93 – 3.83 (m, 1H), 3.48 – 3.40 (m, 2H), 3.12 – 2.99 (m, 4H), 2.96 – 2.80 (m, 4H), 2.49 – 2.40 (m, 2H), 2.00 – 1.85 (m, 3H), 1.70 – 1.61 (m, 1H), 1.50 – 1.38 (m, 2H).
CDK7_959	¹ H NMR (400 MHz, DMSO) δ 8.68 – 8.46 (m, 1H), 8.32 – 8.18 (m, 2H), 7.92 – 7.83 (m, 1H), 7.83 – 7.78 (m, 1H), 7.65 (s, 1H), 7.50 – 7.34 (m, 1H), 7.26 and 7.11 (2d, <i>J</i> = 7.4 Hz, 1H, rotamers), 4.03 – 3.93 (m, 1H), 3.90 (s, 3H), 3.16 – 3.06 (m, 1H), 2.93 – 2.82 (m, 1H), 2.63 – 2.56 (m, 2H), 2.06 – 1.85 (m, 1H), 1.77 – 1.64 (m, 1H), 1.61 – 1.41 (m, 2H).
CDK7_966	¹ H NMR (400 MHz, DMSO) δ 8.63 and 8.56 (2s, 1H, rotamers), 8.32 (s, 1H), 8.06 (s, 1H), 7.89 – 7.70 (m, 2H), 4.15 – 3.92 (m, 1H), 3.62 (s, 3H), 3.05 (s, 3H), 2.68 – 2.54 (m, 2H), 2.01 – 1.87 (m, 2H), 1.83 – 1.67 (m, 1H), 1.63 – 1.45 (m, 3H).
CDK7_1017	¹ H NMR (400 MHz, DMSO) δ 8.52 and 8.45 (m, 1H, rotamers), 7.67 – 7.20 (m, 2H), 6.84 (d, <i>J</i> = 7.6 Hz, 1H), 3.93 – 3.79 (m, 1H), 3.79 – 3.69 (m, 3H), 3.12 – 2.96 (m, 5H), 2.85 – 2.72 (m, 1H), 2.45 – 2.35 (m, 2H), 1.89 (s, 4H), 1.68 – 1.58 (m, 1H), 1.51 – 1.36 (m, 2H).
CDK7_550	¹ H NMR (400 MHz, DMSO) δ 9.17 – 9.07, and 8.50 – 8.34 (2m, 1H, rotamers), 9.06 – 8.05 (m, 1H), 8.41 – 8.23 (m, 1H), 7.76 – 7.56 and 7.14 – 6.99 (2m, 1H,

	rotamers), 7.42 – 7.16 (m, 3H), 4.22 – 4.05 (m, 1H), 3.73 – 3.66 (m, 1H), 3.52 – 3.46 (m, 1H), 3.29 – 2.99 (m, 2H), 2.95 – 2.62 (m, 2H), 1.94 – 1.33 (m, 4H)
CDK7_837	¹ H NMR (400 MHz, DMSO) δ 12.44 – 12.27 (m, 1H), 10.60 – 10.36 (m, 1H), 8.60 – 8.50 (m, 1H), 8.00 – 7.95 (m, 2H), 7.85 – 7.80 (m, 2H), 7.78 – 7.50 (m, 3H), 4.10 – 3.80 (m, 1H), 3.20 – 2.80 (m, 2H), 2.00 – 1.80 (m, 1H), 1.79 – 1.60 (m, 1H), 1.59 – 1.42 (m, 2H), 1.35 – 1.16 (m, 1H), 0.97 – 0.76 (m, 1H).
CDK7_900	¹ H NMR (400 MHz, DMSO) δ 10.65 (s, 1H), 8.25 (s, 1H), 7.89 – 7.64 (m, 1H), 7.15 – 6.92 (m, 3H), 6.66 – 6.43 (m, 1H), 4.77 – 4.53 (m, 2H), 4.16 – 3.85 (m, 1H), 2.92 – 2.52 (m, 4H), 1.69 – 1.16 (m, 5H)
CDK7_997	¹ H NMR (400 MHz, DMSO) δ 12.23 – 11.94 (m, 1H), 9.22 – 9.12 and 8.89 – 8.79 (2m, 1H, rotamers), 8.58 – 8.44 (m, 1H), 7.76 – 7.59 (m, 1H), 7.53 – 7.46 (m, 1H), 7.46 – 7.34 (m, 1H), 7.33 – 7.28 (m, 3H), 7.28 – 7.19 (m, 1H), 4.53 – 4.39 (m, 2H), 4.11 – 3.89 (m, 1H), 3.30 – 3.04 (m, 1H), 3.01 – 2.84 (m, 1H), 2.65 – 2.52 (m, 2H), 2.13 – 2.02 (m, 2H), 1.96 – 1.85 (m, 1H), 1.78 – 1.66 (m, 1H), 1.66 – 1.42 (m, 2H).
CDK7_998	¹ H NMR (400 MHz, DMSO) δ 12.31 – 11.99 (m, 1H), 8.58 – 8.40 (m, 1H), 7.78 – 7.63 (m, 1H), 7.58 – 7.44 (m, 1H), 7.42 – 7.33 (m, 2H), 7.32 – 7.24 (m, 3H), 7.15 – 6.90 (m, 1H), 4.89 – 4.63 (m, 2H), 4.01 – 3.83 (m, 2H), 3.99 – 2.99 (m, 4H), 2.96 – 2.81 (m, 2H), 2.64 – 2.52 (m, 1H), 2.00 – 1.58 (m, 2H), 1.56 – 1.39 (m, 2H).
CDK7_999	¹ H NMR (400 MHz, DMSO) δ 8.64 – 8.51 (m, 1H), 7.89 – 7.80 (m, 1H), 7.72 and 7.59 (2s, 1H, rotamers), 7.37 – 7.29 (m, 1H), 4.27 – 4.19 (m, 3H), 4.16 – 3.94 (m, 1H), 3.32 – 3.11 (m, 1H), 3.07 – 2.88 (m, 1H), 2.69 – 2.53 (m, 2H), 2.00 – 1.88 (m, 2H), 1.81 – 1.69 (m, 1H), 1.64 – 1.47 (m, 2H).
CDK7_1000	¹ H NMR (400 MHz, DMSO) δ 8.86 – 8.53 (m, 1H), 7.88 – 7.76 (m, 1H), 7.71 and 7.60 (2s, 1H, rotamers), 7.33 (s, 1H), 4.27 – 4.20 (m, 3H), 4.09 – 3.94 (m, 2H), 3.26 – 3.18 (m, 1H), 3.16 – 3.09 (m, 1H), 3.00 – 2.05 (m, 1H), 2.64 – 2.53 (m, 2H), 2.01 – 1.87 (m, 1H), 1.79 – 1.76 (m, 1H), 1.59 – 1.44 (m, 2H).
CDK7_1001	¹ H NMR (400 MHz, DMSO) δ 12.07 – 11.86 (m, 1H), 8.59 – 8.47 (m, 1H), 7.78 – 7.63 (m, 1H), 7.57 – 7.44 (m, 1H), 7.43 – 7.30 (m, 1H), 4.03 – 3.69 (m, 2H), 3.23 – 3.12 (m, 1H), 3.02 (s, 3H), 2.96 – 2.86 (m, 1H), 2.58 – 2.51 (m, 2H), 1.99 – 1.87 (m, 1H), 1.79 – 1.63 (m, 1H), 1.57 – 1.42 (m, 2H), 0.94 – 0.84 (m, 2H), 0.77 – 0.68 (m, 2H).
CDK7_1003	¹ H NMR (400 MHz, DMSO) δ 12.53 – 12.44 (m, 1H), 8.58 – 8.49 (m, 1H), 7.76 – 7.69 (m, 1H), 7.67 – 7.54 (m, 3H), 7.39 – 7.30 (m, 2H), 3.99 – 3.94 (m, 3H), 3.92–3.82 (m, 1H), 3.14 – 3.00 (m, 1H), 2.86 – 2.76 (m, 1H), 2.46 – 2.39 (m, 2H), 1.97 – 1.92 (m, 1H), 1.69 – 1.59 (m, 1H), 1.54 – 1.35 (m, 2H).
CDK7_991	¹ H NMR (400 MHz, DMSO) δ 11.42 (s, 1H), 8.55 – 8.44 (m, 1H), 8.28 (s, 1H), 7.73 – 7.55 (m, 1H), 7.10 – 6.94 (m, 2H), 2.91 – 2.78 (m, 8H), 1.88 – 1.74 (m, 1H), 1.71 – 1.60 (m, 1H), 1.52 – 1.36 (m, 1H).
CDK7_924	¹ H NMR (400 MHz, DMSO) δ 8.76 – 8.51 (m, 1H), 7.95 – 7.78 (m, 1H), 7.69 (s, 1H), 7.35 – 7.00 (m, 2H), 4.01 – 3.83 (m, 1H), 3.15 – 3.05 (m, 1H), 2.97 – 2.76 (m, 1H), 2.51 – 2.55 (m, 2H), 2.32 (s, 3H), 2.13 (s, 3H), 2.02 – 1.85 (m, 1H), 1.81 – 1.60 (m, 1H), 1.57 – 1.33 (m, 2H).
CDK7_955	¹ H NMR (400 MHz, DMSO) δ 11.92 – 11.46 (m, 1H), 8.66 – 8.47 (m, 1H), 7.99 (s, 1H), 7.90 – 7.76 (m, 1H), 7.63 (s, 1H), 7.24 – 7.15 (m, 1H), 4.01 – 3.89 (m, 1H), 3.85 – 3.72 (m, 5H), 3.19 – 3.07 (m, 1H), 3.04 – 2.86 (m, 3H), 2.62 – 2.52 (m, 2H), 2.15 – 2.00 (m, 2H), 1.97 – 1.84 (m, 1H), 1.77 – 1.64 (m, 1H), 1.57 – 1.39 (m, 2H).
CDK7_1019	¹ H NMR (400 MHz, DMSO) δ 8.59 – 8.41 (m, 1H), 7.69 – 7.50 (m, 1H), 7.49 – 7.34 (m, 1H), 7.05 (s, 1H), 4.07 – 3.86 (m, 1H), 2.91 – 2.74 (m, 1H), 2.65 –

	2.51 (m, 2H), 2.16 (s, 3H), 1.96 – 1.74 (m, 8H), 1.73 – 1.62 (m, 1H), 1.60 – 1.44 (m, 1H), 1.37 – 1.21 (m, 1H).
CDK7_1020	¹ H NMR (400 MHz, DMSO) δ 12.46 (br.s, 1H), 8.61 – 8.48 (m, 1H), 7.90 – 7.75 (m, 1H), 7.66 and 7.53 (2s, 1H, rotamers), 7.32 – 7.23 (m, 1H), 4.31 – 4.24 (m, 2H), 4.08 – 3.96 (m, 1H), 3.22 – 3.12 (m, 1H), 3.01 – 2.91 (m, 1H), 2.68 – 2.55 (m, 2H), 1.97 – 1.87 (m, 1H), 1.80 – 1.68 (m, 1H), 1.62 – 1.46 (m, 2H), 1.30 (t, <i>J</i> = 7.0 Hz, 3H).
CDK7_1105	¹ H NMR (400 MHz, DMSO) δ 8.64 – 8.50 (m, 1H), 7.83 (d, <i>J</i> =7.4, 1H), 7.61 (s, 1H), 7.25 – 7.16 and 7.03 – 6.94 (2m, 1H, rotamers), 7.12 (s, 1H), 4.00 – 3.86 (m, 1H), 3.42 – 3.32 (m, 1H), 3.15 – 3.05 (m, 1H), 2.95 – 2.79 (m, 2H), 2.58 – 2.52 (m, 1H), 1.99 – 1.88 (m, 1H), 1.76 – 1.62 (m, 1H), 1.57 – 1.41 (m, 2H), 1.10 – 0.97 (m, 2H), 0.93 – 0.81 (m, 2H).
CDK7_1058	¹ H NMR (400 MHz, DMSO) δ 8.67 – 8.57 (m, 1H), 8.56 – 8.47 (m, 1H), 7.91 – 7.87 (m, 1H), 7.77 (s, 2H), 7.64 – 7.57 (m, 1H), 7.51 and 7.37 (2d, <i>J</i> =7.7 Hz, rotamers, 1H), 4.00 – 3.84 (m, 1H), 3.17 – 3.01 (m, 2H), 2.89 – 2.81 (m, 1H), 2.03 – 1.86 (m, 1H), 1.74 – 1.63 (m, 1H), 2.00 – 1.37 (m, 2H), 1.24 (s, 1H), 0.90 – 0.74 (m, 1H).
CDK7_1176	¹ H NMR (400 MHz, DMSO) δ 12.00 – 11.67 (m, 1H), 8.65 – 8.52 (m, 1H), 8.49 – 8.42 (m, 1H), 7.90 – 7.84 (m, 1H), 7.83 – 7.78 (m, 1H), 7.76 – 7.72 (m, 1H), 7.24 – 7.15 (m, 2H), 4.18 – 4.04 (m, 2H), 4.01 – 3.89 (m, 1H), 3.19 – 3.07 (m, 1H), 3.04 – 2.86 (m, 4H), 2.63 – 2.52 (m, 2H), 2.13 – 2.02 (m, 2H), 1.96 – 1.83 (m, 1H), 1.77 – 1.65 (m, 1H), 1.56 – 1.42 (m, 2H).
CDK7_1187	¹ H NMR (400 MHz, DMSO) δ 11.60 – 11.40 (m, 1H), 8.61 – 8.45 (m, 1H), 8.31 (s, 1H, formate), 7.82 – 7.66 (m, 1H), 7.13 – 6.99 (m, 1H), 3.94 – 3.82 (m, 1H), 3.50 – 3.43 (s, 2H), 3.05 (s, 3H), 2.97 – 2.79 (m, 2H), 2.74 – 2.64 (m, 1H), 2.03 – 1.09 (m, 2H), 1.82 – 1.50 (m, 3H), 1.44 – 1.31 (m, 1H), 1.18 – 1.02 (m, 6H).
CDK7_1188	¹ H NMR (400 MHz, DMSO) δ 11.66 – 11.37 (m, 1H), 8.63 – 8.45 (m, 1H), 7.82 – 7.66 (m, 1H), 7.15 – 6.98 (m, 1H), 3.97 – 3.75 (m, 1H), 3.45 – 3.38 (m, 2H), 3.12 – 2.97 (m, 1H), 2.89 – 2.78 (m, 4H), 2.48 – 2.41 (m, 2H), 1.96 – 1.82 (m, 3H), 1.72 – 1.59 (m, 1H), 1.52 – 1.34 (m, 2H), 0.80 – 0.74 (m, 2H), 0.65 – 0.59 (m, 2H).
CDK7_1199	¹ H NMR (400 MHz, DMSO) δ 12.92 – 12.60 (m, 1H), 8.66 – 8.54 (m, 1H), 7.84 – 7.76 (m, 1H), 7.70 – 7.61 (m, 1H), 7.44 – 7.32 (m, 1H), 3.90 (s, 3H), 2.21 – 2.09 (m, 3H), 2.04 – 1.81 (m, 2H), 1.72 – 1.55 (m, 1H), 1.32 – 1.15 (m, 4H), 1.03 – 0.78 (m, 4H).
CDK7_1122	¹ H NMR (400 MHz, DMSO) δ 11.80 – 11.54 (m, 1H), 8.62 – 8.46 (m, 1H), 7.99 (s, 1H), 7.80 – 7.68 (m, 1H), 7.63 (s, 1H), 7.19 – 7.06 (m, 1H), 3.87 – 3.70 (m, 5H), 3.00 – 2.81 (m, 3H), 2.70 – 2.53 (m, 2H), 2.11 – 2.01 (m, 2H), 1.89 (s, 1H), 1.81 – 1.71 (m, 1H), 1.69 – 1.57 (m, 1H), 1.53 – 1.43 (m, 1H), 1.36 – 1.22 (m, 1H), 1.09 – 0.99 (m, 6H).
CDK7_1123	¹ H NMR (400 MHz, DMSO) δ 8.61 (m, 1H), 8.24 (s, 1H), 7.96 – 7.87 (m, 1H), 7.81 (s, 1H), 7.66 (s, 1H), 7.47 – 7.42 and 7.12 – 6.99 (m, 1H, rotamers), 7.41 – 7.35 and 7.30 – 7.23 (m, 1H, rotamers), 4.09 – 3.99 (m, 1H), 3.91 (s, 3H), 3.09 – 3.01 (m, 1H), 2.87 – 2.75 (m, 1H), 1.93 – 1.41 (m, 4H), 1.26 – 1.10 (m, 6H).
CDK7_1021	¹ H NMR (400 MHz, DMSO) δ 12.36 (bs, 1H), 8.62 – 8.43 (m, 1H), 7.91 – 7.69 (m, 1H), 7.08 (s, 1H), 3.95 – 3.59 (m, 3H), 3.12 (s, 3H), 3.08 – 2.95 (m, 1H), 2.88 – 2.64 (m, 3H), 2.44 – 2.31 (m, 2H), 1.96 – 1.78 (m, 1H), 1.73 – 1.58 (m, 1H), 1.54 – 1.32 (m, 2H).
CDK7_1052	¹ H NMR (400 MHz, DMSO) δ 8.65 – 8.55 (m, 1H), 8.08 – 7.97 (m, 1H), 7.86 – 7.76 (m, 1H), 7.71 – 7.62 (m, 1H), 7.61 – 7.52 (m, 1H), 7.33 – 7.26 and

	7.16 – 7.08 (2m, 1 H, rotamers), 7.26 – 7.19 (m, 1H), 6.51 – 6.45 (m, 1H), 3.96 – 3.81 (m, 2H), 3.49 (s, 3H), 3.10 – 3.00 (m, 1H), 2.87 – 2.74 (m, 1H), 2.69 – 2.66 and 2.35 – 2.31 (2m, 1H, rotamers), 2.49 – 2.37 (m, 2H), 2.00 – 1.90 (m, 1H), 1.70 – 1.59 (m, 1H), 1.56 – 1.35 (m, 1H).
CDK7_1060	¹ H NMR (400 MHz, DMSO) δ 12.74 – 12.38 (m, 1H), 8.60 – 8.48 (m, 1H), 7.80 – 7.69 (m, 1H), 7.66 and 7.56 (s, 1H, rotamers), 7.29 (s, 1H), 4.08 – 3.85 (m, 1H), 3.20 – 3.06 (m, 1H), 2.97 – 2.82 (m, 2H), 2.65 (s, 3H), 2.58 – 2.51 (m, 2H), 2.08 – 1.83 (m, 1H), 1.81 – 1.60 (m, 1H), 1.60 – 1.41 (m, 1H).
CDK7_1061	¹ H NMR (400 MHz, DMSO) δ 12.31 – 11.86 (m, 1H), 8.63 – 8.40 (m, 2H), 7.89 – 7.62 (m, 3H), 7.60 – 7.43 (m, 1H), 7.35 – 7.16 (m, 2H), 4.14 – 3.89 (m, 1H), 3.26 – 3.07 (m, 1H), 3.01 – 2.85 (m, 1H), 2.65 – 2.53 (m, 2H), 2.00 – 1.84 (m, 1H), 1.79 – 1.66 (m, 1H), 1.65 – 1.45 (m, 2H).
CDK7_1062	¹ H NMR (400 MHz, DMSO) δ 12.56 – 12.03 (m, 1H), 8.85 – 8.71 (m, 2H), 8.61 – 8.44 (m, 1H), 8.31 (s, 1H), 7.76 – 7.66 (m, 1H), 7.65 – 7.53 (m, 1H), 7.50 (s, 1H) 7.35 – 7.20 (m, 1H), 4.10 – 3.83 (m, 1H), 3.22 – 3.03 (m, 1H), 2.98 – 2.82 (m, 1H), 2.61 – 2.51 (m, 2H), 2.01 – 1.82 (m, 1H), 1.79 – 1.61 (m, 1H), 1.59 – 1.39 (m, 2H).
CDK7_1121	¹ H NMR (400 MHz, DMSO) δ 11.17 (m, 1H), 8.66 – 8.53 (m, 1H), 8.27 (s, 1H), 7.89 – 7.80 (m, 1H), 7.76 (s, 1H), 7.04 – 6.82 (m, 2H), 5.66 – 5.48 (m, 1H), 4.05 – 3.97 (m, 2H), 3.96 – 3.88 (m, 1H), 3.53 – 3.44 (m, 3H), 3.14 – 3.05 (m, 1H), 2.92 – 2.80 (m, 1H), 2.05 – 1.85 (m, 6H), 1.73 – 1.62 (m, 1H), 1.55 – 1.40 (m, 2H).
CDK7_1102	¹ H NMR (400 MHz, DMSO) δ 8.62 – 8.51 (m, 1H), 7.84 – 7.43 (m, 3H), 4.07 – 3.99 (m, 1H), 3.98 – 3.89 (m, 1H), 3.26 – 3.09 (m, 2H), 2.99 – 2.87 (m, 1H), 2.62 – 2.52 (m, 2H), 1.99 – 1.88 (m, 1H), 1.79 – 1.66 (m, 1H), 1.59 – 1.46 (m, 2H), 1.37 – 1.18 (m, 4H).
CDK7_1103	¹ H NMR (400 MHz, DMSO) δ 12.69 – 12.27 (m, 1H), 8.63 – 8.43 (m, 1H), 8.34 – 8.24 (s, 1H), 7.79 – 7.69 (m, 1H), 7.63 and 7.53 (2s, 1H, rotamers), 7.30 – 7.20 (m, 1H), 4.07 – 3.82 (m, 1H), 3.19 – 3.02 (m, 2H), 2.97 – 2.80 (m, 2H), 2.43 – 2.35 (m, 1H), 1.98 – 1.86 (m, 1H), 1.77 – 1.61 (m, 1H), 1.59 – 1.39 (m, 2H), 1.32 – 1.12 (m, 4H).
CDK7_1124	¹ H NMR (400 MHz, DMSO) δ 12.97 – 11.91 (m, 1H), 8.63 – 8.46 (m, 1H), 8.28 (s, 1H), 7.80 – 7.70 (m, 1H), 7.67 – 7.52 (m, 3H), 7.32 (s, 1H), 7.30 – 7.18 (m, 2H), 4.09 – 3.90 (m, 4H), 3.26 – 3.08 (m, 1H), 2.99 – 2.84 (m, 1H), 2.61 – 2.53 (m, 2H), 2.03 – 1.86 (m, 1H), 1.80 – 1.64 (m, 1H), 1.63 – 1.41 (m, 2H).
CDK7_1154	¹ H NMR (400 MHz, DMSO) δ 11.57 – 11.39 (m, 1H), 7.73 – 7.63 (m, 1H), 7.14 – 7.00 (m, 1H), 5.06 – 4.87 (m, 1H), 3.76 (br.s, 1H), 2.94 – 2.78 (m, 3H), 2.65 – 2.56 (m, 1H), 1.95 – 1.88 (m, 3H), 1.80 – 1.43 (m, 12H), 1.36 – 1.18 (m, 2H), 1.07 – 0.97 (m, 6H).
CDK7_1155	¹ H NMR (400 MHz, DMSO) δ 11.56 – 11.36 (m, 1H), 8.59 – 8.44 (m, 1H), 7.74 – 7.61 (m, 1H), 7.13 – 6.98 (m, 1H), 3.82 – 3.68 (m, 1H), 3.48 – 3.39 (m, 3H), 2.95 – 2.80 (m, 3H), 2.66 – 2.56 (m, 1H), 2.26 – 2.15 (m, 1H), 1.99 – 1.92 (m, 2H), 1.81 – 1.42 (m, 10H), 1.35 – 1.16 (m, 3H), 1.08 – 0.98 (m, 6H).
CDK7_1172	¹ H NMR (400 MHz, DMSO) δ 12.14 – 11.98 (m, 1H), 8.66 – 8.57 (m, 1H), 7.86 – 7.79 (m, 1H), 7.59 – 7.53 (m, 1H), 7.33 – 7.21 (m, 2H), 4.64 – 4.44 (m, 2H), 3.90 – 3.78 (m, 1H), 3.08 – 2.93 (m, 3H), 2.87 – 2.75 (m, 1H), 2.46 – 2.36 (m, 2H), 2.20 – 1.96 (m, 2H), 1.91 – 1.81 (m, 1H), 1.71 – 1.57 (m, 1H), 1.54 – 1.34 (m, 2H).
CDK7_1177	¹ H NMR (400 MHz, DMSO) δ 11.89 – 11.68 (m, 1H), 8.66 – 8.51 (m, 1H), 7.87 – 7.76 (m, 1H), 7.23 – 7.13 (m, 1H), 7.09 (s, 1H), 6.81 (s, 1H), 3.98 – 3.82 (m, 1H), 3.81 – 3.66 (m, 2H), 3.13 – 2.93 (m, 5H), 2.90 – 2.79

	(m, 2H), 2.17 – 2.03 (m, 2H), 1.97 – 1.83 (m, 1H), 1.77 – 1.57 (m, 1H), 1.56 – 1.37 (m, 2H), 1.19 – 1.03 (m, 2H).
CDK7_1201	¹ H NMR (400 MHz, DMSO) δ 11.58 – 11.38 (m, 1H), 8.66 – 8.44 (m, 1H), 7.86 – 7.69 (m, 1H), 7.17 – 6.96 (m, 1H), 3.98 – 3.75 (m, 1H), 3.53 – 3.41 (m, 2H), 3.09 – 3.00 (m, 3H), 2.99 – 2.83 (m, 3H), 2.57 – 2.52 (m, 1H), 2.02 – 1.82 (m, 3H), 1.72 – 1.56 (m, 2H), 1.31 – 1.17 (m, 1H), 0.52 – 0.42 (m, 1H), 0.41 – 0.32 (m, 2H), 0.31 – 0.23 (m, 1H).
CDK7_1225	¹ H NMR (400 MHz, DMSO) δ 11.60 – 11.40 (m, 1H), 8.61 – 8.43 (m, 1H), 7.76 – 7.62 (m, 1H), 7.16 – 6.94 (m, 1H), 3.75 (br.s, 1H), 3.47 – 3.38 (m, 2H), 2.89 – 2.73 (m, 4H), 2.65 – 2.56 (m, 1H), 1.94 – 1.83 (m, 1H), 1.81 – 1.70 (m, 1H), 1.68 – 1.55 (m, 1H), 1.53 – 1.42 (m, 1H), 1.37 – 1.19 (m, 2H), 1.06 – 0.99 (m, 6H), 0.81 – 0.72 (m, 2H), 0.69 – 0.57 (m, 2H).
CDK7_1226	¹ H NMR (400 MHz, DMSO) δ 11.59 – 11.40 (m, 1H), 8.58 – 8.47 (m, 1H), 7.74 – 7.62 (m, 1H), 7.12 – 7.01 (m, 1H), 5.91 – 5.75 (m, 1H), 5.15 – 5.06 (m, 1H), 5.05 – 4.98 (m, 1H), 3.76 (br.s, 1H), 3.59 – 3.51 (m, 2H), 3.49 – 3.41 (m, 2H), 2.94 – 2.78 (m, 3H), 2.66 – 2.56 (m, 1H), 2.36 – 2.26 (m, 2H), 2.00 – 1.91 (m, 2H), 1.81 – 1.71 (m, 1H), 1.70 – 1.56 (m, 1H), 1.54 – 1.45 (m, 1H), 1.37 – 1.20 (m, 1H), 1.09 – 0.99 (m, 6H).
CDK7_1227	¹ H NMR (400 MHz, DMSO) δ 11.60 – 11.41 (m, 1H), 8.61 – 8.41 (m, 1H), 7.79 – 7.64 (m, 1H), 7.16 – 7.00 (m, 1H), 3.90 – 3.81 (m, 1H), 3.55 – 3.47 (m, 3H), 2.97 – 2.63 (m, 5H), 2.05 – 1.90 (m, 2H), 1.81 – 1.45 (m, 3H), 1.42 – 1.27 (m, 1H), 1.12 – 0.94 (m, 7H), 0.51 – 0.37 (m, 2H), 0.32 – 0.17 (m, 2H).
CDK7_1229	¹ H NMR (400 MHz, DMSO) δ 11.59 – 11.40 (m, 1H), 8.58 – 8.46 (m, 1H), 7.74 – 7.62 (m, 1H), 7.12 – 7.00 (m, 1H), 4.91 – 4.82 (m, 1H), 3.82 – 3.71 (m, 1H), 2.97 – 2.79 (m, 4H), 2.66 – 2.53 (m, 2H), 1.93 – 1.86 (m, 1H), 1.81 – 1.71 (m, 1H), 1.69 – 1.58 (m, 1H), 1.53 – 1.45 (m, 1H), 1.38 – 1.20 (m, 2H), 1.16 – 1.08 (m, 6H), 1.07 – 0.99 (m, 6H).
CDK7_1254	¹ H NMR (400 MHz, DMSO) δ 11.54 – 11.53 (m, 1H), 8.61 – 8.47 (m, 1H), 7.82 – 7.70 (m, 1H), 7.14 – 6.98 (m, 1H), 3.97 – 3.84 (m, 1H), 3.81 – 3.72 (m, 1H), 3.12 – 3.04 (s, 4H), 2.98 – 2.80 (m, 3H), 2.49 – 2.43 (m, 2H), 2.16 – 2.05 (m, 1H), 1.97 – 1.82 (m, 2H), 1.75 – 1.62 (m, 1H), 1.53 – 1.36 (m, 2H), 1.19 – 1.06 (m, 3H).
CDK7_1259	¹ H NMR (400 MHz, DMSO) δ 11.60 – 11.36 (m, 1H), 8.57 – 8.48 (m, 1H), 7.73 – 7.65 (m, 1H), 7.10 – 7.02 (m, 1H), 3.76 (br.s, 1H), 3.54 – 3.47 (m, 2H), 3.46 – 3.39 (m, 2H), 2.93 – 2.80 (m, 3H), 2.67 – 2.52 (m, 2H), 1.99 – 1.91 (m, 1H), 1.80 – 1.70 (m, 1H), 1.68 – 1.57 (m, 1H), 1.53 – 1.44 (m, 1H), 1.36 – 1.20 (m, 1H), 1.13 – 0.99 (m, 9H).
CDK7_1268	¹ H NMR (400 MHz, DMSO) δ 11.58 – 11.42 (m, 1H), 8.59 – 8.49 (m, 1H), 7.79 – 7.66 (m, 1H), 7.14 – 6.98 (m, 1H), 3.96 – 3.84 (m, 1H), 3.67 – 3.61 (m, 2H), 3.50 – 3.46 (m, 3H), 3.27 (s, 3H), 3.02 – 2.91 and 2.76 – 2.63 (2m, 3H, rotamers), 2.90 – 2.81 (m, 2H), 1.99 – 1.88 (m, 2H), 1.84 – 1.74 (m, 1H), 1.74 – 1.51 (m, 2H), 1.48 – 1.33 (m, 1H), 1.15 – 1.06 (m, 6H).
CDK7_1276	¹ H NMR (400 MHz, DMSO) δ 11.71 – 11.32 (m, 1H), 8.66 – 8.45 (m, 1H), 7.92 – 7.67 (m, 1H), 7.14 – 7.00 (m, 1H), 3.93 – 3.72 (m, 1H), 3.49 – 3.38 (m, 2H), 2.99 – 2.77 (m, 4H), 2.61 – 2.52 (m, 1H), 1.98 – 1.82 (m, 3H), 1.72 – 1.56 (m, 2H), 1.33 – 1.17 (m, 1H), 0.82 – 0.72 (m, 2H), 0.66 – 0.58 (m, 2H), 0.51 – 0.42 (m, 1H), 0.40 – 0.31 (m, 2H), 0.31 – 0.23 (m, 1H).
CDK7_1277	¹ H NMR (400 MHz, DMSO) δ 11.59 – 11.40 (m, 1H), 8.60 – 8.45 (m, 1H), 7.86 – 7.68 (m, 1H), 7.13 – 7.00 (m, 1H), 4.95 – 4.78 (m, 1H), 3.96 – 3.77 (m, 1H), 3.02 – 2.80 (m, 3H), 2.57 – 2.51 (m, 2H), 1.99 – 1.81 (m, 3H), 1.74 – 1.56

	(m, 2H), 1.33 – 1.19 (m, 2H), 1.15 – 1.06 (m, 6H), 0.51 – 0.42 (m, 1H), 0.40 – 0.32 (m, 2H), 0.31 – 0.23 (m, 1H).
--	---

Table 4. Analytical data for exemplary compounds

Code	ESI-MS [M+H]⁺	Code	ESI-MS [M+H]⁺
CDK7_535	416.2	CDK7_997	445.3
CDK7_536	383.1	CDK7_998	459.3
CDK7_537	355.2	CDK7_999	394.3
CDK7_538	431.2	CDK7_1000	394.1
CDK7_539	438.2	CDK7_1001	409.3
CDK7_548	354.1	CDK7_1003	452.2
CDK7_735	414.2	CDK7_991	383.2
CDK7_754	339.2	CDK7_924	474.2
CDK7_766	337.2	CDK7_955	475.2
CDK7_809	362.2	CDK7_1105	419.2
CDK7_810	363.2	CDK7_1058	462.1
CDK7_811	364.1	CDK7_1176	472.2
CDK7_829	394.2	CDK7_1187	437.2
CDK7_830	390.2	CDK7_1122	503.3
CDK7_834	438.2	CDK7_1123	487.3
CDK7_835	435.2	CDK7_1021	423.1
CDK7_836	432.2	CDK7_1052	486.1
CDK7_838	516.2	CDK7_1060	394.2
CDK7_845	393.2	CDK7_1061	389.1
CDK7_846	369.2	CDK7_1062	390.2
CDK7_847	437.2	CDK7_1121	463.2
CDK7_848	461.2	CDK7_1103	420.1
CDK7_849	425.2	CDK7_1124	442.1
CDK7_850	439.2	CDK7_533	387.9
CDK7_851	453.3	CDK7_814	381.1
CDK7_888	414.2	CDK7_843	395.2
CDK7_891	417.2	CDK7_1019	397.2
CDK7_897	415.2	CDK7_1020	384.2
CDK7_898	353.2 (-Ac)	CDK7_1188	435.2
CDK7_901	390.2	CDK7_1199	485.2
CDK7_903	428.3	CDK7_1102	420.1
CDK7_904	428.3	CDK7_1154	491.3
CDK7_921	390.3	CDK7_1155	505.2
CDK7_929	407.2	CDK7_1172	478.1
CDK7_944	381.2	CDK7_1177	475.2
CDK7_951	395.2	CDK7_1201	435.3
CDK7_953	409.2	CDK7_1225	463.2
CDK7_959	459.2	CDK7_1226	477.2
CDK7_966	384.2	CDK7_1227	477.2
CDK7_1017	397.2	CDK7_1229	465.3
CDK7_550	353.2	CDK7_1254	423.2
CDK7_833	421.4	CDK7_1259	451.3
CDK7_837	456.2	CDK7_1268	481.3
CDK7_972	312.2	CDK7_1276	461.2
CDK7_900	393.2	CDK7_1277	463.2

Example 48. Inhibiting enzyme activity of CDK7-CycH, CDK2-CycA, CDK9-CycK, CDK13-CycK.

Measurements were carried out in a 5 µl reaction volume using a 384-well plate (Corning Inc., order no. 4513). The analyzed substances were titrated from 50 µM in increments of 4 at 11 concentration points in a normal strength kinase buffer A (Thermo Fisher Scientific Inc, cat. no. PV3189) supplemented with DTT (Sigma, orde. no. 646563-10X.5ML) and prepared with water (Panreac, order no. 701074). Kinase was diluted in an analogous buffer, mixed with test compounds in plate wells and preincubated for 10 minutes at 25 °C. Following incubation, a mixture of reaction substrate (CDKtide (SignalChem, order no. C06-58) or Histone H1 (SignalChem, order no. H10-54N) or PDKtide (SignalChem, order no. P10-58) and ATP (SignalChem, order no.V915B) prepared on a normal strength kinase buffer A was added. 1% solution of dimethyl sulfoxide (DMSO) (Sigma-Aldrich, order no. D2438) in the reaction volume was used as a negative control. The final concentrations of all reaction components are shown in **Table 5**. The plate was centrifuged at 400 rcf (Eppendorf, 5804 R) for 1 minute and incubated for 40-60 minutes at 25 °C. The amount of ATP consumed in the kinase reaction was then measured using the ADP-Glo™ detection system. Luminescence signal was measured using Spark 20M tablet multifunction reader (Tecan Group Ltd, Switzerland). IC₅₀ value was calculated using Magellan 7.2 (Tecan Group Ltd, Switzerland) by approximating experimental points by four-parameter model with the optimization by Levenberg-Marquardt:

$$Y = D + \frac{A - D}{1 + \left(\frac{X}{C}\right)^B},$$

where A is the upper asymptote; D is the lower asymptote; C is IC₅₀, half-maximal inhibitory concentration of kinase, nM; B is the curvature (slope).

Table 5. Final concentrations of kinase reaction components.

Active kinase	CDK7/Cyclin H/MNAT1	CDK2/Cyclin A	CDK9/Cyclin K	CDK13/Cyclin K
	2.5 ng/µl	0.225 ng/µl	2.5 ng/µl	5 ng/µl
Substrate	CDKtide	Histone H1	PDKtide	CDKtide
	0.05 mg/ml	0.1 mg/ml	0.2 mg/ml	0.2 mg/ml
ATP	10 µM	50 µM	50 µM	10 µM
DMSO	-	1%	1%	1%
DTT	2000 µM	50 µM	50 µM	50 µM

Table 6. Results of biochemical test of inhibition of CDK7/Cyclin H/MNAT1, CDK2/Cyclin A, CDK9/Cyclin K kinase activities. IC₅₀ is shown as a mean value from several measurements, "A" for IC₅₀<5nM, "B" for 5nM<IC₅₀<50nM, "C" for 50nM<IC₅₀<200nM, and "D" for IC₅₀>200nM, "N/A" means that a given compound was not analyzed in the present test.

Compound number	CDK7/Cyclin H/MNAT	CDK2/Cyclin A	CDK9/Cyclin K	CDK13/Cyclin K
CDK7_533	C	D	D	N/A
CDK7_536	A	D	B	N/A
CDK7_548	B	D	B	N/A
CDK7_538	B	D	C	N/A
CDK7_537	B	D	B	N/A
CDK7_539	B	D	C	N/A
CDK7_754	B	D	N/A	N/A
CDK7_550	C	D	N/A	N/A

CDK7_735	A	D	B	N/A
CDK7_535	B	D	C	N/A
CDK7_809	B	D	N/A	N/A
CDK7_810	B	D	N/A	N/A
CDK7_811	B	D	N/A	N/A
CDK7_845	A	D	B	N/A
CDK7_837	C	D	D	N/A
CDK7_766	C	D	N/A	N/A
CDK7_835	B	D	B	N/A
CDK7_846	B	D	B	N/A
CDK7_847	B	D	B	N/A
CDK7_848	B	D	B	N/A
CDK7_849	A	D	C	N/A
CDK7_850	B	D	C	N/A
CDK7_851	B	D	C	N/A
CDK7_897	N/A	D	N/A	N/A
CDK7_898	B	D	N/A	N/A
CDK7_951	A	D	B	N/A
CDK7_834	B	D	B	N/A
CDK7_838	B	D	C	N/A
CDK7_972	B	D	C	N/A
CDK7_961	A	D	C	N/A
CDK7_833	B	D	C	N/A
CDK7_836	B	D	C	N/A
CDK7_888	B	D	N/A	N/A
CDK7_929	A	D	B	N/A
CDK7_944	A	D	B	N/A
CDK7_991	D	N/A	N/A	N/A
CDK7_891	B	D	N/A	N/A
CDK7_921	B	N/A	N/A	N/A
CDK7_966	B	D	D	N/A
CDK7_903	B	D	D	N/A
CDK7_904	A	D	B	N/A
CDK7_1017	B	D	C	N/A
CDK7_959	A	D	C	N/A
CDK7_997	B	D	C	N/A
CDK7_998	B	D	D	N/A
CDK7_999	A	D	B	N/A
CDK7_953	A	D	C	B
CDK7_1000	A	D	B	B
CDK7_1001	A	D	B	N/A
CDK7_1003	B	D	C	N/A
CDK7_1021	B	B	D	N/A
CDK7_1052	A	D	D	N/A
CDK7_1060	A	D	B	N/A
CDK7_1061	B	N/A	B	N/A
CDK7_1062	A	D	B	N/A
CDK7_1105	A	D	B	N/A
CDK7_924	A	D	D	N/A
CDK7_955	B	D	D	N/A

CDK7_1122	B	N/A	D	N/A
CDK7_1124	B	D	D	N/A
CDK7_1123	A	D	D	B
CDK7_1176	B	N/A	N/A	N/A
CDK7_1187	A	D	D	C
CDK7_1020	A	D	B	N/A
CDK7_1034	A	D	B	N/A
CDK7_1019	B	D	D	N/A
CDK7_814	A	D	C	N/A
CDK7_843	B	D	D	N/A
CDK7_960	A	D	B	N/A
CDK7_1058	A	D	B	N/A
CDK7_1102	B	D	B	N/A
CDK7_1103	A	D	C	N/A
CDK7_524	A	D	D	N/A
CDK7_1154	A	D	D	N/A
CDK7_1155	B	D	D	N/A
CDK7_1172	A	D	D	N/A
CDK7_1177	B	D	D	N/A
CDK7_1188	A	D	D	N/A
CDK7_1199	A	D	D	N/A
CDK7_1201	A	D	C	C
CDK7_1225	A	D	D	D
CDK7_1226	A	D	D	N/A
CDK7_1227	A	D	D	N/A
CDK7_1229	A	D	D	D
CDK7_1236	A	D	C	N/A
CDK7_1237	N/A	N/A	N/A	N/A
CDK7_1254	A	D	D	N/A
CDK7_1259	A	D	D	N/A
CDK7_1268	N/A	D	D	N/A
CDK7_1276	A	D	D	C
CDK7_1277	A	D	D	D

Table 7. Results of biochemical test of inhibition of CDK7/Cyclin H/MNAT1, CDK2/Cyclin A, CDK9/Cyclin K, CDK13/Cyclin K kinase activities. IC₅₀ is shown as a mean value from the data of multiple measurement series:

Compound number	CDK7/Cyclin H/MNAT1	CDK2/Cyclin A	CDK9/Cyclin K	CDK13/Cyclin K
	IC ₅₀ , nM			
CDK7_924	<5	7796	431	N/A
CDK7_953	<5	5828	60	20
CDK7_961	<5	3698	69	N/A
CDK7_959	<5	5258	106	N/A
CDK7_1052	<5	12396	479	N/A
CDK7_1123	<5	35219	265	25
CDK7_1187	<5	15651	182	81
CDK7_1199	<5	3394	843	N/A
CDK7_1201	<5	4496	112	65
CDK7_1225	<5	13883	441	203
CDK7_1226	5.6	18646	760	N/A

CDK7_1229	<5	22938	485	242
CDK7_1276	<5	6224	296	152
CDK7_1277	<5	14316	442	425

The results of these experiments confirm production of high-affinity CDK7 inhibitors selective for CDK9, CDK13 and CDK2.

Example 49. Antiproliferative activity against sensitive cell lines *in vitro*.

The antiproliferative activity of CDK7 inhibitors according to the present invention was evaluated in a cell test using continuous cell cultures: MDA-MB-468 (*triple negative breast cancer*, ATCC® HTB-132™) and NCI-H82 (*small cell lung cancer*, ATCC® HTB-175™) using a viable AlamarBlue dye (Thermo Fisher Scientific Inc, order no. DAL1100). The cells were grown in DMEM (PanEco, order no. C330π) and RPMI-1640 (PanEco, order no. C330π), respectively. The basal growth medium was enriched with L-glutamine at a concentration of 2 mM (PanEco, order no. Φ032) and 10% fetal bovine serum (Gibco, order no. 16140-071). The night before measurement, MDA-MB-468 adherent cells were detached from plastic with a solution of trypsin-EDTA (PanEco, order no. Π039π), the solution was then inactivated by a DMEM complete medium, centrifuged at 1000 RPM (Eppendorf, 5804 R), a suspension of cells was prepared in a DMEM complete medium, introduced into 96-well culture plates (Corning Inc, order no. 3599) at 8×10^3 cells in 100 μl of medium per well. The next day, test compounds were dissolved in DMSO and diluted with a DMEM complete medium to a final concentration ranging from 50 μM to 10 pM. Diluted compounds in a volume of 50 μl were introduced into the plate wells with cells in triplicates (the final concentration of DMSO was no more than 1%) and incubated at 37 °C in an incubator with 5% CO₂ for 96 hours. NCI-H82 suspension cell line was prepared on the day of the analysis. 10×10^3 cells in 100 μl of an RPMI-1640 complete medium were added to each well of a 96-well plate. The procedure for sample dilution is as described above. Cells with test compounds were incubated at 37 °C in an incubator with 5% CO₂ for 120 hours. Following incubation, 15 ml of AlamarBlue reagent (Thermo Fisher Scientific Inc., order no. DAL1100) was added to the wells, the contents of the plates were stirred in an orbital shaker (Biosan, Latvia), then further incubated for 5.5–6 hours at 37 °C in an incubator with 5% CO₂. The number of viable cells was detected using the Infinite M200Pro microplate spectrophotometer (Tecan Group Ltd, Switzerland) by way of measuring a fluorescent signal at an excitation wavelength (λEx) of 540 nm and an emission wavelength (λEm) of 590 nm.

IC₅₀ value was calculated using the Magellan software (Tecan Group Ltd, Switzerland) by approximating experimental points by four-parameter model with the optimization by Levenberg-Marquardt:

Table 8. Results of antiproliferative assay on MDA-MB-468 and NCI-H82 cell lines. IC₅₀ is shown as a mean value from several measurements, "A" for IC₅₀<10nM, "B" for 10nM<IC₅₀<100nM, "C" for 100nM<IC₅₀<1000nM, and "D" for IC₅₀>1000nM, "N/A" means that a given compound was not analyzed in the present test.

Compound number	MDA-MB-468	NCI-H82	Compound number	MDA-MB-468	NCI-H82
CDK7_533	C	N/A	CDK7_891	C	N/A
CDK7_536	B	N/A	CDK7_966	C	N/A
CDK7_548	B	N/A	CDK7_904	C	N/A
CDK7_538	C	N/A	CDK7_1017	C	D
CDK7_537	C	N/A	CDK7_959	A	A
CDK7_539	C	N/A	CDK7_924	A	B
CDK7_754	D	N/A	CDK7_953	A	A

CDK7_550	D	N/A	CDK7_1052	C	C
CDK7_735	C	N/A	CDK7_1060	B	B
CDK7_535	C	N/A	CDK7_1061	B	B
CDK7_809	D	N/A	CDK7_1062	B	B
CDK7_810	D	N/A	CDK7_1105	A	B
CDK7_811	C	N/A	CDK7_955	C	D
CDK7_845	A	A	CDK7_1122	C	N/A
CDK7_829	D	N/A	CDK7_1123	A	B
CDK7_830	D	N/A	CDK7_1176	C	N/A
CDK7_837	C	N/A	CDK7_1187	A	A
CDK7_766	D	N/A	CDK7_1020	B	B
CDK7_835	C	N/A	CDK7_1034	A	A
CDK7_846	B	N/A	CDK7_814	B	B
CDK7_847	B	N/A	CDK7_960	B	B
CDK7_848	D	N/A	CDK7_1058	A	A
CDK7_849	B	N/A	CDK7_1102	B	B
CDK7_850	C	N/A	CDK7_1103	B	B
CDK7_851	D	N/A	CDK7_524	A	A
CDK7_898	C	N/A	CDK7_1154	B	B
CDK7_951	B	N/A	CDK7_1155	B	B
CDK7_834	B	N/A	CDK7_1172	B	B
CDK7_838	C	C	CDK7_1188	A	A
CDK7_972	C	N/A	CDK7_1199	A	A
CDK7_997	C	N/A	CDK7_1201	A	A
CDK7_998	D	N/A	CDK7_1225	B	B
CDK7_999	B	C	CDK7_1226	B	B
CDK7_1000	B	B	CDK7_1227	B	B
CDK7_1001	C	N/A	CDK7_1229	B	B
CDK7_1003	C	N/A	CDK7_1236	B	B
CDK7_961	A	A	CDK7_1254	B	B
CDK7_833	C	N/A	CDK7_1259	B	B
CDK7_836	C	N/A	CDK7_1268	B	B
CDK7_888	D	N/A	CDK7_1276	A	A
CDK7_929	A	A	CDK7_1277	A	A
CDK7_944	A	N/A			

Results of these experiments confirm production of inhibitors capable of stopping the growth of triple-negative breast cancer cells and small cell lung cancer cells in nM concentrations.

Example 50. General cytotoxicity against primary HepaRG liver cells *in vitro*.

General cytotoxicity of CDK7 inhibitors according to the present invention was investigated using primary differentiated HepaRG cells (Biopredic Int, order no. HPR101). A week before the measurement, the cells were thawed and maintained in Basal hepatic cell medium (Biopredic Int, order no. MIL700079) supplemented with HepaRG maintenance/metabolism medium supplement with antibiotics (Biopredic Int, order no. ADD620C). The medium in vials with cells was changed every 2-3 days. The night before the test, the cells were plated in wells of 96-well plates (Corning Inc., order no. 3599) at 25×10^3 cells in 100 μ l of medium per well. The next day, test compounds were introduced into plate wells with cells, untitrated at a concentration ranging from 200 μ M to 0.2 nM, and incubated for 72 hours. Cell viability was assessed as described above.

Table 9. Results of general toxicity investigation on primary HepaRG cells. CC₅₀ is shown as a mean value from several measurements, "A" for CC₅₀>2000nM, "B" for

2000nM>CC₅₀>500nM, "C" for 500nM>CC₅₀>100nM, and "D" for CC₅₀<100nM, "N/A" means that a given compound was not analyzed in the present test.

Compound number	HepaRG	Compound number	HepaRG	Compound number	HepaRG
CDK7_533	B	CDK7_997	B	CDK7_1187	A
CDK7_536	B	CDK7_998	A	CDK7_829	A
CDK7_548	B	CDK7_999	B	CDK7_830	A
CDK7_538	B	CDK7_1000	C	CDK7_900	A
CDK7_537	C	CDK7_1001	B	CDK7_1020	A
CDK7_539	A	CDK7_1003	B	CDK7_1034	C
CDK7_754	A	CDK7_961	C	CDK7_1019	A
CDK7_550	A	CDK7_833	B	CDK7_814	B
CDK7_735	A	CDK7_836	B	CDK7_960	C
CDK7_535	A	CDK7_888	A	CDK7_1058	B
CDK7_809	A	CDK7_900	A	CDK7_1102	B
CDK7_810	A	CDK7_929	C	CDK7_1103	C
CDK7_811	B	CDK7_944	D	CDK7_1121	A
CDK7_845	C	CDK7_966	A	CDK7_524	A
CDK7_829	A	CDK7_904	A	CDK7_1154	A
CDK7_830	A	CDK7_1017	A	CDK7_1155	A
CDK7_837	B	CDK7_959	A	CDK7_1172	A
CDK7_766	A	CDK7_924	A	CDK7_1177	A
CDK7_835	A	CDK7_1021	A	CDK7_1188	A
CDK7_846	B	CDK7_1052	A	CDK7_1201	B
CDK7_847	C	CDK7_1060	C	CDK7_1225	A
CDK7_848	A	CDK7_1061	B	CDK7_1226	A
CDK7_849	B	CDK7_1062	C	CDK7_1227	A
CDK7_850	A	CDK7_953	B	CDK7_1229	A
CDK7_851	A	CDK7_1105	C	CDK7_1236	A
CDK7_897	A	CDK7_955	A	CDK7_1254	A
CDK7_898	A	CDK7_1121	A	CDK7_1259	A
CDK7_951	B	CDK7_1122	A	CDK7_1268	A
CDK7_834	C	CDK7_1123	A	CDK7_1276	A
CDK7_838	B	CDK7_1124	A	CDK7_1277	A
CDK7_972	B	CDK7_1176	A		

Results of these experiments confirm production of CDK7 inhibitors that are non-toxic in the human hepatocyte model.

Example 51. Determination of incorporation of bromodeoxyuridine into DNA

Upon adding bromodeoxyuridine (BrdU) to dividing cells, Thymidine is replaced by its analogue, BrdU, and BrdU is incorporated into DNA. Thus, BrdU incorporated into DNA characterizes the intensity of DNA replication and ongoing S phase of the cell cycle. Reduced incorporation of bromodeoxyuridine into DNA under the action of the compounds characterizes a slowdown in the cell cycle and DNA replication.

The ability of the CDK7 inhibitors according to the present invention to inhibit the incorporation of BrdU into DNA was evaluated in a cell test on continuous cell cultures: MDA-MB-468 (*triple negative breast cancer*, ATCC® HTB-132™) using a BrdU detection kit (Roche, order no. 11 669 915 001)). The cells were grown in DMEM medium (PanEco, order no. C330π). The basal growth medium was enriched with L-glutamine at a concentration of 2 mM (PanEco, order no. Φ032) and 10% fetal bovine serum (Gibco, order no. 16140-071). MDA-MB-468

adherent cells were detached from plastic with a solution of trypsin-EDTA (PanEco, order no. П039п), the solution was then inactivated by a DMEM complete medium, centrifuged at 1000 RPM (Eppendorf, 5804 R), a suspension of cells was prepared in a DMEM complete medium, introduced into 96-well culture plates (Corning Inc, order no. 3599) at 4×10^3 cells in 75 μ l of medium per well. Test compounds were dissolved in DMSO and diluted with a DMEM complete medium to a final concentration ranging from 0.5 μ M to 2 pM. Serum-free growth medium was removed from plate wells, and solutions comprising diluted compounds in the volume of 100 μ l in basal growth medium were added. Candidates were introduced into the plate wells with cells in triplicates (the final concentration of DMSO was no more than 1%) and incubated at 37 °C in an incubator with 5% CO₂ for 24 hours. Following incubation, 10 μ l of solution of BrdU at a concentration of 10 μ M was added to wells, the contents of the plates were agitated on an orbital shaker (Biosan, Latvia), and incubated for additional 72 hours at 37 °C in an incubator with 5% CO₂. Incorporated BrdU was then detected according to the guidelines of the kit manufacturer. BrdU-containing medium was removed from plate wells and the plate with cells was dried at 60 °C for 1 hour. To fix the cells, 200 μ l of FixDenat solution was introduced into each well and incubated for 90 minutes at room temperature (+15-25 °C). The fixing solution was removed and 100 μ l of anti-BrdU-POD working solution was introduced, followed by 90 minute incubation at room temperature (+15-25 °C). The anti-BrdU-POD antibody solution was removed from plate wells and the wells were washed 3 times with 200 μ l of washing solution (10 minutes at room temperature (+15-25 °C)). 100 μ l of substrate solution was introduced into each analyzed well and incubated for 3 minutes at room temperature (+15-25 °C) on an orbital shaker (Biosan, Latvia) and for another 5-7 minutes without stirring.

The amount of incorporated BrdU was detected on the Infinite M200Pro microplate spectrophotometer (Tecan Group Ltd, Switzerland) by measuring the luminescent signal. IC₅₀ value was calculated using the Magellan software (Tecan Group Ltd, Switzerland) by approximating experimental points by four-parameter model with the optimization by Levenberg-Marquardt:

Table 10. Results of BrdU incorporation into DNA on MDA-MB-468 cell lines. IC₅₀ is shown as a mean value from the data of multiple measurements, IC₅₀ , nM.

Compound number	MDA-MB-468, IC ₅₀ , nM
CDK7_1187	14
CDK7_1225	24
CDK7_1229	28
CDK7_1276	2

The results of these experiments confirm production of CDK7 inhibitors, the antiproliferative activity of which on the MDA-MB-468 model (**Example 49**) is caused by a stopped cell cycle and slowed-down DNA replication.

Example 52. Determination of passive membrane permeability through PAMPA artificial membrane.

Passive membrane permeability of the compounds described in the present invention was determined using the PAMPA (parallel artificial membrane permeability assay) model at pH 7.4. We used a 96-well plate (Corning® BioCoat® Pre-coated PAMPA Plate System, order no. 353015), composed of acceptor and donor parts with cell membrane-modeling PVDF filter pre-impregnated with a lecithin-phospholipid bilayer. The substance (at a concentration of 10 μ mol) in a 0.01 M phosphate buffer with pH 7.4 was introduced into a donor well of the plate, and pure buffer was introduced into an acceptor well; afterwards, the both parts were combined and

incubated for 20 hours at room temperature so that the compound was redistributed between the two wells and the membrane. Verapamil (USP, USA, order no. 1711202) and propranolol (MCE LLC, USA, cat. no. HY-B0573/CS-2680) were used as control compounds with high and medium permeability. Samples were analyzed by reversed-phase HPLC with UV detection (Agilent 1260 chromatograph with UV detector) by determining their peak areas in the donor and acceptor wells. Permeability (P_e , cm/s) and retention factor (R , %) in the lipid bilayer were calculated using the formula:

$$P_e = (-\ln[1 - C_A(t)/C_{\text{equilibrium}}]) / (A \times (1/V_D + 1/V_A) \times t)$$

$$R = 1 - [C_D(t) \times V_D + C_A(t) \times V_A] / (C_0 + V_D), \text{ where}$$

C_0 is donor-well concentration of the substance at the initial moment of time t (μM);

$C_A(t)$ is acceptor-well concentration of the substance at time t (μM);

V_d is donor-well volume (0.3 ml);

V_a is acceptor-well volume (0.15 ml).

$$C_{\text{equilibrium}} = [C_D(t) \times V_D + C_A(t) \times V_A] / (V_D + V_A)$$

A is filter area (0.3 cm^2);

t is incubation time (72000 s);

Table 11. Results of investigation of passive permeability through PAMPA lipid membrane model. P_e is shown as a mean value from multiple measurements, "A" for $P_e > 1 \cdot 10^{-5} \text{ cm/s}$, "B" for $10^{-5} \text{ cm/s} > P_e > 5 \cdot 10^{-6} \text{ cm/s}$, "C" for $5 \cdot 10^{-6} \text{ cm/s} > P_e > 1 \cdot 10^{-6} \text{ cm/s}$, and "D" for $P_e < 1 \cdot 10^{-6} \text{ cm/s}$, and "N/A" means that a given compound was not analyzed in the present test.

Compound number	PAMPA	Compound number	PAMPA	Compound number	PAMPA
CDK7_597	C	CDK7_904	D	CDK7_1176	C
CDK7_624	D	CDK7_1017	C	CDK7_524	C
CDK7_478	C	CDK7_1034	C	CDK7_1154	B
CDK7_807	B	CDK7_959	D	CDK7_1155	B
CDK7_524	C	CDK7_1000	D	CDK7_1172	C
CDK7_533	D	CDK7_1060	B	CDK7_1187	C
CDK7_536	C	CDK7_814	D	CDK7_1188	C
CDK7_548	C	CDK7_960	D	CDK7_1201	B
CDK7_735	D	CDK7_953	C	CDK7_1225	C
CDK7_811	B	CDK7_1021	D	CDK7_1226	C
CDK7_845	D	CDK7_1105	D	CDK7_1227	B
CDK7_835	D	CDK7_1058	C	CDK7_1229	B
CDK7_951	D	CDK7_1102	C	CDK7_1254	C
CDK7_961	C	CDK7_1121	D	CDK7_1259	C
CDK7_929	D	CDK7_1061	C	CDK7_1276	B
CDK7_944	D	CDK7_955	D	CDK7_1277	B
CDK7_999	D	CDK7_1122	C		
CDK7_903	D	CDK7_1123	D		

Results of these experiments confirm production of CDK7 inhibitors capable of passively penetrating a lipid membrane.

Example 53. *In vitro* microsomal stability study of compounds in human, rat and mouse microsomes.

The procedure for investigating the metabolism of selected compounds was carried out in microsomal fractions of the liver (of rats, mice, humans) in the presence of a phase I metabolism cofactor (NADPH).

The rate of enzymatic decomposition of the compound in liver microsomes in *in vitro* analysis was determined by keeping a reaction mixture comprising 0.5 mg/ml of pooled human, rat or mouse liver microsomes, 1 µM test compound, 1.5 mM β-nicotinamide adenine dinucleotide (Carbosynth, UK, cat. no. NN108711101) and 1.5 mM magnesium chloride in 0.1 M sodium phosphate buffer (pH 7.4) in a solid-state thermostat at 37 °C (XenoTech, USA, order no. H2610, cat. no. R1000, cat. no. M1000). Incubation was carried out in three replicates.

To determine the stability of compounds under sample preparation conditions, an aliquot in two replicates with the test compound and a microsomal fraction devoid of the NADPH cofactor, which was incubated for 40 minutes at 37 °C.

Reactions were stopped at 0, 10, 20, 30 and 40 min time points by way of adding an ice mixture of acetonitrile:methanol (3:1) comprising an internal standard.

Samples were then centrifuged to remove the precipitated protein, supernatants were analyzed using HPLC-MS/MS to determine *in vitro* clearance (CL_{int}) of the compounds. The Agilent 6410 mass spectrometer (Agilent, USA) combined with the Agilent 1260 Infinity II HPLC liquid chromatograph (Agilent, USA) and a cooling autosampler controlled by MassHunter software (Agilent, USA) was used as an HPLC-MS/MS system. Chromatographic analysis was performed on a C18 reversed-phase HPLC column (Agilent, Waters or equivalent) in a gradient elution mode in a system of mobile phases: 0.1% formic acid in water / 0.1% formic acid in acetonitrile, flow rate: 0.5 ml/min, column T = 35°C. Mass spectrometric detection was performed with electrospray ionization in multiple reaction monitoring mode. Precursor ions, fragment ions and collision energies for each substance were selected experimentally.

A graph of dependence of the logarithmic ratio between substance's peak area and internal standard's peak area vs time was built. The dependent coefficient of this line corresponded to the elimination rate constant k , on the basis of which the *in vitro* clearance CL_{int} was calculated:

Константа скорости элиминации(k) = (–градиент)

$$\text{Период полураспада } (t_{1/2}) (\text{мин}) = \frac{0,693}{k}$$

$$V (\text{мкл/мг}) = \frac{\text{объем инкубирования (мкл)}}{\text{белок в инкубации (мг)}}$$

$$\text{внутренний клиренс } (CL_{int}) (\text{мкл} \cdot \text{мин}^{-1} \cdot \text{мг белка}^{-1}) = \frac{V \times 0,693}{t_{1/2}}$$

Английская версия формул:

$$\text{Elimination rate constant } (k) = (-\text{gradient}) \quad \text{Half life } (t_{1/2}) (\text{min}) = \frac{0.693}{k}$$

$$V (\mu\text{L/mg}) = \frac{\text{volume of incubation } (\mu\text{L})}{\text{protein in the incubation (mg)}} \quad \text{Intrinsic Clearance } (CL_{int}) (\mu\text{L/min/mg protein}) = \frac{V \times 0.693}{t_{1/2}}$$

Based on resulting *in vitro* clearance CL_{int} values, the microsomal stability of the selected compounds for humans, rats and mice was concluded.

Results are shown in **Tables 12** and **13**.

Table 12. Results of microsomal stability of selected compounds *in vitro* in humans (HLM). CL_{int} is shown as a mean value from multiple measurements, "A" for $CL_{int} < 20 \mu\text{L}/(\text{min} \cdot \text{mg of protein})$, "B" for $50 \mu\text{L}/(\text{min} \cdot \text{mg of protein}) > CL_{int} > 20 \mu\text{L}/(\text{min} \cdot \text{mg of protein})$, "C" for 100

$\mu\text{l}/(\text{min}\cdot\text{mg of protein}) > \text{Clint} > 50 \mu\text{l}/(\text{min}\cdot\text{mg of protein})$, and "D" for $\text{Clint} > 100 \mu\text{l}/(\text{min}\cdot\text{mg of protein})$).

Compound number	<i>In vitro</i> clearance CLint (HLM)	Compound number	<i>In vitro</i> clearance CLint (HLM)
CDK7_961	A	CDK7_524	A
CDK7_959	A	CDK7_1172	B
CDK7_924	A	CDK7_1177	A
CDK7_953	A	CDK7_1188	A
CDK7_955	A	CDK7_1225	A
CDK7_1122	A	CDK7_1229	B
CDK7_1123	A	CDK7_1254	A
CDK7_1176	A	CDK7_1276	B
CDK7_1187	B	CDK7_1277	B
CDK7_1103	A		

Table 13. Stability results of selected compounds *in vitro* in rat (RLM) and mouse (MLM) microsomes. Clint is shown as a mean value from multiple measurements, "A" for $\text{Clint} < 20 \mu\text{l}/(\text{min}\cdot\text{mg of protein})$, "B" for $50 \mu\text{l}/(\text{min}\cdot\text{mg of protein}) > \text{Clint} > 20 \mu\text{l}/(\text{min}\cdot\text{mg of protein})$, "C" for $100 \mu\text{l}/(\text{min}\cdot\text{mg of protein}) > \text{Clint} > 50 \mu\text{l}/(\text{min}\cdot\text{mg of protein})$, and "D" for $\text{Clint} > 100 \mu\text{l}/(\text{min}\cdot\text{mg of protein})$, and "N/A" means that a given compound was not analyzed in the present test.

Compound number	RLM	MLM
CDK7_953	A	A
CDK7_1187	A	A
CDK7_1188	A	N/A
CDK7_1225	A	A
CDK7_1227	A	N/A
CDK7_1229	A	A
CDK7_1259	A	N/A
CDK7_1276	B	B
CDK7_1277	B	B

Results of these experiments confirm production of CDK7 inhibitors resistant (B) and highly resistant (A) to the activity of human, mouse and rat liver microsomal enzymes.

Example 54. Pharmacokinetics (PK) study.

Dosing and administration

Pharmacokinetic studies of the compounds were carried out on CD-1 mice, males, aged about 8 weeks, weighing 16-29 g. The animals were given a single oral (PO) administration (intragastrically) at a dose of 10 mg/kg in a formulation in the form of a 0.2% solution of hydroxypropyl methylcellulose (HPMC) in a 5% aqueous solution of glucose and a single IV-bolus administration (intravenously) at a dose of 1.0 mg/kg for 1277 and at a dose of 2.0 mg/kg for substances **CDK7_1187**, **CDK7_1225**, **CDK7_1229** and **CDK7_1276** in a formulation comprising 0.2% DMSO in 5% aqueous solution of glucose.

For each substance, 21-30 animals were used for each dose of PO and IV administration. All mice were weighed and randomized before dosing.

3-5 animals were used for each time point. Each animal was sampled at one time point. At least 12 h before the start of the experiment, the animals were deprived of food while maintaining free access to water.

Animals from the PO and IV groups were subjected to blood sampling before administering the substance and after 0.5, 2, 4, 8, 12 and 24 h.

Blood samples

Blood was collected by cardiac puncture using the closed method. Blood was collected in a volume of 0.2 ml into polypropylene tubes comprising 20 μ l of 5% EDTA. Blood plasma was separated by centrifugation at 10000 rpm for 10 minutes at 4 ± 2 °C and transferred into clean labeled tubes.

Blood plasma analysis

Test blood plasma samples of mice from the experimental groups were obtained in a volume of 50 μ l. Afterwards, 200 μ l of a solution of internal standard (tolbutamide (Sigma, T0891)) prepared in acetonitrile, at a concentration of 50 ng/ml, pre-cooled in the refrigerator at 2-8 °C, was added to the samples, stirred in a vortex mixer (G560E, Scientific Industries) for 10 seconds and incubated at 4 °C for 15 minutes for protein precipitation. Samples were then centrifuged (5417R centrifuge, Eppendorf) at 1500 g for 10 minutes while cooling to 4 °C. 150 μ l of the supernatant were transferred to a 96-well plate for HPLC-MS/MS analysis. Quantitative determination of test compounds in blood plasma was carried out by absolute calibration method. The calibration curve was generated by analyzing plasma samples with supplementations of known amounts of the standard of the compound being determined. The lower limit of the quantitative determination of test compounds in blood plasma was 1 ng/ml, the analytical range of the technique was 1 ng/ml to 2000 ng/ml. The technique was partially validated in terms of selectivity, accuracy, precision, linearity, matrix effects, stability.

The Qtrap 5500 mass spectrometer (AB Sciex) in combination with the Agilent 1290 Infinity II HPLC liquid chromatograph (Agilent) and a cooling multisampler (Agilent 1290 Infinity II Multisampler G7167B) was used as an HPLC-MS/MS system. Chromatographic analysis was performed on a C18 reversed-phase HPLC column (YMC-Triart C18, 50*2 mm, 1.9 μ m, YMC CO. Ltd). The mobile phase used was used a mixture of deionized water and acetonitrile supplemented with 0.1% formic acid in gradient mode, flow rate: 0.5 ml/min, column T = 40°C, autosampler T = 8°C, analysis time: 2.8 min. Mass spectrometric detection was performed in multiple reaction monitoring (MRM) mode. Precursor ions, fragment ions and collision energies for each substance were selected experimentally. Two Q1/Q3 transitions were used to record MRM chromatograms of the analyte and internal standard; one of transitions was used for quantitative calculations, the other one was used to confirm that the extraction time corresponds to a given substance. Based on the results of HPLC-MS/MS analysis of calibration samples using the Analyst 1.6.3 software, the calibration dependence of areas of chromatographic analyte peaks normalized to the signal of the internal standard on the concentration of the analyte was built.

Pharmacokinetic analysis

Pharmacokinetic analysis was performed by non-compartmental method using the Phoenix WinNonlin 6.3 software package. Main averaged PK parameters were determined: time to maximum concentration (T_{max}), elimination half-life (T_{1/2}), maximum concentration (C_{max}), area under pharmacokinetic curve (AUC_{last}) from administration time to the last measured concentration, clearance (CL), steady-state volume of distribution (V_{ss}), and also absolute systemic bioavailability value (F). The pharmacokinetic parameters of the compounds according to the present invention are shown in **Tables 14-15**.

Table 14. Pharmacokinetic parameters of compounds in mice plasma following intravenous administration.

Intravenous administration						
PK parameter	Unit	CDK7_1187	CDK7_1225	CDK7_1229	CDK7_1276	CDK7_1277
dose	mg/kg	2	2	2	2	1
Animals in group	pc	5	5	5	3	5
T1/2	h	1.7	1.8	2.0	4.15	2.1
>AUClast	h×ng/ml	314	691	1135	1376	363
Vss	ml/kg	10101	6298	4166	6520	6889
Cl	ml/h/kg	6194	2878	1740	1288	2701

Table 15. Pharmacokinetic parameters of compounds in mice plasma following oral administration.

Oral administration						
Parameter	Unit	CDK7_1187	CDK7_1225	CDK7_1229	CDK7_1276	CDK7_1277
dose	mg/kg	10	10	10	10	10
Animals in group	pc	5	5	5	3	5
T1/2	h	1.7	1.9	2.5	2.0	3.8
Tmax	h	0.5	2.0	0.5	0.5	0.5
Cmax	ng/ml	309	712	1066	1513	893
AUClast	h×ng/ml	1164	3536	5787	7979	3113
F	%	72.7	102	102	116	85.8

The compounds according to the present invention correspond to target pharmacokinetic characteristics: the compounds are well absorbed and have a sufficient oral absorption effect. According to mean value of area under curve (AUClast), absolute bioavailability of the compounds was calculated to be more than 72.7%. In turn, this demonstrates the possibility of oral administration of these compounds to achieve pharmacodynamic effects.

Example 55. Permeability of blood-brain barrier (BBB)

The ability of substances to penetrate the blood-brain barrier and enter the central nervous system (mouse brain) was evaluated by the ratio of their areas under pharmacokinetic curves in rat brain and blood plasma following oral administration, characterizing tissue bioavailability to the rat brain, and expressed as the AUCbrain / AUCplasma coefficient.

Dosing and administration

Pharmacokinetic studies of the compounds were carried out on CD-1 mice, males, aged about 8 weeks, weighing 16-29 g. Animals were subjected a single oral (PO) administration (intragastrically) at a dose of 10 mg/kg in a formulation in the form of a 0.2% solution of hydroxypropyl methylcellulose (HPMC) in a 5% aqueous solution of glucose.

21 animals were used for each substance. All mice were weighed and randomized before dosing.

3 animals were used for each time point. Biomaterials were collected from each animal at one time point. At least 12 h before the start of the experiment, the animals were deprived of food while maintaining free access to water. Blood and brain samples were collected 0.5, 2, 4, 6, 8, 12 and 16 h after administration of the substance.

Blood samples

All manipulations were carried out analogously to **Example 54**.

Brain samples

Perfusion of mice brains was carried out with saline solution via the vascular system. To this end, the chest was opened, the right atrium was incised with microsurgical scissors, and saline

cooled to 4 °C was administered using a 10-ml syringe into the left ventricle until the liver turned white. After washing, the skull was opened and the brain removed, it was weighed and transferred to labeled 5 ml tubes.

Analysis of brain samples

A 4-fold volume of water was added to the brain suspension and homogenized using a homogenizer (Omni Bead Ruptor 24) 2 times for 45 seconds. Analogously, the intact mice brain homogenate was prepared, which was then used to prepare calibration samples and QC samples. The calibration curve was generated by analyzing brain homogenate samples with supplementations of known amounts of the standard of the compound being determined.

Samples were precipitated with cooled acetonitrile at 1:4 ratio, containing the internal standard tolbutamide (Sigma, T0891) at a concentration of 50 ng/ml. Samples were stirred on a vortex mixer (G560E, Scientific Industries) for 10 seconds and kept at 4 °C for 15 minutes to precipitate proteins, then centrifuged (5417R centrifuge, Eppendorf) for 10 minutes at 1500 g. 150 ml of supernatants were collected and transferred to corresponding wells of a 96-well plate for HPLC-MS/MS analysis. Quantitative determination of test compounds in brain homogenates was carried out analogously to the procedure described in **Example 54**. Lower limit of quantitative determination of test compounds in brain homogenate was 0.3 ng/ml.

Pharmacokinetic analysis of data was performed analogously to the procedure described in **Example 54**.

In addition to main PK parameters, the ratio of concentration in the brain and plasma at the studied time points ($AUC_{\text{brain}} / AUC_{\text{plasma}}$) was calculated to assess bioavailability in the brain (permeability through the blood-brain barrier).

Table 16. Pharmacokinetic parameters of compounds in mice brain homogenate following oral administration.

Parameter	Unit	CDK7_1225	CDK7_1229	CDK7_1276	CDK7_1277
dose	mg/kg	10	10	10	10
Animals in group	pc	3	3	3	3
t _{1/2}	h	2.33	2.25	1.99	2.76
T _{max}	h	6.0	0.5	0.5	2.0
C _{max}	ng/ml	93.8	136	335	226
AUC _{last}	h*ng/ml	855	909	1939	1750
AUC _{brain} / AUC _{plasma}	-	0.165	0.182	0.243	0.302

Resulting data demonstrate the ability of compounds according to the present invention to penetrate through the BBB

Example 56. Study of antitumor activity *in vivo*.

The antitumor activity of **CDK7_1187** *in vivo* was evaluated on a subcutaneous xenograft model of the NCI-H69 line (*small cell lung cancer*, ATCC® HTB-119™) transplanted to BALB/c Nude mice.

NCI-H69 cell line was cultured in RPMI-1640 medium (PanEco, Russia) with enrichment of the basal growth medium with L-glutamine at a concentration of 2 mM (PanEco, Russia) and 10% fetal bovine serum (Gibco, Thermo Fisher Scientific, USA).

The study used 20 BALB/c Nude mice (females) obtained from the Federal State Budgetary Scientific Institution "Federal Research Center Institute of Cytology and Genetics, Siberian Branch of Russian Academy of Sciences" (ICG SB RAS).

BALB/c Nude mice were subcutaneously administered into the right lateral side of the body with 200 μ l of suspension containing 5×10^6 NCI-H69 tumor cells and Matrigel (Corning® Matrigel® Basement Membrane Matrix) at 1:1 ratio.

Linear dimensions of the tumor were determined using an electronic caliper. Volume of the tumor node was determined by the formula (1):

$$V = L \times W^2 / 2, \quad (1)$$

where L and W are large and small diameters of the tumor node, respectively.

The efficacy of test product was measured by the index of tumor growth inhibition (TGI, %) calculated according to the formula (2):

$$TGI(V) = \frac{V_c - V_p}{V_c} \times 100\%, \quad (2)$$

where V_c and V_p are mean volumes of the tumor node in the control group and the group of animals that received the test substance, respectively.

After achieving an average tumor volume of 130 mm^3 , animals were divided into 2 groups of 10 animals, volume of the tumor node was used as a randomization criterion. The day of the first administration of the product was designated as day 0.

Test substance **CDK7_1187** was administered intragastrically at a dose of 6 mg/kg daily, for 21 days. Before each administration, **CDK7_1187** was diluted in 5% solution of glucose (Solopharm, Russia) and 0.1% solution of DMSO (Sigma, USA). The control group received a 5% solution of glucose and a 0.1% solution of DMSO in an analogous volume.

Statistical analysis was carried out using the GraphPad Prism 9 software package. Mann-Whitney U test was employed to assess the significance of differences when comparing the volumes of tumor nodes. Differences were identified as significant if $p < 0.05$.

Dynamics of tumor growth is shown in **Fig 1**. TGI indexes, %, are shown in **Table 17**.


Table 17. TGI index, %

Group	Day 9	Day 13	Day 17	Day 21
CDK7_1187	10	36	40	48

Results of the study showed that following administering **CDK7_1187** to BALB/c Nude mice the tumor volume was statistically significantly less than that in the control group of animals on days 17 and 21 of the experiment. The index of tumor growth inhibition at the end of the observation period was 48%.

Claims

1. Compound of formula I:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof,

R₁ is -C(Hal)₃, -CH(Hal)₂, -CH₂Hal, -Hal, -H, -NO₂, -(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -O-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal, -S-(C₁-C₆)alkyl unsubstituted or substituted by one or several atoms of Hal;

R₂, R₃ are each independently H, -(C₁-C₆)alkyl;

or R₂, R₃ together form -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several -(C₁-C₆)alkyls;

L₁, L₂ are each independently a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

A is 6-10 membered aryl unsubstituted or substituted by one or several R₇; 6-10 membered aryl unsubstituted or substituted by one or several R_{7a}, fused with -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8a}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7b}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8b}; 6-10 membered aryl unsubstituted or substituted by one or several R_{7c}, fused with 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{8c}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{7d}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7e}, fused with (C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{8e}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{7f}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8f}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7g}; 5-6 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{7h}, fused with 4-10 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{8h};

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}, -P(O)(C₁-C₆ alkyl)₂, -S(O)₂NH₂, -S(O)₂C₁-C₆ alkyl, -(C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with

1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i} ; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j} , fused with phenyl unsubstituted or substituted by one or several R_{4k} ; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l} , fused with phenyl unsubstituted or substituted by one or several R_{4m} ;

R_{4a} , R_{4b} , R_{4c} , R_{4d} , R_{4e} , R_{4f} , R_{4g} , R_{4h} , R_{4i} , R_{4j} , R_{4k} , R_{4l} , R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl, R_5 , R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a} , R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;


R_7 , R_{7a} , R_{7b} , R_{7c} , R_{7d} , R_{7e} , R_{7f} , R_{7g} , R_{7h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;


R_{8a} , R_{8b} , R_{8c} , R_{8d} , R_{8e} , R_{8f} , R_{8h} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

2. The compound according to claim 1, wherein -A-L₂-R₄ is:







wherein k,m are each independently 0, 1,2 or 3,

X₁, X₂, X₃, X₄, X₅, X₆ are each independently C, N, CH, CH₂, NH, S or O;

Y₁, Y₂, Y₃, Y₄, Y₅ are each independently C, N or CH;

L₂ is a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, -(CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}-, -P(O)(C₁-C₆ alkyl)₂-, -S(O)₂NH₂-, -S(O)₂C₁-C₆ alkyl-, (C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k} 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl, R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,


or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;


R₇, R_{7b}, R_{7d}, R_{7f}, are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8b}, R_{8d}, R_{8f} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

3. The connection according to claims 1-2, wherein -A-L₂-R₄ is





wherein k,m are each independently 0, 1,2 or 3,

L₂ is a chemical bond, -NR₅-, -O-, -S-, -S(O)-, -S(O)₂-, -C(O)-, -C(O)O-, -C(S)-, -C(=NH)-, - (CR_{6a}R_{6b})₁₋₃-, -P(O)(CH₃)-, -O-(CH₂)₁₋₃-, -(CH₂)₁₋₃O-, -NR_{5a}-(CH₂)₁₋₃-, -(CH₂)₁₋₃NR_{5a}-, -C(O)-(CH₂)₁₋₃-, -(CH₂)₁₋₃-C(O)-, -C(O)-NH- or -NH-C(O)-;

R₄ is -H -Hal, (=O), -OH, -C(O)OH, -NR_{4a}R_{4b}-, -P(O)(C₁-C₆ alkyl)₂-, -S(O)₂NH₂-, -S(O)₂C₁-C₆ alkyl-, - (C₁-C₆)alkyl unsubstituted or substituted by one or several R_{4c}; -O-C₁-C₆ alkyl unsubstituted or

substituted by one or several R_{4d}; phenyl unsubstituted or substituted by one or several R_{4e}; 4-7 membered heterocyclyl with 1,2,3 or 4 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4f}; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several R_{4g}; -C₂-C₆ alkenyl unsubstituted or substituted by one or several R_{4h}; 5-6 membered heteroaryl with 1,2,3 or 4 atoms N, unsubstituted or substituted by one or several R_{4i}; 4-7 membered heterocyclyl with 1 or 2 heteroatoms selected from N, S or O, unsubstituted or substituted by one or several R_{4j}, fused with phenyl unsubstituted or substituted by one or several R_{4k}; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S, unsubstituted or substituted by one or several R_{4l}, fused with phenyl unsubstituted or substituted by one or several R_{4m};

R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O), -OH, -NH₂, -Hal, -CN, -C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl, -C₁-C₆ alkyl; -O-C₁-C₆ alkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -C₃-C₆ cycloalkyl unsubstituted or substituted by one or several radicals selected from -Hal, -OH, -NH₂, -CN, phenyl; -S(O)₂C₁-C₆ alkyl, -S(O)₂NH₂, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂, -P(O)(C₁-C₆ alkyl)₂, morpholinyl, thiazolyl, phenyl, R₅, R_{5a} are each independently -H, C₁-C₆ alkyl,

R_{6a}, R_{6b} are each independently -H, C₁-C₆ alkyl, -Hal,

or R_{6a} and R_{6b} together with the C atom they are attached to form C₃-C₆ cycloalkyl unsubstituted or unsubstituted by several -(C₁-C₆)alkyls;

R₇, R_{7b}, R_{7d}, R_{7f}, are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

R_{8b}, R_{8d}, R_{8f} are each independently -Hal, -OH, -NH₂, -CN, (=O), -(C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal; -O (C₁-C₆)alkyl unsubstituted or substituted by one or several -Hal;

Hal is an atom of F, Cl, Br, I.

4. The compounds according to claims 1-3, wherein L₁, L₂ are each independently a chemical bond, -C(O)-, -CH₂-, -CH(CH₃)-, -C(CH₃)₂-, -C(CH₂)₂-, -CF₂-, -S-, -S(O)-, -S(O)₂-, -O-, -NH-, -NCH₃-, -P(O)(CH₃)-, -C(O)O-, -C(O)NH- or -NHC(O)-.

5. The compounds according to claims 1-3, wherein R₁ is -H, -CF₃, -CCl₃, -CHF₂, -CH₂F, -CHCl₂, -CH₂Cl, -NO₂.

6. The compounds according to claims 1-3, wherein R₂, R₃ are each independently -H, methyl, ethyl, propyl, or R₂ and R₃ together with the C atom they are attached to form cyclopropyl, cyclobutyl or cyclopentyl.

7. The compounds according to claims 1-3, wherein R₄ is -H, (=O), -OH, -F, -Cl, -Br, -P(O)(C₁-C₆ alkyl)₂; -S(O)₂NH₂; -S(O)₂-(C₁-C₆)alkyl; -(C₁-C₆)alkyl unsubstituted or substituted by one or several (=O), -Hal, -CN, -S(O)₂-(C₁-C₆)alkyl; -O-(C₁-C₆)alkyl unsubstituted or substituted by one or several (=O), -CN, -Hal; -(C₃-C₆)cycloalkyl unsubstituted or substituted by one or several (=O), -OH, -Hal, -CN; phenyl unsubstituted or substituted by one or several -Hal, -CN, morpholine, -P(O)(CH₃)₂; 5-6 membered heterocyclyl with 1,2 or 3 heteroatoms selected from N or O unsubstituted or substituted by -(C₁-C₆)alkyl, (=O), -Hal, -CN; 5-6 membered heteroaryl with 1,2,3 or 4 heteroatoms selected from N,O or S unsubstituted or substituted by (=O), -(C₁-C₆)alkyl, -(C₃-C₆)cycloalkyl, -Hal, -CN; 5-6 membered heteroaryl with 1 or 2 atoms N, unsubstituted or substituted by one or several (=O), -(C₁-C₆)alkyl, fused with phenyl; -NR_{4a}R_{4b},

R_{4a}, R_{4b} are each independently -H, (=O), morpholinyl, -F, -S(O)₂CH₃, -CN, -one, cyclopropyl, thiazolyl, -C₁-C₃ alkyl unsubstituted or substituted with phenyl, -P(O)(CH₃)₂.

8. The compounds according to claim 7, wherein R₄ is -H, -one, (=O), -P(O)(CH₃)₂, -CH₃, -CH₂CH₃, tert-butyl, -CH₂CF₃, -CH₂CH₂S(O)₂CH₃, -CF₃, -Cl, -OCF₃, -S(O)₂NH₂, -S(O)₂CH₃, phenyl unsubstituted or substituted by one or several -Hal, -CN; methylpiperazinyl, piperazinyl, imidazolyl, thiazolyl, methylpyrazolyl, pyrazolyl, pyridinyl, pyrimidinyl, morpholine, oxanyl, oxolanyl, cyclohexyl, cyclopentyl, cyclopropyl, oxazolyl, methyloxazolyl, dimethyloxazolyl, dihydropyridinyl, methyl dihydropyridinyl, -NR_{4a}R_{4b},
R_{4a}, R_{4b} are each independently -H, (=O), morpholinyl, -F, -S(O)₂CH₃, -CN, -one, cyclopropyl, thiazolyl, -C₁-C₃ alkyl unsubstituted or substituted by phenyl, -P(O)(CH₃)₂.
9. The compounds according to claims 1-3,7, wherein R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f}, R_{4g}, R_{4h}, R_{4i}, R_{4j}, R_{4k}, R_{4l}, R_{4m} are each independently -H, (=O) -CH₃, morpholinyl, -F, -S(O)₂CH₃, -CN, -one, cyclopropyl, thiazolyl, -C₁-C₃ alkyl unsubstituted or substituted by phenyl, -P(O)(CH₃)₂.
10. The compounds according to claims 1-3, wherein R₅ is -H, -CH₃.
11. The compounds according to claims 1-3, wherein R_{6a} and R_{6b} are each independently -H, -F, -CH₃, or R_{6a} and R_{6b} together with the C atom they are attached to are cyclopropyl.
12. The compounds according to claims 1-3, wherein R₇, R_{7a}, R_{7b}, R_{7c}, R_{7d}, R_{7e}, R_{7f}, R_{7g}, R_{7h} are each independently (=O), -Hal, -CH₃; -OCH₃; -CHal₃; -OCHal₃.
13. The compounds according to claims 1-3, wherein R_{8a}, R_{8b}, R_{8c}, R_{8d}, R_{8e}, R_{8f}, R_{8g}, R_{8h} are each independently (=O), -Hal, -CH₃; -OCH₃; -CHal₃; -OCHal₃.
14. The compound according to claim 1, being:
4-(2-aminobenzoyl)-N-[(3S)-6,6-dimethylpiperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_461**);
4-(2-aminobenzoyl)-N-(6,6-dimethylpiperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_461_a**);
4-(2-aminobenzoyl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_549**);
4-(2-aminobenzoyl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_549_a**);
4-benzoyl-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_813**);
4-benzoyl-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_813_a**);
4-[(2-aminophenyl)methyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_540**);
4-[(2-aminophenyl)methyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_540_a**);
4-[1-(2-aminophenyl)ethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_541**);
4-[1-(2-aminophenyl)ethyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_541_a**);
4-[(1S)-1-(2-aminophenyl)ethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_541_b**);
4-[(1R)-1-(2-aminophenyl)ethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_541_c**);
4-[2-(2-aminophenyl)propan-2-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_542**);
4-[2-(2-aminophenyl)propan-2-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_542_a**);
4-[1-(2-aminophenyl)cyclopropyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_543**);

4-[1-(2-aminophenyl)cyclopropyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_543_a);

4-[(2-aminophenyl)difluoromethyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_544);

4-[(2-aminophenyl)difluoromethyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_544_a);

4-[(2-aminophenyl)sulfanyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_545);

4-[(2-aminophenyl)sulfanyl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_545_a);

4-(2-aminobenzenesulfinyl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_546);

4-(2-aminobenzenesulfinyl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_546_a);

4-[(R)-2-aminobenzenesulfinyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_546_b);

4-[(S)-2-aminobenzenesulfinyl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_546_c);

4-(2-aminobenzenesulfonyl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_547);

4-(2-aminobenzenesulfonyl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_547_a);

4-(2-aminophenoxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (CDK7_548);

4-(2-aminophenoxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (CDK7_548_a);

N4-(2-aminophenyl)-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine
(CDK7_550);

N4-(2-aminophenyl)-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine
(CDK7_550_a);

N4-(2-aminophenyl)-N4-methyl-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine
(CDK7_551);

N4-(2-aminophenyl)-N4-methyl-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine
(CDK7_551_a);

N4-phenyl-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (CDK7_812);

N4-phenyl-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (CDK7_812_a);

4-phenoxy-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (CDK7_754);

4-phenoxy-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (CDK7_754_a);

4-benzyl-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (CDK7_766);

4-benzyl-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (CDK7_766_a);

4-[(2-aminophenyl)(methyl)phosphoryl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_552);

4-[(2-aminophenyl)(methyl)phosphoryl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_552_a);

4-[(R)-(2-aminophenyl)(methyl)phosphoryl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_552_b);

4-[(S)-(2-aminophenyl)(methyl)phosphoryl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_552_c);

N4-[2-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_735**);
 N4-[2-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_735_a**);
 4-[2-(dimethylphosphoryl)phenoxy]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_897**);
 4-[2-(dimethylphosphoryl)phenoxy]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_897_a**);
 4-[3-(dimethylphosphoryl)phenoxy]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_892**);
 4-[3-(dimethylphosphoryl)phenoxy]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_892_a**);
 N4-[3-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_888**);
 N4-[3-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_888_a**);
 N4-[2-(dimethylphosphoryl)-5-methylphenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_903**);
 N4-[2-(dimethylphosphoryl)-5-methylphenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_903_a**);
 N4-[2-(dimethylphosphoryl)-4-methylphenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_904**);
 N4-[2-(dimethylphosphoryl)-4-methylphenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_904_a**);
 N4-[2-(dimethylphosphoryl)-5-(trifluoromethyl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_905**);
 N4-[2-(dimethylphosphoryl)-5-(trifluoromethyl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_905_a**);
 N4-[2-(dimethylphosphoryl)-4-(trifluoromethyl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_906**);
 N4-[2-(dimethylphosphoryl)-4-(trifluoromethyl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_906_a**);
 N4-[5-chloro-2-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_907**);
 N4-[5-chloro-2-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_907_a**);
 N4-[4-chloro-2-(dimethylphosphoryl)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_908**);
 N4-[4-chloro-2-(dimethylphosphoryl)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_908_a**);
 N4-[2-(dimethylphosphoryl)-4-(trifluoromethoxy)phenyl]-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_915**);
 N4-[2-(dimethylphosphoryl)-4-(trifluoromethoxy)phenyl]-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_915_a**);
 2-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]amino]benzene-1-sulfonamide (**CDK7_889**);

2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}amino)benzene-1-sulfonamide (**CDK7_889_a**);

N4-(3-methylsulfonylphenyl)-N2-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_890**);

N4-(3-methylsulfonylphenyl)-N2-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2,4-diamine (**CDK7_890_a**);

3-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]amino]benzene-1-sulfonamide (**CDK7_891**);

3-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}amino)benzene-1-sulfonamide (**CDK7_891_a**);

2-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]oxy]benzene-1-sulfonamide (**CDK7_893**);

2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}oxy)benzene-1-sulfonamide (**CDK7_893_a**);

4-(3-methylsulfonylphenoxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_894**);

4-(3-methylsulfonylphenoxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_894_a**);

3-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]oxy]benzene-1-sulfonamide (**CDK7_895**);

3-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}oxy)benzene-1-sulfonamide (**CDK7_895_a**);

4-(2,3-dihydro-1H-indol-7-yloxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_827**);

4-(2,3-dihydro-1H-indol-7-yloxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_827_a**);

4-(1H-indol-7-yloxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_828**);

4-(1H-indol-7-yloxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_828_a**);

4-(1,2,3,4-tetrahydroquinolin-8-yloxy)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_829**);

4-(1,2,3,4-tetrahydroquinolin-8-yloxy)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_829_a**);

N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)-4-(quinolin-8-yloxy)pyrimidin-2-amine (**CDK7_830**);

N-(piperidin-3-yl)-5-(trifluoromethyl)-4-(quinolin-8-yloxy)pyrimidin-2-amine (**CDK7_830_a**);

N-{2-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]amino]phenyl}acetamide (**CDK7_898**);

N-[2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}amino)phenyl]acetamide (**CDK7_898_a**);

N-{2-[(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl]oxy]phenyl}acetamide (**CDK7_899**);

N-[2-({2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}oxy)phenyl]acetamide (**CDK7_899_a**);

N-[(3S)-6,6-dimethylpiperidin-3-yl]-4-(5-phenyl-1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_460**);

N-(6,6-dimethylpiperidin-3-yl)-4-(5-phenyl-1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_460_a**);

N-[(3S)-piperidin-3-yl]-4-(5-phenyl-1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_533);
 N-(piperidin-3-yl)-4-(5-phenyl-1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_533_a);
 4-(4-phenyl-1H-pyrrol-3-yl)-N-[(3S)-piperidin-3-yl]-4-(4-phenyl-1H-pyrrol-3-yl)-5-
 (trifluoromethyl)pyrimidin-2-amine **(CDK7_534)**;
 4-(4-phenyl-1H-pyrrol-3-yl)-N-[piperidin-3-yl]-4-(4-phenyl-1H-pyrrol-3-yl)-5-
 (trifluoromethyl)pyrimidin-2-amine **(CDK7_534_a)**;
 4-(5-benzoyl-1H-pyrrol-3-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_535);
 4-(5-benzoyl-1H-pyrrol-3-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(CDK7_535_a);
 N,N-dimethyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-
 carboxamide **(CDK7_536)**;
 N,N-dimethyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-
 carboxamide **(CDK7_536_a)**;
 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide
(CDK7_537);
 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide
(CDK7_537_a);
 N-phenyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-
 carboxamide **(CDK7_538)**;
 N-phenyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-
 carboxamide **(CDK7_538_a)**;
 4-[5-(4-methylpiperazine-1-carbonyl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
 (trifluoromethyl)pyrimidin-2-amine **(CDK7_539)**;
 4-[5-(4-methylpiperazine-1-carbonyl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
 (trifluoromethyl)pyrimidin-2-amine **(CDK7_539_a)**;
 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxylic acid
(CDK7_922);
 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxylic acid
(CDK7_922_a);
 4-[4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-
 yl]benzonitrile **(CDK7_695)**;
 4-(4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-yl)benzonitrile
(CDK7_695_a);
 3-[4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-
 yl]benzonitrile **(CDK7_696)**;
 3-(4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-yl)benzonitrile
(CDK7_696_a);
 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-
 pyrrolo[2,3-c]pyridin-7-one **(CDK7_814)**;
 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[2,3-
 c]pyridin-7-one **(CDK7_814_a)**;
 3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-
 pyrrolo[3,2-c]pyridin-4-one **(CDK7_815)**;

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[3,2-c]pyridin-4-one (**CDK7_815_a**);

6-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_843**);

6-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_843_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-4-one (**CDK7_844**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-4-one (**CDK7_844_a**);

6-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_845**);

6-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_845_a**);

N-(1H-imidazol-2-yl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_833**);

N-(1H-imidazol-2-yl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_833_a**);

4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-N-(1,3-thiazol-2-yl)-1H-pyrrol-2-carboxamide (**CDK7_834**);

4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-N-(1,3-thiazol-2-yl)-1H-pyrrol-2-carboxamide (**CDK7_834_a**);

N-(1-methyl-1H-pyrazol-4-yl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_835**);

N-(1-methyl-1H-pyrazol-4-yl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_835_a**);

4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-N-(pyridin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_836**);

4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-N-(pyridin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_836_a**);

N-(4-cyanophenyl)-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_837**);

N-(4-cyanophenyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_837_a**);

N-[4-(morpholin-4-yl)phenyl]-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_838**);

N-[4-(morpholin-4-yl)phenyl]-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_838_a**);

N-methyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_846**);

N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_846_a**);

4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-N-(2,2,2-trifluoroethyl)-1H-pyrrol-2-carboxamide (**CDK7_847**);

4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-N-(2,2,2-trifluoroethyl)-1H-pyrrol-2-carboxamide (**CDK7_847_a**);

N-(2-methanesulfonylethyl)-4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrol-2-carboxamide (**CDK7_848**);
 N-(2-methanesulfonylethyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_848_a**);
 4-[5-(morpholin-4-carbonyl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_849**);
 4-[5-(morpholin-4-carbonyl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_849_a**);
 N-(oxan-4-yl)-4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrol-2-carboxamide (**CDK7_850**);
 N-(oxan-4-yl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_850_a**);
 N-(4-hydroxycyclohexyl)-4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrol-2-carboxamide (**CDK7_851**);
 N-(4-hydroxycyclohexyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_851_a**);
 N-(4-cyanocyclohexyl)-4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrol-2-carboxamide (**CDK7_852**);
 N-(4-cyanocyclohexyl)-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_852_a**);
 N-cyclopropyl-4-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H-pyrrol-2-carboxamide (**CDK7_951**);
 N-cyclopropyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_951_a**);
 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_923**);
 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_923_a**);
 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_924**);
 6-(3,5-dimethyl-1,2-oxazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_924_a**);
 3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_925**);
 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_925_a**);
 3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_926**);
 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_926_a**);
 3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_927**);
 3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_927_a**);
 7-methyl-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_928**);

7-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_928_a**);

7-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_929**);

7-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_929_a**);

7-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_930**);

7-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_930_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_931**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_931_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_932**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_932_a**);

8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_933**);

8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_933_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_934**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_934_a**);

8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_935**);

8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_935_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_936**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_936_a**);

8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_937**);

8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_937_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_938**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_938_a**);

8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_939**);

8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_939_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_940**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_940_a**);

8-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_941**);

8-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,8H,9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_941_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_942**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-9H-pyrrolo[2,3-c]azocin-9-one (**CDK7_942_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H-pyrrolo[2,3-c]pyrrol-6-one (**CDK7_943**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H-pyrrolo[2,3-c]pyrrol-6-one (**CDK7_943_a**);

5-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H-pyrrolo[2,3-c]pyrrol-6-one (**CDK7_944**);

5-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H-pyrrolo[2,3-c]pyrrol-6-one (**CDK7_944_a**);

N-[(3S)-piperidin-3-yl]-4-(1H-pyrazol-4-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_945**);

N-(piperidin-3-yl)-4-(1H-pyrazol-4-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_945_a**);

4-(1H-imidazol-4-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_946**);

4-(1H-imidazol-4-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_946_a**);

N-[(3S)-piperidin-3-yl]-4-(1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_947**);

N-(piperidin-3-yl)-4-(1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_947_a**);

N-[(3S)-piperidin-3-yl]-4-(1H-1,2,3-triazol-4-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_948**);

N-(piperidin-3-yl)-4-(1H-1,2,3-triazol-4-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_948_a**);

N-[(3S)-piperidin-3-yl]-4-(1H-1,2,4-triazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_949**);

N-(piperidin-3-yl)-4-(1H-1,2,4-triazol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_949_a**);

N-[(3S)-piperidin-3-yl]-4-(2H-1,2,3,4-tetrazol-5-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_950**);

N-(piperidin-3-yl)-4-(2H-1,2,3,4-tetrazol-5-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_950_a**);

4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,2,3,4-tetrahydroquinoxalin-2-one (**CDK7_900**);

4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,2,3,4-tetrahydroquinoxalin-2-one (**CDK7_900_a**);

4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,2-dihydroquinolin-2-one (**CDK7_901**);

4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,2-dihydroquinolin-2-one (**CDK7_901_a**);

4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,2-dihydroisoquinolin-1-one (**CDK7_921**);

4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,2-dihydroisoquinolin-1-one
(**CDK7_921_a**);

N-cyclopropyl-N-methyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-
pyrrol-2-carboxamide (**CDK7_1001**);

N-cyclopropyl-N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-
pyrrol-2-carboxamide (**CDK7_1001_a**);

N-methyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-N-(1,3-thiazol-2-
yl)-1H-pyrrol-2-carboxamide (**CDK7_1003**);

N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-N-(1,3-thiazol-2-yl)-
1H-pyrrol-2-carboxamide (**CDK7_1003_a**);

N-benzyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-
carboxamide (**CDK7_997**);

N-benzyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-
carboxamide (**CDK7_997_a**);

N-benzyl-N-methyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-
pyrrol-2-carboxamide (**CDK7_998**);

N-benzyl-N-methyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-
carboxamide (**CDK7_998_a**);

4-[5-(1-methyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_999**);

4-[5-(1-methyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_999_a**);

4-[5-(2-methyl-2H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1000**);

4-[5-(2-methyl-2H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1000_a**);

4-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1060**);

4-[5-(5-methyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1060_a**);

4-[5-(1-cyclopropyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1102**);

4-[5-(1-cyclopropyl-1H-1,2,3,4-tetrazol-5-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1102_a**);

4-[5-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1103**);

4-[5-(5-cyclopropyl-1,2,4-oxadiazol-3-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1103_a**);

4-[5-(1-methyl-1H-1,3-benzodiazol-2-yl)-1H-pyrrol-3-yl]-N-[(3S)-piperidin-3-yl]-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1124**);

4-[5-(1-methyl-1H-1,3-benzodiazol-2-yl)-1H-pyrrol-3-yl]-N-(piperidin-3-yl)-5-
(trifluoromethyl)pyrimidin-2-amine (**CDK7_1124_a**);

ethyl 4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-
carboxylate (**CDK7_1020**);

ethyl 4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxylate
(**CDK7_1020_a**);

N,N,1-trimethyl-4-(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-2-carboxamide (**CDK7_1017**);

N,N,1-trimethyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-2-carboxamide (**CDK7_1017_a**);

N-[(3S)-piperidin-3-yl]-4-[5-(pyridin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1061**);

N-(piperidin-3-yl)-4-[5-(pyridin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1061_a**);

N-[(3S)-piperidin-3-yl]-4-[5-(pyrimidin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1062**);

N-(piperidin-3-yl)-4-[5-(pyrimidin-2-yl)-1H-pyrrol-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1062_a**);

4-{5-[2-(dimethylphosphoryl)phenyl]-1H-pyrrol-3-yl}-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1057**);

4-{5-[2-(dimethylphosphoryl)phenyl]-1H-pyrrol-3-yl}-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1057_a**);

7-(1-methyl-1H-pyrazol-4-yl)-3-(2-[(3S)-piperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_955**);

7-(1-methyl-1H-pyrazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_955_a**);

3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1122**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1122_a**);

3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1-methyl-1H-pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1123**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(1-methyl-1H-pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1123_a**);

3-(2-[(6S)-4-azaspiro[2.5]octan-6-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1136**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1136_a**);

3-(2-[(7S)-5-azaspiro[3.5]nonan-7-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1137**);

3-[2-({5-azaspiro[3.5]nonan-7-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1137_a**);

3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-7-(1,3-thiazol-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1138**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(1,3-thiazol-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1138_a**);

7-(3,5-dimethyl-1,2-oxazol-4-yl)-3-(2-[(3S)-6,6-dimethylpiperidin-3-yl]amino)-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1139**);

7-(3,5-dimethyl-1,2-oxazol-4-yl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1139_a**);

5-[3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl]-1-methyl-1,2-dihydropyridin-2-one (**CDK7_1140**);

5-(3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl)-1-methyl-1,2-dihydropyridin-2-one (**CDK7_1140_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-(pyridin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1141**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1141_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-(pyridin-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1142**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1142_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-(pyridin-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1143**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1143_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-[(1-methyl-1H-pyrazol-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1144**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(1-methyl-1H-pyrazol-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1144_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-[(1,3-thiazol-2-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1145**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(1,3-thiazol-2-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1145_a**);

7-[(3,5-dimethyl-1,2-oxazol-4-yl)methyl]-3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1146**);

7-[(3,5-dimethyl-1,2-oxazol-4-yl)methyl]-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1146_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-[1-(1-methyl-1H-pyrazol-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1147**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(1-methyl-1H-pyrazol-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1147_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-[1-(1,3-thiazol-2-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1148**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(1,3-thiazol-2-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1148_a**);

7-[1-(3,5-dimethyl-1,2-oxazol-4-yl)ethyl]-3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1149**);

7-[1-(3,5-dimethyl-1,2-oxazol-4-yl)ethyl]-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1149_a**);

3-(2-[[[(3S)-6,6-dimethylpiperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-7-(oxan-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1150**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxan-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1150_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(oxan-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1151**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxan-4-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1151_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(oxolan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1152**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxolan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1152_a**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxolan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1153**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxolan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1153_a**);

7-cyclopentyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1154**);

7-cyclopentyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1154_a**);

7-(cyclopentylmethyl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1155**);

7-(cyclopentylmethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1155_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[1-(oxan-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1156**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(oxanyl-4-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1156_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[1-(oxolan-3-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1157**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[1-(oxolan-3-yl)ethyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1157_a**);

7-(1-cyclopentylethyl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1158**);

7-(1-cyclopentylethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1158_a**);

7-tert-butyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1159**);

7-tert-butyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1159_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2,2-dimethylpropyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1160**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2,2-dimethylpropyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1160_a**);

7-(3,3-dimethylbutan-2-yl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1161**);

7-(3,3-dimethylbutan-2-yl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1161_a**);

8-(2-{{[(3S)-6,6-dimethylpiperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-4-(1-methyl-1H-pyrazol-4-yl)-2H,3H,4H,5H,6H-pyrrolo[2,3-f][1,4]oxazepin-5-one (**CDK7_1162**);

8-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-4-(1-methyl-1H-pyrazol-4-yl)-2H,3H,4H,5H,6H-pyrrolo[2,3-f][1,4]oxazepin-5-one (**CDK7_1162_a**);

(6S)-3-(2-{{[(3S)-6,6-dimethylpiperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1163**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1163_a**);

(6R)-3-(2-{{[(3S)-6,6-dimethylpiperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1164**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1164_a**);

3-(2-{{[(3S)-6,6-dimethylpiperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-6,6-dimethyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1165**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,6-dimethyl-7-(1-methyl-1H-pyrazol-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1165_a**);

6-cyclopropyl-3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1105**);

6-cyclopropyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1105_a**);

3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1,3-thiazol-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1158**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(1,3-thiazol-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1058_a**);

1-methyl-5-[7-oxo-3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-6-yl]-1,2-dihydropyridin-2-one (**CDK7_1052**);

1-methyl-5-(7-oxo-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-6-yl)-1,2-dihydropyridin-2-one (**CDK7_1052_a**);

N-[(3S)-piperidin-3-yl]-4-(1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_972**);

N-(piperidin-3-yl)-4-(1H-pyrrol-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_972_a**);

7-methyl-3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-4,8-dione (**CDK7_1021**);

7-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-4,8-dione (**CDK7_1021_a**);

6-(oxan-4-yl)-3-(2-{{[(3S)-piperidin-3-yl]amino}}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1121**);

6-(oxan-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1121_a**);

4-(1H-indol-1-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_809**);

4-(1H-indol-1-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_809_a**);

4-(1H-1,3-benzodiazol-1-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_810**);

4-(1H-1,3-benzodiazol-1-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_810_a**);

4-(1H-1,2,3-benzotriazol-1-yl)-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_811**);

4-(1H-1,2,3-benzotriazol-1-yl)-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine
(**CDK7_811_a**);

7-methyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_953**);

7-methyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-
pyrrolo[2,3-c]azepin-8-one (**CDK7_953_a**);

6-(1-methyl-1H-pyrazol-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-
yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_959**);

6-(1-methyl-1H-pyrazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-
1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_959_a**);

6-benzyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-
pyrrolo[2,3-c]pyridin-7-one (**CDK7_961**);

6-benzyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-
c]pyridin-7-one (**CDK7_961_a**);

N,N-dimethyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-imidazol-
2-carboxamide (**CDK7_966**);

N,N-dimethyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-imidazol-2-
carboxamide (**CDK7_966_a**);

N,N-dimethyl-4-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H-pyrrol-3-
carboxamide (**CDK7_991**);

N,N-dimethyl-4-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H-pyrrol-3-
carboxamide (**CDK7_991_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(pyridin-2-yl)-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1176**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(pyridin-2-yl)-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1176_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-methyl-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1187**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-methyl-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1187_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1-methyl-1H-
pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1199**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-6-(1-methyl-1H-
pyrazol-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1199_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-
1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1200**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-6-methyl-1H,6H,7H-
pyrrolo[2,3-c]pyridin-7-one (**CDK7_1200_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-methyl-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1201**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-methyl-
1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1201_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(1,3-thiazol-4-yl)-
1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1202**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(1,3-thiazol-4-yl)-1H,6H,7H-
pyrrolo[2,3-c]pyridin-7-one (**CDK7_1202_a**);

3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-6-(1,3-thiazol-5-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1203**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(1,3-thiazol-5-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1203_a**);

6-(2-methyl-1,3-thiazol-4-yl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1204**);

6-(2-methyl-1,3-thiazol-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1204_a**);

6-(2-methyl-1,3-thiazol-5-yl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1205**);

-(2-methyl-1,3-thiazol-5-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1205_a**);

6-(4-methyl-1,3-thiazol-2-yl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1206**);

6-(4-methyl-1,3-thiazol-2-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1206_a**);

6-(5-methyl-1,3-thiazol-2-yl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1207**);

6-(5-methyl-1,3-thiazol-2-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1207_a**);

6-(cyclopropylmethyl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1208**);

6-(cyclopropylmethyl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1208_a**);

6-cyclobutyl-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1209**);

6-cyclobutyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1209_a**);

6-cyclopentyl-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1210**);

6-cyclopentyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1210_a**);

6-(1-methylcyclopropyl)-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1211**);

6-(1-methylcyclopropyl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1211_a**);

6-cyclohexyl-3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1212**);

6-cyclohexyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1212_a**);

3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-6-(pyridin-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1213**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(pyridin-2-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1213_a**);

3-(2-[[[(3S)-piperidin-3-yl]amino]-5-(trifluoromethyl)pyrimidin-4-yl]-6-(pyridin-3-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1214**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(pyridin-3-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1214_a**);

3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-(pyridin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1215**);

3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-(pyridin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1215_a**);

6-(3-methylpyridin-4-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1216**);

6-(3-methylpyridin-4-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1216_a**);

6-(2-methylpyridin-3-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1217**);

6-(2-methylpyridin-3-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1217_a**);

6-(4-methylpyridin-3-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1218**);

6-(4-methylpyridin-3-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1218_a**);

6-(3-methylpyridin-2-yl)-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1219**);

6-(3-methylpyridin-2-yl)-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1219_a**);

7-cyclopropyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1225**);

7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1225_a**);

7-cyclobutyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1226**);

7-cyclobutyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1226_a**);

7-(cyclopropylmethyl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1227**);

7-(cyclopropylmethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1227_a**);

7-(cyclobutylmethyl)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1228**);

7-(cyclobutylmethyl)-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1228_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1229**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1229_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2,2,2-trifluoroethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1230**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2,2,2-trifluoroethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1230_a**);

4-{5-methyl-3,4,6,13-tetraazatricyclo[8.3.0.0²,⁶]trideca-1(10),2,4,11-tetraen-11-yl}-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1236**);

4-{5-methyl-3,4,6,13-tetraazatricyclo[8.3.0.0²,⁶]trideca-1(10),2,4,11-tetraen-11-yl}-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1236_a**);

4-{5-cyclopropyl-3,4,6,13-tetraazatricyclo[8.3.0.0²,⁶]trideca-1(10),2,4,11-tetraen-11-yl}-N-[(3S)-piperidin-3-yl]-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1237**);

4-{5-cyclopropyl-3,4,6,13-tetraazatricyclo[8.3.0.0²,⁶]trideca-1(10),2,4,11-tetraen-11-yl}-N-(piperidin-3-yl)-5-(trifluoromethyl)pyrimidin-2-amine (**CDK7_1237_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,6,7-trimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1238**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,6,7-trimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1238_a**);

(6S)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1239**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1239_a**);

(6R)-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1240**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,7-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1240_a**);

7-cyclopropyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6,6-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1241**);

7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6,6-dimethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1241_a**);

(6S)-7-cyclopropyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1242**);

7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1242_a**);

(6R)-7-cyclopropyl-3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1243**);

7-cyclopropyl-3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-6-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1243_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-5,5,6-trimethyl-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1244**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-5,5,6-trimethyl-1H,4H,5H,6H,7H-pyrrolo[2,3-c]pyridin-7-one (**CDK7_1244_a**);

6,7-dimethyl-3-(2-{[(3S)-piperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1254**);

6,7-dimethyl-3-{2-[(piperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1254_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-ethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1259**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-ethyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1259_a**);

6-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1,4-diazatricyclo[7.1.1.0³,⁷]undeca-3(7),5-dien-2-one (**CDK7_1261**);

6-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1,4-diazatricyclo[7.1.1.0^{3,7}]undeca-3(7),5-dien-2-one (**CDK7_1261_a**);

7-methyl-3-(2-{[(3S,6S)-6-methylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1264**);

7-methyl-3-{2-[(6-methylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1264_a**);

3-(2-{[(3S,6S)-6-ethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1265**);

3-{2-[(6-ethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-methyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1265_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1268**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1268_a**);

2-[3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl]acetonitrile (**CDK7_1269**);

2-(3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl)acetonitrile (**CDK7_1269_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1270**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1270_a**);

3-(2-{[(3S)-6,6-dimethylpiperidin-3-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1271**);

3-{2-[(6,6-dimethylpiperidin-3-yl)amino]-5-(trifluoromethyl)pyrimidin-4-yl}-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1271_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1272**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(2-methoxyethyl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1272_a**);

2-[3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl]acetonitrile (**CDK7_1273**);

2-{3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-8-oxo-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-7-yl}acetonitrile (**CDK7_1273_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1274**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-(oxetan-3-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1274_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1275**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-[(oxetan-3-yl)methyl]-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1275_a**);

3-(2-{[(6S)-4-azaspiro[2.5]octan-6-yl]amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-cyclopropyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1276**);

3-[2-({4-azaspiro[2.5]octan-6-yl}amino)-5-(trifluoromethyl)pyrimidin-4-yl]-7-cyclopropyl-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1276_a**);

3-(2-{{(6S)-4-azaspiro[2.5]octan-6-yl}amino}-5-(trifluoromethyl)pyrimidin-4-yl)-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1277**);

3-[2-{{4-azaspiro[2.5]octan-6-yl}amino}-5-(trifluoromethyl)pyrimidin-4-yl]-7-(propan-2-yl)-1H,4H,5H,6H,7H,8H-pyrrolo[2,3-c]azepin-8-one (**CDK7_1277_a**).

15. A method for inhibiting the biological activity of cyclin-dependent protein kinases CDK7 in a subject, consisting in contacting cyclin-dependent protein kinases CDK7 with a compound according to any one of claims 1-13.

16. A pharmaceutical composition comprising a therapeutically effective amount of the compound according to any one of claims 1-13, or pharmaceutically acceptable salt, solvate, stereoisomer thereof, and one or more pharmaceutically acceptable excipients.

17. The pharmaceutical composition according to claim 15 intended for the prophylaxis or treatment of a disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7.

18. The pharmaceutical composition according to claim 16 intended for the prophylaxis or treatment of a disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7 is an oncological, infectious inflammatory or non-infectious inflammatory, auto-inflammatory, autoimmune disease.

19. The pharmaceutical composition according to claim 17, wherein the oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma;

an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angiitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis,

pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, ostitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis.

20. A method of treatment of a disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7, comprising administering a therapeutically effective amount of the compound according to any one of claims 1-13 or a pharmacologically acceptable salt, solvate or stereoisomer thereof or the pharmaceutical composition according to claim 15 to a subject in need of such treatment.

21. The method of treatment of a disease or disorder according to claim 19, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7 is an oncological, infectious inflammatory or non-infectious inflammatory, auto-inflammatory, autoimmune disease.

22. The method of treatment of a disease or disorder according to claim 20, wherein the oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma;

an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis, myocarditis, nephritis, pyelonephritis, ostitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis.

23. The use of the compound according to any one of claims 1-13 or a pharmacologically acceptable salt, solvate or stereoisomer thereof or the pharmaceutical composition according to claim 15 for treatment of a disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7 in a subject in need of such treatment.

24. The use according to claim 22, wherein the disease or disorder associated with increased activity of cyclin-dependent protein kinase CDK7 is an oncological, infectious inflammatory or non-infectious inflammatory, auto-inflammatory, autoimmune disease.

25. The use according to claim 23, wherein the oncological disease selected from a group comprising breast cancer, metastatic breast cancer, HER2-positive breast cancer, breast cancer expressing estrogen receptors, breast cancer expressing progesterone receptors, triple negative breast cancer (TNBC), HER2-negative breast cancer expressing estrogen and/or progesterone receptors, HER2-negative breast cancer, small cell lung cancer, metastatic small cell lung cancer, non-small cell lung cancer, ovarian cancer, metastatic ovarian cancer, glioma, astrocytoma, glioblastoma, pancreatic cancer, stomach cancer, colorectal cancer, head and neck cancer, head and neck squamous cell carcinoma, oral cancer, squamous cell carcinoma of oral cavity, hepatobiliary cancer, hepatocellular carcinoma, urothelial cancer, prostate cancer, castration-resistant prostate cancer, thyroid cancer, leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, lymphoma, T-cell lymphoma, B-cell lymphoma, mantle cell lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma, neuroblastoma, skin cancer, melanoma, cervical cancer, uterine corpus cancer, fallopian tube cancer, sarcoma, Ewing sarcoma, soft tissue sarcoma, primary peritoneal cancer, retinoblastoma, multiple myeloma;

an autoimmune disease selected from a group comprising axial spondyloarthritis, antiphospholipid syndrome, vasculitides (ANCA-associated vasculitides including Wegener's granulomatosis, and related forms of angitis, temporal arteritis and polyarteritis nodosa, necrotizing vasculitis), Goodpasture syndrome, autoimmune thyroiditis, Graves' disease, bronchial asthma, cardiomyopathy, Crohn's disease, dermatomyositis, polymyositis, mixed connective tissue disease, glomerulonephritis, Guillain-Barre syndrome, Lyme arthritis, psoriasis, pemphigus, psoriatic arthritis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic lupus erythematosus, sarcoidosis, ulcerative colitis, multiple sclerosis, myasthenia gravis and uveitis;

an autoinflammatory disease selected from a group comprising Behcet's disease, Blau syndrome, chronic recurrent multi-focal osteomyelitis (CRMO), synovitis, acne, pustulosis, hyperostosis, osteitis (SAPHO) syndrome, cryopyrin-associated periodic syndromes (CAPS), deficiency of interleukin-1 receptor antagonist (DIRA), familial Mediterranean fever (FMF), NLRP12-associated autoinflammatory diseases, neonatal onset multisystem inflammatory disease (NOMID), chronic infantile neurological cutaneous and articular (CINCA) syndrome, Majeed syndrome, mevalonate kinase deficiency (hyperimmunoglobulin-D syndrome), periodic fever, aphthous stomatitis, pharyngitis, and cervical adenitis (PFAPA) syndrome, pyogenic arthritis, pyoderma gangrenosum, and acne (PAPA) syndrome, Schnitzler syndrome, Sweet syndrome, systemic juvenile idiopathic arthritis, Still's disease, adult onset Still's disease (AOSD), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS);

an infectious inflammatory or non-infectious inflammatory disease selected from a group comprising allergies (including, but not limited to: delayed-type hypersensitivity reactions, immediate-type hypersensitivity reactions, anaphylactic shock), allograft rejection, graft-versus-host disease, atherosclerosis, arteritis, arthritis, arthrositis, adult respiratory distress syndrome, respiratory tract inflammation (including, but not limited to: bronchiolitis, bronchitis, bronchiectasis, cellular interstitial pneumonia, giant cell interstitial pneumonia, lymphoid interstitial pneumonia, desquamative interstitial pneumonia, hypersensitivity pneumonitis, laryngitis, pneumonitis, pharyngitis, pleurisy, pneumonia, pneumoconiosis, asbestosis, berylliosis, silicosis, talcosis, usual interstitial pneumonitis), tonsillitis, bursitis, phlegmon, chorioamnionitis, blepharitis, iritis, conjunctivitis, dacryoadenitis, chronic cholecystitis, cystic fibrosis, diabetes mellitus, encephalitis, endocarditis, epicondylitis, fasciitis, fibromyalgia, gastritis, gastroenteritis, gingivitis, stomatitis, lymphadenitis, hay fever, inflammatory bowel disease (including, but not limited to: appendicitis, enteritis, enterocolitis, ileitis, necrotizing enterocolitis, proctitis, inflammatory dermatoses), myelitis,

myocarditis, nephritis, pyelonephritis, ostitis, osteomyelitis, optic neuritis, otitis, pancreatitis, mumps, pemphigoid, pericarditis, pernicious anemia, phlebitis, rheumatic polymyalgia, hepatitis, cholangitis, progressive systemic sclerosing cholangitis, reperfusion injury, acute rheumatic fever, rhinitis, sinusitis, synovitis, tendinitis, necrotizing fasciitis, prostatitis, cystitis, epididymitis, omphalitis, oophoritis, orchitis, testitis, transverse myelitis, urethritis, urocystitis, uveitis, cervicitis, salpingitis, endometritis, vaginitis, vulvitis, vulvovaginitis.