

PYRAZOLE DERIVATIVES AS 11-BETA-HSD1 INHIBITORS

This invention relates to chemical compounds, or pharmaceutically-acceptable salts thereof. These compounds possess human 11- β -hydroxysteroid dehydrogenase type 1 enzyme (11 β HSD1) inhibitory activity and accordingly have value in the treatment of disease states including metabolic syndrome and are useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said compounds, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments to inhibit 11 β HSD1 in a warm-blooded animal, such as man.

Glucocorticoids (cortisol in man, corticosterone in rodents) are counter regulatory hormones i.e. they oppose the actions of insulin (Dallman MF, Strack AM, Akana SF et al. 1993; Front Neuroendocrinol 14, 303-347). They regulate the expression of hepatic enzymes involved in gluconeogenesis and increase substrate supply by releasing glycerol from adipose tissue (increased lipolysis) and amino acids from muscle (decreased protein synthesis and increased protein degradation). Glucocorticoids are also important in the differentiation of pre-adipocytes into mature adipocytes which are able to store triglycerides (Bujalska IJ et al. 1999; Endocrinology 140, 3188-3196). This may be critical in disease states where glucocorticoids induced by "stress" are associated with central obesity which itself is a strong risk factor for type 2 diabetes, hypertension and cardiovascular disease (Bjorntorp P & Rosmond R 2000; Int. J. Obesity 24, S80-S85).

It is now well established that glucocorticoid activity is controlled not simply by secretion of cortisol but also at the tissue level by intracellular interconversion of active cortisol and inactive cortisone by the 11-beta hydroxysteroid dehydrogenases, 11 β HSD1 (which activates cortisone) and 11 β HSD2 (which inactivates cortisol) (Sandeep TC & Walker BR 2001 Trends in Endocrinol & Metab. 12, 446-453). That this mechanism may be important in man was initially shown using carbenoxolone (an anti-ulcer drug which inhibits both 11 β HSD1 and 2) treatment which (Walker BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159) leads to increased insulin sensitivity indicating that 11 β HSD1 may well be regulating the effects of insulin by decreasing tissue levels of active glucocorticoids (Walker BR et al. 1995; J. Clin. Endocrinol. Metab. 80, 3155-3159).

Clinically, Cushing's syndrome is associated with cortisol excess which in turn is associated with glucose intolerance, central obesity (caused by stimulation of pre-adipocyte differentiation in this depot), dyslipidaemia and hypertension. Cushing's syndrome shows a number of clear parallels with metabolic syndrome. Even though the metabolic syndrome is not generally associated with excess circulating cortisol levels (Jessop DS et al. 2001; J. Clin. Endocrinol. Metab. 86, 4109-4114) abnormally high 11 β HSD1 activity within tissues would be expected to have the same effect. In obese men it was shown that despite having similar or lower plasma cortisol levels than lean controls, 11 β HSD1 activity in subcutaneous fat was greatly enhanced (Rask E et al. 2001; J. Clin. Endocrinol. Metab. 1418-1421). Furthermore, the central fat, associated with the metabolic syndrome expresses much higher levels of 11 β HSD1 activity than subcutaneous fat (Bujalska IJ et al. 1997; Lancet 349, 1210-1213). Thus there appears to be a link between glucocorticoids, 11 β HSD1 and the metabolic syndrome.

11 β HSD1 knock-out mice show attenuated glucocorticoid-induced activation of gluconeogenic enzymes in response to fasting and lower plasma glucose levels in response to stress or obesity (Kotelevtsev Y et al. 1997; Proc. Natl. Acad. Sci USA 94, 14924-14929) indicating the utility of inhibition of 11 β HSD1 in lowering of plasma glucose and hepatic glucose output in type 2 diabetes. Furthermore, these mice express an anti-atherogenic lipoprotein profile, having low triglycerides, increased HDL cholesterol and increased apo-lipoprotein AI levels. (Morton NM et al. 2001; J. Biol. Chem. 276, 41293-41300). This phenotype is due to an increased hepatic expression of enzymes of fat catabolism and PPAR α . Again this indicates the utility of 11 β HSD1 inhibition in treatment of the dyslipidaemia of the metabolic syndrome.

The most convincing demonstration of a link between the metabolic syndrome and 11 β HSD1 comes from recent studies of transgenic mice over-expressing 11 β HSD1 (Masuzaki H et al. 2001; Science 294, 2166-2170). When expressed under the control of an adipose specific promoter, 11 β HSD1 transgenic mice have high adipose levels of corticosterone, central obesity, insulin resistant diabetes, hyperlipidaemia and hyperphagia. Most importantly, the increased levels of 11 β HSD1 activity in the fat of these mice are similar to those seen in obese subjects. Hepatic 11 β HSD1 activity and plasma corticosterone levels were normal, however, hepatic portal vein levels of corticosterone

were increased 3 fold and it is thought that this is the cause of the metabolic effects in liver.

Overall it is now clear that the complete metabolic syndrome can be mimicked in mice simply by overexpressing 11 β HSD1 in fat alone at levels similar to those in obese man.

11 β HSD1 tissue distribution is widespread and overlapping with that of the glucocorticoid receptor. Thus, 11 β HSD1 inhibition could potentially oppose the effects of glucocorticoids in a number of physiological/pathological roles. 11 β HSD1 is present in human skeletal muscle and glucocorticoid opposition to the anabolic effects of insulin on protein turnover and glucose metabolism are well documented (Whorwood CB et al. 2001; J. Clin. Endocrinol. Metab. 86, 2296-2308). Skeletal muscle must therefore be an important target for 11 β HSD1 based therapy.

Glucocorticoids also decrease insulin secretion and this could exacerbate the effects of glucocorticoid induced insulin resistance. Pancreatic islets express 11 β HSD1 and carbenoxolone can inhibit the effects of 11-dehydrocorticosterone on insulin release (Davani B et al. 2000; J. Biol. Chem. 275, 34841-34844). Thus in treatment of diabetes 11 β HSD1 inhibitors may not only act at the tissue level on insulin resistance but also increase insulin secretion itself.

Skeletal development and bone function is also regulated by glucocorticoid action. 11 β HSD1 is present in human bone osteoclasts and osteoblasts and treatment of healthy volunteers with carbenoxolone showed a decrease in bone resorption markers with no change in bone formation markers (Cooper MS et al 2000; Bone 27, 375-381). Inhibition of 11 β HSD1 activity in bone could be used as a protective mechanism in treatment of osteoporosis.

Glucocorticoids may also be involved in diseases of the eye such as glaucoma. 11 β HSD1 has been shown to affect intraocular pressure in man and inhibition of 11 β HSD1 may be expected to alleviate the increased intraocular pressure associated with glaucoma (Rauz S et al. 2001; Investigative Ophthalmology & Visual Science 42, 2037-2042).

There appears to be a convincing link between 11 β HSD1 and the metabolic syndrome both in rodents and in humans. Evidence suggests that a drug which specifically inhibits 11 β HSD1 in type 2 obese diabetic patients will lower blood glucose by reducing

hepatic gluconeogenesis, reduce central obesity, improve the atherogenic lipoprotein phenotype, lower blood pressure and reduce insulin resistance. Insulin effects in muscle will be enhanced and insulin secretion from the beta cells of the islet may also be increased.

5 Currently there are two main recognised definitions of metabolic syndrome.

1) The Adult Treatment Panel (ATP III 2001 JMA) definition of metabolic syndrome indicates that it is present if the patient has three or more of the following symptoms:

Waist measuring at least 40 inches (102 cm) for men, 35 inches (88 cm) for women;

Serum triglyceride levels of at least 150 mg/dl (1.69 mmol/l);

10 HDL cholesterol levels of less than 40 mg/dl (1.04 mmol/l) in men, less than 50 mg/dl (1.29 mmol/l) in women;

Blood pressure of at least 135/80 mm Hg; and / or Blood sugar (serum glucose) of at least 110 mg/dl (6.1 mmol/l).

2) The WHO consultation has recommended the following definition which does not
15 imply causal relationships and is suggested as a working definition to be improved upon in due course:

The patient has at least one of the following conditions: glucose intolerance, impaired glucose tolerance (IGT) or diabetes mellitus and/or insulin resistance; together with two or more of the following:

20 Raised Arterial Pressure;

Raised plasma triglycerides


Central Obesity

Microalbuminuria

We have found that the compounds defined in the present invention, or a
25 pharmaceutically-acceptable salt thereof, are effective 11β HSD1 inhibitors, and accordingly have value in the treatment of disease states associated with metabolic syndrome. We have also found that the compounds of the invention have improved properties, which would make them better candidates for use as pharmaceuticals. For example, in general the compounds of the invention have good oral bioavailability whilst
30 retaining potency. Therefore this group of compounds would be expected to provide superior oral exposure at a lower dose and thereby be particularly suitable for use in the treatment or prevention of a disease or medical condition treatable by inhibiting 11β HSD1.

The compounds of the invention may also have superior potency and/or advantageous physical properties and/or favourable toxicity profiles and/or favourable metabolic profiles in comparison with other 11 β HSD1 inhibitors known in the art. For example, in general the compounds of the present invention have acceptable free drug levels as measured by plasma protein binding experiments.

Accordingly there is provided a compound of formula (1):



(1)

wherein:

- Q is O, S or a single bond;
- R^1 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, heterocyclyl, aryl C_{1-3} alkyl, heteroaryl C_{1-3} alkyl, C_{3-7} cycloalkyl C_{1-3} alkyl, C_{3-7} cycloalkyl C_{2-3} alkenyl or C_{3-7} cycloalkyl C_{2-3} alkynyl, [each of which is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-3} alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl, C_{1-3} alkoxy, C_{1-3} alkylS(O) $_n$ - (wherein n is 0, 1, 2 or 3), $R^{5'}CON(R^{5'})$ -, $(R^{5'})(R^{5''})NC(O)$ -, $R^{5'}C(O)O$ -, $R^{5'}OC(O)$ -, $(R^{5'})(R^{5''})NC(O)N(R^{5''})$ -, $R^{5'}SO_2N(R^{5''})$ -, and $(R^{5'})(R^{5''})NSO_2$ - (wherein R^5 is C_{1-3} alkyl optionally substituted by 1, 2 or 3 substituents selected from hydroxyl, halo or cyano; and $R^{5'}$ and $R^{5''}$ are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from hydroxyl, halo, C_{1-3} alkoxy, carboxy and cyano or $R^{5'}$ and $R^{5''}$ together with the nitrogen atom to which they are attached form a 4-7 membered saturated ring)];
- R^2 is selected from heterocyclyl, C_{3-7} cycloalkyl(CH $_2$) $_m$ -, and C_{6-12} polycycloalkyl(CH $_2$) $_m$ - (wherein m is 0, 1 or 2 and the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R^6);

R^3 is selected from hydrogen, C_{1-4} alkyl C_{3-5} cycloalkyl and C_{3-5} cycloalkylmethyl (each of which is optionally substituted by 1, 2 or 3 fluoro atoms);

R^2 and R^3 together with the nitrogen atom to which they are attached form a saturated mono, bicyclic or bridged ring system optionally containing 1 or 2 additional ring

5 heteroatoms selected from nitrogen, oxygen and sulphur and which is optionally fused to a saturated, partially saturated or unsaturated monocyclic ring wherein the resulting ring system is optionally substituted by 1, 2, or 3 substituents independently selected from R^7 ; R^6 and R^7 are independently selected from hydroxyl, halo, oxo, carboxy, cyano, trifluoromethyl, R^9 , R^9O- , R^9CO- , $R^9C(O)O-$, $R^9CON(R^9')$ -, $(R^9')(R^9'')NC(O)-$, $(R^9')(R^9''')N-$, $R^9S(O)_a-$ wherein a is 0 to 2, $R^9OC(O)-$, $(R^9')(R^9'')NSO_2-$, $R^9SO_2N(R^9)-$, $(R^9')(R^9'')NC(O)N(R^9''')$ -, phenyl and heteroaryl [wherein the phenyl and heteroaryl groups are optionally fused to a phenyl, heteroaryl or a saturated or partially-saturated 5- or 6-membered ring optionally containing 1, 2 or 3 heteroatoms independently selected from nitrogen, oxygen and sulphur and the resulting ring system is optionally substituted by 1, 2
15 or 3 substituents independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, N- C_{1-4} alkylamino, di-N,N-(C_{1-4} alkyl)amino, N- C_{1-4} alkylcarbonyl, di-N,N-(C_{1-4} alkyl)carbonyl, C_{1-4} alkylS(O)_r-, C_{1-4} alkylS(O)_r C_{1-4} alkyl (wherein r is 0, 1 or 2)];

R^9 is independently selected from C_{1-3} alkyl optionally substituted by hydroxyl, halo,
20 C_{1-4} alkoxy, carboxy or cyano;

R^9' , R^9'' and R^9''' are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by 1,2,or 3 substituents independently selected from hydroxyl, halo, C_{1-4} alkoxy, carboxy and cyano);

A is a phenyl or heteroaryl ring (the phenyl or heteroaryl ring being optionally substituted
25 on ring carbon atoms by 1, 2 or 3 R^{10} groups and on an available ring nitrogen in a heteroaryl group by R^{11});

R^{10} is independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, N- C_{1-4} alkylamino, di-N,N-(C_{1-4} alkyl)amino, N- C_{1-4} alkylcarbonyl, di-N,N-(C_{1-4} alkyl)carbonyl,
30 C_{1-4} alkylS(O)_s-, C_{1-4} alkylS(O)_s C_{1-4} alkyl (wherein s is 0, 1 or 2)];

R^{11} is independently C_{1-3} alkyl optionally substituted by 1, 2 or 3 fluoro atoms;

X is a direct bond, C₃₋₄cycloalkandiyl, C₃₋₄cycloalkanylidene, -C(R¹²)(R¹³)-, -C(R¹²)(R¹³)C(R¹⁴)(R¹⁵)-, -CH₂O- or -CH₂S(O)_t- (wherein t is 0, 1 or 2):

Y is a direct bond, C₃₋₄cycloalkandiyl, C₃₋₄cycloalkanylidene, -C(R¹⁶)(R¹⁷)- or -C(R¹⁸)(R¹⁹)C(R²⁰)(R²¹)-;

5 wherein R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from hydrogen and methyl;

or a pharmaceutically-acceptable salt thereof.

In this specification the term “alkyl” includes both straight and branched chain alkyl groups but references to individual alkyl groups such as “propyl” are specific for the
10 straight chain version only. For example, “C₁₋₄alkyl” includes propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as ‘propyl’ are specific for the straight chain version only and references to individual branched chain alkyl groups such as ‘isopropyl’ are specific for the branched chain version only. A similar convention applies to other radicals therefore “C₁₋₄alkoxyC₁₋₄alkyl” would include
15 1-(C₁₋₄alkoxy)propyl, 2-(C₁₋₄alkoxy)ethyl and 3-(C₁₋₄alkoxy)butyl. The term “halo” refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from “one or more” groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

20 A 4-7 membered saturated ring (for example formed between R^{5'} and R^{5''} and the nitrogen atom to which they are attached) is a monocyclic ring containing the nitrogen atom as the only ring atom.

“Heteroaryl”, unless otherwise specified, is a totally unsaturated, monocyclic ring containing 5 or 6 atoms of which at least 1, 2 or 3 ring atoms are independently chosen
25 from nitrogen, sulphur or oxygen, which may, unless otherwise specified, be carbon-linked. A ring nitrogen atom may be optionally oxidised to form the corresponding N-oxide. Examples and suitable values of the term “heteroaryl” are thienyl, furyl, thiazolyl, pyrazolyl, isoxazolyl, imidazolyl, pyrrolyl, thiadiazolyl, isothiazolyl, triazolyl, pyrimidyl, pyrazinyl, pyridazinyl and pyridyl. Particularly “heteroaryl” refers to thienyl, furyl,
30 thiazolyl, pyridyl, imidazolyl or pyrazolyl.

“Heterocycyl” is a 4-7 saturated, monocyclic ring having 1-3 ring heteroatoms selected from nitrogen, oxygen and sulphur. The ring sulphur may be optionally oxidised to SO₂.

A C₃₋₇cycloalkyl ring is a saturated carbon ring containing from 3 to 7 ring atoms.

5 A C₃₋₄cycloalkandiyl ring is a saturated carbon ring containing 3 or 4 ring atoms. It is a diradical with the radicals on different ring carbon atoms.

A C₃₋₄cycloalkanylidene ring is a saturated carbon ring containing 3 or 4 ring atoms. It is a diradical with the radicals on the same ring carbon atom.

10 A polycycloalkyl ring is a ring system in which either at least 2 rings are fused together or in which 2 ring have one ring atom in common (spiro).

A “saturated mono, bicyclic or bridged ring system optionally containing 1 or 2 additional ring heteroatoms selected from nitrogen, oxygen and sulphur”, unless otherwise specified contains 4-14 ring atoms. Particularly a mono ring contains 4 –7 ring atoms, a bicyclic ring 6-14 ring atoms and a bridged ring system 6-14 ring atoms. Examples of
15 mono rings include piperidinyl, piperazinyl and morpholinyl. Examples of bicyclic rings include decalin and 2,3,3a,4,5,6,7,7a-octahydro-1H-indene.

Bridged ring systems are ring systems in which there are two or more bonds common to two or more constituent rings. Examples of bridged ring systems include 1,3,3-trimethyl-6-azabicyclo[3.2.1]octane, 2-aza-bicyclo[2.2.1]heptane and 7-
20 azabicyclo(2,2,1)heptane, 1- and 2-adamantanyl.

A “saturated, partially saturated or unsaturated monocyclic ring” is, unless otherwise specified, a 4-7 membered ring. Examples include, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl and phenyl.

25 Examples of a “saturated or partially-saturated 5- or 6-membered ring optionally containing 1, 2 or 3 heteroatoms independently selected from nitrogen, oxygen and sulphur” include piperidinyl, piperazinyl and morpholinyl.

Examples of “C₁₋₄alkoxy” include methoxy, ethoxy and propoxy. Examples of “C₁₋₄alkoxyC₁₋₄alkyl” include methoxymethyl, ethoxymethyl, propoxymethyl, 2-methoxyethyl, 2-ethoxyethyl and 2-propoxyethyl. Examples of “C₁₋₄alkylS(O)_n wherein n
30 is 0 to 2” include methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl and ethylsulphonyl. Examples of “C₁₋₄alkylS(O)_qC₁₋₄alkyl” wherein q is 0 to 2” include methylthio, ethylthio, methylsulphinyl, ethylsulphinyl, mesyl, ethylsulphonyl,

methylthiomethyl, ethylthiomethyl, methylsulphinylmethyl, ethylsulphinylmethyl, mesylmethyl and ethylsulphonylmethyl. Examples of "C₁₋₄alkanoyl" include propionyl and acetyl. Examples of "*N*-(C₁₋₄alkyl)amino" include methylamino and ethylamino. Examples of "*N,N*-(C₁₋₄alkyl)₂amino" include *N,N*-dimethylamino, *N,N*-diethylamino and

5 *N*-ethyl-*N*-methylamino. Examples of "C₂₋₄alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C₂₋₄alkynyl" are ethynyl, 1-propynyl and 2-propynyl. Examples of "*N*-(C₁₋₄alkyl)carbamoyl" are methylaminocarbonyl and ethylaminocarbonyl. Examples of "*N,N*-(C₁₋₄alkyl)₂carbamoyl" are dimethylaminocarbonyl and methylethylaminocarbonyl. Examples of "C₃₋₇cycloalkylC₁₋₃alkalkyl" include cyclopropylmethyl, 2-cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl. Examples of "C₃₋

10 ₇cycloalkylC₂₋₃alkalkenyl" include 2-cyclopropylethenyl, 2-cyclopentylethenyl and 2-cyclohexylethenyl. Examples of "C₃₋₇cycloalkylC₂₋₃alkalkynyl" include 2-cyclopropylethynyl, 2-cyclopentylethynyl and 2-cyclohexylethynyl.

Examples of "C₃₋₇cycloalkyl(CH₂)_{*m*}" include cyclopropylmethyl, 2-

15 cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl. Examples of C₆₋₁₂polycycloalkyl(CH₂)_{*m*}- include norbornyl bicyclo[2.2.2]octane(CH₂)_{*m*}-, bicyclo[3.2.1]octane(CH₂)_{*m*}- and 1- and 2-adamantanyl(CH₂)_{*m*}-.

A suitable pharmaceutically-acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic,

20 for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric or maleic acid. In addition a suitable pharmaceutically-acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an

25 ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

Some compounds of the formula (1) may have chiral centres and/or geometric isomeric centres (E- and Z- isomers), and it is to be understood that the invention

30 encompasses all such optical, diastereoisomers and geometric isomers that possess 11βHSD1 inhibitory activity.


The invention relates to any and all tautomeric forms of the compounds of the formula (1) that possess 11 β HSD1 inhibitory activity.

It is also to be understood that certain compounds of the formula (1) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms, which possess 11 β HSD1 inhibitory activity.

The invention also relates to in vivo hydrolysable esters of a compound of the formula (1). In vivo hydrolysable esters are those esters that are broken down in the animal body to produce the parent carboxylic acid.


In one embodiment of the invention are provided compounds of formula (1). In an alternative embodiment are provided pharmaceutically-acceptable salts of compounds of formula (1).

In one aspect of the invention, there is provided a compound of the formula (1a):



wherein R^1 , R^2 and R^3 are as hereinabove defined and R^{10} is selected from hydrogen, C_{1-4} alkyl, trifluoromethyl, C_{1-4} alkoxy and C_{1-4} alkylS-. In another aspect R^{10} is selected from hydrogen, methyl, trifluoromethyl, methoxy and methylthio. In another aspect R^{10} is hydrogen.

In another aspect of the invention, there is provided a compound of the formula (1b):



wherein R^1 , R^2 and R^3 are as hereinabove defined and R^{10} is selected from hydrogen, C_{1-4} alkyl, trifluoromethyl, C_{1-4} alkoxy and C_{1-4} alkylS-. In another aspect R^{10} is selected from hydrogen, methyl, trifluoromethyl, methoxy and methylthio. In another aspect R^{10} is hydrogen.

Particular values of variable groups are as follows. Such values may be used where appropriate with any of the definitions, claims or embodiments defined hereinbefore or hereinafter, for compounds of formula (1). The definitions of R^1 , R^2 and R^3 and variables within those groups may be used for the compound of formula (1a):

Definition of Q

- a) In one aspect, the invention relates to a compound of the formula (I) as hereinabove defined wherein Q is O.
- b) In another aspect Q is S.
- c) In another aspect Q is a single bond.

Definition of R^1

- a) In one aspect R^1 is C_{3-6} cycloalkyl optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-3} alkyl, hydroxy, halo, oxo, cyano, fluoro, trifluoromethyl and C_{1-3} alkoxy.
- b) In another aspect R^1 is C_{3-6} cycloalkyl.
- c) In another aspect R^1 is C_{3-6} cycloalkyl C_{1-2} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-3} alkyl, hydroxy, halo, oxo, cyano, fluoro, trifluoromethyl and C_{1-3} alkoxy.

- d) In another aspect R^1 is C_{3-4} cycloalkyl C_{1-2} alkyl.
- e) In another aspect R^1 is C_{1-4} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-3} alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl and C_{1-3} alkoxy.
- 5 f) In another aspect R^1 is C_{1-4} alkyl.
- g) In another aspect R^1 is propyl optionally substituted by 1 or 2 substituents independently selected from C_{1-3} alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl and C_{1-3} alkoxy.
- h) In another aspect R^1 is propyl.
- 10 Definition of R^2
- a) In one aspect, R^2 is selected from C_{3-7} cycloalkyl $(CH_2)_m$ -, and C_{6-12} polycycloalkyl $(CH_2)_m$ - (wherein m is 0, 1 or 2 and the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R^6) wherein m is 0, 1 or 2.
- b) In another aspect, R^2 is selected from C_{5-7} cycloalkyl $(CH_2)_m$ - and
- 15 C_{8-12} polycycloalkyl $(CH_2)_m$ - (wherein the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R^6) and wherein m is 0, 1 or 2.
- c) In another aspect, R^2 is selected from C_{5-7} cycloalkyl $(CH_2)_m$ -, C_{7-10} bicycloalkyl $(CH_2)_m$ - and C_{10} tricycloalkyl $(CH_2)_m$ - (wherein the cycloalkyl, bicycloalkyl and tricycloalkyl rings are optionally substituted by 1, 2 or 3 substituents
- 20 independently selected from R^6) and wherein m is 0, 1 or 2.
- d) In yet another aspect, R^2 is selected from C_{5-7} cycloalkyl $(CH_2)_m$ -, C_{7-10} bicycloalkyl $(CH_2)_m$ - and adamantyl (wherein the cycloalkyl, bicycloalkyl and tricycloalkyl rings are optionally substituted by 1, 2 or 3 substituents independently selected from R^6) and wherein m is 0, 1 or 2.
- 25 Definition of m
- a) In one aspect, m is 0 or 1.
- Definition of R^3
- a) In one aspect, R^3 is C_{1-4} alkyl.
- b) In another aspect, R^3 is hydrogen, methyl or ethyl.
- 30 c) In another aspect, R^3 is hydrogen.
- d) In another aspect, R^3 is methyl.
- e) In another aspect, R^3 is ethyl.

f) In another aspect, R^3 is cyclopropyl.

Definition of R^2 and R^3 together

- a) In another aspect, R^2 and R^3 together with the nitrogen atom to which they are attached form a saturated 5 or 6-membered mono, 6-12 membered bicyclic or 6-12 membered bridged ring system optionally containing 1 or 2 additional ring heteroatoms selected from nitrogen, oxygen and sulphur and which is optionally fused to a saturated, partially-saturated or aryl monocyclic ring wherein the resulting ring system is optionally substituted by 1, 2, or 3 substituents independently selected from R^7 .

Definition of R^6

- 10 a) In one aspect, R^6 is independently selected from hydroxyl, R^9O- , R^9CO- and $R^9C(O)O-$ wherein R^9 is as hereinabove defined.
- b) In another aspect, R^6 is independently selected from hydroxyl, R^9O- , R^9CO- and $R^9C(O)O-$ wherein R^9 is C_{1-3} alkyl optionally substituted by C_{1-4} alkoxy or carboxy.
- 15 c) In another aspect, R^6 is independently selected from $R^9CON(R^{9'})-$, $R^9SO_2N(R^{9''})-$ and $(R^{9'}) (R^{9''})NC(O)N(R^{9''''})-$; wherein R^9 is as hereinabove defined.
- d) In another aspect, R^6 is independently selected from $R^9CON(R^{9'})-$, $R^9SO_2N(R^{9''})-$ and $(R^{9'}) (R^{9''})NC(O)N(R^{9''''})-$; R^9 is C_{1-3} alkyl optionally substituted by C_{1-4} alkoxy or carboxy; $R^{9'}$, $R^{9''}$ and $R^{9''''}$ are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by C_{1-4} alkoxy or carboxy).
- 20 e) In another aspect, R^6 is independently selected from $(R^{9'}) (R^{9''})NC(O)-$ and $(R^{9'}) (R^{9''})N-$; wherein $R^{9'}$ and $R^{9''}$ are as hereinabove defined.
- f) In another aspect, R^6 is independently selected from $(R^{9'}) (R^{9''})NC(O)-$ and $(R^{9'}) (R^{9''})N-$; wherein $R^{9'}$ and $R^{9''}$ are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by C_{1-4} alkoxy or carboxy.
- 30 g) In one aspect R^6 is selected from methyl, trifluoromethyl, chloro, fluoro, bromo, methoxy, ethoxy, trifluoromethoxy, methanesulfonyl, ethanesulfonyl, methylthio, ethylthio,

amino, N-methylamino, N-ethylamino, N-propylamino, N,N-dimethylamino, N,N-methylethylamino or N,N-diethylamino.

- h) In another aspect, R⁶ is optionally substituted phenyl, pyridyl or pyrimidyl.
- i) In another aspect, R⁶ is optionally substituted pyrid-2-yl, pyrid-3-yl or pyrid-4-yl.

5 Definition of R⁷

- a) In another aspect, R⁷ is independently selected from hydroxyl, halo, oxo, cyano, trifluoromethyl, R⁹ and R⁹O- (wherein R⁹ is as hereinabove defined).
- b) In another aspect, R⁷ is independently selected from hydroxyl, halo, trifluoromethyl, R⁹ and R⁹O- (wherein R⁹ is as hereinabove defined).

10 Definition of R⁹

- a) In one aspect, R⁹ is independently selected from C₁₋₃alkyl.

Definition of R^{9'}, R^{9''} and R^{9'''}

- a) In one aspect, R^{9'}, R^{9''} and R^{9'''} are independently selected from hydrogen and C₁₋₃alkyl.

15 Definition of Y

- a) In one aspect, Y is independently selected from direct bond, -CH₂- and -CH₂CH₂-.
- b) In one aspect, Y is independently selected from -CH₂- and -CH₂CH₂-.
- c) In another aspect Y is a direct bond.

Definition of A

- 20 a) In one aspect A is phenyl optionally substituted by R¹⁰.
- b) In another aspect A is heteroaryl optionally substituted by R¹⁰ and R¹¹.
- c) In another aspect A is thienyl optionally substituted by R¹⁰ and R¹¹.
- d) In another aspect A is pyridyl optionally substituted by R¹⁰ and R¹¹.
- e) In another aspect A is phen-1,4-diyl

25 Definition of R¹⁰

- a) In one aspect, R¹⁰ is independently selected from C₁₋₄alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, difluoromethoxy, halo, C₁₋₄alkoxy and C₁₋₄alkoxyC₁₋₄alkyl.
- b) In another aspect, R¹⁰ is independently selected from methyl, ethyl, hydroxyl, 30 cyano, trifluoromethyl, trifluoromethoxy, difluoromethoxy, halo, methoxy, ethoxy, methoxymethyl and ethoxymethyl.

c) In another aspect, R^{10} is independently selected from methyl, ethyl, cyano, trifluoromethyl, trifluoromethoxy, difluoromethoxy, halo, methoxy, ethoxy.

Definition of R^{11}

a) In one aspect, R^{11} , is independently selected from C_{1-3} alkyl, trifluoromethyl and difluoromethyl.

b) In one aspect, R^{11} , is independently selected from methyl, ethyl, trifluoromethyl and difluoromethyl.

Definition of X

a) In one aspect, X is independently selected from direct bond, $-CH_2-$, $-CHMe-$, $-CMe_2-$, $-CH_2CH_2-$, $-CH_2O-$ and $-CH_2S-$.

b) In one aspect, X is independently selected from $-CH_2-$, $-CHMe-$, $-CMe_2-$, $-CH_2CH_2-$, $-CH_2O-$ and $-CH_2S-$.

c) In another aspect X is independently selected from cyclopropanylidene, cyclobutanylidene, cyclopropane-1,2-diyl and cyclobutan-1,2-diyl.

d) In another aspect X is a direct bond.

In one aspect, R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} and R^{21} are hydrogen.

In one aspect R^1 is optionally substituted by 0 substituents.

In one aspect R^1 is optionally substituted by 1 substituent.

In one aspect R^1 is optionally substituted by 2 substituents.

In one aspect R^1 is optionally substituted by 3 substituents.

In one aspect R^2 is optionally substituted by 0 substituents.

In one aspect R^2 is optionally substituted by 1 substituent.

In one aspect R^2 is optionally substituted by 2 substituents.

In one aspect R^2 is optionally substituted by 3 substituents.

In one aspect R^3 is optionally substituted by 0 substituents.

In one aspect R^3 is optionally substituted by 1 substituent.

In one aspect R^3 is optionally substituted by 2 substituents.

In one aspect R^3 is optionally substituted by 3 substituents.

In one aspect the group formed by R^2 and R^3 together is optionally substituted by 0 substituents.

In one aspect the group formed by R^2 and R^3 together is optionally substituted by 1 substituent.

In one aspect the group formed by R^2 and R^3 together is optionally substituted by 2 substituents.

5 In one aspect the group formed by R^2 and R^3 together is optionally substituted by 3 substituents.

In one aspect A is optionally substituted by 0 substituents.

In one aspect A is optionally substituted by 1 substituent.

In one aspect A is optionally substituted by 2 substituents.

10 In one aspect A is optionally substituted by 3 substituents.

In one aspect the phenyl and heteroaryl groups in R^6 and R^7 are independently optionally substituted by 0 substituents.

In one aspect the phenyl and heteroaryl groups in R^6 and R^7 are independently optionally substituted by 1 substituent.

15 In one aspect the phenyl and heteroaryl groups in R^6 and R^7 are independently are optionally substituted by 2 substituents.

In one aspect the phenyl and heteroaryl groups in R^6 and R^7 are independently are optionally substituted by 3 substituents.

In one aspect of the invention Q is a direct bond and X is a direct bond.

20 Particular classes of compounds of the present invention are disclosed in Table A using combinations of the definitions described hereinabove. For example, 'a' in the column headed R^2 in the table refers to definition (a) given for R^2 hereinabove and 'I' refers to the first definition given for the variables in the compound of formula (I) at the beginning of the description. R^6 and R^7 are optional substituents on R^2 and the group
25 formed by R^2 and R^3 together. R^2 and R^3 may of course be unsubstituted or substituted by the values listed for R^6 and R^7 . A "--" in the column for R^6 and R^7 means that the relevant R^2 group or the group formed by R^2 and R^3 together is unsubstituted. Table A refers to compounds of the formula (1).

Table A

Class	Q	R ¹	R ²	R ⁶	R ³	R ² and R ³ together	R ⁷	X	Y	A
1	I	I	a	I	I	-	-	I	I	I
2	I	a	-	-	-	I	I	I	I	I
3	I	b	a	I	I	-	-	a	a	a
4	a	I	-	-	-	I	I	a	a	a
5	c	I	a	I	I	-	-	a	a	a
6	b	c	-	-	-	a	g	d	c	a
7	b	d	b	g	b	-	-	d	c	b
8	b	e	c	g	b	-	-	d	c	a
9	b	f	d	-	c	-	-	d	c	e
10	b	h	d	-	c	-	-	d	c	e

Particular classes of compounds of the formula (1a) are disclosed in Table B using combinations of the definitions described hereinabove in a similar manner as for Table A.

Table B

Class	R ¹	R ²	R ⁶	R ³	R ² and R ³ together	R ⁷
1a	I	I	I	I	-	-
2a	a	-	-	-	I	I
3a	b	a	I	I	-	-
4a	I	-	-	-	I	I
5a	I	a	I	I	-	-
6a	c	-	-	-	a	g
7a	d	b	g	b	-	-
8a	e	c	g	c	-	-
9a	f	d	-	-	-	-
10a	h	d	-	-	-	-

Particular classes of compounds of the formula (1b) are disclosed in Table B using combinations of the definitions described hereinabove in a similar manner as for Table A.

Table C

Class	R ¹	R ²	R ⁶	R ³	R ² and R ³ together	R ⁷
1b	I	I	I	I	-	-
2b	a	-	-	-	I	I
3b	b	a	I	I	-	-
4b	I	-	-	-	I	I
5b	I	a	I	I	-	-
6b	c	-	-	-	a	g
7b	d	b	g	b	-	-
8b	e	c	g	c	-	-
9b	f	d	-	-	-	-
10b	h	d	-	-	-	-

A further class of compounds is of formula (1) wherein:

- 5 Q is a single bond;
- R¹ is C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₇cycloalkyl, heterocyclyl, arylC₁₋₃alkyl, heteroarylC₁₋₃alkyl, C₃₋₇cycloalkylC₁₋₃alkyl, C₃₋₇cycloalkylC₂₋₃alkenyl or C₃₋₇cycloalkylC₂₋₃alkynyl, [each of which is optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₃alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl,
- 10 C₁₋₃alkoxy, C₁₋₃alkylS(O)_n- (wherein n is 0, 1, 2 or 3), R⁵CON(R^{5'})-, (R^{5'})(R^{5''})NC(O)-, R^{5'}C(O)O-, R^{5'}OC(O)-, (R^{5'})(R^{5''})NC(O)N(R^{5'''})-, R⁵SO₂N(R^{5'})-, and (R^{5'})(R^{5''})NSO₂- (wherein R⁵ is C₁₋₃alkyl optionally substituted by 1, 2 or 3 substituents selected from hydroxyl, halo or cyano; and
- R^{5'} and R^{5''} are independently selected from hydrogen and C₁₋₃alkyl optionally substituted
- 15 by 1, 2 or 3 substituents independently selected from hydroxyl, halo, C₁₋₃alkoxy, carboxy and cyano or R^{5'} and R^{5''} together with the nitrogen atom to which they are attached form a 4-7 membered saturated ring)];

R^2 is selected from heterocyclyl, C_{3-7} cycloalkyl $(CH_2)_m$ -, and C_{6-12} polycycloalkyl $(CH_2)_m$ - (wherein m is 0, 1 or 2 and the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R^6);

R^3 is selected from hydrogen, C_{1-4} alkyl C_{3-5} cycloalkyl and C_{3-5} cycloalkylmethyl (each of which is optionally substituted by 1, 2 or 3 fluoro atoms);

R^2 and R^3 together with the nitrogen atom to which they are attached form a saturated mono, bicyclic or bridged ring system optionally containing 1 or 2 additional ring heteroatoms selected from nitrogen, oxygen and sulphur and which is optionally fused to a saturated, partially saturated or unsaturated monocyclic ring wherein the resulting ring system is optionally substituted by 1, 2, or 3 substituents independently selected from R^7 ;

R^6 and R^7 are independently selected from hydroxyl, halo, oxo, carboxy, cyano, trifluoromethyl, R^9 , R^9O -, R^9CO -, $R^9C(O)O$ -, $R^9CON(R^9)$ -, $(R^9)(R^9)NC(O)$ -, $(R^9)(R^9)N$ -, $R^9S(O)_a$ - wherein a is 0 to 2, $R^9OC(O)$ -, $(R^9)(R^9)NSO_2$ -, $R^9SO_2N(R^9)$ -, $(R^9)(R^9)NC(O)N(R^9)$ -, phenyl and heteroaryl [wherein the phenyl and heteroaryl groups are optionally fused to a phenyl, heteroaryl or a saturated or partially-saturated 5- or 6-membered ring optionally containing 1, 2 or 3 heteroatoms independently selected from nitrogen, oxygen and sulphur and the resulting ring system is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, N - C_{1-4} alkylamino, di- N,N -(C_{1-4} alkyl)amino, N - C_{1-4} alkylcarbamoyl, di- N,N -(C_{1-4} alkyl)carbamoyl, C_{1-4} alkyl $S(O)_r$ -, C_{1-4} alkyl $S(O)_rC_{1-4}$ alkyl (wherein r is 0, 1 or 2)];

R^9 is independently selected from C_{1-3} alkyl optionally substituted by hydroxyl, halo, C_{1-4} alkoxy, carboxy or cyano;

R^9 , R^9 and R^9 are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by 1,2, or 3 substituents independently selected from hydroxyl, halo, C_{1-4} alkoxy, carboxy and cyano);

A is a phenyl or heteroaryl ring (the phenyl or heteroaryl ring being optionally substituted on ring carbon atoms by 1, 2 or 3 R^{10} groups and on an available ring nitrogen in a heteroaryl group by R^{11});

R^{10} is independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, N - C_{1-4} alkylamino, di-

N,N-(C₁₋₄alkyl)amino, N-C₁₋₄alkylcarbonyl, di-N,N-(C₁₋₄alkyl)carbonyl,

C₁₋₄alkylS(O)_s-, C₁₋₄alkylS(O)_sC₁₋₄alkyl (wherein s is 0, 1 or 2)];

R¹¹ is independently C₁₋₃alkyl optionally substituted by 1, 2 or 3 fluoro atoms;

X is a direct bond, C₃₋₄cycloalkandiyl, C₃₋₄cycloalkanylidene, -C(R¹²)(R¹³)-, -

5 C(R¹²)(R¹³)C(R¹⁴)(R¹⁵)-, -CH₂O- or -CH₂S(O)- (wherein t is 0, 1 or 2):

Y is a direct bond, C₃₋₄cycloalkandiyl, C₃₋₄cycloalkanylidene, -C(R¹⁶)(R¹⁷)- or -

C(R¹⁸)(R¹⁹)C(R²⁰)(R²¹)-;

wherein R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from hydrogen and methyl;

10 or a pharmaceutically-acceptable salt thereof.

Yet a further class of compound is of formula (1) wherein:

Q is a single bond;

R¹ is C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₇cycloalkyl, heterocyclyl, arylC₁₋₃alkyl,

heteroarylC₁₋₃alkyl, C₃₋₇cycloalkylC₁₋₃alkyl, C₃₋₇cycloalkylC₂₋₃alkenyl or C₃₋

15 7cycloalkylC₂₋₃alkynyl, [each of which is optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₃alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl,

C₁₋₃alkoxy, C₁₋₃alkylS(O)_n- (wherein n is 0, 1, 2 or 3), R⁵CON(R^{5'})-, (R^{5'})(R^{5''})NC(O)-,

R^{5'}C(O)O-, R^{5'}OC(O)-, (R^{5'})(R^{5''})NC(O)N(R^{5'''})-, R⁵SO₂N(R^{5''})-, and (R^{5'})(R^{5'''})NSO₂-

(wherein R⁵ is C₁₋₃alkyl optionally substituted by 1, 2 or 3 substituents selected from

20 hydroxyl, halo or cyano; and

R^{5'} and R^{5''} are independently selected from hydrogen and C₁₋₃alkyl optionally substituted by 1, 2 or 3 substituents independently selected from hydroxyl, halo, C₁₋₃alkoxy, carboxy and cyano or R^{5'} and R^{5''} together with the nitrogen atom to which they are attached form a 4-7 membered saturated ring)];

25 R² is selected from heterocyclyl, C₃₋₇cycloalkyl(CH₂)_m-, and C₆₋₁₂polycycloalkyl(CH₂)_m- (wherein m is 0, 1 or 2 and the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R⁶);

R³ is selected from hydrogen, C₁₋₄alkyl C₃₋₅cycloalkyl and C₃₋₅cycloalkylmethyl (each of which is optionally substituted by 1, 2 or 3 fluoro atoms);

30 R² and R³ together with the nitrogen atom to which they are attached form a saturated mono, bicyclic or bridged ring system optionally containing 1 or 2 additional ring heteroatoms selected from nitrogen, oxygen and sulphur and which is optionally fused to a

saturated, partially saturated or unsaturated monocyclic ring wherein the resulting ring system is optionally substituted by 1, 2, or 3 substituents independently selected from R^7 ; R^6 and R^7 are independently selected from hydroxyl, halo, oxo, carboxy, cyano, trifluoromethyl, R^9 , R^9O- , R^9CO- , $R^9C(O)O-$, $R^9CON(R^9)-$, $(R^9)(R^9)NC(O)-$, $(R^9)(R^9)N-$, $R^9S(O)_a-$ wherein a is 0 to 2, $R^9OC(O)-$, $(R^9)(R^9)NSO_2-$, $R^9SO_2N(R^9)-$, $(R^9)(R^9)NC(O)N(R^9)-$, phenyl and heteroaryl [wherein the phenyl and heteroaryl groups are optionally fused to a phenyl, heteroaryl or a saturated or partially-saturated 5- or 6-membered ring optionally containing 1, 2 or 3 heteroatoms independently selected from nitrogen, oxygen and sulphur and the resulting ring system is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, $N-C_{1-4}$ alkylamino, di- $N,N-(C_{1-4}$ alkyl)amino, $N-C_{1-4}$ alkylcarbonyl, di- $N,N-(C_{1-4}$ alkyl)carbonyl, C_{1-4} alkylS(O) $_r-$, C_{1-4} alkylS(O) $_rC_{1-4}$ alkyl (wherein r is 0, 1 or 2)]; R^9 is independently selected from C_{1-3} alkyl optionally substituted by hydroxyl, halo, C_{1-4} alkoxy, carboxy or cyano; R^9 , R^9 and R^9 are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by 1, 2, or 3 substituents independently selected from hydroxyl, halo, C_{1-4} alkoxy, carboxy and cyano);

A is a phenyl or heteroaryl ring (the phenyl or heteroaryl ring being optionally substituted on ring carbon atoms by 1, 2 or 3 R^{10} groups and on an available ring nitrogen in a heteroaryl group by R^{11});

R^{10} is independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, $N-C_{1-4}$ alkylamino, di- $N,N-(C_{1-4}$ alkyl)amino, $N-C_{1-4}$ alkylcarbonyl, di- $N,N-(C_{1-4}$ alkyl)carbonyl, C_{1-4} alkylS(O) $_s-$, C_{1-4} alkylS(O) $_sC_{1-4}$ alkyl (wherein s is 0, 1 or 2)];

R^{11} is independently C_{1-3} alkyl optionally substituted by 1, 2 or 3 fluoro atoms; either X is a direct bond and

Y is a direct bond, C_{3-4} cycloalkandiyl, C_{3-4} cycloalkanylidene, $-C(R^{16})(R^{17})-$ or $-C(R^{18})(R^{19})C(R^{20})(R^{21})-$; or

X is a direct bond, C_{3-4} cycloalkandiyl, C_{3-4} cycloalkanylidene, $-C(R^{12})(R^{13})-$, $-C(R^{12})(R^{13})C(R^{14})(R^{15})-$, $-CH_2O-$ or $-CH_2S(O)-$ (wherein t is 0, 1 or 2) and Y is a direct bond;

wherein R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} and R^{21} are independently selected from hydrogen and methyl;

or a pharmaceutically-acceptable salt thereof.

Yet a further class of compound is of formula (1) wherein:

5 Q is a single bond;

R^1 is C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, heterocyclyl, aryl C_{1-3} alkyl, heteroaryl C_{1-3} alkyl, C_{3-7} cycloalkyl C_{1-3} alkyl, C_{3-7} cycloalkyl C_{2-3} alkenyl or C_{3-7} cycloalkyl C_{2-3} alkynyl, [each of which is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-3} alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl,

10 C_{1-3} alkoxy, C_{1-3} alkylS(O) $_n$ - (wherein n is 0, 1, 2 or 3), R^5 CON($R^{5'}$)-, ($R^{5'}$)($R^{5''}$)NC(O)-, $R^{5'}$ C(O)O-, $R^{5'}$ OC(O)-, ($R^{5'}$)($R^{5''}$)NC(O)N($R^{5'''}$)-, R^5 SO₂N($R^{5''}$)-, and ($R^{5'}$)($R^{5''}$)NSO₂- (wherein R^5 is C_{1-3} alkyl optionally substituted by 1, 2 or 3 substituents selected from hydroxyl, halo or cyano; and

$R^{5'}$ and $R^{5''}$ are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by 1, 2 or 3 substituents independently selected from hydroxyl, halo, C_{1-3} alkoxy, carboxy and cyano or $R^{5'}$ and $R^{5''}$ together with the nitrogen atom to which they are attached form a 4-7 membered saturated ring)];

R^2 is selected from heterocyclyl, C_{3-7} cycloalkyl(CH₂) $_m$ -, and C_{6-12} polycycloalkyl(CH₂) $_m$ - (wherein m is 0, 1 or 2 and the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R^6);

R^3 is selected from hydrogen, C_{1-4} alkyl C_{3-5} cycloalkyl and C_{3-5} cycloalkylmethyl (each of which is optionally substituted by 1, 2 or 3 fluoro atoms);

R^2 and R^3 together with the nitrogen atom to which they are attached form a saturated mono, bicyclic or bridged ring system optionally containing 1 or 2 additional ring

25 heteroatoms selected from nitrogen, oxygen and sulphur and which is optionally fused to a saturated, partially saturated or unsaturated monocyclic ring wherein the resulting ring system is optionally substituted by 1, 2, or 3 substituents independently selected from R^7 ;

R^6 and R^7 are independently selected from hydroxyl, halo, oxo, carboxy, cyano, trifluoromethyl, R^9 , R^9 O-, R^9 CO-, R^9 C(O)O-, R^9 CON($R^{9'}$)-, ($R^{9'}$)($R^{9''}$)NC(O)-,

30 ($R^{9'}$)($R^{9''}$)N-, R^9 S(O) $_a$ - wherein a is 0 to 2, R^9 OC(O)-, ($R^{9'}$)($R^{9''}$)NSO₂-, R^9 SO₂N($R^{9''}$)-, ($R^{9'}$)($R^{9''}$)NC(O)N($R^{9'''}$)-, phenyl and heteroaryl [wherein the phenyl and heteroaryl groups are optionally fused to a phenyl, heteroaryl or a saturated or partially-saturated 5- or 6-

membered ring optionally containing 1, 2 or 3 heteroatoms independently selected from nitrogen, oxygen and sulphur and the resulting ring system is optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₄alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, amino, N-C₁₋₄alkylamino, di-

5 N,N-(C₁₋₄alkyl)amino, N-C₁₋₄alkylcarbamoyl, di-N,N-(C₁₋₄alkyl)carbamoyl,

C₁₋₄alkylS(O)_r-, C₁₋₄alkylS(O)_rC₁₋₄alkyl (wherein r is 0, 1 or 2)];

R⁹ is independently selected from C₁₋₃alkyl optionally substituted by hydroxyl, halo, C₁₋₄alkoxy, carboxy or cyano;

10 R^{9'}, R^{9''} and R^{9'''} are independently selected from hydrogen and C₁₋₃alkyl optionally substituted by 1,2,or 3 substituents independently selected from hydroxyl, halo, C₁₋₄alkoxy, carboxy and cyano);

A is a phenyl or heteroaryl ring (the phenyl or heteroaryl ring being optionally substituted on ring carbon atoms by 1, 2 or 3 R¹⁰ groups and on an available ring nitrogen in a heteroaryl group by R¹¹);

15 R¹⁰ is independently selected from C₁₋₄alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C₁₋₄alkoxy, C₁₋₄alkoxyC₁₋₄alkyl, amino, N-C₁₋₄alkylamino, di-N,N-(C₁₋₄alkyl)amino, N-C₁₋₄alkylcarbamoyl, di-N,N-(C₁₋₄alkyl)carbamoyl, C₁₋₄alkylS(O)_s-, C₁₋₄alkylS(O)_sC₁₋₄alkyl (wherein s is 0, 1 or 2)];

R¹¹ is independently C₁₋₃alkyl optionally substituted by 1, 2 or 3 fluoro atoms;

20 X is a direct bond;

Y is a direct bond, C₃₋₄cycloalkandiyl, C₃₋₄cycloalkanylidene, -C(R¹⁶)(R¹⁷)- or -C(R¹⁸)(R¹⁹)C(R²⁰)(R²¹)-; or

X is a direct bond, C₃₋₄cycloalkandiyl, C₃₋₄cycloalkanylidene, -C(R¹²)(R¹³)-, -C(R¹²)(R¹³)C(R¹⁴)(R¹⁵)-, -CH₂O- or -CH₂S(O)_t- (wherein t is 0, 1 or 2) and

25 Y is a direct bond;

wherein R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²⁰ and R²¹ are independently selected from hydrogen and methyl;

or a pharmaceutically-acceptable salt thereof.

30 In another aspect of the invention, suitable compounds of the invention are any one or more of the Examples or a pharmaceutically-acceptable salt thereof.


In another aspect of the invention, suitable compounds of the invention are any one or more of the following or a pharmaceutically-acceptable salt thereof:

- 4-[4-[[[(1S,3R)-5-hydroxy-2-adamantyl]carbamoyl]-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(1-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 5 4-[4-(N-cyclohexyl-N-methyl-carbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(oxan-4-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[5-propylsulfanyl-4-[3-[2-(trifluoromethyl)phenyl]pyrrolidine-1-carbonyl]pyrazol-1-yl]benzoic acid;
- 4-[4-(cyclohexylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(1-adamantylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(cyclohexyl-methyl-carbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[5-cyclopropyl-4-[(4-hydroxy-1-adamantyl)carbamoyl]pyrazol-1-yl]benzoic acid;
- 10 2-[4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid;
- 2-[4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid;
- 4-(4-cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid;
- 3-(4-cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid;
- 3-[4-(adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-ylmethyl]-benzoic acid;
- 4-[4-(adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-ylmethyl]-benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoic acid;
- 15 4-[4-(2-adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-ethylpyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-propan-2-ylpyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-cyclobutylpyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-methyl-pyrazol-1-yl]benzoic acid;
- 4-(5-tert-butyl-4-(cyclohexylcarbamoyl)-1H-pyrazol-1-yl)benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(1-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[5-cyclohexylsulfanyl-4-[(5-hydroxy-2-adamantyl)carbamoyl]pyrazol-1-yl]benzoic acid;
- 4-[5-cyclohexylsulfanyl-4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoic acid;

- 4-[4-(2-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid;
 methyl 4-[4-(1-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate;
 methyl 4-[5-cyclopentylsulfanyl-4-[(1R,3S)-5-hydroxy-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoate;
 4-[4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanylpyrazol-1-yl]benzoic acid;
 4-[4-(cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid;
 5 4-[4-(cyclohexylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[5-cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-methylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[4-(5-methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-
 10 benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-2-(trifluoromethyl)benzoic acid;
 or
 15 4-[4-(adamantan-2-ylcarbamoyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl]benzoic acid.

Another aspect of the present invention provides a process for preparing a compound of formula (1) or a pharmaceutically acceptable salt thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (1)) comprises any one of processes a) or b):


- 20 a) hydrolysis of an ester of formula (2):



(2)

wherein R²² is an alkyl or aryl group; or

b) converting Z in a compound of the formula (3):



(3)

into a carboxy group, wherein Z is an functional group capable of conversion into a
 5 carboxylic acid;

and thereafter if necessary or desirable:

i) converting a compound of the formula (1) into another compound of the formula (1);

ii) removing any protecting groups;

iii) resolving enantiomers;

10 iv) forming a pharmaceutically-acceptable salt thereof.


Examples of conversions of a compound of Formula (1) into another compound of
 Formula (1), well known to those skilled in the art, include functional group interconversions
 such as hydrolysis, hydrogenation, hydrogenolysis, oxidation or reduction, and/or further
 functionalisation by standard reactions such as amide or metal-catalysed coupling, or
 15 nucleophilic displacement reactions.

Suitable conditions for the above processes a) to b) are as follows.

Process a) may be carried out under either acidic or basic conditions dependant on the
 nature of the ester group (R^{22}) but typically may be carried out under basic conditions, for
 example with aqueous sodium hydroxide, using a suitable solvent such as methanol for
 20 example. Typically the reaction is carried out at ambient temperature, however some esters
 may require cleavage using Microwave or conventional heating, for example at temperatures
 between 30-100°C. Examples of suitable values for R^{22} include methyl, ethyl, tert-butyl,
 phenyl, benzyl and paramethoxybenzyl, particularly methyl or ethyl.

Compounds of formula (2) may be made by processes known in the art and typically
 25 by reaction of a compound of Formula (4) with a compound of Formula (5):


27



Such reactions may be carried out by direct treatment of esters with amines or alternatively by hydrolysis of the R^{23} ester group to an intermediate carboxylic acid followed by conventional amide coupling reactions. Methods and standard conditions for orthogonal removal of ester groups are known persons skilled in the art, for example if R^{22} is ethyl and R^{23} is *tert*-butyl, R^{23} may be selectively cleaved under acidic conditions.

Amide formation can be carried out in a suitable solvent such as dichloromethane for example with the addition of a suitable coupling agent (or combination of agents) such as HOBT and EDCI for example, optionally in the presence of a suitable base such as triethylamine or *N,N*-di-*iso*-propylamine for example. Typically the reaction is carried out at ambient or elevated temperature between 0-60°C.

Compounds of formula (4) in which Q is a sulphur, oxygen or nitrogen atom may be made by processes known in the art and typically by reaction of a compound of Formula (6) with displacement of an appropriate leaving group (L):




wherein L is a suitable leaving group, for example halo or triflate (in particular fluoro or chloro), R^{22} is an alkyl or aryl group and R^{24} is either $-OR^{23}$ or NR^2R^3 .

Displacements may be carried out for example using a suitable nucleophilic reagent, for example propane thiol, in a suitable solvent such as DMF for example in the presence of an

appropriate base, for example sodium hexamethyldisilazide (NaHMDS). Typically the reaction is carried out at ambient temperature when L is chloro, however some reactions may require using Microwave or conventional heating, for example at temperatures between 30-100°C.

Compounds of formula (6) may be made by processes known in the art and typically by functional group interconversion of a compound of Formula (7).




(7)

wherein R^{22} is an alkyl or aryl group and R^{24} is either $-OR^{23}$ or NR^2R^3 .

Examples of such processes are known to the art and may be carried out using a combination of reagents such as *tert*-butylnitrite with a cupric halide in a suitable solvent such as acetonitrile for example. Typically the reaction is carried out at elevated temperature when for example, L is chloro, using Microwave or conventional heating, for example at temperatures between 60-100°C.

Compounds of formula (7) may be made by processes known in the art and typically by reaction of a hydrazine of formula (8) with an enol ether of Formula (9).



(8)

(9)


wherein R^{22} is an alkyl or aryl group, R^{24} is either $-OR^{23}$ or NR^2R^3 and R^{25} is an alkyl group. In particular, R^{25} is methyl or ethyl.

Examples of such processes are known to the art and are typically carried out using a suitable solvent such as ethanol for example in the presence of a suitable base such as DIPEA for example. Typically the reaction is carried out at elevated temperature using

Microwave or conventional heating, for example at temperatures between 60-100°C.

Hydrazines of formula (8) are known in the chemical literature or may be prepared using standard conditions known to those skilled in the art.

Compounds of formula (9) may be made by processes known in the art and typically
 5 by functionalisation of a ketonitrile of Formula (10).




(10)

wherein R^{24} is either $-OR_{23}$ or NR^2R^3 .

Examples of such processes are known to the art and are typically carried out using a
 10 combination of reagents, for example neat acetic anhydride and triethyl orthoformate.


Typically the reaction is carried out at elevated temperature using Microwave or conventional heating, for example at temperatures between 60-100°C. Ketonitriles of formula (10) are known in the chemical literature or may be prepared using standard conditions known to those skilled in the art.

15 Compounds of formula (4) in which Q is a single bond may also be prepared by processes known in the art and typically involve the formation of a functionalised keto ester of Formula (11B) wherein X represents either dialkylamino (such as dimethylamino) or lower alkoxy (such as methoxy or ethoxy) and subsequent reaction with hydrazines of Formula (8)



The ketoesters of formula (11B) are known in the chemical literature or may be prepared using standard conditions known to those skilled in the art, including by reaction of an acid chloride with Meldrum's acid (see for example *J. Org. Chem.* **2001**, 26, 6756) in an inert solvent such as dichloromethane in the presence of a base such as pyridine, followed by reaction of the resultant intermediate with an alcohol HOR²³ (see for example *J. Org. Chem.* **1978**, 43, 2087)

Compounds of formula (2) in which Q is a single bond may also be prepared by processes known in the art and typically involve the formation of a functionalised keto amide of Formula (12) wherein X' represents either dialkylamino (such as dimethylamino) or lower alkoxy (such as methoxy or ethoxy) and subsequent reaction with hydrazines of Formula (8)




Ketoamides of formula (13) are known in the chemical literature or may be prepared using standard conditions known to those skilled in the art, including by reaction of an amine with a suitable acid chloride, optionally prepared in situ from the corresponding acid, or by reaction of an enolate anion, generated by treating a ketone with a strong base (such as LDA), with an appropriate isocyanate.

An example of process b) is the conversion of an aryl halide into an aryl carboxylic acid through the use of metal-catalysed carbonylation. Examples of such processes are known to the art and are carried out in a suitable solvent such as ethanol/dioxane for example using a suitable catalyst, or combination of catalysts, for example, Herrmann's catalyst together with Fu's salt in the presence of a suitable source of carbon monoxide, for example, molybdenum hexacarbonyl or gaseous CO typically in the presence of a suitable

base, or combination of bases for example DMAP/DIPEA. Typically the reaction is carried out at elevated temperature using Microwave or conventional heating, for example at temperatures between 100-180°C. It will be appreciated by those skilled in the art that the choice of solvent will depend on the nature of the product isolated, for example alcoholic solvents will tend to lead to isolation of the ester which may be subsequently cleaved on work-up of the reaction to give the appropriate acid. It will also be appreciated by those skilled in the art that compounds of formula (3) may be accessed by all of the methods used to describe the synthesis of compounds of formula (2).

In addition, a compound of formula (3) may be prepared:

by reaction of a suitably functionalised moiety of formula (3A) with an N unsubstituted pyrazole of formula (3B) to provide the precursor of formula (3)



wherein L' is a leaving group capable of being displaced in an SN2 reaction (when Y becomes joined by an sp³ carbon to the ring) or in SNAr reaction (when Y becomes joined by an sp² carbon to the ring; optionally catalysed by appropriate transition metal catalysts, for example Buchwald displacements) and in which optionally involves deprotonation of the pyrazole (3B) by a base such as potassium t-butoxide. L' is a leaving group such as chloro, bromo or iodo.

The reactions described above may be performed under standard conditions known to the person skilled in the art. The intermediates described above are commercially available, are known in the art or may be prepared by known procedures and/or by the procedures shown above.

It will be appreciated that certain of the various substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a

substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated
5 nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by for example,
10 catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those
15 skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, Protective Groups in Organic Synthesis, John Wiley and Sons, 1991). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an
20 acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an
25 alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a
30 benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is,

for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example hydroxylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possess 11 β HSD1 inhibitory activity. These properties may be assessed using the following assay.

Assays

The conversion of cortisone to the active steroid cortisol by 11 β HSD1 oxo-reductase activity, can be measured using a competitive homogeneous time resolved fluorescence assay (HTRF) (CisBio International, R&D, Administration and Europe Office, In Vitro Technologies — HTRF® / Bioassays BP 84175, 30204 Bagnols/Cèze Cedex, France. Cortisol bulk HTRF kit: Cat No. 62CORPEC).

The evaluation of compounds described herein was carried out using a baculovirus expressed N terminal 6-His tagged full length human 11 β HSD1 enzyme(*1). The enzyme was purified from a detergent solubilised cell lysate, using a copper chelate column.

Inhibitors of 11 β HSD1 reduce the conversion of cortisone to cortisol, which is identified by an increase in signal, in the above assay.

Compounds to be tested were dissolved in dimethyl sulphoxide (DMSO) to 10mM and diluted further in assay buffer containing 1% DMSO to 10 fold the final assay concentration. Diluted compounds were then plated into black 384 well plates (Matrix, Hudson NH, USA).

5 The assay was carried out in a total volume of 20µl consisting of cortisone (Sigma, Poole, Dorset, UK, 160nM), glucose-6-phosphate (Roche Diagnostics, 1mM), NADPH (Sigma, Poole, Dorset, 100µM), glucose-6-phosphate dehydrogenase (Roche Diagnostics, 12.5µg/ml), EDTA (Sigma, Poole, Dorset, UK, 1mM), assay buffer (K₂HPO₄/KH₂PO₄, 100mM) pH 7.5, recombinant 11βHSD1 [using an appropriate dilution to give a viable
10 assay window – an example of a suitable dilution may be 1 in 1000 dilution of stock enzyme] plus test compound. The assay plates were incubated for 25 minutes at 37°C after which time the reaction was stopped by the addition of 10µl of 0.5mM glycerolacetic acid plus conjugated cortisol(XL665 or D2). 10µl of anti-cortisol Cryptate was then added and the plates sealed and incubated for 6 hours at room temperature. Fluorescence at 665nm
15 and 620nm was measured and the 665nm:620nm ratio calculated using an Envision plate reader.

These data were then used to calculate IC₅₀ values for each compound (Origin 7.5, Microcal software, Northampton MA, USA) and/or the % inhibition at 30µM of compound.

*1 The Journal of Biological Chemistry, Vol. 26, No 25, pp16653 – 16658

20 Compounds of the present invention typically show an IC₅₀ of less than 30µM, and preferably less than 5 µM.

For example, the following results were obtained:

Ex. No.	IC50 (uM)	Ex. No.	IC50 (uM)	Ex. No.	IC50 (uM)	Ex. No.	IC50 (uM)
3	0.012	17	0.004	24	0.005	38	0.007
6	3.505	20	0.006	27	0.008	41	0.003
4	0.034	22	0.008	33	0.009	43	0.185
8	0.161	23	0.004	36	0.004	44	0.005

The following table displays %inhibition of human 11- β HSD at a test concentration of 30 μ M of compound

	%		%		%		%
Ex. No.	@30uM	Ex. No.	@30uM	Ex. No.	@30uM	Ex. No.	@30uM
1	91	13	96	25	90	37	94
2	93	14	89	26	100	38	91
3	105	15	92	27	95	39	98
4	114	16	91	28	87	40	99
5	90	17	95	29	91	41	91
6	80	18	89	30	91	42	94
7	91	19	90	31	105	43	97
8	94	20	90	32	103	44	94
9	78	21	91	33	97	45	94
10	84	22	98	34	100	46	98
11	82	23	102	35	94	47	90
12	92	24	104	36	89		

The oral bioavailability of the compounds of the invention may be tested as follows:

5 Determination of Bioavailability in PK Studies

Compounds are dosed intravenously at 2mg/kg (2ml/kg) and orally at 5mg/kg (5ml/kg) in a 25% HPBCD in sorrensons buffer pH 5.5 formulation. Blood samples (200ul) are taken Predose, 0.25, 0.5, 1, 2, 3, 4, 5, 6, 8 and 24 h post dose for both routes and plasma prepared by centrifugation. Plasma samples are analysed as below. PK parameters (clearance, 10 volume of distribution, bioavailability, fraction absorbed etc.) are calculated by standard PK methods using suitable PK software (WinNon-Lin).

Bioanalysis of plasma samples

The guidelines described are for the manual preparation of plasma samples following single compound or cassette dosing of project compounds to all PK species used within 15 discovery DMPK. Analysis by open access (LC-MS/MS) or manual approaches (LC-MS) is described.

Contents

1. Materials
2. Generic Extraction Method
3. Example Sample List Using Generic Plate Layout
- 5 4. Open Access Batch Submission and System Checks
5. Acceptance Criteria for Batch Pass

1. Materials

Solvents: Methanol, acetonitrile and DMSO

Water: Purified or HPLC grade

10 1ml shallow 96-well plates OR eppendorf tubes

2ml deep well 96-well plates plus lids

Blank (control) plasma

2. Generic Extraction Method

Solubilise compound(s) to 1mg/ml using DMSO taking into account salt factors if any.

15 The DMSO stock(s) may be used to make all calibration & quality control (QC) samples:

2.i Single compound analysis

2.i.a Preparation of calibration and QC samples:

1. Prepare standard solutions as follows:

Stock diluted ng/ml	Volume methanol ml	Volume stock ml	Standard conc. ng/ml	Post plasma dilution conc. ng/ml
1mg/ml	0.9	0.1	100,000	10,000
100,000	0.5	0.5	50,000	5,000
50,000	0.75	0.5	20,000	2,000
20,000	0.5	0.5	10,000	1,000
10,000	0.5	0.5	5,000	500
5,000	2	0.5	1,000	100
1,000	0.5	0.5	500	50
500	0.75	0.5	200	20
200	0.5	0.5	100	10
100	0.5	0.5	50	5
50	0.5	0.5	10	1

2. Transfer 50ul blank plasma to a well of a 1ml 96-well plate (shallow well)
3. Transfer 5ul of each of the standard solutions to further wells of the plate
4. Add 50ul blank plasma to each of these wells.
5. To generate the QC samples, add three aliquots of 5ul of the 100ng/ml, 1000ng/ml
5 and 10,000ng/ml standard solutions to the plate (3 QCs at each concentration).
6. Add 50ul blank plasma to each of these.
7. Transfer 50ul of each PK sample to the 1ml 96-well plate
8. Add 5ul methanol (- compound) to each of the PK samples
9. Ensure all dose formulations are well mixed by vortex mixing.
10. Dilute intravenous (IV) and oral dose (PO) formulations of expected concentration
10 to 10ug/ml in methanol. (For example, a formulation made to an expected
concentration of 2 mg/ml would be diluted 1:200 to give 10ug/ml solution).
11. Add 6x 50 ul aliquots of plasma to the plate. Add 5 ul of diluted IV formulation to
three of the wells, repeat with PO formulation and remaining 3 wells.
12. Precipitate proteins by adding 100ul acetonitrile containing a project related
15 internal standard (at 1ug/ml) to all calibration, QC, PK and formulation samples.
13. Vortex mix the plate before centrifugation at 4,000g for 10 minutes.
14. Transfer 100ul of the supernatant to the wells of a 2ml 96-well plate (see following
plate map). Care should be taken not to disturb the pellet.
15. Add ~1.5ml of 50:50 Methanol: Water into the last well.
20
16. For analysis on triple quad systems: add 400ul water (HPLC grade) to each sample.
Gently mix.
17. Add 100ul of the 100,000ng/ml stock of each of the standard solutions to the 2ml
plate and add 900ul water. Add a sample of internal standard to a further well (see
25 plate map). *These are for compound tuning (denoted on the plate map as tune
solutions)*
18. For analysis on platform systems: add 100ul water (HPLC grade) to each sample.
Gently mix.
19. Manually tune all compounds using compound solutions prepared to 5,000ng/ml
30 (add 100ul of the 50,000ng/ml standard solutions to 900ul water)

2.ii Cassette dose analysis

2.ii.a Preparation of calibration and QC samples:

Note: For cassette dosing, the amount of methanol required to dilute the 1mg/ml stock will be adjusted according to the number of compounds present.

1. Add 100ul of each 1mg/ml stock required to a vial.
2. Add the required volume of methanol to yield a total volume of 1ml.
3. Perform all further steps as for single compound analysis (steps 2 –16 above).

2.iii In cases where PK samples exceed the Upper limit of Quantification (ULOQ).

1. Prepare a further calibration curve and QC samples as above (steps 1 – 6).
2. Transfer <50ul (e.g. 25ul) of the PK samples that exceed the ULOQ.
3. Add enough control plasma to these samples to yield a final plasma volume of 50ul. Make a note of the dilution made.
4. Transfer 50ul of all remaining PK samples.
5. Prepare all formulation samples and extract all samples as described above. (steps 8 – 16)

Note: Upper concentrations used to generate the calibration curve may be reviewed, however, care must be taken to avoid saturation of the HPLC column or MS equipment. It is for this reason that dilution of PK samples is recommended.

2.iv In cases of poor sensitivity (high Lower Limit of Quantification).

Note: High LLOQ is taken as when most of the plasma concentrations lie below the lower limit of quantification or where the LLOQ is greater the 10ng/ml. The following methods should be applied when either of these scenarios is encountered.

According to a further aspect of the invention there is provided a pharmaceutical composition, which comprises a compound of the Examples, or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous,

intramuscular or intramuscular dosing or as a suppository for rectal dosing). In general, compositions in a form suitable for oral use are preferred.

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

Suitable pharmaceutically-acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol; or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene

sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl p-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

5 Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These
10 compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above.
15 Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these.
20 Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, an esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain
25 sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable
30 aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile

injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol
5 containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

For further information on formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial
10 Board), Pergamon Press 1990.

The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral
15 administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch;
20 Chairman of Editorial Board), Pergamon Press 1990.

We have found that the compounds defined in the present invention, or a pharmaceutically-acceptable salt thereof, are effective 11 β HSD1 inhibitors, and accordingly have value in the treatment of disease states associated with metabolic syndrome.

25 It is to be understood that where the term "metabolic syndrome" is used herein, this relates to metabolic syndrome as defined in 1) and/or 2) or any other recognised definition of this syndrome. Synonyms for "metabolic syndrome" used in the art include Reaven's Syndrome, Insulin Resistance Syndrome and Syndrome X. It is to be understood that where the term "metabolic syndrome" is used herein it also refers to Reaven's Syndrome,
30 Insulin Resistance Syndrome and Syndrome X.

According to a further aspect of the present invention there is provided a compound of formula (1), or a pharmaceutically-acceptable salt thereof, as defined hereinbefore for

use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

Thus according to this aspect of the invention there is provided a compound of formula (1), or a pharmaceutically-acceptable salt thereof, as defined hereinbefore for use as a medicament.

According to another feature of the invention there is provided the use of a compound of formula (1), or a pharmaceutically-acceptable salt thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of an 11 β HSD1 inhibitory effect in a warm-blooded animal, such as man.

Where production of or producing an 11 β HSD1 inhibitory effect is referred to suitably this refers to the treatment of metabolic syndrome. Alternatively, where production of an 11 β HSD1 inhibitory effect is referred to this refers to the treatment of diabetes, obesity, hyperlipidaemia, hyperglycaemia, hyperinsulinemia or hypertension, particularly diabetes and obesity. Alternatively, where production of an 11 β HSD1 inhibitory effect is referred to this refers to the treatment of glaucoma, osteoporosis, tuberculosis, dementia, cognitive disorders or depression.

Alternatively, where production of an 11 β HSD1 inhibitory effect is referred to this refers to the treatment of cognitive disorders, such as improving the cognitive ability of an individual, for example by improvement of verbal fluency, verbal memory or logical memory, or for treatment of mild cognitive disorders. See for example WO03/086410 and references contained therein, and Proceedings of National Academy of Sciences (PNAS), 2001, 98(8), 4717-4721.

Alternatively, where production of an 11 β HSD1 inhibitory effect is referred to this refers to the treatment of, delaying the onset of and/or reducing the risk of atherosclerosis – see for example J. Experimental Medicine, 2005, 202(4), 517-527.

Alternatively, where production of an 11 β HSD1 inhibitory effect is referred to this refers to the treatment of Alzheimers and/or neurodegenerative disorders.

According to a further feature of this aspect of the invention there is provided a method for producing an 11 β HSD1 inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (1), or a pharmaceutically-acceptable salt thereof.

In addition to their use in therapeutic medicine, the compounds of formula (1), or a pharmaceutically-salt thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of 11 β HSD1 in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search for new therapeutic agents.

The inhibition of 11 β HSD1 described herein may be applied as a sole therapy or may involve, in addition to the subject of the present invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. Simultaneous treatment may be in a single tablet or in separate tablets. For example agents than might be co-administered with 11 β HSD1 inhibitors, particularly those of the present invention, may include the following main categories of treatment:

- 1) Insulin and insulin analogues;
- 2) Insulin secretagogues including sulphonylureas (for example glibenclamide, glipizide), prandial glucose regulators (for example repaglinide, nateglinide), glucagon-like peptide 1 agonist (GLP1 agonist) (for example exenatide, liraglutide) and dipeptidyl peptidase IV inhibitors (DPP-IV inhibitors);
- 3) Insulin sensitising agents including PPAR γ agonists (for example pioglitazone and rosiglitazone);
- 4) Agents that suppress hepatic glucose output (for example metformin);
- 5) Agents designed to reduce the absorption of glucose from the intestine (for example acarbose);
- 6) Agents designed to treat the complications of prolonged hyperglycaemia; e.g. aldose reductase inhibitors
- 7) Other anti-diabetic agents including phosphotyrosine phosphatase inhibitors, glucose 6 - phosphatase inhibitors, glucagon receptor antagonists, glucokinase activators, glycogen phosphorylase inhibitors, fructose 1,6 bisphosphatase inhibitors, glutamine:fructose -6-phosphate amidotransferase inhibitors
- 8) Anti-obesity agents (for example sibutramine and orlistat);
- 9) Anti- dyslipidaemia agents such as, HMG-CoA reductase inhibitors (statins, eg pravastatin); PPAR α agonists (fibrates, eg gemfibrozil); bile acid sequestrants (cholestyramine); cholesterol absorption inhibitors (plant stanols, synthetic inhibitors);

ileal bile acid absorption inhibitors (IBATi), cholesterol ester transfer protein inhibitors and nicotinic acid and analogues (niacin and slow release formulations);

- 10 10) Antihypertensive agents such as, β blockers (eg atenolol, inderal); ACE inhibitors (eg lisinopril); calcium antagonists (eg. nifedipine); angiotensin receptor antagonists (eg
5 candesartan), α antagonists and diuretic agents (eg. furosemide, benzthiazide);
- 11) Haemostasis modulators such as, antithrombotics, activators of fibrinolysis and antiplatelet agents; thrombin antagonists; factor Xa inhibitors; factor VIIa inhibitors; antiplatelet agents (eg. aspirin, clopidogrel); anticoagulants (heparin and Low molecular weight analogues, hirudin) and warfarin;
- 10 12) Anti-inflammatory agents, such as non-steroidal anti-inflammatory drugs (eg. aspirin) and steroidal anti-inflammatory agents (eg. cortisone); and
- 13) Agents that prevent the reabsorption of glucose by the kidney (SGLT inhibitors).

15 In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

Examples

The invention will now be illustrated by the following Examples in which, unless stated otherwise:

- 20 (i) temperatures are given in degrees Celsius ($^{\circ}\text{C}$); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25 $^{\circ}\text{C}$ and under an atmosphere of an inert gas such as argon;
- (ii) evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pa; 4.5-30 mmHg) with a bath temperature of up to 60 $^{\circ}\text{C}$;
- (iii) chromatography means flash chromatography on silica gel;
- 25 (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- 30 (vi) where given, NMR data (^1H) is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS), determined at 300 or 400 MHz (unless otherwise stated) using perdeuterio dimethyl sulfoxide

(DMSO-*d*₆) as solvent, unless otherwise stated; peak multiplicities are shown thus: s, singlet; d, doublet; dd, doublet of doublets; dt, doublet of triplets; dm, doublet of multiplets; t, triplet; m, multiplet; br, broad;


(vii) chemical symbols have their usual meanings; SI units and symbols are used;

5 (viii) solvent ratios are given in volume : volume (v/v) terms;

(ix) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionisation (CI) mode using a direct exposure probe; where indicated ionisation was effected by electron impact (EI), fast atom bombardment (FAB) or electrospray (ESP); values for *m/z* are given; generally, only ions which indicate the parent mass are reported;

10 (x) The following abbreviations may be used below or in the process section hereinbefore:


	Et ₂ O	diethyl ether
	DMF	dimethylformamide
	DCM	dichloromethane
	DME	1,2-dimethoxyethane
15	MeOH	methanol
	EtOH	ethanol
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
	DMSO	dimethylsulfoxide
20	HOBT	1-hydroxybenzotriazole
	EDCI (EDAC)	1-ethyl-3-(3-dimethylaminopropyl)carbodi-imide hydrochloride
	DIPEA	diisopropylethylamine
	DMAP	4-dimethylaminopyridine
25	DEAD	diethyl azodicarboxylate
	EtOAc	ethyl acetate
	MgSO ₄	magnesium sulfate
	MTBE	methyl tert-butyl ether
	NaHMDS	sodium hexamethyldisilazide

Example 1**4-[4-((1R,2S,3S,5S)-5-Hydroxy-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid**

5 4-[4-((1R,2S,3S,5S)-5-Hydroxy-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid methyl ester (0.34mmol) (**Intermediate # 1**) was dissolved in methanol (10mL) and treated with 2M sodium hydroxide solution (0.84mL, 1.7mmol). The mixture was stirred at ambient temperature for 24h and then the methanol was removed by evaporation under reduced pressure. The residue was dissolved in water (10mL), acidified to pH4 with 2M HCl and extracted with EtOAc (2x10mL). The combined extracts were washed with water (10mL) and brine (10mL) and dried (MgSO₄) and evaporated to leave the title compound as a white solid. (74mg, 48%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.66 (3H, t), 1.18 - 1.30 (2H, m), 1.42 (2H, d), 1.58-1.80 (6H, m), 1.88 (2H, d), 2.06 (3H, s), 2.62 (2H, t), 3.95 - 4.06 (1H, m), 4.42 (1H, s),
 15 7.71 (2H, d), 7.95 (1H, d), 8.09 (2H, d), 8.16 (1H, s), 13.19 (1H, s)

MS m/z 456 M+H

Example 2**4-[4-(2-Adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid**

20 Methyl 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate (190mg, 0.42mmol) (**Intermediate #2**) was dissolved in methanol (10mL) and treated at

ambient temperature with 2M aqueous sodium hydroxide solution (1.05mL, 2.1mmol). The mixture was stirred at ambient temperature for 18h. and then heated to 65°C for a further 2h. Methanol was removed by evaporation under reduced pressure and the clear solution diluted with water (25ml). 2M HCl was added to pH4 and the mixture extracted with ethyl acetate (2x25mL). The combined extracts were washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated to give the title compound as a white solid.


(174mg, 94%)

¹H NMR (300.073 MHz, DMSO-d₆) δ0.65 (3H, t), 1.17 - 1.29 (2H, m), 1.60 (2H, d), 1.73 (2H, s), 1.83 (6H, s), 1.91-2.05 (4H, m), 2.62 (2H, t), 4.09 (1H, d), 7.75 (2H, d), 8.03 (1H, d), 8.13 (2H, d), 8.20 (1H, s)

MS m/z 440 M+H

Example 3

4-[4-(1-Adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid




Methyl 4-[4-(1-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate (143mg, 0.32mmol) (**Intermediate#3**) was dissolved in methanol and treated at ambient temperature with 2M aqueous sodium hydroxide solution. The mixture was stirred at ambient for 18h. Methanol was removed by evaporation under reduced pressure and the clear solution diluted with water (25ml). 2M HCl was added to pH4 and the mixture extracted with ethyl acetate (2x25mL). The combined extracts were washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated to give the title compound as a white solid (132mg, 94%).

¹H NMR (300.073 MHz, DMSO-d₆) δ0.69 (3H, t), 1.25 (2H, q), 1.66 (6H, s), 2.06 (9H, s), 2.65 (2H, t), 7.51 (1H, s), 7.66 - 7.69 (2H, m), 8.07 - 8.10 (3H, m)

MS m/z 440 M+H

Example 4

4-[4-(N-Cyclohexyl-N-methyl-carbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid




- 5 Methyl 4-[4-(cyclohexyl-methyl-carbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate (**Intermediate#4**) (162mg, 0.39mmol) was dissolved in methanol (10mL) and treated at ambient temperature with 2M aqueous sodium hydroxide solution (0.96mL, 1.95mmol). The mixture was stirred at ambient for 18h. Methanol was removed by evaporation under reduced pressure and the clear solution diluted with water (25ml).
- 10 2M HCl was added to pH4 and the mixture extracted with ethyl acetate (2x25mL). The combined extracts were washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated to give the title compound as a white solid foam (150mg, 96%).
- ¹H NMR (300.073 MHz, DMSO-d₆) δ 0.68 (3H, t), 1.10-1.41 (4H, m), 1.42 - 1.85 (8H, m), 2.58 (2H, t), 2.86 (3H, s), 3.45-3.60 (0.5H, m), 4.21-4.38 (0.5H, m), 7.74 (2H, d), 7.89 (1H, s), 8.09 (2H, d)
- 15

MS m/z 402 M+H

Example 5

4-[4-(Oxan-4-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid




Methyl 4-[4-(oxan-4-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate (**Intermediate#5**) (104mg, 0.26mmol) was dissolved in methanol (5mL) and treated at ambient temperature with 2M aqueous sodium hydroxide solution (0.65mL, 1.29 mmol). The mixture was stirred at ambient for 18h. Methanol was removed by evaporation under reduced pressure and the clear solution diluted with water (25ml). 2M HCl was added to pH4 and the mixture extracted with ethyl acetate (2x25mL). The combined extracts were washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated to give the product as a white solid (89mg, 85%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.68 (3H, t), 1.19 - 1.31 (2H, m), 1.47 - 1.56 (2H, m), 1.77 - 1.81 (2H, m), 2.71 (2H, t), 3.29 - 3.43 (2H, m), 3.85 - 3.89 (2H, m), 3.94 - 4.04 (1H, m), 7.67 (2H, d), 8.03 (1H, d), 8.09 (2H, d), 8.16 (1H, s)

MS m/z 390 M+H

Example 6


4-[5-Propylsulfanyl-4-[3-[2-(trifluoromethyl)phenyl]pyrrolidine-1-carbonyl]pyrazol-1-yl]benzoic acid



Methyl 4-[5-propylsulfanyl-4-[3-[2-(trifluoromethyl)phenyl]pyrrolidine-1-carbonyl]pyrazol-1-yl]benzoate (**Intermediate#6**) (177mg, 0.34mmol) was stirred at ambient temperature for 18h in a mixture of methanol (5mL) and 2M sodium hydroxide (0.855mL, 1.71mmol) for 18h. The reaction mixture was evaporated to remove the methanol. The residue was dissolved in water (20mL) and acidified to pH4 with 2M HCl. The resulting white precipitate was recovered by filtration, washed with water and dried under vacuum to give the title compound as a white solid (116mg, 68%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.64 (3H, q), 1.14 - 1.26 (2H, m), 1.67 - 1.76 (1H, m), 1.89 - 1.98 (2H, m), 2.42 - 2.59 (3H, m), 3.62-3.82 (1H, m), 3.93 - 4.01 (1H, m), 5.24 - 5.43 (1H, m), 7.43 - 7.48 (2H, m), 7.58 - 7.75 (4H, m), 8.00-8.20 (3H, m), 13.18 (1H, s)

MS m/z 503 M+H


Example 7**4-[4-(Cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid**

Methyl 4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate

(363mg, 0.94mmol) (**Intermediate # 11**) was dissolved in methanol (20mL) and treated at ambient temperature with 2M aqueous sodium hydroxide solution (2.35mL). The mixture was stirred at ambient temperature for 18h and then volatiles were removed by evaporation under reduced pressure. The residue was dissolved in water (25ml) and 2M HCl was added until the pH=4. The mixture was extracted with ethyl acetate (2x25mL) and the combined extracts were washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated to give the title product as a white solid. (300mg 85%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.68 (3H, t), 1.14 - 1.39 (7H, m), 1.57 - 1.61 (1H, m), 1.72 (2H, d), 1.84 (2H, d), 2.70 (2H, t), 3.75 - 3.78 (1H, m), 7.65 - 7.69 (2H, m), 7.90 - 7.93 (1H, m), 8.07 - 8.13 (2H, m), 8.14 (1H, s), 13.20 (1H, s)

MS m/z 388 M+H

Example 8**3-[4-(Cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid**

Methyl 3-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate


(175mg, 0.45), (**Intermediate #12**) was dissolved in methanol (10mL), treated with 2M aqueous sodium hydroxide solution (1.125mL) and stirred at ambient temperature for 18h.

Methanol was removed by evaporation under reduced pressure and the clear solution diluted with water (25ml). 2M HCl was added to pH4 and the mixture extracted with ethyl acetate (2x25mL). The combined extracts were washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated to give the title product as a white solid. (115mg, 65%)

¹H NMR (300.073 MHz, DMSO-d₆) δ0.67 (3H, t), 1.14 - 1.39 (7H, m), 1.57 - 1.61 (1H, m), 1.70 (2H, d), 1.82 - 1.85 (2H, m), 2.70 (2H, t), 3.74 - 3.79 (1H, m), 7.66 - 7.71 (1H, m), 7.78 - 7.82 (1H, m), 7.91 (1H, d), 8.03 - 8.06 (2H, m), 8.13 (1H, s), 13.29 (1H, s)
MS m/z 388 M+H

Example 9


4-[4-(Cyclohexylcarbamoyl)-5-propyl-pyrazol-1-yl]benzoic acid



Ethyl 4-[4-(cyclohexylcarbamoyl)-5-propyl-pyrazol-1-yl]benzoate (130mg, 0.34mmol) (**Intermediate # 21**) was dissolved in methanol (10mL) and treated at ambient temperature with 2M aqueous sodium hydroxide solution (2.5mL). The mixture was stirred at ambient temperature for 18h and then methanol was removed by evaporation under reduced pressure. The remaining aqueous solution was acidified with 2M aqueous hydrochloric acid to pH=2. A white solid precipitated out of solution, it was filtered, dried under vacuum and identified as the desired product (120 mg, quantitative reaction).

¹H NMR (400.13 MHz, DMSO-d₆) δ0.73 (3H, t), 1.12 (1H, m), 1.23 - 1.31 (1H, m), 1.28 (3H, m), 1.36 - 1.43 (2H, m), 1.59 - 1.63 (1H, m), 1.73 (2H, d), 1.80 (2H, s), 2.97 (2H, t), 3.73 (1H, m), 7.59 - 7.61 (2H, d), 7.95 (1H, d), 8.09 - 8.11 (2H, d), 8.21 (1H, s), 13.27 (1H, s)

MS m/z 356 M+H

Example 10**4-[4-(Cyclohexyl-methyl-carbamoyl)-5-propyl-pyrazol-1-yl]benzoic acid**


Ethyl 4-[4-(cyclohexyl-methyl-carbamoyl)-5-propyl-pyrazol-1-yl]benzoate

- 5 (Intermediate # 23, 105mg, 0.26mmol) was dissolved in methanol (10mL) and treated at ambient temperature with 2M aqueous sodium hydroxide solution (2.5mL). The mixture was stirred at ambient temperature for 18h and then methanol was removed by evaporation under reduced pressure. The remaining aqueous solution was acidified with 2M aqueous hydrochloric acid to pH=2. A white solid precipitated out of solution, it was filtered, dried
- 10 under vacuum and identified as the desired product (96 mg, quantitative reaction).

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.71 (3H, t), 1.12 - 1.50 (6H, m), 1.59 - 1.77 (6H, m), 2.79 (2H, t), 2.89 (3H, s), 7.45 - 7.65 (1H, m), 7.67 (2H, d), 7.84-7.86 (1H, m), 8.10 (2H, d), 13.24 (1H, s)


MS m/z 370 M+H




15 **Example 11**

4-[4-(cyclohexylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid

1-(4-bromophenyl)-N-cyclohexyl-5-cyclopropyl-pyrazole-4-carboxamide
(Intermediate#25) (111mg, 0.29mmol), molybdenum hexacarbonyl (38mg, 0.14 mmol),
 DMAP (70mg, 0.57mmol), DIPEA (74mg, 0.57mmol) and trans-Di(-acetato)bis[o-(di-o-
 tolylphosphino)benzyl]dipalladium (II) (14mg, 0.015mmol) were added to a microwave
 5 vial and suspended in a mixture of dioxan (2mL) and ethanol (2mL). The vial was capped
 and heated to 150°C for 1h. The reaction mixture was evaporated to dryness and the
 residue dissolved in DCM (10mL). 2M HCl (10ml) was added and the mixture shaken and
 passed through a phase separating filter. The DCM solution was dry loaded onto Ceelite
 and the product recovered by flash chromatography on silica gel (elution gradient 0- 50%
 10 EtOAc in Hexane). Pure fractions were combined and evaporated to give the product as a
 white solid. The white solid was dissolved in methanol (5mL), treated with 2M NaOH
 solution (1mL) and stirred at ambient temperature for 5h. The reaction mixture was
 concentrated under reduced pressure, diluted with water (20mL), washed with ether
 (2x10mL), acidified to pH4 with 2M HCl and extracted with EtOAc (3x10mL). The
 15 combined extracts were washed with water (10mL), dried (MgSO₄) and evaporated to
 afford 4-[4-(cyclohexylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid
 as white solid. (40mg, 40%)


The following Examples were prepared in a similar manner to Example #11, using
 an appropriate bromophenyl starting material:

Structure	Ex	Name	¹ H NMR δ	MS m/e MH ⁺
	12	4-[4-(2-adamantylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid	¹ H NMR (300.073 MHz, DMSO-d ₆) δ 0.41 - 0.46 (2H, m), 0.85 - 0.91 (2H, m), 1.56 (2H, d), 1.73 (2H, s), 1.85 (6H, d), 1.98 (2H, d), 2.08 (2H, d), 2.21 - 2.30 (1H, m), 4.05 (1H, d), 7.58 (1H, d), 7.77 (2H, d), 7.94 (1H, s), 8.09 (2H, d)	406

Structure	Ex	Name	¹ H NMR δ	MS m/e MH ⁺
	13	4-[4-(1-adamantylcarbamoyl)-5-cyclopropylpyrazol-1-yl]benzoic acid	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 0.42 - 0.46 (2H, m), 0.87 - 0.92 (2H, m), 1.67 (6H, s), 2.07 (9H, s), 2.14 - 2.21 (1H, m), 7.17 (1H, s), 7.73 - 7.76 (2H, m), 7.86 (1H, s), 8.07 - 8.10 (2H, m)	406
	14	4-[4-(cyclohexylmethyl-carbamoyl)-5-cyclopropylpyrazol-1-yl]benzoic acid	¹ H NMR (300.073 MHz, DMSO-d ₆) δ 0.45 (2H, s), 0.83 (2H, d), 1.08 - 1.85 (10H, m), 2.06 (1H, s), 2.86 (3H, s), 3.57 (0.4H, s), 4.15 (0.6H, s), 7.68 (1H, s), 7.82 (2H, d), 8.07 (2H, d), 13.0 (1H, s)	368
	15	4-[5-cyclopropyl-4-[(4-hydroxy-1-adamantyl)carbamoyl]pyrazol-1-yl]benzoic acid	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 0.42 - 0.46 (2H, m), 0.85 - 0.90 (2H, m), 1.38 (2H, d), 1.65 (5H, d), 1.75 (2H, d), 1.96 (2H, d), 2.03 (1H, s), 2.08 (2H, s), 2.21 - 2.28 (1H, m), 3.98 (1H, d), 4.40 (1H, s), 7.53 (1H, d), 7.75 - 7.78 (2H, m), 7.94 (1H, s), 8.08 - 8.11 (2H, m), 13.1 (1H	422

Example 16


2-[4-[4-(Cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid



Methyl 2-[4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetate (**Intermediate#33**) (210mg, 0.51mmol) was dissolved in methanol (10mL) and treated at ambient temperature with a 2M solution of sodium hydroxide (1.27mL, 2.53mmol). The mixture was stirred overnight and then methanol was removed by evaporation under reduced pressure. The clear aqueous solution was diluted with water (20mL) and acidified to pH3 with 2M HCl. The resulting white precipitate was extracted into ethyl acetate (2x20mL). The combined extracts were washed with brine (10mL), dried (MgSO₄) and evaporated to give the crude product. The crude product was purified by preparative HPLC using decreasingly polar mixtures of water (containing 0.1% NH₃) and MeCN as eluents. Fractions containing the desired compound were evaporated to dryness to afford 2-[4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid. (84mg, 41%).

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.72 (3H, t), 1.20 - 1.36 (7H, m), 1.60 (1H, d), 1.73 (2H, d), 1.85 (2H, d), 2.69 (2H, t), 3.69 (2H, s), 3.77 (1H, d), 7.42 - 7.47 (4H, m), 7.86 (1H, d), 8.10 (1H, s)


MS m/z (ESI+) (M+H)+ 402

Example 17**2-[4-[4-(2-Adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid**

Methyl 2-[4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetate (**Intermediate #34**) (220mg, 0.47mmol) was dissolved in methanol (10mL) and treated at ambient temperature with a 2M solution of sodium hydroxide (1.17mL, 2.35mmol). The mixture was stirred overnight and then methanol was removed by evaporation under reduced pressure. The clear aqueous solution was diluted with water (20mL) and acidified to pH3 with 2M HCl. The resulting white precipitate was extracted into ethyl acetate (2x20mL). The combined extracts were washed with brine (10mL), dried (MgSO₄) and evaporated to give the crude product. The crude product was purified by preparative HPLC using decreasingly polar mixtures of water (containing 0.1% NH₃) and MeCN as eluents. Fractions containing the desired compound were evaporated to dryness to afford 2-[4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid (66mg, 31%).

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.70 (3H, t), 1.23 - 1.32 (2H, m), 1.62 (2H, d), 1.74 (2H, s), 1.86 (6H, d), 1.99 (4H, d), 2.62 (2H, t), 3.70 (2H, s), 4.11 (1H, d), 7.45 (2H, d), 7.50 - 7.52 (2H, m), 8.02 (1H, d), 8.12 (1H, s)

MS m/z (ESI+) (M+H)⁺ 454




Example 18**4-(4-Cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid**


A solution of 4-(4-Cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid methyl ester (116mg, 0.28mmol) (**Intermediate #40**) and LiOH (47mg, 1.12mmol) in MeOH (2ml) / Water (1ml) was stirred overnight at ambient temperature. The bulk of the MeOH was removed in vacuo and the resulting solution was treated with citric acid (~10ml) then extracted with EtOAc (2 x ~15ml). The organic layers were combined, washed with brine (~10ml), dried (MgSO₄), filtered and evaporated to yield the title compound as a white solid (102mg, 91%).

¹H NMR (700.03 MHz, CDCl₃) δ 0.89 (3H, t), 1.22 - 1.30 (4H, m), 1.41 - 1.46 (2H, m), 1.48 - 1.53 (2H, m), 1.62 - 1.64 (1H, m), 1.71 - 1.73 (1H, m), 1.98 - 2.00 (2H, m), 2.56 (2H, t), 3.99 - 4.04 (1H, m), 5.59 (2H, s), 7.27 (2H, d), 7.37 (1H, d), 8.05 (2H, d), 8.16 (1H, s)

MS m/e MH⁺ 402.

The following examples were made using the above procedure, replacing 4-(4-Cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid methyl ester with a corresponding starting material

Structure	Example #	Name	NMR	[M+H] ⁺
	19	3-(4-Cyclohexylcarbonyl-5-propylsulfanylmethyl)-pyrazol-1-ylmethylbenzoic acid	¹ H NMR (300.072 MHz, CDCl ₃) δ 0.81 – 1.85 (3H, t), 1.11 - 1.31 (3H, m), 1.35 - 1.52 (4H, m), 1.55 - 1.69 (3H, m), 1.87 – 1.98 (2H, m), 2.51 (2H, t), 3.85 – 4.03 (1H, m), 5.51 (2H, s), 7.32 - 7.40 (3H, m), 7.94 - 7.99 (2H, m), 8.09 (1H, s)	402
	20	3-[4-(Adamantan-2-ylcarbonyl)-5-propylsulfanylmethyl]-pyrazol-1-ylmethylbenzoic acid	¹ H NMR (300.072 MHz, CDCl ₃) δ 0.82 (3H, t), 1.39 - 1.51 (2H, m), 1.57 – 1.73 (4H, m), 1.82 (8H, s), 1.96 (2H, s), 2.52 (2H, t), 4.20 – 4.28 (1H, m), 5.52 (2H, s), 7.34 - 7.41 (2H, m), 7.90 (1H, d), 7.93 - 7.99 (2H, m), 8.12 (1H, s)	454
	21	4-[4-(Adamantan-2-ylcarbonyl)-5-propylsulfanylmethyl]-pyrazol-1-ylmethylbenzoic acid	¹ H NMR (300.072 MHz, CDCl ₃) δ 0.81 (3H, t), 1.37 - 1.49 (2H, m), 1.58 - 1.75 (4H, m), 1.82 (8H, s), 1.96 (2H, s), 2.50 (2H, t), 4.20 – 4.28 (1H, m), 5.53 (2H, s), 7.21 (2H, d), 7.89 (1H, d), 7.99 (2H, d), 8.13 (1H, s)	454

Example 22**4-[4-(2-Adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoic acid**

- 2M aqueous sodium hydroxide solution (51.7 mL, 103.32 mmol) was added to methyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoate (**Intermediate#56**) (4.5 g, 10.33 mmol) in methanol (100 mL). The mixture was stirred at 70 °C for 1 hour and then cooled to ambient temperature, concentrated under reduced pressure and diluted with water (100 mL). The reaction mixture was adjusted to pH 3 with 2M HCl. The reaction mixture was extracted with EtOAc (500 mL) and washed sequentially with water (2x100 mL), and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to give a pale yellow solid. The solid was washed with EtOAc (20mL), collected by filtration and dried under vacuum to give 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoic acid (3.89 g, 89 %) as a cream crystalline solid.
- ¹H NMR (400.13 MHz, DMSO-*d*₆) δ 1.19 (9H, s), 1.49 (2H, d), 1.70-1.96 (10H, m), 2.09 (2H, d), 3.98 - 4.01 (1H, m), 7.49 - 7.53 (2H, m), 7.61 (1H, s), 8.06 - 8.09 (2H, m), 8.20 (1H, d), 13.30 (1H, s)
- m/z (ESI+) (M+H)⁺ = 422
- m.p. 308.8°C (onset)

- Example 22 may also be prepared as follows:

Aqueous sodium hydroxide (2M) (2.5 eq) was added portionwise over 5 minutes to a stirred suspension of methyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoate (**Intermediate#56**) (1.0 eq) in methanol (10 vol) at 20°C (exotherm 20 - 27°C). The resulting suspension was heated to 70°C (jacket temperature), (batch refluxes

approx 60-65°C) for 1 hour (complete by LCMS). The orange reaction mixture was cooled to 20°C (solution remained slightly cloudy) and filtered through celite to remove a small amount of solids. The filtrate was then poured into a flange flask and water (25 vol) was added. The mixture was then adjusted to pH 3 with 2M HCl (approx 800-850ml) (turns
5 very thick). The aqueous was then filtered and the pale yellow solid washed with water, sucked dry overnight, and washed with acetonitrile and finally 1:1 acetonitrile/diethyl ether and dried under vacuum at 50°C for 72 hours (weekend) to give 4-[4-(2-adamantylcarbamoyl)-5-tertbutyl-pyrazol-1-yl]benzoic acid (80 %) as a solid.

Approximately 50mg of 4-[4-(2-adamantylcarbamoyl)-5-tertbutyl-pyrazol-1-
10 yl]benzoic acid as prepared above (Form 1) was placed in a vial with a magnetic flea, and approximately 2ml of acetonitrile added. The vial was then sealed tightly with a cap. The slurry was then left to stir in a heated stirrer block with magnetic stirring capabilities at 50°C. After 3 days, the sample was removed from the plate, the cap taken off and the slurry left to dry under ambient conditions before it was analysed by XRPD and DSC.
15 This form (Form 2) was determined to be crystalline by XRPD and seen to be different to the previous form. This material had a melting point of 310.3°C (onset). It had 2 theta peaks measured using CuK α radiation at 18.0 and 17.7.

Approximately 20mg of 4-[4-(2-adamantylcarbamoyl)-5-tertbutyl-pyrazol-1-
yl]benzoic acid (form 1) was placed in a vial with a magnetic flea, and approximately 2ml
20 of methanol added, the vial was then sealed tightly with a cap and left to stir on a magnetic stirrer plate. After 3 days, the sample was removed from the plate, the cap taken off and the slurry left to dry under ambient conditions before it was analysed by XRPD and DSC.
This form (Form 3) was determined to be crystalline by XRPD and seen to be different to previously seen forms. This material had a melting point of 309.4°C (onset). It had 2 theta
25 peaks measured using CuK α radiation at 18.7 and 11.7.


Approximately 20mg of 4-[4-(2-adamantylcarbamoyl)-5-tertbutyl-pyrazol-1-
yl]benzoic acid as Form 1 and 20mg of the Form 3 material was placed in a vial with a
magnetic flea, and approximately 2ml of ethyl acetate added, the vial was then sealed
tightly with a cap and left to stir on a magnetic stirrer plate. After 3 days, the sample was
30 removed from the plate, the cap taken off and the slurry left to dry under ambient
conditions before it was analysed by XRPD and DSC. This form (Form 4) was determined
to be crystalline by XRPD and seen to be different to previously seen forms. This material

(Form 4) had a melting point of 309.1°C (onset).). It had 2 theta peaks measured using CuK α radiation at 16.2 and 20.6

4-[4-(2-Adamantylcarbamoyl)-5-tertbutyl-pyrazol-1-yl]benzoic acid as prepared above (Form 1) was suspended in acetonitrile (7 vol), seeded with 5g of (form 4) and slurried at
5 reflux for 3 days (jacket temperature 85°C). A sample was taken and checked by DSC (shows 2 peaks). The sample was stirred at reflux for a further 3 days (weekend), cooled to 20°C, filtered, washed through with acetonitrile then diethyl ether, sucked dry and dried under vacuum at 50°C for 48 hours to give a pale yellow solid (Form 4) (960g, 90 %).

Example 23

10 4-[4-(2-Adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoic acid



A solution of aqueous 2N sodium hydroxide (8.45 mL, 16.90 mmol) was added to a stirred solution of methyl 4-[4-(2-adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoate (**Intermediate#57**), 1.221 g, 2.82 mmol) in methanol (25 mL) at room
15 temperature. The resulting solution was stirred at 70 °C for 1 hour, and at room temperature overnight.


The reaction mixture was evaporated to dryness and re-dissolved in water (15 mL) and acidified with 2M HCl (10 mL). The reaction mixture was then extracted into EtOAc (75 mL), and washed sequentially with water (10 mL), and saturated brine (10 mL). The
20 organic layer was dried over MgSO₄, filtered and evaporated to afford 4-[4-(2-adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoic acid (1.055 g, 89 %) as a white solid.

m/z (ESI+) (M+H)⁺ = 420 ; HPLC t_R = 2.56 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.51 - 0.53 (2H, m), 0.68 - 0.69 (2H, m), 1.54 - 1.58 (5H, m), 1.73 (2H, s), 1.84 - 1.87 (6H, m), 1.95 - 1.99 (2H, m), 2.06 (2H, d), 4.03 - 4.09 (1H, m), 7.44 (1H, d), 7.67 (2H, d), 8.06 (1H, s), 8.11 (2H, d), 13.16 (1H, s)

Example 24

4-[4-(2-Adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoic acid






A solution of aqueous 2N sodium hydroxide (4.10 mL, 8.19 mmol) was added to a stirred solution of methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoate (**Intermediate#62**) 611 mg, 1.37 mmol) in methanol (15 mL) at room temperature. The resulting solution was stirred at 70 °C for 1 hour.


The reaction mixture was evaporated to dryness and redissolved in water (15 mL) and acidified with 2M HCl (6 mL). The suspension obtained was then filtered. The product recovered was washed with water (10 mL) and dried under vacuum to give 4-[4-(2-adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoic acid (576 mg, 97 %) as a white solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.43 - 1.55 (4H, m), 1.74 - 1.85 (12H, m), 1.94 (2H, s), 2.03 - 2.12 (4H, m), 2.99 - 3.08 (1H, m), 3.98 - 4.03 (1H, m), 7.53 - 7.55 (2H, m), 7.74 (1H, d), 8.09 (1H, s), 8.10 - 8.12 (2H, m), 13.30 (1H, s)

m/z (ESI+) (M+H)⁺ = 434; HPLC t_R = 2.80 min.

The same process as used for Example#24 prepared the following examples from the appropriate intermediate.


Structure	Ex	Name	¹ H NMR δ	MS m/e MH ⁺
	25	4-[4-(2-ethylpyrazol-1-yl)benzoyl]adamantylcarbamate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.03 (3H, t), 1.53 (2H, d), 1.72 (2H, s), 1.78 - 1.85 (6H, m), 1.95 (2H, s), 2.11 (2H, d), 2.97 (2H, q), 4.04 (1H, t), 7.58 - 7.63 (3H, m), 8.09 - 8.13 (2H, m), 8.28 (1H, s)	394
	26	4-[4-(2-isopropylpyrazol-1-yl)benzoyl]adamantylcarbamate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.29 (6H, d), 1.52 (2H, d), 1.73 (2H, s), 1.82 (5H, s), 1.86 (1H, s), 1.98 (2H, s), 2.11 (2H, d), 3.10 - 3.17 (1H, m), 4.02 - 4.05 (1H, m), 7.54 (2H, d), 7.67 (1H, d), 8.07 (1H, s), 8.11 (2H, d), 13.20 (1H, s)	408
	27	4-[4-(2-cyclobutylpyrazol-1-yl)benzoyl]adamantylcarbamate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.52 (2H, d), 1.65 (1H, q), 1.72 (2H, s), 1.80 - 1.85 (7H, m), 1.96 (2H, s), 2.03 - 2.13 (4H, m), 2.20 - 2.28 (2H, m), 3.76 - 3.85 (1H, m), 4.01 - 4.06 (1H, m), 7.52 (2H, d), 7.80 (1H, d), 7.93 (1H, s), 8.07 (2H, d), 13.40 (1H, s)	420

Example 28**4-[4-(2-Adamantylcarbamoyl)-5-methyl-pyrazol-1-yl]benzoic acid**

A solution of 1M sodium hydroxide (24.28 mL, 24.28 mmol) was added to a stirred
5 suspension of N-(2-adamantyl)-1-(4-cyanophenyl)-5-methyl-pyrazole-4-carboxamide
(**Intermediate#66**) (1.25 g, 3.47 mmol) in dioxane (25 mL). The resulting suspension was
stirred at 100 °C for 7 hours. The reaction mixture was concentrated, diluted with water
(40 mL) and filtered through Celite. The filtrates were acidified with 1M citric acid. The
precipitate was recovered by filtration, washed with water (3x20 mL) and dried under
10 vacuum at 50°C. The crude product was purified by preparative HPLC (Phenomenex
Gemini C18 110A (axia) column, 5 μ silica, 30 mm diameter, 100 mm length), using
decreasingly polar mixtures of water (containing 1% formic acid) and MeCN as eluents.
Fractions containing the desired compound were evaporated to dryness to afford 4-[4-(2-
adamantylcarbamoyl)-5-methyl-pyrazol-1-yl]benzoic acid (550mg, 42%) as a pale yellow
15 powder.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.45 - 1.54 (2H, m), 1.70 - 1.88 (8H, m), 1.90 - 2.00
(2H, m), 2.05 - 2.18 (2H, m), 2.56 (3H, s), 4.00 - 4.10 (1H, m), 7.57 (1H, d), 7.67 (2H, d),
8.11 (2H, d), 8.29 (1H, s), 13.25 (1H, s).

m/z (ESI+) (M+H)⁺ = 380;

Example 29**4-(5-tert-Butyl-4-(cyclohexylcarbamoyl)-1H-pyrazol-1-yl)benzoic acid**


1-(4-bromophenyl)-5-tert-butyl-N-cyclohexyl-1H-pyrazole-4-carboxamide

- 5 **(Intermediate# 82)** (132 mg, 0.33 mmol), molybdenum hexacarbonyl (43.1 mg, 0.16 mmol), trans-Di(-acetato)bis[o-(di-o-tolylphosphino)benzyl]dipalladium (II) (15.34 mg, 0.02 mmol), 4-Dimethylaminopyridine (80 mg, 0.65 mmol) and N-Ethyldiisopropylamine (0.113 mL, 0.65 mmol) were suspended in Dioxane (4 mL) and Water (1 mL) and sealed into a microwave tube. The reaction was heated to 150 °C for 1 hour in the microwave
- 10 reactor and cooled to RT. The reaction mixture was diluted with DCM (20mL) and water (10mL) and then adjusted to pH3 with 2M HCl and filtered through celite. The organic layer was separated, dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by preparative reverse phase HPLC using decreasingly polar mixtures of water (containing 0.1% NH₃) and MeCN as eluents.

- 15 Fractions containing the desired compound were evaporated to dryness to afford 4-(5-tert-butyl-4-(cyclohexylcarbamoyl)-1H-pyrazol-1-yl)benzoic acid (18 mg, 14.92 %) as a white solid.




- ¹H NMR (400.13 MHz, DMSO-d₆) δ 1.15 (1H, s), 1.20 (9H, s), 1.25-1.4 (4H, m), 1.58 (1H, s), 1.71 - 1.74 (2H, m), 1.82 (2H, d), 3.18 (1H, s), 7.48 - 7.50 (2H, m), 7.60 (1H, s),
- 20 8.05 - 8.10 (3H, m)
- m/z (ESI+) (M+H)⁺ = 370

Example 30**4-[4-(2-Adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid**


2M aqueous sodium hydroxide (1.671 mL, 3.34 mmol) was added in one portion to methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoate (Intermediate# 84) (330 mg, 0.67 mmol) in methanol (10 mL). The resulting mixture was stirred at 20 °C for 18 hours. The reaction mixture was concentrated and diluted with water (50 mL), and washed with ether (20mL). The aqueous solution was adjusted to pH 3 with 2M HCl and extracted with EtOAc (2x25 mL), and the combined extracts washed sequentially with water (2x20 mL) and saturated brine (20 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford 4-[4-(2-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid (321 mg, 100 %).

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.01 - 1.11 (5H, m), 1.41 (1H, s), 1.50 - 1.57 (4H, m), 1.64 (2H, d), 1.75 (2H, s), 1.85 (6H, s), 1.93-2.05 (4H, m), 2.94 (1H, s), 4.12 (1H, d), 7.72 - 7.76 (2H, m), 8.05 (1H, d), 8.10 - 8.13 (2H, m), 8.18 (1H, s), 13.20 (1H, s)
m/z (ESI+) (M+H)⁺ = 480

The following Examples were prepared in a similar manner to Example #30, using an appropriate ester starting material

Structure	Ex	Name	¹ H NMR δ	MS m/e
	31	4-[4-(1-adamantylcarbamoyl)-5-cyclohexylsulfany-1-pyrazol-1-yl]benzoic acid	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.05 - 1.18 (5H, m), 1.42 (1H, s), 1.56 (4H, d), 1.68 (6H, s), 2.07 (9H, s), 2.98 (1H, s), 7.53 (1H, s), 7.68 - 7.71 (2H, m), 8.09 - 8.11 (3H, m), 13.19 (1H, s)	(M-H)- 478
	32	4-[5-cyclohexylsulfany-1-4-[(5-hydroxy-2-adamantyl)carbamoyl]pyrazol-1-yl]benzoic acid	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.03 - 1.08 (5H, m), 1.41-1.56 (7H, m), 1.67 (4H, d), 1.76 (2H, d), 1.89 (2H, d), 2.08 (3H, s), 2.93 (1H, s), 4.03 (1H, d), 4.43 (1H, s), 7.73 (2H, d), 7.97 (1H, d), 8.10 - 8.12 (2H, m), 8.18 (1H, s), 13.25 (1H, s)	(M+H)+ 496
	33	4-[5-cyclohexylsulfany-1-4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoic acid	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.03 - 1.08 (5H, m), 1.41 - 1.54 (7H, m), 1.93 (6H, d), 2.05 (2H, d), 2.18 (3H, d), 2.96 (1H, d),	(M+H)+ 546

Structure	Ex	Name	¹ H NMR δ	MS m/e
		yl]benzoic acid	4.10 (1H, t), 6.88 (1H, t), 7.73 (2H, d), 7.99 (1H, d), 8.12 (2H, d), 8.18 (1H, s), 13.20 (1H, s)	



Example 34**4-[4-(2-Adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid**


2M aqueous sodium hydroxide (1.694 mL, 3.39 mmol) was added in one portion to methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate (Intermediate # 90) (325 mg, 0.68 mmol) in methanol (10 mL). The resulting mixture was stirred at 20 °C for 18 hours. The reaction mixture was concentrated and diluted with water (50 mL), and washed with ether (20mL). The aqueous solution was adjusted to pH 3 with 2M HCl and extracted with EtOAc (2x25 mL), and the combined extracts washed sequentially with water (2x20 mL) and saturated brine (20 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford 4-[4-(2-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid (307 mg, 97 %).

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.17 - 1.25 (2H, m), 1.41 (4H, d), 1.65 (4H, d), 1.74 (2H, s), 1.85 (6H, s), 1.96 - 2.02 (4H, m), 3.31 - 3.33 (1H, m), 4.12 (1H, d), 7.75 - 7.77 (2H, m), 8.08 (1H, d), 8.10 - 8.15 (2H, m), 8.19 (1H, s), 13.20 (1H, s)


m/z (ESI+) (M+H)⁺ = 466

The following Examples were prepared in a similar manner to Example #34, using an appropriate ester starting material

Structure	Ex#	Name	¹ H NMR δ	MS m/e (M+H) ⁺
	35	methyl 4-[4-(1-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.18 - 1.28 (2H, m), 1.41 - 1.49 (4H, m), 1.64 - 1.71 (8H, m), 2.07 (9H, s), 3.32 - 3.37 (1H, m), 7.54 (1H, s), 7.70 - 7.73 (2H, m), 8.09 - 8.13 (3H, m), 13.20 (1H, s)	466
	36	methyl 4-[5-cyclopentylsulfanyl-4-[[[(1R,3S)-5-hydroxy-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.17 - 1.25 (2H, m), 1.36 - 1.47 (6H, m), 1.65 (6H, d), 1.76 (2H, d), 1.89 (2H, d), 2.08 (3H, s), 3.35 (1H, d), 4.02 - 4.05 (1H, m), 4.43 (1H, s), 7.74 - 7.77 (2H, m), 7.99 (1H, d), 8.10 - 8.13 (2H, m), 8.18 (1H, s)	482

Example 37**4-[4-[[5-(Difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanylpirazol-1-yl]benzoic acid**

- 5 2M aqueous sodium hydroxide (2.050 mL, 4.10 mmol) was added in one portion to methyl 4-[4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanylpirazol-1-yl]benzoate (**Intermediate# 95**) (459 mg, 0.82 mmol) in methanol (10 mL). The resulting mixture was stirred at 20 °C for 18 hours and then a further 4 hours at 55°C. The reaction mixture was concentrated and diluted with water (50 mL), and washed with ether (20mL).
- 10 The aqueous solution was adjusted to pH 3 with 2M HCl and extracted with EtOAc (2x25 mL), and the combined extracts washed sequentially with water (2x20 mL) and saturated brine (20 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford 4-[4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanylpirazol-1-yl]benzoic acid (284 mg, 63.5 %).
- 15 ¹H NMR (400.13 MHz, DMSO-d₆) δ 1.03 - 1.08 (5H, m), 1.41 - 1.54 (7H, m), 1.93 (6H, d), 2.05 (2H, d), 2.18 (3H, d), 2.96 (1H, d), 4.10 (1H, t), 6.88 (1H, t), 7.73 (2H, d), 7.99 (1H, d), 8.12 (2H, d), 8.18 (1H, s), 13.20 (1H, s)
- m/z (ESI+) (M+H)⁺ = 506


Example 38**4-[4-(Cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid**

A solution of aqueous 2N sodium hydroxide (4 mL, 8 mmol) was added to a stirred
5 solution of methyl 4-[4-(cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate (**Intermediate#96**), 233 mg, 0.55 mmol) in methanol (7 mL) at room temperature. The resulting solution was stirred at room temperature overnight.


The reaction mixture was evaporated to dryness and redissolved in water (15 mL)
and acidified with 2M HCl (6 mL). The reaction mixture was then extracted in EtOAc (30
10 mL), and washed sequentially with water (10 mL), and saturated brine (10 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford 4-[4-(cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid (217 mg, 96 %) as a white solid.

m/z (ESI+) (M+H)⁺ = 414

1H NMR (400.13 MHz, DMSO-d₆) δ 1.21 - 1.45 (11H, m), 1.58 - 1.74 (5H, m), 1.84 - 1.87
15 (2H, m), 3.42 - 3.48 (1H, m), 3.76 - 3.82 (1H, m), 7.70 (2H, d), 7.90 - 7.92 (1H, d), 8.11
(2H, d), 8.16 (1H, s), 13.19 (1H, s)

Example 39**4-[4-(Cyclohexylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid**

- 4-[4-(cyclohexylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid was prepared from methyl 4-[4-(cyclohexylcarbamoyl)-5-cyclohexylsulfanylpyrazol-1-yl]benzoate (**Intermediate#97**) by the same process used for Example#38
- m/z (ESI+) $(M+H)^+ = 428$; HPLC $t_R = 2.67$ min.
- 1H NMR (400.13 MHz, DMSO- d_6) δ 1.09 (4H, m), 1.19 - 1.30 (2H, m), 1.33 - 1.41 (5H, m), 1.51 - 1.58 (5H, m), 1.71 - 1.75 (2H, m), 1.84 - 1.87 (2H, m), 3.03 - 3.12 (1H, m), 3.72 - 3.80 (1H, m), 7.71 (2H, d), 7.89 - 7.90 (1H, d), 8.09 - 8.11 (2H, m), 8.16 (1H, s), 13.20 (1H, s)

Example 40**4-[5-Cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoic acid**


4-[5-cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoic acid was prepared from methyl 4-[5-cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate#98**) by the same process used for Example#38.

m/z (ESI+) (M+H)+ = 442

5 1H NMR (400.13 MHz, DMSO-d₆) δ 1.16 - 1.52 (15H, m), 1.58 - 1.67 (3H, m), 1.71 - 1.75 (2H, m), 1.84 - 1.87 (2H, m), 3.30 (1H, m), 3.78 - 3.81 (1H, m), 7.67 - 7.70 (2H, d), 7.90 (1H, d), 8.09 - 8.12 (2H, d), 8.16 (1H, s), 13.18 (1H, s)

Example 41

4-[4-(2-Adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoic acid



10


2M aqueous sodium hydroxide (2.446 mL, 4.89 mmol) was added in one portion to methyl 4-[4-(2-adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoate (**Intermediate# 99**) (430 mg, 0.98 mmol) in methanol (20 mL). The resulting mixture was stirred at 20 °C for 18 hours. The reaction mixture was concentrated, diluted with water (50 mL) and adjusted to pH 3 with 2M HCl. The precipitate was collected by filtration, washed with water (20 mL) and dried under vacuum to afford 4-[4-(2-

15 adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoic acid (383 mg, 92 %) as a white solid.

1H NMR (400.13 MHz, DMSO-d₆) δ 0.94 (3H, t), 1.63 (2H, d), 1.74 (2H, s), 1.86 (6H, d), 1.99 (4H, d), 2.68 (2H, q), 4.11 (1H, t), 7.72 - 7.76 (2H, m), 8.04 (1H, d), 8.10 - 8.13 (2H, m), 8.19 (1H, s), 13.2 (1H, s)

20


m/z (ESI+) (M+H)+ = 426

Example 42**4-[4-(2-Adamantylcarbamoyl)-5-methylsulfanyl-pyrazol-1-yl]benzoic acid**

Prepared from methyl 4-[4-(2-adamantylcarbamoyl)-5-methylsulfanyl-pyrazol-1-yl]benzoate (**Intermediate#100**) by the same process as Example#41.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.61 (2H, d), 1.74 (2H, s), 1.86 (6H, d), 2.00 (4H, d), 2.30 (3H, s), 4.11 (1H, t), 7.72 - 7.75 (2H, m), 8.01 - 8.04 (1H, m), 8.10 - 8.14 (2H, m), 8.19 (1H, s), 13.2 (1H, s)

m/z (ESI+) (M+H)⁺ = 412

Example 43**4-[4-(5-Methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid**

Lithium hydroxide monohydrate (27.5 mg, 0.65 mmol) was added to a suspension of 4-[4-(5-Methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid methyl ester (**Intermediate#105**) (116 mg, 0.22 mmol) in methanol (4 mL) / water (2 mL) at ambient temperature. The resulting suspension was stirred at ambient temperature for 18 hours. The bulk of the organic solvent was removed in vacuo and the


resulting solution was diluted with water (10mL) and washed with ether (10mL). The aqueous layer was acidified to ~pH4 with 2M HCl then extracted with EtOAc (3 x 25ml). The EtOAc layers were combined, washed sequentially with water (5 mL) and saturated brine (10 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford 4-[4-(5-Methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid (113 mg, 100 %) as a white solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.68 (3H, t), 1.22 - 1.31 (2H, m), 1.57 (2H, d), 1.96 (2H, s), 2.00 - 2.16 (7H, m), 2.20 (2H, s), 2.65 (2H, t), 2.87 (3H, s), 4.09 (1H, m), 7.74 (2H, d), 8.05 (1H, d), 8.12 (2H, m), 8.19 (1H, s), 13.20 (1H, s)

MS m/e MH⁺ 518.

Example 44

4-[4-(2-Adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoic acid



2M aqueous sodium hydroxide (1.256 mL, 2.51 mmol) was added in one portion to methyl 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoate (**Intermediate# 106**) (243 mg, 0.50 mmol) in methanol (10 mL). The resulting mixture was stirred at 20 °C for 18 hours.


The reaction mixture was concentrated, diluted with water (50 mL) and adjusted to pH 3 with 2M HCl. The precipitate was collected by filtration, washed with water (20 mL) and dried under vacuum to afford 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoic acid (202 mg, 86 %) as a white solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.70 (3H, t), 1.23 - 1.32 (2H, m), 1.62 (2H, d), 1.74 (2H, s), 1.86 (6H, d), 1.92 - 2.05 (4H, m), 2.65 (2H, t), 3.87 (3H, s), 4.11 (1H, d), 7.23 - 7.25 (1H, m), 7.39 (1H, d), 7.81 (1H, d), 8.09 (1H, d), 8.17 (1H, s), 12.92 (1H, s)

m/z (ESI+) (M+H)⁺ = 470

Example 45

4-[4-(2-Adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoic acid




5 A solution of Sodium hydroxide (1.904 mL, 3.81 mmol) was added in one portion to a stirred solution of ethyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoate (**Intermediate#113**) (353 mg, 0.76 mmol) in methanol (6 mL). The resulting suspension was stirred at 20 °C for 16 hours. The resulting mixture was evaporated to remove the methanol and washed with ether (20 mL). The reaction mixture

10 was acidified with 2M HCl. The precipitate was collected by filtration, washed with water (10 mL) and dried under vacuum to afford 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoic acid (266 mg, 80 %) as a white solid.

¹H NMR (300.073 MHz, dmso) δ 1.16 (s, 9H), 1.50 (d, *J* = 12.6 Hz, 2H), 1.70 (s, 2H), 1.74 - 1.88 (m, 6H), 1.89 - 1.99 (m, 2H), 2.01 - 2.14 (m, 5H), 3.99 (d, 1H), 7.37 (d, 1H),

15 7.66 (s, 1H), 7.86 (d, 1H), 7.94 (s, 1H), 8.05 (d, 1H)


m/z (ESI+) (M+H)⁺ = 436

Example 46**4-[4-(2-Adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-2-(trifluoromethyl)benzoic acid**

5 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-2-(trifluoromethyl)benzoic acid was prepared from ethyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoate (**Intermediate# 117**) by the same process used for Example#45.

1H NMR (400.13 MHz, CDCl₃) δ 1.29 (9H, s), 1.71 - 1.79 (6H, m), 1.91 (6H, s), 2.07 (2H, s), 4.24 (1H, d), 6.20 (1H, d), 7.63 - 7.66 (1H, m), 7.70 (1H, s), 7.79 - 7.80 (1H, m),
 10 8.00 (1H, d)

MS m/z (ESI+) (M+H)⁺ = 490.

Example 47 : 4-[4-(Adamantan-2-ylcarbamoyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl]benzoic acid.


15 A solution of Sodium hydroxide (1.056 mL, 2.11 mmol) was added in one portion to a stirred solution of N-adamantan-2-yl-1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-

pyrazole-4-carboxamide (**Intermediate# 124**) (250 mg, 0.60 mmol), in methanol (10 mL) under air. The resulting solution was stirred at 65°C for 45 hours. The resulting mixture was evaporated to dryness and the residue dissolved in ice/water (25mL) and the mixture was acidified with 2M HCl. The precipitate was collected by filtration, washed with water (25 mL) and dried under vacuum to afford 4-[4-(adamantan-2-ylcarbamoyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl]benzoic acid (243 mg, 93 %) as a white solid, which was used without further purification.

m/z (ESI+) (M+H)+ = 434; HPLC t_R = 2.57 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.49 - 1.53 (2H, m), 1.71 (2H, s), 1.80 (5H, s), 1.84 (1H, s), 1.93 (2H, s), 2.05 (2H, d), 3.98-4.05 (1H, m), 7.63 (2H, d), 8.11 - 8.14 (3H, m), 8.34 (1H, d), 13.30 (1H, s)

Intermediate#1 : 4-[4-((1R,2S,3S,5S)-5-Hydroxy-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid methyl ester




1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid (**Intermediate#7**) (160mg, 0.5mmol), (1S,3S,4S,5R)-4-Amino-adamantan-1-ol (84mg, 0.5mmol), HOBT (81mg, 0.6mmol), and DIPEA (174 μ L, 1mmol) were dissolved in DMF (5mL) and treated at ambient temperature with EDCI (115mg, 0.6mmol). The mixture was stirred at ambient for 18h and then diluted with ethyl acetate (50mL), washed with water (3x20mL) and brine, dried (MgSO₄) and evaporated to leave a brown gum, which was purified by chromatography on silica gel (12 silica 0-100% EtOAc / isohexane) to give the title compound (159mg, 69%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.65 (3H, t), 1.17 - 1.29 (2H, m), 1.42 (2H, d), 1.60-1.80 (6H, m), 1.89 (2H, d), 2.06 (3H, s), 2.62 (2H, t), 3.90 (3H, s), 4.00 (1H, d), 4.43 (1H, s), 7.76 (2H, d), 7.95 (1H, d), 8.13 (2H, d), 8.17 (1H, s)

MS m/z 470 M+H

Intermediate#2 : Methyl 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate




1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid

- 5 **(Intermediate #7)** (160mg, 0.5mmol), 2-adamantylamine hydrochloride (94mg, 0.5mmol), HOBT (81mg, 0.6mmol) and DIPEA (261μL, 1.5mmol) were dissolved in DMF (5mL) and treated at ambient temperature with EDCI (115mg, 0.6mmol). The mixture was stirred at ambient for 18h and then diluted with ethyl acetate (50mL), washed with water (3x20mL) and brine, dried (MgSO₄) and evaporated to leave a white solid, which was
- 10 purified by chromatography on silica gel, (12g silica 0-50% EtOAc /isohexane) to give the title compound as a white solid. (203mg, 92%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.65 (3H, t), 1.17 - 1.29 (2H, m), 1.60 (2H, d), 1.73 (2H, s), 1.83 (6H, s), 1.91-2.05 (4H, m), 2.62 (2H, t), 3.90 (3H, s), 4.09 (1H, d), 7.79 (2H, d), 8.03 (1H, d), 8.13 (2H, d), 8.16 (1H, s)

- 15 MS m/z 454 M+H

Intermediate#3 : Methyl 4-[4-(1-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate




1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid

- 5 **(Intermediate #7)** (160mg, 0.5mmol), 1-adamantylamine (76mg, 0.5mmol), HOBt (81mg, 0.6mmol) and DIPEA (174μL, 1.0mmol) were dissolved in DMF (5mL) and treated at ambient temperature with EDCI (115mg, 0.6mmol). The mixture was stirred at ambient for 18h and then diluted with ethyl acetate (50mL), washed with water (3x20mL) and brine, dried (MgSO₄) and volatiles removed by evaporation. The residue was purified by
- 10 chromatography on silica gel (12g silica 0-50% EtOAc / isohexane) to give the title compound as a colourless gum (143mg, 63%).

MS m/z 454 M+H

Intermediate#4 : Methyl 4-[4-(N-cyclohexyl-N-methyl-carbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate




1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid
(**Intermediate #7**), (160mg, 0.5mmol), N-methylcyclohexylamine (57mg, 0.5mmol), HOBT (81mg, 0.6mmol) and DIPEA (174 μ L, 1.0mmol) were dissolved in DMF (5mL) and treated at ambient temperature with EDCI (115mg, 0.6mmol). The mixture was stirred at ambient
5 for 18h and then diluted with ethyl acetate (50mL), washed with water (3x20mL) and brine, dried (MgSO₄) and volatiles removed by evaporation. The residue was purified by chromatography on silica gel (12g silica 0-70% EtOAc / isohexane) to give the title compound as a colourless gum (162mg, 78%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.67 (3H, t), 1.01-1.41 (4H, m), 1.42-1.85 (8H, m),
10 2.58 (2H, t), 2.86 (3H, s), 3.40-3.60 (0.5H, m), 3.89 (3H, s), 4.20-4.40 (0.5H, m) 7.78 (2H, d), 7.91 (1H, s), 8.12 (2H, d)

MS m/z 416 M+H

Intermediate#5 : Methyl 4-[4-(oxan-4-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate




15

1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid
(**Intermediate #7**), (160mg, 0.5mmol), 4-aminotetrahydropyran (61mg, 0.5mmol), HOBT (81mg, 0.6mmol) and DIPEA (174 μ L, 1.0mmol) were dissolved in DMF (5mL) and treated at ambient temperature with EDCI (115mg, 0.6mmol). The mixture was stirred at ambient
20 for 18h and then diluted with ethyl acetate (50mL), washed with water (3x20mL) and brine, dried (MgSO₄) and volatiles removed by evaporation. The residue was purified by chromatography on silica gel (12g silica, 0-100% EtOAc / isohexane) to give the title compound as a white solid (114mg, 56%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.68 (3H, t), 1.18 - 1.30 (2H, m), 1.47 - 1.60 (2H, m),
25 1.77 - 1.81 (2H, m), 2.72 (2H, t), 3.36 - 3.43 (2H, m), 3.81-3.89 (5H, m), 3.95-4.05 (1H, m), 7.71 (2H, d), 8.04 (1H, d), 8.12 (2H, d), 8.17 (1H, s)

MS m/z 403 M+H


Intermediate#6 : Methyl 4-[5-propylsulfanyl-4-[3-[2-(trifluoromethyl)phenyl]pyrrolidine-1-carbonyl]pyrazol-1-yl]benzoate



- 5 1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid
(**Intermediate #7**) was dissolved in DCM (3mL). 1 drop of DMF and oxalyl chloride (71μL, 0.83mmol) was added. The mixture was stirred at ambient temperature for 2h and then volatiles were removed by evaporation under reduced pressure. The residue was dissolved in DCM (5mL) and added to a solution of 3-[2-
- 10 (trifluoromethyl)phenyl]pyrrolidine HCl (105mg, 0.42mmol) and DIPEA (217μL, 1.25mmol) in DCM (5mL). Water (10mL) was added and the mixture stirred vigorously and passed through a phase separating column. The filtrate was purified by chromatography on silica gel, eluting with 0-20% EtOAc/DCM to give the title compound as a clear colourless oil (177mg, 81%).

15 MS m/z 518 M+H

Intermediate#7 : 1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylic acid




- 20 tert-butyl 1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylate
(**Intermediate#8**) (2.86g, 7.97mmol) was dissolved in DCM (40mL) and TFA (10mL) added, the mixture was stirred at ambient temperature for 3h and then evaporated under reduced pressure to leave a light brown oil. Trituration of the oil with isohexane gave a

light brown solid that was recovered by filtration and dried under vacuum to give the title compound. (2.23g, 89%)

^1H NMR (300.073 MHz, DMSO- d_6) δ 0.67 (3H, t), 1.20 - 1.32 (2H, m), 2.81 (2H, t), 3.90 (3H, s), 7.71 (2H, d), 8.12 (2H, d), 8.17 (1H, s), 12.73 (1H, s)

5 MS m/z 321 M+H

Intermediate#8 : tert-butyl 1-(4-methoxycarbonylphenyl)-5-propylsulfanyl-pyrazole-4-carboxylate



tert-butyl 5-chloro-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylate

10 **(Intermediate#9)** (2.016g, 6.0mmol) was dissolved in butyronitrile (30mL), potassium carbonate (2.48g, 18mmol) and propanethiol (678 μ L, 7.5mmol) were added and the mixture heated to reflux for 5h. Ethyl acetate (150mL) was added and the mixture washed with water (4x25mL), dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica gel (120g silica 0-25% EtOAc / isohexane) to
15 give the title compound as a clear pale yellow oil that slowly crystallised on standing to a white solid. (2.01g, 89%)


^1H NMR (300.073 MHz, DMSO- d_6) δ 0.67 (3H, t), 1.25 (2H, q), 1.54 (9H, s), 2.76 (2H, t), 3.89 (3H, s), 7.70 (2H, d), 8.11 (3H, d)

MS m/z 321 M-tBut

20

25

Intermediate#9 : tert-butyl 5-chloro-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylate




t-Butylnitrite (2.9mL, 24.22mmol) and cupric chloride (4.06g, 30.27mmol) were added to acetonitrile (150mL) and heated to 65°C. Tert-butyl 5-amino-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylate (**Intermediate#10**) was added as a solid giving a vigorous gas evolution. After the addition was complete, heating was continued for a further 15min. The reaction mixture was cooled to ambient, diluted with water (500mL) and extracted with ethyl acetate (3x100mL). The combined extracts were washed with water (2x100mL) and brine (100mL), dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica gel (120g silica column, EtOAc/ Hexane 0-50%) to give the title compound as an oil. (5.68g, 83%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.53 (9H, s), 3.87 - 3.90 (3H, m), 7.78 (2H, d), 8.15 (2H, d), 8.20 (1H, s)

MS m/z 281 M-tBut

Intermediate#10 Tert-butyl 5-amino-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylate




t-Butyl cyanoacetate (14.1g, 100mmol) was dissolved in triethyl orthoformate (24.8mL, 150mmol). Acetic anhydride (9.625mL, 100mL) was added and the mixture heated to 125°C for 3h and then volatiles were removed by evaporation under reduced pressure. The residue was dissolved in ethanol (100mL) and methyl 4-hydrazinylbenzoate

hydrochloride (**Intermediate# 123**) (6.06g, 30mmol) and DIPEA (5.23mL, 30mmol) was added. The mixture was heated to reflux for 5h and then evaporated under reduced pressure to leave a brown oil which was dissolved in ethyl acetate (300mL) and washed with water (2x100mL) and brine (100mL), dried (MgSO₄) and evaporated. The residue was purified by chromatography on silica gel (120g silica, EtOAc/Hexane 0-50%) to give the title compound as a yellow solid. (7.1g)

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.51 (9H, s), 3.88 (3H, s), 6.43 (2H, s), 7.67 (1H, s), 7.73 (2H, s), 8.08 (2H, d)

MS m/z 262 M-tBut

Intermediate # 11 :Methyl 4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate




Propane thiol (88mg, 1.16mmol) was dissolved in DMF (5mL) and treated at ambient temperature with a 1M solution of NaHMDS in THF (1.16mL). After stirring for 15min the clear solution was added to a suspension of methyl 4-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate#15**) (378mg, 1.05mmol) in DMF (10mL). Stirring was continued at ambient temperature for 2h and then the reaction mixture was diluted with ethyl acetate (100mL), washed with water (4x25mL) and dried (MgSO₄). Volatiles were removed by evaporation under reduced pressure to give a clear oil which was purified by chromatography on silica gel, eluting with an ethyl acetate / hexane gradient (0-50%) to give the title compound as a white solid. (363mg, 86%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.67 (3H, t), 1.14 - 1.39 (7H, m), 1.59 (1H, d), 1.71 - 1.74 (2H, m), 1.84 (2H, d), 2.70 (2H, t), 3.74 - 3.79 (1H, m), 3.89 (3H, s), 7.72 (2H, d), 7.92 (1H, d), 8.10 - 8.16 (3H, m)

MS m/z 402 M+H

Intermediate #12: Methyl 3-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoate




Propane thiol (48mg, 0.57mmol) was dissolved in DMF (3mL) and treated at
5 ambient temperature with a 1M solution of NaHMDS in THF (0.63mL, 0.63mmol). After
stirring for 15min a solution of methyl 3-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate # 19**) (205mg, 0.57mmol) in DMF (5mL) was added and
stirring continued for 2h. Ethyl acetate (50mL) was added and the mixture washed with
water (3x20mL), dried (MgSO₄) and evaporated to give an oil which was purified by
10 chromatography on silica gel eluting with an EtOAc/DCM gradient (0-25%) to provide the
title compound as a clear colourless oil. (175mg, 75%)

¹H NMR (300.073 MHz, DMSO-d₆) δ0.67 (3H, t), 1.14 - 1.39 (7H, m), 1.57 - 1.61 (1H, m), 1.71 - 1.74 (2H, m), 1.84 (2H, d), 2.68 - 2.72 (2H, m), 3.71-3.85 (1H, m), 3.89 (3H, s), 7.72 (1H, t), 7.83 - 7.94 (2H, m), 8.05 - 8.08 (2H, m), 8.14 (1H, s)

15 MS m/z 402 M+H

Intermediate#15: methyl 4-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate




t-Butylnitrite (210mg, 2.04 mmol) and cupric chloride (342mg, 2.55mmol) were
added to acetonitrile (15mL) and heated to 65°C. Methyl 4-[5-amino-4-
20 (cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate # 16**) (581mg, 1.7mmol) was
added as a solid in portions giving a vigorous gas evolution. After the addition was
complete the reaction mixture was heated for a further 15min. at 65°C, cooled to ambient,

diluted with water (100mL) and extracted with ethyl acetate (3x50mL). The combined extracts were washed with brine (2x20mL), dried (MgSO₄) and evaporated to give a solid, which was purified by chromatography on silica gel eluting with an EtOAc/Hexane gradient (0-30%) to give the title compound as a white solid (385mg, 65%)

5 ¹H NMR (300.072 MHz, CDCl₃) δ 1.21 (3H, d), 1.31 - 1.45 (2H, m), 1.57 - 1.71 (3H, m), 1.93 - 1.98 (2H, m), 3.87 - 3.99 (4H, m), 6.02 (1H, d), 7.56 - 7.60 (2H, m), 8.08 - 8.14 (3H, m)

MS m/z 362 M+H

Intermediate # 16: Methyl 4-[5-amino-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate



10

2-cyano-N-cyclohexyl-3-ethoxy-prop-2-enamide (**Intermediate #17**) (605mg, 2.73mmol) and methyl 4-hydrazinylbenzoate hydrochloride (**Intermediate# 123**) (552mg, 2.73mmol) were suspended in ethanol (20mL). DIPEA (351mg, 2.73mmol) was added and the mixture heated to 70°C for 1h. The reaction mixture was cooled to ambient and the resulting precipitate recovered by filtration, washed with ether and dried *in vacuo* to give the title compound as a white solid. (618mg 66%)


15

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.13 (1H, d), 1.22 - 1.35 (4H, m), 1.60 (1H, d), 1.76 (4H, d), 3.69 - 3.74 (1H, m), 3.87 (3H, s), 6.56 (2H, s), 7.62 (1H, d), 7.73 - 7.78 (2H, m), 8.04 (1H, s), 8.05 - 8.09 (2H, m)

20

MS m/z 343 M+H

Intermediate #17: 2-cyano-N-cyclohexyl-3-ethoxy-prop-2-enamide




2-cyano-N-cyclohexyl-acetamide (**Intermediate # 18**) (1.35g, 8.09 mmol) was suspended in acetic anhydride (20mL). Triethyl orthoformate (3.91g, 21 mmol) was added

and the mixture heated to reflux for 5h. The reaction mixture was cooled to ambient and volatiles removed by evaporation *in vacuo* to leave a brown oil which was which was purified by chromatography on silica gel eluting with an EtOAc/DCM gradient (0-10%) to give the title compound as a solid (637mg, 35%)

5 ^1H NMR (300.073 MHz, DMSO- d_6) δ 1.01 - 1.35 (8H, m), 1.51-1.82 (5H, m), 3.54 - 3.61 (1H, m), 4.32 (2H, q), 7.57 (1H, d), 8.11 (1H, t)

MS m/z 223 M+H


Intermediate # 18: 2-cyano-N-cyclohexyl-acetamide



10 Cyanoacetic acid (4.26g, 50mmol) and cyclohexylamine (4.96g, 50mmol) were dissolved in DCM (100mL). EDCI (10.51g, 55mmol) was added and the mixture stirred at ambient temperature for 24h. The reaction mixture was washed with water (2x100mL), dried (MgSO₄) and evaporated to leave a yellow solid residue which was purified by chromatography on silica gel eluting with an EtOAc/DCM gradient (0-100%) to provide
15 the title compound as a white solid (5.66g, 68%).

^1H NMR (300.073 MHz, DMSO- d_6) δ 1.06 - 1.32 (5H, m), 1.51 - 1.55 (1H, m), 1.63 - 1.75 (4H, m), 3.46 - 3.57 (1H, m), 3.55 (2H, s), 8.06 - 8.09 (1H, m)

Intermediate # 19: Methyl 3-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate




20 t-Butylnitrite (145mg, 1.38mmol) and cupric chloride (236mg, 1.76mmol) were added to acetonitrile (10mL) and heated to 65°C. Methyl 3-[5-amino-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate #20**) (400mg, 1.38mmol) was added as a solid in portions giving a vigorous gas evolution. After the addition was complete heating was continued for a further 15min. and then cooled to ambient, diluted
25 with water (50mL) and extracted with ethyl acetate (3x25mL). The combined extracts were

washed with water (2x20mL), dried (MgSO₄) and evaporated to give a brown oil which was purified by chromatography on silica gel eluting with an EtOAc/Hexane gradient (0-50%) to provide the title compound as a clear colourless oil. (256mg, 51%)

¹H NMR (400.13 MHz, CDCl₃) δ 1.15 - 1.25 (3H, m), 1.32 - 1.43 (2H, m), 1.58 (1H, d),
5 1.65 - 1.71 (2H, m), 1.94 - 1.98 (2H, m), 3.89 (3H, s), 3.91 - 3.98 (1H, m), 6.04 (1H, d),
7.55 (1H, t), 7.66 - 7.68 (1H, m), 8.08 - 8.10 (2H, m), 8.15 (1H, t)

MS m/z 360 M-H


Intermediate # 20: Methyl 3-[5-amino-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate



10 2-cyano-N-cyclohexyl-3-ethoxy-prop-2-enamide (**Intermediate #17**) (605mg, 2.73mmol) and methyl 3-hydrazinylbenzoate hydrochloride (552mg, 2.73mmol) were suspended in ethanol (20mL). DIPEA (352mg, 2.73mmol) was added and the mixture heated to 70°C for 1h. The reaction mixture was cooled to ambient and concentrated to a small volume. Water (50mL) was added and the resulting solid recovered by filtration,
15 redissolved in ethyl acetate (50mL), dried (MgSO₄) and evaporated to give a brown solid which was slurried with ether and recovered by filtration to give the title compound as light brown solid (443mg, 47%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.20 (5H, d), 1.61 (1H, d), 1.72 - 1.75 (2H, m), 1.80 (2H, d), 3.70 - 3.74 (1H, m), 3.88 (3H, s), 6.47 (2H, s), 7.59 - 7.69 (2H, m), 7.84 - 7.88
20 (1H, m), 7.91 - 7.94 (1H, m), 8.01 (1H, s), 8.12 (1H, t)

MS m/z 343 M+H

Intermediate # 21 Ethyl 4-[4-(cyclohexylcarbamoyl)-5-propyl-pyrazol-1-yl]benzoate

1-(4-chlorophenyl)-N-cyclohexyl-5-propyl-pyrazole-4-carboxamide


(**Intermediate#22**) (159mg, 0.46mmol), Molybdenum hexacarbonyl (61mg, 0.23mmol),

- 5 Herrmann's catalyst (Trans-Di-Mu-Acetato-bis[2-(Di-O-Tolylphosphino)Benzyl]dipalladium(II), 22mg, 0.02mmol), DMAP (113mg, 0.92mmol), DIPEA (161 μ L, 0.92mmol), Fu's salt (Tri-(T-Butyl)Phosphonium Hydrogen tetrafluoroborate Salt, 27mg, 0.09mmol), Dioxane (2mL), Ethanol (2 L) were mixed into a microwave tube. The reaction mixture was then heated at 150°C for 1 hour by microwave.
- 10 LC-MS showed complete conversion to the product. The solvent was evaporated under reduced pressure and the reaction mixture extracted with EtOAc (2x25mL), washed with water (10mL), 2N HCl (10mL) and brine (10mL). The organic phase was dried over MgSO₄ and evaporated under reduced pressure to give a black gum. The residue was purified by column chromatography (12g silicycle column, gradient: 1:0 to 1:1
- 15 hexane/EtOAc) and appropriate fractions combined and concentrated *in vacuo* to yield the desired compound as a white solid (130mg, 74%).

¹H NMR (400.13 MHz, CDCl₃) δ 0.76 - 0.81 (3H, t), 1.09 - 1.21 (3H, m), 1.36 (5H, m), 1.45 - 1.55 (2H, m), 1.61 (1H, m), 1.67 - 1.72 (2H, m), 1.96 - 1.98 (2H, m), 2.89 - 2.93 (2H, m), 3.85 - 3.92 (1H, m), 4.35 (2H, q), 5.62 (1H, d), 7.41 - 7.44 (2H, m), 7.71 (1H, s), 8.10 -

20 8.13 (2H, m)

MS m/z 384 M+H


Intermediate #22: 1-(4-chlorophenyl)-N-cyclohexyl-5-propyl-pyrazole-4-carboxamide

To a suspension of 1-(4-chlorophenyl)-5-propyl-pyrazole-4-carbonyl chloride (commercially available, 302 mg, 1.07mmol) in DCM (5 ml) was added 1 drop of DMF
5 followed by cyclohexylamine (306 μ L, 2.68mmol). The reaction mixture was stirred at room temperature for two hours then stopped.

It was extracted in DCM (10mL), washed with 2N NaOH (5mL), 2N HCl (5mL), water (mL) and brine (5mL). The organic phase was dried over MgSO₄ and evaporated under reduced pressure to give a white solid (200mg, 54%), which was used without
10 further purification.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.73 (3H, t), 1.13 (1H, d), 1.22 - 1.31 (4H, m), 1.35 - 1.44 (2H, m), 1.60 - 1.63 (1H, m), 1.69 - 1.75 (2H, m), 1.80 - 1.82 (2H, m), 2.91 (2H, t), 3.72 - 3.75 (1H, m), 7.48 - 7.52 (2H, m), 7.61 - 7.64 (2H, m), 7.84 (1H, d), 8.13 (1H, s)
MS m/z 346 M+H

Intermediate #23: Ethyl 4-[4-(cyclohexyl-methyl-carbamoyl)-5-propyl-pyrazol-1-yl]benzoate




1-(4-chlorophenyl)-N-cyclohexyl-N-methyl-5-propyl-pyrazole-4-carboxamide
 5 **(Intermediate #24)**, 101mg, 0.28mmol), Molybdenum hexacarbonyl (37mg, 0.14mmol),
 Herrmann's catalyst (Trans-Di-Mu-Acetato-bis[2-(Di-O-
 Tolyphosphino)Benzyl]dipalladium(II), 14 mg, 0.01 mmol), DMAP (69 mg, 0.56 mmol),
 DIPEA (98 μ L, 0.56mmol), Fu's salt (Tri-(T-Butyl)Phosphonium Hydrogen
 tetrafluoroborate Salt, 17mg, 0.06mmol), Dioxane (2mL), Ethanol (2mL) were mixed into
 10 a microwave tube. The reaction mixture was then heated at 150°C for 1 hour by
 microwave.

LC-MS showed complete conversion to the product. The same reaction was
 repeated in another microwave tube and the two reaction mixtures were combined for work
 up and purification. The solvent was evaporated under reduced pressure and the reaction
 15 mixture extracted with EtOAc (2x25mL), washed with water (10mL), 2N HCl (10mL) and
 brine (10mL). The organic phase was dried over MgSO₄ and evaporated under reduced
 pressure to give a black gum. The residue was purified by column chromatography (12g
 silicycle column, gradient: 1:0 to 1:1 hexane/EtOAc) and appropriate fractions combined
 and concentrated *in vacuo* to yield the desired compound as a white solid (105mg, 61%).
 20 ¹H NMR (400.13 MHz, CDCl₃) δ 0.71 - 0.79 (3H, t), 1.02 - 1.10 (1H, m), 1.19 (2H, t), 1.31
 - 1.40 (5H, m), 1.55 - 1.72 (4H, m), 1.75 - 1.78 (2H, m), 2.73 - 2.79 (2H, m), 2.89 (3H,
 m), 4.05 (1H, q), 4.35 (2H, q), 6.99 - 7.11 (1H, m), 7.46 (2H, d), 7.55 (1H, s), 8.10 - 8.13
 (2H, d)

MS m/z 398 M+H

Intermediate #24: 1-(4-chlorophenyl)-N-cyclohexyl-N-methyl-5-propyl-pyrazole-4-carboxamide




5 To a suspension of 1-(4-chlorophenyl)-5-propyl-pyrazole-4-carbonyl chloride (commercially available, 302mg, 1.07mmol) in DCM (5ml) was added 1 drop of DMF followed by N-methylcyclohexylamine (349 μ L, 2.68mmol). The reaction mixture was stirred at room temperature for two hours then stopped.

10 It was extracted in DCM (10mL), washed with 2N NaOH (5mL), 2N HCl (5mL), water (5mL) and brine (5mL). The organic phase was dried over MgSO₄ and evaporated under reduced pressure to give a colourless gum (263mg, 73%), which was used without further purification.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.70 (3H, t), 1.20 - 1.46 (5H, m), 1.58 - 1.63 (5H, m), 1.76 - 1.79 (2H, m), 2.73 - 2.74 (3H, m), 2.89 (4H, m), 7.54 - 7.57 (2H, m), 7.61 - 7.64
15 (2H, m)

MS m/z 360 M+H

Intermediate # 25: 1-(4-bromophenyl)-N-cyclohexyl-5-cyclopropyl-pyrazole-4-carboxamide







1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylic acid (**Intermediate#30**)
5 (237mg, 0.77mmol) was suspended in DCM (5mL). One drop of DMF was added and then oxalyl chloride (200μL, 2.32mmol) was added slowly. The reaction mixture was stirred at ambient temperature for 4h and then evaporated under reduced pressure. The residue was dissolved in DCM (5mL) and added to a solution of cyclohexylamine (97μL, 0.85mmol) and DIPEA (403μL, 2.32mmol) in DCM (5mL). The mixture was stirred at ambient for
10 24h and then water, (10mL) was added and the mixture passed through a phase separating filter. The product was recovered from the filtrate by flash column chromatography (SiO₂, elution gradient 0-100% EtOAc in isohexane). Pure fractions were combined and evaporated to give 1-(4-bromophenyl)-N-cyclohexyl-5-cyclopropyl-pyrazole-4-carboxamide as white solid. (277mg, 92%)

15 ¹H NMR (400.13 MHz, DMSO-d₆) δ 0.39 - 0.43 (2H, m), 0.84 - 0.89 (2H, m), 1.12 - 1.20 (1H, m), 1.24 - 1.36 (4H, m), 1.61 (1H, d), 1.69 - 1.75 (2H, m), 1.87 (2H, d), 2.14 - 2.21 (1H, m), 3.72 - 3.77 (1H, m), 7.73 - 7.79 (3H, m), 7.92 - 7.98 (1H, m), 8.07 - 8.10 (2H, m), 13.17 (1H, s)


MS m/z (ESI+) (M+H)⁺ 390

20 The following intermediates were prepared from 1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylic acid (**Intermediate#30**) in a similar manner to Intermediate #25, using the appropriate amine starting material:

Structure	Int.#	Name	¹ H NMR δ	MS m/e MH ⁺
	26	N-(2-adamantyl)-1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxamide	¹ H NMR (300.073 MHz, DMSO-d ₆) δ 0.41 - 0.46 (2H, m), 0.79 - 0.88 (2H, m), 1.54 (2H, d), 1.72 (2H, s), 1.83 (6H, d), 1.94 (2H, s), 2.05 (2H, d), 2.13 - 2.22 (1H, m), 4.04 (1H, d), 7.52 (1H, d), 7.56 (2H, d), 7.73 (2H, d), 7.89 (1H, s)	440
	27	N-(1-adamantyl)-1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxamide	¹ H NMR (300.073 MHz, DMSO-d ₆) δ 0.41 - 0.46 (2H, m), 0.83 - 0.89 (2H, m), 1.65 (6H, s), 2.05 (9H, s), 2.08 - 2.14 (1H, m), 7.13 (1H, s), 7.51 - 7.56 (2H, m), 7.69 - 7.74 (2H, m), 7.81 (1H, s)	440
	28	1-(4-bromophenyl)-N-cyclohexyl-5-cyclopropyl-N-methyl-pyrazole-4-carboxamide	¹ H NMR (300.073 MHz, DMSO-d ₆) δ 0.35-0.55 (2H,m) 0.79 - 0.81 (2H, m), 1.11 - 1.23 (2H, m), 1.33 (1H, s), 1.55 (2H, s), 1.61 - 1.65 (3H, m), 1.76 (2H, s), 1.98 (1H, s), 2.85 (3H, s), 3.43-3.70 (0.4H, s), 4.31 (0.6H, s), 7.63 (3H, d),	402

Structure	Int.#	Name	¹ H NMR δ	MS m/e MH ⁺
			7.72 (2H, d)	
	29	1-(4-bromophenyl)-5-cyclopropyl-N-(4-hydroxy-1-adamantyl)pyrazole-4-carboxamide	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 0.43 - 0.47 (2H, m), 0.83 - 0.88 (2H, m), 1.38 (2H, d), 1.65 (4H, d), 1.75 (2H, d), 1.95 (2H, d), 2.03 - 2.08 (3H, m), 3.97 (1H, t), 4.39 (1H, s), 7.47 - 7.49 (1H, m), 7.56 - 7.59 (2H, m), 7.72 - 7.76 (2H, m), 7.91 (1H, s)	458

Intermediate#30 1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylic acid




Ethyl 1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylate (**Intermediate# 31**) (423mg, 1.29mmol) was dissolved in methanol (20mL) and treated with 2M aqueous sodium hydroxide solution (3.23mL, 6.46mmol). The mixture was stirred at ambient temperature for 24h and then the methanol was removed by evaporation under reduced pressure. The residue was dissolved in water (50mL), acidified to pH4 with 2M HCl and extracted with EtOAc (3x25mL). The combined extracts were washed with water and

brine, dried (MgSO₄) and evaporated to leave 1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylic acid a white solid. (343mg, 87%).

¹H NMR (300.073 MHz, DMSO-d₆) δ0.46 - 0.57 (2H, m), 0.82 - 0.87 (2H, m), 1.98 - 2.11 (1H, m), 7.55 - 7.62 (2H, m), 7.71 - 7.75 (2H, m), 7.94 (1H, s), 12.32 (1H, s)

5 MS m/z (ESI+) (M+H)⁺ 309

Intermediate# 31 Ethyl 1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylate




Ethyl -2-(cyclopropanecarbonyl)-3-dimethylamino-prop-2-enoate

(Intermediate#32) (316mg, 1.5mmol) was dissolved in ethanol (5mL). 4-

10 bromophenylhydrazine hydrochloride (335mg, 1.5mmol) and DIPEA (264μL, 1.5mmol) were added and the mixture was heated to reflux for 2h. The reaction mixture was cooled to ambient and evaporated under reduced pressure. The residue was dissolved in DCM (10mL) washed with water and poured through a phase separating tube. The product was recovered from the filtrate by flash column chromatography (SiO₂, elution gradient 0-25%
15 EtOAc in isohexane). Pure fractions were combined and evaporated to give ethyl 1-(4-bromophenyl)-5-cyclopropyl-pyrazole-4-carboxylate as an oil that crystallised on standing. (343mg, 87%)

¹H NMR (300.073 MHz, DMSO-d₆) δ0.43 - 0.49 (2H, m), 0.84 - 0.90 (2H, m), 1.29 (3H, t), 2.05 - 2.14 (1H, m), 4.24 (2H, q), 7.55 - 7.60 (2H, m), 7.71 - 7.76 (2H, m), 7.98 (1H, d)


20 MS m/z (ESI+) (M⁺) 335

Intermediate#32 Ethyl -2-(cyclopropanecarbonyl)-3-dimethylamino-prop-2-enoate

Ethyl 3-cyclopropane-3-oxopropionionate (312mg, 2mmol) and N,N-dimethylformamide dimethylacetal (402μL, 3mmol) were dissolved in dioxan (5mL) and heated to reflux for 2h. The reaction mixture was cooled to ambient and evaporated under reduced pressure to leave a yellow oil, which was purified by flash column chromatography (SiO₂, elution gradient 0-100% EtOAc in isohexane). Pure fractions were combined and evaporated to give ethyl -2-(cyclopropanecarbonyl)-3-dimethylamino-prop-2-enoate as an oil. (316mg, 74%)

¹H NMR (300.073 MHz, DMSO-d₆) δ0.69 - 0.80 (4H, m), 1.15 - 1.24 (3H, m), 2.26 - 2.34 (1H, m), 2.94 (6H, s), 4.11 (2H, q), 7.57 (1H, s)

MS m/z (ESI+) (M+H)+ 212

Intermediate #33 : Methyl 2-[4-[4-(cyclohexylcarbonyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetate


1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylic acid (**Intermediate#35**) (220mg, 0.66mmol) was dissolved in DCM (5mL) and 1 drop of DMF added. Oxalyl chloride (176μL, 1.96mmol) was added and the mixture stirred at ambient temperature for 3h. Volatiles were removed by evaporation under reduced pressure and the resulting gum re-dissolved in DCM (2mL) and added at ambient temperature to a solution

of cyclohexylamine (66mg, 0.66mmol) and DIPEA (230 μ L, 1.32mmol) in DCM(5mL). The reaction mixture was stirred at ambient temperature for 2h, diluted with EtOAc (50mL) and washed with water (2x10mL) and brine (10mL), dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by flash chromatography on silica gel (elution gradient 0-100% EtOAc in isohexane) to afford methyl 2-[4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetate as a white solid. (220mg, 88%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.69 (3H, t), 1.20 - 1.32 (7H, m), 1.58 (1H, d), 1.69 - 1.74 (2H, m), 1.83 (2H, d), 2.68 (2H, t), 3.64 (3H, s), 3.74 - 3.82 (1H, m), 3.80 (2H, s), 7.44 (4H, s), 7.84 - 7.89 (1H, m), 8.08 (1H, s)

MS m/z (ESI+) (M+H)⁺ 416

Intermediate #34 : Methyl 2-[4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetate




Prepared from 1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylic acid (**Intermediate#35**) and 2-adamantylamine hydrochloride by the same procedure as used for Intermediate #6.

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.67 (3H, t), 1.15 - 1.31 (2H, m), 1.61 (2H, d), 1.73 (2H, s), 1.83 (6H, s), 1.96 (4H, d), 2.61 (2H, t), 3.64 (3H, s), 3.80 (2H, s), 4.09 (1H, d), 7.40 - 7.52 (4H, m), 8.02 (1H, d), 8.11 (1H, s)


MS m/z (ESI+) (M+H)⁺ 468

Intermediate #35 : 1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylic acid



- tert-butyl 1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylate (**Intermediate#36**) was dissolved in DCM (10mL). TFA (2mL) was added and the mixture was stirred at ambient temperature for 5h. Volatiles were then evaporated under reduced pressure and the residue dried under vacuum to afford 1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylic acid as a light brown oil. (440mg, 98%)
- ¹H NMR (300.073 MHz, DMSO-d₆) δ 0.70 (3H, t), 1.21 - 1.33 (2H, m), 2.79 (2H, t), 3.64 (3H, s), 3.80 (2H, s), 7.44 (4H, s), 8.10 (1H, s)


Intermediate #36 : tert-butyl 1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylate



tert-butyl 5-chloro-1-[4-(methoxycarbonylmethyl)phenyl]pyrazole-4-carboxylate (**Intermediate# 37**) (767mg, 2.19mmol) was dissolved in butyronitrile (10mL). Potassium carbonate (906mg, 6.57mmol) and propanethiol (284μL, 2.74mmol) were added and the mixture heated at 90°C for 18h. The reaction mixture was cooled to ambient and ethyl acetate (50mL) was added. The mixture was washed with water (4x25mL), dried (MgSO₄) and evaporated under reduced pressure. The residue was purified by flash chromatography on silica gel (elution gradient 0-25% EtOAc in isohexane) to afford tert-butyl 1-[4-(methoxycarbonylmethyl)phenyl]-5-propylsulfanyl-pyrazole-4-carboxylate as a yellow oil. (527mg, 62%).

¹H NMR (300.073 MHz, DMSO-d₆) δ 0.70 (3H, t), 1.27 (2H, q), 1.53 - 1.54 (9 s, 2.75 (2H, t), 3.64 (3H, s), 3.80 (2H, s), 7.44 (4H, s), 8.07 - 8.12 (1H, m)
MS m/z (ESI+) (M+H)⁺ 391

Intermediate #37 : tert-butyl 5-chloro-1-[4-(methoxycarbonylmethyl)phenyl]pyrazole-4-carboxylate




t-Butylnitrite (419μL, 3.49mmol) and copper (2) chloride (585mg, 4.37mmol) were added to acetonitrile (10mL) and heated to 65°C. Tert-butyl 5-amino-1-[4-(methoxycarbonylmethyl)phenyl]pyrazole-4-carboxylate (**Intermediate# 38**) (965mg, 2.91mmol) was added as a solution in acetonitrile (5mL) giving a vigorous gas evolution. After the addition was complete heating was continued for a further 15min. The reaction mixture was cooled to ambient, diluted with water (50mL) and extracted with ethyl acetate (3x50mL). The combined extracts were washed with water (2x50mL) and brine (10mL), dried (MgSO₄) and evaporated to give a brown oil which was purified by

flash chromatography on silica gel (elution gradient 0-50% EtOAc in isohexane) to give the title compound as a yellow oil. (785mg, 77%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.52 (9H, s), 3.62 - 3.64 (3H, m), 3.81 (2H, s), 7.48 - 7.54 (4H, m), 8.12 (1H, s)

5 MS m/z (ESI+) (M+H)+ 351


Intermediate# 38: tert-butyl 5-amino-1-[4-(methoxycarbonylmethyl)phenyl]pyrazole-4-carboxylate



t-Butyl cyanoacetate (3.525g, 25mmol) was dissolved in triethyl orthoformate
10 (6.2mL, 37.5mmol). Acetic anhydride (2.41mL, 25mmol) was added and the mixture heated to 125°C for 3h. Volatiles were then removed by evaporation under reduced pressure to leave an orange oil. The crude oil was dissolved in methanol and methyl 2-(4-hydrazinylphenyl)acetate hydrochloride (**Intermediate# 39**) (1.62g, 7.5mmol) and DIPEA (1.3mL, 7.5mmol) were added. The mixture was heated to reflux for 2h and then
15 evaporated to dryness. The residue was dissolved in ethyl acetate (100mL), washed with water (2x50mL) and brine (50mL), dried (MgSO₄) and evaporated to leave a crude product which was purified by flash chromatography on silica gel (elution gradient 0-50% EtOAc in isohexane) to give the title compound as a yellow solid.
(974mg, 39%)


20 ¹H NMR (300.073 MHz, DMSO-d₆) δ 1.51 (9H, d), 3.63 (3H, s), 3.76 (2H, s), 6.20 (2H, s), 7.41 (2H, d), 7.48 (2H, d), 7.60 (1H, s)

MS m/z (ESI+) (M+H)+ 331

Intermediate# 39 : methyl 2-(4-hydrazinylphenyl)acetate hydrochloride

A solution of methyl 4-iodophenylacetate (4.25g, 15.39mmol) was dissolved in DMF (25mL). BOC-carbazate (2.44g, 18.47mmol); 1,10-phenanthroline (278mg, 1.54mmol), copper iodide (147mg, 0.77mmol) and cesium carbonate (7.0g, 21.55mmol) were added and the mixture heated at 120°C for 60min. The reaction mixture was diluted with ethyl acetate (25mL), washed with water (3x10mL) and brine (10mL), dried (MgSO₄) and evaporated to leave a brown oil which was purified by flash chromatography on silica gel (elution gradient 0-50% EtOAc in isohexane) to give a colourless oil. The oil was dissolved in 4M HCl Dioxan (20mL) and stirred at ambient temperature for 3h giving a thick precipitate. Diethyl ether (100mL) was added and the precipitate recovered by filtration, washed with ether (2x20mL) and dried under vacuum to leave methyl 2-(4-hydrazinylphenyl)acetate_hydrochloride as a light brown solid. (1.63g, 49%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 3.59 (5H, s), 6.93 (2H, d), 7.16 (2H, d), 8.21 (1H, s), 10.20 (3H, s)


Intermediate#40: 3-(4-Cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid methyl ester



A solution of 1-(4-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid (**Intermediate#44**) (118mg, 0.35mmol), cyclohexylamine (49μl, 0.42mmol), EDAC (75mg, 0.39mmol) and DMAP (9mg, 0.07mmol) in DCM (5ml) was stirred overnight at ambient temperature. The solvent was evaporated in vacuo and the resulting residue was partitioned between EtOAc (~30ml) and citric acid (~20ml). The layers were separated and the aqueous layer was re-extracted with EtOAc (~10ml). The

combined organic layers were washed with brine (~10ml) then dried (MgSO₄), filtered and evaporated to yield an oil. The oil was purified by column chromatography (4g Si, eluting with 20 to 60% EtOAc in IH) to yield the title compound as a pale yellow oil (116mg, 79%).


- 5 ¹H NMR (300.072 MHz, CDCl₃) δ0.81 (3H, t), 1.13 - 1.25 (3H, m), 1.30 - 1.48 (4H, m), 1.52 - 1.69 (3H, m), 1.89 - 1.94 (2H, m), 2.47 (2H, t), 3.83 (3H, s), 3.88 - 3.98 (1H, m), 5.49 (2H, s), 7.18 (2H, d), 7.24 (1H, d), 7.92 (2H, d), 8.07 (1H, s)
MS m/e MH⁺ 416.

- 10 The following intermediates were made by the above procedure using the corresponding starting materials

Structure	Intermediate #	Name	NMR	[M+H] ⁺
	41	3-(4-Cyclohexylcarbonyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid methyl ester	¹ H NMR (300.072 MHz, CDCl ₃) δ0.82 (3H, t), 1.13 - 1.26 (3H, m), 1.30 - 1.50 (4H, m), 1.50 - 1.70 (3H, m), 1.85 - 1.95 (2H, m), 2.48 (2H, t), 3.84 (3H, s), 3.88 - 3.96 (1H, m), 5.48 (2H, s), 7.31 - 7.34 (3H, m), 7.86 - 7.91 (2H, m), 8.06 (1H, s)	416

	42	3-[4-(Adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-ylmethyl]-benzoic acid methyl ester	1H NMR (300.072 MHz, CDCl ₃) δ 0.81 (3H, t), 1.37 - 1.50 (2H, m), 1.58 - 1.75 (4H, m), 1.82 (8H, s), 1.95 (2H, s), 2.50 (2H, t), 3.84 (3H, s), 4.20 - 4.23 (1H, m), 5.49 (2H, s), 7.33 - 7.35 (2H, m), 7.83 - 7.92 (3H, m), 8.09 (1H, s)	468
	43	4-[4-(Adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-ylmethyl]-benzoic acid methyl ester	1H NMR (300.072 MHz, CDCl ₃) δ 0.80 (3H, t), 1.36 - 1.49 (2H, m), 1.58 - 1.77 (4H, m), 1.82 (8H, s), 1.95 (2H, s), 2.49 (2H, t), 3.83 (3H, s), 4.20 - 4.23 (1H, m), 5.50 (2H, s), 7.19 (2H, d), 7.84 (1H, d), 7.92 (2H, d), 8.10 (1H, s)	468

Intermediate#44: 1-(4-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid




TFA (1.5ml) was added to a stirred solution of 1-(4-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid *tert*-butyl ester (**Intermediate# 46**)

(274mg, 0.70mmol) in DCM (6ml). The reaction was stirred at ambient temperature for 3 hours then evaporated in vacuo. The resulting gum was sonicated with isohexane to yield a solid. The solvent was removed then the white solid was dissolved in EtOAc and evaporated in vacuo to yield the title compound as an off white solid (236mg, 100%).

¹H NMR (300.072 MHz, CDCl₃) δ0.84 (3H, t), 1.35 - 1.50 (2H, m), 2.81 (2H, t), 3.83 (3H, s), 5.53 (2H, s), 7.18 - 7.21 (2H, d), 7.91 - 7.94 (2H, d), 8.05 (1H, s)

MS m/e MH⁺ 335.

Intermediate# 45: 1-(3-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid




Compound prepared in an analogous manner to Intermediate 44, replacing 1-(4-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid *tert*-butyl ester with 1-(3-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid *tert*-butyl ester (**Intermediate# 47**)

¹H NMR (300.072 MHz, CDCl₃) δ0.85 (3H, t), 1.37 - 1.49 (2H, m), 2.82 (2H, t), 3.83 (3H, s), 5.52 (2H, s), 7.33 - 7.34 (2H, m), 7.88 - 7.91 (2H, m), 8.05 (1H, s)

MS m/e [M-H]⁻ 333.

Intermediate# 46: 1-(4-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid *tert*-butyl ester




5-Chloro-1-(4-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester (**Intermediate#48**) (397mg, 1.13mmol) was dissolved in butyronitrile (8mL),

potassium carbonate (470mg, 3.40mmol) and propanethiol (128 μ l, 1.41mmol) were added and the mixture heated to reflux and stirred at this temperature over night. The reaction mixture was transferred to a microwave vial (using ~5ml extra Butyronitrile to ensure complete transfer), a further 3eq of propane thiol was added and the reaction was heated at 180⁰C for 4 hours. EtOAc (~50mL) was added and the mixture washed with water (3 x ~20mL) and brine (~20ml) then dried (MgSO₄), filtered and evaporated to give a yellow oil. Oil purified by column chromatography (12g Si, eluting with 10 to 50% EtOAc in IH) to yield the title compound as a yellow oil (274mg, 62%).

¹H NMR (300.072 MHz, CDCl₃) δ 0.90 (3H, t), 1.40 - 1.52 (2H, m), 1.55 (9H, s), 2.84 (2H, t), 3.90 (3H, s), 5.57 (2H, s), 7.25 (2H, d), 7.96 - 8.00 (3H, m)

MS m/e MH⁺ 391.

Intermediate# 47: 1-(3-Methoxycarbonyl-benzyl)-5-propylsulfanyl-1H-pyrazole-4-carboxylic acid *tert*-butyl ester




Compound prepared in an analogous manner to **Intermediate# 46**, replacing 5-Chloro-1-(4-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester with 5-Chloro-1-(3-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester

¹H NMR (300.072 MHz, CDCl₃) δ 0.91 (3H, t), 1.42 - 1.54 (2H, m), 1.56 (9H, s), 2.84 (2H, t), 3.90 (3H, s), 5.56 (2H, s), 7.37 - 7.41 (2H, m), 7.93 - 7.97 (3H, m)

MS m/e MH⁺ 391.

Intermediate# 48: 5-Chloro-1-(4-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester




t-Butylnitrite (417□l, 3.48mmol) and copper chloride (583mg, 4.35mmol) were added to acetonitrile (20mL) and heated to 65°C. A solution of 5-Amino-1-(4-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester (**Intermediate# 50**) in acetonitrile (~4ml) was added dropwise. After the addition was complete heating was continued for a further 15min. Cooled to ambient, diluted with water (~100mL) and extracted with ethyl acetate (3 x ~40mL). The combined extracts were washed with water (2 x ~20mL) and brine (~20mL), dried (MgSO₄), filtered and evaporated to an oil. Chromatographed (40g silica column, EtOAc/ Hexane 10-50%) to yield the title compound as an off white solid (397mg, 39%).

¹H NMR (300.072 MHz, CDCl₃) δ 1.49 (9H, s), 3.83 (3H, s), 5.33 (2H, s), 7.19 (2H, d), 7.84 (1H, s), 7.93 (2H, d)

MS m/e [M+Na]⁺ 373.

Interemdiate# 49: 5-Chloro-1-(3-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester




Compound prepared in an analogous manner to Intermediate# 48, replacing 5-Amino-1-(4-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester with 5-Amino-1-(3-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester (**Intermediate# 51**)

^1H NMR (300.072 MHz, CDCl_3) δ 1.49 (9H, s), 3.84 (3H, s), 5.32 (2H, s), 7.33 - 7.35 (2H, m), 7.84 (1H, s), 7.88 - 7.93 (2H, m)

MS m/e $[\text{M}-\text{tBu}+\text{H}]^+$ 295.

Intermediate# 50: 5-Amino-1-(4-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester




t-Butyl cyanoacetate (600mg, 4.25mmol) was dissolved in triethyl orthoformate (1.06ml, 6.38mmol), acetic anhydride (0.40ml, 4.25mmol) was added and the mixture heated to 125°C for 2 hours. The solution was evaporated to leave a yellow oil. This oil was redissolved in ethanol (5ml) and treated with 4-Hydrazinomethyl-benzoic acid methyl ester hydrochloride (**Intermediate# 52**) (230mg, 1.06mmol) and DIPEA (184mg, 1.06mmol). The resulting mixture was heated to reflux for 2h and then cooled to ambient temperature and evaporated to leave a brown oil. This oil was dissolved in ethyl acetate (50mL), washed with water (2x20mL) and brine (20mL) then dried (MgSO_4), filtered and evaporated. The residue was chromatographed (40g silica EtOAc/ Hexane 0-50%) to yield the title compound as a yellow solid (214mg, 43%).

^1H NMR (300.072 MHz, CDCl_3) δ 1.55 (9H, s), 3.90 (3H, s), 4.86 (2H, s), 5.18 (2H, s), 7.21 (2H, d), 7.62 (1H, s), 8.00 (2H, d)

MS m/e $[\text{M}-\text{H}]^-$ 330.

Intermediate# 51: 5-Amino-1-(3-methoxycarbonyl-benzyl)-1H-pyrazole-4-carboxylic acid *tert*-butyl ester




Compound prepared in an analogous manner to Intermediate#50, replacing 4-

Hydrazinomethyl-benzoic acid methyl ester hydrochloride with 3-Hydrazinomethyl-benzoic acid methyl ester hydrochloride (**Intermediate# 53**)

^1H NMR (300.072 MHz, CDCl_3) δ 1.47 (9H, s), 3.84 (3H, s), 4.75 (2H, s), 5.11 (2H, s), 7.26 (1H, d), 7.36 (1H, t), 7.55 (1H, s), 7.83 (1H, s), 7.90 (1H, d)

5 MS m/e MH^+ 332.


Intermediate# 52: 4-Hydrazinomethyl-benzoic acid methyl ester hydrochloride



A solution of methyl 4-[[[(2-methylpropan-2-yl)oxycarbonyl-[(2-methylpropan-2-yl)oxycarbonylamino]amino]methyl]benzoate (**Intermediate# 54**) (3.3g, 8.67mmol) in 4M HCl in Dioxane (100mL) was stirred overnight at ambient temperature. The solvent was removed under reduced pressure and the resulting solid was dissolved in hot MeOH. The hot suspension was filtered then evaporated in vacuo to give a solid. This solid was triturated with ether, filtered then dried under high vac to yield the title compound as a yellow solid (1.5g, 80%).

15 ^1H NMR (300.073 MHz, $\text{DMSO}-d_6$) δ 3.85 (3H, s), 4.13 (2H, s), 7.55 (2H, d), 7.93 (2H, d)

Intermediate# 53: 3-Hydrazinomethyl-benzoic acid methyl ester hydrochloride




Compound prepared in an analogous manner to intermediate# 52, replacing 4-[[[(2-methylpropan-2-yl)oxycarbonyl-[(2-methylpropan-2-yl)oxycarbonylamino]amino]methyl]benzoate with 3-[[[(2-methylpropan-2-yl)oxycarbonyl-[(2-methylpropan-2-yl)oxycarbonylamino]amino]methyl]benzoate (**Intermediate# 55**)

^1H NMR (300.073 MHz, $\text{DMSO}-d_6$) δ 3.86 (3H, s), 4.13 (2H, s), 7.52 (1H, t), 7.70 (1H, d), 7.91 (1H, d), 8.02 (1H, s), 8.77 (4H, br s)


MS m/e MH^+ 181.

Intermediate# 54: methyl 4-[[[(2-methylpropan-2-yl)oxycarbonyl]-[(2-methylpropan-2-yl)oxycarbonylamino]amino]methyl]benzoate




NaH (362mg, 9.04mmol) was added to a stirred solution of Di-Tert-Butyl
 5 Hydrazodicarboxylate (2g, 8.61mmol) in anhydrous THF (50ml). The reaction was stirred
 at ambient temperature for 10 minutes then treated with a solution of Methyl 4-
 (Bromomethyl)Benzoate (1.98g, 8.61mmol) in anhydrous THF (20ml). The reaction was
 stirred at ambient temperature for 4 hours. The reaction was partitioned between ether
 (~100ml) and water (~100ml). The layers were separated and the aqueous layer was re-
 10 extracted with ether (~50ml). The ether layers were combined, washed with brine
 (~50ml), dried (MgSO₄), filtered and evaporated to yield the title compound (3.3g, 100%).
¹H NMR (300.073 MHz, DMSO-d₆) δ 1.38 (18H, s), 3.84 (3H, s), 4.52 (2H, br s), 7.42
 (2H, d), 7.90 (2H, d), 9.19 (1H, s)
 MS m/e [M+Na]⁺ 403.

Intermediate# 55: methyl 3-[[[(2-methylpropan-2-yl)oxycarbonyl]-[(2-methylpropan-2-yl)oxycarbonylamino]amino]methyl]benzoate



Compound prepared in an analogous manner to Intermediate # 54, replacing
 Methyl 4-(Bromomethyl)Benzoate with Methyl 3-(Bromomethyl)Benzoate
 20 ¹H NMR (400.13 MHz, DMSO-d₆) δ 1.39 (18H, s), 3.86 (3H, s), 4.50 - 4.58 (2H, m), 7.45
 - 7.52 (1H, m), 7.57 (1H, d), 7.85 - 7.90 (2H, m), 9.25 (1H, s)
 MS m/e [M+Na]⁺ 403.

Intermediate#56: methyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoate



Methyl 4-hydrazinylbenzoate hydrochloride (**Intermediate# 123**) (3.04 g, 15.00 mmol) was added in one portion to (2)-N-(2-adamantyl)-2-(dimethylaminomethylidene)-4,4-dimethyl-3-oxo-pentanamide (**Intermediate# 58**) (4.99 g, 15 mmol) in ethanol (100 mL). 5 drops of acetic acid were added and the resulting solution was stirred at 80 °C for 2 hours. The reaction mixture was concentrated and diluted with EtOAc (500 mL), and washed sequentially with water (200 mL), and saturated brine (200 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoate (4.66 g, 71.3 %) as a yellow solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.19 (9H, s), 1.50 (2H, d), 1.69-1.95 (10H, m), 2.09 (2H, d), 3.91 (3H, s), 3.99 (1H, d), 7.53 - 7.56 (2H, m), 7.62 (1H, s), 8.09 - 8.12 (2H, m), 8.20 (1H, d)

m/z (ESI+) (M+H)⁺ = 436

Intermediate# 56 may also be prepared as follows:


2-(4-(Methoxycarbonyl)phenyl)hydrazinium chloride (**Intermediate# 123**) (1 equiv.) and then acetic acid (0.023 equivs.) were added to a solution of (2Z)-N-(2-adamantyl)-2-(dimethylamino-methylidene)-4,4-dimethyl-3-oxo-pentanamide (**Intermediate# 58**) (1 equiv.) in methanol (200 vols.), under nitrogen. The mixture stirred under reflux for 1.5 hours, cooled, concentrated to below 3.5 vols. and the resulting suspension diluted with

ethyl acetate (96 vols.). The suspension was washed with water (34.4 vols.) giving a solution which was washed with brine (34.4 vols.), dried (MgSO₄) and concentrated to dryness. The crude product was slurried in MTBE (9 vols.) and stirred for 15 minutes. The pale yellow solid was filtered, washed with MTBE (11.4 vols.) and dried under vacuum at 60 °C.

TLC DCM : MeOH, 9 : 1, Product R_f 0.86 (trace impurity R_f 0.68)

mp 193.6 – 194.5 °C

Intermediate#57: methyl 4-[4-(2-adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoate




Methyl 4-hydrazinylbenzoate hydrochloride (**Intermediate# 123**) (0.809 g, 3.99 mmol) was added in one portion to (Z)-N-(2-adamantyl)-3-dimethylamino-2-(1-methylcyclopropanecarbonyl)prop-2-enamide (**Intermediate#59**), 1.320 g, 3.99 mmol) in ethanol (30 mL). 5 drops of acetic acid were added and the resulting solution was stirred at 80 °C for 2 hours. The reaction mixture was concentrated and diluted with EtOAc (100 mL), and washed sequentially with water (50 mL), and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(2-adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoate (1.221 g, 70.5 %) as a cream solid.

m/z (ESI+) (M+H)+ = 434; HPLC t_R = 2.98 min.

1H NMR (400.13 MHz, DMSO- d_6) δ 0.48 - 0.51 (2H, m), 0.67 - 0.69 (2H, m), 1.54 - 1.57 (5H, m), 1.73 (2H, s), 1.83 - 1.86 (6H, m), 1.97 (2H, s), 2.04 - 2.07 (2H, m), 3.90 (3H, s), 4.05 - 4.10 (1H, m), 7.50 (1H, d), 7.71 (2H, d), 8.09 (1H, s), 8.13 (2H, d)

5 **Intermediate# 58: (2)-N-(2-adamantyl)-2-(dimethylaminomethylidene)-4,4-dimethyl-3-oxo-pentanamide**



N,N -Dimethylformamide dimethyl acetal (3.02 mL, 22.71 mmol) was added to a stirred suspension of N -(2-adamantyl)-4,4-dimethyl-3-oxo-pentanamide (**Intermediate#**
10 **60**) (5.25 g, 18.93 mmol) in 1,4-dioxane (50 mL) under nitrogen. The resulting mixture was stirred at 100 °C for 2 hours. The reaction mixture was evaporated to dryness and the resulting pale cream solid was dried under vacuum to afford (2)- N -(2-adamantyl)-2-(dimethylaminomethylidene)-4,4-dimethyl-3-oxo-pentanamide (5.83 g, 93 %).

1H NMR (400.13 MHz, DMSO- d_6) δ 1.13 (9H, s), 1.47 (2H, d), 1.69 - 1.83 (10H, m), 2.03
15 (2H, d), 2.92 (6H, s), 3.90 (1H, d), 7.24 (1H, s), 7.94 (1H, d)
 m/z (ESI+) (M+H)+ = 333


Intermediate#58 may also be prepared as follows:

N,N -Dimethylformamide dimethyl acetal (1.2 equivs.) was added to a solution of N -(2-adamantyl)-4,4-dimethyl-3-oxo-pentanamide (**Intermediate# 60**) (1 equiv.) in 1,4-
20 dioxane (9.6 vols.) under nitrogen. The mixture was heated under reflux for five hours and then cooled to room temperature. The solvent was removed *in vacuo* and the pale yellow solid used directly in the next stage.

TLC Hexane : EtOAc, 1 : 1, Product R_f 0.94 (impurities: R_f 0.06 + 0.66)

mp 143.6 – 147.6 °C

Intermediate#59: (Z)-N-(2-adamantyl)-3-dimethylamino-2-(1-methylcyclopropanecarbonyl)prop-2-enamide




Prepared from N-(2-adamantyl)-3-(1-methylcyclopropyl)-3-oxo-propanamide

(Intermediate#61) by the same process as used for Intermediate #58

¹H NMR (400.13 MHz, DMSO-*d*₆) δ 0.54 - 0.55 (2H, m), 0.91 - 0.94 (2H, m), 1.26 (3H, s), 1.51 (2H, d), 1.69 (2H, s), 1.72 - 1.85 (8H, m), 1.92 (2H, d), 3.00 (6H, s), 3.90 - 3.92 (1H, m), 7.57 (1H, s), 8.08 (1H, s)

(Intermediate# 60): N-(2-adamantyl)-4,4-dimethyl-3-oxo-pentanamide



A 1M solution of solution of lithium bis(trimethylsilyl)amide in THF (22.84 ml, 22.84 mmol) was added to THF (25mL) and cooled under nitrogen to -78°C. A solution of 3,3-dimethyl-2-butanone (2.287 g, 22.84 mmol) in THF (25mL) was added drop wise over a period of 5 minutes. The resulting solution was stirred at -78 °C under nitrogen for 15 minutes. A solution of 2-isocyanatoadamantane (*prepared from 2-adamantylamine hydrochloride by the method of R.Reck & C.Jochims Chem. Ber. 115(1982) p864*) (3.68 g, 20.76 mmol) in THF (20mL) was added over a period of 5 minutes. The resulting solution was stirred at -78 °C for 1 hour and then allowed to warm to 20°C over 1h. The reaction mixture was poured into saturated NH₄Cl (150 mL) and extracted with EtOAc (2 x 100 mL), the organic layer was washed with water (50mL) and brine (50mL) dried over MgSO₄, filtered and evaporated to afford a yellow oil. The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford N-(2-adamantyl)-4,4-dimethyl-3-oxo-pentanamide (4.64 g, 81 %) as a white solid.

116.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.08 - 1.09 (9H, m), 1.50 (2H, d), 1.66 - 1.89 (10H, m), 1.95 - 2.00 (2H, m), 3.53 (1.4H, s), 3.80 - 3.94 (1H, m), 5.30 (0.3H, s), 7.77- 7.87 (1H, m), 14.43 (0.3H, s) (2:1 mixture of keto and enol forms)

m/z (ESI+) (M+H)⁺ = 278

5 Intermediate # 60 may also be prepared as follows:


Aqueous sodium hydroxide solution (3M) (5 vols.) was added to a stirred suspension of 2-adamantylamine hydrochloride (1 equiv.) in water (5 vols.). DCM (5 vols.) was added to the resulting thick suspension and the phases separated. The aqueous was extracted with DCM (4 x 5 vols.) and the combined organics concentrated to give the free amine as a
10 white solid.

Ethyl pivaloylacetate (1 equiv.) was added to a suspension of the free amine in xylenes (6.5 vols.), under nitrogen, and the mixture stirred under reflux for 6.5 hours. The batch was cooled to room temperature and concentrated to dryness. The residue was purged with toluene (3 x 1 vol.) followed by hexane (3 x 1 vol.). The resulting solid was
15 digested in hexane at 50 °C for five minutes and then cooled to room temperature. The white solid was filtered, washed with hexane (2 vols.) and dried in air.

TLC Hexane : EtOAc, 1 : 1, Product R_f 0.66

mp 124.5 – 125.1 °C

Intermediate#61: N-(2-adamantyl)-3-(1-methylcyclopropyl)-3-oxo-propanamide




20

Prepared from 1-(1-methylcyclopropyl)ethanone by the same process as used for Intermediate#60

m/z (ESI+) (M+H)⁺ = 276; HPLC t_R = 2.26 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.76 - 0.78 (2H, m), 1.18 - 1.20 (2H, m), 1.25 (3H, s), 1.50 (2H, d), 1.70 - 1.80 (11H, m), 1.95 (2H, d), 3.82 (1H, d), 7.83 (1H, d)
25

Intermediate#62: methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoate




Methyl 4-hydrazinylbenzoate hydrochloride (**Intermediate# 123**) (0.712 g, 3.51 mmol) was added in one portion to (Z)-N-(2-adamantyl)-2-(cyclopentanecarbonyl)-3-dimethylamino-prop-2-enamide (**Intermediate#67**), 1.21 g, 3.51 mmol) in ethanol (30 mL). 5 drops of acetic acid were added and the resulting solution was stirred at 80 °C for 2 hours and then cooled to ambient giving a precipitate. The reaction mixture was filtered and the product recovered, washed with ethanol (10 mL) and water (10 mL) before being dried under vacuum to give methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoate (0.620 g, 39.4 %) as a white solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.48 - 1.53 (4H, m), 1.71 - 1.85 (12H, m), 1.94 (2H, s), 2.02 - 2.12 (4H, m), 2.98 - 3.09 (1H, m), 3.90 (3H, s), 3.98 - 4.03 (1H, m), 7.57 - 7.60 (2H, m), 7.74 - 7.76 (1H, m), 8.10 - 8.15 (3H, m)


m/z (ESI+) (M+H)⁺ = 448; HPLC t_R = 3.26 min.

The same process as used for Intermediate#62 prepared the following intermediates from the appropriate starting material.

Structure	Int#	Name	¹ H NMR δ	MS m/e MH ⁺
The structure shows a central benzene ring substituted at the 4-position with a methyl benzoate group (-COOCH₃) and at the 1-position with a 2-ethylpyrazol-1-yl group. The pyrazole ring is substituted at the 2-position with an ethyl group (-CH₂CH₃). The pyrazole ring is also substituted at the 5-position with an adamantylcarbamoyl group (-CONH-adamantyl).	63	methyl 4-[4-(2-ethylpyrazol-1-yl)-5-(adamantylcarbamoyl)benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.03 (3H, t), 1.53 (2H, d), 1.72 (2H, s), 1.84 (6H, d), 1.95 (2H, s), 2.11 (2H, d), 2.98 (2H, q), 3.90 (3H, s), 4.04 (1H, t), 7.59 - 7.61 (1H, m), 7.65 - 7.68 (2H, m), 8.12 - 8.15 (2H, m), 8.30 (1H, s)	408
The structure shows a central benzene ring substituted at the 4-position with a methyl benzoate group (-COOCH₃) and at the 1-position with a 2-isopropylpyrazol-1-yl group. The pyrazole ring is substituted at the 2-position with an isopropyl group (-CH(CH₃)₂). The pyrazole ring is also substituted at the 5-position with an adamantylcarbamoyl group (-CONH-adamantyl).	64	methyl 4-[4-(2-isopropylpyrazol-1-yl)-5-(adamantylcarbamoyl)benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.30 (6H, d), 1.52 (2H, d), 1.73 (2H, s), 1.82 - 1.86 (6H, m), 1.97 (2H, s), 2.11 (2H, d), 3.11 - 3.18 (1H, m), 3.91 (3H, s), 4.00 - 4.05 (1H, m), 7.57 (2H, d), 7.67 (1H, d), 8.08 (1H, s), 8.13 (2H, d)	422

Structure	Int#	Name	¹ H NMR δ	MS m/e MH ⁺
	65	methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclobutylpyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.51 - 1.54 (2H, m), 1.65 (1H, q), 1.72 (2H, s), 1.76 - 1.85 (7H, m), 1.96 (2H, s), 2.04 - 2.15 (4H, m), 2.18 - 2.26 (2H, m), 3.82 (1H, q), 3.90 (3H, s), 4.00 - 4.06 (1H, m), 7.59 (2H, d), 7.81 (1H, d), 7.95 (1H, s), 8.11 (2H, d)	434

Intermediate# 66: N-(2-adamantyl)-1-(4-cyanophenyl)-5-methyl-pyrazole-4-carboxamide



Acetic acid (0.031 mL, 0.50 mmol) was added in one portion to (2E)-N-(2-adamantyl)-2-(dimethylaminomethylidene)-3-oxo-butanamide (**Intermediate# 71**) (1.45 g, 4.99 mmol) and 4-cyanophenylhydrazine hydrochloride (0.847 g, 4.99 mmol) in ethanol (40 mL). The resulting suspension was stirred at 80 °C for 3 hours. The reaction mixture was concentrated and diluted with EtOAc (75 mL), and washed sequentially with water (50


mL) and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 20 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford N-(2-adamantyl)-1-(4-cyanophenyl)-5-methyl-pyrazole-4-carboxamide (1.3g, 72%) as an orange solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.45-1.58 (2H, m), 1.70-1.9 (8H, m), 1.92-1.99 (2H, m), 2.05-2.15 (2H, m), 2.57 (3H, s), 4.00-4.06 (1H, m), 7.59 (1H, d), 7.79 (2H, d), 8.05 (2H, d), 8.32 (1H, s)

m/z (ESI+) (M+H)⁺ = 361;



Intermediate#67: (Z)-N-(2-adamantyl)-2-(cyclopentanecarbonyl)-3-dimethylamino-prop-2-enamide






N,N-Dimethylformamide dimethyl acetal (0.587 mL, 4.42 mmol) was added to N-(2-adamantyl)-3-cyclopentyl-3-oxo-propanamide (**Intermediate#72**), (1.023g, 3.53 mmol) in 1,4-dioxane (25 mL). The resulting solution was stirred at 100 °C for 2 hours. The resulting mixture was evaporated to dryness to afford (Z)-N-(2-adamantyl)-2-(cyclopentanecarbonyl)-3-dimethylamino-prop-2-enamide (1.210 g, 99 %).

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.45 - 1.50 (4H, m), 1.54 - 1.68 (8H, m), 1.72 - 1.85 (8H, m), 2.00 (2H, d), 2.97 (6H, s), 3.05 - 3.15 (1H, m), 3.95 (1H, d), 7.42 (1H, s), 8.30 (1H, d)

The same process as used for Intermediate#67 prepared the following intermediates from the appropriate starting material.

Structure	Int#	Name	^1H NMR δ	MS m/e MH^+
	68	(2Z)-N-(2-adamantyl)-2-(dimethylaminomethylidene)-3-oxopentanamide	^1H NMR (300.073 MHz, DMSO- d_6) δ 0.94 (3H, t), 1.48 (2H, d), 1.68 (2H, s), 1.77 - 1.81 (8H, m), 1.95 - 2.00 (2H, m), 2.38 (2H, q), 2.97 (6H, s), 3.92 (1H, d), 7.40 (1H, s), 8.25 (1H, d)	305
	69	(2Z)-N-(2-adamantyl)-2-(dimethylaminomethylidene)-4-methyl-3-oxopentanamide	^1H NMR (400.13 MHz, DMSO- d_6) δ 0.95 - 1.00 (6H, m), 1.50 (2H, d), 1.70 (2H, s), 1.76 - 1.78 (4H, m), 1.79 (2H, s), 1.82 (3H, s), 2.01 (2H, d), 2.93 - 2.99 (6H, m), 3.58 (3H, s), 3.95 (1H, d), 7.40 (1H, s), 8.24 (1H, d)	

Structure	Int#	Name	¹ H NMR δ	MS m/e MH ⁺
	70	(Z)-N-(2-adamantyl)-2-(cyclobutanecarbonyl)-3-dimethylaminoprop-2-enamide	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.51 (2H, d), 1.67 - 1.71 (2H, m), 1.78 - 1.87 (8H, m), 1.94 - 2.00 (4H, m), 2.11 - 2.21 (2H, m), 2.98 (5H, s), 3.29 (3H, s), 3.53 (1H, t), 3.93 (1H, d), 7.37 (1H, s), 8.29 (1H, d)	
	71	(2Z)-N-(2-adamantyl)-2-(dimethylaminomethylidene)-3-oxobutanamide	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.46 - 1.52 (2H, m), 1.65-1.70 (2H, m), 1.72 - 1.85 (8H, m), 1.92-2.00 (2H, m), 2.04 (3H, s), 2.99 (6H, s), 3.91-3.96 (1H, m), 7.44 (1H, s), 8.35 (1H, d)	



Intermediate#72: N-(2-adamantyl)-3-cyclopentyl-3-oxo-propanamide

2-Adamantanamine hydrochloride (1.641 g, 8.74 mmol) was added to 5-
 (cyclopentanecarbonyl)-2,2-dimethyl-1,3-dioxane-4,6-dione (**Intermediate#77**), (2.1 g,
 8.74 mmol) and N-Ethyldiisopropylamine (1.512 mL, 8.74 mmol) in toluene (30 mL). The
 resulting suspension was stirred at 110 °C for 2 hours. The reaction mixture was diluted
 with EtOAc (100 mL), and washed sequentially with 2M HCl (20 mL), and water (2x50
 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude
 product. The crude product was purified by flash silica chromatography, elution gradient 0
 to 100% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford N-(2-
 adamantyl)-3-cyclopentyl-3-oxo-propanamide (1.030 g, 40.7 %) as a brown oil which
 crystallised on standing.


¹H NMR (400.13 MHz, DMSO-d₆) δ 1.46 - 1.59 (7H, m), 1.60 - 1.78 (16H, m), 1.90 - 2.00
 (2H, m), 2.95 - 3.03 (1H, m), 3.84 (0.9H, d), 3.90 (0.1H, d), 7.78 (0.1H, d), 7.93 (0.9H, d),
 14.21 (0.1H, s) 9:1 ketone:enol form

The same process as used for Intermediate#72 prepared the following
 intermediates from the appropriate starting material.

Structure	Int#	Name	¹ H NMR δ
	73	N-(2-adamantyl)-3-oxopentanamide	¹ H NMR (400.13 MHz, CDCl ₃) δ 1.11 (3H, t), 1.67 (2H, d), 1.77 (2H, s), 1.80-1.98 (10H, d), 2.59 (2H, q), 3.46 (2H, s), 4.08 - 4.10 (1H, m), 7.70 (1H, s)
	74	N-(2-adamantyl)-4-methyl-3-oxopentanamide	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 0.98 - 1.06 (6H, m), 1.48 - 1.54 (2H, m), 1.70 - 1.73 (3H, m),

Structure	Int#	Name	¹ H NMR δ
			1.75 - 1.81 (4H, m), 1.79 (4H, d), 1.95 - 2.00 (2H, m), 2.68 - 2.75 (1H, m), 3.46 (1H, s), 3.84 (1H, d), 7.87 (1H, d)
	75	N-(2-adamantyl)-3-cyclobutyl-3-oxopropanamide	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.50 - 1.53 (3H, m), 1.65 - 1.73 (3H, m), 1.74 - 1.81 (8H, m), 1.86 - 1.98 (4H, m), 2.02 - 2.10 (4H, m), 3.29 (1H, s), 3.84 (1H, d), 7.88 (1H, d)
	76	N-(2-adamantyl)-3-oxo-butanamide	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.48 - 1.54 (2H, m), 1.69 - 1.85 (10H, m), 1.92-2.00 (2H, s), 2.13 (3H, s), 3.38 (2H, s), 3.84 (1H, d), 7.95 (1H, d)

Intermediate#77: 5-(cyclopentanecarbonyl)-2,2-dimethyl-1,3-dioxane-4,6-dione







A solution of cyclopentanecarbonyl chloride (1.100 mL, 9.05 mmol) in DCM (5mL) was added dropwise to a stirred solution of isopropylidene malonate (1.304 g, 9.05 mmol), and pyridine (1.464 mL, 18.10 mmol) in DCM (20mL) at 5°C, over a period of 10 minutes under nitrogen. The resulting solution was stirred at 5 °C for 1 hour and then allowed to warm to 20°C and stirred for another hour. The reaction mixture was diluted


with DCM (100 mL), and washed sequentially with 2M HCl (2x50 mL), water (50 mL), and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product, 5-(cyclopentanecarbonyl)-2,2-dimethyl-1,3-dioxane-4,6-dione (2.100 g, 97 %) as a brown oil.

- 5 ¹H NMR (400.13 MHz, DMSO-d₆) δ 1.51 - 1.60 (5H, m), 1.61 - 1.68 (3H, s), 1.70 - 1.79 (4H, m), 1.84 - 1.98 (2H, m), 2.89 - 3.00 (1H, m), 4.04 (1H, s)

The same process as used for Intermediate#77 prepared the following intermediates from the appropriate starting materials.

Structure	Int#	Name	¹ H NMR δ
	78	2,2-dimethyl-5-propanoyl-1,3-dioxane-4,6-dione	Used crude
	79	2,2-dimethyl-5-(2-methylpropanoyl)-1,3-dioxane-4,6-dione	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.02 - 1.08 (1H, m), 1.15 - 1.18 (6H, m), 1.70 - 1.72 (6H, m), 3.92 - 3.99 (1H, m)
	80	5-(cyclobutanecarbonyl)-2,2-dimethyl-1,3-dioxane-4,6-dione	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.68 (6H, s), 1.82 - 1.89 (2H, m), 2.09 (2H, s), 2.05 - 2.13 (2H, m), 2.18 - 2.32 (5H, m), 4.30 - 4.35 (1H, m)
	81	5-acetyl-2,2-dimethyl-1,3-dioxane-4,6-dione	Used crude

Intermediate# 82 : 1-(4-bromophenyl)-5-tert-butyl-N-cyclohexyl-1H-pyrazole-4-carboxamide




Ethyl (2)-2-(dimethylaminomethylidene)-4,4-dimethyl-3-oxo-pentanoate

- 5 **(Intermediate # 83)** (1.24g, 5.88mmol) was dissolved in ethanol (20mL). 4-bromophenylhydrazine HCl (1.32g, 5.88mmol) and DIPEA (1.02mL, 5.88mmol) were added. The mixture was heated to reflux for 2h cooled to ambient and evaporated under reduced pressure. The residue was dissolved in DCM (50mL), washed with water (2x10mL), dried (MgSO₄) and evaporated to afford crude product. The crude product was purified by
- 10 flash silica chromatography, elution gradient 0 to 25% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford 1-(4-bromophenyl)-5-tert-butyl-N-cyclohexyl-1H-pyrazole-4-carboxamide (627mg, 32%)

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.23 (9H, s), 1.29 (3H, t), 4.24 (2H, q), 7.39 - 7.41 (2H, m), 7.69 - 7.72 (2H, m), 7.92 (1H, s)

- 15 m/z (ESI+) (M+H)⁺ = 353

Intermediate # 83 : Ethyl (2)-2-(dimethylaminomethylidene)-4,4-dimethyl-3-oxo-pentanoate




- 20 Ethyl pivaloylacetate (1.72g, 10mmol) and N,N-dimethylformamide dimethylacetal (1.68mL, 12.5mmol) were dissolved in dioxan (20mL) and heated to reflux for 3h. The reaction mixture was cooled to ambient and evaporated under reduced pressure to leave

crude product. The crude product was purified by flash chromatography on silica gel (elution gradient 0 to 100% EtOAc in hexane) to afford Ethyl (2)-(dimethylaminomethylidene)-4,4-dimethyl-3-oxo-pentanoate as a colourless oil that crystallised on standing. (1.24g, 54%)

5 ¹H NMR (300.073 MHz, DMSO-d₆) δ 1.11 (9H, d), 1.18 (3H, t), 2.82 (6H, s), 4.04 (2H, q), 7.31 (1H, s)

m/z (ESI+) (M+H)⁺ = 228

Intermediate# 84 : methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoate



10

1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (191 mg, 1.00 mmol) was added in one portion to 5-(cyclohexylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (**Intermediate# 88**) (300 mg, 0.83 mmol), 2-Adamantanamine hydrochloride (172 mg, 0.92 mmol) and N-Ethyldiisopropylamine
15 (0.432 mL, 2.50 mmol) in DMF (7 mL) at 20°C under nitrogen. The resulting mixture was stirred at 20 °C for 18 hours.

The reaction mixture was diluted with EtOAc (75 mL), and washed sequentially with water (4x25 mL) and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product. The crude product was purified
20 by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoate (349 mg, 85 %) as a white crystalline solid.


¹H NMR (300.073 MHz, DMSO-d₆) δ 1.05 (5H, d), 1.39 - 1.54 (5H, m), 1.62 (2H, d), 1.73 (2H, s), 1.83 (6H, s), 1.94 - 2.01 (4H, m), 2.92 (1H, s), 3.89 (3H, s), 4.09 (1H, d), 7.76 (2H, d), 8.04 (1H, d), 8.12 (2H, d), 8.17 (1H, s)
m/z (ESI+) (M+H)⁺ = 494

- 5 The following Intermediates were prepared in a similar manner to Intermediate #84 from 5-(cyclohexylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (**Intermediate#88**) and an appropriate amine.

Structure	Int.#	Name	¹ H NMR δ	MS m/e MH ⁺
	85	methyl 4-[4-(1-adamantylcarbamoyl)-5-cyclohexylsulfanylpyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.05 - 1.17 (5H, m), 1.42 (1H, s), 1.51 - 1.63 (4H, m), 1.68 (6H, s), 2.02 - 2.12 (9H, m), 2.98 (1H, s), 3.91 (3H, s), 7.53 (1H, s), 7.72 - 7.74 (2H, m), 8.13 (3H, d)	(ESI+) 494
	86	methyl 4-[5-cyclohexylsulfanyl-4-[(1S,3R)-5-hydroxy-2-adamantyl]carbamoylpyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.02 - 1.10 (5H, m), 1.41 - 1.56 (7H, m), 1.67 (4H, d), 1.76 (2H, d), 1.89 (2H, d), 2.08 (3H, s), 2.93 (1H, s), 3.91 (3H, s), 4.03 (1H, d), 4.43 (1H, s), 7.76 - 7.79 (2H, m), 7.97 (1H, d), 8.13 - 8.15 (2H, m), 8.18 (1H, s)	(ESI+) 510

Structure	Int.#	Name	¹ H NMR δ	MS m/e MH ⁺
	87	methyl 4-[[5-(cyclohexylsulfanyl)-4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.00 - 1.10 (5H, m), 1.41 (1H, s), 1.45-1.58 (6H, m), 1.91 - 1.99 (6H, m), 2.05 (2H, d), 2.18 (3H, d), 2.95 (1H, d), 3.91 (3H, s), 4.10 (1H, t), 6.88 (1H, t), 7.75 - 7.79 (2H, m), 7.99 (1H, d), 8.12 - 8.16 (2H, m), 8.19 (1H, s)	(ESI+) 560

Intermediate #88 : 5-cyclohexylsulfanyl-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylic acid




Trifluoroacetic acid (4.72 mL, 61.46 mmol) was added to tert-butyl 5-(cyclohexylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (Intermediate# 89) (2.56 g, 6.15 mmol) in CH₂Cl₂ (40mL). The resulting solution was stirred at 20 °C for 24 hours. The reaction mixture was evaporated to dryness, re-dissolved in dioxan (20 mL) and re-evaporated to dryness to afford 5-(cyclohexylthio)-1-(4-

(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (2.25 g, 100 %) as a white crystalline solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.04 - 1.11 (5H, m), 1.42 (1H, s), 1.51 (2H, s), 1.59 (2H, d), 3.31-3.22 (1H, m), 3.91 (3H, s), 7.70 - 7.73 (2H, m), 8.11 - 8.15 (2H, m), 8.19 (1H, s), 12.66 (1H, s)

MS m/z (ESI+) (M+H)⁺ = 361.

Intermediate # 89 : tert-butyl 5-cyclohexylsulfanyl-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylate



Sodium bis(trimethylsilyl)amide (10.69 mL, 10.69 mmol) was added dropwise to Cyclohexanethiol (1.307 mL, 10.69 mmol) in DMF (35mL) under nitrogen. The resulting solution was stirred at 20 °C for 30 minutes. A solution of tert-butyl 5-chloro-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (**Intermediate# 9**) (3 g, 8.91 mmol) in DMF(10mL) was then added dropwise and the resulting mixture was stirred at 20 °C for 2 hours.


The reaction mixture was diluted with EtOAc (200 mL), and washed sequentially with water (4x50 mL) and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 25% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford tert-butyl 5-(cyclohexylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (2.56 g, 69.0 %) as a colourless oil which crystallised on standing.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.95-1.05 (5H, m), 1.35 (1H, s), 1.42-1.55 (13H, d), 3.12 (1H, d), 3.81 (3H, s), 7.69 - 7.72 (2H, m), 8.11 - 8.14 (2H, m), 8.16 (1H, s)

m/z (ESI+) (M+H)⁺ = 417

Intermediate # 90 : methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate





1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (199 mg, 1.04 mmol) was added in one portion to 5-(cyclopentylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (**Intermediate# 93**) (300 mg, 0.87 mmol), 2-Adamantanamine hydrochloride (179 mg, 0.95 mmol), 1-Hydroxybenzotriazole (140 mg, 1.04 mmol) and N-Ethyldiisopropylamine (0.450 mL, 2.60 mmol) in DMF (7 mL) at 20°C under nitrogen. The resulting mixture was stirred at 20 °C for 18 hours. The reaction mixture was diluted with EtOAc (75 mL), and washed sequentially with water (4x25 mL) and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.


The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(2-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate (336 mg, 81 %) as a white crystalline solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.17 - 1.25 (2H, m), 1.35 - 1.42 (4H, m), 1.65 (4H, d), 1.74 (2H, s), 1.87 (6H, d), 1.96 - 2.02 (4H, m), 3.25 - 3.35 (1H, m), 3.91 (3H, s), 4.12 (1H, t), 7.78 - 7.82 (2H, m), 8.08 (1H, d), 8.13 - 8.16 (2H, m), 8.19 (1H, s)
m/z (ESI+) (M+H)⁺ = 480

The following Intermediates were prepared in a similar manner to Intermediate #90 from 5-(cyclopentylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (**Intermediate# 93**) and an appropriate amine.

Structure	Int#	Name	¹ H NMR δ	MS m/e (M+H) ⁺
	91	methyl 4-[4-(1-adamantylcarbamoyl)-5-cyclopentylsulfanylpirazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.19 - 1.27 (2H, m), 1.40 - 1.48 (4H, m), 1.63 - 1.71 (8H, m), 2.07 (9H, s), 3.31 - 3.37 (1H, m), 3.91 (3H, s), 7.54 (1H, s), 7.74 - 7.77 (2H, m), 8.12 - 8.15 (3H, m)	480
	92	methyl 4-[5-cyclopentylsulfany-4-[(1S,3R)-5-hydroxy-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoate	¹ H NMR (400.13 MHz, DMSO-d ₆) δ 1.17 - 1.24 (2H, m), 1.32-1.48 (6H, m), 1.64 (6H, d), 1.76 (2H, d), 1.89 (2H, d), 2.08 (3H, s), 3.25-3.35 (1H, m), 3.91 (3H, s), 4.03 (1H, t), 4.41 (1H, s), 7.77 - 7.81 (2H, m), 7.98 (1H, d), 8.13 - 8.17 (2H, m), 8.19 (1H, s)	496

Intermediate # 93 : 5-cyclopentylsulfanyl-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylic acid



Trifluoroacetic acid (2.63 mL, 34.29 mmol) was added to tert-butyl 5-(cyclopentylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate
 5 **(Intermediate# 94)** (1.38 g, 3.43 mmol) in CH₂Cl₂ (25 mL) The resulting solution was stirred at 20 °C for 24 hours. The reaction mixture was evaporated to dryness, re-dissolved in dioxan (20 mL) and re-evaporated to dryness to afford 5-(cyclopentylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (1.180 g, 99 %) as a white
 10 crystalline solid.


¹H NMR (400.13 MHz, DMSO-d₆) δ 1.20 - 1.26 (2H, m), 1.37 - 1.47 (4H, m), 1.65 - 1.73 (2H, m), 3.65 - 3.71 (1H, m), 3.86 - 3.96 (3H, m), 7.68 - 7.76 (2H, m), 8.11 - 8.15 (2H, m), 8.19 (1H, s), 12.84 (1H, s)

MS m/z (ESI-) (M-H)⁻ = 345

15

20

Intermediate# 94 : tert-butyl 5-cyclopentylsulfanyl-1-(4-methoxycarbonylphenyl)pyrazole-4-carboxylate




Sodium bis(trimethylsilyl)amide (7.13 mL, 7.13 mmol) was added dropwise to
5 cyclopentanethiol (0.761 mL, 7.13 mmol) in DMF (25mL) under nitrogen. The resulting
solution was stirred at 20 °C for 30 minutes. A solution of tert-butyl 5-chloro-1-(4-
(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (**Intermediate#9**) (2 g, 5.94 mmol)
in DMF(10mL) was then added dropwise and the resulting mixture was stirred at 20 °C for
2 hours. The reaction mixture was diluted with EtOAc (200 mL), and washed sequentially
10 with water (4x50 mL) and saturated brine (50 mL). The organic layer was dried over
MgSO₄, filtered and evaporated to afford crude product. The crude product was purified
by flash silica chromatography, elution gradient 0 to 25% EtOAc in isohexane. Pure
fractions were evaporated to dryness to afford tert-butyl 5-(cyclopentylthio)-1-(4-
(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (1.380 g, 57.7 %) as a colourless oil
15 which crystallised on standing.

¹H NMR (300.073 MHz, DMSO-d₆) δ 1.14 - 1.24 (2H, m), 1.35-1.47 (4H, m), 1.54 (9H, s), 1.60-1.77 (2H, m), 3.54 - 3.58 (1H, m), 3.90 (3H, s), 7.71 (2H, d), 8.11 (2H, d), 8.14 (1H, d)

MS m/z (ESI⁺) (M+H)⁺ = 403.

Intermediate# 95 : Methyl 4-[4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanylpirazol-1-yl]benzoate



1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (250 mg, 1.31 mmol) was added in one portion to 1-(4-methoxycarbonylphenyl)-5-propylsulfanylpirazole-4-carboxylic acid (**Intermediate#7**) (349mg, 1.09mmol), 5-(difluoromethoxy)adamantan-2-amine (**Intermediate#121**) (260 mg, 1.20 mmol), 1-Hydroxybenzotriazole (176 mg, 1.31 mmol) and N-Ethyldiisopropylamine (0.376 mL, 2.18 mmol) in DMF (10 mL). The resulting mixture was stirred at 20 °C for 5 hours.


The reaction mixture was diluted with EtOAc (100 mL), and washed sequentially with water (4x25 mL) and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanyl-pyrazol-1-yl]benzoate (338 mg, 59.8 %) as a white crystalline solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.68 (3H, t), 1.21 - 1.30 (2H, m), 1.51 (2H, d), 1.90 - 1.97 (6H, m), 2.05 (2H, d), 2.15 - 2.20 (3H, m), 2.65 (2H, t), 3.91 (3H, s), 4.09 (1H, d), 6.88 (1H, t), 7.76 - 7.79 (2H, m), 7.98 (1H, d), 8.13 - 8.16 (2H, m), 8.19 (1H, s)

m/z (ESI+) (M+H)⁺ = 520

Intermediate#96: methyl 4-[4-(cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate




To a solution of cyclopentyl mercaptan (0.071 ml, 0.66 mmol) in DMF (2ml) was added a 1N solution of NaHMDS in THF (0.66 ml, 0.66 mmol). The reaction was stirred at ambient temperature for 2 minutes then added to a solution of methyl 4-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate#15**) (200mg , 0.55 mmol) in DMF (3ml).

The reaction mixture was stirred at room temperature for two hours. The reaction mixture was evaporated to dryness and redissolved in DCM (50 mL) and washed with saturated NH₄Cl (10 mL), water (10 mL) and brine (10 mL). It was dried over MgSO₄ and the solvent was evaporated under reduced pressure to give a solid. The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate (233 mg, 98%) as a solid.

m/z (ESI+) (M+H)⁺ = 428


Intermediate# 97: methyl 4-[4-(cyclohexylcarbamoyl)-5-cyclohexylsulfanylpyrazol-1-yl]benzoate



Methyl 4-[4-(cyclohexylcarbamoyl)-5-cyclohexylsulfanylpyrazol-1-yl]benzoate
5 was prepared from cyclohexylthiol and methyl 4-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate#15**) by the same process used for Intermediate#96.

m/z (ESI+) (M+H)⁺ = 442

Intermediate# 98: methyl 4-[5-cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate




Methyl 4-[5-cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate
was prepared from cycloheptylthiol and methyl 4-[5-chloro-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoate (**Intermediate#15**) by the same process used
15 for Intermediate#96.

m/z (ESI+) (M+H)⁺ = 456; HPLC t_R = 3.27 min.

¹H NMR (300.072 MHz, CDCl₃) δ 1.08 - 1.48 (15H, m), 1.56 - 1.72 (5H, m), 1.91 - 1.97 (2H, m), 2.84 - 2.93 (1H, m), 3.90 (3H, s), 3.93 - 4.02 (1H, m), 7.57 - 7.65 (3H, m), 8.08 - 8.12 (2H, d), 8.22 (1H, s)

5 **Intermediate# 99 : methyl 4-[4-(2-adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoate**



N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (396 mg, 2.07 mmol) was added in one portion to 2-Adamantanamine hydrochloride (324 mg, 1.72 mmol), 5-(ethylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (528 mg, 1.72 mmol) (**Intermediate#101**), 1-Hydroxybenzotriazole (279 mg, 2.07 mmol) and N-Ethyldiisopropylamine (0.885 mL, 5.17 mmol) in DMF (10 mL). The resulting mixture was stirred at 20 °C for 18 hours.


15 The reaction mixture was diluted with EtOAc (100 mL), and washed sequentially with water (4x25 mL) and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(2-adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoate (430 mg, 56.8 %) as a white solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.94 (3H, t), 1.62 (2H, d), 1.74 (2H, s), 1.86 (6H, d), 1.96 - 2.02 (4H, m), 2.68 (2H, q), 3.91 (3H, s), 4.11 (1H, t), 7.76 - 7.80 (2H, m), 8.04 (1H, d), 8.13 - 8.16 (2H, m), 8.20 (1H, s)

m/z (ESI+) (M+H)⁺ = 440

Intermediate# 100 : methyl 4-[4-(2-adamantylcarbamoyl)-5-methylsulfanyl-pyrazol-1-yl]benzoate




Prepared from methyl 4-[4-(2-adamantylcarbamoyl)-5-methylsulfanyl-pyrazol-1-yl]benzoate (**Intermediate# 102**) by the same process as Intermediate# 101.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.61 (2H, d), 1.74 (2H, s), 1.86 (6H, s), 1.95 - 2.04 (4H, m), 2.29 (3H, s), 3.91 (3H, s), 4.11 (1H, t), 7.76 - 7.79 (2H, m), 8.02 (1H, d), 8.13 - 8.16 (2H, m), 8.19 (1H, s)

m/z (ESI+) (M+H)⁺ = 426

Intermediate # 101 : 5-(ethylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid




Trifluoroacetic acid (1.324 mL, 17.24 mmol) was added to tert-butyl 5-(ethylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (**Intermediate# 103**) (625 mg, 1.72 mmol) in CH₂Cl₂ (25mL) The resulting solution was stirred at 20 °C for 24 hours.

The reaction mixture was evaporated to dryness and redissolved in dioxan (20 mL), and re- evaporated to dryness to afford 5-(ethylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid (528 mg, 100 %) as a white solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 0.96 (3H, t), 2.88 (2H, q), 3.91 (3H, s), 7.70 - 7.74 (2H, m), 8.12 - 8.15 (2H, m), 8.19 (1H, s), 12.72 (1H, s)

m/z (ESI+) (M+H)⁺ = 307

Intermediate # 102 : 5-(methylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylic acid




Prepared from tert-butyl 1-(4-(methoxycarbonyl)phenyl)-5-(methylthio)-1H-pyrazole-4-carboxylate (**Intermediate# 104**) by the same process as Intermediate# 101.

¹H NMR (400.13 MHz, DMSO-d₆) δ 2.40 (3H, s), 3.90 (3H, d), 7.72 - 7.75 (2H, m), 8.10 - 8.14 (2H, m), 8.17 (1H, s), 12.77 (1H, s)

m/z (ESI+) (M+H)⁺ = 293

Intermediate# 103 : tert-butyl 5-(ethylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate




Sodium bis(trimethylsilyl)amide (3.56 mL, 3.56 mmol) was added dropwise to
5 ethanethiol (0.264 mL, 3.56 mmol) in DMF (10mL) under nitrogen. The resulting solution
was stirred at 20 °C for 30 minutes. tert-butyl 5-chloro-1-(4-(methoxycarbonyl)phenyl)-
1H-pyrazole-4-carboxylate (**Intermediate#9**) (1 g, 2.97 mmol) was added in one portion
and the resulting suspension was stirred at 20 °C for 5 hours. The reaction mixture was
diluted with EtOAc (100 mL), and washed sequentially with water (4x25 mL), and
10 saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated
to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0
to 30% EtOAc in isohexane. Product containing fractions were evaporated to dryness to
afford tert-butyl 5-(ethylthio)-1-(4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate
15 (0.625 g, 58.1 %) as a white crystalline solid.

¹H NMR (400.13 MHz, DMSO-*d*₆) δ 0.96 (3H, t), 1.55 - 1.56 (9H, m), 2.84 (2H, q), 3.91
(3H, s), 7.70 - 7.73 (2H, m), 8.10 - 8.15 (3H, m)

m/z (ESI⁺) (M+H)⁺ = 363

Intermediate# 104 : tert-butyl 1-(4-(methoxycarbonyl)phenyl)-5-(methylthio)-1H-pyrazole-4-carboxylate




Prepared from (**Intermediate# 9**) and sodium methanethiolate by a similar process
 5 to that used for (**Intermediate# 103**).

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.56 (9H, s), 2.38 (3H, s), 3.91 (3H, s), 7.72 - 7.74
 (2H, m), 8.10 - 8.17 (3H, m)

m/z (ESI+) (M+H)⁺ = 349

Intermediate# 105: 4-[4-(5-Methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid methyl ester



1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (127 mg, 0.66
 mmol) was added to 1-(4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-
 carboxylic acid (**Intermediate#7**) (152 mg, 0.47 mmol) 5-Methanesulfonyl-adamantan-2-
 15 ylamine (*Prepared by the method described in Bioorganic & Medicinal Chemistry Letters*
17 (2007) 527-532) (109 mg, 0.47 mmol), 4-Dimethylaminopyridine (11.59 mg, 0.09
 mmol) and Triethylamine (0.132 mL, 0.95 mmol) in DCM (7 mL) at ambient temperature
 under nitrogen. The resulting solution was stirred at ambient temperature for 20 hours.


The reaction mixture was evaporated to dryness and redissolved in EtOAc (50 mL),

and washed sequentially with 1N citric acid (25 mL), water (20 mL), and saturated brine (20 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product. The crude product was purified by flash silica chromatography, elution gradient 20 to 100% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford 4-[4-(5-Methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-benzoic acid methyl ester (116 mg, 46.0 %) as a white solid.

¹H NMR (400.13 MHz, CDCl₃) δ 0.76 (3H, t), 1.31 - 1.40 (2H, m), 1.71 (2H, d), 1.96 (2H, d), 2.11 (2H, s), 2.15 - 2.30 (5H, m), 2.33 (2H, s), 2.54 (2H, t), 2.78 (3H, s), 3.97 (3H, s), 4.30 - 4.40 (1H, m), 7.71 (2H, d), 8.07 (1H, d), 8.20 (2H, d), 8.30 (1H, s)

MS m/e MH⁺ 532.

Intermediate# 106: methyl 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoate




N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (197 mg, 1.03 mmol) was added in one portion to 2-adamantanamine hydrochloride (161 mg, 0.86 mmol), 1-(3-methoxy-4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-carboxylic acid (**Intermediate# 107**) (300 mg, 0.86 mmol) 1-Hydroxybenzotriazole (139 mg, 1.03 mmol) and N-Ethyldiisopropylamine (0.440 mL, 2.57 mmol) in DMF (10 mL). The resulting mixture was stirred at 20 °C for 18 hours.

The reaction mixture was diluted with EtOAc (100 mL), and washed sequentially with water (4x25 mL) and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford methyl 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoate (243 mg, 58.7 %) as colourless oil.

- 5 ¹H NMR (400.13 MHz, DMSO-d₆) δ 0.69 (3H, t), 1.24 - 1.32 (2H, m), 1.62 (2H, d), 1.74 (2H, s), 1.86 (6H, d), 1.95 - 2.02 (4H, m), 2.65 (2H, t), 3.83 (3H, s), 3.88 (3H, s), 4.11 (1H, d), 7.26 - 7.29 (1H, m), 7.44 (1H, d), 7.83 (1H, d), 8.09 (1H, d), 8.18 (1H, s)
m/z (ESI+) (M+H)⁺ = 484

10 **Intermediate# 107: 1-(3-methoxy-4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-carboxylic acid**




- Trifluoroacetic acid (1.194 mL, 15.55 mmol) was added to tert-butyl 1-(3-methoxy-4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-carboxylate (**Intermediate# 108**) (632 mg, 1.55 mmol) in CH₂Cl₂ (15mL) The resulting solution was stirred at 20 °C
15 for 24 hours.

The reaction mixture was evaporated to dryness and re-dissolved in dioxan (20 mL), and re-evaporated to dryness to afford 1-(3-methoxy-4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-carboxylic acid (540 mg, 99 %) as a colourless oil which solidified on standing.

- 20 ¹H NMR (400.13 MHz, DMSO-d₆) δ 0.72 (3H, t), 1.25 - 1.34 (2H, m), 2.85 (2H, t), 3.57 (1H, s), 3.83 (3H, s), 3.87 (3H, s), 7.19 - 7.22 (1H, m), 7.36 - 7.36 (1H, m), 7.82 (1H, d), 8.18 (1H, s), 12.6 (1H, s)
m/z (ESI+) (M+H)⁺ = 351

Intermediate# 108: tert-butyl 1-(3-methoxy-4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-carboxylate




Sodium bis(trimethylsilyl)amide 1M solution in THF (3.60 mL, 3.60 mmol) was added dropwise to 1-propanethiol (0.326 mL, 3.60 mmol) in DMF (10mL) under nitrogen. The resulting solution was stirred at 20 °C for 30 minutes. tert-butyl 5-chloro-1-(3-methoxy-4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (**Intermediate# 109**) (1.1 g, 3.00 mmol) was added as a solution in DMF (5mL) and the resulting mixture was stirred at 20 °C for 5 hours. The reaction mixture was diluted with EtOAc (100 mL), and washed sequentially with water (4x25 mL), and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Product containing fractions were evaporated to dryness to afford tert-butyl 1-(3-methoxy-4-(methoxycarbonyl)phenyl)-5-(propylthio)-1H-pyrazole-4-carboxylate (0.642 g, 52.7 %) as a colourless oil.

¹H NMR (400.13 MHz, DMSO-*d*₆) δ 0.72 (3H, t), 1.29 (2H, q), 1.55 (9H, s), 2.79 (2H, t), 3.83 (3H, s), 3.87 (3H, s), 7.19 - 7.22 (1H, m), 7.36 (1H, d), 7.82 (1H, d), 8.14 (1H, s)
m/z (ESI+) (M+H)⁺ = 407

Intermediate# 109: tert-butyl 5-chloro-1-(3-methoxy-4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate




Copper(II) chloride (0.917 g, 6.82 mmol) was added in one portion to tert-Butyl
5 nitrite (0.649 mL, 5.46 mmol) in acetonitrile (25 mL) at and warmed to 50°C . The
resulting mixture was stirred at 50 °C while tert-butyl 5-amino-1-(3-methoxy-4-
(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (**Intermediate# 110**) (1.58 g, 4.55
mmol) was added in portions as a solid. After the addition was complete the reaction
mixture was stirred at 50 °C for 15 minutes and then cooled to 20°C. The reaction mixture
was diluted with EtOAc (100 mL), and washed sequentially with water (2 x 50 mL), and
saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated
to afford crude product. The crude product was purified by flash silica chromatography,
elution gradient 0 to 50% EtOAc in isohexane. Fractions were evaporated to dryness to
afford tert-butyl 5-chloro-1-(3-methoxy-4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-
carboxylate (1.120 g, 67.1 %) as a yellow oil.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.55 (9H, s), 3.83 (3H, s), 3.88 (3H, s), 7.26 - 7.29
(1H, m), 7.42 (1H, d), 7.84 (1H, d), 8.20 (1H, s)

m/z (ESI+) (M+H)⁺ = 367


Intermediate# 110 :tert-butyl 5-amino-1-(3-methoxy-4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate



Acetic anhydride (2.83 mL, 30.00 mmol) was added in one portion to tert-Butyl
5 cyanoacetate (4.29 mL, 30 mmol), and triethyl orthoformate (7.48 mL, 45.00 mmol). The
resulting mixture was stirred at 125°C for 3 hours and then volatiles were removed by
evaporation under reduced pressure. The resulting oil was dissolved in a ethanol (50 mL),
treated with methyl 4-hydrazinyl-2-methoxybenzoate hydrochloride (**Intermediate# 111**)
(2.094 g, 9.00 mmol) and N-Ethyldiisopropylamine (1.572 mL, 9.00 mmol) and stirred at
80°C for 4h.

The reaction mixture was concentrated and diluted with EtOAc (200 mL), and
washed sequentially with water (2x100 mL), and saturated brine (50 mL). The organic
layer was dried over MgSO₄, filtered and evaporated to afford crude product.
The crude product was purified by flash silica chromatography, elution gradient 0 to 50%
EtOAc in isohexane. Pure fractions were evaporated to dryness to afford tert-butyl 5-
amino-1-(3-methoxy-4-(methoxycarbonyl)phenyl)-1H-pyrazole-4-carboxylate (1.640 g, 52
%) as an orange solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.52 (9H, s), 3.81 (3H, s), 3.89 (3H, s), 6.47 (2H, s),
7.21 - 7.23 (1H, m), 7.30 (1H, d), 7.68 (1H, s), 7.81 (1H, d)
m/z (ESI+) (M+H)⁺ = 348

Intermediate# 111: methyl 4-hydrazinyl-2-methoxybenzoate hydrochloride


tert-butyl 1-(3-methoxy-4-(methoxycarbonyl)phenyl)hydrazinecarboxylate

(Intermediate# 112) (4.86g, 16.40 mmol) was added to a 4M solution of hydrogen

chloride (61.5 mL, 246 mmol) in dioxane. The resulting solution was stirred at 20 °C for 5 hours. The reaction mixture was evaporated to dryness and the crude residue was triturated with Et2O to give a solid which was collected by filtration and dried under vacuum to give methyl 4-hydrazinyl-2-methoxybenzoate hydrochloride (3.50 g, 92 %) as a pale green solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 3.73 (3H, s), 3.81 (3H, s), 6.53 - 6.56 (1H, m), 6.79 (1H, d), 7.66 (1H, d), 8.79 (1H, s), 10.44 (3H, s)

m/z (ESI+) (M+H)⁺ = 197

Intermediate# 112: tert-butyl 1-(3-methoxy-4-(methoxycarbonyl)phenyl)hydrazinecarboxylate

15

Copper(I) iodide (0.213 g, 1.12 mmol) was added to 1,10-Phenanthroline (0.403 g, 2.24 mmol), tert-Butyl carbazate (3.55 g, 26.83 mmol), cesium carbonate (10.20 g, 31.30 mmol) and methyl 4-iodo-2-methoxybenzoate (6.53 g, 22.36 mmol) in DMF (75 mL) under nitrogen. The resulting mixture was stirred at 100 °C for 1 hour. The reaction mixture was diluted with EtOAc (400 mL), and washed sequentially with water (4x100

20


mL), and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product.

The crude product was purified by flash silica chromatography, elution gradient 0 to 50% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford tert-butyl 1-(3-methoxy-4-(methoxycarbonyl)phenyl)hydrazinecarboxylate (4.86 g, 73.4 %) as a solid.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.50 (9H, s), 3.76 (3H, s), 3.80 (3H, s), 5.11 (2H, s), 7.25 - 7.28 (1H, m), 7.39 (1H, d), 7.64 (1H, d)

m/z (ESI+) (M+H)⁺ = 297 .

Intermediate# 113: Ethyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoate




N-(2-adamantyl)-1-(4-chloro-2-methyl-phenyl)-5-tert-butyl-pyrazole-4-carboxamide (**Intermediate#114**) (647 mg, 1.52 mmol), Molybdenum hexacarbonyl (0.102 mL, 0.76 mmol), 4-Dimethylaminopyridine (371 mg, 3.04 mmol), N-Ethyldiisopropylamine (0.529 mL, 3.04 mmol), trans-Di-μ-acetatobis[2-(di-o-tolylphosphino)benzyl]dipalladium(II) (71.4 mg, 0.08 mmol) and Tri-tert-butylphosphine tetrafluoroborate (88 mg, 0.30 mmol) were suspended in ethanol (6 mL) and dioxane (6.00 mL) and sealed into a microwave tube. The reaction was heated to 150 °C for 1 hour in the microwave reactor and cooled to RT. The reaction mixture was evaporated to dryness and redissolved in EtOAc (100 mL), and washed with water (10 mL), filtered, then washed with 2M HCl (10 mL), and saturated brine (10 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product. The crude product was purified by flash silica chromatography, elution gradient 0 to 25% EtOAc in isohexane.

Pure fractions were evaporated to dryness to afford ethyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoate (352 mg, 50.0 %) as a colourless oil.

m/z (ESI+) (M+H)⁺ = 3.22; HPLC t_R = 464 min.

¹H NMR (300.072 MHz, cdcl₃) δ 1.25 (s, 9H), 1.42 (t, 3H), 1.66 - 1.95 (m, 12H), 2.01 - 2.10 (m, 2H), 2.13 (s, 3H), 4.22 (d, 1H), 4.41 (q, 2H), 6.14 (d, 1H), 7.31 (d, 1H), 7.68 (s, 1H), 7.94 (d, 1H), 8.01 (s, 1H)


Intermediate# 114: N-(2-adamantyl)-1-(4-chloro-2-methyl-phenyl)-5-tert-butyl-pyrazole-4-carboxamide



10 N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride
(**Intermediate#115**) (515 mg, 2.68 mmol) was added in one portion to 2-Adamantanamine hydrochloride (420 mg, 2.24 mmol), 5-tert-butyl-1-(4-chloro-2-methylphenyl)-1H-pyrazole-4-carboxylic acid (655 mg, 2.24 mmol), 1-Hydroxybenzotriazole (363 mg, 2.68 mmol) and N-Ethyldiisopropylamine (1.149 mL, 6.71 mmol) in DMF (10 mL). The
15 resulting mixture was stirred at 20 °C for 18 hours. The reaction mixture was diluted with Et₂O (100 mL), and washed sequentially with water (3x25 mL), and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford desired product N-(2-adamantyl)-1-(4-chloro-2-methyl-phenyl)-5-tert-butyl-pyrazole-4-carboxamide (860 mg, 90 %).

20 ¹H NMR (300.072 MHz, cdcl₃) δ 1.26 (s, 9H), 1.51 - 2.18 (m, 17H), 4.22 (d, 1H), 6.12 (d, 1H), 7.17 (d, 1H), 7.25 (d, 1H), 7.31 (s, 1H), 7.65 (s, 1H)

Intermediate# 115: 5-tert-butyl-1-(4-chloro-2-methylphenyl)-1H-pyrazole-4-carboxylic acid




5-tert-butyl-1-(4-chloro-2-methylphenyl)-1H-pyrazole-4-carboxylic acid was prepared from ethyl 5-tert-butyl-1-(4-chloro-2-methylphenyl)-1H-pyrazole-4-carboxylate (**Intermediate#116**) by the same process used for Intermediate#119.

^1H NMR (300.072 MHz, cdcl_3) δ 1.30 (s, 9H), 2.04 (s, 3H), 7.15 (d, 1H), 7.26 (d, 1H), 7.31 (s, 1H), 8.16 (s, 1H)

m/z (ESI+) (M+H) $^+$ = 293

Intermediate# 116: Ethyl 5-tert-butyl-1-(4-chloro-2-methylphenyl)-1H-pyrazole-4-carboxylate




N-Ethyldiisopropylamine (1.553 mL, 8.97 mmol) was added to (4-chloro-2-methylphenyl)hydrazine hydrochloride (1.733 g, 8.97 mmol) and (Z)-ethyl 2-((dimethylamino)methylene)-4,4-dimethyl-3-oxopentanoate (**Intermediate#83**) (2.04 g, 8.97 mmol) in ethanol (30 mL). The resulting solution was stirred at 90 °C for 2 hours. The reaction mixture was evaporated to dryness and redissolved in DCM (50 mL), and washed sequentially with water (2x10mL). The organic layer was dried over MgSO_4 , filtered and evaporated to afford crude product. The crude product was purified by flash silica chromatography, elution gradient 0 to 10% EtOAc in isohexane. Pure fractions were

evaporated to dryness to afford ethyl 5-tert-butyl-1-(4-chloro-2-methylphenyl)-1H-pyrazole-4-carboxylate (0.891 g, 30.9 %) as a orange oil.

^1H NMR (300.072 MHz, cdcl_3) δ 1.29 (s, 9H), 1.38 (t, 3H), 2.03 (s, 3H), 4.31 (q, 2H), 7.14 (d, 1H), 7.23 (d, 1H), 7.29 (d, 1H), 8.04 (s, 1H)

5 m/z (ESI+) (M+H) $^+$ = 321

Intermediate# 117: Ethyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-2-(trifluoromethyl)benzoate



Ethyl 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-2-


10 (trifluoromethyl)benzoate was prepared from N-(2-adamantyl)-1-[4-chloro-3-(trifluoromethyl)phenyl]-5-tert-butyl-pyrazole-4-carboxamide (**Intermediate#118**) by the same process used for Intermediate#113.

^1H NMR (300.072 MHz, CDCl_3) δ 1.27 - 1.33 (9H, m), 1.42 (3H, t), 1.74 - 1.79 (5H, m), 1.85 - 1.91 (7H, m), 2.06 (2H, s), 4.23 (1H, d), 4.44 (2H, q), 6.13 (1H, d), 7.62 (1H, d),

15 7.62 (1H, t), 7.75 - 7.76 (1H, m), 7.92 (1H, d)

m/z (ESI+) (M+H) $^+$ = 518

Intermediate# 118: N-(2-adamantyl)-1-[4-chloro-3-(trifluoromethyl)phenyl]-5-tert-butyl-pyrazole-4-carboxamide




N-(2-adamantyl)-1-[4-chloro-3-(trifluoromethyl)phenyl]-5-tert-butyl-pyrazole-4-carboxamide was prepared from 5-tert-butyl-1-(4-chloro-3-(trifluoromethyl)phenyl)-1H-pyrazole-4-carboxylic acid (**Intermediate#119**) by the same process used for Intermediate#114.

^1H NMR (300.072 MHz, cdCl_3) δ 1.29 (s, 9H), 1.55 - 1.97 (m, 12H), 2.06 (s, 2H), 4.22 (d, 1H), 6.12 (d, 1H), 7.48 (d, 1H), 7.62 (d, 1H), 7.63 (s, 1H), 7.71 (s, 1H)

m/z (ESI+) ($M+H$) $^+$ = 480

Intermediate# 119: 5-tert-butyl-1-(4-chloro-3-(trifluoromethyl)phenyl)-1H-pyrazole-4-carboxylic acid




A solution of Sodium hydroxide (6.67 mL, 13.34 mmol) was added in one portion to a stirred solution of ethyl 5-tert-butyl-1-(4-chloro-3-(trifluoromethyl)phenyl)-1H-pyrazole-4-carboxylate (**Intermediate#120**) (1 g, 2.67 mmol) in methanol (20 mL). The resulting suspension was stirred at 80 °C for 6 hours. The resulting mixture was evaporated to remove the methanol, washed with ether (20 mL). The reaction mixture was acidified

with 2M HCl, extracted with ethyl acetate(2x30mL) The organic layers were combined and washed with water (10mL) and brine (10mL), dried over MgSO₄, filtered and evaporated to afford pure 5-tert-butyl-1-(4-chloro-3-(trifluoromethyl)phenyl)-1H-pyrazole-4-carboxylic acid (0.766 g, 83 %).

¹H NMR (300.072 MHz, cdcl₃) δ 1.36 (s, 9H), 7.47 (d, 1H), 7.63 (d, 1H), 7.70 (s, 1H), 8.12 (s, 1H)

m/z (ESI+) (M+H)⁺ = 345

Intermediate # 120: Ethyl 5-tert-butyl-1-(4-chloro-3-(trifluoromethyl)phenyl)-1H-pyrazole-4-carboxylate




Ethyl 5-tert-butyl-1-(4-chloro-3-(trifluoromethyl)phenyl)-1H-pyrazole-4-carboxylate was prepared from (Z)-ethyl 2-((dimethylamino)methylene)-4,4-dimethyl-3-oxopentanoate (**Intermediate#83**) and (4-chloro-3-(trifluoromethyl)phenyl)hydrazine hydrochloride by the same process used for Intermediate#116.

¹H NMR (300.072 MHz, cdcl₃) δ 1.32 (s, 9H), 1.38 (t, 3H), 4.32 (q, 2H), 7.46 (d, 1H), 7.61 (d, 1H), 7.68 (s, 1H), 7.98 (s, 1H)

m/z (ESI+) (M+H)⁺ = 375

Intermediate# 121: (1R,2S,3S,5S)-5-Difluoromethoxy-adamantan-2-ylamine




((1R,2S,3S,5S)-5-Difluoromethoxy-adamantan-2-yl)-carbamic acid benzyl ester (**Intermediate#122**) (255 mg, 0.73 mmol) and Palladium (10% on carbon) (25 mg, 0.23 mmol) in MeOH (5 mL) were stirred under an atmosphere of hydrogen at 1 atm and ambient temperature for 22 hours. The reaction mixture was filtered through celite and the

solvent was evaporated in vacuo to yield (1R,2S,3S,5S)-5-Difluoromethoxy-adamantan-2-ylamine (130 mg, 82 %) as a clear oil.

¹H NMR (400.13 MHz, CDCl₃) δ 1.38 - 1.44 (5H, m), 1.91 - 2.00 (9H, m), 2.14 (1H, s), 3.05 (1H, s), 6.15 - 6.54 (1H, t)

5 m/z (ESI+) (M+H)⁺ = 218


Intermediate# 122: ((1R,2S,3S,5S)-5-Difluoromethoxy-adamantan-2-yl)-carbamic acid benzyl ester



Solution of 2,2-Difluoro-2-(fluorosulfonyl)acetic acid (0.237 mL, 2.30 mmol) in
 10 acetonitrile (2mL) was added dropwise to a stirred solution of ((1R,2S,3S,5S)-5-Hydroxy-adamantan-2-yl)-carbamic acid benzyl ester (346 mg, 1.15 mmol) and Copper(I) iodide (7.78 μL, 0.23 mmol) in acetonitrile (10 mL) at 45°C, over a period of 1 hour under nitrogen. The resulting solution was stirred at 45°C for 15 minutes. The reaction mixture was evaporated to dryness and redissolved in EtOAc (50 mL), and washed sequentially
 15 with water (50 mL) and saturated brine (25 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product. The crude product was purified by flash silica chromatography, elution gradient 0 to 30% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford ((1R,2S,3S,5S)-5-Difluoromethoxy-adamantan-2-yl)-carbamic acid benzyl ester (269 mg, 66.7 %) as a colourless oil.

20 ¹H NMR (400.13 MHz, DMSO-d₆) δ 1.35 (2H, d), 1.86 (4H, d), 1.94 (4H, t), 2.06 (3H, s), 3.65 (1H, m), 5.04 (2H, s), 6.65 - 7.03 (1H, t), 7.29 - 7.42 (6H, m)

Intermediate# 123: methyl 4-hydrazinylbenzoate hydrochloride



Hydrogen chloride 4M in Dioxan (100 mL, 399.60 mmol) was added to 4-Hydrazinobenzoic acid (15.2 g, 99.90 mmol) in MeOH (200 mL). The resulting suspension was stirred at 90 °C for 5 hours. After cooling to 20°C the precipitate was collected by filtration, washed with Et₂O (100 mL) and dried under vacuum to afford 2-(4-(methoxycarbonyl)phenyl)hydrazinium chloride (16.50 g, 82 %) as a cream crystalline solid.

m/z (ESI-) (M-H)- = 165; HPLC t_R = 1.12 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 3.81 (3H, s), 6.99 - 7.02 (2H, m), 7.86 - 7.90 (2H, m), 8.98 (1H, s), 10.47 (3H, s)

Intermediate# 123 may also be prepared as follows:


Methanolic hydrochloric acid solution (4M) (4 equiv., freshly prepared) was added to a suspension of 4-hydrazinobenzoic acid (1 equiv.) in methanol (12.6 vols.), under nitrogen.

The mixture was stirred under reflux for three hours and then cooled to below 15 °C. The solid was collected by filtration, washed with MTBE (6.5 vols.) and dried in air to give the product as a solid.

TLC DCM : MeOH, 9 : 1, Product R_f 0.87

mp 233.8 – 234.6 °C

Intermediate # 124 : N-adamantan-2-yl-1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxamide




2-Adamantanamine hydrochloride (0.375 g, 2.00 mmol) was added in one portion to 1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxylic acid (**Intermediate# 125**) (0.562 g, 2 mmol), 1-Hydroxybenzotriazole (0.297 g, 2.20 mmol), N-Ethyl-diisopropylamine (1.384 mL, 8.00 mmol) and 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.460 g, 2.40 mmol) in DMF (10 mL) under nitrogen.

The resulting solution was stirred at room-temperature for 16 hours. The reaction mixture was poured onto water (75 mL), extracted with EtOAc (2 x 50 mL), the organic layer was dried over MgSO₄, filtered and evaporated to afford beige solid. The crude product was purified by flash silica chromatography, elution gradient 20 to 60% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford N-adamantan-2-yl-1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxamide (0.480 g, 57.9 %) as a white solid.

m/z (ESI+) (M+H)⁺ = 415; HPLC t_R = 2.82 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.51 (2H, d), 1.70 (2H, s), 1.81 (5H, s), 1.84 (1H, s), 1.92 (2H, s), 2.04 (2H, d), 4.01 (1H, t), 7.73 (2H, d), 8.08 - 8.11 (2H, m), 8.18 (1H, s), 8.34 (1H, d).

Intermediate # 125 : 1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxylic acid




Potassium trimethylsilanolate (3.32 g, 23.28 mmol) was added in one portion to ethyl 1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxylate (**Intermediate# 126**) (2.4 g, 7.76 mmol), in THF (50 mL), under nitrogen. The resulting suspension was stirred at room temperature for 3 hours. The reaction mixture was evaporated to dryness and redissolved in EtOAc (100 mL), and washed sequentially with 0.1M HCl (50 mL),

water (50 mL), and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product. The crude solid was triturated with isohexane to give a solid which was collected by filtration and dried under vacuum to give 1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxylic acid (1.600 g, 73.3 %) as a orange solid. Used directly in next stage.

m/z (ESI+) (M-H)⁻ = 280; HPLC t_R = 1.86 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 7.78 - 7.80 (2H, m), 8.07 - 8.10 (2H, m), 8.26 (1H, s), 13.39 (1H, s)

Intermediate# 126 : Ethyl 1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxylate.




A solution of (Z)-ethyl 2-(ethoxymethylene)-4,4,4-trifluoro-3-oxobutanoate (2.41 g, 10.03 mmol) in ethanol (10 mL) was added dropwise to a stirred solution of 4-hydrazinylbenzonitrile hydrochloride (1.702 g, 10.03 mmol), in ethanol (50 mL) cooled to -10°C. The resulting solution was allowed to stir at room temperature for 16 hours. The reaction mixture was evaporated to dryness and the residue redissolved in EtOAc (100 mL), and washed sequentially with water (50 mL) and saturated brine (50 mL). The organic layer was dried over MgSO₄, filtered and evaporated to afford crude product. The crude product was purified by flash silica (120g) chromatography, elution gradient 10 to 60% EtOAc in isohexane. Pure fractions were evaporated to dryness to afford ethyl 1-(4-cyanophenyl)-5-(trifluoromethyl)-1H-pyrazole-4-carboxylate (2.400 g, 77 %) as a white solid.

m/z (ESI+) (M+H)⁺ = 310; HPLC t_R = 2.51 min.

¹H NMR (400.13 MHz, DMSO-d₆) δ 1.31 (3H, t), 4.33 (2H, q), 7.81 - 7.84 (2H, m), 8.10 - 8.13 (2H, m), 8.38 - 8.38 (1H, m)

Claims

1. A compound of the formula (1):



(1)

wherein:

Q is O, S or a single bond;

R¹ is C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₇cycloalkyl, heterocyclyl, arylC₁₋₃alkyl, heteroarylC₁₋₃alkyl, C₃₋₇cycloalkylC₁₋₃alkyl, C₃₋₇cycloalkylC₂₋₃alkenyl or C₃₋

7cycloalkylC₂₋₃alkynyl, [each of which is optionally substituted by 1, 2 or 3 substituents independently selected from C₁₋₃alkyl, hydroxy, halo, oxo, cyano, trifluoromethyl,

C₁₋₃alkoxy, C₁₋₃alkylS(O)_n- (wherein n is 0, 1, 2 or 3), R⁵CON(R^{5'})-, (R^{5'})(R^{5''})NC(O)-, R⁵C(O)O-, R⁵OC(O)-, (R^{5'})(R^{5''})NC(O)N(R^{5'''})-, R⁵SO₂N(R^{5''})-, and (R^{5'})(R^{5''})NSO₂- (wherein R⁵ is C₁₋₃alkyl optionally substituted by 1, 2 or 3 substituents selected from

hydroxyl, halo or cyano; and

R^{5'} and R^{5''} are independently selected from hydrogen and C₁₋₃alkyl optionally substituted by 1, 2 or 3 substituents independently selected from hydroxyl, halo, C₁₋₃alkoxy, carboxy and cyano or R^{5'} and R^{5''} together with the nitrogen atom to which they are attached form a 4-7 membered saturated ring];

R² is selected from heterocyclyl, C₃₋₇cycloalkyl(CH₂)_m-, and C₆₋₁₂polycycloalkyl(CH₂)_m- (wherein m is 0, 1 or 2 and the rings are optionally substituted by 1, 2 or 3 substituents independently selected from R⁶);


R³ is selected from hydrogen, C₁₋₄alkyl C₃₋₅cycloalkyl and C₃₋₅cycloalkylmethyl (each of which is optionally substituted by 1, 2 or 3 fluoro atoms);

R² and R³ together with the nitrogen atom to which they are attached form a saturated mono, bicyclic or bridged ring system optionally containing 1 or 2 additional ring

- heteroatoms selected from nitrogen, oxygen and sulphur and which is optionally fused to a saturated, partially saturated or unsaturated monocyclic ring wherein the resulting ring system is optionally substituted by 1, 2, or 3 substituents independently selected from R^7 ; R^6 and R^7 are independently selected from hydroxyl, halo, oxo, carboxy, cyano,
- 5 trifluoromethyl, R^9 , R^9O- , R^9CO- , $R^9C(O)O-$, $R^9CON(R^{9'})-$, $(R^{9'})(R^{9''})NC(O)-$, $(R^{9'})(R^{9''})N-$, $R^9S(O)_a-$ wherein a is 0 to 2, $R^9OC(O)-$, $(R^{9'})(R^{9''})NSO_2-$, $R^9SO_2N(R^{9''})-$, $(R^{9'})(R^{9''})NC(O)N(R^{9''''})-$, phenyl and heteroaryl [wherein the phenyl and heteroaryl groups are optionally fused to a phenyl, heteroaryl or a saturated or partially-saturated 5- or 6-membered ring optionally containing 1, 2 or 3 heteroatoms independently selected from
- 10 nitrogen, oxygen and sulphur and the resulting ring system is optionally substituted by 1, 2 or 3 substituents independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, $N-C_{1-4}$ alkylamino, di- $N,N-(C_{1-4}$ alkyl)amino, $N-C_{1-4}$ alkylcarbonyl, di- $N,N-(C_{1-4}$ alkyl)carbonyl, C_{1-4} alkylS(O) $_r-$, C_{1-4} alkylS(O) $_rC_{1-4}$ alkyl (wherein r is 0, 1 or 2)];
- 15 R^9 is independently selected from C_{1-3} alkyl optionally substituted by hydroxyl, halo, C_{1-4} alkoxy, carboxy or cyano; $R^{9'}$, $R^{9''}$ and $R^{9''''}$ are independently selected from hydrogen and C_{1-3} alkyl optionally substituted by 1, 2, or 3 substituents independently selected from hydroxyl, halo, C_{1-4} alkoxy, carboxy and cyano);
- 20 A is a phenyl or heteroaryl ring (the phenyl or heteroaryl ring being optionally substituted on ring carbon atoms by 1, 2 or 3 R^{10} groups and on an available ring nitrogen in a heteroaryl group by R^{11});
- R^{10} is independently selected from C_{1-4} alkyl, hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, halo, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, amino, $N-C_{1-4}$ alkylamino, di-
- 25 $N,N-(C_{1-4}$ alkyl)amino, $N-C_{1-4}$ alkylcarbonyl, di- $N,N-(C_{1-4}$ alkyl)carbonyl, C_{1-4} alkylS(O) $_s-$, C_{1-4} alkylS(O) $_sC_{1-4}$ alkyl (wherein s is 0, 1 or 2)];
- R^{11} is independently C_{1-3} alkyl optionally substituted by 1, 2 or 3 fluoro atoms;
- X is a direct bond, C_{3-4} cycloalkandiyl, C_{3-4} cycloalkanylidene, $-C(R^{12})(R^{13})-$, $-C(R^{12})(R^{13})C(R^{14})(R^{15})-$, $-CH_2O-$ or $-CH_2S(O)-$ (wherein t is 0, 1 or 2);
- 30 Y is a direct bond, C_{3-4} cycloalkandiyl, C_{3-4} cycloalkanylidene, $-C(R^{16})(R^{17})-$ or $-C(R^{18})(R^{19})C(R^{20})(R^{21})-$;


wherein R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} and R^{21} are independently selected from hydrogen and methyl;
or a pharmaceutically-acceptable salt thereof.

2. A compound according to claim 1 wherein Q is S.
3. A compound according to claim 1 wherein Q is a single bond.
4. A compound according to any one of claims 1 to 3 wherein R^2 is selected from C_5 -
7cycloalkyl(CH₂)_m-, C_{7-10} bicycloalkyl(CH₂)_m- and C_{10} tricycloalkyl(CH₂)_m- (wherein the
cycloalkyl, bicycloalkyl and tricycloalkyl rings are optionally substituted by 1, 2 or 3
substituents independently selected from R^6 wherein R^6 is as defined in claim 1) and
wherein m is 0, 1 or 2 and R^3 is hydrogen.
5. A compound according to any one of claims 1 to 4 wherein X is a direct bond.
6. A compound according to any one of claim 1 to 5 wherein Y is selected from direct
bond, -CH₂- and -CH₂CH₂-.
7. A compound according to claim 1 of the formula (1a):



wherein R^1 , R^2 and R^3 are as defined in claim 1 and R^{10} is selected from hydrogen, C_{1-4} alkyl, trifluoromethyl, C_{1-4} alkoxy and C_{1-4} alkylS-; or a pharmaceutically-acceptable salt thereof.

- 5 8. A compound according to claim 1 of the formula (1b):



- wherein R^1 , R^2 and R^3 are as defined in claim 1 and R^{10} is selected from hydrogen, C_{1-4} alkyl, trifluoromethyl, C_{1-4} alkoxy and C_{1-4} alkylS-; or a pharmaceutically-acceptable salt thereof.
- 10

9. A compound according to claim 1 selected from:
- 4-[4-[[[(1S,3R)-5-hydroxy-2-adamantyl]carbamoyl]-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 15 4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(1-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(N-cyclohexyl-N-methyl-carbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(oxan-4-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]benzoic acid;
- 4-[5-propylsulfanyl-4-[3-[2-(trifluoromethyl)phenyl]pyrrolidine-1-carbonyl]pyrazol-1-
- 20 yl]benzoic acid;
- 4-[4-(cyclohexylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(2-adamantylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(1-adamantylcarbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;
- 4-[4-(cyclohexyl-methyl-carbamoyl)-5-cyclopropyl-pyrazol-1-yl]benzoic acid;

- 4-[5-cyclopropyl-4-[(4-hydroxy-1-adamantyl)carbamoyl]pyrazol-1-yl]benzoic acid;
 2-[4-[4-(cyclohexylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid;
 2-[4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]phenyl]acetic acid;
 4-(4-cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid;
 3-(4-cyclohexylcarbamoyl-5-propylsulfanyl-pyrazol-1-ylmethyl)-benzoic acid;
 3-[4-(adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-ylmethyl]-benzoic acid;
 4-[4-(adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-ylmethyl]-benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]benzoic acid;
 5 4-[4-(2-adamantylcarbamoyl)-5-(1-methylcyclopropyl)pyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-cyclopentyl-pyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-ethylpyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-propan-2-ylpyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-cyclobutylpyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-methyl-pyrazol-1-yl]benzoic acid;
 4-(5-tert-butyl-4-(cyclohexylcarbamoyl)-1H-pyrazol-1-yl)benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[4-(1-adamantylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[5-cyclohexylsulfanyl-4-[(5-hydroxy-2-adamantyl)carbamoyl]pyrazol-1-yl]benzoic acid;
 4-[5-cyclohexylsulfanyl-4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoic acid;
 10 4-[4-(2-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid;
 methyl 4-[4-(1-adamantylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoate;
 methyl 4-[5-cyclopentylsulfanyl-4-[(1R,3S)-5-hydroxy-2-adamantyl]carbamoyl]pyrazol-1-yl]benzoate;
 4-[4-[[5-(difluoromethoxy)-2-adamantyl]carbamoyl]-5-propylsulfanylpyrazol-1-yl]benzoic acid;
 4-[4-(cyclohexylcarbamoyl)-5-cyclopentylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[4-(cyclohexylcarbamoyl)-5-cyclohexylsulfanyl-pyrazol-1-yl]benzoic acid;
 15 4-[5-cycloheptylsulfanyl-4-(cyclohexylcarbamoyl)pyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-ethylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[4-(2-adamantylcarbamoyl)-5-methylsulfanyl-pyrazol-1-yl]benzoic acid;
 4-[4-(5-methanesulfonyl-adamantan-2-ylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-

benzoic acid;

4-[4-(2-adamantylcarbamoyl)-5-propylsulfanyl-pyrazol-1-yl]-2-methoxy-benzoic acid;

4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-3-methyl-benzoic acid;

4-[4-(2-adamantylcarbamoyl)-5-tert-butyl-pyrazol-1-yl]-2-(trifluoromethyl)benzoic acid;

5 and

4-[4-(adamantan-2-ylcarbamoyl)-5-(trifluoromethyl)-1H-pyrazol-1-yl]benzoic acid.;

and pharmaceutically-acceptable salts thereof.

10 10. A pharmaceutical composition, which comprises a compound of formula (I), or a pharmaceutically acceptable salt thereof, as claimed in claim 1 in association with a pharmaceutically-acceptable diluent or carrier.


11. A compound of formula (1), or a pharmaceutically acceptable salt thereof, as claimed in claim 1, for use in a method of prophylactic or therapeutic treatment of a
15 warm-blooded animal, such as man.

12. A compound of formula (1), or a pharmaceutically acceptable salt thereof, as claimed in claim 1 for use as a medicament.

20 13. The use of a compound of formula (1), or a pharmaceutically acceptable salt thereof, as claimed in claim 1 in the manufacture of a medicament for use in the production of an 11 β HSD1 inhibitory effect in a warm-blooded animal, such as man.

14. A process for preparing a compound of the formula (1) or a pharmaceutically
25 acceptable salt thereof which process comprises any one of processes a) or b):
a) hydrolysis of an ester of formula (2):

166




(2)

wherein R²² is an alkyl or aryl group; or

b) converting Z in a compound of the formula (3):

5



(3)

into a carboxy group, wherein Z is an functional group capable of conversion into a carboxylic acid;

10 and thereafter if necessary or desirable:

- i) converting a compound of the formula (1) into another compound of the formula (1);
- ii) removing any protecting groups;
- iii) resolving enantiomers;
- iv) forming a pharmaceutically-acceptable salt thereof.

15