


Description

The present invention relates to certain 5-hetero- or aryl-substituted-imidazo[2,1-a]isoquinolines and to their use as platelet activating factor (PAF) receptor antagonists. The invention also relates to pharmaceutical compositions containing a select group of the afore-mentioned compounds as an active ingredient thereof. In addition, the invention relates to the use of a select group of said compounds as anti-tumor agents.

USP 3,887,566 discloses certain 2,3-dihydroimidazo-isoquinolines exhibiting analgesic, anti-inflammatory, anti-bacterial, anti-viral and cardiovascular properties. USP 4,100,165 discloses certain 5-hydroxy-2,3,5,6-tetrahydrofuran imidazo-[2,1-a]isoquinolines containing a pyridyl-, thienyl- or furyl ring in the 5-position, which compounds are useful as anorexics and anti-depressants. USP 4,101,553 discloses certain 5-hydroxy-2,3,5,6-tetrahydrofuran imidazo[2,1-a]isoquinolines containing an optionally substituted aryl group in the 5-position, said compounds useful as anorexics and anti-depressants.

The essence of the present invention is the discovery that certain 5-hetero- or aryl-substituted-imidazo-(2,1-a)-isoquinolines are useful as PAF receptor antagonists and in addition, that a select group thereof are useful as anti-tumor agents.

In one aspect therefore the invention concerns the use of compounds of formula I



wherein each R independently is hydrogen or methyl,

R₁ is in 8- or 9-position and represents hydrogen, chloro, or C₁₋₃alkyl and


R₂ represents thienyl, furyl or a group



wherein each Ro represents independently hydrogen, fluoro, chloro, C₁₋₁₀alkyl or C₁₋₁₀alkoxy or one Ro is hydrogen and the other represents

a) tri-(C₁₋₃alkyl)silyl, trifluoromethyl or phenyl;


b) a group




wherein each R₃ represents independently hydrogen, chloro, fluoro, C₁₋₄alkyl or C₁₋₅alkoxy and X represents

-CH₂O-, -OCH₂-, -CH₂-, -CH₂CH₂-, or -O-;

c) a group




wherein each R_3' represents independently C_1-3 alkoxy and
 Y represents $-O(CH_2)_{1-5}-$, $-(CH_2)_{1-6}-$ or $-CH_2OCH_2-$; or
 d) a group




wherein each R_4 represents independently C_1-4 alkyl, C_5-7 cycloalkyl, allyl or benzyl; or the two R_4 's together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl or thiomorpholinyl; or two R_4 's on adjacent carbon atoms form methylenedioxy, with the proviso that when an R_4 is other than chloro, fluoro, methyl or methoxy it may only be in a meta or para position and their pharmaceutically acceptable acid addition salts where such may exist, in the manufacture of a medicament for use in inhibiting PAF mediated bronchoconstriction and extravasation and PAF mediated endotoxin induced lung injury

Preferred compounds of formula I are those wherein each R is hydrogen, R_1 is as defined above and one R_4 is hydrogen and the other is in meta or para position and is as defined under b) or c) above, and their pharmaceutically acceptable acid addition salts where such exist (compounds I').

Another aspect of the present invention concerns novel compounds of formula I i.e. compounds of formula Ia



wherein R and R_1 are as defined above and
 R_2' represents thienyl, furyl or a group




wherein one RO'' is hydrogen and the other represents tri- $(C_1-3$ alkyl)-silyl or phenyl, or is as defined under b), c) or d) above whereby RO'' substituents may only be in meta or para positions; and their pharmaceutically acceptable acid addition salts where such may exist.

Preferred compounds of formula Ia are those wherein each R is hydrogen, R_1 is in 8-position and is otherwise as defined above and R_2 represents thienyl, furyl or a group




wherein RO''' is in meta or para position and represents tri-(C_{1-3} alkyl)silyl or is as defined under b), c) or d) above, and their pharmaceutically acceptable acid addition salts where such may exist (compound Ia').


A further aspect of the invention concerns the use of compounds of formula Ib



wherein R and R_1 are as defined above and R_2''' represents thienyl, furyl or a group




wherein each RO^IV represents independently, chloro, fluoro, or C_{1-6} alkyl or one RO^IV is hydrogen and the other is C_{2-6} alkyl tri-(C_{1-3} alkyl) silyl, phenyl or is as defined under b) or c) above or is d)' a group




wherein each R_4' is allyl or the two R_4'' s together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl or thiomorpholinyl, with the proviso that when RO^IV is other than chloro, fluoro or methyl it may only be in meta- or para-position and their pharmaceutically acceptable acid addition salts, where such may exist for use in the manufacture of a medicament for use in treating tumors.

Preferred compounds of formula Ib are those wherein R and R_1 are as defined above and R_2''' represents thienyl, furyl or a group



wherein RO^V is in meta or para position and represents C_{2-6} alkyl, tri-(C_{1-3} alkyl)silyl, phenyl, is as defined under b) or c) above or represents d)' a group



5

wherein each R_4'' is allyl or the two R_4'' 's together with the nitrogen atom to which they are attached represent pyrrolidinyl or piperidyl; and their pharmaceutically acceptable acid addition salts, where such may exist (compound lb').

10

Alkyl moieties may be straight chained or branched.

A particular compound group covers those of formula I (and analogously for Ia and Ib) wherein

R_1 is as defined above and

R_2 is thienyl, furyl or a group


15



20

wherein each Ro represents independently hydrogen, fluoro, chloro, C_{1-10} alkyl, C_{1-10} alkoxy or C_{1-10} alkylthio or one is hydrogen and the other is as defined under a) or d) above or b)' a group


25



30

wherein each R_{3p} represents independently hydrogen, chloro, fluoro or C_{1-3} alkoxy and X is as defined above or c)' a group

35



40

wherein each $\text{R}_{3'p}$ represents C_{1-3} alkoxy and Y represents $-\text{OCH}_2-$, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$ or CH_2OCH_2 , and their pharmaceutically acceptable acid addition salts where such may exist.

45

In another compound group of formula I (and analogously for Ia and Ib)

R_1 is in 8- or 9-position and represents hydrogen, fluoro or C_{1-3} alkyl and

R_2 represents thienyl, furyl or a group

50



55


wherein each Ro represents independently hydrogen, fluoro, chloro, C_{1-10} alkyl or C_{1-10} alkoxy or one Ro is hydrogen and the other represents

a) tri- $(\text{C}_{1-3}$ alkyl)silyl, trifluoromethyl or phenyl


b) a group



wherein each R_3 represents independently hydrogen, chloro, fluoro or C_{1-5} alkoxy and X represents $-OCH_2-$, $-O-$, $-CH_2O-$ or $-CH_2CH_2-$ or
c) a group




wherein each $R_{3'}$ represents independently C_{1-3} alkoxy and Y represents $-OCH_2-$, $-CH_2OCH_2-$, $-CH_2-$, $-CH_2CH_2-$ or $-CH_2CH_2CH_2-$ or
d) a group



wherein each R_4 represents independently C_{1-4} alkyl, C_{5-7} cycloalkyl, allyl or benzyl or the two R_4 's together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl or thiomorpholinyl or two R_4 's on adjacent carbon atoms stand for methylenedioxy, with the proviso that when an R_4 is other than chloro, fluoro, methyl or methoxy it may only be in a meta- or para-position.

The compounds of formula I may be prepared by dehydrating a compound of formula V




V


wherein R_1 and R_2 are as defined above and recovering the compound obtained in free form or in the form of a pharmaceutically acceptable acid addition salt where such may exist.

This reaction is carried out in an inert organic solvent and in the presence of an acid catalyst at temperatures e.g. of 35° to 200° C especially between 75° and 120° C. Examples of suitable solvents are aliphatic hydrocarbons e.g. hexane, heptane and the like, aromatic hydrocarbons e.g. benzene, toluene and the like, chlorinated hydrocarbons e.g. chloroform, methylenechloride and the like, aliphatic ethers e.g. diethyl ether, cyclic ethers e.g. tetrahydrofuran or an excess of acid catalyst e.g. acetic acid. Examples of suitable acid catalysts are mineral acids e.g. hydrochloric acid, sulphuric acid, phosphoric acid and the like or organic acids such as arylcarboxylic acids e.g. benzoic acid, alkylsulphonic acids e.g. methanesulphonic acid, aryl sulphonic acids e.g. p-toluenesulphonic acid or preferably alkylcarboxylic acids especially acetic acid. The invention also concerns a process for preparing compounds of formula Ia.

The compound of formula V may be prepared by reacting a compound of formula III




10 with a compound of formula IV



wherein R_1 and R_2 are as defined above and R_5 represents C_{1-4} alkyl and hydrolysing the adduct thus formed and recovering the compound obtained in free form or in the form of a pharmaceutically acceptable acid addition salt.

20 The reaction is carried out under nitrogen atmosphere in an inert solvent such as an aliphatic hydrocarbon e.g. hexane, heptane and the like, an aliphatic ether e.g. diethyl ether or a cyclic ether e.g. tetrahydrofuran at temperatures of e.g. -30° to 50° C preferably -20° to 0° C. Hydrolysis is conducted in conventional manner e.g. employing water, dilute mineral acid, ammonium chloride solution or the like.

25 The compounds of formula III can be prepared by reacting a compound of formula II




35 with R_5Li wherein R_1 and R_5 are as defined above.

The reaction is carried out under a nitrogen atmosphere in an inert solvent such as described above for the reaction of III with IV at a temperature of e.g. 25° to 75° C preferably 30° to 40° C.

Certain compounds of formula V are new and also form part of the invention.

The invention thus also concerns compounds of formula Va



45 wherein R , R_1 and each Ro'' are as defined above and their pharmaceutically acceptable acid addition salts.

The starting materials of formula II and IV are either known and/or may be obtained analogously to known methods in conventional manner.

50 Final products and intermediates may be isolated and purified in conventional manner. Intermediates, where appropriate may be employed directly in the following step without purification.

As is evident to those skilled in the art, the compounds of formulae I and V may exist in racemic or enantiomeric form and the invention is intended to cover all forms.

Enantiomeric forms may be recovered in conventional manner e.g. by resolution of end or intermediate

products or by employing optically active starting materials.

Examples of pharmaceutically acceptable acid addition salts include those with mineral acids e.g. hydrochloric, hydrobromic, phosphonic and sulphuric and organic acids e.g. tartaric, acetic, citric, malic, maleic, methanesulphuric and gluconic which may be prepared in conventional manner.

5 The compounds of formula I are indicated for use as platelet activating factor receptor antagonists as indicated by their ability to inhibit specific binding of [^{3} H]-PAF to platelets according to the Human Platelet PAF Receptor Assay test (Test A) as follows:

Human blood is obtained by venipuncture of healthy, human donors into an anti-coagulant mixture containing 3.15% of trisodium citrate and 20 ug/ml of Prostaglandin I_2 (PGI $_2$) in a ratio of blood to anti-coagulant of 9:1. Platelet rich plasma (PRP) is prepared by centrifugation (250 x g) of the blood for 20 minutes at room temperature. The PRP is then centrifuged (900 x g) for 10 minutes at room temperature and the platelet pellet is washed two times with Tris-Tyrode's (TT) solution having a pH of 7.4 and containing 0.25% bovine serum albumin (BSA), and to which has been added PGI $_2$ at a final concentration of 0.3 μ g/ml. The platelets are resuspended at 350,000 μ l in TT/BSA containing 1.4 mM CaCl $_2$ ·2H $_2$ O and 0.7 mM MgCl $_2$ ·6H $_2$ O. All of the tests are conducted in duplicate and each of the test compounds is evaluated at concentrations of 100, 50, 1 and 0.1 μ M. For each determination, the following solutions are mixed:

500 μ l of the above-described platelets;
10 μ l of [^{3} H]-PAF (40,000 counts per minute (cpm) to a final concentration of 1.5 μ M); and
either
10 μ l of the test compound at 50x the desired final concentration,
10 μ l of vehicle (total bound), or
10 μ l of 1.85×10^{-5} M cold PAF (non-specifically bound).

Each mixture is allowed to incubate at room temperature for one hour, after which time the reaction is terminated by the addition of 500 μ l of ice cold TT/BSA and centrifugation (900 x g) at 4° C for 10 minutes. The resultant supernatant is aspirated into scintillation vials and the pellet is washed with 250 μ l of ice cold TT/BSA and centrifuged (900 x g) at 4° C for 10 minutes. The supernatants are then aspirated into the same scintillation vials as before and 10 ml of Scintiverse II (a liquid scintillation cocktail) is added to and mixed therewith. The pellets are resuspended in 500 μ l of Scintiverse II and mixed well. An additional 2 ml of Scintiverse II is then added to the vials and, after mixing, the vials are counted for 1 minute in a liquid scintillation spectrometer. The amount of specific binding is calculated as the difference in cpm between the total bound [^{3} H]-PAF and non-specifically bound [^{3} H]-PAF. The percent inhibition of specific binding is determined by dividing the cpm specifically bound in the presence of the test compound by the cpm specifically bound in total, multiplying by 100 and then subtracting from 100. An IC $_{50}$ (50% inhibitory concentration) value is generated by evaluating the test compound over the full concentration range.

Furthermore, in view of their potential as PAF receptor antagonists, the compounds of formula I are indicated for use as inhibitors of PAF-mediated bronchoconstriction, which property was evaluated by the PAF-induced Pulmonary Inflation Pressure (PIP) Increase test (Test B) as follows:

Male guinea pigs, weighing between 300 and 400 g, are anesthetized, after which time trachea tube, carotid and jugular catheters are inserted. The test animal is then force ventilated employing a small animal Harvard respirator and the resistance to lung inflation (PIP) is measured utilizing a pressure transducer and recorder. The test compound is administered either orally at 30 minutes up to 4 hours prior to or parenterally e.g. intravenously (jugular) etc. at 5 minutes up to 4 hours prior to the introduction of PAF. The PAF (C $_{18}$ -Sandoz, Hanover) is dissolved in Tris-Tyrode's bovine serum albumin buffer and administered intravenously (jugular) at 100 ng/kg. Any blood pressure measurements taken are recorded from a transducer attached to the carotid catheter. Two responses are noted in the PIP recordings after the PAF is administered: 1) an immediate response which, in PAF-only treated test animals, averages out to between 70% and 80% more than the baseline PIP values. (This early response is also the greatest response and is, therefore, termed maximal PIP); and 2) the long term (at least 30 minutes) PIP response which slowly decreases to baseline. A reading at 15 minutes after the administration of PAF is termed the endpoint PIP. The effect of the test compound on the PIP response is determined by the difference between the percent increase in maximal PIP over baseline for the test animal to which has been administered PAF and the test compound compared to the test animal to which only PAF has been administered.

Still further, the compounds of formula I are indicated for use as inhibitors of PAF-mediated extravasation (the extrusion of plasma from the lumen of the blood vessels into the vessel wall and surrounding tissues) measured as a function of hemoconcentration according to the PAF-induced Extravasation test (Test C) as follows:

Male guinea pigs, weighing between 300 and 400 g, are anesthetized, after which time a femoral

catheter is inserted. The test compound is administered either orally or parenterally one hour up to 4 hours prior to the introduction of PAF. The PAF (C₁₈-Sandoz, Hanover) is dissolved in Tris-Tyrode's bovine serum albumin buffer and administered intravenously (jugular) at 100 ng/kg.

To determine the hematocrit value, which is employed to index hemoconcentration and is defined as the percent of packed red blood cells in a sample of blood which is centrifuged to separate plasma from the cellular components, blood samples are collected in 50 μ l heparinized hematocrit tubes. These samples are taken just prior to the injection of PAF, one minute subsequent to the injection of PAF and every two minutes thereafter until 15 minutes has lapsed subsequent to the injection of PAF. The tubes are then centrifuged and the percent of packed red blood cells (hematocrit) is measured (PAF induces a maximal increase in hematocrit at 5 to 7 minutes subsequent to the injection of PAF). The percent increase in hematocrit over the value prior to the injection of PAF is calculated. The hematocrit values obtained with the test compound are compared to the hemoconcentration values obtained with PAF alone and are expressed as percent inhibition of percent increase in hematocrit.

Yet still further, the compounds of formula I are indicated for use as inhibitors of PAF-mediated, endotoxin-induced lung injury and, analogously, endotoxin-induced-septic shock and adult respiratory distress syndrome. The term endotoxin induced lung injury is used generically for all three indications. The ability of the compounds of formula I to inhibit PAF-mediated, endotoxin-induced lung injury can be measured in accordance with the test presented by S. Chang at the 2nd International Conference on Platelet Activating Factor and Structurally Related Alkyl Ether Lipids in Gatlinburg, Tennessee on October 26-29, 1986.

Based on previous reports that lung tissue and blood PAF increased in endotoxin-treated rats, it was determined that the intraperitoneal administration of 2mg/kg of endotoxin (*S. enteritidis*) causes acute lung injury, as assessed by the extravascular accumulation of water ¹²⁵I-albumin in perfused lungs isolated from rats ninety minutes following *in vivo* endotoxin treatment. Thus, the wet lung/body weight ratio (as an index of lung water content) increases from 5.33 ± 0.48 to 8.26 ± 0.36 and the albumin leak index increases from 0.46 ± 0.09 to 1.01 ± 0.07 . In order to measure the effectiveness of a compound as an inhibitor of endotoxin-induced lung injury, the test compound is administered intraperitoneally prior to the *in vivo* endotoxin treatment.

The ability of the compounds of formula I to inhibit PAF-mediated, endotoxin-induced septic shock can be measured in accordance with the test presented by C.N. Sessler, et al at the Annual Meeting of the American Federation for Clinical Research in New Orleans, Louisiana during January, 1987.

All sheep are prepared for testing employing the Chronic Sheep Lung Lymph preparation which is well documented in the literature, with the modifications that chronic tracheostomies are performed on the test animals and pleural pressure catheters inserted at the time of the initial surgery. All catheters are brought to the outside through stab wounds in the skin, the chest is closed and the test animals are allowed to recover for several days until they appear healthy and lung lymphs are free of blood before experiments are commenced.

In order to measure the effectiveness of a compound as an inhibitor of some of the important hemodynamic effects of endotoxin on septic shock, 1.3 μ g/kg of endotoxin or saline is administered to groups of test animals intravenously over a 30 minute period and 20mg/kg of the test compound or saline is administered intravenously over a five-hour period. The pulmonary arterial pressure (PAP), cardiac output (CO) and partial oxygen pressure (PO₂) are monitored continuously over the five-hour period.

The ability of the compounds of formula I to inhibit PAF-mediated, endotoxin-induced adult respiratory distress syndrome can be measured in accordance with the test presented by B.W. Christman, et al at the Annual Meeting of the American Thoracic Society and American Lung Association on May 10th -13th, 1987.

In order to measure the effectiveness of a compound as an inhibitor of some of the important hemodynamic effects of endotoxin on adult respiratory distress syndrome, 0.5 μ g/kg of *E.coli* endotoxin over a 20 minute period, 20mg/kg/hr of the test compound for 6 hours, or 0.5 μ g/kg of *E.coli* endotoxin 1 hour after commencing 20mg/kg/hr of the test compound for 6 hours, are administered to groups of test animals intravenously. The pulmonary arterial pressure (PAP), dynamic compliance (DC) of the lungs and lung lymph flow (LLF) are monitored continuously over a five-hour period.

The compounds are thus indicated for use in treating PAF mediated bronchoconstriction and extravasation and PAF mediated endotoxin induced lung injury and an indicated suitable daily dosage for this use is from about 10 to 2000 mg preferably 10 to 350 mg suitably administered in divided dosages of 0.25 to 500 mg (esp. 0.25 to 350 mg) up to four times daily or in controlled release form. A typical oral dosage is 50 or 100 mg two or three times a day. A typical oral unit dosage may contain 2.5 to 500 mg preferably 5 to 200 mg especially 10 to 100 mg of active substance.

The compounds of formula V are also indicated for use in the same indications and can be employed at

similar dosages.

The compounds of formula Ib are also indicated for use as anti-tumor agents as evidenced by their ability in inhibiting various lymphoma, sarcoma, myeloma and leukemia cell lines as indicated in the following tests.

Test D) Tumor cell cytotoxicity Test (TCCT).

In flat bottom microtiter plates (Nunc Roskilde, Denmark) were placed Abelson 8.1 tumor cells in DMEM + 10% fetal calf serum and the tumor cell-containing plates were incubated with 1,3, and 5 μ g of the test compound for a period of 6 to 72 hours. The number of viable tumor cells present was determined by measuring the alkaline phosphatase in the following manner. The tumor cell plates were centrifuged (500 x g.) for ten minutes and the supernatant flicked off. Without further washing, 100 μ l of buffer containing 20 μ l of diethanolamine, 2 μ M of MgCl₂ · 6H₂O, 2.5 μ M of p-nitrophenylphosphate and 10 mg Triton X-100 were added. The samples were incubated for 60 minutes at room temperature and the enzymatic reaction was terminated by the addition of 100 μ l of 0.5N NaOH. The absorbance was then measured at 405 nM using a Titertek Multiskan apparatus.

Test E) Influence on Cytotoxicity of ET-18-OCH₃ test (IC-ET test).

Bone marrow cell macrophages (10⁵/well) obtained from [BALB/CX57/BL₆]F₁ mice were incubated with 10 μ g of 1-octadecyl-2-methoxy-3-phosphoryl choline (ET-18-OCH₃) for 24 hours in flat bottom microtiter plates (Nunc Roskilde, Denmark), after which time they are centrifuged and washed once. Abelson 8.1 tumor cells in DMEM + 10% fetal calf serum and 1,3 and 5 μ g of the test compound were then added to the plates. With the cytotoxicity of ET-18-OCH₃ (10 μ g) alone set at 100%, the inhibition or enhancement of the cytotoxic effect, as measured by an alkaline phosphatase assay, was determined and values recorded after 72 hours for 1,3 and 5 μ g of the test substance.

Test F) Meth A fibrosarcoma cells were induced in BALB/C mice by administering methylcholanthrene according to the procedure of Old, et al. (L.J. Old, E.A. Boyse, D.A. Clarke, and E. Carswell, Ann. N.Y. Acad. Sci., 101, 80 (1962). These tumor cells were harvested from the peritoneal cavity 10 to 12 days after administration of methylcholanthrene. Ten CBF₁ mice of 10-12 week age were each implanted with 7.3 x 10⁶ Meth A sarcoma cells to serve as control. A second group of ten CBF₁ mice were each implanted with 7.3 x 10⁶ Meth A sarcoma cells and on day one after implant each mouse was treated p.o. with 5-50 μ g of the test compound per day for a total of twenty or twenty-seven days. Tumor growth and survivors were assayed on days 7, 14, 21 and 28 after tumor implantation. Under these conditions, none of the control group animals survived, whereas all of the animals which were administered 50 μ g per day of the compound of Example 3N) for 28 days not only survived but showed no evidence of the presence of any tumors as well.

The compounds are thus further indicated for use in inhibiting tumors and an indicated suitable dosage for this use is from about 500-2000 mg preferably 1000 - 1500 mg administered in divided dosages of e.g. 125 to 750 mg up to 4 times daily or in controlled release form. A typical oral dosage is 400 mg two or three times a day or 20 mg/kg body weight intravenously over a 24 hours period. A typical oral unit dosage may contain 300 to 600 mg preferably 300 to 500 mg especially 350 to 450 mg of active substances.

A suitable daily dosage will depend on a number of factors such as relative potency of activity of the compound. The preferred compound according to the invention 5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride achieved e.g. the following results in tests A to C, Test A -IC₅₀ 0.06 μ M, Test B ED₅₀ 5.0 mg/kg p.o., Test C -ED₅₀ 4.2 mg/kg p.o. The compound 5-(4'-diallylaminoethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride achieved e.g. the following results in tests D and E : Test D - 99% inh. at 5 μ g, Test E - 97% inh. at 5 μ g.

The invention therefore also concerns a the use of a compound of formula I or a pharmaceutically acceptable salt thereof in the manufacture of medicaments for treating PAF mediated bronchoconstriction and extravasation and PAF-mediated endotoxin induced lung injury.

The invention further concerns the use of a compound of formula Ib or a pharmaceutically acceptable salt thereof in the manufacture of medicaments for use as anti-tumor agents.

The compounds may be employed in free form or in the form of pharmaceutically acceptable acid addition salts where such exist.

The compounds of the invention, in free form or in pharmaceutically acceptable acid addition salt form may be combined with one or more pharmaceutically acceptable carriers and, optionally, one or more other conventional pharmaceutical adjuvants and administered orally in the form of tablets, dispersible powders,

granules, capsules, elixirs, suspensions and the like or parenterally in the form of sterile injectable solutions or suspensions. The compositions may be prepared by conventional means.

Examples of compositions are as follows:

Solid unit dosage forms (weight (mg))				
	tablet		capsule	
Compound of formula I (e.g. of Ex. 7E)	50	-	50	-
Compound of formula Ib (e.g. of Ex 3N)	-	400	-	400
tragacanth	10	10	-	-
lactose (spray-dried)	212.5	197.5	100	250
corn starch	15	25	-	-
talcum	10	15	-	-
magnesium stearate	2.5	2.5	-	-
Total	300.0	650.0	150.0	650.0

Liquid forms (weight (mg))		
Ingredients	sterile injectable suspension	oral liquid suspension
compound of formula I, e.g., the compound of Example 7E)	5	3
sodium carboxymethylcellulose	U.S.P 1	8
methyl cellulose	0.3	-
polyvinylpyrrolidone	2.7	-
lecithin	1.5	-
benzyl alcohol	0.01	-
magnesium aluminum silicate	-	25
flavor	-	q.s.
color	-	q.s.
methyl paraben, U.S.P	-	3
propyl paraben, U.S.P.	-	0.7
polysorbate 80 (e.g., Tween 80),	U.S.P. -	5
sorbitol solution, 70%, U.S.P.	-	1450
buffer agent to adjust pH for desired stability	q.s. for injection	q.s.
water	q.s. to 1 ml	q.s. to 5 ml

The preferred pharmaceutical compositions from the standpoint of preparation and ease of administration are solid compositions, particularly liquid or hard-filled capsules and tablets containing from about 10 to 100 mg of the active ingredient concerning the PAF inhibition use and from about 350 to 450 mg of the active ingredient with respect to tumor inhibition.

Such compositions also form part of the invention.

The following examples, in which all temperatures are in °C illustrate the invention.

Example 1

5-(p-fluorophenyl)-2,3-dihydroimidazo-[2,1-a]isoquinoline ($R_1 = H$, $R_2 = p$ -fluorophenyl).

a) 5-(p-fluorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a] isoquinoline-5-ol (Compound of formula V in wherein $R_1 = H$, $R_2 = p$ -fluorophenyl).

Into a flask, maintained under nitrogen and equipped with a stirrer, condenser, and dropping funnel, are charged 6.26 g (0.04 mol) of 2-(o-tolyl)-2(1H)-imidazoline and 100 ml of dry tetrahydrofuran. The solution is stirred, treated with 84 ml of 1.6M n-butyl lithium (0.13 mol) in hexane and maintained at 35°C for ca. 5 hours. The mixture is treated with 20.7 g (0.16 mol) of ethyl-p-fluorobenzoate in 50 ml of dry tetrahydrofuran and then maintained at 50°C for ca. 5 hours. The reaction is, then cooled in an icebath and treated with

22.8 ml of saturated ammonium chloride. The tetrahydrofuran layer is separated in a separatory funnel, dried with magnesium sulfate, filtered and then concentrated in vacuo. The residue is dissolved in a hot methylene chloride-methanol (1:1) mixture which, upon cooling, yields the desired compound;

5 b) 5-(p-fluorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline

A solution containing 2.8 g (0.01 mol) of the compound prepared in a) above and 40 ml of glacial acetic acid is refluxed, with stirring, under a nitrogen atmosphere for ca. 5 hours. The acetic acid is then removed by evaporation in vacuo and the residue treated with 50 ml of water and then made alkaline with 35 ml of 2N sodium carbonate. The mixture is then extracted with two 100 ml portions of methylene chloride, after which time the organic layer is separated, dried over anhydrous magnesium sulfate, filtered and the solvent removed in vacuo. The crude product obtained is recrystallized employing a mixture of methanol and water to yield the title compound, m.p. 150°-151° C.

15 Example 2

Following essentially the procedure of Example 1a) above, and using in place of the ethyl-p-fluorobenzoate, the appropriate compound of formula IV, the following compounds of formula V are obtained:

- A) 5-[4-diethylamino)methylphenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- B) 5-(p-chlorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- C) 5-phenyl-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- D) 5-(3,4-dimethoxyphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- E) 5-(4-methoxyphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- F) 5-(3,4-methylenedioxyphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 25 G) 5-(3,4-dichlorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- H) 5-(2,4-dichlorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- I) 5-(3-trifluoromethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- J) 5-(2-chlorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- K) 5-(4-methylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 30 L) 5-(3-chlorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- M) 5-(2-fluorophenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- N) 5-(4-piperidinomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- O) 5-(4-pyrrolidinomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- P) 5-(4-morpholinomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 35 Q) 5-(2-thienyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- R) 5-[4'-(2-chloro-4-fluorobenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- S) 5-[4'-(2,6-dichlorobenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- T) 5-[4'-(4-t-butylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 40 U) 5-[4'-(phenylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- V) 5-[4'-(dibenzylaminomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- W) 5-[4'-(dicyclohexylaminomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- X) 5-[4'-(diisopropylaminomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- Y) 5-[4'-(ethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- Z) 5-[4'-(trimethylsilylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 45 AA) 5-[4'-(phenoxyphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- BB) 5-[4'-(diallylaminomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- CC) 5-[4'-(benzyloxyphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- DD) 5-(2'-furyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- EE) 5-[4'-(thiomorpholinomethylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 50 FF) 5-(3'-t-butylphenyl)-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- GG) 5-[4'-(2,6-dichlorophenyloxymethyl)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- HH) 5-[4'-(3,4,5-trimethoxybenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- II) 5-[4'-(2-fluorobenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- JJ) 5-[4'-(4-fluorobenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 55 KK) 5-[4'-(2-chlorobenzyloxy)phenyl]-2,3,4,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- LL) 5-[4'-(2-chloro-6-fluorobenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- MM) 5-[4'-(2,4-dichlorobenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- NN) 5-[4'-(4-t-butylbenzyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;

OO) 5-[4'-(3,4,5-trimethoxyphenylpropyloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
 PP) 5-[4'-(3,4,5-trimethoxyphenylhexyl)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol; and
 QQ) 5-[4'-(3,4,5-trimethoxyphenylethylloxy)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol, respectively.

Example 3

Following essentially the procedure of Example 1b, and using in place of the compound of Example 1a), a suitable amount of a compound of Examples 2A) through 2NN), respectively, there are obtained:

- A) 5-[4'-(diethylamino)methylphenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. 292° - 294° C;
- B) 5-(p-chlorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >250° C;
- C) 5-phenyl-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 143° - 145° C;
- D) 5-(3',4'-dimethoxyphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p., 158° - 160° C;
- E) 5-(4'-methoxyphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 126° - 128° C;
- F) 5-(3',4'-methylenedioxyphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 128° - 130° C;
- G) 5-(3',4'-dichlorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 158° - 160° C;
- H) 5-(2',4'-dichlorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 150° - 152° C;
- I) 5-(3'-trifluoromethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >250° C;
- J) 5-(2'-chlorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 134° - 136° C;
- K) 5-(p-methylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >250° C;
- L) 5-(3'-chlorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >250° C;
- M) 5-(2'-fluorophenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >250° C;
- N) 5-(4'-piperidinomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. 317° - 320° C;
- O) 5-[4'-pyrrolidinomethylphenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 106° - 108° C;
- P) 5-(4'-morpholinomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 160° - 162° C;
- Q) 5-[2-thienyl]-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. 98° - 100° C;
- R) 5-[4'-(2-chloro-4-fluorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 252° - 254° C. (dec.);
- S) 5-[4'-(2,6-dichlorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 277° - 280° C;
- T) 5-(4'-t-butylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >200° C;
- U) 5-(4'-phenylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline, m.p. >200° C;
- V) 5-(4'-dibenzylaminomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. 225° C;
- W) 5-(4'-dicyclohexylaminomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. 273° C (dec.);
- X) 5-(4'-diisopropylaminomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, monohydrate, m.p. 264° C (dec.);
- Y) 5-(4'-ethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 298° - 300° C;
- Z) 5-(4'-trimethylsilylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. >270° C;
- AA) 5-(4'-phenoxyphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. >270° C;
- BB) 5-(4'-diallylaminomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. 228° C;
- CC) 5-(4'-benzyloxyphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 213° - 215° C;
- DD) 5-(2'-furyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. >315° C;
- EE) 5-(4'-thiomorpholinomethylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline dihydrochloride, m.p. 275° - 278° C;
- FF) 5-(3'-t-butylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 290° C;
- GG) 5-[4'-(2,6-dichlorophenylloxymethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 242° C;
- HH) 5-[4'-(3,4,5-trimethoxybenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 205° - 207° C;
- II) 5-[4'-(2-fluorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 253° - 254° C. (dec.);
- JJ) 5-[4'-(4-fluorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 257° - 258° C (dec.);
- KK) 5-[4'-(2-chlorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 272° C. (dec.);

- LL) 5-[4'-(2-chloro-6-fluorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. >250 °C;
- MM) 5-[4'-(2,4-dichlorobenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 254 °C. (dec.);
- 5 NN) 5-[4'-(4-*t*-butylbenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 271 °C. (dec.);
- OO) 5-[4'-(3,4,5-trimethoxyphenylpropyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 215 °-217 °C. (dec.);
- 10 PP) 5-[4'-(3,4,5-trimethoxyphenylhexyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 213 °-215 °C; and
- QQ) 5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 242 °-243 °C, respectively.

Example 4

15 Following essentially the procedure of Example 1a) above, and using in place of 2-(*o*-tolyl)-2(1H)-imidazoline and ethyl-*p*-fluorobenzoate, appropriate compounds of formulae III and IV, there are obtained:

- A) 5-(4'-*t*-butylphenyl)-8-chloro-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- B) 5-(4'-*i*-butylphenyl)-9-chloro-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- C) 5-(4'-*i*-butylphenyl)-9-methyl-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- D) 8-chloro-5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol; and
- E) 2,2-dimethyl-5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol, respectively.

Example 5

25 Following essentially the procedure of Example 1b and using in place of the compound of Example 1a), a suitable amount of the compounds of Examples 4A) through 4C), respectively, there are obtained:

- 30 A) 5-(4'-*t*-butylphenyl)-8-chloro-2,3-dihydroimidazo [2,1-a]isoquinoline hydrochloride, m.p. >250 °C;
- B) 5-(4'-*i*-butylphenyl)-9-chloro-2,3-dihydroimidazo [2,1-a]isoquinoline hydrochloride, m.p. >298 °C;
- C) 5-(4'-*i*-butylphenyl)-9-methyl-2,3-dihydroimidazo [2,1-a]isoquinoline hydrochloride, m.p. >275 °C;
- D) 8-chloro-5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride m.p. 268 ° (dec); and
- 35 E) 2,2-dimethyl-5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, respectively.

Example 6

40 Following essentially the procedure of Example 1a) above, and using in place of the ethyl-*p*-fluorobenzoate, the appropriate compound of formula IV, there are obtained:

- A) 5-[4'-(3,4,5-trimethoxybenzyloxymethyl)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- B) 5-[4'-(3,4-dimethoxybenzyloxy)phenyl]-2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 45 C) 5-[3'-(3,4,5-trimethoxybenzyloxy)phenyl]-2,3,5, 6-tetrahydroimidazo [2,1-a]isoquinolin-5-ol;
- D) 5-[2'-(3,4,5-trimethoxybenzyloxy)phenyl] -2,3,5,6-tetrahydroimidazo [2,1-a]isoquinolin-5-ol;
- E) 5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]2,3,5, 6-tetrahydroimidazo [2,1-a]isoquinolin-5-ol;
- F) 5-[3'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3,5, 6-tetrahydroimidazo [2,1-a]isoquinolin-5-ol;
- G) 5-[4'-(3,4-dimethoxyphenylethyl)phenyl]2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 50 H) 5-[4'-(2-methoxyphenylethyl)phenyl]-2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- I) 5-[4'-(4-methoxybenzyloxy)phenyl]-2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- J) 5-[4'-(3,4-dimethylbenzyloxy)phenyl]-2,3,5, 6-tetrahydroimidazo [2,1-a]isoquinolin-5-ol;
- K) 5-[4'-(3,4,5-trimethoxyphenylpropyl)phenyl]-2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- L) 5-[4'-(2,3,4-trimethoxybenzyloxy)phenyl]-2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol;
- 55 M) 5-[4'-(3-methoxy-4-pentoxypheylethyl)phenyl]-2,3,5,6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol; and
- N) 5-[4'-(3,4,5-trimethoxybenzyl)phenyl]-2,3,5, 6-tetrahydroimidazo[2,1-a]isoquinolin-5-ol; respectively.

Example 7

Following essentially the procedure of Example 1b and using in place of the compound of Example 1a), a suitable amount of the compounds of Examples 6A) through 6N), respectively, there are obtained:

A) 5-[4'-(3,4,5-trimethoxybenzyloxymethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $>240^{\circ}\text{C}(\text{dec.})$;

B) 5-[4'-(3,4-dimethoxybenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, monohydrate, m.p. $>239^{\circ}\text{C}(\text{dec.})$;

C) 5-(3'-(3,4,5-trimethoxybenzyloxy)phenyl)-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 230°C ;

D) 5-[2'-(3,4,5-trimethoxybenzyloxy)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 252°C ;

E) 5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2, 3-dihydroimidazo [2,1-a]isoquinoline hydrochloride, m.p. $243^{\circ} - 245^{\circ}\text{C}$;

F) 5-[3'-(3,4,5-trimethoxyphenylethyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $260^{\circ} - 261^{\circ}\text{C}$;

G) 5-[4'-(3,4-dimethoxyphenylethyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $263^{\circ} - 265^{\circ}\text{C}$;

H) 5-[4'-(2-methoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $>270^{\circ}\text{C}$;

I) 5-[4'-(4-methoxybenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $>250^{\circ}\text{C}$;

J) 5-[4'-(3,4-dimethylbenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $>270^{\circ}\text{C}$;

K) 5-[4'-(3,4,5-trimethoxyphenylpropyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $242^{\circ} - 244^{\circ}\text{C}$;

L) 5-[4'-(2,3,4-trimethoxybenzyloxy)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $>240^{\circ}\text{C}$;

M) 5-(4'-(3-methoxy-4-pentoxyphenylethyl)phenyl)-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $260^{\circ}\text{C}(\text{dec.})$; and


N) 5-[4'-(3,4,5-trimethoxybenzyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. $238^{\circ} - 240^{\circ}\text{C}$,

respectively.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Use of a compound of formula I



wherein each R independently is hydrogen or methyl R₁ is in 8- or 9-position and represents hydrogen, chloro, or C₁₋₃alkyl and R₂ represents thienyl, furyl or a group



wherein each Ro represents independently hydrogen, fluoro, chloro, C₁₋₁₀alkyl or C₁₋₁₀alkoxy or one Ro is hydrogen and the other represents

a) tri-(C₁₋₃alkyl)silyl, trifluoromethyl or phenyl;



Following essentially the procedure of Example 1b and using in place of the compound of Example 1a), a suitable amount of the compounds of Examples 6A) through 6N), respectively, there are obtained:

A) 5-[4'-(3,4,5-trimethoxybenzyloxymethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. >240° C(dec.);

B) 5-[4'-(3,4-dimethoxybenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, monohydrate, m.p.>239° C(dec.);

C) 5-[3'-(3,4,5-trimethoxybenzyloxy)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 230° C;

D) 5-[2'-(3,4,5-trimethoxybenzyloxy)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 252° C ;

E) 5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2, 3-dihydroimidazo [2,1-a]isoquinoline hydrochloride, m.p. 243° - 245° C ;

F) 5-[3'-(3,4,5-trimethoxyphenylethyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 260° - 261° C ;

G) 5-[4'-(3,4-dimethoxyphenylethyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 263° - 265° C;

H) 5-[4'-(2-methoxyphenylethyl)phenyl]2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p.>270° C ;

I) 5-[4'-(4-methoxybenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p.>250° C ;

J) 5-[4'-(3,4-dimethylbenzyloxy)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p.>270° C ;

K) 5-[4'-(3,4,5-trimethoxyphenylpropyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 242° - 244° C ;

L) 5-[4'-(2,3,4-trimethoxybenzyloxy)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. >240° C ;

M) 5-[4'-(3-methoxy-4-pentoxyphenylethyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 260° C(dec.); and


N) 5-[4'-(3,4,5-trimethoxybenzyl)phenyl]-2, 3-dihydroimidazo[2,1-a]isoquinoline hydrochloride, m.p. 238° - 240° C,

respectively.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Use of a compound of formula I



wherein each R independently is hydrogen or methyl R₁ is in 8- or 9-position and represents hydrogen, chloro, or C₁₋₃alkyl and R₂ represents thienyl, furyl or a group




wherein each Ro represents independently hydrogen, fluoro, chloro, C₁₋₁₀alkyl or C₁₋₁₀alkoxy or one Ro is hydrogen and the other represents

a) tri-(C₁₋₃alkyl)silyl, trifluoromethy or phenyl;




b) a group



wherein each R_3 represents independently hydrogen, chloro, fluoro C_{1-4} alkyl or C_{1-5} alkoxy and X represents

$-CH_2O-$, $-OCH_2-$, $-CH_2-$, $-CH_2CH_2-$, or $-O-$;


c) a group



wherein each R_3' represents independently C_{1-3} alkoxy and

Y represents $-(OCH_2)_{1-5}-$, $-(CH_2)_{1-6}-$ or $-CH_2OCH_2-$; or

d) a group




wherein each R_4 represents independently C_{1-4} alkyl, C_{5-7} cycloalkyl, allyl or benzyl; or the two R_4 's together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl or thiomorpholinyl;

or two R_4 's on adjacent carbon atoms form methylenedioxy, with the proviso that when an R_4 is other than chloro, fluoro, methyl or methoxy it may only be in a meta or para position or a pharmaceutically acceptable acid addition salt thereof in the manufacture of a medicament for use in inhibiting PAF mediated bronchoconstriction and extravasation and PAF mediated endotoxin induced lung injury


2. The use according to claim 1 whereby in formula I each R is hydrogen, R_1 is as defined in claim 1 and one R_0 is hydrogen and the other is in meta- or para-position and is as defined under b) or c) in claim 1.

3. Use of a compound of formula Ib




Ib

wherein R and R_1 are as defined in claim 1 and R_2''' represents thienyl, furyl or a group



wherein each Ro^{iv} represents independently chloro, fluoro or C_{1-6} alkyl or one Ro^{iv} is hydrogen and the other is in meta or para position and represents C_{2-6} alkyl, tri- C_{1-3} alkyl)silyl, phenyl, or is as defined under b) or c) in claim 1 or is

d)' a group




wherein each R_4' is allyl or the two R_4' 's together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl, thiomorpholinyl with the proviso that when Ro^{iv} is other than chloro, fluoro or methyl it may be only in meta- or para- position or a pharmaceutically acceptable acid addition salt, in the manufacture of a medicament for use in treating tumors.

4. The use according to claim 3 whereby in formula Ib R and R_1 are as defined in claim 1 and R_2''' represents thienyl, furyl or a group




wherein Ro^v is in meta or para position and represents C_{2-6} alkyl, tri(C_{1-3} alkyl)silyl, phenyl, is as defined under b) or c) in claim 1 or represents

d)'' a group




wherein each R_4'' is allyl or the two R_4'' 's together with the nitrogen atom to which they are attached represent pyrrolidinyl or piperidyl.

5. A compound of formula Ia



Ia

wherein R and R_1 are as defined in claim 1 and R_2' represents thienyl, furyl or a group




wherein one Ro'' is hydrogen and the other represents tri- C_{1-3} alkyl)silyl or phenyl or is as defined under b), c) or d) in claim 1 whereby Ro'' substituents may only be in meta or para positions, or a pharmaceutically acceptable acid addition salt thereof.

6. A compound according to claim 5 wherein each R is hydrogen, R_1 is in 8-position and is as defined in claim 1 and R_2 represents thienyl, furyl or a group




wherein Ro''' is in meta or para position and represents tri- $(C_{1-3}$ alkyl)silyl or is as defined under b), c) or d) in claim 1.


7. A compound of formula Ia according to claim 5 which is 5-[4'-piperidinomethylphenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline; or a pharmaceutically acceptable addition salt thereof.
8. A compound of formula Ia according to claim 5 which is 5-[4'-(3,4,5-trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline; or a pharmaceutically acceptable acid addition salt thereof.
9. A compound of formula I as defined in claim 1 or of formula Ib as defined in claim 3 which 5-(4'-t-butylphenyl)-2,3-dihydroimidazo[2,1-a]isoquinoline; or a pharmaceutically acceptable acid addition salt thereof.
10. The use according to claim 1 or 3 of a compound according to claim 9
11. A pharmaceutical composition comprising a compound according to any one of claims 5 to 9 together with a pharmaceutically acceptable diluent or carrier.
12. A compound according to any one of claims 5 to 9 for use as a pharmaceutical.
13. A process for preparing a compound of formula Ia as wherein R and R_1 are as defined in claim 1 and R_2' is as defined in claim 5 or a pharmaceutically acceptable acid addition salt thereof which comprises dehydrating the corresponding compound of formula V




wherein R and R_1 are as defined in claim 1 and R_2 has the significance indicated in claim 5 for R_2' and recovering the compound obtained in free form or in the form of a pharmaceutically acceptable acid addition salt.

14. A compound of formula Va






wherein each R_3' represents independently C_1-3 alkoxy and
Y represents $-(\text{OCH}_2)_{1-5}-$, $-(\text{CH}_2)_{1-6}-$ or $-\text{CH}_2\text{OCH}_2-$; or
d) a group




wherein each R_4 represents independently C_1-4 alkyl, C_5-7 cycloalkyl, allyl or benzyl; or the two R_4 's together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl or thiomorpholinyl;
or two R_4 's on adjacent carbon atoms form methylenedioxy, with the proviso that when an R_4 is other than chloro, fluoro, methyl or methoxy it may only be in a meta or para position
or a pharmaceutically acceptable acid addition salt thereof, with a pharmaceutically acceptable carrier or diluent.

2. A process of manufacture of a medicament for use in treating tumors which comprises mixing a compound of formula Ib




Ib

wherein R and R_1 are as defined in claim 1 and
 R_2'' represents thienyl, furyl or a group wherein each R_4^{IV} represents independently chloro, fluoro or C_1-6 alkyl or one R_4^{IV} is hydrogen and the other is C_2-6 alkyl, tri- $(\text{C}_1-3$ alkyl)silyl, phenyl, or is as defined under b) or c) in claim 1 or is
d)' a group




wherein each R_4' is allyl or the two R_4'' 's together with the nitrogen atom to which they are attached represent pyrrolidinyl, piperidyl, morpholinyl or thiomorpholinyl, R_4^{IV} is other than, with the proviso that when R_4^{IV} is other than chloro, fluoro or methyl it may be only in meta- or para- position or a pharmaceutically acceptable acid addition salt thereof, with a pharmaceutically acceptable carrier or diluent.

3. A process of manufacture of a medicament which comprises mixing a compound of formula Ia




wherein R and R₁ are as defined in claim 1 and
R₂' represents thienyl, furyl or a group



wherein one RO'' is hydrogen and the other represents tri-(C₁-₃alkyl)-silyl or phenyl or is as defined under b), c) or d) in claim 1 whereby RO'' substituents may only be in meta or para positions, or a pharmaceutically acceptable acid addition salt thereof, with a pharmaceutically acceptable carrier or diluent.

4. A process for the preparation of a compound of formula Ia as defined in claim 3 or a pharmaceutically acceptable acid addition salt thereof which comprises dehydrating the corresponding compound of formula V wherein R and R₁ are as defined in claim 1 and R₂ has the significance indicated in claim 3 for R₂'




wherein R and R₁ are as defined in claim 1 and R₂ has the significance indicated in claim 3 for R₂' and recovering the compound obtained in free form or in the form of a pharmaceutically acceptable acid addition salt.

5. A process according to claim 1, 3 or 4 wherein the compound is 5-[4'-piperidinomethylphenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline or a pharmaceutically acceptable acid addition salt thereof.
6. A process according to claim 1, 3 or 4 wherein the compound is 5-[4'-(3,4,5-trimethoxyphenylethyl)-phenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline or a pharmaceutically acceptable acid addition salt thereof.
7. A process according to claim 1 or 2 wherein the compound is 5-[4'-t-butylphenyl]-2,3-dihydroimidazo[2,1-a]isoquinoline or a pharmaceutically acceptable acid addition salt thereof.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. L'utilisation d'un composé de formule I



dans laquelle

chaque R

signifie indépendamment l'hydrogène ou un méthyle,

R₁

est en position 8 ou 9 et représente l'hydrogène, le chlore ou un alkyle en C₁-C₃ et

R₂


représente un groupe thiényle, furyle ou un groupe



où chaque Ro représente indépendamment l'hydrogène, le fluor, le chlore, un alkyle en C₁-C₁₀ ou un alcoxy en C₁-C₁₀, ou bien un Ro signifie l'hydrogène et l'autre représente

a) un groupe tri-(alkyl en C₁-C₃)silyle, trifluorométhyle ou phényle,


b) un groupe



où chaque R₃ représente indépendamment l'hydrogène, le chlore, le fluor, un alkyle en C₁-C₄ ou un alcoxy en C₁-C₅ et

X représente -CH₂O-, -OCH₂-, -CH₂-, -CH₂CH₂-, ou -O-,


c) un groupe



où chaque R₃' représente indépendamment un alcoxy en C₁-C₃ et

Y représente -(OCH₂)₁₋₅-, -(CH₂)₁₋₆- ou -CH₂OCH₂- ou

d) un groupe




où chaque R₄ représente indépendamment un groupe alkyle en C₁-C₄, cycloalkyle en C₅-C₇, allyle ou benzyle, ou bien les deux R₄ représentent ensemble, avec l'atome d'azote auquel ils sont fixés, un groupe pyrrolidinyle, pipéridyle, morpholinyle ou thiomorpholinyle;
ou bien les deux Ro sur des atomes de carbone adjacents forment un groupe méthylènedioxy, avec la condition que lorsque un Ro a une signification autre que le chlore, le fluor, méthyle ou




méthoxy, il ne puisse être qu'en position méta ou para,
ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé, pour la préparation d'un
médicament destiné à inhiber la bronchoconstriction et l'extravasation provoquées par le PAF et les
lésions pulmonaires induites par des endotoxines par l'intermédiaire du PAF.

2. L'utilisation selon la revendication 1, caractérisée en ce que, dans la formule I, chaque R signifie l'hydrogène, R₁ est tel que défini à la revendication 1 et un R₀ signifie l'hydrogène et l'autre est en position méta ou para et est tel que défini sous b) ou c) à la revendication 1.


3. L'utilisation d'un composé de formule Ib



dans laquelle R et R₁ sont tels que définis à la revendication 1 et R₂''' représente un groupe thiényle, furyle ou un groupe



où
chaque R₀^{iv} représente indépendamment le chlore, le fluor ou un alkyle en C₁-C₆, ou bien un R₀^{iv} signifie l'hydrogène et l'autre est en position méta ou para et représente un groupe alkyle en C₂-C₆, tri(alkyl en C₁-C₃)silyle, phényle ou est tel que défini sous b) ou c) à la revendication 1, ou signifie d) un groupe



où
chaque R₄' signifie un groupe allyle ou bien les deux R₄' représentent ensemble, avec l'atome d'azote auquel ils sont fixés, un groupe pyrrolidinyne, pipéridyle, morpholinyne, thiomorpholinyne, avec la condition que lorsque R₀^{iv} a une signification autre que le chlore, le fluor ou méthyle, il ne puisse être qu'en position méta ou para,
ou un sel d'addition pharmaceutiquement acceptable de ce composé, pour la préparation d'un médicament destiné au traitement des tumeurs.


4. L'utilisation selon la revendication 3, caractérisée en ce que, dans la formule Ib, R et R₁ sont tels que définis à la revendication 1 et R₂''' représente un groupe thiényle, furyle ou un groupe



où R₀^v est en position méta ou para et représente un groupe alkyle en C₂-C₆, tri(alkyl en C₁-C₃)silyle,




phényle, est tel que défini sous b) ou c) à la revendication 1 ou représente d)" un groupe



où chaque R_4'' signifie un groupe allyle ou bien les deux R_4'' représentent ensemble, avec l'atome d'azote auquel ils sont fixés, un groupe pyrrolidinyne ou pipéridyle.

5. Un composé de formule la



dans laquelle

R et R_1 sont tels que définis à la revendication 1 et R_2' représente un groupe thiényne, furyne ou un groupe



où un Ro'' signifie l'hydrogène et l'autre représente un groupe tri-(alkyl en $\text{C}_1\text{-C}_3$)silyle ou phényle ou est tel que défini sous b), c) ou d) à la revendication 1, les substituants Ro'' ne pouvant être qu'en position méta ou para, ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé.

6. Un composé selon la revendication 5, dans lequel chaque R signifie l'hydrogène, R_1 est en position 8 et est tel que défini à la revendication 1 et R_2 représente un groupe thiényne, furyne ou un groupe



où Ro''' est en position méta ou para et représente un groupe tri(alkyl en $\text{C}_1\text{-C}_3$)silyle ou est tel que défini sous b), c) ou d) à la revendication 1.

7. Un composé de formule la selon la revendication 5, qui est la 5-[4'-pipéridinométhylphényl]-2,3-dihydro-imidazo[2,1-a]isoquinoléine, ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé.

8. Un composé de formule la selon la revendication 5, qui est la 5-[4'-(3,4,5-triméthoxyphényléthyl)-phényl]-2,3-dihydro-imidazo[2,1-a]isoquinoléine, ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé.

9. Un composé de formule I tel que défini à la revendication 1 ou de formule Ib tel que défini à la revendication 3, qui est la 5-[4'-tert.-butylphényl]-2,3-dihydro-imidazo[2,1-a]isoquinoléine, ou un sel




d'addition d'acide pharmaceutiquement acceptable de ce composé.

10. L'utilisation selon la revendication 1 ou 3 d'un composé selon la revendication 9.

11. Une composition pharmaceutique, comprenant un composé selon l'une quelconque des revendications 5 à 9 ensemble avec un diluant ou véhicule pharmaceutiquement acceptable.


12. Un composé selon l'une quelconque des revendications 5 à 9, pour l'utilisation comme médicament.

13. Un procédé de préparation d'un composé de formule la où R et R₁ sont tels que définis à la revendication 1 et R₂' est tel que défini à la revendication 5, ou d'un sel d'addition d'acide pharmaceutiquement acceptable de ce composé, qui comprend la déshydratation du composé correspondant de formule V



25 dans laquelle R et R₁ sont tels que définis à la revendication 1 et R₂ a la signification indiquée à la revendication 5 pour R₂', et la récupération du composé obtenu sous forme libre ou sous forme d'un sel d'addition d'acide pharmaceutiquement acceptable.


14. Un composé de formule Va



45 dans laquelle R et R₁ sont tels que définis à la revendication 1 et Ro" est tel que défini à la revendication 5.

Revendications pour les Etats contractants suivants : ES, GR

1. Un procédé de préparation d'un médicament destiné à inhiber la bronchoconstriction et l'extravasation provoquées par le PAF et les lésions pulmonaires induites par des endotoxines par l'intermédiaire du PAF, qui comprend le mélange d'un composé de formule I



dans laquelle

chaque R

R₁

R₂

signifie indépendamment l'hydrogène ou un méthyle,

est en position 8 ou 9 et représente l'hydrogène, le chlore ou un alkyle en C₁-C₃ et


signifie un groupe thiényle, furyle ou un groupe



où chaque Ro représente indépendamment l'hydrogène, le fluor, le chlore, un alkyle en C₁-C₁₀ ou un alcoxy en C₁-C₁₀, ou bien un Ro signifie l'hydrogène et l'autre représente

a) un groupe tri-(alkyle en C₁-C₃)silyle, trifluorométhyle ou phényle,


b) un groupe



où chaque R₃ représente indépendamment l'hydrogène, le chlore, le fluor, un alkyle en C₁-C₄ ou un alcoxy en C₁-C₅ et

X représente -CH₂O-, -OCH₂-, -CH₂-, -CH₂CH₂-, ou -O-,


c) un groupe



où chaque R₃' représente indépendamment un alcoxy en C₁-C₃ et

Y représente -(OCH₂)₁₋₅-, -(CH₂)₁₋₅- ou -CH₂OCH₂- ou

d) un groupe




où chaque R₄ représente indépendamment un groupe alkyle en C₁-C₄, cycloalkyle en C₅-C₇, allyle ou benzyle, ou bien les deux R₄ représentent ensemble, avec l'atome d'azote auquel ils sont fixés,



un groupe pyrrolidinyle, pipéridyle, morpholinyle ou thiomorpholinyle;
 ou bien les deux Ro sur des atomes de carbone adjacents forment un groupe méthylènedioxy,
 avec la condition que lorsque Ro a une signification autre que le chlore, le fluor, méthyle ou méthoxy, il
 ne puisse être qu'en position méta ou para,
 ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé,
 avec un véhicule ou diluant pharmaceutiquement acceptable.


2. Un procédé de préparation d'un médicament pour l'utilisation dans le traitement des tumeurs, qui comprend le mélange d'un composé de formule Ib



dans laquelle R et R₁ sont tels que définis à la revendication 1 et R₂''' représente un groupe thiényl, furyl ou un groupe




où
 chaque Ro^{iv} représente indépendamment le chlore, le fluor ou un alkyle en C₁-C₆, ou bien un Ro^{iv} signifie l'hydrogène et l'autre signifie un groupe alkyle en C₂-C₆, tri(alkyl en C₁-C₃)silyl, phényle, ou est tel que défini sous b) ou c) à la revendication 1 ou signifie
 d) un groupe



où
 chaque R₄' signifie un groupe allyle ou bien les deux R₄' représentent ensemble, avec l'atome d'azote auquel ils sont fixés, un groupe pyrrolidinyle, pipéridyle, morpholinyle ou thiomorpholinyle, avec la condition que lorsque Ro^{iv} a une signification autre que le chlore, le fluor ou méthyle, il ne puisse être qu'en position méta ou para,
 ou un sel d'addition pharmaceutiquement acceptable de ce composé,
 avec un véhicule ou diluant pharmaceutiquement acceptable.

3. Un procédé de préparation d'un médicament, qui comprend le mélange d'un composé de formule Ia





dans laquelle


R et R₁ sont tels que définis à la revendication 1 et R₂' représente un groupe thiényle, furyle ou un groupe



où un Ro'' signifie l'hydrogène et l'autre représente un groupe tri-(alkyl en C₁-C₃)silyle ou phényle ou est tel que défini sous b), c) ou d) à la revendication 1, les substituants Ro'' ne pouvant être qu'en position méta ou para,

ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé, avec un véhicule ou diluant pharmaceutiquement acceptable.

4. Un procédé de préparation d'un composé de formule la tel que défini à la revendication 3 ou d'un sel d'addition d'acide pharmaceutiquement acceptable de ce composé, qui comprend la déshydratation du composé correspondant de formule V




dans laquelle R et R₁ sont tels que définis à la revendication 1 et R₂ a la signification indiquée à la revendication 3 pour R₂', et la récupération du composé obtenu sous forme libre ou sous forme d'un sel d'addition d'acide pharmaceutiquement acceptable.

5. Un procédé selon la revendication 1, 3 ou 4, dans lequel le composé est la 5-[4'-pipéridinométhylphényl]-2,3-dihydro-imidazo[2,1-a]isoquinoléine, ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé.
6. Un procédé selon la revendication 1, 3 ou 4, dans lequel le composé est la 5-[4'-(3,4,5-triméthoxyphényléthyl)phényl]-2,3-dihydro-imidazo[2,1-a]isoquinoléine, ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé.
7. Un procédé selon la revendication 1 ou 2, dans lequel le composé est la 5-[4'-tert.-butylphényl]-2,3-dihydro-imidazo[2,1-a]isoquinoléine, ou un sel d'addition d'acide pharmaceutiquement acceptable de ce composé.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Verwendung einer Verbindung der Formel I






worin jeder Rest R unabhängig Wasserstoff oder ein Methylrest ist,
 R_1 in 8- oder 9-stellung ist und Wasserstoff, Chlor oder einen C_1 - C_3 -Alkylrest bedeutet und
 R_2 einen Thienylrest, Furylrest oder eine Gruppe




bedeutet, worin jeder Rest R_o unabhängig Wasserstoff, Fluor, Chlor, einen C_1 - C_{10} -Alkyl- oder C_1 - C_{10} -Alkoxyrest bedeutet oder worin ein Rest R_o Wasserstoff ist und der andere

- a) einen Tri- $(C_1$ - C_3 -alkyl)silyl-, Trifluormethyl- oder Phenylrest bedeutet;
- b) eine Gruppe



bedeutet, worin jeder Rest R_3 unabhängig Wasserstoff, Chlor, Fluor, einen C_1 - C_4 -Alkyl- oder C_1 - C_5 -Alkoxyrest bedeutet und X $-CH_2O-$, $-OCH_2-$, $-CH_2-$, $-CH_2CH_2-$ oder $-O-$ bedeutet;

- c) eine Gruppe



bedeutet, worin jeder Rest R_3' unabhängig einen C_1 - C_3 -Alkoxyrest bedeutet und Y $-(OCH_2)_{1-5}-$, $-(CH_2)_{1-6}-$ oder $-CH_2OCH_2-$ bedeutet; oder

- d) eine Gruppe




bedeutet, worin jeder Rest R_4 unabhängig einen C_1 - C_4 -Alkyl-, C_5 - C_7 -Cycloalkyl-, Allyl- oder Benzylrest bedeutet oder worin die zwei Reste R_4 zusammen mit dem Stickstoffatom, an das sie gebunden sind, eine Pyrrolidinyl-, Piperidyl-, Morpholinyl- oder Thiomorpholinylgruppe darstellen oder worin zwei Reste R_o an benachbarten Kohlenstoffatomen einen Methylendioxyrest bilden, mit dem Vorbehalt, daß dann, wenn ein Rest R_o etwas anderes als Chlor, Fluor, einen Methyl- oder Methoxyrest bedeutet, er nur in meta- oder para-Stellung sein kann, oder ein pharmazeutisch annehmbares Säureadditionssalz davon zur Herstellung eines Arzneimittels zur Verwendung zur Hemmung von durch PAF vermittelte Bronchienverengung und Extravasation und durch PAF vermittelte endotoxininduzierte Lungenschädigung.

2. Verwendung nach Anspruch 1, wobei in Formel I jeder Rest R Wasserstoff ist, R_1 wie in Anspruch 1 definiert ist und einer der Reste R_o Wasserstoff ist und der andere sich in meta- oder para-Stellung befindet und wie unter b) oder c) in Anspruch 1 definiert ist.



3. Verwendung einer Verbindung der Formel Ib




15
 worin R und R₁ wie in Anspruch 1 definiert sind und
 R₂^{'''} einen Thienylrest, Furylrest oder eine Gruppe



20
 bedeutet, worin jeder Rest Ro^{iv} unabhängig Chlor, Fluor oder einen C₁-C₆-Alkylrest bedeutet oder einer
 der Reste Ro^{iv} Wasserstoff ist und der andere sich in meta- oder para-Stellung befindet und einen C₂-
 C₆-Alkyl-, Tri-(C₁-C₃-alkyl)silyl- oder Phenylrest bedeutet oder wie unter b) oder c) in Anspruch 1
 definiert ist oder

25
 d) eine Gruppe




35
 bedeutet, worin jeder Rest R₄['] ein Allylrest ist oder worin die zwei Reste R₄['] zusammen mit dem
 Stickstoffatom, an das sie gebunden sind, eine Pyrrolidinyl-, Piperidyl-, Morpholinyl-, Thiomorpholi-
 nylgruppe darstellen, mit dem Vorbehalt, daß dann, wenn Ro^{iv} etwas anderes als Chlor, Fluor oder
 eine Methylgruppe ist, es nur in meta- oder para-Stellung sein kann,
 oder ein pharmazeutisch annehmbares Säureadditionssalz zur Herstellung eines Arzneimittels zur
 Verwendung zur Behandlung von Tumoren.

40
 4. Verwendung nach Anspruch 3, worin in Formel Ib R und R₁ wie in Anspruch 1 definiert sind und R₂^{'''}
 einen Thienylrest, Furylrest oder eine Gruppe




50
 darstellt, worin Ro^v in meta- oder para-Stellung ist und einen C₂-C₆-Alkyl-, Tri-(C₁-C₃-alkyl)silyl-,
 Phenylrest bedeutet, wie unter b) oder c) in Anspruch 1 definiert ist oder
 d) eine Gruppe



bedeutet, worin jeder Rest R_4'' ein Allylrest ist oder worin die zwei Reste R_4'' zusammen mit dem Stickstoffatom, an das sie gebunden sind, eine Pyrrolidiny- oder Piperidylgruppe darstellen.

5. Verbindung der Formel Ia



worin R und R_1 wie in Anspruch 1 definiert sind und R_2' einen Thienylrest, Furylrest oder eine Gruppe



bedeutet, worin ein Rest Ro'' Wasserstoff ist und der andere einen Tri-(C_1 - C_3 -alkyl)silyl- oder Phenylrest bedeutet oder wie unter b), c) oder d) in Anspruch 1 definiert ist, wobei sich die Ro'' -Substituenten nur in meta- oder para-Stellung befinden können, oder ein pharmazeutisch annehmbares Säureadditionssalz davon.

6. Verbindung nach Anspruch 5, worin jeder Rest R Wasserstoff ist, R_1 sich in 8-Stellung befindet und wie in Anspruch 1 definiert ist und R_2 einen Thienylrest, Furylrest oder eine Gruppe



bedeutet, worin Ro'' sich in meta- oder para-Stellung befindet und einen Tri-(C_1 - C_3 -alkyl)silylrest bedeutet oder wie unter b), c) oder d) in Anspruch 1 definiert ist.

7. Verbindung der Formel Ia nach Anspruch 5, die 5-[4'-Piperidinomethylphenyl]-2,3-dihydroimidazo[2,1-a]isochinolin oder ein pharmazeutisch annehmbares Säureadditionssalz davon ist.

8. Verbindung der Formel Ia nach Anspruch 5, die 5-[4'-(3,4,5-Trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isochinolin oder ein pharmazeutisch annehmbares Säureadditionssalz davon ist.

9. Verbindung der Formel I wie in Anspruch 1 definiert oder der Formel Ib wie in Anspruch 3 definiert, die 5-(4'-t-Butylphenyl)-2,3-dihydroimidazo[2,1-a]isochinolin oder ein pharmazeutisch annehmbares Säureadditionssalz davon ist.


10. Verwendung nach Anspruch 1 oder 3 einer Verbindung nach Anspruch 9.

11. Pharmazeutische Zusammensetzung enthaltend eine Verbindung nach einem der Ansprüche 5 bis 9 zusammen mit einem pharmazeutisch annehmbaren Verdünnungsmittel oder Träger.




12. Verbindung nach einem der Ansprüche 5 bis 9 zur Verwendung als Pharmazeutikum.

13. Verfahren zur Herstellung einer Verbindung der Formel Ia, worin R und R₁ wie in Anspruch 1 definiert sind und R₂' wie in Anspruch 5 definiert ist oder eines pharmazeutisch annehmbaren Säureadditionssalzes davon, umfassend, daß man die entsprechende Verbindung der Formel V



worin R und R₁ wie in Anspruch 1 definiert sind und R₂ die in Anspruch 5 für R₂' angegebene Bedeutung hat, dehydratisiert und die in freier Form oder in Form eines pharmazeutisch annehmbaren Säureadditionssalzes erhaltene Verbindung gewinnt.


14. Verbindung der Formel Va



worin R und R₁ wie in Anspruch 1 definiert sind und Ro'' wie in Anspruch 5 definiert ist.

Patentansprüche für folgende Vertragsstaaten : ES, GR


1. Verfahren zur Herstellung eines Arzneimittels zur Verwendung zur Hemmung von durch PAF vermittelter Bronchienverengung und Extravasation und PAF vermittelter endotoxininduzierter Lungenschädigung, umfassend, daß man eine Verbindung der Formel I




worin jeder Rest R unabhängig Wasserstoff oder ein Methylrest ist, R₁ in 8- oder 9-Stellung ist und Wasserstoff, Chlor oder einen C₁-C₃-Alkylrest bedeutet und R₂ einen Thienylrest, Furylrest oder eine Gruppe




bedeutet, worin jeder Rest Ro unabhängig Wasserstoff, Fluor, Chlor, einen C₁-C₁₀-Alkyl- oder C₁-C₁₀-Alkoxyrest bedeutet oder worin ein Rest Ro Wasserstoff ist und der andere
a) einen Tri-(C₁-C₃-alkyl)silyl-, Trifluormethyl- oder Phenylrest bedeutet;
b) eine Gruppe



bedeutet, worin jeder Rest R₃ unabhängig Wasserstoff, Chlor, Fluor, einen C₁-C₄-Alkyl- oder C₁-C₅-Alkoxyrest bedeutet und X -CH₂O-, -OCH₂-, -CH₂-, -CH₂CH₂- oder -O- bedeutet;
c) eine Gruppe




bedeutet, worin jeder Rest R₃' unabhängig einen C₁-C₃-Alkoxyrest bedeutet und Y -(OCH₂)₁₋₅-, -(CH₂)₁₋₆- oder -CH₂OCH₂- bedeutet; oder
d) eine Gruppe



bedeutet, worin jeder Rest R₄ unabhängig einen C₁-C₄-Alkyl-, C₅-C₇-Cycloalkyl-, Allyl- oder Benzylrest bedeutet oder worin die zwei Reste R₄ zusammen mit dem Stickstoffatom, an das sie gebunden sind, eine Pyrrolidiny-, Piperidyl-, Morpholinyl- oder Thiomorpholinylgruppe darstellen oder worin zwei Reste Ro an benachbarten Kohlenstoffatomen einen Methylendioxyrest bilden, mit dem Vorbehalt, daß dann, wenn ein Rest Ro etwas anderes als Chlor, Fluor, einen Methyl- oder Methoxyrest bedeutet, er nur in meta- oder para-Stellung sein kann, oder ein pharmazeutisch annehmbares Säureadditionssalz davon mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel vermischt.

2. Verfahren zur Herstellung eines Arzneimittels zur Verwendung zur Behandlung von Tumoren, umfassend, daß man eine Verbindung der Formel Ib




Ib

worin R und R₁ wie in Anspruch 1 definiert sind und R₂''' einen Thienylrest, Furylrest oder eine Gruppe




bedeutet, worin jeder Rest Ro^{iv} unabhängig Chlor, Fluor oder einen C₁-C₆-Alkylrest bedeutet oder einer der Reste Ro^{iv} Wasserstoff ist und der andere einen C₂-C₆-Alkyl-, Tri-(C₁-C₃-alkyl)silyl- oder Phenylrest bedeutet oder wie unter b) oder c) in Anspruch 1 definiert ist oder d) eine Gruppe



bedeutet, worin jeder Rest R₄' ein Allylrest ist oder worin die zwei Reste R₄' zusammen mit dem Stickstoffatom, an das sie gebunden sind, eine Pyrrolidiny-, Piperidyl-, Morpholiny- oder Thiomorpholinygruppe darstellen, mit dem Vorbehalt, daß dann, wenn Ro^{iv} etwas anderes als Chlor, Fluor oder eine Methylgruppe ist, es nur in meta- oder para-Stellung sein kann, oder ein pharmazeutisch annehmbares Säureadditionssalz davon mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel vermischt.

3. Verfahren zur Herstellung eines Arzneimittels, umfassend, daß man eine Verbindung der Formel Ia



Ia


worin R und R₁ wie in Anspruch 1 definiert sind und R₂' einen Thienylrest, Furylrest oder eine Gruppe



bedeutet, worin ein Rest Ro'' Wasserstoff ist und der andere einen Tri-(C₁-C₃-alkyl)silyl- oder Phenylrest bedeutet oder wie unter b), c) oder d) in Anspruch 1 definiert ist, wobei die Ro''-Substituenten nur in meta- oder para-Stellung sein können, oder ein pharmazeutisch annehmbares Säureadditionssalz davon mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel vermischt.



4. Verfahren zur Herstellung einer Verbindung der Formel Ia wie in Anspruch 3 definiert oder eines pharmazeutisch annehmbaren Säureadditionssalzes davon, umfassend, daß man die entsprechende Verbindung der Formel V



worin R und R₁ wie in Anspruch 1 definiert sind und R₂ die in Anspruch 3 die für R₂' angegebene Bedeutung hat, dehydratisiert und die in freier Form oder in Form eines pharmazeutisch annehmbaren Säureadditionssalzes erhaltene Verbindung gewinnt.

5. Verfahren nach Anspruch 1, 3 oder 4, worin die Verbindung 5-[4'-Piperidinomethylphenyl]-2,3-dihydroimidazo[2,1-a]isochinolin oder ein pharmazeutisch annehmbares Säureadditionssalz davon ist.
6. Verfahren nach Anspruch 1, 3 oder 4, worin die Verbindung 5-[4'-(3,4,5-Trimethoxyphenylethyl)phenyl]-2,3-dihydroimidazo[2,1-a]isochinolin oder ein pharmazeutisch annehmbares Säureadditionssalz davon ist.
7. Verfahren nach Anspruch 1 oder 2, worin die Verbindung 5-[4'-t-Butylphenyl]-2,3-dihydroimidazo[2,1-a]isochinolin oder ein pharmazeutisch annehmbares Säureadditionssalz davon ist.