

embodiment has a higher affinity towards human prostate specific membrane antigen than that of a binding protein which has the sequence set forth in SEQ ID NO: 4. In some cases, the prostate specific membrane antigen binding protein of the above embodiment further has a higher affinity towards cynomolgus prostate specific membrane antigen than that of a binding protein which has the sequence set forth in SEQ ID NO: 4.

[0012] One embodiment provides a method for the treatment or amelioration of prostate cancer, the method comprising administration of the PSMA binding protein comprising complementarity determining regions CDR1, CDR2, and CDR3, wherein (a) the amino acid sequence of CDR1 is as set forth in RFMISX₁YX₂MH (SEQ ID NO: 1); (b) the amino acid sequence of CDR2 is as set forth in X₃INPAX₄X₅TDYAEX₆VKG (SEQ ID NO: 2); and (c) the amino acid sequence of CDR3 is as set forth in DX₇YGY (SEQ ID NO: 3), to a subject in need thereof.

[0013] In some embodiments the PSMA binding protein is a single domain antibody. In some embodiments, said single domain antibody is part of a trispecific antibody.

[0013a] Another embodiment of the invention provides a prostate specific membrane ~~antigen binding protein~~single domain antibody, comprising complementarity determining regions CDR1, CDR2, and CDR3, wherein

- (a) the CDR1 comprises the sequence RFMISEYHMH (SEQ ID NO: 5), the CDR2 comprises the sequence TINPAGTTDYAESVKG (SEQ ID NO: 17), and the CDR3 comprises the sequence DSYGY (SEQ ID NO: 15);
- (b) the CDR1 comprises the sequence RFMISPYSMH (SEQ ID NO: 6), the CDR2 comprises the sequence TINPAGTTDYAESVKG (SEQ ID NO: 17), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18);
- (c) the CDR1 comprises the sequence RFMISEYSMH (SEQ ID NO: 16), the CDR2 comprises the sequence TINPAGQTDYAESVKG (SEQ ID NO: 10), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18);
- (d) the CDR1 comprises the sequence RFMISEYSMH (SEQ ID NO: 16), the CDR2 comprises the sequence TINPAKTTDYAESVKG (SEQ ID NO: 9), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18); or

CLAIMS

WHAT IS CLAIMED IS:

1. A prostate specific membrane ~~antigen-binding protein~~single domain antibody, comprising complementarity determining regions CDR1, CDR2, and CDR3, wherein

- (a) the CDR1 comprises the sequence RFMISEYHMH (SEQ ID NO: 5), the CDR2 comprises the sequence TINPAGTTDYAESVKG (SEQ ID NO: 17), and the CDR3 comprises the sequence DSYGY (SEQ ID NO: 15);
- (b) the CDR1 comprises the sequence RFMISPYSMH (SEQ ID NO: 6), the CDR2 comprises the sequence TINPAGTTDYAESVKG (SEQ ID NO: 17), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18);
- (c) the CDR1 comprises the sequence RFMISEYSMH (SEQ ID NO: 16), the CDR2 comprises the sequence TINPAGQTDYAESVKG (SEQ ID NO: 10), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18);
- (d) the CDR1 comprises the sequence RFMISEYSMH (SEQ ID NO: 16), the CDR2 comprises the sequence TINPAKTTDYAESVKG (SEQ ID NO: 9), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18); or
- (e) the CDR1 comprises the sequence RFMISEYHMH (SEQ ID NO: 5), the CDR2 comprises the sequence DINPAGTTDYAESVKG (SEQ ID NO: 8), and the CDR3 comprises the sequence DSYGY (SEQ ID NO: 15).

2. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1, wherein said protein comprises the following formula:

$$f1-r1-f2-r2-f3-r3-f4$$

wherein, r1 is the CDR1; r2 is the CDR2; and r3 is the CDR3; and wherein f₁, f₂, f₃ and f₄ are framework residues selected so that said protein is at least eighty percent identical to the amino acid sequence set forth in SEQ ID NO: 4.

3. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1 or 2, wherein the CDR1 comprises the sequence RFMISEYHMH (SEQ ID NO: 5), the CDR2 comprises the sequence TINPAGTTDYAESVKG (SEQ ID NO: 17), and the CDR3 comprises the sequence DSYGY (SEQ ID NO: 15).

4. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1 or 2, wherein the CDR1 comprises the sequence RFMISPYSMH (SEQ ID NO: 6), the CDR2 comprises the sequence TINPAGTTDYAESVKG (SEQ ID NO: 17), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18).

5. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1 or 2, wherein the CDR1 comprises the sequence RFMISEYSMH (SEQ ID NO: 16), the CDR2 comprises the sequence TINPAGQTDYAESVKG (SEQ ID NO: 10), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18).
6. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1 or 2, wherein the CDR1 comprises the sequence RFMISEYSMH (SEQ ID NO: 16), the CDR2 comprises the sequence TINPAKTTDYAESVKG (SEQ ID NO: 9), and the CDR3 comprises the sequence DGYGY (SEQ ID NO: 18).
7. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1 or 2, wherein the CDR1 comprises the sequence RFMISEYHMH (SEQ ID NO: 5), the CDR2 comprises the sequence DINPAGTTDYAESVKG (SEQ ID NO: 8), and the CDR3 comprises the sequence DSYGY (SEQ ID NO: 15).
8. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 1 or 2, wherein the protein comprises a sequence selected from the group consisting of SEQ ID Nos. 21, 22, 24, 25, 31 and 32.
9. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of claim 8, wherein the protein comprises a sequence of SEQ ID NO: 32.
10. The prostate specific membrane single domain antibody~~antigen-binding protein~~ of any one of claims 1-9, wherein said binding protein binds to one or both of human prostate specific membrane antigen and cynomolgus prostate specific membrane antigen.
- ~~11. The prostate specific membrane antigen-binding protein of any one of claims 1-10, wherein said binding protein binds to human prostate specific membrane antigen and cynomolgus prostate specific membrane antigen with comparable binding affinities.~~
- ~~12. The prostate specific membrane antigen-binding protein of any one of claims 1-11, wherein said binding protein binds to human prostate specific membrane antigen with a higher binding affinity than cynomolgus prostate specific membrane antigen.~~
- ~~13.~~11. A polynucleotide encoding a PSMA binding protein according to any one of claims 1-~~12~~10.
- ~~14.~~12. A vector comprising the polynucleotide of claim ~~13~~11.
- ~~15.~~13. A host cell transformed with the vector according to claim ~~14~~12.
- ~~16.~~14. A pharmaceutical composition comprising (i) a PSMA binding protein according to any one of claims 1-~~12~~10, the polynucleotide according to claim 11~~13~~, the vector according to claim 12~~14~~ or the host cell according to claim ~~15~~13, and (ii) a pharmaceutically acceptable carrier.

~~17~~.15. Use of the PSMA binding protein according to any one of claims 1-~~12~~10 in the manufacture of a medicament for the treatment or amelioration of prostate cancer, wherein the medicament is to be administered to a subject in need thereof.

~~18~~.16. The use of claim ~~17~~15, wherein the subject is human.

~~19~~.17. The use of claim ~~18~~16, wherein an agent is to be further administered in combination with the medicament, and wherein the agent comprises an antibody, a small molecule drug, a hormone, a radiotherapy, a gene therapy, an immunotherapy, an anti-diarrheal agent, an anti-emetic agent, an analgesic, an opioid or a non-steroidal anti-inflammatory agent.

~~20~~.18. A multispecific binding protein comprising the PSMA binding protein according to any one of claims 1-~~11~~10.