

DESCRIPTION



159159

METHOD FOR PRODUCING HYDROXYLATED ADAMANTANE USING CYTOCHROME P450

Technical field

[0001]

The present invention relates to a method for producing an hydroxylated adamantane using cytochrome P450.

Background art

[0002]

The hydroxylated adamantane is known as a monomer of a functional resin such as a photosensitive resin and the like in the photolithography field (Patent Document 1), or an inhibitor of 11 β -hydroxysteroid dehydrogenase type I (11 β HSD1) in the field of medicines (Patent Document 2, page 27). In addition, a compound having an adamantyl group is known also as a DPPIV inhibitor. And, when these compounds are synthesized, a compound wherein a part of substituents of the hydroxylated adamantane is protected is useful as a synthesis intermediate from its reactivity.

[0003]

A process for producing a hydroxylated adamantane derivative using a microorganism has been previously known. For example, a hydroxylated adamantane derivative protected with benzamide can be produced using in vivo conversion of *Beauveria sulfurescens* (Non-patent Document 1). In addition, regarding an adamantane derivative protected with phthalimide, it has been reported that a dihydroxy form can be produced by in vivo conversion of *Sporotrichum sulfurescens* (Non-patent Document 2).

[0004]

Cytochrome P450 (hereinafter, also referred to as P450) is a generic name of protoheme-containing proteins exhibiting a specific absorption band (Soret band) at around 450 nm when it is reduced and carbon monoxide is bound thereto. P450 is bound to a microsome of many animal and plant tissues, fungi, and yeasts, or an internal membrane of mitochondria of a part of animal tissues. Additionally, it exists in a soluble form in certain bacteria



G00002

or fungi.

[0005]

P450 has various substrate specificities, and there are enzymes exhibiting very broad substrate specificity, which can utilize a variety of organic compounds as a substrate. On the other hand, there are enzymes having more restricted substrate specificity, which react only with relatively limited kinds of organic compounds. P450 is involved in biosynthesis of bioactive lipids such as cholesterol, steroid hormone, bile acid and active-type vitamin D, metabolism of bioactive lipids such as heme, fatty acid and eicosanoid, metabolism of exogenous chemical substances including drugs, and the like. As specific function of P450, it is known that P450 catalyzes a variety of reactions such as a hydroxylation reaction, an epoxidation reaction, a dealkylation reaction, a denitrification reaction and the like of xenobiotic in cells expressing the P450.

[0006]

Some P450s derived from microorganisms such as fungi or bacteria are known that they serve for production of industrially useful substances, and a part thereof are actually utilized in industrial production of useful drugs. A representative example is to hydroxylate a 6 β -position of compactin with an actinomycete, *Streptomyces carbophilus*, and obtain pravastatin which is an antihyperlipemic agent, as a product (Non-patent Document 3, Patent Document 3). A process for producing active-type vitamin D₃ by hydroxylating a 1 α -position and a 25-position of vitamin D₃ utilizing an actinomycete, *Pseudonocardia autotrophica* has been also put into practical use.

[0007]

With respect to hydroxylation of adamantane, Non-patent Document 4 describes that adamantane is hydroxylated with P450cam, which is P450 of *Pseudomonas putida*. In addition, Non-patent Document 5 describes that five microorganisms hydroxylating adamantane were screened from 470 microorganisms, and suggests that P450 is involved in hydroxylation of adamantane in *Streptomyces griseoplanus*, which is one of those microorganisms.

[0008]

However, none of documents describe or suggest hydroxylation of a

compound having an adamantane skeleton using P450, for example, hydroxylation of an N-(2-adamantyl)-benzamide derivative using P450.

[Patent Document 1] Japanese Patent Application Laid-Open (JP-A) No.2006-63061

5 [Patent Document 2] WO04/056744

[Patent Document 3] JP-A No.6-70780

[Non-patent Document 1] Journal of Organic Chemistry, 1992, vol.57, p.7209-7212

10 [Non-patent Document 2] The Journal of Organic Chemistry, 1968, vol.33, p.3201-3207

[Non-patent Document 3] Gene, 1995, vol.163, p.81-85

[Non-patent Document 4] Archives of Biochemistry and Biophysics, 1984, vol.228, p.493-502

15 [Non-patent Document 5] Applied Microbiology and Biotechnology, 2006, vo.71, p.502-504

Disclosure of the invention

Problems to be solved by the invention

[0009]

20 Object of the present invention is to provide a high-yield and stable process using P450 for producing a hydroxylated form of a compound having an adamantane skeleton which is a raw material of monohydroxy-2-adamantanamine useful as an intermediate of functional resins and medicines (e.g. N-(5-hydroxy-2-adamantyl)-benzamide derivative).

25

Means to solve the problems


[0010]

30 The present inventors have intensively studied and, as a result, found out a method for hydroxylating an adamantane part of a compound having an adamantane skeleton using CYP109B1 which is one of P450s. More particularly, it was first found out that an N-(5-hydroxy-2-adamantyl)-benzamide derivative can be produced from an N-(2-adamantyl)-benzamide derivative. In addition, it was found out that productivity of an N-(5-hydroxy-2-adamantyl)-benzamide derivative can be
35 increased by using a CYP109B1 mutant, 109FK, or a 109FK mutant, obtained

by altering CYP109B1. In addition, it was found out that the resulting N-(5-hydroxy-2-adamantyl)-benzamide derivative is Anti isomoer at 100 % which is preferable as an intermediate of medicines.

That is, the present invention relates to the following:

- 5 (1) A process for producing an N-(5-hydroxy-2-adamantyl)-benzamide derivative characterized by hydroxylating an N-(2-adamantyl)-benzamide derivative using cytochrome P450.
- (2) The process of (1),
wherein cytochrome P450 is
- 10 a protein consisting of an amino acid sequence of SEQ ID No.: 1 or 3, or
a protein consisting of an amino acid sequence in which one to several amino acids are deleted, substituted, or added in the amino acid sequence, and having hydroxylation activity.
- (3) The process of (1),
- 15 wherein cytochrome P450 is
a protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, L357P, A362T, F41L, I77W, M105I, T234A, F232I, F232L and F232M in the amino acid sequence of SEQ ID No.: 1, or
- 20 a protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3.
- (4) The process of (1), wherein the N-(2-adamantyl)-benzamide derivative is a
- 25 compound of the formula:
- [Formula 1]




wherein R is hydrogen or a hydrophilic group.

- (5) The process of (1), wherein the N-(5-hydroxy-2-adamantyl)-benzamide

derivative is a compound of the formula:


[Formula 2]



wherein R is hydrogen or a hydrophilic group.

5 (6) A process for producing a compound of the formula:

[Formula 3]




characterized by obtaining an N-(5-hydroxy-2-adamantyl)-benzamide derivative by the process of any one of (1) to (5), and

10 deprotecting it.

(7) The process of (6), characterized in that the N-(5-hydroxy-2-adamantyl)-benzamide derivative is not isolated.

(8) A process for producing a compound of the formula (III):

[Formula 4]



15

characterized by obtaining a compound of (I) by the process of (6) or (7), and reacting with a compound of the formula (II): A-R¹-R²-R³-X

wherein A is an optionally substituted cyclic hydrocarbon group or an optionally substituted heterocyclic group, R¹ is a single bond, -C(=O)-, -O- or

20 NR⁴-, R² is a single bond or optionally substituted alkylene, R³ is a single

bond or C(=O)-, X is a hydroxy group, halogen or a leaving group derived from a hydroxy group, and R⁴ is hydrogen or optionally substituted alkyl.

(9) A method for hydroxylating an adamantane part of a compound having an adamantane skeleton, comprising using

5 a protein consisting of an amino acid sequence of SEQ ID No.: 1 or 3, or
a protein consisting of an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence, and having hydroxylation activity.

10 (10) A protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence of SEQ ID No.: 1, and having hydroxylation activity.


(11) A protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, L357P, A362T, F41L, I77W, M105I, T234A, F232I, F232L and F232M in the amino acid sequence of
15 SEQ ID No.: 1.

(12) A protein consisting of the amino acid sequence of SEQ ID No.: 3, or a protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence, and having hydroxylation activity.

20 (13) A protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3.

(14) A compound of the formula:


25 [Formula 5]



a salt or a solvate thereof.

(15) A compound of the formula:

[Formula 6]



a salt or solvate thereof,
wherein R is hydrogen or a hydrophilic group.

Effect of the invention

5 [0011]

The method for hydroxylating an adamantane part of a compound having an adamantane skeleton of the present invention realizes stable production and supply, and enables effective, safe and low-cost production of a hydroxylated form of a compound having an adamantane skeleton.

10

Best mode for carrying out the invention

[0012]

Terms used herein, unless otherwise indicated, are used in a sense normally used in the art. Terms which are particularly used herein will be explained below.

15

[0013]

In the present invention, "cytochrome P450" is utilized as a catalyst for hydroxylating a compound having an adamantane skeleton. For example, it is utilized as a catalyst in a process for producing an N-(5-hydroxy-2-adamantyl)-benzamide derivative by hydroxylating an N-(5-hydroxy-2-adamantyl)-benzamide derivative. Whether or not a protein which is P450 can be utilized as a catalyst for hydroxylating a compound having an adamantane skeleton can be determined, for example, by whether or not an N-(5-hydroxy-2-adamantyl)-benzamide derivative is obtained by acting the protein, or the protein and a reducing system consisting of ferredoxin and reductase supplying an electron to the protein on an N-(2-adamantyl)-benzamide derivative which is a substrate. More particularly, a method for culturing a microorganism capable of expressing P450 in a suitable medium, and detecting the presence or absence of the

25

N-(5-hydroxy-2-adamantyl)-benzamide derivative accumulated in the medium can be used. If necessary, by introducing a reducing system consisting of ferredoxin supplying an electron to P450 and reductase into a microorganism, the presence or absence can be more effectively detected. Detection of the N-(5-hydroxy-2-adamantyl)-benzamide derivative can be performed, for example, by high performance liquid chromatography (HPLC) using the N-(5-hydroxy-2-adamantyl)-benzamide derivative obtained by the present invention as a standard. Specifically, the method described in Examples later can be used. Examples of P450 useful in the present invention include CYP109B1, CYP101, CYP105D and the like, and a derivative thereof.

[0014]

"Derivative" of P450 means a polypeptide having substantially the same biological function or activity as that of P450, that is, in the present invention, hydroxylation activity. A C-terminus of the polypeptide may be either of a carboxyl group, carboxylate, amide or ester. Examples include a fused protein with other protein, a modified entity obtained by adding a modifying group, a mutant obtained by deleting, substituting or adding an amino acid residue(s), and the like.

As "other protein" used in the fused protein, a protein that the fused protein exhibits substantially the same or more hydroxylation activity as that of P450 by fusing is preferable. Examples include R450RhF reductase domain, P450BM3 reductase domain and the like.

Examples of the modifying group of the modified entity include a functional group exhibiting fluorescence property, a functional group which is not involved in formation of a steric structure of a polypeptide, and the like. Alternatively, it may be an amino acid residue other than amino acid residues constituting naturally occurring P450 (e.g. β -alanine and the like). Examples of the functional group exhibiting fluorescence property include eosin, fluorescein isothiocyanate (FITC) and the like. Examples of the functional group which is not involved in formation of a steric structure of a polypeptide include a spacer group, a representative of which is a β -alanine residue and the like. It is preferable that such a functional group is present at a terminal.

Examples of the mutant include a protein consisting of an amino acid sequence in which one to several amino acids are deleted, substituted or

added in the amino acid sequence of P450, and having the function of P450, and the like.

[0015]

Examples of the "amino acid sequence in which one to several amino acids are deleted, substituted or added" include (A) an amino acid sequence in which one or two or more (preferably, around 1 to 30, preferably around 1 to 10, and further preferably several (1 to 5)) amino acids in the amino acid sequence of P450 are deleted, (B) an amino acid sequence in which one or two or more (preferably, around 1 to 30, preferably around 1 to 10, and further preferably several (1 to 5)) amino acids in the amino acid sequence of P450 are substituted with other amino acids, (C) an amino acid sequence in which one or two or more (preferably, around 1 to 30, preferably around 1 to 10, and further preferably several (1 to 5)) amino acids are added in the amino acid sequence of P450, and (D) a protein containing an amino acid sequence which is a combination of them. When an amino acid sequence is deleted, substituted or added, a position of the deletion, substitution or addition is not particularly limited, provided that a mutant of P450 used in the present invention is a polypeptide having activity of catalyzing hydroxylation of a compound having an adamantane skeleton, even though an amino acid(s) is deleted, substituted or added. Examples include a polypeptide having at least 60% or more homology, preferably a polypeptide having 80% or more homology, and further preferably a polypeptide having 95% or more homology with the amino acid sequence of P450.

[0016]

Herein, "CYP109B1" means a protein consisting of the amino acid sequence of SEQ ID No.: 1.

"CYP109B1 mutant" means a protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence of SEQ ID No.: 1, and having hydroxylation activity. Examples include a protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, L357P, A362T, F41L, I77W, M105I, T234A, F232I, F232L and F232M, in the amino acid sequence of SEQ ID No.: 1. "One to several mutations" means around 1 to 5, preferably 1 to 3, and further preferably 1 or 2 mutations. Particularly, a protein consisting of an amino acid sequence having 1 to 5, and

preferably 1 to 3 variations selected from the group consisting of I77F; I77F and L357P; I77F, M105I and T234A; and I77F and F232M, in the amino acid sequence of SEQ ID No.: 1, is preferable.

[0017]

5 "CYP109FK" means a protein consisting of the amino acid sequence of SEQ ID No.: 3.

"CYP109FK mutant" means a protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence of SEQ ID No.: 3, and having hydroxylation
10 activity. Examples include a protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3. "One to several mutations" means around 1 to 5, preferably 1 to 3, and further preferably 1 or 2 mutations.
15 Particularly, a protein consisting of an amino acid sequence having 1 to 5, preferably 1 to 3 and particularly preferably one mutation selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3, is preferable.

20 [0018]

"Compound having an adamantane skeleton" includes any compound as far as it is a compound having an adamantane skeleton. Preferable examples include protected 2-aminoadamantane.


"Protected 2-aminoadamantane" includes any compound as far as it is
25 a compound wherein an amino group of adamantanamine is protected. In this respect, a part other than a part to be hydroxylated, of an adamantane part may have a substituent. The substituent is not particularly limited, but examples thereof include halogen such as F, Cl, Br, I and the like, optionally substituted alkyl (e.g. unsubstituted alkyl and carbamoyl alkyl), hydroxy,
30 alkoxy, nitro, aryl, arylalkyl, carboxy, ester (e.g. alkoxycarbonyl and the like), carbamoyl, alkenyl, alkynyl, carbamoyloxy, protected hydroxy (e.g. alkylhydroxy and the like) and the like. Preferable examples include an N-(2-adamantyl)-benzamide derivative, N-(2-adamantyl)-benzamide,
N-(adamantan-2-yl)-phthalamic acid,
35 N-(2-adamantyl)-p-methoxybenzyloxycarbonylamine and the like.

[0019]

"N-(2-adamantyl)-benzamide derivative" means

N-(2-adamantyl)benzamide or a derivative thereof. The derivative includes a compound wherein a benzene ring part is substituted with halogen, alkyl, hydroxy, alkoxy, nitro, aryl, arylalkyl, carboxy, ester, carbamoyl or the like, a compound wherein a part other than a part to be hydroxylated, of an adamantyl group is substituted with halogen, alkyl, hydroxy, alkoxy, nitro, aryl, arylalkyl, carboxy, ester, carbamoyl or the like, and a compound wherein a benzene ring part and an adamantane part are substituted with the
10 aforementioned substituents. Preferable examples include a compound of the formula:

[Formula 7]



wherein R is hydrogen or a hydrophilic group.

15 [0020]

"Hydrophilic group" is a group having strong affinity for water. That is, it refers to an atomic group which can form a weak bond with a water molecule by electrostatic interaction, a hydrogen bond or the like. Examples include -OH, -COOH, -NH₂, -OSO₃H, -SO₃H, -OPO₃H₂ and the like.


20 Particularly, -COOH is preferable.

[0021]

"N-(5-hydroxy-2-adamantyl)-benzamide derivative" means


N-(monohydroxy-2-adamantyl)-benzamide or a derivative thereof. Examples of the derivative also include a compound wherein a benzene ring is substituted with halogen, alkyl, hydroxy, alkoxy, nitro, aryl, arylalkyl, carboxy, alkoxycarbonyl, carbamoyl or the like. Preferable examples include a compound of the formula:

[Formula 8]



wherein R is hydrogen or a hydrophilic group. Further preferable examples include a compound of the formula:

[Formula 9]




5

wherein R is hydrogen or a hydrophilic group.

[0022]

A compound having an adamantane skeleton obtained by the present invention such as the N-(5-hydroxy-2-adamantyl)-benzamide derivative can be easily converted into monohydroxy-2-adamantanamine, that is, a compound of the formula:

[Formula 10]



by removing a protecting group.

15 [0023]

In addition, in the present invention, the N-(5-hydroxy-2-adamantyl)-benzamide derivative may or may not be isolated in a step of producing an N-(5-hydroxy-2-adamantyl)-benzamide derivative from an N-(2-adamantyl)-benzamide derivative and obtaining

monohydroxy-2-adamantanamine by deprotection. Upon production of the N-(5-hydroxy-2-adamantyl)-benzamide derivative, the derivative can be once purified and deprotected. Alternatively, as described in Example 8, monohydroxy-2-adamantanamine can be obtained by performing deprotection while sterilization is conducted, without isolating the N-(5-hydroxy-2-adamantyl)-benzamide derivative.


[0024]

Further, by reacting monohydroxy-2-adamantanamine obtained above with a compound of the formula (II): $A-R^1-R^2-R^3-X$

wherein A is an optionally substituted cyclic hydrocarbon group or an optionally substituted heterocyclic group, R^1 is a single bond, $-C(=O)-$, $-O-$ or NR^4 , R^2 is a single bond or optionally substituted alkylene, R^3 is a single bond or $C(=O)-$, X is a hydroxy group, halogen or a leaving group derived from a hydroxy group, and R^4 is hydrogen or optionally substituted alkyl,

a compound of the formula (III):


[Formula 11]



can be produced.

[0025]

[Formula 12]



This is a step of reacting a compound of the formula (I) and a compound of the formula (II) to produce a compound of the formula (III).

Examples of a solvent include N-dimethylformamide, dimethyl sulfoxide, aromatic hydrocarbons (e.g. toluene, benzene, xylene and the like), saturated hydrocarbons (e.g. cyclohexane, hexane and the like), halogenated

hydrocarbons (e.g. dichloromethane, chloroform, 1,2-dichloroethane and the like), ethers (e.g. tetrahydrofuran, diethyl ether, dioxane, 1,2-dimethoxyethane and the like), esters (e.g. methyl acetate, ethyl acetate and the like), ketones (e.g. acetone, methyl ethyl ketone and the like), nitriles (e.g. acetonitrile and the like), alcohols (e.g. methanol, ethanol, isopropanol, tert-butanol and the like), water, a mixed solvent thereof and the like. Preferable are halogenated hydrocarbons (e.g. dichloromethane, chloroform, 1,2-dichloroethane and the like), nitriles (e.g. acetonitrile and the like), ethers (e.g. tetrahydrofuran, diethyl ether, dioxane, 1,2-dimethoxyethane and the like), alcohols (e.g. methanol, ethanol, isopropanol, tert-butanol and the like), water and the like.

Without any limitation, preferable are N,N-dimethylformamide, dimethyl sulfoxide, xylene, dichloromethane, chloroform, 1,2-dichloroethane, diethyl ether, dioxane, 1,2-dimethoxyethane, acetonitrile, methanol, ethanol, isopropanol, tert-butanol, toluene, tetrahydrofuran and water.

Further preferable are dichloromethane, methanol, and ethanol.

When R^5 is $-C(=O)-$ and X is a hydroxy group, a condensing agent and a base can be used in the step. As the condensing agent, for example, 1,1-carbonyldiimidazole, dicyclohexylcarbodiimide, water-soluble carbodiimide (1-ethyl-3-(3'-dimethylaminopropyl)carbodiimide) or the like can be used. Examples of the base include metal hydrides (e.g. sodium hydride and the like), metal hydroxides (e.g. sodium hydroxide, potassium hydroxide, lithium hydroxide, barium hydroxide and the like), metal carbonates (e.g. sodium carbonate, calcium carbonate, cesium carbonate and the like), metal alkoxides (e.g. sodium methoxide, sodium ethoxide, potassium tert-butoxide and the like), sodium bicarbonate, metal sodium, organic amines (e.g. triethylamine, diisopropylethylamine, DBU, 2,6-lutidine and the like) and the like.

When R^5 is $-C(=O)-$ and X is halogen, the above base can be used in the step.

When R^3 is $-C(=O)-$, R^4 is optionally substituted alkylene, R^5 is a single bond and X is halogen or a leaving group derived from a hydroxy group, the above base can be used in the step.

Reaction conditions are not particularly limited, but the substances may be stirred at about -20 to 100°C , and preferably about -10 to 80°C for

usually 1 hour to 36 hours, and preferably 1 hour to 24 hours.

The thus obtained Compound (III) is useful as an 11β -hydroxysteroid dehydrogenase inhibitor, a Dipeptidyl Peptidase IV (DPP IV) inhibitor, a Jak3 inhibitor or the like.

5 [0026]

"Halogen" includes fluorine, chlorine, bromine and iodine.

"Alkyl" means C1 to C10 straight or branched alkyl. Examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, isohexyl, n-heptyl, n-octyl,
10 n-nonyl, n-decyl and the like. Preferable is C1 to C6 or C1 to C4 alkyl. Examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl and isohexyl.

An alkyl part of "alkoxy" or "alkoxycarbonyl" has the same meaning as the above alkyl.

15 Examples of "aryl" include monocyclic aromatic hydrocarbon groups (e.g. phenyl) and polycyclic aromatic hydrocarbon groups (e.g. 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl, 9-anthryl, 1-phenanthryl, 2-phenanthryl, 3-phenanthryl, 4-phenanthryl, 9-phenanthryl and the like).

"Arylalkyl" means the above alkyl substituted with the above aryl.

20 "Cyclic hydrocarbon group" includes "cycloalkyl", "cycloalkenyl" and "aryl".

"Cycloalkyl" means C3 to C15 cyclic saturated hydrocarbon group. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, a bridged cyclic hydrocarbon group, a spiro hydrocarbon group and
25 the like.

"Bridged cyclic hydrocarbon group" includes a group obtained by removing one hydrogen from C5 to C8 aliphatic ring in which two or more rings share 2 or more atoms. Examples include bicyclo[2.1.0]pentyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl and bicyclo[3.2.1]octyl,
30 tricyclo[2.2.1.0]heptyl and the like.

"Spiro hydrocarbon group" includes a group obtained by removing one hydrogen from a ring in which two hydrocarbon rings are constructed by sharing one carbon atom. Examples include spiro[3.4]octyl and the like.

35 "Cycloalkenyl" means C3 to C7 cyclic unsaturated aliphatic hydrocarbon group. Examples include cyclopropenyl, cyclobutenyl,

cyclopentenyl, cyclohexenyl, and cycloheptenyl. Preferable are cyclopropenyl, cyclobutenyl, cyclopentenyl, and cyclohexenyl. Cycloalkenyl includes a bridged cyclic hydrocarbon group and a spiro hydrocarbon group having an unsaturated bond in a ring.

5 "Heterocyclic group" includes "heteroaryl and heterocycle".

"Heteroaryl" means a monocyclic aromatic heterocyclic group and a fused aromatic heterocyclic group. The monocyclic aromatic heterocyclic group means a group derived from a 5- to 8- membered aromatic ring optionally containing 1 to 4 of oxygen, sulfur and/or nitrogen atoms in a ring,
10 optionally having a bond at an arbitrary replaceable position. The fused aromatic heterocyclic group means a group in which a 5- to 8- membered aromatic ring optionally containing 1 to 4 of oxygen, sulfur and/or nitrogen atoms in a ring is fused with 1 to 4 of a 5- to 8- membered aromatic carbocycle or other 5- to 8- membered aromatic heterocycle, and optionally having a bond
15 at an arbitrary replaceable position. Examples include furyl (e.g. 2-furyl and 3-furyl), thienyl (e.g. 2-thienyl and 3-thienyl), pyrrolyl (e.g. 1-pyrrolyl, 2-pyrrolyl and 3-pyrrolyl), imidazolyl (e.g. 1-imidazolyl, 2-imidazolyl and 4-imidazolyl), pyrazolyl (e.g. 1-pyrazolyl, 3-pyrazolyl and 4-pyrazolyl), triazolyl (e.g. 1,2,4-triazol-1-yl, 1,2,4-triazol-3-yl and 1,2,4-triazol-4-yl),
20 tetrazolyl (e.g. 1-tetrazolyl, 2-tetrazolyl and 5-tetrazolyl), oxazolyl (e.g. 2-oxazolyl, 4-oxazolyl and 5-oxazolyl), isoxazolyl (e.g. 3-isoxazolyl, 4-isoxazolyl and 5-isoxazolyl), thiazolyl (e.g. 2-thiazolyl, 4-thiazolyl and 5-thiazolyl), thiadiazolyl, isothiazolyl (e.g. 3-isothiazolyl, 4-isothiazolyl and 5-isothiazolyl), pyridyl (e.g. 2-pyridyl, 3-pyridyl and 4-pyridyl), pyridazinyl
25 (e.g. 3-pyridazinyl and 4-pyridazinyl), pyrimidinyl (e.g. 2-pyrimidinyl, 4-pyrimidinyl and 5-pyrimidinyl), furazanyl (e.g. 3-furazanyl), pyrazinyl (e.g. 2-pyrazinyl), oxadiazolyl (e.g. 1,3,4-oxadiazol-2-yl), benzofuryl (e.g. 2-benzo[b]furyl, 3-benzo[b]furyl, 4-benzo[b]furyl, 5-benzo[b]furyl, 6-benzo[b]furyl and 7-benzo[b]furyl), benzothienyl (e.g. 2-benzo[b]thienyl, 3-benzo[b]thienyl, 4-benzo[b]thienyl, 5-benzo[b]thienyl, 6-benzo[b]thienyl and 7-benzo[b]thienyl), benzimidazolyl (e.g. 1-benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl and 5-benzimidazolyl), dibenzofuryl, benzoxazolyl, quinoxalyl (e.g. 2-quinoxalyl, 5-quinoxalyl and 6-quinoxalyl), cinnolinyl (e.g. 3-cinnolinyl, 4-cinnolinyl, 5-cinnolinyl, 6-cinnolinyl,
35 7-cinnolinyl and 8-cinnolinyl), quinazolyl (e.g. 2-quinazolyl, 4-quinazolyl,

5-quinazolyl, 6-quinazolyl, 7-quinazolyl and 8-quinazolyl), quinolyl (e.g. 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl and 8-quinolyl), phthalazinyl (e.g. 1-phthalazinyl, 5-phthalazinyl and 6-phthalazinyl), isoquinolyl (e.g. 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl and 8-isoquinolyl), puryl, pteridinyl (e.g. 2-pteridinyl, 4-pteridinyl, 6-pteridinyl and 7-pteridinyl), carbazolyl, phenanthrydinyl, acridinyl (e.g. 1-acridinyl, 2-acridinyl, 3-acridinyl, 4-acridinyl and 9-acridinyl), indolyl (e.g. 1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl and 7-indolyl), isoindolyl, phenazinyl (e.g. 1-phenazinyl and 2-phenazinyl), phenothiazinyl (e.g. 1-phenothiazinyl, 2-phenothiazinyl, 3-phenothiazinyl and 4-phenothiazinyl) and the like.

"Heterocycle" means a non-aromatic heterocyclic group optionally containing 1 to 4 of oxygen, sulfur and/or nitrogen atoms in a ring, and optionally having a bond at an arbitrary replaceable position. In addition, such a non-aromatic heterocyclic group may be further bridged with C1 to C4 alkyl chain, and may be fused with cycloalkane (5- to 6-membered ring is preferable) or a benzene ring. Heterocycle may be saturated or unsaturated as far as it is non-aromatic. Examples include 1-pyrolinyl, 2-pyrolinyl, 3-pyrolinyl, 1-pyrolidinyl, 2-pyrolidinyl, 3-pyrolidinyl, 1-imidazolynyl, 2-imidazolynyl, 4-imidazolynyl, 1-imidazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 1-pyrazolynyl, 3-pyrazolynyl, 4-pyrazolynyl, 1-pyrazolidinyl, 3-pyrazolidinyl, 4-pyrazolidinyl, piperidino, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 1-piperazinyl, 2-piperazinyl, 2-morpholinyl, 3-morpholinyl, morpholino, tetrahydropyranyl and the like.

"Alkylene" includes a divalent group in which 1 to 6 methylenes are continuous. Examples include methylene, ethylene, trimethylene, tetramethylene, pentamethylene, hexamethylene and the like.

Examples of "leaving group derived from a hydroxy group" include -Oms, -OTs, -OTf, -ONs and the like. Herein, "Ms" represents a methanesulfonyl group, "Ts" represents a paratoluenesulfonyl group, "Tf" represents a trifluoromethanesulfonyl group, and "Ns" represents an orthonitrobenzenesulfonyl group.

[0027]

A in the formula (II) and the formula (III) is preferably an optionally substituted heterocyclic group. Further preferable examples include

optionally substituted heteroaryl and optionally substituted heterocycle. More preferable examples include furan, thiophene, pyrrole, pyrazole, triazole, oxazole, thiazole, isothiazole, pyridine, morpholine, piperidine, piperazine, pyrrolidine, tetrahydrothiophene, benzoxazine, benzofuran, and pyrrolopyridine. Without any limitation, particularly, isoxazole and pyrazole are preferable. Examples of A include an optionally substituted cyclic hydrocarbon group. Preferable is phenyl.

Examples of the substituent include $-OR^5$, $-SR^5$, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocycle, a group of the formula: $-CH=CH-C(R^aR^b)-R^c-R^d$, a group of the formula: $-C(R^eR^f)_m-C(R^aR^b)-R^c-R^d$ and the like.

R^a and R^b are each independently hydrogen, optionally substituted alkyl or halogen, or R^a and R^b may be taken together with an adjacent carbon atom to form an optionally substituted ring,

R^c is $-(CH_2)_n-$, wherein n is a integer of 0 to 3,

R^d is hydrogen, halogen, hydroxy, carboxy, cyano, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocycle, a group of the formula: $-C(=O)-NR^gR^h$ or a group of the formula: $-NR^iR^j$,

R^e and R^f are each independently hydrogen, halogen or optionally substituted alkyl,

R^g and R^h are each independently hydrogen, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocycle, optionally substituted alkylsulfonyl, optionally substituted cycloalkylsulfonyl, optionally substituted arylsulfonyl, optionally substituted heteroarylsulfonyl, optionally substituted heterocyclesulfonyl, optionally substituted alkyloxy, or optionally substituted carbamoyl, or R^g and R^h may be taken together with an adjacent nitrogen atom to form an optionally substituted ring,

Rⁱ and R^j are each independently hydrogen, carboxy, hydroxy, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted cycloalkyl, optionally substituted cycloalkenyl, optionally substituted aryl, optionally substituted heteroaryl, optionally substituted heterocycle, optionally substituted acyl, optionally substituted carbamoyl, optionally substituted thiocarbamoyl, optionally substituted alkylsulfonyl, optionally substituted cycloalkylsulfonyl, optionally substituted arylsulfonyl, optionally substituted heteroarylsulfonyl, optionally substituted heterocyclesulfonyl, optionally substituted alkyloxycarbonyl, optionally substituted cycloalkyloxycarbonyl, optionally substituted aryloxycarbonyl, optionally substituted heteroaryloxycarbonyl, optionally substituted heterocycleoxycarbonyl, optionally substituted alkylcarbonyl, optionally substituted cycloalkylcarbonyl, optionally substituted arylcarbonyl, optionally substituted heteroarylcarbonyl, optionally substituted heterocyclecarbonyl, or optionally substituted sulfamoyl, or Rⁱ and R^j may be taken together with an adjacent nitrogen atom to form an optionally substituted ring,

R⁵ is optionally substituted alkyl, optionally substituted alkenyl, optionally substituted cycloalkyl, optionally substituted aryl, optionally substituted heteroaryl, or optionally substituted heterocycle, and

m are each independently an integer of 1 to 3.

R¹ is preferably a single bond.

R² is preferably a single bond.

R³ is preferably -C(=O)-.

X is preferably a hydroxy group.

Details of compounds of the formula (II) and the formula (III), and a process for preparing them are collectively described in WO2006/132197, WO2007/058346, PCT/JP2007/056538, and Japanese Patent Application No.2007-132259.

[0028]

A method for hydroxylating a compound having an adamantane skeleton of the present invention, for example, a process for producing an N-(5-hydroxy-adamantan-2-yl)-benzamide derivative by hydroxylating an N-(2-adamantyl)-benzamide derivative (hereinafter, referred to as "present process" in some cases) characterized by using P450 having hydroxylation

activity (hereinafter, referred to as "present protein" in some cases) as a catalyst. Specifically, the present process includes a step of acting at least one of

(a) the present protein, or

5 the present protein and a reduction system consisting of ferredoxin and reductase supplying an electron to the protein, or

(b) a microorganism capable of expressing the present protein,

on an N-(2-adamantyl)-benzamide derivative which is a substrate, to selectively hydroxylate an adamantane part.

10 [0029]

(a) Present protein

The present protein can be made by obtaining and utilizing a polynucleotide encoding the protein (hereinafter, referred to as "present gene" in some cases). A method for obtaining the present gene and a process for
15 producing the present protein (a process for making an expression vector or a transformant and the like) will be specifically described below.

[0030]

(1) Method for obtaining present gene

Present gene can be obtained from *Bacillus subtilis* by making a cDNA
20 library by a conventional method.

[0031]

Examples of the process for making a cDNA library include the methods described in *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) (Cold Spring Harbor Laboratory Press), *Current Protocols in*
25 *Molecular Biology* (1994) (Wiley-Interscience), *DNA Cloning 1: Core Techniques, A Practical Approach*, Second Edition (1995) (Oxford University Press) and the like, and a method using a commercially available kit such as SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Invitrogen), ZAP-cDNA Synthesis Kits (Stratagene) and the like.

30 [0032]

Examples of a cDNA obtained by the methods include a DNA encoding a protein consisting of an amino acid sequence of SEQ ID No.: 1, specifically, a DNA consisting of a base sequence of SEQ ID No.: 2 and the like. The cDNA can be used for making an expression plasmid in which the present gene is
35 incorporated into a suitable expression vector. The expression vector, a

method for using the expression plasmid and the like will be described in "Process for producing present protein" later.

[0033]

Alternatively, a DNA can be also prepared by chemically synthesizing a DNA encoding the present protein based on an amino acid sequence. Chemical synthesis of a DNA can be performed by a DNA synthesizer manufactured by Shimadzu Corporation utilizing a thiophosphite method, a DNA synthesizer model 392 manufactured by Perkin Elmer utilizing a phosphoamidite method, or the like.

[0034]

Further, an objective DNA can be also prepared by performing PCR using oligonucleotides as a sense primer and an antisense primer based on an amino acid sequence and a cDNA prepared from a mRNA of cells expressing a mRNA complementary with these DNAs as a template.

[0035]

(2) Process for producing present protein

The present protein can be produced by expressing a polynucleotide encoding the present protein in a host cell, for example, by the methods described in Molecular Cloning: A Laboratory Manual, Second Edition (1989) (Cold Spring Harbor Laboratory Press), Current Protocols in Molecular Biology (1994) (Wiley-Interscience) and the like, as below.

As the host cell, any of a prokaryotic cell, yeast, an animal cell, a plant cell, an insect cell and the like can be used as far as it can express an objective gene. As an expression vector, an expression vector which can be autonomously replicated in the above host cell, or can be incorporated into a chromosome, and contains a promoter at a position suitable for transcription of the present protein is used.

[0036]

(i) Case where prokaryote is used as host

It is preferable that the expression vector of the present protein can be autonomously replicated in a prokaryote and, is composed of a promoter, a ribosome binding sequence, a DNA encoding the present polypeptide and a transcription terminator sequence. A gene for controlling a promoter may be contained.

[0037]

Examples of the expression vector include pBTrp2, pBTac1, pBTac2 (Roche Diagnostics), Bluescript II SK (+), pBluescript II SK (-) (Stratagene), pSKV28, pUC118, pUC19 (Takara Shuzo Co., Ltd.), pKK233-2 (GE Healthcare), pSE280, pSupex, pUB110, pTP5, pC194, pTixFus (Invitrogen), pGEMEX-1 (Promega), pQE-8 (Qiagen), pGEX (GE Healthcare), pET system (Novagen), pMAL-c2 (New England Biolabs), pKYP10 (JP-A No.58-110600), pKYP200 (Agricultural Biological Chemistry, 48, 669 (1984)), pLSA1 (Agricultural Biological Chemistry, 53,277 (1989)), pGEL1 (Proceeding of the National Academy of Sciences USA, 82, 4306 (1985)), pEG400 (Journal of Bacteriology, 172, 2392 (1990)), pTrs30 (FERM BP-5407), pTrs32 (FERM BP-5408), pGHA2 (FERM BP-400), pGKA2 (FERM B-6798), pPA1 (JP-A No.63-233798), pTerm2 (JP-A No.3-22979, US4686191, US4939094 and US5160735) and the like.

[0038]

As the promoter, any promoter can be used as far as it can be expressed in a host cell such as *Escherichia coli* and the like. Examples include promoters derived from *Escherichia coli* and phage such as trp promoter (Ptrp), lac promoter (P lac), PL promoter, PR promoter, PSE promoter and the like, SPO1 promoter, SPO2 promoter, penP promoter and the like. Alternatively, artificially designed and altered promoters such as a promoter in which two Ptrp(s) are connected in series (Ptrpx2), tac promoter, lacT7 promoter, and letI promoter can be also used.

[0039]

It is preferable to use a plasmid in which a length between a Shine-Dalgarno sequence being a ribosome binding sequence and an initiation codon is adjusted at a suitable distance, for example, 6 to 18 bases. A transcription terminator sequence is not necessarily required for expression of the present gene, but it is preferable to arrange a transcription terminator sequence immediately downstream of a structural gene.

[0040]

Examples of the host cell include prokaryotes such as genus *Escherichia*, *Serratia*, *Bacillus*, *Brevibacterium*, *Corynebacterium*, *Microbacterium*, *Pseudomonas* and the like. Examples of genus *Escherichia* include strains of *E. coli* XL1-Blue, XL2-Blue, DH1, MC1000, KY3276, W1485, JM109, HB101, No.49, W3110, NY49, BL21, BL21 (DE3), BL21 (DE3) pLysS, HMS174 (DE3), HMS174 (DE3) pLysS and the like. Examples of genus

Serratia include strains of *S. ficaria*, *S. fonticola*, *S. liquefaciens*, *S. marcescens* and the like. Examples of genus *Bacillus* include strains of *B. subtilis*, *B. amyloliquefaciens* and the like. Examples of genus *Brevibacterium* include strains of *B. ammoniagenes*, *B. immariophilum* (ATCC:14068), *B. saccharolyticum* (ATCC:14066) and the like. Examples of genus *Corynebacterium* include strains of *C. glutamicum* (ATCC:13032), *C. glutamicum* (ATCC14067), *C. glutamicum* (ATCC13869), *C. acetoacidophilum* (ATCC13870) and the like. Examples of genus *Microbacterium* include strains of *M. ammoniophilum* (ATCC:15354) and the like. Examples of genus *Pseudomonas* include strains of *P. mephitica* and the like.

[0041]

As a method for introducing a recombinant vector, any method can be used as far as it is a method for introducing a DNA in the above host cell. Examples include an electroporation method (Nucleic Acids Research, 16, 6127 (1988)), a Calcium Phosphate Method (Proceedings of the National Academy of Sciences USA, 69, 2110 (1972)), a Protoplast Method (JP-A No.63-1483942), methods described in Gene, 17, 107 (1982) and Molecular & General Genetics, 168, 111 (1979) and the like.

[0042]

(ii) Case where yeast is used as host

When yeast is used as a host, examples of an expression vector include Yep13 (ATCC: 37115), Yep24 (ATCC: 37051), YCp50 (ATCC: 37419), pHS19, pHS15 and the like.

As the promoter, any promoter can be used as far as it can be expressed in yeast. Examples include ADH1 (alcohol dehydrogenase) promoter, PHO5 (acidic phosphatase) promoter, PGK1 (phosphoglycerate kinase) promoter, GAPDH (glyceraldehyde 3-phosphate dehydrogenase) promoter, GAL1 (galactose kinase) promoter, GAL10 (UDP galactose 4-epimerase) promoter, MF α 1 (α pheromone) promoter, CUP1 (metallothionein) promoter and the like.

Examples of the host include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Trichosporon pullulans*, *Schwanniomyces alluvius* *Pichia pastoris* and the like.

As the method for introducing a recombinant vector, any method can be used as far as it is a method for introducing a DNA into a host. Examples include an electroporation method (Methods in Enzymology, 194, 182 (1990)),

a Spheroplast Method (Proceedings of the National Academy of Sciences USA, 84, 1929 (1978)), a Lithium Acetate Method (Journal of Bacteriology, 153, 163 (1983)), the methods described in Proceedings of the National Academy of Sciences USA, 75, 1929 (1978)) and the like.

5 [0043]

(iii) Case where animal cell is used as host

When an animal cell is used as a host, examples of an expression vector includes pcDNA1/Amp, pcDNA1, pCDM8, pREP4 (Invitrogen), pAGE107 (Cytotechnology, 3, 133 (1990)), pAGE103 (The Journal of Biochemistry, 101, 1307 (1987)), pAMo, pAMoA (pAMoPRSA) (The Journal of Biological Chemistry, 268, 22782-22787 (1993)), pAS3-3 (JP-A No.2-22705) and the like.

As a promoter, any promoter can be used as far as it can be expressed in a host. Examples include IE (Immediate-early) gene promoter of human cytomegalovirus (hCMV), SV40 early promoter, a long terminal repeat promoter of Moloney Murine Leukemia Virus, a retrovirus promoter, HSP promoter, SR α promoter, a metallothionein promoter and the like. Alternatively, an enhancer of an IE gene of human CMV may be used together with a promoter.

[0044]

20 Examples of an animal cell used in a host include a human-derived strain cell such as HEK298 (human fetal kidney cell, ATCC: CRL-1573), Namalwa (Burkitt's lymphoma, ATCC: CRL-1432), HeLa (uterine cervical cancer cell, ATCC: CCL-2), HBT5637 (leukemia cell, JP-A No.63-299), BALL-1 (leukemia cell) and HCT-15 (large intestine cancer cell), a mouse-derived strain cell such as Sp2/0-Ag14 (mouse myeloma cell, ATCC: CRL-1581) and NSO (mouse myeloma cell), a monkey-derived strain cell such as COS-1 (African Green Monkey kidney cell (SV40 transformant cell), ATCC: CRL-1650) and COS-7 (African Green Monkey kidney cell (SV40 transformant cell), ATCC: CRL-1651), a hamster-derived strain cell such as CHO-K1 (Chinese hamster ovary cell, ATCC: CCL-61) and BHK-21 (C-13) (Sicilian hamster young kidney cell, ATCC: CCL-10), a rat-derived strain such as cell PC12 (adrenal melanocytoma, ATCC: CRL-1721) and YB2/0 (rat myeloma cell, ATCC: CRL-1662) and the like.

[0045]

35 As a method for introducing a recombinant vector, any method can be

used as far as it is a method for introducing a DNA into a host. Examples include an electroporation method (Cytotechnology, 3, 133,(1990)), a calcium phosphate method (JP-A No.2-22705), a lipofection (Proceedings of the National Academy of Sciences USA, 84, 7413 (1987)); Virology, 52, 456 (1973)).

5 [0046]

(iv) Case where plant cell is used as host

When a plant cell or a plant individual is used as a host, a polypeptide can be produced according to the known method (Tissue Culture, 20 (1994), Tissue Culture 21 (1995), Trends in Biotechnology, 15, 45 (1997)). Examples
10 of an expression vector include Ti plasmid, a tobacco mosaic virus vector and the like. As a promoter used in gene expression, any promoter can be used as far as it can be expressed in a plant cell. Examples include a cauliflower mosaic virus (CaMV) 35S promoter, a rice actin 1 promoter and the like. In addition, an efficiency of gene expression can be increased by inserting intron
15 1 of an alcohol dehydrogenase gene of corn or the like between a promoter and a gene to be expressed.

[0047]

Examples of a host include plant cells such as potato, tobacco, corn, rice, rapeseed, soybean, tomato, carrot, wheat, barley, rye, alfalfa, flax and
20 the like.

As a method for introducing a recombinant vector, any method can be used as far as it is a method for introducing a DNA into a host. Examples include a method using Agrobacterium (JP-A No.59-140885, JP-A No.60-70080 and WO 94/00977), an electroporation method (JP-A No.60-251887), a particle
25 gun (gene gun) method (Japanese Patent No.2606856 and Japanese Patent No.2517813) and the like.

[0048]

(v) Case where insect cell is used as host

When an insect cell is used as a host, examples of a transfer vector
30 include pVL1392, pVL1393, pBlueBacIII (Invitrogen) and the like. Examples of a virus for infection include baculovirus which infects Hadeninae insects, Autographa californica nuclear polyhedrosis virus (AcMNPY) Bac-N-Blue DNA and the like. Examples of a method for transforming an insect cell include the methods described in Baculovirus Expression Vector: A Laboratory
35 Manual (1992) (W. H. Freeman and Company), Molecular Cloning: A

Laboratory Manual, Second Edition (1989) (Cold Spring Harbor Laboratory Press), Current Protocols in Molecular Biology (1994) (Wiley-Interscience), BioTechnology, 6, 47 (1988) and the like.

[0049]

- 5 A polypeptide can be expressed by adding a transfer vector containing an objective gene, and a baculovirus DNA for infecting an insect cell to an insect cell culture solution to infect the insect cell with a virus expressing an objective gene obtained by recombination.

10 Examples of the insect cell used as a host include a strain cell derived from *Spodoptera frugiperda* (cabbage armyworm) or *Trichoplusia ni* (nettle), and the like. Examples of the cell derived from *S. frugiperda* include Sf9 (ATCC: CRL-1711, ovary cell), Sf21 (ovary cell) and the like. Examples of the cell strain derived from *T. ni* include High Five, BTI-TN-5B1-4 (ovary cell, Invitrogen) and the like.

- 15 As a method for introducing a recombinant vector, any method can be used as far as it is a method which can introduce the vector into a host. Examples include a calcium phosphate method (JP-A No.2-22705), a lipofection method (Proceedings of the National Academy of Sciences USA, 84, 7413 (1987)) and the like. In addition, as in the animal cell, an
20 electroporation method (Cytotechnology, 3, 133 (1990)) and the like can be also used.

[0050]

(vi) Culturing method

- 25 When a transformant harboring a recombinant vector with a DNA encoding the present protein incorporated therein is a cell such as *Escherichia coli*, yeast, an animal cell a plant cell or the like, a cell is cultured according to a conventional culturing method suitable for various hosts, the protein is made to be produced and accumulated, and the protein is recovered from a transformant or a culture solution, thereby, the protein can be
30 produced. When a transformant is an animal or plant individual, an individual is reared or cultivated according to a conventional growing method suitable for various hosts, the protein is made to be produced and accumulated, and the protein is recovered from the animal or plant individual, thereby, the protein can be produced.

- 35 When a host is an animal individual, for example, a non-human

transgenic animal harboring the present gene is reared, the present protein encoded by the recombinant DNA is made to be produced and accumulated in the animal, and the protein is recovered from the animal individual, thereby, the present protein can be produced. Examples of a place of production and accumulation in an animal individual include milk, saliva, egg, and the like of the animal.

When a host is a plant individual, for example, a transgenic plant harboring a gene encoding the present protein is cultivated, the present protein encoded by the recombinant DNA is made to be produced and accumulated in the plant individual, and the polypeptide is recovered from the plant individual, thereby, the present protein can be produced.

When a host is a prokaryote such as *Escherichia coli* and the like, or a eukaryote such as yeast and the like, for example, a transformant harboring a gene encoding the present protein is cultured in a medium, the present protein encoded by the recombinant DNA is made to be produced and accumulated in a culture solution, and the protein is recovered from the culture solution, thereby, the present protein can be produced.

A method for culturing a transformant harboring a gene encoding the present protein in a medium can be performed according to a conventional method used in culturing a host.

As a medium for culturing the resulting transformant obtained by using a prokaryote such as *Escherichia coli* and the like or a eukaryote such as yeast and the like as a host, any of a natural medium and a synthetic medium may be used as far as it is a medium containing a carbon source, a nitrogen source, inorganic salts or the like which can be utilized by the organism, and which can effectively perform culturing of the transformant.

[0051]

When a transformant is a prokaryote such as *Escherichia coli* and the like or a eukaryote such as yeast and the like, as a medium for culturing the resulting transformant, any of a natural medium and a synthetic medium may be used as far as it is a medium containing a carbon source, a nitrogen source, inorganic salts or the like which can be utilized by a host, and which can effectively perform culturing of the transformant. As a medium for culturing a transformant when a host is *Escherichia coli*, for example, a YT medium containing Bacto triptone, yeast extract and sodium chloride is preferable.

[0052]

As the carbon source, a carbon source which can be utilized by each microorganism can be used. Examples include hydrocarbons such as glucose, fructose, sucrose, molasses containing them, starch, starch hydrolysate and the like; organic acids such as acetic acid, propionic acid and the like, and alcohols such as ethanol, propanol and the like.

As the nitrogen source, ammonia, various ammonium salts of inorganic or organic acids such as ammonium chloride, ammonium sulfate, ammonium acetate, ammonium phosphate and the like, other nitrogen-containing substances, peptone, meat extract, yeast extract, corn steep liquor, casein hydrolysate, soybean cake, soybean cake hydrolysate, various fermentation bacterial cells and digested substances thereof and the like can be used.

As the inorganic salt, primary potassium phosphate, secondary potassium phosphate, magnesium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, copper sulfate, calcium carbonate and the like can be used. Culturing is performed under the aerobic condition, such as shaking culturing, deep bubbling stirring culturing and the like.

[0053]

A culturing temperature is suitably 15 to 40°C, and a culturing time is usually 5 hours to 7 days. pH during culturing is retained at 3.0 to 9.0. pH is adjusted with an inorganic or organic acid, an alkaline solution, urea, calcium carbonate, ammonia or the like. And, if necessary, an antibiotic such as ampicillin, tetracycline, kanamycin and the like may be added to a medium during culturing.

When a microorganism transformed with an expression vector using an inducible promoter as a promoter is cultured, if necessary, an inducer may be added to a medium. For example, when a transformant transformed with an expression vector using a lac promoter is cultured, isopropyl-β-D-thiogalactopyranoside or the like may be added to a medium. When a transformant transformed with an expression vector using a trp promoter is cultured, indoleacrylic acid or the like may be added to a medium. A cell or organ of a plant with a gene introduced therein can be cultured at a large scale using a jar fermenter. As a medium for culturing, a medium generally used such as a Murashige and Skoog (MS) medium, a White medium,

a medium in which a plant hormone such as auxin, cytokinin and the like is added to these medium, or the like, can be used.

[0054]

When a transformant for producing the present protein is an animal cell, as a medium for culturing the cell, a medium generally used such as a RPMI1640 medium (The Journal of the American Medical Association, 199, 519 (1967)), a MEM medium (Science, 130, 432 (1959)), a D-MEM medium (Virology, 8, 396 (1959)), a 199 medium (Proceedings of the Society for the Biological Medicine, 73, 1 (1950)), a medium in which bovine fetal serum (FCS) or the like is added to these medium, or the like, can be used.

Culturing is usually performed for 1 to 7 days under the condition such as pH 6 to 8, 25 to 40°C, the presence of 5% CO₂ and the like. In addition, if necessary, an antibiotic such as kanamycin, penicillin, streptomycin and the like may be added to a medium during culturing.

When a transformant is an insect cell, as a medium for culturing, a TNM-FH medium (Pharmingen), a Sf-900II SFM medium (Invitrogen), ExCell400, ExCell405 (JRH Bioscience), Grace's Insect Medium (Nature, 195, 788 (1962)) and the like can be used.

[0055]

(vii) Process of production

The present protein can be produced by culturing a transformant, and isolating/purifying the present protein from a culture solution. A method for isolating/purifying the present protein can be performed by a conventional method well-known in the art. For example, a method for isolating/purifying an enzyme or a method for purifying glucotransferase of Sandler et al. (Methods in Enzymology, 83, 458) can be used.

When the present protein is produced/accumulated as a soluble polypeptide, a culture solution in which a transformant has been cultured as described above is separated into cells or bacterial cells and a medium, for example, by a method such as centrifugation and the like. When the present protein is in a host cell, after collected cells or bacterial cells are washed with a suitable buffer such as a STE solution and the like, cells or bacterial cells are lysed with sonication, a French press, a Manton-Gaulin homogenizer, a Dymomill or the like, and the present protein can be obtained as a cell-free solution by centrifugation or filtration.

A buffer used for separating/purifying the present protein may contain a suitable amount of a surfactant, for example, sodium laurylsulfate (SDS), N-lauroylsarcosine sodium (Sarcosil) or the like.

A method for separating/purifying an objective protein contained in the resulting crude product can be performed by combining the known per se various separating/purifying methods. Examples of these known methods include a solvent extraction method, a salting out method using ammonium sulfate or the like, a dialysis method, a precipitation method using an organic solvent, a ultrafiltration method, a gel filtration, various chromatography methods such as diethylaminoethyl (DEAE)-sepharose chromatography, anion chromatography or ion-exchange chromatography using a resin such as DIAION HPA-75 (Mitsubishi Chemical Corporation) and the like, cation chromatography using lysine such as S-Sepharose FF (Pharmacia) and the like, hydrophobic chromatography using butyl sepharose or the like, and affinity chromatography, various electrophoreses such as SDS-polyacrylamide gel electrophoresis and isoelectric focusing, and the like.

[0056]

When the present protein is produced/accumulated as an insoluble polypeptide, cells or bacterial cells are separated as described above, and lysed by a suitable method, and a fraction containing the polypeptide is recovered. The recovered sample is solubilized with a solubilizer such as a surfactant such as sodium laurylsulfate (SDS), N-lauroylsarcosine sodium (Sarcosil) and the like. The solubilized solution is diluted or dialyzed to a concentration at which little or no solubilizer is contained, the polypeptide is reconstituted into a normal steric structure, thereafter, a purified specimen can be obtained by a separating/purifying method similar to that described above.

[0057]

Alternatively, the present protein is produced as a fused protein with other protein, and this can be also purified utilizing affinity chromatography using a substance having affinity for the fused protein (Akio Yamakawa, Experimental Medicine, 13, 469-474 (1995)). Examples of an addition protein to be used in a fused protein include Protein A, FLAG and the like (Proceedings of the National Academy of Sciences USA, 86, 8227 (1989), Genes Development, 4, 1288 (1990), JP-A No.5-336963, and JP-A No.6-823021).

When Protein A is used, a fused protein of the present protein and Protein A is produced, and this can be purified by performing affinity chromatography using immunoglobulin G. When a FLAG peptide is used, a fused protein of the present protein and FLAG is produced, and this can be purified by affinity chromatography using an anti-FLAG antibody.

[0058]

The present protein can be produced using an in vitro transcription/translation system according to the known method (Journal of Biomolecular NMR, 6, 129-134 (1995), Science, 242, 1162-1164 (1988), The Journal of Biochemistry, 110, 166-168 (1991)).

The present protein can be chemically synthesized by a chemical synthesis method such as a Fmoc method (fluorenylmethyloxycarbonyl method), a Boc method (t-butyloxycarbonyl method) and the like, or with a commercially available peptide synthesizer, for example, a peptide synthesizer such as APEX396 (Advanced Chemtech), 443A (Applied Biosystems), PS3 (Protein Technologies), 9050 (Perceptive), PSSM-8 (Shimadzu Corporation) and the like.

Structural analysis of the present protein can be performed by a method which is usually used in protein chemistry, for example, the method described in Protein Structural Analysis for Gene Cloning (Hisashi Hirano, published by Tokyo Kagaku Dojin Co., Ltd., 1993). Hydroxylation activity of the present protein can be detected as an amount of production of an N-(5-hydroxy-2-adamantyl)-benzamide derivative by acting the present protein on a compound having an adamantane skeleton, for example, an N-(2-adamantyl)-benzamide derivative which is a substrate. More particularly, a method for culturing a microorganism capable of expressing the present protein in a suitable medium, and detecting an accumulation amount of an N-(5-hydroxy-2-adamantyl)-benzamide derivative accumulated in a medium can be used. Detection of an N-(5-hydroxy-2-adamantyl)-benzamide derivative can be performed, for example, by HPLC using an N-(5-hydroxy-2-adamantyl)-benzamide derivative obtained by the present invention, as a standard product. Specifically, the method described in Example described later can be used.

[0059]

(vii) Process for making variant polypeptide

Deletion or substitution of an amino acid of the present protein can be performed by a site-directed mutagenesis method which is the technique well-known before filing. Deletion or substitution of one to several amino acids can be prepared according to the methods described in Molecular Cloning: A Laboratory Manual, Second Edition (1989) (Cold Spring Harbor Laboratory Press), Current Protocols in Molecular Biology (1994) (Wiley-Interscience), Nucleic Acids Research, 10 6487 (1982), Proceedings of the National Academy of Sciences USA, 79, 6409 (1982)), Gene 34, 315 (1985), Nucleic Acids Research, 13 4431 (1985), Proceedings of the National Academy of Sciences USA, 82, 488 (1985)), Proceedings of the National Academy of Sciences USA, 81, 5662 (1984)), Science 224, 1431 (1984), WO 85/00817, Nature, 316, 601 (1985) and the like.

[0060]

(b) Microorganism capable of expressing present protein

A microorganism which can be used in the present process is not particularly limited as far as it is a microorganism capable of expressing the present protein in the state where it can act with a substrate upon acting with a substrate. Examples include a transformant with the present gene introduced therein and a natural microorganism known to express the present protein such as microorganisms belonging to genus *Bacillus*, preferably *Bacillus subtilis*, genus *Pseudomonas*, genus *Streptomyces* and the like. The microorganism is preferably a transformant in that an amount of the present protein contained in a microorganism can be artificially increased.

[0061]

A host cell which can be used for making various transformants usable in the present process is not particularly limited as far as it can express the present protein in the state where it can act with a substrate. Examples include the known microorganisms which are usually used such as *Escherichia coli* or *Saccharomyces cerevisiae*, and the known cultured cell such as vertebrate cell (e.g. CHO cell or COS cell) and an insect cell. *Escherichia coli* is preferable in that a proliferation rate is great, and handling is easy. In addition, an expression vector usable for making the transformant is not particularly limited as far as it can express the present protein in the state where it can act with a substrate, and can be arbitrarily selected depending on a kind of a host cell used.

The host cell, the expression vector, the transformant, the method for culturing a transformant and the like are described in detail in the above "(2) Process for producing present protein".

[0062]

5 In the present process, when a substrate and a microorganism capable of expressing the present protein are acted, a resting bacterial cell reaction or growing bacterial cell reaction may be used. Each of them can utilize the known method. The resting bacterial cell reaction is a method for culturing a microorganism capable of expressing the present protein, then stopping
10 proliferation of a bacterial cell, and adding a substrate. For example, the method described in Examples 1 to 6 described later can be utilized. The growing bacterial cell reaction is a method for adding a substrate while a microorganism capable of expressing the present protein is cultured. For example, the method described in Example 7 or 8 described later can be
15 utilized.

[0063]

In the present process, a reaction condition when a substrate and at least one of:

- (a) the present protein or
 - 20 the present protein and a reducing system consisting of ferredoxin and reductase supplying an electron to the protein, or
 - (b) a microorganism capable of expressing the present protein
- are reacted, is preferably as follows.

A reaction temperature can be arbitrarily determined by considering
25 an optimal temperature for the present protein, a temperature for stabilizing a substrate and product, and the like. For example, the reaction can be performed at 10 to 45°C, preferably 15 to 40°C, further preferably 20 to 40°C, and particularly preferably 25 to 35°C.

A reaction pH can be arbitrarily determined by considering an optimal
30 pH for the present protein, pH for stabilizing a substrate and product, and the like. For example, the reaction can be performed at 4 to 10, preferably 5 to 9, further preferably 6 to 9, and particularly preferably 7 to 8.

A reaction time can be arbitrarily determined depending on the above various conditions (i.e. a solvent used, a reaction temperature or a reaction
35 pH), or a kind of a substrate, the present protein or a microorganism used.

For example, it is 1 to 100 hours, preferably 5 to 48 hours and more preferably 12 to 36 hours.

An amount of the present protein used is not particularly limited, but the reaction can be performed at a concentration of, for example, 0.5 to 20 $\mu\text{mol/L}$.

[0064]

In addition, in the present process, when a substrate and the present protein or a microorganism capable of expressing the present protein are acted, an inclusion agent may be optionally added. Chemical stability or solubility of a substrate in water can be increased, or substrate inhibition on an enzyme or microorganism can be decreased. Examples of the inclusion agent include hydroxypropyl- β -cyclodextrin, α -cyclodextrin, β -cyclodextrin, γ -cyclodextrin, and methyl- β -cyclodextrin (Me- β -CD).

[0065]

In the present process, when a substrate and the present protein or a microorganism capable of expressing the present protein are acted, a coenzyme (e.g. NADPH or NADH, preferably NADPH) can be optionally coexistent and, further, a regeneration system of the coenzyme can be coexistent. In the present process, by the coexistence of a suitable coenzyme depending on a kind of the present protein used, reactivity thereof can be improved.

[0066]

In addition, since a coenzyme is consumed with progression of a reaction (e.g. NAD(P)H is consumed, and converted into NAD(P)+), reactivity thereof can be improved by the coexistence of a degeneration system of the coenzyme. As the coenzyme regeneration system, various regeneration systems are known for each coenzyme. Examples of a regeneration system of NAD(P)H include a combination of glucose and glucose dehydrogenase; formic acid and formic acid dehydrogenase; and propanol and alcohol dehydrogenase; and the like.

[0067]

An addition amount of these coenzymes or the coenzyme regeneration system which can be coexistent in the present process can be arbitrarily determined depending on a kind of a substrate and the present protein or a microorganism used.

[0068]

A reaction is stopped, and a hydroxylated form of a compound having an adamantane skeleton such as an N-(5-hydroxy-2-adamantyl)-benzamide derivative may be isolated according to a general method for collecting a fermentation product from the reaction product. Specifically, a hydroxylated form of a compound having an adamantane skeleton can be isolated from a medium by solvent extraction, ion-exchange chromatography, active carbon treatment, recrystallization, membrane separation or the like.

[0069]

10 The present invention includes an enzyme group utilized for performing the reaction. Examples include the following proteins.

-A protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence of SEQ ID No.: 1, and having hydroxylation activity.

15 -A protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, L357P, A362T, F41L, I77W, M105I, T234A, F232I, F232L and F232M in the amino acid sequence of SEQ ID No.: 1.

-A protein consisting of the amino acid sequence of SEQ ID No.: 3, or a protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence, and having hydroxylation activity.

20 -A protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3.


These proteins can be obtained by the method described in the above "Process for producing present protein".

[0070]

30 In addition, the present invention includes the following compounds. These compounds are useful for producing an N-(5-hydroxy-2-adamantyl)-benzamide derivative which is a raw material of monohydroxy-2-adamantanamine useful as an intermediate of functional resins or medicines.

35 A compound of the formula:

[Formula 13]




a salt or solvate thereof.

[0071]

5 In addition, the present invention includes the following compounds. These compounds are useful for producing monohydroxy-2-adamantamine useful as an intermediate of functional resins or medicines.

A compound of the formula:

[Formula 14]



10

a salt or solvate thereof.

[0072]

Examples of "salt" include salts of inorganic acids such as hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid and the like; salts of organic acids such as paratoluenesulfonic acid, methanesulfonic acid, oxalic acid, citric acid and the like; salts of organic bases such as ammonium, trimethylammonium, triethylammonium and the like; salts of alkali metals such as sodium, potassium and the like; and salts of alkaline earth metals such as calcium, magnesium and the like.

20

[0073]

"Solvate" is such that the aforementioned compound may be coordinated with an arbitrarily number of solvent molecules. Preferable is a hydrate.

[0074]

The present invention will be explained in more detail by way of Examples, but these are not limit the present invention.

As a gene manipulation procedure, unless otherwise indicated, the method described in Molecular Cloning: A Laboratory Manual, Second Edition (Cold Spring Harbor Laboratory) was used.

Example 1

[0075]

10 Hydroxylation of adamantane by resting bacterial cell reaction using CYP109B1

[Seed culture]

An expression vector (pET-109B1-CamA-CamB) with a CYP109B1 gene (SEQ ID No.: 2) inserted therein was introduced into Escherichia coli (BL21(DE3)) to obtain a transformant. The transformant was inoculated in 2 ml of a LB medium, and cultivated at 37°C on reciprocating shaker at 300 rpm for 24 hours to obtain a seed starter.

[Main culture]

A M9 mix medium (100 ml) (6.78 g/L Na₂HPO₄ (anhydrous), 3 g/L KH₂PO₄, 0.5 g/L NaCl, 1 g/L NH₄Cl, 4 g/L casamino acid, 100 µM CaCl₂) was sterilized and, to this were added 100 µl of 100 mM FeSO₄ and 1 ml of 2 mg/ml thymine. To this was added Overnight Auto Induction System manufactured by Merck (Solution I 2 ml, Solution II 5ml, Solution III 100 µl) and, further, 100 µl of 50 mg/ml carbenicillin, and 100 µl of 80 mg/ml 5-aminolevulinic acid were added to obtain a main culture medium. 500 µl of a seed starter was seeded in a 500 ml wide-mouthed Erlenmeyer flask containing 50 ml of a main culture medium, and this was cultured at 25°C and 120 rpm for 24 hours.

[Biocatalytic reaction]

The culture solution was centrifuged to remove the supernatant to obtain bacterial cells. The bacterial cells were diluted with 25 ml of a potassium phosphate buffer (50 mM potassium phosphate (pH 7.4), 10% glycerol, 1mM EDTA, 2 mM DTT, 1 mM D-glucose) so that a bacterial cell concentration in terms of a culture solution became 1-fold, 2.5-fold, and 5-fold, respectively, and placed into a 500 ml Erlenmeyer flask with baffles to obtain a reaction solution. To this reaction solution was added

N-(adamantan-2-yl)-phthalamidic acid to a final concentration of 5000 µg/ml, pH 7.4 to initiate a reaction. The reaction was performed at 28°C and 180 rpm for 31 hours, an equivalent amount of acetone was added, and this was analyzed by reverse-phase HPLC. As a result, a substrate was converted into N-(5-hydroxy-adamantan-2-yl)-phthalamidic acid at a conversion rate described in Table 1.

[Table 1]

Bacterial cell concentration	1-fold	2.5-fold	5-fold
Substrate 5000 µg/ml	61%	100%	100%

Example 2

[0076]

Hydroxylation of adamantane by resting bacterial cell reaction using CYP109FK

[Making of vector]

Referring to Applied Microbiology and Biotechnology, 2006, vol. 71, p.455-462, an expression vector of a fused protein (109FK) of CYP109B1 and a P450RhF reduction domain was prepared. Specifically, for constructing a 109FK gene (SEQ ID No.: 4), an Overlap Extension PCR method was used. PCR was performed employing pET-109B1-CamA-CamB as a template, and using a primer 1 (SEQ ID No.: 7) and a primer 2 (SEQ ID No.: 8), and PCR was performed by employing a genome of genus Rhodococcus NCIMB9784 strain as a template, and using a primer 3 (SEQ ID No.: 9) and a primer 4 (SEQ ID No.: 10). Then, PCR was performed by employing each PCR product as a template, and using primers 1 and 4, and the PCR product was inserted into a cloning vector (pCR-BluntII-TOPO) (pCR-109FK). A fragment containing a 109FK gene was excised with XbaI and BamHI from the pCR-109FK vector, and inserted into an expression vector pET21a to prepare an expression vector of 109FK (pET-109FK).

[Seed culture]

Into Escherichia coli (BL21(DE3)) was introduced pET-109FK, to obtain a transformant. The transformant was seeded into a 96-well deep well plate containing 500 µl of a LB medium per well, and cultured at 900 rpm and 37°C for 24 hours to obtain a seed starter.

[Main culture]

Using the main culture medium described in Example 1, the seed starter was inoculated in a 96-well deep well plate containing 500 μ l of the main culture medium per well at 5 μ l per well, and this was cultured at 25°C and 900 rpm for 24 hours.

5 [Biocatalytic reaction]

A culture solution of the 96-well deep well plate was centrifuged to remove the supernatant, to obtain bacterial cells. The bacterial cells were diluted with 100 μ l of a potassium phosphate buffer (50 mM potassium phosphate (pH 7.4), 10% glycerol, 1 mM EDTA, 2 mM DTT, 1 mM D-glucose) containing 1 mg/ml N-(adamantan-2-yl)-phthalamidic acid, to start a reaction. A reaction was performed at 28°C and 900 rpm for 24 hours. An equivalent amount of acetone was added, this was analyzed by reverse-phase HPLC and, as a result, 18% of a substrate was converted into N-(5-hydroxy-adamantan-2-yl)-phthalamidic acid.

15

Example 3

[0077]

Hydroxylation of adamantane by resting bacterial cell reaction using CYP109FK mutant

20 [Introduction of random mutation]

For introducing a random mutation into a 109FK gene, an Error Prone PCR method was used. Specifically, the Error Prone PCR was performed employing pET-109FK as a template, using primers 1 and 4, and using GeneMorphII Random Mutagenesis Kit manufactured by Stratagene, according to the protocol, so that a mutation frequency became 0 to 4.5/kb. The resulting PCR fragment was restriction enzyme-treated with XbaI and HindIII, and inserted into pET21a to prepare an expression vector (pET-109FKm).

[Screening]

30 Into Escherichia coli (BL21(DE3)) was introduced pET-109FKm, to obtain a transformant. 2296 colonies were collected as the transformant, each colony was inoculated in a 96-well deep well plate containing 500 μ l of a LB medium per well, and this was cultured at 900 rpm and 37°C for 24 hours to obtain a seed starter.

35 Using the main culture medium described in Example 1, the seed

starter was inoculated in a 96-well deep well plate containing 500 µl of the main culture medium per well, at 5 µl per well, and this was cultured at 25°C and 900 rpm for 24 hours.

A culture solution of the 96-well deep well plate was centrifuged to remove the supernatant, to obtain bacterial cells. The bacterial cells were diluted with 1 ml of a potassium phosphate buffer (50 mM potassium phosphate (pH 7.4), 10% glycerol, 1 mM EDTA, 2 mM DTT, 1 mM D-glucose) containing 1 mg/ml N-(adamantan-2-yl)-phthalamidic acid, to start a reaction. A reaction was performed at 28°C and 900 rpm for 24 hours. An equivalent amount of acetone was then added, and the resultant was analyzed by reverse-phase HPLC. As a result, 91 colonies had activity which was improved 1.50-fold or more than that of a transformant with pET-109FK introduced therein. A plasmid was extracted from colonies, and a mutation position was identified by sequencing. Regarding I77 and F232 among resulting mutations, substitution with other amino acid was performed by site-directed mutation using QuickChange II Site-Directed Mutagenesis Kit manufactured by Stratagene. Activity of each mutant was confirmed by the same method as that of screening and, as a result, it was found out that mutations described in Table 2 improved activity as compared with other mutations (e.g. mutations described in Table 3).

[Table 2]

	Conversion rate	Relative ratio relative to 109FK
109FK	12.9	1.00
I77F	67.8	5.26
I77W	67.8	5.27
M105I	28.0	2.17
A196D	21.2	1.64
F232I	31.7	2.47
F232L	38.7	3.01
F232M	31.7	2.46
T234A	26.3	2.04
T244A	21.8	1.70
V399E	19.0	1.48
K702E	21.9	1.70

[Table 3]

	Conversion rate	Relative ratio relative to 109FK
M272T	10.1	0.78
F232K	2.7	0.21

Example 4

[0078]

- 5 Hydroxylation of adamantane by resting bacterial cell reaction using CYP109B1 mutant by random mutation

[Introduction of random mutation]

For introducing a random mutation into a CYP109B1 gene, an Error Prone PCR method was used. Specifically, the Error Prone PCR was performed employing pET-109B1-CamA-CamB as a template, using a primer 1 and a primer 5 (SEQ ID No.: 11), and using GeneMorphII Random Mutagenesis Kit manufactured by Stratagene, according to the protocol, so that a mutation frequency became 0 to 4.5/kb. The resulting PCR fragment was restriction enzyme-treated with XbaI and KpnI, and inserted into a 7kb fragment of pET-109B1-CamA-CamB which had been restriction enzyme-treated with XbaI and KpnI to prepare an expression vector (pET-109B1m-CamA-CamB).

[Screening]

Into Escherichia coli (BL21(DE3)) was introduced pET-109B1m-CamA-CamB, to obtain a transformant. 880 colonies were collected as the transformant, each colony was inoculated in a 96-well deep well plate containing 500 µl of a LB medium per well, and this was cultured at 900 rpm and 37°C for 24 hours to obtain a seed starter.

Using the main culture medium described in Example 1, the seed starter was inoculated in a 96-well deep well plate containing 500 µl of the main culture medium per well, at 5 µl per well, and this was cultured at 25°C and 900 rpm for 24 hours.

A culture solution of the 96-well deep well plate was centrifuged to remove the supernatant to obtain bacterial cells. The bacterial cells were diluted with 500 µl of a potassium phosphate buffer (50 mM potassium phosphate (pH 7.4), 10% glycerol, 1 mM EDTA, 2 mM DTT, 1 mM D-glucose) containing 1 mg/ml N-(adamantan-2-yl)-phthalamidic acid, to start a reaction.

A reaction was performed at 28°C and 900 rpm for 2 hours. An equivalent amount of acetone was then added, the resultant was analyzed by reverse-phase HPLC. As a result, 16 colonies had activity which was improved 1.23-fold or more than that of a transformant with pET-109FK introduced therein. A plasmid was extracted from colonies, and a mutation position was identified by sequencing. Activity of each mutation was reconfirmed by the same method as that of screening and, as a result, it was found out that mutations described in Table 4 improved activity as compared with other mutations (e.g. mutations described in Table 5).

[Table 4]

	Conversion rate	Relative ratio relative to 109B1
109B1	23.2	1.00
I77F	33.5	1.44
L357P	30.3	1.31
A362T	27.1	1.17
F41L	26.7	1.15

[Table 5]

	Conversion rate	Relative ratio relative to 109B1
E242K	14.3	0.62
Q46R	23.2	1.00

Example 5

[0079]

Hydroxylation of adamantane by resting bacterial cell reaction using CYP109B1 mutant with mutation of CYP109FK mutant introduced therein

A part of mutation of the 109FK gene found out in Example 3 was introduced into a CYP109B1 gene, and activity was measured by the method for screening of Example 4. In this respect, bacterial cells were diluted with 1 ml of a potassium phosphate buffer (50 mM potassium phosphate (pH 7.4), 10% glycerol, 1 mM EDTA, 2 mM DTT, 1 mM D-glucose) containing 3 mg/ml N-(adamantan-2-yl)-phthalamidic acid, and a reaction time was changed to 16 hours. As a result, it was found out that mutations described in Table 6 improved activity.

[Table 6]

	Conversion rate	Relative ratio relative to 109B1
109B1	47.9	1.00
I77F	70.9	1.48
I77W	61.2	1.28
M105I	57.5	1.20
T234A	55.8	1.17

On the other hand, confirmation of activity of the CYP109B1 gene in which mutation was introduced into F232 was performed by the method for screening of Example 4. As a result, it was found out that mutations described in Table 7 improved activity.

[Table 7]

	Conversion rate	Relative ratio relative to 109B1FK
109B1	9.6	1.00
F232I	14.9	1.55
F232L	14.8	1.55
F232M	15.7	1.64

Example 6

[0080]

Hydroxylation of adamantane by resting bacterial cell reaction using CYP109B1 mutant (I77F)

[Seed culture]

An expression vector (pET-109B1 (I77F)-CamA-CamB) with a CYP109B1 mutant (I77F) gene (SEQ ID No.: 5) inserted therein was introduced into Escherichia coli (BL21(DE3)) to obtain a transformant. The transformant was inoculated in 2 ml of LB medium, and cultured with a 300 rpm reciprocating shaker at 37°C for 24 hours to obtain a seed starter.

[Main culture]

Main culture was performed as in Example 1.

[Biocatalytic reaction]

A Biocatalytic reaction was performed as in Example 1. After the reaction, an equivalent amount of acetone was added, the resultant was analyzed by reverse-phase HPLC and, as a result, a substrate was converted

into N-(5-hydroxy-adamantan-2-yl)-phthalamidic acid at a conversion rate in Table 8.

[Table 8]

Bacterial cell concentration	1-fold	2.5-fold	5-fold
Substrate 5000 µg/ml	88%	100%	100%

5 [Purification]

After the reaction solution as a whole was centrifuged, 113 ml of the supernatant was obtained. The supernatant (40 ml) was adsorbed onto 5 ml of HP20SS, and this was 2CV-washed with a 0.1% aqueous formic acid solution, and eluted with 0.1% formic acid, and a 50% aqueous acetone solution to obtain an active fraction. The active fraction was concentrated, and lyophilized to obtain 125 mg of a powder of N-(5-hydroxy-adamantan-2-yl)-phthalamidic acid. A structure was analyzed by NMR and, as a result, since NOE was observed between H of 3.92, and H of 1.71 and 1.62, it was found out that the structure was Anti entity 100%.

1H NMR (DMSO-d₆) δ : 8.40(1H, br. s), 7.74 (1H, br. s), 7.522 (1H, td, J=7.5, 1.1Hz), 7.461 (1H, td, J=7.5, 1.1Hz), 7.40 (1H, br. d, J= ~8Hz), 4.40 (1H, br. s), 3.92 (1H, m), 2.05 (2H, br. s), 1.97 (2H, m), 1.96 (1H, m), 1.71 (2H, br. d), 1.62 (2H, br. d, J=~12Hz), 1.61 (1H, br. s), 1.28 (2H, br. d, J=~12Hz)

20 Example 7

[0081]

Hydroxylation of adamantane by growing bacterial cell reaction using CYP109B1 mutant (I77F)

[Seed culture]

25 A expression vector (pET-109B1 (I77F)-CamA-CamB) with a CYP109B1 mutant (I77F) gene (SWQ ID No.: 5) inserted therein was introduced into Escherichia coli (BL21(DE3)) to obtain a transformant.

One platinum loop of the transformant was seeded in a 500 ml wide-mouthed Erlenmeyer flask containing 50 ml of LB medium, and cultured at 37°C and 180 rpm for 5.5 hours to obtain a seed starter.

[Main culture]

1 L of a L medium (10 g/L triptone, 5 g/L yeast extract, 5 g/L NaCl) was sterilized, to this were added 1 ml of 100 mM FeSO₄, 20 g of glycerol, 30 g of

hydroxypropyl- β -cyclodextrin, 10 ml of 50 mg/ml carbenicillin, 10 ml of 80 mg/ml 5-aminolevulinic acid, 10 ml of 2 mg/ml thymine, 0.1 g of $MgCl_2$, and 0.1 ml of Adecanol LG121, and N-(adamantan-2-yl)-phthalamidic acid which is a substrate was added to a final concentration of 5000 $\mu g/ml$, to obtain a main culture medium. A seed starter (40 ml) was seeded in 3 L-minijar containing 1 L of the main culture medium, and culturing was initiated by controlling at 28°C, 600 rpm and pH 6.5. Six hours after culturing initiation, IPTG was added to a final concentration of 1 mM. During culturing, casamino acid, glycerol, and a substrate were arbitrarily added, and an amount of a finally added substrate reached 30 g/L. Seventy-two hours after reaction initiation, the reaction was analyzed by reverse-phase HPLC and, as a result, 67% of the substrate was converted into N-(5-hydroxy-adamantan-2-yl)-phthalamidic acid.

Example 8

[0082]

Hydroxylation of adamantane by growing bacterial cell reaction using CYP109B1 mutant (I77F) and production of monohydroxy-2-adamantanamine [Production of vector]

An *Escherichia coli* JM109 strain-derived glycerol dehydrogenase gene (SEQ ID No.: 6) was inserted into pCDFDuet to construct an expression vector (pCDFD-GLD).

[Seed culture]

Into *Escherichia coli* (BL21(DE3)) were introduced pCDFD-GLD, and pET-109B1 (I77F)-CamA-CamB described in Example 7, to obtain a transformant. One platinum loop of the transformant was seeded in a 500 ml wide-mouthed Erlenmeyer flask containing 50 ml of LB medium, and cultured at 37°C and 180 rpm for 7 hours to obtain a seed starter.

[Main culture]

1 L of a L medium/M9 (10 g/L triptone, 5 g/L yeast extract, 6.78 g/L Na_2HPO_4 (anhydrous), 3 g/L KH_2PO_4 , 5.5 g/L NaCl, 1 g/L NH_4Cl) was sterilized, to this were added 1 ml of 100 mM $FeSO_4$, 20 g of glycerol, 30 g of hydroxypropyl- β -cyclodextrin, 1 ml of 50 mg/ml carbenicillin, 1 ml of 50 mg/ml streptomycin, 1 ml of 80 mg/ml 5-aminolevulinic acid, and 10 ml of 2 mg/ml thymine. Further, Overnight Auto Induction System manufactured by Merck

(Solution I 20 ml, Solution II 50 ml, Solution III 1 ml) was added, and N-(adamantan-2-yl)-phthalamidic acid which is a substrate was added to a final concentration of 5000 µg/ml, to obtain a main culture medium. A seed starter (40 ml) was seeded in 3 L-minijar containing 1 L of the main culture medium, and culturing was initiated by controlling at 28°C, 500 rpm, and pH 6.5. During culturing, casamino acid, glycerol, and a substrate were arbitrarily added, and an amount of a finally added substrate reached 30 g/L. Seventy-two hours after reaction initiation, the reaction was analyzed by reverse-phase HPLC and, as a result, 90% of the substrate was converted into N-(5-hydroxy-adamantan-2-yl)-phthalamidic acid.


[Purification]

A culture solution was adjusted with hydrochloric acid to pH of 4, and sterilized at 120°C for 15 minutes, and a protecting group was deprotected. A pH was adjusted with an aqueous sodium hydroxide solution to 7, and the resultant was centrifuged to obtain 1.65 L of the supernatant. 1 L of the supernatant was adsorbed on 300 ml of an ammonium ion-type Diaion PK208, and this was eluted with 2 N aqueous ammonia to obtain an active fraction. The active fraction was concentrated, lyophilized, dissolved in 100 ml of water, and adsorbed onto 120 ml of hydroxide ion-type Amberlite IRA410. This was eluted with water, and an active fraction was obtained, concentrated, and lyophilized to obtain 9.64 g of a powder of monohydroxy-2-adamantamine. A purity was measured by the FDLA method and, as a result, a purity of monohydroxy-2-adamantamine was 96.6%.

Example 9

[0083]

[Formula 15]



To a solution of Compound 3 (150 mg) in dimethylformamide (DMF) (5

ml) were added monohydroxy-2-adamantanamine (140 mg),
1-hydroxybenzotriazole (HOBT) (31 mg),
1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (WSC) (174
mg), and triethylamine (TEA) (180 ml) under the nitrogen atmosphere, and
5 the mixture was stirred at room temperature for 14 hours. After completion
of the reaction, 2N hydrochloric acid (30 ml) was added, followed by extraction
with ethyl acetate. The organic layer was sequentially washed with an
aqueous saturated sodium bicarbonate solution and an aqueous saturated
sodium chloride solution, and dried over magnesium sulfate. A solvent was
10 removed, and the residue was purified by silica gel chromatography to obtain
Compound 4 (226mg).

NMR: (CDCl₃): 1.06 (d, J=6.6 Hz, 6H), 1.53-2.20 (m, 14H), 3.72 (s, 3H), 3.98 (d,
J=6.6 Hz, 2H), 6.25-6.30 (m, 1H), 7.71 (s, 1H)

15 Industrial applicability [0084]

The present process can be utilized as a process for producing a
hydroxylated form of a compound having an adamantane skeleton useful as an
intermediate for functional resins and medicines, at low cost and with high
20 yield.

Sequence listing free text

[0085]

SEQ ID No.: 1 represents an amino acid sequence of CYP109B1.

25 [0086]

SEQ ID No.: 2 represents a base sequence of a gene encoding
CYP109B1 used in Example 1.

[0087]

SEQ ID No.: 3 represents an amino acid sequence of CYP109FK.

30 [0088]

SEQ ID No.: 4 represents a base sequence of a gene encoding
CYP109FK.

[0089]

SEQ ID No.: 5 represents a base sequence of a gene encoding a
35 CYP109B1 mutant (I77F).

[0090]

SEQ ID No.: 6 represents a glycerol dehydrogenase gene derived from
Escherichia coli JM109 strain.

Claims

[1] A process for producing an N-(5-hydroxy-2-adamantyl)-benzamide derivative characterized by hydroxylating an N-(2-adamantyl)-benzamide derivative using cytochrome P450.

[2] The process of Claim 1,

wherein cytochrome P450 is

a protein consisting of an amino acid sequence of SEQ ID No.: 1 or 3, or

a protein consisting of an amino acid sequence in which one to several amino acids are deleted, substituted, or added in the amino acid sequence, and having hydroxylation activity.


[3] The process of Claim 1,

wherein cytochrome P450 is

a protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, L357P, A362T, F41L, I77W, M105I, T234A, F232I, F232L and F232M in the amino acid sequence of SEQ ID No.: 1, or

a protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3.

[4] The process of Claim 1, wherein the N-(2-adamantyl)-benzamide derivative is a compound of the formula:




wherein R is hydrogen or a hydrophilic group.

[5] The process of Claim 1, wherein the N-(5-hydroxy-2-adamantyl)-benzamide derivative is a compound of the

formula:


[Formula 2]



wherein R is hydrogen or a hydrophilic group.

5 [6] A process for producing a compound of the formula:

[Formula 3]




characterized by obtaining an N-(5-hydroxy-2-adamantyl)-benzamide derivative by the process of any one of Claims 1 to 5, and
10 deprotecting it.

[7] The process of Claim 6, characterized in that the N-(5-hydroxy-2-adamantyl)-benzamide derivative is not isolated.

[8] A process for producing a compound of the formula (III):

[Formula 4]



15

characterized by obtaining a compound of (I) by the process of Claim 6 or 7, and

reacting with a compound of the formula (II): A-R¹-R²-R³-X

wherein A is an optionally substituted cyclic hydrocarbon group or an
20 optionally substituted heterocyclic group, R¹ is a single bond, -C(=O)-, -O- or

NR⁴·, R² is a single bond or optionally substituted alkylene, R³ is a single bond or C(=O)·, X is a hydroxy group, halogen or a leaving group derived from a hydroxy group, and R⁴ is hydrogen or optionally substituted alkyl.

[9] A method for hydroxylating an adamantane part of a compound having an adamantane skeleton, comprising using

a protein consisting of an amino acid sequence of SEQ ID No.: 1 or 3, or
a protein consisting of an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence, and having hydroxylation activity.

[10] A protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence of SEQ ID No.: 1, and having hydroxylation activity.


[11] A protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, L357P, A362T, F41L, I77W, M105I, T234A, F232I, F232L and F232M in the amino acid sequence of SEQ ID No.: 1.

[12] A protein consisting of the amino acid sequence of SEQ ID No.: 3, or
a protein containing an amino acid sequence in which one to several amino acids are deleted, substituted or added in the amino acid sequence, and having hydroxylation activity.

[13] A protein consisting of an amino acid sequence having one to several mutations selected from the group consisting of I77F, I77W, M105I, A196D, F232I, F232L, F232M, T234A, T244A, V399E and K702E in the amino acid sequence of SEQ ID No.: 3.

[14] A compound of the formula:


[Formula 5]



a salt or a solvate thereof.

[15] A compound of the formula:

[Formula 6]



a salt or solvate thereof,

wherein R is hydrogen or a hydrophilic group.

SEQUENCE LISTING

<110> Shionogi & Co.,Ltd.

5

<120> 08P00078WO

<130> A process for the preparation of adamantane hydroxide with
cytochrome P450

10

<160> 11

<170> PatentIn version 3.3

15

<210> 1

<211> 396

<212> PRT

<213> Bacillus subtilis

20

<220>

<222> (2)..(2)

<223> Xaa is Asp or Asn residue.

<220>

25

<222> (113)..(113)

<223> Xaa is Arg or Gln residue.

<400> 1

30

Met Xaa Val Leu Asn Arg Arg Gln Ala Leu Gln Arg Ala Leu Leu Asn

1

5

10

15

Gly Lys Asn Lys Gln Asp Ala Tyr His Pro Phe Pro Trp Tyr Glu Ser

35

20

25

30

Met Arg Lys Asp Ala Pro Val Ser Phe Asp Glu Glu Asn Gln Val Trp

35

40

45

5

Ser Val Phe Leu Tyr Asp Asp Val Lys Lys Val Val Gly Asp Lys Glu

50

55

60

10

Leu Phe Ser Ser Cys Met Pro Gln Gln Thr Ser Ser Ile Gly Asn Ser

65

70

75

80

15

Ile Ile Asn Met Asp Pro Pro Lys His Thr Lys Ile Arg Ser Val Val

85

90

95

Asn Lys Ala Phe Thr Pro Arg Val Met Lys Gln Trp Glu Pro Arg Ile

20

100

105

110

Xaa Glu Ile Thr Asp Glu Leu Ile Gln Lys Phe Gln Gly Arg Ser Glu

115

120

125

25

Phe Asp Leu Val His Asp Phe Ser Tyr Pro Leu Pro Val Ile Val Ile

130

135

140

30

Ser Glu Leu Leu Gly Val Pro Ser Ala His Met Glu Gln Phe Lys Ala

145

150

155

160

35

Trp Ser Asp Leu Leu Val Ser Thr Pro Lys Asp Lys Ser Glu Glu Ala

165

170

175

Glu Lys Ala Phe Leu Glu Glu Arg Asp Lys Cys Glu Glu Glu Leu Ala

5

180

185

190

Ala Phe Phe Ala Gly Ile Ile Glu Glu Lys Arg Asn Lys Pro Glu Gln

195

200

205

10

Asp Ile Ile Ser Ile Leu Val Glu Ala Glu Glu Thr Gly Glu Lys Leu

210

215

220

15

Ser Gly Glu Glu Leu Ile Pro Phe Cys Thr Leu Leu Leu Val Ala Gly

225

230

235

240

20

Asn Glu Thr Thr Thr Asn Leu Ile Ser Asn Ala Met Tyr Ser Ile Leu

245

250

255

Glu Thr Pro Gly Val Tyr Glu Glu Leu Arg Ser His Pro Glu Leu Met

25

260

265

270

Pro Gln Ala Val Glu Glu Ala Leu Arg Phe Arg Ala Pro Ala Pro Val

275

280

285

30

Leu Arg Arg Ile Ala Lys Arg Asp Thr Glu Ile Gly Gly His Leu Ile

290

295

300

35

5
 10
 15
 20
 25
 30
 35

ggggataaag agttgttttc cagttgcatg cgcagcaga caagctctat tggaaattcc 240
 atcattaaca tggaccgcc gaagcataca aaaatccgtt cagtcgtgaa caaagccttt 300
 actccgcgcg tgatgaagca atgggaaccg agaattcgag aaatcacaga tgaactgatt 360
 caaaaatttc aggggcgcag tgagtttgac cttgttcagc attttcata cccgcttcgc 420
 gttattgtga tatctgagtt gctgggagtg cttcagcgc atatggaaca gtttaaagca 480
 tggctcgate ttctggtcag tacaccgaag gataaaagtg aagaagctga aaaagccttt 540
 ttggaagaac gagataagtg tgaggaagaa ctggccgcgt ttttgccgg catcatagaa 600
 gaaaagcgaa acaaaccgga acaggatatt atttctattt tagtggaagc ggaagaaaca 660
 ggcgagaagc tgccgggtga agagctgatt ccgttttgca cgtgctgct ggtggccgga 720
 aatgaaacca ctacaaacct gatttcaaat gcgatgtaca gcatattaga aacgccaggc 780
 gtttacgagg aactgcgcag ccactctgaa ctgatgcctc aggcagtgga ggaagccttg 840
 cgtttcagag cgccggcccc ggttttgagg cgcattgcca agcgggatac ggagatcggg 900
 gggcacctga ttaaagaagg tgatatggtt ttggcgtttg tggcatcggc aaatcgtgat 960
 gaagcaaagt ttgacagacc gcacatgttt gatatccgcc gccatcccaa tccgcatatt 1020
 gcgtttggcc acggcateca tttttgectt ggggccccgc ttgcccgtct tgaagcaaat 1080
 atcgcgttaa cgtctttgat ttctgctttt cctcatatgg agtgcgtcag taccactccg 1140
 attgaaaaca gtgtgatata cggattaaag agcttccgtg tgaaaatgta a 1191

<210> 3

<211> 727

<212> PRT

5 <213> Artificial sequence

<220>

<223> A protein containing fused CYP109B1 and and P450Rhfr reductase domain

10

<400> 3

Met Asp Val Leu Asn Arg Arg Gln Ala Leu Gln Arg Ala Leu Leu Asn

1

5

10

15

15

Gly Lys Asn Lys Gln Asp Ala Tyr His Pro Phe Pro Trp Tyr Glu Ser

20

25

30

20

Met Arg Lys Asp Ala Pro Val Ser Phe Asp Glu Glu Asn Gln Val Trp

35

40

45

25

Ser Val Phe Leu Tyr Asp Asp Val Lys Lys Val Val Gly Asp Lys Glu

50

55

60

Leu Phe Ser Ser Cys Met Pro Gln Gln Thr Ser Ser Ile Gly Asn Ser

30

65

70

75

80

Ile Ile Asn Met Asp Pro Pro Lys His Thr Lys Ile Arg Ser Val Val

85

90

95

35

Asn Lys Ala Phe Thr Pro Arg Val Met Lys Gln Trp Glu Pro Arg Ile

100

105

110

5

Arg Glu Ile Thr Asp Glu Leu Ile Gln Lys Phe Gln Gly Arg Ser Glu

115

120

125

10

Phe Asp Leu Val His Asp Phe Ser Tyr Pro Leu Pro Val Ile Val Ile

130

135

140

Ser Glu Leu Leu Gly Val Pro Ser Ala His Met Glu Gln Phe Lys Ala

15

145

150

155

160

Trp Ser Asp Leu Leu Val Ser Thr Pro Lys Asp Lys Ser Glu Glu Ala

165

170

175

20

Glu Lys Ala Phe Leu Glu Glu Arg Asp Lys Cys Glu Glu Glu Leu Ala

180

185

190

25

Ala Phe Phe Ala Gly Ile Ile Glu Glu Lys Arg Asn Lys Pro Glu Gln

195

200

205

30

Asp Ile Ile Ser Ile Leu Val Glu Ala Glu Glu Thr Gly Glu Lys Leu

210

215

220

Ser Gly Glu Glu Leu Ile Pro Phe Cys Thr Leu Leu Leu Val Ala Gly

35

225

230

235

240

Asn Glu Thr Thr Thr Asn Leu Ile Ser Asn Ala Met Tyr Ser Ile Leu

245

250

255

5

Glu Thr Pro Gly Val Tyr Glu Glu Leu Arg Ser His Pro Glu Leu Met

260

265

270

10

Pro Gln Ala Val Glu Glu Ala Leu Arg Phe Arg Ala Pro Ala Pro Val

275

280

285

15 Leu Arg Arg Ile Ala Lys Arg Asp Thr Glu Ile Gly Gly His Leu Ile

290

295

300

Lys Glu Gly Asp Met Val Leu Ala Phe Val Ala Ser Ala Asn Arg Asp

20

305

310

315

320

Glu Ala Lys Phe Asp Arg Pro His Met Phe Asp Ile Arg Arg His Pro

325

330

335

25

Asn Pro His Ile Ala Phe Gly His Gly Ile His Phe Cys Leu Gly Ala

340

345

350

30

Pro Leu Ala Arg Leu Glu Ala Asn Ile Ala Leu Thr Ser Leu Ile Ser

355

360

365

35 Ala Phe Pro His Met Glu Cys Val Ser Ile Thr Pro Ile Glu Asn Ser

370

375

380

Val Ile Tyr Gly Leu Lys Ser Phe Arg Val Lys Met Glu Phe Val Leu

5

385

390

395

400

His Arg His Gln Pro Val Thr Ile Gly Glu Pro Ala Ala Arg Ala Val

405

410

415

10

Ser Arg Thr Val Thr Val Glu Arg Leu Asp Arg Ile Ala Asp Asp Val

420

425

430

15

Leu Arg Leu Val Leu Arg Asp Ala Gly Gly Lys Thr Leu Pro Thr Trp

435

440

445

20

Thr Pro Gly Ala His Ile Asp Leu Asp Leu Gly Ala Leu Ser Arg Gln

450

455

460

Tyr Ser Leu Cys Gly Ala Pro Asp Ala Pro Ser Tyr Glu Ile Ala Val

25

465

470

475

480

His Leu Asp Pro Glu Ser Arg Gly Gly Ser Arg Tyr Ile His Glu Gln

485

490

495

30

Leu Glu Val Gly Ser Pro Leu Arg Met Arg Gly Pro Arg Asn His Phe

500

505

510

35

Ala Leu Asp Pro Gly Ala Glu His Tyr Val Phe Val Ala Gly Gly Ile

515

520

525

5 Gly Ile Thr Pro Val Leu Ala Met Ala Asp His Ala Arg Ala Arg Gly

530

535

540

Trp Ser Tyr Glu Leu His Tyr Cys Gly Arg Asn Arg Ser Gly Met Ala

10

545

550

555

560

Tyr Leu Glu Arg Val Ala Gly His Gly Asp Arg Ala Ala Leu His Val

565

570

575

15

Ser Glu Glu Gly Thr Arg Ile Asp Leu Ala Ala Leu Leu Ala Glu Pro

580

585

590

20

Ala Pro Gly Val Gln Ile Tyr Ala Cys Gly Pro Gly Arg Leu Leu Ala

595

600

605

25 Gly Leu Glu Asp Ala Ser Arg Asn Trp Pro Asp Gly Ala Leu His Val

610

615

620

Glu His Phe Thr Ser Ser Leu Ala Ala Leu Asp Pro Asp Val Glu His

30

625

630

635

640

Ala Phe Asp Leu Glu Leu Arg Asp Ser Gly Leu Thr Val Arg Val Glu

645

650

655

35

Pro Thr Gln Thr Val Leu Asp Ala Leu Arg Ala Asn Asn Ile Asp Val

660

665

670

5

Pro Ser Asp Cys Glu Glu Gly Leu Cys Gly Ser Cys Glu Val Ala Val

675

680

685

10 Leu Asp Gly Glu Val Asp His Arg Asp Thr Val Leu Thr Lys Ala Glu

690

695

700

Arg Ala Ala Asn Arg Gln Met Met Thr Cys Cys Ser Arg Ala Cys Gly

15

705

710

715

720

Asp Arg Leu Ala Leu Arg Leu

725

20

<210> 4

<211> 2184

<212> DNA

25

<213> Artificial sequence

<220>

<223> A gene coding a protein containing fused CYP109B1 and P450Rhfr
reductase domain

30

<400> 4

atggatgtgt taaaccgccg gcaagccttg cagcgagcgc tgctcaatgg gaaaaacaaa 60

caggatgcgt atcatccgtt tccatgggtat gaatcgatga gaaaggatgc gcctgtttcc 120

35

	tttgatgaag aaaaccaagt gtggagcggtt tttctttatg atgatgtcaa aaaagttggt	180
	ggggataaag agttgttttc cagttgcatg ccgcagcaga caagctctat tggaaattcc	240
5	atcattaaca tggacccgcc gaagcataca aaaatccggt cagtcgtgaa caaagccttt	300
	actccgcgcg tgatgaagca atgggaaccg agaattcgag aaatcacaga tgaactgatt	360
	caaaaatttc aggggcgcag tgagtttgac cttgttcacg atttttcata cccgcttccg	420
10	gttattgtga tatctgagtt gctgggagtg cttcagcgc atatggaaca gtttaaagca	480
	tggtctgata ttctggtcag tacaccgaag gataaaagtg aagaagctga aaaagccttt	540
15	ttggaagaac gagataagt tgaggaagaa ctggccgcgt tttttgccg catcatagaa	600
	gaaaagcgaa acaaaccgga acaggatatt atttctatgt tagtggaagc ggaagaaaca	660
	ggcgagaagc tgtcgggtga agagctgatt ccgttttgca cgctgctgct ggtggccgga	720
20	aatgaaacca ctacaaacct gatttcaaat gcgatgtaca gcatattaga aacgccaggc	780
	gtttacgagg aactgcgcag ccatactgaa ctgatgcctc aggcagtgga ggaagccttg	840
25	cgtttcagag cgccggcccc ggttttgagg cgcatgtcca agcgggatac ggagatcggg	900
	gggcacctga ttaaagaagg tgatatggtt ttggcgtttg tggcatcggc aaatcgtgat	960
	gaagcaaagt ttgacagacc gcacatgttt gatatccgcc gccatcccaa tccgcatatt	1020
30	gcgtttggcc acggcatcca tttttgctt ggggccccgc ttgccgtct tgaagcaaat	1080
	atcgcgtaa cgtctttgat ttctgctttt cctcatatgg agtgcgtcag tatcactccg	1140
35	attgaaaaca gtgtgatata cggattaaag agcttccgtg tgaaaatgga attcgtgctg	1200

caccgccatc aaccgggtcac catcgagagaa cccgccgccc gggcggtgtc ccgcaccgtc 1260
 accgtcgagc gcctggaccg gatcgccgac gacgtgctgc gcctcgtcct gcgcgacgcc 1320
 5 ggcggaaaga cattgcccac gtggactccc ggcgcccata tcgacctga cctcgggcgcg 1380
 ctgtcgcgcc agtactccct gtgcggcgcg cccgatgcgc cgagctacga gattgccgtg 1440
 10 cacctggatc ccgagagccg cggcggttcg cgtacatcc acgaacagct cgaggtggga 1500
 agcccgtcc ggatgcgcgg cctcgggaac catttcgcgc tcgaccccgg cgcgagcac 1560
 tacgtgttcg tcgcggcgg catcgccatc accccagtc tgccatggc cgaccagcc 1620
 15 cgcgcccggg ggtggagcta cgaactgcac tactcgcgcc gaaaccgttc cgccatggcc 1680
 tatctcgagc gtgtcgccgg gcacggtgac cgggccgccc tgcacgtgtc cgaggaaggc 1740
 20 acccgatcg acctcgccgc cctctcgcc gagcccgccc ccggcgcca gatctacgcg 1800
 tgcgggcccg ggcggctgct cgccggactc gaggacgcga gccggaactg gcccgacggg 1860
 gcgctgcacg tcgagcactt cacctcgcc ctcggcgcc tcgatccgga cgtcgagcac 1920
 25 gccttcgacc tcgaactgcg tgactcgggg ctgaccgtgc ggtcgaacc caccagacc 1980
 gtctcgacg cgttgcgcgc caacaacatc gacgtgccca gcgactgcga ggaaggcctc 2040
 30 tgcggctcgt gcgaggtcgc cgtctcgac ggcgaggtcg accatcgca cacggtgctg 2100
 accaaggccg agcggggcgc gaaccggcag atgatgacct gctgctcgcg tgctgtggc 2160
 gaccggctgg cctgcgact ctga 2184
 35

<210> 5
 <211> 1191
 <212> DNA
 5 <213> Artificial sequence

 <220>
 <223> A gene coding a protein CYP109B1 with I77F mutation

 10 <400> 5
 atggatgtgt taaaccgccg gcaagccttg cagcgagcgc tgctcaatgg gaaaaacaaa 60

 caggatgcgt atcatccgtt tccatgggtat gaatcgatga gaaaggatgc gcctgtttcc 120

 15 tttgatgaag aaaaccaagt gtggagcgtt tttctttatg atgatgtcaa aaaagttgtt 180

 ggggataaag agttgttttc cagttgcatg ccgcagcaga caagctcttt tggaaattcc 240

 atcattaaca tggacccgcc gaagcataca aaaatccgtt cagtcgtgaa caaagccttt 300
 20 actccgcgcg tgatgaagca atgggaaccg agaattcgag aaatcacaga tgaactgatt 360

 caaaaatttc aggggcgcag tgagtttgac cttgttcacg atttttcata cccgttccg 420

 25 gttattgtga tatctgagtt gctgggagtg ccttcagcgc atatggaaca gtttaaagca 480

 tggctctgatc ttctggtcag tacaccgaag gataaaagtg aagaagctga aaaagccttt 540

 ttggaagaac gagataagtg tgaggaagaa ctggccgcgt tttttgccgg catcatagaa 600
 30 gaaaagcgaa acaaaccgga acaggatatt atttctatct tagtggaagc ggaagaaaca 660

 ggcgagaagc tgtccggtga agagctgatt ccgttttgca cgctgctgct ggtggccgga 720

 35 aatgaaacca ctacaaacct gatttcaaat gcgatgtaca gcatattaga aacgccaggc 780

gtttacgagg aactgcgag ccacccctgaa ctgatgcctc aggagtgga ggaagccttg 840
 egtttcagag cgccggcccc ggttttgagg cgcatggcca agcgggatac ggagatcggg 900
 5 gggcacctga ttaaagaagg tgatatgggt ttggcggttg tggcatcggc aaatcgtgat 960
 gaagcaaagt ttgacagacc gcacatgttt gatatecgcc gccatccaa tccgcatatt 1020
 10 gcgtttggcc acggcatcca ttttgcctt ggggccccgc ttgccgtct tgaagcaa 1080
 atcggttaa cgtcttgat ttctgtttt cctcatatgg agtcgctcag tatcactccg 1140
 attgaaaaca gtgtgatata cggattaaag agcttccgtg tgaaaatgta a 1191
 15
 <210> 6
 <211> 1104
 <212> DNA
 20 <213> Escherichia coli
 <400> 6
 atggaccgca ttattcaatc accgggtaaa tacatccagg gcgtgatgt gattaatcgt 60
 25 ctggggaat acctgaagcc gctggcagaa cgctggtag tgggggtga caaatttgtt 120
 ttaggttttg ctcaatccac tgcgagaaa agctttaaag atctggact ggtagtagaa 180
 attgcgcgt ttggcggtga atgttcgaa aatgagatcg accgtctcg tggcatcgcg 240
 30 gagactgcgc agtgtggcgc aattctcgt atcgttggcg gaaaaacct cgatactgcc 300
 aaagcactgg cacatttcac ggtgttcg gtagcgatcg caccgactat cgcctctacc 360
 35 gatgcacgt gcagcgcat gtctgttatc tacaccgatg aggtgagtt tgaccgctat 420

ctgctgttgc caaataaccc gaatatggtc attgtcgaca ccaaaatcgt cgctggcgca 480
 cctgcacgtc tgtttagcggc gggatatcggc gatgcgctgg caacctgggt tgaagcgct 540
 5 gcctgctctc gtagcggcgc gaccaccatg gcgggcgga agtgcacca ggctgcgtg 600
 gcactggctg aactgtgcta caacaccctg ctggaagaag gcgaaaaagc gatgcttgct 660
 10 gccgaacagc atgtagtac tccggcgctg gagcgctga ttgaagcgaa cacctatttg 720
 agcgggtgtg gttttgaaag tggtagctg gctgcggcgc acgcagtga taacggcctg 780
 accgtatcc cggacgcga tcaactattat cacggtgaaa aagtggcatt cggtacgtg 840
 15 acgcagctgg ttctggaaa tgcgccgtg gaggaaatcg aaaccgtagc tgcccttagc 900
 catgcggtag gtttgcaat aactctgct caactggata ttaaagaaga tgtcccgcg 960
 20 aaaatgcgaa ttgtggcaga agcggcatgt gcagaaggtg aaaccattca caacatgcct 1020
 ggcggcgcga cgccagatca ggtttacgcc gctctgctgg tagccgacca gtacggtcag 1080
 cgtttctgc aagagtggga ataa 1104

25

<210> 7

<211> 29

<212> DNA

30 <213> Artificial sequence

<220>

<223> Primer 1(FW T7)

35 <400> 7

ccgcgaaat taatacgact cactatagg

29

<210> 8

5 <211> 55

<212> DNA

<213> Artificial sequence

<220>

10 <223> Primer 2(RV-109-EcoRI-Rhf-2)

<400> 8

cggttgatgg cgggtgcagca cgaattccat ttccacacgg aagctcttta atccg

55

15

<210> 9

<211> 55

<212> DNA

<213> Artificial sequence

20

<220>

<223> Primer 3(FW Rhf-EcoRI-109-2)

<400> 9

25 cggattaaag agcttcctg tgaaaatgga attcgtgctg caccgccatc aaccg

55

<210> 10

<211> 34

30 <212> DNA

<213> Artificial sequence

<220>

<223> Primer 4(RV-Rhf-Link-Red-HindIII+EcoRV)

35

<400> 10

aagatatcaa gctttcagag tcgcagggcc agcc

34

5 <210> 11

<211> 33

<212> DNA

<213> Artificial sequence

10 <220>

<223> Primer 5(PCR cyp109B1 A11)

<400> 11

cctgcagggtt acattttcac acggaagctc ttt

33

15