

METAL COMPLEX AS SENSOR COMPOUND FOR DETECTING GAS ANALYTE

BACKGROUND

[0001] Exemplary embodiments pertain to the art of sensor compositions based on metal complexes.

[0002] Volatile compounds with a double bond form an important group of compounds for detection. In particular, volatile alkenes, such as ethylene, are analytes of considerable importance. In particular, the detection of ethylene is important to industries related to produce and agriculture. Due to its small size and limited chemical functionality, however, ethylene is a challenging chemical analyte to detect. More efficient and sensitive methods of detection than those currently available are desired.

BRIEF DESCRIPTION


[0003] Disclosed is a sensor compound comprising a metal atom and a mercaptoimidazolyl multidentate ligand.

[0004] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the metal atom is Cu(I), Ag(I) or Au(I).

[0005] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the mercaptoimidazolyl multidentate ligand comprises more than one mercaptoimidazolyl group.


[0006] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the mercaptoimidazolyl multidentate ligand further comprises a pyrazolyl group or an indolyl group.

[0007] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the compound has formula (I)



where each instance of R_1 and R_2 can be hydrogen, or a group having one or more carbons, M is Au(I), Ag(I) or Cu(I), and L is a pyrazolyl group, a mercaptoimidazolyl group, or an indolyl group.

[0008] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the compound has formula II



where each instance of R_1 and R_2 can be hydrogen, or a group having one or more carbons.


[0009] Also disclosed is a sensor composition comprising a sensor compound and a metal dichalcogenide, wherein the sensor compound comprises a metal atom and a mercaptoimidazolyl multidentate ligand.

[0010] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the metal atom is Cu(I), Ag(I) or Au(I).

[0011] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the mercaptoimidazolyl ligand comprises more than one mercaptoimidazolyl group.


[0012] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the mercaptoimidazolyl ligand further comprises a pyrazolyl group or an indolyl group.

[0013] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the sensor compound has formula (I)



where each instance of R_1 and R_2 can be hydrogen, or a group having one or more carbons, M is Au, Ag or Cu, and L is a pyrazolyl group, a mercaptoimidazolyl group, or an indolyl.

[0014] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the sensor compound has formula II



where each instance of R_1 and R_2 can be hydrogen, or a group having one or more carbons.


[0015] In addition to one or more of the features described above, or as an alternative to any of the foregoing embodiments, the metal dichalcogenide comprises MoS_2 , WS_2 , MoSe_2 , WSe_2 , MoTe_2 , WTe_2 , and combinations thereof.

DETAILED DESCRIPTION

[0016] A detailed description of one or more embodiments of the disclosed apparatus and method are presented herein by way of exemplification and not limitation.


[0017] Volatile compounds with a double or triple bond is a group of important compounds for detecting and monitoring. The term volatile, as used herein, refers to compounds that are in the gas phase at standard temperature and pressure. Exemplary compounds include NO₂, CO₂, CO, and alkenes such as C₂H₄ (ethylene). As the hormone responsible for initiating the ripening of fruit as well as other processes in plant development, ethylene is an analyte of considerable importance to industries related to produce and agriculture. Due to its small size and limited chemical functionality, ethylene and other volatile alkenes are challenging chemicals to detect. Disclosed herein is a sensor compound and a sensor composition capable of detecting volatile compounds with a double or triple bond such as ethylene and other volatile alkenes. These compounds can be detected at levels down to 100 parts per billion (ppb).

[0018] The sensor compound comprises a metal atom and a multidentate mercaptoimidazolyl ligand, which combine to form a multidentate metal-ligand coordination complex. The mercaptoimidazolyl ligand comprises at least one mercaptoimidazolyl groups which coordinates to the metal atom through the sulfur atoms. In addition to the mercaptoimidazolyl group(s) the ligand may further comprise a pyrazolyl group, an indolyl group, or a combination thereof, which can be referred to as a heteroleptic ligand. If the mercaptoimidazolyl ligand comprises only mercaptoimidazolyl groups it can be referred to as a homoleptic ligand. The sensor compound may have formula (I)



where each instance of R_1 and R_2 can be hydrogen or a group containing one more carbon atoms. In some embodiments each instance of R_1 and R_2 can be hydrogen or an alkyl group having 1 to 5 carbons. L in formula (I) can be a pyrazolyl group, a mercaptoimidazolyl group, or an indolyl group. VOC in formula I is present to show a postulated interaction with the volatile compound having a π bond. Without being bound by theory it is believed that the π bond of the volatile compound coordinates with an empty coordination site on the metal-ligand complex. The coordination alters the electronic configuration of the complex and can impact the electrical properties of the metal-ligand complex.

[0019] A more specific example of a mercaptoimidazolyl metal-ligand complex is shown in formula (II).



In formula (II) there are three mercaptoimidazolyl groups. R_1 and R_2 are defined as in formula (I).

[0020] The metal atom in the mercaptoimidazolyl metal-ligand complex may be a Group 11 element such as Cu(I), Ag(I), or Au(I).

[0021] The sensor compound (or sensor compounds) may be combined with a metal dichalcogenides to form a sensor composition. Metal dichalcogenides include transition metal dichalcogenides which are compounds formed from a 6B Group metal and a chalcogenide (S, Se, and Te). Exemplary metal dichalcogenides include MoS₂, WS₂, MoSe₂, WSe₂, MoTe₂, WTe₂, and combinations thereof. The metal dichalcogenide is in the form of nanosized particles. “Nanosized” as it applies to the metal dichalcogenides refers to the fact that the particles have at least one dimension that is less than or equal to 100 nanometers. The metal dichalcogenides are typically available in a flake form with a thickness of 100 nanometers or less although other physical forms are not excluded such as few-layer or single-layer materials, with the caveat that the physical form has at least one linear dimension that is less than or equal to 100 nanometers.

[0022] The sensor compound and metal dichalcogenide interact and when the volatile compound is bound to the sensor compound the electrical properties of the sensor composition change – either by increasing the conductivity or the resistivity. The change in electrical property is consistent and predictable, allowing reliable detection of the target volatile compound.

[0023] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the present disclosure. As used herein, the singular forms “a”, “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise. It will be further understood that the terms “comprises” and/or “comprising,” when used in this specification, specify the presence of stated features, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, integers, steps, operations, element components, and/or groups thereof.


[0024] While the present disclosure has been described with reference to an exemplary embodiment or embodiments, it will be understood by those skilled in the art that various changes may be made and equivalents may be substituted for elements thereof without departing from the scope of the present disclosure. In addition, many

modifications may be made to adapt a particular situation or material to the teachings of the present disclosure without departing from the essential scope thereof. Therefore, it is intended that the present disclosure not be limited to the particular embodiment disclosed as the best mode contemplated for carrying out this present disclosure, but that the present disclosure will include all embodiments falling within the scope of the claims.

Claims


1. A sensor compound comprising a metal atom and a mercaptoimidazolyl multidentate ligand, wherein the mercaptoimidazolyl multidentate ligand further comprises a pyrazolyl group or an indolyl group, and wherein the metal atom is Cu(I), Ag(I) or Au(I).

2. The sensor compound of claim 1, wherein the compound has formula (I)



where each instance of R₁ and R₂ can be hydrogen, or a group having one or more carbons, M is Au(I), Ag(I) or Cu(I), and L is a pyrazolyl group, a mercaptoimidazolyl group, or an indolyl group.

3. The sensor compound of claim 1, wherein the compound has formula II




where each instance of R₁ and R₂ can be hydrogen, or a group having one or more carbons.

4. A sensor composition comprising a sensor compound and a metal dichalcogenide wherein the sensor compound comprises a metal atom and a mercaptoimidazolyl multidentate ligand.

5. The sensor composition of claim 4, wherein the metal atom is Cu(I), Ag(I) or Au(I).


6. The sensor composition of claim 4 or 5, wherein the mercaptoimidazolyl ligand further comprises a pyrazolyl group or an indolyl group.

7. The sensor composition of claim 4, wherein the sensor compound has formula (I)



where each instance of R₁ and R₂ can be hydrogen, or a group having one or more carbons, M is Au, Ag or Cu, and L is a pyrazolyl group, a mercaptoimidazolyl group, or an indolyl group.

8. The sensor composition of claim 4, wherein the sensor compound has formula II



where each instance of R₁ and R₂ can be hydrogen, or a group having one or more carbons.

9. The sensor composition of any one of claims 4-8, wherein the metal dichalcogenide comprises MoS₂, WS₂, MoSe₂, WSe₂, MoTe₂, WTe₂, and combinations thereof.