

3,4,5-TRISUBSTITUTED ARYL NITRONE COMPOUNDS AND PHARMACEUTICAL COMPOSITIONS CONTAINING THE SAME

BACKGROUND OF THE INVENTION

Field of the Invention

5 This invention relates to novel 3,4,5-trisubstituted aryl nitrone compounds and their use as therapeutic agents for the treatment of inflammation-related conditions in mammals, such as arthritis, and as analytical reagents for detecting free radicals.

10 State of the Art

 Arthritis and related inflammatory disease conditions occur in more than 100 different forms, including rheumatoid arthritis (RA), osteoarthritis (OA), ankylosing spondylitis and systemic lupus erythematosus (SLE). Most forms of arthritis are characterized by some type of chronic inflammation. For example, RA
15 typically involves chronic inflammation of the lining of the joints and/or the internal organs. Such chronic inflammation generally causes pain and swelling in the joints of those afflicted and may result in damage to cartilage, bone, tendons, ligaments and the like, ultimately leading to deformity and disability.

20 Prostaglandins have long been known to be involved in the inflammation process. Accordingly, a number of inhibitors of prostaglandin synthesis have been developed for the treatment of arthritis and related inflammatory disease conditions. Such non-steroidal antiinflammatory drugs (NSAIDS), such as aspirin, ibuprofen, naproxen and indomethacin, typically prevent the production of
25 prostaglandins by inhibiting enzymes in the arachidonic acid/prostaglandin

-- 2 --


pathway, including the enzyme cyclooxygenase (COX). The enzyme COX catalyzes the conversion of arachidonic acid to prostaglandin H₂, the first step in the biosynthesis of prostaglandins such as prostacyclin and thromboxanes. The enzyme COX is now known to exist in two forms. COX-1 is a constitutive form of the enzyme found in most tissues and organs. Among other properties, COX-1 produces relatively small amounts of prostoglandins necessary for maintaining the integrity of the gastrointestinal tract. COX-2, on the other hand, is an inducible form of the enzyme associated with the increased production of prostoglandins during inflammatory conditions. Since many NSAIDS inhibit both forms of COX, they interfere with prostaglandin-regulated processes not associated with the inflammation process. As a result, many NSAIDS cause severe side effects, such as stomach ulcers and renal damage, which limit their effectiveness as therapeutics.

Accordingly, a need exists for novel classes of therapeutic compounds which effectively treat arthritis and other inflammation-related conditions without producing undesired side effects.

SUMMARY OF THE INVENTION

This invention provides novel 3,4,5-trisubstituted aryl nitron compounds which are useful as therapeutics for reducing inflammation in mammals. In particular, the compounds of this invention are useful for treating arthritis and other inflammation-related conditions.

Accordingly, in one of its composition aspects, this invention is directed to compounds of formula I:




I

This Sheet Has Been Replaced




wherein

R¹ is selected from the group consisting of:



each R² is independently selected from a group of the formula:

10



15

R³ is selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl;

R⁴ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

20

R⁵ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

25

R⁶ and R⁷ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R⁶ and R⁷ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

30

R⁸ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

R⁹ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl,



substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R⁸ and R⁹ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

5 R¹⁰ is selected from the group consisting of hydrogen, lower alkyl and lower cycloalkyl; or R¹ and R¹⁰ can be joined to form an alkylene, substituted alkylene, -C(O)-, -S(O)- or -S(O)₂- group;

R¹¹ and R¹² are independently selected from the group consisting of lower alkyl and lower cycloalkyl; or R¹¹ and R¹² can be joined to form an alkylene group having from 2 to 10 carbon atoms;

10 X is oxygen, sulfur, -S(O)- or -S(O)₂-; and

W is oxygen or sulfur; and pharmaceutically-acceptable salts thereof.

Preferably, R³ is hydrogen or lower alkyl. More preferably, R³ is hydrogen or alkyl having 1 to 4 carbon atoms, more preferably 1 or 2 carbon atoms. Still
15 more preferably, R³ is hydrogen.

R⁴ is preferably selected from the group consisting of alkyl, substituted alkyl and cycloalkyl. More preferably, R⁴ is alkyl having 3 to 6 carbon atoms or cycloalkyl having 5 to 6 carbon atoms. Particularly preferred R⁴ groups include
20 methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

R⁵ is preferably selected from the group consisting of alkyl and cycloalkyl.
25 More preferably, R⁵ is lower alkyl. Particularly preferred R⁵ groups include methyl, ethyl, *n*-propyl, isopropyl and *n*-butyl.


R⁶ is preferably selected from the group consisting of alkyl and
alkoxycarbonylalkyl (i.e., ROC(O)-alkyl-, where R is alkyl or cycloalkyl).
30 Particularly preferred R⁶ groups include ethyl, *n*-propyl, isopropyl, *n*-butyl, ethoxycarbonylmethyl and 2-(ethoxycarbonyl)ethyl. R⁷ is preferably hydrogen.

Preferably, R^8 is alkyl or alkoxyalkyl (i.e., RO-alkyl-, where R is alkyl). Particularly preferred R^8 groups include methyl and methoxyethyl. R^9 is preferably hydrogen. Preferably, X is oxygen.

5 Preferably, R^{10} , R^{11} and R^{12} are independently lower alkyl. More preferably, R^{10} , R^{11} and R^{12} are methyl.

W is preferably oxygen.

10 In a preferred embodiment, this invention is directed to a compound of formula II:



wherein

20 R^{13} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl;

R^{14} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.


25 Preferably, R^{13} is lower alkyl.

R^{14} is preferably selected from the group consisting of alkyl, substituted alkyl and cycloalkyl. More preferably, R^{14} is alkyl having 3 to 6 carbon atoms or
30 cycloalkyl having 5 to 6 carbon atoms. Particularly preferred R^{14} groups include methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-



thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

In another preferred embodiment, this invention is directed to a compound
5 of formula III:



wherein

R^{15} and R^{16} are independently selected from the group consisting of
15 hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl; or R^{15} and R^{16} can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R^{17} is selected from the group consisting of alkyl, substituted alkyl,
20 cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.


R^{15} is preferably selected from the group consisting of alkyl and
alkoxycarbonylalkyl (i.e., $ROC(O)$ -alkyl-, where R is alkyl or cycloalkyl).
Particularly preferred R^{15} groups include ethyl, *n*-propyl, isopropyl, *n*-butyl,
25 ethoxycarbonylmethyl and 2-(ethoxycarbonyl)ethyl. R^{16} is preferably hydrogen.

R^{17} is preferably selected from the group consisting of alkyl, substituted
alkyl and cycloalkyl. More preferably, R^{17} is alkyl having 3 to 6 carbon atoms or
cycloalkyl having 5 to 6 carbon atoms. Particularly preferred R^{17} groups include
30 methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-
thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl
and 3,4,5-trimethoxybenzyl.

This Sheet Has Been Replaced



In still another preferred embodiment, this invention is directed to a compound of formula IV:



wherein

R^{18} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl;

R^{19} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; or R^{18} and R^{19} can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R^{20} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.

Preferably, R^{18} is alkyl or alkoxyalkyl (i.e., RO-alkyl-, where R is alkyl).

Particularly preferred R^{18} groups include methyl and methoxyethyl. R^{19} is preferably hydrogen.


R^{20} is preferably selected from the group consisting of alkyl, substituted alkyl and cycloalkyl. More preferably, R^{20} is alkyl having 3 to 6 carbon atoms or cycloalkyl having 5 to 6 carbon atoms. Particularly preferred R^{20} groups include methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

This Sheet Has Been Replaced



Particularly preferred 3,4,5-trisubstituted aryl nitron compounds include those having the formulae shown in Tables I, II and III.

Table I




Number	R ^a	R ^b
1	CH ₃ -	-C(CH ₃) ₃
2	(CH ₃) ₂ CH-	-C(CH ₃) ₃
3	CH ₃ CH ₂ CH ₂ -	-C(CH ₃) ₃
4	CH ₃ -	-CH(CH ₃) ₂
5	CH ₃ -	-C(CH ₃) ₂ CH ₂ OH
6	CH ₃ CH ₂ CH ₂ CH ₂ -	-C(CH ₃) ₃
7	CH ₃ -	4-CF ₃ -Ph-
8	CH ₃ CH ₂ -	-C(CH ₃) ₃
9	CH ₃ -	-CH ₃
10	CH ₃ -	3,4,5-tri(CH ₃ O-)Ph-

This Sheet Has Been Replaced




Table II



10

Number	R ^c	R ^d
11	CH ₃ CH ₂ -	-C(CH ₃) ₃
12	CH ₃ CH ₂ CH ₂ -	-C(CH ₃) ₃
13	CH ₃ CH ₂ CH ₂ CH ₂ -	-C(CH ₃) ₃
14	CH ₃ CH ₂ OC(O)CH ₂ CH ₂ -	-C(CH ₃) ₃
15	CH ₃ CH ₂ OC(O)CH ₂ -	-C(CH ₃) ₃

Table III



30

Number	R ^e	R ^f
16	CH ₃ -	-C(CH ₃) ₃
17	CH ₃ -O-CH ₂ CH ₂ -	-C(CH ₃) ₃
18	CH ₃ -	-CH ₂ CH ₂ CH(SCH ₃)CH ₃
19	CH ₃ -	-CH ₂ CH ₂ CH ₂ -SCH ₃

This Sheet Has Been Replaced



Accordingly, in another of its composition aspects, this invention is directed to each of the individual compounds:

- α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitronone
- 5 α -(4-isobutanoyloxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitronone
- α -(4-*n*-butanoyloxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitronone
- 10 α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-isopropylnitronone
- α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-1-hydroxy-2-methylprop-2-yl
nitronone
- 15 α -(4-*n*-pentanoyloxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitronone
- α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-4-trifluoromethylbenzylnitronone
- α -(4-propionyloxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitronone
- 20 α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-methylnitronone
- α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-3,4,5-trimethoxybenzylnitronone
- 25 α -[4-(ethylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitronone
- α -[4-(*n*-propylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-
butylnitronone
- 30 α -[4-(*n*-butylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-
butylnitronone
- α -[4-(2-ethoxycarbonyl)ethylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-
N-tert-butylnitronone
- 35 α -[4-(2-ethoxycarbonyl)methylaminocarbonyloxy)-3,5-di-*tert*-
butylphenyl]-*N-tert*-butylnitronone
- α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitronone
- 40 α -[4-(2-methoxy)ethoxymethoxy-3,5-di-*tert*-butylphenyl]-*N-tert*-
butylnitronone
- α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-3-(thiomethoxy)but-1-
yl
nitronone
- 45


α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-3-thiomethoxypropyl nitrone

and pharmaceutically acceptable salts thereof.

5

In another of its composition aspects, this invention is directed to pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula I:

10



15

wherein R¹-R⁴ are as defined above.

20

In additional composition aspects, this invention is directed to pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula II, III or IV above.

25

Among other properties, the 3,4,5-trisubstituted aryl nitrone compounds of this invention are believed to inhibit the induction of cyclooxygenase associated with prostaglandin E₂ (PGE₂) synthesis and inflammation. Compounds having such properties are useful for reducing inflammation, including inflammation resulting from arthritis and related inflammatory conditions.

30

Accordingly, in one of its method aspects, this invention is directed to a method for treating a mammal with an inflammation-related condition which method comprises administering to said mammal a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective inflammation-reducing amount of a compound of formula I, II, III or IV above.

In preferred embodiments of this invention, the inflammation-related condition treated in the above methods is rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus, psoriatic arthritis, and the like.

5

DETAILED DESCRIPTION OF THE INVENTION

For the purposes of this invention, the 3,4,5-trisubstituted aryl nitrone compounds of formula I are named using conventional nitrone nomenclature, i.e., the carbon atom of the carbon-nitrogen double bond (C=N) is designated the α -
10 position and substituents on the nitrogen atom of the carbon-nitrogen double bond are given the *N*- prefix.

In some cases, the 3,4,5,-trisubstituted aryl nitrones of this invention may contain one or more chiral centers. Typically, such compounds will be prepared as
15 a racemic mixture. If desired, however, such compounds can be prepared or isolated as pure stereoisomers, i.e., as individual enantiomers or diastereomers, or as stereoisomer-enriched mixtures. All such stereoisomers (and enriched mixtures) of the 3,4,5-trisubstituted aryl nitrones of formula I are included within the scope of this invention. Pure stereoisomers (or enriched mixtures) may be
20 prepared using, for example, optically active starting materials or stereoselective reagents well known in the art. Alternatively, racemic mixtures of such compounds can be separated using, for example, chiral column chromatography, chiral resolving agents and the like.

25 Additionally, all geometric isomers of the nitrone compounds of formula I are included within the scope of this invention including, for example, all isomers (i.e. E and Z isomers) of the carbon-nitrogen double bond of the nitrone functionality.

This Sheet Has Been Replaced



Definitions

When describing the 3,4,5-trisubstituted aryl nitrones, pharmaceutical compositions and methods of this invention, the following terms have the following meanings unless otherwise specified.

5

"Acyl" refers to the group $-C(O)R$ where R is hydrogen, alkyl, aryl or cycloalkyl.

10

"Acylamino" refers to the group $-NRC(O)R$ where each R is independently hydrogen, alkyl, aryl or cycloalkyl.

"Acyloxy" refers to the group $-OC(O)R$ where R is hydrogen, alkyl, aryl or cycloalkyl.

15

"Alkenyl" refers to a monvalent branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of carbon-carbon double bond unsaturation. Preferred alkenyl groups include ethenyl ($-\text{CH}=\text{CH}_2$), *n*-propenyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), isopropenyl ($-\text{C}(\text{CH}_3)=\text{CH}_2$), and the like.

20

"Substituted alkenyl" refers to an alkenyl group having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxycarbonyl, alkoxycarbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

25
30

"Alkoxy" refers to the group $-\text{OR}$ where R is alkyl. Preferred alkoxy groups include, by way of example, methoxy, ethoxy, *n*-propoxy, isopropoxy, *n*-



butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

"Substituted alkoxy" refers to an alkoxy group having from 1 to 5
5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxy carbonyl, alkoxy carbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy,
10 substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

"Alkoxy carbonyl" refers to the group -C(O)OR where R is alkyl or
15 cycloalkyl.

"Alkoxy carbonylamino" refers to the group -NRC(O)OR' where R is
hydrogen, alkyl, aryl or cycloalkyl, and R' is alkyl or cycloalkyl.

"Alkyl" refers to a monovalent branched or unbranched saturated
20 hydrocarbon group preferably having from 1 to about 10 carbon atoms, more preferably from 1 to 8 carbon atoms and still more preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *tert*-butyl, *n*-hexyl, *n*-octyl, *tert*-octyl and the like. The term
"lower alkyl" refers to an alkyl group having from 1 to 6 carbon atoms.

25 "Substituted alkyl" refers to an alkyl group having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxy carbonyl, alkoxy carbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano,
30 cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol,

This Sheet Has Been Replaced



alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

"Alkylene" refers to a divalent branched or unbranched saturated hydrocarbon group preferably having from 1 to 10 carbon atoms and more preferably from 1 to 6 carbon atoms. This term is exemplified by groups such as methylene (-CH₂-), ethylene (-CH₂CH₂-), the propylene isomers (e.g., -CH₂CH₂CH₂- and -CH(CH₃)CH₂-) and the like.

"Substituted alkylene" refers to an alkylene group having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxycarbonyl, alkoxycarbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

"Alkynyl" refers to a monovalent branched or unbranched unsaturated hydrocarbon group preferably having from 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of carbon-carbon triple bond unsaturation. Preferred alkynyl groups include ethynyl (-C≡CH), propargyl (-CH₂C≡CH) and the like.

"Substituted alkynyl" refers to an alkynyl group having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxycarbonyl, alkoxycarbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

"Amino" refers to the group -NH₂.

This Sheet Has Been Replaced



"Substituted amino" refers to the group $-N(R)_2$ where each R is independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, aryl, cycloalkyl, substituted cycloalkyl, and where both R groups are joined to form an alkylene group. When both R groups are hydrogen, $-N(R)_2$ is an amino group.

"Aminocarbonyl" refers to the group $-C(O)NRR$ where each R is independently hydrogen, alkyl, aryl and cycloalkyl, or where the R groups are joined to form an alkylene group.

"Aminocarbonylamino" refers to the group $-NRC(O)NRR$ where each R is independently hydrogen, alkyl, aryl or cycloalkyl, or where two R groups are joined to form an alkylene group.

"Aminocarbonyloxy" refers to the group $-OC(O)NRR$ where each R is independently hydrogen, alkyl, aryl or cycloalkyl, or where the R groups are joined to form an alkylene group.

"Aryl" refers to an unsaturated aromatic carbocyclic group of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed rings (e.g., naphthyl or anthryl). Preferred aryls include phenyl, naphthyl and the like. Unless otherwise constrained by the definition for the individual substituent, such aryl groups can optionally be substituted with from 1 to 5 substituents, preferably 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkenyl, substituted alkenyl, alkoxy, substituted alkoxy, alkoxycarbonyl, alkyl, substituted alkyl, alkynyl, substituted alkynyl, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

"Aryloxy" refers to the group $-OR$ where R is aryl.

This Sheet Has Been Replaced



"Cycloalkyl" refers to a cyclic alkyl group of from 3 to 10 carbon atoms having a single cyclic ring or multiple condensed or bridged rings which can be optionally substituted with from 1 to 3 alkyl groups. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, 1-methylcyclopropyl, 2-methylcyclopentyl, 2-methylcyclooctyl, and the like, or multiple or bridged ring structures such as adamantanyl and the like. The term "lower cycloalkyl" refers to a cycloalkyl group having from 3 to 6 carbon atoms.

10 "Substituted cycloalkyl" refers to a cycloalkyl group having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxy carbonyl, alkoxy carbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, 15 cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

"Cycloalkoxy" refers to the group -OR where R is cycloalkyl. Such 20 cycloalkoxy groups include, by way of example, cyclopentoxy, cyclohexoxy and the like.

"Cycloalkenyl" refers to a cyclic alkenyl group of from 4 to 10 carbon atoms having a single cyclic ring and at least one point of internal unsaturation 25 which can be optionally substituted with from 1 to 3 alkyl groups. Examples of suitable cycloalkenyl groups include, for instance, cyclopent-3-enyl, cyclohex-2-enyl, cyclooct-3-enyl and the like.

"Substituted cycloalkenyl" refers to a cycloalkenyl group having from 1 to 30 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxy carbonyl, alkoxy carbonylamino, amino, substituted amino, aminocarbonyl,

This Sheet Has Been Replaced



aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

5

"Halo" or "halogen" refers to fluoro, chloro, bromo and iodo. Preferred halo groups are either fluoro or chloro.

"Hydroxyl" refers to the group -OH.

10

"Keto" or "oxo" refers to the group =O.

"Nitro" refers to the group -NO₂.

15

"Thioalkoxy" refers to the group -SR where R is alkyl.

"Substituted thioalkoxy" refers to a thioalkoxy group having from 1 to 5 substituents, and preferably from 1 to 3 substituents, selected from the group consisting of acyl, acylamino, acyloxy, alkoxy, substituted alkoxy, alkoxycarbonyl, alkoxycarbonylamino, amino, substituted amino, aminocarbonyl, aminocarbonylamino, aminocarbonyloxy, aryl, aryloxy, azido, carboxyl, cyano, cycloalkyl, substituted cycloalkyl, halogen, hydroxyl, keto, nitro, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioketo, thiol, alkyl-S(O)-, aryl-S(O)-, alkyl-S(O)₂- and aryl-S(O)₂-.

20

"Thioaryloxy" refers to the group -SR where R is aryl.

"Thioketo" refers to the group =S.

25

"Thiol" refers to the group -SH.

30

"Pharmaceutically-acceptable salt" refers to any salt of a compound of this invention which retains its biological properties and which is not biologically or otherwise undesirable. Such salts may be derived from a variety of organic and inorganic counter-ions well known in the art and include, by way of example

5 illustration, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like. The term "pharmaceutically- acceptable cation" refers to a pharmaceutically acceptable

10 cationic counter-ion of an acidic functional group. Such cations are exemplified by sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium cations, and the like.

General Synthetic Procedures


15 The 3,4,5-trisubstituted aryl nitrones of this invention can be prepared from readily available starting materials using the following general methods and procedures. It will be appreciated that where typical or preferred process conditions (i.e., reaction temperatures, times, mole ratios of reactants, solvents, pressures, etc.) are given, other process conditions can also be used unless

20 otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvent used, but such conditions can be determined by one skilled in the art by routine optimization procedures.


25 Additionally, as will be apparent to those skilled in the art, conventional protecting groups may be necessary to prevent certain functional groups from undergoing undesired reactions. The choice of a suitable protecting group for a particular functional group as well as suitable conditions for protection and deprotection are well known in the art. For example, numerous protecting groups, and their introduction and removal, are described in T. W. Greene and G. M. Wuts,

30 *Protecting Groups in Organic Synthesis*, Second Edition, Wiley, New York, 1991, and references cited therein.

This Sheet Has Been Replaced



In a preferred method of synthesis, the 3,4,5-trisubstituted aryl nitrones of this invention are prepared by coupling an aryl carbonyl compound of formula V:



10 wherein R¹, R² and R³ are as defined above, with a hydroxylamine of formula VI:



wherein R⁴ is as defined above, under conventional reaction conditions.

15

This coupling reaction is typically conducted by contacting the aryl carbonyl compound V with at least one equivalent, preferably about 1.1 to about 2 equivalents, of hydroxylamine VI in an inert polar solvent such as methanol, ethanol, 1,4-dioxane, tetrahydrofuran, dimethyl sulfoxide, dimethylformamide and the like. This reaction is preferably conducted at a temperature of from about 0°C to about 100°C for about 1 to about 48 hours. Optionally, a catalytic amount of an acid, such as hydrochloric acid, acetic acid, *p*-toluenesulfonic acid, silica gel and the like, may be employed in this reaction. When R¹ in formula V is -C(O)R³, at least two equivalents of hydroxylamine VI are employed in this coupling reaction.

25 Upon completion of the reaction, the 3,4,5-trisubstituted aryl nitrone of formula I is recovered by conventional methods including precipitation, chromatographic separation, filtration, distillation, sublimation, and the like.

The aryl carbonyl compounds of formula V employed in the above-described coupling reaction are either known compounds or compounds that can be prepared from known compounds by conventional procedures. For example, aryl carbonyl compounds of formula V where R¹ is -C(O)R⁵ are readily prepared

30



by acylation of the corresponding 4-hydroxy derivative. For example, in a preferred embodiment, 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde (available from Aldrich Chemical Co., 1001 W. St. Paul Avenue, Milwaukee, WI, USA 53233-2641) is acetylated by contacting the benzaldehyde with excess acetic anhydride in the presence of an acid catalyst, such as perchloric acid, followed by hydrolysis of the intermediate acetal, to afford 4-acetoxy-3,5-di-*tert*-butylbenzaldehyde. Other carboxylic anhydrides may also be employed in this reaction including, by way of example, propionic anhydride, butyric anhydride, isobutyric anhydride and the like. Alternatively, such compounds can be prepared by acylation of the 4-hydroxy compound with other acylating agents, such as acyl halides, under conventional reaction conditions. The acyl halides employed in this reaction are preferably acyl chlorides or acyl bromides, such as acetyl chloride, acetyl bromide, propionyl chloride, *n*-butyryl chloride, isobutyryl chloride and the like. Typically, this reaction is conducted in the presence of a trialkylamine, such as triethylamine, to neutralize the acid generated during the reaction.

Similarly, the aryl carbonyl compounds of formula V where R^1 is $-C(O)NR^6R^7$ are readily prepared by reaction of the corresponding 4-hydroxy derivative with an isocyanate (i.e., $R^6R^7N=C=O$). For example, in a preferred embodiment, 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde is reacted with ethylisocyanate to afford 4-(ethylaminocarbonyloxy)-3,5-di-*tert*-butylbenzaldehyde. Typically, this reaction is conducted at ambient temperature in an inert diluent, such as N,N-dimethylformamide, in the presence of an excess of a trialkylamine, such as triethylamine and the like. Other isocyanates may be employed in this reaction including, by way of illustration, *n*-propylisocyanate, *n*-butylisocyanate and the like.

Additionally, the aryl carbonyl compounds of formula V where R^1 is $-CHR^9-X-R^8$ are readily prepared by reacting the corresponding 4-hydroxy derivative with a compound of the formula $L-CHR^9-X-R^8$, where L is a leaving group, such as a halogen or a sulfonate ester, and R^8 , R^9 and X are as defined herein. Typically, this reaction is conducted by contacting the 4-hydroxy



derivative with an excess of the alkylating agent in the presence of an equimolar amount of a trialkylamine, such as N,N-diisopropylethylamine, in an inert diluent such as 1,2-dichloroethane. Preferred alkylating agents for use in this reaction include, by way of example, methoxymethyl chloride and 2-methoxyethoxymethyl (MEM) chloride.

The hydroxylamine compounds of formula VI above are also known compounds or compounds which can be prepared from known compounds by conventional procedures. Typically, the hydroxylamine compounds of formula VI are prepared by reducing the corresponding nitro compound (i.e., R^4-NO_2 , wherein R^4 is as defined above) using a suitable reducing agent such as activated zinc/acetic acid, activated zinc/ammonium chloride or an aluminum/mercury amalgam. This reaction is typically conducted at a temperature ranging from about 15°C to about 100°C for about 0.5 to 12 hours, preferably about 2 to 6 hours, in an aqueous reaction media, such as an alcohol/water mixture in the case of the zinc reagents or an ether/water mixture in the case of the aluminum amalgams. Aliphatic nitro compounds (in the form of their salts) can also be reduced to hydroxylamines using borane in tetrahydrofuran. Since some hydroxylamines have limited stability, such compounds are generally prepared immediately prior to reaction with the aryl carbonyl compound of formula V.

Preferred hydroxylamines for use in this invention include, but are not limited to, *N*-isopropylhydroxylamine, *N*-*n*-propylhydroxylamine, *N*-*n*-butylhydroxylamine, *N*-*tert*-butylhydroxylamine, *N*-cyclohexylhydroxylamine and the like.

Pharmaceutical Compositions

When employed as pharmaceuticals, the 3,4,5-trisubstituted aryl nitrones of this invention are typically administered in the form of a pharmaceutical composition. Such compositions can be prepared using procedures well known in the pharmaceutical art and comprise at least one active compound.

This Sheet Has Been Replaced



Generally, the compounds of this invention are administered in a pharmaceutically effective amount. The amount of the compound actually administered will typically be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response
5 of the individual patient, the severity of the patient's symptoms, and the like.

The pharmaceutical compositions of this invention can be administered by any suitable routes including, by way of illustration, oral, topical, rectal,
10 transdermal, subcutaneous, intravenous, intramuscular, intranasal, and the like. Depending on the intended route of delivery, the compounds of this invention are preferably formulated as either oral, topical or injectable compositions.

Pharmaceutical compositions for oral administration can take the form of
15 bulk liquid solutions or suspensions, or bulk powders. More commonly, however, such compositions are presented in unit dosage forms to facilitate accurate dosing. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired
20 therapeutic effect, in association with a suitable pharmaceutical excipient. Typical unit dosage forms include prefilled, premeasured ampules or syringes of the liquid compositions or pills, tablets, capsules or the like in the case of solid compositions. In such compositions, the nitro compound is usually a minor component (from about 0.1 to about 50% by weight or preferably from about 1 to about 40% by
25 weight) with the remainder being various vehicles or carriers and processing aids helpful for forming the desired dosing form.

Liquid forms suitable for oral administration may include a suitable aqueous or nonaqueous vehicle with buffers, suspending and dispensing agents,
30 colorants, flavors and the like. Solid forms may include, for example, any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or



lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

5

Topical compositions are typically formulated as a topical ointment or cream containing the active ingredient(s), generally in an amount ranging from about 0.01 to about 20% by weight, preferably from about 0.1 to about 10% by weight, and more preferably from about 0.5 to about 15% by weight. When formulated as an ointment, the active ingredients will typically be combined with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with, for example, an oil-in-water cream base. Such topical formulations are well-known in the art and generally include additional ingredients to enhance the dermal penetration or stability of the active ingredients or the formulation. All such known topical formulations and ingredients are included within the scope of this invention.

10

15

20

The compounds of this invention can also be administered by a transdermal device. Accordingly, topical administration can be accomplished using a patch either of the reservoir or porous membrane type or of a solid matrix variety.

25

Injectable compositions are typically based upon injectable sterile saline or phosphate-buffered saline or other injectable carriers known in the art. As before, the 3,4,5-trisubstituted aryl nitron compound in such compositions is typically a minor component, often being from about 0.05 to 10% by weight with the remainder being the injectable carrier and the like.

30

The above-described components for orally and topically administrable or injectable compositions are merely representative. Other materials as well as processing techniques and the like are set forth in Part 8 of Remington's Pharmaceutical Sciences, 18th edition, 1990, Mack Publishing Company, Easton, Pennsylvania, 18042, which is incorporated herein by reference.



The compounds of this invention can also be administered in sustained release forms or from sustained release drug delivery systems. A description of representative sustained release materials can be found in the incorporated materials in Remington's Pharmaceutical Sciences.

5

The following formulation examples illustrate representative pharmaceutical compositions of this invention. The present invention, however, is not limited to the following pharmaceutical compositions.

10 Formulation 1 - Tablets

A compound of formula I is admixed as a dry powder with a dry gelatin binder in an approximate 1:2 weight ratio. A minor amount of magnesium stearate is added as a lubricant. The mixture is formed into 240-270 mg tablets (80-90 mg of active nitron compound per tablet) in a tablet press.

15

Formulation 2 - Capsules

A compound of formula I is admixed as a dry powder with a starch diluent in an approximate 1:1 weight ratio. The mixture is filled into 250 mg capsules (125 mg of active nitron compound per capsule).

20

Formulation 3 - Liquid

A compound of formula I (125 mg), sucrose (1.75 g) and xanthan gum (4 mg) are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of microcrystalline cellulose and sodium carboxymethyl cellulose (11:89, 50 mg) in water. Sodium benzoate (10 mg), flavor, and color are diluted with water and added with stirring. Sufficient water is then added to produce a total volume of 5 mL.

25

Formulation 4 - Injection

The compound of formula I is dissolved in a buffered sterile saline injectable aqueous medium to a concentration of approximately 5 mg/mL.

30

-- 26 --

Formulation 5 - Ointment

Stearyl alcohol (250 g) and white petrolatum (250 g) are melted at about 75°C and then a mixture of a compound of formula I (50 g), methylparaben (0.25 g), propylparaben (0.15 g), sodium lauryl sulfate (10 g), and propylene glycol (120 g) dissolved in water (about 370 g) is added and the resulting mixture is stirred until it congeals.

Compound Utility

Among other properties, the 3,4,5-trisubstituted aryl nitrones of this invention have been discovered to inhibit the induction of inducible cyclooxygenase (COX-2) and/or to inhibit the release of physiologically active leukotrienes and/or to be effective in various *in vivo* arthritis models. Accordingly, the compounds and pharmaceutical compositions of this invention find use as therapeutics for treating inflammation-related conditions in mammals including humans.

In particular, the compounds of this invention have been discovered to effectively inhibit the induction of inducible cyclooxygenase (COX-2), the release of which results in prostaglandin E₂ (PGE₂) synthesis. PGE₂ is produced by the enzyme COX as part of the arachidonic acid pathway. The enzyme COX is now known to exist in two forms, COX-1 and COX-2. COX-1 is a constitutive form of the enzyme found in most tissues and organs. COX-2, on the other hand, is an inducible form of the enzyme associated with the production of PGE₂ and inflammation. Without being limited to theory, it is believed that selective inhibition of COX-2 formation will provide therapeutic agents which effectively reduce inflammation with fewer or none of the side effects associated with inhibition of COX-1 and/or COX-2. Since the compounds of this invention have been discovered to inhibit the release PGE₂, such compounds are useful for treating diseases or conditions characterized by an overproduction or a dysregulated production of prostaglandin E₂ including many inflammatory conditions.

Among the inflammation-related conditions which may be treated with the 3,4,5-trisubstituted aryl nitron compounds and pharmaceutical compositions of this invention are various forms of arthritis, including but not limited to, rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, systemic lupus erythematosus, psoriatic arthritis, and the like. Other inflammation-related conditions include, by way of illustration, inflammatory bowel disease (IBD), septic shock, erythema nodosum leprosy, septicemia, uveitis, adult respiratory distress syndrome (ARDS), organ rejection, neuro-inflammatory conditions, cardio-inflammatory conditions and the like.

As discussed above, the compounds described herein are suitable for use in a variety of drug delivery systems. Injection dose levels for treating inflammation-related conditions range from about 0.1 mg/kg/hour to at least 10 mg/kg/hour, all for from about 1 to about 120 hours and especially 24 to 96 hours. A preloading bolus of from about 0.1 mg/kg to about 10 mg/kg or more may also be administered to achieve adequate steady state levels. The maximum total dose is not expected to exceed about 2 g/day for a 40 to 80 kg human patient.

For the treatment of long-term conditions, such as arthritis, the regimen for treatment may stretch over many months or years so oral dosing is preferred for patient convenience and tolerance. With oral dosing, one to five and especially two to four and typically three oral doses per day are representative regimens. Using these dosing patterns, each dose provides from about 0.1 to about 20 mg/kg of the nitron, with preferred doses each providing from about 0.1 to about 10 mg/kg and especially about 1 to about 5 mg/kg.

The compounds of this invention can be administered as the sole active agent or they can be administered in combination with other active agents, such as cyclooxygenase inhibitors, 5-lipoxygenase inhibitors, non-steroidal antiinflammatory drugs (NSAIDs), steroids, peripheral analgesic agents such as zomepirac, diflunisol, and the like, and other active nitron derivatives.

This Sheet Has Been Replaced



The novel 3,4,5-trisubstituted aryl nitrones of this invention also find use as analytical reagents, i.e. as spin traps, for detecting unstable free radicals using electron spin resonance (ESR) spectroscopy and related techniques. When used as analytical reagents, the nitro compounds of this invention are typically contacted with the radical to be studied in solution and an ESR spectrum generated in a conventional manner. In particular, the nitrones of this invention may be used to detect and identify free radicals in biological systems. Any ESR spectrometer, such as a JEOL JES-FE3XG spectrometer, may be employed in these experiments. Typically, the solution containing the spin-trap will be deoxygenated by, for example, bubbling argon or nitrogen through the solution before the ESR experiment is conducted. Preferably, an excess of the nitro compound is used in such ESR experiments.

The actual experimental procedures employed in the spin-trapping experiment will depend on a number of factors, such as the manner of radical production, the inertness of the solvent and reagents with respect to the spin trap, the lifetime of the spin adduct and the like. Spin trapping procedures are well known in the art and the exact procedure employed can be determined by those skilled in the art. Typical procedures and apparatus for conducting spin trapping experiments are described, for example, in C. A. Evans, "Spin Trapping", *Aldrichimica Acta*, (1979), 12(2), 23-29, and references cited therein.

The following synthetic and biological examples are offered to illustrate this invention and are not to be construed in any way as limiting the scope of this invention.

EXAMPLES

In the examples below, the following abbreviations have the following meanings. Abbreviations not defined below have their generally accepted meaning.

bd	=	broad doublet
bs	=	broad singlet

This Sheet Has Been Replaced



-- 29 --

	d	=	doublet
	dd	=	doublet of doublets
	dec	=	decomposed
	dH ₂ O	=	distilled water
5	ELISA	=	enzyme-linked immuno-sorbent assay
	EtOAc	=	ethyl acetate
	EtOH	=	ethanol
	g	=	grams
	h	=	hours
10	Hz	=	hertz
	ip	=	intraperitoneal
	L	=	liter
	m	=	multiplet
	min	=	minutes
15	M	=	molar
	MeOH	=	methanol
	mg	=	milligram
	MHz	=	megahertz
	mL	=	milliliter
20	mmol	=	millimole
	m.p.	=	melting point
	N	=	normal
	po	=	per os, oral
	q	=	quartet
25	quint.	=	quintet
	s	=	singlet
	t	=	triplet
	THF	=	tetrahydrofuran
	tlc	=	thin layer chromatography
30	μg	=	microgram
	μL	=	microliter
	UV	=	ultraviolet

35 In the examples below, all temperatures are in degrees Celsius (unless otherwise indicated). Example A-N describe the synthesis of intermediates useful for preparing nitrones of this invention; Examples 1-19 describe the synthesis of various nitrones; and Examples I-VI describe the testing of such compounds.

-- 30 --

Example A

Synthesis of
N-tert-Butylhydroxylamine

5 Zinc dust (648 g) was added in portions to a cooled mixture of 2-methyl-2-nitropropane (503 g) and ammonium chloride (207 g) in deionized water (6 L) at such a rate so as to maintain the temperature below 18°C. The reaction mixture was stirred mechanically for 15 hours and then filtered. The solid was washed with hot water (1.75 L). The combined filtrate was saturated with potassium carbonate (4.6 Kg) and extracted with ethyl acetate (2 x 1300 mL). The organic
10 solution was dried over anhydrous sodium sulfate, filtered and rotary evaporated to give the title compound (329 g, 75.7% yield) as white crystals. This material was used without further purification.

Spectroscopic data were as follows:

¹H NMR (CDCl₃, 270 MHz) δ = 1.090 (s, 3 CH₃).

15

Example B

Synthesis of
N-Isopropylhydroxylamine

Using the procedure of Example A above and 2-nitropropane, the title
20 compound was prepared. The crude hydroxylamine product was used without further purification.

Example C

Synthesis of
N-Cyclohexylhydroxylamine

25

Using the procedure of Example A above and nitrocyclohexane, the title compound can be prepared. Alternatively, *N*-cyclohexylhydroxylamine hydrochloride may be purchased commercially from Aldrich Chemical Company, Inc., Milwaukee, WI USA and neutralized with a base, such as potassium
30 carbonate, to provide the title compound.

Example D
Synthesis of
4-Acetoxy-3,5-di-*tert*-butylbenzaldehyde

5 3,5-Di-*tert*-butyl-4-hydroxybenzaldehyde (100 g, 0.411 moles) was placed
in a 5 L round-bottomed flask equipped with a mechanical stirrer. Acetic
anhydride (300 mL) was added with stirring followed by 70% perchloric acid
(0.600 mL). The solid immediately dissolved and a blue colored solution formed.
The reaction mixture was stirred overnight under nitrogen. The reaction progress
10 was determined by TLC. After completion of the reaction, the flask was cooled in
an ice bath. While stirring the reaction mixture vigorously, ice water was added in
small portions. The flask became warm during addition of the water. A total of 2
L of water was added. An oil separated and upon continuous stirring the oil
solidified to provide brown lumps. The brown lumps were separated in a Buchner
15 funnel and the solid was washed freely with water to remove acetic acid. The solid
was then dried under vacuum.

To a solution of the solid in ethanol (250 mL) was added concentrated
hydrochloric acid (25 mL). The resulting solution was boiled for ten minutes and
20 then left to cool. While stirring, the reaction mixture was poured into 2 L of water
and an oil separated. Upon continuous stirring, the oil solidified to form brown
lumps. The solid was separated and packed in Buchner funnel and washed freely
with water (about one liter). The solid was then dried on a mechanical pump to
afford the title compound in 99.4% yield as a light brown solid, m.p. 65.2-74.5°C.

25 Spectroscopic data were as follows:

¹H NMR (CDCl₃, 270 MHz): δ = 9.96 (1H, s, carbonyl H), 7.86 (2H, s,
phenyl H), 2.38 (3H, s, 3CH₃), 1.39 (18H, s, 18 CH₃).

30

-- 32 --

Example E

Synthesis of

4-Isobutanoyl-3,5-di-*tert*-butylbenzaldehyde

5 3,5-Di-*tert*-butyl-4-hydroxybenzaldehyde (10 g, 0.041 mol) was placed in a 500-mL round-bottomed flask fitted with a mechanical stirrer. Isobutyric anhydride (40 mL) was added with stirring, followed by 0.200 mL of 40% perchloric acid. The solid immediately dissolved to form a red color solution. The reaction mixture was stirred overnight under nitrogen. The reaction progress was
10 determined by TLC. After completion of reaction, the reaction mixture was cooled in an ice bath and added to 400 mL of vigorously stirred ice water. The mixture became warm and an oil separated. The oil was extracted with methylene chloride and the organic layer was washed with 1.5M NaOH solution (2 X 100 mL), water (2 X 100 mL) and then dried over magnesium sulfate. The solvent was removed *in vacuo* to afford a red oil.
15

To a solution of the red oil in ethanol (200 mL) was added concentrated hydrochloric acid (50 mL). The resulting solution was boiled for ten minutes and the solution was left to cool. While stirring, the mixture was then poured into 2 L
20 of water and an oil separated. Upon continuous stirring, the oil solidified to brown lumps. The solid was separated and packed in Buchner funnel and then washed freely with water to remove acetic acid (about one liter). The solid was then dried on a mechanical pump to afford the title compound (56.7% yield) which was used without further purification.
25

Example F

Synthesis of

4-*n*-Butanoyl-3,5-di-*tert*-butylbenzaldehyde

30 3,5-Di-*tert*-butyl-4-hydroxybenzaldehyde (15 g, 0.062 mol) was placed in a 500-mL round-bottomed flask fitted with a mechanical stirrer. *n*-Butyric anhydride (50 mL) was added with stirring, followed by 0.200 mL of 40% perchloric acid. The solid immediately dissolved to form a blue colored solution. The reaction

-- 33 --

mixture was stirred overnight under nitrogen. The reaction progress was determined by TLC. After completion of reaction, the reaction mixture was cooled in an ice bath and added to 400 mL of vigorously stirred ice water. The mixture became warm and an oil separated. The oil was extracted with methylene chloride and the organic layer was washed with 5% NaOH (4 X 100 mL), brine (1 X 100 mL), water (4 X 100 mL) and dried over MgSO₄. The solvent was removed *in vacuo* to afford a red oil.

To a solution of the red oil in ethanol (200 mL) was added concentrated hydrochloric acid (40 mL). The resulting solution was boiled for ten minutes and left to cool. While stirring, the mixture was then poured into 2 L of water and an oil separated. Upon continuous stirring, the oil did not solidify, so the mixture was extracted with methylene chloride. The organic layer was washed with saturated NaHCO₃ (2 X 100 mL), water (3 X 100 mL) and dried over MgSO₄. The solvent was removed *in vacuo* to afford the title compound as a red oil (77.6% yield) which was used without further purification.

Example G

Synthesis of 4-*n*-Pentanoyl-3,5-di-*tert*-butylbenzaldehyde

3,5-Di-*tert*-butyl-4-hydroxybenzaldehyde (10.0 g, 0.081 mol) was placed in a 500-mL round-bottomed flask fitted with a mechanical stirrer. Valeric anhydride (50 mL) was added with stirring, followed by 0.200 mL of 40% perchloric acid. The solid immediately dissolved to form a green colored solution. The reaction mixture was stirred overnight under nitrogen. The reaction progress was determined by TLC. After completion of reaction, the reaction mixture was cooled in an ice bath and added to 400 mL of vigorously stirred ice water. The mixture became warm and an oil separated. The oil was extracted with methylene chloride and the organic layer was washed with 5% NaOH (3 X 50 mL), brine (3 X 50 mL), water (3 X 100 mL) and dried over magnesium sulfate. The solvent was removed *in vacuo* to afford a thick oil.

-- 34 --

To a solution of the oil in ethanol (200 mL) was added concentrated hydrochloric acid (50 mL). The resulting solution was boiled for ten minutes and left to cool. While stirring, the mixture was then poured into 2 L of water and an oil separated. Upon continuous stirring, the oil did not solidify, so the mixture was extracted with methylene chloride. The organic layer was washed with saturated NaHCO₃ (3 X 50 mL), water (5 X 100 mL) and dried over MgSO₄. The solvent was removed *in vacuo* to afford the title compound as a red oil which was used without further purification.

Example H

Synthesis of

4-(Ethylaminocarbonyloxy)-3,5-di-*tert*-butylbenzaldehyde

To a solution of 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde (5 g; 20 mmol) in DMF (100 mL) was added triethylamine (3.5 mL, 25 mmol) and the solution was stirred for 15 min at room temperature. Ethyl isocyanate (1.95 mL, 25 mmol) was added and the reaction mixture was stirred for 2 h at room temperature until no more hydroxybenzaldehyde was detected by TLC (R_f = 0.78 for product and 0.89 for starting material using 1:1 hexanes/EtOAc). The reaction solution was concentrated by azeotropic removal of DMF with water. The resulting suspension was filtered, washed with water and dried in a vacuum oven to afford 96% of the title compound as a light buff solid.

Spectroscopic data were as follows:

¹H NMR (DMSO-*d*₆, 270 MHz): δ = 9.96 (1H, s, aldehyde CHO), 8.04 (1H, t, carbamate NH), 7.83 (2H, s, phenyl H), 3.09 (2H, m, ethyl CH₂), 1.34 (18H, s, 6 CH₃), 1.07 (3H, t, ethyl CH₃).

¹³C NMR (DMSO-*d*₆, 270 MHz): δ = 193.4, 155.1, 154.2, 145.0, 133.3, 127.8, 35.9, 31.4, 30.5.

-- 35 --

Example I

Synthesis of

4-(*n*-Propylaminocarbonyloxy)-3,5-di-*tert*-butylbenzaldehyde

5 The title compound was prepared in DMF using 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde and *n*-propyl isocyanate according to the procedure described in Example H. The title compound was isolated in 100.0% yield as an off-white solid.

Example J

Synthesis of

4-(*n*-Butylaminocarbonyloxy)-3,5-di-*tert*-butylbenzaldehyde

10 The title compound was prepared in DMF using 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde and *n*-butyl isocyanate according to the procedure described in Example H. The title compound was isolated in 87.5% yield as a buff solid.

Example K

Synthesis of

20 4-(2-Ethoxycarbonyl)ethylaminocarbonyloxy)-3,5-di-*tert*-butylbenzaldehyde

 The title compound was prepared in DMF using 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde and ethyl 3-isocyanatopropionate according to the procedure described in Example H. The title compound was isolated in 100% yield as an off-white solid.

Example L

Synthesis of

30 4-(2-Ethoxycarbonyl)methylaminocarbonyloxy)-3,5-di-*tert*-butylbenzaldehyde

 The title compound was prepared in DMF using 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde and ethyl 2-isocyanatoacetate according to the procedure described in Example H. The title compound was isolated in 100% yield as an off-white solid.

-- 36 --

Example M

Synthesis of
4-Methoxymethoxy-3,5-di-*tert*-butylbenzaldehyde

5 To a solution of 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde (10 g, 40 mmol) in 1,2-dichloroethane (200 mL) was added N,N-diisopropylethylamine (6.97 mL, 40 mmol) and the solution was stirred for 1 h at room temperature. Chloromethoxymethyl ether (3.77 mL, 50 mmol) was added and the reaction mixture was stirred for 1 h at room temperature and then refluxed for 16 h until no more hydroxybenzaldehyde was detected by TLC ($R_f = 0.78$ for product and 0.70 for starting material using 1:1 hexanes/EtOAc). The reaction mixture was cooled to room temperature and washed with water. The organic layer was concentrated and the residue applied to a silica gel column and eluted with mixture of pentane/EtOAc. The title compound was isolated in 100 % yield as a brown oil.

15 Spectroscopic data were as follows:

^1H NMR (DMSO- d_6 , 270 MHz): $\delta = 9.89$ (1H, s, aldehyde CHO), 7.80 (2H, s, phenyl H), 4.92 (2H, s, CH_2), 3.64 (3H, s, CH_3), 1.46 (18H, s, 6 CH_3).

^{13}C NMR (DMSO- d_6 , 270 MHz): $\delta = 192.11$, 160.11, 145.77, 131.65, 128.55, 100.98, 57.66, 35.99 and 30.16.

Example N

Synthesis of
4-(2-Methoxy)ethoxymethoxy-3,5-di-*tert*-butylbenzaldehyde

25 To a solution of 3,5-di-*tert*-butyl-4-hydroxybenzaldehyde (8 g, 33 mmol) in 1,2-dichloroethane (150 mL) was added N,N-diisopropylethylamine (6.97 mL, 40 mmol) and the solution was stirred for 1 h at room temperature. 2-Methoxyethoxymethyl (MEM) chloride (5.7 mL, 50 mmol) was added and the reaction mixture was stirred for 1 h at room temperature and then at reflux for 26 h until no more hydroxybenzaldehyde was detected by TLC ($R_f = 0.92$ for product and 0.70 for starting material using 1:1 hexanes/EtOAc). The reaction mixture was cooled to room temperature and washed with water. The organic layer was

-- 37 --

concentrated and the residue applied to a silica gel column and eluted with mixture of pentane/EtOAc. The title compound was isolated (98 % yield) as a brown oil.

Example 1

Synthesis of

α -(4-Acetoxy-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

4-Acetoxy-3,5-di-*tert*-butylbenzaldehyde (113.16 g, 0.41 mol) was placed in a 2 L round-bottomed flask fitted with a magnetic stirrer. Benzene (500 mL) was added and the mixture was stirred until the solids dissolved. To the resulting red solution was added *tert*-butylhydroxylamine (43.80 g, 0.49 mol) and silica gel (20 g). The mixture was refluxed overnight at which time TLC showed no remaining starting material (R_f = 0.31 for product and 0.80 for starting material using 3:1 hexanes/EtOAc). The benzene was removed *in vacuo* on a rotovap to provide a grey solid. The solid was dissolved in a minimum amount of ethyl acetate and the flask was left to stand in the freezer. The white crystals which formed were separated, washed with hexanes and dried under vacuum to afford 105.78 g of the title compound as a crystalline white solid (74.4% yield), m.p. 227.0-248.9°C.

Spectroscopic data were as follows:

^1H NMR (CDCl_3 , 270 MHz): δ = 8.32 (1H, s, phenyl H), 7.50 (1H, s, nitronyl H), 2.35 (3H, s, 1 CH_3), 1.61 (9H, s, 3 CH_3), 1.37, s, 18 CH_3).

^{13}C NMR (CDCl_3 , 270 MHz): δ = 170.7, 149.2, 142.6, 129.7, 128.4, 127.2, 70.7, 35.5, 31.4, 28.5, 22.6.

Example 2

Synthesis of

α -(4-Isobutanoyl-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

To a solution of 4-isobutanoyl-3,5-di-*tert*-butylbenzaldehyde (7.00 g, 0.023 mol) in benzene (200 mL) was added *tert*-butylhydroxylamine (2.56 g, 0.029 mol) and *p*-toluenesulfonic acid (0.100 g). Using a Dean-Stark trap, the resulting

-- 38 --

mixture was refluxed until no more aldehyde was detected by TLC ($R_f = 0.30$ for product and 0.89 for starting material using 3:1 hexanes/EtOAc). The solvent was removed *in vacuo* and the residue was thoroughly washed with hexanes. The title compound was isolated in 50.8% yield as a white solid, m.p. 197.1-208.1 °C.

5 Spectroscopic data were as follows:

^1H NMR (CDCl_3 , 270 MHz): $\delta = 8.32$ (2H, s, phenyl), 7.51 (1H, s, nitronyl H), 2.87 (1H, q, $J=7.17\text{Hz}$, isobutyryl H), 1.61 (9H, s, 9CH_3), 1.38 (6H, d, $J=7.17$, 6CH_3), 1.36 (18H, s, 18CH_3).

10 ^{13}C NMR (CDCl_3 , 270 MHz): $\delta = 176.0, 150.2, 142.7, 129.9, 128.1, 127.1, 70.7, 35.5, 35.2, 31.3, 28.5, 18.7$.

Example 3

Synthesis of

α -(4-*n*-Butanoyl-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

15

To a solution of 4-*n*-butanoyl-3,5-di-*tert*-butylbenzaldehyde (9.63 g, 0.032 mol) in benzene (200 mL) was added *tert*-butylhydroxylamine (3.52 g, 0.040 mol) and *p*-toluenesulfonic acid (0.100 g). Using a Dean-Stark trap, the resulting solution was refluxed until no more aldehyde was detected by TLC ($R_f = 0.30$ for product and 0.91 for starting material using 3:1 hexanes/EtOAc). The solvent was removed *in vacuo* and the residue was crystallized from EtOAc and hexanes to afford the titled compound (50.2% yield) as a white solid, m.p. 216.9-236.5 °C.

20

Spectroscopic data were as follows:

25 ^1H NMR (CDCl_3 , 270 MHz): $\delta = 8.32$ (2H, s, phenyl H), 7.50 (1H, s, nitronyl H), 2.61 (2H, t, $J=7.67\text{Hz}$, 2CH_2), 1.78 (2H, m, $J=4.70\text{Hz}$, 2CH_2), 1.61 (9H, s, 9CH_3), 1.36 (18H, s, 18CH_3), 1.05 (3H, t, $J=7.42\text{Hz}$, 3CH_3).

^{13}C NMR (CDCl_3 , 270 MHz): $\delta = 173.2, 149.5, 142.7, 129.8, 128.2, 127.1, 70.7, 37.6, 35.5, 31.4, 28.5, 17.7, 13.8$.

30

Example 4

Synthesis of
 α -(4-Acetoxy-3,5-di-*tert*-butylphenyl)-*N*-isobutylnitrone

5 To a solution of 4-acetoxy-3,5-di-*tert*-butylbenzaldehyde (17.0 g, 0.0615 mol) in methanol (250 mL) was added isobutylhydroxylamine (5.36 g, 0.0714 mol) and concentrated hydrochloric acid (10 drops). Using a Dean-Stark trap, the resulting solution was refluxed until no more aldehyde was detected by TLC (R_f = 0.23 for product and 0.84 for starting material using 3:1 hexanes/EtOAc). The solvent was removed in vacuo and the residue was a red oil which when triturated with hexanes it turned into solid. The solid was separated and washed completely with hexanes. The titled compound was isolated in 52.3% yield as a white solid, m.p. 176.8-180.4°C.

Spectroscopic data were as follows:

15 ^1H NMR (CDCl_3 , 270 MHz): δ = 8.3 (2H, s, phenyl H), 7.42 (1H, s, nitronyl H), 4.20 (1H, q, J = 6.50 Hz, 1CH), 2.35 (3H, s, 3CH₃), 1.50 (6H, d, J = 6.43 Hz, 6CH₃), 1.37 (18H, s, 18CH₃).

^{13}C NMR (CDCl_3 , 270 MHz): δ = 170.7, 150.1, 142.7, 131.9, 128.0, 127.0, 67.8, 35.5, 31.3, 22.6, 20.9.

20

Example 5

Synthesis of
 α -(4-Acetoxy-3,5-di-*tert*-butylphenyl)-*N*-1-hydroxy-2-methylprop-2-yl nitrone

25 To a solution of 4-acetoxy-3,5-di-*tert*-butylbenzaldehyde (10.0 g, 0.0362 mol) in benzene (200 mL) was added *N*-(1-hydroxy-2-methylprop-2-yl) hydroxylamine (5.55 g, 0.0543 mol) and *p*-toluenesulfonic acid (0.090 g). Using a Dean-Stark trap, the resulting mixture was refluxed until no more aldehyde was detected by TLC (R_f = 0.20 for product and 0.98 for starting material using 1:1 hexanes/EtOAc). The solvent was removed *in vacuo* and the residue was washed with hot hexanes. The title compound was isolated (83.9 % yield) as a off white solid, m.p. 204.3-204.8°C.

30

Spectroscopic data were as follows:

This Sheet Has Been Replaced



-- 40 --

¹H NMR (CDCl₃, 270 MHz): δ=8.30 (2H, s, phenyl H), 7.46 (1H, s, nitronyl H). 3.79 (2H, s, 2 CH₂), 2.36 (3H, s, 3 CH₃), 1.60, s, 6 CH₃), 1.37 (18H, s, 18CH₃).

¹³C NMR (CDCl₃, 270 MHz): δ =171.1, 150.1, 142.9, 132.3, 127.7, 72.8, 69.8, 35.5, 31.6, 31.3, 23.9, 22.6.

Example 6

Synthesis of α-(4-*n*-Pentanoyl-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

10

To a solution of 4-pentanoyl-3,5-di-*tert*-butylbenzaldehyde (17.48 g, 0.050 mol) in benzene (250 mL) was added *tert*-butylhydroxylamine (4.57 g, 0.0510 mol) and *p*-toluenesulfonic acid (0.080 g). Using a Dean-Stark trap, the resulting mixture was refluxed until no more aldehyde was detected by TLC (R_f = 0.43 for product using 1:1 hexanes/EtOAc). The solvent was removed *in vacuo* and the residue was thoroughly washed with hexanes. The title compound was isolated (56.50 % yield) as a white solid, m.p. 195.5-204.8°C.

15

Spectroscopic data were as follows:

¹H NMR (CDCl₃, 270 MHz): δ= 8.31 (2H, s, phenyl H), 7.50 (1H, s, nitronyl H). 2.63 (2H, t, J=7.80 2CH), 1.85-1.67 (2H, m, 2CH₂), 1.61 (9H, s, 9CH₃), 1.52-1.36 (2H, m, 2CH₂), 1.36 (18H, s, 18CH₃), 0.97 (3H, t, J=7.30, 3CH₃).

20

¹³C NMR (CDCl₃, 270 MHz): δ =173.2, 149.4, 142.7, 129.8, 128.2, 127.1, 70.7, 35.5, 35.4, 31.4, 28.4, 26.2, 22.2, 13.8.

25

Examples 7-10

Using the appropriate starting materials and the procedures described herein, the following additional compounds were prepared:

α-(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-4-trifluoromethylbenzylnitrone (m.p. 217-232°C);

30

α-(4-propionyloxy-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone (m.p. 202-222°C);

α-(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-methylnitrone (m.p. 136-143°C);

and

α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-3,4,5-trimethoxybenzylinitrone
(m.p. 231.7-235°C).

Example 11

Synthesis of

α -[4-(Ethylaminocarbonyloxy-3,5-di-*tert*-butylphenyl)]-*N*-*tert*-butylnitrone

To a solution of 3, 5-di-*tert*-butyl-4-(ethylaminocarbonyloxy)-benzaldehyde (5 g, 16 mmol) in benzene (200 mL) was added *tert*-butylhydroxylamine (2.14 g, 24 mmol). The resulting solution was refluxed for 96 h until no more aldehyde was detected by TLC (R_f = 0.57 for product and 0.78 for starting material using 1:1 hexanes/EtOAc). The solvent was removed *in vacuo* and the residue applied to a silica gel column and eluted with mixture of hexane/EtOAc. The title compound was isolated in 57 % yield as a white solid, m.p. 202.2-212.2°C.

Spectroscopic data were as follows:

¹H NMR (DMSO-*d*₆, 270 MHz): δ = 8.37 (2H, s, phenyl H), 7.86 (1H, t, J = 6.3 Hz, carbamate NH), 7.83 (1H, s, nitronyl H), 3.08 (2H, quintet, J = 6.3 Hz, ethyl CH₂), 1.49 (9H, s, 3 CH₃), 1.31 (18H, s, 6 CH₃), 1.06 (3H, t, J = 6.4 Hz, ethyl CH₃).

¹³C NMR (DMSO-*d*₆, 270 MHz): δ = 155.8, 149.6, 143.30, 129.5, 128.7, 126.9, 70.6, 35.8, 31.6, 28.5 and 15.7.

Example 12

Synthesis of

α -[4-(*n*-Propylaminocarbonyloxy-3,5-di-*tert*-butylphenyl)]-*N*-*tert*-butylnitrone

The title compound was prepared in benzene using 3,5-di-*tert*-butyl-4-(*n*-propylaminocarbonyloxy)benzaldehyde and *tert*-butylhydroxylamine according to the procedure described in Example 11. The title compound was isolated in 96.6% yield as a white solid, m.p. 206.5-214.0°C.

Spectroscopic data were as follows:

-- 42 --

¹H NMR (DMSO-*d*₆, 270 MHz): δ = 8.38 (2H, s, phenyl H), 7.90 (1H, t, J = 5.8 Hz, carbamate NH), 7.84 (1H, s, nitronyl H), 3.02 (2H, q, J = 6.5 Hz, propyl NCH₂), 1.50 (9H, s, 3 CH₃), 1.45 (3H, hextet, J = 7.2 Hz, propyl CH₂), 1.32 (18H, s, 6 CH₃), 0.88 (3H, t, J = 7.3 Hz, propyl CH₃).

5 ¹³C NMR (DMSO-*d*₆, 270 MHz): δ = 155.7, 149.7, 143.3, 129.5, 128.7, 127.0, 70.6, 42.8, 35.8, 31.7, 28.5, 23.1 and 11.8.

Example 13

Synthesis of

10 α-[4-(*n*-Butylaminocarbonyloxy-3,5-di-*tert*-butylphenyl)]-*N*-*tert*-butylnitrone

The title compound was prepared in benzene using 3,5-di-*tert*-butyl-4-(*n*-butylaminocarbonyloxy)benzaldehyde and *tert*-butylhydroxylamine according to the procedure described in Example 11. The title compound was isolated in 93.4%
15 yield as a white solid, m.p. 203.6-205.1 °C.

Spectroscopic data were as follows:

¹H NMR (DMSO-*d*₆, 270 MHz): δ = 8.38 (2H, s, phenyl H), 7.88 (1H, t, J = 5.7 Hz, carbamate NH), 7.84 (1H, s, nitronyl H), 3.05 (2H, q, J = 6.3 Hz, *n*-butyl NCH₂), 1.50 (9H, s, 3 CH₃), 1.45-1.37 (4H, m, *n*-butyl 2 CH₂), 1.31 (18H, s, 6
20 CH₃), 0.87 (3H, t, J = 7.1 Hz, *n*-butyl CH₃).

¹³C NMR (DMSO-*d*₆, 270 MHz): δ = 155.6, 149.7, 143.3, 129.5, 128.7, 127.0, 70.6, 40.7, 35.8, 31.9, 31.6, 28.5, 19.9 and 14.2.

Example 14

Synthesis of

25 α-[4-(2-Ethoxycarbonyl)ethylaminocarbonyloxy-3,5-di-*tert*-butylphenyl]-*N*-*tert*-butylnitrone

The title compound was prepared in benzene using 3,5-di-*tert*-butyl-4-(2-ethoxycarbonyl)ethylaminocarbonyloxy)benzaldehyde and *tert*-
30 butylhydroxylamine according to the procedure described in Example 11. The title compound was isolated in 88.4% yield as a white solid, m.p. 130.5-146.3 °C.

Spectroscopic data were as follows:

¹H NMR (DMSO-*d*₆, 270 MHz): δ = 8.38 (2H, s, phenyl H), 8.01 (1H, t, J = 6.0 Hz, carbamate NH), 7.84 (1H, s, nitronyl H), 4.08 (2H, q, J = 7.0 Hz, ethyl OCH₂), 3.28 (2H, t, J = 6.0 Hz, propionate NCH₂), 2.50 (2H, t, J = 6.0 Hz, propionate COCH₂), 1.49 (9H, s, 3 CH₃), 1.26 (18H, s, 6 CH₃), 1.19 (3H, t, J = 7.0 Hz, ethyl CH₃).

¹³C NMR (DMSO-*d*₆, 270 MHz): δ = 171.5, 155.6, 149.5, 143.3, 129.4, 128.8, 127.0, 70.6, 60.5, 37.2, 35.8, 34.6, 31.7, 28.5, and 14.7.

Example 15

Synthesis of α-[4-(2-Ethoxycarbonyl)methylaminocarbonyloxy-3,5-di-*tert*-butylphenyl]-*N*-*tert*-butylnitrone

The title compound was prepared in benzene using 3,5-di-*tert*-butyl-4-(2-ethoxycarbonyl)methylaminocarbonyloxy)benzaldehyde and *tert*-butylhydroxylamine according to the procedure described in Example 11. The title compound was isolated in 42% yield as a white solid, m.p. 177.2-181.6°C.

Spectroscopic data were as follows:

¹H NMR (DMSO-*d*₆, 270 MHz): δ = 8.38 (2H, s, phenyl H), 8.35 (1H, t, J = 5.7 Hz, carbamate NH), 7.84 (1H, s, nitronyl H), 4.10 (2H, q, J = 7.0 Hz, ethyl OCH₂), 3.83 (2H, d, J = 5.7 Hz, NCH₂), 1.50 (9H, s, 3 CH₃), 1.32 (18H, s, 6 CH₃), 1.19 (3H, t, J = 7.0 Hz, ethyl CH₃).

¹³C NMR (DMSO-*d*₆, 270 MHz): δ = 170.3, 155.9, 143.3, 129.4, 128.9, 127.0, 70.7, 61.0, 42.9, 35.8, 31.7, 28.5, and 14.6.

Example 16

Synthesis of α-(4-Methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

To a solution of 3,5-di-*tert*-butyl-4-methoxymethoxybenzaldehyde (11.42 g, 40 mmol) in benzene (200 mL) was added *tert*-butylhydroxylamine (4.0 g, 50 mmol). The resulting solution was refluxed for 72 h until no more aldehyde was detected by TLC (*R*_f = 0.56 for product and 0.78 for starting material in 1:1 hexanes/EtOAc). The solvent was removed *in vacuo* and the residue was

suspended in hexane/EtOAc. The suspension was filtered, washed with hexanes and dried to afford the title compound (69 % yield) as a white solid, m.p. 202.2-205.9°C.

Spectroscopic data were as follows:

5 ^1H NMR (DMSO- d_6 , 270 MHz): δ = 8.37 (2H, s, phenyl H), 7.79 (1H, s, nitronyl H), 4.84 (2H, s, OCH_2), 3.54 (3H, s, OCH_3), 1.48 (9H, s, 3 CH_3), 1.40 (18H, s, 6 CH_3).

^{13}C NMR (DMSO- d_6 , 270 MHz): δ = 155.3, 143.9, 129.4, 127.8, 127.4, 101.0, 70.5, 57.6, 36.0, 32.3, 28.5.

10

Example 17

Synthesis of

α -[4-(2-Methoxy)ethoxymethoxy-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

15

The title compound was prepared in benzene with 3,5-di-*tert*-butyl-4-(2-methoxy)ethoxymethoxybenzaldehyde and *tert*-butylhydroxylamine using the procedure described in Example 16. The title compound was isolated (70.1 % yield) as a white solid, m.p. 169.6-173.5°C.

Spectroscopic data were as follows:

20 ^1H NMR (DMSO- d_6 , 270 MHz): δ = 8.38 (2H, s, phenyl H), 7.80 (1H, s, nitronyl H), 4.91 (2H, s, OCH_2), 3.87 (2H, m, J = 4.76 Hz, ethoxy OCH_2), 3.54 (2H, m, J = 4.76 Hz, OCH_2), 3.28 (3H, s, OCH_3), 1.49 (9H, s, 3 CH_3), 1.40 (18H, s, 6 CH_3).

25 ^{13}C NMR (DMSO- d_6 , 270 MHz): δ = 155.3, 143.9, 129.4, 127.8, 127.4, 100.1, 71.7, 70.5, 69.3, 58.7, 36.0, 32.3, 28.5.

Examples 18 and 19

Using the appropriate starting materials and the procedures described herein, the following additional compounds were prepared:

30 α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-3-(thiomethoxy)but-1-yl nitrone (m.p. 76.7-80.0°C); and

35 α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-3-thiomethoxypropyl nitrone (m.p. 55-63°C).

Example I

Lack of Inhibition of COX-1 and COX-2

In this experiment, the compound of Example 1 was tested to determine if it inhibited the enzymes cyclooxygenase 1 (COX-1) and cyclooxygenase 2 (COX-2).

The inhibition of COX-1 was measured by incubating arachidonic acid with COX-1 derived from ram seminal vesicles. The inhibition of COX-2 was measured by incubation of arachidonic acid with COX-2 derived from sheep placenta. These assays are further described in Evans et al., *Biochem. Pharmacol.*, 36:2035, 1987. In each assay, the formation of prostaglandin E₂ (PGE₂) was measured by spectrophotometric quantitation of malondialdehyde. The compound of Example 1 was added to incubation solutions at a concentration of 300 μ M. At this concentration, indomethacin, a known inhibitor of COX-1 and COX-2, completely inhibits both enzymes. Under the same conditions, the compound of Example 1 did not significantly affect COX-1 or COX-2 at a concentration of 300 μ M. These results demonstrate that the compound of Example 1 is not an inhibitor of the enzymes cyclooxygenase 1 (COX-1) and cyclooxygenase 2 (COX-2).

Example II

Inhibition of PGE₂

In this experiment, the ability of nitrones of formula I to inhibit induction of prostaglandin E₂ (PGE₂) is demonstrated.

For this assay, the test compounds were solubilized in 500 μ L of EtOH and then adjusted to 10 mL with complete DMEM with 1 μ g/mL lipopolysaccharide (LPS) to obtain a 10mM solution. This solution was then diluted 1/100 to obtain a 100 μ M. As a reference, indomethacin was solubilized in 500 μ L DMSO and then adjusted to 50 mL with complete DMEM with 1 μ g/mL LPS. This 2 mM solution was then diluted 1/200 to obtain a 10 μ M solution. This solution was used as a reference. All dilutions were made with complete DMEM

Macrophages (murine macrophages RAW 264.7) were obtained from American Type Culture Collection, Rockville, MD. They were cultured in 75 cm³ flasks with DMEM (Dulbecco Modified Eagle Medium with 10% Fetal Bovine Serum, penicillin, streptomycin, and glutamine) and seeded into a 96 well plate at 20 x 10⁶ cells/10 mL, 200μL per well. At 80-100% confluence, the macrophages were washed with HBSS (with penicillin, streptomycin, and glutamine) and incubated with the test compound or a reference diluted into DMEM (with 10% FCS, penicillin, streptomycin, and glutamine) with 1μg/mL LPS for 18 hours. The cells were then washed with HBSS and incubated with 30μM arachidonic acid in HBSS for 15 minutes at 37°C. The supernatant was taken to determine PGE₂ levels using conventional procedures. Results were expressed as percent inhibition. Compounds reducing the induction of PGE₂ by at least about 30% compared to the control were considered effective in this assay.

Initially, test compounds were screened at 100μM. Compounds showing activity were screened at lower concentrations to obtain an IC₅₀. For such screening, the stock solution was diluted in a series 1:10 dilutions to obtain 30μM, 10μM, 1μM, 0.1μM solutions. The 10μM solution was diluted 1:3 to obtain a 3μM solution which is then diluted in a series to produce solutions of 0.3μM and 0.03μM. The serial dilutions are used to determine the IC₅₀ of the test compound.

Example III

Carrageenan Footpad Edema Assay

In this example, the ability of compounds of formula I to reduce carrageenan footpad edema in rats is demonstrated. This assay is commonly used to screen and evaluate anti-inflammatory drug candidates. See, for example, C. A. Winter et al., "Carrageenin-induced Edema in Hind Paw of the Rat as an Assay for Anti-inflammatory Drugs," *Proc. Sci. Exp. Biol. Med.* 111, 544-547 (1962) and references cited therein.

In this assay, a carrageenan suspension (0.5%) was prepared by mixing 50 mg of carrageenan (Type IV, λ) in 10 mL of sterile saline solution. Male

Sprague-Dawley rats (150-250 g) were then injected subcutaneously with 100 μ L of the carrageenan suspension in the plantar portion of the right rear paw. The test compound (100 mg in 2 mL) or a vehicle control (2 mL) was then administered by po or ip. The initial foot paw volume was measured immediately before and 3 hours after carrageenan challenge using plethysmography (Ugo-Basile). The difference between the 3-hour and the initial paw volume was for each test group of animals was used to calculate the percent inhibition of edema achieved by the test compound at the test dose compared with vehicle control group. Compounds reducing edema by at least about 10% compared to the vehicle control group were considered to be effective in this assay.

Example IV

Adjuvant Assay

In this example, the ability of compounds of formula I to reduce adjuvant-induced footpad edema in rats is demonstrated. This assay is a model for chronic inflammation. See, for example, B. M. Weichman, "Rat Adjuvant Arthritis: A Model of Chronic Inflammation," *Pharmacological Methods in the Control of Inflammation*, 363-380 (1989) and references cited therein.

In this assay, Male Lewis rats weighing between 180-220 g were lightly anesthetized with an ip injection of 30 mg/kg of sodium pentobarbital (50 mg/mL). Desiccated *Mycobacterium butyrium* (Difco, 20 mg/mL) suspended in mineral oil was injected (50 μ L) at both sides of the base of the tail under the skin. A line was tattooed on both rear paws at 5 mm above the angle of the ankle. The paw volumes, under the line, were measured by volume displacement using a plethysmometer (Ugo Basile) at the day of adjuvant injection (day 0) and on day 14. On day 14, animals with paw volumes equal to the mean of paw volumes $\pm$ SD were randomized into treatment groups. Rats which fell outside $\pm$ 1 SD were not used in the experiment. One group received vehicle (1% methyl cellulose) by the po route and the other group received indomethacin (3 mg/kg suspended in 1% methyl cellulose). Dosing began on day 14, and continued until final assessment on day 21 post-adjuvant injection. A separate group, which did not receive

-- 48 --

adjuvant or test compound, was also monitored as a control. This group has a slightly positive volume increase when paw volumes on day 21 are subtracted from day 0 values due to growth of the rat. Indomethacin (3 mg/kg, po), a known anti-inflammation compound, significantly reduced paw volume as compared to vehicle controls. Compounds reducing paw volume by at least about 30% compared to vehicle control group were considered effective in this test.

Example V

Collagen Arthritis Assay

In this example, the ability of compounds of formula I to reduce collagen footpad edema in rats is demonstrated. This assay is commonly used to screen and evaluate anti-inflammatory drug candidates. See, for example, Larsson et al., *Arthritis & Rheumatism*, 33:693-701, 1990 and references cited therein.

For these experiments, Female DA rats (7-8 weeks of age) were immunized with Type II collagen derived from bovine nasal septum as described in Cremer et al., *J. of Immunology*, 149:1045-1053, 1992. The collagen was dissolved and administered with incomplete Freund's adjuvant. Standard precautions were taken to avoid denaturing the collagen before its administration such as keeping the solution cold during preparation. Rats were immunized in the base of the tail on day 0. Dosing began on day 10 with rats scored on day 21 on a severity scale. Rats are scored for gait (0-3), swelling (metatarsals, ankle, carpals, metacarpals, 0-3) with the highest scores given for the most swelling or impairment. All scoring was done in a blinded manner. Compounds reducing scores by at least about 30% compared to the controls were considered effective in this assay. The arthritis was also evaluated by comparing paw weights at day 21. Compounds reducing paw weight by at least about 30% compared to the controls were also considered effective in this assay.

30

Assay Results:

Each compound of formula I that was tested in the above assays was found to be effective for reducing the induction of PGE₂ and/or effective in the carrageenan, adjuvant and/or collagen assay.

5

Example VI**Electron Spin Resonance (ESR) Study**

Using the following procedures, the nitrones of this invention could be shown to trap free radicals using ESR spin trapping techniques. For additional
10 experimental details, see, for example, K. R. Maples et al., "In Vivo Detection of Free Radical Metabolites", *Free Radicals in Synthesis and Biology* (F. Minisci, ed.) pp. 423-436 (Kluwer Academic Publishers, Boston, 1989); and J. A. DeGray et al., "Biological Spin Trapping", *Electron Spin Resonance* 14:246-300 (1994).
In this experiment, a *t*-butyl hydroperoxide/ferrous iron free radical generating
15 system is used. This free radical generating system produces *t*-butyl-alkoxyl radicals, *t*-butyl-peroxyl radicals, and methyl radicals. If the nitrones of this invention are capable of trapping any of these radicals to form a stable radical adduct, such radical adducts should be detectable by ESR spectroscopy.

20 To 490 μ l of a 100 mM solution of the nitron in water is added 5 μ l of 100 mM ferrous sulfate. The reaction is initiated by the addition of 5 μ l of 100 mM *t*-butyl hydroperoxide. The final concentrations of reagents are 1 mM ferrous iron, 1 mM *t*-butyl hydroperoxide and 98 mM of the nitron compound in water. Once mixed, the solution is quickly transferred into a quartz flat cell and this cell
25 is placed in the cavity of a Bruker ESP 300 ESR spectrometer, and scanned within 5 minutes of mixing. ESR spectrometer settings are: 3480 G center field, 200 G field width, 480 seconds sweep time, 9.76 GHz frequency, 10 dB power, 1.6×10^5 receiver gain, 0.200 G modulation amplitude, 0.320 second time constant, and 270° phase. The resulting ESR spectrum would show that the nitron is effective
30 at trapping free radicals and that such compounds can be used as analytical reagents for ESR applications.


-- 50 --

From the foregoing description, various modifications and changes in the compositions and methods of this invention will occur to those skilled in the art. All such modifications coming within the scope of the appended claims are intended to be included therein.

WHAT IS CLAIMED IS:

1. A compound of formula I:

5




I

10

wherein


R¹ is selected from the group consisting of:

15



each R² is independently selected from a group of the formula:

20



25

R³ is selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl;

R⁴ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

30

R⁵ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;



This Sheet Has Been Replaced



5 R^6 and R^7 are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R^6 and R^7 can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R^8 is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

10 R^9 is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R^8 and R^9 can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

15 R^{10} is selected from the group consisting of hydrogen, lower alkyl and lower cycloalkyl; or R^1 and R^{10} can be joined to form an alkylene, substituted alkylene, -C(O)- -S(O)- or -S(O)₂- group;

R^{11} and R^{12} are independently selected from the group consisting of lower alkyl and lower cycloalkyl; or R^{11} and R^{12} can be joined to form an alkylene group having from 2 to 10 carbon atoms;

20 X is oxygen, sulfur, -S(O)- or -S(O)₂-; and

W is oxygen or sulfur; and pharmaceutically-acceptable salts thereof.

2. The compound of Claim 1 wherein W is oxygen.

25 3. The compound of Claim 2 wherein R^3 is hydrogen or lower alkyl.

4. The compound of Claim 3 wherein R^3 is hydrogen.

30 5. The compound of Claim 4 wherein R^4 is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.



This Sheet Has Been Replaced



6. The compound of Claim 5 wherein R^4 is selected from the group consisting of methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

5

7. The compound of Claim 4 wherein R^5 is selected from the group consisting of alkyl and cycloalkyl.

8. The compound of Claim 7 wherein R^5 is selected from the group consisting of methyl, ethyl, *n*-propyl, isopropyl and *n*-butyl.

10

9. The compound of Claim 4 wherein R^7 is hydrogen and R^6 is selected from the group consisting of alkyl and alkoxycarbonylalkyl.

15

10. The compound of Claim 9 wherein R^6 groups is selected from the group consisting of ethyl, *n*-propyl, isopropyl, *n*-butyl, ethoxycarbonylmethyl and 2-(ethoxycarbonyl)ethyl.

20

11. The compound of Claim 4 wherein X is oxygen; R^9 is hydrogen; and R^8 is alkyl or alkoxyalkyl.

12. The compound of Claim 11 wherein R^8 is selected from the group consisting of methyl and methoxyethyl.

25


13. The compound of Claim 4 wherein R^{10} , R^{11} and R^{12} are independently lower alkyl.

14. The compound of Claim 13 wherein R^{10} , R^{11} and R^{12} are methyl.


30



This Sheet Has Been Replaced



15. A compound of formula II:



wherein

10 R^{13} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl;


R^{14} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.

16. The compound of Claim 15 wherein R^{13} is lower alkyl.

17. The compound of Claim 15 wherein R^{14} is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.

18. The compound of Claim 17 wherein R^{14} is selected from the group consisting of methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

19. A compound of formula III:



This Sheet Has Been Replaced



wherein

R¹⁸ is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl;

5 R¹⁹ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; or R¹⁸ and R¹⁹ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R²⁰ is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.

10

25. The compound of Claim 24 wherein R¹⁹ is hydrogen and R¹⁸ is alkyl or alkoxyalkyl.

15

26. The compound of Claim 25 wherein R¹⁸ is methyl or methoxyethyl.

27. The compound of Claim 24 wherein R²⁰ is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.

20

28. The compound of Claim 27 wherein R²⁰ is selected from the group consisting of methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

25 29. A compound selected from the group consisting of:

α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

α -(4-isobutanoyloxy-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

30

α -(4-*n*-butanoyloxy-3,5-di-*tert*-butylphenyl)-*N*-*tert*-butylnitrone

α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-isopropylnitrone

α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-1-hydroxy-2-methylprop-2-yl
35 ylnitrone



This Sheet Has Been Replaced



α -(4-*n*-pentanoyloxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitrone

α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-4-trifluoromethylbenzylnitrone

5 α -(4-propionyloxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitrone

α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-methylnitrone

10 α -(4-acetoxy-3,5-di-*tert*-butylphenyl)-*N*-3,4,5-trimethoxybenzylnitrone

α -[4-(ethylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

15 α -[4-(*n*-propylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

α -[4-(*n*-butylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

20 α -[4-(2-ethoxycarbonyl)ethylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

α -[4-(2-ethoxycarbonyl)methylaminocarbonyloxy)-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

25 α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N-tert*-butylnitrone

α -[4-(2-methoxy)ethoxymethoxy-3,5-di-*tert*-butylphenyl]-*N-tert*-butylnitrone

30 α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-3-(thiomethoxy)but-1-ynitrone

35 α -(4-methoxymethoxy-3,5-di-*tert*-butylphenyl)-*N*-3-thiomethoxypropylnitrone


and pharmaceutically acceptable salts thereof.


30. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula I:

40




This Sheet Has Been Replaced






wherein


R¹ is selected from the group consisting of:



each R² is independently selected from a group of the formula:



This Sheet Has Been Replaced



R³ is selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl;

R⁴ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

R⁵ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

R⁶ and R⁷ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted



* C L M S *

cycloalkenyl; or R⁶ and R⁷ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R⁸ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

R⁹ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R⁸ and R⁹ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R¹⁰ is selected from the group consisting of hydrogen, lower alkyl and lower cycloalkyl; or R¹ and R¹⁰ can be joined to form an alkylene, substituted alkylene, -C(O)- -S(O)- or -S(O)₂- group;

R¹¹ and R¹² are independently selected from the group consisting of lower alkyl and lower cycloalkyl; or R¹¹ and R¹² can be joined to form an alkylene group having from 2 to 10 carbon atoms;

X is oxygen, sulfur, -S(O)- or -S(O)₂-; and

W is oxygen or sulfur; and pharmaceutically-acceptable salts thereof.

31. The pharmaceutical composition of Claim 30 wherein W is oxygen.

32. The pharmaceutical composition of Claim 31 wherein R³ is hydrogen or lower alkyl.

33. The pharmaceutical composition of Claim 3 wherein R³ is hydrogen.

34. The pharmaceutical composition of Claim 33 wherein R⁴ is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.

35. The pharmaceutical composition of Claim 34 wherein R⁴ is selected from the group consisting of methyl, n-propyl, isopropyl, 1-hydroxy-2-

This Sheet Has Been Replaced



-- 60 --

methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

5 36. The pharmaceutical composition of Claim 33 wherein R⁵ is selected from the group consisting of alkyl and cycloalkyl.

10 37. The pharmaceutical composition of Claim 36 wherein R⁵ is selected from the group consisting of methyl, ethyl, *n*-propyl, isopropyl and *n*-butyl.

 38. The pharmaceutical composition of Claim 33 wherein R⁷ is hydrogen and R⁶ is selected from the group consisting of alkyl and alkoxyacetylalkyl.

15 39. The pharmaceutical composition of Claim 38 wherein R⁶ groups is selected from the group consisting of ethyl, *n*-propyl, isopropyl, *n*-butyl, ethoxycarbonylmethyl and 2-(ethoxycarbonyl)ethyl.

20 40. The pharmaceutical composition of Claim 33 wherein X is oxygen; R⁹ is hydrogen; and R⁸ is alkyl or alkoxyalkyl.

 41. The pharmaceutical composition of Claim 40 wherein R⁸ is selected from the group consisting of methyl and methoxyethyl.

25 42. The pharmaceutical composition of Claim 33 wherein R¹⁰, R¹¹ and R¹² are independently lower alkyl.


30 43. The pharmaceutical composition of Claim 42 wherein R¹⁰, R¹¹ and R¹² are methyl.



This Sheet Has Been Replaced



44. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula II:



wherein

R^{13} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl;

R^{14} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.

45. The pharmaceutical composition of Claim 44 wherein R^{13} is lower alkyl.

46. The pharmaceutical composition of Claim 44 wherein R^{14} is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.

47. The pharmaceutical composition of Claim 46 wherein R^{14} is selected from the group consisting of methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.


48. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula III:



This Sheet Has Been Replaced



5



wherein

10 R^{15} and R^{16} are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl; or R^{15} and R^{16} can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

15 R^{17} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.

20 49. The pharmaceutical composition of Claim 48 wherein R^{16} is hydrogen and R^{15} is selected from the group consisting of alkyl and alkoxy-carbonylalkyl.

50. The pharmaceutical composition of Claim 49 wherein R^{15} is selected from the group consisting of ethyl, *n*-propyl, isopropyl, *n*-butyl, ethoxycarbonylmethyl and 2-(ethoxycarbonyl)ethyl.

25 51. The pharmaceutical composition of Claim 48 wherein R^{17} is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.


30 52. The pharmaceutical composition of Claim 51 wherein R^{17} is selected from the group consisting of methyl, *n*-propyl, isopropyl, 1-hydroxy-2-methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.



This Sheet Has Been Replaced



53. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a pharmaceutically effective amount of a compound of formula IV:



wherein

R^{18} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl;

R^{19} is selected from the group consisting of hydrogen, alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; or R^{18} and R^{19} can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R^{20} is selected from the group consisting of alkyl, substituted alkyl, cycloalkyl and substituted cycloalkyl; and pharmaceutically-acceptable salts thereof.

54. The pharmaceutical composition of Claim 53 wherein R^{19} is hydrogen and R^{18} is alkyl or alkoxyalkyl.

55. The pharmaceutical composition of Claim 54 wherein R^{18} is methyl or methoxyethyl.

56. The pharmaceutical composition of Claim 53 wherein R^{20} is selected from the group consisting of alkyl, substituted alkyl and cycloalkyl.

57. The pharmaceutical composition of Claim 56 wherein R^{20} is selected from the group consisting of methyl, *n*-propyl, isopropyl, 1-hydroxy-2-




This Sheet Has Been Replaced




methylprop-2-yl, *n*-butyl, *tert*-butyl, 3-thiomethylpropyl, 3-(thiomethoxy)but-1-yl, cyclohexyl, 4-trifluoromethylbenzyl and 3,4,5-trimethoxybenzyl.

58. A method for treating a mammal with an inflammation-related condition which method comprises administering to said mammal a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective inflammation-reducing amount of a compound of formula I:



15 wherein
R¹ is selected from the group consisting of:



each R² is independently selected from a group of the formula:



30 R³ is selected from the group consisting of hydrogen, alkyl, cycloalkyl and aryl;



This Sheet Has Been Replaced



R⁴ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

5 R⁵ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

10 R⁶ and R⁷ are independently selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R⁶ and R⁷ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

R⁸ is selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl;

15 R⁹ is selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkenyl, substituted alkenyl, alkynyl, substituted alkynyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl and substituted cycloalkenyl; or R⁸ and R⁹ can be joined to form an alkylene or substituted alkylene group having from 2 to 10 carbon atoms;

20 R¹⁰ is selected from the group consisting of hydrogen, lower alkyl and lower cycloalkyl; or R¹ and R¹⁰ can be joined to form an alkylene, substituted alkylene, -C(O)- -S(O)- or -S(O)₂- group;

25 R¹¹ and R¹² are independently selected from the group consisting of lower alkyl and lower cycloalkyl; or R¹¹ and R¹² can be joined to form an alkylene group having from 2 to 10 carbon atoms;


X is oxygen, sulfur, -S(O)- or -S(O)₂-; and

W is oxygen or sulfur; and pharmaceutically-acceptable salts thereof.

30 59. The method according to Claim 58 wherein the inflammation-related condition is rheumatoid arthritis.



This Sheet Has Been Replaced



60. The method according to Claim 58 wherein the inflammation-related condition is osteoarthritis.

5 61. The method according to Claim 58 wherein the inflammation-related condition is ankylosing spondylitis.

62. The method according to Claim 58 wherein the inflammation-related condition is systemic lupus erythematosus.



This Sheet Has Been Replaced

