

CANCEROUS DISEASE MODIFYING ANTIBODIES

STATEMENT OF COOPERATIVE RESEARCH AGREEMENT

The present invention, as defined by the claims herein, was made by parties to a Joint Research Agreement ("Agreement") between Arius Research Inc. and Takeda
5 Pharmaceutical Company Limited, as a result of activities undertaken within the scope of that Agreement. The Agreement was in effect prior to the date of the invention.

FIELD OF THE INVENTION

This invention relates to the isolation and production of cancerous disease
modifying antibodies (CDMAB) and to the use of these CDMAB in therapeutic and diagnostic
10 processes, optionally in combination with one or more chemotherapeutic agents. The invention further relates to binding assays which utilize the CDMAB of the instant invention.

BACKGROUND OF THE INVENTION

Monoclonal Antibodies as Cancer Therapy: Each individual who presents with cancer is unique and has a cancer that is as different from other cancers as that person's identity.
15 Despite this, current therapy treats all patients with the same type of cancer, at the same stage, in the same way. At least 30 percent of these patients will fail the first line therapy, thus leading to further rounds of treatment and the increased probability of treatment failure, metastases, and ultimately, death. A superior approach to treatment would be the customization of therapy for the particular individual. The only current therapy which lends itself to customization is surgery.
20 Chemotherapy and radiation treatment cannot be tailored to the patient, and surgery by itself, in most cases is inadequate for producing cures.

With the advent of monoclonal antibodies, the possibility of developing methods for customized therapy became more realistic since each antibody can be directed to a single epitope. Furthermore, it is possible to produce a combination of antibodies that are directed to
25 the constellation of epitopes that uniquely define a particular individual's tumor.

Having recognized that a significant difference between cancerous and normal cells is that cancerous cells contain antigens that are specific to transformed cells, the scientific community has long held that monoclonal antibodies can be designed to specifically target transformed cells by binding specifically to these cancer antigens; thus giving rise to the belief

that monoclonal antibodies can serve as "Magic Bullets" to eliminate cancer cells. However, it is now widely recognized that no single monoclonal antibody can serve in all instances of cancer, and that monoclonal antibodies can be deployed, as a class, as targeted cancer treatments. Monoclonal antibodies isolated in accordance with the teachings of the instantly disclosed invention have been shown to modify the cancerous disease process in a manner which is beneficial to the patient, for example by reducing the tumor burden, and will variously be referred to herein as cancerous disease modifying antibodies (CDMAB) or "anti-cancer" antibodies.

At the present time, the cancer patient usually has few options of treatment. The regimented approach to cancer therapy has produced improvements in global survival and morbidity rates. However, to the particular individual, these improved statistics do not necessarily correlate with an improvement in their personal situation.

Thus, if a methodology was put forth which enabled the practitioner to treat each tumor independently of other patients in the same cohort, this would permit the unique approach of tailoring therapy to just that one person. Such a course of therapy would, ideally, increase the rate of cures, and produce better outcomes, thereby satisfying a long-felt need.

Historically, the use of polyclonal antibodies has been used with limited success in the treatment of human cancers. Lymphomas and leukemias have been treated with human plasma, but there were few prolonged remission or responses. Furthermore, there was a lack of reproducibility and there was no additional benefit compared to chemotherapy. Solid tumors such as breast cancers, melanomas and renal cell carcinomas have also been treated with human blood, chimpanzee serum, human plasma and horse serum with correspondingly unpredictable and ineffective results.

There have been many clinical trials of monoclonal antibodies for solid tumors. In the 1980s there were at least four clinical trials for human breast cancer which produced only one responder from at least 47 patients using antibodies against specific antigens or based on tissue selectivity. It was not until 1998 that there was a successful clinical trial using a humanized anti-Her2/neu antibody (Herceptin®) in combination with CISPLATIN. In this trial 37 patients were assessed for responses of which about a quarter had a partial response rate and an additional

quarter had minor or stable disease progression. The median time to progression among the responders was 8.4 months with median response duration of 5.3 months.

Herceptin[®] was approved in 1998 for first line use in combination with Taxol[®]. Clinical study results showed an increase in the median time to disease progression for those who received antibody therapy plus Taxol[®] (6.9 months) in comparison to the group that received Taxol[®] alone (3.0 months). There was also a slight increase in median survival; 22 versus 18 months for the Herceptin[®] plus Taxol[®] treatment arm versus the Taxol[®] treatment alone arm. In addition, there was an increase in the number of both complete (8 versus 2 percent) and partial responders (34 versus 15 percent) in the antibody plus Taxol[®] combination group in comparison to Taxol[®] alone. However, treatment with Herceptin[®] and Taxol[®] led to a higher incidence of cardiotoxicity in comparison to Taxol[®] treatment alone (13 versus 1 percent respectively). Also, Herceptin[®] therapy was only effective for patients who over express (as determined through immunohistochemistry (IHC) analysis) the human epidermal growth factor receptor 2 (Her2/neu), a receptor, which currently has no known function or biologically important ligand; approximately 25 percent of patients who have metastatic breast cancer. Therefore, there is still a large unmet need for patients with breast cancer. Even those who can benefit from Herceptin[®] treatment would still require chemotherapy and consequently would still have to deal with, at least to some degree, the side effects of this kind of treatment.

The clinical trials investigating colorectal cancer involve antibodies against both glycoprotein and glycolipid targets. Antibodies such as 17-1A, which has some specificity for adenocarcinomas, has undergone Phase 2 clinical trials in over 60 patients with only 1 patient having a partial response. In other trials, use of 17-1A produced only 1 complete response and 2 minor responses among 52 patients in protocols using additional cyclophosphamide. To date, Phase III clinical trials of 17-1A have not demonstrated improved efficacy as adjuvant therapy for stage III colon cancer. The use of a humanized murine monoclonal antibody initially approved for imaging also did not produce tumor regression.

Only recently have there been any positive results from colorectal cancer clinical studies with the use of monoclonal antibodies. In 2004, ERBITUX[®] was approved for the second line treatment of patients with EGFR-expressing metastatic colorectal cancer who are refractory to irinotecan-based chemotherapy. Results from both a two-arm Phase II clinical

study and a single arm study showed that ERBITUX[®] in combination with irinotecan had a response rate of 23 and 15 percent respectively with a median time to disease progression of 4.1 and 6.5 months respectively. Results from the same two-arm Phase II clinical study and another single arm study showed that treatment with ERBITUX[®] alone resulted in an 11 and 9 percent response rate respectively with a median time to disease progression of 1.5 and 4.2 months respectively.

Consequently in both Switzerland and the United States, ERBITUX[®] treatment in combination with irinotecan, and in the United States, ERBITUX[®] treatment alone, has been approved as a second line treatment of colon cancer patients who have failed first line irinotecan therapy. Therefore, like Herceptin[®], treatment in Switzerland is only approved as a combination of monoclonal antibody and chemotherapy. In addition, treatment in both Switzerland and the US is only approved for patients as a second line therapy. Also, in 2004, AVASTIN[®] was approved for use in combination with intravenous 5-fluorouracil-based chemotherapy as a first line treatment of metastatic colorectal cancer. Phase III clinical study results demonstrated a prolongation in the median survival of patients treated with AVASTIN[®] plus 5-fluorouracil compared to patients treated with 5-fluorouracil alone (20 months versus 16 months respectively). However, again like Herceptin[®] and ERBITUX[®], treatment is only approved as a combination of monoclonal antibody and chemotherapy.

There also continues to be poor results for lung, brain, ovarian, pancreatic, prostate, and stomach cancer. The most promising recent results for non-small cell lung cancer came from a Phase II clinical trial where treatment involved a monoclonal antibody (SGN-15; dox-BR96, anti-Sialyl-LeX) conjugated to the cell-killing drug doxorubicin in combination with the chemotherapeutic agent TAXOTERE[®]. TAXOTERE[®] is the only FDA approved chemotherapy for the second line treatment of lung cancer. Initial data indicate an improved overall survival compared to TAXOTERE[®] alone. Out of the 62 patients who were recruited for the study, two-thirds received SGN-15 in combination with TAXOTERE[®] while the remaining one-third received TAXOTERE[®] alone. For the patients receiving SGN-15 in combination with TAXOTERE[®], median overall survival was 7.3 months in comparison to 5.9 months for patients receiving TAXOTERE[®] alone. Overall survival at 1 year and 18 months was 29 and 18 percent

respectively for patients receiving SNG-15 plus TAXOTERE® compared to 24 and 8 percent respectively for patients receiving TAXOTERE® alone. Further clinical trials are planned.

Preclinically, there has been some limited success in the use of monoclonal antibodies for melanoma. Very few of these antibodies have reached clinical trials and to date
5 none have been approved or demonstrated favorable results in Phase III clinical trials.

The discovery of new drugs to treat disease is hindered by the lack of identification of relevant targets among the products of 30,000 known genes that could contribute to disease pathogenesis. In oncology research, potential drug targets are often selected simply due to the fact that they are over-expressed in tumor cells. Targets thus identified are then
10 screened for interaction with a multitude of compounds. In the case of potential antibody therapies, these candidate compounds are usually derived from traditional methods of monoclonal antibody generation according to the fundamental principles laid down by Kohler and Milstein (1975, Nature, 256, 495-497, Kohler and Milstein). Spleen cells are collected from mice immunized with antigen (e.g. whole cells, cell fractions, purified antigen) and fused with
15 immortalized hybridoma partners. The resulting hybridomas are screened and selected for secretion of antibodies which bind most avidly to the target. Many therapeutic and diagnostic antibodies directed against cancer cells, including Herceptin® and RITUXIMAB, have been produced using these methods and selected on the basis of their affinity. The flaws in this strategy are two-fold. Firstly, the choice of appropriate targets for therapeutic or diagnostic
20 antibody binding is limited by the paucity of knowledge surrounding tissue specific carcinogenic processes and the resulting simplistic methods, such as selection by overexpression, by which these targets are identified. Secondly, the assumption that the drug molecule that binds to the receptor with the greatest affinity usually has the highest probability for initiating or inhibiting a signal may not always be the case.

25 Despite some progress with the treatment of breast and colon cancer, the identification and development of efficacious antibody therapies, either as single agents or co-treatments, has been inadequate for all types of cancer.

Prior Patents:

U.S. Patent No. 5,750,102 discloses a process wherein cells from a patient's tumor are transfected with MHC genes which may be cloned from cells or tissue from the patient. These transfected cells are then used to vaccinate the patient.

5 U.S. Patent No. 4,861,581 discloses a process comprising the steps of obtaining monoclonal antibodies that are specific to an internal cellular component of neoplastic and normal cells of the mammal but not to external components, labeling the monoclonal antibody, contacting the labeled antibody with tissue of a mammal that has received therapy to kill
10 neoplastic cells, and determining the effectiveness of therapy by measuring the binding of the labeled antibody to the internal cellular component of the degenerating neoplastic cells. In preparing antibodies directed to human intracellular antigens, the patentee recognizes that malignant cells represent a convenient source of such antigens.

U.S. Patent No. 5,171,665 provides a novel antibody and method for its production. Specifically, the patent teaches formation of a monoclonal antibody which has the
15 property of binding strongly to a protein antigen associated with human tumors, e.g. those of the colon and lung, while binding to normal cells to a much lesser degree.

U.S. Patent No. 5,484,596 provides a method of cancer therapy comprising surgically removing tumor tissue from a human cancer patient, treating the tumor tissue to obtain tumor cells, irradiating the tumor cells to be viable but non-tumorigenic, and using these cells to
20 prepare a vaccine for the patient capable of inhibiting recurrence of the primary tumor while simultaneously inhibiting metastases. The patent teaches the development of monoclonal antibodies which are reactive with surface antigens of tumor cells. As set forth at col. 4, lines 45 et seq., the patentees utilize autochthonous tumor cells in the development of monoclonal antibodies expressing active specific immunotherapy in human neoplasia.

25 U.S. Patent No. 5,693,763 teaches a glycoprotein antigen characteristic of human carcinomas and not dependent upon the epithelial tissue of origin.

U.S. Patent No. 5,783,186 is drawn to Anti-Her2 antibodies which induce apoptosis in Her2 expressing cells, hybridoma cell lines producing the antibodies, methods of treating cancer using the antibodies and pharmaceutical compositions including said antibodies.

U.S. Patent No. 5,849,876 describes new hybridoma cell lines for the production of monoclonal antibodies to mucin antigens purified from tumor and non-tumor tissue sources.

U.S. Patent No. 5,869,268 is drawn to a method for generating a human lymphocyte producing an antibody specific to a desired antigen, a method for producing a monoclonal antibody, as well as monoclonal antibodies produced by the method. The patent is particularly drawn to the production of an anti-HD human monoclonal antibody useful for the diagnosis and treatment of cancers.

U.S. Patent No. 5,869,045 relates to antibodies, antibody fragments, antibody conjugates and single-chain immunotoxins reactive with human carcinoma cells. The mechanism by which these antibodies function is two-fold, in that the molecules are reactive with cell membrane antigens present on the surface of human carcinomas, and further in that the antibodies have the ability to internalize within the carcinoma cells, subsequent to binding, making them especially useful for forming antibody-drug and antibody-toxin conjugates. In their unmodified form the antibodies also manifest cytotoxic properties at specific concentrations.

U.S. Patent No. 5,780,033 discloses the use of autoantibodies for tumor therapy and prophylaxis. However, this antibody is an antinuclear autoantibody from an aged mammal. In this case, the autoantibody is said to be one type of natural antibody found in the immune system. Because the autoantibody comes from "an aged mammal", there is no requirement that the autoantibody actually comes from the patient being treated. In addition the patent discloses natural and monoclonal antinuclear autoantibody from an aged mammal, and a hybridoma cell line producing a monoclonal antinuclear autoantibody.

SUMMARY OF THE INVENTION

This application utilizes methodology for producing patient specific anti-cancer antibodies taught in the U.S. 6,180,357 patent for isolating hybridoma cell lines which encode for cancerous disease modifying monoclonal antibodies. These antibodies can be made specifically for one tumor and thus make possible the customization of cancer therapy. Within the context of this application, anti-cancer antibodies having either cell-killing (cytotoxic) or cell-growth inhibiting (cytostatic) properties will hereafter be referred to as cytotoxic. These antibodies can be used in aid of staging and diagnosis of a cancer, and can be used to treat tumor

metastases. These antibodies can also be used for the prevention of cancer by way of prophylactic treatment. Unlike antibodies generated according to traditional drug discovery paradigms, antibodies generated in this way may target molecules and pathways not previously shown to be integral to the growth and/or survival of malignant tissue. Furthermore, the binding
5 affinities of these antibodies are suited to requirements for initiation of the cytotoxic events that may not be amenable to stronger affinity interactions. Also, it is within the purview of this invention to conjugate standard chemotherapeutic modalities, e.g. radionuclides, with the CDMAB of the instant invention, thereby focusing the use of said chemotherapeutics. The CDMAB can also be conjugated to toxins, cytotoxic moieties, enzymes e.g. biotin conjugated
10 enzymes, or hematogenous cells, thereby forming an antibody conjugate.

The prospect of individualized anti-cancer treatment will bring about a change in the way a patient is managed. A likely clinical scenario is that a tumor sample is obtained at the time of presentation, and banked. From this sample, the tumor can be typed from a panel of pre-existing cancerous disease modifying antibodies. The patient will be conventionally staged but
15 the available antibodies can be of use in further staging the patient. The patient can be treated immediately with the existing antibodies, and a panel of antibodies specific to the tumor can be produced either using the methods outlined herein or through the use of phage display libraries in conjunction with the screening methods herein disclosed. All the antibodies generated will be added to the library of anti-cancer antibodies since there is a possibility that other tumors can
20 bear some of the same epitopes as the one that is being treated. The antibodies produced according to this method may be useful to treat cancerous disease in any number of patients who have cancers that bind to these antibodies.

In addition to anti-cancer antibodies, the patient can elect to receive the currently recommended therapies as part of a multi-modal regimen of treatment. The fact that the
25 antibodies isolated via the present methodology are relatively non-toxic to non-cancerous cells allows for combinations of antibodies at high doses to be used, either alone, or in conjunction with conventional therapy. The high therapeutic index will also permit re-treatment on a short time scale that should decrease the likelihood of emergence of treatment resistant cells.

If the patient is refractory to the initial course of therapy or metastases develop,
30 the process of generating specific antibodies to the tumor can be repeated for re-treatment.

Furthermore, the anti-cancer antibodies can be conjugated to red blood cells obtained from that patient and re-infused for treatment of metastases. There have been few effective treatments for metastatic cancer and metastases usually portend a poor outcome resulting in death. However, metastatic cancers are usually well vascularized and the delivery of anti-cancer antibodies by red blood cells can have the effect of concentrating the antibodies at the site of the tumor. Even prior to metastases, most cancer cells are dependent on the host's blood supply for their survival and an anti-cancer antibody conjugated to red blood cells can be effective against *in situ* tumors as well. Alternatively, the antibodies may be conjugated to other hematogenous cells, e.g. lymphocytes, macrophages, monocytes, natural killer cells, etc.

There are five classes of antibodies and each is associated with a function that is conferred by its heavy chain. It is generally thought that cancer cell killing by naked antibodies are mediated either through antibody dependent cellular cytotoxicity or complement dependent cytotoxicity. For example murine IgM and IgG2a antibodies can activate human complement by binding the C-1 component of the complement system thereby activating the classical pathway of complement activation which can lead to tumor lysis. For human antibodies the most effective complement activating antibodies are generally IgM and IgG1. Murine antibodies of the IgG2a and IgG3 isotype are effective at recruiting cytotoxic cells that have Fc receptors which will lead to cell killing by monocytes, macrophages, granulocytes and certain lymphocytes. Human antibodies of both the IgG1 and IgG3 isotype mediate ADCC.

Another possible mechanism of antibody mediated cancer killing may be through the use of antibodies that function to catalyze the hydrolysis of various chemical bonds in the cell membrane and its associated glycoproteins or glycolipids, so-called catalytic antibodies.

There are three additional mechanisms of antibody-mediated cancer cell killing. The first is the use of antibodies as a vaccine to induce the body to produce an immune response against the putative antigen that resides on the cancer cell. The second is the use of antibodies to target growth receptors and interfere with their function or to down regulate that receptor so that its function is effectively lost. The third is the effect of such antibodies on direct ligation of cell surface moieties that may lead to direct cell death, such as ligation of death receptors such as TRAIL R1 or TRAIL R2, or integrin molecules such as alpha V beta 3 and the like.

The clinical utility of a cancer drug is based on the benefit of the drug under an acceptable risk profile to the patient. In cancer therapy survival has generally been the most sought after benefit, however there are a number of other well-recognized benefits in addition to prolonging life. These other benefits, where treatment does not adversely affect survival, include symptom palliation, protection against adverse events, prolongation in time to recurrence or disease-free survival, and prolongation in time to progression. These criteria are generally accepted and regulatory bodies such as the U.S. Food and Drug Administration (F.D.A.) approve drugs that produce these benefits (Hirschfeld *et al.* Critical Reviews in Oncology/Hematology 42:137-143 2002). In addition to these criteria it is well recognized that there are other endpoints that may presage these types of benefits. In part, the accelerated approval process granted by the U.S. F.D.A. acknowledges that there are surrogates that will likely predict patient benefit. As of year-end 2003, there have been sixteen drugs approved under this process, and of these, four have gone on to full approval, i.e., follow-up studies have demonstrated direct patient benefit as predicted by surrogate endpoints. One important endpoint for determining drug effects in solid tumors is the assessment of tumor burden by measuring response to treatment (Therasse *et al.* Journal of the National Cancer Institute 92(3):205-216 2000). The clinical criteria (RECIST criteria) for such evaluation have been promulgated by Response Evaluation Criteria in Solid Tumors Working Group, a group of international experts in cancer. Drugs with a demonstrated effect on tumor burden, as shown by objective responses according to RECIST criteria, in comparison to the appropriate control group tend to, ultimately, produce direct patient benefit. In the pre-clinical setting tumor burden is generally more straightforward to assess and document. In that pre-clinical studies can be translated to the clinical setting, drugs that produce prolonged survival in pre-clinical models have the greatest anticipated clinical utility. Analogous to producing positive responses to clinical treatment, drugs that reduce tumor burden in the pre-clinical setting may also have significant direct impact on the disease. Although prolongation of survival is the most sought after clinical outcome from cancer drug treatment, there are other benefits that have clinical utility and it is clear that tumor burden reduction, which may correlate to a delay in disease progression, extended survival or both, can also lead to direct benefits and have clinical impact (Eckhardt *et al.* Developmental Therapeutics: Successes and Failures of

Clinical Trial Designs of Targeted Compounds; ASCO Educational Book, 39th Annual Meeting, 2003, pages 209-219).

The present invention describes the development and use of AR27A807.14.1 identified by its effect in a cytotoxic assay and in an animal model of human cancer. This invention describes reagents that bind specifically to an epitope or epitopes present on the target molecule, and that also have *in vitro* cytotoxic properties, as a naked antibody, against malignant tumor cells but not normal cells, and which also directly mediate, as a naked antibody, inhibition of tumor growth. A further advance is of the use of anti-cancer antibodies such as this to target tumors expressing cognate antigen markers to achieve tumor growth inhibition, and other positive endpoints of cancer treatment.

In all, this invention teaches the use of the AR27A807.14.1 antigen as a target for a therapeutic agent, that when administered can reduce the tumor burden of a cancer expressing the antigen in a mammal. This invention also teaches the use of CDMAB (AR27A807.14.1), and their derivatives, and antigen binding fragments thereof, and cytotoxicity inducing ligands thereof, to target their antigen to reduce the tumor burden of a cancer expressing the antigen in a mammal. Furthermore, this invention also teaches the use of detecting the AR27A807.14.1 antigen in cancerous cells that can be useful for the diagnosis, prediction of therapy, and prognosis of mammals bearing tumors that express this antigen.

Accordingly, it is an objective of the invention to utilize a method for producing cancerous disease modifying antibodies (CDMAB) raised against cancerous cells derived from a particular individual, or one or more particular cancer cell lines, which CDMAB are cytotoxic with respect to cancer cells while simultaneously being relatively non-toxic to non-cancerous cells, in order to isolate hybridoma cell lines and the corresponding isolated monoclonal antibodies and antigen binding fragments thereof for which said hybridoma cell lines are encoded.

It is an additional objective of the invention to teach cancerous disease modifying antibodies, ligands and antigen binding fragments thereof.

It is a further objective of the instant invention to produce cancerous disease modifying antibodies whose cytotoxicity is mediated through antibody dependent cellular toxicity.

It is yet an additional objective of the instant invention to produce cancerous disease modifying antibodies whose cytotoxicity is mediated through complement dependent cellular toxicity.

5 It is still a further objective of the instant invention to produce cancerous disease modifying antibodies whose cytotoxicity is a function of their ability to catalyze hydrolysis of cellular chemical bonds.

A still further objective of the instant invention is to produce cancerous disease modifying antibodies which are useful for in a binding assay for diagnosis, prognosis, and monitoring of cancer.

10 Other objects and advantages of this invention will become apparent from the following description wherein are set forth, by way of illustration and example, certain embodiments of this invention.

BRIEF DESCRIPTION OF THE FIGURES

15 Figure 1 compares the percentage cytotoxicity and binding levels of the hybridoma supernatants against cell lines HT-29, SW1116, Lovo and CCD-27sk.

Figure 2 represents binding of AR27A807.14.1 to cancer and normal cell lines. The data is tabulated to present the binding as a fold increase above isotype control.

20 Figure 3 demonstrates the effect of AR27A807.14.1 on tumor growth in a prophylactic DLD-1 colon cancer model. The vertical dashed lines indicate the period during which the antibody was administered. Data points represent the mean +/- SEM.

Figure 4 demonstrates the effect of AR27A807.14.1 on body weight in a prophylactic DLD-1 colon cancer model. Data points represent the mean +/- SEM.

DETAILED DESCRIPTION OF THE INVENTION

25 In general, the following words or phrases have the indicated definition when used in the summary, description, examples, and claims.

The term "antibody" is used in the broadest sense and specifically covers, for example, single monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies, de-immunized, murine, chimerized or humanized antibodies), antibody compositions

with polyepitopic specificity, single-chain antibodies, immunoconjugates and antibody fragments (see below).

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations which include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma (murine or human) method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). The "monoclonal antibodies" may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

"Antibody fragments" comprise a portion of an intact antibody, preferably comprising the antigen-binding or variable region thereof. Examples of antibody fragments include less than full length antibodies, Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules; single-chain antibodies, single domain antibody molecules, fusion proteins, recombinant proteins and multispecific antibodies formed from antibody fragment(s).

An "intact" antibody is one which comprises an antigen-binding variable region as well as a light chain constant domain (C_L) and heavy chain constant domains, C_H1, C_H2 and C_H3. The constant domains may be native sequence constant domains (e.g. human native sequence constant domains) or amino acid sequence variant thereof. Preferably, the intact antibody has one or more effector functions.

Depending on the amino acid sequence of the constant domain of their heavy chains, intact antibodies can be assigned to different “classes”. There are five-major classes of intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into “subclasses” (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

Antibody “effector functions” refer to those biological activities attributable to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody. Examples of antibody effector functions include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor; BCR), etc.

“Antibody-dependent cell-mediated cytotoxicity” and “ADCC” refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express Fc γ RIII only, whereas monocytes express Fc γ RI, Fc γ RII and Fc γ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in U.S. Pat. No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in a animal model such as that disclosed in Clynes *et al. PNAS* (USA) 95:652-656 (1998).

“Effector cells” are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express at least Fc γ RIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source thereof, e.g. from blood or PBMCs as described herein.

The terms "Fc receptor" or "FcR" are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors. FcγRII receptors include FcγRIIA (an "activating receptor") and FcγRIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (see review M. in Daëron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9:457-92 (1991); Capel *et al.*, *Immunomethods* 4:25-34 (1994); and de Haas *et al.*, *J. Lab. Clin. Med.* 126:330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer *et al.*, *J. Immunol.* 117:587 (1976) and Kim *et al.*, *Eur. J. Immunol.* 24:2429 (1994)).

"Complement dependent cytotoxicity" or "CDC" refers to the ability of a molecule to lyse a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (e.g. an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, e.g. as described in Gazzano-Santoro *et al.*, *J. Immunol. Methods* 202:163 (1996) may be performed.

The term "variable" refers to the fact that certain portions of the variable domains differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of variable domains are called the framework regions (FRs). The variable domains of native heavy and light chains each comprise four FRs, largely adopting a β-sheet configuration, connected by three hypervariable regions, which form loops connecting, and

in some cases forming part of, the β -sheet structure. The hypervariable regions in each chain are held together in close proximity by the FRs and, with the hypervariable regions from the other chain, contribute to the formation of the antigen-binding site of antibodies (see Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. pp 15-17; 48-53 (1991)). The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody dependent cellular cytotoxicity (ADCC).

The term "hypervariable region" when used herein refers to the amino acid residues of an antibody which are responsible for antigen-binding. The hypervariable region generally comprises amino acid residues from a "complementarity determining region" or "CDR" (e.g. residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md. pp 15-17; 48-53 (1991)) and/or those residues from a "hypervariable loop" (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). "Framework Region" or "FR" residues are those variable domain residues other than the hypervariable region residues as herein defined. Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an $F(ab')_2$ fragment that has two antigen-binding sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and antigen-binding site. This region consists of a dimer of one heavy chain and one light chain variable domain in tight, non-covalent association. It is in this configuration that the three hypervariable regions of each variable domain interact to define an antigen-binding site on the surface of the V_H - V_L dimer. Collectively, the six hypervariable regions confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three hypervariable regions specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site. The Fab fragment also

contains the constant domain of the light chain and the first constant domain (CH I) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

"Single-chain Fv" or "scFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Plückthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a variable heavy domain (V_H) connected to a variable light domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

An antibody “which binds” an antigen of interest is one capable of binding that antigen with sufficient affinity such that the antibody is useful as a therapeutic or diagnostic agent in targeting a cell expressing the antigen. Where the antibody is one which binds the antigenic moiety it will usually preferentially bind that antigenic moiety as opposed to other
5 receptors, and does not include incidental binding such as non-specific Fc contact, or binding to post-translational modifications common to other antigens and may be one which does not significantly cross-react with other proteins. Methods, for the detection of an antibody that binds an antigen of interest, are well known in the art and can include but are not limited to assays such as FACS, cell ELISA and Western blot.

10 As used herein, the expressions “cell”, “cell line”, and “cell culture” are used interchangeably, and all such designations include progeny. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. It will be clear from the context where distinct
15 designations are intended.

“Treatment or treating” refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be
20 prevented. Hence, the mammal to be treated herein may have been diagnosed as having the disorder or may be predisposed or susceptible to the disorder.

The terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth or death. Examples of cancer include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma,
25 and leukemia or lymphoid malignancies. More particular examples of such cancers include squamous cell cancer (e.g. epithelial squamous cell cancer), lung cancer including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver
30 cancer, bladder cancer, hepatoma, breast cancer, colon cancer, rectal cancer, colorectal cancer,

endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, as well as head and neck cancer.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN™); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carnomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitostanol, mepitostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine;

mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, e.g. paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, N.J.) and docetaxel (TAXOTERE®, Aventis, Rhone-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, mice, SCID or nude mice or strains of mice, domestic and farm animals, and zoo, sports, or pet animals, such as sheep, dogs, horses, cats, cows, etc. Preferably, the mammal herein is human.

"Oligonucleotides" are short-length, single- or double-stranded polydeoxynucleotides that are chemically synthesized by known methods (such as phosphotriester, phosphite, or phosphoramidite chemistry, using solid phase techniques such as described in EP 266,032, published 4 May 1988, or via deoxynucleoside H-phosphonate intermediates as described by Froehler *et al.*, *Nucl. Acids Res.*, 14:5399-5407, 1986. They are then purified on polyacrylamide gels.

In accordance with the present invention, "humanized" and/or "chimeric" forms of non-human (e.g. murine) immunoglobulins refer to antibodies which contain specific chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which results in the decrease of a human anti-mouse antibody (HAMA), human anti-chimeric antibody (HACA) or a human anti-human antibody (HAHA) response, compared to the original antibody, and contain the requisite portions

(e.g. CDR(s), antigen binding region(s), variable domain(s) and so on) derived from said non-human immunoglobulin, necessary to reproduce the desired effect, while simultaneously retaining binding characteristics which are comparable to said non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from the complementarity determining regions (CDRs) of the recipient antibody are replaced by residues from the CDRs of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human FR residues. Furthermore, the humanized antibody may comprise residues which are found neither in the recipient antibody nor in the imported CDR or FR sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR residues are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

“De-immunized” antibodies are immunoglobulins that are non-immunogenic, or less immunogenic, to a given species. De-immunization can be achieved through structural alterations to the antibody. Any de-immunization technique known to those skilled in the art can be employed. One suitable technique for de-immunizing antibodies is described, for example, in WO 00/34317 published June 15, 2000.

An antibody which induces “apoptosis” is one which induces programmed cell death by any means, illustrated by but not limited to binding of annexin V, caspase activity, fragmentation of DNA, cell shrinkage, dilation of endoplasmic reticulum, cell fragmentation, and/or formation of membrane vesicles (called apoptotic bodies).

As used herein “antibody induced cytotoxicity” is understood to mean the cytotoxic effect derived from the hybridoma supernatant or antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 which effect is not necessarily related to the degree of binding.

Throughout the instant specification, hybridoma cell lines, as well as the isolated monoclonal antibodies which are produced therefrom, are alternatively referred to by their internal designation, AR27A807.14.1 or Depository Designation, IDAC 180706-03.

As used herein "antibody-ligand" includes a moiety which exhibits binding
5 specificity for at least one epitope of the target antigen, and which may be an intact antibody molecule, antibody fragments, and any molecule having at least an antigen-binding region or portion thereof (i.e., the variable portion of an antibody molecule), e.g., an Fv molecule, Fab molecule, Fab' molecule, F(ab')₂ molecule, a bispecific antibody, a fusion protein, or any
10 genetically engineered molecule which specifically recognizes and binds at least one epitope of the antigen bound by the isolated monoclonal antibody produced by the hybridoma cell line designated as IDAC 180706-03 (the IDAC 180706-03 antigen).

As used herein "cancerous disease modifying antibodies" (CDMAB) refers to monoclonal antibodies which modify the cancerous disease process in a manner which is beneficial to the patient, for example by reducing tumor burden or prolonging survival of tumor
15 bearing individuals, and antibody-ligands thereof.

As used herein "antigen-binding region" means a portion of the molecule which recognizes the target antigen.

As used herein "competitively inhibits" means being able to recognize and bind a determinant site to which the monoclonal antibody produced by the hybridoma cell line
20 designated as IDAC 180706-03, (the IDAC 180706-03 antibody) is directed using conventional reciprocal antibody competition assays. (Belanger L., Sylvestre C. and Dufour D. (1973), Enzyme linked immunoassay for alpha fetoprotein by competitive and sandwich procedures. Clinica Chimica Acta 48, 15).

As used herein "target antigen" is the IDAC 180706-03 antigen or portions
25 thereof.

As used herein, an "immunoconjugate" means any molecule or CDMAB such as an antibody chemically or biologically linked to a cytotoxin, a radioactive agent, enzyme, toxin, an anti-tumor drug or a therapeutic agent. The antibody or CDMAB may be linked to the cytotoxin, radioactive agent, anti-tumor drug or therapeutic agent at any location along the

molecule so long as it is able to bind its target. Examples of immunoconjugates include antibody toxin chemical conjugates and antibody-toxin fusion proteins.

As used herein, a "fusion protein" means any chimeric protein wherein an antigen binding region is connected to a biologically active molecule, e.g., toxin, enzyme, or protein drug.

In order that the invention herein described may be more fully understood, the following description is set forth.

The present invention provides CDMABs (i.e., IDAC 180706-03 CDMAB) which specifically recognize and bind the IDAC 180706-03 antigen.

The CDMAB of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 may be in any form as long as it has an antigen-binding region which competitively inhibits the immunospecific binding of the isolated monoclonal antibody produced by hybridoma IDAC 180706-03 to its target antigen. Thus, any recombinant proteins (e.g., fusion proteins wherein the antibody is combined with a second protein such as a lymphokine or a tumor inhibitory growth factor) having the same binding specificity as the IDAC 180706-03 antibody fall within the scope of this invention.

In one embodiment of the invention, the CDMAB is the IDAC 180706-03 antibody.

In other embodiments, the CDMAB is an antigen binding fragment which may be a Fv molecule (such as a single-chain Fv molecule), a Fab molecule, a Fab' molecule, a F(ab')₂ molecule, a fusion protein, a bispecific antibody, a heteroantibody or any recombinant molecule having the antigen-binding region of the IDAC 180706-03 antibody. The CDMAB of the invention is directed to the epitope to which the IDAC 180706-03 monoclonal antibody is directed.

The CDMAB of the invention may be modified, i.e., by amino acid modifications within the molecule, so as to produce derivative molecules. Chemical modification may also be possible.

Derivative molecules would retain the functional property of the polypeptide, namely, the molecule having such substitutions will still permit the binding of the polypeptide to the IDAC 180706-03 antigen or portions thereof.

These amino acid substitutions include, but are not necessarily limited to, amino acid substitutions known in the art as "conservative".

For example, it is a well-established principle of protein chemistry that certain amino acid substitutions, entitled "conservative amino acid substitutions," can frequently be made in a protein without altering either the conformation or the function of the protein.

Such changes include substituting any of isoleucine (I), valine (V), and leucine (L) for any other of these hydrophobic amino acids; aspartic acid (D) for glutamic acid (E) and vice versa; glutamine (Q) for asparagine (N) and vice versa; and serine (S) for threonine (T) and vice versa. Other substitutions can also be considered conservative, depending on the environment of the particular amino acid and its role in the three-dimensional structure of the protein. For example, glycine (G) and alanine (A) can frequently be interchangeable, as can alanine and valine (V). Methionine (M), which is relatively hydrophobic, can frequently be interchanged with leucine and isoleucine, and sometimes with valine. Lysine (K) and arginine (R) are frequently interchangeable in locations in which the significant feature of the amino acid residue is its charge and the differing pK's of these two amino acid residues are not significant. Still other changes can be considered "conservative" in particular environments.

EXAMPLE 1

Hybridoma Production – Hybridoma Cell Line AR27A807.14.1

The hybridoma cell line AR27A807.14.1 was deposited, in accordance with the Budapest Treaty, with the International Depository Authority of Canada (IDAC), Bureau of Microbiology, Health Canada, 1015 Arlington Street, Winnipeg, Manitoba, Canada, R3E 3R2, on July 18, 2006, under Accession Number 180706-03. In accordance with 37 CFR 1.808, the depositors assure that all restrictions imposed on the availability to the public of the deposited materials will be irrevocably removed upon the granting of a patent. The deposit will be replaced if the depository cannot dispense viable samples.

To produce the hybridoma that produces the anti-cancer antibody AR27A807.14.1, a single cell suspension of frozen human colon adenocarcinoma tumor tissue (Genomics Collaborative, Cambridge, MA) was prepared in PBS. IMMUNEASY™ (Qiagen, Venlo, Netherlands) adjuvant was prepared for use by gentle mixing. Five to seven week old BALB/c mice were immunized by injecting subcutaneously, 2 million cells in 50 microliters of

the antigen-adjuvant. Recently prepared antigen-adjuvant was used to boost the immunized mice intraperitoneally, 2 weeks after the initial immunization, with 2 million cells in 70 microliters. A spleen was used for fusion three days after the last immunization. The hybridomas were prepared by fusing the isolated splenocytes with NSO-1 myeloma partners. The supernatants from the fusions were tested from subclones of the hybridomas.

To determine whether the antibodies secreted by the hybridoma cells are of the IgG or IgM isotype, an ELISA assay was employed. 100 microliters/well of goat anti-mouse IgG + IgM (H+L) at a concentration of 2.4 micrograms/mL in coating buffer (0.1 M carbonate/bicarbonate buffer, pH 9.2-9.6) at 4°C was added to the ELISA plates overnight. The plates were washed thrice in washing buffer (PBS + 0.05 percent Tween). 100 microliters/well blocking buffer (5 percent milk in wash buffer) was added to the plate for 1 hour at room temperature and then washed thrice in washing buffer. 100 microliters/well of hybridoma supernatant was added and the plate incubated for 1 hour at room temperature. The plates were washed thrice with washing buffer and 1/100,000 dilution of either goat anti-mouse IgG or IgM horseradish peroxidase conjugate (diluted in PBS containing 5 percent milk), 100 microliters/well, was added. After incubating the plate for 1 hour at room temperature the plate was washed thrice with washing buffer. 100 microliters/well of TMB solution was incubated for 1-3 minutes at room temperature. The color reaction was terminated by adding 50 microliters/well 2M H₂SO₄ and the plate was read at 450 nm with a Perkin-Elmer HTS7000 plate reader. As indicated in Figure 1, the AR27A807.14.1 hybridoma secreted primarily antibodies of the IgG isotype.

To determine the subclass of antibody secreted by the hybridoma cells, an isotyping experiment was performed using a Mouse Monoclonal Antibody Isotyping Kit (GE Healthcare, Baie d'Urfé, QC). Antibody-containing hybridoma supernatant was added to a test tube (in a 1:10 dilution with TBS-T) with an isotyping strip carrying goat antibody specific for the different types of peptide chain. The tube was agitated for 15 minutes. The strip was then washed twice with TBS-T for 5 minutes with agitation. A peroxidase-labeled, species-specific anti-mouse antibody was added (in a 1:500 dilution with TBS-T) to the test tube for 15 minutes, to detect the monoclonal antibody bound to the goat antibody on the stick. The strip was again washed twice with TBS-T for 5 minutes with agitation. The peroxidase-labeled antibody bound

to the strip was then detected using a peroxidase substrate system. One 30 mg tablet of 4-chloro-1-naphthol was dissolved in 10 mL cold ethanol, and one drop of hydrogen peroxide solution (30 percent v/v) was diluted in 50 mL TBS. The two solutions were combined immediately before use, and 3 mL was added to the isotyping strip for 15 minutes with agitation. The substrate solution was then discarded and the strip was washed three times with 5 mL distilled water with agitation. The typing stick was then removed from the test tube and analysed for results. The anti-cancer antibody AR27A807.14.1 is of the IgG1, kappa isotype.

After two rounds of limiting dilution, hybridoma supernatants were tested for antibodies that bound to target cells in a cell ELISA assay. Three human colon cancer cell lines and 1 human normal skin cell line were tested: HT-29, SW1116, Lovo and CCD-27sk respectively. All cell lines were obtained from the American Type Tissue Collection (ATCC; Manassas, VA). The plated cells were fixed prior to use. The plates were washed thrice with PBS containing $MgCl_2$ and $CaCl_2$ at room temperature. 100 microliters of 2 percent paraformaldehyde diluted in PBS was added to each well for 10 minutes at room temperature and then discarded. The plates were again washed with PBS containing $MgCl_2$ and $CaCl_2$ three times at room temperature. Blocking was done with 100 microliters/well of 5 percent milk in wash buffer (PBS + 0.05 percent Tween) for 1 hour at room temperature. The plates were washed thrice with wash buffer and the hybridoma supernatant was added at 75 microliters/well for 1 hour at room temperature. The plates were washed 3 times with wash buffer and 100 microliters/well of 1/25,000 dilution of goat anti-mouse IgG or IgM antibody conjugated to horseradish peroxidase (diluted in PBS containing 5 percent milk) was added. After 1 hour incubation at room temperature the plates were washed 3 times with wash buffer and 100 microliter/well of TMB substrate was incubated for 1-3 minutes at room temperature. The reaction was terminated with 50 microliters/well 2M H_2SO_4 and the plate read at 450 nm with a Perkin-Elmer HTS7000 plate reader. The results as tabulated in Figure 1 were expressed as the number of folds above background compared to an in-house IgG isotype control that has previously been shown not to bind to the cell lines tested. The antibodies from the hybridoma AR27A807.14 showed no detectable binding to the cell lines tested.

In conjunction with testing for antibody binding, the cytotoxic effect of the hybridoma supernatants (antibody induced cytotoxicity) was tested in the cell lines: HT-29,

SW1116, Lovo and CCD-27sk. Calcein AM was obtained from Molecular Probes (Eugene, OR) and the assay was performed as outlined below. Cells were plated before the assay at the predetermined appropriate density. After 2 days, 75 microliters of supernatant from the hybridoma microtitre plates were transferred to the cell plates and incubated in a 5 percent CO₂ incubator for 5 days. The wells that served as the positive controls were aspirated until empty and 100 microliters of sodium azide (NaN₃, .01 percent, Sigma, Oakville, ON), cycloheximide (CHX, 0.5 micromolar, Sigma, Oakville, ON) or anti-EGFR antibody (c225, IgG1, kappa, 5 micrograms/mL, Cedarlane, Hornby, ON) dissolved in culture medium, was added. After 5 days of treatment, the plates were then emptied by inverting and blotting dry. Room temperature DPBS (Dulbecco's phosphate buffered saline) containing MgCl₂ and CaCl₂ was dispensed into each well from a multichannel squeeze bottle, tapped 3 times, emptied by inversion and then blotted dry. 50 microliters of the fluorescent calcein dye diluted in DPBS containing MgCl₂ and CaCl₂ was added to each well and incubated at 37°C in a 5 percent CO₂ incubator for 30 minutes. The plates were read in a Perkin-Elmer HTS7000 fluorescence plate reader and the data was analyzed in Microsoft Excel. The results are tabulated in Figure 1. Supernatant from the AR27A807.14.1 hybridoma produced specific cytotoxicity of 39 percent on the HT-29 cells, 29 percent on the SW1116 cells and 20 percent on the Lovo cells. This was 63 and 75 percent of the cytotoxicity obtained with the positive controls sodium azide and cycloheximide respectively for HT-29; 67 and 112 percent of the cytotoxicity of cycloheximide and c225 respectively for SW1116, and likewise, 54 percent of the cytotoxicity of cycloheximide for Lovo. There was no observable cytotoxicity to the normal skin cell line CCD-27sk. The known non-specific cytotoxic agents cycloheximide and NaN₃ generally produced cytotoxicity as expected. The anti-EGFR antibody c225 produced cytotoxicity as expected on SW1116.

Results from Figure 1 demonstrate that the cytotoxic effects of AR27A807.14.1 on the different cell lines did not correlate to the level of binding. Although there was no detectable binding to the three colon cell lines under these conditions, there were detectable levels of cytotoxicity. AR27A807.14.1 did not produce cytotoxicity in, nor did it bind to, the CCD-27sk normal skin cell line. The antibody therefore exhibited functional specificity, which was not necessarily related to the degree of binding.

EXAMPLE 2***In vitro* Binding**

AR27A807.14.1 monoclonal antibody was produced by culturing the hybridoma in CL-1000 flasks (BD Biosciences, Oakville, ON) with collections and reseeded occurring
5 twice/week. Standard antibody purification procedures with Protein G Sepharose 4 Fast Flow (Amersham Biosciences, Baie d'Urfé, QC) were followed. It is within the scope of this invention to utilize monoclonal antibodies that are humanized, de-immunized, chimerized or murine.

Binding of AR27A807.14.1 to breast (MDA-MB-231 and MCF-7), colon (HT-29,
10 Lovo, DLD-1, SW620 and SW1116), prostate (PC-3), pancreatic (BxPC-3) and ovarian (OVCAR-3) cancer, and non-cancer cell lines from skin (CCD-27sk) and lung (Hs888.Lu) was assessed by cell ELISA. All cell lines were obtained from the American Type Tissue Collection (ATCC; Manassas, VA). The same procedure was used as outlined in Example 1 with the exception of the use of purified antibody at a concentration of 20 micrograms/mL versus hybridoma supernatant.

15 The results as tabulated in Figure 2 were expressed as the number of folds above background compared to an in-house IgG isotype control that has previously been shown not to bind to the cell lines tested. AR27A807.14 showed strongest binding to the colon cancer cell line HT-29 (6.1-fold), with lower levels of binding seen in the ovarian cancer cell line OVCAR-3 (1.8-fold) and to the normal lung cell line Hs888.Lu (1.8-fold). Under these conditions, no
20 binding was observed to the Lovo or SW1116 colon cancer cell lines or to the normal skin cell line CCD-27sk. From Examples 1 and 2, it is evident that the binding of AR27A807.14.1 to the Lovo and SW1116 colon cancer cell lines is not detectable using cell ELISA. Nevertheless, the antibody was able to induce a cytotoxic effect in these cell lines. The effect may be due to the following factor. It is possible that the Lovo/ SW1116 colon cancer cell line expresses the
25 antigen target at a level below the threshold of detection by this method under these conditions, but that the low level of expression is sufficient to trigger an event leading to delayed tumor growth. The cytotoxic effect of this antibody in a Lovo and SW1116 colon cancer model is a non-obvious finding that could not have been predicted on the basis of binding. From Example 1, AR27A807.14.1 binding to the colon cell line HT-29 was not detectable by cell ELISA whereas

in Example 2 there is clear binding; this may be due to a lower concentration of antibody in the supernatant versus what was tested in the purified form.

EXAMPLE 3

In vivo Tumor Experiments with DLD-1 Cells

5 Example 1 demonstrated that AR27A807.14.1 had anti-cancer properties against several colon cancer cell lines. With reference to Figures 3 and 4, 4 to 6 week old female SCID mice were implanted with 5 million human colon cancer cells (DLD-1) in 100 microlitres saline injected subcutaneously in the scruff of the neck. The mice were randomly divided into 2 treatment groups of 5. On the day after implantation, 20 mg/kg of AR27A807.14.1 test antibody or buffer control was administered intraperitoneally to each cohort in a volume of 300 microlitres after dilution from the stock concentration with a diluent that contained 2.7 mM KCl, 1 mM KH₂PO₄, 137 mM NaCl and 20 mM Na₂HPO₄. The antibody and control samples were then administered once per week for the duration of the study in the same fashion. Tumor growth was measured about every seventh day with calipers. The study was completed after 6 injections of antibody. Body weights of the animals were recorded once per week for the duration of the study. At the end of the study all animals were euthanized according to CCAC guidelines.

15 AR27A807.14.1 reduced tumor growth in the highly aggressive DLD-1 *in vivo* prophylactic model of human colon cancer. Figure 3 presents the results of the tumor volume for mice treated with the antibody AR27A807.14.1, compared to the buffer treated group. At the end of the study, on day 37, AR27A807.14.1 inhibited tumor growth by 65 percent (p=0.0039) compared to the buffer-treated control. From Example 2, it is evident that the binding of AR27A807.14.1 to the DLD-1 cell line is not detectable using cell ELISA. Nevertheless, the antibody was able to reduce the tumor growth in a DLD-1 xenograft model. It is possible that the DLD-1 cell line expresses the antigen target at a level below the threshold of detection by cell ELISA under these conditions, but that the low level of expression is sufficient to trigger an event leading to delayed tumor growth. It is also possible that antigen expression is induced when the DLD-1 cells are placed into the more physiological *in vivo* environment. In either case, the efficacy of this antibody in a DLD-1 colon cancer model is a non-obvious finding that could not have been predicted on the basis of binding.

There were no clinical signs of toxicity throughout the study. Body weight measured at weekly intervals was a surrogate for well-being and failure to thrive. As shown in Figure 4, there was a significant reduction in the mean body weight of both the control (p=0.0033) and the AR27A807.14.1-treated groups (p=0.0421) by day 37. However, there was no significant difference in body weight between the control and the antibody treated groups at the end of the study.

In summary, AR27A807.14.1 was well-tolerated and decreased the tumor burden in this human colon cancer xenograft model.

EXAMPLE 4

Isolation of Competitive Binders

Given an antibody, an individual ordinarily skilled in the art can generate a competitively inhibiting CDMAB, for example a competing antibody, which is one that recognizes the same epitope (Belanger L *et al. Clinica Chimica Acta* 48:15-18 (1973)). One method entails immunizing with an immunogen that expresses the antigen recognized by the antibody. The sample may include but is not limited to tissues, isolated protein(s) or cell line(s). Resulting hybridomas could be screened using a competition assay, which is one that identifies antibodies that inhibit the binding of the test antibody, such as ELISA, FACS or Western blotting. Another method could make use of phage display antibody libraries and panning for antibodies that recognize at least one epitope of said antigen (Rubinstein JL *et al. Anal Biochem* 314:294-300 (2003)). In either case, antibodies are selected based on their ability to displace the binding of the original labeled antibody to at least one epitope of its target antigen. Such antibodies would therefore possess the characteristic of recognizing at least one epitope of the antigen as the original antibody.

EXAMPLE 5

Cloning of the Variable Regions of the AR27A807.14.1 Monoclonal Antibody

The sequences of the variable regions from the heavy (V.H) and light (V.L) chains of monoclonal antibody produced by the AR27A807.14.1 hybridoma cell line can be determined. RNA encoding the heavy and light chains of immunoglobulin can be extracted from the subject hybridoma using standard methods involving cellular solubilization with guanidinium

isothiocyanate (Chirgwin *et al.* Biochem. 18:5294-5299 (1979)). The mRNA can be used to prepare cDNA for subsequent isolation of V.H and V.L genes by PCR methodology known in the art (Sambrook *et al.*, eds., Molecular Cloning, Chapter 14, Cold Spring Harbor laboratories Press, N.Y. (1989)). The N-terminal amino acid sequence of the heavy and light chains can be
5 independently determined by automated Edman sequencing. Further stretches of the CDRs and flanking FRs can also be determined by amino acid sequencing of the V.H and V.L fragments. Synthetic primers can be then designed for isolation of the V.H and V.L genes from AR27A807.14.1 monoclonal antibody, and the isolated gene can be ligated into an appropriate vector for sequencing. To generate chimeric and humanized IgG, the variable light and variable
10 heavy domains can be subcloned into an appropriate vector for expression.

(i) Monoclonal Antibody

DNA encoding the monoclonal antibody (as outlined in Example 1) is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of the
15 monoclonal antibodies). The hybridoma cell serves as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for
20 example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences. Chimeric or hybrid antibodies also may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include
25 iminothiolate and methyl-4-mercaptobutyrimidate.

(ii) Humanized Antibody

A humanized antibody has one or more amino acid residues introduced into it from a non-human source. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization
30 can be performed the method of Winter and co-workers by substituting rodent CDRs or CDR

sequences for the corresponding sequences of a human antibody (Jones *et al.*, Nature 321:522-525 (1986); Riechmann *et al.*, Nature 332:323-327 (1988); Verhoeyen *et al.*, Science 239:1534-1536 (1988); reviewed in Clark, Immunol. Today 21:397-402 (2000)).

5 A humanized antibody can be prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the
10 residues in the functioning of the candidate immunoglobulin sequence, i.e. the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the consensus and import sequence so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in
15 influencing antigen binding.

(iii) Antibody Fragments

Various techniques have been developed for the production of antibody fragments. These fragments can be produced by recombinant host cells (reviewed in Hudson, Curr. Opin. Immunol. 11:548-557 (1999); Little *et al.*, Immunol. Today 21:364-370 (2000)). For
20 example, Fab'-SH fragments can be directly recovered from E. coli and chemically coupled to form F(ab')₂ fragments (Carter *et al.*, Biotechnology 10:163-167 (1992)). In another embodiment, the F(ab')₂ is formed using the leucine zipper GCN4 to promote assembly of the F(ab')₂ molecule. According to another approach, Fv, Fab or F(ab')₂ fragments can be isolated directly from recombinant host cell culture.

25 EXAMPLE 6

A Composition Comprising the Antibody of the Present Invention

The antibody of the present invention can be used as a composition for preventing/treating cancer. The composition for preventing/treating cancer, which comprises the antibody of the present invention, are low-toxic and can be administered as they are in the form
30 of liquid preparations, or as pharmaceutical compositions of suitable preparations to human or

mammals (e.g., rats, rabbits, sheep, swine, bovine, feline, canine, simian, etc.) orally or parenterally (e.g., intravascularly, intraperitoneally, subcutaneously, etc.). The antibody of the present invention may be administered in itself, or may be administered as an appropriate composition. The composition used for the administration may contain a pharmacologically acceptable carrier with the antibody of the present invention or its salt, a diluent or excipient. Such a composition is provided in the form of pharmaceutical preparations suitable for oral or parenteral administration.

Examples of the composition for parenteral administration are injectable preparations, suppositories, etc. The injectable preparations may include dosage forms such as intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, intraarticular injections, etc. These injectable preparations may be prepared by methods publicly known. For example, the injectable preparations may be prepared by dissolving, suspending or emulsifying the antibody of the present invention or its salt in a sterile aqueous medium or an oily medium conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mols) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared is usually filled in an appropriate ampoule. The suppository used for rectal administration may be prepared by blending the antibody of the present invention or its salt with conventional bases for suppositories. The composition for oral administration includes solid or liquid preparations, specifically, tablets (including dragees and film-coated tablets), pills, granules, powdery preparations, capsules (including soft capsules), syrup, emulsions, suspensions, etc. Such a composition is manufactured by publicly known methods and may contain a vehicle, a diluent or excipient conventionally used in the field of pharmaceutical preparations. Examples of the vehicle or excipient for tablets are lactose, starch, sucrose, magnesium stearate, etc.

Advantageously, the compositions for oral or parenteral use described above are prepared into pharmaceutical preparations with a unit dose suited to fit a dose of the active ingredients. Such unit dose preparations include, for example, tablets, pills, capsules, injections (ampoules), suppositories, etc. The amount of the aforesaid compound contained is generally 5 to 500 mg per dosage unit form; it is preferred that the antibody described above is contained in about 5 to about 100 mg especially in the form of injection, and in 10 to 250 mg for the other forms.

The dose of the aforesaid prophylactic/therapeutic agent or regulator comprising the antibody of the present invention may vary depending upon subject to be administered, target disease, conditions, route of administration, etc. For example, when used for the purpose of treating/preventing, e.g., breast cancer in an adult, it is advantageous to administer the antibody of the present invention intravenously in a dose of about 0.01 to about 20 mg/kg body weight, preferably about 0.1 to about 10 mg/kg body weight and more preferably about 0.1 to about 5 mg/kg body weight, about 1 to 5 times/day, preferably about 1 to 3 times/day. In other parenteral and oral administration, the agent can be administered in a dose corresponding to the dose given above. When the condition is especially severe, the dose may be increased according to the condition.

The antibody of the present invention may be administered as it stands or in the form of an appropriate composition. The composition used for the administration may contain a pharmacologically acceptable carrier with the aforesaid antibody or its salts, a diluent or excipient. Such a composition is provided in the form of pharmaceutical preparations suitable for oral or parenteral administration (e.g., intravascular injection, subcutaneous injection, etc.). Each composition described above may further contain other active ingredients. Furthermore, the antibody of the present invention may be used in combination with other drugs, for example, alkylating agents (e.g., cyclophosphamide, ifosfamide, etc.), metabolic antagonists (e.g., methotrexate, 5-fluorouracil, etc.), antitumor antibiotics (e.g., mitomycin, adriamycin, etc.), plant-derived antitumor agents (e.g., vincristine, vindesine, Taxol, etc.), cisplatin, carboplatin, etoposide, irinotecan, etc. The antibody of the present invention and the drugs described above may be administered simultaneously or at staggered times to the patient.

The preponderance of evidence shows that AR27807.14.1 mediates anti-cancer effects through ligation of an epitope present on cancer cell lines. Further it could be shown that the AR27A807.14.1 antibody could be used in detection of cells which express the epitope which specifically binds thereto; utilizing techniques illustrated by, but not limited to FACS, cell

5 ELISA or IHC.

All patents and publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

10 It is to be understood that while a certain form of the invention is illustrated, it is not to be limited to the specific form or arrangement of parts herein described and shown. It will be apparent to those skilled in the art that various changes may be made without departing from the scope of the invention and the invention is not to be considered limited to what is shown and described in the specification.

15 One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. Any oligonucleotides, peptides, polypeptides, biologically related compounds, methods, procedures and techniques described herein are presently representative of the preferred embodiments, are intended to be exemplary and are not intended as limitations on the
20 scope. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention and are defined by the scope of the appended claims. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying
25 out the invention which are obvious to those skilled in the art are intended to be within the scope of the following claims.

CLAIMS

What is claimed is:

Claim 1. The isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03.

Claim 2. A humanized antibody of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or an antigen binding fragment produced from said humanized antibody.

Claim 3. A chimeric antibody of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or an antigen binding fragment produced from said chimeric antibody.

Claim 4. The isolated hybridoma cell line deposited with the IDAC as accession number 180706-03.

Claim 5. A method for initiating antibody induced cytotoxicity of cancerous cells in a tissue sample selected from a human tumor comprising:

providing a tissue sample from said human tumor;

providing the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03, the humanized antibody of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03, the chimeric antibody of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or a CDMAB thereof, which CDMAB is characterized by an ability to competitively inhibit binding of said isolated monoclonal antibody to its target antigen; and

contacting said isolated monoclonal antibody, said humanized antibody, said chimeric antibody or CDMAB thereof with said tissue sample;

wherein binding of said isolated monoclonal antibody, said humanized antibody, said chimeric antibody or CDMAB thereof with said tissue sample induces cytotoxicity.

Claim 6. A CDMAB of the isolated monoclonal antibody of claim 1.

- Claim 7. A CDMAB of the humanized antibody of claim 2.
- Claim 8. A CDMAB of the chimeric antibody of claim 3.
- Claim 9. The isolated antibody or CDMAB thereof, of any one of claims 1, 2, 3, 6, 7 or 8 conjugated with a member selected from the group consisting of cytotoxic moieties, enzymes, radioactive compounds, and hematogenous cells.
- Claim 10. A method of treating a human tumor susceptible to antibody induced cytotoxicity in a mammal, wherein said human tumor expresses at least one epitope of an antigen which specifically binds to the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or a CDMAB thereof, which CDMAB is characterized by an ability to competitively inhibit binding of said isolated monoclonal antibody to its target antigen, comprising administering to said mammal said monoclonal antibody or said CDMAB thereof in an amount effective to result in a reduction of said mammal's tumor burden.
- Claim 11. The method of claim 10 wherein said isolated monoclonal antibody is conjugated to a cytotoxic moiety.
- Claim 12. The method of claim 11 wherein said cytotoxic moiety is a radioactive isotope.
- Claim 13. The method of claim 10 wherein said isolated monoclonal antibody or CDMAB thereof activates complement.
- Claim 14. The method of claim 10 wherein said isolated monoclonal antibody or CDMAB thereof mediates antibody dependent cellular cytotoxicity.
- Claim 15. The method of claim 10 wherein said isolated monoclonal antibody is humanized.
- Claim 16. The method of claim 10 wherein said isolated monoclonal antibody is chimerized.
- Claim 17. A monoclonal antibody capable of specific binding to the same epitope or epitopes as the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03.

Claim 18. A method of treating a human tumor in a mammal, wherein said human tumor expresses at least one epitope of an antigen which specifically binds to the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or a CDMAB thereof, which CDMAB is characterized by an ability to competitively inhibit binding of said isolated monoclonal antibody to its target antigen, comprising administering to said mammal said monoclonal antibody or CDMAB thereof in an amount effective to result in a reduction of said mammal's tumor burden.

Claim 19. The method of claim 18 wherein said isolated monoclonal antibody is conjugated to a cytotoxic moiety.

Claim 20. The method of claim 19 wherein said cytotoxic moiety is a radioactive isotope.

Claim 21. The method of claim 18 wherein said isolated monoclonal antibody or CDMAB thereof activates complement.

Claim 22. The method of claim 18 wherein said isolated monoclonal antibody or CDMAB thereof mediates antibody dependent cellular cytotoxicity.

Claim 23. The method of claim 18 wherein said isolated monoclonal antibody is humanized.

Claim 24. The method of claim 18 wherein said isolated monoclonal antibody is chimerized.

Claim 25. A method of treating a human tumor in a mammal, wherein said human tumor expresses at least one epitope of an antigen which specifically binds to the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or a CDMAB thereof, which CDMAB is characterized by an ability to competitively inhibit binding of said isolated monoclonal antibody to its target antigen, comprising administering to said mammal said monoclonal antibody or CDMAB thereof in conjunction with at least one chemotherapeutic agent in an amount effective to result in a reduction of said mammal's tumor burden.

Claim 26. The method of claim 25 wherein said isolated monoclonal antibody is conjugated to a cytotoxic moiety.

Claim 27. The method of claim 26 wherein said cytotoxic moiety is a radioactive isotope.

Claim 28. The method of claim 25 wherein said isolated monoclonal antibody or CDMAB thereof activates complement.

Claim 29. The method of claim 25 wherein said isolated monoclonal antibody or CDMAB thereof mediates antibody dependent cellular cytotoxicity.

Claim 30. The method of claim 25 wherein said isolated monoclonal antibody is humanized.

Claim 31. The method of claim 25 wherein said isolated monoclonal antibody is chimerized.

Claim 32. A binding assay to determine a presence of cancerous cells in a tissue sample selected from a human tumor, which is specifically bound by the isolated monoclonal antibody produced by hybridoma cell line AR27A807.14.1 having IDAC Accession No. 180706-03, the humanized antibody of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or the chimeric antibody of the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03, comprising:

providing a tissue sample from said human tumor;

providing at least one of said isolated monoclonal antibody, said humanized antibody, said chimeric antibody or CDMAB thereof that recognizes the same epitope or epitopes as those recognized by the isolated monoclonal antibody produced by a hybridoma cell line AR27A807.14.1 having IDAC Accession No. 180706-03;

contacting at least one said provided antibodies or CDMAB thereof with said tissue sample; and

determining binding of said at least one provided antibody or CDMAB thereof with said tissue sample;

whereby the presence of said cancerous cells in said tissue sample is indicated.

Claim 33. Use of monoclonal antibodies for reduction of human tumor burden, wherein said human tumor expresses at least one epitope of an antigen which specifically binds to the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number

180706-03 or a CDMAB thereof, which CDMAB is characterized by an ability to competitively inhibit binding of said isolated monoclonal antibody to its target antigen, comprising administering to said mammal said monoclonal antibody or CDMAB thereof in an amount effective to result in a reduction of said mammal's human tumor burden.

Claim 34. The method of claim 33 wherein said isolated monoclonal antibody is conjugated to a cytotoxic moiety.

Claim 35. The method of claim 34 wherein said cytotoxic moiety is a radioactive isotope.

Claim 36. The method of claim 33 wherein said isolated monoclonal antibody or CDMAB thereof activates complement.

Claim 37. The method of claim 33 wherein said isolated monoclonal antibody or CDMAB thereof mediates antibody dependent cellular cytotoxicity.

Claim 38. The method of claim 33 wherein said isolated monoclonal antibody is humanized.

Claim 39. The method of claim 33 wherein said isolated monoclonal antibody is chimerized.

Claim 40. Use of monoclonal antibodies for reduction of human tumor burden, wherein said human tumor expresses at least one epitope of an antigen which specifically binds to the isolated monoclonal antibody produced by the hybridoma deposited with the IDAC as accession number 180706-03 or a CDMAB thereof, which CDMAB is characterized by an ability to competitively inhibit binding of said isolated monoclonal antibody to its target antigen, comprising administering to said mammal said monoclonal antibody or CDMAB thereof; in conjunction with at least one chemotherapeutic agent in an amount effective to result in a reduction of said mammal's human tumor burden.

Claim 41. The method of claim 40 wherein said isolated monoclonal antibody is conjugated to a cytotoxic moiety.

Claim 42. The method of claim 41 wherein said cytotoxic moiety is a radioactive isotope.

Claim 43. The method of claim 40 wherein said isolated monoclonal antibody or CDMAB thereof activates complement.

Claim 44. The method of claim 40 wherein said isolated monoclonal antibody or CDMAB thereof mediates antibody dependent cellular cytotoxicity.

Claim 45. The method of claim 40 wherein said isolated monoclonal antibody is humanized.

Claim 46. The method of claim 40 wherein said isolated monoclonal antibody is chimerized.

Claim 47. A composition effective for treating a human cancerous tumor comprising in combination:

an antibody or CDMAB of any one of claims 1,2,3,6,7,8, or 17;

a conjugate of said antibody or an antigen binding fragment thereof with a member selected from the group consisting of cytotoxic moieties, enzymes, radioactive compounds, and hematogenous cells; and

a requisite amount of a pharmaceutically acceptable carrier;

wherein said composition is effective for treating said human cancerous tumor.