CANNABINOID RECEPTOR MODULATORS

FIELD OF THE INVENTION

The present invention relates to cannabinoid receptor modulators, particularly, antagonists or inverse agonists of the CB_1 receptor, useful for the treatment of obesity, metabolic disorders, addiction, diseases of the central nervous system, cardiovascular disorders, respiratory disorders, and gastrointestinal disorders, pharmaceutical compositions comprising such compounds, and methods of treatment using the compounds and compositions to treat conditions such as obesity, metabolic disorders, addiction, diseases of the central nervous system, cardiovascular disorders, respiratory disorders, and gastrointestinal disorders.

BACKGROUND OF THE INVENTION


The CB_1 receptor is one of the most abundant neuromodulatory receptors in the brain, and is expressed at high levels in the hippocampus, cortex, cerebellum, and basal ganglia (e.g., Wilson et al., *Science*, **2002**, vol. 296, 678-682). Selective CB_1 receptor antagonists, for example pyrazole derivatives such as rimonabant (e.g., U.S. 6,432,984), can be used to treat various conditions, such as obesity and metabolic syndrome (e.g., Bensaid et al., *Molecular Pharmacology*, **2003** vol. 63, no. 4, pp. 908-914; Trillou et al., *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2002** vol. 284, R345-R353; Kirkham, *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2002** vol. 284, R343-R344), neuroinflammatory disorders (e.g., Adam, et al., *Expert Opin. Ther. Patents*, **2002**, vol. 12, no. 10, 1475-1489; U.S. 6,642,258), cognitive disorders and psychosis (e.g., Adam et al., *Expert Opin. Ther. Pat.*, **2002**, vol. 12, pp. 1475-1489), addiction (e.g., smoking cessation; U.S. Patent Publ. 2003/0087933), gastrointestinal disorders (e.g., Lange et al., *J. Med. Chem.* **2004**, vol. 47, 627-643) and cardiovascular conditions (e.g., Porter et al.,

Pharmacology and Therapeutics, **2001** vol. 90, 45-60; Sanofi-Aventis Publication, Bear Steams Conference, New York, September 14, 2004, pages 19-24).

However, there is still a need for improved cannabinoid agents, particularly cannabinoid receptor modulators (e.g., antagonists or inverse agonists of the CB_1 receptor) with fewer side-effects and improved efficacy. It is therefore an object of the present invention to provide fused bicyclic and spirocyclic cannabinoid receptor modulators useful in the treatment of diseases or conditions mediated by cannabinoid receptors.

SUMMARY OF THE INVENTION

In one embodiment, the present invention is directed to a compound of Formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, or ester thereof, wherein:


m is 0 or 1, n is 1 or 2, and m + n is 1 or 2;

the dashed lines (====) in Formula (I) represent single or double bonds as permitted by valence requirements;

R^1 is selected from the group consisting of $-C(O)-N(R^{10})_2$, -C(O)-O-alkyl, and $-C(O)-R^{14}$;


R^2 is selected from the group consisting of H, unsubstituted alkyl, alkyl substituted with one or more U groups, and -alkylene-$N(R^{10})_2$;

or R^1 and R^2 together with the carbon atoms to which they are shown attached in Formula (I) form a group Q as shown in Formula (IA):




(IA)

wherein Q is selected from the group consisting of:



(a) , (b) , (c) , (d) , (e) , (f) , and



(f) ;

Y^1 is –O- or –N(R^7)-;

Y^2 is –O- or –N(R^8)-;

R^3, R^4, R^5, and R^6 are each independently selected from the group consisting of H, -O-R^9, R^{11}, and –N(R^{16})$_2$;

R^7 is selected from the group consisting of H, alkyl, arylalkyl, alkenyl, -alkylene-N(R^9)$_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, -alkylene-C(O)H, -C(O)-O-R^{11}, and Boc;

R^8 is selected from the group consisting of H, -alkylene-R^{12}, -C(O)-R^{17}, -S(O_2)-R^{11}, -S(O_2)-R^{14}, -C(O)-N(R^{18})$_2$, R^{14}, and Boc;

with the proviso wherein the group -N(R^{18})$_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

R^9 is selected from the group consisting of H, TBS, TIPS, Tf and R^{11};

each R^{10} is independently selected from the group consisting of H, unsubstituted alkyl, alkyl substituted with one or more U groups, -alkylene-R^{12}, -alkylene-R^{13}, -alkylene-R^{14}, -C(O)-R^{14}, -alkylene-O-R^9, R^{14}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, and benzo-fused cycloalkyl;

R^{11} is selected from the group consisting of unsubstituted alkyl, alkyl substituted with one or more U groups, -alkylene-O-alkyl,

-alkylene-O-aryl, unsubstituted aryl, and aryl substituted with one or more X^1 groups;

R^{12} is selected from the group consisting of unsubstituted aryl and aryl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more X^2 groups;

R^{14} is selected from the group consisting of unsubstituted cycloalkyl, cycloalkyl substituted with one or more X^4 groups unsubstituted alkyl, and alkyl substituted with one or more U groups;

each R^{15} is independently selected from the group consisting of H, $-N_3$, halogen, alkenyl, -alkylene-R^{12}, -alkylene-O-R^9, -alkylene-$N(R^{18})_2$, -alkylene-C(O)H, -OH, -CN, -O-alkyl, $-C(O)N(R^{18})_2$, $-N(R^{18})_2$, $-NR^{18}C(O)R^{18}$, $-NR^{18}C(O)_2R^{18}$, $-NR^{18}C(O)N(R^{18})_2$, $-NR^{18}S(O)_2R^{18}$, –O-alkenyl, $-C(O)_2R^{18}$; unsubstituted alkyl, alkyl substituted with one or more U groups, -O-alkylene-$C(O)R^{18}$ or $-C(O)R^{18}$;

with the proviso wherein the group $-N(R^{18})_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

R^{16} is selected from the group consisting of R^9 and $-C(O)-R^{12}$;

R^{17} is selected from the group consisting of unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, -alkylene-R^{12}, -O-R^9, and R^{12};

each R^{18} is independently selected from the group consisting of H, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, R^{12}, R^{13} and R^{14};

with the proviso that when R^{18} is attached to N, then each R^{18} is independently selected from the group consisting of H, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, $-C(O)R^{21}$, R^{12}, R^{13} and R^{14};

R^{19} is selected from the group consisting of H, TBS, TIPS, Tf and R^{21};

each R^{20} is independently selected from the group consisting of H, unsubstituted alkyl, alkyl substituted with one or more U groups,

-alkylene-R^{22}, -alkylene-R^{23}, -alkylene-R^{24}, -C(O)-R^{24}, -alkylene-O-R^{19}, R^{24}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, and benzo-fused cycloalkyl;

R^{21} is selected from the group consisting of unsubstituted alkyl, alkyl substituted with one or more U groups, -alkylene-O-alkyl, -alkylene-O-aryl, unsubstituted aryl, aryl substituted with one or more W^1 groups; unsubstituted heteroaryl, heteroaryl substituted with one or more W^2 groups, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, -O-alkylene-O-R^{24}, -C(O)-O-alkylene-O-R^{24}; -C(O)-alkylene-R^{23}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -$S(O)_2$-R^{24}, and -alkylene-O-alkylene-O-R^{24}, with the proviso that –O-O- cannot be formed with R^{21} and the atom said R^{21} is attached to;

R^{22} is selected from the group consisting of unsubstituted aryl and aryl substituted with one or more W^1 groups;

R^{23} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more W^2 groups;

R^{24} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, unsubstituted alkyl, and alkyl substituted with one or more U groups;

each R^{25} is independently selected from the group consisting of H, R^{22}, R^{23}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -alkylene-OR^{19}, -alkylene-N$R^{19}R^{19}$, -alkylene-SR^{19},-alkylene-R^{23}, -alkylene-R^{22}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, -alkylene-heterocycloalkyl, -alkylene-heterocycloalkyl substituted with one or more W^3 groups, -C(O)-R^{24}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -$S(O)_2$-R^{24}, -C(O)-NH-R^{22} and -C(O)-NH-R^{24};

with the proviso wherein the group -$N(R^{25})_2$, both R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups,

or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

each W^1 is independently selected from the group consisting of halogen, -CN, -OH, -O-$S(O)_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, and –O-alkyl;

each W^2 is independently selected from the group consisting of halogen, unsubstituted aryl, and aryl substituted with one or more Z groups;

each W^3 is independently selected from the group consisting of –OH, alkyl, -alkylene-OH, -O-alkyl, -C(O)-alkyl, -$C(O)NH_2$, -NHC(O)alkyl, -NHC(O)H, -NHC(O)-O-alkyl and -C(O)-O-alkyl; or

two W^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each W^4 is independently halogen or alkyl;

Ar^1 and Ar^2 are independently selected from the group consisting of R^{12} and R^{13};

each X^1 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-$S(O)_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-C(O)N(R^{20})$_2$, -O-alkylene-O-R^{19}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl, -O-alkylene-O-alkylene-OR^{19}, -O-alkylene-C(O)R^{24}, -O-alkylene-C(O)OR^{19},–O-alkyl, -N(R^{25})$_2$, -C(O)alkyl, –C(O)OH, -C(O)O-alkyl, -C(O)O-cycloalkyl, -C(O)N(R^{25})$_2$, -O-alkylene-heterocycloalkyl, -O-alkylene-heterocycloalkyl substituted with one or more W^3 groups, unsubstituted heterocycloalkyl, -heterocycloalkyl substituted with one or more W^3 groups, -O-alkenylene-O-alkylene-O-R^{24}, -O-alkylene-N(R^{25})$_2$, -O-alkylene-C(O)N(R^{25})$_2$, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -$S(O)_2$-R^{24}, and alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an

unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

each X^2 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-C(O)N(R^{20})$_2$, -O-alkylene-O-R^{19}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl,-O-alkylene-O-alkylene-OR^{19}, -O-alkylene-C(O)R^{24}, -O-alkylene-C(O)OR^{19},–O-alkyl, -N(R^{25})$_2$, -C(O)alkyl, –C(O)OH, -C(O)O-alkyl, -C(O)O-cycloalkyl, -C(O)N(R^{25})$_2$, -O-alkylene-heterocycloalkyl, -O-alkylene-heterocycloalkyl substituted with one or more W^3 groups, unsubstituted heterocycloalkyl, -heterocycloalkyl substituted with one or more W^3 groups, -O-alkenylene-O-alkylene-O-R^{24}, -O-alkylene-N(R^{25})$_2$, -O-alkylene-C(O)N(R^{25})$_2$, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S(O)$_2$-R^{24}, and alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

each X^3 is independently selected from the group consisting of –OH, alkyl, -alkylene-OH, -O-alkyl, -C(O)-alkyl, -C(O)NH_2, -NHC(O)alkyl, -NHC(O)H, -NHC(O)-O-alkyl and -C(O)-O-alkyl; or

two X^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each X^4 is independently halogen or alkyl;

each U is independently selected from the group consisting of –OH, -O-alkyl, -O-aryl, -O-alkylene-aryl, -O-alkylene-O-alkyl, -O-alkylene-O-haloalkyl, -O-alkylene-O-aryl, halogen, -CN, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, OTBS, OTIPS and OTf;

and

each Z is independently selected from the group consisting of –OH; -O-alkyl; halogen; alkyl; –CN; -CF_3; cycloalkyl; -alkylene-OH; -alkylene-O-alkyl; -alkylene-O-alkyl substituted with one or more groups selected from the group consisting of –OH, -O-alkyl, halogen, -CN, cycloalkyl, heterocycloalkyl, aryl, heteroaryl; -alkylene-O-alkylene-O-alkyl; -alkylene-O-alkylene-O-aryl; -alkylene-O-aryl; and -alkylene-O-aryl substituted with one or more groups selected from the group consisting of halogen, -CN, -OH, -O-S(O)$_2$-haloalkyl, aryl, heteroaryl and –O-alkyl; or

two Z groups together with the ring carbon atom to which they are attached form a carbonyl group.

In another embodiment, the present invention is directed to a pharmaceutical composition comprising a therapeutically effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable salt, solvate, or ester thereof, and at least one pharmaceutically acceptable carrier.


In yet another embodiment, the present invention also provides for a method of treating, reducing, or ameliorating metabolic syndrome, obesity, waist circumference, lipid profile, insulin sensitivity, neuroinflammatory disorders, cognitive disorders, psychosis, addictive behavior, gastrointestinal disorders, and cardiovascular conditions by administering an effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable salt, solvate, ester, or stereoisomer thereof, to a patient in need thereof.

In another embodiment, the present invention is directed to a method of treating, reducing, or ameliorating a disease or disorder in a patient, such as metabolic syndrome, obesity, waist circumference, lipid profile, insulin sensitivity, neuroinflammatory disorders, cognitive disorders, psychosis, addictive behavior, gastrointestinal disorders, and cardiovascular conditions. The method comprises administering to the patient an effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable salt, solvate, or ester thereof, in combination with one or more cholesterol lowering agents.

DETAILED DESCRIPTION OF THE INVENTION

In one embodiment, the present invention is directed to a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, ester, or stereoisomer thereof, as described herein.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, and stereoisomers thereof have the following Formula (I):



wherein


m is 0 or 1, n is 1 or 2, and m + n is 1 or 2;

the dashed lines (----) in Formula (I) represent single or double bonds as permitted by valence requirements;

R^1 is selected from the group consisting of $-C(O)-N(R^{10})_2$, $-C(O)-O-(C_1-C_6)$alkyl, and $-C(O)-R^{14}$;


R^2 is selected from the group consisting of H, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, and $-(C_1-C_6)$alkylene-$N(R^{10})_2$;

or R^1 and R^2 together with the carbon atoms to which they are shown attached in Formula (I) form a group Q as shown in Formula (IA):




(IA)

wherein Q is selected from the group consisting of:



(a) , (b) , (c) , (d) , (e) , (f) , and



(f) ;

Y^1 is –O- or –N(R^7)-;

Y^2 is –O- or –N(R^8)-;

R^3, R^4, R^5, and R^6 are each independently selected from the group consisting of H, -O-R^9, R^{11}, and –N(R^{16})$_2$;

R^7 is selected from the group consisting of H, (C_1-C_6)alkyl, (C_6-C_{12})aryl(C_1-C_6)alkyl, (C_2-C_6)alkenyl, -(C_1-C_6)alkylene-N(R^9)$_2$, -(C_1-C_6)alkylene-O-R^9, -(C_1-C_6)alkylene-R^{12}, -C(O)-R^{14}, -(C_1-C_6)alkylene-C(O)H, -C(O)-O-R^{11}, and Boc;

R^8 is selected from the group consisting of H, -(C_1-C_6)alkylene-R^{12}, -C(O)-R^{17}, -S(O_2)-R^{11}, -S(O_2)-R^{14}, -C(O)-N(R^{18})$_2$, R^{14}, and Boc;

with the proviso wherein the group -N(R^{18})$_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted (C_3-C_5)heterocycloalkyl group is fused with (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, (C_3-C_7)cycloalkyl or (C_3-C_5)heterocycloalkyl;

R^9 is selected from the group consisting of H, TBS, TIPS, Tf and R^{11};

each R^{10} is independently selected from the group consisting of H, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, -(C_1-C_6)alkylene-R^{12}, -(C_1-C_6)alkylene-R^{13}, -(C_1-C_6)alkylene-R^{14}, -C(O)-R^{14}, -(C_1-C_6)alkylene-O-R^9, R^{14}, unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, and benzo-fused (C_3-C_7)cycloalkyl;

R^{11} is selected from the group consisting of unsubstituted $(C_1\text{-}C_6)$alkyl, $(C_1\text{-}C_6)$alkyl substituted with one or more U groups, $-(C_1\text{-}C_6)$alkylene-O-$(C_1\text{-}C_6)$alkyl, $-(C_1\text{-}C_6)$alkylene-O-$(C_6\text{-}C_{12})$aryl, unsubstituted $(C_6\text{-}C_{12})$aryl, and $(C_6\text{-}C_{12})$aryl substituted with one or more X^1 groups;

R^{12} is selected from the group consisting of unsubstituted $(C_6\text{-}C_{12})$aryl and $(C_6\text{-}C_{12})$aryl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted $(C_2\text{-}C_{10})$heteroaryl and $(C_2\text{-}C_{10})$heteroaryl substituted with one or more X^2 groups;

R^{14} is selected from the group consisting of unsubstituted $(C_3\text{-}C_7)$cycloalkyl, $(C_3\text{-}C_7)$cycloalkyl substituted with one or more X^4 groups unsubstituted $(C_1\text{-}C_6)$alkyl and $(C_1\text{-}C_6)$alkyl substituted with one or more U groups;

each R^{15} is independently selected from the group consisting of H, $-N_3$, halogen, $(C_2\text{-}C_6)$alkenyl, $-(C_1\text{-}C_6)$alkylene-R^{12}, $-(C_1\text{-}C_6)$alkylene-O-R^9, $-(C_1\text{-}C_6)$alkylene-$N(R^{18})_2$, $-(C_1\text{-}C_6)$alkylene-C(O)H, -OH, -CN, $-O-(C_1\text{-}C_6)$alkyl, $-C(O)N(R^{18})_2$, $-N(R^{18})_2$, $-NR^{18}C(O)R^{18}$, $-NR^{18}C(O)_2R^{18}$, $-NR^{18}C(O)N(R^{18})_2$, $-NR^{18}S(O)_2R^{18}$, $-O-(C_2\text{-}C_6)$alkenyl, $-C(O)_2R^{18}$, unsubstituted $(C_1\text{-}C_6)$alkyl, $(C_1\text{-}C_6)$alkyl substituted with one or more U groups, $-O-(C_1\text{-}C_6)$alkylene-$C(O)R^{18}$, or $-C(O)R^{18}$;

with the proviso wherein the group $-N(R^{18})_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl group is fused with $(C_6\text{-}C_{12})$aryl, $(C_2\text{-}C_{10})$heteroaryl, $(C_3\text{-}C_7)$cycloalkyl or $(C_3\text{-}C_5)$heterocycloalkyl;

R^{16} is selected from the group consisting of R^9 and $-C(O)\text{-}R^{12}$;

R^{17} is selected from the group consisting of unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more X^3 groups, $-(C_1\text{-}C_6)$alkylene-R^{12}, -O-R^9, and R^{12};

each R^{18} is independently selected from the group consisting of H, unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more X^3 groups, R^{12}, R^{13} and R^{14};

with the proviso that when R^{18} is attached to N, then each R^{18} is independently selected from the group consisting of H, unsubstituted

(C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -C(O)R^{21}, R^{12}, R^{13} and R^{14};

R^{19} is selected from the group consisting of H, TBS, TIPS, Tf and R^{21};

each R^{20} is independently selected from the group consisting of H, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, -(C_1-C_6)alkylene-R^{22}, -(C_1-C_6)alkylene-R^{23}, -(C_1-C_6)alkylene-R^{24}, -C(O)-R^{24}, -(C_1-C_6)alkylene-O-R^{19}, R^{24}, unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, and benzo-fused (C_3-C_7)cycloalkyl;

R^{21} is selected from the group consisting of unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, -(C_1-C_6)alkylene-O-(C_1-C_6)alkyl, -(C_1-C_6)alkylene-O-(C_6-C_{12})aryl, unsubstituted (C_6-C_{12})aryl, and (C_6-C_{12})aryl substituted with one or more W^1 groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more W^2 groups, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl with one or more W^3 groups, -O-(C_1-C_6)alkylene-O-R^{24}, -C(O)-O-(C_1-C_6)alkylene-O-R^{24}; -C(O)-(C_1-C_6)alkylene -R^{23}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -S(O)$_2$-R^{24}, and -(C_1-C_6)alkylene-O-(C_1-C_6)alkylene-O-R^{24}, with the proviso that –O-O- cannot be formed with R^{21} and the atom said R^{21} is attached to;

R^{22} is selected from the group consisting of unsubstituted (C_6-C_{12})aryl and (C_6-C_{12})aryl substituted with one or more W^1 groups;

R^{23} is selected from the group consisting of unsubstituted (C_2-C_{10})heteroaryl and (C_2-C_{10})heteroaryl substituted with one or more W^2 groups;

R^{24} is is selected from the group consisting of unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more X^4 groups, unsubstituted (C_1-C_6)alkyl and (C_1-C_6)alkyl substituted with one or more U groups;

each R^{25} is independently selected from the group consisting of H, R^{22}, R^{23}, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, -(C_1-C_6)alkylene-OR^{19}, -(C_1-C_6)alkylene-N$R^{19}R^{19}$, -(C_1-C_6)alkylene-SR^{19},- (C_1-C_6)alkylene-R^{23}, -(C_1-C_6)alkylene-R^{22},

unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl, -(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -C(O)-R^{24}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -$S(O)_2$-R^{24}, -C(O)-NH-R^{22} and -C(O)-NH-R^{24};

with the proviso wherein the group -$N(R^{25})_2$ both R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted (C_3-C_5)heterocycloalkyl group is fused with (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, (C_3-C_7)cycloalkyl or (C_3-C_5)heterocycloalkyl;

each W^1 is independently selected from the group consisting of halogen, -CN, -OH, -O-$S(O)_2$-(C_1-C_6)haloalkyl, unsubstituted (C_6-C_{12})aryl, (C_6-C_{12})aryl substituted with one or more Z groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more Z groups, and –O-(C_1-C_6)alkyl;

each W^2 is independently selected from the group consisting of halogen, unsubstituted (C_6-C_{12})aryl, and (C_6-C_{12})aryl substituted with one or more Z groups;

each W^3 is independently selected from the group consisting of –OH, (C_1-C_6)alkyl, -(C_1-C_6)alkylene-OH, -O-(C_1-C_6)alkyl, -C(O)- (C_1-C_6)alkyl, -$C(O)NH_2$, -NHC(O) (C_1-C_6)alkyl, -NHC(O)H, -NHC(O)-O-(C_1-C_6)alkyl and -C(O)-O-(C_1-C_6)alkyl; or

two W^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each W^4 is independently halogen or (C_1-C_6)alkyl;

Ar^1 and Ar^2 are independently selected from the group consisting of R^{12} and R^{13};

each X^1 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-$S(O)_2$-(C_1-C_6)haloalkyl, unsubstituted (C_6-C_{12})aryl, (C_6-C_{12})aryl substituted with one or more Z groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more Z groups, –O-(C_3-C_7)cycloalkyl, –O-(C_3-C_7)cycloalkyl(C_1-C_6)alkyl, -O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)$N(R^{20})_2$,

-O-(C_1-C_6)alkylene-O-R^{19}, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, unsubstituted –O-(C_1-C_6)alkyl, -O-(C_1-C_6)alkyl substituted with one or more U groups, -O-(C_2-C_7)alkenyl, -O-(C_1-C_6)alkylene-O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)R^{24}, -O-(C_1-C_6)alkylene-C(O)OR^{19}, –O-(C_1-C_6)alkyl, -N(R^{25})$_2$, -C(O)(C_1-C_6)alkyl, –C(O)OH, -C(O)O-(C_1-C_6)alkyl, -C(O)O-(C_3-C_7)cycloalkyl, -C(O)N(R^{25})$_2$, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, unsubstituted(C_3-C_5)heterocycloalkyl, -(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -O-(C_2-C_7)alkenylene-O-(C_1-C_6)alkylene-O-R^{24}, -O-(C_1-C_6)alkylene-N(R^{25})$_2$, -O-(C_1-C_6)alkylene-C(O)N(R^{25})$_2$, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S($O)_2$-R^{24}, and (C_2-C_7)alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted (C_3-C_5)heterocycloalkyl group is fused with (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, (C_3-C_7)cycloalkyl or (C_3-C_5)heterocycloalkyl;

each X^2 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S($O)_2$-(C_1-C_6)haloalkyl, unsubstituted (C_6-C_{12})aryl, (C_6-C_{12})aryl substituted with one or more Z groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more Z groups, –O-(C_3-C_7)cycloalkyl, –O-(C_3-C_7)cycloalkyl(C_1-C_6)alkyl, -O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)N(R^{20})$_2$, -O-(C_1-C_6)alkylene-O-R^{19}, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, unsubstituted –O-(C_1-C_6)alkyl, -O-(C_1-C_6)alkyl substituted with one or more U groups, -O-(C_2-C_7)alkenyl, -O-(C_1-C_6)alkylene-O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)R^{24}, -O-(C_1-C_6)alkylene-C(O)OR^{19}, –O-(C_1-C_6)alkyl, -N(R^{25})$_2$, -C(O)(C_1-C_6)alkyl, –C(O)OH, -C(O)O-(C_1-C_6)alkyl, -C(O)O-(C_3-C_7)cycloalkyl, -C(O)N(R^{25})$_2$,

-O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, unsubstituted(C_3-C_5)heterocycloalkyl, -(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -O-(C_2-C_7)alkenylene-O-(C_1-C_6)alkylene-O-R^{24}, -O-(C_1-C_6)alkylene-N(R^{25})$_2$, -O-(C_1-C_6)alkylene-C(O)N(R^{25})$_2$, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S(O)$_2$-R^{24}, and (C_2-C_7)alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted (C_3-C_5)heterocycloalkyl group is fused with (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, (C_3-C_7)cycloalkyl or (C_3-C_5)heterocycloalkyl;

each X^3 is independently selected from the group consisting of –OH, (C_1-C_6)alkyl, -(C_1-C_6)alkylene-OH, -O-(C_1-C_6)alkyl, -C(O)- (C_1-C_6)alkyl, -C(O)NH_2, -NHC(O) (C_1-C_6)alkyl, -NHC(O)H, -NHC(O)-O-(C_1-C_6)alkyl and -C(O)-O-(C_1-C_6)alkyl; or

two X^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each X^4 is independently halogen or (C_1-C_6)alkyl;

each U is independently selected from the group consisting of –OH, -O-(C_1-C_6)alkyl, -O-(C_6-C_{12})aryl, -O-(C_1-C_6)alkylene-(C_6-C_{12})aryl, -O-(C_1-C_6)alkylene-O-(C_1-C_6)alkyl, -O-(C_1-C_6)alkylene-O-(C_1-C_6)haloalkyl, -O-(C_1-C_6)alkylene-O-(C_6-C_{12})aryl, halogen, -CN, (C_3-C_7)cycloalkyl, (C_3-C_5)heterocycloalkyl, (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, OTBS, OTIPS and OTf;


and

each Z is independently selected from the group consisting of –OH; -O-(C_1-C_6)alkyl; halogen; (C_1-C_6)alkyl; –CN; -CF_3; (C_3-C_7)cycloalkyl; -(C_1-C_6)alkylene-OH; -(C_1-C_6)alkylene-O-(C_1-C_6)alkyl; -(C_1-C_6)alkylene-O-(C_1-C_6)alkyl substituted with one or more groups selected from the group consisting of –OH, -O-(C_1-C_6)alkyl, halogen, -CN, (C_3-C_7)cycloalkyl, (C_3-C_5)heterocycloalkyl, (C_6-C_{12})aryl,

$(C_2$-$C_{10})$heteroaryl; -$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkyl; -$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl; -$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl; and -$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl substituted with one or more groups selected from the group consisting of halogen, -CN, -OH, -O-S(O)$_2$-$(C_1$-$C_6)$haloalkyl, $(C_6$-$C_{12})$aryl, $(C_2$-$C_{10})$heteroaryl and –O-$(C_1$-$C_6)$alkyl; or

two Z groups together with the ring carbon atom to which they are attached form a carbonyl group.


The dashed lines (----) in Formula (I) represent bonds which can independently be either a single bond or a double bond. The ring of Formula (I) can contain no double bonds, or one or more double bonds (i.e., one or two double bonds), provided that the resulting compound is stable. Non-limiting examples of compounds of Formula (I) can have one of the following generic formulae, where R^1, R^2, R^{15}, Ar^1, and Ar^2 are as defined herein:




, and

.


When present, each R^{15} group of Formula (I) can be bonded to any carbon atom of Formula (I) which is capable of substitution. As a non-limiting example, the following generic structure:




includes the following structures wherein group R^{15}, when present, can be bonded to any of the ring carbon atoms. For example, when one R^{15} group is present, the above generic structure includes the following:




When two R^{15} groups are present, the generic structure above includes the following:






When the six-membered ring of Formula (I) contains one or more double bonds, each R^{15}, when present, may be attached to any ring carbon atom to which such substitution is possible.

Similarly, it is contemplated that the compounds of Formula (I) include all possible stable stereoisomers. As a non-limiting example, the following generic structure (wherein each R^{15} is H):




includes the following stereoisomers:




, and .

In another embodiment of the compounds of Formula (I), or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, R^1 and R^2 together with the carbon atoms to which they are shown attached in Formula (I) form a group Q as shown in Formula (IA):




(IA)

wherein the group Q is selected from the group consisting of:




(a) , (b) , (c) , (d) , (e) , (f) , and




(f) .


Thus, the compounds of Formula (IA) of the present invention include the following generic formulae:




(II-1) ,




(II-2)




(III-1) ,




(III-2)




(IV) ,




(V) ,




(VI) ,




(VII-1) ,




(VII-2) ,




(VIII-1) ,




(VIII-2) ,



(IX) ,




(X) , and




(XI) .

It will be recognized by one of skill in the art that Formulae (VII-1), (VII-2), (VIII-1), and (VIII-2) can describe equivalent tautomeric forms of the same compound (e.g., when R^5 of Formulae (VIII-1) or (VIII-2) is $-NHR^{16}$, and R^7 of Formulae (VII-1) or (VII-2) is H). Thus, the following structural formulae are considered equivalent:




In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (II):



(II) .

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IIA):



(IIA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IIA), wherein:

Ar^1 and Ar^2 are independently selected from the group consisting of R^{12} and R^{13};

R^7 is selected from the group consisting of H, alkyl, alkenyl, -alkylene-$N(R^9)_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, and -C(O)-O-R^{11};

R^9 is selected from the group consisting of H and alkyl;

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted pyridyl and pyridyl substituted with one or more X^2 groups;

R^{14} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more X^4 groups; and

R^{15} is selected from the group consisting of H, alkyl, alkenyl, -alkylene-R^{12}, and –O-alkenyl.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IIA), wherein:


Ar^1 is phenyl substituted with one or more X^1 groups;

Ar^2 is phenyl substituted with one or more X^1 groups or pyridyl substituted with one or more X^2 groups;

R^7 is H; and


R^{15} is alkyl.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (III):



(III)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IIIA):



(IIIA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IIIA), wherein:

Ar^1 and Ar^2 are independently selected from the group consisting of R^{12} and R^{13};

R^3 and R^4 are each independently H or alkyl;

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups; and

R^{13} is selected from the group consisting of unsubstituted pyridyl and pyridyl substituted with one or more X^2 groups.


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IIIA), wherein:

R^3 and R^4 are each independently H or $-CH_3$;

Ar^1 is unsubstituted phenyl or phenyl substituted with one or more halogens; and


Ar^2 is selected from the group consisting of unsubstituted phenyl, phenyl substituted with one or more halogens, unsubstituted pyridyl, and pyridyl substituted with one or more X^2 groups.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IV):



(IV)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IVA):



(IVA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IVA), wherein:

Ar^1 and Ar^2 are independently selected from the group consisting of R^{12} and R^{13};

R^8 is selected from the group consisting of -alkylene-R^{12}, -C(O)-R^{17}, -$S(O_2)$-R^{14}, -C(O)-$N(R^{18})_2$, and R^{14};

R^9 is selected from the group consisting of H and alkyl;

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted pyridyl and pyridyl substituted with one or more X^2 groups;

R^{14} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more X^4 groups; and

R^{17} is selected from the group consisting of unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, -alkylene-R^{12}, -O-R^9, and R^{12}; and

each R^{18} is independently selected from the group consisting of H, R^{12}, and R^{14}.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IVA), wherein:

R^8 is selected from the group consisting of $-CH_2-R^{12}$, $-CH(CH_3)-R^{12}$, $-C(O)-R^{17}$, $-S(O_2)-R^{14}$, $-C(O)-N(R^{18})_2$, and R^{14};

R^9 is selected from the group consisting of H and alkyl;


R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

R^{14} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more X^4 groups; and

R^{17} is selected from the group consisting of unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, $-CH_2-R^{12}$, $-CH(CH_3)-R^{12}$, $-O-R^9$, and R^{12}; and


each R^{18} is independently H, unsubstituted cycloalkyl, cycloalkyl substituted with one or more X^4 groups, unsubstituted aryl, and aryl substituted with one or more X^1 groups.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (V):



(V)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VA):




(VA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (VA), wherein:

R^7 is H or -alkylene-R^{12}; and


R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VI):



(VI)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VIA):



(VIA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (VIA), wherein:

R^3, R^5, and R^6 are each independently selected from the group consisting of H, -O-R^9, and R^{11};

R^9 is H or alkyl;


R^{11} is selected from the group consisting of alkyl, unsubstituted phenyl, and phenyl substituted with one or more X^1 groups;

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted pyridyl and pyridyl substituted with one or more X^2 groups; and


Ar^2 is selected from the group consisting of R^{12} and R^{13}.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VII):



(VII)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VIIA):




(VIIA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (VIIA), wherein:

R^{16} is–C(O)-R^{12}; and


R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VIII):



(VIII)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (VIIIA):



(VIIIA)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (VIIIA), wherein:

R^3, R^4, and R^5 are each independently H, or $-N(R^{16})_2$;

R^9 is selected from the group consisting of H and R^{11};


R^{11} is selected from the group consisting of alkyl, unsubstituted aryl, and aryl substituted with one or more X^1 groups;

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups; and

R^{16} is selected from the group consisting of R^9 and $-C(O)-R^{12}$.

In another embodiment of the compounds of Formula (I), or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, R^1 is selected from the group consisting of $-C(O)-N(R^{10})_2$, -C(O)-O-alkyl, and $-C(O)-R^{14}$; and R^2 is selected from the group consisting of H, alkyl, alkyl substituted with one or more –OH groups, and -alkylene-$N(R^{10})_2$.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IB):



(IB)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IB), wherein:

R^9 is selected from the group consisting of H and alkyl;


each R^{10} is independently selected from the group consisting of H, alkyl substituted with one or more -OH groups, -alkylene-R^{12}, -alkylene-R^{13}, -alkylene-R^{14}, -alkylene-O-R^9, R^{14}, and benzo-fused cycloalkyl;

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more X^2 groups; and

R^{14} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more X^4 groups.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IC):



(IC)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IC), wherein:

R^1 is –C(O)-N(R^{10})$_2$ or -C(O)-O-alkyl;


R^2 is selected from the group consisting of H, alkyl, alkyl substituted with one or more –OH groups, and -alkylene-N(R^{10})$_2$;

each R^{10} is independently selected from the group consisting of H, -alkylene-R^{12}, and -C(O)-R^{14};

R^{12} is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups; and


R^{15} is H or –OH.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (ID):



(ID)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof have the following Formula (IE):



(IE)

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (ID), wherein:

Y^1 is NH or N-Boc;

R^3 is H or alkyl;

R^4 is H or alkyl;

R^{15} is alkyl;

and


Ar^1 and Ar^2 are selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups.


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (ID), wherein:

each X^1 is independently selected from the group consisting of halogen, -CN, -OH, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-$C(O)N(R^{20})_2$, -O-alkylene-O-R^{19}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl -O-alkylene-O-alkylene-OR^{19}, -O-alkylene-$C(O)R^{24}$, -O-alkylene-$C(O)OR^{19}$ and –O-alkyl.


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (ID), wherein:






each X^1 is independently selected from the group consisting of $-OCH_3$, -OH, -OTf, -CN, $-OCH_2CH_3$, $-OCH(CH_3)_2$,














and


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (ID),



(ID)

or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

Y^1 is NH or N-Boc;

R^3 is H or alkyl;


R^4 is H or alkyl;


R^{15} is alkyl;

Ar^1 and Ar^2 are selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;




and











each X^1 is independently selected from the group consisting of $-OCH_3$, -OH, -OTf, -CN, $-OCH_2CH_3$, $-OCH(CH_3)_2$,

















and


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (IE),



(IE)

or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

Y^1 is NH or N-Boc;


R^3 is H or alkyl;

R^4 is H or alkyl;

R^{15} is alkyl;

and


each X^1 is independently selected from the group consisting of $-OCH_3$, -OH, -OTf, -CN, $-OCH_2CH_3$, $-OCH(CH_3)_2$,




In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein each R^{15} is independently selected from the group consisting of H, $-N_3$, halogen, alkenyl, -alkylene-R^{12}, -alkylene-O-R^9, -alkylene-$N(R^{18})_2$, -alkylene-C(O)H, -OH, -CN, -O-alkyl, |-$C(O)N(R^{18})_2$, $-N(R^{18})_2$, $-NHC(O)R^{18}$, $-NHC(O)_2R^{18}$, $-NR^{18}C(O)N(R^{18})_2$, $-NHS(O)_2R^{18}$, –O-alkenyl, $-C(O)_2R^{18}$, unsubstituted alkyl, alkyl substituted with one or more U groups, -O-alkylene-$C(O)R^{18}$ or $-C(O)R^{18}$; with the proviso wherein the group $-N(R^{18})_2$, both R^{18} groups taken together with the N atom to

which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl.


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein each R^{15} is independently selected from the group consisting of H, -OH, $-OCH_3$, $-OCH_2CH_3$, $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-N_3$, $-NH_2$, $-CO_2H$, $-CO_2CH_3$, $-CH_2OH$, $-CH_2CH_2OH$, $-CH_2CH_2OTBS$, $-CH_2OCH_3$, $-OCH_2CH_2OH$,






In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein: R^3 and R^4 are each independently selected from the group consisting of H, -O-R^9, R^{11}, and $-N(R^{16})_2$.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein each R^3 and R^4 is independently selected from the group consisting of H, $-CH_3$, $-CH_2OH$, $-CH_2OCH_2CH_2OCH_3$, $-CH_2OCH_3$, $-CH_2OCH_2CH_3$, $-CH_2OCH_2CH_2CH_3$, $-CH_2CH_2OH$,




In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein R^7 is selected from the group consisting of H, alkyl, arylalkyl, alkenyl, -alkylene-$N(R^9)_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, -alkylene-C(O)H, -C(O)-O-R^{11}, and Boc.


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein R^7 is H, $-CH_3$, Boc,





In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein Ar^1 is phenyl substituted with one or more X^1 groups; and each X^1 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-$S(O)_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-$C(O)N(R^{20})_2$, -O-alkylene-O-R^{19}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl, -O-alkylene-O-alkylene-OR^{19}, -O-alkylene-$C(O)R^{24}$, -O-alkylene-$C(O)OR^{19}$,–O-alkyl, -$N(R^{25})_2$, -C(O)alkyl, –C(O)OH, -C(O)O-alkyl, -C(O)O-cycloalkyl, -$C(O)N(R^{25})_2$, -O-alkylene-heterocycloalkyl, -O-alkylene-heterocycloalkyl substituted with one or more W^3 groups, unsubstituted heterocycloalkyl, -heterocycloalkyl substituted with one or more W^3 groups, -O-alkenylene-O-alkylene-O-R^{24}, -O-alkylene-$N(R^{25})_2$, -O-alkylene-$C(O)N(R^{25})_2$, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -$S(O)_2$-R^{24}, and alkenyl; with the proviso wherein the group -$N(R^{20})_2$ or -$N(R^{25})_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl.


In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein each X^1 is independently selected from the group consisting of Cl, F, $–CH_3$, $-OCH_3$, $-OCH_2CH_3$, -OH, -OTf, -CN, $-OCH_2CH_3$, $-OCH(CH_3)_2$,












In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein Ar^2 is phenyl substituted with one or more X^1 groups or pyridyl substituted with one or more X^2 groups; X^1 is selected from the group consisting of –OH, -CN, halogen, -OTIPS, -OTf, alkyl, -O-alkyl, -O-alkyl-OH and heteroaryl; andX^2 is selected from the group consisting of halogen and cycloalkyl.

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of Formula (I), wherein Ar^2 is phenyl substituted with one or


more X^1 groups or pyridyl substituted with one or more X^2 groups, X^1 is selected from the group consisting of –OH, -CN, Cl, -OTIPS, $-CH_3$,$-OCH_3$, $-OCH_2CH_3$, -OTf and  ; and X^2 is selected from the group consisting of Cl, Br and  .

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of the following Formulas (IF) or (IFa):




(IIF) or (IIFa) .

In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of the following Formulas (IF) or (IFa):


















(IG) .













In another embodiment, the compounds of the present invention, or pharmaceutically acceptable salts, solvates, esters, or stereoisomers thereof, have the structure of the following Formula (IH):













(IH) .















An even further embodiment of formula I are those compounds with the following structures

















,

,

,

,

,

,

,

,

,

,

,

,

,

,

,


,

,

,

,

,

,

,

,

,

,

,

,
5



,

,

,

,

,

,

,

,

,

,


5



,

,

,

,

,

,

,

,

,

,

,

,

,

,



















5


















O
Me
HN
H
H
H
Cl
O
Cl


,


O
Me
HN
H
H
H
Cl
O
O
Cl


,


O
Me
HN
H
H
H
Cl
O
CF3
Cl


,


O
Me
HN
H
H
H
Cl
O
NH2
O
Cl


,


O
Me
HN
H
H
H
Cl
O
OH
Cl


,


O
Me
HN
H
H
H
Cl
OCHF2
Cl


,


O
Me
HN
H
H
H
Cl
O
Cl


,


O
Me
HN
H
H
H
Cl
O
Cl









,


O
Et
HN
H
H
H
Cl
O
O
O
Cl


,



O
Et
HN
H
H
H
Cl
O
OTBS
Cl


,












5





































5



































































































5


















,

,

,

,

,

,

,

,

,

,

,

,


5






















































5




5





5





,

,

,

,

,

,

,

,

,

,

,

,









5























5








and

The compounds of the present invention, e.g., according to Formula (I), are preferably purified to a degree suitable for use as a pharmaceutically active substance. That is, the compounds of Formula (I) can have a purity of 95 wt% or more (excluding adjuvants such as pharmaceutically acceptable carriers, solvents, etc., which are used in formulating the compound of Formula (I) into a conventional form, such as a pill, capsule, IV solution, etc. suitable for administration into a patient). More preferably, the purity can be 97 wt% or more, even more preferably, 99 wt% or more. A purified compound

of Formula (I) includes a single isomer having a purity, as discussed above, of 95 wt% or more, 97 wt% or more, or 99 wt% or more, as discussed above. For example, the purified compound of Formula (I) can have a purity of 95 wt% or more, 97 wt% or more, or 99 wt% or more.


Alternatively, the purified compound of Formula (I) can include a mixture of isomers, each having a structure according to Formula (I), where the amount of impurity (i.e., compounds or other contaminants, exclusive of adjuvants as discussed above) is 5 wt% or less, 3 wt% or less, or 1 wt% or less. For example, the purified compound of Formula (I) can be an isomeric mixture of compounds of Formula (I), where the ratio of the amounts of the two isomers is approximately 1:1, and the combined amount of the two isomers is 95 wt% or more, 97 wt% or more, or 99 wt% or more.

The variables m and n can represent, respectively, the integers 0 or 1 and 1 or 2, with the proviso that the sum of m and n (i.e., m + n) is 1 or 2. Thus, in one embodiment of the compounds of Formula (I), m is 0 and n is 2, e.g.:



; and

in another embodiment of the compounds of Formula (I), m is 1 and n is 1, e.g.:




R^1 is selected from the group consisting of $-C(O)-N(R^{10})_2$, -C(O)-O-alkyl, and $-C(O)-R^{14}$. The substituents R^{10} and R^{14} are defined as disclosed herein. The term "alkyl" of -C(O)-O-alkyl includes, for example, lower alkyls such as $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$ (n-propyl), $-CH(CH_3)_2$ (i-propyl), $-CH_2CH_2CH_2CH_3$ (n-butyl), $-C(CH_3)_3$ (t-butyl), $-CH(CH_3)-CH_2CH_3$ (sec-butyl), $-CH_2CH(CH_3)_2$ (i-butyl), $-CH_2CH_2CH_2CH_2CH_3$ (n-pentyl), $-CH_2C(CH_3)_3$ (neo-pentyl), etc. Thus, -C(O)-O-alkyl includes, for example, $-C(O)-O-CH_3$, $-C(O)-O-CH_2CH_3$, $-C(O)-O-CH_2CH_2CH_3$, $-C(O)-O-CH(CH_3)_2$, $-C(O)-O-CH_2CH_2CH_2CH_3$, $-C(O)-O-C(CH_3)_3$, $-C(O)-O-CH(CH_3)-CH_2CH_3$,


-C(O)-O-$CH_2CH(CH_3)_2$, -C(O)-O-$CH_2CH_2CH_2CH_2CH_3$, -C(O)-O-$CH_2C(CH_3)_3$, etc. Likewise, -C(O)-N(R^{10})$_2$ includes –C(O)-NH_2, –C(O)-NH(alkyl), -C(O)-N(alkyl)$_2$, -C(O)-NH(alkyl-OH), -C(O)-N(alkyl-OH)$_2$, -C(O)-NH-alkylene-R^{12}, -C(O)-N(alkyl)-alkylene-R^{12}, -C(O)-NH-alkylene-R^{13}, -C(O)-N(alkyl)-alkylene-R^{13}, -C(O)-NH-alkylene-R^{14}, -C(O)-N(alkyl)-alkylene-R^{14}, -C(O)-NH-C(O)-R^{14}, -C(O)-N(alkyl)-C(O)-R^{14}, -C(O)-NH-alkylene-O-R^9, -C(O)-N(alkyl)-alkylene-O-R^9, -C(O)-NH-heterocycloalkyl optionally substituted on the heterocycloalkyl with one or more X^3 groups, -C(O)-N(alkyl)-heterocycloalkyl optionally substituted on the heterocycloalkyl with one or more X^3, -C(O)-NH-(benzo-fused cycloalkyl), and -C(O)-N(alkyl)-(benzo-fused cycloalkyl). The terms "alkyl", "alkylene", "benzo-fused cycloalkyl", and "heterocycloalkyl" are as defined herein. The term "alkyl-OH" above refers to an alkyl substituted with one or more –OH groups, for example the groups described below for R^2. Each R^{10} of –C(O)-N(R^{10})$_2$ can independently include any group defined herein for R^{10}, and is not limited to the specific groups and combinations above.

R^2 is selected from the group consisting of H, alkyl, alkyl substituted with one or more –OH groups, and –alkylene-N(R^{10})$_2$. Alkyl includes, for example, the lower alkyls described above for R^1. Alkyl substituted with one or more or more –OH groups includes, for example, -CH_2-OH, -CH_2CH_2-OH, -$CH(OH)CH_3$, -$CH(OH)CH_2CH_3$, -$CH_2CH(OH)CH_3$, -$CH_2CH_2CH_2$-OH, -$CH(OH)CH_2CH_2$-OH, -$CH(OH)CH(OH)CH_2$-OH, -$C(OH)(CH_3)_2$, -$CH(CH_3)CH_2$-OH, -$CH(OH)CH_2CH_2CH_3$, -$CH_2CH(OH)CH_2CH_3$, -$CH_2CH_2CH(OH)CH_3$, -$CH_2CH_2CH_2CH_2$-OH, - $CH(OH)CH_2CH(OH)CH_3$, etc. Non-limiting examples of the "alkylene" portion of –alkylene-N(R^{10})$_2$ include -CH_2-, -CH_2CH_2-, -$CH_2CH_2CH_2$-, -$CH_2CH_2CH_2CH_2$-, -$CH_2CH_2CH_2CH_2CH_2$-, -$CH_2CH_2CH_2CH_2CH_2CH_2$-, -$CH(CH_3)$-, -$CH(CH_3)CH_2$-, -$CH_2CH(CH_3)$-, -$CH(CH_2CH_3)$-, -$CH(CH_3)CH_2CH_2$-, -$CH_2CH(CH_3)CH_2$-, -$CH_2CH_2CH(CH_3)$-, -$CH(CH_2CH_3)CH_2$-, -$CH_2CH(CH_2CH_3)$-, -$CH(CH_2CH_2CH_3)$-, -$CH(CH_3)CH(CH_3)$-, etc. R^{10} is defined herein, and each R^{10} of -alkylene-N(R^{10})$_2$ is independently selected and may be independently combined in any combination with any of the alkylene groups defined herein.

In an alternative embodiment, R^1 and R^2 together with the carbon atoms to which they are shown attached in Formula (I) form a group Q selected from:



(a) , (b) , (c) , (d) , (e) , (f) , and



(f) ,

where Y^1, Y^2, R^3, R^4, R^5, R^6, R^7, and R^{16} are as defined herein. The various possible bicyclic structures thus formed are described above.

R^3, R^4, R^5, and R^6 are each independently selected from the group consisting of H, -O-R^9, R^{11}, and –N(R^{16})$_2$. R^9 and R^{11} are as defined herein. For example, -O-R^9 can include –OH and –O-R^{11}. –O-R^{11} can include, for example, -O-CH_3, -O-CH_2CH_3, -O-$CH_2CH_2CH_3$, -O-$CH(CH_3)_2$, -O-$CH_2CH_2CH_2CH_3$, -O-$C(CH_3)_3$, -O-$CH(CH_3)CH_2CH_3$, -O-$CH_2CH(CH_3)_2$, -O-$CH_2CH_2CH_2CH_2CH_3$, -O-$CH_2C(CH_3)_3$, -O-phenyl, -O-naphthyl, -O-biphenyl, etc., wherein said phenyl, naphthyl, and biphenyl are unsubstituted or substituted with one or more X^1 groups, wherein said X^1 is as described herein. Likewise, non-limiting examples of R^{11} include the alkyl groups defined above for R^1, as well as unsubstituted or X^1 substituted aryls, e.g., phenyl, naphthyl, biphenyl, etc. R^{16} is defined herein, and each R^{16} of -N(R^{16})$_2$ may be independently selected. Thus, non-limiting examples of -N(R^{16})$_2$ include, for example, -N(R^9)$_2$ wherein each R^9 is independently selected and defined herein and can include H, alkyl, unsubstituted aryl, and aryl substituted with one or more X^1 groups in any combination; and -N(R^9)-C(O)-R^{12} wherein R^9 and R^{12} are independently selected and defined herein. Thus, non-limiting examples of -N(R^9)-C(O)-aryl include -NH-C(O)-aryl, -N(alkyl)-C(O)-aryl, -N(aryl)-C(O)-aryl, wherein each "aryl" is

e.g., unsubstituted or X^1 substituted phenyl, naphthyl, biphenyl, etc., and each "alkyl" is selected from the lower alkyls described for R^1 above.

R^7 is selected from the group consisting of H, arylalkyl, alkyl, alkenyl, -alkylene-$N(R^9)_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, -alkylene-C(O)H, and -C(O)-O-R^{11}. The term "alkyl" includes, for example, the lower alkyl groups described above for R^1. The term "arylalkyl" includes, for example, which may be substituted or unsubstituted, is as defined herein. The term "alkenyl" includes, for example, -CH=CH_2, -CH_2-CH=CH_2, -CH=CH-CH_3, -CH_2-C(CH_3)=CH_2, and -CH_2-CH=$CHCH_3$. The term "-alkylene-R^{12}" includes combinations of the "alkylene" groups defined above, and R^{12} groups as defined herein. For example, "-alkylene-R^{12}" includes e.g., -CH_2-aryl, -CH_2CH_2-aryl, -$CH_2CH_2CH_2$-aryl, -$CH_2CH_2CH_2CH_2$-aryl, -$CH_2CH_2CH_2CH_2CH_2$-aryl, -$CH_2CH_2CH_2CH_2CH_2CH_2$-aryl, -CH(CH_3)-aryl, -CH(CH_3)CH_2-aryl, -CH_2CH(CH_3)-aryl, -CH(CH_2CH_3)-aryl, -CH(CH_3)CH_2CH_2-aryl, -CH_2CH(CH_3)CH_2-aryl, -CH_2CH_2CH(CH_3)-aryl, -CH(CH_2CH_3)CH_2-aryl, -CH_2CH(CH_2CH_3)-aryl, -CH($CH_2CH_2CH_3$)-aryl, -CH(CH_3)CH(CH_3)-aryl, etc., wherein the "aryl" includes for example phenyl, naphthyl, or biphenyl which may be unsubstituted or substituted with one or more X^1 group. Non-limiting examples of the "alkylene" portion of -alkylene-$N(R^9)_2$ include -CH_2-, -CH_2CH_2-, -$CH_2CH_2CH_2$-, -$CH_2CH_2CH_2CH_2$-, -$CH_2CH_2CH_2CH_2CH_2$-, -$CH_2CH_2CH_2CH_2CH_2CH_2$-, -CH(CH_3)-, -CH(CH_3)CH_2-, -CH_2CH(CH_3)-, -CH(CH_2CH_3)-, -CH(CH_3)CH_2CH_2-, -CH_2CH(CH_3)CH_2-, -CH_2CH_2CH(CH_3)-, -CH(CH_2CH_3)CH_2-, -CH_2CH(CH_2CH_3)-, -CH($CH_2CH_2CH_3$)-, -CH(CH_3)CH(CH_3)-, etc. R^9 is as defined herein, and each R^9 of -alkylene-$N(R^9)_2$ is independently selected and may be independently combined in any combination with any of the alkylene groups defined herein. The term "-alkylene-O-R^9" includes, for example, -CH_2-O-R^9, -CH_2CH_2-O-R^9, -$CH_2CH_2CH_2$-O-R^9, -$CH_2CH_2CH_2CH_2$-O-R^9, -$CH_2CH_2CH_2CH_2CH_2$-O-R^9, -$CH_2CH_2CH_2CH_2CH_2CH_2$-O-R^9, -CH(CH_3)-O-R^9, -CH(CH_3)CH_2-O-R^9, -CH_2CH(CH_3)-O-R^9, -CH(CH_2CH_3)-O-R^9, -CH(CH_3)CH_2CH_2-O-R^9, -CH_2CH(CH_3)CH_2-O-R^9, -CH_2CH_2CH(CH_3)-O-R^9, -CH(CH_2CH_3)CH_2-O-R^9, -CH_2CH(CH_2CH_3)-O-R^9, -CH($CH_2CH_2CH_3$)-O-R^9, -CH(CH_3)CH(CH_3)-O-R^9, -CH(OR^9)CH_2-, etc. Non-limiting examples of -alkylene-R^{12} include for example, -CH_2-aryl, -CH_2CH_2-aryl, -$CH_2CH_2CH_2$-aryl,

-$CH_2CH_2CH_2CH_2$-aryl, -$CH_2CH_2CH_2CH_2CH_2$-aryl, -$CH_2CH_2CH_2CH_2CH_2CH_2$-aryl, -$CH(CH_3)$-aryl, -$CH(CH_3)CH_2$-aryl, -$CH_2CH(CH_3)$-aryl, -$CH(CH_2CH_3)$-aryl, -$CH(CH_3)CH_2CH_2$-aryl, -$CH_2CH(CH_3)CH_2$-aryl, -$CH_2CH_2CH(CH_3)$-aryl, -$CH(CH_2CH_3)CH_2$-aryl, -$CH_2CH(CH_2CH_3)$-aryl, -$CH(CH_2CH_2CH_3)$-aryl, -$CH(CH_3)CH(CH_3)$-aryl, etc., wherein the "aryl" includes for example phenyl, naphthyl, or biphenyl which may be unsubstituted or substituted with one or more X^1 group. Non-limiting examples of -C(O)-R^{14} include -C(O)-cyclopropyl, -C(O)-cyclobutyl, -C(O)-cyclopentyl, -C(O)-cyclohexyl, -C(O)-cycloheptyl, wherein said cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, or cycloheptyl portion are unsubstituted or substituted with one or more X^4 groups. Non-limiting examples of -C(O)-O-R^{11} include –C(O)-OH, -C(O)-O-CH_3, -C(O)-O-CH_2CH_3, -C(O)-O-$CH_2CH_2CH_3$, -C(O)-O-$CH(CH_3)_2$, -C(O)-O-$C(CH_3)_3$, -C(O)-O-$CH_2CH_2CH_2CH_3$, -C(O)-O-$CH(CH_3)CH_2CH_3$, -C(O)-O-$CH_2CH(CH_3)_2$, -C(O)-O-phenyl, -C(O)-O-naphthyl, -C(O)-O-biphenyl, etc., wherein said phenyl, naphthyl, and biphenyl may be unsubstituted or substituted with X^1.


R^8 is selected from the group consisting of H, -alkylene-R^{12}, -C(O)-R^{17}, -$S(O_2)$-R^{11}, -$S(O_2)$-R^{14}, -C(O)-$N(R^{18})_2$, and R^{14}. The terms "alkenyl", and "-alkylene-R^{12}", for example, are as defined above. The term "-C(O)-R^{17}" includes, for example -C(O)-heterocycloalkyl, -C(O)-alkylene-R^{12}, -C(O)-O-R^9, and –C(O)-R^{12}. Thus, -C(O)-R^{17} includes, for example, -C(O)-morpholinyl, -C(O)-piperazinyl, -C(O)-piperidinyl, -C(O)-pyrrolidinyl, -C(O)-tetrahydrofuranyl, -C(O)-tetrahydrofuranyl, -C(O)-thiazolinyl, -C(O)-tetrahydropyranyl, etc.; -C(O)-CH_2-aryl, -C(O)-CH_2CH_2-aryl, -C(O)-$CH_2CH_2CH_2$-aryl, -C(O)-$CH_2CH_2CH_2CH_2$-aryl, -C(O)-$CH_2CH_2CH_2CH_2CH_2$-aryl, -C(O)-$CH_2CH_2CH_2CH_2CH_2CH_2$-aryl, -C(O)-$CH(CH_3)$-aryl, -C(O)-$CH(CH_3)CH_2$-aryl, -C(O)-$CH_2CH(CH_3)$-aryl, -C(O)-$CH(CH_2CH_3)$-aryl, -C(O)-$CH(CH_3)CH_2CH_2$-aryl, -C(O)-$CH_2CH(CH_3)CH_2$-aryl, -C(O)-$CH_2CH_2CH(CH_3)$-aryl, -C(O)-$CH(CH_2CH_3)CH_2$-aryl, -C(O)-$CH_2CH(CH_2CH_3)$-aryl, -C(O)-$CH(CH_2CH_2CH_3)$-aryl, -C(O)-$CH(CH_3)CH(CH_3)$-aryl, etc., wherein the term "aryl" includes for example phenyl, naphthyl, or biphenyl which may be unsubstituted or substituted with one or more X^1 group; -C(O)-O-H, -C(O)-O-CH_3, -C(O)-O-CH_2CH_3, -C(O)-O-$CH_2CH_2CH_3$, -C(O)-O-$CH(CH_3)_2$,

-C(O)-O-$CH_2CH_2CH_2CH_3$, -C(O)-O-$C(CH_3)_3$, -C(O)-O-$CH(CH_3)$-CH_2CH_3, -C(O)-O-$CH_2CH(CH_3)_2$, -C(O)-O-$CH_2CH_2CH_2CH_2CH_3$, -C(O)-O-$CH_2C(CH_3)_3$, etc., -C(O)-O-phenyl, -C(O)-O-naphthyl, -C(O)-O-biphenyl, wherein said phenyl, naphthyl, and biphenyl portion may be unsubstituted or substituted with one or more X^1 group; and -C(O)-phenyl, -C(O)-naphthyl, -C(O)-biphenyl, wherein said phenyl, naphthyl, and biphenyl portion may be unsubstituted or substituted with one or more X^1 group. Non-limiting examples of -$S(O_2)$-R^{11} include, e.g., -$S(O_2)$-CH_3, -$S(O_2)$-CH_2CH_3, -$S(O_2)$-$CH_2CH_2CH_3$, -$S(O_2)$-$CH(CH_3)_2$, -$S(O_2)$-$CH_2CH_2CH_2CH_3$, -$S(O_2)$-$C(CH_3)_3$, -$S(O_2)$-$CH(CH_3)$-CH_2CH_3, -$S(O_2)$-$CH_2CH(CH_3)_2$, -$S(O_2)$-$CH_2CH_2CH_2CH_2CH_3$, -$S(O_2)$-$CH_2C(CH_3)_3$, etc., -$S(O_2)$-phenyl, -$S(O_2)$-naphthyl, -$S(O_2)$-biphenyl, wherein said phenyl, naphthyl, and biphenyl portion may be unsubstituted or substituted with one or more X^1 group. Non-limiting examples of -$S(O_2)$-R^{14} include, e.g., -$S(O_2)$-CH_3, -$S(O_2)$-CH_2CH_3, -$S(O_2)$-$CH_2CH_2CH_3$, -$S(O_2)$-$CH(CH_3)_2$, -$S(O_2)$-$CH_2CH_2CH_2CH_3$, -$S(O_2)$-$C(CH_3)_3$, -$S(O_2)$-$CH(CH_3)$-CH_2CH_3, -$S(O_2)$-$CH_2CH(CH_3)_2$, -$S(O_2)$-$CH_2CH_2CH_2CH_2CH_3$, -$S(O_2)$-$CH_2C(CH_3)_3$, etc., and -$S(O_2)$-cyclopropyl, -$S(O_2)$-cyclobutyl, -$S(O_2)$-cyclopentyl, -$S(O_2)$-cyclohexyl, -$S(O_2)$-cycloheptyl, wherein the cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl portion thereof may be unsubstituted or substituted with one or more X^4 group. R^8 also includes R^{14}, defined herein, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl, each of which may be unsubstituted or substituted with one or more X^4 group. The term -C(O)-$N(R^{18})_2$ includes, for example, -C(O)-NHR^{18}, where R^{18} is as defined below.

R^9 is selected from the group consisting of H and R^{11}, wherein R^{11} is defined as described herein.

R^{10} is selected from the group consisting of H, alkyl substituted with one or more –OH group, -alkylene-R^{12}, -alkylene-R^{13}, -alkylene-R^{14}, -C(O)-R^{14}, -alkylene-O-R^9, R^{14}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, and benzo-fused cycloalkyl. The terms "alkyl substituted with one or more –OH group" "-alkylene-R^{12}", and "–C(O)-R^{14}" are defined as described herein. Non-limiting examples of –alkylene-R^{13} include –alkylene-heteroaryls wherein the "alkylene' portion thereof includes, e.g., -CH_2-, -CH_2CH_2-, -$CH_2CH_2CH_2$-,

$-CH_2CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2CH_2CH_2-$, $-CH(CH_3)-$, $-CH(CH_3)CH_2-$, $-CH_2CH(CH_3)-$, $-CH(CH_2CH_3)-$, $-CH(CH_3)CH_2CH_2-$, $-CH_2CH(CH_3)CH_2-$, $-CH_2CH_2CH(CH_3)-$, $-CH(CH_2CH_3)CH_2-$, $-CH_2CH(CH_2CH_3)-$, $-CH(CH_2CH_2CH_3)-$, $-CH(CH_3)CH(CH_3)-$, and the "heteroaryl" portion thereof includes, e.g., azaindolyl, benzimidazolyl, benzofuranyl, benzoazaindolyl, benzothiophenyl, cinnolinyl, furanyl, furazanyl, indolyl, isoquinolyl, phthalazinyl, pyrazinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinolinyl, quinoxalinyl, quinazolinyl, thiophenyl, isoxazolyl, triazolyl, thiazolyl, thiadiazolyl, etc., each of which may be unsubstituted or substituted with one or more X^2 groups. Non-limiting examples of -alkylene-R^{14} include -alkylene-cycloalkyls wherein the "alkylene' portion thereof includes, e.g., $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2CH_2CH_2-$, $-CH(CH_3)-$, $-CH(CH_3)CH_2-$, $-CH_2CH(CH_3)-$, $-CH(CH_2CH_3)-$, $-CH(CH_3)CH_2CH_2-$, $-CH_2CH(CH_3)CH_2-$, $-CH_2CH_2CH(CH_3)-$, $-CH(CH_2CH_3)CH_2-$, $-CH_2CH(CH_2CH_3)-$, $-CH(CH_2CH_2CH_3)-$, $-CH(CH_3)CH(CH_3)-$, and the cycloalkyl portion thereof includes, e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclo[2.2.1]heptyl, adamantyl, etc. each of which may be unsubstituted or substituted with one or more X^4 groups. When R^{10} is R^{14}, non-limiting examples include e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclo[2.2.1]heptyl, adamantyl, etc. each of which may be unsubstituted or substituted with one or more X^4 groups. When R^{10} is unsubstituted heterocycloalkyl or heterocycloalkyl substituted with one or more X^3 groups, said heterocycloalkyl may include morpholinyl, piperazinyl, piperidinyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydrofuranyl, thiazolinyl, tetrahydropyranyl, etc. Non-limiting examples of benzo-fused cycloalkyls include the following structures:



etc. Although the structures above suggest that the benzo-fused cycloalkyl is bonded to the parent structure from the cycloalkyl portion of the group, it is contemplated that the benzo-fused cycloalkyl can be bonded to the parent

structure either from the cycloalkyl portion (i.e., from a saturated or sp^3 ring carbon, or alternatively from the "benzo" portion (i.e., from an unsaturated or sp^2 ring carbon).

R^{11} is selected from the group consisting of unsubstituted alkyl, alkyl substituted with one or more –OH groups, -alkylene-O-alkyl, -alkylene-O-aryl, unsubstituted aryl, and aryl substituted with one or more X^1 groups. The terms "alkyl", "alkylene" and "aryl" are as defined herein.

R^{12} is selected from the group consisting of unsubstituted aryl and aryl substituted with one or more X^1 groups, wherein the term "aryl" is as defined herein.

R^{13} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more X^2 groups. Non-limiting examples of suitable R^{13} groups include, e.g., azaindolyl, benzimidazolyl, benzofuranyl, benzoazaindolyl, benzothiophenyl, cinnolinyl, furanyl, furazanyl, indolyl, isoquinolyl, phthalazinyl, pyrazinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinolinyl, quinoxalinyl, quinazolinyl, thiophenyl, isoxazolyl, triazolyl, thiazolyl, thiadiazolyl, etc., each of which may be unsubstituted or substituted with one or more X^2 groups.

R^{14} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more X^4 groups. Non-limiting examples of suitable "alkyl" groups include those defined above. Non-limiting examples of suitable cycloalkyl groups include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, bicyclo[2.2.1]heptyl, adamantyl, etc. each of which may be unsubstituted or substituted with one or more X^4 groups.

Each R^{15} is independently selected from the group consisting of H, alkyl, alkenyl, -alkylene-R^{12}, -OH, and –O-alkenyl. Non-limiting examples of suitable alkyl, alkenyl, and -alkylene-R^{12} include those defined herein. Non-limiting examples of suitable –O-alkenyl groups include, for example, -O-CH=CH_2, -O-CH_2-CH=CH_2, -O-CH=CH-CH_3, -O-CH_2-C(CH_3)=CH_2, -O-CH_2-CH=$CHCH_3$, etc. There are 0, 1, or 2 R^{15} groups present in the compounds of Formula (I).

R^{16} is selected from the group consisting of R^9 and $-C(O)-R^{12}$. Non-limiting examples of suitable R^9 and $-C(O)-R^{12}$ groups are defined as disclosed herein.

R^{17} is selected from the group consisting of unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, -alkylene-R^{12}, -O-R^9, and R^{12}. Non-limiting examples of suitable heterocycloalkyl, -alkylene-R^{12}, -O-R^9, and R^{12} groups are defined as disclosed herein.

Each R^{18} is independently selected from the group consisting of H, R^{12}, and R^{14}, where R^{12} and R^{14} are as defined above.

Each R^{19} is selected from the group consisting of H and R^{21}, wherein R^{21} is defined as described herein.

Each R^{20} is selected from the group consisting of H, alkyl substituted with one or more -OH or –O-alkyl groups, -alkylene-R^{22}, -alkylene-R^{23}, -alkylene-R^{24}, -C(O)-R^{24}, -alkylene-O-R^{19}, R^{24}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, and benzo-fused cycloalkyl. Non-limiting examples of suitable alkyl substituted with one or more -OH or –O-alkyl groups, -alkylene-R^{22}, -alkylene-R^{23}, -alkylene-R^{24}, -C(O)-R^{24}, -alkylene-O-R^{19}, R^{24}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, and benzo-fused cycloalkyl.groups are defined as disclosed herein.

R^{21} is selected from the group consisting of unsubstituted alkyl, alkyl substituted with one or more –OH groups, -alkylene-O-alkyl, -alkylene-O-aryl, unsubstituted aryl, and aryl substituted with one or more W^1 groups. Non-limiting examples of suitable unsubstituted alkyl, alkyl substituted with one or more –OH groups, -alkylene-O-alkyl, -alkylene-O-aryl, unsubstituted aryl, and aryl substituted with one or more W^1 groups are described herein.

R^{22} is selected from the group consisting of unsubstituted aryl and aryl substituted with one or more W^1 groups. Non-limiting examples of suitable unsubstituted aryl and aryl substituted with one or more W^1 groups are described herein.

R^{23} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more W^2 groups. Non-limiting

examples of suitable unsubstituted heteroaryl and heteroaryl substituted with one or more W^2 groups are described herein.

R^{24} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more W^4 groups. Non-limiting examples of suitable alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more W^4 groups are described herein.

W^1 is independently selected from the group consisting of halogen, -CN, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, and –O-alkyl. Non-limiting examples of suitable halogen, -CN, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, and –O-alkyl are described herein.

W^2 is independently selected from the group consisting of halogen, unsubstituted aryl, and aryl substituted with one or more Z groups. Non-limiting examples of suitable halogens include F, Cl, and Br. Suitable examples of aryl groups include, for example, those described herein.

W^3 is -C(O)-O-alkyl. Non-limiting examples of suitable -C(O)-O-alkyl groups include those defined herein. In addition two W^3 groups together with the ring carbon atom to which they are attached form a carbonyl group. It is contemplated that a heterocycloalkyl group may be independently substituted with one or more –C(O)-O-alkyl groups and/or one or more carbonyl groups (i.e., one, two, three, four, or five W^3 groups).

W^4 is independently halogen or alkyl. Non-limiting examples of suitable halogens include F, Cl, and Br. Non-limiting examples of suitable alkyl groups include those described herein.


Ar^1 and Ar^2 are independently selected from the group consisting of R^{12} and R^{13}. Non-limiting examples of suitable R^{12} and R^{13} groups are defined as disclosed herein.


Aryls substituted with one or more X^1 groups include, for example mono-substituted, di-substituted, tri-substituted, tetra-substituted aryls, etc, wherein each of the substituents are independently selected from X^1. Non-limiting examples include, for example, chlorophenyl, dichlorophenyl, bromophenyl, dibromophenyl, bromo-chlorophenyl, fluorophenyl,

difluorophenyl, chloro-fluorophenyl, bromo-fluorophenyl, cyanophenyl, biphenyl, chlorobiphenyl, dichlorobiphenyl, etc. Similarly, heteroaryls substituted with one or more X^2 groups include mono-substituted, di-substituted, tri-substituted, tetra-substituted heteroaryls, etc, wherein each of the substituents are independently selected from X^2. Suitable aryls and heteroaryls include any of those disclosed herein.


X^1 is independently selected from the group consisting of halogen, -CN, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-$C(O)N(R^{20})_2$, -O-alkylene-O-R^{19}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl -O-alkylene-O-alkylene-OR^{19}, -O-alkylene-$C(O)R^{24}$, -O-alkylene-$C(O)OR^{19}$, and –O-alkyl. Non-limiting examples of suitable halogens includes, for example F, Cl, and Br. Non-limiting examples of suitable -O-S(O)$_2$-haloalkyls include -O-S(O)$_2$-CH_2F, -O-S(O)$_2$-CHF_2, -O-S(O)$_2$-CF_3, -O-S(O)$_2$-CH_2CF_3, -O-S(O)$_2$-CF_2CF_3, -O-S(O)$_2$-CH_2Cl, -O-S(O)$_2$-CH_2Br, etc. Non-limiting examples of suitable –O-alkyl groups include those described herein.


Non-limiting examples of suitable X^1 groups are independently selected from the group consisting of –OCH_3, -OH, -OTf, -CN, -OCH_2CH_3, -$OCH(CH_3)_2$,



































and


Similarly, X^2 is selected from the group consisting of halogen, -CN, unsubstituted aryl, and aryl substituted with one or more Z groups. Non-limiting examples of suitable halogens include F, Cl, and Br. Suitable examples of aryl groups include, for example, those described herein.

X^3 is -C(O)-O-alkyl. Non-limiting examples of suitable -C(O)-O-alkyl groups include those defined herein. In addition two X^3 groups together with the ring carbon atom to which they are attached form a carbonyl group. It is contemplated that a heterocycloalkyl group may be independently substituted with one or more –C(O)-O-alkyl groups and/or one or more carbonyl groups (i.e., one, two, three, four, or five X^3 groups).

X^4 is independently halogen or alkyl. Non-limiting examples of suitable halogens include F, Cl, and Br. Non-limiting examples of suitable alkyl groups include those described herein.

U is independently selected from the group consisting of –OH, -O-alkyl and halogen. Non-limiting examples of suitable alkyl and halogen groups include those described herein.

Z is selected from the group consisting of halogen, alkyl, and –CN. Non-limiting examples of halogen and alkyl include those defined herein.

As used above, and throughout this disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

"Patient" includes both human and animals.

"Mammal" means humans and other mammalian animals.

"Alkyl" means an aliphatic hydrocarbon group which may be straight or branched and comprising about 1 to about 20 carbon atoms in the chain. Preferred alkyl groups contain about 1 to about 12 carbon atoms in the chain. More preferred alkyl groups contain about 1 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkyl chain. "Lower alkyl" means a group having about 1 to about 6 carbon atoms in the chain which may be straight or branched. The term "substituted alkyl" means that the alkyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halogen, alkyl, aryl, cycloalkyl, cyano, hydroxy, alkoxy, alkylthio, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)$_2$, carboxy and –C(O)O-alkyl. Non-limiting examples of suitable alkyl groups include methyl, ethyl, n-propyl, isopropyl and t-butyl.

"Alkylene" means a divalent group obtained by removal of a hydrogen atom from an alkyl group that is defined above. Non-limiting examples of alkylene include methylene, ethylene (i.e., -CH_2CH_2- or -$CH(CH_3)$-) and propylene (e.g., including –$CH_2CH_2CH_2$- and –$CH(CH_3)CH_2$-).

"Alkenyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkenyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 6 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkenyl chain. "Lower alkenyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. The term "substituted alkenyl" means that the alkenyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of halogen, alkyl. aryl, cycloalkyl, cyano, alkoxy and –S(alkyl). Non-limiting examples of suitable alkenyl groups include ethenyl, propenyl, n-butenyl, 3-methylbut-2-enyl, n-pentenyl, octenyl and decenyl.

"Alkenylene" means a divalent group obtained by removal of a hydrogen from an alkenyl group that is defined above. Non-limiting examples of alkenylene include –CH=CH-, -C(CH_3)=CH-, and –CH=CHCH_2-.

"Alkynyl" means an aliphatic hydrocarbon group containing at least one carbon-carbon triple bond and which may be straight or branched and comprising about 2 to about 15 carbon atoms in the chain. Preferred alkynyl groups have about 2 to about 12 carbon atoms in the chain; and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl, are attached to a linear alkynyl chain. "Lower alkynyl" means about 2 to about 6 carbon atoms in the chain which may be straight or branched. Non-limiting examples of suitable alkynyl groups include ethynyl, propynyl, 2-butynyl and 3-methylbutynyl. The term "substituted alkynyl" means that the alkynyl group may be substituted by one or more substituents which may be the same or different, each substituent being independently selected from the group consisting of alkyl, aryl and cycloalkyl.

"Alkynylene" means a difunctional group obtained by removal of a hydrogen from an alkynyl group that is defined above. Non-limiting examples of alkenylene include -C≡C- and –CH_2C≡C-.

"Aryl" means an aromatic monocyclic or multicyclic ring system comprising about 6 to about 14 carbon atoms, preferably about 6 to about 10 carbon atoms. The aryl group can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Non-limiting examples of suitable aryl groups include phenyl and naphthyl.

"Heteroaryl" means an aromatic monocyclic or multicyclic ring system comprising about 5 to about 14 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the ring atoms is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. Preferred heteroaryls contain about 5 to about 6 ring atoms. The "heteroaryl" can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The prefix aza, oxa or thia before the heteroaryl root name means that at least a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. A nitrogen atom of a

heteroaryl can be optionally oxidized to the corresponding N-oxide. Non-limiting examples of suitable heteroaryls include pyridyl, pyrazinyl, furanyl, thienyl, pyrimidinyl, pyridone (including N-substituted pyridones), isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, pyrazolyl, furazanyl, pyrrolyl, pyrazolyl, triazolyl, 1,2,4-thiadiazolyl, pyrazinyl, pyridazinyl, quinoxalinyl, phthalazinyl, oxindolyl, imidazo[1,2-a]pyridinyl, imidazo[2,1-b]thiazolyl, benzofurazanyl, indolyl, azaindolyl, benzimidazolyl, benzothienyl, quinolinyl, imidazolyl, thienopyridyl, quinazolinyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, isoquinolinyl, benzoazaindolyl, 1,2,4-triazinyl, benzothiazolyl and the like. The term "heteroaryl" also refers to partially saturated heteroaryl moieties such as, for example, tetrahydroisoquinolyl, tetrahydroquinolyl, indazolyl, and the like, in which there is at least one aromatic ring.

"Aralkyl", "arylalkyl", or "–alkylene-aryl" means an aryl-alkyl- group in which the aryl and alkyl are as previously described. Preferred aralkyls comprise a lower alkyl group. Non-limiting examples of suitable aralkyl groups include benzyl, 2-phenethyl and naphthalenylmethyl. The bond to the parent moiety is through the alkyl.

"Alkylaryl" means an alkyl-aryl- group in which the alkyl and aryl are as previously described. Preferred alkylaryls comprise a lower alkyl group. A non-limiting example of a suitable alkylaryl group is tolyl. The bond to the parent moiety is through the aryl.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system comprising about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms. Preferred cycloalkyl rings contain about 5 to about 7 ring atoms. The cycloalkyl can be optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined above. Non-limiting examples of suitable monocyclic cycloalkyls include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. Non-limiting examples of suitable multicyclic cycloalkyls include 1-decalinyl, norbornyl, adamantyl and the like, as well as partially saturated species such as, for example, indanyl, tetrahydronaphthyl and the like.


"Cycloalkylalkyl" means a cycloalkyl moiety as defined above linked via an alkyl moiety (defined above) to a parent core. Non-limiting examples of

suitable cycloalkylalkyls include cyclohexylmethyl, adamantylmethyl and the like.

"Halogen" or "halo" means fluorine, chlorine, bromine, or iodine. Preferred are fluorine, chlorine and bromine.

"Haloalkyl" means an alkyl group as defined above wherein one or more hydrogen atoms on the alkyl is replaced by a halo group defined above.


"Ring system substituent" means a substituent attached to an aromatic or non-aromatic ring system which, for example, replaces an available hydrogen on the ring system. Ring system substituents may be the same or different, each being independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, aralkyl, alkylaryl, heteroaralkyl, heteroarylalkenyl, heteroarylalkynyl, alkylheteroaryl, hydroxy, hydroxyalkyl, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halogen, nitro, cyano, carboxy, alkoxycarbonyl, aryloxycarbonyl, aralkoxycarbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, cycloalkyl, heterocyclyl, -C(=N-CN)-NH_2, -C(=NH)-NH_2, -C(=NH)-NH(alkyl), Y_1Y_2N-, Y_1Y_2N-alkyl-, $Y_1Y_2NC(O)$-, $Y_1Y_2NSO_2$- and $-SO_2NY_1Y_2$, wherein Y_1 and Y_2 can be the same or different and are independently selected from the group consisting of hydrogen, alkyl, aryl, cycloalkyl, and aralkyl. "Ring system substituent" may also mean a single moiety which simultaneously replaces two available hydrogens on two adjacent carbon atoms (one H on each carbon) on a ring system. Examples of such moiety are methylenedioxy, ethylenedioxy, -$C(CH_3)_2$- and the like which form moieties such as, for example:

 and .

"Heterocyclyl" means a monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Heterocyclyls may be completely saturated, partially unsaturated, or aromatic. Aromatic heterocyclyls are termed "heteroaryl", as defined above. Preferred

heterocyclyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocyclyl root name means that at least a nitrogen, oxygen or sulfur atom respectively is present as a ring atom. Any -NH in a heterocyclyl ring may exist protected such as, for example, as an -N(Boc), -N(CBz), -N(Tos) group and the like; such protections are also considered part of this invention. The heterocyclyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocyclyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocyclyl rings include saturated heterocyclyls, for example piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,4-dioxanyl, tetrahydrofuranyl, tetrahydrothiophenyl, lactams, lactones, isoxazolyl, 1,2,4-triazolyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-triazolyl and the like. Non-limiting examples of partially unsaturated monocyclic heterocyclyl rings include, for example, thiazolinyl, and the like.

It should be noted that in hetero-atom containing ring systems of this invention, there are no hydroxyl groups on carbon atoms adjacent to a N, O or S, as well as there are no N or S groups on carbon adjacent to another heteroatom. Thus, for example, in the ring:



there is no -OH attached directly to carbons marked 2 and 5.

It should also be noted that the compounds of the present invention include tautomers of the compounds of Formula (I).

"Heterocycloalkyl" means a non-aromatic saturated monocyclic or multicyclic ring system comprising about 3 to about 10 ring atoms, preferably about 5 to about 10 ring atoms, in which one or more of the atoms in the ring system is an element other than carbon, for example nitrogen, oxygen or sulfur, alone or in combination. There are no adjacent oxygen and/or sulfur atoms present in the ring system. Preferred heterocycloalkyls contain about 5 to about 6 ring atoms. The prefix aza, oxa or thia before the heterocycloalkyl root name means that at least a nitrogen, oxygen or sulfur atom respectively

is present as a ring atom. The heterocycloalkyl can be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen or sulfur atom of the heterocycloalkyl can be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Non-limiting examples of suitable monocyclic heterocycloalkyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, 1,3-dioxolanyl, tetrahydrofuranyl, tetrahydrothiophenyl and the like.

"Heteroaralkyl" means a heteroaryl-alkyl- group in which the heteroaryl and alkyl are as previously described. Preferred heteroaralkyls contain a lower alkyl group. Non-limiting examples of suitable aralkyl groups include pyridylmethyl, and quinolin-3-ylmethyl. The bond to the parent moiety is through the alkyl.

"Hydroxyalkyl" means a HO-alkyl- group in which alkyl is as previously defined. Preferred hydroxyalkyls contain lower alkyl. Non-limiting examples of suitable hydroxyalkyl groups include hydroxymethyl and 2-hydroxyethyl.

"Acyl" means an H-C(O)-, alkyl-C(O)- or cycloalkyl-C(O)-, group in which the various groups are as previously described. The bond to the parent moiety is through the carbonyl. Preferred acyls contain a lower alkyl. Non-limiting examples of suitable acyl groups include formyl, acetyl and propanoyl.

"Aroyl" means an aryl-C(O)- group in which the aryl group is as previously described. The bond to the parent moiety is through the carbonyl. Non-limiting examples of suitable groups include benzoyl and 1- naphthoyl.

"Alkoxy" means an alkyl-O- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkoxy groups include methoxy, ethoxy, n-propoxy, isopropoxy and n-butoxy. The bond to the parent moiety is through the ether oxygen.

"Aryloxy" means an aryl-O- group in which the aryl group is as previously described. Non-limiting examples of suitable aryloxy groups include phenoxy and naphthoxy. The bond to the parent moiety is through the ether oxygen.

"Alkylthio" means an alkyl-S- group in which the alkyl group is as previously described. Non-limiting examples of suitable alkylthio groups include methylthio and ethylthio. The bond to the parent moiety is through the sulfur.

"Arylthio" means an aryl-S- group in which the aryl group is as previously described. Non-limiting examples of suitable arylthio groups include phenylthio and naphthylthio. The bond to the parent moiety is through the sulfur.

"Aralkylthio" means an aralkyl-S- group in which the aralkyl group is as previously described. Non-limiting example of a suitable aralkylthio group is benzylthio. The bond to the parent moiety is through the sulfur.

"Alkoxycarbonyl" means an alkyl-O-CO- group. Non-limiting examples of suitable alkoxycarbonyl groups include methoxycarbonyl and ethoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aryloxycarbonyl" means an aryl-O-C(O)- group. Non-limiting examples of suitable aryloxycarbonyl groups include phenoxycarbonyl and naphthoxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Aralkoxycarbonyl" means an aralkyl-O-C(O)- group. Non-limiting example of a suitable aralkoxycarbonyl group is benzyloxycarbonyl. The bond to the parent moiety is through the carbonyl.

"Alkylsulfonyl" means an alkyl-S(O_2)- group. Preferred groups are those in which the alkyl group is lower alkyl. The bond to the parent moiety is through the sulfonyl.

"Arylsulfonyl" means an aryl-S(O_2)- group. The bond to the parent moiety is through the sulfonyl.

The term "substituted" means that one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency under the existing circumstances is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. By "stable compound' or "stable structure" is meant a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

The term "optionally substituted" means optional substitution with the specified groups, radicals or moieties.

The term "purified", "in purified form" or "in isolated and purified form" for a compound refers to the physical state of said compound after being

isolated from a synthetic process or natural source or combination thereof. Thus, the term "purified", "in purified form" or "in isolated and purified form" for a compound refers to the physical state of said compound after being obtained from a purification process or processes described herein or well known to the skilled artisan, in sufficient purity to be characterizable by standard analytical techniques described herein or well known to the skilled artisan.

It should also be noted that any carbon as well as heteroatom with unsatisfied valences in the text, schemes, examples and Tables herein is assumed to have the sufficient number of hydrogen atom(s) to satisfy the valences.

When a functional group in a compound is termed "protected", this means that the group is in modified form to preclude undesired side reactions at the protected site when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for example, T. W. Greene *et al*, *Protective Groups in Organic Synthesis* (1991), Wiley, New York.

When any variable (e.g., aryl, heterocycle, R^9, etc.) occurs more than one time in any constituent or in Formula I, its definition on each occurrence is independent of its definition at every other occurrence.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

Prodrugs and solvates of the compounds of the invention are also contemplated herein. The term "prodrug", as employed herein, denotes a compound that is a drug precursor which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of Formula I or a salt and/or solvate thereof. A discussion of prodrugs is provided in T. Higuchi and V. Stella, *Pro-drugs as Novel Delivery Systems* (1987) 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design,* (1987) Edward B. Roche, ed., American

Pharmaceutical Association and Pergamon Press, both of which are incorporated herein by reference thereto.

"Solvate" means a physical association of a compound of this invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanolates, methanolates, and the like. "Hydrate" is a solvate wherein the solvent molecule is H_2O.

"Effective amount" or "therapeutically effective amount" is meant to describe an amount of compound or a composition of the present invention effective in inhibiting the diseases or conditions noted below, and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect.

The compounds of Formula (I) can form salts which are also within the scope of this invention. Reference to a compound of Formula (I) herein is understood to include reference to salts thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of Formula (I) contains both a basic moiety, such as, but not limited to a pyridine or imidazole, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful. Salts of the compounds of the Formula (I) may be formed, for example, by reacting a compound of Formula (I) with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides,

lactates, maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates,) and the like. Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by P. Stahl *et al*, Camille G. (eds.) *Handbook of Pharmaceutical Salts. Properties, Selection and Use.* (2002) Zurich: Wiley-VCH; S. Berge *et al*, *Journal of Pharmaceutical Sciences* (1977) 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33 201-217; Anderson *et al*, *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as dicyclohexylamines, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, and dibutyl sulfates), long chain halides (e.g. decyl, lauryl, and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

All such acid salts and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.

Pharmaceutically acceptable esters of the present compounds include the following groups: (1) carboxylic acid esters obtained by esterification of the hydroxy groups, in which the non-carbonyl moiety of the carboxylic acid portion of the ester grouping is selected from straight or branched chain alkyl (for example, acetyl, n-propyl, t-butyl, or n-butyl), alkoxyalkyl (for example, methoxymethyl), aralkyl (for example, benzyl), aryloxyalkyl (for example, phenoxymethyl), aryl (for example, phenyl optionally substituted with, for

example, halogen, (C_1-C_4)alkyl, or (C_1-C_4)alkoxy or amino); (2) sulfonate esters, such as alkyl- or aralkylsulfonyl (for example, methanesulfonyl); (3) amino acid esters (for example, L-valyl or L-isoleucyl); (4) phosphonate esters and (5) mono-, di- or triphosphate esters. The phosphate esters may be further esterified by, for example, a (C_1-C_{20}) alcohol or reactive derivative thereof, or by a 2,3-di-(C_6-C_{24})acyl glycerol.

One or more compounds of the invention may also exist as, or optionally converted to, a solvate. Preparation of solvates is generally known. Thus, for example, M. Caira *et al*, *J. Pharmaceutical Sci.*, 93(3), 601-611 (2004) describe the preparation of the solvates of the antifungal fluconazole in ethyl acetate as well as from water. Similar preparations of solvates, hemisolvate, hydrates and the like are described by E. C. van Tonder *et al*, *AAPS PharmSciTech.*, 5(1), article 12 (2004); and A. L. Bingham *et al*, *Chem. Commun.*, 603-604 (2001). A typical, non-limiting, process involves dissolving the inventive compound in desired amounts of the desired solvent (organic or water or mixtures thereof) at a higher than ambient temperature, and cooling the solution at a rate sufficient to form crystals which are then isolated by standard methods. Analytical techniques such as, for example I. R. spectroscopy, show the presence of the solvent (or water) in the crystals as a solvate (or hydrate).

Compounds of Formula (I), and salts, solvates, esters and prodrugs thereof, may exist in their tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds (including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this invention, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for

example, as racemates or with all other, or other selected, stereoisomers. The chiral centers of the present invention can have the S or R configuration as defined by the *IUPAC* 1974 Recommendations. The use of the terms "salt", "solvate", "ester", "prodrug" and the like, is intended to equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or prodrugs of the inventive compounds.

Polymorphic forms of the compounds of Formula I, and of the salts, solvates, esters and prodrugs of the compounds of Formula I, are intended to be included in the present invention.

The compounds of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof according to the invention have pharmacological properties; in particular, the compounds of Formula (I) can be selective CB_1 antagonists. The term "selective" means that the compounds of Formula (I) bind to the CB_1 receptor more strongly than to other cannabinoid receptors.

The compounds of Formula (I) of the present invention, or pharmaceutically acceptable salts, solvates, or esters thereof are useful in treating diseases or conditions including obesity, metabolic disorders, addiction, diseases of the central nervous system, cardiovascular disorders, respiratory disorders, gastrointestinal disorders, achieving weight reduction, lowering waist circumference, treating dyslipidemia, insulin sensitivity, diabetes mellitus, hypertriglyceridemia, eating disorders, alcoholism, inflammation, psychiatric disorders, migraine, nicotine dependence, Parkinson's disease, psychosis, schizophrenia, sleep disorders, attention deficit hyperactivity disorder, male sexual dysfunction, premature ejaculation, premenstrual syndrome, seizure, epilepsy and convulsion, non-insulin dependent diabetes, dementia, major depressive disorder, bulimia nervosa, drug dependence, septic shock, cognitive disorder, endocrine disorders, eczema, emesis, allergy, glaucoma, hemorrhagic shock, hypertension, angina, thrombosis, atherosclerosis, restenosis, hypertension, acute coronary syndrome, angina pectoris, arrhythmia, heart failure, cerebral ischemia, stroke, myocardial infarction, glomerulonephritis, thrombotic and thromboembolytic stroke, peripheral vascular diseases, neurodegenerative disease, osteoporosis, pulmonary disease, autoimmune disease,

hypotension, arthropathy, cancer, demyelinating diseases, Alzheimer's disease, hypoactive sexual desire disorder, bipolar disorder, hyperlipidemia, hypertension, narcotic dependence, Huntington's chorea, pain, multiple sclerosis, anxiety disorder, bone disorders, Paget's disease, rheumatoid arthritis, ulcerative colitis, irritable bowel syndrome, and inflammatory bowel diseases.

The term "pharmaceutical composition" is also intended to encompass both the bulk composition and individual dosage units comprised of more than one (e.g., two) pharmaceutically active agents such as, for example, a compound of the present invention and an additional agent selected from the lists of the additional agents described herein, along with any pharmaceutically inactive excipients. The bulk composition and each individual dosage unit can contain fixed amounts of the afore-said "more than one pharmaceutically active agents". The bulk composition is material that has not yet been formed into individual dosage units. An illustrative dosage unit is an oral dosage unit such as tablets, pills and the like. Similarly, the herein-described method of treating a patient by administering a pharmaceutical composition of the present invention is also intended to encompass the administration of the afore-said bulk composition and individual dosage units.

The compounds of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, can be administered in any suitable form, e.g., alone, or in combination with a pharmaceutically acceptable carrier, excipient or diluent in a pharmaceutical composition, according to standard pharmaceutical practice. The compounds of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, can be administered orally or parenterally, including intravenous, intramuscular, interperitoneal, subcutaneous, rectal, or topical routes of administration.

Pharmaceutical compositions comprising at least one compound of Formula I, or a pharmaceutically acceptable salt, solvate, or ester thereof can be in a form suitable for oral administration, e.g., as tablets, troches, capsules, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, syrups, or elixirs. Oral compositions may be prepared by any

conventional pharmaceutical method, and may also contain sweetening agents, flavoring agents, coloring agents, and preserving agents.

The amount of compound of Formula (I), or a pharmaceutically acceptable salt, solvate, or ester thereof, administered to a patient can be determined by a physician based on the age, weight, and response of the patient, as well as by the severity of the condition treated. For example, the amount of compound of Formula I, or a pharmaceutically acceptable salt, solvate, or ester thereof, administered to the patient can range from about 0.1 mg/kg body weight per day to about 60 mg/kg/d, preferably about 0.5 mg/kg/d to about 40 mg/kg/d.

The compounds of Formula I, or pharmaceutically acceptable salts, solvates, or esters thereof, can also be administered in combination with other therapeutic agents. For example one or more compounds of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, can be administered with one or more additional cholesterol lowering agents.

A non-limiting list of cholesterol lowering agents useful in the present invention include HMG CoA reductase inhibitor compounds such as lovastatin (for example MEVACOR® which is available from Merck & Co.), simvastatin (for example ZOCOR® which is available from Merck & Co.), pravastatin (for example PRAVACHOL® which is available from Bristol Meyers Squibb), atorvastatin, fluvastatin, cerivastatin, CI-981, rivastatin (sodium 7-(4-fluorophenyl)-2,6-diisopropyl-5-methoxymethylpyridin-3-yl)-3,5-dihydroxy-6-heptanoate), rosuvastatin calcium (CRESTOR® from AstraZeneca Pharmaceuticals), pitavastatin (such as NK-104 of Negma Kowa of Japan); HMG CoA synthetase inhibitors, for example L-659,699 ((E,E)-11-[3'R-(hydroxy-methyl)-4'-oxo-2'R-oxetanyl]-3,5,7R-trimethyl-2,4-undecadienoic acid); squalene synthesis inhibitors, for example squalestatin 1; squalene epoxidase inhibitors, for example, NB-598 ((E)-N-ethyl-N-(6,6-dimethyl-2-hepten-4-ynyl)-3-[(3,3'-bithiophen-5-yl)methoxy]benzene-methanamine hydrochloride); sterol (e.g., cholesterol) biosynthesis inhibitors such as DMP-565; nicotinic acid derivatives (e.g., compounds comprising a pyridine-3-carboxylate structure or a pyrazine-2-carboxylate structure, including acid forms, salts, esters, zwitterions and tautomers) such as niceritrol,


nicofuranose and acipimox (5-methyl pyrazine-2-carboxylic acid 4-oxide); clofibrate; gemfibrazol; bile acid sequestrants such as cholestyramine (a styrene-divinylbenzene copolymer containing quaternary ammonium cationic groups capable of binding bile acids, such as QUESTRAN® or QUESTRAN LIGHT® cholestyramine which are available from Bristol-Myers Squibb), colestipol (a copolymer of diethylenetriamine and 1-chloro-2,3-epoxypropane, such as COLESTID® tablets which are available from Pharmacia), colesevelam hydrochloride (such as WelChol® Tablets (poly(allylamine hydrochloride) cross-linked with epichlorohydrin and alkylated with 1-bromodecane and (6-bromohexyl)-trimethylammonium bromide) which are available from Sankyo), water soluble derivatives such as 3,3-ioene, N-(cycloalkyl) alkylamines and poliglusam, insoluble quaternized polystyrenes, saponins and mixtures thereof; inorganic cholesterol sequestrants such as bismuth salicylate plus montmorillonite clay, aluminum hydroxide and calcium carbonate antacids; ileal bile acid transport ("IBAT") inhibitors (or apical sodium co-dependent bile acid transport ("ASBT") inhibitors) such as benzothiepines, for example the therapeutic compounds comprising a 2,3,4,5-tetrahydro-1-benzothiepine 1,1-dioxide structure such as are disclosed in PCT Patent Application WO 00/38727 which is incorporated herein by reference; AcylCoA:Cholesterol O-acyltransferase ("ACAT") Inhibitors such as avasimibe ([[2,4,6-tris(1-methylethyl)phenyl]acetyl]sulfamic acid, 2,6-bis(1-methylethyl)phenyl ester, formerly known as CI-1011), HL-004, lecimibide (DuP-128) and CL-277082 (*N*-(2,4-difluorophenyl)-*N*-[[4-(2,2-dimethylpropyl)phenyl]methyl]-*N*-heptylurea), and the compounds described in P. Chang et al., "Current, New and Future Treatments in Dyslipidaemia and Atherosclerosis", Drugs 2000 Jul;60(1); 55-93, which is incorporated by reference herein; Cholesteryl Ester Transfer Protein ("CETP") Inhibitors such as those disclosed in PCT Patent Application No. WO 00/38721 and U.S. Patent No. 6,147,090, which are incorporated herein by reference; probucol or derivatives thereof, such as AGI-1067 and other derivatives disclosed in U.S. Patents Nos. 6,121,319 and 6,147,250, herein incorporated by reference; low-density lipoprotein (LDL) receptor activators such as HOE-402, an imidazolidinyl-pyrimidine derivative that directly stimulates LDL receptor

activity, described in M. Huettinger et al., "Hypolipidemic activity of HOE-402 is Mediated by Stimulation of the LDL Receptor Pathway", Arterioscler. Thromb. 1993; 13:1005-12, herein incorporated by reference; fish oils containing Omega 3 fatty acids (3-PUFA); natural water soluble fibers, such as psyllium, guar, oat and pectin; plant stanols and/or fatty acid esters of plant stanols, such as sitostanol ester used in BENECOL® margarine; nicotinic acid receptor agonists (e.g., agonists of the HM74 and HM74A receptor which receptor is described in US 2004/0142377, US 2005/0004178, US 2005/0154029, US 6902902, WO 2004/071378, WO 2004/071394, WO 01/77320, US 2003/0139343, WO 01/94385, WO 2004/083388, US 2004/254224, US 2004/0254224, US 2003/0109673 and WO 98/56820) for example those described in WO 2004/033431, WO 2005/011677, WO 2005/051937, US 2005/0187280, US 2005/0187263, WO 2005/077950, WO 2005/016867, and WO 2005/016870; and the substituted azetidinone or substituted β-lactam sterol absorption inhibitors discussed in detail below.

As used herein, "sterol absorption inhibitor" means a compound capable of inhibiting the absorption of one or more sterols, including but not limited to cholesterol, phytosterols (such as sitosterol, campesterol, stigmasterol and avenosterol), 5α-stanols (such as cholestanol, 5α-campestanol, 5α-sitostanol), and/or mixtures thereof, when administered in a therapeutically effective (sterol and/or 5α-stanol absorption inhibiting) amount to a mammal or human.

Substituted Azetidinones of Formula (A)

In one embodiment, substituted azetidinones useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (A) below:



(A)

or pharmaceutically acceptable salts, solvates, or esters of the compounds of Formula (A), wherein, in Formula (A) above:

Ar^1 and Ar^2 are independently selected from the group consisting of aryl and R^4-substituted aryl;

Ar^3 is aryl or R^5-substituted aryl;

X, Y and Z are independently selected from the group consisting of $-CH_2-$, -CH(lower alkyl)- and $-C(\text{lower alkyl})_2-$;

R and R^2 are independently selected from the group consisting of $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$ and $-OC(O)NR^6R^7$;

R^1 and R^3 are independently selected from the group consisting of hydrogen, lower alkyl and aryl;

q is 0 or 1; r is 0 or 1; m, n and p are independently selected from 0, 1, 2, 3 or 4; provided that at least one of q and r is 1, and the sum of m, n, p, q and r is 1, 2, 3, 4, 5 or 6; and provided that when p is 0 and r is 1, the sum of m, q and n is 1, 2, 3, 4 or 5;

R^4 is 1-5 substituents independently selected from the group consisting of lower alkyl, $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$, $-O(CH_2)_{1-5}OR^6$, $-OC(O)NR^6R^7$, $-NR^6R^7$, $-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6SO_2R^9$, $-C(O)OR^6$, $-C(O)NR^6R^7$, $-C(O)R^6$, $-S(O)_2NR^6R^7$, $S(O)_{0-2}R^9$, $-O(CH_2)_{1-10}-C(O)OR^6$, $-O(CH_2)_{1-10}CONR^6R^7$, -(lower alkylene)$COOR^6$, $-CH=CH-C(O)OR^6$, $-CF_3$, -CN, $-NO_2$ and halogen;

R^5 is 1-5 substituents independently selected from the group consisting of $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$, $-O(CH_2)_{1-5}OR^6$, $-OC(O)NR^6R^7$, $-NR^6R^7$, $-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_2R^9$, $-C(O)OR^6$, $-C(O)NR^6R^7$, $-C(O)R^6$, $-SO_2NR^6R^7$, $S(O)_{0-2}R^9$, $-O(CH_2)_{1-10}-C(O)OR^6$, $-O(CH_2)_{1-10}C(O)NR^6R^7$, -(lower alkylene)$C(O)OR^6$ and $-CH=CH-C(O)OR^6$;

R^6, R^7 and R^8 are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and

R^9 is lower alkyl, aryl or aryl-substituted lower alkyl.

Preferably, R^4 is 1-3 independently selected substituents, and R^5 is preferably 1-3 independently selected substituents.

Certain compounds useful in the therapeutic compositions or combinations of the invention may have at least one asymmetrical carbon atom and therefore all isomers, including enantiomers, diastereomers,

stereoisomers, rotamers, tautomers and racemates of the compounds of Formula A-M (where they exist) are contemplated as being part of this invention. The invention includes d and l isomers in both pure form and in admixture, including racemic mixtures. Isomers can be prepared using conventional techniques, either by reacting optically pure or optically enriched starting materials or by separating isomers of a compound of the Formulae A-M. Isomers may also include geometric isomers, e.g., when a double bond is present.

Those skilled in the art will appreciate that for some of the compounds of the Formulae A-M, one isomer may show greater pharmacological activity than other isomers.

Preferred compounds of Formula (A) are those in which Ar^1 is phenyl or R^4-substituted phenyl, more preferably (4-R^4)-substituted phenyl. Ar^2 is preferably phenyl or R^4-substituted phenyl, more preferably (4-R^4)-substituted phenyl. Ar^3 is preferably R^5-substituted phenyl, more preferably (4-R^5)-substituted phenyl. When Ar^1 is (4-R^4)-substituted phenyl, R^4 is preferably a halogen. When Ar^2 and Ar^3 are R^4- and R^5-substituted phenyl, respectively, R^4 is preferably halogen or -OR^6 and R^5 is preferably -OR^6, wherein R^6 is lower alkyl or hydrogen. Especially preferred are compounds wherein each of Ar^1 and Ar^2 is 4-fluorophenyl and Ar^3 is 4-hydroxyphenyl or 4-methoxyphenyl.

X, Y and Z are each preferably -CH_2-. R^1 and R^3 are each preferably hydrogen. R and R^2 are preferably -OR^6 wherein R^6 is hydrogen, or a group readily metabolizable to a hydroxyl (such as -$OC(O)R^6$, -$OC(O)OR^9$ and -$OC(O)NR^6R^7$, defined above).

The sum of m, n, p, q and r is preferably 2, 3 or 4, more preferably 3. Preferred are compounds of Formula (A) wherein m, n and r are each zero, q is 1 and p is 2.


Also preferred are compounds of Formula (A) in which p, q and n are each zero, r is 1 and m is 2 or 3. More preferred are compounds wherein m, n and r are each zero, q is 1, p is 2, Z is -CH_2- and R is -OR^6, especially when R^6 is hydrogen.

Also more preferred are compounds of Formula (A) wherein p, q and n are each zero, r is 1, m is 2, X is $-CH_2-$ and R^2 is $-OR^6$, especially when R^6 is hydrogen.

Another group of preferred compounds of Formula (A) is that in which Ar^1 is phenyl or R^4-substituted phenyl, Ar^2 is phenyl or R^4-substituted phenyl and Ar^3 is R^5-substituted phenyl. Also preferred are compounds in which Ar^1 is phenyl or R^4-substituted phenyl, Ar^2 is phenyl or R^4-substituted phenyl, Ar^3 is R^5-substituted phenyl, and the sum of m, n, p, q and r is 2, 3 or 4, more preferably 3. More preferred are compounds wherein Ar^1 is phenyl or R^4-substituted phenyl, Ar^2 is phenyl or R^4-substituted phenyl, Ar^3 is R^5-substituted phenyl, and wherein m, n and r are each zero, q is 1 and p is 2, or wherein p, q and n are each zero, r is 1 and m is 2 or 3.

Substituted Azetidinones of Formula (B)

In a preferred embodiment, a substituted azetidinone of Formula (A) useful in the compositions, therapeutic combinations and methods of the present invention is represented by Formula (B) (ezetimibe) below:




(B)

or pharmaceutically acceptable salts, solvates, or esters of the compound of Formula (B). The compound of Formula (B) can be in anhydrous or hydrated form. A product containing ezetimibe compound is commercially available as ZETIA® ezetimibe formulation from MSP Pharmaceuticals.

Compounds of Formula (A) can be prepared by a variety of methods well known to those skilled in the art, for example such as are disclosed in U.S. Patents Nos. 5,631,365, 5,767,115, 5,846,966, 6,207,822, 6,627,757, 6,093,812, 5,306,817, 5,561,227, 5,688,785, and 5,688,787, each of which is incorporated herein by reference.

Substituted Azetidinones of Formula (C)

Alternative substituted azetidinones useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (C) below:



(C)

or a pharmaceutically acceptable salt thereof or a solvate thereof, or an ester thereof, wherein, in Formula (C) above:

Ar^1 is R^3-substituted aryl;

Ar^2 is R^4-substituted aryl;

Ar^3 is R^5-substituted aryl;

Y and Z are independently selected from the group consisting of $-CH_2-$, -CH(lower alkyl)- and $-C(\text{lower alkyl})_2-$;

A is selected from -O-, -S-, -S(O)- or $-S(O)_2-$;

R^1 is selected from the group consisting of $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$ and $-OC(O)NR^6R^7$;

R^2 is selected from the group consisting of hydrogen, lower alkyl and aryl; or R^1 and R^2 together are =O;

q is 1, 2 or 3;

p is 0, 1, 2, 3 or 4;

R^5 is 1-3 substituents independently selected from the group consisting of $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$, $-O(CH_2)_{1-5}OR^9$, $-OC(O)NR^6R^7$, $-NR^6R^7$, $-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_2$-lower alkyl, $-NR^6S(O)_2$-aryl, $-C(O)NR^6R^7$, $-COR^6$, $-SO_2NR^6R^7$, $S(O)_{0-2}$-alkyl, $S(O)_{0-2}$-aryl, $-O(CH_2)_{1-10}-C(O)OR^6$, $-O(CH_2)_{1-10}C(O)NR^6R^7$, o-halogeno, m-halogeno, o-lower alkyl, m-lower alkyl, -(lower alkylene)-$C(O)OR^6$, and $-CH=CH-C(O)OR^6$;


R^3 and R^4 are independently 1-3 substituents independently selected from the group consisting of R^5, hydrogen, p-lower alkyl, aryl, $-NO_2$, $-CF_3$ and p-halogeno;

R^6, R^7 and R^8 are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl; and R^9 is lower alkyl, aryl or aryl-substituted lower alkyl.

Methods for making compounds of Formula (C) are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,688,990, which is incorporated herein by reference.

Substituted Azetidinones of Formula (D)

In another embodiment, substituted azetidinones useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (D):




(D)

or a pharmaceutically acceptable salt thereof or a solvate thereof, or an ester thereof, wherein, in Formula (D) above:

A is selected from the group consisting of R^2-substituted heterocycloalkyl, R^2-substituted heteroaryl, R^2-substituted benzo-fused heterocycloalkyl, and R^2-substituted benzo-fused heteroaryl;

Ar^1 is aryl or R^3-substituted aryl;

Ar^2 is aryl or R^4-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the spiro group  ; and

R^1 is selected from the group consisting of:

$-(CH_2)_q-$, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

$-(CH_2)_e-G-(CH_2)_r-$, wherein G is -O-, -C(O)-, phenylene, $-NR^8-$ or $-S(O)_{0-2}-$, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

$-(C_2-C_6$ alkenylene)-; and

$-(CH_2)_f-V-(CH_2)_g-$, wherein V is C_3-C_6 cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R^5 is selected from:

$$-\overset{|}{C}H-,\ -\overset{|}{C}(C_1\text{-}C_6\ \text{alkyl})-,\ -\overset{|}{C}F-,\ -\overset{|}{C}(OH)-,\ -\overset{|}{C}(C_6H_4\text{-}R^9)-,\ -\overset{|}{N}-,\ \text{or}\ -\overset{|}{\overset{+}{N}}O^-;$$

R^6 and R^7 are independently selected from the group consisting of $-CH_2-$, $-CH(C_1\text{-}C_6\ \text{alkyl})-$, $-C(\text{di-}(C_1\text{-}C_6)\ \text{alkyl})$, -CH=CH- and $-C(C_1\text{-}C_6\ \text{alkyl})=CH-$; or R^5 together with an adjacent R^6, or R^5 together with an adjacent R^7, form a -CH=CH- or a $-CH=C(C_1\text{-}C_6\ \text{alkyl})-$ group;

a and b are independently 0, 1, 2 or 3, provided both are not zero; provided that when R^6 is -CH=CH- or $-C(C_1\text{-}C_6\ \text{alkyl})=CH-$, a is 1; provided that when R^7 is -CH=CH- or $-C(C_1\text{-}C_6\ \text{alkyl})=CH-$, b is 1; provided that when a is 2 or 3, the R^6's can be the same or different; and provided that when b is 2 or 3, the R^7's can be the same or different;

and when Q is a bond, R^1 also can be selected from:

$$-M\text{-}Y_d-\overset{\overset{R^{10}}{|}}{\underset{\underset{R^{11}}{|}}{C}}\text{-}Z_h-,\ -X_m-\overset{\overset{R^{12}}{|}}{\underset{\underset{R^{13}}{|}}{(C)_s}}-Y_n-\overset{\overset{R^{10}}{|}}{\underset{\underset{R^{11}}{|}}{(C)_t}}-Z_p-\ \text{or}\ -X_j-\overset{\overset{R^{10}}{|}}{\underset{\underset{R^{11}}{|}}{(C)_v}}-Y_k-S(O)_{0\text{-}2}-;$$

where M is -O-, -S-, -S(O)- or $-S(O)_2-$;

X, Y and Z are independently selected from the group consisting of $-CH_2-$, $-CH(C_1\text{-}C_6\ \text{alkyl})-$ and $-C(\text{di-}(C_1\text{-}C_6)\ \text{alkyl})$;

R^{10} and R^{12} are independently selected from the group consisting of $-OR^{14}$, $-OC(O)R^{14}$, $-OC(O)OR^{16}$ and $-OC(O)NR^{14}R^{15}$;

R^{11} and R^{13} are independently selected from the group consisting of hydrogen, $(C_1\text{-}C_6)$alkyl and aryl; or R^{10} and R^{11} together are =O, or R^{12} and R^{13} together are =O;


d is 1, 2 or 3;


h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4; provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

R^2 is 1-3 substituents on the ring carbon atoms selected from the group consisting of hydrogen, $(C_1$-$C_{10})$alkyl, $(C_2$-$C_{10})$alkenyl, $(C_2$-$C_{10})$alkynyl, $(C_3$-$C_6)$cycloalkyl, $(C_3$-$C_6)$cycloalkenyl, R^{17}-substituted aryl, R^{17}-substituted benzyl, R^{17}-substituted benzyloxy, R^{17}-substituted aryloxy, halogeno, $-NR^{14}R^{15}$, $NR^{14}R^{15}(C_1$-C_6 alkylene)-, $NR^{14}R^{15}C(O)(C_1$-C_6 alkylene)-, $-NHC(O)R^{16}$, OH, C_1-C_6 alkoxy, $-OC(O)R^{16}$, $-C(O)R^{14}$, hydroxy$(C_1$-$C_6)$alkyl, $(C_1$-$C_6)$alkoxy$(C_1$-$C_6)$alkyl, NO_2, $-S(O)_{0\text{-}2}R^{16}$, $-S(O)_2NR^{14}R^{15}$ and $-(C_1$-C_6 alkylene$)C(O)OR^{14}$; when R^2 is a substituent on a heterocycloalkyl ring, R^2 is as defined, or R^2 is =O or  ; and, where R^2 is a substituent on a substitutable ring nitrogen, R^2 is hydrogen, $(C_1$-$C_6)$alkyl, aryl, $(C_1$-$C_6)$alkoxy, aryloxy, $(C_1$-$C_6)$alkylcarbonyl, arylcarbonyl, hydroxy, $-(CH_2)_{1\text{-}6}CONR^{18}R^{18}$,



;

wherein J is -O-, -NH-, $-NR^{18}$- or $-CH_2$-;

R^3 and R^4 are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of $(C_1$-$C_6)$alkyl, $-OR^{14}$, $-OC(O)R^{14}$, $-OC(O)OR^{16}$, $-O(CH_2)_{1\text{-}5}OR^{14}$, $-OC(O)NR^{14}R^{15}$, $-NR^{14}R^{15}$, $-NR^{14}C(O)R^{15}$, $-NR^{14}C(O)OR^{16}$, $-NR^{14}C(O)NR^{15}R^{19}$, $-NR^{14}S(O)_2R^{16}$, $-C(O)OR^{14}$, $-C(O)NR^{14}R^{15}$, $-C(O)R^{14}$, $-S(O)_2NR^{14}R^{15}$, $S(O)_{0\text{-}2}R^{16}$, $-O(CH_2)_{1\text{-}10}$-$C(O)OR^{14}$, $-O(CH_2)_{1\text{-}10}C(O)NR^{14}R^{15}$, $-(C_1$-C_6 alkylene$)$-$C(O)OR^{14}$, $-CH{=}CH$-$C(O)OR^{14}$, $-CF_3$, -CN, $-NO_2$ and halogen;

R^8 is hydrogen, $(C_1$-$C_6)$alkyl, aryl $(C_1$-$C_6)$alkyl, $-C(O)R^{14}$ or $-C(O)OR^{14}$;

R^9 and R^{17} are independently 1-3 groups independently selected from the group consisting of hydrogen, $(C_1$-$C_6)$alkyl, $(C_1$-$C_6)$alkoxy, -C(O)OH, NO_2, $-NR^{14}R^{15}$, OH and halogeno;

R^{14} and R^{15} are independently selected from the group consisting of hydrogen, $(C_1$-$C_6)$alkyl, aryl and aryl-substituted $(C_1$-$C_6)$alkyl;

R^{16} is $(C_1$-$C_6)$alkyl, aryl or R^{17}-substituted aryl;


R^{18} is hydrogen or $(C_1$-$C_6)$alkyl; and

R^{19} is hydrogen, hydroxy or $(C_1$-$C_6)$alkoxy.

Methods for making compounds of Formula (D) are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,656,624, which is incorporated herein by reference.

Substituted Azetidinones of Formula (E)

In another embodiment, substituted azetidinones useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (E):



(E)

or a pharmaceutically acceptable salt thereof or a solvate thereof, or an ester thereof, wherein, in Formula (E) above:

Ar^1 is aryl, R^{10}-substituted aryl or heteroaryl;

Ar^2 is aryl or R^4-substituted aryl;

Ar^3 is aryl or R^5-substituted aryl;

X and Y are independently selected from the group consisting of $-CH_2-$, -CH(lower alkyl)- and $-C(lower\ alkyl)_2-$;

R is $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$ or $-OC(O)NR^6R^7$; R^1 is hydrogen, lower alkyl or aryl; or R and R^1 together are =O;

q is 0 or 1;

r is 0, 1 or 2;

m and n are independently 0, 1, 2, 3, 4 or 5; provided that the sum of m, n and q is 1, 2, 3, 4 or 5;

R^4 is 1-5 substituents independently selected from the group consisting of lower alkyl, $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$, $-O(CH_2)_{1-5}OR^6$, $-OC(O)NR^6R^7$, $-NR^6R^7$, $-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_2R^9$, $-C(O)OR^6$, $-C(O)NR^6R^7$, $-C(O)R^6$, $-S(O)_2NR^6R^7$, $S(O)_{0-2}R^9$, $-O(CH_2)_{1-10}-C(O)OR^6$, $-O(CH_2)_{1-10}C(O)NR^6R^7$, -(lower alkylene)$C(O)OR^6$ and $-CH=CH-C(O)OR^6$;

R^5 is 1-5 substituents independently selected from the group consisting of $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$, $-O(CH_2)_{1-5}OR^6$, $-OC(O)NR^6R^7$, $-NR^6R^7$,

$-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_2R^9$, $-C(O)OR^6$, $-C(O)NR^6R^7$, $-C(O)R^6$, $-S(O)_2NR^6R^7$, $S(O)_{0-2}R^9$, $-O(CH_2)_{1-10}$-$C(O)OR^6$, $-O(CH_2)_{1-10}C(O)NR^6R^7$, $-CF_3$, $-CN$, $-NO_2$, halogen, -(lower alkylene)$C(O)OR^6$ and $-CH=CH-C(O)OR^6$;

R^6, R^7 and R^8 are independently selected from the group consisting of hydrogen, lower alkyl, aryl and aryl-substituted lower alkyl;


R^9 is lower alkyl, aryl or aryl-substituted lower alkyl; and

R^{10} is 1-5 substituents independently selected from the group consisting of lower alkyl, $-OR^6$, $-OC(O)R^6$, $-OC(O)OR^9$, $-O(CH_2)_{1-5}OR^6$, $-OC(O)NR^6R^7$, $-NR^6R^7$, $-NR^6C(O)R^7$, $-NR^6C(O)OR^9$, $-NR^6C(O)NR^7R^8$, $-NR^6S(O)_2R^9$, $-C(O)OR^6$, $-C(O)NR^6R^7$, $-C(O)R^6$, $-S(O)_2NR^6R^7$, $-S(O)_{0-2}R^9$, $-O(CH_2)_{1-10}$-$C(O)OR^6$, $-O(CH_2)_{1-10}C(O)NR^6R^7$, $-CF_3$, $-CN$, $-NO_2$ and halogen.

Methods for making compounds of Formula (E) are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,624,920, which is incorporated herein by reference.

Substituted Azetidinones of Formula (F)

In another embodiment, substituted azetidinones useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formula (F):



(F)

or a pharmaceutically acceptable salt thereof or a solvate thereof, or an ester thereof, wherein:

R^1 is:

$-\overset{|}{C}H-$, $-\overset{|}{C}$(lower alkyl)-, $-\overset{|}{C}F-$, $-\overset{|}{C}(OH)-$, $-\overset{|}{C}(C_6H_5)-$, $-\overset{|}{C}(C_6H_4-R_{15})-$,

$-\overset{|}{N}-$ or $-\overset{|}{\overset{+}{N}}O^-$;

R^2 and R^3 are independently selected from the group consisting of:

$-CH_2-$, -CH(lower alkyl)-, $-C(\text{lower alkyl})_2-$, -CH=CH- and -C(lower alkyl)=CH-; or


R^1 together with an adjacent R^2, or R^1 together with an adjacent R^3, form a -CH=CH- or a -CH=C(lower alkyl)- group;


u and v are independently 0, 1, 2 or 3, provided both are not zero; provided that when R^2 is -CH=CH- or -C(lower alkyl)=CH-, v is 1; provided that when R^3 is -CH=CH- or -C(lower alkyl)=CH-, u is 1; provided that when v is 2 or 3, each R^2 can be the same or different; and provided that when u is 2 or 3, each R^3 can be the same or different;

R^4 is selected from $B-(CH_2)_mC(O)-$, wherein m is 0, 1, 2, 3, 4 or 5; $B-(CH_2)_q-$, wherein q is 0, 1, 2, 3, 4, 5 or 6; $B-(CH_2)_e-Z-(CH_2)_r-$, wherein Z is -O-, -C(O)-, phenylene, $-N(R^8)-$ or $-S(O)_{0-2}-$, e is 0, 1, 2, 3, 4 or 5 and r is 0, 1, 2, 3, 4 or 5, provided that the sum of e and r is 0, 1, 2, 3, 4, 5 or 6; $B-(C_2-C_6$ alkenylene)-; $B-(C_4-C_6$ alkadienylene)-; $B-(CH_2)_t-Z-(C_2-C_6$ alkenylene)-, wherein Z is as defined above, and wherein t is 0, 1, 2 or 3, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6; $B-(CH_2)_f-V-(CH_2)_g-$, wherein V is C_3-C_6 cycloalkylene, f is 1, 2, 3, 4 or 5 and g is 0, 1, 2, 3, 4 or 5, provided that the sum of f and g is 1, 2, 3, 4, 5 or 6; $B-(CH_2)_t-V-(C_2-C_6$ alkenylene)- or $B-(C_2-C_6$ alkenylene)$-V-(CH_2)_t-$, wherein V and t are as defined above, provided that the sum of t and the number of carbon atoms in the alkenylene chain is 2, 3, 4, 5 or 6; $B-(CH_2)_a-Z-(CH_2)_b-V-(CH_2)_d-$, wherein Z and V are as defined above and a, b and d are independently 0, 1, 2, 3, 4, 5 or 6, provided that the sum of a, b and d is 0, 1, 2, 3, 4, 5 or 6; or $T-(CH_2)_s-$, wherein T is a C_3-C_6 cycloalkyl and s is 0, 1, 2, 3, 4, 5 or 6; or

R^1 and R^4 together form the group $B-CH=\overset{|}{C}-$;

B is selected from indanyl, indenyl, naphthyl, tetrahydronaphthyl, heteroaryl or W-substituted heteroaryl, wherein heteroaryl is selected from the group consisting of pyrrolyl, pyridinyl, pyrimidinyl, pyrazinyl, triazinyl, imidazolyl, thiazolyl, pyrazolyl, thienyl, oxazolyl and furanyl, and for nitrogen-containing heteroaryls, the N-oxides thereof, or




W is 1 to 3 substituents independently selected from the group consisting of lower alkyl, hydroxy lower alkyl, lower alkoxy, alkoxyalkyl, alkoxyalkoxy, alkoxycarbonylalkoxy, (lower alkoxyimino)-lower alkyl, lower alkanedioyl, lower alkyl lower alkanedioyl, allyloxy, $-CF_3$, $-OCF_3$, benzyl, R^7-benzyl, benzyloxy, R^7-benzyloxy, phenoxy, R^7-phenoxy, dioxolanyl, NO_2, $-N(R^8)(R^9)$, $N(R^8)(R^9)$-lower alkylene-, $N(R^8)(R^9)$-lower alkylenyloxy-, OH, halogeno, -CN, $-N_3$, $-NHC(O)OR^{10}$, $-NHC(O)R^{10}$, $R^{11}(O)_2SNH-$, $(R^{11}(O)_2S)_2N-$, $-S(O)_2NH_2$, $-S(O)_{0-2}R^8$, tert-butyldimethyl-silyloxymethyl, $-C(O)R^{12}$, $-C(O)OR^{19}$, $-C(O)N(R^8)(R^9)$, $-CH=CHC(O)R^{12}$, -lower alkylene-$C(O)R^{12}$, $R^{10}C(O)$(lower alkylenyloxy)-, $N(R^8)(R^9)C(O)$(lower alkylenyloxy)- and  for substitution on ring carbon atoms, and the substituents on the substituted heteroaryl ring nitrogen atoms, when present, are selected from the group consisting of lower alkyl, lower alkoxy, $-C(O)OR^{10}$, $-C(O)R^{10}$, OH, $N(R^8)(R^9)$-lower alkylene-, $N(R^8)(R^9)$-lower alkylenyloxy-, $-S(O)_2NH_2$ and 2-(trimethylsilyl)-ethoxymethyl;

R^7 is 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, -C(O)OH, NO_2, $-N(R^8)(R^9)$, OH, and halogeno;

R^8 and R^9 are independently selected from H or lower alkyl;

R^{10} is selected from lower alkyl, phenyl, R^7-phenyl, benzyl or R^7-benzyl;

R^{11} is selected from OH, lower alkyl, phenyl, benzyl, R^7-phenyl or R^7-benzyl;

R^{12} is selected from H, OH, alkoxy, phenoxy, benzyloxy,  , $-N(R^8)(R^9)$, lower alkyl, phenyl or R^7-phenyl;

R^{13} is selected from -O-, $-CH_2-$, -NH-, -N(lower alkyl)- or $-NC(O)R^{19}$;

R^{15}, R^{16} and R^{17} are independently selected from the group consisting of H and the groups defined for W; or R^{15} is hydrogen and R^{16} and R^{17},

together with adjacent carbon atoms to which they are attached, form a dioxolanyl ring;


R^{19} is H, lower alkyl, phenyl or phenyl lower alkyl; and

R^{20} and R^{21} are independently selected from the group consisting of phenyl, W-substituted phenyl, naphthyl, W-substituted naphthyl, indanyl, indenyl, tetrahydronaphthyl, benzodioxolyl, heteroaryl, W-substituted heteroaryl, benzo-fused heteroaryl, W-substituted benzo-fused heteroaryl and cyclopropyl, wherein heteroaryl is as defined above.

Methods for making compounds of Formula (F) are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,698,548, which is incorporated herein by reference.


Substituted Azetidinones of Formula (G)

In another embodiment, substituted azetidinones useful in the compositions, therapeutic combinations and methods of the present invention are represented by Formulas (GA) and (GB):



(GA)

and




(GB)


or a pharmaceutically acceptable salt, solvate, or ester thereof, wherein:

A is -CH=CH-, -C≡C- or $-(CH_2)_p-$ wherein p is 0, 1 or 2;

B is



B' is




D is $-(CH_2)_mC(O)-$ or $-(CH_2)_q-$ wherein m is 1, 2, 3 or 4 and q is 2, 3 or 4;

E is C_{10} to C_{20} alkyl or $-C(O)-(C_9$ to $C_{19})$-alkyl, wherein the alkyl is straight or branched, saturated or containing one or more double bonds;

R is hydrogen, C_1-C_{15} alkyl, straight or branched, saturated or containing one or more double bonds, or $B-(CH_2)_r-$, wherein r is 0, 1, 2, or 3;

R^1, R^2, R^3, $R^{1'}$, $R^{2'}$, and $R^{3'}$ are independently selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, carboxy, NO_2, NH_2, OH, halogeno, lower alkylamino, dilower alkylamino, $-NHC(O)OR^5$, $R^6(O)_2SNH-$ and $-S(O)_2NH_2$;

R^4 is




wherein n is 0, 1, 2 or 3;

R^5 is lower alkyl; and

R^6 is OH, lower alkyl, phenyl, benzyl or substituted phenyl wherein the substituents are 1-3 groups independently selected from the group consisting of lower alkyl, lower alkoxy, carboxy, NO_2, NH_2, OH, halogeno, lower alkylamino and dilower alkylamino; or a pharmaceutically acceptable salt, solvate, or ester thereof.

Sterol Absorption Inhibitors of Formula (H)

In another embodiment, sterol absorption inhibitors useful in the compositions and methods of the present invention are represented by Formula (H):







(H)

or a pharmaceutically acceptable salt, solvate, or ester thereof, wherein, in Formula (H) above,

R^{26} is H or OG^1;

G and G^1 are independently selected from the group consisting of

H,  ,  , 

and  ; provided that when R^{26} is H or OH, G is not H;

R, R^a and R^b are independently selected from the group consisting of H, -OH, halogeno, $-NH_2$, azido, (C_1-C_6)alkoxy(C_1-C_6)-alkoxy or $-W-R^{30}$;

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R^{31})-, -NH-C(O)-N(R^{31})- and -O-C(S)-N(R^{31})-;

R^2 and R^6 are independently selected from the group consisting of H, (C_1-C_6)alkyl, aryl and aryl(C_1-C_6)alkyl;

R^3, R^4, R^5, R^7, R^{3a} and R^{4a} are independently selected from the group consisting of H, (C_1-C_6)alkyl, aryl(C_1-C_6)alkyl, -C(O)(C_1-C_6)alkyl and -C(O)aryl;

R^{30} is selected from the group consisting of R^{32}-substituted T, R^{32}-substituted-T-(C_1-C_6)alkyl, R^{32}-substituted-(C_2-C_4)alkenyl, R^{32}-substituted-(C_1-C_6)alkyl, R^{32}-substituted-(C_3-C_7)cycloalkyl and R^{32}-substituted-(C_3-C_7)cycloalkyl(C_1-C_6)alkyl;

R^{31} is selected from the group consisting of H and (C_1-C_4)alkyl;

T is selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, iosthiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;

R^{32} is independently selected from 1-3 substituents independently selected from the group consisting of halogeno, (C_1-C_4)alkyl, -OH, phenoxy, $-CF_3$, $-NO_2$, (C_1-C_4)alkoxy, methylenedioxy, oxo, (C_1-C_4)alkylsulfanyl, (C_1-C_4)alkylsulfinyl, (C_1-C_4)alkylsulfonyl, $-N(CH_3)_2$, $-C(O)-NH(C_1-C_4)$alkyl, $-C(O)-N((C_1-C_4)alkyl)_2$, $-C(O)-(C_1-C_4)$alkyl, $-C(O)-(C_1-C_4)$alkoxy and pyrrolidinylcarbonyl; or

R^{32} is a covalent bond and R^{31}, the nitrogen to which it is attached and R^{32} form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C_1-C_4)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

Ar^1 is aryl or R^{10}-substituted aryl;

Ar^2 is aryl or R^{11}-substituted aryl;

Q is a bond or, with the 3-position ring carbon of the azetidinone, forms the spiro group  ; and

R^1 is selected from the group consisting of

$-(CH_2)_q-$, wherein q is 2-6, provided that when Q forms a spiro ring, q can also be zero or 1;

$-(CH_2)_e-E-(CH_2)_r-$, wherein E is -O-, -C(O)-, phenylene, $-NR^{22}-$ or $-S(O)_{0-2}-$, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

$-(C_2-C_6)$alkenylene-; and

$-(CH_2)_f-V-(CH_2)_g-$, wherein V is C_3-C_6 cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6;

R^{12} is:



R^{13} and R^{14} are independently selected from the group consisting of $-CH_2-$, $-CH((C_1-C_6)$ alkyl)-, $-C((C_1-C_6)$ alkyl)$_2$, -CH=CH- and $-C((C_1-C_6)$ alkyl)=CH-; or

R^{12} together with an adjacent R^{13}, or R^{12} together with an adjacent R^{14}, form a -CH=CH- or a -CH=C(C_1-C_6 alkyl)- group;

a and b are independently 0, 1, 2 or 3, provided both are not zero;

provided that when R^{13} is -CH=CH- or -C(C_1-C_6 alkyl)=CH-, a is 1;

provided that when R^{14} is -CH=CH- or -C(C_1-C_6 alkyl)=CH-, b is 1;

provided that when a is 2 or 3, each R^{13} can be the same or different; and

provided that when b is 2 or 3, each R^{14} can be the same or different;

and when Q is a bond, R^1 also can be:

$$-M\text{-}Y_d\text{-}\overset{\overset{R^{15}}{|}}{\underset{\underset{R^{16}}{|}}{C}}\text{-}Z_h-\ ,\quad -X_m\text{-}\overset{\overset{R^{17}}{|}}{\underset{\underset{R^{18}}{|}}{(C)_s}}\text{-}Y_n\text{-}\overset{\overset{R^{15}}{|}}{\underset{\underset{R^{16}}{|}}{(C)_t}}\text{-}Z_p-\quad \text{or}\quad -X_j\text{-}\overset{\overset{R^{15}}{|}}{\underset{\underset{R^{16}}{|}}{(C)_v}}\text{-}Y_k\text{-}S(O)_{0\text{-}2}-;$$

M is -O-, -S-, -S(O)- or -$S(O)_2$-;

X, Y and Z are independently selected from the group consisting of -CH_2-, -CH(C_1-C_6)alkyl- and -C((C_1-C_6)alkyl)$_2$;

R^{10} and R^{11} are independently selected from the group consisting of 1-3 substituents independently selected from the group consisting of (C_1-C_6)alkyl, -OR^{19}, -OC(O)R^{19}, -OC(O)OR^{21}, -O(CH_2)$_{1\text{-}5}OR^{19}$, -OC(O)$NR^{19}R^{20}$, -$NR^{19}R^{20}$, -$NR_{19}C(O)R^{20}$, -$NR^{19}C(O)OR^{21}$, -$NR^{19}C(O)NR^{20}R^{25}$, -$NR^{19}S(O)_2R^{21}$, -C(O)OR^{19}, -C(O)$NR^{19}R^{20}$, -C(O)R^{19}, -$S(O)_2NR^{19}R^{20}$, $S(O)_{0\text{-}2}R^{21}$, -O(CH_2)$_{1\text{-}10}$-C(O)OR^{19}, -O(CH_2)$_{1\text{-}10}$C(O)$NR^{19}R^{20}$, -(C_1-C_6 alkylene)-C(O)OR^{19}, -CH=CH-C(O)OR^{19}, -CF_3, -CN, -NO_2 and halogen;

R^{15} and R^{17} are independently selected from the group consisting of -OR^{19}, -OC(O)R^{19}, -OC(O)OR^{21} and -OC(O)$NR^{19}R^{20}$;

R^{16} and R^{18} are independently selected from the group consisting of H, (C_1-C_6)alkyl and aryl; or R^{15} and R^{16} together are =O, or R^{17} and R^{18} together are =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1; t is 0 or 1; m, n and p are independently 0-4;

provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6;

provided that when p is 0 and t is 1, the sum of m, s and n is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;

v is 0 or 1;

j and k are independently 1-5, provided that the sum of j, k and v is 1-5;

and when Q is a bond and R^1 is $-X_j-(C)_v-Y_k-S(O)_{0-2}-$ (with R^{15} and R^{16}), Ar^1 can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R^{19} and R^{20} are independently selected from the group consisting of H, (C_1-C_6)alkyl, aryl and aryl-substituted (C_1-C_6)alkyl;

R^{21} is (C_1-C_6)alkyl, aryl or R^{24}-substituted aryl;

R^{22} is H, (C_1-C_6)alkyl, aryl (C_1-C_6)alkyl, $-C(O)R^{19}$ or $-C(O)OR^{19}$;


R^{23} and R^{24} are independently 1-3 groups independently selected from the group consisting of H, (C_1-C_6)alkyl, (C_1-C_6)alkoxy, -C(O)OH, NO_2, $-NR^{19}R^{20}$, -OH and halogeno; and

R^{25} is H, -OH or (C_1-C_6)alkoxy.

Methods for making compounds of Formula (H) are well known to those skilled in the art. Non-limiting examples of suitable methods are disclosed in U.S. Patent No. 5,756,470, which is incorporated herein by reference.

Substituted Azetidinones of Formula (J)

In another embodiment, substituted azetidinones useful in the compositions and methods of the present invention are represented by Formula (J) below:








(J)

or a pharmaceutically acceptable salt, solvate, or ester thereof, wherein in Formula (J):

R^1 is selected from the group consisting of H, G, G^1, G^2, $-SO_3H$ and $-PO_3H$;

G is selected from the group consisting of: H,


,

,

,

and

,

(sugar derivatives)

wherein R, R^a and R^b are each independently selected from the group consisting of H, -OH, halogen, $-NH_2$, azido, (C_1-C_6)alkoxy(C_1-C_6)alkoxy or -W-R^{30};

W is independently selected from the group consisting of -NH-C(O)-, -O-C(O)-, -O-C(O)-N(R^{31})-, -NH-C(O)-N(R^{31})- and -O-C(S)-N(R^{31})-;

R^2 and R^6 are each independently selected from the group consisting of H, (C_1-C_6)alkyl, acetyl, aryl and aryl(C_1-C_6)alkyl;

R^3, R^4, R^5, R^7, R^{3a} and R^{4a} are each independently selected from the group consisting of H, (C_1-C_6)alkyl, acetyl, aryl(C_1-C_6)alkyl, -C(O)(C_1-C_6)alkyl and -C(O)aryl;

R^{30} is independently selected from the group consisting of R^{32}-substituted T, R^{32}-substituted-T-(C_1-C_6)alkyl, R^{32}-substituted-(C_2-C_4)alkenyl, R^{32}-substituted-(C_1-C_6)alkyl, R^{32}-substituted-(C_3-C_7)cycloalkyl and R^{32}-substituted-(C_3-C_7)cycloalkyl(C_1-C_6)alkyl;


R^{31} is independently selected from the group consisting of H and (C_1-C_4)alkyl;

T is independently selected from the group consisting of phenyl, furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzothiazolyl, thiadiazolyl, pyrazolyl, imidazolyl and pyridyl;


R^{32} is independently selected from 1-3 substituents which are each independently selected from the group consisting of H, halogen, (C_1-C_4)alkyl, -OH, phenoxy, -CF_3, -NO_2, (C_1-C_4)alkoxy, methylenedioxy, oxo, (C_1-C_4)alkylsulfanyl, (C_1-C_4)alkylsulfinyl, (C_1-C_4)alkylsulfonyl, -$N(CH_3)_2$, -C(O)-NH(C_1-C_4)alkyl, -C(O)-N(C_1-C_4)alkyl)$_2$, -C(O)-(C_1-C_4)alkyl, -C(O)-(C_1-C_4)alkoxy and pyrrolidinylcarbonyl; or

R^{32} is a covalent bond and R^{31}, the nitrogen to which it is attached and R^{32} form a pyrrolidinyl, piperidinyl, N-methyl-piperazinyl, indolinyl or morpholinyl group, or a (C_1-C_4)alkoxycarbonyl-substituted pyrrolidinyl, piperidinyl, N-methylpiperazinyl, indolinyl or morpholinyl group;

G^1 is represented by the structure:



wherein R^{33} is independently selected from the group consisting of unsubstituted alkyl, R^{34}-substituted alkyl, (R^{35})(R^{36})alkyl-,




R^{34} is one to three substituents, each R^{34} being independently selected from the group consisting of HO(O)C-, HO-, HS-, (CH_3)S-, H_2N-, (NH_2)(NH)C(NH)-, (NH_2)C(O)- and HO(O)CCH(NH_3^+)CH_2SS-;

R^{35} is independently selected from the group consisting of H and NH_2-;

R^{36} is independently selected from the group consisting of H, unsubstituted alkyl, R^{34}-substituted alkyl, unsubstituted cycloalkyl and R^{34}-substituted cycloalkyl;

G^2 is represented by the structure:



wherein R^{37} and R^{38} are each independently selected from the group consisting of $(C_1\text{-}C_6)$alkyl and aryl;

R^{26} is one to five substituents, each R^{26} being independently selected from the group consisting of:

a) H;

b) -OH;

c) $-OCH_3$;

d) fluorine;

e) chlorine;

f) –O-G;

g) $-O\text{-}G^1$;

h) $-O\text{-}G^2$;

i) $-SO_3H$; and

j) $-PO_3H$;

provided that when R^1 is H, R^{26} is not H, –OH, $-OCH_3$ or –O-G;

Ar^1 is aryl, R^{10}-substituted aryl, heteroaryl or R^{10}-substituted heteroaryl;

Ar^2 is aryl, R^{11}-substituted aryl, heteroaryl or R^{11}-substituted heteroaryl;

L is selected from the group consisting of:

a) a covalent bond;


b) $-(CH_2)_q-$, wherein q is 1-6;

c) $-(CH_2)_e\text{-}E\text{-}(CH_2)_r-$, wherein E is –O-, -C(O)-, phenylene, $-NR^{22}-$ or $-S(O)_{0\text{-}2}-$, e is 0-5 and r is 0-5, provided that the sum of e and r is 1-6;

d) $-(C_2\text{-}C_6)$alkenylene-;

e) $-(CH_2)_f\text{-}V\text{-}(CH_2)_g-$, wherein V is $C_3\text{-}C_6$cycloalkylene, f is 1-5 and g is 0-5, provided that the sum of f and g is 1-6; and

f)



wherein M is –O-, -S-, -S(O)- or –$S(O)_2$-;

X, Y and Z are each independently selected from the group consisting of

–CH_2-, -CH(C_1-C_6)alkyl- and –C((C_1-C_6)alkyl)$_2$-;

R^8 is selected from the group consisting of H and alkyl;

R^{10} and R^{11} are each independently selected from the group consisting of 1-3 substituents which are each independently selected from the group consisting of (C_1-C_6)alkyl, -OR^{19}, -$OC(O)R^{19}$, -$OC(O)OR^{21}$, -$O(CH_2)_{1-5}OR^{19}$, -$OC(O)NR^{19}R^{20}$, -$NR^{19}R^{20}$, -$NR^{19}C(O)R^{20}$, -$NR^{19}C(O)OR^{21}$, -$NR^{19}C(O)NR^{20}R^{25}$, -$NR^{19}S(O)_2R^{21}$, -$C(O)OR^{19}$, -$C(O)NR^{19}R^{20}$, -$C(O)R^{19}$, -$S(O)_2NR^{19}R^{20}$, $S(O)_{0-2}R^{21}$, -$O(CH_2)_{1-10}$-$C(O)OR^{19}$, -$O(CH_2)_{1-10}C(O)NR^{19}R^{20}$, -($C_1$-$C_6$ alkylene)-$C(O)OR^{19}$, -CH=CH-$C(O)OR^{19}$, -CF_3, -CN, -NO_2 and halogen;

R^{15} and R^{17} are each independently selected from the group consisting of –OR^{19}, -$OC(O)R^{19}$, -$OC(O)OR^{21}$, - $OC(O)NR^{19}R^{20}$;

R^{16} and $R^{18}$are each independently selected from the group consisting of H, (C_1-C_6)alkyl and aryl; or

R^{15} and R^{16} together are =O, or $R^{17}$and R^{18} together are =O;

d is 1, 2 or 3;

h is 0, 1, 2, 3 or 4;

s is 0 or 1;

t is 0 or 1;


m, n and p are each independently selected from 0-4;

provided that at least one of s and t is 1, and the sum of m, n, p, s and t is 1-6; provided that when p is 0 and t is 1, the sum of m, n and p is 1-5; and provided that when p is 0 and s is 1, the sum of m, t and n is 1-5;


v is 0 or 1;

j and k are each independently 1-5, provided that the sum of j, k and v is 1-5;

Q is a bond, -$(CH_2)_q$-, wherein q is 1-6, or, with the 3-position ring carbon of the azetidinone, forms the spiro group




wherein R^{12} is



R^{13} and R^{14} are each independently selected from the group consisting of $-CH_2-$, $-CH(C_1-C_6$ alkyl)-, $-C((C_1-C_6)$ alkyl$)_2$, -CH=CH- and $-C(C_1-C_6$ alkyl)=CH-; or R^{12} together with an adjacent R^{13}, or R^{12} together with an adjacent R^{14}, form a -CH=CH- or a $-CH=C(C_1-C_6$ alkyl)- group;

a and b are each independently 0, 1, 2 or 3, provided both are not zero; provided that when R^{13} is -CH=CH- or $-C(C_1-C_6$ alkyl)=CH-, a is 1; provided that when R^{14} is -CH=CH- or $-C(C_1-C_6$ alkyl)=CH-, b is 1; provided that when a is 2 or 3, each R^{13} can be the same or different; and provided that when b is 2 or 3, each R^{14} can be the same or different;

and when Q is a bond and L is



then Ar^1 can also be pyridyl, isoxazolyl, furanyl, pyrrolyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, pyrazinyl, pyrimidinyl or pyridazinyl;

R^{19} and R^{20} are each independently selected from the group consisting of H, (C_1-C_6)alkyl, aryl and aryl-substituted (C_1-C_6)alkyl;

R^{21} is (C_1-C_6)alkyl, aryl or R^{24}-substituted aryl;

R^{22} is H, (C_1-C_6)alkyl, aryl (C_1-C_6)alkyl, $-C(O)R^{19}$ or $-C(O)OR^{19}$;


R^{23} and R^{24} are each independently selected from the group consisting of 1-3 substituents which are each independently selected from the group consisting of H, (C_1-C_6)alkyl, (C_1-C_6)alkoxy, -C(O)OH, NO_2, $-NR^{19}R^{20}$, -OH and halogen; and

R^{25} is H, -OH or (C_1-C_6)alkoxy.

Examples of compounds of Formula (J) which are useful in the methods and combinations of the present invention and methods for making such compounds are disclosed in U.S. Patent Application Serial No. 10/166,942, filed June 11, 2002, incorporated herein by reference.

Substituted Azetidinones of Formulae (K)-(M)


An example of a useful substituted azetidinone is one represented by the Formula (K):



(K)


wherein R^1 is defined as above.

A more preferred compound is one represented by Formula (L):



(L).


Another useful compound is represented by Formula (M):



(M)

Other useful substituted azetidinone compounds include N-sulfonyl-2-azetidinones such as are disclosed in U.S. Patent No. 4,983,597, ethyl 4-(2-oxoazetidin-4-yl)phenoxy-alkanoates such as are disclosed in Ram et al., Indian J. Chem. Sect. B. 29B, 12 (1990), p. 1134-7, diphenyl azetidinones and derivatives disclosed in U.S. Patent Publication Nos. 2002/0039774, 2002/0128252, 2002/0128253 and 2002/0137689, 2004/063929, WO 2002/066464, U.S. Patent Nos. 6,498,156 and 6,703,386, each of which is incorporated by reference herein.

Other sterol absorption inhibitors useful in the compositions, therapeutic combinations and methods of the present invention are described in WO 2004/005247, WO 2004/000803, WO 2004/000804, WO 2004/000805, WO 0250027, U.S. published application 2002/0137689, and the compounds described in L. Kværnø et al., Angew. Chem. Int. Ed., **2004**, vol. 43, pp. 4653-4656, all of which are incorporated herein by reference. An illustrative compound of Kværnø et al. is:



The compounds of Formulae A-M can be prepared by known methods, including the methods discussed above and, for example, in WO 93/02048, U.S. 5,306,817 and 5,561,227, herein incorporated by reference, which describe the preparation of compounds wherein -R^1-Q- is alkylene, alkenylene or alkylene interrupted by a hetero atom, phenylene or cycloalkylene; WO 94/17038 and U.S. 5,698,548, herein incorporated by reference, describe the preparation of compounds wherein Q is a spirocyclic group; WO 95/08532, U.S. 5,631,365, U.S. 5,767,115, U.S. 5,846,966, and U.S. R.E. 37,721, herein incorporated by reference, describe the preparation of compounds wherein -R^1-Q- is a hydroxy-substituted alkylene group; PCT/US95/03196, herein incorporated by reference, describes compounds wherein -R^1-Q- is a hydroxy-substituted alkylene attached to the Ar^1 moiety through an -O- or $S(O)_{0-2}$- group; and U.S. Serial No. 08/463,619, filed June 5, 1995, herein incorporated by reference, describes the preparation of compounds wherein -R^1-Q- is a hydroxy-substituted alkylene group attached to the azetidinone ring by a -$S(O)_{0-2}$- group. Each of the above patents or publications are herein incorporated by reference in their entirety.

The daily dose of the sterol absorption inhibitor(s) administered to the subject can range from about 0.1 to about 1000 mg per day, preferably about 0.25 to about 50 mg/day, and more preferably about 10 mg per day, given in a single dose or 2-4 divided doses. The exact dose, however, is determined by the attending clinician and is dependent on the potency of the compound administered, the age, weight, condition and response of the patient.

For administration of pharmaceutically acceptable salts of the above compounds, the weights indicated above refer to the weight of the acid equivalent or the base equivalent of the therapeutic compound derived from the salt.

In another embodiment of the present invention, the compositions or therapeutic combinations described above comprise one or more selective CB_1 receptor antagonist compounds of Formula (I) in combination with one or more cholesterol biosynthesis inhibitors and/or lipid-lowering compounds discussed below.

Generally, a total daily dosage of cholesterol biosynthesis inhibitor(s) can range from about 0.1 to about 160 mg per day, and preferably about 0.2 to about 80 mg/day in single or 2-3 divided doses.

In another alternative embodiment, the compositions, therapeutic combinations or methods of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more bile acid sequestrants (insoluble anion exchange resins), co-administered with or in combination with the compound of Formula (I), or a pharmaceutically acceptable salt, solvate, or ester thereof, and a substituted azetidinone or a substituted β-lactam discussed above.

Bile acid sequestrants bind bile acids in the intestine, interrupting the enterohepatic circulation of bile acids and causing an increase in the faecal excretion of steroids. Use of bile acid sequestrants is desirable because of their non-systemic mode of action. Bile acid sequestrants can lower intrahepatic cholesterol and promote the synthesis of apo B/E (LDL) receptors that bind LDL from plasma to further reduce cholesterol levels in the blood.

Generally, a total daily dosage of bile acid sequestrant(s) can range from about 1 to about 50 grams per day, and preferably about 2 to about 16 grams per day in single or 2-4 divided doses.

In an alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more IBAT inhibitors. The IBAT inhibitors can inhibit bile acid transport to reduce LDL cholesterol levels. Generally, a total daily dosage of IBAT inhibitor(s) can range from about 0.01 to about 1000 mg/day, and preferably about 0.1 to about 50 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and nicotinic acid (niacin) and/or derivatives thereof. Nicotinic acid and its derivatives inhibit hepatic production of VLDL and its metabolite LDL and increases HDL and apo A-1 levels. An example of a suitable nicotinic acid product is NIASPAN® (niacin extended-release tablets) which are available from Kos.

Generally, a total daily dosage of nicotinic acid or a derivative thereof can range from about 500 to about 10,000 mg/day, preferably about 1000 to about 8000 mg/day, and more preferably about 3000 to about 6000 mg/day in single or divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or estes thereof, and one or more AcylCoA:Cholesterol *O*-acyltransferase ("ACAT") Inhibitors, which can reduce LDL and VLDL levels. ACAT is an enzyme responsible for esterifying excess intracellular cholesterol and may reduce the synthesis of VLDL, which is a product of cholesterol esterification, and overproduction of apo B-100-containing lipoproteins. Generally, a total daily dosage of ACAT inhibitor(s) can range from about 0.1 to about 1000 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more Cholesteryl Ester Transfer Protein ("CETP") Inhibitors. CETP is responsible for the exchange or transfer of cholesteryl ester carrying HDL and triglycerides in VLDL. Pancreatic cholesteryl ester hydrolase (pCEH) inhibitors such as WAY-121898 also can be co-administered with or in combination.

Generally, a total daily dosage of CETP inhibitor(s) can range from about 0.01 to about 1000 mg/day, and preferably about 0.5 to about 20 mg/kg body weight/day in single or divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and probucol or derivatives thereof, which can reduce LDL levels.

Generally, a total daily dosage of probucol or derivatives thereof can range from about 10 to about 2000 mg/day, and preferably about 500 to about 1500 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or

pharmaceutically acceptable salts, solvates, or esters thereof, and low-density lipoprotein (LDL) receptor activators.

Generally, a total daily dosage of LDL receptor activator(s) can range from about 1 to about 1000 mg/day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and fish oil. Generally, a total daily dosage of fish oil or Omega 3 fatty acids can range from about 1 to about 30 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can further comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and natural water soluble fibers, such as psyllium, guar, oat and pectin, which can reduce cholesterol levels. Generally, a total daily dosage of natural water soluble fibers can range from about 0.1 to about 10 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and plant sterols, plant stanols and/or fatty acid esters of plant stanols, such as sitostanol ester used in BENECOL® margarine, which can reduce cholesterol levels. Generally, a total daily dosage of plant sterols, plant stanols and/or fatty acid esters of plant stanols can range from about 0.5 to about 20 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and antioxidants, such as probucol, tocopherol, ascorbic acid, β-carotene and selenium, or vitamins such as vitamin B_6 or vitamin B_{12}. Generally, a total daily dosage of antioxidants or vitamins can range from about 0.05 to about 10 grams per day in single or 2-4 divided doses.

In another alternative embodiment, the compositions or treatments of the present invention can comprise at least one compound of Formula (I), or

pharmaceutically acceptable salts, solvates, or esters thereof, and monocyte and macrophage inhibitors such as polyunsaturated fatty acids (PUFA), thyroid hormones including throxine analogues such as CGS-26214 (a thyroxine compound with a fluorinated ring), gene therapy and use of recombinant proteins such as recombinant apo E. Generally, a total daily dosage of these agents can range from about 0.01 to about 1000 mg/day in single or 2-4 divided doses.

Also useful with the present invention are compositions or therapeutic combinations that further comprise hormone replacement agents and compositions. Useful hormone agents and compositions for hormone replacement therapy of the present invention include androgens, estrogens, progestins, their pharmaceutically acceptable salts and derivatives thereof. Combinations of these agents and compositions are also useful.

The dosage of androgen and estrogen combinations vary, desirably from about 1 mg to about 4 mg androgen and from about 1 mg to about 3 mg estrogen. Examples include, but are not limited to, androgen and estrogen combinations such as the combination of esterified estrogens (sodium estrone sulfate and sodium equilin sulfate) and methyltestosterone (17-hydroxy-17-methyl-, (17B)- androst-4-en-3-one) available from Solvay Pharmaceuticals, Inc., Marietta, GA, under the tradename Estratest.

Estrogens and estrogen combinations may vary in dosage from about 0.01 mg up to 8 mg, desirably from about 0.3 mg to about 3.0 mg. Examples of useful estrogens and estrogen combinations include:

(a) the blend of nine (9) synthetic estrogenic substances including sodium estrone sulfate, sodium equilin sulfate, sodium 17 α -dihydroequilin sulfate, sodium 17 α -estradiol sulfate, sodium 17 β -dihydroequilin sulfate, sodium 17 α -dihydroequilenin sulfate, sodium 17 β -dihydroequilenin sulfate, sodium equilenin sulfate and sodium 17 β -estradiol sulfate; available from Duramed Pharmaceuticals, Inc., Cincinnati, OH, under the tradename Cenestin;

(b) ethinyl estradiol (19-nor-17 α -pregna-1,3,5(10)-trien-20-yne-3,17-diol; available by Schering Plough Corporation, Kenilworth, NJ, under the tradename Estinyl;

(c) esterified estrogen combinations such as sodium estrone sulfate and sodium equilin sulfate; available from Solvay under the tradename Estratab and from Monarch Pharmaceuticals, Bristol, TN, under the tradename Menest;

(d) estropipate (piperazine estra-1,3,5(10)-trien-17-one, 3-(sulfooxy)- estrone sulfate); available from Pharmacia & Upjohn, Peapack, NJ, under the tradename Ogen and from Women First Health Care, Inc., San Diego, CA, under the tradename Ortho-Est; and

(e) conjugated estrogens (17 α-dihydroequilin, 17 α-estradiol, and 17 β-dihydroequilin); available from Wyeth-Ayerst Pharmaceuticals, Philadelphia, PA, under the tradename Premarin.

Progestins and estrogens may also be administered with a variety of dosages, generally from about 0.05 to about 2.0 mg progestin and about 0.001 mg to about 2 mg estrogen, desirably from about 0.1 mg to about 1 mg progestin and about 0.01 mg to about 0.5 mg estrogen. Examples of progestin and estrogen combinations that may vary in dosage and regimen include:

(a) the combination of estradiol (estra-1, 3, 5 (10)-triene-3, 17 β-diol hemihydrate) and norethindrone (17 β-acetoxy-19-nor-17 α-pregn-4-en-20-yn-3-one); which is available from Pharmacia & Upjohn, Peapack, NJ, under the tradename Activella;

(b) the combination of levonorgestrel (d(-)-13 β-ethyl-17 α-ethinyl-17 β-hydroxygon- 4-en-3-one) and ethinyl estradial; available from Wyeth-Ayerst under the tradename Alesse, from Watson Laboratories, Inc., Corona, CA, under the tradenames Levora and Trivora, Monarch Pharmaceuticals, under the tradename Nordette, and from Wyeth-Ayerst under the tradename Triphasil;

(c) the combination of ethynodiol diacetate (19-nor-17 α-pregn-4-en-20-yne-3 β, 17-diol diacetate) and ethinyl estradiol; available from G.D. Searle & Co., Chicago, IL, under the tradename Demulen and from Watson under the tradename Zovia;

(d) the combination of desogestrel (13-ethyl-11- methylene-18,19-dinor-17 α-pregn- 4-en- 20-yn-17-ol) and ethinyl estradiol; available from

Organon under the tradenames Desogen and Mircette, and from Ortho-McNeil Pharmaceutical, Raritan, NJ, under the tradename Ortho-Cept;

(e) the combination of norethindrone and ethinyl estradiol; available from Parke-Davis, Morris Plains, NJ, under the tradenames Estrostep and FemHRT, from Watson under the tradenames Microgestin, Necon, and Tri-Norinyl, from Ortho-McNeil under the tradenames Modicon and Ortho-Novum, and from Warner Chilcott Laboratories, Rockaway, NJ, under the tradename Ovcon;

(f) the combination of norgestrel ((±)-13-ethyl-17-hydroxy-18, 19-dinor-17 α-preg-4-en-20-yn-3-one) and ethinyl estradiol; available from Wyeth-Ayerst under the tradenames Ovral and Lo/Ovral, and from Watson under the tradenames Ogestrel and Low-Ogestrel;

(g) the combination of norethindrone, ethinyl estradiol, and mestranol (3-methoxy-19-nor-17 α-pregna-1,3,5(10)-trien-20-yn-17-ol); available from Watson under the tradenames Brevicon and Norinyl;

(h) the combination of 17 β-estradiol (estra-1,3,5(10)-triene-3,17 β-diol) and micronized norgestimate (17 α-17-(Acetyloxyl)-13-ethyl-18,19-dinorpregn-4-en-20-yn-3-one3-oxime); available from Ortho-McNeil under the tradename Ortho-Prefest;

(i) the combination of norgestimate (18,19-dinor-17-pregn-4-en-20-yn-3-one, 17--(acetyloxy)-13-ethyl-,oxime, (17(α)-(+)-) and ethinyl estradiol; available from Ortho-McNeil under the tradenames Ortho Cyclen and Ortho Tri-Cyclen; and

(j) the combination of conjugated estrogens (sodium estrone sulfate and sodium equilin sulfate) and medroxyprogesterone acetate (20-dione, 17-(acetyloxy)-6-methyl-, (6(α))- pregn-4-ene-3); available from Wyeth-Ayerst under the tradenames Premphase and Prempro.

In general, a dosage of progestins may vary from about .05 mg to about 10 mg or up to about 200 mg if microsized progesterone is administered. Examples of progestins include norethindrone; available from ESI Lederle, Inc., Philadelphia, PA, under the tradename Aygestin, from Ortho-McNeil under the tradename Micronor, and from Watson under the tradename Nor-QD; norgestrel; available from Wyeth-Ayerst under the tradename Ovrette; micronized progesterone (pregn-4-ene-3, 20-dione);

available from Solvay under the tradename Prometrium; and medroxyprogesterone acetate; available from Pharmacia & Upjohn under the tradename Provera.

In another alternative embodiment, the compositions, therapeutic combinations or methods of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more obesity control medications. Useful obesity control medications include, but are not limited to, drugs that reduce energy intake or suppress appetite, drugs that increase energy expenditure and nutrient-partitioning agents. Suitable obesity control medications include, but are not limited to, noradrenergic agents (such as diethylpropion, mazindol, phenylpropanolamine, phentermine, phendimetrazine, phendamine tartrate, methamphetamine, phendimetrazine and tartrate); serotonergic agents (such as sibutramine, fenfluramine, dexfenfluramine, fluoxetine, fluvoxamine and paroxtine); thermogenic agents (such as ephedrine, caffeine, theophylline, and selective β3-adrenergic agonists); alpha-blocking agents; kainite or AMPA receptor antagonists; leptin-lipolysis stimulated receptors; phosphodiesterase enzyme inhibitors; compounds having nucleotide sequences of the mahogany gene; fibroblast growth factor-10 polypeptides; monoamine oxidase inhibitors (such as befloxatone, moclobemide, brofaromine, phenoxathine, esuprone, befol, toloxatone, pirlindol, amiflamine, sercloremine, bazinaprine, lazabemide, milacemide and caroxazone); compounds for increasing lipid metabolism (such as evodiamine compounds); and lipase inhibitors (such as orlistat). Generally, a total dosage of the above-described obesity control medications can range from 1 to 3,000 mg/day, desirably from about 1 to 1,000 mg/day and more desirably from about 1 to 200 mg/day in single or 2-4 divided doses.

The compositions, therapeutic combinations or methods of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more blood modifiers which are chemically different from the substituted azetidinone and substituted β-lactam compounds (such as compounds II-XIII above) and the lipid modulating agents discussed above, for example, they

contain one or more different atoms, have a different arrangement of atoms or a different number of one or more atoms than the sterol absorption inhibitor(s) or lipid modulating agents discussed above. Useful blood modifiers include but are not limited to anti-coagulants (argatroban, bivalirudin, dalteparin sodium, desirudin, dicumarol, lyapolate sodium, nafamostat mesylate, phenprocoumon, tinzaparin sodium, warfarin sodium); antithrombotic (anagrelide hydrochloride, bivalirudin, cilostazol, dalteparin sodium, danaparoid sodium, dazoxiben hydrochloride, efegatran sulfate, enoxaparin sodium, fluretofen, ifetroban, ifetroban sodium, lamifiban, lotrafiban hydrochloride, napsagatran, orbofiban acetate, roxifiban acetate, sibrafiban, tinzaparin sodium, trifenagrel, abciximab, zolimomab aritox); fibrinogen receptor antagonists (roxifiban acetate, fradafiban, orbofiban, lotrafiban hydrochloride, tirofiban, xemilofiban, monoclonal antibody 7E3, sibrafiban); platelet inhibitors (cilostazol, clopidogrel bisulfate, epoprostenol, epoprostenol sodium, ticlopidine hydrochloride, aspirin, ibuprofen, naproxen, sulindae, idomethacin, mefenamate, droxicam, diclofenac, sulfinpyrazone, piroxicam, dipyridamole); platelet aggregation inhibitors (acadesine, beraprost, beraprost sodium, ciprostene calcium, itazigrel, lifarizine, lotrafiban hydrochloride, orbofiban acetate, oxagrelate, fradafiban, orbofiban, tirofiban, xemilofiban); hemorrheologic agents (pentoxifylline); lipoprotein associated coagulation inhibitors; Factor VIIa inhibitors (4H-31-benzoxazin-4-ones, 4H-3,1-benzoxazin-4-thiones, quinazolin-4-ones, quinazolin-4-thiones, benzothiazin-4-ones, imidazolyl-boronic acid-derived peptide analogues TFPI-derived peptides, naphthalene-2-sulfonic acid {1-[3-(aminoiminomethyl)-benzyl]-2-oxo-pyrrolidin-3-(S)-yl} amide trifluoroacetate, dibenzofuran-2-sulfonic acid {1-[3-(aminomethyl)-benzyl]-5-oxo-pyrrolidin-3-yl}-amide, tolulene-4-sulfonic acid {1-[3-(aminoiminomethyl)-benzyl]-2-oxo-pyrrolidin-3-(S)-yl}-amide trifluoroacetate, 3,4-dihydro-1H-isoquinoline-2-sulfonic acid {1-[3-(aminoiminomethyl)-benzyl]-2-oxo-pyrrolin-3-(S)-yl}-amide trifluoroacetate); Factor Xa inhibitors (disubstituted pyrazolines, disubstituted triazolines, substituted n-[(aminoiminomethyl)phenyl] propylamides, substituted n-[(aminomethyl)phenyl] propylamides, tissue factor pathway inhibitor (TFPI), low molecular weight heparins, heparinoids, benzimidazolines, benzoxazolinones, benzopiperazinones, indanones, dibasic (amidinoaryl)

propanoic acid derivatives, amidinophenyl-pyrrolidines, amidinophenyl-pyrrolines, amidinophenyl-isoxazolidines, amidinoindoles, amidinoazoles, bis-arlysulfonylaminobenzamide derivatives, peptidic Factor Xa inhibitors).

The compositions, therapeutic combinations or methods of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more cardiovascular agents which are chemically different from the substituted azetidinone and substituted β-lactam compounds (such as compounds II-XIII above) and the lipid modulating agents discussed above, for example, they contain one or more different atoms, have a different arrangement of atoms or a different number of one or more atoms than the sterol absorption inhibitor(s) or PPAR receptor activators discussed above. Useful cardiovascular agents include but are not limited to calcium channel blockers (clentiazem maleate, amlodipine besylate, isradipine, nimodipine, felodipine, nilvadipine, nifedipine, teludipine hydrochloride, diltiazem hydrochloride, belfosdil, verapamil hydrochloride, fostedil); adrenergic blockers (fenspiride hydrochloride, labetalol hydrochloride, proroxan, alfuzosin hydrochloride, acebutolol, acebutolol hydrochloride, alprenolol hydrochloride, atenolol, bunolol hydrochloride, carteolol hydrochloride, celiprolol hydrochloride, cetamolol hydrochloride, cicloprolol hydrochloride, dexpropranolol hydrochloride, diacetolol hydrochloride, dilevalol hydrochloride, esmolol hydrochloride, exaprolol hydrochloride, flestolol sulfate, labetalol hydrochloride, levobetaxolol hydrochloride, levobunolol hydrochloride, metalol hydrochloride, metoprolol, metoprolol tartrate, nadolol, pamatolol sulfate, penbutolol sulfate, practolol, propranolol hydrochloride, sotalol hydrochloride, timolol, timolol maleate, tiprenolol hydrochloride, tolamolol, bisoprolol, bisoprolol fumarate, nebivolol); adrenergic stimulants; angiotensin converting enzyme (ACE) inhibitors (benazepril hydrochloride, benazeprilat, captopril, delapril hydrochloride, fosinopril sodium, libenzapril, moexipril hydrochloride, pentopril, perindopril, quinapril hydrochloride, quinaprilat, ramipril, spirapril hydrochloride, spiraprilat, teprotide, enalapril maleate, lisinopril, zofenopril calcium, perindopril erbumine); antihypertensive agents (althiazide, benzthiazide, captopril, carvedilol, chlorothiazide sodium,

clonidine hydrochloride, cyclothiazide, delapril hydrochloride, dilevalol hydrochloride, doxazosin mesylate, fosinopril sodium, guanfacine hydrochloride, methyldopa, metoprolol succinate, moexipril hydrochloride, monatepil maleate, pelanserin hydrochloride, phenoxybenzamine hydrochloride, prazosin hydrochloride, primidolol, quinapril hydrochloride, quinaprilat, ramipril, terazosin hydrochloride, candesartan, candesartan cilexetil, telmisartan, amlodipine besylate, amlodipine maleate, bevantolol hydrochloride); angiotensin II receptor antagonists (candesartan, irbesartan, losartan potassium, candesartan cilexetil, telmisartan); anti-anginal agents (amlodipine besylate, amlodipine maleate, betaxolol hydrochloride, bevantolol hydrochloride, butoprozine hydrochloride, carvedilol, cinepazet maleate, metoprolol succinate, molsidomine, monatepil maleate, primidolol, ranolazine hydrochoride, tosifen, verapamil hydrochloride); coronary vasodilators (fostedil, azaclorzine hydrochloride, chromonar hydrochloride, clonitrate, diltiazem hydrochloride, dipyridamole, droprenilamine, erythrityl tetranitrate, isosorbide dinitrate, isosorbide mononitrate, lidoflazine, mioflazine hydrochloride, mixidine, molsidomine, nicorandil, nifedipine, nisoldipine, nitroglycerine, oxprenolol hydrochloride, pentrinitrol, perhexiline maleate, prenylamine, propatyl nitrate, terodiline hydrochloride, tolamolol, verapamil); diuretics (the combination product of hydrochlorothiazide and spironolactone and the combination product of hydrochlorothiazide and triamterene).

The compositions, therapeutic combinations or methods of the present invention can comprise at least one compound of Formula (I), or pharmaceutically acceptable salts, solvates, or esters thereof, and one or more antidiabetic medications for reducing blood glucose levels in a human. Useful antidiabetic medications include, but are not limited to, drugs that reduce energy intake or suppress appetite, drugs that increase energy expenditure and nutrient-partitioning agents. Suitable antidiabetic medications include, but are not limited to, sulfonylurea (such as acetohexamide, chlorpropamide, gliamilide, gliclazide, glimepiride, glipizide, glyburide, glibenclamide, tolazamide, and tolbutamide), meglitinide (such as repaglinide and nateglinide), biguanide (such as metformin and buformin), alpha-glucosidase inhibitor (such as acarbose, miglitol, camiglibose, and voglibose), certain peptides (such as amlintide, pramlintide, exendin, and

GLP-1 agonistic peptides), and orally administrable insulin or insulin composition for intestinal delivery thereof. Generally, a total dosage of the above-described antidiabetic medications can range from 0.1 to 1,000 mg/day in single or 2-4 divided doses.

Mixtures of two, three, four or more of any of the pharmacological or therapeutic agents described above can be used in the compositions and therapeutic combinations of the present invention.

Since the present invention relates to treating conditions as discussed above, by treatment with a combination of active ingredients wherein the active ingredients may be administered separately, the invention also relates to combining separate pharmaceutical compositions in kit form. That is, a kit is contemplated wherein two separate units are combined: a pharmaceutical composition comprising at least one selective CB_1 receptor antagonist of Formula (I), or a pharmaceutically acceptable salt, solvate, or ester thereof, and a separate pharmaceutical composition comprising at least one cholesterol lowering compound as described above. The kit will preferably include directions for the administration of the separate components. The kit form is particularly advantageous when the separate components must be administered in different dosage forms (e.g., oral and parenteral) or are administered at different dosage intervals.

In yet another embodiment, the present invention provides a method of treating, reducing, or ameliorating a disease or condition selected from the group consisting of metabolic syndrome, obesity, waist circumference, lipid profile, insulin sensitivity, neuroinflammatory disorders, cognitive disorders, psychosis, addictive behavior, gastrointestinal disorders, vascular conditions, hyperlipidaemia, atherosclerosis, hypercholesterolemia, sitosterolemia, vascular inflammation, stroke, diabetes, and cardiovascular conditions, and/or reduce the level of sterol(s) in a patient in need thereof, comprising administering to said patient an effective amount of at least one compound of Formula (I), or a pharmaceutically acceptable salt, solvate, or ester thereof, and one or more cholesterol lowering compound.

The treatment compositions and therapeutic combinations comprising at least one compound of Formula (I) and at least one cholesterol lowering agent can inhibit the intestinal absorption of cholesterol in mammals can be

useful in the treatment and/or prevention of conditions, for example vascular conditions, such as atherosclerosis, hypercholesterolemia and sitosterolemia, stroke, obesity and lowering of plasma levels of cholesterol in mammals, in particular in mammals.

In another embodiment of the present invention, the compositions and therapeutic combinations of the present invention can inhibit sterol or 5α-stanol absorption or reduce plasma concentration of at least one sterol selected from the group consisting of phytosterols (such as sitosterol, campesterol, stigmasterol and avenosterol) and/or 5α-stanol (such as cholestanol, 5α-campestanol, 5α-sitostanol), cholesterol and mixtures thereof. The plasma concentration can be reduced by administering to a mammal in need of such treatment an effective amount of at least one treatment composition or therapeutic combination comprising at least one selective CB_1 receptor antagonist and at least one cholesterol lowering compound, for example a sterol absorption inhibitor described above. The reduction in plasma concentration of sterols or 5α-stanols can range from about 1 to about 70 percent, and preferably about 10 to about 50 percent. Methods of measuring serum total blood cholesterol and total LDL cholesterol are well known to those skilled in the art and for example include those disclosed in PCT WO 99/38498 at page 11, incorporated by reference herein. Methods of determining levels of other sterols in serum are disclosed in H. Gylling et al., "Serum Sterols During Stanol Ester Feeding in a Mildly Hypercholesterolemic Population", J. Lipid Res. 40: 593-600 (1999), incorporated by reference herein.

The treatments of the present invention can also reduce the size or presence of plaque deposits in vascular vessels. The plaque volume can be measured using (IVUS), in which a tiny ultrasound probe is inserted into an artery to directly image and measure the size of atherosclerotic plaques, in a manner well know to those skilled in the art.

In the Schemes and Experiments below, as well as the specification and claims, the following abbreviations are applicable:

rt	room temperature
THF	tetrahydrofuran
Et_2O	ethyl ether


Me	methyl
Et	ethyl
Bu	Butyl
Bn	benzyl
Pr	propyl
i-Pr	isopropyl
Ac	acetyl
EtOAc	ethyl acetate
$BnOCH_2Cl$	benzylchloromethylether
BuLi	Butyl Lithium
DBAD	Di-*tert*-butyl azodicarboxylate
DBU	1,8-diazabicylco[5.4.0]undec-7-ene
DCE	1,2-dichloroethane
DCM	Dichloromethane
DMF	*N,N* -Dimethylformamide
DMSO	Methyl sulfoxide
HOBT or HOBt	Hydroxybenzotriazole
KHMDS	Potassium bis(trimethylsilyl)amide
LiHMDS or LHMDS:	Lithium bis(trimethylsilyl)amide
$NaB(O_2CCH_3)_3H$	Sodium triacetoxyborohydride
OTf	Triflate or trifluoromethanesulfonate (CF_3SO_2-O-)
PhSeBr	Phenyl selenium bromide
PS	Polymer supported
PS-EDC	Polymer supported dimethyl aminopropyl ethylcarbodiimide hydrochloride
Py	Pyridine
PS-NCO	Polymer supported isocyanate
PS-Tris-NH_2	Polymer supported trisamine
TFA	Trifluoroacetic acid
$Ti(OiPr)_4$	titanium isopropoxide;
TLC	thin layer chromatography
TMSI	Trimethylsilyl iodide or iodotrimethylsilane
HATU	O-(7-Azabenzotriazole-1-yl)-N,N,N'N'-tetramethyluronium hexafluorophosphate
DIPEA	*N,N*-diisopropylethylamine
dppp	1,3-bis(diphenylphosphino)propane
dppf	1,1'-bis(diphenylphosphino)ferrocene
PS-BEMP	2-tert-Butylimino-2-diethylamino-1,3-dimethyl-perhydro-1,3,2-diaza-phosphorine on polystyrene
DMAP:	N,N-dimethylaminopyridine
EDCI:	N(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride
HPLC:	High Performance Liquid Chromatography
PhMe:	Toluene
MeOH:	Methyl alcohol
EtOH:	Ethyl alcohol
Trisyl Azide:	2,4,6-tri-isopropylbenzenesulfonyl azide
DEAD:	Diethylazodicarboxylate
NMP:	N-Methylpyrrolidinone

DMPU: 1,3-Dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone
Dess-Martin reagent: 1,1,1-Tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3(1H)-one
BOM: Benzyloxymethyl
Cbz: Benzyloxycarbonyl
MOM: Methoxymethyl
Tf: Trifluoromethanesufonyl
TIPS: Triisopropylsilyl
TBS or TBDMS: tert-Butyldimethylsilyl

EXAMPLES

Preparation of Di(hetero)aryl-isoindol-1-ones

General Scheme A:



Diaryl-isoindol-1-ones, diheteroaryl-isoindol-1-ones, and aryl-heteroaryl-isoindol-1-ones of Formula (I) can be prepared by a variety of methods, for example by condensing an aryl or heteroaryl substituted diene acid or acid chloride with an aryl or heteroaryl substituted unsaturated amine to form a triene amide, or alternatively condensing a diaryl-, diheteroaryl- or aryl-heteroaryl-substituted diene acid or acid chloride with an unsaturated amine, then cyclizing the resulting triene amide via an intramolecular Diels-

Alder reaction to form a di(hetero)aryl-tetrahydro-isoindol-1-one that can be further modified (e.g., by reduction or alklation, etc.) as desired.


Aldehyde **a** can be reacted with crotyl phosphonate to provide ester **b** which on saponification followed by reaction with oxalyl chloride gives acid chloride **c**. Sonagashira coupling [K. Sonogashira et al, Tet, Lett., 4467, (1975)] of aryl or heteroaryl bromide or iodide or chloride or *O*-triflate **d** with propargyl amine gives alkynyl amine **e**, which is reduced to allyl amine **f** using Lindlar catalyst and hydrogen. Allyl amine **f** is coupled with acid chloride **c** to provide amide **g**. Amide **g** is subjected to Diels-Alder reaction conditions, and then treated with DBU (i.e., 1,8-diazabicyclo[5.4.0]undec-7-ene) to provide cyclization products **h** and **i**. Double bond reduction gives **j** and **k**.

Unless expressly indicated otherwise, when compounds or mixtures in any of the following preparations are purified by chromatography, said chromatography is conventional flash chromatography using a silica gel staionary phase.

Preparation of Compounds **1-4**

Step 1


Scheme 1:



To a suspension of 60% NaH in mineral oil (16 g) in 1L THF was added triethyl 4-phosphonocrotonate (100 g). The resulting solution was stirred for 2 hr, and then a solution of 2,4-dichlorobenzaldehyde (54 g) in 200 mL THF was added thereto. The mixture was stirred at room temperature for 1 hr. The reaction was quenched by the addition of 1L of aq. NH_4Cl and the THF was evaporated. The mixture was extracted with 4x200 ml ethyl acetate and the combined organic layer was washed with water, brine and dried over $MgSO_4$ then evaporated to provide the crude product. This was purified by silica gel chromatography using 5% ethyl acetate-hexanes to give 25.9 g of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid ethyl ester. MS: m/e 271.04 (MH^+)

Step 2


Scheme 2:



To a solution of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid ethyl ester (25.5 g) in 100 mL each of MeOH and THF was added a solution of KOH (16 g) in 100 mL H_2O and the mixture was stirred at room temperature for 2 hr. The mixture was diluted with 300 mL H_2O, acidified with 1N HCl and the precipitated product was isolated to provide 23.3 g of acid. To a suspension of the acid (19.5 g) in 400 mL CH_2Cl_2 at room temperature was added $(COCl)_2$ (20.7 mL) followed by DMF (190 μL) and the mixture was stirred at room temperature for 3 hr. The resultant clear solution was concentrated to provide 5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl chloride. NMR (300 MHz, $CDCl_3$) 7.60 (ddd, J = 15.0, 11.2, 0.8 Hz, 1H), 7.54 (d, J = 8.4 Hz, 1H), 7.42-7.38 (m, 2H), 7.25-7.21 (m, 1H), 6.85 (ddd, J = 15.4, 11.2, 0.8 Hz), 6.20 (d, J = 14.8 Hz).

Step 3


Scheme 3:



To a solution of 4-chloro iodobenzene (25 g) in 400 mL CH_2Cl_2 at room temperature was added propargyl amine (13.5 mL), diisopropyl amine (37 mL), CuI (4 g) and $Pd(PPh_3)_2Cl_2$ (3.7 g). The mixture was stirred overnight at room temperature, diluted with 600 mL EtOAc and filtered through a CELITE pad to remove insoluble materials. The solution was washed with water, brine, dried over $MgSO_4$, concentrated and purified by chromatography to provide 14.1 g of 3-(4-chloro-phenyl)-prop-2-ynylamine. MS: m/e 166.1 (MH^+)

Step 4


Scheme 4:



To a solution of 3-(4-chloro-phenyl)-prop-2-ynylamine (14.0 g) in 200 mL each of MeOH and CH_2Cl_2 was added Et_3N (1.2 mL) and Lindlar catalyst (1.4 g) and the resulting suspension was stirred under a H_2 balloon. The reaction was followed by TLC (i.e., thin layer chromatography) using 6% methanol-dichloromethane as eluent and once completed, filtered through a CELITE pad, concentrated and purified by chromatography to provide 10.5 g of 3-(4-chloro-phenyl)-allylamine. MS: m/e 168.6 (MH^+)

Step 5


Scheme 5:



To a solution of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl chloride (62.5 mmol) in 300 mL CH_2Cl_2 at 0°C was added Et_3N (13.0 mL) followed by DMAP (i.e., 4-diaminomethylpyridine) (760 mg). To this was added a solution of 3-(4-chloro-phenyl)-allylamine (10.5 g) in 30 mL CH_2Cl_2. The mixture was stirred at 0°C for 1 hr. Then 2 mL of MeOH was added, and the solution was stirred for 10 min. The solution was then diluted with aq. $NaHCO_3$ and extracted with EtOAc. Filtration of solids and concentration of EtOAc layers provided a combined yield of 24.6 g of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide. MS: m/e 392.2 (MH^+)


Step 6

Scheme 6:




A solution of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide (13.6 g) in 1:9 MeOH-CH_2Cl_2 was filtered through a pad of alumina and concentrated. This solid was taken up in 300 mL of toluene and heated at 185°C for 10 hr. It was concentrated and stirred with DBU (i.e., 1,8-diazabicyclo[5.4.0]undec-7-ene) (1g) in 200 mL CH_2Cl_2 for 1 hr. The mixture was concentrated and chromatographed with 75:25 ethyl acetate-hexanes to provide 8.5 g of a mixture of cyclization products, **1** and **2**. The products **3** and **4** were also isolated from these reactions.

One of skill in the art will also recognize that when compounds in the any of the preparative procedures described herein are prepared from racemic starting materials, e.g., as in the preparation of compounds **1** and **2** described above, the resulting products (e.g., compounds **1** and **2**) are also racemic mixtures of enantiomers. For example, compound **1**:



1

is understood to represent a racemic mixture of the following enantiomers:




One of skill in the art will also recognize that racemic mixtures of products prepared from racemic starting materials can be separated into the individual enantiomers, e.g., by chiral chromatographic methods. Alternatively, specific enantiomers can be prepared by chiral synthetic methods from chiral starting materials.

The structures disclosed herein are intended to represent relative stereochemistry, i.e., both racemic mixtures and individual enantiomers, rather than absolute stereochemistry, unless there is an express indication that the structure is either intended to represent only a racemic mixture (e.g., by labeling the structure as a "racemic mixture", "D/L", or "+/-") or that the structure is intended to represent a single enantiomer having the indicated absolute stereochemistry (e.g., by expressly labeling the structure with "absolute stereochemistry" or labeling chiral centers with the "R" or "S" designations of the well-known Cahn-Ingold-Prelog system).

Preparation of Compounds **5** and **6**


Scheme 7:



The above mixture of **1** and **2** was taken up in 200 mL EtOAc and stirred with 800 mg of $PtO_2 \cdot H_2O$ under a H_2 balloon for 1.5 hr. The mixture was then filtered through a CELITE pad and concentrated. The solution was taken up in a minimum amount of CH_2Cl_2 and diluted with Et_2O and the precipitated solid was filtered to provide 1.93 g of **6**. The filtrate was concentrated and purified by chromatography to provide 1.99 g of **5**.

Chiral Resolution of **5**

Scheme 8:



Compound **5** (25-50 mg) was sonicated in about 1.5 mL of isopropyl alcohol for one minute. After allowing the solution to stand for 30 minutes, about 2.5 mL of hexanes was added and the mixture gently stirred. Once the mixture was allowed to settle for several minutes, the clear solution was decanted off and, after filtration, injected onto a Chiralpac AD preparative HPLC column (5 cm x 50 cm) and eluted with 15% isopropyl alcohol in hexanes to obtain **5a** and **5b** in roughly equal amounts. This process was repeated until the desired quantity of each isomer was obtained. Detection was at 220 nm, and the flow rate was 100 mL/min.


MS for **5a**: 394.2 (MH^+)

MS for **5b**: 394.2 (MH^+)

Preparation of Compound **7**

Step 1

Scheme 9:




To a solution of 3-(4-chloro-phenyl)-prop-2-ynylamine (2.5 g) in 50 mL of THF at room temperature was added a 1M solution of $LiAlH_4$ in THF (30 mL) and the mixture was heated at reflux for 1 hr. It was cooled to room temperature, and to it was successively added 1.1 mL H_2O, 1.1 mL of 15% aq. NaOH and 3.4 mL H_2O. The precipitated solid was filtered off and the filtrate was concentrated and purified by chromatography to provide 1.45 g of 3-(4-chloro-phenyl)-allylamine. NMR (400 MHz, CDCl3) 7.26-7.21 (m, 4H),

6.42 (dt, J = 16.0, 1.6 Hz, 1H), 6.29-6.22 (m, 1H), 3.44 (dd, J = 5.6, 1.6 Hz, 2H)

Step 2


Scheme 10:



To a solution of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl chloride (7.6 mmol) in 30 mL CH_2Cl_2 at 0°C was added Et_3N (1.6 mL), DMAP (47 mg) followed by a solution of 3-(4-chloro-phenyl)-allylamine (1.4 g) in 10 mL of CH_2Cl_2. The solution turned into a thick slurry after few minutes. It was stirred for 30 min at 0°C, 30 min at room temperature then diluted with H_2O. The solid was filtered and washed with H_2O followed by Et_2O then dried in a vacuum oven to provide 3.8g of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide. MS: m/e 394.02 (MH^+)

Step 3

Scheme 11:




7

A suspension of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide (3.8 g) in 70 mL of toluene was heated in a sealed tube at 185°C for 6 hr, cooled to room temperature and concentrated to provide 2.8 g of **7**. MS: m/e 394.0 (MH^+)

Preparation of Compound **8**


Scheme 12:



To a solution of **7** (2.8 g) in 70 mL of 1:9 MeOH-CH_2Cl_2 was added $PtO_2 \cdot H_2O$ (280 mg) and the mixture was stirred under a H_2 balloon for 40 min. To the suspension was added activated carbon, then the suspension was filtered through a CELITE pad, concentrated and recrystallized from CH_2Cl_2/hexanes to provide 1.4 g of **8**. MS: m/e 394.03 (MH^+)


Preparation of Compounds **9-12**

Scheme 13:



The lactam **5** can be N-alkylated using standard conditions. For example, **5** was N-allylated to give **9** using NaH and allyl iodide. Oxidative cleavage of the double bond followed by reduction or reductive amination gave **11** and **12** respectively as shown in Scheme 12.

Preparation of Compounds **13** and **14**




13 **14**

Compounds **13** and **14** were prepared by the method of Scheme A using the appropriate starting aldehyde **a** and iodide **d**.

It will be recognized by one of ordinary skill in the art that the substituted aryl moieties of the above isoindol-1-ones can be independently replaced with heteroaryl moieties by the appropriate selection of heteroaryl substituted starting materials. For example, a benzaldehyde and/or phenyl iodide could be replaced with the corresponding pyridyl aldehyde or pyridyl halide.

Preparation of Compound **15**

Scheme 14




To a solution of **5** (50 mg) in 5 mL DMF at room temperature was added NaH (1.1 equivalent of a 60% dispersion in mineral oil). The solution was then stirred for 20 min. To the stirred solution was added α-bromo-tolunitrile (75 mg) and Bu_4NI (5 mg). After 3 hrs of stirring, an additional portion of NaH (20 mg) was added and the mixture was stirred overnight at room temperature. The solution was then diluted with EtOAc, washed with water, 1N HCl, brine, dried over $MgSO_4$, filtered and concentrated. The crude product was chromatographed using 10% MeOH in CH_2Cl_2 mixture to provide 40 mg of **15**.

MS: 509.3 (MH^+)

Preparation of Compound **16**

Scheme 15




To a solution of **5** (50 mg) in 5 mL DMF at room temperature was added NaH (6 mg of 60% dispersion in mineral oil). The solution was stirred for 20 min. To the stirred solution was added 3 equivalents of CH_3I and the mixture was stirred overnight at room temperature. An additional portion of NaH (20 mg) was then added, and the mixture was heated at 50°C for 1 day. The solution was then poured into EtOAc and washed with water, 1N HCl and brine, dried over $MgSO_4$, filtered and evaporated then purified by chromatography using 5% MeOH in CH_2Cl_2 to provide 31 mg of **16**.


MS: 406.2 (MH^+)

Preparation of N-Substituted-4,5-diphenyl-isoindoles


General Scheme B:



For example, compounds **19** and **20**, below, were prepared by reacting compound **17** with methylsulfonyl chloride or cyclopropylsulfonyl chloride, respectively.




19







20

Preparation of Compounds **21-36**


The following are representative compounds of structure **18A** prepared by reacting isoindole **17** with the appropriate carboxylic acid using well-known EDCI and HOBt coupling conditions.






18A

Example	R	MS (MH^+)
21		478.3
22		491.3
23		498.3
24		509.3
25		509.3


The following are representative compounds of structure **18C** prepared by reacting isoindole **17** with the appropriate isocyanate.





18C

Compound	R	MS (MH^+)
26		505.3
27		533.3
28		567.3


The following are representative compounds of structure **18D** prepared by reacting isoindole **17** with the appropriate chloroformate.








18D

Compound	R	MS (MH^+)
29	$-CH_3$	438.2
30		534.3
31		500.3

The following are some representative compounds of structure **18E** prepared by reacting isoindole **17** with the appropriate aldehyde or ketone and sodium triacetoxyborohydride as the reducing agent.



18E

Compound	R	MS (MH^+)
32		462.3
33		495.3
34		495.3
35		509.3
36		509.3

It will be recognized by one of ordinary skill in the art that the substituted aryl moieties of the above isoindoles can be independently replaced with heteroaryl moieties by the use of the appropriate heteroaryl substituted starting materials.


It will also be recognized by one of ordinary skill in the art that isoindoles having different substitution patterns of aryl and/or heteroaryl groups on the six-membered ring may be provided by use of the appropriate starting materials. In addition, substitution on the isoindole nitrogen atom may be provided by methods known in the art (e.g., alkylation, acylation, arylation, etc.).

Preparation of Di(hetero)aryl-isoindol-1-ylideneamines

Di(hetero)aryl-isoindol-1-ylideneamines can be prepared by a variety of methods, for example by conversion of an isoindol-1-one to the corresponding amidine, followed by further modification of the amidine, e.g., by acylation.

Preparation of Compounds **37-39**

Scheme 16:




The lactam **5** was converted to amidine **37** which can be coupled with the appropriate carboxylic acid using EDCI/HOBt to provide **37A**.

Step 1:

To a solution of 89 mg of **5** in about 3 mL of dichloromethane was added triethyloxonium tetrafluoroborate (2 eq.) and sodium carbonate (2 eq.) and the mixture was stirred under nitrogen for three days. The reaction mixture was poured onto a pH 7 buffer and extracted with dichloromethane three times. The combined extracts were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. To the residue in methanol (~4 mL) was added ammonium chloride (10 eq.) and the mixture heated to reflux. After 24 hours the mixture was evaporated to dryness and partitioned between dichloromethane and aq. K_2CO_3. The aqueous phase was twice extracted with dichloromethane. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by HPLC (C-18 column, elution with water/acetonitrile w/0.1% formic acid) yielded **37** as the formate salt.

MS: 393.2 (MH^+)

Preparation of **38** and **39**



To 20 mg of **37** (free base) in dichloromethane (1.5 mL) at 0°C was added 4-cyanobenzoic acid (2 eq.), HOBt (2 eq.) and EDCI (2 eq.) and the mixture was stirred under nitrogen while slowly warming to room temperature.

After 24 hours the mixture was poured onto aqueous sodium bicarbonate and extracted three times with ~1% methanol in dichloromethane. The combined extracts were dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (0-50% ethylacetate in hexane) yielded 16 mg of **38**.


Compound **39** was prepared using similar conditions, except that benzoic acid was used instead of 4-cycanobezoic acid.

MS for **38** : 522.3 (MH^+)


MS for **39** : 497.3 (MH^+)

It will be recognized by one of ordinary skill in the art that the substituted aryl moieties of the above isoindol-1-ylideneamines can be independently replaced with heteroaryl moieties by the appropriate selection of heteroaryl substituted isoindol-1-one starting materials, followed by conversion of the isoindol-1-one to an isoindol-1-ylideneamines, for example as shown above. Furthermore, other substitution patterns on the 6-membered ring of the isoindoles may be obtained by modification of the appropriate isoindol-1-one starting material (e.g., use of a 5,6-di(hetero)aryl-isoindol-1-one starting material rather than a 4,5-di(hetero)aryl-isoindol-1-one).


Preparation of Compounds **40-43**




40 MS: 476.3 (MH^+)



41 MS: 492.3 (MH^+)



42 MS: 575.3 (MH^+)




43 MS: 607.3 (MH^+)

Compound **37** was coupled with cyclobutanecarboxylic acid or 4-methylvaleric acid under conditions similar to those used to prepare compounds **38** and **39** using EDCI and HOBt. The above products (**40-43**) were obtained.

It will be recognized by one of ordinary skill in the art that the substituted aryl moieties of the above isoindol-1-ylideneamines can be independently replaced with heteroaryl moieties by the appropriate selection of heteroaryl substituted starting materials, followed by conversion of the isoindol-1-one to an isoindol-1-ylideneamines, for example as shown above. Furthermore, other substitution patterns on the 6-membered ring of the isoindoles may be obtained by modification of the appropriate isoindol-1-one starting material (e.g., use of a 5,6-di(hetero)aryl-isoindol-1-one starting material rather than a 4,5-di(hetero)aryl-isoindol-1-one).

Alternative Preparation of N-Benzyl-4,5-di(hetero)aryl-isoindol-1-ones

Scheme 17:




Alternatively, the group attached to the N atom of the desired 4,5-diphenyl-isoindol-1-one was inserted prior to the Diels-Alder reaction, for example as shown in Scheme 17, above. The propargyl amine (3,4-difluoro-benzyl)-prop-2-ynyl-carbamic acid tert-butyl ester was coupled with either 4-chloro iodobenzene or 2-bromo-5-chloro pyridine under Sonagashira conditions to provide **l-a/l-b**. The resulting alkyne was reduced to the *cis*-olefin (**m-a/m-b**) and the Boc group was cleaved under acidic conditions (**n-a/n-b**). The amine was coupled with 5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl chloride and the Diels-Alder precursor **o-a/o-b** was thermally cyclized to provide **44a/44b** and **45a/45b**. The *trans*-lactam **44a/44b** was isomerized to the *cis*-lactam **46a/46b** using DBU, and the double bond was reduced either with H_2/PtO_2 or with $H_2/Rh(PPh_3)_3Cl$.

Alternative Preparation of Compounds **49** and **50**

Step 1

Scheme 18:




To a solution of 3,4-difluorobenzaldehyde (5 g, 35.2 mmol) in 200 mL dichloroethane was added *mono*-propargyl amine (4.85 mL, 70.7 mmol, 2 eq.) followed by $Na(OAc)_3BH$ (9 g, 42.5 mmol) and the mixture was stirred for 2 days at room temperature. The solution was diluted with 100 mL of dichloromethane, washed with 2x100 mL aq. $NaHCO_3$, 100 mL brine, dried over Na_2SO_4, filtered and concentrated. The crude product was chromatographed with 20:80 ethyl acetate-hexanes to provide 3.8 g of (3,4-difluoro-benzyl)-prop-2-ynyl-amine as an oil.

MS: 182.15 (MH^+)

Step 2

Scheme 19:




A mixture of (3,4-difluoro-benzyl)-prop-2-ynyl-amine (3.7g, 20.4 mmol), triethylamine (4.3 mL, 30.9 mmol, 1.5 eq,) and $(Boc)_2O$ (6.7 g, 30.7 mmol, 1.5 eq.) in 100 mL dichloromethane was stirred overnight at room temperature. The solution was diluted with 200 mL ether, washed with 2x100 mL aq. $NaHCO_3$, 100 mL brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 1:9 ethyl acetate-hexanes to provide 6.6 g of (3,4-difluoro-benzyl)-prop-2-ynyl-carbamic acid tert-butyl ester.

MS: 282.13 (MH^+)

Step 3

Scheme 20:




A mixture of (3,4-difluoro-benzyl)-prop-2-ynyl-carbamic acid tert-butyl ester (3.2 g, 11.4 mmol), 4-chloroiodobenzene (4.1 g, 17.2 mmol, 1.5 eq.), $Pd(PPh_3)_2Cl_2$ (0.4 g. 0.57 mmol, 5 mol%), iPr_2NH (4 mL, 28.5 mmol, 2.5 eq.) and CuI (0.435 g, 2.3 mmol, 0.2 eq.) in 100 mL dichloromethane was stirred overnight at room temperature. The solution was diluted with 300 mL ether and filtered through a CELITE pad to remove the insoluble components. The filtrate was washed with 2x100 mL 1N HCl, 100 mL brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 6% ethyl acetate in hexanes to provide 2.6 g of [3-(4-chloro-phenyl)-prop-2-ynyl]-(3,4-difluoro-benzyl)-carbamic acid tert-butyl ester.

MS: 336.10 ($[M-^tBu]^+$)

Step 4

Scheme 21:




To a solution of [3-(4-chloro-phenyl)-prop-2-ynyl]-(3,4-difluoro-benzyl)-carbamic acid tert-butyl ester (1.05 g, 2.7 mmol) in 20 mL of 1:1 methanol-dichloromethane was added triethylamine (40 μL, 0.1 eq) and Lindlar catalyst (105 mg). The suspension was stirred under a H_2 balloon. After 1 hr, an additional 500 mg of Lindlar catalyst was added and the mixture was stirred for another 1.5 hr. The mixture was filtered through a CELITE pad and concentrated to provide the crude product.

The reaction was carried again with 1.5 g of [3-(4-chloro-phenyl)-prop-2-ynyl]-(3,4-difluoro-benzyl)-carbamic acid tert-butyl ester. The resultant crude product was combined with the crude product prepared in the first batch and chromatographed using 6% ethyl acetate-hexanes as solvent to provide 2.04 g of [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-carbamic acid tert-butyl ester.

MS: 338.11 ($[M-^tBu]^+$)

Step 5

Scheme 22:




A solution of [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-carbamic acid tert-butyl ester (2.04 g) in 20 mL of 4N HCl in dioxane was stirred at room temperature for 1 hr. The solution was poured into 150 mL of aq. K_2CO_3, and extracted with 3x50 mL of ethyl acetate. The combined organic layer was washed with 50 mL brine, dried over Na_2SO_4, filtered and concentrated to provide 1.49 g of [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amine.

MS: 294.12 (MH^+)

Step 6

Scheme 23:




To a solution of [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amine (1.49 g, 5.1 mmol), triethylamine (1.06 mL, 7.6 mmol, 1.5 eq.), DMAP (i.e., dimethylaminopyridine) (62 mg, 0.51 mmol, 0.1 eq.) in 15 mL dichloromethane at 0°C was added a solution of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl chloride (6.3 mmol) in 15 mL dichloromethane and stirred for 1 hr. After aqueous work-up, the crude product was purified by chromatography with 20% ethyl acetate-hexanes to provide 2.05 g of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amide.

MS: 518.11 (MH^+)

Step 7

Scheme 24:




A solution of 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amide (2.04 g) in 40 mL toluene was heated in a sealed tube at 150°C for 2 hr. The solution was concentrated and chromatographed with 15% to 30% ethyl acetate in hexanes to provide 720 mg of **49** and 1.1 g of **50**.

MS for **49**: 518.3 (MH^+)

MS for **50**: 518.3 (MH^+)

Preparation of Compound **51**

Scheme 25:




A solution of **49** (215 mg, 0.41 mmol) and DBU (63 mg, 0.42 mmol, 1 eq.) in 6 mL dichloromethane was stirred at room temperature for 2 hr,

concentrated and chromatographed with 40% ethyl acetate-hexanes to provide 160 mg of **51**.

MS: 519.07 (MH^+)

Preparation of Compound **52**


Scheme 26:




To a solution of **51** (105 mg) in 5 mL ethyl acetate was added 10 mg of PtO_2 and the suspension was stirred under a H_2 balloon for 80 min. The solution was then filtered through a CELITE pad, concentrated and chromatographed with 40% to 70% ethyl acetate-hexane to provide 67 mg of **52**.

MS: 521.03 (MH+)

Preparation of Compounds **53** and **54**



MS: 520.3 (MH^+)








MS: 518.3 (MH^+)

Using a procedure similar to the procedure described above in Scheme 26, compound **50** was reduced to compound **53**. During this reaction, compound **54** was also obtained.

Preparation of Compounds **55-59**


The following compounds (X=N) were also prepared using procedures similar to those described above in Scheme 17.

Compound	MS (MH^+)

Compound	MS (MH^+)
 55	519.1
 56	519.1
 57	519.3
 58	521.3
 59	521.3

Preparation of Compounds **60-62**


Scheme 27:




The *cis*-diaryl compounds were prepared as shown above. To a solution of 4-chlorocinnamaldehyde (1.45 g, 8.70 mmol) in 100 mL of dichloroethane was added 3,4-difluorobenzylamine followed by sodium triacetoxyborohydride (3.7 g, 17.5 mmol, 2 eq.). The mixture was stirred overnight at room temperature, washed with aq. $NaHCO_3$, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 40% ethyl acetate - hexanes to provide 0.84 g of [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amine. The [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amine was coupled with 5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl chloride to provide 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amide. When 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-(3,4-difluoro-benzyl)-amide was subjected to the Diels-Alder reaction conditions shown above, it gave **60** as the major product. Epimerization of the *trans*-lactam gave **61** and reduction of the double bond of **61** gave **62**.

Compound	MS (MH^+)
60	518.18
61	518.06
62	520.05


Preparation of Compounds **63-65**



63




64



65

Substituted N-benzyl-4,5-diphenyl-isoindoles **63-65** were prepared by reducing the corresponding lactams. As a representative example, the preparation of **64** is presented below.

Scheme 28



52 **64**

To a solution of **52** (160 mg, 0.31 mmol) in 3 mL THF at was added 0.92 mL of 1M solution of $LiAlH_4$ in THF and the mixture was heated at reflux for 1 hr. The solution was cooled to room temperature and successively added 30 μL water, 30 μL 15% aq. NaOH solution and 90 μL water. The precipitate was filtered off and the filtrate was concentrated and chromatographed with 30% ethyl acetate-hexane to provide 108 mg of **64**. MS: 506.3 (MH^+)

Compounds **63** and **65** were prepared using similar procedures from the appropriate starting lactam.

63 [MS: 507.3 (MH^+)]

65 [MS: 506.3 (MH^+)].


It will be recognized by one of ordinary skill in the art that the substituted aryl moieties of the above isoindoles can be independently replaced with heteroaryl moieties by the use of the appropriate heteroaryl substituted starting materials.

It will also be recognized by one of ordinary skill in the art that isoindoles having different substitution patterns of aryl and/or heteroaryl groups on the six-membered ring may be provided by use of the appropriate starting materials. In addition, substitution on the isoindole nitrogen atom may be provided by methods known in the art (e.g., alkylation, acylation, arylation, etc.).

Preparation of Compounds **66-71**


The Diels-Alder reaction (e.g., Scheme 45, Step 3) can also be carried out using a Boc protected intermediate such as [5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoyl]-[3-(4-chloro-phenyl)-allyl]-carbamic acid tert-butyl ester as described below in Scheme 29 to give the cyclization products **66** and **67**. The *trans*-lactam **66** can be epimerized to the *cis*-lactam **68**, the double bond reduced and the Boc group cleaved to give **69**. Alternatively, **69** can be prepared by first cleaving the Boc group followed by lactam epimerization and double bond reduction.

Scheme 29:



Step 1

Scheme 30:




To a mixture of 4-bromo-3-chloro anisole (35 g, 0.159 mol), KCl (11.9 g, 0.160 mol, 1 eq.), K_2CO_3 (33 g, 0.238 mol, 1.5 eq) in 400 mL DMF in a sealed tube was added acrolein diethylacetal (73 mL, 0.479 mol, 3 eq.) and Bu_4NOAc (96 g, 0.318 mol, 2 eq.). N_2 was bubbled through the mixture and

$Pd(OAc)_2$ (1.1 g, 4. mmol, 3 mol%) was added. The reaction mixture was heated in an oil bath for 6 hr at 100°C then cooled in an ice-bath. 300 mL of water was then added, followed by 500 mL of 1N HCl. The ice-bath was removed and the mixture was stirred for 30 min. The solution was extracted with ethyl acetate once then with diethyl ether 3 times. The combined organic layer was washed with water, brine, dried over $MgSO_4$, filtered and concentrated to provide the crude product which was recrystallized from hot ethyl acetate-hexanes to provide 15.5 g of crystalline 3-(2-chloro-4-methoxy-phenyl)-propenal. The mother liquor was concentrated and chromatographed with 10% ethyl acetate to provide another 5.6 g of 3-(2-chloro-4-methoxy-phenyl)-propenal.

Step 2

Scheme 31:



To a suspension of NaH (6.3 g, 0.157 mol, 60% dispersion in mineral oil) in 400 mL THF at room temperature was added triethyl phosphonoacetate (29 mL, 0.146 mol) and stirred for 30 min. The solution was cooled in an ice-bath and a solution of 3-(2-chloro-4-methoxy-phenyl)-propenal (22 g, 0.112 mol) in 200 mL THF was added. The mixture was stirred for 1 hr and quenched with the addition of aq. NH_4Cl. The THF was evaporated and the slurry was extracted with diethyl ether. The combined organic layer was washed with water, brine, dried over $MgSO_4$, filtered and concentrated to provide the crude 5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoic acid ethyl ester.


The crude ester product was taken up in 200 mL each of methanol and THF and then cooled to 0°C. To this mixture was added a solution of KOH (19 g, 0.338 mol, 3 eq.) in 200 mL H_2O. The mixture was stirred at room temperature for 1.5 hr and the organic solvent was evaporated. The aqueous layer was diluted with 300 mL H_2O and washed with ether to remove the mineral oil. It was then cooled in an ice-bath and acidified with concentrated

HCl to ~pH 2. The thick precipitate was filtered and rinsed with water, then dried in a vacuum oven at 75°C to provide 26.2 g of 5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoic acid.

MS: 239.24 (MH^+)

Step 3

Scheme 32:




To a suspension of 5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoic acid (6 g, 25.1 mmol) in 100 mL dichloromethane at room temperature was added oxalyl chloride (4.4 mL, 50.4 mmol, 2 eq.) followed by DMF (39 μL, 0.51 mmol, 2 mol %). The mixture was stirred for 1 hr and the resultant clear solution was concentrated and evaporated with toluene to provide an acid chloride.

The acid chloride was dissolved in 150 mL dichloromethane, cooled to 0°C, and then triethylamine (5.3 mL, 38.0 mmol, 1.5 eq.), DMAP (310 mg, 2.54 mmol, 0.1 eq.) followed by a solution of 3-(4-chloro-phenyl)-allylamine (5.1 g, 30.2 mmol, 1.2 eq.) in dichloromethane were added. The mixture was stirred for 1 hr at room temperature, diluted with 700 mL ethyl acetate and washed with 1N HCl, aq. $NaHCO_3$ and brine. It was dried over $MgSO_4$, filtered and concentrated to provide 10.1 g of 5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide.

MS: 388.19 (MH^+)

Step 4


Scheme 33:



To a suspension of 5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide (9.8 g, 25 mmol) and $(Boc)_2O$ (11 g, 50.4 mmol, 2 eq.) in 150 mL dichloromethane was added triethylamine (3.5 mL, 25.1 mmol, 1 eq.) followed by DMAP (3.1 g, 25.4 mmol, 1 eq.). The mixture was stirred for 1 hr at room temperature, diluted with ether, washed with 1N HCl, aq. $NaHCO_3$, brine, dried over $MgSO_4$, filtered and concentrated to provide 9.8 g of [5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoyl]-[3-(4-chloro-phenyl)-allyl]-carbamic acid tert-butyl ester.

Step 5

Scheme 34:




A solution of 5-(2-chloro-4-methoxy-phenyl)-penta-2,4-dienoyl]-[3-(4-chloro-phenyl)-allyl]-carbamic acid tert-butyl ester (9.5 g) in 200 mL toluene was heated in a sealed tube at 150 °C for 3 hr. The solution was concentrated and chromatographed with 15% to 30% ethyl acetate-hexanes to provide 4.8 g of **66** and 2.8 g of **67**.

MS for **66**: 488.3 (MH^+)

MS for **67**: 488.3 (MH^+)

Step 6

Scheme 35:




A solution of **66** (690 mg, 1.41 mmol) in 15 mL of dichloromethane was stirred at room temperature with DBU (215 mg, 1.41 mmol, 1 eq.) for 1 hr, concentrated and chromatographed with 20% to 30% ethyl acetate-hexane to provide a mixture of products. This mixture was taken up in 15 mL of 1:1 methanol - dichloromethane and stirred under a H_2 balloon with 65 mg of PtO_2

for 30 min., then filtered though a CELITE pad, concentrated and chromatographed with 30% ethyl acetate-hexane to provide 34 mg of **68**.

MS: 434.2 ([M-^{t}Bu]$^+$)

Step 7

Scheme 36:




To a flask containing 32 mg of **68** was added 1 mL of TFA (i.e., trifluoroacetic acid). The mixture was stirred for 1 hr at room temperature, then concentrated and chromatographed with 3% methanol - dichloromethane to provide 25 mg of **69**.

MS: 390.2 (MH$^+$)

Step 8

Scheme 37:




To a solution of **66** (2.3 g, 4.71 mmol) in 20 mL dichloromethane at 0°C was added 20 mL of TFA. The mixture was stirred for 50 min, then concentrated to provide the deprotected product.

The deprotected product was stirred overnight with 720 mg of DBU in 20 mL each of dichloromethane and acetonitrile. After overnight stirring, another 25 mL of acetonitrile and 1.4 g of DBU was added and the mixture was stirred for 2.5 hr. The mixture was diluted with ethyl acetate, washed with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 50% to 60% ethyl acetate-hexane then with 5% methanol - dichloromethane to provide 1.88 g of **71**.

MS: 388.2 (MH$^+$)

Alternate Preparation of **69**

Scheme 38:




To a solution of **71** (1.8 g) in 50 mL of 1:9 methanol - dichloromethane was added 90 mg of PtO_2 and the suspension was stirred under a H_2 balloon for 40 min. The mixture was then filtered though a CELITE pad, concentrated and the crude product recrystallized from a hot dichloromethane-hexane mixture to provide 1.1 g of crystalline **69**.

MS: 390.2 (MH^+)

Preparation of Compounds **72-78**

Scheme 39:




The methoxy functionality at the 5-phenyl group can be converted into a variety of other groups. For example, as shown above, the methoxy group of **69** was hydrolyzed to a hydroxyl **72**, which was then converted to the triflate **73**. The triflate **73** was converted to the CN derivative **74** using $Zn(CN)_2$ and

Pd catalyst or it was reduced to **75** using formic acid and a Pd catalyst. The triflate **73** was reacted with Zn reagents to provide compounds **76** and **77** or with boronic acid to give **78.**

Preparation of Compound **72**

Scheme 40:




To a solution of **69** (600 mg, 1.54 mmol) in 15 mL dichloromethane was added BBr_3 (7.7 mL of a 1M solution in dichloromethane). The mixture was stirred for 1 hr at 0°C then stirred with 100 mL water for 20 min. The dichloromethane layer was separated and the aqueous phase was extracted with ethyl acetate. The combined organic layer was washed with water, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 3% to 10% methanol - dichloromethane to provide 535 mg of **72.**

MS: 376.2 (MH^+)

Preparation of Compound **73**


Scheme 41:



To a suspension of **72** (400 mg, 1.1 mmol) in 10 mL acetonitrile was added triethylamine (300 µL, 2.1 mmol, 2 eq.) followed by N-phenyltrifluoromethane sulfonimide (870 mg, 1.5 eq.). The mixture was stirred for 2.5 hr at room temperature and the clear solution was diluted with ethyl acetate, washed twice with aq. $NaHCO_3$, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 3% methanol - dichloromethane to provide 490 mg of **73.**

Preparation of Compound **74**

Scheme 42:




A mixture of **73** (62 mg, 0.12 mmol), Zn $(CN)_2$ (9 mg, 0.08 mmol, 0.6 eq.) and $Pd(PPh_3)_4$ (14 mg, 0.012 mmol, 0.1 eq.) in 1 mL DMF in a sealed tube was heated at 100°C for 1 hr. The mixture was diluted with water, washed twice with water, brine, dried over $MgSO_4$, filtered, concentrated and the crude product was purified by preparative TLC using 80% ethyl acetate - hexane to provide 50 mg of **74.**

MS: 385.21 (MH^+)

Preparation of Compound **75**

Scheme 43:




A mixture of **73** (50 mg, 0.10 mmol), HCO_2H (19 μL, 0.50 mmol, 5 eq.), triethylamine (70 μL, 0.5 mmol, 5 eq.) and $Pd(PPh_3)_4$ (6 mg, 5.2 μmol, 5 mol%.) in 1 mL DMF in a sealed tube was heated at 100°C for 4 hr. The mixture was diluted with ethyl acetate, washed twice with aq. $NaHCO_3$, brine, dried over $MgSO_4$, filtered, concentrated and the the product was purified by preparative TLC using 3% methanol - dichloromethane to provide 29 mg of **75**.

MS: 360.2 (MH^+)

Preparation of Compound **76**

Scheme 44:




A mixture of **73** (50 mg, 0.10 mmol), MeZnCl (0.5 mL of 2M solution, 10 eq.), and $Pd(PPh_3)_4$ (6 mg, 5.2 μmol, 5 mol%.) in 1 mL THF in a sealed tube was heated at 80°C for 4 hr. The mixture was quenched with methanol, diluted with ethyl acetate and washed twice with 1N HCl, brine, dried over $MgSO_4$, filtered and concentrated and the crude product was purified by preparative TLC using 60% ethyl acetate-hexane to provide 26 mg of **76**.

MS: 385.21 (MH^+)

Preparation of Compound **77**

Scheme 45:




A mixture of **73** (30 mg, 0.059 mmol), 2-pyridyl zinc bromide (0.59 mL of 0.5M solution, 5 eq.), and $Pd(PPh_3)_4$ (6.8 mg, 5.8 μmol, 10 mol%.) in 0.5 mL THF in a sealed tube was heated at 100°C for 4 hr. The mixture was diluted with ethyl acetate washed twice with aq. NH_4Cl, brine, dried over $MgSO_4$, filtered, concentrated and the crude product was purified by preparative TLC using 3% methanol - dichloromethane to provide 8 mg of **77**.

MS: 437.2 (MH+)

Preparation of Compound **78**

Scheme 46:




A mixture of **73** (30 mg, 0.059 mmol), phenyl boronic acid (15 mg, 0.13 mmol, 2 eq.), K_2CO_3 (33 mg, 0.24 mmol, 4 eq.) and $Pd(PPh_3)_4$ (6.8 mg, 5.8 μmol, 10 mol%.) in 1 mL of 4:2:1 toluene-EtOH-H_2O in a sealed tube was heated at 100°C for 4 hr. The mixture was diluted with ethyl acetate washed twice with aq. K_2CO_3 brine, dried over $MgSO_4$, filtered, concentrated and the crude product was purified by preparative TLC using 60% ethyl acetate - hexane to provide 25 mg of **78**.

MS: 436.2 (MH^+)

Preparation of Compounds **79-86**


Scheme 47:



The 7a-position of the isoindolone can be functionalized as shown above in Scheme 47 by reacting the enolate with the appropriate electrophiles. For example, a hydroxy functionality was introduced in **66** by generating the potassium enolate followed by reaction with oxaziridine. Double-bond reduction followed by deprotection of the Boc group gave **80**. Alternatively, the double bond was reduced before generating the enolate (e.g., reduction of **66** to form **81**). Compound **81** was subsequently reacted with MeI and TFA to give **82**. Alkylation, e.g., ethylation with KHMDS/18-crown-6 and EtI, gave **83**. In a similar fashion, **81** was converted to **84**, **85** and **86** by reacting the enolate with the appropriate optionally substituted benzyl bromide or allyl iodide.

Preparation of Compound **79**

Scheme 48:




To a solution of **66** (44 mg, 0.090 mmol) in 1 mL THF at -78°C was added a 0.5 M solution of KHMDS (i.e., potassium bis(trimethylsilyl)amide) in toluene (0.27 mL, 1.5 eq.). The solution was stirred for 10 min at -78°C, 10 min at 0°C, and then cooled back to -78°C. To the cooled solution was then added a solution of (1S)-(+)-(camphorsulfonyl)oxaziridine (Davis reagent) (31 mg, 0.135 mmol, 1.5 eq.) in 0.5 mL THF and stirred for 2 hr. The solution was then quenched by the addition of aq. NH_4Cl. Extraction with ethyl acetate followed by chromatographic purification using 1% methanol - dichloromethane gave 30 mg of **79**.

MS: 504.3 (MH^+)

Preparation of Compound **80**

Scheme 49:




A suspension of **79** (30 mg) and PtO_2 (3 mg) in 2 mL ethyl acetate was stirred under H_2 balloon for 2 hr. The mixture was filtered through a CELITE pad, concentrated and stirred with 1 mL each of TFA (i.e., trifluoroacetic acid) and dichloromethane at 0°C for 1 hr. The solution was concentrated and purified by preparative TLC using 3% methanol - dichloromethane to provide 24 mg of **80**.

MS: 406.2 (MH^+)

Preparation of Compound **81**

Scheme 50:




A suspension of **66** (230 mg) and PtO_2 (23 mg) in 5 mL of ethyl acetate was stirred under H_2 balloon for 30 min, filtered and evaporated to provide 230 mg of **81**.

MS: 490.3 (MH^+)

Preparation of Compound **82**

Scheme 51:




To a solution of **81** (30 mg, 0.061 mmol) in 1 mL of THF at -78°C was added a 0.5 M solution of KHMDS (0.15 mL, 0.075 mmol, 1.2 eq.) in toluene and the mixture was stirred for 20 min. To the mixture was added MeI (19 μL, 0.305 mmol, 5 eq.), and it was then stirred for 10 min. and quenched with aqueous NH_4Cl. After extracting the mixture with ethyl acetate, the crude product was purified by preparative TLC using 30% ethyl acetate - hexane to provide 11 mg of the methylated product. The methylated product was stirred with 0.5 mL each of TFA and dichloromethane at 0°C for 1 hr, concentrated and evaporated with toluene to provide 8 mg of **82**.

MS: 404.2 (MH^+)

Preparation of Compound **83**

Scheme 52:



To 50 mg of **81** in 1.5 mL of dry THF was added 18-crown-6 (1.5 eq.). The mixture was degassed and, at -78°C under argon, potassium bis(trimethylsilyl)amide (1.5 eq., 306 μL of a 0.5 M solution in toluene) was added. After stirring for twenty minutes, iodoethane (1.5 eq.) was added. The mixture was quenched with aqueous ammonium chloride and extracted with ethyl acetate three times. The combined organic extracts were washed with brine, dried with magnesium sulfate, filtered and evaporated to dryness. Purification by flash chromatography (0-25% ethyl acetate in hexane) yielded 30 mg of the ethylated product.

To a solution of the ethylated product prepared as described above in 1.5 mL of dichloromethane at 0°C was added 1 mL of trifluoroacetic acid and the mixture was stirred under nitrogen for one hour. The mixture was evaporated to dryness and the residue was dissolved in toluene, evaporated to dryness then dissolved in diethyl ether and again evaporated to dryness yielding 23 mg of **83**.


MS: 418.2 (MH^+)

Intermediate **81** was converted to **84**, **85** and **86** using a procedure similar to the procedure used to prepare **82** except that 4-cyanobenzyl bromide, 3,4-difluorobenzyl bromide, and allyl bromide, respectively, were used instead of methyl iodide.

Compound	MS (MH^+)
84	505.3
85	516.3
86	430.2

Preparation of Compounds **87-91**

Scheme 53:




Using procedures similar to the preparation of compounds **72** and **73**, **82** was converted to **87** and triflate **88**. Also using procedures similar to the transformation of the triflate functionality of **73** to compounds **74**, **75**, and **76**, compound **88** was converted to compounds **89**, **90** and **91**.

Compound	MS (MH^+)
87	390.2
88	522.3
89	399.2
90	374.2
91	388.2

Compound **89** was also resolved using chiral HPLC conditions

Chiral Resolution of **89**




A solution of **89** (~550 mg) in 4 ml of 1:1 iso-propyl alcohol and hexanes was injected into a Chiralpac AD preparative HPLC column (5 cm x 50 cm) and eluted with 15% isopropyl alcohol in hexanes to obtain 149 mg of **89A** and 230 mg of **89B**.

MS for **89A**: 399.2 (MH^+)

MS for **89B**: 399.2 (MH^+)

Preparation of Compounds **92-95**

Scheme 54




Using procedures similar to those used to prepare compounds **66**, **82**, **89**, and **91** and using appropriate starting materials as shown in Scheme 54, compounds **92**, **93**, **94**, and **95** were prepared.

Compound	MS (MH^+)

Compound	MS (MH^+)
93	404.2
94	399.2
95	388.2

Preparation of Compounds **96-107**

Scheme 55:




Step 1

Compound **90** was converted to compound **96** using procedures similar to those shown in Scheme 25.

Step 2


Scheme 56:



To a solution of compound **96** (30 mg, 0.064 mmol), LiCl (35 mg, 0.826 mmol, 6.5 eq.), and 2,4,6-triisopropylbenzenesulfonyl azide (45 mg, 0.15 mmol, 2.3 eq.) in 1 mL DMF under argon atmosphere at room temperature was added DBU (19 μL, 0.13 mmol, 2 eq.). The solution was stirred for 0.5 hr at room temperature and diluted with ethyl acetate. The solution was washed with NH_4Cl (sat.), dried over $MgSO_4$, filtered, and concentrated. The residue was subjected to preparative TLC purification (SiO_2, 3:2 hexane/EtOAc) to afford **97** as a white solid (25 mg, 74%). LCMS: *m/e* 529.3 (MH^+)

Step 3

Scheme 57:




To a solution of compound **97** (171 mg, 0.323 mmol) in 7 mL EtOAc-H_2O (10-1) under argon atmosphere at room temperature was added PMe_3 (650 μL, 1 M in THF, 0.65 mmol, 2 eq.). The solution was stirred for 4 hr at room temperature. The solution was concentrated. The residue was chromatographed with 3% methanol - dichloromethane to afford **98** as an off-white solid (145 mg, 89%).

Step 4

Compound **98** was converted to compound **99** using procedures similar to those shown in Scheme 50.

Step 5


Scheme 58:



A mixture of compound **99** (50 mg, 0.10 mmol), 18-crown-6 (52 mg, 2 eq.), K_2CO_3 (82 mg, 6 eq.), and 1,5-diiodopentane (29 μL, 2 eq.) in 2 mL


MeCN was heated for 19 hr at 90°C. The mixture was filtered. The filtrate was dissolved in EtOAc, washed with NH_4Cl (sat.), dried over $MgSO_4$, filtered, and concentrated. The residue was subjected to preparative TLC purification (SiO_2, 3:2 hexane/EtOAc) to afford 97f (10 mg, 18%). Also isolated was compound **103** (18 mg, 24%).

Step 6



101

LCMS: m/e 473.3 (MH^+)




105

LCMS: m/e 645.4 (MH^+)

Compounds **100** and **103** were converted to compounds **101** and **105** using procedures similar to those shown in Scheme 36.

Scheme 59:




99 **102** **107**

LCMS: m/e 475.3 (MH^+) LCMS: m/e 599.3 (MH^+)


Compounds **102** and **107** were similarly prepared as shown above in Scheme 60.

The amine functionally at the 7a-position can also be introduced in a slightly modified sequence as presented in the following scheme. Intermediate **108**, which was prepared using a procedure similar to the preparation of **66**, was treated with KHMDS followed by trisyl azide to give the azido derivative **109**. Reduction of the azide followed by cleavage of the BOC group afforded the amine **111** which was converted to the amide **112**. The amine can be similarly converted to various analogs such as sulfonamides, carbamates, ureas etc. using standard literature procedures.



Preparation of Compound **109**

Scheme 60:




To 1.0g of **108** in 10mL of dry THF at 0°C was added 1.3 eq. of potassium hexamethyldisilyl amide (0.5M solution in toluene) and the mixture stirred for 20 minutes then cooled to -78°C. A cooled (-78°C) solution of Tris-N_3 (1.5 eq.) in 2.5ml of THF was added followed, after two minutes, by acetic acid (3eq.). The reaction was immediately warmed to room temperature using a warm water bath then stirred for 1.5 hours. The mixture was evaporated to dryness, dissolved in 10mL of dichloromethane and washed with water, aq. $NaHCO_3$, and brine, then dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography with 0% to 15% ethyl acetate – hexanes yielded 864 mg of **109**.

MS : 479.3 (MH^+).

Preparation of Compound **110**

Scheme 61:




To 860 mg of **109** in 22 mL of ethylacetate/water (10:1) at 0°C, after degassing under argon, was added 2eq. of trimethyl phosphine (1.0M solution in THF) and the mixture warmed slowly to room temperature. After 16 hours the mixture was evaporated to dryness, then toluene was added to the residue and the mixture evaporated to dryness. Purification by flash chromatography with 0% to 2.5% methanol - dichloromethane yielded 575 mg of **110**.

MS: 392.2 (MH^+)

Preparation of Compound **111**

Scheme 62:




To 565 mg of **110** in 5 mL of dichloromethane at 0°C was added 2 mL of trifluoroacetic acid and the mixture was stirred under a N_2 atmosphere. After two hours the reaction mixture was poured onto aq. $NaHCO_3$ and extracted with dichloromethane three times. The combined extracts were dried with $MgSO_4$, filtered and evaporated to dryness yielding 465 mg of **111**.

MS: 409.2 (MH^+)

Preparation of Compound **112**

Scheme 63:




To 20 mg of **111** in 1.5 mL pyridine at 0°C was added 3 eq. of acetyl chloride and the mixture was warmed to room temperature while stirring under a N_2 atmosphere. The reaction mixture was poured onto aq. ammonium chloride and extracted with ethylacetate three times. The combined extracts were washed with water twice and brine once, dried with $MgSO_4$, filtered and evaporated to dryness yielding 17mg of **112** after purification by flash chromatography with 0% to 2.5% methanol - dichloromethane.

MS: 451.2 (MH^+)

Preparation of Compounds **113-117**


Scheme 64:



The C_3 position of the lactam can be substituted with various groups by starting with an appropriately functionalized propargyl amine. For example, a methyl group can be introduced at the C_3 position as described above in Scheme 64.

Step 1


Scheme 65:



To a solution of 4-chloro iodobenzene (10 g, 42 mmol), 3-butyn-2-ol (6.6 mL, 84 mmol, 2 eq.), and iPr_2NH (14.8 mL, 105 mmol, 2.5 eq.) in 200 mL dichloromethane was added CuI (1.6 g, 8.4 mmol, 0.2 eq.) followed by $Pd(PPh_3)_2Cl_2$ (1.5 g, 2.1 mmol, 5 mol %). The mixture was stirred for 1.5 hr at room temperature, concentrated and diluted with ether. The insoluble components of the mixture were filtered off and the filtrate was washed with 1N HCl and brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 20% ethyl acetate - hexane to provide 7.6 g of 4-(4-chloro-phenyl)-but-3-yn-2-ol.

Step 2

Scheme 66:




To a solution of 4-(4-chloro-phenyl)-but-3-yn-2-ol (3 g, 16.6 mmol), phthalimide (3.7 g, 25.2 mmol, 1.5 eq.) and PPh_3 (4.8 g, 18.3 mmol, 1.1 eq.) in 100 mL of THF cooled to 0°C with an ice bath was added dropwise DEAD (i.e., diethyl azodicarboxylate) (3.1 mL, 19.9 mmol, 1.2 eq.). The ice-bath was removed and the mixture was stirred overnight at room temperature. The THF was evaporated and the residue was stirred with 200 mL of 1:1 ether-hexane mixture for 1 hr. The precipitate was filtered off and the filtrate was concentrated and chromatographed with 15% ethyl acetate-hexane to provide 4.1 g of 2-[3-(4-chloro-phenyl)-1-methyl-prop-2-ynyl]-isoindole-1,3-dione.

MS: 310.09 (MH^+)

Step 3

Scheme 67:




To a solution of 2-[3-(4-chloro-phenyl)-1-methyl-prop-2-ynyl]-isoindole-1,3-dione (4.1 g, 13.2 mmol) in 20 mL of dichloromethane was added 30 mL of methanol followed by hydrazine (4.2 mL, 133 mmol, 10 eq.). The mixture was stirred for 1 hr, filtered, concentrated and chromatographed with 3% methanol - dichloromethane to provide 1.5 g of 3-(4-chloro-phenyl)-1-methyl-prop-2-ynylamine.

MS: 163.21 (MH^+)

Step 4

Scheme 68:



A mixture of 3-(4-chloro-phenyl)-1-methyl-prop-2-ynylamine (1.5 g, 8.4 mmol), triethylamine (0.12 mL, 0.86 mmol, 0.1 eq.) and Lindlar catalyst (150 mg) in 40 mL of 1:1 methanol - dichloromethane was stirred for 1 hr, filtered through a CELITE pad and concentrated to provide 1.5 g of 3-(4-chloro-phenyl)-1-methyl-allylamine.


Compound 3-(4-chloro-phenyl)-1-methyl-allylamine was converted to 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-1-methyl-allyl]-amide using procedures similar to those described in Scheme 5, above. Compound 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-1-methyl-allyl]-amide was converted to compounds **113** and **114** using the Diels-Alder conditions similar to those described above in Scheme 6. Compound **115** was prepared by the isomerization of compound **113** using procedures similar to those described in Scheme 6. Compound **116** was prepared by the reduction of compound **115** using procedures similar to those

described in Scheme 7. Compound **117** was prepared by the reduction of compound **114** using procedures similar to those described in Scheme 7.

Compound	MS (MH^+)
113	406.2
114	406.2
115	406.2
116	408.1
117	408.2

Preparation of Compounds **118-121**


Scheme 69



Intermediate **113** from the above synthesis was further transformed into 7a-methyl analog **121** as described in the following scheme. This was also further resolved into the single enantiomers **121a** and **121b**.

Preparation of Compounds **118** and **119**

Scheme 70:




To a solution of **113** (500 mg, 1.23 mmol) $(Boc)_2O$ (2.8 g, 12.8 mmol, 10 eq.), Et_3N (345 μL, 2.48 mmol, 2 eq.) in 5 mL each of dichloromethane and acetonitrile was added DMAP (300 mg, 2.46 mmol, 2 eq.) and the mixture was stirred overnight at room temperature. The solution was diluted with ether, washed with 1N HCl, aq. $NaHCO_3$, and brine, dried over $MgSO_4$, filtered, evaporated and chromatographed with 0% to 20% ethyl acetate – hexanes to provide 220 mg of **118** and 330 mg of **119**.

MS for **118**: 450.2 (M^+-^{t}Bu)

MS for **119**: 450.2 (M^+-^{t}Bu)

Compound **118** was converted to compound **120** using a procedure similar to the conversion of **66** to **81**. Conversion of **120** to **121** was achieved using a procedure similar to the conversion of **81** to **82**. Also, **121** was resolved using chiral HPLC conditions as described below to obtain the single enantiomers **121a** and **121b**.

Scheme 71:



About 40 mg of **121** was dissolved in isopropanol and injected onto a 5x50cm CHIRALPAK AD column and eluted with 10% isopropanol in hexane at 100 mL/min, using a 220 nm UV detector. 15 mg each of **121a** (enantiomer 1) and **121b** (enantiomer 2) was obtained. Retention times were 19.8 minutes and 25.4 minutes, respectively


MS for **120**: 510.3 (MH^+)

MS for **121**: 422.2 (MH^+)

MS for **121a**: 422.2 (MH^+)


MS for **121b**: 422.2 (MH^+)

Scheme 72:



In the above preparation of **121**, racemic starting materials were employed and the individual enantiomers were resolved using chiral HPLC conditions. Alternatively, these chiral C_3-substiuted analogs were synthesized as a single enantiomer by starting with 3-butyn-2(S)-ol as shown in Schemes 64 and 69.


Scheme 72a



For example, 3-(4-chloro-phenyl)-1-methyl-allylamine (prepared, e.g., by the method of Scheme 7) was reacted with 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid using the method of, e.g., Scheme 29, to provide [3-(4-chloro-phenyl)-1-methyl-propyl]-[5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl]-carbamic acid tert-butyl ester. [3-(4-Chloro-phenyl)-1-methyl-propyl]-[5-(2,4-dichloro-phenyl)-penta-2,4-dienoyl]-carbamic acid tert-butyl ester was then cyclized (e.g., using the method of Scheme 47) to provide 4-(4-Chloro-phenyl)-5-(2,4-dichloro-phenyl)-3,7a-dimethyl-octahydro-isoindol-1-one.

Preparation of Compounds **122-131**

Scheme 73:




Another example of chiral synthesis of the compounds of the present invention is presented above in Scheme 73 for the preparation of **130** and **131** in an optically active form starting with optically active 3-butyn-2(S)-ol.

Preparation of Compound **122**

Amide **122** was prepared using a procedure similar to that used to prepare the amide from which **113** was prepared, except that optically active 3-butyn-2(S)-ol was used instead of racemic 3-butyn-2-ol.

Preparation of Compound **123**

Scheme 74:




To a suspension of **122** (10 g, 24.8 mmol), $(Boc)_2O$ (10.9 g, 2 eq.) in 100 mL dichloromethane at room temperature was added Et_3N (3.5 mL, 1 eq) followed by DMAP (3.1 g, 1 eq) and the suspension was stirred at room temperature. After two hours another 2 equivalents of $(Boc)_2O$ was added and the mixture was stirred overnight. The solution was concentrated to half of its volume, diluted with ethyl acetate, washed with 1N HCl, aq. $NaHCO_3$ and brine, dried over $MgSO_4$, filtered and concentrated to give a crude product. The crude product was suspended in MeOH-CH_2Cl_2 mixture and diluted with diethyl ether. The solid was filtered off and rinsed with diethyl ether. The filtrate was concentrated and chromatographed with 0% to 10% ethyl acetate – hexanes to provide 9.5 g of **123**.

MS: 502.3 (MH^+)

Preparation of Compounds **124** and **125**

Scheme 74




A solution of **123** (9.5 g) in 100 mL of toluene was heated in a sealed tube at 130°C for 2 hr. The solution was concentrated and chromatographed with 0% to 20% ethyl acetate – hexanes to provide 4.3 g of **124** and 3.1 g of **125**.

MS for **124**: 502.3 (MH^+)

MS for **125**: 502.3 (MH^+)

Preparation of Compound **126**

Scheme 75




A suspension of **124** (4.3 g) and platinum oxide hydrate (215 mg, 5 wt%) in 100 mL of ethyl acetate was stirred under a hydrogen balloon for 30 min, filtered through a CELITE pad and concentrated to provide 4.15 g of **126**.

MS: 504.3 (MH^+)

Preparation of Compound **127**

Scheme 76:




To a solution of **126** (600 mg, 1.19 mmol) in 7 mL of THF at -78°C, degassed under vacuum, was added a THF solution of 0.5M 1M LHMDS (i.e., lithium hexamethyldisilazide) (3.6 mL, 1.8 mmol, 1.5 eq.) and stirred for 20 min. To this solution was added methyl iodide (0.37 mL, 5.94 mmol, 1.5 eq.), stirred for 10 min, quenched with aq. NH_4Cl and THF was evaporated. The mixture was extracted with ethyl acetate, the combined organic layer washed with brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 15% ethyl acetate in hexanes to provide 470 mg of **127**.

MS: 518.3 (MH^+)

Preparation of Compound **128**

Scheme 77:




To a solution of **127** (460 mg, 0.89 mmol) in 5 mL dichloromethane at room temperature was added a dichloromethane solution of 1M BBr_3 (4.5 mmol, 5 eq.) and the mixture was stirred at room temperature for 2 hr. The solution was diluted with water, stirred for few minutes, the organic layer separated and the aqueous layer extracted twice with dichloromethane. The combined organic layer was washed with brine, dried over $MgSO_4$, filtered and concentrated to provide 380 mg **128**.

MS: 404.2 (MH^+)

Preparation of Compound **129**

Scheme 78:




To a solution of **128** (330 mg, 0.82 mmol) in 5 mL each of dichloromethane and acetonitrile was added triethyl amine (0.23 mL, 1.65 mmol, 2 eq) followed by N-phenyltrifluoromethane sulfonimide (440 mg, 1.23 mmol, 1.5 eq.). The mixture was stirred for 1.5 hr, diluted with ethyl acetate washed twice with aq. $NaHCO_3$, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 3% methanol-dichloromethane to provide 370 mg of **129**.

MS: 536.3 (MH^+)

Preparation of Compound **130**

Scheme 79:




A mixture of **129** (150 mg, 0.28 mmol), $Pd_2(dba)_3$ (13 mg, 0.014 mmol, 5 mol%), dppf (i.e., diphenylphosphine-ferrocene complex) (19.5 mg, 0.035 mmol, 12.5 mol%), $Zn(OAc)_2$ (14 mg, 0.088 mmol, 0.3 eq.), Zn dust (5.5 mg, 0.084 mmol, 0.3 eq.), $Zn(CN)_2$ (23 mg, 0.20 mmol, 07 eq) in 2 mL of DMF in a sealed tube was bubbled with argon and heated in an oil-bath at 100°C for 1 hr. The solution was diluted with ethyl acetate, washed with ferrous ammonium sulfate, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 3% methanol-dichloromethane to provide 120 mg **130**.

MS: 413.2 (MH^+)

Preparation of Compound **131**


Scheme 80:



A mixture of **128** (40 mg, 0.10 mmol), K_2CO_3 (69 mg, 0.50 mmol, 5 eq.) and iodomethane (0.04 mL, 050 mmol, 5 eq.) in 2 mL of acetone was heated overnight in a sealed tube at 50°C. The mixture was diluted with water, extracted three times with ethyl acetate, combined organic layer washed with brine, dried over MgSO4, filtered and evaporated to provide 47 mg of **131**.

MS: 432.2 (MH^+)


General Scheme C



Preparation of Compounds **132** and **133**

Step 1


Preparation of 3-(2,4-Dichlorophenyl)-2-methylpropenal



2,4-dichlorobenzaldehyde (26.26 g, 0.15 mol) was dissolved in toluene (200 mL), and then KOH (8.4 g, 1 eq) in water (183 mL) was added followed by $BnEt_3NCl$. The mixture was cooled to 0°C and n-propanal (26.15g, 0.45 mol, 3 eq) in toluene (50 mL) was added dropwise. The mixture was allowed to warm to room temperature and stirred overnight. The organic layer was separated and washed successively with water, brine, dried ($MgSO_4$), and concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (0-1-2-3% EtOAc in hexanes) to give 17.56 g of 3-(2,4-dichlorophenyl)-2-methylpropenal.

Step 2

Preparation of 5-(2,4-Dichlorophenyl)-4-methylpenta-2,4-dienoic acid ethyl ester




Sodium hydride (4.88 g of a 60% dispersion in mineral oil, 1.5 eq, 0.122 mol) was suspended in THF (442 mL). To this suspension was added

a solution of diethoxyphosphorylacetic acid ethyl ester (27.5 g, 0.122 mol, 1.5 eq) in THF (10 mL), dropwise. After 10 minutes a solution of 3-(2,4-dichlorophenyl)-2-methylpropenal (17.56 g, 0.082 mol) in THF (50 mL) was added, dropwise. After stirring for 4 hours the mixture was added to $NH_4Cl_{(sat)}$ and extracted with EtOAc. The combined organic extracts were dried ($MgSO_4$), concentrated under reduced pressure to give a residue that was purified by silica gel chromatography (0-2.5-5% EtOAc in hexanes) to give 20.9 g of 5-(2,4-dichlorophenyl)-4-methylpenta-2,4-dienoic acid ethyl ester.

Step 3


Preparation of 5-(2,4-Dichlorophenyl)-4-methylpenta-2,4-dienoic acid



5-(2,4-Dichlorophenyl)-4-methylpenta-2,4-dienoic acid ethyl ester (2.37 g, 0.083 mol) was suspended in EtOH (50 mL). 2 N NaOH (13 mL, 3 eq) was added and the mixture was heated at 50°C for 1 hour. The mixture was cooled to room temperature and acidified to pH 1. The resulting solid was collected to give 2 g of 5-(2,4-dichlorophenyl)-4-methylpenta-2,4-dienoic acid.

Step 4

5-(2,4-Dichlorophenyl)-4-methylpenta-2,4-dienoic acid [3-(4-chlorophenyl)allyl]amide




5-(2,4-Dichlorophenyl)-4-methylpenta-2,4-dienoic acid (1 g, 0.0039 mol) was suspended in CH_2Cl_2 (10 mL). Oxalyl chloride (0.5 mL, 1.5 eq) was added, followed by 1 drop of DMF. After 2 hours the volatiles were removed, the residue taken up in THF (10 mL), and the mixture cooled to 0°C. A mixture of 3-(4-chlorophenyl)allylamine (0.7 g, 1.1 eq, 0.0042 mol, prepared as shown in Scheme 4) and Et_3N (0.82 mL, 0,0042 mol, 1.5 eq) in THF (5 mL) was added dropwise. After 1 hour, TLC analysis showed complete

conversion of the starting materials and the mixture was then diluted with EtOAc. The organic layer was washed with $NH_4Cl_{(sat)}$, dried ($MgSO_4$), and concentrated under reduced pressure. The resulting residue was purified by silica gel chromatography (0-10-20-30-40% EtOAc in hexanes) to give 1.25 g of 5-(2,4-dichlorophenyl)-4-methylpenta-2,4-dienoic acid [3-(4-chlorophenyl)allyl]amide.


Step 5

Preparation of [3-(4-Chlorophenyl)allyl]-[5-(2,4-dichlorophenyl)-4-methylpenta-2,4-dienoyl]carbamic acid tert-butyl ester




5-(2,4-Dichlorophenyl)-4-methylpenta-2,4-dienoic acid [3-(4-chlorophenyl)allyl]amide (1.25 g, 0.0032 mol) was dissolved in CH_2Cl_2 (20 mL). To this mixture was added, Et_3N (0.443 mL, 1 eq), Boc_2O (1.39 g, 2eq), and DMAP (0.39 g, 1 eq). After stirring for 16 hours, the mixture was diluted with CH_2Cl_2, washed with $NH_4Cl_{(sat)}$, and dried ($MgSO_4$). Silica gel chromatography (5 % EtOAc in hexanes) gave 1.43 g of [3-(4-Chlorophenyl)allyl]-[5-(2,4-dichlorophenyl)-4-methylpenta-2,4-dienoyl]carbamic acid tert-butyl ester.

Step 6




[3-(4-Chlorophenyl)allyl]-[5-(2,4-dichlorophenyl)-4-methylpenta-2,4-dienoyl]carbamic acid tert-butyl ester (1.25 g, 0.00247 mol) was dissolved in toluene (60 mL) and heated at 150°C for 2.5 hours. The mixture was concentrated and the residue purified by silica gel chromatography (5-15-20-40% EtOAc in hexanes) to give, in order of elution, 0.46 g of **132** LCMS: 452 (MH^+-Bu^t), and 0.46 g of **133** LCMS: 452 (MH^+-Bu^t).

General Scheme D




Preparation of Compound 134



134

Compound **132** (120 mg, 0.000236 mol) was dissolved in CH_2Cl_2 (2 mL) and cooled to 0°C. TFA (i.e., trifluoroacetic acid (2 mL)) was added and the mixture was stirred for 30 min. Removal of volatiles gave 105 mg of the compound **134**. LCMS: 406.2, 408.2 (MH^+)


Preparation of Compound **135**



135

Starting from compound **133**, compound **135** was produced using a procedure similar to the procedure used to prepare **134**. LCMS: 406.2, 408.2 (MH^+)


Preparation of Compound **136**



136

Compound **134** (100 mg, 0.000246 mol) was suspended in $MeCN/CH_2Cl_2$ (2mL/2mL), DBU (0.111 mL, 3 eq) was added and the mixture stirred for 4 hours. The mixture was diluted with EtOAc and washed with $NH_4Cl_{(sat)}$, the organic layer was dried ($MgSO_4$) and concentrated under reduced pressure to give 40 mg of **136**. LCMS: 406.2, 408.2 (MH^+)


Preparation of Compound **137**.



137

Compound **136** (30 mg, 0.000074 mol) was dissolved in EtOAc. PtO_2 (3 mg) was added and the mixture stirred under 1 atm of H_2 for 4 hours. The catalyst was removed by filtration through a pad of CELITE and the filtrate was concentrated to give 29 mg of compound **137**. LCMS: 408.2, 410.2 (MH^+)


Preparation of Compound **138**



138

Compound **132** (100 mg, 0.000197 mol) was dissolved in DMF (1 mL), MeI (0.246 mL, 20 eq). NaH (11.8 mg of a 60% dispersion in mineral oil, 1.5 eq) was then added. After stirring for 2 hours, the reaction was quenched with $NH_4Cl_{(sat)}$, extracted with EtOAc, and the combined organic extracts were dried ($MgSO_4$) and concentrated under reduced pressure. The resulting residue was purified using silica gel chromatography to give 80 mg of compound **138**. LCMS: 466.3 (MH^+-Bu^t).


Preparation of Compound **139**



139

Compound **138** (80 mg, 0.000154 mol) was dissolved in CH_2Cl_2 (5 mL) and cooled to 0°C. TFA (5 mL) was added and the mixture was stirred for 30 min. Removal of volatiles followed by silica gel chromatography (0-30-50-70% EtOAc in hexanes) gave 35 mg of compound **139**. LCMS: 420.2 (MH^+)


Preparation of Compounds **140** and **141**



140 **141**

Using procedures similar to those employed for the synthesis of compounds **132** and **133**, compounds **140** (LCMS: 416.2 (MH^+-Bu^t)) and **141** (LCMS: 416.2 (MH^+-Bu^t)) were synthesized.

Preparation of Compound **142**




142

Compound **142** was prepared from compound **140** using a procedure similar to that employed for the synthesis of compound **134**.

Compound **142** LCMS: 372.2 (MH^+).

Preparation of Compound **143**




143

Compound **143** was prepared from compound **141** using a procedure similar to that employed for the synthesis of compound **134**.

Compound **143** LCMS: 372.2 (MH^+).


Preparation of Compound **144**



Compound **144** was prepared from compound **142** using a procedure similar to that employed for the synthesis of compound **136**.

Compound **144** LCMS: 372.2 (MH^+).


Preparation of Compound **145**



Compound **145** was prepared from compound **144** using a procedure similar to that employed for the synthesis of compound **137**.

Compound **145**, LCMS: 374.2 (MH^+).


Preparation of Compound **146**



Compound **146** was prepared from compound **140** using a procedure similar to that employed for the synthesis of compound **138**.

Compound **146** LCMS: 430.2 (MH^+-Bu^t).


Preparation of Compound **147**



Compound **147** was prepared from compound **146** using a procedure similar to that employed for the synthesis of compound **139**.

Compound **147** LCMS: 386.2 (MH^+).


General Scheme E



Preparation of Compounds **148** and **149**

Step 1


Preparation 5-(2,4-Dichlorophenyl)-3-methylpenta-2,4-dienoic acid ethyl ester.



Triethyl 3-methyl-4-phosphonocrotonate (34.24 g, 0.129 mol, 1.3 eq) was dissolved in THF/DMPU (100 mL/200 mL) and the mixture was cooled to -78°C. BuLi (51.8 mL of a 2.5M solution in hexanes, 1.3 eq) was added dropwise. The mixture was stirred for 20 min and then a solution of 2,4-dichlorobenzaldehyde (17.6 g, 0.1 mol) in THF/DMPU (20 mL/40 mL) was added dropwise. The resulting mixture was stirred for 1 hour and then allowed to warm to room temperature. $NH_4Cl_{(sat)}$ was added and the mixture extracted with EtOAc. The organic layers were washed with, water, brine, and dried ($MgSO_4$). Removal of solvent under reduced pressure followed by silica gel chromatography (0-1-2-3% EtOAc in hexanes) gave 23.3g of 5-(2,4-dichlorophenyl)-3-methylpenta-2,4-dienoic acid ethyl ester.

Step 2


Preparation 5-(2,4-Dichlorophenyl)-3-methylpenta-2,4-dienoic acid



5-(2,4-Dichlorophenyl)-3-methylpenta-2,4-dienoic acid ethyl ester (5 g, 0.0184 mol) was dissolved in THF (125 mL). Water (50 mL), MeOH (50 mL), and NaOH (18.44 mL of a 2M solution in water, 2 eq) were then added. After 6 hours, TLC analysis showed complete conversion of the starting materials and the mixture was acidified to pH 1 by addition of 6N HCl. The resulting solid was collected to give 4.24 g of 5-(2,4-dichlorophenyl)-3-methylpenta-2,4-dienoic acid.

Step 3


Preparation of [3-(4-Chlorophenyl)allyl]-[5-(2,4-dichlorophenyl)-3-methylpenta-2,4-dienoyl]carbamic acid tert-butyl ester



5-(2,4-Dichlorophenyl)-3-methylpenta-2,4-dienoic acid (4.24 g, 0.0165 mol) was suspended in CH_2Cl_2 (40 mL), oxalyl chloride (2.12 mL, 1.5 eq) was added followed by 1 drop of DMF. After 2 hours the volatiles were removed, the residue taken up in THF (34 mL), and the mixture cooled to 0°C. A mixture of 3-(4-chlorophenyl)allylamine (3.2 g, 1.1 eq, Scheme 4) and Et_3N (3.6 mL, 1.5 eq) in THF (20 mL) was added dropwise. After 1 hour, TLC analysis showed complete conversion of the starting materials and the mixture was then diluted with EtOAc. The organic layer was washed with $NH_4Cl_{(sat)}$, dried ($MgSO_4$), and concentrated under reduced pressure, and the resulting residue was taken up in CH_2Cl_2 (100 mL). To this mixture was added Et_3N (2.3 mL, 1 eq), Boc_2O (7.17 g, 2eq), and DMAP (2 g, 1 eq). After stirring for 16 hours, the mixture was diluted with CH_2Cl_2, washed with $NH_4Cl_{(sat)}$, and dried ($MgSO_4$). Silica gel chromatography (0-2-4 % EtOAc in


hexanes) gave 5.8 g of [3-(4-chlorophenyl)allyl]-[5-(2,4-dichlorophenyl)-3-methylpenta-2,4-dienoyl]carbamic acid tert-butyl ester.

Step 4




[3-(4-Chlorophenyl)allyl]-[5-(2,4-dichlorophenyl)-3-methylpenta-2,4-dienoyl]carbamic acid tert-butyl ester (5.8 g, 0.011 mol) was dissolved in toluene (350 mL) and heated at 110°C for 3 hours. The mixture was concentrated and the residue purified by silica gel chromatography (0-2.5-5-10-40% EtOAc in hexanes) to give, in order of elution, 2.4 g of compound **148** LCMS: 452 (MH^+-Bu^t), and 2.2 g of compound **149** LCMS: 452 (MH^+-Bu^t).

General Scheme **F**



Preparation of Compound **150**




150

Compound **150** was prepared from compound **148** using a procedure similar to that employed for the synthesis of compound **134**.

Compound **150**, LCMS: 406.2, 408.2 (MH^+).

Preparation of Compound **151**




151

Compound **151** was prepared from compound **149** using a procedure similar to that employed for the synthesis of compound **134**.

Compound **151**, LCMS: 406.2, 408.2 (MH^+).

Preparation of Compound **152**




152


Compound **152** was prepared from compound **150** using a procedure similar to that employed for the synthesis of compound **136**.

Compound **152**, LCMS: 406.2, 408.2 (MH^+).

Preparation of Compounds **153** and **154**



153




154

Compounds **153** and **154** were prepared by the hydrogenation of compound **152** using a procedure similar to that employed for the synthesis of compound **137**.

Compound **153**, LCMS: 408.2, 410.2 (MH^+).

Compound **154**, LCMS: 408.2, 410.2 (MH^+).

Preparation of Compound **155**




155

Compound **155** was prepared from compound **148** using a procedure similar to that employed for the synthesis of compound **138**.

Compound **155**, LCMS: 466.3 (MH^+-Bu^t).

Preparation of Compound **156**




156

Compound **156** was prepared from compound **155** using a procedure similar to that employed for the synthesis of compound **139**.

Compound **156**, LCMS: 420.2 (MH^+).

Preparation of 5,6-Di(hetero)aryl-isoindol-1-ones

General Scheme G:




Substituted 5,6-Di(hetero)aryl-isoindol-1-ones may be prepared by procedures shown in General Scheme G, above. An aryl or heteroaryl alkyne **p** can be coupled by the Sonogashira reaction to an aryl or heteroaryl iodide or bromide **q** to give the alkyne **r**. Palladium catalyzed addition of tributyltin hydride followed by treatment with iodine gives the vinyl iodide **t**. Heck coupling reaction with methyl acrylate followed by hydrolysis gives the dienoic acid **u** which was converted to the allylamide **v** either through the acid chloride or by EDCI (i.e., 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide) coupling. Intramolecular Diels-Alder reaction gave product **w** which can be converted to product **x** by hydrogenation or product **y** by alkylation.

Preparation of Compounds **157** and **158**

Step 1


Scheme 81:



A mixture of 1-chloro-4-ethynyl-benzene (2.71 g, 0.020 mol), 2,4-dichloro-1-iodo-benzene (3.23 mL, 0.024 mol), $PdCl_2(PPh_3)_2$ (0.10 g, 1.4 mmol), CuI (1.44 g, 7.6 mmol), and diisopropylamine (7 mL) in DCM (i.e., dichloromethane) (100 mL) was stirred at room temperature for 16 h. The mixture was filtered through CELITE. The filtrate was concentrated *in vacuo*. The residue was chromatographed (SiO_2, hexane) to afford 2,4-dichloro-1-(4-chloro-phenylethynyl)-benzene as a white solid (3.23 g, 87%).

Step 2


Scheme 82:



Tributyltin hydride (2.7 mL, 10 mmol) was added to a mixture of 2,4-dichloro-1-(4-chloro-phenylethynyl)-benzene (2.01 g, 7.13 mmol) and $PdCl_2(PPh_3)_2$ (0.45 g, 0.64 mmol) in THF (70 mL) at room temperature and stirred for 1 h. Iodine crystals (2.7 g, 11 mmol) were added to the mixture at room temperature and stirred for 30 min. The mixture was then stirred with sodium thiosulfate aqueous solution and KF aqueous solution. The mixture was filtered through CELITE. The organic layer of the filtrate was separated and concentrated *in vacuo*. The residue was chromatographed (SiO_2, hexane) to afford 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-iodo-vinyl]-benzene as a light yellow solid (2.43 g, 83%).

Step 3

Scheme 83:




To a solution of 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-iodo-vinyl]-benzene (270 mg, 0.66 mmol) in DMF (6 mL) was added methyl acrylate

(0.30 mL, 3.3 mmol), Et_3N (0.30 mL, 2.2 mmol), and $Pd(PPh_3)_2Cl_2$ (46 mg, 0.07 mmol). The mixture was stirred at 100°C for 16 h. The mixture was filtered through CELITE. The filtrate was concentrated *in vacuo*. The mixture was diluted with EtOAc and washed with NH_4Cl (sat.). The organic layer was dried ($MgSO_4$) and concentrated *in vacuo*. Flash chromatography of the residue on a silica gel column with EtOAc–hexane (5–95) as eluent gave the methyl ester (98 mg, 40%) as a clear oil.

To a solution of the methyl ester (98 mg, 0.27 mmol) in THF–MeOH (3 mL, 1–1) was added NaOH (1.5 mL, 10%). The mixture was stirred at room temperature for 1 h. The mixture was diluted with water and the organics were evaporated. The aqueous mixture was diluted with water, acidified with 10% HCl, and extracted with CH_2Cl_2. The organic layer was washed with brine, dried ($MgSO_4$), and concentrated *in vacuo* to give 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid (82 mg, 87%) as an off-white solid.

Step 4


Scheme 84:



The 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid was converted to 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid allylamide in a manner similar to the procedure used to prepare amide **g** in General Scheme A.

Step 5

Scheme 85:




The 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid allylamide (55 mg) and dihydroquinone (4 mg) were dissolved in *o*-xylene and heated in a closed pressure tube at 220°C under nitrogen for 16 h. After cooling to room temperature, the solution was diluted with EtOAc and hexanes, washed with 10% NaOH and brine, dried ($MgSO_4$), and concentrated. The residue was chromatographed (SiO_2, 9:1-2:1 CH_2Cl_2/EtOAc) to give **157** as a white solid (12 mg, 22%), LCMS: *m/e* 392 (MH^+), and **158** as a white solid (2 mg, 4 %), LCMS: *m/e* 392 (MH^+).

Preparation of Compounds **159** and **160**

Step 1


Scheme 86:



The 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid was also converted to 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid diallylamide in a manner similar to the procedure used to prepare amide **g** in General Scheme A.

Step 2:

Scheme 87:




The 5-(4-chloro-phenyl)-4-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid diallylamide (39 mg) and dihydroquinone (2 mg) was dissolved in toluene and heated in a closed pressure tube at 175°C under nitrogen for 16 h. After cooling to room temperature, the solution was diluted with EtOAc and hexanes, washed with 10% NaOH and brine, dried ($MgSO_4$), and concentrated. The residue was chromatographed (SiO_2, 4:1-3:2

hexane/EtOAc) to give **159** as an off-white solid (15 mg, 38%), LCMS: *m/e* 432 (MH^+), and **160** as an off-white solid (12 mg, 30%), LCMS: *m/e* 432 (MH^+).

Preparation of Compound **161**


Scheme 88:



Compound **159** (12 mg) and 1 drop of DBU (i.e., 1,8-diazabicyclo[5.4.0]undec-7-ene) were dissolved in CH_2Cl_2 (2 mL) and stirred at room temperature for 1 h. The solution was concentrated. The residue was chromatographed (SiO_2, 4:1-3:2 hexane/EtOAc) to give **161** as an off-white resin (7 mg, 58%), LCMS: *m/e* 432 (MH^+).

Preparation of Compound **162**


Scheme 89:



Compound **157** was converted to **162** in a manner similar to that used to prepare compounds **5** and **6** in Scheme **7**. LCMS: *m/e* 394 (MH^+).

Preparation of Compound **163**


Scheme 90:



To a solution of **157** (31 mg, 0.079 mmol) and *p*-cyanobenzylbromide (46 mg, 0.24 mmol) in DMF (6 mL) was added NaH (31 mg, 0.79 mmol, 60% in mineral oil) at room temperature and the mixture was stirred at room temperature for 16 h. The mixture was concentrated *in vacuo*. The residue was diluted with EtOAc and washed with NH_4Cl (sat.). The organic layer was dried ($MgSO_4$) and concentrated in vacuo. Chromatography of the residue on a preparative silica gel plate with EtOAc–hexane (40–60) as eluent gave **163** (10 mg, 25%) as a white solid. LCMS: *m/e* 507 (MH^+).

Alternative Preparation of 5,6-Diphenyl-isoindol-1-ones


General Scheme H:



An alternative method for preparing substituted 5,6-diphenyl-isoindol-1-ones is shown in General Scheme H, above. The bis aryl or heteroaryl substituted ester **z** may be prepared by a procedure similar to the procedure published in *Acta Chem Scand*, **1993**, *47*, 1112 (herein incorporated by reference in its entirety) and converted to the dienoic acid **ab** by well known procedures as shown in General Scheme H. The dienoic acid **ab** was then converted to the target compounds using procedures similar to those used in General Scheme G.


Preparation of Compounds **164-170**

Scheme 91:







2-(4-Chloro-phenyl)-3-(2,4-dichloro-phenyl)-acrylic acid methyl ester was prepared by a reported procedure (*Acta Chem Scand*, **1993**, *47*, 1112; herein incorporated by reference in its entirety) and converted to the dienoic acid 4-(4-chloro-phenyl)-5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid by well known procedures as shown in Scheme 92. Compounds **164** and **166** were prepared from 4-(4-chloro-phenyl)-5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid using procedures similar to those used to prepare compounds **157**, **158**, and **162**.

General Scheme I:




The amide **ai** was also alkylated to give **aj** which was converted to compounds **168-171** as shown in General Scheme I.

Compound #	Compound	LCMS *m/e* (MH^+)
168		432
169		482
170		507

Compound #	Compound	LCMS *m/e* (MH^+)
171		507

Preparation of 5,6-Di(hetero)aryl-isoindol-1,3-diones or 3,4-Di(hetero)aryl-cyclohex-3-ene Carboxylic Amides


General Scheme J:



Di(hetero)aryl-isoindol-1,3-diones or 3,4-di(hetero)aryl-cyclohex-3-ene carboxylic amides can be prepared by various methods, for example by the intermolecular Diels-Alder reaction of a diaryl-, diheteroaryl-, or aryl-heteroaryl-substituted diene with a pyrrole-2,5-dione or an acrylamide derivative, for example as shown in General Scheme J, above.

Preparation of Compounds **172-175**

Scheme 92:



Compounds **172-175** were prepared by converting the commercially available ketone 1-(4-chloro-phenyl)-propan-2-one to 3-(4-chloro-phenyl)-but-3-en-2-one with Eschenmoser's salt. 3-(4-Chloro-phenyl)-but-3-en-2-one was then converted to trifluoro-methanesulfonic acid 2-(4-chloro-phenyl)-1-methylene-allyl ester with triflic anhydride and a hindered pyridine base. Suzuki coupling gave 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene. Diels-Alder reaction of 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene with the different dienophiles shown above in Scheme 93 gave compounds **172-175**.

Step 1:

1-(4-Chloro-phenyl)-propan-2-one (8.6 g, 51 mmol) and Eschenmoser's salt (12.3 g, 66.3 mmol) were dissolved in glacial acetic acid and heated in a closed pressure tube at 125 °C under nitrogen for 1 h. After cooling to room temperature, the solution was concentrated. The residue was dissolved in


EtOAc, washed with saturated $NaHCO_3$ aqueous solution, dried ($MgSO_4$), concentrated, and chromatographed (SiO_2, 39:1-19:1 hexane/EtOAc) to give 3-(4-chloro-phenyl)-but-3-en-2-one as yellow oil (3.92 g, 43%).

Step 2:

Triflic anhydride (0.23 mL, 1.4 mmol) was added to a mixture of 3-(4-chloro-phenyl)-but-3-en-2-one (189 mg, 1.04 mmol) and 2,6-di-t-butyl-4-methylpyridine (0.32 g, 1.6 mmol) in CH_2Cl_2 (10 mL) and refluxed for 16 h. After cooling to room temperature, the solution was washed with 10% HCl solution and saturated NH_4Cl aqueous solution, dried ($MgSO_4$), concentrated, and chromatographed (SiO_2, hexanes) to give trifluoro-methanesulfonic acid 2-(4-chloro-phenyl)-1-methylene-allyl ester as a clear oil (136 mg, 41%).

Step 3


Scheme 93:



Trifluoro-methanesulfonic acid 2-(4-chloro-phenyl)-1-methylene-allyl ester was converted to 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene by the Suzuki procedure shown above.


Step 4

Scheme 94:



2,4-Dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene was converted to the respective products shown in Scheme 95 above by the same Diels-Alder procedure as is shown in Scheme 88.


Preparation of Compound **176**



176

Substituted 5,6-diphenyl-isoindol-1-dione **176** was prepared using procedures similar to those used to prepare compound **172** except that diene 4-chloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene was used instead of 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene.

Preparation of Compound **177**




177

Compound **171** was prepared using procedures similar to those used to prepare compound **172** except that 4-chloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene was used instead of 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene. (4-Chloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene was prepared using procedures similar to those used to prepare 2,4-dichloro-1-[2-(4-chloro-phenyl)-1-methylene-allyl]-benzene, as shown in Scheme 93, above.)






Preparation of Compounds **178-185**




General Scheme K:



au


Di(hetero)aryl-cyclohex-3-ene carboxylic amides can be prepared by the intermolecular Diels-Alder reaction of a diaryl-, diheteroaryl-, or aryl-heteroaryl-substituted diene with an unsaturated amide, or alternatively by the intermolecular Diels-Alder reaction of a diaryl-, diheteroaryl-, or aryl-heteroaryl-substituted diene with an unsaturated ester which is subsequently converted to an amide, for example as shown in General Scheme K, above. Compounds prepared by the methods shown in General Scheme K are shown in the following Table:

Compound	Structure	LCMS *m/e* (MH^+)
178		416
179		432
180		414
181		501
182		450

Compound	Structure	LCMS *m/e* (MH^+)
183		478
184		470
185		464

Preparation of Di(hetero)aryl-isobenzofuran-1-ones

General Scheme L:




Di(hetero)aryl-isobenzofuran-1-ones can be prepared by a variety of methods. For example, an aryl- or heteroaryl-substituted unsaturated alcohol or acid can be condensed with an aryl- or heteroaryl-substituted diene acid or acid chloride **c** to provide a triene ester **ay** by the method shown above in General Scheme L. The triene ester **ay** can be cyclized via an intramolecular Diels-Alder reaction to form a di(hetero)aryl-tetrahydro-isobenzofuran-1-one **az**, that can be further modified (e.g., by reduction, etc.) to form the saturated di(hetero)aryl-isobenzofuran-1-one **ba**.

Preparation of Compounds **186-191**


Step 1

Scheme 95:



A mixture of but-3-yn-2-ol (7.0 g, 0.10 mol), 2.5-dibromopyridine (15.89 g, 0.067 mol), $PdCl_2(PPh_3)_2$ (2.35 g, 3.4 mmol), CuI (2.55 g, 0.013 mol), and diisopropylamine (300 mL) was stirred at RT for 1 h. The mixture was filtered through CELITE. The filtrate was concentrated in vacuo. The residue was dissolved in a CH_2Cl_2, washed with saturated NH_4Cl aqueous solution, dried ($MgSO_4$), and concentrated. The residue was chromatographed (SiO_2, 7:3 hexane/EtOAc) to afford the alkyne alcohol product 4-(5-bromo-pyridin-2-yl)-but-3-yn-2-ol as a tan solid (13.24 g, 87%).


Scheme 96:



4-(4-Chloro-phenyl)-but-3-yn-2-ol was prepared as shown in Scheme 95.


Step 2

Scheme 97:



The 4-(5-bromo-pyridin-2-yl)-but-3-yn-2-ol prepared in Step 1 (2.26 g, 0.010 mol) was dissolved in EtOAc (50 mL). Lindlar's catalyst (1.13 g) was added and the mixture was shaken on a Parr shaker under 55 psi hydrogen for 16 h. The mixture was filtered and concentrated. The residue was chromatographed (SiO_2, 9/1 to 4/1 hexane/EtOAc) to afford 4-(5-bromo-pyridin-2-yl)-but-3-en-2-ol as a clear oil (1.12 g, 49%).


Scheme 98:



5-(2,4-Dichloro-phenyl)-penta-2,4-dienoic acid 3-(4-chloro-phenyl)-1-methyl-allyl ester was prepared as shown in Scheme 97 using 1 atmosphere hydrogen and 10-20% by weight Lindlar's catalyst.

Step 3


Scheme 99:



The 4-(5-bromo-pyridin-2-yl)-but-3-en-2-ol prepared in Step 2 was converted to 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid 3-(5-bromo-pyridin-2-yl)-1-methyl-allyl ester in a manner similar to the procedure used to

prepare 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid [3-(4-chloro-phenyl)-allyl]-amide in Scheme 5 above.


Scheme 100:



5-(2,4-Dichloro-phenyl)-penta-2,4-dienoic acid 3-(4-chloro-phenyl)-1-methyl-prop-2-ynyl ester was similarly prepared as shown in Scheme 101.


Step 4:

Scheme 101:



The 5-(2,4-dichloro-phenyl)-penta-2,4-dienoic acid 3-(5-bromo-pyridin-2-yl)-1-methyl-allyl ester (248 mg) prepared in Step 3 above was dissolved in *o*-xylene and heated in a pressure tube at 175°C under nitrogen for 1 h. After cooling to room temperature, the solution was concentrated. The residue was chromatographed (SiO_2, 19:1-5:1 hexane/EtOAc) to give the cycloadduct **186** as an off-white solid (141 mg, 49 %). LCMS: *m/e* 452.1 (MH^+). Also isolated were minor cycloadducts **187** (12 mg, 5 %), LCMS: *m/e* 452.1 (MH^+), and **188** (35 mg, 14 %), LCMS: *m/e* 452.1 (MH^+).

Scheme 102:




LCMS: *m/e* 407.2 (MH^+)

Compounds **189-191** were prepared in a manner shown in Scheme 101.

Preparation of Compounds **192** and **193**


Scheme 103:



The lactone **186** (140 mg) and $NiCl_2$ hexahydrate (110 mg) were dissolved in THF (5 mL) and MeOH (1 mL). NaBH4 (18 mg) was added at 0°C. The mixture was stirred at 0°C for 10 min. The mixture was diluted with EtOAc, washed with saturated NH_4Cl aqueous solution, dried ($MgSO_4$), and concentrated. The residue was separated by preparative TLC (SiO_2, 3:1 hexane/EtOAc) to give the reduced product **192** as a white solid (35 mg, 25%). LCMS: *m/e* 454.1 (MH^+). Also isolated was minor reduced product **193** (10 mg, 7%). LCMS: *m/e* 454.1 (MH^+).

Preparation of Compound **194**


Scheme 104:



Compound 192 was prepared by the reduction procedure described in Scheme 12.

Preparation of Compound **195**


Scheme 105:



Compound 193 was prepared by a procedure similar to that shown in Scheme 55, Step 2, except that methyl iodide was used instead of trisyl azide.


Preparation of Compound **195**

Scheme 106:




The bromopyridine **192** (6.5 mg), $Pd(PPh_3)_4$ (2 mg), 2,5-dichlorophenylboronic acid (7 mg), K_2CO_3 (22 mg), ethanol (0.2 mL), water (0.1 mL), and toluene (0.4 mL) was heated at 90°C for 1 h. The reaction mixture was diluted with EtOAc, washed with saturated NH_4Cl aqueous solution, dried ($MgSO_4$), and concentrated. The residue was subjected to preparative TLC purification (SiO_2, 3:2 hexane/EtOAc) to afford **196** as a white solid (6 mg, 81%). LCMS: *m/e* 520.1 (MH^+)

Preparation of Compounds **197** and **198**



Substituted 4-pyridyl-5-phenyl-isobenzofuran-1-ones **197** and **198** were prepared using procedures similar to those used to prepare compound **196** using the appropriate dienoic acid chloride **c** as shown in General Scheme L.


Preparation of Compounds **199-201**



199 **200** **201**


Compounds **199-201** were prepared as shown in Scheme 107, below:


Scheme 107:



Scheme 108:


Similar to the methylation of **126** to provide **127** (see scheme 73), other alkyl groups can be introduced at the alpha-position of lactam carbonyl. For example, an ethyl group can be introduced as described below:





Preparation of Compound **210**

Scheme 108:




A degassed solution of **126** (1.16g, 2.30 mmol) and 18-crown-6 (1.2g, 4.54 mmol, 2 eq.) in 20 ml THF was cooled to -78°C and a 0.5M solution of KHMDS in toluene (9.1ml, 4.55mmol, 2 eq.) was added and stirred for 1 hr. To this was added ethyl iodide (2.8ml, 34.68mmol, 15 eq.) and stirred for 2 hr at -78°C then quenched with aq. NH_4Cl. The THF was evaporated and the aqueous slurry was extracted with 3x ethyl acetate. The combined organic

layers was washed with brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 20% ethyl acetate in hexanes to provide 1.04 g of **210** as white solid.
MS: 476.3 ($M-{}^tBu^+$)

Preparation of Compounds **211**, **212**, **213** and **214**
These analogs were prepared using a similar procedures described under scheme 73 and scheme 80.
MS for **211**: 418.2 (MH^+)
MS for **212**: 550.3 (MH^+)
MS for **213**: 427.2 (MH^+)
MS for **214**: 446.2 (MH^+)

Scheme 109:
The phenolic functionality of **128** (see scheme 73) can be used as a versatile handle to introduce a variety of other groups. For example the phenol can be alkylated with appropriate alkyl halides to provide the ethers as presented below.



P

reparation of Compounds **215, 216** and **217**

These analogs were prepared using a procedure similar to the procedure used for the preparation of **131** (see scheme 80)


MS for **215**: 446.2 (MH^+)

MS for **216**: 458.3 (MH^+)

MS for **217**: 462.3 (MH^+)

Preparation of Compound **218**

Scheme 110:




A mixture of **128** (50mg, 0.12 mmol), trifluoroiodoethane (25μL, 1.27 mmol, 10 eq.) and K_2CO_3 (50mg, 0.36 mmol, 3 eq.) in 1ml DMF in a sealed tube was heated overnight at ~85°C. The mixture was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 3% methanol in dichloromethane to provide 55mg of **218**.

MS: 486.3 (MH^+)

Preparation of Compound **219**

Scheme 111:



A mixture of **128** (30mg, 0.072 mmol), iodoacetamide (67mg, 0.36mmol, 5 eq.) and K_2CO_3 (40mg, 0.29 mmol, 4 eq.) in 1ml DMF in a sealed tube was heated for 2.5h at ~60°C. The mixture was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 4% methanol in dichloromethane to provide 32mg of **219**.

MS: 475.3 (MH^+)

Preparation of Compounds **220** and **221**

Scheme 112:



A mixture of **128** (25mg, 0.044 mmol), (2-bromoethoxy)-*tert*-butyldimethylsilane (106μL, 10 eq.) and K_2CO_3 (28mg, 4 eq.) in 0.5ml DMF in a sealed tube was heated for 2h at ~65°C. The mixture was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 4% methanol in dichloromethane to provide 25mg of **220**.


This was dissolved in 0.5ml THF and stirred with 0.14ml of 1M solution of tetrabutylammonium fluoride in THF (0.14 mmol, 3 eq) for about 3 hr, diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 4% methanol in dichloromethane to provide 20mg of **221**.

MS for **220**: 562.3 (MH^+)

MS for **221**: 448.2 (MH^+)

Preparation of Compound **222**

Scheme 113:




A two-necked round bottom flask containing a solution of **128** (50mg, 0.124 mmol) and K_2CO_3 (68 mg, 0.49 mmol) in 2ml DMF was fitted with a cold finger. The cold finger was cooled by dry ice – acetone mixture and chlorodifluoromethane gas was introduced from a cylinder and allowed to


condense under the cold finger. The RB flask was immersed in an oil bath kept at ~40°C and stirred for 5 hr. The mixture was diluted with ethyl acetate, washed with 3x 1N HCl, brine, dried over $MgSO_4$, filtered and evaporated. The crude product was purified by chromatography using 0% to 3% methanol in dichloromethane to provide 22mg of **222**.

MS: 454.2 (MH^+)

Preparation of Compound **224**

Scheme 114:





224


A mixture of **128** (50mg, 0.124 mmol), Na_2CO_3 (8mg, 0,076 mmol, 0.6 eq.), $[Ir(COD)Cl]_2$ (8 mg, 0.012 mmol, 0.1 eq.) and vinyl acetate (57µl, 0.616 mmol, 5 eq.) in 1ml toluene in a sealed tube was bubbled with argon and heated at ~105°C for 8 hr. The mixture was diluted with ethyl acetate, washed 2x with water, brine, dried over $MgSO_4$, filtered and chromatographed with 0% to 3% methanol in dichloromethane to provide 45mg of **223**. (Refer to *Organic Synthesis*, volume 82, page 55-58 for literature procedure).

To a solution of **223** (45 mg, 0.105 mmol) in 2ml dichloromethane at at 0°C was added a 1M solution of diethyl zinc in hexanes (0.52 mmol, 0.52 mmol, 5 eq.) followed by trifluoroacetic acid (39 µl, 0.52mmol, 5 eq.) and stirred for 10 min. To this was added diiodomethane (42µl, 0.52 mmol, 5 eq.) and mixture stirred overnight at rt. It was quenched with aq. NH_4Cl, extracted with 3x ethyl


acetate and combined organic layer washed with brine, dried over $MgSO_4$, filtered, and chromatographed with 0% to 3% methanol in dichloromethane to provide 38mg of product which contained some unreacted starting materials. This product was once again treated with 10eq. each of diethyl zinc, trifluoroacetic acid and diiodomethane under the above conditions and after work-up purified as above to provide 34 mg of **224**.

MS: 444.2 (MH^+)


Preparation of Compounds **225-229**




225




226



227



228



229

These compounds were prepared starting with **211** and using procedures described for the preparation of compounds under scheme 109.

MS for **225**: 520.3 MS (MH^+)

MS for **226**: 576.3 MS (MH^+)


MS for **227**: 462.3 MS (MH^+)

MS for **228**: 475.3 MS (MH^+)

MS for **229**: 474.3 MS (MH^+)


Scheme 115

Amide analog such as **232** can be prepared from the carboxylic acid **231**. The carboxylic acid **231** in-turn can readily be obtained from the phenol **128**.



Preparation of Compound **230**

Scheme 116:




A mixture of **128** (300mg, 0.742 mmol), methyl bromoacetate (350µl, 3.7mmol, 5 eq.) and K_2CO_3 (410mg, 2.97mmol, 4 eq.) in 5ml DMF was heated for 4h at ~60°C. The mixture was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 4% methanol in dichloromethane then filtered through a pad of basic alumina to provide 305mg of **230**.

MS: 476.3 (MH^+)

Preparation of Compound **231**

Scheme 117:




To a solution of **230** (300mg, 0.63 mmol) in 2ml each of methanol and THF was added 2ml of 1M aq. NaOH solution and the mixture stirred at rt for 2hr. It was diluted with water, acidified with 1N HCl and extracted with 3x ethyl acetate. The combined organic layers was washed with brine, dried over $MgSO_4$, filtered and concentrated to provide 250mg of **231**.

MS: 462.3 (MH^+)

Preparation of Compound **232**

Scheme 118:




To solution of **231** (50mg, 0.108mmol), N-methyl ethanolamine (41mg, 0.546mmol, 5 eq.), HOBt (30mg, 0.22mmol, 2 eq.) in 1ml DMF and 0.5ml dichloromethane was added EDCI (42mg, 0.219mmol, 2 eq.) and the mixture stirred overnight at rt. The solution was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 5% methanol in dichloromethane to provide 52mg of **232**.

MS: 519.3 (MH^+)

Preparation of Compound **234**

Scheme 119:




To solution of **231** (110mg, 0.238mmol), aminoacetaldehyde dimethyl acetal (125mg, 1.19 mmol, 5 eq.), HOBt (65mg, 0.48mmol, 2 eq.) in 1.5ml DMF and 0.5ml dichloromethane was added EDCI (92mg, 0.48mmol, 2 eq.) and the mixture stirred overnight at rt. The solution was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by chromatography using 0% to 5% methanol in dichloromethane to provide 51mg of **233**.

This was stirred with 3ml of 1:1 trifluoroacetic acid and dichloromethane and few drops of water at rt for 3hr. It was diluted with ethyl acetate, washed 2x with water, brine, dried over $MgSO_4$, filtered and concentrated to provide 41mg of aldehyde. This was dissolved in 1ml of 1:1 methanol and dichloromethane, cooled to 0°C and treated with sodium borohydride (4mg, 0.10mmol, 1 eq) for 2 min. It was quenched with aq. ammonium chloride, extracted 3x with ethyl acetate, combined organic layers washed with brine, dried over $MgSO_4$, filtered, evaporated and purified by preparative TLC using 5% methanol in dichloromethane to provide 38mg of **234**.


MS: 505.3 (MH^+)

Preparation of Compounds **235-237**


Scheme 120:



235



236




237

These compounds were prepared using similar procedures described under scheme 115.

MS for **235**: 504.3 (MH^+)
MS for **236**: 476.3 (MH^+)
MS for **237**: 489.3 (MH^+)

Scheme 121:



Preparation of **250**:

To a degassed solution of **120** (400 mg, 0.786 mmol) in 5ml THF at -78°C was added 0.5M solution of KHMDS in toluene (2.4 ml, 1.2 mmol, 1.5 eq.) and stirred for 30 min. To this was added methyl cyanoformate (125 μl, 1.58 mmol, 2 eq.) and stirred for 30 min. The reaction was quenched with ferrous ammonium sulfate, and extracted with 3x ethyl acetate. The combined organic layers was washed with brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 20% ethyl acetate in hexanes to provide 394 mg of **250**.

MS: 566.3 (MH^+)

Preparation of **251**:

To a solution of **250** (0.459 mmol) in 3ml dichloromethane at rt was added 1M BBr_3 solution in dichloromethane (2.3 ml, 2.3 mmol, 5 eq.) and stirred for 2 hr. The reaction was quenched by the addition of water, extracted 3x with dichloromethane. The combined organic layer was washed with brine, dried over $MgSO_4$, filtered and concentrated to provide 177 mg of **251** as a solid.

MS: 452.2 (MH^+)

Preparation of **252-254**:

To a solution of **251** (20 mg, 0.044 mmol), pyrrolidine (11µl, 0.134 mmol, 3 eq), HOBt (9 mg, 0.067 mmol, 1.5 eq) in 0.75 ml dichloromethane at rt was added EDCI (13 mg, 0.068 mmol, 1.5 eq) and stirred overnight at rt. It was diluted with ethyl acetate, washed with 2x 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 3% methanol in dichloromethane to provide 10 mg of **252**.

Using a similar procedure, compounds **253** and **254** were prepared.

MS for **252**: 505.3 (MH^+)

MS for **253**: 491.3 (MH^+)

MS for **254**: 495.3 (MH^+)

Preparation of **255**:

A solution of **250** (390 mg, 0.688 mol) in 1:1 dichloromethane - trifluoroacetic acid was stirred at 0°C for 1 hr and concentrated to provide 320 mg of **255**.

MS: 466.3 (MH^+)

Preparation of **256**:


To a solution of **255** (300 mg, 0.643 mmol) in 3 ml THF at rt was added 2M solution of $LiBH_4$ in THF (1 ml, 2 mmol, 3 eq.) and stirred for 1.5 hr. It was quenched with aq. NH_4Cl, and extracted 3x with ethyl acetate. The combined organic layer was washed with brine, dried over $MgSO_4$, filtered, concentrated

and chromatographed with 0% to 3% methanol in dichloromethane to provide 256 mg of **256**.
MS: 438.2 (MH^+)


Preparation of **257** and **258**:
To a solution of **256** (50 mg, 0.114 mmol) in 1 ml DMF at 0°C was added 60% NaH in mineral oil (4.6 mg, 0.115 mmol, 1 eq) followed by iodomethane (35 µl, 0.562 mol, 5eq).

The mixture was stirred for 2 hr at 0°C and 1 hr at rt. It was diluted with ethyl acetate, washed 3x with water, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 4% methanol in dichloromethane followed by preparative TCL purification of the overlapped fractions in 3% methanol in dichloromethane to obtain 14 mg of **257** and 16 mg of **258**.
MS for **257**: 452.2 (MH^+)
MS for **258**: 466.3 (MH^+)

Scheme 122:
Compound **120** was converted to compounds **259**, **260** and **261** using procedure similar to conversion of **81** to **84** (see scheme 47) using appropriate benzylic bromides.




Scheme 123:



MS: 450.2 (MH^+)

The intermediate **120** alkylated with allylbromide similar to the alkylation of **81** to give **86** then the double bond was reduced using conditions similar to the conversion of **79** to **80** (see scheme 47).

Scheme 124:



Preparation of **263** and **264**:


A solution of **127** (30 mg) in 0.5 ml each of dichloromethane and trifluoroacetic acid was stirred at 0 °C for 40 min, concentrated and chromatographed with 0% to 3% methanol in dichloromethane to provide 21 mg of **263**.

Intermediate **120** was ethylated using similar condition used in scheme 108 then the Boc group was cleaved using the above deprotection conditions to give **264**.

MS for **263**: 418.2 (MH^+)

MS for **264**: 432.2 (MH^+)

Scheme 125:




Preparation of **265**:

A mixture of **211** (70 mg, 0.167 mmol), (R)-2,2-dimethyl-1,3-dioxolan-4-ylmethyl p-toluenesulfonate (145 mg, 0.506 mmol, 3 eq.), K_2CO_3 (93 mg, 0.67 mmol, 4 eq), NaI (25 mg, 0.166 mmol, 1 eq) in 2 ml DMF was heated at 80°C in sealed tube for 2 days. It was diluted with ethyl acetate, washed with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 4% methanol in dichloromethane to provide 82 mg of a mixture of alkylated product and unreacted starting material. This was stirred with 1.5 ml of 1:3 v/v con. HCl and dioxane at rt for 4 hr. The mixture was diluted with ethyl acetate, washed with water, aq. $NaHCO_3$, and brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 5% methanol in dichloromethane to provide 31 mg of **265**.

MS: 492.3 (MH^+)

Using a similar procedure, the following compounds were prepared:




266 MS:492.3 (MH^+)

267 MS: 478.3 (MH^+)

268 MS: 518.3 (MH^+)

Scheme 126:



129 269 270


A mixture of **129** (30 mg, 0.056 mmol) and pyrrolidine (46 µl, 0.56 mmol, 10 eq.) in 1 ml NMP in a sealed tube was heated in a microwave reactor at 200 °C for 60 min. It was diluted with ethyl acetate, washed with water, brine, dried over $MgSO_4$, filtered, concentrated and purified by preparative TLC using 4% methanol in dichloromethane to provide 17 mg of **269**.

Using a similar procedure **270** was prepared.

MS for **269**: 457.3 (MH^+)

MS for **270**: 473.3 (MH^+)

Scheme 127:



Preparation of **271**:

To a solution of **229** (5 mg) in 1 ml of 1:1 v/v dichloromethane-methanol mixture at rt was added excess $NaBH_4$ and stirred at rt for 10 min. It was quenched with water, extracted with ethyl acetate, washed with brine, dried over $MgSO_4$, filtered, concentrated and purified by preparative TLC using 4% methanol in dichloromethane to provide 4 mg of **271**.


MS: 476.3 (MH^+)

Preparation of **272**:

To a solution of **229** (60 mg, 0.126 mmol)in 2 ml THF at -78 °C was added 1.4 M solution of MeMgBr in toluene/THF mixture. The reaction mixture was stirred at -78°C for 30 min then at rt for 1.5 hr. It was quenched with aq. NH_4Cl, extracted 3x with ethyl acetate. The combined organic layer was washed with brine, dried over $MgSO_4$, filtered, concentrated and purified by preparative TLC using 4% methanol in dichloromethane to provide 20 mg of **272**.

MS: 490.3 (MH^+)

Scheme 128:



Preparation of **273** and **274**:


To a solution of **235** (65 mg, 0.129 mmol) in 2 ml THF at -78°C was added $Ti(O^iPr)_4$ (115 µl, 0.388 mmol, 3 eq.) followed by 3M solution of EtMgCl in ether (0.39 ml, 1.17 mmol, 9 eq.). The mixture was stirred for 1 hr at -78°C and 2 hr at rt. It was quenched by the addition of aq. NH_4Cl, stirred for 20 min and diluted with aq. sodium potassium tartrate. The slurry was extracted 3x with ethyl acetate, the combined organic layers washed with aq. sodium potassium tartrate followed by brine. It was dried over $MgSO_4$, filtered, concentrated and purified by preparative TLC using 4% methanol in dichloromethane to provide 33 mg of **273**.

Similarly, **230** was converted to **274**.

MS for **273**: 518.3 (MH^+)

MS for **274**: 504.3 (MH^+)

Scheme 129:




Preparation of **275**:

A mixture of **213** (25 mg, 0.059 mmol), NaN_3 (39 mg, 0.60 mmol) and NH_4Cl (32 mg (0.59 mmol) in 0.5 ml DMF in a sealed tube was heated overnight at 120 °C. The mixture was diluted with 1N HCl and extracted 3x with ethyl acetate. The combined organic layer was washed with brine, dried over $MgSO_4$, filtered and concentrated to provide ~30 mg **275**.

MS: 470.3 (MH^+)

Scheme 130:



Preparation of **276**:


To a solution of **130** (55 mg, 0.133 mmol), in 2 ml ether was added $Ti(O^iPr)_4$ (120µl, 0.401 mmol, 3 eq.) cooled to -78°C and added 3M solution of EtMgBr in ether (0.27 ml, 0.81 mmol, 6 eq.) and stirred for 10 min at -78°C and for 1 hr at rt. To this was added $BF_3.OEt_2$ (101 µl, 0.80 mmol, 6 eq) and stirred for 1 hr at rt. The mixture was poured in to aq. sodium potassium tartrate and extracted 3x with ethyl acetate. The combined organic layer was washed with brine, dried over $MgSO_4$, filtered concentrated and


chromatographed with 0% to 4% methanol in dichloromethane to provide 26 mg of **276**.

MS: 444.2 (MH^+)


Scheme 131:

Using chemistry similar to the alkylation conditions given under scheme 73 and scheme 109, intermediates **126** and **120** were transformed to compounds **277-281**.






Similarly:




Scheme 132:

Using the conditions similar to the transformations described under scheme 109, intermediates **211** and **128** were converted to compounds **282-287**.




Scheme 133:

Using conditions similar to the transformation of intermediate **124** to **221**(see scheme 73 and scheme 109) intermediate **124** was transformed to compounds **288-292.**


BocN
KHMDS
MeI
OMe
Cl
124
Me
288
MS: 460.3 (M-tBu+)
BBr3
HN
OH
289
MS: 402.2 (MH+)
1) BrCH2CH2OTBS, K2CO3, DMF
2) TBAF, THF
290
MS: 446.2 (MH+)
Similarly:
124
291
MS: 416.2 (MH+)
292
MS: 460.3 (MH+)

Scheme134:



Preparation of compound **295**:

To 250 mg of compound **120** was added 18-crown-6 and the mixture twice dissolved in toluene and evaporated to dryness. To the residue in 3 ml of dry THF at -78°C was added two equivalents of potassium hexamethyldisilylamide (as a 0.5M solution in toluene) and the mixture stirred under argon for 1.5 hours after which the reaction mixture was stirred under a balloon of O_2 for two hours. The mixture was then quenched with aqueous sodium sulfite and allowed to warm to room temperature. The mixture was then extracted three times with ethyl acetate and the combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to

dryness. Purification by flash chromatography (0-40% acetone in hexane) yielded 96 mg of compound **293**.

To 45 mg of compound **293** was added 18-crown-6 and the mixture twice dissolved in toluene and evaporated to dryness. To the residue in 3mL of dry THF at -78°C was added two equivalents of potassium hexamethyldisilylamide (as a 0.5M solution in toluene) and the mixture stirred under argon for about one hour after which ten equivalents iodoethane was added and, after two hours, the mixture placed in a freezer overnight. The reaction mixture was quenched with aqueous ammonium chloride and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (0-50% ethyl acetate in hexane) yielded 12 mg of compound **294**.

To 12 mg of compound **294** in 2 mL of dry dichloromethane at 0°C was added 0.5 mL of trifluoroacetic acid and the mixture stirred under nitrogen. After one hour the mixture was evaporated to dryness and twice dissolved in toluene and evaporated to dryness yielding 10 mg of **295**.
MS for **295**: 452.2 (MH^+)

Preparation of **296**:

To 30 mg of **293** in 2 ml of dry DMF at 0 °C was added 2 equivalents of sodium hydride (5mg of 60% NaH in mineral oil) and the mixture stirred under nitrogen. After 15 minutes 5 equivalents of iodomethane was added and the mixture slowly warmed to room temperature. After a further two hours the mixture was quenched with aqueous ammonium chloride and extracted with ethyl acetate three times. The combined organic extracts were washed with brine, dried with magnesium sulfate, filtered and evaporated to dryness. Purification by flash chromatography (0-30% EtOAc in hexane) yielded 15 mg of **296**.
MS: 538.3 (MH^+)

Preparation of **297**:

To 13 mg of **296** in 1.5 ml of dichloromethane at 0 °C was added 1 ml of trifluoroacetic acid and the mixture stirred under nitrogen for two hours.

The reaction mixture was then evaporated to dryness and twice dissolved in toluene and evaporated to dryness. The crude product was purified by reversed phase HPLC yielding 5 mg of **297.**


MS: 438.2 (MH^+)

Preparation of **298:**

To 30 mg of **293** in 2 ml of dichloromethane at 0 °C was added 1 ml of trifluoroacetic acid and the mixture stirred under nitrogen for two hours. The reaction mixture was then evaporated to dryness and twice dissolved in toluene and evaporated to dryness yielding 24mg of **298.**

MS: 424.2 (MH^+)

Scheme 135:



To 100 mg of **120** in 3 ml of dry THF at -78°C was added 2 equivalents of 18-crown-6 and 2 equivalents of potassium hexamethyldisilylamide (as a 0.5M solution in toluene) and the mixture stirred under argon for 15 minutes after which 10 equivalents of allyl iodide was added. After 1.5 hours the mixture was poured onto aqueous ammonium chloride and extracted three

times with ethyl acetate. The combined extracts were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. The crude mixture was purified by flash chromatography (0 to 20% EtOAc in hexane) yielding 61mg of a mixture of compounds **299** and **300**.

To this mixture in 2 ml of dichloromethane at 0°C was added 1 mL of trifluoroacetic acid and the mixture stirred under nitrogen for one hour. The reaction mixture was then evaporated to dryness and twice dissolved in toluene and evaporated to dryness yielding 60 mg of a mixture of compounds **301** and **302**.


To the mixture of **301** and **302** in 3 mL of dioxane and 1mL of water was added 2 equivalents of 2,6-lutidine, 4 equivalents of sodium periodate and 2% mole of osmium tetroxide (as a 2.5% solution in t-butanol) and the mixture stirred under nitrogen. After five hours the mixture was placed in a freezer for 16 hours after which the mixture was poured onto water and extracted three times with dichloromethane. The combined extracts were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness yielding 60 mg of a mixture of compounds **303** and **304**.

To the mixture of **303** and **304** in 5 mL of methanol was added ~10 mg of sodium borohydride and the mixture stirred under nitrogen for 10 minutes then quenched with aqueous ammonium chloride and extracted three times with dichloromethane. The combined extracts were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. The crude product was purified by reversed phase HPLC yielding 15 mg of **305** and 12mg of **306**.

MS for **305**: 452.2 (MH^+)

MS for **306**: 468.3 (MH^+)

Scheme 136:



Preparation of **308**:

To 98 mg of **256**, in 5 mL of dry dichloromethane, was added two equivalents of sodium bicarbonate and 1.2 equivalents of Dess-Martin reagent and the mixture stirred under nitrogen. After one hour aqueous sodium bicarbonate and ether were added to the reaction mixture and allowed to stir for ten minutes. The phases were separated and the aqueous phase extracted with diethyl ether. The combined organic phased were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness yielding 90 mg of aldehyde **307**.

To 90 mg of **307** in 5 mL of dry THF at 0°C was added three equivalents of methyl magnesium bromide (as a 1.4M solution in toluene) and the mixture stirred under nitrogen. After one hour an additional equivalent of methyl magnesium bromide (as a 1.4M solution in toluene) was added and, after a further 20 minutes, the reaction mixture was quenched with aqueous ammonium chloride and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness yielding 100mg of crude product. Purification by flash chromatography (0-2% MeOH in DCM) yielded 6 mg of **308**.
MS: 452.2 (MH^+)


Preparation of **310**:

To 22 mg of **308** in dry dichloromethane were added two equivalents of sodium bicarbonate and 1.2 equivalents of Des Martin reagent and the mixture stirred under nitrogen. After two hours the reaction mixture was poured onto aqueous sodium bicarbonate and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness yielding 26 mg of crude ketone **309**.


To 26 mg of crude ketone **309** in 5 mL of dry THF at 0°C was added five equivalents of methyl magnesium bromide (as a 1.4M solution in toluene) and the mixture stirred under nitrogen. After three hours the reaction mixture was quenched with aqueous ammonium chloride and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (0-2% MeOH in DCM) yielded 1.2 mg of **310**.







MS: 466.3 (MH^+)








Scheme 137:

















Similar to the preparation of amide **112** described under scheme 59 and scheme 63, a variety of amides, sulfonamides, ureas and carbamates were prepared using appropriate acid chlorides, sulfonyl chlorides, isocyanates and chloroformates to give **311A** to **311BB**.

Compound#	Structure	M+H
311A		477.3


311B		491.3
311C		503.3
311D		503.3
311E		505.3
311F		507.3
311G		507.3

311H		507.3
311I		514.3
311J		519.3
311K		520.3
311L		525.3
311M		533.3
311N		538.3

311O		519.3
311P		467.3
311Q		481.3
311R		495.3
311S		509.3
311T		511.3
311U		521.3

311V		480.3
311W		494.3
311X		508.3
311Y		508.3
311Z		520.3
311AA		553.3
311BB		534.3

Scheme 138:



Preparation of **312**:


To **111** in 2mL of dry dichloroethane was added 5 equivalents of propionaldehyde and 5 equivalents of sodium triacetoxy borohydride and the mixture stirred under nitrogen. After about 16 hours the reaction mixture was poured onto aqueous potassium carbonate and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (0-35% ethyl acetate in hexane) yielded 25mg of **312**.
MS: 493.3 (MH^+)

Preparation of **313**:













To 40 mg of **111** in 2mL of dry dichloroethane was added 5 equivalents of acetaldehyde and 5 equivalents of sodium triacetoxy borohydride and the mixture stirred under nitrogen. After 16 hours the reaction mixture was poured onto aqueous potassium carbonate and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. The crude product was purified by reversed phase HPLC yielding 5mg of **313**.



MS: 465.3 (MH^+)

Scheme 139:




Intermediate **120** was converted to amine **314** then to a variety of amino derivatives **315A-315Q** using a procedures similar to the transformation of **108** to **112** (see scheme 59)

Compound #	R	R'	R''	MS (MH^+)
315A	H	H	H	423.2
315B			H	625.3
315C	H		H	525.3
315D			H	601.3
315E	H		H	501.3
315F			H	639.4
315G	H		H	539.3
315H			H	615.3
315I	H		H	515.3

315J			H	607.3
315K	H		H	507.3
315L			H	591.3
315M	H		H	491.3
315N	H		H	477.3
315O	H	Me	H	437.2
315P	H	Et	Et	479.3
315Q	H	n-Pr	n-Pr	507.3


Scheme 140:



To 60 mg of **128** in 1 mL of dry DMF was added four equivalents of potassium carbonate and five equivalents of 2-(chloromethyl)-3,5-dioxahex-1-ene. After bubbling with argon for one minute, the mixture was heated to 65°C in a pressure tube for three hours. The reaction mixture was diluted with ethyl acetate, washed three times with water, once with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (hexane to 95:5 DCM/MeOH) yielded 61 mg of **316**.

MS: 504.3 (MH^+)

Scheme 141:



Preparation of **318**:

To 750 mg of **128** in 10 mL of dry DMF was added ten equivalents of potassium carbonate and ten equivalents of N(2-bromoethyl)phthalimide. After bubbling with argon for one minute, the mixture was heated to 75°C in a pressure tube for eight hours. An additional five equivalents of potassium carbonate and five equivalents of N(2-bromoethyl)phthalamide were added and the mixture heated at 75°C in a pressure tube overnight. The reaction mixture was diluted with ethyl acetate, washed three times with water, once with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (hexane to 95:5 DCM/MeOH) yielded 1.08g of compound **317**.

To 30 mg of compound **317** in 2 mL of dichloromethane was added 300 µL of hydrazine and the mixture stirred under nitrogen overnight. The reaction mixture was then washed three times with water and then extracted with 1N aqueous HCl. The acidic aqueous phase was washed three times with dichloromethane. The acidic aqueous phase was made basic with aqueous potassium carbonate then extracted three times with dichloromethane. The basic extracts were combined, dried with $MgSO_4$, filtered and evaporated to dryness yielding 5 mg of **318**.

MS: 447.2 (MH^+)


Preparation of **319**:

To 40 mg of **318** in 2 mL of dichloromethane and 2mL of aqueous potassium carbonate was added five equivalents of acetyl chloride and the


mixture stirred under nitrogen. After three hours the phases were separated and the aqueous phase extracted twice with dichloromethane. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. The crude product was purified by reversed phase HPLC yielding 12mg of **319**.

MS: 489.3 (MH^+)


Compounds **320-326** were similarly prepared.




320 MS: 503.3 (MH^+)




321 MS: 525.3 (MH^+)




322 MS: 539.3 (MH^+)




323 MS: 519.3 (MH^+)



324 MS: 504.3 (MH^+)




325 MS: 518.3 (MH^+)



326 MS: 461.3 (MH^+)


Scheme 142:

Using procedures similar to those described under scheme 73 and scheme 109, compound **124** was converted to compounds **327-334**.




Scheme 143:

Compounds **335** to **344** were prepared using transformations similar those described under scheme 73 and scheme 109.




Scheme 144:

Compounds **345** to **353** were prepared using transformations similar those described under scheme 39, scheme 73 and scheme 109.




Similarly, the following compounds were prepared:




354

MS: 517.3 (MH^+)




355

MS: 519.3 (MH^+)




356

MS: 417.2 (MH^+)




357

MS: 419.2 (MH^+)



358


MS: 405.2 (MH^+)



359

MS: 449.2 (MH^+)

Scheme 145:




357 360

To 30 mg of **357** in 1 mL of toluene was added 50μL of water, 1.3 equivalents of cyclopropane boronic acid, 10% mole of tricyclohexyl phosphine and 5% mole of palladium acetate. After bubbling with argon for 2 minutes the mixture was heated to 100°C in a pressure tube. After heating overnight an additional 10% mole of tricyclohexyl phosphine and 5% mole of palladium acetate were added and the temperature was raised to 120°C. After a further 16 hours the reaction mixture was poured onto water and extracted three times with ethyl acetate. The combined organic phases were washed with brine, dried with $MgSO_4$, filtered and evaporated to dryness. Purification by flash chromatography (0-45% ethyl acetate in hexane) yielded 22 mg of **360**.


MS: 425.2 (MH^+)

Scheme 146:

Compounds **361** to **370** were prepared using transformations similar those described under general scheme A, scheme 29, scheme 39, scheme 73 and scheme 109.



Scheme 147:




A solution of **114** (2.6 g, 6.39 mmol) in 50 ml of 1:1 v/v dichloromethane-methanol solution at -78°C was bubbled with ozone until the blue color persisted. To excess ozone was bubbled off with nitrogen and 3 ml of dimethyl sulfide was added and stirred for 20 min at -78°C and 30 min. at 0°C. To this mixture was added 3.5 g of K_2CO_3 and stirred for 1.5 hr at rt. The mixture was diluted with water, extracted 3x with ethyl acetate, the combined organic layers washed with brine, dried over $MgSO_4$, filtered and evaporated to give the crude aldehyde. This was dissolved in 30 ml of 2:1 v/v methanol-dichloromethane, cooled 0°C and 240 mg of $NaBH_4$ was added. The mixture was stirred for 10 min then quenched with aq. NH_4Cl. The slurry was extracted 3x with ethyl acetate, combined organic layers washed with brine, dried over $MgSO_4$, filtered and evaporated to give the crude product. The crude product was stored as an ether solution in the refrigerator and the precipitated solids were filtered off to give 1.07g of **371**. The filtrate was concentrated and chromatographed with 0% to 5% methanol in dichloromethane to provide 0.65g of **372**.

MS for **371**: 440.2 (MH^+)

MS for **372**: 440.2 (MH^+)

Scheme 148:

Preparation of **380**




380

Compound **212** (860 mg, 0.000156 mol) was dissolved in DMF/MeOH (17 mL, 1.58 mL), dppp (65 mg, 10 mol %), $Pd(OAc)_2$ (35 mg, 10 mol%), and Et_3N) 0.463 mL, 2 eq) were added. The mixture was put under an atmosphere of CO and heated at 70°C for 16 hours. The mixture was cooled to rt and $NH_4Cl_{(sat)}$ was added, the resulting mixture was extracted with EtOAc (x3), and the combined extracts were washed with water, dried ($MgSO_4$), and

concentrated. The residue was purified by silica gel chromatography (1:3-3:1 EtOAc/Hexanes) to give 675 mg of the title compound. LCMS: 460.3 (MH^+)

Scheme 149:


Preparation of **381**



Compound **381** was prepared from compound **129** using a procedure similar to the procedure used for the preparation of **380**. LCMS: 446.2 (MH^+)

Scheme 150:


Preparation of **382**



Compound **380**, 675 mg (0.00147 mol) was dissolved in MeOH/THF (5.6mL/5.6 mL), NaOH (3.37 mL of a 1 M solution) was added and the mixture stirred overnight. The organic solvents were removed under reduced pressure and the remaining solution diluted with water, after the mixture was acidified to PH2 the resulting solid was collected by filtration to give 573 mg of the title compound. LCMS: 446.2 (MH^+).

Scheme 151:


Preparation of **383**



Compound **383** was prepared from compound **381** using a procedure similar to the procedure used for the preparation of **382**. LCMS: 446.2 (MH^+)

Scheme 152:


Preparation of **384**












Compound **382** (50 mg, 0.112 mmol) was dissolved in DMF (2 mL), DIPEA (94μL, 5eq) and HATU (110 mg, 2.5 eq) was added. The mixture was stirred for 5 hours, diluted with EtOAc and washed with $NH_4Cl_{(sat)}$. The organic layer was dried over $MgSO_4$, concentrated under reduced pressure and purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 $MeCN/H_2O/HCO_2H$) to give 36 mg of the title compound. LCMS: 445.2 (MH^+).















Scheme 153:













Preparation of **385A-385AAA**
















The following compounds were synthesized from compound **383** and the appropriate amine/amine hydrochloride using a procedure similar to the procedure used for the preparation of **384**.

Compound No	R^1  R^2	LCMS MH^+
385A		485.3
385B		533.3
385C		533.3
385D		515.3
385E		499.3
385F		501.3
385G		501.3
385H		503.3
385I		513.3
385J		515.3

385K		515.3
385L		517.3
385M		517.3
385N		525.3
385O		527.3
385P		528.3
385Q		529.3
385R		535.3
385S		542.3
385T		542.3
385U		542.3
385V		542.3
385W		544.3
385X		547.3

385Y		549.3
385Z		549.3
385AA		557.3
385BB		514.3
385CC		528.3
385DD		542.3
385EE		550.3
385FF		499.3
385GG		513.3
385HH		527.3
385II		527.3
385JJ		528.3
385KK		529.3


385LL		550.3
385MM		553.3
385NN		459.3
385OO		473.3
385PP		487.3
385QQ		487.3
385RR		487.3
385SS		489.3
385TT		501.3
385UU		503.3
385VV		527.3
385WW		543.3
385XX		549.3
385YY		475.3
385ZZ		515.3







385AAA		431.2

Scheme 154:

Compounds **386A-386I**


The following compounds were synthesized from compound **382** and the appropriate amine/amine hydrochloride using a procedure similar to the procedure used for the preparation of **384**.



Compound No		LCMS MH+
386A		529.3
386B		529.3
386C		503.3
386D		473.3
386E		489.3
386F		485.3
386G		459.3
386H		503.3
386I		503.3

Scheme 155:

Preparation of **387**






Compound **382** (25 mg, 0.056 mmol) was dissolved in DMF (1 mL), DIPEA (23 µL, 2.5 eq), methoxyacetic acid hydrazide (14.5 mg, 2.5 eq) then HATU (53 mg, 2.5 eq) were added. The mixture was stirred for 3 hours and then diluted with EtOAc. The mixture was washed with $NH_4Cl_{(sat)}$, dried, concentrated, and dissolved in THF (3 mL). 2-tert-Butylimino-2-diethylamino-1,3-dimethyl-perhydro-1,3,2-diaza-phosphorine on polystyrene (PS-BEMP) (140 mg, 5 eq, 2.2 mmol base/g) and TsCl (13.5 mg, 1.2 eq) were added, the resulting mixture was heated at 120°C for 15 minutes via microwave. Once the mixture was cooled to room temperature the mixture was filtered, concentrated, and purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 MeCN/H_2O/HCO_2H) to give 17 mg of the title compound. LCMS: 514.3 (MH^+).

Scheme 156:

Preparation of **388A-388D**

The following compounds were synthesized from compound **382** using a procedure similar to the procedure used for the preparation of **387** substituting the appropriate carboxylic acid hydrazide.




Compound No	R	LCMS MH+
388A		484.3
388B		512.3
388C		526.3
388D		512.3
388E		528.3
388F		498.3
388G		510.3
388H		528.3

Scheme 157:

Preparation of **389A-389B**

The following compounds were synthesized from compound **383** using a procedure similar to the procedure used for the preparation of **387** using the appropriate carboxylic acid hydrazide.




Compound No	R	LCMS MH+
389A		470.3

389B		500.3

Scheme 158:


Preparation of **390**



Compound **387** (55 mg, 0.107 mmol) was dissolved in CH_2Cl_2 and cooled in an ice-water bath. BBr_3 (45 μL, 4.5 eq) was added and the mixture stirred for 4 hours. Water was added slowly and the resulting mixture stirred for 15 minutes. The mixture was extracted with EtOAc, washed with brine, concentrated, and purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 MeCN/H_2O/HCO_2H) to give 25 mg of the title compound. LCMS: 500.3 (MH^+).

Scheme 159:


Preparation of **391**



Compound **391** was synthesized from compound **389B** using a procedure similar to the procedure used for the preparation of **390**. LCMS: 486.3 (MH^+).

Scheme 160:

Preparation of **392**




Step 1

Compound **382** (50 mg, 0.112 mmol) was dissolved in DMF (1 mL), DIPEA (55 μL, 3 eq), aminoethanal dimethyl acetal (37 μg, 3 eq) then HATU (128 mg, 3 eq) were added. The mixture was stirred for 4 hours and then diluted with EtOAc. The mixture was washed with $NH_4Cl_{(sat)}$, dried ($MgSO_4$), concentrated, and dissolved in THF/1M HCl (3mL/3mL). The mixture was stirred for 4 hours then diluted with EtOAc. The mixture was washed with $NaHCO_{3(sat)}$ dried ($MgSO_4$), and concentrated to give 50 mg of product.

Step 2


The product of step 1 (50 mg, 0.0113 mmol) was dissolved in THF (3 mL), Burgess reagent (54 mg, 2 eq) was added and the mixture heated in the microwave for 15 minutes at 120°C. The mixture was concentrated and purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 $MeCN/H_2O/HCO_2H$) to give 5 mg of the title compound. LCMS: 469.3 (MH^+).



Compound **393** was synthesized from compound **383** using a procedure similar to the procedure used for the preparation of **392**. LCMS: 455.3 (MH^+).

Scheme 161:

Preparation of **394**



Step1


Compound **386H** (80 mg, 0.165 mmol) was dissolved in CH_2Cl_2, Dess-Martin Periodinate (88 mg) was added and the mixture stirred for 1 hour. EtOAc was added followed by $NaHCO_{3(sat)}$ and sodium thiosulphate$_{(sat)}$, the mixture was stirred for 1 hour and the organic layer collected, dried ($MgSO_4$), and concentrated to give 74 mg of product.

Step 2

The product of step 1 (74 mg, 0.0147 mmol) was dissolved in THF (3 mL), Burgess reagent (65 mg, 2 eq) was added and the mixture heated in the microwave for 15 minutes at 120°C. The mixture was concentrated and purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 MeCN/H_2O/HCO_2H) to give 7 mg of the title compound. LCMS: 484.3 (MH^+).

Scheme 162:

Preparation of **395**




Compound **383** (55 mg, 0.127 mmol) was dissolved in DMF (2.5 mL), DIPEA (55 μL, 2.5 eq), acetamide oxime (24 mg, 2.5 eq) and HATU (121 mg, 2.5 eq) were added. The mixture was stirred at rt for 3 hours and then heated at 191°C via microwave for 4 minutes. After cooling to rt the mixture was diluted

with EtOAc, washed with $NH_4Cl_{(sat)}$, dried ($MgSO_4$), and concentrated. The residue was purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 $MeCN/H_2O/HCO_2H$) to give 17 mg of the title compound. LCMS: 470.3 (MH^+).

Scheme 163:


Preparation of **396**



Compound **396** was synthesized from compound **382** using a procedure similar to the procedure used for the preparation of **395**. LCMS: 484.3 (MH^+).

Scheme 164:


Preparation of **397**



Compound **397** was synthesized from compound **382** using a procedure similar to the procedure used for the preparation of **395** substituting *N*-hydroxy-2-methoxyacetamidine in place of acetamide oxime. LCMS: 514.3 (MH^+).

Scheme 165:


Preparation of **398**



Compound **398** was synthesized from compound **397** using a procedure similar to the procedure used for the preparation of **390**. LCMS: 500.3 (MH^+).

Scheme 166:


Preparation of **399**



Compound **384** (55 mg, 0.123 mmol) was dissolved in *N,N*-dimethylformamide dimethyl acetal (2 mL) and heated at 120°C for 1.5 hours. The excess reagent was removed under reduced pressure and the residue treated with hydrazine hydrate (7 μL, 1.1 eq) in acetic acid (1 mL) at 90°C for 1.5 hours. The volatiles were removed under reduced pressure and the residue treated with $NaHCO_{3(sat)}$, the mixture was extracted with EtOAc, dried ($MgSO_4$), concentrated, and purified by reverse phase HPLC, C18 (10:90:0.5 to 90:10:0.5 $MeCN/H_2O/HCO_2H$) to give 17 mg of the title compound. LCMS: 469.3 (MH^+).

Scheme 167:


Preparation of **400**



Compound **400** was synthesized from compound **384** using a procedure similar to the procedure used for the preparation of **399** by substituting methyl hydrazine for hydrazine hydrate. LCMS: 483.3 (MH^+).

Scheme 168:

Preparation of **401**




Compound **401** and was synthesized in an analogous manner to compound **153** substituting 3-(4-chlorophenyl)-1(R)-methylallylamine for 3-(4-chlorophenyl)allyl amine. LCMS: 422.2 (MH^+).

Scheme 169:

Preparation of **402** and **403**

Step 1



3-Chloro-4-bromoanisole (10 g, 0.045 mol) was dissolved in THF (200 mL) the mixture was cooled to -78°C. n-BuLi (18 mL of a 2.5 M solution in hexanes , 1 eq) was added over 5 minutes. After the resulting mixture was stirred at -78°C for 30 minutes DMF (5.24 mL, 1.5 eq) was added. The mixture stirred for an additional 30 minutes then allowed to warm to rt. $NH_4Cl_{(sat)}$ was added and the mixture extracted with Ether. The combined


extracts were washed with water, dried ($MgSO_4$), and concentrated to give the product of step1.

Step 2

The product of step 1 was converted to compounds **402** and **403** using analogous procedures to those used for the synthesis of compounds **153** and **154** substituting 3-(4-chlorophenyl)-1(R)-methylallylamine for 3-(4-chlorophenyl)allyl amine. LCMS: 87a, 418.2 (MH^+), 88a, 418.2 (MH^+).

Scheme 170:


Preparation of **404**



Compound **404** was prepared from compound **402** using a procedure similar to the procedure used for the preparation of **72**. LCMS: 404.2 (MH^+)

Scheme 171:


Preparation of **405**



Compound **405** was prepared from compound **403** using a procedure similar to the procedure used for the preparation of **72**. LCMS: 404.2 (MH^+)

Scheme 172:


Preparation of **406**



Compound **406** was prepared from compound **404** in two steps using the procedures described for compound **129** and compound **380**. LCMS: 446.2 (MH^+)

Scheme 173:


Preparation of **407**



Compound **407** was prepared from compound **406** using a procedure similar to the procedure used for the preparation of **382**. LCMS: 432.2 (MH^+)

Scheme 174:


Preparation of **408**



Compound **408** was prepared from compound **404** using a procedure similar to the procedure used for the preparation of **230**. LCMS: 476.3 (MH^+)

Scheme 175:


Preparation of **409**



Compound **409** was prepared from compound **408** using procedures similar to the procedure used for the preparation of **231**. LCMS: 462.3 (MH^+)

Scheme 176:


Preparation of **410**



Compound **410** was prepared from compound **404** using procedures similar to the procedures used for the preparation of **221**. LCMS: 448.2 (MH^+)

Scheme 177:


Preparation of **411**



Compound **411** was prepared from compound **404** using a procedure similar to the procedure used for the preparation of **131**. LCMS: 461.3 (MH^+)


Scheme 178:

Preparation of **412**




Compound **412** and was synthesized in an analogous manner to **137** substituting 3-(4-chlorophenyl)-1(R)-methylallylamine for 3-(4-chlorophenyl)allyl amine. LCMS: 424.2 (MH^+).

Scheme 179:




Step 1:



A mixture of (*S*)-2-methyl-2-propyl sulfinamide (19.4 g, 0.16 mol), benzyloxyacetaldehyde (20 g, 0.133 mol), and anhydrous $CuSO_4$ (12.8 g, 80 mmol) in DCM (170 mL) was stirred at room temperature under N_2 for 6 h. Another 30 g of anhydrous $CuSO_4$ (12.8 g, 80 mmol) was added and the mixture was stirred at room temperature for 16 h. The mixture was filtered through CELITE. The filtrate was concentrated *in vacuo*. The residue was chromatographed on a silica gel cartridge with EtOAc in hexane (0→50%) to afford **420** (27.5 g, 82%).

Racemic **420** was obtained similarly from racemic 2-methyl-2-propyl sulfonamide. From racemic **420**, racemic final products such as racemic **428** were prepared.

Step 2:




A solution of EtMgBr (31.5 mL, 3.0 M in ether, 0.102 mol) was added to a solution of 4-chlorophenylacetylene (15 g, 0.11 mol) in anhydrous ether (370 mL) under argon at 50°C and refluxed for 1 h. The resulting Grignard solution was added to a solution of **420** (13 g, 0.051 mol) in DCM (150 mL) at -78 °C under argon. The mixture was stirred at -78 °C for 5 h and at room temperature for 16 h. The reaction was quenched with NH_4Cl (sat.) and extracted with EtOAc. The organic layer was dried ($MgSO_4$) and concentrated in vacuo. The crude product (24 g) was used for the next reaction.

A solution of HCl in dioxane (93 mL, 4.0 M, 0.372 mol) was added to a solution of the crude product (24 g) from the above reaction in MeOH at 0 °C

and stirred for 1.5 h. The mixture was concentrated *in vacuo*. The residue was dissolved with EtOAc and washed with $NaHCO_3$ (sat.). The organic layer was dried ($MgSO_4$) and concentrated in vacuo. The residue was chromatographed on a silica gel cartridge with EtOAc in hexane (0→70%) to afford a mixture of **422a** and **422b** (7.3 g, 50%). The ratio of **422a** and **422b** is around 4 to 1 based on final product (e.g., **434**) ratios.


Racemic **422** was obtained similarly from racemic **420**. From racemic **422**, racemic final products such as racemic **428** were prepared.

Step 3:




The alkyne **422** was reduced to give the cis-olefin **423** in a similar way as shown in Scheme 4.

Steps 4 and 5:




The cis-olefin **423** was converted to the Diels-Alder precursor **424** and then to the Diels-Alder product **425** in a similar way as shown in Scheme 73.

Step 6:



The Diels-Alder product **425** was ethylated to give **426** in a similar way as shown in Scheme 52.

Step 7:



The intermediate **426** was converted to compound **427** in a similar way as shown in Scheme 77. Neat BBr_3 was used instead of 1 M solution.


LCMS for **427**: 450.2 (MH^+)

Compound **427** was separated to give single enantiomers **427a** and **427b** in a similar way as shown in Scheme 71. Chiralpac OD column was used for the chiral HPLC separation.

LCMS for **427a**: 450.2 (MH^+)

LCMS for **427b**: 450.2 (MH^+)

Step 8:




Compound **427** was converted to single enantiomers **428a** and **428b** and in a similar way as shown in Schemes 75 and 71. Chiralpac OD column was used for the chiral HPLC separation.

LCMS for **428a**: 452.2 (MH^+)


LCMS for **428b**: 452.2 (MH^+)

Step 9:




A solution of **427** (30 mg, 0.066 mmol) in anhydrous THF (1.3 mL) was added to NaH (15 mg, 0.33 mmol) at room temperature and stirred for 0.5 h. Bromoethyl methyl ether (0.20 mL, 0.22 mmol) was added and stirred for 16 h. The reaction was quenched with water and extracted with DCM. The organic layer was dried ($MgSO_4$) and concentrated in vacuo. The residue was chromatographed on a silica gel cartridge with EtOAc in DCM (0→60%) to afford racemic **431** (26 mg, 76%).
LCMS: 508.3 (MH^+)

The following compounds were similarly prepared from **427** using appropriate alkylating agents:




Compound No	R	LCMS
429A	Me	464.3
429B	Et	478.3
429C	n-Pr	492.3

The following compounds were prepared from appropriate precursors (e.g. **430A** from **429A**) in a similar way as shown in Schemes 75.



Compound No	R	LCMS
430A	Me	466.3
430B	Et	480.3
430C	$-CH_2CH_2OMe$	510.3


The Diels-Alder product **425** was converted to give **432** in a similar way as shown in Scheme 52:



LCMS for **432**: 512.3 (MH^+)

Scheme 180:

Synthesis of **433** and **434**




The cis-olefin **16** was converted to racemic **433** in a similar way as shown for the synthesis of **428** above. Racemic **433** was converted to single enantiomer **434** in a similar way as shown for the synthesis of **221** given in scheme 109 followed by chiral HPLC separation as described in scheme 71. Chiralpac OD column was used for the chiral HPLC separation.

LCMS for **433**: 434.2 (MH^+)

LCMS for **434**: 478.3 (MH^+)

Scheme 181:

Synthesis of **436-438:**



Compound **433** was converted to **436**, **437** and **438** in a similar way as shown in Scheme 39.


LCMS for **436**: 485.3 (MH^+)

LCMS for **437**: 484.3 (MH^+)

LCMS for **438**: 525.3 (MH^+)

Scheme 182:

Synthesis of **440**:




The commercially available 2-chloro-4-fluoro-benzaldehydewas converted to **440** in a similar way as shown in Schemes 1, 2, and 64.


LCMS: 420.2 (MH^+)

Scheme 183:

Synthesis of **443-446**




Step 1:



The commercially available 2,4-dimethoxy-benzaldehyde was converted to intermediate **443** in a similar way as shown in Schemes 1, 2, 179.


Step 2:



A mixture of intermediate **443** (241 mg, 0.624 mmol), triisopropylsilyl triflate (0.42 mL, 1.6 mmol), and pyridine (0.25 mL, 3.1 mmol) in DCM (6 mL) was stirred at room temperature for 1.5 h. The solution was washed with

NH_4Cl (sat.). The organic layer was dried ($MgSO_4$) and concentrated in vacuo. The residue was chromatographed on a silica gel cartridge with MeOH in DCM (0→15%) to afford intermediate **444** (294 mg, 87%).


Step 3:



The intermediate **444** was converted to **445** in a similar way as shown in Scheme 39. During the conversion of the triflate to the nitrile group, the triisopropyl group was cleaved to give the phenol.

445, LCMS: 409.2 (MH^+)

Step 4:




Compound **445** was converted to **446** in a similar way as shown for the synthesis of **221** (see scheme 109)

LCMS: 453.2 (MH^+)


Scheme 184:

Preparation of compounds **450-452**:

Using a procedure similar to the conversion of **73** to **78** (see scheme 46) compounds **450-452** were prepared from intermediates **212** and **341** and the appropriate boronic acid or boronate ester.



Scheme 185:




5 Preparation of **453**:

A mixture of **211** (160 mg, 0.382 mmol), Ethyl O-(p-toluenesulfonyl)-L-(-)-lactate (520 mg, 1.91 mmol, 5 eq) (Ref: *Tetrahedron*, **1985**, *vol.41*, page 541-546) and K_2CO_3 (160 mg, 1.16 mmol, 3 eq.) in 3 ml DMF was heated overnight in a sealed tube at 100°C. The mixture was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 5% methanol in dichloromethane to provide 204 mg of **453**.
MS: 518.3 (MH^+)

Preparation of **454**:
To a solution of **453** (100 mg, 0.21 mmol) in 2 ml THF at rt was added 2M solution of $LiBH_4$ in THF (0.5 ml, 1 mmol) and the mixture was stirred at rt for 1 hr. It was poured in to aq. NH_4Cl, extracted 3x with ethyl acetate, the combined organic layers washed with brine, dried over $MgSO_4$, filtered, concentrated and chromatographed with 0% to 5% methanol in dichloromethane to provide 84 mg of **454**.
MS: 476.3 (MH^+)

Preparation of **455**:
A mixture of **211** (40 mg, 0.095 mmol), and ~7 equivalents of methane sulfonyl chloride in 1ml pyridine was stirred overnight. It was diluted with ethyl acetate, washed 3x with 1N HCl, brine, dried over $MgSO_4$, filtered, concentrated and purified by preparative TLC using 4% methanol in dichloromethane as eluent to provide 16 mg of **455**.
MS: 496.3 (MH^+)


Preparation of **461**



461

Compound **461** was prepared from **388H** using the procedure used for compound **390**. LCMS: 514.3 (MH^+).

Preparation of **462**




462

Compound **462** was prepared from compound **384** in a similar manner to compound **399** using dimethylacetamide dimethylacetal in place of dimethylformamide dimethylacetal. LCMS: 483.3 (MH^+).

Scheme 186:

Preparation of **463**




463

Compound **212** (110 mg, 0.2 mmol) was dissolved in PhMe/EtOH/H_2O (1.1 mL/1.1 mL/0.45 mL). Pd (dppf)Cl_2.CH_2Cl_2 (15 mg, 10 mol %), Na_2CO_3 (64 mg, 3 eq), and 1-Methylpyrazole-4-boronic acid pinacol ester (62 mg, 1.5 eq) were added. The mixture was heated at 120°C in a microwave for 20 minutes. $NH_4Cl_{(sat)}$ was added and the mixture extracted with EtOAc. The extracts were dried, concentrated, and then purified (SiO_2, hexane/EtOAc 2:1-1:2) to give the title compound, 28 mg, LCMS: 482.3 (MH^+).


Scheme 187:

Preparation of **464**




Compound **212** (94 mg, 0.171 mmol) and 1-methyl-2-(tributylstannyl)imidazole (317 mg, 5 eq) was dissolved in THF. $Pd(Ph_3P)_4$ (60 mg, 0.3 eq) was added and the mixture heated at 85°C overnight. The mixture was diluted with EtOAc, washed with $NH_4Cl_{(sat)}$, brine, dried, and concentrated. Purification (SiO_2, hexane/EtOAc 2:1-1:2) gave the title compound, 20 mg, LCMS: 482.3 (MH^+).


The following compounds were prepared in a similar manner to **463** using appropriate boronic acid pinacol esters.



LCMS: 468.3 (MH+).




LCMS: 497.3 (MH+).




LCMS: 496.3 (MH+).


The following compounds were prepared in a similar manner to **464** using appropriate tributylstannanes.



LCMS: 550.3 (MH^+)




LCMS: 485.3 (MH+).



LCMS: 485.3 (MH+).


Preparation of **471**



471

Compound **471** was prepared from 5-(4-methoxyphenyl)-2(E), 4(E)-pentadienoic acid in a similar manner to compound **211**. LCMS 384.2 (MH^+)

The following compounds were prepared from **435** in a similar fashion as shown in Scheme 39 using appropriate organoboron or organotin reagents:



LCMS for **480**: 524.3 (MH^+)

LCMS for **481**: 537.3 (MH^+)

LCMS for **482**: 519.3 (MH^+)

LCMS for **483**: 539.3 (MH^+)

LCMS for **484**: 525.3 (MH^+)

LCMS for **485**: 526.3 (MH^+)


LCMS for **486**: 512.3 (MH^+)

LCMS for **487**: 498.3 (MH^+)

LCMS for **488**: 484.3 (MH^+)

Preparation of 4,5-Di(hetero)aryl-octahydro-benzo[1,2,5]thiadiazole 2,2-dioxides and 4,5-Di(hetero)aryl-octahydro-benzoimidazol-2-ones


General Scheme M:



4,5-Di(hetero)aryl-octahydro-benzo[1,2,5]thiadiazole 2,2-dioxides and 4,5-di(hetero)aryl-octahydro-benzoimidazol-2-ones (represented by **bg** and **bh**) can be prepared as shown in General Scheme M, above. Triflate **bc** generated from ketone **bb** by treatment with LDA (i.e., lithium diisopropylamide) followed by triflic anhydride can undergo Stille coupling with vinyl tributyltinhydride to give the diene **bd**. Diels-Alder reaction of diene **bd** with maleic anhydride can give the anhydride **be**. Compound **be** can undergo Curtius rearrangement (e.g., using the procedure of R.Seito et.al Bull. Chem. Soc. Jpn. *59*, 1689 (**1986**); herein incorporated by reference in its entirety) to form the diamine **bf**. The diamine **bf** can be readily converted to compounds **bg** and **bh** by treatment with sulfurylchloride and carbonyldiimidazole, respectively.


Synthesis of Isolactones and Isolactams **bl** and **bp**

General Scheme N:



solactone **bl** and isolactam **bp** can be prepared according to General Scheme N, above. Cinnamyl alcohol **bi** can undergo coupling with a substituted propargylic acid to give the ester **bj**, which under selective hydrogenation conditions can yield **bk**. Standard Diels-Alder cyclization of **bk** and subsequent reduction of the internal double bond can provide compound **bl**. Similarly, **bi** can be converted to the amine **bm**. Amine **bm** can be converted to Diels-Alder precursor **bo** by reaction with a propargylic acid and subsequent selective reduction. Thermal Diels-Alder reaction of **bo** can give the lactam **bp** after reduction of the double bond.


General Scheme:



The synthesis of an embodiment of the formula I involves the application of Diels-Alder reaction. The Diels-Alder precursor **III**^{a} can be readily prepared by coupling the dienoic acid **I**^{a} and the allyl amine **II**^{a}. The amide can be either protected with an appropriate functional group (R^a is a protecting group) or it can be used without any protection (R^a is hydrogen). When **III**^{a} is subjected to the Diels-Alder reaction it gives the cyclization product **IV**^{a}. The yield and selectivity of the products in this step can depend on a variety of reaction conditions such as the solvent used, the temperature employed for the cyclization, the additives used in the reaction medium such as Lewis acids etc. Following the Diels-Alder reaction an R^{15} group can be introduced and the double bond reduced to provide **V**^{a} which can be subjected to the deprotection conditions to cleave the protecting group to give **VI**^{a}. The substitution of Ar^1 or Ar^2 can be further functionalized or transformed. Non-limiting examples of protecting groups for R^a include all those known to protect a nitrogen and examples may be found in Green et al., (Protective Groups in Organic Synthesis by T.W. Greene and P.G. Wuts; 1999, Third edition, John Wiley & Sons, Inc.) Preferred amide protecting groups include but are not limited to tert-butoxycarbonyl (Boc), benzyl oxycarbonyl (Cbz), para-methoxy benzyl, 3,4-dimethoxybenzyl, allyl, trimethylsilyl ethyl (TMSE), methoxy methyl (MOM), benzyloxymethyl (BOM), methoxy, *tert*-butyldimethylsilyl (TBDMS), Triisopropylsilyl (TIPS), methoxy carbonyl and ethoxycarbonyl, etc. Preferred electrophilic reagents include for example optionally substituted alkyl halides

(e.g., methyl iodide, ethyl iodide, propyl iodide, Br-CH_2CH_2-OTBS), optionally substituted benzyl halides (e.g., benzyl bromide, para-cyano benzyl bromide, ortho-cyano benzyl bromide and meta-cyano benzyl bromide). Electrophilic reagents are electron deficient reagents that can react with another molecule by accepting a pair of electrons to form a new bond. The electrophiles (which are also a Lewis acids), can be positively charged, have an atom which carries a partial positive charge, or have an atom which does not have an octet of electrons [see Page 541 of Mechanism and Theory in Organic Chemistry by T.H. Lowry and K.S. Richardson, Third edition, Harper Collins Publishers].


In another embodiment is a process for preparing compounds of formula V^a



V^a


wherein R^a is H or a protecting group and R^3, R^{15}, Ar^1 and Ar^2 are herein defined above, comprising

a) coupling a compound of formula I^a




I^a

with a compound of formula II^a




II^a

to obtain a compound of formula III^a



IIIa


b) cyclizing the compound of formula III^a to obtain a compound of formula IV^a



IV^a

c) generating the enolate and reacting with an electrophilic reagent to introduce R^{15} group to the compound of formula III^a and

d) reducing the double bond of the compound of formula IV^a to obtain a compound of formula V^a



V^a

In an embodiment of the invention is the process to obtain a compound of formula V^a wherein R^a is H, tert-butoxycarbonyl (Boc), benzyl oxycarbonyl (Cbz), para-methoxy benzyl, 3,4-dimethoxybenzyl, allyl or trimethylsilyl ethyl (TMSE), methoxy methyl (MOM), benzyloxymethyl (BOM), methoxy, *tert*-butyldimethylsilyl (TBDMS), Triisopropylsilyl (TIPS), methoxy carbonyl or ethoxycarbonyl and wherein said electrophilic reagent is alkyl-iodide.

ASSAY

Method for evaluating Cannabinioid CB_1 and CB_2 affinity

Competition binding assays for cannabinoid CB_1 and CB_2 affinity were performed by incubating commercially purchased membranes prepared from cells expressing each receptor subtype (8 µg pro) with 0.5 nM ^{3}H-CP55,940, a non-selective cannabinoid agonist, along with concentrations of drug ranging from 0.0001-3 µM in Buffer A (5 mM $MgCl_2$, 2.5 mM EDTA and 013% BSA). Non-specific binding was defined in the presence of 10 µM CP55,940. For saturation studies, concentrations of ^{3}H-CP55,940 ranging from 0.1-5 nM were incubated with membranes in the presence and absence of 10 µM CP55,940. Assays were terminated after incubation for 1 ½ hours by rapid filtration onto 0.3 % polyethylenamine treated GF/C filterplates using a BRANDEL cell harvester. The plates were dried and MICROSCINT

scintillation cocktail was added, after which the bound radioactivity was quantified using a TOPCOUNT scintillation counter.

The dissociation constant (K_d) of ^{3}H-CP55,940 at the CB_1 and CB_2 receptor were determined by plotting specific binding at each concentration of radioligand, and analysis by non-linear regression. For competition studies, the concentration of each drug that inhibited 50 percent of ^{3}H-CP55,940 binding (IC_{50}) was determined by non-linear regression analysis of the radioligand displacement curves. Affinity constants (K_i) were calculated using the equation derived by Cheng and Prusoff (1973), defined as: IC_{50}/1+[conc. ligand / K_d].

GTPγS Binding Protocol

The functional efficacy of compounds to activate second messengers within the cell was determined utilizing the GTPγS binding assay. Guanine nucleotides are phosphorylated within the plasma membrane of the cell following binding and activation by agonists. A radiolabelled derivative of guanine triphosphate (GTP) is utilized in this assay as it cannot be dephosphorylated and therefore accumulates following agonist binding. The simultaneous presence of an antagonist into this system will shift the agonist concentration curve to the right, with increasing concentrations of antagonist producing a greater rightward shift in the dose-response curve of the agonist.


Commercially purchased membranes were incubated with 10 mM GDP to allow sufficient substrate for phosphorylation in the presence of agonist. The membranes were then pre-incubated with increasing concentrations of test compound for 30 minutes to determine if they were capable of stimulating phosphorylation alone. Increasing concentrations of the non-selective cannabinoid agonist WIN55,122 were then added in the presence or absence of each concentration of test compound. The assay was then incubated for 1 hour at room temperature. To complete the assay, ^{35}S-GTPγS was added and the assay incubated for another 30 minutes. Assays were terminated by rapid filtration onto 10 mM sodium phosphate-treated GF/C filterplates using a BRANDEL cell harvester. The plates were dried and Microscint scintillation cocktail was added, after which the bound radioactivity was quantified using a TOPCOUNT scintillation counter.

The stimulation of ^{35}S-GTPγS binding as a function of the concentration of the agonist WIN55,122, in the absence and presence of test compound, was plotted and the EC_{50} determined by nonlinear regression analysis using GraphPad Prism software. A Schild analysis of the rightward shift in the dose response curve of WIN55,122 in the presence of test compound was determined by plotting the concentration of test compound against the negative log of the dose ratio [1-(EC_{50} agonist + test compound/EC50 of agonist alone)]. A linear regression analysis yields the Kb, defined as the X-intercept of the linear equation.

Preferred compounds of Formula I of the present invention, and salts, solvates, or esters thereof, have K_i values of about 200 nM or less. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 100 nM or less. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 50 nM or less. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 20 nM or less. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 10 nM or less. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 5 nM or less. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 10 to about 1 nM. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 10 to about 0.1 nM. In another embodiment, the compounds of Formula (I) of the present invention, and salts, solvates, or esters thereof, have K_i values of about 10 to about 0.01 nM. Examples 40, 42, 169, 170, 174, 178, 180, 181, 183, 182, 185, 213, 221, 227, 228, 260, 282, 387, 397, 438, 451, 454, 463 and 483 have K_i values in the range of about 10 to about 1 nM.

WE CLAIM:

1. A compound of Formula (I):




(I)

or a pharmaceutically acceptable salt, solvate, or ester thereof, wherein:


m is 0 or 1, n is 1 or 2, and m + n is 1 or 2;

R^1 and R^2 together with the carbon atoms to which they are shown attached in Formula (I) form a group Q as shown in Formula (IA):



(IA)

wherein Q is selected from the group consisting of:



(a)

Y^1 is –O- or –N(R^7)-;

R^3, and R^4, are each independently selected from the group consisting of H, -O-R^9, R^{11}, and –N(R^{16})$_2$;

R^7 is selected from the group consisting of H, alkyl, arylalkyl, alkenyl, -alkylene-N(R^9)$_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, -alkylene-C(O)H, -C(O)-O-R^{11}, and Boc;

R^9 is selected from the group consisting of H, TBS, TIPS, Tf and R^{11};

R^{11} is selected from the group consisting of unsubstituted alkyl, alkyl substituted with one or more U groups, -alkylene-O-alkyl, -alkylene-O-aryl, unsubstituted aryl, and aryl substituted with one or more X^1 groups;

R^{12} is selected from the group consisting of unsubstituted aryl and aryl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more X^2 groups;

R^{14} is selected from the group consisting of unsubstituted cycloalkyl, cycloalkyl substituted with one or more X^4 groups unsubstituted alkyl, and alkyl substituted with one or more U groups;

each R^{15} is independently selected from the group consisting of H, $-N_3$, halogen, alkenyl, -alkylene-R^{12}, -alkylene-O-R^9, -alkylene-$N(R^{18})_2$, -alkylene-C(O)H, -OH, -CN, -O-alkyl, $-C(O)N(R^{18})_2$, $-N(R^{18})_2$, $-NR^{18}C(O)R^{18}$, $-NR^{18}C(O)_2R^{18}$, $-NR^{18}C(O)N(R^{18})_2$, $-NR^{18}S(O)_2R^{18}$, –O-alkenyl, $-C(O)_2R^{18}$; unsubstituted alkyl, alkyl substituted with one or more U groups, -O-alkylene-$C(O)R^{18}$,or $-C(O)R^{18}$;

with the proviso wherein the group $-N(R^{18})_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

is selected from the group consisting of R^9 and $–C(O)-R^{12}$;

each R^{18} is independently selected from the group consisting of H, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, R^{12}, R^{13} and R^{14};

with the proviso that when R^{18} is attached to N, then each R^{18} is independently selected from the group consisting of H, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, $-C(O)R^{21}$, R^{12}, R^{13} and R^{14};

R^{19} is selected from the group consisting of H, TBS, TIPS, Tf and R^{21};

each R^{20} is independently selected from the group consisting of H, unsubstituted alkyl, alkyl substituted with one or more U groups, -alkylene-R^{22}, -alkylene-R^{23}, -alkylene-R^{24}, -C(O)-R^{24}, -alkylene-O-R^{19}, R^{24}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, and benzo-fused cycloalkyl;

R^{21} is selected from the group consisting of unsubstituted alkyl, alkyl substituted with one or more U groups, -alkylene-O-alkyl,

-alkylene-O-aryl, unsubstituted aryl, aryl substituted with one or more W^1 groups; unsubstituted heteroaryl, heteroaryl substituted with one or more W^2 groups, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, -O-alkylene-O-R^{24}, -C(O)-O-alkylene-O-R^{24}; -C(O)-alkylene-R^{23}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -$S(O)_2$-R^{24}, and -alkylene-O-alkylene-O-R^{24}, with the proviso that –O-O- cannot be formed with R^{21} and the atom said R^{21} is attached to;

R^{22} is selected from the group consisting of unsubstituted aryl and aryl substituted with one or more W^1 groups;

R^{23} is selected from the group consisting of unsubstituted heteroaryl and heteroaryl substituted with one or more W^2 groups;

R^{24} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, unsubstituted alkyl, and alkyl substituted with one or more U groups;

each R^{25} is independently selected from the group consisting of H, R^{22}, R^{23}, unsubstituted alkyl, alkyl substituted with one or more U groups, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -alkylene-OR^{19}, -alkylene-$NR^{19}R^{19}$, -alkylene-SR^{19},-alkylene-R^{23}, -alkylene-R^{22}, unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more W^3 groups, -alkylene-heterocycloalkyl, -alkylene-heterocycloalkyl substituted with one or more W^3 groups, -C(O)-R^{24}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -$S(O)_2$-R^{24}, -C(O)-NH-R^{22} and -C(O)-NH-R^{24};

with the proviso wherein the group -$N(R^{25})_2$, both R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

each W^1 is independently selected from the group consisting of halogen, -CN, -OH, -O-$S(O)_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, and –O-alkyl;

each W^2 is independently selected from the group consisting of halogen, unsubstituted aryl, and aryl substituted with one or more Z groups;

each W^3 is independently selected from the group consisting of –OH, alkyl, -alkylene-OH, -O-alkyl, -C(O)-alkyl, -C(O)NH_2, -NHC(O)alkyl, -NHC(O)H, -NHC(O)-O-alkyl and -C(O)-O-alkyl; or

two W^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each W^4 is independently halogen or alkyl;

Ar^1 is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

Ar^2 is selected from the group consisting of unsubstituted phenyl, phenyl substituted with one or more X^1 groups, pyridyl-2-yl and pyridyl-2-yl substituted with one or more X^2 groups;

each X^1 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-C(O)N(R^{20})$_2$, -O-alkylene-O-R^{19}, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl, -O-alkylene-O-alkylene-OR^{19}, -O-alkylene-C(O)R^{24}, -O-alkylene-C(O)OR^{19},–O-alkyl, -N(R^{25})$_2$, -C(O)alkyl, –C(O)OH, -C(O)O-alkyl, -C(O)O-cycloalkyl, -C(O)N(R^{25})$_2$, -O-alkylene-heterocycloalkyl, -O-alkylene-heterocycloalkyl substituted with one or more W^3 groups, unsubstituted heterocycloalkyl, -heterocycloalkyl substituted with one or more W^3 groups, -O-alkenylene-O-alkylene-O-R^{24}, -O-alkylene-N(R^{25})$_2$, -O-alkylene-C(O)N(R^{25})$_2$, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S(O)$_2$-R^{24}, and alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

each X^2 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z

groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-$C(O)N(R^{20})_2$, -O-alkylene-O-R^{19}, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl,-O-alkylene-O-alkylene-OR^{19}, -O-alkylene-$C(O)R^{24}$, -O-alkylene-$C(O)OR^{19}$,–O-alkyl, $-N(R^{25})_2$, -C(O)alkyl, –C(O)OH, -C(O)O-alkyl, -C(O)O-cycloalkyl, $-C(O)N(R^{25})_2$, -O-alkylene-heterocycloalkyl, -O-alkylene-heterocycloalkyl substituted with one or more W^3 groups, unsubstituted heterocycloalkyl, -heterocycloalkyl substituted with one or more W^3 groups, -O-alkenylene-O-alkylene-O-R^{24}, -O-alkylene-$N(R^{25})_2$, -O-alkylene-$C(O)N(R^{25})_2$, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, $-S(O)_2$-R^{24}, and alkenyl;

with the proviso wherein the group $-N(R^{20})_2$ or $-N(R^{25})_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl;

each X^3 is independently selected from the group consisting of –OH, alkyl, -alkylene-OH, -O-alkyl, -C(O)-alkyl, $-C(O)NH_2$, -NHC(O)alkyl, -NHC(O)H, -NHC(O)-O-alkyl and -C(O)-O-alkyl; or

two X^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each X^4 is independently halogen or alkyl;

each U is independently selected from the group consisting of –OH, -O-alkyl, -O-aryl, -O-alkylene-aryl, -O-alkylene-O-alkyl, -O-alkylene-O-haloalkyl, -O-alkylene-O-aryl, halogen, -CN, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, OTBS, OTIPS and OTf;

and

each Z is independently selected from the group consisting of –OH; -O-alkyl; halogen; alkyl; –CN; $-CF_3$; cycloalkyl; -alkylene-OH; -alkylene-O-alkyl; -alkylene-O-alkyl substituted with one or more groups selected from the group consisting of –OH, -O-alkyl, halogen, -CN, cycloalkyl, heterocycloalkyl, aryl, heteroaryl; -alkylene-O-alkylene-O-alkyl; -alkylene-O-alkylene-O-aryl;


-alkylene-O-aryl; and -alkylene-O-aryl substituted with one or more groups selected from the group consisting of halogen, -CN, -OH, -O-S(O)$_2$-haloalkyl, aryl, heteroaryl and –O-alkyl; or

two Z groups together with the ring carbon atom to which they are attached form a carbonyl group.

2. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:


m is 0 or 1, n is 1 or 2, and m + n is 1 or 2;

R^1 and R^2 together with the carbon atoms to which they are shown attached in Formula (I) form a group Q as shown in Formula (IA):



(IA)

wherein Q is selected from the group consisting of:



(a)

Y^1 is –O- or –N(R^7)-;

R^3, and R^4, are each independently selected from the group consisting of H, -O-R^9, R^{11}, and –N(R^{16})$_2$;

R^7 is selected from the group consisting of H, (C_1-C_6)alkyl, (C_6-C_{12})aryl(C_1-C_6)alkyl, (C_2-C_6)alkenyl, -(C_1-C_6)alkylene-N(R^9)$_2$, -(C_1-C_6)alkylene-O-R^9, -(C_1-C_6)alkylene-R^{12}, -C(O)-R^{14}, -(C_1-C_6)alkylene-C(O)H, -C(O)-O-R^{11}, and Boc;

R^9 is selected from the group consisting of H, TBS, TIPS, Tf and R^{11};

R^{11} is selected from the group consisting of unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, -(C_1-C_6)alkylene-O-(C_1-C_6)alkyl, -(C_1-C_6)alkylene-O-(C_6-C_{12})aryl, unsubstituted (C_6-C_{12})aryl, and (C_6-C_{12})aryl substituted with one or more X^1 groups;

R^{12} is selected from the group consisting of unsubstituted (C_6-C_{12})aryl and (C_6-C_{12})aryl substituted with one or more X^1 groups;

R^{13} is selected from the group consisting of unsubstituted $(C_2\text{-}C_{10})$heteroaryl and $(C_2\text{-}C_{10})$heteroaryl substituted with one or more X^2 groups;

R^{14} is selected from the group consisting of unsubstituted $(C_3\text{-}C_7)$cycloalkyl, $(C_3\text{-}C_7)$cycloalkyl substituted with one or more X^4 groups unsubstituted $(C_1\text{-}C_6)$alkyl and $(C_1\text{-}C_6)$alkyl substituted with one or more U groups;

each R^{15} is independently selected from the group consisting of H, $\text{-}N_3$, halogen, $(C_2\text{-}C_6)$alkenyl, $\text{-}(C_1\text{-}C_6)$alkylene-R^{12}, $\text{-}(C_1\text{-}C_6)$alkylene-O-R^9, $\text{-}(C_1\text{-}C_6)$alkylene-$N(R^{18})_2$, $\text{-}(C_1\text{-}C_6)$alkylene-C(O)H, -OH, -CN, $\text{-O-}(C_1\text{-}C_6)$alkyl, $\text{-C(O)}N(R^{18})_2$, $\text{-N}(R^{18})_2$, $\text{-NR}^{18}\text{C(O)R}^{18}$, $\text{-NR}^{18}\text{C(O)}_2\text{R}^{18}$, $\text{-NR}^{18}\text{C(O)}N(R^{18})_2$, $\text{-NR}^{18}\text{S(O)}_2\text{R}^{18}$, $\text{–O-}(C_2\text{-}C_6)$alkenyl, $\text{-C(O)}_2\text{R}^{18}$, unsubstituted $(C_1\text{-}C_6)$alkyl, $(C_1\text{-}C_6)$alkyl substituted with one or more U groups, $\text{-O-}(C_1\text{-}C_6)$alkylene-$\text{C(O)R}^{18}$, or -C(O)R^{18};

with the proviso wherein the group $\text{-N}(R^{18})_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl group is fused with $(C_6\text{-}C_{12})$aryl, $(C_2\text{-}C_{10})$heteroaryl, $(C_3\text{-}C_7)$cycloalkyl or $(C_3\text{-}C_5)$heterocycloalkyl;

R^{16} is selected from the group consisting of R^9 and $\text{–C(O)-}R^{12}$;

each R^{18} is independently selected from the group consisting of H, unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more X^3 groups, R^{12}, R^{13} and R^{14};

with the proviso that when R^{18} is attached to N, then each R^{18} is independently selected from the group consisting of H, unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more W^3 groups, $\text{-C(O)}R^{21}$, R^{12}, R^{13} and R^{14};

R^{19} is selected from the group consisting of H, TBS, TIPS, Tf and R^{21};

each R^{20} is independently selected from the group consisting of H, unsubstituted $(C_1\text{-}C_6)$alkyl, $(C_1\text{-}C_6)$alkyl substituted with one or more U groups, $\text{-}(C_1\text{-}C_6)$alkylene-R^{22}, $\text{-}(C_1\text{-}C_6)$alkylene-R^{23}, $\text{-}(C_1\text{-}C_6)$alkylene-R^{24}, $\text{-C(O)-}R^{24}$, $\text{-}(C_1\text{-}C_6)$alkylene-O-R^{19}, R^{24}, unsubstituted $(C_3\text{-}C_5)$heterocycloalkyl, $(C_3\text{-}C_5)$heterocycloalkyl substituted with one or more W^3 groups, and benzo-fused $(C_3\text{-}C_7)$cycloalkyl;

R^{21} is selected from the group consisting of unsubstituted $(C_1\text{-}C_6)$alkyl, $(C_1\text{-}C_6)$alkyl substituted with one or more U groups,

-(C_1-C_6)alkylene-O-(C_1-C_6)alkyl, -(C_1-C_6)alkylene-O-(C_6-C_{12})aryl, unsubstituted (C_6-C_{12})aryl, and (C_6-C_{12})aryl substituted with one or more W^1 groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more W^2 groups, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl with one or more W^3 groups, -O-(C_1-C_6)alkylene-O-R^{24}, -C(O)-O-(C_1-C_6)alkylene-O-R^{24};

-C(O)-(C_1-C_6)alkylene -R^{23}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -S(O)$_2$-R^{24}, and -(C_1-C_6)alkylene-O-(C_1-C_6)alkylene-O-R^{24}, with the proviso that –O-O- cannot be formed with R^{21} and the atom said R^{21} is attached to;

R^{22} is selected from the group consisting of unsubstituted (C_6-C_{12})aryl and (C_6-C_{12})aryl substituted with one or more W^1 groups;

R^{23} is selected from the group consisting of unsubstituted (C_2-C_{10})heteroaryl and (C_2-C_{10})heteroaryl substituted with one or more W^2 groups;

R^{24} is is selected from the group consisting of unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more X^4 groups, unsubstituted (C_1-C_6)alkyl and (C_1-C_6)alkyl substituted with one or more U groups;

each R^{25} is independently selected from the group consisting of H, R^{22}, R^{23}, unsubstituted (C_1-C_6)alkyl, (C_1-C_6)alkyl substituted with one or more U groups, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, -(C_1-C_6)alkylene-OR^{19}, -(C_1-C_6)alkylene-N$R^{19}R^{19}$, -(C_1-C_6)alkylene-SR^{19},- (C_1-C_6)alkylene-R^{23}, -(C_1-C_6)alkylene-R^{22}, unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl, -(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -C(O)-R^{24}, -C(O)-R^{22}, -C(O)-R^{24}, -C(O)-O-R^{22}, -C(O)-O-R^{24}, -NHR^{22}, -NHR^{24}, -S(O)$_2$-R^{24}, -C(O)-NH-R^{22} and -C(O)-NH-R^{24};

with the proviso wherein the group -N(R^{25})$_2$ both R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted (C_3-C_5)heterocycloalkyl group is fused with (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, (C_3-C_7)cycloalkyl or (C_3-C_5)heterocycloalkyl;

each W^1 is independently selected from the group consisting of halogen, -CN, -OH, -O-S(O)$_2$-(C_1-C_6)haloalkyl, unsubstituted (C_6-C_{12})aryl, (C_6-C_{12})aryl substituted with one or more Z groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more Z groups, and –O-(C_1-C_6)alkyl;

each W^2 is independently selected from the group consisting of halogen, unsubstituted (C_6-C_{12})aryl, and (C_6-C_{12})aryl substituted with one or more Z groups;

each W^3 is independently selected from the group consisting of –OH, (C_1-C_6)alkyl, -(C_1-C_6)alkylene-OH, -O-(C_1-C_6)alkyl, -C(O)- (C_1-C_6)alkyl, -C(O)NH_2, -NHC(O) (C_1-C_6)alkyl, -NHC(O)H, -NHC(O)-O-(C_1-C_6)alkyl and -C(O)-O-(C_1-C_6)alkyl; or

two W^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;

each W^4 is independently halogen or (C_1-C_6)alkyl;

Ar^1 is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

Ar^2 is selected from the group consisting of unsubstituted phenyl, phenyl substituted with one or more X^1 groups, pyridyl-2-yl and pyridyl-2-yl substituted with one or more X^2 groups;

each X^1 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S(O)$_2$-(C_1-C_6)haloalkyl, unsubstituted (C_6-C_{12})aryl, (C_6-C_{12})aryl substituted with one or more Z groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more Z groups, –O-(C_3-C_7)cycloalkyl, –O-(C_3-C_7)cycloalkyl(C_1-C_6)alkyl, -O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)N(R^{20})$_2$, -O-(C_1-C_6)alkylene-O-R^{19}, unsubstituted –O-(C_1-C_6)alkyl, -O-(C_1-C_6)alkyl substituted with one or more U groups, -O-(C_2-C_7)alkenyl, -O-(C_1-C_6)alkylene-O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)R^{24}, -O-(C_1-C_6)alkylene-C(O)OR^{19}, –O-(C_1-C_6)alkyl, -N(R^{25})$_2$, -C(O)(C_1-C_6)alkyl, –C(O)OH, -C(O)O-(C_1-C_6)alkyl, -C(O)O-(C_3-C_7)cycloalkyl, -C(O)N(R^{25})$_2$, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl substituted with one or more W^3

groups, unsubstituted(C_3-C_5)heterocycloalkyl, -(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -O-(C_2-C_7)alkenylene-O-(C_1-C_6)alkylene-O-R^{24}, -O-(C_1-C_6)alkylene-N(R^{25})$_2$, -O-(C_1-C_6)alkylene-C(O)N(R^{25})$_2$, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S(O)$_2$-R^{24}, and (C_2-C_7)alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted (C_3-C_5)heterocycloalkyl, (C_3-C_5)heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted (C_3-C_5)heterocycloalkyl group is fused with (C_6-C_{12})aryl, (C_2-C_{10})heteroaryl, (C_3-C_7)cycloalkyl or (C_3-C_5)heterocycloalkyl;

each X^2 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S(O)$_2$-(C_1-C_6)haloalkyl, unsubstituted (C_6-C_{12})aryl, (C_6-C_{12})aryl substituted with one or more Z groups, unsubstituted (C_2-C_{10})heteroaryl, (C_2-C_{10})heteroaryl substituted with one or more Z groups, –O-(C_3-C_7)cycloalkyl, –O-(C_3-C_7)cycloalkyl(C_1-C_6)alkyl, -O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)N(R^{20})$_2$, -O-(C_1-C_6)alkylene-O-R^{19}, unsubstituted –O-(C_1-C_6)alkyl, -O-(C_1-C_6)alkyl substituted with one or more U groups, -O-(C_2-C_7)alkenyl, -O-(C_1-C_6)alkylene-O-(C_1-C_6)alkylene-OR^{19}, -O-(C_1-C_6)alkylene-C(O)R^{24}, -O-(C_1-C_6)alkylene-C(O)OR^{19}, –O-(C_1-C_6)alkyl, -N(R^{25})$_2$, -C(O)(C_1-C_6)alkyl, –C(O)OH, -C(O)O-(C_1-C_6)alkyl, -C(O)O-(C_3-C_7)cycloalkyl, -C(O)N(R^{25})$_2$, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl, -O-(C_1-C_6)alkylene-(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, unsubstituted(C_3-C_5)heterocycloalkyl, -(C_3-C_5)heterocycloalkyl substituted with one or more W^3 groups, -O-(C_2-C_7)alkenylene-O-(C_1-C_6)alkylene-O-R^{24}, -O-(C_1-C_6)alkylene-N(R^{25})$_2$, -O-(C_1-C_6)alkylene-C(O)N(R^{25})$_2$, unsubstituted (C_3-C_7)cycloalkyl, (C_3-C_7)cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S(O)$_2$-R^{24}, and (C_2-C_7)alkenyl;

with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted

$(C_3$-$C_5)$heterocycloalkyl, $(C_3$-$C_5)$heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted $(C_3$-$C_5)$heterocycloalkyl group is fused with $(C_6$-$C_{12})$aryl, $(C_2$-$C_{10})$heteroaryl, $(C_3$-$C_7)$cycloalkyl or $(C_3$-$C_5)$heterocycloalkyl;

each X^3 is independently selected from the group consisting of –OH, $(C_1$-$C_6)$alkyl, -$(C_1$-$C_6)$alkylene-OH, -O-$(C_1$-$C_6)$alkyl, -C(O)- $(C_1$-$C_6)$alkyl, -$C(O)NH_2$, -NHC(O) $(C_1$-$C_6)$alkyl, -NHC(O)H, -NHC(O)-O-$(C_1$-$C_6)$alkyl and -C(O)-O-$(C_1$-$C_6)$alkyl; or

two X^3 groups together with the ring carbon atom to which they are attached form a carbonyl group;


each X^4 is independently halogen or $(C_1$-$C_6)$alkyl;

each U is independently selected from the group consisting of –OH, -O-$(C_1$-$C_6)$alkyl, -O-$(C_6$-$C_{12})$aryl, -O-$(C_1$-$C_6)$alkylene-$(C_6$-$C_{12})$aryl, -O-$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkyl, -O-$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$haloalkyl, -O-$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl, halogen, -CN, $(C_3$-$C_7)$cycloalkyl, $(C_3$-$C_5)$heterocycloalkyl, $(C_6$-$C_{12})$aryl, $(C_2$-$C_{10})$heteroaryl, OTBS, OTIPS and OTf; and

each Z is independently selected from the group consisting of –OH; -O-$(C_1$-$C_6)$alkyl; halogen; $(C_1$-$C_6)$alkyl; –CN; -CF_3; $(C_3$-$C_7)$cycloalkyl; -$(C_1$-$C_6)$alkylene-OH; -$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkyl; -$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkyl substituted with one or more groups selected from the group consisting of –OH, -O-$(C_1$-$C_6)$alkyl, halogen, -CN, $(C_3$-$C_7)$cycloalkyl, $(C_3$-$C_5)$heterocycloalkyl, $(C_6$-$C_{12})$aryl, $(C_2$-$C_{10})$heteroaryl; -$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkyl; -$(C_1$-$C_6)$alkylene-O-$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl; -$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl; and -$(C_1$-$C_6)$alkylene-O-$(C_6$-$C_{12})$aryl substituted with one or more groups selected from the group consisting of halogen, -CN, -OH, -O-$S(O)_2$-$(C_1$-$C_6)$haloalkyl, $(C_6$-$C_{12})$aryl, $(C_2$-$C_{10})$heteroaryl and –O-$(C_1$-$C_6)$alkyl; or


two Z groups together with the ring carbon atom to which they are attached form a carbonyl group.

3. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof having the following Formula (II):




(II)

4. The compound of Claim 3, or a pharmaceutically acceptable salt, solvate or ester thereof having the following Formula (IIA):



(IIA) .

5. The compound of Claim 4, or a pharmaceutically acceptable salt, solvate or ester thereof having the following Formula (IIB):



(IIB) .

6. The compound of Claim 4, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

Ar^1 is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

Ar^2 is selected from the group consisting of unsubstituted phenyl, phenyl substituted with one or more X^1 groups, pyridyl-2-yl and pyridyl-2-yl substituted with one or more X^2 groups;

R^7 is selected from the group consisting of H, alkyl, alkenyl, -alkylene-$N(R^9)_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, -alkylene-C(O)H, and -C(O)-O-R^{11};

R^9 is selected from the group consisting of H and alkyl;

R^{14} is selected from the group consisting of alkyl, unsubstituted cycloalkyl, or cycloalkyl substituted with one or more X^4 groups; and

each R^{15} is independently selected from the group consisting of H, alkyl, alkenyl, -alkylene-R^{12}, and –O-alkenyl.

7. The compound of Claim 6, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:


Ar^1 is phenyl substituted with one or more X^1 groups;

Ar^2 is phenyl substituted with one or more X^1 groups or pyridyl substituted with one or more X^2 groups;

R^7 is H; and


R^{15} is alkyl.

8. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, having the following Formula (III):



(III)

9. The compound of Claim 8, or a pharmaceutically acceptable salt, solvate or ester thereof, having the following Formula (IIIA):



(IIIA) ,

wherein:

Ar^1 is selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

Ar^2 is selected from the group consisting of unsubstituted phenyl, phenyl substituted with one or more X^1 groups, pyridyl-2-yl and pyridyl-2-yl substituted with one or more X^2 groups; and

R^3 and R^4 are each independently H or alkyl;


10. The compound of Claim 9, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

R^3 and R^4 are each independently H or $–CH_3$;

Ar^1 is unsubstituted phenyl or phenyl substituted with one or more halogens; and


Ar^2 is selected from the group consisting of unsubstituted phenyl, phenyl substituted with one or more halogens, unsubstituted pyridyl, and pyridyl substituted with one or more X^2 groups.

11. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, having the following Formula (V):




(V) .

12. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, having the following Formula (ID):



(ID).

13. The compound of Claim 12, or a pharmaceutically acceptable salt, solvate or ester thereof, having the following Formula (IE):



(IE).

14. (Original) The compound of Claim 12, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

Y^1 is NH or N-Boc;

R^3 is H or alkyl;

R^4 is H or alkyl;

R^{15} is alkyl;

and


Ar^1 and Ar^2 are selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups.

15. The compound of Claim 14, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:


each X^1 is independently selected from the group consisting of halogen, -CN, -OH, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-C(O)N(R^{20})$_2$, -O-alkylene-O-R^{19}, unsubstituted, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl,-O-alkylene-O-alkylene-OR^{19}, -O-alkylene-C(O)R^{24}, -O-alkylene-C(O)OR^{19} and –O-alkyl.


16. The compound of Claim 14, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:



each X^1 is independently selected from the group consisting of –OCH_3, -OH, -OTf, -CN, -OCH_2CH_3, -OCH(CH_3)$_2$,


















,

,



,

,


,


,

,


,

and

.

17. A compound of Formula (ID),



(ID)

or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

Y^1 is NH or N-Boc;

R^3 is H or alkyl;






R^4 is H or alkyl;


R^{15} is alkyl;


















Ar^1 and Ar^2 are selected from the group consisting of unsubstituted phenyl and phenyl substituted with one or more X^1 groups;

and


each X^1 is independently selected from the group consisting of $–OCH_3$, -OH, -OTf, -CN, $-OCH_2CH_3$, $-OCH(CH_3)_2$,


,

,

,

,





,

,

,


,

,

,

,

,

,

,

,

,

,

,

and

.

18. A compound of Formula (IE),



(IE)

or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

Y^1 is NH or N-Boc;






R^3 is H or alkyl;




R^4 is H or alkyl;


R^{15} is alkyl;




and



each X^1 is independently selected from the group consisting of $–OCH_3$, -OH, -OTf, -CN, $-OCH_2CH_3$, $-OCH(CH_3)_2$,



 ,  ,  ,  , 


 ,  ,  ,  ,








 ,  ,  ,

 ,  ,

 ,  ,

 ,

 ,  ,

 ,  and  .

19. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:


each R^{15} is independently selected from the group consisting of H, -N_3, halogen, alkenyl, -alkylene-R^{12}, -alkylene-O-R^9, -alkylene-$N(R^{18})_2$, -alkylene-C(O)H, -OH, -CN, -O-alkyl, $-C(O)N(R^{18})_2$, $-N(R^{18})_2$, $-NHC(O)R^{18}$, $-NHC(O)_2R^{18}$, $-NR^{18}C(O)N(R^{18})_2$, $-NHS(O)_2R^{18}$, –O-alkenyl, $-C(O)_2R^{18}$, unsubstituted alkyl, alkyl substituted with one or more U groups, -O-alkylene-$C(O)R^{18}$, or $-C(O)R^{18}$;


with the proviso wherein the group $-N(R^{18})_2$, both R^{18} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl.

20. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

each R^{15} is independently selected from the group consisting of

H, -OH, –OCH_3, -OCH_2CH_3, -CH_3, -CH_2CH_3, -$CH_2CH_2CH_3$, -N_3, -NH_2, -CO_2H, -CO_2CH_3,-CH_2OH, -CH_2CH_2OH, -CH_2CH_2OTBS, -CH_2OCH_3, -OCH_2CH_2OH,






21. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

R^3 and R^4 are each independently selected from the group consisting of H, -O-R^9, R^{11}, and –N(R^{16})$_2$.

22. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

each R^3 and R^4 is independently selected from the group consisting of H, -CH_3, -CH_2OH, -$CH_2OCH_2CH_2OCH_3$, -CH_2OCH_3, -$CH_2OCH_2CH_3$, -$CH_2OCH_2CH_2CH_3$, -CH_2CH_2OH,




23. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein

R^7 is selected from the group consisting of H, alkyl, arylalkyl, alkenyl, -alkylene-N(R^9)$_2$, -alkylene-O-R^9, -alkylene-R^{12}, -C(O)-R^{14}, -alkylene-C(O)H, -C(O)-O-R^{11}, and Boc.

24. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein

R^7 is H, -CH_3, Boc,




25. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein Ar^1 is phenyl substituted with one or more X^1 groups; and


each X^1 is independently selected from the group consisting of halogen, -CN, -O-R^{19}, -OH, -O-S(O)$_2$-haloalkyl, unsubstituted aryl, aryl substituted with one or more Z groups, unsubstituted heteroaryl, heteroaryl substituted with one or more Z groups, –O-cycloalkyl, –O-cycloalkylalkyl, -O-alkylene-OR^{19}, -O-alkylene-C(O)N(R^{20})$_2$, -O-alkylene-O-R^{19}, unsubstituted –O-alkyl, -O-alkyl substituted with one or more U groups, -O-alkenyl, -O-alkylene-O-alkylene-OR^{19}, -O-alkylene-C(O)R^{24}, -O-alkylene-C(O)OR^{19},–O-alkyl, -N(R^{25})$_2$, -C(O)alkyl, –C(O)OH, -C(O)O-alkyl, -C(O)O-cycloalkyl, -C(O)N(R^{25})$_2$, -O-alkylene-heterocycloalkyl, -O-alkylene-heterocycloalkyl substituted with one or more W^3 groups, unsubstituted heterocycloalkyl, -heterocycloalkyl substituted with one or more W^3 groups, -O-alkenylene-O-alkylene-O-R^{24}, -O-alkylene-N(R^{25})$_2$, -O-alkylene-C(O)N(R^{25})$_2$, unsubstituted cycloalkyl, cycloalkyl substituted with one or more W^4 groups, -S(O)-R^{24}, -S(O)$_2$-R^{24}, and alkenyl;


with the proviso wherein the group -N(R^{20})$_2$ or -N(R^{25})$_2$ both R^{20} or R^{25} groups taken together with the N atom to which they are bonded form an unsubstituted heterocycloalkyl, heterocycloalkyl substituted with one or more X^3 groups, or said substituted or unsubstituted heterocycloalkyl group is fused with aryl, heteroaryl, cycloalkyl or heterocycloalkyl.


26. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein:

each X^1 is independently selected from the group consisting of Cl, F, -OCH_3, -OCH_2CH_3, -OH, -OTf, -CN, -OCH_2CH_3, -$OCH(CH_3)_2$,











































27. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein


Ar^2 is phenyl substituted with one or more X^1 groups or pyridyl substituted with one or more X^2 groups;

X^1 is selected from the group consisting of –OH, -CN, halogen, -OTIPS, -OTf, -O-alkyl, -O-alkyl-OH and heteroaryl; and

X^2 is selected from the group consisting of halogen and cycloalkyl.


28. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof, wherein

Ar^2 is phenyl substituted with one or more X^1 groups or pyridyl substituted with one or more X^2 groups,

X^1 is selected from the group consisting of –OH, -CN, Cl, -OTIPS, $-OCH_3$, $-OCH_2CH_3$, -OTf and  ; and


X^2 is selected from the group consisting of Cl, Br and  .

29. The compound of Claim 1 or a pharmaceutically acceptable salt, solvate or ester thereof having the following Formulas (IF) or (IFa):




(IIF)

or




(IIFa) .

30. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof having the following Formula (IG):




(IG) .

31. The compound of Claim 1, or a pharmaceutically acceptable salt, solvate or ester thereof having the following Formula (IH):




(IH) .


32. A compound, or a pharmaceutically acceptable salt, solvate, ester or stereoisomer thereof, selected from the group consisting of:




,




,




,




,




,




,




,



















,

































,




























,



O
N
Cl
Cl
F
F
Cl
,

O
N
Cl
Cl
F
F
Cl
,

O
N
Cl
N
Cl
F
F
Cl
,

O
N
Cl
Cl
F
F
Cl
,

O
BocN
Cl
OMe
Cl
,

O
HN
Cl
OMe
Cl
,

O
HN
Cl
OH
Cl
,

O
HN
Cl
OTf
Cl
,

O
HN
Cl
CN
Cl
,

O
HN
Cl
Cl
,

O
HN
Cl
Me
Cl
,

O
HN
Cl
N
Cl
,

O
HN
Cl
Cl
,

O
OH
HN
Cl
O
Me
Cl
,

O
Me
HN
Cl
OMe
Cl
,

O
Et
HN
Cl
OMe
Cl
,


CN
O
HN
Cl
OMe
Cl
,

O
HN
Cl
O
Me
Cl
,

O
Me
HN
Cl
OH
Cl
,

O
Me
HN
Cl
OTf
Cl
,

O
Me
HN
Cl
CN
Cl
,

O
Me
HN
Cl
Cl
,

O
Me
HN
OMe
Cl
Cl
,

O
Me
HN
CN
Cl
Cl
,

O
Me
HN
Me
Cl
Cl
,

O
NH2
BocN
OMe
Cl
Cl
,

O
N
BocN
OMe
Cl
Cl
,

O
N
HN
OMe
Cl
Cl
,

O
O
N
HN
OMe
Cl
Cl
,

O
O
O
NH
I
BocN
OMe
Cl
Cl
,

O
O
O
Br
O
NH
HN
OMe
Cl
Cl
,



O
BocN
Cl
Cl
Cl
,

O
N3
BocN
Cl
Cl
Cl
,

O
NH2
BocN
Cl
Cl
Cl
,

O
NH2
HN
Cl
Cl
Cl
,

O
O
HN
HN
Cl
Cl
Cl
,

O
HN
Cl
Cl
Cl
,

O
BocN
Cl
Cl
Cl
,

O
Me
HN
Cl
Cl
Cl
,

O
BocN
Cl
OMe
Cl
,

O
Me
BocN
Cl
OMe
Cl
,

O
Me
HN
Cl
OH
Cl
,

O
Me
HN
Cl
OTf
Cl
,

O
Me
HN
Cl
CN
Cl
,

O
Me
HN
Cl
OEt
Cl
,

O
Me
HN
Cl
Cl
Cl
,


O
Me
HN
Cl
Cl
Cl
,

HN
O
Cl
Cl
Cl
,

O
HN
Cl
Cl
Cl
,

O
O
Cl
N
Cl
Br
,

O
O
Cl
Cl
Cl
,

O
O
Cl
N
Cl
Cl
Cl
,

O
O
N
Cl
Cl
Cl
,

O
O
N
Cl
Cl
,

HO
O
Cl
N
Cl
Br
,

O
Et
BocN
H
H
H
Cl
OMe
Cl
,

O
Et
HN
H
H
H
Cl
OH
Cl
,

O
Et
HN
H
H
H
Cl
OTf
Cl
,

O
Et
HN
H
H
H
Cl
CN
Cl
,

O
Et
HN
H
H
H
Cl
OEt
Cl
,

O
Me
HN
H
H
H
Cl
O
Cl
,



O
Me
HN
H
H
H
Cl
O
Cl
,

O
Me
HN
H
H
H
Cl
O
O
Cl
,

O
Me
HN
H
H
H
Cl
O
CF3
Cl
,

O
Me
HN
H
H
H
Cl
O
NH2
O
Cl
,

O
Me
HN
H
H
H
Cl
O
OH
Cl
,

O
Me
HN
H
H
H
Cl
OCHF2
Cl
,

O
Me
HN
H
H
H
Cl
O
Cl
,

O
Me
HN
H
H
H
Cl
O
Cl
,

O
Et
HN
H
H
H
Cl
O
O
O
Cl
,

O
Et
HN
H
H
H
Cl
O
OTBS
Cl
,


O
Et
HN
H
H
H
Cl
O
OH
Cl


,


O
Et
HN
H
H
H
Cl
O
NH2
O
Cl


,


O
Et
HN
H
H
H
Cl
O
O
Cl


,


O
Me
HN
H
H
H
Cl
O
OMe
O
Cl


,


O
Me
HN
H
H
H
Cl
O
OH
O
Cl


,


O
Me
HN
H
H
H
Cl
O
N
OH
O
Cl













,
















O
Me
HN
H
H
H
Cl
O
H
N
OMe
OMe
O
Cl















,

















O
Me
HN
H
H
H
Cl
O
H
N
OH
O
Cl













,















































































































,

,

,

,

,

,

,

,

,

,

,

,


















,

,

,

,

,

,

,

,

,

,

,

,



,

,

,

,

,

,

,

OTBS,

,

,

,

,

,

,

,

,

















,













,













,


































,

,

,

,

,

,

,

,

,

,

,

,














,

,















,

,

,

,

,

,

,

,

,

,

,

,
















,

,


,














,

,

,

,

,

,

,

,

,

,

,

,

,


,

,

,

,

,

,

,

,

,

,

,

,

,


,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,






















,

,

,

,

,

,

,

,

,

,

,

,

















,


,


,


,



,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,



,

,

,

,

,

,

,

,

,

,

,

,

,

,

,

,




,


,



,


,


and



.

33. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

34. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

35. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

36. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

37. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

38. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

39. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

40. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

41. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

42. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

43. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

44. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

45. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

46. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

47. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

48. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

49. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

50. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

51. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

52. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

53. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

54. A compound having the following structural formula:




or a pharmaceutically acceptable salt, solvate, or ester thereof.

55. A compound having the following structural formula:



or a pharmaceutically acceptable salt, solvate, or ester thereof.

56. A compound having the following structural formula:



or a pharmaceutically acceptable salt, solvate, or ester thereof.

57. A composition comprising:

at least one compound of Formula (I) according to Claim 1, or a pharmaceutically acceptable salt, solvate, or ester thereof; and

at least one cholesterol lowering compound.

58. The composition of Claim 57, wherein said at least one cholesterol lowering compound is at least one sterol absorption inhibitor or at least one 5α-stanol absorption inhibitor.

59. The composition of Claim 57, wherein said at least one cholesterol lowering compound is at least one substituted azetidinone compound or substituted β-lactam compound or a pharmaceutically acceptable salt, solvate, or ester thereof.

60. The composition of Claim 57, wherein said at least one cholesterol lowering compound is ezetimibe.