

DESCRIPTION

CRYSTAL OF PYRIMIDINE COMPOUND

Technical Field

[0001]

The present invention relates to a crystal of pyrimidine compound useful as an antitumor agent, and a pharmaceutical composition comprising the crystal.

Background Art

[0002]

HER2 (which is also referred to as "ErbB2") is receptor tyrosine kinase belonging to the ErbB family.

HER2 is considered to be a proto-oncogene. It has been reported that HER2 gene amplification, overexpression, mutation and the like occur in various types of cancers. From non-clinical and clinical research data, it is considered that activation of HER2 and downstream signals plays an important role in the survival and/or proliferation, etc. of cancer cells associated with the genetic abnormality, overexpression and the like of HER2 (Non Patent Literature 1).

Accordingly, an inhibitor capable of regulating the kinase activity of HER2 is assumed to inhibit HER2 and downstream signals in cancer cells having HER2 gene amplification, overexpression or mutation, so as to exhibit antitumor effects on the cancer cells. Therefore, such an inhibitor is considered to be useful for the treatment, life-prolonging, or QOL improvement of cancer patients.

[0003]

It has been reported that brain metastasis occurs in approximately 25% to 40% of lung cancer cases, in approximately 15% to 30% of breast cancer cases, and in certain percentages of other multiple cancer cases (Non Patent Literatures 2 and 3). As a matter of fact, it has been reported that brain metastasis occurs in approximately 20% to 30% of HER2-positive breast cancer cases (Non Patent Literature 4).

[0004]

Compounds having HER2 inhibitory activity, such as Lapatinib and Neratinib, have been approved as therapeutic agents against HER2-positive breast cancer. However, it has been reported that since all of these therapeutic agents are substrates of p-gp or Bcrp, the brain penetration properties of these agents are limited in non-clinical tests (Non Patent Literature 5). In fact, in clinical tests using Lapatinib or Neratinib,

sufficient effects of these agents could not be obtained against brain metastatic cancer (Non Patent Literatures 6, 7, 8, and 9).

From the viewpoint of the control of pathological conditions including brain metastasis nidus, it has been desired to develop a HER2 inhibitor having inhibitory activity against HER2 and also having brain penetration properties.

[0005]

Generally, when a compound is used as an active ingredient in a pharmaceutical product, the chemical and physical stability of the compound is required to maintain stable quality and/or to facilitate storage control. It has been desired to develop a HER2 inhibitor having chemical and physical stability and inhibitory activity against HER2 and also having brain penetration properties.

Citation List

Non Patent Literature

[0006]

Non Patent Literature 1: Cancer Treatment Reviews, 40, pp. 770-780 (2014)

Non Patent Literature 2: Current Oncology, 25, pp. S103-S114 (2018)

Non Patent Literature 3: Breast Cancer Research, 18(1), 8, pp. 1-9 (2016)

Non Patent Literature 4: Journal of Clinical Oncology, 28, pp. 3271-3277 (2010)

Non Patent Literature 5: Journal of Medicinal Chemistry, 59, pp. 10030-10066 (2016)

Non Patent Literature 6: Journal of Medicinal Chemistry, 26, pp. 2999-3005 (2008)

Non Patent Literature 7: Journal of Clinical Oncology, 26, pp. 1993-1999 (2008)

Non Patent Literature 8: Journal of Clinical Oncology, 28, pp. 1301-1307 (2010)

Non Patent Literature 9: Journal of Clinical Oncology, 34, pp. 945-952 (2016)

Summary of Invention

Technical Problem

[0007]


It is an object of the present invention to provide a HER2 inhibitor having chemical and physical stability and inhibitory activity against HER2 and also having brain penetration properties. In particular, it is an object of the present invention to provide a crystal of a HER2 inhibitor that are excellent in stability, have good oral absorbability, and can be obtained with good reproducibility.

Solution to Problem

[0008]

As a result of intensive studies in order to achieve the above objects, the present inventors have found that 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide represented by the following formula (I) (hereinafter also referred to as "compound (I)") has HER2 inhibitory activity and brain penetration properties, and is useful as a therapeutic agent for diseases involving HER2 (particularly malignant tumor) by inhibiting HER2.

[0009]



[0010]

In addition, the present inventors have found that there are two free-form crystals (type I crystal and type II crystal) and crystals with acids (salt crystals or co-crystals) of the compound (I), including crystals with hydrochloric acid (type I crystal with hydrochloric acid, type II crystal with hydrochloric acid, type III crystal with hydrochloric acid), crystals with hydrobromic acid, crystals with 1 equivalent of tartaric acid, crystals with fumaric acid (type I crystal with 0.5 equivalents of fumaric acid, type II crystal with 0.5 equivalents of fumaric acid, type I crystal with 1 equivalent of fumaric acid, type II crystal with 1 equivalent of fumaric acid, type III crystal with 1 equivalent of fumaric acid, type IV crystal with 1 equivalent of fumaric acid, type V crystal with 1 equivalent of fumaric acid), and crystals with succinic acid.

The present inventors have found that among these crystals, a free-form type I crystal, a free-form type II crystal, a type II crystal with 1 equivalent of fumaric acid, a type V crystal with 1 equivalent of fumaric acid have advantageous properties in pharmaceutical manufacturing such as a non-hygroscopic property and excellent solid stability, and also have excellent oral absorbability. This has led to the completion of the present invention.

[0011]

Specifically, the present invention includes the following embodiments.

[1] A crystal having peaks at three or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 5.5° , 6.8° , 9.3° , 13.4° , 15.3° , 16.3° , 18.5° , 19.8° , 22.0° , and 24.5° in a powder X-ray diffraction spectrum, which is a type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:1.

[2] The crystal according to [1], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 5.5° , 6.8° , 9.3° , 13.4° , 15.3° , 16.3° , 18.5° , 19.8° , 22.0° , and 24.5° in a powder X-ray diffraction spectrum.

[3] The crystal according to [1] or [2], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 1.

[4] The crystal according to any one of [1] to [3], which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 178°C .

[5] The crystal according to any one of [1] to [4], wherein the purity of the crystal is 90% by mass or more.

[6] A crystal having peaks at three or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 8.3° , 14.8° , 17.3° , 18.0° , 19.1° , 20.3° , 21.0° , 22.5° , 23.0° , and 26.2° in a powder X-ray diffraction spectrum, which is a type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide.

[7] The crystal according to [6], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 8.3° , 14.8° , 17.3° , 18.0° , 19.1° , 20.3° , 21.0° , 22.5° , 23.0° , and 26.2° in a powder X-ray diffraction spectrum.

[8] The crystal according to [6] or [7], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 3.

[9] The crystal according to any one of [6] to [8], which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 182°C .

[10] The crystal according to any one of [6] to [9], wherein the purity of the crystal is 90% by mass or more.

[11] A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 9.9° , 11.7° , 13.2° , 17.7° , 18.1° , 18.8° , and 20.8° in a powder X-ray diffraction

spectrum, which is a type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide.

[12] The crystal according to [11], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 9.9° , 11.7° , 13.2° , 17.7° , 18.1° , 18.8° , and 20.8° in a powder X-ray diffraction spectrum.

[13] The crystal according to [11] or [12], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 5.

[14] The crystal according to any one of [11] to [13], which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 182°C .

[15] The crystal according to any one of [11] to [14] wherein the purity of the crystal is 90% by mass or more.

[16] A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 6.9° , 9.4° , 10.2° , 13.7° , 21.1° , 23.6° , and 26.5° in a powder X-ray diffraction spectrum, which is a type V crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:1.

[17] The crystal according to [16], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 6.9° , 9.4° , 10.2° , 13.7° , 21.1° , 23.6° , and 26.5° in a powder X-ray diffraction spectrum.

[18] The crystal according to [16] or [17], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 7.

[19] The crystal according to any one of [16] to [18], which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 151°C .

[20] The crystal according to any one of [16] to [19], wherein the purity of the crystal is 90% by mass or more.

[21] A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 6.4° , 10.3° , 12.8° , 15.0° , 20.7° , 23.4° , and 26.6° in a powder X-ray diffraction spectrum, which is a type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:1.

[22] The crystal according to [21], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$)

of 6.4°, 10.3°, 12.8°, 15.0°, 20.7°, 23.4°, and 26.6° in a powder X-ray diffraction spectrum.

[23] The crystal according to [21] or [22], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 9.

[24] The crystal according to any one of [21] to [23], which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 153°C.

[25] The crystal according to any one of [21] to [24], wherein the purity of the crystal is 90% by mass or more.

[26] A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 6.4°, 7.5°, 9.7°, 11.7°, 15.1°, 19.6°, and 23.9° in a powder X-ray diffraction spectrum, which is a type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:0.5.

[27] The crystal according to [26], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 6.4°, 7.5°, 9.7°, 11.7°, 15.1°, 19.6°, and 23.9° in a powder X-ray diffraction spectrum.

[28] The crystal according to [26] or [27], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 11.

[29] The crystal according to any one of [26] to [28], which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 161°C.

[30] The crystal according to any one of [26] to [29], wherein the purity of the crystal is 90% by mass or more.

[31] A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 5.4°, 6.4°, 7.3°, 12.8°, 13.4°, 14.7°, and 15.4° in a powder X-ray diffraction spectrum, which is a type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:0.5.

[32] The crystal according to [31], which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 5.4°, 6.4°, 7.3°, 12.8°, 13.4°, 14.7°, and 15.4° in a powder X-ray diffraction spectrum.

[33] The crystal according to [31] or [32], which is a crystal having a powder X-ray diffraction spectrum shown in Figure 13.

[34] The crystal according to any one of [31] to [33], which has an endothermic peak

determined by simultaneous thermogravimetry-differential thermal analysis at around 162°C.

[35] The crystal according to any one of [31] to [34], wherein the purity of the crystal is 90% by mass or more.

[36] A pharmaceutical composition comprising the crystal according to any one of [1] to [35].

[37] A pharmaceutical composition for oral administration comprising the crystal according to any one of [1] to [35].

[38] An antitumor agent comprising the crystal according to any one of [1] to [35].

[39] The crystal according to any one of [1] to [35] for use as a medicament.

[40] Use of the crystal according to any one of [1] to [35] for the production of an antitumor agent for oral administration.

[41] The crystal according to any one of [1] to [35] for use in the treatment of tumor.

[42] The crystal according to any one of [1] to [35] for use in the treatment of tumor by oral administration thereof.

[43] A method for treating tumor, comprising administering an effective amount of the crystal according to any one of [1] to [12] to a subject in need thereof.

[44] Use of the crystal according to any one of [1] to [35] for the production of a pharmaceutical composition.

[45] Use of the crystal according to any one of [1] to [35] for the production of an antitumor agent.

Effects of Invention

[0012]

A free-form type I crystal, a free-form type II crystal, a type II crystal with 1 equivalent of fumaric acid, a type V crystal with 1 equivalent of fumaric acid of the compound (I) (7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide) of the present invention are superior to other crystal forms in terms of high stability, handleability (lower hygroscopicity), quality controllability, and the like. In addition, the free-form type II crystal and the type II crystal with 1 equivalent of fumaric acid have excellent oral absorbability, and thus they are useful forms when using the compound as an active pharmaceutical ingredient. The type I crystal with 1 equivalent of fumaric acid is in a form that can be isolated as a solid from acetone (solvent) (after isolation, it is dried to form a type V crystal), it is also useful as an intermediate for the production of a type V crystal.

According to the present invention, a novel crystal of compound (I) or a novel crystal of compound (I) with an acid, and a pharmaceutical composition, an antitumor agent, or an antitumor agent for oral administration comprising the novel crystal are provided.

In preferred aspects, a type II crystal with 1 equivalent of fumaric acid, a free-form type II crystal, a free-form type I crystal, a type V crystal with 1 equivalent of fumaric acid, a type I crystal with 0.5 equivalents of fumaric acid, and a type II crystal with 0.5 equivalents of fumaric acid of the compound (I) have at least one of advantageous properties in pharmaceutical manufacturing such as a non-hygroscopic property, ability to be obtained with reproducibility, solid stability, and oral absorbability.

Brief Description of Drawings

[0013]

[Figure 1] Figure 1 shows the powder X-ray diffraction spectrum of the type II crystal of compound (I) with 1 equivalent of fumaric acid obtained in Example 1 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 2] Figure 2 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type II crystal of compound (I) with 1 equivalent of fumaric acid obtained in Example 1 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 3] Figure 3 shows the powder X-ray diffraction spectrum of the type II crystal of compound (I) obtained in Example 2 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 4] Figure 4 shows shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type II crystal of compound (I) obtained in Example 2 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 5] Figure 5 shows the powder X-ray diffraction spectrum of the type I crystal of compound (I) obtained in Example 3 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 6] Figure 6 shows shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type I crystal of compound (I) obtained in Example 3 (the left vertical axis represents the weight (mg) on the TG curve, the right

vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 7] Figure 7 shows the powder X-ray diffraction spectrum of the type V crystal of compound (I) with 1 equivalent of fumaric acid obtained in Example 4 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 8] Figure 8 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type V crystal of compound (I) with 1 equivalent of fumaric acid obtained in Example 4 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 9] Figure 9 shows the powder X-ray diffraction spectrum of the type I crystal of compound (I) with 1 equivalent of fumaric acid obtained in Example 5 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 10] Figure 10 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type I crystal of compound (I) with 1 equivalent of fumaric acid obtained in Example 5 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 11] Figure 11 shows the powder X-ray diffraction spectrum of the type I crystal of compound (I) with 0.5 equivalents of fumaric acid obtained in Example 6 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 12] Figure 12 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type I crystal of compound (I) with 0.5 equivalents of fumaric acid obtained in Example 6 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 13] Figure 13 shows the powder X-ray diffraction spectrum of the type II crystal of compound (I) with 0.5 equivalents of fumaric acid obtained in Example 7 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 14] Figure 14 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type II crystal of compound (I) with 0.5 equivalents of fumaric acid obtained in Example 7 (the left vertical axis represents the weight (mg)

on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 15] Figure 15 shows the powder X-ray diffraction spectrum of the type III crystal of compound (I) with 1 equivalent of fumaric acid obtained in Reference Example 1 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 16] Figure 16 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type III crystal of compound (I) with 1 equivalent of fumaric acid obtained in Reference Example 1 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 17] Figure 17 shows the powder X-ray diffraction spectrum of the type IV crystal of compound (I) with 1 equivalent of fumaric acid obtained in Reference Example 2 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 18] Figure 18 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type IV crystal of compound (I) with 1 equivalent of fumaric acid obtained in Reference Example 2 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 19] Figure 19 shows the powder X-ray diffraction spectrum of the type I crystal of compound (I) with hydrochloric acid obtained in Reference Example 3 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 20] Figure 20 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type I crystal of compound (I) with hydrochloric acid obtained in Reference Example 3 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 21] Figure 21 shows the powder X-ray diffraction spectrum of the type II crystal of compound (I) with hydrochloric acid obtained in Reference Example 4 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 22] Figure 22 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type II crystal of compound (I) with hydrochloric acid obtained in Reference Example 4 (the left vertical axis represents the weight (mg) on the

TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 23] Figure 23 shows the powder X-ray diffraction spectrum of the type III crystal of compound (I) with hydrochloric acid obtained in Reference Example 5 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 24] Figure 24 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the type III crystal of compound (I) with hydrochloric acid obtained in Reference Example 5 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 25] Figure 25 shows the powder X-ray diffraction spectrum of the crystal of compound (I) with hydrobromic acid obtained in Reference Example 6 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 26] Figure 26 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the crystal of compound (I) with hydrobromic acid obtained in Reference Example 6 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 27] Figure 27 shows the powder X-ray diffraction spectrum of the crystal of compound (I) with L-tartaric acid obtained in Reference Example 7 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 28] Figure 28 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the crystal of compound (I) with L-tartaric acid obtained in Reference Example 7 (the left vertical axis represents the weight (mg) on the TG curve, the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).

[Figure 29] Figure 29 shows the powder X-ray diffraction spectrum of the crystal of compound (I) with succinic acid obtained in Reference Example 8 (the vertical axis represents the intensity (counts), and the horizontal axis represents the diffraction angle (2θ)).

[Figure 30] Figure 30 shows the results of simultaneous thermogravimetry-differential thermal analysis (TG-DTA) for the crystal of compound (I) with succinic acid obtained in Reference Example 8 (the left vertical axis represents the weight (mg) on the TG curve,

the right vertical axis represents the heat flux (μV) on the DTA curve, and the horizontal axis represents the temperature ($^{\circ}\text{C}$)).


[Figure 31] Figure 31 shows the antitumor effects of the compound (I) against models involving direct brain transplantation of the Luciferase gene-introduced HER2 expressing cell line (NCI-N87-luc).

[Figure 32] Figure 32 shows the body weight reduction percentage of models involving direct brain transplantation of the Luciferase gene-introduced HER2 expressing cell line (NCI-N87-luc) caused by the compound (I).

Description of Embodiments

[0014]

The present invention relates to a crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide, and a crystal thereof with an acid (crystal of a salt thereof or co-crystal).



(I)

Specifically, the present invention relates to a type II crystal with 1 equivalent of fumaric acid, a free-form type II crystal, a free-form type I crystal, a type V crystal with 1 equivalent of fumaric acid, a type I crystal with 1 equivalent of fumaric acid, a type I crystal with 0.5 equivalents of fumaric acid, a type II crystal with 0.5 equivalents of fumaric acid of the compound (I).

In the present description, type I, type II, type III, type IV, and type V are convenient names for distinguishing the crystal forms, and the crystals according to the present invention are not limited by these names.

[0015]

In the present description, a crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-

3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with an acid means a salt crystal or a co-crystal with an acid. A salt crystal is a crystal in which a compound (I) and an acid molecule are bonded by an ionic bond, and a co-crystal is a crystal in which a compound (I) and an acid molecule are bonded by a nonionic interaction. In the present invention, a crystal of a compound (I) with an acid may be a salt crystal or a co-crystal, and includes the meanings of both. For example, a type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid means a crystal of fumarate of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide or a co-crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid.

[0016]

A crystal represents a solid in which atoms and molecules have regular repeating structures, and is different from an amorphous solid having no repeating structures. Crystalline or amorphous solids can be examined by methods such as powder X-ray diffraction analysis (XRD analysis), differential scanning calorimetry (DSC analysis), simultaneous thermogravimetry-differential thermal analysis (TG-DTA analysis), and single crystal analysis. It is known that a crystal polymorph indicates that molecules are the same but the arrangement of atoms and molecules is different in a crystal, and the peaks obtained by XRD analysis are different among crystal polymorphs. It is also known that each polymorph has different solubility, oral absorbability, stability, and the like.

[0017]

In the present description, the terms "crystal" and "amorphous" are used in the usual meaning.

[0018]

In the present description, the description "compound (I)" simply means 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide, which is used in the meaning including both "amorphous " and "crystal."

In the present description, the description "crystal of compound (I)" is used in the meaning including any of free-form crystals of compound (I) and crystals of compound (I) with an acid (salt crystal of compound (I) and co-crystal of compound (I)).

In the present description, a crystal for which molecules other than the compound

(I) constituting the crystal (other molecules constituting a salt or co-crystal) have not been specified means a free-form crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)).

In the present description, the term "equivalent" means "molar equivalent."

In a crystal of compound (I) with an acid (a salt-form crystal or co-crystal), the equivalent of the acid to the compound (I) can be analyzed, for example, by NMR or ion chromatography.

In the present description, crystals may be hydrates.

[0019]

In addition, the present invention also encompasses a labeled compound (I) or salt thereof, i.e., a compound obtained by substituting one or more atoms of a compound (I) or a salt thereof with radioactive or non-radioactive isotopes.

[0020]

As long as the crystal contains a crystal of compound (I), it may be solely one type of crystal of compound (I) or may be a polymorphic mixture containing another type of crystal of compound (I) in addition to one type of crystal of compound (I). Specifically, it is preferable that the purity of the crystal is 50% by weight or more, that is to say, 50% by weight or more is a single crystal. It is more preferable that the purity of the crystal is 75% by weight or more, that is to say, 75% by weight or more is a single crystal. It is still more preferable that the purity of the crystal is 90% by weight or more, that is to say, 90% by weight or more is a single crystal. It is even more preferable that the purity of the crystal is 95% by weight or more, that is to say, 95% by weight or more is a single crystal. It is particularly preferable that the purity of the crystal is 99% by weight or more, that is to say, 99% by weight or more is a single crystal.

[0021]

In the present description, the chemical purity is the purity measured by high performance liquid chromatography (HPLC), and when it is described as the chemical purity of the compound (I), it means the purity when the compound (I) is measured by high performance liquid chromatography. At that time, the wavelength of the detector used for the purity measurement can be appropriately set. Specifically, the chemical purity of compound (I) is preferably 95% or more, more preferably 98% or more, and particularly preferably 99% or more.

[0022]

For powder X-ray diffraction patterns, the diffraction angle and the overall pattern are important when recognizing the crystal identity due to the nature of data. The

relative intensity of a powder X-ray diffraction pattern may vary slightly depending on the direction of crystal growth, particle size, and measurement conditions, and thus should not be strictly understood.

[0023]

Numerical values obtained from various patterns may have some errors depending on the direction of crystal growth, particle size, measurement conditions, and the like. Therefore, in the present description, the numerical value of a diffraction angle (2θ) in a powder X-ray diffraction pattern may have a measurement error in a range of approximately $\pm 0.2^\circ$.

[0024]

In the present description, "room temperature" generally means from approximately 10°C to approximately 35°C .

[0025]

In addition, the endothermic peak in a simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve generally means a value with a measurement error in a range of approximately $\pm 5.0^\circ\text{C}$ because the measurement temperature may vary depending on the range of temperature increase per minute, the chemical purity of a sample, and the like. Therefore, when the crystal according to the present invention is measured (TG-DTA), an error of the peak (peak top value) of $\pm 5.0^\circ\text{C}$ is taken into consideration. The term "around" used in such case means $\pm 5.0^\circ\text{C}$.

[0026]

One embodiment of the present invention relates to a free-form crystal (type I crystal or type II crystal) of the compound (I), a crystal (type I crystal, type II crystal, type III crystal, type IV crystal, or type V crystal) of the compound (I) with 1 equivalent of fumaric acid, and a crystal (type I crystal or type II crystal) of the compound (I) with 0.5 equivalents of fumaric acid.

[0027]

(Type II crystal of compound (I) with 1 equivalent of fumaric acid)

A type II crystal of compound (I) with 1 equivalent of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 1, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 2.

In addition, in one embodiment of the present invention, the type II crystal of compound (I) with 1 equivalent of fumaric acid has the diffraction angle (2θ) and intensity (cts) shown in Table 2 in powder X-ray diffraction.

[0028]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type

II crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 5.5° , 6.8° , 9.3° , 13.4° , 15.3° , 16.3° , 18.5° , 19.8° , 22.0° , and 24.5° .

The type II crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, still more preferably a crystal having six or more peaks selected from the characteristic peaks, still more preferably a crystal having seven or more peaks selected from the characteristic peaks, even more preferably a crystal having eight or more peaks selected from the characteristic peaks, still even more preferably a crystal having nine or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the peaks.

[0029]

The endothermic peak (peak top value) of the type II crystal of compound (I) with 1 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 173°C to 183°C , in other words, around 178°C .

[0030]

(Free-form type II crystal of compound (I))

A type II crystal of compound (I) has the powder X-ray diffraction spectrum shown in Figure 3, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 4.

In addition, in one embodiment of the present invention, the free-form type II crystal of compound (I) has the diffraction angle (2θ) and intensity (cts) shown in Table 3 in powder X-ray diffraction.

[0031]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type II crystal of compound (I) may have diffraction angles ($2\theta \pm 0.2^\circ$) of 8.3° , 14.8° , 17.3° , 18.0° , 19.1° , 20.3° , 21.0° , 22.5° , 23.0° , and 26.2° .

The type II crystal of compound (I) according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, still more preferably a crystal having six or more peaks selected from the characteristic peaks, still more preferably a crystal having seven or more peaks selected from the characteristic peaks, even more preferably a crystal having eight or more peaks selected from the characteristic peaks, still even more preferably a crystal having nine or more peaks

selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

[0032]

The endothermic peak (peak top value) of the type II crystal of compound (I) determined by simultaneous thermogravimetry-differential thermal analysis may be 177°C to 187°C, in other words, around 182°C.

[0033]

(Free-form type I crystal of compound (I))

A type I crystal of compound (I) has the powder X-ray diffraction spectrum shown in Figure 5, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 6.

In addition, in one embodiment of the present invention, the free-form type I crystal of compound (I) has the diffraction angle (2θ) and intensity (cts) shown in Table 4 in powder X-ray diffraction.

[0034]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type I crystal of compound (I) may have diffraction angles ($2\theta \pm 0.2^\circ$) of 9.9° , 11.7° , 13.2° , 17.7° , 18.1° , 18.8° , and 20.8° .

The type I crystal of compound (I) according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, still more preferably a crystal having six or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type I crystal of compound (I) may have diffraction angles ($2\theta \pm 0.2^\circ$) of 9.9° , 11.7° , 13.2° , 18.8° , and 20.8° .

The type I crystal of compound (I) according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

[0035]

The endothermic peak (peak top value) of the type I crystal of compound (I) determined by simultaneous thermogravimetry-differential thermal analysis may be 178°C to 188°C, in other words, around 183°C.

[0036]

(Type V crystal of compound (I) with 1 equivalent of fumaric acid)

A type V crystal of compound (I) with 1 equivalent of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 7, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 8.

In addition, in one embodiment of the present invention, the type V crystal of compound (I) with 1 equivalent of fumaric acid has the diffraction angle (2θ) and intensity (cts) shown in Table 5 in powder X-ray diffraction.

[0037]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type V crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 6.9° , 9.4° , 10.2° , 13.7° , 21.1° , 23.6° , and 26.5° .

The type V crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, still more preferably a crystal having six or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type V crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 6.9° , 13.7° , 21.1° , 23.6° , and 26.5° .

The type V crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

[0038]

The endothermic peak (peak top value) of the type V crystal of compound (I) with 1 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 146°C to 156°C , in other words, around 151°C .

[0039]

(Type I crystal of compound (I) with 1 equivalent of fumaric acid)

A type I crystal of compound (I) with 1 equivalent of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 9, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 10.

In addition, in one embodiment of the present invention, the type I crystal of

compound (I) with 1 equivalent of fumaric acid has the diffraction angle (2θ) and intensity (cts) shown in Table 6 in powder X-ray diffraction.

[0040]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type V crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 6.4° , 10.3° , 12.8° , 15.0° , 20.7° , 23.4° , and 26.6° .

The type I crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, still more preferably a crystal having five or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

The endothermic peak (peak top value) of the type I crystal of compound (I) with 1 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 148°C to 158°C , in other words, around 153°C .

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type V crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 6.4° , 10.3° , 12.8° , 20.7° , 23.4° , and 26.6° .

The type I crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

The endothermic peak (peak top value) of the type I crystal of compound (I) with 1 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 148°C to 158°C , in other words, around 153°C .

[0041]

(Type I crystal of compound (I) with 0.5 equivalents of fumaric acid)

A type I crystal of compound (I) with 0.5 equivalents of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 11, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 12.

[0042]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type I crystal of compound (I) with 0.5 equivalents of fumaric acid may have diffraction angles

($2\theta \pm 0.2^\circ$) of 6.4° , 7.5° , 9.7° , 11.7° , 15.1° , 19.6° , and 23.9° .

The type I crystal of compound (I) with 0.5 equivalents of fumaric acid according to the present invention is a crystal having four or more peaks selected from the above characteristic peaks. It is preferably a crystal having five or more peaks selected from the characteristic peaks, more preferably a crystal having six or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

[0043]

The endothermic peak (peak top value) of the type I crystal of compound (I) with 0.5 equivalents of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 156°C to 166°C , in other words, around 161°C .

[0044]

(Type II crystal of compound (I) with 0.5 equivalents of fumaric acid)

A type II crystal of compound (I) with 0.5 equivalents of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 13, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 14.

[0045]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type II crystal of compound (I) with 0.5 equivalents of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 5.4° , 6.4° , 7.3° , 12.8° , 13.4° , 14.7° , and 15.4° .

The type II crystal of compound (I) with 0.5 equivalents of fumaric acid according to the present invention is a crystal having four or more peaks selected from the above characteristic peaks. It is preferably a crystal having five or more peaks selected from the characteristic peaks, more preferably a crystal having six or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

[0046]

The endothermic peak (peak top value) of the type II crystal of compound (I) with 0.5 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 157°C to 167°C , in other words, around 162°C .

[0047]

(Type III crystal of compound (I) with 1 equivalents of fumaric acid)

A type III crystal of compound (I) with 1 equivalents of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 15, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 16.

[0048]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type III crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 8.1° , 13.2° , 20.4° , 23.1° , 24.7° , and 26.1° .

The type III crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having three or more peaks selected from the above characteristic peaks. It is preferably a crystal having four or more peaks selected from the characteristic peaks, more preferably a crystal having five or more peaks selected from the characteristic peaks, and particularly preferably a crystal having all of the characteristic peaks.

The exothermic peak (peak top value) of the type III crystal of compound (I) with 1 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 148°C to 158°C , in other words, around 153°C .

[0049]

(Type IV crystal of compound (I) with 1 equivalent of fumaric acid)

A type IV crystal of compound (I) with 1 equivalent of fumaric acid has the powder X-ray diffraction spectrum shown in Figure 17, and also have the simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curve shown in Figure 18.

[0050]

Here, characteristic peaks in the powder X-ray diffraction spectrum of the type IV crystal of compound (I) with 1 equivalent of fumaric acid may have diffraction angles ($2\theta \pm 0.2^\circ$) of 6.0° , 15.7° , and 18.8° .

The type IV crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention is a crystal having all of the above characteristic peaks.

The exothermic peak (peak top value) of the type IV crystal of compound (I) with 1 equivalent of fumaric acid determined by simultaneous thermogravimetry-differential thermal analysis may be 125°C to 135°C , in other words, around 130°C .

[0051]

(Method for producing type II crystal with 1 equivalent of fumaric acid)

In one embodiment of the present description, a type II crystal of compound (I) with 1 equivalent of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of adding fumaric acid and the following solvent to a type I crystal of compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step to obtain a crystal of compound (I) with 1 equivalent of fumaric acid.

Thereafter, the resulting solid was recovered by filtration. It is preferable that

the recovered solid is washed with water and dried.

In step 1, preferably 1.5 to 10 equivalents (more preferably 2 to 5 equivalents and most preferably 3 equivalents) of fumaric acid is added with respect to 1 equivalent of a free-form compound (I).

The solvent in step 1 is added in an amount of preferably 5 to 35 times (v/w) (more preferably 5 to 30 times (v/w) and most preferably 10 times (v/w)).

Examples of the solvent used in step 1 may include single solvents or mixed solvents of alcohols such as methanol, ethanol, isopropanol, and tert-butanol, ethers such as 1,4-dioxane and tetrahydrofuran, ketones such as acetone and methyl ethyl ketone, and aprotic polar solvents such as acetonitrile.

A type I crystal of compound (I) can be obtained in the section of " (Method for producing free-form type I crystal)" or the method described later in Example 3.

The temperature during the reaction in step 2 is preferably room temperature or higher and lower than the boiling point of each solvent as described above, but is not particularly limited and can be appropriately set to preferably 25°C to 70°C (more preferably 40°C to 60 ° C and most preferably 50°C).

The stirring time during the reaction in step 2 is preferably 24 to 124 hours (more preferably 48 to 72 hours and most preferably 60 to 65 hours) for stirring the suspension.
[0052]

(Method for producing free-form type II crystal)

(Method A-1)

In one embodiment of the present description, a free-form type II crystal of compound (I) can be obtained by a method comprising, for example, the following two steps (Method A-1).

Step 1: Step of adding any one of succinic acid, phosphoric acid, and adipic acid and a mixed solvent of good solvent/poor solvent to the compound (I), and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step to obtain a crystal of compound (I) in a solid form.

During suspension and stirring described above, an additional solvent (for example, 1-propanol as a good solvent) may be added in a plurality of times as necessary, depending on the amount of evaporation of the solvent.

Thereafter, the resulting solid was recovered by filtration and dried.

Preferably 0.5 to 2 equivalents (more preferably 0.75 to 1.5 equivalents and particularly preferably 1 equivalent) of the acid in step 1 is added with respect to 1 equivalent of the compound (I).

The mixed solvent of good solvent/poor solvent in step 1 (e.g., 1-propanol/water) is added in an amount of preferably 5 to 50 times (v/w) (more preferably 10 to 30 times (v/w) and particularly preferably 20 times (v/w)) the amount of the free-form compound (I).

The temperature during the reaction in step 2 is preferably 25°C to 70°C (more preferably 40°C to 60°C and particularly preferably 50°C).

The stirring time during the reaction in step 2 is preferably 2 to 7 days (more preferably 3 to 5 days and particularly preferably 4 days) for stirring the suspension.

[0053]

(Method A-2)

In another embodiment of the present description, a free-form type II crystal of compound (I) can be obtained by a method comprising, for example, the following two steps (Method A-2).

Step 1: Step of adding a mixed solvent of good solvent/poor solvent and a seed crystal to the compound (I), and optionally adding any one of succinic acid, phosphoric acid, and adipic acid.

Step 2: Step of stirring the suspension obtained in the above step to obtain a crystal of compound (I) in a solid form.

During suspension and stirring described above, an additional solvent (for example, 1-propanol as a good solvent) may be added in a plurality of times as necessary, depending on the amount of evaporation of the solvent.

Thereafter, the resulting solid was recovered by filtration and dried.

Preferably 0.5 to 2 equivalents (more preferably 0.75 to 1.5 equivalents and particularly preferably 1 equivalent) of succinic acid in step 1 is added with respect to 1 equivalent of the compound (I).

The mixed solvent of good solvent/poor solvent in step 1 (e.g., a mixed solvent of 1-propanol/water) is added in an amount of preferably 1 to 10 times (v/w) (more preferably 2 to 5 times (v/w) and particularly preferably 3 times (v/w)) the amount of the free-form compound (I), and a seed crystal (e.g., a seed crystal obtained in either (i) or (ii)) is added thereto.

The reaction temperature in step 2 is preferably 25°C to 70°C (more preferably 40°C to 60°C and particularly preferably 50°C).

The reaction time in step 2 is preferably 6 to 48 hours (more preferably 12 to 24 hours and most preferably 17.5 to 24 hours).

[0054]

(Method B)

In another embodiment of the present description, a free-form type II crystal of compound (I) can be obtained by a method comprising, for example, the following two steps (Method B).

Step 1: Step of adding a poor solvent to the compound (I) and optionally adding a seed crystal.

Step 2: Step of adding a good solvent to the solution obtained in the above step and stirring the resulting suspension to obtain a crystal of compound (I) in a solid form.

The poor solvent (e.g., diisopropyl ether) in step 1 is added in an amount of preferably 2 to 10 times (v/w) (more preferably 3 to 7 times (v/w) and particularly preferably 5 times (v/w)) the amount of the free-form compound (I).

The reaction time in step 1 is preferably 24 to 96 hours (more preferably 36 to 72 hours and particularly preferably 44.5 hours) for suspension and stirring.

The reaction temperature in step 2 is preferably 25°C to 70°C (more preferably 40°C to 60°C and particularly preferably 50°C).

The good solvent in step 2 (e.g., ethanol) is added in an amount of preferably 0.5 to 3 times (v/w) (more preferably 0.8 to 2 times (v/w) and particularly preferably 1 time (v/w)) the amount of the free-form compound (I).

The stirring time in step 2 is preferably 6 to 72 hours (more preferably 12 to 48 hours and particularly preferably 21 hours), following which the solid is recovered by filtration and dried.

During suspension and stirring described above, an additional solvent (e.g., diisopropyl ether as a poor solvent and/or ethanol as a good solvent) may be added in a plurality of times as necessary according to the amount of evaporation of the solvent.

[0055]

Examples of a combination of a good solvent and a poor solvent used in the production of a free form II crystal of compound (I) include methanol (good solvent) and water (poor solvent), methanol (good solvent) and diisopropyl ether (IPE) (poor solvent), methanol (good solvent) and heptane (poor solvent), ethanol (good solvent) and IPE (poor solvent), ethanol (good solvent) and heptane (poor solvent), 1-propanol (good solvent) and water (poor solvent), 1-propanol (good solvent) and IPE (poor solvent), 1-propanol (good solvent) and heptane (poor solvent), 2-propanol (good solvent) and heptane (poor solvent), acetone (good solvent) and IPE (poor solvent), acetone (good solvent) and heptane (poor solvent), dimethyl sulfoxide (DMSO) (good solvent) and water (poor solvent), dimethylacetamide (DMA) (good solvent) and water (poor solvent), tetrahydrofuran (THF) (good solvent) and IPE (poor solvent), and THF (good solvent) and heptane (poor solvent).

Of these, a preferred combination of a good solvent and a poor solvent is ethyl acetate (good solvent) and n-heptane (poor solvent), ethanol (good solvent) and water (poor solvent), 1-propanol (good solvent) and water (poor solvent), or ethanol (good solvent) and diisopropyl ether (poor solvent). The amount of the poor solvent is preferably 1 to 20 times (v/v) and more preferably 1 to 10 times (v/v) of the good solvent.
[0056]

In the above method, the amount of the seed crystal added can be preferably 0.5% to 30% (w/w), more preferably 1% to 5% (w/w) of the amount of the introduced compound (I).
[0057]

(Method for producing free-form type I crystal)

In one embodiment of the present description, a free-form type I crystal of compound (I) can be obtained by a method comprising, for example, a step of suspending a crude product of the compound (I) in a solvent and optionally adding a seed crystal to obtain a crystal of compound (I) in a solid form.

The seed crystal is optionally added in order to promote the crystallization of the free-form type I crystal, and as the seed crystal, an appropriate amount of a type I crystal of compound (I) or a mixed crystal containing the type I crystal may be added. A type I crystal of compound (I) can be obtained by the above method without adding a seed crystal. However, by adding a seed crystal, the time for obtaining the compound can be shortened. Further, crystallization may be carried out with stirring in order to shorten the crystallization time and control the particle size.

The seed crystal to be added is 0.5% to 30% (w / w) and preferably 1% to 10% (w / w) of the amount of the introduced compound (I).

The temperature can be appropriately set, but is preferably 50°C to 65°C.

The compound (I) can be precipitated at the above dissolution temperature. However, if it is not precipitated at the dissolution temperature, a free-form type I crystal can be obtained by cooling to 25°C.

[0058]

As the solvent, ethyl acetate, ethanol, acetonitrile, acetone, tert-butyl methyl ether, or the like can be used.

[0059]

(Method for producing type V crystal with 1 equivalent of fumaric acid)

In one embodiment of the present description, a type V crystal of compound (I) with 1 equivalent of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of to the compound (I), adding 1 to 2 equivalents of fumaric acid and acetone

as a solvent in an amount 40 to 60 times (v/w) with respect to 1 equivalent of the free-form compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step, filtrating the suspension, and recovering the solid.

Step 3: Step of drying the solid obtained in the above step under reduced pressure conditions (preferably a pressure of 2 kPa or less) at preferably 20°C to 60°C to obtain a crystal of compound (I) with 1 equivalent of fumaric acid.

Fumaric acid in step 1 is more preferably added in an amount of 2 equivalents to 1 equivalent of the free-form compound (I).

Acetone in step 1 is preferably added in an amount 40 to 60 times (v/w) (more preferably 50 times (v/w)) with respect to the free-form compound (I).

The reaction temperature in step 2 is preferably 10°C to 30°C (preferably 25°C).

The reaction time in step 2 is preferably 12 to 96 hours (more preferably 71 hours).

The reaction temperature of step 3 is 20°C to 60°C (e.g., room temperature) for the obtained solid.

The reduced pressure time in step 3 (e.g., a pressure of 2 kPa or less) include drying for preferably 6 to 24 hours (more preferably 7.5 hours).

[0060]

In one embodiment of the present description, a type V crystal of compound (I) with 1 equivalent of fumaric acid can be obtained by a method comprising, for example, the following step.

Step 1: Step of drying a type I crystal with 1 equivalent of fumaric acid under reduced pressure conditions (preferably at a pressure of 2 kPa or less).

The reaction temperature in step 1 is 20°C to 60°C.

[0061]

(Method for producing type I crystal with 1 equivalent of fumaric acid)

In one embodiment of the present description, a type I crystal of compound (I) with 1 equivalent of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of to the compound (I), adding 1 to 2 equivalents of fumaric acid and acetone as a solvent in an amount 40 to 60 times (v/w) with respect to 1 equivalent of the free form compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step at 30°C or less to obtain a crystal of compound (I) with fumaric acid.

Thereafter, the resulting solid was recovered by filtration.

Fumaric acid in step 1 is more preferably added in an amount of 2 equivalents to 1 equivalent of the free-form compound (I).

Acetone in step 1 is preferably added in an amount 40 to 60 times (more preferably 50 times) with respect to the free-form compound (I).

The reaction temperature in step 2 is preferably 10°C to 30°C (preferably 25°C).

The reaction time in step 2 is preferably 12 to 24 hours (more preferably 19.5 hours).

[0062]

(Method for producing type III crystal with 1 equivalent of fumaric acid)

In one embodiment of the present description, a type III crystal of compound (I) with 1 equivalent of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of adding fumaric acid and acetonitrile as a solvent to the compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step in a short time of 1.5 hours or less to obtain a crystal of compound (I) with 1 equivalent of fumaric acid.

Thereafter, the resulting solid was recovered by filtration.

To 1 equivalent of the free-form compound (I) in step 1, preferably 1 to 5 equivalents (more preferably 3 equivalents) of fumaric acid is added.

The solvent (e.g., acetonitrile) in step 1 is preferably added in an amount 25 to 35 times (v/w) (more preferably 30 times (v/w)) with respect to the free-form compound (I).

The reaction temperature in step 2 is preferably 45°C to 55°C (particularly preferably 50°C).

The reaction time in step 2 is preferably 0.5 to 1.5 hours (more preferably 1 hour).

[0063]

(Method for producing type IV crystal with 1 equivalent of fumaric acid)

In one embodiment of the present description, a type IV crystal of compound (I) with 1 equivalent of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of adding fumaric acid and water as a solvent to the compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step in a short time of 2 hours or less to obtain a crystal of compound (I) with 1 equivalent of fumaric acid.

Thereafter, the resulting solid was recovered by filtration.

To 1 equivalent of the free-form compound (I), preferably 1 to 5 equivalents

(more preferably 3 equivalents) of fumaric acid is added in step 1.

Water as the solvent in step 1 is added in an amount preferably 15 to 25 times (more preferably 20 times) with respect to the free-form compound (I).

The reaction temperature in step 2 is preferably 45°C to 55°C (particularly preferably 50°C).

The reaction time in step 2 is preferably 1 to 2 hours (more preferably 1.5 hour).

[0064]

(Method for producing type I crystal with 0.5 equivalent of fumaric acid)

In one embodiment of the present description, a type I crystal of compound (I) with 0.5 equivalents of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of adding fumaric acid and water as a solvent to the compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step at a temperature of preferably room temperature or higher and lower than the boiling point of the solvent to obtain a crystal.

Thereafter, the resulting solid was recovered by filtration and dried.

In step 1, preferably 0.5 to 1 equivalent (more preferably 0.5 to 0.75 equivalents and most preferably 0.5 equivalents) of fumaric acid is added with respect to 1 equivalent of the free-form compound (I).

Water as the solvent in step 1 is added in an amount of preferably 10 to 40 times (v/w) (more preferably 15 to 25 times (v/w) and most preferably 20 times (v/w)) with respect to the free-form compound (I).

The reaction temperature in step 2 is preferably 40°C to 60°C (more preferably 45°C to 55°C and particularly preferably 50°C).

The reaction time in step 2 is preferably 48 to 120 hours (more preferably 72 to 96 hours and most preferably 93.5 hours).

[0065]

(Method for producing type II crystal with 0.5 equivalents of fumaric acid)

In one embodiment of the present description, a type II crystal of compound (I) with 0.5 equivalents of fumaric acid can be obtained by a method comprising, for example, the following two steps.

Step 1: Step of adding fumaric acid and ethanol to the compound (I) and optionally adding a seed crystal.

Step 2: Step of stirring the suspension obtained in the above step at a temperature of preferably room temperature or higher and lower than the boiling point of the solvent to

obtain a crystal.

Thereafter, the resulting solid was recovered by filtration and dried.

To 1 equivalent of the free-form compound (I), preferably 0.5 to 1 equivalent (more preferably 1 equivalent) of fumaric acid is added in step 1.

Ethanol as the solvent in step 1 is added in an amount of preferably 10 to 40 times (v/w) (more preferably 15 to 25 times (v/w) and most preferably 20 times (v/w)) with respect to the free-form compound (I).

The reaction temperature in step 2 is preferably 40°C to 60°C (more preferably 45°C to 55°C and particularly preferably 50°C).

The reaction time in step 2 is preferably 6 to 48 hours (more preferably 12 to 24 hours and most preferably 19.5 hours).

[0066]

A type II crystal with 1 equivalent of fumaric acid, a free-form type II crystal, a free-form type I crystal, a type V crystal with 1 equivalent of fumaric acid, a type I crystal with 0.5 equivalents of fumaric acid, and a type II crystal with 0.5 equivalents of fumaric acid of the compound (I) have at least one of advantageous properties in pharmaceutical manufacturing such as a non-hygroscopic property, ability to be obtained with reproducibility, solid stability, and oral absorbability.

Of these, a type II crystal with 1 equivalent of fumaric acid, a free-form type II crystal, a free-form type I crystal, and a type V crystal with 1 equivalent of fumaric acid of the compound (I) have advantageous properties in pharmaceutical manufacturing such as a non-hygroscopic property, ability to be obtained with reproducibility, solid stability, and oral absorbability, as compared with other crystal forms of the compound (I).

[0067]

The type II crystal of compound (I) with 1 equivalent of fumaric acid is less hygroscopic and has ability to be obtained with stability. It is important for the industrial manufacturing of pharmaceutical products that drug development candidate compounds are less hygroscopic can be stably obtained. In addition, it has properties that are easy to handle as pharmaceutical products, such as solid stability and solubility, and excellent oral absorbability. Therefore, the type II crystal of compound (I) with 1 equivalent of fumaric acid has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0068]

The free-form type II crystal of compound (I) is less hygroscopic. It is important for the industrial manufacturing of pharmaceutical products having stable quality with drug development candidate compounds. In addition, it has properties that

are easy to handle as pharmaceutical products, such as solid stability and solubility, and excellent oral absorbability. Therefore, the free-form crystal of compound (I) has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0069]

The free-form type I crystal of compound (I) is less hygroscopic and has excellent ability to be obtained with stability. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds are less hygroscopic and have ability to be stably obtained. Therefore, the type I crystal of compound (I) has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0070]

The type V crystal of compound (I) with 1 equivalent of fumaric acid is less hygroscopic and has ability to be obtained with stability. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds are less hygroscopic and have ability to be stably obtained. Therefore, the type V crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention have excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0071]

The type I crystal of compound (I) with 1 equivalent of fumaric acid has excellent ability to be stably obtained. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds have ability to be stably obtained. Therefore, the type I crystal of compound (I) with 1 equivalent of fumaric acid has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

Since the type I crystal of compound (I) with 1 equivalent of fumaric acid can be recovered as a solid form, it is also useful as an intermediate for the production of a type V crystal with 1 equivalent of fumaric acid.

[0072]

The type III crystal of compound (I) with 1 equivalent of fumaric acid has excellent ability to be stably obtained. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds have ability to be stably obtained. Therefore, the type III crystal of compound (I) with 1 equivalent of fumaric acid has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0073]

The type IV crystal of compound (I) with 1 equivalent of fumaric acid has excellent ability to be stably obtained. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds have ability to be stably obtained. Therefore, the type IV crystal of compound (I) with 1 equivalent of fumaric acid has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0074]

The type I crystal of compound (I) with 0.5 equivalents of fumaric acid has excellent ability to be stably obtained. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds have ability to be stably obtained. Therefore, the type I crystal of compound (I) with 0.5 equivalents of fumaric acid has excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0075]

The type II crystal of compound (I) with 0.5 equivalents of fumaric acid has excellent ability to be stably obtained. It is also important for the industrial manufacturing of pharmaceutical products with stable quality that drug development candidate compounds have ability to be stably obtained. Therefore, the type II crystal of compound (I) with 0.5 equivalents of fumaric acid according to the present invention have excellent properties required as a pharmaceutical product or active pharmaceutical ingredient.

[0076]

(Crystals with other acids)

Another embodiment of the present invention relates to a crystal of compound (I) with hydrochloric acid (type I crystal, type II crystal, or type III crystal), a crystal of compound (I) with hydrobromic acid, a crystal of compound (I) with 1 equivalent of L-tartaric acid, or a crystal of compound (I) with succinic acid.

The powder X-ray diffraction spectra and simultaneous thermogravimetry-differential thermal analysis (TG-DTA) curves for these crystals are shown in Figures 19 to 30.

Characteristic peaks in the powder X-ray diffraction spectra of these crystals are described in Reference Examples 3 to 7. These crystals are crystals having three or more (preferably four or more, and if present, more preferably 5 or more) peaks selected from characteristic peaks described in the Reference Examples, and in particular, they are crystals having all of the characteristic peaks.

In addition, endothermic peak temperatures determined by simultaneous

thermogravimetry-differential thermal analysis for these crystals are also described in the Reference Examples.

[0077]

Regarding free-form crystals of the compound (I) and crystals of the compound (I) with an acid (salt crystal or co-crystal) as described above, a precipitated crystal can be isolated and purified from a solution in which the crystal is dissolved, mixed, or the like by a known separation and purification means such as filtration, washing with water, and drying under reduced pressure.

[0078]

The free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention has excellent HER2 inhibitory activity. Moreover, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention has excellent selectivity to HER2. Accordingly, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention is useful as an antitumor agent against diseases or malignant tumor having HER2 overexpression, HER2 gene amplification, HER2 mutation, etc. In addition, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention is advantageous in that it has a few side effects.

In the present description, the term "HER2" includes the HER2 of a human or a non-human mammal, and it is preferably human HER2. Furthermore, the term "HER2" includes isoforms.

Since the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention has excellent HER2 inhibitory activity, it is useful as a medicament for treating disease associated with HER2.

The "disease associated with HER2" means disease, in which a reduction in the incidence, or the remission, alleviation and/or complete recovery of the symptoms thereof is achieved by deleting, suppressing and/or inhibiting the function of HER2. Examples of such disease may include malignant tumors, but are not limited thereto. Preferred examples of the disease may include malignant tumors having HER2 overexpression, HER2 gene amplification, or HER2 mutation.

[0079]

The free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention selectively inhibits wild-type HER2, and mutant HER2 having one or more insertion mutations, point mutations, deletion mutations, etc. in the HER2 domain thereof, such as exon 20 insertion mutation.

One embodiment of the present invention provides: the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) having inhibitory activity against wild-type HER2, and mutant HER2 including HER2 having YVMA insertion mutation that is one of exon 20 insertion mutations, or a salt thereof; or a medicament or a pharmaceutical composition each comprising the same.

One embodiment of the present invention provides an inhibitor against wild-type HER2, and mutant HER2 including HER2 having YVMA insertion mutation, etc., wherein the inhibitor comprises the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention.

[0080]

The human HER2 gene is shown in, for example, SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5. The wild-type HER2 protein consists of the amino acid sequence set forth in, for example, SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. The nucleotide sequence information of the human HER2 gene and the amino acid sequence information of the wild-type HER2 protein can be obtained from, for example, Accession No. NM_004448, NM_001289936, NM_001005862, or the like.

[0081]

In several embodiments, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention exhibits inhibitory activity against mutant HER2 comprising one or more mutations from G309A, S310F, R678Q, L755S, L755_T759del, D769H, A775_G776insYVMA, V777L, V842I and R896C, using the amino acid sequence set forth in SEQ ID NO: 2 as a reference. In another embodiment, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention exhibits inhibitory activity against mutant HER2 comprising A775_G776insYVMA, using the amino acid sequence set forth in SEQ ID NO: 2 as a reference.

In several embodiments, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention exhibits inhibitory activity against mutant HER2 comprising one or more mutations from G294A, S295F, R663Q, L740S, L740_T744del, D754H, A760_G761insYVMA, V762L, V827I and R881C, using the amino acid sequence set forth in SEQ ID NO: 4 as a reference. In another embodiment, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention exhibits inhibitory activity against mutant HER2 comprising A760_G761insYVMA, using the amino acid sequence set forth in SEQ ID NO: 4 as a reference.

In several embodiments, the free-form crystals or crystals with acids (salt

crystals or co-crystals) of the compound (I) of the present invention exhibits inhibitory activity against mutant HER2 comprising one or more mutations from G279A, S280F, R648Q, L725S, L725_T729del, D739H, A745_G746insYVMA, V747L, V812I and R866C, using the amino acid sequence set forth in SEQ ID NO: 6 as a reference. In another embodiment, the free-form crystals or crystals with acids (salt crystals or co-crystals) of the compound (I) of the present invention exhibits inhibitory activity against mutant HER2 comprising A745_G746insYVMA, using the amino acid sequence set forth in SEQ ID NO: 6 as a reference.

[0082]

Further, in several embodiments, with regard to a mutation in a certain HER2 isoform, even when the position of the mutation is different from the position of an amino acid shown in SEQ ID NO: 2 due to deletion or insertion of an amino acid(s), it is understood that the mutation is the same as the mutation at a position corresponding to the position of the amino acid shown in SEQ ID NO: 2. Hence, for example, the glycine at position 309 in the HER2 shown in SEQ ID NO: 2 corresponds to glycine at position 294 in HER2 consisting of the amino acid sequence set forth in SEQ ID NO: 4. For example, the term "G309A" means that the glycine at position 309 in the HER2 shown in SEQ ID NO: 2 is mutated to alanine. Since such "G309" is at a position corresponding to the amino acid at position 294 in HER2 consisting of the amino acid sequence set forth in SEQ ID NO: 4, "G294A" in the HER2 consisting of the amino acid sequence set forth in SEQ ID NO: 4 corresponds to "G309A" in the HER2 shown in SEQ ID NO: 2. Besides, the position of an amino acid in SEQ ID NO: 2 that corresponds to a certain amino acid in a certain HER2 isoform can be confirmed by Multiple Alignment of BLAST.

[0083]

Sequence Listing

SEQ ID NO: 1

Accession No.: NM_004448

CDS: 262..4029

```
1 gcttgctccc aatcacagga gaaggaggag gtggaggagg agggctgctt gaggaagtat
61 aagaatgaag ttgtgaagct gagattcccc tccattggga ccggagaaac caggggagcc
121 ccccgggcag ccgcgcgccc cttcccacgg ggccctttac tgcgccgcgc gcccgcccc
181 caccctcgc agcaccgccg gccccgcgcc ctcccagccg ggtccagccg gagccatggg
241 gccggagccg cagtgaacac catggagctg gcggccttgt gccgctgggg gtcctcctc
301 gccctcttgc cccccggagc cgcgagcacc caagtgtgca ccggcacaga catgaagctg
361 cggtctcctg ccagtcccga gaccacactg gacatgtctc gccacctcta ccagggtgc
421 caggtggtgc agggaaacct ggaactcacc tacctgcca ccaatgccag cctgtccttc
481 ctgcaggata tccaggaggt gcagggtac gtgtcatcg ctcaacca agtgaggcag
541 gtccactgc agaggctgcg gattgtgcga ggcaccagc tctttgagga caactatgcc
601 ctggccgtgc tagacaatgg agaccgctg aacaatacca cccctgtcac aggggcctcc
661 ccaggaggcc tgcgggagct gcagcttca agcctcacag agatcttgaa aggaggggtc
721 ttgatccagc ggaaccccc gctctgtac caggacacga tttgtggaa ggacatcttc
781 cacaagaaca accagctggc tctcacactg atagacacca accgctctcg ggcctgccac
841 ccctgttctc cgatgtgtaa gggctccgc tgtggggag agagtctga ggattgtcag
901 agcctgacgc gactgtctg tgccggtggc tgtgccgct gcaaggggcc actgccact
961 gactgtgcc atgagcagtg tgtgccggc tgcacgggc ccaagcactc tgactgcctg
1021 gcctgcctcc acttcaacca cagtggcatc tgtgagctgc actgccagc cctggtcacc
1081 tacaacacag acacgtttga gtccatgcc aatcccgagg gccggtatac attcggcgcc
1141 agctgtgtga ctgcctgtcc ctacaactac ctttctacgg acgtgggatc ctgcaccctc
1201 gtctgcccc tgcacaacca agaggtgaca gcagaggatg gaacacagcg gtgtgagaag
1261 tgcagcaagc cctgtgccc agtgtgtat ggtctgggca tggagcactt gcgagagggtg
1321 agggcagtta ccagtgccaa tatccaggag tttgtggct gcaagaagat ctttgggagc
1381 ctggcatttc tgccggagag ctttgatggg gaccagcct ccaacactgc cccgtccag
1441 ccagagcagc tccaagtgtt tgagactctg gaagagatca caggttacct atacatctca
1501 gcatggccgg acagcctgcc tgacctcagc gtcttccaga acctgcaagt aatccgggga
1561 cgaattctgc acaatggcgc ctactcgtg accctgcaag ggctgggcat cagctggctg
1621 gggctgcgct cactgaggga actgggcagt ggactggccc tcatccacca taacaccac
1681 ctctgcttcg tgcacacggt gccctgggac cagctctttc ggaaccgca ccaagctctg
1741 ctccacactg ccaaccggcc agaggacgag tgtgtgggcg agggcctggc ctgccaccag
```

```

1801 ctgtgcgccc gagggcactg ctgggggtcca gggcccaccc agtgtgtcaa ctgcagccag
1861 ttccttcggg gccaggagtg cgtggaggaa tgccgagtac tgcaggggct ccccgaggag
1921 tatgtgaatg ccaggcactg tttgccgtgc caccctgagt gtcagcccca gaatggctca
1981 gtgacctgtt ttggaccgga ggctgaccag tgtgtggcct gtgcccacta taaggacctt
2041 cccttctgcg tggcccgtg ccccgagcgt gtgaaacctg acctctccta catgcccata
2101 tggaaagtttc cagatgagga gggcgcatgc cagccttgcc ccatcaactg caccacttcc
2161 tgtgtggacc tggatgacaa gggctgcccc gccgagcaga gagccagccc tctgacgtcc
2221 atcatctctg cggtggttgg cattctgctg gtcgtggtct tgggggtggt ctttgggata
2281 ctcataagc gacggcagca gaagatccgg aagtacacga tgcggagact gctgcaggaa
2341 acggagctgg tggagccgct gacacctagc ggagcgtatc ccaaccaggc gcagatgcgg
2401 atcctgaaag agacggagct gaggaagggt aagggtgctt gatctggcgc ttttggcaca
2461 gtctacaagg gcatctggat ccctgatggg gagaatgtga aaattccagt ggccatcaaa
2521 gtgttgaggg aaaacacatc ccccaaagcc aacaaagaaa tcttagacga agcatatcgt
2581 atggctggtg tgggctcccc atatgtctcc cgccttctgg gcatctgcct gacatccacg
2641 gtgcagctgg tgacacagct tatgccctat ggctgcctct tagaccatgt ccgggaaaac
2701 cgcggacgcc tgggctccca ggacctgctg aactgggtga tgcagattgc caaggggatg
2761 agctacctgg aggatgtgcg gctcgtagac agggacttgg ccgctcggaa cgtgctggtc
2821 aagagtccca accatgtcaa aattacagac ttcgggctgg ctggtgctgt ggacattgac
2881 gagacagagt accatgcaga tgggggcaag gtgcccatac agtggatggc gctggagtcc
2941 attctccgcc ggcggttcac ccaccagagt gatgtgtgga gttatggtgt gactgtgtgg
3001 gagctgatga cttttggggc caaaccttac gatgggatcc cagcccggga gatccctgac
3061 ctgctggaag agggggagcg gctgccccag ccccccattt gcaccattga tgtctacatg
3121 atcatggtca aatgttggat gattgactct gaatgtcggc caagattccg ggagttggtg
3181 tctgaattct cccgcatggc cagggacccc cagcgctttg tggcatcca gaatgaggac
3241 ttggggccag ccagtccctt ggacagcacc ttctaccgct cactgctgga ggacgatgac
3301 atgggggacc tggtggtatg tgaggagtat ctggtacccc agcagggtt cttctgtcca
3361 gacctgtccc cgggcgctgg gggcatggtc caccacaggc accgcagctc atctaccagg
3421 agtggcgggtg gggacctgac actagggctg gagccctctg aagaggaggc cccaggtctt
3481 cactggcac cctccgaagg ggctggctcc gatgtatttg atggtgacct gggaatgggg
3541 gcagccaagg ggctgcaaag cctccccaca catgaccca gccctctaca gcggtacagt
3601 gaggaccca cagtaccctt gccctctgag actgatggct acgttgcccc cctgacctgc
3661 agccccagc ctgaatatgt gaaccagcca gatgttcggc cccagcccc ttcgccccga
3721 gagggccctc tgcctgtctc ccgacctgct ggtgccactc tggaaaggcc caagactctc
3781 tccccaggga agaattgggt cgtcaaagac gtttttgctt ttgggggtgc cgtggagaa
3841 cccgagtact tgacaccca gggaggagct gccctcagc cccaccttcc tctgccttc

```

```

3901 agcccagcct tgcacaacct ctattactgg gaccaggacc caccagagcg gggggctcca
3961 cccagcacct tcaaagggac acctacggca gagaaccag agtacctggg tctggacgtg
4021 ccagtgtgaa ccagaaggcc aagtccgcag aagccctgat gtgtcctcag ggagcagggg
4081 aggcctgact tctgctggca tcaagagggt ggagggccct ccgaccactt ccaggggaac
4141 ctgccatgcc aggaacctgt cctaaggaac cttccttcct gcttgagttc ccagatggct
4201 ggaaggggtc cagcctcggt ggaagaggaa cagcactggg gagtctttgt ggattctgag
4261 gccctgccc atgagactct aggggtccagt ggatgccaca gccagcttg gccctttcct
4321 tccagatcct ggggtactgaa agccttaggg aagctggcct gagaggggaa gcggccctaa
4381 gggagtgtct aagaacaaaa gcgacccatt cagagactgt ccctgaaacc tagtactgcc
4441 ccccatgagg aaggaacagc aatgggtgtca gtatccaggc tttgtacaga gtgcttttct
4501 gtttagtttt tacttttttt gttttgtttt tttaaagatg aaataaagac ccagggggag
4561 aatgggtgtt gtatggggag gcaagtgtgg ggggtccttc tccacacca cttgtccat
4621 ttgcaaatat attttggaac acagctaaaa aaaaaaaaaa aaaa

```

SEQ ID NO: 2

Accession No.: NM_004448

```

MELAALCRWG LLLALLPPGA ASTQVCTGTD MKLRLPASPE THLDMLRHLY QGCQVVQGNL 60
ELTYLPTNAS LSFLQDIQEV QGYVLIAHNQ VRQVPLQRLR IVRGTQLFED NYALAVLDNG 120
DPLNNTTPVT GASPGLREL QLRSLTEILK GGVLIQRNPQ LCYQDTILWK DIFHKNNQLA 180
LTLIDTNRSR ACHPCSPMCK GSRCWGESSE DCQSLTRTVC AGGCARCKGP LPTDCCHEQC 240
AAGCTGPKHS DCLACLHFNH SGICELHCPA LVTYNTDTFE SMPNPEGRYT FGASCVTACP 300
YNYLSTDVGS CTLVCPHNPQ EVTAEDGTQR CEKSKPCAR VCYGLGMEHL REVRAVTSAN 360
IQEFAGCKKI FGSLAFLPES FDGPASNTA PLQPEQLQVF ETLEEITGYL YISAWPDSLP 420
DLSVFQNLQV IRGRILHNGA YSLTLQGLGI SWLGLRSLRE LGSGLALHH NTHLCFVHTV 480
PWDQLFRNPH QALLHTANRP EDECVGEGLA CHQLCARGHC WPGGPTQCVN CSQFLRGQEC 540
VEECRVLQGL PREYVNARHC LPCHPECQPQ NSVTCFGPE ADQCVACAHY KDPPFCVARC 600
PSGVKPDLSY MPIWKFPDEE GACQPCPINC THSCVDLDDK GCPAEQRASP LTSIISAVVG 660
ILLVVVLGVV FGILIKRRQQ KIRKYTMRR LQETELVEPL TPGAMPNQA QMRILKETEL 720
RKVKVLGSGA FGTVYKGIWI PDGENVKIPV AIKVLRENTS PKANKEILDE AYVMAGVGSP 780
YVSRLGICL TSTVQLVTQL MPYGCLLDHV RENRGLGSQ DLLNWCMIQIA KGMSYLEDVR 840
LVHRDLAARN VLVKSPNHVK ITDFGLARLL DIDETEHAD GGKVPKWM LESILRRRFT 900
HQSDVWSYGV TWELMTFGA KPYDGIPARE IPDLLEKGER LPQPPICTID VYMIMVKCWM 960
IDSECRPRFR ELVSEFSRMA RDPQRFVVIQ NEDLGASPL DSTFYRSLLE DDDMGDLVDA 1020
EEYLVPQQGF FCPDPAPGAG GMVHHRHRSS STRSGGDLT LGLEPSEEEA PRSPLAPSEG 1080

```

AGSDVFDGDL	GMGAAKGLQS	LPTHDP SPLQ	RYSEDPTVPL	PSETDGYVAP	LTCSPQPEYV	1140
NQPDVRPQPP	SPREGPLPAA	RPAGATLERP	KTLSPGKNGV	VKDVFAFGGA	VENPEYLTPQ	1200
GGAAPQPHPP	PAFSPAFDNL	YYWDQDPPER	GAPPSTFKGT	PTAENPEYLG	LDVPV	1255
SEQ ID NO: 3 Accession No.: NM_001289936 CDS: 583..4305						
1 aagttcctgt gttctttatt ctactctccg ctgaagtcca cacagtttaa attaaagttc 61 ccggattttt gtgggcgcct gccccgcccc tcgtccccct gctgtgtcca tatatcgagg 121 cgataggggtt aaggggaaggc ggacgcctga tgggttaatg agcaaactga agtggttttc 181 atgatctttt ttgagtcgca attgaagtac cacctcccga gggtgattgc ttccccatgc 241 ggggtagaac ctttgctgtc ctgttcacca ctctacctcc agcacagaat ttggcttatg 301 cctactcaat gtgaagatga tgaggatgaa aacctttgtg atgatccact tccacttaat 361 gaatggtggc aaagcaaagc tatattcaag accacatgca aagctactcc ctgagcaaag 421 agtcacagat aaaacggggg caccagtaga atggccagga caaacgcagt gcagcacaga 481 gactcagacc ctggcagcca tgcctgcgca ggcagtgatg agagtgcacat gtactgttgt 541 ggacatgcac aaaagtgaga tacttcaaag attccagaag atatgccccg ggggtcctgg 601 aagccacaag tgtgcaccgg cacagacatg aagctgcggc tccctgccag tcccagagacc 661 cacctggaca tgctccgcca cctctaccag ggctgccagg tgggtgcagg aaacctggaa 721 ctcacctacc tgcccaccaa tgccagcctg tccttctctg aggatatcca ggaggtgcag 781 ggctacgtgc tcatcgctca caaccaagtg aggcagggtcc cactgcagag gctgcggatt 841 gtgcgaggca cccagctctt tgaggacaac tatgccctgg ccgtgctaga caatggagac 901 ccgctgaaca ataccacccc tgtcacaggg gcctccccag gaggcctgcg ggagctgcag 961 cttcgaagcc tcacagagat cttgaaagga ggggtcttga tccagcggaa cccccagctc 1021 tgctaccagg acacgatttt gtggaaggac atcttccaca agaacaacca gctggctctc 1081 aactgatag acaccaaccg ctctcggggc tgccaccctt gttctccgat gtgtaagggc 1141 tcccgtgct ggggagagag ttctgaggat tgtcagagcc tgacgcgcac tgtctgtgcc 1201 ggtggctgtg cccgctgcaa ggggccactg cccactgact gctgcatga gcagtgtgct 1261 gccggctgca cgggccccaa gcactctgac tgcttggcct gcctccactt caaccacagt 1321 ggcatctgtg agctgcactg cccagccctg gtcacctaca acacagacac gtttgagtcc 1381 atgccaatc ccgagggccg gtatacatc gccgccagct gtgtgactgc ctgtccctac 1441 aactaccttt ctacggacgt gggatcctgc accctcgtct gccccctgca caaccaagag 1501 gtgacagcag aggatggaac acagcgggtg gagaagtgca gcaagccctg tgcccagagt 1561 tgctatggtc tgggcatgga gcacttgca gaggtgaggg cagttaccag tgccaatatc						

1621 caggagtttg ctggctgcaa gaagatcttt gggagcctgg catttctgcc ggagagcttt
1681 gatggggacc cagcctccaa cactgccccg ctccagccag agcagctcca agtgtttgag
1741 actctggaag agatcacagg ttacctatac atctcagcat ggccggacag cctgcctgac
1801 ctcagcgtct tccagaacct gcaagtaatc cggggacgaa ttctgcacaa tggcgcttac
1861 tcgctgacct tgcaagggct gggcatcagc tggctggggc tgcgctcact gaggggaactg
1921 ggcagtggac tggccctcat ccaccataac acccacctct gcttcgtgca cacggtgccc
1981 tgggaccagc tctttcgga cccgcaccaa gctctgctcc aactgcca ccggccagag
2041 gacgagtgtg tgggcgaggg cctggcctgc caccagctgt gcgcccaggg gcactgctgg
2101 ggtccagggc ccaccagtg tgtcaactgc agccagttcc ttcggggcca ggagtgcgtg
2161 gaggaatgcc gactactgca ggggtcccc agggagtatg tgaatgccag gcactgtttg
2221 ccgtgccacc ctgagtgtca gccccagaat ggctcagtga cctgttttgg accggaggct
2281 gaccagtgtg tggcctgtgc ccactataag gacctccct tctgcgtggc ccgtgcccc
2341 agcgggtgtg aacctgacct ctctacatg cccatctgga agtttccaga tgaggagggc
2401 gcatgccagc ctgccccat caactgcacc cactcctgtg tggacctgga tgacaagggc
2461 tgccccgccg agcagagagc cagccctctg acgtccatca tctctgcggt ggttggcatt
2521 ctgctggtcg tggctttggg ggtggtcttt gggatcctca tcaagcgacg gcagcagaag
2581 atccggaagt acacgatgcg gagactgctg caggaaacgg agctggtgga gccgtgaca
2641 cctagcggag cgatgccc aaaggcgag atgcggatcc tgaaagagac ggagtgtagg
2701 aaggtgaagg tgcttgatc tggcgctttt ggcacagtct acaagggcat ctggatccct
2761 gatggggaga atgtgaaaat tccagtggcc atcaaagtgt tgagggaaaa cacatcccc
2821 aaagccaaca aagaaatctt agacgaagca tacgtgatgg ctggtgtggg ctccccatat
2881 gtctccgcc ttctgggcat ctgcctgaca tccacggtgc agctggtgac acagcttatg
2941 ccctatggct gcctcttaga ccatgtccgg gaaaaccgag gacgcctggg ctcccaggac
3001 ctgctgaact ggtgtatgca gattgccaag gggatgagct acctggagga tgtgcggctc
3061 gtacacaggg acttgccgc tcggaacgtg ctggtcaaga gtcccaacca tgtcaaaatt
3121 acagacttcg ggctggctcg gctgctggac attgacgaga cagagtacca tgcagatggg
3181 ggcaagggtc ccatcaagtg gatggcgctg gactccattc tccgccggcg gttcaccac
3241 cagagtgatg tgtggagtta tgggtgact gtgtgggagc tgatgacttt tggggccaaa
3301 ccttacgatg ggtacccagc ccgggagatc cctgacctgc tggaaaaggg ggagcggctg
3361 cccagcccc ccatctgcac cattgatgtc tacatgatca tgggtcaaat ttggatgatt
3421 gactctgaat gtcggccaag attccgggag ttggtgtctg aattctccg catggccagg
3481 gacccccagc gctttgtggt catccagaat gaggacttgg gccagccag tcccttggac
3541 agcaccttct accgctcact gctggaggac gatgacatgg gggacctggt ggtatgctgag
3601 gagtatctgg taccacagca gggcttcttc tgtccagacc ctgccccggg cgctgggggc
3661 atggtccacc acaggcaccg cagctcatct accaggagtg gcggtgggga cctgacacta

```

3721 gggctggagc cctctgaaga ggaggccccc aggtctccac tggcacccctc cgaaggggct
3781 ggctccgatg tatttgatgg tgacctggga atggggggcag ccaaggggct gcaaagcctc
3841 cccacacatg accccagccc tctacagcgg tacagtgagg accccacagt acccctgccc
3901 tctgagactg atggctacgt tgccccctg acctgcagcc cccagcctga atatgtgaac
3961 cagccagatg ttcggcccca gcccccttcg ccccagagagg gccctctgcc tgctgcccga
4021 cctgctgggtg ccactctgga aaggcccaag actctctccc cagggaagaa tggggtcgtc
4081 aaagacgttt ttgcctttgg gggtgccgtg gagaaccccg agtacttgac accccagggg
4141 ggagctgccc ctcagcccca ccctcctcct gccttcagcc cagccttcga caacctctat
4201 tactgggacc aggaccacc agagcggggg gctccacca gcaccttcaa agggacacct
4261 acggcagaga acccagagta cctgggtctg gacgtgccag tgtgaaccag aaggccaagt
4321 ccgcagaagc cctgatgtgt cctcagggag cagggaaggc ctgacttctg ctggcatcaa
4381 gaggtgggag ggccctccga ccacttccag gggaacctgc catgccagga acctgtccta
4441 aggaaccttc ctctctgctt gagttcccg atggctggaa ggggtccagc ctcgttgga
4501 gaggaacagc actggggagt ctttgtggat tctgaggccc tgcccaatga gactctaggg
4561 tccagtggat gccacagccc agcttgggcc ttccttcca gatcctgggt actgaaagcc
4621 ttagggaagc tggcctgaga ggggaagcgg ccctaaggga gtgtctaaga acaaaagcga
4681 cccattcaga gactgtccct gaaacctagt actgcccccc atgaggaagg aacagcaatg
4741 gtgtcagtat ccaggctttg tacagagtgc ttttctgttt agtttttact tttttgttt
4801 tgttttttta aagatgaaat aaagaccag ggggagaatg ggtgttgat ggggaggcaa
4861 gtgtgggggg tccttctcca caccacttt gtccatttgc aaatatattt tggaaaacag
4921 ctaaaaaaaaa aaaaaaaaaa

```

SEQ ID NO: 4

Accession No.: NM_001289936

```

MPRGSWKPQV CTGDMKLRL PASPETHLDM LRHLYQGCQV VQGNLELYL PTNASLSFLQ 60
DIQEVQGYVL IAHNQVRQVP LQRLRIVRGT QLFEDNYALA VLDNGDPLNN TTPVTGASPG 120
GLRELQLRSL TEILKGGVLI QRNPQLCYQD TILWKDIFHK NNQLALTLID TNRSRACHPC 180
SPMCKGSRCW GESSEDCQSL TRTVCAGGCA RCKGPLPTDC CHEQCAAGCT GPKHSDCLAC 240
LHFNHSGICE LHCPALVTYN TDTFESMPNP EGRYTFGASC VTACPYNLYS TDVGSCTLVC 300
PLHNQEVTA E DGTQRCEKCS KPCARVCYGL GMEHLREVRA VTSANIQEFA GCKKIFGSLA 360
FLPESFDGDP ASNTAPLQPE QLQVFETLEE ITGYLYISAW PDSLPLDSVF QNLQVIRGRI 420
LHNGAYSLTL QGLGISWLGL RSLRELGSGL ALIHHNTHLC FVHTVPWDQL FRNPHQALLH 480
TANRPEDECV GEGLACHQLC ARGHCWGP GP TQCVNCSQFL RGQECVEECR VLQGLPREYV 540
NARHCLPCHP ECQPQNGSVT CFGPEADQCV ACAHYKDPPF CVARCPSGVK PDLSYMPIWK 600

```

FPDEEGACQP	CPINCTHSCV	DLDDKGCPAE	QRASPLTSII	SAVVGILLV	VLGVVFGILI	660
KRRQKIRKY	TMRLLQETE	LVEPLTPSGA	MPNQAQMRIL	KETELRKVKV	LGSGAFGTVY	720
KGIWIPDGEN	VKIPVAIKVL	RENTSPKANK	EILDEAYVMA	GVGSPYVSRL	LGICLTSTVQ	780
LVTQLMPYGC	LLDHVRENRG	RLGSQDLLNW	CMQIAKGMSY	LEDVRLVHRD	LAARNVLVKS	840
PNHVKITDFG	LARLLDIDET	EYHADGGKVP	IKWMALESIL	RRRFTHQSDV	WSYGVTVWEL	900
MTFGAKPYDG	IPAREIPDLL	EKGERLPQPP	ICTIDVYMIM	VKCWMIDSEC	RPRFRELVSE	960
FSRMARDPQR	FVVIQNEDLG	PASPLDSTFY	RSLLDDDDMG	DLVDAEEYLV	PQQGFFCPDP	1020
APGAGGMVHH	RHRSSSTRSG	GGDLTLGLEP	SEEEAPRSPL	APSEGAGSDV	FDGDLGMGAA	1080
KGLQSLPTH	PSPLQRYSED	PTVPLPSETD	GYVAPLTCSP	QPEYVNQPDV	RPQPPSPREG	1140
PLPAARPAGA	TLERPKTLSP	GKNGVVKDVF	AFGGAVENPE	YLTPQGGAAP	QPHPPPAFSP	1200
AFDNLYYWDQ	DPPERGAPPS	TFKGTPTAEN	PEYLGLDVPV			1240

SEQ ID NO: 5

Accession No.: NM_001005862

CDS: 577..4254

1 aagttcctgt gttctttatt ctactctccg ctgaagtcca cacagttaa attaaagttc
61 ccggattttt gtgggcgcct gccccgcccc tcgtccccct gctgtgtcca tatatcgagg
121 cgataggggtt aaggggaaggc ggacgcctga tgggttaatg agcaaactga agtgttttcc
181 atgatctttt ttgagtcgca attgaagtac cacctcccga gggtgattgc ttccccatgc
241 ggggtagaac ctttgctgtc ctgttcacca ctctacctcc agcacagaat ttggcttatg
301 cctactcaat gtgaagatga tgaggatgaa aacctttgtg atgatccact tccacttaat
361 gaatgggtggc aaagcaaagc tatattcaag accacatgca aagctactcc ctgagcaaag
421 agtcacagat aaaacggggg caccagtaga atggccagga caaacgcagt gcagcacaga
481 gactcagacc ctggcagcca tgacctgcga ggcagtgatg agagtgacat gtactgttgt
541 ggacatgcac aaaagtgagt gtgcaccggc acagacatga agctgcggct ccctgccagt
601 cccgagaccc acctggacat gctccgccac ctctaccagg gctgccaggt ggtgcaggga
661 aacctggaac tcacctacct gccaccaat gccagcctgt ccttcctgca ggatatccag
721 gaggtgcagg gctacgtgct catcgctcac aaccaagtga ggcagggtccc actgcagagg
781 ctgcggattg tgcgaggcac ccagctcttt gaggacaact atgccctggc cgtgctagac
841 aatggagacc cgctgaacaa taccaccctt gtcacagggg cctccccagg aggctgcgg
901 gagctgcagc ttcgaagcct cacagagatc ttgaaaggag gggctttgat ccagcggaac
961 cccagctctt gctaccagga cacgattttg tggaaggaca tcttcacaa gaacaaccag
1021 ctggctctca cactgataga caccaaccgc tctcgggcct gccaccctg ttctccgatg
1081 tgtaagggct cccgctgctg gggagagagt tctgaggatt gtcagagcct gacgcgcact

1141 gtctgtgccg gtggctgtgc ccgctgcaag gggccactgc cactgactg ctgccatgag
1201 cagtgtgctg ccggctgcac gggccccaag cactctgact gcctggcctg cctccacttc
1261 aaccacagtg gcatctgtga gctgcaactgc ccagccctgg tcacctacaa cacagacacg
1321 tttgagtcca tgcccaatcc cgaggggccg tatacattcg gcgccagctg tgtgactgcc
1381 tgtccctaca actacctttc tacggacgtg ggatcctgca ccctcgtctg cccctgcac
1441 aaccaagagg tgacagcaga ggatggaaca cagcgggtgtg agaagtgcag caagccctgt
1501 gcccagagtgt gctatggctt gggcatggag cacttgcgag aggtgagggc agttaccagt
1561 gccaatatcc aggagtttgc tggctgcaag aagatctttg ggagcctggc atttctgccg
1621 gagagctttg atggggaccc agcctccaac actgccccgc tccagccaga gcagctccaa
1681 gtgtttgaga ctctggaaga gatcacaggt tacctataca tctcagcatg gccggacagc
1741 ctgcctgacc tcagcgtctt ccagaacctg caagtaatcc ggggacgaat tctgcacaat
1801 ggcgcctact cgtgaccct gcaagggctg ggcatcagct ggctggggct gcgctcactg
1861 agggaaactgg gcagtggact ggccctcatc caccataaca cccacctctg ctctgtgcac
1921 acggtgccct gggaccagct ctttcggaac ccgcaccaag ctctgctcca cactgccaac
1981 cgccagagg acgagtgtgt gggcgagggc ctggcctgcc accagctgtg cgcccagagg
2041 cactgctggg gtccagggcc caccagtggt gtcaactgca gccagttcct tcggggccag
2101 gagtgcgtgg aggaatgccg agtactgcag gggctcccca gggagtatgt gaatgccagg
2161 cactgtttgc cgtgccacc tgagtgtcag cccagaatg gctcagtgac ctgttttgga
2221 ccggaggctg accagtgtgt ggctgtgcc cactataagg accctccctt ctgcgtggcc
2281 cgtgccccca gcggtgtgaa acctgacctc tcctacatgc ccatctggaa gttccagat
2341 gaggaggcg catgccagcc ttgccccatc aactgcaccc actcctgtgt ggacctggat
2401 gacaagggct gcccgcgga gcagagagcc agccctctga cgtccatcat ctctgcggtg
2461 gttggcattc tgtggctgt ggtcttgggg gtggtctttg ggatcctcat caagcgacgg
2521 cagcagaaga tccggaagta cacgatgcgg agactgctgc aggaaacgga gctggtggag
2581 ccgctgacac ctagcggagc gatgcccac caggcgcaga tgcggatcct gaaagagacg
2641 gagctgagga aggtgaaggt gcttggatct ggcgcttttg gcacagtcta caagggcatc
2701 tggatccctg atggggagaa tgtgaaaatt ccagtggcca tcaaagtgtt gagggaaaac
2761 acatcccca aagccaacaa agaaatctta gacgaagcat acgtgatggc tgggtgtggc
2821 tccccatatg tctccgcct tctgggcatc tgctgacat ccacggtgca gctggtgaca
2881 cagcttatgc cctatggctg cctcttagac catgtccggg aaaaccgcg acgcctgggc
2941 tccaggacc tgctgaactg gtgtatgcag attgccaagg ggatgagcta cctggaggat
3001 gtgcggctcg tacacagga cttggccgct cggaacgtgc tggtaagag tccaacat
3061 gtcaaaatta cagacttcgg gctggctcgg ctgctggaca ttgacgagac agagtacat
3121 gcagatgggg gcaaggtgcc catcaagtgg atggcgctgg agtccattct ccgccggcg
3181 ttcaccacc agagtgatgt gtggagttat ggtgtgactg tgtgggagct gatgactttt

```

3241 ggggccaaac cttacgatgg gatcccagcc cgggagatcc ctgacctgct ggaaaagggg
3301 gagcggtgc cccagcccc catctgcacc attgatgtct acatgatcat ggtcaaattgt
3361 tggatgattg actctgaatg tcggccaaga ttccgggagt tgggtgtctga attctccgc
3421 atggccaggg acccccagcg ctttgtggtc atccagaatg aggacttggg cccagccagt
3481 cccttggaca gcaccttcta ccgctcactg ctggaggacg atgacatggg ggacctggtg
3541 gatgctgagg agtatctggt accccagcag ggcttcttct gtccagacct tggccgggc
3601 gctgggggca tgggccacca caggcaccgc agctcatcta ccaggagtgg cggtggggac
3661 ctgacactag ggctggagcc ctctgaagag gaggcccca ggtctccact ggcacctcc
3721 gaaggggctg gctccgatgt atttgatggt gacctgggaa tgggggcagc caaggggctg
3781 caaagcctcc ccacacatga cccagccct ctacagcggg acagtgagga cccacagta
3841 ccctgcccct ctgagactga tggctacgtt gccccctga cctgcagccc ccagcctgaa
3901 tatgtgaacc agccagatgt tcggccccag ccccttcgc cccgagaggg ccctctgcct
3961 gctgcccgc ctgctggtgc cactctggaa aggcccaaga ctctctcccc agggaagaat
4021 ggggtcgta aagacgtttt tgcctttggg ggtgccgtgg agaacccca gtacttgaca
4081 cccaggggag gagctgcccc tcagccccac cctcctcctg cttcagccc agccttcgac
4141 aacctctatt actgggacca ggaccacca gagcgggggg ctccaccag caccttcaa
4201 gggacaccta cggcagagaa cccagagtac ctgggtctgg acgtgccagt gtgaaccaga
4261 aggccaaagtc cgcagaagcc ctgatgtgtc ctgaggagc aggaaggcc tgacttctgc
4321 tggcatcaag aggtgggagg gccctccgac cacttccagg ggaacctgcc atgccaggaa
4381 cctgtcctaa ggaaccttcc ttcctgcttg agttcccaga tggctggaag ggtccagcc
4441 tcgttggaag aggaacagca ctggggagtc tttgtggatt ctgaggccct gcccaatgag
4501 actctagggg ccagtggatg ccacagccca gcttggccct ttccttcag atcctgggta
4561 ctgaaagcct tagggaagct ggctgagag ggaagcggc cctaagggag tgtctaagaa
4621 caaaagcgac ccattcagag actgtccctg aaacctagta ctgccccca tgaggaagga
4681 acagcaatgg tgtcagtatc caggctttgt acagagtgt tttctgttta gtttttactt
4741 tttttgtttt gtttttttaa agatgaaata aagaccagc gggagaatgg gtgttgtatg
4801 gggaggcaag tgtggggggg cttctccac acccacttg tccatttgca aatatatatt
4861 ggaaaacagc taaaaaaaaa aaaaaaaaaa

```

SEQ ID NO: 6

Accession No.: NM_001005862

MKLRLPASPE THLDMLRHLY QGCQVVQGNL ELTYLPTNAS LSFLQDIQEV QGYVLIAHNQ 60
 VRQVPLQRLR IVRGTQLFED NYALAVLDNG DPLNNTTPVT GASPGLREL QLRSLTEILK 120
 GGVLIQRNPQ LCYQDTILWK DIFHKNNQLA LTLIDTNRSR ACHPCSPMCK GSRCWGESSE 180

DCQSLTRTVC	AGGCARCKGP	LPTDCCHEQC	AAGCTGPKHS	DCLACLHFNH	SGICELHCPA	240
LVTYNTDTFE	SMPNPEGRYT	FGASCVTACP	YNYLSTDVGS	CTLVCPLHNQ	EVTAEDEGTQR	300
CEKCSKPCAR	VCYGLGMEHL	REVRAVTSAN	IQEFAGCKKI	FGSLAFLPES	FDGDPASNTA	360
PLQPEQLQVF	ETLEEITGYL	YISAWPDSL	DLSVFQNLQV	IRGRILHNGA	YSLTLQGLGI	420
SWLGLRSLRE	LGSGLALIIH	NTHLCFVHTV	PWDQLFRNPH	QALLHTANRP	EDECVGEGLA	480
CHQLCARGHC	WGPGPTQCVN	CSQFLRGQEC	VEECRVLQGL	PREYVNARHC	LPCHPECQPQ	540
NGSVTCFGPE	ADQCVACAHY	KDPPFCVARC	PSGVKPDLSY	MPIWKFPDEE	GACQPCPINC	600
THSCVDLDDK	GCPAEQRASP	LTSIISAVVG	ILLVVVLGVV	FGILIKRRQQ	KIRKYTMRR	660
LQETELVEPL	TPSGAMPNQA	QMRILKETEL	RKVKVLGSGA	FGTVYKGIWI	PDGENVKIPV	720
AIKVLRENTS	PKANKEILDE	AYVMAGVGSP	YVSRLGICL	TSTVQLVTQL	MPYGCLLDHV	780
RENRLGSLQ	DLNWCMIQA	KGMSYLEDVR	LVHRDLAARN	VLVKSPNHVK	ITDFGLARLL	840
DIDETEHAD	GGKVPIKWMA	LESILRRRFT	HQSDVWSYGV	TVWELMTFGA	KPYDGIPARE	900
IPDLLEKGER	LPQPPICTID	VYMIMVKCWM	IDSECRPRFR	ELVSEFSRMA	RDPQRFVVIQ	960
NEDLGASPL	DSTFYRSLLE	DDMGDLVDA	EEYLVPQQGF	FCPDAPAGAG	GMVHHRHRSS	1020
STRSGGDLT	LGLEPSEEEA	PRSPLAPSEG	AGSDVFDGDL	GMGAAGLQSQ	LPTHDPSPQL	1080
RYSEDPTVPL	PSETDGYVAP	LTCSPQPEYV	NQPDVRPQPP	SPREGPLPAA	RPAGATLERP	1140
KTLSPGKNGV	VKDVFAGGA	VENPEYLTPQ	GGAAPQPHPP	PAFSPAFDNL	YYWDQDPPER	1200
GAPPSTFKGT	PTAENPEYLG	LDVPV				1225

[0084]

One embodiment of the present invention provides an antitumor agent, comprising the above-described crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)). In addition, one embodiment of the present invention provides a method for treating tumor, comprising administering an effective amount of the above-described crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) to a subject in need thereof. Moreover, one embodiment of the present invention provides use of the above-described crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for the production of an antitumor agent. Furthermore, one embodiment of the present invention provides the above-described crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for use in the treatment of tumor.

[0085]

The crystals of the present invention may be used in postoperative adjuvant

chemotherapy performed to prevent recurrence after surgical removal of a tumor, or may be used in neoadjuvant chemotherapy performed in advance before surgical removal of a tumor.

[0086]

The tumor that is the target of the present invention is not particularly limited. Examples of the tumor may include brain tumor, head and neck cancer, digestive cancer (esophageal cancer, stomach cancer, duodenal cancer, liver cancer, biliary tract cancer (gallbladder and/or bile duct cancer, etc.), pancreatic cancer, colorectal cancer (colon cancer, rectal cancer, etc.), etc.), lung cancer (non-small cell lung cancer, small cell lung cancer, mesothelioma, etc.), breast cancer, genital cancer (ovarian cancer, uterine cancer (cervical cancer, endometrial cancer, etc.), etc.), urinary organ cancer (kidney cancer, bladder cancer, prostate cancer, testicular tumor, etc.), hematopoietic tumor (leukemia, malignant lymphoma, multiple myeloma, etc.), bone and/or soft tissue tumor, and skin cancer. Among these, preferable is lung cancer, breast cancer, stomach cancer, colorectal cancer, bladder cancer, biliary tract cancer or uterine cancer, and more preferable is lung cancer, breast cancer, stomach cancer, bladder cancer, or biliary tract cancer.

[0087]

In one embodiment, the tumor is a brain tumor. The compound of the present invention may be useful for the treatment of the symptoms of brain that is required to pass through the blood-brain barrier. The compound of one embodiment has favorable permeability through the blood-brain barrier for the delivery thereof into the brain, namely, excellent brain penetration properties. As an indicator of the penetration properties of the compound into the brain, the concentration of the compound in the brain or a Kp value (brain-to-plasma drug concentration ratio) is applied.

The brain tumor treated with the compound of the present invention includes metastatic brain tumor and primary brain tumor.

Examples of the brain tumor may include, but are not particularly limited to, metastatic brain tumor (e.g., brain metastasis of lung cancer, breast cancer, stomach cancer, colorectal cancer, bladder cancer, biliary tract cancer, uterine cancer, etc. (preferably, lung cancer, breast cancer, or stomach cancer)), pilocytic astrocytoma, diffuse astrocytoma, oligodendroma and/or oligodendroastrocytoma, anaplastic astrocytoma and/or anaplastic oligodendroglioma, anaplastic oligodendroastrocytoma, glioblastoma, ependymoma, anaplastic ependymoma, ganglioglioma, central neurocytoma, medulloblastoma, germinoma, central nervous system malignant lymphoma, meningioma, neurilemmoma, GH secreting pituitary adenoma, PRL-

secreting pituitary adenoma, ACTH-secreting pituitary adenoma, nonfunctional pituitary adenoma, craniopharyngioma, chordoma, hemangioblastoma, and epidermoid tumor.

[0088]

In the present description, the term "effective amount" of the compound means the amount of the compound of the present invention that induces the biological or medical response of a subject, such as, for example, reduction or inhibition of enzyme or protein activity; or ameliorates symptoms, alleviates conditions, and retards or delays the progression of disease; and the like (therapeutically effective amount).

In the present description, the term "subject" includes mammals and non-mammals. Examples of the mammal may include, but are not limited to, a human, a chimpanzee, an ape, a monkey, a bovine, a horse, sheep, a goat, a swine, a rabbit, a dog, a cat, a rat, a mouse, a guinea pig, a hedgehog, a kangaroo, a mole, a wild pig, a bear, a tiger, and a lion. Examples of the non-mammal may include, but are not limited to, birds, fish, and reptiles. In one embodiment, the subject is a human, and may be a human who has been diagnosed to need the treatment for the symptoms, conditions or disease disclosed in the present description.

[0089]

Upon the use of the compound (I) or a salt crystal or co-crystal thereof as a medicament, various types of dosage forms can be adopted depending on the therapeutic purpose with or without crushing the crystal. Examples of the dosage form may include all of oral agents such as tablets, capsules, granules, fine granules, powders, and dry syrups, suppositories, inhalants, nasal drops, ointments, patches, and injections. Pharmaceutical compositions suitable for these dosage forms can be prepared by commonly used production methods that are known to skilled persons using pharmaceutically acceptable carriers.

[0090]

One embodiment of the present invention provides an antitumor agent comprising the above-described crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)). One embodiment of the present invention also provides a method for treating tumor, comprising administering an effective amount of the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) to a subject in need thereof. One embodiment of the present invention also provides the use of the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for the production of an antitumor agent. One embodiment of the present invention also provides the crystal

of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for use in the treatment of tumor by administration thereof.

One embodiment of the present invention provides an antitumor agent for oral administration comprising the above-described crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)). One embodiment of the present invention also provides a method for treating tumor, comprising orally administering an effective amount of the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) to a subject in need thereof. One embodiment of the present invention also provides the use of the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for the production of an antitumor agent for oral administration. One embodiment of the present invention also provides the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for use in the treatment of tumor by oral administration thereof.

[0091]

One embodiment of the present invention provides a pharmaceutical composition comprising the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)). The pharmaceutical composition in one embodiment of the present invention comprises the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) and a pharmaceutically acceptable carrier. One embodiment of the present invention also provides the use of the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for the production of a pharmaceutical composition. One embodiment of the present invention provides the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) for use as a medicament. One embodiment of the present invention provides a kit comprising (i) a pharmaceutical composition comprising the crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) and (ii) an instruction for use of the pharmaceutical composition.

[0092]

As pharmaceutically acceptable carriers, various types of organic or inorganic carrier substances, which are commonly used as preparation materials, are used. When the compound of the present invention is processed into a solid preparation, examples of

the pharmaceutically acceptable carrier mixed into the compound of the present invention may include an excipient, a binder, a disintegrator, a lubricant, and a coating agent. When the compound of the present invention is processed into a liquid preparation, examples of the pharmaceutically acceptable carrier mixed into the compound of the present invention may include a solvent, a solubilizer, a suspending agent, a tonicity agent, a buffer, and a soothing agent. In addition, preparation additives such as an antiseptic, an antioxidant, a coloring agent, a sweetener, and a stabilizer can also be used, as necessary.

[0093]

Examples of the excipient may include lactose, sucrose, D-mannitol, starch, crystalline cellulose, and calcium silicate.

Examples of the binder may include hydroxypropyl cellulose, methyl cellulose, polyvinylpyrrolidone, candy powder, and hypromellose.

Examples of the disintegrator may include sodium starch glycolate, carmellose calcium, croscarmellose sodium, crospovidone, low-substituted hydroxypropyl cellulose, and partially pregelatinized starch.

Examples of the lubricant may include talc, magnesium stearate, sucrose fatty acid ester, stearic acid, and sodium stearyl fumarate.

Examples of the coating agent may include ethyl cellulose, aminoalkyl methacrylate copolymer RS, hypromellose, and sucrose.

Examples of the solvent may include water, propylene glycol, and physiological saline.

Examples of the solubilizer may include polyethylene glycol, ethanol, α -cyclodextrin, macrogol 400, and polysorbate 80.

Examples of the suspending agent may include carrageenan, crystalline cellulose/carmellose sodium, and polyoxyethylene hydrogenated castor oil.

Examples of the tonicity agent may include sodium chloride, glycerin, and potassium chloride.

Examples of the pH adjuster/buffer may include sodium citrate, hydrochloric acid, lactic acid, phosphoric acid, and sodium dihydrogen phosphate.

Examples of the soothing agent may include procaine hydrochloride and lidocaine.

Examples of the antiseptic may include ethyl paraoxybenzoate, cresol, and benzalkonium chloride.

Examples of the antioxidant may include sodium sulfite, ascorbic acid, and natural vitamin E.

Examples of the coloring agent may include titanium oxide, iron sesquioxide, edible blue No. 1, and copper chlorophyll.

Examples of the corrigent may include aspartame, saccharin, sucralose, l-menthol, and mint flavor.

Examples of the stabilizer may include sodium pyrosulfite, sodium edetate, erythorbic acid, magnesium oxide, and dibutylhydroxytoluene.

[0094]

In the case of preparing a solid preparation for oral administration, an excipient, and as necessary, a binder, a disintegrator, a lubricant, a coloring agent, a corrigent, and the like are added to a crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)), and thereafter, a tablet, a coated tablet, a granule, a powder agent, a capsule, and the like can be produced according to ordinary methods.

In the case of preparing an injection, a pH adjuster, a buffer, a stabilizer, a tonicity agent, a local anesthetic, and the like are added to a crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)), and thereafter, subcutaneous, intramuscular, and intravenous injections can be produced according to ordinary methods.

[0095]

The amount of a crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) to be mixed into the above-described each dosage unit form depends on the symptoms of a patient to whom the crystal should be applied, the dosage form, or the like, and thus, the amount of the compound of the present invention is not constant. In general, it is preferable that the applied dose is set to be approximately 0.05 to 1000 mg per dosage unit form in the case of an oral agent, it is set to be approximately 0.1 to 500 mg per dosage unit form in the case of an injection, and it is set to be approximately 1 to 1000 mg per dosage unit form in the case of a suppository or a topical agent in terms of the free-form compound (I).

[0096]

The daily dose of a crystal of compound (I) (i.e., a free-form crystal of compound (I) or a crystal of compound (I) with an acid (salt crystal or co-crystal)) in a drug having the above-described dosage form is different depending on the symptoms, body weight, age, sex and the like of a patient, and thus, it cannot be generally determined. However, the crystal may be administered to an adult (body weight: 50 kg) at a daily dose of generally approximately 0.05 to 5000 mg, and preferably 0.1 to 1000 mg in terms of the free-form compound (I), and it is preferably administered once a day or in 2 to 3 divided

doses.

EXAMPLES

[0097]

Hereinafter, the present invention will be further described specifically with reference to the Examples below. However, these Examples are not intended to limit the scope of the present invention. Although the present invention has been fully described by way of the Examples, it will be appreciated that various changes and/or modifications may be made by skilled persons. Therefore, such changes and/or modifications are included in the invention unless they are outside the scope of the present invention.

[0098]

In the following Examples regarding compounds, "%" indicates weight percent, unless otherwise specified.

[0099]

Various types of reagents used in the Examples were commercially available products, unless otherwise specified. Silica gel chromatography was carried out using Purif-Pack (registered trademark) SI manufactured by MORITEX Corporation, KP-Sil (registered trademark) Silica Prepacked Column manufactured by Biotage Japan Ltd., or HP-Sil (registered trademark) Silica Prepacked Column manufactured by Biotage Japan Ltd. Basic silica gel column chromatography was carried out using Purif-Pack (registered trademark) NH manufactured by MORITEX Corporation or KP-NH (registered trademark) Prepacked Column manufactured by Biotage Japan Ltd. Preparative thin-layer chromatography was carried out using Kieselgel TM60F254, Art. 5744, manufactured by Merck, or NH₂ Silica Gel 60F254 Plate, manufactured by Wako Pure Chemical Industries, Ltd. For NMR spectral measurement, AL400 (400 MHz; JEOL Ltd. (JEOL)), a Mercury400 (400 MHz; Agilent Technologies) type spectrometer or an Inova400 (400 MHz; Agilent Technologies) type spectrometer equipped with a 400 MNMR probe (Protasis) was used. When tetramethylsilane was contained in a heavy solvent, tetramethylsilane was used as an internal reference, and in other cases, an NMR solvent was used as an internal reference for measurement, and the total δ value was shown in ppm.

[0100]

In addition, LCMS spectral measurement was carried out using ACQUITY SQD (quadrupole type) manufactured by Waters under the following conditions.

Column: XSelect CSH C18 (4.6 × 150 mm, 3.5 μ m) manufactured by Waters
MS detection: ESI positive

UV detection: 246nm
 Column temperature: 40°C
 Column flow rate: 1.0 mL/min
 Mobile phase: water/acetonitrile (0.1% formic acid)
 Injection amount: 5 µL
 Sample cooler: 5°C
 Sample concentration: 1 mg/mL
 Gradient (Table 1)

[0101]

[Table 1]

Time (min)	Water	Acetonitrile
0-0.1	95%	5%
0.1-2.1	95%→5%	5%→95%
2.1-3.1	5%	95%

[0102]

Reverse phase preparative HPLC purification was carried out under the following conditions using a preparative system manufactured by WATERS.

Column: A column prepared by connecting YMC-Actus Triart C18 (20 × 50 mm, 5 µm) manufactured by YMC and YMC-Actus Triart C18 (20 × 10 mm, 5 µm) manufactured by YMC was used.

UV detection: 254 nm
 MS detection: ESI positive
 Column flow rate: 25 mL/min
 Mobile phase: water/acetonitrile (0.1% formic acid)
 Injection amount: 0.1-0.5 mL

[0103]

Abbreviations have the following meanings.

s: Singlet

d: Doublet

t: Triplet

dt: Double triplet

m: Multiplet

DMSO-d6: Deuterated dimethyl sulfoxide

CDCl₃: Deuterated chloroform

THF: Tetrahydrofuran

DMA: N,N-dimethylacetamide

NMP: 1-Methyl-2-pyrrolidinone

DMSO: Dimethyl sulfoxide

[0104]

Powder X-ray diffraction analysis

Powder X-ray diffraction analysis was carried out according to any of the following test conditions after lightly pulverizing an appropriate amount of a test substance in an agate mortar as needed.

[0105]

Apparatus: EMPYREAN (Method A) manufactured by PANalytical

Reflection method (concentration method)

Target: Cu

X-ray tube current: 40 mA

X-ray tube voltage: 45 kV

Scanning range: $2\theta = 5.0^\circ$ to 40.0°

Step: $2\theta = 0.0131^\circ$

Average time/step 8.670 s

Scan speed: $0.0015^\circ/\text{s}$

Divergence slit: 1°

Scattering slit: 2.0 mm

Light receiving slit: 8.0 mm

Apparatus: EMPYREAN (Method B) manufactured by PANalytical

Permeation method

Target: Cu

X-ray tube current: 40 mA

X-ray tube voltage: 45 kV

Scanning range: $2\theta = 2.0^\circ$ to 40.0°

Step: $2\theta = 0.0066^\circ$

Average time/step 8.670 s

Scan speed: $0.0008^\circ/\text{s}$

Divergence slit: $1/2^\circ$

Scattering slit: 2.0 mm

Light receiving slit: None

[0106]

Handling of the apparatuses, including data processing, was in accordance with the methods and procedures instructed for each apparatus. Numerical values obtained from various spectra may vary slightly depending on the direction of crystal growth, particle size, analysis conditions, and the like. Therefore, those numerical values should not be understood exactly as they are.

[0107]

For simultaneous thermogravimetry-differential thermal analysis (TG-DTA), analysis was carried out using 2 to 3 mg of a test substance according to the following test conditions.

Apparatus: TG/DTA7200

Manufactured by Hitachi High-Tech Science Corporation

Sample container: Made of aluminum

Temperature increase rate: Temperature was increased from 25°C to 290°C at 10°C/min.

Atmospheric gas: Air (200 mL/min)

Control substance: Empty container

Handling of the apparatuses, including data processing, was in accordance with the methods and procedures instructed for each apparatus.

[0108]

Synthesis Example 1

Synthesis of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I))

Synthesis Example 1 (1)

Synthesis of tert-Butyl (2S,4R)-4-(4-amino-5-iodo-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylpyrrolidine-1-carboxylate

tert-Butyl (2S,4S)-4-hydroxy-2-methylpyrrolidine-1-carboxylate (19.0 g) and 4-chloro-5-iodo-7H-pyrrolo[2,3-d]pyrimidine (13.1 g) were dissolved in THF (190 mL), and the obtained solution was then cooled to 0°C. Thereafter, triphenylphosphine (37.2 g) and diisopropyl azodicarboxylate (28.1 mL) were added to the reaction solution, and the temperature of the mixture was then increased to room temperature, followed by stirring for 1 hour. Thereafter, the reaction mixture was concentrated under reduced pressure, and the obtained residue was then purified by silica gel chromatography (hexane : ethyl acetate) to obtain the corresponding coupling body. The obtained compound was used in the subsequent reaction without being further purified.

The obtained coupling body, THF (114 mL) and ammonia water (114 mL) were

added into a pressure resistant tube, and the obtained mixture was then stirred at 100°C for 14 hours. Thereafter, the reaction mixture was cooled to room temperature, and was then poured into water (285 mL). The thus obtained mixture was stirred at room temperature for 5 hours. Thereafter, the precipitated solid was collected by filtration, was then washed with water, and was then dried to obtain a product of interest (34.5 g).

¹HNMR (CDCl₃)δ: 8.27(s,1H) 7.15(s,1H) 5.55-5.73(m,2H) 5.12-5.25(m,1H) 3.86-4.18(m,2H) 3.43-3.57(m,1H) 2.59-2.69(m,1H) 1.92-2.03(m,1H) 1.48(s,9H) 1.30-1.40(m,3H)

ESI-MS m/z 444 (MH⁺)

[0109]

Synthesis Example 1(2)

Synthesis of 4-Amino-7-((3R,5S)-1-(tert-butoxycarbonyl)-5-methylpyrrolidin-3-yl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxylic acid

The compound of Synthesis Example 1(1) (28.0 g), 10% palladium carbon catalyst (720 mg), NMP (84 mL), methanol (26 mL), and triethylamine (17.6 mL) were added into a pressure resistant tube, followed by carbon monoxide substitution, and the obtained mixture was stirred at 100°C for 2 hours. Thereafter, the reaction mixture was cooled to room temperature, a 2 M sodium hydroxide aqueous solution (79 mL) was then added thereto, and the obtained mixture was then stirred at 80°C for 2 hours. Thereafter, the reaction mixture was cooled to room temperature, was then filtrated through Celite, and was then washed with methanol. Subsequently, methanol in the filtrate was concentrated under reduced pressure. Water was further added, and the water layer was then washed with tert-butyl methyl ether. A 1 M potassium hydrogen sulfate aqueous solution was added to the water layer to adjust the pH to approximately 3. The precipitated solid was collected by filtration, was then washed with water, and was then dried to obtain a product of interest (23.4 g).

¹HNMR (400MHz, DMSO-d₆)δ: 8.14 (s, 1H) 8.08 (s, 1H) 5.16-4.93(m,1H) 4.07-3.79(m,2H) 3.61-3.45(m,1H) 2.53(m,1H) 2.33-2.02(m,1H) 1.42(s,9H) 1.29(d,J = 6.1Hz,3H) ESI-MS m/z 362 (MH⁺)

[0110]

Synthesis Example 1(3)

Synthesis of tert-Butyl(2S,4R)-4-(4-amino-6-bromo-5-(((R)-1-phenylethyl)carbamoyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)-2-methylpyrrolidine-1-carboxylate

The compound of Synthesis Example 1(2) (1.00 g), (R)-(+)-1-phenylethylamine (0.503 g), diisopropylethylamine (1.79 g), and N,N-dimethylformamide (10 mL) were added, and subsequently, HATU (1.58 g) was added. The obtained mixture was stirred

at room temperature overnight. Thereafter, to the reaction mixture, ethyl acetate and a saturated sodium hydrogen carbonate aqueous solution were added, and the obtained mixture was then extracted with ethyl acetate. The gathered organic layer was washed with water, and then with saturated saline. The resultant was dried over anhydrous sodium sulfate, and was then concentrated under reduced pressure. The obtained residue was purified by silica gel chromatography (hexane : acetone) to obtain an amide form (1.53 g). The obtained compound was used in the subsequent reaction without being further purified.

To the amide form (1.53 g), chloroform (15 mL) was added, and the obtained mixture was then cooled to 0°C. Thereafter, N-bromosuccinimide (0.88 g) was added to the reaction mixture, and the obtained mixture was then stirred at 0°C for 1 hour. Thereafter, the reaction mixture was concentrated under reduced pressure, and the obtained residue was purified by silica gel chromatography (hexane : ethyl acetate) to obtain a product of interest (1.39 g).

¹HNMR (CDCl₃)δ: 8.21 (s, 1H) 7.42-7.28(m,5H) 6.97(d,J = 7.3Hz,1H) 5.36-5.29(m,1H) 5.20-5.07(m,1H) 4.30(t,J = 10.3Hz,1H) 4.04-3.72(m,2H) 3.00-2.86(m,1H) 2.38(dt,J = 14.3,6.0Hz,1H) 1.63(d,J = 7.0Hz,3H) 1.53-1.43(m,12H)

ESI-MS m/z 543,545 (MH⁺)

[0111]

Synthesis Example 1(4)

7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-bromo-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide

To the compound of Synthesis Example 1(3) (600 mg), chloroform(3 mL) was added, and the obtained mixture was then cooled to 0°C. Thereafter, trifluoroacetic acid (4.44 g) was added to the reaction mixture, and the thus obtained mixture was then stirred at room temperature for 1 hour. Thereafter, the reaction mixture was concentrated under reduced pressure, and acetonitrile (5 mL) was then added to the residue. The obtained mixture was concentrated under reduced pressure again to obtain an amine form. The obtained compound was used in the subsequent reaction without being further purified.

To the obtained amine form, acetonitrile (3 mL) was added, and the obtained mixture was then cooled to 0°C. Thereafter, acryloyl chloride (99.9 mg) and diisopropylethylamine (713 mg) were added, and the obtained mixture was then stirred at 0°C for 1 hour. Thereafter, the reaction mixture was concentrated under reduced pressure, and the obtained residue was purified by silica gel chromatography (ethyl acetate : methanol) to obtain a product of interest (281 mg).

¹HNMR (CDCl₃)δ: 8.20(d,J = 7.3Hz,1H) 7.42-7.36(m,4H) 7.32-7.28(m,1H) 7.00-

6.94(m,1H) 6.57-6.33(m,2H) 5.76-5.66(m,1H) 5.36-5.29(m, 1H) 5.14-5.08(m,1H) 4.71(t, J = 9.9Hz,0.7H) 4.42-4.23(m,1.6H) 3.83(t, J = 8.6Hz,0.7H) 3.03-2.92(m,1H) 2.60-2.57(m,0.3H) 2.44-2.40(m, 0.7H) 1.64(d, J = 6.6Hz,3H) 1.56(dd, J = 11.7,6.2Hz,3H)

ESI-MS m/z 497,499 (MH^+)

[0112]

Synthesis Example 1(5)

Synthesis of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I))

The compound of Synthesis Example 1(4) (65 mg), dichlorobis(triphenylphosphine)dipalladium (9.2 mg), copper(I) iodide (5.0 mg), cyclopropylacetylene (13.0 mg), triethylamine (39.7 mg), and N,N-dimethylformamide (1.3 mL) were added, and the inside of the reaction system was then substituted with nitrogen. After that, the mixture was stirred at 70°C for 2.5 hours. Thereafter, to the reaction mixture, ethyl acetate and a saturated ammonium chloride aqueous solution were added, and the obtained mixture was then extracted with ethyl acetate. The gathered organic layer was washed with water, and then with saturated saline. The resultant was dried over anhydrous sodium sulfate, and was then concentrated under reduced pressure. The obtained residue was purified by silica gel chromatography (chloroform : methanol) to obtain a product of interest (50 mg).

1H NMR ($CDCl_3$) δ : 8.22(d, J = 5.1Hz,1H) 7.82(d, J = 7.3Hz,1H) 7.43-7.35(m,4H) 7.30(t, J = 6.8Hz,1H) 6.58-6.34(m,2H) 5.77-5.66(m,1H) 5.35-5.20(m,2H) 4.54(t, J = 10.1Hz,0.7H) 4.35-4.25(m,1.6H) 3.88(t, J = 8.8Hz,0.7H) 2.90-2.78(m,1H) 2.65-2.56(m,0.3H) 2.49-2.40(m,0.7H) 1.63(d, J = 7.0Hz,3H) 1.56-1.45(m,4H) 1.03-0.91(m,2H) 0.84-0.69(m,2H)
ESI-MS m/z 483 (MH^+)

[0113]

Example 1

Production of type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with 1 equivalent of fumaric acid

Type II crystals of the compound (I) with fumaric acid were produced by the two methods described below in Production Method 1 and Production Method 2. A free-form type I crystal of compound (I) was synthesized by the same method as in Example 3 described later.

(i) Production Method 1

Fumaric acid (3 equivalents) was added to 30 mg of a free-form type I crystal of

compound (I), 0.3 mL of tert-butanol was added, the suspension was stirred at 25°C for approximately 124 hours, and then the solid was collected by filtration, recovered, and dried to obtain a crystal of interest.

(ii) Production Method 2

Fumaric acid (1.06 g) and 22 mL of acetone were added to 2.20 g of a free-form type I crystal of compound (I), the suspension was stirred at 50°C for approximately 62 hours, naturally cooled for 30 minutes, and then the solid was collected by filtration and recovered. This was washed with 2-propanol, and then the solid was collected by filtration, recovered, and dried to obtain 2.27 g of a crystal of interest.

[0114]

Powder X-ray diffraction spectrum (Method A): See Figure 1. The powder X-ray diffraction spectral data are shown in Table 2.

[Table 2]

Table 2: Powder X-ray diffraction spectral data of type II crystal of compound (I) with 1 equivalent of fumaric acid

Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]
5.5	1065	18.5	1706	27.3	358
6.8	489	19.8	516	27.9	144
8.1	187	20.5	236	28.1	137
9.3	370	22.0	1123	29.0	154
10.8	54	22.6	101	29.9	183
12.1	107	23.3	129	30.9	106
13.4	564	23.8	151	32.0	73
14.0	319	24.5	929	32.9	104
14.6	171	25.1	676	34.3	174
15.3	771	25.5	774	35.1	125
16.3	602	26.2	432	38.2	105
17.1	101	27.2	455	39.1	114

In addition, the characteristic diffraction angles are as follows.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$):

5.5°, 6.8°, 9.3°, 13.4°, 15.3°, 16.3°, 18.5°, 19.8°, 22.0°, and 24.5°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 2.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 182°C

[0115]

Example 2

Production of free-form type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I))

Free-form type II crystals were produced by the four methods described below in Production Methods 1 to 4.

(i) Production Method 1

Succinic acid (1 equivalent) and 0.3 mL of 1-propanol/water (1:3 v/v) were added to 15 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 50°C for 4 days, and then the solid was collected by filtration, recovered, and dried to obtain a crystal of interest.

(ii) Production Method 2

Succinic acid (1 equivalent) and 1.5 mL of 1-propanol/water (1:3 v/v) were added to 300 mg of the compound (I) obtained according to the method in Synthesis Example 1, a small amount of the crystals obtained in (i) was added as a seed crystal, the mixture was suspended and stirred at 50°C for 5 hours, and then 0.3 mL of 1-propanol was added, the mixture was further suspended and stirred at 50°C for 2 hours, and then 0.3 mL of 1-propanol was added, the mixture was suspended and stirred at 50°C for 17 hours, and then the solid was collected by filtration, recovered, and dried to obtain 84.5 mg of a crystal of interest.

(iii) Production Method 3

Succinic acid (1 equivalent) and 3 mL of 1-propanol/water (1:1 v/v) were added to 1000 mg of the compound (I) obtained according to the method in Synthesis Example 1, a small amount of the crystals obtained in (ii) was added as a seed crystal, the mixture was suspended and stirred at 50°C for 17.5 hours, and then the solid was collected by filtration (washing with water during filtration), recovered, and dried to obtain 487.6 mg of a crystal of interest.

(iv) Production Method 4

Diisopropyl ether (5 mL) was added to 1000 mg of the compound (I) obtained according to the method in Synthesis Example 1, a small amount of the crystals obtained in (iii) was added as a seed crystal, and the mixture was suspended and stirred at 50°C.

After 42.5 hours, as the solvent had evaporated, 5 mL of diisopropyl ether was added. The mixture was suspended and stirred at 50°C for 2 hours, and then 0.5 mL of ethanol was added. The mixture was further suspended and stirred at 50°C for 2 hours, and then 0.5 mL of ethanol was added. The mixture was further suspended and stirred at 50°C for 19 hours, and naturally cooled for 30 minutes, and then the solid was collected by filtration, recovered, and dried to obtain 798.8 mg of a crystal of interest.

[0116]

Powder X-ray diffraction spectrum (Method A): See Figure 3. The powder X-ray diffraction spectral data are shown in Table 3.

[Table 3]

Table 3: Powder X-ray diffraction spectral data of free-form type II crystal of compound (I)

Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]
8.3	1844	19.4	154	27.6	218
9.7	179	20.3	952	28.3	100
10.2	531	21.0	963	28.6	101
11.5	248	21.6	271	29.7	374
12.5	660	21.9	520	30.9	113
13.6	440	22.5	957	31.7	261
14.8	2512	23.0	1442	32.3	148
15.8	530	23.6	559	34.0	229
16.0	105	23.9	980	34.6	159
16.5	641	24.7	418	35.3	67
16.9	117	25.0	367	36.4	59
17.3	2130	25.5	208	37.0	123
18.0	945	26.2	1551	37.7	60
18.3	269	26.8	323	38.6	53
19.1	844	27.2	771	39.0	55

In addition, the characteristic diffraction angles are as follows.

[0117]

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$):

8.3°, 14.8°, 17.3°, 18.0°, 19.1°, 20.3°, 21.0°, 22.5°, 23.0°, and 26.2°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 4.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 182°C

[0118]

Example 3

Production of free-form type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I))

A crude product (100 mg) of the compound (I) was suspended in ethanol (500

μL) and stirred at 50°C overnight. The solution was cooled to 25°C and the obtained suspension was filtrated to obtain a free-form type I crystal of compound (I) (31 mg).

[0119]

Powder X-ray diffraction spectrum (Method A): See Figure 5. The powder X-ray diffraction spectral data are shown in Table 4.

[Table 4]

Table 4: Powder X-ray diffraction spectral data of free-form type I crystal of compound (I)

Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]
9.9	623	21.5	481	30.3	208
10.4	278	22.4	437	31.0	72
11.7	1459	23.3	198	31.8	113
12.0	457	23.9	297	32.2	114
13.2	618	24.3	230	32.5	108
14.6	74	24.9	293	33.0	126
15.7	588	25.5	379	33.6	171
16.0	555	26.1	321	34.7	68
17.7	730	27.1	99	35.5	82
18.1	1070	28.0	183	36.9	80
18.8	1539	28.3	545	37.2	55
19.7	120	28.9	391	38.0	50
20.1	382	29.5	168		
20.8	2893	29.8	105		

In addition, the characteristic diffraction angles are as follows.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 9.9°, 11.7°, 13.2°, 17.7°, 18.1°, 18.8°, and 20.8°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 6.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 183°C

[0120]

Example 4

Production of type V crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-

carboxamide (compound (I))with 1 equivalent of fumaric acid

Fumaric acid (481 mg) and 50 mL of acetone were added to 1000 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 25°C for 71 hours, and then the solid was collected by filtration and recovered. This was dried under room temperature and reduced pressure conditions for 7.5 hours to obtain 695.0 mg of a crystal of interest.

[0121]

Powder X-ray diffraction spectrum (Method A): See Figure 7. The powder X-ray diffraction spectral data are shown in Table 5.

[Table 5]

Table 5: Powder X-ray diffraction spectral data of type V crystal of compound (I) with 1 equivalent of fumaric acid

Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]
6.9	180	18.1	201	27.5	235
8.7	184	18.7	278	28.2	140
9.4	470	20.1	319	29.0	103
10.2	450	21.1	481	31.0	92
11.9	197	21.7	215	32.4	65
13.7	219	23.1	343	33.0	54
14.5	199	23.6	573	34.4	74
15.9	350	24.6	206		
17.5	356	26.5	519		

In addition, the characteristic diffraction angles are as follows.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 6.9°, 9.4°, 10.2°, 13.7°, 21.1°, 23.6°, and 26.5°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 8.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 151°C

[0122]

Example 5

Production of type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I))with 1 equivalent of fumaric acid

Fumaric acid (481 mg) and 50 mL of acetone were added to 1000 mg of the

compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 25°C for 19.5 hours, and then the solid was collected by filtration and recovered to obtain 868.8 mg of a crystal of interest.

[0123]

Powder X-ray diffraction spectrum (Method A): See Figure 9. The powder X-ray diffraction spectral data are shown in Table 6.

[Table 6]

Table 6: Powder X-ray diffraction spectral data of type I crystal of compound (I) with 1 equivalent of fumaric acid

Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]	Peak position [° 2θ]	Net intensity [cts]
6.4	637	17.8	129	26.6	931
8.5	145	18.4	290	28.1	112
8.9	115	19.2	145	28.9	164
10.3	461	19.4	151	30.3	70
11.4	321	20.7	938	30.8	75
12.8	279	21.4	195	31.6	77
14.4	172	22.1	146	32.9	138
15.0	393	22.9	451	33.4	61
15.7	326	23.4	690	34.6	57
16.0	422	24.2	157		
16.9	206	24.8	146		

In addition, the characteristic diffraction angles are as follows.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 6.4°, 10.3°, 12.8°, 15.0°, 20.7°, 23.4°, and 26.6°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 10.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 153°C

[0124]

Example 6

Production of type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with 0.5 equivalent of fumaric acid

Fumaric acid (0.5 equivalents) and 0.3 mL of water were added to 15 mg of the

compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 50°C for 93.5 hours, and then the solid was collected by filtration, recovered, and dried to obtain a crystal of interest.

[0125]

Powder X-ray diffraction spectrum (Method B): See Figure 11.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 6.4° , 7.5° , 9.7° , 11.7° , 15.1° , 19.6° , and 23.9°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 12.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 161°C

[0126]

Example 7

Production of type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with 0.5 equivalents of fumaric acid

Fumaric acid (1 equivalent) and 6 mL of ethanol were added to 300 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 50°C for 19.5 hours and naturally cooled for 3 hours, and then the solid was collected by filtration and recovered. This was washed with water, and then the solid was collected by filtration, recovered, and dried to obtain 224.6 mg of a crystal of interest.

[0127]

Powder X-ray diffraction spectrum: See Figure 13.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 5.4° , 6.4° , 7.3° , 12.8° , 13.4° , 14.7° , and 15.4°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 14.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 162°C

[0128]

Example 8 Analysis of molar ratio of compound (I) and fumaric acid

The molar ratio of compound (I) and fumaric acid and the molar ratio of compound (I) and L-tartaric acid were confirmed by $^1\text{H-NMR}$. The measurement results of the NMR spectrum of the type I crystal of compound (I) with 1 equivalent of fumaric acid, the type II crystal of compound (I) with 1 equivalent of fumaric acid, and the type V crystal of compound (I) with 1 equivalent of fumaric acid are shown below. The measurement results of the NMR spectra of the free form I crystal of compound (I) and the free form II crystal of compound (I) are also shown below.

(Free form I crystal of compound (I))

¹HNMR(400MHz,DMSO-d₆)δ: 8.32(brd,J=7.38Hz,1H) 8.14(s,1H) 7.25-7.50(m,5H) 6.54-6.72(m,1H) 6.13-6.23(m,1H) 5.65-5.76(m,1H) 5.12-5.39(m,2H) 4.33-4.40(m,1H) 4.00-4.22(m,2H) 2.44-2.70(m,2H) 1.73(tt,J=4.99,8.27Hz,1H) 1.51(d,J=6.88Hz,3H) 1.36-1.43(m,3H) 0.78-1.03(m,4H)

(Free form II crystal of compound (I))

¹HNMR(400MHz,DMSO-d₆)δ: 8.32(brd,J=7.38Hz,1H) 8.14(s,1H) 7.27-7.46(m,5H) 6.54-6.72(m,1H) 6.13-6.23(m,1H) 5.65-5.76(m,1H) 5.12-5.39(m,2H) 4.32-4.40(m,1H) 4.00-4.22(m,2H) 2.44-2.69(m,2H) 1.73(tt,J=4.99,8.27Hz,1H) 1.51(d,J=6.88Hz,3H) 1.40-1.43(m,3H) 0.78-1.03(m,4H)

(Type I crystal of compound (I) with 1 equivalent of fumaric acid)

¹HNMR(400MHz,DMSO-d₆)δ: 13.13(hrs,2H) 8.32(brd,J=7.38Hz,1H) 8.14(s,1H) 7.24-7.48(m,5H) 6.46-6.78(m,3H) 6.06-6.30(m,1H) 5.63-5.77(m,1H) 5.11-5.38(m,2H) 4.27-4.44(m,1H) 3.98-4.23(m,2H) 2.42-2.73(m,2H) 1.73(tt,J=5.02,8.30Hz,1H) 1.51(d,J=6.88Hz,3H) 1.32-1.45(m,3H) 0.77-1.02(m,4H)

(Type II crystal of compound (I) with 1 equivalent of fumaric acid)

¹HNMR(400MHz,DMSO-d₆)δ: 13.14(hrs,2H) 8.32(brd,J=7.38Hz,1H) 8.14(s,1H) 7.25-7.49(m,5H) 6.39-6.89(m,3H) 6.07-6.28(m,1H) 5.62-5.77(m,1H) 5.12-5.39(m,2H) 4.27-4.45(m,1H) 3.99-4.24(m,2H) 2.42-2.75(m,2H) 1.74(tt,J=5.02,8.30Hz,1H) 1.51(d,J=6.88Hz,3H) 1.37-1.46(m,3H) 0.78-1.03(m,4H)

(Type V crystal of compound (I) with 1 equivalent of fumaric acid)

¹HNMR(400MHz,DMSO-d₆)δ: 13.13(hrs,2H) 8.32(brd,J=7.38Hz,1H) 8.14(s,1H) 7.21-7.49(m,5H) 6.47-6.80(m,3H) 6.09-6.26(m,1H) 5.63-5.77(m,1H) 5.11-5.38(m,2H) 4.27-4.44(m,1H) 3.98-4.23(m,2H) 2.42-2.72(m,2H) 1.73(tt,J=4.99,8.27Hz,1H) 1.51(d,J=6.88Hz,3H) 1.34-1.46(m,3H) 0.77-1.02(m,4H)

[0129]

Reference Example 1

Production of type III crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with 1 equivalent of fumaric acid

Fumaric acid (3 equivalents) and 9 mL of acetonitrile were added to 300 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 50°C for 1 hour and naturally cooled, and then the solid was collected by filtration and recovered. This was dried to obtain 303.7 mg of a crystal of interest.

[0130]

Powder X-ray diffraction spectrum (Method B): See Figure 15.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 8.1° , 13.2° , 20.4° , 23.1° , 24.7° , and 26.1°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 16.

Exothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 153°C

[0131]

Reference Example 2

Production of type IV crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with 1 equivalent of fumaric acid

Fumaric acid (3 equivalents) and 6 mL of water were added to 300 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 50°C for 1.5 hours and naturally cooled for 3 hours, and then the solid was collected by filtration and recovered. This was washed with water, and then the solid was collected by filtration, recovered, and dried to obtain 311.2 mg of a crystal of interest.

[0132]

Powder X-ray diffraction spectrum (Method B): See Figure 17.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 6.0° , 15.7° , and 18.8°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 18.

Exothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 130°C

[0133]

Reference Example 3

Production of type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with hydrochloric acid

A 2M hydrochloric acid/ethanol solution (15 μL) and 0.3 mL of a liquid mixture of 2-propanol/heptane (1:3 v/v) were added to 15 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred for 4 days and naturally cooled, and then the solid was collected by filtration and recovered. This was dried to obtain a crystal of interest.

[0134]

Powder X-ray diffraction spectrum (Method B): See Figure 19.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 7.0° , 7.4° , 12.6° , 19.0° , and 25.4°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 20.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 51°C and 179°C

[0135]

Reference Example 4

Production of type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with hydrochloric acid

A 2M hydrochloric acid/ethanol solution (15 μ L) and 0.3 mL of a liquid mixture of acetone/heptane (1:3 v/v) were added to 15 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred for 4 days and naturally cooled, and then the solid was collected by filtration and recovered. This was dried to obtain a crystal of interest.

[0136]

Powder X-ray diffraction spectrum (Method B): See Figure 21.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 6.3°, 7.4°, 8.0°, 17.6°, 22.0°, and 23.6°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 22.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 147°C

[0137]

Reference Example 5

Production of type III crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with hydrochloric acid

A 2M hydrochloric acid/ethanol solution (15 μ L) and 0.3 mL of ethyl acetate were added to 15 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred for 4 days and naturally cooled, and then the solid was collected by filtration and recovered. This was dried to obtain a crystal of interest.

[0138]

Powder X-ray diffraction spectrum (Method B): See Figure 23.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 5.3°, 6.5°, 8.1°, 15.3°, and 24.2°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 24.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 156°C

[0139]

Reference Example 6

Production of crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with hydrobromic acid

Hydrobromic acid (1 equivalent) and 0.3 mL of a liquid mixture of 2-propanol/heptane (1:3 v/v) were added to 15 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred for 5 days and naturally cooled, and then the solid was collected by filtration and recovered. This was dried to obtain a crystal of interest.

[0140]

Powder X-ray diffraction spectrum (Method B): See Figure 25.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 7.0° , 7.4° , 13.4° , 15.6° , 19.2° , and 25.6°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 26.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 204°C

[0141]

Reference Example 7

Production of type IV crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with 1 equivalent of L-tartaric acid

L-tartaric acid (466 mg) and 10 mL of acetone were added to 500 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was stirred at 50°C for 40 minutes for dissolution. Thereafter, the dissolved mixture was stirred at room temperature for 17 hours, the suspension was filtrated, and the resulting solid was recovered. This was dried to obtain 506.3 mg of a crystal of interest.

[0142]

Powder X-ray diffraction spectrum (Method A): See Figure 27.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 7.2° , 12.6° , 17.3° , and 23.4°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 28.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 121°C

[0143]

Reference Example 8

Production of crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide (compound (I)) with succinic acid

Succinic acid (3 equivalents) and 0.3 mL of acetonitrile were added to 15 mg of the compound (I) obtained according to the method in Synthesis Example 1, the mixture was suspended and stirred at 50°C for 3 days and naturally cooled, and then the solid was collected by filtration and recovered. This was dried to obtain a crystal of interest.

[0144]

Powder X-ray diffraction spectrum (Method B): See Figure 29.

Characteristic diffraction angle ($2\theta \pm 0.2^\circ$): 6.5° , 10.7° , 13.2° , 14.6° , 18.7° , and 23.9°

Simultaneous thermogravimetry-differential thermal analysis curve: See Figure 30.

Endothermic peak (peak top value) in the simultaneous thermogravimetry-differential thermal analysis curve: Around 148°C and 159°C

[0145]

Test Example 1 Measurement of inhibitory effect (*in vitro*) on HER2 phosphorylation activity

In order to determine conditions for a method of measuring the *in vitro* inhibitory activity of a compound against HER2 phosphorylation activity, based on the report regarding a HER2 kinase reaction using, as a substrate, a peptide having the same sequence (5-FAM-EEPLYWSFPAKKK-CONH₂) as that of ProfilerPro Peptide 22 of PerkinElmer (Xie H et al., PLoS One.2011; 6(7): e21487), ProfilerPro Peptide 22 was used as a substrate. A purified recombinant human HER2 protein used in the present test was purchased from Carna Biosciences, Inc. Upon the measurement of the inhibitory activity of the compound, first, the compound (I) obtained according to the method in Synthesis Example 1 was diluted stepwise with dimethyl sulfoxide (DMSO). Subsequently, the HER2 protein, the substrate peptide (final concentration: 1 μM), manganese chloride (final concentration: 10 mM), ATP (final concentration: 5 μM), and the compound (I) in DMSO solution (final concentration of DMSO: 5%) were added to a buffer for the kinase reaction (13.5 mM Tris (pH 7.5), 2 mM dithiothreitol, and 0.009% Tween 20), and the obtained mixture was then incubated at 25°C for 30 minutes, so that the kinase reaction was carried out. To the reaction solution, EDTA was added to a final concentration of 30 mM, so as to terminate the reaction. Finally, using LabChip (registered trademark) EZ Reader II (PerkinElmer), an unphosphorylated substrate peptide (S) and a phosphorylated peptide (P) were separated and detected according to microchannel capillary electrophoresis. From the peak heights of S and P, the amount of the phosphorylation reaction was obtained, and the concentration of the compound capable of inhibiting the phosphorylation reaction by 50% was defined as an IC₅₀ value (nM). The results are shown in Table 1.

[0146]

Test Example 2 Measurement of inhibitory action (*in vitro*) against HER2 exon 20 insertion mutant (HER2ex20insYVMA) phosphorylation activity

In order to determine conditions for a method of measuring the *in vitro* inhibitory activity of a compound against HER2 exon 20 insertion mutant phosphorylation activity, as in the case of HER2, ProfilerPro Peptide 22 was used as a substrate. A purified recombinant human HER2 exon 20 insertion mutant (A775_G776insYVMA) protein was purchased from SignalChem. Upon the measurement of the inhibitory activity of the compound, first, the compound (I) obtained according to the method in Synthesis Example 1 was diluted stepwise with dimethyl sulfoxide (DMSO). Subsequently, the HER2 exon 20 insertion mutant protein and the compound (I) in DMSO solution (final concentration of DMSO: 5%) were added into a buffer for the kinase reaction (13.5 mM Tris (pH 7.5), 2 mM dithiothreitol, and 0.009% Tween 20), and the obtained mixture was then pre-incubated at 25°C for 30 minutes. Thereafter, the substrate peptide (final concentration: 1 µM), manganese chloride (final concentration: 25 mM), magnesium chloride (final concentration: 20 mM), and ATP (final concentration: 200 µM) were added into the reaction mixture, and the thus obtained mixture was then incubated at 25°C for 220 minutes, so that the kinase reaction was carried out. To the reaction solution, EDTA was added to a final concentration of 30 mM, so as to terminate the reaction. Finally, using LabChip (registered trademark) EZ Reader II (PerkinElmer), an unphosphorylated substrate peptide (S) and a phosphorylated peptide (P) were separated and detected according to microchannel capillary electrophoresis. From the peak heights of S and P, the amount of the phosphorylation reaction was obtained, and the concentration of the compound capable of inhibiting the phosphorylation reaction by 50% was defined as an IC50 value (nM). The results are shown in Table 7.

[0147]

[Table 7]

HER2 inhibitory activity IC50 value (nM)	HER2ex20insYVMA inhibitory activity IC50 value (nM)
3.2	< 0.30

[0148]

From the above results, it was found that the compound (I) has excellent inhibitory activity against phosphorylation of HER2 and against phosphorylation of HER2 exon 20 insertion mutant.

[0149]

Test Example 3 Measurement of growth inhibitory activity against HER2 expressing

cell line

SK-BR-3 cells as a HER2 overexpressing human breast cancer cell line were suspended in a McCoy's 5a medium (manufactured by Life Technologies) supplemented with 10% fetal bovine serum. The cell suspension was seeded in each well of a 384-well flat-bottom microplate, and was then cultured in a 5% carbon dioxide gas-containing culture vessel at 37°C for 1 day. Thereafter, the compound (I) obtained according to the method in Synthesis Example 1 was dissolved in DMSO, and the compound was diluted to 500 times the final concentration in DMSO. The compound in the DMSO solution was diluted with DMSO solution or the medium used in the suspension of the cells, and the obtained solution was then added to each well of the culture plate so that the final concentration of DMSO was 0.2%. The obtained mixture was further cultured in the 5% carbon dioxide gas-containing culture vessel at 37°C for 3 days. After completion of the culture for 3 days in the presence of the compound, the cells were counted using CellTiter-Glo 2.0 (manufactured by Promega), and the growth inhibition percentage was then calculated according to the following equation. The concentration of the compound, in which the growth of the cells can be inhibited by 50%, was defined as IC₅₀ (nM).

Growth inhibitory percentage (%) = (C-T) / (C) x 100

T: Emission intensity from the well to which the test compound was added

C: Emission intensity from the well to which the test compound was not added

The results are shown in the following Table 8.

[0150]

Test Example 4 Measurement of growth inhibitory activity against HER2 exon 20 insertion mutant expressing cell line

Growth inhibitory activity against the HER2 exon 20 insertion mutant was measured using Ba/F3 cells that were a mouse B lymphocyte precursor cell line, into which a human HER2 exon 20 insertion mutant gene had been introduced. The Ba/F3 cells were maintained in an RPMI-1640 medium (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum (FBS), 100 U/mL penicillin, 100 µg/mL streptomycin (Thermo Fisher Scientific) and 1 ng/mL mouse interleukin-3 (mIL-3) (CST). Thereafter, a pCDNA3.1-hyg(+) vector, into which a human HER2 exon 20 insertion mutant gene (A775_G776insYVMA (HER2ex20insYVMA)), Internal Ribosome Binding Sequence (IRES), and a Kusabira orange gene had been incorporated, was introduced into the Ba/F3 cells according to an electroporation method using Amaxa (registered trademark) Cell Line Nucleofector (registered trademark) Kit V. The Ba/F3 cells expressing the HER2 exon 20 insertion mutant (Ba/F3-HER2insYVMA), which were selected with hygromycin B (Nacalai Tesque), exhibited mIL-3-independent growth.

Upon evaluation of cell growth inhibitory activity, the Ba/F3-HER2insYVMA cells were suspended in an RPMI-1640 medium supplemented with 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin. The cell suspension was seeded in each well of a 96-well flat-bottom microplate, and was then cultured in a 5% carbon dioxide gas-containing culture vessel at 37°C for 1 day. The compound (I) obtained according to the method in Synthesis Example 1 was dissolved in DMSO, and was then diluted with DMSO or the medium used in the suspension of the cells. The obtained solution was then added to each well of the culture plate, so that the final concentration of DMSO became 0.2%. The obtained mixture was further cultured in the 5% carbon dioxide gas-containing culture vessel at 37°C for 3 days. After completion of the culture for 3 days in the presence of the compound, the cells were counted using CellTiter-Glo 2.0 (manufactured by Promega), and the growth inhibition percentage was then calculated according to the following equation. The concentration of the compound, in which the growth of the cells can be inhibited by 50%, was defined as IC50 (nM).

$$\text{Growth inhibitory percentage (\%)} = (C-T) / (C) \times 100$$

T: Emission intensity from the well to which the test compound was added

C: Emission intensity from the well to which the test compound was not added

The results are shown in the following Table 8.

[0151]

[Table 8]

SK-BR-3 cell growth inhibitory activity IC50 value (nM)	HER2ex20insYVMA cell growth inhibitory activity IC50 value (nM)
6.6	29

[0152]

From the above results, it was found that the compound (I) has excellent cell growth inhibitory activity even against the HER2 expressing cell line (SK-BR-3) and also, against the HER2 exon 20 insertion mutant expressing cell line (Ba/F3-HER2insYVMA).

[0153]

Test Example 5 Measurement of growth inhibitory activity against HER2 expressing cell line (NCI-N87)

NCI-N87 cells as a HER2 overexpressing human stomach cancer cell line (American Type Culture Collection, Cat No. ATCC (registered trademark) CRL-5822) were suspended in an RPMI1640 medium (Wako Pure Chemical Industries, Ltd.) supplemented with 10% fetal bovine serum. Subsequently, the cell suspension was

seeded in each well of a 96-well flat-bottom microplate, and was then cultured in a 5% carbon dioxide gas-containing culture vessel at 37°C for 1 day. Thereafter, the compound (I) obtained according to the method in Synthesis Example 1 was dissolved in DMSO, and the compound was diluted to 1000 times the final concentration in DMSO. The compound (I) in the DMSO solution was diluted with the medium used in the suspension of the cells, and the obtained solution was then added to each well of the culture plate, so that the final concentration of DMSO became 0.1%. Regarding a control well, DMSO was diluted with the medium used in the suspension of the cells, and the obtained solution was then added to each well of the culture plate, so that the final concentration of DMSO became 0.1%. After addition of a drug solution, the obtained mixture was further cultured in the 5% carbon dioxide gas-containing culture vessel at 37°C for 3 days. After completion of the culture for 3 days in the presence of the compound, the cells were counted using CellTiter-Glo 2.0 (manufactured by Promega) in accordance with the protocols recommended by Promega. The growth inhibition percentage was calculated according to the following equation. The concentration of the compound, in which the growth of the cells can be inhibited by 50%, was defined as IC50 (nM).

$$\text{Growth inhibitory percentage (\%)} = (C - T) / (C) \times 100$$

T: Emission intensity from the well to which the test compound was added

C: Emission intensity from the well to which the test compound was not added

The results are shown in the following Table 9.

[0154]

[Table 9]

NCI-N87 cell growth inhibitory activity IC50 value (nM)
9.9

[0155]

From the above results, it was found that the compound (I) has excellent cell growth inhibitory activity even against the HER2 overexpressing cell line (NCI-N87).

[0156]

Test Example 6 Evaluation of oral absorbability

The compound (I) obtained according to the method in Synthesis Example 1 was suspended or dissolved in 0.5% HPMC aqueous solution and 0.1 N hydrochloric acid, and the obtained suspension or solution was orally administered to BALB/cA mice (CLEA Japan, Inc.) at a dose of 50 mg/kg/day. At 0.5, 1, 2, 4 and 6 hours after

completion of the oral administration, blood was collected from the facial vein over time, so as to obtain plasma. The concentration of the compound in the obtained plasma was measured by LC-MS/MS, and the oral absorbability of the present compound was evaluated.

The results are shown in the following Table 10.

[0157]

[Table 10]

AUC 0 - 6 hr ($\mu\text{M}\cdot\text{hr}$)
31

[0158]

From the above results, it was found that the compound of the present invention was contained in a sufficient concentration in the plasma, so that the compound (I) exhibited favorable oral absorbability.

[0159]

Test Example 7 Evaluation of brain penetration properties

The compound (I) obtained according to the method in Synthesis Example 1 was suspended or dissolved in 0.5% HPMC aqueous solution and 0.1 N hydrochloric acid, and the obtained suspension or solution was orally administered to BALB/cA mice (CLEA Japan, Inc.) at a dose of 50 mg/kg/day. At 0.5 hours after completion of the oral administration, blood was collected from the facial vein, and whole brain was then excised, so as to obtain plasma and brain samples. Water was added to the obtained brain sample in 3 times the volume of the brain sample, and the resultant was then homogenized using an ultrasonic homogenizer, so as to obtain a brain homogenate. The concentration of the compound in the obtained plasma and brain homogenate was measured by LC-MS/MS, and the brain penetration properties of the present compound were evaluated from the brain/plasma concentration of the compound.

The results are shown in the following Table 11.

[0160]

[Table 11]

Compound concentration in plasma (μM)	Compound concentration in brain (μM)	Kp value (Compound concentration in brain/plasma)
12	2.7	0.23

[0161]

From the above results, it was found that the compound (I) exhibited favorable brain penetration properties.

[0162]

Test Example 8 Antitumor effect confirmation test (*in vivo*) on direct brain transplantation models, into which luciferase gene-introduced HER2 expressing cell line (NCI-N87-luc) is directly transplanted

In order to confirm the antitumor effects of a test compound on direct brain transplantation models, NCI-N87-Luc, which was obtained by introducing a luciferase gene into NCI-N87 that was a human stomach cancer tumor cell line purchased from American Type Culture Collection, was used. The NCI-N87-Luc was added into a 10% fetal bovine serum (FBS)-containing RPMI-1640 medium (supplemented with 4.5 g/L glucose, 10 mM HEPES, and 1 mM sodium pyruvate) (Wako Pure Chemical Industries, Ltd.), and this cell line was then cultured in a 5% CO₂ incubator at 37°C.

The NCI-N87-Luc cells were re-suspended in PBS in a concentration of 6.25×10^7 cells/mL.

Using a mouse ear bar, a nude mouse with 6 to 7 weeks old (BALB/cAJcl-nu/nu, CLEA Japan, Inc.) was fixed in a brain stereotaxic apparatus, and the skin on the upper brain portion was disinfected with alcohol cotton and was then excised with a surgical knife.

A microdrill was used to drill a hole in the skull, and then, using a needle, a manipulator, and a syringe pump, 4 µL of the cell suspension was transplanted into the brain at a rate of 0.8 µL/min.

As a reference of the amount of brain tumor, approximately 3 weeks after the transplantation, Total Flux (Photon/sec) was measured in all of the survival cases, using IVIS (PerkinElmer, Inc., model: Lumina II). Based on the obtained results, 6 animals were assigned to each group, using the grouping program of MiSTAT (Ver. 2.00).

The test compound was orally administered to the mice once a day, every day, for 21 days from the following day of the grouping (Days 1 - 21).

For judgment of the presence or absence of effects, the value (Log₁₀) obtained by logarithmic transformation of the total flux on the judgment date was used. The compound (I) obtained according to the method in Synthesis Example 1 as a test compound was administered to the mice at a dose of 25 mg/kg/day.

A graph was prepared with the value obtained by logarithmic transformation (Log₁₀) of the average total flux as a vertical axis, and with the number of days (Day) after the transplantation as a horizontal axis. The transition of the total flux over time in

the drug administration period was observed.

As a control, 0.1 N HCl and 0.5% HPMC aqueous solution were used.

[0163]

The results are shown in the following Figure 31 and Figure 32. The value obtained by logarithmic transformation (Log10) of the total flux on Day 22 in each group was analyzed by a significance test. As a result, it was demonstrated that the aforementioned value of the compound was statistically significantly lower than the value of the control group (significance level (both sides): 5%). For the measurement of the body weight, an animal electronic balance was used. A body weight change percentage (Body weight change; BWCn) from the body weight on the nth day (BWn) was calculated according to the following equation:

$$\text{BWCn (\%)} = [(\text{body weight on } n^{\text{th}} \text{ day}) - (\text{body weight on grouping day})] / [(\text{body weight on grouping day})] \times 100.$$

From the results of this test, it was found that the compound (I) has excellent antitumor effects against the HER2 overexpressing cell line (NCI-N87-luc) transplanted into the nude mice. Moreover, a body weight reduction of -20% or more was not observed in all of the mice had been administered. Accordingly, it was found that there were no serious side effects.

[0164]

Test Example 9

Solid stability test

The free-form type II crystals of the compound (I) and type II crystals thereof with 1 equivalent of fumaric acid obtained in the Examples and Comparative Examples were evaluated for solid stability after storage for 4 weeks.

Storage condition: 60°C (closed system)

Storage period: 4 weeks

Stored amount: Approximately 30 mg

Storage container: Glass bottle

The results are shown in the following Table 12.

[Table 12]

	Free-form type II crystal	Type II crystal with 1 equivalent of fumaric acid
Chemical purity before storage (%)	99.87	99.85
Chemical purity after storage (%)	99.88	99.85

Amount of related substances before storage (%)	0.13	0.15
Amount of related substances after storage (%)	0.12	0.15

Approximately 1 mg of a sample was weighed, dissolved in approximately 1 mL of a liquid mixture of water/acetonitrile (1: 1), and 5 μ L of this solution was accurately measured. Changes in the amount of related substances (the amount of substances detected other compound (I)) were analyzed by HPLC by the following method.

HPLC analysis method (stability test)

The amount of related substances in the sample solution was measured by HPLC analysis. Handling of the apparatuses, including data processing, was in accordance with the methods and procedures instructed for each apparatus.

Column: XSelect CSH C18 (4.6 $\times$ 150 mm, 3.5 μ m) manufactured by Waters

UV detection: 246nm

Column temperature: 40°C

Flow rate: 1.0 mL / min

Sample cooler: 5°C

Sample concentration: 1 mg/mL

Mobile phase A: Liquid mixture of 5 mmol/L ammonium formate buffer (pH 6.5)/acetonitrile (9:1)

Mobile phase B: Acetonitrile

The gradients are shown in Table 13.

[Table 13]

Time (min)	Mobile phase A (vol%)	Mobile phase B (vol%)
0-31	90→30	10→70
31-36	30	70
36-37	30→90	70→10

As a result, no change in the powder X-ray diffraction pattern was observed in either the free-form type II crystal of compound (I) or the type II crystal of the same with 1 equivalent of fumaric acid, and almost no increase in related substances was observed. They were found to be extremely stable crystals.

[0165]

Test Example 10

Dynamic vapor sorption (DVS) test

Dynamic vapor sorption tests were conducted using a type II crystal of compound (I) with 1 equivalent of fumaric acid, a free-form type II crystal of the same, a free-form type I crystal of the same, a type V crystal of the same with 1 equivalent of fumaric acid, and a crystal of the same with 1 equivalent of L-tartaric acid obtained in the Examples and Reference Examples.

The dynamic vapor sorption test was conducted for measurement according to the following conditions.

Approximately 10 mg of each sample was packed in a dedicated quartz holder, and the weight of the sample at each humidity was continuously measured and recorded under the following conditions. Handling of the apparatuses, including data processing, was in accordance with the methods and procedures instructed for each apparatus.

Apparatus: VTI SA+ (manufactured by TA Instruments)

Drying temperature: 60°C

Temperature increase rate: 1°C/min

Equilibrium of drying: It was confirmed that a decrease by 0.01 wt% in 5 minutes did not occur within the range not exceeding 300 minutes.

Measurement temperature: 25 °C

Equilibrium of humidification : It was confirmed that an increase by 0.01 wt% in 5 minutes did not occur within the range not exceeding 120 minutes.

Relative humidity program: Increase by 5% RH from 5% RH to 95% RH and decrease by 5% RH from 95% RH to 5% RH

The weight changes in the range of measurement conditions obtained in these tests are shown in Tables 14 to 18.

[Table 14]

Table 14: Results of dynamic vapor sorption test of type II crystal of compound (I) with 1 equivalent of fumaric acid

Relative humidity (%)	Weight change rate (%)	
	Adsorption	Desorption
5	0.01	0.04
20	0.11	0.13
40	0.18	0.20
60	0.27	0.30

80	0.34	0.34
95	0.38	0.38

[Table 15]

Table 15: Results of dynamic vapor sorption test of free-form type II crystal of compound (I)

	Weight change rate (%)	
Relative humidity (%)	Adsorption	Desorption
5	0.01	0.01
20	0.02	0.03
40	0.05	0.06
60	0.08	0.08
80	0.10	0.11
95	0.12	0.12

[Table 16]

Table 16: Results of dynamic vapor sorption test of free-form type I crystal of compound (I)

	Weight change rate (%)	
Relative humidity (%)	Adsorption	Desorption
5	0.01	0.01
20	0.04	0.04
40	0.10	0.13
60	0.26	0.60
80	0.50	0.69
95	0.72	0.72

[Table 17]

Table 17: Results of dynamic vapor sorption test of type V crystal of compound (I) with 1 equivalent of fumaric acid

	Weight change rate (%)	
Relative humidity (%)	Adsorption	Desorption
5	0.01	0.06
20	0.11	0.16
40	0.21	0.27
60	0.33	0.39
80	0.47	0.51
95	0.59	0.59

[Table 18]

Table 18: Results of dynamic vapor sorption test of crystal of compound (I) with 1 equivalent of L-tartaric acid

	Weight change rate (%)	
Relative humidity (%)	Adsorption	Desorption
5	0.50	0.54
20	1.05	1.16
40	1.60	1.67
60	2.16	2.38
80	3.26	3.64
95	6.41	6.41

[0166]

As shown in Table 14, the type II crystal of compound (I) with 1 equivalent of fumaric acid had a weight increase of 0.38% in a dynamic vapor sorption test under 95% relative humidity, which was less than 1%. The crystal was determined to be almost non-hygroscopic. Similarly, the free-form type II crystal of compound (I), the free-form type I crystal of the same, and the type V crystal of the same with 1 equivalent of fumaric acid also had a mass increase of less than 1% in a dynamic vapor sorption test under 95% relative humidity, which was less than 1%. The crystals were determined to be almost non-hygroscopic (Tables 15 to 17). Meanwhile, as shown in Table 18, the crystal of compound (I) with 1 equivalent of L-tartaric acid had a mass increase of as high as 6.41%

in a dynamic vapor sorption test under 95% relative humidity, which was less than 1%. The crystal was determined to be highly hygroscopic.

Therefore, since the type II crystal of compound (I) with 1 equivalent of fumaric acid, the free-form type II crystal of the same, the free-form type I crystal of the same, and the type V crystal with 1 equivalent of fumaric acid are less hygroscopic than the crystal of compound (I) with 1 equivalent of L-tartaric acid, they are considered to be excellent candidate compounds for drug development in terms of industrial manufacturing of pharmaceutical products with stable quality. The type II crystal of compound (I) with 1 equivalent of fumaric acid, the free-form type II crystal of the same, the free-form type I crystal of the same, and the type V crystal of the same with 1 equivalent of fumaric acid were confirmed to have excellent properties as pharmaceutical products or active pharmaceutical ingredients.

[0167]

Test Example 11

Blood concentration measurement test

Blood concentration measurement tests were conducted for the type I crystal of compound (I), the type II crystal of compound (I), and the type II crystal of compound (I) with 1 equivalent of fumarate obtained in the Examples. Specifically, 0.5% HPMC containing 0.1N hydrochloric acid was used for the type I crystal of compound (I) and 0.5% HPMC was used for the type II crystal of compound (I) and the type II crystal of compound (I) with 1 equivalent of fumaric acid to convert each crystal into the compound (I) (free-form) in terms of molecular weight so as to prepare suspensions of 100 mg/10 mL and 300 mg/10 mL. These administration solutions were orally administered to female rats (CrI:CD(SD), Charles River Laboratories Japan, Inc.) kept under feeding conditions at a volume of 10 mL/kg body weight using an oral administration sonde. After administration, the rats were returned to the rat cage and their conditions were checked. Water and feeding were freely available in the cage. At 0.5, 1, 2, 4, and 6 hours after administration, the rats were anesthetized with isoflurane, and approximately 80 μ L of blood was collected from the jugular vein using a heparin-coated syringe and an injection needle (25 G).

The collected blood was ice-cooled and plasma was separated by centrifugation. After the blood collection was completed, the rats were returned to the rat cage and the state after awakening of anesthesia was checked. After the final blood collection, the depth of isoflurane anesthesia was checked. Then, the limbs were fixed in a supine position with a small animal experiment holding apparatus, and the abdomen was opened such that each rat was euthanized by exsanguination through an incision in the abdominal

aorta and vena cava.

AUC_{0-6hr} (Area under the blood concentration-time curve for 0 to 6 hours after administration calculated by the trapezoidal method), C_{max} (maximum blood concentration), and T_{max} (time to reach maximum blood concentration) were calculated from the concentration of the compound (I) in each plasma measured by the MRM method using LC/MS/MS and quantified from the calibration curve using Phoenix WinNonlin (v6.4.0, Certara USA, Inc.).

[0168]

The results are shown in Table 19. When the dose was increased from 100 mg/kg to 300 mg/kg, for the type I crystal of compound (I) with hydrochloric acid and the type II crystal of compound (I), C_{max} increased 0.8 times and 1.2 times, respectively, and AUC_{0-6hr} increased 1.0 times and 1.5 times, respectively. Meanwhile, for the type II crystal of compound (I) with 1 equivalent of fumaric acid, each of C_{max} and AUC_{0-6hr} increased 1.9 times. In addition, at a dose of 300 mg/kg, C_{max} of the type II crystal of compound (I) with 1 equivalent of fumaric acid was 1.7 times the value of the type I crystal of compound (I) with the acid and 1.4 times the value of the type II crystal of compound (I), and AUC_{0-6hr} of the type II crystal of compound (I) with 1 equivalent of fumaric acid was 2.4 times the values of the type I crystal of compound (I) with the acid and the type II crystal of compound (I). Therefore, the type II crystal of compound (I) with 1 equivalent of fumaric acid according to the present invention was confirmed to show more favorable oral absorbability.

[Table 19]

Dose	Parameter	Oral administration		
		Type I crystal of compound (I) (with hydrochloric acid)	Type II crystal of compound (I)	Type II crystal of compound (I) with 1 equivalent of fumaric acid
100 mg/kg	AUC _{0-6hr} (μM·hr)	7.42	5.79	6.14
	C _{max} (μM)	2.67	1.88	2.74
	T _{max} (hr)	0.75	1.50	1.25
300 mg/kg	AUC _{0-6hr} (μM·hr)	7.11	8.53	11.9
	C _{max} (μM)	2.25	2.25	5.34
	T _{max} (hr)	3.25	3.33	0.5

[0169]

It is to be noted that all documents and publications cited in the present description are incorporated herein by reference in their entirety, regardless of the purpose thereof. Moreover, the present description includes the contents disclosed in the claims, specification and drawings of Japanese Patent Application No. 2020-121520 (filed on July 15, 2020), from which the present application claims priority.

Several embodiments of the present invention are described above. However, these embodiments are provided for illustrative purpose only, and thus, are not intended to limit the scope of the present invention. These novel embodiments can be carried out in various other forms, and various abbreviations, substitutions and alternations can be carried out, unless they are deviated from the spirit of the invention. These embodiments and the modifications thereof are included in the scope or spirit of the invention, and are also included in the invention according to the claims and the scope equivalent thereto.

CLAIMS

1. A crystal having peaks at three or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 5.5° , 6.8° , 9.3° , 13.4° , 15.3° , 16.3° , 18.5° , 19.8° , 22.0° , and 24.5° in a powder X-ray diffraction spectrum, which is a type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:1.
2. The crystal according to claim 1, which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 5.5° , 6.8° , 9.3° , 13.4° , 15.3° , 16.3° , 18.5° , 19.8° , 22.0° , and 24.5° in a powder X-ray diffraction spectrum.
3. The crystal according to claim 1 or 2, which is a crystal having a powder X-ray diffraction spectrum shown in Figure 1.
4. The crystal according to any one of claims 1 to 3, which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 178°C .
5. A crystal having peaks at three or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 8.3° , 14.8° , 17.3° , 18.0° , 19.1° , 20.3° , 21.0° , 22.5° , 23.0° , and 26.2° in a powder X-ray diffraction spectrum, which is a type II crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide.
6. The crystal according to claim 5, which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 8.3° , 14.8° , 17.3° , 18.0° , 19.1° , 20.3° , 21.0° , 22.5° , 23.0° , and 26.2° in a powder X-ray diffraction spectrum.
7. The crystal according to claim 5 or 6, which is a crystal having a powder X-ray diffraction spectrum shown in Figure 3.
8. The crystal according to any one of claims 5 to 7, which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around

182°C.

9. A crystal having peaks at three or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 9.9° , 11.7° , 13.2° , 17.7° , 18.1° , 18.8° , and 20.8° in a powder X-ray diffraction spectrum, which is a type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide.

10. The crystal according to claim 9, which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 9.9° , 11.7° , 13.2° , 17.7° , 18.1° , 18.8° , and 20.8° in a powder X-ray diffraction spectrum.

11. The crystal according to claim 9 or 10, which is a crystal having a powder X-ray diffraction spectrum shown in Figure 5.

12. The crystal according to any one of claims 9 to 11, which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 183°C.

13. A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 6.9° , 9.4° , 10.2° , 13.7° , 21.1° , 23.6° , and 26.5° in a powder X-ray diffraction spectrum, which is a type V crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:1.

14. The crystal according to claim 13, which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 6.9° , 9.4° , 10.2° , 13.7° , 21.1° , 23.6° , and 26.5° in a powder X-ray diffraction spectrum.

15. The crystal according to claim 13 or 14, which is a crystal having a powder X-ray diffraction spectrum shown in Figure 7.

16. The crystal according to any one of claims 13 to 15, which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at

around 151°C.

17. A crystal having peaks at four or more diffraction angles ($2\theta \pm 0.2^\circ$) selected from 6.4° , 10.3° , 12.8° , 15.0° , 20.7° , 23.4° , and 26.6° in a powder X-ray diffraction spectrum, which is a type I crystal of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide with fumaric acid, wherein a molar ratio of 7-((3R,5S)-1-acryloyl-5-methylpyrrolidin-3-yl)-4-amino-6-(cyclopropylethynyl)-N-((R)-1-phenylethyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carboxamide to fumaric acid is 1:1.

18. The crystal according to claim 17, which has peaks at diffraction angles ($2\theta \pm 0.2^\circ$) of 6.4° , 10.3° , 12.8° , 15.0° , 20.7° , 23.4° , and 26.6° in a powder X-ray diffraction spectrum.

19. The crystal according to claim 17 or 18, which is a crystal having a powder X-ray diffraction spectrum shown in Figure 9.

20. The crystal according to any one of claims 17 to 19, which has an endothermic peak determined by simultaneous thermogravimetry-differential thermal analysis at around 153°C.

21. A pharmaceutical composition, comprising the crystal according to any one of claims 1 to 20.

22. A pharmaceutical composition for oral administration, comprising the crystal according to any one of claims 1 to 20

23. An antitumor agent, comprising the crystal according to any one of claims 1 to 20.