

Description

This invention relates to a novel fatty acid-lactic acid ester and, more particularly, to a lactic acid ester of α -branched fatty acid which possesses an excellent hair conditioning effect and can be used as toiletries.

Description of the Background:


Since lactic acid esters of fatty acids function to produce an antiseptic effect, emulsion formation, detergency, and the like, and can be absorbed by proteins, they are used in foods, shampoos, cosmetics, or the like. Stearoyl lactic acid salts (sodium or potassium salt) which are linear fatty acid esters, for example, are known to be capable of ensuring soft-baked bread, and of providing an anti-aging effect. These compounds are used as a bread substrate improver. In addition to these lactic acid esters of a linear fatty acid, lactic acid esters having an acyl group of iso-stearic acid with a methyl group as a branched chain, which are produced as by-products in the production of dimer acids, are also used for shampoos and cosmetics as agents producing the effects of detergency, emulsification, and conditioning.

In this way, the above-mentioned lactic acid ester of iso-stearic acid, which is a by-product of dimer acid production processes, or the like, are presently used as hair cosmetics such as shampoos and rinse, relying on its hair conditioning effect due to its ability to absorb proteins. The conditioning effect of these esters, however, is not yet satisfactory in view of the tangling feeling when hair is combed or an incompletely finished feeling after washing due to the absence of dampishness.

The inventors have conducted extensive studies to obtain a compound which provides an excellent hair conditioning effect, and found that lactic acid esters of fatty acids having a branched chain at the α -position, represented by formula (I) which follows, exhibit this hair conditioning effect. Such a finding has led to the completion of this invention.

SUMMARY OF THE INVENTION

Accordingly, an object of this invention is to provide a fatty acid-lactic acid ester represented by the formula:



in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5.


Another object of the invention is to provide a shampoo composition comprising as essential components the fatty acid-lactic acid ester of the above formula (I) and an anionic surface active agent.

Still another object of the invention is to provide a rinse composition comprising as essential components the fatty acid-lactic acid ester of the above formula (I) and a quarternary ammonium salt.

Other objects, features, and advantages of this invention will appear more fully from the following description.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS


In fatty acid-lactic acid esters of this invention represented by the formula (I), the acyl group of



may be 2-butyloctanoyl group, 2-butylnonadecanoyl group, 2-hexyldecanoyl group, 2-heptylundecanoyl group, 2-octyldecanoyl group, 2-decyltetradecanoyl group, 2-decylheptadecanoyl group, 2-tetradecyl-

tadecanoyl group, 2-hexadecyleicosanoyl group, or the like. Among these, those containing 14 to 32 carbon atoms are preferable.

Further, particularly preferable fatty acid-lactic acid esters are those containing an acyl group of the formula:



in which n has a value of 5 to 15 and m has a value of 3 to 14.

The n value in formula (I) represents the average numbers of lactic radicals $\{\text{OCHCH}_3\text{CO}\}$, and is preferably 1 to 3.

Fatty acid-lactic acid esters of this invention can be prepared according to methods known in the art. For example, they are easily prepared by reacting a fatty acid of a suitable structure and carbon atom content with lactic acid in the presence of an alkali such as sodium hydroxide, potassium hydroxide, sodium carbonate, or the like, at 140 to 220°C for 4 to 10 hours while eliminating formed water, and, as required, by converting the free acids thus obtained into various kinds of salts. In this reaction, an n mole amount of lactic acid [n is the number in formula (I)] is charged to the reaction system and reacted with one mole of fatty acid, using usually an alkali of one mole.

A 50% or 90% aqueous lactic acid solution available in the market may be used as a starting raw material for this reaction. Such an aqueous lactic acid solution, if heated to a temperature of 180 to 220°C, is readily converted by dehydration-condensation to condensed lactic acid. Such condensed lactic acid can also be used as a raw material.

As a fatty acid, which is another starting raw material for deriving the esters of this invention, commercially available fatty acids can be used such as, for example, 2-heptylundecanoic acid (Trade Name: Diadol 18GA, manufactured by Mitsubishi Chemical Industries, Ltd.) which is obtained by oxidation of Guerbet alcohol, 2-(1,3,3-trimethylbutyl)-5,7,7-trimethyloctanoic acid (Tradename: Iso-stearic acid, manufactured by Nissan Chemical Co., Ltd.) derived by oxidation of an aldol condensate, and the like. Fatty acids to be used, however, are not limited to those mentioned above. Various fatty acids with a variety of carbon atom contents and prepared by diversified methods can be used as raw materials.

The fatty acid-lactic acid esters of this invention can be formulated in shampoo compositions and hair rinse compositions. Examples of preferred formulations are as follows:

Shampoo compositions

- | | |
|-----------------------------------|--------------------|
| (A) Fatty acid-lactic acid esters | 0.1 - 5% by weight |
| (B) Anionic surface active agents | 5 - 50% by weight |

Examples of anionic surface active agents to be used as component (B) of these compositions are salts of alkylbenzene sulfonic acids having alkyl groups of an average 10 to 16 carbon atom content, salts of sulfuric acid esters of higher alcohols having an average 8 to 18 carbon atom content, salts of sulfuric acid esters of polyoxyalkylene (an average 1 to 10 mole adduct of C_2 or C_3 alkylene oxide) adducts of higher alcohols having an average 8 to 18 carbon atom content, salts of α -olefin sulfonates with an average 8 to 18 carbon atom content, salts of alkyl phosphates with an average 8 to 18 carbon atom content, and the like. A counter ion of these anionic surface active agents is selected from the group consisting of alkali metals, alkaline earth metals, ammonium, and alkanol amines of a C_2 or C_3 carbon atom content.


Known components for conventional shampoos may be formulated in the shampoo compositions of this invention as required in the amount not impairing the effect of this invention. Such known components include, for example, amphoteric or nonionic surface active agents, solubilizing agents such as propylene glycol, glycerol, urea, and the like, viscosity controlling agents such as ethyl alcohol, isopropyl alcohol, hydroxyethyl cellulose, methyl cellulose, higher alcohols, and the like, perfumes, coloring agents, ultraviolet absorbers, antioxidants, antiseptics, pearly agents, and the like.

Hair rinse compositions

- | | |
|-----------------------------------|--------------------|
| (A) Fatty acid-lactic acid esters | 0.1 - 5% by weight |
|-----------------------------------|--------------------|

(C) Salts of quaternary ammonium 0.1 - 10% by weight

Preferable salts of quaternary ammonium to be used as the component (C) are compounds represented by formula (II):



in which one or two radicals among from $R_3 - R_6$ represent long-chained alkyl or hydroxyalkyl groups each having 8 to 22 carbon atoms, with the remaining radicals being alkyl or hydroxyalkyl groups each having 1 to 3 carbon atoms, or benzyl groups, and X represents a halogen atom or an alkyl sulfate group of a 1 or 2 carbon atom content. Various additives commonly used for hair treatment agents may be added to this hair rinse composition to the extent that its intended effect may not be impaired. Such additives include oils or fats such as fatty acids, higher alcohol, oil or fat esters of lanolin, liquid paraffin, and the like, surface active agents such as nonionic surface active agents, amphoteric surface active agents, and the like, disinfectants, vehicles, perfumes, coloring agents, and the like. In particular, the addition of hydrocarbons with molecular weights of 100 to 700, such as liquid and solid paraffin, in the amount of 0.1 to 10% will enhance softness of hair treated by these hair rinse compositions.

Fatty acid-lactic acid esters prepared as illustrated above not only possess excellent dehydration resistance in themselves, but also, when applied to hair or skin, produce an easiness or readiness of handling, and improve such conditioning effects as slipperiness and hair curling formation or retention performance. The compounds of this invention, therefore, may find wide application in the toiletry field as ingredients for shampoos, rinses, treatments, creams, and the like.

Other features of the invention will become apparent in the course of the following description of the exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

Example 1

To a 1 liter 4-necked flask were charged 287 gm of 2-heptylundecanoic acid, 148 gm of 90% lactic acid, and 45 gm of sodium carbonate. Following a temperature increase while feeding nitrogen gas, an esterification reaction was effected at 180°C for 5 hours to obtain 415 gm of an amber, slightly viscous liquid of sodium 2-heptylundecanoyl acid lactate [the compound in which $R_1 = C_9H_{19}$, $R_2 = C_7H_{15}$, $n = 1.5$, and $X = Na$ in formula (I)].

Analysis of this compound revealed that it had a saponification value of 216.8 mg/KOH (after having been saponified for 3 hours), an acid value of 62.1 mg/KOH, a hydroxyl value of 35.5 mg/KOH, an Na content of 4.7%, a viscosity of 580 cp (at 100°C), and a specific gravity of 0.948 (at 100°C).

Examples 2 - 7

In these examples, the raw materials listed in Table 1 were reacted under the reaction conditions noted to give the compounds summarized Table 2.

Table 1

Examples	Amounts of Raw Materials (gm)			Reaction Temp. (°C)	Reaction Time (hour)
	Fatty acid	Lactic acid	Na ₂ CO ₃		
No. 2	2-butyloctanoic acid 200	149	36	200	5
No. 3	2-(1,3,3-trimethyl butyl)-5,7,7-trimethyl octanoic acid 284	199	45	200	5
No. 4	2-octyldecanoic acid 292	148	46	180	5
No. 5	2-heptylundecanoic acid 284	268 *	45	180	5
No. 6	2-decyldodecanoic acid 456	147	53	180	5
No. 7	2-decyloctadecanoic acid 562	147	53	180	5

* 50% lactic acid

Table 2

Esters prepared in Example	Analytical Values			Degree of Condensation (n)	Yield (gm)
	Saponification Value	Acid Value	Hydroxyl Value		
No. 2	282.2	60.4	19.3	1.5	317
No. 3	187.3	65.1	41.8	2.0	460
No. 4	211.6	66.2	40.7	1.5	423
No. 5	218.3	72.9	43.7	1.5	408
No. 6	137.7	53.5	45.4	1.5	601
No. 7	123.7	37.8	26.1	1.5	654

Example 8

The shampoo compositions in Table 3 were prepared according to a conventional method. Each shampoo composition was subjected to testing, in which human hair weighing 30 gm retaining 20 gm of water at 40° C therein was washed with 1 gm of the shampoo composition. Lathering, cleansing ability, and the like of the composition were evaluated by expert panelists. The results are shown in Table 3. In this test, the following standards were applied to the composition evaluation.

Slipperiness of lather during washing:

AAA: lather is very slippery

BBB: lather is fairly slippery and does not adhere to the fingers

CCC: slipperiness of lather is normal

DDD: slipperiness of lather is inadequate causing the hair to tangle around the fingers and be brittle.

Feeling of lather during washing:

AAA: lather is abundant and creamy
BBB: lather is abundant, but not creamy
CCC: the amount of lather is adequate, but lather is coarse
DDD: lather is light and coarse

5 Slipperiness of hair after having been dried:

AAA: hair is very slippery
BBB: hair is moderately slippery
CCC: hair is slightly slippery
DDD: hair is not slippery but is rather brittle feelings

10 Glossiness of hair after having been dried:

AAA: glossiness of hair is clearly observed
BBB: hair appears glossy
CCC: hair appears slightly glossy
DDD: hair is not glossy

15 In the tables below, the figures of various ingredients for various compositions are expressed in % by weight. The compositions the numbers of which are asterisked are those of the present invention, and others represent comparative compositions.

25

30

35

40

45

50

55

Table 3

5		Composition Nos.								
		1	2*	3*	4*	5*	6*	7*	8*	9
	triethanolamine lauryl sulfate	20	20	20	20	20	20	20	20	20
10	diethanol amide of coconut oil fatty acid	3	3	3	3	3	3	3	3	3
15	Ester prepared in Example 1	-	2	-	-	-	-	-	-	-
	Ester prepared in Example 2	-	-	2	-	-	-	-	-	-
20	Ester prepared in Example 3	-	-	-	2	-	-	-	-	-
	Ester prepared in Example 4	-	-	-	-	2	-	-	-	-
25	Ester prepared in Example 5	-	-	-	-	-	2	-	-	-
	Ester prepared in Example 6	-	-	-	-	-	-	2	-	-
30	Ester prepared in Example 7	-	-	-	-	-	-	-	2	-
35	stearoyl lactylate (comparative product)	-	-	-	-	-	-	-	-	2
	ion exchanged water	77	75	75	75	75	75	75	75	75
40	<u>Lathering Characteristics</u>									
	<u>Slipperiness</u>	DDD	BBB	CCC	BBB	AAA	AAA	AAA	AAA	CCC
45	<u>Creaminess</u>	DDD	BBB	CCC	CCC	BBB	BBB	BBB	BBB	DDD
	<u>Finish</u>									
	<u>Slipperiness</u>	DDD	AAA	BBB	BBB	AAA	AAA	AAA	AAA	CCC
50	<u>Gloss</u>	DDD	BBB	BBB	BBB	AAA	BBB	BBB	AAA	DDD

55 Example 9

Shampoo compositions noted in Table 4 were prepared and were evaluated in the same manner as those in Example 8. The results are shown in Table 4. In this table and tables that follow, "POE" stands for

polyoxyethylene and the number in parenthesis immediately following "POE" designates the average number of moles of ethylene oxide in the polyoxyethylene.

Table 4

		Composition Nos.								
		10	11*	12*	13*	14*	15*	16*	17*	18
10	POE(3) sodium lauryl sulfate	20	20	20	20	20	20	20	20	20
15	diethanol amide of coconut oil fatty acid	3	3	3	3	3	3	3	3	3
	Ester prepared in Example 1	-	2	-	-	-	-	-	-	-
	Ester prepared in Example 2	-	-	2	-	-	-	-	-	-
	Ester prepared in Example 3	-	-	-	2	-	-	-	-	-
25	Ester prepared in Example 4	-	-	-	-	2	-	-	-	-
	Ester prepared in Example 5	-	-	-	-	-	2	-	-	-
30	Ester prepared in Example 6	-	-	-	-	-	-	2	-	-
	Ester prepared in Example 7	-	-	-	-	-	-	-	2	-
35	stearoyl lactylate (comparative product)	-	-	-	-	-	-	-	-	2
40	ion exchanged water	77	75	75	75	75	75	75	75	75
45	<u>Lathering Characteristics</u>									
	<u>Slipperiness</u>	DDD	BBB	BBB BBB	BBB	AAA	AAA	AAA	AAA	CCC
	<u>Creaminess</u>	DDD	BBB	CCC	CCC	BBB	BBB	BBB	BBB	DDD
50	<u>Finish</u>									
	<u>Slipperiness</u>	DDD	AAA	BBB	BBB	BBB	AAA	AAA	AAA	CCC
	<u>Gloss</u>	DDD	BBB	BBB	BBB	BBB	BBB	BBB	AAA	DDD

Example 10

Shampoo compositions in Table 5 were prepared and evaluated in the same manner as in Example 8. The results are shown in Table 5.

5	Table 5								
10		Composition Nos.							
15		19	20*	21*	22*	23*	24*	25	26
20	triethanolamine monolauryl phosphate	20	20	20	20	20	20	20	20
25	diethanol amide of coconut oil fatty acid	3	3	3	3	3	3	3	3
30	Ester prepared in Example 1	-	2	-	-	-	-	-	-
35	Ester prepared in Example 2	-	-	2	-	-	-	-	-
40	Ester prepared in Example 3	-	-	-	2	-	-	-	-
45	Ester prepared in Example 4	-	-	-	-	2	-	-	-
50	Ester prepared in Example 5	-	-	-	-	-	2	-	-
55	stearoyl lactylate (comparative product)	-	-	-	-	-	-	2	-
	methyl-branched isostearoyl lactylate (comparative product)	-	-	-	-	-	-	-	2
	ion exchanged water	77	75	75	75	75	75	75	75
	<u>Lathering Characteristics</u>								
	Slipperiness	DDD	AAA	BBB	BBB	AAA	AAA	CCC	CCC
	Creaminess	DDD	AAA	BBB	CCC	BBB	BBB	DDD	CCC
	<u>Finish</u>								
	Slipperiness	DDD	AAA	BBB	BBB	AAA	BBB	CCC	CCC
	Gloss	DDD	AAA	BBB	CCC	AAA	BBB	DDD	DDD

Example 11

Hair rinse compositions of formulations shown in Table 6 were prepared. Hair bundles each weighing 30 gm were washed using a plane shampoo and rinsed in the same manner as in Example 8. To each hair bundle the hair rinse compositions of 1 gm each were applied, which was rinsed again, and dried. Each bundle was subjected to organoleptic evaluation to determine product satisfaction by expert panelist. The results shown in Table 6 were obtained using the following standards for evaluating the compositions.

Slippery feelings:

AAA: slippery

BBB: slightly slippery

CCC: not slippery

Dampishness:

AAA: felt wet or dampish

BBB: slightly wet or dampish

CCC: not wet or dampish

Softness:

AAA: soft

BBB: slightly soft

CCC: not soft

Table 6

	Composition Nos.								
	27	28*	29*	30*	31*	32*	33*	34*	35
stearyl tri-methyl ammonium chloride	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2
stearyl alcohol	2	2	2	2	2	2	2	2	2
hydroxyethyl cellulose	-	-	-	-	-	-	0.3	0.3	-
Ester prepared in Example 1	-	1	-	-	-	-	-	-	-
Ester prepared in Example 2	-	-	1	-	-	-	-	-	-
Ester prepared in Example 3	-	-	-	1	-	-	-	-	-
Ester prepared in Example 4	-	-	-	-	1	-	-	-	-
Ester prepared in Example 5	-	-	-	-	-	1	-	-	-
Ester prepared in Example 6	-	-	-	-	-	-	0.5	-	-
Ester prepared in Example 7	-	-	-	-	-	-	-	0.5	-
stearyl lactylate (comparative product)	-	-	-	-	-	-	-	-	1
ion exchanged water	96.8	95.8	95.8	95.8	95.8	95.8	96.0	96.0	95.8
Slipperiness	CCC	AAA	AAA	AAA	AAA	AAA	AAA	AAA	BBB
Dampishness	CCC	AAA	CCC	BBB	AAA	AAA	AAA	AAA	CCC
Softness	CCC	AAA	BBB	BBB	AAA	AAA	AAA	AAA	BBB

Example 12

Creams with formulations shown in Table 7 were prepared and applied to arms of expert panelists to evaluate the same according to the following standards. The results are shown in Table 7.

Stickiness:

AAA: neat and refreshed

BBB: slightly sticky

CCC: sticky

Wetness or dampishness

AAA: dampish

BBB: slightly dampish

CCC: not dampish

Table 7

	Composition Nos.						
	36	37*	38*	39*	40*	41*	42
liquid paraffin	13	13	13	13	13	13	13
Tween 60	5	5	5	5	5	5	5
Leodol MS 60	1	1	1	1	1	1	1
hydrogenated castor oil	2	2	2	2	2	2	2
Ester prepared in Example 1	-	2	-	-	-	-	-
Ester prepared in Example 2	-	-	2	-	-	-	-
Ester prepared in Example 3	-	-	-	2	-	-	-
Ester prepared in Example 4	-	-	-	-	2	-	-
Ester prepared in Example 5	-	-	-	-	-	2	-
stearoyl lactylate (comparative product)	-	-	-	-	-	-	2
ion exchanged water	79	77	77	77	77	77	77
Feelings when creams were applied							
Stikilessness	CCC	AAA	BBB	BBB	AAA	AAA	CCC
Dampishness	CCC	AAA	AAA	AAA	AAA	AAA	CCC

Example 13

Shampoo compositions of Table 8 were prepared and evaluated in the same manner as in Example 8. The results are shown in Table 8.

Table 8

6		Composition Nos.			
		43	44 *	45 *	46 *
10	POE (2.5) laurylether sulfate	15	15	15	15
	coconut oil fatty acid diethanol amide	5	5	5	5
15	Ester prepared in Example 4	0.01	0.1	1.0	2.0
	Emanon 3299R (polyoxyethylene distearate)	1.2	1.2	1.2	1.2
	citric acid	0.5	0.5	0.5	0.5
	methyl paraben	0.1	0.1	0.1	0.1
25	perfume	small amount	small amount	small amount	small amount
	coloring agent	small amount	small amount	small amount	small amount
30	ion exchanged water	balance	balance	balance	balance
35	<u>Feelings of lather</u>				
	Slippery feeling	BBB	AAA	AAA	BBB
	Creamy feeling	CCC	AAA	AAA	BBB
40	<u>Finishness</u>				
	Slipperiness	CCC	AAA	AAA	BBB
	Gloss	BBB	AAA	AAA	AAA
45					

Example 14

Shampoo compositions of Table 9 were prepared and evaluated in the same manner as in Example 8. The results are shown on Table 9.

Table 9

5		Composition Nos.			
		47	48 *	49 *	50 *
10	triethanolamine lauryl sulfate	20	20	20	20
	coconut oil fatty acid diethanol amide	5	5	5	5
15	Ester prepared in Example 4	0.01	0.1	1.0	2.0
	Methorose 60 SH1000 (methyl cellulose)	0.5	0.5	0.5	0.5
20	urea	6	6	6	6
	perfume	small amount	small amount	small amount	small amount
25	coloring agent	small amount	small amount	small amount	small amount
	ion exchanged water	balance	balance	balance	balance
30	<u>Feelings of lather</u>				
	Slippery feeling	BBB	BBB	AAA	BBB
35	Creamy feeling	CCC	BBB	AAA	BBB
	<u>Finishness</u>				
	Slipperiness	CCC	AAA	AAA	BBB
40	Gloss	CCC	AAA	AAA	AAA

45
Example 15

The Hair rinse compositions of Table 10 were prepared and evaluated in the same manner as in Example 11. The results are shown on Table 10.

50


55

Table 10

	Composition Nos.			
	51	52 *	53 *	54 *
stearyl trimethyl ammonium chloride	1.2	1.2	1.2	1.2
stearyl alcohol	2	2	2	2
Ester prepared in Example 4	0.01	0.1	1.0	2.0
Cellosize QP52000 (hydroxyethyl cellulose)	0.3	0.3	0.3	0.3
perfume	small amount	small amount	small amount	small amount
coloring agent	small amount	small amount	small amount	small amount
ion exchanged water	balance	balance	balance	balance
Slippery feeling	CCC	BBB	BBB	BBB
Dampishness	DDD	CCC	BBB	BBB
Softness	DDD	CCC	BBB	BBB

Claims


1. A fatty acid-lactic acid ester represented by the formula:



in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5.

2. A shampoo composition comprising:

(A) 0.1 to 5% by weight of a fatty acid-lactic acid ester represented by the formula:




in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5; and

(B) 5 to 50% by weight of an anionic surface active agent.

3. A hair rinse composition comprising:

(A) 0.1 to 5% by weight of a fatty acid-lactic acid ester represented by the formula:



in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5; and


(C) 0.1 to 10% by weight of a quaternary ammonium salt.

Table 10

	Composition Nos.			
	51	52 *	53 *	54 *
stearyl trimethyl ammonium chloride	1.2	1.2	1.2	1.2
stearyl alcohol	2	2	2	2
Ester prepared in Example 4	0.01	0.1	1.0	2.0
Cellose QP52000 (hydroxyethyl cellulose)	0.3	0.3	0.3	0.3
perfume	small amount	small amount	small amount	small amount
coloring agent	small amount	small amount	small amount	small amount
ion exchanged water	balance	balance	balance	balance
Slippery feeling	CCC	BBB	BBB	BBB
Dampishness	DDD	CCC	BBB	BBB
Softness	DDD	CCC	BBB	BBB

Claims


1. A fatty acid-lactic acid ester represented by the formula:



in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5.

2. A shampoo composition comprising:

(A) 0.1 to 5% by weight of a fatty acid-lactic acid ester represented by the formula:




in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5; and

(B) 5 to 50% by weight of an anionic surface active agent.

3. A hair rinse composition comprising:

(A) 0.1 to 5% by weight of a fatty acid-lactic acid ester represented by the formula:




in which R_1 and R_2 independently represent alkyl groups of a C_{4-18} carbon atom content, X represents a hydrogen atom, alkali metal, alkaline earth metal, ammonium group, alkanol ammonium group, or lower alkyl ammonium group, and n represents a value of 1 to 5; and

(C) 0.1 to 10% by weight of a quaternary ammonium salt.

Revendications

1. Ester d'acide lactique d'un acide gras, représenté par la formule :




dans laquelle :

- R₁ et R₂ représentent indépendamment des groupes alkyle ayant une teneur de 4 à 18 atomes de carbone ;
- X représente un atome d'hydrogène, un métal alcalin, un métal alcalino-terreux, un groupe ammonium, un groupe alcool ammonium, ou un groupe alkyl inférieur ammonium ; et
- n représente une valeur de 1 à 5.

2. Composition de shampoing comprenant :

(A) 0,1 à 5% en poids d'un ester d'acide lactique d'un acide gras, représenté par la formule :




dans laquelle :

- R₁ et R₂ représentent indépendamment des groupes alkyle ayant une teneur de 4 à 18 atomes de carbone ;
- X représente un atome d'hydrogène, un métal alcalin, un métal alcalino-terreux, un groupe ammonium, un groupe alcool ammonium, ou un groupe alkyl inférieur ammonium ; et
- n représente une valeur de 1 à 5 ; et

(B) 5 à 50% en poids d'un agent tensio-actif anionique.

3. Composition de rinçage capillaire comprenant :

(A) 0,1 à 5% en poids d'un ester d'acide lactique d'un acide gras, représenté par la formule :



dans laquelle :


- R₁ et R₂ représentent indépendamment des groupes alkyle ayant une teneur de 4 à 18 atomes de carbone ;
- X représente un atome d'hydrogène, un métal alcalin, un métal alcalino-terreux, un groupe ammonium, un groupe alcool ammonium ou un groupe alkyl inférieur ammonium ; et
- n représente une valeur de 1 à 5 ; et

(C) 0,1 à 10% poids d'un sel d'ammonium quaternaire.

Patentansprüche

1. Fettsäure-milchsäureester, dargestellt durch die Formel:






in der R_1 und R_2 unabhängig für Alkylgruppen mit einem C_4-18 -Kohlenstoffatomgehalt stehen, X für Wasserstoffatom, Alkalimetall, Erdalkalimetall, Ammoniumgruppe, Alkanolammoniumgruppe oder Niederalkylammoniumgruppe steht und n für eine Zahl von 1 bis 5 steht.

2. Schampoozusammensetzung, umfassend:


(A) 0,1 bis 5 Gew.% eines Fettsäure-milchsäureesters, dargestellt durch die Formel:



in der R_1 und R_2 unabhängig für Alkylgruppen mit einem C_4-18 -Kohlenstoffatomgehalt stehen, X für Wasserstoffatom, Alkalimetall, Erdalkalimetall, Ammoniumgruppe, Alkanolammoniumgruppe oder Niederalkylammoniumgruppe steht und n für eine Zahl von 1 bis 5 steht; und
(B) 5 bis 50 Gew.% eines anionischen oberflächenaktiven Mittels.

3. Haarspülzusammensetzung, umfassend

(A) 0,1 bis 5 Gew.% eines Fettsäure-milchsäureesters, dargestellt durch die Formel:



in der R_1 und R_2 unabhängig für Alkylgruppen mit einem C_4-18 -Kohlenstoffatomgehalt stehen, X für Wasserstoffatom, Alkalimetall, Erdalkalimetall, Ammoniumgruppe, Alkanolammoniumgruppe oder Niederalkylammoniumgruppe steht und n für eine Zahl von 1 bis 5 steht; und
(C) 0,1 bis 10 Gew.% eines quaternären Ammoniumsalzes.

