

GLUTAMINASE

The present invention relates to a new salt-resistant and thermotolerant glutaminase from *Lactobacillus* and a process for the production of such salt-resistant thermotolerant glutaminase from *Lactobacillus*.

Hydrolysed proteins have been known for use as seasonings in the form of soy sauce, which traditionally has been prepared by enzymatic hydrolysis requiring a long period of time for preparation. In producing soy sauce, protein-containing plant materials are inoculated with *Aspergilli* and the solid culture is fermented for 2 days to make koji. During this time, enzymes are produced which are able to hydrolyse proteins into peptides and amino acids in the moromi stage. The koji is mixed with a solution of common salt and water to give moromi, which is fermented for 4 weeks to 8 months by the activity of micro-organisms such as halophilic lactic acid bacteria and yeasts, and the soya sauce is obtained by removing the solids portion from the fermented moromi. Such fermentation is known as a process for producing glutamic acid-rich flavouring foods by enzymatically degrading proteins materials in the presence of high concentration of salt. In this case, the presence of salt is intended to prevent decomposition and putrefaction resulting from microbial contamination.

Glutaminase is known as playing an important role in such processes, particularly in the moromi stage, to enzymatically convert glutamine to glutamic acid (or
5 glutamate) by deamination. This enzyme is regarded as a key enzyme that controls the savoury and umami taste of such kind of fermented products like soy sauce, for example. In the traditional soy sauce or biohydrolysate manufacturing processes, glutaminase comes from
10 *Aspergillus* strains inoculated in the koji. However glutaminase from these moulds are reportedly being inhibited by salt concentrations above 10-15% (w/v) which are the practical conditions for soy sauce production, in the moromi stage. It is thus necessary
15 to use a salt-resistant glutaminase in order to achieve efficient glutamic acid production.

Furthermore, enzymatic degradation of protein at high temperature is very effective for raising the rate of
20 proteins degradation on the one hand, and preventing contamination by micro-organisms on the other hand. Since glutamine can be converted to tasteless pyroglutamic acid under increased temperature conditions in the absence of glutaminase the necessity
25 of a thermotolerant glutaminase is desirable.

The main reason for the lower glutamic acid content in the enzymatic degradation method is that glutaminase is

significantly inhibited by either high salt concentration or high temperature.

Accordingly it is thus important to have a glutaminase
5 that is salt-resistant and thermotolerant.

Glutaminase from *Candida famate* (Japanese Patent Publication N° 38748/1994) and from *Cryptococcus albidus* (Japanese Patent Publication N° 48759/1974) are
10 disclosed as having salt resistance and thermostability. However the relative activity in the presence of 18% salt of both enzymes is as low as about 50 to 60 % with respect to 100% activity in the absence of salt.

15

US 6063409 discloses a glutaminase from *Cryptococcus nodaensis* which exhibits higher salt tolerance and better temperature resistance.

20 Besides the fact that all these glutaminases present more or less the desired features of salt-resistance and thermotolerance, they all originates from micro-organism that generally are pathogen for human and are therefore not recognised as safe nor as food-grade. The
25 use of cultures of such micro-organisms is impossible without extended purification in order to recover enzymes. These kind of manipulation are cumbersome and would make impossible the use of the micro-organism

directly, either dead or alive, in a food product or at any step of the manufacture process of a food product.

Accordingly, the object of the present invention is to
5 provide a glutaminase that exhibits salt-resistance and thermo-tolerance and that comes from a micro-organism that has a long history of use in food process and that is therefore food-grade.

10 To this end, the present invention provides a new salt-resistant and thermotolerant glutaminase from *Lactobacillus* which has a temperature dependence such that it retains more than 80% of its activity between 45°C and 55°C, a salt dependence such that its activity
15 in the presence of up to 15% NaCl is from 80 to 180 % of that without NaCl and a thermal stability such that its energy of activation is about 80 to 150 kJ.mol⁻¹, preferably 90 to 120 kJ.mol⁻¹, more preferably 100 to 110 kJ.mol⁻¹.

20

In the drawings :

Figure 1 shows the pH dependence of the glutaminase where the vertical axis define the relative activity
25 (R.A.) expressed as percentage of the maximum activity in the indicated pH range.

Figure 2 shows the temperature dependence of the glutaminase where the vertical axis define the relative

activity (R.A.) expressed as percentage of the maximum activity in the temperature range $T(^{\circ}\text{C})$.

Figure 3 shows the salt dependence of the glutaminase
5 where the vertical axis define the relative activity (R.A.) expressed as percentage of the maximum activity in the NaCl concentration range in percent w/v(NaCl).

Figure 4 exhibits the thermal stability of the
10 glutaminase and shows :

A - the change of the residual activity (R.A.) as a function of incubation time $t(\text{min.})$ at (+) 45°C , (o) 55°C , (Δ) 60°C , (■) 65°C and (●) 75°C

15 B - Arrhenius plot of the neperian logarithm of rate constant of thermal inactivation $k(\text{min}^{-1})$ as a function of the reciprocal of temperature ($1/\text{K}$).

20 The present invention is described in more detail hereafter.

The present salt-resistant thermotolerant glutaminase may be produced by a process which comprises
25 cultivating a micro-organism belonging to the genus *Lactobacillus*, preferably *Lactobacillus rhamnosus* having the ability to produce the salt-resistant thermotolerant glutaminase thereby producing the salt-

resistant thermotolerant glutaminase and then recovering it.

A *Lactobacillus rhamnosus* strain having the required ability was deposited by way of example under the
5 Budapest Treaty at the Collection National de Cultures de Microorganismes (CNCM), Institut Pasteur, 28 rue du Docteur Roux, 75724 Paris Cedex 15, France, on April 5, 2000 under the reference CNCM I-2433.

10 Usually, the cultivation process is advantageously carried out in a liquid medium in jar, flask or bioreactor, for example. The temperature of the medium may range from 35°C to 55°C, and the cultivation time may be of 6 hours or more.

15 The culture medium may include as carbon source, monosaccharides and oligosaccharides such as glucose, maltose, dextrans and other carbon source from vegetal origin such as wheat, corn, wheat gluten or soya for example. The culture medium may include as nitrogen
20 source includes peptides, proteins and amino acids coming from from vegetal origin such as wheat, corn, wheat gluten or soya as well, for example.

The culture medium obtained after cultivation may be used as such for enzyme activity characterisation but
25 may also be subjected to solid/liquid separation and purification. The culture medium may also be added into a protein containing medium for the glutaminase to deaminates the glutamine contained. However since the salt-resistant thermotolerant glutaminase according to

the present invention is membrane-bound a further step may be necessary to release the enzyme from the cells in order to further characterise it and/or to increase its activity. Thus, conventional enzyme extraction methods may be applied. For example, it is possible to use mechanical grinding and disruption, sonication or ultrasonication, use of detergents, lysis of the cells using lysozyme or other kind of method allowing release of the enzyme from the membrane. Hence, by these techniques, a crude enzyme solution of the salt-resistant thermotolerant glutaminase may be prepared. In order to obtain a more purified solution, the crude enzyme preparation may be subjected to a suitable combination of chromatographic adsorption-elution techniques using various kind of ion exchangers, for example DEAE cellulose, TEAE cellulose, QAE Sephadex or Hydroxyapatite, gel filtration using Sephadex or Superdex whereby a purified enzyme preparation may be obtained.

Another aspect of the present invention concerns the use of a mixture containing a salt-resistant thermotolerant glutaminase from *Lactobacillus* in a process for producing a biohydrolysate. Specifically, a culture medium containing the salt-resistant thermotolerant glutaminase according to the present invention may be added in the step of mixing or in the step of fermentation and/or ageing of a process for producing biohydrolysates starting from proteinaceous material such as soy sauce process or

wheat gluten hydrolysate. Such culture medium may be the cell-containing medium obtained after cultivation of *Lactobacillus rhamnosus* (CNCM I-2433). This cell-containing medium or cell-fractions-containing medium
5 may also be concentrated by liquid-solid separation means in order to recover cell-containing residue. Advantageously, the medium used may also be a mixture in which the enzyme has been released from the cells by sonication, lysis, or any appropriate means.

10 Indeed, thermally denatured proteinaceous materials (soybeans, wheat gluten, ...) are mixed with thermally denatured starchy materials (rice, wheat, corn, ...) and after the water content of the mixture is adjusted to 30-45%, a koji mould is inoculated into it
15 and cultivated in an usual manner to give solid koji which is then mixed with brine and subjected to fermentation and ageing. However, the step of hydrolysing of a protein-containing material may be performed using either endogenous enzymes from koji or
20 technical added enzymes in the presence of a *Lactobacillus* providing a glutaminase activity, preferably a *Lactobacillus rhamnosus*, more preferably *Lactobacillus rhamnosus* (CNCM I-2433).

An other aspect of the present invention concerns the
25 use of a mixture containing a salt-resistant thermotolerant glutaminase from *Lactobacillus* in a process for producing a biohydrolysate. Advantageously, the biohydrolysate is a soy sauce or wheat gluten hydrolysate. The salt-resistant thermotolerant

glutaminase according to the present invention may be use at the step of hydrolysing and/or at the step of ageing in the biohydrolysate process production.

The new salt-resistant thermotolerant glutaminase
5 (purified or cell-bound) according to the present invention may thus be added in a process for producing food and drink containing peptides and/or amino acids as a part or a majority of tasting components. In such a process protein materials or protein-containing
10 materials are hydrolysed with at least one proteolytic enzyme or a proteolytic enzyme-containing material, such as a koji for example. Due to its high salt-resistance and thermotolerance, when the present enzyme is used in a process including protein hydrolysis at
15 high temperature and high salt concentration in producing glutamic acid-rich seasoning, for example, foods significantly reinforced with glutamic acid as a tasting component may be obtained efficiently.

20 **Glutaminase activity test**

Glutaminase activity is tested by determining the formation of L-glutamate from L-glutamine using a colorimetric method (kit Boehringer Mannheim, n°139
092, Darmstadt, Germany). The reaction mixture,
25 containing 0.5 ml of 60 mM L-glutamine in 50 mM sodium phosphate buffer pH 7.0 is preincubated at 37°C for 3 min and the reaction is started by adding 0.5 ml of the enzyme containing preparation to be tested. The incubation is carried out at 37°C for 20 to 30 min. The

reaction is stopped by cooling the samples on ice. The amount of glutamate produced is subsequently enzymatically determined.

One unit (U) of glutaminase activity is defined as the amount of enzyme that liberates 1 μ mol of glutamate per min under the above conditions.

Measurement of pH-dependence

The pH-dependence of glutaminase activity was determined over the pH-range 4-9. Glutaminase activity was tested using 100 mM buffer indicating various pHs and is illustrated in Figure 1. After incubation of the enzyme solution with a final concentration of glutamine of 30 mM in 100 mM buffer for 30 min at 37°C, glutaminase activity was determined as described in the section « glutaminase activity test ».

Buffer used : pH 4 to 6.5 : 100 mM citrate/ 200 mM sodium phosphate

pH 7 and 7.5 : 100 mM sodium phosphate

pH 8 and 9 : 100 mM Tris/ HCl

Each experiment point is the average of two determinations and the activities are expressed as percentages of the maximum activity observed in the pH range 4-9.

The optimum pH value for the glutaminase according to the present invention is about pH 7.0. The enzyme is less active at lower pH values since around 62 % of the activity is lost at pH about 6.0. On the contrary, the enzyme has greater activity in alkaline than acidic pH

because about 74 and 69 % of activity still remains at pH 8 and 9 respectively.

Temperature dependence

- 5 The temperature dependence of glutaminase activity was determined by incubating the enzyme with substrate for 30 min at different temperatures in the range of 25 to 80°C as shown in Figure 2. Glutaminase activity was determined as described in « glutaminase assays ».
- 10 The optimal temperature for glutaminase activity is around 50°C and the enzyme retains more than 90% of its activity at about 45°C corresponding to the temperature of the cultivation conditions. It is observed that about 50% of activity still remains at 70°C.

15

Salt dependence

- The salt dependence of glutaminase was determined by adding 0-20% (w/v) NaCl to the reaction medium. Relative activities of the enzyme were calculated and
- 20 expressed as percentage of the activity observed without salt (Figure 3). Glutaminase activity is increased in the presence of NaCl concentrations lower than 10%, in particular 1.8-fold improved at 2.5% of NaCl. Furthermore, in the presence of 15% of salt the
- 25 enzyme activity is about 90% of that without salt.

Thermal stability

Thermal stability of glutaminase was determined between
5 45 and 75°C (Figure 4).

Following heat treatment at 45°C, there is no thermal
inactivation. Treatments at $T \geq 55^\circ\text{C}$ induce a
progressive inactivation of the enzyme and the
glutaminase is completely inactivated on incubation at
10 75°C for about 15 min (Figure 4A).

The apparent inactivation rate constant k for thermal
inactivation was plotted against the reciprocal of
absolute temperature (Figure 4B). As may be seen, the
15 temperature dependence of the inactivation constant
followed the Arrhenius equation. The energy of
activation E_a calculated from the slope of the
Arrhenius plot was about 90 to 110 $\text{kJ}\cdot\text{mol}^{-1}$. It is in
line with those obtained for most thermotolerant
20 enzymes.

The results clearly show that the thermal stability of
glutaminase decreases linearly with increasing
temperature from 55 to 75°C.

25

Catalytic parameters

Glutaminase-catalysed deamination of glutamine was
tested in 50 mM sodium phosphate buffer, pH 7.0, at

37°C using 12 different concentrations of glutamine ranging from 0.5 to 80 mM. Substrate Inhibition was not significant in the glutamine concentration range used. The catalytic behaviour of glutaminase with glutamine
5 can be described by Michaelis-Menten equation (2). The catalytic parameters were calculated by weighted linear regression using Lineweaver-Burk inversion and by non-linear regression of Michaelis-Menten eqn (2) :

$$10 \quad v = \frac{V_{\max} * [S]}{K_m + [S]} \quad (2)$$

Where v and $V_{\max}$ are the initial and maximal velocity of the reaction, $[S]$ is the substrate concentration and K_m is the Michaelis-Menten constant.

15

The result are in harmony with Michaelis-Menten kinetics. From the intercept on the abscissa of Lineweaver-Burk plot, the K_m value for the enzymatic reaction was estimated to be 5.7 mM and $V_{\max} = 108$ U/l
20 of hydrolysate. The non-linear regression of eqn (2) yielded K_m and $V_{\max}$ values of 4.8 ± 0.4 mM and 101 ± 2 U/l, respectively.

So we can conclude that the K_m value for the glutaminase of the present invention is from about 4 to
25 6 mM and V_m is from about 100 to 110 U/l of hydrolysate.

EXAMPLE I**Cultivation of *Lactobacillus***

A culture of *Lactobacillus rhamnosus* (CNCM I-2433)
5 containing $5 \cdot 10^8$ cfu/ml is grown in a medium containing
22% (w/w) wheat gluten (Roquette, Lestern, France),
0.1% alcalase (Novozymes, Bagsvaerd, Denmark), 4% (w/w)
Flavorzyme L1000 (Novozymes, Bagsvaerd, Denmark) and 30
mM sodium acetate buffer pH 5.8 for 12h at 45°C. The
10 broth is then cooled at 4°C and immediately subjected
to analysis.

EXAMPLE II**Enzyme preparation**

15 After cultivation, the medium is washed 3 times with 50
mM sodium phosphate buffer pH 7.0 and finally
reconstituted in the same buffer containing 10% (w/v)
sucrose and 1 mM PMSF. Because the glutaminase is
membrane-bound, the hydrolysate was subjected to 1
20 mg/ml lysozyme for 1h at 37°C. This solution was
stored at 4°C and used for enzyme assays.

EXAMPLE III**Production of an improved soy sauce.**

25 0.2 kg of Koji mould was inoculated into a mixture of
1.2 kg of defatted and crushed soybeans, 0.3 kg of
wheat gluten, 1.2 kg of salt and 3.4 kg of water to
form a moromi according to a conventional process for
producing soy sauce koji. In a separate step, the

medium obtained according to example I was centrifuged at 2000xg for 10 min. The resulting supernatant was discarded and the cell-containing residue was added to the Moromi above. The mixture thus obtained was
5 subjected to a traditional moromi fermentation process and in that way, the salt-resistant thermotolerant glutaminase accumulated in the cell-containing residue of medium-example I deaminates the glutamine present in Moromi during the time required to achieve a
10 satisfactory yield of glutamate and to produce a mature soy sauce.

CLAIMS

1. A salt-resistant thermotolerant glutaminase from a micro-organism belonging to the genus *Lactobacillus*.

5

2. A salt-resistant thermotolerant glutaminase according claim 1, in which the *Lactobacillus* is *Lactobacillus rhamnosus*.

10 3. A salt-resistant thermotolerant glutaminase according to claim 1, in which the *Lactobacillus rhamnosus* is *Lactobacillus rhamnosus* (CNCM I-2433).

4. A salt-resistant thermotolerant glutaminase
15 according to claim 1, which has a temperature dependence such that it retains more than 80% of its activity between 45°C and 55°C, a salt dependence such that its activity in the presence of up to 15% NaCl is from 80 to 180 % of that without NaCl and a thermal
20 stability such that its energy of activation is about 80 to 150 kJ.mol⁻¹.

5. A Process for producing a salt-resistant thermotolerant glutaminase, which comprises cultivating
25 a micro-organism belonging to the genus *Lactobacillus* having the ability to produce a salt-resistant thermotolerant glutaminase, thereby producing the salt-resistant thermotolerant glutaminase and then recovering it.

6. A Process according to claim 5, in which the *Lactobacillus* is *Lactobacillus rhamnosus* (CNCM I-2433).
- 5 7. Use of a mixture containing a salt-resistant thermotolerant glutaminase according to claim 1 in a process for producing a biohydrolysate.
8. Use according to claim 7, in which the mixture is a
10 *Lactobacillus* cell-containing medium or cell-fractions-containing medium.
9. Use according to claim 7, in which the
15 biohydrolysate is a soy sauce or a wheat gluten hydrolysate.