

ANTICOCCIDIAL COMPOSITIONS

This invention relates to anticoccidial compositions, such as feedstuffs or premixes, destined for consumption by avian species, particularly poultry such as chickens or turkeys, and which compositions contain as the active ingredients a combination of an antibiotic and a urea derivative.

Coccidiosis has long constituted a problem of major commercial significance in the rearing of poultry. If untreated, the severe forms of the disease lead to poor weight gain, reduced feed efficiency and high mortality. Although many agents have been discovered which are, to a greater or lesser extent, effective against the Eimeria - the genera of protozoans principally responsible for coccidiosis - it is unfortunately the case that the Eimeria show a propensity to develop drug resistant strains, see Advances in Pharmacology and Chemotherapy, II, 221-293 (1973). There is thus a continuing need for the provision of new chemotherapeutic regimes capable of controlling this disease.

Polyether antibiotics are known to be effective in the control of coccidiosis, see for example U.S. Patent No. 3,501,563, as are carbanilide derivatives, see for example U.S. Patents Nos. 2,731,382 and 3,284,433. However, heretofore, it has not been appreciated that a combination of these substances exhibits surprisingly enhanced activity (i.e. synergism and potentiation) against coccidiosis-producing strains of Eimeria.

Accordingly, the present invention provides an anticoccidial composition for consumption by an avian species; characterized in that the composition comprises as active ingredients a polyether antibiotic and a carbanilide derivative associated with a carrier or feedstuff.

The polyether antibiotics are a class of antibiotics produced by the Streptomyces genus of microorganisms. They comprise a multiplicity of cyclic ethers in their structures. The class is reviewed in Kirk-Othmer: Encyclopedia of Chemical Technology, Vol. 3, Third Edition (John Wiley & Sons, Ind., 1978), page 47 et seq.; in Annual Reports in Medicinal Chemistry Volume 10 (Academic Press, N.Y. 1975), page 246 et seq.; and in J. Chrom. Lib., Vol. 15 (Elsevier Scientific Publishing Co., N.Y. 1978), page 488 et seq.


Like other products of fermentation origin, many of the polyether antibiotics comprise more than one factor. The various factors are all usable in the present invention. Further, many of these antibiotics readily form ethers, esters or salts, which are either active as such or are converted in vivo to the basic antibiotic. Such derivatives can also be employed in the present invention. All that is necessary is that an active moiety of a polyether antibiotic be delivered in vivo.

The polyether antibiotic for use in the composition of the invention is one of the following: monensin (factors A, B, and C), laidlomycin, grisorixin, lenoremycin, salinomycin, narasin, lonomycin,

alborixin, antibiotic A204, A32887 (K41), etheromycin, lasalocid (factors A, B, C, D, and E), isolasalocid A, lysocellin, mutalomycin, and antibiotic A23187.


5 Preferred polyether antibiotics include mutalomycin, monensin, narasin, lasalocid, salinomycin A204 and lonomycin, and especially monensin, narasin, lasalocid, salinomycin and A-204. Monensin is the presently preferred polyether antibiotic for use in the invention, particularly in the commercially available form, i.e. that form consisting of factor A and a minor amount of factor B.

10 The term "carbanilide derivative" as used herein refers to that particular class of urea derivatives containing the moiety:




20 where X is oxygen or sulfur and where each phenyl ring may be substituted. The term also embraces simple complexes of these ureas, for example the complex formed between 4',4-dinitrocarbanilide and 2-hydroxy-4,6-dimethylpyrimidine and known as "nicarbazin".

25 Preferred carbanilide derivatives are those which possess the structural formula (I):



30

where R^1 , R^2 and R^3 are the same or different and can each represent hydrogen; halogen; cyano; amino; nitro; C_{1-6} alkyl; C_{1-4} alkoxy; C_{2-4} alkanoylamino; C_{1-4} alkylthio; C_{1-4} haloalkyl; C_{1-4} haloalkoxy; C_{1-4} haloalkylthio;
 5 C_{1-6} alkoxycarbonyl; C_{2-4} haloalkenyloxy; C_{1-4} alkoxycarbonylthio; C_{1-4} alkylsulfonyl; C_{1-4} haloalkylsulfonyl; fluorosulfonyl; phenoxy optionally substituted by halogen, C_{1-4} haloalkyl or nitro; phenoxycarbonyl optionally substituted by C_{1-4} alkyl; phenyl; or phenyl-
 10 sulfonyl optionally substituted by nitro, acetamido, isopropylideneamino or a group of the formula



where R^9 is C_{1-4} haloalkylthio or C_{2-4} haloalkenyloxy; R^4 and R^5 are the same or different and can each represent hydrogen or C_{1-4} alkyl;
 20 R^6 , R^7 and R^8 are the same or different and can each represent hydrogen; halo; cyano; amino; C_{2-4} alkanoylamino; N- C_{1-4} alkyl C_{2-4} alkanoylamino; nitro; C_{1-6} alkyl; C_{1-4} alkoxy; C_{1-4} alkylthio, C_{1-4} haloalkyl; C_{1-4} haloalkoxy, C_{1-4} haloalkylthio, C_{2-4} haloalkenyloxy; C_{1-4} haloalkylsulfonyl; C_{1-6} alkoxycarbonyl, aminosulfonyl or
 25 benzoyl; provided that at least one of R^1 , R^2 , R^3 , R^6 , R^7 and R^8 is other than hydrogen, and complexes of such compounds with 2-hydroxy-4,6-dimethylpyrimidine.
 30

It is preferred that there be substituents at the para-positions of the phenyl rings of the carbanilide moieties, particularly for electronegative substituents, meta-substitution is less preferred and ortho-substitution least preferred. In general, compounds having amino substitution are not very active and the acyl derivatives of such compounds are preferably utilized.

X is preferably oxygen.

Many of the carbanilides of value in the present invention are known, some also being described as possessing anticoccidial activity.

The following list illustrates specific carbanilides of value in the invention and includes, where known and or appropriate, relevant literature references:

Group A

- A.1. 3,3'-bis(trifluoromethyl)-4,4'-dichlorocarbanilide (CAS registry #370-50-3; C.A. 68:2655v)
- A.2. 3,3',5,5'-tetrakis(trifluoromethyl)carbanilide (CAS registry #3824-74-6; C.A. 66: P646444, C.A. 68: 2655v, and C.A. 71: P91052y)
- A.3. 3-chloro-4'-fluorocarbanilide (C.A. 65: 12792c)
- A.4. 2-chloro-2'-fluorocarbanilide
- A.5. 3,4',5-tris(trifluoromethyl)carbanilide (CAS registry #23747-71-9; C.A. 71: P91056c)
- A.6. 3,4,5-trichlorocarbanilide


- A.7. 2-chloro-5-(trifluoromethyl)-4'-(ethoxycarbonyl)carbanilide
- A.8. 2,6-dimethyl-4'-(N-methylacetamido)carbanilide
- 5 A.9. 2-methoxy-4'-acetamidocarbanilide
- A.10. 3-(trifluoromethyl)-4'-iodo-thiocarbanilide
- A.11. 2-fluoro-4'-(aminosulfonyl)carbanilide
- A.12. 2-methoxy-4'-isopropylcarbanilide
- 10 A.13. 2-methyl-2',5'-diethoxycarbanilide
- A.14. 4-ethyl-2'-methoxycarbanilide
- A.15. 2-methyl-5-chloro-2',5'-dimethoxycarbanilide
- A.16. 2,4,4'-trimethyl-3'-nitrocarbanilide
- 15 A.17. 2-amino-3-nitro-5-(trifluoromethyl)-2',4'-dimethylcarbanilide
- A.18. 2-amino-3,4'-dinitro-5-(trifluoromethyl)-2'-chlorocarbanilide
- 20 A.19. 2-amino-3,5'-dinitro-5-(trifluoromethyl)-2'-fluorocarbanilide
- A.20. 2-amino-3-nitro-5-(trifluoromethyl)-2'-(ethoxycarbonyl)carbanilide
- 25 A.21. 2-ethyl-6-sec-butylcarbanilide
- A.22. 2-isopropyl-2',4',6'-trimethylcarbanilide
- A.23. 2-amino-3-nitro-3',5-bis(trifluoromethyl)-4'-chlorocarbanilide
- 30 A.24. 2-ethyl-6-sec-butyl-4'-n-butoxycarbanilide

- A.25. 2-amino-3-nitro-5-(trifluoromethyl)-2',4',5'-trichlorocarbanilide
- A.26. 2-amino-3,3'-dinitro-5-(trifluoromethyl)-4'-chlorocarbanilide
- 5 A.27. 2-amino-3-nitro-5-(trifluoromethyl)-2'-methyl-4'-bromocarbanilide
- A.28. 2-amino-3-nitro-5-(trifluoromethyl)-2',6'-dibromo-4'-fluorocarbanilide
- 10 A.29. 2-amino-3-nitro-2',5-bis(trifluoromethyl)-4'-chlorocarbanilide
- A.30. 3,3',5,5'-tetrakis(trifluoromethyl)thiocarbanilide (CAS registry #1060-92-0; C.A. 66: 64644H)
- 15 A.31. 2,4-dimethoxy-4'-(ethoxycarbonyl)carbanilide
- A.32. 4-(ethoxycarbonyl)-2'-methyl-6'-ethylcarbanilide
- A.33. 2,2'-dinitro-4,4'-bis(trifluoromethyl)carbanilide (CAS registry #16588-81-1; C.A. 68: 2655v)
- 20 A.34. 3,3',4,4',5,5'-hexachlorocarbanilide
- A.35. 3-nitro-4-chloro-4'-(trifluoromethyl)carbanilide
- 25 A.36. 4-chloro-3,4'-bis(trifluoromethyl)carbanilide (CAS registry #23747-70-8; C.A. 71: P91056c)
- A.37. 4,4'-dinitro-2,2'-bis(trifluoromethyl)carbanilide (CAS registry #16588-84-4; C.A. 68: 2655v)
- 30

- A.38. 4,4'-bis(trifluoromethyl)carbanilide (CAS registry #1960-88-9; C.A. 71: 91056c)
- A.39. 3-bromo-3',5'-dimethylcarbanilide
- 5 A.40. 2,5-dichloro-4'-methyl-N²-ethylcarbanilide
- A.41. 2,5-dichloro-2',4'-difluorocarbanilide
- A.42. 2-amino-3-nitro-3',5,5'-tris(trifluoromethyl)-carbanilide
- 10 A.43. 2,6-diethyl-4'-(ethoxycarbonyl)carbanilide
- A.44. 3-ethyl-3'-chloro-4'-methyl-N²-ethylcarbanilide
- A.45. 2,6-dimethyl-4'-(ethoxycarbonyl)carbanilide
- A.46. 4-methoxy-3'-acetamidocarbanilide
- 15 A.47. 2-methoxy-4'-(n-butoxycarbonyl)carbanilide
- A.48. 4-(isobutoxycarbonyl)carbanilide
- A.49. 2,4'-bis(methoxycarbonyl)carbanilide
- A.50. 2',4-dichloro-3-nitro-3'-(trifluoromethyl)-carbanilide
- 20 A.51. 3,4,4',5-tetrachloro-3'-(trifluoromethyl)-carbanilide (C.A. 63: P440f)
- A.52. 4-chloro-3-nitro-3',5'-bis(trifluoromethyl)-carbanilide
- 25 A.53. 2,4,6-trimethyl-4'-(ethoxycarbonyl)carbanilide
- A.54. 2-(trifluoromethyl)-2'-ethyl-6'-isopropyl-carbanilide
- 30 A.55. 4-chloro-3,3',5'-tris(trifluoromethyl)carbanilide (CAS registry #4528-83-0; C.A. 71: 91052Y and 66: 64644h)

- A.56. 3,4,4',5-tetrachloro-3'-nitrocarbanilide
- A.57. 2,6-dimethyl-4'-benzoylcarbanilide
- A.58. 3,4-dimethyl-2'-ethoxycarbanilide
- 5 A.59. 2-chloro-4,4'-bis(methylthio)carbanilide
- A.60. 2-methyl-2'-ethoxycarbanilide
- A.61. 4-chloro-2-methoxythiocarbanilide
- A.62. 4,4'-dinitro-N,N'-dimethylcarbanilide (CAS
10 registry #34594-37-3; C.A. 83: 36180m and
C.A. 85: 20476t)
- A.63. 4-(trifluoromethyl)-4'-nitrocarbanilide (CAS
registry #23747-76-4 U.S. 3,867,544)
- 15 A.64. 3,3',5,5'-tetrakis(trifluoromethyl)-N,N'-
dimethylcarbanilide
- A.65. 4-phenoxy-4'-nitrocarbanilide

20 A second group of carbanilides of particular
value in the present invention are those carbanilides
described in U.S. Patent No. 3,284,433. These com-
pounds may be represented by the formula



wherein each Y is independently selected from the group
consisting of chlorine, bromine and nitro radicals; and
(2) compounds of the same formula as (II) except that a
30 hydrogen of at least one of the phenyl rings of (II)

is replaced by a substituent selected from the group consisting of chlorine, bromine, nitro and C₁₋₆ alkyl radicals.

5 Representative compounds of this group include the following:

Group B

- | | | |
|----|-------|---|
| | B.1. | 4-nitro-4'-(4-chlorophenoxy) carbanilide |
| | B.2. | 4-nitro-4'-(3,4-dichlorophenoxy) carbanilide |
| 10 | B.3. | 4-nitro-3'-chloro-4'-(4-chlorophenoxy) carbanilide |
| | B.4. | 4-nitro-3',5'-dichloro-4'-(4-chlorophenoxy) - carbanilide |
| 15 | B.5. | 4-nitro-3'-chloro-4'-(3,4-dichlorophenoxy) - carbanilide |
| | B.6. | 4-nitro-4'-(2,4-dichlorophenoxy) carbanilide |
| | B.7. | 4-nitro-4'-(4-nitrophenoxy) carbanilide |
| 20 | B.8. | 4-nitro-3'-nitro-4'-(4-chlorophenoxy) - carbanilide |
| | B.9. | 4-nitro-3'-methyl-4'-(4-chlorophenoxy) - carbanilide |
| 25 | B.10. | 4-nitro-2'-nitro-4'-(4-chlorophenoxy) - carbanilide |
| | B.11. | 4-nitro-4'-(2,4-dinitrophenoxy) carbanilide |
| | B.12. | 4-nitro-3'-chloro-4'-(2-tert-butyl-4-chloro- phenoxy) carbanilide |
| 30 | B.13. | 4-nitro-4'-(2-methyl-4-chlorophenoxy) - carbanilide |

- B.14. 4-nitro-3'-bromo-4'-(4-bromophenoxy) carbanilide
- B.15. 4-nitro-3'-bromo-4'-(4-chlorophenoxy) -
carbanilide
- 5 B.16. 4-nitro-3'-chloro-4'-(4-bromophenoxy) -
carbanilide
- B.17. 3,3',4'-trichloro-4-(4-chlorophenoxy) -
carbanilide
- 10 B.18. 3,4'-dichloro-4-(4-chlorophenoxy) carbanilide
- B.19. 3,4-dichloro-4'-(4-chlorophenoxy) carbanilide
- B.20. 2-methyl-4-nitro-3'-chloro-4'-(4-chloro-
phenoxy) carbanilide

15 A third group of carbanilides of value in the
present invention are those carbanilides described in
German Patent Specification No. 2,334,355. These
compounds include the following:

Group C

- 20 C.1. 3,3',5'-tris(trifluoromethyl)-4-methoxy-
carbanilide
- C.2. 3-(trifluoromethoxy)-3',5'-bis(trifluoro-
methyl) carbanilide
- 25 C.3. 4-(trifluoromethoxy)-3',5'-bis(trifluoro-
methyl) carbanilide
- C.4. 2,3',5'-tris(trifluoromethyl)-4-(trifluoro-
methoxy) carbanilide
- 30 C.5. 4-(methylthio)-3',5'-bis(trifluoromethyl)-
carbanilide

- C.6. 4-(methylthio)-3',5'-bis(trifluoromethyl)-
thiocarbanilide
- 5 C.7. 3-chloro-4-(methylthio)-3',5'-bis(trifluoro-
methyl) carbanilide
- C.8. 3-(trifluoromethylthio)-3',5'-bis(trifluoro-
methyl) carbanilide
- C.9. 4-(trifluoromethylthio)-3',5'-bis(trifluoro-
methyl) carbanilide
- 10 C.10. 2-chloro-4-(trifluoromethylthio)-3',5'-
bis(trifluoromethyl) carbanilide
- C.11. 4-(trifluoromethylthio)-3'-(trifluoromethyl)-
5'-nitrocarbanilide
- 15 C.12. 4-(trifluoromethylthio)-2',5'-bis(trifluoro-
methyl)-4'-nitrocarbanilide
- C.13. 4,4'-bis(trifluoromethylthio) carbanilide
- C.14. 4-(trifluoromethylthio)-4'-(chloromethyl-
sulfonyl) carbanilide
- 20 C.15. 3-(difluoromethylthio)-3',5'-bis(trifluoro-
methyl) carbanilide
- C.16. 4-(ethylthio)-3',5-bis(trifluoromethyl)-
carbanilide
- 25 C.17. 3-chloro-4-(ethylthio)-3',5'-bis(trifluoro-
methyl) carbanilide
- C.18. 4-ethoxy-3,3',5'-tris(trifluoromethyl) car-
banilide
- 30 C.19. 3-(trifluoromethyl)-4-ethoxy-4'-(trifluoro-
methylthio) carbanilide

- C.20. 3-(trifluoromethyl)-4-ethoxy-3'-(trifluoromethylthio)carbanilide
- 5 C.21. 4-methyl-3-(2-chloroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- C.22. 4-(1,2-dichlorovinyl)-3'-(trifluoromethyl)-4'-chlorocarbanilide
- C.23. 4-(1,2-dichlorovinyl)-3',5'-bis(trifluoromethyl)carbanilide
- 10 C.24. 4-(2,2,2-trichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- C.25. 2-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- 15 C.26. 3-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- C.27. 3-(2,2-dichloro-1,1-difluoroethoxy)-4-bromo-3',5'-bis(trifluoromethyl)carbanilide
- 20 C.28. 4-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- C.29. 4-(2,2-dichloro-1,1-difluoroethoxy)-3'-(trifluoromethyl)-4'-chlorocarbanilide
- 25 C.30. 3-methoxy-4-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- C.31. 3-methyl-4-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl)carbanilide
- 30 C.32. 3-methyl-4-(2,2-dichloro-1,1-difluoroethoxy)-4'-isopropylcarbanilide

- C.33. 3-nitro-4-(2,2-dichloro-1,1-difluoroethoxy)-
3',5'-bis(trifluoromethyl) carbanilide
- 5 C.34. 2-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl) carbanilide
- C.35. 4-(2,2-dichloro-1,1-difluoroethoxy)-3,3',5'-
tris(trifluoromethyl) carbanilide
- C.36. 3-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl) carbanilide
- 10 C.37. 4-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl) carbanilide
- C.38. 4-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl) thiocarbanilide
- 15 C.39. 4-(2-chloro-1,1,2-trifluoroethoxy)-3'-
(trifluoromethyl)-5'-nitrocarbanilide
- C.40. 4,4'-bis(2-chloro-1,1,2-trifluoroethoxy)-
carbanilide
- 20 C.41. 4-(2-chloro-1,1,2-trifluoroethoxy)-4'-
(trifluoromethylthio) carbanilide
- C.42. 4-(2-chloro-1,1,2-trifluoroethoxy)-3'-
(trifluoromethyl)-4'-chlorocarbanilide
- C.43. 4-(2-chloro-1,1,2-trifluoroethoxy)-3,3'-
25 bis(trifluoromethyl)-5'-nitrocarbanilide
- C.44. 4-(2-chloro-1,1,2-trifluoroethoxy)-3,3',5'-
tris(trifluoromethyl) carbanilide
- C.45. 3-(1,1,2-trifluoroethoxy)-3',5'-bis(tri-
30 fluoromethyl) carbanilide
- C.46. 3-(1,1,2,2-tetrafluoroethoxy)-3',5'-bis(tri-
fluoromethyl) carbanilide

- C.47. 3-methyl-4-(1,1,2,2-tetrafluoroethoxy)-
3'-(chlorodifluoromethyl)carbanilide
- C.48. 3-methyl-4-(1,1,2,2-tetrafluoroethoxy)-3'-
(trifluoromethyl)-4'-chlorocarbanilide
- 5 C.49. 3-(trifluoromethyl)-4-(1,1,2,2-tetrafluoro-
ethoxy)-4'-bromocarbanilide
- C.50. 3-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3',5'-bis(trifluoromethyl)carbanilide
- 10 C.51. 4-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3',5'-bis(trifluoromethyl)carbanilide
- C.52. 3-chloro-4-(1,1,2,3,3,3-hexafluoro-n-
propoxy)-3',5'-bis(trifluoromethyl)-
carbanilide
- 15 C.53. 3-methyl-4-(1,1,2,3,3,3-hexafluoro-n-
propoxy)-3',5'-bis(trifluoromethyl)-
carbanilide
- C.54. 4-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3,3',5'-tris(trifluoromethyl)carbanilide
- 20 C.55. 4-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3,3'-bis(trifluoromethyl)-5'-nitro-
carbanilide
- 25 C.56. 4-(methoxycarbonylthio)-3',5'-bis(tri-
fluoromethyl)carbanilide
- C.57. 4-(4-chlorophenoxy)-4'-(trifluoromethyl-
thio)thiocarbanilide
- 30 C.58. 4-(4-chlorophenoxy)-3',5'-bis(tri-
fluoromethyl)carbanilide


- C.59. 3-chloro-4-(4-chlorophenoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 5 C.60. 4-(4-chlorophenoxy)-3'-(trifluoromethyl)-5'-nitrocarbanilide
- C.61. 4-(3,5-bis(trifluoromethyl)) phenoxy-3',5'-bis(trifluoromethyl) carbanilide
- C.62. 4-(4-methylphenoxy carbonyl)-3',5'-bis(trifluoromethyl) carbanilide
- 10 C.63. 4-(4-methylphenoxy carbonyl)-4'-(trifluoromethylthio) carbanilide
- C.64. 3,5-bis(trifluoromethyl)-2',4',6'-trichlorocarbanilide
- 15 C.65. 3,5-bis(trifluoromethyl)-2',4',5'-trichlorocarbanilide
- C.66. 3,3'-bis(trifluoromethyl)-5'-nitrocarbanilide
- 20 C.67. 3,3',5-tris(trifluoromethyl)-5'-nitrocarbanilide
- C.68. 2',3,5,6'-tetrakis(trifluoromethyl)-4'-nitrocarbanilide
- 25 C.69. 3-(chlorodifluoromethyl)-3',5'-bis(trifluoromethyl) carbanilide
- C.70. 3-(1,1,2,2-tetrafluoroethyl)-3',5'-bis(trifluoromethyl) carbanilide
- 30 C.71. 4-phenyl-3',5'-bis(trifluoromethyl)-carbanilide

- C.72. 3-(fluorosulfonyl)-3',5'-bis(trifluoromethyl)carbanilide
- C.73. 4-(fluorosulfonyl)-4'-(trifluoromethylthio)carbanilide
- 5 C.74. 4-(fluorosulfonyl)-3'-(trifluoromethyl)-5'-nitrocarbanilide
- C.75. 4-(chloromethylsulfonyl)-3',5'-bis(trifluoromethyl)carbanilide
- 10 C.76. 2-(ethylsulfonyl)-3',5,5'-tris(trifluoromethyl)carbanilide
- C.77. 2-(ethylsulfonyl)-3',5-bis(trifluoromethyl)-5'-nitrocarbanilide
- 15 C.78. 4-(trifluoromethylsulfonyl)-3',5'-bis(trifluoromethyl)carbanilide
- C.79. 4-(trifluoromethylsulfonyl)-3'-(trifluoromethyl)-4'-methoxycarbanilide
- 20 C.80. 4-(trifluoromethylsulfonyl)-4'-(trifluoromethylthio)carbanilide
- C.81. 4-(4-nitrophenylsulfonyl)-4'-(trifluoromethylthio)carbanilide
- 25 C.82. 4-(4-acetamidophenylsulfonyl)-4'-(trifluoromethylthio)carbanilide
- C.83. 4-(4-(isopropylideneamino)phenylsulfonyl)-4'-(trifluoromethylthio)carbanilide
- 30 C.84. 4,4-sulfonylbis(3'-(trifluoromethylthio)-carbanilide)

- C.85. 4,4-sulfonylbis(4'-(1,2-dichlorovinyl-oxy)carbanilide)
- C.86. 4,4-sulfonylbis(3'-(1,1,2,2-tetrafluoroethoxy)carbanilide)
- C.87. 4,4-sulfonylbis(4'-(2-chloro-1,1,2-trifluoroethoxy)carbanilide)
- C.88. 4,4-sulfonylbis(4'-(trifluoromethylthio)carbanilide)

The presently preferred carbanilide derivative for use in the anticoccidial formulations of the invention is Nicarbazin which as previously stated is a complex of 4,4'-dinitrocarbanilide and 2-hydroxy-4,6-dimethylpyrimidine. However, it should be noted that it is the carbanilide portion which exhibits the anticoccidial activity (see Science 122, 244 (1955)).

Many carbanilides are known; however, there exists a class of novel carbanilide derivatives which the Applicants have found to be of particular value in the invention. These carbanilides have the structural formula:



where R^{10} , R^{11} and R^{12} are the same or different and can each represent hydrogen, C_{1-4} alkyl, nitro, halo, C_{1-4} haloalkyl, cyano, or C_{1-6} alkoxy carbonyl, provided that at least one of R^{10} , R^{11} and R^{12} is other than hydrogen.

Interestingly, although normally free amino substitution is not advantageous in connection with the carbanilides of the invention, despite the presence of the free amino groups, these novel compounds are particularly effective in accordance with the invention.

Examples of this type of compound are:

2-amino-3-nitro-5-(trifluoromethyl)-2',4'-dimethylcarbanilide

2-amino-3,3'-dinitro-5-(trifluoromethyl)-4'-chlorocarbanilide

2-amino-3-nitro-5-(trifluoromethyl)-2',4',5'-trichlorocarbanilide

2-amino-3-nitro-5-(trifluoromethyl)-2'-(ethoxycarbonyl)carbanilide

2-amino-3,5'-dinitro-5-(trifluoromethyl)-2'-fluorocarbanilide

2-amino-3,4'-dinitro-5-(trifluoromethyl)-2'-chlorocarbanilide

2-amino-3-nitro-3',5-bis(trifluoromethyl)-4'-chlorocarbanilide


2-amino-3-nitro-3',5,5'-tris(trifluoromethyl)-carbanilide

2-amino-3-nitro-2',5-bis(trifluoromethyl)-4'-chlorocarbanilide

2-amino-3-nitro-5-trifluoromethyl-2'-methyl-4'-bromocarbanilide and

2-amino-3-nitro-5-(trifluoromethyl)-2',6'-dibromo-4'-fluorocarbanilide.

This novel class of ureas, as well as the other carbanilides of the invention, can be prepared by conventional synthetic procedures such as those specified in German Patent Specification No. 2,334,355. For instance, the compounds may be prepared by reaction of the appropriate *o*-phenylenediamine with the corresponding phenyl isocyanate as depicted below:



This reaction is preferably carried out at a temperature within the range of 10° to 130°C. and in the presence of a tertiary organic base such as pyridine or triethylamine. Any suitable inert organic solvent such as benzene, toluene, chlorobenzene, dioxane or methylene chloride may be utilized.

The following non-limiting Examples illustrate the synthesis of this novel class of carbanilides.

EXAMPLE 1

2-AMINO-3-NITRO-5-(TRIFLUOROMETHYL)-2',4'-DIMETHYLCARBANILIDE

5 3-Nitro-5-(trifluoromethyl)-o-phenylene-
diamine (2.2 grams; 0.01 mole) and 2,4-dimethylphenyl
isocyanate (1.5 grams; 0.01 mole) were taken up in
40 ml. of methylene chloride, and 1 ml. of pyridine
was added. The reaction mixture was maintained at
10 23°C. for 16 hours. The yellow precipitate was fil-
tered and dried, m.p. >300°C. Elemental analysis showed
the following:
Calculated for $C_{16}H_{15}F_3N_4O_3$: C, 52.17; H, 4.08; N, 15.22.
Found: C, 53.01; H, 4.05; N, 16.44.

15 Additional carbanilides were prepared by
substantially the same procedures as that of Example 1.

EXAMPLE 2

20 2-Amino-3,3'-dinitro-5-(trifluoromethyl)-4'-
chlorocarbanilide was prepared by reacting 3-nitro-5-
(trifluoromethyl)-o-phenylenediamine (2.2 grams; 0.01 mole)
and 3-nitro-4-chlorophenyl isocyanate (2.0 grams;
0.01 mole). The product, 3.3 grams, melted at 214-216°C.
Elemental analysis showed the following:
25 Calculated for $C_{14}H_9ClF_3N_5O_5$: C, 40.06; H, 2.16; N, 16.96.
Found: C, 40.33; H, 1.88; N, 16.45.

EXAMPLE 3

30 2-Amino-3-nitro-5-(trifluoromethyl)-2',4',5'-
trichlorocarbanilide was prepared by reacting 3-nitro-5-
(trifluoromethyl)-o-phenylenediamine (2.2 grams; 0.1 mole)

and 2,4,5-trichlorophenyl isocyanate (2.2 grams; 0.01 mole). The product melted at 225-227°C. Elemental analysis showed the following:

Calculated for $C_{14}H_8Cl_3F_3N_4O_3$: C, 37.91; H, 1.82; N, 12.63.

Found: C, 37.83; H, 1.55; N, 12.40.

EXAMPLE 4

2-Amino-3-nitro-5-(trifluoromethyl)-2'-(ethoxycarbonyl)carbanilide was prepared by reacting 3-nitro-5-(trifluoromethyl)-o-phenylenediamine (1.1 grams; 0.005 mole) and 2-(ethoxycarbonyl)phenyl isocyanate (1.0 grams; 0.005 mole). The product, 1.4 grams, melted at 186-188°C. Elemental analysis showed the following:

Calculated for $C_{17}H_{15}F_3N_4O_5$: C, 49.52; H, 3.67; N, 13.59.

Found: C, 49.75; H, 3.59; N, 13.50.

EXAMPLE 5

2-Amino-3,5'-dinitro-5-(trifluoromethyl)-2'-fluorocarbanilide was prepared from 3-nitro-5-(trifluoromethyl)-o-phenylenediamine (1.1 grams; 0.005 mole) and 2-fluoro-5-nitrophenyl isocyanate (0.8 gram; 0.005 mole). The product, 1.2 grams, melted at 218-220°C. Elemental analysis showed the following:

Calculated for $C_{14}H_9F_4N_5O_5$: C, 41.70; H, 2.25; N, 17.37.

Found: C, 41.91; H, 1.98; N, 17.27.

EXAMPLE 6

2-Amino-3,4'-dinitro-5-(trifluoromethyl)-2'-chlorocarbanilide was prepared by reacting 3-nitro-5-(trifluoromethyl)-o-phenylenediamine (1.1 grams; 0.005 mole) and 2-chloro-4-nitrophenyl isocyanate (1.0 gram; 0.005 mole). The product, 1.2 grams, melted at 220-222°C. Elemental analysis showed the following:

Calculated for $C_{14}H_9ClF_3N_5O_5$: C, 40.06; H, 2.16; N, 16.67.
Found: C, 40.11; H, 2.11; N, 16.48.

EXAMPLE 7

5 2-Amino-3-nitro-3',5-bis(trifluoromethyl)-4'-
chlorocarbanilide was prepared by reacting 2-amino-3-
nitro-5-(trifluoromethyl)-o-phenylenediamine (2.2 grams;
0.01 mole) and 4-chloro-3-(trifluoromethyl)phenyl iso-
cyanate (2.2 grams; 0.01 mole). The product, 1.8 grams,
10 melted at 214-216°C. Elemental analysis showed the
following:

Calculated for $C_{15}H_9ClF_6N_4O_3$: C, 40.70; H, 2.05; N, 12.66.
Found: C, 40.90; H, 2.04; N, 12.67.

EXAMPLE 8

15 2-Amino-3-nitro-3',5,5'-tris(trifluoromethyl)-
carbanilide was prepared by reacting 3-nitro-5-(tri-
fluoromethyl)-o-phenylenediamine (1.1 grams; 0.005 mole)
and (3,5-bis(trifluoromethyl)phenyl)carbamoyl chloride
(1.4 grams; 0.005 mole). Elemental analysis of the
20 product showed the following:

Calculated for $C_{16}H_9F_9N_4O_3$: C, 40.35; H, 1.92; N, 11.76.
Found: C, 40.52; H, 1.82; N, 11.78.

EXAMPLE 9

25 2-Amino-3-nitro-2',5-bis(trifluoromethyl)-4'-
chlorocarbanilide was prepared by reacting 3-nitro-5-
(trifluoromethyl)-o-phenylenediamine (2.2 grams; 0.01
mole) and 4-chloro-2-(trifluoromethyl)phenyl isocyanate
(2.2 grams; 0.01 mole). The product melted at 245-247°C.
30 Elemental analysis showed the following:

Calculated for $C_{15}H_9ClF_6N_4O_3$: C, 40.70; H, 2.05; N, 12.66.
Found: C, 40.94; H, 1.88; N, 12.55.

EXAMPLE 10

2-Amino-3-nitro-5-(trifluoromethyl)-2'-methyl-4'-bromocarbanilide was prepared by reacting 3-nitro-5-(trifluoromethyl)-o-phenylenediamine (2.2 grams; 0.01 mole) and 2-methyl-4-bromophenyl isocyanate (2.1 grams; 0.01 mole). The product melted at 230-231°C. Elemental analysis showed the following:
Calculated for $C_{15}H_{11}BrF_3N_4O_3$: C, 41.59; H, 2.79; N, 12.93.
Found: C, 41.40; H, 2.65; N, 12.71.

EXAMPLE 11

2-Amino-3-nitro-5-(trifluoromethyl)-2',6'-dibromo-4'-fluorocarbanilide was prepared by reacting 3-nitro-5-(trifluoromethyl)-o-phenylenediamine (2.2 grams; 0.01 mole) and 2,6-dibromo-4-fluorophenyl isocyanate. The product melted at 249-251°C. Elemental analysis showed the following:
Calculated for $C_{14}H_8Br_2F_4N_4O_3$: C, 32.59; H, 1.56; N, 10.86.
Found: C, 32.64; H, 1.39; N, 10.76.

The preferred ratio of carbanilide to polyether in the anticoccidial compositions of the invention is from 40:1 to 1:20, more preferably from 10:1 to 1:10, for instance 2:1 to 1:2, by weight. Although the compositions of the invention are effective in controlling coccidiosis in all avian species, obviously from the economic viewpoint, the present invention only has commercial significance in coccidiosis treatment in poultry such as chickens or turkeys.

Since coccidiosis affects the intestinal tract, the compositions of the present invention are in a form suitable for oral administration. The polyether antibiotics are generally of low solubility in water, even in the sodium or other salt form. Therefore, the present invention is practiced by administering the polyether/carbanilide combination to the poultry in a feedstuff rather than in drinking water. Furthermore, it is the practice of the industry to supply poultry with only one source of feed, constituting the entire food supply of the poultry. Therefore, in a preferred practice of the present invention, the anticoccidial combinations are supplied in a total feed, with concentrations adjusted accordingly. Those skilled in the art, however, will recognize that concentrations are to be adjusted upward, should it be desired to supply poultry with multiple sources of food only one of which contains the combination of the present invention.

The active ingredients of the compositions of the invention can be employed in a wide range of concentrations in the feed administered to poultry. In general, for those components which are known anticoccidials, the maxima to be employed in accordance with the present invention are the same as the maxima for anticoccidial treatment by the individual components. The lower limits in accordance with the present invention are generally less than for control by the individual components, especially where the components are being used to minimize side effects of either individual component. Generally, compositions in accordance with the invention will not contain less than 15 ppm nor more than 25% by weight of active ingredients.

Representative amounts of selected polyether antibiotics in the final feed destined for consumption are as follows:

- 5 from about 20 to about 120 ppm of
 monensin;
- from about 25 to about 100 ppm of
- narasin;
- from about 35 to about 125 ppm of
- lasalocid;
- 10 from about 25 to about 100 ppm of
- salinomycin;
- from about 5 to about 15 ppm of A-204;
- from about 25 to about 100 ppm of
- lonomycin;
- 15 from about 25 to about 100 ppm of X-206;
- from about 50 to about 200 ppm of
- nigericin; and
- from about 10 to about 50 ppm of
- dianemycin.

20 The carbanilide is generally employed in a concentration
 of from about 10 to about 250 ppm, preferably from
 about 25 to about 100 ppm in the poultry feed. Amounts
 will be adjusted downward where more than one polyether,
 or more than one carbanilide, is employed. Thus,
25 compositions of the invention in the form of feedstuffs
 for direct consumption by the avian species to be treated
 will normally contain from 15 to 450 ppm of the active
 ingredients.

30 In a preferred embodiment of the present
 invention, compositions comprise a polyether antibiotic
 and nicarbazin or 4,4'-dinitrocarbanilide as the sole
 anticoccidial agents.

Compositions containing a mixture of nicarbazin or 4,4'-dinitrocarbanilide and monensin, especially with the monensin in the commercially available form consisting of factor A and a minor amount of factor B, constitute a particularly preferred embodiment of the invention. A preferred feedstuff for direct feeding of poultry contains from about 25 to about 75 ppm of monensin and from about 25 to about 80 ppm of nicarbazin or from about 25 to about 100 ppm of 4,4'-dinitrocarbanilide.

Since the polyether antibiotics alone are active as anticoccidials, the compositions of the invention are useful regardless of the exact concentration of the carbanilide. As noted above, certain of the carbanilides, alone, exhibit anticoccidial activity. Therefore, preferred embodiments of the present invention are those wherein the carbanilide is employed in a concentration that potentiates the anticoccidial activity of the polyether (where the carbanilide is, itself, lacking anticoccidial activity); or in a concentration that is synergistic with the anticoccidial activity of the polyether (where the carbanilide also exhibits anticoccidial activity).

The Group A compounds include both compounds exhibiting no independent anticoccidial activity, at typical rates, as well as compounds exhibiting independent anticoccidial activity. Each of the Group A compounds has been found to potentiate or synergize, respectively, the activity of a representative polyether antibiotic, monensin. The potentiating and synergizing effect of the Group A compounds is shown in Tables I and IV, below.

The Group B compounds are taught in U.S. Patent No. 3,284,433 to exhibit anticoccidial activity. Therefore, they are preferably employed in amounts which are synergistic as to the anticoccidial activity of the polyethers.

The Group C compounds are taught in German Patent Specification No. 2,334,355 to exhibit anticoccidial activity. They are likewise preferably employed in amounts which are synergistic as to the anticoccidial activity of the polyethers. Data on the effects of the Group C compounds is shown in Tables I, II, and III, below. Simple range-finding experiments as to any of the carbanilides to be employed in the present invention will enable those skilled in the art to determine the preferred potentiating or synergistic amounts of the respective carbanilide.

Poultry feedstuffs of all types and formulae in the poultry industry may be used in administering the combinations of the present invention. The following formulae are exemplary only.

Broiler Starter

	<u>Ingredients</u>	<u>Percent</u>
	Corn, Yellow, Ground	50.0
25	Soybean Oil Meal, Solvent Extracted Dehulled (50%)	30.9
	Animal Fat	6.5
	Fish Meal with Solubles (60%)	5.0
	Corn Distillers Dried Solubles	4.0
	Dicalcium Phosphate, Feed Grade	1.8
30	Calcium Carbonate (Ground Limestone)	0.8
	Vitamin Premix TK-01 (1.03) <u>1/</u>	0.5
	Salt (NaCl)	0.3
	Trace Mineral Premix TK-01 (1.02) <u>2/</u>	0.1
	Methionine Hydroxy Analog	0.1
	Total	<u>100.0</u>

Broiler Grower

	<u>Ingredients</u>	<u>Percent</u>
	Corn, Yellow, Ground	57.7
5	Soybean Meal, Solvent, Extracted, Dehulled (50%)	31.7
	Animal Fat (Beef tallow)	6.0
	Dicalcium Phosphate, Feed Grade	2.7
	Calcium Carbonate (Ground Limestone)	0.9
	Vitamin Premix TK-01 (1.03) ¹ /	0.5
	Salt (NaCl)	0.2
	Methionine Hydroxy Analog	0.2
10	Trace Mineral Premix TK-01 (1.02) ² /	0.1
	Total	100.0

Chick Starter, Light Breeds

	<u>Ingredients</u>	<u>Percent</u>
	Corn, Yellow, Ground	56.3
15	Soybean Meal, Solvent Extracted, Dehulled (50%)	17.9
	Wheat Middlings	10.0
	Corn Distillers Dried Solubles	5.0
	Fish Meal with Solubles	5.0
	Alfalfa Meal, Dehydrated (17%)	2.5
	Dicalcium Phosphate, Feed Grade	1.3
20	Calcium Carbonate	0.9
	Vitamin Premix ¹	0.5
	Salt (NaCl)	0.3
	Methionine Hydroxy Analog	0.2
	Trace Mineral Premix ²	0.1
	Total	100.0

25

30

Pullet Grower

	<u>Ingredients</u>	<u>Percent</u>
	Corn, Yellow, Ground	73.5
5	Soybean Meal, Solvent Extracted, Dehulled (50%)	21.9
	Dicalcium Phosphate, Feed Grade	2.5
	Calcium Carbonate	1.0
	Vitamin Premix ¹	0.5
	Salt (NaCl)	0.3
	Methionine Hydroxy Analog	0.2
	Trace Mineral Premix ²	0.1
10	Total	100.0

Pullet Developer

	<u>Ingredients</u>	<u>Percent</u>
	Corn, Yellow, Ground	67.5
15	Oats, Ground Whole	15.0
	Soybean Meal, Solvent Extracted, Dehulled (50%)	13.4
	Dicalcium Phosphate, Feed Grade	2.1
	Calcium Carbonate	1.0
	Vitamin Premix ¹	0.5
	Methionine Hydroxy Analog	0.3
	Salt (NaCl)	0.2
20	Trace Mineral Premix ²	0.1
	Total	100.0

25

30

Turkey Starter

	<u>Ingredients</u>	<u>Percent</u>
	Soybean Meal, Solvent Extracted, Dehulled	40.7
5	Corn, Yellow, Ground	39.7
	Fish Meal with Solubles	5.0
	Beef Tallow	5.0
	Corn Distillers Dried Solubles	2.5
	Alfalfa Meal, Dehydrated (17%)	2.5
	Dicalcium Phosphate, Feed Grade	2.5
	Calcium Carbonate	1.2
10	Vitamin Premix ¹	0.5
	Salt (NaCl)	0.2
	Trace Mineral Premix ²	0.1
	Methionine Hydroxy Analog	0.1
	Total	100.0

Turkey Finisher

	<u>Ingredients</u>	<u>Percent</u>
	Corn, Yellow, Ground	71.2
	Soybean Meal, Solvent Extracted, Dehulled (50%)	9.9
	Corn Distillers Dried Solubles	5.0
	Alfalfa Meal, Dehydrated (17%)	5.0
20	Animal Fat	3.0
	Fish Meal With Solubles	2.5
	Dicalcium Phosphate, Feed Grade	1.7
	Calcium Carbonate	0.5
	Vitamin Premix ¹	0.5
	Salt (NaCl)	0.4
	Methionine Hydroxy Analog	0.2
	Trace Mineral Premix ²	0.1
25	Total	100.0

1/ Vitamin premix provides 3000 IU of vitamin A, 900 ICU of vitamin D, 40 mg. of vitamin E, 0.7 mg. of vitamin K, 1000 mg. of choline, 70 mg. of niacin, 4 mg. of pantothenic acid, 4 mg. of riboflavin, 0.10 mg. of vitamin B₁₂, 0.10 mg. of biotin and 125 mg. of ethoxy-quin per kg. of complete feed.

2/ Trace mineral premix provides 75 mg. of manganese, 50 mg. of zinc, 25 mg. of iron and 1 mg. of iodine per kg. of complete feed.

The foregoing compositions are typical of feedstuffs actually administered to poultry. Premixes are commonly used in the poultry industry, and the compositions of the present invention can also take this form. Such premixes typically comprise the polyether antibiotic, the selected carbanilide, and a solid substance capable of being ingested by the poultry, such as corn meal, rice hulls, crushed limestone, soybean meal, soya grits, distillers' dried grains, citrus meal, wheat middlings and clays. Such a premix is mixed or blended with other substances to constitute either an intermediate premix or the finished feed.

A representative premix in accordance with the present invention is as follows:

monensin: 45 grams activity
a selected carbanilide in accordance with the
present invention: 45 grams
rice hulls q.s. ad 454 grams

This formulation is mixed thoroughly and can be added to one ton of finished feed to provide a composition in accordance with the invention in the form of a feed suitable for consumption by poultry. Premixes in accordance with the invention will normally contain from 1 to 25% by weight of active ingredients.

The present invention was evaluated in chickens as follows: one-week-old broiler chicks were fed a medicated or control ration, typically for one day, prior to infection (by gavage) with oocysts of a coccidiosis-causing organism, generally 750,000 oocysts

of Eimeria acervulina and 75,000 oocysts of Eimeria tenella. The chicks were maintained on their respective rations for an additional period of time, typically six days. There were five chicks in a group and generally
5 two replicates per treatment. Anticoccidial efficacy was determined by lesion scores. To determine lesion scores, the birds were sacrificed and the severity of lesions scored on a 0-4 scale, with lesion-free birds scored as 0, extremely severe infections scored as 4,
10 and intermediate degrees of infection scored as 1, 2 or 3. The scores of all birds which received a given treatment were averaged.

Most of the compounds were tested in a preliminary test "A"; in this test, monensin was tested
15 alone at 50 ppm and the carbanilide compound was tested in combination with monensin, each at 50 ppm. In a secondary test "B", monensin and the carbanilide compound were each tested alone, at 50 ppm, and the two were tested in combination, each at 50 ppm. Additional
20 tests at more varied concentrations were also conducted with certain of the compounds.

The results are reported in the following tables. The concentration of monensin and the carbanilides is expressed as the parts per million (ppm)
25 of the total feed provided to the chicks.

TABLE I

	<u>Compound Number</u>	<u>LESION SCORES</u>					
		<u>Eimeria acervulina</u>			<u>Eimeria tenella</u>		
		<u>monensin</u>			<u>monensin</u>		
		<u>0</u>	<u>50</u>		<u>0</u>	<u>50</u>	
5	A.3.	A 0	2.8	1.6	2.0	1.9	
		50		0.7		0.9	
		B 0	3.0	1.2	2.0	1.2	
		50	3.0	1.2	2.5	0.7	
10	A.4.	A 0	2.8	1.6	2.0	0.9	
		50		0		0	
		B 0	3.0	1.2	2.0	1.2	
		50	3.0	0.9	2.4	0.6	
15	A.5.	A 0	2.4	1.2	2.5	1.4	
		50		0.3		0.4	
		B 0	3.0	1.2	2.0	1.2	
		50	3.0	1.1	2.2	0	
	A.6.	A 0	2.8	1.6	2.0	0.9	
		50		1.6		0.6	
20		B 0	3.0	1.2	2.0	1.2	
		50	2.9	0.8	2.5	0.5	
	A.7.	A 0	2.5	1.6	2.4	2.3	
		50		0.9		0.4	
		B 0	3.0	1.2	2.0	1.2	
25		50	3.0	1.4	1.8	0.6	
	A.8.	A 0	3.0	1.2	2.1	0.9	
		50		0.4		0.2	
		B 0	3.3	1.3	3.0	2.2	
		50	2.6	0.8	3.0	1.0	
30							

TABLE I (cont'd)

	Compound Number		LESION SCORES			
			<u>Eimeria acervulina</u> monensin		<u>Eimeria tenella</u> monensin	
			0	50	0	50
5	A.9.	A 0	3.0	1.2	2.1	0.9
		50		0.7		0.9
		B 0	3.3	1.3	3.0	2.2
		50	2.8	0.8	2.9	1.3
10	A.10.	A 0	2.7	1.7	3.0	1.0
		50		0.5		0.6
		B 0	3.3	1.3	3.0	2.2
		50	2.4	0.9	2.7	1.7
15	A.11.	A 0	3.6	1.2	3.0	1.2
		50		0.6		0.8
		B 0	3.3	1.3	3.0	2.2
		50	2.6	1.4	2.8	1.1
20	A.12.	A 0	2.8	1.1	2.3	1.0
		50		1.1		0.6
		B 0	3.3	1.3	3.0	2.2
		50	2.5	0.9	2.9	0.8
25	A.13.	A 0	3.0	1.2	2.1	0.9
		50		0.9		0.5
		B 0	3.3	1.3	3.0	2.2
		50	2.9	1.4	3.0	1.2
30	A.14.	A 0	3.0	1.2	2.1	0.9
		50		1.2		0.6
		B 0	3.3	1.3	3.0	2.2
		50	2.6	1.1	3.0	1.0
	C.8.	A 0	2.7	1.7	3.0	1.0
		50		1.2		0.4
		B 0	3.3	1.3	3.0	2.2
		50	2.3	1.1	3.0	0.3

TABLE I (cont'd)

	Compound Number		LESION SCORES			
			<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>	
			<u>monensin</u>		<u>monensin</u>	
			<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>
5	A.15.	A 0	3.2	1.5	2.4	1.9
		50		1.8		1.0
		B 0	3.3	1.3	3.0	2.2
		50	2.6	1.6	2.6	1.1
10	A.16.	A 0	3.0	1.1	2.6	1.0
		50		0.9		0.5
		B 0	3.3	1.3	3.0	2.2
		50	2.8	1.0	2.8	1.1
	A.17.	A 0	2.8	1.4	2.6	0.8
15		50		0.9		0.9
		B 0	2.6	1.0	2.5	1.5
		50	2.4	0.6	2.6	1.3
	A.18.	A 0	2.8	1.4	2.6	0.8
		50		0.1		0.3
20		B 0	2.6	1.0	2.5	1.5
		50	2.7	0.6	2.0	0.7
	A.19.	A 0	2.8	1.4	2.6	0.8
		50		0.4		0.4
		B 0	2.6	1.0	2.5	1.5
25		50	2.7	0.4	2.4	0.7
	A.20.	A 0	2.8	1.4	2.6	0.8
		50		0.5		0.8
		B 0	2.6	1.0	2.5	1.5
		50	2.4	0.4	1.4	0.1
30						

TABLE I (cont'd)

	Compound Number		LESION SCORES			
			<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>	
			<u>monensin</u>		<u>monensin</u>	
			<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>
5	A.21.	A 0	2.5	0.9	2.4	1.7
		50		1.3		1.0
		B 0	2.6	1.0	2.5	1.5
		50	2.3	1.0	2.2	0.7
10	A.22.	A 0	3.2	2.3	3.0	1.8
		50		1.3		0.8
		B 0	2.6	1.0	2.5	1.5
		50	2.7	1.3	2.2	0.6
15	A.23.	A 0	2.6	0.6	2.7	0.9
		50		0.5		0.6
		B 0	2.6	1.0	2.5	1.5
		50	2.8	0.6	2.0	0.4
20	A.24.	A 0	2.5	0.9	2.4	1.7
		50		0.9		0.6
		B 0	2.6	1.0	2.5	1.5
		50	2.8	0.8	2.2	0.7
25	A.25.	A 0	2.7	0.6	2.7	0.9
		50		1.1		0.3
		B 0	2.6	1.0	2.5	1.5
		50	2.9	0.9	1.8	0.6
30	A.26.	A 0	2.6	0.6	2.7	0.9
		50		0.7		0.4
		B 0	2.6	1.0	2.5	1.5
		50	2.8	1.3	2.3	0.8

TABLE I (cont'd)

5	Compound Number		LESION SCORES			
			<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>	
			<u>monensin</u>		<u>monensin</u>	
			<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>
5	A.27.	A 0	2.6	0.6	2.7	0.9
		50		0.3		0.7
	B	0	2.6	1.2	2.6	1.5
		50	2.8	0.9	2.6	1.2
10	A.28.	A 0	2.6	0.6	2.7	0.9
		50		0.4		0.9
	B	0	2.6	1.2	2.6	1.5
		50	2.7	0.8	2.2	0.7
15	A.29.	A 0	2.6	0.6	2.7	0.9
		50		0.8		0.4
	B	0	2.6	1.2	2.6	1.5
		50	1.7	1.4	1.2	0.6
20	A.30.	A 0	2.5	0.9	2.4	1.7
		50		0.8		0.4
	B	0	2.6	1.2	2.6	1.5
		50	2.9	0.2	2.0	0.2
25	A.31.	A 0	2.7	1.6	2.5	1.6
		50		0.3		0.7
	B	0	2.6	1.2	2.6	1.5
		50	2.8	0.8	2.7	1.2
30	A.32.	A 0	2.7	1.6	2.5	1.6
		50		0.4		0.6
	B	0	2.6	1.2	2.6	1.5
		50	2.8	0.7	2.1	0.9
30	B	0	2.4	1.2	2.2	1.8
		50	2.5	1.0	2.9	1.0

TABLE I (cont'd)

		LESION SCORES				
		<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>		
		<u>monensin</u>		<u>monensin</u>		
<u>Compound</u>		<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>	
<u>Number</u>						
5	A.33.	B 0	2.8	1.2	2.7	1.6
		50	2.8	0.5	2.6	1.0
	A.34.	B 0	2.8	1.2	2.7	1.6
		50	2.9	0.3	2.2	0.3
10	A.35.	B 0	2.8	1.2	2.7	1.6
		50	2.8	0.2	3.0	0.9
	A.36.	B 0	2.8	1.2	2.7	1.6
		50	2.2	0	2.6	0.1
	A.37.	B 0	2.8	1.2	2.7	1.6
		50	2.0	0.4	2.9	1.2
15	A.38.	B 0	2.8	1.2	2.7	1.6
		50	2.6	0.4	2.9	1.3
	A.39.	A 0	3.6	1.6	2.8	1.2
		50		1.4		0.5
		B 0	3.6	2.0	2.9	1.3
		50	3.1	0.5	3.0	0.8
20	A.40.	A 0	3.6	1.6	2.8	1.2
		50		1.3		0.3
		B 0	3.6	2.0	2.9	1.3
		50	2.6	0.8	2.6	0.8
25						

TABLE I (cont'd)

		LESION SCORES			
		<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>	
		<u>monensin</u>		<u>monensin</u>	
	<u>Compound</u>		<u>0</u>	<u>50</u>	
	<u>Number</u>				
5	A.41.	A 0	3.6	1.6	2.8
		50		1.3	0.2
		B 0	3.6	2.0	2.9
		50	2.1	0	2.2
10	A.42.	A 0	2.7	1.6	2.6
		50		0.3	0.7
		B 0	3.6	2.0	2.9
		50	3.2	2.5	2.1
	A.43.	A 0	2.7	1.6	2.6
		50		1.3	0.4
15		B 0	3.2	1.1	1.9
		50	3.1	0.8	2.3
	A.44.	A 0	3.6	1.6	2.8
		50		1.2	0.4
		B 0	3.2	1.1	1.9
		50	2.9	0.5	2.6
20	A.45.	A 0	3.0	0.9	2.8
		50		0.8	0.3
		B 0	3.2	1.1	1.9
		50	2.7	1.0	2.1
25					0.2

TABLE I (cont'd)

5	Compound Number		LESION SCORES			
			<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>	
			<u>monensin</u>		<u>monensin</u>	
			<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>
10	A.46.	A 0	3.0	0.7	2.8	0.2
		50		1.0		0.2
		B 0	3.2	1.1	1.9	0.8
		50	2.9	0.9	2.1	0.2
	A.47.	A 0	2.8	1.0	3.0	1.2
		50		1.4		0.7
15		B 0	3.2	1.1	1.9	0.8
		50	2.7	0.9	2.4	0.5
	A.48.	A 0	2.8	1.0	3.0	1.2
		50		0.9		0.7
		B 0	3.2	1.1	1.9	0.8
		50	3.3	1.0	2.3	0.4
20	A.49.	A 0	3.0	0.7	2.8	0.2
		50		0.6		0.4
		B 0	3.2	1.1	1.9	0.8
		50	3.1	0.3	2.8	0.2
	A.50.	B 0	2.8	1.0	1.7	0.5
		50	3.0	0	2.0	0

25

30

TABLE I (cont'd)

		LESION SCORES			
		<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>	
		<u>monensin</u>		<u>monensin</u>	
		<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>
5	<u>Compound</u>				
	<u>Number</u>				
	A.51.	B 0	2.8	1.0	1.7 0.5
		50	1.3	0	1.1 0
10	A.52.	B 0	2.8	1.0	1.7 0.5
		50	2.8	0	2.1 0.2
	A.53.	A 0	2.7	1.6	2.5 1.6
		50		0.8	0.9
15		B 0	2.4	1.2	2.2 1.8
		50	2.8	0.8	2.6 1.4
	A.54.	A 0	3.1	2.0	2.4 0.6
		50		1.7	0.6
20		B 0	2.4	1.2	2.2 1.8
		50	2.6	0.7	2.9 0.4
	A.55.	B 0	2.4	1.2	2.2 1.8
		50	2.9	0.8	2.5 0.1
25	A.56.	B 0	2.4	1.2	2.2 1.3
		50	2.8	0.6	2.8 0.2
	A.57.	A 0	2.7	1.6	2.5 1.6
		50		0.7	0.9
30		B 0	2.4	1.2	2.2 1.3
		50	3.0	1.2	2.8 0.9
	A.58.	A 0	2.7	1.6	2.5 1.6
		50		0.9	0.7
		B 0	2.4	1.2	2.2 1.8
		50	2.7	0.7	2.7 0.8

TABLE I (cont'd)

		LESION SCORES				
		<u>Eimeria acervulina</u>		<u>Eimeria tenella</u>		
		<u>monensin</u>		<u>monensin</u>		
<u>Compound Number</u>		<u>0</u>	<u>50</u>	<u>0</u>	<u>50</u>	
5	A.59.	A 0	2.9	0.9	2.2	1.7
		50		0.5		1.4
		B 0	2.4	1.2	2.2	1.8
		50	2.9	0.7	2.9	1.1
10	A.60.	A 0	2.7	1.6	2.5	1.6
		50		0.6		0.7
		B 0	2.7	1.2	2.1	2.1
		50	2.9	0.8	2.7	1.6
15	A.61.	A 0	3.0	1.5	2.8	1.9
		50		0.8		1.7
		B 0	2.0	0.2	2.8	1.3
		50	2.0	0.7	3.1	2.2
20	A.62.	B 0	2.0	0.2	2.8	1.3
		50	0.4	0	2.4	0.8
	A.63.	B 0	2.0	0.2	2.8	1.3
		50	2.6	0.1	2.9	0.6
	A.64.	B 0	2.0	0.2	2.8	1.3
		50	0.7	0	2.4	0.8

TABLE II

LESION SCORES

COMPOUND NO.	LESION SCORES									
	<i>Eimeria acervulina</i> monensin					<i>Eimeria tenella</i> monensin				
C.1.		0	50	100		0	50	100		
	0	3.1	1.3	0.2		2.0	0.5	0		
	50	2.8	0.8			2.0	0.3			
	100	2.1				2.1				
		0	20	40	80	160	0	20	40	80
C.5.	0	2.9	2.6	1.2	0.4	0	2.6	2.9	2.2	0.6
	40	3.0		2.3			2.7		2.5	
	80	2.7			0.2		2.6			0.5
	160	2.4					1.9			
		0	20	40	80	160	0	20	40	80
	0	2.9	2.6	1.2	0.4	0	2.6	2.9	2.2	0.6
	40	3.0		1.0			1.4		1.7	
	80	2.4			0.3		2.9			1.3
	160	3.0					2.7			

TABLE II (cont'd)

COMPOUND NO.	LESION SCORES											
	<i>Eimeria acervulina</i> monensin						<i>Eimeria tenella</i> monensin					
C.6.		0	50	100	0	0	0	50	100			
	0	3.1	1.3	0.2			2.0	0.5	0			
	50	3.3	1.3				2.0	0.8				
	100	2.7					2.4					
		0	20	40	50	80	160	0	20	40	50	80
C.8.	0	2.9	2.6	1.2	1.8	0.4	0	2.6	2.9	2.2	1.6	0.6
	100											
	200	1.7					2.0					
		0	50	100			0	50	100			
	0	3.1	1.3	0.2			2.0	0.5	0			
C.11.	50		0.7					0.1				
	100											
		0	50	100			0	50	100			
	0	3.1	1.3	0.2			2.0	0.5	0			
	50	2.6	0.2				1.9	0				
	100	0.2					0.2					

TABLE IV (cont'd)

COMPOUND NO.	LESION SCORES											
	<i>Eimeria acervulina</i> monensin						<i>Eimeria tenella</i> monensin					
C.13.	0	3.1	0	50	100	0	0	50	100	0	50	100
	0	3.1	1.3	0.2	0	2.0	0.5	0	0	0	0.5	0
	50	0	0	0	0	0.1	0	0	0	0.1	0	0
	100	0	0	0	0	0.2	0	0	0	0.2	0	0
	0	2.9	2.6	1.8	0.7	0.8	0	20	40	80	160	0
C.58.	0	2.9	2.6	1.8	0.7	0.8	0	20	40	80	160	0
	10	2.6	1.8	0.7	0.8	0	2.6	2.9	2.2	0.6	0.2	0
	20	2.4	0.7	0.8	0	0	3.0	1.4	0	0	0	0
	40	0.8	0	0	0	0	2.8	0.7	0	0	0	0
	0	3.1	1.3	0.2	0	2.0	0.5	0	0	0	0.5	0
C.62.	0	3.1	1.3	0.2	0	2.0	0.5	0	0	0	0.5	0
	50	2.9	1.3	0.2	0	0.7	2.2	2.9	2.2	0.6	0.2	0
	100	2.9	1.3	0.2	0	0	0	50	100	0	0.4	0
	0	3.1	1.3	0.2	0	0.7	2.2	2.9	2.2	0.6	0.2	0
	50	2.9	1.3	0.2	0	0	0	50	100	0	0.4	0

LESION SCORES

TABLE III

LESION SCORES

COMPOUND NO.	<u>Elmeria acervulina*</u>										<u>Elmeria tenella*</u>									
	<u>monensis</u>										<u>monensis</u>									
C.1.	0	2.9	2.8	1.7	0.2	0	0	2.9	2.7	1.3	0.3	0.1	0	0	2.9	2.4	2.0	2.9	2.6	
	40	3.1		1.0	0.1									40	3.1		1.0			
	80	2.7		1.1	0.1									80	2.7		1.1			
	120	2.9												120	2.9					
	240	2.8												240	2.8					
C.5.	0	2.9	2.8	1.7	0.2	0	0	2.9	2.7	1.3	0.3	0.1	0	0	2.9	2.4	2.0	2.9	2.6	
	40	3.1		1.0	0.1									40	3.1		1.0			
	80	2.7		1.1	0.1									80	2.7		1.1			
	120	2.9												120	2.9					
	240	2.8												240	2.8					
C.58.	0	2.9	2.8	1.7	0.2	0	0	2.9	2.7	1.3	0.3	0.1	0	0	2.9	2.4	2.0	2.9	2.6	
	40	3.1		1.0	0.1									40	3.1		1.0			
	80	2.7		1.1	0.1									80	2.7		1.1			
	120	2.9												120	2.9					
	240	2.8												240	2.8					
C.76.	0	2.9	2.8	1.7	0.2	0	0	2.9	2.7	1.3	0.3	0.1	0	0	2.9	2.4	2.0	2.9	2.6	
	40	3.1		1.0	0.1									40	3.1		1.0			
	80	2.7		1.1	0.1									80	2.7		1.1			
	120	2.9												120	2.9					
	240	2.8												240	2.8					

* Infected with 1,000,000 oocysts of Elmeria acervulina and 250,000 oocysts of Elmeria tenella.

TABLE IV

LESION SCORES										
		<u>Eimeria acervulina*</u>				<u>Eimeria tenella*</u>				
		<u>Eimeria maxima*</u>								
<u>Compound</u>		<u>(Intestinal)</u>				<u>(Cecal)</u>				
<u>Number</u>		<u>Monensin</u>				<u>Monensin</u>				
		0	25	50	100	0	25	50	100	
5	A.1.	0	4.9	2.8	.3	0	3.3	3.3	.9	0
		25	3.7	.3	.07		3.2	.67	.13	
		50	2.1	0	0		2.7	0	.07	
		100	1.3				.6			
10		0	25	50	100	0	25	50	100	
	A.2.	0	5.93	3.7	2.1	0	2.83	3.1	1.2	.47
		25	1.72	.33	.13		1.78	.4	0	
		50	.45	0	0		.55	.2	0	
		100	0			0				

* Infected with 500,000 oocysts of Eimeria acervulina, 60,000 oocysts of Eimeria maxima, and 40,000 oocysts of Eimeria tenella. Test conducted with three replicates.

Demonstration of the synergistic effect possessed by the preferred carbanilides of the invention, i.e. nicarbazin or 4,4'-dinitrocarbanilide alone, in combination with the polyether was obtained as follows: one-week-old broiler chicks were allotted to five-bird cages and were fed a medicated or control ration, typically for one day, prior to infection with oocysts of a coccidiosis-causing organism. The chicks were maintained on their respective rations for a period of time, typically seven days. Generally, there were from three to six replicates per treatment. Anticoccidial efficacy was typically determined by the lesion scores, but other measures of efficacy were employed in many of the tests. In determining lesion scores, the birds were sacrificed and the severity of lesions scored on a 0-4 scale, with lesion-free birds scored as 0, extremely severe infections scored as 4, and intermediate degrees of infection scored as 1, 2, or 3. The scores of all birds which received a given treatment were averaged.

In those evaluations where data is reported with superscript letters, data not followed by a common letter are significantly different ($P < .05$).

The results of evaluations follows.

Test 1: Eimeria acervulina (strain FS-254), inoculated with 200,000 oocysts.

Lesion Scores

		monensin	
	ppm	0	100
	0	3.16 ^c	1.58 ^b
nicarbazin	125	1.60 ^b	0 ^a

Test 2: Eimeria tenella (FS-226), inoculated with 100,000 oocysts.

Lesion Scores

5		ppm	monensin	
			0	100
		0	3.16 ^c	1.09 ^b
	nicarbazin	125	1.24 ^b	0.18 ^a

Test 3: Eimeria acervulina (strain FS-254), inoculated with 40,000 oocysts

10

Lesion Scores

15		ppm	monensin		
			0	60	100
		0	3.36 ^e	0.95 ^c	0.80 ^c
	nicarbazin	60	1.05 ^{cd}	0 ^a	0 ^a
		100	0.50 ^b		

Test 4: Eimeria acervulina (strain FS-254), inoculated with 200,000 oocysts.

20

Lesion Scores

		ppm	monensin	
			0	100
		0	3.45 ^c	1.70 ^b
	nicarbazin	75	2.95 ^c	0 ^a

25 Test 5: Eimeria tenella (strain FS-257), inoculated with 100,000 oocysts

Lesion Scores

30		ppm	monensin	
			0	100
		0	3.46 ^c	1.40 ^b
	nicarbazin	125	1.20 ^b	0.15 ^a

Test 6: Eimeria acervulina (strain FS-254), inoculated with 1,000,000 oocysts

Lesion Scores

5		ppm	monensin		
			0	60	100
		0	3.45 ^d	1.85 ^c	1.15 ^b
	nicarbazin	60	2.95 ^d	0.50 ^a	0.11 ^a
		100	3.00 ^d		

10 Test 7: Combination of Eimeria acervulina and Eimeria maxima (culture FS-266), 500,000 oocysts.

Lesion Scores*

15		ppm	monensin			
			0	40	80	120
		0	5.2 ^c	1.8 ^b	0.5 ^a	0.1 ^a
	nicarbazin	40	1.8 ^b	0.4 ^a	0.4 ^a	
		80	1.1 ^{ab}	0.3 ^a		

20 *Lesion scores were determined at three locations, anterior, mid-, and posterior portions of the small intestine, scored on 0-4 in each section and expressed as the total.

25 Test 8: Combination of Eimeria acervulina (strain FS-254; 250,000 oocysts, and Eimeria tenella (strain FS-257), 50,000 oocysts. One group of birds was sacrificed at 5 days; the other group was used to evaluate oocyst production and then sacrificed at 7 days.

30

Intestinal lesion scores at 5-days (Eimeria acervulina)

		monensin				
	ppm	0	20	40	60	80
	0	3.70f			1.50 ^{cde}	1.13 ^{bcd}
nicarbazin	20		1.63 ^{de}	1.20 ^{bcd}	1.40 ^{cde}	0.40 ^a
	40		1.00 ^{bc}	0.70 ^{ab}	0.60 ^{ab}	
	60	1.83 ^e	1.10 ^{bcd}	0.70 ^{ab}	0.40 ^a	
	80	1.60 ^{cde}	0.80 ^{ab}			
	100	1.50 ^{cde}				
					100	1.50 ^{cde}

Cecal lesion scores at 7-days (Eimeria tenella)

	ppm	monensin				
		0	20	40	60	80
	0	3.05 ^f			2.90 ^f	2.00 ^e
nicarbazin	20		2.86 ^f	1.90 ^{de}	1.20 ^{cd}	0.55 ^{abc}
	40		1.15 ^c	0.92 ^{bc}	0.75 ^{abc}	0 ^a
	60	2.05 ^e	1.00 ^{bc}	0.65 ^{bc}	0.30 ^{ab}	
	80	0.75 ^{abc}	0.65 ^{abc}	0.10 ^a		
	100	0.66 ^{abc}				
						100
						1.21 ^{cd}

Average oocyst passage/Bird(x10⁶)*

		monensin					
	ppm	<u>0</u>	<u>20</u>	<u>40</u>	<u>60</u>	<u>80</u>	<u>100</u>
	0	64.8			40.9	53.3	1.1
nicarbazin	20		64.6	55.6	7.5	2.0	
	40		17.1	22.9	8.7	0.7	
	60		7.3	4.4	0.9		
	80		1.4	4.7			
	100		18.7				

*for a 48-hour period, beginning on the 5th day following inoculation and continuing through to sacrifice.

Test 9: Eimeria acervulina (strain FS-273) inoculated with 1,150,000 oocysts.

Lesion Scores

5		ppm	narasin			
			0	25	50	100
		0	3.6 ^f	3.3 ^f	2.4 ^e	1.6 ^d
	nicarbazin	25	3.6 ^f	3.2 ^f	0.6 ^b	
		50	3.6 ^f	1.1 ^c	0.1 ^a	
		100	1.8 ^d			

10

Average survivor weight gain in grams

15		ppm	narasin			
			0	25	50	100
		0	126.7 ^a	159.6 ^c	193.3 ^{de}	183.4 ^d
	nicarbazin	25	142.7 ^b	188.3 ^d	212.9 ^{fg}	
		50	159.1 ^c	203.7 ^{ef}	216.8 ^{fg}	
		100	188.3 ^d			
	(noninfected, nonmedicated controls = 219 grams)					

20

Average feed/gain

25		ppm	narasin			
			0	25	50	100
		0	2.23 ^d	1.96 ^{bc}	1.62 ^a	1.60 ^a
	nicarbazin	25	2.04 ^c	1.64 ^a	1.49 ^a	
		50	1.89 ^b	1.50 ^a	1.50 ^a	
		100	1.61 ^a			
	(noninfected nonmedicated controls = 1.49 ^a)					

30

Comprehensive anticoccidial indices*

	ppm	narasin			
		0	25	50	100
nicarbazin	0	1.38(0)	1.68(22)	2.09(52)	2.17(58)
	25	1.49(8)	1.91(39)	2.52(84)	
	50	1.62(17)	2.39(75)	2.66(94)	
	100	2.16(57)			

(noninfected, nonmedicated controls = 2.74 (100))

*Index =

[growth and survival ratio] - [average lesion score/X].
 Where: $X = 4 / [1.25 \times \text{ave. growth and survival ratio of noninfected nonmedicated controls}]$. Growth and survival ratio = [pen weight at termination/pen weight at initiation], adjusted for mortality due to causes other than coccidiosis. Average of five replicates per treatment. The number in parentheses is the percent of optimum anticoccidial activity = [index of infected medicated group - index of infected controls] / [index of noninfected nonmedicated group - index of infected controls] x 100.

Test 10: Eimeria acervulina (strain FS-254), inoculated with 250,000 oocysts.

Lesion Scores

5		monensin			
		<u>ppm</u>	<u>0</u>	<u>50</u>	<u>100</u>
		0	2.38 ^e	1.36 ^{cd}	0.73 ^{abc}
	nicarbazin	60	1.80 ^{de}	0.44 ^{ab}	0.17 ^a
		120	1.00 ^{bc}		

10 Average oocyst passage/Bird (x10⁶)*

			monensin		
		<u>ppm</u>	<u>0</u>	<u>50</u>	<u>100</u>
		0	37.4	14.8	16.0
	nicarbazin	60	44.5	0.7	2.1
15		120	44.5		

*for a 24-hour period, beginning on the 5th day following inoculation and continuing through to sacrifice on the 6th day.

Test 11: Eimeria tenella (strain FS-286), inoculated with 125,000 oocysts.

Percent mortality attributable to coccidiosis

	monensin				
	0	25	50	100	
	0 32 ^d	24 ^{cd}	8 ^{ab}	0 ^a	
nicarbazin	25 16 ^{bc}	0 ^a	0 ^a		
	50 4 ^{ab}	0 ^a	0 ^a		
	100 0 ^a				

noninfected nonmedicated controls = 0^a

Average survivor weight gain in grams

	monensin				
	0	25	50	100	
	0 153.4 ^a	165.8 ^{ab}	198.7 ^{cd}	220.3 ^{de}	
nicarbazin	25 185.6 ^{bc}	242.5 ^{ef}	227.9 ^e		
	50 224.2 ^e	246.4 ^{ef}	244.4 ^{ef}		
	100 237.8 ^{ef}				

noninfected nonmedicated controls = 258.2^f

Average feed/gain

	monensin				
	0	25	50	100	
	0 *	1.64 ^{cd}	1.68 ^d	1.53 ^{bc}	
25 nicarbazin	25 1.79 ^d	1.41 ^a	1.44 ^a		
	50 1.50 ^{abc}	1.43 ^a	1.42 ^a		
	100 1.44 ^{ab}				

noninfected nonmedicated controls = 1.48^{ab}

*no data because of mortality which occurred in all replicates

Average cecal lesion score per bird

		monensin			
		<u>0</u>	<u>25</u>	<u>50</u>	<u>100</u>
		0 3.9 ^e	3.8 ^{de}	3.3 ^d	0.8 ^{ab}
5	nicarbazin	25 3.8 ^{de}	2.1 ^c	0.8 ^{ab}	
		50 3.3 ^d	0.4 ^a	0.3 ^a	
		100 1.3 ^b			

Comprehensive anticoccidial indices*

		monensin			
		<u>0</u>	<u>25</u>	<u>50</u>	<u>100</u>
10		0 0.80(0)	1.08(13)	1.75(43)	2.56(80)
	nicarbazin	25 1.34(24)	2.54(79)	2.69(85)	
		50 2.07(58)	2.88(94)	2.93(96)	
		100 2.70(86)			

15 noninfected nonmedicated controls = 3.01 (100)

*for method of calculation, see Test 9, above.

Test 12: Eimeria tenella (strain FS-226-A-204R, a strain propagated in the presence of 15 ppm A-204 for 13 generations prior to use in this experiment), inoculated with 130,000 oocysts.

Mortality attributable to coccidiosis

		narasin		
		<u>0</u>	<u>25</u>	<u>50</u>
25		0 13.3 ^b	6.7 ^a	6.7 ^a
	nicarbazin	25 0 ^a	0 ^a	0 ^a
		50 0 ^a	0 ^a	0 ^a

noninfected nonmedicated controls = 0^a

30

Average survivor weight gain in grams

		narasin		
		<u>0</u>	<u>25</u>	<u>50</u>
5	nicarbazin	0 172.2 ^a	243.4 ^c	239.1 ^c
		25 198.4 ^b	238.1 ^c	234.1 ^c
		50 231.7 ^c	238.7 ^c	244.9 ^c
		noninfected nonmedicated controls = 239.7 ^c		

Average feed/gain

		narasin		
		<u>0</u>	<u>25</u>	<u>50</u>
10		0 1.88 ^b	1.51 ^a	1.48 ^a
	nicarbazin	25 1.75 ^b	1.57 ^a	1.51 ^a
		50 1.57 ^a	1.56 ^a	1.49 ^a
		noninfected nonmedicated controls = 1.58 ^a		

Average cecal lesion score per bird

		narasin		
		<u>0</u>	<u>25</u>	<u>50</u>
20	nicarbazin	0 3.8 ^d	2.7 ^c	1.5 ^b
		25 3.5 ^d	0.6 ^a	0.2 ^a
		50 2.5 ^c	0.3 ^a	0 ^a

Comprehensive anticoccidial indices*

		narasin		
		<u>0</u>	<u>25</u>	<u>50</u>
25	nicarbazin	0 1.36(0)	2.24(55)	2.40(65)
		25 1.92(35)	2.85(93)	2.80(90)
		50 2.43(67)	2.86(94)	2.92(98)

noninfected nonmedicated controls = 2.96 (100)

*for method of calculation, see Test 9, above.

Test 13: Eimeria acervulina (strain FS-254), inoculated with 1,000,000 oocysts.

Average survivor weight gain in grams

		narasin		
		0	25	50
5		0 147.3 ^a	186.7 ^c	199.0 ^{cd}
	nicarbazin	25 157.3 ^{ab}	196.9 ^{cd}	205.5 ^{de}
		50 162.6 ^b	215.9 ^e	200.7 ^d
		noninfected nonmedicated controls = 217.1 ^e		

10

Average feed/gain

		narasin		
		0	25	50
15		0 2.09 ^c	1.82 ^b	1.62 ^a
	nicarbazin	25 1.81 ^b	1.57 ^a	1.51 ^a
		50 1.96 ^{bc}	1.54 ^a	1.53 ^a
		noninfected nonmedicated controls = 1.59 ^a		

Average intestinal lesion score per bird

		narasin		
		0	25	50
20		0 3.4 ^d	2.6 ^c	2.1 ^c
	nicarbazin	25 3.6 ^d	1.4 ^b	0.3 ^a
		50 3.3 ^d	0.6 ^a	0.1 ^a

25

30

Comprehensive anticoccidial indices*

narasin

		<u>0</u>	<u>25</u>	<u>50</u>
		0 1.58(0)	2.02(38)	2.23(57)
5	nicarbazin	25 1.61(3)	2.36(68)	2.62(91)
		50 1.76(16)	2.59(88)	2.60(89)
		noninfected nonmedicated controls = 2.73 (100)		
		*for method of calculation, see Test 9, above.		

10 Test 14: Eimeria acervulina (strain FS-273), inoculated with 780,000 oocysts.

Average survivor weight gain in grams

salinomycin

		<u>0</u>	<u>25</u>	<u>50</u>
		0 155.3 ^a	170.2 ^{ab}	199.7 ^{cd}
15	nicarbazin	25 164.5 ^{ab}	210.1 ^{de}	204.3 ^{de}
		50 180.0 ^{bc}	209.7 ^{de}	209.6 ^{de}
		noninfected nonmedicated controls = 226.8 ^e		

Average feed/gain

salinomycin

		<u>0</u>	<u>25</u>	<u>50</u>
		0 1.80 ^d	1.72 ^{bcd}	1.63 ^{abcd}
	nicarbazin	25 1.78 ^{cd}	1.56 ^{abcd}	1.50 ^{ab}
		50 1.68 ^{abcd}	1.53 ^{abc}	1.43 ^a
25		noninfected nonmedicated controls = 1.42 ^a		

Average intestinal lesion score per bird

		salinomycin		
		<u>0</u>	<u>25</u>	<u>50</u>
5	nicarbazin	0 3.9 ^d	3.7 ^d	2.9 ^{bc}
		25 3.9 ^d	2.1 ^b	0.2 ^a
		50 3.7 ^{cd}	0.1 ^a	0 ^a

Comprehensive anticoccidial indices*

		salinomycin		
		<u>0</u>	<u>25</u>	<u>50</u>
10		0 1.54(0)	1.70(15)	2.04(45)
	nicarbazin	25 1.60(6)	2.23(61)	2.54(89)
		50 1.77(21)	2.57(91)	2.60(94)

noninfected nonmedicated controls = 2.67 (100)

*for method of calculation, see Test 9, above.

15 Test 15: Eimeria acervulina (strain FS-273), inoculated with 780,000 oocysts.

Average survivor weight gain in grams

		lonomycin		
		<u>0</u>	<u>25</u>	<u>50</u>
20		0 145.7 ^a	162.4 ^{abc}	179.5 ^{cde}
	nicarbazin	25 154.0 ^{ab}	184.3 ^{de}	198.6 ^{ef}
		50 171.8 ^{bcd}	195.9 ^{ef}	212.7 ^f

noninfected nonmedicated controls = 210.9^f

25

30

Average feed/gain

		lonomycin		
		<u>0</u>	<u>25</u>	<u>50</u>
5	nicarbazin	0 1.89 ^d	1.68 ^{bc}	1.57 ^{ab}
		25 1.80 ^{cd}	1.55 ^{ab}	1.49 ^a
		50 1.68 ^{bc}	1.46 ^a	1.46 ^a

noninfected nonmedicated controls = 1.43^a

Average intestinal lesion score per bird

		lonomycin		
		<u>0</u>	<u>25</u>	<u>50</u>
10	nicarbazin	0 3.3 ^d	3.1 ^d	1.9 ^c
		25 2.8 ^d	2.1 ^c	0.7 ^b
		50 2.3 ^c	0.5 ^b	0 ^a

noninfected nonmedicated controls = 0

15

Comprehensive anticcoccidial indices*

		lonomycin		
		<u>0</u>	<u>25</u>	<u>50</u>
20	nicarbazin	0 1.61(0)	1.77(16)	2.12(48)
		25 1.76(15)	2.14(50)	2.48(83)
		50 1.97(35)	2.49(83)	2.62(96)

noninfected nonmedicated controls = 2.66(100)

*for method of calculation, see Test 9, above.

25

30

Test 16: Eimeria tenella (strain FS-226-A204R), inoculated with 130,00 oocysts.

Mortality attributable to coccidiosis

A-204

5

		<u>0</u>	<u>5</u>	<u>10</u>
	0	13.3 ^b	6.7 ^a	0 ^a
nicarbazin	50	0 ^a	0 ^a	0 ^a
	100	0 ^a	0 ^a	0 ^a

noninfected nonmedicated controls = 0^a

10

Average survivor weight gain in grams

A-204

		<u>0</u>	<u>5</u>	<u>10</u>
	0	200.9 ^{ab}	190.2 ^a	225.7 ^{bc}
nicarbazin	50	235.5 ^c	243.4 ^c	241.6 ^c
	100	232.8 ^c	227.2 ^{bc}	236.2 ^c

noninfected nonmedicated controls = 256.3^c

Average feed/gain

A-204

		<u>0</u>	<u>5</u>	<u>10</u>
	0	1.75 ^c	1.74 ^{bc}	1.63 ^{abc}
nicarbazin	50	1.50 ^a	1.49 ^a	1.48 ^a
	100	1.54 ^{ab}	1.67 ^{abc}	1.51 ^a

noninfected nonmedicated controls = 1.48^a

25

30

Average cecal lesion score per bird

A-204

		<u>0</u>	<u>5</u>	<u>10</u>
	0	3.8 ^e	3.1 ^e	2.3 ^d
5	nicarbazin 50	1.9 ^d	0.5 ^{bc}	0 ^a
	100	0.8 ^c	0.1 ^{ab}	0 ^a

Comprehensive anticoccidial indices*

A-204

		<u>0</u>	<u>5</u>	<u>10</u>
10		0 1.42(0)	1.71(20)	2.25(57)
	nicarbazin 50	2.42(70)	2.70(90)	2.79(96)
	100	2.53(78)	2.65(86)	2.72(91)
	noninfected nonmedicated controls = 2.85 (100)			
	*for method of calculation, see Test 9, above.			

15

The following test was conducted in two-week-old straight run turkeys.

Test 17: Eimeria meleagritidis (strain FS-230-MR, pass #7)

Mortality attributable to coccidiosis*

		monensin					
		<u>0</u>	<u>40</u>	<u>60</u>	<u>80</u>	<u>100</u>	<u>120</u>
20		0 6.2		6.2	0	0	0
	nicarbazin 40		0	0			
	60	6.2	0	0			
25		80 6.2					
	100	0					
	120	0					

noninfected nonmedicated controls = 0

*There were no significant differences among treatments $P < .05$.

30

Average survivor weight gain in grams

monensin

	0	40	60	80	100	120
	100.7 ^{ab}	135.8 ^{de}	118.8 ^{bcd}	124.8 ^{cd}	134.1 ^{de}	140.3 ^{de}
nicarbazin	40					
	60	98.8 ^{ab}	164.6 ^f			
	80	95.7 ^a	169.3 ^f			
	100	99.2 ^{ab}				
	120	112.4 ^{abc}				

noninfected nonmedicated controls = 221.6^g

		Average feed/gain				
		monensin				
		0	40	60	80	100
		2.42 ^{fgh}		2.17 ^{def}	2.16 ^{def}	2.07 ^{de}
nicarbazin	40		2.04 ^{cde}	1.80 ^{bc}		120 ^{bcd}
	60	2.63 ^h	1.90 ^{bcd}	1.77 ^b		2.00
	80	2.55 ^{gh}				
	100	2.53 ^{gh}				
	120	2.29 ^{efg}				
		noninfected nonmedicated controls = 1.77 ^a				

Growth and survival ratios*

		monensin					
		<u>0</u>	<u>40</u>	<u>60</u>	<u>80</u>	<u>100</u>	<u>120</u>
		0 1.35(0)		1.42(11)	1.53(30)	1.59(39)	1.61(44)
nicarbazin 40			1.59(40)	1.71(59)			
60		1.34(0)	1.64(48)	1.71(60)			
80		1.33(0)					
100		1.42(12)					
120		1.48(22)					

noninfected nonmedicated controls = 1.95 (100)

*Growth and survival ratio (GSR) = pen weight at termination/pen weight at initiation, adjusted for mortality due to causes other than coccidiosis.

Data expressed as the average of four replicates/treatment.

Numbers in parentheses are the % of optimum anticoccidial activity = [GSR of infected medicated group - GSR of infected controls]/[GSR of noninfected nonmedicated group - GSR of infected controls] x 100

The following additional tests were conducted in chickens, as described above.

Test 18: Eimeria acervulina (strain FS-254), inoculated with 1,000,000 oocysts.

5

<u>Lesion scores</u>					
		lasalocid			
	ppm	0	25	50	100
	0	2.7 ^{ef}	2.4 ^{cdef}	2.7 ^{def}	1.6 ^{bc}
10	nicarbazin 25	2.8 ^f	2.3 ^{bcdef}	1.0 ^{ab}	
	50	2.0 ^{bcde}	1.7 ^{bcd}	0.1 ^a	
	100	2.0 ^{bcde}			

Average oocyst passage/Bird (x10⁶)*

		lasalocid			
	ppm	0	25	50	100
15	0	51 ^{ab}	92 ^{ab}	93 ^{ab}	112 ^{ab}
	nicarbazin 25	152 ^b	10 ^{ab}	12 ^{ab}	
	50	83 ^{ab}	15 ^{ab}	5 ^a	
	100	44 ^{ab}			

20 *for a 24-hour period, 120-144 hours post inoculation.

Test 19: Eimeria tenella (strain FS-257), inoculated with 200,000 oocysts.

25

<u>Lesion scores</u>					
		lasalocid			
	ppm	0	25	50	100
	0	4.0 ^c	4.0 ^c	3.9 ^c	3.6 ^c
30	nicarbazin 25	3.9 ^c	3.4 ^c	2.5 ^b	
	50	3.6 ^c	1.9 ^{ab}	1.6 ^a	
	100	2.0 ^{ab}			

Average oocyst passage/Bird ($\times 10^6$) *

		lasalocid			
	<u>ppm</u>	<u>0</u>	<u>25</u>	<u>50</u>	<u>100</u>
5		25 ^{ab}	51 ^c	49 ^c	20 ^{ab}
	0				
	nicarbazin 25	31 ^b	13 ^{ab}	<1 ^a	
	50	5 ^a	0 ^a	0 ^a	
	100	0 ^a			

*for a 24-hour period, 144-168 hours
postinoculation

10

Test 20: Combination of Eimeria acervulina (strain FS-254), 300,000 oocysts, and Eimeria tenella (strain FS-287), 88,000 oocysts.

<u>Lesion Scores</u>				
	<u>Intestinal</u>		<u>Cecal</u>	
	<u>(Eimeria acervulina)</u>		<u>(Eimeria tenella)</u>	
	monensin			
	0	25	50	100
ppm				
0	2.1 ^{fg} /3.9 ^g	1.9 ^{efg} /3.9 ^g	1.7 ^{efg} /2.7 ^{efg}	0.7 ^{abcd} /1.7 ^{bcde}
4,4'-dinitro- carbanilide			0 ^a /0.1 ^a	
50				
100	0.3 ^{ab} /1.1 ^{abcd}			

Test 21: Eimeria tenella (strain FS-283), 125,000 oocysts.

Lesion Scores

		monensin			
	<u>ppm</u>	<u>25</u>	<u>50</u>	<u>100</u>	
5	0	2.8 ^{fg}	2.1 ^{cdefg}	0.7 ^{ab}	
	4,4'-dinitro-				
	carbanilide	50	1.1 ^{abcd}		
	100	2.4 ^{defg}			

10 Test 22: Combination of Eimeria acervulina (strain FS-280), 430,000 oocysts, and Eimeria tenella (strain FS-260), 43,000 oocysts.

15

20

25

30

Lesion Scores

Intestinal Cecal
(Eimeria acervulina)/(Eimeria tenella)

monensin

	0		25		50		75		100		125	
	ppm		ppm		ppm		ppm		ppm		ppm	
4,4'-dinitro- carbanilide	0	2.6 ^d /4.0 ^f	2.3 ^d /3.9 ^f	0.7 ^b /1.2 ^{cd}	1.5 ^c /2.5 ^e	0.2 ^a /0.4 ^{ab}	0.2 ^a /0.3 ^a	0 ^a /0 ^a	0.3 ^a /0.7 ^{abc}			
	25	2.6 ^d /3.9 ^f	0.8 ^b /2.2 ^e	0.2 ^a /0.3 ^a	0 ^a /0 ^a	0 ^a /0 ^a						
	50	0.9 ^b /3.6 ^f	0 ^a /0.2 ^a	0 ^a /0 ^a								
	75	0.2 ^a /2.3 ^e	0 ^a /0 ^a	0 ^a /0 ^a								
	100	0 ^a /1.6 ^d	0 ^a /0.1 ^a									
	125	0 ^a /1.0 ^{bcd}										

Test 23: Eimeria acervulina (750,000 oocysts)

A32887 (K41)

Lesion Scores

5	ppm	<u>0</u>	<u>25</u>	<u>50</u>	<u>100</u>
	0	3.07	3.20	1.68	0.48
Nicarbazin	25	2.43	0		
	50	0			
	100	0			

10

Test 24: Eimeria tenella (75,000 oocysts)

A32887 (K41)

Lesion Scores

15	ppm	<u>0</u>	<u>25</u>	<u>50</u>	<u>100</u>
	0	1.73	2.0	0.9	0.22
Nicarbazin	25	1.87	0.2		
	50	0.6		0	
	100	0.67			


20

25

30


CLAIMS

- 5 1. An anticoccidial composition for consumption by an avian species; characterized in that the composition comprises as active ingredients a polyether antibiotic selected from monensin (factors A, B, and C), laidlomycin, grisorixin, lenoremycin, salinomycin, narasin, lonomycin, alborixin, antibiotic A204, A32887
10 (K41), etheromycin, lasalocid (factors A, B, C, D, and E), isolasalocid A, lysocellin, mutalomycin, or antibiotic A23187 and a coccidiostatically active carbanilide derivative, associated with a carrier or feedstuff.
2. An anticoccidial composition as claimed in claim 1,
15 wherein the polyether antibiotic is monensin, narasin, lasalocid, salinomycin or A-204.
3. An anticoccidial composition as claimed in claim 2 wherein the polyether antibiotic is monensin.
4. An anticoccidial composition as claimed in claim 2
20 wherein the polyether antibiotic is narasin.
5. An anticoccidial composition as claimed in any one of claims 1 to 4, wherein the ratio of carbanilide to polyether is in the range of from 40:1 to 1:20.
6. An anticoccidial composition as claimed in claim 5
25 wherein the ratio is in the range of from 10:1 to 1:10.
7. An anticoccidial composition as claimed in claim 5 wherein the ratio is in the range of from 2:1 to 1:2.
8. An anticoccidial composition as claimed in any one of claims 1 to 7 wherein the carbanilide possesses the structural
30 formula (I):



35 where R^1 , R^2 and R^3 are the same or different and can each represent hydrogen; halogen; cyano; amino; nitro;

C_{1-6} alkyl; C_{1-4} alkoxy; C_{2-4} alkanoylamino; C_{1-4} alkylthio;
 C_{1-4} haloalkyl; C_{1-4} haloalkoxy; C_{1-4} haloalkylthio;
 C_{1-6} alkoxycarbonyl; C_{2-4} haloalkenyloxy; C_{1-4} alkoxycar-
 bonylthio; C_{1-4} alkylsulfonyl; C_{1-4} haloalkylsulfonyl;
 5 fluorosulfonyl; phenoxy optionally substituted by
 halogen, C_{1-4} haloalkyl or nitro; phenoxycarbonyl
 optionally substituted by C_{1-4} alkyl; phenyl; or phenyl-
 sulfonyl optionally substituted by nitro, acetamido,
 isopropylideneamino or a group of the formula



10
 15 where R^9 is C_{1-4} haloalkylthio or C_{2-4} haloalkenyloxy;
 R^4 and R^5 are the same or different and can each
 represent hydrogen or C_{1-4} alkyl;
 R^6 , R^7 and R^8 are the same or different and can each
 represent hydrogen; halo; cyano; amino; C_{2-4} alkanoyl-
 amino; N- C_{1-4} alkyl C_{2-4} alkanoylamino; nitro; C_{1-6} alkyl;
 20 C_{1-4} alkoxy; C_{1-4} alkylthio, C_{1-4} haloalkyl; C_{1-4} halo-
 alkoxy, C_{1-4} haloalkylthio, C_{2-4} haloalkenyloxy; C_{1-4} halo-
 alkylsulfonyl; C_{1-6} alkoxycarbonyl, aminosulfonyl or
 benzoyl;
 25 provided that at least one of R^1 , R^2 , R^3 , R^6 , R^7 and
 R^8 are other than hydrogen,
 and complexes of such compounds with 2-hydroxy-4,6-
 dimethylpyrimidine.

30 9. An anticoccidial composition as claimed
 in claim 8, wherein the compound of formula (I) is

- 3,3'-bis(trifluoromethyl)-4,4'-dichlorocarbanilide
- 3,3',5,5'-tetrakis(trifluoromethyl)carbanilide
- 5 3-chloro-4'-fluorocarbanilide
- 2-chloro-2'-fluorocarbanilide
- 3,4',5-tris(trifluoromethyl)carbanilide
- 3,4,5-trichlorocarbanilide
- 10 2-chloro-5-(trifluoromethyl)-4'-(ethoxycarbonyl)carbanilide
- 2,6-dimethyl-4'-(N-methylacetamido)carbanilide
- 2-methoxy-4'-acetamidocarbanilide
- 15 3-(trifluoromethyl)-4'-iodo-thiocarbanilide
- 2-fluoro-4'-(aminosulfonyl)carbanilide
- 2-methoxy-4'-isopropylcarbanilide
- 2-methyl-2',5'-diethoxycarbanilide
- 20 4-ethyl-2'-methoxycarbanilide
- 2-methyl-5-chloro-2',5'-dimethoxycarbanilide
- 2,4,4'-trimethyl-3'-nitrocarbanilide
- 2-amino-3-nitro-5-(trifluoromethyl)-2',4'-dimethylcarbanilide
- 25 2-amino-3,4'-dinitro-5-(trifluoromethyl)-2'-chlorocarbanilide
- 2-amino-3,5'-dinitro-5-(trifluoromethyl)-2'-fluorocarbanilide
- 30 2-amino-3-nitro-5-(trifluoromethyl)-2'-(ethoxycarbonyl)carbanilide

- 2-ethyl-6-sec-butylcarbanilide
- 2-isopropyl-2',4',6'-trimethylcarbanilide
- 2-amino-3-nitro-3',5-bis(trifluoromethyl)-
4'-chlorocarbanilide
- 5 2-ethyl-6-sec-butyl-4'-n-butoxycarbanilide
- 2-amino-3-nitro-5-(trifluoromethyl)-2',4',5'-
trichlorocarbanilide
- 10 2-amino-3,3'-dinitro-5-(trifluoromethyl)-4'-
chlorocarbanilide
- 2-amino-3-nitro-5-(trifluoromethyl)-2'-
methyl-4'-bromocarbanilide
- 15 2-amino-3-nitro-5-(trifluoromethyl)-2',6'-
dibromo-4'-fluorocarbanilide
- 2-amino-3-nitro-2',5-bis(trifluoromethyl)-
4'-chlorocarbanilide
- 20 3,3',5,5'-tetrakis(trifluoromethyl)thio-
carbanilide
- 2,4-dimethoxy-4'-(ethoxycarbonyl)carbanilide
- 4-(ethoxycarbonyl)-2'-methyl-6'-ethylcarbanilide
- 25 2,2'-dinitro-4,4'-bis(trifluoromethyl)car-
banilide
- 3,3',4,4',5,5'-hexachlorocarbanilide
- 3-nitro-4-chloro-4'-(trifluoromethyl)car-
banilide
- 30 4-chloro-3,4'-bis(trifluoromethyl)carbanilide

- 4,4'-dinitro-2,2'-bis(trifluoromethyl)carbanilide
- 4,4'-bis(trifluoromethyl)carbanilide
- 5 3-bromo-3',5'-dimethylcarbanilide
- 2,5-dichloro-4'-methyl-N²-ethylcarbanilide
- 2,5-dichloro-2',4'-difluorocarbanilide
- 2-amino-3-nitro-3',5,5'-tris(trifluoromethyl)-carbanilide
- 10 2,6-diethyl-4'-(ethoxycarbonyl)carbanilide
- 3-ethyl-3'-chloro-4'-methyl-N²-ethylcarbanilide
- 2,6-dimethyl-4'-(ethoxycarbonyl)carbanilide
- 15 4-methoxy-3'-acetamidocarbanilide
- 2-methoxy-4'-(n-butoxycarbonyl)carbanilide
- 4-(isobutoxycarbonyl)carbanilide
- 2,4'-bis(methoxycarbonyl)carbanilide
- 20 2',4-dichloro-3-nitro-3'-(trifluoromethyl)-carbanilide
- 3,4,4',5-tetrachloro-3'-(trifluoromethyl)-carbanilide
- 25 4-chloro-3-nitro-3',5'-bis(trifluoromethyl)-carbanilide
- 2,4,6-trimethyl-4'-(ethoxycarbonyl)carbanilide
- 2-(trifluoromethyl)-2'-ethyl-6'-isopropyl-carbanilide
- 30 4-chloro-3,3',5'-tris(trifluoromethyl)carbanilide

- 3,4,4',5-tetrachloro-3'-nitrocarbanilide
2,6-dimethyl-4'-benzoylcarbanilide
3,4-dimethyl-2'-ethoxycarbanilide
5 2-chloro-4,4'-bis(methylthio)carbanilide
2-methyl-2'-ethoxycarbanilide
4-chloro-2-methoxythiocarbanilide
4,4'-dinitro-N,N'-dimethylcarbanilide
10 4-(trifluoromethyl)-4'-nitrocarbanilide
3,3',5,5'-tetrakis(trifluoromethyl)-N,N'-
dimethylcarbanilide
4-phenoxy-4'-nitrocarbanilide
15 4-nitro-4'-(4-chlorophenoxy)carbanilide
4-nitro-4'-(3,4-dichlorophenoxy)carbanilide
4-nitro-3'-chloro-4'-(4-chlorophenoxy)
carbanilide
20 4-nitro-3',5'-dichloro-4'-(4-chlorophenoxy)-
carbanilide
4-nitro-3'-chloro-4'-(3,4-dichlorophenoxy)-
carbanilide
25 4-nitro-4'-(2,4-dichlorophenoxy)carbanilide
4-nitro-4'-(4-nitrophenoxy)carbanilide
4-nitro-3'-nitro-4'-(4-chlorophenoxy)-
carbanilide
30 4-nitro-3'-methyl-4'-(4-chlorophenoxy)-
carbanilide

- 4-nitro-2'-nitro-4'-(4-chlorophenoxy) -
carbanilide
- 4-nitro-4'-(2,4-dinitrophenoxy) carbanilide
- 5 4-nitro-3'-chloro-4'-(2-tert-butyl-4-chloro-
phenoxy) carbanilide
- 4-nitro-4'-(2-methyl-4-chlorophenoxy) -
carbanilide
- 10 4-nitro-3'-bromo-4'-(4-bromophenoxy) carbanilide
- 4-nitro-3'-bromo-4'-(4-chlorophenoxy) -
carbanilide
- 4-nitro-3'-chloro-4'-(4-bromophenoxy) -
carbanilide
- 15 3,3',4'-trichloro-4-(4-chlorophenoxy) -
carbanilide
- 3,4'-dichloro-4-(4-chlorophenoxy) carbanilide
- 3,4-dichloro-4'-(4-chlorophenoxy) carbanilide
- 20 2-methyl-4-nitro-3'-chloro-4'-(4-chloro-
phenoxy) carbanilide
- 3,3',5'-tris(trifluoromethyl)-4-methoxy-
carbanilide
- 25 3-(trifluoromethoxy)-3',5'-bis(trifluoro-
methyl) carbanilide
- 4-(trifluoromethoxy)-3',5'-bis(trifluoro-
methyl) carbanilide
- 30 2,3',5'-tris(trifluoromethyl)-4-(trifluoro-
methoxy) carbanilide

- 4-(methylthio)-3',5'-bis(trifluoromethyl)-
carbanilide
- 4-(methylthio)-3',5'-bis(trifluoromethyl)-
thiocarbanilide
- 5 3-chloro-4-(methylthio)-3',5'-bis(trifluoro-
methyl)carbanilide
- 3-(trifluoromethylthio)-3',5'-bis(trifluoro-
methyl)carbanilide
- 10 4-(trifluoromethylthio)-3',5'-bis(trifluoro-
methyl)carbanilide
- 2-chloro-4-(trifluoromethylthio)-3',5'-
bis(trifluoromethyl)carbanilide
- 15 4-(trifluoromethylthio)-3'-(trifluoromethyl)-
5'-nitrocarbanilide
- 4-(trifluoromethylthio)-2',5'-bis(trifluoro-
methyl)-4'-nitrocarbanilide
- 20 4,4'-bis(trifluoromethylthio)carbanilide
- 4-(trifluoromethylthio)-4'-(chloromethyl-
sulfonyl)carbanilide
- 3-(difluoromethylthio)-3',5'-bis(trifluoro-
methyl)carbanilide
- 25 4-(ethylthio)-3',5-bis(trifluoromethyl)-
carbanilide
- 3-chloro-4-(ethylthio)-3',5'-bis(trifluoro-
methyl)carbanilide
- 30 4-ethoxy-3,3',5'-tris(trifluoromethyl)car-
banilide

- 3-(trifluoromethyl)-4-ethoxy-4'-(trifluoromethylthio) carbanilide
- 5 3-(trifluoromethyl)-4-ethoxy-3'-(trifluoromethylthio) carbanilide
- 4-methyl-3-(2-chloroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 4-(1,2-dichlorovinyloxy)-3'-(trifluoromethyl)-4'-chlorocarbanilide
- 10 4-(1,2-dichlorovinyloxy)-3',5'-bis(trifluoromethyl) carbanilide
- 4-(2,2,2-trichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 15 2-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 3-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 20 3-(2,2-dichloro-1,1-difluoroethoxy)-4-bromo-3',5'-bis(trifluoromethyl) carbanilide
- 4-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 25 4-(2,2-dichloro-1,1-difluoroethoxy)-3'-(trifluoromethyl)-4'-chlorocarbanilide
- 3-methoxy-4-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 3-methyl-4-(2,2-dichloro-1,1-difluoroethoxy)-3',5'-bis(trifluoromethyl) carbanilide
- 30

- 3-methyl-4-(2,2-dichloro-1,1-difluoroethoxy)-
4'-isopropylcarbanilide
- 3-nitro-4-(2,2-dichloro-1,1-difluoroethoxy)-
3',5'-bis(trifluoromethyl)carbanilide
- 5 2-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl)carbanilide
- 4-(2,2-dichloro-1,1-difluoroethoxy)-3,3',5'-
tris(trifluoromethyl)carbanilide
- 10 3-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl)carbanilide
- 4-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl)carbanilide
- 15 4-(2-chloro-1,1,2-trifluoroethoxy)-3',5'-
bis(trifluoromethyl)thiocarbanilide
- 4-(2-chloro-1,1,2-trifluoroethoxy)-3'-
(trifluoromethyl)-5'-nitrocarbanilide
- 20 4,4'-bis(2-chloro-1,1,2-trifluoroethoxy)-
carbanilide
- 4-(2-chloro-1,1,2-trifluoroethoxy)-4'-
(trifluoromethylthio)carbanilide
- 25 4-(2-chloro-1,1,2-trifluoroethoxy)-3'-
(trifluoromethyl)-4'-chlorocarbanilide
- 4-(2-chloro-1,1,2-trifluoroethoxy)-3,3'-
bis(trifluoromethyl)-5'-nitrocarbanilide
- 30 4-(2-chloro-1,1,2-trifluoroethoxy)-3,3',5'-
tris(trifluoromethyl)carbanilide

- 3-(1,1,2-trifluoroethoxy)-3',5'-bis(tri-
fluoromethyl)carbanilide
- 3-(1,1,2,2-tetrafluoroethoxy)-3',5'-bis(tri-
fluoromethyl)carbanilide
- 5 3-methyl-4-(1,1,2,2-tetrafluoroethoxy)-
3'-(chlorodifluoromethyl)carbanilide
- 3-methyl-4-(1,1,2,2-tetrafluoroethoxy)-3'-
(trifluoromethyl)-4'-chlorocarbanilide
- 10 3-(trifluoromethyl)-4-(1,1,2,2-tetrafluoro-
ethoxy)-4'-bromocarbanilide
- 3-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3',5'-bis(trifluoromethyl)carbanilide
- 15 4-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3',5'-bis(trifluoromethyl)carbanilide
- 3-chloro-4-(1,1,2,3,3,3-hexafluoro-n-
propoxy)-3',5'-bis(trifluoromethyl)-
carbanilide
- 20 3-methyl-4-(1,1,2,3,3,3-hexafluoro-n-
propoxy)-3',5'-bis(trifluoromethyl)-
carbanilide
- 4-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3,3',5'-tris(trifluoromethyl)carbanilide
- 25 4-(1,1,2,3,3,3-hexafluoro-n-propoxy)-
3,3'-bis(trifluoromethyl)-5'-nitro-
carbanilide
- 30 4-(methoxycarbonylthio)-3',5'-bis(tri-
fluoromethyl)carbanilide

- 4-(4-chlorophenoxy)-4'-(trifluoromethyl-
thio) thiocarbanilide
- 4-(4-chlorophenoxy)-3',5'-bis(tri-
fluoromethyl) carbanilide
- 5 3-chloro-4-(4-chlorophenoxy)-3',5'-
bis(trifluoromethyl) carbanilide
- 4-(4-chlorophenoxy)-3'-(trifluoromethyl)-
5'-nitrocarbanilide
- 10 4-(3,5-bis(trifluoromethyl))phenoxy-3',5'-
bis(trifluoromethyl) carbanilide
- 4-(4-methylphenoxy carbonyl)-3',5'-
bis(trifluoromethyl) carbanilide
- 15 4-(4-methylphenoxy carbonyl)-4'-(tri-
fluoromethylthio) carbanilide
- 3,5-bis(trifluoromethyl)-2',4',6'-
trichlorocarbanilide
- 20 3,5-bis(trifluoromethyl)-2',4',5'-tri-
chlorocarbanilide
- 3,3'-bis(trifluoromethyl)-5'-nitro-
carbanilide
- 25 3,3',5-tris(trifluoromethyl)-5'-nitro-
carbanilide
- 2',3,5,6'-tetrakis(trifluoromethyl)-4'-
nitrocarbanilide
- 3-(chlorodifluoromethyl)-3',5'-bis(tri-
fluoromethyl) carbanilide
- 30 3-(1,1,2,2-tetrafluoroethyl)-3',5'-
bis(trifluoromethyl) carbanilide

- 4-phenyl-3',5'-bis(trifluoromethyl)-
carbanilide
- 3-(fluorosulfonyl)-3',5'-bis(trifluoro-
methyl) carbanilide
- 5 4-(fluorosulfonyl)-4'-(trifluoromethyl-
thio) carbanilide
- 4-(fluorosulfonyl)-3'-(trifluoromethyl)-
5'-nitrocarbanilide
- 10 4-(chloromethylsulfonyl)-3',5'-bis(tri-
fluoromethyl) carbanilide
- 2-(ethylsulfonyl)-3',5,5'-tris(trifluoro-
methyl) carbanilide
- 15 2-(ethylsulfonyl)-3',5-bis(trifluoro-
methyl)-5'-nitrocarbanilide
- 4-(trifluoromethylsulfonyl)-3',5'-
bis(trifluoromethyl) carbanilide
- 20 4-(trifluoromethylsulfonyl)-3'-(tri-
fluoromethyl)-4'-methoxycarbanilide
- 4-(trifluoromethylsulfonyl)-4'-(tri-
fluoromethylthio) carbanilide
- 25 4-(4-nitrophenylsulfonyl)-4'-(tri-
fluoromethylthio) carbanilide
- 4-(4-acetamidophenylsulfonyl)-4'-
(trifluoromethylthio) carbanilide
- 30 4-(4-(isopropylideneamino) phenylsul-
fonyl)-4'-(trifluoromethylthio) car-
banilide

carbanilide)

4,4-sulfonylbis(4'-(1,2-dichlorovinyl)-oxy)-
carbanilide)

5

4,4-sulfonylbis(3-(1,1,2,2-tetrafluoroethoxy)-
carbanilide)

4,4-sulfonylbis(4'-(2-chloro-2,2,3-trifluoro-
ethoxy)carbanilide) or

10

4,4-sulfonylbis(4'-(trifluoromethylthio)-
carbanilide).

10. An anticoccidial composition as claimed
in any one of claims 1 to 9, wherein the carbanilide
derivative is Nicarbazin.

15 11. A composition according to claim 1 to 10
wherein the carbanilide is present in a concentration of
from about 10 to about 250 ppm.

12. A composition according to claim 11,
wherein the concentration is from about 25 to about 100
ppm.

20

13. A composition according to any one of
claims 1 to 12 which contains from 15 ppm to 25% by
weight of active ingredients.

25 14. A composition according to any one of
claims 1, 2 or 4 to 13 which contains from about 25 to
about 100 ppm of narasin.

15. A composition according to any one of
claims 1 to 14 which comprises a total feed.

16 15. A composition as claimed in any one of
claims 1 to 15 for use in a method of controlling
coccidiosis in poultry, which method comprises supplying
the composition as at least part of their feed.

The text of the specification has been reproduced by photocopying the applicants original
typescript. It may contain a few amendments which are difficult to read. The original
typescript containing these amendments may be inspected on the premises of the Patent Office