



159159

A PHARMACEUTICAL COMPOSITION FOR TREATING DEPRESSION AND METHOD FOR PREPARATION THEREOF

FIELD OF THE INVENTION

[0001] The present invention relates to a pharmaceutical composition. In particular, the present invention relates to a pharmaceutical composition for treating depression as the main effect. The present invention further relates to a preparation method of the pharmaceutical composition for treating depression as the main goal.

BACKGROUND OF THE INVENTION

[0002] Depression is a common disease. According to statistics, about 25% females in the global population had been experiencing depression in their lives, and about 10% males had been experiencing depression (referring to *Morden Psychology* written by Ch'un-Hsing Chang). World Health Organization (WHO) published, "The incidence of depression in the world is about 11%. At present, about 340 million psychological depressed patients are in the world, and the number is increased. The investigation is found that the depression will be increased to be the number two common disease in the world from now on to 20 years later."

[0003] At present, anti-depression pharmaceuticals in the domestic and overseas markets consist mainly selective serotonin reuptake inhibitors (SSRIs), such as Prozac (fluoxetine hydrochloride), Paxil (Paroxetine or paroxetine hydrochloride) and Zoloft (sertraline hydrochloride), etc. These pharmaceuticals function by increasing the component and the content of serotonin in the human body to decrease and release the symptoms of depression. This kind of pharmaceuticals all have side effects of different



G00002

levels. The research is published that these pharmaceuticals have the ability to correct chemical imbalance in the human body, but more often than not, they are still unable to calm the patients. In recent years, whether the depression pharmaceuticals, such as Prozac, are harmful had become a serious social problem, whereas Paxil was even found to be harmful in 1996. Paxil has been recalled continually from the market since 2001. In June 2004, the New York State Attorney General accused GlaxoSmithKline Company of the Great British of beguilingly concealing the research report of the linkage between Paxil and "increased risk of suicidal behavior and tendencies in adolescents." In light of the current situation, the search for a new generation of pharmaceuticals with less side effects and more pronounced/potent anti-depression qualities has become the center of attention of the entire pharmaceutical world.

[0004] It is therefore attempted by the applicant to deal with the above situation encountered in the prior art.

SUMMARY OF THE INVENTION

[0005] In order to overcome the insufficiency of the modern technology, the purpose of the present invention provides a herbal pharmaceutical composition for anti-depression as the main effect. It can be used as pharmaceuticals or health food for improving the depression.

[0006] According to one aspect of the present invention, a pharmaceutical composition for treating depression is provided. The pharmaceutical composition comprises one of the following compositions of raw materials: (A) 4 ~ 18 parts by weight of a ginseng and 3 ~ 14 parts by weight of a liquorice; (B) a ginseng water extract or a ginseng ethanol extract extracted from 4 ~ 18 parts by weight of the ginseng and a liquorice water extract or a

liquorice ethanol extract extracted from 3 ~ 14 parts by weight of the liquorice; (C) 4 ~ 18 parts by weight of the ginseng and the liquorice water extract or the liquorice ethanol extract extracted from 3 ~ 14 parts by weight of the liquorice; and (D) the ginseng water extract or the ginseng ethanol extract extracted from 4 ~ 18 parts by weight of the ginseng and 3 ~ 14 parts by weight of the liquorice.

[0007] Preferably, the pharmaceutical composition further comprises: (E) 3 ~ 14 parts by weight of a jujuba; or (F) a jujuba water extract or a jujuba ethanol extract extracted from 3 ~ 14 parts by weight of the jujuba.

[0008] Preferably, the pharmaceutical composition further comprises: 9 parts by weight of the ginseng and 6 parts by weight of the liquorice; or the ginseng water extract or the ginseng ethanol extract extracted from 9 parts by weight of the ginseng and the liquorice water extract or the liquorice ethanol extract extracted from 6 parts by weight of the liquorice.

[0009] Preferably, the pharmaceutical composition further comprises: 6 parts by weight of the jujuba; or the jujuba water extract or the jujuba ethanol extract extracted from 6 parts by weight of the jujuba.

[0010] According to another aspect of the present invention, a pharmaceutical composition for treating depression is provided. The pharmaceutical composition comprises: a ginseng water extract or a ginseng ethanol extract extracted from 3 ~ 10 parts by weight of a ginseng; and 0.2 ~ 0.8 part by weight of a glycyrrhizic acid or a glycyrrhetic acid.

[0011] Preferably, the ginseng ethanol extract comprises 20 ~ 40% of a ginsenoside, and the glycyrrhizic acid or the glycyrrhetic acid has 80 ~ 98% purity.

[0012] Preferably, the pharmaceutical composition further comprises a jujuba water extract or a jujuba ethanol extract extracted from 0.05 ~ 0.2 part by weight of a jujuba.

[0013] Preferably, the pharmaceutical composition according to claim 7, wherein the jujuba ethanol extract comprises 0.5 ~ 3% of a jujuba cyclic adenosine monophosphate (cAMP).

[0014] Preferably, the pharmaceutical composition comprises 5 parts by weight of the ginseng ethanol extract, 0.4 part by weight of the glycyrrhetic acid and 0.1 part by weight of the jujuba ethanol extract, wherein the ginseng ethanol extract comprises 30% of the ginsenosides, the purity of the glycyrrhetic acid is 90%, and the jujuba ethanol extract comprises 1% of the jujuba cAMP.

[0015] According to another aspect of the present invention, a preparation method of a pharmaceutical composition is provided. The preparation method comprises steps of: (1) decocting 4 ~ 18 parts by weight of the ginseng in 60 ~ 77% of an ethanol solution, then purifying by a chromatography to obtain a first extract; (2) decocting 3 ~ 14 parts by weight of the liquorice in a water, then concentrating and drying to obtain a second extract; and (3) mixing and pulverizing the first extract obtained from the step (1) and the second extract obtained from the step (2), then sifting to obtain the pharmaceutical composition.

[0016] Preferably, the preparation method further comprises the step of: decocting 3 ~ 14 parts by weight of the jujuba in 60 ~ 75% of the ethanol solution, then purifying by the chromatography to obtain a third extract, and adding the third extract thereof into the pharmaceutical composition.

[0017] According to another aspect of the present invention, a preparation method of a pharmaceutical composition is provided. The preparation method comprises a step of: mixing and pulverizing 0.2 ~ 0.8 part by weight of the glycyrrhizic acid or the glycyrrhetic acid having 80 ~ 98% purity and 3 ~ 10 parts by weight of the ginseng extract having 20 ~ 40% of the ginsenoside to obtain the pharmaceutical composition.

[0018] Preferably, the preparation method further comprises the step of: compounding 0.05 ~ 0.2 part by weight of a jujuba extract having 1% of a jujuba cyclic adenosine monophosphate (cAMP) with a β -cyclodextrin to obtain a jujuba extract compound, and adding the jujuba extract compound thereof into the pharmaceutical composition.

[0019] Concretely speaking, there are only 2 to 3 pharmaceuticals, the ginseng, liquorice and/or jujuba in the pharmaceutical composition of the present invention.

[0020] Ginseng: The ginseng contains adenylate cyclase (AC) for stimulating adenosine, and the phosphodiesterase inhibitor. Both of the adenylate cyclase and phosphodiesterase inhibitor have synergism and collectively increase the cAMP in the cells. The phenylalanine is promoted by the ginseng to increase the synthesis of dopamine (DA) and norepinephrine (NE) through the blood-brain barrier, and thus the concentrations of the dopamine and norepinephrine are increased.

[0021] Liquorice: The glycyrrhizic acid and glycyrrhetic acid in liquorice are strong cAMP phosphodiesterase inhibitors. The cAMP degradation is decreased by inhibiting cAMP phosphodiesterase, and then the usage of cAMP in the central nervous system is increased.

[0022] Jujuba: The jujuba contains a large amount of cAMP-like materials. The extrinsic non-hydrated cAMP can be participated in the metastasis of cAMP in the body and be analogized the enzyme's function, and the cAMP in the cells is increased.

[0023] The ginseng, liquorice and jujuba in the pharmaceutical composition of the present invention are paired and acted collectively by stimulating the adenylate cyclase to increase the concentration of cAMP in brain cells, and by inhibiting the cAMP phosphodiesterase to decrease the degradation of cAMP and increase the usage of cAMP. The concentration and activity of the increased cAMP can increase the synthesis and release of neurotransmitters, such as norepinephrine, etc. (referring to *Volume One, Principles of Neurosciences* regarding the related description of the cAMP to the synthesis of catecholamine (CA)). This process is the mechanism of the modern pharmacology for anti-depression in this composition.

[0024] In other words, in order to accomplish the purpose of the present invention, the preferred parts by weight of compositions of the present invention are described as follows.

[0025] 1. Composition 1: 4 ~ 18 parts by weight of the ginseng and 3 ~ 14 parts by weight of the liquorice.

[0026] The preferred composition of the medicine prepared by the raw materials of the weight ratio is described as follows: 9 parts by weight of the ginseng and 6 parts by weight of the liquorice.

[0027] 2. Composition 2: 4 ~ 18 parts by weight of the ginseng, 3 ~ 14 parts by weight of the liquorice, and 3 ~ 14 parts by weight of the jujuba.

[0028] The preferred composition of the medicine prepared by the raw materials of the weight ratio is described as follows: 9 parts by weight of the ginseng, 6 parts by weight of the liquorice, and 6 parts by weight of the jujuba.

[0029] 3. Composition 3: 3 ~ 10 parts by weight of the ginseng ethanol extract (containing 20 ~ 40% of the ginsenoside), 0.2 ~ 0.8 part by weight of the glycyrrhetic acid (80 ~ 98% purity), and 0.05 ~ 0.2 part by weight of the jujuba ethanol extract (containing 0.5 ~ 3% of the jujuba cAMP).

[0030] In Composition 3, the preferred composition of the medicine prepared by the raw materials of the weight ratio is described as follows: 5 parts by weight of the ginseng ethanol extract (containing 30% of the ginsenoside), 0.4 part by weight of the glycyrrhetic acid (90% purity), and 0.1 part by weight of the jujuba ethanol extract (containing 1% of the jujuba cAMP).

[0031] In order to prepare the pharmaceutical composition of the present invention, the pulverized substance of the ginseng and liquorice is directly used according to the dictated weight ratio of the composition, and the pharmaceutical composition is directly prepared. Another pharmaceutical composition is prepared by adding the jujuba dry powder on the basis of this pharmaceutical composition.

[0032] In addition, according to the component weight ratio of the composition, either a dry powder of the raw material is adopted, and the water extract or the ethanol extract of the other component is added to prepare the pharmaceutical composition of the present invention, or a water extract or an ethanol extract of the raw material is adopted, and the dry powder of the other component is added to prepare the pharmaceutical composition of the present invention.

[0033] The preparation method of the pharmaceutical composition of the present invention includes:

[0034] Method 1:

[0035] 1. decocting 4 ~ 18 parts by weight of the ginseng in 60 ~ 77% concentration of the ethanol solution, separating and purifying by chromatography to obtain the first extract;

[0036] 2. decocting 4 ~ 18 parts by weight of the liquorice in the water, concentrating and drying to obtain the second extract; and

[0037] 3. mixing and sifting the first extract obtained from the step 1 and the second extract obtained from the step 2 to obtain the pharmaceutical composition 1 of the present invention.

[0038] The preferred composition of the medicine is 9 parts by weight of the ginseng and 6 parts by weight of the liquorice in the above method.

[0039] Method 2:

[0040] Three (3) ~ 14 parts by weight of the jujuba (the preferred composition of the medicine is 6 parts by weight) is further added and decocted in the ethanol solution in Method 1, is then separated and purified by chromatography, and is compounded with the β -cyclodextrin to obtain the jujuba extract compound. The jujuba extract compound is mixed and pulverized with the first extract and the second extract to obtain the pharmaceutical composition 2 of the present invention.

[0041] Method 3:

[0042] 1. Compounding 0.05 ~ 0.2 part by weight of the jujuba extract containing 1% of the jujuba cAMP with the β -cyclodextrin to obtain the jujuba extract compound.

[0043] 2. Mixing the jujuba extract compound, 0.2 ~ 0.8 part by weight of the glycyrrhetic acid having 90% purity and 3 ~ 10 parts by weight of the ginseng extract having 30% purity to obtain the pharmaceutical Composition 3 of the present invention.

[0044] The preferred parts by weight of each composition in the above method are: 0.1 part by weight of the jujuba extract having 1% of the jujuba cAMP (compounded with 9 parts by weight of the β -cyclodextrin), 5 parts by weight of the ginseng extract having 30% of the ginsenoside, and 0.4 part by weight of the glycyrrhetic acid having 90% purity.

[0045] The resolving scheme of the herbal pharmaceutical composition of the present invention is to cooperate with the treating mechanism of modern medicine and pharmacology with regards to depression, so as to investigate and develop a herbal pharmaceutical composition for the treatment of depression as the main goal based on the principles of Chinese medicine. The characteristics are that all the raw materials are pharmaceutical that doubles as food, the combinations of pharmaceuticals are simple (only 2 ~ 3 pharmaceuticals), the function and mechanism are definite (conforming with the function and mechanism of modern pharmacology), the effects and ingredients can be quantified, and the curative effect is significant and safe. This kind of plant derived pharmaceuticals that doubles as food has no toxicity or side effects. It can be used as pharmaceuticals or health food for treating depression and be taken on a long term basis.

[0046] The pharmaceutical composition of the present invention can be administrated as the unit dose formula, and the way of administration can be intestinal or non-intestinal, such as oral administration, etc. The media include tablet, capsule, pill, roll, powder, solution, suspension, emulsion, and

particle, etc. It can be prepared as immediate release, sustained release, controlled release, and microsphere delivery system. In order to prepare the unit delivery in tablet form, each carrier for one skilled in the art can be widely used. The examples regarding to the carriers are the dilutents and the absorbents, *i.e.* starch, dextrin, calcium sulphate, lactose, mannitol, sucrose, sodium chloride, glucose, urea, calcium carbonate, kaolin, microcrystalline cellulose, and aluminum silicate, etc. The further examples regarding to the carriers are the wetting agents and the bonding agents, *i.e.* water, glycerol, polyethylene glycol, ethanol, propanol, starch slurry, dextrin, syrup, honey, glucose solution, arabic mucilage, gelatin, sodium carboxymethylcellulose, lac, methyl cellulose, potassium phosphate, and poly vinyl pyrrolidone, etc. The further examples regarding to the carriers are the lysis agents, *i.e.* dried starch, alginate, agar, laminaran, sodium hydrogencarbonate, citric acid, calcium carbonate, polyoxyethylenesorbitanalkylester, sodium dodecyl-sulfonate, methyl cellulose, and ethyl cellulose, etc. The further examples regarding to the carriers are the lysis inhibitors, *i.e.* sucrose, tristearyl glycerol, cocoa butter, and hydrogenated oil, etc. The further examples regarding to the carriers are the absorbefacients, *i.e.* quaternary ammonium salt, and sodium dodecyl-sulfonate, etc. The further examples regarding to the carriers are the lubricants, *i.e.* talcum powder, silicon dioxide, corn starch, stearate, boric acid, liquid paraffin, and polyethylene glycol, etc. The tablet is further produced as the coating tablet, *i.e.* sugar coating tablet, film coating tablet, intestine-dissolving coating tablet, bi-layer tablet, and multi-layer tablet. In order to prepare the unit delivery in Chinese medicine pill form, each carrier for one skilled in the art can be widely used. The examples regarding to the carrier are the dilutents and the absorbents, *i.e.* glucose, sucrose, starch, cocoa butter,

hydrogenated vegetable oil, poly vinyl pyrrolidone, Gelucire, kaolin, talcum powder, etc. The further examples regarding to the carrier are the bonding agents, *i.e.* arabic gum, tragacanth gum, gelatin, ethanol, honey, liquid sugar, rice slurry, and batter, etc. The further examples regarding to the carrier are the lysis agents, *i.e.* agar, dried starch, alginate, sodium dodecyl-sulfonate, methyl cellulose, and ethyl cellulose, etc. In order to prepare the unit delivery in suppository form, each carrier for one skilled in the art can be widely used. The examples regarding to the carrier are polyethylene glycol, lecithin, cocoa butter, high alcohol, high alcohol ester, gelatin, semisynthetic glyceride, etc. In order to prepare the unit delivery in capsule form, the pharmaceutical composition or the extract of the present invention are mixed with each carrier described above, and the mixtures obtained from these methods are added into the hard gelatin capsules or the soft capsules. The pharmaceutical composition and the extract of the present invention can be prepared as the microcapsule, and be suspended in aqueous medium to form the suspension. This can be applied to be added into hard capsules.

[0047] Furthermore, if necessary, coloring agents, spices, flavor enhancers, sweeteners, and other materials can be added into the pharmaceutical composition.

[0048] The above objectives and advantages of the present invention will become more readily apparent to those ordinarily skilled in the art after reviewing the following detailed descriptions and accompanying drawings, in which:

BRIEF DESCRIPTION OF THE DRAWINGS

[0049] Fig. 1 is a flowchart diagram showing a preparation method of a pharmaceutical composition in accordance with a first preferred Embodiment of the present invention;

[0050] Fig. 2 is a flowchart diagram showing a preparation method of a pharmaceutical composition in accordance with a second preferred Embodiment of the present invention; and

[0051] Fig. 3 is a flowchart diagram showing a preparation method of a pharmaceutical composition in accordance with a third preferred Embodiment of the present invention.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

[0052] The present invention will now be described more specifically with reference to the following Embodiments. It is to be noted that the following descriptions of preferred Embodiments of this invention are presented herein for purpose of illustration and description only; it is not intended to be exhaustive or to be limited to the precise form disclosed.

[0053] Embodiment 1

[0054] Please refer to Fig. 1, which is a flowchart diagram showing a preparation method of a pharmaceutical composition in accordance with a first preferred Embodiment of the present invention. Fig. 1 adopts the method which one is skilled in the art. Nine (9) kg of the ginseng (101) is decocted in ethanol solution of 75% purity, and then is separated and purified by column chromatography to obtain the first extract (102). The first extract has 40% of ginsenoside. Six (6) kg of the liquorice (103) is decocted in the water solution, and then filtered, concentrated, and dried to obtain the second extract (104). The first extract is mixed with the second extract, and then is pulverized to

obtain the first pharmaceutical composition of the present invention (105, 106, 107, and 108).

[0055] Embodiment 2

[0056] Please refer to Fig. 2, which is a flowchart diagram showing a preparation method of a pharmaceutical composition in accordance with a second preferred Embodiment of the present invention. In Fig. 2, 9 kg of the ginseng (201) is decocted in 60% of ethanol solution, and then separated and purified by column chromatography to obtain the first extract (202). Six (6) kg of the liquorice (203) is decocted in water solution, and then filtered, concentrated, and dried to obtain the second extract (204). Six (6) kg of the jujuba (205) is decocted in 75% of ethanol solution, and then separated and purified by column chromatography to obtain the third extract. The third extract is compounded with 9 parts by weight of the β -cyclodextrin to obtain the extract compound (206). The first extract, the second extract, and the extract compound of the third extract are mixed and pulverized to obtain the second pharmaceutical composition of the present invention (207, 208, 209, and 210).

[0057] Embodiment 3

[0058] Please refer to Fig. 3, which is a flowchart diagram showing a preparation method of a pharmaceutical composition in accordance with a third preferred Embodiment of the present invention. One (1) g of the jujuba extract (having 1% of jujuba cAMP) (302) is compounded with 9 g of β -cyclodextrin to obtain 10 g of extract compound (303). Ten (10) g of extract compound, 50 g of ginseng extract (having 30% of ginsenoside) (301), and 4 g of glycyrrhetic acid (90% purity) (304) are mixed and pulverized to obtain

the third pharmaceutical composition of the present invention (305, 306, 307, and 308).

[0059] Embodiment 4

[0060] Four (4) kg of ginseng and 3 kg of liquorice are pulverized into dry powder. After the obtained dry powder is mixed by the preparation method adopted by one skilled in the art, the pharmaceutical bonding agent, such as honey, etc. is added to prepare the honey pills.

[0061] Embodiment 5

[0062] This pharmaceutical composition is prepared by adopting the same method that of Embodiment 4. The difference is adopting 18 kg of ginseng and 14 kg of liquorice.

[0063] Embodiment 6

[0064] Four (4) kg of commercial ginseng and 3 kg of liquorice are pulverized into dry powder. After the obtained dry powder is mixed by the preparation method adopted by one skilled in the art, 0.2 kg of jujuba ethanol extract and pharmaceutical carriers, *i.e.* starch and dextrin, are added to prepare the pills.

[0065] Embodiment 7

[0066] Eighteen (18) kg of the commercial ginseng water extract and 14 kg of the liquorice are pulverized into dry powder. After the obtained dry powder is mixed by the preparation method adopted by one skilled in the art, the pharmaceutical carriers, *i.e.* starch and dextrin, are added to prepare the pills.

[0067] Embodiment 8

[0068] Four (4) kg of ginseng and 14 kg of liquorice water extract obtained by the method of Embodiment 1 (10) are pulverized into dry powder. After

the obtained dry powder is mixed by the preparation method adopted by one skilled in the art, then 14 kg of 70% of jujuba water extract obtained by the method of the Embodiment 2 (20), and the pharmaceutical carriers, *i.e.* starch and dextrin, are added to prepare the pills.

[0069] Embodiment 9

[0070] After 3 kg of the commercial ginseng ethanol extract having 40% of ginsenoside and 0.2 kg of glycyrrhetic acid are mixed to obtain a mixture, the pharmaceutical pills are prepared by the method adopted by one skilled in the art.

[0071] Embodiment 10

[0072] After 4 kg of the commercial ginseng ethanol extract containing 20% of ginsenoside and 0.8 kg of glycyrrhizic acid are mixed to obtain a mixture, the pharmaceutical soft capsules are prepared by the method adopted by person skilled in the art.

[0073] EXPERIMENT The anti-depression experiment of the present invention

[0074] Experiment 1: "Mouse tail-hanging" experiment

[0075] Experimental animal: ICR mice

[0076] Experimental pharmaceuticals: The pharmaceutical of the Embodiment 3 of the present invention is provided by Beijing Wonner Biotech Ltd. Co., the depression-relieving pill is the product of Zhengzhou Yumi Medicines Co. Ltd., and Paroxetine (Paxil) is the product of Zhong Mei Tianjin Smith Kline pharmaceuticals Co. Ltd.

[0077] Experimental method:

[0078] I. Group division: 1. Large dose of the Embodiment 3 medicine of the present invention (188.5 mg/kg), 2. middle dose of the Embodiment 3

medicine of the present invention (94.25 mg/kg), 3. small dose of the Embodiment 3 medicine of the present invention (47.125 mg/kg), 4. depression-reliving pill (650 mg/kg), 5. Paroxetine (16.7 mg/kg), and 6. physiological saline. (Ten (10) mice are in each group.)

[0079] II. Administration of drug: The abovementioned pharmaceutical water solutions are fed into the stomach according to 0.2 ml/10 g body weight, 2 times per day for a total of 7 days. After 1 hour of the last administration of drug, the mouse tail-hanging experiment is proceeded.

[0080] III. Mouse tail-hanging experiment: The mouse's tail (near to the tail end for 1 cm) is taped on the 5 cm of the wood strip of the high mountain platform and hanged up for 6 minutes. The time of non-movement of the mouse for the last 5 minutes is recorded.

[0081] Experimental result:

[0082] The variance analysis calculation and the p-value compared with the control of the experimental result are calculated by using SPSS 11.5 analytic software.

Group	Animal number	Time of non-movement (s)	p-value
Physiological saline (control)	10	122.66±33.53	
Depression-reliving pill	10	88.21±52.50	0.081
Paroxitine	10	54.98±46.92	0.01
Large dose of the Embodiment 3 medicine of the present invention	10	60.41±36.42	0.02
Middle dose of the Embodiment 3 medicine of the present invention	10	72.68±55.37	0.013
Small dose of the Embodiment 3 medicine of the present invention	10	84.35±27.56	0.053

[0083] Conclusion: According to the above experiment, it is shown that the time of non-movement after the mouse tail-hanging experiment is decreased in all of the large, middle and small doses of the Embodiment 3 medicine of the present invention, and has significant difference compared with the physiological saline (control). Therefore, the Embodiment 3 of the present invention is inferred to have anti-depression function.

[0084] Experiment 2: Body temperature decrease experiment induced by reserpine

[0085] Experimental animal: ICR mice

[0086] Experimental pharmaceuticals: The pharmaceutical of the Embodiment 3 of the present invention is provided by Beijing Wonner Biotech Ltd. Co., the depression-relieving pill is the product of Zhengzhou Yumi Medicines Co. Ltd., and Paroxitine (Paxil) is the product of Zhong Mei Tianjin Smith Kline pharmaceuticals Co. Ltd.

[0087] Experimental method:

[0088] I. Group division: 1. Large dose of the Embodiment 3 medicine of the present invention (188.5 mg/kg), 2. middle dose of the Embodiment 3 medicine of the present invention (94.25 mg/kg), 3. small dose of the Embodiment 3 medicine of the present invention (47.125 mg/kg), 4. depression-reliving pill (650 mg/kg), 5. Paroxetine (16.7 mg/kg), and 6. physiological saline. (Ten (10) mice are in each group.)

[0089] II. Administration of drug: the abovementioned pharmaceutical water solutions are fed into the stomach according to 0.2 ml/10 g body weight, 2 times per day for a total of 7 days.

[0090] III. After the last administration of drug, the anal temperature (abbreviated as anal temp.) is determined, and then 2 mg reserpine per kg of the body weight is taken by intraperitoneal injection. After injecting the reserpine for 2, 3, 4, 5, 6 and 7 hours respectively, the anal temperature of the mice are determined.

[0091] Experimental result:

[0092] The variance analysis calculation and the p-value compared with the control of the experimental result are calculated by using SPSS 11.5 analytic software.

Group	Animal number	Decreased	p-value	Decreased	p-value	Decreased	p-value
		anal temp. for 2 hr (°C)		anal temp. for 3 hr (°C)		anal temp. for 4 hr (°C)	
Physiological saline (control)	10	2.63±0.56		2.33±0.85		2.84±0.84	
Paroxetine	10	1.29±0.47	0.001	1.08±0.35	0.001	1.55±0.64	0.001
Depression-reliving pill	10	2.03±0.55	0.003	2.67±0.48	0.201	2.88±0.65	0.882
Large dose of the Embodiment 3 medicine of the present invention	10	1.82±0.38	0.001	1.77±0.51	0.038	2.92±0.51	0.767
Middle dose of the Embodiment 3 medicine of the present invention	10	0.90±0.44	0.001	0.48±0.36	0.001	0.85±0.21	0.001
Small dose of the Embodiment 3 medicine of the present invention	10	2.63±0.43	0.815	2.04±0.77	0.275	1.45±0.55	0.001

Group	Animal number	Decreased	p-value	Decreased	p-value	Decreased	p-value
		anal temp. for 5 hr (°C)		anal temp. for 6 hr (°C)		anal temp. for 7 hr (°C)	
Physiological saline (control)	10	2.97±0.51		2.60±0.57		3.05±0.67	
Paroxetine	10	1.44±0.32	0.001	2.51±0.47	0.720	2.76±0.59	0.272
Depression-reliving pill	10	2.49±0.60	0.033	2.71±0.46	0.660	3.45±0.65	0.131
Large dose of the Embodiment 3 medicine of the present invention	10	2.88±0.44	0.683	2.43±0.64	0.499	2.30±0.57	0.006
Middle dose of the Embodiment 3 medicine of the present invention	10	1.29±0.56	0.001	1.08±0.59	0.001	1.08±0.39	0.001
Small dose of the Embodiment 3 medicine of the present invention	10	2.28±0.48	0.003	2.68±0.61	0.750	2.29±0.59	0.005

[0093] Conclusion: According to the above results, it is shown that all of the large, middle, and small doses of the Embodiment 3 of the present invention have the function against the decrease of the mice anal temperature

induced by reserpine. The middle dose has significant difference compared with the physiological saline (control). Therefore, the Embodiment 3 of the present invention is inferred to have anti-depression function.

[0094] While the invention has been described in terms of what is presently considered to be the most practical and preferred Embodiments, it is to be understood that the invention needs not be limited to the disclosed Embodiments. On the contrary, it is intended to cover various modifications and similar arrangements included within the spirit and scope of the appended claims, which are to be accorded with the broadest interpretation so as to encompass all such modifications and similar structures.

WHAT IS CLAIMED IS:

1. A pharmaceutical composition for treating a depression, wherein the pharmaceutical composition comprises one of the following compositions of raw materials:

(A) 4 ~ 18 parts by weight of a ginseng and 3 ~ 14 parts by weight of a liquorice;

(B) a ginseng water extract or a ginseng ethanol extract extracted from 4 ~ 18 parts by weight of the ginseng and a liquorice water extract or a liquorice ethanol extract extracted from 3 ~ 14 parts by weight of the liquorice;

(C) 4 ~ 18 parts by weight of the ginseng and the liquorice water extract or the liquorice ethanol extract extracted from 3 ~ 14 parts by weight of the liquorice; and

(D) the ginseng water extract or the ginseng ethanol extract extracted from 4 ~ 18 parts by weight of the ginseng and 3 ~ 14 parts by weight of the liquorice.

2. The pharmaceutical composition according to claim 1, further comprising:

(E) 3 ~ 14 parts by weight of a jujuba; or

(F) a jujuba water extract or a jujuba ethanol extract extracted from 3 ~ 14 parts by weight of the jujuba.

3. The pharmaceutical composition according to claim 1, further comprising:

9 parts by weight of the ginseng and 6 parts by weight of the liquorice; or

the ginseng water extract or the ginseng ethanol extract extracted from 9 parts by weight of the ginseng and the liquorice water extract or the liquorice ethanol extract extracted from 6 parts by weight of the liquorice.

4. The pharmaceutical composition according to claim 2, further comprising:

6 parts by weight of the jujuba; or

the jujuba water extract or the jujuba ethanol extract extracted from 6 parts by weight of the jujuba.

5. A pharmaceutical composition for treating a depression, comprising:

a ginseng water extract or a ginseng ethanol extract extracted from 3 ~ 10 parts by weight of a ginseng; and

0.2 ~ 0.8 part by weight of a glycyrrhizic acid or a glycyrrhetic acid.

6. The pharmaceutical composition according to claim 5, wherein the ginseng ethanol extract comprises 20 ~ 40% of a ginsenoside, and the glycyrrhizic acid or the glycyrrhetic acid has 80 ~ 98% purity.

7. The pharmaceutical composition according to claim 5 or 6, further comprising a jujuba water extract or a jujuba ethanol extract extracted from 0.05 ~ 0.2 part by weight of a jujuba.

8. The pharmaceutical composition according to claim 7, wherein the jujuba ethanol extract comprises 0.5 ~ 3% of a jujuba cyclic adenosine monophosphate (cAMP).

9. The pharmaceutical composition according to claim 8, comprising 5 parts by weight of the ginseng ethanol extract, 0.4 part by weight of the glycyrrhetic acid and 0.1 part by weight of the jujuba ethanol extract, wherein the ginseng ethanol extract comprises 30% of the ginsenosides, the purity of the glycyrrhetic acid is 90%, and the jujuba ethanol extract comprises 1% of the jujuba cAMP.

10. A preparation method of a pharmaceutical composition of claim 1, comprising steps of:

(1) decocting 4 ~ 18 parts by weight of the ginseng in 60 ~ 77% of an ethanol solution, then purifying by a chromatography to obtain a first extract;

(2) decocting 3 ~ 14 parts by weight of the liquorice in a water, then concentrating and drying to obtain a second extract; and

(3) mixing and pulverizing the first extract obtained from the step (1) and the second extract obtained from the step (2), then sifting to obtain the pharmaceutical composition.

11. The preparation method according to claim 10, further comprising the step of:

decocting 3 ~ 14 parts by weight of the jujuba in 60 ~ 75% of the ethanol solution, then purifying by the chromatography to obtain a third extract, and adding the third extract thereof into the pharmaceutical composition.

12. A preparation method of a pharmaceutical composition of claim 5, comprising a step of:

mixing and pulverizing 0.2 ~ 0.8 part by weight of the glycyrrhizic acid or the glycyrrhetic acid having 80 ~ 98% purity and 3 ~ 10 parts by weight of the ginseng extract having 20 ~ 40% of the ginsenoside to obtain the pharmaceutical composition.

13. The preparation method according to claim 12, further comprising the step of:

compounding 0.05 ~ 0.2 part by weight of a jujuba extract having 1% of a jujuba cyclic adenosine monophosphate (cAMP) with a β -cyclodextrin to obtain a jujuba extract compound, and adding the jujuba extract compound thereof into the pharmaceutical composition.