

METHODS, COMPOSITIONS, AND SYSTEMS FOR CAPTURING PROBES AND/OR BARCODES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application claims the benefit of U.S. Provisional Application Serial Nos. 63/128,796, filed December 21, 2020; 63/148,825, filed February 12, 2021; and 63/252,323, filed October 5, 2021, the entire contents of each application are incorporated by reference herein.

BACKGROUND

10 Cells within a tissue have differences in cell morphology and/or function due to varied analyte levels (e.g., gene and/or protein expression) within the different cells. The specific position of a cell within a tissue (e.g., the cell's position relative to neighboring cells or the cell's position relative to the tissue microenvironment) can affect, e.g., the cell's
15 morphology, differentiation, fate, viability, proliferation, behavior, signaling, and cross-talk with other cells in the tissue.

 Spatial heterogeneity has been previously studied using techniques that typically provide data for a handful of analytes in the context of intact tissue or a portion of a tissue (e.g., tissue section), or provide significant analyte data from individual, single cells, but fails
20 to provide information regarding the position of the single cells from the originating biological sample (e.g., tissue).

 Spatial analysis of an analyte within a biological sample may require determining the sequence of the analyte sequence or a complement thereof and the sequence of the spatial barcode or a complement thereof to identify the location of the analyte. The biological
25 sample may be placed on a solid support to improve specificity and efficiency when being analyzed for identification or characterization of an analyte, such as DNA, RNA or other genetic material, within the sample.

SUMMARY

30 The present disclosure features methods, compositions, devices, and systems for determining the location and abundance of an analyte in a biological sample. Determining the spatial location and abundance of analytes (e.g., proteins, DNA, or RNA) within a biological sample leads to better understanding of spatial heterogeneity in various contexts, such as disease models. Described herein are methods for capturing probes and/or barcodes to a

capture domain. In some instances, the techniques disclosed herein facilitate downstream processing, such as sequencing of the probes and/or barcodes bound to a capture domain.

In some examples, the methods, compositions, devices, and systems disclosed herein utilize RNA-templated ligation (RTL) for analyzing an analyte (e.g., RNA) in a biological sample. In some examples, RTL is used in combination with a “sandwich process,” wherein the analyte is transferred from a first substrate to a second substrate for further downstream processing. In some examples, analyte capture agents are used for analyzing an analyte (e.g., protein) in a biological sample. In some examples, the methods disclosed herein allow spatial analysis of two different types of analytes.

10 In some instances, disclosed herein is a method for analyzing an analyte in a biological sample mounted on a first substrate comprising: hybridizing a first probe oligonucleotide and a second probe oligonucleotide to the analyte, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the analyte, and wherein the second
15 probe oligonucleotide comprises a capture probe binding domain; coupling the first probe oligonucleotide and the second probe oligonucleotide, thereby generating a connected probe; aligning the first substrate with a second substrate comprising an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array comprises a plurality of capture probes, wherein a capture probe of the plurality of
20 capture probes comprises: (i) a spatial barcode and (ii) a capture domain; when the biological sample is aligned with at least a portion of the array, (i) releasing the connected probe from the analyte and (ii) migrating the connected probe from the biological sample to the array; and hybridizing the connected probe to the capture domain.

In some instances, the first probe oligonucleotide and the second probe
25 oligonucleotide are on a contiguous nucleic acid sequence. In some instances, the first probe oligonucleotide is on the 3' end of the contiguous nucleic acid sequence. In some instances, the second probe oligonucleotide is on the 5' end of the contiguous nucleic acid sequence. In some instances, the adjacent sequences abut one another. In some instances, the adjacent sequences are at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more nucleotides away from one another.

30 In some instances, the methods further include generating an extended first probe oligonucleotide, wherein the extended first probe oligonucleotide comprises a sequence complementary to a sequence between the sequence hybridized to the first probe oligonucleotide and the sequence hybridized to the second probe oligonucleotide.

In some instances, the methods further include generating an extended second probe oligonucleotide using a polymerase, wherein the extended second probe oligonucleotide comprises a sequence complementary to a sequence between the sequence hybridized to the first probe oligonucleotide and the sequence hybridized to the second probe oligonucleotide.

5 In some instances, the methods further include hybridizing a third probe oligonucleotide to the first probe oligonucleotide and the second probe oligonucleotide.

In some instances, the third probe oligonucleotide comprises: a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, or 100% complementary to a portion of the first probe oligonucleotide that
10 hybridizes to the third probe oligonucleotide; and a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, or 100% complementary to a portion of the second probe oligonucleotide that hybridizes to the third probe oligonucleotide.

In some instances, the coupling the first probe oligonucleotide and the second probe
15 oligonucleotide comprises ligating via a ligase the first probe oligonucleotide and the second probe oligonucleotide. In some instances, the coupling the first probe oligonucleotide and the second probe oligonucleotide comprises ligating via a ligase: (a) the first probe oligonucleotide and the extended second probe oligonucleotide; or (b) the extended first probe oligonucleotide and the second probe oligonucleotide. In some instances, the ligase is
20 selected from a splintR ligase, a single stranded DNA ligase, or a T4 DNA ligase.

In some instances, the methods further include amplifying the connected probe prior to the releasing step, optionally wherein the amplifying comprises rolling circle amplification.

In some instances, the aligning step comprises: (i) mounting the first substrate on a
25 first member of a support device, the first member configured to retain the first substrate; (ii) mounting the second substrate on a second member of the support device; (iii) applying a reagent medium to the first substrate and/or the second substrate, wherein the reagent medium comprises a permeabilization agent and optionally an agent for releasing the connected probe; and (iv) operating an alignment mechanism of the support device to move
30 the first member and/or the second member such that at least a portion of the biological sample is aligned with at least a portion of the array, and such that the portion of the biological sample and the portion of the array contact the reagent medium.

In some instances, the alignment mechanism is coupled to the first member, the second member, or both the first member and the second member. In some instances, the

alignment mechanism comprises a linear actuator, optionally wherein: the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member, and/or the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member, and/or the linear actuator is configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec, and/or the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.

In some instances, during the releasing step (d), a separation distance is maintained between the first substrate and the second substrate, optionally wherein the separation distance is less than 50 microns, optionally wherein the separation distance is between 2-25 microns, optionally wherein the separation distance is measured in a direction orthogonal to the surface of the first substrate that supports the biological sample, and/or at least the portion of the biological sample is vertically aligned with the at least portion of the array.

In some instances, at least one of the first substrate and the second substrate further comprise a spacer, wherein after the first and second substrate being mounted on the support device, the spacer is disposed between the first substrate and second substrate and is configured to maintain the reagent medium within a chamber formed by the first substrate, the second substrate, and the spacer, and maintain a separation distance between the first substrate and the second substrate, the spacer positioned to at least partially surround an area on the first substrate on which the biological sample is disposed and/or the array disposed on the second substrate, wherein the area of the first substrate, the spacer, and the second substrate at least partially encloses a volume comprising the biological sample.

In some instances, the chamber comprises a partially or fully sealed chamber, and/or the second substrate comprises the spacer, and/or the first substrate comprises the spacer, and/or the applying the reagent medium to the first substrate and/or the second substrate comprises applying the reagent medium to a region of the spacer, the region outside an enclosed area of the second substrate, the enclosed area formed by the spacer.

In some instances, as the first substrate and/or the second substrate are moved via the alignment mechanism, the first substrate is at an angle relative to the second substrate such that a dropped side of the first substrate and a portion of the second substrate contact the reagent medium, optionally wherein: the dropped side of the first substrate urges the reagent medium toward the opposite direction, and/or the alignment mechanism further moves the first substrate and/or the second substrate to maintain an approximately parallel arrangement

of the first substrate and the second substrate and a separation distance between the first substrate and the second substrate, optionally when the approximately parallel arrangement and the separation distance are maintained, the spacer fully encloses and surrounds the at least portion of the biological sample and the at least portion of the array, and the spacer
 5 forms the sides of the chamber which hold a volume of the reagent medium.

In some instances, the methods further include contacting the biological sample with a reagent medium comprising a permeabilization agent and an agent for releasing the connected probe, thereby permeabilizing the biological sample and releasing the connected probe from the analyte. In some instances, the agent for releasing the connected probe
 10 comprises a nuclease. In some instances, the nuclease comprises an RNase, optionally wherein the RNase is selected from RNase A, RNase C, RNase H, or RNase I. In some instances, the permeabilization agent comprises a protease. In some instances, the protease is selected from trypsin, pepsin, elastase, or proteinase K. In some instances, the reagent medium further comprises a detergent. In some instances, the detergent is selected from
 15 sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100TM, or Tween-20TM. In some instances, the reagent medium comprises less than 5 w/v% of a detergent selected from SDS and sarkosyl. In some instances, the reagent medium comprises at least 5% w/v% of a detergent selected from SDS and sarkosyl. In some instances, the reagent medium does not comprise sodium dodecyl sulfate (SDS) or sarkosyl.

20 In some instances, the reagent medium further comprises polyethylene glycol (PEG).

In some instances, the biological sample and the array are contacted with the reagent medium for about 1 to about 60 minutes. In some instances, the biological sample and the array are contacted with the reagent medium for about 30 minutes.

In some instances, the methods further include determining (i) all or a part of the
 25 sequence of the connected probe, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof, optionally wherein the method further comprises using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample. In some instances, the determining comprises sequencing (i) all or a part of the sequence of the connected probe, or a
 30 complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof.

In some instances, the sequence of the connected probe comprises the sequence of the spatial barcode or the reverse complement thereof, and a sequence corresponding to the analyte in the biological sample or reverse complement thereof.

In some instances, at least 50% of the connected probe released from the portion of the biological sample aligned with the portion of the array are captured by capture probes of the portion of the array. In some instances, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 5 96%, at least 97%, at least 98%, at least 99%, or 100% of the connected probe are detected in spots directly under the biological sample.

In some instances, the capture probe comprises a poly(T) sequence. In some instances, the capture probe comprises a sequence specific to the analyte. In some instances, the capture probe further comprises one or more functional domains, a unique molecular 10 identifier (UMI), a cleavage domain, and combinations thereof.

In some instances, the analyte is RNA. In some instances, the analyte is mRNA.

In some instances, the methods include analyzing a different analyte in the biological sample. In some instances, the analyzing of the different analyte comprises: contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent 15 of the plurality of analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the different analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence; and hybridizing the capture handle sequence to the capture domain. In some instances, the methods further include determining (i) all or part of 20 the sequence of the capture agent barcode domain; and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some instances, the methods further include using the determined sequence of (i), and (ii) to analyze the different analyte in the biological sample.

In some instances, the releasing step further releases the capture agent barcode 25 domain from the different analyte. In some instances, the different analyte is a protein analyte. In some instances, the protein analyte is an extracellular protein. In some instances, the protein analyte is an intracellular protein. In some instances, the analyte binding moiety is an antibody. In some instances, the analyte capture agent comprises a linker. In some instances, the linker is a cleavable linker. In some instances, the cleavable linker is a photo-cleavable linker, a UV-cleavable linker, or an enzyme cleavable linker. In some instances, the 30 cleavable linker is a photo-cleavable linker.

In some instances, also disclosed herein is a method for analyzing an analyte in a biological sample mounted on a first substrate comprising: contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of

analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence; aligning the first substrate with a second substrate comprising an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises: (i) a spatial barcode and (ii) a capture domain; when the biological sample is aligned with at least a portion of the array, (i) releasing the capture agent barcode domain from the analyte and (ii) passively or actively migrating the capture agent barcode domain from the biological sample to the array; and coupling the capture handle sequence to the capture domain. In some instances, the methods further include determining (i) all or a part of the capture agent barcode domain, or a complement thereof; and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some instances, the methods further include using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample.

In some instances, the aligning step comprises: (i) mounting the first substrate on a first member of a support device, the first member configured to retain the first substrate; (ii) mounting the second substrate on a second member of the support device; (iii) applying a reagent medium to the first substrate and/or the second substrate, wherein the reagent medium comprises a permeabilization agent and optionally an agent for releasing the connected probe; and (iv) operating an alignment mechanism of the support device to move the first member and/or the second member such that at least a portion of the biological sample is aligned with at least a portion of the array, and such that the portion of the biological sample and the portion of the array contact the reagent medium.

In some instances, the alignment mechanism is coupled to the first member, the second member, or both the first member and the second member. In some instances, the alignment mechanism comprises a linear actuator, optionally wherein: the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member, and/or the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member, and/or the linear actuator is configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec, and/or the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.

In some instances, during the releasing step (d), a separation distance is maintained between the first substrate and the second substrate, optionally wherein the separation distance is less than 50 microns, optionally wherein the separation distance is between 2-25 microns, optionally wherein the separation distance is measured in a direction orthogonal to the surface of the first substrate that supports the biological sample, and/or at least the portion of the biological sample is vertically aligned with the at least portion of the array.

In some instances, at least one of the first substrate and the second substrate further comprise a spacer, wherein after the first and second substrate being mounted on the support device, the spacer is disposed between the first substrate and second substrate and is configured to maintain the reagent medium within a chamber formed by the first substrate, the second substrate, and the spacer, and maintain a separation distance between the first substrate and the second substrate, the spacer positioned to at least partially surround an area on the first substrate on which the biological sample is disposed and/or the array disposed on the second substrate, wherein the area of the first substrate, the spacer, and the second substrate at least partially encloses a volume comprising the biological sample.

In some instances, the chamber comprises a partially or fully sealed chamber, and/or the second substrate comprises the spacer, and/or the first substrate comprises the spacer, and/or the applying the reagent medium to the first substrate and/or the second substrate comprises applying the reagent medium to a region of the spacer, the region outside an enclosed area of the second substrate, the enclosed area formed by the spacer.

In some instances, as the first substrate and/or the second substrate are moved via the alignment mechanism, the first substrate is at an angle relative to the second substrate such that a dropped slide of the first substrate and a portion of the second substrate contact the reagent medium, optionally wherein: the dropped side of the first substrate urges the reagent medium toward the opposite direction, and/or the alignment mechanism further moves the first substrate and/or the second substrate to maintain an approximately parallel arrangement of the first substrate and the second substrate and a separation distance between the first substrate and the second substrate, optionally when the approximately parallel arrangement and the separation distance are maintained, the spacer fully encloses and surrounds the at least portion of the biological sample and the at least portion of the array, and the spacer forms the sides of the chamber which hold a volume of the reagent medium.

In some instances, the releasing comprises contacting the biological sample and the array with a reagent medium comprising a nuclease. In some instances, the coupling of the capture handle sequence to the capture domain comprises hybridization. In some instances,

the nuclease comprises an RNase. In some instances, the RNase is selected from RNase A, RNase C, RNase H, and RNase I. In some instances, the reagent medium further comprises a permeabilization agent. In some instances, the releasing further comprises simultaneously permeabilizing the biological sample and releasing the capture agent barcode domain from the analyte binding moiety. In some instances, the permeabilization agent comprises a protease. In some instances, the protease is selected from trypsin, pepsin, elastase, or Proteinase K. In some instances, the reagent medium further comprises a detergent. In some instances, the detergent is selected from sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100™, or Tween-20™. In some instances, the reagent medium comprises less than 5 w/v% of a detergent selected from SDS and sarkosyl. In some instances, the reagent medium comprises as least 5% w/v% of a detergent selected from SDS and sarkosyl. In some instances, the reagent medium does not comprise sodium dodecyl sulfate (SDS) or sarkosyl.

In some instances, the reagent medium further comprises PEG.

In some instances, the biological sample and the array are contacted with the reagent medium for about 1 to about 60 minutes.

In some instances, the biological sample and the array are contacted with the reagent medium for about 30 minutes. In some instances, the methods further include analyzing a different analyte in the biological sample. In some instances, the analyzing of the different analyte comprises: hybridizing a first probe oligonucleotide and a second probe oligonucleotide to the different analyte, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the different analyte, and wherein the second probe oligonucleotide comprises a capture probe binding domain; coupling the first probe oligonucleotide and the second probe oligonucleotide, thereby generating a connected probe comprising the capture probe binding domain; and hybridizing the capture probe binding domain of the connected probe to the capture domain. In some instances, the methods further include determining (i) all or part of the sequence of the connected probe, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some instances, the methods further include using the determined sequence of (i), and (ii) to analyze the different analyte in the biological sample. In some instances, the releasing step further releases the connected probe from the different analyte.

In some instances, the different analyte is RNA. In some instances, the different analyte is mRNA.

In some instances, the capture probe comprises a poly(T) sequence. In some instances, the capture probe comprises a sequence specific to the capture handle sequence. In some instances, the capture probe further comprises one or more functional domains, a unique molecular identifier (UMI), a cleavage domain, and combinations thereof.

5

In some instances, the analyte binding moiety is an antibody. In some instances, the analyte capture agent comprises a linker. In some instances, the linker is a cleavable linker. In some instances, the cleavable linker is a photo-cleavable linker, a UV-cleavable linker, or an enzyme cleavable linker.

10 In some instances, the biological sample is a tissue sample. In some instances, the tissue sample is a solid tissue sample. In some instances, the solid tissue sample is a tissue section. In some instances, the tissue sample is a fixed tissue sample. In some instances, the fixed tissue sample is a formalin fixed paraffin embedded (FFPE) tissue sample. In some instances, the FFPE tissue is deparaffinized and decrosslinked prior to step (a). In some
15 instances, the fixed tissue sample is a formalin fixed paraffin embedded cell pellet. In some instances, the tissue sample is a fresh frozen tissue sample.

In some instances, the tissue sample is fixed and stained prior to step (a).

In some instances, the methods further include analyzing a second analyte in a second biological sample on a third substrate. In some instances, the analyzing the second analyte
20 comprises: hybridizing a third probe oligonucleotide and a fourth probe oligonucleotide to the second analyte, wherein the third probe oligonucleotide and the fourth probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the second analyte, and wherein the fourth probe oligonucleotide comprises a second capture probe binding domain; coupling the third probe oligonucleotide and the fourth probe
25 oligonucleotide, thereby generating a second connected probe; aligning the third substrate with the second substrate comprising the array, such that at least a portion of the second biological sample is aligned with at least a portion of the array; when the second biological sample is aligned with at least a portion of the array, (i) releasing the second connected probe from the second analyte and (ii) migrating the second connected probe from the second
30 biological sample to the array; and hybridizing the second connected probe to the capture domain.

In some instances, the second analyte is RNA. In some instances, the RNA is mRNA.

In some instances, analyzing a second analyte in a second biological sample on a third substrate comprises: contacting the second biological sample with a plurality of second

analyte capture agents, wherein a second analyte capture agent of the plurality of second analyte capture agents comprises a second analyte binding moiety and a second capture agent barcode domain, wherein the second analyte binding moiety specifically binds to the second analyte, and wherein the second capture agent barcode domain comprises a second analyte binding moiety barcode and a second capture handle sequence; aligning the third substrate with a second substrate comprising an array, such that at least a portion of the second biological sample is aligned with at least a portion of the array; when the second biological sample is aligned with at least a portion of the array, (i) releasing the second capture agent barcode domain from the second analyte and (ii) migrating the second capture agent barcode domain from the second biological sample to the array; and coupling the second capture handle sequence to the capture domain.

In some instances, the second analyte is a protein analyte. In some instances, the protein analyte is an extracellular protein. In some instances, the protein analyte is an intracellular protein.

In some instances, the methods further include determining (i) all or a part of the sequence of the connected probe, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof, optionally wherein the method further comprises using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample.

In some instances, the second biological sample is a tissue sample, optionally wherein the tissue sample is a solid tissue sample. In some instances, the tissue sample is a tissue section. In some instances, the tissue sample is a fixed tissue sample. In some instances, the fixed tissue sample is a formalin fixed paraffin embedded (FFPE) tissue sample. In some instances, the FFPE tissue is deparaffinized and decrosslinked. In some instances, the fixed tissue sample is a formalin fixed paraffin embedded cell pellet. In some instances, the tissue sample is a fresh tissue sample or a frozen tissue sample. In some instances, the tissue sample is fixed and stained.

In some instances, the methods disclosed herein further include analyzing: a third analyte in a third biological sample on a fourth substrate; a fourth analyte in a fourth biological sample on a fifth substrate; a fifth analyte or more in a fifth biological sample or more on a sixth substrate or more.

Also disclosed herein is a system or a kit for analyzing an analyte in a biological sample. In some instance, the system or the kit includes (a) a support device configured to retain a first substrate and a second substrate, wherein the biological sample is placed on the

first substrate, and wherein the second substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises: (i) a spatial barcode and (ii) a capture domain; (b) (b1) a first probe oligonucleotide and a second probe oligonucleotide, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the analyte, wherein the second probe oligonucleotide comprises a capture probe binding domain, and wherein the first probe oligonucleotide and the second probe oligonucleotide are capable of being ligated together to form a connected probe; or (b2) a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence; (c) a reagent medium comprising a permeabilization agent and optionally an agent for releasing the connected probe; and (d) instructions for performing any one of the methods disclosed herein. In some instances, the permeabilization agent is pepsin or proteinase K. In some instances, the agent for releasing the connected probe is an RNase, optionally wherein the RNase is RNase A, RNase C, RNase H, or RNase I. In some instances, the system or the kit further comprises an alignment mechanism on the support device to align the first substrate and the second substrate. In some instances, the alignment mechanism comprises a linear actuator, wherein the first substrate comprises a first member and the second substrate comprises a second member, and optionally wherein: the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member, and/or the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member, and/or the linear actuator is configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec, and/or the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.

All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, patent application, or item of information was specifically and individually indicated to be incorporated by reference. To the extent publications, patents, patent applications, and items of information incorporated by reference contradict the disclosure contained in the

specification, the specification is intended to supersede and/or take precedence over any such contradictory material.

Where values are described in terms of ranges, it should be understood that the description includes the disclosure of all possible sub-ranges within such ranges, as well as specific numerical values that fall within such ranges irrespective of whether a specific numerical value or specific sub-range is expressly stated.

The term “each,” when used in reference to a collection of items, is intended to identify an individual item in the collection but does not necessarily refer to every item in the collection, unless expressly stated otherwise, or unless the context of the usage clearly indicates otherwise.

Various embodiments of the features of this disclosure are described herein. However, it should be understood that such embodiments are provided merely by way of example, and numerous variations, changes, and substitutions can occur to those skilled in the art without departing from the scope of this disclosure. It should also be understood that various alternatives to the specific embodiments described herein are also within the scope of this disclosure.

DESCRIPTION OF DRAWINGS

The following drawings illustrate certain embodiments of the features and advantages of this disclosure. These embodiments are not intended to limit the scope of the appended claims in any manner. Like reference symbols in the drawings indicate like elements.

FIG. 1 is a schematic diagram showing an example of a barcoded capture probe, as described herein.

FIG. 2 is a schematic illustrating a cleavable capture probe, wherein the cleaved capture probe can enter into a non-permeabilized cell and bind to target analytes within the sample.

FIG. 3 is a schematic diagram of an exemplary multiplexed spatially-barcoded feature.

FIG. 4 is a schematic diagram of an exemplary analyte capture agent.

FIG. 5 is a schematic diagram depicting an exemplary interaction between a feature-immobilized capture probe **524** and an analyte capture agent **526**.

FIGs. 6A, 6B, and 6C are schematics illustrating how streptavidin cell tags can be utilized in an array-based system to produce a spatially-barcoded cells or cellular contents.

FIG. 7 is a schematic illustrating a side view of a diffusion-resistant medium, e.g., a lid.

FIGs. 8A and **8B** are schematics illustrating expanded **FIG. 8A** and side views **FIG. 8B** of an electrophoretic transfer system configured to direct transcript analytes toward a spatially-barcoded capture probe array.

FIG. 9 is a schematic illustrating an exemplary workflow protocol utilizing an electrophoretic transfer system.

FIG. 10A shows a schematic of an example analytical workflow in which electrophoretic migration of analytes is performed after permeabilization.

FIG. 10B shows a schematic of an example analytical workflow in which electrophoretic migration of analytes and permeabilization are performed simultaneously.

FIG. 11A shows an example perpendicular, single slide configuration for use during electrophoresis.

FIG. 11B shows an example parallel, single slide configuration for use during electrophoresis.

FIG. 11C shows an example multi-slide configuration for use during electrophoresis.

FIG. 12A shows an exemplary workflow for using ligated probes to capture intracellular analytes.

FIG. 12B shows an exemplary schematic illustrating the tissue sample sandwiched between a substrate and a spatially-barcoded capture probe array, wherein the ligated probes are transferred to the spatially-barcoded capture probe array.

FIG. 13 shows exemplary images of spatially-resolved information of gene expression after analyzing the ligated probes in FFPE mouse brain tissue.

FIG. 14 shows exemplary images of spatially-resolved information of gene expression after analyzing the ligated probes in FFPE mouse kidney tissue.

FIG. 15 shows exemplary images of spatially-resolved information of gene expression after analyzing the ligated probes in FFPE mouse brain tissue.

FIG. 16A shows an exemplary workflow for using ligated probes and analyte capture agents to capture intracellular analytes.

FIG. 16B shows an exemplary schematic illustrating the tissue sample sandwiched between a substrate and a spatially-barcoded capture probe array, wherein the ligated probes and capture agent barcode domains are transferred to the spatially-barcoded capture probe array.

FIG. 17A shows an exemplary schematic illustrating the use of an RNase cleavable linker to release a capture agent barcode domain from an analyte binding moiety of an analyte capture agent.

5 **FIG. 17B** shows an exemplary schematic illustrating the use of a UV cleavable linker to release a capture agent barcode domain from an analyte binding moiety of an analyte capture agent.

FIGs. 18A and 18B shows exemplary images of spatially-resolved information of gene expression after analyzing the ligated probes in FFPE mouse brain tissue.

10 **FIG. 19A** shows an exemplary workflow of spatial analysis assays using fresh frozen tissue samples.

FIG. 19B shows an exemplary workflow of spatial analysis assays using formalin-fixation and paraffin-embedded (FFPE) tissue samples.

FIG. 20 shows a table providing data for multiplexed transfer of analytes from multiple biological samples.

15 **FIG. 21A** shows heat maps and clustering images of mouse thymus, mouse testes, and human breast cancer samples using non-sandwiching techniques.

FIG. 21B shows heat maps and clustering images of mouse thymus, mouse testes, and human breast cancer samples using single-sample-sandwiching techniques.

20 **FIGs. 22A and 22B** show heat maps and clustering images mouse and human samples using multiple-sample-sandwiching techniques.

FIG. 23 shows an exemplary schematic diagram depicting a sandwiching process.

FIG. 24 shows a table providing data from an experiment comparing a non-sandwich control and a sandwich configuration permeabilization condition.

25 **FIG. 25** shows an exemplary comparison between a non-sandwich control and a sandwich configuration permeabilization condition.

FIG. 26 shows exemplary images of spatial clustering analysis and analysis of hippocampal transcript Hpca, comparing non-sandwich control and sandwich configuration permeabilization conditions.

30 **FIG. 27A** shows heat maps and clustering images of human brain comparing a non-sandwich control and a sandwich configuration condition.

FIG. 27B shows heat maps and clustering images of human breast comparing a non-sandwich control and a sandwich configuration condition.

FIG. 28A shows an exemplary image of spatially-resolved information of gene expression of protein tyrosine phosphatase receptor type C (Ptprc) RNA using ligated probes in FFPE mouse spleen tissue under sandwich configuration conditions.

5 **FIG. 28B** shows an exemplary image of spatially-resolved information of protein expression of CD45R using analyte capture agents in FFPE mouse spleen tissue under sandwich configuration conditions.

FIG. 28C shows an exemplary image of spatially-resolved information of gene expression of Ptprc RNA using ligated probes in FFPE spleen tissue under non-sandwich configuration control conditions.

10 **FIG. 28D** shows an exemplary image of spatially-resolved information of protein expression of CD45R using analyte capture agents in FFPE mouse spleen tissue under non-sandwich configuration control conditions.

FIG. 29A shows an exemplary image of spatially-resolved information of gene expression of tyrosine hydroxylase (Th) RNA using ligated probes in FFPE mouse brain
15 tissue under sandwich configuration conditions.

FIG. 29B shows an exemplary image of spatially-resolved information of protein expression of tyrosine hydroxylase protein (TH) using analyte capture agents in FFPE mouse brain tissue under sandwich configuration conditions.

FIG. 29C shows an exemplary image of spatially-resolved information of gene
20 expression of Th RNA using ligated probes in FFPE mouse brain tissue under non-sandwich configuration control conditions.

FIG. 29D shows an exemplary image of spatially-resolved information of protein expression of TH using analyte capture agents in FFPE mouse brain tissue under non-sandwich configuration control conditions.

25 **FIG. 30A** shows an exemplary image of spatially-resolved information of gene expression of Ter119 RNA using ligated probes in FFPE mouse embryo torso tissue under sandwich configuration conditions.

FIG. 30B shows an exemplary image of spatially-resolved information of protein expression of Ter119 using analyte capture agents in FFPE mouse embryo torso tissue under
30 sandwich configuration conditions.

FIG. 30C shows an exemplary image of spatially-resolved information of gene expression of Ter119 RNA using ligated probes in FFPE mouse embryo torso tissue under non-sandwich configuration conditions.

FIG. 30D shows an exemplary image of spatially-resolved information of protein expression of Ter119 using analyte capture agents in FFPE mouse embryo torso tissue under non-sandwich configuration conditions.

FIG. 31A shows an exemplary image of spatially-resolved information of gene expression of Ter119 RNA using ligated probes in FFPE mouse embryo head and upper torso tissue under sandwich configuration conditions.

FIG. 31B shows an exemplary image of spatially-resolved information of protein expression of Ter119 using analyte capture agents in FFPE mouse embryo head and upper torso tissue under sandwich configuration conditions.

FIG. 31C shows an exemplary image of spatially-resolved information of gene expression of Ter119 RNA using ligated probes in FFPE mouse embryo head and upper torso tissue under non-sandwich configuration conditions.

FIG. 31D shows an exemplary image of spatially-resolved information of protein expression of Ter119 using analyte capture agents in FFPE mouse embryo head and upper torso tissue under non-sandwich configuration conditions.

FIGs. 32A, 33A, 34A, 35A, and 36A show exemplary images of spatially-resolved information of gene expression of Mpped1 (**FIG. 32A**), Tnncl (**FIG. 33A**), Fgf15 (**FIG. 34A**), Epyc (**FIG. 35A**), and Serpina1e (**FIG. 36A**) RNA using ligated probes in FFPE mouse embryo head and upper torso tissue using sandwich configuration conditions.

FIGs. 32B, 33B, 34B, 35B, and 36B show exemplary images of spatially-resolved information of protein expression of Mpped1 (**FIG. 32B**), Tnncl (**FIG. 33B**), Fgf15 (**FIG. 34B**), Epyc (**FIG. 35B**), and Serpina1e (**FIG. 36B**) using analyte capture agents in FFPE mouse embryo head and upper torso tissue under sandwich configuration conditions.

FIGs. 32C, 33C, 34C, 35C, and 36C show exemplary images of spatially-resolved information of gene expression of Mpped1 (**FIG. 32C**), Tnncl (**FIG. 33C**), Fgf15 (**FIG. 34C**), Epyc (**FIG. 35C**), and Serpina1e (**FIG. 36C**) RNA using ligated probes in FFPE mouse embryo torso tissue under sandwich configuration conditions.

FIGs. 32D, 33D, 34D, 35D, and 36D show exemplary images of spatially-resolved information of gene expression of Mpped1 (**FIG. 32D**), Tnncl (**FIG. 33D**), Fgf15 (**FIG. 34D**), Epyc (**FIG. 35D**), and Serpina1e (**FIG. 36D**) RNA using ligated probes in FFPE mouse embryo head and upper torso tissue under non-sandwich configuration conditions.

FIGs. 32E, 33E, 34E, 35E, and 36E show exemplary images of spatially-resolved information of protein expression of Mpped1 (**FIG. 32E**), Tnncl (**FIG. 33E**), Fgf15 (**FIG.**

34E), Epyc (FIG. 35E), and Serpina1e (FIG. 36E) using analyte capture agents in FFPE mouse embryo head and upper torso tissue under non-sandwich configuration conditions.

FIGs. 32F, 33F, 34F, 35F, and 36F show exemplary images of spatially-resolved information of gene expression of Mpped1 (FIG. 32F), Tnnc1 (FIG. 33F), Fgf15 (FIG. 34F), Epyc (FIG. 35F), and Serpina1e (FIG. 36F) RNA using ligated probes in FFPE mouse embryo torso tissue under non-sandwich configuration conditions.

FIGs. 32G, 33G, 34G, 35G, and 36G show exemplary images of spatially-resolved information of protein expression of Mpped1 (FIG. 32G), Tnnc1 (FIG. 33G), Fgf15 (FIG. 34G), Epyc (FIG. 35G), and Serpina1e (FIG. 36G) using analyte capture agents in FFPE mouse embryo torso tissue under non-sandwich configuration conditions.

FIGs. 32H, 33H, 34H, 35H, and 36H show exemplary images of spatially-resolved information of gene expression of Mpped1 (FIG. 32H), Tnnc1 (FIG. 33H), Fgf15 (FIG. 34H), Epyc (FIG. 35H), and Serpina1e (FIG. 36H) RNA using ligated probes in FFPE mouse embryo head and upper torso tissue under non-sandwich configuration conditions in which only RNA was detected.

FIGs. 32I, 33I, 34I, 35I, and 36I show exemplary images of spatially-resolved information of gene expression of Mpped1 (FIG. 32I), Tnnc1 (FIG. 33I), Fgf15 (FIG. 34I), Epyc (FIG. 35I), and Serpina1e (FIG. 36I) RNA using ligated probes in FFPE mouse embryo torso tissue under non-sandwich configuration conditions in which only RNA was detected.

FIG. 37A shows a perspective view of an example sample handling apparatus in a closed position.

FIG. 37B shows a perspective view of the example sample handling apparatus in an open position.

FIG. 38A shows an exemplary sandwiching process where a first substrate, including a biological sample, and a second substrate are brought into proximity with one another.

FIG. 38B shows a fully formed sandwich configuration creating a chamber formed from the one or more spacers, the first substrate, and the second substrate.

FIG. 39A shows the first substrate angled over (superior to) the second substrate.

FIG. 39B shows that as the first substrate lowers, and/or as the second substrate rises, the dropped side of the first substrate may contact the drop of the reagent medium.

FIG. 39C shows a full closure of the sandwich between the first substrate and the second substrate with the spacer contacting both the first substrate and the second substrate.

FIGs. 40A-40E show an example workflow for an angled sandwich assembly.

FIG. 41A shows a side view of the angled closure workflow.

FIG. 41B shows a top view of the angled closure workflow.

DETAILED DESCRIPTION

I. Introduction

Spatial analysis methodologies, systems, and compositions described herein can provide a vast amount of analyte and/or expression data for a variety of analytes within a biological sample at high spatial resolution, while retaining native spatial context. Spatial analysis methods and compositions can include, e.g., the use of a capture probe including a spatial barcode (e.g., a nucleic acid sequence that provides information as to the location or position of an analyte within a cell or a tissue sample (e.g., mammalian cell or a mammalian tissue sample) and a capture domain that is capable of binding to an analyte (e.g., a protein and/or a nucleic acid) produced by and/or present in a cell. Spatial analysis methods and compositions can also include the use of a capture probe having a capture domain that captures an intermediate agent for indirect detection of an analyte. For example, the intermediate agent can include a nucleic acid sequence (e.g., a barcode) associated with the intermediate agent. Detection of the intermediate agent is therefore indicative of the analyte in the cell or tissue sample.

Non-limiting aspects of spatial analysis methodologies and compositions are described in U.S. Patent Nos. 10,774,374, 10,724,078, 10,480,022, 10,059,990, 10,041,949, 10,002,316, 9,879,313, 9,783,841, 9,727,810, 9,593,365, 8,951,726, 8,604,182, 7,709,198, U.S. Patent Application Publication Nos. 2020/239946, 2020/080136, 2020/0277663, 2020/024641, 2019/330617, 2019/264268, 2020/256867, 2020/224244, 2019/194709, 2019/161796, 2019/085383, 2019/055594, 2018/216161, 2018/051322, 2018/0245142, 2017/241911, 2017/089811, 2017/067096, 2017/029875, 2017/0016053, 2016/108458, 2015/000854, 2013/171621, WO 2018/091676, WO 2020/176788, Rodriques et al., Science 363(6434):1463-1467, 2019; Lee et al., Nat. Protoc. 10(3):442-458, 2015; Trejo et al., PLoS ONE 14(2):e0212031, 2019; Chen et al., Science 348(6233):aaa6090, 2015; Gao et al., BMC Biol. 15:50, 2017; and Gupta et al., Nature Biotechnol. 36:1197-1202, 2018; the Visium Spatial Gene Expression Reagent Kits User Guide (e.g., Rev C, dated June 2020), and/or the Visium Spatial Tissue Optimization Reagent Kits User Guide (e.g., Rev C, dated July 2020), both of which are available at the 10x Genomics Support Documentation website, and can be used herein in any combination. Further non-limiting aspects of spatial analysis methodologies and compositions are described herein.

Some general terminology that may be used in this disclosure can be found in Section (I)(b) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

Typically, a “barcode” is a label, or identifier, that conveys or is capable of conveying information (e.g., information about an analyte in a sample, a bead, and/or a capture probe).

- 5 A barcode can be part of an analyte, or independent of an analyte. A barcode can be attached to an analyte. A particular barcode can be unique relative to other barcodes. For the purpose of this disclosure, an “analyte” can include any biological substance, structure, moiety, or component to be analyzed. The term “target” can similarly refer to an analyte of interest.

- 10 Analytes can be broadly classified into one of two groups: nucleic acid analytes, and non-nucleic acid analytes. Examples of non-nucleic acid analytes include, but are not limited to, lipids, carbohydrates, peptides, proteins, glycoproteins (N-linked or O-linked), lipoproteins, phosphoproteins, specific phosphorylated or acetylated variants of proteins, amidation variants of proteins, hydroxylation variants of proteins, methylation variants of proteins, ubiquitylation variants of proteins, sulfation variants of proteins, viral proteins (e.g.,
15 viral capsid, viral envelope, viral coat, viral accessory, viral glycoproteins, viral spike, etc.), extracellular and intracellular proteins, antibodies, and antigen binding fragments. In some embodiments, the analyte(s) can be localized to subcellular location(s), including, for example, organelles, e.g., mitochondria, Golgi apparatus, endoplasmic reticulum, chloroplasts, endocytic vesicles, exocytic vesicles, vacuoles, lysosomes, etc. In some
20 embodiments, analyte(s) can be peptides or proteins, including without limitation antibodies and enzymes. Additional examples of analytes can be found in Section (I)(c) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. In some embodiments, an analyte can be detected indirectly, such as through detection of an intermediate agent, for example, a connected probe (e.g., a ligation product) or an analyte
25 capture agent (e.g., an oligonucleotide-conjugated antibody), such as those described herein.

- A “biological sample” is typically obtained from the subject for analysis using any of a variety of techniques including, but not limited to, biopsy, surgery, and laser capture microscopy (LCM), and generally includes cells and/or other biological material from the subject. In some embodiments, a biological sample can be a tissue section. In some
30 embodiments, a biological sample can be a fixed and/or stained biological sample (e.g., a fixed and/or stained tissue section). Non-limiting examples of stains include histological stains (e.g., hematoxylin and/or eosin) and immunological stains (e.g., fluorescent stains). In some embodiments, a biological sample (e.g., a fixed and/or stained biological sample) can

be imaged. Biological samples are also described in Section (I)(d) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

In some embodiments, a biological sample is permeabilized with one or more permeabilization reagents. For example, permeabilization of a biological sample can facilitate
5 analyte capture. Exemplary permeabilization agents and conditions are described in Section (I)(d)(ii)(13) or the Exemplary Embodiments Section of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

Array-based spatial analysis methods involve the transfer of one or more analytes from a biological sample to an array of features on a substrate, where each feature is
10 associated with a unique spatial location on the array. Subsequent analysis of the transferred analytes includes determining the identity of the analytes and the spatial location of the analytes within the biological sample. The spatial location of an analyte within the biological sample is determined based on the feature to which the analyte is bound (e.g., directly or indirectly) on the array, and the feature's relative spatial location within the array.

15 A "capture probe" refers to any molecule capable of capturing (directly or indirectly) and/or labelling an analyte (e.g., an analyte of interest) in a biological sample. In some embodiments, the capture probe is a nucleic acid or a polypeptide. In some embodiments, the capture probe includes a barcode (e.g., a spatial barcode and/or a unique molecular identifier (UMI)) and a capture domain). In some embodiments, a capture probe can include a
20 cleavage domain and/or a functional domain (e.g., a primer-binding site, such as for next-generation sequencing (NGS)).

In some instance, the capture domain is designed to detect one or more specific analytes of interest. For example, a capture domain can be designed so that it comprises a sequence that is complementary or substantially complementary to one analyte of interest.
25 Thus, the presence of a single analyte can be detected. Alternatively, the capture domain can be designed so that it comprises a sequence that is complementary or substantially complementary to a conserved region of multiple related analytes. In some instances, the multiple related analytes are analytes that function in the same or similar cellular pathways or that have conserved homology and/or function. The design of the capture probe can be
30 determined based on the intent of the user and can be any sequence that can be used to detect an analyte of interest. In some embodiments, the capture domain sequence can therefore be random, semi-random, defined or combinations thereof, depending on the target analyte(s) of interest.

FIG. 1 is a schematic diagram showing an exemplary capture probe, as described herein. As shown, the capture probe **102** is optionally coupled to a feature **101** by a cleavage domain **103**, such as a disulfide linker. The capture probe can include a functional sequence **104** that are useful for subsequent processing. The functional sequence **104** can include all or a part of sequencer specific flow cell attachment sequence (e.g., a P5 or P7 sequence), all or a part of a sequencing primer sequence, (e.g., a R1 primer binding site, a R2 primer binding site), or combinations thereof. The capture probe can also include a spatial barcode **105**. The capture probe can also include a unique molecular identifier (UMI) sequence **106**. While **FIG. 1** shows the spatial barcode **105** as being located upstream (5') of UMI sequence **106**, it is to be understood that capture probes wherein UMI sequence **106** is located upstream (5') of the spatial barcode **105** is also suitable for use in any of the methods described herein. The capture probe can also include a capture domain **107** to facilitate capture of a target analyte. In some embodiments, the capture probe comprises one or more additional functional sequences that can be located, for example between the spatial barcode **105** and the UMI sequence **106**, between the UMI sequence **106** and the capture domain **107**, or following the capture domain **107**. The capture domain can have a sequence complementary to a sequence of a nucleic acid analyte. The capture domain can have a sequence complementary to a connected probe described herein. The capture domain can have a sequence complementary to a capture handle sequence present in an analyte capture agent. The capture domain can have a sequence complementary to a splint oligonucleotide. Such splint oligonucleotide, in addition to having a sequence complementary to a capture domain of a capture probe, can have a sequence of a nucleic acid analyte, a sequence complementary to a portion of a connected probe described herein, and/or a capture handle sequence described herein.

The functional sequences can generally be selected for compatibility with any of a variety of different sequencing systems, e.g., Ion Torrent Proton or PGM, Illumina sequencing instruments, PacBio, Oxford Nanopore, etc., and the requirements thereof. In some embodiments, functional sequences can be selected for compatibility with non-commercialized sequencing systems. Examples of such sequencing systems and techniques, for which suitable functional sequences can be used, include (but are not limited to) Ion Torrent Proton or PGM sequencing, Illumina sequencing, PacBio SMRT sequencing, and Oxford Nanopore sequencing. Further, in some embodiments, functional sequences can be selected for compatibility with other sequencing systems, including non-commercialized sequencing systems.

In some embodiments, the spatial barcode **105** and functional sequences **104** is common to all of the probes attached to a given feature. In some embodiments, the UMI sequence **106** of a capture probe attached to a given feature is different from the UMI sequence of a different capture probe attached to the given feature.

5 **FIG. 2** is a schematic illustrating a cleavable capture probe, wherein the cleaved capture probe can enter into a non-permeabilized cell and bind to analytes within the sample. The capture probe **201** contains a cleavage domain **202**, a cell penetrating peptide **203**, a reporter molecule **204**, and a disulfide bond (-S-S-). **205** represents all other parts of a capture probe, for example a spatial barcode and a capture domain.

10 **FIG. 3** is a schematic diagram of an exemplary multiplexed spatially-barcoded feature. In **FIG. 3**, the feature **301** can be coupled to spatially-barcoded capture probes, wherein the spatially-barcoded probes of a particular feature can possess the same spatial barcode, but have different capture domains designed to associate the spatial barcode of the feature with more than one target analyte. For example, a feature may be coupled to four
15 different types of spatially-barcoded capture probes, each type of spatially-barcoded capture probe possessing the spatial barcode **302**. One type of capture probe associated with the feature includes the spatial barcode **302** in combination with a poly(T) capture domain **303**, designed to capture mRNA target analytes. A second type of capture probe associated with the feature includes the spatial barcode **302** in combination with a random N-mer capture
20 domain **304** for gDNA analysis. A third type of capture probe associated with the feature includes the spatial barcode **302** in combination with a capture domain complementary to a capture handle sequence of an analyte capture agent of interest **305**. A fourth type of capture probe associated with the feature includes the spatial barcode **302** in combination with a capture domain that can specifically bind a nucleic acid molecule **306** that can function in a
25 CRISPR assay (e.g., CRISPR/Cas9). While only four different capture probe-barcoded constructs are shown in **FIG. 3**, capture-probe barcoded constructs can be tailored for analyses of any given analyte associated with a nucleic acid and capable of binding with such a construct. For example, the schemes shown in **FIG. 3** can also be used for concurrent analysis of other analytes disclosed herein, including, but not limited to: (a) mRNA, a lineage tracing construct, cell surface or intracellular proteins and metabolites, and gDNA; (b)
30 mRNA, accessible chromatin (e.g., ATAC-seq, DNase-seq, and/or MNase-seq) cell surface or intracellular proteins and metabolites, and a perturbation agent (e.g., a CRISPR crRNA/sgRNA, TALEN, zinc finger nuclease, and/or antisense oligonucleotide as described herein); (c) mRNA, cell surface or intracellular proteins and/or metabolites, a barcoded

labelling agent (e.g., the MHC multimers described herein), and a V(D)J sequence of an immune cell receptor (e.g., T-cell receptor). In some embodiments, a perturbation agent can be a small molecule, an antibody, a drug, an aptamer, a miRNA, a physical environmental (e.g., temperature change), or any other known perturbation agents. See, e.g., Section (II)(b) (e.g., subsections (i)-(vi)) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Generation of capture probes can be achieved by any appropriate method, including those described in Section (II)(d)(ii) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

In some embodiments, more than one analyte type (e.g., nucleic acids and proteins) from a biological sample can be detected (e.g., simultaneously or sequentially) using any appropriate multiplexing technique, such as those described in Section (IV) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

In some embodiments, detection of one or more analytes (e.g., protein analytes) can be performed using one or more analyte capture agents. As used herein, an “analyte capture agent” refers to an agent that interacts with an analyte (e.g., an analyte in a biological sample) and with a capture probe (e.g., a capture probe attached to a substrate or a feature) to identify the analyte. In some embodiments, the analyte capture agent includes: (i) an analyte binding moiety (e.g., that binds to an analyte), for example, an antibody or antigen-binding fragment thereof; (ii) analyte binding moiety barcode; and (iii) a capture handle sequence. As used herein, the term “analyte binding moiety barcode” refers to a barcode that is associated with or otherwise identifies the analyte binding moiety. As used herein, the term “analyte capture sequence” or “capture handle sequence” refers to a region or moiety configured to hybridize to, bind to, couple to, or otherwise interact with a capture domain of a capture probe. In some embodiments, a capture handle sequence is complementary to a capture domain of a capture probe. In some cases, an analyte binding moiety barcode (or portion thereof) may be able to be removed (e.g., cleaved) from the analyte capture agent.

FIG. 4 is a schematic diagram of an exemplary analyte capture agent **402** comprised of an analyte-binding moiety **404** and an analyte-binding moiety barcode domain **408**. The exemplary analyte-binding moiety **404** is a molecule capable of binding to an analyte **406** and the analyte capture agent is capable of interacting with a spatially-barcoded capture probe. The analyte-binding moiety can bind to the analyte **406** with high affinity and/or with high specificity. The analyte capture agent can include an analyte-binding moiety barcode domain **408**, a nucleotide sequence (e.g., an oligonucleotide), which can hybridize to at least a portion or an entirety of a capture domain of a capture probe. The analyte-binding moiety

barcode domain 408 can comprise an analyte binding moiety barcode and a capture handle sequence described herein. The analyte-binding moiety **404** can include a polypeptide and/or an aptamer. The analyte-binding moiety **404** can include an antibody or antibody fragment (e.g., an antigen-binding fragment).

5 **FIG. 5** is a schematic diagram depicting an exemplary interaction between a feature-immobilized capture probe **524** and an analyte capture agent **526**. The feature-immobilized capture probe **524** can include a spatial barcode **508** as well as functional sequences **506** and UMI **510**, as described elsewhere herein. The capture probe can also include a capture domain **512** that is capable of binding to an analyte capture agent **526**. The analyte capture agent **526** can include a functional sequence **518**, analyte binding moiety barcode **516**, and a capture handle sequence **514** that is capable of binding to the capture domain **512** of the capture probe **524**. The analyte capture agent can also include a linker **520** that allows the capture agent barcode domain **516** to couple to the analyte binding moiety **522**.

15 **FIGs. 6A, 6B, and 6C** are schematics illustrating how streptavidin cell tags can be utilized in an array-based system to produce a spatially-barcoded cell or cellular contents. For example, as shown in **FIG. 6A**, peptide-bound major histocompatibility complex (MHC) can be individually associated with biotin (β 2m) and bound to a streptavidin moiety such that the streptavidin moiety comprises multiple pMHC moieties. Each of these moieties can bind to a TCR such that the streptavidin binds to a target T-cell via multiple MCH/TCR binding interactions. Multiple interactions synergize and can substantially improve binding affinity. Such improved affinity can improve labelling of T-cells and also reduce the likelihood that labels will dissociate from T-cell surfaces. As shown in **FIG. 6B**, a capture agent barcode domain **601** can be modified with streptavidin **602** and contacted with multiple molecules of biotinylated MHC **603** such that the biotinylated MHC **603** molecules are coupled with the streptavidin conjugated capture agent barcode domain **601**. The result is a barcoded MHC multimer complex **605**. As shown in **FIG. 6B**, the capture agent barcode domain sequence **601** can identify the MHC as its associated label and also includes optional functional sequences such as sequences for hybridization with other oligonucleotides. As shown in **FIG. 6C**, one example oligonucleotide is capture probe **606** that comprises a complementary sequence (e.g., rGrGrG corresponding to C C C), a barcode sequence and other functional sequences, such as, for example, a UMI, an adapter sequence (e.g., comprising a sequencing primer sequence (e.g., R1 or a partial R1 ("pR1"), R2), a flow cell attachment sequence (e.g., P5 or P7 or partial sequences thereof)), etc. In some cases, capture probe **606** may at first be associated with a feature (e.g., a gel bead) and released from the feature. In other

embodiments, capture probe **606** can hybridize with a capture agent barcode domain **601** of the MHC-oligonucleotide complex **605**. The hybridized oligonucleotides (Spacer C C C and Spacer rGrGrG) can then be extended in primer extension reactions such that constructs comprising sequences that correspond to each of the two spatial barcode sequences (the spatial barcode associated with the capture probe, and the barcode associated with the MHC-oligonucleotide complex) are generated. In some cases, one or both of these corresponding sequences may be a complement of the original sequence in capture probe **606** or capture agent barcode domain **601**. In other embodiments, the capture probe and the capture agent barcode domain are ligated together. The resulting constructs can be optionally further processed (e.g., to add any additional sequences and/or for clean-up) and subjected to sequencing. As described elsewhere herein, a sequence derived from the capture probe **606** spatial barcode sequence may be used to identify a feature and the sequence derived from spatial barcode sequence on the capture agent barcode domain **601** may be used to identify the particular peptide MHC complex **604** bound on the surface of the cell (e.g., when using MHC-peptide libraries for screening immune cells or immune cell populations).

Additional description of analyte capture agents can be found in Section (II)(b)(ix) of WO 2020/176788 and/or Section (II)(b)(viii) U.S. Patent Application Publication No. 2020/0277663.

There are at least two methods to associate a spatial barcode with one or more neighboring cells, such that the spatial barcode identifies the one or more cells, and/or contents of the one or more cells, as associated with a particular spatial location. One method is to promote analytes or analyte proxies (e.g., intermediate agents) out of a cell and towards a spatially-barcoded array (e.g., including spatially-barcoded capture probes). Another method is to cleave spatially-barcoded capture probes from an array and promote the spatially-barcoded capture probes towards and/or into or onto the biological sample.

In some cases, capture probes may be configured to prime, replicate, and consequently yield optionally barcoded extension products from a template (e.g., a DNA or RNA template, such as an analyte or an intermediate agent (e.g., a connected probe (e.g., a ligation product) or an analyte capture agent), or a portion thereof), or derivatives thereof (see, e.g., Section (II)(b)(vii) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663 regarding extended capture probes). In some cases, capture probes may be configured to form a connected probe (e.g., a ligation product) with a template (e.g., a DNA or RNA template, such as an analyte or an intermediate agent, or portion thereof), thereby creating ligation products that serve as proxies for a template.

As used herein, an “extended capture probe” refers to a capture probe having additional nucleotides added to the terminus (e.g., 3’ or 5’ end) of the capture probe thereby extending the overall length of the capture probe. For example, an “extended 3’ end” indicates additional nucleotides were added to the most 3’ nucleotide of the capture probe to extend the length of the capture probe, for example, by polymerization reactions used to extend nucleic acid molecules including templated polymerization catalyzed by a polymerase (e.g., a DNA polymerase or a reverse transcriptase). In some embodiments, extending the capture probe includes adding to a 3’ end of a capture probe a nucleic acid sequence that is complementary to a nucleic acid sequence of an analyte or intermediate agent specifically bound to the capture domain of the capture probe. In some embodiments, the capture probe is extended using reverse transcription. In some embodiments, the capture probe is extended using one or more DNA polymerases. The extended capture probes include the sequence of the capture probe and the sequence of the spatial barcode of the capture probe.

In some embodiments, extended capture probes are amplified (e.g., in bulk solution or on the array) to yield quantities that are sufficient for downstream analysis, e.g., via DNA sequencing. In some embodiments, extended capture probes (e.g., DNA molecules) act as templates for an amplification reaction (e.g., a polymerase chain reaction).

Additional variants of spatial analysis methods, including in some embodiments, an imaging step, are described in Section (II)(a) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Analysis of captured analytes (and/or intermediate agents or portions thereof), for example, including sample removal, extension of capture probes, sequencing (e.g., of a cleaved extended capture probe and/or a cDNA molecule complementary to an extended capture probe), sequencing on the array (e.g., using, for example, in situ hybridization or in situ ligation approaches), temporal analysis, and/or proximity capture, is described in Section (II)(g) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Some quality control measures are described in Section (II)(h) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

Spatial information can provide information of biological and/or medical importance. For example, the methods and compositions described herein can allow for: identification of one or more biomarkers (e.g., diagnostic, prognostic, and/or for determination of efficacy of a treatment) of a disease or disorder; identification of a candidate drug target for treatment of a disease or disorder; identification (e.g., diagnosis) of a subject as having a disease or disorder; identification of stage and/or prognosis of a disease or disorder in a subject;

identification of a subject as having an increased likelihood of developing a disease or disorder; monitoring of progression of a disease or disorder in a subject; determination of efficacy of a treatment of a disease or disorder in a subject; identification of a patient subpopulation for which a treatment is effective for a disease or disorder; modification of
5 treatment of a subject with a disease or disorder; selection of a subject for participation in a clinical trial; and/or selection of a treatment for a subject with a disease or disorder.

Spatial information can provide information of biological importance. For example, the methods and compositions described herein can allow for: identification of transcriptome and/or proteome expression profiles (e.g., in healthy and/or diseased tissue); identification of
10 multiple analyte types in close proximity (e.g., nearest neighbor analysis); determination of up- and/or down-regulated genes and/or proteins in diseased tissue; characterization of tumor microenvironments; characterization of tumor immune responses; characterization of cells types and their co-localization in tissue; and identification of genetic variants within tissues (e.g., based on gene and/or protein expression profiles associated with specific disease or
15 disorder biomarkers).

Typically, for spatial array-based methods, a substrate functions as a support for direct or indirect attachment of capture probes to features of the array. A “feature” is an entity that acts as a support or repository for various molecular entities used in spatial analysis. In some embodiments, some or all of the features in an array are functionalized for analyte capture.
20 Exemplary substrates are described in Section (II)(c) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. Exemplary features and geometric attributes of an array can be found in Sections (II)(d)(i), (II)(d)(iii), and (II)(d)(iv) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

Generally, analytes and/or intermediate agents (or portions thereof) can be captured
25 when contacting a biological sample with a substrate including capture probes (e.g., a substrate with capture probes embedded, spotted, printed, fabricated on the substrate, or a substrate with features (e.g., beads, wells) comprising capture probes). As used herein, “contact,” “contacted,” and/or “contacting,” a biological sample with a substrate refers to any contact (e.g., direct or indirect) such that capture probes can interact (e.g., bind covalently or
30 non-covalently (e.g., hybridize)) with analytes from the biological sample. Capture can be achieved actively (e.g., using electrophoresis) or passively (e.g., using diffusion). Analyte capture is further described in Section (II)(e) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

In some cases, spatial analysis can be performed by attaching and/or introducing a molecule (e.g., a peptide, a lipid, or a nucleic acid molecule) having a barcode (e.g., a spatial barcode) to a biological sample (e.g., to a cell in a biological sample). In some embodiments, a plurality of molecules (e.g., a plurality of nucleic acid molecules) having a plurality of barcodes (e.g., a plurality of spatial barcodes) are introduced to a biological sample (e.g., to a plurality of cells in a biological sample) for use in spatial analysis. In some embodiments, after attaching and/or introducing a molecule having a barcode to a biological sample, the biological sample can be physically separated (e.g., dissociated) into single cells or cell groups for analysis. Some such methods of spatial analysis are described in Section (III) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663.

In some cases, spatial analysis can be performed by detecting multiple oligonucleotides that hybridize to an analyte. In some instances, for example, spatial analysis can be performed using RNA-templated ligation (RTL). Methods of RTL have been described previously. See, e.g., Credle *et al.*, *Nucleic Acids Res.* 2017 Aug 21;45(14):e128. Typically, RTL includes hybridization of two oligonucleotides to adjacent sequences on an analyte (e.g., an RNA molecule, such as an mRNA molecule). In some instances, the oligonucleotides are DNA molecules. In some instances, one of the oligonucleotides includes at least two ribonucleic acid bases at the 3' end and/or the other oligonucleotide includes a phosphorylated nucleotide at the 5' end. In some instances, one of the two oligonucleotides includes a capture domain (e.g., a poly(A) sequence, a non-homopolymeric sequence). After hybridization to the analyte, a ligase (e.g., SplintR ligase) ligates the two oligonucleotides together, creating a connected probe (e.g., a ligation product). In some instances, the two oligonucleotides hybridize to sequences that are not adjacent to one another. For example, hybridization of the two oligonucleotides creates a gap between the hybridized oligonucleotides. In some instances, a polymerase (e.g., a DNA polymerase) can extend one of the oligonucleotides prior to ligation. After ligation, the connected probe (e.g., a ligation product) is released from the analyte. In some instances, the connected probe (e.g., a ligation product) is released using an endonuclease (e.g., RNase A, RNase C, RNase H, or RNase I). The released connected probe (e.g., a ligation product) can then be captured by capture probes (e.g., instead of direct capture of an analyte) on an array, optionally amplified, and sequenced, thus determining the location and optionally the abundance of the analyte in the biological sample.

During analysis of spatial information, sequence information for a spatial barcode associated with an analyte is obtained, and the sequence information can be used to provide

information about the spatial distribution of the analyte in the biological sample. Various methods can be used to obtain the spatial information. In some embodiments, specific capture probes and the analytes they capture are associated with specific locations in an array of features on a substrate. For example, specific spatial barcodes can be associated with specific array locations prior to array fabrication, and the sequences of the spatial barcodes can be stored (e.g., in a database) along with specific array location information, so that each spatial barcode uniquely maps to a particular array location.

Alternatively, specific spatial barcodes can be deposited at predetermined locations in an array of features during fabrication such that at each location, only one type of spatial barcode is present so that spatial barcodes are uniquely associated with a single feature of the array. Where necessary, the arrays can be decoded using any of the methods described herein so that spatial barcodes are uniquely associated with array feature locations, and this mapping can be stored as described above.

When sequence information is obtained for capture probes and/or analytes during analysis of spatial information, the locations of the capture probes and/or analytes can be determined by referring to the stored information that uniquely associates each spatial barcode with an array feature location. In this manner, specific capture probes and captured analytes are associated with specific locations in the array of features. Each array feature location represents a position relative to a coordinate reference point (e.g., an array location, a fiducial marker) for the array. Accordingly, each feature location has an “address” or location in the coordinate space of the array.

Some exemplary spatial analysis workflows are described in the Exemplary Embodiments section of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. See, for example, the Exemplary embodiment starting with “In some non-limiting examples of the workflows described herein, the sample can be immersed...” of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663. See also, e.g., the Visium Spatial Gene Expression Reagent Kits User Guide (e.g., Rev C, dated June 2020), and/or the Visium Spatial Tissue Optimization Reagent Kits User Guide (e.g., Rev C, dated July 2020).

In some embodiments, spatial analysis can be performed using dedicated hardware and/or software, such as any of the systems described in Sections (II)(e)(ii) and/or (V) of WO 2020/176788 and/or U.S. Patent Application Publication No. 2020/0277663, or any of one or more of the devices or methods described in Sections *Control Slide for Imaging*, *Methods of Using Control Slides and Substrates for*, *Systems of Using Control Slides and Substrates for*

Imaging, and/or Sample and Array Alignment Devices and Methods, Informational labels of WO 2020/123320.

Suitable systems for performing spatial analysis can include components such as a chamber (e.g., a flow cell or sealable, fluid-tight chamber) for containing a biological sample.

- 5 The biological sample can be mounted for example, in a biological sample holder. One or more fluid chambers can be connected to the chamber and/or the sample holder via fluid conduits, and fluids can be delivered into the chamber and/or sample holder via fluidic pumps, vacuum sources, or other devices coupled to the fluid conduits that create a pressure gradient to drive fluid flow. One or more valves can also be connected to fluid conduits to
10 regulate the flow of reagents from reservoirs to the chamber and/or sample holder.

The systems can optionally include a control unit that includes one or more electronic processors, an input interface, an output interface (such as a display), and a storage unit (e.g., a solid state storage medium such as, but not limited to, a magnetic, optical, or other solid state, persistent, writeable and/or re-writeable storage medium). The control unit can

- 15 optionally be connected to one or more remote devices via a network. The control unit (and components thereof) can generally perform any of the steps and functions described herein. Where the system is connected to a remote device, the remote device (or devices) can perform any of the steps or features described herein. The systems can optionally include one or more detectors (e.g., CCD, CMOS) used to capture images. The systems can also
20 optionally include one or more light sources (e.g., LED-based, diode-based, lasers) for illuminating a sample, a substrate with features, analytes from a biological sample captured on a substrate, and various control and calibration media.

The systems can optionally include software instructions encoded and/or implemented in one or more of tangible storage media and hardware components such as application

- 25 specific integrated circuits. The software instructions, when executed by a control unit (and in particular, an electronic processor) or an integrated circuit, can cause the control unit, integrated circuit, or other component executing the software instructions to perform any of the method steps or functions described herein.

In some cases, the systems described herein can detect (e.g., register an image) the
30 biological sample on the array. Exemplary methods to detect the biological sample on an array are described in PCT Application No. 2020/061064 and/or U.S. Patent Application Serial No. 16/951,854.

Prior to transferring analytes from the biological sample to the array of features on the substrate, the biological sample can be aligned with the array. Alignment of a biological

sample and an array of features including capture probes can facilitate spatial analysis, which can be used to detect differences in analyte presence and/or level within different positions in the biological sample, for example, to generate a three-dimensional map of the analyte presence and/or level. Exemplary methods to generate a two- and/or three-dimensional map of the analyte presence and/or level are described in PCT Application No. 2020/053655 and spatial analysis methods are generally described in WO 2020/061108 and/or U.S. Patent Application Serial No. 16/951,864.

In some cases, a map of analyte presence and/or level can be aligned to an image of a biological sample using one or more fiducial markers, e.g., objects placed in the field of view of an imaging system which appear in the image produced, as described in the *Substrate Attributes* Section, *Control Slide for Imaging* Section of WO 2020/123320, PCT Application No. 2020/061066, and/or U.S. Patent Application Serial No. 16/951,843. Fiducial markers can be used as a point of reference or measurement scale for alignment (e.g., to align a sample and an array, to align two substrates, to determine a location of a sample or array on a substrate relative to a fiducial marker) and/or for quantitative measurements of sizes and/or distances.

The sandwich process is described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

II. Methods, Compositions, Devices, and Systems for Capturing Analytes and Derivatives Thereof

(a) Introduction

Provided herein are methods, devices, compositions, and systems for analyzing the location and abundance of a nucleic acid or protein analyte in a biological sample. In some instances, the methods include aligning (i.e., sandwiching) a first substrate having the biological sample with a second substrate that includes a plurality of capture probes, thereby “sandwiching” the biological sample between the two substrates. Upon interaction of the biological sample with the substrate having a plurality of probes (in either instance), the location and abundance of a nucleic acid or protein analyte in a biological sample can be determined, as provided herein. This methods includes an advantage in that steps provided herein prior to analyte or analyte-derived molecule by the capture probe, most—if not all—steps can be performed on a substrate that does not have capture probes, thereby providing a method that is cost effective.

The methods and systems provided herein can be applied to an analyte or an analyte-derived molecule(s). As used herein, an analyte derived molecule includes, without limitation, a connected probe (e.g., a ligation product) from an RNA-templated ligation (RTL) assay, a product of reverse transcription (e.g., an extended capture probe), and an analyte binding moiety barcode (e.g., a binding moiety barcode that identifies that analyte binding moiety (e.g., an antibody)). In some embodiments, the analyte or analyte derived molecules comprise RNA and/or DNA. In some embodiments, the analyte or analyte derived molecules comprise one or more proteins.

In some instances, the methods, devices, compositions, and systems disclosed herein provide efficient release of an analyte or analyte derived molecule from a biological sample so that it can be easily captured or detected using methods disclosed herein.

In some instances, the methods, devices, compositions, and systems disclosed herein allow for detection of analytes or analyte derived molecules from different biological samples using a single array comprising a plurality of capture probes. As such, in some instances, the methods, devices, compositions, and systems allow for serial capture of analytes or analyte derived molecules from multiple samples. The analytes or analyte derived molecules can then be demultiplexed using biological-sample-specific index sequences to identify its biological sample origin.

Embodiments of the methods, devices, compositions, and systems disclosed herein are provided below.

(A) Exemplary Biological Samples

The biological sample as used herein can be any suitable biological sample described herein or known in the art. In some embodiments, the biological sample is a tissue. In some embodiments, the tissue sample is a solid tissue sample. In some embodiments, the biological sample is a tissue section. In some embodiments, the tissue is flash-frozen and sectioned. Any suitable methods described herein or known in the art can be used to flash-freeze and section the tissue sample. In some embodiments, the biological sample, e.g., the tissue, is flash-frozen using liquid nitrogen before sectioning. In some embodiments, the sectioning is performed using cryosectioning. In some embodiments, the methods further comprises a thawing step, after the cryosectioning. In some embodiments, the biological sample, e.g., the tissue sample is fixed, for example in methanol, acetone, PFA or is formalin-fixed and paraffin-embedded (FFPE). In some embodiments, the biological sample comprises intact cells. In some embodiments, the biological sample is a cell pellet, e.g., a fixed cell pellet, e.g., an FFPE cell pellet. FFPE samples are used in some instances in the RTL methods

disclosed herein. A limitation of direct RNA capture for fixed samples is that the RNA integrity of fixed (e.g., FFPE) samples can be lower than a fresh sample, thereby making it more difficult to capture RNA directly, e.g., by capture of a common sequence such as a poly(A) tail of an mRNA molecule. However, by utilizing RTL probe oligonucleotides that hybridize to RNA target sequences in the transcriptome, one can avoid a requirement for RNA analytes to have both a poly(A) tail and target sequences intact. Accordingly, RTL probes can be utilized to beneficially improve capture and spatial analysis of fixed samples.

The biological sample, e.g., tissue sample, can be stained, and imaged prior, during, and/or after each step of the methods described herein. Any of the methods described herein or known in the art can be used to stain and/or image the biological sample. In some embodiments, the imaging occurs prior to deaminating the sample. In some embodiments, the biological sample is stained using an H&E staining method. In some embodiments, the tissue sample is stained and imaged for about 10 minutes to about 2 hours (or any of the subranges of this range described herein). Additional time may be needed for staining and imaging of different types of biological samples.

The tissue sample can be obtained from any suitable location in a tissue or organ of a subject, e.g., a human subject. In some instances, the sample is a human sample. In some instances, the sample is a human breast tissue sample. In some instances, the sample is a human brain tissue sample. In some embodiments, the sample is an embryo sample. The embryo sample can be a non-human embryo sample. In some instances, the sample is a mouse embryo sample.

(B) Exemplary First and Second Substrates

In some instances, the biological sample is placed (e.g., mounted or otherwise immobilized) on a first substrate. The first substrate can be any solid or semi-solid support upon which a biological sample can be mounted. In some instances, the first substrate is a slide. In some instances, the slide is a glass slide. In some embodiments, the substrate is made of glass, silicon, paper, hydrogel, polymer monoliths, or other material known in the art. In some embodiments, the first substrate is comprised of an inert material or matrix (e.g., glass slides) that has been functionalized by, for example, treating the substrate with a material comprising reactive groups which facilitate mounting of the biological sample.

In some embodiments, the first substrate does not comprise a plurality (e.g., array) of capture probes, each comprising a spatial barcode.

A substrate, e.g., a first substrate and/or a second substrate, can generally have any suitable form or format. For example, a substrate can be flat, curved, e.g., convexly or concavely curved. For example, a first substrate can be curved towards the area where the interaction between a biological sample, e.g., tissue sample, and a first substrate takes place.

- 5 In some embodiments, a substrate is flat, e.g., planar, chip, or slide. A substrate can contain one or more patterned surfaces within the first substrate (e.g., channels, wells, projections, ridges, divots, etc.).

- A substrate, e.g., a first substrate and/or second substrate, can be of any desired shape. For example, a substrate can be typically a thin, flat shape (e.g., a square or a rectangle). In
10 some embodiments, a substrate structure has rounded corners (e.g., for increased safety or robustness). In some embodiments, a substrate structure has one or more cut-off corners (e.g., for use with a slide clamp or cross-table). In some embodiments wherein a substrate structure is flat, the substrate structure can be any appropriate type of support having a flat surface (e.g., a chip or a slide such as a microscope slide).

- 15 First and/or second substrates can optionally include various structures such as, but not limited to, projections, ridges, and channels. A substrate can be micropatterned to limit lateral diffusion of analytes (e.g., to improve resolution of the spatial analysis). A substrate modified with such structures can be modified to allow association of analytes, features (e.g., beads), or probes at individual sites. For example, the sites where a substrate is modified with
20 various structures can be contiguous or non-contiguous with other sites.

- In some embodiments, the surface of a first and/or second substrate is modified to contain one or more wells, using techniques such as (but not limited to) stamping, microetching, or molding techniques. In some embodiments in which a first and/or second substrate includes one or more wells, the first substrate can be a concavity slide or cavity
25 slide. For example, wells can be formed by one or more shallow depressions on the surface of the first and/or second substrate. In some embodiments, where a first and/or second substrate includes one or more wells, the wells can be formed by attaching a cassette (e.g., a cassette containing one or more chambers) to a surface of the first substrate structure.

- In some embodiments where the first and/or second substrate is modified to contain
30 one or more structures, including but not limited to, wells, projections, ridges, features, or markings, the structures can include physically altered sites. For example, a first and/or second substrate modified with various structures can include physical properties, including, but not limited to, physical configurations, magnetic or compressive forces, chemically functionalized sites, chemically altered sites, and/or electrostatically altered sites. In some

embodiments where the first substrate is modified to contain various structures, including but not limited to wells, projections, ridges, features, or markings, the structures are applied in a pattern. Alternatively, the structures can be randomly distributed.

In some embodiments, a first substrate includes one or more markings on its surface, e.g., to provide guidance for aligning at least a portion of the biological sample with a plurality of capture probes on the second substrate during a sandwich process disclosed herein. For example, the first substrate can include a sample area indicator identifying the sample area. In some embodiments, during a sandwiching process described herein the sample area indicator on the first substrate is aligned with an area of the second substrate comprising a plurality of capture probes. In some embodiments, the first and/or second substrate can include a fiducial mark. In some embodiments, the first and/or second substrate does not comprise a fiducial mark. In some embodiments, the first substrate does not comprise a fiducial mark and the second substrate comprises a fiducial mark. Such markings can be made using techniques including, but not limited to, printing, sand-blasting, and depositing on the surface.

In some embodiments, imaging can be performed using one or more fiducial markers, i.e., objects placed in the field of view of an imaging system which appear in the image produced. Fiducial markers are typically used as a point of reference or measurement scale. Fiducial markers can include, but are not limited to, detectable labels such as fluorescent, radioactive, chemiluminescent, and colorimetric labels. The use of fiducial markers to stabilize and orient biological samples is described, for example, in Carter et al., *Applied Optics* 46:421-427, 2007), the entire contents of which are incorporated herein by reference. In some embodiments, a fiducial marker can be a physical particle (e.g., a nanoparticle, a microsphere, a nanosphere, a bead, a post, or any of the other exemplary physical particles described herein or known in the art).

In some embodiments, a fiducial marker can be present on a first substrate to provide orientation of the biological sample. In some embodiments, a microsphere can be coupled to a first substrate to aid in orientation of the biological sample. In some examples, a microsphere coupled to a first substrate can produce an optical signal (e.g., fluorescence). In some embodiments, a quantum dot can be coupled to the first substrate to aid in the orientation of the biological sample. In some examples, a quantum dot coupled to a first substrate can produce an optical signal.

In some embodiments, a fiducial marker can be an immobilized molecule with which a detectable signal molecule can interact to generate a signal. For example, a marker nucleic

acid can be linked or coupled to a chemical moiety capable of fluorescing when subjected to light of a specific wavelength (or range of wavelengths). Although not required, it can be advantageous to use a marker that can be detected using the same conditions (e.g., imaging conditions) used to detect a labelled cDNA.

5 In some embodiments, a fiducial marker can be randomly placed in the field of view. For example, an oligonucleotide containing a fluorophore can be randomly printed, stamped, synthesized, or attached to a first substrate (e.g., a glass slide) at a random position on the first substrate. A tissue section can be contacted with the first substrate such that the oligonucleotide containing the fluorophore contacts, or is in proximity to, a cell from the
10 tissue section or a component of the cell (e.g., an mRNA or DNA molecule). An image of the first substrate and the tissue section can be obtained, and the position of the fluorophore within the tissue section image can be determined (e.g., by reviewing an optical image of the tissue section overlaid with the fluorophore detection). In some embodiments, fiducial markers can be precisely placed in the field of view (e.g., at known locations on a first
15 substrate). In this instance, a fiducial marker can be stamped, attached, or synthesized on the first substrate and contacted with a biological sample. Typically, an image of the sample and the fiducial marker is taken, and the position of the fiducial marker on the first substrate can be confirmed by viewing the image.

 In some embodiments, a fiducial marker can be an immobilized molecule (e.g., a
20 physical particle) attached to the first substrate. For example, a fiducial marker can be a nanoparticle, e.g., a nanorod, a nanowire, a nanocube, a nanopyramid, or a spherical nanoparticle. In some examples, the nanoparticle can be made of a heavy metal (e.g., gold). In some embodiments, the nanoparticle can be made from diamond. In some embodiments, the fiducial marker can be visible by eye.

25 A wide variety of different first substrates can be used for the foregoing purposes. In general, a first substrate can be any suitable support material. Exemplary first substrates include, but are not limited to, glass, modified and/or functionalized glass, hydrogels, films, membranes, plastics (including e.g., acrylics, polystyrene, copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, TeflonTM, cyclic
30 olefins, polyimides etc.), nylon, ceramics, resins, Zeonor, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses, optical fiber bundles, and polymers, such as polystyrene, cyclic olefin copolymers (COCs), cyclic olefin polymers (COPs), polypropylene, polyethylene polycarbonate, or combinations thereof.

Among the examples of first substrate materials discussed above, polystyrene is a hydrophobic material suitable for binding negatively charged macromolecules because it normally contains few hydrophilic groups. For nucleic acids immobilized on glass slides, by increasing the hydrophobicity of the glass surface the nucleic acid immobilization can be increased. Such an enhancement can permit a relatively more densely packed formation (e.g., provide improved specificity and resolution).

In another example, a first substrate can be a flow cell. Flow cells can be formed of any of the foregoing materials, and can include channels that permit reagents, solvents, features, and analytes to pass through the flow cell. In some embodiments, a hydrogel embedded biological sample is assembled in a flow cell (e.g., the flow cell is utilized to introduce the hydrogel to the biological sample). In some embodiments, a hydrogel embedded biological sample is not assembled in a flow cell. In some embodiments, the hydrogel embedded biological sample can then be prepared and/or isometrically expanded as described herein.

Exemplary substrates similar to the first substrate (e.g., a substrate having no capture probes) and/or the second substrate are described in Section (I) above and in WO 2020/123320, which is hereby incorporated by reference in its entirety.

(b) Capturing Nucleic Acid Analytes using RNA-Templated Ligation

In some embodiments, the methods, devices, compositions, and systems described herein utilize RNA-templated ligation to detect the analyte. As used herein, spatial “RNA-templated ligation,” or “RTL” or simply “templated ligation” is a process wherein individual probe oligonucleotides (e.g., a first probe oligonucleotide, a second probe oligonucleotide) in a probe pair hybridize to adjacent sequences of an analyte (e.g., an RNA molecule) in a biological sample (e.g., a tissue sample). The RTL probe oligonucleotides are then coupled (e.g., ligated) together, thereby creating a connected probe (e.g., a ligation product). RNA-templated ligation is disclosed in PCT Publ. No. WO 2021/133849 A1 and US Publ. No. US 2021/0285046 A1, each of which is incorporated by reference in its entirety.

An advantage to using RTL is that it allows for enhanced detection of analytes (e.g., low expressing analytes) because both probe oligonucleotides must hybridize to the analyte in order for the coupling (e.g., ligating) reaction to occur. As used herein, “coupling” refers to an interaction between two probe oligonucleotides that results in a single connected probe that comprises the two probe oligonucleotides. In some instances, coupling is achieved through ligation. In some instances, coupling is achieved through extension of one probe

oligonucleotide to the second probe oligonucleotide followed by ligation. In some instances, coupling is achieved through hybridization (e.g., using a third probe oligonucleotide that hybridized to each of the two probe oligonucleotides) followed by extension of one probe oligonucleotide or gap filling of the sequence between the two probe oligonucleotides using the third probe oligonucleotide as a template.

The connected probe (e.g., ligation product) that results from the coupling (e.g., ligation) of the two probe oligonucleotides can serve as a proxy for the target analyte, as such an analyte derived molecule. Further, it is appreciated that probe oligonucleotide pairs can be designed to cover any gene of interest. For example, a pair of probe oligonucleotides can be designed so that each analyte, e.g., a whole exome, a transcriptome, a genome, can conceivably be detected using a probe oligonucleotide pair.

In some instances, disclosed herein are methods for analyzing an analyte in a biological sample comprising (a) hybridizing a first probe oligonucleotide and a second probe oligonucleotide to the analyte, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the analyte, and wherein the second probe oligonucleotide comprises a capture probe binding domain; (b) coupling the first probe oligonucleotide and the second probe oligonucleotide, thereby generating a connected probe (e.g., a ligation product) comprising the capture probe binding domain; (c) contacting the biological sample with a reagent medium comprising a permeabilization agent and an agent for releasing the connected probe (e.g., a ligation product), thereby (i) permeabilizing the biological sample and (ii) releasing the connected probe (e.g., a ligation product) from the analyte; and (d) hybridizing the capture probe binding domain of the connected probe (e.g., a ligation product) to a capture domain of a capture probe, wherein the capture probe comprises: (i) a spatial barcode and (ii) a capture domain.

Also provided herein are methods for analyzing an analyte in a biological sample mounted on a first substrate including (a) hybridizing a first probe oligonucleotide and a second probe oligonucleotide to the analyte, wherein the first probe oligonucleotide and the second probe oligonucleotide each include a sequence that is substantially complementary to adjacent sequences of the analyte, and wherein the second probe oligonucleotide includes a capture probe binding domain; (b) coupling (e.g., ligating) the first probe oligonucleotide and the second probe oligonucleotide, thereby generating a connected probe (e.g., a ligation product) including the capture probe binding domain; (c) aligning the first substrate with a second substrate including an array, such that at least a portion of the biological sample is

aligned with at least a portion of the array, wherein the array includes a plurality of capture probes, wherein a capture probe of the plurality of capture probes includes: (i) a spatial barcode and (ii) a capture domain; (d) when the biological sample is aligned with at least a portion of the array, (i) releasing the connected probe (e.g., a ligation product) from the analyte and (ii) passively or actively migrating the connected probe (e.g., a ligation product) from the biological sample to the array; and (e) hybridizing the capture probe binding domain of the connected probe (e.g., a ligation product) to the capture domain.

In some embodiments, the process of transferring the connected probe (e.g., a ligation product) from the first substrate to the second substrate is referred to as a “sandwich” process.

The sandwich process is described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety. Described herein are methods in which an array with capture probes located on a substrate and a biological sample located on a different substrate, are contacted such that the array is in contact with the biological sample (e.g., the substrates are sandwiched together). In some embodiments, the array and the biological sample can be contacted (e.g., sandwiched), without the aid of a substrate holder. In some embodiments, the array and biological sample substrates can be placed in a substrate holder (e.g., an array alignment device) designed to align the biological sample and the array. For example, the substrate holder can have placeholders for two substrates. In some embodiments, an array including capture probes can be positioned on one side of the substrate holder (e.g., in a first substrate placeholder). In some embodiments, a biological sample can be placed on the adjacent side of the substrate holder in a second placeholder. In some embodiments, a hinge can be located between the two substrate placeholders that allows the substrate holder to close, e.g., make a sandwich between the two substrate placeholders. In some embodiments, when the substrate holder is closed the biological sample and the array with capture probes are contacted with one another under conditions sufficient to allow analytes present in the biological sample to interact with the capture probes of the array. For example, dried permeabilization reagents can be placed on the biological sample and rehydrated. A permeabilization solution can be flowed through the substrate holder to permeabilize the biological sample and allow analytes in the biological sample to interact with the capture probes. Additionally, the temperature of the substrates or permeabilization solution can be used to initiate or control the rate of permeabilization. For example, the substrate including the array, the substrate including the biological sample, or both substrates can be held at a low temperature to slow diffusion and permeabilization efficiency. Once sandwiched, in some embodiments, the substrates can be heated to initiate

permeabilization and/or increase diffusion efficiency. Transcripts that are released from the permeabilized tissue can diffuse to the array and be captured by the capture probes. The sandwich can be opened, and cDNA synthesis can be performed on the array.

In some embodiments, the methods as disclosed herein include hybridizing of one or more probe oligonucleotide probe pairs (e.g., RTL probes) to adjacent or nearby sequences of a target analyte (e.g., RNA; e.g., mRNA) of interest. In some embodiments, the probe oligonucleotide pairs include sequences that are complementary or substantially complementary to an analyte. For example, in some embodiments, each probe oligonucleotide includes a sequence that is complementary or substantially complementary to an mRNA of interest (e.g., to a portion of the sequence of an mRNA of interest). In some embodiments, each target analyte includes a first target region and a second target region. In some embodiments, the methods include providing a plurality of first probe oligonucleotides and a plurality of second probe oligonucleotides, wherein a pair of probe oligonucleotides for a target analyte comprises both a first and second probe oligonucleotide. In some embodiments, a first probe oligonucleotide hybridizes to a first target region of the analyte, and the second probe oligonucleotide hybridizes to a second, adjacent or nearly adjacent target region of the analyte.

In some instances, the probe oligonucleotides are DNA molecules. In some instances, the first probe oligonucleotide is a DNA molecule. In some instances, the second probe oligonucleotide is a DNA molecule. In some instances, the first probe oligonucleotide comprises at least two ribonucleic acid bases at the 3' end. In some instances, the second probe oligonucleotide comprises a phosphorylated nucleotide at the 5' end.

RTL probes can be designed using methods known in the art. In some instances, probe pairs are designed to cover an entire transcriptome of a species (e.g., a mouse or a human). In some instances, RTL probes are designed to cover a subset of a transcriptome (e.g., a mouse or a human). In some instances, the methods disclosed herein utilize about 500, about 1000, about 2000, about 3000, about 4000, about 5000, about 6000, about 7000, about 8000, about 9000, about 10,000, about 15,000, about 20,000, or more probe pairs.

In some embodiments, one of the probe oligonucleotides of the pair of probe oligonucleotides for RTL includes a poly(A) sequence or a complement thereof. In some instances, the poly(A) sequence or a complement thereof is on the 5' end of one of the probe oligonucleotides. In some instances, the poly(A) sequence or a complement thereof is on the 3' end of one of the probe oligonucleotides. In some embodiments, one probe oligonucleotide of the pair of probe oligonucleotides for RTL includes a degenerate or UMI sequence. In

some embodiments, the UMI sequence is specific to a particular target or set of targets. In some instances, the UMI sequence or a complement thereof is on the 5' end of one of the probe oligonucleotides. In some instances, the UMI sequence or a complement thereof is on the 3' end of one of the probe oligonucleotides.

5 In some instances, the first and second target regions of an analyte are directly adjacent to one another. In some embodiments, the complementary sequences to which the first probe oligonucleotide and the second probe oligonucleotide hybridize are 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, about 15, about 20, about 25, about 30, about 35, about 40, about 45, about 50, about 55, about 60, about 65, about 70, about 75, about 80, about 85, about 90, about 95,
10 about 100, about 125, or about 150 nucleotides away from each other. Gaps between the probe oligonucleotides may first be filled prior to coupling (e.g., ligation), using, for example, dNTPs in combination with a polymerase such as polymerase mu, DNA polymerase, RNA polymerase, reverse transcriptase, VENT polymerase, Taq polymerase, and/or any combinations, derivatives, and variants (e.g., engineered mutants) thereof. In some
15 embodiments, when the first and second probe oligonucleotides are separated from each other by one or more nucleotides, deoxyribonucleotides are used to extend and couple (e.g., ligate) the first and second probe oligonucleotides.

 In some instances, the first probe oligonucleotide and the second probe oligonucleotide hybridize to an analyte on the same transcript. In some instances, the first
20 probe oligonucleotide and the second probe oligonucleotide hybridize to an analyte on the same exon. In some instances, the first probe oligonucleotide and the second probe oligonucleotide hybridize to an analyte on different exons. In some instances, the first probe oligonucleotide and the second probe oligonucleotide hybridize to an analyte that is the result of a translocation event (e.g., in the setting of cancer). The methods provided herein make it
25 possible to identify alternative splicing events, translocation events, and mutations that change the hybridization rate of one or both probe oligonucleotides (e.g., single nucleotide polymorphisms, insertions, deletions, point mutations).

 In some embodiments, the first and/or second probe as disclosed herein includes at least two ribonucleic acid bases at the 3' end; a functional sequence; a phosphorylated
30 nucleotide at the 5' end; and/or a capture probe binding domain. In some embodiments, the functional sequence is a primer sequence.

 The "capture probe binding domain" is a sequence that is complementary to a particular capture domain present in a capture probe. In some embodiments, the capture probe binding domain includes a poly(A) sequence. In some embodiments, the capture probe

binding domain includes a poly-uridine sequence, a poly-thymidine sequence, or a combination thereof. In some embodiments, the capture probe binding domain includes a random sequence (e.g., a random hexamer or octamer). In some embodiments, the capture probe binding domain is complementary to a capture domain in a capture probe that detects a particular target(s) of interest. In some embodiments, a capture probe binding domain blocking moiety that interacts with the capture probe binding domain is provided. In some embodiments, a capture probe binding domain blocking moiety includes a sequence that is complementary or substantially complementary to a capture probe binding domain. In some embodiments, a capture probe binding domain blocking moiety prevents the capture probe binding domain from binding the capture probe when present. In some embodiments, a capture probe binding domain blocking moiety is removed prior to binding the capture probe binding domain (e.g., present in a connected probe (e.g., a ligation product)) to a capture probe. In some embodiments, a capture probe binding domain blocking moiety comprises a poly-uridine sequence, a poly-thymidine sequence, or a combination thereof.

Hybridization of the probe oligonucleotides to the target analyte can occur at a target having a sequence that is 100% complementary to the probe oligonucleotide(s). In some embodiments, hybridization can occur at a target having a sequence that is at least (e.g. at least about) 80%, at least (e.g. at least about) 85%, at least (e.g. at least about) 90%, at least (e.g. at least about) 95%, at least (e.g. at least about) 96%, at least (e.g. at least about) 97%, at least (e.g. at least about) 98%, or at least (e.g. at least about) 99% complementary to the probe oligonucleotide(s). After hybridization, in some embodiments, the first probe oligonucleotide is extended. After hybridization, in some embodiments, the second probe oligonucleotide is extended. For example, in some instances a first probe oligonucleotide hybridizes to a target sequence upstream for a second oligonucleotide probe, whereas in other instances a first probe oligonucleotide hybridizes to a target sequence downstream of a second probe oligonucleotide.

In some embodiments, methods disclosed herein include a wash step after hybridizing the first and the second probe oligonucleotides. The wash step removes any unbound oligonucleotides and can be performed using any technique known in the art. In some embodiments, a pre-hybridization buffer is used to wash the sample. In some embodiments, a phosphate buffer is used. In some embodiments, multiple wash steps are performed to remove unbound oligonucleotides. For example, it is advantageous to decrease the amount of unhybridized probes present in a biological sample as they may interfere with downstream applications and methods.

In some embodiments, after hybridization of probe oligonucleotides (e.g., first and the second probe oligonucleotides) to the target analyte, the probe oligonucleotides (e.g., the first probe oligonucleotide and the second probe oligonucleotide) are coupled (e.g., ligated) together, creating a single connected probe (e.g., a ligation product) that is complementary to the target analyte. Ligation can be performed enzymatically or chemically, as described herein. For example, the first and second probe oligonucleotides are hybridized to the first and second target regions of the analyte, and the probe oligonucleotides are subjected to a nucleic acid reaction to ligate them together. For example, the probes may be subjected to an enzymatic ligation reaction using a ligase (e.g., T4 RNA ligase (Rnl2), a SplintR ligase, or a T4 DNA ligase). See, e.g., Zhang L., et al.; Archaeal RNA ligase from thermococcus kodakarensis for template dependent ligation RNA Biol. 2017; 14(1): 36–44 for a description of KOD ligase. A skilled artisan will understand that various reagents, buffers, cofactors, etc. may be included in a ligation reaction depending on the ligase being used.

In some embodiments, the first probe oligonucleotide and the second probe oligonucleotides are on a contiguous nucleic acid sequence. In some embodiments, the first probe oligonucleotide is on the 3' end of the contiguous nucleic acid sequence. In some embodiments, the first probe oligonucleotide is on the 5' end of the contiguous nucleic acid sequence. In some embodiments, the second probe oligonucleotide is on the 3' end of the contiguous nucleic acid sequence. In some embodiments, the second probe oligonucleotide is on the 5' end of the contiguous nucleic acid sequence.

In some embodiments, the method further includes hybridizing a third probe oligonucleotide to the first probe oligonucleotide and the second probe oligonucleotide such that the first probe oligonucleotide and the second probe oligonucleotide abut each other. In some embodiments, the third probe oligonucleotide comprises a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% complementary to a portion of the first probe oligonucleotide that hybridizes to the third probe oligonucleotide. In some embodiments, the third probe oligonucleotide comprises a sequence that is 100% complementary to a portion of the first probe oligonucleotide that hybridizes to the third probe oligonucleotide. In some embodiments, the third probe oligonucleotide comprises a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% complementary to a portion of the second probe oligonucleotide that hybridizes to the third probe oligonucleotide. In some embodiments, the third probe oligonucleotide comprises a sequence that is 100%

complementary to a portion of the second probe oligonucleotide that hybridizes to the third probe oligonucleotide.

In some embodiments, a method for identifying a location of an analyte in a biological sample exposed to different permeabilization conditions includes (a) contacting the biological sample with a substrate, wherein the substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises a capture domain; (b) contacting the biological sample with a first probe oligonucleotide and a second probe oligonucleotide, wherein the first probe oligonucleotide and the second probe oligonucleotide are substantially complementary to adjacent sequences of the analyte, and wherein the second probe oligonucleotide comprises a capture probe-binding domain that is capable of binding to a capture domain of the capture probe; (c) hybridizing the first probe oligonucleotide and the second probe oligonucleotide to adjacent sequences of the analyte; (d) coupling (e.g., ligating) the first probe oligonucleotide and the second probe oligonucleotide, thereby creating a connected probe (e.g., a connected probe (e.g., a ligation product)) that is substantially complementary to the analyte; (e) releasing the connected probe (e.g., a ligation product) from the analyte; (f) hybridizing the capture probe-binding domain of the connected probe (e.g., a ligation product) to the hybridization domain of the capture probe; (g) hybridizing a padlock oligonucleotide to the connected probe (e.g., a ligation product) bound to the capture domain (e.g., such that the padlock oligonucleotide is circularized), wherein the padlock oligonucleotide comprises: (i) a first sequence that is substantially complementary to a first portion of the connected probe (e.g., a ligation product), (ii) a backbone sequence, and (iii) a second sequence that is substantially complementary to a second portion of the connected probe (e.g., a ligation product); and (i) ligating and amplifying the circularized padlock oligonucleotide (e.g., using rolling circle amplification using the circularized padlock oligonucleotide as a template), thereby creating an amplified circularized padlock oligonucleotide, and using the amplified circularized padlock oligonucleotide to identify the location of the analyte in the biological sample.

In some embodiments, the method further includes amplifying the connected probe (e.g., a ligation product) prior to the releasing step. In some embodiments, the entire connected probe (e.g., a ligation product) is amplified. In some embodiments, only part of the connected probe (e.g., a ligation product) is amplified. In some embodiments, amplification is isothermal. In some embodiments, amplification is not isothermal. Amplification can be performed using any of the methods described herein such as, but not limited to, a strand-displacement amplification reaction, a rolling circle amplification reaction, a ligase chain

reaction, a transcription-mediated amplification reaction, an isothermal amplification reaction, and/or a 10 loop-mediated amplification reaction. In some embodiments, amplifying the connected probe (e.g., a ligation product) creates an amplified connected probe (e.g., a ligation product) that includes (i) all or part of sequence of the connected probe (e.g., a ligation product) specifically bound to the capture domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof.

In some embodiments, the method further includes determining (i) all or a part of the sequence of the connected probe (e.g., a ligation product), or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some embodiments, the method further includes using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample.

In some embodiments, after coupling (e.g., ligation) of the first and second probe oligonucleotides to create a ligation product, the connected probe (e.g., a ligation product) is released from the analyte. To release the connected probe (e.g., a ligation product), an endoribonuclease (e.g., RNase A, RNase C, RNase H, or RNase I) is used. An endoribonuclease such as RNase H specifically cleaves RNA in RNA:DNA hybrids. In some embodiments, the connected probe (e.g., a ligation product) is released enzymatically. In some embodiments, an endoribonuclease is used to release the probe from the analyte. In some embodiments, the endoribonuclease is one or more of RNase H. In some embodiments, the RNase H is RNase H1 or RNase H2.

In some embodiments, the releasing of the connected probe (e.g., a ligation product) includes contacting the biological sample with a reagent medium comprising a permeabilization agent and an agent for releasing the connected probe (e.g., a ligation product), thereby permeabilizing the biological sample and releasing the connected probe (e.g., a ligation product) from the analyte. In some embodiments, the agent for releasing the connected probe (e.g., a ligation product) comprises a nuclease. In some embodiments, the nuclease is an endonuclease. In some embodiments, the nuclease is an exonuclease. In some embodiments, the nuclease includes an RNase. In some embodiments, the RNase is selected from RNase A, RNase C, RNase H, or RNase I.

In some embodiments, the reagent medium comprises polyethylene glycol (PEG). In some embodiments, the PEG is from about PEG 2K to about PEG 16K. In some embodiments, the PEG is PEG 2K, 3K, 4K, 5K, 6K, 7K, 8K, 9K, 10K, 11K, 12K, 13K, 14K, 15K, or 16K. In some embodiments, the PEG is present at a concentration from about 2% to

25%, from about 4% to about 23%, from about 6% to about 21%, or from about 8% to about 20% (v/v).

In some embodiments, the reagent medium includes a wetting agent.

In some instances, after creation of the connected probe (e.g., a ligation product), the methods disclosed herein include simultaneous treatment of the biological sample with a permeabilization agent such as proteinase K (to permeabilize the biological sample) and a releasing agent such as an endonuclease such as RNase H (to release the connected probe (e.g., a ligation product) from the analyte). In some instances, the permeabilization step and releasing step occur at the same time. In some instances, the permeabilization step occurs before the releasing step. In some embodiments, the permeabilization agent comprises a protease. In some embodiments, the protease is selected from trypsin, pepsin, elastase, or Proteinase K. In some embodiments, the protease is an endopeptidase. Endopeptidases that can be used include but are not limited to trypsin, chymotrypsin, elastase, thermolysin, pepsin, clostripain, glutamyl endopeptidase (GluC), ArgC, peptidyl-asp endopeptidase (ApsN), endopeptidase LysC and endopeptidase LysN. In some embodiments, the endopeptidase is pepsin.

In some embodiments, the reagent medium further includes a detergent. In some embodiments, the detergent is selected from sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100TM, or Tween-20TM. In some embodiments, the reagent medium includes less than 5 w/v% of a detergent selected from sodium dodecyl sulfate (SDS) and sarkosyl. In some embodiments, the reagent medium includes at least 5% w/v% of a detergent selected from SDS and sarkosyl. In some embodiments, the reagent medium does not include SDS or sarkosyl.

In some embodiments, the biological sample and the array are contacted with the reagent medium for about 1 to about 60 minutes (e.g., about 1 to about 55 minutes, about 1 to about 50 minutes, about 1 to about 45 minutes, about 1 to about 40 minutes, about 1 to about 35 minutes, about 1 to about 30 minutes, about 1 to about 25 minutes, about 1 to about 20 minutes, about 1 to about 15 minutes, about 1 to about 10 minutes, about 1 to about 5 minutes, about 5 to about 60 minutes, about 5 to about 55 minutes, about 5 to about 50 minutes, about 5 to about 45 minutes, about 5 to about 40 minutes, about 5 to about 35 minutes, about 5 to about 30 minutes, about 5 to about 25 minutes, about 5 to about 20 minutes, about 5 to about 15 minutes, about 5 to about 10 minutes, about 10 to about 60 minutes, about 10 to about 55 minutes, about 10 to about 50 minutes, about 10 to about 45 minutes, about 10 to about 40 minutes, about 10 to about 35 minutes, about 10 to about 30

minutes, about 10 to about 25 minutes, about 10 to about 20 minutes, about 10 to about 15 minutes, about 15 to about 60 minutes, about 15 to about 55 minutes, about 15 to about 50 minutes, about 15 to about 45 minutes, about 15 to about 40 minutes, about 15 to about 35 minutes, about 15 to about 30 minutes, about 15 to about 25 minutes, about 15 to about 20 minutes, about 20 to about 60 minutes, about 20 to about 55 minutes, about 20 to about 50 minutes, about 20 to about 45 minutes, about 20 to about 40 minutes, about 20 to about 35 minutes, about 20 to about 30 minutes, about 20 to about 25 minutes, about 25 to about 60 minutes, about 25 to about 55 minutes, about 25 to about 50 minutes, about 25 to about 45 minutes, about 25 to about 40 minutes, about 25 to about 35 minutes, about 25 to about 30 minutes, about 30 to about 60 minutes, about 30 to about 55 minutes, about 30 to about 50 minutes, about 30 to about 45 minutes, about 30 to about 40 minutes, about 30 to about 35 minutes, about 35 to about 60 minutes, about 35 to about 55 minutes, about 35 to about 50 minutes, about 35 to about 45 minutes, about 35 to about 40 minutes, about 40 to about 60 minutes, about 40 to about 55 minutes, about 40 to about 50 minutes, about 40 to about 45 minutes, about 45 to about 60 minutes, about 45 to about 55 minutes, about 45 to about 50 minutes, about 50 to about 60 minutes, about 50 to about 55 minutes, or about 55 to about 60 minutes). In some embodiments, the biological sample and the array are contacted with the reagent medium for about 30 minutes.

In some embodiments, the connected probe (e.g., a ligation product) includes a capture probe binding domain, which can hybridize to a capture probe (e.g., a capture probe immobilized, directly or indirectly, on a substrate). In some embodiments, the capture probe includes a spatial barcode and the capture domain. In some embodiments, the capture probe binding domain of the connected probe (e.g., a ligation product) specifically binds to the capture domain of the capture probe.

In some embodiments, methods provided herein include mounting a biological sample on a first substrate, then aligning the first substrate with a second substrate including an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array includes a plurality of capture probes. After hybridization of the connected probe (e.g., a ligation product) to the capture probe, downstream methods as disclosed herein can be performed.

In some embodiments, at least 50% of connected probes (e.g., a ligation products) released from the portion of the biological sample aligned with the portion of the array are captured by capture probes of the portion of the array. In some embodiments, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at

least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% of connected probe (e.g., a ligation products) are detected in spots directly under the biological sample.

In some embodiments, the capture probe includes a poly(T) sequence. In some embodiments, capture probe includes a sequence specific to the analyte. In some
5 embodiments, the capture probe includes a functional domain. In some embodiments, the capture probe further includes one or more functional domains, a unique molecular identifier (UMI), a cleavage domain, and combinations thereof. In some embodiments, the capture probe binding domain includes a poly(A) sequence. In some embodiments, the capture probe binding domain includes a sequence complementary to a capture domain of a capture probe
10 that detects a target analyte of interest. In some embodiments, the analyte is RNA. In some embodiments, the analyte is mRNA.

In some embodiments, the connected probe (e.g., a ligation product) (e.g., the analyte derived molecule) includes a capture probe binding domain, which can hybridize to a capture probe (e.g., a capture probe immobilized, directly or indirectly, on a substrate). Methods
15 provided herein include contacting a biological sample with a substrate, wherein the capture probe is affixed to the substrate (e.g., immobilized to the substrate, directly or indirectly). After hybridization of the connected probe (e.g., a ligation product) to the capture probe, downstream methods as disclosed herein (e.g., sequencing, in situ analysis such as RCA) can be performed.

In some embodiments, the method further includes analyzing a different analyte in the
20 biological sample. In some embodiments, the analysis of the different analyte includes (a) further contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents includes an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically
25 binds to the different analyte, and wherein the capture agent barcode domain includes an analyte binding moiety barcode and an capture handle sequence that is complementary to a capture domain of a capture probe; and (b) hybridizing the analyte capture sequence to the capture domain.

An exemplary embodiment of a workflow for analysis of protein and RNA analytes is
30 shown in **FIG. 16A**. As shown in FIG. 16A, a fixed tissue sample mounted on a first substrate (e.g., a slide-mounted tissue sample) is decrosslinked, followed by hybridization of probe pairs to nucleic acid analyte analytes. Also as shown in FIG. 16A, a first and second probe of a probe pair is connected, e.g., ligated. The sample is optionally washed (e.g., with a buffer), prior to incubation with an analyte capture agent (e.g., an antibody) that specifically

binds a different analyte, e.g., a protein analyte. The analyte capture agent comprises a capture agent barcode domain. In some embodiments, the analyte capture agent is an antibody with an oligonucleotide tag, the oligonucleotide tag comprising a capture agent barcode domain. In some embodiments, the connected probes (e.g., the ligation products) and antibody oligonucleotide tags are released from the tissue under sandwich conditions as described herein. For the sandwich conditions, the tissue-mounted slide can be aligned with an array and permeabilized with a reagent medium in the sandwich configuration as described herein (see, e.g., **FIG. 16B**). In some embodiments, the reagent medium comprises RNase and a permeabilization agent (e.g., Proteinase K). Permeabilization releases the connected probe (e.g., a ligation product) and capture agent barcode domain, for capture onto a second substrate comprising an array with a plurality of capture probes (see, e.g., **FIG. 16B**). After capture of the connected probe and capture agent barcode domain, the tissue slide can be removed (e.g., the sandwich can be “opened” or “broken”).

In some embodiments, following opening of the sandwich, the capture probes can be extended, sequencing libraries can be prepared and sequenced, and the results can be analyzed computationally.

In some embodiments, the method further includes determining (i) all or part of the sequence of the capture agent barcode domain; and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some embodiments, the method further includes using the determined sequence of (i), and (ii) to analyze the different analyte in the biological sample. In some embodiments, the releasing step further releases the capture agent barcode domain from the different analyte. In some embodiments, the different analyte is a protein analyte. In some embodiments, the protein analyte is an extracellular protein. In some embodiments, the protein analyte is an intracellular protein.

25

(c) Capturing Analytes for Spatial Detection using Analyte Capture Agents

In some embodiments, the methods, compositions, devices, and systems provided herein utilize analyte capture agents for spatial detection. An “analyte capture agent” refers to a molecule that interacts with a target analyte (e.g., a protein) and with a capture probe. Such analyte capture agents can be used to identify the analyte. In some embodiments, the analyte capture agent can include an analyte binding moiety and a capture agent barcode domain. In some embodiments, the analyte capture agent includes a linker. In some embodiments, the linker is a cleavable linker. In some embodiments, the cleavable linker is a photo-cleavable linker, a UV-cleavable linker, or an enzyme cleavable linker.

An analyte binding moiety is a molecule capable of binding to a specific analyte. In some embodiments, the analyte binding moiety comprises an antibody or antibody fragment. In some embodiments, the analyte binding moiety comprises a polypeptide and/or an aptamer. In some embodiments, the analyte is a protein (e.g., a protein on a surface of a cell or an intracellular protein).

A capture agent barcode domain can include a capture handle sequence which can hybridize to at least a portion or an entirety of a capture domain of a capture probe. In some embodiments, the capture handle sequence is complementary to a portion or entirety of a capture domain of a capture probe. In some embodiments, the capture handle sequence includes a poly (A) tail. In some embodiments, the capture handle sequence includes a sequence capable of binding a poly (T) domain. In some embodiments, the capture agent barcode domain comprises an analyte binding moiety barcode and a capture handle sequence. The analyte binding moiety barcode refers to a barcode that is associated with or otherwise identifies the analyte binding moiety, and the capture handle sequence can hybridize to a capture probe. In some embodiments, the capture handle sequence specifically binds to the capture domain of the capture probe. Other embodiments of an analyte capture agent useful in spatial analyte detection are described herein.

Provided herein are methods for analyzing an analyte in a biological sample including (a) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents includes an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain includes an analyte binding moiety barcode and an capture handle sequence; (b) contacting the biological sample with a reagent medium including an agent for releasing the capture agent barcode domain from the analyte binding moiety, thereby releasing the capture agent barcode domain from the analyte binding moiety; and (c) hybridizing the capture handle sequence to a capture domain of a capture probe, wherein the capture probe includes (i) a spatial barcode and (ii) a capture domain.

Also provided herein are methods for analyzing an analyte in a biological sample mounted on a first substrate including (a) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents includes an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain includes an analyte binding moiety barcode and an capture handle sequence;

(b) aligning the first substrate with a second substrate comprising an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array includes a plurality of capture probes, wherein a capture probe of the plurality of capture probes includes (i) a spatial barcode and (ii) a capture domain; (c) when the biological sample is aligned with at least a portion of the array, (i) releasing the capture agent barcode domain from the analyte and (ii) passively or actively migrating the capture agent barcode domain from the biological sample to the array; and (d) coupling the capture handle sequence to the capture domain.

Also provided herein are methods for analyzing an analyte in a biological sample mounted on a first substrate including (a) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents includes an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain includes an analyte binding moiety barcode and an capture handle sequence; (b) aligning the first substrate with a second substrate comprising an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array includes a plurality of capture probes, wherein a capture probe of the plurality of capture probes includes (i) a spatial barcode and (ii) a capture domain; (c) when the biological sample is aligned with at least a portion of the array, (i) releasing the capture agent barcode domain from the analyte and (ii) passively or actively migrating the capture agent barcode domain from the biological sample to the array; and (d) hybridizing the capture handle sequence to the capture domain.

In some embodiments, the process of transferring the connected probe (e.g., a ligation product) from the first substrate to the second substrate is referred to as a “sandwich process”. The sandwich process is described above and in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

In some embodiments, the method further includes determining (i) all or a part of the capture agent barcode domain, or a complement thereof; and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some embodiments, the method further includes using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample.

In some embodiments, an analyte capture agent is introduced to a biological sample, wherein the analyte binding moiety specifically binds to a target analyte, and then the biological sample can be treated to release the capture agent barcode domain from the

biological sample. In some embodiments, the capture agent barcode domain can then migrate and bind to a capture domain of a capture probe, and the capture agent barcode domain can be extended to generate a spatial barcode complement at the end of the capture agent barcode domain. In some embodiments, the spatially-tagged capture agent barcode domain can be
5 denatured from the capture probe, and analyzed using methods described herein.

In some embodiments, the releasing includes contacting the biological sample and the array with a reagent medium including a nuclease. In some embodiments, the nuclease includes an RNase. In some embodiments, the RNase is selected from RNase A, RNase C, RNase H, and RNase I. In some embodiments, the reagent medium further includes a
10 permeabilization agent. In some embodiments, the releasing further includes simultaneously permeabilizing the biological sample and releasing the capture agent barcode domain from the analyte. In some embodiments, the permeabilization agent further includes a protease. In some embodiments, the protease is selected from trypsin, pepsin, elastase, or Proteinase K.

In some embodiments, the capture agent barcode domain is released from the analyte
15 binding moiety by using a different stimulus that can include, but is not limited to, a proteinase (e.g., Proteinase K), an RNase, and UV light.

In some embodiments, the reagent medium further includes a detergent. In some embodiments, the detergent is selected from sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100™, or Tween-20™. In some embodiments, the reagent medium
20 includes less than 5 w/v% of a detergent selected from sodium dodecyl sulfate (SDS) and sarkosyl. In some embodiments, the reagent medium includes at least 5% w/v% of a detergent selected from SDS and sarkosyl. In some embodiments, the reagent medium does not include SDS or sarkosyl.

In some embodiments, the biological sample and the array are contacted with the
25 reagent medium for about 1 to about 60 minutes (e.g., about 1 to about 55 minutes, about 1 to about 50 minutes, about 1 to about 45 minutes, about 1 to about 40 minutes, about 1 to about 35 minutes, about 1 to about 30 minutes, about 1 to about 25 minutes, about 1 to about 20 minutes, about 1 to about 15 minutes, about 1 to about 10 minutes, about 1 to about 5 minutes, about 5 to about 60 minutes, about 5 to about 55 minutes, about 5 to about 50 minutes, about 5 to about 45 minutes, about 5 to about 40 minutes, about 5 to about 35 minutes, about 5 to about 30 minutes, about 5 to about 25 minutes, about 5 to about 20 minutes, about 5 to about 15 minutes, about 5 to about 10 minutes, about 10 to about 60 minutes, about 10 to about 55 minutes, about 10 to about 50 minutes, about 10 to about 45 minutes, about 10 to about 40 minutes, about 10 to about 35 minutes, about 10 to about 30
30 minutes).

minutes, about 10 to about 25 minutes, about 10 to about 20 minutes, about 10 to about 15 minutes, about 15 to about 60 minutes, about 15 to about 55 minutes, about 15 to about 50 minutes, about 15 to about 45 minutes, about 15 to about 40 minutes, about 15 to about 35 minutes, about 15 to about 30 minutes, about 15 to about 25 minutes, about 15 to about 20 minutes, about 20 to about 60 minutes, about 20 to about 55 minutes, about 20 to about 50 minutes, about 20 to about 45 minutes, about 20 to about 40 minutes, about 20 to about 35 minutes, about 20 to about 30 minutes, about 20 to about 25 minutes, about 25 to about 60 minutes, about 25 to about 55 minutes, about 25 to about 50 minutes, about 25 to about 45 minutes, about 25 to about 40 minutes, about 25 to about 35 minutes, about 25 to about 30 minutes, about 30 to about 60 minutes, about 30 to about 55 minutes, about 30 to about 50 minutes, about 30 to about 45 minutes, about 30 to about 40 minutes, about 30 to about 35 minutes, about 35 to about 60 minutes, about 35 to about 55 minutes, about 35 to about 50 minutes, about 35 to about 45 minutes, about 35 to about 40 minutes, about 40 to about 60 minutes, about 40 to about 55 minutes, about 40 to about 50 minutes, about 40 to about 45 minutes, about 45 to about 60 minutes, about 45 to about 55 minutes, about 45 to about 50 minutes, about 50 to about 60 minutes, about 50 to about 55 minutes, or about 55 to about 60 minutes). In some embodiments, the biological sample and the array are contacted with the reagent medium for about 30 minutes.

Also provided herein are methods further including analyzing a different analyte in the biological sample. In some embodiments, the analysis of the different analyte includes (a) hybridizing a first probe oligonucleotide and a second probe oligonucleotide to the different analyte, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the different analyte, and wherein the second probe oligonucleotide comprises a capture probe binding domain; (b) ligating the first probe oligonucleotide and the second probe oligonucleotide, thereby generating a connected probe (e.g., a ligation product) comprising the capture probe binding domain; and (c) hybridizing the capture probe binding domain of the connected probe (e.g., a ligation product) to the capture domain.

In some embodiments, the method further includes determining (i) all or part of the sequence of the connected probe (e.g., a ligation product), or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof. In some embodiments, the method further includes using the determined sequence of (i), and (ii) to analyze the different analyte in the biological sample. In some embodiments, the releasing step further releases the connected probe (e.g., a ligation product) from the different analyte.

In some embodiments, the different analyte is RNA. In some embodiments, the different analyte is mRNA.

5 In some embodiments, the capture probe comprises a poly(T) sequence. In some embodiments, the capture probe comprises a sequence complementary to the capture handle sequence. In some embodiments, the capture probe comprises a functional domain. In some embodiments, the capture probe further comprises one or more functional domains, a unique molecular identifier (UMI), a cleavage domain, and combinations thereof.

10 In some embodiments, the biological sample is a tissue sample. In some embodiments, the tissue sample is a tissue section. In some embodiments, the tissue sample is a fixed tissue sample. In some embodiments, the fixed tissue sample is a formalin fixed paraffin embedded (FFPE) tissue sample. In some embodiments, the FFPE tissue is deparaffinized and decrosslinked prior to step (a) of any one of the methods provided herein. In some embodiments, the fixed tissue sample is a formalin fixed paraffin embedded cell pellet. In some embodiments, the tissue sample is a fresh tissue sample or a frozen tissue sample. In some embodiments, the tissue sample is fixed and stained prior to step (a) of any one of the methods provided herein.

15 In some instances, RTL is performed between two oligonucleotides that each are affixed to an analyte binding moiety (i.e., a protein-binding moiety). Generally, the methods of RTL in this setting is as follows. In some embodiments, provided herein is a method of determining a location of at least one analyte in a biological sample including: (a) hybridizing a first analyte-binding moiety to a first analyte in the biological sample, wherein the first analyte-binding moiety is bound to a first oligonucleotide, wherein the first oligonucleotide comprises: (i) a functional sequence; (ii) a first barcode; and (iii) a first bridge sequence; (b) hybridizing a second analyte-binding moiety to a second analyte in the biological sample, wherein the second analyte-binding moiety is bound to a second oligonucleotide; wherein the second oligonucleotide comprises: (i) capture probe binding domain sequence, (ii) a second barcode; and (ii) a second bridge sequence; (c) contacting the biological sample with a third oligonucleotide; (d) hybridizing the third oligonucleotide to the first bridge sequence of the first oligonucleotide and second bridge sequence of the second oligonucleotide; (e) ligating the first oligonucleotide and the second oligonucleotide, creating a connected probe (e.g., a ligation product); (f) contacting the biological sample with a substrate, wherein a capture probe is affixed to the substrate, wherein the capture probe comprises a spatial barcode and the capture domain; and (g) allowing the capture probe binding domain sequence of the

second oligonucleotide to specifically bind to the capture domain. In some instances, the connected probe (e.g., a ligation product) is cleaved from the analyte binding moieties.

In some instances, two analytes (e.g., two different proteins) in close proximity in a biological sample are detected by a first analyte-binding moiety and a second analyte-binding moiety, respectively. In some embodiments, a first analyte-binding moiety and/or the second analyte-binding moiety is an analyte capture agent (e.g., any of the exemplary analyte capture agents described herein). In some embodiments, the first analyte-binding moiety and/or the second analyte-binding moiety is a first protein. In some embodiments, the first analyte-binding moiety and/or the second analyte-binding moiety is an antibody. For example, the antibody can include, without limitation, a monoclonal antibody, recombinant antibody, synthetic antibody, a single domain antibody, a single-chain variable fragment (scFv), and or an antigen-binding fragment (Fab). In some embodiments, the first analyte-binding moiety binds to a cell surface analyte (e.g., any of the exemplary cell surface analytes described herein). In some embodiments, binding of the analyte is performed metabolically. In some embodiments, binding of the analyte is performed enzymatically. In some embodiments, the methods include a secondary antibody that binds to a primary antibody, enhancing its detection.

In some embodiments, the first analyte-binding moiety and the second analyte-binding moiety each bind to the same analyte. In some embodiments, the first analyte-binding moiety and/or second analyte-binding moiety each bind to a different analyte. For example, in some embodiments, the first analyte-binding moiety binds to a first polypeptide and the second analyte-binding moiety binds to a second polypeptide.

In some embodiments of any of the methods of determining a location of at least one analyte in a biological sample, a first and/or a second oligonucleotide are bound (e.g., conjugated or otherwise attached using any of the methods described herein) to a first analyte-binding moiety and/or a second analyte-binding moiety, respectively.

In some embodiments of any of the methods of determining a location of at least one analyte in a biological sample as described herein, a second oligonucleotide is bound (e.g., conjugated or otherwise attached using any of the methods described herein) to a second analyte-binding moiety. For example, the second oligonucleotide can be covalently linked to the second analyte-binding moiety. In some embodiments, the second oligonucleotide is bound to the second analyte-binding moiety via its 5' end. In some embodiments, the second oligonucleotide includes a free 3' end. In some embodiments the second oligonucleotide is

bound to the second analyte-binding moiety via its 3' end. In some embodiments, the second oligonucleotide includes a free 5' end.

In some embodiments, the oligonucleotides are bound to the first and/or second analyte-binding moieties via a linker (e.g., any of the exemplary linkers described herein). In some embodiments, the linker is a cleavable linker. In some embodiment, the linker is a linker with photo-sensitive chemical bonds (e.g., photo-cleavable linkers). In some embodiments, the linker is a cleavable linker that can undergo induced dissociation.

FIGs. 17A-17B show exemplary schematics illustrating methods to release the capture agent barcode domain from the analyte binding moiety. **FIG. 17A** shows the use of an RNase cleavable linker to release a capture agent barcode domain from an analyte binding moiety of an analyte capture agent. In particular, a target (indicated by the circle in **FIG. 17A**) interacts with an antibody. After an interaction is established, an enzyme (e.g., an RNase) cleaves a linker, freeing a barcode from the antibody. The barcode, which is specific to the antibody, is captured on a slide comprising a plurality of probes. Similarly, **FIG. 17B** shows the use of a UV cleavable linker to release a capture agent barcode domain from an analyte binding moiety of an analyte capture agent.

In some embodiments, the oligonucleotides are bound (e.g., attached via any of the methods described herein) to an analyte-binding domain via a 5' end.

In some embodiments, a barcode is used to identify the analyte-binding moiety to which it is bound. The barcode can be any of the exemplary barcodes described herein. In some embodiments, the first and/or second oligonucleotide include a capture probe binding domain sequence. For example, a capture probe binding domain sequence can be a poly(A) sequence when the capture domain sequence is a poly(T) sequence.

In some embodiments, a third oligonucleotide (e.g., a splint oligonucleotide) hybridizes to both the first and second oligonucleotides and enables ligation of the first oligonucleotide and the second oligonucleotide. In some embodiments, a ligase is used. In some aspects, the ligase includes a DNA ligase. In some aspects, the ligase includes a RNA ligase. In some aspects, the ligase includes T4 DNA ligase. In some embodiments, the ligase is a SplintR ligase.

30

(d) Sandwich Processes

In some embodiments, one or more analytes from the biological sample are released from the biological sample and migrate to a substrate comprising an array of capture probes for attachment to the capture probes of the array. In some embodiments, the release and

migration of the analytes to the substrate comprising the array of capture probes occurs in a manner that preserves the original spatial context of the analytes in the biological sample. In some embodiments, the biological sample is mounted on a first substrate and the substrate comprising the array of capture probes is a second substrate. In some embodiments, the alignment of the first substrate and the second substrate is facilitated by a sandwiching process. Accordingly, described herein are methods, compositions, devices, and systems for sandwiching together the first substrate as described herein with a second substrate having an array with capture probes.

FIG. 23 is a schematic diagram depicting an exemplary sandwiching process between a first substrate comprising a biological sample (e.g., a tissue section **2302** on a slide **2303**) and a second substrate comprising a spatially barcoded array, e.g., a slide **2304** that is populated with spatially-barcoded capture probes **2306**. During the exemplary sandwiching process, the first substrate is aligned with the second substrate, such that at least a portion of the biological sample is aligned with at least a portion of the array (e.g., aligned in a sandwich configuration). As shown, the second substrate (e.g., slide **2304**) is in a superior position to the first substrate (e.g., slide **2303**). In some embodiments, the first substrate (e.g., slide **2303**) may be positioned superior to the second substrate (e.g., slide **2304**). In some embodiments, the first and second substrates are aligned to maintain a gap or separation distance **2307** between the two substrates. When the first and second substrates are aligned, one or more analytes are released from the biological sample and actively or passively migrate to the array for capture. In some embodiments, the migration occurs while the aligned portions of the biological sample and the array are contacted with a reagent medium **2305**. The released one or more analytes may actively or passively migrate across the gap **2307** via the reagent medium **2305** toward the capture probes **2306**, and be captured by the capture probes **2306**.

In some embodiments, the separation distance **2307** between first and second substrates is maintained between 2 microns and 1 mm (e.g., between 2 microns and 800 microns, between 2 microns and 700 microns, between 2 microns and 600 microns, between 2 microns and 500 microns, between 2 microns and 400 microns, between 2 microns and 300 microns, between 2 microns and 200 microns, between 2 microns and 100 microns, between 2 microns and 25 microns, between 2 microns and 10 microns), measured in a direction orthogonal to the surface of first substrate that supports sample. In some embodiments, the separation distance **2307** between first and second substrates is less than 50 microns. In some instances, the distance is 2 microns. In some instances, the distance is 2.5 microns. In

some instances, the distance is about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 microns. In some embodiments, second substrate is placed in direct contact with the sample on the first substrate ensuring no diffusive spatial resolution losses. In some embodiments, the separation distance is measured in a direction orthogonal to a surface of the first substrate that supports the biological sample.

In some embodiments, the sandwiching process may be facilitated by a device, sample holder, sample handling apparatus, or system described in, e.g., US. Patent Application Pub. No. 20210189475, PCT/US2021/036788, or PCT/US2021/050931.

In some embodiments, the first and second substrates are placed in a substrate holder (e.g., an array alignment device) configured to align the biological sample and the array. In some embodiments, the device comprises a sample holder. In some embodiments, the sample holder includes a first member and a second member that receive a first substrate and a second substrate, respectively. The device can include an alignment mechanism that is connected to at least one of the members and aligns the first and second members. Thus, the devices of the disclosure can advantageously align the first substrate and the second substrate and any samples, barcoded probes, or permeabilization reagents that may be on the surface of the first and second substrates.

In some embodiments, the sandwiching process comprises: mounting the first substrate on a first member of a support device, the first member configured to retain the first substrate; mounting the second substrate on a second member of the support device, the second member configured to retain the second substrate, applying a reagent medium to the first substrate and/or the second substrate, the reagent medium comprising a permeabilization agent, operating an alignment mechanism of the support device to move the first member and/or the second member such that a portion of the biological sample is aligned (e.g., vertically aligned) with a portion of the array of capture probes and within a threshold distance of the array of capture probes, and such that the portion of the biological sample and the capture probe contact the reagent medium, wherein the permeabilization agent releases the analyte from the biological sample.

In some embodiments of a sample holder, the sample holder can include a first member including a first retaining mechanism configured to retain a first substrate comprising a sample. The first retaining mechanism can be configured to retain the first substrate disposed in a first plane. The sample holder can further include a second member including a second retaining mechanism configured to retain a second substrate disposed in a second plane. The sample holder can further includes an alignment mechanism connected to

one or both of the first member and the second member. The alignment mechanism can be configured to align the first and second members along the first plane and/or the second plane such that the sample contacts at least a portion of the reagent medium when the first and second members are aligned and within a threshold distance along an axis orthogonal to the second plane. The alignment mechanism may be configured to move the second member along the axis orthogonal to the second plane and/or move the first member along an axis orthogonal to the first plane.

In some embodiments, the alignment mechanism includes a linear actuator. In some embodiments, the alignment mechanism includes one or more of a moving plate, a bushing, a shoulder screw, a motor bracket, and a linear actuator. The moving plate may be coupled to the first member or the second member. The alignment mechanism may, in some cases, include a first moving plate coupled to the first member and a second moving plate coupled to the second member. In some embodiments, the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member. For example, the moving plate may be coupled to the second member and adjust the separation distance along a z axis (e.g., orthogonal to the second substrate) by moving the moving plate up in a superior direction toward the first substrate. In some embodiments, the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member. The movement of the moving plate may be accomplished by the linear actuator configured to move the first member and/or the second member at a velocity. The velocity may be controlled by a controller communicatively coupled to the linear actuator. In some embodiments, the linear actuator is configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec (e.g., at least 0.1 mm/sec to 2 mm/sec). In some aspects, the velocity may be selected to reduce or minimize bubble generation or trapping within the reagent medium. In some embodiments, the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs (e.g., between 0.1-4.0 pounds of force).

In some aspects, the velocity of the moving plate (e.g., closing the sandwich) may affect bubble generation or trapping within the reagent medium. It may be advantageous to minimize bubble generation or trapping within the reagent medium during the “sandwiching” process, as bubbles can interfere with the migration of analytes through the reagent medium to the array. In some embodiments, the closing speed is selected to minimize bubble generation or trapping within the reagent medium. In some embodiments, the closing speed

is selected to reduce the time it takes the flow front of the reagent medium from an initial point of contact with the first and second substrate to sweep across the sandwich area (also referred to herein as “closing time”). In some embodiments, the closing speed is selected to reduce the closing time to less than about 1100 milliseconds (ms). In some embodiments, the closing speed is selected to reduce the closing time to less than about 1000 ms. In some embodiments, the closing speed is selected to reduce the closing time to less than about 900 ms. In some embodiments, the closing speed is selected to reduce the closing time to less than about 750 ms. In some embodiments, the closing speed is selected to reduce the closing time to less than about 600 ms. In some embodiments, the closing speed is selected to reduce the closing time to about 550 ms or less. In some embodiments, the closing speed is selected to reduce the closing time to about 370 ms or less. In some embodiments, the closing speed is selected to reduce the closing time to about 200 ms or less. In some embodiments, the closing speed is selected to reduce the closing time to about 150 ms or less.

FIG. 37A is a perspective view of an example sample handling apparatus **3700** (also referred to herein as a support device, a sample holder, and an array alignment device) in a closed position in accordance with some example implementations. As shown, the sample handling apparatus **3700** includes a first member **3704**, a second member **3710**, optionally an image capture device **3720**, a first substrate **3706**, optionally a hinge **3715**, and optionally a mirror **3716**. The hinge **3715** may be configured to allow the first member **3704** to be positioned in an open or closed configuration by opening and/or closing the first member **3704** in a clamshell manner along the hinge **3715**.

FIG. 37B is a perspective view of the example sample handling apparatus **3700** in an open position in accordance with some example implementations. As shown, the sample handling apparatus **3700** includes one or more first retaining mechanisms **3708** configured to retain one or more first substrates **3706**. In the example of **FIG. 37B**, the first member **3704** is configured to retain two first substrates **3706**, however the first member **3704** may be configured to retain more or fewer first substrates **3706**.

In some aspects, when the sample handling apparatus **3700** is in an open position (as in **FIG. 37B**), the first substrate **3706** and/or the second substrate **3712** may be loaded and positioned within the sample handling apparatus **3700** such as within the first member **3704** and the second member **3710**, respectively. As noted, the hinge **3715** may allow the first member **3704** to close over the second member **3710** and form a sandwich configuration (e.g., the sandwich configuration shown in **FIG. 23**).

In some aspects, after the first member **3704** closes over the second member **3710**, an alignment mechanism (not shown) of the sample handling apparatus **3700** may actuate the first member **3704** and/or the second member **3710** to form the sandwich configuration for the permeabilization step (e.g., bringing the first substrate **3706** and the second substrate **3712** closer to each other and within a threshold distance for the sandwich configuration). The alignment mechanism may be configured to control a speed, an angle, or the like of the sandwich configuration.

In some embodiments, the biological sample (e.g., tissue sample **2302** of **FIG. 23**) may be aligned within the first member **3704** (e.g., via the first retaining mechanism **3708**) prior to closing the first member **3704** such that a desired region of interest of the sample **2302** is aligned with the barcoded array of the second substrate (e.g., the slide **2304**), e.g., when the first and second substrates are aligned in the sandwich configuration. Note that element numbers “**37XX**” refer to elements from **FIGs. 37A** and **37B** and element numbers “**23XX**” refer to elements in **FIG. 23**, wherein “**XX**” is any two digits. Such alignment may be accomplished manually (e.g., by a user) or automatically (e.g., via an automated alignment mechanism). After or before alignment, spacers may be applied to the first substrate **3706** and/or the second substrate **3712** to maintain a minimum spacing between the first substrate **3706** and the second substrate **3712** during sandwiching. In some aspects, the reagent medium (e.g., reagent medium **2305**) may be applied to the first substrate **3706** and/or the second substrate **3712**. The first member **3704** may then close over the second member **3710** and form the sandwich configuration. Analytes (including derivatives such as RTL ligation products and/or analyte capture agents) **2308** may be captured by the capture probes **2306** and may be processed for spatial analysis.

In some embodiments, during the permeabilization step, the image capture device **3720** may capture images of the overlap area between the tissue **2302** and the capture probes **2306**. If more than one first substrates **3706** and/or second substrates **3712** are present within the sample handling apparatus **3700**, the image capture device **3720** may be configured to capture one or more images of one or more overlap areas. Further details on support devices, sample holders, sample handling apparatuses, or systems for implementing a sandwiching process are described in, e.g., PCT Publ. No. WO 2021/0189475 and PCT/US2021/050931, each of which are incorporated by reference in their entirety.

Analytes within a biological sample may be released through disruption (e.g., permeabilization, digestion, etc.) of the biological sample or may be released without *disruption*. Various methods of permeabilizing (e.g., any of the permeabilization reagents

and/or conditions described herein) a biological sample are described herein, including for example including the use of various detergents, buffers, proteases, and/or nucleases for different periods of time and at various temperatures. Additionally, various methods of delivering fluids (e.g., a buffer, a permeabilization solution) to a biological sample are described herein including the use of a substrate holder (e.g., for sandwich assembly, sandwich configuration, as described herein)

Provided herein are methods for delivering a fluid to a biological sample disposed on an area of a first substrate and an array disposed on a second substrate.

In some embodiments and with reference to **FIG. 23**, the sandwich configuration described herein between a first substrate comprising a biological sample (e.g., slide **2303**) and a second substrate comprising a spatially barcoded array (e.g., slide **2304** with barcoded capture probes **2306**) may include a reagent medium (e.g., a liquid reagent medium, e.g., a permeabilization solution **2305** or other target molecule release and capture solution) to fill a gap (e.g., gap **2307**). It may be desirable that the reagent medium be free from air bubbles between the slides to facilitate transfer of target molecules with spatial information. Additionally, air bubbles present between the slides may obscure at least a portion of an image capture of a desired region of interest. Accordingly, it may be desirable to ensure or encourage suppression and/or elimination of air bubbles between the two substrates (e.g., slide **2303** and slide **2304**) during a permeabilization step.

In some aspects, it may be possible to reduce or eliminate bubble formation between the slides using a variety of filling methods and/or closing methods.

Workflows described herein may include contacting a drop of the reagent medium (e.g., liquid reagent medium, e.g., a permeabilization solution **2305**) disposed on a first substrate or a second substrate with at least a portion of the second substrate or first substrate, respectively. In some embodiments, the contacting comprises bringing the two substrates into proximity such that the sample on the first substrate is aligned with the barcode array of capture probes on the second substrate.

In some embodiments, the drop includes permeabilization reagents (e.g., any of the permeabilization reagents described herein). In some embodiments, the rate of permeabilization of the biological sample is modulated by delivering the permeabilization reagents (e.g., a fluid containing permeabilization reagents) at various temperatures.

In the example sandwich maker workflows described herein, the reagent medium (e.g., liquid reagent medium, permeabilization solution **2305**) may fill a gap (e.g., the gap **2307**) between a first substrate (e.g., slide **2303**) and a second substrate (e.g., slide **2304** with

barcoded capture probes **2306**) to warrant or enable transfer of target molecules with spatial information. Described herein are examples of filling methods that may suppress bubble formation and suppress undesirable flow of transcripts and/or target molecules or analytes. Robust fluidics in the sandwich making described herein may preserve spatial information by
 5 reducing or preventing deflection of molecules as they move from the tissue slide to the capture slide.

FIG. 38A shows an exemplary sandwiching process **3800** where a first substrate (e.g., slide **3803**), including a biological sample **3802** (e.g., a tissue section), and a second substrate (e.g., slide **3804** including spatially barcoded capture probes **3806**) are brought into proximity
 10 with one another. As shown in **FIG. 38A** a liquid reagent drop (e.g., permeabilization solution **3805**) is introduced on the second substrate in proximity to the capture probes **3806** and in between the biological sample **3802** and the second substrate (e.g., slide **3804** including spatially barcoded capture probes **3806**). The permeabilization solution **3805** may release analytes that can be captured by the capture probes **3806** of the array. As further
 15 shown, one or more spacers **3810** may be positioned between the first substrate (e.g., slide **3803**) and the second substrate (e.g., slide **3804** including spatially barcoded capture probes **3806**). The one or more spacers **3810** may be configured to maintain a separation distance between the first substrate and the second substrate. While the one or more spacers **3810** is shown as disposed on the second substrate, the spacer may additionally or alternatively be
 20 disposed on the first substrate.

In some embodiments, the one or more spacers **3810** is configured to maintain a separation distance between first and second substrates that is between about 2 microns and 1 mm (e.g., between about 2 microns and 800 microns, between about 2 microns and 700 microns, between about 2 microns and 600 microns, between about 2 microns and 500
 25 microns, between about 2 microns and 400 microns, between about 2 microns and 300 microns, between about 2 microns and 200 microns, between about 2 microns and 100 microns, between about 2 microns and 25 microns, or between about 2 microns and 10 microns), measured in a direction orthogonal to the surface of first substrate that supports the sample. In some instances, the separation distance is about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13,
 30 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 microns. In some embodiments, the separation distance is less than 50 microns. In some embodiments, the separation distance is less than 25 microns. In some embodiments, the separation distance is less than 20 microns. The separation distance may include a distance of at least 2 μm .

FIG. 38B shows a fully formed sandwich configuration creating a chamber **3650** formed from the one or more spacers **3610**, the first substrate (e.g., the slide **303**), and the second substrate (e.g., the slide **304** including spatially barcoded capture probes **306**) in accordance with some example implementations. In the example of **FIG. 38B**, the liquid reagent (e.g., the permeabilization solution **305**) fills the volume of the chamber **3650** and may create a permeabilization buffer that allows analytes, RTL ligation products, and analyte capture agents to diffuse from the biological sample **302** toward the capture probes **306** of the second substrate (e.g., slide **3804**). In some aspects, flow of the permeabilization buffer may deflect transcripts and/or molecules from the biological sample **3802** and may affect diffusive transfer of analytes for spatial analysis. A partially or fully sealed chamber **3850** resulting from the one or more spacers **3810**, the first substrate, and the second substrate may reduce or prevent flow from undesirable convective movement of transcripts and/or molecules over the diffusive transfer from the biological sample **3802** to the capture probes **3806**.

In some instances, the first substrate and the second substrate are arranged in an angled sandwich assembly as described herein. For example, during the sandwiching of the two substrates (e.g., the slide **3803** and the slide **3804**), an angled closure workflow may be used to suppress or eliminate bubble formation.

FIGs. 39A-39C depict a side view and a top view of an exemplary angled closure workflow **3900** for sandwiching a first substrate (e.g., slide **3903**) having a biological sample **3902** and a second substrate (e.g., slide **3904** having capture probes **3906**) in accordance with some example implementations.

FIG. 39A depicts the first substrate (e.g., the slide **3903** including biological sample **3902**) angled over (superior to) the second substrate (e.g., slide **3904**). As shown, a drop of the reagent medium (e.g., permeabilization solution) **3905** is located on the spacer **3910** toward the right-hand side of the side view in **FIG. 39A**. While **FIG. 39A** depicts the reagent medium on the right hand side of side view, it should be understood that such depiction is not meant to be limiting as to the location of the reagent medium on the spacer.

FIG. 39B shows that as the first substrate lowers, and/or as the second substrate rises, the dropped side of the first substrate (e.g., a side of the slide **3903** angled toward the second substrate) may contact the drop of the reagent medium **3905**. The dropped side of the first substrate may urge the reagent medium **3905** toward the opposite direction (e.g., towards an opposite side of the spacer **3910**, towards an opposite side of the first substrate relative to the dropped side). For example, in the side view of **FIG. 39B** the reagent medium **3905** may be urged from right to left as the sandwich is formed.

In some embodiments, the first substrate and/or the second substrate are further moved to achieve an approximately parallel arrangement of the first substrate and the second substrate.

FIG. 39C depicts a full closure of the sandwich between the first substrate and the second substrate with the spacer **3910** contacting both the first substrate and the second substrate and maintaining a separation distance and optionally the approximately parallel arrangement between the two substrates. As shown in the top view of **FIG. 39C**, the spacer **3910** fully encloses and surrounds the biological sample **3902** and the capture probes **3906**, and the spacer **3910** forms the sides of chamber **3950** which holds a volume of the reagent medium **3905**.

It should be understood that while **FIGs. 39A-39C** depict the first substrate (e.g., the slide **3903** including biological sample **3902**) angled over (superior to) the second substrate (e.g., slide **3904**) and the second substrate comprising the spacer **3910**, it should be understood that an exemplary angled closure workflow can include the second substrate angled over (superior to) the first substrate and the first substrate comprising the spacer **3910**.

FIGs. 40A-40E depict an example workflow **4000** for an angled sandwich assembly in accordance with some example implementations. As shown in **FIG. 40A**, a substrate **4012** (e.g., comprising a first substrate such as slide **2303** or a second substrate such as slide **2304** comprising spatially barcoded capture probes **2306**, as shown in **FIG. 23**) may be positioned and placed on a base **4004** (e.g., a first member or a second member of a sample holder disclosed herein) with a side of the substrate **4012** supported by a spring **4015**. The spring **4015** may extend from the base **4004** in a superior direction and may be configured to dispose the substrate **4012** along a plane angled differently than the base **4004**. The angle of the substrate **4012** may be such that a drop of reagent medium **4005** (e.g., drop of liquid reagent medium) placed on the surface of the substrate **4012** (e.g., a surface of a spacer attached to the substrate) will not fall off the surface (e.g., due to gravity). The angle may be determined based on a gravitational force versus any surface force to move the drop away from and off the substrate **4012**.

FIG. 40B depicts a drop **4005** of reagent medium placed on the substrate **4012**. As shown, the drop **4005** is located on the side of the substrate **4012** contacting the spring **4015** and is located in proximity and above (superior to) the spring **4015**.

As shown in **FIG. 40C**, another substrate **4006** may be positioned above (superior to) the substrate **4012** and at an angle substantially parallel with the base **4004**. For example, in cases wherein substrate **4012** is a second substrate disclosed herein (e.g., slide **2304** from

FIG. 23 comprising spatially barcoded capture probes), substrate **4006** may be a first substrate disclosed herein (e.g., slide **2303**). In cases wherein substrate **4012** is a first substrate disclosed herein (e.g., slide **2303**), substrate **4006** may be a second substrate (e.g., slide **2304** comprising spatially barcoded capture probes).

5 In some cases, another base (not shown) supporting substrate **4006** (e.g., a first member or a second member of a sample holder disclosed herein) may be configured to retain substrate **4006** at the angle substantially parallel to the base **4004**.

 As shown in **FIG. 40D**, substrate **4006** may be lowered toward the substrate **4012** such that a dropped side of the substrate **4006** contacts the drop **4005** first. In some aspects, 10 the dropped side of the substrate **4006** may urge the drop **4005** toward the opposite side of the substrate **4006**. In some embodiments, the substrate **4012** may be moved upward toward the substrate **4006** to accomplish the contacting of the dropped side of the substrate **4006** with the drop **4005**.

FIG. 40E depicts a full sandwich closure of the substrate **4006** and the substrate **4012** 15 with the drop of reagent medium **4005** positioned between the two sides. In some aspects and as shown, as the substrate **4006** is lowered onto the drop **4005** and toward the substrate **4012** (and/or as the substrate **4012** is raised up toward the substrate **4006**), the spring **4015** may compress and the substrate **4012** may lower to the base **4004** and become substantially parallel with the substrate **4006**.

20 **FIG. 41A** is a side view of the angled closure workflow **4100** in accordance with some example implementations. **FIG. 41B** is a top view of the angled closure workflow **4100** in accordance with some example implementations. As shown at **4105** and in accordance with **FIGs. 40C-40D**, the drop of reagent medium **4105** is positioned to the side of the substrate **4112** contacting the spring **4115**.

25 At step **4110**, the dropped side of the angled substrate **4106** contacts the drop of reagent medium **4105** first. The contact of the substrate **4106** with the drop of reagent medium **4105** may form a linear or low curvature flow front that fills uniformly with the slides closed.

 At step **4115**, the substrate **4106** is further lowered toward the substrate **4112** (or the 30 substrate **4112** is raised up toward the substrate **4106**) and the dropped side of the substrate **4106** may contact and may urge the liquid reagent toward the side opposite the dropped side and creating a linear or low curvature flow front that may prevent or reduce bubble trapping between the slides. As further shown, the spring **4115** may begin to compress as the substrate **4106** is lowered.

At step **4120**, the drop of reagent medium **4105** fills the gap (e.g., the gap **2307** as shown in **FIG. 23**) between the substrate **4106** and the substrate **4112**. The linear flow front of the liquid reagent may form by squeezing the drop **4105** volume along the contact side of the substrate **4112** and/or the substrate **4106**. Additionally, capillary flow may also contribute to filling the gap area. As further shown in step **4120**, the spring **4115** may be fully compressed such that the substrate **4106**, the substrate **4112**, and the base **4104** are substantially parallel to each other.

In some aspects, an angled closure workflow disclosed herein (e.g., FOREMAN HUB **FIGs. 39A-39C, 40A-40E, and 41A-41B**) may be performed by a sample handling apparatus (e.g., as described in PCT/US2021/050931, which is hereby incorporated by reference in its entirety).

Further details on angled closure workflows, and devices and systems for implementing an angled closure workflow, are described in PCT/US2021/036788 and PCT/US2021/050931, which are hereby incorporated by reference in their entirety.

Additional configurations for reducing or eliminating bubble formation, and/or for reducing unwanted fluid flow, are described in PCT/US2021/036788, which is hereby incorporated by reference in its entirety.

In some embodiments, the reagent medium comprises a permeabilization agent. Suitable agents for this purpose include, but are not limited to, organic solvents (e.g., acetone, ethanol, and methanol), cross-linking agents (e.g., paraformaldehyde), detergents (e.g., saponin, Triton X-100™, Tween-20™, or sodium dodecyl sulfate (SDS)), and enzymes (e.g., trypsin, proteases (e.g., proteinase K)). In some embodiments, the detergent is an anionic detergent (e.g., SDS or N-lauroylsarcosine sodium salt solution). Exemplary permeabilization reagents are described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

In some embodiments, the reagent medium comprises a lysis reagent. Lysis solutions can include ionic surfactants such as, for example, sarkosyl and sodium dodecyl sulfate (SDS). More generally, chemical lysis agents can include, without limitation, organic solvents, chelating agents, detergents, surfactants, and chaotropic agents. Exemplary lysis reagents are described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

In some embodiments, the reagent medium comprises a protease. Exemplary proteases include, e.g., pepsin, trypsin, pepsin, elastase, and proteinase K. Exemplary

proteases are described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

In some embodiments, the reagent medium comprises a detergent. Exemplary detergents include sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100TM, and Tween-20TM. Exemplary detergents are described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

In some embodiments, the reagent medium comprises a nuclease. In some embodiments, the nuclease comprises an RNase. In some embodiments, the RNase is selected from RNase A, RNase C, RNase H, and RNase I. In some embodiments, the reagent medium comprises one or more of sodium dodecyl sulfate (SDS), proteinase K, pepsin, N-lauroylsarcosine, RNase, and a sodium salt thereof.

The sample holder is compatible with a variety of different schemes for contacting the aligned portions of the biological sample and array with the reagent medium to promote analyte capture. In some embodiments, the reagent medium is deposited directly on the second substrate (e.g., forming a reagent medium that includes the permeabilization reagent and the feature array), and/or directly on the first substrate. In some embodiments, the reagent medium is deposited on the first and/or second substrate, and then the first and second substrates aligned in the sandwich configuration such that the reagent medium contacts the aligned portions of the biological sample and array. In some embodiments, the reagent medium is introduced into the gap **2307** while the first and second substrates are aligned in the sandwich configuration.

In certain embodiments a dried permeabilization reagent is applied or formed as a layer on the first substrate or the second substrate or both prior to contacting the sample and the feature array. For example, a reagent can be deposited in solution on the first substrate or the second substrate or both and then dried. Drying methods include, but are not limited to spin coating a thin solution of the reagent and then evaporating a solvent included in the reagent or the reagent itself. Alternatively, in other embodiments, the reagent can be applied in dried form directly onto the first substrate or the second substrate or both. In some embodiments, the coating process can be done in advance of the analytical workflow and the first substrate and the second substrate can be stored pre-coated. Alternatively, the coating process can be done as part of the analytical workflow. In some embodiments, the reagent is a permeabilization reagent. In some embodiments, the reagent is a permeabilization enzyme, a buffer, a detergent, or any combination thereof. In some embodiments, the permeabilization enzyme is pepsin. In some embodiments, the reagent is a dried reagent (e.g., a reagent free

from moisture or liquid). In some instances, the substrate that includes the sample (e.g., a histological tissue section) is hydrated. The sample can be hydrated by contacting the sample with a reagent medium, e.g., a buffer that does not include a permeabilization reagent. In some embodiments, the hydration is performed while the first and second substrates are aligned in a sandwich configuration.

In some instances, the aligned portions of the biological sample and the array are in contact with the reagent medium **2305** for about 1 minute. In some instances, the aligned portions of the biological sample and the array are in contact with the reagent medium **2305** for about 5 minutes. In some instances, the aligned portions of the biological sample and the array are in contact with the reagent medium **1205** in the gap **1207** for about 1 minute, about 5 minutes, about 10 minutes, about 12 minutes, about 15 minutes, about 18 minutes, about 20 minutes, about 25 minutes, about 30 minutes, about 36 minutes, about 45 minutes, or about an hour. In some instances, the aligned portions of the biological sample and the array are in contact with the reagent medium **2305** for about 1-60 minutes. In some instances, the aligned portions of the biological sample and the array are in contact with the reagent medium **2305** for about 30 minutes.

In some embodiments, following initial contact between sample and a permeabilization agent, the permeabilization agent can be removed from contact with sample (e.g., by opening sample holder) before complete permeabilization of sample. For example, in some embodiments, only a portion of sample is permeabilized, and only a portion of the analytes in sample may be captured by feature array. In some instances, the reduced amount of analyte captured and available for detection can be offset by the reduction in lateral diffusion that results from incomplete permeabilization of sample. In general, the spatial resolution of the assay is determined by the extent of analyte diffusion in the transverse direction (i.e., orthogonal to the normal direction to the surface of sample). The larger the distance between the sample on the first substrate and the feature array on the second substrate, the greater the extent of diffusion in the transverse direction, and the concomitant loss of resolution. Analytes liberated from a portion of the sample closest to the feature array have a shorter diffusion path, and therefore do not diffuse as far laterally as analytes from portions of the sample farthest from the feature array. As a result, in some instances, incomplete permeabilization of the sample (by reducing the contact interval between the permeabilization agent and the sample) can be used to maintain adequate spatial resolution in the assay.

In some instances, the device is configured to control a temperature of the first and second substrates. In some embodiments, the temperature of the first and second members is lowered to a first temperature that is below room temperature (e.g., 25 degrees Celsius) (e.g., 20 degrees Celsius or lower, 15 degrees Celsius or lower, 10 degrees Celsius or lower, 5 degrees Celsius or lower, 4 degrees Celsius or lower, 3 degrees Celsius or lower, 2 degrees Celsius or lower, 1 degree Celsius or lower, 0 degrees Celsius or lower, -1 degrees Celsius or lower, -5 degrees Celsius or lower). In some embodiments, the device includes a temperature control system (e.g., heating and cooling conducting coils) to control the temperature of the sample holder. Alternatively, in other embodiments, the temperature of the sample holder is controlled externally (e.g., via refrigeration or a hotplate). In a first step, the second member, set to or at the first temperature, contacts the first substrate, and the first member, set to or at the first temperature, contacts the second substrate, thereby lowering the temperature of the first substrate and the second substrate to a second temperature. In some embodiments, the second temperature is equivalent to the first temperature. In some embodiments, the first temperature is lower than room temperature (e.g., 25 degrees Celsius). In some embodiments, the second temperature ranges from about -10 degrees Celsius to about 4 degrees Celsius. In some embodiments, the second temperature is below room temperature (e.g., 25 degrees Celsius) (e.g., 20 degrees Celsius or lower, 15 degrees Celsius or lower, 10 degrees Celsius or lower, 5 degrees Celsius or lower, 4 degrees Celsius or lower, 3 degrees Celsius or lower, 2 degrees Celsius or lower, 1 degree Celsius or lower, 0 degrees Celsius or lower, -1 degrees Celsius or lower, -5 degrees Celsius or lower).

In an exemplary embodiment, the second substrate is contacted with the permeabilization reagent. In some embodiments, the permeabilization reagent is dried. In some embodiments, the permeabilization reagent is a gel or a liquid. Also in the exemplary embodiment, the biological sample is contacted with buffer. Both the first and second substrates are placed at lower temperature to slow down diffusion and permeabilization efficiency. Alternatively, in some embodiments, the sample can be contacted directly with a liquid permeabilization reagent without inducing an unwanted initiation of permeabilization due to the substrates being at the second temperature. In some embodiments, the low temperature slows down or prevents the initiation of permeabilization. In a second step, keeping the sample holder and substrates at a cold temperature (e.g., at the first or second temperatures) continues to slow down or prevent the permeabilization of the sample. In a third step, the sample holder (and consequently the first and second substrates) is heated up to initiate permeabilization. In some embodiments, the sample holder is heated up to a third

temperature. In some embodiments, the third temperature is above room temperature (e.g., 25 degrees Celsius) (e.g., 30 degrees Celsius or higher, 35 degrees Celsius or higher, 40 degrees Celsius or higher, 50 degrees Celsius or higher, 60 degrees Celsius or higher). In some embodiments, analytes that are released from the permeabilized tissue of the sample
5 diffuse to the surface of the second substrate and are captured on the array (e.g., barcoded probes) of the second substrate. In a fourth step, the first substrate and the second substrate are separated (e.g., pulled apart) and temperature control is stopped.

In some embodiments, where either the first substrate or substrate second (or both) includes wells, a permeabilization solution can be introduced into some or all of the wells,
10 and then the sample and the feature array can be contacted by closing the sample holder to permeabilize the sample. In certain embodiments, a permeabilization solution can be soaked into a hydrogel film that is applied directly to the sample, and/or soaked into features (e.g., beads) of the array. When the first and second substrates are aligned in the sandwich configuration, the permeabilization solution promotes migration of analytes from the sample
15 to the array.

In certain embodiments, different permeabilization agents or different concentrations of permeabilization agents can be infused into array features (e.g., beads) or into a hydrogel layer as described above. By locally varying the nature of the permeabilization reagent(s), the process of analyte capture from the sample can be spatially adjusted.

20 In some instances, migration of the analyte from the biological sample to the second substrate is passive (e.g., via diffusion). Alternatively, in certain embodiments, migration of the analyte from the biological sample is performed actively (e.g., electrophoretic, by applying an electric field to promote migration). In some instances, first and second substrates can include a conductive epoxy. Electrical wires from a power supply can connect
25 to the conductive epoxy, thereby allowing a user to apply a current and generate an electric field between the first and second substrates. In some embodiments, electrophoretic migration results in higher analyte capture efficiency and better spatial fidelity of captured analytes (e.g., on a feature array) than random diffusion onto matched substrates without the application of an electric field (e.g., via manual alignment of the two substrates). Exemplary
30 methods of electrophoretic migration, including those illustrated in **FIGs. 7-11C**, are described in WO 2020/176788, including at FIGs. 13-15, 24A-24B, and 25A-25C, which is hereby incorporated by reference in its entirety.

Loss of spatial resolution can occur when analytes migrate from the sample to the feature array and a component of diffusive migration occurs in the transverse (e.g., lateral)

direction, approximately parallel to the surface of the first substrate on which the sample is mounted. To address this loss of resolution, in some embodiments, a permeabilization agent deposited on or infused into a material with anisotropic diffusion can be applied to the sample or to the feature array. The first and second substrates are aligned by the sample holder and brought into contact. A permeabilization layer that includes a permeabilization solution infused into an anisotropic material is positioned on the second substrate.

In some embodiments, the feature array can be constructed atop a hydrogel layer infused with a permeabilization agent. The hydrogel layer can be mounted on the second substrate, or alternatively, the hydrogel layer itself may function as the second substrate. When the first and second substrates are aligned, the permeabilization agent diffuses out of the hydrogel layer and through or around the feature array to reach the sample. Analytes from the sample migrate to the feature array. Direct contact between the feature array and the sample helps to reduce lateral diffusion of the analytes, mitigating spatial resolution loss that would occur if the diffusive path of the analytes was longer.

Spatial analysis workflows can include a sandwiching process described herein, e.g., a process as described in **FIG. 23**. In some embodiments, the workflow includes provision of the first substrate comprising the biological sample. In some embodiments, the workflow includes, mounting the biological sample onto the first substrate. In some embodiments wherein the biological sample is a tissue sample, the workflow include sectioning of the tissue sample (e.g., cryostat sectioning). In some embodiments, the workflow includes a fixation step. In some instances, the fixation step can include fixation with methanol. In some instances, the fixation step includes formalin (e.g., 2% formalin).

In some embodiments, the biological sample on the first substrate is stained using any of the methods described herein. In some instances, the biological sample is imaged, capturing the stain pattern created during the stain step. In some instances, the biological sample then is destained prior to the sandwiching process.

The biological sample can be stained using known staining techniques, including, without limitation, Can-Grunwald, Giemsa, hematoxylin and eosin (H&E), hematoxylin, Jenner's, Leishman, Masson's trichrome, Papanicolaou, Romanowsky, silver, Sudan, Wright's, and/or Periodic Acid Schiff (PAS) staining techniques. PAS staining is typically performed after formalin or acetone fixation. In some embodiments, the biological sample can be stained using a detectable label (e.g., radioisotopes, fluorophores, chemiluminescent compounds, bioluminescent compounds, and dyes) as described elsewhere herein. In some embodiments, a biological sample is stained using only one type of stain or one technique. In

some embodiments, staining includes biological staining techniques such as H&E staining. In some embodiments, staining includes biological staining using hematoxylin. In some embodiments, staining includes identifying analytes using fluorescently-conjugated antibodies, e.g., by immunofluorescence. In some embodiments, a biological sample is stained using two or more different types of stains, or two or more different staining techniques. For example, a biological sample can be prepared by staining and imaging using one technique (e.g., H&E staining and brightfield imaging), followed by staining and imaging using another technique (e.g., IHC/IF staining and fluorescence microscopy) for the same biological sample. In some instances, a biological sample on the first substrate is stained.

- 10 In some instances, methods for immunofluorescence include a blocking step. The blocking step can include the use of blocking probes to decrease unspecific binding of the antibodies. The blocking step can optionally further include contacting the biological sample with a detergent. In some instances, the detergent can include Triton X-100™. The method can further include an antibody incubation step. In some embodiments, the antibody incubation step effects selective binding of the antibody to antigens of interest in the biological sample. In some embodiments, the antibody is conjugated to an oligonucleotide (e.g., an oligonucleotide-antibody conjugate as described herein). In some embodiments, the antibody is not conjugated to an oligonucleotide. In some embodiments, the method further comprises an antibody staining step. The antibody staining step can include a direct method of immunostaining in which a labelled antibody binds directly to the analyte being stained for. Alternatively, the antibody staining step can include an indirect method of immunostaining in which a first antibody binds to the analyte being stained for, and a second, labelled antibody binds to the first antibody. In some embodiments, the antibody staining step is performed prior to sandwich assembly. In some embodiments wherein an oligonucleotide-antibody conjugate is used in the antibody incubation step, the method does not comprise an antibody staining step.

- 30 In some instances, the methods include imaging the biological sample. In some instances, imaging occurs prior to sandwich assembly. In some instances, imaging occurs while the sandwich configuration is assembled. In some instances, imaging occurs during permeabilization of the biological sample. In some instances, image are captured using high resolution techniques (e.g., having 300 dots per square inch (dpi) or greater). For example, images can be captured using brightfield imaging (e.g., in the setting of hematoxylin or H&E stain), or using fluorescence microscopy to detect adhered labels. In some instances, high resolution images are captured temporally using e.g., confocal microscopy. In some

instances, a low resolution image is captured. A low resolution image (e.g., images that are about 72dpi and normally have an RGB color setting) can be captured at any point of the workflow, including but not limited to staining, destaining, permeabilization, sandwich assembly, and migration of the analytes. In some instances, a low resolution image is taken
5 during permeabilization of the biological sample.

In some embodiments, the location of the one or more analytes in a biological sample are determined by immunofluorescence. In some embodiments, one or more detectable labels (e.g., fluorophore-labeled antibodies) bind to the one or more analytes that are captured (hybridized to) by a probe on the first slide and the location of the one or more analytes is
10 determined by detecting the labels under suitable conditions. In some embodiments, one or more fluorophore-labeled antibodies are used to conjugate to a moiety that associates with a probe on the first slide or the analyte that is hybridized to the probe on the first slide. In some instances, the location(s) of the one or more analytes is determined by imaging the fluorophore-labeled antibodies when the fluorophores are excited by a light of a suitable
15 wavelength. In some embodiments, the location of the one or more analytes in the biological sample is determined by correlating the immunofluorescence data to an image of the biological sample. In some instances, the tissue is imaged throughout the permeabilization step.

In some instances, the biological samples can be destained. In some instances,
20 destaining occurs prior to permeabilization of the biological sample. By way of example only, H&E staining can be destained by washing the sample in HCl. In some instances, the hematoxylin of the H&E stain is destained by washing the sample in HCl. In some embodiments, destaining can include 1, 2, 3, or more washes in HCl. In some embodiments, destaining can include adding HCl to a downstream solution (e.g., permeabilization solution).

Between any of the methods disclosed herein, the methods can include a wash step
25 (e.g., with SSC (e.g., 0.1x SSC)). Wash steps can be performed once or multiple times (e.g., 1x, 2x, 3x, between steps disclosed herein). In some instances, wash steps are performed for about 10 seconds, about 15 seconds, about 20 seconds, about 30 seconds, or about a minute. In some instances, three washes occur for 20 seconds each. In some instances, the wash step
30 occurs before staining the sample, after destaining the sample, before permeabilization the sample, after permeabilization the sample, or any combination thereof.

In some instances, after the sandwiching process the first substrate and the second substrate are separated (e.g., such that they are no longer aligned in a sandwich configuration, also referred to herein as opening the sandwich). In some embodiments, subsequent analysis

(e.g., cDNA synthesis, library preparation, and sequences) can be performed on the captured analytes after the first substrate and the second substrate are separated.

In some embodiments, the process of transferring the ligation product or methylated-adaptor-containing nucleic acid from the first substrate to the second substrate is referred to interchangeably herein as a “sandwich process,” “sandwiching process,” or “sandwiching”. The sandwich process is further described in PCT Patent Application Publication No. WO 2020/123320, PCT/US2021/036788, and PCT/US2021/050931, which are incorporated by reference in its entirety.

10 (e) Use of Multiplexed Sandwich Makers

This disclosure also provides methods, compositions, devices, and systems for using a single capture probe-containing to detect analytes from different biological samples (e.g., tissues) on different slides using serial sandwich processes. Thus, only one capture probe-containing array is necessary for the methods disclosed herein. In this way, as described
15 herein, analytes from different samples or tissues can be captured serially and demultiplexed by sample-specific index sequences.

The methods include generating a connected probe (e.g., a ligation product) in multiple biological samples (i.e., a first sample, a second sample, a third sample, etc.). Generation of a connected probe (e.g., a ligation product) has been described above, and, the
20 same methods are used herein to generate a connected probe (e.g., a ligation product) from analytes that are either protein analytes or nucleic acid (i.e., mRNA) analytes. That is, in some instances, the multiplexed sandwich maker methods disclosed herein can be used to detect protein analytes. In other instances, the multiplexed sandwich maker methods disclosed herein can be used to detect nucleic acid (i.e., mRNA) analytes.

25 The methods, compositions, devices, and systems include utilizing an analyte capture agent in multiple biological samples (i.e., a first sample, a second sample, a third sample, etc.). Using analyte capture agents for spatial detection has been described above, and, the same methods are used herein to use an analyte capture agent to identify analytes in a biological sample. In some embodiments, the multiplexed sandwich maker methods disclosed
30 herein can be used to detect protein analytes.

As discussed below, each connected probe (e.g., a ligation product) that is generated or analyte capture agent includes a sample index sequence, which is a nucleotide sequence that is associated with a particular sample of origin in the multiplex sandwich methods. After generation of each connected probe (e.g., a ligation product) or analyte capture agent, each

sample is serially sandwiched to an array or slide having a plurality of capture probes that can detect and hybridize to a capture probe binding domain from the connected probe (e.g., a ligation product) or analyte capture agent. During the sandwiching process, the indexed connected probe or analyte capture agent actively or passively migrates from the sample to the array for capture by a capture probe. Then the sandwich is opened, and the next sample is sandwiched with the array. In some embodiments, the array is washed prior to sandwiching with the next sample. Additional samples or tissues (e.g., 2 or more) can then be sandwiched with the array or slide having a plurality of capture probes, wherein connected probes (e.g., ligation products) or analyte capture agents from the additional samples or tissues can be transferred to the array in a similar manner. Because each sample includes a unique sample index, the sample of origin for each connected probe (e.g., a ligation product) or analyte capture agent that is captured on the array can be identified. In addition, the location of the connected probe (e.g., a ligation product) can be identified. In some embodiments, the location of the analyte capture agent can be identified. In some instances, the location is identified using fiducial markers on the gene expression slide (i.e., array) so that location of the ligation probe on the array mirrors the location of the sample on the sample slide. Exemplary fiducial markers are described in PCT Patent Application Publication No. WO 2020/123320, which is incorporated by reference in its entirety.

Such methods, compositions, devices, and systems allow for detection of analytes in multiple samples using only one gene expression slide and can be performed on any slide sample described herein. For example, in some instances, the biological sample is a formalin-fixed, paraffin-embedded (FFPE) sample, a frozen sample, or a fresh sample. In some instances, the biological sample is contacted with one or more stains. In some instances, the one or more stains include hematoxylin and eosin. In some instances, cell markers are detected using methods known in the art (e.g., using one or more optical labels such as fluorescent labels, radioactive labels, chemiluminescent labels, calorimetric labels, or colorimetric labels. In some instances, the biological sample is imaged before generating a connected probe (e.g., a ligation product) and before transferring the connected probe (e.g., a ligation product) to the gene expression slide.

The multiplex sandwich methods, compositions, devices, and systems described herein allow for detection of different types of samples and different analytes. For example in some instances, the samples used in the multiplex sandwich methods are from different species. In some instances, the samples used in the multiplex sandwich methods are from the same species but different individuals in the same species. In some instances, the samples

used in the multiplex sandwich methods are from the same individual organism. In some instances, the samples are from different tissues or cell types. In some instances, the samples are from the same tissues or cell types. In some instances, the samples are from the same subject taken at different time points (e.g., before and after treatment). It is appreciated that the samples can be from any source so long as ligated products having sample index sequences unique to each sample are generated.

Multiple samples can be used in the methods described herein. For example, in some instances, at least two samples are used. In some instances, more than two samples (e.g., at least 3, 4, 5, 6, 7, 8, 9, 10, or more) samples are used in the methods disclosed herein. It is appreciated that each sample can be from different sources (e.g., different species, different organisms). In some embodiments, from each sample, the same gene is detected and identified. In some embodiments, for each sample, different genes are detected and identified.

In order to differentiate one sample from another, probe oligonucleotide for each sample in a multiplexed setting can include one or more unique sequences to identify the origin of the connected probe (e.g., a ligation product). In some instances, the unique sequence is a sample index sequence. In some instances, probe oligonucleotides for each sample include one or more (e.g., at least 1, 2, 3, 4, 5, or more) unique sample index sequence to identify the origin of the connected probe (e.g., a ligation product).

In some instances, the sample index is about 5 nucleotides to about 50 nucleotides long (e.g., about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50) nucleotides long. In some embodiments, the sample index is about 5-15 nucleotides long. In some embodiments, the sample index is about 10-12 nucleotides long. Both synthetic and/or naturally-occurring nucleotides can be used to generate a sample index sequence. It is appreciated that any sequence can be designed so long as it is unique among other sample index sequences and optionally that it can be distinguished from any sequence in the genome of the sample.

A sample index sequence can be located anywhere on the connected probe (e.g., a ligation product) so long as it does not affect (1) hybridization of the probe oligonucleotides to the analyte, (2) ligation of the probe oligonucleotides to generate the connected probe (e.g., a ligation product), and (3) hybridization of the capture probe binding domain to the capture probe on an array. For example, in some instances, the sample index sequence can be located on the first probe oligonucleotide (e.g., the left hand probe; i.e., as shown in **FIG. 12A**). In some instances, the sample index is located on the flap of the first probe oligonucleotide that

does not hybridize to the analyte. In some instances, the sample index sequence can be located on the second probe oligonucleotide (e.g., the right hand probe; i.e., as shown in **FIG. 12A**). In some instances, the sample index is located on the flap of the second probe oligonucleotide that does not hybridize to the analyte.

5

(f) Systems and Kits

Also disclosed herein are systems and kits used for any one of the methods disclosed herein. In some instances, the system or kit is used for analyzing an analyte in a biological sample. In some instances, the system or kit includes a support device configured to retain a first substrate and a second substrate, wherein the biological sample is placed on the first substrate, and wherein the second substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises: (i) a spatial barcode and (ii) a capture domain. In some instances, the system or kit includes a first probe oligonucleotide and a second probe oligonucleotide, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the analyte, wherein the second probe oligonucleotide comprises a capture probe binding domain, and wherein the first probe oligonucleotide and the second probe oligonucleotide are capable of being ligated together to form a connected probe. In some instances, the system or kit includes a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence. In some instances, the system or kit further includes a reagent medium comprising a permeabilization agent and optionally an agent for releasing the connected probe. In some instances, the system or kit includes instructions for performing any one of the methods described herein.

In some instances, the permeabilization agent is pepsin or proteinase K. In some instances, the agent for releasing the connected probe is an RNase, optionally wherein the RNase is RNase H.

In some instances, the system or kit further includes an alignment mechanism on the support device to align the first substrate and the second substrate. In some instances, the alignment mechanism comprises a linear actuator and the first substrate comprises a first member and the second substrate comprises a second member. The linear actuator can be configured to move the second member along an axis orthogonal to the plane of the first

member and/or the second member. The linear actuator can be configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member. The linear actuator can be configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec. Finally, in some instances, the linear actuator can be configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.

EXAMPLES

10 **EXAMPLE 1 – Methods for capturing connected probes with sandwich process**

a. Methods of capturing connected probes that hybridize to analytes in a mouse brain and mouse kidney samples using probe pairs that hybridize to a fraction of the mouse transcriptome.

In a non-limiting example, mouse brain and mouse kidney FFPE sections on standard slides (for sandwich conditions) or gene expression (GEx) slides (for non-sandwich control conditions) were deparaffinized, H&E stained, and imaged. Next, the tissue samples were hematoxylin-destained with three HCl solution washes. The sections were then decrosslinked by incubating at 70°C for 1 hour in TE pH 9.0. TE was removed and the tissues were incubation in 1x PBS-Tween for 15 minutes.

Individual probe oligonucleotides (e.g., a first probe oligonucleotide, a second probe oligonucleotide) of probe pairs were hybridized to adjacent sequences of an analyte (e.g., an RNA molecule) in the mouse brain tissue. The RTL probe oligonucleotides were then ligated together, thereby creating a connected probe (e.g., a ligation product) (**FIG. 12A**). The connected probe (e.g., a ligation product) included a capture probe binding domain. The probes were designed to hybridize to part of the mouse transcriptome (e.g., using 5000 total probe pairs) or to hybridize to each transcript in the mouse transcriptome.

After ligation of the RTL probe oligonucleotides, the connected probes were released from the tissue using various conditions: (A) sandwich process with RNase in the reagent medium; (B) sandwich process with RNase + Proteinase K in the reagent medium; or (C) without sandwich methods (i.e., a control). For sandwich conditions A and B, the tissue mounted standard slides were aligned with a GEx slide and permeabilized in the sandwich configuration as described herein (see, e.g., **FIG. 12B**). For the non-sandwich condition C, the tissue mounted on the GEx slides were permeabilized directly on the GEx array.

Following permeabilization, the capture probes were extended, sequencing libraries were prepared and sequenced, and the results were analyzed computationally.

Table 1 and **Table 2** below show results of the mouse brain 5000 probe pair experiment.

5 **[Table 1]**

Conditions	Valid Barcodes	Valid UMIs	Fraction targeted reads usable	Reads Mapped Confidently to Transcriptome	Fraction Reads in Spots Under Tissue	Fraction reads unmapped
RNAse only (30mins) mouse brain	97.5%	100.0%	82.8%	92.0%	91.9%	4.8%
RNAse + Proteinase K (30mins) mouse brain	97.6%	100.0%	86.1%	91.1%	96.4%	5.6%
RNAse + Proteinase K + Sarkosyl (30mins) mouse brain	97.0%	100.0%	83.6%	89.2%	96.0%	6.7%
Control mouse brain (RTL but no sandwich methods)	97.1%	100.0%	63.8%	84.2%	77.9%	11.6%

[Table 2]

Conditions	Median panel genes detected at 260 panel reads per spot	Median panel UMI counts at 250 reads per spot	Median panel genes directed at 1000 panel reads per spot	Median panel UMI counts at 1000 panel reads per spot
RNAse only (30mins) mouse brain	90	123	134	201

RNAse + Proteinase K (30mins) mouse brain	115	169	192	330
RNAse + Proteinase K + Sarkosyl (30mins) mouse brain	47	63	57	79
Control mouse brain (RTL but no sandwich methods)	106	149	192	330

As shown in **Table 1** and **Table 2**, the combination of RNAse + Proteinase K using sandwiching methods resulted in increased useable fraction targeted reads, increased reads in spots under tissues, and increased median panel genes and UMI counts.

5 **FIG. 13** shows results from the mouse brain 5000 probe experiment. As seen in **FIG. 13**, analyte detection and clustering was observed on a large scale (e.g., using 5000 probe pairs). Further, individual gene (e.g., *Hpca*) detection was achieved. Taken together, these data demonstrate the ability to image clusters of genes and even individual genes using these sandwich methods.

10 **Table 3** and **Table 4** below show results of the captured ligated probes on mouse kidney in a setting of using probes designed to hybridize to part of the mouse transcriptome (e.g., using partial (i.e., n=5000) total probe pairs).

[Table 3]

Conditions	Valid Barcodes	Valid UMIs	Fraction targeted reads usable	Reads Mapped Confidently to Transcriptome	Fraction Reads in Spots Under Tissue	Fraction reads unmapped
RNAse only (30mins) mouse kidney	96.3%	100.0%	79.3%	88.3%	91.7%	8.8%

RNAse + Proteinase K (30mins) mouse kidney	97.9%	100.0%	84.8%	93.3%	92.8%	3.1%
RNAse + Proteinase K + Sarkosyl (30mins) mouse kidney	96.9%	100.0%	79.9%	91.6%	89.0%	4.9%
Control mouse kidney	96.4%	100.0%	65.3%	85.9%	78.3%	9.1%

[Table 4]

Conditions	Median panel genes detected at 260 panel reads per spot	Median panel UMI counts at 250 reads per spot	Median panel genes detected at 1000 panel reads per spot	Median panel UMI counts at 1000 panel reads per spot
RNAse only (30mins) mouse kidney	99	167	166	335
RNAse + Proteinase K (30mins) mouse kidney	127	208	283	642
RNAse + Proteinase K + Sarkosyl (30mins) mouse kidney	79	121	129	226

Control	103	164	203	411
mouse				
kidney				

As shown in **Table 3** and **Table 4**, the combination of RNase + Proteinase K using sandwiching methods resulted in increased useable fraction targeted reads, increased reads in spots under tissues, and increased median panel genes and UMI counts.

5 **FIG. 14** shows results from the mouse kidney 5000 probe experiment. As seen in **FIG. 14**, analyte detection and clustering was observed on a large scale (e.g., using 5000 probe pairs). Further, individual gene (e.g., *Gatm*) detection was achieved. Taken together, these data demonstrate the ability to image clusters of genes and even individual genes using these sandwich methods.

10 In addition, as shown in **Table 5**, other FFPE samples from human brain, human lung, mouse spleen, mouse thymus, and mouse testes were tested using either 3000 probe pairs for human samples or 5000 probe pairs for mouse samples. The samples were sandwiched, and probes were released using a combination of RNase and Proteinase K.

[Table 5]

Sample	Fraction Reads in Spots under Tissue	Median Panel Genes Detected at 1000 Panel Reads per Spot	Median Panel UMI Counts at 1000 Panel Reads per Spot
Human Brain (Control – no sandwich)	97.1%	247	315
Human Brain – RNase + Proteinase K (30mins)	75.6%	322	564
Human Lung (Control – no sandwich)	42.6%	569	1494
Human Lung – RNase + Proteinase K (30mins)	95.3%	862	1254
Mouse Spleen (Control – no sandwich)	74.1%	264	799
Mouse Spleen – RNase + Proteinase K (30mins)	90.9%	317	919
Mouse Thymus (Control – no sandwich)	75.8%	293	639

Mouse Thymus – RNase + Proteinase K (30mins)	76.1%	359	918
Mouse Testes (Control – no sandwich)	93.0%	385	921
Mouse Testes – RNase + Proteinase K (30mins)	98.6%	352	941

b. Methods of capturing connected probes that hybridize to analytes in a mouse brain samples using probe pairs that hybridize to probe pairs designed to cover the entire mouse transcriptome.

5

The methods of Example 1a were repeated in a second set of mouse brain samples, except that probe pairs that cover the entire mouse transcriptome were used (21,000 probe pairs). **Table 6** shows results of capture of connected probes using probe pairs designed to cover the entire mouse transcriptome.

10 **[Table 6]**

Sample	Valid Barcodes	Valid UMIs	Fraction Targeted reads usable	Reads mapped confidently to transcriptome	Fraction Reads in Spots under Tissue	Fraction Reads Unmapped	Median Panel Genes detected at 10000 panel reads per spot	Median Panel UMI counts at 10000 panel reads per spot
Control mouse brain (RTL but no sandwich methods)	97.2%	100.0%	57.9%	67.5%	88.2%	2.6%	2542	4746
RNase + Proteinase K (30mins)	97.2%	100.0%	88.8%	94.7%	96.3%	1.8%	3484	7753

mouse								
brain								

As shown in **FIG. 15**, analyte detection was observed in both sandwich and non-sandwich methods, demonstrating the ability to detect analytes in both settings. Further, individual gene (e.g., Rbfox3; Hpca) detection was achieved. Taken together, these data demonstrate the ability to image clusters of genes and even individual genes using these sandwich methods.

Table 7 shows results of capture of connected probes using probe pairs designed to cover human transcriptome (3000 probe panel) or mouse transcriptome (5000 probe panel) with different types of tissue samples.

10 **[Table 7]**

Tissue type	Sample	Fraction Reads in Spots Under Tissue	Median panel genes detected at 1000 panel reads per spot	Median panel UMI counts at 1000 panel reads per spot
Human brain	Control – no sandwich	97.1%	247	315
	RNase + Proteinase K	75.6%	322	564
Human lung	Control – no sandwich	42.6%	569	1494
	RNase + Proteinase K	95.3%	862	1254
Mouse spleen	Control – no sandwich	74.1%	264	799
	RNase + Proteinase K	90.9%	317	919
Mouse thymus	Control – no sandwich	75.8%	293	639
	RNase + Proteinase K	76.1%	359	918
Mouse testes	Control – no sandwich	93.0%	385	921
	RNase + Proteinase K	98.6%	352	941

EXAMPLE 2 – Methods for spatial analysis of RNA and protein with sandwich process

In a non-limiting example, methods for spatial analysis of RNA and protein were performed. In brief, mouse brain FFPE sections on standard slides (for sandwich conditions)

- or gene expression (GEx) slides (for non-sandwich control conditions) were decrosslinked according to the steps in Example 1a. Probe hybridization using mouse-specific probe sets to the mouse partial transcriptome (n = 5000 probe pairs) was performed, followed by RTL probe ligation as described in Example 1. The sections were then incubated with totalseq B
- 5 oligo-tagged antibodies for NeuN and Gfap, followed by washing. The antibodies were tagged with oligonucleotides that have a sequence complementary to a capture probe capture domain of a GEx slide and a barcode sequence that uniquely identifies the antibody. The connected probes and antibody oligonucleotide tags were released from the tissue under sandwich conditions as described herein or under non-sandwich conditions. For both
- 10 conditions, the reagent medium included RNase and Proteinase K. For the sandwich conditions, the tissue-mounted slides were aligned with a GEx slide and permeabilized in the sandwich configuration as described herein (see, e.g., FIG. 16B). For the non-sandwich condition, the tissue mounted on the GEx slide was permeabilized directly on the GEx array. Following permeabilization, the capture probes were extended, sequencing libraries were
- 15 prepared and sequenced, and the results were analyzed computationally.

Table 8 shows mRNA analysis results. **Table 9** shows antibody analysis results.

[Table 8]

Sample	Valid Barcode s	Valid UMIs	Fraction Targeted reads usable	Reads mapped confidently to transcriptome	Fraction Reads in Spots under Tissue	Fraction Reads Unmapped	Median Panel Genes detected at 10000 panel reads per spot	Median Panel UMI counts at 10000 panel reads per spot
Control mouse brain (RTL but no sandwich methods)	98.9%	100.0 %	69.9	73.5%	96.2%	12.2%	216	366
RNAse + Proteinase	99.0%	100.0 %	91.3%	97.1%	95.0%	2.2%	391	814

e K (30mins) mouse brain								
-----------------------------------	--	--	--	--	--	--	--	--

[Table 9]

Sample	Valid Barcodes	Valid UMIs	Fraction Targeted reads usable	Reads mapped confidently to transcriptome
Control mouse brain (RTL but no sandwich methods)	88%	61%	12%	70%
RNAse + Proteinase K (30mins) mouse brain	74%	68%	26%	93%

As shown in **FIG. 18A-18B**, mRNA and protein analyte detection was observed in both sandwich and non-sandwich methods, demonstrating the ability to detect analytes in both settings. Further, individual gene (e.g., RbFox3/NeuN; Gfap) detection was achieved. Taken together, these data demonstrate the ability to image clusters of protein or mRNA analytes and even individual protein or mRNA analytes using these sandwich methods.

10 **EXAMPLE 3 – Methods for spatial transcriptomics analysis utilizing sandwich process**

In a non-limiting example, the sandwich process can be utilized for a downstream analytical step in spatial transcriptomics analysis workflows as described herein.

FIG. 19A shows an exemplary workflow of spatial analysis assays using fresh frozen tissue samples. For example, a tissue section is (a) fixed and stained (e.g., hematoxylin and eosin staining, fluorescent antibody staining); (b) imaged to evaluate the quality of the antibody staining; (c) destaining and unmounting the tissue section; (d) permeabilizing the tissue section with Proteinase K and sarkosyl prior to performing the sandwich process; and (e) performing a reverse transcription protocol and generating an analyte library.

FIG. 19B shows an exemplary workflow of spatial analysis assays using formalin-fixation and paraffin-embedded (FFPE) tissue samples. For example, a FFPE tissue section is (a) deparaffinized, wherein the paraffin-embedding material is removed; (b) fixed and stained (e.g., hematoxylin and eosin staining, fluorescent antibody staining); (b) imaged to evaluate the quality of the antibody staining; (c) destaining and unmounting the tissue section; (d) permeabilizing the tissue section with RNase, Proteinase K and sarkosyl prior to performing the sandwich process; and (e) performing a reverse transcription protocol and generating an analyte library.

10 **EXAMPLE 4 – Methods for spatial transcriptomics analysis utilizing serial sandwich process**

In a non-limiting example, the sandwich process can be repeated using serial biological sections with transfer to the same substrate comprising a plurality of capture probes.

15 To demonstrate the ability to capture analytes from multiple biological samples, connected probes were generated on separate slides, with each slide comprising a different biological sample. Briefly, in Biological Sample 1, a mouse sample was used. In Biological Sample 2, a human sample was used. In particular, mouse thymus or mouse testes biological samples were used as Biological Sample 1, and human lung or human breast cancer samples were used for Biological Sample 2. Probes were designed to hybridize to mouse-specific mRNA analytes or human-specific analytes. In some embodiments, partial transcriptome probe panels were used. In some embodiments, mouse transcriptome probe panels were used. In some embodiments, human transcriptome probe pairs were used.

25 Individual probe oligonucleotides (e.g., a first probe oligonucleotide, a second probe oligonucleotide) in a probe pair hybridized to adjacent sequences of an analyte (e.g., an RNA molecule) in each biological sample. The RTL probe oligonucleotides were then ligated together, thereby creating a connected probe. After ligation of the RTL probe oligonucleotides, RNase H was added to release the ligated product from the analyte, and proteinase K was added to the samples to permeabilize the biological samples.

30 During the permeabilization step, each sample (i.e., Biological Sample 1 and Biological Sample 2) was serially added to the second substrate (i.e., the substrate comprising capture probes that capture the connected probes. The poly(A) sequence on the connected probe hybridized to the poly(T) sequence of the capture probe domain. After hybridization of the connected probes from Biological Sample 1, Biological Sample 1 was

removed from the capture probe-containing substrate. The capture-probe-containing substrate was washed briefly with 1x SSC, dried quickly, and this sandwich-transfer method was repeated for Biological Sample 2.

As shown in **FIG 20**, human RTL connected probes were detected, demonstrating that serial transfer of multiple probes can be achieved. As controls, biological samples (i.e., mouse thymus, mouse testes, and human breast cancer) in which no sandwiching methods were used (**FIG. 21A**), and biological samples in which a single sandwiching transfer was used (**FIG. 21B**). In each control, as shown in the heat maps and clustering images (**FIGs. 21A-21B**), connected probes were captured and detected on the array.

FIGs. 22A-22B show heat maps for each sample type, demonstrating the location and saturation of the sequenced reads in the second substrate. **FIGs. 22A-22B** also show the clustering and spatiality of reads under the tissue in an overlay utilizing the loupe file fiducials. It is assumed that regions of the tissue that have similar gene expression will be clustered together and represented by the same color in these images. The overlay of the clustered reads suggest that reads from both mouse and human tissues were captured. The pattern of the reads under tissue seen in the heat map are represented for each tissue type in the overlaid clustered reads.

The results demonstrate the ability to serially capture analytes from different biological samples on the same array. The results also demonstrate the ability to demultiplex the samples computationally.

EXAMPLE 5 – Efficient analyte capture from slide-mounted fresh frozen mouse brain sections onto spatial array slides

In a non-limiting example, analyte capture onto spatially barcoded arrays and subsequent sequencing was demonstrated under sandwich and non-sandwich conditions. For the test (sandwiching) condition, archived tissue-mounted standard glass slides containing hematoxylin/eosin stained fresh frozen mouse brain sections were used. For control (non-sandwich) condition, gene expression (GEx) array slides with hematoxylin/eosin stained fresh frozen mouse brain sections mounted directly onto the array area were used. Under both conditions, tissue sections were subjected to a hematoxylin destaining step. Slides processed according to the “sandwiching” condition were briefly dried at 37°C, then mounted in an instrument along with a GEx slide and a permeabilization buffer comprising sarkosyl and proteinase K. Upon sandwich closure in the instrument, the tissue sections were permeabilized for 1 minute. For the tissue-mounted GEx slides processed according to the

non-sandwich condition, sections were permeabilized for 5 minutes using the same permeabilization buffer without sandwiching. For both conditions, following permeabilization, captured poly A-containing mRNA transcripts on the GEx slides were reverse transcribed into cDNA, followed by standard sequencing library preparation and sequencing.

Results depicting median genes per spot and median UMI counts per spot are shown in **FIG. 24**.

Visual heat map results showing Log10 UMIs are shown in **FIG. 25**. Spatial patterns of the Log10 UMI counts were similar across the sandwich and non-sandwich conditions.

Spatial clustering analysis (top row **2605**) and analysis of hippocampal transcript Hpcr (bottom row **2610**) are depicted in **FIG. 26**. Spatial patterns were comparable across the sandwich and non-sandwich conditions.

EXAMPLE 6 – Efficient analyte capture from slide-mounted FFPE human brain and human breast sections onto spatial array slides

In a non-limiting example, human brain and human breast FFPE sections on standard slides (for sandwich conditions) or gene expression (GEx) slides (for non-sandwich control conditions) were deparaffinized, H&E stained, and imaged. Next, the tissue samples were hematoxylin-destained with three HCl solution washes and then decrosslinked according to the steps in Example 1a.

Probe hybridization was performed, followed by RTL probe ligation as described in Example 1. The sections were then incubated with totalseq B oligo-tagged antibodies for NeuN and Gfap, followed by washing. The antibodies were tagged with oligonucleotides that have a sequence complementary to a capture probe capture domain of a GEx slide and a barcode sequence that uniquely identifies the antibody. The connected probes and antibody oligonucleotide tags were released from the tissue under sandwich conditions as described herein or under non-sandwich conditions. For both conditions, the reagent medium included RNase and Proteinase K. For the sandwich conditions, the tissue-mounted slides were aligned with a GEx slide and permeabilized in the sandwich configuration as described herein (see, e.g., **FIG. 16B**). For the non-sandwich condition, the tissue mounted on the GEx slide was permeabilized directly on the GEx array. Following permeabilization, the capture probes were extended, sequencing libraries were prepared and sequenced, and the results were analyzed computationally.

Visual heat map results showing Log10 UMIs for human brain and human breast tissues are shown in **FIG. 27A** and **27B**, respectively. Spatial patterns of the Log10 UMI counts were similar across the sandwich and non-sandwich conditions. Spatial clustering analysis of the human brain and human breast tissues are depicted in **FIG. 27A** and **27B**, respectively. These data demonstrate that spatial clustering was comparable across the sandwich and non-sandwich conditions.

EXAMPLE 7 – Efficient multiplexed analyte capture from slide-mounted mouse spleen, brain, head, and torso sections onto spatial array slides

In a non-limiting example, FFPE mouse spleen samples, FFPE mouse brain samples, FFPE mouse embryo torso samples, and FFPE mouse embryo head sections samples were placed onto standard slides (for sandwich conditions) or spatial expression (GEx) slides (as non-sandwich conditions). GEx slides included an array of spatially barcoded capture probes. Briefly, tissues were sectioned and were mounted on slides and dried overnight in a desiccator. The following day, the tissues were heated to 60°C, followed by deparaffinization and rehydration. H&E staining was performed and tissues were imaged. Tissues were destained using HCl and decrosslinked for 1 hour in citrate buffer (pH 6.0) at 95°C. After decrosslinking, tissues were incubated overnight with whole mouse transcriptome RTL (templated ligation) probe sets at 50°C. The following day, tissues were washed to remove un-hybridized probes, then treated with ligase to ligate together the RTL probes. After another wash step, the tissues were blocked with antibody blocking buffer. Tissues were incubated overnight with a library of conjugated antibodies (e.g., a library comprising a plurality of analyte capture agents, each comprising an antigen-specific antibody conjugated to an oligonucleotide). The following day, tissues were subjected to sandwiching or non-sandwich conditions as follows.

Tissues placed on standard slides for the sandwiching conditions were washed with PBS-T, subjected to an eosin retain, and washed with SSC. These tissues were subjected to sandwiching conditions described herein. Briefly, the tissue slides were mounted in an instrument along with a GEx slide and a reagent solution including RNase and proteinase K in buffer. Upon sandwich closure in the instrument, the tissue sections were permeabilized for 30 minutes, allowing the ligation products and oligonucleotides from analyte capture agents to migrate to the GEx slide for capture by the capture probes. Following permeabilization and capture, the GEx slides were removed from the instrument.

Tissues placed on the GEx slides for non-sandwiching conditions were washed with PBST, and washed with SSC. The tissues were subjected to a 30 minute probe release step with RNase, followed by a 1 hour permeabilization step with a permeabilization buffer including Proteinase K and detergent. Accordingly, the ligation products and analyte capture agents were captured by the capture probes of the GEx slide.

Regardless of condition, GEx slides were washed twice with 2x SSC and subjected to probe extension, denaturation, and pre-amplification followed by amplification and sequencing of the templated ligation and analyte capture agent libraries.

After sequencing, the quality, sensitivity, and detection under each conditions (sandwiching and non-sandwiching conditions) were evaluated. As shown in **Table 10**, the quality, sensitivity, and detection of globally-detected transcripts (i.e., mRNA) and proteins were comparable across the sandwich and non-sandwich conditions.

[Table 10]

	Metric	Sandwiching Conditions	Non-Sandwiching Conditions
Templated Ligation Quality	Valid barcodes	99.00%	98.90%
	Fraction reads on target	85.60%	82.10%
	Fraction reads usable	79.80%	78.70%
	Fraction reads in spots	81.60%	81.00%
	Fraction reads unmapped	0.90%	1.00%
Templated Ligation Sensitivity	Median genes (20K prps)	4856	4705
	Median UMIs (20K prps)	16966	15156
Protein Detection using Analyte Capture Agents	Fraction reads usable	75.20%	67.10%
	Fraction reads in spot	78.70%	70.10%
	Fraction unknown	3.70%	3.60%
	Median UMIs per spot (5K reads usable per spot)	4632	4114
	Correlation of selected Templated Ligation/Analyte Capture Agent	0.77	0.76

Images were generated to evaluate overlap of gene expression and gene product (e.g., protein) profiles in mouse spleen tissue and mouse brain tissue for individual biomarkers. For instance, CD45, which is expressed in both spleen and brain, was evaluated. As shown in **FIG. 28A** (sandwiching conditions) and **FIG. 28C** (non-sandwiching conditions), protein tyrosine phosphatase receptor type C (Ptprc; e.g., Ensembl: ENSMUSG00000026395) mRNA expression was detected in a spleen tissue sample. Further, gene product (e.g., protein) CD45R was also detected both in sandwiching conditions (**FIG. 28B**) and in non-sandwiching conditions (**FIG. 28D**). CD45R is the protein name of Ptprc, and it was determined whether there was overlap of mRNA expression of Ptprc and protein expression of CD45R. As shown in **FIGs. 28A-28B** and **FIGs. 28C-28D**, both sandwiching (77% correlation) and non-sandwiching conditions (76% correlation), respectively, demonstrated overlap of transcript and protein, indicating that transcript (i.e., mRNA) and protein detection (1) was identified in similar areas of the tissues, and the (2) was comparable across the sandwich and non-sandwich conditions in mouse spleen samples.

Mouse brain samples were also evaluated. In particular, the quality, sensitivity, and detection under each conditions (sandwiching and non-sandwiching conditions) were evaluated globally in mouse brain samples. As shown in **Table 11**, the quality, sensitivity, and detection of globally-detected transcripts (i.e., mRNA) and proteins were comparable across the sandwich and non-sandwich conditions in mouse brain samples.

[Table 11]

	Metric	Sandwiching Conditions	Non-Sandwiching Conditions
Templated Ligation Quality	Valid barcodes	99.00%	98.90%
	Fraction reads on target	92.00%	87.30%
	Fraction reads usable	88.80%	72.30%
	Fraction reads in spots	91.80%	79.10%
	Fraction reads unmapped	1.10%	1.90%
Templated Ligation Sensitivity	Median genes (10K prps)	4405	3185
	Median UMIs (10K prps)	9386	5553
Protein Detection using Analyte	Fraction reads usable	80.70%	62.30%
	Fraction reads in spot	84.40%	65.10%

Capture Agents	Fraction unknown	3.70%	3.70%
	Median UMIs per spot (5K reads usable per spot)	3232	3221
	Correlation of selected Templated Ligation/Analyte Capture Agent	0.63	0.63

Similar to the mouse spleen sample images described above, individual gene expression and protein products were evaluated in mouse brain samples. As shown in **FIG. 29A** (sandwiching conditions) and **FIG. 29C** (non-sandwiching conditions), tyrosine hydroxylase (Th; e.g., Ensembl: ENSMUSG00000000214) mRNA expression was detected in brain. Further, the gene product (e.g., protein) TH was also detected in sandwiching conditions (**FIG. 29B**) and non-sandwiching conditions (**FIG. 29D**). Since TH is the protein made by Th, it was determined whether there was overlap of mRNA expression of Th and protein expression of TH. As shown in **FIGs. 29A-29B** and **FIGs. 29C-29D**, both sandwiching (63% correlation) and non-sandwiching conditions (63% correlation), respectively, saw overlap of transcript and protein, demonstrating that transcript (i.e., mRNA) and protein detection was comparable across the sandwich and non-sandwich conditions in mouse brain samples.

Additional experiments using the same methods (i.e., testing sandwiching conditions versus non-sandwiching control conditions while detecting both RNA and protein) were performed on embryo torso and head sections. In addition to conditions in which both RNA and protein were detected, a third condition was included as a control. This third condition (Conditions 3 in **Tables 12** and **13**) detected the presence and abundance of only RNA. In each condition, RNA was detected using templated ligation as previously described herein. For Conditions 1 and 2 as shown in **Tables 12** and **13** below, protein was also detected using analyte capture agent methods described herein. The quality, sensitivity, and detection under each condition (sandwiching versus non-sandwiching conditions) were evaluated in mouse embryo torso and head/upper torso samples. As shown in **Tables 12** and **13**, the quality, sensitivity, and detection of globally-detected transcripts (i.e., mRNA) and proteins were comparable across the sandwich and non-sandwich conditions in mouse embryo torso and head samples. Further, the quality, sensitivity, and detection of globally-detected transcripts (i.e., mRNA) was roughly the same between Conditions 1 and 3, demonstrating that both protein capture and sandwiching methods did not interfere with RNA capture using templated ligation methods.

[Table 12] – Mouse Embryo Torso Sample Data

	Metric	Condition 1: Sandwiching Conditions Detecting both RNA and Protein	Condition 2: Non-Sandwiching Conditions Detecting both RNA and Protein	Condition 3: Non-Sandwiching Conditions Detecting RNA
Templated Ligation Quality	Valid barcodes	98.9%	99.0%	98.5%
	Fraction reads on target	88.5%	88.5%	87.9%
	Fraction reads usable	81.5%	77.7%	84.8%
	Fraction reads in spots	84.0%	79.9%	88.4%
	Fraction reads unmapped	1.0%	1.0%	1.4%
Templated Ligation Sensitivity	Median genes (10K prps)	4144	4356	3562
	Median UMIs (10K prps)	8767	8176	6121
Protein Detection using Analyte Capture Agents	Fraction reads usable	77.3%	72.6%	-
	Fraction reads in spot	80.8%	75.9%	-
	Fraction unknown	3.7%	3.7%	-
	Median UMIs per spot (5K reads usable per spot)	3358	3264	-
	Correlation of selected Templated Ligation/Analyte Capture Agent	0.90	0.80	-

[Table 13] – Mouse Embryo Head/Upper Torso Sample Data

	Metric	Condition 1: Sandwiching Conditions Detecting both RNA and Protein	Condition 2: Non-Sandwiching Conditions Detecting both RNA and Protein	Condition 3: Non-Sandwiching Conditions Detecting RNA
Templated Ligation Quality	Valid barcodes	99.0%	99.0%	98.5%
	Fraction reads on target	89.1%	89.4%	89.0%
	Fraction reads usable	86.7%	82.9%	82.2%
	Fraction reads in spots	89.5%	85.2%	85.2%
	Fraction reads unmapped	1.0%	1.0%	1.5%

Templated Ligation Sensitivity	Median genes (10K prps)	4400	4708	3562
	Median UMIs (10K prps)	9077	8431	6571
Protein Detection using Analyte Capture Agents	Fraction reads usable	84.3%	77.9%	-
	Fraction reads in spot	88.2%	81.5%	-
	Fraction unknown	3.7%	3.8%	-
	Median UMIs per spot (5K reads usable per spot)	3358	3349	-
	Correlation of selected Templated Ligation/Analyte Capture Agent	0.72	0.59	-

In addition to performing global expression analysis on each group, individual targets were analyzed for the location and abundance of transcript (i.e., mRNA) and protein of a single target in the mouse embryo torso and head/upper torso samples. **FIGs. 30A-30B** show mRNA (**FIG. 30A**) and protein (**FIG. 30B**) detection of lymphocyte antigen 76 (Ter119) (e.g., NCBI Gene ID: 104231), of mouse embryo torso samples in Condition 1 (i.e., in sandwiching conditions). **FIGs. 30C-30D** show mRNA (**FIG. 30C**) and protein (**FIG. 30D**) detection of Ter119 of mouse embryo torso samples in Condition 2 (i.e., in non-sandwiching conditions). **FIGs. 31A-31B** show mRNA (**FIG. 31A**) and protein (**FIG. 31B**) detection of Ter119 of mouse embryo head samples in Condition 1 (i.e., in sandwiching conditions). **FIGs. 31C-31D** show mRNA (**FIG. 31C**) and protein (**FIG. 31D**) detection of Ter119 of mouse embryo head/upper torso samples in Condition 2 (i.e., in non-sandwiching conditions). As shown in **FIGs. 30A-31D**, Ter119 mRNA and protein was readily detected with considerable overlap of mRNA and protein detection in both sandwiching conditions and non-sandwiching conditions, demonstrating the adaptability and reproducibility of the methods regardless of condition.

Additional single biomarkers were analyzed in the mouse embryo torso and head samples. As shown in **FIGs. 32A-32I**, metallophosphoesterase domain containing 1 (Mpped1; e.g., Ensembl: ENSMUSG00000041708) mRNA and protein was analyzed. Mpped1 was readily detected in the brain region of mouse embryo head samples. **FIG. 32A** shows detection of Mpped1 RNA in a mouse embryo head sample using sandwiching conditions (Condition 1 of **Tables 12 and 13**). **FIG. 32B** shows detection of Mpped1 protein in a mouse embryo head sample using sandwiching conditions (Condition 1 of **Tables 12 and 13**). However, Mpped1 mRNA was not readily detected in a mouse embryo torso sample.

Consistent with these observations, Mpped1 has metallophosphoesterase activity, which could have a role in brain development. Consistent with the sandwiching conditions data, Mpped1 RNA and protein was detected in non-sandwiching conditions in the head (**FIG. 32D** (mRNA) and **FIG. 32E** (protein)) but not in the torso (**FIG. 32F** (mRNA) and **FIG. 32G** (protein)). Similarly, in non-sandwiching conditions in which only RNA was detected, Mpped1 RNA was detected in the head (**FIG. 32H**) but not in the torso (**FIG. 32I**). As such, regardless of conditions, both gene and protein expression, down to the single biomarker level, can be detected concurrently in the same sample.

Four additional biomarkers—troponin C1, slow skeletal and cardiac type (Tnncl, e.g., Ensembl: ENSMUSG00000091898); fibroblast growth factor 15 (Fgf15, e.g., Ensembl: ENSMUSG00000031073); epiphy can (Epyc, e.g., Ensembl: ENSMUSG00000019936); and serine (or cysteine) peptidase inhibitor, clade A, member 1E (Serpinal e, e.g., Ensembl: ENSMUSG00000072849)—were examined in head/upper torso and torso mouse embryo samples under Conditions 1, 2, and 3 from **Tables 12** and **13**. See **FIGs. 33A-36I**. Tnncl is involved in muscle contraction regulation and its expression was readily detected in all samples under each Condition. See **FIGs. 33A-33I**. Fgf15 functions in retinal neurogenesis and as a cell fate determination factor. Indeed, its expression was found in the eye of the embryo in each head sample, but was not detected in the torso. See **FIGs. 34A-34I**. Epyc functions in bone formation and in cartilage structure and was detected in each sample (head and torso). See **FIGs. 35A-35I**. Finally, Serpinal e is active in the liver as it functions in alpha-1 antitrypsin protein production. As shown in **FIGs. 36A-36I**, Serpinal e was detected in the torso of mouse embryos but was not detected in the head/upper torso samples. Consistent among each of these biomarker images was that sandwiching methods readily detected individual mRNA or protein biomarkers compared to non-sandwiching methods. As such, using sandwiching or non-sandwiching conditions, both gene and protein expression, down to the single biomarker level, can be detected concurrently across multiple tissue types using the methods described herein.

WHAT IS CLAIMED IS:

1. A method for analyzing an analyte in a biological sample mounted on a first substrate, the method comprising:
 - (a) hybridizing a first probe oligonucleotide and a second probe oligonucleotide to the analyte, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the analyte, and wherein the second probe oligonucleotide comprises a capture probe binding domain;
 - (b) coupling the first probe oligonucleotide and the second probe oligonucleotide, thereby generating a connected probe;
 - (c) aligning the first substrate with a second substrate comprising an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises: (i) a spatial barcode and (ii) a capture domain;
 - (d) when the biological sample is aligned with at least a portion of the array, (i) releasing the connected probe from the analyte and (ii) migrating the connected probe from the biological sample to the array; and
 - (e) hybridizing the connected probe to the capture domain.
2. The method of claim 1, wherein the first probe oligonucleotide and the second probe oligonucleotide are on a contiguous nucleic acid sequence.
3. The method of claim 2, wherein the first probe oligonucleotide is on the 3' end of the contiguous nucleic acid sequence.
4. The method of claims 2 or 3, wherein the second probe oligonucleotide is on the 5' end of the contiguous nucleic acid sequence.
5. The method of any one of claims 1-4, wherein the adjacent sequences abut one another.
6. The method of any one of claims 1-4, wherein the adjacent sequences are at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more nucleotides away from one another.

7. The method of claim 6, further comprising generating an extended first probe oligonucleotide, wherein the extended first probe oligonucleotide comprises a sequence complementary to a sequence between the sequence hybridized to the first probe oligonucleotide and the sequence hybridized to the second probe oligonucleotide.
- 5
8. The method of claim 6, further comprising generating an extended second probe oligonucleotide using a polymerase, wherein the extended second probe oligonucleotide comprises a sequence complementary to a sequence between the sequence hybridized to the first probe oligonucleotide and the sequence hybridized to the second probe oligonucleotide.
- 10
9. The method of any one of the preceding claims, further comprising hybridizing a third probe oligonucleotide to the first probe oligonucleotide and the second probe oligonucleotide.
- 15
10. The method of claim 9, wherein the third probe oligonucleotide comprises:
- (i) a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, or 100% complementary to a portion of the first probe oligonucleotide that hybridizes to the third probe oligonucleotide; and
 - 20 (ii) a sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%, or 100% complementary to a portion of the second probe oligonucleotide that hybridizes to the third probe oligonucleotide.
- 25
11. The method of any one of the preceding claims, wherein the coupling the first probe oligonucleotide and the second probe oligonucleotide comprises ligating via a ligase the first probe oligonucleotide and the second probe oligonucleotide.
- 30
12. The method of any one of claims 7-11, wherein the coupling the first probe oligonucleotide and the second probe oligonucleotide comprises ligating via a ligase:
- (i) the first probe oligonucleotide and the extended second probe oligonucleotide;
 - or
 - (ii) the extended first probe oligonucleotide and the second probe oligonucleotide.

13. The method of claim 11 or 12, wherein the ligase is selected from a splintR ligase, a single stranded DNA ligase, or a T4 DNA ligase.
- 5 14. The method of any one of the preceding claims, further comprising amplifying the connected probe prior to the releasing step, optionally wherein the amplifying comprises rolling circle amplification.
15. The method of claim 14, wherein the amplifying comprises rolling circle amplification.
- 10 16. The method of any one of claims 1-15, wherein the releasing step (d) comprises contacting the biological sample with a reagent medium comprising a permeabilization agent and an agent for releasing the connected probe, thereby permeabilizing the biological sample and releasing the connected probe from the analyte.
- 15 17. The method of claim 16, wherein the aligning comprises:
- (i) mounting the first substrate on a first member of a support device, the first member configured to retain the first substrate;
 - (ii) mounting the second substrate on a second member of the support device;
 - 20 (iii) applying the reagent medium to the first substrate and/or the second substrate; and
 - (iv) operating an alignment mechanism of the support device to move the first member and/or the second member such that at least a portion of the biological sample is aligned with at least a portion of the array, and such that the portion of the
 - 25 biological sample and the portion of the array contact the reagent medium.
18. The method of claim 17, wherein the alignment mechanism is coupled to the first member, the second member, or both the first member and the second member.
- 30 19. The method of claim 17 or 18, wherein the alignment mechanism comprises a linear actuator, optionally wherein:
- the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member, and/or

the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member, and/or

the linear actuator is configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec, and/or

the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.

20. The method of any one of the preceding claims, wherein during the releasing step (d), a separation distance is maintained between the first substrate and the second substrate, optionally wherein the separation distance is less than 50 microns, optionally wherein the separation distance is between 2-25 microns, optionally wherein the separation distance is measured in a direction orthogonal to the surface of the first substrate that supports the biological sample, and/or at least the portion of the biological sample is vertically aligned with the at least portion of the array.

21. The method of any one of the preceding claims, wherein at least one of the first substrate and the second substrate further comprise a spacer, wherein after the first and second substrate being mounted on the support device, the spacer is disposed between the first substrate and second substrate and is configured to maintain the reagent medium within a chamber formed by the first substrate, the second substrate, and the spacer, and maintain a separation distance between the first substrate and the second substrate, the spacer positioned to at least partially surround an area on the first substrate on which the biological sample is disposed and/or the array disposed on the second substrate, wherein the area of the first substrate, the spacer, and the second substrate at least partially encloses a volume comprising the biological sample.

22. The method of claim 21, wherein:
the chamber comprises a partially or fully sealed chamber, and/or
the second substrate comprises the spacer, and/or
the first substrate comprises the spacer, and/or

the applying the reagent medium to the first substrate and/or the second substrate comprises applying the reagent medium to a region of the spacer, the region outside an enclosed area of the second substrate, the enclosed area formed by the spacer.

- 5 23. The method of any one of claims 17-22, wherein as the first substrate and/or the second substrate are moved via the alignment mechanism, the first substrate is at an angle relative to the second substrate such that a dropped slide of the first substrate and a portion of the second substrate contact the reagent medium,
optionally wherein:
10 the dropped side of the first substrate urges the reagent medium toward the opposite direction, and/or
 the alignment mechanism further moves the first substrate and/or the second substrate to maintain an approximately parallel arrangement of the first substrate and the second substrate and a separation distance between the first substrate and
15 the second substrate, optionally when the approximately parallel arrangement and the separation distance are maintained, the spacer fully encloses and surrounds the at least portion of the biological sample and the at least portion of the array, and the spacer forms the sides of the chamber which hold a volume of the reagent medium.
20
24. The method of any one of claims 16-23, wherein the agent for releasing the connected probe comprises a nuclease.
- 25 25. The method of claim 24, wherein the nuclease comprises an RNase, optionally wherein the RNase is selected from RNase A, RNase C, RNase H, or RNase I.
26. The method of any one of claims 16-25, wherein the permeabilization agent comprises a protease.
- 30 27. The method of claim 26, wherein the protease is selected from trypsin, pepsin, elastase, or proteinase K.
28. The method of any one of claim 16-27, wherein the reagent medium further comprises a detergent.

29. The method of claim 28, wherein the detergent is selected from sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100™, or Tween-20™.
- 5 30. The method of any one of claims 16-29, wherein the reagent medium comprises less than 5 w/v% of a detergent selected from SDS and sarkosyl.
31. The method of any one of claims 16-29, wherein the reagent medium comprises at least 5% w/v% of a detergent selected from SDS and sarkosyl.
- 10 32. The method of any one of claims 16-28, wherein the reagent medium does not comprise sodium dodecyl sulfate (SDS) or sarkosyl.
33. The method of any one of claims 16-32, wherein the reagent medium further comprises polyethylene glycol (PEG).
- 15 34. The method of any one of claims 16-33, wherein the biological sample and the array are contacted with the reagent medium for about 1 to about 60 minutes.
- 20 35. The method of any one of claims 16-34, wherein the biological sample and the array are contacted with the reagent medium for about 30 minutes.
36. The method of any one of the preceding claims, further comprising determining (i) all or a part of the sequence of the connected probe, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof, optionally wherein the method further comprises using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample.
- 25 37. The method of claim 36, wherein the determining comprises sequencing (i) all or a part of the sequence of the connected probe, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof.
- 30

38. The method of claim 37, wherein the sequence of the connected probe comprises the sequence of the spatial barcode or the reverse complement thereof, and a sequence corresponding to the analyte in the biological sample or reverse complement thereof.
- 5 39. The method of any one of the preceding claims, wherein at least 50% of the connected probe released from the portion of the biological sample aligned with the portion of the array are captured by capture probes of the portion of the array.
- 10 40. The method of any one of the preceding claims, wherein at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% of the connected probe are detected in spots directly under the biological sample.
- 15 41. The method of any one of the preceding claims, wherein the capture probe comprises a poly(T) sequence.
- 20 42. The method of any one of the preceding claims, wherein the capture probe comprises a sequence complementary to the first probe oligonucleotide or the second probe oligonucleotide.
- 25 43. The method of any one of the preceding claims, wherein the capture probe further comprises one or more functional domains, a unique molecular identifier (UMI), a cleavage domain, and combinations thereof.
44. The method of any one of the preceding claims, wherein the analyte is RNA.
45. The method of claim 44, wherein the RNA is mRNA.
- 30 46. The method of any one of the preceding claims, further comprising analyzing a different analyte in the biological sample.
47. The method of claim 46, wherein the analyzing the different analyte comprises:

- 5 (a) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the different analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence; and
- (b) hybridizing the capture handle sequence to the capture domain.
- 10 48. The method of claim 47, further comprising determining (i) all or part of the sequence of the capture agent barcode domain; and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof.
- 15 49. The method of claim 48, further comprising using the determined sequence of (i), and (ii) to analyze the different analyte in the biological sample.
50. The method of any one of claims 46-49, further comprising a releasing step, wherein the capture agent barcode domain is released from the different analyte.
- 20 51. The method of any one of claims 46-50, wherein the different analyte is a protein analyte.
52. The method of claim 51, wherein the protein analyte is an extracellular protein.
53. The method of claim 52, wherein the protein analyte is an intracellular protein.
- 25 54. The method of any one of claims 47-53, wherein the analyte binding moiety is an antibody.
55. The method of any one of claims 47-54, wherein the analyte capture agent comprises a linker.
- 30 56. The method of claim 55, wherein the linker is a cleavable linker.
57. The method of claim 56, wherein the cleavable linker is a photo-cleavable linker, a UV-cleavable linker, or an enzyme cleavable linker.

58. The method of claim 56 or 57, wherein the cleavable linker is a photo-cleavable linker.

5 59. A method for analyzing an analyte in a biological sample mounted on a first substrate, the method comprising:

- 10 (a) contacting the biological sample with a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence;
- 15 (b) aligning the first substrate with a second substrate comprising an array, such that at least a portion of the biological sample is aligned with at least a portion of the array, wherein the array comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises: (i) a spatial barcode and (ii) a capture domain;
- 20 (c) when the biological sample is aligned with at least a portion of the array, (i) releasing the capture agent barcode domain from the analyte and (ii) passively or actively migrating the capture agent barcode domain from the biological sample to the array; and
- (d) coupling the capture handle sequence to the capture domain.

25 60. The method of claim 59, further comprising determining (i) all or a part of the capture agent barcode domain, or a complement thereof; and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof.

30 61. The method of claim 60, further comprising using the determined sequence of (i) and (ii) to determine the location and abundance of the analyte in the biological sample.

62. The method of any one of claims 59-61, wherein the determining comprises sequencing (i) all or a part of the sequence of the capture agent barcode domain, or a complement thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement thereof.

35

63. The method of any one of claims 59-62, wherein the releasing step (c) comprises contacting the biological sample and the array with a reagent medium comprising a nuclease.
- 5 64. The method of any one of claims 59-63, wherein the releasing step (c) comprises contacting the biological sample and the array with a reagent medium comprising a permeabilization agent.
65. The method of claim 63 or 64, wherein the aligning comprises:
- 10 (i) mounting the first substrate on a first member of a support device, the first member configured to retain the first substrate;
- (ii) mounting the second substrate on a second member of the support device,
- (iii) applying the reagent medium to the first substrate and/or the second substrate, and
- 15 (iv) operating an alignment mechanism of the support device to move the first member and/or the second member such that at least a portion of the biological sample is aligned with at least a portion of the array, and such that the portion of the biological sample and the portion of the array contact the reagent medium.
- 20 66. The method of claim 65, wherein the alignment mechanism is coupled to the first member, the second member, or both the first member and the second member.
67. The method of claim 65 or 66, wherein the alignment mechanism comprises a linear actuator, optionally wherein:
- 25 the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member, and/or
- the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member, and/or
- the linear actuator is configured to move the first member, the second member,
- 30 or both the first member and the second member at a velocity of at least 0.1 mm/sec, and/or
- the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.

68. The method of any one of claims 59-67, wherein during the releasing step (d), a separation distance is maintained between the first substrate and the second substrate, optionally wherein the separation distance is less than 50 microns, optionally wherein the separation distance is between 2-25 microns, optionally wherein the separation distance is measured in a direction orthogonal to the surface of the first substrate that supports the biological sample, and/or at least the portion of the biological sample is vertically aligned with the at least portion of the array.
69. The method of any one of claims 59-68, wherein at least one of the first substrate and the second substrate further comprise a spacer, wherein after the first and second substrate being mounted on the support device, the spacer is disposed between the first substrate and second substrate and is configured to maintain the reagent medium within a chamber formed by the first substrate, the second substrate, and the spacer, and maintain a separation distance between the first substrate and the second substrate, the spacer positioned to at least partially surround an area on the first substrate on which the biological sample is disposed and/or the array disposed on the second substrate, wherein the area of the first substrate, the spacer, and the second substrate at least partially encloses a volume comprising the biological sample.
70. The method of claim 69, wherein:
- the chamber comprises a partially or fully sealed chamber, and/or
 - the second substrate comprises the spacer, and/or
 - the first substrate comprises the spacer, and/or
 - the applying the reagent medium to the first substrate and/or the second substrate comprises applying the reagent medium to a region of the spacer, the region outside an enclosed area of the second substrate, the enclosed area formed by the spacer.
71. The method of any one of claims 65-70, wherein as the first substrate and/or the second substrate are moved via the alignment mechanism, the first substrate is at an angle relative to the second substrate such that a dropped slide of the first substrate and a portion of the second substrate contact the reagent medium, optionally wherein:

the dropped side of the first substrate urges the reagent medium toward the opposite direction, and/or

5 the alignment mechanism further moves the first substrate and/or the second substrate to maintain an approximately parallel arrangement of the first substrate and the second substrate and a separation distance between the first substrate and the second substrate, optionally when the approximately parallel arrangement and the separation distance are maintained, the spacer fully encloses and surrounds the at least portion of the biological sample and the at least portion of the array, and the spacer forms the sides of the chamber which hold a volume of the reagent
10 medium.

72. The method of any one of claims 59-71, wherein the coupling of the capture handle sequence to the capture domain comprises hybridization.

15 73. The method of claim 71 or 72, wherein the nuclease comprises an RNase.

74. The method of claim 73, wherein the RNase is selected from RNase A, RNase C, RNase H, and RNase I.

20 75. The method of claim 63, wherein the reagent medium further comprises a permeabilization agent.

76. The method of any one of claims 63-75, wherein the releasing step further comprises simultaneously permeabilizing the biological sample and releasing the capture agent
25 barcode domain from the analyte binding moiety.

77. The method of claims 63-75, wherein the permeabilization agent comprises a protease.

78. The method of claim 76, wherein the protease is selected from trypsin, pepsin, elastase, or
30 Proteinase K.

79. The method of any one of claim 63-78, wherein the reagent medium further comprises a detergent.

80. The method of claim 79, wherein the detergent is selected from sodium dodecyl sulfate (SDS), sarkosyl, saponin, Triton X-100™, or Tween-20™.
81. The method of any one of claims 63-79, wherein the reagent medium comprises less than 5 w/v% of a detergent selected from SDS and sarkosyl, or wherein the reagent medium comprises as least 5% w/v% of a detergent selected from SDS and sarkosyl.
82. The method of any one of claims 63-78, wherein the reagent medium does not comprise sodium dodecyl sulfate (SDS) or sarkosyl.
83. The method of any one of claims 63-82, wherein the reagent medium further comprises PEG.
84. The method of any one of claims 59-83, wherein the biological sample and the array are contacted with the reagent medium for about 1 to about 60 minutes.
85. The method of any one of claims 59-84, wherein the biological sample and the array are contacted with the reagent medium for about 30 minutes.
86. The method of any one of claims 59-85, wherein the capture probe comprises a poly(T) sequence.
87. The method of any one of claims 59-86, wherein the capture probe comprises a sequence complementary to the capture handle sequence.
88. The method of any one of claims 59-76, wherein the capture probe further comprises one or more functional domains, a unique molecular identifier (UMI), a cleavage domain, and combinations thereof.
89. The method of any one of claims 59-88, wherein the analyte binding moiety is an antibody.
90. The method of any one of claims 59-89, wherein the analyte capture agent comprises a linker.

91. The method of claim 90, wherein the linker is a cleavable linker.

92. The method of claim 91, wherein the cleavable linker is a photo-cleavable linker, a UV-
5 cleavable linker, or an enzyme cleavable linker.

93. The methods of any one of the preceding claims, wherein the biological sample is a tissue
sample.

10 94. The method of claim 93, wherein the tissue sample is a solid tissue sample.

95. The method of claim 94, wherein the solid tissue sample is a tissue section.

96. The method of any one of the preceding claims, wherein the biological sample is a fixed
15 tissue sample.

97. The method of claim 96, wherein the fixed tissue sample is a formalin fixed paraffin
embedded (FFPE) tissue sample.

20 98. The method claim 97, wherein the FFPE tissue is deparaffinized and decrosslinked prior
to step (a).

99. The method of claim 96, wherein the fixed tissue sample is a formalin fixed paraffin
embedded cell pellet.

25

100. The method of claim 93, wherein the tissue sample is a fresh frozen tissue sample.

101. The method of claim 93, wherein the tissue sample is fixed and stained prior to step
(a).

30

102. The method of any one of the preceding claims, further comprising analyzing a
second analyte in a second biological sample on a third substrate.

103. The method of claim 102, wherein the analyzing the second analyte comprises:

- 5 (a) hybridizing a third probe oligonucleotide and a fourth probe oligonucleotide to the second analyte, wherein the third probe oligonucleotide and the fourth probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the second analyte, and wherein the fourth probe oligonucleotide comprises a second capture probe binding domain;
- (b) coupling the third probe oligonucleotide and the fourth probe oligonucleotide, thereby generating a second connected probe;
- 10 (c) aligning the third substrate with the second substrate comprising the array, such that at least a portion of the second biological sample is aligned with at least a portion of the array;
- (d) when the second biological sample is aligned with at least a portion of the array, (i) releasing the second connected probe from the second analyte and (ii) migrating the second connected probe from the second biological sample
- 15 to the array; and
- (e) hybridizing the second connected probe to the capture domain.

104. The method of claim 102 or 103, wherein the second analyte is RNA.

20 105. The method of claim 104, wherein the RNA is mRNA.

106. The method of claim 102, wherein analyzing a second analyte in a second biological sample on a third substrate comprises:

- 25 (a) contacting the second biological sample with a plurality of second analyte capture agents, wherein a second analyte capture agent of the plurality of second analyte capture agents comprises a second analyte binding moiety and a second capture agent barcode domain, wherein the second analyte binding moiety specifically binds to the second analyte, and wherein the second capture agent barcode domain comprises a second analyte binding moiety barcode and a second capture handle
- 30 sequence;
- (b) aligning the third substrate with a second substrate comprising an array, such that at least a portion of the second biological sample is aligned with at least a portion of the array;

- (c) when the second biological sample is aligned with at least a portion of the array,
(i) releasing the second capture agent barcode domain from the second analyte and
(ii) migrating the second capture agent barcode domain from the second biological
sample to the array; and
5 (d) coupling the second capture handle sequence to the capture domain.
107. The method of claim 106, wherein the second analyte is a protein analyte.
108. The method of claim 107, wherein the protein analyte is an extracellular protein.
109. The method of claim 107, wherein the protein analyte is an intracellular protein.
110. The method of any one of claims 102-105, further comprising determining (i) all or a
part of the sequence of the connected probe, or a complement thereof, and (ii) all or a part
15 of the sequence of the spatial barcode, or a complement thereof, optionally wherein the
method further comprises using the determined sequence of (i) and (ii) to determine the
location and abundance of the second analyte in the biological sample.
111. The method of any one of claims 102, or 106-109, further comprising determining (i)
20 all or a part of the sequence of the second capture handle sequence, or a complement
thereof, and (ii) all or a part of the sequence of the spatial barcode, or a complement
thereof, optionally wherein the method further comprises using the determined sequence
of (i) and (ii) to determine the location and abundance of the second analyte in the
biological sample.
112. The method of any one of claims 102-111, wherein the second biological sample is a
second tissue sample, optionally wherein the second tissue sample is a second solid tissue
sample.
113. The method of claim 112, wherein the second tissue sample is a second tissue section.
114. The method of claim 112 or 113, wherein the second tissue sample is a second fixed
tissue sample.

115. The method of claim 114, wherein the second fixed tissue sample is a second formalin fixed paraffin embedded (FFPE) tissue sample.
116. The method of claim 115, wherein the second FFPE tissue is deparaffinized and
5 decrosslinked prior to step (a).
117. The method of any one of claims 114-116, wherein the second fixed tissue sample is a second formalin fixed paraffin embedded cell pellet.
- 10 118. The method of claim 112, wherein the second tissue sample is a second fresh tissue sample or a second frozen tissue sample.
119. The method of any one of claims 112-117, wherein the second tissue sample is fixed and stained prior to step (a).
15
120. The method of any one of claims 102-119, further comprising analyzing:
(a) a third analyte in a third biological sample on a fourth substrate;
(b) a fourth analyte in a fourth biological sample on a fifth substrate;
(c) a fifth analyte or more in a fifth biological sample or more on a sixth
20 substrate or more.
121. A system or kit for analyzing an analyte in a biological sample, the system or the kit comprising:
(a) a support device configured to retain a first substrate and a second
25 substrate, wherein the biological sample is placed on the first substrate, and wherein the second substrate comprises a plurality of capture probes, wherein a capture probe of the plurality of capture probes comprises: (i) a spatial barcode and (ii) a capture domain;
(b)
30 (b1) a first probe oligonucleotide and a second probe oligonucleotide, wherein the first probe oligonucleotide and the second probe oligonucleotide each comprise a sequence that is substantially complementary to adjacent sequences of the analyte, wherein the second probe oligonucleotide comprises a capture probe binding domain, and wherein the first probe

oligonucleotide and the second probe oligonucleotide are capable of being ligated together to form a connected probe;

or

- 5 (b2) a plurality of analyte capture agents, wherein an analyte capture agent of the plurality of analyte capture agents comprises an analyte binding moiety and a capture agent barcode domain, wherein the analyte binding moiety specifically binds to the analyte, and wherein the capture agent barcode domain comprises an analyte binding moiety barcode and an capture handle sequence;
- 10 (c) a reagent medium comprising a permeabilization agent and optionally an agent for releasing the connected probe; and
- (d) instructions for performing the method of any one of the preceding claims.

15 122. The system or kit of claim 121, wherein the permeabilization agent is pepsin or proteinase K.

123. The system or kit of claim 121 or 122, wherein the agent for releasing the connected probe is an RNase, optionally wherein the RNase is selected from RNase A, RNase C, 20 RNase H, or RNase I.

124. The system or kit of any one of claims 121-123, further comprising an alignment mechanism on the support device to align the first substrate and the second substrate.

25 125. The system or kit of claim 124, wherein the alignment mechanism comprises a linear actuator, wherein the first substrate comprises a first member and the second substrate comprises a second member, and optionally wherein:

the linear actuator is configured to move the second member along an axis orthogonal to the plane of the first member and/or the second member, and/or

30 the linear actuator is configured to move the first member along an axis orthogonal to the plane of the first member and/or the second member, and/or

the linear actuator is configured to move the first member, the second member, or both the first member and the second member at a velocity of at least 0.1 mm/sec, and/or

the linear actuator is configured to move the first member, the second member, or both the first member and the second member with an amount of force of at least 0.1 lbs.