

DESCRIPTION
NITROGENATED HETEROCYCLIC COMPOUND

Technical Field

5 [0001]

The present invention relates to a nitrogen-containing heterocyclic compound having a PDE2A selective inhibitory action, which is useful as an agent for the prophylaxis or treatment of schizophrenia, Alzheimer's disease and the like.

10 **Background of the Invention**

[0002]

Cyclic nucleotide phosphodiesterases (PDEs) are enzymes that regulate the cellular levels of the second messengers, cyclic adenosine monophosphate (cAMP) and cyclic guanosine
15 monophosphate (cGMP), by controlling their rates of degradation. PDEs are a superfamily of enzymes encoded by 21 genes and subdivided into 11 distinct families according to structural and functional properties. The PDE enzymes selectively catalyze the hydrolysis of the 3'-ester bond of
20 cAMP and/or cGMP, forming the inactive 5'-monophosphate. On the basis of substrate specificity, the PDE families can be further classified into three groups: i) the cAMP-PDEs (PDE4, PDE7 and PDE8), ii) the cGMP-PDEs (PDE5, PDE6 and PDE9), and
iii) the dual-substrate PDEs (PDE1, PDE2, PDE3, PDE10 and
25 PDE11).

[0003]

cAMP and cGMP are involved in the regulation of virtually every physiological process such as pro-inflammatory mediator production and action, ion channel function, muscle relaxation,
30 learning and memory formation, differentiation, apoptosis, lipogenesis, glycogenolysis and gluconeogenesis. Especially, in neurons, these second messengers have an important role in the regulation of synaptic transmission as well as in neuronal differentiation and survival (Non-Patent Document 1).
35 Regulation of these processes by cAMP and cGMP are accompanied

by activation of protein kinase A (PKA) and protein kinase G (PKG), which in turn phosphorylate a variety of substrates, including transcription factors, ion channels and receptors that regulate a variety of physiological processes.

5 Intracellular cAMP and cGMP concentrations seem to be temporally, spatially, and functionally compartmentalized by regulation of adenylate and guanylate cyclases in response to extracellular signaling and their degradation by PDEs (Non-Patent Document 2). PDEs provide the only means of degrading
10 the cyclic nucleotides cAMP and cGMP in cells, thus PDEs play an essential role in cyclic nucleotide signaling. Thereby, PDEs could be promising targets for various therapeutic drugs.
[0004]

Phosphodiesterase 2A (PDE2A) is a dual substrate enzyme
15 that hydrolyzes both cAMP and cGMP. It is organized into four domains, N-terminus, GAF-A, GAF-B, and catalytic domains, and functions as a homodimer. PDE2A catalytic activity is allosterically stimulated by cGMP binding. GAF-B domain binds with a high affinity and a high selectivity to cGMP. A
20 conformational change is caused by the cGMP binding in the PDE2A homodimer which causes an increase in the catalytic activity of the enzyme (Non-Patent Document 3-6). In contrast, there are as yet no known in vivo examples that cAMP stimulates PDE2A catalytic activity, even though it can also
25 bind to the GAF-B domain with a 30-100-fold lower affinity than cGMP (Non-Patent Document 6 and 7). PDE2A activity may become functionally significant under conditions in which cellular cGMP concentrations are elevated, which shows a physiological role for GAF domain-regulation of the enzyme.
30 [0005]

PDE2A is expressed in a wide variety of tissues and highly in the brain. The protein was originally purified from heart, liver, adrenal gland, platelets, endothelial cells, and macrophages (Non-Patent Document 8-13). In the brain, the
35 PDE2A mRNA levels are the highest in the caudate lobe, nucleus

accumbens, cortex (frontal, parietal and temporal) and the hippocampus, and are at least 10-fold lower expression in other brain regions (Non-Patent Document 14). This suggests that PDE2A may control intraneuronal cAMP and cGMP levels in
5 areas that are important for learning and memory formation.
[0006]

Inhibition of PDE2A results in increased cAMP and cGMP levels that could improve cognitive function. In both cortical neurons and hippocampal slices, a PDE2A inhibitor potentially
10 increased cGMP concentrations in the presence of guanylate cyclase activators and also increased cAMP concentrations in the presence of forskolin (Non-Patent Document 15). The PDE2A inhibitor was also found to potentially increase the induction of long-term potentiation (LTP) in hippocampal slices in response
15 to a weak tetanizing stimulus. This effect on LTP in slices suggests that PDE2A inhibition has positive effects on learning and memory in vivo (Non-Patent Document 15). In fact, the same PDE2A inhibitor increased retention on both novel object and social recognition tasks in rats, and improved
20 object memory and object recognition task in 3-, 12-, and 24-month old rats. It also attenuated the extradimensional (ED) shift deficit on extradimensional-intradimensional (ED/ID) cognitive task in subchronic PCP-treated rats (Non-Patent Document 15-17). These results suggest that PDE2A inhibition
25 could facilitate learning and memory processes through potentiation of cAMP and cGMP-regulated signaling cascades.
[0007]

Increased cGMP levels by PDE2A inhibition could also influence anxiety and stress-related events. PDE2A inhibitors
30 decreased oxidative stress and induced the expression of NADPH oxidase subunits in oxidative stress inducer-treated mice. It improved anxiety-like behavior in elevated plus maze, open-field, and hole-board tests through the NADPH oxidase pathway (Non-Patent Document 18). In addition, PDE2A inhibitors also
35 produced anxiolytic effects on behavior in non-stressed mice

in the elevated plus-maze and hole-board tests (Non-Patent Document 19). PDE2A may be a novel pharmacological target for treatment of not only cognitive deficit, but also anxiety in neuropsychiatric and neurodegenerative disorders.


5 [0008]

These unique distribution and functions in the brain indicate that PDE2A represents an important novel target for the treatment of neuropsychiatric and neurodegenerative disorders, in particular schizophrenia and Alzheimer's disease.

10 [0009]

Patent Document 1 discloses, as a nitrogen-containing heterocyclic compound, a compound represented by the formula (I):

[0010]



15

[0011]

wherein

Z¹ and Z² are each independently N or CR (at least one is CR, and R is halogen, -OH, -CN, fluoro-substituted C₁₋₂ alkyl, -O-(C₁₋₂) fluoro-substituted alkyl, -S-C₁₋₂ fluoro-substituted alkyl or the like);

W is -O-, -NH-, -N(C₁₋₄ alkyl)-, -S-, -S(O)-, -S(O)₂- or the like;

R⁶ is H, C₁₋₄ alkyl or fluoro-substituted C₁₋₄ alkyl, or two R⁶, which are bonded to the single carbon atom, in combination optionally form =O;

R¹ is carbocycle or heterocycle, each of which is optionally substituted by halo, -CN, C₁₋₄ alkyl or the like;

R² is carbocycle or heterocycle, each of which is optionally substituted by halo, -CN, C₁₋₄ alkyl or the like;

P is 1, 2 or 3;

X² is C(=O), -C(=O)-O-*, -C(=O)-NH-*, -C(=O)-NHCR⁴R⁵-* or the like (* means the binding position to R¹);

R⁴ and R⁵ are each independently H, C₁₋₄ alkyl, -CF₃ or (C₁₋₃ alkyl)-CF₃,

5 which is a sirtuin modulator, and useful for treatment of Alzheimer's disease, Parkinson's disease.

However, the structure of the present invention is different from that of the above-mentioned compound.

Document List

10 Patent Document

[0012]

Patent Document 1: WO 2011/059839

Non-Patent Document

[0013]

15 Non-Patent Document 1: Nat. Rev. Drug Discov. 2006, vol. 5: 660-670

Non-Patent Document 2: Circ. Res. 2007, vol. 100: 950-966

Non-Patent Document 3: J. Biol. Chem. 1971, vol. 246: 3841-3846

20 Non-Patent Document 4: J. Biol. Chem. 1973, vol. 248: 1334-1340

Non-Patent Document 5: PNAS 2005, vol. 99: 13260-13265

Non-Patent Document 6: British J. Pharmacol. 2010, vol. 161: 1645-1660

25 Non-Patent Document 7: J. Biol. Chem. 2004, vol. 279: 37928-37938

Non-Patent Document 8: J. Biol. Chem. 1982, vol. 257: 1973-1979

30 Non-Patent Document 9: J. Biol. Chem. 1983, vol. 258: 12526-12533

Non-Patent Document 10: Phosphodiesterase Inhibitors, Academic Press: 21-40

Non-Patent Document 11: Rev. Physiol. Biochem. Pharmacol. 1999, vol. 135: 67-104

35 Non-Patent Document 12: Cell Signal 2004, vol. 16: 365-374

Non-Patent Document 13: J. Histochem. Cytochem. 2009, vol. 57: 933-949

Non-Patent Document 14: Neuropharmacology 2010, vol. 59: 367-374

5 Non-Patent Document 15: Neuropharmacology 2004, vol. 47: 1081-1092

Non-Patent Document 16: Mol. Neurobiol. 2010, vol. 41: 129-137

Non-Patent Document 17: Neuropharmacology 2012, vol. 62: 1182-1190

10 Non-Patent Document 18: J. Pharmacol. Exp. Ther. 2008, vol. 326: 369-379

Non-Patent Document 19: J. Pharmacol. Exp. Ther. 2009, vol. 331: 690-699

Summary of the Invention

15 Problems to be Solved by the Invention

[0014]

The present invention aims to provide a compound having a PDE2A selective inhibitory action, which is useful as a prophylactic or therapeutic drug for schizophrenia,

20 Alzheimer's disease and the like.

Means of Solving the Problems

[0015]


The present inventors have conducted intensive studies, and have found that a compound represented by the formula (1) shown below unexpectedly has a superior PDE2A selective inhibitory action, and therefore, is useful as a prophylactic or therapeutic drug for schizophrenia, Alzheimer's disease and the like, and completed the present invention based on these findings.

30 [0016]

Accordingly, the present invention provides the following:

[1] A compound represented by the formula (1):

[0017]



[0018]

wherein

R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl
5 group,

R² and R³ are each independently a hydrogen atom or an
optionally substituted C₁₋₆ alkyl group, or R² and R³ in
combination optionally form, together with the adjacent carbon
atom, an optionally substituted ring,

10 X is a bond or an optionally substituted C₁₋₆ alkylene group,
A is an optionally substituted cyclic group,

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom,
a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
15 substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
alkoxy group, an optionally substituted amino group, a cyano
group or an optionally substituted cyclic group, or a nitrogen
atom,

Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom,
20 a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
alkoxy group, an optionally substituted amino group, a cyano
group or an optionally substituted cyclic group, or a nitrogen
25 atom,

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom,
a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
30 alkoxy group, an optionally substituted amino group, a cyano

group, an optionally substituted C₁₋₆ alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom, and B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms,

5 provided that

2,3-dihydro-1-methyl-2-oxo-N-2-thiazolyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

2,3-dihydro-1-methyl-N-(4-methyl-2-thiazolyl)-2-oxo-6-[3-

10 (trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-(tetrahydro-2-oxo-3-furanyl)-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

15 2,3-dihydro-1-methyl-N-2-oxazolyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-1H-1,2,4-triazol-3-yl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

20 4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-4-pyridinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-4-pyrimidinyl-6-[3-

25 (trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-5-pyrimidinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

30 2,3-dihydro-1-methyl-2-oxo-N-3-pyridazinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,

2,3-dihydro-1-methyl-N-[6-(4-morpholinyl)-2-pyridinyl]-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

35 4(1H)carboxamide,

N-(4,5-dimethyl-2-thiazolyl)-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 2,3-dihydro-1-methyl-N-(5-methyl-2-thiazolyl)-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 N-[5-[(dimethylamino)carbonyl]-4-methyl-thiazolyl]-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 10 N-(4,6-dimethyl-2-pyridinyl)-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 2,3-dihydro-1-methyl-2-oxo-N-2-pyrazinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 15 2,3-dihydro-1-methyl-2-oxo-N-2-pyrimidinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 N-[2-(1-azetidynyl)-4-pyridinyl]-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 20 [3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
 2,3-dihydro-1-methyl-N-[6-(4-morpholinylmethyl)-2-pyridinyl]-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide, and
 25 N-[6-(1-azetidynyl)-2-pyridinyl]-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide are excluded,
 or a salt thereof (hereinafter to be referred as compound (1)).
 [2] The compound or salt of the above-mentioned [1], wherein
 30 R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group or a cyano group, and
 35 Z^3 is CR^{Z3} wherein R^{Z3} is a hydrogen atom, a halogen atom, an

optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group.

[3] The compound or salt of the above-mentioned [1], wherein R¹ is a hydrogen atom.

[4] The compound or salt of the above-mentioned [1], wherein R² and R³ are both hydrogen atoms.

[5] The compound or salt of the above-mentioned [1], wherein Z¹, Z² and Z³ are respectively CR^{Z1}, CR^{Z2} and CR^{Z3}.

[6] The compound or salt of the above-mentioned [5], wherein R^{Z1} and R^{Z3} are both hydrogen atoms.

[7] The compound or salt of the above-mentioned [5], wherein R^{Z2} is an optionally substituted C₁₋₆ alkoxy group, an optionally substituted C₁₋₆ alkyl group or an optionally substituted C₃₋₈ cycloalkyl group.

[8] The compound or salt of the above-mentioned [1], wherein X is an optionally substituted C₁₋₆ alkylene group.

[9] The compound or salt of the above-mentioned [1], wherein A is an optionally substituted phenyl group.

[10] The compound or salt of the above-mentioned [1], wherein R¹ is a hydrogen atom,

R² and R³ are both hydrogen atoms,

X is an optionally substituted C₁₋₆ alkylene group,

A is an optionally substituted phenyl group,

Z¹ is CH,

Z² is a group represented by CR^{Z2} wherein R^{Z2} is an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₁₋₆ alkoxy group or an optionally substituted C₃₋₈ cycloalkyl group, and

Z³ is CH.

[11] 7-Methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide or a salt thereof.

[12] N-((1S)-1-(3-Fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide or a salt thereof.

[13] 7-Cyclopropyl-N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide or a salt thereof.

[14] 7-Cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide or a salt thereof.

10 [15] N-(1-(3-Fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide or a salt thereof.


[16] A medicament comprising the compound or salt of the above-mentioned [1].

15 [17] The medicament of the above-mentioned [16], which is a phosphodiesterase 2A inhibitor.

[18] The medicament of the above-mentioned [16], which is an agent for the prophylaxis or treatment of schizophrenia.

[19] A method of inhibiting phosphodiesterase 2A, which
20 comprises administering an effective amount of a compound represented by the formula (1):

[0019]



[0020]

25 wherein

R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group,


R² and R³ are each independently a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, or R² and R³ in

30 combination optionally form, together with the adjacent carbon

atom, an optionally substituted ring,
X is a bond or an optionally substituted C₁₋₆ alkylene group,
A is an optionally substituted cyclic group,
Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom,
5 a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
alkoxy group, an optionally substituted amino group, a cyano
group or an optionally substituted cyclic group, or a nitrogen
10 atom,
Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom,
a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
15 alkoxy group, an optionally substituted amino group, a cyano
group or an optionally substituted cyclic group, or a nitrogen
atom,
Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom,
a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
20 optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
alkoxy group, an optionally substituted amino group, a cyano
group, an optionally substituted C₁₋₆ alkyl-carbonyl group or an
optionally substituted cyclic group, or a nitrogen atom, and
25 B is a nitrogen-containing aromatic heterocycle containing 1
or 2 nitrogen atoms,
or a salt thereof, to a mammal.

[20] A method for the prophylaxis or treatment of
schizophrenia, which comprises administering an effective
30 amount of a compound represented by the formula (1):

[0021]



[0022]

wherein

R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl
5 group,

R² and R³ are each independently a hydrogen atom or an
optionally substituted C₁₋₆ alkyl group, or R² and R³ in
combination optionally form, together with the adjacent carbon
atom, an optionally substituted ring,

10 X is a bond or an optionally substituted C₁₋₆ alkylene group,
A is an optionally substituted cyclic group,

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom,
a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
15 substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
alkoxy group, an optionally substituted amino group, a cyano
group or an optionally substituted cyclic group, or a nitrogen
atom,

Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom,
20 a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
alkoxy group, an optionally substituted amino group, a cyano
group or an optionally substituted cyclic group, or a nitrogen
25 atom,


Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom,
a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
optionally substituted C₂₋₆ alkenyl group, an optionally
substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
30 alkoxy group, an optionally substituted amino group, a cyano

alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z^3 is a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom,
5 a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C_{1-6} alkyl-carbonyl group or an
10 optionally substituted cyclic group, or a nitrogen atom, and
B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms,
or a salt thereof, for the production of an agent for the prophylaxis or treatment of schizophrenia.

15 [22] A compound represented by the formula (1):

[0025]



[0026]

wherein

20 R^1 is a hydrogen atom or an optionally substituted C_{1-6} alkyl group,

R^2 and R^3 are each independently a hydrogen atom or an optionally substituted C_{1-6} alkyl group, or R^2 and R^3 in combination optionally form, together with the adjacent carbon
25 atom, an optionally substituted ring,

X is a bond or an optionally substituted C_{1-6} alkylene group,

A is an optionally substituted cyclic group,

Z^1 is a group represented by CR^{Z^1} wherein R^{Z^1} is a hydrogen atom,
a halogen atom, an optionally substituted C_{1-6} alkyl group, an
30 optionally substituted C_{2-6} alkenyl group, an optionally

substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

5 Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C₁₋₆ alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom, and B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms,

20 or a salt thereof, for use in the prophylaxis or treatment of schizophrenia.

Effect of the Invention

[0027]

25 According to the present invention, the compound having a PDE2A selective inhibitory action, which is useful as an agent for the prophylaxis or treatment of schizophrenia, Alzheimer's disease and the like, can be provided.

Brief Description of the Drawings

30 [0028]

Fig. 1 shows a graph that a test compound has an improvement effect for MK-801 induced disorder in Contextual fear conditioning test (Experimental Example 2).

[0029]

35 [Detailed Description of the Invention]

The present invention is explained in detail in the following.

The "halogen atom" in the present specification means, unless otherwise specified, a fluorine atom, a chlorine atom,
5 a bromine atom or an iodine atom.

[0030]

The "optionally substituted hydrocarbon group" in the present specification means, unless otherwise specified, for example, an "optionally substituted C₁₋₁₀ alkyl", an "optionally
10 substituted C₂₋₁₀ alkenyl", an "optionally substituted C₂₋₁₀ alkynyl", an "optionally substituted C₃₋₈ cycloalkyl", an "optionally substituted C₃₋₈ cycloalkenyl", an "optionally substituted C₆₋₁₄ aryl", an "optionally substituted C₇₋₁₄ aralkyl" or the like.

15 [0031]

The "C₁₋₁₀ alkyl" in the present specification means, unless otherwise specified, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, nonyl, decyl or
20 the like. The "C₁₋₆ alkyl" in the present specification means C₁₋₆ alkyl from among the above-mentioned "C₁₋₁₀ alkyl". The "C₁₋₅ alkyl" in the present specification means C₁₋₅ alkyl from among the above-mentioned "C₁₋₁₀ alkyl".

[0032]

25 The "C₂₋₁₀ alkenyl" in the present specification means, unless otherwise specified, for example, vinyl, propenyl, isopropenyl, 2-buten-1-yl, 4-penten-1-yl, 5-hexen-1-yl or the like. The "C₂₋₆ alkenyl" in the present specification means C₂₋₆ alkenyl from among the above-mentioned "C₂₋₁₀ alkenyl".

30 [0033]

The "C₂₋₁₀ alkynyl" in the present specification means, unless otherwise specified, for example, 2-butyne-1-yl, 4-pentyne-1-yl, 5-hexyne-1-yl or the like. The "C₂₋₆ alkynyl" in the present specification means C₂₋₆ alkynyl from among the
35 above-mentioned "C₂₋₁₀ alkynyl".

[0034]

The "C₃₋₈ cycloalkyl" in the present specification means, unless otherwise specified, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl
5 or the like.

[0035]

The "C₃₋₈ cycloalkenyl" in the present specification means, unless otherwise specified, for example, cyclopropenyl(e.g., 2-cyclopropen-1-yl), cyclobutenyl(e.g., 2-cyclobuten-1-yl),
10 cyclopentenyl(e.g., 1-cyclopenten-1-yl, 2-cyclopenten-1-yl, 3-cyclopenten-1-yl), cyclohexenyl(e.g., 1-cyclohexen-1-yl, 2-cyclohexen-1-yl, 3-cyclohexen-1-yl) or the like.

[0036]

The "C₆₋₁₄ aryl" in the present specification means,
15 unless otherwise specified, for example, phenyl, 1-naphthyl, 2-naphthyl, 2-biphenyl, 3-biphenyl, 4-biphenyl, 2-anthryl or the like. The C₆₋₁₄ aryl may be partially saturated, and examples of the partially saturated C₆₋₁₄ aryl include tetrahydronaphthyl and the like.

20 [0037]

The "C₇₋₁₄ aralkyl" in the present specification means, unless otherwise specified, for example, benzyl, phenethyl, 1-methyl-2-phenylethyl, diphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2,2-diphenylethyl, 3-phenylpropyl, 4-
25 phenylbutyl, 5-phenylpentyl, 2-biphenylmethyl, 3-biphenylmethyl, 4-biphenylmethyl or the like.

[0038]

The "optionally substituted hydroxy" in the present specification means, unless otherwise specified, for example,
30 "hydroxy", "optionally substituted C₁₋₆ alkoxy", "optionally substituted heterocycl-yl-oxy", "optionally substituted C₆₋₁₄ aryloxy", "optionally substituted C₇₋₁₄ aralkyloxy" or the like.

[0039]

The "C₁₋₆ alkoxy" in the present specification means,
35 unless otherwise specified, for example, methoxy, ethoxy,

propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, hexyloxy or the like.

[0040]

The "C₁₋₆ alkoxy-C₁₋₆ alkoxy" in the present specification
5 means, unless otherwise specified, for example, methoxymethoxy, methoxyethoxy, ethoxymethoxy, ethoxyethoxy or the like.

[0041]

The "heterocyclyl-oxy" in the present specification means hydroxyl substituted by the below-mentioned "heterocyclic
10 group". Preferable examples of the heterocyclyl-oxy include tetrahydropyranyloxy, thiazolyloxy, pyridyloxy, pyrazolyloxy, oxazolyloxy, thienyloxy, furyloxy and the like.

[0042]

The "C₆₋₁₄ aryloxy" in the present specification means,
15 unless otherwise specified, for example, phenoxy, 1-naphthyloxy, 2-naphthyloxy or the like.

[0043]

The "C₇₋₁₄ aralkyloxy" in the present specification means, unless otherwise specified, for example, benzyloxy,
20 phenethyloxy or the like.

[0044]

The "optionally substituted sulfanyl" in the present specification means, unless otherwise specified, for example, "sulfanyl", "optionally substituted C₁₋₆ alkylsulfanyl",
25 "optionally substituted heterocyclyl-sulfanyl", "optionally substituted C₆₋₁₄ arylsulfanyl", "optionally substituted C₇₋₁₄ aralkylsulfanyl" or the like.

[0045]

The "C₁₋₆ alkylsulfanyl" in the present specification
30 means, unless otherwise specified, for example, methylsulfanyl, ethylsulfanyl, propylsulfanyl, isopropylsulfanyl, butylsulfanyl, sec-butylsulfanyl, tert-butylsulfanyl or the like.

[0046]

35 The "heterocyclyl-sulfanyl" in the present specification

means sulfanyl substituted by the below-mentioned
"heterocyclic group". Preferable examples of the heterocyclyl-
sulfanyl include tetrahydropyranylsulfanyl, thiazolylsulfanyl,
pyridylsulfanyl, pyrazolylsulfanyl, oxazolylsulfanyl,
5 thienylsulfanyl, furylsulfanyl and the like.

[0047]

The "C₆₋₁₄ arylsulfanyl" in the present specification
means, unless otherwise specified, for example, phenylsulfanyl,
1-naphthylsulfanyl, 2-naphthylsulfanyl or the like.

10 [0048]

The "C₇₋₁₄ aralkylsulfanyl" in the present specification
means, unless otherwise specified, for example, benzylsulfanyl,
phenethylsulfanyl or the like.

[0049]

15 The "heterocyclic group" in the present specification
means, unless otherwise specified, for example, a 5- to 14-
membered (monocyclic, bicyclic or tricyclic) heterocyclic
group containing, as a ring-constituting atom besides carbon
atom, 1 or 2 kinds of 1 to 4 hetero atoms selected from a
20 nitrogen atom, a sulfur atom and an oxygen atom,
preferably (i) a 5- to 14-membered (preferably 5- to 10-
membered) aromatic heterocyclic group, (ii) a 5- to 10-
membered nonaromatic heterocyclic group or the like. Of these,
a 5- or 6-membered aromatic heterocyclic group is preferable.
25 Specifically, for example, aromatic heterocyclic groups such
as thienyl (e.g., 2-thienyl, 3-thienyl), furyl (e.g., 2-furyl,
3-furyl), pyridyl (e.g., 2-pyridyl, 3-pyridyl, 4-pyridyl),
thiazolyl (e.g., 2-thiazolyl, 4-thiazolyl, 5-thiazolyl),
oxazolyl (e.g., 2-oxazolyl, 4-oxazolyl, 5-oxazolyl), pyrazinyl,
30 pyrimidinyl (e.g., 2-pyrimidinyl, 4-pyrimidinyl), pyrrolyl
(e.g., 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl), imidazolyl (e.g.,
1-imidazolyl, 2-imidazolyl, 4-imidazolyl), pyrazolyl (e.g., 1-
pyrazolyl, 3-pyrazolyl, 4-pyrazolyl), triazolyl (e.g., 1-
triazolyl, 2-triazolyl), tetrazolyl, pyridazinyl (e.g., 3-
35 pyridazinyl, 4-pyridazinyl), isothiazolyl (e.g., 3-

isothiazolyl, 4-isothiazolyl, 5-isothiazolyl), isoxazolyl
(e.g., 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl), indolyl
(e.g., 1-indolyl, 2-indolyl, 3-indolyl), 2-benzothiazolyl, 2-
benzoxazolyl, benzimidazolyl (e.g., 1-benzimidazolyl, 2-
5 benzimidazolyl), benzo[b]thienyl (e.g., 2-benzo[b]thienyl, 3-
benzo[b]thienyl), benzo[b]furyl (e.g., 2-benzo[b]furanyl, 3-
benzo[b]furanyl), quinolyl (e.g., 2-quinolyl, 3-quinolyl, 4-
quinolyl, 5-quinolyl, 8-quinolyl), isoquinolyl (e.g., 1-
isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl),
10 pyrazolopyridinyl (e.g., pyrazolo[1,5-a]pyridin-3-yl) and the
like;

for example, nonaromatic heterocyclic groups such as
pyrrolidinyl (e.g., 1-pyrrolidinyl, 2-pyrrolidinyl, 3-
pyrrolidinyl), oxazolidinyl (e.g., 2-oxazolidinyl),
15 imidazolinyl (e.g., 1-imidazolinyl, 2-imidazolinyl, 4-
imidazolinyl), piperidyl (e.g., piperidino, 2-piperidyl, 3-
piperidyl, 4-piperidyl), piperazinyl (e.g., 1-piperazinyl, 2-
piperazinyl), morpholinyl (e.g., 2-morpholinyl, 3-morpholinyl,
morpholino), thiomorpholinyl (e.g., 2-thiomorpholinyl, 3-
20 thiomorpholinyl, thiomorpholino), tetrahydrofuryl,
tetrahydropyranyl and the like can be mentioned.

[0050]

The "optionally substituted cyclic group" in the present
specification means, unless otherwise specified, for example,
25 "optionally substituted C₆₋₁₄ aryl", "optionally substituted C₃₋₈
cycloalkyl", "optionally substituted C₃₋₈ cycloalkenyl",
"optionally substituted heterocyclic group" or the like.

[0051]

The "optionally substituted ring" in the present
30 specification means, unless otherwise specified, for example,
an optionally substituted ring corresponding to the above-
mentioned "optionally substituted cyclic group".

[0052]

The "C₆₋₁₄ aromatic hydrocarbon" in the present
35 specification means a ring corresponding to the above-

mentioned "C₆₋₁₄ aryl".

The "C₃₋₈ cycloalkane" in the present specification means a ring corresponding to the above-mentioned "C₃₋₈ cycloalkyl".

The "C₃₋₈ cycloalkene" in the present specification means
5 a ring corresponding to the above-mentioned "C₃₋₈ cycloalkenyl".

The "heterocycle" in the present specification means a ring corresponding to the above-mentioned "heterocyclic group".
[0053]

The "nitrogen-containing aromatic heterocycle containing
10 1 to 2 nitrogen atoms" means, unless otherwise specified, for example, a nitrogen-containing aromatic heterocycle containing, as a ring-constituting atom besides carbon atom and 1 to 2 nitrogen atoms, one or two kinds of 1 to 3 hetero atoms selected from a sulfur atom and an oxygen atom. Specifically,
15 for example, pyridine ring, pyridazine ring, pyrimidine ring, pyrazine ring and the like can be mentioned.

[0054]

The "C₁₋₆ alkyl-carbonyl" in the present specification means, unless otherwise specified, for example, acetyl,
20 isobutanoyl, isopentanoyl or the like.

[0055]

The "C₁₋₆ alkoxy-carbonyl" in the present specification means, unless otherwise specified, for example, methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-
25 butoxycarbonyl or the like.

[0056]

The "C₃₋₈ cycloalkyl-carbonyl" in the present specification means, unless otherwise specified, for example, cyclopentylcarbonyl, cyclohexylcarbonyl or the like.

30 [0057]

The "C₆₋₁₄ aryl-carbonyl" in the present specification means, unless otherwise specified, for example, benzoyl, 1-naphthoyl, 2-naphthoyl or the like.

[0058]

35 The "C₇₋₁₄ aralkyl-carbonyl" in the present specification

means, unless otherwise specified, for example, phenylacetyl,
2-phenylpropanoyl or the like.

[0059]

The "C₆₋₁₄ aryloxy-carbonyl" in the present specification
5 means, unless otherwise specified, for example,
phenoxycarbonyl, naphthyloxycarbonyl or the like.

[0060]

The "C₇₋₁₄ aralkyloxy-carbonyl" in the present
specification means, unless otherwise specified, for example,
10 benzyloxycarbonyl, phenethyloxycarbonyl or the like.

[0061]

The "nitrogen-containing heterocyclyl-carbonyl" in the
present specification means, unless otherwise specified, for
example, pyrrolidinylcarbonyl, piperidinocarbonyl or the like.

15 [0062]

The "C₁₋₆ alkylsulfonyl" in the present specification
means, unless otherwise specified, for example, methylsulfonyl,
ethylsulfonyl or the like.

[0063]

20 The "C₆₋₁₄ arylsulfonyl" in the present specification
means, unless otherwise specified, for example, phenylsulfonyl,
1-naphthylsulfonyl, 2-naphthylsulfonyl or the like.

[0064]

The "C₁₋₆ alkylsulfinyl" in the present specification
25 means, unless otherwise specified, for example, methylsulfinyl,
ethylsulfinyl or the like.

[0065]

The "C₆₋₁₄ arylsulfinyl" in the present specification
means, unless otherwise specified, for example, phenylsulfinyl,
30 1-naphthylsulfinyl, 2-naphthylsulfinyl or the like.

[0066]

The "optionally esterified carboxyl" in the present
specification means, unless otherwise specified, for example,
carboxyl, C₁₋₆ alkoxy-carbonyl, C₆₋₁₄ aryloxy-carbonyl, C₇₋₁₄
35 aralkyloxy-carbonyl or the like.

[0067]

The "optionally halogenated C₁₋₆ alkyl" in the present specification means, unless otherwise specified, the above-mentioned "C₁₋₆ alkyl" optionally substituted by 1 to 5 of the
5 above-mentioned "halogen atom". For example, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, isobutyl, trifluoromethyl and the like can be mentioned.

[0068]

The "optionally halogenated C₁₋₆ alkoxy" in the present
10 specification means, unless otherwise specified, the above-mentioned "C₁₋₆ alkoxy" optionally substituted by 1 to 5 of the above-mentioned "halogen atom". For example, methoxy, ethoxy, isopropoxy, tert-butoxy, trifluoromethoxy and the like can be mentioned.

15 [0069]

The "mono- or di-C₁₋₆ alkyl-amino" in the present specification means, unless otherwise specified, amino mono- or di-substituted by the above-mentioned "C₁₋₆ alkyl". For example, methylamino, ethylamino, propylamino, dimethylamino,
20 diethylamino and the like can be mentioned.

[0070]

The "mono- or di-C₆₋₁₄ aryl-amino" in the present specification means, unless otherwise specified, amino mono- or di-substituted by the above-mentioned "C₆₋₁₄ aryl". For
25 example, phenylamino, diphenylamino, 1-naphthylamino, 2-naphthylamino and the like can be mentioned.

[0071]

The "mono- or di-C₇₋₁₄ aralkyl-amino" in the present specification means, unless otherwise specified, amino mono-
30 or di-substituted by the above-mentioned "C₇₋₁₄ aralkyl". For example, benzylamino, phenethylamino and the like can be mentioned.

[0072]

The "N-C₁₋₆ alkyl-N-C₆₋₁₄ aryl-amino" in the present
35 specification means, unless otherwise specified, amino

substituted by the above-mentioned "C₁₋₆ alkyl" and the above-mentioned "C₆₋₁₄ aryl". For example, N-methyl-N-phenylamino, N-ethyl-N-phenylamino and the like can be mentioned.

[0073]

5 The "N-C₁₋₆ alkyl-N-C₇₋₁₄ aralkyl-amino" in the present specification means, unless otherwise specified, amino substituted by the above-mentioned "C₁₋₆ alkyl" and the above-mentioned "C₇₋₁₄ aralkyl". For example, N-methyl-N-benzylamino, N-ethyl-N-benzylamino and the like can be mentioned.

10 [0074]

 The "C₁₋₆ alkyl-carbonylamino" in the present specification means, unless otherwise specified, amino substituted by the above-mentioned "C₁₋₆ alkyl-carbonyl". For example, acetyl amino, propionylamino and the like can be

15 mentioned.

[0075]

 The "mono- or di-C₁₋₆ alkyl-carbamoyl" in the present specification means, unless otherwise specified, carbamoyl mono- or di-substituted by the above-mentioned "C₁₋₆ alkyl".

20 For example, methylcarbamoyl, ethylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl and the like can be mentioned.

[0076]

 The "mono- or di-C₆₋₁₄ aryl-carbamoyl" in the present

25 specification means, unless otherwise specified, carbamoyl mono- or di-substituted by the above-mentioned "C₆₋₁₄ aryl". For example, phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl and the like can be mentioned.

[0077]

30 The "mono- or di-C₃₋₈ cycloalkyl-carbamoyl" in the present specification means, unless otherwise specified, carbamoyl mono- or di-substituted by the above-mentioned "C₃₋₈ cycloalkyl". For example, cyclopropylcarbamoyl and the like can be mentioned.

35 [0078]

The "mono- or di-C₇₋₁₄ aralkyl-carbamoyl" in the present specification means, unless otherwise specified, carbamoyl mono- or di-substituted by the above-mentioned "C₇₋₁₄ aralkyl". For example, benzylcarbamoyl and the like can be mentioned.

5 [0079]

The "N-C₁₋₆ alkyl-N-C₆₋₁₄ aryl-carbamoyl" in the present specification means, unless otherwise specified, carbamoyl substituted by the above-mentioned "C₁₋₆ alkyl" and "C₆₋₁₄ aryl". For example, (n-butyl)(phenyl) carbamoyl and the like can be

10 mentioned.

[0080]

The "mono- or di-5- to 7-membered heterocyclyl-carbamoyl" in the present specification means, unless otherwise specified, carbamoyl mono- or di-substituted by a 5- to 7-membered

15 heterocyclic group. Here, examples of the 5- to 7-membered heterocyclic group include a heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 or 2 kinds of 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom can be mentioned. Preferable examples

20 of the "mono- or di-5- to 7-membered heterocyclyl-carbamoyl" include 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl and the like.

[0081]

25 As the "mono- or di-C₁₋₆ alkyl-sulfamoyl" in the present specification, unless otherwise specified, sulfamoyl mono- or di-substituted by the above-mentioned "C₁₋₆ alkyl" can be used. For example, methylsulfamoyl, ethylsulfamoyl, dimethylsulfamoyl, diethylsulfamoyl and the like can be

30 mentioned.

[0082]

As the "mono- or di-C₆₋₁₄ aryl-sulfamoyl" in the present specification, unless otherwise specified, sulfamoyl mono- or di-substituted by the above-mentioned "C₆₋₁₄ aryl" can be used.

35 For example, phenylsulfamoyl, diphenylsulfamoyl, 1-

naphthylsulfamoyl, 2-naphthylsulfamoyl and the like can be mentioned.

[0083]

The "mono- or di-C₇₋₁₄ aralkyl-sulfamoyl" in the present
5 specification means, unless otherwise specified, sulfamoyl
mono- or di-substituted by the above-mentioned "C₇₋₁₄ aralkyl".
For example, benzylsulfamoyl and the like can be mentioned.

[0084]

The "C₁₋₆ alkyl-carbonyloxy" in the present specification
10 means, unless otherwise specified, for example,
methylcarbonyloxy, ethylcarbonyloxy or the like.

[0085]

The "C₁₋₆ alkylene" in the present specification means,
unless otherwise specified, for example, -CH₂-, -CH₂CH₂-, -
15 CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂CH₂CH₂-,
-CH(CH₃)-, -C(CH₃)₂-, -CH(CH₂CH₃)-, -C(CH₂CH₃)₂-, -CH(CH(CH₃)₂)-,
-CH(CH₃)CH₂-, -CH₂CH(CH₃)-, -CH(CH₂CH₃)CH₂-, -CH₂CH(CH₂CH₃)- or
the like. The "C₁₋₅ alkylene" in the present specification
means C₁₋₅ alkylene from among the above-mentioned "C₁₋₆
20 alkylene".

[0086]

Examples of the "optionally substituted C₁₋₁₀ (or C₁₋₆ or
C₁₋₅) alkyl", "optionally substituted C₂₋₁₀ (or C₂₋₆) alkenyl",
"optionally substituted C₂₋₁₀ (or C₂₋₆) alkynyl", "optionally
25 substituted C₁₋₆ alkoxy", "optionally substituted C₁₋₆
alkylsulfanyl" and "optionally substituted C₁₋₆ alkylene" in the
present specification include "C₁₋₁₀ (or C₁₋₆ or C₁₋₅) alkyl", "C₂₋₆
alkenyl", "C₂₋₁₀ (or C₂₋₆) alkynyl", "C₂₋₁₀ (or C₂₋₆) alkoxy", "C₁₋₆
alkylsulfanyl" and "C₁₋₆ alkylene", each of which optionally
30 has, at substitutable position(s), 1 to 5 substituents
selected from

- (1) a halogen atom;
- (2) hydroxy;
- (3) amino;
- 35 (4) nitro;

- (5) cyano;
- (6) a heterocyclic group optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, oxo, amino, nitro, cyano, an optionally halogenated C₁₋₆ alkyl, mono- or di-
5 C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl,
10 mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;
- (7) mono- or di-C₁₋₆ alkyl-amino;
- (8) mono- or di-C₆₋₁₄ aryl-amino;
- (9) mono- or di-C₇₋₁₄ aralkyl-amino;
- 15 (10) N-C₁₋₆ alkyl-N-C₆₋₁₄ aryl-amino;
- (11) N-C₁₋₆ alkyl-N-C₇₋₁₄ aralkyl-amino;
- (12) C₃₋₈ cycloalkyl;
- (13) optionally halogenated C₁₋₆ alkoxy;
- (14) C₁₋₆ alkylsulfanyl;
- 20 (15) C₁₋₆ alkylsulfinyl;
- (16) C₁₋₆ alkylsulfonyl;
- (17) optionally esterified carboxyl;
- (18) C₁₋₆ alkyl-carbonyl;
- (19) C₃₋₈ cycloalkyl-carbonyl;
- 25 (20) C₆₋₁₄ aryl-carbonyl;
- (21) carbamoyl;
- (22) thiocarbamoyl;
- (23) mono- or di-C₁₋₆ alkyl-carbamoyl;
- (24) mono- or di-C₆₋₁₄ aryl-carbamoyl;
- 30 (25) N-C₁₋₆ alkyl-N-C₆₋₁₄ aryl-carbamoyl;
- (26) mono- or di-5- to 7-membered heterocyclyl-carbamoyl;
- (27) C₁₋₆ alkyl-carbonylamino optionally substituted by carboxyl;
- (28) C₆₋₁₄ aryloxy optionally substituted by 1 to 3 substituents
35 selected from a halogen atom, hydroxy, amino, nitro, cyano, an

optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified
5 carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;
(29) a C₆₋₁₄ aryl optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, amino, nitro, cyano, an
10 optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified
carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-
15 carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;
(30) heterocyclyl-oxy;
(31) sulfamoyl;
(32) mono- or di-C₁₋₆ alkyl-sulfamoyl;
20 (33) mono- or di-C₆₋₁₄ aryl-sulfamoyl;
(34) C₇₋₁₄ aralkyloxy optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, amino, nitro, cyano, an optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈
25 cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-
30 sulfamoyl;
(35) C₁₋₆ alkyl-carbonyloxy;
(36) C₁₋₆ alkoxy-carbonyl;
and the like. When the number of the substituents is plural, the respective substituents may be the same or different.

35 [0087]

Examples of the "optionally substituted C₃₋₈ cycloalkyl", "optionally substituted C₃₋₈ cycloalkenyl", "optionally substituted C₆₋₁₄ aryl", "optionally substituted C₇₋₁₄ aralkyl", "optionally substituted heterocyclic group", "optionally substituted heterocyclyl-oxy", "optionally substituted C₆₋₁₄ aryloxy", "optionally substituted C₇₋₁₄ aralkyloxy", "optionally substituted heterocyclyl-sulfanyl", "optionally substituted C₆₋₁₄ arylsulfanyl" and "optionally substituted C₇₋₁₄ aralkylsulfanyl" in the present specification include "C₃₋₈ cycloalkyl", "C₃₋₈ cycloalkenyl", "C₆₋₁₄ aryl", "C₇₋₁₄ aralkyl", "heterocyclic group", "heterocyclyl-oxy", "C₆₋₁₄ aryloxy", "C₇₋₁₄ aralkyloxy", "heterocyclyl-sulfanyl", "C₆₋₁₄ arylsulfanyl" and "C₇₋₁₄ aralkylsulfanyl", each of which optionally have, at substitutable position(s), 1 to 5 substituents selected from

- (1) a halogen atom;
- (2) hydroxy;
- (3) amino;
- (4) nitro;
- (5) cyano;
- (6) optionally substituted C₁₋₆ alkyl;
- (7) optionally substituted C₂₋₆ alkenyl;
- (8) optionally substituted C₂₋₆ alkynyl;
- (9) C₆₋₁₄ aryl optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, amino, nitro, cyano, an optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;
- (10) C₆₋₁₄ aryloxy optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, amino, nitro, cyano, an optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆

alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;

(11) C₇₋₁₄ aralkyloxy optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, amino, nitro, cyano, an optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;

(12) a heterocyclic group optionally substituted by 1 to 3 substituents selected from a halogen atom, hydroxy, amino, nitro, cyano, an optionally halogenated C₁₋₆ alkyl, mono- or di-C₁₋₆ alkyl-amino, C₆₋₁₄ aryl, mono- or di-C₆₋₁₄ aryl-amino, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkoxy-C₁₋₆ alkoxy, C₁₋₆ alkylsulfanyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyl, optionally esterified carboxyl, carbamoyl, thiocarbamoyl, mono- or di-C₁₋₆ alkyl-carbamoyl, mono- or di-C₆₋₁₄ aryl-carbamoyl, sulfamoyl, mono- or di-C₁₋₆ alkyl-sulfamoyl and mono- or di-C₆₋₁₄ aryl-sulfamoyl;

(13) mono- or di-C₁₋₆ alkyl-amino;

(14) mono- or di-C₆₋₁₄ aryl-amino;

(15) mono- or di-C₇₋₁₄ aralkyl-amino;

(16) N-C₁₋₆ alkyl-N-C₆₋₁₄ aryl-amino;

(17) N-C₁₋₆ alkyl-N-C₇₋₁₄ aralkyl-amino;

(18) C₃₋₈ cycloalkyl;

(19) optionally substituted C₁₋₆ alkoxy;

(20) optionally substituted C₁₋₆ alkylsulfanyl;

(21) C₁₋₆ alkylsulfinyl;

(22) C₁₋₆ alkylsulfonyl;

- (23) optionally esterified carboxyl;
- (24) C₁₋₆ alkyl-carbonyl;
- (25) C₃₋₈ cycloalkyl-carbonyl;
- (26) C₆₋₁₄ aryl-carbonyl;
- 5 (27) carbamoyl;
- (28) thiocarbamoyl;
- (29) mono- or di-C₁₋₆ alkyl-carbamoyl;
- (30) mono- or di-C₆₋₁₄ aryl-carbamoyl;
- (31) N-C₁₋₆ alkyl-N-C₆₋₁₄ aryl-carbamoyl;
- 10 (32) mono- or di-5- to 7-membered heterocyclyl-carbamoyl;
- (33) sulfamoyl;
- (34) mono- or di-C₁₋₆ alkyl-sulfamoyl;
- (35) mono- or di-C₆₋₁₄ aryl-sulfamoyl;
- (36) C₁₋₆ alkyl-carbonylamino optionally substituted by
- 15 carboxyl;
- (37) heterocyclyl-oxy;
- (38) C₁₋₆ alkyl-carbonyloxy;
- (39) C₁₋₆ alkoxy-carbonyl;
- and the like. When the number of the substituents is plural,
- 20 the respective substituents may be the same or different.
- [0088]

- The "optionally substituted amino" in the present specification means, unless otherwise specified, amino optionally substituted by 1 or 2 substituents selected from
- 25 (1) optionally substituted C₁₋₆ alkyl;
 - (2) optionally substituted C₂₋₆ alkenyl;
 - (3) optionally substituted C₂₋₆ alkynyl;
 - (4) optionally substituted C₃₋₈ cycloalkyl;
 - (5) optionally substituted C₆₋₁₄ aryl;
 - 30 (6) optionally substituted C₁₋₆ alkoxy;
 - (7) optionally substituted acyl;
 - (8) optionally substituted heterocyclic group (preferably furyl, pyridyl, thienyl, pyrazolyl, thiazolyl, oxazolyl);
 - (9) sulfamoyl;
 - 35 (10) mono- or di-C₁₋₆ alkyl-sulfamoyl;

(11) mono- or di-C₆₋₁₄ aryl-sulfamoyl;
or the like. In addition, when the "optionally substituted
amino" is amino substituted by two substituents, these
substituents may be same or different, and these substituents
5 may form, together with the adjacent nitrogen atom, a
nitrogen-containing heterocycle. Examples of the "nitrogen-
containing heterocycle" include a 5- to 7-membered nitrogen-
containing heterocycle optionally containing, as a ring-
constituting atom besides carbon atom, at least one nitrogen
10 atom, and further, 1 or 2 hetero atoms selected from an oxygen
atom, a sulfur atom and a nitrogen atom. Preferable examples
of the nitrogen-containing heterocycle include pyrrolidine,
imidazolidine, pyrazolidine, piperidine, piperazine,
morpholine, thiomorpholine, thiazolidine, oxazolidine and the
15 like.

[0089]

The "optionally substituted aminocarbonyl" in the present
specification means, unless otherwise specified, aminocarbonyl
wherein the "optionally substituted amino" moiety is the
20 above-mentioned "optionally substituted amino".

[0090]

The "optionally substituted acyl" in the present
specification means, unless otherwise specified, a group
represented by the formula: -COR¹⁸, -CO-OR¹⁸, -SO₂R¹⁸, -SOR¹⁸, -
25 PO(OR¹⁸)(OR¹⁹), -CO-NR^{18a}R^{19a} and -CS-NR^{18a}R^{19a} wherein R¹⁸ and R¹⁹
are the same or different and each is a hydrogen atom, an
optionally substituted hydrocarbon group or an optionally
substituted heterocyclic group, R^{18a} and R^{19a} are the same or
different and each is a hydrogen atom, an optionally
30 substituted hydrocarbon group or an optionally substituted
heterocyclic group, R^{18a} and R^{19a} may form, together with the
adjacent nitrogen atom, an optionally substituted nitrogen-
containing heterocycle, or the like.

[0091]

35 Examples of the "nitrogen-containing heterocycle" of the

"optionally substituted nitrogen-containing heterocycle"
formed by R^{18a} and R^{19a} together with the adjacent nitrogen atom
include a 5- to 7-membered nitrogen-containing heterocycle
optionally containing, as a ring-constituting atom besides
5 carbon atom, at least one nitrogen atom, and further, 1 or 2
hetero atoms selected from an oxygen atom, a sulfur atom and a
nitrogen atom. Preferable examples of the nitrogen-containing
heterocycle include pyrrolidine, imidazolidine, pyrazolidine,
piperidine, piperazine, morpholine, thiomorpholine,
10 thiazolidine, oxazolidine and the like.

[0092]

The nitrogen-containing heterocycle may have 1 or 2
substituents at substitutable position(s). Examples of such
substituent include hydroxy, an optionally halogenated C₁₋₆
15 alkyl, C₆₋₁₄ aryl, C₇₋₁₄ aralkyl and the like. When the number of
the substituents is 2, these substituents may be the same or
different.

[0093]

Preferable examples of the "optionally substituted acyl"
20 include
formyl;
carboxyl;
carbamoyl;
C₁₋₆ alkyl-carbonyl;
25 C₁₋₆ alkoxy-carbonyl;
C₃₋₈ cycloalkyl-carbonyl;
C₆₋₁₄ aryl-carbonyl;
C₇₋₁₄ aralkyl-carbonyl;
C₆₋₁₄ aryloxy-carbonyl;
30 C₇₋₁₄ aralkyloxy-carbonyl;
mono- or di-C₁₋₆ alkyl-carbamoyl;
mono- or di-C₆₋₁₄ aryl-carbamoyl;
mono- or di-C₃₋₈ cycloalkyl-carbamoyl;
mono- or di-C₇₋₁₄ aralkyl-carbamoyl;
35 C₁₋₆ alkylsulfonyl;

C₆₋₁₄ arylsulfonyl optionally substituted by nitro;
 nitrogen-containing heterocyclyl-carbonyl;
 C₁₋₆ alkylsulfinyl;
 C₆₋₁₄ arylsulfinyl;
 5 thiocarbamoyl;
 sulfamoyl;
 mono- or di-C₁₋₆ alkyl-sulfamoyl;
 mono- or di-C₆₋₁₄ aryl-sulfamoyl;
 mono- or di-C₇₋₁₄ aralkyl-sulfamoyl;
 10 and the like.

[0094]

The definition of each symbol in the formula (1) is explained in detail in the following.

R¹ is a hydrogen atom or an optionally substituted C₁₋₆
 15 alkyl group.

R¹ is preferably a hydrogen atom or a C₁₋₆ alkyl group.

R¹ is more preferably a hydrogen atom or a C₁₋₃ alkyl group (preferably methyl).

R¹ is particularly preferably a hydrogen atom.

20 [0095]


R² and R³ are each independently a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, or R² and R³ in combination optionally form, together with the adjacent carbon atom, an optionally substituted ring.

25 R² and R³ are preferably each independently a hydrogen atom or a C₁₋₆ alkyl group (e.g., methyl).

R² and R³ are more preferably both hydrogen atoms.

Examples of the "optionally substituted ring" formed by R² and R³ in combination together with the adjacent carbon atom
 30 include a ring represented by

[0096]



[0097]

wherein R²⁰ is an optionally substituted C₁₋₆ alkyl group, and the like.

Examples of the substituent for "ring" of the "optionally substituted ring" formed by R² and R³ in combination together with the adjacent carbon atom include those similar to the substituent for the "heterocyclic group" of the above-mentioned "optionally substituted heterocyclic group".

[0098]

10 X is a bond or an optionally substituted C₁₋₆ alkylene group.

The "C₁₋₆ alkylene group" of the "optionally substituted C₁₋₆ alkylene group" for X is preferably a C₁₋₅ alkylene group, more preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)-
15 or -CH(CH(CH₃)₂)-.

In another embodiment, the "C₁₋₆ alkylene group" of the "optionally substituted C₁₋₆ alkylene group" for X is more preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)-, -CH(CH(CH₃)₂)-, -CH(C(CH₃)₃)-, -CH(CH₂CH₂CH₂CH₃)-, -
20 CH(CH(CH₂CH₃)₂)- or -CH₂CH₂-.

In yet another embodiment, the "C₁₋₆ alkylene group" of the "optionally substituted C₁₋₆ alkylene group" for X is more preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)-, -CH(CH(CH₃)₂)-, -CH(C(CH₃)₃)-, -CH(CH₂CH₂CH₂CH₃)- or -
25 CH(CH(CH₂CH₃)₂)-.

X is preferably an optionally substituted C₁₋₆ alkylene group (preferably an optionally substituted C₁₋₅ alkylene group).

X is more preferably a C₁₋₆ alkylene group (preferably a C₁₋₅ alkylene group, more preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)- or -CH(CH(CH₃)₂)-) optionally substituted by 1 to 3 substituents selected from

- (1) hydroxy,
- (2) cyano,
- (3) carbamoyl,
- 35 (4) C₁₋₆ alkoxy (e.g., methoxy),

- (5) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
- (6) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
- (7) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
- (8) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- 5 (9) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from optionally halogenated C₁₋₆ alkoxy (e.g., methoxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)), and
- (10) a heterocyclic group (preferably a 5- or 6-membered
- 10 aromatic heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl).

In another embodiment, X is more preferably a C₁₋₆ alkylene group (preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)-, -CH(CH(CH₃)₂)-, -CH(C(CH₃)₃)-, -CH(CH₂CH₂CH₂CH₃)-, -CH(CH(CH₂CH₃)₂)- or -CH₂CH₂-) optionally substituted by 1 to 5 substituents selected from

- (1) a halogen atom (e.g., a fluorine atom),
- (2) hydroxy,
- 20 (3) amino,
- (4) cyano,
- (5) carbamoyl,
- (6) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl, dimethylcarbamoyl),
- 25 (7) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) C₁₋₆ alkoxy (e.g., methoxy),
- (8) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
- 30 (9) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),
- (10) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
- (11) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
- (12) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- (13) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3
- 35 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom, a bromine atom),
(ii) hydroxy, and
(iii) optionally halogenated C₁₋₆ alkoxy (e.g., methoxy)
(preferably optionally substituted by 1 to 3 halogen atoms
5 (e.g., a fluorine atom)),
(14) a heterocyclic group (preferably a 4- to 10-membered
heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl,
pyrrolidinyl, triazolyl, oxetanyl, tetrahydrofuryl,
tetrahydropyranyl, dihydroisoindolyl) optionally substituted
10 by 1 to 3 substituents selected from
(i) C₁₋₆ alkyl (e.g., methyl),
(ii) hydroxy, and
(iii) oxo,
(15) C₃₋₈ cycloalkyloxy (e.g., cyclopentyloxy),
15 (16) optionally halogenated heterocyclyl-oxy (e.g.,
pyridyloxy) (preferably optionally substituted by 1 to 3
halogen atoms (e.g., a bromine atom)),
(17) C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy),
(18) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-
20 butyl(diphenyl)silyloxy),
(19) C₁₋₆ alkyl-carbonyl (e.g., acetyl), and
(20) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl),
or a bond.

In yet another embodiment, X is more preferably a C₁₋₆
25 alkylene group (preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -
CH(CH₂CH₂CH₃)-, -CH(CH(CH₃)₂)-, -CH(C(CH₃)₃)-, -CH(CH₂CH₂CH₂CH₃) -
or -CH(CH(CH₂CH₃)₂)) optionally substituted by 1 to 5
substituents selected from
(1) a halogen atom (e.g., a fluorine atom),
30 (2) hydroxy,
(3) amino,
(4) cyano,
(5) carbamoyl,
(6) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl,
35 dimethylcarbamoyl),

- (7) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from
- (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) C₁₋₆ alkoxy (e.g., methoxy),
- 5 (8) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
- (9) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),
- (10) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
- (11) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
- (12) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- 10 (13) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from
- (i) a halogen atom (e.g., a fluorine atom, a bromine atom),
 - (ii) hydroxy, and
 - (iii) optionally halogenated C₁₋₆ alkoxy (e.g., methoxy)
- 15 (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
- (14) a heterocyclic group (preferably a 4- to 10-membered heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl, pyrrolidinyl, triazolyl, oxetanyl, tetrahydrofuryl,
- 20 tetrahydropyranyl, dihydroisoindolyl) optionally substituted by 1 to 3 substituents selected from
- (i) C₁₋₆ alkyl (e.g., methyl),
 - (ii) hydroxy, and
 - (iii) oxo,
- 25 (15) C₃₋₈ cycloalkyloxy (e.g., cyclopentyloxy),
- (16) optionally halogenated heterocyclyl-oxy (e.g., pyridyloxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a bromine atom)),
- (17) C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy),
- 30 (18) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-butyl(diphenyl)silyloxy),
- (19) C₁₋₆ alkyl-carbonyl (e.g., acetyl), and
- (20) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl).

X is still more preferably a C₁₋₆ alkylene group

35 (preferably a C₁₋₅ alkylene group, more preferably -CH(CH₃)- or

-CH(CH(CH₃)₂)-) optionally substituted by 1 to 3 substituents selected from

(1) hydroxy, and

(2) C₁₋₆ alkoxy (e.g., methoxy).

5 [0099]

A is an optionally substituted cyclic group.

The "optionally substituted cyclic group" for A is "optionally substituted C₆₋₁₄ aryl", "optionally substituted C₃₋₈ cycloalkyl", "optionally substituted C₃₋₈ cycloalkenyl" or
10 "optionally substituted heterocyclic group". The "cyclic group" is optionally fused, for example, with a C₆₋₁₄ aromatic hydrocarbon, a C₃₋₈ cycloalkane, a C₃₋₈ cycloalkene, a heterocycle or the like.

The "cyclic group" of the "optionally substituted cyclic
15 group" for A is preferably a C₆₋₁₄ aryl group or a 5- or 6-membered aromatic heterocyclic group, more preferably a C₆₋₁₀ aryl group or a 5- or 6-membered nitrogen-containing aromatic heterocyclic group, still more preferably phenyl, pyridazinyl, pyridyl, pyrimidinyl, isoxazolyl or pyrazolyl, particularly
20 preferably phenyl.

A is preferably a C₆₋₁₀ aryl group or a 5- or 6-membered nitrogen-containing aromatic heterocyclic group, each of which is optionally substituted, more preferably an optionally substituted phenyl group.

25 A is more preferably a C₆₋₁₀ aryl group or a 5- or 6-membered aromatic heterocyclic group (preferably a C₆₋₁₀ aryl group or a 5- or 6-membered nitrogen-containing aromatic heterocyclic group, more preferably phenyl, pyridazinyl, pyridyl, pyrimidinyl, isoxazolyl or pyrazolyl, particularly
30 preferably phenyl), each of which is optionally substituted by 1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom),

(2) hydroxy,

(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, isopropyl)
35 (preferably optionally substituted by 1 to 3 halogen atoms

(e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3 substituents selected from

5 (i) a halogen atom (e.g., a fluorine atom), and

(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl), and

(7) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, dihydropyranyl,

10 tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl).

In another embodiment, A is more preferably a C₆₋₁₀ aryl group, a 4- to 10-membered heterocyclic group or a C₃₋₈ cycloalkyl group (preferably phenyl, pyridazinyl, pyridyl, 15 pyrimidinyl, isoxazolyl, pyrazolyl, oxetanyl, pyrrolidinyl, tetrahydrofuryl, imidazolyl, triazolyl, thienyl, tetrahydropyranyl, imidazo[1,2-a]pyridyl, benzodioxolyl, pyrazolo[1,5-a]pyridyl, dihydroisoindolyl or cyclopropyl, more preferably phenyl), each of which is optionally substituted by 20 1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom),

(2) hydroxy,

(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, 25 isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from

30 (i) a halogen atom (e.g., a fluorine atom), and

(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by optionally halogenated C₁₋₆ alkoxy (e.g., methoxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a 35 fluorine atom)),

- (7) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl, pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl),
5 (8) C₃₋₈ cycloalkyl (e.g., cyclopropyl), and
(9) oxo.

In yet another embodiment, A is more preferably a C₆₋₁₀ aryl group or a 5- to 10-membered aromatic heterocyclic group (preferably phenyl, pyridazinyl, pyridyl, pyrimidinyl,
10 isoxazolyl, pyrazolyl, imidazolyl, triazolyl, thienyl, imidazo[1,2-a]pyridyl or pyrazolo[1,5-a]pyridyl, more preferably phenyl), each of which is optionally substituted by 1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a
15 bromine atom),

(2) hydroxy,

(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

20 (4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

25 (6) C₆₋₁₄ aryl (e.g., phenyl),

(7) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl, pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), and

30 (8) C₃₋₈ cycloalkyl (e.g., cyclopropyl).

In yet another embodiment, A is more preferably a C₆₋₁₀ aryl group (preferably phenyl), each of which is optionally substituted by 1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a
35 bromine atom),


- (2) hydroxy,
(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
5 (4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),
(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from
(i) a halogen atom (e.g., a fluorine atom), and
(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
10 (6) C₆₋₁₄ aryl (e.g., phenyl),
(7) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl, pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), and
15 (8) C₃₋₈ cycloalkyl (e.g., cyclopropyl).

A is particularly preferably a C₆₋₁₀ aryl group (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from

- (1) a halogen atom (e.g., a fluorine atom), and
20 (2) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom).

Preferable examples of the group represented by


[0100]



25 [0101]

include a group represented by

[0102]



[0103]


wherein

R^{21} is a hydrogen atom, an optionally substituted C_{1-5} alkyl group, a C_{3-8} cycloalkyl group, a carbamoyl group or a cyano group, and

R^{22} is a hydrogen atom or a substituent.

Another preferable examples of the group represented by


[0104]



[0105]

include a group represented by

[0106]



[0107]

wherein

R^{21} is a hydrogen atom, an optionally substituted C_{1-5} alkyl group, a C_{3-8} cycloalkyl group, a carbamoyl group, a cyano group or an optionally substituted heterocyclic group,


R^{22} is a hydrogen atom or a substituent, and

p is an integer of 1 to 5.

The "substituent" for R^{22} corresponds to the "substituent" of the "optionally substituted cyclic group" for A.

In another embodiment, another preferable examples of the group represented by


[0108]



[0109]

include a group represented by

10 [0110]



[0111]

wherein


R^{21} is a hydrogen atom, an optionally substituted C_{1-6} alkyl group, a C_{3-8} cycloalkyl group, an optionally substituted carbamoyl group, a cyano group, an optionally substituted heterocyclic group, a C_{1-6} alkyl-carbonyl group, a C_{3-8} cycloalkyl-carbonyl group or an optionally substituted C_{6-14} aryl group,

20 R^{22} is a hydrogen atom or a substituent, and p is an integer of 1 to 5.

The "substituent" for R^{22} corresponds to the "substituent" of the "optionally substituted cyclic group" for A.

25 The group represented by


[0112]



[0113]

is more preferably

[0114]



5

[0115]

wherein

R^{21} is

- (1) cyano
- 10 (2) carbamoyl
- (3) C_{1-5} alkyl (e.g., methyl, ethyl, isopropyl, propyl)
optionally substituted by 1 to 3 substituents selected from
 - (i) hydroxy,
 - (ii) C_{1-6} alkoxy (e.g., methoxy),
 - 15 (iii) mono- or di- C_{1-6} alkyl-amino (e.g., dimethylamino),
 - (iv) C_{1-6} alkylsulfanyl (e.g., methylsulfanyl), and
 - (v) C_{1-6} alkylsulfonyl (e.g., methylsulfonyl),
 - (4) C_{3-8} cycloalkyl (e.g., cyclopropyl), or
 - (5) a heterocyclic group (preferably a 5- or 6-membered
20 aromatic heterocyclic group) (e.g., pyridyl, isoxazolyl,
pyrazolyl) optionally substituted by 1 to 3 C_{1-6} alkyl (e.g.,
methyl),

R^{22} is

- (1) a halogen atom (e.g., a fluorine atom),
- 25 (2) hydroxy,
- (3) optionally halogenated C_{1-6} alkyl (e.g., methyl, isopropyl)

(preferably optionally substituted by 1 to 3 halogen atoms
(e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3
5 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and


(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl), or

(7) a heterocyclic group (preferably a 5- or 6-membered
10 heterocyclic group) (e.g., pyrazolyl, dihydropyranyl,
tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl
(e.g., methyl), and

p is an integer of 1 to 3.


In another embodiment, the group represented by
15 [0116]



[0117]

is more preferably

[0118]



[0119]

wherein

R²¹ is

(1) cyano,

25 (2) carbamoyl,

(3) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl,
dimethylcarbamoyl),

- (4) C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl, propyl, butyl, tert-butyl, 1-ethylpropyl) optionally substituted by 1 to 5 substituents selected from
- (i) a halogen atom (e.g., a fluorine atom),
 - 5 (ii) hydroxy,
 - (iii) amino,
 - (iv) cyano,
 - (v) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
 - (vi) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy)
 - 10 optionally substituted by 1 to 3 substituents selected from
 - (a) a halogen atom (e.g., a fluorine atom), and
 - (b) C₁₋₆ alkoxy (e.g., methoxy),
 - (vii) C₃₋₈ cycloalkyloxy (e.g., cyclopentyloxy),
 - (viii) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
 - 15 (ix) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
 - (x) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),
 - (xi) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
 - (xii) carbamoyl,
 - (xiii) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g.,
 - 20 dimethylcarbamoyl),
 - (xiv) a heterocyclic group (e.g., oxetanyl, pyrrolidinyl, pyrazolyl, triazolyl, tetrahydropyranyl, dihydroisoindolyl) optionally substituted by 1 to 3 substituents selected from
 - (a) hydroxy, and
 - 25 (b) oxo,
 - (xv) C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy),
 - (xvi) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-butyl(diphenyl)silyloxy), and
 - (xvii) optionally halogenated heterocyclyl-oxy (e.g.,
 - 30 pyridyloxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a bromine atom)),
- (5) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- (6) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl,
- 35 oxetanyl, tetrahydrofuryl, tetrahydropyranyl) optionally

substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl),
(7) C₁₋₆ alkyl-carbonyl (e.g., acetyl),
(8) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl), or
(9) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3
5 substituents selected from

(i) a halogen atom (e.g., a fluorine atom, a bromine atom),
and

(ii) hydroxy,

R²² is

10 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a
bromine atom),

(2) hydroxy,

(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, isopropyl)
(preferably optionally substituted by 1 to 3 halogen atoms
15 (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally
substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

20 (ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl),


(7) a heterocyclic group (preferably a 4- to 6-membered
heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl,
pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl)
25 optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or

(8) C₃₋₈ cycloalkyl (e.g., cyclopropyl), and

p is an integer of 1 to 3.

The group represented by


[0120]



[0121]

is still more preferably a group represented by

[0122]



[0123]

wherein

5 R^{21} is a C_{1-5} alkyl group (e.g., methyl, isopropyl) optionally substituted by hydroxyl or C_{1-6} alkoxy (e.g., methoxy),

R^{22} is

(1) a halogen atom (e.g., a fluorine atom), or

(2) optionally halogenated C_{1-6} alkoxy (e.g., methoxy)

10 (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)), and

p is an integer of 1 or 2.

[0124]

Z^1 is a group represented by CR^{Z1} wherein R^{Z1} is a
15 hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group or a cyano group, or a nitrogen atom.

20 In another embodiment, Z^1 is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally
25 substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom.

Z^1 is preferably a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group,
30 an optionally substituted C_{1-6} alkoxy group or a cyano group.

In another embodiment, Z^1 is preferably a group represented by CR^{Z^1} wherein R^{Z^1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{1-6} alkoxy group, a cyano group or an optionally substituted cyclic group, more preferably a group represented by CR^{Z^1} wherein R^{Z^1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{1-6} alkoxy group, a cyano group, an optionally substituted C_{6-14} aryl group, an optionally substituted heterocyclic group or an optionally substituted C_{3-8} cycloalkyl group.

Z^1 is more preferably a group represented by CR^{Z^1} wherein R^{Z^1} is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- (3) a C_{1-6} alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) C_{6-14} aryl (e.g., phenyl),
- (4) a C_{2-6} alkenyl group (e.g., vinyl),
- (5) a C_{1-6} alkoxy group (e.g., methoxy), or
- (6) a cyano group.

In another embodiment, Z^1 is more preferably a group represented by CR^{Z^1} wherein R^{Z^1} is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- (3) a C_{1-6} alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) C_{6-14} aryl (e.g., phenyl),
- (4) a C_{2-6} alkenyl group (e.g., vinyl),
- (5) a C_{1-6} alkoxy group (e.g., methoxy),
- (6) a cyano group,
- (7) a C_{6-14} aryl group (e.g., phenyl) optionally substituted by

1 to 3 C₁₋₆ alkoxy (e.g., methoxy),
(8) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, thienyl, furyl, isoxazolyl, pyridyl, dihydropyranyl, tetrahydropyranyl, 5 tetrahydropyridyl, piperidyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or
(9) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl).

Z¹ is still more preferably
a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom.
10 [0125]

Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally
15 substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom.

Z² is preferably a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted
20 C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, a cyano group or an optionally substituted cyclic group.

Z² is more preferably a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, a cyano group, an optionally
25 substituted C₆₋₁₄ aryl group or an optionally substituted C₃₋₈ cycloalkyl group.

Z² is particularly preferably
a group represented by CR^{Z2} wherein R^{Z2} is
(1) a hydrogen atom,
30 (2) a halogen atom (e.g., a bromine atom, an iodine atom),
(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
(4) a C₂₋₆ alkenyl group (e.g., vinyl),
35 (5) a cyano group,

- (6) a C₆₋₁₄ aryl group (e.g., phenyl), or
(7) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl).

Z² is preferably a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted
5 C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₁₋₆ alkoxy group, a cyano group or an optionally substituted cyclic group.

In another embodiment, Z² is preferably a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen
10 atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₁₋₆ alkoxy group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom.

Z² is more preferably a group represented by CR^{Z2} wherein
15 R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₁₋₆ alkoxy group, a cyano group, an optionally substituted C₆₋₁₄ aryl group or an optionally substituted C₃₋₈ cycloalkyl group.

In another embodiment, Z² is more preferably a group
20 represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₁₋₆ alkoxy group, a cyano group, an optionally substituted C₆₋₁₄ aryl group, an optionally substituted heterocyclic group or an
25 optionally substituted C₃₋₈ cycloalkyl group, or a nitrogen atom.

Z² is particularly preferably
a group represented by CR^{Z2} wherein R^{Z2} is
(1) a hydrogen atom,
(2) a halogen atom (e.g., a bromine atom, an iodine atom),
30 (3) optionally halogenated C₁₋₆ alkyl group (e.g., methyl, ethyl, isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
(4) a C₂₋₆ alkenyl group (e.g., vinyl),
(5) a C₁₋₆ alkoxy group (e.g., methoxy),
35 (6) a cyano group,

- (7) a C₆₋₁₄ aryl group (e.g., phenyl), or
(8) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl).

In another embodiment, Z² is particularly preferably a group represented by CR^{Z2} wherein R^{Z2} is

- 5 (1) a hydrogen atom,
(2) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom, an iodine atom),
(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl, isopropyl) optionally substituted by 1 to 3 substituents selected from
10 (i) a halogen atom (e.g., a fluorine atom), and
(ii) hydroxy,
(4) a C₂₋₆ alkenyl group (e.g., vinyl),
(5) an optionally halogenated C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) (preferably optionally substituted by 1 to
15 3 halogen atoms (e.g., a fluorine atom)),
(6) a cyano group,
(7) a C₆₋₁₄ aryl group (e.g., phenyl),
(8) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl), or
(9) a heterocyclic group (e.g., azetidiny, pyrazolyl)
20 optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or a nitrogen atom.

Z² is most preferably

a group represented by CR^{Z2} wherein R^{Z2} is

- (1) a C₁₋₆ alkoxy group (e.g., methoxy), or
25 (2) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl).

In another embodiment, Z² is most preferably a group represented by CR^{Z2} wherein R^{Z2} is an optionally substituted C₁₋₆ alkoxy group, an optionally substituted C₁₋₆ alkyl group or an optionally substituted C₃₋₈ cycloalkyl group.

- 30 In another embodiment, Z² is most preferably a group represented by CR^{Z2} wherein R^{Z2} is
(1) a C₁₋₆ alkoxy group (e.g., methoxy),
(2) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl), or
(3) a C₁₋₆ alkyl group (e.g., methyl).

35 [0126]

Z^3 is a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom.

In another embodiment, Z^3 is a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C_{1-6} alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom.

Z^3 is preferably a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom or an optionally substituted C_{1-6} alkyl group, or a nitrogen atom.

In another embodiment, Z^3 is preferably a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{1-6} alkoxy group or an optionally substituted C_{1-6} alkyl-carbonyl group, or a nitrogen atom.

Z^3 is more preferably a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom or a C_{1-6} alkyl group (e.g., methyl), or a nitrogen atom.

In another embodiment, Z^3 is more preferably a group represented by CR^{Z^3} wherein R^{Z^3} is

- (1) a hydrogen atom,
- (2) a C_{1-6} alkyl group (e.g., methyl, isopropyl) optionally substituted by 1 to 3 hydroxy groups,
- (3) a C_{1-6} alkoxy group (e.g., methoxy), or
- (4) a C_{1-6} alkyl-carbonyl group (e.g., acetyl),

or a nitrogen atom.

Z^3 is still more preferably a group represented by CR^{Z^3}

wherein R^{Z3} is a hydrogen atom.

[0127]

B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms.

5 B is preferably a 6-membered nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms.

B is more preferably a pyridine ring or a pyrazine ring.

In another embodiment, B is more preferably a pyridine ring, a pyrimidine ring or a pyrazine ring.

10 B is particularly preferably a pyridine ring.

[0128]

Preferable examples of compound (1) include the following compounds.

[Compound A-1]

15 compound (1) wherein

R^1 is a hydrogen atom or an optionally substituted C_{1-6} alkyl group;

R^2 and R^3 are a hydrogen atom or an optionally substituted C_{1-6} alkyl group;

20 X is an optionally substituted C_{1-6} alkylene group (preferably an optionally substituted C_{1-5} alkylene group);

A is a C_{6-10} aryl group or a 5- or 6-membered nitrogen-containing aromatic heterocyclic group, each of which is optionally substituted;

25 Z^1 is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{1-6} alkoxy group or a cyano group;

Z^2 is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom,
30 a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, a cyano group, an optionally substituted C_{6-14} aryl group or an optionally substituted C_{3-8} cycloalkyl group;

Z^3 is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom
35 or an optionally substituted C_{1-6} alkyl group, or a nitrogen

atom; and

B is a 6-membered nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms.

[0129]

5 [Compound A-2]

compound (1) wherein

R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group;

R² and R³ are a hydrogen atom or an optionally substituted C₁₋₆
10 alkyl group;

X is an optionally substituted C₁₋₆ alkylene group (preferably an optionally substituted C₁₋₅ alkylene group);

A is a C₆₋₁₀ aryl group or a 5- or 6-membered nitrogen-containing aromatic heterocyclic group, each of which is
15 optionally substituted;

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₁₋₆ alkoxy group or a cyano group;

20 Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₁₋₆ alkoxy group, a cyano group, an optionally substituted C₆₋₁₄ aryl group or an optionally substituted C₃₋₈
25 cycloalkyl group;

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, or a nitrogen atom; and

B is a 6-membered nitrogen-containing aromatic heterocycle
30 containing 1 or 2 nitrogen atoms.

[0130]

[Compound A-3]

compound (1) wherein

R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl
35 group;

R^2 and R^3 are a hydrogen atom or an optionally substituted C_{1-6} alkyl group;
 X is an optionally substituted C_{1-6} alkylene group (preferably an optionally substituted C_{1-5} alkylene group);
5 A is a C_{6-10} aryl group, a 4- to 10-membered heterocyclic group or a C_{3-8} cycloalkyl group, each of which is optionally substituted;
 Z^1 is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an
10 optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{1-6} alkoxy group or a cyano group;
 Z^2 is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally
15 substituted C_{1-6} alkoxy group, a cyano group, an optionally substituted C_{6-14} aryl group or an optionally substituted C_{3-8} cycloalkyl group, or a nitrogen atom;
 Z^3 is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom or an optionally substituted C_{1-6} alkyl group, or a nitrogen
20 atom; and
 B is a 6-membered nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms.
[0131]
[Compound A-4]
25 compound (1) wherein
 R^1 is a hydrogen atom,
 R^2 and R^3 are both hydrogen atoms,
 X is an optionally substituted C_{1-6} alkylene group,
 A is an optionally substituted phenyl group,
30 Z^1 is CH ,
 Z^2 is a group represented by CR^{Z2} wherein R^{Z2} is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{1-6} alkoxy group or an optionally substituted C_{3-8} cycloalkyl group,
and
35 Z^3 is CH .

[0132]

[Compound B-1]

compound (1) wherein

R¹ is a hydrogen atom or a C₁₋₆ alkyl group;

5 R² and R³ are both hydrogen atoms;

X is a C₁₋₆ alkylene group (preferably a C₁₋₅ alkylene group, more preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)- or -CH(CH(CH₃)₂)-) optionally substituted by 1 to 3 substituents selected from

10 (1) hydroxy,

(2) cyano,

(3) carbamoyl,

(4) C₁₋₆ alkoxy (e.g., methoxy),

(5) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),

15 (6) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),

(7) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),

(8) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(9) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from optionally halogenated C₁₋₆ alkoxy

20 (e.g., methoxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)), and

(10) a heterocyclic group (preferably a 5- or 6-membered aromatic heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g.,
25 methyl);

A is a C₆₋₁₀ aryl group or a 5- or 6-membered aromatic heterocyclic group (preferably a C₆₋₁₀ aryl group or a 5- or 6-membered nitrogen-containing aromatic heterocyclic group, more preferably phenyl, pyridazinyl, pyridyl, pyrimidinyl,
30 isoxazolyl or pyrazolyl, particularly preferably phenyl), each of which is optionally substituted by 1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom),

(2) hydroxy,

35 (3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, isopropyl)

(preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and


(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl), and

(7) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, dihydropyranyl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl);

[preferably the group represented by


[0133]



[0134]

is a group represented by

[0135]



[0136]

wherein

R²¹ is a hydrogen atom, an optionally substituted C₁₋₅ alkyl group, a C₃₋₈ cycloalkyl group, a carbamoyl group or a cyano group, and

25 R²² is a hydrogen atom or a substituent]

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is

(1) a hydrogen atom,

(2) a halogen atom (e.g., a chlorine atom),
(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
- 5 (ii) a C₆₋₁₄ aryl group (e.g., phenyl),
- (4) a C₂₋₆ alkenyl group (e.g., vinyl),
- (5) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (6) a cyano group;

Z² is a group represented by CR^{Z2} wherein R^{Z2} is

- 10 (1) a hydrogen atom,
 - (2) a halogen atom (e.g., a bromine atom, an iodine atom),
 - (3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
 - 15 (4) a C₂₋₆ alkenyl group (e.g., vinyl),
 - (5) a cyano group,
 - (6) a C₆₋₁₄ aryl group (e.g., phenyl), or
 - (7) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl);
- Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom
20 or a C₁₋₆ alkyl group, or a nitrogen atom; and
B is a pyridine ring or a pyrazine ring.

[0137]

[Compound B-2]

compound (1) wherein


- 25 R¹ is a hydrogen atom or a C₁₋₆ alkyl group (preferably methyl);
- R² and R³ are both hydrogen atoms;

X is a C₁₋₆ alkylene group (preferably a C₁₋₅ alkylene group, more preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)- or -CH(CH(CH₃)₂)-) optionally substituted by 1 to 3

30 substituents selected from

- (1) hydroxy,
- (2) cyano,
- (3) carbamoyl,
- (4) C₁₋₆ alkoxy (e.g., methoxy),
- 35 (5) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),


(6) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
 (7) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
 (8) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
 (9) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3
 5 substituents selected from optionally halogenated C₁₋₆ alkoxy
 (e.g., methoxy) (preferably optionally substituted by 1 to 3
 halogen atoms (e.g., a fluorine atom)), and
 (10) a heterocyclic group (preferably a 5- or 6-membered
 aromatic heterocyclic group) (e.g., pyridyl, isoxazolyl,
 10 pyrazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g.,
 methyl);
 A is a C₆₋₁₀ aryl group or a 5- or 6-membered aromatic
 heterocyclic group (preferably a C₆₋₁₀ aryl group or a 5- or 6-
 membered nitrogen-containing aromatic heterocyclic group, more
 15 preferably phenyl, pyridazinyl, pyridyl, pyrimidinyl,
 isoxazolyl or pyrazolyl, particularly preferably phenyl), each
 of which is optionally substituted by 1 to 3 substituents
 selected from
 (1) a halogen atom (e.g., a fluorine atom),
 20 (2) hydroxy,
 (3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, isopropyl)
 (preferably optionally substituted by 1 to 3 halogen atoms
 (e.g., a fluorine atom)),
 (4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),
 25 (5) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3
 substituents selected from
 (i) a halogen atom (e.g., a fluorine atom), and
 (ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
 (6) C₆₋₁₄ aryl (e.g., phenyl), and
 30 (7) a heterocyclic group (preferably a 5- or 6-membered
 heterocyclic group) (e.g., pyrazolyl, dihydropyranyl,
 tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl
 (e.g., methyl);
 [preferably the group represented by
 35 [0138]



[0139]

is a group represented by

[0140]



5

[0141]

wherein

R^{21} is

- (1) cyano,
- 10 (2) carbamoyl,
- (3) C_{1-5} alkyl (e.g., methyl, ethyl, isopropyl, propyl)
optionally substituted by 1 to 3 substituents selected from
 - (i) hydroxy,
 - (ii) C_{1-6} alkoxy (e.g., methoxy),
 - 15 (iii) mono- or di- C_{1-6} alkyl-amino (e.g., dimethylamino),
 - (iv) C_{1-6} alkylsulfanyl (e.g., methylsulfanyl), and
 - (v) C_{1-6} alkylsulfonyl (e.g., methylsulfonyl),
 - (4) C_{3-8} cycloalkyl (e.g., cyclopropyl), or
 - (5) a heterocyclic group (preferably a 5- or 6-membered
20 aromatic heterocyclic group) (e.g., pyridyl, isoxazolyl,
pyrazolyl) optionally substituted by 1 to 3 C_{1-6} alkyl (e.g.,
methyl),

R^{22} is

- (1) a halogen atom (e.g., a fluorine atom),
- 25 (2) hydroxy,
- (3) optionally halogenated C_{1-6} alkyl (e.g., methyl,
isopropyl) (preferably optionally substituted by 1 to 3

halogen atoms (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3 substituents selected from

5 (i) a halogen atom (e.g., a fluorine atom), and

(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl), or

(7) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, dihydropyranyl,

10 tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), and

p is an integer of 1 to 3],

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is

(1) a hydrogen atom,

15 (2) a halogen atom (e.g., a chlorine atom),

(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₆₋₁₄ aryl (e.g., phenyl),

20 (4) a C₂₋₆ alkenyl group (e.g., vinyl),

(5) a C₁₋₆ alkoxy group (e.g., methoxy), or

(6) a cyano group;

Z² is a group represented by CR^{Z2} wherein R^{Z2} is

(1) a hydrogen atom,

25 (2) a halogen atom (e.g., a bromine atom, an iodine atom),

(3) an optionally halogenated C₁₋₆ alkyl group (e.g., methyl, ethyl, isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

(4) a C₂₋₆ alkenyl group (e.g., vinyl),

30 (5) a C₁₋₆ alkoxy group (e.g., methoxy),

(6) a cyano group,

(7) a C₆₋₁₄ aryl group (e.g., phenyl), or

(8) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl);

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom

35 or a C₁₋₆ alkyl group (e.g., methyl), or a nitrogen atom; and

B is a pyridine ring or a pyrazine ring.

[0142]

[Compound B-3]

compound (1) wherein

5 R¹ is a hydrogen atom or a C₁₋₆ alkyl group (preferably methyl);
R² and R³ are each independently a hydrogen atom or a C₁₋₆ alkyl
group (preferably methyl);

X is a C₁₋₆ alkylene group (preferably C₁₋₆ alkylene group, more
preferably -CH₂-, -CH(CH₃)-, -CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)-, -

10 CH(CH(CH₃)₂)-, -CH(C(CH₃)₃)-, -CH(CH₂CH₂CH₂CH₃)-, -

CH(CH(CH₂CH₃)₂)- or -CH₂CH₂-) optionally substituted by 1 to 5
substituents selected from

(1) a halogen atom (e.g., a fluorine atom),

(2) hydroxy,

15 (3) amino,

(4) cyano,

(5) carbamoyl,

(6) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl,
dimethylcarbamoyl),

20 (7) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally
substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₁₋₆ alkoxy (e.g., methoxy),

(8) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),

25 (9) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),

(10) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),

(11) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),

(12) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(13) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3

30 substituents selected from

(i) a halogen atom (e.g., a fluorine atom, a bromine atom),

(ii) hydroxy, and


(iii) optionally halogenated C₁₋₆ alkoxy (e.g., methoxy)

(preferably optionally substituted by 1 to 3 halogen atoms


35 (e.g., a fluorine atom)),

(14) a heterocyclic group (preferably a 4- to 10-membered heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl, pyrrolidinyl, triazolyl, oxetanyl, tetrahydrofuryl, tetrahydropyranyl, dihydroisoindolyl) optionally substituted
5 by 1 to 3 substituents selected from
 (i) C₁₋₆ alkyl (e.g., methyl),
 (ii) hydroxy, and
 (iii) oxo,
(15) C₃₋₈ cycloalkyloxy (e.g., cyclopentyloxy),
10 (16) optionally halogenated heterocyclyl-oxy (e.g., pyridyloxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a bromine atom)),
(17) C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy),
(18) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-
15 butyl(diphenyl)silyloxy),
(19) C₁₋₆ alkyl-carbonyl (e.g., acetyl), and
(20) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl),
or a bond;
A is a C₆₋₁₀ aryl group, a 4- to 10-membered heterocyclic group
20 or a C₃₋₈ cycloalkyl group (preferably phenyl, pyridazinyl, pyridyl, pyrimidinyl, isoxazolyl, pyrazolyl, oxetanyl, pyrrolidinyl, tetrahydrofuryl, imidazolyl, triazolyl, thienyl, tetrahydropyranyl, imidazo[1,2-a]pyridyl, benzodioxolyl, pyrazolo[1,5-a]pyridyl, dihydroisoindolyl or cyclopropyl, more
25 preferably phenyl), each of which is optionally substituted by 1 to 3 substituents selected from
 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom),
 (2) hydroxy,
30 (3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl) (e.g., a fluorine atom) (preferably optionally substituted by 1 to 3 halogen atoms,
 (4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),
 (5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally
35 substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and
(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
(6) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by
optionally halogenated C₁₋₆ alkoxy (e.g., methoxy) (preferably
5 optionally substituted by 1 to 3 halogen atoms (e.g., a
fluorine atom)),
(7) a heterocyclic group (preferably a 4- to 6-membered
heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl,
pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl)
10 optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl),
(8) C₃₋₈ cycloalkyl (e.g., cyclopropyl), and
(9) oxo;
[preferably the group represented by
[0143]



is a group represented by
[0145]



wherein

R²¹ is

- (1) cyano,
- (2) carbamoyl,
- 25 (3) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl,
dimethylcarbamoyl),

(4) C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl, propyl, butyl, tert-butyl, 1-ethylpropyl) optionally substituted by 1 to 5 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom),
- 5 (ii) hydroxy,
- (iii) amino,
- (iv) cyano,
- (v) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- (vi) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy)
- 10 optionally substituted by 1 to 3 substituents selected from
 - (a) a halogen atom (e.g., a fluorine atom), and
 - (b) C₁₋₆ alkoxy (e.g., methoxy),
 - (vii) C₃₋₈ cycloalkyloxy (e.g., cyclopentyloxy),
 - 15 (viii) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
 - (ix) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
 - (x) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),
 - (xi) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
 - (xii) carbamoyl,
 - 20 (xiii) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., dimethylcarbamoyl),
 - (xiv) a heterocyclic group (e.g., oxetanyl, pyrrolidinyl, pyrazolyl, triazolyl, tetrahydropyranyl, dihydroisoindolyl) optionally substituted by 1 to 3
 - 25 substituents selected from
 - (a) hydroxy, and
 - (b) oxo,
 - (xv) C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy),
 - (xvi) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-butyl(diphenyl)silyloxy), and
 - 30 (xvii) optionally halogenated heterocyclyl-oxy (e.g., pyridyloxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a bromine atom)),
- (5) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- 35 (6) a heterocyclic group (preferably a 4- to 6-membered

heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl, oxetanyl, tetrahydrofuryl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl),

(7) C₁₋₆ alkyl-carbonyl (e.g., acetyl),

5 (8) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl), or

(9) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom, a bromine atom),

10 (ii) hydroxy,

(iii) optionally halogenated C₁₋₆ alkoxy (e.g., methoxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

R²² is

15 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom),

(2) hydroxy,

(3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, isopropyl) (preferably optionally substituted by 1 to 3
20 halogen atoms (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

25 (ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl),

(7) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., azetidyl, pyrazolyl, pyrrolyl, pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl)

30 optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or

(8) C₃₋₈ cycloalkyl (e.g., cyclopropyl), and

p is an integer of 1 to 3],

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is

35 (1) a hydrogen atom,

- (2) a halogen atom (e.g., a chlorine atom),
 - (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - 5 (ii) C₆₋₁₄ aryl (e.g., phenyl),
 - (4) a C₂₋₆ alkenyl group (e.g., vinyl),
 - (5) a C₁₋₆ alkoxy group (e.g., methoxy),
 - (6) a cyano group,
 - (7) a C₆₋₁₄ aryl group (e.g., phenyl) optionally substituted by
 - 10 1 to 3 C₁₋₆ alkoxy (e.g., methoxy),
 - (8) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, thienyl, furyl, isoxazolyl, pyridyl, dihydropyranyl, tetrahydropyranyl, tetrahydropyridyl, piperidyl) optionally substituted by 1 to 3
 - 15 C₁₋₆ alkyl (e.g., methyl), or
 - (9) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl);
- Z² is a group represented by CR^{Z2} wherein R^{Z2} is
- (1) a hydrogen atom,
 - (2) a halogen atom (e.g., a fluorine atom, a chlorine atom, a
 - 20 bromine atom, an iodine atom),
 - (3) C₁₋₆ alkyl (e.g., methyl, ethyl, isopropyl) optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) hydroxy,
 - 25 (4) a C₂₋₆ alkenyl group (e.g., vinyl),
 - (5) an optionally halogenated C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
 - (6) a cyano group,
 - 30 (7) a C₆₋₁₄ aryl group (e.g., phenyl),
 - (8) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl), or
 - (9) a heterocyclic group (e.g., azetidiny, pyrazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or a nitrogen atom;
 - 35 Z³ is a group represented by CR^{Z3} wherein R^{Z3} is

- (1) a hydrogen atom,
- (2) a C₁₋₆ alkyl group (e.g., methyl, isopropyl) optionally substituted by 1 to 3 hydroxy,
- (3) a C₁₋₆ alkoxy group (e.g., methoxy), or
- 5 (4) a C₁₋₆ alkyl-carbonyl group (e.g., acetyl),
or a nitrogen atom; and

B is a pyridine ring, a pyrimidine ring or a pyrazine ring.

[0147]

[Compound B-4]

10 compound (1) wherein

R¹ is a hydrogen atom or a C₁₋₆ alkyl group (preferably methyl);
R² and R³ are each independently a hydrogen atom or a C₁₋₆ alkyl
group (preferably methyl);

X is a C₁₋₆ alkylene group (preferably -CH₂-, -CH(CH₃)-, -
15 CH(CH₂CH₃)-, -CH(CH₂CH₂CH₃)-, -CH(CH(CH₃)₂)-, -CH(C(CH₃)₃)-, -
CH(CH₂CH₂CH₂CH₃)-, -CH(CH(CH₂CH₃)₂)- or -CH₂CH₂-) optionally
substituted by 1 to 5 substituents selected from

- (1) a halogen atom (e.g., a fluorine atom),
- (2) hydroxy,
- 20 (3) amino,
- (4) cyano,
- (5) carbamoyl,
- (6) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., methylcarbamoyl,
dimethylcarbamoyl),
- 25 (7) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally
substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) C₁₋₆ alkoxy (e.g., methoxy),
- (8) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
- 30 (9) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),
- (10) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
- (11) mono- or di-C₁₋₆ alkyl-amino (e.g., dimethylamino),
- (12) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
- (13) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3
35 substituents selected from

(i) a halogen atom (e.g., a fluorine atom, a bromine atom),
 (ii) hydroxy, and
 (iii) optionally halogenated C₁₋₆ alkoxy (e.g., methoxy)
 (preferably optionally substituted by 1 to 3 halogen atoms
 5 (e.g., a fluorine atom)),
 (14) a heterocyclic group (preferably a 4- to 10-membered
 heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl,
 pyrrolidinyl, triazolyl, oxetanyl, tetrahydrofuryl,
 tetrahydropyranyl, dihydroisoindolyl) optionally substituted
 10 by 1 to 3 substituents selected from
 (i) C₁₋₆ alkyl (e.g., methyl),
 (ii) hydroxy, and
 (iii) oxo,
 (15) C₃₋₈ cycloalkyloxy (e.g., cyclopentyloxy),
 15 (16) optionally halogenated heterocyclyl-oxy (e.g.,
 pyridyloxy) (preferably optionally substituted by 1 to 3
 halogen atoms (e.g., a bromine atom)),
 (17) C₁₋₆ alkyl-carbonyloxy (e.g., acetoxy),
 (18) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-
 20 butyl(diphenyl)silyloxy),
 (19) C₁₋₆ alkyl-carbonyl (e.g., acetyl), and
 (20) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl),
 or a bond;
 A is a C₆₋₁₀ aryl group, a 4- to 10-membered heterocyclic group
 25 or a C₃₋₈ cycloalkyl group (preferably phenyl, pyridazinyl,
 pyridyl, pyrimidinyl, isoxazolyl, pyrazolyl, oxetanyl,
 pyrrolidinyl, tetrahydrofuryl, imidazolyl, triazolyl, thienyl,
 tetrahydropyranyl, imidazo[1,2-a]pyridyl, benzodioxolyl,
 pyrazolo[1,5-a]pyridyl, dihydroisoindolyl or cyclopropyl, more
 30 preferably phenyl), each of which is optionally substituted by
 1 to 3 substituents selected from
 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a
 bromine atom),
 (2) hydroxy,
 35 (3) optionally halogenated C₁₋₆ alkyl (e.g., methyl, ethyl,

isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally

5 substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by optionally halogenated C₁₋₆ alkoxy (e.g., methoxy) (preferably

10 optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

(7) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl, pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl)

15 optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl),

(8) C₃₋₈ cycloalkyl (e.g., cyclopropyl), and

(9) oxo;

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is

(1) a hydrogen atom,

20 (2) a halogen atom (e.g., a chlorine atom),

(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₆₋₁₄ aryl (e.g., phenyl),

25 (4) a C₂₋₆ alkenyl group (e.g., vinyl),

(5) a C₁₋₆ alkoxy group (e.g., methoxy),

(6) a cyano group,

(7) a C₆₋₁₄ aryl group (e.g., phenyl) optionally substituted by 1 to 3 C₁₋₆ alkoxy (e.g., methoxy),

30 (8) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, thienyl, furyl, isoxazolyl, pyridyl, dihydropyranyl, tetrahydropyranyl, tetrahydropyridyl, piperidyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or

35 (9) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl);

Z² is a group represented by CR^{Z2} wherein R^{Z2} is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom, an iodine atom),
- 5 (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl, isopropyl) optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) hydroxy,
- (4) a C₂₋₆ alkenyl group (e.g., vinyl),
- 10 (5) an optionally halogenated C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),
- (6) a cyano group,
- (7) a C₆₋₁₄ aryl group (e.g., phenyl),
- 15 (8) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl), or
- (9) a heterocyclic group (e.g., azetidiny, pyrazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or a nitrogen atom;

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is

- 20 (1) a hydrogen atom,
- (2) a C₁₋₆ alkyl group (e.g., methyl, isopropyl) optionally substituted by 1 to 3 hydroxy,
- (3) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (4) a C₁₋₆ alkyl-carbonyl group (e.g., acetyl),
- 25 or a nitrogen atom; and

B is a pyridine ring, a pyrimidine ring or a pyrazine ring.

[0148]


[Compound B-5]

compound (1) wherein

- 30 R¹ is a hydrogen atom or a C₁₋₆ alkyl group (preferably methyl);
- R² and R³ are each independently a hydrogen atom or a C₁₋₆ alkyl group (preferably methyl);

the group represented by


[0149]



[0150]

is a group represented by

[0151]



5

[0152]

wherein

R^{21} is

- (1) cyano,
- 10 (2) carbamoyl,
- (3) mono- or di- C_{1-6} alkyl-carbamoyl (e.g., methylcarbamoyl, dimethylcarbamoyl),
- (4) C_{1-6} alkyl (e.g., methyl, ethyl, isopropyl, propyl, butyl, tert-butyl, 1-ethylpropyl) optionally substituted by 1 to 5
- 15 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom),
 - (ii) hydroxy,
 - (iii) amino,
 - (iv) cyano,
 - 20 (v) C_{3-8} cycloalkyl (e.g., cyclopropyl),
 - (vi) C_{1-6} alkoxy (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 substituents selected from
 - (a) a halogen atom (e.g., a fluorine atom), and
 - 25 (b) C_{1-6} alkoxy (e.g., methoxy),
 - (vii) C_{3-8} cycloalkyloxy (e.g., cyclopentyloxy),
 - (viii) mono- or di- C_{1-6} alkyl-amino (e.g., dimethylamino),

(ix) C₁₋₆ alkylsulfanyl (e.g., methylsulfanyl),
 (x) C₁₋₆ alkylsulfinyl (e.g., methylsulfinyl),
 (xi) C₁₋₆ alkylsulfonyl (e.g., methylsulfonyl),
 (xii) carbamoyl,
 5 (xiii) mono- or di-C₁₋₆ alkyl-carbamoyl (e.g., dimethylcarbamoyl),
 (xiv) a heterocyclic group (e.g., oxetanyl, pyrrolidinyl, pyrazolyl, triazolyl, tetrahydropyranyl, dihydroisoindolyl) optionally substituted by 1 to 3
 10 substituents selected from
 (a) hydroxy, and
 (b) oxo,
 (xv) C₁₋₆ alkyl-carboxyloxy (e.g., acetoxy),
 (xvi) C₁₋₆ alkyl(diaryl)silyloxy (e.g., tert-
 15 butyl(diphenyl)silyloxy), and
 (xvii) optionally halogenated heterocycl-yl-oxy (e.g., pyridyloxy) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a bromine atom)),
 (5) C₃₋₈ cycloalkyl (e.g., cyclopropyl),
 20 (6) a heterocyclic group (preferably a 4- to 6-membered heterocyclic group) (e.g., pyridyl, isoxazolyl, pyrazolyl, oxetanyl, tetrahydrofuryl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl),
 (7) C₁₋₆ alkyl-carbonyl (e.g., acetyl),
 25 (8) C₃₋₈ cycloalkyl-carbonyl (e.g., cyclopropylcarbonyl), or
 (9) C₆₋₁₄ aryl (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from
 (i) a halogen atom (e.g., a fluorine atom, a bromine atom), and
 30 (ii) hydroxy,
 R²² is
 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom),
 (2) hydroxy,
 35 (3) optionally halogenated C₁₋₆ alkyl (e.g., methyl,

isopropyl) (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)),

(4) C₂₋₆ alkenyl (e.g., prop-1-en-2-yl),

(5) C₁₋₆ alkoxy (e.g., methoxy, ethoxy, isopropoxy)

5 optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₃₋₈ cycloalkyl (e.g., cyclopropyl),

(6) C₆₋₁₄ aryl (e.g., phenyl),

(7) a heterocyclic group (preferably a 4- to 6-membered

10 heterocyclic group) (e.g., azetidiny, pyrazolyl, pyrrolyl, pyrrolidinyl, triazolyl, dihydropyranyl, tetrahydropyranyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or

(8) C₃₋₈ cycloalkyl (e.g., cyclopropyl), and

15 p is an integer of 1 to 3,

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is

(1) a hydrogen atom,

(2) a halogen atom (e.g., a chlorine atom),

(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally

20 substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) C₆₋₁₄ aryl (e.g., phenyl),

(4) a C₂₋₆ alkenyl group (e.g., vinyl),

(5) a C₁₋₆ alkoxy group (e.g., methoxy),

25 (6) a cyano group,

(7) a C₆₋₁₄ aryl group (e.g., phenyl) optionally substituted by 1 to 3 C₁₋₆ alkoxy (e.g., methoxy),

(8) a heterocyclic group (preferably a 5- or 6-membered heterocyclic group) (e.g., pyrazolyl, thienyl, furyl,

30 isoxazolyl, pyridyl, dihydropyranyl, tetrahydropyranyl, tetrahydropyridyl, piperidyl) optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or

(9) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl);

Z² is a group represented by CR^{Z2} wherein R^{Z2} is

35 (1) a hydrogen atom,

(2) a halogen atom (e.g., a fluorine atom, a chlorine atom, a bromine atom, an iodine atom),
 (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl, isopropyl) optionally substituted by 1 to 3 substituents selected from
 5 (i) a halogen atom (e.g., a fluorine atom), and
 (ii) hydroxy,
 (4) a C₂₋₆ alkenyl group (e.g., vinyl),
 (5) an optionally halogenated C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) (preferably optionally substituted by 1 to
 10 3 halogen atoms (e.g., a fluorine atom)),
 (6) a cyano group,
 (7) a C₆₋₁₄ aryl group (e.g., phenyl),
 (8) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl), or
 (9) a heterocyclic group (e.g., azetidiny, pyrazolyl)
 15 optionally substituted by 1 to 3 C₁₋₆ alkyl (e.g., methyl), or a nitrogen atom;
 Z³ is a group represented by CR^{Z3} wherein R^{Z3} is
 (1) a hydrogen atom,
 (2) a C₁₋₆ alkyl group (e.g., methyl, isopropyl) optionally
 20 substituted by 1 to 3 hydroxy,
 (3) a C₁₋₆ alkoxy group (e.g., methoxy), or
 (4) a C₁₋₆ alkyl-carbonyl group (e.g., acetyl),
 or a nitrogen atom; and
 B is a pyridine ring, a pyrimidine ring or a pyrazine ring.
 25 [0153]
 [Compound C-2]
 compound (1) wherein
 R¹ is a hydrogen atom;
 R² and R³ are both hydrogen atoms;
 30 X is a C₁₋₆ alkylene group (preferably a C₁₋₅ alkylene group, more preferably -CH(CH₃)- or -CH(CH(CH₃)₂)-) optionally substituted by 1 to 3 substituents selected from
 (1) hydroxy, and
 (2) C₁₋₆ alkoxy (e.g., methoxy);
 35 A is a C₆₋₁₀ aryl group (e.g., phenyl) optionally substituted by

1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom), and

(2) C₁₋₆ alkoxy (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom);

5 [preferably the group represented by


[0154]



[0155]

is a group represented by

10 [0156]



[0157]

wherein

R²¹ is a C₁₋₅ alkyl group (e.g., methyl, isopropyl)

15 optionally substituted by hydroxy or C₁₋₆ alkoxy (e.g., methoxy),

R²² is

(1) a halogen atom (e.g., a fluorine atom), or

(2) optionally halogenated C₁₋₆ alkoxy (e.g., methoxy)

20 (preferably optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom)), and

p is an integer of 1 or 2];

Z¹ is CH;

Z² is a group represented by CR²² wherein R²² is

25 (1) a C₁₋₆ alkoxy group (e.g., methoxy), or

(2) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl);

Z³ is CH; and

B is a pyridine ring.

[0158]

[Compound C-3]

compound (1) wherein

5 R^1 is a hydrogen atom;

R^2 and R^3 are both hydrogen atoms;

X is a C_{1-6} alkylene group (preferably a C_{1-5} alkylene group, more preferably $-CH(CH_3)-$ or $-CH(CH(CH_3)_2)-$) optionally substituted by 1 to 3 substituents selected from

10 (1) hydroxy, and

(2) C_{1-6} alkoxy (e.g., methoxy);


A is a C_{6-10} aryl group (e.g., phenyl) optionally substituted by 1 to 3 substituents selected from

(1) a halogen atom (e.g., a fluorine atom), and

15 (2) C_{1-6} alkoxy (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom);

[preferably the group represented by


[0159]



20 [0160]

is a group represented by

[0161]



[0162]

25 wherein

R^{21} is a C_{1-5} alkyl group (e.g., methyl, isopropyl)

optionally substituted by hydroxy or C_{1-6} alkoxy (e.g.,

methoxy),

R²² is

(1) a halogen atom (e.g., a fluorine atom), or

(2) an optionally halogenated C₁₋₆ alkoxy group (e.g.,
5 methoxy) (preferably optionally substituted by 1 to 3
halogen atoms (e.g., a fluorine atom)),

p is an integer of 1 or 2];

Z¹ is CH;

Z² is a group represented by CR²² wherein R²² is

10 (1) a C₁₋₆ alkoxy group (e.g., methoxy),

(2) a C₃₋₈ cycloalkyl group (e.g., cyclopropyl), or

(3) a C₁₋₆ alkyl group (e.g., methyl);

Z³ is CH; and

B is a pyridine ring.

15 [0163]

Specific examples of compound (1) include compounds of
Examples 1 - 280.

[0164]

Examples of the salt of the compound represented by the
20 formula (1) include metal salts, ammonium salts, salts with an
organic base, salt with an inorganic acid, salts with an
organic acid, salts with a basic or acidic amino acid, and the
like.

[0165]

25 Preferable examples of the metal salt include alkaline
metal salts such as sodium salt, potassium salt and the like;
alkaline earth metal salts such as calcium salt, magnesium
salt, barium salt and the like; aluminum salt, and the like.

[0166]

30 Preferable examples of the salt with an organic base
include salts with trimethylamine, triethylamine, pyridine,
picoline, 2,6-lutidine, ethanolamine, diethanolamine,
triethanolamine, cyclohexylamine, dicyclohexylamine, N,N'-
dibenzylethylenediamine and the like.

35 [0167]

Preferable examples of the salt with an inorganic acid include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid and the like.

[0168]

5 Preferable examples of the salt with an organic acid include salts with formic acid, acetic acid, trifluoroacetic acid, phthalic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic
10 acid and the like.

[0169]

Preferable examples of the salt with a basic amino acid include salts with arginine, lysine, ornithine and the like. Preferable examples of the salt with an acidic amino acid
15 include salt with aspartic acid, glutamic acid and the like.

[0170]

Among the above-mentioned salts, pharmaceutically acceptable salts are preferable.

[0171]

20 The prodrug of compound (1) means a compound which is converted to compound (1) with a reaction due to an enzyme, gastric acid and the like under the physiological condition in the living body, that is, a compound which is converted to compound (1) by enzymatic oxidation, reduction, hydrolysis and
25 the like; a compound which is converted to compound (1) by hydrolysis and the like due to gastric acid, and the like.

[0172]

Examples of the prodrug of compound (1) include a compound obtained by subjecting an amino group in compound (1)
30 to an acylation, alkylation or phosphorylation (e.g., a compound obtained by subjecting an amino group in compound (1) to an eicosanoylation, alanylation, pentylaminocarbonylation, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylation, tetrahydrofuranylation, pyrrolidylmethylation,
35 pivaloyloxymethylation or tert-butylation); a compound

obtained by subjecting a hydroxy group in compound (1) to an acylation, alkylation, phosphorylation or boration (e.g., a compound obtained by subjecting a hydroxy group in compound (1) to an acetylation, palmitoylation, propanoylation, pivaloylation, succinylation, fumarylation, alanylation or dimethylaminomethylcarbonylation); a compound obtained by subjecting a carboxyl group in compound (1) to an esterification or amidation (e.g., a compound obtained by subjecting a carboxyl group in compound (1) to a C₁₋₆ alkyl esterification, phenyl esterification, carboxymethyl esterification, dimethylaminomethyl esterification, pivaloyloxymethyl esterification, ethoxycarbonyloxyethyl esterification, phthalidyl esterification, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl esterification, cyclohexyloxycarbonylethyl esterification or methylamidation) and the like. Among them, a compound esterified by C₁₋₆ alkyl (e.g., methyl, ethyl, tert-butyl and the like) are preferably used. These compounds can be produced from compound (1) according to a method known per se.

[0173]

The prodrug of compound (1) may also be one which is converted into compound (1) under a physiological condition, such as those described in IYAKUJIN no KAIHATSU (Development of Pharmaceuticals), Vol.7, Design of Molecules, p.163-198, 1990, Published by HIROKAWA SHOTEN.

[0174]

Each symbol of the compound in the following Reaction Schemes is as defined above, unless otherwise specified. Each compound described in the following Reaction Schemes may be in the form of a salt as long as it does not inhibit the reaction. Examples of the salt include those similar to the salt of compound (1).

[0175]

The compound obtained in each step can be used directly for the next reaction as the reaction mixture or as a crude

product, or can also be isolated according to a conventional method from the reaction mixture, and can also be easily purified according to a separation means (e.g., recrystallization, distillation, chromatography etc.).

5 [0176]

The production methods of the compound of the present invention are described in the following.

Compound (1) (compound (1a) and compound (1b) in the following Reaction Scheme 1) can be produced according to a method known per se, for example, the production method shown
10 in Reaction Scheme 1 to Reaction Scheme 7 or a method analogous thereto.

In each of the following production methods, each starting compound used for the production of compound (1) may
15 be in the form of a salt. Examples of the salt include those similar to the salt of compound (1).

[0177]

Each starting compound to be used for the production of compound (1) can be used directly for the next reaction as the
20 reaction mixture or as a crude product, or can also be isolated according to a conventional method from the reaction mixture, and can also be purified according to a separation means (e.g., extraction, concentration, neutralization, filtration, distillation, recrystallization, chromatography
25 etc.). Examples of the solvent used for the above-mentioned recrystallization include water, alcohols, ethers, hydrocarbons, amides, halogenated hydrocarbons, nitriles, ketones, esters, sulfoxides, organic acids and the like. These solvents may be used alone, or two or more kinds of solvents
30 may be mixed at a suitable ratio, for example, 1:1 - 1:10. In addition, the compounds in the Reaction Schemes may be commercially available, or can be produced according to a method known per se or a method analogous thereto.

[0178]

35 When compound (1) and intermediate for the production of

compound (1) have a convertible functional group (e.g., a carboxyl group, an amino group, a hydroxy group, a carbonyl group, a mercapto group, a C₁₋₆ alkoxy-carbonyl group, a C₆₋₁₄ aryloxy-carbonyl group, a C₇₋₁₆ aralkyloxy-carbonyl group, a
5 sulfo group, a halogen atom, an optionally halogenated C₁₋₆ alkylsulfonyloxy group, a cyano group, an aminocarbonyl group, a boryl group etc.), various compounds can be produced by converting such functional group according to a method known per se or a method analogous thereto.

10 Carboxyl group can be converted, for example, by reactions such as esterification, reduction, amidation, conversion reaction to optionally protected amino group and the like.

Amino group can be converted, for example, by reactions
15 such as amidation, sulfonylation, nitrosation, alkylation, arylation, imidation and the like.

Hydroxy group can be converted, for example, by reactions such as esterification, carbamoylation, sulfonylation, alkylation, fluorination, arylation, oxidation, halogenation
20 and the like.

Carbonyl group can be converted, for example, by reactions such as reduction, oxidation, fluorination, imination (including oximation, hydrazone formation),
(thio)ketalization, alkylidenation, thiocarbonylation and the
25 like.

Mercapto group can be converted, for example, by reactions such as alkylation, oxidation and the like.

C₁₋₆ alkoxy-carbonyl group, C₆₋₁₄ aryloxy-carbonyl group and C₇₋₁₆ aralkyloxy-carbonyl group can be converted, for
30 example, by reactions such as reduction, hydrolysis and the like.

Sulfo group can be converted, for example, by reactions such as sulfonamidation, reduction and the like.

Halogen atom can be converted, for example, by various
35 nucleophilic substitution reactions, various coupling

reactions and the like.

Optionally halogenated C₁₋₆ alkylsulfonyloxy group can be converted, for example, by various nucleophilic substitution reactions, various coupling reactions and the like.

5 Cyano group can be converted, for example, by reactions such as reduction, hydrolysis and the like.

Aminocarbonyl group can be converted, for example, by reactions such as dehydration, reduction and the like.

Boryl group can be converted, for example, by oxidation,
10 various coupling reactions and the like.

In each of the above-mentioned reactions, when the compound is obtained in a free form, it may be converted to a salt according to a conventional method. When it is obtained as a salt, it may be converted to a free form or other salt
15 according to a conventional method.

The conversion of these functional group can be carried out according to a method known per se, for example, the method described in Comprehensive Organic Transformations, Second Edition, Wiley-VCH, Richard C. Larock, or the like.
20 [0179]

In each reaction in the production method of compound (1) and each reaction of the synthesis of the starting materials, when the starting compound has an amino group, a carboxyl group, a hydroxy group, a carbonyl group or a mercapto group
25 as a substituent, a protecting group generally used in peptide chemistry and the like may be introduced into these substituents. By removing the protecting group as necessary after the reaction, the object compound can be obtained.

Examples of the amino-protecting group include a formyl
30 group; and a C₁₋₆ alkyl-carbonyl group (e.g., acetyl, ethylcarbonyl etc.), a phenylcarbonyl group, a C₁₋₆ alkyl-oxy carbonyl group (e.g., methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl (Boc) etc.), an allyloxycarbonyl (Alloc) group, a phenyloxycarbonyl group, a fluorenylmethoxycarbonyl
35 (Fmoc) group, a C₇₋₁₀ aralkyl-carbonyl group (e.g.,

benzylcarbonyl etc.), a C₇₋₁₀ aralkyl-oxycarbonyl group (e.g., benzyloxycarbonyl (Z) etc.), a C₇₋₁₀ aralkyl group (e.g., benzyl etc.), a 2-(trimethylsilyl)ethoxymethyl (SEM) group, a trityl group, a phthaloyl group, an N,N-dimethylaminomethylene group,
5 an allyl group and the like, each of which optionally has substituent(s), and the like. As these substituents, a phenyl group, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.), a C₁₋₆ alkyl-carbonyl group (e.g., methylcarbonyl, ethylcarbonyl, butylcarbonyl etc.), a nitro group and the like
10 are used, and the number of the substituents is about 1 to 3.

Examples of the carboxyl-protecting group include a C₁₋₆ alkyl group (e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl etc.), an allyl group, a benzyl group, a phenyl group, a trityl group, a trialkylsilyl group and the like,
15 each of which optionally has substituent(s). As these substituents, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.), a formyl group, a C₁₋₆ alkyl-carbonyl group (e.g., acetyl, ethylcarbonyl, butylcarbonyl etc.), a nitro group and the like are used, and the number of the
20 substituents is about 1 to 3.

Examples of the hydroxyl-protecting group include a C₁₋₆ alkyl group (e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, tert-butyl etc.), a C₇₋₁₀ aralkyl group (e.g., benzyl etc.), a formyl group, a C₁₋₆ alkyl-carbonyl group (e.g., acetyl,
25 ethylcarbonyl etc.), a benzoyl group, a C₇₋₁₀ aralkyl-carbonyl group (e.g., benzylcarbonyl etc.), a tetrahydropyranyl group, a furanyl group, a silyl group and the like, each of which optionally has substituent(s). As these substituents, a halogen atom (e.g., fluorine, chlorine, bromine, iodine etc.),
30 a C₁₋₆ alkyl group (e.g., methyl, ethyl, n-propyl etc.), a phenyl group, a C₇₋₁₀ aralkyl group (e.g., benzyl etc.), a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, n-propoxy etc.), a nitro group and the like are used, and the number of the substituents is about 1 to 4.

35 Examples of the protected carbonyl group include cyclic

acetal (e.g., 1,3-dioxane), noncyclic acetal (e.g., di-C₁₋₆ alkylacetal) and the like.

Examples of the mercapto-protecting group include a C₁₋₆ alkyl group, a phenyl group, a trityl group, a C₇₋₁₀ aralkyl group (e.g., benzyl), a C₁₋₆ alkyl-carbonyl group, a benzoyl group, a C₇₋₁₀ aralkyl-carbonyl group (e.g., benzylcarbonyl), a C₁₋₆ alkoxy-carbonyl group, a C₆₋₁₄ aryloxy-carbonyl group (e.g., phenyloxycarbonyl), a C₇₋₁₄ aralkyloxy-carbonyl group (e.g., benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl), a 2-tetrahydropyranyl group, a C₁₋₆ alkylamino-carbonyl group (e.g., methylaminocarbonyl, ethylaminocarbonyl) and the like. These protecting groups are optionally substituted by 1 to 3 substituents selected from a halogen atom, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group and a nitro group.

These protecting groups can be introduced and removed by a method known per se, for example, the method described in Greene's Protective Groups in Organic Synthesis, 4th Edition, Wiley-Interscience, Theodora W. Greene, Peter G. M. Wuts or the like. Specifically, a method using acid, base, ultraviolet rays, hydrazine, phenylhydrazine, sodium N-methyldithiocarbamate, tetrabutylammonium fluoride, palladium acetate, trialkylsilyl halide (e.g., trimethylsilyl iodide, trimethylsilyl bromide) and the like, a reduction method, and the like can be employed.

[0180]

When compound (1) is present as a configurational isomer, a diastereomer, a conformer and the like, each can be isolated by a known means. When compound (1) has an optical isomer, racemates can be resolved by a general optical resolution means, whereby an optically active forms ((+) form, (-) form) can be obtained.

[0181]

When compound (1) has an optical isomer, a stereoisomer, a positional isomer, a rotamer or a tautomer, these are also encompassed in compound (1), and can be obtained as a single

product according to synthesis and separation methods known per se.

For example, the method of optical resolution may be a method known per se, such as a fractional recrystallization method, a chiral column method, a diastereomer method etc.

1) Fractional recrystallization method

A method wherein a salt of a racemate with an optically active compound (e.g., (+)-mandelic acid, (-)-mandelic acid, (+)-tartaric acid, (-)-tartaric acid, (+)-1-phenethylamine, (-)-1-phenethylamine, cinchonine, (-)-cinchonidine, brucine etc.) is formed, which is separated by a fractional recrystallization method, and if desired, a neutralization step to give a free optical isomer.

2) Chiral column method

A method wherein a racemate or a salt thereof is applied to a column (a chiral column) for separation of an optical isomer to allow separation. In the case of a liquid chromatography, for example, a mixture of the optical isomers is applied to a chiral column such as ENANTIO-OVM (manufactured by Tosoh Corporation), CHIRAL series (manufactured by Daicel Chemical Industries, Ltd.) and the like, and developed with water, various buffers (e.g., phosphate buffer, etc.) and organic solvents (e.g., ethanol, methanol, isopropanol, acetonitrile, trifluoroacetic acid, diethylamine, etc.) as an eluent, solely or in admixture to separate the optical isomer. In the case of a gas chromatography, for example, a chiral column such as CP-Chirasil-DeX CB (manufactured by GL Sciences Inc.) and the like is used to allow separation.

3) Diastereomer method

A method wherein a racemic mixture is prepared into a diastereomeric mixture by chemical reaction with an optically active reagent, which is made into a single substance by a typical separation means (e.g., a fractional recrystallization method, a chromatography method etc.) and the like, and is subjected to a chemical treatment such as hydrolysis and the

like to separate an optically active reagent moiety, whereby an optical isomer is obtained. For example, when compound (1) contains hydroxy group, or primary or secondary amino group within a molecule, the compound and an optically active
5 organic acid (e.g., MTPA [α -methoxy- α -(trifluoromethyl)phenylacetic acid],
(-)-menthoxyacetic acid etc.) and the like are subjected to condensation reaction to give diastereomers of the ester compound or the amide compound, respectively. When compound (1)
10 has a carboxyl group, the compound and an optically active amine or an optically active alcohol reagent are subjected to condensation reaction to give diastereomers of the amide compound or the ester compound, respectively. The separated diastereomer is converted to an optical isomer of the original
15 compound by acid hydrolysis or base hydrolysis.

[0182]

The solvent, acid and base recited in the production methods of the compound of the present invention are explained in the following.

20 Examples of the "solvent" include "alcohols", "ethers", "hydrocarbons", "amides", "halogenated hydrocarbons", "nitriles", "ketones", "esters", "sulfoxides", "water" and the like.

Examples of the "alcohols" include methanol, ethanol, 1-
25 propanol, 2-propanol, tert-butyl alcohol and the like.

Examples of the "ethers" include diethyl ether, diisopropyl ether, diphenyl ether, tetrahydrofuran, 1,4-dioxane, 1,2-dimethoxyethane, tert-butyl methyl ether and the like.

30 Examples of the "hydrocarbons" include benzene, toluene, cyclohexane, hexane, petroleum ether and the like.

Examples of the "amides" include N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, hexamethylphosphoric triamide and the like.

35 Examples of the "halogenated hydrocarbons" include

dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, chlorobenzene, benzotrifluoride and the like.

Examples of the "nitriles" include acetonitrile, propionitrile and the like.

5 Examples of the "ketones" include acetone, ethyl methyl ketone and the like.

Examples of the "esters" include ethyl acetate, tert-butyl acetate and the like.

10 Examples of the "sulfoxides" include dimethyl sulfoxide and the like.

Examples of the "acid" include "organic acids", "mineral acids", "Lewis acids" and the like.

Examples of the "organic acids" include formic acid, acetic acid, propionic acid, trifluoroacetic acid, citric acid,
15 methanesulfonic acid, p-toluenesulfonic acid and the like.

Examples of the "mineral acids" include hydrochloric acid, sulfuric acid and the like.

Examples of the "Lewis acids" include boron trichloride, boron tribromide and the like.

20 Examples of the "base" include "inorganic bases", "basic salts", "aromatic amines", "tertiary amines", "alkali metal hydrides", "alkali metals", "metal amides", "alkyl metals", "aryl metals", "metal alkoxides" and the like.

Examples of the "inorganic bases" include sodium
25 hydroxide, potassium hydroxide, lithium hydroxide, barium hydroxide and the like.

Examples of the "basic salts" include sodium carbonate, potassium carbonate, cesium carbonate, sodium hydrogen carbonate, sodium acetate, ammonium acetate and the like.

30 Examples of the "aromatic amines" include pyridine, 2,6-lutidine and the like.

Examples of the "tertiary amines" include triethylamine, tripropylamine, tributylamine, diisopropylethylamine, cyclohexyldimethylamine, 4-dimethylaminopyridine, N,N-
35 dimethylaniline, N-methylpiperidine, N-methylpyrrolidine, N-

methyldmorpholine, 1,8-diazabicyclo[5,4,0]undec-7-ene and the like.

Examples of the "alkali metal hydrides" include sodium hydride, potassium hydride and the like.

5 Examples of the "alkali metals" include sodium, lithium, potassium and the like.

Examples of the "metal amides" include sodium amide, lithium diisopropylamide, lithium hexamethyldisilazide and the like.

10 Examples of the "alkyl metals" include butyllithium, sec-butyllithium, tert-butyllithium and the like.

Examples of the "aryl metals" include phenyllithium and the like.

Examples of the "metal alkoxides" include sodium
15 methoxide, sodium ethoxide, sodium tert-butoxide, potassium tert-butoxide and the like.


[0183]

Compound (1) can be produced, for example, according to the method shown in the following Reaction Scheme 1 or a
20 method analogous thereto.

[0184]

Reaction Scheme 1

[0185]



[0186]

wherein R^4 is an optionally substituted C_{1-6} alkyl group, R is an optionally substituted C_{1-6} alkyl group or an amino-
 5 protecting group, L^1 and L^2 are each independently a leaving group, and the other symbols are as defined above.

[0187]

Examples of the amino-protecting group for R include a formyl group; a C_{1-6} alkyl-carbonyl group (e.g., acetyl,
 10 ethylcarbonyl and the like), a phenylcarbonyl group, a C_{1-6} alkyl-oxycarbonyl group (e.g., methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl (Boc) and the like), an allyloxycarbonyl (Alloc) group, a phenyloxycarbonyl group, a fluorenylmethoxycarbonyl (Fmoc) group, a C_{7-10} aralkyl-carbonyl
 15 group (e.g., benzyl carbonyl and the like), a C_{7-10} aralkyl-oxycarbonyl group (e.g., benzyloxycarbonyl (Z) and the like), a C_{7-10} aralkyl group (e.g., benzyl and the like), a 2-

(trimethylsilyl)ethoxymethyl (SEM) group, a trityl group, a phthaloyl group, an N,N-dimethylaminomethylene group and an allyl group, each optionally having substituent(s), and the like. The amino-protecting group is preferably a 2,4-
5 dimethoxybenzyl (DMB) group or a 2-(trimethylsilyl)ethoxymethyl (SEM) group.

Examples of the leaving group for L¹ include a halogen atom, an optionally halogenated C₁₋₆ alkylsulfonyloxy group, a C₆₋₁₀ arylsulfonyloxy group optionally substituted by C₁₋₆ alkyl
10 group(s), and the like. The leaving group is preferably a halogen atom or an optionally halogenated C₁₋₆ alkylsulfonyloxy group.

Examples of the leaving group for L² include a halogen atom, an optionally halogenated C₁₋₆ alkylsulfonyloxy group, an
15 optionally halogenated C₁₋₆ alkoxy group, an optionally substituted aryloxy group, a 1-H-imidazolyl group and the like. The leaving group is preferably a halogen atom, a 4-nitrophenoxy group or a trichloromethoxy group.
[0188]

20 Compounds (2), (3), (6), (10), (11), (13) and (14) may be commercially available, or can also be produced according to a method known per se or a method analogous thereto.

[0189]

<Step 1> Compound (4) can be produced by reacting compound (2)
25 with compound (3).

The amount of compound (3) to be used is generally about 0.5-20 mol, preferably about 1-10 mol, per 1 mol of compound (2).

This reaction is generally carried out in the presence of
30 a base. Examples of the base include tertiary amines, basic salts, metal hydrogen complex compounds, metal alkoxides, metal amides, alkyl metals, aryl metals and the like. The amount of the base to be used is generally about 1-100 mol, preferably about 1-10 mol, per 1 mol of compound (2).

35 This reaction is advantageously carried out in a solvent

inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include alcohols, ethers, amides, sulfoxides, tertiary amines, mixed solvents thereof and the like.

5 While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally 0-300°C, preferably 20-150°C.

10 [0190]

<Step 2> Compound (5) can be produced from compound (4).

The conversion of the nitro group to an amino group can be carried out according to a method known per se, for example, the method described in Reductions in Organic Chemistry,
15 Second Edition, The American Chemical Society, 1996, or a method analogous thereto, for example, a hydrogenation reaction, a reaction using a metal or metal salt, or the like. The next intramolecular amidation reaction mostly proceeds under a condition of the reduction of the nitro group. Where
20 necessary, this amidation reaction can be promoted by raising the reaction temperature or adding a suitable acid or base.

Examples of the acid include organic acids, mineral acids and the like. Examples of the base include tertiary amines, aromatic amines, alkali metal hydrides, basic salts and the
25 like. The amount of the acid or base to be used is generally about 0.01-100 mol, preferably about 0.01-10 mol, per 1 mol of compound (4).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited
30 as long as the reaction proceeds. Preferable examples thereof include alcohols, ethers, esters, organic acids, amides, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably
35 0.5-24 hr.

The reaction temperature is generally 0-200°C, preferably 20-100°C.

[0191]

<Step 3> Compound (7) can be produced by reacting compound (5)
5 with compound (6).

The amount of compound (6) to be used is generally about 0.1-10 mol, preferably about 0.3-5 mol, per 1 mol of compound (5).

This reaction may be carried out in the presence of a
10 base. Examples of the base include tertiary amines, aromatic amines, alkali metal hydrides, basic salts and the like. The amount of the base to be used is generally about 1-100 mol, preferably about 1-10 mol, per 1 mol of compound (5).

This reaction is advantageously carried out in a solvent
15 inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, aromatic amines, amides, tertiary amines, ethers, nitriles, esters, hydrocarbons, mixed solvents thereof and the like.

20 While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally -50-150°C, preferably -20-100°C.

25 [0192]

<Step 4> Compound (1a) can be produced by reacting compound (7) with compound (8).

The amount of compound (8) to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (7).

30 This reaction is generally carried out in the presence of a base. Examples of the base include tertiary amines, aromatic amines, alkali metal hydrides, basic salts, metal alkoxides, inorganic bases and the like. The amount of the base to be used is generally about 1-100 mol, preferably about 1-10 mol,
35 per 1 mol of compound (7).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, aromatic amines, amides, sulfoxides, tertiary amines, ethers, nitriles, esters, hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

10 The reaction temperature is generally -20-150°C, preferably 0-100°C.

[0193]

<Step 5> Compound (1a) can also be produced by reacting compound (5) with compound (9).

15 The amount of compound (9) to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (5).

This reaction may be carried out in the presence of a base. Examples of the base include tertiary amines, aromatic amines, alkali metal hydrides, basic salts, metal amides, alkyl metals, aryl metals and the like. The amount of the base to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (5).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, ethers, nitriles, amides, sulfoxides, tertiary amines, aromatic amines, esters, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally -20-150°C, preferably 0-100°C.

[0194]

35 <Step 6> Compound (12) can be produced by reacting compound

(10) with compound (11).

The amount of compound (11) to be used is generally about 0.5-10 mol, preferably about 1-3 mol, per 1 mol of compound (10).

5 This reaction is generally carried out in the presence of a base. Examples of the base include tertiary amines, aromatic amines, basic salts, alkali metal hydrides, metal alkoxides, metal amides, alkyl metals, aryl metals and the like. The amount of the base to be used is generally about 1-10 mol,
10 preferably about 1-3 mol, per 1 mol of compound (10).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include amides, sulfoxides, ethers, nitriles, hydrocarbons,
15 mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally 0-200°C, preferably
20 20-150°C.

[0195]

<Step 7> Compound (15) can be produced by reacting compound (12) with compound (13).

The amount of compound (13) to be used is generally about
25 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (12).

This reaction is generally carried out in the presence of a base. Examples of the base include tertiary amines, basic salts, alkali metal hydrides, metal alkoxides, metal amides, alkyl metals, aryl metals and the like. The amount of the base
30 to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (13).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
35 include sulfoxides, amides, tertiary amines, aromatic amines,

alcohols, ethers, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

5 The reaction temperature is generally 0-300°C, preferably 50-200°C.

[0196]

<Step 8> Compound (15) can also be produced by reacting compound (5) with compound (14).

10 The amount of compound (14) to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (5).

This reaction is generally carried out in the presence of a base. Examples of the base include metal amides, alkyl metals, aryl metals, alkali metal hydrides, metal alkoxides, 15 inorganic bases, basic salts, tertiary amines, aromatic amines and the like. The amount of the base to be used is generally 1-10 mol, preferably 1-3 mol, per 1 mol of compound (5).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited 20 as long as the reaction proceeds. Preferable examples thereof include amides, sulfoxides, ethers, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 25 0.5-24 hr.

The reaction temperature is generally -100-200°C, preferably -80-150°C.

[0197]

<Step 9> Compound (16) can be produced by reacting compound 30 (15) with compound (6).

This reaction is carried out in the same manner as in Step 3.

[0198]

<Step 10> Compound (1b) can be produced by reacting compound 35 (16) with compound (8).

This reaction is carried out in the same manner as in Step 4.

[0199]

<Step 11> Compound (1b) can also be produced by reacting
5 compound (15) with compound (9).

This reaction is carried out in the same manner as in Step 5.

[0200]

<Step 12> Compound (1a) can also be produced by subjecting
10 compound (1b) wherein R is an amino-protecting group to a deprotection reaction.

The removal of the protecting group of compound (1b) can be carried out according to a method known per se, for example, the method described in Greene's Protective Groups in Organic
15 Synthesis, 4th Edition, Wiley-Interscience, Theodora W. Greene, Peter G. M. Wuts or the like, or a method analogous thereto. For the removal of the protecting group, for example, a method using an acid, base or the like, a hydrogenation reaction or the like are employed.


20 [0201]

Compound (5) in Reaction Scheme 1 (compound (5a) in the following Reaction Scheme 2 or compound (5) in Reaction Scheme 3) can be produced, for example, according to the method described in Reaction Scheme 2 or Reaction Scheme 3 or a
25 method analogous thereto.

[0202]

Reaction Scheme 2

[0203]



[0204]

wherein R⁵ is an optionally substituted C₁₋₆ alkyl group, R⁶ and R⁷ are each a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, and the other symbols are as defined above.

[0205]

Compounds (17), (18) and (23) may be commercially available, or can also be produced according to a method known per se or a method analogous thereto.

[0206]

<Step 13> Compound (19) can be produced by reacting compound (17) with compound (18).

The amount of compound (18) to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (17).

This reaction is generally carried out in the presence of a base. Examples of the base include tertiary amines, aromatic amines, alkali metal hydrides, basic salts, metal alkoxides, inorganic bases and the like. The amount of the base to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (17).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include aromatic amines, halogenated hydrocarbons, amides, sulfoxides, tertiary amines, ethers, nitriles, esters,

hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

5 The reaction temperature is generally -20-150°C, preferably 0-100°C.

[0207]

<Step 14> Compound (20) can be produced by reacting compound (19) with a nitrating reagent.

10 Examples of the nitrating reagent include metal nitrates such as sodium nitrate, potassium nitrate and the like, acetyl nitrate, dinitrogen pentoxide, nitronium salts, fuming nitric acid, nitric acid, mixed acids (a mixture of sulfuric acid and fuming nitric acid or nitric acid), and mixtures thereof.

15 The amount of the nitrating reagent to be used is generally about 0.8-100 mol, preferably about 1.0-20 mol, per 1 mol of compound (19). When fuming nitric acid, nitric acid, a mixed acid or the like is used as a nitrating reagent, it may be used in an excess amount as a reaction solvent.

20 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include alcohols, ethers, hydrocarbons, amides, halogenated hydrocarbons, nitriles, sulfoxides, organic acids, mineral
25 acids, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 1-120 hr, preferably 1-96 hr.

30 The reaction temperature is -20-150°C, preferably 0-80°C.
[0208]

<Step 15> Compound (21) can be produced by subjecting compound (20) to a deprotection reaction.

The removal of the protecting group of compound (20) can be carried out according to a method known per se, for example,
35 the method described in Greene's Protective Groups in Organic

Synthesis, 4th Edition, Wiley-Interscience, Theodora W. Greene, Peter G. M. Wuts or the like, or a method analogous thereto, for example, a method using an acid or base, hydrogenation reaction or the like.

5 [0209]

<Step 16> Compound (22) can be produced from compound (21).

The conversion of the nitro group to an amino group can be carried out according to a method known per se, for example, the method described in Reductions in Organic Chemistry,
10 Second Edition, The American Chemical Society, 1996 or a method analogous thereto, for example, a hydrogenation reaction, a reaction using a metal or metal salt, or the like.

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited
15 as long as the reaction proceeds. Preferable examples thereof include alcohols, ethers, amides, esters, organic acids, mineral acids, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably
20 0.5-24 hr.

The reaction temperature is generally 0-200°C, preferably 20-100°C.

[0210]

<Step 17> Compound (24) can be produced by reacting compound
25 (22) with compound (23).

The amount of compound (23) to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (22).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited
30 as long as the reaction proceeds. Preferable examples thereof include water, alcohols, ethers, esters, amides, nitriles, aromatic amines, organic acids, mineral acids, halogenated hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent
35 and solvent to be used, it is generally 0.1-48 hr, preferably

0.5-24 hr.

The reaction temperature is generally -20-150°C, preferably 0-100°C.

[0211]

5 <Step 18> Compound (5a) can be produced by subjecting compound (24) to a reduction reaction.

Examples of the reducing agent include alkali metal hydrides, metal hydrogen complex compounds, borane complexes and the like. The amount of the reducing agent to be used is
10 generally about 1-100 mol, preferably about 1-10 mol, per 1 mol of compound (24).

Alternatively, this reduction reaction can also be carried out by a hydrogenation reaction. In this case, for example, a catalyst such as palladium on carbon, palladium
15 hydroxide on carbon, palladium black, platinum on carbon, platinum dioxide, Raney nickel, Raney cobalt and the like are can be used. The amount of the catalyst to be used is generally about 5-1000 wt%, preferably about 10-300 wt%, per 1 mol of compound (24).

20 The hydrogenation reaction can also be carried out by using various hydrogen sources instead of gaseous hydrogen. Examples of the hydrogen source include formic acid, ammonium formate, triethylammonium formate, sodium phosphinate, hydrazine and the like. The amount of the hydrogen source to
25 be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (24).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
30 include halogenated hydrocarbons, alcohols, ethers, esters, aromatic hydrocarbons, saturated hydrocarbons, amides, organic acids, water, mixed solvents thereof and the like.


While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably
35 0.5-50 hr.

The reaction temperature is generally -20-100°C,
preferably 0-80°C.

[0212]

Reaction Scheme 3

5 [0213]



[0214]

wherein R⁸ is an optionally substituted C₁₋₆ alkyl group or an
optionally substituted benzyl group, and the other symbols are
10 as defined above.

[0215]

Compounds (25) and (28) may be commercially available, or
can also be produced according to a method known per se or a
method analogous thereto.

15 [0216]

<Step 19> Compound (26) can be produced by reacting compound
(21) with compound (25).

The amount of compound (25) to be used is generally about
1-10 mol, preferably about 1-5 mol, per 1 mol of compound (21).

20 This reaction may be carried out in the presence of a
base to promote the reaction. Examples of the base include
tertiary amines, aromatic amines, inorganic bases, basic salts
and the like. The amount of the base to be used is generally
about 1-10 mol, preferably about 1-5 mol, per 1 mol of
25 compound (25).

This reaction is advantageously carried out in a solvent

inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include aromatic amines, amides, alcohols, ethers, halogenated hydrocarbons, esters, nitriles, sulfoxide, hydrocarbons, mixed
5 solvents thereof and the like.

The reaction time is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally -30-120°C, preferably -10-100°C.

10 [0217]

<Step 20> Compound (27) can be produced from compound (26).

This reaction is carried out in the same manner as in Step 16.

[0218]

15 <Step 21> Compound (29) can be produced by reacting compound (27) with compound (28).

This reaction is carried out in the same manner as in Step 13.

[0219]

20 <Step 22> Compound (30) can be produced by reacting compound (29) in the presence of a base.

Examples of the base include basic salts, alkali metal hydrides, metal alkoxides, metal amides, alkyl metals, aryl metals, aromatic amines, tertiary amines and the like. The
25 amount of the base to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (29).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
30 include ethers, aromatic hydrocarbons, saturated hydrocarbons, amides, halogenated hydrocarbons, nitriles, ketones, esters, sulfoxides, aromatic amines, tertiary amines, mixed solvents thereof and the like.

The reaction time is generally 0.1-48 hr, preferably 0.3-
35 24 hr.

The reaction temperature is generally -70-250°C,
preferably -20-100°C.

[0220]

<Step 23> Compound (5) can be produced from compound (30).

5 This reaction is carried out in the same manner as in
Step 15.


[0221]

Among compound (8) and compound (9) in Reaction Scheme 1,
compounds (8a) and (9a) can be produced, for example,
10 according to the method shown in Reaction Scheme 4 or a method
analogous thereto.

[0222]

Reaction Scheme 4

[0223]



[0224]

wherein R^9 is a hydrogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group or an optionally substituted C_{2-6} alkynyl group, R^{10} is an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group or an optionally esterified carboxyl group, R^{11} is an optionally substituted C_{1-6} alkyl group, R^{12} is a hydroxy group, an optionally substituted C_{1-3} alkoxy group or an optionally substituted benzyl group, R' is a substituent defined by R^9 and R^{10} , L^3 is a hydrogen atom or a leaving group, L^4 and L^5 are

each a leaving group, R^9M^1 or ring AM^1 is an organic metal reagent for introducing a substituent defined by R^9 or ring A, and the other symbols are as defined above.

[0225]

5 Examples of the leaving group for L^3 include an N,O-dimethylhydroxylamino group, an optionally substituted C_{1-6} alkoxy group, a halogen atom and the like.

 Examples of the leaving group for L^4 include a halogen atom, an optionally halogenated C_{1-6} alkylsulfonyloxy group and
10 the like.

 Examples of the leaving group for L^5 include a halogen atom, an optionally halogenated C_{1-6} alkylsulfonyloxy group, a C_{6-10} arylsulfonyloxy group optionally substituted by C_{1-6} , and the like.

15 [0226]

 Compounds (31), (32), (35), (36), (37), (38), (39), (42) and (45) may be commercially available, or can be produced according to a method known per se or a method analogous thereto.

20 [0227]

<Step 24> Compound (33) wherein R' is R^{10} can be produced by reacting compound (31) wherein L^3 is a leaving group with compound (32).

 The amount of compound (32) to be used is generally about
25 0.5-10 mol, preferably about 1-5 mol, per 1 mol of compound (31).

 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
30 include ethers, hydrocarbons, mixed solvents thereof and the like.

 While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

35 The reaction temperature is generally -78-150°C,

preferably -30-80°C.

[0228]

<Step 25> Compound (34) wherein R' is R¹⁰ can be produced by reacting compound (31) wherein L³ is a hydrogen atom with
5 compound (32).

The amount of compound (32) to be used is generally about 0.5-10 mol, preferably about 1-5 mol, per 1 mol of compound (31).

This reaction is advantageously carried out in a solvent
10 inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include ethers, hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent
15 and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally -78-150°C,
preferably -30-80°C.

[0229]

<Step 26> Compound (34) can be produced by subjecting compound
20 (33) to a reduction reaction.

Examples of the reducing agent include alkali metal hydrides, metal hydrogen complex compounds, borane complexes and the like. The amount of the reducing agent to be used is
25 generally about 0.25-50 mol, preferably about 0.5-5 mol, per 1 mol of compound (33).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
30 include alcohols, ethers, aromatic hydrocarbons, saturated hydrocarbons, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably 0.5-50 hr.

35 The reaction temperature is generally -78-100°C,

preferably -30-80°C.

[0230]

<Step 27> Compound (33) can be produced by subjecting compound (34) to an oxidation reaction.

5 This reaction is carried out using an oxidizing agent according to a conventional method.

 Examples of the oxidizing agent include metal salts or metal oxides such as chromium(VI) oxide, pyridinium chlorochromate, manganese dioxide and the like, and organic
10 oxidizing agents such as o-iodoxybenzoic acid (IBX), 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one (Dess-Martin periodinane) and the like. The amount of the oxidizing agent to be used is generally about 1-100 mol, preferably about 1-50 mol, per 1 mol of compound (34).

15 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, aromatic hydrocarbons, saturated hydrocarbons, nitriles, esters, ethers, sulfoxides,
20 water, mixed solvents thereof and the like.

 While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.1-24 hr.

 The reaction temperature is generally -78-150°C,
25 preferably -78-100°C.

[0231]

<Step 28> Compound (33) wherein R' is R⁹ can be produced by reacting compound (35) wherein L³ is a leaving group with compound (36).

30 This reaction is carried out in the same manner as in Step 24.

[0232]

<Step 29> compound (34) wherein R' is R⁹ can be produced by reacting compound (35) wherein L³ is a hydrogen atom with
35 compound (36).

This reaction is carried out in the same manner as in Step 24.

[0233]

<Step 30> Compound (39) can be produced by reacting compound
5 (37) with zinc cyanide (38) in the presence of a palladium catalyst.

The amount of the zinc cyanide to be used is generally about 0.5-10 mol, preferably about 1-5 mol, per 1 mol of compound (37).

10 Examples of the palladium catalyst include palladium(II) acetate, tris(dibenzylideneacetone)dipalladium(0), tetrakis(triphenylphosphine)palladium(0), di-tert-butylphosphine palladium(0),
bis(triphenylphosphine)palladium(II) chloride, [1,1'-
15 bis(diphenylphosphino)ferrocene]palladium(II) chloride and the like. The amount of the palladium catalyst to be used is generally about 0.005-1 mol, preferably about 0.01-1 mol, per 1 mol of compound (37).

This reaction is generally under an inert gas (e.g.,
20 argon gas or nitrogen gas) atmosphere or stream in the presence of a phosphine ligand. Examples of the phosphine ligand include tert-butylphosphine, triphenylphosphine, 1,1'-bis(diphenylphosphino)ferrocene and the like. The amount of the phosphine ligand to be used is generally about 1-10 mol,
25 preferably about 1-5 mol, per 1 mol of the palladium catalyst. Where necessary, a zinc metal may be added to the reaction system. The amount of the zinc metal to be used is generally about 0.005-10 mol, preferably about 0.01-5 mol, per 1 mol of compound (37).

30 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include amides, ethers, hydrocarbons, mixed solvents thereof and the like.

35 While the reaction time varies depending on the reagent

and solvent to be used, it is generally 0.5-100 hr, preferably 1-48 hr.

The reaction temperature is generally 0-200°C, preferably 20-150°C.

5 [0234]

<Step 31> Compound (33) wherein R' is R⁹ can be produced by reacting (39) with compound (36).

This reaction is carried out in the same manner as in Step 24.

10 [0235]

<Step 32> Compound (41) can be produced by reacting compound (37) with an organotin reagent (40) in the presence of a palladium catalyst.

Examples of the palladium catalyst include palladium(II) acetate, tris(dibenzylideneacetone)dipalladium(0), tetrakis(triphenylphosphine)palladium(0), di-tert-butylphosphine palladium(0), bis(triphenylphosphine)palladium(II) chloride, [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) chloride and the like. The amount of the palladium catalyst to be used is generally about 0.005-1 mol, preferably about 0.01-1 mol, per 1 mol of compound (37).

The amount of the organotin reagent (40) to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (37).

This reaction is generally under an inert gas (e.g., argon gas or nitrogen gas) atmosphere or stream in the presence of a phosphine ligand. Examples of the phosphine ligand include tert-butylphosphine, triphenylphosphine, 1,1'-bis(diphenylphosphino)ferrocene and the like. The amount of the phosphine ligand to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of the palladium catalyst.

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof

include hydrocarbons, amides, ethers, sulfoxides, nitriles, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.5-100 hr, preferably
5 1-48 hr.

The reaction temperature is generally 0-200°C, preferably 20-150°C.

[0236]

<Step 33> Compound (33) wherein R' is a methyl group can be
10 produced by subjecting compound (41) to an acid hydrolysis reaction.

Examples of the acid include mineral acids, organic acids and the like. The amount of the acid to be used is generally about 1-100 mol, preferably about 1-10 mol, per 1 mol of
15 compound (41).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include hydrocarbons, alcohols, ethers, nitriles, water, mixed
20 solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally 0-150°C, preferably
25 20-100°C.

[0237]

<Step 34> Compound (43a) or compound (43b) can be produced by reacting compound (33) with compound (42).

(i) Compound (43a) can be produced by subjecting compound (33)
30 to a dehydration and condensation reaction with compound (42).

The amount of compound (42) to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (33).

When compound (42) is in the form of a salt, this reaction is generally carried out in the presence of a base.
35 Examples of the base include tertiary amines, aromatic amines,

basic salts, inorganic bases, alkali metal hydrides, metal alkoxides and the like. The amount of the base to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (42).

5 This reaction can also be promoted by adding a dehydrating agent such as molecular sieves and the like, p-toluenesulfonic acid, zinc chloride, phosphoryl chloride, boron trifluoride, titanium tetrachloride, acetic acid, trifluoroacetic acid or the like to the reaction system, by
10 removing water generated in the reaction system using Dean-Stark and the like, or by a combination thereof.

 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
15 include alcohols, ethers, hydrocarbons, esters, mixed solvents thereof and the like.

 While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

20 The reaction temperature is generally 0-150°C, preferably 20-100°C.

(ii) Compound (43b) can be produced by subjecting compound (33) to a reductive amination reaction with compound (42) (e.g., described in 4th ed., Jikken Kagaku Koza, vol. 20,
25 pages 282-284 and 366-368 (Japan Chemical Society); J. Am. Chem. Soc., vol. 93, pages 2897-2904, 1971; Synthesis, page 135, 1975, or the like).

 In this reaction, compound (43b) is produced by subjecting the imine compound, which is produced by the
30 dehydration reaction of compound (33) with compound (42), to a reduction reaction.

 The amount of compound (42) to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of compound (33).

 The dehydration reaction can also be promoted by adding a
35 dehydrating agent such as molecular sieves and the like, p-

toluenesulfonic acid, zinc chloride, phosphoryl chloride, boron trifluoride, titanium tetrachloride, acetic acid, trifluoroacetic acid or the like to the reaction system, by removing water generated in the reaction system using Dean-
5 Stark and the like, or by a combination thereof.

The reduction reaction is generally carried out using a reducing agent according to a conventional method. Examples of the reducing agent include metal hydrides such as aluminium hydride, diisobutylaluminium hydride, tributyltin hydride and
10 the like, metal hydrogen complex compounds such as sodium cyanoborohydride, sodium triacetoxymorohydride, sodium borohydride, lithium aluminium hydride and the like, borane complexes such as borane tetrahydrofuran complex, borane dimethyl sulfide complex, picoline-borane complex and the like,
15 alkyl boranes such as hexylborane, disiamyl borane and the like, and the like.

The amount of the reducing agent to be used is generally about 0.25-10 mol, preferably about 0.5-5 mol, per 1 mol of compound (33).

20 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, alcohols, ethers, nitriles, esters, hydrocarbons, amides, organic acids, mixed solvents
25 thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably 0.5-24 hr.

The reaction temperature is generally -20-200°C,
30 preferably 0-100°C.

[0238]

<Step 35> Compound (8a) can be produced from compound (43a) or compound (43b).

(i) This reaction is carried out, for example, using a
35 catalyst such as palladium-carbon, palladium hydroxide,

palladium black, platinum dioxide, Raney nickel, Raney cobalt and the like. The amount of the catalyst to be used is generally about 1-1000 wt%, preferably about 5-300 wt%, relative to compound (43a) or compound (43b).

5 This reaction is also carried out using various hydrogen sources instead of gaseous hydrogen. Examples of the hydrogen source include formic acid, ammonium formate, triethylammonium formate, sodium phosphinate, hydrazine and the like. The amount of the hydrogen source to be used is generally about 1-
10 10 mol, preferably about 1-5 mol, per 1 mol of compound (43a) or compound (43b).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
15 include halogenated hydrocarbons, alcohols, ethers, esters, organic acids, amides, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably 0.5-50 hr.

20 The reaction temperature is generally -20-150°C, preferably 0-100°C.

(ii) When R^{13} is a hydroxy group or an optionally substituted C_{1-3} alkoxy group, compound (8a) can be produced, for example, by subjecting compound (43a) or compound (43b) to a reduction
25 reaction.

Examples of the reducing agent include borane complexes, metal hydrogen complex compounds and the like.

The amount of the reducing agent to be used is generally about 0.25-100 mol, preferably about 0.5-10 mol, per 1 mol of
30 compound (43a) or compound (43b).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include ethers, alcohols, water, mixed solvents thereof and
35 the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally -50-150°C,
5 preferably -20-100°C.

[0239]

<Step 36> Compound (44) can be produced from compound (34).

(i) Compound (44) can be produced, for example, by reacting compound (34) with diphenylphosphoryl azide (DPPA) under the
10 Mitsunobu reaction conditions (e.g., described in Synthesis, pages 1-27, 1981, Tetrahedron Lett., vol.36, pages 6373-6374, 1995, Tetrahedron Lett., vol.38, pages 5831-5834, 1997, or the like).

The amount of the DPPA to be used is generally about 1-10
15 mol, preferably about 1-5 mol, per 1 mol of compound (34).

The amount of the azodicarboxylate and phosphine to be used for this reaction is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (34), respectively.

This reaction is advantageously carried out in a solvent
20 inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include ethers, aromatic hydrocarbons, saturated hydrocarbons, amides, halogenated hydrocarbons, nitriles, ketones, esters, sulfoxides, mixed solvent thereof and the like.

25 The reaction time is generally 0.1-100 hr, preferably 0.5-48 hr.

The reaction temperature is generally -20-200°C, preferably 0-100°C.

(ii) Compound (44) can also be produced, for example, by
30 reacting compound (34) with DPPA in the presence of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).

The amount of the DPPA to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (34).

The amount of the DBU to be used is generally about 1-10
35 mol, preferably about 1-5 mol, per 1 mol of compound (34).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include aromatic hydrocarbons, saturated hydrocarbons and the like, ethers, amides, esters, nitriles, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

10 The reaction temperature is generally 0-150°C, preferably 20-100°C.

[0240]

<Step 37> Compound (46) can be produced from compound (34).

This reaction is generally carried out using
15 methanesulfonyl chloride, p-toluenesulfonyl chloride or the like, in the presence of a base, according to a conventional method.

Examples of the base include tertiary amines, aromatic amines, basic salts and the like. The amount of the base to be
20 used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (34).

The amount of the methanesulfonyl chloride, p-toluenesulfonyl chloride or the like to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of
25 compound (34).

This reaction can also be carried out by employing the conversion reaction known per se of alcohol to halide using triphenylphosphine/carbon tetrachloride, thionyl chloride or the like.

30 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include ethers, aromatic amines, tertiary amines, halogenated hydrocarbons, hydrocarbons, mixed solvents thereof and the like.
35

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.1-24 hr.

The reaction temperature is generally -30-100°C,
5 preferably -10-at 60°C.

[0241]

<Step 38> Compound (46) wherein L⁵ is a halogen atom can be produced by reacting compound (45) with a halogenating agent.

Examples of the halogenating agent include N-
10 bromosuccinimide, N-chlorosuccinimide, bromine, chlorine, thionyl chloride and the like. The amount of the halogenating agent to be used is generally about 1-50 mol, preferably about 1-10 mol, per 1 mol of compound (45).

When N-bromosuccinimide or N-chlorosuccinimide is used as
15 a halogenating agent, this reaction is generally carried out by heating together with a radical initiator such as azobisisobutyronitrile, benzoyl peroxide or the like, by light irradiation, or by a combination thereof. The amount of the radical initiator to be used is generally about 0.001-5 mol,
20 preferably about 0.01-1 mol, per 1 mol of compound (45).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, aromatic hydrocarbons,
25 saturated hydrocarbons, esters, nitriles, ethers, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably 0.5-50 hr.

30 The reaction temperature is generally 0-150°C, preferably 20-100°C.

[0242]

<Step 39> Compound (44) can be produced by reacting compound (46) with an azidating agent.

35 Examples of the azidating agent include sodium azide,

cesium azide, tetrabutylammonium azide and the like. The amount of the azidating agent to be used is generally about 1-50 mol, preferably about 1-10 mol, per 1 mol of compound (46).

This reaction can be promoted by adding a crown ether.
5 Examples of the crown ether include 18-crown-6 and the like. The amount of the crown ether to be used is generally about 0.01-20 mol, preferably about 0.1-5 mol, per 1 mol of compound (46).

This reaction is advantageously carried out in a solvent
10 inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include amides, nitriles, ketones, sulfoxides, esters, ethers, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent
15 and solvent to be used, it is generally 0.1-100 hr, preferably 0.5-50 hr.

The reaction temperature is generally 0-150°C, preferably 20-100°C.

[0243]

20 <Step 40> Compound (8a) can be produced from compound (44).

The conversion of the azide group to an amino group can be carried out according to a method known per se, for example, the method described in Comprehensive Organic Transformations, Second Edition, Wiley-VCH, Richard C. Larock or the like, or a
25 method analogous thereto, for example, a hydrogenation reaction, Staudinger reaction using a triphenylphosphine, or the like.

[0244]

<Step 41> Compound (9a) can be produced by reacting compound
30 (8a) with phosgene or an equivalent thereof.

Examples of the equivalent of phosgene include bis(trichloromethyl)carbonate, trichloromethyl chloroformate, 1,1'-carbonyldiimidazole and the like. The amount of the phosgene or an equivalent thereof to be used is generally
35 about 0.1-100 mol, preferably about 0.3-10 mol, per 1 mol of

compound (8a).

This reaction is generally carried out in the presence of a base. Examples of the base include tertiary amines, aromatic amines, basic salts and the like. The amount of the base to be used is generally 1-100 mol, preferably 1-10 mol, per 1 mol of compound (8a).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, ethers, hydrocarbons, nitriles, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.5-24 hr.

The reaction temperature is generally -50-200°C, preferably -20-150°C.


[0245]

Among compound (8) in Reaction Scheme 1, compounds (8b), (8c) and (8d) in Reaction Scheme 5 can be produced, for example, according to the method shown in Reaction Scheme 5 or a method analogous thereto.

[0246]

Reaction Scheme 5

[0247]



[0248]

wherein R¹³ is an amino-protecting group, R¹⁴ is an optionally substituted C₁₋₆ alkyl group, R¹⁵ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, and the other symbols
5 are as defined above.

[0249]

Compound (33a) may be commercially available, or can be produced according to a method known per se or a method analogous thereto.

10 [0250]

<Step 42> Compound (8b) can be produced by subjecting compound (33a) to Strecker reaction.

This reaction is generally carried out by condensing compound (33a) with ammonia or an equivalent thereof and
15 hydrogen cyanide or an equivalent thereof to give the corresponding α -aminonitrile (8b).

Examples of the equivalent of ammonia include ammonium chloride, ammonium carbonate, benzyl amine and the like. The amount of the reagent to be used is generally about 1-50 mol, preferably about 1-10 mol, per 1 mol of compound (33a).
20

Examples of the equivalent of hydrogen cyanide include sodium cyanide, potassium cyanide, trimethylsilyl cyanide and the like. The amount of the reagent to be used is generally about 1-50 mol, preferably about 1-10 mol, per 1 mol of
25 compound (33a).

This reaction may be carried out by adding a Lewis acid such as titanium(IV) tetraisopropoxide and the like. The amount of the Lewis acid to be used is generally about 0.05-50 mol, preferably about 0.1-10 mol, per 1 mol of compound (33a).

30 This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include alcohols, halogenated hydrocarbons, ethers, hydrocarbons, nitriles, water, mixed solvents thereof and the
35 like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably 0.5-48 hr.

The reaction temperature is generally -78-200°C,
5 preferably -78-100°C.

[0251]

<Step 43> Compound (43) can be produced by protecting the amino group of compound (8b).

The introduction of protecting group in this step can be
10 carried out according to a method known per se, for example, the method described in Greene's Protective Groups in Organic Synthesis, 4th Edition, Wiley-Interscience, Theodora W. Greene, Peter G. M. Wuts or the like.

[0252]

15 <Step 44> Compound (47) can be produced from compound (43).

This reaction can be carried out according to a method known per se, for example, the method described in Synthesis, vol.12, pages 949-950, 1989, or a method analogous thereto.

This reaction is advantageously carried out in a solvent
20 inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include sulfoxides, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably
25 0.5-24 hr.

The reaction temperature is generally -30-100°C, preferably 0-50°C.

[0253]

<Step 45> Compound (48) can be produced by subjecting compound
30 (47) to a hydrolysis reaction.

This reaction is carried out using an acid or base according to a conventional method.

Examples of the acid include mineral acids, organic acids and the like. Examples of the base include inorganic bases,
35 basic salts and the like. The amount of the acid or base to be

used is generally about 0.5-100 mol, preferably about 1-20 mol, per 1 mol of compound (47).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited
5 as long as the reaction proceeds. Preferable examples thereof include alcohols, ethers, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably
10 0.1-48 hr.

The reaction temperature is generally -10-200°C, preferably 0-150°C.

[0254]

<Step 46> Compound (49) can be produced from compound (33a).

15 This reaction can be carried out according to a method known per se, for example, the method described in ORGANIC PREPARATIONS AND PROCEDURES INT., vol.36, pages 391-443, 2004, or a method analogous thereto.

[0255]

20 <Step 47> Compound (8c) can be produced by subjecting compound (49) to a hydrolysis reaction.

This reaction is carried out in the same manner as in Step 45.

[0256]

25 <Step 48> Compound (48) can be produced by protecting the amino group of compound (8c).

This reaction is carried out in the same manner as in Step 43.

[0257]

30 Compound (8d) can be produced, for example, from compound (48) successively according to an esterification reaction known per se, a reduction reaction known per se, and a deprotection reaction known per se of the protecting group.

[0258]

35 Among compound (33) in Reaction Scheme 4, compound (33c)


in Reaction Scheme 6 can be produced, for example, according to the method shown in Reaction Scheme 6 or a method analogous thereto.

[0259]

5 Reaction Scheme 6

[0260]

[0260]



[0261]

10 wherein R¹⁶ is an optionally substituted C₁₋₆ alkyl group, and the other symbols are as defined above.

[0262]

Compound (33a) and compound (33b) may be commercially available, or can be produced according to a method known per se or a method analogous thereto.

[0263]

<Step 49> Compound (52) can be produced by subjecting compound (33b) to a bromination.

Examples of the brominating agent include bromine, phenyltrimethylammonium tribromide, N-bromosuccinimide and the like. The amount of the brominating agent to be used is generally about 0.5-2 mol, preferably about 0.8-1.5 mol, per 1 mol of compound (33b).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include organic acids, alcohols, ethers, amides, halogenated hydrocarbons, hydrocarbons, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-100 hr, preferably 0.1-48 hr.

The reaction temperature is generally -30-200°C,
5 preferably 0-100°C.

[0264]

<Step 50> Compound (33c) can be produced from compound (52).

(i) This reaction is carried out, for example, by reacting compound (52) with a desirable alcohol ($R^{16}OH$) in the presence
10 of silver(I) carbonate or silver(I) oxide and boron trifluoride diethyl ether complex.

The amount of the silver(I) carbonate or silver(I) oxide to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (52).

15 The amount of the boron trifluoride diethyl ether complex to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (52).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited
20 as long as the reaction proceeds. Preferable examples thereof include alcohols corresponding to $R^{16}OH$, ethers, halogenated hydrocarbons, hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent
25 and solvent to be used, it is generally 0.1-100 hr, preferably 0.1-48 hr.

The reaction temperature is generally -30-150°C,
preferably 0-80°C.

(ii) This reaction can also be carried out by reacting
30 compound (52) with the desired alcohol ($R^{16}OH$) in the presence of a base.

Examples of the base include tertiary amines, basic salts, metal hydrogen complex compounds, metal alkoxides, metal amides, alkyl metals, aryl metals and the like. The amount of
35 the base to be used is generally about 1-10 mol, preferably

about 1-5 mol, per 1 mol of compound (52).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof
5 include alcohols corresponding to $R^{16}OH$, ethers, amides, sulfoxides, halogenated hydrocarbons, hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably
10 0.5-24 hr.

The reaction temperature is generally 0-150°C, preferably 20-100°C.

[0265]

<Step 51> Compound (53) can be produced by subjecting compound
15 (33a) to Wittig reaction.

This reaction is carried out, for example, by reacting compound (33a) with phosphorus ylide prepared by methyl(triphenyl)phosphonium salt and a base.

Examples of the base include metal alkoxides, alkyl
20 metals, alkali metal hydrides, metal amides and the like. The amount of the base to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (33a).

The amount of the methyl(triphenyl)phosphonium salt to be used is generally about 1-10 mol, preferably about 1-5 mol,
25 per 1 mol of compound (33a).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include ethers, halogenated hydrocarbons, hydrocarbons,
30 sulfoxides, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.1-24 hr.

The reaction temperature is generally -78-100°C,
35 preferably 0-100°C.

[0266]

<Step 52> Compound (54) can be produced by reacting compound (53) with an oxidizing agent.

Examples of the oxidizing agent include 3-chloroperbenzoic acid, peracetic acid and the like.

The amount of the oxidizing agent to be used is generally about 1-10 mol, preferably about 1-5 mol, per 1 mol of compound (53).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, organic acids, esters, ethers, hydrocarbons, nitriles, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably 0.1-24 hr.

The reaction temperature is generally -78-150°C, preferably -20-80°C.

[0267]

<Step 53> Compound (55) can be produced by reacting compound (54) with the alcohol ($R^{16}OH$) in the presence of a base.

Examples of the base include metal alkoxides prepared from $R^{16}OH$, basic salts, metal hydrogen complex compounds, tertiary amines and the like. The amount of the base to be used is generally about 1-100 mol, preferably about 1-10 mol, per 1 mol of compound (54).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include alcohols corresponding to $R^{16}OH$, amides, halogenated hydrocarbons, ethers, hydrocarbons, nitriles, water, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.1-48 hr, preferably

0.1-24 hr.

The reaction temperature is generally -30-150°C,
preferably 0-100°C.

[0268]

5 <Step 54> Compound (33c) can be produced by subjecting
compound (55) to an oxidation reaction.

This reaction is carried out in the same manner as in
Step 27.

[0269]

10 Among compound (33) in Reaction Scheme 4, compound (33e)
in Reaction Scheme 7 can be produced, for example, according
to the method shown in Reaction Scheme 7 or a method analogous
thereto.

[0270]

15 Reaction Scheme 7

[0271]



[0272]

wherein R¹⁷ is an optionally substituted C₁₋₆ alkyl group, and
20 the other symbols are as defined above.

[0273]

Compound (33d) may be commercially available, or can be
produced according to a method known per se or a method
analogous thereto.

25 [0274]

<Step 55> Compound (33e) can be produced by reacting compound
(33d) with the alcohol (R¹⁷OH) in the presence of a
palladium(II) catalyst.

Examples of the palladium(II) catalyst include
30 bis(acetonitrile)palladium chloride, palladium chloride,
palladium acetate and the like. The amount of the

palladium(II) catalyst to be used is generally about 0.005-1 mol, preferably about 0.01-1 mol, per 1 mol of compound (33d).

The amount of the alcohol ($R^{17}OH$) to be used is generally about 1-10 mol, preferably about 1-3 mol, per 1 mol of
5 compound (33d).

This reaction is advantageously carried out in a solvent inert to the reaction. The solvent is not particularly limited as long as the reaction proceeds. Preferable examples thereof include halogenated hydrocarbons, alcohols corresponding to
10 $R^{17}OH$, ethers, hydrocarbons, mixed solvents thereof and the like.

While the reaction time varies depending on the reagent and solvent to be used, it is generally 0.5-48 hr, preferably 1-24 hr.

15 The reaction temperature is generally 0-100°C, preferably 20-80°C.
[0275]

Compound (1) obtained in each reaction scheme can be isolated and purified by known separation and purification
20 means such as concentration, concentration under reduced pressure, solvent extraction, crystallization, recrystallization, phase transfer, chromatography and the like. In addition, each material compound used in each reaction scheme can be isolated and purified by those similar to the
25 above-mentioned known separation and purification means. The material compound may be used directly in the next step as the reaction mixture without isolation.

When compound (1) has isomers such as an optical isomer, a stereoisomer, a regioisomer and a rotamer and the like, such
30 isomers and a mixture thereof are also encompassed in compound (1). For example, when compound (1) has an optical isomer, the optical isomer resolved from racemate is also encompassed in compound (I). These isomers can be obtained as single products according to synthetic methods known per se, separation
35 methods known per se (e.g., concentration, solvent extraction,

column chromatography, recrystallization etc.), optical resolutions (e.g., fractional recrystallization method, chiral column method, diastereomer method and the like).

Compound (1) may be a crystal, and the crystal form may
5 be single or a mixture of crystal forms, both of which are encompassed in compound (1). The crystal can be produced according to a crystallization method known per se.

The compound (1) may be a solvate (e.g., hydrate) or a non-solvate (e.g., non-hydrate etc.) and both are encompassed
10 in compound (1).

The compounds labeled with isotopes (e.g., ^3H , ^{14}C , ^{35}S , ^{125}I etc.) and the like are also encompassed in compound (1).

A deuterium conversion form wherein ^1H is converted to $^2\text{H(D)}$ is also encompassed in compound (1).

15 Compound (1) labeled or substituted with an isotope can be used as, for example, a tracer (PET tracer) used for Positron Emission Tomography (PET), and therefore, it is useful in the fields of medical diagnosis and the like.
[0276]

20 Since the compound of the present invention has a superior PDE2A inhibitory action, shows low toxicity (e.g., phototoxicity, acute toxicity, chronic toxicity, genetic toxicity, reproductive toxicity, cardiac toxicity, drug interactions, carcinogenicity and the like, particularly
25 phototoxicity), and is superior in stability (particularly metabolic stability), pharmacokinetics (absorption, distribution, metabolism, excretion etc.) and high solubility, it is useful as a medicament. The compound of the present invention has a PDE2A inhibitory action to mammals (e.g.,
30 mouse, rat, hamster, rabbit, cat, dog, bovine, horse, sheep, monkey, human etc.), and can be used for the prophylaxis or treatment of the following diseases and symptoms:

- (1) psychotic disorder (e.g., brief psychotic disorder, shared psychotic disorder),
- 35 (2) psychosis induced by alcohol, amphetamine, cannabis,

cocaine, hallucinogens, obesity, inhalants, opioids or phencyclidine,

- (3) delusional disorder,
- (4) anxiety disorder,
- 5 (5) movement disorder,
- (6) mood disorder,
- (7) major depressive disorder,
- (8) a major depressive disorder superimposed on a psychotic disorder (including delusional disorder and schizophrenia),
- 10 (9) major depressive episode of the mild, moderate or severe type,
- (10) manic or mixed mood episode,
- (11) hypomanic mood episode,
- (12) depressive episode with atypical features,
- 15 (13) depressive episode with melancholic features,
- (14) depressive episode with catatonic features,
- (15) mood episode with postpartum onset;
- (16) post-stroke depression,
- (17) dysthymic disorder,
- 20 (18) minor depressive disorder,
- (19) autism;
- (20) drug addiction,
- (21) neurodegenerative disorder,
- (22) neurodegeneration associated with cerebral trauma,
- 25 (23) neurodegeneration associated with stroke,
- (24) neurodegeneration associated with cerebral infarct,
- (25) neurodegeneration associated with hypoglycemia,
- (26) neurodegeneration associated with epileptic seizure,
- (27) neurodegeneration associated with neurotoxin poisoning,
- 30 (28) multi-system atrophy,
- (29) Alzheimer's disease,
- (30) dementia,
- (31) multi-infarct dementia,
- (32) alcoholic dementia or other drug-related dementia,
- 35 (33) dementia associated with intracranial tumors or cerebral

- trauma,
- (34) dementia associated with Huntington's disease or Parkinson's disease,
 - (35) AIDS-related dementia,
 - 5 (36) frontotemporal dementia,
 - (37) delirium,
 - (38) amnestic disorder,
 - (39) post-traumatic stress disorder,
 - (40) mental retardation,
 - 10 (41) learning disorder (e.g., reading disorder, mathematics disorder, or a disorder of written expression),
 - (42) attention-deficit/hyperactivity disorder;
 - (43) age-related cognitive decline,
 - (44) premenstrual dysphoric disorder,
 - 15 (45) post-psychotic depressive disorder of schizophrenia,
 - (46) bipolar disorder (including bipolar I disorder and bipolar II disorder),
 - (47) cyclothymic disorder,
 - (48) Parkinson's disease,
 - 20 (49) Huntington's disease,
 - (50) paranoia,
 - (51) schizophrenia (e.g., paranoid schizophrenia, disorganized schizophrenia, catatonic schizophrenia, undifferentiated schizophrenia, residual schizophrenia),
 - 25 (52) schizophreniform disorder,
 - (53) schizoaffective disorder of the delusional type or the depressive type,
 - (54) personality disorder of the paranoid type,
 - (55) personality disorder of the schizoid type,
 - 30 (56) obesity,
 - (57) metabolic syndrome,
 - (58) non-insulin dependent diabetes (NIDDM),
 - (59) glucose intolerance,
 - (60) pneumonia, and/or
 - 35 (61) osteoarthritis.

[0277]

In particular, the compound of the present invention is useful for the prophylaxis or treatment of schizophrenia and Alzheimer's disease.

5 [0278]

Since the compound of the present invention is superior in metabolic stability, it can be expected to have an excellent therapeutic effect on the above-mentioned diseases even in a low dose.

10 Since the compound of the present invention has low toxicity, a pharmaceutical composition containing the compound of the present invention (hereinafter to be referred to as the "medicament of the present invention") is obtained as, for example, tablet (including sugar-coated tablet, film-coated
15 tablet, sublingual tablet, orally disintegrating tablet, buccal tablet and the like), pill, powder, granule, capsule (including soft capsule, microcapsule), troche, syrup, liquid, emulsion, suspension, controlled-release preparation (e.g., immediate-release preparation, sustained-release preparation,
20 sustained-release microcapsule), aerosol, films (e.g., orally disintegrable films, oral cavity mucosa patch film), injection (e.g., subcutaneous injection, intravenous injection, intramuscular injection, intraperitoneal injection), drip infusion, transdermal absorption type preparation, ointment,
25 lotion, adhesive preparation, suppository (e.g., rectal suppository, vaginal suppository), pellet, nasal preparations, pulmonary preparation (inhalant), eye drop and the like by using the compound of the present invention alone or along with a pharmacologically acceptable carrier according to a
30 method known per se as a production method of pharmaceutical preparations (e.g., the method described in the Japanese Pharmacopoeia etc.). It can be safely administered orally or parenterally (e.g., intravenous, intramuscular, subcutaneous, intraorgan, intranasal, intradermal, instillation,
35 intracerebral, rectal, vaginal, intraperitoneal, intratumor,

tumor proximal administration, administration to a lesion and the like).

[0279]

Examples of the pharmacologically acceptable carrier
5 include various organic or inorganic carrier substances conventionally used as preparation materials, which are added as excipient, lubricant, binder or disintegrant for solid preparations; as solvent, solubilizing agent, suspending agent, isotonicity agent, buffer or soothing agent for liquid
10 preparation, and the like. Where necessary, preparation additives such as preservative, antioxidant, colorant, sweetener, absorbent, humectant and the like can also be appropriately used in suitable amounts.

[0280]

15 Examples of the excipient include lactose, sucrose, D-mannitol, D-sorbitol, starch, pregelatinized starch, cornstarch, dextrin, microcrystalline cellulose, low-substituted hydroxypropyl cellulose, sodium carboxymethylcellulose, gum arabic, pullulan, light anhydrous
20 silicic acid, synthetic aluminum silicate, magnesium aluminate metasilicate and the like.

Examples of the lubricant include magnesium stearate, calcium stearate, talc, colloidal silica and the like.

[0281]

25 Examples of the binder include pregelatinized starch, microcrystalline cellulose, sucrose, gum arabic, D-mannitol, trehalose, dextrin, pullulan, hydroxypropyl cellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone, starch, saccharose, gelatin, methylcellulose, carboxymethylcellulose,
30 sodium carboxymethylcellulose and the like.

[0282]

Examples of the disintegrant include lactose, sucrose, starch, carboxymethylcellulose, calcium carboxymethylcellulose, croscarmellose sodium, sodium carboxymethyl starch, light
35 anhydrous silicic acid, low-substituted hydroxypropyl

cellulose and the like.

[0283]

Examples of the solvent include water for injection, physiological saline, Ringer's solution, alcohol, propylene glycol, polyethylene glycol, macrogol, sesame oil, corn oil, olive oil, cottonseed oil and the like.

[0284]

Examples of the solubilizing agent include polyethylene glycol, propylene glycol, D-mannitol, trehalose, benzyl benzoate, ethanol, tris(hydroxymethyl)aminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate, sodium salicylate, sodium acetate and the like.

[0285]

Examples of the suspending agent include surfactants such as stearyltriethanolamine, sodium lauryl sulfate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzethonium chloride, glyceryl monostearate and the like; and hydrophilic polymers such as polyvinyl alcohol, polyvinylpyrrolidone, sodium carboxymethylcellulose, methylcellulose, hydroxymethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, polysorbate, polyoxyethylene hydrogenated castor oil and the like.

[0286]

Examples of the isotonicity agent include glucose, D-sorbitol, sodium chloride, glycerol, D-mannitol and the like.

Examples of the buffer include buffers such as phosphates, acetates, carbonates, citrates and the like, and the like.

Examples of the soothing agent include benzyl alcohol and the like.

Examples of the preservative include p-hydroxybenzoate, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, sorbic acid and the like.

Preferred examples of the antioxidant include sulfites, ascorbic acid, α -tocopherols and the like.

[0287]

Examples of the colorant include water-soluble edible tar pigments (e.g., Food Color Red Nos. 2 and 3, Food Color Yellow Nos. 4 and 5, Food Color Blue Nos. 1 and 2), water insoluble lake pigments (e.g., aluminum salts of the above-mentioned
5 water-soluble edible tar pigment), natural pigments (e.g., beta-carotene, chlorophyll, red iron oxide) and the like.

Examples of the sweetening agent include saccharin sodium, dipotassium glycyrrhizinate, aspartame, stevia and the like.
[0288]

10 While the content of the compound of the present invention in the medicament of the present invention varies depending on the dosage form, dose of the compound of the present invention, and the like, it is, for example, about 0.1 to 100 wt%, preferably about 0.1-95 wt%.

15 [0289]

While the dose of the compound of the present invention varies depending on the subject of administration, administration route, target disease, symptom and the like, for example, for oral administration to a schizophrenia
20 patient (adult, about 60 kg weight), it is generally about 0.1 - about 20 mg/kg body weight, preferably about 0.2 - about 10 mg/kg body weight, more preferably about 0.5 - about 10 mg/kg body weight, which is desirably administered once to several times (e.g., once to 3 times) a day depending on the symptom.

25 [0290]

The compound of the present invention can be administered as a single active substance, or can be administered in combination with other medicaments such as other drugs used in the treatment of psychotic disorder (particularly
30 schizophrenia and bipolar disorder), obsessive-compulsive disorder, major depression, Parkinson's disease, Alzheimer's disease, cognitive disorder, memory loss and the like, (hereinafter to be abbreviated as concomitant drug).

Examples of the concomitant drug include nicotinic $\alpha 7$
35 agonists, nicotinic $\alpha 7$ partial agonists, nicotinic $\alpha 7$ positive

allosteric modulators, PDE2 inhibitors, PDE4 inhibitors, PDE5 inhibitors, PDE10 inhibitors, other PDE inhibitors, calcium channel blockers, muscarinic m1 and m2 modulators, adenosine receptor modulators, ampakines, Glycine transporter 1
5 inhibitors, NMDA-R modulators, mGluR modulators, dopamine modulators, serotonin modulators, selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, norepinephrine and dopamine reuptake inhibitors, triple reuptake inhibitors, cannabinoid modulators, cholinesterase
10 inhibitors (e.g., donepezil, rivastigmine, galantamine) and the like.

In addition, examples of the concomitant drug include, but are not limited to, other suitable schizophrenia drugs (e.g., Haloperidol, Clozapine, Olanzapine, Risperidone,
15 Aripiprazole, Ziprasidone, Paliperidone, Quetiapine fumarate etc.), bipolar disorder drug (e.g., Lithium, Olanzapine, Aripiprazole, Valproic acid etc.), Parkinson's disease drugs (e.g., Levodopa, Bromocriptine, Pergolide, Pramipexole, Tolcapone, Procyclidine, Trihexyphenidyl, Benztropine etc.),
20 agents used in the treatment of major depression (e.g., Amitriptyline, Imipramine, Desipramine, Nortriptyline, Paroxetine, Fluoxetine, Sertraline, Bupropion, Escitalopram, Mirtazapine, Venlafaxine, Duloxetine etc.), agents used in the treatment of Alzheimer's disease (e.g., Galantamine, Tacrine,
25 Donepezil, Rivastigmine, Memantine, Neotropin, Selegiline, Estrogen, Clioquinol etc.), agents used in the treatment of dementia (e.g., Thioridazine, Haloperidol, Risperidone, Tacrine, Donepezil, Rivastigmine etc.), agents used in the treatment of epilepsy (e.g., Phenytoin, Phenobarbital,
30 Carbamazepine, Valproic acid, Ethosuximide, Gabapentin, Solfeton, Felbatol etc.), agents used in the treatment of multiple sclerosis (e.g., Tolterodine, Oxybutynin, Oxycodone, Interferon beta-1b, Interferon beta-1a, Azathioprine, Methotrexate, Glatiramer etc.), agents used in the treatment
35 of Huntington's disease (e.g., Amitriptyline, Imipramine,

Desipramine, Nortriptyline, Paroxetine, Fluoxetine, Sertraline, Tetrabenazine, Haloperidol, Chlorpromazine, Thioridazine, Sulpiride, Quetiapine, Clozapine, Risperidone etc.), agents useful in the treatment of diabetes [e.g., PPAR ligands (e.g. 5 agonists or antagonists such as Rosiglitazone, Troglitazone, Pioglitazone etc.), insulin secretagogues (e.g., sulfonylurea drugs such as Glyburide, Glimepiride, Chlopropamide, Tolbutamide, Glipizide etc., and non-sulfonyl secretagogues), α -glucosidase inhibitors (e.g., Acarbose, Miglitol, Voglibose 10 etc), insulin sensitizers (e.g., PPAR- γ agonists (e.g., the glitazones); biguanides, PTP-1B inhibitors, DPP-IV inhibitors, 11beta-HSD inhibitors etc.), hepatic glucose output lowering compounds (e.g., glucagon antagonists and metformin(e.g., Glucophage, Glucophage XR etc.)), insulin and insulin 15 derivatives (including both long and short acting forms and formulations of insulin)], antiobesity drugs [e.g., β -3 agonists, CB-1 agonists, neuropeptide Y5 inhibitors, Ciliary Neurotrophic Factor and derivatives (e.g., Axokine), appetite suppressants (e.g., Sibutramine), lipase inhibitors (e.g., 20 Orlistat) etc.].

[0291]

The dosage form of concomitant drugs is not particularly limited, and is acceptable as long as the compound of the present invention is combined with concomitant drugs at the 25 time of administration. Examples of such dosage forms are as follows:

- (1) Administration of a single formula obtained simultaneous formulation of the compound of the present invention with a concomitant drug,
- 30 (2) Simultaneous administration via the same administration route for two kinds of formulas obtained by independent formulations of the compound of the present invention and a concomitant drug,
- (3) Administrations at different times via the same 35 administration route for two kinds of formulas obtained by

independent formulations of the compound of the present invention and a concomitant drug,

(4) Simultaneous administration via different administration routes for two kinds of formulas obtained by independent
5 formulations of the compound of the present invention and a concomitant drug,

(5) Administrations at different times via different administration routes for two kinds of formulas obtained by independent formulations of the compound of the present
10 invention and a concomitant drug (e.g., administration in the order of the compound of the present invention and then a concomitant drug, or administration in the reversed order).

These forms of administration are summarized below and abbreviated as a combination drug of the present invention.

15 [0292]

When administering the combination drug of the present invention, the concomitant drug and the compound of the present invention can be administered simultaneously. Alternatively, the compound of the present invention can be
20 administered after a concomitant drug is administered, or a concomitant drug can be administered after the compound of the present invention is administered. When administering at different times, the time difference depends upon the active ingredients to be administered, drug forms and methods of
25 administration.

For example, when the concomitant drug or a pharmaceutical composition thereof is administered first, the compound of the present invention or a pharmaceutical composition thereof can be administered within 1 min. to 3
30 days, preferably within 10 min to 1 day and more preferably within 15 min to 1 hour after the concomitant drug or a pharmaceutical composition thereof is administered. When the compound of the present invention or a pharmaceutical composition thereof is administered first, the concomitant
35 drug or a pharmaceutical composition thereof can be

administered within 1 min to 1 day, preferably within 10 min to 6 hours and more preferably within 15 min to 1 hour after the compound of the present invention or a pharmaceutical composition thereof is administered.

5 [0293]

If there are no problems with side effects of the concomitant drugs, any dosages can be set. A dosage as a concomitant drug varies depending on dosages, administration subjects, administration routes, target diseases, symptoms,
10 etc. For example, in the case of oral administration in patients with schizophrenia (adults, body weight of approximately 60 kg), a dosage range is generally about 0.1 to 20 mg/kg body weight, preferably from about 0.2 to 10 mg/kg body weight and more preferably from about 0.5 to 10 mg/kg
15 body weight. It is preferable that this dosage is administered once daily to several times daily (e.g., once to 3 times).

If the compound of the present invention is used in combination with a concomitant drug, the respective dosages can be reduced within a safe range with consideration of the
20 opposite effects of the respective drugs.

[0294]

The combination drug of the present invention exhibits low toxicity. For example, the compound of the present invention or (and) the concomitant drug can be combined with a
25 pharmaceutically acceptable carrier according to the known method to prepare a pharmaceutical composition such as tablets (including sugar-coated tablets and film-coated tablets), powder agents, granular agents, capsules (including soft capsules), liquids, injection solutions, suppositories,
30 sustained-release agents, etc. These compositions can be administered safely orally or non-orally (e.g., including local, rectal, venous routes etc.).

[0295]

The pharmaceutically acceptable carriers that can be used
35 for manufacturing the combination drug of the present

invention can be the same as those used in the medicament of the present invention as mentioned above.

[0296]

A mixing ratio between the compound of the present invention and the concomitant drug in the combination drug of the present invention can be selected appropriately based on the administration subjects, administration routes, diseases and the like.

The concomitant drug in the combination drug of the present invention can be combined at an appropriate proportion if two or more drugs are combined.

A dosage of the concomitant drug can be selected appropriately based on the dosages used clinically. In addition, a mixing ratio between the compound of the present invention and the concomitant drug can be selected appropriately based on the administration subjects, administration routes, target diseases, symptoms, combinations, etc. For example, if the administration subject is humans, a concomitant drug may be used in an amount ranging from about 0.01 to 100 parts by weight relative to 1 part by weight of the compound of the present invention.

[0297]

For example, the content of the compound of the present invention in the combination drug of the present invention varies with the form of formulations. Generally, it is present in a range from about 0.01 to 99.9 wt%, preferably from about 0.1 to about 50 wt% and more preferably from about 0.5 to about 20 wt% relative to the entire formula.

[0298]

The content of the concomitant drug in the combination drug of the present invention varies with the form of formulations. Generally it is present in a range from about 0.01 to 99.9 wt%, preferably from about 0.1 to about 50 wt% and more preferably from about 0.5 to about 20 wt% relative to the entire formula.

[0299]

The content of an additive such as carriers in the combination drug of the present invention varies with the form of formulations. Generally it is present in a range from about
5 1 to 99.99 wt% and preferably from about 10 to about 90 wt% relative to the entire formula.

[0300]

When the compound of the present invention and a concomitant drug are formulated independently, the same
10 contents can be applied.

[0301]

Since the dosages may fluctuate under various conditions as mentioned above, a dosage less than the dosages may be sufficient or it may be necessary to administer at a dosage
15 exceeding the range.

Example

[0302]

The present invention is explained in detail in the following by referring to Examples, Formulation Examples and
20 Experimental Examples and which are merely exemplified and not to be construed as limitative, and the invention may be changed within the scope of the present invention. In the following Examples, the "room temperature" generally means about 10°C to about 35°C. The ratios indicated for mixed
25 solvents are volume mixing ratios, unless otherwise specified. % means wt%, unless otherwise specified.

In silica gel column chromatography, NH means use of aminopropylsilane-bonded silica gel. In HPLC (high performance liquid chromatography), C18 means use of octadecyl-bound
30 silica gel. The ratios of elution solvents are volume mixing ratios, unless otherwise specified.

¹H NMR (proton nuclear magnetic resonance spectrum) was measured by Fourier-transform type NMR. Peaks with very mild protons such as a hydroxyl group, an amino group and the like
35 are not described.

In the following Reference Examples and Examples, mass spectrum (MS), nuclear magnetic resonance spectrum (NMR) and melting point were measured by the following apparatus.

MS (mass spectrum) was measured by LC/MS (liquid chromatograph mass spectrometer). As the ionization method, API (Atmospheric Pressure Ionization, atmospheric pressure chemical ionization) method or ESI (Electron Spray Ionization) method was used. The data indicate measured value (found) found. Generally, a molecular ion peak is observed. In the case of a compound having an amino group ($-NH_2$), a peak after elimination of NH_3 may be observed as a fragment ion. In the case of a salt, a molecular ion peak or fragment ion peak of free form is generally observed.

[0303]

Example 1

N-(1-(4-methoxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0304]

A) methyl N-(3-nitropyridin-2-yl)glycinate

To a solution of methyl glycinate hydrochloride (1.98 g) and 2-chloro-3-nitropyridine (2.50 g) in N,N-dimethylformamide (16.0 mL) was added triethylamine (5.50 mL). The reaction mixture was stirred overnight at 80°C, and the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the ethyl acetate solution was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (2.25 g).

MS (API+): $[M+H]^+$ 212.0.

[0305]

B) 3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(3-nitropyridin-2-yl)glycinate (50.0 mg) in ethanol (3.00 mL) was added 10% palladium-carbon

(containing 50% water, 50.0 mg). The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, and the catalyst was filtered off. The filtrate was stirred overnight at 70°C, and the solvent was evaporated under reduced
5 pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate) to give the title compound (23.5 mg).

MS (API+): [M+H]⁺ 150.2.

[0306]

10 C) N-hydroxy-1-(4-methoxyphenyl)propan-1-imine

To a solution of 1-(4-methoxyphenyl)propan-1-one (25.0 g) in ethanol (300 mL) were added hydroxylamine hydrochloride (11.6 g) and triethylamine (17.0 g). The reaction mixture was heated at reflux for 3 hr, and the solvent was evaporated
15 under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (26.7 g).

20 MS (API+): [M+H]⁺ 180.0.

[0307]

D) 1-(4-methoxyphenyl)propan-1-amine

To a solution of N-hydroxy-1-(4-methoxyphenyl)propan-1-imine (12.4 g) in ethanol (300 mL) was added 10% palladium-
25 carbon (containing 50% water, 8.19 g). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (NH,
30 hexane/ethyl acetate) to give the title compound (7.56 g).

¹H NMR (300 MHz, DMSO-d₆) δ 0.76 (3H, t, J = 7.4 Hz), 1.30-1.64 (2H, m), 1.87 (2H, brs), 3.63 (1H, t, J = 6.7 Hz), 3.72 (3H, s), 6.74-6.91 (2H, m), 7.15-7.29 (2H, m).

[0308]

35 E) 1-(4-methoxyphenyl)propan-1-amine hydrochloride

To a solution of 1-(4-methoxyphenyl)propan-1-amine (6.74 g) in ethyl acetate (20 mL) was added 1 M hydrogen chloride/diethyl ether solution (50 mL). The reaction mixture was diluted with ethyl acetate, and the precipitated solid was collected by filtration. The obtained solid was washed with ethyl acetate, and dried under reduced pressure to give the title compound (7.46 g).

¹H NMR (300 MHz, DMSO-d₆) δ 0.73 (3H, t, J = 7.4 Hz), 1.66-1.87 (1H, m), 1.89-2.10 (1H, m), 3.76 (3H, s), 3.96-4.11 (1H, m), 6.91-7.03 (2H, m), 7.36-7.50 (2H, m), 8.51 (3H, brs).

[0309]

F) N-(1-(4-methoxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (23.5 mg) in a mixed solvent of tetrahydrofuran (5 mL) and N,N-dimethylacetamide (1 mL) were added triethylamine (65.9 μL) and bis(trichloromethyl)carbonate (46.8 mg) at 0°C. The reaction mixture was stirred at 0°C for 2 hr, triethylamine (65.9 μL) and 1-(4-methoxyphenyl)propan-1-amine hydrochloride (159 mg) were added thereto at 0°C. The reaction mixture was stirred at room temperature for 2 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, ethyl acetate/hexane), and crystallized from hexane/isopropyl ether to give the title compound (6.3 mg).

MS (API-): [M-H]⁻ 339.0.

¹H NMR (300 MHz, CDCl₃) δ 0.79-1.00 (3H, m), 1.76-1.97 (2H, m), 3.78 (3H, s), 4.68 (2H, s), 4.83 (1H, q, J = 7.2 Hz), 6.86 (2H, d, J = 8.7 Hz), 6.96 (1H, dd, J = 7.7, 5.1 Hz), 7.08-7.37 (3H, m), 7.97 (1H, dd, J = 5.1, 1.3 Hz), 9.69 (1H, brs), 10.35 (1H, d, J = 7.6 Hz).

[0310]

Example 2

2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0311]

5 A) 4-(trifluoromethoxy)benzoyl chloride

To a suspension of 4-(trifluoromethoxy)benzoic acid (25.0 g) in toluene (300 mL) was added thionyl chloride (17.3 g) at room temperature. To the reaction mixture was added N,N-dimethylacetamide (1 mL) solution, the mixture was stirred at
10 60°C for 2 hr, and the solvent was evaporated under reduced pressure to give the title compound (25.1 g).

¹H NMR (300 MHz, DMSO-d₆) δ 7.39-7.52 (2H, m), 8.01-8.11 (2H, m).

[0312]

15 B) N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide

To a solution of 4-(trifluoromethoxy)benzoyl chloride (25.0 g) in toluene (240 mL) were added N-methoxymethanamine hydrochloride (13.0 g) and N-ethyl-N-(propan-2-yl)propan-2-amine (36.0 g) at room temperature. The reaction mixture was
20 stirred at 60°C for 1 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate. The residue was purified by silica gel column chromatography (NH, ethyl acetate), and the solvent was
25 evaporated under reduced pressure to give the title compound (16.1 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.27 (3H, s), 3.55 (3H, s), 7.35-7.51 (2H, m), 7.64-7.83 (2H, m).

[0313]

30 C) 1-(4-(trifluoromethoxy)phenyl)propan-1-one

To a solution of N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide (16.1 g) in tetrahydrofuran (200 mL) was added 1 M ethylmagnesium bromide/tetrahydrofuran solution (97 mL) under ice-cooling. The reaction mixture was
35 stirred at room temperature for 3 hr, 0.5 M hydrochloric acid

was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with 0.5 M hydrochloric acid and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under
5 reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (12.7 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.09 (3H, t, J = 7.2 Hz), 3.07 (2H, q, J = 7.2 Hz), 7.45-7.56 (2H, m), 8.04-8.15 (2H, m).

10 [0314]

D) N-hydroxy-1-(4-(trifluoromethoxy)phenyl)propan-1-imine

To a solution of 1-(4-(trifluoromethoxy)phenyl)propan-1-one (12.7 g) in ethanol (120 mL) were added hydroxylamine hydrochloride (4.86 g) and triethylamine (7.08 g). The
15 reaction mixture was heated at reflux for 4 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give
20 the title compound (13.8 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.04 (3H, t, J = 7.6 Hz), 2.72 (2H, q, J = 7.6 Hz), 7.32-7.44 (2H, m), 7.71-7.81 (2H, m), 11.31 (1H, s).

[0315]

25 E) 1-[4-(trifluoromethoxy)phenyl]propan-1-amine

To a solution of N-hydroxy-1-(4-(trifluoromethoxy)phenyl)propan-1-imine (13.8 g) in ethanol (130 mL) was added 10% palladium-carbon (containing 50% water, 3.0 g). The reaction mixture was stirred overnight at room
30 temperature under hydrogen atmosphere, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure to give the title compound (11.3 g).

^1H NMR (400 MHz, DMSO- d_6) δ 0.77 (3H, t, J = 7.2 Hz), 1.48-1.63 (2H, m), 3.76 (1H, t, J = 6.8 Hz), 7.27 (2H, d, J = 8.4 Hz),
35 7.45 (2H, d, J = 8.4 Hz).

[0316]

F) 1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride

To a solution of 1-(4-(trifluoromethoxy)phenyl)propan-1-amine (11.3 g) in a mixed solvent of ethyl acetate (70 mL) and
5 methanol (10 mL) was added 4 M hydrogen chloride/ethyl acetate solution (40 mL). The reaction mixture was stirred for 20 min, the solvent was evaporated under reduced pressure, and diethyl ether was added thereto. The precipitated solid was collected by filtration, washed with diethyl ether, and dried under
10 reduced pressure to give the title compound (10.12 g).

¹H NMR (300 MHz, DMSO-d₆) δ 0.76 (3H, t, J = 7.4 Hz), 1.70-2.11 (2H, m), 4.20 (1H, dd, J = 9.1, 5.7 Hz), 7.39-7.53 (2H, m), 7.61-7.71 (2H, m), 8.63 (3H, brs).

[0317]

15 G) 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step F of Example 1.

MS (API+): [M+H]⁺ 395.2.

20 ¹H NMR (300 MHz, CDCl₃) δ 0.96 (3H, t, J = 7.4 Hz), 1.76-1.96 (2H, m), 4.67 (2H, s), 4.90 (1H, q, J = 7.2 Hz), 6.99 (1H, dd, J = 7.6, 4.9 Hz), 7.12-7.25 (3H, m), 7.30-7.42 (2H, m), 7.99 (1H, dd, J = 5.1, 1.7 Hz), 9.91 (1H, s), 10.46 (1H, d, J = 7.2 Hz).

25 [0318]

Example 3

N-(1-(4-methoxyphenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0319]

30 A) methyl N-(5-methyl-3-nitropyridin-2-yl)glycinate

The title compound was obtained in the same manner as in Step A of Example 1.

MS (API+): [M+H]⁺ 226.1.

[0320]

35 B) 7-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

hydrochloride

To an ethanol solution (100 mL) of methyl N-(5-methyl-3-nitropyridin-2-yl)glycinate (1.16 g) was added 10% palladium-carbon (containing 50% water, 1.00 g). The reaction mixture
5 was stirred at room temperature for 2 hr under hydrogen atmosphere, and the catalyst was filtered off. The filtrate was stirred at 70°C for 4 hr, 5-10% hydrochloric acid/methanol (10 mL) solution was added thereto, and the mixture was stirred overnight at 70°C. The solvent was evaporated under
10 reduced pressure, and the obtained residue was crystallized from hexane/ethyl acetate to give the title compound (1.02 g). MS (API+): [M+H]⁺ 164.2.

[0321]

C) N-(1-(4-methoxyphenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
15

The title compound was obtained in the same manner as in Step F of Example 1.

MS (API+): [M+H]⁺ 355.1.

¹H NMR (300 MHz, DMSO-d₆) δ 0.82 (3H, t, J = 7.2 Hz), 1.67-1.83
20 (2H, m), 2.24 (3H, s), 3.72 (3H, s), 4.39 (2H, d, J = 0.8 Hz), 4.70 (1H, q, J = 7.1 Hz), 6.83-6.92 (2H, m), 7.12 (1H, d, J = 1.5 Hz), 7.17-7.26 (2H, m), 7.81-7.88 (1H, m), 10.11 (1H, d, J = 7.6 Hz), 10.78 (1H, brs).

[0322]

25 Example 4

7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one hydrochloride (200 mg) in a mixed solvent
30 of tetrahydrofuran (10 mL) and N,N-dimethylacetamide (10 mL) were added triethylamine (838 µL) and a solution of bis(trichloromethyl)carbonate (297 mg) in tetrahydrofuran (3 mL) at 0°C. The reaction mixture was stirred at 0°C for 3 hr, triethylamine (698 µL) and a solution of 1-[4-
35 (trifluoromethoxy)phenyl]propan-1-amine hydrochloride (1.28 g)

in a mixed solvent of tetrahydrofuran (3 mL) and N,N-dimethylacetamide (5 mL) were added thereto at 0°C. The reaction mixture was stirred at room temperature for 1 hr, water was added thereto, and the mixture was extracted with
5 ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate) to give a crude product. The crude product was
10 purified by HPLC (C18, mobile phase: water/acetonitrile (containing 10 mM NH₄HCO₃)), the obtained fraction was concentrated under reduced pressure, and the residue was crystallized from hexane/ethyl acetate to give the title compound (56.3 mg).

15 MS (API+): [M+H]⁺ 409.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.76-0.95 (3H, m), 1.78 (2H, quin, J = 7.2 Hz), 2.25 (3H, s), 4.38 (2H, s), 4.80 (1H, q, J = 7.2 Hz), 7.13 (1H, d, J = 1.9 Hz), 7.28-7.37 (2H, m), 7.38-7.48 (2H, m), 7.86 (1H, d, J = 1.5 Hz), 10.21 (1H, d, J = 7.2
20 Hz), 10.81 (1H, brs).

[0323]

Example 5

optically active 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-
25 b]pyrazine-4(1H)-carboxamide

Racemic 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (56.3 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel
30 Chemical Industries, Ltd., mobile phase: hexane/2-propanol = 850/150) to give the title compound having a shorter retention time (10.2 mg).

MS (API+): [M+H]⁺ 409.1.

¹H NMR (300 MHz, CDCl₃) δ 0.95 (3H, t, J = 7.4 Hz), 1.85 (2H, quin, J = 7.3 Hz), 2.31 (3H, s), 4.66 (2H, s), 4.89 (1H, q, J

=6.9 Hz), 7.03 (1H, d, J = 1.5 Hz), 7.16 (2H, d, J = 8.0 Hz), 7.30-7.39 (2H, m), 7.83 (1H, d, J = 1.1 Hz), 9.31 (1H, s), 10.33 (1H, d, J = 7.2 Hz).

[0324]

5 Example 6

optically active 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (56.3 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/2-propanol = 850/150) to give the title compound having a longer retention
15 time (9.00 mg).

MS (API+): [M+H]⁺ 409.1.

¹H NMR (300 MHz, CDCl₃) δ 0.95 (3H, t, J = 7.4 Hz), 1.85 (2H, quin, J = 7.3 Hz), 2.31 (3H, s), 4.66 (2H, s), 4.89 (1H, q, J = 6.9 Hz), 7.04 (1H, d, J = 1.5 Hz), 7.16 (2H, d, J = 8.0 Hz),
20 7.29-7.38 (2H, m), 7.83 (1H, d, J = 1.1 Hz), 9.49 (1H, s), 10.33 (1H, d, J = 7.2 Hz).

[0325]

Example 7

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0326]

A) 4-ethenylphenyl trifluoromethyl ether

To a suspension of potassium 2-methylpropan-2-olate (6.10 g) in tetrahydrofuran (80 mL) was added
30 methyl(triphenyl)phosphonium iodide (22.1 g) over 5 min under ice-cooling. The reaction mixture was stirred for 30 min, and a solution of 4-(trifluoromethoxy)benzaldehyde (8.00 g) in tetrahydrofuran (20 mL) was added slowly thereto. The reaction mixture was stirred at room temperature for 1.5 hr, water was
35 added thereto, and the mixture was extracted with ethyl

acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum
5 ether) to give the title compound (5.30 g).

^1H NMR (300 MHz, CDCl_3) δ 5.30 (1H, d, $J = 10.8$ Hz), 5.75 (1H, d, $J = 17.7$ Hz), 6.72 (1H, dd, $J = 17.7, 10.8$ Hz), 7.19 (2H, d, $J = 7.8$ Hz), 7.44 (2H, d, $J = 6.9$ Hz).

[0327]

10 B) 2-(4-(trifluoromethoxy)phenyl)oxirane

To a solution of 4-ethenylphenyl trifluoromethyl ether (5.30 g) in chloroform (50 mL) was added 3-chloroperbenzoic acid (11.4 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and washed successively
15 with aqueous sodium thiosulfate solution and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/dichloromethane) to give the title compound (4.20 g).

20 ^1H NMR (400 MHz, CDCl_3) δ 2.76 (1H, dd, $J = 5.6, 2.4$ Hz), 3.16 (1H, dd, $J = 5.2, 4.0$ Hz), 3.87 (1H, dd, $J = 4.0, 2.8$ Hz), 7.20 (2H, d, $J = 8.0$ Hz), 7.31 (2H, d, $J = 8.0$ Hz).

[0328]

C) 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanol

25 To a solution of 2-(4-(trifluoromethoxy)phenyl)oxirane (4.20 g) in *N,N*-dimethylacetamide (20 mL) was added sodium methoxide (5.56 g) at room temperature. The reaction mixture was stirred at 60°C for 3 hr. The reaction mixture was cooled to room temperature, and ethyl acetate was added thereto. The
30 mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane/ethyl acetate) to give the title compound (2.75 g).

35 ^1H NMR (300 MHz, CDCl_3) δ 2.87 (1H, d, $J = 2.7$ Hz), 3.33-3.45

(4H, m), 3.55 (1H, dd, J = 9.6, 3.0 Hz), 4.92 (1H, d, J = 8.7 Hz), 7.22 (2H, d, J = 8.1 Hz), 7.43 (2H, d, J = 8.7 Hz).

[0329]

D) 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanone

5 To a solution of 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanol (2.70 g) in dichloromethane (50 mL) was added 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one (6.28 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and
10 the insoluble substance was filtered off. The filtrate was washed successively with sodium thiosulfate and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane)
15 to give the title compound (2.00 g).

¹H NMR (400 MHz, CDCl₃) δ 3.51 (3H, s), 4.68 (2H, s), 7.30 (2H, d, J = 8.0 Hz), 8.02 (2H, d, J = 9.2 Hz).

[0330]

E) 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanamine

20 To a solution of 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanone (2.00 g) in ethanol (80 mL) were added potassium carbonate (4.70 g) and hydroxylamine hydrochloride (1.18 g) at room temperature. The reaction mixture was heated at reflux overnight, and the solvent was
25 evaporated under reduced pressure. To the residue was added ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (1.50 g) of N-hydroxy-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanimine. To a solution of the
30 obtained crude product (1.50 g) of N-hydroxy-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanimine in methanol (50 mL) was added 10% palladium-carbon (containing 50% water, 0.20 g) at room temperature. The reaction mixture was stirred overnight
35 at room temperature under hydrogen atmosphere. The insoluble

substance was filtered off, and the filtrate was concentrated under reduced pressure to give the title compound (1.3 g).

¹H NMR (300 MHz, CDCl₃) δ 3.72-3.40 (4H, m), 3.49 (1H, dd, J = 9.0, 3.0 Hz), 4.22 (1H, dd, J = 8.1, 3.6 Hz), 7.20 (2H, d, J = 8.1 Hz), 7.42 (2H, d, J = 8.1 Hz).

[0331]

F) 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

To a solution of 3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.00 g) in a mixed solvent of N,N-dimethylacetamide (60 mL) and pyridine (10 mL) was added slowly 4-nitrophenyl chloroformate (1.62 g) under ice-cooling. The reaction mixture was stirred at room temperature for 1.5 hr, and water was added thereto. The precipitated solid was collected by filtration, and washed with water to give the title compound (2.09 g).

MS (API+): [M+H]⁺ 315.0.

[0332]

G) N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (157 mg) in N,N-dimethylacetamide (5 mL) were added 2-methoxy-1-[4-(trifluoromethoxy)phenyl]ethanamine (118 mg) and triethylamine (191 µL) at room temperature. The reaction mixture was stirred at room temperature for 3 hr, aqueous sodium hydrogen carbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with 1 M hydrochloric acid and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (102 mg).

MS (API+): [M+H]⁺ 411.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.55-3.71 (2H, m),

4.33-4.50 (2H, m), 4.99-5.13 (1H, m), 7.11 (1H, dd, J = 7.9, 4.9 Hz), 7.27-7.39 (3H, m), 7.42-7.53 (2H, m), 8.00 (1H, dd, J = 4.9, 1.7 Hz), 10.51 (1H, d, J = 7.4 Hz), 10.83 (1H, brs).
[0333]

5 Example 8

optically active 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
10 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (907 mg)
was resolved by HPLC (column: CHIRALPAK IC, 50 mmID×500 mmL,
manufactured by Daicel Chemical Industries, Ltd., mobile
phase: hexane/ethanol = 400/600) to give the title compound
having a shorter retention time (429 mg).

15 MS (API+): [M+H]⁺ 395.1.

¹H NMR (300 MHz, DMSO-d₆) δ 0.88 (3H, t, J = 7.3 Hz), 1.70-1.90
(2H, m), 4.41 (2H, s), 4.81 (1H, q, J = 7.0 Hz), 7.11 (1H, dd,
J = 7.7, 5.1 Hz), 7.26-7.38 (3H, m), 7.38-7.51 (2H, m), 8.02
(1H, dd, J = 4.9, 1.5 Hz), 10.35 (1H, d, J = 7.2 Hz), 10.83
20 (1H, brs).

[0334]

Example 9

optically active 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-
25 b]pyrazine-4(1H)-carboxamide

Racemic 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (907 mg)
was resolved by HPLC (column: CHIRALPAK IC, 50 mmID×500 mmL,
manufactured by Daicel Chemical Industries, Ltd., mobile
30 phase: hexane/ethanol = 400/600) to give the title compound
having a longer retention time (439 mg).

MS (API+): [M+H]⁺ 395.1.

¹H NMR (300 MHz, DMSO-d₆) δ 0.88 (3H, t, J = 7.3 Hz), 1.70-1.89
(2H, m), 4.41 (2H, s), 4.81 (1H, q, J = 6.9 Hz), 7.10 (1H, dd,
35 J = 7.7, 5.1 Hz), 7.25-7.37 (3H, m), 7.38-7.51 (2H, m), 8.02

(1H, dd, J = 5.1, 1.7 Hz), 10.34 (1H, d, J = 7.5 Hz), 10.82 (1H, brs).

[0335]

Example 10

5 8-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0336]

A) methyl N-(4-methyl-3-nitropyridin-2-yl)glycinate

The title compound was obtained in the same manner as in
10 Step A of Example 1.

MS (API+): [M+H]⁺ 226.1.

[0337]

B) 8-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To an ethanol solution (120 mL) of methyl N-(4-methyl-3-nitropyridin-2-yl)glycinate (1.31 g) was added 10% palladium-carbon (containing 50% water, 1.00 g). The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, and the catalyst was filtered off. The filtrate was stirred overnight at 80°C, and the solvent was evaporated
15 under reduced pressure. The obtained residue was crystallized from hexane/ethyl acetate to give the title compound (870 mg).
20 MS (API+): [M+H]⁺ 164.2.

[0338]

C) 8-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
25 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Steps F-G of Example 7.

MS (API+): [M+H]⁺ 409.1.

¹H NMR (300 MHz, CDCl₃) δ 0.95 (3H, t, J = 7.4 Hz), 1.75-1.95
30 (2H, m), 2.32 (3H, s), 4.54-4.74 (2H, m), 4.89 (1H, q, J = 7.2 Hz), 6.82-6.91 (1H, m), 7.16 (2H, d, J = 7.7 Hz), 7.29-7.40 (2H, m), 7.90 (1H, d, J = 5.1 Hz), 8.33 (1H, brs), 10.33 (1H, d, J = 7.4 Hz).

[0339]

35 Example 11

6-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0340]

A) 4-nitrophenyl 6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

The title compound was obtained in the same manner as in Step A of Example 1, Step B of Example 10 and Step F of Example 7.

MS (API+): [M+H]⁺ 164.2.

[0341]

B) 6-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 409.1.

¹H NMR (300 MHz, CDCl₃) δ 0.96 (3H, t, J = 7.5 Hz), 1.77-1.97 (2H, m), 2.47 (3H, s), 4.65 (2H, s), 4.90 (1H, q, J = 7.0 Hz), 6.83 (1H, d, J = 7.9 Hz), 7.17 (2H, d, J = 7.9 Hz), 7.31-7.39 (2H, m), 7.50-7.58 (1H, m), 8.79 (1H, brs), 10.65 (1H, d, J = 7.2 Hz).

[0342]

Example 12

N-(2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0343]

A) 2-methyl-1-[4-(trifluoromethoxy)phenyl]propan-1-amine hydrochloride

The title compound was obtained in the same manner as in Steps C-F of Example 2.

MS (API+), found: 217.1.

[0344]

B) N-(2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 409.1.

¹H NMR (300 MHz, CDCl₃) δ 0.89-1.02 (6H, m), 1.92-2.18 (1H, m),
4.57-4.73 (2H, m), 4.76-4.92 (1H, m), 6.96-7.06 (1H, m), 7.09-
7.22 (3H, m), 7.24-7.35 (2H, m), 8.03 (1H, dd, J = 5.1, 1.7 Hz),
5 8.74 (1H, brs), 10.60 (1H, d, J = 7.9 Hz).

[0345]

Example 13

N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 [0346]

A) ethyl amino (4-(trifluoromethoxy)phenyl)acetate

To a solution of 1-bromo-4-(trifluoromethoxy)benzene (10
g) in tetrahydrofuran (200 mL) was added 1.6 M n-
butyllithium/hexane solution (31.1 mL) at -78°C, and the
15 mixture was stirred at the same temperature for 50 min under
nitrogen atmosphere. Ethyl 2-chloro-2-oxoacetate (6.23 g) was
added thereto at the same temperature, and the reaction
mixture was stirred overnight at room temperature. The
reaction mixture was added to 1 M hydrochloric acid under ice-
20 cooling, and the mixture was extracted with ethyl acetate. The
extract was washed successively with water and saturated brine,
and dried over anhydrous magnesium sulfate, and the solvent
was evaporated under reduced pressure. The residue was
purified by silica gel column chromatography (ethyl
25 acetate/hexane) to give a crude product (2.50 g) of ethyl oxo-
(4-(trifluoromethoxy)phenyl)acetate. The title compound (590
mg) was obtained in the same manner as in Steps D-E of Example
2 from the obtained crude product

MS (API+): [M+H]⁺ 264.1.

30 [0347]

B) ethyl ((tert-butoxycarbonyl) amino)(4-
(trifluoromethoxy)phenyl)acetate

To a solution of ethyl amino (4-
(trifluoromethoxy)phenyl)acetate (590 mg) in tetrahydrofuran
35 (20 mL) was added tert-butyl dicarbonate (430 mg). The

reaction mixture was stirred at room temperature for 2 days, and the solvent was evaporated under reduced pressure to give the title compound (952 mg).

¹H NMR (300 MHz, CDCl₃) δ 1.22 (3H, t, J = 7.2 Hz), 1.53 (9H, s), 3.99-4.33 (2H, m), 5.32 (1H, d, J = 6.8 Hz), 5.63 (1H, brs), 7.19 (2H, d, J = 7.9 Hz), 7.41 (2H, d, J = 8.7 Hz).

[0348]

C) tert-butyl (2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

10 To a suspension of lithium aluminium hydride (41.8 mg) in tetrahydrofuran (5 mL) was added ethyl ((tert-butoxycarbonyl)amino)(4-(trifluoromethoxy)phenyl)acetate (100 mg) under ice-cooling, and the mixture was stirred at the same temperature for 30 min. Anhydrous magnesium sulfate was added thereto, and
15 then one drop of water and ethyl acetate were successively added thereto, and the insoluble substance was filtered off using Celite. The filtrate was concentrated under reduced pressure to give the title compound (54.8 mg).

MS (API-): [M-H]⁻ 320.1.

20 [0349]

D) N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Tert-butyl (2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (52 mg) was added 2 M
25 hydrogen chloride/ethanol solution (2 mL). The reaction mixture was stirred at 60°C for 2 min, and the solvent was evaporated under reduced pressure. To the residue were added N,N-dimethylformamide (2 mL), triethylamine (0.045 mL) and 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
30 carboxylate (50.9 mg) at room temperature. The reaction mixture was stirred overnight at room temperature, saturated brine was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent
35 was evaporated under reduced pressure. The residue was

purified by silica gel column chromatography (ethyl acetate/hexane) to give a crude product containing N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide. The obtained
5 crude product was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated sodium carbonate aqueous solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous
10 magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (32.2 mg).

MS (API+): [M+H]⁺ 397.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.51-3.80 (2H, m), 4.29-4.53 (2H, m), 5.09 (1H, t, J = 5.1 Hz), 7.11 (1H, dd, J = 7.7, 5.1 Hz),
15 7.28-7.35 (4H, m), 7.40-7.51 (2H, m), 8.01 (1H, dd, J = 4.9, 1.5 Hz), 10.39 (1H, d, J = 7.2 Hz), 10.79 (1H, brs).

[0350]

Example 14

N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
20

[0351]

A) tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

1 M Methylmagnesium bromide/tetrahydrofuran solution
25 (2.20 mL) was diluted with tetrahydrofuran (5 mL), ethyl ((tert-butoxycarbonyl) amino)(4-(trifluoromethoxy)phenyl)acetate (200 mg) was added thereto at 0°C. The reaction mixture was stirred at the same temperature for 2 hr, saturated aqueous ammonium chloride solution was
30 added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to
35 give the title compound (78 mg).

MS (API-): [M-H]⁻ 348.2.

[0352]

B) N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step D of Example 13

MS (API+): [M+H]⁺ 425.0.

¹H NMR (300 MHz, CDCl₃) δ 1.16 (3H, s), 1.37 (3H, s), 1.70 (1H, s), 4.55-4.76 (2H, m), 4.91 (1H, d, J = 8.3 Hz), 7.02 (1H, dd, J = 7.9, 4.9 Hz), 7.13-7.22 (3H, m), 7.37-7.46 (2H, m), 8.07 (1H, dd, J = 4.9, 1.5 Hz), 8.51 (1H, s), 10.93 (1H, d, J = 8.3 Hz).

[0353]

Example 15

N-(cyclopropyl(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0354]

A) 1-cyclopropyl-N-hydroxy-1-(4-(trifluoromethoxy)phenyl)methanimine

The title compound was obtained in the same manner as in Step D of Example 2.

MS (API+): [M+H]⁺ 246.1.

[0355]

B) 1-cyclopropyl-1-(4-(trifluoromethoxy)phenyl)methanamine

To a solution of 1-cyclopropyl-N-hydroxy-1-(4-(trifluoromethoxy)phenyl)methanimine (4.2 g) in ethanol (100 mL) was added 10% palladium-carbon (containing 50% water, 1.823 g). The reaction mixture was stirred at room temperature for 1 hr under hydrogen atmosphere, 2 M hydrochloric acid/ethanol solution (1 mL) was added thereto, and the mixture was stirred overnight under hydrogen atmosphere. The insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. To the residue was added 4 M hydrogen chloride/ethyl acetate solution (45 mL), and the

solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether, and the precipitated solid was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)). To the obtained fraction was added
5 saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to give 1-cyclopropyl-1-(4-(trifluoromethoxy)phenyl)methanamine (0.924 g).

10 MS (API+), found: 215.2.

[0356]

C) 1-cyclopropyl-1-(4-(trifluoromethoxy)phenyl)methanamine hydrochloride

The title compound was obtained in the same manner as in
15 Step F of Example 2.

MS (API+), found: 215.1.

[0357]

D) N-(cyclopropyl(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 407.0.

¹H NMR (300 MHz, CDCl₃) δ 0.35-0.55 (2H, m), 0.55-0.69 (2H, m), 1.14-1.31 (1H, m), 4.43-4.55 (1H, m), 4.59-4.77 (2H, m), 7.00
25 (1H, dd, J = 7.9, 4.9 Hz), 7.11-7.22 (3H, m), 7.38-7.48 (2H, m), 8.01 (1H, dd, J = 5.1, 1.7 Hz), 8.38 (1H, brs), 10.44 (1H, d, J = 7.5 Hz).

[0358]

Example 16

30 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)butyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0359]

A) 1-(4-(trifluoromethoxy)phenyl)butan-1-amine

To a solution of 1-butyl-N-hydroxy-1-(4-(trifluoromethoxy)phenyl)methanimine (4.2 g) in ethanol (100
35

mL) was added 10% palladium-carbon (containing 50% water, 1.823 g). The reaction mixture was stirred at room temperature for 1 hr under hydrogen atmosphere, 2 M hydrogen chloride/ethanol solution (1 mL) was added thereto, and the mixture was stirred overnight under hydrogen atmosphere. The insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. To the residue was added 4 M hydrogen chloride/ethyl acetate solution (45 mL), and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether, and the precipitated solid was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)). To the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and concentrated under reduced pressure to give 1-(4-(trifluoromethoxy)phenyl)butan-1-amine (0.524 g).

MS (API+), found: 217.2.

[0360]

B) 1-(4-(trifluoromethoxy)phenyl)butan-1-amine hydrochloride

The title compound was obtained in the same manner as in Step F of Example 2.

MS (API+), found: 233.2.

[0361]

C) 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)butyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 409.0.

¹H NMR (300 MHz, CDCl₃) δ 0.86-1.01 (3H, m), 1.23-1.51 (2H, m), 1.68-1.94 (2H, m), 4.58-4.75 (2H, m), 4.96 (1H, q, J = 7.5 Hz), 7.00 (1H, dd, J = 7.7, 5.1 Hz), 7.10-7.22 (3H, m), 7.31-7.38 (2H, m), 8.00 (1H, dd, J = 4.9, 1.5 Hz), 8.99 (1H, s), 10.40 (1H, d, J = 7.5 Hz).

[0362]

Example 17

1-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0363]

5 A) 1-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of 3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.40 g) in N,N-dimethylformamide (15 mL) was slowly added 1 M potassium 1,1,1,3,3,3-hexamethyldisilazane/tetrahydrofuran solution (6.6 mL) under ice-cooling. The reaction mixture was
10 stirred at 0°C for 30 min, and methyl iodide (2.00 g) was slowly added thereto at the same temperature. The reaction mixture was stirred overnight at room temperature, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl
15 acetate. The extract was washed successively with water and saturated brine, and the solvent was evaporated under reduced pressure to give the title compound (1.10 g).

¹H NMR (400 MHz, CDCl₃) δ 3.33 (3H, s), 4.18 (2H, s), 6.73 (1H, dd, J = 8.0, 5.2 Hz), 7.06 (1H, d, J = 7.2 Hz), 7.78 (1H, d, J
20 = 4.4 Hz).

[0364]

B) 4-nitrophenyl 1-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

To a solution of 1-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (0.40 g) in a mixed solvent of N,N-dimethylacetamide (2 mL) and pyridine (1 mL) was added 4-nitrophenyl chloroformate (0.75 g). The reaction mixture was
25 stirred at 45°C for 30 min, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound
30 (0.30 g).

¹H NMR (300 MHz, CDCl₃) δ 3.43 (3H, s), 4.67 (2H, s), 7.28 (1H, dd, J = 8.1, 4.8 Hz), 7.42-7.47 (3H, m), 8.26-8.33 (3H, m).

[0365]

C) 1-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
5 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

¹H NMR (300 MHz, CDCl₃) δ 0.97 (3H, t, J = 7.5 Hz) 1.83-1.93 (2H, m), 3.36 (3H, s), 4.69 (2H, s), 4.91 (1H, q, J = 6.9 Hz),
10 7.09 (1H, dd, J = 7.8, 5.1 Hz), 7.18 (2H, d, J = 8.4 Hz), 7.27-7.37 (3H, m), 8.02 (1H, d, J = 4.8 Hz), 10.31 (1H, d, J = 7.5 Hz).

[0366]

Example 18

15 N-(1-(4-(difluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0367]

A) 4-(N-hydroxypropanimidoyl)phenol

The title compound was obtained in the same manner as in
20 Step C of Example 1.

MS (API+): [M+H]⁺ 166.1.

[0368]

B) 4-(1-aminopropyl)phenol

The title compound was obtained in the same manner as in
25 Step D of Example 1.

¹H NMR (300 MHz, DMSO-d₆) δ 0.74 (3H, t, J = 7.5 Hz), 1.36-1.62 (2H, m), 3.56 (1H, t, J = 6.6 Hz), 6.59-6.72 (2H, m), 7.02-7.14 (2H, m).

[0369]

30 C) tert-butyl (1-(4-hydroxyphenyl)propyl)carbamate

To a solution of 4-(1-aminopropyl)phenol (10.4 g) in tetrahydrofuran (200 mL) were added triethylamine (11.5 mL) and di-tert-butyl dicarbonate (19.1 mL). The reaction mixture was stirred at room temperature for 2 hr, water was added
35 thereto, and the mixture was extracted with ethyl acetate. The

extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The obtained solid was triturated with hexane/ethyl acetate, and washed with hexane/ethyl acetate to
5 give the title compound (13.8 g).

MS (API-): $[M-H]^-$ 250.2.

[0370]

D) tert-butyl (1-(4-(difluoromethoxy)phenyl)propyl)carbamate

Under chlorodifluoromethane atmosphere, to a solution of
10 tert-butyl (1-(4-hydroxyphenyl)propyl)carbamate (833 mg) and benzyltriethylammonium chloride (226 mg) in tetrahydrofuran (20 mL) was slowly added 8 M aqueous sodium hydroxide solution under ice-cooling. The reaction mixture was stirred at room temperature for 2 hr, water was added thereto, and the mixture
15 was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (832 mg).

20 ^1H NMR (300 MHz, DMSO- d_6) δ 0.81 (3H, t, J = 7.3 Hz), 1.35 (9H, s), 1.49-1.71 (2H, m), 4.26-4.45 (1H, m), 6.90-7.47 (6H, m).

[0371]

E) tert-butyl 1-(4-(difluoromethoxy)phenyl)propan-1-amine hydrochloride

25 To a solution of (1-(4-(difluoromethoxy)phenyl)propyl)carbamate (830 mg) in ethyl acetate (5 mL) was added 4 M hydrogen chloride/ethyl acetate (10 mL) solution. The reaction mixture was stirred at room temperature for 3 hr, and the solvent was evaporated under
30 reduced pressure to give the title compound (643 mg).

^1H NMR (300 MHz, DMSO- d_6) δ 0.75 (3H, t, J = 7.3 Hz), 1.67-2.10 (2H, m), 4.06-4.24 (1H, m), 6.99-7.62 (5H, m), 8.51 (3H, brs).

[0372]

F) N-(1-(4-(difluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
35

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): $[M+H]^+$ 377.1.

^1H NMR (300 MHz, DMSO-d_6) δ 0.86 (3H, t, $J = 7.3$ Hz), 1.67-1.91 (2H, m), 4.41 (2H, s), 4.77 (1H, q, $J = 7.0$ Hz), 6.91-7.50 (7H, m), 7.94-8.10 (1H, m), 10.30 (1H, d, $J = 7.5$ Hz), 10.82 (1H, brs).

[0373]

Example 19

10 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0374]

A) methyl N-(3-nitro-5-(trifluoromethyl)pyridin-2-yl)glycinate

15 To a solution of methyl glycinate hydrochloride (13.9 g) and 2-chloro-3-nitro-5-(trifluoromethyl)pyridine (5.0 g) in ethanol (100 mL) was added triethylamine (15.4 mL). The reaction mixture was heated at reflux for 5 hr, and the solvent was evaporated under reduced pressure. The residue was
20 triturated with water, and the solid was collected by filtration, and washed with ethanol to give the title compound (5.09 g).

^1H NMR (400 MHz, CDCl_3) δ 3.82 (3H, s), 4.43 (2H, d, $J = 5.2$ Hz), 8.63 (1H, d, $J = 2.0$ Hz), 8.68 (1H, d, $J = 2.4$ Hz), 8.74
25 (1H, s).

[0375]

B) 7-(trifluoromethyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(3-nitro-5-(trifluoromethyl)pyridin-2-yl)glycinate (5.09 g) in ethanol
30 (200 mL) was added tin(II) chloride dihydrate (20.6 g). The reaction mixture was stirred at 80°C for 2 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate/ethanol, the pH of the solution was
35 adjusted to 9 with saturated aqueous ammonia solution, and the

insoluble substance was filtered off using Celite. The filtrate was concentrated under reduced pressure to give the title compound (1.10 g).

¹H NMR (400 MHz, DMSO-d₆) δ 4.04 (2H, s), 7.05 (1H, d, J = 1.6 Hz), 7.61 (1H, s), 7.93 (1H, s), 10.61 (1H, brs).

[0376]

C) 4-nitrophenyl 2-oxo-7-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

To a solution of 7-(trifluoromethyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (300 mg) in a mixed solvent of N,N-dimethylacetamide (2 mL) and pyridine (0.5 mL) was added 4-nitrophenyl chloroformate (418 mg). The reaction mixture was stirred overnight at 45°C, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (220 mg).

¹H NMR (400 MHz, CDCl₃) δ 4.63 (2H, s), 7.45 (2H, d, J = 8.8 Hz), 7.57 (1H, s), 8.31 (2H, d, J = 9.2 Hz), 8.44 (1H, s).

[0377]

D) 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 4-nitrophenyl 2-oxo-7-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (220 mg) in N,N-dimethylformamide (3 mL) were added 1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride (191 mg) and triethylamine (0.3 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give a crude product. The crude product was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate.

The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (72 mg).

MS (API+): [M+H]⁺ 463.2.

5 ¹H NMR (400 MHz, CDCl₃) δ 0.97 (3H, t, J = 4.0 Hz) 1.84-1.92 (2H, m), 4.71 (2H, s), 4.91 (1H, q, J = 6.8 Hz), 7.17 (1H, d, J = 8.0 Hz), 7.33-7.36 (3H, m), 8.28 (1H, s), 9.62 (1H, s), 10.33 (1H, d, J = 7.2 Hz).

[0378]

10 Example 20

2-oxo-7-phenyl-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (140 mg), phenylboronic acid (49 mg) and tetrakis(triphenylphosphine)palladium(0) (37 mg) in 1,2-dimethoxyethane (4 mL) was added 2 M aqueous sodium carbonate solution (2 mL) at room temperature. The reaction mixture was stirred overnight at reflux, and extracted with 20 ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by preparative TLC (petroleum ether/ethyl acetate) to give the title compound (72 mg).

25 MS (API+): [M+H]⁺ 471.3.

¹H NMR (400 MHz, CDCl₃) δ 0.98 (3H, t, J = 7.2 Hz), 1.87-1.93 (2H, m), 4.71 (1H, d, J = 1.2 Hz), 4.92 (1H, q, J = 6.8 Hz), 7.18 (2H, d, J = 7.6 Hz), 7.36-7.38 (3H, m), 7.41-7.45 (1H, m), 7.47-7.51 (2H, m), 7.55-7.57 (2H, m), 8.24 (1H, d, J = 2.0 Hz), 30 8.79 (1H, s), 10.41 (1H, d, J = 7.2 Hz).

[0379]

Example 21

6-bromo-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide

35 [0380]

A) ethyl N-(3,5-dibromopyrazin-2-yl)glycinate

To a solution of 3,5-dibromopyrazin-2-amine (16.0 g) and cesium carbonate (26.7 g) in N,N-dimethylformamide (90 mL) was slowly added 2-chloroethyl acetate (8.1 mL) at room
5 temperature. The reaction mixture was stirred overnight at 65°C under nitrogen atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated
10 under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (8.6 g).

¹H NMR (400 MHz, CDCl₃) δ 1.31 (3H, t, J = 7.2 Hz), 4.16 (2H, d, J = 5.6 Hz), 4.25 (2H, q, J = 7.2 Hz), 5.75 (1H, brs), 8.04
15 (1H, s).

[0381]

B) 7-bromo-1-(2,4-dimethoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one

To a solution of ethyl N-(3,5-dibromopyrazin-2-yl)glycinate (4.20 g) and 2,4-dimethoxybenzyl amine (2.20 g)
20 in dimethyl sulfoxide (5 mL) was added diisopropylethylamine (5 mL) at room temperature. The reaction mixture was stirred at 135°C for 2.5 hr using microwave, and cooled to room temperature, water was added thereto, and the mixture was
25 extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from methanol to give the title compound (1.2 g).

30 ¹H NMR (300 MHz, CDCl₃) δ 3.77 (3H, s), 3.81 (3H, s), 4.29 (2H, s), 4.92 (1H, s), 5.22 (2H, s), 6.38-6.42 (2H, m), 7.10 (1H, d, J = 8.1 Hz), 7.67 (1H, s).

[0382]

C) 6-bromo-4-(2,4-dimethoxybenzyl)-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-

35

b]pyrazine-1(2H)-carboxamide

Under nitrogen atmosphere, to a solution of 7-bromo-1-(2,4-dimethoxybenzyl)-3,4-dihydropyrazino[2,3-b]pyrazin-2(1H)-one (1.7 g) in tetrahydrofuran (45 mL) was slowly added
5 triphosgene (0.81 g) at room temperature. The reaction mixture was stirred at 50°C for 2.5 hr, and the solvent was evaporated under reduced pressure. The residue was diluted with tetrahydrofuran (30 mL), and the mixture was slowly added to a solution of 1-(4-(trifluoromethoxy)phenyl)propan-1-amine
10 hydrochloride (1.26 g) and triethylamine (2.15 mL) in tetrahydrofuran (45 mL) at room temperature. The reaction mixture was stirred at 60°C for 30 min, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl
15 acetate) to give the title compound (2.5 g).

¹H NMR (400 MHz, CDCl₃) δ 0.93 (3H, t, J = 7.2 Hz), 1.79-1.91 (2H, m), 3.77 (3H, s), 3.79 (3H, s), 4.72 (2H, dd, J = 18.8, 14.8 Hz), 4.86 (1H, q, J = 7.2 Hz), 5.23 (2H, dd, J = 16.8, 14.4 Hz), 6.38-6.41 (2H, m), 7.15-7.20 (3H, m), 7.27-7.33 (2H,
20 m), 7.88 (1H, s), 9.69 (1H, d, J = 7.2 Hz).

[0383]

D) 6-bromo-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide

To 6-bromo-4-(2,4-dimethoxybenzyl)-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide (300 mg) was added
25 trifluoroacetic acid (3 mL) at room temperature. The reaction mixture was stirred at 75°C for 1 hr under nitrogen atmosphere, water was added thereto. The pH of the mixture was adjusted to 10 with 2 M aqueous sodium hydroxide solution, and the mixture
30 was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
35 column chromatography (petroleum ether/ethyl acetate) to give

the title compound (40 mg).

MS (API-): [M-H]⁻ 471.9.

¹H NMR (400 MHz, CDCl₃) δ 0.94 (3H, t, J = 7.2 Hz), 1.83-1.91 (2H, m), 4.71 (2H, s), 4.87 (1H, q, J = 7.2 Hz), 7.17 (2H, d, J = 8.4 Hz), 7.31 (2H, d, J = 8.4 Hz), 8.00 (1H, s), 8.42 (1H, s), 9.70 (1H, d, J = 6.8 Hz).

[0384]

Example 22

3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide
[0385]

A) 4-(2,4-dimethoxybenzyl)-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide
To a solution of 6-bromo-4-(2,4-dimethoxybenzyl)-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide (800 mg) in a mixed solvent of tetrahydrofuran (10 mL) and methanol (10 mL) was added 10% palladium-carbon (containing 50% water, 120 mg). The reaction mixture was stirred overnight at 30°C under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (610 mg).

¹H NMR (400 MHz, CDCl₃) δ 0.95 (3H, t, J = 6.4 Hz), 1.85-1.91 (2H, m), 3.76 (3H, s), 3.79 (3H, s), 4.77 (2H, s), 4.90 (1H, q, J = 7.2 Hz), 5.27 (2H, s), 6.35 (1H, dd, J = 11.2, 3.2 Hz), 6.43 (1H, d, J = 3.2 Hz), 6.94 (1H, d, J = 11.2 Hz), 7.17 (2H, d, J = 10.8 Hz), 7.34 (2H, d, J = 10.8 Hz), 7.81 (1H, d, J = 3.6 Hz), 7.93 (1H, d, J = 3.6 Hz), 10.12 (1H, d, J = 10.0 Hz).
[0386]

B) 3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide

The title compound was obtained in the same manner as in Step D of Example 21.

MS (API-): [M-H]⁻ 394.0.

¹H NMR (400 MHz, CDCl₃) δ 0.96 (3H, t, J = 7.6 Hz), 1.84-1.93 (2H, m), 4.72 (2H, s), 4.89 (1H, q, J = 7.6 Hz), 7.17 (2H, d, J = 8.8 Hz), 7.33 (2H, d, J = 8.8 Hz), 7.91 (1H, d, J = 2.8 Hz), 7.94 (1H, d, J = 2.8 Hz), 9.01 (1H, s), 10.09 (1H, d, J = 7.2 Hz).

[0387]

Example 23

6-methyl-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide

[0388]

A) 4-(2,4-dimethoxybenzyl)-6-methyl-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide

Under nitrogen atmosphere, to a solution of 6-bromo-4-(2,4-dimethoxybenzyl)-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide (1.0 g) and (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (150 mg) in 1,4-dioxane (15 mL) was added 1 M dimethylzinc(II)/hexane solution (4 mL) at room temperature. The reaction mixture was stirred at 80°C for 30 min under nitrogen atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by preparative TLC (petroleum ether/ethyl acetate) to give the title compound (730 mg).

¹H NMR (400 MHz, CDCl₃) δ 0.93 (3H, t, J = 7.2 Hz), 1.83-1.88 (2H, m), 2.39 (3H, s), 3.76 (3H, s), 3.79 (3H, s), 4.73 (2H, s), 4.89 (1H, q, J = 7.2 Hz), 5.27 (2H, s), 6.36 (1H, dd, J = 8.0, 2.4 Hz), 6.41 (1H, d, J = 2.4 Hz), 7.03 (1H, d, J = 8.8 Hz), 7.16 (2H, d, J = 8.0 Hz), 7.32 (2H, d, J = 8.0 Hz), 7.66 (1H, s), 10.0 (1H, d, J = 7.2 Hz).

[0389]

B) 6-methyl-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide

To 4-(2,4-dimethoxybenzyl)-6-methyl-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide (730 mg) was added trifluoroacetic acid (10 mL) at room temperature. The reaction mixture was stirred at 70°C for 15 min, and water was added thereto. The pH of the mixture was adjusted to 10 or more with 4 M aqueous sodium hydroxide solution, and the mixture was extracted with dichloromethane. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (150 mg).

MS (API+): [M+H]⁺ 410.2.

¹H NMR (400 MHz, CDCl₃) δ 0.94 (3H, t, J = 7.2 Hz), 1.82-1.90 (2H, m), 2.45 (3H, s), 4.70 (2H, s), 4.88 (1H, q, J = 7.2 Hz), 7.17 (2H, d, J = 8.4 Hz), 7.32 (2H, d, J = 8.4 Hz), 7.77 (1H, s), 8.56 (1H, s), 9.98 (1H, d, J = 6.8 Hz).

[0390]

Example 24

N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (548 mg) was resolved by HPLC (column: CHIRALPAK IC, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 400/600) to give the title compound having a shorter retention time (265 mg).

MS (API+): [M+H]⁺ 411.2.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.54-3.73 (2H, m), 4.31-4.53 (2H, m), 4.97-5.14 (1H, m), 7.11 (1H, dd, J = 7.9, 4.9 Hz), 7.25-7.39 (3H, m), 7.40-7.54 (2H, m), 8.00 (1H, dd, J = 4.9, 1.5 Hz), 10.51 (1H, d, J = 7.5 Hz), 10.83 (1H, brs).

5 [0391]

Example 25

N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (548 mg) was resolved by HPLC (column: CHIRALPAK IC, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 400/600) to give the title compound having a longer retention
15 time (273 mg).

MS (API+): [M+H]⁺ 411.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.55-3.71 (2H, m), 4.33-4.51 (2H, m), 4.99-5.14 (1H, m), 7.11 (1H, dd, J = 7.7, 5.1 Hz), 7.26-7.37 (3H, m), 7.42-7.53 (2H, m), 8.00 (1H, dd, J
20 = 4.9, 1.9 Hz), 10.51 (1H, d, J = 7.5 Hz), 10.83 (1H, brs).

[0392]

Example 26

7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

25 [0393]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

The title compound was obtained in the same manner as in Step A of Example 19.

¹H NMR (400 MHz, CDCl₃) δ 3.81 (3H, s), 4.37 (2H, d, J = 7.6 Hz), 8.45 (1H, s), 8.54 (1H, d, J = 2.8 Hz), 8.71 (1H, d, J = 2.8 Hz).

[0394]

B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

The title compound was obtained in the same manner as in.
35 Step B of Example 19

¹H NMR (400 MHz, DMSO-d₆) δ 4.00 (2H, s), 7.17 (1H, d, J = 1.6 Hz), 7.50 (1H, brs), 7.77 (1H, d, J = 1.6 Hz), 10.66 (1H, s).
[0395]

C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-
5 dihydropyrido[2,3-b]pyrazin-2(1H)-one

Under nitrogen atmosphere, to a solution of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (12.5 g) in a mixed solvent of N,N-dimethylformamide (200 mL) and dimethyl sulfoxide (600 mL) was slowly added 1 M potassium 1,1,1,3,3,3-hexamethyldisilazane/tetrahydrofuran solution (63.6 mL) under
10 ice-cooling. The reaction mixture was stirred at 0°C for 30 min, and [2-(chloromethoxy)ethyl](trimethyl)silane (11.3 mL) was slowly added thereto at the same temperature. The reaction mixture was stirred at room temperature for 2 hr, and the
15 solvent was evaporated under reduced pressure. To the residue was added aqueous sodium carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and the solvent was evaporated under reduced pressure. The residue was
20 purified by silica gel column chromatography (dichloromethane/ethyl acetate) to give the title compound (6.86 g).

¹H NMR (400 MHz, DMSO-d₆) δ 0.01 (9H, s), 0.90 (2H, t, J = 7.6 Hz), 3.59 (2H, t, J = 8.0 Hz), 4.07 (2H, d, J = 1.6 Hz), 5.31
25 (2H, s), 7.55 (1H, d, J = 1.6 Hz), 7.92 (1H, d, J = 2.0 Hz).
[0396]

D) 7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-1-
((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

30 Under nitrogen atmosphere, to a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (4.05 g) in tetrahydrofuran (150 mL) was slowly added a solution of triphosgene (2.37 g) in tetrahydrofuran (15 mL) at room temperature. The reaction
35 mixture was stirred at 40°C for 2 hr, and the solvent was

evaporated under reduced pressure. The residue was diluted with tetrahydrofuran (40 mL), and the mixture was slowly added to a solution of 1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride (3.07 g) and triethylamine (3.03 g) in
5 tetrahydrofuran (50 mL) at room temperature. The reaction mixture was stirred overnight at 60°C, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (4.08 g).
10 ¹H NMR (400 MHz, CDCl₃) δ 0.01 (9H, s), 0.93 (3H, t, J = 9.2 Hz), 1.80-1.86 (2H, m), 3.65 (1H, t, J = 8.0 Hz), 4.66 (2H, dd, J = 17.6, 7.6 Hz), 4.87 (1H, q, J = 6.8 Hz), 5.27 (2H, dd, J = 16.8, 10.8 Hz), 7.16 (2H, d, J = 8.0 Hz), 7.31 (2H, d, J = 8.8 Hz), 7.95 (1H, d, J = 1.6 Hz), 8.24 (1H, d, J = 1.6 Hz), 9.86
15 (1H, d, J = 7.2 Hz).

[0397]

E) 7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To 7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
20 1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (4.08 g) were added trifluoroacetic acid (18 mL) and water (2 mL) at room temperature. The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under
25 reduced pressure. To the residue was added aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was
30 purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (2.64 g).

MS (API⁺): [M+H]⁺ 521.1.

¹H NMR (400 MHz, CDCl₃) δ 0.95 (3H, t, J = 7.2 Hz), 1.81-1.87 (2H, m), 4.67 (1H, d, J = 2.0 Hz), 4.87 (1H, q, J = 6.8 Hz),
35 7.17 (2H, d, J = 8.0 Hz), 7.32 (2H, d, J = 8.4 Hz), 7.44 (1H,

d, J = 2.0 Hz), 8.19 (1H, d, J = 2.0 Hz), 9.20 (1H, s), 10.07 (1H, d, J = 7.2 Hz).

[0398]

Example 27

5 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-vinyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (520 mg), 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (231 mg) and
10 tetrakis(triphenylphosphine) palladium(0) (116 mg) in 1,2-dimethoxyethane (12 mL) was added 2 M aqueous sodium carbonate solution (4 mL) at room temperature. The reaction mixture was stirred overnight at reflux, and extracted with ethyl acetate.
15 The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (372 mg).

20 MS (API+): [M+H]⁺ 421.1.

¹H NMR (400 MHz, CDCl₃) δ 0.96 (3H, t, J = 7.2 Hz), 1.83-1.90 (2H, m), 4.80 (1H, d, J = 0.8 Hz), 4.90 (1H, q, J = 6.8 Hz), 5.38 (1H, d, J = 10.8 Hz), 5.76 (1H, d, J = 17.6 Hz), 6.65 (1H, dd, J = 18.0, 10.8 Hz), 7.17 (2H, d, J = 7.6 Hz), 7.24 (1H, d, J = 1.6 Hz), 7.34 (2H, d, J = 8.8 Hz), 8.00 (1H, d, J = 1.6 Hz), 9.12 (1H, s), 10.38 (1H, d, J = 7.2 Hz).

[0399]

Example 28

N-(1-(biphenyl-4-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
30

[0400]

A) 4-(1-(((2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-yl)carbonyl) amino)propyl)phenyl trifluoromethanesulfonate

To a solution of N-(1-(4-hydroxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.00 g) in
35

pyridine (30 mL) was slowly added trifluoromethanesulfonic anhydride (0.621 mL) under ice-cooling. The reaction mixture was stirred at room temperature for 3 hr, and the solvent was evaporated under reduced pressure. To the residue was added
5 water, the mixture was washed with 0.5 M hydrochloric acid, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.34 g).

10 MS (API+): [M+H]⁺ 459.1.

[0401]

B) N-(1-(biphenyl-4-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 4-(1-(((2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-yl)carbonyl) amino)propyl)phenyl
15 trifluoromethanesulfonate (125 mg) in a mixed solvent of water (1.5 mL) and 1,2-dimethoxyethane (3 mL) were added phenylboronic acid (36.5 mg), potassium carbonate (150 mg) and tetrakis(triphenylphosphine) palladium(0) (15.7 mg) at room
20 temperature. The reaction mixture was stirred at 150°C for 20 min under nitrogen atmosphere using microwave, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was
25 purified by silica gel column chromatography (hexane/ethyl acetate) to give a crude product. The crude product was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the
30 mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (29.8 mg).

MS (API+): [M+H]⁺ 387.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.90 (3H, t, J = 7.3 Hz), 1.73-1.94
35 (2H, m), 4.43 (2H, s), 4.82 (1H, q, J = 6.9 Hz), 7.11 (1H, dd,

J = 7.5, 4.9 Hz), 7.26-7.52 (6H, m), 7.56-7.71 (4H, m), 8.03 (1H, dd, J = 4.9, 1.5 Hz), 10.36 (1H, d, J = 7.9 Hz), 10.82 (1H, brs).

[0402]

5 Example 29

7-ethyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-vinyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (177 mg) in methanol (25 mL) was added 10% palladium-carbon (containing 50% water, 18 mg) under nitrogen atmosphere. The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (135 mg).

MS (API+): [M+H]⁺ 423.1.

¹H NMR (400 MHz, CDCl₃) δ 0.95 (3H, t, J = 7.2 Hz), 1.26 (3H, t, J = 7.6 Hz), 1.82-1.89 (2H, m), 2.63 (2H, q, J = 7.6 Hz), 4.66 (2H, dd, J = 20.0, 18.4 Hz), 4.90 (1H, q, J = 7.2 Hz), 7.04 (1H, d, J = 1.6 Hz), 7.16 (1H, d, J = 8.4 Hz), 7.34 (2H, d, J = 8.8 Hz), 7.85 (1H, d, J = 1.6 Hz), 9.32 (1H, brs), 10.38 (1H, d, J = 7.2 Hz).

[0403]

30 Example 30

N-(1-(6-methoxypyridazin-3-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0404]

A) 6-methoxypyridazine-3-carbonitrile

35 To a solution of 3-chloro-6-methoxypyridazine (5 g) and

zinc cyanide (4.87 g) in N,N-dimethylformamide (100 mL) were added 1,1'-bis(diphenylphosphino)ferrocene (1.917 g) and tris(dibenzylideneacetone)dipalladium(0) (1.584 g). The reaction mixture was stirred at 100°C for 2 days under nitrogen
5 atmosphere, saturated brine was added thereto, and the mixture was extracted with ethyl acetate and tetrahydrofuran. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
10 column chromatography (ethyl acetate/hexane) to give the title compound (3.20 g).

¹H NMR (300 MHz, CDCl₃) δ 4.19-4.28 (3H, m), 7.03-7.15 (1H, m), 7.64-7.73 (1H, m).

[0405]

15 B) 1-(6-methoxypyridazin-3-yl)propan-1-one

To a solution of 6-methoxypyridazine-3-carbonitrile (1.9 g) in tetrahydrofuran (30 mL) was added 3 M ethylmagnesium bromide/diethyl ether solution (5.62 mL) under ice-cooling. The reaction mixture was stirred at 0°C for 1 hr under nitrogen
20 atmosphere, 1 M hydrochloric acid was added thereto at the same temperature, and the mixture was stirred at room temperature for further 5 min. The reaction mixture was neutralized with saturated aqueous sodium hydrogen carbonate solution, and extracted with ethyl acetate. The extract was
25 washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.323 g).

30 ¹H NMR (300 MHz, CDCl₃) δ 1.25 (3H, t, J = 7.2 Hz), 3.35 (2H, q, J = 7.4 Hz), 4.23 (3H, s), 7.06 (1H, d, J = 9.4 Hz), 8.05 (1H, d, J = 9.0 Hz).

[0406]

C) 1-(6-methoxypyridazin-3-yl)propan-1-amine

35 The title compound was obtained in the same manner as in

Steps D-E of Example 2.

MS (API+): [M+H]⁺ 168.2.

D) N-(1-(6-methoxypyridazin-3-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 343.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.86 (3H, t, J = 7.3 Hz), 1.70-1.99 (2H, m), 4.01 (3H, s), 4.43 (2H, s), 5.01 (1H, q, J = 7.2 Hz),
10 7.11 (1H, dd, J = 7.9, 4.9 Hz), 7.18 (1H, d, J = 9.0 Hz), 7.32 (1H, dd, J = 7.9, 1.5 Hz), 7.62 (1H, d, J = 9.0 Hz), 8.02 (1H, dd, J = 4.9, 1.5 Hz), 10.42 (1H, d, J = 7.5 Hz), 10.81 (1H, s).
[0407]

Example 31

15 N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0408]

A) 3-fluoro-N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide

To a solution of 3-fluoro-4-(trifluoromethoxy)benzoic
20 acid (20 g), N,O-dimethylhydroxylamine hydrochloride (10.5 g) and triethylamine (24.9 mL) in N,N-dimethylformamide (300 mL) were added 1-hydroxybenzotriazole monohydrate (16.4 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (20.5 g) at room temperature. The reaction mixture was stirred
25 overnight at room temperature, saturated aqueous sodium hydrogen carbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give
30 the title compound (22.9 g).

¹H NMR (300 MHz, CDCl₃) δ 3.38 (3H, s), 3.56 (3H, s), 7.30-7.42 (1H, m), 7.51-7.65 (2H, m).

[0409]

B) 1-(3-fluoro-4-(trifluoromethoxy)phenyl)propan-1-amine
35 hydrochloride

The title compound was obtained in the same manner as in Steps C-F of Example 2.

MS (API+), found: 221.1.

C) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-
5 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 413.1.

¹H NMR (300 MHz, DMSO-d₆) δ 0.77-0.95 (3H, m), 1.81 (2H, quin,
10 J = 7.3 Hz), 4.35-4.47 (2H, m), 4.81 (1H, q, J = 7.2 Hz), 7.11
(1H, dd, J = 7.7, 5.1 Hz), 7.27 (1H, d, J = 8.3 Hz), 7.32 (1H,
dd, J = 7.9, 1.5 Hz), 7.44-7.58 (2H, m), 8.02 (1H, dd, J = 4.9,
1.5 Hz), 10.30 (1H, d, J = 7.2 Hz), 10.81 (1H, s).

[0410]

15 Example 32

N-(1-(2-methoxy-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in
Step A of Example 31, Steps C-F of Example 2 and Step G of
20 Example 7.

MS (API+): [M+H]⁺ 425.1.

¹H NMR (300 MHz, CDCl₃) δ 0.92 (3H, t, J = 7.5 Hz), 1.71-1.95
(2H, m), 3.88 (3H, s), 4.68 (2H, s), 5.05-5.26 (1H, m), 6.71
(1H, s), 6.77 (1H, d, J = 9.4 Hz), 6.94-7.03 (1H, m), 7.13 (1H,
25 brs), 7.22 (1H, d, J = 8.3 Hz), 8.02 (1H, d, J = 4.9 Hz),
10.43 (1H, brs).

[0411]

Example 33

N-(1-(4-hydroxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
30 b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in
Step G of Example 7.

MS (API+): [M+H]⁺ 327.1.

[0412]

35 Example 34

N-(1-(2-fluoro-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Steps A-B of Example 30, Steps D-E of Example 2 and Step G of
5 Example 7.

MS (API+): [M+H]⁺ 413.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.89 (3H, t, J = 7.2 Hz), 1.70-1.88 (2H, m), 4.40 (2H, s), 5.00 (1H, q, J = 7.4 Hz), 7.11 (1H, dd, J = 7.9, 4.9 Hz), 7.22 (1H, d, J = 8.7 Hz), 7.27-7.42 (2H, m),
10 7.49 (1H, t, J = 8.5 Hz), 8.02 (1H, dd, J = 4.9, 1.9 Hz), 10.43 (1H, d, J = 7.5 Hz), 10.81 (1H, s).

[0413]

Example 35

7-cyclopropyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
15 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-methyl-N-nitrosourea (577 mg) in diethyl ether (10 mL) was slowly added 3 M potassium hydroxide solution (6 mL) under ice-cooling. The reaction mixture was stirred at the same temperature for 30 min, and the organic
20 layer was dried over anhydrous sodium sulfate, and slowly added to a solution of 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-vinyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (118 mg) in dichloromethane (10 mL) under ice-cooling. Then, palladium(II) acetate (25 mg) was added thereto, the reaction mixture was
25 stirred at room temperature for 1 hr, and some drops of acetic acid was added thereto. The insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by HPLC (C18, mobile phase:
30 water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound
35 (43 mg).

MS (API+): [M+H]⁺ 435.1.

¹H NMR (400 MHz, CDCl₃) δ 0.69-0.73 (2H, m), 0.95 (3H, t, J = 7.2 Hz), 1.00-1.05 (2H, m), 1.82-1.90 (3H, m), 4.60-4.71 (2H, m), 4.89 (1H, q, J = 6.8 Hz), 6.85 (1H, d, J = 2.0 Hz), 7.16
5 (1H, d, J = 7.6 Hz), 7.34 (2H, d, J = 8.4 Hz), 7.81 (1H, d, J = 2.0 Hz), 9.31 (1H, s), 10.30 (1H, d, J = 7.2 Hz).

[0414]

Example 36

7-isopropyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-
10 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0415]

A) 2-oxo-7-(prop-1-en-2-yl)-N-(1-[4-(trifluoromethoxy)phenyl]propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 The title compound was obtained in the same manner as in Example 27.

¹H NMR (400 MHz, CDCl₃) δ 0.97 (3H, t, J = 3.9 Hz), 1.84-1.93 (2H, m), 2.17 (3H, s), 4.69 (1H, d, J = 3.9 Hz), 4.92 (1H, q, J = 6.9 Hz), 5.19 (1H, s), 5.41 (1H, s), 7.18 (2H, d, J = 7.8
20 Hz), 7.24 (1H, d, J = 2.1 Hz), 7.34-7.38 (2H, m), 8.11 (1H, d, J = 2.1 Hz), 8.82 (1H, s), 10.41 (1H, d, J = 7.5 Hz).

[0416]

B) 7-isopropyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

25 The title compound was obtained in the same manner as in Example 29.

MS (API+): [M+H]⁺ 437.2.

[0417]

Example 37

30 6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0418]

A) methyl N-(6-chloro-3-nitropyridin-2-yl)glycinate

To a solution of methyl glycinate hydrochloride (15.0 g)
35 and 2,6-dichloro-3-nitropyridine (21.0 g) in ethanol (100 mL)

was added triethylamine (30.9 mL). The reaction mixture was heated at reflux overnight, and the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water
5 and saturated brine, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (16.3 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.67 (3H, s), 4.26 (2H, d, J = 5.6
10 Hz), 6.87 (1H, d, J = 8.8 Hz), 8.48 (1H, d, J = 8.8 Hz), 8.98 (1H, t, J = 5.2 Hz).

[0419]

B) 6-chloro-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(6-chloro-3-nitropyridin-2-
15 yl)glycinate (13.3 g) in ethanol (650 mL) was added tin(II) chloride dihydrate (98.0 g). The reaction mixture was heated at reflux overnight, and the precipitated solid was collected by filtration, and dried under reduced pressure to give the title compound (9.4 g).

20 ¹H NMR (400 MHz, DMSO-d₆) δ 3.95 (2H, s), 6.56 (1H, d, J = 8.0 Hz), 6.91 (1H, d, J = 7.6 Hz), 7.22 (1H, s), 10.49 (1H, s).

[0420]

C) 6-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

25 The title compound was obtained in the same manner as in Step C of Example 26.

¹H NMR (400 MHz, DMSO-d₆) δ 0.01 (9H, s), 0.90 (2H, t, J = 8.4 Hz), 3.58 (2H, t, J = 8.0 Hz), 4.09 (2H, s), 5.29 (2H, s),
30 6.74 (1H, d, J = 7.6 Hz), 7.34 (1H, d, J = 8.0 Hz), 7.39 (1H, brs).

[0421]

D) 6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

35 Under nitrogen atmosphere, to a solution of 6-chloro-1-

((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (7.9 g) and triethylamine (7.63 g) in tetrahydrofuran (150 mL) was slowly added triphosgene (5.98 g) at room temperature. The reaction mixture was stirred at 45°C for 2 hr, 1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride (7.72 g) was added thereto, and the mixture was stirred at 60°C for further 12 hr. The reaction mixture was concentrated under reduced pressure, and the obtained residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (7.0 g).
¹H NMR (400 MHz, DMSO-d₆) δ 0.02 (9H, s), 0.90-1.01 (5H, m), 1.86-1.92 (2H, m), 3.65 (2H, t, J = 11.2 Hz), 4.68 (2H, t, J = 6.4 Hz), 4.88-4.90 (1H, m), 5.30 (2H, d, J = 4.0 Hz), 7.07 (1H, d, J = 11.2 Hz), 7.19 (2H, d, J = 10.8 Hz), 7.35-7.39 (2H, m), 7.67 (1H, d, J = 11.6 Hz), 9.80 (1H, d, J = 9.2 Hz).

[0422]

E) 6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step E of Example 26.

MS (API+): [M+H]⁺ 429.1.

¹H NMR (300 MHz, CDCl₃) δ 0.97 (3H, t, J = 7.2 Hz), 1.84-1.93 (2H, m), 4.64 (2H, d, J = 2.0 Hz), 4.85-4.90 (1H, m), 6.98 (1H, d, J = 7.6 Hz), 7.17-7.19 (3H, m), 7.36 (2H, d, J = 8.4 Hz), 9.81 (1H, s), 9.96 (1H, d, J = 7.2 Hz).

[0423]

Example 38

N-(1-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step B of Example 28.

MS (API+): [M+H]⁺ 391.2.

[0424]

Example 39

2-oxo-N-(1-(4-(prop-1-en-2-yl)phenyl)propyl)-2,3-

dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step B of Example 28.

MS (API+): [M+H]⁺ 351.2.

5 [0425]

Example 40

N-(1-(4-isopropylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 2-oxo-N-(1-(4-(prop-1-en-2-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (46 mg) in methanol (3 mL) was added palladium-carbon-ethylene diamine complex (15 mg). The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, and the insoluble substance was filtered off. The filtrate was concentrated under reduced pressure, the residue was dissolved in methanol (3 mL), and 10% palladium-carbon (containing 50% water, 15 mg) was added thereto. The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (22.0 mg).

MS (API+): [M+H]⁺ 353.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.85 (3H, t, J = 7.3 Hz), 1.18 (6H, d, J = 6.8 Hz), 1.68-1.85 (2H, m), 2.75-2.94 (1H, m), 4.41 (2H, s), 4.73 (1H, q, J = 7.0 Hz), 7.09 (1H, dd, J = 7.9, 4.9 Hz), 7.15-7.25 (4H, m), 7.31 (1H, dd, J = 7.9, 1.5 Hz), 8.01 (1H, dd, J = 4.9, 1.5 Hz), 10.29 (1H, d, J = 7.5 Hz), 10.81 (1H, brs).

35 [0426]

Example 41

N-(1-(4-(3,6-dihydro-2H-pyran-4-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in
5 Step B of Example 28.

MS (API+): [M+H]⁺ 393.2.

[0427]

Example 42

N-(2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-
10 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0428]

A) amino (4-(trifluoromethoxy)phenyl)acetonitrile
hydrochloride

4-(Trifluoromethoxy)benzaldehyde (10 g) was dissolved in
15 2 M ammonia/methanol solution (105 mL), and titanium(IV)
tetraisopropoxide (15.4 mL) was added thereto under ice-
cooling. The reaction mixture was stirred at the same
temperature for 10 min, trimethylsilanecarbonitrile (7.83 g)
was added thereto, and the mixture was stirred overnight at
20 room temperature. Then, saturated aqueous sodium hydrogen
carbonate solution was added thereto, the insoluble substance
was filtered off using Celite, and the filtrate was extracted
with ethyl acetate. The extract was washed successively with
water and saturated brine, and dried over anhydrous magnesium
25 sulfate, and the solvent was evaporated under reduced pressure.
To the residue was added 4 M hydrogen chloride/ethyl acetate
solution (20 mL) under ice-cooling, and the solvent was
evaporated under reduced pressure to give the title compound
(10.0 g).

30 MS (API+), found: 200.0.

¹H NMR (300 MHz, DMSO-d₆) δ 5.91-6.12 (1H, m), 7.47-7.62 (2H,
m), 7.74-7.95 (2H, m), 9.61 (3H, brs).

[0429]

B) 2-amino-2-(4-(trifluoromethoxy)phenyl)acetamide

35 To a mixture of amino (4-

(trifluoromethoxy)phenyl)acetonitrile hydrochloride (3 g), potassium carbonate (4.92 g) and dimethyl sulfoxide (30 mL) was added 35% aqueous hydrogen peroxide (3.12 mL) under ice-cooling. The reaction mixture was stirred overnight at room temperature, saturated aqueous sodium thiosulfate solution was added thereto, and the mixture was stirred at room temperature for further 5 min. The reaction mixture was added to saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (2.15 g). MS (API+): [M+H]⁺ 235.1.

[0430]

C) N-(2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 410.1.

¹H NMR (300 MHz, DMSO-d₆) δ 4.28-4.50 (2H, m), 5.45 (1H, d, J = 6.8 Hz), 7.12 (1H, dd, J = 7.9, 4.9 Hz), 7.24-7.41 (4H, m), 7.51-7.60 (2H, m), 7.87 (1H, s), 8.01 (1H, dd, J = 4.9, 1.5 Hz), 10.66 (1H, d, J = 6.8 Hz), 10.81 (1H, s).

[0431]

Example 43

N-(cyano(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 392.1.

¹H NMR (300 MHz, DMSO-d₆) δ 4.46 (2H, s), 6.30 (1H, d, J = 7.5 Hz), 7.13 (1H, dd, J = 7.9, 4.9 Hz), 7.33 (1H, dd, J = 7.9, 1.5 Hz), 7.47 (2H, d, J = 8.3 Hz), 7.64-7.74 (2H, m), 7.93-8.00 (1H, m), 10.67 (1H, d, J = 7.5 Hz), 10.86 (1H, s).

[0432]

Example 44

7-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0433]

A) 2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-7-carbonitrile

To a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.00 g), zinc cyanide (579 mg) and zinc (81 mg) in N,N-dimethylacetamide (10 mL) was added di-tert-butylphosphine palladium(0) (126 mg). The reaction mixture was stirred overnight at 110°C under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (598 mg).
¹H NMR (400 MHz, CDCl₃) δ 0.01 (9H, s), 0.97 (2H, t, J = 10.8 Hz), 3.65 (2H, t, J = 11.0 Hz), 4.31 (2H, d, J = 1.6 Hz), 5.34 (2H, s), 7.57 (1H, d, J = 2.0 Hz), 8.14 (1H, s).

[0434]

B) 7-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step D of Example 26.

¹H NMR (300 MHz, CDCl₃) δ 0.01 (9H, s), 0.91-1.00 (5H, m), 1.84-1.94 (2H, m), 3.66 (2H, t, J = 8.1 Hz), 4.66-4.79 (2H, m), 4.88 (1H, q, J = 6.9 Hz), 5.29-5.37 (2H, m), 7.19 (2H, d, J = 8.1 Hz), 7.34 (2H, d, J = 8.7 Hz), 7.89 (1H, d, J = 2.1 Hz), 8.34 (1H, d, J = 1.8 Hz), 10.01 (1H, d, J = 7.2 Hz).

[0435]

C) 7-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step E of Example 26.

MS (API-): [M-H]⁻ 418.0.

¹H NMR (400 MHz, CDCl₃) δ 0.96 (3H, t, J = 7.2 Hz), 1.85-1.92

(2H, m), 4.65-4.75 (2H, m), 4.87 (1H, q, J = 7.6 Hz), 7.17 (2H, d, J = 8.0 Hz), 7.31-7.34 (2H, m), 7.36 (1H, d, J = 2.0 Hz), 8.29 (1H, d, J = 2.0 Hz), 9.79 (1H, s), 10.18 (1H, d, J = 7.6 Hz).

5 [0436]

Example 45

2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-6-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 [0437]

A) 2-nitroacetamide

An aqueous solution of nitroethyl acetate (45.0 g) in 28% ammonia (300 mL) was stirred at room temperature for 24 hr, and the solvent was evaporated under reduced pressure. To the
15 residue was added water, the pH of the mixture was adjusted to 3 with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the
20 title compound (35 g).

¹H NMR (400 MHz, DMSO-d₆) δ 5.28 (2H, s), 7.64 (1H, s), 7.86 (1H, s).

[0438]

B) 3-nitro-6-(trifluoromethyl)pyridin-2-ol

25 To a solution of 2-nitroacetamide (16.9 g) and (3E)-4-ethoxy-1,1,1-trifluorobut-3-en-2-one (30.0 g) in ethanol (700 mL) was slowly added sodium ethoxide (22.1 g) at room temperature. The reaction mixture was stirred at 90°C for 2 hr, the pH of the mixture was adjusted to 2 with 1 M hydrochloric
30 acid, and the mixture was stirred at room temperature for additional 30 min. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced
35 pressure to give the title compound (5.5 g).

¹H NMR (400 MHz, CDCl₃) δ 7.46 (1H, d, J = 8.0 Hz), 8.69 (1H, d, J = 8.0 Hz), 11.02 (1H, brs).

[0439]

C) 2-chloro-3-nitro-6-(trifluoromethyl)pyridine

5 To 3-nitro-6-(trifluoromethyl)pyridin-2-ol (5.5 g) was added phosphorous pentachloride (25.0 g) at room temperature. The reaction mixture was stirred at 170°C for 2.5 hr, cooled to room temperature, and slowly poured into ice-water. The mixture was extracted with ethyl acetate, and the extract was
10 washed successively with saturated brine and saturated aqueous sodium hydrogen carbonate solution, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (5.0 g).

¹H NMR (300 MHz, CDCl₃) δ 7.84 (1H, d, J = 8.1 Hz), 8.38 (1H, d, J = 8.1 Hz).
15

[0440]

D) methyl N-(6-trifluoromethyl-3-nitropyridin-2-yl)glycinate

The title compound was obtained in the same manner as in Step A of Example 1.

20 ¹H NMR (300 MHz, CDCl₃) δ 3.81 (3H, s), 4.41 (2H, d, J = 5.4 Hz), 7.07 (1H, d, J = 8.4 Hz), 8.53 (1H, brs), 8.62 (1H, d, J = 8.4 Hz).

[0441]

E) 6-(trifluoromethyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-
25 one

To a solution of methyl N-(6-trifluoromethyl-3-nitropyridin-2-yl)glycinate (3.37 g) in ethanol (75 mL) was added tin(II) chloride dihydrate (14.2 g). The reaction mixture was stirred at 80°C for 2 hr, and the solvent was
30 evaporated under reduced pressure. The residue was dissolved in ethyl acetate, the pH of the solution was adjusted to 9 with saturated aqueous ammonia solution, and the insoluble substance was filtered off using Celite. The filtrate was concentrated under reduced pressure, and the obtained residue
35 was purified by silica gel column chromatography (petroleum

ether/ethyl acetate) to give the title compound (1.04 g).

^1H NMR (400 MHz, $\text{CDCl}_3+\text{CD}_3\text{OD}$) δ 4.19 (2H, s), 6.97 (1H, d, J = 7.6 Hz), 7.02 (1H, d, J = 7.6 Hz).

[0442]

- 5 F) 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-6-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step D of Example 26.

- 10 MS (API+): $[\text{M}+\text{H}]^+$ 463.1.

^1H NMR (400 MHz, CDCl_3) δ 0.95 (3H, t, J = 7.2 Hz), 1.79-1.93 (2H, m), 4.66-4.75 (2H, m), 4.85 (1H, q, J = 7.2 Hz), 7.17 (2H, d, J = 8.0 Hz), 7.28 (1H, d, J = 8.0 Hz), 7.32-7.36 (3H, m), 7.63 (1H, brs), 10.12 (1H, d, J = 7.6 Hz).

- 15 [0443]

Example 46

2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-6-vinyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

- To a solution of 6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.10 g), 2-ethenyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (539 mg) and (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (100 mg) in a mixed solvent of 1,2-dimethoxyethane (15 mL) and water (5 mL) was added cesium carbonate (1.67 g) at room temperature. The reaction mixture was stirred at 120°C for 4 hr under nitrogen atmosphere using microwave, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (500 mg).

MS (API+): $[\text{M}+\text{H}]^+$ 421.2.

- 35 ^1H NMR (300 MHz, CDCl_3) δ 0.98 (3H, t, J = 7.2 Hz), 1.84-1.94

(2H, m), 4.70 (2H, s), 4.90 (1H, d, J = 7.2 Hz), 5.41 (1H, d, J = 10.8 Hz), 5.91 (1H, d, J = 17.1 Hz), 6.68 (1H, dd, J = 17.4, 7.8 Hz), 6.98 (1H, d, J = 8.1 Hz), 7.16-7.22 (3H, m), 7.37 (2H, d, J = 8.4 Hz), 10.14 (1H, s), 10.54 (1H, d, J = 7.2 Hz).

[0444]

Example 47

6-benzyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 To a solution of 6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (420 mg), 2-benzyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (181 mg) and (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (50 mg)
15 in a mixed solvent of 1,2-dimethoxyethane (15 mL) and water (5 mL) was added cesium carbonate (645 mg) at room temperature. The reaction mixture was stirred at 100°C for 40 min under nitrogen atmosphere using microwave, water was added thereto, and the mixture was extracted with ethyl acetate. The extract
20 was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous
25 sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (100 mg).

MS (API+): [M+H]⁺ 485.2.

30 ¹H NMR (300 MHz, CDCl₃) δ 0.88 (3H, t, J = 7.2 Hz), 1.60-1.68 (2H, m), 4.07 (2H, s), 4.65 (2H, s), 4.80 (1H, d, J = 7.2 Hz), 6.89 (1H, d, J = 8.1 Hz), 7.08-7.19 (8H, m), 7.25-7.34 (2H, m), 9.69 (1H, s), 10.34 (1H, d, J = 7.5 Hz).

[0445]

35 Example 48

2-oxo-N-(1-(4-(tetrahydro-2H-pyran-4-yl)phenyl)propyl)-2,3-dihydro
pyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-(1-(4-(3,6-dihydro-2H-pyran-4-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (42.5 mg) in methanol (3 mL) was added 10% palladium-carbon (containing 50% water, 15 mg). The reaction mixture was stirred at room temperature for 4 hr under hydrogen atmosphere, the insoluble substance was filtered off,
10 and the filtrate was concentrated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate.
15 The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (27.9 mg).

MS (API+): [M+H]⁺ 395.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.85 (3H, t, J = 7.3 Hz), 1.52-1.86
20 (6H, m), 2.62-2.82 (1H, m), 3.34-3.50 (2H, m), 3.86-4.00 (2H, m), 4.41 (2H, s), 4.74 (1H, q, J = 6.9 Hz), 7.09 (1H, dd, J = 7.9, 4.9 Hz), 7.15-7.28 (4H, m), 7.31 (1H, dd, J = 7.7, 1.7 Hz), 8.01 (1H, dd, J = 4.9, 1.5 Hz), 10.29 (1H, d, J = 7.5 Hz), 10.80 (1H, brs).

25 [0446]

Example 49

N-(1-(6-methoxypyridin-3-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0447]

30 A) 6-methoxynicotinonitrile

To a solution of 6-chloronicotinonitrile (10.0 g) in methanol (100 mL) was added sodium methoxide (7.80 g). The reaction mixture was heated at reflux overnight, and the solvent was evaporated under reduced pressure. To the residue
35 was added water, and the mixture was extracted with ethyl

acetate. The extract was washed with saturated brine, and dried over sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (8.8 g).

¹H NMR (400 MHz, DMSO-d₆) δ 4.00 (3H, s), 6.83 (1H, dd, J= 8.8, 0.8 Hz), 7.78 (1H, dd, J= 8.6, 2.4 Hz), 8.50 (1H, d, J = 1.4 Hz).

[0448]

B) 6-methoxynicotine acid

To a solution of 6-methoxynicotinonitrile (5.00 g) in ethanol (100 mL) was added 2 M aqueous potassium hydroxide solution (20 mL). The reaction mixture was heated at reflux for 2 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, the pH of the mixture was adjusted to 4-5 with 2 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (4.2 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.92 (3H, s), 6.86-6.92 (1H, m), 8.11-8.15 (1H, m), 8.73 (1H, d, J = 2.0 Hz), 13.0 (1H, s).

[0449]

C) N,6-dimethoxy-N-methylnicotinamide

To a solution of 6-methoxynicotine acid (4.2 g) and N-methoxymethylamine hydrochloride (3.8 g) in dichloromethane (100 mL) were added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (7.6 g) and 4-dimethylaminopyridine (0.32 g). The reaction mixture was stirred at room temperature for 10 min, diisopropylethylamine (10.0 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (3.6 g).

¹H NMR (400 MHz, CDCl₃) δ 3.58 (3H, s), 3.77 (3H, s), 3.98 (3H, s), 6.72-6.77 (1H, m), 7.97-8.00 (1H, m), 8.62-8.65 (1H, m).
[0450]

D) 1-(6-methoxypyridin-3-yl)propan-1-one

5 To a solution of N,6-dimethoxy-N-methylnicotinamide (3.6 g) in tetrahydrofuran (50 mL) was added 3 M ethylmagnesium bromide/tetrahydrofuran solution (18 mL) at -40°C. The reaction mixture was stirred at the same temperature for 2 hr, saturated aqueous ammonium chloride solution was added thereto,
10 and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (2.9 g).

¹H NMR (400 MHz, CDCl₃) δ 1.23 (3H, t, J = 7.2 Hz), 2.95 (2H, q, J = 7.2 Hz), 4.00 (3H, s), 6.78 (1H, dd, J = 8.4, 0.4 Hz),
15 8.15 (1H, dd, J = 8.8, 2.4 Hz), 8.80 (1H, d, J = 2.4 Hz).
[0451]

E) 1-(6-methoxypyridin-3-yl)propan-1-amine

To a solution of 1-(6-methoxypyridin-3-yl)propan-1-one
20 (2.9 g) in ethanol (50 mL) were added hydroxylamine hydrochloride (1.5 g) and triethylamine (2.1 g). The reaction mixture was heated at reflux overnight, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The
25 extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. To a solution of the obtained solid (800 mg) in methanol (10 mL) was added Raney nickel (160 mg). The reaction mixture was stirred at 50°C for 3 hr under hydrogen
30 atmosphere (50 psi), the insoluble substance was filtered off using Celite, and the filtrate was concentrated under reduced pressure to give the title compound (600 mg).

¹H NMR (400 MHz, DMSO-d₆) δ 0.76 (3H, t, J = 7.6 Hz), 1.45-1.61 (2H, m), 3.35 (2H, brs), 3.64-3.67 (1H, m), 3.81 (3H, s), 6.74
35 (1H, dd, J = 13.6, 8.4 Hz), 7.67 (1H, dd, J = 8.4, 2.4 Hz),

8.03 (1H, dd, J= 8.4, 2.0 Hz).

[0452]

F) N-(1-(6-methoxypyridin-3-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 To a solution of 1-(6-methoxypyridin-3-yl)propan-1-amine (100 mg) in N,N-dimethylformamide (10 mL) were added 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (223 mg) and triethylamine (181 mg). The reaction mixture was stirred overnight at room temperature, and the
10 solvent was evaporated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was
15 dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (80 mg).

MS (API+): [M+H]⁺ 342.1.

¹H NMR (300 MHz, CDCl₃) δ 0.97 (3H, t, J = 7.5 Hz), 1.83-1.95 (2H, m), 3.92 (3H, s), 4.62-4.75 (2H, m), 4.87 (1H, q, d, J =
20 7.2 Hz), 6.72 (1H, d, J = 8.4 Hz), 6.99 (1H, dd, J = 7.8, 5.1 Hz), 7.20 (1H, dd, J = 7.8, 1.5 Hz), 7.55 (1H, dd, J = 8.7, 2.4 Hz), 8.00 (1H, dd, J = 4.8, 1.5 Hz), 8.14 (1H, d, J = 2.7 Hz), 9.16 (1H, s), 10.40 (1H, d, J = 7.2 Hz).

[0453]

25 Example 50

N-(1-(5-methoxypyrimidin-2-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0454]

A) 5-methoxypyrimidine-2-carbonitrile

30 To a solution of 2-chloro-5-methoxypyrimidine (4.4 g) and 1,4-diazabicyclo[2.2.2]octane (6.7 g) in a mixed solvent of dimethyl sulfoxide (114 mL) and water (18 mL) was added potassium cyanide (4.00 g). The reaction mixture was stirred at 70°C for 5 hr, water was added thereto, and the mixture was
35 extracted with ethyl acetate. The extract was washed with

saturated brine, and dried over sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (3.5 g).

¹H NMR (400 MHz, CDCl₃) δ 4.02 (3H, s), 8.47 (2H, s).

5 [0455]

B) 1-(5-methoxypyrimidin-2-yl)propan-1-one

The title compound was obtained in the same manner as in Step B of Example 30.

¹H NMR (300 MHz, CDCl₃) δ 1.25 (3H, t, J = 7.2 Hz), 3.23 (2H, q, J = 7.2 Hz), 4.02 (3H, s), 8.53 (2H, s).

[0456]

C) N-benzyl-1-(5-methoxypyrimidin-2-yl)propan-1-amine

To a solution of 1-(5-methoxypyrimidin-2-yl)propan-1-one (500 mg) and benzyl amine (308 mg) in 1,2-dichloroethane (10 mL) was added sodium triacetoxyborohydride (953.5 mg). The reaction mixture was stirred overnight at 25°C, dichloromethane was added thereto, and washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (700 mg).

¹H NMR (300 MHz, CDCl₃) δ 0.85 (3H, t, J = 7.5 Hz), 1.77-1.83 (2H, m), 3.61 (2H, dd, J = 15.3, 12.9 Hz), 3.82 (1H, t, J = 6.6 Hz), 3.94 (3H, s), 7.20-7.36 (5H, m), 8.40 (2H, s).

[0457]

25 D) 1-(5-methoxypyrimidin-2-yl)propan-1-amine

To a solution of N-benzyl-1-(5-methoxypyrimidin-2-yl)propan-1-amine (700 mg) in methanol (10 mL) was added 10% palladium-carbon (containing 50% water, 100 mg). The reaction mixture was stirred at room temperature for 5 hr under hydrogen atmosphere (20 psi), the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure to give the title compound (500 mg).

¹H NMR (300 MHz, CDCl₃) δ 0.92 (3H, t, J = 7.5 Hz), 1.69-1.79 (2H, m), 3.94 (3H, s), 3.92-4.02 (1H, m), 8.38 (2H, s).

35 [0458]

E) N-(1-(5-methoxypyrimidin-2-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

5 MS (API+): [M+H]⁺ 343.2.

¹H NMR (400 MHz, CDCl₃) δ 0.95 (3H, t, J = 7.2 Hz), 1.96-2.09 (2H, m), 3.91 (3H, s), 4.60 (2H, q, J = 18.0 Hz), 5.17 (1H, q, J = 6.8 Hz), 6.95-6.98 (1H, dd, J = 8.0, 4.8 Hz), 7.14 (1H, dd, J = 7.6, 1.6 Hz), 8.03 (1H, dd, J = 4.8, 1.6 Hz), 8.40 (2H, s),
10 9.46 (1H, s), 10.63 (1H, d, J = 7.2 Hz).

[0459]

Example 51

6-ethyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 The title compound was obtained in the same manner as in Example 29.

MS (API+): [M+H]⁺ 423.1.

¹H NMR (400 MHz, CDCl₃) δ 0.97 (3H, t, J = 7.2 Hz), 1.27 (3H, t, J = 7.6 Hz), 1.83-1.93 (2H, m), 2.71-2.77 (2H, m), 4.66 (2H, s),
20 s), 4.87-4.92 (1H, m), 6.82 (1H, d, J = 7.6 Hz), 7.13-7.18 (3H, m), 7.33-7.37 (2H, m), 9.91 (1H, s), 10.65 (1H, d, J = 7.6 Hz).

[0460]

Example 52

6-methoxy-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0461]

A) methyl N-(6-methoxy-3-nitropyridin-2-yl)glycinate

To a solution of ethyl N-(6-chloro-3-nitropyridin-2-yl)glycinate (1.50 g) in methanol (40 mL) was added sodium
30 methoxide (469 mg). The reaction mixture was stirred at 70°C for 2 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the
35 title compound (1.4 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.67 (3H, s), 3.84 (3H, s), 4.29 (2H, d, J = 6.0 Hz), 6.22 (1H, t, J = 4.4 Hz), 8.34 (1H, d, J = 8.8 Hz), 9.11 (1H, s).

[0462]

- 5 B) 6-methoxy-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of methyl N-(6-methoxy-3-nitropyridin-2-yl)glycinate (1.40 g) in ethanol (40 mL) was added tin(II) chloride dihydrate (8.00 g). The reaction mixture was stirred
10 overnight at 80°C, and the solvent was evaporated under reduced pressure to give a crude product (9.4 g) of 6-methoxy-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one. To a solution of the crude product (4.00 g) of 6-methoxy-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one and diisopropylethylamine (10 mL) in
15 tetrahydrofuran (100 mL) was slowly added triphosgene (602 mg) at room temperature. The reaction mixture was stirred at 45°C for 2 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in tetrahydrofuran (50 mL), and 1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride
20 (602 mg) was added thereto. The mixture was stirred overnight at 65°C, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the
25 solvent was evaporated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was
30 dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (90 mg).

MS (API+): [M+H]⁺ 425.1.

¹H NMR (400 MHz, CDCl₃) δ 0.97 (3H, t, J = 7.6 Hz), 1.82-1.91 (2H, m), 3.77 (3H, s), 4.56-4.68 (2H, m), 4.85-4.90 (1H, m),
35 6.46 (1H, d, J = 8.4 Hz), 7.16-7.22 (3H, m), 7.36 (2H, d, J

=8.4 Hz), 9.61 (1H, d, J = 7.2 Hz), 9.64 (1H, s).

[0463]

Example 53

N-(1-(4-(cyclopropylmethoxy)phenyl)propyl)-2-oxo-2,3-

5 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0464]

A) tert-butyl 1-(4-

(cyclopropylmethoxy)phenyl)propyl)carbamate

To a solution of tert-butyl 1-(4-

10 hydroxyphenyl)propyl)carbamate (303 mg), cyclopropylmethanol
(104 mg) and tributylphosphine (394 mg) in tetrahydrofuran (10
mL) was added 1,1'-(azodicarbonyl)dipiperidine (457 mg). The
reaction mixture was stirred at room temperature for 1 hr, and
hexane was added thereto. The precipitated solid was filtered
15 off, and the filtrate was concentrated under reduced pressure.
The residue was purified by silica gel column chromatography
(hexane/ethyl acetate) to give the title compound (292 mg).

MS (API+): [M+H]⁺ 306.2.

[0465]

20 B) N-(1-(4-(cyclopropylmethoxy)phenyl)propyl)-2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of the hydrochloride (217 mg) (obtained
from tert-butyl 1-(4-

(cyclopropylmethoxy)phenyl)propyl)carbamate (292 mg) in the

25 same manner as in Step E of Example 18) in N,N-
dimethylacetamide (6 mL) were added 4-nitrophenyl 2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (282 mg) and
triethylamine (0.456 mL). The reaction mixture was stirred at
room temperature for 16 hr, and the solvent was evaporated
30 under reduced pressure. To the residue was added aqueous
sodium hydrogen carbonate solution, and the mixture was
extracted with ethyl acetate. The extract was washed with
water, and dried over anhydrous sodium sulfate, and the
solvent was evaporated under reduced pressure. The obtained
35 solid was triturated with ethyl acetate/methanol, collected by

filtration, and washed with ethyl acetate to give the title compound (136 mg).

MS (API+): [M+H]⁺ 381.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.20-0.37 (2H, m), 0.48-0.64 (2H, m), 0.83 (3H, t, J = 7.3 Hz), 1.09-1.32 (1H, m), 1.64-1.88 (2H, m), 3.77 (2H, d, J = 6.8 Hz), 4.41 (2H, d, J = 0.8 Hz), 4.70 (1H, q, J = 7.0 Hz), 6.81-6.93 (2H, m), 7.09 (1H, dd, J = 7.9, 4.9 Hz), 7.14-7.24 (2H, m), 7.30 (1H, dd, J = 7.9, 1.5 Hz), 7.99 (1H, dd, J = 4.9, 1.5 Hz), 10.23 (1H, d, J = 7.5 Hz), 10.80 (1H, s).

[0466]

Example 54

2-oxo-N-(1-(4-(trifluoromethyl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0467]

A) N-hydroxy-1-(4-(trifluoromethyl)phenyl)propan-1-imine

The title compound was obtained in the same manner as in Step C of Example 1.

¹H NMR (300 MHz, CDCl₃) δ 1.06-1.23 (3H, m), 2.54-2.90 (2H, m), 7.47-7.78 (4H, m), 8.14-8.53 (1H, m).

[0468]

B) 1-(4-(trifluoromethyl)phenyl)propan-1-amine

To a solution of N-hydroxy-1-(4-(trifluoromethyl)phenyl)propan-1-imine (2.15 g) in ethanol (100 mL) was added 20% palladium hydroxide-carbon (200 mg). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (988 mg).

¹H NMR (300 MHz, CDCl₃) δ 0.87 (3H, t, J = 7.3 Hz), 1.62-1.77 (2H, m), 3.89 (1H, t, J = 6.8 Hz), 7.44 (2H, d, J = 7.9 Hz), 7.58 (2H, d, J = 7.9 Hz).

[0469]

C) 2-oxo-N-(1-(4-(trifluoromethyl)phenyl)propyl)-2,3-

dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 379.2.

5 ¹H NMR (300 MHz, CDCl₃) δ 0.97 (3H, t, J = 7.3 Hz), 1.89 (2H, quin, J = 7.3 Hz), 4.67 (2H, s), 4.93 (1H, q, J = 6.8 Hz), 7.01 (1H, dd, J = 7.7, 5.1 Hz), 7.19 (1H, dd, J = 7.5, 1.5 Hz), 7.44 (2H, d, J = 8.3 Hz), 7.58 (2H, d, J = 8.3 Hz), 8.01 (1H, dd, J = 4.9, 1.5 Hz), 9.16 (1H, brs), 10.48 (1H, d, J = 7.2
10 Hz).

[0470]

Example 55

2-oxo-N-((pyridin-2-yl)(4-(trifluoromethoxy)phenyl)methyl)-
2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 [0471]

A) N-methoxy-N-methylpyridine-2-carboxamide

The title compound was obtained in the same manner as in Step A of Example 31.

MS (API+): [M+H]⁺ 167.1.

20 [0472]

B) pyridin-2-yl(4-(trifluoromethoxy)phenyl)methanone

To a solution of N-methoxy-N-methylpyridine-2-carboxamide (2.7 g) in tetrahydrofuran (60 mL) was added 0.5 M (4-(trifluoromethoxy)phenyl)magnesium bromide/tetrahydrofuran
25 solution (32.5 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, saturated aqueous ammonium chloride solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate,
30 and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (2.69 g).

MS (API+): [M+H]⁺ 268.0.

[0473]

35 C) 1-(pyridin-2-yl)-1-(4-(trifluoromethoxy)phenyl)methanamine

hydrochloride

The title compound was obtained in the same manner as in Steps C-E of Example 1.

MS (API+): [M+H]⁺ 269.0.

5 [0474]

D) 2-oxo-N-((pyridin-2-yl)(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

10 MS (API+): [M+H]⁺ 444.2.

¹H NMR (300 MHz, CDCl₃) δ 4.53-4.74 (2H, m), 6.25 (1H, d, J = 6.8 Hz), 6.99 (1H, dd, J = 7.9, 4.9 Hz), 7.09-7.25 (4H, m), 7.29 (1H, d, J = 7.9 Hz), 7.43-7.53 (2H, m), 7.66 (1H, td, J = 7.7, 1.9 Hz), 8.08 (1H, dd, J = 5.1, 1.7 Hz), 8.64 (1H, dd, J = 15 4.9, 0.8 Hz), 9.24 (1H, brs), 11.41 (1H, d, J = 6.8 Hz).

[0475]

Example 56

N-((3,5-dimethyl-1,2-oxazol-4-yl)(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-
20 b]pyrazine-4(1H)-carboxamide

[0476]

A) (3,5-dimethyl-1,2-oxazol-4-yl)(4-(trifluoromethoxy)phenyl)methanone

The title compound was obtained in the same manner as in
25 Step A of Example 31 and Step B of Example 55.

MS (API+): [M+H]⁺ 286.0.

[0477]

B) 1-(3,5-dimethyl-1,2-oxazol-4-yl)-N-hydroxy-1-(4-(trifluoromethoxy)phenyl)methanimine

30 The title compound was obtained in the same manner as in Step C of Example 1.

MS (API+): [M+H]⁺ 301.1.

[0478]

C) 1-(3,5-dimethyl-1,2-oxazol-4-yl)-1-(4-(trifluoromethoxy)phenyl)methanamine
35

To a solution of 1-(3,5-dimethyl-1,2-oxazol-4-yl)-N-hydroxy-1-(4-(trifluoromethoxy)phenyl)methanimine (460 mg) in tetrahydrofuran (15 mL) was added 1.1 M borane-tetrahydrofuran complex/tetrahydrofuran solution (4.18 mL) at room temperature.

5 The reaction mixture was stirred overnight at 80°C under nitrogen atmosphere, 1 M hydrochloric acid was added thereto, and the mixture was stirred at room temperature for 15 min. Then, to the reaction mixture was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was

10 extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (398 mg).

15 MS (API+), found: 270.1.

[0479]

D) N-((3,5-dimethyl-1,2-oxazol-4-yl)(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 462.1.

¹H NMR (300 MHz, CDCl₃) δ 2.12 (3H, s), 2.38 (3H, s), 4.62-4.83 (2H, m), 6.25 (1H, d, J = 8.3 Hz), 7.02 (1H, dd, J = 7.7, 5.1

25 Hz), 7.15-7.23 (3H, m), 7.27-7.36 (2H, m), 7.91 (1H, dd, J = 5.1, 1.3 Hz), 8.38 (1H, brs), 10.79 (1H, d, J = 8.3 Hz).

[0480]

Example 57

N-((1-methyl-1H-pyrazol-4-yl)(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30

The title compound was obtained in the same manner as in Step A of Example 31, Step B of Example 55, Steps C-E of Example 1 and Step G of Example 7.

35 MS (API+): [M+H]⁺ 447.2.

¹H NMR (300 MHz, CDCl₃) δ 3.86 (3H, s), 4.60-4.77 (2H, m), 6.18 (1H, d, J = 7.2 Hz), 6.99 (1H, dd, J = 7.7, 5.1 Hz), 7.08-7.24 (4H, m), 7.33-7.48 (3H, m), 7.94 (1H, d, J = 4.9 Hz), 9.00 (1H, brs), 10.68 (1H, d, J = 7.5 Hz).

5 [0481]

Example 58

N-(2-(dimethylamino)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0482]

10 A) tert-butyl (cyano(4-(trifluoromethoxy)phenyl)methyl)carbamate

To a solution of amino (4-(trifluoromethoxy)phenyl)acetonitrile hydrochloride (3 g) and triethylamine (3.31 mL) in tetrahydrofuran (50 mL) was added
15 di-tert-butyl dicarbonate (3.03 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, further di-tert-butyl dicarbonate (500 mg) was added at room temperature, and the mixture was stirred for 2 days. The reaction mixture was filtered through silica gel pad, and
20 eluted with ethyl acetate. The filtrate was concentrated under reduced pressure to give the title compound (4.14 g).

¹H NMR (300 MHz, CDCl₃) δ 1.49 (9H, s), 5.01-5.24 (1H, m), 5.82 (1H, brs), 7.22-7.34 (2H, m), 7.49-7.60 (2H, m).
[0483]

25 B) tert-butyl (2-amino-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

To a solution of cobalt(II) chloride (328 mg) and tert-butyl (cyano(4-(trifluoromethoxy)phenyl)methyl)carbamate (400 mg) in methanol (8 mL) was added sodium borohydride (478 mg)
30 under ice-cooling. The reaction mixture was stirred for 40 min under ice-cooling, 1 M hydrochloric acid (10 mL) was added thereto, and the mixture was stirred at room temperature for 10 min. To the reaction mixture was added 8 M aqueous sodium hydroxide solution (5 mL), the insoluble substance was
35 filtered off using Celite, and the filtrate was extracted with

ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (360 mg).

5 MS (API+): [M+H]⁺ 321.1.

[0484]

C) tert-butyl (2-(dimethylamino)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

To a solution of tert-butyl (2-amino-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (180 mg) and 37%
10 aqueous formaldehyde solution (109 mg) in methanol (5 mL) was added borane-2-picoline complex (180 mg) under ice-cooling. The reaction mixture was stirred overnight at room temperature under nitrogen atmosphere, 1 M hydrochloric acid (5 mL) was
15 added thereto at room temperature, and the mixture was stirred at the same temperature for 10 min. The reaction mixture was neutralized with saturated aqueous sodium hydrogen carbonate solution, and extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried
20 over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (118 mg).

MS (API+): [M+H]⁺ 349.1.

[0485]

25 D) N²,N²-dimethyl-1-(4-(trifluoromethoxy)phenyl)ethane-1,2-diamine dihydrochloride

To tert-butyl (2-(dimethylamino)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (117 mg) was added 4
M hydrogen chloride/ethyl acetate solution (5 mL). The
30 reaction mixture was stirred at room temperature for 10 min, and the solvent was evaporated under reduced pressure to give the title compound (128 mg).

MS (API+), found: 249.1.

[0486]

35 E) N-(2-(dimethylamino)-1-(4-(trifluoromethoxy)phenyl)ethyl)-

2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 424.1.

5 ¹H NMR (300 MHz, DMSO-d₆) δ 2.20 (6H, s), 2.35-2.78 (2H, m), 4.29-4.52 (2H, m), 4.89 (1H, brs), 7.10 (1H, dd, J = 7.5, 4.9 Hz), 7.25-7.35 (3H, m), 7.45 (2H, d, J = 8.3 Hz), 7.99 (1H, dd, J = 4.9, 1.9 Hz), 10.38 (1H, d, J = 6.0 Hz), 10.81 (1H, s).

[0487]

10 Example 59

N-(2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0488]

A) 2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethanone

15 To a solution of 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone (5 g) in tetrahydrofuran (150 mL) was added sodium thiomethoxide (1.24 g) under ice-cooling. The reaction mixture was stirred at room temperature for 1 hr, filtered through NH silica gel pad, and eluted with
20 ethyl acetate/hexane. The filtrate was concentrated under reduced pressure to give the title compound (4.23 g).

¹H NMR (300 MHz, CDCl₃) δ 2.14 (3H, s), 3.74 (2H, s), 7.27-7.36 (2H, m), 7.99-8.10 (2H, m).

[0489]

25 B) 2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethanamine

The title compound was obtained in the same manner as in Step C of Example 1 and Step C of Example 56.

MS (API+), found: 235.0.

[0490]

30 C) N-(2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺ 427.1.

35 ¹H NMR (300 MHz, CDCl₃) δ 2.05 (3H, s), 2.97 (2H, d, J = 6.0

Hz), 4.69 (2H, s), 5.24 (1H, q, J = 6.4 Hz), 7.01 (1H, dd, J = 7.7, 5.1 Hz), 7.12-7.25 (3H, m), 7.34-7.46 (2H, m), 8.03 (1H, dd, J = 4.9, 1.5 Hz), 9.55 (1H, brs), 10.76 (1H, d, J = 7.2 Hz).

5 [0491]

Example 60

N-(2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-(2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (54.0 mg) in ethyl acetate (10 mL) was added 3-chloroperbenzoic acid (68.7 mg) at room temperature. The reaction mixture was stirred overnight at room temperature, saturated aqueous sodium thiosulfate solution was added thereto, and the mixture was stirred for 5 min. Then, to the reaction mixture was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained solid was triturated with hexane/ethyl acetate, collected by filtration, and washed with hexane/ethyl acetate to give the title compound (38.2 mg).

25 MS (API+): [M+H]⁺ 459.2.

¹H NMR (300 MHz, DMSO-d₆) δ 2.91 (3H, s), 3.63 (1H, dd, J = 14.7, 4.5 Hz), 4.00 (1H, dd, J = 14.5, 9.2 Hz), 4.26-4.56 (2H, m), 5.46 (1H, td, J = 8.5, 3.8 Hz), 7.11 (1H, dd, J = 7.9, 4.9 Hz), 7.30 (1H, dd, J = 7.9, 1.5 Hz), 7.33-7.41 (2H, m), 7.53-7.61 (2H, m), 7.98 (1H, dd, J = 4.9, 1.5 Hz), 10.62 (1H, d, J = 7.9 Hz), 10.82 (1H, s).

[0492]

Example 61

N-(1-(5-methoxypyridin-2-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0493]

A) methyl 5-methoxypyridine-2-carboxylate

To a solution of 2-bromo-5-methoxypyridine (21 g) and triethylamine (55 g) in a mixed solvent of N,N-dimethylformamide (100 mL) and methanol (100 mL) were added 1,3-bis(diphenylphosphino)-propane (7.0 g) and palladium(II) acetate (3.6 g). The reaction mixture was stirred at 100°C for 24 hr under carbon monoxide atmosphere (5.0 MPa), and diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (10.2 g).

¹H NMR (300 MHz, CDCl₃) δ 3.93 (3H, s), 3.99 (3H, s), 7.25-7.29 (1H, m), 8.12 (1H, d, J = 8.4 Hz), 8.40 (1H, d, J = 3.0 Hz).

[0494]

B) 5-methoxypyridine-2-carboxylic acid

To a solution of methyl 5-methoxypyridine-2-carboxylate (4.0 g) in methanol (50 mL) was added a solution of lithium hydroxide monohydrate (5.0 g) in water (10 mL). The reaction mixture was stirred at room temperature for 12 hr, the pH of the mixture was adjusted to 4-5 with 2 M hydrochloric acid, and the precipitate was collected by filtration. The obtained solid was washed with water, and dried to give the title compound (2.4 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.93 (3H, s), 7.49-7.51 (1H, m), 8.03 (1H, d, J = 7.2 Hz), 8.36 (1H, d, J = 2.8 Hz), 12.85 (1H, s).

[0495]

C) N-(1-(5-methoxypyridin-2-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Steps C-E of Example 49 and Step G of Example 7.

MS (API+): [M+H]⁺ 342.1.

¹H NMR (400 MHz, CDCl₃) δ 0.79 (3H, t, J = 7.2 Hz), 1.77-1.85 (2H, m), 3.81 (3H, s), 4.43 (2H, s), 4.82 (1H, q, J = 6.8 Hz), 7.09 (1H, dd, J = 8.0, 5.2 Hz), 7.28-7.36 (3H, m), 8.00 (1H, d, J = 4.0 Hz), 8.26 (1H, d, J = 2.8 Hz), 10.35 (1H, d, J = 7.6 Hz).

[0496]

Example 62

6-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 [0497]

A) 6-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.00 g), zinc cyanide (126 mg) and zinc (200 mg) in N,N-dimethylformamide (10 mL) were added tris(dibenzylideneacetone)dipalladium(0) (100 mg) and 1,1'-bis(diphenylphosphino)ferrocene (100 mg). The reaction mixture was stirred at 120°C for 6 hr under nitrogen atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (320 mg).

¹H NMR (300 MHz, CDCl₃) δ 0.00 (9H, s), 0.92-1.00 (5H, m), 1.84-1.98 (2H, m), 3.61-3.67 (2H, m), 4.66-4.78 (2H, m), 4.85-4.90 (1H, m), 5.30-5.38 (2H, m), 7.20 (2H, d, J = 8.4 Hz), 7.36 (2H, d, J = 8.8 Hz), 7.46 (1H, d, J = 8.4 Hz), 7.74 (1H, d, J = 8.4 Hz), 9.82 (1H, d, J = 7.6 Hz).

[0498]

B) 6-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To 6-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (320 mg) was added
5 trifluoroacetic acid (3 mL) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and the solvent was evaporated under reduced pressure. To the residue was added aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was
10 washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate
15 solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give the title compound (110 mg).

MS (API-): [M-H]⁻ 418.2.

20 ¹H NMR (400 MHz, DMSO-d₆) δ 0.94 (3H, t, J = 7.2 Hz), 1.76-1.84 (2H, m), 4.34-4.43 (2H, m), 4.80-4.85 (2H, m), 7.31-7.36 (3H, m), 7.48 (2H, d, J = 8.4 Hz), 7.71 (1H, d, J = 8.0 Hz), 9.59 (1H, d, J = 7.2 Hz), 11.27 (1H, s).

[0499]

25 Example 63

7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0500]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

30 To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride (10.6 g) and triethylamine (11.8 mL) at room temperature. The reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue
35 was added water, and the mixture was extracted with ethyl

acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the
5 title compound (11.1 g).

MS (API+): [M+H]⁺ 338.0.

[0501]

B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (6.00 g) in tetrahydrofuran (300 mL) was added 5%
10 platinum-carbon (500 mg). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol
15 (200 mL), the solution was heated at reflux for 16 hr, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (4.48 g).

MS (API+): [M+H]⁺ 276.0.

20 [0502]

C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a suspension of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.82 g) in N,N-dimethylformamide (160 mL)
25 was slowly added 1 M potassium hexamethyldisilazide/tetrahydrofuran solution (17.4 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, 2-(chloromethoxy)ethyltrimethylsilane (3.62 mL) was slowly added thereto, and the mixture was stirred at room temperature for
30 20 hr. To the reaction mixture was added water, the insoluble substance was filtered off using Celite, and the filtrate was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated
35 under reduced pressure. The residue was purified by silica gel

column chromatography (NH, hexane/ethyl acetate) to give the title compound (2.71 g).

MS (API+): $[M+H]^+$ 406.0.

[0503]

5 D) 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanone

To a solution of 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone (6.12 g) in methanol (150 mL) were added silver(I) carbonate (7.75 g) and boron trifluoride diethyl ether complex (3.29 mL) at room
10 temperature. The reaction mixture was stirred at 50°C for 20 hr, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated
15 brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (4.78 g).

MS (API+): $[M+H]^+$ 235.1.

20 [0504]

E) 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanamine hydrochloride

To a solution of 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanone (4.78 g) in ethanol (120 mL)
25 were added hydroxylamine hydrochloride (2.84 g) and triethylamine (5.69 mL) at room temperature. The reaction mixture was stirred at room temperature for 5.5 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl
30 acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a crude product of N-hydroxy-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanimine. To a solution of the
35 obtained crude product in ethanol (160 mL) was added 10%

palladium-carbon (containing 50% water, 350 mg). The reaction mixture was stirred at room temperature for 16 hr under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. To a
5 solution of the residue in ethyl acetate (10 mL) was added 4 M hydrogen chloride/ethyl acetate solution (100 mL), the mixture was stirred at room temperature for 2 hr, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (4.2 g).

10 ¹H NMR (300 MHz, DMSO-d₆) δ 3.34 (3H, s), 3.58-3.74 (2H, m), 4.58 (1H, dd, J = 6.6, 5.5 Hz), 7.46 (2H, d, J = 8.7 Hz), 7.65 (2H, d, J = 8.3 Hz), 8.57 (3H, brs).

[0505]

F) 7-iodo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-
15 oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.11 g) in tetrahydrofuran (42.0 mL) was
20 added a solution of bis(trichloromethyl)carbonate (647 mg) in tetrahydrofuran (6.30 mL) at room temperature under nitrogen atmosphere. The reaction mixture was stirred at 40°C for 1 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue was added
25 tetrahydrofuran, and the solvent was evaporated under reduced pressure. The operation (addition of tetrahydrofuran and then evaporation) was repeated three times. The residue was diluted with tetrahydrofuran (21.0 mL), and the mixture was added to a mixture of 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanamine
30 hydrochloride (889 mg) and triethylamine (1.15 mL) in tetrahydrofuran (14.0 mL) at room temperature. The reaction mixture was stirred overnight at 60°C, and the solvent was evaporated under reduced pressure. To the residue was added ethyl acetate, washed successively with water and saturated
35 brine, and dried over anhydrous sodium sulfate, and the

solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (1.17 g).

MS (API+): [M+H]⁺ 667.1.

5 [0506]

G) 7-hydroxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 To a solution of 7-iodo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (100 mg), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (79.0 mg) and potassium acetate (60.7 mg) in N,N-dimethylformamide (1.50 mL) was added (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (11.2 mg). The reaction mixture was stirred overnight at 80°C under argon atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was
20 washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (100 mg) of N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-
25 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide. To a solution of the obtained crude product (100 mg) in tetrahydrofuran (1.50 mL) was added 2 M aqueous sodium hydroxide solution (0.300 mL) at 0°C. The reaction mixture was stirred at 0°C for 30 min, and 35% aqueous hydrogen peroxide
30 (53.0 µL) was added thereto. The reaction mixture was stirred at room temperature for 1.5 hr, ice water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was
35 evaporated under reduced pressure. The residue was purified by

silica gel chromatography (hexane/ethyl acetate) to give the title compound (43.3 mg).

MS (API+): [M+H]⁺ 557.2.

[0507]

5 H) 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-hydroxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (43.3 mg) in N,N-dimethylformamide (1.00 mL) were added potassium carbonate (10.8 mg) and iodomethane (7.33 µL). The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (28.6 mg).

MS (API+): [M+H]⁺ 571.2.

[0508]

I) 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (28.6 mg) were added trifluoroacetic acid (0.710 mL) and water (79.5 µL). The reaction mixture was stirred at room temperature for 3 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (1.35 mL), and 8 M ammonia/methanol solution (270 µL) was added thereto. The

reaction mixture was stirred at room temperature for 2 hr, and the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was washed with hexane/ethyl acetate to give the title compound (18.8 mg).

MS (API+): [M+H]⁺ 441.2.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.54-3.68 (2H, m), 3.83 (3H, s), 4.30-4.49 (2H, m), 4.98-5.11 (1H, m), 6.96 (1H, d, J = 2.6 Hz), 7.28-7.36 (2H, m), 7.40-7.52 (2H, m), 7.76 (1H, d, J = 2.6 Hz), 10.02 (1H, d, J = 7.5 Hz), 10.76 (1H, brs).

[0509]

Example 64

7-methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (200 mg) was resolved by SFC (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol = 860/140), and crystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (78.3 mg).

MS (API+): [M+H]⁺ 441.2.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.52-3.70 (2H, m), 3.83 (3H, s), 4.26-4.52 (2H, m), 4.97-5.13 (1H, m), 6.96 (1H, d, J = 2.6 Hz), 7.32 (2H, d, J = 8.3 Hz), 7.46 (2H, d, J = 8.7 Hz), 7.76 (1H, d, J = 2.6 Hz), 10.02 (1H, d, J = 7.5 Hz), 10.77 (1H, brs).

[0510]

Example 65

7-methoxy-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

Racemic 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (200 mg) was resolved by SFC

5 (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol = 860/140), and crystallized from hexane/ethyl acetate to give the title compound having a longer retention time (60.0 mg).

10 MS (API+): [M+H]⁺ 441.2.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.53-3.67 (2H, m), 3.83 (3H, s), 4.27-4.51 (2H, m), 4.96-5.16 (1H, m), 6.96 (1H, d, J = 3.0 Hz), 7.32 (2H, d, J = 8.3 Hz), 7.41-7.54 (2H, m), 7.76 (1H, d, J = 2.6 Hz), 10.02 (1H, d, J = 7.2 Hz), 10.76 (1H, 15 brs).

[0511]

Example 66

N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-

20 carboxamide

[0512]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride 25 (10.6 g) and triethylamine (11.8 mL) at room temperature. The reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and 30 saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (11.1 g).

MS (API+): [M+H]⁺ 338.0.

35 [0513]

B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (6.00 g) in tetrahydrofuran (300 mL) was added 5% platinum-carbon (500 mg). The reaction mixture was stirred
5 overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol (200 mL), the solution was heated at reflux for 16 hr, and the solvent was evaporated under reduced pressure. The residue was
10 washed with diisopropyl ether to give the title compound (4.48 g).

MS (API+): [M+H]⁺ 276.0.

[0514]

C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-
15 dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a suspension of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.82 g) in N,N-dimethylformamide (160 mL) was slowly added 1 M potassium
hexamethyldisilazide/tetrahydrofuran solution (17.4 mL) at 0°C.
20 The reaction mixture was stirred at 0°C for 1 hr, 2-(chloromethoxy)ethyltrimethylsilane (3.62 mL) was slowly added thereto, and the mixture was stirred at room temperature for 20 hr. To the reaction mixture was added water, the insoluble substance was filtered off using Celite, and the filtrate was
25 extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the
30 title compound (2.71 g).

MS (API+): [M+H]⁺ 406.0.

[0515]

D) 3-fluoro-N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide

To a solution of 3-fluoro-4-(trifluoromethoxy)benzoic
35 acid (20.0 g), N,O-dimethylhydroxylamine hydrochloride (10.5

g) and triethylamine (24.9 mL) in N,N-dimethylformamide (300 mL) were added 1-hydroxybenzotriazole monohydrate (16.4 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (20.5 g) at room temperature. The reaction mixture was stirred
5 overnight at room temperature, saturated aqueous sodium hydrogen carbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated
10 under reduced pressure to give the title compound (22.9 g).
¹H NMR (300 MHz, CDCl₃) δ 3.38 (3H, s), 3.56 (3H, s), 7.30-7.42 (1H, m), 7.51-7.65 (2H, m).

[0516]

E) 1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethanone

15 To a solution of 3-fluoro-N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide (3.52 g) in tetrahydrofuran (60 mL) was slowly added 1 M methylmagnesium bromide/tetrahydrofuran solution (39.5 mL) at 0°C. The reaction mixture was stirred at room temperature for 4 hr, and
20 poured into saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to
25 give the title compound (2.33 g).
¹H NMR (300 MHz, DMSO-d₆) δ 2.62 (3H, s), 7.68-7.80 (1H, m), 7.86-7.96 (1H, m), 8.04 (1H, dd, J = 11.0, 2.1 Hz).

[0517]

F) 2-bromo-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethanone

30 To a solution of 1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethanone (4.51 g) in acetic acid (50 mL) was slowly added a solution of bromine (1.12 mL) in acetic acid (5 mL) at room temperature. The reaction mixture was stirred at 50°C for 1.5 hr, and the solvent was evaporated
35 under reduced pressure. Aqueous sodium hydrogen carbonate

solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated aqueous sodium hydrogen carbonate solution, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced
5 pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (6.01 g).

^1H NMR (300 MHz, DMSO- d_6) δ 4.99 (2H, s), 7.73-7.84 (1H, m), 7.92-8.00 (1H, m), 8.12 (1H, dd, J = 11.0, 2.0 Hz).

10 [0518]

G) 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanone

To a solution of 2-bromo-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethanone (6.01 g) in methanol (60 mL) were added silver(I) carbonate (7.53 g) and boron trifluoride
15 diethyl ether complex (3.10 mL). The reaction mixture was stirred at 60°C for 4 hr under nitrogen atmosphere, and the insoluble substance was filtered off, and washed with ethyl acetate. The filtrate was concentrated under reduced pressure, and to the residue were added ethyl acetate and brine. The
20 precipitated solid was filtered off, and the organic layer and aqueous layer of the filtrate were separated. The aqueous layer was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
25 column chromatography (hexane/ethyl acetate) to give the title compound (3.57 g).

^1H NMR (300 MHz, DMSO- d_6) δ 3.36 (3H, s), 4.81 (2H, s), 7.70-7.81 (1H, m), 7.83-7.92 (1H, m), 8.01 (1H, dd, J = 10.9, 2.3 Hz).

30 [0519]

H) 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanamine hydrochloride

To a mixture of 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanone (3.57 g) and hydroxylamine hydrochloride
35 (2.01 g) in ethanol (50 mL) was added triethylamine (3.99 mL).

The reaction mixture was stirred at room temperature for 16 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine,
5 and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (3.89 g) of 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-N-hydroxy-2-methoxyethanimine. To a solution of the obtained crude product (3.89 g) in ethanol (60 mL) was added 20% palladium
10 hydroxide-carbon (1.00 g). The reaction mixture was stirred at room temperature for 5 hr under hydrogen atmosphere, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethyl acetate (10 mL), and 4 M hydrogen chloride/ethyl
15 acetate solution (10 mL) was added thereto. The solvent was evaporated under reduced pressure, and the precipitated solid was washed with diisopropyl ether to give the title compound (2.86 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.32 (3H, s), 3.60-3.78 (2H, m),
20 4.53-4.66 (1H, m), 7.46-7.55 (1H, m), 7.62-7.84 (2H, m), 8.70 (3H, brs).

[0520]

I) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-
25 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.00 g) in tetrahydrofuran (37.4 mL) was slowly added a solution of bis(trichloromethyl)carbonate (586
30 mg) in tetrahydrofuran (5.60 mL) at room temperature under nitrogen atmosphere. The reaction mixture was stirred at 40°C for 3 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue was added tetrahydrofuran, and the solvent was evaporated under reduced
35 pressure. The operation (addition of tetrahydrofuran and then

evaporation) was repeated three times. The residue was diluted with tetrahydrofuran (18.7 mL), and the mixture was slowly added to a mixture of 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanamine hydrochloride (858 mg) and triethylamine (1.04 mL) in tetrahydrofuran (12.4 mL) at room temperature. The reaction mixture was stirred overnight at 60°C, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (1.58 g).

MS (API+): [M+H]⁺ 685.1.

[0521]

J) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-hydroxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.58 g), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (1.19 g) and potassium acetate (935 mg) in N,N-dimethylformamide (23.1 mL) was added (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (173 mg). The reaction mixture was stirred overnight at 80°C under argon atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (2.02 g) of N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide. To a solution of the obtained

crude product (2.02 g) in tetrahydrofuran (29.6 mL) was added
2 M aqueous sodium hydroxide solution (5.91 mL) at 0°C. The
reaction mixture was stirred at 0°C for 30 min, and 35% aqueous
hydrogen peroxide (1.04 mL) was added thereto. The reaction
5 mixture was stirred at room temperature for 2 hr, and ice
water was added thereto. The mixture was acidified with 1 M
hydrochloric acid, and extracted with ethyl acetate. The
extract was washed successively with water and saturated brine,
and dried over anhydrous sodium sulfate, and the solvent was
10 evaporated under reduced pressure. The residue was purified by
silica gel chromatography (hexane/ethyl acetate) to give the
title compound (1.24 g).

MS (API+): [M+H]⁺ 575.2.

[0522]

15 K) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-
7-methoxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-(1-(3-fluoro-4-
(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-hydroxy-2-oxo-1-
20 ((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide (1.24 g) in N,N-dimethylformamide
(27.7 mL) were added potassium carbonate (300 mg) and
iodomethane (204 µL). The reaction mixture was stirred at room
temperature for 3 hr, water was added thereto, and the mixture
25 was extracted with ethyl acetate. The extract was washed with
saturated brine, and dried over anhydrous sodium sulfate, and
the solvent was evaporated under reduced pressure. The residue
was purified by silica gel chromatography (hexane/ethyl
acetate) to give the title compound (646 mg).

30 MS (API+): [M+H]⁺ 589.2.

[0523]

L) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-
7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
carboxamide

35 To N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-

methoxyethyl)-7-methoxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (646 mg) were added trifluoroacetic acid (15.8 mL) and water (1.77 mL). The
5 reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (30.1 mL), and 8 M ammonia/methanol solution (5.91 mL) was added thereto. The reaction mixture was stirred at room temperature for 2 hr, and
10 the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The obtained solid was washed with
15 hexane/ethyl acetate to give the title compound (402 mg).
MS (API+): [M+H]⁺ 459.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.29 (3H, s), 3.54-3.70 (2H, m), 3.83 (3H, s), 4.29-4.49 (2H, m), 5.00-5.13 (1H, m), 6.97 (1H, d, J = 2.6 Hz), 7.29 (1H, d, J = 8.7 Hz), 7.41-7.59 (2H, m),
20 7.75 (1H, d, J = 2.6 Hz), 10.01 (1H, d, J = 7.5 Hz), 10.76 (1H, brs).

[0524]

Example 67-I

N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (388 mg) was resolved by SFC
30 (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol = 860/140), and crystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (139 mg) as crystals.
35 MS (API+): [M+H]⁺ 459.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.29 (3H, s), 3.55-3.70 (2H, m), 3.83 (3H, s), 4.28-4.51 (2H, m), 4.99-5.14 (1H, m), 6.97 (1H, d, J = 3.0 Hz), 7.29 (1H, d, J = 8.7 Hz), 7.41-7.59 (2H, m), 7.75 (1H, d, J = 2.6 Hz), 10.02 (1H, d, J = 7.5 Hz), 10.76 (1H, 5 brs).

[0525]

Example 67-II

N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
10 b]pyrazine-4(1H)-carboxamide

[0526]

A) ethyl (5-bromopyridin-3-yl)carbamate

To a mixture of 5-bromopyridin-3-amine (35.4 g) and pyridine (19.8 mL) in tetrahydrofuran (600 mL) was slowly
15 added ethyl chloroformate (23.4 mL) at 0°C. The reaction mixture was stirred overnight at room temperature under nitrogen atmosphere, and poured into water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate,
20 and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether (400 mL), the mixture was stirred at room temperature for 20 min, and the obtained solid was collected by filtration to give the title compound (29.0 g).

25 MS (API+): [M+H]⁺ 245.1.

[0527]

B) ethyl (5-bromo-2-nitropyridin-3-yl)carbamate

To a solution of ethyl (5-bromopyridin-3-yl)carbamate (19.0 g) in conc. sulfuric acid (37.2 mL) was slowly added
30 fuming nitric acid (26.1 mL) at 0°C. The reaction mixture was stirred at room temperature for 72 hr under nitrogen atmosphere. The reaction mixture was slowly poured into ice, the pH of the mixture was adjusted to 9 with 28% aqueous ammonia, and the mixture was extracted with ethyl acetate. The
35 extract was washed with saturated brine, and dried over

anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. To the residue was added ethyl acetate/hexane, the mixture was stirred for 15 min, and the resulting solid was collected by filtration, and washed with
5 hexane to give the title compound (13.9 g).

MS (API+): [M+H]⁺ 290.1.

[0528]

C) 5-methoxy-2-nitropyridin-3-amine

To a solution of ethyl (5-bromo-2-nitropyridin-3-yl)carbamate (31.1 g) in methanol (900 mL) was added 28%
10 sodium methoxide/methanol solution (83.0 g) at room temperature. The reaction mixture was stirred at 65°C for 4 hr under nitrogen atmosphere, and the solvent was evaporated to about 150 mL under reduced pressure. To the residue was added
15 saturated aqueous ammonium chloride solution, the mixture was stirred at room temperature for 20 min, and the solvent was evaporated to about 100 mL under reduced pressure. The resulting solid was collected by filtration, and washed with water to give the title compound (16.6 g).

20 MS (API+): [M+H]⁺ 170.2.

[0529]

D) 2-chloro-N-(5-methoxy-2-nitropyridin-3-yl)acetamide

To a solution of 5-methoxy-2-nitropyridin-3-amine (24.7 g) in N,N-dimethylformamide (740 mL) was slowly added
25 chloroacetyl chloride (23.2 mL) in tetrahydrofuran (40 mL) solution at 0°C. The reaction mixture was stirred at room temperature for 3 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue was added saturated aqueous sodium hydrogen carbonate solution,
30 and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether/hexane, and the mixture was stirred at room temperature
35 for 30 min. The obtained solid was collected by filtration,

and washed with diisopropyl ether/hexane to give the title compound (34.1 g).

MS (API+): [M+H]⁺ 246.0.

[0530]

5 E) N-(2-amino-5-methoxypyridin-3-yl)-2-chloroacetamide

A mixture of 2-chloro-N-(5-methoxy-2-nitropyridin-3-yl)acetamide (34.1 g), ammonium chloride (44.6 g), iron (27.1 g), ethanol (823 mL) and water (206 mL) was stirred 75°C for 40 min, and the solvent was evaporated under reduced pressure. To
10 the residue were added tetrahydrofuran (500 mL) and saturated aqueous sodium hydrogen carbonate solution (300 mL), and the reaction mixture was stirred for 15 min. The insoluble substance was filtered off through Celite, and to the filtrate was added saturated brine, and the mixture was extracted with
15 ethyl acetate. The extract was washed with saturated brine, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether/hexane, and the obtained solid was collected by filtration, and washed with diisopropyl ether/hexane to give the title compound (26.3 g).

20 MS (API+): [M+H]⁺ 216.1.

[0531]

F) benzyl (3-((chloroacetyl)amino)-5-methoxypyridin-2-yl)carbamate

To a mixture of N-(2-amino-5-methoxypyridin-3-yl)-2-
25 chloroacetamide (2.74 g) and pyridine (5.14 mL) in tetrahydrofuran (85 mL) was slowly added benzyl chloroformate (2.72 mL) at 0°C. The reaction mixture was stirred at room temperature for 1 hr, ethyl acetate was added thereto, and the mixture was washed with a mixture of aqueous sodium hydrogen
30 carbonate solution and brine. The organic layer was again washed with saturated brine, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and the obtained solid was washed with diisopropyl ether to give the title
35 compound (3.79 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.82 (3H, s), 4.36 (2H, s), 5.13 (2H, s), 7.30-7.43 (5H, m), 7.87 (1H, d, J = 3.0 Hz), 7.95 (1H, d, J = 2.6 Hz), 9.46 (1H, s), 9.64 (1H, s).

[0532]

- 5 G) benzyl 7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

To a solution of benzyl (3-((chloroacetyl)amino)-5-methoxypyridin-2-yl)carbamate (200 mg) in N,N-dimethylformamide (11 mL) was added cesium carbonate (279 mg) at room temperature, and reaction mixture was stirred at 50°C for 30 min. The reaction mixture was diluted with ethyl acetate, and washed with saturated brine, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (115 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 3.81 (3H, s), 4.36 (2H, s), 5.20 (2H, s), 6.92 (1H, d, J = 2.6 Hz), 7.27-7.43 (5H, m), 7.84 (1H, d, J = 2.6 Hz), 10.72 (1H, s).

[0533]

- 20 H) 7-methoxy-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of benzyl 7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (0.722 g) in tetrahydrofuran (46.1 mL) was added 10% palladium-carbon (containing 50% water, 0.049 g), and the reaction mixture was stirred at room temperature for 15 hr under hydrogen atmosphere. To the reaction mixture was added methanol at 50°C to dissolve the precipitated solid, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The obtained solid was washed with diisopropyl ether to give the title compound (0.380 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.68 (3H, s), 3.82 (2H, d, J = 1.9 Hz), 6.32 (1H, s), 6.66 (1H, d, J = 2.6 Hz), 7.37 (1H, d, J = 2.6 Hz), 10.32 (1H, s).

[0534]

- 35 I) N,2-dimethoxy-N-methylacetamide

To a mixture of N,O-dimethylhydroxylamine hydrochloride (24.7 g) and potassium carbonate (63.7 g) in acetonitrile (330 mL) was added 2-methoxyacetyl chloride (25.0 g) at the temperature of 10°C or lower. The reaction mixture was stirred overnight at room temperature, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was distilled (boiling point: 60°C/0.8 KPa) to give the title compound (25.8 g).

¹H NMR (300 MHz, CDCl₃) δ 3.20 (3H, s), 3.47 (3H, d, J = 0.8 Hz), 3.69 (3H, s), 4.22 (2H, s).

[0535]

J) 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanone

To a mixture of 4-bromo-2-fluoro-1-(trifluoromethoxy)benzene (25.0 g) and N,2-dimethoxy-N-methylacetamide (15.4 g) in tetrahydrofuran (400 mL) was slowly added 1.6 M n-butyllithium/hexane solution (72.4 mL) at -78°C under nitrogen atmosphere. The reaction mixture was stirred at -78°C for 20 min under nitrogen atmosphere, and neutralized with 0.1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (15.7 g).

¹H NMR (300 MHz, CDCl₃) δ 3.50 (3H, s), 4.63 (2H, s), 7.42 (1H, ddd, J = 8.6, 7.3, 1.5 Hz), 7.75-7.81 (1H, m), 7.83 (1H, dd, J = 10.2, 1.9 Hz).

[0536]

K) 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-N-hydroxy-2-methoxyethanimine

To a solution of 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanone (7.56 g) in ethanol (250 mL) were added hydroxylamine hydrochloride (4.17 g) and triethylamine (8.36 mL) at room temperature. The reaction mixture was stirred at

room temperature for 16 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried
5 over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (7.88 g).

MS (API+): [M+H]⁺ 268.0.

10 [0537]

L) tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)carbamate

To a solution of 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-N-hydroxy-2-methoxyethanimine (7.88 g) in ethanol (200 mL) was
15 added 10% palladium-carbon (containing 50% water, 1.00 g). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. To a tetrahydrofuran solution (200 mL) of the
20 residue were added di-tert-butyl dicarbonate (7.53 mL) and triethylamine (6.17 mL), and the mixture was stirred at room temperature for 16 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous
25 magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.25 g).

¹H NMR (300 MHz, CDCl₃) δ 1.42 (9H, brs), 3.35 (3H, s), 3.49-
30 3.67 (2H, m), 4.78 (1H, brs), 5.34 (1H, brs), 7.08-7.29 (3H, m).

[0538]

M) tert-butyl ((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)carbamate

35 Racemic tert-butyl (1-(3-fluoro-4-

(trifluoromethoxy)phenyl)-2-methoxyethyl)carbamate (12.48 g) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 950/50) to give the title compound
5 having a shorter retention time (5.73 g).

MS (API+), found: 254.0.

[0539]

N) (1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanamine hydrochloride

10 A mixture of tert-butyl ((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)carbamate (5.73 g) and 4 M hydrogen chloride/ethyl acetate solution (100 mL) was stirred at room temperature for 3 hr. The solvent was evaporated under reduced pressure, and the residue was washed
15 with diisopropyl ether to give the title compound (3.22 g).

MS (API+), found: 254.0.

[0540]

O) N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
20 b]pyrazine-4(1H)-carboxamide

To a mixture of 7-methoxy-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (2.49 g) and diisopropylethylamine (7.13 mL) in tetrahydrofuran (120 mL) was slowly added 4-nitrophenyl chloroformate (3.64 g) at 0°C. The reaction mixture was
25 stirred at room temperature for 3 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue were added diisopropyl ether (200 mL) and saturated brine (150 mL), and the mixture was stirred at room temperature for 20 min. The resulting solid was collected by
30 filtration, washed with water and diisopropyl ether to give 4-nitrophenyl 7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (4.20 g). A mixture of 4-nitrophenyl 7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (4.20 g), diisopropylethylamine (11.9 mL) and (S)-
35 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanamine

hydrochloride (4.43 g) in N,N-dimethylformamide (130 mL) was stirred at room temperature for 16 hr, and the solvent was evaporated under reduced pressure. To the residue was added aqueous sodium hydrogen carbonate solution, and the mixture
5 was extracted with a mixed solvent of ethyl acetate and tetrahydrofuran. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (NH, hexane/ethyl acetate) to give
10 (S)-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (5.01 g). To a solution of the obtained (S)-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
15 b]pyrazine-4(1H)-carboxamide (5.00 g) in ethanol (200 mL) was added activated carbon (shirasagi) (10 g), and the mixture was stirred at room temperature for 30 min. The insoluble substance was filtered off, and washed with acetone. The filtrate was concentrated under reduced pressure to give (S)-
20 N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (4.17 g). To a solution of the obtained (S)-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
25 carboxamide (4.17 g) in acetone (35 mL) was slowly added heptane (70 mL) at 52°C. The mixture was stirred at 52°C for 1 hr, heptane (35 mL) was slowly added thereto at 52°C, and the mixture was gradually cooled to room temperature. The mixture was stirred overnight at room temperature, and then at 5°C for
30 1 hr. The resulting solid was collected by filtration, and washed with acetone/heptane to give the title compound (3.15 g) as crystals.

MS (API+): [M+H]⁺ 459.2.

¹H NMR (300 MHz, DMSO-d₆) δ 3.29 (3H, s), 3.56-3.70 (2H, m),
35 3.83 (3H, s), 4.30-4.52 (2H, m), 4.95-5.16 (1H, m), 6.96 (1H,

d, J = 2.6 Hz), 7.29 (1H, d, J = 8.3 Hz), 7.43-7.57 (2H, m), 7.75 (1H, d, J = 3.0 Hz), 10.02 (1H, d, J = 7.2 Hz), 10.78 (1H, brs).

[0541]

5 Example 68

N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (388 mg) was resolved by SFC (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol = 860/140) to give the title compound having
15 a longer retention time (145 mg).

MS (API+): [M+H]⁺ 459.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.29 (3H, s), 3.57-3.69 (2H, m), 3.83 (3H, s), 4.29-4.50 (2H, m), 4.97-5.15 (1H, m), 6.97 (1H, d, J = 2.6 Hz), 7.29 (1H, d, J = 8.7 Hz), 7.41-7.59 (2H, m),
20 7.75 (1H, d, J = 2.6 Hz), 10.02 (1H, d, J = 7.5 Hz), 10.76 (1H, brs).

[0542]

Example 69

7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0543]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride
30 (10.6 g) and triethylamine (11.8 mL) at room temperature. The reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and
35 saturated brine, and dried over anhydrous magnesium sulfate,

and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (11.1 g).

MS (API+): $[M+H]^+$ 338.0.

5 [0544]

B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (6.00 g) in tetrahydrofuran (300 mL) was added 5% platinum-carbon (500 mg). The reaction mixture was stirred
10 overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol (200 mL), the mixture was heated at reflux for 16 hr, and the solvent was evaporated under reduced pressure. The residue was
15 washed with diisopropyl ether to give the title compound (4.48 g).

MS (API+): $[M+H]^+$ 276.0.

[0545]

C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-
20 dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a suspension of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.82 g) in N,N-dimethylformamide (160 mL) was slowly added 1 M potassium
hexamethyldisilazide/tetrahydrofuran solution (17.4 mL) at 0°C.
25 The reaction mixture was stirred at 0°C for 1 hr, 2-(chloromethoxy)ethyltrimethylsilane (3.62 mL) was slowly added thereto, and the mixture was stirred at room temperature for 20 hr. To the reaction mixture was added water, the insoluble substance was filtered off using Celite, and the filtrate was
30 extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the
35 title compound (2.71 g).

MS (API+): [M+H]⁺ 406.0.

[0546]

D) 2-methoxy-1-(4-(trifluoromethyl)phenyl)ethanone

To a solution of 2-bromo-1-(4-(trifluoromethyl)phenyl)ethanone (2.50 g) in methanol (50 mL) were added silver(I) carbonate (2.99 g) and boron trifluoride diethyl ether complex (1.42 mL) at room temperature. The reaction mixture was stirred overnight at 50°C, the insoluble substance was filtered off, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.84 g).

¹H NMR (300 MHz, CDCl₃) δ 3.51 (3H, s), 4.70 (2H, s), 7.75 (2H, d, J = 8.3 Hz), 8.06 (2H, d, J = 7.9 Hz).

[0547]

E) 2-methoxy-1-(4-(trifluoromethyl)phenyl)ethanamine hydrochloride

To a solution of 2-methoxy-1-(4-(trifluoromethyl)phenyl)ethanone (1.84 g) in ethanol (100 mL) were added hydroxylamine hydrochloride (1.17 g) and triethylamine (2.35 mL) at room temperature. The reaction mixture was stirred at room temperature for 4.5 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a crude product of N-hydroxy-2-methoxy-1-(4-(trifluoromethyl)phenyl)ethanimine. To a solution of the obtained crude product in ethanol (100 mL) was added 10% palladium-carbon (containing 50% water, 200 mg). The reaction mixture was stirred overnight at room

temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. A solution of the residue in ethyl acetate (5 mL) was added to 4 M hydrogen chloride/ethyl acetate solution (20 mL). The reaction mixture was stirred at room temperature for 30 min, and the solvent was evaporated under reduced pressure. The residue was crystallized from ethyl acetate to give the title compound (1.81 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.26-3.41 (3H, m), 3.59-3.78 (2H, m), 4.58-4.73 (1H, m), 7.66-7.91 (4H, m), 8.69 (3H, brs).

[0548]

F) 7-iodo-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.53 g) in tetrahydrofuran (60 mL) was added a solution of bis(trichloromethyl)carbonate (0.896 g) in tetrahydrofuran (8.0 mL) at room temperature. The reaction mixture was stirred at 40°C for 5 hr, and the solvent was evaporated under reduced pressure. The residue was diluted with tetrahydrofuran (10.0 mL), and the mixture was added to a mixture of 2-methoxy-1-(4-(trifluoromethyl)phenyl)ethanamine hydrochloride (1.16 g) and triethylamine (1.58 mL) in tetrahydrofuran (60 mL) at room temperature. The reaction mixture was stirred at 60°C for 16 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (2.18 g).

MS (API+): [M+H]⁺ 651.1.

[0549]

G) 7-hydroxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-

2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.19 g), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (941 mg), potassium acetate (742 mg) in N,N-dimethylformamide (18.3 mL) was added (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (137 mg). The reaction mixture was stirred overnight at 80°C under argon atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (1.95 g) of N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide. To a solution of the obtained crude product (1.19 g) in tetrahydrofuran (18.3 mL) was added 2 M aqueous sodium hydroxide solution (3.67 mL) at 0°C. The reaction mixture was stirred at 0°C for 30 min, and 35% aqueous hydrogen peroxide (642 µL) was added thereto. The reaction mixture was stirred at room temperature for 2 hr, ice water was added thereto, the mixture was acidified with 1 M hydrochloric acid, and extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (796 mg).

MS (API+): [M+H]⁺ 541.2.

[0550]

H) 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-

dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-hydroxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (796 mg) in N,N-dimethylformamide (18.9 mL) were added potassium carbonate (205 mg) and iodomethane (139 μ L). The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (665 mg). MS (API+): [M+H]⁺ 555.2.

[0551]

I) 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (665 mg) were added trifluoroacetic acid (17.3 mL) and water (1.93 mL). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (32.9 mL), and 8 M ammonia/methanol solution (6.46 mL) was added thereto. The reaction mixture was stirred at room temperature for 10 min, and the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (474 mg). MS (API+): [M+H]⁺ 425.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.59-3.68 (2H, m),

3.83 (3H, s), 4.25-4.55 (2H, m), 5.00-5.17 (1H, m), 6.97 (1H, d, $J = 3.0$ Hz), 7.56 (2H, d, $J = 8.3$ Hz), 7.64-7.73 (2H, m), 7.77 (1H, d, $J = 2.6$ Hz), 10.07 (1H, d, $J = 7.5$ Hz), 10.77 (1H, s).

5 [0552]

Example 70

optically active 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 Racemic 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (444 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol =
15 600/400), and crystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (197 mg).
MS (API+): $[M+H]^+$ 425.1.

^1H NMR (300 MHz, DMSO- d_6) δ 3.28 (3H, s), 3.61-3.68 (2H, m), 3.83 (3H, s), 4.29-4.50 (2H, m), 5.03-5.16 (1H, m), 6.97 (1H,
20 d, $J = 2.6$ Hz), 7.56 (2H, d, $J = 8.3$ Hz), 7.69 (2H, d, $J = 8.3$ Hz), 7.77 (1H, d, $J = 2.6$ Hz), 10.07 (1H, d, $J = 7.2$ Hz), 10.77 (1H, brs).

[0553]

Example 71

25 optically active 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (444 mg) was resolved by HPLC
30 (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 600/400), and crystallized from hexane/ethyl acetate to give the title compound having a longer retention time (189 mg).
35 MS (API+): $[M+H]^+$ 425.1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.61-3.67 (2H, m), 3.83 (3H, s), 4.30-4.50 (2H, m), 5.04-5.16 (1H, m), 6.97 (1H, d, J = 2.6 Hz), 7.56 (2H, d, J = 8.3 Hz), 7.69 (2H, d, J = 8.3 Hz), 7.77 (1H, d, J = 2.6 Hz), 10.07 (1H, d, J = 7.5 Hz),
5 10.77 (1H, s).

[0554]

Example 72-I

7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
10 b]pyrazine-4(1H)-carboxamide

[0555]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride
15 (10.6 g) and triethylamine (11.8 mL) at room temperature. The reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and
20 saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (11.1 g).

MS (API+): [M+H]⁺ 338.0.

25 [0556]

B) methyl N-(5-cyclopropyl-3-nitropyridin-2-yl)glycinate

A mixture of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (1.00 g), cyclopropylboronic acid (0.510 g), palladium(II) acetate (0.100 g), tricyclohexylphosphine (0.250
30 g) and tripotassium phosphate (1.89 g) in toluene (30 mL) was stirred at 90°C for 16 hr under argon atmosphere, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (0.130 g).

35 MS (API+): [M+H]⁺ 252.1.

[0557]

C) 7-cyclopropyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a mixture of methyl N-(5-cyclopropyl-3-nitropyridin-2-yl)glycinate (73 mg) in a mixed solvent of ethanol (10 mL) and
5 tetrahydrofuran (2 mL) was added 10% palladium-carbon
(containing 50% water, 10 mg). The reaction mixture was
stirred at room temperature for 2.5 hr under hydrogen
atmosphere, the catalyst was filtered off, and the filtrate
was concentrated under reduced pressure. The residue (64 mg)
10 was dissolved in ethanol (8 mL), the reaction mixture was
heated at reflux for 16 hr, and the solvent was evaporated
under reduced pressure. The residue was washed with ethyl
acetate to give the title compound (39.9 mg).

MS (API+): [M+H]⁺ 190.1.

15 [0558]

D) 5-cyclopropyl-3-nitropyridin-2-amine

A mixture of 5-bromo-3-nitropyridin-2-amine (2.27 g),
cyclopropylboronic acid (1.79 g), palladium(II) acetate (0.117
g), tricyclohexylphosphine (0.292 g), tripotassium phosphate
20 (5.53 g), toluene (30 mL), 1,2-dimethoxyethane (10 mL) and
water (10 mL) was stirred overnight at 100°C under nitrogen
atmosphere, and the insoluble substance was filtered off using
Celite. The filtrate was washed successively with water and
saturated brine, and dried over anhydrous magnesium sulfate,
25 and the solvent was evaporated under reduced pressure. The
residue was purified by silica gel column chromatography
(hexane/ethyl acetate) to give the title compound (2.04 g).

MS (API+): [M+H]⁺ 180.2.

[0559]

30 E) 7-cyclopropylpyrido[2,3-b]pyrazin-2(1H)-one

To a solution of 5-cyclopropyl-3-nitropyridin-2-amine
(6.89 g) in ethanol (300 mL) was added 10% palladium-carbon
(containing 50% water, 1.64 g). The reaction mixture was
stirred at room temperature for 5 hr under hydrogen atmosphere,
35 the catalyst was filtered off, and the filtrate was

concentrated under reduced pressure. The residue was suspended in water (150 mL), 9 M aqueous glyoxylic acid solution (6.41 mL) was added thereto at room temperature. The reaction mixture was stirred overnight at room temperature, and the
5 resulting solid was collected by filtration, and washed with water and diisopropyl ether to give the title compound (4.52 g).

[0560]

F) 7-cyclopropyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

10 To a solution of 7-cyclopropylpyrido[2,3-b]pyrazin-2(1H)-one (4.52 g) in ethanol (150 mL) was added sodium borohydride (1.83 g) at room temperature. The reaction mixture was stirred at 32°C for 1 hr, and concentrated to about half of the volume under reduced pressure. To the residue was added water, and
15 the resulting solid was collected by filtration, and washed with diisopropyl ether to give the title compound (3.45 g).

MS (API+): [M+H]⁺190.2.

[0561]

G) 7-cyclopropyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

20 To a solution of 7-cyclopropylpyrido[2,3-b]pyrazin-2(1H)-one (40.0 mg) in methanol (30 mL) was added 20% palladium hydroxide-carbon (15.0 mg). The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated
25 under reduced pressure to give the title compound (30.0 mg).

MS (API+): [M+H]⁺190.2.

[0562]

H) 4-nitrophenyl 7-cyclopropyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

30 To a mixture of 7-cyclopropyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (35.0 mg) in a mixed solvent of N,N-dimethylacetamide (1.5 mL) and pyridine (0.3 mL) was added 4-nitrophenyl chloroformate (44.7 mg). The reaction mixture was stirred at room temperature for 2 hr, water was added thereto,
35 and the mixture was extracted with ethyl acetate. The extract

was washed successively with 1 M hydrochloric acid, water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the
5 title compound (24.0 mg).

MS (API+): $[M+H]^+$ 355.1.

[0563]

I) methyl 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetate

10 To a mixture of 4-(trifluoromethoxy)benzaldehyde (19.0 g) and ammonium carbonate (25.9 g) in a mixed solvent of ethanol (114 mL) and water (45.6 mL) was slowly added an aqueous solution (71.1 mL) of potassium cyanide (8.14 g) at 50°C. The reaction mixture was stirred at 60°C for 3 hr, and cooled to
15 room temperature, and the ethanol was evaporated under reduced pressure. The pH of the residue was adjusted to 1 with conc. hydrochloric acid at 0°C, and the resulting solid was filtered off, and washed with water. To an aqueous solution (100 mL) of potassium hydroxide (23.6 g) was added the solid obtained by
20 the above-mentioned operation at room temperature, and the reaction mixture was stirred at 90°C for 3 days. The reaction mixture was cooled to room temperature, and neutralized with conc. hydrochloric acid. The resulting solid was filtered off, and washed with water to give a crude product (13.3 g) of 2-
25 amino-2-(4-(trifluoromethoxy)phenyl)acetic acid. To a solution of the obtained crude product (13.3 g) in tetrahydrofuran (113 mL) were added di-tert-butyl dicarbonate (19.7 mL) and 2 M aqueous sodium hydroxide solution (85 mL) at room temperature. The reaction mixture was stirred overnight at room temperature,
30 and poured into water, and the mixture was washed with diethyl ether. The pH of the aqueous layer was adjusted to 3 with 1 M hydrochloric acid at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent
35 was evaporated under reduced pressure to give a crude product

(11.3 g) of 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetic acid. To a solution of the obtained crude product (11.3 g) in N,N-dimethylformamide (84 mL) were added methyl iodide (2.53 mL) and potassium carbonate (5.59 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and poured into water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.20 g).

MS (API-): [M-H]⁻ 348.1.

[0564]

J) tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

To a solution of methyl 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetate (5.00 g) in tetrahydrofuran (71.6 mL) was slowly added 1 M methylmagnesium bromide/tetrahydrofuran solution (57.3 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr under argon atmosphere, saturated aqueous ammonium chloride solution was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.99 g).

MS (API-): [M-H]⁻ 348.2.

[0565]

K) 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride

To tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate (2.50 g) was added 4 M hydrogen chloride/ethyl acetate solution (71.6 mL). The

reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether, and the resulting crystals were collected by filtration to give the title
5 compound (2.01 g).

MS (API+), found: 250.1.

[0566]

L) 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
10 b]pyrazine-4(1H)-carboxamide

To a solution of 4-nitrophenyl 7-cyclopropyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (84 mg) in N,N-dimethylformamide (5 mL) were added 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride (88 mg) and
15 triethylamine (0.099 mL) at room temperature. The reaction mixture was stirred at room temperature for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate,
20 and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and recrystallized from hexane/ethyl acetate to give the title compound (52 mg).

MS (API+): [M+H]⁺ 465.2.

25 ¹H NMR (300 MHz, DMSO-d₆) δ 0.61-0.71 (2H, m), 0.94-1.05 (5H, m), 1.17 (3H, s), 1.91-2.04 (1H, m), 4.25-4.46 (2H, m), 4.68 (1H, d, J = 8.3 Hz), 4.76 (1H, s), 6.94 (1H, d, J = 2.3 Hz), 7.22-7.31 (2H, m), 7.38-7.48 (2H, m), 7.87 (1H, d, J = 2.3 Hz), 10.50 (1H, d, J = 8.3 Hz), 10.67 (1H, brs).

30 [0567]

Example 72-II

7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

35 [0568]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride (10.6 g) and triethylamine (11.8 mL) at room temperature. The
5 reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate,
10 and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (11.1 g).

MS (API+): $[M+H]^+$ 338.0.

[0569]

15 B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (6.00 g) in tetrahydrofuran (300 mL) was added 5% platinum-carbon (500 mg). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the
20 catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol (200 mL), the reaction mixture was heated at reflux for 16 hr, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title
25 compound (4.48 g).

MS (API+): $[M+H]^+$ 276.0.

[0570]

C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

30 To a suspension of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.82 g) in N,N-dimethylformamide (160 mL) was slowly added 1 M potassium hexamethyldisilazide/tetrahydrofuran solution (17.4 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, 2-
35 (chloromethoxy)ethyltrimethylsilane (3.62 mL) was slowly added

thereto, and the mixture was stirred at room temperature for 20 hr. To the reaction mixture was added water, the insoluble substance was filtered off using Celite, and the filtrate was extracted with ethyl acetate. The extract was washed
5 successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (2.71 g).

10 MS (API+): $[M+H]^+$ 406.0.
[0571]

D) methyl 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetate

To a mixture of 4-trifluoromethoxybenzaldehyde (19.0 g)
15 and ammonium carbonate (25.9 g) in a mixed solvent of ethanol (114 mL) and water (45.6 mL) was slowly added an aqueous solution (71.1 mL) of potassium cyanide (8.14 g) at 50°C. The reaction mixture was stirred at 60°C for 3 hr, and cooled to room temperature, and the ethanol was evaporated under reduced
20 pressure. The pH of the residue was adjusted to 1 with conc. hydrochloric acid at 0°C, and the resulting solid was filtered off, and washed with water. To an aqueous solution (100 mL) of potassium hydroxide (23.6 g) was added the solid obtained by the above-mentioned operation at room temperature, the
25 reaction mixture was stirred at 90°C for 3 days. The reaction mixture was cooled to room temperature, and neutralized with conc. hydrochloric acid. The resulting solid was collected by filtration, and washed with water to give a crude product (13.3 g) of 2-amino-2-(4-(trifluoromethoxy)phenyl)acetic acid.
30 To a solution of the obtained crude product (13.3 g) in tetrahydrofuran (113 mL) were added di-tert-butyl dicarbonate (19.7 mL) and 2 M aqueous sodium hydroxide solution (85 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, and poured into water, and the mixture
35 was washed with diethyl ether. The pH of the aqueous layer was

adjusted to 3 with 1 M hydrochloric acid at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (11.3 g) of 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetic acid. To a solution of the obtained crude product (11.3 g) in N,N-dimethylformamide (84 mL) were added methyl iodide (2.53 mL) and potassium carbonate (5.59 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and poured into water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.20 g).

MS (API-): [M-H]⁻ 348.1.

[0572]

E) tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

To a solution of methyl 2-((tert-butoxycarbonyl)amino)-2-(4-(trifluoromethoxy)phenyl)acetate (5.00 g) in tetrahydrofuran (71.6 mL) was slowly added 1 M methylmagnesium bromide/tetrahydrofuran solution (57.3 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr under argon atmosphere, saturated aqueous ammonium chloride solution was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.99 g).

MS (API-): [M-H]⁻ 348.2.

[0573]

F) 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol

hydrochloride

To tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate (2.50 g) was added 4 M hydrogen chloride/ethyl acetate solution (71.6 mL). The
5 reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether, and the resulting crystals were collected by filtration to give the title compound (2.01 g).

10 MS (API+), found: 250.1.
[0574]

G) N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-
15 b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.40 g) in tetrahydrofuran (120 mL) was slowly added a solution of bis(trichloromethyl)carbonate (1.99
20 g) in tetrahydrofuran (20 mL) at room temperature under nitrogen atmosphere. The reaction mixture was stirred at 40°C for 5 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue was added tetrahydrofuran, and the solvent was evaporated under reduced
25 pressure. The operation (addition of tetrahydrofuran and then evaporation) was repeated three times. The residue was diluted with tetrahydrofuran (120 mL), and slowly added to a mixture of 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride (2.64 g) and triethylamine (3.51 mL) in
30 tetrahydrofuran (20 mL) at room temperature. The reaction mixture was stirred overnight at 60°C, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under
35 reduced pressure. The residue was purified by silica gel

chromatography (NH, hexane/ethyl acetate) to give the title compound (5.32 g).

MS (API+): [M+H]⁺ 681.1.

[0575]

5 H) 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A mixture of N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (5.32 g), cyclopropylboronic acid (1.34 g), palladium(II) acetate (0.176 g), tricyclohexylphosphine (0.438 g) and tripotassium phosphate
10 (4.98 g) in a mixed solvent of toluene (150 mL) and water (6.01 mL) was stirred under argon atmosphere at 80°C for 3 hr, at 100°C for 3 hr, at room temperature for 3 days, and at 100°C for 2.5 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed
15 successively with water and saturated brine, the aqueous layers were combined, and extracted with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl
20 acetate) to give the title compound (3.80 g).

MS (API+): [M+H]⁺ 595.3.

[0576]

I) 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
30

To 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (3.80 g) were slowly added
35 trifluoroacetic acid (120 mL) and water (14 mL). The reaction

mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (220 mL), and 7 M ammonia/methanol solution (18.3 mL) was slowly added thereto.
5 The reaction mixture was stirred at room temperature for 2 hr, ethyl acetate was added thereto, and the mixture was washed successively with water and saturated brine. The aqueous layer was neutralized with 1 M hydrochloric acid, and extracted with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure.
10 The residue was dissolved in ethyl acetate, and the solution was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give the title compound
15 (2.65 g).

MS (API+): [M+H]⁺ 465.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.60-0.71 (2H, m), 0.94-1.04 (5H, m), 1.18 (3H, s), 1.90-2.05 (1H, m), 4.25-4.47 (2H, m), 4.68 (1H, d, J = 8.3 Hz), 4.78 (1H, s), 6.94 (1H, d, J = 2.3 Hz),
20 7.23-7.33 (2H, m), 7.38-7.48 (2H, m), 7.88 (1H, d, J = 2.3 Hz), 10.51 (1H, d, J = 7.9 Hz), 10.68 (1H, brs).

[0577]

Example 73

7-cyclopropyl-N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (2.65 g) was resolved by HPLC
30 (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 850/150), and recrystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (1.01 g).

MS (API+): [M+H]⁺ 465.1.

35 ¹H NMR (300 MHz, DMSO-d₆) δ 0.62-0.71 (2H, m), 0.95-1.05 (5H,

m), 1.17 (3H, s), 1.91-2.04 (1H, m), 4.25-4.47 (2H, m), 4.68 (1H, d, J = 7.9 Hz), 4.77 (1H, s), 6.94 (1H, d, J = 1.9 Hz), 7.22-7.32 (2H, m), 7.38-7.47 (2H, m), 7.87 (1H, d, J = 2.3 Hz), 10.50 (1H, d, J = 8.3 Hz), 10.67 (1H, s).

5 [0578]

Example 74

7-cyclopropyl-N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 Racemic 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (2.65 g) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol =
15 850/150), and recrystallized from hexane/ethyl acetate to give the title compound having a longer retention time (0.792 g).
MS (API+): [M+H]⁺ 465.1.
¹H NMR (300 MHz, DMSO-d₆) δ 0.62-0.71 (2H, m), 0.94-1.05 (5H, m), 1.17 (3H, s), 1.91-2.05 (1H, m), 4.25-4.46 (2H, m), 4.68
20 (1H, d, J = 8.3 Hz), 4.77 (1H, s), 6.94 (1H, d, J = 1.9 Hz), 7.23-7.32 (2H, m), 7.38-7.47 (2H, m), 7.87 (1H, d, J = 2.3 Hz), 10.50 (1H, d, J = 8.3 Hz), 10.67 (1H, s).

[0579]

Example 75-I

25 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0580]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

30 To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride (10.6 g) and triethylamine (11.8 mL) at room temperature. The reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue
35 was added water, and the mixture was extracted with ethyl

acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the
5 title compound (11.1 g).

MS (API+): $[M+H]^+$ 338.0.

[0581]

B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (6.00 g) in tetrahydrofuran (300 mL) was added 5%
10 platinum-carbon (500 mg). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol
15 (200 mL), the solution was heated at reflux for 16 hr, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (4.48 g).

MS (API+): $[M+H]^+$ 276.0.

20 [0582]

C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a suspension of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.82 g) in N,N-dimethylformamide (160 mL)
25 was slowly added 1 M potassium hexamethyldisilazide/tetrahydrofuran solution (17.4 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, 2-(chloromethoxy)ethyltrimethylsilane (3.62 mL) was slowly added thereto, and the mixture was stirred at room temperature for
30 20 hr. To the reaction mixture was added water, the insoluble substance was filtered off using Celite, and the filtrate was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated
35 under reduced pressure. The residue was purified by silica gel

column chromatography (NH, hexane/ethyl acetate) to give the title compound (2.71 g).

MS (API+): $[M+H]^+$ 406.0.

[0583]

5 D) 3-fluoro-N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide

To a solution of 3-fluoro-4-(trifluoromethoxy)benzoic acid (20 g), N,O-dimethylhydroxylamine hydrochloride (10.5 g) and triethylamine (24.9 mL) in N,N-dimethylformamide (300 mL) were added 1-hydroxybenzotriazole monohydrate (16.4 g) and 1-
10 (3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (20.5 g) at room temperature. The reaction mixture was stirred overnight at room temperature, and then at 50°C for 1.5 hr. To the reaction mixture was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted
15 with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. Toluene was added to the residue because the residue contained a small amount of N,N-dimethylformamide, and the solvent was
20 evaporated under reduced pressure to give the title compound (22.0 g).

MS (API+): $[M+H]^+$ 268.1.

[0584]

E) 3-fluoro-4-(trifluoromethoxy)benzaldehyde

25 To a solution of 3-fluoro-N-methoxy-N-methyl-4-(trifluoromethoxy)benzamide (10.0 g) in tetrahydrofuran (101 mL) was added 1.5 M diisobutylaluminum hydride/toluene solution (27.4 mL) at 0°C. The reaction mixture was stirred overnight at room temperature, saturated aqueous ammonium
30 chloride solution and 6 M hydrochloric acid was successively added, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
35 column chromatography (hexane/ethyl acetate) to give the title

compound (3.55 g).

^1H NMR (300 MHz, DMSO- d_6) δ 7.71-7.97 (2H, m), 7.97-8.10 (1H, m), 10.02 (1H, d, J = 1.9 Hz).

[0585]

5 F) 2-amino-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetic acid

To a mixture of 3-fluoro-4-(trifluoromethoxy)benzaldehyde (3.50 g) and ammonium carbonate (4.36 g) in a mixed solvent of ethanol (21 mL) and water (8 mL) was slowly added an aqueous solution (13 mL) of potassium cyanide (1.37 g) at 50°C. The
10 mixture was stirred at 60°C for 4 hr, and cooled to room temperature, and the ethanol was evaporated under reduced pressure. The pH of the residue was adjusted to 1 with conc. hydrochloric acid at 0°C, and the mixture was left overnight at freezer. The resulting solid was collected by filtration, and
15 washed with water. To an aqueous solution (100 mL) of potassium hydroxide (3.96 g) was added the obtained solid at room temperature, and the mixture was stirred at 90°C for 60 hr, cooled to room temperature, and neutralized with conc. hydrochloric acid. The resulting solid was collected by
20 filtration, and washed with water to give the title compound (937 mg).

^1H NMR (300 MHz, DMSO- d_6) δ 4.34 (1H, s), 7.22-7.42 (1H, m), 7.45-7.66 (2H, m).

[0586]

25 G) 2-((tert-butoxycarbonyl) amino)-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetic acid

To a solution of 2-amino-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetic acid (937 mg) in tetrahydrofuran (7.40 mL) were added di-tert-butyl dicarbonate
30 (1.29 mL) and 2 M aqueous sodium hydroxide solution (5.55 mL) at room temperature. The reaction mixture was stirred at room temperature for 3 days, and poured into water, and the mixture was washed with diethyl ether. The pH of the aqueous layer was adjusted to 3 with 1 M hydrochloric acid at 0°C, and the
35 mixture was extracted with ethyl acetate. The extract was

washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (750 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.39 (9H, s), 5.02-5.35 (1H, m),
5 7.15-7.64 (3H, m), 7.64-7.85 (1H, m).

[0587]

H) methyl 2-((tert-butoxycarbonyl) amino)-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetate

To a solution of 2-((tert-butoxycarbonyl) amino)-2-(3-
10 fluoro-4-(trifluoromethoxy)phenyl)acetic acid (750 mg) in N,N-dimethylformamide (5.5 mL) were added methyl iodide (0.159 mL) and potassium carbonate (352 mg) at room temperature. The reaction mixture was stirred at room temperature for 4 hr under nitrogen atmosphere, and poured into water, and the
15 mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title
20 compound (630 mg).

MS (API-): [M-H]⁻ 366.1.

[0588]

I) tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)carbamate

To a solution of methyl 2-((tert-butoxycarbonyl) amino)-
2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetate (630 mg) in tetrahydrofuran (9 mL) was slowly added 1 M methyllmagnesium bromide/tetrahydrofuran solution (6.86 mL) at 0°C under argon atmosphere. The reaction mixture was stirred at 0°C for 4 hr
30 under argon atmosphere, saturated aqueous ammonium chloride solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was
35 purified by silica gel column chromatography (hexane/ethyl

acetate) to give the title compound (540 mg).

MS (API-): [M-H]⁻ 366.2.

[0589]

J) 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methylpropan-2-ol hydrochloride

To tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)carbamate (540 mg) was added 4 M hydrogen chloride/ethyl acetate solution (4 mL). The reaction mixture was stirred at room temperature for 30 min, and the solvent was evaporated under reduced pressure to give the title compound (510 mg).

MS (API+), found: 268.1.

[0590]

K) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (586 mg) in tetrahydrofuran (28 mL) was added bis(trichloromethyl)carbonate (343 mg) at room temperature. The reaction mixture was stirred at 40°C for 5 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue was added tetrahydrofuran, and the solvent was evaporated under reduced pressure. The operation (addition of tetrahydrofuran and then evaporation) was repeated three times. The residue was diluted with tetrahydrofuran (14 mL), and the mixture was slowly added to a solution of 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methylpropan-2-ol hydrochloride (439 mg) and triethylamine (0.604 mL) in tetrahydrofuran (9 mL) at room temperature. The reaction mixture was stirred at 60°C for 16 hr, ethyl acetate was added thereto, washed successively with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent

was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (590 mg).

MS (API+): [M+H]⁺ 699.1.

5 [0591]

L) 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 To a mixture of N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (590 mg), cyclopropylboronic acid (145 mg) and tripotassium phosphate
15 (628 mg) in a mixed solvent of toluene (16.2 mL) and water (0.648 mL) were added palladium(II) acetate (37.9 mg) and tricyclohexylphosphine (95 mg) at room temperature. The reaction mixture was stirred overnight at 100°C under argon atmosphere, the insoluble substance was filtered off, and the
20 filtrate was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to
25 give the title compound (160 mg).

MS (API+): [M+H]⁺ 613.2.

[0592]

M) 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
30

To 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (160 mg) were slowly added
35 trifluoroacetic acid (4 mL) and water (0.4 mL) at room

temperature. The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (3 mL), and 8 M ammonia/methanol solution
5 (0.5 mL) was slowly added thereto. The reaction mixture was stirred at room temperature for 30 min, and the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous
10 magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (27.0 mg) and the title compound (100 mg) containing an impurity.

15 MS (API+): [M+H]⁺ 483.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.62-0.70 (2H, m), 0.94-1.04 (5H, m), 1.19 (3H, s), 1.91-2.04 (1H, m), 4.25-4.47 (2H, m), 4.68 (1H, d, J = 8.3 Hz), 4.85 (1H, s), 6.94 (1H, d, J = 1.9 Hz), 7.22-7.30 (1H, m), 7.35-7.43 (1H, m), 7.43-7.53 (1H, m), 7.87
20 (1H, d, J = 1.9 Hz), 10.49 (1H, d, J = 7.9 Hz), 10.67 (1H, brs).

[0593]

Example 75-II

7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-
25 b]pyrazine-4(1H)-carboxamide

To a mixture of 4-nitrophenyl 7-cyclopropyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (1.06 g) and 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methylpropan-
30 2-ol hydrochloride (1.09 g) in N,N-dimethylformamide (30 mL) was added triethylamine (1.25 mL) at room temperature. The reaction mixture was stirred at room temperature for 16 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water
35 and saturated brine, and dried over anhydrous magnesium

sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (1.12 g).

MS (API+): [M+H]⁺ 483.2.

5 ¹H NMR (300 MHz, CDCl₃) δ 0.67-0.76 (2H, m), 0.99-1.09 (2H, m), 1.16 (3H, s), 1.37 (3H, s), 1.62 (1H, s), 1.83-1.96 (1H, m), 4.52-4.72 (2H, m), 4.85 (1H, d, J = 8.3 Hz), 6.83 (1H, d, J = 1.9 Hz), 7.13-7.34 (3H, m), 7.88 (1H, d, J = 1.9 Hz), 8.57 (1H, s), 10.81 (1H, d, J = 8.3 Hz).

10 [0594]

Example 76-I

optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 7-Cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (100 mg) containing an impurity, which was obtained in Step M of Example 75, was purified by HPLC (C18, mobile phase: water (10 mM, containing
20 NH₄HCO₃)/acetonitrile), and the obtained fraction was concentrated under reduced pressure. Then, the residue was optically resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 850/150) to give the title compound
25 having a shorter retention time (24 mg).

MS (API+): [M+H]⁺ 483.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.62-0.70 (2H, m), 0.94-1.04 (5H, m), 1.19 (3H, s), 1.91-2.03 (1H, m), 4.25-4.46 (2H, m), 4.68 (1H, d, J = 8.3 Hz), 4.86 (1H, s), 6.94 (1H, d, J = 1.9 Hz),
30 7.22-7.30 (1H, m), 7.35-7.42 (1H, m), 7.43-7.54 (1H, m), 7.87 (1H, d, J = 1.9 Hz), 10.49 (1H, d, J = 7.9 Hz), 10.68 (1H, brs).

[0595]

Example 76-II

35 optically active 7-cyclopropyl-N-(1-(3-fluoro-4-

(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

7-Cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.25 g) was optically resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 850/150), and recrystallized from acetone/heptane to give the title compound having a shorter retention time (474 mg).

MS (API+): [M+H]⁺ 483.2.

¹H NMR (300 MHz, CDCl₃) δ 0.67-0.76 (2H, m), 0.99-1.09 (2H, m), 1.16 (3H, s), 1.37 (3H, s), 1.64 (1H, s), 1.84-1.95 (1H, m), 4.52-4.73 (2H, m), 4.86 (1H, d, J = 8.3 Hz), 6.84 (1H, d, J = 1.9 Hz), 7.13-7.32 (6H, m), 7.88 (1H, d, J = 1.9 Hz), 8.81 (1H, brs), 10.82 (1H, d, J = 8.3 Hz).

[0596]

Example 77

optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

7-Cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (100 mg) containing an impurity, which was obtained in Step M of Example 75, was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), and optically resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 850/150) to give the title compound having a longer retention time (29 mg).

MS (API+): [M+H]⁺ 483.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.62-0.70 (2H, m), 0.95-1.04 (5H, m), 1.19 (3H, s), 1.92-2.03 (1H, m), 4.26-4.46 (2H, m), 4.68 (1H, d, J = 8.3 Hz), 4.86 (1H, s), 6.94 (1H, d, J = 2.3 Hz), 7.22-7.29 (1H, m), 7.35-7.42 (1H, m), 7.43-7.53 (1H, m), 7.87

(1H, d, J = 2.3 Hz), 10.49 (1H, d, J = 8.3 Hz), 10.68 (1H, s).
[0597]

Example 78

2-oxo-N-(1-(4-(2,2,2-trifluoroethoxy)phenyl)propyl)-2,3-
5 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0598]

A) tert-butyl (1-(4-(2,2,2-trifluoroethoxy)phenyl)propyl)carbamate

To a mixture of tert-butyl (1-(4-
10 hydroxyphenyl)propyl)carbamate (302 mg) and 2,2,2-trifluoroethyl trifluoromethanesulfonate (418 mg) in N,N-dimethylformamide (10 mL) was added potassium carbonate (498 mg). The reaction mixture was stirred at 80°C for 16 hr, and the solvent was evaporated under reduced pressure. To the
15 residue was added ethyl acetate. The organic layer was washed with water, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (362 mg).

MS (API-): [M-H]⁻332.1.

20 [0599]

B) 2-oxo-N-(1-(4-(2,2,2-trifluoroethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

1-(4-(2,2,2-Trifluoroethoxy)phenyl)propylamine hydrochloride was obtained from tert-butyl (1-(4-(2,2,2-trifluoroethoxy)phenyl)propyl)carbamate in the same manner as
25 in Step E of Example 18. The title compound was obtained in the same manner as in Step B of Example 53.

MS (API+): [M+H]⁺409.1.

[0600]

30 Example 79

N-(2-(methylsulfinyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from N-(2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-
35 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same

manner as in Example 60.

MS (API+): [M+H]⁺ 443.1.

[0601]

Example 80

5 2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-imidazol-4-yl)propyl)-
2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0602]

A) 1-trityl-1H-imidazol-4-carbaldehyde

A mixture of 1H-imidazol-4-carbaldehyde (3.00 g),
10 triethylamine (7.00 g) and N,N-dimethylformamide (40 mL) was
cooled to 0°C, trityl chloride (10.5 g) was added thereto. The
reaction mixture was stirred at room temperature for 18 hr,
water was added thereto, and the solid was collected by
filtration. The obtained solid was washed with water and
15 diethyl ether to give the title compound (10.0 g).

¹H NMR (400 MHz, DMSO-d₆) δ 7.11-7.14 (6H, m), 7.40-7.48 (9H,
m), 7.67 (1H, d, J = 0.8 Hz), 7.80 (1H, d, J = 1.2 Hz), 9.23
(1H, s).

[0603]

20 B) 1-(1-trityl-1H-imidazol-4-yl)propan-1-ol

To a mixture of 1-trityl-1H-imidazol-4-carbaldehyde (1.00
g) and tetrahydrofuran (100 mL) was slowly added 2.96 M
ethylmagnesium bromide/tetrahydrofuran solution (1.50 mL) at
0°C. The reaction mixture was stirred at room temperature for
25 18 hr, saturated aqueous ammonium chloride solution was added
thereto, and the mixture was extracted with ethyl acetate. The
extract was washed successively with water and saturated brine,
and dried over anhydrous sodium sulfate, and the solvent was
evaporated under reduced pressure to give the title compound
30 (1.00 g).

¹H NMR (400 MHz, DMSO-d₆) δ 0.78 (3H, t, J=7.6 Hz), 1.54-1.59
(1H, m), 1.68-1.72 (1H, m), 4.36 (1H, t, J = 6.0 Hz), 4.81 (1H,
d, J = 5.6 Hz), 6.65 (1H, s), 7.02-7.12 (6H, m), 7.26 (1H, d,
J = 1.2 Hz), 7.35-7.43 (9H, m).

35 [0604]

C) 1-(1-trityl-1H-imidazol-4-yl)propan-1-one

A mixture of 1-(1-trityl-1H-imidazol-4-yl)propan-1-ol (1.00 g), manganese dioxide (2.36 g) and dioxane (20 mL) was heated at reflux for 2 hr. The reaction mixture was cooled to
5 room temperature, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure to give the title compound (1.00 g).

¹H NMR (400 MHz, DMSO-d₆) δ 1.03 (3H, t, J = 7.6 Hz), 2.86 (2H, q, J = 7.6 Hz), 7.10-7.15 (6H, m), 7.35-7.80 (10H, m), 7.54
10 (1H, d, J = 1.2 Hz).

[0605]

D) 2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-imidazol-4-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A mixture of 1-(1-trityl-1H-imidazol-4-yl)propan-1-one
15 (1.00 g) and 4 M hydrogen chloride/ethyl acetate solution (40 mL) was stirred at room temperature for 2 hr, and the solvent was evaporated under reduced pressure. To the residue was added N,N-dimethylformamide (10 mL), potassium carbonate (829 mg) and 2,2,2-trifluoroethyl trifluoromethanesulfonate (950
20 mg) was added thereto. The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated
25 under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give 1-(1-(2,2,2-trifluoroethyl)-1H-imidazol-4-yl)propan-1-one containing an impurity. To the obtained residue were added hydroxylamine hydrochloride (202 mg), triethylamine (294 mg)
30 and ethanol (20 mL), and the reaction mixture was heated at reflux for 4 hr. The reaction mixture was cooled to room temperature, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed
35 successively with water and saturated brine, and dried over

anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. To the residue were added methanol (50 mL) and Raney nickel (1.00 g), and the reaction mixture was stirred at 50°C for 4 hr under hydrogen atmosphere (50 psi).
5 The reaction mixture was cooled to room temperature, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. To the residue were added 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (491 mg), triethylamine (303 mg) and N,N-
10 dimethylformamide (20 mL), and the reaction mixture was stirred overnight. The reaction mixture was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), and the obtained solid was crystallized from ethyl acetate to give the title compound (76 mg).
15 MS (API+): [M+H]⁺383.1.

[0606]

Example 81

2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-1,2,4-triazol-3-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 [0607]

A) methyl 1-trityl-1H-1,2,4-triazol-3-carboxylate

The title compound was obtained from methyl 1H-1,2,4-triazol-3-carboxylate in the same manner as in Step A of Example 80.

25 ¹H NMR (400 MHz, DMSO-d₆) δ 3.84 (3H, s), 7.02-7.12 (6H, m), 7.30-7.42 (9H, m), 8.39 (1H, s).

[0608]

B) 1-(1-(2,2,2-trifluoroethyl)-1H-1,2,4-triazol-3-yl)propan-1-one

30 A mixture of 3.0 M ethylmagnesium bromide/tetrahydrofuran solution (13.2 mL) and triethylamine (11.0 g) was stirred at room temperature for 30 min. The reaction mixture was cooled to -20°C, a solution of methyl 1-trityl-1H-1,2,4-triazol-3-carboxylate (7.30 g) in tetrahydrofuran (150 mL) was added
35 thereto, and the mixture was stirred at -20°C for 1 hr. To the

reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under
5 reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (4.8 g).

¹H NMR (400 MHz, DMSO-d₆) δ 1.04 (3H, t, J = 7.6 Hz), 2.97 (2H, q, J = 7.6 Hz), 7.02-7.10 (6H, m), 7.35-7.43 (9H, m), 8.36 (1H,
10 s).

[0609]

C) 2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-1,2,4-triazol-3-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(1-(2,2,2-trifluoroethyl)-1H-1,2,4-triazol-3-yl)propan-1-one in the same
15 manner as in Step D of Example 80.

MS (API+): [M+H]⁺384.1.

[0610]

Example 82

20 N-(1-(4-isopropoxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0611]

A) tert-butyl (1-(4-isopropoxyphenyl)propyl)carbamate

To a mixture of tert-butyl (1-(4-hydroxyphenyl)propyl)carbamate (628 mg) and 2-iodopropane
25 (0.324 mL) in N,N-dimethylformamide (15 mL) was added potassium carbonate (518 mg). The reaction mixture was stirred at room temperature for 16 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was
30 washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (309 mg).

35 ¹H NMR (300 MHz, CDCl₃) δ 0.87 (3H, t, J = 7.3 Hz), 1.32 (6H, d,

J = 6.0 Hz), 1.41 (9H, s), 1.66-1.83 (2H, m), 4.35-4.61 (2H, m), 4.73 (1H, brs), 6.80-6.87 (2H, m), 7.11-7.19 (2H, m).

[0612]

B) N-(1-(4-isopropoxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To tert-butyl (1-(4-isopropoxyphenyl)propyl)carbamate (305 mg) was added 4 M hydrogen chloride/ethyl acetate solution (5 mL), and the mixture was stirred at room temperature for 1 hr. The solvent was evaporated under reduced pressure to give 1-(4-isopropoxyphenyl)propan-1-amine hydrochloride. To a mixture of 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (297 mg) and 1-(4-isopropoxyphenyl)propan-1-amine hydrochloride in N,N-dimethylformamide (10 mL) was added triethylamine (0.395 mL), and the mixture was stirred at room temperature for 16 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and crystallized from hexane/ethyl acetate to give the title compound (178 mg). MS (API-): [M-H]⁻367.2.

[0613]

Example 83

N-(1-(4-chlorophenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0614]

A) 1-(4-chlorophenyl)propan-1-amine

To a solution of 1-(4-chlorophenyl)propan-1-one (1.69 g) in ethanol (60 mL) were added hydroxylamine hydrochloride (1.39 g) and triethylamine (2.79 mL) at room temperature. The reaction mixture was stirred at room temperature for 16 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with

ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give 1-(4-chlorophenyl)-N-hydroxypropan-1-imine. To a
5 solution of 1-(4-chlorophenyl)-N-hydroxypropan-1-imine in tetrahydrofuran (100 mL) was added 1.1 M borane-tetrahydrofuran complex/tetrahydrofuran solution (27.3 mL), and the mixture was stirred at 80°C for 16 hr. To the reaction mixture was added 1 M hydrochloric acid, and the mixture was
10 extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the
15 title compound (520 mg).

¹H NMR (300 MHz, CDCl₃) δ 0.85 (3H, t, J = 7.3 Hz), 1.58-1.74 (2H, m), 3.80 (1H, t, J = 7.0 Hz), 7.21-7.40 (4H, m).

[0615]

B) N-(1-(4-chlorophenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
20 b]pyrazine-4(1H)-carboxamide

To a mixture of 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (300 mg) and 1-(4-chlorophenyl)propan-1-amine (211 mg) in N,N-dimethylformamide (10 mL) was added triethylamine (0.279 mL),
25 and the mixture was stirred at room temperature for 16 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated
30 under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and recrystallized from hexane/ethyl acetate to give the title compound (205 mg).

MS (API+): [M+H]⁺345.1.

35 [0616]

Example 84

2-oxo-N-(1-(2-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0617]

5 A) N-methoxy-N-methyl-2-(trifluoromethoxy)benzamide

To a mixture of 2-(trifluoromethoxy)benzoic acid (5.00 g), N,O-dimethylhydroxylamine hydrochloride (2.63 g), and triethylamine (7.36 g) in N,N-dimethylformamide (50 mL) were added 1-hydroxybenzotriazole monohydrate (4.94 g) and 1-(3-
10 dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (2.63 g). The reaction mixture was stirred at 10°C for 16 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was
15 evaporated under reduced pressure to give the title compound (5.21 g).

¹H NMR (400 MHz, CDCl₃) δ 3.36 (3H, s), 3.45 (3H, s), 7.27-7.37 (2H, m), 7.39-7.50 (2H, m).

[0618]

20 B) N-hydroxy-1-(2-(trifluoromethoxy)phenyl)propan-1-imine

The title compound was obtained from N-methoxy-N-methyl-2-(trifluoromethoxy)benzamide in the same manner as in Steps C-D of Example 2.

¹H NMR (400 MHz, CDCl₃) δ 1.05 (3H, t, J = 7.6 Hz), 2.77 (2H, q, J = 7.6 Hz), 7.27-7.35 (2H, m), 7.36-7.44 (2H, m), 8.28 (1H, s).

[0619]

C) 1-(2-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride

To a solution of N-hydroxy-1-(2-(trifluoromethoxy)phenyl)propan-1-imine (2.31 g) in methanol
30 (100 mL) was added Raney nickel (1.00 g). The reaction mixture was stirred at 50°C for 4 hr under hydrogen atmosphere (50 psi), the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved
35 in a mixed solvent of ethyl acetate (70 mL) and methanol (10

mL), 4 M hydrogen chloride/ethyl acetate solution (40 mL) was added thereto. The reaction mixture was stirred at 15°C for 2 hr, and the solvent was evaporated under reduced pressure. The obtained solid was washed with tert-butyl methyl ether to give
5 the title compound (1.36 g).

MS (API+): [M+H]⁺219.9.

[0620]

D) 2-oxo-N-(1-(2-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 The title compound was obtained from 1-(2-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺395.1.

[0621]

15 Example 85

N-(1-(4-methylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0622]

A) methyl 4-(difluoromethyl)benzoate

20 To a solution of methyl 4-formylbenzoate (821 mg) in toluene (25 mL) was added N,N-diethylaminosulfur trifluoride (1.98 mL). The reaction mixture was stirred at room temperature for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was
25 washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and crystallized from hexane/ethyl acetate to give the title
30 compound (619 mg).

¹H NMR (300 MHz, CDCl₃) δ 3.95 (3H, s), 6.46-6.92 (1H, m), 7.59 (2H, d, J = 8.3 Hz), 8.13 (2H, d, J = 8.7 Hz).

[0623]

B) 4-(difluoromethyl)-N-methoxy-N-methylbenzamide

35 To a solution of methyl 4-(difluoromethyl)benzoate (615

mg) in a mixed solvent of tetrahydrofuran (15 mL) and methanol (5 mL) was added 2 M aqueous sodium hydroxide solution (3.0 mL), and the mixture was stirred at room temperature for 3 hr. The reaction mixture was neutralized with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. A mixture of the obtained residue, N,O-dimethylhydroxylamine hydrochloride (387 mg), triethylamine (0.921 mL), 1-hydroxybenzotriazole monohydrate (607 mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (760 mg) in N,N-dimethylformamide (15 mL) was stirred at room temperature for 20 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (540 mg).

^1H NMR (300 MHz, CDCl_3) δ 3.38 (3H, s), 3.54 (3H, s), 6.47-6.89 (1H, m), 7.56 (2H, d, $J = 8.3$ Hz), 7.77 (2H, d, $J = 8.7$ Hz).
[0624]

C) 1-(4-(difluoromethyl)phenyl)propan-1-one

The title compound was obtained from 4-(difluoromethyl)-N-methoxy-N-methylbenzamide in the same manner as in Step C of Example 2.

^1H NMR (300 MHz, CDCl_3) δ 1.24 (3H, t, $J = 7.2$ Hz), 3.03 (2H, q, $J = 7.2$ Hz), 6.48-6.90 (1H, m), 7.61 (2H, d, $J = 8.3$ Hz), 8.05 (2H, d, $J = 8.7$ Hz).
[0625]

D) 1-(4-methylphenyl)propan-1-amine

To a solution of 1-(4-(difluoromethyl)phenyl)propan-1-one (360 mg) in ethanol (10 mL) were added hydroxylamine hydrochloride (272 mg) and triethylamine (0.545 mL). The

reaction mixture was stirred at room temperature for 16 hr,
and the solvent was evaporated under reduced pressure. To the
residue was added water, and the mixture was extracted with
ethyl acetate. The extract was washed successively with water
5 and saturated brine, and dried over anhydrous magnesium
sulfate, and the solvent was evaporated under reduced pressure.
To the residue were added ethanol (20 mL) and 20% palladium
hydroxide-carbon (40 mg). The reaction mixture was stirred at
room temperature for 16 hr under hydrogen atmosphere, the
10 catalyst was filtered off, and the filtrate was concentrated
under reduced pressure to give the title compound (280 mg).
¹H NMR (300 MHz, CDCl₃) δ 0.72 (3H, t, J = 7.2 Hz), 1.66-1.94
(2H, m), 2.31 (3H, s), 3.73 (1H, dd, J = 8.7, 5.7 Hz), 7.08-
7.23 (4H, m), 8.26 (2H, brs).

15 [0626]

E) N-(1-(4-methylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-
methylphenyl)propan-1-amine in the same manner as in Step B of
20 Example 83.

MS (API+): [M+H]⁺325.2.

[0627]

Example 86

2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl)propyl)-
25 2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0628]

A) methyl 1H-pyrazol-4-carboxylate

A mixture of 1H-pyrazol-4-carboxylic acid (4.00 g) and 4
M hydrogen chloride/methanol solution (150 mL) was stirred
30 overnight at room temperature, and the solvent was evaporated
under reduced pressure to give the title compound (5.0 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.72 (3H, s), 8.09 (2H, s), 11.62
(1H, s).

[0629]

35 B) methyl 1-trityl-1H-pyrazol-4-carboxylate

The title compound was obtained from methyl 1H-pyrazole-4-carboxylate in the same manner as in Step A of Example 80. ¹H NMR (400 MHz, DMSO-d₆) δ 3.79 (3H, s), 7.11-7.14 (6H, m), 7.31-7.33 (9H, s) 7.93 (1H, s), 8.04 (1H, s).

5 [0630]

C) N-methoxy-N-methyl-1-trityl-1H-pyrazole-4-carboxamide

A mixture of methyl 1-trityl-1H-pyrazol-4-carboxylate (8.00 g) and lithium hydroxide (2.70 g) in water/tetrahydrofuran/methanol (1:2:2) (100 mL) was stirred at
10 60°C for 3 hr, and the solvent was evaporated under reduced pressure. To the residue were added N,N-dimethylformamide (50 mL), N,O-dimethylhydroxylamine hydrochloride (2.10 g), triethylamine (6.80 g) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (16.8 g) at
15 room temperature. The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with aqueous sodium carbonate solution, water and saturated brine, and dried over anhydrous sodium sulfate, and
20 the solvent was evaporated under reduced pressure to give the title compound (5.7 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.29 (3H, s), 3.63 (3H, s), 7.13-7.16 (6H, m), 7.31-7.33 (9H, s) 7.98 (1H, s), 8.11 (1H, s).

[0631]

25 D) 2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from N-methoxy-N-methyl-1-trityl-1H-pyrazol-4-carboxamide in the same manner as in Step C of Example 2 and Step D of Example 80.

30 MS (API+): [M+H]⁺383.1.

[0632]

Example 87

N-(1-(3-methyl-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

35 The title compound was obtained from 1-bromo-3-methyl-4-

(trifluoromethoxy)benzene in the same manner as in Steps A-B of Example 30, Step D of Example 2, Step C of Example 56 and Step G of Example 7.

MS (API+): [M+H]⁺409.1.

5 [0633]

Example 88

N-(2-(dimethylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 [0634]

A) tert-butyl (2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

A mixture of 2-amino-2-(4-(trifluoromethoxy)phenyl)acetamide (2.15 g), di-tert-butylidicarbonate (2.41 g) and tetrahydrofuran (50 mL) was stirred overnight at room temperature. The solvent was evaporated under reduced pressure to give the title compound (2.99 g).

¹H NMR (300 MHz, CDCl₃) δ 1.41 (9H, s), 5.24 (1H, brs), 5.67 (1H, brs), 5.83 (1H, d, J = 5.7 Hz), 5.91 (1H, brs), 7.16-7.24 (2H, m), 7.37-7.49 (2H, m).

[0635]

B) tert-butyl (2-(dimethylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

25 A mixture of 2 M aqueous sodium hydroxide solution (20 mL), tert-butyl (2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (1.5 g) and methanol (15 mL) was stirred at 70°C for 2 days. The reaction mixture was neutralized with 1 M hydrochloric acid at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a pale yellow solid (1.26 g). A mixture of the obtained solid (400 mg),
35 triethylamine (0.166 mL), 1-hydroxybenzotriazole monohydrate

(219 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (274 mg), dimethylamine hydrochloride (117 mg) and N,N-dimethylformamide (5 mL) was stirred overnight at 80°C. The reaction mixture was poured into saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (254 mg). ¹H NMR (300 MHz, CDCl₃) δ 1.41 (9H, s), 2.90 (3H, s), 2.99 (3H, s), 5.56 (1H, d, J = 7.5 Hz), 6.08 (1H, d, J = 7.5 Hz), 7.14-7.23 (2H, m), 7.36-7.46 (2H, m).

[0636]

C) N-(2-(dimethylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A mixture of tert-butyl (2-(dimethylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (270 mg) and 2 M hydrogen chloride/methanol solution (5 mL) was stirred at room temperature for 30 min, and the solvent was evaporated under reduced pressure. To the residue were added N,N-dimethylformamide (15 mL), triethylamine (226 mg) and 4-nitrophenyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (234 mg), and the mixture was stirred overnight at room temperature. To the reaction mixture was added saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with 1 M aqueous sodium hydroxide solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) and then silica gel column chromatography (ethyl acetate/methanol) to give the title compound (83 mg).

MS (API+): [M+H]⁺438.1.

[0637]

Example 89

N-(2-(methylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-
5 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Steps A-C of Example 88.

MS (API+): [M+H]⁺424.1.

[0638]

10 Example 90

N-(2-isopropoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-
2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0639]

A) 2-isopropoxy-1-(4-(trifluoromethoxy)phenyl)ethanamine

15 The title compound was obtained in the same manner as in Step D of Example 63, Step C of Example 1 and Step B of Example 54.

MS (API+): [M+H]⁺264.1.

[0640]

20 B) N-(2-isopropoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-isopropoxy-1-(4-(trifluoromethoxy)phenyl)ethanamine in the same manner as in Step B of Example 83.

25 MS (API+): [M+H]⁺439.2.

[0641]

Example 91

N-(2-(cyclopentyloxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-
2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 [0642]

A) 2-(cyclopentyloxy)-1-(4-(trifluoromethoxy)phenyl)ethanamine

The title compound was obtained in the same manner as in Step D of Example 63, Step C of Example 1 and Step B of Example 54.

35 MS (API+): [M+H]⁺290.2.

[0643]

B) N-(2-(cyclopentyloxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-(cyclopentyloxy)-1-(4-(trifluoromethoxy)phenyl)ethanamine in the same manner as in Step B of Example 83.

MS (API+): [M+H]⁺465.2.

[0644]

Example 92

10 N-(3-amino-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0645]

A) tert-butyl (3-amino-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

15 To a solution of 3-((tert-butoxycarbonyl)amino)-3-(4-(trifluoromethoxy)phenyl)propanoic acid (700 mg) in N,N-dimethylformamide (25.0 mL) were added 1H-benzotriazol-1-ol ammonium salt (335 mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (431 mg). The reaction mixture was stirred overnight at room temperature, water was added thereto, and the precipitated solid was collected by filtration, and washed with water to give the title compound (740 mg).

MS (API-): [M-H]⁻347.2.

25 [0646]

B) 3-amino-3-(4-(trifluoromethoxy)phenyl)propanamide hydrochloride

To tert-butyl (3-amino-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate (740 mg) was added 4 M hydrogen chloride/ethyl acetate solution (10 mL). The reaction mixture was stirred at room temperature for 1 hr, and the precipitated solid was collected by filtration, and washed with ethyl acetate to give the title compound (413 mg).

MS (API+), found: 249.1.

35 [0647]

C) N-(3-amino-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-amino-3-(4-(trifluoromethoxy)phenyl)propanamide hydrochloride in the same
5 manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺424.1.

[0648]

Example 93

N-(3-(dimethylamino)-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
10 b]pyrazine-4(1H)-carboxamide

[0649]

A) tert-butyl (3-(dimethylamino)-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

15 To a solution of 3-((tert-butoxycarbonyl)amino)-3-(4-(trifluoromethoxy)phenyl)propanoic acid (600 mg) in N,N-dimethylformamide (25 mL) were added 1-hydroxybenzotriazole monohydrate (353 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (437 mg), N-methylmethanamine
20 (1.12 mL) and triethylamine (725 µL). The reaction mixture was stirred overnight at room temperature, water was added thereto, and the precipitated solid was collected by filtration, and washed with water to give the title compound (600 mg).

MS (API-): [M-H]⁻375.1.

25 [0650]

B) 3-amino-N,N-dimethyl-3-(4-(trifluoromethoxy)phenyl)propanamide hydrochloride

The title compound was obtained from tert-butyl (3-(dimethylamino)-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate in the same manner
30 as in Step B of Example 92.

MS (API+), found: 277.1.

[0651]

C) N-(3-(dimethylamino)-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
35

b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-amino-N,N-dimethyl-3-(4-(trifluoromethoxy)phenyl)propanamide hydrochloride in the same manner as in Step H of Example 72-I.

5 MS (API+): [M+H]⁺452.2.

[0652]

Example 94

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 The title compound was obtained from 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethylamine hydrochloride in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺425.2.

[0653]

15 Example 95

3-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0654]

A) methyl 2-((3-nitropyridin-2-yl) amino)propanoate

20 The title compound was obtained from methyl 2-aminopropanoate hydrochloride in the same manner as in Step A of Example 19.

MS (API+): [M+H]⁺225.8.

[0655]

25 B) 3-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl 2-((3-nitropyridin-2-yl) amino)propanoate (18.0 g) in ethanol (500 mL) were added iron powder (17.9 g) and conc. hydrochloric acid (2 mL). The reaction mixture was heated at reflux for 16 hr, the insoluble
30 substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), and the obtained fraction was concentrated under reduced pressure. To the residue was added conc. hydrochloric acid (1 mL), and
35 the mixture was concentrated under reduced pressure. To the

residue was added saturated aqueous sodium hydrogen carbonate solution, the reaction mixture was stirred at 15°C for 16 hr, and the solid was collected by filtration. The obtained solid was washed with water, and dried under reduced pressure to
5 give the title compound (1.2 g).

MS (API+): [M+H]⁺163.8.

[0656]

C) 3-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 The title compound was obtained from 3-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner as in Step F of Example 1.

MS (API+): [M+H]⁺409.1.

[0657]

15 Example 96

N-(1-(4-(difluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(difluoromethyl)phenyl)propan-1-one in the same manner as in
20 Steps A-B of Example 83.

MS (API+): [M+H]⁺361.2.

[0658]

Example 97

N-(2-cyano-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
25

[0659]

A) tert-butyl (3-amino-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

The title compound was obtained in the same manner as in
30 Step A of Example 92.

MS (API-): [M-H]⁻347.1.

[0660]

B) tert-butyl (2-cyano-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

35 To a solution of tert-butyl (3-amino-3-oxo-1-(4-

(trifluoromethoxy)phenyl)propyl)carbamate (425 mg) in tetrahydrofuran (30.0 mL) was added thionyl chloride (281 μ L). The reaction mixture was stirred overnight at 60°C, and the solvent was evaporated under reduced pressure. To the residue
5 was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
10 column chromatography (hexane/ethyl acetate) to give the title compound (251 mg).

MS (API-): [M-H]⁻329.1.

[0661]

C) 3-amino-3-(4-(trifluoromethoxy)phenyl)propanenitrile
15 hydrochloride

The title compound was obtained from tert-butyl (2-cyano-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate in the same manner as in Step B of Example 92.

¹H NMR (300 MHz, DMSO-d₆) δ 3.23-3.44 (2H, m), 4.80 (1H, dd, J = 8.3, 6.0 Hz), 7.52 (2H, d, J = 7.9 Hz), 7.72-7.84 (2H, m), 9.01 (3H, brs).

[0662]

D) N-(2-cyano-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

25 The title compound was obtained from 3-amino-3-(4-(trifluoromethoxy)phenyl)propanenitrile hydrochloride in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺406.2.

[0663]

30 Example 98

2-oxo-N-(1-(3-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0664]

A) 1-(3-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride

35 The title compound was obtained from 3-

(trifluoromethoxy)benzoic acid in the same manner as in Step A of Example 84, Steps C-D of Example 2 and Step C of Example 84. ¹H NMR (400 MHz, DMSO-d₆) δ 0.75 (3H, t, J = 7.2 Hz), 1.71-1.85 (1H, m), 1.92-2.05 (1H, m), 4.17-4.28 (1H, m), 7.34-7.42 (1H, m), 7.51-7.61 (3H, m), 8.66 (3H, brs).

[0665]

B) 2-oxo-N-(1-(3-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(3-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺395.2.

[0666]

Example 99

N-(1-(4-cyclopropylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0667]

A) 1-(4-cyclopropylphenyl)propan-1-one

A mixture of 1-(4-bromophenyl)propan-1-one (1.07 g), cyclopropylboronic acid (0.558 g), (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (183 mg), tripotassium phosphate (2.12 g) in a mixed solvent of 1,2-dimethoxyethane (15 mL) and water (5 mL) was stirred at 85°C for 20 hr under nitrogen atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (679 mg).

MS (API+): [M+H]⁺174.9.

[0668]

B) N-(1-(4-cyclopropylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-

cyclopropylphenyl)propan-1-one in the same manner as in Steps A-B of Example 83.

MS (API+): [M+H]⁺351.2.

[0669]

5 Example 100

N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0670]

A) 1-(4-(trifluoromethoxy)phenyl)prop-2-en-1-ol

10 To a solution of 4-(trifluoromethoxy)benzaldehyde (12.0 g) in tetrahydrofuran (200 mL) was slowly added 1 M vinylmagnesium bromide/tetrahydrofuran solution (69.5 mL) under nitrogen atmosphere, at -78°C. The reaction mixture was allowed to be slowly warmed to room temperature, and stirred
15 at room temperature for 16 hr under nitrogen atmosphere. To the reaction mixture was added saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was
20 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound (11.8 g).

¹H NMR (400 MHz, DMSO-d₆) δ 5.02-5.13 (2H, m), 5.26 (1H, dt, J = 17.2, 1.6 Hz), 5.64 (1H, d, J = 4.4 Hz), 5.87-5.99 (1H, m),
25 7.31 (2H, d, J = 8.0 Hz), 7.44 (2H, dd, J = 6.8, 1.6 Hz).

[0671]

B) 1-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one

To a solution of 1-(4-(trifluoromethoxy)phenyl)prop-2-en-1-ol (9.80 g) in dichloromethane (150 mL) was added
30 manganese(IV) dioxide (39.1 g). The reaction mixture was stirred at room temperature for 2 days, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give
35 the title compound (2.75 g).

¹H NMR (400 MHz, CDCl₃) δ 5.97 (1H, dd, J = 10.8, 1.6 Hz), 6.46 (1H, dd, J = 17.2, 1.6 Hz), 7.13 (1H, dd, J = 17.2, 10.8 Hz), 7.31 (2H, dd, J = 8.8, 0.8 Hz), 8.00 (2H, dd, J = 6.8, 2.0 Hz).
[0672]

5 C) 3-methoxy-1-(4-(trifluoromethoxy)phenyl)propan-1-one

A mixture of 1-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one (2.75 g), methanol (407 mg) and bis(acetonitrile)dichloropalladium(II) (328 mg) in dichloromethane (30 mL) was stirred at room temperature for 16
10 hr under nitrogen atmosphere, dichloromethane was added thereto, and the insoluble substance was filtered off. The filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) to give the title compound
15 (2.80 g).

¹H NMR (400 MHz, CDCl₃) δ 3.22 (2H, t, J = 6.4 Hz), 3.38 (3H, s), 3.82 (2H, t, J = 6.4 Hz), 7.29 (2H, d, J = 8.4 Hz), 8.02 (2H, dd, J = 6.8, 2.0 Hz).
[0673]

20 D) 3-methoxy-1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride

The title compound was obtained from 3-methoxy-1-(4-(trifluoromethoxy)phenyl)propan-1-one in the same manner as in Steps D-F of Example 2.

25 MS (API⁺): [M+H]⁺250.0.
[0674]

E) N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-methoxy-1-(4-(trifluoromethoxy)phenyl)propan-1-amine hydrochloride in the
30 same manner as in Step L of Example 72-I.

MS (API⁺): [M+H]⁺425.2.
[0675]

Example 101

35 2-oxo-N-(2-(pyrrolidin-1-yl)-1-(4-

(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0676]

A) 2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine
5 (trifluoromethoxy)phenyl)ethanamine

To a mixture of 4-(trifluoromethoxy)phenacyl bromide (3.00 g) and triethylamine (1.61 g) in tetrahydrofuran (15 mL) was added pyrrolidine (0.902 g). The reaction mixture was stirred at room temperature for 1 hr, water was added thereto,
10 and the mixture was extracted with dichloromethane. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give 2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone (3.20 g). To a solution of 2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone (3.20 g) in ethanol (30 mL)
15 were added hydroxylamine hydrochloride (0.805 g) and triethylamine (1.18 g). The reaction mixture was stirred at 78°C for 16 hr, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column
20 chromatography (petroleum ether/ethyl acetate) to give N-hydroxy-2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanimine (1.12 g). To a solution of N-hydroxy-2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanimine (1.12 g) in methanol (50
25 mL) was added Raney nickel (300 mg) at room temperature. The reaction mixture was stirred at 50°C for 3 hr under hydrogen atmosphere (50psi), the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure to give the title compound (1.01 g).

30 MS (API+): [M+H]⁺274.9.

[0677]

B) tert-butyl (2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

To a mixture of 2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine (1.01 g) and triethylamine
35 (trifluoromethoxy)phenyl)ethanamine (1.01 g) and triethylamine

(558 mg) in methanol (15 mL) was added di-tert-butyl dicarbonate (1.20 g) at room temperature. The reaction mixture was stirred at room temperature for 4 hr, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate) and then HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% NH_4HCO_3)) to give the title compound (560 mg). ^1H NMR (400 MHz, CDCl_3) δ 1.39 (9H, s), 1.71-1.85 (4H, m), 2.41-2.69 (5H, m), 2.73-2.86 (1H, m), 4.51-4.71 (1H, m), 5.58-5.78 (1H, m), 7.16 (2H, d, $J = 8.0$ Hz), 7.32 (2H, d, $J = 8.4$ Hz).

[0678]

C) 2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine hydrochloride

To tert-butyl (2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (560 mg) was added 4 M hydrogen chloride/ethyl acetate solution (20 mL) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and the solvent was evaporated under reduced pressure to give the title compound (310 mg). ^1H NMR (400 MHz, CD_3OD) δ 2.02-2.15 (4H, m), 3.35-3.58 (4H, m), 3.97 (2H, d, $J = 6.8$ Hz), 5.00 (1H, t, $J = 6.8$ Hz), 7.49 (2H, d, $J = 8.0$ Hz), 7.80 (2H, d, $J = 8.8$ Hz).

[0679]

D) 2-oxo-N-(2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine hydrochloride in the same manner as in Step L of Example 72-I.

MS (API+): $[\text{M}+\text{H}]^+450.2$.

[0680]

Example 102

2-oxo-N-(2-(2-oxopyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

[0681]

A) 5-methoxy-3,4-dihydro-2H-pyrrole

A mixture of pyrrolidine (7.30 g) and dimethyl sulfate
5 (12.1 g) was stirred at 60°C for 18 hr, water was added thereto,
and the pH of the mixture was adjusted to 8-9 with saturated
potassium carbonate aqueous solution. The reaction mixture was
extracted with tert-butyl methyl ether, dried over anhydrous
sodium sulfate, and the solvent was evaporated under reduced
10 pressure to give the title compound (7.2 g).

¹H NMR (400 MHz, CDCl₃) δ 1.89-1.99 (2H, m), 2.30-2.40 (2H, m),
3.52-3.60 (2H, m), 3.71 (3H, s).

[0682]

B) 1-(2-oxo-2-(4-(trifluoromethoxy)phenyl)ethyl)pyrrolidin-2-
15 one

A mixture of 5-methoxy-3,4-dihydro-2H-pyrrole (5.60 g)
and 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone (3.00 g) in
N,N-dimethylformamide (20 mL) was stirred at 50-60°C for 5 hr,
water was added thereto, and the mixture was extracted with
20 ethyl acetate. The extract was washed with saturated brine,
and dried over anhydrous sodium sulfate, and the solvent was
evaporated under reduced pressure. The residue was purified by
silica gel column chromatography (petroleum ether/ethyl
acetate) to give the title compound (2.32 g).

25 ¹H NMR (400 MHz, CDCl₃) δ 2.01-2.15 (2H, m), 2.48 (2H, t, J =
8.0 Hz), 3.50 (2H, t, J = 7.2 Hz), 4.70 (2H, s), 7.31 (2H, d,
J = 8.0 Hz), 8.03 (2H, d, J = 8.8 Hz).

[0683]

C) 1-(2-amino-2-(4-(trifluoromethoxy)phenyl)ethyl)pyrrolidin-
30 2-one

A mixture of 1-(2-oxo-2-(4-
(trifluoromethoxy)phenyl)ethyl)pyrrolidin-2-one (2.32 g) and
hydroxylamine hydrochloride (1.12 g) in pyridine (20 mL) was
stirred at 12°C for 16 hr, water was added thereto, and the
35 mixture was extracted with ethyl acetate. The extract was

washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give 1-(2-(hydroxyimino)-2-(4-(trifluoromethoxy)phenyl)ethyl)pyrrolidin-2-one (2.19 g). A
5 mixture of 1-(2-(hydroxyimino)-2-(4-(trifluoromethoxy)phenyl)ethyl)pyrrolidin-2-one (1.00 g) and 10% palladium-carbon (dry, 100 mg) in methanol (50 mL) was stirred at 25°C for 18 hr under hydrogen atmosphere. The insoluble substance was filtered off through Celite, and the
10 filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (dichloromethane/methanol) to give the title compound (700 mg).
MS (API+): [M+H]⁺288.9.

D) 2-oxo-N-(2-(2-oxopyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-
15 b]pyrazine-4(1H)-carboxamide

The title compound was obtained in the same manner as in Step B of Example 83.

MS (API+): [M+H]⁺464.2.

20 [0684]

Example 103

3-(((2-oxo-2,3-dihydropyrido[2,3-b]pyrazin-4(1H)-yl)carbonyl)amino)-3-(4-(trifluoromethoxy)phenyl)propyl acetate

The title compound was obtained from tert-butyl (3-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate
25 in the same manner as in Step E of Example 18 and Step L of Example 72-I.

MS (API+): [M+H]⁺453.2.

[0685]

30 Example 104

N-(3-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 3-(((2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-yl)carbonyl)amino)-3-(4-(trifluoromethoxy)phenyl)propyl acetate (115 mg) in methanol (5
35

mL) was added potassium carbonate (45.7 mg). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was recrystallized from hexane/ethyl acetate to give the title compound (89.4 mg).

10 MS (API+): [M+H]⁺410.8.
[0686]

Example 105

N-(1-(4-(azetidin-1-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 [0687]

A) 1-(4-(azetidin-1-yl)phenyl)propan-1-one

A mixture of 1-(4-bromophenyl)propan-1-one (639 mg), azetidine (0.303 mL), tris(dibenzylideneacetone)dipalladium(0) (137 mg), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (174 mg) and sodium tert-butoxide (432 mg) in toluene (15 mL) was stirred at 85°C for 20 hr under nitrogen atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (414 mg).

25 ¹H NMR (300 MHz, CDCl₃) δ 1.20 (3H, t, J = 7.3 Hz), 2.42 (2H, quin, J = 7.3 Hz), 2.90 (2H, q, J = 7.2 Hz), 3.99 (4H, t, J = 7.3 Hz), 6.32-6.39 (2H, m), 7.82-7.90 (2H, m).

[0688]

B) N-(1-(4-(azetidin-1-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

35 The title compound was obtained from 1-(4-(azetidin-1-

yl)phenyl)propan-1-one in the same manner as in Steps D-E of Example 2 and Step B of Example 83.

MS (API-): [M-H]⁻364.2.

[0689]

5 Example 106

N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.90 g) was resolved by HPLC (column: CHIRALPAK IC, 46 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 400/600) to give the title compound having a shorter retention time (815 mg).

MS (API+): [M+H]⁺425.2.

[0690]

Example 107

N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (1.90 g) was resolved by HPLC (column: CHIRALPAK IC, 46 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 400/600) to give the title compound having a longer retention time (772 mg).

MS (API+): [M+H]⁺425.1.

30 [0691]

Example 108

N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0692]

35 A) tert-butyl (2-(difluoromethoxy)-1-(4-

(trifluoromethoxy)phenyl)ethyl)carbamate

To a solution of tert-butyl 2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (4.00 g) in acetonitrile (120 mL) were added copper(I) iodide (3.57 g) and
5 difluoro(fluorosulfonyl)acetic acid (1.93 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, saturated aqueous ammonium chloride solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and
10 dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (673 mg).

MS (API-): [M-H]⁻370.1.

15 [0693]

B) 2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethanamine hydrochloride

The title compound was obtained from tert-butyl 2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate in the same manner as
20 in Step B of Example 92.

MS (API+): [M+H]⁺272.1.

[0694]

C) N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
25

The title compound was obtained from 2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethanamine hydrochloride in the same manner as in Step L of Example 72-I.

30 MS (API+): [M+H]⁺447.1.

[0695]

Example 109

N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

35 The title compound was obtained from 2-bromo-1-(4-

(trifluoromethyl)phenyl)ethanone in the same manner as in Steps D-E of Example 63 and Step L of Example 72-I.

MS (API+): [M+H]⁺395.1.

[0696]

5 Example 110

N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-

10 carboxylate and 2-methoxy-1-(4-(trifluoromethyl)phenyl)ethan-1-amine in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺409.2.

[0697]

Example 111

15 N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate and 2-methoxy-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethan-1-amine

20 in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺429.1.

[0698]

Example 112

N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
25 carboxamide

The title compound was obtained from 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate and 2-methoxy-1-(3-fluoro-4-

30 (trifluoromethoxy)phenyl)ethan-1-amine in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺443.1.

[0699]

Example 113

35 N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-

oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-
4(1H)-carboxamide
[0700]

A) methyl N-(6-chloro-3-nitropyridin-2-yl)glycinate

5 The title compound was obtained in the same manner as in
Step A of Example 37.

¹H NMR (300 MHz, DMSO-d₆) δ 3.67 (3H, s), 4.27 (2H, d, J = 6.0
Hz), 6.87 (1H, d, J = 8.7 Hz), 8.48 (1H, d, J = 8.7 Hz), 8.96
(1H, t, J = 5.7 Hz).

10 [0701]

B) methyl N-(3-nitro-6-(1H-pyrazol-1-yl)pyridin-2-yl)glycinate

To a solution of methyl N-(6-chloro-3-nitropyridin-2-
yl)glycinate (1.02 g) in N,N-dimethylacetamide (10 mL) was
added 1H-pyrazole (844 mg) at room temperature. The reaction
15 mixture was stirred at 120°C for 16 hr, water was added thereto,
and the mixture was extracted with ethyl
acetate/tetrahydrofuran. The extract was washed successively
with water and saturated brine, and dried over anhydrous
sodium sulfate, and the solvent was evaporated under reduced
20 pressure. The residue was purified by silica gel column
chromatography (hexane/ethyl acetate) to give the title
compound (293 mg).

MS (API+): [M+H]⁺278.1.

[0702]

25 C) 6-(1H-pyrazol-1-yl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-
one

The title compound was obtained from methyl 2-((3-nitro-
6-(1H-pyrazol-1-yl)pyridin-2-yl) amino)acetate in the same
manner as in Step C of Example 72-I.

30 MS (API-): [M-H]⁻214.1.

[0703]

D) 4-nitrophenyl 2-oxo-6-(1H-pyrazol-1-yl)-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

The title compound was obtained from 6-(1H-pyrazol-1-yl)-
35 3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner

as in Step F of Example 7.

MS (API+): [M+H]⁺381.1.

[0704]

E) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-
5 2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-
4(1H)-carboxamide

The title compound was obtained from 4-nitrophenyl 2-oxo-
6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
carboxylate and 2-methoxy-1-(3-fluoro-4-
10 (trifluoromethoxy)phenyl)ethan-1-amine in the same manner as
in Step L of Example 72-I.

MS (API+): [M+H]⁺495.2.

[0705]

Example 114

15 7-cyclopropyl-N-(2-methoxy-1-(4-
(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

The title compound was obtained from N-(1-(4-
(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-iodo-2-oxo-1-((2-
20 (trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide in the same manner as in Step L
of Example 75 and Step I of Example 63.

MS (API+): [M+H]⁺451.2.

[0706]

25 Example 115

7-cyclopropyl-N-(2-methoxy-1-(4-
(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-methoxy-1-(4-
30 (trifluoromethyl)phenyl)ethan-1-amine hydrochloride in the
same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺435.2.

[0707]

Example 116

35 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-

pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
carboxamide

The title compound was obtained from 4-nitrophenyl 2-oxo-
6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
5 carboxylate and 2-methoxy-1-(4-(trifluoromethyl)phenyl)ethan-
1-amine hydrochloride in the same manner as in Step L of
Example 72-I.

MS (API+): [M+H]⁺477.2.

[0708]

10 Example 117

N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-
methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-
15 methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide (118 mg) was resolved by SFC
(column: CHIRALCEL OJH, 20 mmID×250 mmL, manufactured by
Daicel Chemical Industries, Ltd., mobile phase: carbon
dioxide/methanol=860/140) to give the title compound having a
20 shorter retention time (31.0 mg).

MS (API+): [M+H]⁺495.2.

[0709]

Example 118

N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-
25 methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-
methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide (118 mg) was resolved by SFC
30 (column: CHIRALCEL OJH, 20 mmID×250 mmL, manufactured by
Daicel Chemical Industries, Ltd., mobile phase: carbon
dioxide/methanol=860/140) to give the title compound having a
longer retention time (41.8 mg).

MS (API+): [M+H]⁺495.1.

35 [0710]

Example 119

7-cyclopropyl-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 Racemic 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (245 mg) was resolved by SFC (column: CHIRALCEL OJH, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=860/140) to give the title compound having a shorter retention time (98.6 mg).

MS (API+): [M+H]⁺451.2.

[0711]

Example 120

15 7-cyclopropyl-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 Racemic 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (245 mg) was resolved by SFC (column: CHIRALCEL OJH, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=860/140) to give the title compound having a longer retention time (102 mg).

25 MS (API+): [M+H]⁺451.2.

[0712]

Example 121

N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (184 mg) was resolved by SFC (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=860/140) to give

the title compound having a shorter retention time (70.1 mg).
MS (API+): [M+H]⁺429.1.

[0713]

Example 122

5 N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
10 carboxamide (184 mg) was resolved by SFC (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=860/140) to give the title compound having a longer retention time (73.6 mg).
MS (API+): [M+H]⁺429.1.

15 [0714]

Example 123

N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (203 mg) was resolved by SFC (column: CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon
25 dioxide/methanol=860/140) to give the title compound having a shorter retention time (61.0 mg).

MS (API+): [M+H]⁺443.1.

[0715]

Example 124

30 N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-
35 4(1H)-carboxamide (203 mg) was resolved by SFC (column:

CHIRALPAK IA, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=860/140) to give the title compound having a longer retention time (78.8 mg).

5 MS (API+): [M+H]⁺443.1.

[0716]

Example 125

7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-

10 carboxamide

The title compound was obtained from 2-methoxy-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethan-1-amine hydrochloride in the same manner as in Step L of Example 72-I.

MS (API+): [M+H]⁺469.2.

15 [0717]

Example 126

optically active N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 Racemic N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (243 mg) was resolved by SFC (column: CHIRALPAK IC, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=770/230) to give the title compound having a shorter retention time (81.1 mg).

MS (API+): [M+H]⁺447.1.

[0718]

Example 127

30 optically active N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (243 mg) was resolved by SFC

35

(column: CHIRALPAK IC, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=770/230) to give the title compound having a longer retention time (81.4 mg).

5 MS (API+): [M+H]⁺447.1.

[0719]

Example 128

7-cyclopropyl-N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-
2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
10 carboxamide

Racemic 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (129 mg) was resolved by HPLC (column: CHIRALPAK IC, 50 mmID×500 mmL,
15 manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 600/400) to give the title compound having a shorter retention time (58.2 mg).

MS (API+): [M+H]⁺469.2.

[0720]

20 Example 129

7-cyclopropyl-N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (129 mg) was resolved by HPLC (column: CHIRALPAK IC, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 600/400) to give the title compound
30 having a longer retention time (60.7 mg).

MS (API+): [M+H]⁺469.2.

[0721]

Example 130

N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-
35 6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-

carboxamide

Racemic N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (142 mg) was
5 resolved by SFC (column: CHIRALCEL OJH, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=860/140) to give the title compound having a shorter retention time (53.5 mg).
MS (API+): [M+H]⁺477.2.

10 [0722]

Example 131

N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 Racemic N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (142 mg) was resolved by SFC (column: CHIRALCEL OJH, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile
20 phase: carbon dioxide/methanol=860/140) to give the title compound having a longer retention time (57.3 mg).
MS (API+): [M+H]⁺477.2.

[0723]

Example 132

25 N-(1-(4-bromophenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0724]

A) 1-(4-bromophenyl)-N-hydroxy-2-methoxyethanimine

The title compound was obtained from 2-bromo-1-(4-bromophenyl)ethanone in the same manner as in Step D of
30 Example 63 and Step D of Example 2.

MS (API+): [M+H]⁺244.1.

[0725]

B) 1-(4-bromophenyl)-2-methoxyethanamine

35 A mixture of 1-(4-bromophenyl)-N-hydroxy-2-

methoxyethanimine (7.5 g), 1.1 M borane-tetrahydrofuran complex/tetrahydrofuran solution (55.9 mL) and tetrahydrofuran (300 mL) was stirred at 80°C for 2 days, 1 M aqueous sodium hydroxide solution was added thereto, and the mixture was
5 stirred at 80°C for 3 hr. To the reaction mixture was added saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound
10 containing an impurity (8.44 g).

MS (API+): [M+H]⁺230.1.

[0726]

C) N-(1-(4-bromophenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 The title compound was obtained from 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate and 1-(4-bromophenyl)-2-methoxyethanamine in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺419.1.

20 [0727]

Example 133

N-(2-(2-methoxyethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

25 The title compound was obtained from 4-(trifluoromethoxy)phenacyl bromide in the same manner as in Step D of Example 63, Step D of Example 2, Step B of Example 15 and Step G of Example 7.

MS (API+): [M+H]⁺469.2.

30 [0728]

Example 134

N-(2-methoxy-1-(4-(1H-pyrazol-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0729]

35 A) 1-(4-bromophenyl)-2-methoxyethanone

The title compound was obtained from 2-bromo-1-(4-bromophenyl)ethanone in the same manner as in Step D of Example 63.

¹H NMR (300 MHz, CDCl₃) δ 3.50 (3H, s), 4.65 (2H, s), 7.55-7.68
5 (2H, m), 7.75-7.87 (2H, m).

[0730]

B) 2-(4-bromophenyl)-2-(methoxymethyl)-1,3-dioxolane

A mixture of 1-(4-bromophenyl)-2-methoxyethanone (3.00 g),
p-toluenesulfonic acid monohydrate (0.249 g), ethylene glycol
10 (1.63 g) and toluene (50 mL) was stirred overnight using Dean-Stark at 140°C. The reaction mixture was filtered through silica gel pad (NH), and the solvent was evaporated under reduced pressure to give the title compound (3.97 g).

¹H NMR (300 MHz, CDCl₃) δ 3.39 (3H, s), 3.57 (2H, s), 3.76-3.92
15 (2H, m), 4.02-4.19 (2H, m), 7.35-7.43 (2H, m), 7.43-7.52 (2H, m).

[0731]

C) 1-(4-(2-(methoxymethyl)-1,3-dioxolane-2-yl)phenyl)-1H-pyrazole

20 A mixture of 2-(4-bromophenyl)-2-(methoxymethyl)-1,3-dioxolane (3.5 g), 1H-pyrazole (0.960 g), quinolin-8-ol (0.372 g), copper(I) iodide (0.244 g), potassium carbonate (3.54 g) and dimethyl sulfoxide (50 mL) was stirred overnight at 140°C under nitrogen atmosphere, water was added thereto, and the
25 mixture was extracted with ethyl acetate. The extract was washed successively with saturated aqueous ammonium chloride solution and saturated brine, dried over anhydrous magnesium sulfate, and filtered through NH silica gel pad. The filtrate was concentrated under reduced pressure, and the residue was
30 purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.53 g).

MS (API+): [M+H]⁺261.1.

[0732]

D) 2-methoxy-1-(4-(1H-pyrazol-1-yl)phenyl)ethanone

35 A mixture of 1-(4-(2-(methoxymethyl)-1,3-dioxolane-2-

yl)phenyl)-1H-pyrazole (2.6 g), 1 M hydrochloric acid (15 mL) and tetrahydrofuran (30 mL) was stirred overnight at room temperature, saturated brine was added thereto, and the mixture was extracted with ethyl acetate. The extract was
5 washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (2.15 g).
¹H NMR (300 MHz, CDCl₃) δ 3.52 (3H, s), 4.71 (2H, s), 6.48-6.56 (1H, m), 7.78 (1H, d, J = 1.9 Hz), 7.79-7.87 (2H, m), 8.02 (1H,
10 d, J = 2.6 Hz), 8.03-8.10 (2H, m).

[0733]

E) N-(2-methoxy-1-(4-(1H-pyrazol-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-methoxy-1-(4-(1H-pyrazol-1-yl)phenyl)ethanone in the same manner as in Step D
15 of Example 2, Step B of Example 15 and Step G of Example 7.

MS (API+): [M+H]⁺407.2.

[0734]

Example 135

20 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-3-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0735]

A) methyl N-(5-bromo-3-nitropyridin-2-yl)glycinate

25 The title compound was obtained from 5-bromo-2-chloro-3-nitropyridine in the same manner as in Step A of Example 19.

MS (API+): [M+H]⁺290.0.

[0736]

B) 7-bromo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

30 A mixture of methyl N-(5-bromo-3-nitropyridin-2-yl)glycinate (18.0 g) and 5% platinum-carbon (1.80 g) in tetrahydrofuran (414 mL) was stirred at room temperature for 1 hr under hydrogen atmosphere. The catalyst was filtered off, and the solvent was evaporated under reduced pressure. The
35 residue was dissolved in ethanol (200 mL), the solution was

stirred overnight at 80°C. The reaction mixture was cooled to room temperature, and the solid was collected by filtration, and washed with ethyl acetate to give the title compound (9.60 g).

5 ¹H NMR (300 MHz, DMSO-d₆) δ 3.94 (2H, d, J = 1.7 Hz), 6.99 (1H, s), 7.01 (1H, d, J = 2.1 Hz), 7.65 (1H, d, J = 2.3 Hz), 10.46 (1H, s).

[0737]

C) 7-bromo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-
10 2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]-pyrazine-4(1H)-carboxamide

The title compound was obtained from 7-bromo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner as in Steps C-D of Example 26.

15 MS (API+): [M+H]⁺619.2.

[0738]

D) N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-3-yl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]-
20 pyrazine-4(1H)-carboxamide

To a mixture of 7-bromo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]-pyrazine-4(1H)-carboxamide (100 mg) and 1-methyl-3-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolane-1H-pyrazole (101 mg),
25 tripotassium phosphate (103 mg) in a mixed solvent of toluene (1.51 mL) and water (0.10 mL) was added bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (11.4 mg) at room temperature, the reaction mixture was stirred at 100°C
30 for 3 hr under argon atmosphere. The reaction mixture was cooled to room temperature, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (72 mg).

35 MS (API+): [M+H]⁺621.3.

[0739]

E) N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-3-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 The title compound was obtained from N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-3-yl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]-pyrazine-4(1H)-carboxamide in the same manner as in Step I of Example 63.

10 MS (API+): [M+H]⁺491.2.

[0740]

Example 136

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
15 carboxamide

The title compound was obtained from 7-bromo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]-pyrazine-4(1H)-carboxamide (100 mg) and 1-methyl-4-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolane-1H-pyrazole in the same
20 manner as in Step D of Example 135 and Step I of Example 63.

MS (API+): [M+H]⁺491.2.

[0741]

Example 137

25 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-5-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 7-bromo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]-pyrazine-4(1H)-carboxamide (100 mg) and 1-methyl-5-(4,4,5,5-tetramethyl-[1,3,2]-dioxaborolane-1H-pyrazole in the same
30 manner as in Step D of Example 135 and Step I of Example 63.

MS (API+): [M+H]⁺491.2.

35 [0742]

Example 138

7-isopropoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 The title compound was obtained from 2-iodopropane in the same manner as in Steps H-I of Example 63.

MS (API+): [M+H]⁺469.1.

[0743]

Example 139

10 7-(difluoromethoxy)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0744]

A) 7-(difluoromethoxy)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a mixture of 7-hydroxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (163 mg) and N-benzyl-N,N,N-triethylammonium chloride (20.4 mg) in tetrahydrofuran (2.00 mL) was added 8 M aqueous sodium hydroxide solution (36.3 μ L) at 0°C. The reaction mixture was stirred overnight at room temperature under chloro(difluoro)methane atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (80.9 mg).

30 MS (API+): [M+H]⁺607.2.

[0745]

B) 7-(difluoromethoxy)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-

35

b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 7-(difluoromethoxy)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-

5 b]pyrazine-4(1H)-carboxamide in the same manner as in Step I of Example 63.

MS (API+): [M+H]⁺477.2.

[0746]

Example 140

10 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-oxo-6,7-dihydropteridine-8(5H)-carboxamide

[0747]

A) methyl 2-((2-chloro-5-nitropyrimidin-4-yl)aminoacetate

The title compound was obtained from 2,4-dichloro-5-nitropyrimidine in the same manner as in Step A of Example 1.

15 MS (API-): [M-H]⁻245.0.

[0748]

B) 7,8-dihydropteridin-6(5H)-one

To a solution of methyl 2-((2-chloro-5-nitropyrimidin-4-yl)aminoacetate (7.23 g) in ethanol (293 mL) was added 5% palladium-carbon (containing 50% water, 3.00 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, the catalyst was filtered off, and the solvent was evaporated under reduced pressure. To a solution of the residue in ethanol (293 mL) was added acetic acid (8.39 mL) at room temperature. The reaction mixture was stirred at 80°C for 3 days, and cooled to room temperature, and the solid was collected by filtration. The obtained solid was suspended in saturated aqueous sodium bicarbonate solution, and the solid was collected by filtration, and washed with water to give the title compound (3.91 g).

¹H NMR (300 MHz, DMSO-d₆) δ 4.02 (2H, d, J = 1.5 Hz), 7.66 (1H, s), 7.70 (1H, brs), 8.05 (1H, s), 10.49 (1H, brs).

35 [0749]

D) N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-oxo-6,7-dihydropteridin-8(5H)-carboxamide

The title compound was obtained from 7,8-dihydropteridin-6(5H)-one in the same manner as in Steps F-G of Example 7.

5 MS (API+): [M+H]⁺412.2.

[0750]

Example 141

7-ethoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 The title compound was obtained from iodoethane in the same manner as in Steps H-I of Example 63.

MS (API+): [M+H]⁺455.2.

[0751]

Example 142

15 7-methyl-N-(oxetan-3-yl(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0752]

A) diethyl (4-(trifluoromethoxy)benzyl)malonate

A mixture of 1-(bromomethyl)-4-(trifluoromethoxy)benzene
20 (10.0 g), diethyl malonate (6.91 g), sodium ethoxide (14.7 g) and ethanol (500 mL) was stirred at 80°C for 2 days. To the reaction mixture was added saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over
25 anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (7.46 g).

MS (API-): [M-H]⁻333.1.

30 [0753]

B) 2-(4-(trifluoromethoxy)benzyl)propane-1,3-diol

To a mixture of diethyl (4-(trifluoromethoxy)benzyl)malonate (5.3 g) and tetrahydrofuran (100 mL) was added lithium aluminium hydride (2.41 g) at 0°C.
35 The reaction mixture was stirred at 0°C for 30 min, 1 M

hydrochloric acid was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give
5 a residue (4.32 g). The same reaction was repeated. The obtained residue (total 5.3 g) was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.9 g).

^1H NMR (300 MHz, CDCl_3) δ 1.89–2.12 (1H, m), 2.34 (2H, brs),
10 2.65 (2H, d, J = 7.5 Hz), 3.60–3.72 (2H, m), 3.75–3.87 (2H, m), 7.08–7.17 (2H, m), 7.17–7.24 (2H, m).

[0754]

C) 3-(4-(trifluoromethoxy)benzyl)oxetane

To a mixture of 2-(4-(trifluoromethoxy)benzyl)propane-
15 1,3-diol (2.37 g) and tetrahydrofuran (100 mL) was added 1.6 M n-butyllithium/hexane solution (6.51 mL) at 0°C. The reaction mixture was stirred at 0°C for 30 min under nitrogen atmosphere, p-toluenesulfonyl chloride (1.99 g) was added thereto at room temperature, and the mixture was stirred at room temperature
20 for 1 hr under nitrogen atmosphere. To the reaction mixture was added 1.6 M n-butyllithium/hexane solution (7.10 mL) at room temperature, and the mixture was stirred at 80°C for 4 hr under nitrogen atmosphere. To the reaction mixture was added saturated aqueous ammonium chloride solution, and the mixture
25 was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (0.950 g).

30 ^1H NMR (300 MHz, CDCl_3) δ 3.03 (2H, d, J = 7.9 Hz), 3.15–3.43 (1H, m), 4.46 (2H, t, J = 6.0 Hz), 4.80 (2H, dd, J = 7.5, 6.0 Hz), 7.13 (4H, s).

[0755]

D) 3-(bromo(4-(trifluoromethoxy)phenyl)methyl)oxetane

35 A mixture of 3-(4-(trifluoromethoxy)benzyl)oxetane (950

mg), N-bromosuccinimide (947 mg), 2,2'-azobis(2-methylpropionitrile) (67.2 mg) and benzotrifluoride (20 mL) was stirred overnight at 80°C. To the reaction mixture was added hexane, the insoluble substance was filtered off, and
5 the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (406 mg).
¹H NMR (300 MHz, CDCl₃) δ 3.68-3.87 (1H, m), 4.17 (1H, t, J = 6.4 Hz), 4.54-4.66 (2H, m), 4.92 (1H, dd, J = 7.4, 6.6 Hz),
10 5.32 (1H, d, J = 11.1 Hz), 7.14-7.24 (2H, m), 7.35-7.43 (2H, m).

[0756]

E) 3-(azido(4-(trifluoromethoxy)phenyl)methyl)oxetane

A mixture of 3-(bromo(4-(trifluoromethoxy)phenyl)methyl)oxetane (230 mg), sodium azide
15 (80 mg), water (3 mL) and acetone (12 mL) was stirred overnight at room temperature, saturated aqueous sodium hydrogen carbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with
20 saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (260 mg).

¹H NMR (300 MHz, CDCl₃) δ 3.18-3.41 (1H, m), 4.33 (1H, t, J = 6.2 Hz), 4.54-4.71 (2H, m), 4.80-4.89 (2H, m), 7.21-7.27 (2H,
25 m), 7.29-7.36 (2H, m).

[0757]

F) 7-methyl-N-(oxetan-3-yl(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A mixture of 3-(azido(4-(trifluoromethoxy)phenyl)methyl)oxetane (260 mg), 5%
30 palladium-carbon (containing 50% water, 30 mg) and tetrahydrofuran (5 mL) was stirred at room temperature for 1 hr under hydrogen atmosphere. Then, to the reaction mixture was added methanol (5 mL), the reaction mixture was stirred
35 overnight at room temperature under hydrogen atmosphere. The

insoluble substance was filtered off through Celite, and the filtrate was concentrated under reduced pressure to give oxetan-3-yl(4-(trifluoromethoxy)phenyl)methanamine (209 mg). The title compound was obtained from the obtained oxetan-3-yl(4-(trifluoromethoxy)phenyl)methanamine in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺437.1.

[0758]

Example 143

10 N-(1-(3-fluoro-4-(1H-pyrazol-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0759]

A) 2-bromo-1-(4-bromo-3-fluorophenyl)ethanone

15 To a solution of 1-(4-bromo-3-fluorophenyl)ethanone (2.8 g) (obtained from 4-bromo-3-fluorobenzonitrile in the same manner as in Step B of Example 30) in tetrahydrofuran (150 mL) was added phenyltrimethylammonium tribromide (4.35 g) at 0°C. The reaction mixture was stirred at room temperature for 2 days, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.58 g).

20 ¹H NMR (300 MHz, CDCl₃) δ 4.33-4.42 (2H, m), 7.59-7.77 (3H, m).

[0760]

25 B) N-(1-(3-fluoro-4-(1H-pyrazol-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-bromo-1-(4-bromo-3-fluorophenyl)ethanone in the same manner as in Steps A-D of Example 134, Step D of Example 2, Step B of Example 15 and Step G of Example 7.

MS (API+): [M+H]⁺425.2.

[0761]

Example 144

35 N-(2-methoxy-1-(5-(1H-pyrazol-1-yl)pyridin-2-yl)ethyl)-7-

methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
carboxamide

The title compound was obtained from 5-bromopicolinic acid
in the same manner as in Step A of Example 31, Steps D-F of
5 Example 66, Steps A-D of Example 134, Step D of Example 2,
Step B of Example 15 and Step G of Example 7.

MS (API+): [M+H]⁺408.2.

[0762]

Example 145

10 N-((1R)-2-hydroxy-2-methyl-1-(4-
(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-
(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
15 b]pyrazine-4(1H)-carboxamide (30.1 mg) was resolved by HPLC
(column: CHIRALPAK IC, 50 mmID×500 mmL, manufactured by Daicel
Chemical Industries, Ltd., mobile phase: hexane/ethanol =
800/200) to give the title compound having a shorter retention
time (10.0 mg).

20 MS (API+): [M+H]⁺425.2.

[0763]

Example 146

N-((1S)-2-hydroxy-2-methyl-1-(4-
(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
25 b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-
(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide (30.1 mg) was resolved by HPLC
(column: CHIRALPAK IC, 50 mmID×500 mmL, manufactured by Daicel
30 Chemical Industries, Ltd., mobile phase: hexane/ethanol =
800/200) to give the title compound having a longer retention
time (9.4 mg).

MS (API+): [M+H]⁺425.2.

[0764]

35 Example 147

7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0765]

A) 1-(4-(1H-pyrrol-1-yl)phenyl)propan-1-one

5 To a mixture of 1H-pyrrole (1.38 g), 1-(4-bromophenyl)propan-1-one (3.66 g), trans-N,N'-bismethyl-1,2-cyclohexane (489 mg) and tripotassium phosphate (7.66 g) in toluene (20 mL) was added copper(I) iodide (164 mg). The reaction mixture was heated at reflux for 16 hr under argon
10 atmosphere, ethyl acetate and water were added thereto. The insoluble substance was filtered off, and the organic layer and aqueous layer of the filtrate were separated, and the aqueous layer was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was
15 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.32 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.10 (3H, t, J = 7.3 Hz), 3.06 (2H, q, J = 7.2 Hz), 6.28-6.37 (2H, m), 7.46-7.57 (2H, m), 7.68-
20 7.80 (2H, m), 7.96-8.10 (2H, m).

[0766]

B) 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)propan-1-one in the same manner as in Step C of
25 Example 1, Step B of Example 54 and Step G of Example 7.

MS (API+): [M+H]⁺390.2.

[0767]

Example 148

30 7-methyl-2-oxo-N-(1-(4-(pyrrolidin-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)propan-1-one in the same manner as in Step C of Example 1, Step B of Example 54 and Step G of Example 7.

35 MS (API+): [M+H]⁺394.3.

[0768]

Example 149

7-(2-hydroxypropan-2-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0769]

A) methyl 4-((2-methoxy-1-(4-trifluoromethoxy)phenyl)ethyl) carbamoyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-7-carboxylate

10 A mixture of 7-bromo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (537 mg) and 1,1'-bis(diphenylphosphino)ferrocene dichloropalladium(II) dichloromethanecomplex (70.8 mg), triethylamine (0.242 mL) and methanol (1.5 mL) in N,N-dimethylformamide (15 mL) was stirred at 80°C for 3 hr under carbon monoxide atmosphere (5 atm). The reaction mixture was cooled, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
20 column chromatography (hexane/ethyl acetate) to give the title compound (403 mg).

MS (API+): [M+H]⁺599.3.

[0770]

C) methyl 4-((2-methoxy-1-(4-trifluoromethoxy)phenyl)ethyl) carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-7-carboxylate

The title compound was obtained from methyl 4-((2-methoxy-1-(4-trifluoromethoxy)phenyl)ethyl) carbamoyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-7-carboxylate in the same
30 manner as in Step I of Example 63.

MS (API+): [M+H]⁺469.1.

[0771]

D) 7-(2-hydroxypropan-2-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

To a solution of methyl 4-((2-methoxy-1-(4-trifluoromethoxy)phenyl)ethyl) carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-7-carboxylate (40 mg) in
5 tetrahydrofuran (0.85 mL) was added 1 M methylmagnesium bromide/tetrahydrofuran solution (0.43 mL) at 0°C. The reaction mixture was stirred at 0°C for 2 hr under argon atmosphere, 1 M methylmagnesium bromide/tetrahydrofuran solution (0.85 mL) was added thereto. The reaction mixture was
10 stirred at 0°C for 30 min, saturated aqueous ammonium chloride solution was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The
15 residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (27 mg).
MS (API+): [M+H]⁺469.2.

[0772]

Example 150

20 7-(azetidin-1-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0773]

A) 7-(azetidin-1-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a mixture of 7-bromo-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (300 mg), azetidine (32.6 µL),
30 and cesium carbonate (249 mg) in toluene (4.84 mL) were added (9,9-dimethyl-9H-xanthene-4,5-diyl)bis(diphenylphosphine) (28.0 mg) and tris(dibenzylideneacetone)dipalladium(0) (29.6
35 mg), the reaction mixture was stirred overnight at 110°C under

nitrogen atmosphere. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue
5 was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (19.3 mg).

MS (API+): [M+H]⁺596.3.

[0774]

B) 7-(azetidin-1-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
10

The title compound was obtained from 7-(azetidin-1-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Step I
15 of Example 63.

MS (API+): [M+H]⁺466.2.

[0775]

Example 151

20 N-(1-(imidazo[1,2-a]pyridin-7-yl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from imidazo[1,2-a]pyridine-7-carboxylic acid in the same manner as in Step A of Example 31, Step B of Example 30, Step D of Example 2, Step
25 B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺365.2.

[0776]

Example 152

7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)cyclopropyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
30

[0777]

A) 1-(4-(trifluoromethoxy)phenyl)cyclopropaneamine

To a mixture of 4-(trifluoromethoxy)benzonitrile (2.02 g) and titanium(IV) tetraisopropoxide (3.48 mL) in diethyl ether
35 (50 mL) was slowly added 3 M ethylmagnesium

bromide/tetrahydrofuran solution (7.91 mL) over 50 min at -78°C. The reaction mixture was stirred at room temperature for 1 hr, boron trifluoride diethyl ether complex (3.01 mL) was added thereto. The reaction mixture was stirred at room temperature
5 for 16 hr, 1 M hydrochloric acid and diethyl ether were added thereto. Then, 2 M aqueous sodium hydroxide solution was added thereto, and the mixture was extracted with diethyl ether. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the
10 title compound (1.51 g).

MS (API+): [M+H]⁺218.1.

[0778]

B) 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)cyclopropyl)-2,3-dihydropyrido[2,3-
15 b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(trifluoromethoxy)phenyl)cyclopropaneamine in the same manner as in Step G of Example 7.

MS (API+): [M+H]⁺407.1.

20 [0779]

Example 153

7-fluoro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0780]

25 A) methyl N-(5-fluoro-3-nitropyridin-2-yl)glycinate

To a solution of methyl glycinate hydrochloride (1.39 g) and 2-chloro-5-fluoro-3-nitropyridine (2.00 g) in ethanol (35.9 mL) was added triethylamine (3.79 mL). The reaction mixture was stirred overnight at 80°C, and the solvent was
30 evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column
35 chromatography (hexane/ethyl acetate) to give the title

compound (93.0 mg).

MS (API+): [M+H]⁺230.1.

[0781]

B) 7-fluoro-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

5 To a solution of methyl N-(5-fluoro-3-nitropyridin-2-yl)glycinate (90.0 mg) in tetrahydrofuran (6.66 mL) was added 5% platinum-carbon (10.0 mg). The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated
10 under reduced pressure. The residue was dissolved in ethanol (5.00 mL), and the mixture was stirred at 87°C for 1 hr. To the reaction mixture was added 2 M hydrogen chloride/methanol solution (1.00 mL), the mixture was stirred overnight at 87°C, and the solvent was evaporated under reduced pressure. The
15 residue was diluted with ethyl acetate, and the mixture was neutralized with saturated aqueous sodium hydrogen carbonate solution, and extracted with ethyl acetate. The extract was washed with saturated brine, and the solvent was evaporated under reduced pressure to give the title compound (39.1 mg).

20 MS (API+): [M+H]⁺168.1.

[0782]

C) 7-fluoro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a mixture of 7-fluoro-3,4-dihydropyrido[2,3-b]pyrazin-
25 2(1H)-one (39.1 mg) and triethylamine (99.0 µL) in a mixed solvent of tetrahydrofuran (2.00 mL) and N,N-dimethylacetamide (2.00 mL) was added a solution of bis(trichloromethyl)carbonate (69.4 mg) in tetrahydrofuran (1.00 mL) at 0°C. The reaction mixture was stirred at 0°C for
30 2 hr, a mixture of triethylamine (165 µL) and 2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethanamine hydrochloride (318 mg) in a mixed solvent of tetrahydrofuran (1.00 mL) and N,N-dimethylacetamide (2.00 mL) was added thereto at 0°C. The reaction mixture was stirred overnight at room temperature,
35 water was added thereto, and the mixture was extracted with

ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography
5 (hexane/ethyl acetate), and crystallized from hexane/ethyl acetate to give the title compound (10.1 mg).

MS (API+): [M+H]⁺429.1.

[0783]

Example 154

10 N-(2-methoxy-1-(4-(1H-pyrrol-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0784]

A) 2-bromo-1-(4-iodophenyl)ethanone

The title compound was obtained from 1-(4-iodophenyl)ethanone in the same manner as in Step F of Example
15 66.

¹H NMR (300 MHz, DMSO-d₆) δ 4.91 (2H, s), 7.72-7.80 (2H, m), 7.91-8.00 (2H, m).

[0785]

20 B) 1-(4-iodophenyl)-2-methoxyethanone

The title compound was obtained from 2-bromo-1-(4-iodophenyl)ethanone in the same manner as in Step D of Example
63.

¹H NMR (300 MHz, DMSO-d₆) δ 3.35 (3H, s), 4.75 (2H, s), 7.63-
25 7.73 (2H, m), 7.88-8.00 (2H, m).

[0786]

C) 1-(4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethanone

The title compound was obtained from 1H-pyrrole and 1-(4-iodophenyl)-2-methoxyethanone in the same manner as in Step A
30 of Example 147.

¹H NMR (300 MHz, DMSO-d₆) δ 3.37 (3H, s), 4.79 (2H, s), 6.29-6.41 (2H, m), 7.50-7.58 (2H, m), 7.71-7.82 (2H, m), 7.95-8.05 (2H, m).

[0787]

35 D) N-(2-methoxy-1-(4-(1H-pyrrol-1-yl)phenyl)ethyl)-7-methyl-2-

oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethanone in the same manner as in Step C of Example 1, Step B of Example 54 and Step G of Example 7.

5 MS (API+): [M+H]⁺406.2.

[0788]

Example 155

7-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 [0789]

A) methyl N-(5-chloro-3-nitropyridin-2-yl)glycinate

To a solution of methyl N-(5-bromo-3-nitropyridin-2-yl)glycinate (400 mg) in 1-methylpyrrolidin-2-one (13.8 mL) was added copper(I) chloride (410 mg). The reaction mixture
15 was stirred at 150°C for 2.5 hr under microwave irradiation, and water was added thereto. The insoluble substance was filtered off, and the filtrate was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and
20 the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and crystallized from diisopropyl ether to give the title compound (181 mg).

MS (API+): [M+H]⁺246.0.

25 [0790]

B) 7-chloro-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-chloro-3-nitropyridin-2-yl)glycinate (181 mg) in tetrahydrofuran (12.5 mL) was added 5% platinum-carbon (20.0 mg). The reaction mixture was stirred
30 at room temperature for 2 hr under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol (10.0 mL), and the solution was stirred overnight at 87°C. To the reaction mixture was added 2 M hydrogen chloride/methanol
35 solution (1.00 mL), and the mixture was stirred at 87°C for 10

min, and the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was neutralized with saturated aqueous sodium hydrogen

5 extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (132 mg).

MS (API+): [M+H]⁺184.0.

[0791]

10 C) 7-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 7-chloro-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner as in Step C of Example 153.

15 MS (API+): [M+H]⁺445.1.

[0792]

Example 156

N-(2-methoxy-1-(4-(pyrrolidin-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethanone in the same manner as in Step C of Example 1, Step E of Example 2 and Step G of Example 7.

MS (API+): [M+H]⁺410.3.

[0793]

25 Example 157

N-(2-methoxy-1-(4-(1H-pyrrol-1-yl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethanone in the same manner as in Step C
30 of Example 1, Step E of Example 2 and Step G of Example 7.

MS (API+): [M+H]⁺392.2.

[0794]

Example 158

N-(2-methoxy-1-(4-(pyrrolidin-1-yl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
35

The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethanone in the same manner as in Step C of Example 1, Step B of Example 54 and Step G of Example 7.
MS (API+): [M+H]⁺396.2.

5 [0795]

Example 159

N-(1-(2,2-difluoro-1,3-benzodioxol-5-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0796]

10 A) 1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)propan-1-ol

To a solution of 2,2-difluorobenzo[d][1,3]dioxol-5-carbaldehyde (2.35 g) in tetrahydrofuran (50 mL) was slowly added 3 M ethylmagnesium bromide/tetrahydrofuran solution (6.30 mL) at 0°C. The reaction mixture was stirred at room
15 temperature for 16 hr, saturated ammonium chloride solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography
20 (hexane/ethyl acetate) to give the title compound (2.36 g).
¹H NMR (300 MHz, DMSO-d₆) δ 0.81 (3H, t, J = 7.3 Hz), 1.51-1.67 (2H, m), 4.42-4.53 (1H, m), 5.27 (1H, d, J = 4.5 Hz), 7.10-7.18 (1H, m), 7.27-7.36 (2H, m).
[0797]

25 B) 1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)propan-1-one

To a solution of 1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)propan-1-ol (2.36 g) in toluene (50 mL) was added manganese dioxide (2.36 g). The reaction mixture was stirred at room
30 temperature for 3 days, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (0.492 g).
¹H NMR (300 MHz, DMSO-d₆) δ 1.08 (3H, t, J = 7.2 Hz), 3.05 (2H, q, J = 7.1 Hz), 7.55 (1H, d, J = 8.3 Hz), 7.91 (1H, dd, J =
35 8.5, 1.7 Hz), 7.96 (1H, d, J = 1.7 Hz).

[0798]

C) N-(1-(2,2-difluoro-1,3-benzodioxol-5-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)propan-1-one in the same manner as in Step C of Example 1, Step E of Example 2 and Step G of Example 7.

MS (API+): [M+H]⁺391.1.

[0799]

10 Example 160

7-methyl-2-oxo-N-(2-(1H-pyrazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0800]

15 A) 2-(1H-pyrazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone

A mixture of 1H-pyrazole (0.722 g), 60% sodium hydride (459 mg) and N,N-dimethylformamide (50 mL) was stirred at 0°C for 10 min, 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone (2.5 g) was added thereto, and the mixture was stirred at 0°C for 20 min under nitrogen atmosphere. To the reaction mixture was added saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (598 mg).

MS (API+): [M+H]⁺271.1.

[0801]

30 B) 7-methyl-2-oxo-N-(2-(1H-pyrazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-(1H-pyrazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone in the same manner as in Step D of Example 2, Step B of Example 15 and Step G of

Example 7.

MS (API+): [M+H]⁺461.2.

[0802]

Example 161

5 optically active 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
10 carboxamide (244 mg) was resolved by HPLC (column: CHIRALPAK IC, 4.6 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: ethanol) to give the title compound having a shorter retention time (93.7 mg).

MS (API+): [M+H]⁺390.2.

15 [0803]

Example 162

7-methyl-2-oxo-N-(1-(5-(1H-pyrazol-1-yl)-2-thienyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 5-bromothiophene-2-
20 carboxylic acid in the same manner as in Step A of Example 31, Step C of Example 134, Step B of Example 30, Step D of Example 2, Step B of Example 15 and Step G of Example 7.

MS (API+): [M+H]⁺397.0.

[0804]

25 Example 163

optically active 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
30 carboxamide (244 mg) was resolved by HPLC (column: CHIRALPAK IC, 4.6 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: ethanol) to give the title compound having a longer retention time (93.0 mg).

35 MS (API+): [M+H]⁺390.2.

[0805]

Example 164

7-methyl-2-oxo-N-(2-(1H-1,2,4-triazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone in the same manner as in Step A of Example 160, Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺462.2.

[0806]

Example 165

7-methyl-2-oxo-N-(1-(pyrazolo[1,5-a]pyridin-5-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0807]

A) diethyl pyrazolo[1,5-a]pyridine-3,5-dicarboxylate

To a mixture of (aminooxy)(hydroxy)sulfane dioxide (15.0 g) and water (60 mL) was added saturated aqueous sodium hydrogen carbonate solution at 0°C to adjust the pH of the mixture to 9-10, and a mixture of ethyl isonicotinate (10.0 g) and methanol (250 mL) was added thereto at 0°C. The reaction mixture was stirred overnight at 70°C, N,N-dimethylformamide was added thereto, and the solvent was evaporated under reduced pressure. To the residue were added potassium carbonate (27.4 g), N,N-dimethylformamide (250 mL) and ethyl propynoate (7.14 g) at 0°C, and the reaction mixture was stirred at room temperature for 3 hr. To the reaction mixture was added saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.01 g).

MS (API+): [M+H]⁺263.1.

[0808]

B) pyrazolo[1,5-a]pyridine-5-carboxylic acid

A mixture of diethyl pyrazolo[1,5-a]pyridine-3,5-dicarboxylate (6.7 g), conc. sulfuric acid (50 mL) and water (50 mL) was stirred overnight at 100°C. The reaction mixture
5 was cooled to room temperature, and the pH of the mixture was adjusted to about 3 with 8 M aqueous sodium hydroxide solution. The obtained solid was collected by filtration, and washed with water to give the title compound (3.15 g).

MS (API-): [M-H]⁻161.1.

10 [0809]

C) 7-methyl-2-oxo-N-(1-(pyrazolo[1,5-a]pyridin-5-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from pyrazolo[1,5-a]pyridine-5-carboxylic acid in the same manner as in Step A
15 of Example 31, Step B of Example 30, Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺365.2.

[0810]

Example 166

20 7-methyl-N-(2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0811]

A) (2-oxo-2-(4-(trifluoromethoxy)phenyl)ethyl)triphenylphosphonium bromide
25

To a solution of triphenylphosphine (6.29 g) in diethyl ether (20 mL) was slowly added a solution of 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone (6.68 g) in diethyl ether (10 mL) at room temperature. The reaction mixture was stirred
30 at room temperature for 1 hr, and the resulting solid was collected by filtration, washed with diethyl ether to give the title compound (9.47 g).

MS (API+), found: 465.3.

[0812]

35 B) 2-(oxetan-3-ylidene)-1-(4-(trifluoromethoxy)phenyl)ethanone

A mixture of (2-oxo-2-(4-(trifluoromethoxy)phenyl)ethyl)triphenylphosphonium bromide (6.54 g) and potassium tert-butoxide (1.35 g) in tetrahydrofuran (50 mL) was stirred at 0°C for 15 min. The
5 reaction mixture was stirred at room temperature for 30 min, oxetan-3-one (721 mg) was added thereto, and the mixture was stirred overnight at 70°C under argon atmosphere,. The reaction mixture was cooled, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
10 column chromatography (hexane/ethyl acetate) to give the title compound (2.18 g).

¹H NMR (300 MHz, CDCl₃) δ 5.39-5.48 (2H, m), 5.64-5.72 (2H, m), 6.77-6.85 (1H, m), 7.28-7.36 (2H, m), 7.93-8.02 (2H, m).

[0813]

15 C) 2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone

To a solution of 2-(oxetan-3-ylidene)-1-(4-(trifluoromethoxy)phenyl)ethanone (1.03 g) in ethanol (35 mL) was added 10% palladium-carbon (containing 50% water, 200 mg). The reaction mixture was stirred at room temperature for 1 hr
20 under hydrogen atmosphere, the insoluble substance was filtered off, and the solvent was evaporated under reduced pressure. To a solution of the residue (1.07 g) in acetonitrile (20.4 mL) was added Dess-Martin periodinane (1.90 g) at room temperature. The reaction mixture was stirred at
25 room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (900 mg).

MS (API+): [M+H]⁺261.1.

30 [0814]

D) N-hydroxy-2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethanimine

The title compound was obtained from 2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone in the same manner as in
35 Step C of Example 1.

MS (API+): [M+H]⁺276.1.

[0815]

E) 7-methyl-N-(2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-hydroxy-2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethanimine (275 mg) in ethanol (5 mL) was added 5% palladium-carbon (containing 50% water, 120 mg) at room temperature. The reaction mixture was stirred at room temperature for 5 hr under hydrogen atmosphere, and 20% palladium hydroxide-carbon (30 mg) was added thereto. The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the solvent was evaporated under reduced pressure to give 2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine (260 mg). To a solution of the obtained 2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine in N,N-dimethylformamide (5.06 mL) were added 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4-carboxylate (166 mg) and triethylamine (141 µL) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) and then silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (92 mg).

MS (API+): [M+H]⁺451.2.

[0816]

Example 167

7-(difluoromethyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0817]

A) 6-hydroxy-5-nitronicotinic acid

To a mixture of 2-chloro-5-methyl-3-nitropyridine (25.0 g) in conc. sulfuric acid (540 mL) was slowly added sodium dichromate dihydrate (64.8 g). The reaction mixture was stirred at room temperature for 4 hr, poured into crushed ice, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained solid was triturated with diisopropyl ether/hexane, and washed with diisopropyl ether/hexane to give the title compound (4.00 g).
¹H NMR (300 MHz, DMSO-d₆) δ 8.37 (1H, d, J = 2.6 Hz), 8.64 (1H, d, J = 2.6 Hz), 13.33 (2H, brs).

[0818]

B) 6-chloro-N-methoxy-N-methyl-5-nitronicotinamide

To a mixture of 6-hydroxy-5-nitronicotinic acid (552 mg) and thionyl chloride (6.57 mL) was added N,N-dimethylformamide (two drops). The reaction mixture was stirred overnight at 70°C, and the solvent was evaporated under reduced pressure. To a solution of the residue in N,N-dimethylacetamide (15.0 mL) was added N,O-dimethylhydroxylamine hydrochloride (322 mg) at 0°C. The reaction mixture was stirred at 0°C for 2 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (647 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 3.28-3.36 (3H, m), 3.58 (3H, s), 8.77 (1H, d, J = 2.1 Hz), 8.94 (1H, d, J = 2.1 Hz).

[0819]

C) methyl 2-((5-(methoxy (methyl) carbamoyl)-3-nitropyridin-2-yl) amino)acetate

The title compound was obtained from 6-chloro-N-methoxy-N-methyl-5-nitronicotinamide in the same manner as in Step A

of Example 1.

¹H NMR (300 MHz, DMSO-d₆) δ 3.28 (3H, s), 3.63 (3H, s), 3.67 (3H, s), 4.37 (2H, d, J = 5.7 Hz), 8.68-8.80 (2H, m), 9.00-9.12 (1H, m).

5 [0820]

D) methyl 2-((5-formyl-3-nitropyridin-2-yl) amino)acetate

To a solution of methyl 2-((5-(methoxy (methyl) carbamoyl)-3-nitropyridin-2-yl) amino)acetate (250 mg) in tetrahydrofuran (8.38 mL) was slowly added 1.5 M
10 diisobutylaluminium hydride/toluene solution (1.12 mL) at -78°C. The reaction mixture was stirred at -78°C for 1 hr under argon atmosphere, and methanol (170 µL) was added thereto. The reaction mixture was stirred for 5 min, silica gel was added thereto, and the solvent was evaporated under reduced pressure.
15 The residue was purified by column chromatography (hexane/ethyl acetate) to give the title compound (72 mg).

MS (API-): [M-H]⁻238.1.

[0821]

E) methyl 2-((5-(difluoromethyl)-3-nitropyridin-2-yl) amino)acetate
20 amino)acetate

To a solution of methyl 2-((5-formyl-3-nitropyridin-2-yl) amino)acetate (4.71 g) in acetonitrile (150 mL) was added N,N-diethylaminosulfur trifluoride (20.8 mL) at room temperature. The reaction mixture was stirred at room temperature for 2 hr,
25 N,N-diethylaminosulfur trifluoride (20.8 mL) was added thereto at room temperature. The reaction mixture was stirred at room temperature for 1 hr, saturated aqueous sodium hydrogen carbonate solution was slowly added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was
30 washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (2.78 g).

35 ¹H NMR (300 MHz, CDCl₃) δ 3.81 (3H, s), 4.43 (2H, d, J = 5.3

Hz), 6.44-6.88 (1H, m), 8.51-8.61 (2H, m), 8.65 (1H, brs).

[0822]

F) 7-(difluoromethyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

5 The title compound was obtained from methyl 2-((5-(difluoromethyl)-3-nitropyridin-2-yl) amino)acetate in the same manner as in Step B of Example 135.

¹H NMR (300 MHz, DMSO-d₆) δ 3.99 (2H, s), 6.63-7.12 (1H, m), 7.02 (1H, s), 7.27 (1H, s), 7.74-7.82 (1H, m), 10.55 (1H, s).

10 [0823]

G) 7-(difluoromethyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 7-(difluoromethyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner as in Step F of Example 1.

MS (API+): [M+H]⁺461.1.

[0824]

Example 168

20 7-methyl-2-oxo-N-(tetrahydro-2H-pyran-4-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0825]

A) N-methoxy-N-methyltetrahydro-2H-pyran-4-carboxamide

25 To a solution of tetrahydro-2H-pyran-4-carboxylic acid (800 mg) in N,N-dimethylformamide (60.0 mL) were added 1-hydroxybenzotriazole monohydrate (1.17 g), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.44 g), N,O-dimethylhydroxylamine hydrochloride (600 mg) and triethylamine (1.73 mL). The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced

35

pressure to give the title compound (831 mg).

MS (API+): [M+H]⁺174.3.

[0826]

B) tetrahydro-2H-pyran-4-yl(4-

5 (trifluoromethoxy)phenyl)methanone

To a solution of 1-bromo-4-(trifluoromethoxy)benzene (2.78 g) in tetrahydrofuran (76.8 mL) was slowly added 1.6 M n-butyllithium/hexane solution (7.20 mL) at -78°C. The reaction mixture was stirred at -78°C for 30 min under nitrogen
10 atmosphere, a solution of N-methoxy-N-methyltetrahydro-2H-pyran-4-carboxamide (665 mg) in tetrahydrofuran (1.00 mL) was added thereto at -78°C. The reaction mixture was stirred at room temperature for 2 hr under nitrogen atmosphere, saturated aqueous ammonium chloride solution was added thereto, and the
15 mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (812 mg).

20 MS (API+): [M+H]⁺275.1.

[0827]

C) N-hydroxy-1-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)methanimine

To a mixture of tetrahydro-2H-pyran-4-yl(4-(trifluoromethoxy)phenyl)methanone (690 mg) and hydroxylamine hydrochloride (713 mg) in ethanol (16.8 mL) was added triethylamine (1.42 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, and the solvent was evaporated under reduced pressure. The residue was
30 diluted with ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl
35 acetate) to give the title compound (566 mg).

MS (API+): [M+H]⁺290.1.

[0828]

D) 1-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)methanamine

5 To a solution of N-hydroxy-1-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)methanimine (566 mg) in tetrahydrofuran (25.0 mL) was added 1.1 M borane-tetrahydrofuran complex/tetrahydrofuran solution (5.33 mL) at room temperature. The reaction mixture was stirred overnight
10 at 70°C, 1 M aqueous sodium hydroxide solution (19.6 mL) was added thereto, and the mixture was stirred at 60°C for 3 hr. To the reaction mixture was added saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium
15 sulfate, and the solvent was evaporated under reduced pressure to give the title compound (547 mg).

MS (API+): [M+H]⁺276.1.

[0829]

E) 7-methyl-2-oxo-N-(tetrahydro-2H-pyran-4-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
20

The title compound was obtained from 1-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)methanamine in the same manner as in Step B of Example 83.

25 MS (API+): [M+H]⁺465.2.

[0830]

Example 169

N-(2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
30

[0831]

A) 2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone

To a solution of diisopropylamine (3.36 mL) in
35 tetrahydrofuran (100 mL) was slowly added 1.6 M n-

butyllithium/hexane solution (15 mL) at -78°C. The reaction mixture was stirred at -78°C for 30 min under argon atmosphere, and 4-trifluoromethoxyacetophenone (3.19 mL) was added thereto at -78°C. The reaction mixture was stirred at -78°C for 1 hr
5 under argon atmosphere, and tetrahydro-4H-pyran-4-one (1.85 mL) was added thereto at -78°C. The reaction mixture was stirred at -78°C for 1 hr under argon atmosphere, saturated aqueous ammonium chloride solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was
10 washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (4.34 g).

15 MS (API-): [M-H]⁻303.1.

[0832]

B) N-(2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 The title compound was obtained from 2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone in the same manner as in Steps C-D of Example 1 and Step L of Example 72-I.

MS (API+): [M+H]⁺495.1.

25 [0833]

Example 170

7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 [0834]

A) tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methanone

To a solution of 1-bromo-4-(trifluoromethoxy)benzene (7.6 g) in tetrahydrofuran (50 mL) was slowly added 1.6 M n-butyllithium/hexane solution (19.7 mL) at -78°C. The reaction
35 mixture was stirred at -78°C for 40 min under nitrogen

atmosphere, and N-methoxy-N-methyltetrahydrofuran-2-carboxamide (1.673 g) (obtained from tetrahydrofuran-2-carboxylic acid in the same manner as in Step A of Example 31) was added thereto at -78°C, and the reaction mixture was
5 stirred at room temperature for 1 hr under nitrogen atmosphere. To the reaction mixture was added saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was
10 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.49 g).

MS (API+): [M+H]⁺261.1.

[0835]

15 B) 7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methanone in the same manner as
20 in Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺451.2.

[0836]

Example 171

25 N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)pentyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 4-(trifluoromethoxy)benzaldehyde in the same manner as in
Example 174.

30 MS (API+): [M+H]⁺453.2.

[0837]

Example 172

N-(3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-
35 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0838]

A) methyl 3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethylpropanoate

A mixture of methyl 3-hydroxy-2,2-dimethylpropanoate
5 (18.3 g), tert-butyl(chloro)diphenylsilane (40.0 g), 1H-imidazole (12.3 g) and N,N-dimethylformamide (300 mL) was stirred overnight at room temperature, and the solvent was evaporated under reduced pressure. To the residue was added saturated brine, and the mixture was extracted with ethyl
10 acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (50.6 g).
15 ¹H NMR (300 MHz, CDCl₃) δ 1.03 (9H, s), 1.20 (6H, s), 3.64 (2H, s), 3.67 (3H, s), 7.31-7.47 (6H, m), 7.58-7.68 (4H, m).

[0839]

B) 3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethylpropanoic acid

20 A mixture of methyl 3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethylpropanoate (50.6 g), 8 M aqueous sodium hydroxide solution (150 mL) and tetrahydrofuran (150 mL) was stirred at 100°C for 3 hr, and the pH of the mixture was adjusted to about 1 with 6 M hydrochloric acid, and the mixture was extracted
25 with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (50.1 g).

MS (API-): [M-H]⁻355.2.

30 [0840]

C) N-(3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-((tert-
35 butyl(diphenyl)silyl)oxy)-2,2-dimethylpropanoic acid in the

same manner as in Step A of Example 170, Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺691.4.

[0841]

5 Example 173

N-(2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0842]

10 A) tert-butyl (2-(1,3-dioxoisoindolin-2-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

A mixture of tert-butyl (2-amino-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (762 mg) and isobenzofuran-1,3-dione (423 mg) in toluene (20 mL) was heated at reflex using Dean-Stark for 16 hr, and the solvent was evaporated under reduced pressure. The obtained solid was triturated with ethyl acetate, collected by filtration, and washed with ethyl acetate to give the title compound (399 mg). MS (API+), found: 351.1.

20 [0843]

B) 2-(2-amino-2-(4-(trifluoromethoxy)phenyl)ethyl)isoindoline-1,3-dione hydrochloride

The title compound was obtained from tert-butyl (2-(1,3-dioxoisoindolin-2-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate in the same manner as in Step E of Example 18.

¹H NMR (300 MHz, DMSO-d₆) δ 3.89-4.01 (1H, m), 4.03-4.16 (1H, m), 4.67 (1H, t, J = 6.9 Hz), 7.41-7.51 (2H, m), 7.64-7.74 (2H, m), 7.87 (4H, d, J = 2.8 Hz), 8.65 (3H, brs).

30 [0844]

C) N-(2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-(2-amino-2-(4-(trifluoromethoxy)phenyl)ethyl)isoindoline-1,3-dione

hydrochloride in the same manner as in Step L of Example 72-I.
MS (API+): [M+H]⁺526.2.

[0845]

Example 174

5 N-(2-cyclopropyl-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0846]

A) tert-butyl (cyano(4-(trifluoromethoxy)phenyl)methyl)carbamate

10 To 2 M ammonia/methanol solution (112 mL) of 4-(trifluoromethoxy)benzaldehyde (10.6 g) was added titanium(IV) tetraisopropoxide (16.3 mL) at 0°C. The reaction mixture was stirred at 0°C for 15 min, trimethylsilyl cyanide (11.2 mL) was added thereto, and the reaction mixture was stirred at room
15 temperature for 20 hr, and concentrated to about 1/4 of volume under reduced pressure, saturated aqueous sodium hydrogen carbonate solution was added thereto, the insoluble substance was filtered off using Celite, and the filtrate was extracted with ethyl acetate. The extract was washed successively with
20 water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To a solution of the residue in tetrahydrofuran (300 mL) were added di-tert-butyl dicarbonate (12.94 mL) and triethylamine (9.32 mL). The reaction mixture was stirred at 45°C for 16 hr,
25 water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography
30 (hexane/ethyl acetate) to give the title compound (9.39 g).
MS (API+): [M+H]⁺317.1.

[0847]

B) tert-butyl (2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

35 To a mixture of tert-butyl (cyano(4-

(trifluoromethoxy)phenyl)methyl)carbamate (3.00 g) and potassium carbonate (1.31 g) in dimethyl sulfoxide (70 mL) was added 35% aqueous hydrogen peroxide (1.66 mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hr, 5 water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography 10 (hexane/ethyl acetate) to give the title compound (2.44 g). ¹H NMR (300 MHz, CDCl₃) δ 1.41 (9H, s), 5.11-5.28 (1H, m), 5.48 (1H, brs), 5.68 (1H, brs), 5.78 (1H, brs), 7.18-7.25 (2H, m), 7.40-7.47 (2H, m).

[0848]

15 C) tert-butyl (2-(methoxy (methyl) amino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

To a solution of tert-butyl (2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (2.44 g) in methanol (70 mL) was added 8 M aqueous sodium hydroxide solution (1.8 20 mL), the reaction mixture was heated at reflux for 20 hr. The solvent was evaporated under reduced pressure, the pH of the residue was adjusted to 3-4 with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried 25 over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. A mixture of the obtained residue, N,O-dimethylhydroxylamine hydrochloride (0.783 g), triethylamine (1.32 mL), 1-hydroxybenzotriazole monohydrate (1.45 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide 30 hydrochloride (1.82 g) in N,N-dimethylformamide (30 mL) was stirred at room temperature for 3 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was 35 evaporated under reduced pressure. The residue was purified by

silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.39 g).

¹H NMR (300 MHz, CDCl₃) δ 1.42 (9H, s), 3.19 (3H, s), 3.51 (3H, s), 5.63-5.91 (2H, m), 7.14-7.22 (2H, m), 7.37-7.45 (2H, m).

5 [0849]

D) tert-butyl (2-cyclopropyl-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate

To a solution of tert-butyl (2-(methoxy(methyl)amino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (685 mg) in
10 tetrahydrofuran (15 mL) was added 0.7 M cyclopropylmagnesium bromide/tetrahydrofuran solution (7.76 mL) at 0°C. The reaction mixture was stirred at room temperature for 20 hr, 1 M hydrochloric acid was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed
15 successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (486 mg).

20 ¹H NMR (300 MHz, CDCl₃) δ 0.75-0.86 (1H, m), 0.90-1.05 (2H, m), 1.08-1.17 (1H, m), 1.40 (9H, s), 1.79-1.90 (1H, m), 5.48 (1H, d, J = 6.4 Hz), 5.92-6.04 (1H, m), 7.18-7.25 (2H, m), 7.33-7.41 (2H, m).

[0850]

25 E) N-(2-cyclopropyl-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A mixture of tert-butyl (2-cyclopropyl-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (200 mg) and 4 M
30 hydrogen chloride/ethyl acetate solution (5 mL) was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. To a mixture of the residue and 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (160 mg) in N,N-dimethylformamide (5 mL) was added
35 triethylamine (0.213 mL). The reaction mixture was stirred at

room temperature for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was
5 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and recrystallized from hexane/ethyl acetate to give the title compound (143 mg).

MS (API+): [M+H]⁺435.1.

10 [0851]

Example 175

N-(2-cyclopropyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 To a mixture of N-(2-cyclopropyl-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (126 mg) in a mixed solvent of methanol (3 mL) and tetrahydrofuran (1 mL) was added sodium borohydride (13.2 mg) at 0°C. The reaction mixture was stirred
20 at room temperature for 3 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was
25 crystallized from hexane/ethyl acetate to give the title compound (87.2 mg).

MS (API+): [M+H]⁺ 437.2.

[0852]

Example 176

30 7-methyl-2-oxo-N-(2-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0853]

A) 4-(2-azido-2-(4-(trifluoromethoxy)phenyl)ethyl)tetrahydro-
35 2H-pyran

To a solution of 2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethanone (304 mg) in pyridine (5 mL) was added thionyl chloride (102 μ L) at 0°C. The reaction mixture was stirred at 0°C for 10 min, water was added thereto
5 at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. A mixture of the residue (286 mg) and 10% palladium-carbon (containing 50% water, 60 mg) in ethanol
10 (5 mL) was stirred at room temperature for 2 hr under hydrogen atmosphere. The insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. To a solution of the residue (290 mg) in tetrahydrofuran (5 mL) were added triphenylphosphine (315 mg), diphenylphosphoryl
15 azide (259 μ L) and 1.9 M diisopropyl azodicarboxylate/toluene solution (632 μ L) at room temperature. The reaction mixture was stirred at room temperature for 30 min, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl
20 acetate) to give the title compound (274 mg).

^1H NMR (300 MHz, CDCl_3) δ 1.20-1.44 (2H, m), 1.52-1.72 (4H, m), 1.73-1.87 (1H, m), 3.30-3.43 (2H, m), 3.89-4.01 (2H, m), 4.48-4.59 (1H, m), 7.19-7.46 (4H, m).

[0854]

25 B) 7-methyl-2-oxo-N-(2-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A mixture of 4-(2-azido-2-(4-(trifluoromethoxy)phenyl)ethyl)tetrahydro-2H-pyran (274 mg)
30 and 10% palladium-carbon (containing 50% water, 92 mg) in ethanol (4.4 mL) was stirred at room temperature for 1 hr under hydrogen atmosphere. The catalyst was filtered off, and the solvent was evaporated under reduced pressure to give 2-(tetrahydro-2H-pyran-4-yl)-1-(4-
35 (trifluoromethoxy)phenyl)ethanamine (254 mg). To a mixture of

4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (144 mg) in N,N-dimethylformamide (2.19 mL) were added 2-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethanamine (254 mg) obtained above
5 and triethylamine (122 μ L) at room temperature. The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was
10 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (184 mg).

MS (API+): $[M+H]^+$ 479.2.

[0855]

15 Example 177

2-oxo-N-(2-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-(2-(methoxy (methyl) amino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-
20 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (290 mg) in tetrahydrofuran (7 mL) was added 1 M methylmagnesium bromide/tetrahydrofuran solution (7.0 mL) at 0°C. The reaction mixture was stirred at room temperature for 20 hr, 1 M hydrochloric acid was added thereto, and the mixture was
25 extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and
30 recrystallized from hexane/ethyl acetate to give the title compound (174 mg).

MS (API+): $[M+H]^+$ 409.1.

[0856]

Example 178

35 8-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-

oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-chloro-4-methoxy-3-nitropyridine in the same manner as in Steps A-B of Example 1 and Steps C-D of Example 19.

5 MS (API+): [M+H]⁺441.2.

[0857]

Example 179

N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 The title compound was obtained from 2-oxo-N-(2-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Example 175.

MS (API+): [M+H]⁺ 411.1.

15 [0858]

Example 180

2-oxo-N-(2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0859]

20 A) 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethanol

To a solution of 1-bromo-4-(trifluoromethoxy)benzene (10.1 g) in tetrahydrofuran (90 mL) was slowly added 1.6 M n-butyllithium/hexane solution (28.7 mL) at -78°C. The reaction mixture was stirred at -78°C for 30 min, a solution of 2,2,2-trifluoro-1-(piperidin-1-yl)ethanone (9.07 g) in
25 tetrahydrofuran (10 mL) was slowly added thereto. The reaction mixture was stirred at -78°C for 1.5 hr, saturated aqueous ammonium chloride solution was added thereto, and the mixture was extracted with diethyl ether. The extract was dried over
30 anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product of 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethanone. To a solution of the obtained crude product (11.9 g) in ethanol (80 mL) was slowly added sodium borohydride (3.49 g) under ice-cooling. The
35 reaction mixture was stirred at room temperature for 16 hr,

and the ethanol was evaporated under reduced pressure. To the residue was added 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed successively with 1 M hydrochloric acid and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (9.57 g).

MS (API-): [M-H]⁻259.0.

10 [0860]

B) 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethyl methanesulfonate

To a mixture of 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethanol (9.57 g) and triethylamine (7.69 mL) in tetrahydrofuran (100 mL) was slowly added methanesulfonyl chloride (3.42 mL) under ice-cooling. The reaction mixture was stirred at room temperature for 1.5 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (10.7 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.33 (3H, s), 6.62 (1H, q, J = 6.8 Hz), 7.48-7.58 (2H, m), 7.68-7.78 (2H, m).

25 [0861]

C) 1-(1-azido-2,2,2-trifluoroethyl)-4-(trifluoromethoxy)benzene

To a solution of 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethyl methanesulfonate (10.7 g) in N,N-dimethylformamide (100 mL) was added sodium azide (5.15 g) at room temperature. The reaction mixture was stirred at 100°C for 24 hr, and then at room temperature for 2 days. To the reaction mixture was added water, and the mixture was extracted with diethyl ether. The extract was washed successively with water and saturated brine, and dried over

anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel chromatography (hexane/ethyl acetate) to give the title compound (5.12 g).

5 ¹H NMR (300 MHz, CDCl₃) δ 4.94 (1H, q, J = 6.8 Hz), 7.26-7.33 (2H, m), 7.43-7.53 (2H, m).

[0862]

D) 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethanamine

To a solution of 1-(1-azido-2,2,2-trifluoroethyl)-4-
10 (trifluoromethoxy)benzene (4.92 g) in methanol (30 mL) was added 10% palladium-carbon (containing 50% water, 1 g). The reaction mixture was stirred at room temperature for 2 hr under hydrogen atmosphere, the insoluble substance was filtered off, and the filtrate was concentrated under reduced
15 pressure. To the residue was added brine, and the mixture was extracted with diethyl ether. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, solvent was evaporated under reduced pressure to give the title compound (4.31 g).

20 ¹H NMR (300 MHz, DMSO-d₆) δ 2.57 (2H, brs), 4.49-4.68 (1H, m), 7.34-7.44 (2H, m), 7.58-7.68 (2H, m).

[0863]

E) 2-oxo-N-(2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-
25 b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethanamine in the same manner as in Steps F-I of Example 63.

MS (API+): [M+H]⁺435.1.

30 [0864]

Example 181

7-methyl-2-oxo-N-(2-(pyridin-3-yloxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

35 [0865]

A) 2-((5-bromopyridin-3-yl)oxy)-1-(4-(trifluoromethoxy)phenyl)ethanone

To a solution of 5-bromopyridin-3-ol (5 g) in N,N-dimethylformamide (100 mL) was added 60% sodium hydride (1.72 g) at 0°C. The reaction mixture was stirred at 0°C for 30 min, 2-bromo-1-(4-(trifluoromethoxy)phenyl)ethanone (9.76 g) was added thereto at 0°C. The reaction mixture was stirred at room temperature for 3 hr, and added to saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and the obtained solid was washed with hexane to give the title compound (7.78 g).

MS (API+): [M+H]⁺376.0.

[0866]

B) 7-methyl-2-oxo-N-(2-(pyridin-3-yloxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-((5-bromopyridin-3-yl)oxy)-1-(4-(trifluoromethoxy)phenyl)ethanone in the same manner as in Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺488.2.

[0867]

Example 182

N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-((5-bromopyridin-3-yl)oxy)-1-(4-(trifluoromethoxy)phenyl)ethanone in the same manner as in Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺ 411.1.

[0868]

Example 183

N-(1-(3,5-difluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 [0869]

A) 5-(1-ethoxyvinyl)-1,3-difluoro-2-(trifluoromethoxy)benzene

A mixture of 5-bromo-1,3-difluoro-2-(trifluoromethoxy)benzene (5.00 g), tributyl(1-ethoxyvinyl)stannane (7.17 g),
10 bis(triphenylphosphine)dichloropalladium(II) (0.38 g) and toluene (100 mL) was stirred overnight at 100°C under nitrogen atmosphere, and 1 M hydrochloric acid (20 mL) was added thereto. The reaction mixture was stirred at room temperature for 3 hr, saturated brine was added thereto, and the mixture
15 was extracted with ethyl acetate. The extract was washed with saturated brine, dried over anhydrous magnesium sulfate, and purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the title compound (4.24 g).
¹H NMR (300 MHz, CDCl₃) δ 1.42 (3H, t, J = 7.0 Hz), 3.91 (2H, q, J = 6.8 Hz), 4.31 (1H, d, J = 3.4 Hz), 4.67 (1H, d, J = 3.0 Hz), 7.21-7.35 (2H, m).

[0870]

B) 1-(3,5-difluoro-4-(trifluoromethoxy)phenyl)ethanone

A mixture of 5-(1-ethoxyvinyl)-1,3-difluoro-2-(trifluoromethoxy)benzene (4.24 g), 6 M hydrochloric acid (15
25 mL), acetone (50 mL) and water (15 mL) was stirred at room temperature for 5 hr, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was
30 evaporated under reduced pressure to give the title compound (3.51 g).

¹H NMR (300 MHz, CDCl₃) δ 2.60 (3H, s), 7.56-7.67 (2H, m).

[0871]

C) N-(1-(3,5-difluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-
35

4(1H)-carboxamide

The title compound was obtained from 1-(3,5-difluoro-4-(trifluoromethoxy)phenyl)ethanone in the same manner as in Step A of Example 143, Step D of Example 63, Step D of Example 2, Step B of Example 15 and Step G of Example 7.

MS (API+): [M+H]⁺ 461.1.

[0872]

Example 184

N-(3-hydroxy-2,2-dimethyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a mixture of N-(3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (90 mg) and acetic acid (23.5 mg) in tetrahydrofuran (5 mL) was added 1 M tetra-n-butylammonium fluoride/tetrahydrofuran solution (0.391 mL). The reaction mixture was stirred overnight at room temperature, and 1 M tetra-n-butylammonium fluoride/tetrahydrofuran solution (0.782 mL) was added. The reaction mixture was stirred at room temperature for 2 hr, and then overnight at 50°C. To the reaction mixture was added saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (43.2 mg).

MS (API+): [M+H]⁺453.2.

[0873]

Example 185

N-(2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0874]

A) methyl (4-(trifluoromethoxy)phenyl)acetate

To a solution of (4-(trifluoromethoxy)phenyl)acetic acid

(5.00 g) in methanol (31.3 mL) was added conc. sulfuric acid (237 μ L). The reaction mixture was stirred at 70°C for 3 hr, and concentrated under reduced pressure to about 10 mL. To the residue was added water, and the mixture was extracted with
5 ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (5.23 g).

10 ^1H NMR (300 MHz, CDCl_3) δ 3.63 (2H, s), 3.71 (3H, s), 7.17 (2H, d, $J = 7.9$ Hz), 7.27–7.37 (2H, m).

[0875]

B) 2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol

To a solution of methyl (4-
15 (trifluoromethoxy)phenyl)acetate (5.23 g) in tetrahydrofuran (200 mL) was slowly added 1 M methylmagnesium bromide/tetrahydrofuran solution (55.8 mL). The reaction mixture was stirred overnight at room temperature, 1 M methylmagnesium bromide/tetrahydrofuran solution (55.8 mL) was
20 added thereto, and the mixture was stirred at room temperature for 2 hr. To the reaction mixture was added saturated aqueous ammonium chloride solution at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and
25 the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.65 g).

^1H NMR (300 MHz, CDCl_3) δ 1.23 (6H, s), 1.30 (1H, s), 2.77 (2H, s), 7.11–7.19 (2H, m), 7.21–7.29 (2H, m).

30 [0876]

C) 1-(2-fluoro-2-methylpropyl)-4-(trifluoromethoxy)benzene

To 2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol (2.55 g) was added N,N-diethylaminosulfur trifluoride (2.88 mL) at 0°C, and the reaction mixture was stirred overnight at
35 room temperature. The reaction mixture was diluted with

diethyl ether, water was added thereto at 0°C, and the mixture was extracted with diethyl ether. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane) to give the title compound (1.59 g).

¹H NMR (300 MHz, CDCl₃) δ 1.30 (3H, s), 1.37 (3H, s), 2.83-2.97 (2H, m), 7.09-7.18 (2H, m), 7.20-7.28 (2H, m).

10 [0877]

D) 1-(1-bromo-2-fluoro-2-methylpropyl)-4-(trifluoromethoxy)benzene

A mixture of 1-(2-fluoro-2-methylpropyl)-4-(trifluoromethoxy)benzene (1.59 g), N-bromosuccinimide (1.59 g) and 2,2'-azobis(2-methylpropionitrile) (113 mg) in trifluoromethylbenzene (33.7 mL) was stirred at 80°C for 2 hr. To the reaction mixture was added hexane, the insoluble substance was filtered off, and the filtrate was concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane) to give the title compound (1.66 g).

¹H NMR (300 MHz, CDCl₃) δ 1.48 (3H, d, J = 3.8 Hz), 1.55 (3H, d, J = 4.9 Hz), 4.92 (1H, d, J = 17.3 Hz), 7.18 (2H, d, J = 8.3 Hz), 7.49 (2H, d, J = 8.7 Hz).

25 [0878]

E) 1-(1-azido-2-fluoro-2-methylpropyl)-4-(trifluoromethoxy)benzene

A mixture of 1-(1-bromo-2-fluoro-2-methylpropyl)-4-(trifluoromethoxy)benzene (1.56 g) and sodium azide (1.65 g) in N,N-dimethylformamide (33.1 mL) was stirred overnight at 100°C, water was added thereto, and the mixture was extracted with diethyl ether. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (1.13 g).

¹H NMR (300 MHz, CDCl₃) δ 1.27-1.34 (3H, m), 1.35-1.42 (3H, m), 4.55 (1H, d, J = 13.9 Hz), 7.18-7.26 (2H, m), 7.39 (2H, d, J = 8.7 Hz).

[0879]

- 5 F) 2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-1-amine

A mixture of 1-(1-azido-2-fluoro-2-methylpropyl)-4-(trifluoromethoxy)benzene (1.13 g) and 10% palladium-carbon (containing 50% water, 110 mg) in methanol (7 mL) was stirred
10 at room temperature for 1 hr under hydrogen atmosphere, and the catalyst was filtered off. To the filtrate was added saturated brine, and the mixture was extracted with diethyl ether. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and
15 the solvent was evaporated under reduced pressure to give the title compound (971 mg).

MS (API+): [M+H]⁺252.1.

[0880]

- G) N-(2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-
20 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-1-amine in the same manner as in Step E of Example 168.

MS (API+): [M+H]⁺427.1.

25 [0881]

Example 186

2-oxo-N-(1-(4-(2H-1,2,3-triazol-2-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0882]

- 30 A) 1-(4-(2H-1,2,3-triazol-2-yl)phenyl)propan-1-one

A mixture of 2-di-tert-butylphosphino-3,4,5,6-tetramethyl-2',4',6'-triisopropyl-1,1'-biphenyl (181 mg) and tris(dibenzylideneacetone)dipalladium(0) (172 mg) in toluene (4 mL) was stirred at 120°C for 3 min under argon atmosphere.
35 The mixture was added to a mixture of 2H-1,2,3-triazole (389

mg), 1-(4-bromophenyl)propan-1-one (1 g) and tripotassium phosphate (1.99 g) in toluene (15 mL) at room temperature under argon atmosphere. The reaction mixture was heated at reflux for 16 hr under argon atmosphere, water was added
5 thereto, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (858 mg).

10 ¹H NMR (300 MHz, DMSO-d₆) δ 1.11 (3H, t, J = 7.2 Hz), 3.09 (2H, q, J = 7.2 Hz), 8.16 (4H, s), 8.21 (2H, s).

[0883]

B) 2-oxo-N-(1-(4-(2H-1,2,3-triazol-2-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 The title compound was obtained from 1-(4-(2H-1,2,3-triazol-2-yl)phenyl)propan-1-one in the same manner as in Step C of Example 1, Step B of Example 54 and Step B of Example 53. MS (API+): [M+H]⁺378.2.

[0884]

20 Example 187

N-(2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-1-amine in the same manner
25 as in Step E of Example 168.

MS (API+): [M+H]⁺441.2.

[0885]

Example 188

30 N-(1-(3-fluoro-4-(pyrrolidin-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0886]

A) 2-bromo-1-(4-bromo-3-fluorophenyl)ethanone

35 The title compound was obtained from 1-(4-bromo-3-

fluorophenyl)ethanone in the same manner as in Step F of Example 66.

^1H NMR (300 MHz, DMSO- d_6) δ 4.96 (2H, s), 7.77 (1H, dd, J = 8.3, 1.9 Hz), 7.90-8.00 (2H, m).

5 [0887]

B) 1-(4-bromo-3-fluorophenyl)-2-methoxyethanone

The title compound was obtained from 2-bromo-1-(4-bromo-3-fluorophenyl)ethanone in the same manner as in Step D of Example 63.

10 ^1H NMR (300 MHz, CDCl_3) δ 3.50 (3H, s), 4.62 (2H, s), 7.56-7.76 (3H, m).

[0888]

C) 2-(4-bromo-3-fluorophenyl)-2-(methoxymethyl)-1,3-dioxolane

A mixture of 1-(4-bromo-3-fluorophenyl)-2-methoxyethanone
15 (1.01 g), p-toluenesulfonic acid monohydrate (77 mg) and 1,2-ethanediol (506 mg) in toluene (20 mL) heated at reflux using Dean-Stark for 5 hr, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (NH, hexane/ethyl acetate) to give the
20 title compound (1.00 g).

^1H NMR (300 MHz, DMSO- d_6) δ 3.24 (3H, s), 3.54 (2H, s), 3.73-3.87 (2H, m), 3.95-4.09 (2H, m), 7.21 (1H, dd, J = 8.2, 2.0 Hz), 7.33 (1H, dd, J = 9.8, 2.1 Hz), 7.70 (1H, dd, J = 8.2, 7.3 Hz).

25 [0889]

D) N-(1-(3-fluoro-4-(pyrrolidin-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-(4-bromo-3-fluorophenyl)-2-(methoxymethyl)-1,3-dioxolane in the same
30 manner as in Step A of Example 147, Step C of Example 1, Step B of Example 54 and Step B of Example 53.

MS (API+): $[M+H]^+$ 428.2.

[0890]

35 Example 189

N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-(4-bromo-3-fluorophenyl)-2-(methoxymethyl)-1,3-dioxolane in the same manner as in Step A of Example 147, Step C of Example 1, Step B of Example 54 and Step B of Example 53.

MS (API+): [M+H]⁺424.2.

[0891]

10 Example 190

8-(2-hydroxypropan-2-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0892]

15 A) 2-chloro-3-nitroisonicotinic acid

The title compound was obtained from 2-chloro-4-methyl-3-nitropyridine in the same manner as in Step A of Example 167.

MS (API-): [M-H]⁻201.0.

[0893]

20 B) methyl 2-chloro-3-nitroisonicotinate

To a mixture of 2-chloro-3-nitroisonicotinic acid (2.03 g) and thionyl chloride (21.9 mL) was added N,N-dimethylformamide (two drops). The reaction mixture was stirred at 70°C for 15 min, and the solvent was evaporated under reduced pressure. To the residue was added methanol (10 mL) at 0°C, and the reaction mixture was stirred at 0°C for 10 min, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (2.15 g).

30 ¹H NMR (300 MHz, CDCl₃) δ 3.97 (3H, s), 7.86 (1H, d, J = 5.1 Hz), 8.68 (1H, d, J = 4.9 Hz).

[0894]

C) methyl 4-((2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-8-carboxylate

The title compound was obtained from methyl 2-chloro-3-nitroisonicotinate in the same manner as in Steps A-B of Example 1 and Steps C-D of Example 19.

MS (API+): [M+H]⁺469.2.

5 [0895]

D) 8-(2-hydroxypropan-2-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of methyl 4-((2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-8-carboxylate (234 mg) in tetrahydrofuran (5 mL) was added 1 M methylmagnesium bromide/tetrahydrofuran solution (10 mL) at room temperature. The reaction mixture was stirred at room temperature for 20 hr,
15 1 M hydrochloric acid was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
20 column chromatography (hexane/ethyl acetate), and recrystallized from hexane/ethyl acetate to give the title compound (47 mg).

MS (API+): [M+H]⁺ 469.2.

[0896]

25 Example 191

8-acetyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of methyl 4-((2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-8-carboxylate (234 mg) in tetrahydrofuran (5 mL) was added 1 M methylmagnesium bromide/tetrahydrofuran solution (10 mL) at room temperature. The reaction mixture was stirred at room temperature for 20 hr,
1 M hydrochloric acid was added thereto at 0°C, and the mixture
35 was extracted with ethyl acetate. The extract was washed

successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and

5 recrystallized from hexane/ethyl acetate to give the title compound (22 mg).

MS (API+): [M+H]⁺ 453.1.

[0897]

Example 192

10 8-(hydroxymethyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0898]

A) 4-((2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-8-carboxylic acid

To a mixture of methyl 4-((2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-8-carboxylate (468 mg) in
20 tetrahydrofuran (10 mL) and methanol (2.0 mL) was added 1 M aqueous sodium hydroxide solution (2.0 mL). The reaction mixture was stirred for 1 hr, and neutralized with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and
25 saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (452 mg).

MS (API+): [M+H]⁺ 455.1.

[0899]

30 B) 8-(hydroxymethyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 4-((2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamoyl)-2-oxo-1,2,3,4-tetrahydropyrido[2,3-b]pyrazine-8-carboxylic acid (445 mg) in
35

tetrahydrofuran (10 mL) were added isobutyl chloroformate (0.153 mL) and triethylamine (0.205 mL). The reaction mixture was stirred at room temperature for 1 hr, and the insoluble substance was filtered off. To the filtrate was added an
5 aqueous solution (1 mL) of sodium borohydride (74.1 mg). The reaction mixture was stirred for 1 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was
10 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and recrystallized from hexane/ethyl acetate to give the title compound (13 mg).

MS (API+): [M+H]⁺ 441.2.

15 [0900]

Example 193

N-(2-ethyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)butyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
[0901]

20 A) tert-butyl (2-oxo-1-(4-(trifluoromethoxy)phenyl)butyl)carbamate

To a solution of tert-butyl (2-(methoxy(methyl)amino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)carbamate (355 mg) in tetrahydrofuran (20 mL) was added 3 M ethylmagnesium
25 bromide/diethyl ether solution (5.0 mL) at room temperature. The reaction mixture was stirred at room temperature for 20 hr, 1 M hydrochloric acid was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over
30 anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (254 mg).

¹H NMR (300 MHz, CDCl₃) δ 1.00 (3H, t, J = 7.3 Hz), 1.40 (9H, s), 2.29-2.49 (2H, m), 5.30 (1H, d, J = 6.0 Hz), 5.91-6.01 (1H,

m), 7.17-7.24 (2H, m), 7.30-7.37 (2H, m).

[0902]

B) tert-butyl (2-ethyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)butyl)carbamate

5 To a solution of tert-butyl (2-oxo-1-(4-(trifluoromethoxy)phenyl)butyl)carbamate (320 mg) in tetrahydrofuran (10 mL) was added 3 M ethylmagnesium bromide/diethyl ether solution (5.0 mL) at 0°C. The reaction mixture was stirred at room temperature for 20 hr, 1 M
10 hydrochloric acid was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
15 column chromatography (hexane/ethyl acetate) to give the title compound (170 mg).

MS (API-): [M-H]⁻ 376.2.

[0903]

C) N-(2-ethyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)butyl)-
20 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To tert-butyl (2-ethyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)butyl)carbamate (170 mg) was added 4 M hydrogen chloride/ethyl acetate solution (5 mL) at room temperature. The reaction mixture was stirred at room
25 temperature for 20 min, and the solvent was evaporated under reduced pressure. To a mixture of the residue and 4-nitrophenyl 2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (128 mg) in N,N-dimethylformamide (5 mL) was added triethylamine (0.170 mL). The reaction mixture was stirred at
30 room temperature for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by
35 silica gel column chromatography (hexane/ethyl acetate), and

recrystallized from hexane/ethyl acetate to give the title compound (161 mg).

MS (API+): [M+H]⁺ 453.2.

[0904]

5 Example 194

N-(2-amino-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of N-(2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (285 mg) in ethanol (15 mL) was added hydrazine monohydrate (0.211 mL) at room temperature. The reaction mixture was stirred at 80°C for 3 hr, and the solvent was evaporated under reduced pressure. To the residue was added 1 M aqueous sodium hydroxide solution, and the mixture was purified by styrene/divinyl benzene synthetic adsorbent (HP-20, mobile phase: water-acetonitrile). The residue was purified by preparative HPLC (C18, mobile phase: water (10 mM, containing NH₄HCO₃)/acetonitrile), and the obtained fraction was concentrated under reduced pressure. The obtained solid was triturated with hexane/ethyl acetate, collected by filtration, and washed with hexane/ethyl acetate to give the title compound (151 mg).

MS (API+): [M+H]⁺396.1.

[0905]

25 Example 195

N-((4-bromo-2-fluorophenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0906]

30 A) 4-bromo-2-fluoro-N-methoxy-N-methylbenzamide

The title compound was obtained from 4-bromo-2-fluorobenzoic acid in the same manner as in Step A of Example 31.

MS (API+): [M+H]⁺262.0.

35 [0907]

B) (4-bromo-2-fluorophenyl) (4-(trifluoromethoxy)phenyl)methanone

To a solution of 4-bromo-2-fluoro-N-methoxy-N-methylbenzamide (10.8 g) in tetrahydrofuran (150 mL) was added
5 0.5 M (4-(trifluoromethoxy)phenyl)magnesium bromide/tetrahydrofuran solution (165 mL) at room temperature. The reaction mixture was stirred overnight at 70°C under nitrogen atmosphere, saturated aqueous ammonium chloride solution was added thereto, and the mixture was extracted with
10 ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (11.6 g).

15 MS (API+): [M+H]⁺363.0.
[0908]

C) N-((4-bromo-2-fluorophenyl) (4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 The title compound was obtained from (4-bromo-2-fluorophenyl) (4-(trifluoromethoxy)phenyl)methanone in the same manner as in Step D of Example 2, Step B of Example 132 and Step B of Example 83.

MS (API+): [M+H]⁺ 553.1.
25 [0909]

Example 196

N-((4-bromo-2-hydroxyphenyl) (4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 The title compound was obtained from 4-bromo-2-fluorobenzoic acid in the same manner as in Step A of Example 31, Step B of Example 195, Step D of Example 2, Step B of Example 132 and Step B of Example 83.

MS (API+): [M+H]⁺551.1.
35 [0910]

Example 197

N-(2-((5-bromopyridin-3-yl)oxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 The title compound was obtained in the same manner as in Step A of Example 181, Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺566.1.

[0911]

10 Example 198

N-((2-fluorophenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 A mixture of N-((4-bromo-2-fluorophenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (60 mg), 5% palladium-carbon (containing 50% water, 15 mg) and methanol (5 mL) was stirred for 2 hr under hydrogen atmosphere. The catalyst was filtered off, and the filtrate was concentrated under reduced pressure to give the title compound (35.6 mg).

20 MS (API+): [M+H]⁺475.2.

[0912]

Example 199

N-((2-hydroxyphenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 The title compound was obtained from N-((4-bromo-2-hydroxyphenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Example 198.

30 MS (API+): [M+H]⁺473.2.

[0913]

Example 200

7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-

35 (trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

A diastereomeric mixture (110 mg) of 7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide was resolved by
5 HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/2-propanol = 70/30) to give the title compound having a shortest retention time (28 mg).

MS (API+): [M+H]⁺451.2.

10 [0914]

Example 201

7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 A diastereomeric mixture (110 mg) of 7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/2-propanol = 70/30) to give the title compound having a second
20 shortest retention time (16 mg).

MS (API+): [M+H]⁺451.2.

[0915]

Example 202

25 7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A diastereomeric mixture (110 mg) of 7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide was resolved by
30 HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/2-propanol = 70/30) to give the title compound having a third shortest retention time (33 mg).

35 MS (API+): [M+H]⁺451.2.

[0916]

Example 203

7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

A diastereomeric mixture (110 mg) of 7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/2-propanol = 70/30) to give the title compound having a longest retention time (14 mg).

MS (API+): [M+H]⁺451.2.

[0917]

Example 204

7-methoxy-2-oxo-N-(1-(4-(pyrrolidin-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(4-(1H-pyrrol-1-yl)phenyl)propan-1-amine in the same manner as in Steps I-L of Example 66.

MS (API-): [M-H]⁻408.2.

[0918]

Example 205

N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0919]

A) methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate

To a solution of 2-chloro-5-iodo-3-nitropyridine (12.0 g) in ethanol (200 mL) were added methyl glycinate hydrochloride (10.6 g) and triethylamine (11.8 mL) at room temperature. The reaction mixture was heated at reflux for 20 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and

saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (11.1 g).

5 MS (API+): [M+H]⁺338.0.

[0920]

B) 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl N-(5-iodo-3-nitropyridin-2-yl)glycinate (6.00 g) in tetrahydrofuran (300 mL) was added 5%
10 platinum-carbon (500 mg). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. The residue was dissolved in ethanol (200 mL), the solution was heated at reflux for 16 hr, and the
15 solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (4.48 g).

MS (API+): [M+H]⁺276.0.

[0921]

20 C) 7-iodo-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a suspension of 7-iodo-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (3.82 g) in N,N-dimethylformamide (160 mL) was slowly added 1 M potassium
25 hexamethyldisilazide/tetrahydrofuran solution (17.4 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, 2-(chloromethoxy)ethyltrimethylsilane (3.62 mL) was slowly added thereto, and the mixture was stirred at room temperature for 20 hr. To the reaction mixture was added water, the insoluble
30 substance was filtered off using Celite, and the filtrate was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
35 column chromatography (NH, hexane/ethyl acetate) to give the

title compound (2.71 g).

MS (API+): [M+H]⁺406.0.

[0922]

D) methyl 2-((tert-butoxycarbonyl) amino)-2-(4-
5 (trifluoromethoxy)phenyl)acetate

To a mixture of 4-trifluoromethoxybenzaldehyde (19.0 g) and ammonium carbonate (25.9 g) in a mixed solvent of ethanol (114 mL) and water (45.6 mL) was slowly added an aqueous solution (71.1 mL) of potassium cyanide (8.14 g) at 50°C. The
10 reaction mixture was stirred at 60°C for 3 hr, and cooled to room temperature, and the ethanol was evaporated under reduced pressure. The pH of the residue was adjusted to 1 with conc. hydrochloric acid at 0°C, and the resulting solid was collected by filtration, and washed with water. The obtained solid was
15 added to an aqueous solution (100 mL) of potassium hydroxide (23.6 g) at room temperature, and the reaction mixture was stirred at 90°C for 3 days. The reaction mixture was cooled to room temperature, and neutralized with conc. hydrochloric acid. The resulting solid was collected by filtration, and washed
20 with water to give a crude product (13.3 g) of 2-amino-2-(4-(trifluoromethoxy)phenyl)acetic acid. To a solution of the obtained crude product (13.3 g) in tetrahydrofuran (113 mL) were added di-tert-butyl dicarbonate (19.7 mL) and 2 M aqueous sodium hydroxide solution (85 mL) at room temperature. The
25 reaction mixture was stirred overnight at room temperature, and poured into water, and the mixture was washed with diethyl ether. The pH of the aqueous layer was adjusted to 3 with 1 M hydrochloric acid at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine,
30 and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (11.3 g) of 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetic acid. To a solution of the obtained crude product (11.3 g) in N,N-dimethylformamide (84
35 mL) were added methyl iodide (2.53 mL) and potassium carbonate

(5.59 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and poured into water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.20 g).

MS (API-): [M-H]⁻348.1.

10 [0923]

E) tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

To a solution of methyl 2-((tert-butoxycarbonyl)amino)-2-(4-(trifluoromethoxy)phenyl)acetate (5.00 g) in tetrahydrofuran (71.6 mL) was slowly added 1 M methylmagnesium bromide/tetrahydrofuran solution (57.3 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr under argon atmosphere, saturated aqueous ammonium chloride solution was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.99 g).

25 MS (API-): [M-H]⁻348.2.

[0924]

F) 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride

To tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate (2.50 g) was added 4 M hydrogen chloride/ethyl acetate solution (71.6 mL). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether, and the resulting crystals were collected by filtration to give the title

compound (2.01 g).

MS (API+), found: 250.1.

[0925]

G) N-(2-hydroxy-2-methyl-1-(4-

5 (trifluoromethoxy)phenyl)propyl)-7-iodo-2-oxo-1-((2-
(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

To a solution of 7-iodo-1-((2-
(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-
10 b]pyrazin-2(1H)-one (500 mg) in tetrahydrofuran (18.9 mL) was
slowly added a solution of bis(trichloromethyl)carbonate (293
mg) in tetrahydrofuran (2.83 mL) at room temperature under
nitrogen atmosphere. The reaction mixture was stirred at 40°C
for 3 hr under nitrogen atmosphere, and the solvent was
15 evaporated under reduced pressure. To the residue was added
tetrahydrofuran, and the solvent was evaporated under reduced
pressure. The operation (addition of tetrahydrofuran and then
evaporation) was repeated three times. The residue was diluted
with tetrahydrofuran (9.42 mL), and the mixture was slowly
20 added to a mixture of 1-amino-2-methyl-1-(4-
(trifluoromethoxy)phenyl)propan-2-ol hydrochloride (423 mg)
and triethylamine (521 μ L) in tetrahydrofuran (6.28 mL) at
room temperature. The reaction mixture was stirred at 60°C
overnight, and the solvent was evaporated under reduced
25 pressure. To the residue was added water, and the mixture was
extracted with ethyl acetate. The extract was washed with
saturated brine, and dried over anhydrous sodium sulfate, and
the solvent was evaporated under reduced pressure. The residue
was purified by silica gel column chromatography (hexane/ethyl
30 acetate) to give the title compound (729 mg).

MS (API+): $[M+H]^+$ 681.1.

[0926]

H) 7-hydroxy-N-(2-hydroxy-2-methyl-1-(4-

(trifluoromethoxy)phenyl)propyl)-2-oxo-1-((2-
35 (trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

To a mixture of N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (699 mg), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi-1,3,2-dioxaborolane (537 mg) and potassium acetate (415 mg) in N,N-dimethylformamide (10.3 mL) was added (1,1-bis(diphenylphosphino)ferrocene)dichloropalladium(II) (77.0 mg). The reaction mixture was stirred overnight at 80°C under argon atmosphere, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (986 mg) of N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-7-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide. To a solution of the obtained crude product (986 mg) in tetrahydrofuran (14.5 mL) was added 2 M aqueous sodium hydroxide solution (2.90 mL) at 0°C. The reaction mixture was stirred at 0°C for 30 min, 35% aqueous hydrogen peroxide (508 µL) was added thereto. The reaction mixture was stirred at room temperature for 2 hr, ice water was added thereto, and the mixture was acidified with 1 M hydrochloric acid, and extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (340 mg).

MS (API+): [M+H]⁺571.2.

[0927]

I) N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

To a solution of 7-hydroxy-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (340 mg) in N,N-dimethylformamide (7.64 mL) were added potassium carbonate (83.0 mg) and iodomethane (56.1 μ L). The reaction mixture was stirred overnight at room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) and then HPLC (C18, mobile phase: water (10 mM, containing NH_4HCO_3)/acetonitrile), and the obtained fraction was concentrated under reduced pressure to give the title compound (91.0 mg).

MS (API+): $[\text{M}+\text{H}]^+585.3$.

[0928]

J) N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (90.9 mg) were added trifluoroacetic acid (2.24 mL) and water (251 μ L). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was dissolved in N,N-dimethylformamide (4.27 mL), and 8 M ammonia/methanol solution (838 μ L) was added thereto. The reaction mixture was stirred at room temperature for 10 min, and the solvent was evaporated under reduced pressure. The residue was diluted with ethyl acetate, and the mixture was washed successively with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated

under reduced pressure. The obtained solid was washed with hexane/ethyl acetate to give the title compound (48.8 mg).

MS (API+): [M+H]⁺455.3.

¹H NMR (300 MHz, DMSO-d₆) δ 0.98 (3H, s), 1.18 (3H, s), 3.84 (3H, s), 4.25-4.49 (2H, m), 4.67 (1H, d, J = 8.3 Hz), 4.77 (1H, s), 6.96 (1H, d, J = 2.6 Hz), 7.27 (2H, d, J = 7.9 Hz), 7.42 (2H, d, J = 8.7 Hz), 7.76 (1H, d, J = 2.6 Hz), 10.19 (1H, d, J = 8.3 Hz), 10.73 (1H, brs).

[0929]

10 Example 206

7-methyl-2-oxo-N-(tetrahydrofuran-3-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from tetrahydrofuran-3-carboxylic acid in the same manner as in Step A of Example 170, Step D of Example 2, Step B of Example 132 and Step G of Example 7.

MS (API+): [M+H]⁺451.2.

[0930]

20 Example 207

N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (43.8 mg) was resolved by SFC (column: CHIRALPAK ADH, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=800/200), and crystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (5.4 mg).

MS (API+): [M+H]⁺455.2.

¹H NMR (300 MHz, DMSO-d₆) δ 0.98 (3H, s), 1.18 (3H, s), 3.84 (3H, s), 4.20-4.50 (2H, m), 4.67 (1H, d, J = 8.3 Hz), 4.77 (1H, s), 6.96 (1H, d, J = 3.0 Hz), 7.27 (2H, d, J = 7.9 Hz), 7.42

(2H, d, J = 8.7 Hz), 7.76 (1H, d, J = 2.6 Hz), 10.19 (1H, d, J = 8.3 Hz), 10.73 (1H, brs).

[0931]

Example 208

5 N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (43.8 mg) was
10 resolved by SFC (column: CHIRALPAK ADH, 20 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: carbon dioxide/methanol=800/200), and crystallized from hexane/ethyl acetate to give the title compound having a
15 longer retention time (5.2 mg).

MS (API+): [M+H]⁺455.1.

[0932]

Example 209

7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
20 carboxamide

[0933]

A) 1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethanone

The title compound was obtained from 2-bromo-1-(3-fluoro-4-(trifluoromethyl)phenyl)ethanone in the same manner as in
25 Step D of Example 63.

MS (API-): [M-H]⁻235.0.

[0934]

B) 1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethanamine
30 hydrochloride

The title compound was obtained from 1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethanone in the same manner as in Step E of Example 63.

MS (API+), found: 238.1.

35 [0935]

C) N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethanamine hydrochloride in the same manner as in Step F of Example 63.

MS (API+): [M+H]⁺669.1.

[0936]

D) 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Step H of Example 72-II.

MS (API+): [M+H]⁺583.2.

[0937]

E) 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Step I of Example 72-II.

MS (API+): [M+H]⁺453.1.

[0938]

Example 210

7-methyl-2-oxo-N-(3,3,3-trifluoro-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0939]

A) (4-(trifluoromethoxy)phenyl)acetaldehyde

To a mixture of 2-(4-(trifluoromethoxy)phenyl)ethanol

(2.00 g) and sodium hydrogen carbonate (8.15 g) in acetonitrile (150 mL) was added Dess-Martin periodinane (8.23 g) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, saturated brine was added thereto, and the mixture was
5 extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
10 column chromatography (ethyl acetate) to give the title compound (1.72 g).

¹H NMR (300 MHz, CDCl₃) δ 3.72 (2H, d, J = 1.9 Hz), 6.84-7.39 (4H, m), 9.77 (1H, t, J = 2.1 Hz).

[0940]

15 B) 1,1,1-trifluoro-3-(4-(trifluoromethoxy)phenyl)propan-2-ol

To a mixture of (4-(trifluoromethoxy)phenyl)acetaldehyde (800 mg) and trimethyl(trifluoromethyl)silane (2.79 g) in tetrahydrofuran (10 mL) was added 1 M tetra-n-butylammonium fluoride/tetrahydrofuran solution (7.84 mL) at 0°C. The
20 reaction mixture was stirred at 0°C for 2 hr, 1 M hydrochloric acid was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was
25 purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (761 mg).

¹H NMR (300 MHz, CDCl₃) δ 2.13 (1H, d, J = 5.7 Hz), 2.80-2.94 (1H, m), 2.97-3.11 (1H, m), 4.01-4.31 (1H, m), 7.04-7.24 (2H, m), 7.27-7.36 (2H, m).

30 [0941]

C) tert-butyl(dimethyl)((1,1,1-trifluoro-3-(4-(trifluoromethoxy)phenyl)propan-2-yl)oxy)silane

A mixture of 1,1,1-trifluoro-3-(4-(trifluoromethoxy)phenyl)propan-2-ol (750 mg), tert-
35 butyldimethylsilyl trifluoromethanesulfonate (1.09 g), 2,6-

dimethylpyridine (586 mg) and tetrahydrofuran (5 mL) was stirred at 0°C for 2 hr. The reaction mixture was added to saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed successively with 1 M hydrochloric acid and saturated aqueous sodium hydrogen carbonate solution, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (434 mg).

¹H NMR (300 MHz, CDCl₃) δ -0.48 (3H, s), -0.05 (3H, s), 0.72-0.79 (9H, m), 2.80 (1H, dd, J = 13.8, 10.0 Hz), 3.03 (1H, dd, J = 13.8, 2.8 Hz), 3.93-4.25 (1H, m), 7.07-7.20 (2H m), 7.21-7.25 (2H, m).

[0942]

D) ((3-bromo-1,1,1-trifluoro-3-(4-(trifluoromethoxy)phenyl)propan-2-yl)oxy)(tert-butyl)dimethylsilane

A mixture of tert-butyl(dimethyl)((1,1,1-trifluoro-3-(4-(trifluoromethoxy)phenyl)propan-2-yl)oxy)silane (165 mg), 2,2'-(diazene-1,2-diyl)bis(2-methylpropanenitrile) (6.98 mg), 1-bromopyrrolidine-2,5-dione (98 mg) and benzotrifluoride (5 mL) was stirred overnight at 130°C, were added 1-bromopyrrolidine-2,5-dione (227 mg) and 2,2'-(diazene-1,2-diyl)bis(2-methylpropanenitrile) (14.0 mg), and the reaction mixture was stirred at 130°C for 6 hr. Separately, a mixture of tert-butyl(dimethyl)((1,1,1-trifluoro-3-(4-(trifluoromethoxy)phenyl)propan-2-yl)oxy)silane (200 mg), 2,2'-(diazene-1,2-diyl)bis(2-methylpropanenitrile) (42.3 mg), 1-bromopyrrolidine-2,5-dione (458 mg) and benzotrifluoride (10 mL) was stirred at 120°C for 2 days. These mixtures were combined, hexane was added thereto, and the insoluble substance was filtered off. The filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give a

pale yellow oil (385 mg). To a solution of this oil (380 mg) in N,N-dimethylformamide (15 mL) was added sodium azide (264 mg) at room temperature. The reaction mixture was stirred overnight at 100°C, saturated brine was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate to give an orange oil (280 mg). To a solution of this oil (280 mg) in methanol (5 mL) was added 10% palladium-carbon (containing 50% water, 20 mg). The reaction mixture was stirred overnight under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. To the residue were added triethylamine (92 mg), 4-nitrophenyl-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (100 mg) and N,N-dimethylformamide (5 mL). The reaction mixture was stirred overnight at room temperature, saturated aqueous sodium hydrogen carbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by HPLC (C18, mobile phase: water/acetonitrile (containing 0.1% TFA)), to the obtained fraction was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous magnesium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (23.2 mg). MS (API+): [M+H]⁺479.1.

[0943]

Example 211

N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0944]

A) N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 The title compound was obtained from 1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethanamine in the same manner as in Steps I-K of Example 66.

MS (API+): [M+H]⁺570.3.

[0945]

10 B) N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 To a solution of N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (112 mg) in dichloromethane (3.72 mL) was added boron trifluoride diethyl ether complex (250 µL) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, and the solvent was evaporated under reduced pressure. To the
15 residue was added water, and the mixture was extracted with diethyl ether. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was dissolved in methanol (5.00 mL), N-ethyl-N-isopropylpropan-2-amine (344 µL) was added thereto at room temperature. The reaction
20 mixture was stirred at 50°C for 1 hr, the solvent was evaporated under reduced pressure, and the obtained solid was washed with hexane/ethyl acetate to give the title compound (58.8 mg).

30 MS (API+): [M+H]⁺440.2.

[0946]

Example 212

6-(4-methoxyphenyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
35

[0947]

A) methyl 2-((6-chloro-3-nitropyridin-2-yl) amino)acetate

The title compound was obtained from 2,6-dichloro-3-nitropyridine in the same manner as in Step A of Example 1.

5 ¹H NMR (300 MHz, DMSO-d₆) δ 3.67 (3H, s), 4.27 (2H, d, J = 6.0 Hz), 6.87 (1H, d, J = 8.7 Hz), 8.48 (1H, d, J = 8.7 Hz), 8.96 (1H, t, J = 5.7 Hz).

[0948]

B) 6-chloro-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

10 The title compound was obtained from methyl 2-((6-chloro-3-nitropyridin-2-yl) amino)acetate in the same manner as in Step B of Example 66.

¹H NMR (300 MHz, DMSO-d₆) δ 3.94 (2H, s), 6.55 (1H, d, J = 7.9 Hz), 6.91 (1H, d, J = 7.9 Hz), 7.19 (1H, s), 10.47 (1H, s).

15 [0949]

C) 6-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

The title compound was obtained from 6-chloro-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner as in
20 Step C of Example 63.

MS (API+): [M+H]⁺314.3.

[0950]

D) 6-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
25

The title compound was obtained from 6-chloro-1-((2-(trimethylsilyl)ethoxy)methyl)-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one in the same manner as in Step F of Example 63.

30 MS (API+): [M+H]⁺575.2.

[0951]

E) 6-(4-methoxyphenyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

35 To a solution of 6-chloro-N-(2-methoxy-1-(4-

(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (69.0 mg) in 1,2-dimethoxyethane (1.5 mL) were added 4-methoxyphenylboronic acid (36.5 mg), 1 M potassium carbonate aqueous solution (0.24 mL) and bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (8.5 mg), and the mixture was heated at 120°C for 1 hr using microwave generator (Biotage, Initiator Sixty). To a reaction solution were added water (1 mL) and ethyl acetate (2 mL), and the mixture was stirred, and filtered through phase separating filter, and the organic layer was concentrated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device. To the residue was added trifluoroacetic acid (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was evaporated using air spraying device. To the residue was added 8 M ammonia/methanol solution (1 mL), the mixture was stirred at room temperature for 1 hr, and the solvent was evaporated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device to give the title compound (36.7 mg).

MS (API+): [M+H]⁺517.1.

[0952]

Example 213

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(pyridin-3-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from pyridin-3-ylboronic acid in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺488.1.

[0953]

Example 214

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-phenyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from phenylboronic acid
5 in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺487.1.

[0954]

Example 215

6-(2-methoxyphenyl)-N-(2-methoxy-1-(4-
10 (trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-methoxyphenylboronic acid in the same manner as in Step E of Example 212.

15 MS (API+): [M+H]⁺517.1.

[0955]

Example 216

6-(3-methoxyphenyl)-N-(2-methoxy-1-(4-
(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
20 b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-methoxyphenylboronic acid in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺517.2.

25 [0956]

Example 217

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(pyridin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 The title compound was obtained from pyridin-4-ylboronic acid in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺488.1.

[0957]

Example 218

35 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(2-

thienyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 2-thienylboronic acid in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺493.1.

5 [0958]

Example 219

6-(3-furyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-furylboronic acid
10 in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺477.1.

[0959]

Example 220

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(3-
15 thienyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3-thienylboronic acid in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺493.1.

[0960]

20 Example 221

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1H-pyrazol-5-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole in the same
25 manner as in Step E of Example 212.

MS (API+): [M+H]⁺491.1.

[0961]

Example 222

30 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1H-pyrazol-3-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-methyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole in the same
35 manner as in Step E of Example 212.

MS (API+): [M+H]⁺491.1.

[0962]

Example 223

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-
5 1H-pyrazol-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
carboxamide

The title compound was obtained from 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole in the same manner as in Step E of Example 212.

10 MS (API+): [M+H]⁺491.1.

[0963]

Example 224

6-(3,5-dimethyl-1,2-oxazol-4-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
15 b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3,5-dimethyl-1,2-oxazol-4-ylboronic acid in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺506.1.

20 [0964]

Example 225

N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
carboxamide

25 The title compound was obtained from N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-iodo-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Steps J-L of Example 66.

30 MS (API+): [M+H]⁺443.1.

[0965]

Example 226

optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-
35 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (347 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300) to give the title compound having a shorter retention time (107 mg).

MS (API+): [M+H]⁺453.1.

[0966]

10 Example 227

optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (347 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300) to give the title compound having a longer retention time (83 mg).

MS (API+): [M+H]⁺453.1.

[0967]

Example 228

N-(2-methoxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0968]

A) methyl 2-(4-(trifluoromethoxy)phenyl)acetate

To a solution of 2-(4-(trifluoromethoxy)phenyl)acetic acid (5.00 g) in methanol (32 mL) was added conc. sulfuric acid (0.237 mL) at room temperature. The reaction mixture was stirred at 70°C for 4 hr, and concentrated to about 10 mL under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous

magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (4.00 g).

MS (API-): [M-H]⁻ 233.0.

[0969]

5 B) 2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol

To a solution of methyl 2-(4-(trifluoromethoxy)phenyl)acetate (4.00 g) in tetrahydrofuran (57 mL) was added 1 M methylmagnesium bromide/tetrahydrofuran solution (42.7 mL) at 0°C. The reaction mixture was stirred at
10 room temperature for 4 hr, saturated aqueous ammonium chloride solution was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The
15 residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.26 g).
¹H NMR (300 MHz, DMSO-d₆) δ 1.06 (6H, s), 2.67 (2H, s), 4.36 (1H, s), 7.31 (4H, s).

[0970]

20 C) 1-(2-methoxy-2-methylpropyl)-4-(trifluoromethoxy)benzene

To a solution of 2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol (6.72 g) in N,N-dimethylformamide (140 mL) was added 60% sodium hydride (5.74 g) at room temperature. The reaction mixture was stirred at
25 room temperature for 30 min, methyl iodide (8.97 mL) was slowly added thereto at room temperature, and the mixture was stirred at room temperature for 18 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous magnesium
30 sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (6.01 g).
¹H NMR (300 MHz, DMSO-d₆) δ 1.05 (6H, s), 2.75 (2H, s), 3.16 (3H, s), 7.17-7.37 (4H, m).

35 [0971]

D) 1-(1-bromo-2-methoxy-2-methylpropyl)-4-(trifluoromethoxy)benzene

The title compound was obtained from 1-(2-methoxy-2-methylpropyl)-4-(trifluoromethoxy)benzene in the same manner
5 as in Step D of Example 142.

¹H NMR (300 MHz, DMSO-d₆) δ 1.13 (3H, s), 1.25 (3H, s), 3.20 (3H, s), 5.33-5.42 (1H, m), 7.18-7.38 (2H, m), 7.62-7.70 (2H, m).

[0972]

10 E) 1-(1-azido-2-methoxy-2-methylpropyl)-4-(trifluoromethoxy)benzene

To a solution of 1-(1-bromo-2-methoxy-2-methylpropyl)-4-(trifluoromethoxy)benzene (2.10 g) in N,N-dimethylformamide (43 mL) were added sodium azide (2.09 g) and 18-crown-6 (6.8
15 g). The reaction mixture was stirred at 100°C for 18 hr, water was added thereto, and the mixture was extracted with diethyl ether. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound
20 (1.66 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.00-1.11 (6H, m), 3.20 (3H, s), 4.83-4.90 (1H, m), 7.33-7.40 (2H, m), 7.42-7.56 (2H, m).

[0973]

F) N-(2-methoxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
25

The title compound was obtained from 1-(1-azido-2-methoxy-2-methylpropyl)-4-(trifluoromethoxy)benzene in the same manner as in Steps E-F of Example 2 and Step L of Example
30 72-I.

MS (API+): [M+H]⁺ 453.2.

[0974]

Example 229

optically active N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
35

b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide (55.2 mg) was resolved by HPLC

5 (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300) to give the title compound having a shorter retention time (13.7 mg).

MS (API+): [M+H]⁺440.2.

10 [0975]

Example 230

optically active N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 Racemic N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (55.2 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol =
20 700/300) to give the title compound having a longer retention time (15.4 mg).

MS (API+): [M+H]⁺440.2.

[0976]

Example 231

25 optically active 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
30 b]pyrazine-4(1H)-carboxamide (426 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300) to give the title compound having a shorter retention time (177 mg).

35 MS (API+): [M+H]⁺435.1.

¹H NMR (300 MHz, CDCl₃) δ 0.66-0.75 (2H, m), 0.98-1.08 (2H, m), 1.83-1.95 (1H, m), 3.40 (3H, s), 3.69 (2H, d, J = 5.3 Hz), 4.64 (2H, s), 5.16-5.26 (1H, m), 6.83 (1H, d, J = 2.3 Hz), 7.45-7.62 (4H, m), 7.84 (1H, d, J = 1.9 Hz), 8.84-8.97 (1H, m),
5 10.55 (1H, d, J = 7.2 Hz).

[0977]

Example 232

optically active N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-

10 b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide (471 mg) was resolved by HPLC

(column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel
15 Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300), and recrystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (208 mg).

MS (API+): [M+H]⁺443.1.

[0978]

20 Example 233

optically active 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
25 b]pyrazine-4(1H)-carboxamide (426 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300), and recrystallized from hexane/ethyl acetate to give
30 the title compound having a longer retention time (179 mg).

MS (API+): [M+H]⁺435.1.

¹H NMR (300 MHz, CDCl₃) δ 0.66-0.75 (2H, m), 0.98-1.08 (2H, m), 1.81-1.94 (1H, m), 3.40 (3H, s), 3.69 (2H, d, J = 5.3 Hz), 4.64 (2H, s), 5.14-5.26 (1H, m), 6.83 (1H, d, J = 1.9 Hz),
35 7.45-7.64 (4H, m), 7.84 (1H, d, J = 1.9 Hz), 9.01 (1H, s),

10.55 (1H, d, J = 7.2 Hz).

[0979]

Example 234

optically active N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (471 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300) to give the title compound having a longer retention time (199 mg).

MS (API+): [M+H]⁺443.1.

[0980]

Example 235

6-(3,6-dihydro-2H-pyran-4-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 3,6-dihydro-2H-pyran-4-boronic acid pinacol ester in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺493.1.

[0981]

Example 236

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-methyl-1,2,3,6-tetrahydropyridine-4-boronic acid pinacol ester in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺506.1.

[0982]

Example 237

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-

(1,2,3,6-tetrahydropyridin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate in the same manner as in Step E of Example 212.

MS (API+): [M+H]⁺492.1.

[0983]

Example 238

10 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 6-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-(trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (46.0 mg) in 1,2-dimethoxyethane (1.0 mL) was added water (0.16 mL), trimethylboroxine (50.0 mg), cesium carbonate (52.0 mg) and bis(di-tert-butyl(4-dimethylaminophenyl)phosphine)dichloropalladium(II) (5.6 mg), and the mixture was heated at 110°C for 1 hr using microwave generator (Biotage, Initiator Sixty). To the reaction mixture were added water (1 mL) and ethyl acetate (2 mL), and the mixture was stirred, and filtered through phase separating filter, and the organic layer was concentrated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device. To the residue was added trifluoroacetic acid (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was evaporated using air spraying device. To the residue was added 8 M ammonia/methanol solution (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was evaporated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying

15
20
25
30
35

device to give the title compound (16.5 mg).

MS (API+): [M+H]⁺425.0.

[0984]

Example 239

5 N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(pyridin-2-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 6-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-
10 (trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (46.0 mg) in N,N-dimethylformamide (0.5 mL) were added (2-pyridine)cyclic triolborate lithium (51.1 mg), cesium carbonate (78.1 mg), bis(di-tert-butyl(4-
15 dimethylaminophenyl)phosphine)dichloropalladium(II) (5.6 mg) and copper(I) chloride (4.0 mg), and the mixture was heated at 100°C for 1 hr using microwave generator (Biotage, Initiator Sixty). To the reaction mixture were added water (1 mL) and ethyl acetate (2 mL), and the mixture was stirred, and
20 filtered through phase separating filter, and the organic layer was concentrated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device. To the
25 residue was added trifluoroacetic acid (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was evaporated using air spraying device. To the residue was added 8 M ammonia/methanol solution (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was
30 evaporated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device to give the title compound (6.3 mg).

35 MS (API+): [M+H]⁺488.1.

[0985]

Example 240

6-(2-furyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-
2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 The title compound was obtained from sodium (2-furan)cyclic-triolborate in the same manner as in Example 239.
MS (API+): [M+H]⁺477.1.

[0986]

Example 241

10 6-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 To a solution of 6-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-1-((2-
15 (trimethylsilyl)ethoxy)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (46.0 mg) in toluene (1.0 mL) were added water (0.2 mL), cyclopropylboronic acid (20.6 mg), cesium carbonate (78.1 mg), palladium(II) acetate (3.6 mg) and tricyclohexylphosphine (6.7 mg), and the mixture was heated at
20 100°C for 30 min using microwave generator (Biotage, Initiator Sixty). To the reaction mixture were added water (1 mL) and ethyl acetate (2 mL), and the mixture was stirred, and filtered through phase separating filter, and the organic layer was concentrated using air spraying device. The residue
25 was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device. To the residue was added trifluoroacetic acid (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was
30 evaporated using air spraying device. To the residue was added 8 M ammonia/methanol solution (1 mL), and the mixture was stirred at room temperature for 1 hr, and the solvent was evaporated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10
35 mM aqueous ammonium bicarbonate solution), and the solvent was

evaporated using air spraying device to give the title compound (18.3 mg).

MS (API+): [M+H]⁺451.1.

[0987]

5 Example 242

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(tetrahydro-2H-pyran-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 6-(3,6-dihydro-2H-pyran-4-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (10 mg) in methanol (2 mL) were added ammonium formate (50 mg) and 5% palladium-carbon-ethylene diamine complex (50 mg), and the mixture was stirred at 60°C for 2 hr. The reaction mixture was filtered, and concentrated using air spraying device. The residue was purified by HPLC (column: YMC Triart C18, mobile phase: acetonitrile/10 mM aqueous ammonium bicarbonate solution), and the solvent was evaporated using air spraying device to give the title compound (5.5 mg).

20 MS (API+): [M+H]⁺495.1.

[0988]

Example 243

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methylpiperidine-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Example 242.

30 MS (API+): [M+H]⁺508.1.

[0989]

Example 244

N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(piperidin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-

carboxamide

The title compound was obtained from N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1,2,3,6-tetrahydropyridin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide in the same manner as in Example 242.

MS (API+): [M+H]⁺494.1.

[0990]

Example 245

6-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[0991]

A) 2-chloro-3-methyl-5-nitropyridine 1-oxide

To a mixture of 2-chloro-3-methyl-5-nitropyridine (5.00 g) and hydrogen peroxide urea (5.72 g) in acetonitrile (37.6 mL) was slowly added trifluoroacetic anhydride (8.06 mL) at 0°C. The reaction mixture was stirred at 0°C for 2 hr, 9% (w/w) aqueous sodium dithionite solution (60 mL) was added thereto, and the mixture was stirred at room temperature for 20 min. The reaction mixture was extracted with ethyl acetate. The extract was washed successively with saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.90 g).

¹H NMR (300 MHz, DMSO-d₆) δ 2.33 (3H, s), 7.98 (1H, d, J = 0.8 Hz), 8.70 (1H, dd, J = 1.7, 0.8 Hz).

[0992]

B) 2,6-dichloro-3-methyl-5-nitropyridine

A mixture of 2-chloro-3-methyl-5-nitropyridine 1-oxide (8.17 g) and phosphorus oxychloride (101 mL) was stirred at 90°C for 30 min, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title

compound (7.63 g).

¹H NMR (300 MHz, CDCl₃) δ 2.48 (3H, s), 8.14 (1H, d, J = 0.8 Hz).

[0993]

5 C) methyl 2-(6-chloro-5-methyl-3-nitropyridin-2-yl)acetate

The title compound was obtained from 2,6-dichloro-3-methyl-5-nitropyridine in the same manner as in Step A of Example 1.

MS (API+): [M+H]⁺260.0.

10 [0994]

D) methyl 2-(6-methoxy-5-methyl-3-nitropyridin-2-yl)acetate

A mixture of methyl 2-(6-chloro-5-methyl-3-nitropyridin-2-yl)acetate (2.63 g) and sodium methoxide (2.93 g) in methanol (50.6 mL) was stirred overnight at room temperature, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.12 g).

MS (API+): [M+H]⁺256.1.

[0995]

20 E) 6-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from methyl 2-(6-methoxy-5-methyl-3-nitropyridin-2-yl)acetate in the same manner as in Step B of Example 1 and Steps C-D of Example 19.

MS (API+): [M+H]⁺455.1.

[0996]

Example 246

7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 [0997]

A) 2-nitroacetamide ammonia salt

A mixture of 2-nitroethyl acetate (17.5 g) and 28% aqueous ammonia solution (88 mL) was stirred at room

temperature for 4 days, and the solvent was evaporated under reduced pressure to give the title compound (15.0 g).

^1H NMR (300 MHz, D_2O) δ 6.43 (2H, s).

[0998]

- 5 B) methyl 2-((5-methoxy-6-methyl-3-nitropyridin-2-yl)amino)acetate

To a solution of 28% sodium methoxide/methanol solution (16.7 g) in methanol (87 mL) was slowly added a mixture of 1-methoxypropan-2-one (7.63 g) and ethyl formate (7.68 mL) at -
10 $^{\circ}\text{C}$. The reaction mixture was stirred at -10°C for 1 hr under argon atmosphere, and then overnight at room temperature, and the solvent was evaporated under reduced pressure. To a solution of the residue (12.0 g) and 2-nitroacetamide ammonia salt (12.9 g) in water (54.1 mL) were added 3.4 M aqueous
15 piperidine acetate solution (26.7 mL) and acetic acid (5.85 mL) at room temperature. The reaction mixture was stirred overnight at 50°C , water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate,
20 and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give 5-methoxy-6-methyl-3-nitropyridin-2-ol (1.06 g) containing an impurity. To a solution of the 5-methoxy-6-methyl-3-nitropyridin-2-ol
25 obtained above (153 mg) in pyridine (4.15 mL) was added trifluoromethanesulfonic anhydride (211 μL) at 0°C . The reaction mixture was stirred at 0°C for 2 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with 1 M hydrochloric acid,
30 saturated aqueous sodium hydrogen carbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To a solution of the residue (262 mg) in N,N-dimethylacetamide (4.15 mL) were added glycinemethyl ester hydrochloride (125
35 mg) and triethylamine (289 μL) at room temperature. The

reaction mixture was stirred overnight at 70°C, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated
5 under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (70 mg).

MS (API+): [M+H]⁺256.1.

[0999]

10 C) 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from methyl 2-((5-methoxy-6-methyl-3-nitropyridin-2-yl) amino)acetate in the
15 same manner as in Step B of Example 1 and Steps C-D of Example 19.

MS (API+): [M+H]⁺455.1.

[1000]

Example 247

20 N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 4-nitrophenyl 7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
25 carboxylate and 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride in the same manner as in Step C of Example 19.

MS (API+): [M+H]⁺469.1.

[1001]

30 Example 248

N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 4-nitrophenyl 6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
35

carboxylate and 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride in the same manner as in Step C of Example 19.

MS (API+): [M+H]⁺469.1.

5 [1002]

Example 249

N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 1-(3-methoxy-1-(4-(trifluoromethoxy)phenyl))propylamine hydrochloride in the same manner as in Step C of Example 19.

MS (API+): [M+H]⁺439.1.

[1003]

Example 250

15 N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

[1004]

A) methyl 2-((5-methyl-3-nitropyridin-2-yl) amino)acetate

20 To a mixture of methyl glycinate hydrochloride (8.32 g) and 2-chloro-5-methyl-3-nitropyridine (7.62 g) in N,N-dimethylformamide (100 mL) was added triethylamine (18.5 mL). The reaction mixture was stirred at 80°C for 16 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with water, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (6.99 g).

¹H NMR (300 MHz, DMSO-d₆) δ 2.24 (3H, s), 3.64 (3H, s), 4.28 (2H, d, J = 5.7 Hz), 8.27-8.37 (2H, m), 8.60 (1H, t, J = 5.7 Hz).

[1005]

35 B) 7-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl 2-((5-methyl-3-nitropyridin-2-yl) amino)acetate (6.99 g) in ethanol (100 mL) was added 10% palladium-carbon (containing 50% water, 2.00 g). The reaction mixture was stirred overnight at room temperature under
5 hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. To the residue was added ethanol (100 mL), the reaction mixture was heated at reflux for 5 hr, and the solvent was evaporated to about 1/3 of the volume under reduced pressure. To the residue
10 was added diisopropyl ether, and the precipitated solid was collected by filtration. The solid was washed with diisopropyl ether, and dried under reduced pressure to give the title compound (4.56 g).

¹H NMR (300 MHz, DMSO-d₆) δ 2.09 (3H, s), 3.85 (2H, d, J = 1.5
15 Hz), 6.45 (1H, s), 6.72-6.80 (1H, m), 7.39-7.48 (1H, m), 10.31 (1H, s).

[1006]

C) 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

20 To a mixture of 7-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (1.00 g) in a mixed solvent of N,N-dimethylacetamide (40 mL) and pyridine (15 mL) was added 4-nitrophenyl chloroformate (1.48 g) at room temperature. The reaction mixture was stirred at room temperature for 16 hr, and water
25 was added thereto. The precipitated solid was collected by filtration, washed with water, and dried under reduced pressure to give the title compound (1.63 g).

MS (API+): [M+H]⁺ 329.1.

[1007]

30 D) methyl 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetate

To a mixture of 4-trifluoromethoxybenzaldehyde (19.0 g) and ammonium carbonate (25.9 g) in a mixed solvent of ethanol (114 mL) and water (45.6 mL) was slowly added an aqueous
35 solution (71.1 mL) of potassium cyanide (8.14 g) at 50°C. The

reaction mixture was stirred at 60°C for 3 hr, and cooled to room temperature, and the ethanol was evaporated under reduced pressure. The pH of the residue was adjusted to 1 with conc. hydrochloric acid at 0°C, and the resulting solid was collected
5 by filtration, and washed with water. To an aqueous solution (100 mL) of potassium hydroxide (23.6 g) was added the solid obtained above at room temperature, and the reaction mixture was stirred at 90°C for 2 days. The reaction mixture was cooled to room temperature, and neutralized with conc.
10 hydrochloric acid. The resulting solid was collected by filtration, and washed with water to give a crude product (13.3 g) of 2-amino-2-(4-(trifluoromethoxy)phenyl)acetic acid. To a solution of the obtained crude product (13.3 g) in tetrahydrofuran (113 mL) were added di-tert-butyl dicarbonate
15 (19.7 mL) and 2 M aqueous sodium hydroxide solution (85 mL) at room temperature. The reaction mixture was stirred overnight at room temperature, and poured into water, and the mixture was washed with diethyl ether. The pH of the aqueous layer was adjusted to 3 with 1 M hydrochloric acid at 0°C, and the
20 mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a crude product (11.3 g) of 2-((tert-butoxycarbonyl) amino)-2-(4-(trifluoromethoxy)phenyl)acetic
25 acid. To a solution of the obtained crude product (11.3 g) in N,N-dimethylformamide (84 mL) were added methyl iodide (2.53 mL) and potassium carbonate (5.59 g) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, and poured into water, and the mixture was extracted with ethyl
30 acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (8.20 g).

35 MS (API-): [M-H]⁻348.1.

[1008]

E) tert-butyl(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate

To a solution of methyl 2-((tert-butoxycarbonyl)amino)-2-(4-(trifluoromethoxy)phenyl)acetate (5.00 g) in tetrahydrofuran (71.6 mL) was slowly added 1 M methylmagnesium bromide/tetrahydrofuran solution (57.3 mL) at 0°C under argon atmosphere. The reaction mixture was stirred at 0°C for 1 hr under argon atmosphere, saturated aqueous ammonium chloride solution was added thereto at 0°C, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (3.99 g). MS (API-): [M-H]⁻348.2.

[1009]

F) 1-amino-2-methyl-1-(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride

To tert-butyl(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)carbamate (2.50 g) was added 4 M hydrogen chloride/ethyl acetate solution (71.6 mL). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether, and the resulting crystals were collected by filtration to give the title compound (2.01 g). MS (API+), found: 250.1.

[1010]

G) N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a suspension of 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (263 mg) in N,N-dimethylformamide (4.00 mL) were added 1-amino-2-methyl-1-

(4-(trifluoromethoxy)phenyl)propan-2-ol hydrochloride (251 mg) and triethylamine (245 μ L) at room temperature. The reaction mixture was stirred overnight at room temperature, and poured into water, and the mixture was extracted with ethyl acetate. 5 The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and the obtained solid was washed with diisopropyl ether to give the title 10 compound (298 mg).

MS (API+): [M+H]⁺439.1.

¹H NMR (300 MHz, DMSO-d₆) δ 0.94-1.03 (3H, m), 1.18 (3H, s), 2.27 (3H, s), 4.27-4.36 (1H, m), 4.37-4.47 (1H, m), 4.68 (1H, d, J = 8.3 Hz), 4.78 (1H, s), 7.13 (1H, d, J = 1.5 Hz), 7.27 15 (2H, d, J = 7.9 Hz), 7.42 (2H, d, J = 8.7 Hz), 7.87 (1H, d, J = 1.1 Hz), 10.54 (1H, d, J = 8.3 Hz), 10.77 (1H, brs).

[1011]

Example 251

7-methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide 20

Racemic 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (80 mg) was 25 resolved by HPLC (column: CHIRALPAK IA, 50 mmID $\times$ 500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30, subsequently changed to 50/50) to give the title compound having a shorter retention time (23 mg).

30 MS (API+): [M+H]⁺455.1.

[1012]

Example 252

7-methoxy-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide 35

Racemic 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (80 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30, subsequently changed to 50/50) to give the title compound having a longer retention time (18 mg).

MS (API+): [M+H]⁺455.1.

10 [1013]

Example 253

N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

15 Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (33 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30, subsequently changed to 50/50) to give the title compound having a shorter retention time (15 mg).

MS (API+): [M+H]⁺469.1.

[1014]

25 Example 254

N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (33 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30, subsequently changed to 50/50) to give the title compound having a longer retention time (15

mg).

MS (API+): [M+H]⁺469.1.

[1015]

Example 255

5 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-
2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

The title compound was obtained from 5-(trifluoromethyl)pyridine-2-carboxylic acid in the same manner as in Step A of Example 84, Steps C-F of Example 2 and Example
10 Step L of Example 72-I.

MS (API+): [M+H]⁺394.1.

[1016]

Example 256

7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
15

The title compound was obtained from 1-amino-1-(4-(trifluoromethyl)phenyl)-2-methylpropan-2-ol hydrochloride in the same manner as in Steps K-M of Example 75.

20 MS (API+): [M+H]⁺ 449.2.

[1017]

Example 257

optically active N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide
25

Racemic N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (300 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile
30 phase: hexane/ethanol = 70/30, subsequently changed to 50/50) to give the title compound having a shorter retention time (140 mg).

MS (API+): [M+H]⁺439.1.

35 [1018]

Example 258

optically active N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

5 Racemic N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (300 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile
10 phase: hexane/ethanol = 70/30, subsequently changed to 50/50) to give the title compound having a longer retention time (134 mg).

MS (API+): [M+H]⁺439.1.
[1019]

15 Example 259

6-methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 Racemic N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (100 mg) was
20 resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30) to give the title compound
25 having a shorter retention time (29.2 mg).

MS (API+): [M+H]⁺455.1.
[1020]

Example 260

6-methoxy-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

 Racemic N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (100 mg) was
35 resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL,

manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30) to give the title compound having a longer retention time (33.8 mg).

MS (API+): [M+H]⁺455.1.

5 [1021]

Example 261

N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (85 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile
15 phase: hexane/ethanol = 800/200) to give the title compound having a shorter retention time (29 mg).

MS (API+): [M+H]⁺469.2.

[1022]

Example 262

20 N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-
25 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (85 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 800/200) to give the title compound having a longer retention time (23 mg).

30 MS (API+): [M+H]⁺469.2.

[1023]

Example 263

N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-
35 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (270 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL,
5 manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 70/30, subsequently changed to 50/50), crystallized from hexane/diisopropyl ether to give the title compound having a shorter retention time (116 mg).

MS (API+): [M+H]⁺439.1.

10 ¹H NMR (300 MHz, DMSO-d₆) δ 0.95-1.03 (3H, m), 1.18 (3H, s), 2.27 (3H, s), 4.26-4.37 (1H, m), 4.38-4.47 (1H, m), 4.60-4.93 (2H, m), 7.13 (1H, d, J = 1.5 Hz), 7.27 (2H, d, J = 7.9 Hz), 7.42 (2H, d, J = 9.0 Hz), 7.87 (1H, d, J = 1.5 Hz), 10.54 (1H, d, J = 8.3 Hz), 10.77 (1H, s).

15 [1024]

Example 264

N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 Racemic N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (85 mg) was resolved by HPLC (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile
25 phase: hexane/ethanol = 70/30, subsequently changed to 50/50), and crystallized from hexane/diisopropyl ether to give the title compound having a longer retention time (103 mg).

MS (API+): [M+H]⁺439.1.

[1025]

30 Example 265

optically active 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
35

b]pyrazine-4(1H)-carboxamide (478 mg) was resolved by HPLC (column: CHIRALPAK IA, 400 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300), and recrystallized from hexane/ethyl acetate to
5 give the title compound having a longer retention time (170 mg).

MS (API+): [M+H]⁺449.1.

[1026]

Example 266

10 optically active 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (478 mg) was resolved by HPLC
15 (column: CHIRALPAK IA, 400 mmID×250 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 700/300), and recrystallized from hexane/ethyl acetate to give the title compound having a shorter retention time (156
20 mg).

MS (API+): [M+H]⁺449.1.

[1027]

Example 267

N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
25 carboxamide

[1028]

A) methyl 2-bromo-2-(4-(trifluoromethyl)phenyl)acetate

The title compound was obtained from (4-(trifluoromethyl)phenyl)acetic acid in the same manner as in
30 Steps A and Step D of Example 185.

¹H NMR (300 MHz, DMSO-d₆) δ 3.75 (3H, s), 5.98-6.21 (1H, m), 7.79 (4H, s).

[1029]

35 B) methyl azido(4-(trifluoromethyl)phenyl)acetate

To a solution of methyl 2-bromo-2-(4-(trifluoromethyl)phenyl)acetate (11.8 g) in acetonitrile (265 mL) was added sodium azide (1.55 g) at 0°C. The reaction mixture was stirred at room temperature for 4 hr, and added to
5 water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title
10 compound (10.8 g).

¹H NMR (300 MHz, CDCl₃) δ 3.80 (3H, s), 5.08 (1H, s), 7.53 (2H, d, J = 8.3 Hz), 7.68 (2H, d, J = 8.3 Hz).

[1030]

C) methyl amino (4-(trifluoromethyl)phenyl)acetate

15 To a solution of methyl azido(4-(trifluoromethyl)phenyl)acetate (10.6 g) in methanol (408 mL) was added 10% palladium-carbon (containing 50% water, 1.10 g). The reaction mixture was stirred at room temperature for 6 hr under hydrogen atmosphere, the catalyst was filtered off, and
20 the filtrate was concentrated under reduced pressure to give the title compound (9.13 g).

MS (API+): [M+H]⁺234.1.

[1031]

D) methyl ((tert-butoxycarbonyl) amino)(4-

25 (trifluoromethyl)phenyl)acetate

The title compound was obtained from methyl 2-amino-2-(4-(trifluoromethoxy)phenyl)acetate in the same manner as in Step G of Example 75.

MS (API-): [M-H]⁻332.1.

30 [1032]

E) tert-butyl (2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)carbamate

The title compound was obtained in the same manner as in Step B of Example 250.

35 MS (API-): [M-H]⁻332.2.

[1033]

F) 1-amino-2-methyl-1-(4-(trifluoromethyl)phenyl)propan-2-ol
hydrochloride

The title compound was obtained in the same manner as in
5 Step B of Example 92.

MS (API+), found: 234.1.

[1034]

G) N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-
7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
10 carboxamide

The title compound was obtained from 1-amino-2-methyl-1-
(4-(trifluoromethyl)phenyl)propan-2-ol hydrochloride in the
same manner as in Steps I-L of Example 66.

MS (API+): [M+H]⁺439.2.

15 [1035]

Example 268

optically active N-(2-hydroxy-2-methyl-1-(4-
(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

20 Racemic N-(2-hydroxy-2-methyl-1-(4-
(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (221 mg) was
resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL,
manufactured by Daicel Chemical Industries, Ltd., mobile
25 phase: hexane/ethanol = 50/50) to give the title compound
having a shorter retention time (71.7 mg).

MS (API+): [M+H]⁺439.2.

[1036]

Example 269

30 optically active N-(2-hydroxy-2-methyl-1-(4-
(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-
dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic N-(2-hydroxy-2-methyl-1-(4-
(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-
35 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (221 mg) was

resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 50/50) to give the title compound having a longer retention time (78.3 mg).

5 MS (API+): [M+H]⁺439.2.

[1037]

Example 270

optically active 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 Racemic 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (296 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 450/550) to give the title compound having a shorter retention time (136 mg).

MS (API+): [M+H]⁺ 394.1.

[1038]

20 Example 271

optically active 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

Racemic 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (296 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 450/550) to give the title compound having a longer retention time (134 mg).

30 MS (API+): [M+H]⁺ 394.1.

[1039]

Example 272

N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-

4(1H)-carboxamide

[1040]

A) methyl 2-((5-methyl-3-nitropyridin-2-yl) amino)acetate

To a mixture of methyl glycinate hydrochloride (8.32 g) and 2-chloro-5-methyl-3-nitropyridine (7.62 g) in N,N-dimethylformamide (100 mL) was added triethylamine (18.5 mL). The reaction mixture was stirred at 80°C for 16 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The extract was washed with water, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (6.99 g).

¹H NMR (300 MHz, DMSO-d₆) δ 2.24 (3H, s), 3.64 (3H, s), 4.28 (2H, d, J = 5.7 Hz), 8.27-8.37 (2H, m), 8.60 (1H, t, J = 5.7 Hz).

[1041]

B) 7-methyl-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of methyl 2-((5-methyl-3-nitropyridin-2-yl) amino)acetate (6.99 g) in ethanol (100 mL) was added 10% palladium-carbon (containing 50% water, 2.00 g). The reaction mixture was stirred overnight at room temperature under hydrogen atmosphere, the catalyst was filtered off, and the filtrate was concentrated under reduced pressure. To the residue was added ethanol (100 mL), the reaction mixture was heated at reflux for 5 hr, and the solvent was evaporated to about 1/3 of volume under reduced pressure. To the residue was added diisopropyl ether, and the precipitated solid was collected by filtration, washed with diisopropyl ether, and dried under reduced pressure to give the title compound (4.56 g).

¹H NMR (300 MHz, DMSO-d₆) δ 2.09 (3H, s), 3.85 (2H, d, J = 1.5 Hz), 6.45 (1H, s), 6.72-6.80 (1H, m), 7.39-7.48 (1H, m), 10.31 (1H, s).

[1042]

C) 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

To a mixture of 7-methyl-3,4-dihydropyrido[2,3-b]pyrazin-
5 2(1H)-one (1.00 g) in N,N-dimethylacetamide (40 mL) and
pyridine (15 mL) was added 4-nitrophenyl chloroformate (1.48
g) at room temperature. The reaction mixture was stirred at
room temperature for 16 hr, and water was added thereto. The
precipitated solid was collected by filtration, washed with
10 water, and dried under reduced pressure to give the title
compound (1.63 g).

MS (API+): [M+H]⁺ 329.1.

[1043]

D) 2-amino-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetonitrile
15 3-Fluoro-4-(trifluoromethoxy)benzaldehyde (10 g) was
dissolved in 2 M ammonia/methanol solution (96 mL),
titanium(IV) tetraisopropoxide (15.5 mL) was added thereto
under ice-cooling. The reaction mixture was stirred at the
same temperature for 15 min, trimethylsilanecarbonitrile (9.61
20 mL) was added thereto, and the mixture was stirred at room
temperature for 20 hr. The solvent was evaporated under
reduced pressure, and to the residue were added ethyl acetate
and saturated aqueous sodium hydrogen carbonate solution. The
insoluble substance was filtered off using Celite, and the
25 filtrate was extracted with ethyl acetate. The extract was
washed successively with water and saturated brine, and dried
over anhydrous magnesium sulfate, and the solvent was
evaporated under reduced pressure. The residue was purified by
silica gel column chromatography (ethyl acetate/hexane) to
30 give the title compound (9.4 g).

¹H NMR (300 MHz, CDCl₃) δ 2.01 (2H, brs), 4.93 (1H, s), 7.29-
7.53 (3H, m).

[1044]

E) tert-butyl (cyano(3-fluoro-4-
35 (trifluoromethoxy)phenyl)methyl)carbamate

To a solution of 2-amino-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetonitrile (9.4 g) in tetrahydrofuran (200 mL) was added di-tert-butyl dicarbonate (10.3 mL) and triethylamine (7.27 mL) at room temperature. The
5 reaction mixture was stirred at 40°C for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
10 column chromatography (ethyl acetate/hexane) to give the title compound (8.47 g).

MS (API+): [M+H]⁺ 335.1.

[1045]

F) tert-butyl (2-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-oxoethyl)carbamate
15

To a mixture of tert-butyl (cyano(3-fluoro-4-(trifluoromethoxy)phenyl)methyl)carbamate (9.4 g), potassium carbonate (3.89 g) and dimethyl sulfoxide (200 mL) was added 35% aqueous hydrogen peroxide (4.66 mL) at room temperature.
20 The reaction mixture was stirred at room temperature for 4 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure.
25 The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (3.84 g).

MS (API+), found: 253.0.

[1046]

G) methyl 2-((tert-butoxycarbonyl)amino)-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetate
30

To a solution of tert-butyl (2-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-oxoethyl)carbamate (3.84 g) in methanol (100 mL) was added 8 M aqueous sodium hydroxide solution (2.8 mL) at room temperature. The reaction mixture
35 was heated at reflux overnight, and the methanol was

evaporated under reduced pressure. The residue was neutralized with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent
5 was evaporated under reduced pressure. To a mixture of the residue (3.85 g) and potassium carbonate (1.81 g) in N,N-dimethylformamide (50 mL) was added methyl iodide (0.750 mL) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, water was added thereto, and the mixture
10 was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title
15 compound (1.90 g).

MS (API+), found: 268.0.

[1047]

H) tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)carbamate

20 To a solution of methyl 2-((tert-butoxycarbonyl)amino)-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetate (1.90 g) in tetrahydrofuran (60 mL) was slowly added 3 M methylmagnesium bromide/diethyl ether solution (5.17 mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hr, 1 M
25 hydrochloric acid at 0°C was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
30 column chromatography (hexane/ethyl acetate) to give the title compound (1.53 g).

¹H NMR (300 MHz, CDCl₃) δ 1.06 (3H, s), 1.29-1.50 (13H, m), 4.38-4.52 (1H, m), 5.47-5.60 (1H, m), 7.07-7.30 (3H, m).

[1048]

35 I) 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-

methylpropan-2-ol hydrochloride

To tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)carbamate (3.01 g) was added 4 M hydrogen chloride/ethyl acetate solution (30 mL). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (2.22 g).

MS (API+), found: 268.1.

10 [1049]

J) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 4-nitrophenyl 7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (328 mg) and 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methylpropan-2-ol hydrochloride (334 mg) in N,N-dimethylformamide (10 mL) was added triethylamine (0.335 mL) at room temperature. The reaction mixture was stirred at room temperature for 16 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and recrystallized from hexane/ethyl acetate to give the title compound (357 mg).

MS (API+): [M+H]⁺ 457.1.

¹H NMR (300 MHz, CDCl₃) δ 1.16 (3H, s), 1.37 (3H, s), 1.62 (1H, s), 2.32 (3H, s), 4.54-4.74 (2H, m), 4.85 (1H, d, J = 8.3 Hz), 7.02 (1H, d, J = 1.1 Hz), 7.14-7.21 (1H, m), 7.21-7.30 (2H, m), 7.86-7.95 (1H, m), 8.78 (1H, s), 10.84 (1H, d, J = 8.3 Hz).

[1050]

Example 273

optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-

b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (349 mg) was resolved by HPLC

5 (column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 90/10, subsequently changed to 70/30), and recrystallized from acetone/heptane to give the title compound having a shorter retention time (140 mg).

10 MS (API+): [M+H]⁺ 457.1.

[1051]

Example 274

optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-

15 b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide (349 mg) was resolved by HPLC

(column: CHIRALPAK IA, 50 mmID×500 mmL, manufactured by Daicel
20 Chemical Industries, Ltd., mobile phase: hexane/ethanol = 90/10, subsequently changed to 70/30), and recrystallized from acetone/heptane to give the title compound having a longer retention time (107 mg).

MS (API+): [M+H]⁺ 457.1.

25 [1052]

Example 275

N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

30 [1053]

A) ethyl (5-bromopyridin-3-yl)carbamate

To a mixture of 5-bromopyridin-3-amine (35.4 g) and pyridine (19.8 mL) in tetrahydrofuran (600 mL) was slowly added ethyl chloroformate (23.4 mL) at 0°C. The reaction
35 mixture was stirred overnight at room temperature under

nitrogen atmosphere, and poured into water, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To the
5 residue was added diisopropyl ether (400 mL), the mixture was stirred at room temperature for 20 min, and the obtained solid was collected by filtration to give the title compound (29.0 g).

MS (API+): [M+H]⁺ 245.1.

10 [1054]

B) ethyl (5-bromo-2-nitropyridin-3-yl)carbamate

To a solution of ethyl (5-bromopyridin-3-yl)carbamate (19.0 g) in conc. sulfuric acid (37.2 mL) was slowly added fuming nitric acid (26.1 mL) at 0°C. The reaction mixture was
15 stirred at room temperature for 72 hr under nitrogen atmosphere. The reaction mixture was slowly poured into ice, the pH of the mixture was adjusted to 9 with 28% aqueous ammonia, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over
20 anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. To the residue was added ethyl acetate/hexane, the mixture was stirred for 15 min, and the resulting solid was collected by filtration, and washed with hexane to give the title compound (13.9 g).

25 MS (API+): [M+H]⁺ 290.1.

[1055]

C) 5-methoxy-2-nitropyridin-3-amine

To a solution of ethyl (5-bromo-2-nitropyridin-3-yl)carbamate (31.1 g) in methanol (900 mL) was added 28%
30 sodium methoxide/methanol solution (83.0 g) at room temperature. The reaction mixture was stirred at 65°C for 4 hr under nitrogen atmosphere, and the solvent was evaporated to about 150 mL under reduced pressure. To the residue was added saturated aqueous ammonium chloride solution, and the mixture
35 was stirred at room temperature for 20 min, and the solvent

was evaporated to about 100 mL under reduced pressure. The resulting solid was collected by filtration, and washed with water to give the title compound (16.6 g).

MS (API+): [M+H]⁺ 170.2.

5 [1056]

D) 2-chloro-N-(5-methoxy-2-nitropyridin-3-yl)acetamide

To a solution of 5-methoxy-2-nitropyridin-3-amine (24.7 g) in N,N-dimethylformamide (740 mL) was slowly added a solution of chloroacetyl chloride (23.2 mL) in tetrahydrofuran
10 (40 mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hr under nitrogen atmosphere, and the solvent was evaporated under reduced pressure. To the residue was added saturated aqueous sodium hydrogen carbonate solution, and the mixture was extracted with ethyl acetate. The extract
15 was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether/hexane, and the mixture was stirred at room temperature for 30 min. The obtained solid was collected by filtration,
20 and washed with diisopropyl ether/hexane to give the title compound (34.1 g).

MS (API+): [M+H]⁺ 246.0.

[1057]

E) N-(2-amino-5-methoxypyridin-3-yl)-2-chloroacetamide

25 A mixture of 2-chloro-N-(5-methoxy-2-nitropyridin-3-yl)acetamide (34.1 g), ammonium chloride (44.6 g), iron (27.1 g), ethanol (823 mL) and water (206 mL) was stirred at 75°C for 40 min, and the solvent was evaporated under reduced pressure. To the residue were added tetrahydrofuran (500 mL) and
30 saturated aqueous sodium hydrogen carbonate solution (300 mL), and the reaction mixture was stirred for 15 min. The insoluble substance was filtered off through Celite, to the filtrate was added saturated brine, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine,
35 and the solvent was evaporated under reduced pressure. To the

residue was added diisopropyl ether/hexane, and the obtained solid was collected by filtration, and washed with diisopropyl ether/hexane to give the title compound (26.3 g).

MS (API+): [M+H]⁺ 216.1.

5 [1058]

F) benzyl (3-((chloroacetyl)amino)-5-methoxypyridin-2-yl)carbamate

To a mixture of N-(2-amino-5-methoxypyridin-3-yl)-2-chloroacetamide (2.74 g), pyridine (5.14 mL) and
10 tetrahydrofuran (85 mL) was slowly added benzyl chloroformate (2.72 mL) at 0°C. The reaction mixture was stirred at room temperature for 1 hr, and ethyl acetate was added thereto. The mixture was washed with a mixture of saturated aqueous sodium hydrogen carbonate solution and saturated brine, and the
15 solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate), and the obtained solid was washed with diisopropyl ether to give the title compound (3.79 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.82 (3H, s), 4.36 (2H, s), 5.13
20 (2H, s), 7.30–7.43 (5H, m), 7.87 (1H, d, J = 3.0 Hz), 7.95 (1H, d, J = 2.6 Hz), 9.46 (1H, s), 9.64 (1H, s).

[1059]

G) benzyl 7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate

25 To a solution of benzyl (3-((chloroacetyl)amino)-5-methoxypyridin-2-yl)carbamate (200 mg) in N,N-dimethylformamide (11 mL) was added cesium carbonate (279 mg) at 50°C, and the reaction mixture was stirred at 50°C for 30 min. The reaction mixture was diluted with ethyl acetate, and
30 the mixture was washed with saturated brine, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (115 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 3.81 (3H, s), 4.36 (2H, s), 5.20
35 (2H, s), 6.92 (1H, d, J = 2.6 Hz), 7.27–7.43 (5H, m), 7.84 (1H,

d, $J = 2.6$ Hz), 10.72 (1H, s).

[1060]

H) 7-methoxy-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one

To a solution of benzyl 7-methoxy-2-oxo-2,3-
5 dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (0.722 g) in
tetrahydrofuran (46.1 mL) was added 10% palladium-carbon
(containing 50% water, 0.049 g), and the reaction mixture was
stirred at room temperature for 15 hr under hydrogen
atmosphere. To the reaction mixture was added methanol at 50°C
10 to dissolve the precipitated solid, the catalyst was filtered
off, and the filtrate was concentrated under reduced pressure.
The obtained solid was washed with diisopropyl ether to give
the title compound (0.380 g).

MS (API+): $[M+H]^+$ 180.2.

15 [1061]

I) 2-amino-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetonitrile

3-Fluoro-4-(trifluoromethoxy)benzaldehyde (10 g) was
dissolved in 2 M ammonia/methanol solution (96 mL), and
titanium(IV) tetraisopropoxide (15.5 mL) was added thereto
20 under ice-cooling. The reaction mixture was stirred at the
same temperature for 15 min, trimethylsilanecarbonitrile (9.61
mL) was added thereto, and the mixture was stirred at room
temperature for 20 hr. The solvent was evaporated under
reduced pressure, and to the residue were added ethyl acetate
25 and saturated aqueous sodium hydrogen carbonate solution. The
insoluble substance was filtered off using Celite, and the
filtrate was extracted with ethyl acetate. The extract was
washed successively with water and saturated brine, and dried
over anhydrous magnesium sulfate, and the solvent was
30 evaporated under reduced pressure. The residue was purified by
silica gel column chromatography (ethyl acetate/hexane) to
give the title compound (9.4 g).

^1H NMR (300 MHz, CDCl_3) δ 2.01 (2H, brs), 4.93 (1H, s), 7.29-
7.53 (3H, m).

35 [1062]

J) tert-butyl (cyano(3-fluoro-4-(trifluoromethoxy)phenyl)methyl)carbamate

To a solution of 2-amino-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetonitrile (9.4 g) in
5 tetrahydrofuran (200 mL) were added di-tert-butyl dicarbonate (10.3 mL) and triethylamine (7.27 mL) at room temperature. The reaction mixture was stirred at 40°C for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over
10 anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (8.47 g).

MS (API+): [M+H]⁺ 335.1.

15 [1063]

K) tert-butyl (2-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-oxoethyl)carbamate

To a mixture of tert-butyl (cyano(3-fluoro-4-(trifluoromethoxy)phenyl)methyl)carbamate (9.4 g), potassium
20 carbonate (3.89 g) and dimethyl sulfoxide (200 mL) was added 35% aqueous hydrogen peroxide (4.66 mL) at room temperature. The reaction mixture was stirred at room temperature for 4 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water
25 and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (3.84 g).

MS (API+), found: 253.0.

30 [1064]

L) methyl 2-((tert-butoxycarbonyl) amino)-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetate

To a solution of tert-butyl (2-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-oxoethyl)carbamate (3.84 g) in
35 methanol (100 mL) was added 8 M aqueous sodium hydroxide

solution (2.8 mL) at room temperature. The reaction mixture was heated at reflux overnight, and the methanol was evaporated under reduced pressure. The residue was neutralized with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To a mixture of the residue (3.85 g) and potassium carbonate (1.81 g) in N,N-dimethylformamide (50 mL) was added methyl iodide (0.750 mL) at room temperature. The reaction mixture was stirred at room temperature for 2 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.90 g).

MS (API+), found: 268.0.

[1065]

M) tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)carbamate

To a solution of methyl 2-((tert-butoxycarbonyl) amino)-2-(3-fluoro-4-(trifluoromethoxy)phenyl)acetate (1.90 g) in tetrahydrofuran (60 mL) was slowly added 3 M methylmagnesium bromide/diethyl ether solution (5.17 mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hr, 1 M hydrochloric acid at 0°C was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (1.53 g).

¹H NMR (300 MHz, CDCl₃) δ 1.06 (3H, s), 1.29-1.50 (13H, m), 4.38-4.52 (1H, m), 5.47-5.60 (1H, m), 7.07-7.30 (3H, m).

[1066]

N) 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methylpropan-2-ol hydrochloride

To tert-butyl (1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)carbamate (3.01 g) was added 4 M hydrogen chloride/ethyl acetate solution (30 mL). The reaction mixture was stirred at room temperature for 1 hr, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (2.22 g).

MS (API+), found: 268.1.

[1067]

O) N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

To a solution of 7-methoxy-3,4-dihydropyrido[2,3-b]pyrazin-2(1H)-one (233 mg) in tetrahydrofuran (15 mL) were added 4-nitrophenyl chloroformate (315 mg) and N,N-diisopropylethylamine (0.341 mL) at room temperature. The reaction mixture was stirred at room temperature for 24 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. To a solution of the residue in N,N-dimethylformamide (15 mL) were added 1-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methylpropan-2-ol hydrochloride (395 mg) and triethylamine (0.544 mL) at room temperature. The reaction mixture was stirred at room temperature for 20 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (492 mg).

MS (API+): [M+H]⁺473.1.

¹H NMR (300 MHz, CDCl₃) δ 1.15 (3H, s), 1.38 (3H, s), 1.61 (1H, s), 3.89 (3H, s), 4.52-4.74 (2H, m), 4.84 (1H, d, J = 8.3 Hz), 6.79 (1H, d, J = 2.6 Hz), 7.14-7.31 (3H, m), 7.75 (1H, d, J =
5 2.6 Hz), 8.44-8.60 (1H, m), 10.49 (1H, d, J = 8.3 Hz).

[1068]

Example 276

optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
10 b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide (369 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel
15 Chemical Industries, Ltd., mobile phase: hexane/ethanol = 850/150), and recrystallized from acetone/heptane to give the title compound having a shorter retention time (148 mg).

MS (API+): [M+H]⁺ 473.1.

¹H NMR (300 MHz, CDCl₃) δ 1.15 (3H, s), 1.37 (3H, s), 1.64 (1H, s), 3.89 (3H, s), 4.53-4.72 (2H, m), 4.85 (1H, d, J = 8.3 Hz),
20 6.81 (1H, d, J = 2.6 Hz), 7.13-7.31 (3H, m), 7.75 (1H, d, J = 2.6 Hz), 8.90 (1H, brs), 10.50 (1H, d, J = 8.3 Hz).

[1069]

Example 277

25 optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide

Racemic N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
30 b]pyrazine-4(1H)-carboxamide (369 mg) was resolved by HPLC (column: CHIRALPAK AD, 50 mmID×500 mmL, manufactured by Daicel Chemical Industries, Ltd., mobile phase: hexane/ethanol = 850/150), and recrystallized from acetone/heptane to give the title compound having a longer retention time (144 mg).

35 MS (API+): [M+H]⁺473.1.

¹H NMR (300 MHz, CDCl₃) δ 1.15 (3H, s), 1.37 (3H, s), 1.63 (1H, s), 3.89 (3H, s), 4.52-4.74 (2H, m), 4.85 (1H, d, J = 8.3 Hz), 6.81 (1H, d, J = 2.6 Hz), 7.11-7.31 (3H, m), 7.75 (1H, d, J = 2.6 Hz), 8.78 (1H, brs), 10.50 (1H, d, J = 8.3 Hz).

5 [1070]

Example 278

N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide

10 The title compound was obtained from 4-nitrophenyl 7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate and 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanamine hydrochloride in the same manner as in Step D of Example 19.

15 MS (API+): [M+H]⁺ 473.1.

[1071]

Example 279

N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
20 carboxamide

The title compound was obtained from 4-nitrophenyl 6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate and 1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethanamine hydrochloride in the same manner as in Step
25 D of Example 19.

MS (API+): [M+H]⁺ 473.1.

[1072]

Example 280

N-(2-cyano-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-
30 carboxamide

[1073]

A) 3-amino-3-(3-fluoro-4-(trifluoromethoxy)phenyl)propanoic acid

35 To a solution of 3-fluoro-4-

(trifluoromethoxy)benzaldehyde (2.08 g) in ethanol (40 mL) were added malonic acid (1.35 g) and ammonium acetate (1.16 g) at room temperature, and the mixture was stirred at 80°C for 20 hr. The reaction mixture was filtered, and the obtained solid
5 was washed with ethanol to give the title compound (1.48 g).

MS (API+): [M+H]⁺ 268.1.

[1074]

B) tert-butyl (3-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-3-oxopropyl)carbamate

10 To a solution of 3-amino-3-(3-fluoro-4-(trifluoromethoxy)phenyl)propanoic acid (1.48 g) in tetrahydrofuran (50 mL) were added di-tert-butyl dicarbonate (1.54 mL) and 2 M aqueous sodium hydroxide solution (4.0 mL), and the mixture was stirred at room temperature for 20 hr. The
15 reaction mixture was neutralized with 1 M hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. A mixture of the residue,
20 1H-benzotriazol-1-ol ammonium salt (1.01 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.27 g) in N,N-dimethylformamide (35 mL) was stirred at room temperature for 24 hr, water was added thereto, and the mixture was extracted with ethyl acetate. The extract was
25 washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was crystallized from hexane/ethyl acetate to give the title compound (1.38 g).

30 MS (API+): [M-H]⁻ 365.0.

[1075]

C) tert-butyl (2-cyano-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethyl)carbamate

To a solution of tert-butyl (3-amino-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-3-oxopropyl)carbamate (1.0 g) in
35

toluene/tetrahydrofuran (1:1, 75 mL) was added thionyl chloride (0.498 mL) at room temperature. The reaction mixture was stirred at 60°C for 20 hr, and the solvent was evaporated under reduced pressure. To the residue was added water, and
5 the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to
10 give the title compound (380 mg).

MS (API+): [M+H]⁺ 349.1.

[1076]

D) 3-amino-3-(3-fluoro-4-

(trifluoromethoxy)phenyl)propanenitrile hydrochloride

15 A mixture of tert-butyl (2-cyano-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethyl)carbamate (431 mg) and 4 M hydrogen chloride/ethyl acetate (5 mL) was stirred at room temperature for 30 min. The solvent was evaporated under reduced pressure, and the residue was washed with diisopropyl
20 ether to give the title compound (328 mg).

MS (API+), found: 249.0.

[1077]

E) N-(2-cyano-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-

25 carboxamide

To a solution of 4-nitrophenyl 7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxylate (275 mg) and 3-amino-3-(3-fluoro-4-(trifluoromethoxy)phenyl)propanenitrile hydrochloride (296 mg) in N,N-dimethylformamide (10 mL) was
30 added triethylamine (0.334 mL) at room temperature. The reaction mixture was stirred at room temperature for 20 hr, and then at 60°C for 24 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The extract was washed successively with water and saturated brine,
35 and dried over anhydrous magnesium sulfate, and the solvent

was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (hexane/ethyl acetate) to give the title compound (349 mg).

MS (API+): $[M+H]^+$ 454.1.

5 [1078]

The compounds of Examples according to the above-mentioned method or a method analogous thereto are shown in the following tables. Mass in the tables means measured value.

[1079]

10 Table 1-1

Example No.	IUPAC Name	Structure	Mass (M+1)	Mass (M-1)
1	N-(1-(4-methoxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide			339.0
2	2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.2	
3	N-(1-(4-methoxyphenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		355.1	
4	7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.2	
5	optically active 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
6	optically active 7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
7	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		411.1	
8	optically active 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.1	
9	optically active 2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.1	
10	8-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
11	6-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
12	N-(2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	












[1080]

Table 1-2

13	N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		397.1	
14	N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.0	
15	N-(cyclopropyl(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		407.0	
16	2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)butyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.0	
17	1-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide			
	¹ H NMR (300 MHz, CDCl ₃) δ 0.97 (3H, t, J = 7.5 Hz) 1.83-1.93 (2H, m), 3.36 (3H, s), 4.69 (2H, s), 4.91 (1H, q, J = 6.9 Hz), 7.09 (1H, dd, J = 7.8, 5.1 Hz), 7.18 (2H, d, J = 8.4 Hz), 7.27-7.37 (3H, m), 8.02 (1H, d, J = 4.8 Hz), 10.31 (1H, d, J = 7.5 Hz).			
18	N-(1-(4-(difluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		377.1	
19	2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		463.2	
20	2-oxo-7-phenyl-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		471.3	
21	6-bromo-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide		471.9	
22	3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide		394.0	
23	6-methyl-3-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-3,4-dihydropyrazino[2,3-b]pyrazine-1(2H)-carboxamide		410.2	

[1081]

Table 1-3

24	N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		411.2	
25	N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		411.1	
26	7-iodo-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		521.1	
27	2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-7-vinyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		421.1	
28	N-(1-(biphenyl-4-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		387.2	
29	7-ethyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		423.1	
30	N-(1-(6-methoxypyridazin-3-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		343.2	
31	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		413.1	
32	N-(1-(2-methoxy-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.1	
33	N-(1-(4-hydroxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		327.1	
34	N-(1-(2-fluoro-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		413.2	

[1082]

Table 1-4

35	7-cyclopropyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		435.1	
36	7-isopropyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		437.2	
37	6-chloro-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		429.1	
38	N-(1-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		391.2	
39	2-oxo-N-(1-(4-(prop-1-en-2-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		351.2	
40	N-(1-(4-(isopropylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		353.2	
41	N-(1-(4-(3,6-dihydro-2H-pyran-4-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		393.2	
42	N-(2-amino-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		410.1	
43	N-(cyano(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		392.1	
44	7-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide			418.0
45	2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-6-(trifluoromethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		463.1	
46	2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-6-vinyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		421.2	













[1083]

Table 1-5

47	6-benzyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		485.2	
48	2-oxo-N-(1-(4-(tetrahydro-2H-pyran-4-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.2	
49	N-(1-(6-methoxypyridin-3-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		342.1	
50	N-(1-(5-methoxypyrimidin-2-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		343.2	
51	6-ethyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		423.1	
52	6-methoxy-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.1	
53	N-(1-(4-(cyclopropylmethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		381.2	
54	2-oxo-N-(1-(4-(trifluoromethyl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		379.2	
55	2-oxo-N-(pyridin-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		444.2	
56	N-((3,5-dimethyl-1,2-oxazol-4-yl)(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		462.1	
57	N-((1-methyl-1H-pyrazol-4-yl)(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		447.2	













[1084]

Table 1-6

58	N-(2-(dimethylamino)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		424.1	
59	N-(2-(methylsulfanyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		427.1	
60	N-(2-(methylsulfonyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		459.2	
61	N-(1-(5-methoxypyridin-2-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		342.1	
62	6-cyano-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide			418.2
63	7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		441.2	
64	7-methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		441.2	
65	7-methoxy-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		441.2	
66	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		459.1	
67	N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		459.1	
68	N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		459.1	
69	7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.1	













[1085]

Table 1-7

70	optically active 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.1	
71	optically active 7-methoxy-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.1	
72	7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		465.2	
73	7-cyclopropyl-N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		465.1	
74	7-cyclopropyl-N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		465.1	
75	7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		483.2	
76	optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		483.2	
77	optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		483.2	
78	2-oxo-N-(1-(4-(2,2,2-trifluoroethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
79	N-(2-(methylsulfinyl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	
80	2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-imidazol-4-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		383.1	
81	2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-1,2,4-triazol-3-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		384.1	

[1086]

Table 1-8

82	N-(1-(4-isopropoxyphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		367.2	
83	N-(1-(4-chlorophenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		345.1	
84	2-oxo-N-(1-(2-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.1	
85	N-(1-(4-methylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		325.2	
86	2-oxo-N-(1-(1-(2,2,2-trifluoroethyl)-1H-pyrazol-4-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		383.1	
87	N-(1-(3-methyl-4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
88	N-(2-(dimethylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		438.1	
89	N-(2-(methylamino)-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		424.1	
90	N-(2-isopropoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.2	
91	N-(2-(cyclopentyloxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		465.2	
92	N-(3-amino-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		424.1	
93	N-(3-(dimethylamino)-3-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		452.2	

[1087]

Table 1-9

94	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.2	
95	3-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	
96	N-(1-(4-(difluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		361.2	
97	N-(2-cyano-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		406.2	
98	2-oxo-N-(1-(3-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.2	
99	N-(1-(4-cyclopropylphenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		351.2	
100	N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.2	
101	2-oxo-N-(2-(pyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		450.2	
102	2-oxo-N-(2-(2-oxopyrrolidin-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		464.2	
103	3-(((2-oxo-2,3-dihydropyrido[2,3-b]pyrazin-4(1H)-yl)carbonyl)amino)-3-(4-(trifluoromethoxy)phenyl)propyl acetate		453.2	
104	N-(3-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		410.8	
105	N-(1-(4-(azetidin-1-yl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide			364.2

[1088]

Table 1-10

106	N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.2	
107	N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.1	
108	N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		447.1	
109	N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		395.1	
110	N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.2	
111	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		429.1	
112	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	
113	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		495.2	
114	7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
115	7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		435.2	
116	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		477.2	
117	N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		495.2	

[1089]

Table 1-11

118	N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		495.1	
119	7-cyclopropyl-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
120	7-cyclopropyl-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
121	N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		429.1	
122	N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		429.1	
123	N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	
124	N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	
125	7-cyclopropyl-N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	
126	optically active N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		447.1	
127	optically active N-(2-(difluoromethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		447.1	
128	7-cyclopropyl-N-((1R)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	
129	7-cyclopropyl-N-((1S)-1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	













[1090]

Table 1-12

130	N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		477.2	
131	N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1H-pyrazol-1-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		477.2	
132	N-(1-(4-bromophenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		419.1	
133	N-(2-(2-methoxyethoxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	
134	N-(2-methoxy-1-(4-(1H-pyrazol-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		407.2	
135	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-3-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		491.2	
136	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		491.2	
137	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-(1-methyl-1H-pyrazol-5-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		491.2	
138	7-isopropoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.1	
139	7-(difluoromethoxy)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		477.2	
140	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-oxo-6,7-dihydropteridine-8(5H)-carboxamide		412.2	
141	7-ethoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.2	

[1091]

Table 1-13

142	7-methyl-N-(oxetan-3-yl(4-(trifluoromethoxy)phenyl)methyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		437.1	
143	N-(1-(3-fluoro-4-(1H-pyrazol-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.2	
144	N-(2-methoxy-1-(5-(1H-pyrazol-1-yl)pyridin-2-yl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		408.2	
145	N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.2	
146	N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.2	
147	7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		390.2	
148	7-methyl-2-oxo-N-(1-(4-(pyrrolidin-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		394.3	
149	7-(2-hydroxypropan-2-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	
150	7-(azetidin-1-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		466.2	
151	N-(1-(imidazo[1,2-a]pyridin-7-yl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		365.2	
152	7-methyl-2-oxo-N-(1-(4-(trifluoromethoxy)phenyl)cyclopropyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		407.1	
153	7-fluoro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		429.1	

[1092]

Table 1-14

154	N-(2-methoxy-1-(4-(1H-pyrrol-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		406.2	
155	7-chloro-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		445.1	
156	N-(2-methoxy-1-(4-(pyrrolidin-1-yl)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		410.3	
157	N-(2-methoxy-1-(4-(1H-pyrrol-1-yl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		392.2	
158	N-(2-methoxy-1-(4-(pyrrolidin-1-yl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		396.2	
159	N-(1-(2,2-difluoro-1,3-benzodioxol-5-yl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		391.1	
160	7-methyl-2-oxo-N-(2-(1H-pyrazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		461.2	
161	optically active 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		390.2	
162	7-methyl-2-oxo-N-(1-(5-(1H-pyrazol-1-yl)-2-thienyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		397.0	
163	optically active 7-methyl-2-oxo-N-(1-(4-(1H-pyrrol-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		390.2	
164	7-methyl-2-oxo-N-(2-(1H-1,2,4-triazol-1-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		462.2	
165	7-methyl-2-oxo-N-(1-(pyrazolo[1,5-a]pyridin-5-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		365.2	

[1093]

Table 1-15

166	7-methyl-N-(2-(oxetan-3-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
167	7-(difluoromethyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		461.1	
168	7-methyl-2-oxo-N-(tetrahydro-2H-pyran-4-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		465.2	
169	N-(2-(4-hydroxytetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		495.1	
170	7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
171	N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)pentyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.2	
172	N-(3-((tert-butyl(diphenyl)silyl)oxy)-2,2-dimethyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		691.4	
173	N-(2-(1,3-dioxo-1,3-dihydro-2H-isindol-2-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		526.2	
174	N-(2-cyclopropyl-2-oxo-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		435.1	
175	N-(2-cyclopropyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		437.2	
176	7-methyl-2-oxo-N-(2-(tetrahydro-2H-pyran-4-yl)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		479.2	
177	2-oxo-N-(2-oxo-1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		409.1	













[1094]

Table 1-16

178	8-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		441.2	
179	N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		411.1	
180	2-oxo-N-(2,2,2-trifluoro-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		435.1	
181	7-methyl-2-oxo-N-(2-(pyridin-3-yloxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		488.2	
182	N-(2-hydroxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		411.1	
183	N-(1-(3,5-difluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		461.1	
184	N-(3-hydroxy-2,2-dimethyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.2	
185	N-(2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		427.1	
186	2-oxo-N-(1-(4-(2H-1,2,3-triazol-2-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		378.2	
187	N-(2-fluoro-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		441.2	
188	N-(1-(3-fluoro-4-(pyrrolidin-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		428.2	
189	N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		424.2	

[1095]

Table 1-17

190	8-(2-hydroxypropan-2-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	
191	8-acetyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.1	
192	8-(hydroxymethyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		441.2	
193	N-(2-ethyl-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)butyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.2	
194	N-(2-amino-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		396.1	
195	N-((4-bromo-2-fluorophenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		553.1	
196	N-((4-bromo-2-hydroxyphenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		551.1	
197	N-(2-((5-bromopyridin-3-yl)oxy)-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		566.1	
198	N-((2-fluorophenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		475.2	
199	N-((2-hydroxyphenyl)(4-(trifluoromethoxy)phenyl)methyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		473.2	
200	7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
201	7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	

[1096]

Table 1-18

202	7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
203	7-methyl-2-oxo-N-(tetrahydrofuran-2-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
204	7-methoxy-2-oxo-N-(1-(4-(pyrrolidin-1-yl)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide			408.2
205	N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.3	
206	7-methyl-2-oxo-N-(tetrahydrofuran-3-yl(4-(trifluoromethoxy)phenyl)methyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.2	
207	N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.2	
208	N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
209	7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.1	
210	7-methyl-2-oxo-N-(3,3,3-trifluoro-2-hydroxy-1-(4-(trifluoromethoxy)phenyl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		479.1	
211	N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		440.2	
212	6-(4-methoxyphenyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		517.1	
213	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(pyridin-3-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		488.1	

[1097]

Table 1-19

214	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-phenyl-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		487.1	
215	6-(2-methoxyphenyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		517.1	
216	6-(3-methoxyphenyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		517.2	
217	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(pyridin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		488.1	
218	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(2-thienyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		493.1	
219	6-(3-furyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		477.1	
220	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(3-thienyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		493.1	
221	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1H-pyrazol-5-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		491.1	
222	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1H-pyrazol-3-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		491.1	
223	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1H-pyrazol-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		491.1	
224	6-(3,5-dimethyl-1,2-oxazol-4-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		506.1	
225	N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	

[1098]

Table 1-20

226	optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.1	
227	optically active 7-cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.1	
228	N-(2-methoxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		453.2	
229	optically active N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		440.2	
230	optically active N-(1-(3-fluoro-4-(1H-pyrrol-1-yl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		440.2	
231	optically active 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		435.1	
232	optically active N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	
233	optically active 7-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethyl)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		435.1	
234	optically active N-(1-(3-fluoro-4-(trifluoromethyl)phenyl)-2-methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		443.1	
235	6-(3,6-dihydro-2H-pyran-4-yl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		493.1	
236	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		506.1	
237	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(1,2,3,6-tetrahydropyridin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		492.1	

[1099]

Table 1-21

238	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		425.0	
239	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(pyridin-2-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		488.1	
240	6-(2-furyl)-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		477.1	
241	6-cyclopropyl-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		451.1	
242	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(tetrahydro-2H-pyran-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		495.1	
243	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-(1-methylpiperidin-4-yl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		508.1	
244	N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-2-oxo-6-(piperidin-4-yl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		494.1	
245	6-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
246	7-methoxy-N-(2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
247	N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.1	
248	N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.1	
249	N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.1	

[1100]

Table 1-22

250	N-(2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.1	
251	7-methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
252	7-methoxy-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
253	N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.1	
254	N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.1	
255	7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		394.1	
256	7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		449.2	
257	optically active N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.1	
258	N-(3-methoxy-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.1	
259	6-methoxy-N-((1S)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
260	6-methoxy-N-((1R)-2-methoxy-1-(4-(trifluoromethoxy)phenyl)ethyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		455.1	
261	N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	

[1101]

Table 1-23

262	N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		469.2	
263	N-((1S)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.1	
264	N-((1R)-2-hydroxy-2-methyl-1-(4-(trifluoromethoxy)phenyl)propyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.1	
265	optically active 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		449.1	
266	optically active 7-cyclopropyl-N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		449.1	
267	N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.2	
268	optically active N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.2	
269	optically active N-(2-hydroxy-2-methyl-1-(4-(trifluoromethyl)phenyl)propyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		439.2	
270	optically active 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		394.1	
271	optically active 7-methyl-2-oxo-N-(1-(5-(trifluoromethyl)pyridin-2-yl)propyl)-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		394.1	
272	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		457.1	
273	optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		457.1	

[1102]

Table 1-24

274	optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		457.1	
275	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		473.1	
276	optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		473.1	
277	optically active N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-methylpropyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		473.1	
278	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-7-methoxy-6-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		473.1	
279	N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-2-methoxyethyl)-6-methoxy-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		473.1	
280	N-(2-cyano-1-(3-fluoro-4-(trifluoromethoxy)phenyl)ethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-4(1H)-carboxamide		454.1	

[1103]

5 Experimental Example 1

PDE Enzyme Inhibition Assay

Human PDE2A3 enzyme was generated from Sf9 cells transfected with the full-length gene. The extracted enzyme from Sf9 cells was purified by His-tag affinity column and gel filtration. The enzyme was stored at -70°C until use. PDE activity was measured using a SPA (Scintillation Proximity Assay) (GE Healthcare). To evaluate the inhibitory activity, 10 µl of serial diluted compounds were incubated with 20 µl of PDE enzyme in assay buffer (50 mM HEPES-NaOH, 8.3 mM MgCl₂, 1.7 mM EGTA, 0.1 % BSA (pH 7.4)) for 30 min. at room temperature. Final concentration of DMSO in the assay was 1 percent as

compounds were tested in duplicate in 96-well half-area plates (Corning) or 384-well OptiPlate (PerkinElmer). To start the reaction, 10 μ l of substrate [^{3} H] cGMP (77 nM, PerkinElmer) was added for a final assay volume of 40 μ l. After 60 min
5 incubation at room temperature, 20 μ l of 20 mg/ml yttrium SPA beads containing Zinc sulphate was added to terminate the PDE reaction. After being settled for further 1 hour, the assay plates were counted in a scintillation counter (PerkinElmer) to allow calculation of inhibition rate. Inhibition rate was
10 calculated on the basis of 0% control wells with enzyme and DMSO, and 100% control wells without enzyme. The results are shown in Table 2.

[1104]

Table 2

Example No.	Inhibition Rate (%) (1 μ M)
5	95
9	96
14	99
15	94
25	99
31	96
35	95
52	93
54	95
60	97
64	99
66	101
67	101
70	98
73	99
76	100
93	96
107	99
117	99
120	99
121	99
123	98
129	99
130	99
134	97
136	99
143	98
146	98
152	97
160	99
162	97
167	99
200	100
207	98
222	99
226	100
229	99
231	100
232	100
251	100
253	100
257	99
259	99
261	100
263	100
266	100
268	99
273	98
276	101

[1105]

Experimental Example 2

The improvement effect on MK-801 ((5S,10R)-5-methyl-10,11-dihydro-5H-5,10-epiminodibenzo[a,d][7]annulene)-induced disorder in contextual fear conditioning test.
[1106]

5 Experimental Animals

Male C57BL/6 mice were purchased from CLEA Japan, Inc. They were used for an experiment after a habituation period of at least one week after carrying them in the animal experimental facility. They were bred in the animal
10 experimental facility with environment in which 12-hours light-dark cycle was performed, humidity and temperature were controlled, and free water drinking and feeding were allowed. Handling of experimental animals and experimental procedure of this study were approved by the Experimental Animal Ethics
15 Committee of Takeda Pharmaceutical Company Limited.
[1107]

Used Drugs

A test compound was suspended in 0.5% methylcellulose solution, and orally administered. MK-801 (maleate) (Sigma-
20 Aldrich, St Louis, MO) was dissolved in saline, and subcutaneously administered. All drugs were administered to mice at a dose of 10 mL/kg.
[1108]

The improvement effect on MK-801-induced disorder in
25 contextual fear conditioning test.

Contextual fear conditioning test is widely used as memory and learning test system depending on the hippocampus and amygdala. The improvement effect of test compounds on MK-801-induced disorder was investigated by contextual fear
30 conditioning test. Administrations of 0.5% methylcellulose and saline to the control groups were carried out 60 minutes and 30 minutes before tests, respectively. Administrations of 0.5% methylcellulose and MK-801 (maleate) to the vehicle groups were carried out 60 minutes and 30 minutes before tests,
35 respectively. Test compounds and MK-801 (maleate) were

administered 60 minutes and 30 minutes before tests. In experiments, a chamber with a floor grid for electric shock and a freezing measuring device (O'Hara & Co., Ltd.) were used. The chamber was put in a soundproof box to block the noise
5 from outside, and tests were performed. In the training session on day 1 of the experiment, after habituation for 3 minutes in the chamber, two electric footshocks were given to mice at 1-minute interval. In the test session on day 2 of the experiment, mice were placed in the same chamber for 7 minutes,
10 and freezing was measured during that period. The freezing was calculated as a percentage of freezing rates during measuring period with use of an automatic analysis software manufactured by the O'Hara & Co., Ltd.. All data were shown as the mean + standard error (n = 10-11). A comparison between two groups
15 was tested by Student's t-test (*p $\leq$ 0.05, comparison with control group. #p $\leq$ 0.05, comparison with group treated with MK-801 alone.).

The compounds in Figure 1 correspond to the following compounds.

- 20 Compound A: Example 67-II
Compound B: Example 73
Compound C: Example 76
Compound D: Example 263
Compound E: Example 273
25 Compound F: Example 64

The improvement effect on MK-801-induced deficits was shown by orally administering each of test compounds (A-F) to mice 60 minutes before the tests.

[1109]

30 Formulation Example 1

(1) Compound of the Example 1	10.0 g
(2) Lactose	70.0 g
(3) Cornstarch	50.0 g
(4) Soluble starch	7.0 g
35 (5) Magnesium stearate	3.0 g

After 10.0 g of the compound in Embodiment 1 and 3.0 g of magnesium stearate are granulated in 70 ml aqueous solution of soluble starch (7.0 g as soluble starch) and then dried, and the resulting mixture is mixed with 70.0 g of lactose and 50.0
5 g of cornstarch (lactose, cornstarch, soluble starch and magnesium stearate are all products in compliance with Japanese Pharmacopoeia 14th Edition). The mixture is compressed to obtain a tablet.

Industrial Applicability

10 [1110]


According to the present invention, the compound having a PDE2A selective inhibitory action, which is useful as an agent for the prophylaxis or treatment of schizophrenia, Alzheimer's disease and the like, can be provided.

15 [1111]

This application is based on patent application Nos. 100374/2012 and 283470/2012 filed in Japan, the contents of which are hereby incorporated by reference.

CLAIMS

1. A compound represented by the formula (1):



5 wherein

R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group,

R² and R³ are each independently a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, or R² and R³ in combination optionally form, together with the adjacent carbon atom, an optionally substituted ring,

X is a bond or an optionally substituted C₁₋₆ alkylene group,

A is an optionally substituted cyclic group,

Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally

substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C₁₋₆ alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom, and

5 B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms, provided that

2,3-dihydro-1-methyl-2-oxo-N-2-thiazolyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

10 4(1H)carboxamide,

2,3-dihydro-1-methyl-N-(4-methyl-2-thiazolyl)-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-(tetrahydro-2-oxo-3-furanyl)-6-

15 [3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

2,3-dihydro-1-methyl-N-2-oxazolyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

20 2,3-dihydro-1-methyl-2-oxo-N-1H-1,2,4-triazol-3-yl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-4-pyridinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

25 4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-4-pyrimidinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-5-pyrimidinyl-6-[3-

30 (trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

2,3-dihydro-1-methyl-2-oxo-N-3-pyridazinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-

4(1H)carboxamide,

35 2,3-dihydro-1-methyl-N-[6-(4-morpholinyl)-2-pyridinyl]-2-oxo-

6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
N-(4,5-dimethyl-2-thiazolyl)-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
5 4(1H)carboxamide,
2,3-dihydro-1-methyl-N-(5-methyl-2-thiazolyl)-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
N-[5-[(dimethylamino)carbonyl]-4-methyl-thiazolyl]-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
10 N-(4,6-dimethyl-2-pyridinyl)-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
2,3-dihydro-1-methyl-2-oxo-N-2-pyrazinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
15 2,3-dihydro-1-methyl-2-oxo-N-2-pyrimidinyl-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
N-[2-(1-azetidiny)-4-pyridinyl]-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide,
2,3-dihydro-1-methyl-N-[6-(4-morpholinyl methyl)-2-pyridinyl]-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide, and
25 N-[6-(1-azetidiny)-2-pyridinyl]-2,3-dihydro-1-methyl-2-oxo-6-[3-(trifluoromethyl)phenyl]-pyrido[2,3-b]pyrazine-4(1H)carboxamide are excluded,
30 or a salt thereof.

2. The compound or salt of claim 1, wherein
R²¹ is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an

- optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group or a cyano group, and Z³ is CR^{Z³} wherein R^{Z³} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group.
- 10 3. The compound or salt of claim 1, wherein R¹ is a hydrogen atom.
4. The compound or salt of claim 1, wherein R² and R³ are both hydrogen atoms.
- 15 5. The compound or salt of claim 1, wherein Z¹, Z² and Z³ are respectively CR^{Z¹}, CR^{Z²} and CR^{Z³}.
6. The compound or salt of claim 5, wherein R^{Z¹} and R^{Z³} are both hydrogen atoms.
- 20 7. The compound or salt of claim 5, wherein R^{Z²} is an optionally substituted C₁₋₆ alkoxy group, an optionally substituted C₁₋₆ alkyl group or an optionally substituted C₃₋₈ cycloalkyl group.
8. The compound or salt of claim 1, wherein X is an optionally substituted C₁₋₆ alkylene group.
- 30 9. The compound or salt of claim 1, wherein A is an optionally substituted phenyl group.
10. The compound or salt of claim 1, wherein R¹ is a hydrogen atom,
- 35 R² and R³ are both hydrogen atoms,

X is an optionally substituted C₁₋₆ alkylene group,

A is an optionally substituted phenyl group,

Z¹ is CH,

Z² is a group represented by CR^{Z2} wherein R^{Z2} is an optionally
5 substituted C₁₋₆ alkyl group, an optionally substituted C₁₋₆
alkoxy group or an optionally substituted C₃₋₈ cycloalkyl group,
and

Z³ is CH.

10 11. 7-Methoxy-N-((1S)-2-methoxy-1-(4-
(trifluoromethoxy)phenyl)ethyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide or a salt thereof.

12. N-((1S)-1-(3-Fluoro-4-(trifluoromethoxy)phenyl)-2-
15 methoxyethyl)-7-methoxy-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide or a salt thereof.

13. 7-Cyclopropyl-N-((1S)-2-hydroxy-2-methyl-1-(4-
(trifluoromethoxy)phenyl)propyl)-2-oxo-2,3-dihydropyrido[2,3-
20 b]pyrazine-4(1H)-carboxamide or a salt thereof.

14. 7-Cyclopropyl-N-(1-(3-fluoro-4-(trifluoromethoxy)phenyl)-
2-hydroxy-2-methylpropyl)-2-oxo-2,3-dihydropyrido[2,3-
b]pyrazine-4(1H)-carboxamide or a salt thereof.

25

15. N-(1-(3-Fluoro-4-(trifluoromethoxy)phenyl)-2-hydroxy-2-
methylpropyl)-7-methyl-2-oxo-2,3-dihydropyrido[2,3-b]pyrazine-
4(1H)-carboxamide or a salt thereof.


30 16. A medicament comprising the compound or salt of claim 1.

17. The medicament of claim 16, which is a phosphodiesterase
2A inhibitor.

35 18. The medicament of claim 16, which is an agent for the

prophylaxis or treatment of schizophrenia.

19. A method of inhibiting phosphodiesterase 2A, which comprises administering an effective amount of a compound
5 represented by the formula (1):




wherein

- R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group,
- 10 R² and R³ are each independently a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, or R² and R³ in combination optionally form, together with the adjacent carbon atom, an optionally substituted ring,
- X is a bond or an optionally substituted C₁₋₆ alkylene group,
- 15 A is an optionally substituted cyclic group,
- Z¹ is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆
- 20 alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,
- Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an
- 25 optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,
- 30 Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom,

Z² is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z³ is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom, a halogen atom, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₂₋₆ alkenyl group, an optionally substituted C₂₋₆ alkynyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C₁₋₆ alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom, and
 15 B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms, or a salt thereof, to a mammal.

21. Use of a compound represented by the formula (1):



wherein


R¹ is a hydrogen atom or an optionally substituted C₁₋₆ alkyl group,
 R² and R³ are each independently a hydrogen atom or an optionally substituted C₁₋₆ alkyl group, or R² and R³ in combination optionally form, together with the adjacent carbon atom, an optionally substituted ring,
 X is a bond or an optionally substituted C₁₋₆ alkylene group,
 A is an optionally substituted cyclic group,

Z^1 is a group represented by CR^{Z^1} wherein R^{Z^1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z^2 is a group represented by CR^{Z^2} wherein R^{Z^2} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

Z^3 is a group represented by CR^{Z^3} wherein R^{Z^3} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C_{1-6} alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom, and B is a nitrogen-containing aromatic heterocycle containing 1 or 2 nitrogen atoms, or a salt thereof, for the production of an agent for the prophylaxis or treatment of schizophrenia.

22. A compound represented by the formula (1):



wherein

R^1 is a hydrogen atom or an optionally substituted C_{1-6} alkyl

group,

R^2 and R^3 are each independently a hydrogen atom or an optionally substituted C_{1-6} alkyl group, or R^2 and R^3 in combination optionally form, together with the adjacent carbon

5 atom, an optionally substituted ring,

X is a bond or an optionally substituted C_{1-6} alkylene group,

A is an optionally substituted cyclic group,

Z^1 is a group represented by CR^{Z1} wherein R^{Z1} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an

10 optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group or an optionally substituted cyclic group, or a nitrogen atom,

15 Z^2 is a group represented by CR^{Z2} wherein R^{Z2} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano

20 group or an optionally substituted cyclic group, or a nitrogen atom,

Z^3 is a group represented by CR^{Z3} wherein R^{Z3} is a hydrogen atom, a halogen atom, an optionally substituted C_{1-6} alkyl group, an optionally substituted C_{2-6} alkenyl group, an optionally

25 substituted C_{2-6} alkynyl group, an optionally substituted C_{1-6} alkoxy group, an optionally substituted amino group, a cyano group, an optionally substituted C_{1-6} alkyl-carbonyl group or an optionally substituted cyclic group, or a nitrogen atom, and

B is a nitrogen-containing aromatic heterocycle containing 1

30 or 2 nitrogen atoms,

or a salt thereof, for use in the prophylaxis or treatment of schizophrenia.