USE OF AT-1 RECEPTOR ANTAGONIST OR AT-2 RECEPTOR MODULATOR FOR TREATING DISEASES ASSOCIATED WITH AN INCREASE OF AT-1 OR AT-2 RECEPTORS

[0001] Angiotensinogen, an α2-macroglycoprotein, is cleaved by the enzyme renin into the decapeptide angiotensin I, which is itself only very slightly active biologically. In the next step of the cascade, two further amino acids are cleaved off by the action of the enzyme angiotensin converting enzyme (ACE), which is mainly bound in the endothelium, with the formation of angiotensin II. The latter is regarded as being one of the most powerful natural vasoconstrictors.

[0002] Angiotensin II interacts with specific receptors on the surface of the target cell. Success has by now been achieved in identifying receptor subtypes which are, for example, designated AT_1 receptors and AT_2 receptors. Studies on the renin-angiotensin system, particularly in relation to hypertension, have increased almost exponentially over the last decade. As a result, the number of receptors for Angiotensin II have now been identified and some of them have been cloned and analysed. Recently, considerable efforts have been made to identify the substances which bind to the AT_1 receptor, with active compounds of this nature frequently being termed angiotensin II antagonists. As a consequence of the inhibition of the AT_1 receptor, these antagonists can, for example, be employed as antihypertensives or for treating congestive heart failure.

[0003] The AT_1 and AT_2 receptors have also been studied for their distribution and biological properties and have been shown, despite a 30% homology, to have a very different distribution and activity.

[0004] The AT_1 receptor, which plays a major part in blood pressure regulation, has been found in the adrenal cortex, kidney, uterus etc. At a cellular level it has been found on fibroblasts, macrophages and smooth muscle cells (SMC).

[0005] In contrast, the AT_2 receptor has been found mainly in foetal tissues but also in adult especially in pathological tissue such as in ischaemic heart disease. Here it has been located on fibroblasts and endothelial cells.

[0006] The aim of the studies described hereafter is to evaluate the distribution of AT_1 and AT_2 receptors in the human lung using essentially the immunocytochemical and the in situ hybridisation methodologies.

[0007] Previously, specific antibodies have been made against epitopes of the AT_1 receptor but no such specific tools existed for the AT_2 receptor. Histological studies therefore depended on radio-labelled receptor antagonists together with autoradiography which only gave a relatively crude tissue localisation. In the last two years, however, specific well-characterised antibodies to the human receptor have become available and have therefore been used in the studies relied upon hereafter.

Methods and Materials

1. Antibody and in situ hybridisation (ISH) probe specificity and titration

[0008] In order to confirm the specificity of the immunocytochemical (ICC) and ISH studies, paraffin embedded blocks of normal human adrenal (cortex and medulla) are obtained from the archives of the Pathology Dept., University Hospital, Ghent, Belgium, and used as test material. AT_1 receptors are known to be predominantly located in the adrenal cortex and AT_2 in the medulla.

[0009] For the human lung studies, control material is obtained from autopsies where the patients have died from causes other than lung disease e.g. fatal accidents. Some of this material comes from Ghent as above, some from the archives of the Pathology Dept. of The Pennsylvania Hospital, Philadelphia, USA. Tissue containing small airways is selected as they appear to be more sensitive to injury.

[0010] All samples have been fixed in 10% buffered formalin as rapidly as possible post-mortem, dehydrated and embedded in paraffin wax. 3-5 μm sections are cut and mounted on silane coated glass slides.

[0011] In order to minimise any variations in the processsing, pairs of sequential sections are mounted on each slide, one for AT_1 and the other for AT_2 antibody exposure. Up to 20 slides are treated at the same time so that, with the exception of the primary antibody, al reagents including the chromogen are identical. Section thickness is therefore the only variable which could not be totally controlled.

2. Antibodies

[0012] Two AT_1 receptor antibodies, from Santa Cruz Inc., San Diego, CA, USA, (clones N 10 and 306) are tested and found to give identical adrenal cortex receptor distribution.

[0013] The major part of the study is done with an AT_2 receptor antibody available from Santa Cruz (clone C18) which is tested and is found that it gives the identical staining pattern in adrenal medulla and lung as the first one.

[0014] All the antibodies have been rigorously tested by the commercial and individual suppliers for specificity and cross-reactivity.

3. ISH probes

[0015] PCR (Polymerase Chain Reaction) products were prepared and used as follows:

Oligonucleotides specific to human angiotensin II receptor type I (GenBank accession number M93394) and angiotensin II receptor type II (GenBank accession number U15592) were designed using the Oligo 5.0 software programme to homologous regions from both sequences (Table 1). cDNA from human bone tissue was prepared following standard methods (Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning: a Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989). For PCR the following oligonucleotide primer pairs were used: angiotensin II receptor type I, 5'-CTggCTgACTTATgCTTTTTACTgACT-3' and 5'-gATgCAggTgACTTTggCTACA-3'; (PCR product size 236 base pairs) and for angiotensin II receptor type II, 5'-ATTTACTCCTTTTggCTACTCTTCCTC-3' and 5'-ggTCACgggTTATCCTgTTCTTC-3' (PCR product size 489 base pairs). PCR amplifications were performed with 10 ng of template cDNA using a MJ Research PCR Cycle machine and the following PCR cycles : 1) 94 °C/2 min, 2) 94 °C/10 sec, 60 °C/30 sec, 72 °C/15 sec for 35 cycles, using High Fidelity Taq polymerase (Boehringer Mannheim) with components provided in the manufacturer's kit. Products of the PCR amplifications were identified by electrophoresis through a 0.8% agarose/TBE gel (Sambrook J, Fritsch EF, Maniatis T. Molecular Cloning: a Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989). To confirm the identity of the PCR amplification products, the DNA was eluted from the gel and cloned into the A/T cloning vector pMOSBlue (Amersham). Colonies containing a DNA insert of the correct size (Table 1) were fully sequenced on both strands to confirm their identity.

[0016] Each probe, both sense and anti-sense for the AT_1 and anti-sense only as described above, are labelled with fluorescin (FITC) and the presence of mRNA in the cells detected following hybridisation with the probe and use of a mouse anti-FITC probe plus the alkaline phosphatase anti-alkaline phosphatase (APAAP) detection system. This labelling technique enhanced the detection of very low copy numbers.

4. Immunocytochemistry

[0017] For all the antibodies the procedure is as follows. 5 μm sections are first treated by antigen retrieval techniques, together with microwaving, in citrate buffer (pH 6.0).

[0018] Exposure is for 20 minutes and the slides left to cool down in the buffer. Where the antibodies are goat polyclonal, the peroxidase anti-peroxidase (PAP) method with diaminobenzidine (DAB) as chromogen is utilised, otherwise for rabbit polyclonals the APAAP system with new Fuchsin is used. The sections are first blocked with 1% bovine serum albumen (BSA) for 30 minutes to block non-specific receptors, followed by incubation with the primary antibody for 30 minutes at room temperature. Each antibody is titrated out and the optimal dilutions are as follows:

Antibodies Used		Lung Tissue	Adrenal
AT_1	Clone N 10	1:200	1:500
	Clone N 306	1:200	1:500
AT_2	Clone C 18	1:150	1:500

[0019] As a negative control, for the rabbit polyclonal a negative serum from Dako (Prosan, Ghent, Belgium) is used, for the goat polyclonal the primary antibody is omitted.

ISH

[0020] 5 μm sections are deparaffinised and then exposed to 'in situ' hybridisation following well establised techniques. The sections are first treated with pre-hybridisation solution for 20 minutes at 55°C. They are then washed before exposure to the probes overnight at 55°C. After further washing, they are treated with a mouse anti-FITC antibody and the APAAP detection system for location of the specific message. For the AT_1 receptor, the sense probe is the negative control, for the AT_2 receptor, the probe is omitted.

Image analysis

[0021] In order to measure the stain intensity quantitatively, the slides are viewed in the Leica MR500 and the amount of stain per unit area of tissue recorded in pixels following Leica's instruction. This is done to establish the ratios of AT_1

to AT_2 receptor in different regions. These are a) airways epithelium, b) the sub-epithelial interstium, c) SMC around blood vessels and d) mucous glands.

RESULTS

Adrenal distribution studies

[0022] AT_1 Distribution: All two antibodies give the following distribution pattern in the adrenal cortex, that is distinctive staining around the smooth muscle cells surrounding the blood vessels and also on the interstitial network on and around fibroblasts as predicted. There is no staining of endothelial cells.

[0023] AT_2 Distribution: The antibody is tested against adrenal medulla where strong staining of adrenal pheochromocytoma cells is seen.

[0024] ISH: Both antisense probes give a similar picture.

Control Lung

[0025] AT_1 Distribution : Very clear staining of the interstitial cells underlying the airways epithelium (sub-epithelial) is seen and also the margins of the smooth muscle cell (SMC) surrounding the blood vessels. In addition macrophages are also positive.

[0026] AT_2 Distribution: This receptor appears to be strongly associated with the airway epithelial cells, with dense staining of the brush border. Positive cells are also seen in some of the mucous glands, on some vascular endothelial cells and on fibroblasts, chondrocytes and macrophages. There is no staining of SMC.

[0027] ISH: Again the probes give a similar picture. In particular the AT_2 probe give a strong signal on endothelial cells and on some mucous glands only.

Image analysis

[0028] The distribution of the protein and therefore the receptor as represented by stain intensity i.e. pixels per μm^2 of tissue is given in the following table.

Antibody	Airway Epithelial	Sub-Epithelial	Glands	SMC
AT_1	0.00	8	0.00	10
AT_2	5	0-1	5	0.00

[0029] The presence of Angiotensin II receptors in adrenal cortex and medulla has previously been demonstrated by both biochemical and histological means. The data are obtained with both commercially available and privately supplied antibodies both confirming these findings but also establishing the reliability of the instant immunocytochemical and ISH methodology.

[0030] In view of the results of these studies, the distribution of the AT_1 and AT_2 receptors in normal and diseased lung tissues has to be compared, in order to determine the specific AT_1 and AT_2 distributions and ratios.

[0031] The presence of AT_1 receptors in the lung has previously been shown biochemically and now their exact cellular location has been demonstrated. This information is vital for establishing the proportions of AT_1 and AT_2 receptors in different regions of the lung under normal and pathological conditions.

[0032] The data on the distribution especially of the AT_2 receptor is totally new as no previous data exist as to their presence or distribution. Several important points arise from this study.

[0033] First the presence of the AT_2 receptor on the bronchial epithelial cells of the small airways. As it is already known this receptor is considered to be anti-fibrotic, anti-proliferative and pro-apoptotic. Therefore up- or down-regulation on epithelial cells has profound effects on epithelial cell replacement, development of hyperplasia and even a role in the development of lung cancer.

[0034] Secondly, the presence of considerable amounts of the protein on the brush border of the epithelial cells could well be associated with the amount of mucous secretion as some of the epithelial cells of the mucous glands have also been shown to carry the receptor. From the ISH data, some glands contain a high level of mRNA.

[0035] Finally, the presence of the AT_2 receptor on vascular endothelial cells has now been confirmed both by immunocytochemistry and in situ hybridisation. Its distribution to be both infrequent and not on all the cells of one particular vessel hasbeen found.

[0036] These studies confirm and extend existing data on the presence and distribution of Angiotensin II type AT_1

and AT_2 in the human lung. The presence of the AT_2 receptor on the small airway epithelial cells has considerable consequences for the understanding of diseases arising from alteration in the function of these cells.

[0037] Very little is known on the localisation of the AT receptors in the human lung and about their distribution, up or down regulation and ratio in normal and diseased lung tissue. For this study, samples are collected from normal lung, and from patients with chronic obstructive pulmonary disease (COPD) ± hypertension.

Lung samples

[0038] These have been obtained from the following groups of patients all clinically defined.

NN	(n=3)	-	non-smokers, normal tissue
C	(n=5)	-	smokers but otherwise normal
COPD	(n=8)	-	COPD positive, normal BP
COPD/H	(n=3)	-	COPD plus hypertension
H	(n=4)	-	smokers with raised BP

[0039] The airways are carefully dissected out of the lungs immediately after removal of the lung from the patient for tumour resection or other reasons. The region is carefully selected to be free of any cancerous tissue. The small blocks are then fixed in 4% paraformaldehyde for 2 h at room temperature before embedding in paraffin wax. This is to ensure optimum structural integrity and retention of antigenic activity.

Methods for establishing receptor localisation

[0040] a) Immunocytochemistry (ICC). 2 x AT_1 antibodies are tested, Santa Cruz (clones N-10 and 306). One AT_2 antibody has also been tested, Santa Cruz (clone C 18).

Digital images show:

[0041] Distribution of the AT_2 receptor in bronchiolar epithelial cells including the brush border and on the mucous gland cells.

[0042] Adjacent section stained for AT_1 receptor illustrating the very different distribution on smooth muscle cells, fibroblasts/stroma and macrophages.

Analysis of human lung from patients

[0043] All material obtained to date has been sectioned and stained by ICC for both AT_1 and AT_2 receptor localisation with appropriate negative controls. Image analysis has been started with readings from one section per patient to date (this is a slow process as base lines have to rigidly adhered to). Measurements have been made of the epithelial v subepithelial and blood vessel. This analysis is being done in a "blinded" manner so that no comment can be made other than the fact that some "patients" clearly have levels well away from the average. Discrepancies due to varying section thickness and staining have been minimised by doing the AT_1 and AT_2 ICC simultaneously with sequential sections. Thus the same batch of chromogen could also be used.

[0044] The findings of a lung epithelial localisation for the AT_2 receptor has a number of consequences. As this receptor has been shown to be both anti-proliferative, anti-fibrotic and pro-apoptotic its up-regulation is anticipated to have consequences for a number of lung diseases; e.g.that smoking alone has any influence. It is a role in such fibrotic conditions as adult respiratory distress syndrome (ARDS), or even in reducing the proliferative capacity of the epithelium in lung cancer.

Results

Receptor localisation

[0045] All the antibodies and riboprobes are tested on normal adrenal cortex and medulla where both receptor types are known to be present in relative abundance.

[0046] The process is repeated on normal lung tissue where we are able to detect both AT_1 and AT_2 receptors and their mRNA. The localisation is as follows:

AT_1 - on smooth muscle cells, fibroblasts/stroma, macrophages. This is fairly predictable except the intensity and number of receptors in normal lung is quite high.
AT_2 - on bronchial epithelial cells (especially the brush border), on mucous glands (some). In addition, on the vascular endothelial cells, fibroblasts, macrophages and cartilage cells.

[0047] This is a totally unexpected and novel finding which needs investigating further. The epithelial cell and mucous gland location is confirmed by both protein and mRNA content, the brush border location relates to mucous secretion.

DISTRIBUTION OF ANGIOTENSIN AT_1 AND AT_2 IN THE LUNGS OF PATIENTS WITH CHRONIC BRONCHITIS COMPARED WITH CONTROLS

[0048]

Group	Epithelial		Sub-epithelial		Ratio
	AT_1	AT_2	AT_1	AT_2	AT_1(Epithelia)/AT_2(Sub-epithelial)
Control (1) Smokers - N=4	0.02	7.15	4.07	0.01	0.56
Control (2) Non-Smokers N=5	0.02	9.49	6.45	1.3	0.67
Control (3) Smokers Hypertension	0.1	7.97	11.02	0.04	1.38
COPD Smokers No hypertension N=7	0.10	6.84	19.64	0.11	2.87
COPD Smokers Hypertension N=3	0.30	6.01	6.12	0.05	1.01

[0049] The Leica equipment (image analyser MR 500) translates the stain intensity into a grey scale (0-250), each unit being one pixel. This allows one to quantitate the data accurately.

[0050] These data are based on image analysis of the patient. Data is expressed as pixels per unit area of positively stained tissue and the mean of five fields per slide.

[0051] As can be taken from these results, the ratio of AT_1 distribution in the sub-epithelial and AT_2 in epithelial in normal lung tissue is significantly below 1, whereas this ratio is close to or above 1 in diseased lung tissues. The epithelial forms the inner lining of the trachea and the main bronchii. The increase in the AT_1/AT_2 receptor ratio in the bronchial sub-epithelial region of the lung from chronic bronchitic patients compared with control is mainly due to raised levels of the AT_1 receptor found on the fibroplasts and macrophages surronding the airways epithelium and reflects the increased levels of inflammation and fibrosis seen in COPD.

[0052] The above results clearly demonstrate that AT_1 receptors which modulate angiotensin II are located in sub-epithelial lung tissue and especially the distribution in corresponding lung tissue is increased. Accordingly, the inhibition of angiotensin II by means of AT_1 receptor antagonists leads to decrease in airways obstruction.

[0053] Furthermore, the experiments show that the distribution of AT_2 in epithelial lung tissue, especially in corresponding diseased tissue, e.g. mainly on the bronchial epithelial cells, and also in structural cells of the alveola, e.g. on mucous glands of the alveola, is increased. As AT_2 receptors are anti-proliferative, anti-fibrotic and pro-apoptotic, their modulation is useful for the treatment of specific forms of lung conditions and diseases, especially for the treatment of adults respiratory distress syndrome (ARDS) and for reducing the proliferative capacity of the epithelium in lung and breast cancer, furthermore, for the treatment of sepsis syndrome, lung injury forms, such as pneumonia, aspiration of gastric content, chest trauma, shock, burns, fat embolia, cardiopulmonary bypass, O_2 toxicity, haemorhagic pancreatitis, interstitial and bronchoalveolar inflammation, proliferation of epithelial and interstitial cells, collagen accumulation, fibrosis.

Receptor distribution in normal breast tissue and in patients with breast cancer

[0054] The breast specimens included in this study are randomly recruited from the files of the Pathology Dept, University Hospital, Ghent. All have been fixed in formalin, processed into paraffin wax and a diagnosis of their pathological status made by the departmental pathologists. Sixteen cases have been included: 14 invasive ductal carcinomas, one invasive colloid carcinoma and one invasive lobular carcinoma.

[0055] Immunocytochemistry is carried out on paraffin wax embedded tissue sections using the polyclonal antibodies for AT1 and AT2 used in the lung study together with the streptavidin-biotin-peroxidase complex method as before.

[0056] In order to provide a possible model for testing receptor antagonists, cell lines originating from human breast tissues are also studied for their receptor content. This can provide a useful in vitro working model for further biochemical and cytological studies.

Results

[0057] The data obtained clearly demonstrate the presence of angiotensin II type 1 and 2 receptors in normal human breast tissue with AT2 being found on the cuboidal epithelium lining the ducts and AT1 predominantly present on the ductal myoepithelial cells. All staining was abolished by omitting the primary antibody from the incubation.

[0058] In all cases the connective tissue was positive for the AT1 receptor along with no (11/16) to weak (5/16) staining of the cancer cells. For the AT2 receptor, the opposite was observed with all carcinomas being positive and almost no stromal reactivity (1/16).

[0059] Results from the cell lines tested were very interesting with a different staining pattern being seen in each one. The short term cultures of normal mammary epithelial cells were strongly positive for the AT1 receptor but stained only weakly for the AT2 receptor.

Breast carcinomas

[0060]

Patient	AT_1			AT_2	
	Breast Carcinoma	Carcinoma	Stroma	Carcinoma	Stroma
1	invasive, interm. differentiated ductal carcinoma - grade 2 Bloom-Richardson classification	-	++	++	-
2	invasive, interm. differentiated ductal adenocarcinoma	-	+++	++ focal	-
3	invasive, poorly differentiated ductal carcinoma - ductal carcinoma *in situ*	-	++	+	-
4	invasive, interm. differentiated ductal carcinoma - extensive ductal carcinoma *in situ*	-	+	++	-
5	invasive colloid carcinoma	-	++	++	-
6	invasive, interm. differentiated ductal carcinoma - ductal carcinoma *in situ*	+	+	+++	-
7	invasive, well differentiated ductal carcinoma - large ductal carcinoma *in situ*	-	++	++	-
8	invasive, poorly differentiated ductal carcinoma - ductal carcinoma *in situ*	+	++	+++	-
9	invasive, well differentiated ductal carcinoma	-	+	++ focal	-
10	invasive, poorly differentiated ductal carcinoma - ductal carcinoma *in situ*	-	+	+	-
11	invasive, poorly differentiated ductal carcinoma - multifocal carcinomas *in situ*	-	++	++	-
12	invasive, poorly differentiated ductal carcinoma	-	+++	++	+
13	invasive, poorly differentiated invasive carcinoma	+	++	++	-

(continued)

Patient	AT_1			AT_2	
	Breast Carcinoma	Carcinoma	Stroma	Carcinoma	Stroma
14	invasive, poorly differentiated ducatl carcinoma - several ducal carcinomas *in situ*	-	++focal	++	-
15	invasive, poorly differentiated ductal carcinoma - several ductal carcinomas *in situ*	+	+++	+	-
16	invasive, lobular carcinoma - cribriform intraductal carcinoma	+	+	+	-

Conclusions

[0061] As can be seen from these results, the presence of the AT1 receptor on normal epithelial cells appears to be very different to the lung. However, the epithelial cell type is myoepithelial whereas the AT2 receptor in both tissues is found on cuboidal epithelial cells. At the same time, the positive stromal staining for the AT1 receptor is the same for both tissues. The latter is clearly linked to the presence of fibroblasts in the extracellular matrix.

[0062] The presence of the AT2 receptor on the carcinoma cells and in such a widespread and reproducible manner is surprising. The cell types described here carrying the specific receptors provide an ideal *in vitro* model for such studies.

[0063] These experiments clearly demonstrate the surprising effect that in the instant model using breast carcinoma cells, the AT_1 receptors are mainly distributed in the stroma while the AT_2 receptors could mainly be ascertained in the carcinoma cells.

[0064] All these surprising results clearly demonstrate that any AT_1 receptor antagonist or AT_2 receptor modulator may be used for the treatment of conditions or diseases associated with the increase of AT_1 receptors in the sub-epithelial area or increase of AT_2 receptors in the epithelia, especially for the treatment of obstructive airways diseases. Obstructive airways diseases a classification of respiratory diseases which are characterized by decreased airway size and increased airway secretion, resulting in reduced alveolar ventilation. Obstructive airways diseases comprise reversible and irreversible conditions and are selected, for example, from chronic obstructive pulmonary disease, such as bronchitis, e.g. chronic bronchitis and emphysema, likewise from asthma, cystic fibrosis, interstitial lung disease, invasive lung cancer, pulmonary vascular disease, and increased resistance to airflow during forced expiration. Any such treatment may also, but not necessarily, be associated with the treatment of hypertension as well as both non-smokers and smokers.


[0065] These surprising results clearly demonstrate that any AT_2 receptor modulator may be used for the treatment of conditions or diseases associated with an increase of AT_2 receptors in epithelial lung tissue, especially for the treatment of specific forms of lung conditions and diseases, especially for the treatment of adults respiratory distress syndrome (ARDS) and for reducing the proliferative capacity of the epithelium in invasive lung cancer, , furthermore, for the treatment of sepsis syndrome, lung injury forms, such as pneumonia, aspiration of gastric content, chest trauma, shock, burns, fat embolia, cardiopulmonary bypass, O_2 toxicity, haemorhagic pancreatitis, interstitial and bronchoalveolar inflammation, proliferation of epithelial and interstitial cells, collagen accumulation, fibrosis.

[0066] AT_1 receptor antagonists or AT_2 receptor modulator are agents that modify the host's biological response to tumor cells with resulting therapeutic benefit. The increased AT_1 receptor expression in mammary ductal myoepithelium and of the AT_2 receptor in mammary cuboidal epithelium demonstrate that any AT_1 receptor antagonist or AT_2 receptor modulator may be used for treatment of invasive breast carcinoma. These included scirrhous, infiltrative, papillary, ductal, medullary and lobular breast cancers as well as metastasis in the lungs, pleura, skeleton and liver. Treatment should be considered as adjuvant therapy in combination with surgery, radiotherapy or as palliative therapy with hormonal therapy or other biological response modifiers such as interferons, interleukins, tumor necrosis factors, monoclonal antibodies etc..

[0067] While clinical examination and mammography suggest breast cancer, it is only the examination of the tissue biopsy which allow to make the diagnosis. The distribution pattern of AT_1 and AT_2 receptors can be used as marker for hyperplasia (location of AT_1 receptors) and for invasive cancer (location of AT_2 receptors) and therefore for the diagnostic of the malignancy of the tumor.

AT_1 receptor antagonists include compounds having differing structural features. For example, mention may be made of the compounds which are listed in the European Patent Application having the publication No. 443983 (EP 443983), in particular in the compound claims and the final products of the working examples, the subject-matter of which claims is hereby incorporated into the present application by reference to this publication.

[0068] Preference is given to (S)-N-(1-carboxy-2-methylprop-1-yl)-N-pentanoyl-N-[2'(1 H-tetrazol-5-yl)biphenyl-4-yl-methyl]amine [Valsartan] of the formula




(I)

and its pharmaceutically utilizable salts.

[0069] Furthermore, the compounds which are listed in European Patent Application having the publication No. 253310 (EP 253310), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0070] Preference is given to the compound [Losartan] of the following formula



and its pharmaceutically utilizable salts.

[0071] Furthermore, the compounds listed in the European Patent Application having the publication No. 403159 (EP 403159), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0072] Preference is given to the compound [Eprosartan] of the following formula



and its pharmaceutically utilizable salts.

[0073] Furhermore, the compounds listed in the PCT Patent Application having the publication No. WO 91/14679, in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0074] Preference is given to the compound [Irbesartan] of the following formula



and its pharmaceutically utilizable salts.

[0075] Furthermore, the compounds listed in the European Patent Application having the publication No. EP 420237 (EP 420237), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0076] Preference is given to the compound [E-1477] of the following formula



and its pharmaceutically utilizable salts.

[0077] Furthermore, the compounds listed in the European Patent Application having the publication No. 502314 (EP 502314), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0078] Preference is given to the compound [Telmisartan] of the following formula



and its pharmaceutically utilizable salts.

[0079] Furthermore, the compounds listed in the European Patent Application having the publication No. 459136 (EP 459136), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0080] Preference is given to the compound [Candesartan] of the following formula



and its pharmaceutically utilizable salts.

[0081] Furthermore, the compounds listed in European Patent Application having the publication No. 504888 (EP 504888), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0082] Preference is given to the compound [SC-52458] of the following formula



and its pharmaceutically utilizable salts.

[0083] Furthermore, the compounds listed in the European Patent Application having the publication No. 514198 (EP 514198), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0084] Preference is given to the compound [Saprisartan] of the following formula



and its pharmaceutically utilizable salts.

[0085] Furthermore, the compounds listed in the European Patent Application having the publication No. 475206 (EP 475206), in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.


[0086] Preference is given to the compound of the following formula



and its pharmaceutically utilizable salts.

[0087] Furthermore, the compounds listed in the PCT Patent Application having the publication No. WO 93/20816, in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.

[0088] Preference is given to the compound [ZD-8731] of the following formula



and its pharmaceutically utilizable salts.

[0089] AT_2 receptor ligands (modulators) include compounds having differing structural features. For example, mention may be made of the compounds which are listed in WO 94/13651, in particular in the compound claims and the final products of the working examples, the subject-matter of which claims is hereby incorporated into the present application by reference to this publication.

[0090] Furthermore, the compounds listed in WO 94/13642, in particular in the compound claims and the final products of the working examples, are hereby incorporated into the present application by reference to this publication.

[0091] AT_1 receptor antagonists or AT_2 receptor ligands, respectively, which, for example, possess at least one basic centre can form acid addition salts. These are formed, for example, using strong inorganic acids, such as mineral acids, e.g. sulfuric acid, a phosphoric acid or a hydrohalic acid, using strong organic carboxylic acids, such as C_1-C_4alkanecarboxylic acids which are unsubstituted or substituted, for example, by halogen, e.g. acetic acid, such as saturated or unsaturated dicarboxylic acids, e.g. oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, e.g. ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, e.g. aspartic or glutamic acid, or such as benzoic acid, or using organic sulfonic acids, such as C_1-C_4alkanesulfonic acids or arylsulfonic acids which are unsubstituted or substituted, for example, by halogen, e.g. methanesulfonic acid or p-toluenesulfonic acid. Examples of suitable salts with bases are metal salts, such as alkali metal or alkaline earth metal

salts, e.g. sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkyl amine, e.g. ethyl-, tert-butyl-, diethyl-, diisopropyl-, triethyl-, tributyl- or dimethylpropyl-amines, or a mono-, di- or tri-hydroxy lower alkyl amine, e.g. mono-, di- or tri-ethanolamine. Furthermore, corresponding internal salts can be formed.

[0092] WO 00/02543 A2 describes a method of treatment and a pharmaceutical composition for treating various diseases, wherein a combination of valsartan or a pharmaceutically acceptable salt thereof and a calcium channel blocker or a pharmaceutically acceptable salt thereof is used as a combination of active agents.

[0093] WO 00/44378 A1 relates to the use of an angiotensin II receptor antagonist or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for the treatment of acute MI and for the secondary prevention of acute MI.

[0094] WO 97/49394 A2 is concerned with solid oral dosage forms comprising a) an active agent selected from valsartan and optionally hydrochlorothiazide, and b) pharmaceutically acceptable additives suitable for the preparation of solid oral dosage forms by compression methods.

[0095] US 3 146 168 relates to the manufacture of pharmaceutical preparations having cellulose crystallite aggregates or certain derivatives of cellulose crystallite aggregates present therein.

[0096] WO 01/97805 A2 relates to solid oral dosage forms comprising pharmacologically effective amounts of valsartan or a pharmaceutically acceptable salt thereof or hydrate thereof, wherein the compositions are at least 1.2 times more bioavailable than a conventional valsartan capsule.

[0097] The present invention is featured in claim 1. Preferred embodiments of the invention are set out in the subclaims and in this description.

[0098] The tablets of the present invention, such as coated tablets, tablets, are prepared in a manner which is known per se, for example using conventional mixing, granulation, coating, solubulizing or lyophilizing processes. Thus, the tablets can be obtained by combining the active compound with solid excipients, if desired granulating a mixture which has been obtained, and processing the mixture or granulate into tablets or coated tablet cores after having added suitable auxiliary substances.

[0099] The dosage of the active compound can depend on a variety of factors, such as mode of administration, homeothermic species, age and/or individual condition. Normally, in the case of oral administration, an approximate daily dose of from about 10 mg to about 360 mg, for example in the case of Valsartan e.g. of about 40 mg, 80 mg, 160 mg or 320 mg, is to be estimated for a patient of approximately 75 kg in weight.

[0100] The tablets of the present invention may be used for the treatment of diseases and conditions e.g. as disclosed hereinbefore.

[0101] WO 97/49394 discloses compressed solid oral dosage forms, e.g., by compaction, of valsartan (optionally in salt form) optionally combined with hydrochlorothiazide (HCTZ). In WO 97/49394 the preferred range of cellulose is given as 10 to 30%, e.g., 21 %, for valsartan/HCTZ compositions and 5% Valsartan alone. The preferred range of crosslinked polyvinylpyrolidone (Crospovidone) is given as 10 to 20%, e.g., 13 %.

[0102] After exhaustive testing it has been found surprisingly that it is possible to improve the bioavailability characteristics of known solid formulations of valsartan by increasing the proportion of microcristalline cellulose. It has also been found surprisingly that it is possible to improve the quality, e.g., better weight uniformity and better compression for the tablets, of said known solid formulations of Valsartan by decreasing the proportion of crosslinked PVP crospovidone.

[0103] Thus, the present invention relates to a tablet comprising valsartan as the active agent and more than 30% of microcristalline cellulose by weight based on the total weight of the core components of said solid oral dosage form, e.g., 31 to 65%, e.g., 50%.

[0104] In the present invention, the weight ratio of valsartan to microcristalline cellulose is preferably from 2.5 : 1 to 0.3 : 1, e.g., 2 : 1 to 1 : 1, e.g., 1.4: 1.

[0105] In a further embodiments the tablet of the invention comprises less than 13% of crospovidone, e.g., 2 to 10%, by weight based on the total weight of the core components of the tablet.

[0106] Preferably, the weight ratio of valsartan to Crospovidone is from 7 : 1 to 3 : 1, e.g., 6 : 1 to 4 : 1, e.g., 5.3 : 1.

[0107] Preferably, the weight ratio of microcristalline cellulose to crospovidone is from 7 : 1 to 1 : 1, e.g., 4 : 1 to 2 : 1, e.g., 3.6: 1.

[0108] The tablet according to the invention may comprise from 20 to 360 mg of valsartan, e.g., 40, 80, 160, 320mg. With this range of dosages treatment flexibility and efficacy, e.g., in blood pressure reduction, may be increased.

[0109] In a further aspect, the invention relates to a solid oral dosage form comprising

20 to 65% of valsartan
31 to 65% of microcristalline cellulose
2 to 13% of crospovidone.

[0110] A typical composition may comprise :

20 to 65% of valsartan
31 to 50% of microcristalline cellulose
2 to 10% of crospovidone
1 to 10 % of magnesium stearate
0.5 to 5 % of colloidal anhydrous silica.

[0111] If desired 1 to 10% by weight of the core composition, e.g., 5 to 10%, of Cutina, or 1 to 10% by weight of the core composition, e.g., 5 to 10% of stearic acid may be added.

[0112] Preferably tablets of the invention is in the form of a compressed tablet.

[0113] In a further embodiment the invention relates to a tablet, e.g., a compressed tablet, comprising more than 250 mg and up to 360 mg, e.g., 320 mg, of Valsartan as an active agent.

[0114] Other excipients as lubricants and glidants commonly used in solid oral formulations may be used and reference is made to the extensive literature on suitable substances, see in particular Fiedler's "Lexicon der Hilfstoffe", 4th Edition, ECV Aulendorf 1996 and "Handbook of Pharmaceutical Excipients" Wade and Weller Ed.(1994).

[0115] The tablets according to the present invention may be in the form of dragées in which case the tablet is provided with a coating typically a sugar, shellac or other film coating entirely conventional in the art. Attention is drawn to the numerous known methods of coating employed in the art, e.g. spray coating in a fluidized bed, e.g. by the known methods using apparatus available from Aeromatic, Glatt, Wurster or Hüttlin, in a perforated pan by the Accela Cota method, or to the submerged sword coating method. The additives commonly used in confectioning are employed in such methods. For example, coatings which may be used are those disclosed in WO 97/49394, Opadry and the like.

[0116] The tablets of the present invention are useful in the known indications of the particular active agent incorporated therein.

[0117] The exact dose of active agent and the particular formulation to be adminstered depend on a number of factors, e.g. the condition to be treated, the desired duration of the treatment and the rate of release of the active agent. For example, the amount of the active agent required and the release rate thereof may be determined on the basis of known in vitro or in vivo techniques, determining how long a particular active agent concentration in the blood plasma remains at an acceptable level for a therapeutic effect.

[0118] For example, the composition of the invention in clinical trials has a comparable bioavailability to the commercial form of Diovan®.

[0119] Preferably the dissolution rate of solid form according to the present invention is above 90% over 30 minutes.

[0120] For example dosages in the range of 10 mg to 360 mg of valsartan per day for a 755 kilogram mammal, e.g., humans, and in standard animal models, may be used. An excellent tolerability of valsartan provided by the compositions may be observed in standard animal tests and in clinical trials.

[0121] The tablets of the invention may be produced by working up components as in WO 97/49394), e.g. as defined hereinabove, in appropriate amounts, to form unit dosage forms.

[0122] For example there is a process of making the tablets as hereinabove described comprising the steps of

i) grinding the active agent and pharmaceutically acceptable additives,

ii) subjecting a mixture of the ground active agent and additives to compression to form a coprimate (the compacted mass)

iii) converting the coprimate to form a granulate and

iv) compressing the granulate to form the solid oral dosage form.

[0123] The process is carried out in the absence of water, i.e. it is a dry compression method. The process may be carried out under ambient conditions of temperature and humidity; it is not necessary to ensure that the process is carried out in a dry atmosphere.

[0124] The initial grinding step i) may be carried out according to conventional milling methods or micronisation methods.

[0125] The active agent and the additives can be milled either individually or together to particle sizes from about 0.1 micrometers (μ) to about 1500 μ, e.g., 1.0 μ to 900 μ, e.g., 60μ to 600 μ. At least 90 % of the crystals of both the active agent and the additives are present in these ranges. Particles of this size are obtained by conventional comminution methods, e.g. grinding in an air jet mill, hammer and screen mill, fine impact mill, ball mill or vibrator mill.

[0126] Micronisation is preferably effected by known methods using an ultrasonic disintegrator, e.g. of the BRANSON Sonifier type, or by stirring a suspension with a high speed agitator, for example with a stirrer of the HOMOREX type.

[0127] The ground particles may optionally at this stage be sieved and mixed according to known methods.

[0128] Compression to form a coprimate requires the compaction of the dry ground components. Compaction may be carried out using a slugging technique or preferably, roller compaction. Roller compaction apparatus is conventional and essentially utilises two rollers which roll towards each other. A hydraulic ram forces one of the rollers against the other to exert a compacting force against the ground particles fed into the roller compactor via a screw conveyor system.

[0129] A compaction force of between 25 and 65 kN, e.g., 25 and 45 kN may be used. Within this range of compaction forces it has surprisingly been found that for each particular formulation a minimum compaction force should be used in order to obtain a solid oral dosage form wherein the granulate disintegrates into discrete primary particles at a desirable rate, e.g. disintegration occurs approximately six times faster for a solid oral dosage form compressed above a minimum compaction force. Such a rapid disintegration rate is unusual for tablets and is similar to the disintegration rate of a capsule formulation. The particular minimum compaction force is dependent on the active agent content in any given formulation and therefore also depends on the amount and nature of the additives present.

[0130] Given this information, the skilled addressee would clearly be able to determine the minimum compaction force for other formulations using routine experimentation and without undue burden.

[0131] The roller speed may be set at between 1 and 20 rpm and preferably 9 to 15 rpm. After passing through the rollers the compacted mass (the coprimate) resembles a thin ribbon in segments.

[0132] The coprimate may be screened and or milled to produce the granulate. Screening in its simplest form involves the passing of the coprimate emerging from the rollers through a seive under mechanical pressure. More preferably, the coprimate is screened using an oscillating mill, e.g. a MGI 624 Frewitt (Key International Inc.).

[0133] The compression of the granulates to tablet cores can be carried out in a conventional tabletting machine, e.g. in an EK-0 Korsch eccentric tabletting machine or a rotary compression machine, e.g., at a compression greater than 2 kN. The tablet cores may vary in shape and be, for example, round, oval, oblong, cylindrical or any other suitable shape, and may also vary in size depending on the concentration of the therapeutic agents. A characteristic of tablets according to the invention is their small size having regard to the amount of active agent contained therein.

[0134] In a preferred embodiment of the invention tablets obtained by the compression method described above are slightly oval. The edges of the tablets may be bevelled or rounded.

[0135] In a particularly preferred embodiment of the invention a solid oral dosage form is compressed in the form of a tablet having an oblong shape in which the ratio of dimensions length:width:height is, e.g., 2.5 - 5.0: 0.9 - 2.0: 1.0, and preferably in which the base and top face of the tablet independently of one another are planar or convexly curved about the longitudinal axis; the side faces are planar, the end faces can be of any shape and the edges are optionally bevelled or rounded.

[0136] In a particularly preferred embodiment of the invention a solid oral dosage form is compressed, from the granulate, in the form of a tablet of oblong shape in which the length is approximately 10.0 to 15.0 mm, the width approximately 5.0 to 6.0 mm and the height approximately 3.0 to 4.0 mm.

[0137] In another particularly preferred embodiment of the invention a solid oral dosage form is compressed from granulates in the form of a tablet of oblong shape in which the length is approximately 15.0 to 18.0 mm, the width approximately 6.0 to 9.0 mm and the height approximately 3.5 to 5.0 mm.

[0138] In yet another preferred embodiment of the invention there is provided a tablet which is essentially disc-shaped with the upper and lower faces having a slightly convex surface. Preferably the tablet has a diameter of about 8 to 8.5 mm and a depth of about 3 to 3.5 mm, or a diameter of about 16 mm and a depth of about 6 mm. The tablets may occupy a volume from about 0.1 cm^3 to about 1 cm^3, e.g., 0.1 cm^3 to about 0.45cm^3, e.g., 0.2 to 0.3 cm^3, e.g about 0.125 cm^3 or 0.25 cm^3.

[0139] They may furthermore be transparent, colourless or coloured and also marked so as to impart to this product an individual appearance and to make them instantly recognizable. The use of dyes can serve to enhance the appearance as well as to identify the compositions. Dyes suitable for use in pharmacy typically include carotinoids, iron oxides or chlorophyll.

[0140] The following examples illustrate the above-described invention; however, it is not intended to restrict the scope of this invention in any manner.

Formulation Example 1:

Film-Coated Tablets:

[0141]

Components	Composition Per Unit (mg)	Standards
Granulation		
Valsartan [= active ingredient]	80.00	
Microcrystalline cellulose/ Avicel PH 102	54.00	NF, Ph. Eur
Crospovidone	20.00	NF, Ph. Eur
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	0.75	Ph. Eur/ NF
Magnesium stearate	2.5	NF, Ph. Eur
Blending		
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	0.75	Ph. Eur/ NF
Magnesium stearate	2.00	NF, Ph. Eur
Coating		
Purified water *)	-	
DIOLACK pale red 00F34899	7.00	
Total tablet mass	167.00	

*) Removed during processing.

[0142] The film-coated tablet is manufactured e.g. as follows:

[0143] A mixture of valsartan, microcrystalline cellulose, crospovidone, part of the colloidal anhydrous silica/colloidal silicon dioxide/Aerosile 200, silicon dioxide and magnesium stearate is premixed in a diffusion mixer and then sieve through a screnning mill. The resulting mixture is again pre-mixed in a diffusion mixer, compacted in a roller compacter and then sieve through a screening mill. To the resulting mixture, the rest of the colloidal anhydrous silica/colloidal silicon dioxide/Aerosile 200 are added and the final blend is made in a diffusion mixer. The whole mixture is compressed in a rotary tabletting machine and the tabletts are coated with a film by using Diolack pale red in a perforated pan.

Formulation Example 2:

Film-coated tablets:

[0144]

Components	Composition Per Unit (mg)	Standards
Granulation		
Valsartan [= active ingredient]	160.00	
Microcrystalline cellulose/ Avicel PH 102	108.00	NF, Ph. Eur
Crospovidone	40.00	NF, Ph. Eur
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	1.50	Ph. Eur/ NF
Magnesium stearate	5.00	NF, Ph. Eur
Blending		
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	1.50	Ph. Eur/ NF
Magnesium stearate	4.00	NF, Ph. Eur
Coating		
Opadry Light Brown 00F33172	10.00	
Total tablet mass	330.00	

The film-coated tablet is manufactured e.g. as described in Formulation Example 1.

Formulation Example 3:

Film-Coated Tablets:

[0145]

Components	Composition Per Unit (mg)	Standards
Core: Internal phase		
Valsartan [= active ingredient]	40.00	
Silica, colloidal anhydrous (Colloidal silicon dioxide) [= Glidant]	1.00	Ph. Eur, USP/NF
Magnesium stearate [= Lubricant]	2.00	USP/NF
Crospovidone [Disintegrant]	20.00	Ph. Eur
Microcrystalline cellulose [= Binding agent]	124.00	USP/NF
External phase		
Silica, colloidal anhydrous, (Colloidal silicon dioxide) [= Glidant]	1.00	Ph. Eur, USP/NF
Magnesium stearate [Lubricant]	2.00	USP/NF
Film coating		
Opadry® brown OOF 16711*)	9.40	
Purified Water**)	-	
Total tablet mass	199.44	

*) The composition of the Opadry® brown OOF16711 coloring agent is tabulated below.

**) Removed during processing

Opadry® Composition:

[0146]

Ingredient	Approximate % Composition
Iron oxide, black (C.I. No. 77499, E 172)	0.50
Iron oxide, brown (C.I. No. 77499, E 172	0.50
Iron oxide, red (C.I. No. 77491, E 172)	0.50
Iron oxide, yellow (C.I. No. 77492, E 172)	0.50
Macrogolum (Ph. Eur)	4.00

(continued)

Ingredient	Approximate % Composition
Titanium dioxide (C.I. No. 77891, E 171)	14.00
Hypromellose (Ph. Eur)	80.00

[0147] The film-coated tablet is manufactured e.g. as described in Formulation Example 1.

Reference Example 1:

Capsules:

[0148]

Components	Composition Per Unit (mg)
Valsartan [= active ingredient]	80.00
Microcrystalline cellulose	25.10
Crospovidone	13.00
Povidone	12.50
Magnesium stearate	1.30
Sodium lauryl sulphate	0.60
Shell	
Iron oxide, red (C.I. No. 77491, EC No. E 172)	0.123
Iron oxide, yellow (C.I. No. 77492, EC No. E 172)	0.123
Iron oxide, black (C.I. No. 77499, EC No. E 172)	0.245
Titanium dioxide	1.540
Gelatin	74.969
Total tablet mass	209.50

[0149] The tablet is manufactured e.g. as follows:

Granulation/Drying

[0150] Valsartan and microcrystallin cellulose are spray-granulated in a fluidised bed granulator with a granulating solution consisting of povidone and sodium lauryl sulphate dissolved in purified water. The granulate obtained is dried in a fluidiesd bed dryer.

Milling/Blending

[0151] The dried granulate is milled together with crospovidone and magnesium stearate. The mass is then blended in a conical srew type mixer for approximately 10 minutes.

Encapsulation

[0152] Teh empty hard gelatin capsules are filled with the blended bulk granules under controlled temperature and humidity conditions. The filed capsules are dedustee, visually inspected, weightchecked and quarantied until by Quality assurance department.

Reference Example 2:

Capsules:

[0153]

Components	Composition Per Unit (mg)
Valsartan [= active ingredient]	160.00
Microcrystalline cellulose	50.20
Crospovidone	26.00
Povidone	25.00
Magnesium stearate	2.60
Sodium lauryl sulphate	1.20
Shell	
Iron oxide, red (C.I. No. 77491, EC No. E 172)	0.123
Iron oxide, yellow (C.I. No. 77492, EC No. E 172)	0.123
Iron oxide, black (C.I. No. 77499, EC No. E 172)	0.245
Titanium dioxide	1.540
Gelatin	74.969
Total tablet mass	342.00

[0154] The formulation is manufactured e.g. as described in Formulation Example 4.

Reference Example 3:

Hard Gelatine Capsule:

[0155]

Components	Composition Per Unit (mg)
Valsartan [= active ingredient]	80.00
Sodium laurylsulphate	0.60
Magnesium stearate	1.30
Povidone	12.50
Crospovidone	13.00
Microcrystalline cellulose	21.10
Total tablet mass	130.00

Examples 4 to 8:

[0156]

Example	4	5	6	7	8
Components	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)
Granulation					
Valsartan Drug Substance	80.000	160.000	40.000	320.000	320.000
Microcrystalline Cellulose (NF, Ph.Eur.)/ Avicel PH 102	54.000	108.000	27.000	216.000	216.000
Crospovidone (NF, Ph.Eur.)	15.000	30.000	7.500	80.000	60.000
Colloidal Anhydrous Silica (Ph. Eur.)/ Colloidal Silicon Dioxide (NF)/ Aerosil 200	1.500	3.000	0.750	3.000	6.000
Magnesium Stearate (NF, Ph.Eur.)	3.000	6.000	1.500	10.000	12.000
Blending					
Colloidal Anhydrous Silica (Ph. Eur.)/ Colloidal Silicon Dioxide (NF)/ Aerosil 200	---	---	---	3.000	-
Magnesium Stearate. NF. Ph.Eur,	1.500	3.000	0.750	8.000	6.000
Core Weight/mg	155.000	310.000	77.500	640.000	620.000

(continued)

Example	4	5	6	7	8
Components	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)	COMPOSITION PER UNIT (mg)
Granulation					
Coating	-	-	3.800	15.000	16.000

Example 9: Dissolution of film-coated tablets (FCT)

[0157] The dissolution acceptance criteria are Q= 75% in 30 min (Paddle 50 rpm, phosphate buffer pH 6.8)

	40mg	Strength	320mg	Strength
Dissolution FCT				
	- level	+ level	- level	+ level
Mean [%]	98	97	93	89
rel. std. [%]	1.06	2.48	2.12	2.34
min. [%]	96	94	90	86
Max. [%]	99	99	95	92

Claims:

1. A tablet comprising 20 to 65% of valsartan in free form and more than 30% of microcrystalline cellulose by weight based on the total weight of the core components of said tablet, wherein the following tablets are disclaimed:

Valsartan Tablet Formulation 80 mg + Amlodipine 5 mg (Rollercompaction)

Dosage (mg)	80 mg Valsartan + 5 mg Amlodipine
Diameter (mm)	9
Shape	round
Breaking line	without
Tablet-weight (mg)	215

	Dosage Strength	Function of the Excipient in the Formulation	80 mg Valsartan + 5 mg Amlodipine
I.	Compactate		mg:
1.	Valsartan DS	drug substance	80.0
2.	Amlodipine DS	drug substance	5.0
3.	Avicel PH 102	filler	104.0
4.	PVPP-XL	disintegrant	20.0
5.	Aerosil 200	glidant	0.75
6.	Magnesiumstearate	lubricant	2.5
II.	Outer Phase		
7.	Aerosil 200	glidant	0.75
8.	Magnesiumstearate	lubricant	2.0
	total		215.0

;

Film-Coated Tablets:

Components	Composition Per Unit (mg)	Standards
Granulation		
Valsartan [=active ingredient]	80.00	
Microcrystalline cellulose/ Avicel PH 102	54.00	NF, Ph. Eur
Crospovidone	20.00	NF, Ph. Eur
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	0.75	Ph. Eur/NF
Magnesium stearate	2.5	NF, Ph. Eur
Blending		
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	0.75	Ph. Eur/NF
Magnesium stearate	2.00	NF, Ph. Eur
Coating		
DIOLACK pale red 00F34899	7.00	
Total tablet mass	167.00	

Film-Coated Tablets:

Components	Composition Per Unit (mg)	Standards
Granulation		
Valsartan [=active ingredient]	160.00	
Microcrystalline cellulose/ Avicel PH 102	108.00	NF, Ph. Eur
Crospovidone	40.00	NF, Ph. Eur
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	1.50	Ph. Eur/NF
Magnesium stearate	5.00	NF, Ph. Eur
Blending		
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	1.50	Ph. Eur/NF
Magnesium stearate	4.00	NF, Ph. Eur
Coating		
Opadry Light Brown 00F33172	10.00	
Total tablet mass	330.00	

;

and

Film-Coated Tablets:

Components	Composition Per Unit (mg)	Standards
Core: Internal phase		
Valsartan [=active ingredient]	40.00	
Silica, colloidal anhydrous (Colloidal silicon dioxide) [=Glidant]	1.00	Ph. Eur, USP/NF
Magnesium stearate [=Lubricant]	2.00	USP/NF
Crospovidone [Disintegrant]	20.00	Ph. Eur
Microcrystalline cellulose [Binding agent]	124.00	USP/NF
External Phase		
Silica, colloidal anhydrous, (Colloidal silicon dioxide) [=Glidant]	1.00	Ph. Eur, USP/NF
Magnesium stearate [Lubricant]	2.00	USP/NF
Film Coating		
Opadry® brown OOF 16711	9.40	
Total tablet mass	199.44	

.

2. A tablet according to claim 1 comprising up to 65% of microcrystalline cellulose.

3. A tablet according to claim 2 comprising 50 to 65 % of microcrystalline cellulose.

4. A tablet according to any one of claims 1 to 3 comprising less than 13% of crospovidone.

5. A tablet according to any one of claims 1 to 4, wherein the active ingredient consists of valsartan.

6. A tablet according to any one of claims 1 to 5 comprising 20 to 360 mg valsartan.

7. A tablet according to any one of claims 1, 2, 5 and 6 comprising
20 to 65% of valsartan
31 to 65% of microcrystalline cellulose and
2 to 13% of crospovidone.

8. A tablet according to any one of claims 1 to 7 comprising more than 250 mg and up to 360 mg valsartan.

9. A tablet according to any one of claims 1 to 8, wherein the tablet is a coated tablet.