


Description

This invention relates to certain new herbicidal acrylonitrile derivatives, their preparation, herbicidal compositions containing such derivatives and to a method of combating undesired plant growth using such derivatives and compositions.

The preparation and herbicidal use of certain acrylonitrile derivatives has been described in DE-A 2 330 913. Such acrylonitriles have an α -carbonyl linkage with a substituent selected from optionally substituted phenyl, naphthyl, thienyl and furyl groups, the optional substituents being restricted to halogen, nitro, alkoxy and alkyl.

The present invention relates to a new class of herbicidal aryl- or heteroaryl-oxybenzoylacrylonitrile derivatives.

Accordingly, the present invention provides a compound of the general formula I



in which R^1 and R^2 each independently represents an alkyl group, Z represents a nitrogen atom or a group CH which is unsubstituted or substituted by a substituent X , m is 0 or an integer from 1 to 4, the or each Y independently represents a halogen atom, n is 0 or an integer from 1 to 5, and the or each X independently represents a halogen atom or a group selected from alkyl, haloalkyl, cyanoalkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, alkynyloxy, alkylthio, alkenylthio, alkynylthio, cyano, nitro, alkylsulphonyl, alkylsulphinyl and sulphonamido.

As used herein the term alkyl, alkenyl or alkynyl in respect of a radical or moiety refers to a straight or branched chain radical or moiety. Suitably an alkyl radical or moiety has from 1 to 12 carbon atoms, preferably from 1 to 6 and especially from 1 to 4 carbon atoms. Alkenyl and alkynyl radicals or moieties suitably have from 2 to 12 carbon atoms, preferably from 2 to 6, especially from 2 to 4, carbon atoms.

Preferably R^1 and R^2 each independently represents an alkyl group of 1 to 4 carbon atoms: and most preferably R^1 and R^2 are the same and each represents a methyl group.

Z preferably represents an unsubstituted group CH , the compound of general formula I then being a diphenyl ether derivative.


When m is greater than 1, the substituents Y may be the same or different and may suitably be selected from chlorine, bromine and fluorine atoms. Preferably either m or 0 or m is 1 and Y represents a chlorine atom.

When n is greater than 1, the substituents X may be the same or different and may suitably be selected from halogen atoms, especially fluorine, chlorine and bromine atoms, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{1-6} cyanoalkyl, C_{1-6} alkoxy groups, C_{1-6} alkylthio groups, C_{1-6} alkylsulphinyl groups, C_{1-6} alkylsulphonyl groups, cyano groups and nitro groups. Preferably n is 1 or 2 and the or each substituent X independently represents a fluorine atom, a chlorine atom, a bromine atom, a C_{1-4} alkyl group, a C_{1-4} haloalkyl group, especially trifluoromethyl, a C_{1-4} cyanoalkyl group, especially cyanomethyl, a C_{1-4} alkoxy group, especially methoxy, a C_{1-4} alkylthio group, especially methylthio, a C_{1-4} alkylsulphinyl group, especially methylsulphinyl, a C_{1-4} alkylsulphonyl group, especially methylsulphonyl, a cyano group or a nitro group.

The compounds of general formula I may either be prepared from a corresponding diphenyl ether or pyridyl phenyl ether derivative by reaction in one or more steps to introduce the dialkylaminoacrylonitrile moiety, or be prepared by introducing a suitable phenyl or pyridyl moiety into an appropriate benzoyldialkylamino acrylonitrile derivative.

According to a further aspect of the invention, we therefore provide a process for the preparation of a compound of general formula I, which comprises reacting a compound of general formula II

5




10 wherein X, Y, Z, n and m are as defined above and W represents a halogen atom or a cyanomethyl group, with, in the case where W is halogen, a dialkylaminoacrylonitrile under basic conditions, or, in the case where W is cyanomethyl, either with a dialkylformamide dialkyl acetal or with a trialkylorthoformate followed by reaction with a dialkylamine,

or

15 reacting a compound of general formula III


20



wherein n , X and Z are as hereinbefore defined and L represents a leaving group with a compound of general formula IV

25

30



35 wherein m , Y, R^1 and R^2 are as hereinbefore defined, under basic conditions, and, if desired or required, converting a resulting compound of general formula I into another of general formula I.

When W in a compound of general formula II represents a cyanomethyl group, the reaction to form the compound of formula I is carried out using a dialkylformamide dialkyl acetal, preferably dimethyl formamide dimethyl acetal. Suitably the reaction is carried out in the presence of an inert organic solvent, suitable solvents being halogenated hydrocarbons, for example dichloromethane, ethers, such as diethyl ether, or esters, such as ethyl acetate; mixtures of solvents may also be utilised. Preferably the reaction is carried out at a temperature in the range of from 0 to 50 °C, suitably at ambient temperature.

Alternatively, when W in a compound of general formula II represents a cyanomethyl group, the reaction may be carried out in a two stage procedure by reacting the cyanomethyl carbonyl compound II with a trialkylorthoformate, preferably trimethylorthoformate, followed, optionally after isolation of the alkoxyaryl acrylonitrile so formed, by reaction with excess dialkylamine, preferably dimethylamine in a suitable solvent. Suitable solvents are, for example, halogenated hydrocarbons, such as dichloromethane, alcohols, such as ethanol, ethers such as diethyl ether, and esters; mixtures of solvents are also suitable.

50 The reaction may be carried out at a temperature in the range of from -10 ° to 100 °C. Preferably the first stage is carried out at elevated temperature, for example at a temperature in the range of from 60 to 100 °C, suitably at approximately 80 °C, while the second stage is carried out at a lower temperature, suitably a temperature below 60 °C and conveniently ambient temperature. Preferably the reaction is carried out without isolation of the intermediate acrylonitrile.

55 When W in a compound of general formula II represents a halogen atom, preferably chlorine, the compound of formula II is reacted with a dialkylaminoacrylonitrile, preferably dimethylaminoacrylonitrile preferably in the presence of a solvent, for example an ether solvent, and in the presence of a base, for example a tertiary amine. A preferred ether solvent is dioxan and a preferred tertiary amine base is

triethylamine. The reaction is preferably carried out at a temperature in the range of from 50 to 120 °C, conveniently at the reflux temperature of the solvent employed.


Compounds of the general formula I may also be prepared by reacting a compound of general formula III with a compound of general formula IV. Preferably the reaction is carried out under basic conditions and in the presence of a suitable solvent. Suitably the reaction is carried out in the presence of an alkali metal or alkaline earth metal carbonate, for example potassium carbonate. Suitable solvents are organic solvents that are inert to the reaction media and conveniently may be, for example, dimethyl formamide; again mixtures of solvents may be used. Preferably the reaction is carried out under anhydrous conditions. The reaction may conveniently be effected at a temperature in the range of from 0 to the reflux temperature of the reaction mixture, but is preferably carried out at reflux.

A leaving group represented by L may be any group that will, under the reaction conditions, cleave from the reactant III thus promoting reaction at a specified site. The group L in a compound of general formula III is suitably a halogen atom, preferably chlorine or fluorine.

Following either method of preparation the resulting compound of general formula I may be isolated and purified using conventional techniques, for example by solvent extraction, evaporation and recrystallisation or by chromatography on silica.

Compounds of general formula II are believed to be novel intermediates.

Compounds of general formula II in which W represents a cyanomethyl group may be prepared from the alkyl ester of the corresponding diaryl or aryl heteroaryl ether. Thus a process for the preparation of a compound of formula II as defined above comprises reacting a compound of formula V:



Wherein X, Y, Z, n and m are as defined above and R represents an alkyl group, with acetonitrile under basic conditions.

Preferably the basic conditions are provided by an alkali metal amide, such as sodamide or potassium amide, suitably in liquid ammonia, or sodium or potassium hydride in a solvent such as dimethylformamide. Alternatively the basic conditions may be provided by an alkoxide such as sodium or potassium methoxide with excess acetonitrile as solvent. Suitable temperatures for the reaction are in the range of from -70 °C to 100 °C and are usually determined by the basic conditions employed.

Compounds of formula V may be prepared by esterification of the corresponding acid with an alcohol, such as methanol, in the presence of an acidic catalyst, for example concentrated sulphuric acid, dry hydrogen chloride gas or p-toluene sulphonic acid. The reactant acids may suitably be prepared by oxidation of the corresponding diaryl/aryl heteroaryl ether aldehyde using, for example, chromium trioxide in the presence of acetic acid or, more preferably, sulphuric acid in an inert solvent such as an aromatic solvent, eg. benzene, or a ketone, eg. acetone, or manganese dioxide in an inert solvent such as a halogenated hydrocarbon, eg. dichloromethane, an ester, eg. ethyl acetate, or a ketone. Suitable temperatures are in the range of from -20 to 50 °C, preferably -5 to 20 °C. The ether aldehydes may themselves conveniently be prepared by condensation of an optionally substituted phenol and a m-halobenzaldehyde, for example in the presence of an alkali metal alkoxide, such as sodium methoxide, followed by treatment with a copper catalyst such as cuprous chloride in pyridine and an aromatic hydrocarbon, such as xylene, as described in U.K. patent specification no. 2050168. The reaction may alternatively be carried out in the presence of an alkali metal carbonate, for example sodium or potassium carbonate, followed by treatment with cuprous oxide and/or copper powder in dimethylformamide or quinoline. The reaction may suitably be carried out at a temperature in the range of from 20 to 150 °C and conveniently the reaction is carried out at the reflux temperature of the reaction mixture.


Alternatively, the compounds of general formula V may be prepared by condensation of an optionally substituted phenol with a meta-halobenzoic acid ester, suitably using conditions similar to those described above for the preparation of the diaryl/aryl heteroaryl ether aldehydes.

Compounds of general formula II in which W represents a halogen atom may be derived from the corresponding ether carboxylic acid, for example by reaction of the acid with thionyl chloride. The acid may be derived as described above from a phenolic compound and a m-halobenzaldehyde or m-halobenzoic

acid ester.

Compounds of general formula III are known compounds and may be prepared using conventional techniques.

The compounds of general formula IV are believed to be novel and may be prepared by reacting a
5 compound of general formula VI



in which Y_m is as defined above and W represents a cyanomethyl group,
with a dialkylformamide dialkyl acetal, preferably dimethylformamide dimethylacetal, under the same
conditions as described above for the reaction of compounds of general formula II in which W is a
20 cyanomethyl group, with a dialkylformamide dialkyl acetal.

The compounds of the invention have been found to have surprisingly high herbicidal activity with a wide spectrum of activity against broadleaved weeds and grasses especially when applied post-emergence.

Accordingly, the invention further provides a herbicidal composition comprising a compound of formula I as defined above in association with at least one carrier, and a method of making such a composition
25 which comprises bringing a compound of formula I into association with at least one carrier.

The invention also provides the use of such a compound or composition according to the invention as a herbicide. Further, in accordance with the invention there is provided a method of combating undesired plant growth at a locus by treating the locus with a compound or composition according to the invention. Application to the locus may be pre-emergence or post-emergence, preferably post-emergence. The
30 dosage of active ingredient used may, for example, be in the range of from 0.01 to 10 kg/ha, suitably 0.05 to 4kg/ha. The locus may, for example, be the soil or plants in a crop area, typical crops being cereals such as wheat, barley and rice.

A carrier in a composition according to the invention is any material with which the active ingredient is formulated to facilitate application to the locus to be treated, which may for example be a plant, seed or
35 soil, or to facilitate storage, transport or handling. A carrier may be a solid or a liquid, including a material which is normally gaseous but which has been compressed to form a liquid, and any of the carriers normally used in formulating herbicidal compositions may be used. Preferably compositions according to the invention contain 0.5 to 95% by weight of active ingredient.

Suitable solid carriers include natural and synthetic clays and silicates, for example natural silicas such
40 as diatomaceous earths; magnesium silicates, for example talcs; magnesium aluminium silicates, for example attapulgites and vermiculites; aluminium silicates, for example kaolinities, montmorillonites and micas; calcium carbonate; calcium sulphate; ammonium sulphate; synthetic hydrated silicon oxides and synthetic calcium or aluminium silicates; elements, for example carbon and sulphur; natural and synthetic resins, for example coumarone resins, polyvinyl chloride, and styrene polymers and copolymers; solid
45 polychlorophenols; bitumen; waxes; and solid fertilisers, for example superphosphates.

Suitable liquid carriers include water; alcohols, for example isopropanol and glycols; ketones, for example acetone, methyl ethyl ketone, methyl isobutyl ketone and cyclohexanone; ethers; aromatic or
50 araliphatic hydrocarbons, for example benzene, toluene and xylene; petroleum fractions, for example kerosine and light mineral oils; chlorinated hydrocarbons, for example carbon tetrachloride, perchloroethylene and trichloroethane. Mixtures of different liquids are often suitable.

Agricultural compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small mounts of a carrier which is a surface-active agent facilitates this process of dilution. Thus preferably at least one carrier in a composition
according to the invention is a surface-active agent. For example the composition may contain at least two
55 carriers, at least one of which is a surface-active agent.

A surface-active agent may be an emulsifying agent, a dispersing agent or a wetting agent; it may be nonionic or ionic. Examples of suitable surface-active agents include the sodium or calcium salts of polyacrylic acids and lignin sulphonic acids; the condensation of fatty acids or aliphatic amines or amides

containing at least 12 carbon atoms in the molecule with ethylene oxide and/or propylene oxide; fatty acid esters of glycerol, sorbitol, sucrose or pentaerythritol; condensates of these with ethylene oxide and/or propylene oxide; condensation products of fatty alcohol or alkyl phenols, for example *p*-octylphenol or *p*-octylcresol, with ethylene oxide and/or propylene oxide; sulphates or sulphonates of these condensation products; alkali or alkaline earth metal salts, preferably sodium salts, of sulphuric or sulphonic acid esters containing at least 10 carbon atoms in the molecule, for example sodium lauryl sulphate, sodium secondary alkyl sulphates, sodium salts of sulphonated castor oil, and sodium alkylaryl sulphonates such as dodecylbenzene sulphonate; and polymers of ethylene oxide and copolymers of ethylene oxide and propylene oxide.

The compositions of the invention may for example be formulated as wettable powders, dusts, granules, solutions, emulsifiable concentrates, emulsions, suspension concentrates and aerosols. Wettable powders usually contain 25, 50 or 75% w of active ingredient and usually contain in addition to solid inert carrier, 3-10% w of a dispersing agent and, where necessary, 0-10% w of stabiliser(s) and/or other additives such as penetrants or stickers. Dusts are usually formulated as a dust concentrate having a similar composition to that of a wettable powder but without a dispersant, and are diluted in the field with further solid carrier to give a composition usually containing $\frac{1}{2}$ -10% w of active ingredient. Granules are usually prepared to have a size between 10 and 100 BS mesh (1.676 - 0.152 mm), and may be manufactured by agglomeration or impregnation techniques. Generally, granules will contain $\frac{1}{2}$ -75% w active ingredient and 0-10% w of additives such as stabilisers, surfactants, slow release modifiers and binding agents. The so-called "dry flowable powders" consist of relatively small granules having a relatively high concentration of active ingredient. Emulsifiable concentrates usually contain, in addition to a solvent and, when necessary, co-solvent, 10-50% w/v active ingredient, 2-20% w/v emulsifiers and 0-20% w/v of other additives such as stabilisers, penetrants and corrosion inhibitors. Suspension concentrates are usually compounded so as to obtain a stable, non-sedimenting flowable product and usually contain 10-75% w active ingredient, 0.5-15% w of dispersing agents, 0.1-10% w of suspending agents such as protective colloids and thixotropic agents, 0-10% w of other additives such as defoamers, corrosion inhibitors, stabilisers, penetrants and stickers, and water or an organic liquid in which the active ingredient is substantially insoluble; certain organic solids or inorganic salts may be present dissolved in the formulation to assist in preventing sedimentation or as anti-freeze agents for water.

Aqueous dispersions and emulsions, for example compositions obtained by diluting a wettable powder or a concentrate according to the invention with water, also lie within the scope of the invention. The said emulsions may be of the water-in-oil or of the oil-in-water type, and may have a thick 'mayonnaise'-like consistency.

The composition of the invention may also contain other ingredients, for example other compounds possessing herbicidal, insecticidal or fungicidal properties.

The following Examples illustrate the invention, Examples 1 to 5 relate to the preparation of starting materials, Examples 6 to 44 to the preparation of novel intermediates of general formula II and Examples 45 to 91 to the preparation of compounds of general formula I.

All structures were confirmed by mass spectroscopy and 300'H nmr.

Example 1

Preparation of 3-(3'- α,α,α -trifluoromethylphenoxy)benzaldehyde

A solution of sodium (2.65g) in methanol (45ml) was added to a stirred solution of 3-(α,α,α -trifluoromethyl)phenol (17.8g) in xylene (125ml) under a nitrogen atmosphere. The mixture was heated to reflux and xylene distilled out, with continuous replacement by fresh, dry xylene until the total distillate was 200ml. The mixture was cooled, cuprous chloride (3.0g) and pyridine (50ml) were added and the mixture heated to reflux. A solution of 3-bromobenzaldehyde (18.5g) in xylene (25ml) was added dropwise and the mixture refluxed for 10 hours. After cooling, the mixture was poured into water (250ml) and acidified with dilute hydrochloric acid. The aqueous solution was extracted with diethyl ether. The ethereal extracts were washed with aqueous sodium hydrogen carbonate and brine and dried over anhydrous MgSO_4 . After filtration and evaporation, the residual liquid was chromatographed on a silica gel column to give 3-(3'- α,α,α -trifluoromethylphenoxy)benzaldehyde as a colourless oil (9g, 34%).

Analysis		
Calculated for $C_{14}H_9O_2F_3$:	C 63.2;	H 3.4%
Found	C 63.6;	H 3.5%

Example 2Preparation of 3-(3'- α,α,α -trifluoromethylphenoxy) benzoic acid

A solution of oxidation mixture (45ml), (derived from 41.2g CrO_3 in 120 ml water and 35 ml concentrated sulphuric acid) was added to a solution of 3-(3'- α,α,α -trifluoromethylphenoxy)benzaldehyde (24.3g) in acetone (50ml), keeping the temperature below 25°C. After stirring at room temperature for 1 hour, ice-water (500g) was added and the white solid filtered off to give the title product (25g, 97%) of m.p. 109-110°C.

Analysis:		
Calculated for $C_{14}H_9O_3F_3$:	C 59.6;	H 3.2%
Found	C 59.9;	H 3.3%

Example 3Preparation of Methyl-3-(3'- α,α,α -trifluoromethylphenoxy)benzoate

A solution of 3-(3'- α,α,α -trifluoromethylphenoxy)benzoic acid (23g) in methanol (200ml) with concentrated H_2SO_4 (0.5ml) was refluxed for 6 hours. The methanol was evaporated in vacuo and the residue dissolved in diethyl ether. The ethereal solution was washed with aqueous $NaHCO_3$ and brine, dried over anhydrous $MgSO_4$, filtered and evaporated to give the title ester (22g, 91%) as a yellow oil.

Analysis:		
Calculated for $C_{15}H_{11}O_3F_3$:	C 60.8;	H 3.7%
Found	C 61.6;	H 4.0%

Example 4Preparation of Ethyl-3-(4'-fluorophenoxy) benzoate

A solution of sodium methoxide (from 2.65g sodium and 45ml of methanol) was added to a solution of 4-fluorophenol (12.3g) in xylene (100ml). The resulting solution was evaporated to dryness in vacuo. Fresh xylene (100ml) was added to the sodium phenolate and the xylene re-evaporated. Pyridine (50ml) and xylene (100ml) were added to the dry sodium phenolate, followed by cuprous chloride (3g). The solution was heated to reflux with stirring and a solution of 3-bromobenzoate (22.9g) in xylene (25ml) was added dropwise. The resultant mixture was refluxed a further 6 hours. After cooling, the reaction mixture was poured into water, acidified with dilute HCl and the aqueous phase extracted with diethyl ether. After drying over anhydrous magnesium sulphate, the ether was evaporated. The residual oil was purified on a silica gel column using methylene chloride as eluant to give the title compound (21g) as a pale yellow oil.

Analysis:		
Calculated for $C_{15}H_{13}O_3F$:	C 69.2;	H 5.0%
Found:	C 68.5;	H 4.8%

Example 5Preparation of Methyl-3-(5'- α,α,α -trifluoromethyl-2-pyridyloxy)benzoate

- 5 Methyl-3-hydroxybenzoate (9.1g) in dry dimethylformamide (20ml) is added dropwise to a suspension of sodium hydride (2.6g of 60% dispersion in oil) in dry dimethyl formamide (60ml), the temperature being maintained below 20 °C. After stirring for a further 30 minutes, 2-chloro-5- α,α,α -trifluoromethyl pyridine (10.9g) in dry dimethyl formamide (20ml) was added dropwise and the resulting solution stirred overnight at room temperature. The reaction mixture was then poured into water and the product extracted with diethyl ether. The ethereal extracts were dried over MgSO_4 , filtered and evaporated. The residual oil was chromatographed on a silica gel column using dichloromethane as eluant to give the title compound as a colourless oil (11.5g, 65%).

15

Analysis:			
Calculated for $\text{C}_{14}\text{H}_{10}\text{NO}_3\text{F}_3$	C 56.5;	H 3.4;	N 4.7%
Found	C 56.0;	H 3.6;	N 5.1%

20

Example 6Preparation of 3-(3'- α,α,α -trifluoromethylphenoxy)benzoyl acetonitrile

- 25 To a suspension of sodamide (prepared from 1.2g sodium metal in 60 ml of liquid ammonia) was added a solution of dry acetonitrile (2.1g) in dry diethylether (5ml) over 5 minutes. After a further 5 minutes methyl-3-(3'- α,α,α -trifluoromethyl)phenoxy)benzoate (7.4g) was added as quickly as possible. After stirring for a further 1 hour at a temperature between -30 to -15 °C, the ammonia was removed by warming on a water bath, whilst at the same time, diethyl ether was added to maintain the reaction flask volume of approximately 100ml. The ethereal solution was then cautiously poured onto ice (50g). The aqueous layer was separated, washed with ether and acidified with 6N HCl. Purification of the filtered solid on a silica gel column using 5% v/v diethyl ether/dichloromethane as eluant gave the title compound (5g, 66%) as a white solid of m.p. 60-61 °C.

35

Analysis:			
Calculated for $\text{C}_{16}\text{H}_{10}\text{O}_2\text{NF}_3$	C 62.9;	H 3.3;	N 4.6%
Found	C 63.0;	H 3.3;	N 4.8%

40

Example 7Preparation of 3-phenoxybenzoylacetonitrile

- 45 A solution of sodium methoxide (from 1.2g sodium in 25ml methanol) was added dropwise over 2 hours to a refluxing solution of methyl-3-phenoxybenzoate (11g) in acetonitrile (100ml) under a nitrogen atmosphere while distilling off methanol and excess acetonitrile. After complete removal of methanol the reaction mixture was poured onto ice-water (200g) and acidified with 50% HCl. The solid was filtered off and dried to give the title compound (7.5g, 65%) as a white solid of m.p. 95-96 °C.

50

Analysis:			
Calculated for $\text{C}_{15}\text{H}_{11}\text{O}_2\text{N}$	C 75.9;	H 4.6;	N 5.9%
Found	C 75.7;	H 4.7;	N 5.9%

55

Example 8Preparation of 3-(5'- α,α,α -trifluoromethyl-2-pyridyloxy)benzoylacetonitrile

- 5 Dry acetonitrile (2.8g) was added to a stirred suspension of sodium hydride (2.7g of 60% dispersion in oil) in dry dimethyl formamide (50ml). The mixture was heated to 60°C for 1 hour. Methyl-3-(5'- α,α,α -trifluoromethyl-2-pyridyloxy)benzoate (10g) was added and the reaction mixture left to cool and stirred at room temperature overnight. The mixture was poured onto ice and the product extracted with diethyl ether. The aqueous phase was acidified with dilute HCl and extracted with diethyl ether. The ethereal extracts
10 were dried over MgSO_4 , filtered and evaporated. The residual oil was chromatographed on a silica gel column using dichloromethane as eluant to give the title compound (4.8g, 47%) as a white solid of m.p. 85-86°C.

15

Analysis:			
Calculated for $\text{C}_{15}\text{H}_9\text{N}_2\text{O}_2\text{F}_3$	C 58.8;	H 2.9;	N 9.2%
Found	C 58.5;	H 2.9;	N 9.0%

20

The compounds of general formula II listed in Table 1 below, were prepared by methods similar to those described in Examples 6,7 and 8 above.

25

30

35


40

45

50

55

Table 1



Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
9	2-Cl, 4-CF ₃	CH	-	56.5 55.5	2.6 2.7	4.1 4.0	not recorded
10	4-Cl	CH	-	66.3 66.6	3.7 3.9	5.2 5.7	96-97
11	4-CH ₃	CH	-	76.5 76.3	5.2 5.2	5.6 5.7	77-78

Table 1 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
12	3,5-diCl	CH	-	58.8 58.5	2.9 3.1	4.5 4.1	123-124
13	4-OCH ₃	CH	-	71.9 72.2	4.9 4.8	5.2 5.4	78-79
14	3,4-diCl	CH	-	58.8 59.1	2.9 3.0	4.6 4.6	99-100
15	4-isopropyl	CH	-	77.4 77.1	6.1 6.0	5.0 5.3	77-79
16	2,4-diCH ₃	CH	-	77.0 76.5	5.7 5.6	5.3 5.6	116-117
17	5-Cl	N	-	61.8 61.6	3.3 3.3	10.3 10.3	106-107
18	2-CH ₃	CH	-	76.5 73.3	5.2 4.9	5.6 6.0	55-57

Table 1 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
19	4-SCH ₃	CH	-	67.8 67.4	4.6 4.6	4.9 5.0	103-104
20	2,4-diCl	CH	-	58.8 58.3	2.9 3.0	4.6 4.8	102-104
21	4-F	CH	-	70.6 70.3	3.9 4.0	5.5 5.9	104-105
22	3,4-diF	CH	-	65.9 65.5	3.3 3.3	5.1 5.7	89-90
23	2,5-diCH ₃	CH	-	77.0 77.5	5.7 5.8	5.3 5.3	109-110
24	2-F	CH	-	70.6 70.1	3.9 3.2	5.5 5.8	60-64
25	3-F	CH	-	70.6 67.0	3.9 3.8	5.5 5.9	62-64

Table 1 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
26	4-CN	CH	-	73.3 72.8	3.8 3.8	10.7 10.7	127-129
27	3-Cl	N	-	61.7 61.7	3.3 3.6	10.3 10.2	129-130
28	5-Br	N	-	53.0 52.8	2.8 2.9	8.8 8.9	118-119
29	3-CF ₃ , 4-Cl	CH	-	56.6 56.6	2.7 2.8	4.1 4.2	99-101
30	4-C ₂ H ₅	CH	-	77.0 77.9	5.7 5.3	5.3 5.9	67-69
31	4-Br	CH	-	57.0 56.5	3.2 3.2	4.4 4.4	101-103
32	4-tC ₄ H ₉	CH	-	77.8 71.3	6.5 6.0	4.8 4.8	oil

Table 1 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
33	4-secC ₄ H ₉	CH	-	77.8 74.4	6.5 6.4	4.8 4.4	oil
34	2-CH ₃ , 4-Cl	CH	-	67.4 66.2	4.2 4.0	4.9 4.7	125-126
35	3-CF ₃ , 4-F	CH	-	59.4 59.0	2.8 3.2	4.3 4.6	68-70
36	2-Cl	CH	-	66.4 64.9	3.7 3.9	5.2 5.0	50-51
37	3-CH ₃ , 4-Cl	CH	-	67.4 67.5	4.2 4.2	4.9 4.6	89-90
38	3-Cl	CH	-	66.4 66.1	3.7 3.8	5.2 5.4	87-88
39	3-OCH ₃	CH	-	71.9 71.4	4.9 4.9	5.2 5.4	55-56

Table 1 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
40	2,4-diF	CH	-	65.9 64.9	3.3 3.2	5.1 5.0	oil
41	2-Cl, 4-F	CH	-	62.3 61.9	3.1 3.2	4.8 5.1	95-96
42	3,5-diCH ₃	CH	-	77.0 76.9	5.7 5.7	5.3 5.7	75-76
43	3-CF ₃	CH	5-Cl	56.6 55.7	2.6 2.7	4.1 4.0	oil
44	3-CH ₂ CN, 5-CF ₃	N	-	59.1 59.0	2.9 3.1	12.2 12.0	not recorded

Example 45Preparation of 2-[3-(3'- α,α,α -trifluoromethyl)phenoxy]benzoyl]-3-dimethylaminoacrylonitrile

- 5 To a solution of 3-(3'- α,α,α -trifluoromethyl)phenoxy)benzoyl acetonitrile (3g) in dichloromethane (50ml) was added dimethylformamide dimethylacetal (6ml). After stirring overnight at room temperature, the dichloromethane was evaporated. The residue was purified on a silica gel column using 5% methanoldichloromethane (v/v) as eluant to give the title compound (3g, 85%) as a white solid of m.p. 84-85 ° C.

10

Analysis:			
Calculated for $C_{19}H_{15}N_2O_2F_3$:	C 63.3;	H 4.2;	N 7.8%
Found:	C 63.4;	H 4.5;	N 7.8%

15

Example 46Preparation of 2-(3-phenoxybenzoyl)-3-dimethylaminoacrylonitrile

20

- 3-Phenoxybenzoylacetoneitrile (4g) and trimethylorthoformate (30ml) was stirred and heated to 80 ° C for 2 hours. After cooling to 0 ° C, an excess of dimethylamine in ethanol solution was added and the mixture stirred for 2 hours at room temperature. The volatiles were then removed in vacuo and the residue chromatographed on a silica gel column using 5% methanol-dichloromethane (v/v) as eluant to give the title compound (3.2g, 65%) as a white solid of m.p. 80-82 ° C.

25

Analysis:			
Calculated for $C_{18}H_{15}O_2N_2$	C 73.9;	H 5.5;	N 9.6%
Found	C 73.5;	H 5.5;	N 9.7%

30

Example 47Preparation of 2-(3-phenoxybenzoyl)-3-dimethylaminoacrylonitrile

- A mixture of 3-phenoxybenzoyl chloride (6.9g), 3-dimethylaminoacrylonitrile (2g) and triethylamine (4g) in dry dioxan (50ml) and refluxed for 6 hours under a nitrogen atmosphere. After standing at room temperature overnight, the dioxan was evaporated in vacuo and dichloromethane added to the residue. The extracts were washed with water, 2N NaOH, brine and finally dried over $MgSO_4$. After filtration and evaporation of the dichloromethane, the residue was purified on a silica gel column using 5% methanol-dichloromethane as eluant to give the title compound (2.2g, 36%) as a white solid of m.p. 80-82 ° C (identical to that described in Example 46).

Example 48Preparation of 2-[3-(5'- α,α,α -trifluoromethyl-2-pyridyloxy)benzoyl]-3-dimethylaminoacrylonitrile


- Dimethylformamide-dimethyl acetal (3ml) was added to a solution of 3-(5'- α,α,α -trifluoromethyl-2-pyridyloxy)benzoyl acetonitrile (1g) in dichloromethane (50ml). After stirring at room temperature overnight, the methylene chloride was evaporated and the residual oil chromatographed on a silica gel column using 5% methanol-dichloromethane (v/v) as eluant to give the title product (0.9g, 85%) as a viscous colourless oil.

55

Analysis:			
Calculated for $C_{18}H_{14}N_3O_2F_3$:	C 59.8;	H 3.8;	N 11.6%
Found:	C 57.8;	H 3.7;	N 11.2%

The compounds of general formula I listed in Table 2, were prepared by methods similar to those described in Examples 45 to 48 above.

Table 2



Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
49	2-Cl, 4-CF ₃	CH	-	57.8 57.4	3.5 3.5	7.1 6.9	68-70
50	4-Cl	CH	-	66.3 66.4	4.6 4.6	8.6 8.6	101-102
51	4-CH ₃	CH	-	74.5 74.4	5.9 6.0	9.2 9.4	120-121

Table 2 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
52	3,5-diCl	CH	-	59.6 58.8	3.9 4.5	7.7 7.6	101-102
53	4-OCH ₃	CH	-	70.8 70.7	5.6 5.7	8.7 8.8	117-118
54	3,4-diCl	CH	-	59.6 60.6	3.9 4.1	7.7 8.0	102-103
55	4-isopropyl	CH	-	75.5 74.3	6.6 6.6	8.4 8.5	81-83
56	2,4-diCH ₃	CH	-	75.0 74.1	6.3 6.2	8.8 8.6	120-121
57	5-Cl	N	-	62.4 62.5	4.3 4.9	12.8 12.9	119-120
58	2-CH ₃	CH	-	74.5 71.5	5.9 5.8	9.2 8.9	79-81

Table 2 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
59	4-SCH ₃	CH	-	67.5 67.3	5.3 5.3	8.3 8.3	88-90
60	2,4-diCl	CH	-	60.0 60.8	3.9 4.1	7.8 7.7	88-90
61	4-F	CH	-	69.7 67.8	4.8 5.0	9.0 9.0	oil
62	3,4-diF	CH	-	65.8 60.3	4.3 4.2	8.5 7.9	oil
63	2,5-diCH ₃	CH	-	75.0 74.0	6.3 6.6	8.7 8.4	oil
64	2-F	CH	-	69.7 68.6	4.8 4.5	9.0 9.1	oil
65	3-F	CH	-	69.7 63.7	4.8 4.6	9.0 8.7	oil

Table 2 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
66	4-CN	CH	-	71.9 71.2	4.7 4.7	13.2 13.3	121-122
67	3-Cl	N	-	62.3 62.1	4.3 4.3	12.8 12.8	oil
68	5-Br	N	-	54.8 54.6	3.7 3.8	11.3 11.0	120-122
69	3-CF ₃ , 4-Cl	CH	-	57.9 56.9	3.6 3.7	7.1 7.0	92-94
70	4-C ₂ H ₅	CH	-	75.0 73.7	6.3 6.3	8.8 8.8	93-95
71	4-Br	CH	-	58.2 57.9	4.0 4.1	7.5 7.7	111-113
72	4-tC ₄ H ₉	CH	-	75.9 75.8	6.9 7.1	8.1 8.0	94-96

Table 2 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
73	4-secC ₄ H ₉	CH	-	75.9 78.3	6.9 7.6	8.1 7.6	oil
74	2-CH ₃ ,4-Cl	CH	-	67.1 66.1	5.0 5.0	8.2 8.2	120-122
75	3-CF ₃ ,4-F	CH	-	60.3 58.4	3.7 4.0	7.4 7.2	oil
76	2-Cl	CH	-	66.3 67.5	4.6 4.8	8.6 8.4	72-74
77	3-CH ₃ ,4-Cl	CH	-	67.1 67.0	5.0 5.0	8.3 8.2	108-109
78	3-Cl	CH	-	66.3 66.9	4.6 4.8	8.6 8.5	oil
79	3-OCH ₃	CH	-	70.8 69.5	5.6 5.6	8.7 8.7	oil

Table 2 (cont.)

Example No.	Xn	Z	Ym	Elemental Analysis (%)			Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	N (Calc. Found)	
80	2,4-diF	CH	-	65.8	4.3	8.5	36-40
81	2-Cl,4-F	CH	-	66.3	4.4	8.5	103-105
82	3,5-diCH ₃	CH	-	62.8	4.1	8.1	
				64.5	4.2	8.3	
				75.0	6.2	8.7	69-70
				74.3	6.4	8.7	
83	3-CF ₃	CH	5-Cl	57.8	3.5	7.1	oil
				55.5	3.3	7.0	
84	3-CH ₂ CN, 5-CF ₃	N	-	60.0	3.7	14.0	65-67
				58.4	4.1	13.0	

Example 85Preparation of 2-[3-(4'-methylsulphonylphenoxy)benzoyl]-3-dimethylaminoacrylonitrile

A mixture of 2-[3-(4'-methylmercaptophenoxy)benzoyl]-3-dimethylaminoacrylonitrile (2g, from Example 59) and meta-chloroperoxybenzoic acid (1.2g) in methylene chloride (50ml) was stirred at room temperature overnight. The mixture was then washed with 10% aqueous sodium hydroxide, dried over magnesium sulphate and evaporated. The residual solid was chromatographed on a silica gel column using methanol as eluant to give the title compound (1.6g) as a white solid of m.p. 157-159 °C.

Analysis:

Calculated for $C_{19}H_{18}O_3N_2S$	C 64.4;	H 5.1;	N 7.9%
Found	C 64.0;	H 5.1;	N 7.9%

Example 86Preparation of 2-[3-(4'-methylsulphonylphenoxy)benzoyl]-3-dimethylaminoacrylonitrile

A mixture of 2-[3-(4'-methylmercaptophenoxy)benzoyl]-3-dimethylaminoacrylonitrile (2.5g, from Example 59) and meta-chloroperoxybenzoic acid (2.9g) in methylene chloride (50ml) was stirred at room temperature for 3 hours. The mixture was then filtered and the methylene chloride evaporated. The residual solid was chromatographed on a silica-gel column using 20% (v/v) diethyl ether-methylene chloride as eluant to give the title compound (1.3g) as a white solid of m.p. 45-50 °C.

Analysis:

Calculated for $C_{19}H_{18}O_4N_2S$	C 61.6;	H 4.9;	N 7.6%
Found	C 58.9;	H 5.0;	N 7.4%

Example 87Preparation of 2-[3-(4'-nitrophenoxy)benzoyl]-3-dimethylaminoacrylonitrile


A mixture of anhydrous potassium carbonate (1.3g), p-fluoronitrobenzene (1.5g) and 2-(3'-hydroxybenzoyl)-3-dimethylaminoacrylonitrile (2.1g) in dry dimethylformamide (50ml) was stirred and refluxed for 3 hours. After cooling, the mixture was poured into water (100ml) and the aqueous solution extracted with diethyl ether. After drying the diethyl ether extracts over anhydrous magnesium sulphate, the solvent was evaporated off and the residue chromatographed on a silica gel column using 10% diethyl ether-dichloromethane (v/v) as eluant to give the title compound (1.8g) as a pale yellow solid of m.p. 117-118 °C.

Analysis:

Calculated for $C_{18}H_{15}N_3O_4$	C 64.1;	H 4.4;	N 12.5%
Found	C 63.7;	H 4.5;	N 12.3%

The compounds of general formula I listed in Table 3 below, were prepared by methods similar to those described in Example 87 above.

Table 3



Example No.	Xn	Z	Ym	Elemental Analysis (%)		Melting Point (°C)
				C (Calc. Found)	H (Calc. Found)	
88	2-NO ₂ , 4-Cl	CH	-	58.2	3.8	114-115
89	2-NO ₂ , 4-Br	CH	-	58.3	3.9	oil
90	3-Cl, 5-CF ₃	N	-	52.0	3.4	146-147
91	5-CF ₃ , 6-Cl	N	-	52.6	3.7	50-52
				54.6	3.3	
				54.4	3.4	
				54.6	3.3	
				54.5	3.4	

Example 92 Herbicidal Activity

To evaluate their herbicidal activity, compounds according to the invention were tested using as representative range of plants: maize, Zea mays (Mz); rice, Oryza sativa (R); barnyard grass, Echinochloa crusgalli (BG); oat, Avena sativa (O); linseed, Linum usitatissimum (L); mustard, Sinapsis alba (M); sugar

beet, Beta vulgaris (SB) and soya bean, Glycine max (S).

The tests fall into two categories, pre-emergence and post-emergence. The pre-emergence tests involved spraying a liquid formulation of the compound onto the soil in which the seeds of the plant species mentioned above had recently been sown. The post-emergence tests involved two types of test, viz., soil drench and foliar spray tests. In the soil drench tests the soil in which the seedling plants of the above species were growing was drenched with a liquid formulation containing a compound of the invention, and in the foliar spray tests the seedling plants were sprayed with such a formulation.

The soil used in the tests was a prepared horticultural loam.

The formulations used in the tests were prepared from solutions of the test compounds in acetone containing 0.4% by weight of an alkylphenol/ethylene oxide condensate available under the trade mark TRITON X-155. These acetone solutions were diluted with water and the resulting formulations applied at dosage levels corresponding to 5 kg or 1 kg of active material per hectare in a volume equivalent to 600 litres per hectare in the soil spray and foliar spray test, and at a dosage of level equivalent to 10 kilograms of active material per hectare in a volume equivalent to approximately 3,000 litres per hectare in the soil drench tests.

In the pre-emergence tests untreated sown soil and in the post-emergence tests untreated soil bearing seedling plants were used as controls.

The herbicidal effects of the test compounds were assessed visually twelve days after spraying the foliage and the soil, and thirteen days after drenching the soil and were recorded on a 0-9 scale. A rating 0 indicates growth as untreated control, a rating 9 indicates death. An increase of 1 unit on the linear scale approximates to a 10% increase in the level of effect.

The results of the tests are set out in Table 4 below. There are included in Table 4 for comparison purposes the following acrylonitrile derivatives described in DE-A 2330913:

Compound A	2-benzoyl-3-dimethylaminoacrylonitrile
Compound B	2-(3-methoxybenzoyl)-3-dimethylaminoacrylonitrile

Table 4

Compound of Ex. No.		Soil drench 10/kg/ha						Dosage kg/ha						Foliar spray						Pre-emergence						
		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S	
Compound A		6	3	6	5	5	6	5	7	5	3	1	5	2	4	5	6	5	6	4	8	6	5	6	7	5
										1	0	0	1	0	3	3	3	3	2	0	3	1	2	3	6	0
Compound B		6	5	6	6	6	8	8	8	5	4	0	5	0	5	8	7	5	7	6	9	7	7	9	9	6
										1	0	0	1	0	3	5	6	3	4	0	1	2	1	4	4	2

Table 4 (continued)

Compound of Ex. No.	Soil drench 10/kg/ha						Dosage kg/ha						Foliar spray						Pre-emergence					
	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
45	4	3	6	6	2	3	4	0	5	7	3	9	7	8	9	9	6	2	3	4	5	4	8	0
									1	4	2	7	5	7	9	9	5	1	1	2	4	3	6	0
46	7	7	7	6	6	6	7	1	5	5	4	8	3	6	7	8	6	5	4	9	5	5	9	1
									1	3	2	7	2	6	7	7	5	5	4	8	4	3	8	1
48	8	8	8	9	8	9	9	7	5	5	4	8	8	8	8	9	7	8	6	8	7	8	9	3
									1	2	0	6	3	6	7	8	6	6	3	7	6	3	7	0
49	4	4	6	6	1	3	4	0	5	6	3	8	5	9	8	8	7	2	1	8	5	8	9	7
									1	4	1	7	3	7	8	8	6	1	0	7	2	5	9	3
50	5	4	6	5	2	4	4	2	5	6	5	8	7	6	9	9	6	3	4	5	5	3	8	2
									1	3	2	7	4	6	9	9	6	1	1	3	3	1	6	7

Table 4 (continued)

Compound of Ex. No.		Soil drench 10/kg/ha						Dosage kg/ha						Foliar spray						Pre-emergence						
		Mz	R	BG	O	L	M	SB	S	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S			
51		4	5	6	5	4	3	3	3	5	6	4	8	6	7	8	7	7	1	4	2	4	3	6	1	0
										1	5	3	8	5	7	7	6	7	0	0	0	3	1	4	0	0
52		3	1	5	4	2	2	3	0	5	4	3	8	7	8	9	9	7	0	0	2	1	3	5	6	0
										1	3	1	7	5	7	9	8	5	0	0	0	0	1	4	3	0
53		5	5	5	6	5	4	4	2	5	6	4	8	7	7	8	8	7	4	5	5	5	2	5	4	0
										1	4	3	7	4	6	8	7	7	1	3	4	3	2	3	2	0
54		2	0	5	5	0	0	0	0	5	5	3	8	8	9	9	7	0	0	0	0	2	3	4	0	
										1	4	2	7	4	7	9	9	6	0	0	0	1	3	3	0	
55		5	0	4	4	0	3	0	2	5	6	1	8	4	7	9	7	5	0	0	5	2	4	8	4	0
										1	4	1	6	2	6	8	5	4	0	0	5	1	3	7	3	0

Table 4 (continued)

Compound of Ex. No.		Soil drench 10/kg/ha						Dosage		Foliar spray						Pre-emergence										
		Mz	R	BG	O	L	M	SB	S	kg/ha	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
56		4	4	6	5	1	0	3	2	5	5	2	7	3	7	7	8	6	0	0	3	3	4	6	4	2
										1	4	2	6	3	5	6	6	5	0	0	1	1	2	6	2	1
57		7	5	8	7	4	9	9	4	5	4	0	9	5	7	9	9	6	6	4	8	7	3	8	9	0
										1	3	0	9	4	6	9	8	5	5	2	7	4	1	4	8	0
58		5	4	6	5	4	5	2	0	5	5	4	8	5	7	9	9	6	4	3	8	5	7	8	8	4
										1	4	2	6	3	6	8	9	5	2	3	7	3	4	8	5	0
59		2	0	4	6	1	4	6	0	5	3	2	7	3	5	8	6	4	0	0	2	3	4	7	7	0
										1	2	0	6	2	4	6	6	2	0	0	1	1	2	7	6	0
85		4	1	2	5	0	6	4	0	5	3	2	6	4	4	7	7	4	0	0	3	4	3	8	8	0
										1	1	1	6	3	4	7	7	3	0	0	1	1	2	6	6	0

Table 4 (continued)

Compound																									
of		Soil drench 10/kg/ha						Dosage			Foliar spray						Pre-emergence								
Ex. No.	Mz	R	BG	O	L	M	SB	S	kg/ha	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
86	5	2	4	5	0	4	4	0	5	2	0	7	3	4	7	7	3	3	0	3	3	2	6	5	0
									1	0	0	3	0	2	6	5	2	0	0	0	0	1	3	3	0
60	2	4	4	4	2	2	4	2	5	4	4	7	5	6	9	8	5	0	0	5	2	4	7	8	2
									1	3	3	7	4	6	8	7	4	0	0	4	2	3	7	8	1
61	6	5	7	7	5	8	9	4	5	7	3	8	6	7	9	9	7	5	5	8	7	5	8	9	2
									1	4	1	7	4	5	9	9	6	4	3	7	5	3	7	8	0
62	6	5	6	6	4	6	9	4	5	7	2	8	7	7	9	9	7	4	4	4	7	6	7	8	1
									1	4	1	7	4	5	8	8	6	2	2	6	5	2	4	8	0
63	5	4	5	5	3	4	5	2	5	4	1	6	5	6	8	8	6	3	2	5	4	4	7	8	2
									1	4	0	4	3	4	7	8	5	1	0	4	2	2	5	8	0

Table 4 (continued)

Compound of Ex. No.		Soil drench 10/kg/ha						Dosage kg/ha						Foliar spray						Pre-emergence						
		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S	
64		5	4	6	5	5	9	4	4	5	4	2	8	5	7	9	9	6	4	4	6	7	6	8	9	1
										1	3	1	6	2	6	8	9	6	4	2	6	3	2	7	5	0
65		6	6	7	6	6	9	9	3	5	4	2	9	5	7	9	9	6	6	6	8	7	7	8	9	2
										1	2	0	7	4	5	9	9	5	5	4	6	5	4	6	8	0
66		6	2	6	7	4	5	9	2	5	5	1	6	6	7	8	9	6	5	2	7	7	4	4	9	2
										1	2	0	3	1	3	8	8	4	4	0	5	2	1	1	8	0
67		5	4	6	5	5	7	4	0	5	4	0	5	2	6	7	5	5	4	3	5	5	3	5	7	0
										1	2	0	1	0	4	5	4	5	3	0	3	2	1	3	4	0
68		6	6	7	7	5	7	7	3	5	5	3	8	5	6	9	8	6	4	4	7	5	4	8	8	0
										1	2	1	7	3	5	7	5	4	1	1	4	4	2	7	7	0

5
10
15
20
25
30
35
40
45
50
55

Compound of Ex. No.		Soil drench 10/kg/ha					Dosage kg/ha	Foliar spray					Pre-emergence													
		Mz	R	BG	O	L		M	SB	S	R	BG	O	L	M	SB	S									
69		2	0	4	4	3	4	6	2	5	5	2	9	5	8	9	9	8	0	0	1	2	5	5	0	
										1	3	1	6	3	8	9	9	7	0	0	0	1	2	3	0	
70		7	4	7	7	4	5	3	2	5	5	2	8	5	7	9	8	7	0	0	3	3	5	4	0	
										1	3	1	6	3	6	8	8	6	0	0	3	1	0	4	1	0
88		5	4	5	5	2	4	5	0	5	6	2	7	5	6	8	6	6	0	0	2	1	3	5	8	0
										1	4	1	5	3	3	7	5	6	0	0	1	0	1	4	8	0
87		6	4	6	6	4	4	2	0	5	6	1	6	5	6	7	8	5	4	3	6	5	3	6	8	0
										1	5	0	5	3	4	6	5	5	1	1	4	4	1	4	4	0
89		5	5	6	5	1	3	3	0	5	4	2	6	2	4	7	7	5	0	0	2	0	2	4	4	0
										1	4	2	5	2	3	7	7	4	0	0	0	0	0	3	3	0

Compound

Ex. No.	of	Soil drench 10/kg/ha						Dosage			Foliar spray						Pre-emergence									
		Mz	R	BG	O	L	M	SB	S	kg/ha	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
71		6	5	7	6	2	5	5	0	5	4	3	8	5	7	9	8	7	2	2	4	5	4	7	9	0
									1	4	4	2	6	3	6	9	6	6	1	0	3	4	3	7	8	0
72		0	0	2	3	0	3	0	0	5	4	3	6	6	7	9	5	6	0	0	2	1	3	7	5	0
									1	3	3	1	2	3	5	9	4	3	0	0	1	0	1	4	0	0
73		0	0	0	0	0	0	0	0	5	4	1	6	3	5	8	5	6	0	0	2	0	2	3	5	0
									1	2	2	1	3	2	4	7	5	5	0	0	1	0	0	2	3	0
74		5	3	6	4	0	5	6	0	5	4	3	7	5	6	9	8	7	0	0	4	2	4	5	9	3
									1	2	2	1	5	4	6	9	8	5	0	0	4	1	2	4	9	2
75		5	4	6	6	2	5	4	2	5	7	3	8	7	8	9	9	8	3	2	6	5	4	8	9	3
									1	5	5	2	6	4	7	9	9	7	1	0	3	3	3	5	8	2

Table 4 (continued)

Compound of																									
Ex. No.	Soil drench 10/kg/ha						Dosage						Foliar spray						Pre-emergence						
	Mz	R	BG	O	L	M	SB	S	kg/ha	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
76	6	4	7	6	4	4	5	0	5	5	2	8	3	7	9	8	6	4	4	6	5	5	8	8	1
									1	3	1	7	1	6	9	7	6	4	3	5	3	3	4	8	0
77	4	3	5	4	0	3	4	0	5	4	3	7	4	7	8	8	6	0	0	3	4	4	8	8	0
									1	3	2	5	3	6	8	7	5	0	0	2	1	3	8	6	0
78	5	5	7	6	3	4	6	0	5	5	3	8	4	8	9	8	7	1	2	5	6	5	8	8	0
									1	3	0	6	3	6	8	8	5	0	0	4	4	3	8	8	0
79	5	6	7	7	5	8	6	2	5	5	3	8	4	7	9	8	6	3	4	7	6	5	8	9	0
									1	2	0	7	3	6	9	7	6	2	2	5	4	4	8	8	0
90	7	5	7	8	5	8	8	3	5	5	3	8	5	7	9	8	7	5	5	7	7	6	9	5	0
									1	4	2	6	4	6	9	8	7	4	3	4	6	3	9	3	0

Table 4 (continued)

Compound of		Soil drench 10/kg/ha			Dosage			Foliar spray			Pre-emergence														
Ex. No.	Mz	R	BC	O	L	M	SB	S	kg/ha	Mz	R	BC	O	L	M	SB	S	Mz	R	BC	O	L	M	SB	S
91	5	3	5	4	1	5	5	0	5	5	1	8	4	8	9	9	5	0	0	5	3	2	7	8	2
									1	3	0	5	1	7	9	8	4	0	0	2	0	1	4	4	0
80	7	5	7	7	5	5	8	0	5	7	4	8	3	7	9	9	7	6	6	5	5	2	4	8	0
									1	5	3	6	0	6	8	6	6	5	4	4	3	1	2	4	0
81	7	5	7	7	4	4	8	2	5	5	3	8	6	8	9	9	7	4	3	7	5	5	7	9	2
									1	4	1	7	4	7	8	8	7	3	0	5	2	3	7	8	0
82	5	4	6	6	2	2	4	2	5	4	4	8	6	7	8	9	7	0	1	5	4	4	7	8	1
									1	3	3	5	4	6	8	8	5	0	0	4	3	4	7	6	1
83	0	0	0	0	0	0	0	0	5	5	2	3	3	7	9	8	3	1	0	1	0	0	2	3	1
									1	2	0	3	1	6	7	3	2	0	0	0	0	0	0	0	0

Table 4 (continued)


Compound of Ex. No.	Soil dranch 10/kg/ha						Dosage		Foliar spray						Pre-emergence										
	Mz	R	BG	O	L	M	SB	S	kg/ha	Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
84	7	5	6	7	2	8	7	0	5	4	0	6	3	5	8	8	8	2	0	4	2	3	6	9	3
									1	3	0	2	1	2	7	7	4	0	0	0	0	2	3	2	0



Claims


Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. A compound of the general formula I




in which R^1 and R^2 each independently represents an alkyl group, Z represents a nitrogen atom or a group CH which is unsubstituted or substituted by a substituent X , m is 0 or an integer from 1 to 4, the or each Y independently represents a halogen atom, n is 0 or an integer from 1 to 5, and the or each X independently represents a halogen atom or a group selected from alkyl, haloalkyl, cyanoalkyl, alkenyl, alkynyl, alkenyloxy, alkynyloxy, alkoxy, alkylthio, alkenylthio, alkynylthio, cyano, nitro, alkylsulphonyl, alkylsulphinyl and sulphonamido.

2. A compound as claimed in claim 1, wherein R^1 and R^2 are the same and each represents a methyl group.
3. A compound as claimed in claim 1 or claim 2, wherein Z represents an unsubstituted group CH .
4. A compound as claimed in any one of claims 1 to 3, wherein m is 1 and Y represents a chlorine atom, or m is 0.
5. A compound as claimed in any one of the preceding claims wherein n is 1 or 2.
6. A compound as claimed in claim 5, wherein the or each X is independently selected from halogen atoms, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{1-6} cyanoalkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylthio groups, C_{1-6} alkylsulphinyl groups, C_{1-6} alkylsulphonyl groups, cyano groups and nitro groups.
7. A compound as claimed in claim 6, wherein the or each X is independently selected from fluorine, chlorine, and bromine atoms and C_{1-4} alkyl, trifluoromethyl, cyanomethyl, methoxy, methylthio, methylsulphinyl, methylsulphonyl, cyano and nitro groups.
8. A process for the preparation of a compound of general formula I as claimed in claim 1, which comprises reacting a compound of general formula II:




in which X , Y , Z , n and m are as defined in claim 1, and W represents a halogen atom or a cyanomethyl group, with, in the case where W is halogen, a dialkylaminoacrylonitrile under basic conditions, or, in the case where W is cyanomethyl, either with a dialkylformamide dialkyl acetal, or with a trialkylorthoformate followed by reaction with a dialkylamine or reacting a compound of general formula III



(III)

wherein $\underline{n}$, X and Z are as defined in claim 1 and L represents a leaving group with a compound of general formula IV



(IV)

wherein $\underline{m}$, Y, R^1 and R^2 are as defined in claim 1, under basic conditions, and, if desired or required, converting a resulting compound of general formula I into another compound of general formula I.

9. A process as claimed in claim 8, wherein W is cyanomethyl and the compound of general formula II is reacted with a dialkylformamide dimethyl acetal in an inert solvent at a temperature in the range of from 0 to 50 °C.

10. A herbicidal composition which comprises a compound as claimed in any one of claims 1 to 7 together with at least one carrier.


11. A composition as claimed in claim 10, comprising at least two carriers, at least one of which is a surface-active agent.

12. A method of combating undesired plant growth at a locus, which comprises treating the locus with a compound as claimed in any one of claims 1 to 7, or a composition as claimed in claim 10 or claim 11.

13. The use of a compound as claimed in any one of claims 1 to 7, or a composition as claimed in claim 10 or claim 11, as a herbicide.

Claims for the following Contracting State : ES

1. A method of combating undesired plant growth at a locus, which comprises treating the locus with a compound of the general formula I:




(I)


in which R^1 and R^2 each independently represents an alkyl group, Z represents a nitrogen atom or a group CH which is unsubstituted or substituted by a substituent X, $\underline{m}$ is 0 or an integer from 1 to 4, the or each Y independently represents a halogen atom, $\underline{n}$ is 0 or an integer from 1 to 5, and the or each X independently represents a halogen atom or a group selected from alkyl, haloalkyl, cyanoalkyl, alkenyl, alkynyl, alkenyloxy, alkynyloxy, alkoxy, alkylthio, alkenylthio, alkynylthio, cyano, nitro, alkylsulphonyl, alkylsulphinyl and sulphonamido.




2. A method as claimed in Claim 1, wherein R^1 and R^2 are the same and each represents a methyl group.
3. A method as claimed in Claim 1 or Claim 2, wherein Z represents an unsubstituted group CH.
- 5 4. A method as claimed in any one of Claims 1-3, wherein $\underline{m}$ is 1 and Y represents a chlorine atom, or $\underline{m}$ is 0.
5. A method as claimed in any one of the preceding claims, wherein $\underline{n}$ is 1 or 2.
- 10 6. A method as claimed in Claim 5, wherein the or each X is independently selected from halogen atoms, C_{1-6} alkyl groups, C_{1-6} haloalkyl groups, C_{1-6} cyanoalkyl groups, C_{1-6} alkoxy groups, C_{1-6} alkylthio groups, C_{1-6} alkylsulphinyl groups, C_{1-6} alkylsulphonyl groups, cyano groups and nitro groups.
- 15 7. A method as claimed in Claim 6, wherein the or each X is independently selected from fluorine, chlorine, and bromine atoms and C_{1-4} alkyl, trifluoromethyl, cyanomethyl, methoxy, methylthio, methylsulphinyl, methylsulphonyl, cyano and nitro groups.
8. A method as claimed in any one of the preceding claims, wherein said compound of general formula I is applied in association with at least one carrier.
- 20 9. A method as claimed in Claim 8, wherein said compound of general formula I is applied in association with at least two carriers, at least one of which is a surface-active agent.
10. A process for the preparation of a compound of general formula I as defined in Claim 1, which comprises reacting a compound of general formula II:
- 25



in which X, Y, Z, $\underline{n}$ and $\underline{m}$ are as defined in Claim 1, and W represents a halogen atom or a cyanomethyl group, with, in the case where W is halogen, a dialkylamino-acrylonitrile under basic conditions, or, in the case where W is cyanomethyl, either with a dialkylformamide dialkyl acetal, or with a trialkylorthoformate followed by reaction with a dialkylamine; or reacting a compound of general formula III:



wherein $\underline{n}$, X and Z are as defined in Claim 1 and L represents a leaving group with a compound of general formula IV:



10 wherein m , Y , R^1 and R^2 are as defined in Claim 1, under basic conditions, and, if desired or required, converting a resulting compound of general formula I into another compound of general formula I.

11. A process as claimed in Claim 10, wherein W is cyanomethyl and the compound of general formula II is reacted with a dialkylformamide dimethyl acetal in an inert solvent at a temperature in the range of
15 from 0 to 50 °C.

12. A herbicidal composition which comprises a compound of general formula I as defined in any one of Claims 1-7 together with at least one carrier.


20 13. A composition according to Claim 12, comprising at least two carriers, at least one of which is a surface-active agent.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

25

1. Eine Verbindung mit der allgemeinen Formel I



30 worin R^1 und R^2 jeweils unabhängig voneinander eine Alkylgruppe bedeuten, Z ein Stickstoffatom oder eine Gruppe CH bedeutet, die unsubstituiert oder durch einen Substituenten X substituiert ist, m den Wert 0 oder eine ganze Zahl von 1 bis 4 hat, der oder jeder Rest Y unabhängig voneinander ein Halogenatom darstellt, n den Wert 0 oder eine ganze Zahl von 1 bis 5 aufweist und der oder jeder Rest X unabhängig voneinander ein Halogenatom oder eine unter Alkyl, Halogenalkyl, Cyanoalkyl, Alkenyl, Alkynyl, Alkenyloxy, Alkinyloxy, Alkoxy, Alkylthio, Alkenylthio, Alkynylthio, Cyano, Nitro, Alkylsulfonyl, Alkylsulfinyl und Sulfonamido ausgewählte Gruppe darstellt.

45 2. Verbindung nach Anspruch 1, worin R^1 und R^2 die gleiche Bedeutung aufweisen und jeweils eine Methylgruppe darstellen.

3. Verbindung nach Anspruch 1 oder 2, worin Z eine unsubstituierte CH -Gruppe bedeutet.


50 4. Verbindung nach einem der Ansprüche 1 bis 3, worin m den Wert 1 aufweist und Y ein Chloratom darstellt, oder worin m den Wert 0 aufweist.

5. Verbindung nach einem der vorstehenden Ansprüche, worin n den Wert 1 oder 2 hat.

55 6. Verbindung nach Anspruch 5, worin der oder jeder Rest X unabhängig voneinander ausgewählt ist unter Halogenatomen, C_1 - C_6 -Alkylgruppen, C_1 - C_6 -Halogenalkylgruppen, C_1 - C_6 -Cyanoalkylgruppen, C_1 - C_6 -Alkoxygruppen, C_1 - C_6 -Alkylthiogruppen, C_1 - C_6 -Alkylsulfinylgruppen, C_1 - C_6 -Alkylsulfonylgruppen, Cyanogruppen und Nitrogruppen.



7. Verbindung nach Anspruch 6, worin der oder jeder Rest X unabhängig voneinander ausgewählt ist unter Fluor-, Chlor- und Bromatomen und C₁₋₄-Alkyl-, Trifluormethyl-, Cyanomethyl-, Methoxy-, Methylthio-, Methylsulfinyl-, Methylsulfonyl-, Cyano- und Nitrogruppen.
8. Verfahren zur Herstellung einer Verbindung der allgemeinen Formel I, wie in Anspruch 1 beansprucht, welches ein Umsetzen einer Verbindung der allgemeinen Formel II:




worin X, Y, Z, n und m wie in Anspruch 1 definiert sind und W ein Halogenatom oder eine Cyanomethylgruppe darstellt, für den Fall, daß W ein Halogen bedeutet, mit einem Dialkylaminoacrylnitril unter basischen Bedingungen, oder, für den Fall, daß W Cyanomethyl bedeutet, entweder mit einem Dialkylformamid-dialkylacetal oder mit einem Trialkylorthoformiat, gefolgt von einer Umsetzung mit einem Dialkylamin,

oder ein Umsetzen einer Verbindung der allgemeinen Formel III:



worin n, X und Z wie in Anspruch 1 definiert sind und L eine Leaving-Gruppe darstellt, mit einer Verbindung der allgemeinen Formel IV:



worin m, Y, R¹ und R² wie in Anspruch 1 definiert sind, unter basischen Bedingungen umfaßt, und, falls erwünscht oder benötigt, ein Überführen einer erhaltenen Verbindung der allgemeinen Formel I in eine andere Verbindung der allgemeinen Formel I umfaßt.

9. Verfahren nach Anspruch 8, worin W für Cyanomethyl steht und die Verbindung der allgemeinen Formel II mit einem Dialkylformamid-dimethylacetal in einem inerten Lösungsmittel bei einer Temperatur im Bereich von 0 bis 50 °C umgesetzt wird.
10. Herbizide Zusammensetzung, welche eine Verbindung, wie in einem der Ansprüche 1 bis 7 beansprucht, zusammen mit wenigstens einem Träger umfaßt.
11. Zusammensetzung nach Anspruch 10, umfassend wenigstens zwei Träger, von denen wenigstens einer ein grenzflächenaktives Mittel ist.
12. Verfahren zur Bekämpfung von unerwünschtem Pflanzenwuchs an einem Ort, welches ein Behandeln des Ortes mit einer Verbindung, wie in einem der Ansprüche 1 bis 7 beansprucht, oder mit einer Zusammensetzung, wie in Anspruch 10 oder Anspruch 11 beansprucht, umfaßt.



13. Verwendung einer Verbindung, wie in einem der Ansprüche 1 bis 7 beansprucht, oder einer Zusammensetzung, wie in Anspruch 10 oder Anspruch 11 beansprucht, als ein Herbizid.


Patentansprüche für folgenden Vertragsstaat : ES

5

1. Verfahren zur Bekämpfung von unerwünschtem Pflanzenwuchs an einem Ort, welches ein Behandeln des Ortes mit einer Verbindung der allgemeinen Formel I:

10

15



20

25

umfaßt, worin R^1 und R^2 jeweils unabhängig voneinander eine Alkylgruppe bedeuten, Z ein Stickstoffatom oder eine Gruppe CH bedeutet, die unsubstituiert oder durch einen Substituenten X substituiert ist, m den Wert 0 oder eine ganze Zahl von 1 bis 4 hat, der oder jeder Rest Y unabhängig voneinander ein Halogenatom darstellt, n den Wert 0 oder eine ganze Zahl von 1 bis 5 aufweist und der oder jeder Rest X unabhängig voneinander ein Halogenatom oder eine unter Alkyl, Halogenalkyl, Cyanoalkyl, Alkenyl, Alkynyl, Alkenyloxy, Alkynyloxy, Alkoxy, Alkylthio, Alkenylthio, Alkynylthio, Cyano, Nitro, Alkylsulfonyl, Alkylsulfinyl und Sulfonamido ausgewählte Gruppe darstellt.

30

2. Verfahren nach Anspruch 1, worin R^1 und R^2 die gleiche Bedeutung aufweisen und jeweils eine Methylgruppe darstellen.

3. Verfahren nach Anspruch 1 oder 2, worin Z eine unsubstituierte CH-Gruppe bedeutet.

4. Verfahren nach einem der Ansprüche 1 bis 3, worin m den Wert 1 aufweist und Y ein Chloratom darstellt, oder worin m den Wert 0 aufweist.

35

5. Verfahren nach einem der vorstehenden Ansprüche, worin n den Wert 1 oder 2 hat.

40

6. Verfahren nach Anspruch 5, worin der oder jeder Rest X unabhängig voneinander ausgewählt ist unter Halogenatomen, C_{1-6} -Alkylgruppen, C_{1-6} -Halogenalkylgruppen, C_{1-6} -Cyanoalkylgruppen, C_{1-6} -Alkoxygruppen, C_{1-6} -Alkylthiogruppen, C_{1-6} -Alkylsulfinylgruppen, C_{1-6} -Alkylsulfonylgruppen, Cyanogruppen und Nitrogruppen.

45

7. Verfahren nach Anspruch 6, worin der oder jeder Rest X unabhängig voneinander ausgewählt ist unter Fluor-, Chlor- und Bromatomen und C_{1-4} -Alkyl-, Trifluormethyl-, Cyanomethyl-, Methoxy-, Methylthio-, Methylsulfinyl-, Methylsulfonyl-, Cyano- und Nitrogruppen.


50

8. Verfahren nach einem der vorstehenden Ansprüche, worin die Verbindung der allgemeinen Formel I in Assoziation mit wenigstens einem Träger aufgebracht wird.

9. Verfahren nach Anspruch 8, worin die Verbindung der allgemeinen Formel I in Assoziation mit wenigstens zwei Trägern aufgebracht wird, von denen wenigstens einer ein grenzflächenaktives Mittel ist.

55

10. Verfahren zur Herstellung einer Verbindung der allgemeinen Formel I, wie in Anspruch 1 definiert, welches ein Umsetzen einer Verbindung der allgemeinen Formel II:




10 worin X, Y, Z, n und m wie in Anspruch 1 definiert sind und W ein Halogenatom oder eine Cyanomethylgruppe darstellt, für den Fall, daß W ein Halogen bedeutet, mit einem Dialkylaminoacrylnitril unter basischen Bedingungen, oder, für den Fall, daß W Cyanomethyl bedeutet, entweder mit einem Dialkylformamid-dialkylacetal oder mit einem Trialkylorthoformiat, gefolgt von einer Umsetzung mit einem Dialkylamin,

15 oder ein Umsetzen einer Verbindung der allgemeinen Formel III:



25 worin n, X und Z wie in Anspruch 1 definiert sind und L eine Leaving-Gruppe darstellt, mit einer Verbindung der allgemeinen Formel IV:



35 worin m, Y, R¹ und R² wie in Anspruch 1 definiert sind, unter basischen Bedingungen umfaßt, und, falls erwünscht oder benötigt, ein Überführen einer erhaltenen Verbindung der allgemeinen Formel I in eine andere Verbindung der allgemeinen Formel I umfaßt.


- 40 11. Verfahren nach Anspruch 10, worin W für Cyanomethyl steht und die Verbindung der allgemeinen Formel II mit einem Dialkylformamid-dimethylacetal in einem inerten Lösungsmittel bei einer Temperatur im Bereich von 0 bis 50 °C umgesetzt wird.
- 45 12. Herbizide Zusammensetzung, welche eine Verbindung, wie in einem der Ansprüche 1 bis 7 beansprucht, zusammen mit wenigstens einem Träger umfaßt.
13. Zusammensetzung nach Anspruch 12, umfassend wenigstens zwei Träger, von denen wenigstens einer ein grenzflächenaktives Mittel ist.




Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. Un composé de la formule générale I



- 15 dans laquelle R^1 et R^2 représentent chacun, de façon indépendante, un groupe alkyle, Z représente un atome d'azote ou un groupe CH qui est non substitué ou substitué par un substituant X, m est 0 ou un nombre entier de 1 à 4, le ou chaque Y représente indépendamment un atome d'halogène, n est zéro ou un nombre entier de 1 à 5 et le X ou chaque X représente indépendamment un atome d'halogène ou un groupe choisi parmi les groupes alkyle, haloalkyle, cyanoalkyle, alcényle, alcynyle, alcoxy, alcényloxy, alcynyloxy, alkylthio, alcénylthio, alcynylthio, cyano, nitro, alkylsulfonyle, alkylsulfinyle et sulfonamido.
2. Un composé tel que revendiqué dans la revendication 1, dans lequel R^1 et R^2 sont identiques et représentent chacun un groupe méthyle.
3. Un composé tel que revendiqué dans la revendication 1 ou la revendication 2, dans lequel Z représente un groupe non-substitué CH.
4. Un composé tel que revendiqué dans l'une quelconque des revendications 1 ou 3, dans lequel m est 1 et Y représente un atome de chlore, ou bien m est 0.
5. Un composé tel que revendiqué dans l'une quelconque des revendications précédentes, dans lequel n est 1 ou 2.
- 35 6. Un composé tel que revendiqué dans la revendication 5, dans lequel le ou chaque X est choisi, de façon indépendante, parmi les atomes d'halogène, les groupes alkyle en C_1 à C_6 , les groupes haloalkyle en C_1 à C_6 , les groupes cyanoalkyle en C_1 à C_6 , les groupes alcoxy en C_1 à C_6 , les groupes alkylthio en C_1 à C_6 , les groupes alkylsulfinyle en C_1 à C_6 , les groupes alkylsulfonyle en C_1 à C_6 , les groupes cyano et les groupes nitro.
- 40 7. Un composé tel que revendiqué dans la revendication 6, dans lequel le ou chaque X est choisi de façon indépendante, parmi les atomes de fluor, de chlore et de brome, ainsi que dans les groupes alkyl en C_1 à C_4 , trifluorométhyle, cyanométhyle, méthoxy, méthylthio, méthylsulfinyle, méthylsulfonyle, cyano et nitro.
- 45 8. Un procédé pour la préparation d'un composé de la formule générale I, tel que revendiqué dans la revendication 1, procédé selon lequel on fait réagir un composé de la formule générale II:




dans laquelle X, Y, Z, n et m sont tels que définis dans la revendication 1 et W représente un atome




d'halogène ou un groupe cyanométhyle, avec, dans le cas où W est un halogène, un dialkylamino acrylonitrile sous des conditions basiques, ou bien, dans le cas où W est du cyanométhyle, soit avec un dialkyl formamide dialkyl acétal, soit avec un orthoformiate de trialkyle suivi par une réaction avec une dialkylamine

ou bien

on fait réagir un composé de la formule générale III



dans laquelle n , X et Z sont tels que définis dans la revendication 1 et L représente un groupe partant avec un composé de la formule générale IV




dans laquelle m , Y, R^1 et R^2 sont tels que définis dans la revendication 1, sous les conditions basiques, et, si on le désire ou si cela est nécessaire, on convertit un composé résultant de la formule générale I en un autre composé de la formule générale I.

9. Un procédé tel que revendiqué dans la revendication 8, dans lequel W est du cyanométhyle et le composé de la formule générale II est mis à réagir avec un dialkyl formamide diméthyl acétal dans un solvant inerte avec une température dans la gamme de 0 à 50 °C.
10. Une composition herbicide qui comporte un composé tel que revendiqué dans l'une quelconque des revendications 1 à 7, conjointement avec au moins un support.
11. Une composition telle que revendiquée dans la revendication 10, comportant au moins deux supports dont l'un est un agent tensioactif.
12. Un procédé pour combattre la croissance des plantes non désirée en un endroit, procédé selon lequel on traite cet endroit avec un composé tel que revendiqué dans l'une quelconque des revendications 1 à 7, ou bien une composition telle que revendiquée dans la revendication 10 ou la revendication 11.
13. L'utilisation d'un composé tel que revendiqué dans l'une quelconque des revendications 1 à 7 ou bien d'une composition telle que revendiquée dans la revendication 10 ou la revendication 11, en tant qu'herbicide.


Revendications pour l'Etat contractant suivant : ES

1. Un procédé pour combattre la croissance des plantes indésirables en un lieu, procédé selon lequel on traite ce lieu avec un composé de la formule générale I




10 dans laquelle R¹ et R² représentent chacun, de façon indépendante, un groupe alkyle, Z représente un atome d'azote ou un groupe CH qui est non substitué ou substitué par un substituant X, m est 0 ou un nombre entier de 1 à 4, le ou chaque Y représente indépendamment un atome d'halogène, n est zéro ou un nombre entier de 1 à 5 et le X ou chaque X représente indépendamment un atome d'halogène ou un groupe choisi parmi les groupes alkyle, haloalkyle, cyanoalkyle, alcényle, alcynyle, alcoxy, alcényloxy, alcynyloxy, alkylthio, alcénylthio, alcynylthio, cyano, nitro, alkylsulfonyl, alkylsulfinyle et sulfonamido.

2. Un procédé tel que revendiqué dans la revendication 1, dans lequel R¹ et R² sont identiques et représentent chacun un groupe méthyle.
- 20 3. Un procédé tel que revendiqué dans la revendication 1 ou la revendication 2, dans lequel Z représente un groupe non-substitué CH.
4. Un procédé tel que revendiqué dans l'une quelconque des revendications 1 ou 3, dans lequel m est 1 et Y représente un atome de chlore, ou bien m est 0.
- 25 5. Un procédé tel que revendiqué dans l'une quelconque des revendications précédentes, dans lequel n est 1 ou 2.
- 30 6. Un procédé tel que revendiqué dans la revendication 5, dans lequel le ou chaque X est choisi, de façon indépendante, parmi les atomes d'halogène, les groupes alkyle en C₁ à C₆, les groupes haloalkyle en C₁ à C₆, les groupes cyanoalkyle en C₁ à C₆, les groupes alcoxy en C₁ à C₆, les groupes alkylthio en C₁ à C₆, les groupes alkylsulfinyle en C₁ à C₆, les groupes alkylsulfonyl en C₁ à C₆, les groupes cyano et les groupes nitro.
- 35 7. Un procédé tel que revendiqué dans la revendication 6, dans lequel le ou chaque X est choisi de façon indépendante, parmi les atomes de fluor, de chlore et de brome, ainsi que dans les groupes alkyle en C₁ à C₄, trifluorométhyle, cyanométhyle, méthoxy, méthylthio, méthylsulfinyle, méthylsulfonyl, cyano et nitro.
- 40 8. Un procédé tel que revendiqué dans l'une quelconque des revendications précédentes dans lequel ledit composé de la formule générale I est appliqué en association avec au moins un support.
9. Un procédé tel que revendiqué dans la revendication 8, dans lequel ledit composé de la formule générale I est appliqué en association avec au moins deux supports, dont au moins l'un est un agent tensioactif.
- 45 10. Un procédé pour la préparation d'un composé de la formule générale I tel que défini dans la revendication 1, dans lequel on fait réagir un composé de la formule générale II:






dans laquelle X, Y, Z, n et m sont tels que définis dans la revendication 1 et W représente un atome d'halogène ou un groupe cyanométhyle, avec, dans le cas où W est un halogène, un dialkylamino acrylonitrile sous des conditions basiques, ou bien, dans le cas où W est du cyanométhyle, soit avec un dialkylformamide dialkyl acétal, soit avec un orthoformiate de trialkyle suivi par une réaction avec une dialkylamine
ou bien
on fait réagir un composé de la formule générale III



dans laquelle n, X et Z sont tels que définis dans la revendication 1 et L représente un groupe partant avec un composé de la formule générale IV



dans laquelle m, Y, R¹ et R² sont tels que définis dans la revendication 1, sous les conditions basiques, et, si on le désire ou si cela est nécessaire, on convertit un composé résultant de la formule générale I en un autre composé de la formule générale I.

11. Un procédé tel que revendiqué dans la revendication 10, dans lequel W est du cyanométhyle et le composé de la formule générale II est mis à réagir avec un dialkyl formamide diméthyl acétal dans un solvant inerte avec une température dans la gamme de 0 à 50 °C.
12. Une composition herbicide qui comporte un composé de la formule générale I tel que défini dans l'une quelconque des revendications 1 à 7, conjointement avec au moins un support.
13. Une composition selon la revendication 12, comportant au moins deux supports dont l'un est un agent tensioactif.