

DESCRIPTION

METHOD OF IMPROVING SWEETNESS QUALITIES OF STEVIA EXTRACT

TECHNICAL FIELD

5 [0001]

The present invention relates to a method for improving the sweetness of stevia extract.

[0002]

10 More specifically, a first aspect of the present invention relates to a technology for improving the sweetness quality of rebaudioside A, which is a main sweetness component of stevia extract (e.g., a sweetener or orally administered composition comprising rebaudioside A, and a method for applying sweetness to an orally administered composition).

15 [0003]

A second aspect of the present invention relates to a technology for improving the sweetness quality of stevia extract itself (e.g., a sweetness quality improving method, a sweetness quality improving agent, and a sweetener or orally administered composition in which the sweetness quality has been improved by the above sweetness quality improving method).

[0004]

25 A third aspect of the present invention relates to a technology for accelerating the sweetness expression of stevia extract in the oral cavity (e.g., a method for accelerating sweetness expression, and a sweetener or orally administered composition in which the sweetness expression has been accelerated by the above method).

30 BACKGROUND ART

[0005]

Hitherto, high-intensity sweeteners have been widely used to impart sweetness to foods, beverages, medicines and quasi drugs, as well as to adjust the taste thereof. In particular, due to the recent rise in health concerns, non- or low-calorie high-

35

intensity sweeteners, and high-intensity sweeteners with low caries-producing effects are widely utilized.

[0006]

5 Examples of high-intensity sweeteners include synthetic sweeteners such as aspartame, sucralose, neotame, acesulfame potassium, and the like; and natural sweeteners such as stevia extract, siraitia grosvenori extract, thaumatin, and the like. Of these, natural sweeteners such as stevia extract have become favored as a result of the recent trend toward natural products.

10 [0007]

 Natural sweeteners, however, exhibit a sweetness quality different from that of sugar, exhibit a rough taste in their sweetness, and leave a lingering sweetness (a remaining aftertaste) in the mouth. In addition to the above, stevia
15 extract, in particular, has drawbacks such as a characteristic bitterness and a lack of sweetness due to its slow sweetness expression in the oral cavity.

[0008]

 The term "sweetness quality" used herein refers to the
20 quality of the sweetness itself. In view of an overall evaluation of sweetness, considering its richness, body, fullness, and low distinctiveness, a quality that is closer to that of sugar is regarded as being better quality. The term "rough taste" used herein refers to a bitter, astringent, sour, stimulating taste,
25 and the like, other than sweetness.

[0009]

 A number of proposals have been made to overcome the above drawbacks inherent in high-intensity sweeteners such as stevia extract.

30 [0010]

 For example, it has been proposed to use, as an agent for improving the taste of a high-intensity sweetener, nigerooligosaccharide (Patent Document 1), cellooligosaccharide (Patent Document 2), a solvent extract of roasted coffee (Patent
35 Document 3), and the like. Further, as a taste improving agent,

which improves the long-lasting aftertaste of aspartame, the use of sodium chloride (Patent Document 4), the combined use of potassium aluminum sulfate and naringin (Patent Document 5), and the like, have been proposed.

5 [0011]

Methods involving the use of a sweet flavor (a sugar flavor) such as maltol to improve the sweetness quality of a high-intensity sweetener have also been proposed. Sugar flavors are prepared by using an aromatic component having a granulated-
10 sugar-like aroma, such as maltol, ethyl maltol or the like, as a main component, together with a minor constituent such as sugar lactone, cyclotene, or the like, which is associated with brown sugar and maple syrup; and a natural starting material such as molasses extract, honey extract, and the like (Non-Patent
15 Document 1).

[0012]

For example, Patent Document 6 suggests using a sweet flavor such as maltol or furaneol, in addition to a high-intensity sweetener such as acesulfame potassium or sucralose, in
20 order to improve the watery taste of sugarless candies that are composed mainly of sugar alcohol. Patent Document 7 teaches a feed additive for livestock comprising a mixture of a stevia extract and a flavoring agent containing a sweet flavor such as maltol or furaneol. Patent Document 7 further teaches that the
25 addition of the feed additive to feed for livestock, including cattle and swine, can make the feed more palatable, and mitigate diarrhea in livestock. None of the above documents, however, discloses or suggests a method for accelerating the sweetness expression of stevia extract. These documents even fail to
30 disclose that stevia extract suffers from slow sweetness expression.

[0013]

Patent Document 8 teaches that a sweet flavor such as maltol is effective for improving the sweetness quality of
35 aspartame (by expressing fullness and body. However, Patent

Document 8 discloses in Example 1 that the sweet flavor is effective only for aspartame, and has no effect on other high-intensity sweeteners such as stevia extract, acesulfame potassium, and the like.

5 [0014]

Methods for improving the sweetness quality of a high-intensity sweetener that involve the combined use of several sweeteners have also been proposed. Patent Document 9, for example, discloses using at least one sweetener selected from
10 siraitia grosvenorii extract, stevia extract, licorice extract, and amacha extract, in combination with a syrup containing erythritol and siraitia grosvenorii glycoside in order to improve its sweetness quality. Patent Document 10 teaches that the further addition of stevia and citric acid to guava tea extract
15 and siraitia grosvenorii extract can be used to prepare a healthy tea having a refreshing sweetness and sweet-sour taste. However, Patent Documents 9 and 10 only disclose using siraitia grosvenorii extract (or siraitia grosvenorii glycoside) in an amount equal to or more than the amount of stevia extract, and
20 neither of these documents discloses or suggests using rebaudioside A, which is a main sweetness component of stevia extract, and mogroside V, which is a main sweetness component of siraitia grosvenorii extract, at a specific weight ratio, in order to improve the sweetness quality.

25 [0015]

Thaumatococcus itself can be used as a sweetener. However, because thaumatococcus exhibits a flavor-enhancing effect and a bitterness-masking effect, it is regarded as a high-intensity sweetener that can also be used as a flavor enhancer, for example.
30 As a method for improving taste quality using thaumatococcus, Patent Document 11 discloses a method for preparing a dressing, etc., having natural sweetness and good flavor, by using, in addition to thaumatococcus, at least one member selected from stevia extract, licorice extract, siraitia grosvenorii extract, amacha extract,
35 and the like. Patent Document 12 teaches a method for producing a

sweetness composition comprising thaumatin together with at least one member selected from licorice extract, stevia extract, and siraitia grosvenorii extract. However, Patent Documents 11 and 12 nowhere mention combining thaumatin with rebaudioside A, which is
5 a main sweetness component of stevia extract, and mogroside V, which is a main sweetness component of siraitia grosvenorii extract.

Patent Document 1: Japanese Unexamined Patent
Publication No. H10-234331

10 Patent Document 2: Japanese Unexamined Patent
Publication No. 2002-223721

Patent Document 3: Japanese Unexamined Patent
Publication No. 2006-81544

15 Patent Document 4: Japanese Unexamined Patent
Publication No. S61-212257

Patent Document 5: Japanese Unexamined Patent
Publication No. S52-90667

Patent Document 6: Japanese Unexamined Patent
Publication No. 2006-340650

20 Patent Document 7: Japanese Unexamined Patent
Publication No. 2001-95502

Patent Document 8: Japanese Unexamined Patent
Publication No. H2-276553

25 Patent Document 9: Japanese Unexamined Patent
Publication No. H11-46701

Patent Document 10: Japanese Unexamined Patent
Publication No. 2003-274911

30 Patent Document 11: Japanese Unexamined Patent
Publication No. 2000-166506

Patent Document 12: Japanese Unexamined Patent
Publication Application No. S53-104771

Non-Patent Document 1: Saishin Koryo no Jiten (New
Encyclopedia of Flavors) (Asakura Publishing Co., Ltd.), pp. 535-
537, May 10, 2000, first edition

DISCLOSURE OF THE INVENTION

Problems to be Solved by the Invention

[0016]

As mentioned above, many attempts have been made to
5 improve the sweetness quality of high-intensity sweeteners in
order to obtain a high-intensity sweetener having a sweetness
quality as close as possible to that of sugar. However, none of
these attempts were sufficient as a method for overcoming stevia
extract drawbacks, in particular, a rough taste in sweetness
10 (bitterness), a long-lasting sweetness in the mouth (a remaining
aftertaste), and a lack of sweetness due to slow sweetness
expression in the oral cavity.

[0017]

In view of the above, the present invention aims to
15 provide a method for improving the sweetness quality of stevia
extract or rebaudioside A, which is a main sweetness component
thereof.

[0018]

More specifically, a first object of the present
20 invention is to provide a method for improving the sweetness
quality of rebaudioside A, which is a main sweetness component of
stevia extract, and to provide orally administered compositions,
such as sweeteners, foods, and beverages, in which the sweetness
quality has been improved by the method.

25 [0019]

A second object of the present invention is to provide
a method for improving the sweetness quality of stevia extract
itself; a sweetness quality improving agent that is useful in the
method; and orally administered compositions, such as sweeteners,
30 foods, beverages, and the like, in which the sweetness quality
has been improved by the sweetness quality improving method.

[0020]

A third object of the present invention is to provide a
method for accelerating the sweetness expression of stevia
35 extract in the oral cavity; and compositions, such as sweeteners,

foods and beverages, in which the sweetness expression in the oral cavity has been accelerated by this method.

Means for Solving the Problems

5 [0021]

The present inventors conducted extensive research to achieve the first object, and found the following. That is, by adjusting the compounding ratio of rebaudioside A, which is a main sweetness component of stevia extract, to mogroside V, which is a main sweetness component of siraitia grosvenorii extract, to a range of 95:5 to 60:40 (weight ratio), in particular, from 94:6 to 75:25 (weight ratio), the aforesaid drawbacks of stevia extract regarding sweetness quality can be eliminated, making the production of a composition having an excellent sweetness quality possible. In addition, the use of the mixture as a sweetener enables excellent sweetness to be applied to various products such as foods, medicines and quasi drugs, which are orally administered or used for the oral cavity.

[0022]

20 The present inventors further conducted extensive research to achieve the second object, and found the following. That is, the addition of at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate, to stevia extract can eliminate the drawbacks of the sweetness quality of stevia extract; in particular, a lingering sweetness in the mouth (a remaining aftertaste) can be improved.

[0023]

30 The present inventors yet further conducted extensive research to achieve the third object, and found the following. That is, the addition of a sweet flavor such as maltol or the like to stevia extract at a weight ratio of 200:1 to 6,000:1, preferably 300:1 to 6,000:1, can eliminate the sweetness quality drawbacks of stevia extract, in particular, the slow sweetness expression in the oral cavity; with the accelerated sweetness

35

expression of stevia extract, a lack of sweetness and lingering sweetness (a remaining aftertaste) can be improved.

[0024]

The present invention was accomplished based on the
5 above findings, and encompasses the following aspects.

[0025]

(I) Improvement in Sweetness Quality of Rebaudioside A

(I-1) Sweetener Comprising Rebaudioside A

(I-1-1) A sweetener comprising rebaudioside A and
10 mogroside V at a weight ratio of 95:5 to 60:40 (rebaudioside A:mogroside V).

[0026]

(I-1-2) The sweetener according to (I-1-1), which
further comprises 0.01 to 3 parts by mass of thaumatin per total
15 100 parts by mass of rebaudioside A and mogroside V.

[0027]

(I-2) Composition Orally Administered or Used for the Oral Cavity

(I-2-1) A composition orally administered or used for
an oral cavity having sweetness and containing rebaudioside A and
20 mogroside V at a weight ratio of 95:5 to 60:40 (rebaudioside A:mogroside V).

[0028]

(I-2-2) The composition according to (I-2-1), which
comprises rebaudioside A and mogroside V in a total amount of 10
25 ppm to 5,000 ppm.

[0029]

(I-2-3) The composition according to (I-2-1) or (I-2-2),
which further comprises 0.01 to 3 parts by mass of thaumatin per
total 100 parts by mass of rebaudioside A and mogroside V.

30 [0030]

(I-3) Application of Sweetness to Compositions Orally
Administered or Used for the Oral Cavity

(I-3-1) A method for applying sweetness to a
composition orally administered or used for an oral cavity
35 comprising the step of: adding rebaudioside A and mogroside V to

a composition orally administered or used for the oral cavity so that the weight ratio of rebaudioside A to mogroside V becomes 95:5 to 60:40.

[0031]

5 (I-3-2) The method according to (I-3-1), wherein rebaudioside A and mogroside V are added to the composition orally administered or used for the oral cavity so that the total content of rebaudioside A and mogroside V becomes 10 ppm to 5,000 ppm.

10 [0032]

(I-3-3) The method according to (I-3-1) or (I-3-2), wherein thaumatin is added to the composition orally administered or used for the oral cavity so that the content of the thaumatin becomes 0.01 to 3 parts by mass per total 100 parts by mass of rebaudioside A and mogroside V.

15 [0033]

(I-3-4) The method according to any one of (I-3-1) to (I-3-3), wherein the composition orally administered or used for the oral cavity is a food or beverage, or a medicine or quasi drug that is orally administered or used for the oral cavity.

20 [0034]

(II) Improvement in Sweetness Quality of Stevia Extract

(II-1) Agent for Improving Taste of Stevia Extract

(II-1-1) A sweetness quality improving agent comprising at least one active ingredient selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrins, magnesium chloride and magnesium sulfate.

[0035]

(II-2) Method for Improving Sweetness of Stevia Extract

30 (II-2-1) A method for improving sweetness quality of a stevia extract comprising the step of: adding a sweetness quality improving agent containing at least one active ingredient selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrins, magnesium chloride and magnesium sulfate to a stevia extract.

35

[0036]

(II-2-2) The method for improving the sweetness quality according to (II-2-1), comprising the step of: adding a sweetness quality improving agent of (II-1-1) to stevia extract or a stevia
5 extract-containing composition that is orally administered or used for an oral cavity.

[0037]

(II-2-3) The method for improving the sweetness quality according to (II-2-2), wherein the sweetness quality improving
10 agent is used so that the content of each of the active ingredients per part by mass of the stevia extract falls within the following ranges:

Gentiooligosaccharide: 5 to 100 parts by mass, and preferably 5 to 50 parts by mass;

15 Naringin: 0.01 to 1 part by mass, and preferably 0.01 to 0.5 part by mass;

Highly branched cyclic dextrin: 0.1 to 50 parts by mass, and preferably 0.5 to 30 parts by mass;

Magnesium chloride: 0.05 to 5 parts by mass, and
20 preferably 0.05 to 3 parts by mass; and

Magnesium sulfate: 0.05 to 5 parts by mass, and preferably 0.05 to 3 parts by mass.

[0038]

(II-2-4) The method according to (II-2-2) or (II-2-3),
25 wherein the composition orally administered or used for the oral cavity is a food or beverage, or a medicine or quasi drug that is orally administered or used for the oral cavity.

[0039]

(II-3) Stevia Extract-Containing Sweetener

30 (II-3-1) A sweetener comprising a stevia extract and at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate.

[0040]

35 (II-3-2) The sweetener according to (II-3-1), which

comprises the following ranges of at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate, per part by mass of stevia extract:

5 Gentiooligosaccharide: 5 to 100 parts by mass, and preferably 5 to 50 parts by mass;

 Naringin: 0.01 to 1 part by mass, and preferably 0.01 to 0.5 part by mass;

 Highly branched cyclic dextrin: 0.1 to 50 parts by mass,
10 and preferably 0.5 to 30 parts by mass;

 Magnesium chloride: 0.05 to 5 parts by mass, and preferably 0.05 to 3 parts by mass; and

 Magnesium sulfate: 0.05 to 5 parts by mass, and preferably 0.05 to 3 parts by mass.

15 [0041]

(II-4) Composition Orally Administered or Used for the Oral Cavity

(II-4-1) A composition orally administered or used for the oral cavity having sweetness and containing stevia extract
20 and at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate.

[0042]

(II-4-2) The composition according to (II-4-1), which
25 comprises the following ranges of at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate per part by mass of stevia extract contained in the composition orally administered or used for the oral cavity:

30 Gentiooligosaccharide: 5 to 100 parts by mass, and preferably 5 to 50 parts by mass;

 Naringin: 0.01 to 1 part by mass, and preferably 0.01 to 0.5 part by mass;

 Highly branched cyclic dextrin: 0.1 to 50 parts by mass,
35 and preferably 0.5 to 30 parts by mass;

Magnesium chloride: 0.05 to 5 parts by mass, and preferably 0.05 to 3 parts by mass, and

Magnesium sulfate: 0.05 to 5 parts by mass, and preferably 0.05 to 3 parts by mass.

5 [0043]

(III) Acceleration of Sweetness Expression of Stevia Extract

(III-1) Method for Accelerating Sweetness Expression of Stevia Extract in an Oral Cavity

(III-1-1) A method for accelerating sweetness
10 expression of a stevia extract in an oral cavity comprising the step of: combining 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor.

[0044]

(III-1-2) The method according to (III-1-1), wherein
15 the sweet flavor is at least one member selected from the group consisting of maltol, ethyl maltol, sotolon, cyclotene, furaneol, and homofuraneol.

[0045]

(III-1-3) The method according to (III-1-1), wherein
20 the sweet flavor is maltol.

[0046]

(III-2) Sweetener

(III-2-1) A sweetener comprising 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor.

25 [0047]

(III-2-2) The sweetener according to (III-2-1), wherein the sweet flavor is at least one member selected from the group consisting of maltol, ethyl maltol, sotolon, cyclotene, furaneol, and homofuraneol.

30 [0048]

(III-2-3) The sweetener according to (III-2-1), wherein the sweet flavor is maltol.

[0049]

(III-3) Composition Orally Administered or Used for the Oral
35 Cavity

(III-3-1) A composition that is orally administered or used for an oral cavity having sweetness and comprising 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor.

5 [0050]

(III-3-2) The composition according to (III-3-1), wherein the sweet flavor is at least one member selected from the group consisting of maltol, ethyl maltol, sotolon, cyclotene, furaneol, and homofuraneol.

10 [0051]

(III-3-3) The composition according to (III-3-1), wherein the sweet flavor is maltol.

[0052]

(III-3-4) The composition according to any one of (III-3-1) to (III-3-3), which comprises 10 ppm to 5,000 ppm of stevia extract.

Effect of the Invention

[0053]

20 The sweetener comprising rebaudioside A, which is the first aspect of the present invention, has an excellent sweetness quality that is similar to sugar. This sweetener can apply excellent sweetness to various compositions. Further, the additional use of thaumatin together with rebaudioside A and
25 mogroside V can produce a sweetener having an excellent and more sugar-like sweetness quality.

[0054]

The second aspect of the present invention can provide a novel usage of gentioooligosaccharides, naringins, highly
30 branched cyclic dextrans, magnesium chloride and magnesium sulfate, i.e., a usage as a sweetness quality improving agent of stevia extract. According to this aspect, the use of the above sweetness quality improving agent can improve the sweetness quality of stevia extract; in particular, lingering sweetness (a
35 remaining aftertaste) can be improved. The present invention can

thereby provide a sweetener having an excellent sweetness quality by improving the known drawbacks of stevia extract (e.g. bitterness, remaining aftertaste, and slow sweetness expression).
[0055]

5 The third aspect of the present invention can improve the known drawbacks of stevia extract, such as the slow rate of sweetness expression in the oral cavity, and accelerate the sweetness expression, thereby eliminating the drawbacks, such as a lack of sweetness, and lingering sweetness (a remaining
10 aftertaste).

BEST MODE FOR CARRYING OUT THE INVENTION

[0056]

(I) Improvement in Sweetness Quality of Rebaudioside A

15 (I-1) Sweetener

One of the features of the sweetener of the present invention is that it comprises rebaudioside A and mogroside V at a weight ratio of 95:5 to 60:40.

[0057]

20 Rebaudioside A is a known steviol glycoside, which is a main sweetness component contained in stevia extract. Rebaudioside A has a degree of sweetness (sweetness intensity) 150 to 450 times that of sugar. The rebaudioside A targeted by the present invention includes enzyme-treated rebaudioside A
25 obtained by transferring glucose, fructose or like saccharide to rebaudioside A using α -glucosyltransferase or the like.

[0058]

30 The rebaudioside A used in the present invention may be, but is not limited to, purified rebaudioside A, and may be a mixture with other steviol glycosides (such as stevioside, rebaudioside B, rebaudioside C, rebaudioside D, rebaudioside, dulcoside A, rubusoside, and steviolbioside). When a mixture is used, the content of rebaudioside A is preferably 70% by mass or more per the total amount of the mixture. This is because the
35 greater the content of steviol glycosides such as stevioside

other than rebaudioside A becomes, the less the effects on the present invention, in particular, the improvement in sweetness quality due to thaumatin tend to become.

[0059]

5 The rebaudioside A used in the present invention may be prepared by subjecting leaves or stems of *Stevia Rebaudiana Bertoni* in the composite family to extraction and purification. However, commercially available products may also be conveniently used. Examples of such commercially available products include
10 Rebaudio J-100 (produced by Morita Kagaku Kogyo Co., Ltd.), Chrysanta 99-P (produced by Dainippon Ink and Chemicals, Incorporated), etc. These products contain 80% by mass or more of rebaudioside A.

[0060]

15 Mogroside V is a known triterpene-based glucoside, which is a main sweetness component contained in siraitia grosvenorii extract. Mogroside V has a degree of sweetness about 300 times that of sugar.

[0061]

20 Similar to rebaudioside A, mogroside V may be, but is not limited to, a purified product and may be a mixture with other triterpene-based glucosides (such as mogrol, mogroside IE₁, mogroside IA₁, mogroside IIE, mogroside III, mogroside IVa, mogroside IVE, siamenside, 11-oxo-mogroside, and 5 α ,6 α -
25 epoxymogroside). When a mixture is used, the content of mogroside V is preferably 5% or more per the total amount of the mixture.

[0062]

 The mogroside V used in the present invention may be prepared by subjecting the fruits of siraitia grosvenori in the
30 gourd family to extraction and purification. However, commercially available products may also be conveniently used. One example of such commercially available products is LUO HAN GUO CONCENTRATE EXTRACT POWDER (Siraitia Grosvenori Extract Powder: produced by San-Ei Gen F.F.I., Inc.). The product
35 contains 5% by mass of mogroside V.

[0063]

As described above, the sweetener of the present invention contains rebaudioside A and mogroside V in a weight ratio of 95:5 to 60:40. The proportion of rebaudioside A to mogroside V is preferably 94:6 to 75:25, and more preferably 90:10 to 75:25 (weight ratio). When the content of mogroside V exceeds 40 parts by mass per 60 parts by mass of rebaudioside A, the sweetness quality cannot be improved. When the content of mogroside V exceeds 50 parts by mass per 50 parts by mass of rebaudioside A, i.e., the content of mogroside V exceeds that of rebaudioside A in a sweetener, the sweetness quality tends to be inferior to that of a sweetener in which rebaudioside A is used singly.

[0064]

The sweetener of the present invention may contain thaumatin in addition to rebaudioside A and mogroside V. By using thaumatin in combination with mogroside V, the sweetness quality of rebaudioside A can be further improved, and a sweetness quality similar to that of sugar can be attained.

[0065]

Thaumatococcus daniellii BENTH, which is a perennial plant in the Marantaceae family. Thaumatococcus daniellii is a sweet component having a molecular weight of about 21,000 and containing protein as its main component. Thaumatococcus daniellii has a very high sweetness intensity, i.e., 3,000 to 8,000 times that of sugar.

[0066]

The content of thaumatin is generally 0.01 to 3 parts by mass, preferably 0.05 to 2 parts by mass, and more preferably 0.1 to 1 part by mass per total 100 parts by mass of rebaudioside A and mogroside V contained in the sweetener.

[0067]

The thaumatin can be prepared by subjecting the aril of Thaumatococcus daniellii fruit to water extraction and purification. However, commercially available products may also

be conveniently used. One example of such commercially available products is SAN SWEET T (produced by San-Ei Gen F.F.I., Inc.). SAN SWEET T consists of 100% by mass of thaumatin.

[0068]

5 There is no limitation to the form of the sweetener of the present invention, and the sweetener may take any form including a powder, a granule, a tablet, a capsule and like solid states; or a syrup, an emulsion, a liquid, a gel and like semi-solid or liquid states.

10 [0069]

 The sweetener of the present invention can be prepared by mixing rebaudioside A and mogroside V, and optionally with thaumatin, at the ratio described above. Depending on the form, pharmacologically acceptable carriers or carriers that may be
15 added to foods and beverages may also be added during the preparation. Examples of such carriers include lactose, glucose, fructose, sugar, high-fructose corn syrup and like saccharides; sorbitol, erythritol, lactitol, maltitol, mannitol, xylitol, isomalt and like sugar alcohols; isomalto-oligosaccharide,
20 galacto-oligosaccharide, fructo-oligosaccharide and like oligosaccharides; dextrin, cellulose, gum arabic, cornstarch and like polysaccharides; water; etc.

[0070]

 Furthermore, flavors, colorants, acidulants,
25 preservatives and other components that are ordinarily added to foods, beverages or medicines may be added. It is also possible to suitably add aspartame, sucralose, acesulfame potassium, saccharin or salts thereof (sodium saccharin, calcium saccharin, etc.) and like artificial sweeteners; or licorice extract, amacha
30 extract (extract from leaf of *Hydrangea macrophylla* var. *thunbergii*), brazzein, neohesperidine dihydrochalcone, serendipity berry extract (monellin), tenryocha extract and like natural sweeteners.

[0071]

35 The total amount of rebaudioside A and mogroside V, or

the total amount of rebaudioside A, mogroside V and thaumatin in the sweetener is suitably adjusted so that it falls within the range of 0.1 to 100 weight%.

[0072]

5 (I-2) Composition Orally Administered or Used in the Oral Cavity

The composition orally administered or used for the oral cavity of the present invention is a composition that has a sweet taste and contains rebaudioside A and mogroside V at a weight ratio of 95:5 to 60:40. The compounding ratio of
10 rebaudioside A to mogroside V is preferably 94:6 to 75:25 (weight ratio), and more preferably 90:10 to 75:25 (weight ratio).

[0073]

The composition targeted by the present invention is a composition that is orally administered or used for the oral
15 cavity that needs sweetness. Examples of such compositions include foods and beverages; orally administered medicines; medicines for the oral cavity; and oral care products such as dentifrices and mouthwashes (including medicines or quasi drugs). Among these, foods and beverages are preferable.

20 [0074]

Specific examples of foods and beverages include carbonated beverages, fruit-juice beverages, coffee beverages, black tea beverages, lactic acid beverages, lactic acid bacterium beverages, soft drinks, milk beverages, alcoholic beverages and
25 like beverages; okaki (small rice cracker), senbei (rice cracker), okoshi (dry cake made of rice or millet and starch syrup), manju (steamed azuki bean jam-filled bun), candy and other various Japanese sweets; cookies, biscuits, crackers, pies, sponge cakes, kasutera (Castilla cakes), doughnuts, waffles, butter cream,
30 custard cream, cream puffs, chocolate, chocolate confectioneries, caramel candies, jelly, pancakes, breads and other Western-style confectioneries; chewing gum, bubble gum and like gums; potato chips and other various dry snacks; ice cream, popsicles, sherbet and other ice confectioneries; flour paste, peanut paste, fruit
35 paste and other pastes; pickles; miso, powdered miso, soy sauce,

powdered soy sauce, moromi, fish sauce, sauce, ketchup, mayonnaise, solid bouillon, sauces used for yakiniku, base for stew, base for soup, base for lightly pickled foods and other seasonings; yogurt, pudding, Bavarian cream and like milk products.

[0075]

Specific examples of orally administered medicines include troche tablets, ampuled liquid medicines, granules, pulvis (powders), tablets, and capsules. Specific examples of medicines for the oral cavity include sprays, ointments, creams, pastes and patches. Specific examples of oral care products include liquid dentifrices, toothpastes, mouthwashes, and breath fresheners.

[0076]

These compositions contain rebaudioside A and mogroside V with a compounding ratio that falls within the above-mentioned range, wherein the total content thereof is 10 ppm to 5,000 ppm. The total content of rebaudioside A and mogroside V depends on the type of composition targeted, but is preferably 25 ppm to 3,000 ppm, and more preferably 50 ppm to 2,000 ppm. It is preferable that thaumatin be added in an amount of 0.01 to 3 parts by mass, preferably 0.05 to 2 parts by mass, and more preferably 0.1 to 1 part by mass per total 100 parts by mass of rebaudioside A and mogroside V in the composition. This renders further improved sweetness to the composition.

[0077]

Rebaudioside A and mogroside V, and optionally thaumatin, may be added to a composition that is orally administered or used for the oral cavity at any step of its production.

[0078]

(I-3) Application of Sweetness to Compositions Orally Administered or Used for the Oral Cavity

Sweetness can be applied to a composition that is orally administered or used for the oral cavity of the present

invention by adding rebaudioside A and mogroside V to the compositions targeted in (I-2) so that their weight ratio (rebaudioside A to mogroside V) falls within the range of 95:5 to 60:40. The compounding ratio of rebaudioside A to mogroside V is preferably 94:6 to 75:25 (weight ratio) and more preferably 90:10 to 75:25 (weight ratio).

[0079]

The amounts of rebaudioside A and mogroside V necessary to provide these compositions with excellent sweetness similar to that of sugar vary depending on the type of composition, etc; however, the total amount of rebaudioside A and mogroside V is generally 10 ppm to 5,000 ppm, preferably 25 ppm to 3,000 ppm, and more preferably 50 ppm to 2,000 ppm.

[0080]

In addition to rebaudioside A and mogroside V, thaumatin may be added. In this case, the content of thaumatin is generally 0.01 to 3 parts by mass, preferably 0.05 to 2 parts by mass, and more preferably 0.1 to 1 part by mass per total 100 parts by mass of rebaudioside A and mogroside V in the composition. By adding thaumatin, in addition to rebaudioside A and mogroside V, further improved sweetness can be applied to the objective composition that is orally administered or used for the oral cavity.

[0081]

In the method for applying sweetness of the present invention, rebaudioside A and mogroside V, or rebaudioside A, mogroside V and thaumatin, may be independently added to the targeted composition that is orally administered or used for the oral cavity so that they have the above-mentioned compounding ratio. Alternatively, the sweeteners of the present invention described in (I-1) may be added to a targeted composition that is orally administered or used for the oral cavity so that the amounts of rebaudioside A, mogroside V and thaumatin correspond to the above-mentioned compounding ratio.

[0082]

(II) Improvement in Sweetness Quality of Stevia Extract

(II-1) Agent for Improving Taste of Stevia Extract

The sweetness quality improving agent of the present invention consists of or comprises, as an active ingredient, at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrins, magnesium chloride and magnesium sulfate.

[0083]

The stevia extract targeted by the sweetness quality improving agent of the present invention comprises, as a main component, steviol glycoside that is obtained by subjecting the leaves or stems of *Stevia Rebaudiana Bertoni* in the composite family to extraction using water or ethanol. Examples of commercially available products include Rebaudio J-100, Rebaudio A9-90, and Steviron TK (these are produced by Morita Kagaku Kogyo Co., Ltd.); Chrysanta 99-P and Chrysanta AX-P (these are produced by Dainippon Ink and Chemicals, Incorporated); Steviafin H and Steviafin HN2 (these are produced by Nippon Paper Chemicals Co., Ltd.); Stevia ST-AB and Histevia 500 (these are produced by Ikeda Tohka Industries Co., Ltd.); etc.

[0084]

The stevia extract targeted by the present invention includes enzyme-treated stevia obtained by transferring glucose, fructose or like saccharide to a stevia extract using α -glucosyltransferase, etc.

[0085]

Gentiooligosaccharide, which is one of the active ingredients of the sweetness quality improving agent of the present invention, is one type of β -gluco-oligosaccharide. Gentiooligosaccharide is a low-calorie saccharide and is characteristically bitter. The gentiooligosaccharide used in the present invention can be produced by making β -glucosidase to act upon glucose or β -gluco-oligosaccharide; however, commercially available gentiooligosaccharide-containing syrups (for example, Gentose#45: Produced by Nihon Shokuhin Kako Co., Ltd., containing

45% gentiooligosaccharide) may also be obtained.

[0086]

Gentiooligosaccharide may be used in an amount of generally 5 to 100 parts by mass, preferably 5 to 50 parts by mass, and more preferably 10 to 30 parts by mass per part by mass of stevia extract. When the content of gentiooligosaccharide is less than 5 parts by mass, a satisfactory reduction in the lingering sweetness of stevia extract cannot be obtained. In contrast, when the content of gentiooligosaccharide exceeds 100 parts by mass, the bitterness attributable to gentiooligosaccharide is unfavorably increased.

[0087]

Naringin is a flavanone glucoside that is contained, in considerable amounts, in the peels of grapefruit, hassaku orange and like citrus fruits, and causes the bitterness or pungency of citrus fruits. The naringin used in the present invention can be obtained by subjecting the peels or fruit juice of citrus fruits to extraction and purification using water or ethanol; however, commercially available products such as SAN FIX NARINGIN (produced by San-Ei Gen F.F.I., Inc.) may also be obtained.

[0088]

It is also possible to use enzyme-treated naringin that is obtained by adding glucose to naringin by making cyclodextrin glucosyltransferase to affect upon a mixture of naringin and dextrin.

[0089]

The content of naringin is generally 0.01 to 1 part by mass, preferably 0.01 to 0.5 part by mass, and more preferably 0.02 to 0.2 part by mass per part by mass of stevia extract. When the content of naringin is less than 0.01 part by mass, a satisfactory reduction of the lingering sweetness of stevia extract cannot be attained; however, when it exceeds 1 part by mass, the bitterness attributable to naringin is unfavorably increased.

[0090]

A highly branched cyclic dextrin is a high-molecular-weight dextrin having a molecular weight of about 30,000 to 1,000,000. The highly branched cyclic dextrin has characteristics, such as being easily dissolvable in cold water, and aging less easily than ordinary dextrans. The dextrin used in the present invention can be prepared by liquefying a starch essentially consisting of waxy corn starch using α -amylase, and then saccharifying the resulting mixture using a branching enzyme (EC2.4.1.18). However, commercially available products such as Cluster Dextrin (produced by Nihon Shokuhin Kako Co., Ltd.) are also usable.

[0091]

Highly branched cyclic dextrin may be added in an amount of 0.1 to 50 parts by mass, preferably 0.5 to 30 parts by mass, and more preferably 1 to 20 parts by mass per part by mass of stevia extract. When the content of highly branched cyclic dextrin is less than 0.1 part by mass, satisfactory reduction in the lingering sweetness of stevia extract cannot be attained. When the content thereof exceeds 50 parts by mass, it becomes too powdery and is thus unfavorable.

[0092]

Magnesium chloride, which is an inorganic compound, has deliquescence and is extremely easily dissolvable in water. An aqueous solution of magnesium chloride is called bittern. Magnesium sulfate, which is also an inorganic compound, has a bitter taste, a pleasant cooling sensation, and a salty taste. Magnesium sulfate is used as a laxative, etc.

[0093]

Magnesium chloride and magnesium sulfate can each be used in an amount of 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by mass per part by mass of stevia extract. When the content thereof is less than 0.05 part by mass, satisfactory reduction in the lingering sweetness of stevia extract cannot be attained. When the content thereof exceeds 5 parts by mass, the bitterness attributable to

magnesium chloride and magnesium sulfate is unfavorably increased.
[0094]

The sweetness quality improving agent of the present invention may consist of only one of the above components or may
5 contain two or more components in combination.
[0095]

By adding one or more pharmacologically acceptable carriers or carriers that may be added to foods and beverages to the active ingredients described above, the sweetness quality
10 improving agent of the present invention may be prepared into an arbitrary form such as a powder, a granule, a tablet, a capsule and like solids; or syrup, emulsion, liquid, gel and like semi-solids or liquids.
[0096]

15 There is no limitation to the carriers as long as they do not hinder the effects of the present invention. Examples of carriers include lactose, glucose, fructose, sugar, high-fructose corn syrup and like saccharides; sorbitol, erythritol, lactitol, maltitol, mannitol, xylitol, isomalt and like sugar alcohols;
20 isomalto-oligosaccharide, galacto-oligosaccharide, fructo-oligosaccharide and like oligosaccharides; dextrin, cellulose, gum arabic, cornstarch and like polysaccharides; water; ethanol; etc.
[0097]

25 To the sweetness quality improving agent of the present invention, within the range that does not hinder the effects of the present invention, sucralose, aspartame, acesulfame potassium, siraitia grosvenori extract, thaumatin, licorice, saccharin or salts thereof (sodium saccharin, calcium saccharin, etc.);
30 neotame, alitame and like high intensity sweeteners; L-alanine and like amino acids; sodium gluconate, potassium gluconate, succinic acid and like acidulants; and dextrin, cellulose, gum arabic, cornstarch, inositol, caffeine, vitamins, colorant, and flavor may be added.
35 [0098]

The contents of gentioooligosaccharide, naringin, highly branched cyclic dextrin, magnesium chloride, and magnesium sulfate in the sweetness quality improving agent of the present invention are suitably adjusted so that the total amount thereof
5 falls within the range of 0.1 to 100 weight%.

[0099]

The sweetness quality improving agent of the present invention is formed into a solid, semi-solid or liquid preparation as described above by a known method, and used to
10 improve the sweetness quality of various compositions that are orally administered or used for the oral cavity (foods and beverages, medicines orally administered or used for the oral cavity, quasi drugs used for the oral cavity, oral care products, etc.) that consist of or comprise stevia extract.

15 [0100]

(II-2) Method for Improving Sweetness of Stevia Extract

The present invention provides a method for improving the sweetness quality of stevia extract.

[0101]

20 The method can be conducted by combining stevia extract with the sweetness quality improving agent described in (II-1). In other words, the method can be conducted by making stevia extract coexist with the sweetness quality improving agent of the present invention. More specifically, the method can be conducted
25 by adding the sweetness quality improving agent of the present invention to stevia extract or a stevia extract-containing composition that is orally administered or used for the oral cavity, or by adding stevia extract and the sweetness quality improving agent of the present invention to a composition that is
30 orally administered or used for the oral cavity.

[0102]

The composition of the present invention is a composition that is orally administered or used for the oral cavity that needs sweetness. Examples of such compositions
35 include foods and beverages; orally administered medicines;

medicines for the oral cavity; and oral care products such as dentifrices and mouthwashes (including medicines or quasi drugs). Among these, foods and beverages are preferable.

[0103]

5 Specific examples of foods and beverages include carbonated beverages, fruit-juice beverages, coffee beverages, black tea beverages, lactic acid beverages, lactic acid bacterium beverages, soft drinks, milk beverages, alcoholic beverages and like beverages; okaki (rice cracker), senbei, okoshi (dry cake
10 made of rice or millet and starch syrup), manju (steamed azuki bean jam-filled bun), candy and other various Japanese sweets; cookies, biscuits, crackers, pies, sponge cakes, kasutera (Castilla cakes), doughnuts, waffles, butter cream, custard cream, cream puffs, chocolate, chocolate confectioneries, caramel candy,
15 jelly, pancakes, breads and other Western-style confectioneries; chewing gum, bubble gum and like gums; potato chips and other dry snacks; ice cream, popsicles, sherbet and other ice confectioneries; flour paste, peanut paste, fruit paste and other pastes; pickles; miso, powdered miso, soy sauce, powdered soy
20 sauce, moromi, fish sauce, sauce, ketchup, mayonnaise, solid bouillon, sauces used for yakiniku, base for stew, base for soup, base for lightly pickled foods and other seasonings; and yogurt, pudding, Bavarian cream and like milk products.

[0104]

25 Specific examples of orally administered medicines include troche tablets, ampuled liquid medicines, granules, pulvis (powders), tablets, and capsules. Specific examples of medicines for the oral cavity include sprays, ointments, creams, pastes and patches. Specific examples of oral care products
30 include liquid dentifrices, toothpastes, mouthwashes, and breath fresheners.

[0105]

 The amount of stevia extract contained in these compositions varies depending on the type of composition, but is
35 generally 10 ppm to 5,000 ppm, preferably 25 ppm to 3,000 ppm,

and more preferably 50 to 2,000 ppm.

[0106]

The sweetness quality improving agent of the present invention may be used so that the content of each active ingredient, relative to stevia extract or a stevia extract-containing composition, falls within the following ranges:

Gentiooligosaccharide: generally 5 to 100 parts by mass, preferably 5 to 50 parts by mass, and more preferably 10 to 30 parts by mass per part by mass of stevia extract;

Naringin: 0.01 to 1 part by mass, preferably 0.01 to 0.5 part by mass, and more preferably 0.02 to 0.2 part by mass per part by mass of stevia extract;

Highly branched cyclic dextrin: 0.1 to 50 parts by mass, preferably 0.5 to 30 parts by mass, and more preferably 1 to 20 parts by mass per part by mass of stevia extract;

Magnesium chloride: 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by mass per part by mass of stevia extract; and

Magnesium sulfate: 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by mass per part by mass of stevia extract.

[0107]

(II-3) Stevia Extract-Containing Sweetener

One of the features of the sweetener of the present invention is that it contains stevia extract in combination with the sweetness quality improving agent of the present invention described in (II-1).

[0108]

The compounding ratio of the stevia extract to the sweetness quality improving agent of the present invention in the sweetener may be selected so that the content of each component in the sweetness quality improving agent of the present invention falls within the following ranges:

Gentiooligosaccharide: generally 5 to 100 parts by mass, preferably 5 to 50 parts by mass, and more preferably 10 to 30

parts by mass per part by mass of stevia extract;

Naringin: 0.01 to 1 part by mass, preferably 0.01 to 0.5 part by mass, and more preferably 0.02 to 0.2 part by mass per part by mass of stevia extract;

5 Highly branched cyclic dextrin: 0.1 to 50 parts by mass, preferably 0.5 to 30 parts by mass, and more preferably 1 to 20 parts by mass per part by mass of stevia extract;

 Magnesium chloride: 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by
10 mass per part by mass of stevia extract; and

 Magnesium sulfate: 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by mass per part by mass of stevia extract.

[0109]

15 There is no limitation to the form of the sweetener of the present invention, and the sweetener may take any form including a powder, a granule, a tablet, a capsule and like solid states; or syrup, emulsion, liquid, gel and like semi-solid or liquid states.

20 [0110]

 The sweetener of the present invention can be prepared by mixing stevia extract with the sweetness quality improving agent of the present invention so that their compounding ratio falls within the ranges mentioned above. During the preparation,
25 depending on the form, pharmacologically acceptable carriers or carriers that may be added to foods and beverages may also be added. Examples of such carriers include lactose, glucose, fructose, sugar, high-fructose corn syrup and like saccharides; sorbitol, erythritol, lactitol, maltitol, mannitol, xylitol,
30 isomalt and like sugar alcohols; isomalto-oligosaccharide, galacto-oligosaccharide, fructo-oligosaccharide and like oligosaccharides; dextrin, cellulose, gum arabic, cornstarch and like polysaccharides; water; etc.

[0111]

35 Furthermore, flavors, colorants, acidulants,

preservatives and other components that are ordinarily added to foods, beverages or medicines may be added. If desired, it is also possible to add aspartame, saccharin, sucralose, acesulfame potassium and like synthetic sweeteners; or licorice extract, amacha extract, brazzein, neohesperidine dihydrochalcone, serendipity berry extract (monellin), tenryocha extract and like natural sweeteners.

[0112]

The total amount of the stevia extract and the sweetness quality improving agent of the present invention in the sweetener is suitably adjusted so that it falls within the range of 0.1 to 100 weight%.

[0113]

(II-4) Composition Orally Administered or Used for the Oral Cavity

The composition orally administered or used for the oral cavity of the present invention has a sweet taste and comprises stevia extract and at least one member selected from the group consisting of gentioooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate.

[0114]

The composition targeted by the present invention is a composition that is orally administered or used for the oral cavity that needs sweetness. Examples of such compositions include foods and beverages; orally administered medicines; medicines for the oral cavity; oral care products such as dentifrices and mouthwashes (including medicines or quasi drugs) described in (II-2). Among these, foods and beverages are preferable.

[0115]

The amount of stevia extract contained in these compositions varies depending on the type of composition, but is generally 10 ppm to 5,000 ppm, preferably 25 ppm to 3,000 ppm, and more preferably 50 to 2,000 ppm.

[0116]

As described in (II-2), the content of each component in the composition of the present invention per part by mass of stevia extract falls within the following ranges:

5 Gentiooligosaccharide: 5 to 100 parts by mass, preferably 5 to 50 parts by mass, and more preferably 10 to 30 parts by mass;

 Naringin: 0.01 to 1 part by mass, preferably 0.01 to 0.5 part by mass, and more preferably 0.02 to 0.2 part by mass;

10 Highly branched cyclic dextrin: 0.1 to 50 parts by mass, preferably 0.5 to 30 parts by mass, and more preferably 1 to 20 parts by mass;

 Magnesium chloride: 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by
15 mass;

 Magnesium sulfate: 0.05 to 5 parts by mass, preferably 0.05 to 3 parts by mass, and more preferably 0.1 to 2 parts by mass.

[0117]

20 Stevia extract and gentiooligosaccharide, naringin, highly branched cyclic dextrin, magnesium chloride, magnesium sulfate, etc., may be added in any step of producing the composition that is orally administered or used for the oral cavity.

25 [0118]

 The composition of the present invention thus prepared solves the problem of the sweetness quality of stevia extract (such as bitterness, lingering sweetness, and slow sweetness expression) and eliminates lingering sweetness. Accordingly, the
30 composition of the present invention has an excellent sweetness quality that is similar to that of sugar.

[0119]

(III) Acceleration of Sweetness Expression of Stevia Extract

(III-1) Method for Accelerating Sweetness Expression of Stevia
35 Extract

Among the problems of stevia extract relating to its sweetness quality, the present invention provides a method for improving slow sweetness expression in the oral cavity.

[0120]

5 The method can be conducted by adding stevia extract in an amount of 200 to 6,000 parts by mass per part by mass of sweet flavor. The amount of stevia extract added is preferably 300 to 6,000 parts by mass, more preferably 400 to 6,000 parts by mass, and particularly preferably 600 to 4,000 parts by mass per part
10 by mass of sweet flavor.

[0121]

 The stevia extract targeted by the present invention comprises, as a main component, steviol glycoside that is obtained by subjecting the leaves or stems of *Stevia Rebaudiana*
15 *Bertoni* in the composite family to extraction using water or ethanol. Examples of commercially available products include Rebaudio J-100, Rebaudio A9-90, and Steviron TK (these are produced by Morita Kagaku Kogyo Co., Ltd.); Chrysanta 99-P and Chrysanta AX-P (these are produced by Dainippon Ink and Chemicals,
20 Incorporated); Steviafin H and Steviafin HN2 (these are produced by Nippon Paper Chemicals Co., Ltd.); Stevia ST-AB and Histevia 500 (these are produced by Ikeda Tohka Industries Co., Ltd.); etc.

[0122]

 The stevia extract targeted by the present invention
25 includes enzyme-treated stevia that is obtained by transferring glucose, fructose or a like saccharide to stevia extract using α -glucosyltransferase, etc.

[0123]

 The sweet flavor used in the present invention is an
30 aroma component that is not sweet itself, but has an aroma similar to that of granulated sugar. Specific examples thereof include maltol, ethyl maltol, sotolon, cyclotene, furaneol and homofuraneol. These sweet flavors may be used singly or in combination. Among various possible combinations, it is
35 preferable to use maltol singly or in combination with other

sweet flavors. The sweet flavor may be used singly by diluting it with a solvent, or in combination with other flavors, such as a citrus-based flavor. Furthermore, minor components such as sugar lactone that is reminiscent of brown sugar or maple syrup, and cyclotene; and natural raw materials such as molasses extract and honey extract, may also be added.

[0124]

The proportion of the stevia extract to sweet flavor used to achieve the effect of the present invention, i.e., accelerating the sweetness expression of stevia extract in the oral cavity, depends on the type of the sweet flavor used, but is generally 200 to 6,000 parts by mass, preferably 300 to 6,000 parts by mass, more preferably 400 to 6,000 parts by mass, and particularly preferably 600 to 4,000 parts by mass per part of sweet flavor. When the content of stevia extract per part by mass of sweet flavor is less than 200 parts by mass, the sugar-like aroma of the sweet flavor becomes too conspicuous and an unfavorable lingering sweetness (a remaining aftertaste) increases. In contrast, when the content of stevia extract per part by mass of sweet flavor exceeds 6,000 parts by mass, the satisfactory effect of the present invention cannot be attained.

[0125]

The sweetness expression of the stevia extract in the oral cavity can be accelerated by adding sweet flavor to stevia extract itself, a sweetener that contains stevia extract as an active ingredient, or a composition that is orally administered or used for the oral cavity that has a sweet taste and contains stevia extract, with the proportion of stevia extract to sweet flavor falling within the above-described range.

[0126]

The sweetener targeted by the present invention may suitably contain, in addition to stevia extract, lactose, glucose, fructose, sugar, high-fructose corn syrup and like saccharides; sorbitol, erythritol, lactitol, maltitol, mannitol, xylitol, isomalt and like sugar alcohols; isomalto-oligosaccharide,

galacto-oligosaccharide, fructo-oligosaccharide and like
oligosaccharides; dextrin, cellulose, gum arabic, cornstarch and
like polysaccharides; amino acid or a salt thereof; aspartame,
sucralose, acesulfame potassium, neotame and like synthetic
5 sweeteners; thaumatin, licorice extract, amacha extract, brazzein,
serendipity berry extract (monellin), tenryocha extract and like
natural sweeteners.

[0127]

The composition targeted by the present invention is a
10 composition that is orally administered or used for the oral
cavity that contains stevia extract. Examples of such
compositions include foods and beverages; orally administered
medicines; medicines for the oral cavity; and dentifrices,
mouthwashes and like oral care products (including medicines and
15 quasi drugs). Among these, foods and beverages are preferable,
and beverages are particularly preferable.

[0128]

Specific examples of foods and beverages include
carbonated beverages, fruit-juice beverages, coffee beverages,
20 black tea beverages, lactic acid beverages, lactic acid bacterium
beverages, soft drinks, milk beverages, alcoholic beverages and
like beverages; okaki (small rice cracker), senbei (rice cracker),
okoshi (dry cake made of rice or millet and starch syrup), manju
(steamed azuki bean jam-filled bun), candy and other various
25 Japanese sweets; cookies, biscuits, crackers, pies, sponge cakes,
kasutera (Castilla cakes), doughnuts, waffles, butter cream,
custard cream, cream puffs, chocolate, chocolate confectioneries,
caramel candy, jelly, pancakes, breads and other Western-style
confectioneries; chewing gum, bubble gum and like gums; potato
30 chips and other dry snacks; ice cream, popsicles, sherbet and
other ice confectioneries; flour paste, peanut paste, fruit paste
and other pastes; pickles; miso, powdered miso, soy sauce,
powdered soy sauce, moromi, fish sauce, sauce, ketchup,
mayonnaise, solid bouillon, sauces used for yakiniku, base for
35 stew, base for soup, base for lightly-pickled foods and other

seasonings; yogurt, pudding, Bavarian cream and like milk products.

[0129]

Specific examples of orally administered medicines
5 include troche tablets, ampuled liquid medicines, granules, pulvis (powders), tablets, and capsules. Specific examples of medicines for the oral cavity include sprays, ointments, creams, pastes and patches. Specific examples of oral care products include liquid dentifrices, toothpastes, mouthwashes, and breath
10 fresheners.

[0130]

Stevia extract and sweet flavor may be added at the same time or individually in the process of producing these compositions.

15 [0131]

The amount of stevia extract added to these compositions depends on the type of composition, etc., but is generally 10 ppm to 5,000 ppm, preferably 25 ppm to 3,000 ppm, and more preferably 50 ppm to 2,000 ppm. To stevia extract of
20 such amounts, sweet flavor may be added so that their proportion falls within the above-mentioned ranges. More specifically, for example, when the amount of stevia extract added to the composition is 600 ppm, the amount of sweet flavor added is generally 0.1 to 3 ppm, and preferably 0.1 to 2 ppm.

25 [0132]

(III-2) Sweetener

The sweetener of the present invention contains stevia extract and the sweet flavor described earlier. One of the features of the sweetener of the present invention is that it
30 contains 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor. The amount of stevia extract per part by mass of sweet flavor is preferably 300 to 6,000 parts by mass, more preferably 400 to 6,000 parts by mass, and particularly preferably 600 to 4,000 parts by mass.

35 [0133]

There is no limitation to the form of the sweetener of the present invention, and the sweetener may take any form including a powder, a granule, a tablet, a capsule and like solid states; or syrup, emulsion, liquid, gel and like semi-solid or liquid states.

[0134]

The sweetener of the present invention can be prepared by mixing stevia extract and sweet flavor so that their compounding ratio falls within the range described above. During the preparation, depending on the form, pharmacologically acceptable carriers or carriers that may be added to foods and beverages may also be added. Examples of such carriers include lactose, glucose, fructose, sugar, high-fructose corn syrup and like saccharides; sorbitol, erythritol, lactitol, maltitol, mannitol, xylitol, isomalt and like sugar alcohols; isomalto-oligosaccharide, galacto-oligosaccharide, fructo-oligosaccharide and like oligosaccharides; dextrin, cellulose, gum arabic, cornstarch and like polysaccharides; water; etc.

[0135]

Furthermore, flavors, colorants, acidulants, preservatives and other components that are ordinarily added to foods, beverages or medicines may be added. If desired, it is also possible to add aspartame, sucralose, acesulfame potassium and like artificial sweeteners; or licorice extract, amacha extract, brazzein, neohesperidine dihydrochalcone, serendipity berry extract (monellin), tenryocha extract and like natural sweeteners.

[0136]

The total amount of sweet flavor and stevia extract in the sweetener is suitably adjusted so as to fall within the range of 0.1 to 100% by mass.

[0137]

(III-3) Composition Orally Administered or Used For the Oral Cavity

The composition orally administered or used for the

oral cavity of the present invention has a sweet taste and contains stevia extract and sweet flavor as described in (III-1), wherein the amount of the stevia extract is 200 to 6,000 parts by mass per part by mass of sweet flavor. The amount of stevia
5 extract per part by mass of sweet flavor is preferably 300 to 6,000 parts by mass, more preferably 400 to 6,000 parts by mass, and particularly preferably 600 to 4,000 parts by mass.

[0138]

The composition targeted by the present invention is a
10 composition that is orally administered or used for the oral cavity that has sweetness. Examples of such compositions include those mentioned in (III-1), such as foods and beverages; orally administered medicines; medicines for the oral cavity; dentifrices, mouthwashes and like oral care products (including
15 medicines or quasi drugs). Among these, foods and beverages are preferable.

[0139]

The content of stevia extract in these compositions depends on the type of composition, but is generally 10 ppm to
20 5,000 ppm, preferably 25 ppm to 3,000 ppm, and more preferably 50 ppm to 2,000 ppm. To stevia extract of such amounts, sweet flavor may be added so that their compounding ratio falls within the above-mentioned ranges. More specifically, for example, when the amount of stevia extract added to the composition is 600 ppm, the
25 amount of sweet flavor added is generally 0.1 to 3 ppm, and preferably 0.1 to 2 ppm.

[0140]

The stevia extract and sweet flavor may be added in any step of producing a composition that is orally administered or
30 used for the oral cavity.

[0141]

The composition of the present invention thus prepared solves the problem of the sweetness quality in stevia extract, i.e., slow sweetness expression, and provides an excellent
35 sweetness quality similar to that of sugar.

EXAMPLES

[0142]

The present invention is explained in detail below with
5 reference to the Examples described below. However, the scope of
the present invention is not limited to these Examples. The term
"parts" indicates "parts by mass" and "%" indicates "% by mass"
unless otherwise specified. The "*" symbol in the specification
indicates a product of San-Ei Gen F.F.I., Inc., and "†" indicates
10 a registered trademark of San-Ei Gen F.F.I., Inc.

[0143]

Unless otherwise specified, in the Examples below,
"stevia extract" means "stevia extract (Rebaudio J-100, produced
by Morita Kagaku Kogyo Co., Ltd.) that contains 95% by mass of
15 rebaudioside A, and "siraitia grosvenorii extract" means siraitia
grosvenori extract that contains 15% by mass of mogroside V. The
siraitia grosvenorii extract was prepared by freeze-drying a
water extract of dried siraitia grosvenorii. Examples 1 and 2
were conducted using the "stevia extract" and "siraitia
20 grosvenori extract" described above as the "rebaudioside A" and
"mogroside V" respectively. In Examples 1 and 2, the compounding
ratios and amounts of rebaudioside A and mogroside V are
calculated based on the amount of rebaudioside A itself and that
of mogroside V itself.

25 [0144]

(I) Improvement in Sweetness Quality of Rebaudioside A

Example 1: Evaluation of Sweetness Quality (Without Thaumatin)

(1) Degree of Sweetness (Intensity of Sweetness) of Rebaudioside
A and Mogroside V

30 The degree of sweetness of rebaudioside A and that of
mogroside V depends on the pH. Table 1 shows the concentrations
(%) of rebaudioside A and mogroside V in aqueous solutions (pH3,
pH4.5 and pH7) prepared by dissolving each of rebaudioside A or
mogroside V singly into water to attain the same degree of
35 sweetness (intensity of sweetness) as that of a 5% aqueous sugar

solution. Citric acid and trisodium citrate were used to prepare pH3 and pH4.5 solutions, and sodium bicarbonate was used to prepare pH7 solutions.

[0145]

5 Table 1

Concentrations of rebaudioside A and mogroside V having the same degree of sweetness as that of a 5% aqueous sugar solution

	pH3	pH4.5	pH7
Rebaudioside A singly	0.0228%	0.0219%	0.02%
Mogroside V singly	0.03%	0.03%	0.0255%

[0146]

10 (2) Preparation of Aqueous Test Solutions

Rebaudioside A and mogroside V having the degrees of sweetness described above were mixed at the ratios (weight ratios) shown in Table 2. Water was added to the mixture so as to obtain a degree of sweetness (intensity of sweetness) similar to that of a 5% aqueous sugar solution, preparing eight types of aqueous solutions containing rebaudioside A and/or mogroside V. The pH of each aqueous solution was adjusted to pH3 or pH4.5 using citric acid and trisodium citrate, or to pH7 using sodium bicarbonate.

20 [0147]

(3) Evaluation of Sweetness Quality

Ten panelists were asked to evaluate the sweetness quality of the aqueous solutions that were prepared above. Specifically, the sweetness quality of an aqueous solution (Control) containing only rebaudioside A (i.e., rebaudioside A:mogroside V = 100:0) was set to 0 points. As the sweetness quality of an aqueous test solution came closer to that of a 5% aqueous sugar solution, points were added one by one, and when the sweetness quality became most similar to that of a 5% aqueous sugar solution, the sweetness quality was given 5 points. In the same manner, using the sweetness quality of the Control as

reference (0 points), the sweetness quality score was lowered one point at a time as the sweetness quality of an aqueous test solution became further from that of a 5% aqueous sugar solution. The aqueous test solution having a sweetness quality that was the furthest from that of the 5% aqueous sugar solution was given -5 points. Table 2 shows the average scores of the ten panelists.

[0148]

Table 2

Rebaudioside A: Mogroside V (Weight Ratio)	pH3	pH4.5	pH7
100:0 (Control)	0	0	0
87:13	+2.2	+1.8	+2.0
75:25	+1.5	+1.4	+2.0
64:36	+0.3	+0.3	-0.1
53:47	-0.7	-0.2	-0.5
43:57	-1.3	-0.9	-1.5
16:84	-3.4	-2.8	-3.1
0:100	-4.0	-3.4	-3.8

10 [0149]

As seen by the results, the sweetness quality of rebaudioside A improved and became closer to that of sugar by adding 36 parts or less of mogroside V per 64 parts of rebaudioside A. In contrast, the sweetness quality of rebaudioside A started becoming further from that of sugar when 47 parts or more of mogroside V was added per 53 parts of rebaudioside A.

[0150]

Example 2: Evaluation of Sweetness Quality (with Thaumatin Added)

20 (1) Preparation of Aqueous Test Solutions

Rebaudioside A and mogroside V were mixed in the same manner as in Example 1 with the compounding ratio (weight ratio) shown in Table 3. Water was added to the mixture so as to obtain a degree of sweetness (intensity of sweetness) similar to that of a 5% aqueous sugar solution, and eight types of aqueous solutions containing rebaudioside A and/or mogroside V were prepared. Among these aqueous test solutions, thaumatin was added to each aqueous

solution, except for an aqueous solution consisting of 100 parts by mass of rebaudioside A (Control, rebaudioside A:mogroside V = 100:0), so that its concentration became 1 ppm. The pH of each aqueous solution was adjusted to 3 and 4.5 using citric acid and trisodium citrate, and to 7 using sodium bicarbonate.

[0151]

(2) Evaluation of Sweetness Quality

In the same manner as in Example 1, ten panelists were asked to evaluate the sweetness quality of each of the aqueous test solutions prepared above.

[0152]

Table 3

Rebaudioside A: Mogroside V (Weight Ratio)	Thaumatococin Added	pH3	pH4.5	pH7
100:0 (Control)	-	0	0	0
87:13	1 ppm	+3.5	+3.4	+3.4
75:25	1 ppm	+4.0	+2.2	+2.7
64:36	1 ppm	+1.2	+1.1	+2.4
53:47	1 ppm	-0.6	+0.4	+0.6
43:57	1 ppm	-1.1	-1.1	-0.6
16:84	1 ppm	-2.0	-2.1	-1.8
0:100	1 ppm	-2.8	-2.9	-2.2

[0153]

As seen by the results, the sweetness quality of rebaudioside A improved and became closer to that of sugar by adding 47 parts or less of mogroside V per 53 parts of rebaudioside A, and preferably 36 parts or less of mogroside V per 64 parts of rebaudioside A, in the presence of 1 ppm thaumatococin. The results also indicate that the sweetness quality started becoming further from that of sugar when 57 parts or more of mogroside V was added per 43 parts of rebaudioside A. A comparison of the results of Example 1 (Table 2) and those shown in Table 3 also make it clear that the sweetness quality was further improved and became closer to that of sugar when mogroside V and rebaudioside A were used in combination with thaumatococin.

[0154]

Examples 3 and 4: Sweeteners Containing Rebaudioside A

According to the formulae shown in Table 4, various components were mixed to prepare three types of sweeteners

5 (Comparative Example, and Examples 3 and 4). The proportion of rebaudioside A to mogroside V contained in the sweeteners of Examples 3 and 4 was 86:14 (weight ratio). These sweeteners were adjusted to have a degree of sweetness (sweetness intensity relative to sugar) of about 100 times that of sugar.

10 [0155]

Table 4

(Unit: Part by mass)

	Comparative Example	Example 3	Example 4
Isomalt	58.0	22.2	22.1
Stevia Extract (Rebaudioside A)	42.0 (39.9)	37.8 (35.91)	37.8 (35.91)
Siraitia Grosvenorii Extract (Mogroside V)	-	40.0 (6.00)	40.0 (6.00)
Thaumatococcus	-	-	0.1
Total	100.0	100.0	100.0

[0156]

Each of the obtained sweeteners (0.05 g) was dissolved
15 in water (100 mL). Ten panelists were asked to evaluate the taste (speed of sweetness expression, lingering sweetness, and presence or absence of bitterness) of the test solutions. Table 5 shows the results of the evaluation. The ten panelists were also asked to compare the taste (speed of sweetness expression, lingering
20 sweetness, and presence or absence of bitterness) with that of a Control solution obtained by dissolving 5 g of sugar in 100 mL of water (a 5% aqueous sugar solution: Control). Table 5 shows the results.

[0157]

Table 5

Comparative Example	Slower sweetness expression than Control. Lingering sweetness (remaining aftertaste) and distinctive bitterness.
Example 3	Compared to the Comparative Example, the sweetness expression was faster and became closer to that of sugar. The lingering sweetness observed in the Comparative Example was reduced. The bitterness observed in the Comparative Example was reduced (bitterness was masked).
Example 4	Compared to the Comparative Example, the sweetness expression was faster and became closer to that of sugar. The lingering sweetness observed in the Comparative Example was reduced, and the aftertaste was improved. The bitterness observed in the Comparative Example did not exist (bitterness was masked).

5 [0158]

The results indicate that, by using rebaudioside A together with mogroside V, the speed of sweetness expression felt in the oral cavity became closer to that of sugar, and the lingering sweetness was reduced. By also using thaumatin in
10 combination with rebaudioside A and mogroside V, the lingering sweetness could be further reduced and the bitterness was masked, so that the sweetness quality became closer to that of sugar.

[0159]

Example 5: Coffee Beverages

15 (1) Preparation of Sweeteners

According to the formulae shown in Table 6, five types of sweeteners (Control, Comparative Example, and Examples 5-1 to 5-3) were prepared. The proportion of rebaudioside A to mogroside V in the sweeteners of Examples 5-1 to 5-3 was adjusted to 86:14
20 (weight ratio).

[0160]

Table 6

(Unit: Part by mass)

	Sweeteners				
	Control	Comparative Example	Example 5-1	Example 5-2	Example 5-3
Sugar	5.0	-	-	-	1.7
Stevia Extract (Rebaudioside A)	-	0.021 (0.01995)	0.019 (0.01805)	0.019 (0.01805)	0.013 (0.01235)
Siraitia Grosvenorii Extract (Mogroside V)	- (-)	- (-)	0.02 (0.003)	0.02 (0.003)	0.013 (0.00195)
Thaumatococcus containing preparation (Thaumatococcus)	- (-)	- (-)	- (-)	0.03 (0.000045)	0.03 (0.000045)

- 5 * Thaumatococcus containing preparation: A preparation that contained 0.15% of thaumatococcus (NEO SAN MARUKU[†] AG: produced by San-Ei Gen F.F.I., Inc.)

[0161]

10 (2) Preparation of Coffee Beverages

According to the formulae shown in Table 7, powder mixtures of the sweeteners (Table 6) and an emulsifier were added to water. Each mixture was dissolved while stirring at 80°C for 10 minutes and then cooled to room temperature. Cow's milk, sodium bicarbonate, and coffee were sequentially added to each mixture. The pH was adjusted to 6.8, and water was added thereto to make the total amount 100 parts. The resulting mixtures were heated to 75°C, and homogenized using a homogenizer. Coffee flavor was added to the mixtures. After being placed in a container, each mixture was subjected to retort-sterilization at 121°C for 20 minutes.

[0162]

Table 7

(Unit: Part by Mass)

Coffee (L = 20, Brix = 3)	41.0
Cow's milk	10.0
Sweeteners (Control, Comparative Example, and Examples 5-1 to 5-3)	Table 6
Emulsifier (Homogen [†] No. 1379*)	0.14
pH Controlling agent (Sodium bicarbonate)	q.s. (pH6.8)
Flavor (Coffee Flavor No. 90154*)	0.01
Water	Balance
Total	100.0

5

[0163]

(3) Evaluation of Coffee Beverages (Sensory Evaluation)

Ten panelists were asked to taste the coffee beverages obtained above (depending on the sweeteners used, the coffee beverages were categorized into Control, Comparative Example, Examples 5-1, 5-2, and 5-3) to evaluate their taste (speed of sweetness expression, lingering sweetness, and presence or absence of bitterness). Table 8 shows the evaluation results. The evaluation was conducted by using the taste (speed of sweetness expression, lingering sweetness, and presence or absence of bitterness) of a coffee beverage prepared using only sugar as the sweetener (Control) as reference and comparing the taste of each test sample with that of the Control.

10

15

[0164]

Table 8

Coffee Beverages	Evaluations
Comparative Example	Slower sweetness expression than the Control. Lingering sweetness and distinctive bitterness.
Example 5-1	Compared to the Comparative Example, the sweetness expression was faster and became closer to that of sugar. The lingering sweetness observed in the Comparative Example was reduced. The bitterness observed in the Comparative Example was reduced (bitterness was masked).
Example 5-2	Compared to the Comparative Example, the sweetness expression was faster and became closer to that of sugar. The lingering sweetness observed in the Comparative Example was reduced, and the aftertaste was improved. The bitterness observed in the Comparative Example did not exist (bitterness was masked).
Example 5-3	The sweetness expression was as fast as that of the Control. Lingering sweetness and bitterness did not exist and the sweetness quality was almost the same as that of the Control.

5 [0165]

Example 6: Beverages Containing Orange Juice

(1) Preparation of Sweeteners

According to the formulae shown in Table 9, five types of sweeteners (Control, and Examples 6-1 to 6-4) were prepared.

10 The proportion of rebaudioside A to mogroside V in the sweeteners of Examples 6-1 to 6-4 was adjusted to 79:21 (weight ratio).

[0166]

Table 9

(Unit: Part by mass)

	Sweeteners				
	Control	Example 6-1	Example 6-2	Example 6-3	Example 6-4
Sugar	7.0	-	-	3.5	3.5
Stevia Extract (Rebaudioside A)	-	0.033 (0.03135)	0.033 (0.03135)	0.013 (0.01235)	0.013 (0.01235)
Siraitia Grosvenorii Extract (Mogroside V)	- (-)	0.056 (0.0084)	0.056 (0.0084)	0.02 (0.003)	0.02 (0.003)
Thaumatococcus-containing preparation (Thaumatococcus)	- (-)	- (-)	0.05 (0.000075)	- (-)	0.05 (0.000075)

- 5 * Thaumatococcus-containing preparation: A preparation that contained 0.15% of thaumatococcus (NEO SAN MARUKU[†] AG: produced by San-Ei Gen F.F.I., Inc.)

[0167]

10 (2) Preparation of Beverages Containing Orange Juice

According to the formulae shown in Table 10, 5-fold concentrated orange juice, citric acid (anhydrous) fine particles, trisodium citrate fine particles and each of the sweeteners of Table 9 were mixed. The resulting mixtures were dissolved in
15 water while heating and stirring. After heating the mixtures to 93°C, orange flavor was added thereto, water was added to make the total amount 100 parts, and each of the resulting mixtures was placed in a container.

[0168]

20 Table 10

(Unit: Part by mass)

5-Fold concentrated orange juice	6.6
Citric acid (anhydrous) fine particles*	0.15
Trisodium citrate fine particles*	0.03
Sweeteners (Control, and Examples 6-1 to 6-4)	Table 9
Flavor (Orange Flavor No. 2410*)	0.10
Water	Balance
Total	100.0

[0169]

(3) Evaluation of Beverages Containing Orange Juice (Sensory Evaluation)

Twelve panelists were asked to taste the obtained beverages containing orange juice (depending on the sweeteners used, the beverages containing orange juice were categorized into Control, Comparative Example, and Examples 6-1, 6-2, 6-3, and 6-4) to evaluate their taste (speed of sweetness expression, lingering sweetness, and presence or absence of bitterness). Table 11 shows the evaluation results. The evaluation was conducted using the taste (speed of sweetness expression, lingering sweetness, and presence or absence of bitterness) of a beverage containing orange juice prepared using only sugar as the sweetener (Control) as reference and comparing the taste of each test sample with that of the Control.

[0170]

Table 11

Beverages Containing Orange Juice	Evaluations
Example 6-1	Fast sweetness expression similar to that of the Control, but a slightly bitter aftertaste.
Example 6-2	Fast sweetness expression similar to that of the Control. The bitterness observed in Example 6-1 was reduced, and the aftertaste was improved.
Example 6-3	The sweetness expression was as fast as that of the Control. The sweet quality was similar to that of Example 6-2, but there was a slightly bitter aftertaste.
Example 6-4	The sweetness expression was as fast as that of the Control. The sweetness quality was almost the same as that of the Control.

[0171]

Example 7: Apple Jelly

(1) Preparation of Sweeteners

According to the formulae shown in Table 12, four types of sweeteners (Control, Comparative Example, and Examples 7-1 and 7-2) were prepared. The proportion of rebaudioside A to mogroside

V in the sweeteners of Examples 7-1 and 7-2 was adjusted to 89:11 (weight ratio).

[0172]

Table 12

5

(Unit: Part by mass)

	Control	Comparative Example	Example 7-1	Example 7-2
Sugar	5.0	2.0	2.0	2.0
Stevia Extract (Rebaudioside A)	-	0.0135 (0.012825)	0.012 (0.0114)	0.012 (0.0114)
Siraitia Grosvenorii Extract (Mogroside V)	-	- (-)	0.009 (0.00135)	0.009 (0.00135)
Thaumatococcus containing preparation (Thaumatococcus)	-	- (-)	- (-)	0.05 (0.000075)

* Thaumatococcus containing preparation: A preparation that contained 0.15% of thaumatococcus (NEO SAN MARUKU[†] AG: produced by San-Ei Gen F.F.I., Inc.)

10 [0173]

(2) Preparation of Apple Jelly

According to the formulae shown in Table 13, a high-fructose corn syrup was added to water, then a powder mixture of each of the sweeteners of Table 12 and a gelling agent was added while stirring, and the resulting mixture was dissolved while stirring at 80°C for 10 minutes. Five-fold concentrated apple juice (clarified), trisodium citrate fine particles, citric acid (anhydrous) medium-size particles and apple flavor were added to the mixture, and water was added to make the total amount 100 parts. Each of the resulting mixtures was placed in a container, and subjected to sterilization at 85°C for 30 minutes.

[0174]

Table 13

(Unit: Part by mass)

High-fructose corn syrup	15.0
Sweeteners (Control, Comparative Example, and Examples 7-1 and 7-2)	Table 12
Five-fold concentrated apple juice (clarified)	4.2
Gelling agent (GEL UP [†] WM-100*)	1
Citric acid (anhydrous) medium-size particles	0.25
Trisodium citrate fine particles	0.03
Flavor (Apple Flavor No. 2414*)	0.1
Water	Balance
Total	100.0

5

[0175]

(3) Evaluation of Apple Jelly (Sensory Evaluation)

Twelve panelists were asked to taste the obtained apple jelly (depending on the sweeteners used, the apple jelly was categorized into Control, Comparative Example, and Examples 7-1 and 7-2) to evaluate the taste (lingering sweetness, rough taste (presence of bitterness and/or astringency)). Table 14 shows the results. The evaluation was conducted using the taste (lingering sweetness, and presence of rough taste) of an apple jelly prepared using only sugar and a high-fructose corn syrup as sweeteners (Control) as reference and comparing the taste of each test sample with that of the Control.

10

15

[0176]

Table 14

Apple Jelly	Evaluations
Comparative Example	There was a sweet aftertaste, and a rough aftertaste (bitterness and astringency)
Example 7-1	The sweet aftertaste observed in the Comparative Example was reduced. There was a rough aftertaste (bitterness and astringency).
Example 7-2	The sweet aftertaste and rough aftertaste observed in the Comparative Example and Example 7-1 were reduced. The sweet quality was almost the same as that of the Control (sugar).

[0177]

Example 8: Seasonings (Seasonings for Preparing Lightly Pickled Foods)

5 (1) Preparation of Sweeteners

According to the formulae shown in Table 15, three types of sweeteners (Control, and Examples 8-1 and 8-2) were prepared. The proportion of rebaudioside A to mogroside V in the sweeteners of Examples 8-1 and 8-2 was adjusted to 94:6 (weight
10 ratio).

[0178]

Table 15

(Unit: Part by mass)

	Control	Example 8-1	Example 8-2
High-fructose corn syrup	13.0	-	-
Stevia Extract (Rebaudioside A)	-	0.073 (0.06935)	0.073 (0.06935)
Siraitia Grosvenorii Extract (Mogroside V)	-	0.032 (0.0048)	0.032 (0.0048)
Thaumatococcus preparation (Thaumatococcus)	-	- (-)	0.05 (0.000075)

* Thaumatococcus-containing preparation: A preparation that contained
15 0.15% of thaumatococcus (NEO SAN MARUKU[†] AG: produced by San-Ei Gen
F.F.I., Inc.)

[0179]

20 (2) Preparation of Seasonings (Seasonings for Preparing Lightly Pickled Foods)

According to the formulae shown in Table 16, each of the sweeteners of Table 15 and other components were added to water, followed by stirring and dissolving the resulting mixture to produce seasonings for preparing lightly pickled foods.

[0180]

Table 16

(Unit: Part by mass)

Sweetener	Table 15
Light soy sauce	5.0
Fermented alcohol beverage (Acidity of 4.2%)	12.0
Dietary salt	6.0
L-Sodium glutamate	0.5
Disodium succinate	0.2
Seasoning (SAN-LIKE TASTE BASE A*)	0.4
Flavor(SAN-AROMA RED PEPPER SV-3541(N)*)	0.004
Water	Balance
Total	100.0

5

[0181]

(3) Evaluation (Sensory Evaluation) for Seasonings (Seasonings for Preparing Lightly Pickled Foods)

Daikon radish was soaked in each of the obtained
 10 seasonings (depending on the sweetener used, they are categorized
 into Control, and Examples 8-1 and 8-2) in the proportion of 1:1
 (weight ratio) daikon radish to seasoning. After one day of
 soaking, the daikon radish (lightly pickled daikon radish) was
 removed from the seasoning. Ten panelists were asked to taste
 15 the resulting pickles. Table 17 shows the evaluation results.
 The evaluation was conducted using daikon radish (lightly pickled
 daikon radish) soaked in a seasoning prepared using only a high-
 fructose corn syrup as the sweetener (Control) as reference, and
 the taste of each test sample was compared with that of the
 20 Control.

[0182]

Table 17

Seasoning	Evaluation of the taste of daikon radish soaked in each seasoning (lightly pickled daikon radish)
Example 8-1	The soy sauce flavor was improved compared to that of the Control (seasoning). There was a slightly rough aftertaste.
Example 8-2	The rough aftertaste observed in Example 8-1 had disappeared. There was an increased salty taste.

[0183]

As is clear from the results of Examples 5 to 8, the use of a sweetener that contains rebaudioside A and mogroside V at a weight ratio of 75:25 to 94:6 can render an excellent sweetness, i.e., a sweetness quality similar to that of sugar, to foods, beverages and like orally consumed compositions. It was also confirmed that, by adding thaumatin to a sweetener having the above-mentioned ratio, a more improved sweetness can be obtained that is closer to the sweetness quality of sugar.

[0184]

(II) Improvement in Sweetness Quality of Stevia Extract

Examples 9 to 13: Zero-Calorie Carbonated Beverages

(1) Preparation of carbonated beverages

According to the formulae shown in Table 18, lemon juice, stevia extract, citric acid (anhydrous), trisodium citrate, colorant, and lemon flavoring, in addition to at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate, were added to water and dissolved therein, to prepare syrups for carbonated beverages.

[0185]

After cooling, 80 mL of carbonated water was added to 20 mL of syrup, making the total amount 100 mL. Each of the resulting liquids was placed in a container and sterilized at 70°C for 20 minutes, to prepare a carbonated beverage.

[0186]

Table 18

(Part by mass)

	Example					Comp. Ex.
	9	10	11	12	13	
Lemon juice	4.2	4.2	4.2	4.2	4.2	4.2
Stevia extract	0.033	0.033	0.033	0.033	0.033	0.033
Citric acid (anhydrous)	0.2	0.2	0.2	0.2	0.2	0.2
Trisodium citrate	0.1	0.1	0.1	0.1	0.1	0.1
Additives	Gentiooligosaccharide ¹⁾	0.4	-	-	-	-
	Naringin	-	0.0025	-	-	-
	Magnesium chloride	-	-	0.05	-	-
	Magnesium sulfate	-	-	-	0.05	-
	Highly branched cyclic dextrin ²⁾	-	-	-	-	0.07
Colorant (SAN YELLOW [®] No.2 SFU*)	0.009	0.009	0.009	0.009	0.009	0.009
Flavor (Lemon flavor)	0.1	0.1	0.1	0.1	0.1	0.1
Water	Balance	Balance	Balance	Balance	Balance	Balance
Total	20.0	20.0	20.0	20.0	20.0	20.0

- 5 1) Gentose#45: Produced by Nihon Shokuhin Kako Co., Ltd., containing 45% or more gentiooligosaccharide.
2) Cluster Dextrin: Produced by Nihon Shokuhin Kako Co., Ltd.

[0187]

10 (2) Evaluation Results

Ten panelists were asked to taste each carbonated beverage prepared above. Table 19 shows the evaluation results.

[0188]

Table 19

Carbonated Beverage	Additive Used	Sensory Evaluation Result
Comparative Example	-	There was a sweet aftertaste. There was insufficient richness (body), and the taste was not delicious.
Example 9	Gentiooligosaccharide ¹⁾	The sweet aftertaste observed in the Comparative Example was reduced. Richness was added to the sweetness.
Example 10	Naringin	The sweet aftertaste observed in the Comparative Example was reduced. The juiciness was increased.
Example 11	Magnesium chloride	The sweet aftertaste observed in the Comparative Example was reduced. There was a refreshing aftertaste.
Example 12	Magnesium sulfate	The sweet aftertaste observed in the Comparative Example was reduced. There was an increased sparkling effect of the carbonated beverages.
Example 13	Highly branched cyclic dextrin ²⁾	The sweet aftertaste observed in the Comparative Example was reduced. Richness was added to the sweetness, and there was a refreshing aftertaste.

1) Gentose#45: Produced by Nihon Shokuhin Kako Co., Ltd., containing 45% or more gentiooligosaccharide.

5 2) Cluster Dextrin: Produced by Nihon Shokuhin Kako Co., Ltd.

[0189]

As described above, by adding at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrans, magnesium chloride and magnesium sulfate to a stevia extract-containing carbonated beverage, the lingering sweetness of the stevia extract can be

10

reduced and the richness (body), juiciness, and sparkling effect of carbonated beverages can be increased.

[0190]

Example 14

5 Aqueous solutions containing various sweeteners (stevia extract, siraitia grosvenori extract, licorice extract, thaumatin, aspartame, and neotame) in the concentrations shown in Table 20 were prepared. The concentration of each sweetener was adjusted so as to exhibit the same degree of sweetness as that of a 6%
10 aqueous sugar solution.

[0191]

Table 20

Sweetener	Concentration (%)
Stevia extract	0.03
Siraitia grosvenorii extract ¹⁾	0.05
Licorice extract ²⁾	0.05
Thaumatococcus	0.0024
Aspartame	0.038
Neotame ³⁾	0.0008

1) LUO HAN GUO CONCENTRATE EXTRACT POWDER (containing 15% mogroside V: Produced by San-Ei Gen F.F.I., Inc.)

15 2) Licotin P1 (produced by Ikeda Tohka Industries Co., Ltd.)

3) Mirasee (containing 2% neotame, produced by Dainippon Sumitomo Pharma Co., Ltd.)

[0192]

20 Subsequently, various additives (gentiooligosaccharide, naringin, highly branched cyclic dextrin, magnesium chloride, and magnesium sulfate) were added to each of the above aqueous solutions so that the final concentrations of the additives became as shown in Table 21, thus preparing 30 kinds of aqueous
25 solutions.

[0193]

Table 21

Additives	Concentration (%)
Gentiooligosaccharide ¹⁾	0.4
Naringin	0.0025
Highly branched cyclic dextrin ²⁾	0.07
Magnesium chloride	0.05
Magnesium sulfate	0.05

1) gentiooligosaccharide syrup (Gentose#45: Produced by Nihon Shokuhin Kako Co., Ltd.,)

5 2) Cluster Dextrin: Produced by Nihon Shokuhin Kako Co., Ltd.

[0194]

Twelve panelists were asked to taste the resulting 30 kinds of aqueous solutions to evaluate the improvement in the sweetness quality of each sweetener attributable to each additive. The evaluation was conducted by scoring the improvement in sweetness quality based on the criteria described below in comparison with an aqueous solution containing only each of the sweeteners (Control).

15 [0195]

Criteria

Improvement in sweetness quality: 2 points

Slight improvement in sweetness quality: 1 point

No improvement in sweetness quality: 0 point

20 [0196]

Table 22 shows the average scores by the twelve panelists.

[0197]

Table 22

	Gentiooligo saccharide	Naringin	Highly branched cyclic dextrin	Magnesium chloride	Magnesium sulfate
Stevia extract	1.7	1.3	1.8	1.7	1.8
Siraitia grosvenorii extract	1.0	0.7	0.7	1.5	0.5
Licorice extract	0.7	0.7	0.7	1.3	1.0
Thaumatococcus	0.8	0.7	0.5	1.0	1.3
Aspartame	0.3	0.3	0.3	0.8	0.7
Neotame	0.8	1.0	0.8	0.8	0.8

[0198]

5 As shown in Table 22, all of the additives
(gentiooligosaccharide, naringin, highly branched cyclic dextrin,
magnesium chloride, and magnesium sulfate) reduced the remaining
aftertaste of sweetness in stevia extract, and increased the
richness (body) thereof. Among these, gentiooligosaccharide,
10 magnesium chloride, magnesium sulfate, and highly branched cyclic
dextrin exhibited remarkably excellent improvements in the
sweetness quality of stevia extract.

[0199]

15 However, these additives did not always improve the
sweetness quality of other sweeteners (siraitia grosvenorii
extract, licorice extract, thaumatococcus, aspartame, and neotame).

[0200]

(III) Acceleration of Sweetness Expression in Stevia Extract

Example 15

20 Maltol was added to a 400-ppm aqueous solution of
stevia extract in the proportions shown in Table 23 to prepare an
aqueous solution containing stevia extract and maltol. A 400-ppm
aqueous solution of stevia extract was prepared as a Control.

[0201]

25 Eight panelists were asked to taste the resulting
solutions to select the solutions that exhibited accelerated
sweetness expression as compared to the Control. The results are

shown in Table 23 together with the proportions of maltol.

[0202]

Table 23

Stevia extract (ppm)	400	400	400	400	400	400	400	400
Maltol (ppm)	0.01	0.1	0.2	0.5	1	2	5	10
Proportion (Stevia extract: Maltol)	40000:1	40000:1	20000:1	800:1	400:1	200:1	80:1	40:1
The number of panelists who answered that the expression of sweetness became faster and more favorable compared to the Control	2	7	7	6	5	4	2	0

5 [0203]

As shown in Table 23, more than half of the panelists answered that aqueous solutions containing at least 200 to 4,000 parts of stevia extract per part of maltol were more favorable in exhibiting quick sweetness expression in an oral cavity than an aqueous stevia extract solution (Control). An increased number of panelists answered that when the proportion of stevia extract was less than 200 parts per part of maltol, the smell of maltol became conspicuous and an unfavorable lingering sweetness (a remaining aftertaste) increased.

15 [0204]

Example 16

Maltol was added to an aqueous solution of enzyme-treated stevia extract (SK Sweet FZ, produced by Nippon Paper Chemicals Co., Ltd., 600 ppm) in the proportions shown in Table 24 to prepare an aqueous solution containing enzyme-treated stevia extract and maltol. As a Control, a 600-ppm aqueous solution of enzyme-treated stevia extract was prepared. The 600-ppm aqueous solution of enzyme-treated stevia extract had almost the same degree of sweetness as the 400-ppm aqueous solution of stevia extract used in Example 15.

[0205]

Eight panelists were asked to taste the stevia extracts to select the aqueous solutions that exhibited accelerated sweetness expression in an oral cavity compared to the Control (an aqueous solution of enzyme-treated stevia extract). Table 24 shows the results together with the proportions of maltol.

[0206]

Table 24

Enzyme-treated stevia extract (ppm)	600	600	600	600	600	600	600	600
Maltol (ppm)	0.01	0.1	0.2	0.5	1	2	5	10
Proportion (Stevia extract: Maltol)	60000:1	60000:1	30000:1	1200:1	600:1	300:1	120:1	60:1
The number of panelists who answered that the sweetness expression became faster	3	4	6	8	7	4	3	1

10 [0207]

As shown in Table 24, more than half of the panelists answered that aqueous solutions containing at least 300 to 6,000 parts of enzyme-treated stevia extract per part of maltol were favorable in exhibiting a faster sweetness expression in an oral cavity than an aqueous enzyme-treated stevia extract solution (Control). An increased number of panelists answered that when the proportion of enzyme-treated stevia extract was less than 300 parts per part of maltol, the smell of maltol became conspicuous and an unfavorable lingering sweetness (a remaining aftertaste) increased.

[0208]

The results of Example 15 (Table 23) and Example 16 (Table 24) indicate that adding 200 to 6,000 parts, and preferably 300 to 6,000 parts, of stevia extract per part of maltol is effective for accelerating the sweetness expression of stevia extract.

[0209]

Example 17: Carbonated Beverages Containing Lemon Juice

(1) Preparation of carbonated beverages

Lemon juice, citric acid (anhydrous) fine particles,
5 trisodium citrate fine particles, stevia extract, 1% maltol and
lemon flavoring were mixed in the proportion of the formula of
Example 17 shown in Table 25. Water was added to the mixture so
that the total amount of the water and mixed components became 20
parts, and the components were dissolved while stirring.
10 Subsequently, carbonated water was added thereto to make the
total amount 100 parts and the resulting mixture was placed in a
container, obtaining a carbonated beverage containing lemon juice
(Example 17). For comparison (Comparative Example), a carbonated
beverage containing lemon juice was prepared in the same manner
15 as Example 17 except that 1% maltol was not added.

[0210]

Table 25

	Example 17	Comparative Example
Lemon juice	4.2	4.2
Stevia extract	0.033	0.033
1% Maltol	0.003	-
Citric acid (anhydrous) fine particles*	0.2	0.2
Trisodium citrate fine particles*	0.1	0.1
Flavor (Lemon flavoring)	0.1	0.1
Total with water	20.0	20.0
Carbonated water	80.0	80.0
Total	100.0	100.0

[0211]

20 (2) Evaluation of Carbonated Beverages (Sensory Evaluation)

Ten panelists were asked to taste the resulting
carbonated beverages (Example 17 and the Comparative Example) to
evaluate the sweetness expression speed of the carbonated
beverage of Example 17. The evaluation was conducted by comparing
25 the sweetness expression speed of the carbonated beverage with
that of the Comparative Example.

[0212]

As a result of the sensory evaluation, the Comparative Example, which was prepared without adding maltol, exhibited a slow sweetness expression, and lingering sweetness was observed.

- 5 In contrast, the carbonated beverage of Example 17, to which 0.3 ppm of maltol was added per 330 ppm of stevia extract, exhibited an accelerated sweetness expression, and the lingering sweetness was prevented. The carbonated beverage of Example 17 also achieved a sweetness quality similar to that of sugar.

10

INDUSTRIAL APPLICABILITY

[0213]

- 15 The present invention provides sweeteners, orally administered compositions and compositions used for the oral cavity (foods, beverages, medicines, and quasi drugs (including oral care products)) that contain a natural sweetener having an excellent sweetness quality.

Claims

1. A sweetener comprising rebaudioside A and mogroside V at a weight ratio of 95:5 to 60:40 (rebaudioside A:mogroside V).

5 2. The sweetener according to Claim 1, which further comprises 0.01 to 3 parts by mass of thaumatin per total 100 parts by mass of rebaudioside A and mogroside V.

10 3. A composition orally administered or used for an oral cavity having sweetness and containing rebaudioside A and mogroside V at a weight ratio of 95:5 to 60:40.

15 4. The composition according to Claim 3, which comprises rebaudioside A and mogroside V in a total amount of 10 ppm to 5,000 ppm.

20 5. The composition according to Claim 3, which further comprises 0.01 to 3 parts by mass of thaumatin per total 100 parts by mass of rebaudioside A and mogroside V.

 6. A method for applying sweetness to a composition orally administered or used for an oral cavity comprising the step of:

25 adding rebaudioside A and mogroside V to a composition orally administered or used for the oral cavity so that the weight ratio of rebaudioside A to mogroside V becomes 95:5 to 60:40.

30 7. The method according to Claim 6, wherein rebaudioside A and mogroside V are added to the composition orally administered or used for the oral cavity so that the total content of rebaudioside A and mogroside V becomes 10 ppm to 5,000 ppm.

35 8. The method according to Claim 6, wherein thaumatin

is added to the composition orally administered or used for the oral cavity so that the content of the thaumatin becomes 0.01 to 3 parts by mass per total 100 parts by mass of rebaudioside A and mogroside V.

5

9. The method according to Claim 6, wherein the composition orally administered or used for the oral cavity is a food or beverage, or a medicine or quasi drug that is orally administered or used for the oral cavity.

10

10. A sweetness quality improving agent comprising at least one active ingredient selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrins, magnesium chloride and magnesium sulfate.

15

11. A method for improving sweetness quality of a stevia extract comprising the step of:

adding a sweetness quality improving agent comprising at least one active ingredient selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrins, magnesium chloride and magnesium sulfate to a stevia extract.

12. A sweetener comprising a stevia extract and at least one member selected from the group consisting of gentiooligosaccharides, naringins, highly branched cyclic dextrins, magnesium chloride and magnesium sulfate.

13. A method for accelerating sweetness expression of a stevia extract in an oral cavity comprising the step of:

combining 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor.

14. A sweetener comprising 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor.

15. A composition that is orally administered or used for an oral cavity having sweetness and comprising 200 to 6,000 parts by mass of stevia extract per part by mass of sweet flavor.