

FAST DIAGNOSIS AND PERSONALIZED TREATMENTS FOR ACNE**STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH**

This invention was made with Government support of Grant Nos.

5 UH2AR057503 and R01GM099530, awarded by the National Institutes of Health.

The Government has certain rights in this invention.

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Provisional Patent Application No.

10 61/612,290, filed on March 17, 2012, which is incorporated by reference herein in its entirety for all purposes.

BACKGROUND OF THE INVENTION

Acne is a skin condition that causes pimples or "zits." This includes
15 whiteheads, blackheads, and red, inflamed patches of skin (such as cysts). Acne occurs when tiny pores on the surface of the skin become clogged. Each pore opens to a follicle. A follicle contains a hair and an oil gland. The oil released by the gland helps remove old skin cells and keeps your skin soft. When glands produce too much oil, the pores can become blocked. Dirt, bacteria, and cells build up. The
20 blockage is called a plug or comedone. If the top of the plug is white, it is called a whitehead. If the top of the plug is dark, it is called a blackhead. If the plug breaks open, swelling and red bumps occur. Acne that is deep in your skin can cause hard, painful cysts. This is called cystic acne.

Acne is most common in teenagers, but anyone can get acne. 85% of
25 teenagers have acne. Hormonal changes may cause the skin to be more oily. Acne

tends to run in families. It may be triggered by hormonal changes related to puberty, menstrual periods, pregnancy, birth control pills, or stress; greasy or oily cosmetic and hair products; certain drugs (such as steroids, testosterone, estrogen, and phenytoin); or high levels of humidity and sweating.

5 Various treatments exist for the treatment of acne. In general, acne treatments work by reducing oil production, speeding up skin cell turnover, fighting bacterial infection, reducing the inflammation or doing all four. These types of acne treatments include over-the-counter topical treatments, antibiotics, oral contraceptives and cosmetic procedures. Acne lotions may dry up the oil, kill
10 bacteria and promote sloughing of dead skin cells. Over-the-counter (OTC) lotions are generally mild and contain benzoyl peroxide, sulfur, resorcinol, salicylic acid or sulfur as their active ingredient. Studies have found that using topical benzoyl peroxide along with oral antibiotics may reduce the risk of developing antibiotic resistance. Antibiotics may cause side effects, such as an upset stomach, dizziness
15 or skin discoloration. These drugs also increase your skin's sun sensitivity and may reduce the effectiveness of oral contraceptives. For deep cysts, antibiotics may not be enough. Isotretinoin (Amnesteem, Claravis, Sotret) is a powerful medication available for scarring cystic acne or acne that doesn't respond to other treatments. However, isotretinoin has many side effects, such as dry skin, depression, severe
20 stomach pain, and muscle/joint/back pain, and can cause birth defects in babies whose mothers use isotretinoin. Oral contraceptives, including a combination of norgestimate and ethinyl estradiol (Ortho Tri-Cyclen, Previfem, others), can improve acne in women. However, oral contraceptives may cause other side effects, such as headaches, breast tenderness, nausea, and depression. Chemical peels and
25 microdermabrasion may be helpful in controlling acne. These cosmetic procedures,

which have traditionally been used to lessen the appearance of fine lines, sun damage, and minor facial scars, are most effective when used in combination with other acne treatments. They may cause temporary, severe redness, scaling and blistering, and long-term discoloration of the skin.

5 In addition to the negative side-effects caused by the currently available treatments, there is no treatment available that is personalized to patients to target specific bacteria causing acne on an individual level. Additionally, it will be useful for dermatologists to know which strains are dominant on the skin of a patient at the time of diagnosis in order to personalize acne treatments. Thus, there exists a need
10 in the art for methods of personalized diagnoses and treatment of acne.

BRIEF SUMMARY OF THE INVENTION

The present invention is directed to methods of diagnosis and personalized treatment in patients afflicted with acne.

15 In one embodiment, the invention provides a method for determining whether an individual possesses acne comprising: obtaining a skin sample from an individual; isolating bacterial DNA from said sample; amplifying 16S ribosomal DNA in said sample; sequencing said amplified DNA products; and typing the individual's DNA based on one or more of the ten major ribotypes (RTs) of *P. acnes* strains, RT1-
20 RT10 (SEQ ID NOs 1-10), wherein said typing occurs by determining whether said individual possesses one or more of RT1-RT10 and wherein said individual is diagnosed as having acne if said individual possesses RT4, RT5, RT7, RT8, RT9, or RT10. For example, said individual may be diagnosed as having acne if said individual possesses RT4 (SEQ ID NO:4), RT5 (SEQ ID NO:5), or RT8 (SEQ ID
25 NO:8).

In another embodiment, the invention provides a method for diagnosing different types of acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; amplifying 16S ribosomal DNA in said sample; sequencing said amplified DNA products; and typing the subject's DNA based on
5 one or more of the five major microbiome types of *P. acnes* strains, wherein said subject is diagnosed as having acne if said subject is typed to microbiome IV or V.

In yet another embodiment, the invention provides a method for rapidly diagnosing acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; using one or more primer sets to amplify said DNA;
10 and analyzing said amplified DNA for the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 29-32 and 82-434, wherein said subject is diagnosed as having acne if the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 29-32 and 82-434 exists. For example, said amplified DNA may be analyzed for the presence of a sequence
15 having at least 99% homology with at least one of SEQ ID NOs 29-32 and 82-434 and wherein said subject is diagnosed as having acne if the presence of a sequence having at least 99% homology with at least one of SEQ ID NOs 29-32 and 82-434 exists. As another example, said amplified DNA may be analyzed for the presence of at least one of SEQ ID NOs 29-32 and 82-434 and wherein said subject is
20 diagnosed as having acne if the presence of at least one of SEQ ID NOs 29-32 and 82-434 exists.

In another embodiment, the invention provides a method for rapidly diagnosing acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; using one or more primer sets to amplify said DNA;

using one or more probes to detect said amplified DNA; and analyzing said probe signals for the presence of Locus 1 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 29 and 82-97), Locus 2 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 30 and 98-186), Locus 3 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 31 and 187-423), and/or Locus 4 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 32 and 424-434), wherein said subject is diagnosed as having acne if one or more of Loci 1-4 are present. For example, the signals may be analyzed for the presence of Locus 1, Locus 2, Locus 3, and/or Locus 4 based upon at least 99% homology or 100% homology.

In the foregoing methods, a primer of said primer sets may be selected from the group consisting of SEQ ID NOs 11, 12, 17, and 18 (for Locus 1), SEQ ID NOs 13, 14, 20, and 21 (for Locus 2), SEQ ID NOs 15, 16, 23, and 24 (for Locus 3), and SEQ ID NOs 26 and 27 (for Locus 4). In the foregoing methods, said probes may be SEQ ID NO:19 (for Locus 1), SEQ ID NO:22 (for Locus 2), SEQ ID NO:25 (for Locus 3), and SEQ ID NO:28 (for Locus 4).

In yet another embodiment, the invention provides a vaccine for the prevention and/or treatment of acne caused by *P. acnes* comprising a heat inactivated *P. acnes* strain, an attenuated protein of said strain, or combination thereof, wherein said strain is an RT4 strain, an RT5 strain, an RT7 strain, an RT8 strain, an RT9 strain, or an RT10 strain.

In yet another embodiment, the invention provides a vaccine for the prevention and/or treatment of acne caused by *P. acnes* comprising a heat inactivated *P. acnes* strain, an attenuated protein of said strain, or combination

thereof identified to be specific to a subject based on 16S rDNA sequence analysis of the strains of *P. acnes* affecting said subject.

With regard to the vaccines, said heat inactivated *P. acnes* strain, attenuated protein, or combination thereof may be specific for at least one of unique genomic loci, regions, or sequences identified for the strains of *P. acnes*. Said heat
5 inactivated *P. acnes* strain, attenuated protein, or combination thereof may be specific for at least one of Locus 1 (SEQ ID NOs 29 and 82-97), Locus 2 (SEQ ID NOs 30 and 98-186), Locus 3 (31 and 187-423), and Locus 4 (32 and 424-434).

In yet another embodiment, the invention provides a method for the
10 personalized treatment of acne comprising determining the strains of *P. acnes* affecting a subject and treating said subject with an active ingredient directed to at least one detected strain of *P. acnes*, wherein the active ingredient comprises a drug targeting specific strains of *P. acnes*, wherein the targeting drug comprises small molecules, antisense molecules, siRNA, biologics, antibodies, and combinations
15 thereof targeting genomic elements specific for strains of *P. acnes* associated with acne.

In yet another embodiment, the invention provides a method for treating acne comprising: administering an effective amount of a probiotic that comprises at least one strain of *P. acnes* that is associated with healthy or normal skin based on its 16S
20 rDNA. Said strain may be an RT6 strain. Said strain may have at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54, such as at least 99% homology or 100% homology.

In yet another embodiment, the invention provides a method for treating acne comprising: administering an effective amount of a metabolite produced by a strain
25 of *P. acnes* that is associated with healthy or normal skin, wherein said metabolite is

selected from the group comprising bacterial culture supernatant, cell lysate, proteins, nucleic acids, lipids, and other bacterial molecules. Said strain may be an RT6 strain. Said strain may have at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54, such as at least 99% homology or 100% homology.

5 In yet another embodiment, the invention provides a method for treating acne in a subject comprising: administering an effective amount of a drug specifically targeting RT4, RT5, RT7, RT8, RT9, or RT10, when said subject is determined to possess RT4, RT5, RT7, RT8, RT9, or RT10, respectively. The earlier-described methods may be performed prior to administration of said drug. Said drug may be a
10 small molecule, antisense molecule, siRNA, biologic, antibody, or combination thereof.

 In yet another embodiment, the invention provides a composition comprising at least one strain of *P. acnes* that is associated with healthy or normal skin. Said strain may be an RT6 strain. Said strain may have at least 95% homology to SEQ
15 ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54, such as at least 99% homology or 100% homology.

 In yet another embodiment, the invention provides a method for diagnosing IB-3-based acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; using one or more primer sets to amplify said DNA;
20 and analyzing said amplified DNA for the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 55-81, wherein said subject is diagnosed as having IB-3-based acne if the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 55-81 exists.

 In yet another embodiment, the invention provides a method for the
25 personalized treatment of acne comprising determining the strain(s) of acne affecting

a subject and administering to said subject an effective amount of at least one phage specifically directed to said strain(s). For example, the subject may be treated with phage directed against an RT4 strain, an RT5 strain, an RT7 strain, and RT8 strain, an RT9 strain, and/or an RT10 strain.

5 In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type I comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID
10 NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:40), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

 In yet another embodiment, the invention provides a method for treating an
15 individual suffering from acne of microbiome type I with IB-3 strain comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL082M00 (SEQ ID NO:47) and PHL071N05 (SEQ ID NO:41).

 In yet another embodiment, the invention provides a method for treating an
20 individual suffering from acne of microbiome type II comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL060L00 (SEQ ID NO:34), PHL112N00 (SEQ ID NO:35), and PHL085M01 (SEQ ID NO:44).

 In yet another embodiment, the invention provides a method for treating an
25 individual suffering from acne of microbiome type III or dominant RT8 comprising

administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type IV comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type V comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43),

PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

In yet another embodiment, the invention provides a method for treating a Propionibacterium humerusii-associated malady comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL085N00 (SEQ ID NO:46), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), and PHL010M04 (SEQ ID NO:38).

In yet another embodiment, the invention provides a kit for diagnosing acne in a subject, wherein said kit comprises: at least one primer selected from the group comprising SEQ ID NOs 11-18, 20, 21, 23, 24, 26, and 27; and instructions for use.

In yet another embodiment, the invention provides a kit for diagnosing acne in a subject, wherein said kit comprises: at least one primer selected from the group comprising SEQ ID NOs 11-18, 20, 21, 23, 24, 26, and 27; at least one probe selected from the group comprising SEQ ID NOs 19, 22, 25, and 28; and instructions for use.

BRIEF DESCRIPTION OF THE FIGURES

This application file contains at least one drawing executed in color. Copies of this application with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

Figure 1 shows that *P. acnes* dominates the microbiota of pilosebaceous units, accounting for 87% of the clones. *P. acnes* was dominant in pilosebaceous units in both acne patients and individuals with normal skin. By 16S rDNA

sequencing, *P. acnes* sequences accounted for 87% of all the clones. Species with a relative abundance greater than 0.35% are listed in order of relative abundance. Species distribution from a metagenomic shotgun sequencing of pooled samples from normal individuals confirmed the high abundance of *P. acnes* in pilosebaceous units, as shown on the far right column.

Figure 2 shows that the rank abundance of *P. acnes* ribotypes shows a distribution similar to that seen at the higher taxonomic levels. A few highly-abundant ribotypes and a large number of rare ribotypes were observed in the samples. Some ribotypes were highly enriched in acne patients. Only the top 30 most abundant ribotypes are reflected in Figure 2.

Figure 3 shows that the most abundant *P. acnes* ribotypes in pilosebaceous units were also abundant at other body sites. The major ribotypes found in acne patients and normal individuals were compared to the datasets from the HMP and Grice *et al.* (2009). The top three ribotypes are the most abundant ones in different datasets. The excess RT4 and RT5 seen in the dataset by Grice *et al.* (2009) was due to one subject, HV4, whose *P. acnes* strain population was dominated by these two ribotypes at every skin site sampled. After removal of this subject, the ribotype distribution is similar to the HMP samples and the normal skin samples studied. RT6 is also found abundant in the HMP dataset, which were collected from healthy individuals.

Figure 4 shows that *P. acnes* population structures differ in acne and normal skin. *P. acnes* populations from samples were clustered using principal coordinates analysis of the weighted UniFrac distance matrix for the top ten most abundant ribotypes. The principal coordinate 1 (P1) explains 43.64% of the variation and P2

explains 20.07% of the variation. The analysis was performed using QIIME (Caporaso *et al.* 2010).

Figure 5 shows the distribution of the top ten most abundant *P. acnes* ribotypes in acne patients and individuals with normal skin. Each column represents the percentage of the top ten ribotypes identified in each subject. The average *P. acnes* clone number per subject was 262 and the average clone number of top ten ribotypes was 100. Five major microbiome types at the *P. acnes* strain level were observed in the data. Types IV and V were mostly found in acne patients. Two samples (one from acne, one from normal skin) with fewer than 50 *P. acnes* 16S rDNA sequences are not displayed.

Figure 6 shows the distribution of the top ten most abundant *P. acnes* ribotypes in all samples without separating the two groups of acne and normal skin. Each column represents the percentage of the top ten ribotypes identified in each sample. When all samples were clustered, the same five major microbiome types at the *P. acnes* strain level were observed, indicating that microbiome classification does not depend on the states of the disease. Only three out of 99 samples were clustered differently compared to the one shown in Figure 5 (marked with asterisks). Two samples, one from acne and one from normal skin, with fewer than 50 *P. acnes* 16S rDNA sequences are not shown.

Figure 7 shows that the same five major microbiome types were observed in multiple datasets. Samples from the study, HMP, and Grice *et al.* (2009) were clustered together based on the top ten most abundant *P. acnes* ribotypes. In total, 284 samples were included. Each column represents the percentage of the top ten ribotypes identified in each sample. Both HMP samples and samples from Grice *et al.* (2009) were collected from healthy individuals, therefore the percentage of

microbiome types IV and V are under-represented in the analysis. Samples with fewer than ten sequences of the top ten ribotypes were not included.

Figure 8 indicates that the genome comparison of 71 *P. acnes* strains showed that the genomes of RT4 and RT5 are distinct from others. Two chromosomal regions, loci 1 and 2, are unique to clade IA-2 and one other genome HL086PA1. Clade IA-2 consists of mainly RT4 and RT5 that were highly enriched in acne. The presence of a plasmid (locus 3) is also characteristic of RT4 and RT5. Each row represents a *P. acnes* genome colored according to the ribotypes. Rows are ordered by the phylogeny calculated based on the SNPs in the *P. acnes* core genome. Only the topology is shown. The clades were named based on their *recA* types (IA, IB and II). Columns represent predicted open reading frames (ORFs) in the genomes and are ordered by ORF positions along the finished genome HL096PA1, which encodes a 55 Kb plasmid. Only the first 300 ORFs on the chromosome (on the left) and all the ORFs on the plasmid (on the right) are shown. The colored plasmid regions represent genes on contigs that match exclusively to the HL096PA1 plasmid region. The genes that fall on contigs that clearly extend beyond the plasmid region are likely to be chromosomally located and are colored in grey. Acne index for the ribotypes was calculated based on the percentage of clones of each ribotype found in acne as shown in column 5 in Table 1.

Figure 9 shows the phylogenetic tree constructed based on the 96,887 SNPs in *P. acnes* core genome, which shows that the 71 genomes cluster into distinct clades, consistent with *recA* types that have been used to classify *P. acnes* strains. The 16S ribotypes of the genomes represent the relationship of the lineages to a large extent. At one end of the tree, clades IA-2 and IB-1 mainly consist of the ribotypes enriched in acne, and at the other end of the tree, RT6 in clade II was

mainly found in healthy subjects. Bootstrap test with 1,000 replicates were performed. The distances between the branches were calculated based on the SNPs in the core genome and do not represent the non-core regions of each genome. The enlarged branches were colored according to the 16S ribotypes as shown in Figure 8.

Figure 10 provides a genome comparison of 71 *P. acnes* strains and shows that the genomes of RT4 and RT5 are distinct from others. All of the predicted open reading frames (ORFs) encoded on the chromosome are shown. Each row represents a *P. acnes* genome colored according to the ribotypes. Rows are ordered by the phylogeny calculated based on the SNPs in *P. acnes* core genome. Only the topology is shown. Columns represent ORFs in the genomes and are ordered by their positions along the finished genome HL096PA1. Loci 1 and 2, which are unique to mainly RT4 and RT5 strains, and locus 4, which is unique to mainly RT8 strains, can be seen in the figure.

Figure 11 provides a sequence coverage comparison between the chromosome and the plasmid region in all genomes harboring a putative plasmid, which shows that the copy number of plasmid ranges from 1 to 3 per genome. The X-axis represents the DNA sequences along the chromosome based on the coordinates of the finished genome HL096PA1, followed by plasmid sequences. The Y-axis represents the sequence coverage. The genomes were in the same order as in Figure 8, except HL056PA1 (as a negative control).

Figure 12 reflects that quantitative PCR (qPCR) confirmed that the copy number of plasmid in each genome is 1-3 as predicted from sequence coverage comparison. Pak and RecA are housekeeping genes located on the chromosome and TadA is a conserved gene in the Tad locus located on the plasmid. The copy

number ratio between TadA and Pak ranges from 1 to 3 in genomes, while the ratio between RecA and Pak is 1 in all the genomes. The TadA gene in HL078PA1 and HL045PA1 had amplification in late cycles in qPCR. Conventional PCR confirmed the amplification of TadA in these two strains, while other strains without the plasmid showed no amplification (data not shown).

Figure 13 shows a power law regression for new genes (n) discovered with the addition of new genome sequences (N). Circles are the medians of n for 200 simulations. Error bars indicate the standard deviations for the 200 simulations.

Figure 14 shows a power law regression for total genes (n) accumulated with the addition of new genome sequences (N). Circles are the medians of n for 200 simulations. Error bars indicate the standard deviations for the 200 simulations.

Figure 15 shows the proportion of the 123,223 SNPs in the core regions specific to recA types I, II and III.

Figure 16 shows the phylogenetic tree of 82 *P. acnes* strains constructed based on the 123,223 SNPs in the core regions (2.20 Mb). The distances between strains were calculated as nucleotide substitution rates at all SNP sites, colored according to the scale bar. The strains from the same individuals (SSIs) belonging to the same lineages were marked with "+".

Figure 17 shows the pan-genomes of types IA (A), IB (B) and II (C) strains. Circles are the medians of n for 200 simulations. Error bars are standard deviations for the 200 simulations.

Figure 18 shows the SNP distribution in core regions. Figure 18a shows SNP frequencies (percentage of polymorphic sites) of the genes in the core regions. Figure 18b provides K-S statistics for genes that had higher SNP frequencies with more than two standard deviations (SD). Figure 18c reflects non-synonymous

mutation frequencies of the genes in the core regions. Figure 18d provides K-S statistics for genes that had higher non-synonymous mutation frequencies with more than 2 standard deviations.

Figure 19 provides the distances between *P. acnes* strains in the same lineage (Figure 19a) and in different lineages (Figure 19b).

Figure 20 reflects that *P. acnes* strains within each lineage share unique non-core genomic regions. Rows represent 82 *P. acnes* genomes and columns represent 314 non-core regions that are longer than 500 bp. The genomes and the non-core regions were clustered based on similarity, respectively. The width of each block plotted is not proportional to the genomic length of each non-core region. The presence of a non-core region is colored in yellow, and the absence is colored in blue. The color schemes used for RT and clades are the same as in Figure 16.

Figure 21 provides CRISPR spacer sequences in RT2 and RT6 strains. A total of 48 CRISPR spacer sequences were found in 11 *P. acnes* genomes, 29 of which were unique. Some CRISPR spacers were found in multiple strains. For example, spacer 2 (S2) was shared by HL060PA1 and HL082PA2. Spacer 17 (S17) was shared by J139, ATCC11828, HL110PA3, HL110PA4, HL042PA3 and HL202PA1. Spacer 18 (S18) was shared by J139, ATCC11828, HL110PA3, HL110PA4, and HL202PA1. The tree was from Figure 16 constructed based on the 123,223 SNPs in the core regions.

Figure 22 reflects genes with putative lipase activity in the *P. acnes* genomes. Figure 22a gives a summary of 13 genes with putative lipase activity based on the annotations of KPA171202 and SK137 genomes. Figure 22b reflects Insertions/deletions and frameshift observed in ORF HMPREF0675-4856.

Figure 23 reflects fast detection of acne associated *P. acnes* strains using multiplex PCR targeting loci 1, 2, and 3.

Figure 24 shows the relative abundances of Locus 1 and Locus 2 as compared to the housekeeping gene Pak.

5 **Figure 25** reflects qPCR triplex amplification plots for clinical samples #1 (Figure 25A) and #2 (Figure 25B) showing amplification of *P. acnes* Locus 1, Locus 3, and Pak.

Figure 26 shows the evolutionary relationships/phylogenetic tree of 32 phages.

10 **Figure 27** shows a diagram of the methods of the invention for the diagnosis and personalization of therapy for acne.

Figure 28 shows a flow chart of the methods of the invention for the diagnosis and personalization of therapy for acne.

15 **Figure 29** provides *P. acnes* phage genomes and annotations. Genome organizations of all 15 phages are shown. Hatched arrows in previously published genomes represent newly annotated ORFs proposed. Italicized legend entries refer to newly-annotated or revised ORFs.

20 **Figure 30** provides a phylogenetic tree of 29 sequenced phage genomes constructed based upon the 6,148 SNPs in the core regions. Branches with bootstrap values less than 80 (based on 200 resamplings) were collapsed.

25 **Figure 31** provides phylogenetic trees based on the genome sequences. Figure 31a provides a phylogenetic tree constructed based on the entire genome sequences of all 16 phages. With the exception of PHL112N00, the phylogenetic relationships among the phages remain the same as using the core regions only, shown in Figure 30. Figure 31b shows the phylogenetic tree that was constructed

using only the left-arm of the genomes, which are highly conserved among the phages. Figure 31s shows the phylogenetic tree that was constructed using only the right-arm coding regions. Groups I and II from Figure 30 are also indicated in the trees. Branches with bootstrap values less than 80 (based on 5,000 resamplings) were collapsed.

Figure 32 shows the phylogentic trees constructed based upon the nucleotide sequences of amidase (Figure 32a) and head protein (Figure 32b) from all phages, including the sequences from Lood *et al.* The phylogenetic relationships among the phages from the previous study remain the same in these trees. Groups I and II remain the same as in the genome shown in Figure 30.

Figure 33 reflects multiple alignments generated for genomes from Groups I and II of closely-related phages. Sites of nucleotide variations are mapped to a member from each group. The density of variable sites in each 50-nt window of the genome is indicated in red, with 100% density indicating that all 50 sites in the window vary between the group members. Figure 33a provides variations among Group I phages (PHL010M04, PHL066M04, PHL073M02) mapped to the PHL010M04 genome. Figure 33b provides variation among Group II phages (PHL115M02, PHL085M01, PHL085N00, PHL037M02) mapped to the PHL115M02 genome. Gray arrows represent ORFs in each genome.

Figure 34 shows host range and specificity of *P. acnes* phages. The susceptibility/resistance of 66 *P. acnes* strains, three *P. humerusii* strains, and one *P. granulosum* strain against 15 newly sequenced phages is shown. Dendrograms on the top and to the left represent the respective phylogenetic trees of the phages and *P. acnes* strains (only topology is shown). "S" indicates that the tested *Propionibacterium* strain was susceptible to the tested phage. Numbers in red

represent the fold increase in resistance of the *Propionibacteria* strains against phages relative to *P. acnes* strain ATCC6919.

Figure 35 provides a correlation between *P. acnes* resistance to phages and the presence of matched CRISPR spacers. The colored pixels in each cell represent the CRISPR spacers encoded in each *P. acnes* strain (shown in rows). Each red pixel means that this spacer has an exact protospacer match in the corresponding phage (shown in columns). Each orange pixel means that this spacer has a partially matched protospacer (one to two mismatches) in the corresponding phage. Gray pixels mean no matched protospacers. Pink cells indicate the bacterial resistance to the phages.

Figure 36 reflects that each of the 15 sequenced phages was aligned to all 8 CRISPR spacer arrays identified in the *P. acnes* strains to identify protospacer sequences in each phage genome that have an exact match (red) or up to two mismatches (orange). Plus- and minus-strand protospacers are shown above and below the genomes, respectively.

Figure 37 reflects sequence conservation in protospacers and PAMs. The protospacers that match exactly to the CRISPR spacers encoded in strain HL042PA3 and their associated PAM sequences are shown. Sequence conservation among the protospacer motifs from the phages that HL042PA3 is resistant to is shown in (A) and susceptible to is shown in (B).

DETAILED DESCRIPTION

In one embodiment, the invention provides a method for determining whether an individual possesses acne comprising: obtaining a skin sample from an individual; isolating bacterial DNA from said sample; amplifying 16S ribosomal DNA in said sample; sequencing said amplified DNA products; and typing the individual's DNA based on one or more of the ten major ribotypes (RTs) of *P. acnes* strains, RT1-RT10 (SEQ ID NOs 1-10), wherein said typing occurs by determining whether said individual possesses one or more of RT1-RT10 and wherein said individual is diagnosed as having acne if said individual possesses RT4, RT5, RT7, RT8, RT9, or RT10. For example, said individual may be diagnosed as having acne if said individual possesses RT4 (SEQ ID NO:4), RT5 (SEQ ID NO:5), or RT8 (SEQ ID NO:8).

In another embodiment, the invention provides a method for diagnosing different types of acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; amplifying 16S ribosomal DNA in said sample; sequencing said amplified DNA products; and typing the subject's DNA based on one or more of the five major microbiome types of *P. acnes* strains, wherein said subject is diagnosed as having acne if said subject is typed to microbiome IV or V.

In yet another embodiment, the invention provides a method for rapidly diagnosing acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; using one or more primer sets to amplify said DNA; and analyzing said amplified DNA for the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 29-32 and 82-434, wherein said subject is diagnosed as having acne if the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 29-32 and 82-434 exists. For

example, said amplified DNA may be analyzed for the presence of a sequence having at least 99% homology with at least one of SEQ ID NOs 29-32 and 82-434 and wherein said subject is diagnosed as having acne if the presence of a sequence having at least 99% homology with at least one of SEQ ID NOs 29-32 and 82-434 exists. As another example, said amplified DNA may be analyzed for the presence of at least one of SEQ ID NOs 29-32 and 82-434 and wherein said subject is diagnosed as having acne if the presence of at least one of SEQ ID NOs 29-32 and 82-434 exists.

In another embodiment, the invention provides a method for rapidly diagnosing acne comprising: obtaining a skin sample from a subject; isolating bacterial DNA from said sample; using one or more primer sets to amplify said DNA; using one or more probes to detect said amplified DNA; and analyzing said probe signals for the presence of Locus 1 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 29 and 82-97), Locus 2 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 30 and 98-186), Locus 3 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 31 and 187-423), and/or Locus 4 (at least one sequence having at least 95% homology to at least one of SEQ ID NOs 32 and 424-434), wherein said subject is diagnosed as having acne if one or more of Loci 1-4 are present. For example, the signals may be analyzed for the presence of Locus 1, Locus 2, Locus 3, and/or Locus 4 based upon at least 99% homology or 100% homology.

In the foregoing methods, a primer of said primer sets may be selected from the group consisting of SEQ ID NOs 11, 12, 17, and 18 (for Locus 1), SEQ ID NOs 13, 14, 20, and 21 (for Locus 2), SEQ ID NOs 15, 16, 23, and 24 (for Locus 3), and SEQ ID NOs 26 and 27 (for Locus 4). In the foregoing methods, said probes may be

SEQ ID NO:19 (for Locus 1), SEQ ID NO:22 (for Locus 2), SEQ ID NO:25 (for Locus 3), and SEQ ID NO:28 (for Locus 4).

In yet another embodiment, the invention provides a vaccine for the prevention and/or treatment of acne caused by *P. acnes* comprising a heat
5 inactivated *P. acnes* strain, an attenuated protein of said strain, or combination thereof, wherein said strain is an RT4 strain, an RT5 strain, an RT7 strain, an RT8 strain, an RT9 strain, or an RT10 strain.

In yet another embodiment, the invention provides a vaccine for the prevention and/or treatment of acne caused by *P. acnes* comprising a heat
10 inactivated *P. acnes* strain, an attenuated protein of said strain, or combination thereof identified to be specific to a subject based on 16S rDNA sequence analysis of the strains of *P. acnes* affecting said subject.

With regard to the vaccines, said heat inactivated *P. acnes* strain, attenuated protein, or combination thereof may be specific for at least one of unique genomic
15 loci, regions, or sequences identified for the strains of *P. acnes*. Said heat inactivated *P. acnes* strain, attenuated protein, or combination thereof may be specific for at least one of Locus 1 (SEQ ID NOs 29 and 82-97), Locus 2 (SEQ ID NOs 30 and 98-186), Locus 3 (31 and 187-423), and Locus 4 (32 and 424-434).

In yet another embodiment, the invention provides a method for the
20 personalized treatment of acne comprising determining the strains of *P. acnes* affecting a subject and treating said subject with an active ingredient directed to at least one detected strain of *P. acnes*, wherein the active ingredient comprises a drug targeting specific strains of *P. acnes*, wherein the targeting drug comprises small molecules, antisense molecules, siRNA, biologics, antibodies, and combinations

thereof targeting genomic elements specific for strains of *P. acnes* associated with acne.

In yet another embodiment, the invention provides a method for treating acne comprising: administering an effective amount of a probiotic that comprises at least one strain of *P. acnes* that is associated with healthy or normal skin based on its 16S rDNA. Said strain may be an RT6 strain. Said strain may have at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54, such as at least 99% homology or 100% homology.

In yet another embodiment, the invention provides a method for treating acne comprising: administering an effective amount of a metabolite produced by a strain of *P. acnes* that is associated with healthy or normal skin, wherein said metabolite is selected from the group comprising bacterial culture supernatant, cell lysate, proteins, nucleic acids, lipids, and other bacterial molecules. Said strain may be an RT6 strain. Said strain may have at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54, such as at least 99% homology or 100% homology.

In yet another embodiment, the invention provides a method for treating acne in a subject comprising: administering an effective amount of a drug specifically targeting RT4, RT5, RT7, RT8, RT9, or RT10, when said subject is determined to possess RT4, RT5, RT7, RT8, RT9, or RT10, respectively. The earlier-described methods may be performed prior to administration of said drug. Said drug may be a small molecule, antisense molecule, siRNA, biologic, antibody, or combination thereof.

In yet another embodiment, the invention provides a composition comprising at least one strain of *P. acnes* that is associated with healthy or normal skin. Said strain may be an RT6 strain. Said strain may have at least 95% homology to SEQ

ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54, such as at least 99% homology or 100% homology.

In yet another embodiment, the invention provides a method for diagnosing IB-3-based acne comprising: obtaining a skin sample from a subject; isolating
5 bacterial DNA from said sample; using one or more primer sets to amplify said DNA; and analyzing said amplified DNA for the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 55-81, wherein said subject is diagnosed as having IB-3-based acne if the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 55-81 exists.

10 In yet another embodiment, the invention provides a method for the personalized treatment of acne comprising determining the strain(s) of acne affecting a subject and administering to said subject an effective amount of at least one phage specifically directed to said strain(s). For example, the subject may be treated with phage directed against an RT4 strain, an RT5 strain, an RT7 strain, and RT8 strain,
15 an RT9 strain, and/or an RT10 strain.

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type I comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33),
20 PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:40), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type I with IB-3 strain comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL082M00 (SEQ ID NO:47) and

5 PHL071N05 (SEQ ID NO:41).

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type II comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL060L00 (SEQ ID NO:34),

10 PHL112N00 (SEQ ID NO:35), and PHL085M01 (SEQ ID NO:44).

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type III or dominant RT8 comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01

15 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34),

PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37),

PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

20

In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type IV comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33),

25 PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID

NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

5 In yet another embodiment, the invention provides a method for treating an individual suffering from acne of microbiome type V comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

15 In yet another embodiment, the invention provides a method for treating a *Propionibacterium humerusii*-associated malady comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL085N00 (SEQ ID NO:46), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), and PHL010M04 (SEQ ID NO:38).

20 In yet another embodiment, the invention provides a kit for diagnosing acne in a subject, wherein said kit comprises: at least one primer selected from the group comprising SEQ ID NOs 11-18, 20, 21, 23, 24, 26, and 27; and instructions for use.

In yet another embodiment, the invention provides a kit for diagnosing acne in a subject, wherein said kit comprises: at least one primer selected from the group

comprising SEQ ID NOs 11-18, 20, 21, 23, 24, 26, and 27; at least one probe selected from the group comprising SEQ ID NOs 19, 22, 25, and 28; and instructions for use.

5 Nucleotide, polynucleotide, or nucleic acid sequence will be understood to mean both a double-stranded or single-stranded DNA in the monomeric and dimeric forms and the transcription products of said DNAs.

Homologous nucleotide sequence means a nucleotide sequence having at least a percentage identity with the bases of a nucleotide sequence according to the invention of at least 80%, preferably 90%, 95%, 96%, 97%, 98%, 99% or 100%.

10 This percentage is statistical and the differences between two nucleotide sequences may be determined at random or over the whole of their length.

The invention comprises the polypeptides encoded by a nucleotide sequence according to the invention, including a polypeptide whose sequence is represented by a fragment. Herein, the terms polypeptide, peptide, and protein are
15 interchangeable.

Polypeptides allow monoclonal or polyclonal antibodies to be prepared which are characterized in that they specifically recognize the polypeptides. The invention relates to mono- or polyclonal antibodies or their fragments, or chimeric antibodies, characterized in that they are capable of specifically recognizing a polypeptide.

20 Polypeptides used in vaccine compositions according to the invention may be selected by techniques known to the person skilled in the art such as, for example, depending on the capacity of said polypeptides to stimulate the T cells, which is translated, for example, by their proliferation or the secretion of interleukins, and which leads to the production of antibodies directed against said polypeptides.

25 Vaccine combinations will preferably be combined with a pharmaceutically

acceptable vehicle and, if need be, with one or more adjuvants of the appropriate immunity. Pharmaceutically acceptable vehicle means a compound or a combination of compounds that does not provoke secondary reactions and which allows, for example, the facilitation of the administration of the active compound, an increase in its duration of life and/or its efficacy in the body, an increase in its solubility in solution, or an improvement in its conservation.

Applicants identified ten major lineages of *Propionibacterium acnes* and five major microbiome types in the human pilosebaceous unit ("pore"), where acne arises. Some of the *P. acnes* lineages and microbiome types are highly enriched in acne patients and some are associated with healthy skin. The unique genomic components of each major lineage, including a linear plasmid that is unique to acne-associated lineages, have been identified. This information is used to, for example: (1) for a method/kit to isolate bacterial DNA/RNA from pilosebaceous units for downstream analysis; (2) rapidly and accurately detect/diagnose/identify the microbiome type of the affected subject and the major strains of *P. acnes* present in the pores of the affected subject; (3) develop vaccines against acne-associated *P. acnes* strains; (4) develop probiotics using the strains associated with healthy skin in topical creams, solutions, and the like; (5) develop drugs, including small molecules, biologics, and antibodies targeting the genetic elements and biological pathways unique to the *P. acnes* strains associated with acne, and (6) to develop bacteriophage-based strain specific therapy to treat acne.

Once the microbiome type of a subject affected with acne is diagnosed, several approaches described below may be used formulate an effective treatment plan. For example, if the subjects have microbiome types IV or V, or are dominated by *P. acnes* RT10 strains, it is less likely that antibiotic treatment will succeed

because these strains are antibiotic resistant. However, other method treatments remain available, such as retinoids.

According to one embodiment of the invention, in a case where the subject has the virulent ribotypes, including RT4, RT5, and RT8, target specific drugs including small molecules, biologics, and antibodies may be more effective treatments. In a preferred embodiment of the invention, such a patient may be treated with antibodies targeting the genetic elements and biological pathways that are unique to *P. acnes* strains associated with acne.

According to another embodiment of the invention, in a case where the dominant *P. acnes* strains affecting the subject do not harbor a set of CRISPR/Cas, the additional treatment of phage therapy may be more effective.

The present invention also pertains to alternative treatment strategies for acne treatment to balance the relative abundance of *P. acnes* strains by promoting the growth of health-associated strains.

The present invention pertains to methods and kits to isolate bacterial DNA/RNA from pores of affected subjects for downstream genetic analysis. More specifically, the present invention pertains to protocols for the extraction of bacterial genomic DNA and RNA from microcomedone samples. In one particular embodiment of the invention, Biore® Deep Cleansing Pore Strips may be used to sample the bacteria from a subject. Genomic DNA may be extracted according to methods known in the art. For example, the QIAamp DNA Micro Kit (Qiagen) is a commercially available kit that may be used to extract genomic DNA from the supernatant obtained by lysing cells/microcomedones using a beadbeater.

The present invention also pertains to fast and accurate methods and kits for the detection and/or diagnosis of microbiome types in affected subjects. The

microbiome typing/microbiome-specific treatment is based on ten major lineages of *P. acnes* strains and five major microbiome types in the human pilosebaceous unit found through a comprehensive metagenomic analysis using full length 16S rDNA sequencing.

5 Indeed, samples were PCR-amplified using 16S rDNA specific primers with the following sequences: 27f-MP 5'AGRGTTCGATCMTGGCTCAG-3' and 1492r-MP 5'-TACGGYTACCTTGTTAYGACTT-3'. Optionally, following gel purification, the 1.4 Kb product is excised and further purified using, for example, a Qiagen QIAquick Gel Extraction Kit. The purified product is cloned into OneShot E
10 coli. cells using, for example, a TOPO TA cloning kit from Invitrogen. Sequencing is done with a universal forward, universal reverse, and for a subset, internal 16S rDNA primer 907R with sequences of TGTAACGACGGCCAGT (forward), CAGGAAACAGCTATGACC (reverse), and CCGTCAATTCCTTTRAGTTT (907R). Sequence reactions were loaded on ABI 3730 machines from ABI on 50 cm arrays
15 with a long read run module.

Each lineage of *P. acnes* has unique genomic loci, regions, and sequences. Accordingly, specific primers may be generated to target the lineage-specific genomic regions to detect the presence or absence of each lineage, as well as the relative amount of each lineage using methods known in the art, such as PCR/qPCR.
20 This occurs within several hours of obtaining the samples. Prior to Applicants' invention, this required much more time – often weeks using culture-based methods. According to one embodiment of the invention, affected subjects are grouped for microbiome specific treatments based on these diagnoses.

According to the methods of the present invention, unique genomic loci 1, 2,
25 and 3 for strains of ribotypes 4 and 5 have been shown to be associated with acne.

Using specific primers targeting for loci 1, 2 and 3, lineages that contain these loci can be distinguished from lineages that lack these loci. In addition, using PCR/qPCR techniques, the relative abundance of each strain may also be detected. Analysis of a mock community has shown that isolates with loci 1, 2 and 3 in an abundance of 7.5% or higher in the microbiome may be detected using these techniques. Given the sensitivity of qPCR, lower abundance levels to a few DNA copies may also be detectable.

It has previously been reported that heat inactivation of *P. acnes* may be an effective means of developing *P. acnes*-based vaccines. See T. Nakatsuji *et al.*, 128(10) J. Invest. Dermatol. 2451-2457 (Oct. 2008). In one aspect of the present invention, vaccines are developed against acne-associated *P. acnes* strains. In another aspect of the present invention, personalized vaccines are developed against acne-associated *P. acnes* strains. In yet another aspect of the present invention, vaccines are developed against acne-associated *P. acnes* strains using inactive *P. acnes* strains or heat attenuated proteins. Strains suitable for use as vaccines may be identified based on 16S rDNA sequencing, indentifying lineages of *P. acnes* strains associated with acne, and the unique genomic loci, regions, and sequences for each lineage to specifically target strains of *P. acnes* associated with acne and not those strains associated with healthy skin.

According to methods described above, it has been discovered that *P. acnes* strains with ribotypes 4, 5, 7, 8, 9, and 10 are highly associated with acne. In one embodiment of the present invention, a vaccine is raised against these individual strains separately or in combination. Similarly, the genes in loci 1, 2, and 3 may be targets for vaccination because these loci are unique to ribotypes 4 and 5, and are not found in commensal strains. Locus 4, which is unique to ribotype 8 may also

serve as a potential target for vaccine therapy. The list of genes encoded in loci 1, 2, 3, and 4 are shown in Table 2.

The present invention also pertains to probiotics developed using *P. acnes* strains associated with healthy skin in medicines, compositions, topical creams, solutions, or other cosmetic products. Probiotics have, in the past, been used in topical creams. PROBIOTIC LAB™ announced that mixture of 14 specific strains of bacteria was used for treatment of cystic acne (<http://www.probiotic-lab.com/aboutusprobioticlab.html>). Probiotic skin care/DERMBIOTIX has a product line - Probiotic Collagen Complex (PC3), which is claimed to have targeted anti-aging benefits to the skin. However, this is not targeted to acne treatment. Probiotic Collagen Complex (PC3) infuses the skin with the positive bacteria required to effectively combat and eradicate excess negative bacteria caused by external factors (<http://www.dermbiotix.com>). However, prior to the present invention there existed no skin probiotic product reported for acne treatment using *P. acnes* strains associated with healthy/normal skin. In one aspect of the present invention, skin probiotics are developed for acne treatment using *P. acnes* strains associated with healthy/normal skin. In another aspect of the present invention, skin probiotics are developed for acne treatment using *P. acnes* strains associated with healthy/normal skin based on the 16S rDNA sequencing.

In one particular embodiment of the present invention the RT6 lineage of *P. acnes* and associated with healthy skin is used as a topical product. In yet another embodiment of the present invention the RT6 lineage of *P. acnes* is used by inoculating this isolate on the human skin in order to compete off the acne associated strains. In another embodiment, molecules, including proteins, nucleic

acids, lipids, and other metabolites, supernatant of cultures, and/or cell lysate of these strains may be used at probiotics.

The present invention also pertains to drugs targeting acne associated *P.*

acnes strains. This is based upon multiple genome comparison of *P. acnes* in

5 combination with 16S rDNA metagenomic analysis, thereby identifying certain strains

and genomic variations associated with acne. Drugs intended to target acne

associated *P. acnes* include custom designed small molecules, antisense molecules,

siRNA molecules, biologics, and antibodies targeting genomic elements specific for

strains which are associated with acne. Antisense RNA, antibodies, or small

10 molecules can be designed targeting loci 1, 2, 3, and 4. Strains with ribotypes 4, 5,

and 10 are antibiotic resistant. Thus, there is a need in the art for new antibiotics

targeting ribotypes 4, 5, and 10.

The present invention also pertains to personalized phage therapy for subjects affected with acne comprising phages specific to certain strains of *P. acnes*.

15 Certain companies provide phage therapy for acne patients, such as the Phage

Therapy Center™, [http://www.phagetherapycenter.com/pii/PatientServlet?](http://www.phagetherapycenter.com/pii/PatientServlet?command=static_home)

[command=static_home](http://www.phagetherapycenter.com/pii/PatientServlet?command=static_home)). However, such companies provide no information on the

bacterial host specificity of the phages used for the therapy. *P. acnes* is commensal

and some strains play a protective role for hosts. In one embodiment of the

20 invention, personalized phage therapies include a selections of phages targeting *P.*

acnes strains that have been shown to lack a protective role for subjects affected by

acne. In yet another embodiment of the invention, personalized phage therapy may

be developed according to their bacterial host specificity of the phages to target

specific strains of *P. acnes*, leaving health associated strains intact. In addition, it is

25 possible to identify the structure of *P. acnes* lineages of the affected subjects and

use that structure to predict resistance to phage infection or plasmid conjugation to better target specific phage therapies. For example, *P. acnes* lineages RT2 and RT6 have a CRISPR/Cas structure, indicating they have resistance against certain phage infection and plasmid conjugation. Table 5 shows the sensitivity and resistance of specific *P. acnes* strains to specific *P. acnes* phages.

The invention is described in more detail in the following illustrative examples. Although the examples may represent only selected embodiments of the invention, the following examples are illustrative only and in no way limiting.

EXAMPLES

EXAMPLE 1 - ANALYSIS OF PROPIONIBACTERIUM ACNES STRAIN POPULATIONS IN THE HUMAN SKIN MICROBIOME ASSOCIATED WITH ACNE

The human skin microbiome plays important roles in skin health and disease. However, prior to Applicants' invention the bacterial population structure and diversity at the strain level was poorly understood. The inventors compared the skin microbiome at the strain level and genome level of *Propionibacterium acnes*, a dominant skin commensal, between 49 acne patients and 52 healthy individuals by sampling the pilosebaceous units on their noses. Metagenomic analysis demonstrated that while the relative abundances of *P. acnes* were similar, the strain population structures were significantly different in the two cohorts. Certain strains were highly associated with acne and other strains were enriched in healthy skin. By sequencing 66 novel *P. acnes* strains and comparing 71 *P. acnes* genomes, the inventors identified potential genetic determinants of various *P. acnes* strains in association with acne or health. The analysis indicates that acquired DNA sequences and bacterial immune elements may play roles in determining virulence

properties of *P. acnes* strains and some may be targets for therapeutic interventions. This study demonstrates a previously-unreported paradigm of commensal strain populations that explains the pathogenesis of human diseases. It underscores the importance of strain level analysis of the human microbiome to define the role of commensals in health and disease.

Background

The diversity of the human microbiota at the strain level and its association with human health and disease are largely unknown. However, many studies had shown that microbe-related human diseases are often caused by certain strains of a species, rather than the entire species being pathogenic. Examples include methicillin-resistant *Staphylococcus aureus* (MRSA) (Chambers and Deleo, 2009; Chen *et al.*, 2010; Hansra and Shinkai) and *Escherichia coli* O157 (Chase-Topping *et al.*, 2008; Tarr *et al.*, 2005). *Acne vulgaris* (commonly called acne) is one of the most common skin diseases with a prevalence of up to 85% of teenagers and 11% of adults (White, 1998). Although the etiology and pathogenesis of acne are still unclear, microbial involvement is considered one of the main mechanisms contributing to the development of acne (Bojar and Holland, 2004; Cunliffe, 2002). In particular, *Propionibacterium acnes* has been hypothesized to be an important pathogenic factor (Webster, 1995). Antibiotic therapy targeting *P. acnes* has been a mainstay treatment for more than 30 years (Leyden, 2001). However, despite decades of study, it remained unclear as to how *P. acnes* contributes to acne pathogenesis while being a major commensal of the normal skin flora (Bek-Thomsen *et al.*, 2008; Cogen *et al.*, 2008; Costello *et al.*, 2009; Dominguez-Bello *et al.*, 2010; Fierer *et al.*, 2008; Gao *et al.*, 2007; Grice *et al.*, 2009). Whether *P. acnes* protects

the human skin as a commensal bacterium or functions as a pathogenic factor in acne, or both, remained to be elucidated.

Thus, Applicants compared the skin microbiome at the strain level and genome level in 49 acne patients and 52 normal individuals using a combination of metagenomics and genome sequencing. First, for each sample, 16S ribosomal DNA (rDNA) was amplified, approximately 400 clones were sequenced, and an average of 311 nearly full length 16S rDNA sequences were analyzed. The population structure of *P. acnes* strains was determined in each sample. Second, each *P. acnes* strain was assigned an "acne index" by calculating its prevalence in acne patients based on the 16S rDNA metagenomic data. The *P. acnes* strains associated with the acne patient group were identified, as well as the strains enriched in the individuals with normal skin. This metagenomic approach is fundamentally different than prior approaches in determining disease associations; it is more powerful and less biased than traditional methods by bypassing the biases and selection in strain isolation and culturing. Lastly, 66 novel *P. acnes* strains were sequenced and 71 *P. acnes* genomes compared covering the major lineages of *P. acnes* found in the skin microbiota. By combining a metagenomic study of the skin microbiome and genome sequencing of this major skin commensal, Applicants' study provided insight into bacterial genetic determinants in acne pathogenesis and emphasizes the importance of strain level analysis of the human microbiome to understand the role of commensals in health and disease.

Results

***P. acnes* Dominates The Pilosebaceous Unit**

Applicants characterized the microbiome in pilosebaceous units ("pores") on the nose collected from 49 acne patients and 52 individuals with normal skin. Nearly

full length 16S rDNA sequences were obtained using Sanger method, which permitted analyzing the *P. acnes* at the strain level. After quality filtering, the final dataset contained 31,461 16S rDNA sequences ranging from position 29 to position 1483. 27,358 of the sequences matched to *P. acnes* with greater than 99% identity.

- 5 The data demonstrated that *P. acnes* dominates the microbiota of pilosebaceous units, accounting for 87% of the clones (Figure 1). Other commonly found species in pilosebaceous units included *Staphylococcus epidermidis*, *Propionibacterium humerusii*, and *Propionibacterium granulosum*, each representing 1%-2.3% of the total clones. A total of 536 species level operational taxonomic units (SLOTUs)
- 10 belonging to 42 genera and six phyla were identified in the samples (Table S1).

Table S1. Six phyla and 42 genera found in pilosebaceous units.

Phylum	Genus	Phylum	Genus
Actinobacteria	Actinobaculum	Bacteroidetes	Chryseobacterium
	Corynebacterium		Niastella
	Gordonia		Parabacteroides
	Kocuria		Prevotella
	Microbacterium		Caulobacteraceae
	Propionibacterium	Proteobacteria	Citrobacter
Firmicutes	Anaerococcus		Cupriavidus
	Anoxybacillus		Delftia
	Bacillus		Diaphorobacter
	Enterococcus		Haemophilus
	Erysipelothrix		Klebsiella
	Finegoldia		Massilia
	Gemella		Neisseriaceae
	Lactobacillus		Novosphingobium
	Paenibacillus		Pelomonas
	Peptoniphilus		Phyllobacterium
	Peptostreptococcaceae		Ralstonia
	Ruminococcaceae		Shigella
	Staphylococcus		Sphingomonas
	Streptococcus		Stenotrophomonas
Fusobacteria	Fusobacterium	Cyanobacteria	Streptophyta

To bypass the potential biases due to PCR amplification and due to uneven numbers of 16S rDNA gene copies among different species, a metagenomic shotgun

sequencing of the total DNA pooled from the pilosebaceous unit samples of 22 additional normal individuals was performed. Microbial species were identified by mapping metagenomic sequences to reference genomes. The results confirmed that *P. acnes* was the most abundant species (89%) (Figure 1). This is consistent with the results obtained from 16S rDNA sequencing (87%).

For the 16S rRNA sequence, positions 27 to 1492 were PCR amplified. Yet, when analyzing the sequence only positions 29-1483 are studied. The numbering of positions is based on the *E. coli* system of nomenclature. Thus, the sequences between 29-1483 are important for determining the ribotype (there are many ribotypes, not just 10). As for the top 10 ribotypes, sequences between positions 529-1336 of the 16A rRNA are sufficient.

Different *P. acnes* Strain Populations in Acne

There was no statistically significant difference in the relative abundance of *P. acnes* when comparing acne patients and normal individuals. It was then examined whether there were differences at the strain level of *P. acnes* by extensively analyzing the *P. acnes* 16S rDNA sequences. Herein, each unique 16S rDNA sequence as a 16S rDNA allele type is called a ribotype (RT). The most abundant *P. acnes* sequence was defined as ribotype 1 (RT1) (SEQ ID NO:1). All other defined ribotypes have 99% or greater sequence identity to RT1. Similar to the distributions seen at higher taxonomical levels (Bik *et al.*), at the strain level a few ribotypes were highly abundant in the samples with a significant number of rare ribotypes (Figure 2). After careful examination of the sequence chromatograms and manual correction of the sequences, a total of 11,009 ribotypes were assigned to the *P. acnes* 16S rDNA sequences. Most of the minor ribotypes were singletons. On average, each individual harbored 3 ± 2 *P. acnes* ribotypes with three or more clones. Based on the

genome sequences described below, all the sequenced *P. acnes* strains have three identical copies of 16S rDNA genes (see note below). This allowed the *P. acnes* strain populations in individuals based on the 16S rDNA sequences to be compared. The top ten major ribotypes with more than 60 clones and found in multiple subjects are shown in Table 1:

Table 1 – Top ten most abundant ribotypes found in pilosebaceous units

Ribotype	Nucleotide changes from RT1	Number of subjects	Number of clones	Percentage of clones from acne patients ^a	Percentage of clones from normal individuals ^b	<i>p</i> -value ^c
RT1	-	90	5536	48%	52%	0.84
RT2	T854C	48	1213	51%	49%	0.36
RT3	T1007C	60	2104	40%	60%	0.092
RT4	G1058C, A1201C	23	275	84%	16%	0.049
RT5	G1058C	15	205	99%	1%	0.00050
RT6	T854C, C1336T	11	262	1%	99%	0.025
RT7	G529A	10	188	99%	1%	0.12
RT8	G1004A, T1007C	5	239	100%	0%	0.024
RT9	G1268A	4	68	99%	1%	0.29
RT10	T554C, G1058C	5	61	100%	0%	0.024

^aThe percentage was calculated after the number of clones of each ribotype was normalized by the total number of clones in acne patients (acne index).

^bThe percentage was calculated after the number of clones of each ribotype was normalized by the total number of clones in normal individuals.

^cMann-Whitney-Wilcoxon rank sum test.

Analysis of the top ten ribotypes showed both disease-specific and health-specific associations. The three most abundant ribotypes (RT1, RT2 and RT3) were fairly evenly distributed among acne and normal individuals. However, the next seven major ribotypes were significantly skewed in their distributions (Table 1). Ribotypes 4, 5, 7, 8, 9, and 10 were found predominantly in acne patients, with four of these six statistically significantly enriched in acne ($p < 0.05$, Wilcoxon test). Ribotypes 4, 5, and 10 contain a nucleotide substitution G1058C in the 16S rDNA sequences, which has previously been shown to confer increased resistance to

tetracycline (Ross *et al.*, 1998; Ross *et al.*, 2001). However, only a small percentage of the subjects in our study harboring these ribotypes had been treated with antibiotics (Figure 3), therefore enrichment of these three ribotypes in the acne group was not correlated with antibiotic treatment. This is consistent with previous studies, which showed that previous use of antibiotics was not always associated with the presence of antibiotic resistant strains and that some patients who were not previously treated with antibiotics harbored strains already resistant to antibiotics (Coates *et al.*, 2002; Dreno *et al.*, 2001). One ribotype, RT6, although detected in only 11 subjects, was strongly associated with normal skin ($p=0.025$, Wilcoxon test) (Table 1). Its relative abundance in the normal group was similar to that found in the healthy cohort data from the Human Microbiome Project (HMP) (see Figure 3). The percentage of positive subjects (11/52) was similar as well. Three of the 14 HMP subjects had RT6 found in the anterior nares, and one additional subject had RT6 in the left retroauricular crease.

Based on the distributions of the top ten ribotypes, statistical analysis using several different tests showed significant differences in *P. acnes* population structure between acne and normal skin (Figure 4). This is consistent with a principal coordinate analysis, where acne samples and normal skin samples were separated by mostly principal coordinates 1 and 2 (Figure 4), explaining 44% and 20% of the variation, respectively.

To examine whether different individuals share similar *P. acnes* population structures, the samples were clustered based on the relative abundances of the top ten ribotypes. Five main microbiome types were observed at the *P. acnes* strain level (microbiome types I to V). Types IV and V, which are dominated by *P. acnes* RT4 and RT5, respectively, were mainly found in acne patients (Figures 5 and 6).

The same five main microbiome types were observed in the HMP data and the data from Grice *et al.* (Grice *et al.*, 2009) (see Figure 7).

Genome Sequence Analysis of 71 *P. acnes* Strains

All of the top ten most abundant ribotypes differ from RT1 by only one or two nucleotide changes in the 16S rDNA sequence (Table 1). To determine whether such small changes in the 16S rDNA sequence reflect the lineages and evolutionary history at the genome level, 66 *P. acnes* isolates representing major ribotypes 1, 2, 3, 4, 5, 6, and 8 as well as two minor ribotypes, 16 and 532, were selected for genome sequencing. The genomes of these 66 isolates were fully sequenced and assembled to high quality drafts or complete genomes with 50X coverage or more. Five other *P. acnes* genomes, KPA171202 (Bruggemann *et al.*, 2004), J165, J139, SK137, and SK187, were publicly-available and were included in the analysis. A phylogenetic tree based on 96,887 unique single nucleotide polymorphism (SNP) positions in the core genome obtained from these 71 *P. acnes* genomes was constructed. Most of the genomes with the same ribotypes clustered together. The tree indicates that the 16S rDNA ribotypes do represent the relationship of the lineages to a large extent and that 16S rDNA sequence is a useful molecular marker to distinguish major *P. acnes* lineages (Figures 8 and 9).

Genetic Elements Detected in *P. acnes*

A comparative genome analysis among all 71 genomes grouped by ribotypes was performed. The analysis revealed genetic elements by which acne-associated strains could contribute to acne pathogenesis and the elements by which health-associated strains could contribute to maintaining skin health. Specifically, now known are the unique genome regions of RT4 and RT5, which had a strong association with acne, and RT6, which was found enriched in normal skin. Three

distinct regions, loci 1, 2, and 3, were found almost exclusively in strains that belong to clade IA-2 in the phylogenetic tree. Clade IA-2 consists of mainly RT4 and RT5 (Figures 8 and 10). Loci 1 and 2 are located on the chromosome. Locus 1 contains prophage-related genes and appears to be a genomic island. Locus 2 has plasmid integration sites and may be derived from a plasmid sequence. Locus 3 appears to be on a large mobile genetic element, likely a plasmid. The plasmid is approximately 55 Kb long and has inverted terminal repeats according to the finished genome HL096PA1. The sequence data suggest that the plasmid is linear and possibly originated from a phage (Hinnebusch and Tilly (1993)). All but one of the fifteen genomes of RT4 and RT5 have at least 60% of the genes of the plasmid represented, and all of them have regions homologous to the inverted terminal repeats in the plasmid, suggesting that they harbor the same or a similar linear plasmid (Figure 8). The copy number of the plasmid in the genomes ranges from 1 to 3 based on genome sequencing coverage, which was confirmed by quantitative PCR (Figures 11 and 12).

The fact that acne-enriched RT4 and RT5 strains carry a linear plasmid and two unique loci of genomic islands indicates that these plasmid and chromosomal regions play a role in acne pathogenesis. In fact, the linear plasmid encodes a tight adhesion (Tad) locus, which has been suggested to play a role in virulence in other organisms (Kachlany *et al.*, 2000; Schreiner *et al.*, 2003). The complete Tad locus is found in all but one of the fifteen genomes of RT4 and RT5, and is only occasionally found in other ribotypes. Additionally, in locus 2, a Sag gene cluster is encoded, which has been shown to contribute to hemolytic activity in pathogens (Fuller *et al.*, 2002; Humar *et al.*, 2002; Nizet *et al.*, 2000). Figure 6 summarizes the genes that are mostly unique to RT4 and RT5, several of which play essential roles in virulence

in other organisms. Some of these genes encoded in RT4 and RT5 increase virulence, promote stronger adherence to the human host, or induce a pathogenic host immune response.

In the genome comparison analysis, it was found that all the genomes of RT2
5 and RT6 encode Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR). Among the sequenced genomes, RT2 and RT6 are the only ribotypes encoding CRISPR. CRISPR have been shown to confer protective "immunity" against viruses, phages, and plasmids (Horvath and Barrangou, 2010; Makarova *et al.*, 2011). The CRISPR locus encoded in *P. acnes* consists of a series of cas genes
10 - cas3, cse1, cse2, cse4, cas5e, cse3, cas1, and cas2, which are homologous to the CRISPR locus reported in *E. coli* (Figure S10) and the CRISPR4 locus in *Streptococcus thermophilus* (Horvath and Barrangou, 2010).

CRISPR arrays are composed of a cluster of identical repetitive sequences separated by spacer sequences of similar length but with different nucleotide
15 sequences. It has been found that spacer sequences are identical or with one or two mismatches to phage or plasmid DNA sequences. A total of 39 spacer sequences were found in eight *P. acnes* strains, 25 of which were unique as shown in Table 2.

Table 2 – CRISPR spacer sequences found in the genomes of RT2 and RT6

Ribotype	Strain	Spacer number	Spacer sequence	BLAST result	Match found
RT2	HL001PA1	1	CATGGCCTGCACACCAGGCGCTTTAGCACCT	No hits	
		2	CATGGCCTGCACACCAGGCGCTTTAGCACCT	No hits	
		3	CATGGCCTGCACACCAGGCGCTTTAGCACCT	No hits	
		4	GGCGTATGACGAGTTGTGGTCGGCGTTTCCTC	<i>P. acnes</i> phage PA6 gp15 (minor tail protein)	
		5	CGGTGTTAACGGCTTGCTGGCTTGGATGGAG	No hits	
RT2	HL060PA1	1	CGCCTACCGTCAGCTGACTCACGCCTCCGCGTT	No hits	
		2	TCACACCAGTCATCAGCGTCATAGCTCTCTCGG	No hits	
RT2	HL082PA2	1	GGCTCAGCCCTGCCGATGCCTACGCCAAATGG	<i>C. leptum</i> DSM 753 CLOLEP_00129 (cell wall-associated hydrolases (invasion-associated proteins))	Locus 3
		2	TCACACCAGTCATCAGCGTCATAGCTCTCTCGG	No hits	
RT2	HL103PA1	1	CACCGGGCCCATCCGGTCGGCTCCTGAAAGG	<i>C. leptum</i> DSM 753 CLOLEP_00135	Locus 3
RT2	HL106PA1	1	GATCGAGTTGGCTGAGTCGAAGGTGTTGCGGTT	<i>P. acnes</i> phage PA6 gp16 (conserved protein) <i>P. acnes</i> phage PAD20 gp16	
		2	CTGCTCATCGCTCAGCTCCTGCGCTCATCACA	No hits	
		3	CTGCGCCAACAGCCGCATCTGATCCGAATACGG	<i>P. acnes</i> phage PA6 gp3 (phage portal protein)	
		4	CGCAGCAATCTCAGAAGGCCACAACAAGTTCGT	<i>P. acnes</i> phage PA6 gp7 (conserved protein) <i>P. acnes</i> phage PAD20 gp7 <i>P. acnes</i> phage PAS50 gp7	
		5	CAAATCACCAAGCCCAACACGCCGCCACCACC	No hits	
		6	TGTCACCGATTCAATGTATCTATGAGTGGTGTA	No hits	
		7	TTGGGTGGGTGAGGTGGGTCGTCAGTCATGAG	No hits	
		8	GTCGATGTCGAGATTGGCCTGGGGTCCATGTC	<i>Clostridium leptum</i> DSM 753 CLOLEP_00142	Locus 3
		9	ACGTCGTGAACGTACCCCTTGACGGAGACGGCA	No hits	
RT2	J139	1	CGAGGGCTACCACGTGGTCGATTGGACTGTCG	<i>C. leptum</i> DSM 753 CLOLEP_00167 <i>P. acnes</i> SK137 HMPREF0675_3193 (domain of unknown function)	Locus 2
		2	CAGGCGCTCCACTCCCTCGCCCTGGCCACCAAC	No hits	
RT6	HL110PA3 HL110PA4	1	CTATGTGGACAGTGTTGGTTACTGTGGGGGGAA	<i>P. acnes</i> phage PA6 intergenic region between gp45 and gp46	
		2	GCACTGCACCGATATCGTCGTGGCTGTCACTTG	No hits	
		3	CCCAGACAACCTCGACAACCTGTTTCAGGGGATG	<i>P. acnes</i> phage PAS50 gp25	
		4	CATGGCTAGCCCGGATTTTGGCTGCTGAGCG	<i>P. acnes</i> phage PA6 gp34 (multidrug resistance protein-like transporters) <i>P. acnes</i> phage PAD20 gp34 (DNA helicase)	
		5	CGGCCCTGCGGCAGATTTTGTGCGTTGAATCC	<i>P. acnes</i> phage PA6 gp14 (tape measure protein) <i>P. acnes</i> phage PAD20 gp14 (tape measure protein) <i>P. acnes</i> phage PAS50 gp14 (tape measure protein)	
		6	CGGGCAGAGGATGTGTTGCTCTTCCTGGATGG	<i>P. acnes</i> phage PA6 gp32 (CHC2 zinc finger) <i>P. acnes</i> phage PAD20 gp32 (DNA primase) <i>P. acnes</i> phage PAS50 gp32 (DNA primase)	
		7	GTTACGCTGGAACCCCAATGAACACGCGAGAA	<i>P. acnes</i> phage PAD42 major head protein gene <i>P. acnes</i> phage PAD20 major head protein gene <i>P. acnes</i> phage PAD9 major head protein gene <i>P. acnes</i> phage PAS40 major head protein gene <i>P. acnes</i> phage PAS12 major head protein gene <i>P. acnes</i> phage PAS10 major head protein gene <i>P. acnes</i> phage PAD21 major head protein gene <i>P. acnes</i> phage PAS2 major head protein gene <i>P. acnes</i> phage PA6 gp6 (Phage capsid family) <i>P. acnes</i> phage PAS50 gp6 major head protein gene	
		8	CGAGGGCTACCACGTGGTCGATTGGACTGTCG	<i>C. leptum</i> DSM 753 CLOLEP_00167 <i>P. acnes</i> SK137 HMPREF0675_3193 (Domain of unknown function)	Locus 2
		9	CAGGCGCTCCACTCCCTCGCCCTGGCCACCAAC	No hits	

Abbreviations: BLAST, Basic Local Alignment Search Tool; CRISPR, Clustered Regularly Interspaced Short Palindromic Repeat; *C. leptum*, *Clostridium leptum*; *P. acnes*, *Propionibacterium acnes*; RT, ribotype.

As expected, most of the identifiable spacers target to known *P. acnes* phage sequences. However, among the unique CRISPR spacer sequences, one matched locus 2 on the chromosome and three matched the plasmid region (locus 3) in *P. acnes* genomes of mainly RT4 and RT5. This suggests that these loci may have
5 been acquired by RT4 and RT5 strains, while the genomes of RT2 and RT6 may be capable of protecting against the invasion of the plasmids or other foreign DNA through the CRISPR mechanism.

Discussion

The foregoing study of the human skin microbiome associated with acne
10 provides the first portrait of the microbiota of pilosebaceous units at the bacterial strain level. Since *P. acnes* is the major skin commensal bacterium found in both acne and healthy skin, this strain-level analysis is important to help understand the role of *P. acnes* in acne pathogenesis and in skin health. A strong association between strains of RT4 and RT5 with acne and a strong association between strains
15 of RT6 and healthy skin, each with unique genetic elements, has been shown. Other *P. acnes* strains, including ribotypes 7, 8, 9, and 10, or interactions among different strains, may also contribute to the development of the disease. In addition, host factors, such as hormone level, sebum production, and physical changes in the pilosebaceous unit, may also play a role in acne pathogenesis.

20 The foregoing metagenomic approach in revealing the association of *P. acnes* strains with the disease or health is more powerful than previous studies using traditional methods (Lomholt and Kilian, 2010; McDowell *et al.*, 2011). Because the skin microbiota of each individual and each skin site may harbor “good,” “neutral,” and “bad” strains at the same time, which may have different growth rates under in
25 vitro culturing conditions, culturing a few isolates from a disease lesion or healthy

skin site may not provide an accurate and unbiased measurement of the association of the strains with the disease or health. The sampling technique and disease associations in the foregoing study did not depend on sampling locations, on the presence of lesions in the sampling field, or on inherently biased culture techniques.

5 While sampling lesional skin intentionally may yield interesting results, these results would not be capable of defining the disease associations that unbiased sampling can. The metagenomic approach employed in the foregoing study to identify underlying strain differences in acne may also be applied to the study of other disease/health associations with commensal or pathogenic bacteria.

10 **Materials and Methods**

Subjects

Subjects with acne and subjects with normal skin were recruited from various clinics in Southern California including private practice, managed care, and public hospital settings, as well as outside of dermatology clinics, to best represent the

15 diversity of populations and history of medical care. The subject data are available at dbGaP (http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000263.v1.p1). The diagnosis of acne was made by board-certified dermatologists. The presence of acne was graded on a scale of 0 to

5 relating closely to the Global Acne Severity Scale (Dreno *et al.*, 2011). Grades

20 were recorded for both the face and the nose separately where zero represents normal skin and 5 represents the most severe inflammatory cystic acne. In acne patients, the grades of the face ranged from 1 to 5 with an average of 2.1, and the grades of the nose ranged from 0 to 2 with an average of 0.3. The presence of scarring was also noted. Subjects with normal skin were determined by board-

25 certified dermatologists and were defined as people who had no acneiform lesions

on the face, chest, or back. They were also excluded if they had other skin problems that the investigators felt would affect sampling or the microbial population on the skin. Among the 101 subjects, 59 were female (31 acne patients and 28 normal subjects) and 42 were male (18 acne patients and 24 normal subjects). The average
5 age of the acne cohort was 22.2 and the average age of the normal cohort was 29.6. There was no significant difference in ethnicity between the acne and normal populations. The subjects responded to a written questionnaire, administered by a physician or a well-trained study coordinator who went over each question with the subjects. Most of the subjects had not been treated for acne in the past or were not
10 being treated when samples were collected (Figure 3). Only nine out of 78 subjects, who provided treatment information, were being treated for acne when samples were taken. Among the nine subjects, two were being treated with antibiotics, five were being treated with topical retinoids, one was being treated with both antibiotics and retinoids, and one did not list the treatment. Subjects were asked for acne treatment
15 history in the past (anytime in their life). Eighteen out of 73 subjects, who provided treatment history, had been treated for acne in the past. Among them, seven had been treated with antibiotics, eight had been treated with retinoids, two had been treated with both antibiotics and retinoids, and one did not list the treatment. All subjects provided written informed consent. All protocols and consent forms were
20 approved by both the UCLA and Los Angeles Biomedical Research Institute IRBs. The study was conducted in adherence to the Helsinki Guidelines.

Samples

Skin microcomedone (white head or black head) samples were taken from the nose of the subjects using Bioré Deep Cleansing Pore Strips (Kao Brands Company,
25 Cincinnati, OH) following the instruction of the manufacturer. Clean gloves were

used for each sampling. After being removed from the nose, the strip was immediately placed into a 50 mL sterile tube and kept on ice or at 4 °C. The cells were lysed within four hours in most of the cases.

Metagenomic DNA Extraction, 16S rDNA Amplification, Cloning, and Sequencing

Individual microcomedones were isolated from the adhesive nose strip using sterile forceps. Genomic DNA was extracted using QIAamp DNA Micro Kit (Qiagen, Valencia, CA). 16S rDNA was amplified and cloned according to the protocol by HMP, which is described in detail in Supplementary Information. Nearly full length sequences were obtained by Sanger method.

16S rDNA Sequence Analysis

Base calling and quality was determined with Phred (Ewing and Green, 1998; Ewing *et al.*, 1998). Bidirectional reads were assembled and aligned to a core set of NAST-formatted sequences (rRNA16S.gold) using AmosCmp16Spipeline and NAST-ier (Haas *et al.*, 2011). Suspected chimeras were identified using ChimeraSlayer and WigeonN (Haas *et al.*, 2011). 16S rDNA sequences were extensively manually examined. Chromatograms were visually inspected at all bases with a Phred quality score < 30. Appropriate corrections were applied. QIIME (Caporaso *et al.*, 2010) was used to cluster the sequences into OTUs.

***P. acnes* Isolation and Genotyping**

Colonies with the macroscopic characteristics of *P. acnes* were picked from each sample plate and were passed twice. The ribotype of each isolate was determined by PCR amplification and sequencing of the full length of the 16S rDNA gene by Sanger method.

Whole Genome Shotgun Sequencing, Assembly, and Annotation

Genome HL096PA1 was sequenced using Roche/454 FLX and was assembled using a combination of PHRAP/CONSED (Gordon *et al.*, 1998) and GSMAPPER (Roche, Branford, CT) with extensive manual editing in CONSED. The remaining 65 genomes were sequenced using Illumina/Solexa GAllx (Illumina, San Diego, CA). Sequence datasets were processed by quality trimming and were assembled using Velvet (Zerbino and Birney, 2008). Coding sequences were predicted using GeneMark (Borodovsky and McIninch, 1993) and GLIMMER (Salzberg *et al.*, 1998). The final gene set was processed through a suite of protein categorization tools consisting of Interpro, psort-b and KEGG. A more detailed protocol can be found at http://hmpdacc.org/doc/sops/reference_genomes/annotation/WUGC_SOP_DACC.pdf.

Comparative Genome Analysis

Seventy-one *P. acnes* genome sequences were compared using Nucmer (Kurtz *et al.*, 2004). Phylogenetic analysis was performed using MEGA5 (Tamura *et al.*, 2007). CRISPRFinder (Grissa *et al.*, 2007) was used to identify the CRISPR repeat-spacer sequences.

Supplementary Information

16S rDNA sequence of KPA171202

All sequenced *P. acnes* genomes encode three copies of 16S rRNA genes, which are identical within each isolate, except KPA171202. Based on the KPA171202 genome (Bruggemann *et al.*, 2004), one copy of the 16S rRNA gene has one nucleotide difference from the other two identical copies of RT1. However, this mutation was never observed in the 16S rDNA dataset. Multiple clones of 16S rDNA gene from KPA171202 were amplified, cloned, and sequenced and a

sequence harboring this mutation was not found. Thus, KPA171202 also has three identical copies of 16S rDNA.

Comparison of *P. acnes* Strain Distribution to Other Human Microbiome Datasets

To determine whether the *P. acnes* ribotypes and their relative abundances measured in this study are unique to pilosebaceous units, a similar analysis to the microbiome 16S rDNA data from the Human Microbiome Project (HMP) and the data from Grice *et al.* (2009) were applied. Both datasets were obtained from healthy subjects. The relative abundance of the major ribotypes in healthy subjects from the study was largely similar to that found in these two datasets despite the fact that they were sampled from different anatomical sites (Figure 3). RT6 (6.3%) was found to be more abundant than RT4 and RT5 combined (2.8%) in the HMP data, similar to those found in the normal cohort where RT6 represents 4.8% and RT4 and RT5 combined represent 1.2% of the clones. The same five main microbiome types were observed in the two datasets (Figure 7).

Genome Clustering and Phylogenetic Tree

The *recA* gene has been widely used to classify *P. acnes* strains into four known types: IA, IB, II, and III (McDowell *et al.*, 2008; McDowell *et al.*, 2005). The phylogenetic tree of the 71 genomes based on the SNPs in the core genome matched the *recA* types perfectly except one isolate, HL097PA1. Most of the genomes with ribotypes 1, 4, 5, and 532 were grouped to *recA* Type IA clade, which can be further divided into subclades IA-1 and IA-2. Clade IA-2 is composed of mostly RT4 and RT5. RT4 and most of RT5 genomes seem to belong to the same lineage with very similar genome sequences. All the isolates with ribotypes 3, 8, and 16, who share the mutation of T1007C in the 16S rDNA gene, were grouped to *recA* Type IB clade. Most of the RT3 genomes form a subclade IB-2 and RT8 genomes

form a subclade by themselves, IB-1, which was highly associated with acne.

Notably, RT2 and RT6, who share T854C mutation, have a more distant phylogenetic relationship to other ribotypes, and were grouped to the recA Type II clade. This is consistent with previous studies (Lomholt and Kilian, 2010; McDowell
5 *et al.*, 2005). *P. acnes* isolates with recA type III were not found in the samples.

The associations of *P. acnes* lineages with health and disease states were further analyzed. There was a clear shift of the association strength of the clades with acne along the phylogenetic tree (Figure 9). The three sequenced ribotypes identified as being strongly associated with acne (RT4, RT5, and RT8) were found at
10 one end of the tree in clades IA-2 and IB-1, while the RT6 identified as being associated with normal skin was at the other end of the tree at the tip of clade II (Figure 9).

Antibiotic Resistance

P. acnes ribotypes 4, 5, and 10 have a single nucleotide substitution G1058C
15 in the 16S rDNA sequences, which has previously been shown to confer increased resistance to tetracycline (Ross *et al.*, 1998a; Ross *et al.*, 2001). In addition to the substitution in the 16S rDNA sequences, it was determined that all the strains of RT4 and RT5 that were sequenced have a nucleotide substitution in the 23S rDNA
sequences, which confers increased resistance to a different class of antibiotics,
20 erythromycin and clindamycin (Ross *et al.*, 1997; Ross *et al.*, 1998b). It was experimentally confirmed that these isolates, except two that were unculturable, were resistant to tetracycline, erythromycin, and clindamycin.

It was also examined whether the enrichment of these ribotypes in the acne group could be due to antibiotic treatment. However, in the study only a small

percentage of the subjects harboring ribotypes 4, 5, or 10 were treated with antibiotics (Table S2).

Table S2 – Past and current treatments of the subjects

Group	Acne		Normal	
Number of subjects	49		52	
	with RT4, RT5, or RT10	without RT4, RT5, and RT10	with RT4, RT5, or RT10	without RT4, RT5, and RT10
Number of subjects in each subgroup	20	29	9	43
Subjects reported on current treatment	14	25	8	31
no treatment	10	21	8	30
antibiotics	0	2	0	0
retinoids	3	2	0	0
antibiotics and retinoids	0	0	0	1
unknown	1	0	0	0
Subjects reported on past treatment	12	22	8	31
no treatment	5	16	6	28
antibiotics	2	4	0	1
retinoids	4	1	1	2
antibiotics and retinoids	1	0	1	0
unknown	0	1	0	0

- 5 Eighteen of the 29 subjects who harbored any of these three ribotypes gave reports on both past and current treatments. Among them, 50% (9/18) of the subjects were never treated; 33% (6/18) were treated with retinoids; 11% (2/18) were treated with antibiotics in the past, and 5.6% (1/18) were treated with both antibiotics and retinoids in the past. The theory of selection by antibiotic treatment is not
- 10 favored by this study. Previous surveys of antibiotic resistant strains in acne patients demonstrated that previous use of antibiotics did not always result in the presence of resistant strains and that some patients without previous use of antibiotics harbored

resistant strains (Coates *et al.*, 2002; Dreno *et al.*, 2001). Observations in this study are consistent with previous studies.

CRISPR Spacer Sequences

Although more similar to the GC content of *P. acnes* genomes, four unique
5 spacer sequences found in strains of RT2 and RT6 have the best matches to the
genome of *Clostridium leptum*, a commensal bacterium in the gut microbiota (Table
2). On the 55Kb plasmid harbored in HL096PA1 and other RT4 and RT5 genomes,
there is also a large cluster of 35 genes that are identical to the genes found in *C.*
leptum, including the Tad locus.

10 Materials and Methods

Metagenomic DNA Extraction, PCR Amplification, Cloning and 16S rDNA Sequencing

Metagenomic DNA extraction

Individual microcomedones were isolated from the adhesive nose strip using
sterile forceps and placed in a 2 mL sterile microcentrifuge tube filled with ATL buffer
15 (Qiagen) and 0.1 mm diameter glass beads (BioSpec Products, Inc., Bartlesville,
OK). Cells were lysed using a beadbeater for 3 minutes at 4,800 rpm at room
temperature. After centrifugation at 14,000 rpm for 5 minutes, the supernatant was
retrieved and used for genomic DNA extraction using QIAamp DNA Micro Kit
(Qiagen). The manufacturer protocol for extracting DNA from chewing gum was
20 used. Concentration of the genomic DNA was determined by NanoDrop 1000
Spectrophotometer.

16S rDNA PCR amplification, cloning and sequencing

Most of the metagenomic samples were amplified in triplicate using 16S rDNA
specific primers with the following sequences: 27f-MP 5'-
25 AGRGTTTGATCMTGGCTCAG-3' and 1492r-MP 5'-

TACGGYTACCTTGTTAYGACTT-3'. PCR reactions contained 0.5 U/ μ L Platinum Taq DNA Polymerase High Fidelity (Invitrogen), 1X Pre-mix E PCR buffer from Epicentre Fail-Safe PCR system, 0.12 μ M concentration of each primer 27f-MP and 1492r-MP, and Sigma PCR grade water. One microliter of DNA (ranging from 0.2 - 10 ng total) was added to each reaction. The G-Storm GS4 thermocycler conditions were as following: initial denaturation of 96°C for 5 minutes, and 30 cycles of denaturation at 94°C for 30 seconds, annealing at 57°C for 1 minute, and extension at 72°C for 2 minutes, with a final extension at 72°C for 7 minutes. Following amplification, an A-tailing reaction was performed by the addition of 1 U of GOTaq DNA Polymerase directly to the amplification reaction and incubation in the thermocycler at 72°C for 10 minutes.

The three PCR amplification reactions from each source DNA were pooled and gel purified (1.2% agarose gel stained with SYBR Green fluorescent dye). The 1.4 Kb product was excised and further purified using the Qiagen QIAquick Gel Extraction kit. The purified product was cloned into OneShot E. coli cells using TOPO TA cloning kit from Invitrogen.

White colonies were picked into a 384-well tray containing terrific broth, glycerol, and kanamycin using a Qpix picking robot. Each tray was prepared for sequencing using a magnetic bead prep from Agilent and sequenced with 1/16th Big Dye Terminator from ABI. Sequencing was done with a universal forward, universal reverse, and for a subset, internal 16S rDNA primer 907R with sequences of TGTAACGACGGCCAGT (forward), CAGGAAACAGCTATGACC (reverse), and CCGTCAATTCCTTTRAGTTT (907R). Sequence reactions were loaded on ABI 3730 machines from ABI on 50 cm arrays with a long read run module.

A slightly different PCR and cloning protocol without automation was used for several initial samples as described below. 16S rDNA was amplified using universal primers 8F (5'-AGAGTTTGATYMTGGCTCAG-3') and 1510R (5'-TACGGYTACCTTGTTACGACTT-3') (Gao *et al.*, 2007). Thermocycling conditions were as following: initial denaturation step of 5 minutes at 94°C, 30 cycles of denaturation at 94°C for 45 seconds, annealing at 52°C for 30 seconds and elongation at 72°C for 90 seconds, and a final elongation step at 72°C for 20 minutes.

PCR products were purified using DNA Clean and Concentrator Kit (Zymo Research). Subsequently, the 16S rDNA amplicons were cloned into pCR 2.1-TOPO vector (Invitrogen). One-Shot TOP-10 Chemically Competent *E. coli* cells (Invitrogen) were transformed with the vectors and plated on selective media. Individual positive colonies were picked and inoculated into selective LB liquid medium. After 14 hours of incubation, the plasmids were extracted and purified using PrepEase MiniSpin Plasmid Kit (USB Corporation) or Zyppy Plasmid Miniprep Kit (Zymo Research). The clones were sequenced bidirectionally using Sanger sequencing method with 1/8th chemistry using ABI 3730 sequencer (Applied Biosystems Inc.).

***P. acnes* Isolation and Culturing**

Sample culture plate

Microcomedones on the inner surface of the nose strip were mashed and scraped using a sterile loop (Fisherbrand, Pittsburgh, PA), and plated onto a blood agar plate (Teknova Brucella Agar Plate with Hemin and Vitamin K, Teknova, Hollister, CA). The plates were incubated at 37°C for 5 - 7 days anaerobically using the AnaeroPack System (Mitsubishi Gas Chemical Company, Tokyo, Japan).

Isolation and culturing of individual strains

Colonies with the macroscopic characteristics of *P. acnes* were picked from each sample plate and were streaked onto A-media plates (Pancreatic Digase of Casine, Difco yeast extract, glucose, KH₂PO₄, MgSO₄, Difco Agar, and water).

- 5 These first-pass plates were then incubated anaerobically at 37°C for 5-7 days. As the second pass, single isolated colonies were picked from the first-pass plates and streaked onto new A-Media plates. These plates were then incubated anaerobically at 37°C for 5-7 days. The colonies on these plates were picked for culturing, genotyping, and genome sequencing in the subsequence steps.

10 Genotyping of the *P. acnes* Isolates

Each isolate was analyzed by PCR amplification of the 16S rDNA gene. The ribotypes were determined based on the full length sequences. Isolates with desired ribotypes were selected for future culturing and genome sequencing.

Genomic DNA extraction of *P. acnes* isolates

- 15 Isolates were grown in 5 mL of Clostridial medium under anaerobic conditions at 37°C for 5 - 7 days. Cultures were pelleted by centrifugation and washed with 3 mL phosphate buffer saline (PBS). The same protocol used for the metagenomic DNA extraction was used for extracting the genomic DNA of the isolates.

Metagenomic Shotgun Sequencing and Analysis

- 20 Metagenomic DNA samples from microcomedone samples from 22 individuals with normal skin were pooled and sequenced using Roche/454 FLX. The average read length was 236 bp. The sequencing was limited with 13,291 sequence reads. Sequence reads were aligned against the NCBI's non-redundant database using BLAST. Species assignment was based on 97% identity and 100% of the
25 read length aligned.

Assembly, Alignment and Editing of 16S rDNA Sequences

Assembly and alignment

Base calling and quality were determined with Phred (Ewing and Green, 1998; Ewing *et al.*, 1998) using default parameters. Bidirectional reads were
5 assembled and aligned to a core set of NAST-formatted sequences (rRNA16S.gold) using AmosCmp16Spipeline and NAST-ier, which are from the Microbiome Utilities Portal of the Broad Institute (<http://microbiomeutil.sourceforge.net/>). These tools in turn use Amoscmp (Pop *et al.*, 2004), Mummer (Kurtz *et al.*, 2004), Lucy (Chou and Holmes, 2001), BLAST (Altschul *et al.*, 1990) and CdbTools
10 (<http://compbio.dfci.harvard.edu/tgi/software/>). Suspected chimeras were identified using ChimeraSlayer and WigeeN (Haas *et al.*, 2011). Sequences with at least 90% bootstrap support for a chimeric breakpoint (ChimeraSlayer) or containing a region that varies at more than the 99% quantile of expected variation (WigeeN) were removed from further analysis.

Quality screening

For diversity analysis of the *P. acnes* population, sequences with at least 99% identity over 1,400 nucleotides to *P. acnes* KPA171202 (Bruggemann *et al.*, 2004) 16S rDNA were trimmed to positions 29-1483 (numbering based on the *E. coli* system of nomenclature (Brosius *et al.*, 1978)). Sequences without full coverage
20 over this region were excluded from further strain level analysis. Chimera screening, as described above, resulted in removal of less than 0.35% of the sequences. This may be an under-estimation of the chimeras, since the majority of sequences differ by only 1 or 2 nucleotides. Low quality sequences were excluded, defined as more than 50 nucleotides between positions 79 and 1433 with Phred quality scores of less
25 than 15. To allow detailed strain-level analysis, the data were extensively manually

edited. Chromatograms were visually inspected at all bases with a Phred quality score < 30, and appropriate corrections were applied. For analysis at the species level, the 16S rDNA sequences were not manually edited. Chimera screening of assembled sequences resulted in removal of less than 0.65% of the sequences.

- 5 Aligned sequences were trimmed to *E. coli* equivalent positions 29-1483 (Brosius *et al.*, 1978). Sequences without full coverage over this region were excluded from further analysis.

Sequence Editing

- Nearly 62,000 Sanger sequence reads representing the 26,446 assembled *P. acnes* sequences were mapped to the RT1 sequence in CONSED (Gordon, 2003; Gordon *et al.*, 1998). Comprehensive semi-manual editing of the large number of sequences was made feasible by their very high pairwise similarities: a median of only one nucleotide change from RT1 per sequence (three nucleotide changes prior to editing). Editing was facilitated by the use of scripts and the custom navigation feature of CONSED allowing single click jumps to sites requiring inspection.
- 15

- Chromatograms were inspected for all low quality (Phred < 30) bases that differed from RT1, and corrected as needed, including many commonly occurring sequence errors. In order to minimize the effect of base mis-incorporation and chimera, specific base differences from RT1 occurring in less than 4 sequences (frequency < 0.00015) were considered unreliable and reverted to the corresponding RT1 base.
- 20
- Ribotypes were assigned for the resulting sequences based on 100% identity.

16S rDNA Sequence Analysis

OTUs and taxonomy assignments

- QIIME (Caporaso *et al.*, 2010b) was used to cluster the sequences into OTUs using 99% identity cutoff, furthest neighbor, and UCLUST (Edgar, 2010).
- 25

Representative sequences (most abundant) were selected and aligned using PYNAST (Caporaso *et al.*, 2010a) to the greengenes database. Taxonomy was assigned using RDP method (Cole *et al.*, 2009). The alignment was filtered with the lanemask provided by greengenes, and a phylogenetic tree was built using FastTree (Price *et al.*, 2009).

Wilcoxon Test on the Top Ten Ribotypes

For each sample, the number of clones of each of the top ten ribotypes was normalized by the total number of *P. acnes* clones of the sample. The normalized counts were used to test the significance in enrichment between the acne group and the normal group. The function `wilcox_test` in the R program (<http://www.R-project.org>) was used to calculate the p-values.

Microbiome type assignments

Microbiome types were assigned based on the largest clades seen when samples were clustered using thetacy similarity in MOTHUR (Schloss *et al.*, 2009) (Figures 5 and 6) or hierarchical clustering (Eisen *et al.*, 1998) (Figure 7).

Assigning Ribotypes to Datasets of HMP and Grice et al. 2009

Sequences were assigned to a ribotype if they met the following criteria. First, there was a single best match. Second, it covered the range required to discriminate between the top 45 ribotypes (58-1388). Third, there were no Ns at discriminatory positions. Lastly, there were no more than ten non-discriminatory differences.

The HMP 16S rDNA Sanger sequence dataset was downloaded with permission from the HMP Data Analysis and Coordination Center. It has 8,492 *P. acnes* sequences from 14 subjects and nine body sites (retroauricular crease, anterior nares, hard palate, buccal mucosa, throat, palatine tonsils, antecubital fossa, saliva, and subgingival plaque). More details on the dataset can be found at

http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000228.v2.p1. In this dataset, low quality bases (Phred quality < 20) were converted to Ns, and 26% of the sequences were not assigned due to excessive Ns or Ns at ribotype discriminatory sites. Less than 1% was
5 unresolved due to equal best matches or greater than ten mismatches to RT1.

The dataset from Grice *et al.* (2009) is available at NCBI (GenBank accession numbers GQ000001 to GQ116391). It has 22,378 *P. acnes* sequences from ten subjects and 21 skin sites (buttock, elbow, hypothenar palm, volar forearm, antecubital fossa, axillary vault, gluteal crease, inguinal crease, interdigital web
10 space, nare, plantar heel, popliteal fossa, toe web space, umbilicus, alar crease, back, external auditory canal, glabella, manubrium, occiput, and retroauricular crease). Three percent of the sequences were unassigned due to greater than ten mismatches to RT1, and 1.6% was unassigned due to equal best matches.

For comparison purpose, the unedited 16S rDNA sequences were assigned
15 to ribotypes by the same method described above and the result is shown in Figure 3. Less than 0.6% of the sequences were unassigned due to greater than ten mismatches to RT1, and 1.7 % was unassigned due to equal best matches.

Whole Genome Shotgun Sequencing, Assembly and Annotation of 66 *P. acnes* Isolates

20 *Genome HL096PA1*

The genome was sequenced using Roche/454 FLX at the UCLA Genotyping and Sequencing Core. A total of 590,054 sequence reads were generated with an average read length of 230 bp. Of these, 433,896 were assembled into two contigs, a circular main chromosome of 2,494,190 bp and a linear plasmid of 55,585 bp.
25 Assembly was accomplished by a combination of PHRAP/CONSED (Gordon *et al.*,

1998) and GSMAPPER (Roche) with extensive manual editing in CONSED. GeneMark v2.6r (Borodovsky and McIninch, 1993) and GLIMMER v2.0 (Salzberg *et al.*, 1998) were used to performed ab initio protein coding gene prediction. tRNAScan-SE 1.23 was used for tRNA identification and RNAmmer was used for
5 predicting ribosomal RNA genes (5S, 16S, and 23S). Genome annotation results were based on automated searches in public databases, including Pfam (<http://pfam.jouy.inra.fr/>), KEGG (<http://www.genome.jp/kegg>), and COG (<http://www.ncbi.nlm.nih.gov/COG/>). Manual inspection of the annotation was also performed.

10 **Genomes of the other 65 Isolates**

The genomes were sequenced using Illumina/Solexa Genome Analyzer IIx and annotated by the Genome Center of Washington University at St. Louis.

Assembly

Each genomic DNA sample was randomly sheared and an indexed library
15 was constructed using standard Illumina protocols. Twelve uniquely tagged libraries were pooled and run on one lane of a GAIIx flowcell and paired end sequences were generated. Following deconvolution of the tagged reads into the separate samples, datasets were processed using BWA (Li and Durbin, 2009) quality trimming at a q10 threshold. Reads trimmed to less than 35bp in length were discarded and the
20 remaining reads were assembled using oneButtonVelvet, an optimizer program that runs the Velvet assembler (Zerbino and Birney, 2008) numerous times over a user supplied k-mer range while varying several of the assembler parameters and optimizing for the assembly parameter set which yields the longest N50 contig length.

Annotation

Coding sequences were predicted using GeneMark v3.3 (Borodovsky and McIninch, 1993) and GLIMMER v2.13 (Salzberg *et al.*, 1998). Intergenic regions not spanned by GeneMark and GLIMMER were aligned using BLAST against NCBI's non-redundant database and predictions were generated based on protein alignments. tRNA genes were determined using tRNAscan-SE 1.23 and non-coding RNA genes were determined by RNAmmer-1.2 and Rfam v8.0. The final gene set was processed through a suite of protein categorization tools consisting of Interpro, psort-b and KEGG. The gene product naming comes from the BER pipeline (JCVI). A more detailed standard operating protocol (SOP) can be found at http://hmpdacc.org/doc/sops/reference_genomes/annotation/WUGC_SOP_DACC.pdf.

71 *P. acnes* Genome Analysis and Comparison**Identification of the Core Regions of *P. acnes* Genomes**

The "core" regions were defined as genome sequences that are present in all 71 genomes. *P. acnes* KPA171202 was used as the reference genome. Each of the other 70 genome sequences (a series of contigs in most of the genomes and two complete genomes) was mapped to the reference genome using Nucmer (Kurtz *et al.*, 2004). All the 70 ".coords" output files of Nucmer program were analyzed to identify overlap regions based on the KPA171202 coordinates using a Perl script. Finally, "core" sequences were extracted based on the genome sequence of KPA171202 with the coordinates calculated above. On average, 90% (ranging from 88% to 92%) of the genomes were included in the core regions.

Identification of SNPs in the Core Regions

Single nucleotide polymorphisms (SNPs) were identified by using “show-snps” utility option of the Nucmer program (Kurtz *et al.*, 2004) with the default settings.

Genome sequence of *P. acnes* KPA171202 was used as the reference genome. All the 70 “.snps” output files of Nucmer program were analyzed to identify unique SNP positions based on the KPA171202 coordinates using a Perl script. The SNPs in the core regions were further analyzed to construct a phylogenetic tree.

Phylogenetic Tree Construction

The 71 concatenated sequences of the 96,887 SNP nucleotides in the core regions were used to construct a phylogenetic tree of the *P. acnes* genomes. The evolutionary distance of the core regions among the genomes was inferred using the Neighbor-Joining method (Saitou and Nei, 1987). The bootstrap tree inferred from 1,000 replicates was taken. Branches corresponding to partitions reproduced in less than 80% bootstrap replicates were collapsed. Figure 8 shows only the topology. In Figure 9, the tree was drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the p-distance method and are in the units of the number of nucleotide differences per site. This tree shows the comparison based on only the core regions. The distance does not represent the true evolutionary distance between different genomes, since the non-core regions of each genome were not considered here. All positions containing gaps and missing data were eliminated. Evolutionary analysis was conducted using MEGA5 (Tamura *et al.*, 2007).

Gene Content Comparison

In order to assess the conservation of gene content across the 71 genomes, protein coding genes in all the genomes were clustered using UCLUST (Edgar, 2010) by first sorting by decreasing length then clustering each sequence to an existing seed sequence if it had at least 90% nucleotide identity over its entire length, otherwise it became a new seed. For visualization, the data were reformatted to columns and rows representing genes and genomes, respectively. One or more copies of the genes in a genome were treated as present. Gene columns were sorted by their position based on the coordinates of the HL096PA1 genome, a fully finished genome with a 55Kb plasmid. Genome rows were sorted by their positions in the SNP-based Neighbor Joining tree described above.

Identification of CRISPR/Cas

CRISPRFinder (Grissa *et al.*, 2007) was used to identify the CRISPR repeat-spacer sequences. The annotation of HL110PA3 was used for BLAST alignment in order to identify the presence of CRISPR/Cas structure and CRISPR repeat-spacer sequences in strains of HL001PA1, HL060PA1, HL082PA2, HL103PA1, HL106PA1, HL110PA4 and J139. Each spacer sequence was annotated by BLAST alignment against NCBI's non-redundant nucleotide database and the reference genomic sequences database (refseq_genomic).

Sequence Coverage Analysis

MAQ (Li *et al.*, 2008) was used to map the raw sequence reads from Illumina/Roche platform to the reference genomes. Briefly, "map" command was used for mapping, and "assemble" command was used for calling the consensus sequences from read mapping, then "cnd2win" command was used to extract information averaged in a tilling window. A window size of 1,000 bp was used.

Randomly selected 1 million reads were used for mapping. This accounted for approximately 40X coverage for all the genomes except HL096PA2, HL096PA3, HL097PA1 and HL099PA1, which had approximately 55X to 75X coverage. BWA (Li and Durbin, 2010) was used to map the raw sequence reads from Roche/454 platform to the reference genome HL096PA1. The average coverage was calculated in 1,000 bp window.

Quantitative PCR

Quantitative PCR (qPCR) targeting *TadA* on the plasmid (Locus 3) and housekeeping genes *Pak* and *RecA* on the chromosome was performed using the genomic DNA extracted from the *P. acnes* isolates. LightCycler 480 High Resolution Melting Master kit was used (Roche Diagnostics GmbH, Mannheim, Germany). Each 10 µL reaction solution was consisted of 5 µL master mix (2X concentrate), 1 µL 25 mM MgCl₂, 0.5 µL 4 µM forward and reverse primers, and DNA template. Four qPCR runs were performed on Roche LightCycler 480. Primer sequences for *TadA* are 5'-GATAATCCGTTTCGACAAGCTG-3' (forward) and 5'-ACCCACCACGATGATGTTT-3' (reverse). Primer sequences for *pak* are 5'-CGACGCCTCCAATAACTTCC-3' (forward) and 5'-GTCGGCCTCCTCAGCATC-3' (reverse). Primer sequences for *recA* are 5'-CCGGAGACAACGACAGGT-3' (forward) and 5'-GCTTCCTCATACCACTGGTCATC-3' (reverse). All samples were run in duplicates in each qPCR run, except the second run, which was not duplicated. Thermocycling conditions were as following: initial activation step of 10 minutes at 95oC; 50 amplification cycles with each consisting of 10 seconds at 95oC, 15 seconds at 65oC in the first cycle with a stepwise 0.5oC decrease for each succeeding cycle, and 30 seconds at 72oC; and final melting curve step starting at 65oC and ending at 99oC with a ramp rate of 0.02 oC/s and acquisition rate of 25/oC.

DNA concentration standards were run in duplicates. Copy number ratios of genes were calculated based on the concentrations of the genes on the plasmid and chromosome.

Data Availability

5 16S rDNA sequences have been deposited at GenBank under the project ID 46327. Whole genome shotgun sequences and annotations of the *P. acnes* strains have been deposited at GenBank under the accession numbers ADWB000000000, ADWC000000000, ADWF000000000, ADWH000000000, ADWI000000000, ADXP000000000, ADXQ000000000, ADXR000000000, ADXS000000000, 10 ADXT000000000, ADXU000000000, ADXW000000000, ADXX000000000, ADXY000000000, ADXZ000000000, ADYA000000000, ADYB000000000, ADYC000000000, ADYD000000000, ADYE000000000, ADYF000000000, ADYG000000000, ADYI000000000, ADYJ000000000, ADYK000000000, ADYL000000000, ADYM000000000, ADYN000000000, ADYO000000000, ADYP000000000, 15 ADYQ000000000, ADYR000000000, ADYS000000000, ADYT000000000, ADYU000000000, ADYV000000000, ADYW000000000, ADYX000000000, ADYY000000000, ADYZ000000000, ADZA000000000, ADZB000000000, ADZC000000000, ADZD000000000, ADZE000000000, ADZF000000000, ADZG000000000, ADZH000000000, ADZI000000000, ADZJ000000000, ADZK000000000, 20 ADZL000000000, ADZM000000000, ADZN000000000, ADZO000000000, ADZP000000000, ADZQ000000000, ADZR000000000, ADZS000000000, ADZT000000000, ADZV000000000, ADZW000000000, CP003293, and CP003294.

EXAMPLE 2 – PAN-GENOME AND COMPARATIVE GENOME ANALYSIS OF PROPIONIBACTERIUM ACNES

Propionibacterium acnes is a major human skin bacterium. To understand whether different strains have different virulent properties and thus play different roles in health and diseases, the genomes of 82 *P. acnes* strains, most of which were isolated from acne or healthy skin, were compared. Lineage-specific genetic elements were identified that may explain the phenotypic and functional differences of *P. acnes* as a commensal in health and as a pathogen in diseases. By analyzing a large number of sequenced strains, an improved understanding of the genetic landscape and diversity of the organism at the strain level and at the molecular level is provided.

Introduction

Propionibacterium acnes is a major commensal of the human skin. It contributes to maintaining the skin health by inhibiting the invasion of common pathogens, such as *Staphylococcus aureus* and *Streptococcus pyogenes*. It does so by hydrolyzing triglycerides and releasing free fatty acid that contributes to the acidic pH of the skin surface (1). On the other hand, *P. acnes* has been historically linked to acne vulgaris, a chronic inflammatory disease of the pilosebaceous unit affecting more than 85% of adolescents and young adults (2). A metagenomic study previously demonstrated that *P. acnes* was a dominant bacterium in the pilosebaceous unit in both healthy individuals and acne patients (3, 4). At the strain level, however, the population structures of *P. acnes* were different between the two groups. These findings suggested that microbe-related human diseases are often caused by certain strains of a species rather than the entire species, in line with the studies of other diseases (5, 6).

P. acnes has been classified into three distinct types. Studies by Johnson and Cummins (7) first revealed two distinct phenotypes of *P. acnes*, known as types I and II, that could be distinguished based on serological agglutination tests and cell wall sugar analysis. McDowell *et al.* (8) differentiated types I and II *P. acnes* by
5 monoclonal antibody typing. Furthermore, their phylogenetic analysis of *P. acnes* strains based on the nucleotide sequences of the *recA* gene and a more variable hemolysin/cytotoxin gene (*tly*) demonstrated that types I and II represent distinct lineages. Their investigations also revealed that strains within the type I lineage could be further split into two clades, known as types IA and IB (8, 9). An additional
10 phylogenetic group of *P. acnes*, known as type III was described later (10). Recent studies based on multilocus sequence typing (MLST) further sub-divided *P. acnes* into closely related clusters, some of which were associated with various diseases including acne (11-13).

The first complete genome sequence of *P. acnes*, KPA171202, a type IB
15 strain, provided insights on the pathogenic potential of this Gram-positive bacterium (14). The genome is 2.56 M bp with 60% of GC content. It encodes 2,333 open reading frames (ORFs) including multiple gene products involved in degrading host molecules, such as sialidases, neuraminidases, endoglycoceramidases, lipases, and pore-forming factors. However, the sequence of a single genome does not reflect
20 the genetic landscape of the organism and how genetic variations among strains determine their various phenotypes and pathogenic properties.

To better understand the human microbiome variations at the strain level, as part of the Human Microbiome project (HMP) (15, 16), previously generated were the reference genome sequences of 66 *P. acnes* strains selected from a collection of
25 over 1,000 strains isolated from a cohort of healthy subjects and acne patients (4).

These 66 strains represent the major lineages of *P. acnes* found on the human skin, including types IA, IB, and II. To cover all the main *P. acnes* lineages in the analysis, three additional *P. acnes* strains were sequenced, including the first available type III *P. acnes* genome. Thirteen *P. acnes* genomes sequenced by other research groups (14, 17-22) were also available at the time of analysis. With a total of 82 genomes, performed was a comparative genome analysis to characterize the pan-genome of *P. acnes*, the phylogenetic relationships among different lineages, the microevolution of the strains in the same individual microbiome, and the genetic elements specific to each lineage and their associations with health and disease.

Results

***P. acnes* Strains and General Genome Features**

To understand the genomic diversity of this important skin commensal at the strain level, the genomes of 69 sequenced *P. acnes* strains were analyzed. Among them, 67 *P. acnes* strains were isolated from the skin of healthy individuals and acne patients (3, 4), and two *P. acnes* strains, HL201PA1 and HL202PA1, were isolated from refractory endodontic lesions (23) (Table 2-1).

Table 2-1 – General features of the 82 *P. acnes* genomes

Genome	Strain name	Origin	Ribotype	recA type	Sequencing method	Genome size (Mb)	GC %	Number of ORFs
1	HL001PA1	Skin	2	II	Illumina	2.49	60.0	2,661
2	HL002PA1	Skin	3	IB	Illumina	2.48	60.1	2,549
3	HL002PA2	Skin	1	IA	Illumina	2.48	60.1	2,594
4	HL002PA3	Skin	1	IA	Illumina	2.48	60.1	2,665
5	HL005PA1	Skin	4	IA	Illumina	2.53	60.2	2,724
6	HL005PA2	Skin	1	IA	Illumina	2.48	60.0	2,645
7	HL005PA3	Skin	1	IA	Illumina	2.48	60.1	2,570
8	HL005PA4	Skin	3	IB	Illumina	2.47	60.0	2,607
9	HL007PA1	Skin	4	IA	Illumina	2.53	60.2	2,691
10	HL013PA1	Skin	3	IB	Illumina	2.48	60.0	2,618
11	HL013PA2	Skin	1	IA	Illumina	2.48	60.1	2,588
12	HL020PA1	Skin	1	IA	Illumina	2.48	60.1	2,554
13	HL025PA1	Skin	1	IB	Illumina	2.54	60.1	2,581
14	HL025PA2	Skin	3	IB	Illumina	2.48	60.0	2,610
15	HL027PA1	Skin	3	IB	Illumina	2.53	60.1	2,711
16	HL027PA2	Skin	1	IA	Illumina	2.48	60.1	2,629
17	HL030PA1	Skin	1	IB	Illumina	2.54	60.0	2,662
18	HL030PA2	Skin	3	IB	Illumina	2.51	60.1	2,647
19	HL036PA1	Skin	532	IA	Illumina	2.48	60.1	2,575
20	HL036PA2	Skin	532	IA	Illumina	2.48	60.1	2,585
21	HL036PA3	Skin	1	IA	Illumina	2.48	60.1	2,601
22	HL037PA1	Skin	3	IB	Illumina	2.48	60.1	2,617
23	HL038PA1	Skin	4	IA	Illumina	2.54	60.2	2,663
24	HL042PA3	Skin	6	II	Roche/454	2.53	60.1	2,610
25	HL043PA1	Skin	5	IA	Illumina	2.53	60.2	2,598
26	HL043PA2	Skin	5	IA	Illumina	2.53	60.2	2,688
27	HL045PA1	Skin	4	IA	Illumina	2.53	60.2	2,692
28	HL046PA1	Skin	3	IB	Illumina	2.48	60.0	2,599
29	HL046PA2	Skin	1	IA	Illumina	2.53	60.1	2,692
30	HL050PA1	Skin	3	IB	Illumina	2.48	60.0	2,652
31	HL050PA2	Skin	1	II	Illumina	2.48	60.1	2,581
32	HL050PA3	Skin	3	IB	Illumina	2.48	60.0	2,558
33	HL053PA1	Skin	4	IA	Illumina	2.53	60.2	2,632
34	HL053PA2	Skin	8	IB	Illumina	2.51	60.1	2,664
35	HL056PA1	Skin	4	IA	Illumina	2.48	60.1	2,581
36	HL059PA1	Skin	16	IB	Illumina	2.48	60.1	2,570
37	HL059PA2	Skin	16	IB	Illumina	2.48	60.0	2,535
38	HL060PA1	Skin	2	II	Illumina	2.48	60.1	2,601
39	HL063PA1	Skin	1	IA	Illumina	2.40	60.1	2,520
40	HL063PA2	Skin	3	IB	Illumina	2.53	60.0	2,669
41	HL067PA1	Skin	3	IB	Illumina	2.53	60.1	2,633
42	HL072PA1	Skin	5	IA	Illumina	2.53	60.1	2,594
43	HL072PA2	Skin	5	IA	Illumina	2.53	60.1	2,672
44	HL074PA1	Skin	4	IA	Illumina	2.53	60.2	2,723
45	HL078PA1	Skin	1	IA	Illumina	2.58	60.1	2,785
46	HL082PA1	Skin	8	IB	Illumina	2.50	60.1	2,648
47	HL082PA2	Skin	2	II	Illumina	2.51	60.0	2,644
48	HL083PA1	Skin	1	IA	Illumina	2.48	60.1	2,575
49	HL083PA2	Skin	3	IB	Illumina	2.48	60.0	2,633
50	HL086PA1	Skin	8	IB	Illumina	2.53	60.1	2,610
51	HL087PA1	Skin	3	IB	Illumina	2.48	60.1	2,584
52	HL087PA2	Skin	1	IA	Illumina	2.48	60.1	2,572
53	HL087PA3	Skin	3	IB	Illumina	2.52	60.1	2,619
54	HL092PA1	Skin	8	IB	Illumina	2.50	60.1	2,590
55	HL096PA1	Skin	5	IA	Roche/454	2.49	60.0	2,393
56	HL096PA2	Skin	5	IA	Illumina	2.56	60.1	2,638
57	HL096PA3	Skin	1	IA	Illumina	2.56	60.0	2,651
58	HL097PA1	Skin	5	IC	Illumina	2.52	60.2	2,617
59	HL099PA1	Skin	4	IA	Illumina	2.58	60.1	2,735
60	HL100PA1	Skin	1	IA	Illumina	2.48	60.1	2,582
61	HL103PA1	Skin	2	II	Illumina	2.48	60.1	2,546
62	HL106PA1	Skin	2	II	Illumina	2.48	60.1	2,533
63	HL106PA2	Skin	1	IA	Illumina	2.48	60.1	2,567
64	HL110PA1	Skin	8	IB	Illumina	2.51	60.1	2,667
65	HL110PA2	Skin	8	IB	Illumina	2.50	60.1	2,614
66	HL110PA3	Skin	6	II	Illumina	2.54	60.1	2,806
67	HL110PA4	Skin	6	II	Illumina	2.54	60.1	2,724
68	HL201PA1	Refractory endodontic lesion	6	III	Illumina	2.48	60.1	2,629
69	HL202PA1	Refractory endodontic lesion	Not assigned	II	Illumina	2.56	60.0	2,821
70	KPA171202	Plaque	1	IB	Sanger	2.56	60.0	2,297
71	J139	Skin	2	II	Roche/454	2.48	60.0	2,384
72	J165	Skin	1	IA	Roche/454	2.50	60.0	2,403
73	SK137	Skin	1	IA	Roche/454	2.50	60.0	2,352
74	SK187	Skin	3	IB	Roche/454	2.51	59.0	2,381
75	SK182	Skin	1	IA	Roche/454	2.48	60.0	2,338
76	266	Pleuropulmonary infection	1	IA	Roche/454	2.50	60.0	2,412
77	6609	Skin	1	IB	Solid	2.58	60.0	2,358
78	ATCC11828	Abscess	2	II	Solid	2.49	60.0	2,260
79	P.acn17	Corneal scrape	3	IB	Solid	2.52	60.0	2,266
80	P.acn31	Aqueous humour	3	IB	Solid	2.50	60.0	2,247
81	P.acn33	Aqueous humour	3	IB	Solid	2.49	60.0	2,236
82	PRP-38	Skin	5	IC	Solid	2.51	60.0	2,233

These 69 strains cover all the known *P. acnes* lineages isolated to date. The strains were classified based on their 16S ribosomal RNA (rRNA) sequences. Each unique 16S rRNA sequence was defined as a ribotype (RT). All the sequenced *P. acnes* genomes had three identical copies of 16S rRNA. Based on the metagenomic study of the skin microbiome associated with acne (4), among the top ten major ribotypes, RT1, RT2, and RT3 were the most abundant and found in both healthy individuals and acne patients with no significant differences. RT4, RT5, and RT8, however, were enriched in acne patients, while RT6 was mostly found in healthy individuals. The 69 strains included 19 RT1 strains, five RT2 strains, 15 RT3 strains, eight RT4 strains, seven RT5 strains, four RT6 strains, six RT8 strains, four strains of minor ribotypes, and one type III strain. The average genome size was 2.50 Mb (ranging from 2.46 to 2.58 Mb) and the GC content was 60%. On average, each genome encoded 2,626 ORFs (ranging from 2,393 to 2,806) (Table 2-1).

The analysis included 13 additional *P. acnes* genomes that were publicly available (14,17-22) (Table 2-1). The average genome size of these 13 *P. acnes* strains was 2.51 Mb (ranging from 2.48 to 2.56 Mb) and the GC content was 60%, encoding 2,319 ORFs on average (ranging from 2,233 to 2,412). These 13 genomes include six RT1 strains, two RT2 strains, four RT3 strains, and one RT5 strain, however, no genomes of RT4, RT6, RT8 and type III strains were available. The sequencing effort significantly increased the number of genomes for each *P. acnes* lineage as well as the number of lineages covered.

***P. acnes* Pan-Genome**

To determine the genetic landscape of *P. acnes*, the pan-genome based on the 82 *P. acnes* genomes was estimated. The number of new genes that would be discovered by sequencing additional *P. acnes* genomes by using a power law regression analysis, $n=kN^\gamma$ (24), was estimated (Figure 13). The analysis identified that α was 0.788. The average number of new genes added by a novel genome was three when the 82nd genome was added. The number of *P. acnes* pan-genes that would be accumulated by sequencing additional *P. acnes* genomes by using a power law regression analysis, $n=kN^\gamma$, was then estimated (Figure 14). The exponent γ was 0.067, and *P. acnes* had 3,136 pan-genes ($N=82$). Based on these results, the pan-genome of *P. acnes* is defined as open, as the exponent α was less than one and γ was greater than zero (24). However, since α was close to one and γ was close to zero, it is believed that this organism evolved tightly without large expansions.

Phylogenetic Relationships Among the *P. acnes* Genomes

A genome comparison of the 82 *P. acnes* strains revealed that 2.20 Mb (88% of the average genome) was shared by all the *P. acnes* genomes, which are referred to herein as the “core regions.” Within the core regions, 123,223 unique single nucleotide polymorphisms (SNPs) were detected among the strains. Twenty seven percent of the SNPs were unique to type I, 22% were unique to type II, and 22% were unique to type III (Figure 15). A phylogenetic tree based on the 123,223 SNPs in the core regions was constructed (Figure 16). The tree showed that the *recA* type classification of the strains was consistent with the major clades based on the genomes. The *recA* type IA, IB, and II strains were all clustered together within each type, respectively, except HL097PA1 and PRP-38. The only *recA* type III strain,

HL201PA1, formed a separate branch from type I and type II strains. The tree also showed that the 16S rRNA ribotypes of the strains were consistent with the phylogenetic relationships inferred from the genome sequences. Most of the RT1 strains were clustered in clade IA-1, while all the RT4 and most of the RT5 strains were clustered in clade IA-2. All six RT8 strains were clustered together in clade IB-1. All the RT3 and RT16 strains were clustered together in clade IB-2 except SK187. HL030PA1 and KPA171202 were clustered together with 6609, as a distinct IB-3 clade. HL097PA1 and PRP-38 were clustered together and were classified as a novel type IC recently named by McDowell *et al.* (22). All the RT2 strains were clustered in clade II, distant from clade I, together with RT6 strains. HL202PA1, which is a RT6 strain and was isolated from an oral site, was not much different from the skin RT6 isolates and was clustered together. The sequence types of all the strains were assigned based on two published MLST schemes (11,13) and are shown in Table S1. The phylogenetic tree based on the core genome regions demonstrated that 16S ribotyping can be used for *P. acnes* strain identification and classification. It provides a much higher resolution than recA typing, and in the meantime, it is much simpler and faster with only one gene required than MLST, which is a laborious process generally requiring 7-9 genes.

The large number of genome sequences that were generated permitted analyzing the *P. acnes* pan-genome at the clade level. Clades IA, IB and II had 36, 33 and 12 genomes, respectively. Based on the power law regression analyses described above, it was determined that at the clade level *P. acnes* also has an open pan-genome for recA type IA clade, type IB clade and type II clade with limited expansions (Figure 17). The expansion rates were not significantly different among

the clades and were similar to the one at the species level. This suggests that all the major lineages of *P. acnes* had evolved at a similar rate.

SNP Distribution in the Core Genome Regions

To understand whether there are “hot spots” for mutation and/or recombination in the *P. acnes* genomes, it was determined whether the SNPs were randomly distributed throughout the genomes or were enriched in particular regions. The frequency of SNPs in each protein coding gene in the core regions was calculated. The average rate of polymorphic sites in the core regions was 5.3%, *i.e.*, 5.3 unique SNPs in every 100 bp. This rate is comparable to the ones found in multiple gut bacterial genomes (25). Among the 1,888 genes encoded in the core regions, 55 genes had higher SNP frequencies with more than two standard deviations (SD), and 47 genes with more than three SD (Figure 18a). Using the Kolmogorov–Smirnov (K-S) test, it was demonstrated that these 102 highly mutated genes were not randomly distributed throughout the genome ($P < 0.01$) (Figure 18b). This suggests that *P. acnes* has an evolutionary risk management strategy. Based on the Clusters of Orthologous Groups (COG) categories, the functions of these 102 genes showed a similar distribution as those of all 1,888 genes in the core regions. There was no enrichment of a particular functional category in these frequently mutated genes.

It was further determined whether the mutations in the core regions were under selection by calculating the ratio of non-synonymous (NS) vs. synonymous (S) SNPs for the 1,888 genes. The average rate of NS mutations was 38%. Among the 1,888 genes, 54 genes had higher NS mutation rates with more than two SD and 13 genes with more than three SD (Figure 18c). These 67 genes were randomly distributed in the genome and not particularly enriched in certain regions ($P > 0.05$)

with the K-S test) (Figure 18d). Most of the 102 genes with higher SNP frequencies did not overlap with these 67 genes, suggesting that independent evolutionary events might lead to these gene alternations. Only ten genes had both high SNP frequencies and high NS mutation rates, all annotated as hypothetical proteins.

5 **Evolutionary Relationships of the Strains Isolated from the Same Individuals**

The large number of *P. acnes* strains isolated from the cohort of acne patients and healthy individuals allowed the investigation of whether the *P. acnes* strains in hair follicles from the same individual were clonal. Based on previous metagenomic analysis, it was demonstrated that most individuals harbored multiple *P. acnes* strains from different lineages (4). However, it was not known whether the strains of the same lineage in the same individual were derived from the same ancestor. Genome sequences of the strains isolated from the same samples makes it possible to examine whether the Strains from the Same Individuals (SSIs) evolved from the same origin via clonal expansion. The 69 sequenced *P. acnes* strains included 49 SSIs: 13 duets (*i.e.*, 13 pairs of strains isolated from 13 individuals), five trios, and two quartets. Twenty three SSIs were clustered in the same clades, nine in clade IA-1, four in clade IA-2, two in clade IB-1, six in clade IB-2 and two in clade II. The distance (substitution rate at the 123,223 SNP sites in the core regions) between each pair of SSIs was calculated (Figure 16). The average distance of the SSIs in clade IA-1 was 0.0014, while that of strains from different individuals in clade IA-1 was 0.0064 ($P < 0.001$). Consistent results were observed in other clades including IA-2, IB-1, IB-2, and II (Figure 19a). This demonstrated that the SSIs in the same lineage were significantly more similar to each other than the strains isolated from different individuals, suggesting that they were clonal in each individual. Among the RT4 and RT5 strains within clade IA-2, however, the average distance between SSIs

(0.0004) was not significantly different from the average distance between strains from different individuals (0.0017) ($P=0.072$). Moreover, the average distance between RT4/RT5 strains from different individuals (0.0017) was similar to the average distance between the SSIs in clade IA-1 (0.0014), and even shorter than the average distances between the SSIs in clades IB-1 (0.0059), IB-2 (0.0019) and II (0.0022) (Fig. S3A). This suggests that although isolated from different individuals, these RT4 and RT5 strains seemed to be clonal and had evolved from the same recent ancestor. A similar relationship between the two RT5 strains in clade IC were observed, where HL097PA1 and PRP-38, isolated from different individuals, were closely related to each other with a distance of 0.0012. The metagenomic study has demonstrated a strong association of strains of RT4 and RT5 with acne (4). The clonality of these strains isolated from different individuals suggests that RT4 and RT5 strains may be transmitted among individuals. This finding is consistent with the previous clinical report that antibiotic-resistant *Propionibacteria* were transmissible between acne-prone individuals including dermatologists (26), as most of the antibiotic-resistant *P. acnes* strains belong to RT4 and RT5 (4,13). The analysis of SSIs further supports the theory that RT4 and RT5 strains may be a pathogenic factor in acne.

To determine whether the distances between strain pairs from the same individuals but belonging to different lineages were different from random strain pairs, the distances of any pair of SSIs from different clades were calculated. The average distance of the SSIs between clades IA-1 and IA-2 (*i.e.*, HL005PA3 vs. HL005PA1, HL005PA2 vs. HL005PA1, HL096PA3 vs. HL096PA1, and HL096PA3 vs. HL096PA2) was 0.039, similar to that of the isolates from different individuals (0.040). Similar results were obtained for all other clade pair comparisons (Figure 19b).

These results demonstrated that the SSIs from different clades were similarly different from each other as to the strains from different individuals. This analysis suggests that in each individual microbiome *P. acnes* strains undergo clonal expansion in the same population, while multiple strain populations can often co-exist in the same community with little recombination.

Non-core Genome Regions in Type I Strains

By comparing the genome sequences of the 82 *P. acnes* strains, non-core genome regions were identified that were not shared by all 82 strains. The total length of the non-core regions was approximately 0.90 Mb. The average GC content of the non-core regions was slightly lower than that of the core regions, $58\% \pm 6.9\%$, suggesting that part of the non-core regions might be originated from other species via horizontal gene transfer.

Different lineages of *P. acnes* strains have distinct non-core regions. Using hierarchical clustering of the non-core regions, it was shown that the strains of the same ribotypes were clustered together with distinct separations among the clades (Figure 20). Among the non-core regions, the genetic elements specific to each lineage were identified, which may explain the phenotypic and functional differences of the strains in health and disease. In clade IA-2, genomic loci 1, 2 and 3 were identified, which were unique to mainly the RT4 and RT5 strains (4). These loci appear to be originated from mobile elements, encode several virulent genes, and may contribute to the virulence of these strains. In the meantime, the genomic island-like cluster 2 (GI2) (18) was uniquely absent in most of the strains in this clade. Clade IB-1 consisted of all RT8 strains, which were also highly associated with acne based on our metagenomic study (4). They all have a unique genomic island (locus 4), which is 20 Kb long and encodes a series of nonribosomal peptide synthetases

(NRPS), which may contribute to increased virulence of these strains. Most RT3 and RT16 strains belong to clade IB-2 and have fewer non-core regions than the strains in other clades. This may be explained by the lack of entire rearrangement hot spot (RHS) family proteins, which function in genomic rearrangements as previously implicated in *Escherichia coli* (27). Clade IB-3 consisted of three strains, including KPA171202. Three of the four genomic islands described previously (18), GI1, GI3 and GI4, were unique to this clade and were absent in all other strains. This analysis suggests that KPA171202, although was the first sequenced complete genome of *P. acnes*, did not seem to be a common skin *P. acnes* strain representing one of the major lineages. This result is consistent with previous studies using MLST (11-13). Strains of clade IC belong to RT5. They also contain locus 3, a linear plasmid, which is highly homologous to the locus 3 in the RT4 and RT5 strains of clade IA-2 and encodes a tight adhesion locus originated from *Clostridium leptum* (4). In general, although strains in different lineages had a similar genome size with similar gain and loss of genetic materials, they harbor distinct genetic elements which may give rise to their different virulent properties.

Non-core genome regions in type II strains

Strains in clade II, mainly RT2 and RT6, were more distantly related to the strains in clade I. Based on the metagenomic study, strains in clade II were not associated with acne, as RT2 was evenly distributed between acne patients and healthy individuals, while RT6 was significantly enriched in healthy individuals (4). Compared to type I strains, the genomes of RT2 and RT6 strains lack several regions, which are approximately 92 Kb long in total and encode 107 ORFs. RT2 and RT6 genomes have additional genomic regions with a similar size encoding 93 ORFs (Figure 20). Based on the COG classification, there were no significant

differences in the distribution of the functional categories between the 107 type I specific ORFs and 93 type II specific ORFs.

The most unique genomic feature of RT2 and RT6 strains is represented by the clustered regularly interspaced short palindromic repeats (CRISPR)/Cas locus

(4). CRISPR/Cas system provides acquired bacterial immunity against viruses and plasmids by targeting nucleic acids in a sequence-specific manner (28). All the sequenced strains of RT2 and RT6 encoded a complete set of CRISPR/Cas genes and at least one repeat and spacer sequence, while none of the other ribotype strains did. Based on its complete genome sequence (20), strain ATCC11828 appeared to be an exception, having only terminal sequence but no spacer sequence. However, using PCR and sequencing it was determined that ATCC11828 has one repeat-spacer sequence (Table S2).

Table S2. CRISPR spacer sequences found in the genomes of ATCC 11828, HL042PA3 and HL202PA1.

Strain	Ribotype	Spacer	Protospacer	BLAST result	Match found in
ATCC11828	RT2	1	CATCTGCCAACGAGCGAGAGTGGCGGGTGTTC	No hits	Locus 2
		2	CGAGGGCTACCACTGGTTCGATTTGGACTGTG	<i>Clostridium leptum</i> DSM 753, CLOLEP_00167, <i>Propionibacterium acnes</i> SK137, HMPREF0875_3193 (Domain of unknown function)	
		3	CAGGCGCTCCACTCCCTCGCCCTGGCCCAAC	No hits	
HL042PA3	RT6	1	CTGACTGGTTTGGTTCATAGTCTTCTGACAG	<i>Propionibacterium acnes</i> phage PA8 gp14 (tape measure protein) <i>Propionibacterium acnes</i> phage PAD20 gp14 (Tape measure protein) <i>Propionibacterium acnes</i> phage PAS50 gp14 (Tape measure protein) <i>Propionibacterium acnes</i> phage PA8 gp15 (Minor tail protein) <i>Propionibacterium acnes</i> phage PAD20 gp15 (Minor tail protein) <i>Propionibacterium acnes</i> phage PAS50 gp15 (Minor tail protein)	Locus 2
		2	TCACAGGCCACGCGAGGCACATCACCTTATTAG	<i>Propionibacterium acnes</i> phage PAD20 gp15 (Minor tail protein) <i>Propionibacterium acnes</i> phage PAS50 gp15 (Minor tail protein)	
		3	CTCCCCCTCTCCCCGGGAGGAAAAGCAGACCA	<i>Propionibacterium acnes</i> phage PAS50 gp15 (Minor tail protein)	
		4	CGAGGGCTACCACTGGTTCGATTTGGACTGTG	<i>Clostridium leptum</i> DSM 753, CLOLEP_00167, <i>Propionibacterium acnes</i> SK137, HMPREF0875_3193 (Domain of unknown function)	
HL202PA1	RT6	1	CGAGGGCTACCACTGGTTCGATTTGGACTGTG	<i>Clostridium leptum</i> DSM 753, CLOLEP_00167, <i>Propionibacterium acnes</i> SK137, HMPREF0875_3193 (Domain of unknown function)	Locus 2
		2	CAGGCGCTCCACTCCCTCGCCCTGGCCCAAC	No hits	

A total of 48 spacer sequences were found in the 11 RT2 and RT6 strains, 29 of which were unique. In other bacterial species, it has been established experimentally and computationally that the spacers at the leader-proximal end are more diversified, while the spacers at the leader-distal end are more conserved among strains. The evolutionary relationships among the RT2 and RT6 strains

based on their shared spacer sequences were analyzed. HL060PA1 and HL082PA2, which were clustered tightly in clade II, shared the same spacer S2 (Figure 21).

J139, ATCC11828, HL110PA4, HL110PA3, and HL202PA1 shared the same spacers S17 and S18 (Figure 21). These results suggest that these groups of

5 strains probably evolved from the same ancestors before having acquired additional spacers. The relationships among the strains based on shared CRISPR spacers are consistent with the phylogenetic relationships calculated based on the SNPs in the core regions. In addition, multiple type II strains harbored spacer sequences that match to the sequences in loci 2 and 3, which were unique to mainly acne-
10 associated RT4 and RT5 strains (4). The sequences in loci 2 and 3 appeared to be originated from *C. leptum* and encode potential virulence factors (Fig. S4). These loci may have been acquired by RT4 and RT5 strains, while the genomes of RT2 and RT6 that encode these spacers may be capable of eliminating the invasion of foreign DNA through the CRISPR mechanism (4).

15 The large number of high quality draft genome sequences enabled detection not only large genomic variations, but also small but essential genomic alterations. It was previously reported that type II strains showed decreased lipase activity (10).

Lipase functions in hydrolyzing triglycerides and releasing free fatty acids, which is thought to be essential in *P. acnes* virulence. Based on the genome annotation, 13
20 genes were identified with a potential function of lipase (Figure 22a). Among them, detected insertions/deletions ranging from one nucleotide to 13 nucleotides may explain the decreased lipase activity in type II strains. Two triacylglycerol lipases were encoded in tandem in *P. acnes* genomes, HMPREF0675-4855 and HMPREF0675-4856 (according to the annotation of SK137). All the type II strains
25 and IB-3 strains had a deletion of the "TATA-box" 20 bp upstream of the start codon

of the second lipase gene, HMPREF0675-4856 (Figure 22b). In addition, there was a one-nucleotide deletion at the position of 124G of the second lipase gene, leading to a frameshift and the introduction of a premature stop codon. These two deletions may potentially explain the decreased lipase activity and hence decreased virulence in acne observed in type II strains in previous studies (4, 10).

Non-core genome regions in the type III strain

Type III strains are rarely found on the skin surface. A type III *P. acnes* strain isolated from refractory endodontic lesion, HL201PA1, was sequenced. This first available type III genome permitted the identification of the genetic elements specific to this lineage. Compared to type I and type II strains, the genome of HL201PA1 lacks a few regions with a total length of 43 Kb (Figure 20). There were 42 ORFs encoded in these regions, including anaerobic dimethyl sulfoxide reductase (PPA0516-PPA0517), iron(III) dicitrate transport system permease (PPA0792-PPA0793), 3-isopropylmalate dehydratase (PPA1361-PPA1363), and maltose transport system permease (PPA1553-PPA1554).

Discussion

High-throughput genome sequencing and comparative analysis of a large number of related strains have been used to study the spread and microevolution of several pathogens at the strain level, including methicillin-resistant *Staphylococcus aureus* (29), *Streptococcus pneumoniae* (30), and *Vibrio cholerae* in Haiti outbreak (31), demonstrating the power of comparative genome analysis of multiple strains in improving our understanding of the bacterial pathogens. However, this approach has been rarely applied to study commensal species to understand their varied virulent potentials among different strains and their roles in both health and diseases.

This study presents a comparative genome analysis of a major skin commensal, *P. acnes*, based on a large number of sequenced strains. This collection of strains not only includes strains associated with either healthy skin or acne, but also a large number of strain pairs that were isolated from the same individuals. This allowed the comparison of phylogenetic relationships and microevolution of the *P. acnes* strains associated with health vs. disease as well as of the strains in the same individual microbiome.

By comparing 82 *P. acnes* genomes, it was shown that all *P. acnes* strains had a similar genome size with a similar GC content, encoding 2,577 ORFs on average (Table 1). Although *P. acnes* has an open pan-genome, unlike many other open-genome species (24), it has limited genome expansion with only a few new genes added per genome (Figures 13 and 14). The rate of genome expansion is similar within the major lineages (Figure 17). There was limited recombination among different *P. acnes* strains, and thus 16S rRNA ribotypes can be used as a proxy for *P. acnes* strain identification and classification (Figure 16). Compared to other typing methods, 16S ribotyping has a much higher resolution than recA typing and is much easier and faster than the traditional MLST method. This method can be applied in a high-throughput manner by combining with next-generation sequencing, and thus allows one to detect the microbiome variations at the strain level (4). This is advantageous and important, as identifying and understanding the strain level variations of the human microbiome is medically important.

The genomes of the sets of strains isolated from the same individual samples were compared (Figures 16 and 19). This collection of genome data is unique and no such kind of study has been performed to investigate the microevolution of a human commensal within an individual microbiome. It was found that while multiple

P. acnes strain populations co-existed in the same individual microbiome, clonal expansions occurred in each population with little recombination among different populations. Within each lineage, the strains isolated from the same individuals were significantly more similar to each other than strains from different individuals except the disease associated strains, RT4 and RT5 strains (Figure 19). Although isolated from different individuals, they appeared to be clonal and have evolved from the same virulent ancestor strain (Figure 16). This supports the observation that these strains were transmissible (26) and that they may play a role in acne pathogenesis (4). This finding is important and will help control the spread of antibiotic resistant strains and develop new targeted therapy for acne.

By analyzing the non-core regions, the genomic elements and alterations specific to each lineage were identified (Figure 20). These lineage-specific elements may render the strains different physiological and functional properties and thus lead to their different roles as a commensal in health or as a pathogen in diseases.

Among the acne associated strains, RT4 and RT5 strains encode three distinct loci originated from mobile elements, and RT8 strains encode a distinct region containing a set of NRPS. The virulent genes encoded in these strain-specific regions may explain the associations of these strains with acne and help the development of new drugs targeting against these strains. RT2 and RT6 strains, which were not associated with acne and were enriched in healthy skin, respectively, all encode CRISPR/cas elements. The CRISPR mechanism may prevent these strains from acquiring virulent genes from invading foreign mobile elements. In addition, these strains contain genomic variations in lipases that may alter lipid metabolism and reduce their virulence (Figure 22).

In conclusion, by characterizing the genetic landscape and diversity of *P. acnes* with a large number of genomes, genomic evidence that may explain the diverse phenotypes of *P. acnes* strains and a new insight into the dual role of this commensal in human skin health and disease is provided. The findings from this comparative genome analysis provide new perspectives on the strain diversity and evolution of commensals in the human microbiome. As many current microbiome studies focus on the associations of microbial communities with health and diseases, this study underscores the importance of understanding the commensal microbiome at the strain level (25). The findings from this study also shed light on new strain-specific therapeutics for acne and other *P. acnes* associated diseases.

Materials and Methods

***P. acnes* Strains**

Among the 69 *P. acnes* strains that were sequenced, 67 were isolated from the skin microcomedone samples from acne patients and healthy individuals (4).

The other two strains, HL201PA1 and HL202PA1, were isolated from refractory endodontic lesions (23), provided by Dr. David Beighton at the King's College London.

Whole Genome Shotgun Sequencing, Assembly, and Annotation

The genome of HL042PA3 was sequenced using Roche/454 FLX and assembled using a combination of PHRAP/CONSED (32) and GSMAPPER (Roche). HL201PA1 and HL202PA1 were sequenced using Illumina MiSeq (250 bp, paired-end) and assembled using Velvet (33). The remaining 66 genomes were sequenced previously as described (4). Coding sequences were predicted using GeneMark (34) and GLIMMER (35).

Computation of the Core Regions, Non-Core Regions and the Pan-Genome

The core regions were defined as genome sequences that were present in all 82 genomes, while the non-core regions were defined as genome sequences that were not present in all the genomes. KPA171202 was used as the reference

5 genome. Each of the other 81 genome sequences (a series of contigs in most of the genomes and ten complete genomes) was mapped to the reference genome using Nucmer (36). All the 81 ".coords" output files of Nucmer program were analyzed to identify overlap regions based on the KPA171202 coordinates using a Perl script.

Core sequences were then extracted based on the genome sequence of

10 KPA171202 with the coordinates calculated above.

The unique regions from each genome were added to the reference genome to make a "revised" reference genome, which contained the original sequence plus the unique genome sequences. This process was repeated for all the genomes until all the unique regions from all genomes were included in the pan-genome.

15 Lastly, core regions were subtracted from the pan-genome. The remaining regions were defined as non-core regions, which are not shared by all the strains. Protein coding sequences were predicted by GeneMark.hmm using KPA171202 as a reference file.

Identification of SNPs in the Core Regions

20 Single nucleotide polymorphisms (SNPs) were identified by using "show-snps" utility option of the Nucmer program with the default settings (36). Genome sequence of KPA171202 was used as the reference genome. All the 81 ".snps" output files of Nucmer program were analyzed to identify unique SNP positions based on the KPA171202 coordinates using a Perl script.

Phylogenetic Tree Construction

The 82 concatenated sequences of the 123,223 SNP nucleotides in the core region were used to construct a phylogenetic tree of the *P. acnes* genomes. MEGA5 (37) was used to calculate the distance based on the SNPs in the core regions using the Neighbor-Joining method and the p-distance method. The bootstrap tree inferred from 200 replicates was taken.

Sequence Type Analysis Based on MLST Schemes

The sequence types of the 82 isolates were determined based on the MLST schemes published previously (11-13). The MLST gene sequences were aligned using BLAST against all the alleles used in the two MLST schemes.

Identification of CRISPR/Cas

CRISPRFinder (38) was used to identify the CRISPR repeat-spacer sequences. The annotation of HL110PA3 was used for BLAST alignment in order to identify the presence of CRISPR/Cas structure and CRISPR repeat-spacer sequences in strains of HL001PA1, HL060PA1, HL042PA3, HL082PA2, HL103PA1, HL106PA1, HL110PA4, HL202PA1, J139 and ATCC11828. Each spacer sequence was annotated by BLAST (39) against NCBI's non-redundant nucleotide database and the reference genomic sequences database (refseq_genomic).

Hierarchical Clustering Analysis of the Non-Core Regions

Among the 1,685 non-core fragments (895,905 bp in total), 314 non-core fragments with a length of >500 bp (747,189 bp in total, corresponding to 83% of all the non-core regions) were extracted and used in hierarchical clustering of the non-core regions. Cluster 3.0 program (40) and average linkage method was used. The clustering matrix was composed of 314 rows and 82 columns, in which 1 denotes

presence of the non-core region and 0 denotes absence of the non-core region.

Java TreeView program (41) was used to display the clustering result.

References

1. Grice EA, Segre JA. 2011. The skin microbiome. *Nat Rev Microbiol* 9:244-253.
- 5 2. White GM. 1998. Recent findings in the epidemiologic evidence, classification, and subtypes of acne vulgaris. *J Am Acad Dermatol* 39:S34-37.
3. <http://precedings.nature.com/documents/5305/version/1>
4. Fitz-Gibbon ST, Tomida S, Chiu B, Nguyen L, Du C, Liu M, Elashoff D, Erfe MC, Loncaric A, Kim J, Modlin RL, Miller JF, Sodergren E, Craft N, Weinstock
10 GM, Li H. *Propionibacterium acnes* strain populations in the human skin microbiome associated with acne. *Journal of Investigative Dermatology* (in press).
5. Chambers HF, Deleo FR. 2009. Waves of resistance: *Staphylococcus aureus* in the antibiotic era. *Nat Rev Microbiol* 7:629-641.
- 15 6. Chase-Topping M, Gally D, Low C, Matthews L, Woolhouse M. 2008. Super-shedding and the link between human infection and livestock carriage of *Escherichia coli* O157. *Nat Rev Microbiol* 6:904-912.
7. Johnson JL, Cummins CS. 1972. Cell wall composition and deoxyribonucleic acid similarities among the anaerobic coryneforms, classical propionibacteria, and strains of *Arachnia propionica*. *J Bacteriol* 109:1047-1066.
20
8. McDowell A, Valanne S, Ramage G, Tunney MM, Glenn JV, McLorinan GC, Bhatia A, Maisonneuve JF, Lodes M, Persing DH, Patrick S. 2005. *Propionibacterium acnes* types I and II represent phylogenetically distinct groups. *J Clin Microbiol* 43:326-334.

9. Valanne S, McDowell A, Ramage G, Tunney MM, Einarsson GG, O'Hagan S, Wisdom GB, Fairley D, Bhatia A, Maisonneuve JF, Lodes M, Persing DH, Patrick S. 2005. CAMP factor homologues in *Propionibacterium acnes*: a new protein family differentially expressed by types I and II. *Microbiology* 151:1369-1379.
10. McDowell A, Perry AL, Lambert PA, Patrick S. 2008. A new phylogenetic group of *Propionibacterium acnes*. *J Med Microbiol* 57:218-224.
11. Lomholt HB, Kilian M. 2010. Population genetic analysis of *Propionibacterium acnes* identifies a subpopulation and epidemic clones associated with acne. *PLoS One* 5:e12277.
12. McDowell A, Gao A, Barnard E, Fink C, Murray PI, Dowson CG, Nagy I, Lambert PA, Patrick S. 2011. A novel multilocus sequence typing scheme for the opportunistic pathogen *Propionibacterium acnes* and characterization of type I cell surface-associated antigens. *Microbiology* 157:1990-2003.
13. McDowell A, Barnard E, Nagy I, Gao A, Tomida S, Li H, Eady A, Cove J, Nord CE, Patrick S. 2012. An Expanded Multilocus Sequence Typing Scheme for *Propionibacterium acnes*: Investigation of 'Pathogenic', 'Commensal' and Antibiotic Resistant Strains. *PLoS One* 7:e41480.
14. Bruggemann H, Henne A, Hoster F, Liesegang H, Wiezer A, Strittmatter A, Hujer S, Durre P, Gottschalk G. 2004. The complete genome sequence of *Propionibacterium acnes*, a commensal of human skin. *Science* 305:671-673.
15. The Human Microbiome Project Consortium. 2012. A framework for human microbiome research. *Nature* 486:215-221.
16. The Human Microbiome Project Consortium. 2012. Structure, function and diversity of the healthy human microbiome. *Nature* 486:207-214.

17. Nelson KE, Weinstock GM, Highlander SK, Worley KC, Creasy HH, Wortman JR, Rusch DB, Mitreva M, Sodergren E, Chinwalla AT, Feldgarden M, Gevers D, Haas BJ, Madupu R, Ward DV, Birren BW, Gibbs RA, Methe B, Petrosino JF, Strausberg RL, Sutton GG, White OR, Wilson RK, Durkin S, Giglio MG, Gujja S, Howarth C, Kodira CD, Kyrpides N, Mehta T, Muzny DM, Pearson M, Pepin K, Pati A, Qin X, Yandava C, Zeng Q, Zhang L, Berlin AM, Chen L, Hepburn TA, Johnson J, McCorrison J, Miller J, Minx P, Nusbaum C, Russ C, Sykes SM, Tomlinson CM, Young S, Warren WC, Badger J, Crabtree J, Markowitz VM, Orvis J, Cree A, Ferriera S, Fulton LL, Fulton RS, Gillis M, Hemphill LD, Joshi V, Kovar C, Torralba M, Wetterstrand KA, Abouelleil A, Wollam AM, Buhay CJ, Ding Y, Dugan S, FitzGerald MG, Holder M, Hostetler J, Clifton SW, Allen-Vercoe E, Earl AM, Farmer CN, Liolios K, Surette MG, Xu Q, Pohl C, Wilczek-Boney K, Zhu D. 2010. A catalog of reference genomes from the human microbiome. *Science* 328:994-999.
18. Brzuszkiewicz E, Weiner J, Wollherr A, Thurner A, Hupeden J, Lomholt HB, Kilian M, Gottschalk G, Daniel R, Mollenkopf HJ, Meyer TF, Bruggemann H. 2011. Comparative genomics and transcriptomics of *Propionibacterium acnes*. *PLoS One* 6:e21581.
19. Hunyadkurti J, Feltoti Z, Horvath B, Nagymihaly M, Voros A, McDowell A, Patrick S, Urban E, Nagy I. 2011. Complete genome sequence of *Propionibacterium acnes* type IB strain 6609. *J Bacteriol* 193:4561-4562.
20. Horvath B, Hunyadkurti J, Voros A, Fekete C, Urban E, Kemeny L, Nagy I. 2012. Genome sequence of *Propionibacterium acnes* type II strain ATCC 11828. *J Bacteriol* 194:202-203.

21. Voros A, Horvath B, Hunyadkurti J, McDowell A, Barnard E, Patrick S, Nagy I.
2012. Complete genome sequences of three *Propionibacterium acnes*
isolates from the type IA(2) cluster. J Bacteriol 194:1621-1622.
22. McDowell A, Hunyadkurti J, Horvath B, Voros A, Barnard E, Patrick S, Nagy I.
5 2012. Draft genome sequence of an antibiotic-resistant *Propionibacterium*
acnes strain, PRP-38, from the novel type IC cluster. J Bacteriol 194:3260-
3261.
23. Niazi SA, Clarke D, Do T, Gilbert SC, Mannocci F, Beighton D. 2010.
10 *Propionibacterium acnes* and *Staphylococcus epidermidis* isolated from
refractory endodontic lesions are opportunistic pathogens. J Clin Microbiol
48:3859-3869.
24. Tettelin H, Riley D, Cattuto C, Medini D. 2008. Comparative genomics: the
bacterial pan-genome. Curr Opin Microbiol 11:472-477.
25. Schloissnig S, Arumugam M, Sunagawa S, Mitreva M, Tap J, Zhu A, Waller A,
15 Mende DR, Kultima JR, Martin J, Kota K, Sunyaev SR, Weinstock GM, Bork P.
2013. Genomic variation landscape of the human gut microbiome. Nature
493:45-50.
26. Ross JI, Snelling AM, Carnegie E, Coates P, Cunliffe WJ, Bettoli V, Tosti G,
Katsambas A, Galvan Perez Del Pulgar JI, Rollman O, Torok L, Eady EA,
20 Cove JH. 2003. Antibiotic-resistant acne: lessons from Europe. The British
journal of dermatology 148:467-478.
27. Jackson AP, Thomas GH, Parkhill J, Thomson NR. 2009. Evolutionary
diversification of an ancient gene family (rhs) through C-terminal displacement.
BMC Genomics 10:584.

28. Horvath P, Barrangou R. 2010. CRISPR/Cas, the immune system of bacteria and archaea. *Science* 327:167-170.
29. Harris SR, Feil EJ, Holden MT, Quail MA, Nickerson EK, Chantratita N, Gardete S, Tavares A, Day N, Lindsay JA, Edgeworth JD, de Lencastre H, Parkhill J, Peacock SJ, Bentley SD. 2010. Evolution of MRSA during hospital transmission and intercontinental spread. *Science* 327:469-474.
30. Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, van der Linden M, McGee L, von Gottberg A, Song JH, Ko KS, Pichon B, Baker S, Parry CM, Lambertsen LM, Shahinas D, Pillai DR, Mitchell TJ, Dougan G, Tomasz A, Klugman KP, Parkhill J, Hanage WP, Bentley SD. 2011. Rapid pneumococcal evolution in response to clinical interventions. *Science* 331:430-434.
31. Chin CS, Sorenson J, Harris JB, Robins WP, Charles RC, Jean-Charles RR, Bullard J, Webster DR, Kasarskis A, Peluso P, Paxinos EE, Yamaichi Y, Calderwood SB, Mekalanos JJ, Schadt EE, Waldor MK. 2011. The origin of the Haitian cholera outbreak strain. *The New England journal of medicine* 364:33-42.
32. Gordon D, Abajian C, Green P. 1998. Consed: a graphical tool for sequence finishing. *Genome Res* 8:195-202.
33. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821-829.
34. Borodovsky M, McIninch J. 1993. Recognition of genes in DNA sequence with ambiguities. *Biosystems* 30:161-171.
35. Salzberg SL, Delcher AL, Kasif S, White O. 1998. Microbial gene identification using interpolated Markov models. *Nucleic Acids Res* 26:544-548.

36. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* 5:R12.
37. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731-2739.
38. Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52-57.
39. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403-410.
40. de Hoon MJ, Imoto S, Nolan J, Miyano S. 2004. Open source clustering software. *Bioinformatics* 20:1453-1454.
41. Saldanha AJ. 2004. Java Treeview--extensible visualization of microarray data. *Bioinformatics* 20:3246-3248.

EXAMPLE 3 – Microbial DNA extraction from skin samples**Skin microcomedone sampling**

Skin microcomedone (white head or black head) samples were taken from the skin of the subjects using a specialized adhesive tape. The skin was moistened with
5 water before the adhesive tape was put on. The tape was left on the skin for 15-20 minutes until it became dry. Clean gloves were used for each sampling. After being taken off from the skin, the tape was placed into a 50 mL sterile tube. This can be applied to many skin sites, such as the nose, forehead, chin, and back.

Bacterial DNA extraction

10 Microcomedones were individually picked or scraped off from the adhesive tape using sterile forceps and placed in a 2 mL sterile microcentrifuge tube filled with Buffer ATL (Qiagen) and 0.1mm diameter glass beads (BioSpec Products, Inc., Bartlesville, OK). Cells were lysed using a beadbeater for 3 minutes at 4,800 rpm at room temperature. After centrifugation at 14,000 rpm for 5 minutes, the supernatant
15 was retrieved and used for genomic DNA extraction using QIAamp DNA Micro Kit (Qiagen). The manufacturer protocol for extracting DNA from chewing gum was used. Concentration of the genomic DNA was determined by a spectrometer.

EXAMPLE 4 – Microbiome type detection**Detailed protocol for accurate detection of the skin microbiome type based on 16S rDNA****5 PCR amplification, cloning, and sequencing**

16S rDNA was amplified using primers 8F (5'-AGAGTTTGATYMTGGCTCAG-3') and 1510R (5'-TACGGYTACCTTGTTACGACTT-3'). Thermocycling conditions were as following: initial denaturation step of 5 minutes at 94°C, 30 cycles of denaturation at 94°C for 45 seconds, annealing at 52°C for 30 seconds and
10 elongation at 72°C for 90 seconds, and a final elongation step at 72°C for 20 minutes. PCR products were purified using column-based method. Subsequently, the 16S rDNA amplicons were cloned into pCR 2.1-TOPO vector (Invitrogen). One-Shot TOP-10 Chemically Competent *E. coli* cells (Invitrogen) were transformed with the vectors and plated on selective media. Individual positive colonies were picked and
15 inoculated into selective LB liquid medium. After 14 hours of incubation, the plasmids were extracted and purified, either using column-based plasmid extraction kit or traditional methods. The clones were sequenced bidirectionally using Sanger sequencing method. The microbiome type of each individual was determined based on the 16S rDNA sequence data of the top 10 major ribotypes. See Figure 5 and
20 SEQ ID NOs 1-10.

Detailed protocols for fast detection of the skin microbiome type based on PCR and qPCR

By sequencing and annotating 69 novel *P. acnes* genomes and by comparing
25 a total of 82 *P. acnes* genomes, several genomic loci which are unique to acne associated *P. acnes* strains were identified, *i.e.*, Loci 1-4. See Figure 8. The genomic sequences of Loci 1, 2, 3, and 4 from all sequences *P. acnes* strains and

their sequence similarities, which range from 95% to 100%, are listed as SEQ ID NOs 15-18, respectively.

Detection Method 1

To rapidly detect the presence or absence of RT4 and RT5 strains in patients, multiplex PCR targeting Loci 1, 2, and 3 was designed and performed on genomic DNA extracted from *P. acnes* strains and skin samples. Figure 23 shows that Loci 1, 2, and 3 were amplified from various *P. acnes* strains as predicted based on the genome data.

The PCR primer sequences are shown in Table 1:

Target locus	Forward primer	Reverse primer
Locus 1	GGTATCCACCGAGATGGAAG (SEQ ID NO:11)	GTGGTCCCAGGTGACATTCT (SEQ ID NO:12)
Locus 2	CGACATCGACGTTTCATCTG (SEQ ID NO:13)	GTGTTCTCCTCGTGCTGGTT (SEQ ID NO:14)
Locus 3	GATAATCCGTTTCGACAAGCTG (SEQ ID NO:15)	ACCCACCACGATGATGTTT (SEQ ID NO:16)
Pak	CGACGCCTCCAATAACTTCC	GTCGGCCTCCTCAGCATC

Table 1 – Primers specific for loci 1, 2, 3 and Pak (housekeeping gene)

Additional primers targeting these loci can be designed based on the genome sequences of loci 1-4 (SEQ ID NOs 15-18, respectively). Each 20µL reaction contains 12.7µL molecular grade H₂O, 2µL 10X High Fidelity Buffer, 0.6µL 50mM MgSO₄, 0.4µL 10nM dNTP, 0.8µL of each primer (final primer concentrations is 400nM), 0.1µL Platinum Taq DNA Polymerase High Fidelity (All reagents from Invitrogen) and 1µL gDNA template (approx. 40ng gDNA). The thermocycling conditions are as following: initial denaturation step of 10 minutes at 95°C; 35 cycles with each consisting of 45 seconds at 95°C, 30 seconds at 65°C and 45 seconds at 72°C; and final elongation step of 10 minutes at 72°C.

To quantitatively measure the abundance of acne associated *P. acnes* strains in skin samples, quantitative PCR (qPCR) targeting Loci 1, 2, and 3 were performed on genomic DNA extracted from *P. acnes* strains. See Figure 12. LightCycler 480 High Resolution Melting Master kit was used (Roche Diagnostics GmbH, Mannheim, Germany). Each 10 μ L reaction contains 5 μ L of Master Mix (2X concentrate), 1 μ L of 25mM MgCl₂, 0.5 μ L of 4 μ M forward and reverse primers, 1 μ L to 3.5 μ L of DNA template (approximately 2.5ng DNA), and molecular grade H₂O, up to the volume. The thermocycling conditions were as following: initial activation step of 10 minutes at 95 °C; 40 amplification cycles with each consisting of 10 seconds at 95°C, 15 seconds at 65 °C during first cycle, but with a stepwise 0.5°C decrease for each succeeding cycle and 30 seconds at 72°C; and finally, a melting curve step, starting at 65°C and ending at 99°C, with ramp rate of 0.02°C/s and acquisition rate of 25 per °C.

The protocol was tested using mock samples, where different strains of *P. acnes* were mixed in different proportions to mimic the strain population distributions in real samples. See Table 2.

Mock sample	Microbiome type	Dominant isolate	Percentage of dominant isolate	Abundance of Locus 1	Abundance of Locus 2
1A	I	HL036PA3	80	7.5%	7.5%
1B	I	HL078PA1	80	7.5%	87.5%
1C	I	HL106PA2	80	87.5%	87.5%
2	II	HL103PA1	80	8.0%	10.0%
3	III	HL087PA3	80	8.0%	10.0%
4	IV	HL038PA1	70	79.0%	82.0%
5A	V	HL072PA2	80	6.7%	8.9%
5B	V	HL096PA1	80	86.7%	88.9%
6	minor	HL110PA4	70	12.0%	15.0%
8A	minor	HL086PA1	70	80.0%	83.3%
8B	minor	HL092PA1	70	10.0%	13.3%

Table 2 – Mock samples mimicking the microbiome types observed in human skin samples

The concentration of each locus was quantified from standards derived from Locus 1, Locus 2, and Pak PCR amplicons. The copy number of each gene was quantified from genomic DNA standards that were derived from TadA (in Locus 3) and Pak amplicons using conventional PCR. See Figure 24.

5 Detection Method 2

P. acnes TaqMan qPCR Assay

Primer and Probe Design

Primers and probes for detecting Loci 1, 2, 3, and 4 in *P. acnes* strains and clinical samples were designed as listed in Table 3:

Targeted region	Primer/Probe	Primer/Probe name	Sequence (5'-3')	Amplicon size (bp)
Locus1	forward primer	Locus1_F	GAAGAATCCCGCTCCATTTC (SEQ ID NO:17)	107
	reverse primer	Locus1_R	CCTTTCTTGTAGCCGAGCAG (SEQ ID NO:18)	
	probe	Locus1_Probe	56-FAMATTGTCACC/ZEN/TGGGACCACCGTAAAC/3IABkFQ (SEQ ID NO:19)	
Locus2	forward primer	Locus2_F	CGTGATCCTGATCGACTGTG (SEQ ID NO:20)	103
	reverse primer	Locus2_R	GCTCCACAACCTCGAGTGC (SEQ ID NO:21)	
	probe	Locus2_Probe	CAGGCCGTTGATCGTGAGCTGA (SEQ ID NO:22)	
Locus3	forward primer	Locus3_F	TGCTGATAATCCGTTTCGACA (SEQ ID NO:23)	104
	reverse primer	Locus3_R	ACGACGTCGAAAACAACCTCC (SEQ ID NO:24)	
	probe	Locus3_Probe	5TET/CTCTACCGA/ZEN/AGCTCTTGCCGCAT/3IABkFQ (SEQ ID NO:25)	
Locus4	forward primer	Locus4_F	ATCGCCGTCGACAGGTAGT (SEQ ID NO:26)	103
	reverse primer	Locus4_R	CCGAGATTCTGCGCCTAGT (SEQ ID NO:27)	
	probe	Locus4_Probe	CGGTGCCCTTGCTGAGGTACA (SEQ ID NO:28)	
Pak	forward primer	Pak_F	GCAACCCGACATCCTCATTA	101
	reverse primer	Pak_R	AGTCGAAGAAGTCGCTCAGG	
	probe	Pak_Probe	VIC/CGTTCTACAGCCACCACGACGG/TAMRA	

Table 3 – Primer and probe sequences used for identification of *P. acnes* loci

A triplex Taqman qPCR was designed and tested using *Propionibacterium* specific primers to target Locus 1, Locus 3, and an internal control, Pak, present in all *P. acnes*.

Locus 1, Locus 3, and Pak Amplification

Benchtop amplification was carried out to assess specificity of designed primers and to determine optimum cycling conditions prior to multiplexing.

Amplification was carried out using a BioRad C1000 thermal cycler. Singleplex PCR

5 reactions contained 0.2µM target specific primers, 10X Platinum Taq buffer (Invitrogen), 1.0 mM MgCl₂, 0.2 mM each dNTP, 0.5U/µl Platinum Taq polymerase, 1 µl DNA template and made up to a final volume of 10 µl. Cycling was as follows: initial denaturation 94°C for 5 minutes, followed by 30 cycles of denaturation at 94°C for 30 seconds, annealing at 60°C for 30 seconds and extension at 72°C for 30
10 seconds, and one final extension cycle at 72°C for 5 minutes. Amplification products were analysed electrophoretically on a 1% agarose/TBE gel to check for correct amplification of target and cross-species reactivity with primer targets.

Taqman Triplex PCR

Triplex qPCR was carried out using an Applied Biosystems 7900HT

15 instrument. 1-2 µl of sample DNA were added to mastermix containing X2 QuantiTect Multiplex PCR Master Mix (Qiagen), 0.2 µM primers; Locus1_F, Locus1_R, Pak_F, Pak_R; 0.2 µM probes; Locus1_Probe and Pak_Probe, and 0.1 µM primers Locus3_F and Locus3_R, and 0.1 µM Locus3_Probe. The reaction mix was made up to a final volume of 20 µl with sterile PCR grade water. The PCR
20 program consisted of one cycle at 50°C for 2 minutes, followed by one cycle at 95°C for 15 minutes to allow for activation of the multiplex mastermix, then 45 cycles of 94°C for 60 seconds and 57°C for 90 seconds. Each run contained calibrators of extracted *P. acnes* DNA from culture, as well as no-template controls (NTC) and water controls. qPCR was run with a passive reference, ROX, supplied in the

Quantitect Multiplex PCR mastermix. Data were analysed using the SDS v2.4 software.

Assay Calibration, Sensitivity, and Specificity

Calibrations curves for *P. acnes* targets Locus 1 and Pak were constructed by plotting mean Ct values for a series of log dilutions of quantified genomic DNA standards extracted from *P. acnes* from pure culture. Genome equivalents were estimated. Five replicates of *P. acnes* calibrators were used to calculate mean Ct values and standard deviations. These data were used to determine sensitivity of the assay and the limits of detection (LOD). Calibration plots were used to determine the number of *P. acnes* genomes in clinical samples, with one copy of Locus1 and Pak targets per genome. DNA concentration and copy number were determined and serial ten-fold dilutions of the purified product were used as standards for construction of the Locus3 calibration plot. Strains that display possible combinations of the presence and absence of Locus1 and Locus 3 were used for Locus1 and Pak calibration: HL038PA1 (Locus1+, Locus3+), HL083PA1 (Locus1+, Locus3-), HL078PA1 (Locus1-, Locus3+) and HL063PA1 (Locus1-, Locus3-). The assay was validated using sequenced *P. acnes* strains from pure culture with known loci before being applied to clinical samples. The specificity of the assay for each target was tested using other bacterial species including skin commensals and other Propionibacteria.

Assay Validation

A total of 24 sequenced *P. acnes* strains (HL063PA1, HL078PA1, HL083PA1, HL038PA1, HL037PA1, HL082PA1, HL020PA1, HL001PA1, HL046PA2, HL043PA1, HL086PA1, HL110PA3, HL110PA4, HL007PA1, HL087PA3, HL027PA1, HL056PA1, HL067PA1, HL074PA1, HL045PA1, HL053PA1, HL005PA1, HL072PA1, HL043PA2) including possible combinations with and without Locus 1 and Locus 3 were used to validate the triplex qPCR assays. The qPCR triplex assay successfully identified Locus 1 and Locus 3 in strains previously shown by whole genome sequencing to harbor these loci.

10 Application to Clinical Samples

Genomic DNA extracted from two clinical samples, #1 and #2, were analyzed using the Taqman qPCR triplex assay. Amplification plots revealed the presence of *P. acnes* (Pak) in both samples (Figure 6). Locus1 and Locus 3 targets were also detected in both samples with a much larger percentage of *P. acnes* Locus1 positive and Locus3 positive strains present in 1µl of sample #1 compared to sample #2.

EXAMPLE 5 – Acne Vaccine

Strains with 16S rDNA ribotypes (RTs) 4, 5, 7, 8, 9, and 10 were identified as highly associated with acne. Vaccines can be raised against these strains. See T. Nakatsuji *et al.*, 128(10) J. Invest. Dermatol. 2451-2457 (Oct. 2008).

EXAMPLE 6 – Probiotic development utilizing the strains associated with healthy skin in topical creams, solutions, and the like for cosmetic and other products

5 RT6 is mostly found in healthy skin. These strains can be used as probiotics in topical products for acne prevention and treatment. Four RT6 strains, including HL110PA3, HL110PA4, HL042PA3, and HL202PA1, were isolated and sequenced.

In addition, bacterial culture supernatant and/or cell lysate, including bacterial metabolites, can be used in creams, solutions, and other cosmetic products to
10 prevent the growth of strains associated with acne. Sequences sharing at least 95% homology with SEQ ID NOs 51-54 may be used for the development of probiotics and the like.

15 **EXAMPLE 7 – Drug development targeting specific strains that are associated with acne**

Identification of the Core and Non-Core Regions of *P. acnes*

The “core” genome regions of *P. acnes* were defined as genome sequences that are present in all of the 82 genomes, while the “non-core” regions were defined
20 as genome sequences that are NOT present in all the genomes. See S. Tomida *et al.*, Pan-genome and Comparative Genome Analyses of *Propionibacterium acnes* Reveal Its Genomic Diversity in the Healthy and Diseased Human Skin Microbiome (in press); see also Example 2. Non-core regions specific to strains of RTs 4 and 5, e.g., loci 1, 2, and 3, were identified, as mentioned previously. Non-core regions
25 specific to strains of RT8 (noted as Locus 4) were also identified as well as several other strains such as HL078PA1, HL030PA2, HL063PA2, P.acn17, HL097PA1, and PRP38. See Figure 20. The genomic sequence of Locus 4 is set forth as SEQ ID NO:18. The genes in loci 1-4 below (Tables 4-1, 4-2, and 4-3) that are mostly

unique to acne associated strains RT4, RT5, and RT8 are listed below. Non-core sequences are also set forth. The genes encoded in these loci are drug targets.

Locus in Fig.3a	ID	Description
Locus 1	GM131	ABC transporter ATP-binding protein
Locus 1	GM132/GM133 ²	Site-specific recombinase
Locus 1	GM134	Site-specific recombinase
Locus 1	GM135	Hypothetical protein
Locus 1	GM136	Hypothetical protein
Locus 1	GM137	N-acetylmuramoyl-L-alanine amidase
Locus 2	GM171	Hypothetical protein
Locus 2	GM172	Hypothetical protein
Locus 2	GM173	Single-strand binding family protein
Locus 2	GM174	CobQ/CobB/MinD/ParA nucleotide binding domain protein
Locus 2	GM175	Hypothetical protein
Locus 2	GM176	Hypothetical protein
Locus 2	GM177	Hypothetical protein
Locus 2	GM178	Hypothetical protein
Locus 2	GM179	Hypothetical protein
Locus 2	GM180	Hypothetical protein
Locus 2	GM181	CAAX amino protease family protein
Locus 2	GM182	Hypothetical protein
Locus 2	GM103	YeaO-like protein
Locus 2	GM184	Hypothetical protein
Locus 2	GM185	SagB-type dehydrogenase domain protein
Locus 2	GM186	Hypothetical protein
Locus 2	GM187	ABC transporter, ATP-binding protein
Locus 2	GM188	ABC-2 type transporter
Locus 2	GM189	Hypothetical protein
Locus 2	GM196	Hypothetical protein

5

Table 4-1 – List of genes encoded in loci 1 and 2, specific to RT4 and 5

Locus	ID	Description
Locus 3	PAGK_2319	hypothetical protein
Locus 3	PAGK_2320	hypothetical protein
Locus 3	PAGK_2321	hypothetical protein
Locus 3	PAGK_2322	plasmid stabilization system protein
Locus 3	PAGK_2323	hypothetical protein
Locus 3	PAGK_2324	hypothetical protein
Locus 3	PAGK_2325	hypothetical protein
Locus 3	PAGK_2326	CobQ/CobB/MinD/ParA nucleotide binding domain
Locus 3	PAGK_2327	hypothetical protein
Locus 3	PAGK_2328	hypothetical protein
Locus 3	PAGK_2329	hypothetical protein
Locus 3	PAGK_2330	hypothetical protein
Locus 3	PAGK_2331	hypothetical protein (similar to PPA1279)
Locus 3	PAGK_2332	plasmid partition protein ParA
Locus 3	PAGK_2333	hypothetical protein
Locus 3	PAGK_2334	hypothetical protein
Locus 3	PAGK_2335	hypothetical protein
Locus 3	PAGK_2336	putative ribbon-helix-helix protein, copG family
Locus 3	PAGK_2337	putative ribonuclease E
Locus 3	PAGK_2338	hypothetical protein (similar to PPA1294)
Locus 3	PAGK_2339	hypothetical protein (similar to PPA1295)
Locus 3	PAGK_2340	putative permease
Locus 3	PAGK_2341	hypothetical protein (similar to PPA1297)
Locus 3	PAGK_2342	hypothetical protein (similar to PPA1298)
Locus 3	PAGK_2343	hypothetical protein (similar to PPA1299)
Locus 3	PAGK_2344	hypothetical protein (similar to CLOLEP_00122)
Locus 3	PAGK_2345	hypothetical protein (similar to CLOLEP_00123)
Locus 3	PAGK_2346	hypothetical protein (similar to CLOLEP_00124)
Locus 3	PAGK_2347	hypothetical protein (similar to CLOLEP_00125)
Locus 3	PAGK_2348	hypothetical protein (similar to CLOLEP_00126)
Locus 3	PAGK_2349	hypothetical protein (similar to CLOLEP_00127)
Locus 3	PAGK_2350	hypothetical protein
Locus 3	PAGK_2351	hypothetical protein (similar to CLOLEP_00129)
Locus 3	PAGK_2352	hypothetical protein (similar to CLOLEP_00130)
Locus 3	PAGK_2353	hypothetical protein (similar to CLOLEP_00131)
Locus 3	PAGK_2354	hypothetical protein (similar to CLOLEP_00132)
Locus 3	PAGK_2355	hypothetical protein (similar to CLOLEP_00134)
Locus 3	PAGK_2356	hypothetical protein (similar to CLOLEP_00135)
Locus 3	PAGK_2357	hypothetical protein (similar to CLOLEP_00141)
Locus 3	PAGK_2358	hypothetical protein (similar to CLOLEP_00142)
Locus 3	PAGK_2359	hypothetical protein (similar to CLOLEP_00143)
Locus 3	PAGK_2360	hypothetical protein (similar to CLOLEP_00144, RcpC)
Locus 3	PAGK_2361	hypothetical protein (similar to CLOLEP_00145, TadZ)
Locus 3	PAGK_2362	hypothetical protein (similar to CLOLEP_00146, TadA)
Locus 3	PAGK_2363	hypothetical protein (similar to CLOLEP_00147, TadB)
Locus 3	PAGK_2364	hypothetical protein (similar to CLOLEP_00148, TadC)
Locus 3	PAGK_2365	hypothetical protein (similar to CLOLEP_00149, Flp-1)
Locus 3	PAGK_2366	hypothetical protein (similar to CLOLEP_00151, TadE)
Locus 3	PAGK_2367	hypothetical protein (similar to CLOLEP_00152, TadE)
Locus 3	PAGK_2368	hypothetical protein (similar to CLOLEP_00153, TadE)
Locus 3	PAGK_2369	hypothetical protein (similar to CLOLEP_00154)
Locus 3	PAGK_2370	hypothetical protein (similar to CLOLEP_00157)
Locus 3	PAGK_2371	hypothetical protein (similar to CLOLEP_00158)
Locus 3	PAGK_2372	hypothetical protein (similar to CLOLEP_00159)
Locus 3	PAGK_2373	hypothetical protein (similar to CLOLEP_00160)
Locus 3	PAGK_2374	hypothetical protein
Locus 3	PAGK_2375	hypothetical protein (similar to CLOLEP_00162)
Locus 3	PAGK_2376	hypothetical protein (similar to CLOLEP_00163)
Locus 3	PAGK_2377	hypothetical protein (similar to CLOLEP_00164)
Locus 3	PAGK_2378	hypothetical protein (similar to CLOLEP_00166)
Locus 3	PAGK_2379	repA
Locus 3	PAGK_2380	CobQ/CobB/MinD/ParA nucleotide binding domain
Locus 3	PAGK_2381	hypothetical protein
Locus 3	PAGK_2382	hypothetical protein
Locus 3	PAGK_2383	Yag1E
Locus 3	PAGK_2384	hypothetical protein
Locus 3	PAGK_2385	hypothetical protein
Locus 3	PAGK_2386	hypothetical protein
Locus 3	PAGK_2387	hypothetical protein
Locus 3	PAGK_2388	hypothetical protein
Locus 3	PAGK_2389	hypothetical protein
Locus 3	PAGK_2390	hypothetical protein
Locus 3	PAGK_2391	hypothetical protein
Locus 3	PAGK_2392	ResA

Table 4-2 – List of genes encoded in Locus 3, a linear plasmid and specific to RT4 and 5

Locus	ID	Description
Locus 4	HMPREF9576_00292	tRNA adenylyltransferase
Locus 4	HMPREF9576_00293	conserved hypothetical protein
Locus 4	HMPREF9576_00294	conserved domain protein
Locus 4	HMPREF9576_00295	response regulator receiver domain protein
Locus 4	HMPREF9576_00296	histidine kinase
Locus 4	HMPREF9576_00297	hypothetical protein
Locus 4	HMPREF9576_00298	hypothetical protein
Locus 4	HMPREF9576_00299	hypothetical protein
Locus 4	HMPREF9576_00300	hypothetical protein
Locus 4	HMPREF9576_00301	hypothetical protein
Locus 4	HMPREF9576_00302	drug resistance MFS transporter, drug:H ⁺ antiporter-2 (14 Spanner) (DHA2) family protein
Locus 4	HMPREF9576_00303	hypothetical protein
Locus 4	HMPREF9576_00304	conserved domain protein
Locus 4	HMPREF9576_00305	beta-ketoacyl synthase, N-terminal domain protein
Locus 4	HMPREF9576_00306	hypothetical protein
Locus 4	HMPREF9576_00307	acetyltransferase, GNAT family
Locus 4	HMPREF9576_00308	putative (3R)-hydroxymyristoyl-ACP dehydratase
Locus 4	HMPREF9576_00309	putative acyl carrier protein
Locus 4	HMPREF9576_00310	putative 3-ketoacyl-(acyl-carrier-protein) reductase
Locus 4	HMPREF9576_00311	ornithine cyclodeaminase/mu-crystallin family protein
Locus 4	HMPREF9576_00312	pyridoxal-phosphate dependent enzyme
Locus 4	HMPREF9576_00313	lantibiotic dehydratase, C-terminus
Locus 4	HMPREF9576_00314	aminotransferase, class I/II
Locus 4	HMPREF9576_00315	acyl carrier domain protein
Locus 4	HMPREF9576_00316	AMP-binding enzyme
Locus 4	HMPREF9576_00317	malonyl CoA-acyl carrier protein transacylase family protein
Locus 4	HMPREF9576_00318	ABC-2 type transporter
Locus 4	HMPREF9576_00319	ABC transporter, ATP-binding protein
Locus 4	HMPREF9576_00320	hypothetical protein

Table 4-3 – List of genes encoded in Locus 4, RT8 specific region

EXAMPLE 8 – Targeted phage therapy

- 5 Bacteriophages play an important role in regulating the composition and dynamics of microbial communities, including the human skin microbiota. Bacteriophages of *Propionibacterium acnes*, a major skin commensal, were previously isolated and used as a typing system to distinguish different serotypes of *P. acnes*. However, molecular characterization of these phages had been lacking.
- 10 Recent efforts in genome sequencing have improved our understanding of *P. acnes* phages and their interactions with bacterial hosts.

Bacteriophages are the most abundant organisms on earth (Mc Grath & van Sinderen, 2007) and are believed to outnumber bacteria by 10:1 in many diverse ecosystems (Rohwer, 2003). As important components of microbial communities,

15 bacteriophages are a reservoir of diversity-generating elements (Rohwer & Thurber,

2009) and regulate both the abundances (Suttle, Chan, & Cottrell, 1990) and diversity of microbial hosts by predation (Rodriguez-Valera *et al.*, 2009). The human skin is inhabited by hundreds of microbial species, including bacteria, fungi, and viruses (Grice & Segre, 2011). The homeostasis of this ecosystem is important to its function as a barrier against the invasion and colonization of pathogens on the skin. However, much remains to be learned about the nature and driving forces of the dynamics among the microorganisms in the skin microbial community. In particular, the relative abundances and interactions between bacteriophages and their bacterial hosts on the skin remained to be elucidated.

The microbial community in the pilosebaceous unit of the skin is dominated by *Propionibacterium acnes*, which accounts for approximately 90% of the microbiota (“(Nature Precedings Paper),” n.d.). *P. acnes* has been suggested as a pathogenic factor in the development of acne vulgaris (Bojar & Holland, 2004; Leyden, 2001), one of the most common human skin diseases. Above-detailed studies classified *P. acnes* strains into ribotypes (RT) based on their 16S ribosomal RNA (rRNA) sequences, and demonstrated that *P. acnes* strain population structure in pilosebaceous units differs between healthy skin and acne affected skin.

P. acnes bacteriophages exist on the human skin. In 1968, Zierdt *et al.* (Zierdt, Webster, & Rude, 1968) isolated such a phage, named phage 174, from spontaneous plaques of a *P. acnes* isolate (at the time known as *Corynebacterium acnes*). Phage 174 was able to lyse nearly all *P. acnes* strains tested in the study [10]. Subsequently, more *P. acnes* phages were isolated which exhibited varied life cycles that range from lytic to temperate [11, 12]. However, in the last decades, the study of *P. acnes* bacteriophages had been limited to the development of phage

typing systems to distinguish the different serotypes of *P. acnes* [13, 14]_ENREF_4 , and extensive molecular characterization of the phages has been lacking.

Recent genomic sequencing of *P. acnes* bacteriophages (Farrar *et al.*, 2007; Lood & Collin, 2011; Marinelli *et al.*, 2012) have provided new insight into *P. acnes* phage diversity. *P. acnes* phages are similar to mycobacteriophages both morphologically and genetically, but have a much smaller genome. Currently 14 phage genome sequences are available. Sequencing additional phage isolates is needed to further characterize the diversity. Despite these recent sequencing efforts, the genome-level diversity of *P. acnes* phages in the human skin microbiome and their interactions with *P. acnes* and other Propionibacteria remain to be elucidated. *P. acnes* phages have diverse host specificities among different lineages of *P. acnes* strains [14]. Phage host specificity is important in determining how these phages regulate the composition and dynamics of *P. acnes* populations in the community. On the other hand, certain *P. acnes* strains may also influence phage populations through their anti-viral mechanisms, such as the bacterial immune system based on the transcription of clustered, regularly-interspaced short, palindromic repeat (CRISPR) sequence arrays. The CRISPR arrays contain oligonucleotide 'spacers' derived from phage DNA or plasmid DNA. In a manner analogous to RNA interference, the transcribed, single-stranded CRISPR RNA elements interact with CRISPR-associated (Cas) proteins to direct the degradation of DNA targets containing complementary 'protospacer' sequences from foreign DNA [16]. While characterizing the genome diversity of *P. acnes*, Applicants discovered that *P. acnes* strains of RT2 and RT6 harbor CRISPR arrays. The CRISPR mechanism may play a role in defending against phage or plasmid invasion.

To better understand the interactions between bacteria and bacteriophages in the human skin microbiome and their contributions to skin health and disease, the diversity and host specificity of *P. acnes* phages isolated from acne patients and healthy individuals was investigated. The genomes of 15 phage isolates were investigated and screened against a panel of 69 sequenced Propionibacteria strains to determine their host range and specificity.

Phage Isolation and General Genome Features

To characterize the genetic diversity and the abundance of *P. acnes* phages in the skin microbiome, 203 skin samples of pilosebaceous units from 179 individuals were collected, including 109 samples from normal individuals and 94 from acne patients. All of the samples were cultured for *P. acnes* under anaerobic conditions. Phage plaques in 49 samples were observed: 35 from normal individuals and 14 from acne patients. *P. acnes* phages were found more frequently in samples from normal individuals than from acne patients with statistical significance ($p=0.005$, Fisher's exact test). Among the 93 phage isolates that were obtained from these samples, five phages from acne patients and ten from normal individuals were selected for whole genome sequencing using 454 or Illumina platforms (Table 3-1).

Table 3-1 – Phage Genome Information and Sequencing Statistics

Phage Name	Genome Length (bp)	GC%	Total Input Reads	Total Input Bases	Coverage	Annotated ORFs	Presence of 11-nt Overhang
PHL111M01	29,140	54.33	5,453	2,865,116	98x	46	yes
PHL060L00	29,514	54.05	10,000	12,015,904	407x	47	yes
PHL112N00	29,266	54.48	10,000	12,270,460	419x	47	yes
PHL113M01	29,200	54.10	4,228	2,237,790	77x	45	yes
PHL114L00	29,464	54.21	10,000	12,270,609	416x	47	yes
PHL010M04	29,511	53.99	3,185	1,686,535	57x	46	not verified
PHL066M04	29,512	53.99	4,478	2,364,379	80x	46	yes
PHL073M02	29,503	53.99	4,700	2,529,669	86x	46	not verified
PHL071N05	29,467	53.92	6,143	3,059,098	104x	46	verified
PHL067M10	29,377	54.26	4,486	2,313,471	79x	46	yes
PHL115M02	29,453	53.82	8,914	4,687,391	159x	46	yes
PHL085M01	29,451	53.82	10,000	12,552,610	426x	46	yes
PHL037M02	29,443	53.78	5,895	3,093,818	105x	46	yes
PHL085N00*	29,454	53.83	20,000	3,904,239	133	45	yes
PHL082M00*	29,491	54.38	20,000	3,289,741	112	44	yes
Average	29,383	54.05	6,729	5,688,219	193.37	46	

*sequenced on the Illumina MiSeq platform. All other genomes were sequenced on the

454 platform

All phage genomes were assembled, completed, and annotated (Figure 29).

The genomes of these 15 phages have comparable sizes (29.1 - 29.5 Kb) and GC

5 content (53.8 – 54.5 %), similar to the published *P. acne* phage genomes. On average, 44 open reading frames (ORFs) were predicted in each genome.

Consistent with the genome organization previously reported [11, 15], the ORFs were arranged compactly within the left and right arm regions of each genome. The left arm and right arm of the genomes can be distinguished by their opposite

10 directions of transcription. The sequence identity between any pair of genomes is moderately high, ranging from 78.2 to 99.9% (Table 3-1).

***P. acnes* Phages are Diverse with Subgroups of Highly Related Strains with Distinct Sites of Genetic Variations**

To investigate the genome diversity of *P. acnes* phages, all 29 sequenced
5 phage genomes were compared, including Applicants' 15 phage genomes and the
14 published ones (Farrar *et al.*, 2007; Lood & Collin, 2011; Marinelli *et al.*, 2012).
The core genomic regions shared by all 29 genomes have a combined length of
24,475 bp (83% of the average genome length) and contain 6,812 single-nucleotide
polymorphisms (SNPs). A phylogenetic tree constructed from these 6,812 SNPs
10 (Figure 30) shows that most of the phage genomes isolated from all studies to date
show comparable divergence from each other with an average distance of 0.301
(substitution rate at the SNP sites). However, also found were two groups of phages,
named as Group I and Group II (Figure 30), that are closely related with much
shorter phylogenetic distances. The same results were obtained when the entire
15 genome sequences (including core and non-core regions) were used in the
calculation of phylogenetic relationships (Figure 31).

We next determined whether the newly-sequenced phages belong to the
phylogenetic groups discovered before. Lood *et al.* previously surveyed the
phylogenetic diversity of *P. acnes* phages based on the nucleotide sequences
20 encoding head proteins or amidases of phage isolates [12]. Three major
phylogenetic groups were reported. Applicants' data were combined with the data
from Lood *et al.* and Applicants reconstructed the phylogenetic trees of head protein
and amidase gene sequences. The updated phylogenetic trees reproduced the
relationships among the strains from the previous study (Figure 32). However,
25 Applicants' phages were grouped into separate clades. Moreover, by including the
gene sequences from the current study, the longest phylogenetic distance among all
studied phages was increased from 0.077 to 0.102 for the head protein gene and

from 0.140 to 0.182 for the amidase gene. Although these distances are still considerably shorter than those of the closest outgroups (0.939, head protein from mycobacteriophage Che9d; 0.764, *P. acnes* KPA171202 amidase) [12], the analysis suggests that *P. acnes* phage diversity is broader than previously described.

5 Some of the *P. acnes* phages appear to be closely related strains as previously shown [12]. Among the 29 sequenced genomes, two groups of closely related strains were observed (Figure 30). Group I consists of PHL066M04, PHL010M04, and PHL073M02, which are separated by an average phylogenetic distance of 0.002 at the genome level. Group II consists of PHL085M01,
10 PHL085N00, PHL115M02, and PHL037M02, with an average phylogenetic distance of 0.004. These two groups are statistically robust when core regions or the entire genome or only the left-arm or rightarm coding regions were used in calculating the phylogeny (Figures 30 and 31).

 Whether the genetic variations among Group I phages or Group II phages
15 were located in particular regions of the genomes was investigated. The sites of sequence variation among Group I phages lie primarily within the region encoding a putative type II holin and a peptidoglycan amidase (Gp20 and Gp21 as annotated in PA6, Figure 33). These endolysins permeabilize the membrane and degrade the extracellular peptidoglycan layer to release new phage particles from the bacterial
20 host. The majority of the sequence variations in these two genes among the Group I phage genomes are mostly synonymous and do not appear to affect the functions of the proteins. The genomes of PHL010M02 and PHL066M02 differ at only 11 sites, 9 of which occur in predicted coding regions.

Genetic variations among Group II members reside in a region encoding homologs of Gp16, Gp17, and Gp18 in PA6 (Figure 33). These proteins' location near the 3' end of the left arm between structural proteins and lysis proteins suggests that they may be late-acting genes involved in viral protein processing and packaging.

Alternative Annotations of *P. acnes* Phage Genomes

The large number of newly-sequenced *P. acnes* phage strains provided an opportunity to validate and refine initial annotations of the *P. acnes* phage genomes. Based on the analysis, several alternative annotations of the phage genomes were confirmed.

All 15 phage genomes that were sequenced support an alternative annotation of the Gp22/23 locus, which was previously annotated as two ORFs, Gp22 and Gp23, encoded on the plus-strand in the PA6 genome. Homologs of the PA6 genes Gp22 and Gp23 were not consistently identified in the genomes, as many of the homologs have inconsistent start and stop codon positions at the expected plus-strand locations of these genes. However, on the minus-strand, all genomes appear to encode a single ORF with a length of 513-522 bp. This annotation is consistent with the annotation reported by Marinelli *et al.* (Marinelli *et al.*, 2012), which is referred to herein as Gp22/23 (Figure 29). While no known function was assigned to Gp22 or Gp23 of the original plus-strand annotation, the minus-strand annotation of Gp22/23 in PHL112N00 and PHL111M01 showed modest similarity to a zinc finger protein from *Arthroderma gypseum* (E-values 1.0e-4 and 5.7e-4, respectively). The PHL111M01 annotation also showed similarity to a polyprenyl diphosphate synthase from *Streptomyces albus* (E-value 2.6e-5) and a polyketide synthase from the *Frankia* genus (E-value 2.5e-4). The minus-strand Gp22/23 ORFs from most

genomes are homologs of each other, except those in PHL067M10 and PHL114L00. The Gp22/23 ORFs in these two genomes form a separate group and share little nucleotide similarity to the other Gp22/23 ORFs despite being present at the same locus in the genome. The observation that this ORF appears in all the phage
5 genomes on the minus strand suggests that this region may be part of the right arm. This is consistent with previous reports of a plus-strand transcriptional terminator that separates Gp22/23 from the rest of the left arm in PA6, PAD20, and PAS50 [11].

Homologs of the PA6 ORFs Gp42, Gp45, and Gp46, which occur in the right-arm of the genome near the ~1 kb non-coding region, were not consistently identified.

10 The expected locations of each of these right-arm ORFs in the phage genomes frequently contained numerous stop codons and showed limited homology to corresponding regions of the PA6 genome. This is consistent with the generally high degree of nucleotide variation near the non-coding region and suggests that these ORFs may represent genes that are differentially present among different phage
15 strains.

The sequencing data demonstrated that the ends of the phage genomes are flanked by 11-nucleotide single-stranded overhangs (Table 3-1). In the sequencing data of 10 phage genomes, 1-3 reads that span both the 3' and 5' ends of the genomes were found. The genome ends in these reads are consistently separated
20 by a sequence that matches the 11-nt single-stranded extension previously reported (Marinelli *et al.*, 2012). However, based on the sequencing data, the presence of overhangs in three of the 15 genomes: PHL010M04, PHL073M02, and PHL071N05, were not shown. It is possible that they were simply not detected, as overhang-containing reads were rarely observed in general (2.3 overhang reads per 10,000
25 reads). Nevertheless, the data do suggest that the phage DNA could be circularized

at some point in their life cycle, as previously proposed [11]. The absence of the overhang sequence in reads that map to only one end of the genome may be an artifact of sample processing, as T4 DNA polymerase is used to 'polish' fragmented library DNA by digesting 3' single-strand extensions and extending the complement of 5' single-strand extensions (Roche Diagnostics, 2009). If so, it is surmisable that the overhang may exist on the 3' ends of the genome.

Host Range and Specificity of *P. acnes* Phages

To investigate the host range and specificity of *P. acnes* phages, the 15 sequenced phages were screened against a panel of 69 Propionibacterium strains, including 65 *P. acnes* strains, three *P. humerusii* strains, and one *P. granulosum* strain. Except for the *P. acnes* strains KPA171202 and ATCC11828, all of these Propionibacterium strains were isolated from the same cohort of subjects sampled for phages. The genomes of all 65 *P. acnes* strains and three *P. humerusii* strains were sequenced. A phylogenetic tree of the 65 *P. acnes* strains based on the SNPs in their core genomic regions was constructed (Figure 33, left dendrogram). Based on the previously established typing of *P. acnes* strains by their RecA gene sequences [7], the bacterial collection included all major lineages of *P. acnes* found on the human skin, with multiple strains representing each type: IA-1, IA-2, IB-1, IB-2, IB-3 and II. The susceptibility/resistance of each of the 69 bacterial strains against each of our 15 sequenced phages was determined using a crossstreak method. In total, 1,035 bacterium-phage interactions were determined. Each experiment was repeated at least five times. For the bacterial strains that showed resistance to phages, the fold changes in efficiency of plaquing (EOP) was determined relative to the *P. acnes* strain ATCC6919, which is known to be susceptible to all tested strains.

It was found that the susceptibility/resistance to phage is correlated with the *P. acnes* lineages. Five of the 69 *Propionibacterium* strains showed a 100-fold or greater increase in resistance against at least one phage. *P. acnes* strains of types IA-1, IA-2, IB-1, and IB-2 were all susceptible to all tested phages. However, two
5 strains of type IB-3 (KPA171202 and HL030PA1) were highly resistant to some of the phages (Figure 34). Type IB-3 strains encode components of a type III restriction modification system (genes PPA1611 and PPA1612 in KPA171202). This may explain their resistance to phages. KPA171202 encodes a cryptic prophage in the genome [17]. However, the sequence of the prophage is not related to any of
10 the sequenced *P. acnes* phages, therefore, the presence of the cryptic prophage is unlikely to explain the resistance to phages. Three type II strains were also highly resistant to some of the phages. This is consistent with previous observations that strains of this type were more frequently resistant to phages [14].

On the other hand, the susceptibility/resistance of *P. acnes* strains to phages
15 did not correlate with phage lineages ($r=0.1343$, $p\text{-value}=0.115$, Mantel test). Even the host ranges among closely-related phage strains in Group I or Group II are different (Figure 34). One example is PHL066M04, a Group I phage that showed little similarity to other phages in the same group, but had a similar bacterium-phage interaction pattern to the Group II phages PHL115M02 and PHL037M02. These
20 results suggest that bacterial factors may play an important role in determining the phage host range and specificity.

To determine whether these phages are specific to only *P. acnes* or if they are capable of interacting with other *Propionibacteria*, included were one *P. granulosum* strain and three *P. humerusii* strains in the bacterium-phage interaction experiment.
25 *P. granulosum* is a common skin commensal with approximately 1.1% abundance in

the pilosebaceous unit [7]. *P. humerusii* is a newly-defined species [18]. In the study cohort, *P. humerusii* is one of the major species found on the skin with an abundance of 1.9% in the pilosebaceous unit [7]. It is closely related to *P. acnes* with >98% identity in the 16S rRNA gene sequence [18]. While the *P. granulosum* strain showed strong resistance to all the phages tested, two *P. humerusii* strains, HL037PA2 and HL037PA3, were susceptible to all the phages. The third *P. humerusii* strain, HL044PA1, was lysed by ten of the 15 phages tested. This suggests that the host range of *P. acnes* phages is not limited to *P. acnes* but also includes *P. humerusii* and possibly other closely-related *Propionibacterium* species.

10 **Resistance to Bacteriophages Does not Correlate with the Presence of Matching CRISPR Spacers in *P. acnes* Strains**

Among the 65 *P. acnes* isolates, eight strains belong to RT2 and RT6 (RecA type II) and encode CRISPR/Cas genes, which function as a bacterial adaptive immune mechanism against foreign DNA. These RT2 and RT6 strains each have one to nine spacers, 33 nucleotides long, in their CRISPR arrays. In total, they encode 42 spacers, 28 of which are unique.

Whether the CRISPR/Cas mechanism can explain phage susceptibility/resistance in the RT2 and RT6 strains was investigated. Protospacers in the 15 phage genomes that match a spacer sequence from the RT2 and RT6 *P. acnes* strains were identified. Up to two mismatches were allowed in the sequence alignments. In all phages, protospacers that match the spacers in at least two *P. acnes* strains were identified (Figure 35). These protospacers are all single-copied in the phage genomes and are located primarily on the left arm (Figure 36). Their locations are generally conserved among all other phage genomes harboring the same protospacer sequences.

The susceptibility/resistance patterns of the eight RT2 and RT6 *P. acnes* strains showed little correlation with either the number of spacers in each array that had protospacer matches ($r=0.207$) or whether at least one match could be found against the CRISPR array in general ($r=0.202$). Susceptibility/resistance to phages also did not correlate with the pattern with which any specific spacer matched (maximum absolute correlation 0.051).

Phages can escape the CRISPR defense mechanism by mutating sites involved in protospacer recognition. The short nucleotide motif downstream of the protospacer, known as the protospacer-adjacent motif (PAM), is highly conserved among targets of CRISPR/Cas systems [19]. Mutations in these nucleotides have been found to disrupt CRISPR-mediated resistance despite complete complementarity in the protospacer sequence [20-22]. To determine whether the lack of correlation between bacterial susceptibility/resistance and the presence of matching spacer sequences is due to mutations within the PAM sequence, the PAMs of the nine protospacers that have exact matches to the spacer sequences encoded in HL042PA3 were examined. Six of these protospacers come from phages that HL042PA3 was resistant to, while the other three protospacers are from the phages that were able to lyse HL042PA3. Among the six protospacers, sequence conservation at several sites within their 33-nucleotide length and within the ten downstream nucleotides expected to contain the PAMs were observed (Figure 37). This suggests that these protospacer motifs are conserved and can be targeted by HL042PA3 CRISPRs. However, these same nucleotide positions are also conserved in the three protospacers from other phages (PHL113M01, PHL112N00, and PHL085M01) that were able to lyse HL042PA3 (Figure 37). Thus, the conservation of protospacer motifs including the PAMs cannot explain the lack of

correlation between bacterial susceptibility/resistance and the presence of matching spacer sequences

In summary, the data demonstrate that encoding CRISPR spacers that match against the genome of an invading phage is not sufficient for an effective defense, suggesting that transcriptional and/or translational regulation of CRISPR RNA and Cas gene expression may also be required for CRISPR-mediated resistance. Interactions between these bacteria and phages may also depend on additional phage and bacterial components involved in phage binding, entry, replication, or release.

A diverse group of *P. acnes* bacteriophages that reside on the human skin has been revealed. Most of the sequenced phages show moderately high genetic similarity with certain strains forming closely-related groups. These phages show various patterns of interaction with *P. acnes* and *P. humerusii* strains, but these patterns do not correlate with phage phylogeny. It was determined that resistance or susceptibility to phages correlated well with *P. acnes* lineages. Types IA-1, IA-2, IB-1, and IB-2 were all susceptible to all tested phages, while certain strains of types IB-3 and II were resistant to some phages. Phage resistance in type II *P. acnes* strains does not correlate with the presence of CRISPR spacers that match to phage protospacers, suggesting that additional mechanisms, such as regulation of the CRISPR/Cas system and/or other antiviral mechanisms, are needed in conferring the phage resistance.

This study suggests an important regulatory role of *P. acnes* bacteriophages in the skin microbiome. The strain-specific host ranges demonstrate the ability of these phages to regulate particular subsets of the *P. acnes* population and *P. humerusii* population. Among these subsets of *Propionibacterium* populations,

phages may also disseminate genes that potentially modify virulence, as suggested by Lood and Collin [11], or competitiveness, as it was suggested that gp22/23 encoded in some phages may be potentially involved in the production of polyketide antimicrobials. Both the selective lysis and modification of *P. acnes* strains by phages potentially regulates the relative abundances of the commensal and pathogenic strains of *P. acnes* on the skin. This delicate balance between commensals and pathogens can be especially important for skin health and disease at sites where *P. acnes* dominates. Based on the metagenomic shotgun sequencing data, it is estimated that the ratio between *P. acnes* phage and *P. acnes* in the pilosebaceous unit is 1:20 [7], which is far different from the phage:bacteria ratios estimated in environmental microbial communities, where viruses typically outnumber bacteria [23]. This suggests that the human host also plays a role in selecting and regulating the composition and diversity of the microbiome.

Materials and Methods

Propionibacteria Culture

P. acnes, *P. humerusii*, and *P. granulosum* strains were cultured under anaerobic conditions in Clostridial media (Oxoid) at 37°C for 4-6 days.

Propionibacterium cultures were used to prepare top agar overlays for phage culture on A media plates (12 g/L pancreatic digest of casein, 12 g/L Difco yeast extract, 22.2 mM D-glucose, 29.4 mM g/L potassium phosphate monobasic, 8 mM magnesium sulphate heptahydrate, 20 g/L Difco agar).

Phage Isolation and DNA Extraction

Plaques found on skin sample culture plates were isolated by puncturing the agar with a pipet tip and resuspending in 50 µL SM buffer (0.1 M sodium chloride, 8 mM magnesium sulfate heptahydrate, 1M Tris-HCl, pH 7.5, 2% gelatin, 1 mM

calcium chloride). The phage resuspension was spread onto A media plates with top agar containing *P. acnes* strain ATCC 6919. After incubation at 37° C for 2 days, phages were eluted with 8 mL SM buffer at room temperature, filtered with 0.22µM PES filter (Millipore), and stored at 4° C. Phage titers were determined by plaque assay.

Phage DNA extraction was performed using the Lambda Mini Kit (Qiagen) with the following modifications. Phage particles were precipitated in Buffer L2 by centrifugation at 20,000 g for 1 hour. Extracted DNA was eluted with Buffer QF and precipitated with isopropanol overnight at -20° C before centrifugation.

10 Phage Genome Sequencing and Annotation

Phage genomes were sequenced in multiplex using the Roche GS FLX Titanium or Illumina MiSeq platforms. De novo assembly of reads was performed with MIRA [24], and the resulting contigs were manually finished in Consed [25]. For phages covered by more than 20,000 reads, assembly was performed on a randomly-selected subset of 10,000 reads for 454 data or 20,000 reads for MiSeq data. Fully assembled phage genomes were annotated using Genemark.hmm [26] and Glimmer v3.02 [27].

Genome Sequence Alignment and Phylogenetic Tree Construction

Sequences present in all 16 phage genomes were defined as core regions of the phage genome. To identify these core regions, alignments were first generated between the PA6 genome and each of the other 15 phage genomes using Nucmer [28]. This yielded 15 sets of starting and ending coordinates describing intervals within the PA6 genome that align with any given phage genome. The core regions were then calculated for all phages by determining the overlapping intervals between all of the 15 coordinate sets. The core region sequences were concatenated for the

subsequent multiple sequence alignments. Single nucleotide polymorphisms (SNPs) on the core regions were identified by using the “show-snps” option of Nucmer with the default setting. Using MEGA5 [29], phylogenetic trees were constructed by the Neighbor Joining method on p-distances based on SNP sites. Bootstrapping was
5 based on 200 replicates.

Multiple sequence alignments of full-length phage genomes, left and right arm coding regions, head protein sequences, and amidase sequences were each generated with MAFFT [30] or Muscle [31]. Phylogenetic trees were constructed in Seaview [32] based on the BioNJ method applied to the Jukes-Cantor distances
10 between the sequences. All trees were bootstrapped for 5,000 replicates.

Determination of Variation Sites

Multiple sequence alignments of Group I and Group II phages were generated using MAFFT [30]. In each of these alignments, the positions of all mismatches and gaps (discrepancies) were recorded relative to a reference genome that was chosen
15 at random. Contiguous gaps in the reference genome were counted as a single discrepancy. The reference genome was divided into 50- nucleotide windows, and the discrepancy density of each sequence window was calculated as the total number of discrepancies it contained. Densities were plotted in Artemis [33].

To determine the single-nucleotide variations within each strain, all read data
20 for each phage, including reads not initially included in the de novo genome assembly, were mapped to their corresponding genomes using Mira. as the sites in each phage genome assembly.

Bacterial Resistance Test

The susceptibilities/resistances of *Propionibacterium* strains against 15
25 phages were determined using a modified cross-streak assay. The bacterial strains

were cultured and streaked in parallel across A media plates (5-6 isolates on each plate, ~1 cm apart, along with ATCC 6919 as a control). Approximately 5 µL of 10⁶ pfu/mL phage suspension was applied onto each streak, and then the plates were incubated at 37° C anaerobically for 2 days. At least five replicates of each cross-streak experiment were performed to determine whether the strains were susceptible or resistant judged based on lysis. The resistance of the bacterial strains was further quantified by assaying the efficiency of plaquing of the phages relative to *P. acnes* strain ATCC 6919, calculated as the following:

$$\text{Resistance} = \frac{1}{\text{Efficiency of Plaquing}} = \frac{\text{Titer of Phage Strain X on ATCC6919}}{\text{Titer of Phage Strain X on Bacterial Strain Y}}$$

A 100-fold or greater increase in efficiency of plaquing was considered to be evidence of resistance.

Phage Interaction Correlation

To determine whether genetically similar phages have similar host range and specificity, the correlation between their phylogenetic and phenotypic relationships was calculated, the latter based on results from the bacterial resistance test. Each column in the bacterial resistance table, which represents the host range of a given phage, was converted to binary form by assigning 1 to instances of resistance and 0 to instances of susceptibility. The Euclidean distance between each column was used to calculate a phenotype distance matrix between all phages. A phylogenetic distance matrix among the phage genomes was calculated using MEGA5 [29]. Using the ade4 package [34] in R, a Mantel test was performed on the phenotype and phylogenetic distance matrices to determine the correlation between the two. 10,000 permutations were performed.

CRISPR Search

CRISPR spacer sequences were identified in *P. acnes* genomes using CRISPRfinder [35]. The extracted spacer sequences were aligned against all phage sequences using BLASTn. Protospacers with up to two mismatches were identified.

5 Results

The genomes of 15 *P. acnes* phages isolated from human skin were sequenced. The phage genomes showed moderately high sequence similarity and were comparable in size and organization. Based on a comparison of the genomes, most phages diverge from each other, while some of them form closely-related
10 groups that were not described previously. When tested against a panel of 69 *Propionibacterium* strains, these phages lysed all *P. acnes* strains except some strains from type IB-3 and II. Some of the phages were also able to lyse *Propionibacterium humerusii* strains. It was found that bacterial susceptibility/resistance to phages had no significant correlation with phage
15 phylogeny or the presence of the CRISPR spacers in type II *P. acnes* strains that match the protospacers in the phage genomes.

Conclusions

With 15 new phage genomes, it was determined that the diversity of *P. acnes* phages is broader than previously described with novel groups added. The host
20 range and specificity are different among the phages, but are not correlated with the phylogeny of phage genomes. It was also found that encoding CRISPR spacers that match to phage genomes is not sufficient to confer *P. acnes* resistance to phages. This study provides new insight into the potential application of phages in treating acne and other *P. acnes* associated diseases.

References Cited in Example 8

1. Mc Grath S, van Sinderen D (eds): Bacteriophage: Genetics and Molecular Biology. Norfolk, UK: Caister Academic Press; 2007.
2. Rohwer F: Global phage diversity. Cell 2003, 113:141.
- 5 3. Rohwer F, Thurber RV: Viruses manipulate the marine environment. Nature 2009, 459:207-212.
4. Suttle CA, Chan AM, Cottrell MT: Infection of Phytoplankton by Viruses and Reduction of Primary Productivity. Nature 1990, 347:467-469.
5. Rodriguez-Valera F, Martin-Cuadrado AB, Rodriguez-Brito B, Pasic L, Thingstad
10 TF, Rohwer F, Mira A: Explaining microbial population genomics through phage predation. Nature reviews Microbiology 2009, 7:828-836.
6. Grice EA, Segre JA: The skin microbiome. Nature reviews Microbiology 2011, 9:244-253.
7. <http://precedings.nature.com/documents/5305/version/1>
- 15 8. Bojar RA, Holland KT: Acne and Propionibacterium *acnes*. Clin Dermatol 2004, 22:375-379.
9. Leyden JJ: The evolving role of Propionibacterium *acnes* in acne. Semin Cutan Med Surg 2001, 20:139-143.
10. Zierdt CH, Webster C, Rude WS: Study of the anaerobic corynebacteria. International Journal of Systematic Bacteriology 1968, 18:33-47.
- 20 11. Lood R, Collin M: Characterization and genome sequencing of two Propionibacterium *acnes* phages displaying pseudolysogeny. BMC genomics 2011, 12:198.
12. Lood R, Morgelin M, Holmberg A, Rasmussen M, Collin M: Inducible
25 Siphoviruses in superficial and deep tissue isolates of Propionibacterium

- acnes*. BMC microbiology 2008, 8:139.
13. Jong EC, Ko HL, Pulverer G: Studies on bacteriophages of *Propionibacterium acnes*. Med Microbiol Immunol 1975, 161:263-271.
14. Webster GF, Cummins CS: Use of bacteriophage typing to distinguish
- 5 *Propionibacterium acnes* types I and II. Journal of clinical microbiology 1978, 7:84-90.
15. Farrar MD, Howson KM, Bojar RA, West D, Towler JC, Parry J, Pelton K, Holland KT: Genome sequence and analysis of a *Propionibacterium acnes* bacteriophage. Journal of bacteriology 2007, 189:4161-4167.
- 10 16. Horvath P, Barrangou R: CRISPR/Cas, the immune system of bacteria and archaea. Science 2010, 327:167-170.
17. Brzuszkiewicz E, Weiner J, Wollherr A, Thurmer A, Hupeden J, Lomholt HB, Kilian M, Gottschalk G, Daniel R, Mollenkopf HJ, et al: Comparative genomics and transcriptomics of *Propionibacterium acnes*. PloS one 2011, 6:e21581.
- 15 18. Butler-Wu SM, Sengupta DJ, Kittichotirat W, Matsen FA, 3rd, Bumgarner RE: Genome sequence of a novel species, *Propionibacterium humerusii*. Journal of bacteriology 2011, 193:3678.
19. Mojica FJ, Diez-Villasenor C, Garcia-Martinez J, Almendros C: Short motif sequences determine the targets of the prokaryotic CRISPR defence system.
- 20 Microbiology 2009, 155:733-740.
20. Westra ER, van Erp PB, Kunne T, Wong SP, Staals RH, Seegers CL, Bollen S, Jore MM, Semenova E, Severinov K, et al: CRISPR immunity relies on the consecutive binding and degradation of negatively supercoiled invader DNA by Cascade and Cas3. Molecular cell 2012, 46:595-605.
- 25 21. Semenova E, Jore MM, Datsenko KA, Semenova A, Westra ER, Wanner B, van

der Oost J, Brouns SJ, Severinov K: Interference by clustered regularly interspaced short palindromic repeat (CRISPR) RNA is governed by a seed sequence. *Proceedings of the National Academy of Sciences of the United States of America* 2011, 108:10098-10103.

- 5 22. Semenova E, Nagornyykh M, Pyatnitskiy M, Artamonova, II, Severinov K: Analysis of CRISPR system function in plant pathogen *Xanthomonas oryzae*. *FEMS microbiology letters* 2009, 296:110-116.
23. Srinivasiah S, Bhavsar J, Thapar K, Liles M, Schoenfeld T, Wommack KE: Phages across the biosphere: contrasts of viruses in soil and aquatic
10 environments. *Research in microbiology* 2008, 159:349-357.
24. Chevreux B, Pfisterer T, Drescher B, Driesel AJ, Muller WE, Wetter T, Suhai S: Using the miraEST assembler for reliable and automated mRNA transcript assembly and SNP detection in sequenced ESTs. *Genome research* 2004, 14:1147-1159.
- 15 25. Gordon D, Abajian C, Green P: Consed: a graphical tool for sequence finishing. *Genome research* 1998, 8:195-202.
26. Lukashin AV, Borodovsky M: GeneMark.hmm: new solutions for gene finding. *Nucleic acids research* 1998, 26:1107-1115.
27. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL: Improved microbial
20 gene identification with GLIMMER. *Nucleic acids research* 1999, 27:4636-4641.
28. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL: Versatile and open software for comparing large genomes. *Genome biology* 2004, 5:R12.
- 25 29. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S: MEGA5:

molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular biology and evolution* 2011, 28:2731-2739.

30. Katoh K, Misawa K, Kuma K, Miyata T: MAFFT: a novel method for rapid
5 multiple sequence alignment based on fast Fourier transform. *Nucleic acids research* 2002, 30:3059-3066.

31. Edgar RC: MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research* 2004, 32:1792-1797.

32. Gouy M, Guindon S, Gascuel O: SeaView version 4: A multiplatform
10 graphical user interface for sequence alignment and phylogenetic tree building. *Molecular biology and evolution* 2010, 27:221-224.

33. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B: Artemis: sequence visualization and annotation. *Bioinformatics* 2000, 16:944-945.

15 34. Dray S, Dufour AB: The ade4 package: Implementing the duality diagram for ecologists. *Journal of Statistical Software* 2007, 22:1-20.

35. Grissa I, Vergnaud G, Pourcel C: CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic acids research* 2007, 35:W52-57.

20 (Nature Precedings Paper). (n.d.). Retrieved from

<http://precedings.nature.com/documents/5305/version/1>

Bojar, R. a, & Holland, K. T. (2004). Acne and *Propionibacterium acnes*. *Clinics in dermatology*, 22(5), 375–9. doi:10.1016/j.clindermatol.2004.03.005

Farrar, M. D., Howson, K. M., Bojar, R. a, West, D., Towler, J. C., Parry, J., Pelton,

25 K., *et al.* (2007). Genome sequence and analysis of a *Propionibacterium*

acnes bacteriophage. Journal of bacteriology , 189(11), 4161–7.

doi:10.1128/JB.00106-07

Grice, E. a, & Segre, J. a. (2011). The skin microbiome. Nature reviews.

Microbiology, 9(4), 244–53. doi:10.1038/nrmicro2537

5 Leyden, J. J. (2001). The evolving role of *Propionibacterium acnes* in acne.

Seminars in cutaneous medicine and surgery, 20(3), 139–43.

doi:10.1053/sder.2001.28207

Lood, R., & Collin, M. (2011). Characterization and genome sequencing of two

Propionibacterium acnes phages displaying pseudolysogeny. BMC genomics,

10 12(1), 198. doi:10.1186/1471-2164-12-198

Marinelli, L. J., Fitz-gibbon, S., Hayes, C., Bowman, C., Inkeles, M., Loncaric, A.,

& Russell, D. A. (2012). *Propionibacterium acnes* Bacteriophages Display

Limited Genetic Diversity and Broad Killing Activity against Bacterial Skin

Isolates. mBio, 3(5), 1–13. doi:10.1128/mBio.00279-12.Editor

15 Mc Grath, S., & van Sinderen, D. (eds). (2007). Bacteriophage: Genetics and

Molecular Biology. Norfolk, UK: Caister Academic Press.

Roche Diagnostics. (2009). GS FLX Titanium General Library Preparation Method

Manual. Mannheim: Roche Diagnostics GmbH.

Rodriguez-Valera, F., Martin-Cuadrado, A.-B., Rodriguez-Brito, B., Pasić, L.,

20 Thingstad, T. F., Rohwer, F., & Mira, A. (2009). Explaining microbial

population genomics through phage predation. Nature reviews. Microbiology,

7(11), 828–36. doi:10.1038/nrmicro2235

Rohwer, F. (2003). Global phage diversity. Cell, 113(2), 141.

Zierdt, C. H., Webster, C., & Rude, W. S. (1968). Study of the anaerobic

25 corynebacteria. International Journal of Systematic Bacteriology, 18, 33–47.

All the strains of RT4, RT5, and RT8 show sensitivity to all of the phages shown in Table 5. Therefore, acne patients may be treated with phage by using phage strains that are listed in Table 5:

5

ReeA Type	Rbotype	CESPR/Cs	PHL11M01	PHL11M01	PHL02M00	PHL09M00	PHL07M00	PHL07M05	PHL12M00	PHL07M02	PHL08M00	PHL13M02	PHL08M01	PHL14M00	PHL07M02	PHL09M04	PHL05M04
IA	HL036PA1	512	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL036PA2	532	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL036PA3	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL046PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL002PA3	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL002PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL005PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL005PA3	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL020PA1	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL027PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL100PA1	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL087PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL013PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL063PA1	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL072PA1	5	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL072PA2	5	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL066PA2	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL099PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL083PA1	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL038PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL074PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL005PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL056PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL053PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL045PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL007PA1	4	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL096PA1	5	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL043PA1	5	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL043PA2	5	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL078PA1	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL086PA1	8	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL082PA1	8	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL110PA2	8	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL053PA2	8	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL092PA1	8	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL110PA1	8	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL025PA1	1	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL030PA2	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL063PA2	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL037PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL059PA1	16	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL059PA2	16	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
IB	HL025PA2	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL005PA4	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL067PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL002PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL027PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL046PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL083PA2	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL013PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL050PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL050PA3	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL087PA1	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL087PA3	3	-	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	KPA171202	1	-	S	>10 ¹	S	>10 ¹	S	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹
	HL030PA1	1	-	>10 ¹	S	S	>10 ¹	S	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹
	HL050PA2	1	-	S	S	S	>10 ¹	S	S	S	S	S	S	S	S	S	S
II	HL060PA1	2	+	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL082PA2	2	+	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL001PA1	2	+	S	>10 ¹	S	>10 ¹	S	S	S	S	S	S	>10 ⁶	S	S	S
	HL106PA1	2	+	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	ATCC 11828	2	+	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL110PA3	6	+	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL110PA4	6	+	S	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL042PA3	6	+	S	>10 ⁶	>10 ⁶	S	>10 ⁷	>10 ⁷	S	>10 ⁷	>10 ⁷	S	>10 ⁷	>10 ⁷	>10 ⁷	>10 ⁷
	HL037PA2	-	-	N	S	S	S	S	S	S	S	S	S	S	S	S	S
	HL047PA2	-	-	N	S	S	S	S	S	S	S	S	S	S	S	S	S
-	HL044PA1	-	-	S	S	S	>10 ³	S	>10 ⁴	>10 ³	>10 ³	>10 ³	S	S	S	S	>10 ²
	HL078PG1	-	-	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹	>10 ¹

S susceptible
 10¹ fold increase in resistance

Table 5 – Host range and specificity of *P. acnes* phages

Strains in the IB-3 lineage show resistance against most of the tested phages.

Therefore, patients with those strains may not benefit as much from phage therapy.

SEQ ID NOs 55-81 include four unique genomic sequences for strains in the IB-3 lineage and for several other strains, such as IB-3-s1 (IB-3 and SK187), IB-3-s2 (IB-3 and HL025PA1), IB-3-s3 (IB-3 and HL201PA1), IB-3-s4 (IB-3 and HL201PA1). The sequence similarities range from 95% to 100%. Primers targeting these sequences can be used to estimate and predict the effectiveness of phage therapy.

Figure 26 shows the phylogenetic tree of the 32 phages including the 18 sequenced phages. There are phage strains highly similar to each other, such as the ones in Groups I and II. This suggests that the same phages can be found in different individuals and supports that a particular phage strain can be used as a common treatment agent for different individuals. SEQ ID NOs 33-50 reflect the genome sequences of the 18 sequenced phages, including the 15 phages shown in Table 5.

Potential therapeutic phage for patients with microbiome type I include:

PHL113M01, PHL111M01, PHL082M00, PHL060L00, PHL067M10, PHL071N05, PHL112N00, PHL037M02, PHL085N00, PHL115M02, PHL085M01, PHL114L00, PHL073M02, PHL010M04, and PHL066M04.

Potential therapeutic phage for patients with microbiome type I with IB-3 strain include:

PHL082M00 and PHL071N05.

Potential therapeutic phage for patients with microbiome type II include:

PHL113M01, PHL060L00, PHL112N00, and PHL085M01.

Potential therapeutic phage for patients with microbiome type III or dominant RT8 include:

PHL113M01, PHL111M01, PHL082M00, PHL060L00, PHL067M10, PHL071N05, PHL112N00, PHL037M02, PHL085N00, PHL115M02, PHL085M01, PHL114L00, PHL073M02, PHL010M04, and PHL066M04.

Potential therapeutic phage for patients with microbiome type IV include:

PHL113M01, PHL111M01, PHL082M00, PHL060L00, PHL067M10,
 PHL071N05, PHL112N00, PHL037M02, PHL085N00, PHL115M02, PHL085M01,
 PHL114L00, PHL073M02, PHL010M04, and PHL066M04.

5 Potential therapeutic phage for patients with microbiome type V include:

PHL113M01, PHL111M01, PHL082M00, PHL060L00, PHL067M10,
 PHL071N05, PHL112N00, PHL037M02, PHL085N00, PHL115M02, PHL085M01,
 PHL114L00, PHL073M02, PHL010M04, and PHL066M04.

Specific Interactions Between *Propionibacterium humerusii* and *P. acnes***10 Phages**

Some of the *P. acnes* phage strains can lyse a closely related
Propionibacterium species, *P. humerusii*, which has been hypothesized to be
 associated with infection in prostheses. *P. acnes* phage strains that can lyse *P.*
humerusii strains can be potentially used as a therapeutic agent for *P. humerusii*
 15 associated diseases.

Potential therapeutic phage for *P. humerusii* associated diseases include:

PHL113M01, PHL111M01, PHL082M00, PHL067M10, PHL071N05,
 PHL085N00, PHL085M01, PHL114L00, PHL073M02, and PHL010M04.

20

ORFs in Phage Genomes That Show Identity of 85% or Less to Their PA6 Homolog

phage_gene name	nucleotide differences	percent difference relative to PA6	PA6 ORF length
PA6_gp10			372
PHL111M01_gp9	56	0.849462366	
PHL112N00_gp9	60	0.838709677	
PHL114L00_gp10	60	0.838709677	
PHL010M04_gp11	56	0.849462366	
PHL082M00_gp10	60	0.838709677	
PA6_gp19			747
PHL114L00_gp19	120	0.83935743	
PHL010M04_gp20	138	0.815261044	
PHL066M04_gp20	138	0.815261044	
PHL073M02_gp20	138	0.815261044	
PA6_gp44			309
PHL060L00_gp47	83	0.731391586	
PHL010M04_gp43	75	0.757281553	
PHL066M04_gp43	74	0.760517799	
PHL073M02_gp43	74	0.760517799	
PHL067M10_gp43	74	0.760517799	
PHL082M00_gp43	66	0.786407767	
PA6_gp33			357
PHL115M02_gp37	54	0.848739496	
PHL085M01_gp32	54	0.848739496	
PHL037M02_gp31	54	0.848739496	
PHL085N00_gp32	54	0.848739496	
PA6_gp45			183
PHL111M01_gp44	33	0.819672131	
PA6_gp21			402
PHL112N00_gp20	71	0.823383085	
PHL010M04_gp22	85	0.788557214	
PHL066M04_gp22	84	0.791044776	
PHL073M02_gp22	78	0.805970149	
PHL071N05_gp21	61	0.848258706	
PHL067M10_gp22	63	0.843283582	
PHL115M02_gp26	65	0.838308458	
PHL085M01_gp22	65	0.838308458	
PHL037M02_gp21	65	0.838308458	
PHL085N00_gp22	65	0.838308458	
PA6_gp40			228
PHL111M01_gp41	37	0.837719298	
PHL060L00_gp42	37	0.837719298	
PHL112N00_gp39	54	0.763157895	
PHL113M01_gp43	38	0.833333333	
PHL114L00_gp40	40	0.824561404	
PHL010M04_gp40	50	0.780701754	
PHL066M04_gp40	50	0.780701754	
PHL073M02_gp40	50	0.780701754	
PHL071N05_gp39	39	0.828947368	
PHL067M10_gp40	37	0.837719298	
PHL115M02_gp45	43	0.811403509	
PHL085M01_gp40	43	0.811403509	
PHL037M02_gp39	43	0.811403509	
PHL085N00_gp40	43	0.811403509	
PHL082M00_gp39	42	0.815789474	

phage_gene name	nucleotide differences	percent difference relative to PA6	PA6 ORF length
PA6_gp22_23			504
PHL114L00_gp22	187	0.628968254	
PHL010M04_gp23	110	0.781746032	
PHL066M04_gp23	110	0.781746032	
PHL073M02_gp23	110	0.781746032	
PHL067M10_gp23	186	0.630952381	
PA6_gp29			567
PHL060L00_gp30	88	0.844797178	
PHL112N00_gp27	105	0.814814815	
PHL114L00_gp28	98	0.827160494	
PA6_gp35			471
PHL114L00_gp33	76	0.838641189	
PA6_gp41			540
PHL111M01_gp42	109	0.798148148	
PHL060L00_gp43	104	0.807407407	
PHL112N00_gp40	118	0.781481481	
PHL113M01_gp44	107	0.801851852	
PHL114L00_gp41	124	0.77037037	
PHL071N05_gp40	110	0.796296296	
PHL067M10_gp41	119	0.77962963	
PHL115M02_gp46	112	0.792592593	
PHL085M01_gp41	112	0.792592593	
PHL037M02_gp40	112	0.792592593	
PHL085N00_gp41	112	0.792592593	
PHL082M00_gp40	111	0.794444444	
PA6_gp47			180
PHL115M02_gp50	35	0.805555556	
PHL085M01_gp45	35	0.805555556	
PHL037M02_gp44	35	0.805555556	
PHL111M01_gp45	43	0.761111111	
PHL071N05_gp44.1	32	0.822222222	
PHL114L00_gp45.1	38	0.788888889	
PHL113M01_gp47	31	0.827777778	
PHL085N00_gp45	35	0.805555556	
PA6_gp24			393
PHL114L00_gp23	75	0.809160305	
PHL010M04_gp24	66	0.832061069	
PHL066M04_gp24	66	0.832061069	
PHL073M02_gp24	66	0.832061069	
PHL067M10_gp24	61	0.844783715	
PHL115M02_gp29	65	0.834605598	
PHL085M01_gp24	65	0.834605598	
PHL037M02_gp23	65	0.834605598	
PHL085N00_gp24	65	0.834605598	
PA6_gp30			564
PHL111M01_gp29	115	0.796099291	
PHL113M01_gp31	105	0.813829787	
PHL082M00_gp29	116	0.794326241	

phage_gene name	nucleotide differences	percent difference relative to PA6	PA6 ORF length
PA6_gp18			264
PHL112N00_gp17	40	0.848484848	
PHL082M00_gp18	41	0.84469697	
PA6_gp37			948
PHL112N00_gp34	168	0.82278481	
PHL114L00_gp35	163	0.828059072	
PA6_gp36			411
PHL111M01_gp36	80	0.805352798	
PHL112N00_gp33	76	0.815085158	
PHL113M01_gp38	82	0.800486618	
PHL114L00_gp34	86	0.790754258	
PHL010M04_gp35	82	0.800486618	
PHL066M04_gp35	82	0.800486618	
PHL073M02_gp35	82	0.800486618	
PHL071N05_gp34	82	0.800486618	
PHL115M02_gp40	68	0.834549878	
PHL085M01_gp35	68	0.834549878	
PHL037M02_gp34	68	0.834549878	
PHL085N00_gp35	68	0.834549878	
PHL082M00_gp34	85	0.793187348	

EXAMPLE 9 – Drug Development

Based on the foregoing, it is now known that some *P. acnes* strains are associated with acne. Therefore, at the time of diagnosis, it will be useful for dermatologists to know which strains are dominant on the skin of the patient. In order to do this, at first one needs to extract bacterial DNA from the skin sample of the patient. The method/kit to isolate bacterial DNA from the skin for downstream analysis detailed above can be implemented in practice. After bacterial DNA is extracted, the fast and accurate detection/diagnosis method/kit to identify the microbiome type of the patients, detailed above, can be implemented for diagnosis.

Once the microbiome type of the patient is diagnosed, several approaches can be used to treat the patient.

For example, if the patient has microbiome types IV or V, or is dominated by *P. acnes* RT10 strains, it is less likely antibiotic treatment would succeed, because these strains are antibiotic resistant. These patients should be treated using other therapies, such as retinoids or the methods. In the case that the patient has the virulent ribotypes, including RT4, RT5, and RT8, drugs targeting specifically to RT4, RT5, and RT8, can be used. For example, small molecules, antisense molecules, siRNA, biologics, antibodies, or combinations thereof targeting the genetic elements and biological pathways unique to the *P. acnes* strains associated with acne, detailed above, can be used.

EXAMPLE 10 – Additional Therapies

In the case that the dominant *P. acnes* strains in the patient do not harbor a set of CRISPR/Cas, additional treatment of phage therapy based on the foregoing can be used. For example, bacteriophage-based strain-specific therapy to treat acne can be employed. An alternative treatment strategy is to balance the relative abundance of the *P. acnes* strains by promoting the growth of health-associated strains. The strains associated with health can be used as probiotics. These can be topical creams, solutions, or other cosmetic products.

For prevention purposes, vaccine can be developed against virulent strains of *P. acnes*.

Longitudinal studies determine whether the microbiome types change over time and whether certain strains persist on subjects after treatment.

Inoculation experiments, inoculating virulent and healthy strains, determine whether *P. acnes* strain population changes.

Specific interactions between *P. acnes* strains and phages may be studied.

Immune responses in human cells against different strains of *P. acnes* may
5 also be measured.

The following publications are incorporated herein by reference in their entireties for all purposes, as are all other publications referenced herein and the Sequence Listing:

10 E. Grice *et al.*, 324 Science 1190-1192 (2009).

WHAT IS CLAIMED IS:

1. A method for determining whether an individual possesses acne comprising:

obtaining a skin sample from an individual;

isolating bacterial DNA from said sample;

5 amplifying 16S ribosomal DNA in said sample;

sequencing said amplified DNA products; and

typing the individual's DNA based on one or more of the ten major

ribotypes (RTs) of *P. acnes* strains, RT1-RT10 (SEQ ID NOs 1-

10),

10 wherein said typing occurs by determining whether said individual possesses one or more of RT1-RT10 and wherein said individual is diagnosed as having acne if said individual possesses RT4, RT5, RT7, RT8, RT9, or RT10.

2. The method according to Claim 1, wherein said individual is diagnosed as

15 having acne if said individual possesses RT4 (SEQ ID NO:4), RT5 (SEQ ID NO:5), or RT8 (SEQ ID NO:8).

20

25

3. A method for diagnosing different types of acne comprising:

obtaining a skin sample from a subject;
isolating bacterial DNA from said sample;
amplifying 16S ribosomal DNA in said sample;

5 sequencing said amplified DNA products; and
typing the subject's DNA based on one or more of the five major
microbiome types of *P. acnes* strains,
wherein said subject is diagnosed as having acne if said subject is typed to
microbiome IV or V.

10

4. A method for rapidly diagnosing acne comprising:

obtaining a skin sample from a subject;
isolating bacterial DNA from said sample;
using one or more primer sets to amplify said DNA; and

15 analyzing said amplified DNA for the presence of a sequence having
at least 95% homology with at least one of SEQ ID NOs 29-32
and 82-434,

wherein said subject is diagnosed as having acne if the presence of a
sequence having at least 95% homology with at least one of SEQ ID NOs 29-
20 32 and 82-434 exists.

5. The method according to Claim 4, wherein said amplified DNA is analyzed for the presence of a sequence having at least 99% homology with at least one of SEQ ID NOs 29-32 and 82-434 and wherein said subject is diagnosed as having acne if the presence of a sequence having at least 99% homology with at least one of SEQ ID NOs 29-32 and 82-434 exists.

6. The method according to Claim 4, wherein said amplified DNA is analyzed for the presence of at least one of SEQ ID NOs 29-32 and 82-434 and wherein said subject is diagnosed as having acne if the presence of at least one of SEQ ID NOs 29-32 and 82-434 exists.

7. A method for rapidly diagnosing acne comprising:

obtaining a skin sample from a subject;

isolating bacterial DNA from said sample;

using one or more primer sets to amplify said DNA;

5 using one or more probes to detect said amplified DNA; and

analyzing said probe signals for the presence of Locus 1 (at least one

sequence having at least 95% homology to at least one of SEQ

ID NOs 29 and 82-97), Locus 2 (at least one sequence having at

least 95% homology to at least one of SEQ ID NOs 30 and 98-

10 186), Locus 3 (at least one sequence having at least 95%

homology to at least one of SEQ ID NOs 31 and 187-423),

and/or Locus 4 (at least one sequence having at least 95%

homology to at least one of SEQ ID NOs 32 and 424-434),

wherein said subject is diagnosed as having acne if one or more of Loci 1-4

15 are present.

8. The method according to Claim 7, wherein said signals are analyzed for the presence of Locus 1, Locus 2, Locus 3, and/or Locus 4 based upon at least 99% homology.

20

9. The method according to Claim 7, wherein said signals are analyzed for the presence of Locus 1, Locus 2, Locus 3, and/or Locus 4 based upon 100% homology.

10. The method according to any one of Claims 4-9, wherein a primer of said primer sets is selected from the group consisting of SEQ ID NOs 11, 12, 17, and 18 (for Locus 1), SEQ ID NOs 13, 14, 20, and 21 (for Locus 2), SEQ ID NOs 15, 16, 23, and 24 (for Locus 3), and SEQ ID NOs 26 and 27 (for Locus 4).
11. The method according to any one of Claims 4-9, wherein a primer of said primer sets is selected from the group consisting of SEQ ID NOs 11, 12, 17, and 18 (for Locus 1), SEQ ID NOs 13, 14, 20, and 21 (for Locus 2), SEQ ID NOs 15, 16, 23, and 24 (for Locus 3), and SEQ ID NOs 26 and 27 (for Locus 4) and wherein said probes are SEQ ID NO:19 (for Locus 1), SEQ ID NO:22 (for Locus 2), SEQ ID NO:25 (for Locus 3), and SEQ ID NO:28 (for Locus 4).
12. A vaccine for the prevention and/or treatment of acne caused by *P. acnes* comprising a heat inactivated *P. acnes* strain, an attenuated protein of said strain, or combination thereof, wherein said strain is an RT4 strain, an RT5 strain, an RT7 strain, an RT8 strain, an RT9 strain, or an RT10 strain.
13. A vaccine for the prevention and/or treatment of acne caused by *P. acnes* comprising a heat inactivated *P. acnes* strain, an attenuated protein of said strain, or combination thereof identified to be specific to a subject based on 16S rDNA sequence analysis of the strains of *P. acnes* affecting said subject.

14. The vaccine according to Claim 12 or Claim 13, wherein said heat inactivated *P. acnes* strain, attenuated protein, or combination thereof is specific for at least one of unique genomic loci, regions, or sequences identified for the strains of *P. acnes*.

5

15. The vaccine according to Claim 12 or Claim 13, wherein said heat inactivated *P. acnes* strain, attenuated protein, or combination thereof is specific for at least one of Locus 1 (SEQ ID NOs 29 and 82-97), Locus 2 (SEQ ID NOs 30 and 98-186), Locus 3 (31 and 187-423), and Locus 4 (32 and 424-434).

10

16. A method for the personalized treatment of acne comprising determining the strains of *P. acnes* affecting a subject and treating said subject with an active ingredient directed to at least one detected strain of *P. acnes*, wherein the active ingredient comprises a drug targeting specific strains of *P. acnes*,
15 wherein the targeting drug comprises small molecules, antisense molecules, siRNA, biologics, antibodies, and combinations thereof targeting genomic elements specific for strains of *P. acnes* associated with acne.

15

17. A method for treating acne comprising:

20

administering an effective amount of a probiotic that comprises at least one strain of *P. acnes* that is associated with healthy or normal skin based on its 16S rDNA.

18. The method according to Claim 17, wherein said strain is an RT6 strain.

25

19. The method according to Claim 17, wherein said strain has at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.
- 5 20. The method according to Claim 17, wherein said strain has at least 99% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.
21. The method according to Claim 17, wherein said strain has 100% homology
10 to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.
22. A method for treating acne comprising:
administering an effective amount of a metabolite produced by a strain
of *P. acnes* that is associated with healthy or normal skin,
15 wherein said metabolite is selected from the group comprising bacterial culture supernatant, cell lysate, proteins, nucleic acids, lipids, and other bacterial molecules.
23. The method according to Claim 22, wherein said strain is an RT6 strain.
20
24. The method according to Claim 22, wherein said strain has at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.

25. The method according to Claim 22, wherein said strain has at least 99% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.

5 26. The method according to Claim 22, wherein said strain has 100% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.

27. A method for treating acne in a subject comprising:
administering an effective amount of a drug specifically targeting RT4,
10 RT5, RT7, RT8, RT9, or RT10, when said subject is determined to possess RT4, RT5, RT7, RT8, RT9, or RT10, respectively.

28. The method according to Claim 27, further comprising:
performing the method according to Claim 1, Claim 2, Claim 4, or
15 Claim 7 prior to administration of said drug.

29. The method according to Claim 27 or Claim 28, wherein said drug is a small molecule, antisense molecule, siRNA, biologic, antibody, or combination thereof.

20 30. A composition comprising at least one strain of *P. acnes* that is associated with healthy or normal skin.

31. The composition according to Claim 30, wherein said strain is an RT6 strain.

32. The composition according to Claim 30, wherein said strain has at least 95% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.
- 5 33. The composition according to Claim 30, wherein said strain has at least 99% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.
34. The composition according to Claim 30, wherein said strain has 100% homology to SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, or SEQ ID NO:54.
- 10
35. A method for diagnosing IB-3-based acne comprising:
- obtaining a skin sample from a subject;
- 15 isolating bacterial DNA from said sample;
- using one or more primer sets to amplify said DNA; and
- analyzing said amplified DNA for the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 55-81, wherein said subject is diagnosed as having IB-3-based acne if the presence of a sequence having at least 95% homology with at least one of SEQ ID NOs 55-81 exists.
- 20
36. A method for the personalized treatment of acne comprising determining the strain(s) of acne affecting a subject and administering to said subject an effective amount of at least one phage specifically directed to said strain(s).
- 25

37. The method according to Claim 36, wherein said subject is treated with phage directed against an RT4 strain, an RT5 strain, an RT7 strain, and RT8 strain, an RT9 strain, and/or an RT10 strain.
- 5 38. A method for treating an individual suffering from acne of microbiome type I comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:40), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).
- 10
- 15 39. A method for treating an individual suffering from acne of microbiome type I with IB-3 strain comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL082M00 (SEQ ID NO:47) and PHL071N05 (SEQ ID NO:41).
- 20 40. A method for treating an individual suffering from acne of microbiome type II comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL060L00 (SEQ ID NO:34), PHL112N00 (SEQ ID NO:35), and PHL085M01 (SEQ ID NO:44).

41. A method for treating an individual suffering from acne of microbiome type III or dominant RT8 comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

42. A method for treating an individual suffering from acne of microbiome type IV comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47), PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

43. A method for treating an individual suffering from acne of microbiome type V comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47),
5 PHL060L00 (SEQ ID NO:34), PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL112N00 (SEQ ID NO:35), PHL037M02 (SEQ ID NO:45), PHL085N00 (SEQ ID NO:46), PHL115M02 (SEQ ID NO:43), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), PHL010M04 (SEQ ID NO:38), and PHL066M04 (SEQ ID NO:39).

44. A method for treating a *Propionibacterium humerusii*-associated malady comprising administering to said individual an effective amount of a phage, wherein said phage is selected from the group consisting of: PHL113M01 (SEQ ID NO:36), PHL111M01 (SEQ ID NO:33), PHL082M00 (SEQ ID NO:47),
15 PHL067M10 (SEQ ID NO:42), PHL071N05 (SEQ ID NO:41), PHL085N00 (SEQ ID NO:46), PHL085M01 (SEQ ID NO:44), PHL114L00 (SEQ ID NO:37), PHL073M02 (SEQ ID NO:40), and PHL010M04 (SEQ ID NO:38).

45. A kit for diagnosing acne in a subject, wherein said kit comprises:
20 at least one primer selected from the group comprising SEQ ID NOs 11-18, 20, 21, 23, 24, 26, and 27; and instructions for use.

46. A kit for diagnosing acne in a subject, wherein said kit comprises:

at least one primer selected from the group comprising SEQ ID NOs

11-18, 20, 21, 23, 24, 26, and 27;

5 at least one probe selected from the group comprising SEQ ID NOs 19,

22, 25, and 28; and

instructions for use.