19. ~~Use of a~~A compound according to any one of Claims 1 to 17 or a pharmaceutically acceptable salt thereof for use in the treatment of a disorder that is selected from schizophrenia, cognitive impairment, attention deficit hyperactivity disorder (ADHD), impulsivity, compulsive gambling, overeating, autism spectrum disorder, mild cognitive impairment (MCI), age-related cognitive decline, dementia, restless leg syndrome (RLS), Parkinson's disease, Huntington's chorea, anxiety, depression, major depressive disorder (MDD), treatment-resistant depression (TRD), bipolar disorder, chronic apathy, anhedonia, chronic fatigue, post-traumatic stress disorder, seasonal affective disorder, social anxiety disorder, post-partum depression, serotonin syndrome, substance abuse and drug dependence, drug abuse relapse, Tourette's syndrome, tardive dyskinesia, drowsiness, excessive daytime sleepiness, cachexia, inattention, sexual dysfunction, migraine, systemic lupus erythematosus (SLE), hyperglycemia, atherosclerosis, dislipidemia, obesity, diabetes, sepsis, post-ischemic tubular necrosis, renal failure, hyponatremia, resistant edema, narcolepsy, hypertension, congestive heart failure, postoperative ocular hypotonia, sleep disorders, and pain.

20. The ~~use~~compound according to claim 19, where the disorder is selected from schizophrenia, cognitive impairment, mild cognitive impairment (MCI), age-related cognitive decline, dementia, and Parkinson's disease.