

CLAIMS

What is claimed is:

1. A method for providing a clotting factor VIII dosing regimen comprising:
collecting two blood samples from a patient after an infusion of clotting factor VIII;
determining a clotting factor VIII clearance for the patient based on the two blood samples;
determining if a patient has a half-life greater than a predetermined threshold;
determining, via a processor, an estimated pharmacokinetic profile of a patient using a Bayesian model of pharmacokinetic profiles of sampled patients, the estimated pharmacokinetic profile based upon at least one of a body weight or an age of the patient such that:
a first weighting factor is applied to the Bayesian model of pharmacokinetic profiles of sampled patients if the half-life of the patient is greater than the predetermined threshold, and
a second weighting factor, less than the first weighting factor, is applied to the Bayesian model of pharmacokinetic profiles of sampled patients if the half-life of the patient is less than the predetermined threshold;
determining, via the processor, a dosing regimen for a specified dosing interval including (i) a dosage and (ii) an estimated clotting factor VIII clearance for the patient over a time period based at least upon the estimated pharmacokinetic profile; and
displaying the dosing regimen on a client device.
2. The method of Claim 1, further comprising adjusting, via the processor, the estimated pharmacokinetic profile of the patient upon previous treatments of the patient.
3. The method of Claim 1, wherein the specified dosing interval is 48 hours or 72 hours.
4. The method of Claim 1, wherein the minimum threshold level is less than 20%.

5. The method of Claim 1, wherein the dosage is determined such that the estimated clotting factor VIII clearance in the patient over the time period does not fall below the minimum threshold level.

6. The method of Claim 1, wherein the estimated clotting factor VIII clearance in the patient is based upon at least one of a minimum threshold level, the dosage, or the specified dosing interval.

7. The method of Claim 1, wherein the Bayesian model includes a two-compartment model having a first compartment corresponding to a time to metabolize the clotting factor VIII and a second compartment corresponding to a dose for achieving a certain amount of the clotting factor VIII within the patient.

8. An apparatus for providing a therapeutic plasma protein dosing regimen to a client device, the apparatus comprising:

a model generator configured to create a Bayesian model of pharmacokinetic profiles of sampled patients, the Bayesian model including a (i) therapeutic plasma protein clearance and (ii) a volume of distribution relationship for a therapeutic plasma protein based upon at least one of patient age or body weight; and

a pharmacokinetic server configured to:

determine an approximate pharmacokinetic profile of a patient based upon the Bayesian model, a half-life of the therapeutic plasma protein within the patient, and at least one of an age of the patient or a weight of the patient;

determine the therapeutic plasma protein dosing regimen including a dosage and a therapeutic plasma protein level over a time period based upon the approximate pharmacokinetic profile of the patient;

modify the therapeutic plasma protein dosing regimen in response to receiving a dosing interval for applying the dosage to the patient; and

transmit the modified therapeutic plasma protein dosing regimen to the client device.

9. The apparatus of Claim 8, wherein the dosing interval is a two-day dosing interval, and wherein the pharmacokinetic server is configured to further modify the therapeutic plasma protein dosing regimen in response to receiving a three-day dosing interval in place of the two-day dosing interval.

10. The apparatus of Claim 8, wherein the pharmacokinetic server is configured to transmit a drug dosing tool to the client device, the drug dosing tool being configured to determine the therapeutic plasma protein dosing regimen and the modified therapeutic plasma protein dosing regimen.

11. The apparatus of Claim 8, wherein the pharmacokinetic server is further configured to modify the therapeutic plasma protein dosing regimen based on daily activities of the patient.

12. The apparatus of Claim 8, wherein the pharmacokinetic server is further configured to transmit the modified therapeutic plasma protein dosing regimen to an infusion pump for administering the therapeutic plasma protein to the patient.

13. The apparatus of Claim 8, wherein the approximate pharmacokinetic profile is a first approximate pharmacokinetic profile determined for a first therapeutic plasma protein treatment of the patient, and wherein the pharmacokinetic server is further configured to determine a second approximate pharmacokinetic profile for the patient for a second therapeutic plasma protein treatment of the patient based on the modified therapeutic plasma protein dosing regimen.

14. The apparatus of Claim 8, wherein the volume of distribution relationship for the therapeutic plasma protein is a relationship for at least one of clotting factor VIII and modified forms of clotting factor VIII.

15. A machine-accessible device having instructions stored thereon that are configured, when executed, to cause a machine to at least:

- prompt a user to enter at least one of a patient weight or age;

- use a Bayesian model of pharmacokinetic profiles of sampled patients to determine an approximate pharmacokinetic profile of a patient based upon the Bayesian model, a half-life of the therapeutic plasma protein within the patient, and the at least one of entered patient weight or age, the Bayesian model including (i) a therapeutic plasma protein clearance and (ii) a volume of distribution relationship for a therapeutic plasma protein based upon the at least one of entered patient weight or age;

- determine a dosing regimen for the patient based upon the approximate pharmacokinetic profile of the patient, the dosing regimen including a dosage and a dosage interval;

- modify the dosing regimen in response to receiving another dosing interval for applying the dosage to the patient; and

- enable the dosing regimen and a time-varying therapeutic plasma protein level based on the dosing regimen to be displayed to a user.

16. The machine-accessible device of Claim 15, further comprising instructions stored thereon that are configured when executed to cause the machine to:

- determine a first dosing regimen for a two-day dosing interval;

- determine a second dosing regimen for a three-day dosing interval; and

- enable the display of the first dosing regimen in conjunction with the second dosing regimen.

17. The machine-accessible device of Claim 15, further comprising instructions stored thereon that are configured when executed to cause the machine to display a graphical representation of a time-varying amount of the therapeutic plasma protein within the patient, including at least one indication of a dose of the therapeutic plasma protein being provided to the patient.

18. The machine-accessible device of Claim 15, further comprising instructions stored thereon that are configured when executed to cause the machine to display a graphical feature than enables a user to change at least one of:

- (i) a minimum concentration threshold;
- (ii) the dosage interval; or
- (iii) the dosage of the therapeutic plasma protein.

19. The machine-accessible device of Claim 18, further comprising instructions stored thereon that are configured when executed to cause the machine to modify the dosing regimen in response to receiving a change of any one of the items (i), (ii), or (iii).

20. The machine-accessible device of Claim 19, further comprising instructions stored thereon that are configured when executed to cause the machine to display a graphical representation of a change in the amount of the therapeutic plasma protein within the patient over time based on the change of any one of the items (i), (ii), or (iii).

21. The machine-accessible device of Claim 19, further comprising instructions stored thereon that are configured when executed to cause the machine to receive a minimum concentration threshold and to display an amount of time the therapeutic plasma protein level is below the minimum concentration threshold.

22. The machine-accessible device of Claim 15, further comprising instructions stored thereon that are configured when executed to cause the machine to:

receive patient measurement blood data laboratory including a concentration of the therapeutic plasma protein within the patient after a time from when the therapeutic plasma protein was administered to the patient; and

modify the approximate pharmacokinetic profile based on the patient measurement blood laboratory data.

23. A drug dosing tool, the tool comprising:
- interface circuits configured to receive two blood samples collected from a patient after an infusion of clotting factor VIII;
 - one or more processors coupled to memory, the one or more processors configured to:
 - determine a first elimination rate constant of clotting factor VIII for the patient based on the two blood samples;
 - determine a second elimination rate constant using a Bayesian model of pharmacokinetic (PK) profiles of sampled patients each of which are stored in the memory, wherein the second elimination rate is associated an elimination rate of the clotting factor VIII of a population defined by the sampled patients, the population having a similar age and body weight of the patient;
 - determining a relationship between the first elimination rate constant and the second elimination rate constant and updating the Bayesian model based on the relationship;
 - determine an estimated PK profile of the patient using the updated Bayesian model;
 - determine a dosing regimen for a specified dosing interval including (i) a dosage and (ii) an estimated clotting factor VIII clearance for the patient over a time period based at least upon the estimated pharmacokinetic profile; and
 - control an infusion pump for administration of the clotting factor VIII based on the estimated pharmacokinetic profile.