

Title: Newcastle disease virus infectious clones, vaccines and diagnostic assays.

The invention relates to Newcastle disease virus infections of poultry.

Newcastle disease virus (NDV) is one of the most diverse and deadly avian pathogens. The almost simultaneous
5 occurrence of Newcastle disease as an apparent new disease in several different geographical locations and the great variation in type and severity of the disease has caused some problems with nomenclature.

The disease has been termed pseudo fowl pest, pseudo
10 poultry plague, avian pest, avian distemper and avian pneumoencephalitis. The importance of the disease is primarily due to the development of the poultry industry during the 20th Century into a highly efficient international industry which is dependent on intensive trade between
15 countries.

It is generally assumed that the first outbreaks of Newcastle disease occurred in 1926 in Java, Indonesia, and in Newcastle-upon-Tyne, England (Kraneveld, 1926; Doyle, 1927). The name Newcastle disease was coined by Doyle as a temporary
20 name to avoid a descriptive name that might be confused with other diseases. It later became clear that other less severe diseases were caused by viruses indistinguishable from NDV. In the US a relatively mild respiratory disease was termed avian pneumoencephalitis and was shown to be caused by NDV
25 (Beach, 1944). Within a few years, numerous NDV isolations that caused extremely mild or no disease in chickens were made around the world.

The following methods have been implicated in the spread of the disease: 1) movement of live birds, feral birds, game
30 birds, racing pigeons and commercial poultry; 2) movement of people and equipment; 3) movement of poultry products; 4) airborne spread; 5) contaminated poultry feed; 6) contaminated water; 7) incompletely inactivated or

heterogeneous vaccines. According to the OIE, Newcastle disease is a disease of poultry caused by a virus of avian paramyxovirus serotype 1 (APMV-1) which has an intracerebral pathogenicity index (ICPI) in day-old chicks of 0.7 or greater. Virulent virus can also be confirmed by the presence of multiple basic amino acids at the C-terminus of the F2 protein and F (phenylalanine) at residue 117, the N-terminus of the F1 protein. Failure to demonstrate this amino acid sequence would require characterisation by ICPI tests. The word 'poultry' refers to domestic fowl, turkeys, guinea fowl, ducks, geese, quails, pigeons, pheasants, partridges and ratites that are reared or kept in captivity for breeding, the production of meat or eggs for consumption, or for restocking supplies of game.

According to Alexander (1988) three panzootics of Newcastle disease have occurred since the first recognition of the disease. The first represented the initial outbreaks of the disease and appears to have arisen in South-East Asia. Isolated outbreaks, such as the one in England in 1926, were chance introductions ahead of the mainstream which slowly moved through Asia to Europe.

A second panzootic appears to have begun in the Middle East in the late 1960's and reached most countries by 1973. The more rapid spread of the second panzootic was probably caused by the major revolution of the poultry industry with considerable international trade.

A third panzootic primarily affected domesticated birds such as pigeons and doves (Vindevogel and Duchatel, 1988). The disease apparently arose in the Middle East in the late 1970's. By 1981 it had reached Europe and then spread rapidly to all parts of the world, largely as a result of contact between birds at races and shows and the international trade in such birds.

Nowadays, Newcastle disease is still widespread in many countries of Asia, Africa, the Americas, and Europe. Only the countries of Oceania appear to be relatively free from the disease (Spradbrow, 1988).

NDV belongs to the order *Mononegavirales*, family *Paramyxoviridae*, subfamily *Paramyxovirinae*, genus *Rubulavirus*. Apart from NDV, generally called avian paramyxovirus type-1, eight other serotypes, designated avian
5 paramyxovirus type-2 to -9, can be distinguished on the basis of their antigenic relatedness in hemagglutination-inhibition tests and serum neutralisation tests (Alexander, 1993).

Despite the consistency of the serological grouping there are some cross-relationships between viruses of the
10 different serotypes.

The genome of NDV is a single-stranded RNA molecule of negative polarity, complementary to the messenger RNA's which code for the virus proteins. The RNA genome is approximately 15,200 nt in size and codes for the following gene products
15 (listed from the 3' end to the 5' end of the genomic RNA): nucleocapsid protein (NP), phosphoprotein (P), matrix protein (M), fusion protein (F), hemagglutinin-neuraminidase (HN), and large polymerase protein (L) (Chambers et al., 1986).

The RNA is complexed with the NP, P and L proteins to
20 form a ribonucleocapsid particle (RNP) that is surrounded by an envelope that is lined at the inside by the M protein. The envelope contains the F and HN proteins which are involved in attachment and penetration of the host cell.

Replication of NDV is similar to the strategy used by
25 other paramyxovirinae. The initial step is attachment of the virus to the host cell receptors, mediated by the HN protein. Fusion of the viral envelope with the host cell membrane is dependent on the action of both the HN and F proteins and results in the release of the RNP into the cytoplasm where
30 virus replication takes place.

The viral RNA-dependent RNA polymerase (which is part of the RNP) produces complementary transcripts that act as mRNA's and are used by the cell's translation machinery for the synthesis of virus proteins. Due to the accumulation of
35 NP protein, the RNA polymerase complex switches from

transcription to replication, resulting in the synthesis of full-length genomic and antigenomic RNA molecules.

Newly formed RNP's are encapsidated at the cellular membrane by the action of the M protein and the F and HN proteins which have accumulated in the cellular plasma membrane. Newly formed virus particles are released from the infected cell by a budding mechanism. For more detailed information about NDV replication see Peeples (1988). For a recent review of the molecular biology of paramyxovirinae see Lamb and Kolakofsky (1996).

Apart from commercial domestic poultry (i.e. chickens, turkeys, pheasants, guinea fowl, ducks, geese, pigeons), a wide range of captive, semi-domestic and free-living birds, including migratory waterfowl, is susceptible to NDV and can be primary infection sources (Kaleta and Baldauf, 1988).

The pathogenicity of NDV strains differs greatly with the host. The most resistant species appear to be aquatic birds while the most susceptible are gregarious birds forming temporary or permanent flocks. Chickens are highly susceptible but ducks and geese may be infected and show few or no clinical signs, even with strains which are lethal for chickens.

Newcastle Disease is complicated in that different isolates and strains of the virus may induce enormous variation in the severity of the disease. Beard and Hanson (1984) grouped NDV strains and isolates into different pathotypes that relate to disease signs that may be seen in fully susceptible chickens: 1) viscerotropic velogenic NDV, which produces acute lethal infections in which hemorrhagic lesions are prominent in the gut; and neurotropic velogenic NDV, which produces high mortality preceded by respiratory and neurological signs, but no gut lesions; 2) mesogenic NDV, which produces low mortality, acute respiratory disease and nervous signs in some birds; 3) lentogenic NDV, which produces mild or inapparent respiratory infections or even asymptomatic enteric NDV, avirulent viruses that appear to replicate primarily in the intestinal tract. Some overlap

between the signs associated with the different groups has been reported.

The virus enters the body via the respiratory and the intestinal tract or via the eye. In the trachea, the virus is spread by ciliary action and by cell-to-cell spread. After initial multiplication at the introduction site, virus is carried during episodes of viraemia to spleen, liver, kidney and lungs. Viruses of some strains reach vital organs like liver and kidney very rapidly so that the birds may die before disease symptoms are overt.

Most viruses reach the central nervous system via the blood before significant amounts of antibody exist. A long, asymptomatic carrier state presumed to occur in psittacines constitutes a potential threat to the poultry industry. A long term carrier state of both lentogenic and velogenic virus may also exist in chickens (Heuschele and Easterday, 1970).

During the replication of NDV it is necessary for the precursor glycoprotein Fo to be cleaved to F1 and F2 for the progeny virus to be infectious (Rott and Klenk, 1988). This posttranslational cleavage is mediated by host cell proteases. If cleavage fails to take place, non-infectious virus particles are produced and viral replication cannot proceed. The Fo protein of virulent viruses can be cleaved by a wide range of proteases, but Fo proteins in viruses of low virulence are restricted in their sensitivity and these viruses can only grow *in vivo* in certain host cell types and in general cannot be grown *in vitro*.

Lentogenic viruses only replicate in areas with trypsin-like enzymes such as the respiratory and intestinal tract, whereas virulent viruses can replicate in a range of tissues and organs resulting in fatal systemic infection.

Amino acid sequencing of the Fo precursor has shown that low-virulence viruses have a single arginine (R) that links the F2 and F1 chains, whereas virulent strains possess additional basic amino acids forming two pairs such as

K/R-X-K/R-R-F at the site of cleavage. Furthermore, the F2 chain of virulent strains generally starts with a phenylalanine residue whereas that of nonvirulent strains generally starts with a leucine.

For a few strains of NDV the HN protein is also produced as a precursor that requires cleavage to be biologically active (Garten et al., 1980; Millar et al., 1988).

Besides cleavability of the F and HN proteins, other viral factors may contribute to pathogenicity. Madansky and Bratt (1978, 1981a, 1981b) have shown that alterations in transcription and translation could modulate growth and cell-to-cell spread of the virus and/or cytopathogenicity.

The initial immune response to infection with NDV is cell mediated and may be detectable as early as 2-3 days after infection with live vaccine strains. This presumably explains the early protection against challenge that has been recorded in vaccinated birds before a measurable antibody response is seen (Gough and Alexander, 1973).

At about 1 week after infection, circulating antibodies may protect the host from re-infection. In the early phase IgM is involved, followed by IgG. Titres and protection peak after about 3 weeks and gradually decline if there is no boosting. This means that for older birds, re-vaccinations are necessary.

Only live vaccines administered by the respiratory route, stimulate antibody in all mucosal surfaces as well as in serum. Inactivated vaccine, even when applied via the intramuscular route, does not elicit local resistance in the respiratory tract, despite high concentrations of serum antibody.

This stresses the importance of live vaccines capable of presenting viral antigen to the upper respiratory tract to induce both local and systemic immunity. Small droplets penetrate into the lower respiratory tract thereby provoking

a mainly humoral immune response, while coarse droplets stimulate local immunity in the upper respiratory tract.

Therefor, aerosols with a wide range of droplet sizes generate the best overall local and humoral immunity.

It should be noted, however, that despite intensive vaccination with current vaccines creating high levels of antibody titers, virus can still be recovered from mucous surfaces.

The identification of Newcastle disease in the USA led to the use of inactivated vaccines (Hofstad, 1953). The observation that some of the enzootic viruses produced only mild disease resulted first in the development of the mesogenic live vaccine Roakin (Beaudette et al., 1949) and, subsequently, in the development of the milder Hitchner B1 (Hitchner and Johnson, 1948) and LaSota (Goldhaft, 1980) strains, which are now the most widely used live vaccines.

NDV live vaccines can be divided into two groups, lentogenic and mesogenic. Mesogenic strains are suitable only for secondary vaccination of birds due to their greater virulence. The immune response increases as the pathogenicity of the live vaccine increases. Therefore, to obtain the desired level of protection without serious reaction, currently vaccination programs are used that involve sequential use of progressively more virulent vaccines, or live vaccines followed by inactivated vaccines.

One of the main advantages of live vaccines is that they may be administered by inexpensive mass application techniques. A common method of application is via drinking water. However, drinking water application must be carefully monitored as the virus may be inactivated by excessive heat and light and by virucidal impurities in the water.

Mass application of live vaccines by sprays and aerosols is also very popular due to the ease with which large numbers of birds can be vaccinated in a short time. It is important to achieve the correct particle size by controlling the conditions under which the particles are generated.

Currently used live vaccines have several disadvantages. The vaccine may still cause disease signs, depending upon environmental conditions and the presence of complicating infections. Therefore, it is important to use extremely mild virus for primary vaccination and, as a result, multiple vaccinations are usually needed. Furthermore, maternally derived antibodies may prevent successful primary vaccination with lentogenic live vaccines.

Inactivated vaccines are usually produced from infectious allantoic fluid which is treated with formalin or betapropiolactone to kill the virus and mixed with a suitable adjuvant. Inactivated vaccines are administered by injection, either intramuscularly or subcutaneously. Inactivated vaccines are expensive to produce and to apply.

However, inactivated oil-emulsion vaccines are not as adversely affected by maternal immunity as live vaccines and they can be used in day-old chicks. Advantages of inactivated vaccines are the low level of adverse reactions in vaccinated birds, the high level of protective antibodies, and the long duration of protection. None of the above vaccines can serologically be differentiated from wild-type NDV.

The development of recombinant viral vaccines has been of interest to the poultry industry for a number of years. The concept is to insert genes of critical immunising epitopes of a disease agent of interest into a nonessential gene of a vector virus. Vaccination with the recombinant virus thus results in immunisation against both the vector virus as well as the disease agent of interest.

Several types of viruses have been evaluated as potential live viral vaccines for poultry. Two avian viruses that have received most attention are fowlpox virus (FPV) and herpesvirus of turkeys (HVT). Fowlpox virus is a DNA virus that has a large genome and hence is considered to have ample room to carry foreign genes.

When attenuated, FPV does not cause clinical disease and is commonly used as a vaccine in chickens. HVT is also a DNA

virus and is classified as serotype III of the Marek's disease virus (MDV) family. HVT is non-pathogenic for chickens yet cross-protective against MDV and is commonly used to vaccinate chickens against Marek's disease.

5 It has been shown that protection against Newcastle disease can be induced by using recombinant HVT or FPV vaccines (Morgan et al., 1992, 1993; Heckert et al., 1996; Boursnell et al., 1990; Taylor et al., 1990).

10 However, the onset of protection against Newcastle disease following vaccination with such recombinant vaccines that express either the NDV F protein or both the F and HN proteins was severely delayed compared to that following vaccination with a conventional live or inactivated NDV vaccine, possibly because the recombinant vaccines do not
15 provide a wide enough immunological spectre of antigenically relevant NDV epitopes other than those found on the NDV protein that is expressed by the recombinant vaccine or are not properly presented to the immune system.

20 Furthermore, local (mucosal, respiratory or enteric) protection was not effectively induced in birds vaccinated with the recombinants. This is a serious drawback since vaccines used for primary vaccination against respiratory diseases must induce local immunity to prevent infection and spread of virulent viruses that infect chickens reared under
25 field conditions.

Antibodies against NDV which are capable of protecting the host can be measured in virus neutralisation tests. However, since the neutralisation response appears to parallel the haemagglutination inhibition (HI) response, the
30 latter test is frequently used to assess the protective response, especially after vaccination.

Antibodies against both the F and HN proteins can neutralise NDV. However, antibodies against the F protein appear to induce greater neutralisation than those directed
35 against HN in *in vivo* and *in vitro* tests (Meulemans et al., 1986).

The presence of specific antibodies to NDV in the serum of a bird gives little information on the infecting strain of NDV and therefore has limited diagnostic value.

5 The omnipresence of lentogenic NDV strains in birds in most countries and the almost universal use of live vaccines that can not be distinguished, at least not serologically from wild-type NDV, means that the mere demonstration of infection is rarely adequate cause for control measures to be
10 imposed. Since field disease may be an unreliable measure of the true virulence of the virus, it is necessary to further characterise the virus that is found.

At present, the only method of Newcastle disease diagnosis which allows characterisation of the infecting
15 strain, is virus isolation followed by pathogenicity testing. At present, three *in vivo* tests are used for this purpose: 1) mean death time (MDT) in eggs; 2) Intracerebral pathogenicity index (ICPI) in one-day-old chickens; 3) Intravenous pathogenicity index (IVPI) in 6-week-old birds.

20 These tests suffer from a number of drawbacks, such as the availability of animals, poor reproducibility, and the relatively long duration of the tests. Last but not least, these tests do not allow a simple serological identification of poultry vaccinated with a vaccine or infected with a wild-
25 type strain.

As an alternative to *in vivo* tests, the polymerase chain reaction (PCR) has been successfully used to distinguish between virulent and non-virulent i and non-virulent isolates (Staüber et al., 1995; Kant et al., 1997), however, again
30 serological differentiation is not possible.

The raising of poultry and trade of their products is now organised on a international basis, frequently under management of multinational companies. The threat of Newcastle disease has proven a great restraint on such trade.

35 Successful control of Newcastle disease will only be approached when all countries report outbreaks. However,

international agreements are not simple due to enormous variation in the extent of disease surveillance in different countries. Some countries do not vaccinate and would not want any form of NDV introduced in domestic poultry because
5 vaccinated poultry cannot be distinguished from those infected with wild-type NDV.

Others only allow the use of specific live vaccines and consider other vaccines as unacceptably virulent. Yet other countries have the continued presence of circulating highly
10 virulent virus, which is not recognised as such because overt disease is masked by vaccination.

In many countries legislation exists to control Newcastle disease outbreaks that may occur. National control measures are directed at prevention of introduction and
15 spread. Most countries have restrictions on trade in poultry products, eggs, and live poultry. Most countries have established quarantine procedures for importation, especially for psittacine birds.

Some countries have adopted eradication policies with
20 compulsory slaughter of infected birds, their contacts, and products. Others require prophylactic vaccination of birds even in the absence of outbreaks, while some have a policy of ring vaccination around outbreaks to establish a buffer zone.

Clearly, there is a need for better vaccines and for
25 better diagnostic methods which can be used to control Newcastle disease. Due both to large differences in the dose that is received by individual birds during mass application of live vaccines and to variation in levels of maternal immunity in young chickens, post-vaccination reactions with
30 live vaccines are inevitable. This is one of the main concerns of farmers in countries where vaccination is compulsory.

Furthermore, many vaccines are mixtures of sub-populations. When cloned, these sub-populations may differ
35 significantly from each other in immunogenicity and pathogenicity (Hanson, 1988).

However, the largest drawback of currently used live vaccines and inactivated vaccines is the fact that vaccinated animals cannot be distinguished from infected animals with currently used screening techniques such as

5 heamagglutination-inhibition or virus neutralisation tests.

Virulent field-virus may still spread in vaccinated flocks since disease symptoms are masked by vaccination. Since virus isolation and characterisation of virulence by *in vivo* techniques is not feasible on a large scale, there is a great need for new and effective attenuated live vaccines which can be serologically discriminated from field-viruses.

10

Such vaccines, called NDV marker vaccines (and accompanying diagnostic methods and kits) , which should provide the fullest possible immunological spectre of antigenically relevant NDV epitopes, and yet should be serologically distinct from wild-type NDV are not yet available.

15

20 The invention provides a method to modify an avian-paramyxovirus genome by genetic modification, provides genetically modified avian-paramyxovirus and an avian-paramyxovirus marker vaccine.

The advent of modern molecular biological techniques has allowed the genetic modification of many RNA viruses, including negative-strand RNA viruses. This technique is often referred to as "reverse genetics". One first provides a (full-length) cDNA copy of the viral RNA, after which one transcribes this DNA in susceptible cells to produce

25

30 infectious RNA which can again replicate to produce infectious virus particles.

In general, by previous modification of the cDNA with standard molecular biological techniques, it is possible to obtain a genetically modified RNA virus. However, this has never materialised for NDV or other avian-paramyxoviruses, it has even not yet been possible to generate minigenome

35

fragments or plasmids of avian-paramyxovirus genomic fragments to study replicative events of avian-paramyxovirus, thereby creating an understanding on how to construct infectious copy virus.

5 Surprisingly, although in this description it has now been fully established that the genome of avian-paramyxovirus is the smallest of all paramyxovirus genomes sequenced up to now, especially the 5' terminal end sequence of the NDV genome is much longer than previously had been established
10 and was expected by comparison with other *Paramyxoviridae*. The invention now for the first time provides a full sequence of an avian-paramyxovirus genome and provides full-length or minigenomic length cDNA of such a virus.

The invention herewith provides avian-paramyxovirus cDNA
15 at least comprising a nucleic acid sequence corresponding to the 5'-terminal end of the genome of avian-paramyxovirus allowing generating an infectious copy of avian-paramyxovirus, said cDNA preferably comprising a full-length cDNA. However, the invention also provides cDNA at least
20 comprising a nucleic acid sequence corresponding to the 5'-terminal end of the genome of avian-paramyxovirus thereby allowing generating an replicating avian-paramyxovirus minigenome. Such minigenomes can advantageously be used to transcribe RNA and/or express protein from modified nucleic
25 acid sequences. The invention provides a cDNA according to the invention at least partly derived from Newcastle Disease Virus, for example wherein said Newcastle Disease Virus is a , lentogenic virus, preferably derived from a vaccine strain, such as LaSota strain ATCC VR-699.

30 The invention furthermore provides a cDNA according to the invention additionally provided with a modification, such as a deletion, insertion, mutation, reversion, or otherwise in a nucleic acid. For example a cDNA is provided wherein said modification comprises a nucleic acid encoding a
35 modified protease cleavage site, for example wherein said

cleavage site is a protease cleavage site of the fusion (F) protein.

In yet another embodiment, the invention provides a cDNA according to the invention wherein said modification
5 comprises a nucleic acid encoding a hybrid viral protein, such as a hybrid hemagglutinin-neuraminidase (HN) protein as described in the experimental part of the invention. The invention also provides a cDNA according to the invention wherein said modification comprises a deletion in a nucleic
10 acid encoding a viral protein, such as a matrix (M) protein.

The invention additionally provides a cDNA according to the invention additionally provided with a nucleic acid encoding an heterologous antigen, preferably wherein said antigen is derived from a poultry pathogen, as for example
15 described below. An RNA, and protein derived thereof, obtained from a cDNA according to the invention is also provided.

In recent years, a number of non-segmented negative-strand RNA viruses has been fully characterised and
20 fundamental work on the replication and expression of their genomes has culminated in the ability to generate infectious virus entirely by transfecting cells with cloned cDNA of said virus (reviewed by Conzelmann, 1996).

To date, infectious virus of non-segmented negative-strand RNA viruses has been generated from cloned cDNA of for
25 example rabies virus (Schnell et al., 1994, Conzelmann; EP0702085A1). (Schnell et al., 1994; EP0702085A1), vesicular stomatitis virus (Lawson et al., 1995; Whelan et al., 1995), Sendai virus (Garcin et al., 1995), measles virus (Radecke et
30 al., 1995; Schneider et al., 1997; EP0780475A1), human respiratory syncytial virus (Collins et al., 1995), rinderpest virus (Baron and Barrett, 1997), and human parainfluenza virus type 3 (Hoffman and Banerjee, 1997, Conzelmann; P0702085A1), (Schnell et al., 1994; EP0702085A1).

35 However, all of above infectious copy viruses are capable of growing both in vivo as well as in vitro in hosts,

tissues or cells of various origin, allowing easy cDNA transfection and replication and generation of infectious virus particles on a suitable cell line.

Such possibility does not exist for NDV, certainly not for lentogenic NDV strains which can provide a vaccine. Virulence of such an NDV strain is associated with its ability to replicate in a wide range of cells, reflected by the fact that virulent strains can easily replicate *in vitro* and *in vivo*, whereas vaccine strains can only replicate *in vivo*.

Thus, with NDV a catch 22 situation is apparent. While attempts to generate an infectious copy virus from for example infectious cDNA may possibly result in infectious virus, such virus is in general not suitable for use as a vaccine because the thus generated infectious virus is by default too virulent to be used as vaccine; the fact that it can be generated and replicated after transfection of cDNA on a cell line reflects its easy cleavability of the Fo protein into F1 and F2, as discussed above a hallmark of virulence of a NDV.

Using a vaccine strain as parent material for the cDNA would not solve this problem; a vaccine strain, especially of a lentogenic type does not contain an easily cleavable Fo protein, rendering it impossible for first generation virus to continue to replicate. The cell used for transfection will simply not be susceptible to support one or more rounds of replication of vaccine-type virus with a non-cleaved Fo protein.

The invention now elegantly provides a solution for this problem, and therewith provides infectious copy NDV, for example for use in a vaccine.

The invention provides a method to generate infectious copy Newcastle Disease Virus comprising transfecting cells, capable of expressing viral NP, P and L proteins for complexing with viral RNA with cloned full-length or genomic-length cDNA of said virus and further comprising incubating

said cells in growth medium comprising proteolytic activity allowing cleavage of the Fo protein of said virus.

In our system, co-transfection of a plasmid expressing NP could be omitted. NP is probably expressed from the full length cDNA because the NP gene is the first gene after the 5' end of the antigenomic RNA. Since eukaryotic mRNA's are usually monocistronic, expression of distal genes is not expected. However it is possible to generate full-length cDNA in which the relative positions of the NDV genes are changed. If the first gene of such a cDNA is the P or L gene, it is not necessary to express the corresponding gene product from a co-transfected plasmid.

As an alternative to using full-length cDNA, it is possible to use two or more subgenomic cDNA's which generate replication competent subgenomic RNA's and which together express the full complement of avian-paramixovirus proteins. Even if the RNA's are packaged separately, the resulting virus-like particles can be used for successive rounds of replication by means of co-infection and complementation of gene functions.

In a preferred embodiment, the invention provides a method wherein said proteolytic activity is derived of an enzyme, such as a trypsin-like enzyme, or is derived of a composition comprising said proteolytic activity. In a much preferred embodiment, said growth medium comprises allantoic fluid comprising proteolytic activity. Cleavage of the Fo protein is required for the generation of infectious virus. It is possible to generate infectious virus from lentogenic strain without the addition of exogenous proteolytic activity. By inoculating the supernatant of transfected cells into the allantoic cavity of embryonated eggs, the proteolytic activity which is present in the allantoic fluid is able to cleave the FO protein to generate the fusion-competent F1-F2 complex. Virions with such an activated F protein are able to infect susceptible cells and replication in cells which express the desired proteolytic activity yields infectious progeny. As an alternative to providing the

desired proteolytic activity to the supernatant of transfected cells, it is for example possible to use a cell that is permissive for NDV and which already expresses said proteolytic activity. Such a cell line is used to produce
5 infectious lentogenic NDV without the addition of exogenous proteolytic activity. Such a cell line can also be generated by stable transfecting a cell line with a gene that specifies said activity. Furthermore, it is possible to generate a stable transfected cell line that expresses the wild-type F
10 protein in the virus envelope, thereby providing infectious particles (themselves not provided with genomic information encoding wild-type F protein) with means to enter a cell. Rescue of infectious lentogenic virus is also possible by infection of transfected cells with an NDV helpervirus. An
15 essential requirement for such a helpervirus would be that it can be selected against, for instance by means of neutralizing antibodies which eliminate the helpervirus but which do not react with the lentogenic virus.

Finally, one may construct a stably transfected cell line
20 that expresses one, two, or all of the three essential NDV proteins, NP, P, and L. Such cell lines require the co-expression of only a subset of the three essential proteins or no co-expression at all for supporting generating infectious copy virus.

25 In a preferred embodiment, the invention provides a method wherein said cells used for transfecting are derived of chicken primary or secondary cells or cell-lines. The description provides for example CER or CEF cells, which, as most in vitro cells in general, lack the appropriate
30 proteases which are required to cleave the Fo protein of NDV, for example of strain LaSota. However, cells derived from for example other birds can also be used.

The invention further provides a method to generate infectious copy Newcastle Disease Virus comprising
35 transfecting cells with cloned full-length or genomic-length cDNA of said virus as for example identified in figure 3 and

further comprising incubating said cells in growth medium comprising proteolytic activity allowing cleavage of the Fo protein of said virus, further comprising recovering infectious virus by culturing said cells and inoculating
5 material derived from said cultured cells into the allantoic cavity of embryonated eggs. Said material for example comprises (harvested or freeze-thawed) cells or cell debris or supernatant derived from said cell culture.

For example, the description describes a method to
10 recover infectious virus, wherein the supernatant of transfected CEF monolayers was inoculated into the allantoic cavity of embryonated eggs. Four days later the allantoic fluid was harvested, analyzed in a haemagglutination assay, and passaged further in eggs.

15 In addition, the invention provides a method further comprising passaging said infectious copy Newcastle Disease Virus by harvesting allantoic fluid and re-inoculating embryonated eggs.

In a preferred embodiment of the invention, a method is
20 provided wherein said virus is a lentogenic virus, for example derived from an avirulent field-case of NDV or from a vaccine strain of NDV, such as the LaSota strain of NDV. Furthermore, a method is provided to modify an avian-paramyxovirus genome by means of genetic modification which
25 allows the introduction of one or more mutations, deletions, and/or insertions or other modifications. For example, method is provided to attenuate or modify the virulence of avian-paramyxovirus by modifying cDNA, for example encoding a viral protein, such as the V protein, and cloning said modified
30 cDNA into full-length cDNA and generating infectious copy virus from said full-length cDNA, thereby generating new NDV strains or new attenuated live vaccines with improved properties.

Apart from attenuation by modification of gene products
35 it is also possible to attenuate avian-paramyxovirus by modification of nucleotide sequences which are involved in transcription and/or replication. Such modifications result

in attenuated strains which express wild type like F proteins which are cleavable both *in vitro* and *in vivo* in a wide range of cells and as a result are more immunogenic than the classical vaccine strains.

5 In a preferred embodiment, the invention provides a method to attenuate or modify the virulence of an avian paramyxovirus such as a Newcastle Disease Virus, comprising modifying a protease cleavage site of a viral protein by modifying cDNA encoding said cleavage site, further
10 comprising cloning said cDNA into genomic length cDNA of e.g. Newcastle disease virus and generating infectious copy Newcastle Disease virus. Said cleavage site is for example a protease cleavage site in the F or HN protein of Newcastle Disease Virus. Attenuation is in general restricted to
15 reduction of virulence, however, it is now also possible to use a relatively a-virulent strain of NDV and provide the progeny of such a strain with increased virulence, for example by providing it with an increased tendency to replicate in a specified cell-type. It is now thus possible
20 to assign distinct virulence attributes to NDV.

The invention provides a method to antigenically modify avian paramyxovirus such as a Newcastle Disease Virus, comprising modifying cDNA encoding at least a part of a viral protein harbouring at least one immunodominant epitope,
25 further comprising cloning said cDNA into genomic length cDNA of Newcastle disease virus and generating infectious copy Newcastle Disease virus.

For example, the invention provides a method to (further) modify NDV, for example using a method to produce
30 an infectious copy of NDV (vaccine) which has been provided, a method to produce a recombinant marker NDV vaccine is provided, a marker vaccine that contains the fullest possible or needed immunological spectrum of antigenically relevant NDV epitopes, and yet is serologically distinct from wild-type NDV because a distinct, serologically relevant epitope
35 or marker has been removed by recombinant techniques. The

invention provides a method to modify the antigenic make-up of avian paramyxovirus such as NDV, thus allowing the generation of e.g a live NDV marker vaccine which can be serologically distinguished from avian paramyxovirus field strains.

In one embodiment, the invention provides infectious copy NDV wherein the HN protein of NDV has been modified by recombining cDNA encoding a part of said HN protein with cDNA encoding a part of HN protein derived from an avian-paramyxovirus, for example type 2 or type 4. Said hybrid HN protein serves as a serological marker for the infectious copy NDV strain thus obtained or can serve to change the tropism of the avian paramyxovirus to other cells and/or tissues. These, so called, marker strains as provided by the invention allow the generation of vaccines which are an invaluable tool to assess the prevalence of NDV in commercial flocks around the world. Furthermore, the large-scale application of such marker vaccines will lead to the complete eradication of NDV by a process of intensive screening and stamping out of infected flocks.

Furthermore, a method is provided to generate an infectious copy NDV strain which expresses one or more antigens from other pathogens and which can be used to vaccinate against multiple diseases. Such an infectious copy NDV virus for example comprises a heterologous cDNA encoding a heterologous protein obtained from for example Avian Influenza (AI) (Haemagglutinin (H5 and H7) and Neuraminidase), Avian leukosis virus (ALV) (env protein (gp85)), Chicken anemia virus (CAV) (VP1+VP2), Marek's disease virus (MDV) (glycoprotein B (gB), gH), Infectious laryngotracheitis virus (ILT) (gB, gH, gD), Infectious bursal disease virus (IBDV) (VP2 and VP3), Turkey rhinotracheitis virus (TRT) (fusion (F) protein), Avian paramyxovirus-2,-3,-6 (PMV) (F-protein, Haemagglutinin neuraminidase (HN), or others, Infectious bronchitis virus (IBV) (peplomer protein, nucleoprotein), Reoviruses (sigma protein), Adenoviruses Pneumoviruses, Salmonella enteritidis, Campylobacter

jejuni, Escherichia coli, Bordetella avium (formerly
Alcaligenes faecalis), Haemophilus paragallinarum, Pasteurella
multocida, Ornithobacterium rhinotracheale, Riemerella
(formerly Pasteurella) anatipestifer, Mycoplasmata (M.
5 gallisepticum, M. synoviae, M. mereagris, M. iowae), or
Aspergilli (A. flavus, A. fumigatus).

The invention herewith provides avian-paramyxovirus or
strains derived thereof which can be used as a vaccine vector
for the expression of antigens from other poultry pathogens.

- 10 Several properties make NDV an ideal vaccine vector for
vaccination against respiratory or intestinal diseases. 1)
NDV can be easily cultured to very high titres in embryonated
eggs. 2) Mass culture of NDV in embryonated eggs is
relatively cheap. 3) NDV vaccines are relatively stable and
15 can be simply administered by mass application methods such
as by drinking water or by spraying or aerosol formation. 4)
The natural route of infection of NDV is by the respiratory
and/or intestinal tract which are also the major natural
routes of infection of many other poultry pathogens. 5) NDV
20 can induce local immunity despite the presence of circulating
maternal antibody.

It has been shown that NDV has potent antineoplastic, as
well as immune-stimulating properties (for a review see
Schirrmacher et al., 1998) [Schirrmacher, V., Ahlert, T.,
25 Steiner, H.-H., Herold-Mende, C., Gerhards, R. and Hagmüller
E. (1998) Immunization with virus-modified tumor cells.
Seminars in Oncology 25: 677-696]. Although NDV does not seem
to be able to replicate productively in normal human cells, a
selective NDV-mediated killing of human cancer cells was
30 noted. After local NDV therapy, viral oncolysis and complete
remissions of human tumor xenografts were observed in nude
mice. This has led to the use of NDV for tumor therapy.
However, a problem is that such application may be restricted
to local treatment.

35 NDV infection induces interferons, chemokines, and other
potentially important gene products, and introduces

pleiotropic immune-stimulatory properties into tumor cells. This concept has been used for the production of autologous tumor cell vaccines consisting of fresh operative specimens that have been infected with NDV. This type of vaccine is
5 called autologous tumor vaccine-NDV or ATV-NDV (Schirrmacher et al., 1998). The NDV-infected cells are inactivated by gamma-irradiation which prevents cell division but which still allows replication of NDV in the cytoplasm of infected cells. After inoculation of patients with ATV-NDV, T-cells
10 are recruited through NDV-induced chemokines. Some of these T-cells may express a T-cell receptor that can interact with peptides from tumor-associated antigens in complex with major histocompatibility complex class I molecules at the cell surface. This interaction results in the induction of a
15 cytotoxic T-cell response which results in specific killing of autologous tumor cells.

The invention provides that the repertoire and amount of chemokines and immune stimulatory proteins induced by NDV infection are modulated. The present invention provides a
20 method for generating recombinant NDV that has been modified to incorporate and express (a) heterologous gene(s). Such recombinant NDV may be used to modify the repertoire and amount of immune-stimulatory proteins in infected cells. In one embodiment, the invention provides a recombinant NDV that
25 incorporates and expresses genes encoding human interferons, chemokines or other immune stimulatory proteins. Said recombinant NDV is used for the production of ATV-NDV which is more potent than conventional ATV-NDV. For example: cytokines IFN- α , - β , TNF- α , IL-1, IL-6; chemokines RANTES,
30 IP-10; other genes such as HSP, ACTH, endorphin, iNOS, EPA/TIMP, NF κ B.) The pleiotropic immune-stimulatory properties of NDV may also be used as an adjuvant for vaccination of animals and humans against infectious diseases. In one embodiment of the invention, foreign genes
35 encoding (a) relevant antigen(s) of (an) infectious agent(s) are introduced in the NDV genome and the simultaneous expression of the antigen(s) and the immune-stimulatory

proteins by infected cells may induce a potent immune response against the infectious agent. In another embodiment of the invention, the immune-stimulating properties of NDV may be further enhanced by using NDV recombinants that simultaneously express antigens and specific immune-stimulatory proteins. In a preferred embodiment, the invention is used to generate an AIDS (acquired immune-deficiency syndrome) vaccine by using NDV recombinants that express relevant antigens of human immune-deficiency virus (HIV), either alone or in combination with immune-stimulatory proteins.

NDV are also used as an adjuvant for vaccination of animals and humans against infectious diseases. In one embodiment of the invention, heterologous or foreign genes encoding (a) relevant antigen(s) of (an) infectious agent(s) are introduced in the NDV genome and the simultaneous expression of the antigen(s) and the immune-stimulatory proteins by infected cells may induce a potent immune response against the infectious agent. In another embodiment of the invention, the immune-stimulating properties of NDV are further enhanced by using NDV recombinants that simultaneously express antigens and specific immune-stimulatory proteins. In a preferred embodiment, the invention is used to generate an AIDS (acquired immune-deficiency syndrome) vaccine by using NDV recombinants that express relevant antigens of human immune-deficiency virus (HIV), either alone or in combination with immune-stimulatory proteins.

Also, a method is provided to generate a conditional lethal NDV deletion mutant which can be used as self-restricted non-transmissible (carrier)vaccine. An NDV deletion mutant was generated which is unable to express the matrix (M) protein which is involved in budding of NDV at the inner cell membrane. The invention provides for example a phenotypically complemented NDV strain that is unable to express the M protein which is still able to infect cells and spread by means of cell-to-cell transmission. However, the

mutant virus is unable to generate infectious progeny on non-complementing cells. This shows that phenotypically complemented NDV deletion mutants can be used as safe self-restricted vaccines which are unable to spread into the environment. Such a non-transmissible vaccine combines the most important advantage of live vaccines, i.e., efficacy, with the most important advantage of killed vaccines, i.e., safety.

The invention provides Newcastle Disease Virus, or strains derived thereof, for example by passaging or further cultivation in embryonated eggs or appropriate cells, that is derived from infectious copy virus obtainable by a method provided by the invention.

For example, NDV is provided that has been modified in at least one way to generate infectious copy Newcastle Disease Virus which is attenuated, modified in virulence, antigenically modified, expressing a heterologous antigen or are non-transmissible, or combinations thereof.

Herewith the invention provides NDV vaccines, characterised for example by carrying distinct virulence attributes or distinct antigenic characteristics, be it for marker vaccine purposes and/or for expressing heterologous antigens derived from other pathogens, be it in transmissible and/or non-transmissible form.

Such a vaccine can be a killed or a live vaccine. Preferably, such a vaccine is a live vaccine, however, killed vaccines as provided by the invention are beneficial under those circumstances where a live vaccine is not or only little applicable, for example because of trade restrictions or other conditions set by disease controlling authorities.

The invention herewith also provides a diagnostic method, and corresponding test kit, to detect antibodies against said serologically relevant immunodominant epitope or marker, therewith providing methods and means to execute a method for control and/or eradication of NDV and/or other poultry diseases in poultry. The invention provides new and effective vaccines which can be serologically discriminated

from field-viruses and old-type vaccines. Such new vaccines, called NDV marker vaccines, provide the fullest possible immunological spectrum of antigenically relevant NDV epitopes, and yet are serologically distinct from wild-type NDV by applying accompanying diagnostic methods and kits.

The invention provides a method for distinguishing unvaccinated animals or animals vaccinated with a NDV vaccine according to the invention from animals infected with wild-type NDV or vaccinated with an unmodified mesogenic or lentogenic NDV-vaccine strain comprising taking a least one sample (such as serum, blood, eggs or eye fluid) from said animal and determining in said sample the presence of antibodies directed against an immunodominant epitope or marker expressed by said wild-type or unmodified NDV but not by a vaccine according to the invention.

The invention provides a method wherein said antibodies are directed against the HN or F protein of NDV, for example a hybrid protein as described in the experimental part as this description. The invention provides for example a diagnostic method wherein said animal is selected from the group composed of poultry, preferably of chickens.

The invention also provides a diagnostic kit for use in a method to serologically distinguish between animals. In one embodiment of the invention, a simple and rapid haemagglutination-inhibition (HI) test is used to distinguish between vaccinated animals and infected animals. Animals vaccinated with a marker vaccine in which the complete globular head of HN of NDV has been replaced with the corresponding part of HN of another serotype will not induce antibodies to HN of NDV and therefore will not inhibit haemagglutination of erythrocytes by NDV virions.

By using marker vaccine virions in the HI test, antibodies against the hybrid HN protein is detected and may be used as a measure for the efficacy of vaccination. As an alternative, an ELISA that detects antibodies against the F protein of NDV is used to measure the efficacy of vaccination.

Apart from the HI test, an ELISA can be used to determine the presence of antibodies against HN of NDV. The antigen to be used in such a test is for example HN of NDV that is expressed by recombinant DNA techniques or a conserved peptide from HN of NDV.

5 A blocking ELISA may also be used. In this case one or more monoclonal antibodies against conserved epitopes of HN of NDV are used to determine whether competing antibodies are present in samples from vaccinated animals. The ELISA tests
10 can advantageously be used if the marker vaccine contains a chimeric HN protein only or when a few epitopes of HN of NDV are replaced.

The invention is further explained in the experimental part of this description without limiting the invention
15 thereto.

EXPERIMENTAL PART

20 MATERIALS AND METHODS

Standard cloning procedures were carried out according to Sambrook et al. (1989) unless stated otherwise. All constructions involving DNA fragments which were generated by
25 means of the polymerase chain reaction (PCR) were verified by sequence analysis. In the primer sequences given below, the underlined nucleotides correspond to NDV sequences and the position within the NDV genome is indicated. The nucleotide sequence of restriction sites which were used for cloning are
30 indicated in boldface.

Cells and viruses

CER cells (Smith et al., 1976) were grown in GMEM/EMEM
35 (1:1) containing 5% foetal calf serum and 2% of an antibiotic mix that contained 1000 U/ml Penicillin, 1000 µg/ml Streptomycin, 20 µg/ml Fungizone, 500 µg/ml Polymixin B, and

10 mg/ml Kanamycin. QT35 cells (Moscovici et al., 1977; Cho, 1982) were grown in medium supplied by GibcoBRL/Life Technologies (cat.no. 041-91536; proprietary composition of Fort Dodge) supplemented with 5% FCS and 2% antibiotic mix.

- 5 QM5 cells (Antin and Ordahl, 1991) were grown in M199 medium supplemented with 10 % tryptose phosphate broth, 10% FCS and 2% antibiotic mix.

- NDV strain LaSota was obtained from ATCC (ATCC VR-699) and was passaged two times in embryonated eggs. Before we
10 started with the construction and cloning of cDNA, the virus was plaque purified by three rounds of plaque purification on primary chicken embryo fibroblasts (CEF). To this end the virus was titrated on CEF cells cultured in GMEM/EMEM (1:1) containing 5% foetal calf serum, 2% antibiotic mix, 5%
15 allantoic fluid, 30 mM MgCl₂, 200 µg/ml DEAE-dextran (Sigma) and 0.8% agar Nobel (Difco). Virus from the third round of

- plaque purification (designated clone E13-1) was grown in embryonated eggs and four days after inoculation the
20 allantoic fluid was harvested and stored in aliquots at -70°C. The fowlpox recombinant virus fpEFLT7pol (Britton et al., 1996; hereafter called FPV-T7), which expresses T7 RNA polymerase, was a kind gift of Dr. Michael Skinner and was grown on QT35 cells.

25

Isolation of viral RNA

- All manipulations were carried out in RNase-free glassware or plastics and all solutions were made up with
30 RNase-free water which was treated with 1% diethyl-pyrocabonate (DEPC) and sterilized by autoclaving. Virus was pelleted from allantoic fluid by centrifugation at 21,000 rpm for 70 min in a Beckman SW40 rotor at 4°C. The pellet was resuspended in homogenization buffer (50 mM Tris-HCl pH 7.5, 50 mM NaCl, 5 mM EDTA, 0.5% SDS) and treated with
35 Proteinase K (200 µg/ml) for 90 min at 37°C during constant agitation. The lysate was extracted two times with an equal

volume of phenol/chloroform (1:1) pH 5.4 and once with an equal volume of chloroform. The viral RNA was precipitated from the aqueous phase by the addition of 0.1 volume of 3M NaOAc pH 5.3 and 2.5 volumes of 100% ethanol. The precipitate
5 was collected by centrifugation, washed once with 70% ethanol, resuspended in water, and stored in aliquots at -70°C.

Reverse transcription

10

Viral RNA (1.5 µg) was mixed with 500 ng of primer in a volume of 12 µl and incubated for 10 min at 70°C. Four µl of 5x RT buffer (250mM Tris-HCl, pH 8.3, 375 mM KCl, 15 mM MgCl₂; GibcoBRL/Life Technologies), 2 µl 0.1M DTT and 2 µl
15 10mM dNTP's (2.5 mM each) was added and the mixture was incubated for 2 min at 42°C. Reverse transcription was performed in a final volume of 20 µl by the addition of 200 Units of reverse transcriptase (Superscript II; GibcoBRL/Life Technologies) followed by incubation for 60 min at 42°C.

20

Polymerase Chain Reaction (PCR)

All PCR reactions which were used to determine the 3' and 5' end of the NDV genome (see below) were carried out by
25 using Taq DNA Polymerase (Perkin Elmer). For the cloning of individual NDV genes or large subgenomic cDNA's, either the proofreading DNA polymerase Pwo, or mixtures of Taq and Pwo (Expand High Fidelity Kit or Expand Long Template Kit) were used according to the instructions of the supplier
30 (Boehringer Mannheim). All samples were incubated for 2 min at 94°C before the start of the indicated number of PCR cycles. After the indicated number of PCR cycles, the samples were incubated at the elongation temperature for at least 3x the duration of the elongation time of the PCR cycle. PCR
35 fragments were purified directly by using the High Pure PCR Product Purification Kit (Boehringer Mannheim) or after

agarose gelelectrophoresis by using the QiaexII extraction kit (Qiagen) essentially as described by the suppliers.

Sequence analysis

5

All sequences were determined by using the PRISM Ready Reaction Dye Deoxy Terminator Cycle Sequencing Kit (Perkin Elmer). Reaction mixtures (5 μ l) were subjected to 25 cycles of linear amplification (10 sec at 94°C, 5 sec at 50°C, and 4 min at 60°C) in a GeneAmp2400 thermocycler. Subsequently, the reaction mixtures were precipitated with ethanol, washed once with 70% ethanol, resuspended in 15 μ l TSR buffer (Perkin Elmer) and heated for 2 min at 94°C before being loaded on an Applied Biosystems AB310 automatic sequencer.

15

The nucleotide sequences of the primers which were used to sequence the complete genome of NDV strain LaSota were either derived from published sequences or from sequences established during this sequencing project. The primers are shown in Table 1.

20

Cloning and sequencing of the 3' and 5' termini of the genome of NDV strain LaSota

25

The nucleotide sequence of the 3' and 5' termini of the NDV genome were determined by using RACE procedures (rapid amplification of cDNA ends). NDV RNA was used in a reverse transcription reaction in a final volume of 20 μ l by using primer p360 (5'-GGCGATGTAATCAGCCTAGTGCTT-3'; nt 14756-14779) which was derived from the published sequence of the L-gene of NDV (Yusoff et al., 1987). The single-stranded cDNA (2.5 μ l of the RT mixture) was added to 8 pmol anchorprimer ALG3 (5'-CACGAATTCATCGATTCTGGATCCTTC-3') and ligated overnight at room temperature in 20 μ l of a reaction mixture containing 50 mM Tris-HCl, pH 8.0, 10 mM MgCl₂, 10 μ g/ml BSA, 25% PEG, 1 mM HCC, 20 μ M ATP and 10 units of T4 RNA ligase (New England Biolabs) as described by Tessier et al. (1986). One μ l of the

35

ligation reaction was used as template in a PCR reaction by using primers p375 (5'-CAATGAATTCAAAGGATATTACAGTAACT-3'; nt 14964-14983) and ALG4 (5'-GAAGGATCCAGAATCGATAG-3'). The latter primer is complementary to anchorprimer ALG3. The PCR conditions (40 cycles) were as follows: 1 min at 94°C, 1 min at 55°C, and 2 min at 72°C. The PCR products were purified and cloned in T-vector pBluescriptII-TSK (Ichihara and Kurosawa, 1993). Alternatively, the purified PCR products were treated with Klenow DNA polymerase I to create blunt ends and cloned in the HincII-site of plasmid pGEM4Z (Promega). Thirteen independent clones (8x pBluescriptII-TSK and 5x pGEM4Z) were sequenced to determine the nucleotide sequence of the 5' end of the genome of NDV strain LaSota. The nucleotide sequence of the 3' end was determined by two independent methods. In method I, primer ALG3 was ligated to the 3' end of the viral RNA by using T4 RNA ligase as described by Schütze et al. (1995). The reaction mixture (final volume 10 µl) contained 2.5 µg NDV RNA, 100 pmol ALG3, 1 µl 10x T4 RNA ligase buffer (500mM Tris-HCl, pH 7.8, 100 mM MgCl₂, 100 mM DTT, 10 mM ATP), 1 µl DMSO, 1 µl 10 uM hexamine-cobaltchloride, 1 µl RNasin (Promega) and 10 units of T4 RNA ligase (New England Biolabs). The mixture was incubated overnight at room temperature and 5 µl of the ligation reaction was used as template in a reverse transcription reaction by using ALG4 as primer. One µl of the RT-reaction was used in a PCR reaction by using primers ALG4 and p376 (5'-GAGCCTTAAGGAGCTGCTCGTACTGATC-3'; nt 137-164) which was derived from the published sequence of the 3' end of NDV (Ishida et al., 1986). The PCR conditions were as described above for the 5' RACE. In method II, the 3' and 5' ends of the viral NDV RNA were ligated to each other by using T4 RNA ligase using the same conditions as described above for method I. Five µl of the ligation mixture was used as template in a reverse transcription reaction by using primer p360. One µl of the RT-reaction was used in a PCR reaction by using primers p375 and p376 and PCR conditions described above for the 5' RACE. The PCR products were treated with

Klenow DNA polymerase I to create blunt ends and cloned in the HincII-site of plasmid pGEM4Z (Promega). Ten independent clones (4 from method I and 6 from method II) were sequenced to determine the nucleotide sequence of the 3' end of the genome of NDV strain LaSota.

Construction of transcription vector

A low-copy-number transcription vector was constructed by using plasmid pOK12 (Vieira and Messing, 1991) as the basic replicon. Plasmid pOK12 was digested with PvuII and the DNA fragment containing the replication origin and the Kanamycin-resistance gene was isolated. This DNA fragment was ligated to an Eco47III-AflIII fragment (the AflIII site was made blunt by using Klenow DNA polymerase I) from transcription vector 2.0 (a generous gift of Dr. Andrew Ball; Pattnaik et al., 1992). From the resulting plasmid an XbaI-NheI fragment was deleted to eliminate as much unique restriction sites as possible. The resulting plasmid was designated pOLTV5 (Fig. 1). Transcription vector pOLTV5 contains the T7 DNA dependent RNA polymerase promoter followed by unique StuI and SmaI restriction sites, the autocatalytic ribozyme from hepatitis delta virus (HDV) and the transcription termination signal from bacteriophage T7. DNA fragments cloned between the StuI and SmaI restriction sites can be transcribed either in vitro or in vivo by using T7 RNA polymerase. After transcription, the 5' end of the resulting transcripts contains two G residues encoded by the plasmid. Due to the autocatalytic action of the HDV ribozyme, the 3' end of the transcripts corresponds to the exact terminal nucleotide of the cloned DNA fragment (Pattnaik et al., 1992).

Construction of minigenome plasmids

5 In order to examine the requirements for replication and transcription of NDV, minigenome plasmids were constructed which contained the 3' and 5' terminal regions of NDV flanking a reporter gene that replaced all NDV genes (Fig. 2). DNA fragments corresponding to the 3' and 5' terminal
10 regions of NDV were generated by means of PCR by using Pwo DNA polymerase (30 cycles; 15 sec at 94°C, 30 sec at 50°C, and 30 sec at 72°C) and using plasmids containing the 3' and 5' RACE fragments as templates (see above).
The 3' region (nt 1-119) was generated by using primers 3UIT
15 (5'-ACCAAACAGAGAATCCGTGAGTTACGA-3', nt 1-27) and SEAP3 (5'-ATCGATACTGGTCAGCATGCTGGCAGAAGGCTTTCTCG-3', nt 102-119). The 5' region (nt 14973-15186) was generated by using primers SEAP5 (5'-GCATGCTGACCAGTATCGATATTACAGTAACTGTGACT-3', nt 14973-14990) and 5NDV (5'-ACCAAACAAAGATTTGGTGAATGACGA-3', nt
20 15158-15186). The two DNA fragments were joined in an overlap PCR (the overlap is shown in italics in the primer sequences shown above) by using primers 3UIT and 5NDV. The resulting DNA fragment, which is a fusion of the 3' and 5' end of NDV separated by 20 nucleotides, was phosphorylated by treatment
25 with T4 polynucleotide kinase and cloned in both orientations in transcription plasmid pOLTV5 (Fig. 1) which was cleaved with StuI and SmaI and dephosphorylated with calf intestinal phosphatase (Boehringer Mannheim). Finally, the SEAP-gene (encoding secreted alkaline phosphatase) was recovered from
30 plasmid pSEAP-Basic (Clontech) by digestion with SphI and ClaI, and cloned between the SphI and ClaI sites between the 3' and 5' ends of NDV. The resulting plasmids were designated pOLTV535 and pOLTV553, respectively. In vivo or in vitro transcription using T7 RNA polymerase of plasmid pOLTV535
35 gives rise to antigenomic RNA ([+]-RNA), whereas

transcription of plasmid pOLTV553 gives rise to genomic RNA ([-]-RNA).

Plasmids pOLTV535N0 to -N5 and pOLTV553N0 to -N5 were generated by inserting self-complementary oligonucleotides in the ClaI-site located between the SEAP-gene and the 5' end of NDV in pOLTV535 and pOLTV553, respectively (see Fig. 2). The oligonucleotides used were: N0, 5'-CGCGAGCTCG-3'; N1, 5'-CGCGAGSCTCG-3'; N2, 5'-CGCGAGCGCTCG-3'; N3, 5'-CGCGAGCWGCTCG-3'; N4, 5'-CGCGAGCATGCTCG-3'; N5, 5'-CGCGAGCASTGCTCG-3' (W = A or T; S = C or G).

Modification of the T7 promoter in plasmids pOLTV535 and pOLTV553

To generate in vitro or in vivo transcripts containing the authentic 5' and 3' terminal ends of NDV, the T7 promoter in plasmids pOLTV535 and pOLTV553 was modified such that transcription would start at the first nucleotide of the 3' or 5' terminal end of NDV.

Primers were designed which contained, 1) a BglI-restriction site, 2) the sequence of the T7 promoter (shown in italics) which was modified such that the two G residues at the end of the T7 promoter were replaced by an A residue, and 3) the 3' (nt 1-21) or 5' end (nt 15164-15186) of NDV. Primers BGL3F2 (5'-GATATGGCCATTCAGGCTTAATACGACTCACTATAACCAAACAGAGAATCCGTGAG-3') and SEAP3 (see above) were used to generate a DNA fragment containing the modified T7 promoter and the entire 3' end of NDV up to the start of the SEAP gene in pOLTV535.

Similarly, a DNA fragment was generated that containing the modified T7 promoter and the entire 5' end of NDV up to the end of the SEAP gene in pOLTV553 by using primers BGL5F2 (5'-GATATGGCCATTCAGGCTTAATACGACTCACTATAACCAAACAAGATTGCGTGAATG-3') and SEAP5. The resulting fragments were digested with BglI and SphI (3' end) or BglI and ClaI (5' end), respectively, and used to replace the BglI-SphI fragment in

pOLTV535, or the BglI-ClaI fragment in pOLTV553. The resulting plasmids were designated pOLTV735 and pOLTV753, respectively. Plasmids pOLTV735N3 and pOLTV753N3 were generated by inserting a self-complementary oligonucleotide (5'-CGCGAGCGCTCG-3'; W = A or T) in the ClaI-site located between the SEAP-gene and the 5' end of NDV in pOLTV735 and pOLTV753, respectively.

Construction of SEAP-reporter plasmids

10

Plasmid pCIneoSEAP was constructed by cloning an XhoI-ClaI fragment (ClaI-site was made blunt by using Klenow DNA polymerase I) containing the SEAP-gene from plasmid pSEAP-Basic (Clontech) between the XhoI and SmaI sites of the eukaryotic expression vector pCIneo (Promega). The latter plasmid contains the human cytomegalovirus (hCMV) promoter in addition to the bacteriophage T7 promoter. In order to examine and quantitate SEAP expression by transcripts generated from the T7 promoter only, another plasmid was constructed which lacked the hCMV promoter. To this end the hCMV promoter was deleted from pCIneo by partial digestion with HindIII followed by complete digestion with BglII. The DNA fragment (nt 756-5469 according to numbering of Clontech) from which the hCMV promoter was deleted was isolated, treated with T4 DNA polymerase to generate blunt ends and recircularized by using T4 DNA ligase. The resulting plasmid was designated pCIneoD. Finally, the SEAP gene was recovered from pSEAP-Basis as an MluI-AccI fragment and cloned in pCIneoD between the MluI and ClaI sites. The resulting plasmid was designated pCIneoD SEAP.

Transfections

Cells were seeded in 24-well culture dishes, grown overnight to 60-80% confluency, and infected at a m.o.i. of 1 with FPV-T7 for 1h at 37°C. The cells were transfected with 0,5 µg minigenome plasmid DNA by using 3 µl of LipofectAMINE™

and OptiMem essentially as described by the supplier (GibcoBRL/Life Technologies). After incubation for 4h (CER cells) or 16H (QM5 cells) at 37°C the cells were either infected with NDV (Dutch virulent isolate nr. 152508; 200 µl per well) for 1h at a m.o.i. of 5, or left uninfected. The inoculum was aspirated and replaced by 1ml of complete medium and the cells were further incubated at 37°C.

For co-transfections, cells were grown in 6-well culture dishes and infected with FPV-T7 as described above. The cells were co-transfected with 0.25 µg minigenome plasmid DNA, 0.4 µg pCIneoNP, 0.2 µg pCIneoP and 0.2 µg pCIneoL(c) or pCIneo by using either 8 µl of LipofectAMINE or 9 µl of FuGene[®]6 (Boehringer Mannheim). In order to generate infectious virus, the minigenome plasmid was replaced by a transcription plasmid that contained the full-length NDV cDNA.

Quantitation of SEAP activity

The amount of SEAP which was secreted into the medium of transfected cells was measured in disposable 96-well plates by using the Phospha-Light[™] Chemiluminescent Reporter Assay for Secreted Alkaline Phosphatase kit essentially as described by the supplier (Tropix). Chemiluminescence was quantitated by using a liquid scintillation counter (Wallac 1450 microbeta PLUS).

Cloning and sequencing of cDNA's spanning the entire genome of NDV strain LaSota.

To clone and sequence the entire genome of NDV strain LaSota, large subgenomic cDNA clones were generated by means of RT-PCR and cloned in pGEM-T. First strand cDNA synthesis was performed by using primer 3UIT as described above, and 1 µl of the RTreaction was used in a PCR reaction by using the Expand Long Template PCR kit (Boehringer Mannheim). The PCR consisted of 5 cycles of 10 sec at 94°C, 30 sec at 58°C, and 6 min at 68°C, followed by 10 cycles of 10 sec at 94°C, 30

sec at 58°C, and 6 min at 68°C, in which the elongation time at 68°C was increased by 20 sec per cycle. The PCR fragments were cloned in pGEM-T by using the pGEM-T cloning kit essentially as described by the supplier (Promega). Ligation mixtures were transformed into *E.coli* strain SURE II (Stratagene). Two independent RT-PCR reactions (A and B) were performed and each yielded a similar set of cDNA clones. The nucleotide sequence of the subgenomic cDNA clones was determined by using NDV-specific primers (Table 1) and by primers flanking the inserts. After comparison of the nucleotide sequence of the A and B series of clones, remaining ambiguities were resolved by sequencing relevant regions of a third independent series of cDNA's (C series). The nucleotide sequence of NDV strain LaSota is shown in Fig.3.

Construction of a full length genomic cDNA clone of NDV.

The full-length NDV cDNA was assembled in transcription plasmid pOLTV5 by using pOLTV535 as the starting plasmid. The DNA fragments were joined at overlaps by using common restriction enzymes as detailed in Fig. 4B. In a series of cloning steps, a plasmid (designated p535-DI) was constructed containing nucleotides 1-3521 and 12355-15186 separated by a ClaI-site that was generated by joining the ClaI-sites at position 3521 and 12355. In another series of cloning steps, a plasmid (designated pGEM-B) was constructed which contained part of the NDV genome including nucleotides 3521-12355 (ClaI-fragment). To facilitate cloning, the latter ClaI-fragment was tagged with the Chloramphenicol resistance (Cm) gene from plasmid pACYC184 (Chang and Cohen, 1978). To this end the Cm-gene was recovered from pACYC184 by means of PCR by using primers CAT-F (5' GCGTACGTCTAGACTGGTGTCCCTGTTGATACCGG-3') and CAT-R (5'-GCTCTAGACGTACGACCCTGCCCTGAACCGACG-3'). The PCR was carried out with Pwo DNA polymerase and consisted of 30 cycles of 30 sec at 94°C, 45 sec at 60°C, and 60 sec at 72°C. The

resulting PCR fragment was digested with BsiWI and cloned in the unique BsiWI site of pGEM-B, yielding pGEM-B(CAT). The ClaI fragment from pGEM-B(CAT) was cloned in the unique ClaI site of p535-DI, yielding pNDFL(CAT). Finally, the Cm-gene was removed from this plasmid by digestion with BsiWI followed by religation and transformation of E.coli strain DH5a. The resulting plasmid was designated pNDFL+ and contains the entire NDV cDNA sequence cloned between the T7 promoter and the HDV ribozyme in transcription plasmid pOLTV5.

Cloning and expression of individual NDV genes.

DNA fragments containing each of the NDV LaSota genes were generated by means of RT-PCR and cloned in pCIneo. After cloning, all fragments were sequenced by using primers flanking the inserts and by gene-specific primers.

NP-gene: Primer 386 (5'-GAGCAATCGAAGTCGTACGGGTAGAAGGTG-3', nt 40-69) was used for reverse transcription. Primers 365 (5'-GTGTGAATTCCGAGTGCGAGCCCGAAG-3'; nt 77-94) and 892 (5'-TTGCATGCCTGCAGGTCAGTACCCCCAGTC-3'; nt 1577-1593) were used for PCR by using Pwo DNA polymerase. The following PCR profile (30 cycles) was used; 30 sec at 95°C, 40 sec at 65°C, and 45 sec at 72°C. The resulting DNA fragment was digested with EcoRI and cloned in pCIneo between the EcoRI and SmaI sites. Expression of NP was verified in an immunoperoxidase monolayer assay (IPMA) as described by Peeters et al (1992) by using monoclonal antibody 38 (Russell et al., 1983).

P-gene: Primer pRT1 (5'-CAAAGAATTCAGAAAAAAGTACGGGTAGAA-3'; nt 1794-1814) was used for reverse transcription. Primers pRT1 and p2 (5'-GCAGTCTAGATTAGCCATTCACTGCAAGGCGC-3'; nt 3053-3071) were used for PCR by using Pwo DNA polymerase. The following PCR profile (30 cycles) was used; 30 sec at 95 °C, 40 sec at 65°C, and 60 sec at 72°C. The resulting DNA fragment was digested with EcoRI and XbaI and cloned in pCIneo between the EcoRI and XbaI sites. Expression of P was verified in an IPMA by using monoclonal antibody 688 (Russell et al., 1983).

M-gene: Primer 3UIT (5'-ACCAAACAGAGAATCCGTGAGTTACGA-3'; nt 1-27) was used for reverse transcription. Primers NDV5M (5'-GGGTGCTAGCGGAGTGCCCCAATTGTGCCAA-3'; nt 3268-3288) and NDV3M (5'-TCTCCCCGGGGCAGCTTATTTCTTAAAAGGAT-3'; nt 4368-4389) were used for PCR by using the Expand High Fidelity kit. The PCR consisted of 10 cycles of 15 sec at 95°C, 30 sec at 55°C, and 2 min at 68°C, followed by 15 cycles in which the elongation time at 68°C was increased for 20 sec per cycle. The resulting DNA fragment was treated with T4 DNA polymerase to create blunt ends, digested with *NheI*, and cloned in pCIneo between the *NheI* and *SmaI* sites. Expression of the M protein was verified in an IPMA by using monoclonal antibody 424 (Russell et al., 1983).

F-gene: Primer 3UIT (see above) was used for reverse transcription. Primers NDV5F (5'-ACGGGCTAGCGATTCTGGATCCCGGTTGG-3'; nt 4508-4526) and NDV3F (5'-ACTACCCGGGAAACCTTCGTTCTCAT-3'; nt 6212-31) were used for PCR by using the Expand High Fidelity kit using the conditions described above for the M-gene. The resulting DNA fragment was treated with T4 DNA polymerase to create blunt ends, digested with *NheI*, and cloned in pCIneo between the *NheI* and *SmaI* sites. Expression of the F protein was verified in an IPMA by using monoclonal antibody 8E12A8C3 (ID-DLO, department of Avian Virology).

HN-gene: Primer 3UIT was used for reverse transcription. Primers NDV5HN (5'-GTAGGCTAGCAAGAGAGGCCGCCCTCAAT-3'; nt 6335-6354) and NDV3HN (5'-CGAGCCCGGGCCGGCATTTCGGTTTGATTCTTG-3'; nt 8205-8227) were used for PCR by using the Expand High Fidelity kit using the conditions described above for the M-gene. The resulting DNA fragment was treated with T4 DNA polymerase to create blunt ends and after digestion with *XmaI* it was cloned in pCIneo between the blunted (Klenow DNA polymerase) *NheI* site and the *XmaI* site. Expression of the HN protein was verified in an IPMA by using monoclonal antibody 86 (Russell et al., 1983).

L-gene: The L-gene was recovered from cDNA clone pGEM-L7a (Figure 4A) by digestion with *SacII* and *SalI*. Before

digestion with SalI, the SacII site was made blunt by treatment with T4 DNA polymerase. The resulting fragment was cloned in pCIneo between the blunted (Klenow DNA polymerase) NheI site and the SalI site. The 5' untranslated region between the T7 promoter and the ATG startcodon of the L-gene contained 2 out-of-frame ATG codons which might interfere with correct expression of the L protein. Therefore, a new plasmid was constructed in which the first ATG was missing and in which the second ATG was changed to AAG by means of PCR mutagenesis, as follows. Primers 5LE(E) 5'-CAATGGAATTCAAGGCAAAACAGCTCAAGGTAAATAATACGGG-3'; nt 8332-8374) and 3LE(B) 5'-GTGAATCTAGGAATGCCGGATCCGTACGAATGC-3'; nt 8847-8870) were used in a PCR reaction using plasmid pGEM-L7a (Fig. 4) as a template. The PCR was carried out by using Pwo DNA polymerase and consisted of 30 cycles of 30 sec at 94°C, 45 sec at 60°C, and 60 sec at 72°C. The resulting DNA fragment was digested with EcoRI and XbaI and cloned in pCIneo between the EcoRI and XbaI sites, generating plasmid pCIneoL(N). Subsequently, the BsiWI-SalI fragment from pGEM-L7a, which contains the remaining part of the L-gene (nt 8852-15046), was cloned in pCIneoL(N) between the BsiWI and SalI sites, generating plasmid pCIneoL(c). Since antibodies against the L-protein are not available, expression of L could not be checked by immunochemistry.

25

Introduction of a genetic tag in the F-gene.

To show unambiguously that infectious virus can be generated from cloned full-length cDNA, a genetic tag was introduced in the F gene by means of PCR mutagenesis. To this end, the F-gene was cloned by using two overlapping PCR fragments. The first PCR fragment was generated by using primers NDV5F (see above) and primer F5R (5'-AAAGCGCGCTGTCTCCTCCCTCCAGATGTAGTCAC-3'; nt 4859-4894). The residues shown in bold are changes which were introduced in the primer in order to change the amino acid sequence of the

proteolytic cleavage site between F1 and F2 from that of the NDV LaSota strain (GGRQGR | L) to that of the consensus cleavage site for virulent NDV strains (GRRQRR | F). The second PCR fragment was generated by using primers F3F (5'-
5 GGAGGAGACAGCGGCGCTTTATAGGCGCCATTATTGG-3'; nt 4875-4911) and IV09 (5'-CTCTGTCGACACAGACTACCAGAACTTTCAC-3'; nt 6246-6266). The PCR was performed with Pwo DNA polymerase and consisted of 25 cycles of 15 sec at 94°C, 30 sec at 55°C, and 2 min at 72°C. The two overlapping PCR fragments (the overlap is shown
10 in italics in the primer sequences) were joined in a second PCR by using primers NDV5F and IV09 and by using the same PCR conditions. The resulting fragment, which contains the entire ORF of the F gene and which encodes a virulent consensus cleavage site, was digested with NheI and SalI and cloned in
15 pCIneo between the NheI and SalI sites, yielding pCIneoF^{wt}. The StuI-NotI fragment (nt 4646-4952) from pCIneoF^{wt} was used to replace the corresponding fragment in plasmid p535-S which had been constructed by inserting the ClaI-ScaI (nt 3521-10311) from pGEM-B in p535-DI between the ClaI and ScaI sites
20 (see Fig. 4C). The resulting plasmid was designated p535-S[F^{wt}c]. A PCR fragment containing the Cm-resistance gene from pACYC184 (see above) was cloned as an XbaI fragment into the unique XbaI site (position 6172 in the NDV sequence) of plasmid p535-S[F^{wt}c], yielding plasmid p535-S[F^{wt}c]Cm.
25 Subsequently, the Cm-tagged ApaI-SpeI fragment (nt 2285-8094) of this plasmid was used to replace the corresponding fragment of the full-length cDNA clone pNDFL+. Finally, the Cm-gene was removed from this plasmid by digestion with XbaI followed by recircularization using T4 DNA ligase. The
30 resulting plasmid, which contains the genetically tagged full-length NDV cDNA, was designated pNDFL+[F^{wt}].

Generation of stably transformed cell lines that express individual NDV genes

5 Plasmids pCIneoNP, pCIneoP, pCIneoM, pCIneoF, pCIneoF^{wt}, and pCIneoHN were used for the generation of stably transformed cell lines that express these proteins individually. The day before transfection, CER cells were seeded in 6 cm culture dishes and incubated overnight to give
10 a confluency of 60-80%. The cells were transfected with 2 µg of plasmid DNA by using 12 µl of LipofectAmine and OptiMem essentially as described by the supplier (GibcoBRL/Life Technologies). After 48h the cells were trypsinized and dilutions were seeded in 10 cm culture dishes in medium
15 containing 500 µg/ml of G418 (Boehringer Mannheim). Every 3 days the medium was replaced by fresh medium containing increasing (in steps of 100 µg/ml) amounts of G418 until a concentration of 800 µg/ml was reached. Cells were kept in medium containing 800 µg/ml G418 and three weeks after
20 transfection individual colonies were picked and transferred to 96-well culture dishes. The cloned cell lines were examined for the expression of the respective NDV gene by using an IPMA as described above for transient expression studies.

25 Cell lines that constitutively expressed NP, P, M or F could be identified and isolated. We were unable, however, to generate cell lines that expressed the HN protein. Perhaps constitutive expression of HN is toxic to the cells.

30

Generation of stably transformed cell lines that express T7 RNA polymerase

The gene encoding T7 RNA polymerase was recovered from
35 plasmid pRT7NT (René van Gennip, ID-DLO, Department of Mammalian Virology) by digestion with EcoRI and SalI. The

resulting fragment contains the T7 RNA polymerase gene located behind the baculovirus p10 promoter. The DNA fragment was cloned in plasmid pCIneo0 between the EcoRI and Sali sites, generating plasmid pCIneo107. Plasmid pCIneo0 lacks the T7 promoter and was derived from pCIneo by cleavage with NheI followed by partial cleavage with ScaI, filling in the sticky ends with Klenow DNA polymerase and recircularization by using T4 DNA ligase. The baculovirus sequences were removed from pCIneo107 by digestion with EcoRI and PacI, followed by T4 DNA polymerase treatment to generate blunt ends and recircularization. The resulting plasmid was designated pCIneo007. Expression of T7 RNA polymerase was verified by co-transfection of cells with pCIneo007 and pPRh01. The latter plasmid contains the E2 protein of classical swine fever virus cloned behind a T7 promoter and containing an internal ribosome entry site (René van Gennip, personal communication). Expression of E2 was determined in an IPMA by using monoclonal antibody V4 (Wensvoort et al., 1986). Stably transformed CER cell lines expressing T7 RNA polymerase were generated and isolated as described above except that 10 cm culture dishes were used and the cells were transfected with 5 µg of pCIneo007 DNA and 25 µl of LipofectAmine. To examine individual cell lines for the expression T7 RNA polymerase, they were transfected with plasmid pPRh01 and expression of E2 (which is dependent on T7 RNA polymerase) was determined in an IPMA by using monoclonal antibody V4. Several cell lines which expressed T7 RNA polymerase were identified. One cell line, designated CER-C9, was used for subsequent experiments.

30

Cloning and expression of HN-genes and hybrid-HN genes

Primer 3UIT was used to synthesize single-stranded cDNA of NDV and avian paramyxovirus serotype-2 and -4 (APMV2 and APMV4) as described above. All subsequent PCR reactions were performed by using 25 cycles of 15 sec at 94°C, 30 sec at 55°C and 2 min at 72°C.

35

The entire coding region of the HN-gene of APMV2 was recovered by means of PCR by using primers IV03 (5'-GGGGGAATTCCCCATTCAATGAAGGGTCTAC-3') and IV05 (5'-GATCCCCGGGTCTTAAACCAGGCTTCGCAATG-3') which were derived from the sequence of the HN gene of APMV2 (GenBank accession number D14030). The entire coding region of the HN-gene of APMV4 was recovered by means of PCR by using primers IV06 (5'-GGGGGAATTCTGGTAGGGTGGGGAAGGTAGC-3') and IV08 (5'-ATTGCCCGGGGGGTAATAATCAGGATCTCAG-3') which were derived from the sequence of the HN-gene of APMV4 (GenBank accession number D14031). The resulting PCR fragments were digested (either directly or after subcloning in pGEM-T), with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmids were designated pCIneoHN2 and pCIneoHN4, respectively.

Hybrids between the HN-gene of NDV strain LaSota and the HN-genes of APMV2 and -4 were constructed by means of overlap PCR as follows. The N-terminal part (aa 1-141) of the HN-gene of NDV strain LaSota was amplified with Pwo DNA polymerase by using primers IV01B (5'-GTAGGAATTCAAGAGAGGCCGCCCTCAAT-3'; nt 6325-6354) and IV10 (5'-AATGAGTTCTTTGCCTATCCCCC-3'; nt 6811-6834). The C-terminal part of the HN-gene of APMV2 (aa 142-580) was amplified with Pwo DNA polymerase by using primers IV11B (5'-GGGGGGATAGGCAAAGAACTCATTCAAGGAGATGCATCTGCAGGC-3') and IV05. The resulting PCR fragments were joined in an overlap PCR (overlap shown in italics) by using primers IV01B and IV05 and by using the Expand High Fidelity enzyme mix. The resulting PCR fragment was digested (either directly or after subcloning in pGEM-T) with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmid which contains a hybrid HN-gene consisting of aa 1-141 of NDV and aa 142-580 of APMV2 was designated pCIneoHN1/2^{M1}.

The C-terminal part of the HN-gene of APMV4 (aa 143-569) was amplified by using primers IV14B (5'-GGGGGGATAGGCAAAGAACTCATTGTAGATGATGCATCTGCAGGCCTAAATTTCC-3')

and IV08. This fragment was joined with the N-terminal part of the HN-gene of NDV (see above) in an overlap PCR by using primers IV01B and IV08. The resulting PCR fragment was digested (either directly or after subcloning in pGEM-T) with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmid which contains a hybrid HN-gene consisting of aa 1-141 of NDV and aa 143-569 of APMV4 was designated pCIneoHN1/4¹⁴¹.

In analogy to the constructions described above, hybrid HN genes were constructed which consisted of aa 1-143 of NDV and aa 144-580 of APMV2, or aa 1-143 of NDV and aa 145-569 of APMV4. For these constructions PCR fragments were obtained by using the following pairs of primers; NDV aa 1-143, primer IV01B and IV13 (5'-ATCTACAATGAGTTCTTTGCCTATC-3'; nt 6816-6840); APMV2 aa 144-580, primer IV14B (5'-GGGGGGATAGGCAAAGAACTCATTGTAGATGATGCATCTGCAGGCCTAAATTTCC-3') and IV05; APMV4 aa 145-569, primer IV15B (5'-GGGGGGATAGGCAAAGAACTCATTGTAGATCAAACAGCTGACTACACAGCAG-3') and IV08. The PCR fragments were digested (either directly or after subcloning in pGEM-T) with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmids were designated pCIneo1/2¹⁴³ and pCIneo1/4¹⁴³, respectively. To examine expression of the HN proteins, CER cells or QM5 cells were infected with FPV-T7 for 1h at a m.o.i. of 1, transfected with plasmids pCIneoHN, pCIneoHN2, pCIneoHN4, pCIneoHN1/2¹⁴¹, pCIneoHN1/2¹⁴³, pCIneoHN1/4¹⁴³ and pCIneoHN1/4¹⁴³, and 24h after transfection the monolayers were overlaid with a 1% suspension of chicken erythrocytes in PBS for 45 min at roomtemperature. Subsequently, the monolayers were carefully washed three times with PBS and adhesion of erythrocytes to transfected cells was examined microscopically. To examine induction of cell fusion after co-expression of the HN and F protein, CER cells or QM5 cells were co-transfected with pCIneoF^{wt} together with either pCIneo-HN1, pCIneoHN2, pCIneoHN4, pCIneoHN1/2¹⁴¹,

pCIneoHN1/4¹⁴¹, pCIneoHN1/2¹⁴³ or pCIneoHN1/4¹⁴³. After incubation for 2 to 3 days, the monolayers were washed with PBS, stained for 15 min with a Giemsa solution (1:30 dilution in water), and examined microscopically.

5

Cloning of hybrid-HN genes in full length genomic NDV cDNA

A synthetic linker, designated HN12, was inserted between the NotI and SpeI sites of pGEM-T (Promega) by using
10 oligonucleotides HN12a (5'- GGCCGCATATTCTAGAGTTAACGACTTA-3') and HN12b (5'-CTAGTAAGTCGTTAAGTCTAGAAATATGC-3'). A synthetic linker, designated HN14, was inserted between the NotI and SpeI sites of pGEM-T by using oligonucleotides HN14a (5'- GGCCGCATATTCTAGAGTTAACGA-3') and HN14b (5'-
15 CTAGTCGTTAAGTCTAGAAATATGC-3'). The resulting plasmids were designated pGEM-HN12 and pGEM-HN14, respectively. These plasmids were digested with NotI and XbaI and used to clone the NotI-SpeI fragment (nt 3390-7488) from plasmid p535-S[F^{wt}c]Cm. The resulting plasmids were designated pGEM-
20 HN1/2NS and pGEM-HN1/4NS, respectively. The HN genes of these plasmids were replaced by the hybrid HN genes from plasmids pCIneoHN1/2¹⁴³ and pCIneoHN1/4¹⁴³, respectively (see section: Cloning and expression of HN-genes and hybrid-HN genes). To this end, pCIneoHN1/2¹⁴³ and pCIneoHN1/4¹⁴³ were digested with
25 NheI and SmaI and the resulting fragments (containing the hybrid HN1/2¹⁴³ and hybrid HN1/4¹⁴³ genes) were cloned between the NheI and HpaI site of plasmids pGEM-HN1/2NS and pGEM-HN1/4NS, resulting in pGEM+HN12 and pGEM+HN14, respectively. The latter plasmids were used to introduce the hybrid HN
30 genes into the full length genomic cDNA clone of NDV. To this end, plasmids pGEM+HN12 and pGEM+HN14 were digested with NotI and SpeI and the fragment containing either the HN12 or HN14 gene was used to replace the corresponding fragment of pNDFL+, yielding pNDFL+HN1/2¹⁴³ Cm and pNDFL+HN1/4¹⁴³ Cm,
35 respectively. The Cm-gene was removed from these plasmids by digestion with XbaI followed by recircularization using T4 DNA ligase. In order to comply with the "rule-of-six", a

linker was inserted into the unique SpeI site of these plasmids by using self-complementary oligonucleotides. Linker H2 (5'-CTAGCGAGCGCTCG -3') was inserted in plasmid pNDFL+HN1/2¹⁴³ and linker H3 (5'-CTAGCGAGCGCTCG-3) was

inserted in pNDFL+HN1/4¹⁴³, yielding plasmids pNDFL+HN1/2¹⁴³(H2) and pNDFL+HN1/4¹⁴³(H3), respectively.

Elimination of a specific epitope in the HN protein of NDV La Sota

A specific epitope, i.e. amino acids 346 to 354 (PDEQDYQIR), in the HN protein of NDV LaSota that is recognized by MAb 4D6 (Long et al., 1986; Meulemans et al., 1986), was eliminated by replacing this sequence by the corresponding sequence of the HN proteins of either APMV-2 (NRTDIQQT I) or APMV-4 (PDPLQDQIL). To this end, plasmid pCIneoHN (see section: Cloning and expression of individual NDV genes) was used as template to create overlapping PCR fragments. For the APMV-2 sequence the first PCR fragment was generated by using primers IV01 (5'-GTAGACGCGTAAGAGAGGCGCCCTCAAT-3') and primer 3HN2 (5' GATAGTTTGCTGTATATCAGTCCGATTGCATGTGTCATTGTATC-GCTTGTATATCAC-3'). The second PCR was generated by using the primers 5HN2 (5'-AATCGGACTGATATACAGCAAACTATCATGGCCAAGTCTT-CGTATAAGCCTGGAGCC-3') and NDV3-HN (5'-CGAGCCCGGGCCGGCATT-CGGTTTGATTCTTG- 3'). The resulting fragments were combined and used as template for a third PCR by using the primers IV01B (5'-GTAGGAATTCAAGAGAGGCGCCCTCAAT-3') and primer NDV3-HN. For the APMV-4 sequence the first PCR fragment was generated by using primers IV01 and primer 3HN4 (5' TAAGATC-TGATCTTGCAGCGGGTCAGGGCATGTGTCATTGTATCGCTTGTATATCAC-3'). The second PCR was generated by using the primers 5HN4 (5'-CCTGACCGCTGCAAGATCAGATCTTAATGGCCAAGTCTTCGTATAAGCCTGGAGCC-3') and NDV3-HN. The resulting fragments were combined and used as template for a third PCR by using the primers IV01B and NDV3-HN. Primers 3HN2 / 5HN2 and 3HN4/5HN4 are partly

complementary and contain the genetic codes for the HN2 sequence (NRTDIQQTI) and HN4 sequence (PDPLQDQIL), respectively. The PCR reactions were performed by using the Expand Long Template PCR kit (Boehringer Mannheim). The PCR
5 consisted of 30 cycles of 10 sec 94°C, 30 sec 58 °C and 2 min at 68 °C, followed by 1 cycle of 4 min 68 °C. The PCR fragments were digested with EcoNI and Bsu36I, and cloned between the EcoNI and Bsu36I sites of pCIneoHN. The resulting plasmids were designated pCIneoHN1(HN2e) and
10 pCIneoHN1(HN4e) respectively. Transient expression studied indicated that the modified HN proteins were correctly expressed and transported to the cell surface as judged from haemadsorption studies using chicken erythrocytes. Furthermore, MAb 6D4 which is directed against a linear
15 epitope of HN of NDV and which consists of (or at least includes, amino acids 346-354) did not react with the modified HN proteins.

Plasmids pCIneoHN1(HN2e) and pCIneoHN1(HN4e) were digested with NarI and SpeI and the fragments containing the modified
20 HN genes were cloned between the NarI and SpeI sites of pGEM-HN1/2NS and pGEM-HN1/4NS, respectively. The resulting plasmids, designated pGEM-HN1(HN2e) and pGEM-HN1(HN4e), were digested with NotI and SpeI, and used to replace the NotI-SpeI fragment in pNDFL+. The resulting plasmids were
25 designated pNDFL-HN(HN2e)Cm and pNDFL-HN(HN4e)Cm, respectively. The Cm-gene was removed from these plasmids by digestion with XbaI followed by religation. The resulting plasmids were designated pNDFL-HN(HN2e) and pNDFL-HN(HN4e), respectively.

30

RESULTS

Nucleotide sequence of the 3' and 5' terminal ends of the genome of NDV strain LaSota

35

The sequence of a putative 3' end of the NDV genome has been published (Ishida et al., 1986) albeit from another NDV

strain (D26) than the one used here (LaSota). Yusoff et al. (1987) have published a sequence of the L-gene and a relatively large non-coding region behind the L-gene of NDV strain Beaudette C. However, as shown herein, this sequence did not include the full terminal 5' end of the viral genome which makes it impossible to generate infectious copy NDV. The 3' and 5' terminal ends of the genome of negative-strand RNA viruses fulfil an essential function in replication and transcription (Lamb and Kolakofsky, 1996). Thus, in order to generate a full-length NDV cDNA which can be used to generate infectious virus by means of reverse genetics (Conzelmann, 1996), it is absolutely essential to include the correct 3' and 5' ends of the viral genome. Therefore, we determined the exact nucleotide sequence of both the 3' and 5' ends of the genomic RNA of NDV strain LaSota by using 3' and 5' RACE procedures (rapid amplification of cDNA ends). The 5' end was recovered by means of PCR after ligation of a single-stranded anchorprimer (ALG3) to single-stranded cDNA which was generated by reverse transcription of the 5' end of the genomic RNA. By using a primer (ALG4) that is complementary to the anchorprimer and an NDV-specific primer, PCR products were generated which contained the 5' end. To clone the 3' end of NDV, the single-stranded anchorprimer ALG3 was ligated to the 3' end of viral RNA by using T4 RNA ligase and amplified by means of PCR by using primer ALG4 and an NDV-specific primer (method I). Alternatively, the 3' and 5' ends of the NDV RNA were ligated to each other by using T4 RNA ligase and the resulting concatenated RNA was used for RT-PCR by using NDV-specific primers that flanked the ligation point (method II). The 3' and 5' RACE products were cloned in T-vector pBluescriptII-TSK (Ichihara and Kurosawa, 1993) or in pGEM4Z and several independent clones were isolated and sequenced. The results are compiled in Table 2. To enable the direct comparison of the 3' and 5' terminal ends, the sequences are shown in the form of DNA and the 3' end of the genomic strand is represented as the 5' end of the antigenomic strand. At the genomic RNA level the sequence of

the 3'-end reads 3'-UGGUUUGUCUCUUAG whereas the sequence of the 5'-end reads UUUAGAAACAAACCA-5'. The sequence of the 3' end is almost similar to the published 3' terminal sequence of NDV strain D26 (Ishida et al., 1986). However, the
5 sequence of the 5' end showed that NDV strain LaSota contains 64 additional nucleotides in comparison with the published sequence of the L-gene of NDV strain Beaudette C (Yusoff et al., 1987). (Figure 6.)

10 Replication of NDV minigenomes by helpervirus

To determine whether the 3' and 5' ends of NDV are functional in replication and transcription, minigenomes were constructed which consisted of the 3' end of NDV (nt 1-119),
15 a reporter gene encoding secreted alkaline phosphatase (SEAP), and the 5' end of NDV (nt 14973-15186) (Fig. 2). These minigenomes were cloned in both orientations in transcription vector pOLTV5, generating plasmids pOLTV535 and pOLTV553, respectively (for details of the construction see
20 Materials and Methods). Plasmid pOLTV5 (Fig. 1) contains the T7 RNA polymerase promoter followed by unique StuI and SmaI restriction sites, the autocatalytic ribozyme from hepatitis delta virus (HDV) and the transcription termination signal from bacteriophage T7 (Pattnaik et al., 1992). In vivo or in
25 vitro transcription using T7 RNA polymerase of plasmid pOLTV535 gives rise to antigenomic RNA (or [+]-RNA), whereas transcription of plasmid pOLTV553 gives rise to genomic RNA (or [-]-RNA) (Fig. 5).

To examine whether the minigenome RNA's generated by plasmids
30 pOLTV535 and pOLTV553 could be replicated and expressed by using NDV as helpervirus, we used CER cells which expressed T7 RNA polymerase either constitutively (CER-C9 cells, see Materials and Methods), or after infection with fowlpox recombinant fpEFLT7pol (Britton et al., 1995; hereafter
35 called FPV-T7) that expresses T7 RNA polymerase. CER-C9 cells and FPV-T7 infected CER cells were transfected with the minigenome plasmids pOLTV535 or pOLTV553 and after incubation

for 3h at 37°C the cells were either infected with NDV for 1h, or left uninfected. Approximately 24h after transfection, a sample was taken from the medium and assayed for SEAP activity. The results showed that SEAP expression was very high in FPV-T7 infected cells which were transfected with pOLTV535. This is not surprising since transcription by T7 RNA polymerase generates antigenomic [+] -RNA which is capped by fowlpox enzymes and which is efficiently translated by the host cell. In cells transfected with pOLTV553, transcription by T7 RNA polymerase generates genomic [-] -RNA which must be converted into [+] -RNA by helpervirus in order to be translated into SEAP protein (cf. Fig. 5). In both cases, no increase in SEAP expression could be observed in NDV infected cells in comparison to non-infected cells. On the contrary, SEAP-expression in NDV infected cells was consistently approximately two times lower than in uninfected cells (results not shown). For pOLTV535-transfected cells this may be explained by the already very high level of SEAP expression by transcripts generated by T7 RNA polymerase. However, in pOLTV553-transfected cells, where efficient expression of SEAP is dependent on the conversion of genomic [-] -RNA into antigenomic [+] -RNA or mRNA by the viral polymerase complex, we would have expected an increase in SEAP expression after NDV infection. We could think of two reasons why the minigenomes could not be expressed and replicated by NDV. First, the size of the minigenome RNA's does not conform to the so-called "rule-of-six" (Calain and Roux, 1993; Kolakofsky et al., 1998). According to this rule, paramyxovirus genomes are only replicated efficiently when they are a multiple 6 nt in length. Second, the two extra G residues which are present at the 5' end of the minigenome RNA's might interfere with correct replication and/or transcription by the viral polymerase complex. To find out whether replication of the genomes was dependent on the rule-of-six, we inserted a series of short self-complementary oligonucleotides which increased 1 nt in size in the unique ClaI-site in plasmids

pOLTV535 and pOLTV553 (Fig. 2). The resulting plasmids (pOLTV535N0 to -N5 and pOLTV553N0 to -N5) differ in size from 1 to 6 nt and therefore one of them should generate a minigenome RNA which conforms to the rule-of-six. The plasmids were used to transfect CER cells or FPV-T7 infected CER-C9 cells as described above. The results showed that only plasmids pOLTV535N3 and pOLTV553N3 gave rise to an enhanced SEAP activity after NDV infection. The length of the minigenome RNA's generated from these plasmids by T7 RNA polymerase were calculated to be $6n+2$. Since two extra G residues are present at the 5' end of the minigenome RNA's, these results suggest that only the size of the RNA sequence which is located between the authentic 3' and 5' ends of the minigenome RNA's is relevant for the rule-of-six. This was verified by constructing minigenome plasmids in which the transcription start of T7 RNA polymerase was changed so that the first nucleotide which was incorporated into RNA was the first nucleotide of the 3' or 5' end of NDV (see Materials and Methods). Transfection of these plasmids indicated that only minigenome RNA's generated by plasmids pOLTV735N3 and pOLTV753N3 are replicated by helpervirus (results not shown). These findings again indicate that replication of NDV is strictly dependent on the rule-of-six. Furthermore, these findings indicate that the presence of two extra G residues at the 5'-end of the minigenome RNA's does not interfere with correct replication. Similar results have been obtained with minigenome plasmids (or DI plasmids) from other paramyxoviridae (Pattnaik et al., 1992; Harty and Palese, 1995).

30

Packaging of NDV minigenomes by helpervirus

To determine whether minigenome RNA's could be packaged by NDV helpervirus, the medium of the transfected cells was transferred to fresh monolayers and after 1h of adsorption, the monolayers were washed three times with PBS and further incubated in complete medium. After 24h of incubation, the

35

SEAP activity in the medium was measured. The results showed that SEAP activity was present only in cells which had been treated with the medium from cells transfected with minigenome plasmid pOLTV553N3 (Table 4). This finding indicates that minigenome RNA's can be packaged into NDV envelopes and that these particles are able to infect cells. Furthermore, these results show that packaging is dependent on replication which indicates that only RNA molecules which are complexed with the viral NP, P and L proteins are packaged into virus-like particles.

Replication of NDV minigenomes by plasmids expressing the NP, P, and L proteins

To determine whether the minigenome RNA's could also be replicated by plasmids encoding the essential NP, P, and L proteins, we performed co-transfection experiments in cells infected with FPV-T7. Cells were transfected with a combination of plasmids consisting of the minigenome plasmid and plasmids pCIneoNP, -P, and -L(c), respectively. As a negative control, pCIneoL(c), which encodes the essential L protein, was replaced by the vector plasmid pCIneo. The results (Table 5) indicated that indeed plasmids encoding NP, P, and L are able to replicate minigenome RNA's. The results furthermore show that, similar to minigenome replication by helpervirus, also replication by the NP, P, and L proteins is dependent on the rule-of-six.

Nucleotide sequence of the complete genome of NDV strain

LaSota

Sub-genomic cDNA fragments spanning the entire NDV genome were constructed by means of RT-PCR (Fig. 4). To keep the number of PCR errors to a minimum, a proofreading enzyme-mix (Expand Long Template; Boehringer Mannheim) was used in combination with a limited number of PCR cycles (15 cycles). Primer 3UIT which is complementary to the 3' end of NDV RNA

was used for reverse transcription, and gene-specific primers were used for PCR. To identify possible PCR errors, three

independent RT reactions were performed and used to generate
5 three independent sets of subgenomic cDNA's. The cDNA's,
which varied in size from approximately 4 to 7 kb, were
cloned in pGEM-T. The nucleotide sequence of two sets of
cDNA's was determined by using primers which were either
deduced from published NDV sequences, or by primers derived
10 from the NDV sequence that was deduced during this sequencing
project (Table 1). Remaining ambiguities were resolved by
sequencing the relevant regions of the third set of cDNA
clones. The genome of NDV strain LaSota consists of 15186 nt
(Fig.--3), which makes it the smallest of all paramyxovirus
15 genomes from which the entire sequence has been established
to date (Kolakofsky et al., 1998).

Construction of a full-length NDV cDNA clone in transcription
plasmid pOLTV5

20

To construct a full-length cDNA clone of NDV strain
LaSota, overlapping cDNA clones spanning the entire NDV
genome were joined at shared restriction sites according to
the strategy shown in Fig. 4. The entire NDV cDNA was
25 assembled in the minigenome plasmid pOLTV535 (see above) which
is derived from transcription plasmid pOLTV5.
As can be seen in Fig. 4B, the last step in the assembly of
the complete NDV cDNA was the cloning of an approximately 8.8
kb ClaI (nt 3521-12355) fragment from pGEM-B into p535-DI
30 which contained the NDV sequences flanking the ClaI site at
either side (i.e., nt 1-3521 and 12355-15186, respectively).
This step proved to be quite difficult since we repeatedly
failed in generating the correct clones. Therefore, the ClaI
fragment of pGEM-B was tagged with the chloramphenicol-
35 resistance (Cm) gene from plasmid pACYC184. The ClaI fragment
harboring the Cm-gene was isolated and cloned in the ClaI-
site of p535-DI and transformants were selected for

resistance against both Cm. Since transformants grew poorly, the antibiotic selection was reduced to 15 µg/ml Cm and 10 µg/ml Km and the incubation temperature was reduced from 37°C to 32°C. Finally, the Cm-gene was removed from this plasmid by digestion with BsiWI followed by recircularization by using T4 DNA ligase. After transformation of E.coli, cells harboring the desired plasmid were identified phenotypically by screening for Km-resistance and Cm-sensitivity. The resulting plasmid which consisted of the full-length NDV cDNA cloned between the SmaI and StuI sites of transcription plasmid pOLTV5 was designated pNDFL+.

Generation of infectious NDV from full-length cDNA

To generate infectious NDV entirely from cloned cDNA, plasmid pNDFL+ was used in co-transfection experiments with pCIneoNP, -P, and -L(c), as described above for the minigenome plasmids. Transfection of CER and CEF cells was monitored by using minigenome plasmid pOLTV553N3 and by measuring SEAP expression. As a negative control, pCIneoL(c) was replaced by pCIneo. After co-transfection, the cells were incubated for 3 to 6 days in medium containing 5% allantoic fluid. The addition of allantoic fluid is necessary because CER or CEF cells lack the appropriate proteases which are required to cleave the F protein of NDV strain LaSota. Cleavage of the F protein is absolutely required for cell-to-cell spread and for the generation of infectious virus. After 3 days of incubation, we performed an immunological staining of the fixed monolayers by using a monoclonal antibody against the F protein. The results showed that cells that were stained with the antibody were only present in monolayers which had been co-transfected with pNDFL(+), pCIneoNP, -P, and L(c). These results indicated that genome replication and expression was occurring in these cells. No staining cells were observed when pCIneoL(c) was replaced by pCIneo in the co-transfection experiments.

To recover infectious virus, the supernatant of transfected CEF monolayers was injected into the allantoic cavity of embryonated eggs. Four days later the allantoic fluid was

5 harvested, analyzed in a haemagglutination assay, and passaged further in eggs. The results showed that only the supernatant of cells transfected with a combination of pNDFL+ and pCIneoNP, -P, and -L(c) yielded a positive reaction in the haemagglutination assay. Allantoic fluid which showed a
10 positive haemagglutination reaction was subsequently analyzed in a haemagglutination inhibition assay by using monoclonal antibodies 7B7, 8C11, 5A1, 7D4, and 4D6 (Long et al., 1986) which can be used to differentiate between different NDV strains. The results of this assay indicated that the NDV
15 strain which was recovered from the inoculated eggs showed the same reactivity as the original LaSota strain. The virus which was recovered from the inoculated eggs was designated NDFL to distinguish it from the original LaSota strain.

20 Generation of genetically modified NDV from full-length cDNA

To show unambiguously that the co-transfection system could be used to recover infectious virus from cloned full-length NDV cDNA, a genetic tag was introduced in plasmid
25 pNDFL(+). To this end, the amino acid sequence of the protease cleavage site in the Fo protein was changed from that of the LaSota strain (GGRQGR | L) to the consensus sequence of virulent NDV strains (GRRQRR | F) by means of PCR mutagenesis (for details see Materials and Methods). The
30 resulting plasmid, pNDFL+[F^{wt}], was used to generate virus by using the co-transfection system described above. Infectious virus, designated NDFL[F^{wt}], was recovered from the allantoic fluid of embryonated eggs which had been inoculated with the medium of co-transfected CEF cells. In an HI test,
35 all MAb's including 7D4, which is specific for the LaSota strain, showed the same reactivity with the newly generated virus as with the original LaSota strain. The nucleotide

sequence of the region encoding the protease cleavage site of the F protein was determined by means of RT-PCR. The results showed that the nucleotide sequence contained the exact nucleotide changes which were introduced in the mutagenic primer which was used to modify the original LaSota sequence. This finding shows that the virus was derived from plasmid pNDFL+[F^{wt}] and demonstrates that (genetically modified) NDV can be generated entirely from cloned full-length NDV cDNA.

10 The protease cleavage site of the Fo protein of NDV is a key determinant for virulence

It is generally assumed that the amino acid sequence of the protease cleavage site of the Fo protein is a key determinant for virulence of different NDV strains. The generation of a genetically modified LaSota strain in which the amino acid sequence of the protease cleavage site was changed from a lentogenic (non-virulent) to that of a velogenic (virulent) NDV strain offered the unique opportunity to test this assumption. Therefore, we determined the intracerebral pathogenicity index (ICPI) of the newly generated virus NDFL[F^{wt}] and compared it with that of strain NDFL and of the original LaSota strain (clone E13-1). The results showed that the ICPI of strain NDFL[F^{wt}] was 1.3 which is far above the value for strains NDFL (ICPI=0.0) and clone E13-1 (ICPI=0.3). These results show that, as expected, the virulence of NDV is largely determined by the amino acid sequence of the protease cleavage site of the Fo protein.

30 Introduction of serological marker

The envelope glycoproteins F and HN of NDV are the most immunogenic proteins of the virus. After infection, both the F and HN protein elicit a strong neutralizing antibody response. The induction of such a neutralizing antibody response is the basis of successful vaccination by non-virulent NDV strains (such as the widely used LaSota strain).

However, the antibody response against NDV vaccine strains cannot be distinguished from the antibody response against virulent NDV field strains. Thus, infections with virulent

5 field virus cannot be traced by serological methods. This situation is undesirable since field virus infections are masked by vaccination and clinical signs which are caused by field strains may be overlooked or are even attributed to the vaccine. Since successful differentiation between vaccination
10 and infection is essential for eradication of NDV, we set out to develop genetically modified NDV strains which can be used for vaccination and which can be serologically distinguished from NDV field strains (so called marker vaccines).

In order to develop an NDV marker vaccine, the virus has to
15 be genetically modified such that one or several immunodominant epitopes of one of the (major) antigens are either deleted or modified. Deletion of part(s) of an essential protein may lead to the loss of the biological function of that protein. Therefore, we chose to modify one
20 of the immunodominant envelope proteins of NDV in such a way that the biological function of the protein was retained whereas the antibody repertoire against the modified protein differed from that against the original protein. For reasons specified below, we chose for one embodiment of the invention
25 to modify the HN protein of NDV. Infection of NDV is initiated by fusion of the virion envelope with the plasma membrane of the host cell. For this process, both the F protein and the HN protein are required. It has been shown that the F and HN proteins physically interact and that this
30 interaction is required for membrane fusion (Deng et al., 1995). Furthermore, it has been shown that the interaction is type specific, i.e., the F and HN proteins must be derived from the same virus in order to shown fusion activity. The interacting domain of the HN protein of NDV has been
35 localized to the so-called stalk- or stem-region of the protein, comprising the first 92 amino acid residues of the ectodomain of the HN protein (Deng et al., 1995). Hybrid HN

proteins consisting of aa 1-141 of NDV and aa 141-572 of human parainfluenza virus type-3 (hPIV3) were shown to retain fusion activity when co-expressed with the NDV F protein. These finding suggests that genetically modified NDV strains which harbour a hybrid HN protein which consists of the stem region of NDV followed by the globular head of the HN protein of a different avian paramyxovirus serotype may be viable. Furthermore, such strains would elicit an anti-HN antibody response which is different from that of NDV. Since the neutralizing antibody response against the F protein is sufficient to allow efficient protection against challenge virus infection, such genetically modified NDV strains meet the two essential requirements of a marker vaccine, i.e., protection against disease and serological differentiation.

Hybrid HN genes were constructed which consisted of a fusion of either aa 1-141 of NDV and aa 142-580 of avian paramyxovirus type-2 (APMV2) (designated HN1/2¹⁴¹) or aa 1-143 of NDV and aa 144-580 of APMV2 (designated HN1/2¹⁴³). Similary, hybrid HN genes were constructed which consisted either of aa 1-141 of NDV and aa 143-569 of AMPV4 (designated HN1/4¹⁴¹) or aa 1-143 of NDV and aa 145-569 of APMV4 (designated HN1/4¹⁴³). The hybrid genes were cloned in the eukaryotic expression vector pCIneo and used in co-transfection experiments with a plasmid harboring the NDV F protein. To this end, the F protein was modified such that the amino acid sequence of the proteolytic cleavage site between F2 and F1 was changed from the LaSota sequence to that of the consensus sequence of virulent NDV strains (F^{wt}, see Materials and Methods section). Co-transfection experiments in CER cells and QM5 cells indicated that both HN1/2¹⁴¹ and HN1/2¹⁴³ as well as HN1/4¹⁴¹ and HN1/4¹⁴³ induced cell fusion when co-expressed with the F^{wt} protein. These results indicated that the complexes between the hybrid HN proteins and the F protein were biologically active. The hybrid HN proteins HN1/2¹⁴³ and HN1/4¹⁴³ were used to replace the original HN gene in the full-length cDNA clone pNDFL+, yielding pNDFL-HN1/2¹⁴³ and pNDFL-HN1/4¹⁴³. The latter two

plasmids were subsequently used for the generation of infectious virus by using the co-transfection system described above. Viable recombinant viruses (designated NDFL-HN1/2¹⁴³ and NDFL-HN1/4¹⁴³) could be isolated from the
5 allantoic fluid of embryonated eggs which had been inoculated with the supernatant of transfected monolayers.

The presence of the hybrid HN gene in each of two recombinants was verified by means of RT-PCR. Haemagglutination inhibition tests showed that monoclonal
10 antibodies and polyvalent antisera against NDV were unable to inhibit agglutination of chicken erythrocytes by the recombinant viruses NDFL-HN1/2¹⁴³ and NDFL-HN1/4¹⁴³. These results indicate that strains NDFL-HN1/2¹⁴³ and NDFL-HN1/4¹⁴³ may be used as vaccines that can be serologically
15 distinguished from classical NDV vaccines.

Expression of a heterologous protein from recombinant NDV

To examine whether foreign genes can be inserted into
20 the NDV genome, we constructed a recombinant virus that carried the SEAP reporter gene. The SEAP gene was derived from plasmid pOLTV535 and was modified to include the typical transcriptional stop and start boxes of NDV. A DNA fragment containing the SEAP gene followed by the transcriptional stop
25 and start boxes was inserted into the XmnI site (nt 109) in plasmid pNDFL+[F^{wt}]. Infectious virus, designated NDFL-AP, was generated by means of the co-transfection system, and the presence of the SEAP gene was verified by means of RT-PCR. Cells infected with strain NDFL-AP expressed very high levels
30 of the SEAP protein. By using the specific activity of the SEAP protein, we calculated that x% of the proteins expressed in cells infected with NDFL-AP consisted of SEAP protein. These results show that heterologous genes can be expressed to very high levels from recombinant NDV.

Generation of an NDV deletion mutant on a trans-complementing cell line

In order to abrogate expression of the M protein of NDV,
5 a large part of the M gene was deleted by digestion of
pNDFL+[F^{wt}] with BsaAI (nt 3087) followed by partial digestion
with HindIII (nt 4252). After filling in the HindIII end with
Klenow DNA polymerase, the fragment was recircularized by
10 using T4 DNA ligase and used to transform E. coli. The
resulting plasmid, designated pNDFL+[F^{wt}]dM, was used to
generate virus by means of the co-transfection system in
trans-complementing CER-M cells that expressed the NDV M
protein. The supernatant of transfected monolayers was
15 passaged three times on CER-M cells and analyzed for the
presence of virus. Virus was obtained as evidenced by the
fact that the culture supernatant of the third passage
yielded positive results in hemagglutination (HA) and
hemagglutination-inhibition (HI) tests. The virus was
designated NDFL-dM. When NDFL-dM was used to infect
20 monolayers of CEF cells, the virus was still able to spread
by cell-to-cell transmission as seen in an IPMA by using a
monoclonal antibody against the F protein. As expected,
expression of the M protein could not be demonstrated in an
IPMA by using a monoclonal antibodies against the M protein.
25 When the supernatant was used to infect either CEF cells or
CER-M cells, we were unable to show the presence of
replicating virus in these monolayers by means of IPMA. This
finding indicates that infectious virus could not be
generated in non-complementing CEF cells. This finding was
30 confirmed by the observation that inoculation of embryonated
eggs with supernatant from infected CEF cells did not result
in the generation of progeny virus when tested in HA or HI
tests.

The need for better NDV vaccines, and especially the
35 need for NDV marker vaccines, prompted us to develop a
reverse genetics system which would allow the genetic
modification of NDV. In this document we describe the

generation of infectious NDV entirely from cloned full-length cDNA. We show that the virulence of NDV can be dramatically changed by modifying only 3 nucleotides which determine the specificity of the protease cleavage site of the F protein.

5 In this case the protease cleavage site was changed from that of the LaSota strain to that of the consensus cleavage site of virulent NDV strains. By generating this genetically modified NDV strain we deliver the formal proof that the cleavability of the F protein is the key determinant (but not
10 the only determinant) for virulence of NDV. By using the same reverse genetics approach, the cleavage site can be modified, at will, to any other amino acid sequence. This may lead to the generation of a series of NDV strains which display a spectrum of virulence levels.

15

in vivo

As already mentioned above, it has been shown that, besides the cleavability of the F and HN proteins, other viral
20 factors may contribute to pathogenicity. Alterations in transcription and translation can modulate growth and cell-to-cell spread of the virus and/or cytopathogenicity. The availability of an infectious cDNA of NDV allows for the systematic modification of sequences which are involved in
25 transcription and replication. This may lead to the design of new NDV vaccines which sport optimal immunogenicity to virtually non-existing virulence.

Safety is one of the most important properties of live vaccines. However, for many live vaccines, including NDV,
30 immunogenicity is often inversely related to virulence. Therefore, further attenuation of live vaccines without losing immunogenicity is one of the most desired alterations for which genetic modification could be used. In this respect it is worthwhile mentioning that it has been shown that
35 elimination of expression of the V protein of Sendai virus resulted in a markedly reduced *in vivo* pathogenicity for mice

(Kato et al., 1997). Similar to Sendai virus, NDV also generates a V protein by a mechanism known as RNA editing (Steward et al., 1993). It is predictable that elimination of expression of the V protein of NDV may also result in an attenuated phenotype *in vivo*.

Apart from changing the virulence of NDV, we show that it is possible to modify the antigenic make-up of NDV in such a way that strains can be generated which can be serologically discriminated from NDV field strains. These, so called,

marker vaccines are an invaluable tool to assess the prevalence of NDV in commercial flocks around the world.

Furthermore, the large-scale application of such marker vaccines may ultimately lead to the complete eradication of NDV by a process of intensive screening and stamping out of infected flocks. In this document we show that foreign genes can be inserted into the genome of NDV. These foreign genes can be expressed to very high levels in infected cells. This shows that NDV can be used as a vaccine vector for the expression of antigens from other (poultry) pathogens.

Several properties make NDV an ideal vaccine vector for vaccination against respiratory or intestinal diseases. 1) NDV can be easily cultured to very high titres in embryonated eggs. 2) Mass culture of NDV in embryonated eggs is relatively cheap. 3) NDV vaccines are relatively stable and can be simply administered by mass application methods such as addition to drinking water or by spraying or aerosol formation. 4) The natural route of infection of NDV is by the respiratory and/or intestinal tract which are also the major natural routes of infection of many other poultry pathogens. 5) NDV can induce local immunity despite the presence of circulating maternal antibody.

Finally, we show that viable NDV deletion mutants can be generated by using trans-complementing cell lines. An NDV deletion mutant was generated which is unable to express the matrix (M) protein which is involved in budding of NDV at the inner cell membrane. We show that a phenotypically

complemented NDV strain that is unable to express the M protein is still able to infect cells and spread by means of cell-to-cell transmission. However, the mutant virus is unable to generate infectious progeny on non-complementing
5 cells. This finding shows that phenotypically complemented NDV deletion mutants may be used as safe self-restricted vaccines which are unable to spread into the environment. Such a non-transmissible vaccine combines the most important advantage of live vaccines, i.e., efficacy, with the most important advantage of killed vaccines, i.e., safety.

LEGENDS

Fig.1.

5

Transcription vector pOLTV5 is a derivative of the transcription vector described by Pattnaik et al. (1992). See text for details of the construction. The plasmid contains the T7 DNA-dependent RNA polymerase promoter (shown in boldface) followed by unique StuI and SmaI restriction sites and the autocatalytic ribozyme from hepatitis delta virus (HDV). DNA fragments can be cloned between the StuI and SmaI sites and can be transcribed either in vitro or in vivo by using T7 RNA polymerase. The 5' end of the resulting transcripts contains two extra G-residues which are not encoded by the insert. Due to the action of the ribozyme, the 3' end of the transcripts exactly corresponds to the last nucleotide of the insert.

20 Fig. 2.

Structure of the minigenome plasmids pOLTV535 (Fig. 2A) and pOLTV553 (Fig. 2B). The minigenome plasmids are based on transcription plasmid pOLTV5 (cf. Fig. 1) and contain the 3'-region (nt 1-119) and 5'-region (nt 14970-15186) of NDV strain LaSota flanking the gene encoding secreted alkaline phosphatase (SEAP). Transcription of pOLTV535 by T7 RNA polymerase yields antigenomic RNA (or [+] -RNA) whereas transcription of pOLTV553 yields genomic RNA (or [-] -RNA). The start (S) and end (E) boxes, which are viral transcription initiation and termination signals, are indicated. The start codon of the SEAP gene is underlined. The sequences of the insertions (N0-N5) in the ClaI-site which generate minigenome plasmids which each differ 1 nt in length (pOLTV535N0-N5 and pOLTV553N0-N5, respectively) are also shown.

Fig. 3.

5 Nucleotide sequence of the genome of NDV strain LaSota and deduced amino acid sequence of the NDV genes. The sequence shown corresponds to the antigenomic strand and is shown in the 5' to 3' direction in the form of ssDNA. The sequence shown in this figure is that of the consensus
10 sequence which was determined by completely sequencing two independent sets of overlapping subgenomic cDNA's which span the entire NDV genome. Remaining ambiguities (probably as a result of PCR errors) were resolved by sequencing relevant regions of a third independent set of clones.

15 The sequence of the full length cDNA clone pNDFL+ which was assembled from overlapping subgenomic cDNA clones (see Fig. 4), differs from that of the consensus NDV sequence at the following positions (consensus sequence between parentheses): nt 1755, G (A); nt 3766, A (G); nt 5109, G (A); nt 6999, T
20 (C); nt 7056, G (A); nt 9337, G (A); nt 9486, A (T); nt 10195, T (C); nt 13075, A (G). These differences results in 3 amino acid changes (consensus sequence between parentheses): F protein, R¹⁸⁹ (Q); HN protein S²⁰⁰ (P) L-protein N³⁶⁹ (I).

25 Fig. 4.

(A) Overall strategy used for the assembly of the full-length NDV cDNA from subgenomic overlapping cDNA clones. The cDNA was assembled in plasmid pOLTV535 which already
30 contained the 3' and 5' ends of NDV strain LaSota (cf. Fig. 2). The resulting plasmid, designated pNDFL+, was used for the generation of infectious NDV.

(B) Detailed cloning procedure for the assembly of the full-length NDV cDNA from subgenomic overlapping cDNA clones. Cm
35 denotes the chloramphenicol-resistance gene which was temporarily introduced as a phenotypical tag (see text for details).

(C) Detailed cloning procedure for the generation of genetically modified full-length NDV cDNA. The modification consists of 3 nucleotide changes which were introduced in the F gene and which result in the modification of the amino acid sequence of the proteolytic cleavage site of the F protein (see text for details).

Fig. 5.

10

(A) pOLTV535-series.

Transcription by means of T7 RNA polymerase yields antigenomic RNA (or [+] -RNA) which can be directly translated into SEAP protein by the cell. After infection of cells by helpervirus (or after co-transfection of plasmids encoding NP, P, and L), the antigenomic RNA is used by the viral polymerase complex for the synthesis of genomic RNA (or [-] -RNA). The genomic RNA is then used by the viral polymerase complex for the synthesis of both mRNA (by using the specific transcription start [S] and end [E] boxes) and antigenomic RNA.

25

(B) pOLTV553-series.

Transcription by means of T7 RNA polymerase yields genomic RNA (or [-] -RNA) which cannot be translated into SEAP protein. After infection of cells by helpervirus (or after co-transfection of plasmids encoding NP, P, and L), the genomic RNA is used by the viral polymerase complex for the synthesis of both mRNA (by using the specific transcription start [S] and end [E] boxes) and antigenomic RNA.

30

Figure 6.

- Alignment of nucleic acid sequences 5' terminal ends of NDV LaSota and other paramyxoviruses are given as sequence comparison of NDV across the four members of the *Rubulavirus* genus, three members of the *Paramyxovirus* genus, and three members of the *Morbillivirus* genus. The sequences are presented from the L gene end box to the 5' end (3' - 5' cDNA).
- 10 NDV, new castle disease virus; hPIV2, human parainfluenza virus 2; MuV, mumps virus; SV5 and SV41, simmian virus 5 and 41 respectively; SeV, sendai virus; bPIV3, and hPIV3 bovine and human parainfluenza virus respectively; CDV, canine distemper virus; MeV measles virus
- 15 RPV, rinderpest virus. Nucleotide (nt) sequences of the entire genomes were obtained as follows (accession no.): NDV (AF077761); hPIV2 (X57559); MuV (AB000388); SV5 (AF052755); SV41 (X64275); bPIV3 (D84095); hPIV3 (Z11575); CDV (L13194); MeV (X16565); RPV (Z30697).

REFERENCES

- Alexander, D.J. (1993) Paramyxovirus infections. In
5 Virus infections of birds. McFerran, J.B. and McNulty,
M.S. (eds), pp 321-340, Elsevier Science Publishers
B.V., Amsterdam.
- Antin, P.B and Ordahl, C.P. (1991) Isolation and
characterization of an avian myogenic cell line. *Dev.*
10 *Biol.* 143: 111-121.
- Baron, M.D. and Barrett, T. (1997) Rescue of rinderpest
virus from cloned cDNA. *J. Virol.* 71: 1265-1271.
- Beach, J.R. (1944) The neutralization in vitro of avian
pneumoencephalitis virus by Newcastle disease immune
15 serum. *Science* 100: 361-362.
- Beard, C.W. and Hanson, R.P. (1984) Newcastle disease.
In M.S. Hofstad et al. (eds) *Disease of Poultry*, 8th
Ed., pp. 452-470. Iowa State University Press, Ames.
- Beaudette, F.R., Bivins, J.A. and Miller, B.R. (1949)
20 Newcastle disease immunization with live virus. *Cornell*
Vet. 39: 302-334.
- Boursnell, M.E.G., Green, P.F., Samson, A.C.R.,
Campbell, J.I.A., Deuter, A., Peters, R.W., Millar,
N.S., Emmerson, P.T. and Binns, M.M. (1990) A
25 recombinant fowlpox virus expressing the hemagglutinin-
neuraminidase gene of Newcastle disease virus (NDV)
protects chickens against challenge by NDV. *Virology*
178: 297-300.
- Britton, P., Green, P., Kottier, S., Mawditt, K.L.,
30 Penzes, Z., Cavanagh, D. and Skinner, M. (1996)
Expression of bacteriophage T7 RNA polymerase in avian
and mammalian cells by a recombinant fowlpox virus. *J.*
Gen. Virol. 77: 963-967.

- Calain, P. and Roux, L. (1993) The rule of six, a basic feature for efficient replication of Sendai virus defective interfering RNA. *J. Virol.* 67: 4822-4830.
- 5 Chambers, P., Millar, N.S., Bingham, R.W. and Emmerson, P.T. (1986) Molecular cloning of complementary DNA to Newcastle disease virus and nucleotide sequence analysis of the junction between the genes encoding the haemagglutinin-neuraminidase and the large protein. *J. Gen. Virol.* 67: 475-486
- 10 Chang, A.C.Y. and Cohen, S.N. (1978) Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. *J. Bacteriol.* 134: 1141-1156.
- 15 Cho, B.R. (1982) Cytopathic effects and focus formation by reticuloendotheliosis viruses in a quail fibroblast cell line. *Avian Diseases* 27: 261
- Collins, P.L., Hil, M.G., Camargo, E., Grosfeld, H., Chanock, R.M. and Murphy, B.R. (1995) Production of
- 20 infectious human respiratory syncytial virus from cloned cDNA confirms an essential role for transcription elongation factor from the 5' proximal open reading frame of the M2 mRNA in gene expression and provides a capability for vaccine development. *Proc. Natl. Acad. Sci. USA* 92: 11563-11567.
- 25 Conzelmann, K.-K. (1996) Genetic manipulation of non-segmented negative-strand RNA viruses. *J. Gen. Virol.* 77: 381-389.
- Cowen, B.S. and Braune, M.O. (1988) The propagation of
- 30 avian viruses in a continuous cell line (QT35) of Japanese quail origin. *Avian Diseases* 32: 282-297.
- Deng, R., Wang, Z., Mirza, A.M. and Iorio, R.M. (1995) Localization of a domain on the paramyxovirus attachment protein required for the promotion of cellular fusion by

its homologous fusion protein spike. *Virology* 209: 457-496.

Doyle, T.M. (1927) A hitherto unrecorded disease of fowls due to a filter-passing virus. *J. Comp. Pathol.*

5 *Ther.* 40: 144-169.

Garcin, D., Pelet, T., Calain, P., Roux, L., Curran, J. and Kolakofsky, D. (1995) A highly recombinogenic system for the recovery of infectious Sendai paramyxovirus from
10 cDNA: generation of a novel copy-back non-defective interfering virus. *EMBO J.* 14: 6087-6094.

Garten, W., Berk, W., nagai, Y., Rott, R. and Klenk, H.-D. (1980) Mutational changes of the protease
suceptibility of glycoprotein F of Newcastle disease
15 virus: Effects on pathogenicity. *J. Gen. Virol.* 50: 135-147.

Goldhaft, T.M. (1980) Historical note on the origin of the LaSota strain of Newcastle disease virus. *Avian Dis.*
24: 297-301.

20 Gough, R.E. and Alexander, D.J. (1973) The speed of resistance to challenge induced in chickens vaccinated by different routes with a B1 strain of live NDV. *Vet. Rec.* 92: 563-564.

Hanson, R.P. (1988) Heterogeneity within strains of
25 Newcastle disease virus: key to survival. In D.J. Alexander (ed.), *Newcastle Disease*, pp. 113-130. Kluwer Academic Publ., Boston.

Harty, R.N. and Palese, P. (1995) Mutations within noncoding terminal sequences of model RNA's of Sendai
30 virus: Influence on reporter gene expression. *J. Virol.* 69: 5128-5131.

Heckert, R.A., Riva, J., Cook, S., McMillen, J. and Schwartz, R.D. (1996) Onset of protective immunity in
chicks after vaccination with a recombinant herpesvirus
35 of turkeys vaccine expressing Newcastle disease virus

fusion and hemagglutinin-neuraminidase antigens.
Avian Dis. 40: 770-777.

- Heuschele, W.P. and Easterday, B.C. (1970) Local immunity and persistence of virus in the tracheas of
5 chickens following infection with Newcastle disease virus. II. Immunofluorescent and histopathological studies. *J. Inf. Dis.* 121: 497-504.

- Hitchner, S.B. and Johnson, E.P. (1948) A virus of low
10 virulence for immunizing fowls against Newcastle disease (avian pneumoencephalitis). *Vet. Med.* 43: 525-530.

Hoffman, M.A. and Banerjee, A.K. (1997) An infectious clone of human parainfluenza virus type 3. *J. Virol.* 71: 4272-4277.

- 15 Hofstad, M.S. (1953) Immunization of chickens against newcastle disease by formalin-inactivated vaccine. *Am. J. Vet. Res.* 14: 586-589.

- Ichihara, Y., and Kurosawa, Y. (1993) Construction of new T vectors for direct cloning of PCR products. *Gene*
20 130: 153-154.

- Ishida, N., Taira, H., Omata, T., Mizumoto, K., Hattori, S., Iwasaki, K. and Kawakita, M. (1986) Sequence of 2617 nucleotides from the 3' end of Newcastle disease virus genome RNA and the predicted amino acid sequence of the
25 viral NP protein. *Nucl. Acids Res.* 14: 6551-6564.

Kaleta, E.F. and Baldauf, C. (1988) Newcastle Disease in free-living and pet birds. In D.J. Alexander (ed.), *Newcastle Disease*, pp. 197-246. Kluwer Academic Publ., Boston.

- 30 Kant, A., Koch, G., van Roozelaar, D.J., balk, F. and ter Huurne, A. (1997) Differentiation of virulent and non-virulent strains of Newcastle disease virus within 24 hours by polymerase chain reaction. *Avian Pathol.* 26: 837-849.

- 35 Kolakofsky, D., Pelet, T., Garcin, D., Hausmann, S., Curran, J. and Roux, L. (1998) Paramyxovirus RNA

- synthesis and the requirement for hexamer genome length: the rule of six revisited. *J. Virol.* 72: 891-899.
- Kraneveld, F.C. (1926) A poultry disease in the Dutch East Indies. *Ned. Indisch Bl. Diergeneesk.* 38: 448-450.
- 5 Lamb, R.A. and Lolakofsky, D. (1996) *Paramyxoviridae: the viruses and their replication.* in: *Fundamental Virology* (Fields et al., eds), Chapter 20, p577-604, Lipincott-Raven Publishers, Philadelphia.
- 10 Lawson, N.D., Stillman, E.A., Whitt, M.A. and Rose, J.K. (1995) Recombinant vesicular stomatitis virus from DNA. *Proc. Natl. Acad. Sci. USA* 92: 4477-4481.
- Long, L., Portetelle, D, Ghysdael, J., Gonze, M., Burny, A. and Meulemans, G. (1986) Monoclonal antibodies to
- 15 haemagglutinin-neuraminidase and fusion glycoproteins of Newcastle disease virus: relationship between glycosylation and reactivity. *J. Virol.* 57: 1198-1202.
- Madansky, C.H. and Bratt, M.A. (1978) Noncytopathic mutants of Newcastle disease virus. *J. Virol.* 26: 724-729.
- 20 Madansky, C.H. and Bratt, M.A. (1981a) Noncytopathic mutants of Newcastle disease virus are defective in virus-specific RNA synthesis. *J. Virol.* 37: 317-327.
- Madansky, C.H. and Bratt, M.A. (1981b) Relationships among virus spread, cytopathogenicity, and virulence as
- 25 revealed by the noncytopathic mutants of Newcastle disease virus. *J. Virol.* 40: 691-702.
- Meulemans, G., Gonze, M., Carlier, M.C., Petit, P., Burny, A. and Long, L. (1986) Protective effects of HN and F glycoprotein-specific monoclonal antibodies on
- 30 experimental Newcastle disease. *Avian Pathol.* 15: 761-768.
- Millar, N.S., Chambers, P. and Emmerson, P.T. (1988) Nucleotide sequence of the fusion and haemagglutinin-neuraminidase gene of Newcastle disease virus, strain
- 35 Ulster: Molecular basis for variations in pathogenicity between strains. *J. Gen. Virol.* 69: 613-620.

- Morgan, R.W., Gelb Jr., J., Schreurs, C.S., Lütticken, D., Rosenberger, J.K. and Sondermeijer, P. (1992) Protection of chickens from Newcastle and Marek's diseases with a recombinant herpesvirus of turkeys vaccine expressing the Newcastle disease virus fusion protein. *Avian Dis.* 36: 858-870.
- Morgan, R.W., Gelb Jr., J., Pope, C.R. and Sondermeijer, P. (1993) Efficacy in chickens of a herpesvirus of turkeys recombinant vaccine containing the fusion gene of Newcastle disease virus: onset of protection and effect of maternal antibodies.
- Moscovici, C., Moscovici, M.G., Jimenez, H., Lai, M.M., Haymann, M.J. and Vogt, P.K. (1977) Continuous tissue culture cell lines derived from chemically induced tumors of Japanese quail. *Cell* 11: 95-103.
- Pattnaik, A.K., Ball, L.A., LeGrone, A.W. and Wertz, G.W. (1992) Infectious defective interfering particles of VSV from transcripts of a cDNA clone. *Cell* 69: 1011-1020.
- Peeples, M.E. (1988) Newcastle disease virus replication. In D.J. Alexander (ed.), *Newcastle Disease*, pp. 45-78. Kluwer Academic Publ., Boston.
- Peeters, B., N. de Wind, M. Hooisma, F. Wagenaar, A. Gielkens, and R. Moormann. (1992) Pseudorabies virus envelope glycoproteins gp50 and gII are essential for virus penetration, but only gII is involved in membrane fusion. *J. Virol.* 66: 894-905.
- Radecke, F., Spielhofer, P., Schneider, H., Kaelin, K., Huber, M., Dötsch, C., Christiansen, G. and Billeter, M.A. (1995) Rescue of measles virus from cloned DNA. *EMBO J.* 14: 5773-5784.
- Rott, R. and Klenk, H.-D. (1988) Molecular basis of infectivity and pathogenicity of Newcastle disease virus. In D.J. Alexander (ed.), *Newcastle Disease*, pp. 98-112. Kluwer Academic Publ., Boston.

- Russell, P.H., Griffiths, P.C., Goswami, K.K.A., Alexander, D.J., Cannon, M.J. and Russell, W.C. (1983) The characterization of monoclonal antibodies to Newcastle disease virus. *J. Gen. Virol.* 64: 2069-2072.
- 5 Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) Molecular cloning, a laboratory manual. Cold Spring Harbor Laboratories, Cold Spring Harbor, NY.
- Schneider, H., Spielhofer, P., Kaelin, K., Dötsch, C., Radecke, F., Sutter, G. and Billeter, M.A. (1997) Rescue of measles virus using a replication-deficient vaccinia-T7 vector. *J. Virol. meth.* 64: 57-64.
- 10 Schnell, M.J., Mebatsion, T. and Conzelmann, K.-K. (1994) Infectious rabies viruses from cloned cDNA. *EMBO J.* 13: 4195-4203.
- 15 Schütze, H., Enzmann, P.-J., Kuchling, R., Mundt., E., Niemann, H. and Mettenleiter, T.C. (1995) Complete genomic sequence of the fish rhabdovirus infectious haematopoietic necrosis virus. *J. Gen. Virol.* 76: 2519-2527.
- 20 Smith, A.L., Tignor, G.H., Mifune, K., and Motohashi, T. (1977) Isolation and assay of rabies serogroup viruses in CER cells. *Intervirology* 8: 92-99.
- Spradbrow, P.B. (1988) Geographical distribution. In D.J. Alexander (ed.), Newcastle Disease, pp. 247-255. Kluwer Academic Publ., Boston.
- 25 Staüber, N., Brechtbühl, K., Bruckner, L. and Hofmann, M.A. (1995) Detection of Newcastle disease virus in poultry vaccines using the polymerase chain reaction and direct sequencing of amplified DNA. *Vaccine* 13: 360-364
- 30 Steward, M., Vipond, I.B., Millar, N.S. and Emmerson, P.T. (1993) RNA editing in Newcastle disease virus. *J. Gen. Virol.* 74: 2539-2547.
- Taylor, J., Edbauer, C., Rey-Senelonge, A., Bouquet, J., Norton, E., Goebel, S., Desmettre, P. and Paoletti, E. (1990) Newcastle disease virus fusion protein expressed
- 35

in a fowlpox virus recombinant confers protection in chickens. *J. Virol.* 64: 1441-1450.

Tessier, D.C., Brousseau, R., and Vernet, T. (1986) Ligation of single-stranded oligodeoxyribonucleotides by T4 RNA ligase. *Anal. Biochem.* 158: 171-178.

Vieira, J., and Messing, J. (1991) New pUC-derived cloning vectors with different selectable markers and DNA replication origins. *Gene* 100: 189-194.

10 Vindevogel, H. and Duchatel, J.P. (1988) Panzootic Newcastle disease virus in pigeons. In D.J. Alexander (ed.), *Newcastle Disease*, pp. 184-196. Kluwer Academic Publ., Boston.

Whelan, S.P.J., Ball, L.A., Barr, J.N. and Wertz, G.T.W. 15 (1995) Efficient recovery of infectious vesicular stomatitis virus entirely from cDNA clones. *Proc. Natl. Acad. Sci. USA* 92: 8388-8392.

Wensvoort, G., Terpstra, C., Bonstra, J., Bloemraad, M. and Van Zaane, D. (1986) Production of monoclonal 20 antibodies against swine fever virus and their use in laboratory diagnosis. *Vet. Microbiol.* 12: 101-108.

Yusoff, K., Millar, N.S., Chambers, P., and Emmerson, P.T. (1987) Nucleotide sequence analysis of the L gene of Newcastle disease virus: homologies with Sendai and vesicular 25 stomatitis viruses. *Nucl. Acids Res.* 15: 3961-3976.

Table 1.

3'UIT	ACCAAACAGAGAATCCGTGAGTTA	1-24
P368+	GTGATGAGGAACCATGTTGC	368-387
P800+	GTCCGCATCTTCTTGTTAG	800-819
P1201+	GAGACTTGGAGTAGAGTACG	1201-1220
P1279+	AGCAGCAATGAAGGGCCTGG	1279-1298
P1356+	AAATCGGAGTCCTCACTGGG	1356-1375
P1683+	CTCTATATGACCACACCCTC	1664-1683
PRT1	CAAAGAATTGAGAAAAAGTACGG GTAGAAG	1785-1814
P2357+	GGAAACAGTCAGGAAAGACC	2358-2377
P2599+	TAAGTAAAGTTGACTATCAG	2599-2618
P2852+	GGCACTTAATAAACTTTCGC	2852-2871
P3496+	GAATGAAGAAGCCACTGTCTG	3496-3515
P3587+	CGGAGATCTTGTTGAGTTGG	3589-3608
P4267+	CATTATCCAAGCAGGTACCC	4270-4299
NDV5-F	ACGGGCTAGCGATTCTGGATCCCG GTTGG	4498-4526
P4731+ (LS)	AAGCTCCTCCCGAATCTGCC	4733-4752
P4958+	AGCTCTGATACAAGCCAAAC	4960-4979
P5266+ (LS)	CTGGTGGGAATATGGATTAC	5267-5286
P5591+ (LS)	AGTAACGTTCCCTATGTCCC	5593-5612
P5616+	GTATTTATTCTGCTTGAGC	5616-5635
P6000	AATACCCCTTGATCAGATGAGAGCC	6166-6190
NDV5-HN	GTAGGCTAGCAAGAGAGGCCGCC CTCAAT	6325-6354
P6693+ (L)	CATTGTTAAAACTGAGACC	6695-6714
P7110+ (L)	ATCGGAAGTCTTGCAGTGTG	7112-7131
P7501+ (L)	TGGTGGGAAACGCATCCAGC	7503-7522
P7900+ (LS)	AAGACTTAATCCTACGTCTG	7902-7921
P8590+	AACTCGGAAGGGCAGTACAC	8592-8611
L9000	TTTGTCACCTCTGAACCTTGTCATT	9008-9031
P9359+	CAATGATATAGCAGAATCCG	9361-9380

Table 1 continued

P9371+	GCAGAATCCGTGACTCATGC	9371-9411
P9390+	ATAGCTACTGTATTCTCTGG	9392-9411
P9686+	TCACACGATATCATGTTGAG	9686-9705
P9799+	CACACCCTAACGATAATTGG	9801-9820
P10198+	ATAAGAAACGTATCACTGAC	10200- 10219
P10601+	TTGTCGCGTTGCCTGTATGG	10603- 10622
P11006+	GCAGACATACTTTGACTCTG	11008- 11027
P11393+	TCCCTTATTGTCTGGAGTGC	11395- 11414
P11798+	TGATACGATAGAACTCGTAG	11800- 11819
L12000	CATATGTCGCCACATGTGAAGGCT	12008- 12031
P12373+	CAACCAGGACATATGATGAG	12375- 12394
P12796+	TCGACTGTTCTTACCAACTC	12798- 12817
P12978+	CACACCAACTTGCAGATACG	12978- 12997
P13236+	GAGTATCTACTGTCGGATGC	13238- 13257
P13601+	ATACTTGTTTCAGAGGAATAG	13603- 13622
P13943+	GACCTGACCTCAGATAAAGC	13946- 13965
P14002+	TATCATTGCTGCATTGTGAC	14004- 14023

Table 1 continued

P360	GGCGATGTAATCAGCCTAGTGCTT	14756- 14779
P14812+	ACTAAGGACATACTTGAAGC	14812- 14831
P230-	CCGGGACTTCTACTTTTAAG	230-211
P998-	TTTGATATCGCCTGAGAGG	998-979
P1898-	AAAGGTGGCCATGTTGTCC	1898-1879
P2617-	TGATAGTCAACTTTACTTAC	2617-2598
P3328-	GCAGAATCAAAGTACAGCCC	3330-3311
P3610-	CTTGCCAACTCAACAAGATC	3612-3593
P3990-	GATTAGCATAGTATCCACTG	3992-3973
NDV3-M	TCTCCCCGGGGCAGCTTATTTCTT AAAAGGAT	4400-4368
P4593-	GACAGATGCAACTCAGTACC	4625-4606
P4618- (LS)	ATGCAACTCAGTACCAGCGC	4620-4601
P5390-	GTAGAGTTACCTGTATACCC	5411-5392
NDV3-F	ACTACCCGGGAAACCTTCGTTCCT CAT	6238-6212
P6710- (LS)	TCTCAGTTTTTAACAATGCC	6712-6693
P7093- (LS)	GTTGATGGAACGCAGAGTAG	7095-7076
P7522- (LS)	CTGCTGGATGCGTTTCCCAC	7524-7505
P367	AGGGACCTCAATACTAGCCAGTTC	8692-8666
P9905-	CTCTATCAAGAGGCGATTAG	9907-9888
P10320-	TAAGACAGTACTTTTGCAGG	10322- 10303
P10684-	GATGCAACTGTGTCAACACC	10687- 10706
P11122-	AATTGGGCAGGAGTCAGAAC	11124- 11105
P11510-	TGCCTCCATGATAGCATGCG	11512- 11493

Table 1 continued

P11903-	ATTGCTTGGAAGATGGAACC	11905- 11886
P12717-	TGTCATACACTATTATGGCG	12719- 12700
P13141	CAAAGAGTACCGTGACAGACAGC ATAACC	13172- 13143
P13281-	GACATGATAGAGCTCACCTG	13302- 13283
P14101-	ACGGAATGCATGGCAATCAG	14163- 14144
P14522-	GCTCACCAAACCTCTCTGCAC	14524- 14505
P14687-	AGGATCTGTCTCGTGCACTG	14709- 14690
P377	TTTCCTTAAGTTTGGTAATACCTA GGAC	14888- 14861
P359	CACCAAGTCGACAATTGGCCAGAA AAGGAG	15046- 15017
5NDV	ACCAAACAAAGATTTGGTGAATGA CGA	15186- 15159

Table 2. Sequence of 3' and 5' terminal ends of the genome of NDV strain La Sota

5 A. Sequence of 3' terminal end (shown as 5' end of antigenomic DNA strand)

	method I.	clone	sequence
10		04	ACCAAACAGAGAATC
		05	ACCAAACAGAGAATC
		13	ACCAAACAGAGAATC
		21	ACCAAACAGAGAATC
15	method II.	clone	sequence
		26	ACCAAACAGAGAATC
		28	ACCAAACAGAGAATC
		30	ACCAAACAGAGAATC
20		31	GCCAAACAGAGAATC
		32	ACCAAACAGAGAATC
		33	ACCAAACAGAGAATC
		Consensus	<u>ACCAAACAGAGAATC</u>

25

B. Sequence of 5' terminal end (shown as DNA)

30	pBluescriptII-TSK clones	clone	sequence
		r3101-13	ACCAAACAAAGATTT
		r3101-14	ACCAAACAAAGATTT
		r3101-15	ACCAAACAAAGATTT
35		r2601-17	ACCAAACAAAGATTT
		r2601-18	ACCAAACAAAGATTT

(table 2 continued)

	r2601-19	ACCAAACAAAGATTT
5	r2601-20	AACAAGGTGAAGATA
	r2601-21	ACCAAACAAAGATTT
	pGEM4Z clones	clone sequence
10	r3101-16	ACCAAACAAAGATTT
	r3101-17	ACCAAACAAAGATTT
	r3101-18	ACCAAACAAAGATTT
	r3101-19	ACCAAACAAAGATTT
	r3101-22	ACCAAACAAAGATTT
15	Consensus	<u>ACCAAACAAAGATTT</u>

Table 3. Minigenome replication by NDV helpervirus

5 A. SEAP activity (cps) after transfection of CER-C9 cells
with the pOLTV535 and pOLTV553-series of plasmids.

	plasmid	+NDV	-NDV	ratio
	pOLTV535N0	3.5×10^4	7.1×10^4	0.49
10	pOLTV535N1	5.9	12.1	0.49
	pOLTV535N2	2.4	6.2	0.39
	pOLTV535N3	7.6	5.2	1.46
	pOLTV535N4	1.8	4.1	0.44
	pOLTV535N5	1.5	3.0	0.50
15				
	pOLTV553N0	5.5×10^3	9.6×10^3	0.57
	pOLTV553N1	9.6	27.6	0.35
	pOLTV553N2	2.4	3.5	0.68
	pOLTV553N3	15.1	9.5	1.59
20	pOLTV553N4	3.4	7.9	0.43
	pOLTV553N5	2.9	4.8	0.60

B. SEAP activity (cps) after transfection of FPV-T7 infected CER cells with the pOLTV553 -series of plasmids.

25	Plasmid	+NDV	-NDV	ratio
	pOLTV553N0	7.2×10^4	8.3×10^4	0.86
	pOLTV553N1	8.4	12.0	0.70
	pOLTV553N2	8.9	12.6	0.71
30	pOLTV553N3	27.4	8.6	3.19
	pOLTV553N4	9.7	10.4	0.93
	pOLTV553N5	8.5	8.1	1.05

Table 4.

Transfer of SEAP activity (cps) after treatment of CER cells with the supernatant of FPV-T7 infected CER cells which had been transfected with the pOLTV553-series of plasmids and which had been superinfected with NDV (see Table 3).

Plasmid		
10	pOLTV553N0	2.4×10^3
	pOLTV553N1	6.2
	pOLTV553N2	2.0
	pOLTV553N3	20.6
15	pOLTV553N4	2.0
	pOLTV553N5	2.1

Table 5.

SEAP activity (cps) after co-transfection of CER cells with the pOLTV553-series of plasmids and plasmids pCIneoNP, pCIneoP and pCIneoL(c) (or pCIneo as a negative control).

	Plasmid ratio	NP, P & L	NP, P & pCIneo	
10	pOLTV553N0	3.1×10^4	2.7×10^3	11.7
	pOLTV553N1	4.1	5.2	7.9
	pOLTV553N2	3.1	3.1	10.0
	pOLTV553N3	35.9	3.6	100.8
15	pOLTV553N4	1.9	4.6	4.1
	pOLTV553N5	1.0	4.1	2.5

CLAIMS

1. Avian-paramyxovirus cDNA at least comprising a nucleic acid sequence corresponding to the 5'-terminal end of the genome of avian-paramyxovirus allowing generating an infectious copy of avian-paramyxovirus.
- 5 2. A cDNA according to claim 1 comprising a full-length cDNA.
3. A cDNA at least comprising a nucleic acid sequence corresponding to the 5'-terminal end of the genome of avian-paramyxovirus allowing generating an replicating avian-paramyxovirus minigenome.
- 10 4. A cDNA according to claims 1, 2 or 3 at least partly derived from Newcastle Disease Virus.
5. A cDNA according to claim 4 wherein said Newcastle Disease Virus is a lentogenic virus, preferably derived from a vaccine strain.
- 15 6. A cDNA according to claim 5 wherein said vaccine strain is a LaSota strain ATCC VR-699.
7. A cDNA according to anyone of claims 1 to 6 additionally provided with a modification in a nucleic acid.
8. A cDNA according to claim 7 wherein said modification
- 20 comprises a nucleic acid encoding a modified protease cleavage site.
9. A cDNA according to claim 8 wherein said cleavage site is a protease cleavage site of the fusion (F) protein..
10. A cDNA according to claim 7 wherein said modification
- 25 comprises a nucleic acid encoding a hybrid viral protein.
11. A cDNA according to claim 10 wherein said protein is a hemagglutinin-neuraminidase (HN) protein.
12. A cDNA according to claim 7 wherein said modification comprises a deletion in a nucleic acid encoding a viral
- 30 protein.
13. A cDNA according to claim 12 wherein said viral protein is a matrix (M) protein.



14. A cDNA according to anyone of claims 1 to 13 additionally provided with a nucleic acid encoding an heterologous antigen.
15. A cDNA according to claim 14 wherein said antigen is derived from a poultry pathogen.
16. A cDNA according to claim 14 or 15 additionally provided with a nucleic acid encoding an immune-stimulatory protein or part thereof.
17. An RNA obtained from a cDNA according to anyone of claims 1 to 16.
18. A method for generating infectious copy avian-paramyxovirus comprising transfecting at least one cell with cDNA according to anyone of claims 1 to 16.
19. A method according to claim 18 wherein said cell is at least capable of expressing viral nucleocapsid (NP), phospho-(P) or large polymerase (L) protein.
20. A method according to claim 18 or 19 further comprising allowing cleavage of the fusion protein of said virus.
21. A method according to anyone of claims 18 to 20 further comprising incubating said cell in growth medium comprising proteolytic activity.
22. A method according to claim 21 wherein said growth medium comprises allantoic fluid comprising said proteolytic activity.
23. A method according to anyone of claims 18 to 22 wherein said cell is derived from a chicken cell.
24. An infectious copy avian-paramyxovirus obtainable by a method according to any one of claims 18 to 23.
25. A vaccine comprising a virus according to claim 24.
26. A live vaccine according to claim 25.
27. A vaccine according to claim 25 or 26 wherein said infectious copy avian-paramyxovirus is at least partly derived from a Newcastle Disease Virus (NDV).
28. A method for distinguishing unvaccinated animals or animals vaccinated with a NDV vaccine according to claim 27 from animals infected with wild-type NDV or vaccinated with



an unmodified mesogenic or lentogenic NDV strain comprising taking a least one sample from said animal and determining in said sample the presence of antibodies directed against an immunodominant epitope or marker expressed by said wild-type or unmodified NDV but not by said vaccine.

5

29. A method according to claim 28 wherein said antibodies are directed against the HN or F protein of NDV.

30. A method according to claim 28 or 29 wherein said animal is selected from the group composed of poultry, preferably of chickens.

10

31. A diagnostic kit for use in a method according to any one of claims 28 to 30.



1/22

Transcription-vector pOLTV5




Fig. 1

Fig. 3 Nucleotide sequence of NDV strain LaSota

```

1 accaaacagagaatccgtgagttacgataaaaaggcgaaggagcaattgaagtcgcacgggtagaaggtgtgaatctcgag 80
81 tgcgagcccgaagcacaaactcgagaaagccttctgccaac ATG TCT TCC GTA TTT GAT GAG TAC GAA 148
1 M S S V F D E Y E 9
149 CAG CTC CTC GCG GCT CAG ACT CGC CCC AAT GGA GCT CAT GGA GGG GGA GAA AAA GGG AGT 208
10 Q L L A A Q T R P N G A H G G G E K G S 29
209 ACC TTA AAA GTA GAC GTC CCG GTA TTC ACT CTT AAC AGT GAT GAC CCA GAA GAT AGA TGG 268
30 T L K V D V P V F T L N S D D P E D R W 49
269 AGC TTT GTG GTA TTC TGC CTC CGG ATT GCT GTT AGC GAA GAT GCC AAC AAA CCA CTC AGG 328
50 S F V V F C L R I A V S E D A N K P L R 69
329 CAA GGT GCT CTC ATA TCT CTT TTA TGC TCC CAC TCA CAG GTA ATG AGG AAC CAT GTT GCC 388
70 Q G A L I S L L C S H S Q V M R N H V A 89
389 ATT GCA GGG AAA CAG AAT GAA GCC ACA TTG GCC GTG CTT GAG ATT GAT GGC TTT GCC AAC 448
90 I A G K Q N E A T L A V L E I D G F A N 109
449 GGC ACG CCC CAG TTC AAC AAT AGG AGT GGA GTG TCT GAA GAG AGA GCA CAG AGA TTT GCG 508
110 G T P Q F N N R S G V S E E R A Q R F A 129
509 ATG ATA GCA GGA TCT CTC CCT CGG GCA TGC AGC AAC GGA ACC CCG TTC GTC ACA GCC GGG 568
130 M I A G S L P R A C S N G T P F V T A G 149
569 GCA GAA GAT GAT GCA CCA GAA GAC ATC ACC GAT ACC CTG GAG AGG ATC CTC TCT ATC CAG 628
150 A E D D A P E D I T D T L E R I L S I Q 169
629 GCT CAA GTA TGG GTC ACA GTA GCA AAA GCC ATG ACT GCG TAT GAG ACT GCA GAT GAG TCG 688
170 A Q V W V T V A K A M T A Y E T A D E S 189
689 GAA ACA AGG CGA ATC AAT AAG TAT ATG CAG CAA GGC AGG GTC CAA AAG AAA TAC ATC CTC 748
190 E T R R I N K Y M Q Q G R V Q K K Y I L 209
749 TAC CCC GTA TGC AGG AGC ACA ATC CAA CTC ACG ATC AGA CAG TCT CTT GCA GTC CGC ATC 808
210 Y P V C R S T I Q L T I R Q S L A V R I 229
809 TTT TTG GTT AGC GAG CTC AAG AGA GGC CGC AAC ACG GCA GGT GGT ACC TCT ACT TAT TAT 868
230 F L V S E L K R G R N T A G G T S T Y Y 249
869 AAC CTG GTA GGG GAC GTA GAC TCA TAC ATC AGG AAT ACC GGG CTT ACT GCA TTC TTC TTG 928
250 N L V G D V D S Y I R N T G L T A F F L 269
929 ACA CTC AAG TAC GGA ATC AAC ACC AAG ACA TCA GCC CTT GCA CTT AGT AGC CTC TCA GGC 988
270 T L K Y G I N T K T S A L A L S S L S G 289
989 GAC ATC CAG AAG ATG AAG CAG CTC ATG CGT TTG TAT CGG ATG AAA GGA GAT AAT GCG CCG 1048
290 D I Q K M K Q L M R L Y R M K G D N A P 309
1049 TAC ATG ACA TTA CTT GGT GAT AGT GAC CAG ATG AGC TTT GCG CCT GCC GAG TAT GCA CAA 1108
310 Y M T L L G D S D Q M S F A P A E Y A Q 329

```

4/22

Fig. 3 continued

1109 CTT TAC TCC TTT GCC ATG GGT ATG GCA TCA GTC CTA GAT AAA GGT ACT GGG AAA TAC CAA 1168
 330 L Y S F A M G M A S V L D K G T G K Y Q 349

1169 TTT GCC AGG GAC TTT ATG AGC ACA TCA TTC TGG AGA CTT GGA GTA GAG TAC GCT CAG GCT 1228
 350 F A R D F M S T S F W R L G V E Y A Q A 369

1229 CAG GGA AGT AGC ATT AAC GAG GAT ATG GCT GCC GAG CTA AAG CTA ACC CCA GCA GCA ATG 1288
 370 Q G S S I N E D M A A E L K L T P A A M 389

1289 AAG GGC CTG GCA GCT GCT GCC CAA CGG GTC TCC GAC GAT ACC AGC AGC ATA TAC ATG CCT 1348
 390 K G L A A A A Q R V S D D T S S I Y M P 409

1349 ACT CAA CAA GTC GGA GTC CTC ACT GGG CTT AGC GAG GGG GGG TCC CAA GCT CTA CAA GGC 1408
 410 T Q Q V G V L T G L S E G G S Q A L Q G 429

1409 GGA TCG AAT AGA TCG CAA GGG CAA CCA GAA GCC GGG GAT GGG GAG ACC CAA TTC CTG GAT 1468
 430 G S N R S Q G Q P E A G D G E T Q F L D 449

1469 CTG ATG AGA GCG GTA GCA AAT AGC ATG AGG GAG GCG CCA AAC TCT GCA CAG GGC ACT CCC 1528
 450 L M R A V A N S M R E A P N S A Q G T P 469

1529 CAA TCG GGG CCT CCC CCA ACT CCT GGG CCA TCC CAA GAT AAC GAC ACC GAC TGG GGG TAT 1588
 470 Q S G P P P T P G P S Q D N D T D W G Y 489

1589 TGA tggacaaaaccagcctgcttccacaaaacatcccaatgccctcaccgtagtcgaccctcgatttgcggtct 1667
 490 * 490

1668 atatgaccacaccctcaaaacaaacatccccctctttcctccctccccctgctgtacaactccgcacgcctagataccac 1747

1748 aggcacaatgcggctcactaacaatcaaaacagagccgaggggaattagaaaaagtagcgggtagaagagggatattcaga 1827

1828 gatcagggcaagtctcccgagtctctgctctctcctctacctgatagaccaggacaaac ATG GCC ACC TTT ACA 1901
 1 M A T F T 5

1902 GAT GCA GAG ATC GAC GAG CTA TTT GAG ACA AGT GGA ACT GTC ATT GAC AAC ATA ATT ACA 1961
 6 D A E I D E L F E T S G T V I D N I I T 25

1962 GCC CAG GGT AAA CCA GCA GAG ACT GTT GGA AGG AGT GCA ATC CCA CAA GGC AAG ACC AAG 2021
 26 A Q G K P A E T V G R S A I P Q G K T K 45

2022 GTG CTG AGC GCA GCA TGG GAG AAG CAT GGG AGC ATC CAG CCA CCG GCC AGT CAA GAC AAC 2081
 46 V L S A A W E K H G S I Q P P A S Q D N 65

2082 CCC GAT CGA CAG GAC AGA TCT GAC AAA CAA CCA TCC ACA CCC GAG CAA ACG ACC CCG CAT 2141
 66 P D R Q D R S D K Q P S T P E Q T T P H 85

2142 GAC AGC CCG CCG GCC ACA TCC GCC GAC CAG CCC CCC ACC CAG GCC ACA GAC GAA GCC GTC 2201
 86 D S P P A T S A D Q P P T Q A T D E A V 105

2202 GAC ACA CAG TTC AGG ACC GGA GCA AGC AAC TCT CTG CTG TTG ATG CTT GAC AAG CTC AGC 2261
 106 D T Q F R T G A S N S L L L M L D K L S 125

2262 AAT AAA TCG TCC AAT GCT AAA AAG GGC CCA TGG TCG AGC CCC CAA GAG GGG AAT CAC CAA 2321
 126 N K S S N A K K G P W S S P Q E G N H Q 145

2322 CGT CCG ACT CAA CAG CAG GGG AGT CAA CCC AGT CGC GGA AAC AGT CAG GAA AGA CCG CAG 2381
 146 R P T Q Q Q G S Q P S R G N S Q E R P Q 165

Fig. 3 continued

2382 AAC CAA GTC AAG GCC GCC CCT GGA AAC CAG GGC ACA GAC GTG AAC ACA GCA TAT CAT GGA 2441
 166 N Q V K A A P G N Q G T D V N T A Y H G 185

2442 CAA TGG GAG GAG TCA CAA CTA TCA GCT GGT GCA ACC CCT CAT GCT CTC CGA TCA AGG CAG 2501
 186 Q W E E S Q L S A G A T P H A L R S R Q 205

2502 AGC CAA GAC AAT ACC CTT GTA TCT GCG GAT CAT GTC CAG CCA CCT GTA GAC TTT GTG CAA 2561
 206 S Q D N T L V S A D H V Q P P V D F V Q 225

2562 GCG ATG ATG TCT ATG ATG GAG GCG ATA TCA CAG AGA GTA AGT AAG GTT GAC TAT CAG CTA 2621
 226 A M M S M M E A I S Q R V S K V D Y Q L 245

2622 GAT CTT GTC TTG AAA CAG ACA TCC TCC ATC CCT ATG ATG CGG TCC GAA ATC CAA CAG CTG 2681
 246 D L V L K Q T S S I P M M R S E I Q Q L 265

2682 AAA ACA TCT GTT GCA GTC ATG GAA GCC AAC TTG GGA ATG ATG AAG ATT CTG GAT CCC GGT 2741
 266 K T S V A V M E A N L G M M K I L D P G 285

2742 TGT GCC AAC ATT TCA TCT CTG AGT GAT CTA CGG GCA GTT GCC CGA TCT CAC CCG GTT TTA 2801
 286 C A N I S S L S D L R A V A R S H P V L 305

2802 GTT TCA GGC CCT GGA GAC CCC TCT CCC TAT GTG ACA CAA GGA GGC GAA ATG GCA CTT AAT 2861
 306 V S G P G D P S P Y V T Q G G E M A L N 325

2862 AAA CTT TCG CAA CCA GTG CCA CAT CCA TCT GAA TTG ATT AAA CCC GCC ACT GCA TGC GGG 2921
 326 K L S Q P V P H P S E L I K P A T A C G 345

2922 CCT GAT ATA GGA GTG GAA AAG GAC ACT GTC CGT GCA TTG ATC ATG TCA CGC CCA ATG CAC 2981
 346 P D I G V E K D T V R A L I M S R P M H 365

2982 CCG AGT TCT TCA GCC AAG CTC CTA AGC AAG TTA GAT GCA GCC GGG TCG ATC GAG GAA ATC 3041
 366 P S S S A K L L S K L D A A G S I E E I 385

3042 AGG AAA ATC AAG CGC CTT GCT CTA AAT GGC TAA ttactactgccacacgtagcgggtccctgtccactc 3110
 386 R K I K R L A L N G * 396

3111 ggcacacacggaatctgcacgagttccccccgcagacccaaggtccaactctccaagcggaatctctctcgcttc 3190

3191 ctcagccccactgaatggtcgcgtaaccgtaattaatctagctacatttaagattaagaaaaatacgggtagaattgga 3270

3271 gtgcccccaattgtgccaag ATG GAC TCA TCT AGG ACA ATT GGG CTG TAC TTT GAT TCT GCC CAT 3334
 1 M D S S R T I G L Y F D S A H 15

3335 TCT TCT AGC AAC CTG TTA GCA TTT CCG ATC GTC CTA CAA GGC ACA GGA GAT GGG AAG AAG 3394
 16 S S S N L L A F P I V L Q G T G D G K K 35

3395 CAA ATC GCC CCG CAA TAT AGG ATC CAG CGC CTT GAC TTG TGG ACT GAT AGT AAG GAG GAC 3454
 36 Q I A P Q Y R I Q R L D L W T D S K E D 55

3455 TCA GTA TTC ATC ACC ACC TAT GGA TTC ATC TTT CAA GTT GGG AAT GAA GAA GCC ACT GTC 3514
 56 S V F I T T Y G F I F Q V G N E E A T V 75

3515 GGC ATG ATC GAT GAT AAA CCC AAG CGC GAG TTA CTT TCC GCT GCG ATG CTC TGC CTA GGA 3574
 76 G M I D D K P K R E L L S A A M L C L G 95

3575 AGC GTC CCA AAT ACC GGA GAC CTT ATT GAG CTG GCA AGG GCC TGT CTC ACT ATG ATA GTC 3634
 96 S V P N T G D L I E L A R A C L T M I V 115

Fig. 3 continued

3635 ACA TGC AAG AAG AGT GCA ACT AAT ACT GAG AGA ATG GTT TTC TCA GTA GTG CAG GCA CCC 3694
 116 T C K K S A T N T E R M V F S V V Q A P 135

3695 CAA GTG CTG CAA AGC TGT AGG GTT GTG GCA AAC AAA TAC TCA TCA GTG AAT GCA GTC AAG 3754
 136 Q V L Q S C R V V A N K Y S S V N A V K 155

3755 CAC GTG AAA GCG CCA GAG AAG ATT CCC GGG AGT GGA ACC CTA GAA TAC AAG GTG AAC TTT 3814
 156 H V K A P E K I P G S G T L E Y K V N F 175

3815 GTC TCC TTG ACT GTG GTA CCG AAG AAG GAT GTC TAC AAG ATC CCA GCT GCA GTA TTG AAG 3874
 176 V S L T V V P K K D V Y K I P A A V L K 195

3875 GTT TCT GGC TCG AGT CTG TAC AAT CTT GCG CTC AAT GTC ACT ATT AAT GTG GAG GTA GAC 3934
 196 V S G S S L Y N L A L N V T I N V E V D 215

3935 CCG AGG AGT CCT TTG GTT AAA TCT TTG TCT AAG TCT GAC AGC GGA TAC TAT GCT AAC CTC 3994
 216 P R S P L V K S L S K S D S G Y Y A N L 235

3995 TTC TTG CAT ATT GGA CTT ATG ACC ACC GTA GAT AGG AAG GGG AAG AAA GTG ACA TTT GAC 4054
 236 F L H I G L M T T V D R K G K K V T F D 255

4055 AAG CTG GAA AAC AAT ATA AGG AGC CTT GAT CTA TCT GTC GGG CTC AGT GAT GTG CTC GGG 4114
 256 K L E K K I R S L D L S V G L S D V L G 275

4115 CCT TCC GTG TTG GTA AAA GCA AGA GGT GCA CCG ACT AAG CTT TTG GCA CCT TTC TTC TCT 4174
 276 P S V L V K A R G A R T K L L A P F F S 295

4175 AGC AGT GGG ACA GCC TGC TAT CCC ATA GCA AAT GCT TCT CCT CAG GTG GCC AAG ATA CTC 4234
 296 S S G T A C Y P I A N A S P Q V A K I L 315

4235 TGG AGT CAA ACC GCG TGC CTG CGG AGC GTT AAA ATC ATT ATC CAA GCA GGT ACC CAA CGC 4294
 316 W S Q T A C L R S V K I I I Q A G T Q R 335

4295 GCT GTC GCA GTG ACC GCC GAC CAC GAG GTT ACC TCT ACT AAG CTG GAG AAG GGG CAC ACC 4354
 336 A V A V T A D H E V T S T K L E K G H T 355

4355 CTT GCC AAA TAC AAT CCT TTT AAG AAA TAA gctgcgtctctgagattgcgctccgcccactcaccagat 4424
 356 L A K Y N P F K K * 365

4425 catcatgacacaaaaaactaatctgtcttgattatttacagttagtttacctgtctatcaagttagaaaaaacacgggta 4504

4505 gaagattctggatcccggttgccgcccctccaggtgcaag ATG GGC TCC AGA CCT TCT ACC AAG AAC CCA 4573
 1 M G S R P S T K N P 10

4574 GCA CCT ATG ATG CTG ACT ATC CGG GTT GCG CTG GTA CTG AGT TGC ATC TGT CCG GCA AAC 4633
 11 A P M M L T I R V A L V L S C I C P A N 30

4634 TCC ATT GAT GGC AGG CCT CTT GCA GCT GCA GGA ATT GTG GTT ACA GGA GAC AAA GCC GTC 4693
 31 S I D G R P L A A A G I V V T G D K A V 50

4694 AAC ATA TAC ACC TCA TCC CAG ACA GGA TCA ATC ATA GTT AAG CTC CTC CCG AAT CTG CCC 4753
 51 N I Y T S S Q T G S I I V K L L P N L P 70

4754 AAG GAT AAG GAG GCA TGT GCG AAA GCC CCC TTG GAT GCA TAC AAC AGG ACA TTG ACC ACT 4813
 71 K D K E A C A K A P L D A Y N R T L T T 90

4814 TTG CTC ACC CCC CTT GGT GAC TCT ATC CGT AGG ATA CAA GAG TCT GTG ACT ACA TCT GGA 4873
 91 L L T P L G D S I R R I Q E S V T T S G 110

Fig. 3 continued

4874 GGG GGG AGA CAG GGG CGC CTT ATA GGC GCC ATT ATT GGC GGT GTG GCT CTT GGG GTT GCA 4933
 111 G G R Q G R L I G A I I G G V A L G V A 130
 4934 ACT GCC GCA CAA ATA ACA GCG GCC GCA GCT CTG ATA CAA GCC AAA CAA AAT GCT GCC AAC 4993
 131 T A A Q I T A A A A L I Q A K Q N A A N 150
 4994 ATC CTC CGA CTT AAA GAG AGC ATT GCC GCA ACC AAT GAG GCT GTG CAT GAG GTC ACT GAC 5053
 151 I L R L K E S I A A T N E A V H E V T D 170
 5054 GGA TTA TCG CAA CTA GCA GTG GCA GTT GGG AAG ATG CAG CAG TTT GTT AAT GAC CAA TTT 5113
 171 G L S Q L A V A V G K M Q Q F V N D Q F 190
 5114 AAT AAA ACA GCT CAG GAA TTA GAC TGC ATC AAA ATT GCA CAG CAA GTT GGT GTA GAG CTC 5173
 191 N K T A Q E L D C I K I A Q Q V G V E L 210
 5174 AAC CTG TAC CTA ACC GAA TTG ACT ACA GTA TTC GGA CCA CAA ATC ACT TCA CCT GCT TTA 5233
 211 N L Y L T E L T T V F G P Q I T S P A L 230
 5234 AAC AAG CTG ACT ATT CAG GCA CTT TAC AAT CTA GCT GGT GGA AAT ATG GAT TAC TTA TTG 5293
 231 N K L T I Q A L Y N L A G G N M D Y L L 250
 5294 ACT AAG TTA GGT GTA GGG AAC AAT CAA CTC AGC TCA TTA ATC GGT AGC GGC TTA ATC ACC 5353
 251 T K L G V G N N Q L S S L I G S G L I T 270
 5354 GGT AAC CCT ATT CTA TAC GAC TCA CAG ACT CAA CTC TTG GGT ATA CAG GTA ACT CTA CCT 5413
 271 G N P I L Y D S Q T Q L L G I Q V T L P 290
 5414 TCA GTC GGG AAC CTA AAT AAT ATG CGT GCC ACC TAC TTG GAA ACC TTA TCC GTA AGC ACA 5473
 291 S V G N L N N M R A T Y L E T L S V S T 310
 5474 ACC AGG GGA TTT GCC TCG GCA CTT GTC CCC AAA GTG GTG ACA CAG GTC GGT TCT GTG ATA 5533
 311 T R G F A S A L V P K V V T Q V G S V I 330
 5534 GAA GAA CTT GAC ACC TCA TAC TGT ATA GAA ACT GAC TTA GAT TTA TAT TGT ACA AGA ATA 5593
 331 E E L D T S Y C I E T D L D L Y C T R I 350
 5594 GTA ACG TTC CCT ATG TCC CCT GGT ATT TAT TCC TGC TTG AGC GGC AAT ACG TCG GCC TGT 5653
 351 V T F P M S P G I Y S C L S G N T S A C 370
 5654 ATG TAC TCA AAG ACC GAA GGC GCA CTT ACT ACA CCA TAC ATG ACT ATC AAA GGT TCA GTC 5713
 371 M Y S K T E G A L T T P Y M T I K G S V 390
 5714 ATC GCC AAC TGC AAG ATG ACA ACA TGT AGA TGT GTA AAC CCC CCG GGT ATC ATA TCG CAA 5773
 391 I A N C K M T T C R C V N P P G I I S Q 410
 5774 AAC TAT GGA GAA GCC GTG TCT CTA ATA GAT AAA CAA TCA TGC AAT GTT TTA TCC TTA GGC 5833
 411 N Y G E A V S L I D K Q S C N V L S L G 430
 5834 GGG ATA ACT TTA AGG CTC AGT GGG GAA TTC GAT GTA ACT TAT CAG AAG AAT ATC TCA ATA 5893
 431 G I T L R L S G E F D V T Y Q K N I S I 450
 5894 CAA GAT TCT CAA GTA ATA ATA ACA GGC AAT CTT GAT ATC TCA ACT GAG CTT GGG AAT GTC 5953
 451 Q D S Q V I I T G N L D I S T E L G N V 470
 5954 AAC AAC TCG ATC AGT AAT GCT TTG AAT AAG TTA GAG GAA AGC AAC AGA AAA CTA GAC AAA 6013
 471 N N S I S N A L N K L E E S N R K L D K 490
 6014 GTC AAT GTC AAA CTG ACT AGC ACA TCT GCT CTC ATT ACC TAT ATC GTT TTG ACT ATC ATA 6073
 491 V N V K L T S T S A L I T Y I V L T I I 510

Fig. 3 continued

6074 TCT CTT GTT TTT GGT ATA CTT AGC CTG ATT CTA GCA TGC TAC CTA ATG TAC AAG CAA AAG 6133
 511 S L V F G I L S L I L A C Y L M Y K Q K 530

6134 GCG CAA CAA AAG ACC TTA TTA TGG CTT GGG AAT AAT ACT CTA GAT CAG ATG AGA GCC ACT 6193
 531 A Q Q K T L L W L G N N T L D Q M R A T 550

6194 ACA AAA ATG TGA acacagatgaggaacgaaggtttccctaataagtaatttgtgtgaaagttctggtagtctgtcag 6269
 551 T K M * 554

6270 ttcagagagttaagaaaaactaccggtttagatgaccaaaggacgatatacgggtagaacggtaagagagggccgcccc 6349

6350 tcaattgcgagccagggttcacaacctccgttctaccggttcaccgacaacagtcctcaatc ATG GAC CGC GCC 6423
 1 M D R A 4

6424 GTT AGC CAA GTT GCG TTA GAG AAT GAT GAA AGA GAG GCA AAA AAT ACA TGG CGC TTG ATA 6483
 5 V S Q V A L E N D E R E A K N T W R L I 24

6484 TTC CGG ATT GCA ATC TTA TTC TTA ACA GTA GTG ACC TTG GCT ATA TCT GTA GCC TCC CTT 6543
 25 F R I A I L F L T V V T L A I S V A S L 44

6544 TTA TAT AGC AAG GGG GCT AGC ACA CCT AGC GAT CTT GTA GGC ATA CCG ACT AGG ATT TCC 6603
 45 L Y S M G A S T P S D L V G I P T R I S 64

6604 AGG GCA GAA GAA AAG ATT ACA TCT ACA CTT GGT TCC AAT CAA GAT GTA GTA GAT AGG ATA 6663
 65 R A E E K I T S T L G S N Q D V V D R I 84

6664 TAT AAG CAA GTG GCC CTT GAG TCT CCG TTG GCA TTG TTA AAT ACT GAG ACC ACA ATT ATG 6723
 85 Y K Q V A L E S P L A L L N T E T T I M 104

6724 AAC GCA ATA ACA TCT CTC TCT TAT CAG ATT AAT GGA GCT GCA AAC AAC AGT GGG TGG GGG 6783
 105 N A I T S L S Y Q I N G A A N N S G W G 124

6784 GCA CCT ATC CAT GAC CCA GAT TAT ATA GGG GGG ATA GGC AAA GAA CTC ATT GTA GAT GAT 6843
 125 A P I H D P D Y I G G I G K E L I V D D 144

6844 GCT AGT GAT GTC ACA TCA TTC TAT CCC TCT GCA TTT CAA GAA CAT CTG AAT TTT ATC CCG 6903
 145 A S D V T S F Y P S A F Q E H L N F I P 164

6904 GCG CCT ACT ACA GGA TCA GGT TGC ACT CGA ATA CCC TCA TTT GAC ATG AGT GCT ACC CAT 6963
 165 A P T T G S G C T R I P S F D M S A T H 184,

6964 TAC TGC TAC ACC CAT AAT GTA ATA TTG TCT GGA TGC AGA GAT CAC TCA CAT TCA TAT CAG 7023
 185 Y C Y T H N V I L S G C R D H S H S Y Q 204

7024 TAT TTA GCA CTT GGT GTG CTC CGG ACA TCT GCA ACA GGG AGG GTA TTC TTT TCT ACT CTG 7083
 205 Y L A L G V L R T S A T G R V F F S T L 224

7084 CGT TCC ATC AAC CTG GAC GAC ACC CAA AAT CGG AAG TCT TGC AGT GTG AGT GCA ACT CCC 7143
 225 R S I N L D D T Q N R K S C S V S A T P 244

7144 CTG GGT TGT GAT ATG CTG TGC TCG AAA GTC ACG GAG ACA GAG GAA GAA GAT TAT AAC TCA 7203
 245 L G C D M L C S K V T E T E E E D Y N S 264

7204 GCT GTC CCT ACG CGG ATG GTA CAT GGG AGG TTA GGG TTC GAC GGC CAG TAC CAC GAA AAG 7263
 265 A V P T R M V H G R L G F D G Q Y H E K 284

7264 GAC CTA GAT GTC ACA ACA TTA TTC GGG GAC TGG GTG GCC AAC TAC CCA GGA GTA GGG GGT 7323
 285 D L D V T T L F G D W V A N Y P G V G G 304

Fig. 3 continued

7324 GGA TCT TTT ATT GAC AGC CGC GTA TGG TTC TCA GTC TAC GGA GGG TTA AAA CCC AAT TCA 7383
 305 G S F I D S R V W F S V Y G G L K P N S 324

7384 CCC AGT GAC ACT GTA CAG GAA GGG AAA TAT GTG ATA TAC AAG CGA TAC AAT GAC ACA TGC 7443
 325 P S D T V Q E G K Y V I Y K R Y N D T C 344

7444 CCA GAT GAG CAA GAC TAC CAG ATT CGA ATG GCC AAG TCT TCG TAT AAG CCT GGA CGG TTT 7503
 345 P D E Q D Y Q I R M A K S S Y K P G R F 364

7504 GGT GGG AAA CGC ATA CAG CAG GCT ATC TTA TCT ATC AAG GTG TCA ACA TCC TTA GGC GAA 7563
 365 G G K R I Q Q A I L S I K V S T S L G E 384

7564 GAC CCG GTA CTG ACT GTA CCG CCC AAC ACA GTC ACA CTC ATG GGG GCC GAA GGC AGA ATT 7623
 385 D P V L T V P P N T V T L M G A E G R I 404

7624 CTC ACA GTA GGG ACA TCT CAT TTC TTG TAT CAA CGA GGG TCA TCA TAC TTC TCT CCC GCG 7683
 405 L T V G T S H F L Y Q R G S S Y F S P A 424

7684 TTA TTA TAT CCT ATG ACA GTC AGC AAC AAA ACA GCC ACT CTT CAT AGT CCT TAT ACA TTC 7743
 425 L L Y P M T V S N K T A T L H S P Y T F 444

7744 AAT GCC TTC ACT CGG CCA GGT AGT ATC CCT TGC CAG GCT TCA GCA AGA TGC CCC AAC TCG 7803
 445 N A F T R P G S I P C Q A S A R C P N S 464

7804 TGT GTT ACT GGA GTC TAT ACA GAT CCA TAT CCC CTA ATC TTC TAT AGA AAC CAC ACC TTG 7863
 465 C V T G V Y T D P Y P L I F Y R N H T L 484

7864 CGA GGG GTA TTC GGG ACA ATG CTT GAT GGT GTA CAA GCA AGA CTT AAC CCT GCG TCT GCA 7923
 485 R G V F G T M L D G V Q A R L N P A S A 504

7924 GTA TTC GAT AGC ACA TCC CGC AGT CGC ATT ACT CGA GTG AGT TCA AGC AGT ACC AAA GCA 7983
 505 V F D S T S R S R I T R V S S S S T K A 524

7984 GCA TAC ACA ACA TCA ACT TGT TTT AAA GTG GTC AAG ACT AAT AAG ACC TAT TGT CTC AGC 8043
 525 A Y T T S T C F K V V K T N K T Y C L S 544

8044 ATT GCT GAA ATA TCT AAT ACT CTC TTC GGA GAA TTC AGA ATC GTC CCG TTA CTA GTT GAG 8103
 545 I A E I S N T L F G E F R I V P L L V E 564

8104 ATC CTC AAA GAT GAC GGG GTT AGA GAA GCC AGG TCT GGC TAG ttgagtcaattataaaggaggttg 8169,
 565 I L K D D G V R E A R S G * 578

8170 aaagatggcattgtatcacctatcttctgacatcaagaatcaaaccgaatgccggcgctgctcgaattccatgttgc 8249

8250 cagttgaccacaatcagccagtgtcatgcatcagattaagccttgtcattaatctcttgattaagaaaaaatgtaagt 8329

8330 ggcaatgagatacaaggcaaaacagctcatggttaataatacgggtaggac ATG GCG AGC TCC GGT CCT GAA 8401
 1 M A S S G P E 7

8402 AGG GCA GAG CAT CAG ATT ATC CTA CCA GAG CCA CAC CTG TCT TCA CCA TTG GTC AAG CAC 8461
 8 R A E H Q I I L P E P H L S S P L V K H 27

8462 AAA CTA CTC TAT TAC TGG AAA TTA ACT GGG CTA CCG CTT CCT GAT GAA TGT GAC TTC GAC 8521
 28 K L L Y Y W K L T G L P L P D E C D F D 47

8522 CAC CTC ATT CTC AGC CGA CAA TGG AAA AAA ATA CTT GAA TCG GCC TCT CCT GAT ACT GAG 8581
 48 H L I L S R Q W K K I L E S A S P D T E 67

Fig. 3 continued

8582	AGA	ATG	ATA	AAA	CTC	GGA	AGG	GCA	GTA	CAC	CAA	ACT	CTT	AAC	CAC	AAT	TCC	AGA	ATA	ACC	8641
68	R	M	I	K	L	G	R	A	V	H	Q	T	L	N	H	N	S	R	I	T	87
8642	GGA	GTG	CTC	CAC	CCC	AGG	TGT	TTA	GAA	CAA	CTG	GCT	AAT	ATT	GAG	GTC	CCA	GAT	TCA	ACC	8701
88	G	V	L	H	P	R	C	L	E	Q	L	A	N	I	E	V	P	D	S	T	107
8702	AAC	AAA	TTT	CGG	AAG	ATT	GAG	AAG	AAG	ATC	CAA	ATT	CAC	AAC	ACG	AGA	TAT	GGA	GAA	CTG	8761
108	N	K	F	R	K	I	E	K	K	I	Q	I	H	N	T	R	Y	G	E	L	127
8762	TTC	ACA	AGG	CTG	TGT	ACG	CAT	ATA	GAG	AAG	AAA	CTG	CTG	GGG	TCA	TCT	TGG	TCT	AAC	AAT	8821
128	F	T	R	L	C	T	H	I	E	K	K	L	L	G	S	S	W	S	N	N	147
8822	GTC	CCC	CGG	TCA	GAG	GAG	TTC	AGC	AGC	ATT	CGT	ACG	GAT	CCG	GCA	TTC	TGG	TTT	CAC	TCA	8881
148	V	P	R	S	E	E	F	S	S	I	R	T	D	P	A	F	W	F	H	S	167
8882	AAA	TGG	TCC	ACA	GCC	AAG	TTT	GCA	TGG	CTC	CAT	ATA	AAA	CAG	ATC	CAG	AGG	CAT	CTG	ATG	8941
168	K	W	S	T	A	K	F	A	W	L	H	I	K	Q	I	Q	R	H	L	M	187
8942	GTG	GCA	GCT	AAG	ACA	AGG	TCT	GCG	GCC	AAC	AAA	TTG	GTG	ATG	CTA	ACC	CAT	AAG	GTA	GGC	9001
188	V	A	A	K	T	R	S	A	A	N	K	L	V	M	L	T	H	K	V	G	207
9002	CAA	GTC	TTT	GTC	ACT	CCT	GAA	CTT	GTC	GTT	GTG	ACG	CAT	ACG	AAT	GAG	AAC	AAG	TTC	ACA	9061
208	Q	V	F	V	T	P	E	L	V	V	V	T	H	T	N	E	N	K	F	T	227
9062	TGT	CTT	ACC	CAG	GAA	CTT	GTA	TTG	ATG	TAT	GCA	GAT	ATG	ATG	GAG	GGC	AGA	GAT	ATG	GTC	9121
228	C	L	T	Q	E	L	V	L	M	Y	A	D	M	M	E	G	R	D	M	V	247
9122	AAC	ATA	ATA	TCA	ACC	ACG	GCG	GTG	CAT	CTC	AGA	AGC	TTA	TCA	GAG	AAA	ATT	GAT	GAC	ATT	9181
248	N	I	I	S	T	T	A	V	H	L	R	S	L	S	E	K	I	D	D	I	267
9182	TTG	CGG	TTA	ATA	GAC	GCT	CTG	GCA	AAA	GAC	TTG	GGT	AAT	CAA	GTC	TAC	GAT	GTT	GTA	TCA	9241
268	L	R	L	I	D	A	L	A	K	D	L	G	N	Q	V	Y	D	V	V	S	287
9242	CTA	ATG	GAG	GGA	TTT	GCA	TAC	GGA	GCT	GTC	CAG	CTA	CTC	GAG	CCG	TCA	GGT	ACA	TTT	GCA	9301
288	L	M	E	G	F	A	Y	G	A	V	Q	L	L	E	P	S	G	T	F	A	307
9302	GGA	GAT	TTC	TTC	GCA	TTC	AAC	CTG	CAG	GAG	CTT	AAA	GAC	ATT	CTA	ATT	GGC	CTC	CTC	CCC	9361
308	G	D	F	F	A	F	N	L	Q	E	L	K	D	I	L	I	G	L	L	P	327
9362	AAT	GAT	ATA	GCA	GAA	TCC	GTG	ACT	CAT	GCA	ATC	GCT	ACT	GTA	TTC	TCT	GGT	TTA	GAA	CAG	9421
328	N	D	I	A	E	S	V	T	H	A	I	A	T	V	F	S	G	L	E	Q	347
9422	AAT	CAA	GCA	GCT	GAG	ATG	TTG	TGT	CTG	TTG	CGT	CTG	TGG	GGT	CAC	CCA	CTG	CTT	GAG	TCC	9481
348	N	Q	A	A	E	M	L	C	L	L	R	L	W	G	H	P	L	L	E	S	367
9482	CGT	ATT	GCA	GCA	AAG	GCA	GTC	AGG	AGC	CAA	ATG	TGC	GCA	CCG	AAA	ATG	GTA	GAC	TTT	GAT	9541
368	R	I	A	A	K	A	V	R	S	Q	M	C	A	P	K	M	V	D	F	D	387
9542	ATG	ATC	CTT	CAG	GTA	CTG	TCT	TTC	TTC	AAG	GGA	ACA	ATC	ATC	AAC	GGG	TAC	AGA	AAG	AAG	9601
388	M	I	L	Q	V	L	S	F	F	K	G	T	I	I	N	G	Y	R	K	K	407
9602	AAT	GCA	GGT	GTG	TGG	CCG	CGA	GTC	AAA	GTG	GAT	ACA	ATA	TAT	GGG	AAG	GTC	ATT	GGG	CAA	9661
408	N	A	G	V	W	P	R	V	K	V	D	T	I	Y	G	K	V	I	G	Q	427
9662	CTA	CAT	GCA	GAT	TCA	GCA	GAG	ATT	TCA	CAC	GAT	ATC	ATG	TTG	AGA	GAG	TAT	AAG	AGT	TTA	9721
428	L	H	A	D	S	A	E	I	S	H	D	I	M	L	R	E	Y	K	S	L	447
9722	TCT	GCA	CTT	GAA	TTT	GAG	CCA	TGT	ATA	GAA	TAT	GAC	OCT	GTC	ACC	AAC	CTG	AGC	ATG	TTC	9781
448	S	A	L	E	F	E	P	C	I	E	Y	D	P	V	T	N	L	S	M	F	467

Fig. 3 continued

9782 CTA AAA GAC AAG GCA ATC GCA CAC CCC AAC GAT AAT TGG CTT GCC TCG TTT AGG CGG AAC 9841
 468 L K D K A I A H P N D N W L A S F R R N 487

9842 CTT CTC TCC GAA GAC CAG AAG AAA CAT GTA AAA GAA GCA ACT TCG ACT AAT CGC CTC TTG 9901
 488 L L S E D Q K K H V K E A T S T N R L L 507

9902 ATA GAG TTT TTA GAG TCA AAT GAT TTT GAT CCA TAT AAA GAG ATG GAA TAT CTG ACG ACC 9961
 508 I E F L E S N D F D P Y K E M E Y L T T 527

9962 CTT GAG TAC CTT AGA GAT GAC AAT GTG GCA GTA TCA TAC TCG CTC AAG GAG AAG GAA GTG 10021
 528 L E Y L R D D N V A V S Y S L K E K E V 547

10022 AAA GTT AAT GGA CGG ATC TTC GCT AAG CTG ACA AAG AAG TTA AGG AAC TGT CAG GTG ATG 10081
 548 K V N G R I F A K L T K K L R N C Q V M 567

10082 GCG GAA GGG ATC CTA GCC GAT CAG ATT GCA CCT TTC TTT CAG GGA AAT GGA GTC ATT CAG 10141
 568 A E G I L A D Q I A P F F Q G N G V I Q 587

10142 GAT AGC ATA TCC TTG ACC AAG AGT ATG CTA GCG ATG AGT CAA CTG TCT TTT AAC AGC AAT 10201
 588 D S I S L T K S M L A M S Q L S F N S N 607

10202 AAG AAA CGT ATC ACT GAC TGT AAA GAA AGA GTA TCT TCA AAC CGC AAT CAT GAT CCG AAA 10261
 608 K K R I T D C K E R V S S N R N H D P K 627

10262 AGC AAG AAC CGT CGG AGA GTT GCA ACC TTC ATA ACA ACT GAC CTG CAA AAG TAC TGT CTT 10321
 628 S K N R R R V A T F I T T D L Q K Y C L 647

10322 AAT TGG AGA TAT CAG ACA ATC AAA TTG TTC GCT CAT GCC ATC AAT CAG TTG ATG GGC CTA 10381
 648 N W R Y Q T I K L F A H A I N Q L M G L 667

10382 CCT CAC TTC TTC GAA TGG ATT CAC CTA AGA CTG ATG GAC ACT ACG ATG TTC GTA GGA GAC 10441
 668 P H F F E W I H L R L M D T T M F V G D 687

10442 CCT TTC AAT CCT CCA AGT GAC CCT ACT GAC TGT GAC CTC TCA AGA GTC CCT AAT GAT GAC 10501
 688 P F N P P S D P T D C D L S R V P N D D 707

10502 ATA TAT ATT GTC AGT GCC AGA GGG GGT ATC GAA GGA TTA TGC CAG AAG CTA TGG ACA ATG 10561
 708 I Y I V S A R G G I E G L C Q K L W T M 727

10562 ATC TCA ATT GCT GCA ATC CAA CTT GCT GCA GCT AGA TCG CAT TGT CGT GTT GCC TGT ATG 10621
 728 I S I A A I Q L A A A R S H C R V A C M 747

10622 GTA CAG GGT GAT AAT CAA GTA ATA GCA GTA ACG AGA GAG GTA AGA TCA GAC GAC TCT CCG 10681
 748 V Q G D N Q V I A V T R E V R S D D S P 767

10682 GAG ATG GTG TTG ACA CAG TTG CAT CAA GCC AGT GAT AAT TTC TTC AAG GAA TTA ATT CAT 10741
 768 E M V L T Q L H Q A S D N F F K E L I H 787

10742 GTC AAT CAT TTG ATT GGC CAT AAT TTG AAG GAT CGT GAA ACC ATC AGG TCA GAC ACA TTC 10801
 788 V N H L I G H N L K D R E T I R S D T F 807

10802 TTC ATA TAC AGC AAA CGA ATC TTC AAA GAT GGA GCA ATC CTC AGT CAA GTC CTC AAA AAT 10861
 808 F I Y S K R I F K D G A I L S Q V L K N 827

10862 TCA TCT AAA TTA GTG CTA GTG TCA GGT GAT CTC AGT GAA AAC ACC GTA ATG TCC TGT GCC 10921
 828 S S K L V L V S G D L S E N T V M S C A 847

Fig. 3 continued

10922 AAC ATT GCC TCT ACT GTA GCA CGG CTA TGC GAG AAC GGG CTT CCC AAA GAC TTC TGT TAC	10981
848 N I A S T V A R L C E N G L P K D F C Y	867
10982 TAT TTA AAC TAT ATA ATG AGT TGT GTG CAG ACA TAC TTT GAC TCT GAG TTC TCC ATC ACC	11041
868 Y L N Y I M S C V Q T Y F D S E F S I T	887
11042 AAC AAT TCG CAC CCC GAT CTT AAT CAG TCG TGG ATT GAG GAC ATC TCT TTT GTG CAC TCA	11101
888 N N S H P D L N Q S W I E D I S F V H S	907
11102 TAT GTT CTG ACT CCT GCC CAA TTA GGG GGA CTG AGT AAC CTT CAA TAC TCA AGG CTC TAC	11161
908 Y V L T P A Q L G G L S N L Q Y S R L Y	927
11162 ACT AGA AAT ATC GGT GAC CCG GGG ACT ACT GCT TTT GCA GAG ATC AAG CGA CTA GAA GCA	11221
928 T R N I G D P G T T A F A E I K R L E A	947
11222 GTG GGA TTA CTG AGT CCT AAC ATT ATG ACT AAT ATC TTA ACT AGG CCG CCT GGG AAT GGA	11281
948 V G L L S P N I M T N I L T R P P G N G	967
11282 GAT TGG GCC AGT CTG TGC AAC GAC CCA TAC TCT TTC AAT TTT GAG ACT GTT GCA AGC CCA	11341
968 D W A S L C N D P Y S F N F E T V A S P	987
11342 AAT ATT GTT CTT AAG AAA CAT ACG CAA AGA GTC CTA TTT GAA ACT TGT TCA AAT CCC TTA	11401
988 N I V L K K H T Q R V L F E T C S N P L	1007
11402 TTG TCT GGA GTG CAC ACA GAG GAT AAT GAG GCA GAA GAG AAG GCA TTG GCT GAA TTC TTG	11461
1008 L S G V H T E D N E A E E K A L A E F L	1027
11462 CTT AAT CAA GAG GTG ATT CAT CCC CGC GTT GCG CAT GCC ATC ATG GAG GCA AGC TCT GTA	11521
1028 L N Q E V I H P R V A H A I M E A S S V	1047
11522 GGT AGG AGA AAG CAA ATT CAA GGG CTT GTT GAC ACA ACA AAC ACC GTA ATT AAG ATT GCG	11581
1048 G R R K Q I Q G L V D T T N T V I K I A	1067
11582 CTT ACT AGG AGG CCA TTA GGC ATC AAG AGG CTG ATG CGG ATA GTC AAT TAT TCT AGC ATG	11641
1068 L T R R P L G I K R L M R I V N Y S S M	1087
11642 CAT GCA ATG CTG TTT AGA GAC GAT GTT TTT TCC TCC AGT AGA TCC AAC CAC CCC TTA GTC	11701
1088 H A M L F R D D V F S S S R S N H P E V	1107
11702 TCT TCT AAT ATG TGT TCT CTG ACA CTG GCA GAC TAT GCA CGG AAT AGA AGC TGG TCA CCT	11761
1108 S S N M C S L T L A D Y A R N R S W S P	1127
11762 TTG ACG GGA GGC AGG AAA ATA CTG GGT GTA TCT AAT CCT GAT ACG ATA GAA CTC GTA GAG	11821
1128 L T G G R K I L G V S N P D T I E L V E	1147
11822 GGT GAG ATT CTT AGT GTA AGC GGA GGG TGT ACA AGA TGT GAC AGC GGA GAT GAA CAA TTT	11881
1148 G E I L S V S G G C T R C D S G D E Q F	1167
11882 ACT TGG TTC CAT CTT CCA AGC AAT ATA GAA TTG ACC GAT GAC ACC AGC AAG AAT CCT CCG	11941
1168 T W F H L P S N I E L T D D T S K N P P	1187
11942 ATG AGG GTA CCA TAT CTC GGG TCA AAG ACA CAG GAG AGG AGA GCT GCC TCA CTT GCA AAA	12001
1188 M R V P Y L G S K T Q E R R A A S L A K	1207
12002 ATA GCT CAT ATG TCG CCA CAT GTA AAG GCT GCC CTA AGG GCA TCA TCC GTG TTG ATC TGG	12061
1208 I A H M S P H V K A A L R A S S V L I W	1227
12062 GCT TAT GGG GAT AAT GAA GTA AAT TGG ACT GCT GCT CTT ACG ATT GCA AAA TCT CGG TGT	12121
1228 A Y G D N E V N W T A A L T I A K S R C	1247

Fig. 3 continued

12122 AAT GTA AAC TTA GAG TAT CTT CGG TTA CTG TCC CCT TTA CCC ACG GCT GGG AAT CTT CAA 12181
 1248 N V N L E Y L R L L S P L P T A G N L Q 1267
 12182 CAT AGA CTA GAT GAT GGT ATA ACT CAG ATG ACA TTC ACC CCT GCA TCT CTC TAC AGG TGT 12241
 1268 H R L D D G I T Q M T F T P A S L Y R C 1287
 12242 CAC CTT ACA TTC ACA TAT CCA ATG ATT CTC AAA GGC TGT TCA CTG AAG AAG GAG TCA AAG 12301
 1288 H L T F T Y P M I L K G C S L K K E S K 1307
 12302 AGG GGA ATG TGG TTT ACC AAC AGA GTC ATG CTC TTG GGT TTA TCT CTA ATC GAA TCG ATC 12361
 1308 R G M W F T N R V M L L G L S L I E S I 1327
 12362 TTT CCA ATG ACA ACA ACC AGG ACA TAT GAT GAG ATC ACA CTG CAC CTA CAT AGT AAA TTT 12421
 1328 F P M T T T R T Y D E I T L H L H S K F 1347
 12422 AGT TGC TGT ATC AGA GAA GCA CCT GTT GCG GTT CCT TTC GAG CTA CTT GGG GTG GTA CCG 12481
 1348 S C C I R E A P V A V P F E L L G V V P 1367
 12482 GAA CTG AGG ACA GTG ACC TCA AAT AAG TTT ATG TAT GAT CCT AGC CCT GTA TCG GAG GGA 12541
 1368 E L R T V T S N K F M Y D P S P V S E G 1387
 12542 GAC TTT GCG AGA CTT GAC TTA GCT ATC TTC AAG AGT TAT GAG CTT AAT CTG GAG TCA TAT 12601
 1388 D F A R L D L A I F K S Y E L N L E S Y 1407
 12602 CCC ACG ATA GAG CTA ATG AAC ATT CTT TCA ATA TCC AGC GGG AAG TTG ATT GGC CAG TCT 12661
 1408 P T I E L M N I L S I S S G K L I G Q S 1427
 12662 GTG GTT TCT TAT GAT GAA GAT ACC TCC ATA AAG AAT GAC GCC ATA ATA GTG TAT GAC AAT 12721
 1428 V V S Y D E D T S I K N D A I I V Y D N 1447
 12722 ACC CGA AAT TGG ATC AGT GAA GCT CAG AAT TCA GAT GTG GTC CGC CTA TTT GAA TAT GCA 12781
 1448 T R N W I S E A Q N S D V V R L F E Y A 1467
 12782 GCA CTT GAA GTG CTC CTC GAC TGT TCT TAC CAA CTC TAT TAC CTG AGA GTA AGA GGC CTA 12841
 1468 A L E V L L D C S Y Q L Y Y L R V R G L 1487
 12842 GAC AAT ATT GTC TTA TAT ATG GGT GAT TTA TAC AAG AAT ATG CCA GGA ATT CTA CTT TCC 12901
 1488 D N I V L Y M G D L Y K N M P G I L L S 1507
 12902 AAC ATT GCA GCT ACA ATA TCT CAT CCC GTC ATT CAT TCA AGG TTA CAT GCA GTG GGC CTG 12961
 1508 N I A A T I S H P V I H S R L H A V G L 1527
 12962 GTC AAC CAT GAC GGA TCA CAC CAA CTT GCA GAT ACG GAT TTT ATC GAA ATG TCT GCA AAA 13021
 1528 V N H D G S H Q L A D T D F I E M S A K 1547
 13022 CTA TTA GTA TCT TGC ACC CGA CGT GTG ATC TCC GGC TTA TAT TCA GGA AAT AAG TAT GAT 13081
 1548 L L V S C T R R V I S G L Y S G N K Y D 1567
 13082 CTG CTG TTC CCA TCT GTC TTA GAT GAT AAC CTG AAT GAG AAG ATG CTT CAG CTG ATA TCC 13141
 1568 L L F P S V L D D N L N E K M L Q L I S 1587
 13142 CGG TTA TGC TGT CTG TAC ACG GTA CTC TTT GCT ACA ACA AGA GAA ATC CCG AAA ATA AGA 13201
 1588 R L C C L Y T V L F A T T R E I P K I R 1607
 13202 GGC TTA ACT GCA GAA GAG AAA TGT TCA ATA CTC ACT GAG TAT TTA CTG TCG GAT GCT GTG 13261
 1608 G L T A E E K C S I L T E Y L L S D A V 1627

14/22

Fig. 3 continued

13262	AAA CCA TTA CTT AGC CCC GAT CAA GTG AGC TCT ATC ATG TCT CCT AAC ATA ATT ACA TTC	13321
1628	K P L L S P D Q V S S I M S P N I I T F	1647
13322	CCA GCT AAT CTG TAC TAC ATG TCT CGG AAG AGC CTC AAT TTG ATC AGG GAA AGG GAG GAC	13381
1648	P A N L Y Y M S R K S L N L I R E R E D	1667
13382	AGG GAT ACT ATC CTG GCG TTG TTG TTC CCC CAA GAG CCA TTA TTA GAG TTC CCT TCT GTG	13441
1668	R D T I L A L L F P Q E P L L E F P S V	1687
13442	CAA GAT ATT GGT GCT CGA GTG AAA GAT CCA TTC ACC CGA CAA CCT GCG GCA TTT TTG CAA	13501
1688	Q D I G A R V K D P F T R Q P A A F L Q	1707
13502	GAG TTA GAT TTG AGT GCT CCA GCA AGG TAT GAC GCA TTC ACA CTT AGT CAG ATT CAT CCT	13561
1708	E L D L S A P A R Y D A F T L S Q I H P	1727
13562	GAA CTC ACA TCT CCA AAT CCG GAG GAA GAC TAC TTA GTA CGA TAC TTG TTC AGA GGG ATA	13621
1728	E L T S P N P E E D Y L V R Y L F R G I	1747
13622	GGG ACT GCA TCT TCC TCT TGG TAT AAG GCA TCT CAT CTC CTT TCT GTA CCC GAG GTA AGA	13681
1748	G T A S S S W Y K A S H L L S V P E V R	1767
13682	TGT GCA AGA CAC GGC AAC TCC TTA TAC TTA GCT GAA GGG AGC GGA GCC ATC ATG AGT CTT	13741
1768	C A R H G N S L Y L A E G S G A I M S L	1787
13742	CTC GAA CTG CAT GTA CCA CAT GAA ACT ATC TAT TAC AAT ACG CTC TTT TCA AAT GAG ATG	13801
1788	L E L H V P H E T I Y Y N T L F S N E M	1807
13802	AAC CCC CCG CAA CGA CAT TTC GGG CCG ACC CCA ACT CAG TTT TTG AAT TCG GTT GTT TAT	13861
1808	N P P Q R H F G P T P T Q F L N S V V Y	1827
13862	AGG AAT CTA CAG GCG GAG GTA ACA TGC AAA GAT GGA TTT GTC CAA GAG TTC CGT CCA TTA	13921
1828	R N L Q A E V T C K D G F V Q E F R P L	1847
13922	TGG AGA GAA AAT ACA GAG GAA AGT GAC CTG ACC TCA GAT AAA GCA GTG GGG TAT ATT ACA	13981
1848	W R E N T E E S D L T S D K A V G Y I T	1867
13982	TCT GCA GTG CCC TAC AGA TCT GTA TCA TTG CTG CAT TGT GAC ATT GAA ATT CCT CCA GGG	14041
1868	S A V P Y R S V S L L H C D I E I P P G	1887
14042	TCC AAT CAA AGC TTA CTA GAT CAA CTA GCT ATC AAT TTA TCT CTG ATT GCC ATG CAT TCT	14101
1888	S N Q S L L D Q L A I N L S L I A M H S	1907
14102	GTA AGG GAG GGC GGG GTA GTA ATC ATC AAA GTG TTG TAT GCA ATG GGA TAC TAC TTT CAT	14161
1908	V R E G G V V I I K V L Y A M G Y Y F H	1927
14162	CTA CTC ATG AAC TTG TTT GCT CCG TGT TCC ACA AAA GGA TAT ATT CTC TCT AAT GGT TAT	14221
1928	L L M N L F A P C S T K G Y I L S N G Y	1947
14222	GCA TGT CGA GGA GAT ATG GAG TGT TAC CTG GTA TTT GTC ATG GGT TAC CTG GGC GGG CCT	14281
1948	A C R G D M E C Y L V F V M G Y L G G P	1967
14282	ACA TTT GTA CAT GAG GTG GTG AGG ATG GCA AAA ACT CTG GTG CAG CGG CAC GGT ACG CTC	14341
1968	T F V H E V V R M A K T L V Q R H G T L	1987
14342	TTG TCT AAA TCA GAT GAG ATC ACA CTG ACC AGG TTA TTC ACC TCA CAG CGG CAG CGT GTG	14401
1988	L S K S D E I T L T R L F T S Q R Q R V	2007
14402	ACA GAC ATC CTA TCC AGT CCT TTA CCA AGA TTA ATA AAG TAC TTG AGG AAG AAT ATT GAC	14461
2008	T D I L S S P L P R L I K Y L R K N I D	2027

Fig. 3 continued

14462 ACT GCG CTG ATT GAA GCC GGG GGA CAG CCC GTC CGT CCA TTC TGT GCG GAG AGT CTG GTG 14521
 2028 T A L I E A G G Q P V R P F C A E S L V 2047

14522 AGC ACG CTA GCG AAC ATA ACT CAG ATA ACC CAG ATT ATC GCT AGT CAC ATT GAC ACA GTT 14581
 2048 S T L A N I T Q I T Q I I A S H I D T V 2067

14582 ATC CGG TCT GTG ATA TAT ATG GAA GCT GAG GGT GAT CTC GCT GAC ACA GTA TTT CTA TTT 14641
 2068 I R S V I Y M E A E G D L A D T V F L F 2087

14642 ACC CCT TAC AAT CTC TCT ACT GAC GGG AAA AAG AGG ACA TCA CTT ATA CAG TGC ACG AGA 14701
 2088 T P Y N L S T D G K K R T S L I Q C T R 2107

14702 CAG ATC CTA GAG GTT ACA ATA CTA GGT CTT AGA GTC GAA AAT CTC AAT AAA ATA GGC GAT 14761
 2108 Q I L G V T I L G L R V E N L N K I G D 2127

14762 ATA ATC AGC CTA GTG CTT AAA GGC ATG ATC TCC ATG GAG GAC CTT ATC CCA CTA AGG ACA 14821
 2128 I I S L V L K G M I S M E D L I P L R T 2147

14822 TAC TTG AAG CAT AGT ACC TGC CCT AAA TAT TTG AAG GCT GTC CTA GGT ATT ACC AAA CTC 14881
 2148 Y L K H S T C P K Y L K A V L G I T K L 2167

14882 AAA GAA ATG TTT ACA GAC ACT TCT GTA TTG TAC TTG ACT CGT GCT CAA CAA AAA TTC TAC 14941
 2168 K E M F T D T S V L Y L T R A Q Q K F Y 2187

14942 ATG AAA ACT ATA GGC AAT GCA GTC AAA GGA TAT TAC AGT AAC TGT GAC TCT TAA cgaaaatc 15003
 2188 M K T I G N A V K G Y Y S N C D S * 2205

15004 acatattaataggctccttttttggccaattgtattcttgttgatttaatcatattatgtagaaaaaagttgaaccctg 15083

15084 actccttaggactcgaattcgaactcaaataaatgtcttaaaaaaagggttgcgacaaattattcttgagtgtagtctcgt 15163

15164 cattcaccaaactctttgtttggt 15186




Fig. 4A

17/22

Construction of genome-length cDNA clones of
NDV "LaSota" E13-1 in pOLTV535




Fig. 4B




Fig. 4B continued

20/22




Fig. 5A




Fig. 5B

<i>Mobillivirus</i>	CDV	1:	-----	*	20	*	40	*	60	
	MeV	2:	-----							
	RDV	3:	-----							
	bPIV3	4:	-----							
<i>Paramyxovirus</i>	hPIV3	5:	-----							
	Sev	6:	-----							8
	NDV	7:	TTAGAAAAAGTTGAACCTGACTCCTTAGGACTCGAATTCGAACTCAATAAATGCTCTTAAA							64
	hPIV2	8:	-----							
<i>Rubulavirus</i>	MuV	9:	-----							
	SV41	10:	-----							
	SV5	11:	-----							
<i>Mobillivirus</i>	CDV	1:	-----	*	80	*	100	*	120	
	MeV	2:	-----							54
	RDV	3:	-----							51
	bPIV3	4:	-----							51
<i>Paramyxovirus</i>	hPIV3	5:	-----							55
	Sev	6:	-----							55
	NDV	7:	AAACTTACAAGAAGACAAGAAAAATTAAAGGATACATATCTCTTAAACTCTTGTCTGGT							68
	hPIV2	8:	AAAGTTGCGCACAAATTATCTTGTAGTGTCTCGTCTCATTCCACCAATCTTGTCTGGT							124
<i>Rubulavirus</i>	MuV	9:	-----							32
	SV41	10:	-----							35
	SV5	11:	-----							30
			-----							42
			-----							TGGT

Fig. 6

200007339-5
IPOS

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/45, 15/86, 7/00, A61K 48/00, C12Q 1/70	A1	(11) International Publication Number: WO 99/66045 (43) International Publication Date: 23 December 1999 (23.12.99)
(21) International Application Number: PCT/NL99/00377 (22) International Filing Date: 17 June 1999 (17.06.99) (30) Priority Data: ① 98202054.7 ✓ 19 June 1998 (19.06.98) ✓ EP ✓ (71) Applicant (for all designated States except US): STICHTING DIENST LANDBOUWKUNDIG ONDERZOEK [NL/NL]; Bornsesteeg 53, NL-6708 PD Wageningen (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): PEETERS, Bernardus, Petrus, Hubertus [NL/NL]; Golfpark 18, NL-8241 AB Lelystad (NL); DE LEEUW, Olav, Sven [NL/NL]; Johan Westerveldstraat 4, NL-1318 CC Almere (NL); KOCH, Guus [NL/NL]; Wijkbaan 12, NL-8222 AG Lelystad (NL). ④ GIELKENS, Arnold, Leonard, Josef [NL/NL]; Boeier 04-76, NL-8242 CL Lelystad (NL). (74) Agent: OTTEVANGERS, S., U.; Vereenigde Octrooibureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. 1 PD 1 appl. 4 Invs.	
(54) Title: NEWCASTLE DISEASE VIRUS INFECTIOUS CLONES, VACCINES AND DIAGNOSTIC ASSAYS ✓ (57) Abstract The invention relates to the process for generating infectious Newcastle disease virus (NDV) entirely from cloned full-length cDNA and to the use of vaccines and diagnostic assays generated with and derived from said process. The process offers the possibility to modify the NDV genome by means of genetic modification and allows the introduction of mutations, deletions, and/or insertions. The process can be used to modify the virulence of NDV, thereby generating new attenuated live vaccines with enhanced properties. The process can be used to modify the antigenic make-up of NDV, thus allowing the generation of live NDV marker vaccines which can be serologically distinguished from NDV field strains.		